

---



---

**Errata**


---



---

**Erratum: Spherical proton emitters**  
**[Phys. Rev. C 56, 1762 (1997)]**

Sven Åberg, Paul B. Semmes, and Witold Nazarewicz

[S0556-2813(98)01811-1]

PACS number(s): 23.50.+z, 21.10.Jx, 21.10.Tg, 24.10.Eq, 99.10.+g

In Table II, the columns marked WKB and WKB1 are reversed. This means that the Wentzel-Kramers-Brillouin (WKB) calculations with the usual approximation for the normalization constant [see Eq. (25)] agree quite well with the corresponding distorted-wave Born approximation (DWBA) calculations; WKB calculations with the more sophisticated treatment of the normalization constant [Eq. (24)] in fact agree less well with the DWBA results.

Also, the entries for the WKB half-lives for the  $0h_{11/2}$  orbital in  $^{161}_{75}\text{Re}_{86}$  should be 93 ms and 130 ms instead of 100 ms and 150 ms as listed. The original numbers resulted from taking an outdated  $Q_p$  value.

Finally, a clarification: The form of the spin-orbit potential used was

$$V_{l\cdot s}(r) = V_{so} \frac{1}{r} \frac{df}{dr} \mathbf{L} \cdot \boldsymbol{\sigma},$$

where  $f(r) = 1/1 + \exp(r - R/a)$ . Our ansatz relating  $V_{so}$  and  $V_0$  is  $V_{so} = 0.2 V_0 \text{ fm}^2$ .

We thank Patrick Talou for his probing questions.