

**Erratum: Covariant confinement model for the study of the properties of light mesons  
[Phys. Rev. C 60, 025202 (1999)]**

L. S. Celenza, Bo Huang, Huangsheng Wang, and C. M. Shakin  
(Published 24 August 1999)

[S0556-2813(99)04710-X]

PACS number(s): 12.39.Ki, 14.40.-n, 14.65.Bt, 99.10.+g

The sentence introducing Eq. (6.7) and the equation should read:

Further

$$\Gamma_{S,ab}^{-+}(P^0,k) = 1 + \frac{1}{(2\pi)^2} \int k'^2 dk' \int_{-1}^1 dx \frac{\tilde{A}_1(k,k',x)}{E_a(\vec{k}') E_b(\vec{k}')} \frac{1}{B(\vec{k})} \frac{\Gamma_{S,ab}^{-+}(P^0,k') V^C(\vec{k} - \vec{k}')} {P^0 + E_a(\vec{k}') + E_b(\vec{k}')} , \quad (6.7)$$

where  $\tilde{A}_1(k,k',x)$  is obtained from  $A_1(k,k',x)$  by the replacements  $E_a(\vec{k}) \rightarrow -E_a(\vec{k})$ ,  $E_b(\vec{k}) \rightarrow -E_b(\vec{k})$ ,  $E_a(\vec{k}') \rightarrow -E_a(\vec{k}')$ , and  $E_b(\vec{k}') \rightarrow -E_b(\vec{k}')$ , with the result that  $\tilde{A}_1(k,k',x) = A_1(k,k',x)$ .