

Time dependence of strange baryon freeze-out in relativistic heavy ion collisions

R. Bellwied,¹ H. Caines,² and T. J. Humanic²

¹*Department of Physics and Astronomy, Wayne State University, Detroit, Michigan 48202*

²*Department of Physics, The Ohio State University, Columbus, Ohio 43210*

(Received 3 April 2000; published 16 October 2000)

We investigate chemical and thermal freeze-out time dependencies of strange baryon production for CERN SPS heavy ion collisions in the framework of a dynamical hadronic transport code. We show that the Λ yield changes considerably after hadronization in the case of Pb+Pb collisions, whereas for smaller system sizes (e.g., S+S) the direct particle production dominates over production from inelastic rescattering. Chemical freeze-out times for strange baryons in Pb+Pb are smaller than for nonstrange baryons, but they are still sufficiently long for hadronic rescattering to contribute significantly to the final Λ yield.

PACS number(s): 25.75.-q

Recent model calculations, which were successfully applied to CERN SPS data, implied that most measured particle distributions in relativistic heavy ion collisions can be described with statistical models [1]. The underlying kinematics requires at least thermal equilibration at freeze-out. All kinetic spectra, such as the transverse momentum and the rapidity distributions, can be explained with a single freeze-out temperature, if one assumes a certain expansion velocity after hadronization. Attempts to describe not only the kinetic spectra but also the particle abundances and particle ratios with a common temperature were successful, but led to a different temperature. Whereas the slopes of all momentum spectra yield a T around 130 MeV after inclusion of the expansion velocity, the particle ratios lead to a T of about 180 MeV for a common chemical potential. Two questions arise: is the chemical freeze-out decoupled from the thermal freeze-out and is the system chemically equilibrated at chemical freeze-out? In a systematic study of measured particle ratios and spectra in the framework of a thermal model, Heinz has shown that for all CERN and AGS data the chemical freeze-out seems to be decoupled from the thermal freeze-out [2].

Rafelksi has recently postulated that, on the basis of strange particle ratios measured in Pb+Pb at CERN, one has to assume a phase transition from QGP to hadron gas, and that at the time of hadronization the system is not in chemical equilibrium [3]. To describe the kinetic spectra and particle yields this model requires that the chemical composition of the fireball remains unchanged between hadronization and thermal freeze-out. That means all interactions during this time interval have to be elastic. This model is in agreement with the fact that all systems from e^+e^- collisions up to AA collisions seem to yield the same chemical freeze-out temperature [1]. Thus, the shapes of the particle spectra might change due to rescattering, but the particle abundances and ratios remain constant from hadronization on, a theory which is commonly referred to as “sudden hadronization.” This process explicitly prohibits hadrochemical equilibration through rescattering. Only elastic processes can be employed to explain the difference between the temperature calculated from particle ratios and the temperature calculated from kinetic particle spectra. Thus, the ratios effectively reflect the temperature at hadronization, whereas the kinetic freeze-out

temperature describes the actual freeze-out (last elastic final state interaction). The fact that the hadronization temperature is about 180 MeV, close to the critical temperature, can be viewed as an indicator that the system actually crossed a phase transition. However, the fact that all inelastic scattering processes cease at hadronization time is a very strong constraint. Hadronic population ratios are, in this theory, the result of the hadronization mechanism and not caused by interactions during the hadronic rescattering phase. The main argument employed is that the relaxation times of all relevant hadronic channels is well above $\tau=3$ fm/c at which time the temperature of the system has dropped below $T=185$ MeV which leads to a small probability of chemical equilibration after hadronization. Also based on results of Hanbury Brown–Twiss (HBT) measurements, Stock argues that the time between hadronization and thermal freeze-out might simply be too short to develop a significant inelastic cross section contribution [4]. The required time interval for hadronic processes to adjust the strangeness content is long and the chemical rates are small since the production of pairs of strange hadrons carries a large energy penalty factor. In Stock’s scenario it takes upward of 3 fm/c to equilibrate strangeness in the hadronic phase, a time span that, he postulates, is not available owing to the rapid expansion prevailing at hadronization time.

Surprisingly, a partonic cascade approach, which is very different from the model described above, yields results similar to a thermal sudden hadronization theory. Geiger and Ellis [5] have shown that at large incident energies the hadronization process can determine the final particle ratios based on its combined nonperturbative mechanisms and it can even lead to direct chemical equilibrium. The final multihadronic state materializes into maximal entropy (equilibrium) straight out of the partonic phase. Although certain remnants of the initial particle structure functions remain, the hadron yield near midrapidity stems mostly from the initial parton cascade processes. In this theory, the partonic phase exhibits $\epsilon \geq 2$ GeV/fm³ until about 2 fm/c. Hadronization occurs after a formation time of about 1 fm/c and a mixed phase of hadrons and partons ends at $\tau \approx 20$ fm/c, when ϵ is less than 0.2 GeV/fm³. Although the concept of sudden hadronization is lost in this theory, due to the required long mixed phase between partons and hadrons, the results are comparable to a thermal model without inelastic final state




FIG. 1. Particle abundances as a function of time between hadronization and kinematic freeze-out.

interactions. In addition there are nonequilibrium processes which may cause a rapid hadronization of the QGP [6]. In particular nucleation theories that assume supercooling of a rapidly expanding quark gluon plasma show a hadroproduction rate comparable to thermal hadronization theories [7]. Spieles *et al.* nicely documented the similarities and differences between a static thermal hadronic source and nonequilibrium hadronization of a QGP droplet [8]. While the parton models require a mixed phase for the proper kinematic expansion of the system, they do not require any final state interactions to describe the measured particle abundances.

Kapusta and Mekjian [9] derived estimates for the dynamical equilibration (relaxation) times of quark flavors in a quark gluon gas and deduced predictions for several equilibrium abundance ratios. These ratios are in good agreement with the data, which led Stock [4] to postulate that the equilibrated ratios formed in prehadronization reactions do not change during the final state interactions.

The main motivation for this paper is the fact that most dynamic transport codes, which in the past were successfully applied to describe data from CERN and the AGS, behave very differently from either the parton cascade or the thermal sudden hadronization approach. These models employ measured inelastic scattering cross sections for different particle species to describe the hadronic transport from hadronization to kinetic freeze-out. Many of the relevant particle species are susceptible to number changing interactions, in particular inelastic meson interactions lead to sizable contributions to the hyperon production cross section. Both strangeness creation interactions (e.g., $\pi N \rightarrow K \Lambda$) and strangeness exchange interactions (e.g., $KN \rightarrow \pi \Lambda$) should affect strange baryon production.

Pratt and Haglin have recently pointed out that number changing cross sections after hadronization seem to be relevant for the description of the pion abundance at CERN [10]. They have shown that hadrons interact several times before the freeze-out temperatures are reached if one assumes that binary modeling starts at $T=160$ MeV in a chemically and thermally equilibrated system. It is interesting to note that the chemical equilibration times for strange and non strange quarks shown by Stock and Pratt and Haglin are very comparable. An independent study by Humanic, using an early version of the transport code which is the basis of this paper, shows that pions interact on average ten times between hadronization and freeze-out [11]. The time from hadronization to freeze-out is quite long (about 15 fm/




FIG. 2. Time dependence of channel contributions to the Λ cross section in (a) Pb+Pb and (b) S+S.




FIG. 3. Proof of detailed balance in a nonexpanding system of fixed maximum radius ($r=10$ fm).

c) in particular for the lightest mesons. If these transport models are correct then we should be able to simulate a quantitative dynamic evolution of all particle abundances through the rescattering phase.

HADRONIC PARTICLE RATIO SIMULATIONS

In the following we attempt to prove that in particular the abundance of the singly strange baryon, the Λ , is affected by a series of inelastic rescattering processes by the time thermal freeze-out is accomplished. We have employed the dy-




FIG. 4. Relation between chemical freeze-out time and kinetic freeze-out time for measurable Λ particles.




FIG. 5. Chemical freeze-out time for measurable Λ particles.

namic transport code described in Ref. [11] to model the initial state of the system at hadronization and the subsequent hadronic rescattering to freeze-out. This code has been shown to well represent the features and dynamic dependences of the transverse mass spectra and HBT observables for CERN SPS Pb+Pb data [11]. In addition to the already implemented cross sections for pions, kaons, nucleons and their associated resonances [12], we have augmented the code to include elastic and inelastic rescattering of Λ baryons. Certainly the application of vacuum cross sections has to be considered an approximation at the relevant particle and energy densities. Attempts to parametrize the density dependence of measured interaction cross sections are underway [13], but for the time being most event generators, in particular cascade programs, quite successfully apply free measured cross sections to describe relativistic heavy ion measurements. We expect the inelastic and elastic cross sections to increase at higher densities, in which case our calculations of the effect of rescattering between hadronization and thermal freeze-out should be considered a lower limit.

The initial conditions in our model are described by a common temperature and spatial extension. All particles hadronize at a proper time of 1 fm/c. A Bjorken-type geometry was used to simulate the dynamic evolution of the fireball from hadronization time. No initial radial flow is needed for the calculations to agree with data [11]. Calculations are carried out assuming initial parameter values and multiplicities for each type of particle. In the last stage of the calculation, the freeze-out and decay momenta and space-times are used to produce single-particle and two-particle observables such as pion, kaon, nucleon, and lambda multiplicities, transverse momentum and rapidity distributions, and two-boson correlation functions. The values of the initial parameters of the calculation are constrained to give observables which agree with available measured hadronic observables. For the




FIG. 6. (a) Kinetic freeze-out time for the different components of the Λ spectrum. (b) Comparison between the kinetic freeze-out time distributions of measurable Λ 's, π 's, and protons.

calculations studied in the present work, measured observables were obtained from the NA44 [14,15] and NA49 [16–18] experiments. Initial resonance multiplicity fractions are taken from the HELIOS experiment [19].

Figure 1 shows the effect of rescattering on all four relevant particle abundances (π , K , nucleons, and Λ). In the case of the nucleons, the nucleon resonance production and subsequent decay leads to no change in the number of nucleons and is thus considered an elastic process, whereas in the pion case, generation from resonance decay is considered a number changing process (e.g., $N\pi \rightarrow \Delta\pi \rightarrow N\pi\pi$). The same is true for vector meson production (ρ, ϕ, ω) and decay into pions. Figure 1 shows that the nucleon abundance is close to constant, but the meson and strange baryon abundances display a steady increase. One of the large contributing elastic reactions is $\pi\pi \rightarrow \rho \rightarrow \pi\pi$ which is fast and keeps the number of ρ 's in equilibrium but does not change the pion number. Reactions that change the overall pion number such as $\pi\pi \rightarrow \eta' \rightarrow \pi\pi\pi$ are generally slower and thus do not have a large effect on the chemical state formed at hadronization. In addition reactions that conserve the net number of strange quarks occur rapidly, e.g., $KN \rightarrow \Lambda\pi$, but reactions that change the number of strange quarks, e.g., $\pi N \rightarrow K\Lambda$, are slow as they require a strange and an antistrange hadron to either interact or to be produced jointly.

Figure 2 shows the relative contributions of all relevant reaction channels to the Λ abundance as a function of time for the Pb+Pb and the S+S systems at the SPS. In the Pb case the direct Λ production at hadronization yields only around 50% (≈ 30 Λ 's) of the total yield at kinetic freeze-out. The additional enhancement can be attributed to Λ production well after hadronization. In comparison, in the smaller S+S system the rescattering contributes only about 10% of the total Λ yield. It should be noted that, in either

case, the contribution from the strangeness creating, endothermic reaction ($\pi+N \rightarrow K+\Lambda$) is considerably higher than the one from the simple exothermic strangeness exchange reaction ($K+N \rightarrow \pi+\Lambda$). We attribute this effect to the large pion density in the rescattering volume.

The plots also show that Λ production and annihilation through inelastic rescattering stops after a certain time (chemical freeze-out). It turns out, though, that this is not due to our model reaching strangeness chemical equilibrium but rather based on the rapid expansion and the resulting drop in particle density. After around 30 fm/c the particle density is simply too small to support further inelastic scatterings which produce strange baryons. To prove that our model properly invokes detailed balance in our cross sections we chose a fixed maximum radius of 10 fm and plotted the Λ production and annihilation channels as a function of time. Figure 3 shows that, for the case of a nonexpanding system, the total Λ yield becomes constant as the annihilation yield equals that of the production at long times. Thus, chemical equilibration is possible if the expansion velocity is sufficiently small. However, our calculations agree with other thermal model calculations in predicting that the Pb+Pb system at CERN SPS energies does not reach chemical equilibrium before freeze-out. Based on Fig. 1 we can conclude that all particle ratios that include the Λ yield will undergo significant changes during rescattering, a fact which is incompatible with the notion of sudden hadronization. If we force sudden hadronization in our code by simply turning off all inelastic rescattering modes, our model calculations lead to a much smaller HBT radius in both the $\pi\pi$ and KK channels compared to the actually measured radii.

Figure 4 shows the relation between kinetic and chemical freeze-out time for every Λ measurable in the Pb+Pb system after kinetic freeze-out. It is apparent that only about half of




FIG. 7. Transverse mass spectrum for Λ from our calculations. The extracted slope parameter is seen to agree with data (NA49).

the emitted Λ 's show a chemical freeze-out time consistent with the hadronization time, which is required for sudden hadronization ($t_{\text{chem}}=0, t_{\text{kin}}=\text{any}$). Many Λ 's (about 25%) are actually produced in their final rescattering step ($t_{\text{chem}}=t_{\text{kin}}$). The one-dimensional projection of the chemical freeze-out time distribution for the Λ 's, as shown in Fig. 5, reveals that the chemical freeze-out occurs fast but it is not "sudden." 90% of the Λ 's are chemically frozen out after 5 fm/c, but of those about 40% freeze-out after hadronization. Details of the according kinetic freeze-out times are displayed in Fig. 6. Figure 6(a) shows the difference in kinetic freeze-out time between the direct and the produced Λ component. The production after rescattering significantly enhances the average freeze-out time and alters the shape of the distribution. But the Λ freeze-out is still peaked significantly earlier than the pion and proton freeze-out in our model [11] as seen in Fig. 6(b). The main reason is that the inelastic and elastic cross sections for Λ induced reactions are much smaller than the proton or pion interaction cross sections. This trend will continue for multistrange baryons and it should lead to a very early decoupling of the Ω from the fireball [20]. For multistrange baryons the notion of sudden hadronization seems thus more sensible.

Figure 7 shows the transverse mass spectrum of the frozen out Λ 's. The model is in perfect agreement with the data measured by NA49 [21], as it is for the pions and nucleons.

CONCLUSIONS

Based on our calculations we conclude that neither the Λ yield nor any particle ratio including Λ 's are suitable thermometers for the hadronization temperature. These measurements may still be indirect QGP indicators in any transport code description, though, simply through determination of

the required initial hadronization conditions (initial temperature and energy density). This is a departure from the original wisdom that only leptonic probes can act as real indicators of the initial conditions.

We have shown that the Λ abundance is strongly affected by final state interactions and is not reproducible under the strict assumption of sudden hadronization, which requires no inelastic rescattering after hadronization. This does not preclude a very short mixed phase and there is no attempt in this paper to describe the prehadronization phase or hadronization itself. Our model simply estimates the considerable contributions between hadronization and thermal freeze-out. The probability of inelastic scattering is sufficiently large that even during the rather short time lapse between hadronization and thermal freeze-out the yields change considerably. Thus, conclusions drawn from the strange baryon ratios at CERN concerning a QGP phase transition are probably an oversimplification regarding the actual interactions inside the fireball.

Our simulations agree with other thermal models that the chemical properties are frozen well before the kinetic spectra but we also demonstrate that hadronization and chemical freeze-out mostly do not coincide for strange baryons produced in Pb+Pb. Thus the hadronization should occur above $T=180$ MeV which was deduced from the ratios as the temperature for chemical freeze-out. Indeed, in our calculation the system hadronizes at 213 MeV, well above the critical temperature. It is also evident that both, the Λ chemical and thermal freeze-out, occur much earlier than the proton and pion freeze-out. Recent detailed simulations of multistrange baryon ratios and spectra, seem to indicate that this early decoupling follows a trend as a function of the strangeness content. This effect can be attributed to differences in the rescattering probabilities [22]. In particular the apparent lack of flow in the Ω transverse momentum spectrum could be correlated with the very high Ω particle ratios [23]. Both point at a uniquely different production mechanism for the Ω . In Dumitru's calculations the Ω 's undergo on average two collisions after hadronization, whereas the number increases to 3.5 and 5 for Ξ and Λ , respectively [22]. This early decoupling of the Ω led van Hecke *et al.* to postulate that this is the reason for the lower emission temperatures for Ω 's as measured at the SPS [20]. Our calculation indicates that even the Λ thermally freezes out after a few fm/c, due to the rather small elastic scattering cross section during the rescattering phase. It is our goal to extend our transport code to include the multiply strange baryons to determine more quantitatively the dynamic differences between strange and multistrange baryons.

The complete measurement of strange particle production at RHIC, from the kaon to the Ω , will be an important exercise. Our SPS study shows that final state interactions have to taken into account in order to properly describe singly-strange particle production. Thus, kaon and Λ measurements alone might not lead to conclusive proof of the formation of a QGP. However, plans for detailed measurements of strangeness production up to and including the Ω are well underway at RHIC and we are looking forward to this exciting new era in relativistic heavy ion physics.

- [1] F. Becattini, J. Phys. G **23**, 1933 (1997).
- [2] U. Heinz, Nucl. Phys. **A661**, 140c (1999); J. Phys. G **25**, 263 (1999).
- [3] J. Rafelski, J. Phys. G **25**, 295 (1999).
- [4] H. Stock, Phys. Lett. B **456**, 277 (1999).
- [5] K. Geiger and J. Ellis, Phys. Rev. D **54**, 949 (1996).
- [6] L. Csernai and I. Mishustin, Phys. Rev. Lett. **74**, 5005 (1995).
- [7] E. Zabrodin *et al.*, Phys. Lett. B **423**, 373 (1998).
- [8] C. Spieles *et al.*, Eur. Phys. J. C **2**, 351 (1998).
- [9] J. Kapusta and A. Z. Mekjian, Phys. Rev. D **33**, 1304 (1986).
- [10] S. Pratt and K. Haglin, Phys. Rev. C **59**, 3304 (1999).
- [11] T. Humanic, Phys. Rev. C **57**, 866 (1998).
- [12] M. Prakash *et al.*, Phys. Rep. **227**, 321 (1993).
- [13] D. Zschesche, P. Papazoglou, S. Schramm, J. Schaffner-Bielich, H. Stöcker, and W. Greiner, nucl-th/0001055; M. Hofmann *et al.*, Phys. Lett. B **478**, 161 (2000).
- [14] I. G. Bearden, Phys. Rev. Lett. **78**, 2080 (1997).
- [15] A. Franz *et al.*, Nucl. Phys. **A610**, 240c (1996).
- [16] P. G. Jones *et al.*, Nucl. Phys. **A610**, 188c (1996).
- [17] K. Kadija *et al.*, Nucl. Phys. **A610**, 248c (1996).
- [18] H. Appelshauser, Ph.D. thesis, University of Frankfurt, 1996.
- [19] U. Goerlach *et al.*, Nucl. Phys. **A544**, 109c (1992).
- [20] H. van Hecke *et al.*, Phys. Rev. Lett. **81**, 5764 (1998).
- [21] S. Margetis *et al.*, NA49 Collaboration, J. Phys. G **25**, 189 (1999).
- [22] A. Dumitru *et al.*, Phys. Lett. B **460**, 411 (1999).
- [23] P. Csizmadia and Péter Lévai, Phys. Rev. C **61**, 031903(R) (2000).