

Form factors and photoproduction amplitudes

R. M. Davidson*

Department of Physics, Applied Science and Astronomy, Rensselaer Polytechnic Institute, Troy, New York 12180-3590

Ron Workman†

Center for Nuclear Studies and Department of Physics, The George Washington University, Washington, D.C. 20052

(Received 13 July 2000; published 22 January 2001)

We examine the use of phenomenological form factors in tree level amplitudes for meson photoproduction. Two common recipes are shown to be fundamentally incorrect. An alternate form consistent with gauge invariance and crossing symmetry is proposed.

DOI: 10.1103/PhysRevC.63.025210

PACS number(s): 25.20.Lj, 13.60.Le, 11.40.-q, 11.80.Cr

I. INTRODUCTION

Studies of electromagnetic interactions with extended particles, such as mesons and baryons, often involve the use of (off-shell) form factors to account for internal structure not explicit in models, or to regularize quantities which would otherwise be divergent. The general structures of the πNN [1] and γNN [2] three-point vertices allowed based on the symmetries of the strong and electromagnetic interactions have been known for decades. Of particular importance for electromagnetic interactions is gauge invariance as expressed by the Ward-Takahashi identities. [3] There is vast literature on the subject of form factors in meson production ranging from attempts to satisfy the half-off-shell Ward-Takahashi identities while at the same time using as much on-shell information as possible [4,5], to explicit model calculations [6,7] which test the various prescriptions. As a rule, these model calculations demonstrate that there are kinematical regions where the prescriptions can give drastically different results than the exact model calculations.

While the most consistent way to approach this problem is through the use of field-theoretic models, it is not clear which field-theory approach to use in the resonance region. Although chiral perturbation theory can be used near threshold [8], it is expected to converge slowly, or not at all, at higher energies. Thus, many recent fits to meson photoproduction data have been less ambitious, and have assumed that a phenomenological approach would be adequate for the extraction of resonance contributions. These fits have generally used tree-level diagrams which require a high-energy cutoff. While the functional form of these form factors has usually been chosen on the basis of convenience and simplicity, the resulting amplitudes should obey the constraints imposed by gauge invariance, crossing and unitarity. Most efforts have been directed toward the maintenance of gauge invariance. Crossing and unitarity are often ignored in modifications applied to tree-level diagrams, whereas unitarity is generally built into dynamical models.

In the following, we will concentrate on the modification of tree-level diagrams. The effect of a minimal-substitution

prescription, applied to the most general (pion-nucleon) vertex, was discussed in detail by Ohta [9] and extended by others [10–12]. This work showed that a particular invariant amplitude (A_2) would be unaltered by a wide range of form factors and meson-nucleon vertices satisfying the Ward-Takahashi [3] identity. The possibility of having a form-factor modification of A_2 , accompanied by a gauge-invariance restoring contact term was subsequently considered in Ref. [10]. The recipe of Ref. [10] is simple to apply and has been widely adopted for use in single- and multiple-channel fits to meson-photoproduction data [13].

We find a flaw in the arguments used to derive this modification of the amplitude A_2 . It is, however, straightforward to find a modification of A_2 which *does* satisfy the constraints of gauge invariance and crossing symmetry. We then apply this technique to η' photoproduction to overcome the shortcomings of Refs. [14,15], namely, the violation of crossing symmetry. We also show how the use of our prescription can alter the conclusions of a phenomenological analysis.

II. REVIEW OF THE A_2 PROBLEM

We begin by recalling the basic results given in Ref. [11]. There the discussion was simplified by taking the simplest (pseudoscalar) πNN vertex in the specific process $\gamma p \rightarrow n \pi^+$. The contribution from the Born diagrams corresponding to s -, t -, and u -channel exchanges is

$$\begin{aligned} \epsilon \cdot M_{fi} = & \sqrt{2} g e \bar{u}_n \gamma_5 \frac{(p_1 + k) \cdot \gamma + m}{s - m^2} \left[\epsilon \cdot \gamma - \frac{\kappa_p}{2m} \epsilon \cdot \gamma k \cdot \gamma \right] u_p \\ & + 2 \sqrt{2} g e \bar{u}_n \frac{q \cdot \epsilon}{t - \mu^2} \gamma_5 u_p \\ & - \sqrt{2} g e \bar{u}_n \frac{\kappa_n}{2m} \epsilon \cdot \gamma k \cdot \gamma \frac{(p_2 - k) \cdot \gamma + m}{u - m^2} \gamma_5 u_p, \quad (1) \end{aligned}$$

where k and q represent the photon and pion four-momenta, and p_1 and p_2 are the respective proton and neutron four-momenta. The quantities m and μ are the nucleon and pion masses, ϵ is the photon polarization vector, g is the pseudoscalar $\pi^0 pp$ coupling constant, and κ_p and κ_n are the proton

*Electronic address: davidr@rpi.edu

†Electronic address: rworkman@gwu.edu

and neutron anomalous magnetic moments. Note that in Refs. [10,11], g denoted the pseudoscalar π^+np coupling constant.

If a strong form factor is inserted at the πNN vertex of each Born term, gauge invariance is clearly lost. The following term violates gauge invariance:

$$\begin{aligned} \epsilon \cdot \bar{M} = & \sqrt{2}ge\bar{u}_n\gamma_5 \left[\frac{2p_1 \cdot \epsilon}{s-m^2} F(s, m^2, \mu^2) \right. \\ & \left. + \frac{2q \cdot \epsilon}{t-\mu^2} F(m^2, m^2, t) \right] u_p, \end{aligned} \quad (2)$$

where $F(s, u, t)$ is a general form factor. From Ohta's [9] relations, the term required to restore gauge invariance is

$$\begin{aligned} \sqrt{2}ge\bar{u}_n\gamma_5 \left[\frac{2p_1 \cdot \epsilon}{s-m^2} [F(m^2, m^2, \mu^2) - F(s, m^2, \mu^2)] \right. \\ \left. + \frac{2q \cdot \epsilon}{t-\mu^2} [F(m^2, m^2, \mu^2) - F(m^2, m^2, t)] \right] u_p, \end{aligned} \quad (3)$$

with $F(m^2, m^2, \mu^2) = 1$. This precisely cancels the form factor effect on the terms in Eq. (2). Writing this in terms of invariant amplitudes

$$\epsilon \cdot M_{fi} = \bar{u}_n \sum_{j=1}^4 A_j M_j u_p, \quad (4)$$

with the explicitly gauge invariant representation

$$\begin{aligned} M_1 &= -\gamma_5 \epsilon \cdot \gamma k \cdot \gamma, \\ M_2 &= 2\gamma_5 (\epsilon \cdot p_1 k \cdot p_2 - \epsilon \cdot p_2 k \cdot p_1), \\ M_3 &= \gamma_5 (\epsilon \cdot \gamma k \cdot p_1 - \epsilon \cdot p_1 k \cdot \gamma), \\ M_4 &= \gamma_5 (\epsilon \cdot \gamma k \cdot p_2 - \epsilon \cdot p_2 k \cdot \gamma), \end{aligned} \quad (5)$$

noting that such a representation is only possible for a gauge invariant amplitude, we have

$$A_1 = \frac{\sqrt{2}geF(s, m^2, \mu^2)}{s-m^2} (1 + \kappa_p) + \frac{\sqrt{2}geF(m^2, u, \mu^2)}{u-m^2} \kappa_n, \quad (6)$$

$$\begin{aligned} A_2 &= \frac{2\sqrt{2}ge}{(s-m^2)(t-\mu^2)}, \\ A_3 &= \frac{\sqrt{2}geF(s, m^2, \mu^2)}{s-m^2} \frac{\kappa_p}{m}, \\ A_4 &= \frac{\sqrt{2}geF(m^2, u, \mu^2)}{u-m^2} \frac{\kappa_n}{m}. \end{aligned}$$

Thus, one sees that the A_2 amplitude is unaltered by form factors, within Ohta's prescription for retaining gauge invariance. The result holds for a more general pion-nucleon ver-

text and any charge channel. Of course, this result does not prevent the addition of terms which are individually gauge invariant.

Since form factors were added to provide a cutoff for the Born term contributions, this result is somewhat disappointing, as the A_2 contribution is not damped. In addition, explicit one-loop meson calculations [7,8] show that the A_2 amplitude is modified. In Ref. [10], a form factor for A_2 was inserted and the addition of a contact interaction was proposed to restore gauge invariance. We give this recipe below, show why it has a problem, and then propose an alternate form. The result is then generalized using crossing symmetry.

III. THE HABERZETTL FORM FACTOR

In Ref. [10] a gauge invariant term, $\hat{F}A_2$, was added and subtracted from a set of Born terms with form factors at the strong vertices. This resulted in the replacement $A_2 \rightarrow \hat{F}A_2$ above, with a remaining gauge-invariance-violating term

$$\begin{aligned} \epsilon \cdot M_{\text{viol}} = & \sqrt{2}ge\bar{u}_n\gamma_5 \left[\frac{2p_1 \cdot \epsilon}{s-m^2} [F(s, m^2, \mu^2) - \hat{F}] \right. \\ & \left. + \frac{2q \cdot \epsilon}{t-\mu^2} (F(m^2, m^2, t) - \hat{F}) \right] u_p, \end{aligned} \quad (7)$$

which was to be cancelled by a *contact* interaction.

By requiring this additional term to be free of poles, i.e., a contact term, we can constrain the functional form of \hat{F} . Denoting $F(s, \mu^2, m^2) \equiv F_1(s)$, with similar abbreviations for $F_2(u)$ and $F_3(t)$, subject to the constraints

$$F_1(m^2) = F_2(m^2) = F_3(\mu^2) = 1, \quad (8)$$

the general form for \hat{F} used in Ref. [10] was

$$\hat{F} = a_1 F_1(s) + a_2 F_2(u) + a_3 F_3(t), \quad (9)$$

subject to $a_1 + a_2 + a_3 = 1$. This form has been used in many recent analyses of photonic and multichannel reactions off the nucleon [13]. However, it is straightforward to demonstrate that this recipe results in a term with pole contributions, and is therefore incorrect because it cannot be cancelled by a contact term. Taking $a_2 = 1$, and inserting \hat{F} into the relation for M_{viol} , one can easily see the problem. Choosing, for example, the point $s = m^2$ does not fix the value of u , which is only constrained by the condition $t + u = \mu^2 + m^2$. As a consequence, when $s = m^2$ it is not necessarily the case that $F_2(u) = 1$. As a result, the pole terms are not cancelled, except for zero photon energy, as noted by the authors. To ensure cancellation of the poles, the needed constraint is, in this case,

$$\hat{F}(s = m^2, t) = \hat{F}(s, t = \mu^2) = 1, \quad (10)$$

which does not, however, imply that $\hat{F}(s, t) = 1$ for all s and t . For example, the following choice¹

$$\hat{F} = F_1(s) + F_3(t) - F_1(s)F_3(t) \quad (11)$$

satisfies the above constraint. Note that this method can be easily adapted to charged kaon production.

So far, we have only considered constraints provided by gauge invariance and the pole structure of the amplitude. However, crossing symmetry is also important. As the $p\pi^-$ channel is closely related to the $n\pi^+$ channel via crossing, let us therefore consider the isospin channel $T^{(-)}$ defined by²

$$T^{(-)} = \frac{1}{2\sqrt{2}}(T^{n\pi^+} - T^{p\pi^-}). \quad (12)$$

Inserting strong form factors, the matrix element for the electric Born terms is

$$\begin{aligned} \epsilon \cdot M = & \frac{eg}{2} F_1(s) \bar{u}_2 \gamma_5 \frac{(p_1 + k) \cdot \gamma + m}{s - m^2} \gamma \cdot \epsilon u_1 \\ & - \frac{eg}{2} F_2(u) \bar{u}_2 \gamma \cdot \epsilon \frac{(p_2 - k) \cdot \gamma + m}{u - m^2} \gamma_5 u_1 \\ & + 2eg F_3(t) \bar{u}_2 \gamma_5 \frac{q \cdot \epsilon}{t - \mu^2} u_1. \end{aligned} \quad (13)$$

After some rearrangement, this may be written as

$$\begin{aligned} \epsilon \cdot M = & \frac{eg_-}{2} u_2 M_1 \left[\frac{F_1(s)}{s - m^2} - \frac{F_2(u)}{u - m^2} \right] u_1 \\ & + eg \hat{F}(s, u, t) \bar{u}_2 \frac{M_2}{(t - \mu^2)} \left[\frac{1}{s - m^2} - \frac{1}{u - m^2} \right] u_1 \\ & + \epsilon \cdot \Delta M, \end{aligned} \quad (14)$$

where $M_{1,2}$ are the gauge invariant operators defined in Eq. (5) and

$$\begin{aligned} \epsilon \cdot \Delta M = & -\frac{eg_-}{2} \bar{u}_2 \gamma_5 \left[\frac{2p_1 \cdot \epsilon}{s - m^2} [\hat{F}(s, u, t) - F_1(s)] \right. \\ & + \frac{4q \cdot \epsilon}{t - \mu^2} [\hat{F}(s, u, t) - F_3(t)] \\ & \left. - \frac{2p_2 \cdot \epsilon}{u - m^2} [\hat{F}(s, u, t) - F_2(u)] \right] u_1. \end{aligned} \quad (15)$$

As $A_{1,2}^{(-)}$ are odd under crossing, i.e.,

$$A_{1,2}^{(-)}(s, u, t) = -A_{1,2}^{(-)}(u, s, t), \quad (16)$$

we see that $\hat{F}(s, u, t) = \hat{F}(u, s, t)$ and $F_2(u) = F_1(u)$, that is, $F_2(u)$ is obtained from $F_1(s)$ by the replacement $s \rightarrow u$. In




FIG. 1. The total cross section for $\gamma p \rightarrow K^+ \Lambda$ versus the photon lab energy E_γ . Plotted are the unmodified Born terms (solid), Ohta's prescription [9] (dotted), and our form-factor modification (see text). Data from Ref. [16].

order to cancel the poles in $\epsilon \cdot \Delta M$, so this term may be cancelled by a contact term, we need

$$\hat{F}(m^2, u, t) = \hat{F}(s, m^2, t) = \hat{F}(s, u, \mu^2) = 1. \quad (17)$$

Although not unique, one crossing symmetric form for \hat{F} that satisfies the above constraints is

$$\begin{aligned} \hat{F}(s, u, t) = & F_1(s) + F_1(u) + F_3(t) - F_1(s)F_1(u) \\ & - F_1(s)F_3(t) - F_1(u)F_3(t) + F_1(s)F_1(u)F_3(t). \end{aligned} \quad (18)$$

IV. PHENOMENOLOGY

The use of different form factor schemes can have a significant effect on phenomenological fits to data. This is particularly clear in kaon photoproduction where the rapid increase of a pointlike Born contribution to this quantity is well known. The pointlike Born term contribution to the total cross section (solid) is compared to the data [16] in Fig. 1 using a $K\Lambda N$ coupling of 7.5 [17]. An attempt to cure this, by modifying individual multipoles, was described in Ref. [18]. Using the minimal substitution prescription of Ohta [9] (dotted), which damps all but the A_2 amplitude, one still obtains a Born contribution which grows too rapidly. Here, we use

$$F_1(s) = [1 + (s - m^2)^2 / \Lambda^4]^{-1}, \quad (19)$$

with a cutoff mass Λ of 1 GeV. Finally, we show in Fig. 1 the result arising from our modification of the Habermehl

¹It should be noted that this choice is not unique.

²The isoscalar channel $T^{(0)}$ is discussed below.




FIG. 2. The total cross section for $\gamma p \rightarrow p \eta'$ versus the photon lab energy, E_γ . The data are from SAPHIR [21] (diamonds), AHHM [20] (squares), and ABBHHM [19] (crosses). The solid line is the total cross section obtained from the Born terms with $g_{\eta'} = 1.9$ and without any form factors. The dashed line is from the work of Ref. [14] where the Born terms are multiplied by an overall form factor depending only on s , thus violating crossing symmetry. The dotted line is the best Born model fit to the data using the noncrossing symmetric model [15].

form (dashed). In addition to $F_1(s)$ given above, we also need \hat{F} and $F_3(t)$. We take \hat{F} to be of the form of Eq. (11) with

$$F_3(t) = [1 + (t - \mu^2)^2 / \Lambda^4]^{-1}, \quad (20)$$

where μ is the kaon mass and we again take $\Lambda = 1$ GeV.

Finally, we consider the fits to η' photoproduction performed in Refs. [14,15]. In these studies, the Born terms were added and compared to possible resonance contributions. The use of a coupling, $g_{\eta'}$, within the range of values, $1 \leq g_{\eta'} \leq 6$, estimated in this work [14] results in an (undamped) Born contribution exceeding the data [19–21] by a large factor, as is demonstrated by the solid line in Fig. 2 (here, $g_{\eta'} = 1.9$). There is more than one way to interpret this result. One possibility is that $g_{\eta'}$ is much smaller than estimated in Ref. [14]. A different possibility is that $g_{\eta'}$ is in the range given above, but the *effective* $g_{\eta'}$ is much smaller than $g_{\eta'}$ at the nucleon pole. In other words, as the proton and eta-prime are extended objects, form factors enter at the three-point vertices, and as these data are far from the nucleon pole, form factor effects can be significant. Finally, there are almost certainly other reaction mechanisms such as resonance exchanges, vector meson exchanges, and possibly contact terms.

In the original analysis of the η' data [14], the Born terms were multiplied by an overall form factor given by Eq. (19) with $\Lambda = 1.1$ GeV. In this model, the dominant contribution to the model was from the *background* of the D13(2080)

resonance, and the Born terms played a minor role. The resulting total cross section from the Born terms is shown by the dashed (nearly zero) line in Fig. 2. On the other hand, taking $g_{\eta'} = 3.5$ and $\Lambda = 2$ GeV, it was shown that one could qualitatively reproduce the total cross section with just Born terms (dotted line) [15]. However, the procedure of multiplying the Born terms by an overall form factor violates crossing symmetry and the amplitude has the wrong residue structure, and therefore is theoretically unacceptable. Other inconsistencies of this procedure have been discussed in Refs. [6,7]. Thus, it is not clear what conclusions, if any, can be drawn from the dashed and dotted curves in Fig. 2.

Using Ohta's method [9], one obtains a crossing symmetric amplitude, but as above the A_2 amplitude is not modified. Using our modification of the Habermehl form, the electric Born terms are

$$A_1 = e g_{\eta'} \left[\frac{F_1(s)}{s - m^2} + \frac{F_1(u)}{u - m^2} \right],$$

$$A_2 = \frac{2 e g_{\eta'}}{(s - m^2)(u - m^2)} \hat{F}(s, u), \quad (21)$$

where $F_1(m^2) = 1$. In this case, the constraints on $\hat{F}(s, u)$ are

$$\hat{F}(s, m^2) = \hat{F}(m^2, u) = 1, \quad (22)$$

from the condition to remove the poles, and $\hat{F}(s, u) = \hat{F}(u, s)$ from crossing symmetry. Note that this form is also valid for π^0 and η production, and the electric Born terms for isoscalar production of pions, defined by

$$T^{(0)} = \frac{1}{2\sqrt{2}} (T^{n\pi^+} + T^{p\pi^-}), \quad (23)$$

can be obtained from Eq. (21) using the replacement $g'_{\eta} \rightarrow g/2$. With this form, the Born terms in η' production can be sufficiently damped at high energies even for the largest estimated value of g'_{η} . For example, taking

$$\hat{F} = F_1(s) + F_1(u) - F_1(s)F_1(u), \quad (24)$$

with $F_1(s)$ given by Eq. (19), and using the upper limit for $g_{\eta'}$ with a cutoff of 700 MeV, the total cross section from the Born terms levels off at about $0.2 \mu\text{b}$ at a photon energy of 2 GeV. Furthermore, and this is the essential point, it is *not possible* to reproduce the total cross section shape here with only Born terms and reasonable values for Λ (500 to 2000 MeV), in contrast to the noncrossing symmetry model of Ref. [15]. Thus, the use of the present, more restrictive, form factor prescription narrows the possible interpretations of experimental data.

V. CONNECTION WITH CONTACT TERMS

It is known in field theory [22] that off-shell form factors are not unique and are closely related to contact terms. For different representations of the fields, the form factors and

contact terms differ, but the differences cancel out in the calculation of a physical scattering amplitude. In the case of dispersion relations [1,2], the nonuniqueness of the off-shell form factors is reflected in the a priori unknown number of needed subtractions [23]. For the phenomenological amplitudes presented here, it is also possible to shift form factor effects to contact terms. To illustrate this, recall that $F_1(m^2)=1$ and we may write, for example, for the A_1 term in Eq. (21),

$$A_1 = e g_{\eta'} \left[\frac{1}{s-m^2} + \frac{1}{u-m^2} + \frac{[F_1(s)-1]}{s-m^2} + \frac{[F_1(u)-1]}{u-m^2} \right]. \quad (25)$$

The first two terms above are recognized as the undamped Born terms, while the last two terms correspond to contact terms since the pole is cancelled as s or $u \rightarrow m^2$. As previously noted, the A_2 amplitude is not modified in Ohta's method [9], but his method does not forbid the inclusion of an additional gauge invariant contact term. Our modification of Haberzettl's procedure is equivalent to adding the gauge invariant contact term

$$A_2 \bar{u}_2 M_2 u_1 (\hat{F} - 1), \quad (26)$$

to the amplitude of Ref. [11] [Eq. (6) of this work]. Here, M_2 is the gauge invariant operator defined in Eq. (5) and A_2 is given by the second equation of Eqs. (6).

Finally, let us note that most authors assume that the form factors are real, and the phenomenological results presented here are also obtained using real form factors. In general, however, arguments based on unitarity [1,2,23] show that the off-shell form factors must be complex above the one-pion threshold. On the other hand, there is some justification for

using real form factors if one adopts a K -matrix approach to the scattering amplitude [24].

VI. CONCLUSIONS

We have found a flaw in the gauging procedure of Ref. [10] and have noted that the method used in Refs. [14,15] is theoretically unacceptable as it violates crossing symmetry. We have determined the general constraints \hat{F} must satisfy for the gauge-violating term to be a contact term, which then can be cancelled by a contact term. This method was generalized to be consistent with crossing symmetry and applied to η' photoproduction. This model is theoretically superior to previous works [14,15] and has the important phenomenological consequence that the total cross section cannot be interpreted as arising solely from the Born terms, in contrast to Ref. [15]. Other phenomenological applications are being investigated.

ACKNOWLEDGMENTS

We are grateful to Nimai C. Mukhopadhyay for many useful discussions regarding eta-prime photoproduction and the need for a crossing symmetric model. We thank Paul Stoler for a critical reading of the manuscript. This work was supported in part by the U. S. Department of Energy Grants No. DE-FG02-99ER41110 and DE-FG02-88ER40448. R.W. gratefully acknowledges a contract from Jefferson Lab under which this work was done. Jefferson Lab is operated by the Southeastern Universities Research Association under the U.S. Department of Energy Contract No. DE-AC05-84ER40150. R.M.D. thanks the Center for Nuclear Studies for supporting a visit in which part of this work was completed.

-
- [1] E. Kazes, *Nuovo Cimento* **13**, 1226 (1959).
 [2] A. Bincer, *Phys. Rev.* **118**, 855 (1960).
 [3] J. C. Ward, *Phys. Rev.* **78**, 182 (1950); Y. Takahashi, *Nuovo Cimento* **6**, 371 (1957).
 [4] F. A. Berends and G. B. West, *Phys. Rev.* **188**, 2538 (1969).
 [5] F. Gross and D. Riska, *Phys. Rev. C* **36**, 1928 (1987).
 [6] H. W. L. Naus and J. H. Koch, *Phys. Rev. C* **39**, 1907 (1989).
 [7] J. W. Bos, S. Scherer, and J. H. Koch, *Nucl. Phys.* **A547**, 488 (1992).
 [8] V. Bernard, N. Kaiser, and U.-G. Meissner, *Phys. Lett. B* **282**, 448 (1992); **378**, 337 (1996).
 [9] K. Ohta, *Phys. Rev. C* **40**, 1335 (1989).
 [10] H. Haberzettl, C. Bennhold, T. Mart, and T. Feuster, *Phys. Rev. C* **58**, R40 (1998).
 [11] R. L. Workman, H. W. L. Naus, and S. J. Pollock, *Phys. Rev. C* **45**, 2511 (1992).
 [12] S. Wang and M. K. Banerjee, *Phys. Rev. C* **54**, 2883 (1996).
 [13] T. Feuster and U. Mosel, *Phys. Rev. C* **59**, 460 (1999); T. Mart and C. Bennhold, *ibid.* **61**, 012201(R) (1999); T. Mart, *ibid.* **62**, 038201 (2000); F. X. Lee, T. Mart, C. Bennhold, H. Haberzettl, and L. E. Wright, *nucl-th/9907119*.
 [14] J.-F. Zhang, N. C. Mukhopadhyay, and M. Benmerrouche, *Phys. Rev. C* **52**, 1134 (1995).
 [15] N. C. Mukhopadhyay, J.-F. Zhang, R. M. Davidson, and M. Benmerrouche, *Phys. Lett. B* **410**, 73 (1997).
 [16] Data from the SAID database, URL <http://gwddac.phys.gwu.edu>. This source also contains representative fits from several different groups.
 [17] This value for the KAN coupling is within the range of values found or used in R. A. Adelseck, C. Bennhold, and L. E. Wright, *Phys. Rev. C* **32**, 1681 (1985), and Ref. [10].
 [18] H. Tanabe, M. Kohno, and C. Bennhold, *Phys. Rev. C* **39**, 741 (1989).
 [19] R. Erbe *et al.*, The ABBHHM Collaboration, *Phys. Rev.* **175**, 1669 (1968).
 [20] W. Struczinski *et al.*, The AHHM Collaboration, *Nucl. Phys.* **B108**, 45 (1976).
 [21] R. Plötzke *et al.*, *Phys. Lett. B* **444**, 555 (1998).
 [22] S. Scherer and H. Fearing, *Phys. Rev. C* **51**, 359 (1995).
 [23] R. M. Davidson and G. I. Poulis, *Phys. Rev. D* **54**, 2228 (1996).
 [24] S. Kondratyuk and O. Scholten, *Phys. Rev. C* **59**, 1070 (1999).