

## Attenuation of the intensity within a superdeformed band

A. J. Sargeant,<sup>1</sup> M. S. Hussein,<sup>1</sup> M. P. Pato,<sup>1</sup> N. Takigawa,<sup>2</sup> and M. Ueda<sup>1,\*</sup>

<sup>1</sup>*Nuclear Theory and Elementary Particle Phenomenology Group, Instituto de Física, Universidade de São Paulo, C.P. 66318, 05315-970 São Paulo, SP, Brazil*

<sup>2</sup>*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

(Received 29 January 2001; published 8 January 2002)

The attenuation of the intraband intensity of a superdeformed band, which results from mixing with normally deformed configurations, is calculated using reaction theory. It is found that the sharp increase of the attenuation is mostly due to the tunneling through a spin dependent barrier and not to the chaotic nature of the normally deformed states.

DOI: 10.1103/PhysRevC.65.024302

PACS number(s): 21.10.Re, 24.60.Dr, 24.60.Lz

It is now well established that the intensities of  $E2$  gamma transitions within a superdeformed (SD) rotational band show cascades down to low angular momentum [1–7]. These cascades exhibit the distinct feature that the intensity remains constant until a certain spin is reached whereafter the intensity drops to zero within a few transitions. The sharp drop in intensity is commonly referred to as the decay out of a superdeformed band and is believed to arise from mixing of the SD states with normally deformed (ND) states of identical spin.

The earliest theoretical work to implement such an interpretation [8–11] used a statistical model of the coupling between the SD and ND states. More recently, Refs. [12,13] used a framework originally developed for the study of compound nuclear reactions to derive formulas for the intensity in a more rigorous fashion (the expressions for the intensity in Refs. [8–11] are deduced from probability arguments). Reference [13] concluded that Refs. [8–11] are valid in the nonoverlapping resonance region. References [8–11] further calculate the spin dependence of the relevant parameters (the electromagnetic widths of the SD and ND states, the level density of the ND states, and spin dependence of the barrier separating the SD and ND wells), which Refs. [12,13] do not. Two features common to Refs. [8–13] are (i) the use of the Gaussian orthogonal ensemble (GOE) to simulate the ND states, (ii) the use of the “golden rule” to extract a width for the SD states due to mixing with the ND states.

Here, as in Refs. [12,13] we exploit the similarity between the decay out of superdeformed bands and compound nuclear reactions to write the intensity as the sum of average and fluctuation contributions. However we use an energy average in place of the ensemble average used in Refs. [12,13]. The energy average approach allows the inclusion of the following features, which are more difficult to incorporate into an ensemble average.

(i) A hierarchy of complexity in the ND spectrum may be introduced.

(ii) A statistical model different from the GOE may be used to simulate the ND states, as was proposed in Refs. [14,15].

(iii) A width for the SD states due to mixing with the ND states arises naturally without appealing to the “golden rule” whose range of validity has been found to be restricted [16,17].

In Fig. 1 we show a schematic plot of the energy of ND and SD bands as a function of spin. The observable in which attenuation is seen is the total intensity of two consecutive  $E2$  photons in the cascade down the SD band. Let  $|J\rangle$  denote an SD configuration with spin  $J$ . The relative intensity of the two-step transition  $|J+2\rangle \xrightarrow{\gamma_1} |J\rangle \xrightarrow{\gamma_2} |J-2\rangle$  (relative to the intensity of the same two-step transition in the absence of mixing with other configurations) is given by

$$F_J = \frac{1}{2\pi\Gamma_{J+2}^\gamma} \int_{-\infty}^{\infty} dE_{\gamma_1} \int_{-\infty}^{\infty} dE_{\gamma_2} |\langle J-2|T(E_{J-2}+E_{\gamma_2}) \times |J+2\rangle|^2 \delta(E-E_{J-2}-E_{\gamma_2}) \\ = \frac{1}{2\pi\Gamma_{J+2}^\gamma} \int_{-\infty}^{\infty} dE |\langle J-2|T(E)|J+2\rangle|^2, \quad (1)$$

where  $E \equiv E_{J+2} - E_{\gamma_1} = E_J$  takes account of the Hamiltonian of the electromagnetic field,  $E_J$  being the energy of  $|J\rangle$  and  $E_{\gamma_1}$  and  $E_{\gamma_2}$  the energies of the two consecutive photons. The electromagnetic width of  $|J+2\rangle$  is  $\Gamma_{J+2}^\gamma$  making




FIG. 1. Schematic diagram illustrating the decay out of a superdeformed band.

\*Present address: Institute of Physics, University of Tsukuba, Tennoudai 1-1-1, Tsukuba, Ibaraki 305-8571, Japan.

$2\pi\Gamma_{J+2}^\gamma$  the intensity when there is no mixing with the ND states and thus no flux loss from the SD band. Note that in Eq. (1) we ignore the widths of the initial and final states for the purpose of calculating the relative intensity.

The transition amplitude is given by

$$\langle J-2|T(E)|J+2\rangle = \gamma_{J+2}\langle J|G(E)|J\rangle_{\gamma_j}. \quad (2)$$

Here  $\gamma_{J+2}$  is the electromagnetic decay amplitude of  $|J+2\rangle$  defined such that  $\Gamma_{J+2}^\gamma = \gamma_{J+2}^2$ :  $\gamma_J$  and  $\Gamma_J^\gamma = \gamma_J^2$  are the corresponding decay amplitude and width of  $|J\rangle$  whilst the Green's function is given by

$$G(E) = (E - H)^{-1}. \quad (3)$$

The total nuclear Hamiltonian, which takes the coupling to the electromagnetic field into account, is denoted by  $H$ .

The projected Green's function  $\langle J|G(E)|J\rangle$  may be expressed in terms of its Lorentzian energy average  $\langle J|G^{\text{av}}(E)|J\rangle = \langle J|G(E + iI/2)|J\rangle$  (energy averaging interval  $I$ ) plus a fluctuation part [18,19]

$$\langle J|G|J\rangle = \langle J|G^{\text{av}}|J\rangle + \langle J|G^{\text{fluc}}|J\rangle, \quad (4)$$

where by definition the energy average of  $\langle J|G^{\text{fluc}}|J\rangle$  is zero. Thus Eq. (1) for the relative intensity may be written

$$F_J = F_J^{\text{av}} + F_J^{\text{fluc}}, \quad (5)$$

where

$$F_J^{\text{av}} = \frac{\Gamma_J^\gamma}{2\pi} \int_{-\infty}^{\infty} dE |\langle J|G^{\text{av}}|J\rangle|^2 \quad (6)$$

and

$$F_J^{\text{fluc}} = \frac{\Gamma_J^\gamma}{2\pi} \int_{-\infty}^{\infty} dE (|\langle J|G^{\text{fluc}}|J\rangle|^2)^{\text{av}}. \quad (7)$$

In this paper we focus our discussion on  $F_J^{\text{av}}$ . It can be shown that

$$\langle J|G^{\text{av}}(E)|J\rangle = \frac{1}{E - E_J + i\Gamma_J^\gamma/2 - W_{JJ}(E)}. \quad (8)$$

The derivation of Eq. (8) for  $\langle J|G^{\text{av}}|J\rangle$  and an expression for  $\langle J|G^{\text{fluc}}|J\rangle$  using projection operator techniques will be reported in a subsequent paper.

The form of  $W_{JJ}(E)$  depends on the specific model for the Hamiltonian that is employed. It is our aim to study whether or not the chaotic nature [as classified by random matrix theory (RMT)] of the ND states is decisive in explaining the observed attenuation. In order to isolate the statistical aspects of the calculation we use two different models distinguished by whether the tunneling interaction mixes  $|J\rangle$  randomly with the ND states (*model A*) or whether it couples more strongly to certain ND states than others (*model B*). In the latter we shall make the most extreme assumption that  $|J\rangle$  couples to only one ND state.

*Model A* is represented by the matrix

$$H \rightarrow \begin{pmatrix} E_J & V_{Jn} \\ V_{Jn} & E_n \delta_{n'n} \end{pmatrix} - \frac{i}{2} \begin{pmatrix} \Gamma_J^\gamma & 0 \\ 0 & \Gamma_N^\gamma \delta_{n'n} \end{pmatrix}, \quad n=1, \dots, N, \quad (9)$$

where  $E_n$  denotes the energies of the  $N$  ND states with which  $|J\rangle$  mixes due to the real tunneling interaction  $V_{Jn}$ . Here  $\Gamma_N^\gamma$  is an electromagnetic width, which we assume to be common to the ND states. With these definitions  $W_{JJ}(E)$  becomes

$$W_{JJ}(E) = \sum_{n=1}^N \frac{[V_{Jn}]^2}{E - E_n + i(\Gamma_N^\gamma + I)/2}. \quad (10)$$

The energies  $E_n$  are constructed using the deformed Gaussian orthogonal ensemble (DGOE) [20]. The DGOE allows a smooth interpolation from Poisson to GOE statistics by varying a mixing parameter  $\lambda$  from 0 to 1. Thus the  $E_n$  are the eigenvalues of a random Hamiltonian  $h$ , which is real symmetric and whose matrix elements are taken to be Gaussian distributed random numbers with zero mean and variances

$$\langle h_{nn}^2 \rangle = \frac{2}{N}, \quad \langle h_{n'n}^2 \rangle = \frac{\lambda^2}{N}, \quad n' \neq n. \quad (11)$$

The random tunneling interaction is taken to have zero mean and variance

$$\langle V_{Jn}^2 \rangle = v_J^2. \quad (12)$$

We assume that  $E_J$  lies in the middle of the  $N$  ND states, that is,  $E_J = 0$ . Following Refs. [8,9], [12,13] we introduce a spreading width  $\Gamma_J^\downarrow$  through the golden rule,

$$\Gamma_J^\downarrow = 2\pi v_J^2. \quad (13)$$

We maintain doubts about the meaningfulness of Eq. (13) regarding its interpretation as a width [17]. For practical purposes, however, it is a change of variable from  $v_J$  to  $\Gamma_J^\downarrow$ . All quantities of dimensions energy are to be understood to have units of  $D_J$ , where  $D_J$  denotes the mean spacing of the ND states around  $E_J$ . Thus the  $E_n$  [and the  $E_q$  and  $\mathcal{V}_{dq}$  in Eq. (15)] that are generated from the DGOE are to be understood to have been unfolded such that the  $E_n$  and  $E_q$  have mean spacing equal to unity. Thus we may write  $W_{JJ}(E)$  as

$$W_{JJ}(E) = \frac{\Gamma_J^\downarrow}{2\pi} \sum_{n=1}^N \frac{g_n^2}{E - E_n + i(\Gamma_N^\gamma + I)/2}, \quad (14)$$

where the  $g_n$ ,  $n=1, \dots, N$  are Gaussian distributed random numbers with zero mean and unit variance.

*Model B* is represented by the matrix

$$H \rightarrow \begin{pmatrix} E_J & V_{Jd} & 0 \\ V_{Jd} & E_d & \mathcal{V}_{dq} \\ 0 & \mathcal{V}_{dq} & E_q \delta_{q'q} \end{pmatrix} - \frac{i}{2} \begin{pmatrix} \Gamma_J^\gamma & 0 & 0 \\ 0 & \Gamma_N^\gamma & 0 \\ 0 & 0 & \Gamma_N^\gamma \delta_{q'q} \end{pmatrix}, \quad (15)$$

$$q = 1, \dots, N-1.$$

We assume here that  $|J\rangle$  couples to only one state of normal deformation  $|d\rangle$ , which has energy  $E_d$  with strength  $V_{Jd}$ . This special state is subsequently mixed with other ND configurations with energies  $E_q$  by a residual interaction  $\mathcal{V}_{dq}$ . Now  $W_{JJ}(E)$  becomes

$$W_{J,J}(E) = [V_{Jd}]^2 \sum_{n=1}^N \frac{[c_d(n)]^2}{E - E_n + i(\Gamma_N^\gamma + I)/2}, \quad (16)$$

where  $c_d(n)$  denotes component  $d$  of the  $n$ th eigenvector of the submatrix of the first term of Eq. (15) obtained by excluding the first row and the first column. Now the  $E_q$  are eigenvalues of a random Hamiltonian  $h$ , which is real symmetric and whose matrix elements are taken to be Gaussian distributed random numbers with zero mean and variances,

$$\langle h_{qq}^2 \rangle = \frac{2}{N}, \quad \langle h_{q'q}^2 \rangle = \frac{\lambda^2}{N}, \quad q' \neq q. \quad (17)$$

The residual interaction is also taken to have zero mean and variance,

$$\langle \mathcal{V}_{dq}^2 \rangle = \frac{\lambda^2}{N}. \quad (18)$$

We put  $E_d = E_J = 0$ . Introducing

$$\Gamma_J^\downarrow = 2\pi \frac{[V_{Jd}]^2}{N}, \quad (19)$$

we can write Eq. (16) as

$$W_{JJ}(E) = \frac{\Gamma_J^\downarrow}{2\pi} \sum_{n=1}^N \frac{N[c_d(n)]^2}{E - E_n + i(\Gamma_N^\gamma + I)/2}. \quad (20)$$

Thus comparing Eq. (14) with Eq. (20) we see that, although the meaning of  $\Gamma_J^\downarrow$  is different for the two models, the difference between *model A* and *model B* boils down how much the distribution  $[c_d(n)]^2$  differs from that of  $g_n^2$ . This difference is not trivial as  $[c_d(n)]^2$  has a dramatic  $\lambda$  dependence [see Fig. (1) in Ref. [15]]. The inclusion of the factor  $1/N$  in Eq. (19) makes clear that *model A* and *model B* are only comparable when  $V_{Jd}^2$  is of the order  $NV_J^2$ .

*Model A* is precisely equivalent to that of Refs. [12,13] when  $\lambda = 1$ . The real part of Eq. (15) used in *model B* is equivalent to what is used in Refs. [14,15], however, we calculate the average intensity integrated over the energy  $F_J^{\text{av}}$ , whereas Refs. [14,15] calculate a tunneling probability that is more closely related to  $W_{JJ}(E)$ .

Note that  $F_J^{\text{av}}$  can be written as

$$P_J^{\text{av}} = \frac{\Gamma_J^\gamma}{2\pi} \int_{-\infty}^{\infty} dE \frac{1}{[E - E_J - \Delta_J^\downarrow(E)]^2 + \frac{\Gamma_J^{\gamma^2}}{4} \left(1 + \frac{\Gamma_J^\downarrow(E)}{\Gamma_J^\gamma}\right)^2}. \quad (21)$$

where




FIG. 2. Calculated attenuation factor  $F_J^{\text{av}}$  as a function of  $\lambda$  for some values of  $b_J = \Gamma_J^\downarrow / \Gamma_J^\gamma$ . The thin lines were calculated using *model A* and the thick lines were calculated using *model B*.

$$\Delta_J^\downarrow(E) = \text{Re} W_{JJ}(E) \quad (22)$$

and

$$\Gamma_J^\downarrow(E) = -2 \text{Im} W_{JJ}(E). \quad (23)$$

Ignoring the shift  $\Delta_J^\downarrow(E)$  altogether and assuming that the width  $\Gamma_J^\downarrow(E)$  has the energy independent value  $\Gamma_J^\downarrow$  one obtains the principal result of Ref. [12] that

$$F_J^{\text{av}} \approx \frac{\Gamma_J^\gamma}{\Gamma_J^\gamma + \Gamma_J^\downarrow}. \quad (24)$$

We now present numerical calculations of  $F_J^{\text{av}}$  with  $N = 50$ ,  $\Gamma_J^\gamma = 0.01D_J$ . An ensemble average was performed over 100 realizations in Fig. 2 and over 1000 realizations in




FIG. 3. Attenuation factor  $F_J^{\text{av}}$  as a function of  $b_J = \Gamma_J^\downarrow / \Gamma_J^\gamma$  for some values of  $\lambda$ , calculated using *model B*.

Fig. 3. The effect of increasing  $\Gamma_N^\gamma$ , identical to that obtained by increasing the energy averaging interval  $I$ , is to broaden  $\Gamma_J^\downarrow(E)$  [Eq. (23)] and thus push  $F_J^{\text{av}}$  closer to the approximation given by Eq. (24). This is in line with what it is reported in [21], in which Eq. (24) is obtained in the limit  $\Gamma_N^\gamma/\Gamma_J^\gamma \rightarrow \infty$  for their two-level model.

In our calculations we put  $\Gamma_N^\gamma + I = 3D_J$ . With this choice one may describe what Ref. [13] calls the overlapping resonance region. Reference [13] gives the impression that the relative intensity is independent of  $\Gamma_J^\gamma$ . Whilst we agree with [13] that the ratios  $b_J = \Gamma_J^\downarrow/\Gamma_J^\gamma$  and  $\Gamma_N^\gamma/D$  are of principal importance in understanding the decay out it can be seen from Eq. (21) that  $F_J^{\text{av}}$  is only independent of  $\Gamma_J^\gamma$  if  $\Delta_J^\downarrow(E)$  and  $\Gamma_J^\downarrow(E)$  are constant.

Figure 2 shows the dependence of  $F_J^{\text{av}}$  on  $\lambda$ , the strength of the mixing amongst the ND states, for several values of  $b_J$  for both *model A* and *model B*. For *model A* the variation of  $F_J^{\text{av}}$  with  $\lambda$  is rather slight compared to *model B*. This is because the  $\lambda$  dependence of *model A* is contained in the eigenvalues  $E_n$ , which are unfolded to have unit mean spacing. *Model B* has a further and more significant  $\lambda$  dependence contained in the eigenvectors  $c_d(n)$ . For *model B*,  $F_J^{\text{av}}$  decreases with decreasing  $\lambda$  to a value that is limited by the value of  $\Gamma_N^\gamma + I$ . Note that  $F_J^{\text{av}}$  can change at most by a factor of about 5 by varying  $\lambda$ .

Figure 3 shows  $F_J^{\text{av}}$  as function of  $b_J$  for some values of  $\lambda$  calculated using *model B*. The calculations for *model A* are not shown as they can barely be distinguished from the calculation for  $\lambda = 1$  using *model B*. The effect of changing  $\lambda$  is

to move the value of  $b_J$  (and hence  $J$ ) at which the decay out occurs. Thus from Fig. 3 we conclude that the decay out is slightly hindered by increasing  $\lambda$ .

Regarding Refs. [14,15], which report an increase in the tunneling probability of several orders of magnitude with increasing  $\lambda$ , we do not consider ourselves at odds with this work since, as already mentioned above, we do not calculate the same quantity. A further difference between *model B* of this paper and Refs. [14,15] is that their author places  $|d\rangle$  at the position  $N/4$  thus making the distribution  $c_d(n)$  asymmetric. This would correspond in our calculation to making the difference  $E_J - E_d$  nonzero (we see no reason not to set  $E_J = E_d$ ).

We postpone an investigation of the roles of  $\Gamma_N^\gamma$  and  $I$  to a subsequent paper. The results of Ref. [13] indicate that  $F_J^{\text{fluc}}$ , Eq. (7), is important when  $\Gamma_N^\gamma$  is a small fraction of  $D_J$  (non-overlapping resonance region).

It was already found from phenomenological analysis some years ago [8,9,3] that  $F_J^{\text{av}}$  falls exponentially with decreasing spin. We conclude here that the chaotic nature of the ND states, as classified by  $\lambda$ , cannot account for such behavior. The exponential drop in the intraband intensity must be due to the spin dependence of the tunneling matrix element contained in  $b_J$ . The calculation of  $b_J$  is not trivial and we refer the reader to [22,23], which continue the work of Refs. [8–11] for some recent calculations.

A.J.S. thanks J. A. Tostevin for his comments on an early version of this paper. This work was supported by FAPESP.

- 
- [1] P. J. Twin *et al.*, Phys. Rev. Lett. **57**, 811 (1986).  
 [2] A. Atac *et al.*, Nucl. Phys. **A557**, 109c (1993).  
 [3] T. L. Khoo *et al.*, Nucl. Phys. **A557**, 83c (1993).  
 [4] R. Krücken *et al.*, Phys. Rev. Lett. **73**, 3359 (1994).  
 [5] R. Krücken *et al.*, Phys. Rev. C **54**, 1182 (1996).  
 [6] R. Kühn *et al.*, Phys. Rev. C **55**, R1002 (1997).  
 [7] R. Krücken *et al.*, Phys. Rev. C **55**, R1625 (1997).  
 [8] E. Vigezzi, R. A. Broglia, and T. Døssing, Phys. Lett. B **249**, 163 (1990).  
 [9] E. Vigezzi, R. A. Broglia, and T. Døssing, Nucl. Phys. **A520**, 179c (1990).  
 [10] Y. R. Shimizu, F. Barranco, R. A. Broglia, T. Døssing, and E. Vigezzi, Phys. Lett. B **274**, 253 (1992).  
 [11] Y. R. Shimizu, E. Vigezzi, T. Døssing, and R. Broglia, Nucl. Phys. **A557**, 99c (1993).  
 [12] H. A. Weidenmüller, P. von Brentano, and B. R. Barrett, Phys. Rev. Lett. **81**, 3603 (1998).  
 [13] Jian-zhong Gu and H. A. Weidenmüller, Nucl. Phys. **A660**, 197 (1999).  
 [14] S. Åberg, Phys. Rev. Lett. **82**, 299 (1999).  
 [15] S. Åberg, Nucl. Phys. **A649**, 392c (1999).  
 [16] N. Frazier, B. A. Brown, and V. Zelevinsky, Phys. Rev. C **54**, 1665 (1996).  
 [17] A. J. Sargeant, M. S. Hussein, M. P. Pato, and M. Ueda, Phys. Rev. C **61**, 011302(R) (2000).  
 [18] M. Kawai, A. K. Kerman, and K. W. McVoy, Ann. Phys. (N.Y.) **75**, 156 (1973).  
 [19] H. Feshbach, *Theoretical Nuclear Physics: Nuclear Reactions* (Wiley, New York, 1992), Chap. IV, Sec. 8.  
 [20] M. S. Hussein and M. P. Pato, Phys. Rev. Lett. **70**, 1089 (1993).  
 [21] C. A. Stafford and B. R. Barrett, Phys. Rev. C **60**, 051305(R) (1999).  
 [22] Y. R. Shimizu, M. Matsuo, and K. Yoshida, Nucl. Phys. **A682**, 464 (2001).  
 [23] K. Yoshida, M. Matsuo, and Y. R. Shimizu, nucl-th/0010074.