# Elliptic flow from a transversally thermalized fireball

U. Heinz and S. M. H. Wong

Physics Department, The Ohio State University, Columbus, Ohio 43210 (Received 6 May 2002; published 18 July 2002)

The agreement of elliptic flow data at RHIC at central rapidity with the hydrodynamic model has led to the conclusion of very rapid thermalization. This conclusion is based on the intuitive argument that hydrodynamics, which assumes instantaneous local thermalization, produces the largest possible elliptic flow values and that the data seem to saturate this limit. We here investigate the question whether incompletely thermalized viscous systems may actually produce more elliptic flow than ideal hydrodynamics. Motivated by the extremely fast primordial longitudinal expansion of the reaction zone, we investigate a toy model that exhibits thermalization only in the transverse directions but undergoes collisionless free-streaming expansion in the longitudinal direction. For collisions at RHIC energies, elliptic flow results from the model are compared with those from hydrodynamics. With the final particle yield and  $k_{\perp}$  distribution fixed, the transversally thermalized model is shown not to be able to produce the measured amount of elliptic flow. This investigation provides further support for very rapid local kinetic equilibration at RHIC. It also yields interesting novel results for the elliptic flow of massless particles such as direct photons.

DOI: 10.1103/PhysRevC.66.014907

PACS number(s): 25.75.Ld, 24.10.Nz, 24.85.+p

#### I. INTRODUCTION

Recently the STAR collaboration showed [1-3] that the  $k_{\perp}$  and centrality dependence of the measured elliptic flow coefficient  $v_2$  of pions and protons at the Relativistic Heavy Ion Collider (RHIC) largely agreed with those generated from the hydrodynamic model simulation of heavy ion collisions [4-7]. This was subsequently confirmed by data from the PHENIX [8] and PHOBOS [9] collaborations.  $v_2$  is the second coefficient of an azimuthal Fourier expansion of the transverse momentum spectrum around the beam axis [10] and, for collisions between identical nuclei, is the lowest nonzero anisotropic flow coefficient at midrapidity. Given that the main prerequisite of the hydrodynamic model is complete local thermal equilibrium, which requires very intense rescattering among the matter constituents, it was argued that hydrodynamics should give the largest possible elliptic flow, and the observation that the data saturate the hydrodynamic limit was taken as evidence that thermalization must be very fast at RHIC for central and semicentral collisions [6,7,11].

Elliptic flow requires reinteractions within the produced matter as a mechanism for transferring the initial spatial deformation of the reaction zone in noncentral collisions onto momentum space. It is thus plausible to expect that the largest elliptic flow signal is produced in the hydrodynamic limit, i.e., in the limit of infinite rescattering rates [6,7,11]; however, a proof for this hypothesis has not yet been found. In hydrodynamics, collective flow is generated by pressure gradients, and flow anisotropies such as elliptic flow require anisotropic pressure gradients in the plane transverse to the beam direction. The longitudinal pressure plays a less visible role: the work done by the longitudinal pressure reduces the transverse energy  $dE_T/dy$  and thereby, for given initial conditions for the energy density deposited in the reaction zone, the amount of transverse flow at freeze-out. If there were no longitudinal pressure, more of the initially deposited energy would go into the transverse directions, leading to flatter transverse momentum spectra. It is, at least in principle, conceivable that this could also lead, at fixed radial flow, to larger anisotropies of the transverse collective flow. If this were indeed the case and, at fixed slope of the angle-averaged spectra, larger values of  $v_2$  could be generated in this way than within the usual hydrodynamic, the early thermalization argument would break down since the RHIC data [1–3] would no longer saturate the theoretical prediction from such a model.

To explore this hypothetical possibility, we study in this paper a toy model which assumes that, at high collision energies, during the earliest collision stages the longitudinal momenta of the produced particles do not thermalize, but that the strong initial longitudinal expansion is instead dominated by collisionless free-streaming. The initial transverse momenta are much smaller and assumed to thermalize quickly. This results in a system with local transverse, but zero longitudinal pressure. Due to the masslessness of the partons created in the reaction zone the trace of the energy momentum tensor vanishes; in the absence of longitudinal pressure  $P_{\parallel}$ , the transverse pressure  $P_{\perp}$  must thus be related to the initial energy density  $\varepsilon = T^{00}$  by  $\varepsilon = 2P_{\perp}$  (instead of the usual  $\varepsilon = 3P$ ). This results in a much stiffer equation of state for the transverse dynamics (see Sec. II B), giving more flow and possibly stronger flow anisotropies. Of course, if the transverse momenta thermalize, the longitudinal momenta should do so eventually, too. We assume in our model that when this happens the flow anisotropies have already almost reached their asymptotic values [12].

Similar to the hydrodynamic simulations [4–7], we assume that the longitudinal free-streaming dynamics is boost invariant [14] and correspondingly concentrate on the central rapidity region. Starting from a kinetic description of the longitudinally free streaming but transversally thermalized gluonic system [which we call "transversally thermalized model" (TTHM)], we derive a set of macroscopic evolution equations (TTHM equations) which we solve with appropriate initial conditions. The impact-parameter dependence of the initial conditions is handled in the same way as in the hydrodynamic simulations. When comparing our TTHM solutions to those from the ideal hydrodynamic model (HDM) we retune the initial conditions in such a way that for central collisions (b=0 fm) roughly the same multiplicities and spectral slopes at midrapidity are obtained. We then compare the momentum-space anisotropy  $v_2$  in the two approaches, as a function of transverse momentum  $k_{\perp}$  and impact parameter *b*.

# II. CONSTRUCTING A TRANSVERSALLY THERMALIZED SYSTEM

### A. The phase-space distribution function

We consider a gluon dominated system at a short time after nuclear contact. We assume that at that time the transverse gluon momenta are already thermalized but that the system is still free streaming along the beam direction. The system is required to possess longitudinal boost invariance, reflecting a boost-invariant primary-particle production mechanism [14]. For the easiest implementation we assume that the collision energy is so high that the colliding nuclei are Lorentz contracted to two infinitesimally thin sheets in z direction and that all produced particles point back to z=t=0. Their longitudinal momenta and coordinates thus satisfy  $z=u_zt$  or  $k_0z=k_zt$  where  $u_z=k_z/k_0$ . This automatically identifies [14] the rapidity

$$y = \frac{1}{2} \ln \left( \frac{k_0 + k_z}{k_0 - k_z} \right)$$
(2.1)

of the produced particles with their space-time rapidity

$$\eta = \frac{1}{2} \ln \left( \frac{t+z}{t-z} \right). \tag{2.2}$$

We thus make the ansatz

$$f(\mathbf{k}, \mathbf{x}, t) = \delta(k_0 z - k_z t) \widetilde{h}(\mathbf{k}, \mathbf{x}, t).$$
(2.3)

Longitudinal free streaming is implemented by the condition

$$\left(k_0\frac{\partial}{\partial t} + k_z\frac{\partial}{\partial z}\right)f(\mathbf{k},\mathbf{x},t) = 0.$$
(2.4)

Since the  $\delta$ -function factor solves this equation trivially, this implies a constraint for  $\tilde{h}$ . The latter is most conveniently expressed by using an alternate coordinate system (we hereafter make explicit use of the masslessness of the produced gluon degrees of freedom, by setting  $m_{\perp} = \sqrt{m^2 + k_{\perp}^2} \rightarrow k_{\perp}$ ),

$$t = \tau \cosh \eta, \quad z = \tau \sinh \eta,$$
 (2.5)

$$k_0 = k_\perp \cosh y, \quad k_z = k_\perp \sinh y. \tag{2.6}$$

In these coordinates the ansatz Eq. (2.3) becomes

$$f(\mathbf{k},\mathbf{x},t) = \frac{\delta(\eta - y)}{k_{\perp}\tau\cosh(\eta - y)}\widetilde{h}(\mathbf{k}_{\perp},\mathbf{x}_{\perp},\eta - y,\tau), \quad (2.7)$$

where the dependence of  $\tilde{h}$  on only  $\mathbf{k}_{\perp}$  and the difference  $\eta - y$  results from the requirement of longitudinal boost invariance. We implement transverse thermalization by the ansatz

$$\widetilde{h}(\mathbf{k}_{\perp}, \mathbf{x}_{\perp}, \eta - y, \tau) = \frac{\tau_0 k_{\perp}}{\gamma_{\perp}} \cosh(\eta - y) \frac{1}{e^{k \cdot u/T} - 1},$$
(2.8)

where  $u^{\mu} = \gamma_{\perp}(\cosh \eta, \mathbf{v}_{\perp}, \sinh \eta)$  is the flow vector with Bjorken longitudinal flow velocity  $v_z = \tanh \eta [14]$  and transverse flow velocity  $\mathbf{v}_{\perp} = \mathbf{v}_{\perp}(\mathbf{x}_{\perp}, \tau)$   $(\gamma_{\perp}^{-1} = \sqrt{1 - v_{\perp}^2})$ .  $T = T(\mathbf{x}_{\perp}, \tau)$  is the temperature characterizing the thermalized transverse momentum spectrum. In the local rest frame *T* is a function of the rest frame coordinates  $\mathbf{x}_{\perp}^*$  and  $\tau^* = t^*$ , which after the standard [15] successive transverse and longitudinal boosts to the laboratory frame, with  $\mathbf{v}_{\perp}$  and  $v_z = \tanh \eta$ , respectively, are related to the laboratory variables by

$$t^* = \gamma_{\perp}(t \cosh \eta - z \sinh \eta - \mathbf{v}_{\perp} \cdot \mathbf{x}_{\perp}) = \gamma_{\perp}(\tau - \mathbf{v}_{\perp} \cdot \mathbf{x}_{\perp}) = u \cdot x,$$
(2.9)

$$\begin{aligned} x_{\perp\parallel}^{*} &= \gamma_{\perp} [x_{\perp\parallel} - v_{\perp} (t \cosh \eta - z \sinh \eta)] \\ &= \gamma_{\perp} (x_{\perp\parallel} - v_{\perp} \tau) = -m \cdot x, \end{aligned}$$
(2.10)

$$x_{\perp\perp}^* = x_{\perp\perp} . \tag{2.11}$$

The additional  $\parallel$  and  $\perp$  subscripts denote the components of  $\mathbf{x}_{\perp}$  parallel and perpendicular to  $\mathbf{v}_{\perp}$ , respectively, and the vector  $m^{\mu}$  is defined in Eq. (2.15) below. The sequence of boosts described here agrees with the standard choice [15] and gives the simplest structure for the set of four-vectors needed for the decomposition of the energy momentum tensor (see Sec. II B). As one can see from Eqs. (2.9)–(2.11), the dependence of T on  $\mathbf{x}_{\perp}^*$  and  $t^*$  in the local rest frame translates into a dependence on  $\mathbf{x}_{\perp}$  and  $\tau$  in the laboratory frame.

By virtue of the factor  $\delta(\eta - y)$ , the argument of the Bose distribution in Eq. (2.8) reduces in the local rest frame to  $k \cdot u = k_{\perp}$ , showing explicitly that the system is only transversally thermalized. Combining Eqs. (2.7) and (2.8) and exploiting the  $\delta$  function, the distribution function simplifies to

$$f(\mathbf{k}, \mathbf{x}, t) = \frac{\tau_0}{\tau} \,\delta(\eta - y) h(\mathbf{k}_{\perp}, \mathbf{x}_{\perp}, t),$$
$$h(\mathbf{k}_{\perp}, \mathbf{x}_{\perp}, t) = \frac{1}{\gamma_{\perp}} \frac{1}{e^{k \cdot u/T} - 1}.$$
(2.12)

In the new coordinates, f satisfies the relativistic transport equation

$$\begin{pmatrix} k_{\perp} \cosh(\eta - y) \frac{\partial}{\partial \tau} - \frac{k_{\perp}}{\tau} \sinh(\eta - y) \frac{\partial}{\partial \eta} + \mathbf{k}_{\perp} \cdot \nabla_{\perp} \end{pmatrix} f(\mathbf{k}, \mathbf{x}, t)$$

$$= \mathcal{C}(\mathbf{k}, \mathbf{x}, t),$$
(2.13)

where the collision term C is responsible for keeping the transverse momenta thermalized while not changing any of the longitudinal momenta. Due to the implemented free-streaming properties we have

$$\left(\cosh(\eta - y)\frac{\partial}{\partial \tau} - \sinh(\eta - y)\frac{1}{\tau}\frac{\partial}{\partial \eta}\right) \left(\frac{\tau}{\tau_0}\delta(\eta - y)\right) = 0$$

such that Eq. (2.13) reduces to the following kinetic equation for the transverse distribution function:

$$\left(k_{\perp}\frac{\partial}{\partial\tau} + \mathbf{k}_{\perp} \cdot \boldsymbol{\nabla}_{\perp}\right) h(\mathbf{k}_{\perp}, \mathbf{x}_{\perp}, \tau) = \mathcal{C}(\mathbf{k}, \mathbf{x}, t). \quad (2.14)$$

The space-time evolution of  $h(\mathbf{k}_{\perp}, \mathbf{x}_{\perp}, \tau)$  is entirely due to the collisions and collective transverse expansion, but decoupled from the boost invariant longitudinal expansion. The collisions must be sufficiently dominant over the transverse expansion to maintain the equilibrium form of  $h(\mathbf{k}_{\perp}, \mathbf{x}_{\perp}, \tau)$ given above.

We are aiming at a macroscopic description on the basis of the differential conservation laws for energy and momentum, similar to hydrodynamics. This requires the construction of  $T^{\mu\nu}$  from the distribution functions. Due to the different microscopic physics in the longitudinal and transverse directions implemented in our model, there are now more vectors required to construct a complete set of tensors with respect to which  $T^{\mu\nu}$  should be decomposed. The set of fourvector fields that we need are

$$u^{\mu}(\mathbf{x}_{\perp}, \eta, \tau) = \gamma_{\perp}(\cosh \eta, \mathbf{v}_{\perp}, \sinh \eta),$$
  

$$n^{\mu}(\mathbf{x}_{\perp}, \eta, \tau) = (\sinh \eta, 0, 0, \cosh \eta),$$
  

$$m^{\mu}(\mathbf{x}_{\perp}, \eta, \tau) = \gamma_{\perp}(v_{\perp} \cosh \eta, \hat{\mathbf{v}}_{\perp}, v_{\perp} \sinh \eta). \quad (2.15)$$

They arise from the following local rest frame vectors by successive boosts with  $\mathbf{v}_{\perp}$  and  $v_z$  as described above:

$$u^{\mu}(\mathbf{x}_{\perp}, \eta, \tau) = (1, 0, 0, 0),$$
  

$$n^{\mu}(\mathbf{x}_{\perp}, \eta, \tau) = (0, 0, 0, 1),$$
  

$$m^{\mu}(\mathbf{x}_{\perp}, \eta, \tau) = (0, \hat{\mathbf{v}}_{\perp}, 0).$$
(2.16)

One is timelike,  $u^2=1$ , while the other two are spacelike,  $n^2=m^2=-1$ . They are mutually orthogonal:  $u \cdot n = u \cdot m$  $= n \cdot m = 0$ . With these vectors, the distribution can be expressed in a form that is explicitly boost invariant under longitudinal boosts,

$$f(\mathbf{k}, \mathbf{x}, t) = \frac{\tau_0}{\tau} \frac{\delta(k \cdot n)(k \cdot u - v_\perp k \cdot m)}{e^{k \cdot u/T} - 1}.$$
 (2.17)

#### B. Energy-momentum tensor and conservation laws

Based on the structure of Eq. (2.17) for the phase-space distribution function we want to derive a macroscopic set of dynamical equations based on the conservation laws for en-

ergy and momentum. In terms of the distribution function f, the energy-momentum tensor is given by

$$T^{\mu\nu}(\mathbf{x},t) = \nu_g \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \frac{k^{\mu}k^{\nu}}{k^0} f(\mathbf{k},\mathbf{x},t), \qquad (2.18)$$

where  $\nu_g = 2 \times 8 = 16$  is the degeneracy factor for color and helicity of the gluons. From the preceding section we know that  $f(\mathbf{k}, \mathbf{x}, t)$  depends on the three four-vectors  $u^{\mu}$ ,  $n^{\mu}$ , and  $m^{\mu}$ . The most general form of the tensor  $T^{\mu\nu}$  thus reads

$$T^{\mu\nu} = A u^{\mu} u^{\nu} + B n^{\mu} n^{\nu} + C m^{\mu} m^{\nu} + D (u^{\mu} n^{\nu} + n^{\mu} u^{\nu}) + F (u^{\mu} m^{\nu} + m^{\mu} u^{\nu}) + G (n^{\mu} m^{\nu} + m^{\mu} n^{\nu}) + H g^{\mu\nu}.$$
(2.19)

Since the gluons are massless, the trace of  $T^{\mu\nu}$  vanishes,

$$T^{\mu}_{\ \mu} = A - B - C + 4H = 0. \tag{2.20}$$

In the local rest frame we have

$$T_{00}^* = A + H = \varepsilon,$$
  
 $T_{xx}^* = T_{yy}^* = -H = P_{\perp},$   
 $T_{zz}^* = B - H = P_z,$  (2.21)

where  $\varepsilon$  is the energy density and  $P_{\perp}$  and  $P_z$  are the transverse and longitudinal pressures, respectively. In a general frame, all the Lorentz scalar coefficients in Eq. (2.19) can be found by contracting  $T^{\mu\nu}$  with all pairwise combinations of the four-vectors  $u^{\mu}$ ,  $n^{\mu}$ , and  $m^{\mu}$ . We will denote these contractions by  $(uTn) \equiv u_{\mu}T^{\mu\nu}n_{\nu}$ , etc. The details of this calculation are given in Appendix A where we find

$$(uTu) = A + H = \frac{\tau_0}{\tau} \nu_g \frac{\pi^2 T^4}{60} = \varepsilon,$$
 (2.22)

$$(uTm) = -F = -\frac{\tau_0}{\tau} \nu_g \frac{\pi^2 T^4}{120} \nu_\perp = -\frac{\nu_\perp}{2} \varepsilon, \quad (2.23)$$

$$(mTm) = C - H = \frac{\tau_0}{\tau} \nu_g \frac{\pi^2 T^4}{120} = \frac{1}{2} \varepsilon,$$
 (2.24)

$$(uTn) = -D = 0, (2.25)$$

$$(nTn) = B - H = 0 = P_z,$$
 (2.26)

$$(nTm) = G = 0.$$
 (2.27)

From this and Eq. (2.21) we see that

$$B = H = -P_{\perp}, \quad A = \varepsilon + P_{\perp}, \quad (2.28)$$

and the tracelessness condition gives

$$\varepsilon + P_{\perp} + P_{\perp} - \frac{1}{2} \ \varepsilon - 3P_{\perp} = \frac{1}{2} \ \varepsilon - P_{\perp} = 0.$$
 (2.29)

Therefore the equation of state is

$$=2P_{\perp}$$
. (2.30)

Combining this with the expression for (mTm) one finds C = 0. The expression for the energy momentum tensor is then

ε

$$T^{\mu\nu} = (\varepsilon + P_{\perp})u^{\mu}u^{\nu} - P_{\perp}n^{\mu}n^{\nu} - P_{\perp}g^{\mu\nu} + P_{\perp}v_{\perp}(u^{\mu}m^{\nu} + m^{\mu}u^{\nu}).$$
(2.31)

For completeness and later use we also give the expression for the gluon density in the local comoving frame,

$$n_g = u \cdot j = \frac{\tau_0}{\tau} \nu_g \frac{\zeta(3)T^3}{2\pi^2}, \qquad (2.32)$$

where  $j^{\mu}(x)$  is the gluon number current,

$$j^{\mu}(\mathbf{x},t) = \nu_g \int \frac{d^3 \mathbf{k}}{(2\pi)^3} \frac{k^{\mu}}{k^0} f(\mathbf{k},\mathbf{x},t).$$
(2.33)

We now proceed to derive the macroscopic evolution equations. We first rewrite the conservation laws

$$\partial_{\nu}T^{\mu\nu} = 0 \tag{2.34}$$

in  $(\eta, \tau)$  coordinates

$$\cosh \eta \left( \frac{\partial T^{0\mu}}{\partial \tau} + \frac{1}{\tau} \frac{\partial T^{z\mu}}{\partial \eta} \right) - \sinh \eta \left( \frac{\partial T^{z\mu}}{\partial \tau} + \frac{1}{\tau} \frac{\partial T^{0\mu}}{\partial \eta} \right) + \nabla_{\perp}^{i} T^{i\mu}$$
$$= 0, \qquad (2.35)$$

where the sum over *i* goes over the two transverse directions. When writing these out explicitly in terms of  $\varepsilon$ ,  $P_{\perp}$ , and the longitudinal and transverse flow velocities, using Eqs. (2.31) and (2.16), one finds that all explicit dependence on  $\eta$  disappears. Boost-invariant initial conditions (i.e.  $\eta$ -independent initial expressions for  $\varepsilon$  and  $P_{\perp}$  and the structure (2.16) for the vectors  $u^{\mu}$ ,  $n^{\mu}$ , and  $m^{\mu}$ ) are thus preserved by the macroscopic evolution equations.

For the further analysis we can thus concentrate on the dynamics in the central transverse plane at  $z = \eta = 0$ . There we have

$$u^{\mu} = \gamma_{\perp}(1, \mathbf{v}_{\perp}, 0), \quad \frac{\partial u^{\mu}}{\partial \eta} = \gamma_{\perp}(0, 0, 0, 1), \quad (2.36)$$

$$n^{\mu} = (0,0,0,1), \quad \frac{\partial n^{\mu}}{\partial \eta} = (1,0,0,0),$$
 (2.37)

$$m^{\mu} = \gamma_{\perp}(v_{\perp}, \hat{\mathbf{v}}_{\perp}, 0), \frac{\partial m^{\mu}}{\partial \eta} = \gamma_{\perp}(0, \hat{\mathbf{v}}_{\perp}, v_{\perp}). \quad (2.38)$$

Using the  $\eta$  independence of  $\varepsilon$  and  $P_{\perp}$  we find

$$\frac{\partial T^{z\mu}}{\partial \eta} = \gamma_{\perp}^{2} (\varepsilon + P_{\perp}) (\delta^{0\mu} + v_{\perp}^{i} \delta^{i\mu}) - P_{\perp} \delta^{0\mu} + \gamma_{\perp}^{2} P_{\perp} (2v_{\perp}^{2} \delta^{\mu 0} + (1 + v_{\perp}^{2})v_{\perp}^{i} \delta^{i\mu}) = T^{00} \delta^{0\mu} + T^{0i} \delta^{i\mu} = T^{0\mu}, \quad (2.39)$$

and the equations of motion (2.35) at  $z = \eta = 0$  reduce to

$$\frac{\partial T^{00}}{\partial \tau} + \frac{T^{00}}{\tau} + \boldsymbol{\nabla}_{\perp}^{j} T^{j0} = 0, \qquad (2.40)$$

$$\frac{\partial T^{0i}}{\partial \tau} + \frac{T^{0i}}{\tau} + \boldsymbol{\nabla}_{\perp}^{j} T^{ji} = 0.$$
 (2.41)

Using the equation of state (2.30) to eliminate  $\varepsilon$ , these three equations completely determine the three unknown functions  $P_{\perp}$  and  $\mathbf{v}_{\perp}$ . Since gluons possess no conserved charges, no further current conservation laws need to be considered.

#### **III. HYDRODYNAMICS**

For later comparison we here shortly review the parallel procedure for ideal hydrodynamics based on a locally fully thermalized phase-space distribution function

$$f(\mathbf{k}, \mathbf{x}, t) = \frac{1}{e^{k \cdot u/T} - 1}.$$
(3.1)

In this case the energy-momentum tensor has the ideal fluid decomposition

$$T^{\mu\nu} = (\varepsilon + P)u^{\mu}u^{\nu} - Pg^{\mu\nu}, \qquad (3.2)$$

and for a longitudinally boost-invariant system  $u^{\mu}$  has the same form as in Eq. (2.15). The equation of state is in this case

$$\varepsilon = 3P = \nu_g \frac{\pi^2 T^4}{30},\tag{3.3}$$

and the gluon density in the local rest frame is

$$n_g = \nu_g \frac{\zeta(3)T^3}{\pi^2}.$$
 (3.4)

Note that the energy per particle

$$\frac{\varepsilon}{n_g} = \frac{\pi^4}{30\zeta(3)} T \approx 2.7T \tag{3.5}$$

agrees exactly with the corresponding expression in the TTHM, see Eqs. (2.22) and (2.32).

The pressure is now locally isotropic. With longitudinally boost-invariant ( $\eta$ -independent) initial conditions for  $\varepsilon$  and P the ansatz Eq. (2.16) for the flow velocity is preserved in time by the equations of motion  $\partial_{\nu}T^{\mu\nu} = 0$  [14], and the latter become

$$\frac{\partial T^{00}}{\partial \tau} + \frac{T^{00}}{\tau} + \boldsymbol{\nabla}_{\perp}^{j} T^{j0} = -\frac{P}{\tau}$$
(3.6)

$$\frac{\partial T^{0i}}{\partial \tau} + \frac{T^{0i}}{\tau} + \boldsymbol{\nabla}_{\perp}^{j} T^{ji} = 0.$$
(3.7)

The main difference between these hydrodynamical equations and our TTHM toy model equations (2.40) and (2.41) is the pressure term on the right hand side of Eq. (3.6) which is absent in Eq. (2.40). It indicates the longitudinal work done by the isotropic pressure in hydrodynamics; in the TTHM model there is no longitudinal pressure that could do work in the longitudinal direction.

### **IV. INITIAL CONDITIONS**

We consider Au+Au collisions at RHIC energies. The initial energy density distribution in the transverse plane as a function of impact parameter **b** is taken to be proportional to the density of binary nucleon collisions in the transverse plane, calculated from the normalized nuclear thickness function,

$$T_A(\mathbf{x}_{\perp}) = \int \rho_A(\mathbf{x}_{\perp}, z) dz, \quad \int d^2 x_{\perp} T_A(\mathbf{x}_{\perp}) = A. \quad (4.1)$$

For the nuclear number density we took the Woods-Saxon parametrization

$$\rho_A(\mathbf{x}_{\perp}) = \frac{\rho_0}{e^{(r-R_0)/\xi} + 1} \tag{4.2}$$

with  $R_0 = 1.14A^{1/3}$  fm and  $\xi = 0.54$  fm. For collisions between the nuclei of masses *A* and *B* the number of binary nucleon collisions per unit area in the transverse plane is

$$\frac{dN(\mathbf{x}_{\perp},\mathbf{b})}{d^2 x_{\perp}} = \sigma_0 T_A(\mathbf{x}_{\perp}) T_B(\mathbf{b} - \mathbf{x}_{\perp}), \qquad (4.3)$$

where  $\sigma_0 = 40$  mb is the total inelastic nucleon-nucleon cross section. We assume that the generated energy density is proportional to this density. We performed two classes of calculations: in the first class we used the same maximum initial energy density as the authors of Refs. [6,7] [ $\varepsilon_0 = e_0$  $\equiv 23.0$  GeV/fm<sup>3</sup> for central (b=0 fm)] collisions at an initial time  $\tau_0 = 0.6$  fm/c) in order to compare the HDM and TTHM results for similar initial conditions. In the second class of simulations we retuned the initial conditions in TTHM such that the same multiplicity density dN/dy and the same slope for the transverse momentum spectrum at midrapidity as in HDM is obtained. Both sets of results will be discussed in the following section.

## V. ELLIPTIC FLOW IN A TRANSVERSALLY THERMALIZED SYSTEM

#### A. Freeze-out prescription

Just as the HDM, our TTHM model describes the time evolution of macroscopic thermodynamic quantities that must be converted to particle spectra before one can compare with experiments. In order to do so we use the well-known Cooper-Frye prescription [16] that gives the particle spectrum in terms of an integral over the phase-space distribution function along a so-called freeze-out hypersurface  $\Sigma(x)$ ,

$$E\frac{dN}{d^3k} = \frac{dN}{dyk_{\perp}dk_{\perp}d\phi} = \frac{\nu_g}{(2\pi)^3} \int_{\Sigma} k \cdot d^3\Sigma(x)f(x,k).$$
(5.1)



FIG. 1.  $v_2$  as a function of impact parameter at two freeze-out energy densities  $\varepsilon_{fo}$ . The solid lines are from our TTHM model while the dashed lines are from hydrodynamic simulations with identical initial conditions.

 $d^{3}\Sigma_{\mu}(x)$  is the four-vector integral measure normal to the hypersurface, and in the TTHM case f(x,k) is a transversally thermalized distribution function of the form (2.12), with flow velocity  $u^{\mu}$  evaluated along the freeze-out hypersurface  $\Sigma$  and temperature *T* calculated via Eq. (2.22) from the energy density on  $\Sigma$ . Longitudinal boost invariance dictates freeze-out along a surface of fixed longitudinal proper time  $\tau_{f}(\mathbf{x}_{\perp})$ , and we can write

$$\Sigma_{\mu}(x) = (\tau_{\rm f}(\mathbf{x}_{\perp}) \cosh \eta, \mathbf{x}_{\perp}, \tau_{\rm f}(\mathbf{x}_{\perp}) \sinh \eta).$$
(5.2)

This gives

$$k \cdot d^3 \Sigma = (k_\perp \cosh(y - \eta) - \mathbf{k}_\perp \cdot \nabla_\perp \tau_{\rm f}) \tau_{\rm f} d \, \eta d^2 x_\perp \,.$$
 (5.3)

Of course, the  $\eta$  integration is trivial due to the factor  $\delta(\eta - y)$  in the distribution function in Eq. (2.12).

#### B. TTHM vs HDM for identical initial conditions

In Fig. 1 we compare results for the elliptic flow from the TTHM and HDM models for identical initial conditions, taken from [6,7]. Since the temperature parameter T has a different meaning in our model than usual (only the transverse momenta are thermalized), we enforce "freeze-out" (i.e., we stop the dynamical evolution and calculate the spectra and momentum space anisotropies) along a surface of constant energy density  $\varepsilon_{\rm f}$ . We show results for two values of this parameter both of which lie in the quark-hadron transition region. Since hadronization certainly involves longitudinal momentum exchange, it will violate our assumption of longitudinal free-streaming, and we should definitely not follow the TTHM dynamics beyond the hadronization point. Of course, by truncating the dynamics at such high freeze-out energy densities we forfeit the possibility of comparing our results directly with experiment. However, a large fraction of the elliptic flow should already have developed at this point



FIG. 2. Gluon  $k_{\perp}$  distribution for central collisions (b=0 fm) and two different values for the freeze-out energy density, for identical initial conditions. The solid lines are for our model while the dashed lines are for hydrodynamics.

[5,17], and we can still make a meaningful comparison between the hydrodynamic evolution of a fully thermalized quark-gluon plasma and that of a longitudinally freestreaming gluonic system with only transverse thermalization, by comparing the gluon spectra at this common "final" energy density  $\varepsilon_{\rm f}$ .

Figure 1 shows  $v_2$  as a function of the impact parameter b.  $v_2(b) = \langle \cos(2\phi) \rangle (b)$  is computed from the  $k_{\perp}$ -integrated gluon spectrum at freeze-out as

$$v_{2}(b) = \frac{\int k_{\perp} dk_{\perp} d\phi \cos(2\phi) \frac{dN}{dyk_{\perp} dk_{\perp} d\phi}(b)}{\int k_{\perp} dk_{\perp} d\phi \frac{dN}{dyk_{\perp} dk_{\perp} d\phi}(b)}$$
$$= \frac{\int d\phi \cos(2\phi) \frac{dN}{dyd\phi}(b)}{\int d\phi \frac{dN}{dyd\phi}(b)}.$$
(5.4)

Figure 2 shows that, for identical initial conditions, the TTHM dynamics generates a larger momentum-space asymmetry  $v_2$  than hydrodynamics. We also see that  $v_2$  has not yet fully saturated at these values of  $\varepsilon_f$ ; the reason for this can be found in Table I which shows that, with the very hard equation of state and initial conditions used here, the above values of  $\varepsilon_f$  are reached quite early, before the initial spatial deformation has been fully eliminated (see also Fig. 1 in Ref. [18]).

The larger values of  $v_2$  from TTHM, however, come also with much flatter transverse momentum spectra, as shown in Fig. 2. Part of this flattening is due to a higher freeze-out temperature in the TTHM. By comparing Eqs. (2.22) and (3.3) one sees at the same energy density  $\varepsilon_f$  the temperatures in the two models are related by  $T_{f,T} = (2\tau_f/\tau_0)^{1/4}T_{f,H}$ . With

TABLE I. Rapidity densities at y=0 and final times  $\tau_f$  for central collisions with identical initial conditions (see text) in the TTHM and HDM simulations.

	TTHM		HDM	
$\epsilon_{\rm f}~(GeV/fm^3)$	dN/dy	$ au_{\mathrm{f}}(\mathrm{fm}/c)$	dN/dy	$ au_{ m f}({ m fm}/c)$
0.6	780	6.95	944	5.97
0.8	780	6.48	940	5.36

the  $\tau_{\rm f}$  values given in Table I, the TTHM freeze-out temperature is about a factor 2 higher than the HDM one. However, the inverse slopes of the spectra shown in Fig. 2 differ by more than this factor 2 (the difference is closer to a factor 2.5–2.6). This implies that TTHM also creates somewhat stronger radial flow than HDM which, in view of its harder equation of state, is not unexpected.

The initial conditions for the hydrodynamic model (in particular, the initial energy density) were tuned [6] to reproduce the measured final multiplicity densities  $dN_{ch}/dy$  and spectral slopes in central Au+Au collisions at RHIC; clearly, with these same initial conditions, the transversally thermalized model TTHM will no longer be anywhere close to these data. The larger  $v_2$  values from TTHM in Fig. 1 therefore cannot be compared with the RHIC data, and the comparison with HDM in Fig. 1 is misleading. For a meaningful comparison the initial conditions for TTHM should first be adjusted in such a way that at least the multiplicity densities and spectral slopes for central collisions are similar in both models. This will be our next step.

Before doing so, let us shortly comment on the rapidity densities dN/dy and collision durations listed in Table I. For identical initial conditions, the TTHM gives considerably lower multiplicity densities. This is mostly due to the higher freeze-out temperature in the TTHM, which implies that each gluon carries more energy [see Eq. (3.5)], so the same freeze-out energy density  $\varepsilon_{\rm f}$  translates into fewer gluons. This argument overestimates the difference between the models, though; the actual difference is smaller since in the HDM the calculation of dN/dy involves an integration over space-time rapidity  $\eta$  which can be interpreted as taking an average over fireballs with different flow rapidities  $\eta \neq y$  and correspondingly reduced (redshifted) effective temperatures  $T_{\rm eff} = T_{\rm f}/\cosh(\eta - y)$ .

Longitudinal boost invariance implies that the total transverse energy per unit rapidity,  $dE_T/dy$ , is independent of y. Due to the absence of longitudinal pressure, no longitudinal work is done in the TTHM and  $dE_T/dy$  is a constant of motion. This is different from the HDM where work done by the longitudinal pressure reduces  $dE_T/dy$  with time. (Both statements were checked numerically.) Table I shows that, within numerical accuracy, the gluon multiplicity per unit rapidity, dN/dy, is also constant in time. While in ideal hydrodynamics (HDM) this is a simple consequence of entropy conservation (for boost-invariant systems dS/dy is a constant of motion [14], and for massless particles the entropy per particle is independent of temperature), there is no such simple reason for this observation in the TTHM which is far



FIG. 3. The freeze-out time  $\tau_f(x,y)$  as a function of the transverse coordinates *x* and *y*, respectively, for a freeze-out energy density  $\varepsilon_f = 0.6 \text{ GeV/fm}^3$ . The solid (dashed) lines are for TTHM (HDM), respectively. The upper set of curves is for central collisions (*b*=0 fm) where the freeze-out surface is azimuthally symmetric. The lower set of curves corresponds to *b*=7 fm where we present cuts along the *x* and *y* axes, as indicated, to show the spatial azimuthal anisotropy at freeze-out. The impact parameter **b** points in *x* direction.

from local equilibrium. Detailed checks revealed that, within numerical accuracy, in the TTHM not only the transverse energy per particle,  $E_T/N = (dE_T/dy)/(dN/dy)$ , but, in fact, also the entire  $k_{\perp}$  spectrum of the gluons is completely time-independent.

If the system were transversally homogeneous and expanded only longitudinally, this would not be surprising: in the absence of transverse gradients the conservation law Eq. (2.40) for  $T^{00}$  simplifies to

$$\frac{\partial(\tau T^{00})}{\partial \tau} = 0 \tag{5.5}$$

with  $T^{00} = \varepsilon \cosh^2 \eta$ . In TTHM a constant product  $\varepsilon \tau$  implies a constant temperature [see Eq. (2.22)], which, in the absence of transverse flow, then leads to a time-independent transverse momentum spectrum. In our case, however, the system expands in the transverse direction and cools. Our studies show that, in the longitudinal rest frame, the resulting loss in thermal transverse energy per particle is exactly compensated by the gain in transverse collective motion energy, in a way that exactly preserves the shape of the transverse momentum distribution. Although one should think that there must be a simple reason for this intriguing behavior, we have not been able to find a simple analytical proof and can thus only present the numerical evidence.

The longer freeze-out times in the TTHM simulations (see Table I) are a reflection of the lack of longitudinal pressure. In the HDM the pressure performs longitudinal work and thus causes a more rapid decrease of the energy density with time than in the TTHM. This is further elucidated in Fig. 3 where we show the freeze-out time  $\tau_f(x,y)$  as a function of position in the transverse plane. One sees that, for identical initial conditions, the HDM simulations lead to universally earlier freeze-out. This figure also shows that for noncentral Pb+Pb collisions (b=7 fm) the source at freeze-out is still larger in the y direction perpendicular to the reaction plane than in the reaction plane; the initial out-of-plane deforma-



FIG. 4. The gluon  $k_{\perp}$  distribution from hydrodynamics (dashed line, with the parameters as given in the text) and TTHM (other lines). The TTHM curves correspond to the following sets of parameters. Dotted line:  $\tau_0 = 6 \text{ fm/}c$ ,  $\varepsilon_0 = e_0/15$ . Dot-dashed line:  $\tau_0 = 7 \text{ fm/}c$ ,  $\varepsilon_0 = e_0/18$ . Solid line:  $\tau_0 = 7.8 \text{ fm/}c$ ,  $\varepsilon_0 = e_0/20$ . Here  $e_0 = 23.0 \text{ GeV/fm}^3$  is the initial central energy density for b = 0 Au + Au collisions in the HDM.

tion of the reaction zone thus has not yet fully disappeared when the dynamical evolution was stopped. This explains why the elliptic flow had not yet saturated, see Fig. 1 and Fig. 9 below.

#### C. Retuning the initial conditions for TTHM

Our TTHM model does not include the hadronization of gluons and hence does not allow us to compute hadron spectra; hence we cannot directly compare the gluon multiplicity density dN/dy and their  $k_{\perp}$  distribution from TTHM at  $\varepsilon_{\rm f}$  to any data. However, the relation of these quantities to the observable hadron spectra after hadronization is not expected to depend on their dynamical history prior to reaching the freeze-out point  $\boldsymbol{\epsilon}_f.$  Thus, if we want to create TTHM solutions which are likely to lead, after hadronization, to hadron spectra with similar normalization and shape as the data, we can use the gluon rapidity densities and  $k_{\perp}$  spectra from existing hydrodynamic calculations with initial conditions that were tuned to real data in [6], evaluate them at the same value of  $\varepsilon_{\rm f}$  where we stop the TTHM evolution, and retune the TTHM initial conditions such that they reproduce these HDM gluon spectra. With the TTHM spectra thus tied down to the final state in central (b=0 fm) collisions from the HDM, we can then repeat our comparison of elliptic flow in the two models under more realistic boundary conditions.

For the common decoupling point we chose in both TTHM and HDM the value  $\varepsilon_f = 0.35$  GeV/fm<sup>3</sup>. This may appear somewhat low for still using only gluon degrees of freedom [with an ideal gas equation of state (EOS) in the HDM case], but it has the advantage that then the hydrodynamic model with the ideal massless EOS used here produces an elliptic flow  $v_2$  of roughly the same magnitude as that obtained for pions in Ref. [6] using a more realistic equation of state. Given the good agreement of those HDM calulations with the data [6], we can thus pretend to be comparing directly to data when comparing the TTHM results to our HDM reference.

With this freeze-out energy density, our HDM reference gives the gluon  $k_{\perp}$  (or  $m_{\perp}$ ) spectrum shown in Fig. 4, which



FIG. 5. Variation of the gluon  $k_{\perp}$  distribution with various TTHM model parameters. The solid and dashed lines repeat those from Fig. 2 for  $\varepsilon_f = 0.6$  GeV/fm<sup>3</sup> for comparison. The dot-dashed line is obtained by starting the system earlier at  $t_0 = 0.1$  fm/c. The long-dashed line has a lower initial energy density  $\varepsilon_i = \varepsilon_0/4$ . In each case the remaining parameters are held fixed at the values used in Fig. 2.

integrates to a gluon rapidity density at y=0 of  $dN/dy \approx 942$ . Given the large difference between the HDM and TTHM spectra for identical initial conditions shown in Fig. 2, it is perhaps not surprising that we found it rather difficult to reproduce the HDM spectrum within the TTHM: one needs to find a way to considerably lower the radial flow (since the freeze-out temperature is fixed by  $\varepsilon_{\rm f}$  and is much higher in TTHM than in HDM) and at the same time increase the normalization of the spectrum in order to raise dN/dy.

There are essentially only three parameters in TTHM that we can play with:  $\varepsilon_0$ ,  $\tau_0$  and  $\varepsilon_f$ . Examples of the effects of varying them are shown in Fig. 5. The lines from Fig. 2 corresponding to  $\varepsilon_f = 0.6 \text{ GeV/fm}^3$  are repeated here for comparison. From Fig. 2 and Table I we learned already that changing  $\varepsilon_{\rm f}$  has no influence on dN/dy and little effect on  $dN/dyk_{\perp}dk_{\perp}$ . This leaves us only with changes of the initial conditions  $\varepsilon_0$  and  $\tau_0$ . The dot-dashed curve in Fig. 5 shows that decreasing  $\tau_0$  from 0.6 to 0.1 fm/c goes in the wrong direction, by reducing dN/dy without having much effect on the slope of the  $k_{\perp}$  spectrum. Reducing the initial energy density  $\varepsilon_0$  by a factor of 4 produces the long-dashed line in the figure. Due to the lower initial energy density there is less time until freeze-out to produce radial flow, and the resulting spectrum is steeper, as desired; unfortunately, its normalization dN/dy decreases, too. Combining the insights from these two trial runs, we see that making the TTHM spectrum sufficiently steep requires still much smaller initial energy densities  $\varepsilon_0$ , combined with much larger starting times  $\tau_0$  for the TTHM transverse dynamical evolution, in order to increase dN/dy. This implies a long transverse thermalization time, which poses intrinsic consistency problems to be addressed later.

Following this route, we find that the HDM gluon spectrum can be roughly reproduced with either one of the following three sets of parameters:  $\tau_0 = 6 \text{ fm/}c$  with  $\varepsilon_0 = e_0/15$ ,  $\tau_0 = 7 \text{ fm/}c$  and  $\varepsilon_0 = e_0/18$ , or  $\tau_0 = 7.8 \text{ fm/}c$  and  $\varepsilon_0 = e_0/20$ . The corresponding spectra are shown in Fig. 4. These parameter sets are not unique and the agreement with the HDM spectra is not perfect, but what is clear is that a



FIG. 6.  $v_2$  as a function of impact parameter from TTHM (solid) and HDM (dashed) simulations tuned to produce identical angleaveraged spectra  $dN/dyk_{\perp}dk_{\perp}$ . The parameters and notations used here are the same as those in Fig. 4.

large reduction of  $\varepsilon_0$  by at least 1/15 is essential in TTHM to obtain the much deeper HDM slope.

#### D. Elliptic flow from TTHM with retuned initial conditions

Figure 6 again shows the impact-parameter dependence of  $v_2$ , but now with retuned TTHM initial conditions as described in the preceding section. With the much lower  $\varepsilon_0$  forced upon us by the given slope of the final  $k_{\perp}$  spectrum, there is much less time to generate transverse flow. Figure 6 shows that this leads not only to reduced radial flow, as reflected in the steeper single-particle spectrum for central collisions (see Fig. 4), but also cuts down on the elliptic flow  $v_2$  which now remains significantly below the HDM level. As the latter is representative of the data [3,11], we conclude that a model with only transversally thermalized momenta in the early collision stages cannot produce as much elliptic flow as required by experiment.

In order to further strengthen this argument let us look at the  $k_{\perp}$  dependence of  $v_2$  and study (see Fig. 7)

$$v_{2}(k_{\perp};b) = \frac{\int d\phi \cos(2\phi) \frac{dN(b)}{dyk_{\perp}dk_{\perp}d\phi}}{\int d\phi \frac{dN(b)}{dyk_{\perp}dk_{\perp}d\phi}}$$
(5.6)

as well as its impact-parameter-averaged (minimum bias [1]) value (see Fig. 8)



FIG. 7.  $v_2$  as a function of  $k_{\perp}$  for two impact parameters (as indicated) from TTHM (solid) and HDM (dashed) simulations giving identical angle-averaged spectra  $dN/dyk_{\perp}dk_{\perp}$ .



FIG. 8. Impact-parameter-averaged  $v_2$  as a function of  $k_{\perp}$  from the two models. The dashed line is from HDM, the solid line from TTHM with initial conditions tuned to produce the same angleaveraged transverse momentum spectrum at the same value of  $\varepsilon_f$ .

$$v_{2}(k_{\perp}) = \frac{\int bdb \int d\phi \cos(2\phi) \frac{dN(b)}{dyk_{\perp}dk_{\perp}d\phi}}{\int bdb \int d\phi \frac{dN(b)}{dyk_{\perp}dk_{\perp}d\phi}}.$$
 (5.7)

We here select for our comparison with HDM one of the above three sets of initial parameters for TTHM, namely,  $\tau_0 = 7 \text{ fm/}c$  and  $\varepsilon_0 = e_0/18$ , corresponding to the dot-dashed line in Fig. 4.

Figures 7 and 8 show that the  $k_{\perp}$  slope of  $v_2$  is much lower than in the HDM. This is true not only for the impactparameter-averaged elliptic flow shown in Fig. 8, but holds universally also for all impact parameters; two examples are plotted in Fig. 7. For  $v_2(k_{\perp})$  from minimum bias events accurate data were published in Refs. [1–3], which follow essentially the dashed HDM line in Fig. 8 and definitely exclude our TTHM.

In Fig. 9 we show the time evolution of the momentum anisotropy [5]

$$\varepsilon_p = \frac{\langle T^{xx} - T^{yy} \rangle}{\langle T^{xx} + T^{yy} \rangle},\tag{5.8}$$

where the angular brackets indicate an average over the transverse plane. At freeze-out, this momentum anisotropy is translated into elliptic flow  $v_2$ , where the coefficient between the two variables depends on the particle rest mass [5]. Fig-



FIG. 9. Asymmetry measure  $\varepsilon_p$  as a function of the duration of the time evolution at b=7 fm with similar final particle spectra from both models. Again the solid line is from TTHM and the dashed line from HDM.



FIG. 10.  $v_2$  as a function of  $k_{\perp}$  for massless and massive bosons from the two models. The solid lines are for massless bosons while the dashed and dot-dashed lines are for bosons with m = 100 MeV and m = 1 GeV, respectively.

ure 9 shows that with the hard equation of state for noninteracting massless gluons and RHIC-type initial conditions freeze-out actually happens before the momentum anisotropy saturates [17] (and before the initial spatial anisotropy has fully disappeared). This is even more true for the TTHM (with its even stiffer EOS of  $P_{\perp} = \frac{1}{2}\varepsilon$ ) than for the HDM (where  $P = \frac{1}{3}\varepsilon$ ): the TTHM freezes out significantly earlier, even though about the same amount of transverse flow is generated.

### E. The low- $k_{\perp}$ limit of $v_2(k_{\perp})$ for massless particles

A careful inspection of Figs. 7 and 8 shows that at small transverse momenta the shape of the  $k_{\perp}$ -differential elliptic flow  $v_2(k_{\perp})$  is different for TTHM and HDM. Figure 10 further shows that the small- $k_{\perp}$  behavior of the elliptic flow coefficient differs for massive and massless bosons. In Appendix B we present a detailed analytical treatment of the limiting behavior of  $v_2(k_{\perp})$  for small transverse momenta. For massive particles, the elliptic flow vanishes quadratically at  $k_{\perp} \rightarrow 0$ ,

$$v_2(k_\perp) = O(k_\perp^2)$$
 (massive particles), (5.9)

going over to an almost linear rise at higher  $k_{\perp}$  (which eventually turns over to saturate at  $v_2=1$  as  $k_{\perp} \rightarrow \infty$  [7]). The transition from the quadratic rise at low  $k_{\perp}$  to the quasilinear behavior at intermediate  $k_{\perp}$  occurs around  $k_{\perp} \sim m$ . In detail the transition between these regimes is complex and depends on the ratio m/T of the rest mass to the freeze-out temperature (see Fig. 10): for particles with m/T < 1 the elliptic approaches the curve for massless particles while the elliptic flow of heavier particles with  $m/T \ge 1$  remains always significantly below that of massless particles.

For massless bosons, the singularity of the Bose distribution at zero momentum leads to a qualitative change in the low-momentum limit of the elliptic flow. For the TTHM, which lacks a thermal spread of the longitudinal momenta, the elliptic flow of massless bosons (gluons or photons) approaches a positive constant with finite positive slope,

$$v_2(k_\perp) = a + bk_\perp + O(k_\perp^2)$$
 (TTHM, massless bosons).  
(5.10)

In the HDM, the longitudinal thermal momentum smearing weakens the Bose singularity and causes  $v_2$  to still vanish at zero transverse momentum, albeit just barely so: as  $k_{\perp} \rightarrow 0$ ,  $v_2(k_{\perp})$  vanishes with infinite slope,

$$v_2(k_\perp) = \frac{\text{const}}{\ln \frac{T}{k_\perp}}$$
 (HDM, massless bosons). (5.11)

It would be interesting to verify this prediction of the hydrodynamic model for photons that are directly emitted from the expanding fireball.

#### VI. CONCLUSIONS

We have tried to construct a macroscopic dynamical description of the fireball expansion in relativistic heavy-ion collisions, which relaxes the usual assumption of complete local thermal equilibrium that underlies the popular and successful hydrodynamical approach. This addresses, on a macroscopic level, the observation that existing parton-based microscopic kinetic models [20–22] seem unable to generate sufficiently fast thermalization to reach the hydrodynamic limit. We have tested a toy model where transverse momenta are thermalized very quickly while the system is streaming freely in the longitudinal direction (i.e., without longitudinal momentum transfer). The initial longitudinal motion was assumed to be boost invariant, and the model preserves this property dynamically at later times.

Starting from the kinetic transport equation, we derived the corresponding macroscopic equations of motion (TTHM) and solved them in parallel with the usual hydrodynamic (HDM) equations. Since our TTHM model is parton based and lacks a description of hadronization, we decided to compare the two approaches just before the onset of hadronization. To make phenomenologically meaningful comparisons we used a HDM reference that had previously been shown to yield a good description of RHIC data when allowed to evolve until hadron freeze-out.

The TTHM turns out to have an extremely stiff equation of state,  $P_{\perp} = \frac{1}{2}\varepsilon$ , which causes severe phenomenological problems: in order to avoid the creation of too much radial transverse flow, which would render the single-particle spectra much flatter than observed, while preserving the total multiplicity, we had to start the TTHM evolution very late, at already quite low energy densities. This implies a very long transverse thermalization time scale (and, of course, an even longer longitudinal one). While this cannot *a priori* be excluded theoretically, it causes consistency problems with our implementation which assumes that transverse collective dynamics sets in only after we start the TTHM evolution (i.e., only after thermalization). This is not realistic: even before complete thermalization of the transverse momenta the system will start to evolve in the transverse directions, since nothing confines it. This preequilibrium transverse dynamics will likely have a collective component affecting the transverse momentum spectra, necessitating an even further shift of the starting point for TTHM evolution.

We have not followed this issue any further, since there is also a severe problem when comparing the TTHM predictions with the elliptic flow data, which in our opinion eliminates it as a viable model. For the same radial flow, constrained by the slope of the single-particle spectra, TTHM provides only about half as much elliptic flow as the hydrodynamical model and thus significantly less than that required by the RHIC data [1-3,8]. This is the central result from the present study, and it strengthens the previously made argument [6,11] that the large elliptic flow measured at RHIC can only be reproduced by models that assume very rapid thermalization of the momentum spectra in all three dimensions. Partial thermalization, such as the effectively only two-dimensional thermalization in the TTHM model, is not enough to generate the observed elliptic flow. This leaves us with the continued challenge of identifying the microscopic mechanisms within QCD which can cause such rapid three-dimensional thermalization.

#### ACKNOWLEDGMENTS

S.W. thanks Peter Kolb for the FORTRAN code for calculating the nuclear overlap function that was used to initialize the TTHM time evolution. This work was supported by the U.S. Department of Energy under Contract No. DE-FG02-01ER41190.

# APPENDIX A: DECOMPOSING THE ENERGY-MOMENTUM TENSOR

To calculate the coefficients in the decomposition in Eq. (2.19) of the energy-momentum tensor we find it easiest to use the form of Eq. (2.12) for the distribution function in Eq. (2.18). Assuming massless particles (gluons), we write the four-vector  $k^{\mu}$  as

$$k^{\mu} = k_{\perp}(\cosh y, \cos \phi, \sin \phi, \sinh y). \tag{A1}$$

We integrate over  $d^3k/k_0 = dyk_{\perp}dk_{\perp}d\phi$  in Eq. (2.18) by first doing the trivial  $\eta$  integration and then integrating over  $k_{\perp}$ . After integration over  $\eta$  the exponent of the Bose factor in Eq. (2.12) reduces to

$$\frac{k \cdot u}{T} = \frac{\gamma_{\perp} k_{\perp}}{T} (1 - \mathbf{\hat{k}}_{\perp} \cdot \mathbf{v}_{\perp}) = \frac{\gamma_{\perp} k_{\perp}}{T} [1 - v_{\perp} \cos(\phi - \phi_s)],$$
(A2)

where  $\phi - \phi_s$  is the angle between  $\mathbf{k}_{\perp}$  (azimuthal angle  $\phi$ ) and  $\mathbf{v}_{\perp}$  (azimuthal angle  $\phi_s$ ). The  $k_{\perp}$  integration is then easily done using

$$\int_{0}^{\infty} \frac{k_{\perp}^{n} dk_{\perp}}{e^{\alpha k_{\perp}} - 1} = \frac{n! \zeta(n+1)}{\alpha^{n+1}}.$$
 (A3)

In our case  $\alpha = \gamma_{\perp} [1 - v_{\perp} \cos(\phi - \phi_s)]/T$  and n = 3, and the relevant zeta function is  $\zeta(4) = \pi^4/90$ . In this way we arrive at

PHYSICAL REVIEW C 66, 014907 (2002)

$$T^{\mu\nu} = \frac{\tau_0}{\tau} \frac{\nu}{\gamma_{\perp}^5} \frac{\pi^2 T^4}{60} \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \frac{M^{\mu\nu}}{\left[1 - v_{\perp}\cos(\phi - \phi_s)\right]^4}$$
(A4)

with

1

$$M^{\mu\nu} = \begin{pmatrix} \cosh^2 \eta & \cos\phi \cosh\eta & \sin\phi \cosh\eta & \cosh\eta & \sinh\eta \\ \cos\phi \cosh\eta & \cos^2\phi & \cos\phi \sin\phi & \cos\phi \sinh\eta \\ \sin\phi \cosh\eta & \cos\phi \sin\phi & \sin^2\phi & \sin\phi \sinh\eta \\ \cosh\eta \sinh\eta & \cos\phi \sinh\eta & \sin\phi \sinh\eta & \sinh^2\eta \end{pmatrix}.$$
 (A5)

The remaining azimuthal integration is done by shifting the integration variable to  $\theta = \phi - \phi_s$ , using

$$\cos \phi = \cos \theta \cos \phi_s - \sin \theta \sin \phi_s,$$
  
$$\sin \phi = \sin \theta \cos \phi_s + \cos \theta \sin \phi_s \qquad (A6)$$

in the numerator  $M^{\mu\nu}$ , and exploiting the formula

$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{1}{(1 - v_{\perp} \cos \theta)^n} = \frac{1}{(1 - v_{\perp}^2)^{n/2}} P_{n-1} \left(\frac{1}{\sqrt{1 - v_{\perp}^2}}\right)$$
$$= \gamma_{\perp}^n P_{n-1}(\gamma_{\perp}), \qquad (A7)$$

where the  $P_n$  are Legendre polynomials. We need

$$\gamma_{\perp}^{2} P_{1}(\gamma_{\perp}) = \gamma_{\perp}^{3}, \quad \gamma_{\perp}^{3} P_{2}(\gamma_{\perp}) = \gamma_{\perp}^{5} (1 + \frac{1}{2} v_{\perp}^{2}),$$
$$\gamma_{\perp}^{4} P_{3}(\gamma_{\perp}) = \gamma_{\perp}^{7} (1 + \frac{3}{2} v_{\perp}^{2}).$$
(A8)

 $M^{00}$ ,  $M^{03} = M^{30}$ , and  $M^{33}$  have no  $\phi$  dependence and the corresponding angular integrals thus give

$$\int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \frac{1}{\left[1 - v_{\perp} \cos(\phi - \phi_s)\right]^4} = \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{1}{\left[1 - v_{\perp} \cos\theta\right]^4} = \gamma_{\perp}^4 P_3(\gamma_{\perp}) = \gamma_{\perp}^7 \left(1 + \frac{3}{2} v_{\perp}^2\right).$$
(A9)

For  $M^{01} = M^{10}$  and  $M^{13} = M^{31}$  we need

$$\int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \frac{\cos\phi}{\left[1 - v_{\perp}\cos(\phi - \phi_s)\right]^4}$$
$$= \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\cos\theta\cos\phi_s - \sin\theta\sin\phi_s}{(1 - v_{\perp}\cos\theta)^4}$$
$$= \frac{\cos\phi_s}{v_{\perp}} \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{1 - (1 - v_{\perp}\cos\theta)}{(1 - v_{\perp}\cos\theta)^4}$$
$$= \frac{v_x}{v_{\perp}^2} [\gamma_{\perp}^4 P_3(\gamma_{\perp}) - \gamma_{\perp}^3 P_2(\gamma_{\perp})] = \frac{1}{2} \gamma_{\perp}^7 v_x (4 + v_{\perp}^2).$$
(A10)

In the last step on the first line we dropped the vanishing term  $\sim \sin \theta$  and also added and subtracted a term  $1/v_{\perp}$  in the numerator. In the second line we used  $\cos \phi_s = v_x/v_{\perp}$ . For the remaining components we proceed similarly. The energy-momentum tensor thus takes the form

$$T^{\mu\nu} = \frac{\tau_0}{\tau} \nu_g \frac{\pi^2 T^4}{60} \gamma_{\perp}^2 \begin{pmatrix} (1 + \frac{3}{2} v_{\perp}^2) \cosh^2 \eta & \frac{1}{2} v_x (4 + v_{\perp}^2) \cosh \eta & \frac{1}{2} v_y (4 + v_{\perp}^2) \cosh \eta & (1 + \frac{3}{2} v_{\perp}^2) \cosh \eta \sin \eta \\ \frac{1}{2} v_x (4 + v_{\perp}^2) \cosh \eta & \frac{1}{2} (1 + 4 v_x^2 - v_y^2) & \frac{5}{2} v_x v_y & \frac{1}{2} v_x (4 + v_{\perp}^2) \sinh \eta \\ \frac{1}{2} v_y (4 + v_{\perp}^2) \cosh \eta & \frac{5}{2} v_x v_y & \frac{1}{2} (1 - v_x^2 + 4 v_y^2) & \frac{1}{2} v_y (4 + v_{\perp}^2) \sinh \eta \\ (1 + \frac{3}{2} v_{\perp}^2) \cosh \eta \sinh \eta & \frac{1}{2} v_x (4 + v_{\perp}^2) \cosh \eta & \frac{1}{2} v_y (4 + v_{\perp}^2) \sinh \eta \end{pmatrix}.$$

Contracting this result once with each of the vectors  $u^{\mu}, n^{\mu}$ , and  $m^{\mu}$  we obtain

$$T^{\mu\nu}u_{\nu} = \frac{\tau_{0}}{\tau} \nu_{g} \frac{\pi^{2}T^{4}}{60} \gamma_{\perp}^{3} \begin{pmatrix} (1 - \frac{1}{2}v_{\perp}^{2} - \frac{1}{2}v_{\perp}^{4})\cosh\eta \\ \frac{3}{2}v_{x}(1 - v_{\perp}^{2}) \\ \frac{3}{2}v_{y}(1 - v_{\perp}^{2}) \\ (1 - \frac{1}{2}v_{\perp}^{2} - \frac{1}{2}v_{\perp}^{4})\sinh\eta \end{pmatrix},$$

$$T^{\mu\nu}m_{\nu} = -\frac{\tau_{0}}{\tau}\nu_{g}\frac{\pi^{2}T^{4}}{60}\gamma_{\perp}^{3} \begin{pmatrix} (1-v_{\perp}^{2})v_{\perp}\cosh\eta\\v_{x}(1-v_{\perp}^{4})/2v_{\perp}\\v_{y}(1-v_{\perp}^{4})/2v_{\perp}\\(1-v_{\perp}^{2})v_{\perp}\sinh\eta \end{pmatrix},$$

$$T^{\mu\nu}n_{\nu} = 0.$$
 (A12)

A second contraction then yields the coefficients A through H in Eq. (2.19) as given in Eqs. (2.22)-(2.27).

## APPENDIX B: THE SMALL TRANSVERSE MOMENTUM LIMIT OF ELLIPTIC FLOW

In this appendix we derive the small- $k_{\perp}$  limit of the elliptic flow coefficient  $v_2$  for TTHM and HDM, for massive and massless particles. Suppressing the impact-parameter dependence which is irrelevant here, we write Eq. (5.6) in the form

$$v_2(k_\perp) = \frac{\mathcal{N}}{\mathcal{D}},\tag{B1}$$

where the numerator  $\mathcal{N}$  and denominator  $\mathcal{D}$  are obtained from the Cooper-Frye formula Eq. (5.1). For massive particles we must generalize Eq. (5.3) by writing the first term as  $m_{\perp} \cosh(y - \eta)$  and use

$$k \cdot u = \gamma_{\perp} (m_{\perp} \cosh(y - \eta) - \mathbf{k}_{\perp} \cdot \mathbf{v}_{\perp}). \tag{B2}$$

For the TTHM the distribution function is given by Eq. (2.12), and we obtain for particles at midrapidity y=0

$$\begin{cases} \mathcal{N} \\ \mathcal{D} \end{cases} = \frac{\nu_g \tau_0}{(2\pi)^2} \int \frac{d^2 x_\perp}{\gamma_\perp} \int_{-\infty}^{\infty} d\eta \,\delta(\eta) \begin{cases} \mathcal{N}_{\phi} \\ \mathcal{D}_{\phi} \end{cases}, \qquad (B3)$$

where

$$\begin{pmatrix} \mathcal{N}_{\phi} \\ \mathcal{D}_{\phi} \end{pmatrix} = \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \begin{cases} \cos(2\phi) \\ 1 \end{cases} \frac{m_{\perp} \cosh \eta - \mathbf{k}_{\perp} \cdot \nabla_{\perp} \tau_{f}}{e^{\gamma_{\perp} (m_{\perp} \cosh \eta - \mathbf{k}_{\perp} \cdot \mathbf{v}_{\perp})/T} - 1}.$$
(B4)

Here  $\gamma_{\perp}$  and  $\tau_{\rm f}$  are functions of the position in the transverse plane,  $\gamma_{\perp} = \gamma_{\perp}(\mathbf{x}_{\perp})$  and  $\tau_{\rm f} = \tau_{\rm f}(\mathbf{x}_{\perp})$ . For the HDM we instead use Eq. (3.1) and get

$$\begin{cases} \mathcal{N} \\ \mathcal{D} \end{cases} = \frac{\nu_g}{(2\pi)^2} \int d^2 x_{\perp} \tau_{\rm f} \int_{-\infty}^{\infty} d\eta \begin{cases} \mathcal{N}_{\phi} \\ \mathcal{D}_{\phi} \end{cases}.$$
 (B5)

The main difference (except for the different weighting of the integration over the transverse plane  $d^2x_{\perp} = rdrd\phi_s$ ) is the nontrivial  $\eta$  integration in the HDM case.

#### 1. Massive particles

For nonzero rest mass,  $m \neq 0$ , we can expand in both cases the  $\phi$  integrand for small  $k_{\perp}$ . Keeping only terms up to first order in  $k_{\perp}$  we find

$$\begin{cases} \mathcal{N}_{\phi} \\ \mathcal{D}_{\phi} \end{cases} \approx \frac{m \cosh \eta}{e^{\gamma_{\perp} m \cosh \eta/T} - 1} \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \begin{cases} \cos(2\phi) \\ 1 \end{cases} \\ \times \left( 1 - \frac{\mathbf{k}_{\perp} \cdot \nabla_{\perp} \tau_{\rm f}}{m \cosh \eta} \right) \left( 1 + \frac{\mathbf{k}_{\perp} \cdot \mathbf{v}_{\perp}}{T} \frac{\gamma_{\perp}}{1 - e^{-\gamma_{\perp} m \cosh \eta/T}} \right).$$

$$(B6)$$

Higher orders of  $k_{\perp}$  come with higher inverse powers of *T* or *m* cosh  $\eta$ ; the corresponding  $\eta$  integrals thus are all finite (trivially so in the TTHM), and the  $k_{\perp}$  expansion is well behaved.

As in Appendix A we write  $\mathbf{k}_{\perp} \cdot \mathbf{v}_{\perp} = k_{\perp} v_{\perp} \cos(\phi - \phi_s)$ =  $k_{\perp} v_{\perp} \cos \theta$  and use  $-\pi \le \theta \le \pi$  as integration variable. Then, using Eq. (A6),

$$\hat{\mathbf{k}}_{\perp} \cdot \boldsymbol{\nabla}_{\perp} \tau_{\mathrm{f}} = (\partial_r \tau_{\mathrm{f}}) \cos \theta + (\partial_{\phi} \tau_{\mathrm{f}}) \sin \theta, \qquad (B7)$$

where  $\partial_r \tau_f = \hat{\mathbf{v}}_{\perp} \cdot \nabla_{\perp} \tau_f$  and  $\partial_{\phi} \tau_f = \hat{\mathbf{z}} \cdot (\hat{\mathbf{v}}_{\perp} \times \nabla_{\perp} \tau_f)$ . Here  $\hat{\mathbf{z}}$  is the unit vector along the beam direction, and  $\hat{\mathbf{v}}_{\perp}$  coincides with the radial unit vector,  $\hat{\mathbf{v}}_{\perp} = (\cos \phi_s, \sin \phi_s)$ . We also need

$$\cos(2\phi) = \cos(2\phi_s)(\cos^2\theta - \sin^2\theta) - 2\sin(2\phi_s)\sin\theta\cos\theta.$$
(B8)

After these manipulations the angular integrations in Eq. (B6) are easily performed. Due to symmetric integration over the full circle, only terms containing even powers of  $\cos \theta$  and/or  $\sin \theta$  survive. We find

$$\mathcal{N}_{\phi} = O(k_{\perp}^2), \quad \mathcal{D}_{\phi} = a + bk_{\perp} + O(k_{\perp}^2), \tag{B9}$$

where *a* and *b* are integrable functions of  $\mathbf{x}_{\perp}$  and  $\eta$ . As a result, for massive particles  $v_2(k_{\perp}) = O(k_{\perp}^2)$ , i.e., the elliptic flow coefficient vanishes with zero slope when  $k_{\perp} \rightarrow 0$ , as first observed by Danielewicz [19].

#### 2. Massless bosons in the TTHM

For massless bosons at y=0 the angular integrals  $\mathcal{N}_{\phi}$ ,  $\mathcal{D}_{\phi}$  in Eq. (B4) reduce to

$$\begin{cases} \mathcal{N}_{\phi} \\ \mathcal{D}_{\phi} \end{cases} = \frac{T}{\gamma_{\perp}} \int_{-\pi}^{\pi} \frac{d\phi}{2\pi} \begin{cases} \cos(2\phi) \\ 1 \end{cases} \frac{1 - \frac{\hat{\mathbf{k}}_{\perp} \cdot \mathbf{V}_{\perp} \tau_{\mathrm{f}}}{\cosh \eta}}{1 - \frac{\hat{\mathbf{k}}_{\perp} \cdot \mathbf{v}_{\perp}}{\cosh \eta}} \\ \times \frac{\frac{\gamma_{\perp} k_{\perp}}{T} (\cosh \eta - \hat{\mathbf{k}}_{\perp} \cdot \mathbf{v}_{\perp})}{e^{\gamma_{\perp} k_{\perp} / T (\cosh \eta - \hat{\mathbf{k}}_{\perp} \cdot \mathbf{v}_{\perp}) - 1}}.$$
(B10)

We can try to expand the last factor for small  $k_{\perp}$  by using

$$\frac{x}{e^{x}-1} = 1 - \frac{x}{2} + O(x^{2}).$$
(B11)

However, since in this expression the expansion parameter is  $x = \gamma_{\perp} k_{\perp} / T(\cosh \eta - \hat{\mathbf{k}}_{\perp} \cdot \mathbf{v}_{\perp})$ , each additional power of  $k_{\perp}$  brings in another factor  $\cosh \eta$ . This is no problem for the TTHM with its factor  $\delta(\eta)$  under the  $\eta$  integral, but for the HDM the  $\eta$  integrals of the expansion coefficients diverge, rendering the expansion meaningless. For the HDM thus the  $\eta$  integration must be performed *before* we expand for small  $k_{\perp}$  (see the following section).

For the TTHM, on the other hand, we can continue along this direction. Using the same manipulations as in the preceding section, we find up to linear order in  $k_{\perp}$ 

$$\begin{cases} \mathcal{N}_{\phi} \\ \mathcal{D}_{\phi} \end{cases} \approx \frac{T}{\gamma_{\perp}} \int_{-\pi}^{\pi} \frac{d\,\theta}{2\,\pi} \begin{cases} \cos(2\,\phi_s)\cos(2\,\theta) - \sin(2\,\phi_s)\sin(2\,\theta) \\ 1 \end{cases} \\ \times \frac{1 - (\partial_r \tau_{\rm f})\cos\,\theta - (\partial_\phi \tau_{\rm f})\sin\,\theta}{1 - v_{\perp}\cos\,\theta} \\ \times \left(1 - \frac{\gamma_{\perp}k_{\perp}}{2T}(1 - v_{\perp}\cos\,\theta)\right). \end{cases}$$
(B12)

We use Eq. (A7) to derive the following table of integrals:

$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{1}{1 - v_{\perp} \cos \theta} = \gamma_{\perp} ,$$
$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\cos \theta}{1 - v_{\perp} \cos \theta} = \frac{\gamma_{\perp} - 1}{v_{\perp}} ,$$
$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\cos^2 \theta}{1 - v_{\perp} \cos \theta} = \frac{\gamma_{\perp} - 1}{v_{\perp}^2} ,$$

$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\sin^2\theta}{1 - v_{\perp}\cos\theta} = \frac{\gamma_{\perp} - 1}{\gamma_{\perp}v_{\perp}^2},$$
$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\cos(2\theta)}{1 - v_{\perp}\cos\theta} = \frac{1}{\gamma_{\perp}} \left(\frac{\gamma_{\perp} - 1}{v_{\perp}}\right)^2,$$
$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\sin^2\theta\cos\theta}{1 - v_{\perp}\cos\theta} = \frac{1}{2v_{\perp}} \left(\frac{\gamma_{\perp} - 1}{\gamma_{\perp}v_{\perp}}\right)^2,$$
$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \frac{\cos(2\theta)\cos\theta}{1 - v_{\perp}\cos\theta} = \frac{1}{\gamma_{\perp}v_{\perp}} \left(\frac{\gamma_{\perp} - 1}{v_{\perp}}\right)^2, \quad (B13)$$

and find

$$\mathcal{N}_{\phi} = T \left( \frac{\gamma_{\perp} - 1}{\gamma_{\perp} \upsilon_{\perp}} \right)^2 \left[ \cos(2\phi_s) \left( 1 - \frac{\partial_r \tau_f}{\upsilon_{\perp}} \right) + \sin(2\phi_s) \frac{\partial_{\phi} \tau_f}{\gamma_{\perp} \upsilon_{\perp}} \right] + O(k_{\perp}^2),$$

$$\mathcal{D}_{\phi} = T \left( 1 - \frac{\gamma_{\perp} - 1}{\gamma_{\perp} v_{\perp}} (\partial_r \tau_{\mathbf{f}}) \right) - \frac{k_{\perp}}{2} + O(k_{\perp}^2). \quad (B14)$$

Now both the numerator and denominator approach nonzero constants as  $k_{\perp} \rightarrow 0$ . Since the fireball center freezes out later than its edge, the term  $\sim (\partial_r \tau_f)$ , after integration over the transverse plane, contributes with a negative sign, and the  $k_{\perp}$ -independent first terms in  $\mathcal{N}_{\phi}$  and  $\mathcal{D}_{\phi}$  are correspondingly positive. As a result, the small- $k_{\perp}$  expansion of the elliptic flow coefficient takes the form

$$v_{2}(k_{\perp}) = \frac{\mathcal{N}}{\mathcal{D}} = \frac{a}{1 - bk_{\perp}} + O(k_{\perp}^{2}) = a + abk_{\perp} + O(k_{\perp}^{2}),$$
(B15)

where *a* and *b* are positive constants arising from the integration over the transverse profiles of the freeze-out time and flow velocity. Thus, as  $k_{\perp} \rightarrow 0$ ,  $v_2$  approaches a positive constant value with nonzero positive slope, as confirmed by Figs. 7–9.

#### 3. Massless bosons from the HDM

In the last step we discuss the corresponding limit for massless bosons from the HDM, in order to understand the surprising behavior shown in the upper part of Fig. 9. As already mentioned, in this case the  $\eta$  integration must be performed *before* expanding in  $k_{\perp}$ . Doing this integration first, we rewrite Eq. (B5) as

$$\begin{cases} \mathcal{N} \\ \mathcal{D} \end{cases} = \frac{\nu_g}{(2\pi)^2} \int r dr d\phi_s \frac{\tau_{\rm f}(r,\phi_s)}{\gamma_{\perp}(r,\phi_s)} \\ \times \int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \begin{cases} \cos(2\phi_s)\cos(2\theta) - \sin(2\phi_s)\sin(2\theta) \\ 1 \end{cases} \\ \times \int_{-\infty}^{\infty} d\eta \frac{\xi(\cosh\eta - \hat{\mathbf{k}}_{\perp} \cdot \nabla_{\perp}\tau_{\rm f})}{e^{\xi(\cosh\eta - \hat{\mathbf{k}}_{\perp} \cdot \nu_{\perp}) - 1}}, \qquad (B16) \end{cases}$$

where all the dependence on the magnitude of  $k_{\perp}$  is hidden in the variable

$$\xi = \frac{\gamma_{\perp} k_{\perp}}{T}.$$
 (B17)

Let us denote the result of the  $\eta$  integral by  $I_{\eta}$ . Expanding the Bose-Einstein distribution in a power series and performing the  $\eta$  integration term by term we find

$$I_{\eta} = \xi \sum_{n=1}^{\infty} e^{nv_{\perp}\xi\cos\theta} \int_{-\infty}^{\infty} d\eta e^{-n\xi\cosh\eta}$$
$$\times [\cosh\eta - (\partial_{r}\tau_{\rm f})\cos\theta - (\partial_{\phi}\tau_{\rm f})\sin\theta]$$
$$= 2\xi \sum_{n=1}^{\infty} e^{nv_{\perp}\xi\cos\theta} \{K_{1}(n\xi) - [\cos\theta(\partial_{r}\tau_{\rm f})$$
$$+ \sin\theta(\partial_{\phi}\tau_{\rm f})]K_{0}(n\xi)\}, \qquad (B18)$$

where the  $K_{\nu}$  are modified Bessel functions of the second kind. The integration over the momentum-space angle  $\theta$  can now be performed, too, yielding modified Bessel functions of the first kind  $I_{\nu}$ ,

$$\int_{-\pi}^{\pi} \frac{d\theta}{2\pi} \cos(\nu\theta) e^{z\cos\theta} = I_{\nu}(z), \qquad (B19)$$

where  $z = nv_{\perp}\xi$ . After some simple manipulations using trigonometric identities we find for the elliptic flow coefficient

$$v_{2}(k_{\perp}) = \frac{\sum_{n=1}^{\infty} \int r dr d\phi_{s} \frac{\tau_{f}(r,\phi_{s})}{\gamma_{\perp}(r,\phi_{s})} \mathcal{N}_{n}(r,\phi_{s};k_{\perp})}{\sum_{n=1}^{\infty} \int r dr d\phi_{s} \frac{\tau_{f}(r,\phi_{s})}{\gamma_{\perp}(r,\phi_{s})} \mathcal{D}_{n}(r,\phi_{s};k_{\perp})},$$
(B20)

where

$$\mathcal{N}_{n} = \cos(2\phi_{s}) \left\{ \xi I_{2}(nv_{\perp}\xi) K_{1}(n\xi) - \frac{\partial_{r}\tau_{f}}{2} \xi [I_{1}(nv_{\perp}\xi) + I_{3}(nv_{\perp}\xi)] K_{0}(n\xi) \right\} + \sin(2\phi_{s}) \frac{\partial_{\phi}\tau_{f}}{2} \xi [I_{1}(nv_{\perp}\xi) - I_{3}(nv_{\perp}\xi)] K_{0}(n\xi),$$

$$\mathcal{D}_n = \xi I_0(nv_\perp \xi) K_1(n\xi) - (\partial_r \tau_f) \xi I_1(nv_\perp \xi) K_0(n\xi).$$
(B21)

These expressions generalize the result given in [7] to the general case of a freeze-out time  $\tau_f$ , which depends on the transverse position  $\mathbf{x}_{\perp}$ .

We can now study the small-momentum limit of  $v_2$  by letting  $\xi \rightarrow 0$ . Let us first consider the Boltzmann approximation that corresponds to keeping only the terms  $\mathcal{N}_1$  and  $\mathcal{D}_1$  in Eq. (B20). Using the small- $\xi$  expansions of the Bessel functions one finds that in this approximation

$$\lim_{k_{\perp}\to 0} v_2^{\text{Boltz}}(k_{\perp}) \sim \frac{k_{\perp}^2}{T^2} \ln \frac{k_{\perp}}{T}.$$
 (B22)

For massless Boltzmann particles, the elliptic flow from the HDM thus goes to zero with zero slope, but by a factor  $\ln(k_{\perp}/T)$  more slowly than for massive particles, see Sec. 1 of Appendix B.

For bosons we must sum over all *n* in Eq. (B20). For  $k_{\perp} \rightarrow 0$ , i.e.,  $\xi \rightarrow 0$ , the sum over *n* becomes the discrete representation of an integral: defining

$$f_{\nu_{\perp}}(\zeta) \equiv I_{\nu}(\nu_{\perp}\zeta) K_{\mu}(\zeta), \qquad (B23)$$

we have

$$\xi \sum_{n=1}^{\infty} I_{\nu}(nv_{\perp}\zeta) K_{\mu}(n\zeta) = \xi \sum_{n=1}^{\infty} f_{v_{\perp}}(n\zeta) \rightarrow \int_{\xi/2}^{\infty} f_{v_{\perp}}(\zeta) d\zeta.$$
(B24)

By expanding the Bessel functions for large arguments one sees that for large  $\zeta$ 

$$f_{v_{\perp}}(\zeta) \approx \frac{1}{2\sqrt{v_{\perp}}\zeta} e^{-(1-v_{\perp})\zeta}, \qquad (B25)$$

such that the integral in Eq. (B24) clearly converges at the upper end. The  $k_{\perp}$  dependence of  $v_2$  can now be extracted by studying the dependence of this integral on its lower limit  $\xi/2 = \gamma_{\perp}k_{\perp}/2T$ . To do so we employ the following trick: We split the integral at an arbitrary intermediate scale  $\lambda$ ,

$$\begin{split} \int_{\xi/2}^{\infty} f_{v_{\perp}}(\zeta) d\zeta &= \int_{\xi/2}^{\lambda} f_{v_{\perp}}(\zeta) d\zeta + \int_{\lambda}^{\infty} f_{v_{\perp}}(\zeta) d\zeta \\ &= \int_{\xi/2}^{\lambda} f_{v_{\perp}}(\zeta) d\zeta + F(\lambda; v_{\perp}), \end{split} \tag{B26}$$

where  $F(\lambda; v_{\perp})$  is a finite number whose dependence on  $\lambda$  cancels against the  $\lambda$  dependence of the first integral, irrespective of how we choose  $\lambda$ . By taking  $\xi/2 < \lambda \ll 1$  we can evaluate the first integral analytically by making use of the expansion of the Bessel functions for small arguments, keeping only the leading terms. In the limit  $\xi \rightarrow 0$  we find in this way

$$\int_{\xi/2}^{\lambda} I_2(v_{\perp}\zeta) K_1(\zeta) d\zeta \rightarrow \frac{\lambda^2 v_{\perp}^2}{16},$$

$$\int_{\xi/2}^{\lambda} I_1(v_{\perp}\zeta) K_0(\zeta) d\zeta \rightarrow \frac{v_{\perp}}{16} \xi^2 \ln(a\xi) + C(\lambda),$$

$$\int_{\xi/2}^{\lambda} I_3(v_{\perp}\zeta) K_0(\zeta) d\zeta \rightarrow 0,$$

$$\int_{\xi/2}^{\lambda} I_0(v_{\perp}\zeta) K_1(\zeta) d\zeta \rightarrow \ln\frac{2\lambda}{\xi} + C'(\lambda), \quad (B27)$$

- STAR Collaboration, K. H. Ackermann *et al.*, Phys. Rev. Lett. 86, 402 (2001).
- [2] STAR Collaboration, C. Adler *et al.*, Phys. Rev. Lett. 87, 182 301 (2001).
- [3] STAR Collaboration, R. Snellings, Nucl. Phys. A698, 193c (2002).
- [4] J. Y. Ollitrault, Phys. Rev. D 46, 229 (1992).
- [5] P. F. Kolb, J. Sollfrank, and U. Heinz, Phys. Lett. B 459, 667 (1999); Phys. Rev. C 62, 054909 (2000).
- [6] P. F. Kolb, P. Huovinen, U. Heinz, and H. Heiselberg, Phys. Lett. B 500, 232 (2001); P. F. Kolb, U. Heinz, P. Huovinen, K. J. Eskola, and K. Tuominen, Nucl. Phys. A696, 197 (2001).
- [7] P. Huovinen, P. F. Kolb, U. Heinz, P. V. Ruuskanen, and S. A. Voloshin, Phys. Lett. B 503, 58 (2001).
- [8] PHENIX Collaboration, R. A. Lacey, Nucl. Phys. A698, 559c (2002); PHENIX Collaboration, K. Adcox *et al.*, nucl-ex/0204005.
- [9] PHOBOS Collaboration, I. C. Park, Nucl. Phys. A698, 564c (2002).
- [10] S. Voloshin and Y. Zhang, Z. Phys. C 70, 665 (1996).
- [11] U. Heinz and P. Kolb, Nucl. Phys. A702, 269 (2002); hep-ph/0204061.
- [12] This assumption is not unreasonable because flow anisotropies require anisotropic pressure gradients for building up, but at the same time they reduce the latter dynamically [13]. The

where *a* is a  $\lambda$ -independent constant while *C*, *C'* depend on  $\lambda$  but combine with the corresponding constants  $F(\lambda; v_{\perp})$  to finite,  $\lambda$ -independent constants. Inserting these results into Eqs. (B20) and (B21) we finally obtain

$$\lim_{k_{\perp}/T \to 0} v_2(k_{\perp}) = \frac{\text{const}}{\ln \frac{T}{k_{\perp}}},$$
 (B28)

which approaches zero with infinite slope. This is exactly the behavior seen in the upper panel of Fig. 9 for massless bosons.

necessary spatial transverse anisotropies thus exist only during the very early collision stages, and we here assume that the time needed to eliminate them is shorter than the longitudinal thermalization time.

- [13] H. Sorge, Phys. Rev. Lett. 78, 2309 (1997); 82, 2048 (1999).
- [14] J. D. Bjorken, Phys. Rev. D 27, 140 (1983).
- [15] P. V. Ruuskanen, Acta Phys. Pol. B 18, 551 (1988).
- [16] F. Cooper and G. Frye, Phys. Rev. D 10, 186 (1974).
- [17] As shown in Fig. 1 of [18], the equation of state of an ideal massless gluon gas is so stiff that at RHIC energies the fireball freezes out dynamically (i.e., driven by strong transverse flow) before the spatial deformation of the reaction zone in noncentral collisions has fully disappeared. Therefore, with such an equation of state, elliptic flow does not fully saturate before decoupling. Nonetheless, the same figure shows [18] that more than half of the maximally possible elliptic flow develops before the decoupling energy density is reached.
- [18] P. F. Kolb, J. Sollfrank, P. V. Ruuskanen, and U. Heinz, Nucl. Phys. A661, 349 (1999).
- [19] P. Danielewicz, Phys. Rev. C 51, 716 (1995).
- [20] K. Geiger, Phys. Rep. 258, 237 (1995).
- [21] S. M. H. Wong, Nucl. Phys. A607, 442 (1996); Phys. Rev. C 54, 2588 (1996); 56, 1075 (1997); Nucl. Phys. A638, 527c (1998); Phys. Rev. C 61, 064903 (2000).
- [22] D. Molnar and M. Gyulassy, Nucl. Phys. A697, 495 (2002).