

Centrality dependence of direct photons in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV

Rainer J. Fries,¹ Berndt Müller,² and Dinesh K. Srivastava³

¹*School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455, USA*

²*Department of Physics, Duke University, Durham, North Carolina 27708, USA*

³*Variable Energy Cyclotron Center, 1/AF Bidhan Nagar, Kolkata 700 064, India*

(Received 26 July 2005; published 24 October 2005)

We calculate the spectra of high-energy photons emitted in relativistic Au + Au collisions for various centralities and compare to data recently collected at the Relativistic Heavy Ion Collider by the PHENIX Collaboration. Our results for photons from primary hard scatterings and photons from interactions of jets with the medium are consistent with the measurements of neutral pion and direct photon production in $p + p$ collisions and give a good description of direct photon spectra measured in Au + Au collisions. The contribution of photons from jet-to-photon conversion in the medium can be as large as the photon yield from hard scatterings in the momentum range $p_T \approx 2-6$ GeV/c. We show that this novel mechanism is not ruled out by any existing data.

DOI: [10.1103/PhysRevC.72.041902](https://doi.org/10.1103/PhysRevC.72.041902)

PACS number(s): 25.75.-q, 12.38.Mh

High-energy nuclear collisions are studied to understand the properties of strongly interacting matter at the highest energy densities and to search for the transition of hadronic matter to a quark-gluon plasma (QGP) [1]. Electromagnetic probes are excellent tools to gather information about the dense and hot medium created in such collisions. The mean free path of photons is much larger than the typical size of the emerging fireball, allowing them to escape nearly unperturbed once created in the system [2]. Since photons are emitted throughout the history of the fireball, one expects to obtain information that is complementary to that gained from the measurement of hadrons, which are subject to strong final-state interactions and reflect the latest stage of the collision.

Thermal radiation of photons can occur both from a quark-gluon plasma [3,4] and from a hot hadronic gas [5,6]. Because the photon spectrum reflects the thermal distribution of the emitting matter, it can serve as a thermometer. However, high-energy photons are most likely to be created either in hard processes between partons in the colliding nuclei [7] or in final-state interactions of high-energy jets [8]. Photons from jets naturally carry information about the medium they traverse and hence can serve as a medium probe.

The latter process was introduced in our earlier work [8] where we predicted the existence of a medium-induced component in the photon spectrum in the intermediate p_T range. In this novel mechanism, a hard quark passing through the medium converts into a photon carrying most of its transverse momentum. Recently Zakharov has developed the related concept of modified photon bremsstrahlung of jets in the medium [9]. An up-to-date calculation of photon emission including such jet-medium interactions can be found in the work of Turbide *et al.* [10]. The general arguments given for photons in this work also apply to the emission of lepton pairs initiated by quark jets in the plasma [11].

Relativistic nuclear collisions at different centralities result in the formation of systems of different sizes, initial temperatures, and lifetimes. Assuming that the yield of processes involving large momentum transfer (hard processes) scale as the number of collisions, N_{coll} , while those with small

momentum transfer (soft processes) scale as the number of participants, N_{part} , one can predict the relative importance of the two contributions as a function of centrality (see, e.g. [12]). The recently published data on single photon production in $p + p$ collisions [13] at $\sqrt{s_{NN}} = 200$ GeV [13] and the centrality dependence of the photon yield in Au + Au collisions measured by PHENIX at the Relativistic Heavy Ion Collider (RHIC) [14] provide our first chance to obtain an understanding of the different photon sources.

Here, we shall address the question of whether the data allow us to draw conclusions about the relative contributions from different photon production mechanisms to the direct photon spectrum in Au + Au collisions. The data are still limited by (statistical and systematic) uncertainties, but the results represent a remarkable achievement because of the experimental challenge to subtract the background of photons from secondary hadronic decays, especially neutral pions. Fortunately, in Au + Au collisions at RHIC this background is reduced at intermediate and high transverse momenta $p_T > 2$ GeV/c by the large final-state suppression (commonly called “jet quenching”) for pions and other hadrons [15,16]. Thus, although the low- p_T part of the photon spectrum remains clouded by the large hadronic background, this is an invitation to look for electromagnetic signals of a QGP at intermediate and high p_T [8,17].

Let us briefly recall the photon sources that are important for $p_T > 2$ GeV/c. We define prompt photons as those coming from hard Compton and annihilation processes (e.g., $q + g \rightarrow q + \gamma$ and $q + \bar{q} \rightarrow g + \gamma$) and those emitted from jets after the hard interaction as bremsstrahlung (e.g., $q \rightarrow q + \gamma$). A baseline for these processes can be obtained by studying $p + p$ and $p + \bar{p}$ interactions. Besides an obvious scaling from $p + p$ collisions to Au + Au collisions with the number N_{coll} , and an isospin correction resulting from the presence of $p + n$ and $n + n$ collisions, Compton and annihilation spectra are expected to be altered only marginally by initial-state nuclear effects. Neutral pion spectra measured in $d + Au$ collisions indicate that such effects from shadowing and k_T broadening are small [18]. The hard process itself is pointlike on typical

scales of the medium and therefore remains unaffected. Direct photon spectra measured in $d + \text{Au}$ could help to quantify the role of initial-state effects.

Bremsstrahlung emission from jets, in contrast, takes place on more extended distances and thus can be affected by final-state interactions. Energy-loss effects would then be noticeable as a suppression of bremsstrahlung [19], while secondary scattering in the medium could initiate additional bremsstrahlung or modify bremsstrahlung by coherence effects [9]. In our work we assume that the total photon yield from jets can be described by the superposition of the usual vacuum bremsstrahlung with an additional contribution from jets annihilating and Compton scattering in the medium as described in Ref. [8].

Photons from secondary hard scatterings in the pre-equilibrium phase have been studied elsewhere and could contribute to photons at intermediate and high p_T [20]. To minimize possible effects from neglecting pre-equilibrium emission, we assume a very early thermalization time, $\tau_0 \approx 0.15 \text{ fm}/c$, in central collisions, not much larger than the formation time of minijets. An early thermalization of the matter created in $\text{Au} + \text{Au}$ collisions at RHIC is supported by several observations, especially the large elliptic flow. We also assume that the matter is chemically equilibrated at the initial time. Secondary collisions are then either between thermalized partons or between hard scattered partons and the medium, which are both taken into account here. Similar strategies for dealing with the uncertainties of pre-equilibrium emission have been invoked elsewhere [21].

We now recall the specifics of the jet-medium interaction process. We denote the momenta of the jet, the thermal parton, and the photon by \mathbf{p}_{jet} , \mathbf{p}_{th} , and \mathbf{p}_γ , respectively. The leading order QCD Compton and annihilation cross sections are peaked in the forward and backward directions. In the laboratory frame we have $|\mathbf{p}_{\text{jet}}| \gg |\mathbf{p}_{\text{th}}| \sim T$, where T is the temperature of the plasma. For high-energy photons (i.e. $|\mathbf{p}_\gamma| \gg T$), this implies that $|\mathbf{p}_\gamma| \approx |\mathbf{p}_{\text{jet}}|$. This justifies calling the process a conversion of a jet (or, more precisely, a quark initiating a jet) into a photon with similar momentum [8]. Only quark jets contribute to this process, because the gluon-photon scattering cross section is not peaked at forward angles.

The rate of photon production by annihilation and Compton scattering of jets in the medium can be approximated as [8]

$$\frac{E_\gamma dN}{d^3\mathbf{p}_\gamma d^4x} = \frac{\alpha\alpha_s}{4\pi^2} \sum_q e_q^2 f_q(\mathbf{p}_\gamma, x) T^2 \left[\ln \frac{4E_\gamma T}{m_{\text{th}}^2} + C \right], \quad (1)$$

where $C = -1.916$, $m_{\text{th}}^2 = g^2 T^2/6$, $\alpha_s = g^2/(4\pi)$ is the strong coupling, and α is the electromagnetic coupling; q denotes all light quark and antiquark species with charge e_q , and f_q is the distribution of minijets of flavor q . It is worth emphasizing that the conversion property of the process is reflected in (1) by the fact that the photon spectrum is directly proportional to the jet spectrum f_q . Calculations without some of the simplifying approximations [8] that lead to the expression in Eq. (1) produce only small modifications of the photon yield [10].

To calculate the yield from thermal emission and jet conversion, we need to model the space-time evolution of the fireball. We want to compare with data taken at midrapidity; hence we assume the boost-invariant Bjorken scenario for the longitudinal expansion. In addition we neglect the marginal transverse expansion of the plasma on the jet-photon conversion part (i.e., during the short time jets are propagating through the QGP). But the transverse expansion is fully taken into account for the thermal contribution.

We integrate over the conversion vertex x in (1) in terms of the space-time rapidity η , the radial coordinate \mathbf{r} (with $r = |\mathbf{r}|$) at which the jet is emitted, and the proper time τ , so that $d^4x = \tau d\tau d\eta d^2r$. We parametrize momenta \mathbf{p} by their rapidity y and transverse momentum \mathbf{p}_T ($p_T = |\mathbf{p}_T|$). Starting from the yield $dN_{\text{jet}}/(d^2p_T dy)$ of quarks initiating a jet, the distribution of hard partons in a boost-invariant scenario is given by [8,22]

$$f_q(\mathbf{p}, x) = \frac{(2\pi)^3}{g_q \pi R_T^2 \tau p_T} \frac{dN_{\text{jet}}}{d^2p_T dy} \rho(\mathbf{r}) \delta(y - \eta), \quad (2)$$

where R_T is the radius of the fireball, $g_q = 6$ is the spin-color degeneracy factor of parton q , and ρ is a transverse profile function normalized to one. For simplicity, we will take the profile to be azimuthally symmetric with area density $\rho(r) = 2(1 - r^2/R_T^2)\Theta(R_T - r)$ and transverse radius $R_T = 1.2(N_{\text{part}}/2)^{1/3}$.

The number of participants, N_{part} , is a function of centrality, but we neglect the azimuthal asymmetry in noncentral collisions. This approximation does not allow us to predict the azimuthal distribution of photons, but it introduces only a small error for the yield integrated over the azimuthal angle. The jet spectrum for an arbitrary impact parameter b is obtained by scaling

$$\frac{dN_{\text{jet}}}{d^2p_T dy}(b) = \frac{T_{AA}(b)}{T_{AA}(0)} \times \frac{dN_{\text{jet}}}{d^2p_T dy}(b=0). \quad (3)$$

This expression shows that the photon yield resulting from jet-photon conversion scales as a product of the nuclear thickness and a term dependent on the initial conditions, such as temperature and size, and on the history of the evolution of the fireball. A measurement of the photon spectra at different centralities could thus help reveal this information, as the primary hard photons will scale with the nuclear thickness T_{AA} alone.

We use a leading order perturbative calculation of jets and of direct Compton and annihilation photons [7], utilizing CTEQ5L parton distributions [23]. For gold nuclei EKS98 corrections of the nuclear parton distributions [24] are taken into account. Higher order perturbative effects are accounted for by a phenomenological K factor, which is known to depend weakly on the transverse momentum p_T at RHIC energies [25] and is here assumed to be constant.

Vacuum photon bremsstrahlung is obtained by convoluting the jet distributions with the photon fragmentation functions given in [7] ($\Lambda = 200 \text{ MeV}$), and we use KKP fragmentation functions [26] for the yield of neutral pions. We calculate the π^0 and photon cross sections at midrapidity for $p + p$ and compare these to the values measured by PHENIX. This allows us to fix the K factors for jet production,




FIG. 1. Invariant cross section $E d\sigma/d^3p$ for π^0 and γ production in $p + p$ collisions at $\sqrt{s} = 200$ GeV as a function of transverse momentum p_T at $y = 0$. Data points are from the PHENIX experiment [13,28]. We also show a full NLO calculation [27].

K_{jet} , and direct Compton and annihilation photons, K_γ , independently. We obtain a good description of the $p + p$ data with $K_{\text{jet}} = 1.8$ and $K_\gamma = 1.5$ (see Fig. 1). These values agree with those reported in Ref. [10] on the basis of a next-to-leading order (NLO) perturbative QCD calculation and our results also match the NLO calculation by Gordon and Vogelsang [27].

Once the K factors are fixed we apply them to calculate the yield of direct photons in Au + Au collisions. The prompt photon yield in Au + Au is close to that in $p + p$ scaled with T_{AA} , owing to small shadowing corrections. Note, however, that there is an effect from the admixture of $p + n$ and $n + n$ collisions with different photon yields. Therefore a simple scaling of $p + p$ results, as often invoked by the experimental collaborations for comparison, introduces a small but systematic bias of the photon yield from hard processes toward larger values.

Next we use the jet yields in Au + Au to calculate the photons from jet-photon conversions using (1) and (2). For the fireball we assume a thermally and chemically equilibrated plasma at some early time τ_0 with initial temperature T_0 . The product $T^3\tau$ is conserved during the isentropic longitudinal expansion, and its value can be fixed by the total hadron multiplicity dN_h/dy . The initial conditions are determined by imposing the thermalization condition $\tau_0 \approx 1/(3T_0)$. Table I lists the initial times and temperatures inferred from the

TABLE I. Number of participants N_{part} , nuclear thickness factor T_{AA} , initial time τ_0 , and initial temperature T_0 for different centrality bins.

	Centrality				
	0–10%	10–20%	20–30%	30–40%	40–50%
N_{part}	326.2	234.5	166.0	114.0	75.0
T_{AA} [1/mb]	22.75	14.35	8.00	5.23	2.86
τ_0 [fm/c]	0.151	0.166	0.182	0.201	0.221
T_0 [MeV]	0.434	0.395	0.361	0.328	0.297

charged hadron multiplicity $dN/d\eta$ measured by PHENIX [15] for different centrality bins. Nuclear thickness factors and number of participants are also given.

The transverse profile of the initial temperature is fixed by assuming that the energy density scales with the square of the nuclear thickness factors, so that $T_i(r) = T_0[2(1 - r^2/R_T^2)]^{1/4}$. This has been found to work well for the study of the centrality dependence of hadronic spectra [29] as well as lepton pair production [30] and should be sufficient for our purpose. Starting from the initial time τ_0 , we integrate (1) to either the time τ_f , when the medium has cooled to the critical temperature $T_f = 160$ MeV, or to the time when the jet reaches the boundary of the fireball, whichever comes first.

For the calculation of thermal photons we assume that the thermally and chemically equilibrated QGP, which undergoes a boost-invariant longitudinal and an azimuthally symmetric transverse expansion [31], converts to a hadronic gas below T_c . The production of photons from the deconfined phase is calculated using the complete leading order emission rate [4]. The latest results of Turbide *et al.* [6] for photon emission from hadronic matter are used.

Figures 2, 3, 4, and 5 present our results for photon yields at midrapidity in Au + Au collisions as a function of transverse momentum in four different centrality bins: 0–10%, 10–20%, 20–30%, and 40–50%. Both prompt photons and photons from jet-plasma interactions are shown, and the sum is compared to PHENIX data [14]. For completeness we also show our result for thermal photons. The agreement with data is generally quite good for all centrality bins. This is also true for the other centrality bins, up to the 60–70% bin, which are not shown here.

As already pointed out in Ref. [8], the jet-photon conversion spectrum falls off with p_T faster than the spectrum of prompt photons, a consequence of the additional hard momentum transfer with a propagator $\sim 1/p_T^2$. Interestingly the jet-photon conversion yield is as large as both the primary hard photon




FIG. 2. Photon yield $dN/(d^2p_T dy)$ as a function of p_T for $y = 0$ in central (0–10%) Au + Au collisions at $\sqrt{s} = 200$ GeV. We show primary hard photons (dashed curve), jet-photon conversion (solid and labeled curve), and the sum of both (uppermost solid curve). Thermal photons are also shown (solid and labeled curve). Data are from the PHENIX Collaboration [14].




FIG. 3. The same as Fig. 2 but for the 10–20% centrality bin.

yield and the thermal yield around $p_T = 2$ GeV/c in central Au + Au collisions. Unfortunately, the quality of data in this region is still poor. Between 4 and 6 GeV/c, where the data provide a better constraint, jet-medium photons still add roughly a 50% contribution to that of primary hard photons.

It has been argued that photons from primary hard processes alone are sufficient to describe the Au + Au data [14,21] and that no additional contribution is needed. The first statement is correct, to a certain extent, as can be seen in Fig. 2. However, its value is limited by the size of the experimental error bars. We have demonstrated here that a calculation taking into account jet-photon conversion in a consistent manner also leads to a result that is compatible with present data. Caution should be exercised when isospin effects are omitted in the discussion.

Energy-loss effects for jets before they convert into photons have been investigated by Turbide *et al.* [10]. The net effect on the yield of photons from jet-medium interactions is found to be small, about 20%. This is because most photons are emitted at a time when the plasma is still hot. Jets have then traveled only a short distance through the plasma and have not lost a significant amount of energy.

The centrality dependence of the data is well described by our calculations. Whereas direct photons scale with the number




FIG. 4. The same as Fig. 2 but for the 20–30% centrality bin.




FIG. 5. The same as Fig. 2 but for the 40–50% centrality bin.

of collisions, N_{coll} , photons from jet-medium interactions exhibit a slightly stronger scaling. The relative importance of this process thus decreases in more peripheral collisions. This can also be seen in Fig. 6, where we plot the nuclear modification factor

$$R_{AA} = \frac{dN_{AA}/dy|_{p_T > 6 \text{ GeV}/c}}{N_{\text{coll}} dN_{pp}/dy|_{p_T > 6 \text{ GeV}/c}}, \quad (4)$$

calculated from the photon spectra integrated for $p_T > 6$ GeV/c versus centrality. We show our calculation and a version taking into account an effective 20% energy loss of jets before conversion into photons and compare these with PHENIX data [14]. Owing to the large error bars the data are compatible with 1 for central and midperipheral collisions, in accordance with a binary collision scaling from $p + p$ collisions. Our calculation is compatible with the data as well and predicts a slight rise of R_{AA} with centrality. Our results show that the importance of the contribution from jet-photon conversion to the total photon spectrum grows below




FIG. 6. The nuclear modification factor R_{AA} for direct photons and $p_T > 6$ GeV/c as defined in (4) as a function of the number of participants N_{part} (solid line). We also show a calculation effectively taking into account energy loss of jets before conversion into photons (dashed line). Data are taken from the PHENIX Collaboration [14].

$p_T = 6$ GeV/c. A high-statistics measurement of R_{AA} in this region would be very useful.

In summary, we have calculated the photon spectrum in Au + Au collisions at RHIC energies resulting from primary hard photons, jets interacting with the medium, and thermal radiation. The calculations are consistent with neutral pion and photon spectra measured in $p + p$. We obtain a good description of the p_T dependence and centrality dependence of photon production in Au + Au. The contribution from jet-photon conversion can be as large as 100% (50%) that of

primary hard photons at 2 (5) GeV/c in accordance with data measured by PHENIX.

At the Large Hadron Collider, the importance of the jet-photon conversion process will be significantly enhanced. It has been shown that it will be the dominant contribution below $p_T \approx 12$ GeV/c [8,10].

This work was supported by DOE Grant Nos. DE-FG02-87ER40328 and DE-FG02-05ER41367.

-
- [1] J. W. Harris and B. Müller, *Annu. Rev. Nucl. Part. Sci.* **46**, 71 (1996).
- [2] E. L. Feinberg, *Nuovo Cim. A* **34**, 391 (1976).
- [3] J. Kapusta, P. Lichard, and D. Seibert, *Phys. Rev. D* **44**, 2774 (1991); **47**, 4171(E) (1993); R. Baier, H. Nakkagawa, A. Niegawa, and K. Redlich, *Z. Phys. C* **53**, 433 (1992); P. K. Roy, D. Pal, S. Sarkar, D. K. Srivastava, and B. Sinha, *Phys. Rev. C* **53**, 2364 (1996); P. Aurenche, F. Gelis, H. Zaraket, and R. Kobes, *Phys. Rev. D* **58**, 085003 (1998).
- [4] P. Arnold, G. D. Moore, and L. G. Yaffe, *J. High Energy Phys.* **12** (2001) 009.
- [5] L. Xiong, E. Shuryak, and G. E. Brown, *Phys. Rev. D* **46**, 3798 (1992); V. V. Goloviznin and K. Redlich, *Phys. Lett.* **B319**, 520 (1993); C. Song, *Phys. Rev. C* **47**, 2861 (1993).
- [6] S. Turbide, R. Rapp, and C. Gale, *Phys. Rev. C* **69**, 014903 (2004).
- [7] J. F. Owens, *Rev. Mod. Phys.* **59**, 465 (1987).
- [8] R. J. Fries, B. Müller, and D. K. Srivastava, *Phys. Rev. Lett.* **90**, 132301 (2003).
- [9] B. G. Zakharov, *JETP Lett.* **80**, 1 (2004).
- [10] S. Turbide, C. Gale, S. Jeon, and G. D. Moore, *Phys. Rev. C* **72**, 014906 (2005).
- [11] D. K. Srivastava, C. Gale, and R. J. Fries, *Phys. Rev. C* **67**, 034903 (2003).
- [12] See, e.g., S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. C* **71**, 034908 (2005); **71**, 049901(E) (2005), and references therein.
- [13] S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. D* **71**, 071102 (2005).
- [14] S. S. Adler *et al.* (PHENIX Collaboration), A. L. Landers, and J. A. Tancs, *Phys. Rev. Lett.* **94**, 23201 (2005).
- [15] S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **91**, 072301 (2003); *Phys. Rev. C* **69**, 034910 (2004).
- [16] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **91**, 172302 (2003).
- [17] C. Gale, T. C. Awes, R. J. Fries, and D. K. Srivastava, *J. Phys. G* **30**, S1013 (2004).
- [18] S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **91**, 072303 (2003).
- [19] S. Jeon, J. Jalilian-Marian, and I. Sarcevic, *Phys. Lett.* **B562**, 45 (2003).
- [20] R. Baier, M. Dirks, K. Redlich, and D. Schiff, *Phys. Rev. D* **56**, 2548 (1997); D. K. Srivastava and K. Geiger, *Phys. Rev. C* **58**, 1734 (1998); S. A. Bass, B. Müller, and D. K. Srivastava, *Phys. Rev. Lett.* **93**, 162301 (2004).
- [21] D. d'Enterria and D. Peressounko, *nucl-th/0503054*.
- [22] Z. Lin and M. Gyulassy, *Phys. Rev. C* **51**, 2177 (1995); **52**, 440(E) (1995).
- [23] H. L. Lai *et al.* (CTEQ Collaboration), *Eur. Phys. J. C* **12**, 375 (2000).
- [24] K. J. Eskola, V. J. Kolhinen, and C. A. Salgado, *Eur. Phys. J. C* **9**, 61 (1999).
- [25] G. G. Barnafoldi, G. I. Fai, P. Levai, G. Papp, and Y. Zhang, *J. Phys. G* **27**, 1767 (2001).
- [26] B. A. Kniehl, G. Kramer, and B. Pötter, *Nucl. Phys.* **B582**, 514 (2000).
- [27] L. E. Gordon and W. Vogelsang, *Phys. Rev. D* **48**, 3136 (1993); **49**, 170 (1994).
- [28] S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **91**, 241803 (2003).
- [29] D. K. Srivastava, *Phys. Rev. C* **64**, 064901 (2001); J.-Y. Ollitrault, *Phys. Lett.* **B273**, 31 (1991).
- [30] I. Kvasnikova, C. Gale, and D. K. Srivastava, *Phys. Rev. C* **65**, 064903 (2002).
- [31] D. K. Srivastava, *Phys. Rev. C* **71**, 034905 (2005).