

Photonuclear sum rules and the tetrahedral configuration of ${}^4\text{He}$

Doron Gazit,¹ Nir Barnea,¹ Sonia Bacca,² Winfried Leidemann,^{3,4,*} and Giuseppina Orlandini^{3,4,*}

¹*The Racah Institute of Physics, The Hebrew University, 91904 Jerusalem, Israel*

²*Gesellschaft für Schwerionenforschung, Planckstr. 1, D-64291 Darmstadt, Germany*

³*Department of Physics, George Washington University, Washington DC 20052, USA*

⁴*Istituto Nazionale di Fisica Nucleare, Gruppo Collegato di Trento, Italy*

(Received 15 September 2006; published 14 December 2006)

Three well-known photonuclear sum rules (SR), i.e., the Thomas-Reiche-Kuhn, the bremsstrahlungs and the polarizability SR are calculated for ${}^4\text{He}$ with the realistic nucleon-nucleon potential Argonne V18 and the three-nucleon force Urbana IX. The relation between these sum rules and the corresponding energy weighted integrals of the cross section is discussed. Two additional equivalences for the bremsstrahlungs SR are given, which connect it to the proton-neutron and neutron-neutron distances. Using them, together with our result for the bremsstrahlungs SR, we find a deviation from the tetrahedral symmetry of the spatial configuration of ${}^4\text{He}$. The possibility to access this deviation experimentally is discussed.

DOI: [10.1103/PhysRevC.74.061001](https://doi.org/10.1103/PhysRevC.74.061001)

PACS number(s): 21.45.+v, 25.20.Dc, 27.10.+h

Photonuclear sum rules (SR) are related to moments of different order of the photonuclear cross section and reflect important electromagnetic properties of nuclei [1,2]. In fact they can often be expressed in terms of simple ground-state properties in a model-independent or quasi-model-independent way. Well-known examples are the Thomas-Reiche-Kuhn (TRK) sum rule [3], which gives information about the importance of exchange effects in nuclear dynamics via the so-called TRK enhancement factor κ^{TRK} , the bremsstrahlungs sum rule (BSR) [4–7], which is connected to the nuclear charge radius and to the mean distance between protons [8], and the polarizability sum rule (PSR) [9], related to the electric nuclear polarizability. These SR are all assuming that the dominant contribution to the cross section comes from unretarded electric dipole (E1UR) transitions. Two- and three-body studies [10,11] indeed confirm that other contributions are much smaller. Much discussed is also the Gerasimov-Drell-Hearn (GDH) sum rule [12], which is related to the nuclear anomalous magnetic moment.

In this work we consider the TRK, BSR, and PSR of ${}^4\text{He}$ within a realistic nuclear potential model consisting of two- and three-body forces (AV18 and UIX [13,14]). (The GDH sum rule is trivial for ${}^4\text{He}$: it vanishes, because the ${}^4\text{He}$ total angular momentum is equal to zero.) We also investigate the related moments by integrating explicitly the properly weighted total photoabsorption cross section, which we have calculated for the same potential model [15]. The aim of this study is to show that in some cases sum rules can allow access experimentally to two-body properties of the nuclear ground state, like the proton-proton, neutron-neutron, and proton-neutron distances. In the case of ${}^4\text{He}$ this allows us to test the validity of the configuration tetrahedral symmetry of this nucleus and at the same time to “measure” the amount of symmetry breaking. This work aims also to provide a guideline

for experiments, where only lower bounds for the SR can be determined, as well as to give an idea of the reliability of the SR approach to heavier systems, where the direct theoretical determination of the cross section, and therefore its integration, is presently out of reach. The advantage to perform this kind of study in ${}^4\text{He}$, compared to analogous ones in the two- [16] and three-body systems [17], is that ${}^4\text{He}$ is a rather dense and compact nucleus, resembling heavier systems more closely. Only now that realistic theoretical results for the photonuclear cross section are available such a study is possible and one can put the extrapolation of the results to heavier systems on safer grounds.

We start by recalling the formalism of the photonuclear SR. The various moments of the photonuclear cross section are defined as

$$m_n(\bar{\omega}) \equiv \int_{\omega_{\text{th}}}^{\bar{\omega}} d\omega \omega^n \sigma_{\gamma}^{\text{E1UR}}(\omega), \quad (1)$$

where ω is the photon energy and ω_{th} and $\bar{\omega}$ indicate threshold energy and upper integration limit, respectively. With $\sigma_{\gamma}^{\text{E1UR}}(\omega)$ we indicate the unretarded dipole cross section given by

$$\sigma_{\gamma}^{\text{E1UR}}(\omega) = \mathcal{G}\omega R(\omega), \quad (2)$$

where $\mathcal{G} = 4\pi^2\alpha/3$ with α denoting the fine structure constant and the nucleus total spin, respectively. The response function $R(\omega)$ is given by

$$R(\omega) = \sum_n |\langle n|\mathbf{D}|0\rangle|^2 \delta(\omega - E_n + E_0), \quad (3)$$

where $|0/n\rangle$ and $E_{0/n}$ are the nuclear ground-/excited-state wave functions and energies, respectively, and \mathbf{D} is the unretarded dipole operator $\mathbf{D} = \sum_{i=1}^A \mathbf{r}_i \tau_i^3/2$, where A is the number of nucleons and τ_i^3 and \mathbf{r}_i are the third component of the isospin operator and the coordinate of the i th particle in the center-of-mass frame, respectively.

Assuming that $\sigma_{\gamma}^{\text{E1UR}}(\omega)$ converges to zero faster than ω^{-n-1} and applying the closure property of the eigenstates

*On leave of absence from the Department of Physics, University of Trento, I-38050 Povo (Trento) Italy.

of the Hamiltonian H , one has the following SR for $n = 0, -1, -2$:

$$\Sigma^{\text{TRK}} \equiv m_0(\infty) = \frac{\mathcal{G}}{2} \langle 0 | [\mathbf{D}, [H, \mathbf{D}]] | 0 \rangle \quad (4)$$

$$\Sigma^{\text{BSR}} \equiv m_{-1}(\infty) = \mathcal{G} \langle 0 | \mathbf{D} \cdot \mathbf{D} | 0 \rangle \quad (5)$$

$$\Sigma^{\text{PSR}} \equiv m_{-2}(\infty) = \mathcal{G} \sum_n (E_n - E_0)^{-1} |\langle n | \mathbf{D} | 0 \rangle|^2. \quad (6)$$

Working out the expressions in Eqs. (4)–(6), one finds that those moments are related to interesting properties of the system under consideration. In fact the TRK sum rule is also given by the well-known relation [3]

$$\Sigma^{\text{TRK}} = \mathcal{G} \frac{3NZ}{2mA} (1 + \kappa^{\text{TRK}}), \quad (7)$$

where N and Z are the neutron and proton numbers, respectively, m is the nucleon mass, and κ^{TRK} is the so-called TRK enhancement factor defined as

$$\kappa^{\text{TRK}} \equiv \frac{mA}{3NZ} \langle 0 | [\mathbf{D}, [V, \mathbf{D}]] | 0 \rangle. \quad (8)$$

From this expression it is evident that κ^{TRK} embodies the exchange effects of the nuclear potential V [the double commutator in Eq. (8) vanishes for systems like atoms, where no exchange effects are present].

In the literature one finds a few interesting equivalences for the bremsstrahlungs sum rule. Rewriting the dipole operator as $\mathbf{D} = (NZ/A)\mathbf{R}_{pN}$, where \mathbf{R}_{pN} denotes the distance between the proton and neutron centers of mass, one has [5]

$$\Sigma^{\text{BSR}} = \mathcal{G} \left(\frac{NZ}{A} \right)^2 \langle 0 | R_{pN}^2 | 0 \rangle. \quad (9)$$

In Ref. [6] Foldy demonstrated that

$$\Sigma^{\text{BSR}} = \mathcal{G} \frac{NZ}{A-1} \langle r_p^2 \rangle, \quad (10)$$

where $\langle r_p^2 \rangle$ is the mean square (m.s.) point proton radius

$$\langle r_p^2 \rangle \equiv \frac{1}{Z} \langle 0 | \sum_{i=1}^Z r_i^2 | 0 \rangle. \quad (11)$$

However, this relation is valid only under the assumption that the ground-state wave function is symmetric in the space coordinates of the nucleons.

In Ref. [7] it was found that, in the framework of the oscillator shell model, one has

$$\Sigma^{\text{BSR}} = \mathcal{G} (Z^2 \langle r_p^2 \rangle - Z^2 \langle r_p'^2 \rangle), \quad (12)$$

where $\langle r_p'^2 \rangle$ is the m.s. distance of protons with respect to the proton center of mass \mathbf{R}_P

$$\langle r_p'^2 \rangle \equiv \frac{1}{Z} \langle 0 | \sum_{i=1}^Z (\mathbf{r}_i - \mathbf{R}_P)^2 | 0 \rangle. \quad (13)$$

Later, in Ref. [8], it was shown that the validity of Eq. (12) is not limited to the oscillator shell model, but it is a model-independent relation, which can also be written as

$$\Sigma^{\text{BSR}} = \mathcal{G} \left[Z^2 \langle r_p^2 \rangle - \frac{Z(Z-1)}{2} \langle r_{pp}^2 \rangle \right], \quad (14)$$

where $\langle r_{pp}^2 \rangle$ is the m.s. proton-proton distance

$$\langle r_{pp}^2 \rangle \equiv \frac{1}{Z(Z-1)} \langle 0 | \sum_{i,j=1}^Z (\mathbf{r}_i - \mathbf{r}_j)^2 | 0 \rangle. \quad (15)$$

For the BSR two additional relations exist that are easy to prove but that, to our knowledge, have not been considered in the literature, i.e.,

$$\Sigma^{\text{BSR}} = \mathcal{G} \left[N^2 \langle r_n^2 \rangle - \frac{N(N-1)}{2} \langle r_{nn}^2 \rangle \right] \quad (16)$$

and

$$\Sigma^{\text{BSR}} = \mathcal{G} \frac{NZ}{2} (\langle r_{pn}^2 \rangle - \langle r_p^2 \rangle - \langle r_n^2 \rangle), \quad (17)$$

where $\langle r_n^2 \rangle$ is the m.s. point neutron radius and $\langle r_{\alpha\beta}^2 \rangle$ are the m.s. nucleon-nucleon (NN) distances, i.e.,

$$\langle r_{nn}^2 \rangle \equiv \frac{1}{N(N-1)} \langle 0 | \sum_{i,j=1}^N (\mathbf{r}_i - \mathbf{r}_j)^2 | 0 \rangle, \quad (18)$$

$$\langle r_{pn}^2 \rangle \equiv \frac{1}{NZ} \langle 0 | \sum_{i=1}^Z \sum_{j=1}^N (\mathbf{r}_i - \mathbf{r}_j)^2 | 0 \rangle. \quad (19)$$

It is interesting to note that (i) because Eq. (9) is invariant under the exchange of protons with neutrons, Eq. (16) represents the obvious corresponding expression for neutrons of Eq. (14), and (ii) Eqs. (14), (16), and (17), as well as Eq. (12), express Σ^{BSR} via the one-body quantity $\langle r_\alpha^2 \rangle$ and various two-body quantities. Finally, regarding the polarizability sum rule, one has

$$\Sigma^{\text{PSR}} = 2\pi^2 \alpha_D, \quad (20)$$

where α_D denotes the nuclear polarizability in the E1UR approximation.

In this work we calculate moments and sum rules in different ways. On the one hand, we obtain the moments by integrating our recent result for the ^4He total photoabsorption cross section [15]. This is an *ab initio* calculation where we use the AV18 NN potential and the UIX three-nucleon force. The results have been obtained by the Lorentz integral transform (LIT) method [18]. The necessary equations have been solved via hyperspherical harmonics expansions (EHH approach [19,20]). On the other hand, we obtain the SR in a more direct way, as explained in the following.

The LIT, an integral transform with a Lorentzian kernel, is defined as follows:

$$\mathcal{L}(\epsilon, \Gamma) = \int d\omega \frac{R(\omega)}{(\omega - \epsilon)^2 + \Gamma^2}. \quad (21)$$

One way to evaluate the LIT is by using the Lanczos technique as described in Ref. [21]. In fact the LIT can be re-expressed as

$$\mathcal{L}(\epsilon, \Gamma) = \frac{1}{\Gamma} \langle 0 | \mathbf{D} \cdot \mathbf{D} | 0 \rangle \text{Im} \left\{ \langle \phi_0 | \frac{1}{z - H} | \phi_0 \rangle \right\} \quad (22)$$

with $z = E_0 + \epsilon + i\Gamma$ and

$$|\phi_0\rangle = \frac{\mathbf{D}|0\rangle}{\sqrt{\langle 0 | \mathbf{D} \cdot \mathbf{D} | 0 \rangle}}. \quad (23)$$

It is evident that the LIT depends on the matrix element

$$x_{00}(z) = \langle \phi_0 | \frac{1}{z - H} | \phi_0 \rangle, \quad (24)$$

which can be expressed [22] as a continued fraction containing the Lanczos coefficients

$$a_i = \langle \phi_i | H | \phi_i \rangle, \quad b_i = \|b_i | \phi_i \rangle\|, \quad (25)$$

where the ϕ_i form the Lanczos orthonormal basis $\{|\phi_i\rangle, i = 0, \dots, n\}$. Therefore the implementation of the Lanczos algorithm leads to $\mathcal{L}(\epsilon, \Gamma)$ (for details see Ref. [21]). Although the inversion of the LIT [23] gives access to $R(\omega)$, and thus to the moments of Eq. (1), the normalization of the Lanczos ‘‘pivot’’ $|\text{LP}\rangle = \mathbf{D}|0\rangle$ and the Lanczos coefficients allow us to obtain the SR of Eqs. (4)–(6). In fact one has:

$$\Sigma^{\text{PSR}} = \mathcal{G} x_{00}(E_0), \quad (26)$$

$$\Sigma^{\text{BSR}} = \mathcal{G} \langle \text{LP} | \text{LP} \rangle, \quad (27)$$

$$\Sigma^{\text{TRK}} = (a_0 - E_0) \Sigma^{\text{BSR}}. \quad (28)$$

We use an HH basis, therefore the ground state $|0\rangle$, the Lanczos ‘‘pivot’’ $|\text{LP}\rangle$, and the Lanczos coefficients a_n are given in terms of HH expansions. Although for the ground state the expansion is characterized by an even hyperspherical grand-angular quantum number K and total isospin $T = 0$, $|\text{LP}\rangle$ has to be expanded on $\{K' = K + 1, T = 1\}$ states (we neglect the AV18 isospin mixing, which is very small, as shown for the ${}^4\text{He}$ ground state in Ref. [24]). The rate of convergence of the various SR results from Eqs. (26)–(28) is given in Table I as a function of the hyperspherical grand-angular quantum number K . One observes sufficiently good convergence patterns for Σ^{TRK} and Σ^{BSR} . For the latter we have performed an additional test of the convergence, calculating Σ^{BSR} directly as mean value of the operator $\mathbf{D} \cdot \mathbf{D}$ on the ground state [Eq. (5)]. In this way the expansion of $|\text{LP}\rangle$ on $\{K' = K + 1, T = 1\}$ states is avoided. We obtain practically identical results.

From Table I one sees that the convergence of Σ^{PSR} is slower when the three-nucleon force (3NF) is included. In Ref. [15] a related problem was found for the cross section itself. In fact there it was shown that the cross-section peak slightly shifts toward lower energies with increasing K and Fig. 3 of Ref. [15] shows that such a shift leaves the area under the peak almost unaltered. Therefore one would indeed expect that the sum rule Σ^{PSR} , which has the strongest inverse energy

TABLE I. Convergence in K of the SR for AV18+UIX potentials. The converged AV18 results are also shown.

K	Σ^{PSR} $10^{-2} \text{ (mb MeV}^{-1}\text{)}$	Σ^{BSR} (mb)	Σ^{TRK} 10^2 (mb MeV)
		AV18+UIX	
8	6.230	2.398	1.430
10	6.277	2.396	1.448
12	6.331	2.394	1.451
14	6.382	2.401	1.458
16	6.434	2.406	1.460
18	6.473	2.410	1.462
		AV18	
	7.681	2.696	1.383

TABLE II. Convergence in the upper integration limit $\bar{\omega}$ of the various moments defined in Eq. (1).

$\bar{\omega}$ (MeV)	$m_{-2}(\bar{\omega})$ $10^{-2} \text{ (mb MeV}^{-1}\text{)}$	$m_{-1}(\bar{\omega})$ (mb)	$m_0(\bar{\omega})$ 10^2 (mb MeV)
		AV18+UIX	
135	6.55	2.27	.944
300	6.55	2.37	1.14

weighting, is more sensitive to such a shift than the other two SR. In Ref. [15] an extrapolated cross section $\sigma_\gamma^{K \rightarrow \infty}$ was obtained from a Padè approximation. We have used this to determine from Eq. (1) the various moments for $\bar{\omega} = 300$ and 135 MeV. These results are listed in Table II.

For Σ^{TRK} one sees that the SR is not yet exhausted at 300 MeV. In fact more than 20% of the strength is still missing. At pion threshold, one has only about 2/3 of Σ^{TRK} . As discussed in Ref. [25] for the triton case, the rather strong contribution from higher energies seems to be connected to the strong short-range repulsion of the AV18 potential. As to the TRK enhancement factor we obtain $\kappa^{\text{TRK}} = 1.31$ for AV18 and 1.44 for AV18+UIX. These numbers are somewhat larger than older results obtained either with a variational wave function and AV14+UVII potential ($\kappa^{\text{TRK}} = 1.29$ [26]) or with more approximated wave functions and various soft and hard core NN potentials ($\kappa^{\text{TRK}} = 0.9 - 1.30$ [27–29]).

For Σ^{BSR} , as expected, the contribution at high energy is much smaller than for Σ^{TRK} , in fact at $\bar{\omega} = 300$ MeV we find a missing sum rule strength of less than 2% only. For Σ^{PSR} the strength beyond 300 MeV is even more negligible. Actually in this case the explicit integration leads to an even higher result than the sum rule evaluation of Eq. (27). The seeming contradiction is explained by the already-discussed fact that Σ^{PSR} is not yet convergent, and for the explicit integration the extrapolated cross section of Ref. [15]) is used. Indeed, integrating the $K = 18$ cross section we obtain a value of 6.46 mb, which is consistent with the corresponding sum rule result of 6.47 mb, within the numerical error of the calculation.

For the AV18+UIX force the value of the polarizability α_D that we deduce from the extrapolated Σ^{PSR} is 0.0655 fm^3 . The AV18 result, which already shows a good convergence for $K = 16$, is 0.0768 fm^3 . This means that the 3NF reduces the polarizability by 15%. It would be very interesting to measure this nuclear polarizability by Compton scattering, as a test of the importance of the three-body force on such a classical low-energy observable. We find that Σ^{PSR} is the SR that is affected most by the 3NF. In fact Σ^{BSR} is reduced by only 10% and one has an opposite effect on Σ^{TRK} with a 5% increase. The quenching or enhancement of SR due to the 3NF is the reflection of its effects on the cross section, i.e., a decrease of the peak and an increase of the tail. Altogether one can say that the 3NF appears to make the nucleus stiffer with respect to the application of a static electric field, i.e., it leads to a reduction of the dipole transition strengths at low energies and thus to a lower polarizability and bremsstrahlungs weighted sum rule. However, the high-energy strength is increased, presumably by increased short-range correlations, resulting in an enhanced TRK sum rule.

Here we add a few words about the very old question, already discussed in Ref. [4], of the “existence” of the SR [finiteness of $m_n(\bar{\omega})$ for $\bar{\omega} \rightarrow \infty$], which is connected to the high-energy falloff of the E1UR cross section. Because we find a rather good consistency between the SR and the moment values, we can state with a rather high degree of confidence that Σ^{TRK} (and consequently the other two SR) “exist.” Therefore we can try to extract some information about the high-energy behavior of the cross section and hence about the “existence” of SR with higher n . With an ω^{-p} ansatz for the falloff of the cross section above pion threshold, and requiring that Σ^{TRK} is 146 mb MeV (see Table I), one gets a rather weak energy falloff, i.e., $p \simeq 1.5$. This value is also consistent with Σ^{BSR} . In fact, adding such a tail contribution to $m_{-1}(135)$ one gets $\Sigma^{\text{BSR}} = 2.39$ mb, to be compared with 2.41 mb in Table I. The value of p might be somewhat different for other potentials, but probably it will not change much. Therefore we can rather safely conclude that higher-order SR do not exist for realistic nuclear potential models.

Finally we return to discuss the BSR, because it presents a very interesting aspect. As already mentioned Σ^{BSR} contains information about one- and two-body densities via $\langle r_\alpha^2 \rangle$ and $\langle r_{\alpha\beta}^2 \rangle$, respectively. This means that a measurement of Σ^{BSR} and the knowledge of the experimental m.s. radius allow us to determine $\langle r_{\alpha\beta}^2 \rangle$ via Eqs. (14), (16), and (17). In this way one gets information about the internal configuration of ${}^4\text{He}$ as it is explained in the following.

In his derivation of Eq. (10), Foldy assumed a totally symmetric ${}^4\text{He}$ spatial wave function, which corresponds to a configuration where the four nucleons are located at the four vertices of a tetrahedron. For such a configuration one has $\langle r_p^2 \rangle = \langle r_n^2 \rangle = \langle r^2 \rangle$ and $\langle r_{pp}^2 \rangle = \langle r_{nn}^2 \rangle = \langle r_{np}^2 \rangle$ with $Q_T \equiv \langle r_{\alpha\beta}^2 \rangle / \langle r^2 \rangle = 8/3$. Foldy’s assumption is a very good approximation for ${}^4\text{He}$, but other spatial symmetries (mixed symmetry, antisymmetric) are also possible. What can be learned from our Σ^{BSR} result with respect to this question? For ${}^4\text{He}$, which is a $T = 0$ system one can safely assume that $\langle r_p^2 \rangle = \langle r^2 \rangle = \langle r_n^2 \rangle$ (isospin mixing is tiny [24]). Using in Eqs. (14), (16), and (17) the AV18+UIX value of $\langle r^2 \rangle = 2.04 \text{ fm}^2$ [15] (which coincides with the experimental one [30], corrected for the proton charge radius) and Σ^{BSR} from Table I, we obtain $\langle r_{pp}^2 \rangle = \langle r_{nn}^2 \rangle = 5.67 \text{ fm}^2$ and $\langle r_{pn}^2 \rangle = 5.34 \text{ fm}^2$, i.e., two values that differ by about 6%. The ratios $Q_{pp(nn)} \equiv \langle r_{pp(nn)}^2 \rangle / \langle r^2 \rangle$ and $Q_{np} \equiv \langle r_{np}^2 \rangle / \langle r^2 \rangle$ are not much different from the correspondent value Q_T of a classical tetrahedral configuration. We obtain $Q_{pp} = 2.78$ and $Q_{np} = 2.62$ instead of $Q_T = 2.67$. One notices that $Q_{pp(nn)} - Q_T \simeq 2(Q_{np} - Q_T)$. This reflects the different numbers of proton-proton and neutron-neutron pairs (2) with respect to proton-neutron pairs (4). Using Eq. (9) one can also derive the distance between the proton and neutron centers of mass. One has $R_{pn} = 1.58 \text{ fm}$ instead of 1.65 fm for the tetrahedral configuration.

Notice that with $\langle r^2 \rangle = 2.04 \text{ fm}^2$ the “tetrahedral” BSR would be 2.62 mb, the same value that one obtains using Eq. (10). This value is 9% larger than our result. The distortions that we find from our 9% smaller BSR are the consequence of the different effects of the potential on isospin triplet and isospin singlet pairs.

We conclude that when considered in its body frame ${}^4\text{He}$ should look like a slightly *deformed* tetrahedron. Of course, this statement has to be interpreted in a quantum mechanical sense, regarding the mean-square values of the nucleon-nucleon distances on the two-body density. It is clear that one cannot measure this *deformation* “directly,” because it is not a deformation of the one-body charge density (the ${}^4\text{He}$ charge density has only a monopole). However, such a *deformation* is accessible experimentally in an indirect way via the measurements of the charge radius and of Σ^{BSR} .

This leads to the question how exactly Σ^{BSR} can be measured in a photonuclear experiment. Two points have to be addressed: (i) the contributions of $E1$ retardation and higher multipoles that are not contained in Σ^{BSR} but that will contribute to the experimental cross section and (ii) the contribution of the high-energy tail. To this end it is instructive to consider the results from the three-nucleon photodisintegration. Regarding point (i), additional effects of the $E1$ retardation and higher multipoles have been calculated in Ref. [31] for the AV18+UIX potentials in a Faddeev calculation. Using those results one finds that $E1$ retardation and higher multipoles increase $m_{-1}(135)$ by about 1% only. In fact according to Gerasimov [32] there is a large cancellation of $E1$ retardation and other multipoles. There is no reason for a larger effect in ${}^4\text{He}$. On the contrary, because the leading isovector magnetic dipole ($M1$) transition is suppressed in this nucleus (it is zero for an S wave) one can expect an even smaller contribution. As to point (ii) considering the falloff of the triton E1UR cross section around pion threshold from Ref. [31] one obtains a result very similar to our ${}^4\text{He}$ case, namely $p \simeq 1.5$. However, including the other multipole contributions one gets a considerably smaller value, namely $p \simeq 1.1$. This is no contradiction with the small increase of 1% for Σ^{BSR} , because the inverse energy-weighted cross section integrated from 100 to 135 MeV gives only a rather small contribution to the sum rule. However, one would overestimate Σ^{BSR} taking the full cross section. Thus we suggest that the tail contribution to an experimental BSR should be estimated using the theoretically established falloff ω^{-p} with $p \simeq 1.5$.

In conclusion, we have studied three well-known photonuclear sum rules for ${}^4\text{He}$ within a realistic two- and three-nucleon potential model. Two new equivalences for the BSR sum rules have allowed us to deduce information about two-body properties of the nuclear ground state, like the proton-proton, neutron-neutron, and proton-neutron distances. In particular we have tested the validity of the configuration tetrahedral symmetry of this nucleus and have found that this symmetry is slightly broken. We have suggested an experimental way to access this symmetry breaking via a measurement of the BSR that could be performed in one of the existing or planned low to intermediate energy photonuclear facility.

D.G., N.B., W.L., and G.O. thank the Institute for Nuclear Theory at the University of Washington for its hospitality and the Department of Energy for partial support during the completion of this work. The work of D.G. and N.B. was supported by the Israel Science Foundation (grant 361/05).

- [1] E. Lipparini and S. Stringari, *Phys. Rep.* **175**, 103 (1989).
- [2] G. Orlandini and M. Traini, *Rep. Prog. Phys.* **54**, 257 (1991).
- [3] R. Ladenburg and F. Reiche, *Naturwiss* **11**, 584 (1923); W. Kuhn, *Z. Phys.* **33**, 408 (1925); L. H. Thomas, *Naturwiss* **13**, 627 (1925).
- [4] J. S. Levinger and H. A. Bethe, *Phys. Rev.* **78**, 115 (1950).
- [5] D. M. Brink, *Nucl. Phys.* **4**, 215 (1957).
- [6] L. L. Foldy, *Phys. Rev.* **107**, 1303 (1957).
- [7] A. Dellafiore and D. M. Brink, *Nucl. Phys.* **A286**, 474 (1977).
- [8] A. Dellafiore and E. Lipparini, *Nucl. Phys.* **A388**, 639 (1982).
- [9] J. L. Friar, *Ann. Phys. (NY)* **95**, 170 (1975).
- [10] H. Arenhövel and M. Sanzone, *Few-Body Syst. Suppl.* **3**, 1 (1991).
- [11] J. Golak, R. Skibinski, W. Glöckle, H. Kamada, A. Nogga, H. Witala, V. D. Efros, W. Leidemann, G. Orlandini, and E. L. Tomusiak, *Nucl. Phys.* **A707**, 365 (2002).
- [12] S. B. Gerasimov, *Yad. Fiz.* **2**, 598 (1965) [*Sov. J. Nucl. Phys.* **2**, 430 (1966)]; S. D. Drell and A. C. Hearn, *Phys. Rev. Lett.* **16**, 908 (1966).
- [13] R. B. Wiringa, V. G. J. Stoks, and R. Schiavilla, *Phys. Rev. C* **51**, 38 (1995).
- [14] B. S. Pudliner, V. R. Pandharipande, J. Carlson, S. C. Pieper, and R. B. Wiringa, *Phys. Rev. C* **56**, 1720 (1997).
- [15] D. Gazit, S. Bacca, N. Barnea, W. Leidemann, and G. Orlandini, *Phys. Rev. Lett.* **96**, 112301 (2006).
- [16] H. Arenhövel and W. Fabian, *Nucl. Phys.* **A292**, 429 (1977).
- [17] V. D. Efros, W. Leidemann, and G. Orlandini, *Phys. Lett.* **B408**, 1 (1997).
- [18] V. D. Efros, W. Leidemann, and G. Orlandini, *Phys. Lett.* **B338**, 130 (1994).
- [19] N. Barnea, W. Leidemann, and G. Orlandini, *Phys. Rev. C* **61**, 054001 (2000); *Nucl. Phys.* **A693**, 565 (2001).
- [20] N. Barnea, V. D. Efros, W. Leidemann, and G. Orlandini, *Few-Body Syst.* **35**, 155 (2004).
- [21] M. A. Marchisio, N. Barnea, W. Leidemann, and G. Orlandini, *Few-Body Syst.* **33**, 259 (2003).
- [22] E. Dagotto, *Rev. Mod. Phys.* **66**, 763 (1994), and references therein.
- [23] V. D. Efros, W. Leidemann, and G. Orlandini, *Few-Body Syst.* **26**, 251 (1999); D. Andreasi, W. Leidemann, C. Reiss, and M. Schwamb, *Eur. Phys. J. A* **24**, 361 (2005).
- [24] A. Nogga, H. Kamada, W. Glöckle, and B. R. Barrett, *Phys. Rev. C* **65**, 054003 (2002); M. Viviani, A. Kievsky, and S. Rosati, *ibid.* **71**, 024006 (2005).
- [25] V. D. Efros, W. Leidemann, G. Orlandini, and E. L. Tomusiak, *Phys. Lett.* **B484**, 223 (2000).
- [26] R. Schiavilla, A. Fabrocini, and V. R. Pandharipande, *Nucl. Phys.* **A473**, 290 (1987).
- [27] W. T. Weng, T. T. S. Kuo, and G. E. Brown, *Phys. Lett.* **B46**, 329 (1973).
- [28] W. Heinze, H. Arenhövel, and G. Horlacher, *Phys. Lett.* **B76**, 379 (1978).
- [29] M. Gari, H. Hebach, B. Sommer, and J. G. Zabolitzky, *Phys. Rev. Lett.* **41**, 1288 (1978).
- [30] E. Borie and G. A. Rinker, *Phys. Rev. A* **18**, 324 (1978).
- [31] J. Golak, R. Skibinski, W. Glöckle, H. Kamada, A. Nogga, H. Witala, V. D. Efros, W. Leidemann, G. Orlandini, and E. L. Tomusiak, *Nucl. Phys.* **A707**, 365 (2002).
- [32] S. B. Gerasimov, *Phys. Lett.* **13**, 240 (1964).