

$\pi B_8 B_8$ and $\sigma B_8 B_8$ couplings from a chiral quark potential model

M. T. Fernández-Caramés and P. González

Departamento de Física Teórica e IFIC, Universidad de Valencia-CSIC, E-46100 Burjassot, Valencia, Spain

A. Valcarce

Departamento de Física Fundamental, Universidad de Salamanca, E-37008 Salamanca, Spain

(Received 21 February 2008; published 30 May 2008)

From an $SU(2) \otimes SU(2)$ chiral quark potential model incorporating spontaneous chiral symmetry breaking the asymptotic π and σ exchange pieces of the NN potential are generated. From them the πNN and σNN coupling constants can be extracted. The generalization to $SU(3) \otimes SU(3)$ allows for a determination of $\pi B_8 B_8$ and $\sigma B_8 B_8$ coupling constants according to exact $SU(3)$ hadron symmetry. The implementation of the values of the couplings at $Q^2 = 0$ provided by QCD sum rules and/or phenomenology makes also feasible the extraction of the meson-baryon-baryon form factors. In this manner a quite complete knowledge of the couplings may be attained.

DOI: [10.1103/PhysRevC.77.054003](https://doi.org/10.1103/PhysRevC.77.054003)

PACS number(s): 12.39.Jh, 12.39.Fe, 14.20.Gk, 21.30.Fe

I. INTRODUCTION

The meson-baryon-baryon (mBB') couplings play a central role in hadron physics concerning the baryon-baryon (BB') interactions as well as the formation and decay of baryon resonances. To study these couplings effective hadron Lagrangians involving the mesons and baryons under consideration are postulated. All the complexity of the mBB' vertices is assumed to be taken into account through running couplings depending on Q^2 , the transfer momentum in the vertex. This dependency is usually parametrized in terms of a form factor and a coupling constant defined as the value of the running coupling at a particular Q^2 , usually on-shell $Q^2 = M_m^2$. Therefore, one can calculate physical processes and compare to data to extract these values. Thus the πNN coupling constant is obtained from πN and/or NN -scattering data [1]. For couplings involving baryons and/or mesons for which scattering or decay data are not so complete or unavailable one can also rely on symmetry to derive predictions, see for instance Ref. [2].

From a more fundamental point of view hadrons are made up of quarks. Hence hadron structures and decays as well as hadron-hadron interactions should come out from quark dynamics as dictated by quantum chromodynamics (QCD). Due to the technical difficulty to achieve this objective at present quark models of hadrons incorporating QCD-motivated symmetries and dynamics have been successfully applied to generate the baryon-baryon interactions, and consistently the baryon spectrum, in the light (u, d) [3] as well as in the light+strange (u, d, s) quark sectors [4–6]. These models, sometimes less precise than effective hadronic treatments, offer the advantage of providing a consistent unified description of all baryon-baryon processes from the same Hamiltonian at the quark level. This confers them in principle a great predictive power once the model parameters are tightly constrained from some selected set of existing data.

We shall make use of this power to predict, within a nonrelativistic chiral quark model framework, $mB_8 B_8$ (B_8 : baryon of the flavor octet) coupling constants in terms of meson-quark-quark (mqq) couplings. More precisely we shall

generate, from a mqq Lagrangian incorporating the effect of spontaneous chiral symmetry breaking (SCSB), the quark-quark meson exchange potentials and from them, through a Born-Oppenheimer (BO) approximation, the asymptotic baryon-baryon-meson exchange interactions. The use of justified harmonic oscillator baryon wave functions (in terms of quarks) will allow us to perform analytic calculations. By comparing the resulting interactions to the ones postulated at the effective hadronic level we shall identify the meson-baryon-baryon coupling constants. This procedure has been applied in the literature to πNN coupling [7]. Here we shall be more precise in the extraction of coupling constants and form factors and we extend its application to the σ meson and the other B_8 baryons, making feasible the comparison of our results to the ones obtained with alternative methods based on quark or hadron degrees of freedom.

The contents are organized as follows. In Sec. II we shall center on the light quark sector where spontaneously broken chiral $SU(2) \otimes SU(2)$ symmetry serves as an underlying general guide to generate the quark-quark meson-exchange potentials. We shall revisit the calculation of the πNN coupling in terms of the πqq one and apply the same procedure to the σNN case. We shall also comment on the possibility of applying our method to Δ and nucleon resonances. Then in Sec. III we consider the extension, via chiral $SU(3) \otimes SU(3)$, to the $SU(3)$ octet of baryons. Finally in Sec. IV we summarize our main results and conclusions.

II. LIGHT QUARK SECTOR: $SU(2) \otimes SU(2)$.

Our starting point is the chiral Lagrangian

$$\mathcal{L}_{\text{ch}} = -g_{\text{ch}} \bar{q} [\mathbb{1}_{SU(2)} \sigma + i \gamma_5 \vec{\tau} \vec{\pi}] q, \quad (1)$$

where q has components u and d and g_{ch} is the chiral mqq coupling constant ($m : \pi, \sigma$). The spontaneous chiral symmetry breaking gives rise to a vertex form factor $F(Q^2)$ [8]. From the parametrization given in Refs. [9,10] we propose the Lorentz invariant form (note that for the purpose of the

derivation of static potentials $Q^2 = -\vec{Q}^2$),

$$F(Q^2) = \left(\frac{\Lambda^2}{\Lambda^2 - Q^2} \right)^{1/2}, \quad (2)$$

where Λ is an effective cutoff parameter fitted from data. We should keep in mind that this parametrization of $F(Q^2)$ makes only sense for $Q^2 < \Lambda^2$.

From the form proposed [$F(Q^2 = 0) = 1$] it is clear that g_{ch} represents the value of the mqq coupling, $g_{\text{ch}}F(Q^2)$, at $Q^2 = M_m^2$ in the limit $M_m^2 = 0$. To deal with coupling constants defined on the physical meson masses ($M_\pi^2 \neq 0 \neq M_\sigma^2$) we identify (it is implicitly assumed that $M_\pi^2, M_\sigma^2 < \Lambda^2$)

$$g_{\pi qq} \equiv g_{\text{ch}} \left(\frac{\Lambda^2}{\Lambda^2 - M_\pi^2} \right)^{1/2} \quad (3)$$

$$g_{\sigma qq} \equiv g_{\text{ch}} \left(\frac{\Lambda^2}{\Lambda^2 - M_\sigma^2} \right)^{1/2}, \quad (4)$$

as the values of the coupling $g_{\text{ch}}F(Q^2)$ at $Q^2 = M_\pi^2$ and $Q^2 = M_\sigma^2$, respectively.

Let us point out that the use of the experimental pion mass may have required the introduction of an explicit symmetry breaking term proportional to σ in the Lagrangian but this term has no further effect in the analysis we perform. Regarding the σ mass it will be taken as a parameter to be fitted from data around the value provided by the SCSB relation [11],

$$M_\sigma^2 - M_\pi^2 = 4M_q^2, \quad (5)$$

where M_q denotes the constituent quark mass at $Q^2 = 0$.

In terms of $g_{\pi qq}$ we write the pion Lagrangian as

$$\mathcal{L}_{\pi qq} = -g_{\pi qq} \bar{q} i\gamma_5 \vec{\tau} \vec{\pi} q \quad (6)$$

with a vertex form factor $F_{\pi qq}(Q^2)$ given by

$$F_{\pi qq}(Q^2) \equiv \left(\frac{\Lambda^2 - M_\pi^2}{\Lambda^2 - Q^2} \right)^{1/2}. \quad (7)$$

Analogously, the σ Lagrangian reads

$$\mathcal{L}_{\sigma qq} = -g_{\sigma qq} \bar{q} \mathbb{1}_{\text{SU}(2)} \sigma q \quad (8)$$

with a vertex form factor $F_{\sigma qq}(Q^2)$ given by

$$F_{\sigma qq}(Q^2) \equiv \left(\frac{\Lambda^2 - M_\sigma^2}{\Lambda^2 - Q^2} \right)^{1/2}. \quad (9)$$

Note that both form factors are of the same type $F_{mqq}(Q^2) \equiv (\Lambda^2 - M_m^2 / \Lambda^2 - Q^2)^{1/2}$ so that $F_{mqq}(Q^2 = M_m^2) = 1$.

A. mqq and induced mBB potentials

From \mathcal{L}_{mqq} and the form factors the static OPE and OSE central potentials (to the respective lowest order in Q^2) can be obtained through a nonrelativistic reduction of the

corresponding Feynman diagram amplitudes. They are

$$V_{\text{OPE}}^{ij}(\vec{r}_{ij}) = \frac{1}{3} \frac{g_{\pi qq}^2}{4\pi} \frac{M_\pi^2}{4M_i M_j} M_\pi \times \left[Y(M_\pi r_{ij}) - \frac{\Lambda^3}{M_\pi^3} Y(\Lambda r_{ij}) \right] (\vec{\sigma}_i \cdot \vec{\sigma}_j)(\vec{\tau}_i \cdot \vec{\tau}_j), \quad (10)$$

$$V_{\text{OSE}}^{ij}(\vec{r}_{ij}) = -\frac{g_{\sigma qq}^2}{4\pi} M_\sigma \left[Y(M_\sigma r_{ij}) - \frac{\Lambda}{M_\sigma} Y(\Lambda r_{ij}) \right]. \quad (11)$$

Here i and j are numbers denoting quarks, $M_{i,j} = M_q$, $\vec{\sigma}_{i,j}(\vec{\tau}_{i,j})$ are the spin (isospin) Pauli operators, r_{ij} is the interquark distance and the function Y is defined as,

$$Y(x) = \frac{e^{-x}}{x}. \quad (12)$$

Once the potentials at the quark level have been derived we use them to obtain the baryon-baryon potentials. From V_{OmE}^{ij} the asymptotic baryon-baryon meson exchange static potential is defined as

$$(V_q)_{\text{OmE}}^{B_a B_b \rightarrow B_c B_d}(R \rightarrow \infty) \equiv \lim_{R \rightarrow \infty} \langle \Psi_{B_c B_d} | \sum_{\substack{i \in B_a, B_c \\ j \in B_b, B_d}} V_{\text{OmE}}^{ij} | \Psi_{B_a B_b} \rangle, \quad (13)$$

where Ψ_{B_i, B_j} stands for the two-baryon wave function and R for the interbaryon distance and the integration is over the quark coordinates.

We concentrate on $B_a B_b \rightarrow B_c B_d$ interactions involving baryons with the same mass. Then the asymptotic two-baryon wave function will be expressed in the center-of-mass system as

$$\Psi_{B_a B_b} = \Phi_{B_a}(1, 2, 3; +\vec{R}/2) \Phi_{B_b}(4, 5, 6; -\vec{R}/2), \quad (14)$$

where (1, 2, 3) and (4, 5, 6) denote the quarks forming the baryons and $\pm\vec{R}/2$ the baryons position and

$$\Phi_{B_i} = (\Phi_{B_i})_{\text{spatial}} (\Phi_{B_i})_{\text{spin-flavor}} (\Phi_{B_i})_{\text{color}} \quad (15)$$

is the one-baryon wave function expressed as the direct product of its spatial, spin-flavor and color parts. For the sake of simplicity the baryon spatial wave function will be chosen of harmonic oscillator type

$$(\Phi_B)_{\text{spatial}}(1, 2, 3; +\vec{R}/2) = \prod_{i=1}^3 \left(\frac{1}{\pi b^2} \right)^{3/4} \exp[-(\vec{r}_i - \vec{R}/2)^2 / 2b^2] \quad (16)$$

with an harmonic oscillator parameter, b , related to the size of the baryon.

B. Parameters

To fix the parameters at the quark level: M_q , M_π , M_σ , g_{ch} , Λ , and b , we rely on the efficient description of NN data provided by the chiral quark cluster model (CQCM) [3]. Such a model contains, apart from OPE and OSE potentials derived from \mathcal{L}_{mqq} , a confinement plus a residual one-gluon exchange (OGE) interactions. We should realize, though, that

TABLE I. Quark model parameters [3].

M_q (MeV)	b (fm)	M_σ (fm ⁻¹)	M_π (fm ⁻¹)	Λ (fm ⁻¹)
313	0.518	3.42	0.7	4.2

the precise fitting of the parameters in the CQCM relies on a RGM calculation so the two-baryon wave function as well as the baryon-baryon potentials are different from the ones obtained via the Born-Oppenheimer (BO) approach we follow, Eq. (13). We take this into account in an effective manner by keeping the same values for $M_q, M_\pi, M_\sigma, \Lambda$, and b , and fitting a new value for g_{ch} to reproduce, with our BO approach, the experimental value of the pion-nucleon-nucleon coupling constant (see next section). The values of the parameters used henceforth are listed in Table I.

C. πNN

From Eqs. (10), (13), and (16) the asymptotic OPE central potential for $NN \rightarrow NN$, corresponding to the diagram of Fig. 1 (there are nine equivalent ones), can be obtained. It reads (the calculation has been explicitly done in Ref. [7])

$$(V_q)_{\text{OPE}}^{NN \rightarrow NN}(R \rightarrow \infty) = \frac{g_{\pi qq}^2}{4\pi} \left[9 \langle (\vec{\sigma}_3 \cdot \vec{\sigma}_6)_{NN} \cdot (\vec{\tau}_3 \cdot \vec{\tau}_6)_{NN} \rangle e^{\frac{M_\pi^2 b^2}{2}} \right] \frac{M_\pi^2}{4M_q^2} \frac{1}{3} \frac{e^{-M_\pi R}}{R}. \tag{17}$$

To extract the πNN coupling we have to compare this potential with the one derived from a postulated hadronic Lagrangian. Assuming, for instance, a pseudoscalar coupling we can write a Lagrangian

$$\mathcal{L}_{\pi NN} = -(g_{\pi NN})_{Q^2=M_\pi^2} \bar{N} i \gamma_5 \vec{\tau} \vec{\pi} N, \tag{18}$$

with a vertex form factor $G_{\pi NN}(Q^2)$ so that $G_{\pi NN}(Q^2 = M_\pi^2) = 1$. Note that we have indicated explicitly the on-shell character of the coupling $g_{\pi NN}$ through the subindex. To derive from this Lagrangian a Yukawa-like pion exchange potential, monopole- or dipole-type form factors are usually assumed. Concerning the asymptotic potential both give the same result. We shall use in parallel with the form factor at the quark level




FIG. 1. Asymptotic NN OPE or OSE interactions at quark level.




FIG. 2. Asymptotic NN OPE or OSE interactions at baryon level.

a form

$$G_{\pi NN}(Q^2) \equiv \left(\frac{\Lambda_{\pi NN}^2 - M_\pi^2}{\Lambda_{\pi NN}^2 - Q^2} \right)^{\frac{1}{2}}, \tag{19}$$

valid for $Q^2 < \Lambda_{\pi NN}^2$ and $M_\pi^2 < \Lambda_{\pi NN}^2$ ($\Lambda_{\pi NN}$ is a cutoff parameter to be fitted).

Note that if we had preferred to refer the Lagrangian to the value of the coupling at $Q^2 = 0$, i.e., to $(g_{\pi NN})_{Q^2=0}$, then we would have a different form factor such that

$$(g_{\pi NN})_{Q^2=M_\pi^2} G_{\pi NN}(Q^2) = (g_{\pi NN})_{Q^2=0} \left(\frac{\Lambda_{\pi NN}^2}{\Lambda_{\pi NN}^2 - Q^2} \right)^{\frac{1}{2}}, \tag{20}$$

where the second term on the right-hand side represents the form factor normalized at $Q^2 = 0$.

From $\mathcal{L}_{\pi NN}$ and $G_{\pi NN}(Q^2)$ the nonrelativistic reduction of the one-pion exchange diagram, Fig. 2, to the lowest order in Q^2 , provides us with the pion exchange static potential at the baryonic level. The asymptotic behavior of its central part is given by

$$(\mathcal{V}_B)_{\text{OPE}}^{NN \rightarrow NN}(R \rightarrow \infty) = \frac{(g_{\pi NN}^2)_{Q^2=M_\pi^2} \langle (\vec{\sigma}_N \cdot \vec{\sigma}_N) (\vec{\tau}_N \cdot \vec{\tau}_N) \rangle}{4\pi} \frac{M_\pi^2}{4M_N^2} \frac{1}{3} \frac{e^{-M_\pi R}}{R}. \tag{21}$$

Note that $(\mathcal{V}_B)_{\text{OPE}}^{NN \rightarrow NN}(R \rightarrow \infty)$ depends only on the coupling constant and not on the form factor. Then no information on the form factors at the baryon level can be extracted from it. Regarding the coupling constant we can make use of the relation $(25/9) \langle (\vec{\sigma}_N \cdot \vec{\sigma}_N) (\vec{\tau}_N \cdot \vec{\tau}_N) \rangle = 9 \langle (\vec{\sigma}_3 \cdot \vec{\sigma}_6)_{NN} \cdot (\vec{\tau}_3 \cdot \vec{\tau}_6)_{NN} \rangle$ [7] to compare Eqs. (17) and (21). From this comparison we extract

$$(g_{\pi NN}^2)_{Q^2=M_\pi^2} = g_{\pi qq}^2 \frac{M_N^2}{M_q^2} \frac{25}{9} e^{\frac{M_\pi^2 b^2}{2}}. \tag{22}$$

Having chosen $M_q = 313$ MeV so that $M_N = 3M_q$ we can re-express

$$(g_{\pi NN}^2)_{Q^2=M_\pi^2} = 25 g_{\pi qq}^2 e^{\frac{M_\pi^2 b^2}{2}}. \tag{23}$$

1. Pseudovector couplings

Alternatively to $g_{\pi qq}$ and $g_{\pi NN}$ we could have used pseudovector couplings $f_{\pi qq}$ and $f_{\pi NN}$ so that

$$\mathcal{L}_{\pi qq}^{PV} = -\frac{f_{\pi qq}}{M_\pi} \bar{q} i \gamma_5 \gamma_\mu \vec{q} \partial^\mu \vec{\pi} q \quad (24)$$

with the vertex form factor $F_{\pi qq}(Q^2)$ and

$$\mathcal{L}_{\pi NN}^{PV} = -\frac{(f_{\pi NN})_{Q^2=M_\pi^2}}{M_\pi} \bar{N} i \gamma_5 \gamma_\mu \vec{q} \partial^\mu \vec{\pi} N. \quad (25)$$

with the vertex form factor $G_{\pi NN}(Q^2)$.

It turns out that \mathcal{L}^{PV} give rise to exactly the same potentials as \mathcal{L} , to the lowest Q^2 order, under the identifications

$$\frac{f_{\pi qq}^2}{M_\pi^2} = \frac{g_{\pi qq}^2}{4M_q^2} \quad (26)$$

$$\frac{(f_{\pi NN}^2)_{Q^2=M_\pi^2}}{M_\pi^2} = \frac{(g_{\pi NN}^2)_{Q^2=M_\pi^2}}{4M_N^2}. \quad (27)$$

If we now substitute these relations in Eq. (22) we get

$$(f_{\pi NN}^2)_{Q^2=M_\pi^2} = f_{\pi qq}^2 \frac{25}{9} e^{\frac{M_\pi^2 b^2}{2}}. \quad (28)$$

The corresponding constant f_{ch} is consistently defined as

$$f_{\text{ch}} = \left(\frac{\Lambda^2 - M_\pi^2}{\Lambda^2} \right)^{1/2} f_{\pi qq} = \lim_{M_\pi \rightarrow 0} f_{\pi qq}. \quad (29)$$

2. Coupling constants and form factor values

From a standard experimental value $(f_{\pi NN}^2)_{Q^2=M_\pi^2}/4\pi \simeq 0.079$ (see Ref. [1] and references therein) or $(g_{\pi NN}^2)_{Q^2=M_\pi^2}/4\pi \simeq 14.6$ we fit the pion-quark-quark coupling constant

$$f_{\pi qq}^2/4\pi = 0.027 \quad (30)$$

or

$$g_{\pi qq}^2/4\pi = 0.55 \quad (31)$$

and

$$|g_{\text{ch}}| \equiv |f_{\text{ch}}| \frac{2M_q}{M_\pi} = 2.6. \quad (32)$$

Let us emphasize that this value for g_{ch} differs less than a 10% from the one obtained via QCD sum rules (QCDSR) $(g_{\text{ch}})_{\text{QCDSR}} \simeq 2.83$ [12].

Regarding $\Lambda_{\pi NN}$, the cutoff parameter, its range of values can be estimated. In Ref. [13] a fit to data was attained from a Gaussian form factor $e^{Q^2/\Lambda_{\text{HM}}^2}$ with Λ_{HM} varying from 2.6 to 4.2 fm^{-1} . This form factor is normalized at $Q^2 = 0$. By requiring its low Q^2 behavior $(1 + Q^2/\Lambda_{\text{HM}}^2)$ to be the same than that of our form factor normalized at $Q^2 = 0$ in Eq. (20), $(1 + Q^2/2\Lambda_{\pi NN}^2)$, we get $\Lambda_{\text{HM}}^2 = 2\Lambda_{\pi NN}^2$. Hence the resulting range for $\Lambda_{\pi NN}$ is

$$\Lambda_{\pi NN} = 1.84\text{--}2.97 \text{ fm}^{-1}. \quad (33)$$

From Eqs. (19) and (20) this range can be translated in an interval of values for the coupling at $Q^2 = 0$:

$$\frac{(f_{\pi NN}^2)_{Q^2=0}}{4\pi} = 0.068\text{--}0.075, \quad (34)$$

$$\frac{(g_{\pi NN}^2)_{Q^2=0}}{4\pi} = 12.5\text{--}13.8.$$

These values are in perfect agreement with the phenomenological analysis done in Ref. [14] (let us comment that for the form factor used in this reference the range of values for the cutoff parameter is the same as for Λ_{HM}). The preferred value in Ref. [14] is $(f_{\pi NN}^2)_{Q^2=0}/4\pi \simeq 0.073$, which corresponds to $(g_{\pi NN}^2)_{Q^2=0}/4\pi \simeq 13.5$. It is worthwhile to point out that this corresponds quite approximately to the $M_\pi \rightarrow 0$ limit of our expression $(f_{\pi NN}^2)_{Q^2=M_\pi^2} = f_{\pi qq}^2 (25/9) e^{\frac{M_\pi^2 b^2}{2}}$, i.e.,

$$\frac{(f_{\pi NN}^2)_{Q^2=M_\pi^2 \rightarrow 0}}{4\pi} = \frac{f_{\text{ch}}^2}{4\pi} \frac{25}{9} = 0.072, \quad (35)$$

indicating the quite approximate Goldstone boson character of the pion.

D. σNN

By proceeding in exactly the same way for the σ exchange we obtain from Eqs. (11), (13), and (16) at the quark level

$$(V_q)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty) = -\frac{g_{\sigma qq}^2}{4\pi} 9 e^{\frac{M_\sigma^2 b^2}{2}} \frac{e^{-M_\sigma R}}{R}. \quad (36)$$

At the baryonic level from the Lagrangian

$$\mathcal{L}_{\sigma NN} = -(g_{\sigma NN})_{Q^2=M_\sigma^2} \bar{N} \mathbb{1}_{\text{SU}(2)\sigma} N, \quad (37)$$

with a vertex form factor

$$G_{\sigma NN}(Q^2) = \left(\frac{\Lambda_{\sigma NN}^2 - M_\sigma^2}{\Lambda_{\sigma NN}^2 - Q^2} \right)^{\frac{1}{2}}, \quad (38)$$

we get

$$(V_B)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty) = -\frac{(g_{\sigma NN})_{Q^2=M_\sigma^2}}{4\pi} \frac{e^{-M_\sigma R}}{R}. \quad (39)$$

From their comparison

$$(g_{\sigma NN})_{Q^2=M_\sigma^2} = 9 g_{\sigma qq}^2 e^{\frac{M_\sigma^2 b^2}{2}}. \quad (40)$$

Let us note that once the value of $g_{\pi qq}$ has been fitted our model predicts, from Eqs. (3) and (4), the value of $g_{\sigma qq}$ through

$$\frac{g_{\sigma qq}^2}{g_{\pi qq}^2} = \frac{\Lambda^2 - M_\pi^2}{\Lambda^2 - M_\sigma^2}. \quad (41)$$

Equivalently, from Eqs. (23) and (40)

$$\frac{(g_{\sigma NN})_{Q^2=M_\sigma^2}}{(g_{\pi NN})_{Q^2=M_\pi^2}} = \frac{9}{25} \frac{\Lambda^2 - M_\pi^2}{\Lambda^2 - M_\sigma^2} e^{\frac{(M_\sigma^2 - M_\pi^2)b^2}{2}}. \quad (42)$$

1. Coupling constant and form factor values

From Eq. (42) we predict

$$\frac{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}}{4\pi} = 68.2 \quad (43)$$

and from Eq. (40)

$$\frac{g_{\sigma qq}^2}{4\pi} = 1.59. \quad (44)$$

As mentioned above our asymptotic comparison does not give any information on the cutoff parameter $\Lambda_{\sigma NN}$. Nonetheless we can combine our result for the coupling constant with the value of the coupling at $Q^2 = 0$ provided by QCDSR to get an insight into it. From Ref. [15]

$$\frac{(g_{\sigma NN})_{Q^2=0}}{g_q^\sigma} = 3.9 \pm 1.0. \quad (45)$$

If we tentatively identify g_q^σ with our $g_{\sigma qq}$ ($= 4.47$) we get

$$\frac{(g_{\sigma NN}^2)_{Q^2=0}}{4\pi} = 17.4 \pm 4.5, \quad (46)$$

and using the relation $(g_{\sigma NN}^2)_{Q^2=0} = (g_{\sigma NN}^2)_{Q^2=M_\sigma^2} G_{\sigma NN}^2(Q^2 = 0)$ we extract

$$\Lambda_{\sigma NN} = \left[\frac{M_\sigma^2}{1 - \frac{(g_{\sigma NN}^2)_{Q^2=0}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}}} \right]^{\frac{1}{2}} = 3.97 \pm 0.18 \text{ fm}^{-1}. \quad (47)$$

It is again interesting to consider the limit $M_\sigma \rightarrow 0$ of Eq. (40)

$$\frac{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2 \rightarrow 0}}{4\pi} = \frac{9g_{\text{ch}}^2}{4\pi} = 4.8, \quad (48)$$

or

$$\frac{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2 \rightarrow 0}}{g_{\text{ch}}^2} = 9 \quad (49)$$

and compare it to the interval of values of the coupling at $Q^2 = 0$ from Eq. (46). As can be seen the $M_\sigma \rightarrow 0$ value from Eq. (48) is out of this interval. This might be interpreted as reflecting the non-Goldstone boson nature of the σ .

E. $\pi N \Delta$

Strictly speaking our procedure to extract the couplings makes sense only when the lowest-order expansion in Q^2 we follow is simultaneously valid at the baryonic level ($E_B \simeq M_B$) and at the quark level ($E_q \simeq M_q$). We do not expect this to be true for light-quark baryons in general because the quarks can move relativistically inside them. However the structure of the ground states, the N [and Λ , Σ , and Ξ when considering SU(3)], is well described by nonrelativistic constituent quark models through the effective parameters in the quark-quark potential. Then we expect our procedure to make sense for them. Regarding other baryon states like Δ and $N(1440)$ one should be more cautious as we illustrate next.

To extract the $\pi N \Delta$ coupling constant we consider the $NN \rightarrow N \Delta$ interaction. According to the harmonic oscillator

model we are using the spatial wave function of Δ has exactly the same structure than the N one. However the real Δ differs from the N . To implement the bigger size for Δ predicted by nonrelativistic spectroscopic models we consider the possibility of a slightly different value for the size parameter. Thus the Δ spatial wave function we shall use is

$$(\Phi_\Delta)_{\text{spatial}}(4, 5, 6; +\vec{R}/2) = \prod_{i=4}^6 \left(\frac{1}{\pi b_\Delta^2} \right)^{\frac{3}{4}} \exp[-(\vec{r}_i - \vec{R}/2)^2/2b_\Delta^2] \quad (50)$$

with a baryon size parameter, b_Δ . One should also keep in mind that the mass of the Δ is a 30% bigger than the mass of the N . In our harmonic oscillator quark model this means that the quarks in the Δ have more potential and kinetic energy (virial theorem) than the quarks in the N . According to our comments above this could give rise to corrections in the expression of the static potential at the quark level. Moreover, due to the $\Delta - N$ mass difference the positions of the baryons in the initial and final states should not be the same. Therefore we should not expect an accurate prediction for the coupling constant in this case.

To derive such prediction we first calculate at the quark level, from Eqs. (13), (16), and (50) (we use for easiness the pseudovector form of the coupling), the asymptotic $NN \rightarrow N \Delta$ pion exchange static central potential (Fig. 3). It is

$$(V_q)_{\text{OPE}}^{NN \rightarrow N \Delta}(R \rightarrow \infty) = \frac{f_{\pi qq}^2}{4\pi} \left[\frac{160}{9\sqrt{2}} 2\sqrt{2} \frac{b_\mu^6 b_+^3 b_e^3}{(b^{15} b_\Delta^9)^{\frac{1}{2}}} e^{-\frac{M_\pi^2 b_\Delta^2}{2} R} \right] \frac{1}{3} \frac{e^{-M_\pi R}}{R}, \quad (51)$$

where the factor $160/9\sqrt{2}$ corresponds to $\langle N \Delta | (\vec{\sigma}_3 \cdot \vec{\sigma}_6) (\vec{\tau}_3 \cdot \vec{\tau}_6) | NN \rangle$ for total spin and isospin equal to 1, multiplied by 9, the number of equivalent diagrams, and b_μ , b_+ and b_e are defined as

$$\begin{aligned} \frac{1}{b_\mu^2} &= \frac{1}{2b^2} + \frac{1}{2b_\Delta^2} \\ \frac{1}{b_\pm^2} &= \pm \frac{1}{b^2} + \frac{1}{b_\mu^2} \\ \frac{1}{b_e^2} &= \frac{1}{2b_+^2} - \frac{b_+^2}{2b_-^4}. \end{aligned} \quad (52)$$




FIG. 3. Asymptotic $NN \rightarrow N \Delta$ OPE interaction at quark level.

Note that when $b_\Delta = b$ we also have $b_e = b_\mu = \sqrt{2}b_+ = b$, hence reproducing, except for the spin-isospin factor, the NN result.

At the baryonic level we have the Lagrangians $\mathcal{L}_{\pi NN}^{PV}$ as given by Eq. (25) and

$$\mathcal{L}_{\pi N\Delta}^{PV} = -\frac{(f_{\pi N\Delta})_{Q^2=M_\pi^2}}{M_\pi} \bar{N} \vec{T} \partial \vec{\pi} \Delta, \quad (53)$$

with a form factor $G_{\pi N\Delta}(Q^2)$ which we shall choose as,

$$G_{\pi N\Delta}(Q^2) \equiv \left(\frac{\Lambda_{\pi N\Delta}^2 - M_\pi^2}{\Lambda_{\pi N\Delta}^2 - Q^2} \right)^{\frac{1}{2}}, \quad (54)$$

In Eq. (53) Δ corresponds to a Rarita-Schwinger 3/2 spinor field and \vec{T} is the isospin nucleon- Δ transition operator. The corresponding pion exchange static central potential behaves asymptotically as

$$(\mathcal{V}_B)_{\text{OPE}}^{NN \rightarrow N\Delta}(R \rightarrow \infty) = \frac{(f_{\pi NN})_{Q^2=M_\pi^2} (f_{\pi N\Delta})_{Q^2=M_\pi^2}}{4\pi} \times [(\vec{\sigma}_N \cdot \vec{S})(\vec{\tau}_N \cdot \vec{T})] \frac{1}{3} \frac{e^{-M_\pi R}}{R}, \quad (55)$$

where \vec{S} is the spin nucleon- Δ transition operator and $\langle (\vec{\sigma}_N \cdot \vec{S})(\vec{\tau}_N \cdot \vec{T}) \rangle = 8/3$ for total spin and isospin equal to 1. Thus we identify

$$\frac{(f_{\pi NN})_{Q^2=M_\pi^2} (f_{\pi N\Delta})_{Q^2=M_\pi^2}}{4\pi} = \frac{f_{\pi qq}^2}{4\pi} \left[\frac{20}{3\sqrt{2}} 2\sqrt{2} \frac{b_\mu^6 b_+^3 b_e^3}{(b^{15} b_\Delta^9)^{\frac{1}{2}}} e^{\frac{M_\pi^2 b_e^2}{2}} \right]. \quad (56)$$

By using Eq. (28) we get

$$\frac{(f_{\pi N\Delta})_{Q^2=M_\pi^2}}{(f_{\pi NN})_{Q^2=M_\pi^2}} = \left[\frac{6\sqrt{2}}{5} 2\sqrt{2} \frac{b_\mu^6 b_+^3 b_e^3}{(b^{15} b_\Delta^9)^{\frac{1}{2}}} e^{\frac{M_\pi^2 (b_e^2 - b^2)}{2}} \right] \quad (57)$$

so that for $b_\Delta = b$ one obtains the usual spin-isospin relation $(f_{\pi N\Delta})_{Q^2=M_\pi^2} / (f_{\pi NN})_{Q^2=M_\pi^2} = 6\sqrt{2}/5$.

For b_Δ in the interval $[b, 1.2b]$ we predict $(f_{\pi N\Delta})_{Q^2=M_\pi^2} / 4\pi = [0.23, 0.20]$ to be compared to $(f_{\pi N\Delta})_{Q^2=M_\pi^2} / 4\pi \simeq 0.37$, estimated from the Δ decay to $N\pi$. This discrepancy seems to confirm our initial expectations. If instead $f_{\pi qq}$ we had written an effective $(f_{\pi qq})_{N\Delta}$ as a manner to take into account the $\Delta - N$ mass difference effect, then the needed value to reproduce the experimental number would have been $(f_{\pi qq})_{N\Delta} / f_{\pi qq} = [1.28, 1.37]$ for $b_\Delta \in [b, 1.2b]$, i.e., $(f_{\pi qq})_{N\Delta}$ should be a 30% bigger than $f_{\pi qq}$.

For nucleon resonances in general and in particular for $N^*(1440)$ we expect the calculation of the coupling constants to be much more uncertain. As a matter of fact the importance of relativistic corrections in the description of the structure and decay of $N^*(1440)$ in terms of quarks have been long emphasized in the literature (see for instance Ref. [16]). Furthermore, the nature of the $N^*(1440)$ may involve more than a simple $3q$ structure and the coupling of $q\bar{q}$ pairs to the meson structure can be relevant. Therefore the calculation of $\pi NN^*(1440)$ and $\sigma NN^*(1440)$ coupling constants carried out

in a preceding article within the same framework [17] should be considered too simplistic. A less approximative calculation for the pion case [involving also $\pi N\Delta$, $\pi N\Delta(1600)$, and $\pi NN(1535)$] has been carried out in Ref. [18] with Poincaré covariant constituent quark models with instant, point, and front forms of relativistic kinematics; from the persistent deviation from data of the calculated results the authors suggest the presence of sizable $qqqq\bar{q}$ components in the baryon wave functions.

III. LIGHT AND STRANGE QUARK SECTOR: SU(3) \otimes SU(3)

The generalization of the chiral Lagrangian to SU(3) \otimes SU(3) is straightforward. It is expressed as

$$\tilde{\mathcal{L}}_{\text{ch}} = -\tilde{g}_{\text{ch}} \bar{q} \left(\sum_{a=0}^8 \sigma_a \lambda_a + i \sum_{a=0}^8 \gamma_5 \pi_a \lambda_a \right) q, \quad (58)$$

where q has components u, d and s , σ_0 , and π_0 stand for the scalar and pseudoscalar meson singlets, and σ_i and π_i ($i = 1 \dots 8$) are the scalar and pseudoscalar meson octets.

To derive a potential involving the exchange of σ , the SU(2) singlet, we shall assume the ideal mixing

$$\begin{aligned} \sigma_0 &= \sqrt{2/3} \sigma + \sqrt{1/3} (s\bar{s}) \\ \sigma_8 &= \sqrt{1/3} \sigma - \sqrt{2/3} (s\bar{s}). \end{aligned} \quad (59)$$

When substituting these expressions in Eq. (58) the piece containing the σ and the $\vec{\pi}$ read,

$$\tilde{\mathcal{L}}_{\text{ch}(\pi,\sigma)} = -\tilde{g}_{\text{ch}} \bar{q} [\sigma(\sqrt{2/3}\lambda_0 + \sqrt{1/3}\lambda_8) + i\gamma_5 \vec{\tau} \vec{\pi}] q, \quad (60)$$

where $\lambda_0 \equiv \sqrt{2/3} \mathbb{1}_{\text{SU}(3)}$ and $\lambda_8 \equiv \sqrt{3} Y$, being Y the hypercharge. It is then clear that for $q = u, d$ ($Yu = 1/3 = Yd$) one formally recovers the SU(2) \otimes SU(2) Lagrangian: $\mathbb{1}_{\text{SU}(2)} \equiv 2/3 \mathbb{1}_{\text{SU}(3)} + Y$.

Again we take into account SCSB through a vertex form factor $\tilde{F}(Q^2)$

$$\tilde{F}(Q^2) = \left(\frac{\tilde{\Lambda}^2}{\tilde{\Lambda}^2 - Q^2} \right)^{1/2}. \quad (61)$$

Note also that the SCSB relation, $M_\sigma^2 - M_\pi^2 = 4M_q^2$, is preserved because it is derived for a nonstrange σ [11].

A. Parameters

One should realize that the values of the couplings $\tilde{g}_{\text{ch}} \tilde{F}(Q^2)$ (or equivalently the on-shell couplings $\tilde{g}_{\pi qq}, \tilde{g}_{\sigma qq}$ and the cutoff parameter $\tilde{\Lambda}$) and the size parameter \tilde{b} in SU(3) \otimes SU(3) need not be the same as in SU(2) \otimes SU(2). This is easily understandable by thinking for instance of the extra contribution to the NN interaction coming from η, η' , and a_0 in SU(3) \otimes SU(3). This contribution is taken into account in an effective manner in SU(2) \otimes SU(2), where no η, η' , and a_0 are present, through the fitted values of $g_{\pi qq}, g_{\sigma qq}, \Lambda$, and b . Fortunately we can correlate the variations of b and Λ (or \tilde{b}

and $\tilde{\Lambda}$) through the relation Eq. (42),

$$\begin{aligned} \frac{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}}{(g_{\pi NN}^2)_{Q^2=M_\pi^2}} &= \frac{9}{25} \frac{\Lambda^2 - M_\pi^2}{\Lambda^2 - M_\sigma^2} e^{\frac{(M_\sigma^2 - M_\pi^2)b^2}{2}} \\ &= \frac{9}{25} \frac{\tilde{\Lambda}^2 - M_\pi^2}{\tilde{\Lambda}^2 - M_\sigma^2} e^{\frac{(M_\sigma^2 - M_\pi^2)\tilde{b}^2}{2}}. \end{aligned} \quad (62)$$

Then from the selected experimental value $(g_{\pi NN}^2)_{Q^2=M_\pi^2}/4\pi \simeq 14.6$ and from our prediction $(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}/4\pi = 68.2$ we get, for a typical value of $\tilde{\Lambda} \simeq 5.2 \text{ fm}^{-1}$ ($\simeq 1.0 \text{ GeV}$) (note that it has to be higher than any mass of the scalar or pseudoscalar meson octets) an harmonic oscillator parameter $\tilde{b} \simeq 0.6 \text{ fm}$. Then from equivalent relations to Eqs. (23) and (40) we get

$$\frac{\tilde{g}_{\pi qq}^2}{4\pi} = 0.54 \quad (63)$$

$$\frac{\tilde{g}_{\sigma qq}^2}{4\pi} = 0.93, \quad (64)$$

and

$$\tilde{g}_{\text{ch}} = g_{\text{ch}}. \quad (65)$$

For the sake of completeness we give the B_8 spatial wave function in the $SU(3)_{\text{flavor}}$ limit,

$$\begin{aligned} (\Phi_{B_8})_{\text{spatial}}(1, 2, 3; +\vec{R}/2) \\ = \prod_{i=1}^3 \left(\frac{1}{\pi \tilde{b}^2} \right)^{\frac{3}{4}} \exp[-(\vec{r}_i - \vec{R}/2)^2/2\tilde{b}^2]. \end{aligned} \quad (66)$$

B. $\sigma B_8 B_8$

According to our preceding discussions the σqq Lagrangian will be written as

$$\tilde{\mathcal{L}}_{\sigma qq} = -\tilde{g}_{\sigma qq} \bar{q} \mathbb{1}_{SU(2)} \sigma q, \quad (67)$$

with a vertex form factor

$$\tilde{F}_{\sigma qq}(Q^2) \equiv \left(\frac{\tilde{\Lambda}^2 - M_\sigma^2}{\tilde{\Lambda}^2 - Q^2} \right)^{1/2}. \quad (68)$$

From this Lagrangian it is clear that the only difference when calculating the asymptotic potential at the quark level for the several $B_8 B_8$'s has to do with the number of the pairs of light quarks (u, d) in them allowing for the exchange of the σ , i.e., with the number of equivalent diagrams, Figs. 1 and 4. This number is 9 for NN , 4 for $\Lambda\Lambda$ and $\Sigma\Sigma$, and 1 for $\Xi\Xi$. Thus

$$\begin{aligned} (\tilde{V}_q)_{\text{OSE}}^{\Lambda\Lambda \rightarrow \Lambda\Lambda}(R \rightarrow \infty) &= (\tilde{V}_q)_{\text{OSE}}^{\Sigma\Sigma \rightarrow \Sigma\Sigma}(R \rightarrow \infty) \\ &= \frac{4}{9} (\tilde{V}_q)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty) \end{aligned} \quad (69)$$

$$(\tilde{V}_q)_{\text{OSE}}^{\Xi\Xi \rightarrow \Xi\Xi}(R \rightarrow \infty) = \frac{1}{9} (\tilde{V}_q)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty).$$

At the baryonic level we write the Lagrangian as

$$\mathcal{L}_{\sigma B_8 B_8} \equiv -(g_{\sigma B_8 B_8})_{Q^2=M_\sigma^2} \bar{B}_8 \sigma B_8, \quad (70)$$




FIG. 4. Asymptotic $\Lambda\Lambda$, $\Sigma\Sigma$, and $\Xi\Xi$ OPE and OSE interactions at quark level. Thin lines stand for light (u, d) quarks and thick lines for strange (s) quarks.

with a vertex form factor

$$G_{\sigma B_8 B_8}(Q^2) = \left(\frac{\Lambda_{\sigma B_8 B_8}^2 - M_\sigma^2}{\Lambda_{\sigma B_8 B_8}^2 - Q^2} \right)^{1/2}. \quad (71)$$

so that $G_{\sigma B_8 B_8}(Q^2 = M_\sigma^2) = 1$ and where the $(g_{\sigma B_8 B_8})_{Q^2=M_\sigma^2}$ are expressed in conventional notation

$$\begin{aligned} (g_{\sigma NN})_{Q^2=M_\sigma^2} &\equiv \sqrt{2/3} g_{s,1} + 1/3 g_{s,+} (4\alpha_s - 1) \\ (g_{\sigma \Lambda\Lambda})_{Q^2=M_\sigma^2} &\equiv \sqrt{2/3} g_{s,1} - 2/3 g_{s,+} (1 - \alpha_s) \\ (g_{\sigma \Sigma\Sigma})_{Q^2=M_\sigma^2} &\equiv \sqrt{2/3} g_{s,1} + 2/3 g_{s,+} (1 - \alpha_s) \\ (g_{\sigma \Xi\Xi})_{Q^2=M_\sigma^2} &\equiv \sqrt{2/3} g_{s,1} - 1/3 g_{s,+} (1 + 2\alpha_s), \end{aligned} \quad (72)$$

being $g_{s,1} \equiv (g_{\sigma_0 B_8 B_8})_{Q^2=M_\sigma^2}$ the scalar $SU(3)$ singlet coupling constant, $g_{s,+} \equiv g_{s,D} + g_{s,F}$ the sum of the D (symmetric) and the F (antisymmetric) scalar coupling constants in $SU(3)$ and $\alpha_s \equiv g_{s,F}/g_{s,+}$ the $F/(F + D)$ ratio of the scalar octet.

From this baryonic Lagrangian the following relations between the asymptotic potentials come out (from Fig. 2 and similar figures with N substituted by Λ, Σ , or Ξ)

$$\begin{aligned} (\mathcal{V}_B)_{\text{OSE}}^{\Lambda\Lambda \rightarrow \Lambda\Lambda}(R \rightarrow \infty) \\ = \frac{(g_{\sigma \Lambda\Lambda}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} (\mathcal{V}_B)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty) \\ (\mathcal{V}_B)_{\text{OSE}}^{\Sigma\Sigma \rightarrow \Sigma\Sigma}(R \rightarrow \infty) \\ = \frac{(g_{\sigma \Sigma\Sigma}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} (\mathcal{V}_B)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty) \\ (\mathcal{V}_B)_{\text{OSE}}^{\Xi\Xi \rightarrow \Xi\Xi}(R \rightarrow \infty) \\ = \frac{(g_{\sigma \Xi\Xi}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} (\mathcal{V}_B)_{\text{OSE}}^{NN \rightarrow NN}(R \rightarrow \infty). \end{aligned} \quad (73)$$

From the comparison of the asymptotic potentials at the quark and baryon levels we immediately get relations between the coupling constants

$$\frac{(g_{\sigma \Lambda\Lambda}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} = \frac{(g_{\sigma \Sigma\Sigma}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} = \frac{4}{9} \quad (74)$$

$$\frac{(g_{\sigma\Xi\Xi}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} = \frac{1}{9}.$$

We should emphasize that these ratios are preserved in the limit $M_\sigma \rightarrow 0$, i.e.,

$$\frac{(g_{\sigma\Lambda\Lambda}^2)_{Q^2=M_\sigma^2 \rightarrow 0}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2 \rightarrow 0}} = \frac{(g_{\sigma\Sigma\Sigma}^2)_{Q^2=M_\sigma^2 \rightarrow 0}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2 \rightarrow 0}} = \frac{4}{9} \quad (75)$$

$$\frac{(g_{\sigma\Xi\Xi}^2)_{Q^2=M_\sigma^2 \rightarrow 0}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2 \rightarrow 0}} = \frac{1}{9}.$$

It is interesting to compare these $M_\sigma \rightarrow 0$ ratios with the average QCDSR predictions at $Q^2 = 0$ in the SU(3) limit. From Ref. [19] we take

$$\begin{aligned} \frac{(g_{\sigma\Lambda\Lambda}^2)_{Q^2=0}}{(g_{\sigma NN}^2)_{Q^2=0}} &= (0.43)^2 = 0.19 \\ \frac{(g_{\sigma\Sigma\Sigma}^2)_{Q^2=0}}{(g_{\sigma NN}^2)_{Q^2=0}} &= (0.91)^2 = 0.83 \\ \frac{(g_{\sigma\Xi\Xi}^2)_{Q^2=0}}{(g_{\sigma NN}^2)_{Q^2=0}} &= (0.08)^2 = 0.006. \end{aligned} \quad (76)$$

As can be checked our ratios differ by a factor 2 (1/2) for $\Lambda - N$ and $\Sigma - N$ ($\Xi - N$) from the QCDSR ones. We may interpret this again as a reflection of the non-Goldstone character of the σ meson.

1. Coupling constants and form factors values

From Eq. (74) and from the calculated value $(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}/4\pi = 68.2$ we predict the coupling constants

$$\frac{(g_{\sigma\Lambda\Lambda}^2)_{Q^2=M_\sigma^2}}{4\pi} = \frac{(g_{\sigma\Sigma\Sigma}^2)_{Q^2=M_\sigma^2}}{4\pi} = 30.3 \quad (77)$$

$$\frac{(g_{\sigma\Xi\Xi}^2)_{Q^2=M_\sigma^2}}{4\pi} = 7.6.$$

Concerning the $F/(F+D)$ ratio we have from Eq. (74) $(g_{\sigma\Lambda\Lambda})_{Q^2=M_\sigma^2} = (g_{\sigma\Sigma\Sigma})_{Q^2=M_\sigma^2}$ what implies from Eq. (72)

$$\alpha_s = 1, \quad (78)$$

or

$$g_{S,D} = 0. \quad (79)$$

Regarding $g_{s,1}$ and $g_{s,+}$, we can use the fact that from the ideal mixing we have assumed $g_{(s\bar{s})NN} = -1/\sqrt{3} g_{s,1} + \sqrt{2} g_{s,F} - \sqrt{2}/3 g_{s,D} = 0$. Then from Eq. (79) we immediately obtain $g_{s,+} = 1/\sqrt{6} g_{s,1}$. If we substitute this relation and Eq. (78) in the first expression of Eq. (72) we get $(g_{\sigma NN})_{Q^2=M_\sigma^2} = 3g_{s,+}$, from where

$$\frac{g_{s,+}^2}{4\pi} = 7.6 \quad (80)$$

$$\frac{g_{s,1}^2}{4\pi} = 45.5, \quad (81)$$

satisfying

$$\frac{g_{s,1}^2}{g_{s,+}^2} = 6. \quad (82)$$

With respect to the cutoff parameters $\Lambda_{\sigma B_8 B_8}$ we can tentatively use our on-shell couplings ratios, Eq. (74), altogether with the QCDSR ones at $Q^2 = 0$ detailed above, Eq. (76), to establish a range of variation for them. Explicitly we can write

$$\left[\frac{(g_{\sigma B_8 B_8}^2)_{Q^2=0}}{(g_{\sigma NN}^2)_{Q^2=0}} \right] = \left[\frac{(g_{\sigma B_8 B_8}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} \right] \frac{\left(\frac{\Lambda_{\sigma B_8 B_8}^2 - M_\sigma^2}{\Lambda_{\sigma B_8 B_8}^2} \right)}{\left(\frac{\Lambda_{\sigma NN}^2 - M_\sigma^2}{\Lambda_{\sigma NN}^2} \right)}, \quad (83)$$

or equivalently

$$\Lambda_{\sigma B_8 B_8} = \Lambda_{\sigma NN} \left[\frac{M_\sigma^2}{\Lambda_{\sigma NN}^2 (1 - x_{B_8}) + x_{B_8} M_\sigma^2} \right]^{\frac{1}{2}}, \quad (84)$$

where

$$x_{B_8} \equiv \frac{\left[\frac{(g_{\sigma B_8 B_8}^2)_{Q^2=0}}{(g_{\sigma NN}^2)_{Q^2=0}} \right]}{\left[\frac{(g_{\sigma B_8 B_8}^2)_{Q^2=M_\sigma^2}}{(g_{\sigma NN}^2)_{Q^2=M_\sigma^2}} \right]}. \quad (85)$$

By using the average value $\Lambda_{\sigma NN} = 3.97 \text{ fm}^{-1}$, Eq. (47), we get

$$\begin{aligned} \Lambda_{\sigma NN} &= 3.97 \text{ fm}^{-1} \\ \Lambda_{\sigma\Lambda\Lambda} &= 4.48 \text{ fm}^{-1} \\ \Lambda_{\sigma\Sigma\Sigma} &= 4.75 \text{ fm}^{-1} \\ \Lambda_{\sigma\Xi\Xi} &= 3.45 \text{ fm}^{-1}. \end{aligned} \quad (86)$$

Thus we could use $\Lambda_{\sigma B_8 B_8} \simeq 4.0 \text{ fm}^{-1}$ as an average value for the whole baryon octet.

C. $a_0 B_8 B_8$

The results obtained for σ can be extrapolated to the a_0 meson in a straightforward way. Let us recall that in the additive constituent quark pattern one has degenerate masses $M_{a_0} = M_\sigma$ (see Ref. [11] for an explanation of the nondegeneracy). Then by writing the a_0 on-shell couplings in SU(3) language

$$\begin{aligned} (g_{a_0 NN})_{Q^2=M_{a_0}^2} &= g_{s,+} \\ (g_{a_0 \Sigma\Sigma})_{Q^2=M_{a_0}^2} &= 2g_{s,+} + \alpha_s \\ (g_{a_0 \Xi\Xi})_{Q^2=M_{a_0}^2} &= g_{s,+} + (2\alpha_s - 1), \end{aligned} \quad (87)$$

and imposing the degeneracy we predict from our value for $g_{s,+}$ [Eq. (80)]

$$\frac{(g_{a_0 NN}^2)_{Q^2=M_{a_0}^2}}{4\pi} = \frac{(g_{a_0 \Xi\Xi}^2)_{Q^2=M_{a_0}^2}}{4\pi} = 7.6 \quad (88)$$

$$\frac{(g_{a_0 \Sigma\Sigma}^2)_{Q^2=M_{a_0}^2}}{4\pi} = 30.3.$$

D. $\pi B_8 B_8$

To avoid mass factors we use the pseudovector coupling for the pion, i.e., the Lagrangian

$$\tilde{\mathcal{L}}_{\pi qq}^{PV} = -\frac{\tilde{f}_{\pi qq}}{M_\pi} \bar{q} i \gamma_5 \gamma_\mu \vec{\tau} \partial^\mu \vec{\pi} q \quad (89)$$

with the vertex form factor

$$\tilde{F}_{\pi qq}(Q^2) \equiv \left(\frac{\tilde{\Lambda}^2 - M_\pi^2}{\tilde{\Lambda}^2 - Q^2} \right)^{1/2} \quad (90)$$

to derive the asymptotic pion exchange central potentials. To perform the calculation we choose a total (spin, isospin) in each case. By considering the 1S_0 partial wave, for instance, we take (0, 1) for NN and $\Xi\Xi$ and (0, 0) for $\Sigma\Sigma$ interactions. Then we have

$$\begin{aligned} (V_q)_{\text{OPE}}^{NN \rightarrow NN}(R \rightarrow \infty) &= \frac{\tilde{f}_{\pi qq}^2}{4\pi} \left[-\frac{75}{9} e^{\frac{M_\pi^2 R^2}{2}} \right] \frac{1}{3} \frac{e^{-M_\pi R}}{R} \\ (V_q)_{\text{OPE}}^{\Sigma\Sigma \rightarrow \Sigma\Sigma}(R \rightarrow \infty) &= \frac{\tilde{f}_{\pi qq}^2}{4\pi} \left[\frac{32}{3} e^{\frac{M_\pi^2 R^2}{2}} \right] \frac{1}{3} \frac{e^{-M_\pi R}}{R} \quad (91) \\ (V_q)_{\text{OPE}}^{\Xi\Xi \rightarrow \Xi\Xi}(R \rightarrow \infty) &= \frac{\tilde{f}_{\pi qq}^2}{4\pi} \left[-\frac{3}{9} e^{\frac{M_\pi^2 R^2}{2}} \right] \frac{1}{3} \frac{e^{-M_\pi R}}{R}. \end{aligned}$$

However, at the baryonic level the Lagrangians can be expressed as

$$\mathcal{L}_{\pi NN}^{PV} = -\frac{(f_{\pi NN})_{Q^2=M_\pi^2}}{M_\pi} \bar{N} i \gamma_5 \gamma_\mu \vec{\tau} \partial^\mu \vec{\pi} N \quad (92)$$

$$\mathcal{L}_{\pi\Sigma\Sigma}^{PV} = -\frac{(f_{\pi\Sigma\Sigma})_{Q^2=M_\pi^2}}{M_\pi} (\vec{\Sigma} \times \vec{\Sigma}) \gamma_5 \gamma_\mu \partial^\mu \vec{\pi} \quad (93)$$

$$\mathcal{L}_{\pi\Xi\Xi}^{PV} = -\frac{(f_{\pi\Xi\Xi})_{Q^2=M_\pi^2}}{M_\pi} \bar{\Xi} i \gamma_5 \gamma_\mu \vec{\tau} \partial^\mu \vec{\pi} \Xi, \quad (94)$$

with form factors

$$\begin{aligned} G_{\pi NN}(Q^2) &= \left(\frac{\Lambda_{\pi NN}^2 - M_\pi^2}{\Lambda_{\pi NN}^2 - Q^2} \right)^{\frac{1}{2}} \\ G_{\pi\Sigma\Sigma}(Q^2) &= \left(\frac{\Lambda_{\pi\Sigma\Sigma}^2 - M_\pi^2}{\Lambda_{\pi\Sigma\Sigma}^2 - Q^2} \right)^{\frac{1}{2}} \quad (95) \end{aligned}$$

$$G_{\pi\Xi\Xi}(Q^2) = \left(\frac{\Lambda_{\pi\Xi\Xi}^2 - M_\pi^2}{\Lambda_{\pi\Xi\Xi}^2 - Q^2} \right)^{\frac{1}{2}}, \quad (96)$$

and where in conventional SU(3) notation

$$\begin{aligned} (f_{\pi NN})_{Q^2=M_\pi^2} &= f_{p,+} \\ (f_{\pi\Sigma\Sigma})_{Q^2=M_\pi^2} &= 2f_{p,+} + \alpha_p \\ (f_{\pi\Xi\Xi})_{Q^2=M_\pi^2} &= -f_{p,+} + (1 - 2\alpha_p), \end{aligned}$$

having introduced $f_{p,+} \equiv (f_{p,D} + f_{p,F})$ as the sum of the symmetric and antisymmetric pseudoscalar octet coupling constants and $\alpha_p \equiv f_{p,F}/(f_{p,D} + f_{p,F})$ as the $F/(F+D)$ ratio of the pseudoscalar octet.

By using the same total (spin, isospin) channels as above the corresponding asymptotic pion exchange central potentials

are

$$\begin{aligned} (\mathcal{V}_B)_{\text{OPE}}^{NN \rightarrow NN}(R \rightarrow \infty) &= \frac{(f_{\pi NN}^2)_{Q^2=M_\pi^2}}{4\pi} [-3] \frac{1}{3} \frac{e^{-M_\pi R}}{R} \\ (\mathcal{V}_B)_{\text{OPE}}^{\Sigma\Sigma \rightarrow \Sigma\Sigma}(R \rightarrow \infty) &= \frac{(f_{\pi\Sigma\Sigma}^2)_{Q^2=M_\pi^2}}{4\pi} [6] \frac{1}{3} \frac{e^{-M_\pi R}}{R} \quad (97) \\ (\mathcal{V}_B)_{\text{OPE}}^{\Xi\Xi \rightarrow \Xi\Xi}(R \rightarrow \infty) &= \frac{(f_{\pi\Xi\Xi}^2)_{Q^2=M_\pi^2}}{4\pi} [-3] \frac{1}{3} \frac{e^{-M_\pi R}}{R}. \end{aligned}$$

Then from the comparison of the asymptotic potentials at the baryon level and quark level the following relations come out

$$\begin{aligned} (f_{\pi NN}^2)_{Q^2=M_\pi^2} &= \tilde{f}_{\pi qq}^2 \frac{25}{9} e^{\frac{M_\pi^2 R^2}{2}} \\ (f_{\pi\Sigma\Sigma}^2)_{Q^2=M_\pi^2} &= \tilde{f}_{\pi qq}^2 \frac{16}{9} e^{\frac{M_\pi^2 R^2}{2}} \quad (98) \\ (f_{\pi\Xi\Xi}^2)_{Q^2=M_\pi^2} &= \tilde{f}_{\pi qq}^2 \frac{1}{9} e^{\frac{M_\pi^2 R^2}{2}}, \end{aligned}$$

and

$$\begin{aligned} \frac{(f_{\pi\Sigma\Sigma}^2)_{Q^2=M_\pi^2}}{(f_{\pi NN}^2)_{Q^2=M_\pi^2}} &= \frac{16}{25} \\ \frac{(f_{\pi\Xi\Xi}^2)_{Q^2=M_\pi^2}}{(f_{\pi NN}^2)_{Q^2=M_\pi^2}} &= \frac{1}{25}. \quad (99) \end{aligned}$$

1. Coupling constants and form factors values

From Eqs. (99) and the standard value $(f_{\pi NN}^2)_{Q^2=M_\pi^2}/4\pi = 0.079$ we predict the numerical values

$$\begin{aligned} \frac{(f_{\pi\Sigma\Sigma}^2)_{Q^2=M_\pi^2}}{4\pi} &= 0.051 \\ \frac{(f_{\pi\Xi\Xi}^2)_{Q^2=M_\pi^2}}{4\pi} &= 0.0032, \quad (100) \end{aligned}$$

or

$$\begin{aligned} \frac{(g_{\pi\Sigma\Sigma}^2)_{Q^2=M_\pi^2}}{4\pi} &= 9.4 \\ \frac{(g_{\pi\Xi\Xi}^2)_{Q^2=M_\pi^2}}{4\pi} &= 0.6, \quad (101) \end{aligned}$$

and

$$\alpha_p = 0.4. \quad (102)$$

This compares quite well with a derived value of $\alpha_p \simeq 0.365 \pm 0.007$ from the F/D ratio extracted from semileptonic decays of baryons [20].

With respect to the couplings at $Q^2 = 0$ we can rely on the quite approximate Goldstone boson character of the pion and assume they are given by $(f_{\pi B_8 B_8}^2)_{Q^2=0} \simeq (f_{\pi B_8 B_8}^2)_{Q^2=M_\pi^2 \rightarrow 0}$. As in this limit the ratios between the couplings are the same than the on-shell ones obtained above, Eq. (99), we can use them altogether with $(f_{\pi NN}^2)_{Q^2=M_\pi^2 \rightarrow 0}/4\pi = 0.072$, Eq. (35),

TABLE II. Pion and sigma coupling constants to quarks in our SU(2) \otimes SU(2) and SU(3) \otimes SU(3) models.

m	SU(2) \otimes SU(2)	SU(3) \otimes SU(3)
(π, σ)	$g_{\text{ch}}^2/4\pi = 0.535$	$\tilde{g}_{\text{ch}}^2/4\pi = 0.535$
π	$g_{\pi qq}^2/4\pi = 0.55$	$\tilde{g}_{\pi qq}^2/4\pi = 0.54$
σ	$g_{\sigma qq}^2/4\pi = 1.59$	$\tilde{g}_{\sigma qq}^2/4\pi = 0.93$

to predict

$$\frac{(f_{\pi\Sigma\Sigma}^2)_{Q^2=0}}{4\pi} \simeq 0.046 \quad (103)$$

$$\frac{(f_{\pi\Xi\Xi}^2)_{Q^2=0}}{4\pi} \simeq 0.0029,$$

or

$$\frac{(g_{\pi\Sigma\Sigma}^2)_{Q^2=0}}{4\pi} \simeq 8.5 \quad (104)$$

$$\frac{(g_{\pi\Xi\Xi}^2)_{Q^2=0}}{4\pi} \simeq 0.5.$$

Moreover the preservation of the ratios implies the equality of the form factors. From $(f_{\pi NN}^2)_{Q^2=0} \simeq (f_{\pi NN}^2)_{Q^2=M_\pi^2 \rightarrow 0}$ and $(f_{\pi NN}^2)_{Q^2=M_\pi^2}$ we deduce

$$\Lambda_{\pi B_8 B_8} \simeq 2.35 \text{ fm}^{-1}. \quad (105)$$

IV. SUMMARY

From a SU(2) \otimes SU(2) chiral quark Lagrangian incorporating spontaneous chiral symmetry breaking, and its generalization to SU(3) \otimes SU(3), asymptotic meson exchange $B_8 B_8 \rightarrow B_8 B_8$ interaction potentials are derived in the Born-Oppenheimer approximation. The comparison with the corresponding potentials from a SU(2) or SU(3) invariant hadronic Lagrangian allows for the expression of the $\pi B_8 B_8$ and $\sigma B_8 B_8$ coupling constants in terms of the elementary πqq and σqq ones. By using the πNN coupling constant as an input the πqq one gets fixed. From it the rest of coupling constants (σqq , $\pi B_8 B_8$, and $\sigma B_8 B_8$) are predicted. Their $M_m \rightarrow 0$ limits are also of interest to be compared with the values of the couplings at $Q^2 = 0$ provided by phenomenological analyses or QCDSR. The similar value obtained for πNN indicates the quite approximate Goldstone boson nature of the pion. On the contrary, $\sigma B_8 B_8$ couplings are significantly different as might be expected.

TABLE III. Predicted pairs $[(g_{m B_8 B_8}^2/4\pi)_{Q^2=M_m^2}, \Lambda_{m B_8 B_8}(\text{fm}^{-1})]$ from the chiral quark potential model, for exact SU(3) symmetry ($M_\Lambda = M_\Sigma = M_\Xi = M_N = 939$ MeV). The superindex * indicates the πNN coupling constant value used as input.

m	$m NN$	$m \Lambda \Lambda$	$m \Sigma \Sigma$	$m \Xi \Xi$
π	(14.6*, 2.35)		(9.4, 2.35)	(0.6, 2.35)
σ	(68.2, 3.97)	(30.3, 4.48)	(30.3, 4.75)	(7.6, 3.45)

Further information about the couplings can be extracted under the assumption that our on-shell model predictions and the $Q^2 = 0$ values from external analyses can be managed jointly. Though this assumption is debatable it allows to get some insight into the cutoff parameters $\Lambda_{m B_8 B_8}$ at the baryonic level.

We summarize in Table II the pion and sigma coupling constants to quarks in our models. In Table III the values obtained for the coupling constants and the form factors parameters at the baryon level are listed. By making use of the a_0 - σ degeneracy in our quark model we have also predicted $a_0 B_8 B_8$ on-shell couplings. Concerning other diagonal $m B B$ couplings such as $f_0 B_8 B_8$, $\eta B_8 B_8$, and $\eta' B_8 B_8$, to which our formalism could be also applied, the situation gets complicated by the presence of the strange quark and/or antiquark that may give rise to relevant SU(3) breaking effects out of the scope of our symmetry treatment.

Let us finally add that in our model Goldberger-Treiman relations of the form $(g_A)_{\pi B_8 B_8}/2f_\pi = (f_{\pi B_8 B_8})_{Q^2=M_\pi^2}/M_\pi$, where g_A stands for the axial coupling constant and f_π for the pion decay constant, can be immediately applied. In our nonrelativistic description $(g_A)_{\pi NN} = 5/3$, a value considerably larger than the experimental one, 1.267. Consequently $f_\pi = 116$ MeV, which is 20% bigger than the experimental value of 93 MeV. Regarding these discrepancies it has been shown [18] that a relativistic treatment, beyond our present approach, could correct them to a good extent.

ACKNOWLEDGMENTS

This work was partially funded by Ministerio de Educación y Ciencia and EU FEDER under Contract No. FPA2007-65748, by Junta de Castilla y León under Contract No. SA016A17, and by the Spanish Consolider-Ingenio 2010 Program CPAN (CSD2007-00042).

- [1] T. E. O. Ericson, Nucl. Phys. **A543**, 409c (1992); D. V. Bugg, Eur. Phys. J. C **33**, 505 (2004).
- [2] V. G. J. Stocks and Th. A. Rijken, Nucl. Phys. **A613**, 311 (1997).
- [3] A. Valcarce, H. Garcilazo, F. Fernández, and P. González, Rep. Prog. Phys. **68**, 965 (2005) and references therein.
- [4] Z. Y. Zhang *et al.*, Nucl. Phys. **A625**, 59 (1997).
- [5] K. Shimizu, S. Takeuchi, and A. Buchmann, Prog. Theor. Phys. Suppl. **137**, 43 (2000).

- [6] Y. Fujiwara, Y. Suzuki, and C. Nakamoto, Prog. Part. Nucl. Phys. **58**, 439 (2007).
- [7] G. Q. Liu, M. Swift, and A. W. Thomas, Nucl. Phys. **A556**, 331 (1993).
- [8] D. I. Diakonov and V. Yu. Petrov, Nucl. Phys. **B272**, 457 (1986).
- [9] F. Fernández and E. Oset, Nucl. Phys. **A455**, 720 (1986).
- [10] A. M. Kusainov, V. G. Neudatchin, and I. T. Obukhovskiy, Phys. Rev. C **44**, 2343 (1991).
- [11] M. D. Scadron, Phys. Rev. D **26**, 239 (1982).

- [12] W. Y. Hwang, Z. Phys. C **75**, 701 (1997).
[13] E. M. Henley and G. A. Miller, Phys. Lett. **B251**, 453 (1991).
[14] A. W. Thomas and K. Holinde, Phys. Rev. Lett. **63**, 2025 (1989).
[15] G. Erkol, R. G. E. Timmermans, and Th. A. Rijken, Phys. Rev. C **72**, 035209 (2005).
[16] J. Carlson, J. Kogut, and V. R. Pandharipande, Phys. Rev. D **27**, 233 (1983); J. L. Basdevant and S. Boukaraa, Z. Phys. C **30**, 103 (1986); F. Cano, P. González, S. Noguera, and B. Desplanques, Nucl. Phys. **A603**, 257 (1996); H. Garcilazo and A. Valcarce, Phys. Rev. C **68**, 035207 (2003).
[17] B. Juliá-Díaz, A. Valcarce, P. González, and F. Fernández, Phys. Rev. C **66**, 024005 (2002).
[18] B. Juliá-Díaz, D. O. Riska, and F. Coester, Phys. Rev. C **70**, 045204 (2004).
[19] G. Erkol, R. G. E. Timmermans, M. Oka, and Th. A. Rijken, Phys. Rev. C **73**, 044009 (2006).
[20] F. E. Close and R. G. Roberts, Phys. Lett. **B316**, 165 (1993).