

Equation of state for β -stable hot nuclear matter

Ch. C. Moustakidis and C. P. Panos

Department of Theoretical Physics, Aristotle University of Thessaloniki, 54124 Thessaloniki, Greece

(Received 20 October 2008; revised manuscript received 19 February 2009; published 20 April 2009)

We provide an equation of state for hot nuclear matter in β equilibrium by applying a momentum-dependent effective interaction. We focus on the study of the equation of state of high-density and high-temperature nuclear matter, containing leptons (electrons and muons) under the chemical equilibrium condition in which neutrinos have left the system. The conditions of charge neutrality and equilibrium under the β -decay process lead first to the evaluation of proton and lepton fractions and then to the evaluation of internal energy, free energy, and pressure, and in total to the equation of state of hot nuclear matter. Thermal effects on the properties and equation of state of nuclear matter are assessed and analyzed in the framework of the proposed effective interaction model. Special attention is given to the study of the contribution of the components of β -stable nuclear matter to the entropy per particle, a quantity of great interest in the study of structure and collapse of supernova.

DOI: [10.1103/PhysRevC.79.045806](https://doi.org/10.1103/PhysRevC.79.045806)

PACS number(s): 26.60.-c, 21.65.-f, 21.30.Fe, 24.10.Pa

I. INTRODUCTION

The equation of state (EOS) of hot nuclear matter determines the structure inside a supernova [1] and a hot neutron star [2–4] and affects the state of matter, such as its chemical composition. In addition, the equation of state plays an important role in the study of the supernova explosion, as well as in determining the evolution of a neutron star at the birth stage. The profiles of a neutron star as the density, temperature, and proton fraction during the cooling, which affect the reaction rate of neutrino process inside the star, are determined through the equation of state.

There exist many calculations for hot nuclear matter with applications to the properties of hot neutron stars and supernova [1–40]. Baym *et al.* provided an EOS of neutron matter [5] and Bethe *et al.* [6] an EOS for the gravitational collapse of stars. Friedman and Pandharipande [7] performed variational calculations of the EOS of hot and cold nuclear and neutron matter. Lattimer and Swesty carried out calculations of the EOS for stellar collapse, using the compressible liquid-drop model for nuclei [8]. M. Prakash *et al.* [2] investigated the structure of neutron stars shortly after their birth, by applying various nuclear models. Takatsuka *et al.* [9,10] have performed detailed calculations for supernova matter, within the framework of the finite-temperature Hartree-Fock approach, with effective nucleon-nucleon interaction. Recently, Das *et al.* [11] calculated the EOS of dense supernova matter within the finite-temperature Brueckner-Goldstone approach with effective two-body Sussex interaction.

The present work is based on the previous work of Prakash *et al.* [2]. More specifically, to study the properties and EOS of hot nuclear matter, we apply a momentum-dependent effective interaction model (MDIM), which is able to reproduce the results of more microscopic calculations of dense matter at zero temperature and can be extended to finite temperature [2,39,40].

The aim of this work is to apply a momentum-dependent interaction model for the study of the hot nuclear matter EOS under β equilibrium. The present model has the additional property, compared to the previous ones, that the temperature

not only affects the kinetic part of the energy density but also influences the interaction part of the energy density. In that way, we are able to simultaneously study thermal effects not only on the kinetic part of the symmetry energy and symmetry free energy but also on the interaction part of the above quantities. This is important in the sense that the density-dependent behaviors of the symmetry energy and symmetry free energy influence strongly the values of the proton fraction and, as a consequence, the composition of the hot β -stable nuclear matter under consideration.

Using this method, we will show that the thermal energy (and also the related quantities) depends sensitively on the momentum dependence of the nuclear interaction. We concentrate our study on the properties of hot nuclear matter in the density range $n_0 < n < 6n_0$ (where $n_0 = 0.16 \text{ fm}^{-3}$ is the saturation density) and temperature range $0 < T < 30 \text{ MeV}$, taking into account that nuclear matter consists of neutrons, protons, electrons, and muons with their relative concentrations determined from the conditions of charge neutrality and equilibrium under the β -decay process in the absence of neutrino trapping.

The article is organized as follows. In Sec. II, the model and relative formulas are discussed and analyzed. Results are reported and discussed in Sec. III, whereas the summary of the work is given in Sec. IV.

II. THE MODEL

We start by outlining the momentum-dependent interaction model, then we define the thermodynamic quantities of nuclear matter, and finally we analyze the β -equilibrium conditions, the contribution of leptons to pressure and energy, and the total equation of state of nuclear matter.

A. Momentum-dependent interaction model

The schematic potential model, employed here, is designed to reproduce the results of the microscopic calculations of both

nuclear and neutron-rich matter at zero temperature and can be extended to finite temperature [2]. The energy density of the asymmetric nuclear matter (ANM) is given by the relation

$$\epsilon(n_n, n_p, T) = \epsilon_{\text{kin}}^n(n_n, T) + \epsilon_{\text{kin}}^p(n_p, T) + V_{\text{int}}(n_n, n_p, T), \quad (1)$$

where n_n (n_p) is the neutron (proton) density and the total baryon density is $n = n_n + n_p$. The contributions of the kinetic parts are

$$\begin{aligned} & \epsilon_{\text{kin}}^n(n_n, T) + \epsilon_{\text{kin}}^p(n_p, T) \\ &= 2 \int \frac{d^3k}{(2\pi)^3} \frac{\hbar^2 k^2}{2m} (f_n(n_n, k, T) + f_p(n_p, k, T)), \end{aligned} \quad (2)$$

where f_τ (for $\tau = n, p$) is the Fermi-Dirac distribution function with the form

$$f_\tau(n_\tau, k, T) = \left[1 + \exp\left(\frac{e_\tau(n_\tau, k, T) - \mu_\tau(n_\tau, T)}{T}\right) \right]^{-1}. \quad (3)$$

The nucleon density n_τ is evaluated from the following integral

$$\begin{aligned} n_\tau &= 2 \int \frac{d^3k}{(2\pi)^3} f_\tau(n_\tau, k, T) \\ &= 2 \int \frac{d^3k}{(2\pi)^3} \left[1 + \exp\left(\frac{e_\tau(n_\tau, k, T) - \mu_\tau(n_\tau, T)}{T}\right) \right]^{-1}. \end{aligned} \quad (4)$$

In Eq. (3), $e_\tau(n_\tau, k, T)$ is the single-particle energy (SPE) and $\mu_\tau(n_\tau, T)$ stands for the chemical potential of each species. The SPE has the form

$$e_\tau(n_\tau, k, T) = \frac{\hbar^2 k^2}{2m} + U_\tau(n_\tau, k, T), \quad (5)$$

where the single-particle potential $U_\tau(n_\tau, k, T)$ is obtained by the functional derivative of the interaction part of the energy density with respect to the distribution function f_τ . Including the effect of finite-range forces between nucleons, to avoid acausal behavior at high densities, the potential contribution is

parametrized as follows [2]:

$$\begin{aligned} V_{\text{int}}(n_n, n_p, T) &= \frac{1}{3} A n_0 \left[\frac{3}{2} - \left(\frac{1}{2} + x_0 \right) I^2 \right] u^2 \\ &+ \frac{\frac{2}{3} B n_0 \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma+1}}{1 + \frac{2}{3} B' \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma-1}} \\ &+ u \sum_{i=1,2} \left[C_i (\mathcal{J}_n + \mathcal{J}_p) \right. \\ &\left. + \frac{(C_i - 8Z_i)}{5} I (\mathcal{J}_n - \mathcal{J}_p) \right], \end{aligned} \quad (6)$$

where

$$\mathcal{J}_\tau = 2 \int \frac{d^3k}{(2\pi)^3} g(k, \Lambda_i) f_\tau(n_\tau, k, T). \quad (7)$$

In Eq. (6), I is the asymmetry parameter [$I = (n_n - n_p)/n$] and $u = n/n_0$, with n_0 denoting the equilibrium symmetric nuclear matter density, $n_0 = 0.16 \text{ fm}^{-3}$. The asymmetry parameter I is related to the proton fraction Y_p by the equation $I = (1 - 2Y_p)$. The parameters A, B, σ, C_1, C_2 , and B' , which appear in the description of symmetric nuclear matter, are determined so that $E(n = n_0) - mc^2 = -16 \text{ MeV}$, $n_0 = 0.16 \text{ fm}^{-3}$, and the incompressibility $K = 240 \text{ MeV}$. The additional parameters x_0, x_3, Z_1 , and Z_2 , which are used to determine the properties of asymmetric nuclear matter, are treated as parameters constrained by empirical knowledge [2]. The parametrizations used in the present model have only a modest microscopic foundation. Nonetheless, they have the merit of being able to closely approximate more physically motivated calculations as presented in Fig. 1. More precisely, in Fig. 1, we compare the energy per baryon [for symmetric nuclear matter, Fig. 1(a), and pure neutron matter, Fig. 1(b)] calculated by the present schematic model (MDIM) with those of existing state-of-the-art calculations by Wiringa *et al.* [41] and Pandharipande *et al.* [42].

The first two terms of the right-hand side of Eq. (6) arise from the local contact nuclear interaction and lead to power density contributions, as in the standard Skyrme




FIG. 1. Energy per baryon of symmetric (a) and pure neutron matter (b) of the present model (MDIM) in comparison with those originated from realistic calculations. More details for the models UV14+TNI, UV14+UVII and AV14+UVII in Ref. [41] and for the models A18+UIX and A18+du+UIX* in Ref. [42].




FIG. 2. (a) Contribution of the various terms V_A , V_B , V_C and the total potential energy density V_{int} as a function of the baryon density. (b) Momentum-dependent term V_C as a function of the baryon density at temperature $T = 0$, $T = 10$, and $T = 30$ MeV.

equation of state. The first one concerns attractive interaction, while the second one is repulsive, and both are assumed to be temperature independent. The third term describes the effects of finite-range interactions according to the chosen function $g(k, \Lambda_i)$, and is the temperature-dependent part of the interaction. This interaction is attractive and important at low momentum, but it weakens and disappears at very high momentum. The function $g(k, \Lambda_i)$, suitably chosen to simulate finite-range effects, has the form [2]

$$g(k, \Lambda) = \left[1 + \left(\frac{k}{\Lambda_i} \right)^2 \right]^{-1}, \quad (8)$$

where the finite-range parameters are $\Lambda_1 = 1.5k_F^0$ and $\Lambda_2 = 3k_F^0$, and k_F^0 is the Fermi momentum at the saturation point n_0 .

The main origin of the momentum dependence in Brueckner theory is the nonlocality of the exchange interaction. Following the discussion of Bertsch *et al.* [43], a single-particle potential $U(n)$ which depends only on the baryon density is oversimplified. Furthermore, it is well known that nuclear interaction has strong exchange effects that give rise to a momentum dependence in the single-particle potential, and, as a consequence, it has an effect on the energy density functional. The question here is how best to parametrize momentum dependence in modeling the potential $U(n, k)$. A promising approach might be to adopt the relativistic mean field model, where $U(n, k) = U_v n + \frac{U_s n}{\sqrt{1+k^2/m^2}}$. This potential exhibits a strong momentum dependence for small k which diminishes to zero at high momentum. To perform extensive studies in heavy ion collision studies, Gale *et al.* [44] have proposed the following parametrization for the momentum part of the single particle:

$$U(n, k) \sim C \frac{n}{n_0} \frac{1}{1 + (\mathbf{k} - \langle \mathbf{k}' \rangle)^2 / \Lambda^2}.$$

This has a proper falloff at high k and Galilean invariance is ensured by measuring k with respect to the average of the particles in the neighborhood, $\langle \mathbf{k}' \rangle$. For static nuclear matter, we have $\langle \mathbf{k}' \rangle = 0$.

The present model, which is a generalization of that proposed by Gale *et al.* [44], has been successfully applied in heavy ion collisions and astrophysical studies over the years [2–4, 27, 28, 33, 45, 46].

To clarify the relative contribution of the three terms of the potential energy density mentioned above, we plot them as a function of the baryon density in Fig. 2(a). In this figure, we have that

$$\begin{aligned} V_A &= \frac{1}{3} A n_0 \left[\frac{3}{2} - \left(\frac{1}{2} + x_0 \right) I^2 \right] u^2, \\ V_B &= \frac{\frac{2}{3} B n_0 \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma+1}}{1 + \frac{2}{3} B' \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma-1}}, \\ V_C &= u \sum_{i=1,2} \left[C_i (\mathcal{J}_n^i + \mathcal{J}_p^i) + \frac{(C_i - 8Z_i)}{5} I (\mathcal{J}_n^i - \mathcal{J}_p^i) \right]. \end{aligned} \quad (9)$$

As mentioned above, V_A corresponds to an attractive interaction, whereas V_B corresponds to repulsive interaction and dominates for high values of n ($n > 0.6 \text{ fm}^{-3}$). Both of these terms are temperature independent. The third term, V_C (plotted for $T = 0$), contains the momentum-dependent part of the interaction and corresponds to attractive interaction; its main contribution is to compete with the repulsive interaction of V_B for high values of n and as a consequence avoid acausal behavior of the EOS at high densities. V_C consists of two finite-range terms, one corresponding to a long-range attraction and the other to a short-range repulsion.

Thermal effects on the momentum-dependent term V_C are displayed in Fig. 2(b). The contribution of V_C is plotted for various values of T . It is therefore concluded that thermal effects are more pronounced for high values of T ($T > 10$ MeV), leading to a less attractive contribution. More precisely, we find that for small values of n (i.e., $n = 0.15 \text{ fm}^{-3}$) V_C increases (compared to the cold case $T = 0$) by 3–20% for $T = 10$ –30. For higher values of n , the increase is even less.

An additional test for the present model is to compare the single-particle potential $U_\tau(n_\tau, k, T)$ [or $U_\tau(n, I, k, T)$] originated from the present version of the momentum-dependent

interaction with other calculations. The single-particle potential $U_\tau(n, I, k, T)$ (protons or neutrons), obtained from the functional derivative of the interaction part of the energy density [Eq. (6)] with respect to the distribution function f_τ , has the general form [40]

$$U_\tau(n, I, k, T) = U_\tau^A(n, I) + U_\tau^B(n, I) + U_\tau^{\text{MD}}(n, I, k, T). \quad (10)$$

It is of interest to see that the single-particle potentials are separated into two parts. The first one, $U_\tau^A(n, I) + U_\tau^B(n, I)$ is a function only of the baryon density n and the isospin asymmetry parameter I . The second one, $U_\tau^{\text{MD}}(n, I, k, T)$ has an additional dependence on T and k . Actually, $U_\tau^{\text{MD}}(n, I, k, T)$ is mainly responsible for the trend of the effective mass and also the effective mass splitting. Additionally, it is connected with the effect of the temperature on the interacting part of the energy density [40].

The single-particle potential in symmetric nuclear matter has been calculated microscopically for several Hamiltonians by Wiringa [47]. These Hamiltonians include nucleon-nucleon potentials fit to scattering data and three-nucleon potentials fit to binding energies of few-body nuclei and saturation properties of nuclear matter. The potential was parametrized using the ansatz

$$U(n, k) = \alpha(n) + \frac{\beta(n)}{1 + \left(\frac{k}{\Lambda(n)}\right)^2}, \quad (11)$$

where the density-dependent parameters $\alpha(n)$, $\beta(n)$, and $\Lambda(n)$, for three types of Hamiltonian, are listed in Table I of Ref. [47].

Furthermore, the single-particle potential has been derived by Li *et al.* [48]. The derivation is based upon the Bonn meson-exchange model for the nucleon-nucleon interaction and the Dirac-Brueckner approach for nuclear matter. The potential, named DBHF, has been parametrized as

$$U(n, k) = \alpha n + \beta n^\gamma + \delta \ln^2[\epsilon(\hbar ck)^2 + 1]n^\sigma. \quad (12)$$

The parameters α , β , γ , δ , ϵ , and σ are listed in Table I of Ref. [48].

A comparison with the results of UV14+TNI, UV14+UVII, AV14+UVII, and DBHF interactions (see Fig. 3) show that $U(n, k)$ (for $T = 0$) obtained from the present model is very reasonable for at least up to $k = 3 \text{ fm}^{-1}$. The agreement is not as good for high values of k , especially

compared with the DBHF interaction; but as has been pointed out by Li *et al.* [48], the parametrization in Eq. (12) is bad for large k , since it continues to grow with increasing k , whereas the exact potential becomes independent of k for large momenta. In conclusion, the present results show that the momentum-dependent interaction model, which has been applied in the present work, provides a reliable representation of $U(n, k)$ for a wide range of density and momentum.

The energy density of asymmetric nuclear matter at density n and temperature T , in a good approximation, is expressed as

$$\epsilon(n, T, I) = \epsilon(n, T, I = 0) + \epsilon_{\text{sym}}(n, T, I), \quad (13)$$

where

$$\begin{aligned} \epsilon_{\text{sym}}(n, T, I) &= nI^2 E_{\text{sym}}^{\text{tot}}(n, T) \\ &= nI^2 (E_{\text{sym}}^{\text{kin}}(n, T) + E_{\text{sym}}^{\text{int}}(n, T)). \end{aligned} \quad (14)$$

In Eq. (14), the nuclear symmetry energy $E_{\text{sym}}^{\text{tot}}(n, T)$ is separated in two parts corresponding to the kinetic contribution $E_{\text{sym}}^{\text{kin}}(n, T)$ and the interaction one $E_{\text{sym}}^{\text{int}}(n, T)$.

From Eqs. (13) and (14) and setting $I = 1$, we find that the nuclear symmetry energy $E_{\text{sym}}^{\text{tot}}(n, T)$ is given by

$$E_{\text{sym}}^{\text{tot}}(n, T) = \frac{1}{n} [\epsilon(n, T, I = 1) - \epsilon(n, T, I = 0)]. \quad (15)$$

Thus, from Eq. (15) and by a suitable choice of the parameters x_0 , x_3 , Z_1 , and Z_2 , we can obtain various forms for the density dependence of the symmetry energy $E_{\text{sym}}^{\text{tot}}(n, T)$.

It is well known that the need to explore different forms for $E_{\text{sym}}^{\text{tot}}(n, T)$ stems from the uncertain behavior at high density [2]. The high-density behavior of symmetry energy is the least known property of dense matter [49–51], with different nuclear models giving contradictory predictions. Thus, in relativistic mean field (RMF) models, the symmetry energy increases strongly with the density of nuclear matter [52], whereas in many realistic potential models of nuclear matter in the variational approach [7,53], the symmetry energy saturates and then bends over at higher densities.

Recently, the density dependence of the symmetry energy in the EOS of isospin asymmetric nuclear matter has been studied using isoscaling of the fragment yields and the antisymmetrized molecular dynamic calculation [54]. It was observed that the experimental data at low densities are consistent with the form of symmetry energy, $E_{\text{sym}}(u) \approx 31.6u^{0.69}$, in close




FIG. 3. Comparison of the single-particle potential of symmetric nuclear matter from the present model (MDIM) with the microscopic calculations of Wiringa [47] and Li *et al.* [48], for densities $n = 0.1$, $n = 0.3$, and $n = 0.5 \text{ fm}^{-3}$.

agreement with those predicted by the results of variational many-body calculations. In Ref. [54] it was suggested also that the heavy ion studies favor a dependence of the form $E_{\text{sym}}(u) \approx 31.6u^\gamma$, where $\gamma = 0.6$ – 1.05 . This constrains the form of the density dependence of the symmetry energy at higher densities, ruling out an extremely “stiff” and “soft” dependence [54].

Additionally, Chen *et al.* [55] also showed, using the isospin-dependent Boltzmann-Uehling-Uhlenback transport model calculations, that a stiff density dependence of the symmetry energy parametrized as $E_{\text{sym}}(u) \approx 31.6u^{1.05}$ explains well the isospin diffusion data [56] from NSCL-MSU (National Superconducting Cyclotron Laboratory at Michigan State University).

In this paper, since we are interested mainly in the study of thermal effects on the nuclear symmetry energy, we choose a specific form for it, enabling us to reproduce accurately the results of many other theoretical studies [57,58]. In Ref. [57], the authors carried out a systematic analysis of the nuclear symmetry energy in the formalism of the relativistic Dirac-Brueckner-Hartree-Fock approach, using the Bonn one-boson-exchange potential. In a very recent work [58], the authors applied a similar method as in Ref. [57] for the microscopic predictions of the equation of state in asymmetric nuclear matter. In that case, $E_{\text{sym}}(u)$ was obtained employing the simple parametrization $E_{\text{sym}}(u) = Cu^\gamma$ with $\gamma = 0.7$ – 1.0 and $C \approx 32$ MeV. The authors concluded that a value of γ close to 0.8 gives a reasonable description of their predictions, although the use of different functions in different density regions may be best for an optimal fit [58]. The results of Refs. [57,58] are well reproduced by parametrizing the nuclear symmetry energy according to the formula

$$E_{\text{sym}}^{\text{tot}}(n, T=0) = \underbrace{13u^{2/3}}_{\text{Kinetic}} + \underbrace{17F(u)}_{\text{Interaction}}. \quad (16)$$

For the function $F(u)$, which parametrizes the interaction part of the symmetry energy, we apply the form

$$F(u) = u. \quad (17)$$

The parameters x_0 , x_3 , Z_1 , and Z_2 are chosen so that Eq. (15), for $T=0$, reproduces the results of Eq. (16) for the function $F(u) = u$.

In one of our previous papers [59], the potential part of the symmetry energy are parametrized in the generalized form $F(u) = u^c$, and the obtained nuclear equations of state are applied to the systematic study of the global properties of a neutron star (masses, radii, and compositions). We obtained a linear relation between the parameter c and the radius and the maximum mass of the neutron star [59]. Additionally, we found that a linear relation between the radius and the derivative of the symmetry energy near the saturation density n_0 holds [59].

It is worthwhile to point out that the above parametrization of the interacting part of the nuclear symmetry energy is used extensively for the study of neutron star properties [2,60], as well as for the study of the collisions of neutron-rich heavy ions at intermediate energies [61,62]. For a very recent review of the applications of the proposed momentum-dependent effective

interaction model and its specific parametrizations, see Ref. [4] (and references therein).

B. Thermodynamic description of hot nuclear matter

To study the properties of nuclear matter at finite temperature, we need to introduce the Helmholtz free energy F . The differential of the total free energy F_{tot} (the total free energy of baryons contained in volume V) and total internal energy E_{tot} (the total internal energy of baryons contained in volume V) are given by [63,64]

$$dF_{\text{tot}} = -S_{\text{tot}}dT - PdV + \sum_i \mu_i dN_i, \quad (18)$$

$$dE_{\text{tot}} = TdS_{\text{tot}} - PdV + \sum_i \mu_i dN_i, \quad (19)$$

where S_{tot} is the total entropy of the baryons, while μ_i and N_i are the chemical potential and the number of particles of each species, respectively.

It is easy to prove that the free energy per particle F is written as [63,64]

$$F(n, T, I) = E(n, T, I) - TS(n, I, T). \quad (20)$$

In Eq. (20), E is the internal energy per particle, $E = \epsilon/n$, and S is the entropy per particle, $S = s/n$. From Eq. (20) is also concluded that for $T=0$, the free energy F and the internal energy E coincide.

The entropy density s has the same functional form as that of a noninteracting gas system, given by the equation

$$s_\tau(n, I, T) = -2 \int \frac{d^3k}{(2\pi)^3} [f_\tau \ln f_\tau + (1 - f_\tau) \ln(1 - f_\tau)]. \quad (21)$$

The total internal energy E_{tot} is useful for studying isentropic processes. In that description of a thermodynamic system, the pressure and the chemical potential are defined as [63,64]

$$P = - \left(\frac{\partial E_{\text{tot}}}{\partial V} \right)_{S, N_i} = n^2 \left(\frac{\partial \epsilon/n}{\partial n} \right)_{S, N_i}, \quad (22)$$

$$\mu_i = \left(\frac{\partial E_{\text{tot}}}{\partial N_i} \right)_{S, V, N_{j \neq i}} = \left(\frac{\partial \epsilon}{\partial n_i} \right)_{S, V, n_{j \neq i}}.$$

Now we are going to study the properties and the EOS of nuclear matter by considering an isothermal process. In that, the pressure and the chemical potential are connected with the derivative of the total free energy F_{tot} . More specifically, they are defined as

$$P = - \left(\frac{\partial F_{\text{tot}}}{\partial V} \right)_{T, N_i} = n^2 \left(\frac{\partial f/n}{\partial n} \right)_{T, N_i}, \quad (23)$$

$$\mu_i = \left(\frac{\partial F_{\text{tot}}}{\partial N_i} \right)_{T, V, N_{j \neq i}} = \left(\frac{\partial f}{\partial n_i} \right)_{T, V, n_{j \neq i}},$$

where f is the free energy density.

The pressure P can be calculated also from the equations [63,64]

$$VP = TS_{\text{tot}} - E_{\text{tot}} + \sum_i \mu_i N_i, \quad \text{or} \quad (24)$$

$$P = Ts - \epsilon + \sum_i \mu_i n_i.$$

It is also possible to calculate the entropy per particle $S(n, T)$ by differentiating the free energy density f with respect to the temperature

$$S(n, T) = - \left(\frac{\partial f/n}{\partial T} \right)_{V, N_i}. \quad (25)$$

The comparison of the two entropies, from Eqs. (21) and (25), provides a test of the approximation used in the present work.

It is easy to show by applying Eq. (23) that (for a proof, see Refs. [32,60])

$$\begin{aligned} \mu_n &= F + u \left(\frac{\partial F}{\partial u} \right)_{Y_p, T} - Y_p \left(\frac{\partial F}{\partial Y_p} \right)_{n, T}, \\ \mu_p &= \mu_n + \left(\frac{\partial F}{\partial Y_p} \right)_{n, T}, \\ \hat{\mu} &= \mu_n - \mu_p = - \left(\frac{\partial F}{\partial Y_p} \right)_{n, T}. \end{aligned} \quad (26)$$

We can define the symmetry free energy per particle $F_{\text{sym}}(n, T)$ by the following parabolic approximation (see also Refs. [32, 33]):

$$\begin{aligned} F(n, T, I) &= F(n, T, I = 0) + I^2 F_{\text{sym}}(n, T) \\ &= F(n, T, I = 0) + (1 - 2Y_p)^2 F_{\text{sym}}(n, T), \end{aligned} \quad (27)$$

where

$$F_{\text{sym}}(n, T) = F(n, T, I = 1) - F(n, T, I = 0). \quad (28)$$

It is worthwhile to notice that the above approximation is not valid from the beginning, but one needs to check the validity of the parabolic law in the present model before using it. As we see later, that law is well satisfied as well as the parabolic law holding for the energy.

Now, by applying Eq. (27) to Eq. (26), we obtain the key relation

$$\hat{\mu} = \mu_n - \mu_p = 4(1 - 2Y_p)F_{\text{sym}}(n, T). \quad (29)$$

This equation is similar to that obtained for cold nuclear matter by replacing $E_{\text{sym}}(n)$ with $F_{\text{sym}}(n, T)$.

C. β equilibrium, lepton contribution, and equation of state

Stable high-density nuclear matter must be in chemical equilibrium for all types of reactions, including the weak interactions, while β decay and electron capture take place simultaneously

$$n \longrightarrow p + e^- + \bar{\nu}_e, \quad p + e^- \longrightarrow n + \nu_e. \quad (30)$$

Both types of reactions change the electron per nucleon fraction Y_e and thus affect the equation of state. Here, we assume that neutrinos generated in those reactions have left

the system. The absence of neutrino trapping has a dramatic effect on the EOS and mainly induces a significant change on the values of the proton fraction Y_p [9,10]. The absence of neutrinos implies that

$$\hat{\mu} = \mu_n - \mu_p = \mu_e. \quad (31)$$

When the energy of electrons is large enough (i.e., greater than the muon mass), it is energetically favorable for the electrons to convert to muons:

$$e^- \longrightarrow \mu^- + \bar{\nu}_\mu + \nu_e. \quad (32)$$

Denoting the muon chemical potential by μ_μ , the chemical equilibrium established by the above process and its inverse is given by

$$\mu_e = \mu_\mu.$$

Taking into account that the threshold for muons occurs for $\mu_\mu = m_\mu c^2 \simeq 105.7$ MeV, one may expect muons to appear roughly at nuclear density $n = 0.16$ fm $^{-3}$.

Thus, in total, we consider that nuclear matter contains neutrons, protons, electrons, and muons. They are in a β equilibrium, where the following relations hold:

$$\mu_n = \mu_p + \mu_e, \quad \mu_e = \mu_\mu. \quad (33)$$

Furthermore, they obey the charge neutrality condition, i.e.,

$$n_p = n_e + n_\mu. \quad (34)$$

The lepton (electrons and muons) density is given by the expression

$$n_l = \frac{2}{(2\pi)^3} \int \frac{d\mathbf{k}}{1 + \exp \left[\frac{\sqrt{\hbar^2 k^2 c^2 + m_l^2 c^4} - \mu_l}{T} \right]}. \quad (35)$$

One can solve self-consistently Eqs. (29) and (33)–(35) to calculate the proton fraction Y_p , the lepton fractions Y_e and Y_μ , as well as the electron chemical potential μ_e as a function of the baryon density n , for various values of temperature T .

The next step is to calculate the energy and pressure of leptons given by the formulas

$$\epsilon_l(n_l, T) = \frac{2}{(2\pi)^3} \int \frac{\sqrt{\hbar^2 k^2 c^2 + m_l^2 c^4} d\mathbf{k}}{1 + \exp \left[\frac{\sqrt{\hbar^2 k^2 c^2 + m_l^2 c^4} - \mu_l}{T} \right]}, \quad (36)$$

$$\begin{aligned} P_l(n_l, T) &= \frac{1}{3} \frac{2(\hbar c)^2}{(2\pi)^3} \int \frac{1}{\sqrt{\hbar^2 k^2 c^2 + m_l^2 c^4}} \\ &\quad \times \frac{k^2 d\mathbf{k}}{1 + \exp \left[\frac{\sqrt{\hbar^2 k^2 c^2 + m_l^2 c^4} - \mu_l}{T} \right]}. \end{aligned} \quad (37)$$

The chemical potentials of electrons and muons are equal, and according to Eqs. (29) and (33) they are

$$\begin{aligned} \mu_e &= \mu_\mu = \mu_p - \mu_n = 4(1 - 2Y_p(n, T))F_{\text{sym}}(n, T) \\ &= 4I(n, T)F_{\text{sym}}(n, T). \end{aligned} \quad (38)$$

The EOS of hot nuclear matter in β equilibrium (considering that it consists of neutrons, protons, electrons, and muons) can




FIG. 4. Difference $F(n, T, Y_p) - F(n, T, Y_p = 1/2)$ as a function of $(1 - 2Y_p)^2$ at temperatures (a) $T = 10$ and (b) $T = 30$ MeV, for three baryon densities.

be obtained by calculating the total energy density ϵ_{tot} as well as the total pressure P_{tot} . The total energy density is given by

$$\epsilon_{\text{tot}}(n, T, I) = \epsilon_b(n, T, I) + \sum_{l=e, \mu} \epsilon_l(n, T, I), \quad (39)$$

where $\epsilon_b(n, T, I)$ and $\epsilon_l(n, T, I)$ are the contributions of baryons and leptons, respectively. The total pressure is

$$P_{\text{tot}}(n, T, I) = P_b(n, T, I) + \sum_{l=e, \mu} P_l(n, T, I), \quad (40)$$

where $P_b(n, T, I)$ is the contribution of the baryons [see Eq. (24)], i.e.,

$$P_b(n, T, I) = T \sum_{\tau=p, n} s_{\tau}(n, T, I) + \sum_{\tau=n, p} n_{\tau} \mu_{\tau}(n, T, I) - \epsilon_b(n, T, I), \quad (41)$$

while $P_l(n, T, I)$ is the contribution of the leptons [see Eq. (37)]. From Eqs. (39) and (40) we can construct the isothermal curves for energy and pressure and finally derive the isothermal behavior of the EOS of hot nuclear matter under β equilibrium.

III. RESULTS AND DISCUSSION

The schematic potential model, which has been applied in the present work, has been designed to reproduce the results of the more microscopic calculations of both nuclear and neutron-rich matter up to high values of baryon density (see Fig. 1). The behavior of the high-density EOS is of great significance to the determination of the structure of hot protoneutron stars and cold neutron stars. The model has the additional advantage that with the appropriate parametrization, it is able to reproduce different forms of the density dependence of the nuclear symmetry energy.

In view of the above discussion, we calculate the EOS of hot asymmetric nuclear matter by applying a momentum-dependent effective interaction model describing the baryons

interaction. We consider that nuclear matter contains neutrons, protons, electrons, and muons under β equilibrium and charge neutrality. The key quantities in our calculations are the proton fraction Y_p and the asymmetry free energy defined in Eq. (28).

To check the validity of the parabolic approximation (27), we plot in Fig. 4 the difference $F(n, T, I = 1) - F(n, T, I = 0)$ as a function of $(1 - 2Y_p)^2$ at temperatures $T = 10$ and 30 MeV for three baryon densities, i.e., $n = 0.2, 0.3$, and 0.4 fm^{-3} . It is seen that in a good approximation an almost linear relation holds between $F(n, T, I = 1) - F(n, T, I = 0)$ and $(1 - 2Y_p)^2$. A similar behavior of $F_{\text{sym}}(n, T)$ is found by Xu *et al.* [33], applying an isospin and momentum-dependent interaction model.

It is worth presenting the calculation recipe of our work. The outline of our approach is the following. For a fixed baryon density n , temperature T , and asymmetry parameter I , Eq. (4) may be solved iteratively to calculate the quantity

$$\eta_{\tau}(n_{\tau}, T) = \frac{\mu_{\tau}(n_{\tau}, T) - \tilde{U}_{\tau}(n_{\tau}, T)}{T}, \quad (42)$$

where

$$\tilde{U}_{\tau}(n_{\tau}, T) = U_{\tau}(n_{\tau}, k, T) - \tilde{U}_{\tau}(n_{\tau}, k). \quad (43)$$

Knowledge of $\eta_{\tau}(n, T)$ allows the evaluation of $\tilde{U}_{\tau}(n_{\tau}, T)$, which then may be employed to infer the chemical potential from

$$\mu_{\tau}(n_{\tau}, T) = T \eta_{\tau}(n_{\tau}, T) + \tilde{U}_{\tau}(n_{\tau}, T), \quad (44)$$

required as an input for the calculation of the Fermi-Dirac distribution function $f_{\tau}(n_{\tau}, k, T)$. The knowledge of $f_{\tau}(n_{\tau}, k, T)$ permits the calculation of the bulk quantities of asymmetric nuclear matter.

$F_{\text{sym}}(n, T)$, for various values of temperature T , was derived with a least-squares fit to the numerical values according to Eq. (28) and has the form

$$\begin{aligned} F_{\text{sym}}(u; T = 0) &= 13u^{2/3} + 17u, \\ F_{\text{sym}}(u; T = 5) &= 3.653 + 28.018u - 1.512u^2 \\ &\quad + 0.185u^3 - 0.001u^4, \end{aligned}$$




FIG. 5. Helmholtz free energy $F(n, T, I)$ of β -stable matter vs the baryon density n , for various values of T (in MeV).

$$\begin{aligned}
 F_{\text{sym}}(u; T = 10) &= 5.995 + 26.157u - 0.827u^2 \\
 &\quad + 0.068u^3 - 0.002u^4, \\
 F_{\text{sym}}(u; T = 20) &= 13.200 + 21.267u + 0.800u^2 \\
 &\quad - 0.193u^3 + 0.014u^4, \\
 F_{\text{sym}}(u; T = 30) &= 21.087 + 17.626u + 1.645u^2 \\
 &\quad - 0.289u^3 + 0.018u^4, \quad (45)
 \end{aligned}$$

where the case with $T = 0$ is included as well. In that case, F_{sym} coincides with E_{sym} .

In Fig. 5, we present the behavior of the free energy, corresponding to hot β -stable nuclear matter, as a function of the baryon density n , for various values of the temperature T . It is obvious that the thermal effects are more pronounced for low values of the density n .

In Fig. 6, we plot the calculated free energy for symmetric nuclear matter and pure neutron matter of the proposed momentum-dependent interaction model in comparison with the values of the free energy calculated by Friedman and




FIG. 7. Internal energy $E(n, T)$ of β -stable matter as a function of the baryon density n for various values of T .

Pandharipande (FP model) [7]. In the FP model, the EOS of hot and cold nuclear and neutron matter has been calculated in the framework of a variational calculation, where a realistic nuclear interaction containing two- and three-nucleon body nucleon-nucleon interactions has been used. In the case of symmetric nuclear matter, the results of the two models are very similar up to values $n = 0.4\text{--}0.5 \text{ fm}^{-3}$ depending on the values of T . This agreement is expected in the sense that some of the parameters of the model applied in the present work are determined from constraints provided by the empirical properties of symmetric nuclear matter at the equilibrium density $n_0 = 0.16 \text{ fm}^{-3}$.

However, there is an obvious disagreement in the case of pure nuclear matter: in the two models, the free energy exhibits a different trend, especially for higher values of n . This disagreement will be explained below.

In Fig. 7, we display the internal energy per particle $E(n, T) = \epsilon(n, T)/n$ given by Eq. (1) for various values of temperature. Thermal effects, as expected, are more pronounced for low values of the baryon density n and less important for high values of n .





FIG. 6. Free energy of (a) symmetric nuclear matter (for $T = 5$ and $T = 20$ MeV) and (b) pure neutron matter (for $T = 3$ and $T = 20$ MeV) of the proposed model (MDIM) in comparison with the free energy calculated by the Friedman and Pandharipande model (FP) [7].




FIG. 8. Internal energy (for $T = 0$ MeV) for symmetric nuclear matter (SNM) and pure neutron matter (PNM) calculated with the MDIM in comparison with the FP model.

Figure 8 displays the internal energy of symmetric nuclear matter and pure neutron matter, for $T = 0$, calculated by the MDIM and FP models. In accordance with the case of the free energy, there is a very good agreement in symmetric nuclear matter, but an obvious disagreement is exhibited in pure neutron matter. The explanation of the agreement in the first case is the same as in the case of the free energy. The disagreement is due to the completely different behavior of the two models of the nuclear symmetry energy, presented in Fig. 9. In our model, the parameters x_0 , x_3 , Z_1 , and Z_2 chosen so that Eq. (15), for $T = 0$, reproduces the results of Eq. (16) for the function $F(u) = u$. Consequently, $E_{\text{sym}}(n)$ shows an increasing trend as shown in Fig. 9. In contrast, in the FP model, $E_{\text{sym}}(n)$ is a slightly increasing function of n for low n and then a decreasing function of n for $n > 0.5 \text{ fm}^{-3}$.

In addition, we plot the nuclear symmetry energy extracted from experimental results and presented in Ref. [54], where $E_{\text{sym}}(u)$ is parametrized according to the relation $E_{\text{sym}}(u) \approx 31.6u^{0.69}$ as well as experimental results extracted from




FIG. 9. Nuclear symmetry energy calculated with the MDIM in comparison with the FP model as well as the results of Refs. [54,58] and [55].




FIG. 10. Thermal energy $E_{\text{thermal}}(n, T) = E(n, T) - E(n, T = 0)$ of β -stable matter vs the baryon density n , for various values of T .

Ref. [55], where $E_{\text{sym}}(u)$ is given by $E_{\text{sym}}(u) \approx 31.6u^{1.05}$. The important point to be noted is that both cases clearly favor a stiff density dependence of the symmetry energy at higher densities, ruling out the very stiff and very soft predictions. These results can thus be employed to constrain the form of the density dependence of the symmetry energy at supranormal densities relevant for neutron star studies [54]. In the same figure, the theoretical predictions of Ref. [58] are presented, where $E_{\text{sym}}(u)$ is parametrized by $E_{\text{sym}}(u) \approx 32u^{0.8}$.

The results of Ref. [54] are in a good agreement with the present model up to $n = 0.3 \text{ fm}^{-3}$, while the theoretical predictions of Ref. [58] are very close to the present model up to very high values of the baryon density n .

However, our motivation here is not to perform a systematic comparison of various models but to just present the similarities and deviations existing among them. The deviations, concerning the symmetry energy behavior of the two models (MDIM and FP) are well reflected in the behavior of the free energy and internal energy of pure neutron matter as shown in Figs. 6 and 8.

In Fig. 10, we plot the thermal energy per particle

$$E_{\text{thermal}}(n, T) = E(n, T) - E(n, T = 0)$$

of β -stable matter as a function of the baryon density n for various values of temperature T . The most striking feature of $E_{\text{thermal}}(n, T)$ is that for small values of T , the thermal contribution to the internal energy is almost independent of the density n . For high values of T the situation is different, and $E_{\text{thermal}}(n, T)$, for fixed values of T , is a decreasing function of n .

$E_{\text{thermal}}(n, T)$ can be decomposed to separate contributions of the kinetic and potential energies as follows:

$$E_{\text{thermal}}(n, T) = E_{\text{thermal}}^{\text{kin}}(n, T) + E_{\text{thermal}}^{\text{pot}}(n, T).$$

We find that for small values of the baryon density (i.e., $n = 0.2 \text{ fm}^{-3}$) the contribution to $E_{\text{thermal}}(n, T)$ of the potential energy $E_{\text{thermal}}^{\text{pot}}(n, T)$ is 20–10% for $T = 5$ –30 MeV. For medium values of n (0.4 fm^{-3}), it is 43–20%, and for higher values (0.6 fm^{-3}), it is 70–30% for $T = 5$ –30 MeV. Hence, it




FIG. 11. Pressure P of asymmetric nuclear matter for $Y_p = 0.1$ and 0.3 at (a) $T = 10$ and (b) 30 MeV. The full lines give the results calculated from Eq. (24), while the squares represent results obtained by differentiating $F(n, T)$ [Eq. (23)].

is concluded that the potential part of the energy (as a result of the momentum dependence of the interaction) contributes significantly to the thermal energy, mainly for small values of T (for fixed values of n) and for large values of n (for fixed values of T).

At this point, it is worthwhile to compare the results for the pressure obtained by applying Eqs. (24) and (23). Thus, in Fig. 11 we plot P of asymmetric nuclear matter for $Y_p = 0.1$ and 0.3 at $T = 10$ and 30 MeV. The full lines give the results calculated from Eq. (24), while the squares give results obtained by differentiating $F(n, T)$ [Eq. (23)]. The two calculations for the pressure are in excellent agreement. This agreement provides a test of the calculations performed in the present model.

It is of interest also to study the effect of temperature on the baryon pressure defined by Eq. (24). A related quantity is the thermal pressure $P_{\text{thermal}}(n, T)$ defined as

$$P_{\text{thermal}}(n, T) = P(n, T) - P(n, T = 0).$$

$P_{\text{thermal}}(n, T)$ as a function of n , for various values of T is seen in Fig. 12. $P_{\text{thermal}}(n, T)$, in all cases, is an increasing function of the baryon density.

The proton fraction affects the reaction rate of neutrino processes inside that star. If a neutron star has a large proton fraction, the cooling rate may drastically change through the high neutrino emissivity due to the direct URCA process. This process can occur if the proton fraction in the matter of a cold neutron star exceeds the critical value of 0.11 – 0.15 and would lead to the rapid cooling of the neutron star. Thus, it is important to calculate the proton fraction as a function of the baryon density and investigate the temperature effects on that.

Figure 13 displays the fractions of protons, electrons, and muons as functions of the density, for various values of T . The proton fraction is an increasing function of T , and this effect is more pronounced for $T > 10$ MeV. The proton fraction Y_p was derived also with a least-squares fit to the numerical results obtained from our calculations, leading to the following

relations (for $n > 0.15 \text{ fm}^{-3}$):

$$\begin{aligned} Y_p(n; T = 0) &= -0.050 + 0.633n - 0.521n^2 + 0.184n^3, \\ Y_p(n; T = 5) &= -0.046 + 0.625n - 0.514n^2 + 0.179n^3, \\ Y_p(n; T = 10) &= -0.032 + 0.570n - 0.436n^2 + 0.139n^3, \\ Y_p(n; T = 20) &= 0.021 + 0.378n - 0.163n^2 + 0.004n^3, \\ Y_p(n; T = 30) &= 0.109 - 0.062n + 0.908n^2 \\ &\quad - 1.270n^3 + 0.580n^4. \end{aligned} \quad (46)$$

In Fig. 14, we plot the Fermi distribution function $f_{p,n}(n, T)$ for both neutrons and protons for various values of T . We observe that the diffuseness of $f_p(n, T)$ is larger than that of $f_n(n, T)$. We give an explanation (see also Ref. [9]): the ratio of T to the Fermi kinetic energy ϵ_{F_i} is a measure of the thermal effect. Thus by comparing the two ratios, we have (see also the Appendix)

$$\frac{T/\epsilon_{Fp}}{T/\epsilon_{Fn}} = \frac{k_F^{n,2}}{k_F^{p,2}} = \frac{Y_n^{3/2}}{Y_p^{3/2}}.$$




FIG. 12. Thermal pressure $P_{\text{thermal}}(n, T) = P(n, T) - P(n, T = 0)$ of β -stable matter vs the baryon density n , for various values of T .




FIG. 13. Fractions of protons Y_p , electrons Y_e and muons Y_μ of β -stable matter as functions of the baryon density n , for various values of T .

But, because $Y_n > Y_p$, we conclude that we expect the diffuseness to be larger for the proton distribution than for the neutron one, depending of course on the specific value of the ratio Y_p/Y_n . As we will see later, this fact influences the

values of the contributions of protons and neutrons to the total entropy per particle. The entropy, according to relation (21), is an increasing function of the diffuseness. Thus, the contribution of each species to the total value of the entropy depends strongly on the diffuseness of the corresponding Fermi distribution function.

We can provide a second test, concerning the accuracy of the present calculations, by comparing the entropy per baryon calculated by applying Eqs. (21) and (25). Thus, in Fig. 15, we plot S of asymmetric nuclear matter with $Y_p = 0.2$ at $T = 10, 20$, and 30 MeV. The full lines give the entropy calculated from Eq. (21), while the squares give results obtained by differentiating $F(n, T)$ with respect to T [Eq. (25)]. It is obvious that there is again very good agreement of the results, especially for low values of T and high values of n .

In Fig. 16, we plot the contribution of the proton S_p , the neutron S_n , and the total entropy per baryon S . It is obvious that there is a strong effect of T on the values of the entropies mainly for low density values. The main part of the contribution comes from neutrons, whereas the contribution of protons is three times less. It is worthwhile to notice that in spite of $Y_p \sim (1/20 - 3/10)Y_n$, the approximate relation $S_p \sim (1/4 - 3/7)S_n$ holds. This feature is understood by the




FIG. 14. Fermi-Dirac distribution function $f_\tau(n, T)$ for protons and neutrons ($\tau = p, n$ respectively), for $n = 0.2 \text{ fm}^{-3}$, $n = 0.4 \text{ fm}^{-3}$ and $n = 0.6 \text{ fm}^{-3}$ and various values of T .




FIG. 15. Entropy per particle S of asymmetric nuclear matter with $Y_p = 0.2$ at $T = 10, 20, 30$ MeV. The full lines give the entropy calculated from Eq. (21), while the squares give results obtained by differentiating $F(n, T)$ (Eq. (25)).

previous discussion that $f_p(n, T)$ is diffused more broadly than $f_n(n, T)$, so the larger the diffuseness, the larger is the entropy contribution (see also Ref. [9]).

In Fig. 17, we plot the contributions of the electronic S_e , muonic S_μ , and total (leptonic) S_l to the entropy per baryon. The contribution to the entropy of S_e depends slightly on the density for fixed values of T . Our present results are very close to those found by Onsi *et al.* [19], where they employed the analytical approximate formula for the electron entropy density s_e

$$s_e = \frac{1}{3} \frac{\mu_e^2}{(\hbar c)^3} T, \quad \mu_e = \hbar c (3\pi^2 Y_e n)^{1/3}. \quad (47)$$

According to this formula, the contribution of electrons to the entropy per baryon has the form

$$S_e = s_e/n \sim \left(\frac{Y_e^2}{n} \right)^{1/3} T. \quad (48)$$




FIG. 16. Contributions to the total entropy per particle of protons (S_p) (up triangles) neutrons (S_n) (upside down triangles) and the total entropy (S_b) (squares).




FIG. 17. Contributions to the total entropy per particle of electrons (S_e) (up triangles) and muons (S_μ) (upside down triangles), and the total S_l (squares).

The quantity $(\frac{Y_e^2}{n})^{1/3}$ is a function slightly dependent on the density n , so that for a fixed value of T the contribution S_e is almost constant. The muonic contribution to the entropy, for fixed T , increases slightly as a function of the density.

In Fig. 18, we present the EOS of the β -stable hot nuclear matter by taking into account and analyzing the contribution to the total pressure of each component. The main contribution to the total pressure originates from the baryons, whereas the contribution of the leptons is about a few percent compared to P_b . It is worthwhile to notice that thermal effects are not important for the calculation of P_e , but only for P_μ , especially for small values of n ($n < 0.4 \text{ fm}^{-3}$). We found that thermal effects produce a slightly stiffer EOS for cold nuclear matter. The above EOS can be applied to the evaluation of the bulk properties of hot neutron stars (mass and radius).

The study of hot nuclear matter in the absence of neutrino trapping is the first step in studying the properties of hot neutron stars and supernova matter. Next, one can study




FIG. 18. Pressures of baryons P_b , leptons P_l (electrons+muons) and the total pressure P vs the baryon density, n for various values of T .

the more realistic case of neutrino-trapped matter in β equilibrium. In this case, the β -equilibrium conditions in matter are altered from the case in which neutrinos have left the system, and thus the composition of matter is affected. The proton fraction increases dramatically and influences significantly the properties of nuclear matter. Such a work is in progress.

IV. SUMMARY

The evaluation of the EOS of hot nuclear matter is an important problem in nuclear physics and astrophysics. EOS is the basic ingredient in the study of supernova explosions as well as in determining the properties of hot neutron stars. The motivation of the present work is to apply a momentum-dependent interaction model for the study of the hot nuclear matter EOS under β equilibrium in order to be able to study simultaneously thermal effects, not only on the kinetic part of the symmetry energy and symmetry free energy, but also on the interaction part of the above quantities. We calculate the proton fraction, as well as the lepton fractions, by applying the constraints for chemical equilibrium and charge neutrality. The free energy, the internal energy, and the pressure are calculated as functions of baryon density and for various values of temperature. We also concentrate on the evaluation of thermal effects on the internal energy and baryon pressure. Special attention is dedicated to the study of the contribution of the components of β -stable nuclear matter on the entropy per particle, a quantity of great interest in the study of structure and collapse of supernova. We present and analyze the contribution of each component. Finally, we present the EOS of β -stable hot nuclear matter, by taking into account and analyzing the contributions to the total pressure of each component. This EOS can be applied to the evaluation of the gross properties of hot neutron stars, i.e., mass and radius (work in progress).

ACKNOWLEDGMENTS

One of the authors (Ch.C.M) would like to thank Professor Tatsuyauki Takatsuka for valuable comments and correspondence.

APPENDIX

The energy density of baryons [Eq. (1)], at $T = 0$, is given by

$$\begin{aligned} \epsilon(n, I, T = 0) = & \frac{3}{10} E_F^0 n_0 u^{5/3} [(1+I)^{5/3} + (1-I)^{5/3}] \\ & + \frac{1}{3} A n_0 \left[\frac{3}{2} - \left(\frac{1}{2} + x_0 \right) I^2 \right] u^2 \\ & + \frac{\frac{2}{3} B n_0 \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma+1}}{1 + \frac{2}{3} B' \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma-1}} \\ & + \frac{3}{2} n_0 u \sum_{i=1,2} \left[C_i + \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^3 \\ & \times \left(\frac{[(1+I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} - \tan^{-1} \frac{[(1+I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} \right) \\ & + \frac{3}{2} n_0 u \sum_{i=1,2} \left[C_i - \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^3 \\ & \times \left(\frac{[(1-I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} - \tan^{-1} \frac{[(1-I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} \right). \end{aligned} \quad (A1)$$

The pressure of the baryons, at $T = 0$, defined as

$$P = n^2 \frac{d(\epsilon/n)}{dn},$$

is given by

$$\begin{aligned} P(n, I, T = 0) = & \frac{1}{5} n_0 E_F^0 u^{5/3} [(1+I)^{5/3} + (1-I)^{5/3}] + \frac{1}{3} n_0 u^2 A \left[\frac{3}{2} - \left(\frac{1}{2} + x_0 \right) I^2 \right] \\ & + \frac{\frac{2}{3} B \sigma n_0 u^{\sigma+1} \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] (1 + \frac{2}{3\sigma} B' u^{\sigma-1} \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right])}{(1 + \frac{2}{3} B' \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] u^{\sigma-1})^2} \\ & + \frac{n_0 u^2}{2} \sum_{i=1,2} \left[C_i + \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^2 \frac{(1+I)^{1/3}}{u^{2/3}} \left(1 - \frac{1}{1 + \frac{(1+I)^{2/3} u^{2/3}}{\left(\frac{\Lambda_i}{k_F^0} \right)^2}} \right) \\ & + \frac{n_0 u^2}{2} \sum_{i=1,2} \left[C_i - \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^2 \frac{(1-I)^{1/3}}{u^{2/3}} \left(1 - \frac{1}{1 + \frac{(1-I)^{2/3} u^{2/3}}{\left(\frac{\Lambda_i}{k_F^0} \right)^2}} \right). \end{aligned} \quad (A2)$$

In Eq. (A2), E_F^0 is the Fermi energy of symmetric nuclear matter corresponding to equilibrium density n_0 and is given by

$$E_F^0 = \frac{(\hbar k_F^0)^2}{2m}, \quad k_F^0 = \left(3\pi^2 \frac{n_0}{2}\right)^{1/3}. \quad (\text{A3})$$

The Fermi momenta of protons and neutrons are

$$k_F^p = (3\pi^2 x n)^{1/3} = \left(3\pi^2 \frac{1-I}{2} n\right)^{1/3},$$

$$k_F^n = (3\pi^2 (1-x)n)^{1/3} = \left(3\pi^2 \frac{1+I}{2} n\right)^{1/3}.$$

The chemical potentials of protons and neutrons, at $T = 0$, are given by

$$\mu_n = E + u \left(\frac{\partial E}{\partial u} \right)_{Y_p} - Y_p \left(\frac{\partial E}{\partial Y_p} \right)_n,$$

$$\mu_p = \mu_n + \left(\frac{\partial E}{\partial Y_p} \right)_n,$$

$$\hat{\mu} = \mu_n - \mu_p = - \left(\frac{\partial E}{\partial Y_p} \right)_n,$$

where

$$\begin{aligned} \left(\frac{\partial E}{\partial u} \right)_{Y_p} = & \frac{1}{5} E_F^0 u^{-1/3} [(1+I)^{5/3} + (1-I)^{5/3}] + \frac{1}{3} A \left[\frac{3}{2} - \left(\frac{1}{2} + x_0 \right) I^2 \right] \\ & + \frac{\frac{2}{3} B u^{\sigma-1} \sigma \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] \left(1 + \frac{2}{3\sigma} B' u^{\sigma-1} \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] \right)}{\left(1 + \frac{2}{3} B' u^{\sigma-1} \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] \right)^2} \\ & + \frac{1}{2} \sum_{i=1,2} \left[C_i + \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^2 \frac{(1+I)^{1/3}}{u^{2/3}} \left(1 - \frac{1}{1 + \frac{(1+I)^{2/3} u^{2/3}}{\left(\frac{\Lambda_i}{k_F^0} \right)^2}} \right) \\ & + \frac{1}{2} \sum_{i=1,2} \left[C_i - \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^2 \frac{(1-I)^{1/3}}{u^{2/3}} \left(1 - \frac{1}{1 + \frac{(1-I)^{2/3} u^{2/3}}{\left(\frac{\Lambda_i}{k_F^0} \right)^2}} \right), \end{aligned} \quad (\text{A5})$$

$$\begin{aligned} \left(\frac{\partial E}{\partial Y_p} \right)_n = & \frac{1}{2} E_F^0 u^{2/3} [(1+I)^{2/3} - (1-I)^{2/3}] - \frac{1}{3} A u \left(\frac{1}{2} + x_0 \right) I \\ & - \frac{\frac{2}{3} B u^{\sigma} \left(\frac{1}{2} + x_3 \right) I}{\left(1 + \frac{2}{3} B' u^{\sigma-1} \left[\frac{3}{2} - \left(\frac{1}{2} + x_3 \right) I^2 \right] \right)^2} \\ & + \frac{3}{2} \sum_{i=1,2} \left[\frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^3 \left(\frac{[(1+I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} - \tan^{-1} \frac{[(1+I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} \right) \\ & - \frac{3}{2} \sum_{i=1,2} \left[\frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^3 \left(\frac{[(1-I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} - \tan^{-1} \frac{[(1-I)u]^{1/3}}{\frac{\Lambda_i}{k_F^0}} \right) \\ & + \frac{1}{2} \sum_{i=1,2} \left[C_i + \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^2 \frac{u^{1/3}}{(1+I)^{2/3}} \left(1 - \frac{1}{1 + \frac{(1+I)^{2/3} u^{2/3}}{\left(\frac{\Lambda_i}{k_F^0} \right)^2}} \right) \\ & - \frac{1}{2} \sum_{i=1,2} \left[C_i - \frac{C_i - 8Z_i}{5} I \right] \left(\frac{\Lambda_i}{k_F^0} \right)^2 \frac{u^{1/3}}{(1-I)^{2/3}} \left(1 - \frac{1}{1 + \frac{(1-I)^{2/3} u^{2/3}}{\left(\frac{\Lambda_i}{k_F^0} \right)^2}} \right). \end{aligned} \quad (\text{A6})$$

- [1] H. A. Bethe, Rev. Mod. Phys. **62**, 801 (1990); Annu. Rev. Nucl. Part. Sci. **38**, 1 (1988).
- [2] Madappa Prakash, I. Bombaci, Manju Prakash, P. J. Ellis, J. M. Lattimer, R. Knorren, Phys. Rep. **280**, 1 (1997).
- [3] *Isospin Physics in Heavy-Ion Collisions at Intermediate Energies*, edited by Bao-An Li and W. Udo Schröder (Nova Science, New York, 2001).
- [4] B. A. Li, L. W. Chen, and C. M. Ko, Phys. Rep. **464**, 113 (2008).
- [5] G. Baym, H. A. Bethe, and C. J. Pethick, Nucl. Phys. **A175**, 225 (1971).
- [6] H. A. Bethe, G. E. Brown, J. Applegate, and J. M. Lattimer, Nucl. Phys. **A324**, 487 (1979).
- [7] B. Friedman and V. R. Pandharipande, Nucl. Phys. **A361**, 502 (1981).
- [8] J. M. Lattimer and F. D. Swesty, Nucl. Phys. **A535**, 331 (1991).
- [9] T. Takatsuka, S. Nishizaki, and J. Hiura, Prog. Theor. Phys. **92**, 779 (1994).
- [10] T. Takatsuka, Prog. Theor. Phys. **95**, 901 (1996).
- [11] C. Das, R. Sahu, and A. Mishra, Phys. Rev. C **75**, 015807 (2007).
- [12] Bernard ter Haar and R. Malfliet, Phys. Rev. Lett. **56**, 1237 (1986).
- [13] I. Bombaci, Madappa Prakash, Manju Prakash, P. J. Ellis, J. M. Lattimer, and G. E. Brown, Nucl. Phys. **A583**, 623 (1995).
- [14] W. A. Küpper, G. Wegmann, and E. R. Hilf, Ann. Phys. (NY) **88**, 454 (1974).
- [15] J. M. Lattimer and D. G. Ravenhall, Astrophys. J. **223**, 314 (1978).
- [16] M. F. El Eid and W. Hillebrandt, Astron. Astrophys. Suppl. Ser. **42**, 215 (1980).
- [17] P. Lamb, Mon. Not. R. Astron. Soc. **188**, 565 (1979).
- [18] H. M. Antia, B. Banerjee, and S. M. Chitre, Astrophys. Space Sci. **69**, 471 (1980).
- [19] M. Onsi, H. Przysiecki, and J. M. Pearson, Phys. Rev. C **50**, 460 (1994).
- [20] K. Sumiyoshi and H. Toki, Astrophys. J. **422**, 700 (1994).
- [21] J. M. Lattimer, C. J. Pethick, D. G. Ravenhall, and D. Q. Lamb, Nucl. Phys. **A432**, 646 (1985).
- [22] H. Kanzawa, K. Oyamatsu, K. Sumiyoshi, and M. Takano, Nucl. Phys. **A791**, 232 (2007).
- [23] M. Modarres, J. Phys. G: Nucl. Part. Phys. **23**, 923 (1997); **21**, 351 (1995); **19**, 1349 (1993).
- [24] R. Manka, I. Bednarek, and G. Przybyla, Phys. Rev. C **62**, 015802 (2000).
- [25] W. Zuo, Z. H. Li, A. Li, and G. C. Lu, Phys. Rev. C **69**, 064001 (2004).
- [26] L. W. Chen, F. S. Zhang, Z. H. Lu, W. F. Li, Z. Y. Zhu, and H. R. Ma, J. Phys. G: Nucl. Part. Phys. **27**, 1799 (2001).
- [27] V. K. Mishra, G. Fai, L. P. Csernai, and E. Osnes, Phys. Rev. C **47**, 1519 (1993).
- [28] L. P. Csernai, G. Fai, C. Gale, and E. Osnes, Phys. Rev. C **46**, 736 (1992).
- [29] S. J. Lee and A. Z. Mekjian, Phys. Rev. C **63**, 044605 (2001); A. Z. Mekjian, S. J. Lee, and L. Zamick, *ibid.* **72**, 044305 (2005); A. Z. Mekjian, S. J. Lee, and L. Zamick, Phys. Lett. **B621**, 239 (2005); S. J. Lee and A. Z. Mekjian, Phys. Rev. C **77**, 054612 (2008).
- [30] H. Müller and B. D. Serot, Phys. Rev. C **52**, 2072 (1995).
- [31] M. Baldo and L. S. Ferreira, Phys. Rev. C **59**, 682 (1999).
- [32] G. F. Burgio, M. Baldo, O. E. Nicotra, and H. J. Schulze, Astrophys. Space Sci. **308**, 387 (2007); O. E. Nicotra, M. Baldo, G. F. Burgio, and H. J. Schulze, Astron. Astrophys. **451**, 213 (2006).
- [33] J. Xu, L. W. Chen, B. A. Li, and H. R. Ma, Phys. Rev. C **75**, 014607 (2007).
- [34] P. Wang, Phys. Rev. C **61**, 054904 (2000).
- [35] P. K. Jena and L. P. Singh, Phys. Rev. C **70**, 045803 (2004).
- [36] J. N. De, N. Rudra, S. Pal, and S. K. Samaddar, Phys. Rev. C **53**, 780 (1996); T. Sil, B. K. Agrawal, J. N. De, and S. K. Samaddar, *ibid.* **63**, 054604 (2001); S. K. Samaddar, J. N. De, X. Viñas, and M. Centelles, *ibid.* **76**, 041602(R) (2007).
- [37] L. Tolos, B. Frieman, and A. Schwenk, Nucl. Phys. **A806**, 105 (2008).
- [38] M. Abd-Alla, H. S. Ragab, and M. Y. M. Hassan, Acta Phys. Pol. B **24**, 1519 (1993).
- [39] Ch. C. Moustakidis, Phys. Rev. C **76**, 025805 (2007).
- [40] Ch. C. Moustakidis, Phys. Rev. C **78**, 054323 (2008).
- [41] R. B. Wiringa, V. Fiks, and A. Fabrocini, Phys. Rev. C **38**, 1010 (1988).
- [42] A. Akmal, V. R. Pandharipande, and D. G. Ravenhall, Phys. Rev. C **58**, 1804 (1998).
- [43] G. F. Bertsch and S. Das Gupta, Phys. Rep. **160**, 189 (1988).
- [44] C. Gale, G. F. Bertsch, and S. Das Gupta, Phys. Rev. C **35**, 1666 (1987).
- [45] M. Prakash, T. T. S. Kuo, and S. Das Gupta, Phys. Rev. C **37**, 2253 (1988); G. M. Welke, M. Prakash, T. T. S. Kuo, S. Das Gupta, and C. Gale, *ibid.* **38**, 2101 (1988); C. Gale, G. M. Welke, M. Prakash, S. J. Lee, and S. Das Gupta, *ibid.* **41**, 1545 (1990).
- [46] B. A. Li, C. B. Das, S. Das Gupta, and C. Gale, Phys. Rev. C **69**, 011603(R) (2004); C. B. Das, S. Das Gupta, C. Gale, and B. A. Li, *ibid.* **67**, 034611 (2003); L. W. Chen, C. M. Ko, and B. A. Li, *ibid.* **69**, 054606 (2004); B. A. Li, C. B. Das, S. Das Gupta, and C. Gale, Nucl. Phys. **A735**, 563 (2004); B. A. Li, L. W. Chen, G. C. Yong, and W. Zuo, Phys. Lett. **B634**, 378 (2006); B. A. Li and A. W. Steiner, Phys. Lett. **B642**, 436 (2006).
- [47] R. B. Wiringa, Phys. Rev. C **38**, 2967 (1988).
- [48] G. Q. Li and R. Machleidt, Phys. Rev. C **48**, 2707 (1993).
- [49] M. Kutschera, Phys. Lett. **B340**, 1 (1994); Z. Phys. A **348**, 263 (1994); Acta Phys. Pol. B **29**, 25 (1998); S. Kubis and M. Kutschera, Nucl. Phys. **A720**, 189 (2003).
- [50] B. A. Li, Phys. Rev. Lett. **88**, 192701 (2002).
- [51] C. Fuchs and H. H. Wolter, Eur. Phys. J. A **30**, 5 (2006).
- [52] N. K. Glendenning, *Compact Stars: Nuclear Physics, Particle Physics, and General Relativity* (Springer-Verlag, New York, 1997).
- [53] R. B. Wiringa, V. Fiks, and A. Fabrocini, Phys. Rev. C **38**, 1010 (1988).
- [54] D. V. Shetty, S. J. Yennello, and G. A. Souliotis, Phys. Rev. C **75**, 034602 (2007); **76**, 024606 (2007).
- [55] L. W. Chen, C. M. Ko, and B. A. Li, Phys. Rev. Lett. **94**, 032701 (2005).
- [56] M. B. Tsang *et al.*, Phys. Rev. Lett. **92**, 062701 (2004).
- [57] C. H. Lee, T. T. S. Kuo, G. Q. Li, and G. E. Brown, Phys. Rev. C **57**, 3488 (1998).

- [58] F. Sammarruca and P. Liu, arXiv:0806.1936 [nucl-th] (unpublished, 2008).
- [59] V. P. Psonis, Ch. C. Moustakidis, and S. E. Massen, Mod. Phys. Lett. A **22**, 1233 (2007).
- [60] M. Prakash, *The Equation of State and Neutron Star*, lectures delivered at the Winter School held in Puri, India (unpublished, 1994).
- [61] B. A. Li, C. M. Ko, and Z. Ren, Phys. Rev. Lett. **78**, 1644 (1997).
- [62] V. Baran, M. Colonna, V. Greco, and M. Di Toro, Phys. Rep. **410**, 335 (2005).
- [63] D. L. Goodstein, *States of Matter* (Dover, New York, 1985).
- [64] A. L. Fetter and J. D. Walecka, *Quantum Theory of Many-Particle Systems* (Dover, Mineola, New York, 2003).