

Breaking of factorization of two-particle correlations in hydrodynamics

Fernando G. Gardim,¹ Frédérique Grassi,¹ Matthew Luzum,^{2,3,4} and Jean-Yves Ollitrault²

¹Instituto de Física, Universidade de São Paulo, C.P. 66318, 05315-970, São Paulo-SP, Brazil

²CNRS, URA2306, IPhT, Institut de physique théorique de Saclay, F-91191 Gif-sur-Yvette, France

³McGill University, 3600 University Street, Montreal QC H3A 2T8, Canada

⁴Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA

(Received 7 November 2012; published 26 March 2013)

The system formed in ultrarelativistic heavy-ion collisions behaves as a nearly perfect fluid. This collective behavior is probed experimentally by two-particle azimuthal correlations, which are typically averaged over the properties of one particle in each pair. In this Rapid Communication, we argue that much additional information is contained in the detailed structure of the correlation. In particular, the correlation matrix exhibits an approximate factorization in transverse momentum, which is taken as strong evidence for the hydrodynamic picture, while deviations from the factorized form are taken as a signal of intrinsic, “nonflow” correlations. We show that hydrodynamics in fact predicts factorization breaking as a natural consequence of initial-state fluctuations and averaging over events. We derive the general inequality relations that hold if flow dominates, and which are saturated if the matrix factorizes. For transverse momenta up to 5 GeV, these inequalities are satisfied in data, but not saturated. We find factorization breaking in event-by-event ideal hydrodynamic calculations that is at least as large as in data and argue that this phenomenon opens a new window on the study of initial fluctuations.

DOI: [10.1103/PhysRevC.87.031901](https://doi.org/10.1103/PhysRevC.87.031901)

PACS number(s): 25.75.Ld, 24.10.Nz, 25.75.Gz

Introduction. In relativistic heavy-ion-collision experiments a large second Fourier harmonic is observed in two-particle correlations as a function of relative azimuthal angle [1–4]. This has long been considered a sign of significant collective behavior [5], or “elliptic flow,” indicating the existence of a strongly interacting, low-viscosity fluid [6]. However, only recently has it been realized that *all* such correlations observed between particles separated by a large relative pseudorapidity could be explained by this collective behavior [7–15], at least for the bulk of the system.

One significant piece of evidence for this view was the recent observation of the factorization [16–19] of two-particle correlations into a product of a function of properties of only one of the particles times a function of the properties of the second. Specifically, for pairs of particles in various bins of transverse momentum p_T , factorization of each Fourier harmonic was tested as [16]

$$V_{n\Delta}(p_T^a, p_T^b) \equiv \langle \cos n(\phi^a - \phi^b) \rangle \stackrel{?}{=} v_n(p_T^a) v_n(p_T^b), \quad (1)$$

where the brackets indicate an average over pairs of particles (a and b) coming from the same event as well as an average over a set of collision events, and ϕ^a (ϕ^b) is the azimuthal angle of particle a (b). The left-hand side is a (symmetric) function of two variables p_T^a and p_T^b and in general may not factorize into a product of a function v_n of each variable individually. The fact that this factorization holds at least approximately, then, is a nontrivial observation about the structure of the correlation.

While most known sources of nonflow correlations do not factorize at low p_T [20], a type of factorization comes naturally in a pure hydrodynamic picture where particles are emitted independently. They thus have no intrinsic correlations with other particles, carrying only information about their orientation with respect to the system as a whole. This causes the two-particle probability distribution in a single collision event to factorize [21] into a product of one-particle

distributions,

$$\frac{dN_{\text{pairs}}}{d^3 p^a d^3 p^b} \stackrel{(\text{flow})}{=} \frac{dN}{d^3 p^a} \frac{dN}{d^3 p^b}. \quad (2)$$

Inspired by this fact, it has often been stated [19,22,23] that the factorization test in Eq. (1) should work perfectly in hydrodynamics. The observed approximate factorization was hailed as a success for the flow interpretation of correlations, while small deviations from the factorized form was interpreted as a gradual breakdown of the hydrodynamic description with increasing transverse momentum, and of an increasing contribution from other sources of correlations.

In this work, we show that factorization as in Eq. (1) is not necessarily present even in an ideal hydrodynamic system governed by Eq. (2) because of event-by-event fluctuations [13,24,25]. These stem from quantum fluctuations: the collision takes place over a very short timescale and takes a snapshot of the wave function of incoming nuclei. In the presence of fluctuations, we show that the correlation matrix satisfies general inequalities, which are saturated by Eq. (1). We test these inequalities on ALICE data and point out where breaking of factorization occurs. We then illustrate with a full event-by-event hydrodynamic calculation that the same deviation seen in experiment is also present in ideal hydrodynamics.

Hydrodynamics and two-particle correlations. We begin by recalling the discussion originally found in Ref. [26]. In a pure hydrodynamic picture, particles are emitted independently from the fluid at the end of the system evolution according to some underlying one-particle probability distribution. One can write any such distribution as a Fourier series in the azimuthal angle ϕ of the particles

$$\frac{2\pi}{N} \frac{dN}{d\phi} = \sum_{n=-\infty}^{\infty} V_n(p_T, \eta) e^{-in\phi}, \quad (3)$$

where $V_n = \{e^{in\phi}\}$ is the n th complex Fourier flow coefficient, and curly brackets indicate an average over the probability density in a single event. Writing $V_n = v_n e^{in\Psi_n}$, where v_n is the (real) anisotropic flow coefficient and Ψ_n is the corresponding phase, and using $V_{-n} = V_n^*$ (where V_n^* is the complex conjugate of V_n), this can be rewritten as

$$\frac{2\pi}{N} \frac{dN}{d\phi} = 1 + 2 \sum_{n=1}^{\infty} v_n(p_T, \eta) \cos n[\phi - \Psi_n(p_T, \eta)]. \quad (4)$$

Note that, for this form to describe an arbitrary distribution, both v_n and Ψ_n may depend on transverse momentum p_T and pseudorapidity η .

In this picture, the relation in Eq. (2) holds, and a complex Fourier harmonic of the two-particle correlation factorizes in each event as

$$\{e^{in(\phi^a - \phi^b)}\} = \{e^{in\phi^a}\} \{e^{-in\phi^b}\} = V_n^a V_n^{b*} = v_n^a v_n^b e^{in(\Psi_n^a - \Psi_n^b)}. \quad (5)$$

This factorization only holds in a single hydro event. Both the magnitudes and the phases of anisotropic flow fluctuate event to event [13,24,25]. The experimental quantity, Eq. (1), is then obtained by averaging over events:

$$V_{n\Delta}(p_T^a, p_T^b) = \langle V_n^a V_n^{b*} \rangle = \langle v_n^a v_n^b e^{in(\Psi_n^a - \Psi_n^b)} \rangle. \quad (6)$$

Due to parity symmetry, only the real part remains after this average, hence the cosine in Eq. (1).

From this relation alone, one can make the following general statements about the event-averaged correlation matrix: the diagonal elements must be positive, and the off-diagonal elements must satisfy a Cauchy-Schwarz inequality,

$$V_{n\Delta}(p_T^a, p_T^a) \geq 0, \quad (7)$$

$$V_{n\Delta}(p_T^a, p_T^b)^2 \leq V_{n\Delta}(p_T^a, p_T^a) V_{n\Delta}(p_T^b, p_T^b). \quad (8)$$

Factorization, Eq. (1), implies that the second inequality is saturated, i.e., equality is achieved. Thus, while flow does not necessarily imply factorization, any violation of these inequalities is an unambiguous indication of the presence of nonflow correlations.

An inspection of published data from the ALICE Collaboration [16] shows that these inequalities are indeed violated in certain regimes [27]. For $n = 3$, diagonal elements $V_{3\Delta}(p_T^a, p_T^a)$ are negative above 5 GeV for 0%–10% centrality, and above 4 GeV for 40%–50% centrality. This is a clear indication that there are nonflow correlations at high p_T . For instance, the correlation between back-to-back jets typically yields a relative angle $\Delta\phi \sim \pi$, thus producing a negative $V_{3\Delta}$ at high p_T . For $n = 1$, diagonal elements are negative not only at high p_T (with a slightly higher threshold than for $n = 3$), but also for p_T between 1 and 1.5 GeV. This is believed to be caused by the correlation from global momentum conservation [19,28], but it is interesting to note that its effect can be noticed by a simple inspection of elements.

In order to check the validity of the second inequality (8), we introduce the ratio

$$r_n \equiv \frac{V_{n\Delta}(p_T^a, p_T^b)}{\sqrt{V_{n\Delta}(p_T^a, p_T^a) V_{n\Delta}(p_T^b, p_T^b)}}, \quad (9)$$

which is defined when diagonal elements $V_{n\Delta}(p_T^a, p_T^a)$ and $V_{n\Delta}(p_T^b, p_T^b)$ are both positive and lies between -1 and $+1$ if Eq. (8) holds. Factorization corresponds to the limit $r_n = \pm 1$. Figure 1 displays r_2 and r_3 as a function of p_T^a and p_T^b for Pb-Pb collisions at 2.76 TeV, 0%–10% centrality. ALICE results for r_2 satisfy the inequalities (8) at all p_T . When both particles are below 1.5 GeV, the inequality is saturated, $r_2 = 1$, within errors. As soon as one of the particles is above 1.5 GeV, however, r_2 is smaller than unity, and the difference with unity increases with the difference $p_T^a - p_T^b$. Results for




FIG. 1. (Color online) Ratio of nondiagonal to diagonal correlations, defined by Eq. (9), plotted on an interleaved p_T^a, p_T^b axis: p_T^b is constant between each long hash on the x axis. Filled stars show ALICE data for 0%–10% central Pb-Pb collisions at 2.76 TeV [16]. Open circles show ideal hydrodynamic calculations for 0%–10% central Au-Au collisions at 200 GeV.

r_3 are qualitatively similar below 5 GeV, with larger error bars. However, r_3 is closer to 1 than r_2 between 2 and 3 GeV. The values of r_n for midcentral collisions (40%–50% centrality, not shown) are comparable to the values for central collisions, although r_2 is slightly closer to 1.

The ALICE Collaboration concluded from their analysis that factorization holds approximately for $n > 1$ and p_T below 4 GeV. However, their results actually show evidence for a slight breaking of factorization for $n = 2$, as soon as one of the particles has $p_T > 1.5$ GeV. Even though factorization is broken, the general inequalities implied by flow are satisfied for $n = 2$ and $n = 3$ below 5 GeV for central collisions. It is therefore worth investigating in more detail to what extent the breaking of factorization which is seen experimentally can be understood within hydrodynamics.

First, we recall under which conditions factorization holds in hydrodynamics. It implies that the Cauchy-Schwarz inequality (8) is saturated. By inspection of Eq. (6), this in turn implies that the complex flow vectors V_n^a and V_n^b are linearly dependent. This is true only under the following assumptions:

- (i) By parity symmetry, $\Psi_n^a - \Psi_n^b = 0$ in each event. That is, Ψ_n does not depend on p_T , which removes the exponential from the right-hand side of Eq. (5).
- (ii) $v_n(p_T)$ changes from event to event by only a global factor, with no p_T -dependent fluctuations. $v_n(p_T)$ on the right-hand side of Eq. (1) then represents the rms value over events.

In general, fluctuations ensure that these conditions are not met exactly, and the factorization of Eq. (1) will not be perfect. Within hydrodynamics, the ratio r_n in Eq. (9) has a simple interpretation. Inserting Eq. (6) into Eq. (9), one obtains

$$r_n = \frac{\langle V_n^{a*} V_n^b \rangle}{\sqrt{\langle |V_n^a|^2 \rangle \langle |V_n^b|^2 \rangle}}. \quad (10)$$

The ratio r_n thus represents the linear correlation between the complex flow vectors at momenta p_T^a and p_T^b . Since in each event, V_n^a is a smooth function of p_T^a , one expects that the correlation is stronger when $p_T^a \simeq p_T^b$ and decreases as the difference between p_T^a and p_T^b increases as a result of the decoherence induced by initial fluctuations. ALICE data confirm this qualitative expectation.

Note that even in a single hydrodynamic event, factorization holds in the complex form (5) but is broken if one takes the real part before averaging over particle pairs, as in Eq. (1). The ratio r_n in Eq. (10) is then $\cos n(\Psi_n^a - \Psi_n^b)$, which is smaller than unity as soon as the flow angle Ψ_n depends on p_T .

The question then becomes: how large are factorization-breaking effects in hydrodynamics, and do they have the same properties as seen in data? If purely hydrodynamic calculations give the same result as experiment, then the observed breaking of factorization may not indicate the presence of nonflow correlations.

Ideal hydrodynamic calculations. To illustrate these concepts we perform calculations using the NeXSpheRIO model [29]. This model solves the equations of relativistic ideal hydrodynamics with fluctuating initial conditions given by

the NeXuS event generator [30]. It has proven successful in reproducing results from the BNL Relativistic Heavy Ion Collider (RHIC), in particular the structure of two-particle angular correlations in Au-Au collisions at the top RHIC energy [9]. It has recently been shown to reproduce the whole set of measured anisotropic flow data [31–33]. Our calculations are therefore performed for Au-Au collisions at the top RHIC energy, not for Pb-Pb collisions at the energy of the CERN Large Hadron Collider (LHC), as would be appropriate for a direct quantitative comparison with ALICE data. Our results are merely meant as a proof of concept and as a prediction for measurements at RHIC. Note that the main source of fluctuations (namely, the finite number of nucleons within the nucleus) is identical in both cases.

We run 30 000 NeXuS events, which are then sorted into 10% centrality bins defined by the number of participant nucleons, and then evolved hydrodynamically. Anisotropic flow is calculated accurately in every event [34]. The ratio r_n is displayed in Fig. 1 for $n = 2$ and $n = 3$. Deviations from the factorization limit $r = 1$ are already seen at low momentum but become larger as the difference between p_T^a and p_T^b increases, as expected from the general arguments above. Surprisingly, the breaking of factorization appears *larger* in hydrodynamics than in experiment.

The ALICE collaboration has studied factorization by performing a global fit of the measured correlation $V_{n\Delta}(p_T^a, p_T^b)$ by the right-hand side of Eq. (1), where $v_n(p_T)$ is a fit parameter [16]. The ratio of the measured correlation to the best fit differs from unity if factorization is broken. We can apply the same procedure to our hydrodynamic results. The result is shown in Fig. 2. Again, hydrodynamic calculations and experimental data show similar trends, with the noticeable difference that the breaking of factorization is significantly *stronger* in ideal hydrodynamics than in data.

Several effects can explain this discrepancy. First, the average p_T is significantly larger at LHC than at RHIC [35], so that it might be more natural to compare, e.g., 4 GeV at RHIC to 5 GeV at LHC, rather than doing the comparison at the same p_T . The second effect is viscosity, which is neglected in our calculation. Shear viscosity, in particular, tends to damp the effect of initial fluctuations [36]. It is therefore natural that it will also decrease the breaking of factorization induced by initial fluctuations. A similar observation is that the linear correlation between the initial eccentricity and the final anisotropic flow is stronger in viscous hydrodynamics [37] than in ideal hydrodynamics [34].

Conclusions. We have demonstrated that the detailed structure of two-particle angular correlations contains much more information than traditional analyses of anisotropic flow, where the correlation is averaged over one of the particles [38]. Even though such two-dimensional analyses are much more demanding in terms of statistics than traditional analyses, they bring new, independent insight into the underlying physics of flow fluctuations.

In particular, we have shown that quantum fluctuations in the wave function of incoming nuclei result in a decoherence in the angular correlations produced by collective flow, which becomes increasingly important as the difference between particle momenta increases. Due to this effect, factorization




FIG. 2. (Color online) Ratio of the left-hand side to the right-hand side of Eq. (1). Filled stars show ALICE data for 0%–10% central Pb-Pb collisions at 2.76 TeV [16]. Open circles show ideal hydrodynamic calculations for 0%–10% central Au-Au collisions at 200 GeV.

of angular correlations is broken even if collective flow is the only source of correlations. Our numerical calculations show that factorization breaking can be as strong in hydrodynamics as in experimental data, thereby suggesting that all correlations below $p_T \sim 5$ GeV (for central Pb-Pb collisions at LHC near midrapidity) may actually be dominated by flow. The sensitivity of this decoherence phenomenon to viscosity has not yet been investigated, but we anticipate that factorization should be restored as viscosity increases, thus potentially offering a new means of constraining the viscosity from data. On the other hand, thermal fluctuations should be considered along with viscosity [39] and may also contribute to factorization breaking.

Decoherence also provides a natural explanation for the important observation that event-by-event fluctuations reduce elliptic flow at high p_T [40], thus improving agreement between hydrodynamics and experimental data. Indeed, v_2 at high p_T is inferred from azimuthal correlations between a high- p_T particle and all other particles—mostly low- p_T particles—and these azimuthal correlations are reduced due to the decoherence phenomenon. Note that the other main explanation for the reduction of v_2 at high p_T , viscosity,

typically relies on the assumption of a quadratic momentum dependence of the viscous correction to the distribution function at freeze-out δf , which may not be correct [41].

In this paper, we have focused on the transverse momentum dependence of the correlations. The rapidity dependence of the correlation is also worth investigating. In particular, it was recently observed that azimuthal correlations decrease as a function of the relative pseudorapidity [42], at variance with common lore that correlations due to flow are essentially independent of rapidity. While standard models of initial conditions do predict a mild rapidity dependence of azimuthal correlations [43,44], longitudinal fluctuations [45] could also produce a decoherence effect similar to the one studied here. The detailed structure of two-particle correlations as a function of both particle momenta thus opens a new window on the study of flow fluctuations.

Acknowledgments. We thank Yojiro Hama for useful discussions. This work is funded by FAPESP under Projects 09/50180-0 and 09/16860-3, by the FAPESP/CNRS Grant 2011/51854-0, and by CNPq under Project 301141/2010-0. M.L. is supported by the European Research Council under the Advanced Investigator Grant ERC-AD-267258.

-
- [1] K. H. Ackermann *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **86**, 402 (2001).
 [2] S. S. Adler *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **91**, 182301 (2003).
 [3] B. B. Back *et al.* (PHOBOS Collaboration), *Phys. Rev. C* **72**, 051901 (2005).
 [4] K. Aamodt *et al.* (ALICE Collaboration) *Phys. Rev. Lett.* **105**, 252302 (2010).
 [5] J.-Y. Ollitrault, *Phys. Rev. D* **46**, 229 (1992).
 [6] P. Romatschke and U. Romatschke, *Phys. Rev. Lett.* **99**, 172301 (2007).
 [7] A. P. Mishra, R. K. Mohapatra, P. S. Saumia, and A. M. Srivastava, *Phys. Rev. C* **77**, 064902 (2008).
 [8] P. Sorensen, [arXiv:0808.0503](https://arxiv.org/abs/0808.0503).
 [9] J. Takahashi, B. M. Tavares, W. L. Qian, R. Andrade, F. Grassi, Y. Hama, T. Kodama, and N. Xu, *Phys. Rev. Lett.* **103**, 242301 (2009).
 [10] Y. Hama, R. P. G. Andrade, F. Grassi, and W.-L. Qian, *Nonlinear Phenom. Complex Syst.* **12**, 466 (2009).
 [11] R. P. G. Andrade, F. Grassi, Y. Hama, and W.-L. Qian, *Phys. Lett. B* **712**, 226 (2012).

- [12] P. Sorensen, *J. Phys. G* **37**, 094011 (2010).
- [13] B. Alver and G. Roland, *Phys. Rev. C* **81**, 054905 (2010); **82**, 039903(E) (2010).
- [14] P. Staig and E. Shuryak, *Phys. Rev. C* **84**, 034908 (2011); **84**, 044912 (2011).
- [15] M. Luzum, *Phys. Lett. B* **696**, 499 (2011).
- [16] K. Aamodt *et al.* (ALICE Collaboration), *Phys. Lett. B* **708**, 249 (2012).
- [17] B. Alver *et al.* (PHOBOS Collaboration), *Phys. Rev. C* **81**, 034915 (2010).
- [18] S. Chatrchyan *et al.* (CMS Collaboration), *Eur. Phys. J. C* **72**, 2012 (2012).
- [19] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. C* **86**, 014907 (2012).
- [20] D. Kikola, L. Yi, S. I. Esumi, F. Wang, and W. Xie, *Phys. Rev. C* **86**, 014901 (2012).
- [21] P. M. Dinh, N. Borghini, and J.-Y. Ollitrault, *Phys. Lett. B* **477**, 51 (2000).
- [22] A. Adare, *J. Phys. G* **38**, 124091 (2011).
- [23] J. Jia, *J. Phys. G* **38**, 124012 (2011).
- [24] M. Miller and R. Snellings, [arXiv:nucl-ex/0312008](https://arxiv.org/abs/nucl-ex/0312008).
- [25] B. Alver *et al.* (PHOBOS Collaboration), *Phys. Rev. Lett.* **98**, 242302 (2007).
- [26] M. Luzum, *J. Phys. G* **38**, 124026 (2011).
- [27] J.-Y. Ollitrault and F. G. Gardim, [arXiv:1210.8345](https://arxiv.org/abs/1210.8345).
- [28] E. Retinskaya, M. Luzum, and J.-Y. Ollitrault, *Phys. Rev. Lett.* **108**, 252302 (2012).
- [29] Y. Hama, T. Kodama, and O. Socolowski, Jr., *Braz. J. Phys.* **35**, 24 (2005).
- [30] H. J. Drescher, M. Hladik, S. Ostapchenko, T. Pierog, and K. Werner, *Phys. Rep.* **350**, 93 (2001).
- [31] F. G. Gardim, F. Grassi, M. Luzum, and J.-Y. Ollitrault, *Phys. Rev. Lett.* **109**, 202302 (2012).
- [32] F. G. Gardim, F. Grassi, Y. Hama, M. Luzum, and J.-Y. Ollitrault, *Phys. Rev. C* **83**, 064901 (2011).
- [33] R. D. de Souza, J. Takahashi, T. Kodama, and P. Sorensen, *Phys. Rev. C* **85**, 054909 (2012).
- [34] F. G. Gardim, F. Grassi, M. Luzum, and J.-Y. Ollitrault, *Phys. Rev. C* **85**, 024908 (2012).
- [35] M. Floris, *J. Phys. G* **38**, 124025 (2011).
- [36] B. Schenke, S. Jeon, and C. Gale, *Phys. Rev. Lett.* **106**, 042301 (2011).
- [37] H. Niemi, G. S. Denicol, H. Holopainen, and P. Huovinen, [arXiv:1212.1008](https://arxiv.org/abs/1212.1008).
- [38] M. Luzum and J.-Y. Ollitrault, [arXiv:1209.2323](https://arxiv.org/abs/1209.2323).
- [39] J. I. Kapusta, B. Muller, and M. Stephanov, *Phys. Rev. C* **85**, 054906 (2012).
- [40] R. P. G. Andrade, F. Grassi, Y. Hama, T. Kodama, and W. L. Qian, *Phys. Rev. Lett.* **101**, 112301 (2008).
- [41] K. Dusling, G. D. Moore, and D. Teaney, *Phys. Rev. C* **81**, 034907 (2010).
- [42] Y. Pandit (STAR Collaboration), [arXiv:1209.0244](https://arxiv.org/abs/1209.0244).
- [43] P. Bozek, W. Broniowski, and J. Moreira, *Phys. Rev. C* **83**, 034911 (2011).
- [44] K. Dusling, F. Gelis, T. Lappi, and R. Venugopalan, *Nucl. Phys. A* **836**, 159 (2010).
- [45] L. Pang, Q. Wang, and X.-N. Wang, *Phys. Rev. C* **86**, 024911 (2012).