

Application of the inverse Hamiltonian method to Hartree-Fock-Bogoliubov calculations

Y. Tanimura,¹ K. Hagino,¹ and P. Ring²¹*Department of Physics, Tohoku University, Sendai 980-8578, Japan*²*Physik Department, Technische Universität München, D-85748 Garching, Germany*

(Received 17 April 2013; published 8 July 2013)

We solve the Hartree-Fock-Bogoliubov (HFB) equations for a spherical mean field and a pairing potential with the inverse Hamiltonian method, which we have developed for the solution of the Dirac equation. This method is based on the variational principle for the inverse Hamiltonian and is applicable to Hamiltonians that are not bound from above or below. We demonstrate that the method works well not only for the Dirac, but also for the HFB equations.

DOI: [10.1103/PhysRevC.88.017301](https://doi.org/10.1103/PhysRevC.88.017301)

PACS number(s): 03.65.Ge, 21.10.Pc, 21.10.Gv, 21.60.Jz

Pairing correlations between nucleons play an important role in open shell nuclei [1,2]. Hartree-Fock-Bogoliubov (HFB) theory is a powerful method that treats these correlations in a self-consistent way in the framework of a single generalized Slater determinant of independent quasiparticles [2–4]. The method has been widely used in recent years for the study of the structure of neutron-rich nuclei far from stability up to the neutron drip line, where the coupling to the continuum has a strong influence.

The HFB equations are a set of coupled differential equations. In many cases they have been solved by an expansion in terms of a finite set of basis functions, as, for instance, the eigenfunctions of a harmonic oscillator or a Woods-Saxon potential. Although this method has been used successfully for many investigations in the literature it has its limitations. (i) The convergence with the number of basis functions depends on the parameters of the basis and the optimization of these parameters is often complicated; (ii) in many cases, in particular for two-dimensional (2D) and three-dimensional (3D) calculations in heavy nuclei, the dimension of the matrices becomes large. Since the CPU time for one diagonalization grows with the cube of this dimension this is connected with numerical efforts. (iii) In each step of the iteration, the corresponding matrix elements of the two-body interaction in terms of the basis functions requires, in addition, CPU time. (iv) The treatment of the continuum is connected with specific difficulties in particular in the case of neutron halos. In summary, these methods do not exploit fully the advantages of zero-range interactions as, for instance, the Skyrme energy density functional. These advantages can only be exploited fully in coordinate space. For this reason the Orsay group has introduced in the 1980s for the solution of the Hartree Fock (HF) equations the imaginary time method [5,6]. Starting with an initial HF wave function $|\Psi^{(0)}\rangle$ the solution is obtained in iterative steps with infinitesimal step size τ as

$$|\Psi^{(n+1)}\rangle \propto e^{-\tau\hat{H}}|\Psi^{(n)}\rangle. \quad (1)$$

Obviously this method is limited to Hamiltonians H with a spectrum bounded from below, as the nonrelativistic HF equation. An important fact is that the spectrum of the HFB equation is bound neither from above nor from below [see Fig. 1(a)] because of the coupling between particle

creation and annihilation parts in the quasiparticle operators of the Bogoliubov transformation [2]. This inhibits a direct application of the imaginary time method to HFB, which has been successfully employed in self-consistent mean field calculations in the coordinate space representation [6,7]. That is, if the imaginary time evolution is naively applied, the iterative solution inevitably dives into the quasiparticle negative continuum. To avoid this problem the two-basis method has been introduced in Refs. [7,8] where, in each step of the iteration, the HFB equations are solved by the expansion of the quasiparticle wave functions in an HF basis calculated by the imaginary time method on a 3D mesh in coordinate space. In Ref. [9], the HFB equation on a 3D mesh was solved in the canonical basis.

It should be mentioned that the diagonalization of the huge matrices in the basis expansion method can be avoided by the gradient method introduced in Ref. [10], which is also applicable to the solution of the HFB equation with a spectrum not bound from below. Here the wave function in the next step of the iteration is expressed in terms of the Thouless theorem [2]

$$|\Psi^{(n+1)}\rangle \propto \exp\left(-\tau \sum_{k < k'} H_{kk'}^{20} \alpha_k^\dagger \alpha_{k'}^\dagger\right) |\Psi^{(n)}\rangle, \quad (2)$$

where $|\Psi^{(n)}\rangle$ is the vacuum with respect to the quasiparticle operators α_k^\dagger . For infinitesimal τ only two-quasi-particle states with positive energy $E_k + E_{k'}$ are admixed in the next step of the iteration. This method has been successfully used in the literature for the solution of the HFB equations in an oscillator basis [11] and it has been also applied for a variation after projection (VAP) in Refs. [12,13]. Of course, the matrix elements $H_{kk'}^{20}$ have to be calculated here in the corresponding quasiparticle basis. Therefore the locality of the Skyrme interaction and the quasilocality of the kinetic energy in coordinate space cannot be exploited in this method.

Gradient methods have also the advantage in that constraints can be introduced easily. As discussed in Ref. [14], the calculation of the gradients can be restricted without difficulties to the hypersurfaces on the energy manifolds imposed by the gradients. Multidimensional constraints are needed for the evaluation of potential energy surfaces for




FIG. 1. (Color online) Spectra of (a) a quasiparticle Hamiltonian H itself and (b) the inverse of the Hamiltonian $1/(H - W)$. λ is the chemical potential. The bound states of positive and negative energies are indicated by solid and open circles, respectively. The continuum states are represented by the thick solid lines. The energy shift W is taken between the positive and negative spectra. The eigenvalues are labeled by an integer k such that $E_{-k} = -E_k$.

fission as well as for the application of methods going beyond the mean field approximation [2]. They lead to additional complications in methods based on a diagonalization in a basis.

We therefore propose in this report for the solution of the HFB equations a different method based on gradients keeping in mind that one is confronted with the same problem in the solution of the Dirac equation because the corresponding spectrum has the Dirac sea states down to the negative infinity as well as the positive energy states up to the positive infinity. This leads to a breakdown of variational calculations, which has long been known in the field of relativistic quantum chemistry under the name of “variational collapse,” and there has been a number of prescriptions proposed to avoid it [15–19]. Recently, newly developed methods for iterative solutions of a Dirac equation are introduced by Zhang *et al.* [20,21] and by Hagino and Tanimura [22] in the nuclear physics context. In Ref. [22], based on the idea of Hill and Krauthausser [18], a novel method has been developed, which is called the “inverse Hamiltonian method,” for relativistic mean field calculations in the coordinate space representation. In this method a variational principle is applied to the Hermitian operator $1/(H_{\text{Dirac}} - W)$ instead of the Hamiltonian H_{Dirac} itself. Here W is a real number which is set between the Fermi sea and the Dirac sea. In contrast to some other methods [16,17,20,21], it is relatively straightforward to apply our method not only to the Dirac equation but also to other eigenvalue problems with unbound operators, such as HFB. In this paper, we apply the inverse Hamiltonian method to an HFB equation in the coordinate space representation and show that the equation can be solved successfully without the variational collapse.

In HFB calculations one usually needs to obtain several lowest positive energy quasiparticle states only. That is, one only needs the states ψ_1, ψ_2, \dots associated with eigenvalues E_1, E_2, \dots [see Fig. 1(a)]. As is seen in Fig. 1(b), these states come to the top of the spectrum of $1/(H - W)$ if W is set between the positive and negative spectra. A variational principle for $1/(H - W)$ is that a maximization of $\langle (H - W)^{-1} \rangle$ leads to the desired quasiparticle state wave functions [18]. Our method maximizes $\langle (H - W)^{-1} \rangle$ based on

the relation [22]

$$|\psi^{(n+1)}\rangle \propto \exp\left(\frac{\Delta T}{H - W}\right) |\psi^{(n)}\rangle, \quad (3)$$

where $|\psi^{(0)}\rangle$ is an arbitrary wave function that is not an eigenfunction of the Hamiltonian H and W is a real constant between E_1 and E_{-1} . All the states below W damp out and only ψ_1 which is just above W survives in the limit $n \rightarrow \infty$.

In practice, the wave function $\psi^{(n+1)}$ is evolved for a small step ΔT from $\psi^{(n)}$ as

$$|\psi^{(n+1)}\rangle \propto \left(1 + \frac{\Delta T}{H - W}\right) |\psi^{(n)}\rangle = |\psi^{(n)}\rangle + \Delta T |\phi^{(n)}\rangle. \quad (4)$$

To this end, we have to solve a large sparse linear equation

$$(H - W)|\phi^{(n)}\rangle = |\psi^{(n)}\rangle \quad (5)$$

to invert the Hamiltonian. We here employ an iterative method for linear systems, that is, the conjugate gradient normal residual (CGNR) method [23]. This is one of the Krylov subspace methods for sparse linear systems [23,24]. CGNR solves a linear system $Ax = b$ by applying the conjugate gradient method to an equivalent system $A^\dagger Ax = A^\dagger b$.

When the mean field and pairing potentials are local and spherical, a quasi-particle wave function is given by the form

$$\psi_k(\mathbf{r}) = \frac{1}{r} \begin{pmatrix} U_k(r) \mathcal{Y}_{\ell j m}(\theta, \phi) \\ V_k(r) \mathcal{Y}_{\ell j m}(\theta, \phi) \end{pmatrix}. \quad (6)$$

Here $\mathcal{Y}_{\ell j m}$ is a spherical spinor defined by $\mathcal{Y}_{\ell j m} = \sum_{m, m'} (\ell m \frac{1}{2} m' | j m) Y_{\ell m} \chi_{m'}$, where $Y_{\ell m}$ and $\chi_{m'}$ are the spherical harmonics and spin wave function, respectively. The HFB equation in the coordinate space then reduces to a the radial equation

$$\begin{pmatrix} h - \lambda & \Delta(r) \\ \Delta(r) & -h + \lambda \end{pmatrix} \begin{pmatrix} U_k(r) \\ V_k(r) \end{pmatrix} = E_k \begin{pmatrix} U_k(r) \\ V_k(r) \end{pmatrix}, \quad (7)$$

where h is the mean field Hamiltonian, λ is the chemical potential, and $\Delta(r)$ is the pairing potential. Following the authors of Refs. [25,26], we use a phenomenological Woods-Saxon-type potentials, which simulates medium-heavy neutron-rich nuclei around ^{84}Ni , for the mean field and the pairing potentials. The potential $v(r)$ in the mean field Hamiltonian h and the pairing potential $\Delta(r)$ are thus taken as

$$v(r) = v_0 f(r) + v_{\ell s} \frac{1}{r} \frac{df}{dr} \boldsymbol{\ell} \cdot \mathbf{s}, \quad (8)$$

$$\Delta(r) = \Delta_0 f(r), \quad (9)$$

$$f(r) = \frac{1}{1 + e^{(r-R_0)/a}}, \quad (10)$$

with $v_0 = -38.5$ MeV, $v_{\ell s} = 14$ MeV \cdot fm 2 , $R_0 = 5.63$ fm, and $a = 0.66$ fm [25,26]. The strength of the pairing potential Δ_0 is determined so that the average pairing gap $\bar{\Delta}$ defined by [25]

$$\bar{\Delta} = \frac{\int_0^\infty r^2 dr \Delta(r) f(r)}{\int_0^\infty r^2 dr f(r)} \quad (11)$$

is equal to 1.0 MeV. The chemical potential λ is fixed to $\lambda = -0.5$ MeV in the present calculation. We solve Eq. (7) by discretizing the radial coordinate r with mesh size Δr , and

imposing the box boundary condition. The second derivative of ψ at the i th mesh point is approximated by a three-point difference: $\psi_i'' = (\psi_{i+1} - 2\psi_i + \psi_{i-1})/(\Delta r)^2$.

Let us now apply the inverse Hamiltonian method and numerically solve the HFB equation, Eq. (7). We also solve the equation exactly by directly diagonalizing the coordinate space Hamiltonian. The parameters of the inverse Hamiltonian method are set $W = 0.1$ MeV and $\Delta T = 10$ MeV. The excited states are also calculated simultaneously by orthogonalizing a set of wave functions at every step of the iteration. The radial coordinate is discretized up to $r_{\max} = 30$ fm with $\Delta r = 0.1$ fm. The initial quasiparticle wave functions are taken to be a Gaussian form

$$\begin{pmatrix} U_k^{(0)}(r) \\ V_k^{(0)}(r) \end{pmatrix} = N_k \begin{pmatrix} r^{\ell+1} e^{-r^2/b_k^2} \\ r^{\ell+1} e^{-r^2/b_k^2} \end{pmatrix}, \quad (12)$$

where ℓ is the orbital angular momentum and N_k is an appropriate normalizing constant. The width parameter of the Gaussian b_k is taken as $b_k = 2.0 \times 1.05^{k-1}$ fm, ($k = 1, 2, \dots$).

Let us first discuss the convergence properties of the energy $\langle H \rangle$ and the expectation value of the inverse of Hamiltonian $\langle (H - W)^{-1} \rangle$ for the lowest $s_{1/2}$ quasiparticle state. In Fig. 2 we show the evolution of the two quantities as functions of the number of iteration steps. As was observed in Ref. [22] for a Dirac equation, $\langle (H - W)^{-1} \rangle$ converges monotonically up to a certain value as the iteration step increases. At the same time, $\langle H \rangle$ converges to the lowest $s_{1/2}$ eigenvalue $E = 0.424$ MeV.

In Table I, we show quasiparticle energies and occupation probabilities v_k^2 for the three lowest $s_{1/2}$ states in comparison to the exact values which are obtained by diagonalizing the Hamiltonian. The occupation probabilities are defined in terms of quasiparticle wave function by

$$v_k^2 = \int_0^\infty dr |V_k(r)|^2. \quad (13)$$




FIG. 2. (Color online) Convergence properties of (a) the energy expectation value $\langle H \rangle$ and (b) the expectation value of the inverse of Hamiltonian $\langle (H - W)^{-1} \rangle$ for the lowest $s_{1/2}$ quasiparticle state. The energy shift and the step size of T are taken to be $W = 0.1$ MeV and $\Delta T = 10$ MeV, respectively.

TABLE I. A comparison between the exact calculations and the inverse Hamiltonian method for the three lowest $s_{1/2}$ quasiparticle energies E and occupation probabilities v_k^2 . The exact values are calculated by diagonalizing the real space Hamiltonian.

E (MeV)		v_k^2	
Exact	Inv. H method	Exact	Inv. H method
0.42414	0.42414	0.5574	0.5574
1.0383	1.0383	3.972×10^{-2}	3.972×10^{-2}
2.3063	2.3063	9.689×10^{-3}	9.689×10^{-3}

The agreement is perfect both in the energies and the occupation probabilities for the digits shown in the table. Figure 3 shows comparisons of the wave functions of the three $s_{1/2}$ states. The dashed lines show the exact wave functions, whereas the solid lines show the wave functions obtained with the inverse Hamiltonian method. The left and right panels show the upper component $U_k(r)$ and the lower component $V_k(r)$ of a quasiparticle wave function, respectively. As seen in Fig. 3, the inverse Hamiltonian method reproduces the wave functions almost identically to the exact ones for both the bound state and the excited continuum states. We also obtained the other s -wave states with an accuracy as high as the lower states shown in Table I and Fig. 3.

We checked the performance of the inverse Hamiltonian method for other angular momentum quantum numbers and




FIG. 3. (Color online) Comparisons of wave functions for the three lowest $s_{1/2}$ states. Figures 3(a), 3(c), and 3(e) show the upper components $[U_k(r)]$ while Figs. 3(b), 3(d), and 3(f) show the lower components $[V_k(r)]$ of the quasiparticle wave function, respectively. The exact wave functions are shown with the dashed lines and the ones obtained by the inverse Hamiltonian method are drawn with the solid lines.

confirmed that the method solves the HFB equation as accurately as for the $s_{1/2}$ states. It is apparent that the inverse Hamiltonian method gives practically the exact solutions of the HFB equation in the coordinate space representation and is safe against the variational collapse.

In summary, we have discussed the numerical performance of the inverse Hamiltonian method for an HFB calculation. While the method has been developed for solving Dirac equations, we have shown that it can almost exactly solve a coordinate space HFB equation as well with spherical mean field and pairing potentials without variational collapse. This indicates that the inverse Hamiltonian method provides an alternative coordinate space method for 3D HFB calculations, in addition to the existing methods such as the two-basis

method and the canonical basis method. The inverse Hamiltonian method has an advantage in that it can also be applied in a straightforward manner to relativistic Bogoliubov calculations on 3D mesh, for which the Dirac sea spectrum in a mean-field Hamiltonian prevents a direct application of the two-basis method in combination with the imaginary time method. An obvious future work is the application of the method to self-consistent HFB calculations on 3D mesh. A work in this direction is now in progress.

This work was supported by a grant-in-aid by JSPS under Program No. 24-3429 and the Japanese Ministry of Education, Culture, Sports, Science, and Technology by a grant-in-aid for scientific research under Program No. (C) 22540262.

-
- [1] D. M. Brink and R. A. Broglia, *The Nuclear Superfluidity Pairing in Finite Systems* (Cambridge University Press, Cambridge, England, 2005).
- [2] P. Ring and P. Schuck, *The Nuclear Many-Body Problem* (Springer-Verlag, Berlin, 1980).
- [3] M. Bender, P.-H. Heenen, and P.-G. Reinhard, *Rev. Mod. Phys.* **75**, 121 (2003).
- [4] J. Dobaczewski, W. Nazarewicz, T. R. Werner, J. F. Berger, C. R. Chinn, and J. Dechargé, *Phys. Rev. C* **53**, 2809 (1996).
- [5] K. T. R. Davies, H. Flocard, S. Krieger, and M. S. Weiss, *Nucl. Phys. A* **342**, 111 (1980).
- [6] P. Bonche, H. Flocard, and P.-H. Heenen, *Comput. Phys. Commun.* **171**, 49 (2005).
- [7] P. Bonche, H. Flocard, P.-H. Heenen, S. J. Krieger, and M. S. Weiss, *Nucl. Phys. A* **443**, 39 (1985).
- [8] J. Terasaki, P.-H. Heenen, H. Flocard, and P. Bonche, *Nucl. Phys. A* **600**, 371 (1996).
- [9] N. Tajima, *Phys. Rev. C* **69**, 034305 (2004).
- [10] H.-J. Mang, B. Samadi, and P. Ring, *Z. Phys. A* **279**, 325 (1976).
- [11] J. L. Egido, J. Lessing, V. Martin, and L. M. Robledo, *Nucl. Phys. A* **594**, 70 (1995).
- [12] J. L. Egido and P. Ring, *Nucl. Phys. A* **388**, 19 (1982).
- [13] M. Anguiano, J. L. Egido, and L. M. Robledo, *Nucl. Phys. A* **696**, 70 (2001).
- [14] J. L. Egido, H.-J. Mang, and P. Ring, *Nucl. Phys. A* **334**, 1 (1980).
- [15] H. Wallmeier and W. Kutzelnigg, *Chem. Phys. Lett.* **78**, 341 (1981).
- [16] H. Wallmeier and W. Kutzelnigg, *Phys. Rev. A* **28**, 3092 (1983).
- [17] R. E. Stanton and S. Havriliak, *J. Chem. Phys.* **81**, 1910 (1984).
- [18] R. N. Hill and C. Krauthauser, *Phys. Rev. Lett.* **72**, 2151 (1994).
- [19] P. Falsaperla, G. Fonte, and J. Z. Chen, *Phys. Rev. A* **56**, 1240 (1997).
- [20] Y. Zhang, H. Liang, and J. Meng, *Int. J. Mod. Phys. E* **19**, 55 (2010).
- [21] Y. Zhang, H. Liang, and J. Meng, *Chin. Phys. Lett.* **26**, 092401 (2009).
- [22] K. Hagino and Y. Tanimura, *Phys. Rev. C* **82**, 057301 (2010).
- [23] Y. Saad, *Iterative Methods for Sparse Linear Systems* (Society for Industrial and Applied Mathematics, Philadelphia, PA, 2003).
- [24] H. A. van der Vorst, *Iterative Krylov Methods for Large Linear Systems* (Cambridge University Press, Cambridge, England, 2003).
- [25] M. Kohno, Y. Fujiwara, and Y. Akaishi, *Phys. Rev. C* **68**, 034302 (2003); I. Hamamoto and B. R. Mottelson, *ibid.* **69**, 064302 (2004).
- [26] K. Hagino and H. Sagawa, *Phys. Rev. C* **71**, 044302 (2005).