

Electrical conductivity of the quark-gluon plasma and soft photon spectrum in heavy-ion collisions

Yi Yin*

Physics Department, University of Illinois at Chicago, Chicago, Illinois 60607, USA

and Physics Department, Brookhaven National Laboratory, Upton, New York 11973, USA

(Received 24 December 2013; revised manuscript received 24 August 2014; published 6 October 2014)

I extract the electrical conductivity σ_0 of the quark gluon plasma (QGP) and study the effects of magnetic field and chiral anomaly on soft photon azimuthal anisotropy, v_2 , based on the thermal photon spectrum at $0.4 \text{ GeV} < p_\perp < 0.6 \text{ GeV}$ at the Brookhaven Relativistic Heavy Ion Collider energy. As a basis for my analysis, I derive the behavior of retarded photon self-energy of a strongly interacting neutral plasma in hydrodynamic regime in the presence of magnetic field and chiral anomaly. By evolving the resulting soft thermal photon production rate over the realistic hydrodynamic background and comparing the results with the data from the PHENIX Collaboration, I found that the electrical conductivity at QGP temperature is in the range: $0.4 < \sigma_0/(e^2 T) < 1.1$, which is comparable with recent studies on lattice. I also compare the contribution from the magnetic field and chiral anomaly to soft thermal photon v_2 with the data. I argue that at the CERN Large Hadron Collider, the chiral magnetic wave would give negative contribution to photon v_2 .

DOI: 10.1103/PhysRevC.90.044903

PACS number(s): 25.75.Cj

I. INTRODUCTION

Photons produced in heavy-ion collisions contain rich information on the properties of quark-gluon plasma (QGP). The number of photons emitted per unit time per unit volume, from a plasma in thermal equilibrium, to leading order in α_{EM} , is given by [1]

$$\omega \frac{d\Gamma_\gamma}{d^3\mathbf{p}} = -\frac{1}{(2\pi)^3} \frac{\text{Im}[P_T^{ij} G_{ij}^{R,\text{EM}}(\omega, \mathbf{p})]}{e^{\omega/T} - 1} \Big|_{\omega=|\mathbf{p}|}. \quad (1)$$

where $P_T^{ij} \equiv \delta^{ij} - \hat{\mathbf{p}}^i \hat{\mathbf{p}}^j$ is the projection operator. Here $G_{R,\text{EM}}^{ij}(\omega, \mathbf{p})$ (i.e., retarded photon self-energy) denotes the retarded Green's function of the charge current operator $J_\mu^{\text{EM}} \equiv \sum_f q_f \bar{\psi}^f \gamma_\mu \psi^f$. On the other hand, the low energy and low momentum behavior of $G_{R,\text{EM}}^{ij}(\omega, \mathbf{p})$ for any interacting system in the thermal equilibrium is completely fixed by hydrodynamics. Indeed, if the conductivity tensor of the system is isotropic, i.e., $\sigma^{ij} = \sigma_0 \delta^{ij}$, the thermal emission rate of soft photons is fully parametrized by σ_0 :

$$\omega \frac{d\Gamma_\gamma}{d^3\mathbf{p}} \Big|_{\omega=|\mathbf{p}|} = \frac{\alpha_{\text{EM}}}{\pi^2 e^2} \frac{\sigma_0 \omega}{e^{\omega/T} - 1}. \quad (2)$$

Recently, results of low p_\perp direct thermal photon spectrum at the Brookhaven Relativistic Heavy Ion Collider (RHIC) have been reported by PHENIX Collaboration [2,3]. The lowest p_\perp bin in those results is $0.4 \text{ GeV} \leq p_\perp \leq 0.6 \text{ GeV}$. It is now well accepted that a near perfect fluid is created in heavy-ion collisions. The smallness of η/s as inferred from relativistic hydrodynamic simulations implies that sQGP enjoys a wider hydrodynamic regime, to the order of πT . Therefore, for photon produced at energy $0.4 \text{ GeV} \leq p_\perp \leq 0.6 \text{ GeV}$, the hydrodynamic expression, e.g., Eq. (2), does apply. I then could use Eq. (2) to extract σ_0 . By evolving Eq. (2) with the temperature-flow background as generated by

solutions of relativistic hydrodynamic equations (cf. Sec. III), I found at typical QGP temperature (cf. Fig. 1)

$$0.4 < \frac{\sigma}{e^2 T} < 1.1. \quad (3)$$

To the extent of my knowledge, this is the first direct estimation of the electrical conductivity of QGP based on soft photon production with realistic hydrodynamics simulation.¹

On the other hand, Eq. (2) indicates that photon azimuthal anisotropy, v_2 should be small in low p_\perp regime as the effects due to the background elliptic flow are highly suppressed. However, the PHENIX results on soft photon v_2 [3] suggest that direct photon v_2 does not tend to vanish at low p_\perp limit but saturates at some positive value (cf. Fig. 2). The nonzero soft photon v_2 implies that Eq. (2) *does* receive sizable modifications in QGP.²

One possible source of such modifications is the magnetic field created by the spectator charges of ultrarelativistic heavy ions which can be as large as $eB \sim m_\pi^2$, and it points to the perpendicular direction of the reaction plane [7]. In Ref. [8], the effects of magnetic field were considered to explain photon v_2 at $p_\perp > 1 \text{ GeV}$ as measured by the PHENIX Collaboration [9]. In the present paper, I will study the effects of magnetic field and chiral anomaly on soft photon v_2 . As a basis for my analysis, I will derive the behavior of retarded Green's function

¹The conductivity can be related to the diffusive constant via Einstein relation. The heavy quark diffusive constant in QGP was studied in Ref. [4] based on charm spectrum R_{AA} and charm elliptic flow. Recently, there is encouraging progress on constraining light quark diffusive constant of QGP at crossover regime by applying fluctuating hydrodynamics in Bjorken expansion to the study of charge density fluctuations in QCD matter [5]. However, I am unaware of any work on directly extracting the conductivity and light quark diffusive constant with the realistic hydrodynamic simulation.

²The corrections due to nonequilibrium would also contribute to soft photon v_2 . However, according to the simulation of Ref. [6], the resulting soft photon v_2 is of the order $0.01 \sim 0.02$.

*yyin@bnl.gov




FIG. 1. (Color online) $\langle \sigma/e^2 T \rangle_{QGP}$ (red dots), the average of $\sigma/e^2 T$ in QGP as computed from Eq. (26) using photon production at p_\perp bin $0.4 \text{ GeV} < p_\perp < 0.6 \text{ GeV}$ for three different centrality bins (0–20 %, 20–40 %, 40–60 %) [2]. Blue dotted lines are corresponding to the upper and lower bound for $\sigma/e^2 T$ as quoted in the abstract and in Eq. (3). Dashed horizontal lines are corresponding to the range of $\sigma/(e^2 T)$ as estimated by lattice simulation in Ref. [18] with $C_{EM} = (2/3)e^2$.

$G_{ij}^R(\omega, \mathbf{p})$ in the hydrodynamic regime in a neutral strongly coupled plasma in the presence of homogeneous magnetic field and chiral anomaly. As triangle anomaly leads to additional terms in the constitute relation of hydrodynamics [10], the resulting $G_{ij}^R(\omega, \mathbf{p})$ has a much richer structure. This opens the possibility to distinguish the effects of chiral anomaly to photon v_2 . Furthermore, as it is not unexpected that the magnetic field will give positive contribution to photon v_2 , the phenomenologically important question is how sizable the effects of magnetic field are at heavy-ion collisions. To answer this question, a realistic hydrodynamic simulation of photon production is needed. By evolving the modified soft photon rate in realistic hydrodynamic background, I found that if the life time of magnetic field $\tau_B > 2 \sim 3 \text{ fm}$, the contribution due to magnetic field to the soft photon v_2 is comparable to the experiment results.

This paper is organized as follows. Section II presents the derivation of the behavior of retarded Green's function $G_{ij}^R(\omega, \mathbf{p})$ in the hydrodynamic region in the presence of homogeneous magnetic field and chiral anomaly. Results are summarized in Eqs. (14) and (15). Though they are the direct consequences of the constitute relation of anomalous hydrodynamics and the linear response theory, to the best of my knowledge, Eq. (15) is new in literature. In Sec. III, I extract the electrical conductivity with the realistic hydrodynamic evolution. The results are comparable with recent lattice measurement. In Sec. IV, I investigate the relation between magnetic field and photon v_2 . I summarize and conclude in Sec. V.

II. RETARDED GREEN'S FUNCTION IN THE HYDRODYNAMIC REGIME AND SOFT PHOTON PRODUCTION

In this section, I will work out explicitly the behavior of the retarded Green's function $G_{ij}^R(\omega, \mathbf{p})$ of a neutral ($n_{V,A} = 0$) strongly coupled plasma in the presence of a homogeneous magnetic field \mathbf{B}_0 and chiral anomaly in the hydrodynamic region. For that purpose, it is sufficient to consider a plasma with only one flavor with EM charge q^f . I start with the constitute relation for the spatial part of the vector current $J_V^\mu = q_f^{-1} J_{EM}^\mu$ and axial current J_A^μ in a static and homogeneous flow background [10]:

$$J_V^l = C_{anom} \mu_A B^l + q_f^{-1} \sigma^{lm} E_m - D^{lm} \nabla_m n_V, \quad (4a)$$

$$J_A^l = C_{anom} \mu_V B^l - D^{lm} \nabla_m n_A, \quad (4b)$$

where $l, m = 1, 2, 3$ run over spatial components and μ_V, μ_A denote the chemical potentials of vector charge and axial charge, respectively. In Eq. (4), the anomaly coefficient

$$C_{anom} = \frac{q_f N_c}{2\pi^2} \quad (5)$$




FIG. 2. (Color online) PHENIX results of direct photon v_2 [3]. (Figure is reproduced from Ref. [26].)

is defined through the divergence of the axial current

$$\partial_\mu j_A^\mu = C_{\text{anom}} q_f \mathbf{E} \cdot \mathbf{B}. \quad (6)$$

$C_{\text{anom}} \mu_{A,V} \mathbf{B}$ terms in Eq. (4) are completely induced by the chiral anomaly [10] and are directly related to chiral magnetic effects and charge separation effects [7,11] (see Ref. [12] for a recent review). σ^{ij}, D^{ij} in Eq. (4) are the conductivity and diffusive tensors in the presence of magnetic field \mathbf{B}_0 , respectively, and are related by the Einstein relation

$q_f^2 \chi D^{ij} = \sigma^{ij}$, where χ is the susceptibility. Due to external magnetic field \mathbf{B}_0 , σ^{ij}, D^{ij} , in general, is anisotropic [cf. Eq. (12)].

To determine the behavior of $G_{ij}^R(\omega, \mathbf{p})$ in the hydrodynamic region through the linear response theory, I now perturb the system by imposing a space-time dependent vector potential $\delta \mathbf{A} \propto e^{-i\omega t + i\mathbf{p} \cdot \mathbf{x}}$. Due to Eq. (4) and $\mathbf{E} = -\partial_t \mathbf{A}$, for a neutral plasma, the change of current $\delta \mathbf{j}_{V,A}$ in response to $\delta \mathbf{A}$ now reads

$$\begin{pmatrix} \delta j_V^l(\omega, \mathbf{p}) \\ \delta j_A^l(\omega, \mathbf{p}) \end{pmatrix} = (-ip_m) \begin{pmatrix} D^{ml} & D_5^{ml} \\ D_5^{ml} & D^{ml} \end{pmatrix} \begin{pmatrix} \delta n_V(\omega, \mathbf{p}) \\ \delta n_A(\omega, \mathbf{p}) \end{pmatrix} + i\omega \begin{pmatrix} \sigma^{lm} \\ 0 \end{pmatrix} q_f^{-1} \delta A_m(\omega, \mathbf{p}). \quad (7)$$

Here I relate $\delta \mu_{V,A}, \delta n_{V,A}$ by $\chi \delta \mu_{V,A} = \delta n_{V,A}$, where χ is the susceptibility. I expect that in the chirally symmetric phase, the susceptibility for the axial charge and the vector charge are approximately identical. For future convenience, I have also introduced

$$D_5^{ij} \equiv \frac{iv_\chi}{p} \hat{\mathbf{p}}^i \hat{\mathbf{B}}_0^j, \quad \sigma_5^{ij} \equiv q_f^2 \chi D_5^{ij} = \frac{iC_{\text{anom}} B_0}{p} \hat{\mathbf{p}}^i \hat{\mathbf{B}}_0^j, \quad (8)$$

where the speed of chiral magnetic wave [13] reads

$$v_\chi \equiv \frac{C_{\text{anom}} B_0}{\chi}. \quad (9)$$

From the anomaly equation (6) and conservation of vector charge $\partial_\mu j_V^\mu = 0$, I also have

$$\begin{pmatrix} \omega + ip_l p_m D^{ml} & ip_l p_m D_5^{ml} \\ ip_l p_m D_5^{ml} & \omega + ip_l p_m D^{ml} \end{pmatrix} \begin{pmatrix} \delta n_V(\omega, \mathbf{p}) \\ \delta n_A(\omega, \mathbf{p}) \end{pmatrix} = i\omega p_l \begin{pmatrix} \sigma^{lm} \\ \sigma_5^{lm} \end{pmatrix} q_f^{-1} \delta A_m. \quad (10)$$

Now solving for $\delta n_{V,A}$ in terms of $\delta \mathbf{A}$ in Eq. (10) and put them back in the expression of \mathbf{j}_V in Eq. (7), one arrives at

$$\delta j_V^l = i\omega \left[(p_r D^{rl}, p_r D_5^{rl}) \begin{pmatrix} i\omega - p_i p_j D^{ij} & -p_i p_j D_5^{ij} \\ -p_i p_j D_5^{ij} & i\omega - p_i p_j D^{ij} \end{pmatrix}^{-1} \begin{pmatrix} p_q \sigma^{qm} \\ p_q \sigma_5^{qm} \end{pmatrix} + \sigma^{lm} \right] q_f^{-1} \delta A_m. \quad (11)$$

The tensor σ^{ij} in the presence of \mathbf{B}_0 may be decomposed as [14]

$$\sigma^{ij} = \sigma_0 \delta^{ij} - \sigma_{B,T} (\delta^{ij} - \hat{\mathbf{B}}_0^i \hat{\mathbf{B}}_0^j) + \sigma_{B,L} \hat{\mathbf{B}}_0^i \hat{\mathbf{B}}_0^j = q_f^2 \chi [D_0 \delta^{ij} - D_{B,T} (\delta^{ij} - \hat{\mathbf{B}}_0^i \hat{\mathbf{B}}_0^j) + D_{B,L} \hat{\mathbf{B}}_0^i \hat{\mathbf{B}}_0^j], \quad (12)$$

where $\hat{\mathbf{B}}_0$ is the directional vector of \mathbf{B}_0 . Here σ_0 denotes the conductivity in the absence of magnetic field, $\sigma_{B,L}, \sigma_{B,T}$ denote the change of conductivity in the longitudinal and transverse direction of magnetic field \mathbf{B}_0 .³ According to linear response theory,

$$\langle J_V^\mu(\omega, \mathbf{p}) \rangle_A = -G_{R,\text{EM}}^{\mu\nu}(\omega, \mathbf{p}) (q_f^{-1} A_\nu), \quad (13)$$

I then obtain

$$G_{R,\text{EM}}^{ij}(\omega, \mathbf{p}) = F_L(\omega, \mathbf{p}) \hat{\mathbf{p}}^i \hat{\mathbf{p}}^j + F_T(\omega, \mathbf{p}) P_T^{ij} + F_{pB}(\omega, \mathbf{p}) (\hat{\mathbf{p}}^i \hat{\mathbf{B}}_0^j + \hat{\mathbf{p}}^j \hat{\mathbf{B}}_0^i) + F_{BB}(\omega, \mathbf{p}) \hat{\mathbf{B}}_0^i \hat{\mathbf{B}}_0^j, \quad (14)$$

where form factors in Eq. (14) are given by

$$F_T = -i\omega(\sigma_0 - \sigma_{B,T}), \quad (15a)$$

$$F_L = \frac{-i\omega(\sigma_0 - \sigma_{B,T})}{2} \left\{ \frac{\omega - v_\chi p(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0) + ip^2(D_{B,T} + D_{B,L})(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2}{\Omega(\omega, \mathbf{p}, \hat{\mathbf{B}}_0)} + (\hat{\mathbf{B}}_0 \rightarrow -\hat{\mathbf{B}}_0) \right\}, \quad (15b)$$

$$F_{pB} = \frac{-i\omega\sigma_0 - \sigma_{B,T}}{2} \left\{ \frac{v_\chi p - ip^2(D_{B,T} + D_{B,L})(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2}{\Omega(\omega, \mathbf{p}, \hat{\mathbf{B}}_0)} + (\hat{\mathbf{B}}_0 \rightarrow -\hat{\mathbf{B}}_0) \right\}, \quad (15c)$$

$$F_{BB} = -\frac{i\omega}{2} \left\{ \frac{iv_\chi^2 \chi + (\sigma_{B,T} + \sigma_{B,L})[\omega + v_\chi p(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0) + ip^2(D_0 - D_{B,T})]}{\Omega(\omega, \mathbf{p}, \hat{\mathbf{B}}_0)} + (\hat{\mathbf{B}}_0 \rightarrow -\hat{\mathbf{B}}_0) \right\}, \quad (15d)$$

³If the Hall conductivity σ_H is nonzero, one could add an additional term $i\sigma_H \epsilon_{ijk} \hat{\mathbf{B}}_0^k$ to σ_{ij} .

where

$$\Omega(\omega, \mathbf{p}, \hat{\mathbf{B}}_0) = \omega - v_\chi p(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0) + ip^2[(D_0 - D_{B,T}) + (D_{B,T} + D_{B,L})(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2]. \quad (16)$$

In Eq. (15), the contributions due to right-handed chiral fermions have been written down explicitly in the brackets $\{\dots\}$ while those due to left-handed chiral fermions are easily obtained by replacing $\hat{\mathbf{B}}_0$ with $-\hat{\mathbf{B}}_0$ as denoted by $(\hat{\mathbf{B}}_0 \rightarrow -\hat{\mathbf{B}}_0)$ in Eq. (15).

As one can check, in the absence of \mathbf{B}_0 that $\sigma_{B,T}, \sigma_{B,L}, v_\chi = 0, F_{BB}, F_{pB}$ vanish and I recover the well-known results

$$\begin{aligned} \lim_{\mathbf{B}_0 \rightarrow 0} F_T(\omega, p) &= -i\omega\sigma_0, \\ \lim_{\mathbf{B}_0 \rightarrow 0} F_L(\omega, p) &= \frac{-i\omega^2\sigma_0}{\omega + iD_0p^2}. \end{aligned} \quad (17)$$

Returning to Eqs. (14) and (15), I see immediately that $G_{ij}^R(\omega, \mathbf{p})$ has poles when $\Omega(\omega, \mathbf{p}, \pm \hat{\mathbf{B}}_0) = 0$. The corresponding dispersion relation is

$$\begin{aligned} \omega(\mathbf{p}) &= \pm v_\chi p(\hat{\mathbf{B}}_0 \cdot \hat{\mathbf{p}}) - ip^2 \\ &\times [(D_0 - D_{B,T}) + (D_{B,L} + D_{B,T})(\hat{\mathbf{B}}_0 \cdot \hat{\mathbf{p}})^2]. \end{aligned} \quad (18)$$

For $\hat{\mathbf{B}}_0 \cdot \hat{\mathbf{p}} = 0$, Eq. (18) describes the conventional diffusive modes while for $\hat{\mathbf{B}}_0 \cdot \hat{\mathbf{p}} \neq 0$, Eq. (18) describes a propagating hydrodynamical mode, namely, chiral magnetic wave [13]. I point out here that due to chiral magnetic wave poles, zero frequency limit and zero momentum limit of $G_{R,EM}^{ij}(\omega, \mathbf{p})$ may not commute with each other. Special care may be needed when apply Kubo formula to a plasma in the presence of magnetic field and chiral anomaly.

To determine photon production rate in the hydrodynamic region, I only need to know the imaginary part of $F_T(\omega, \mathbf{p}), F_{BB}(\omega, \mathbf{p})$ along the light-cone $\omega = |\mathbf{p}|$:

$$\begin{aligned} \text{Im}(P_T^{ij} G_{ij}^R(\omega, \mathbf{p}))|_{\omega=|\mathbf{p}|} \\ = 2\text{Im}[F_T(\omega = |\mathbf{p}|)] + \text{Im}[F_{BB}(\omega = |\mathbf{p}|)][1 - (\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2]. \end{aligned} \quad (19)$$

Keeping terms of the lowest order in ω in Eq. (19), I then have, in the presence of magnetic field and chiral anomaly, that

$$\begin{aligned} \text{Im}(P_T^{ij} G_{ij}^R(\omega, \mathbf{p}))|_{\omega=|\mathbf{p}|} \\ = \frac{\alpha_{EM}}{2\pi^2 e^2} \left\{ (2\sigma_0 - \sigma_{B,T}) + (1 - (\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2 \right. \\ \left. \times [(\sigma_{B,T} + \sigma_{B,L}) + v_\chi^2(\sigma_0 - \sigma_{B,T})] \frac{1 + v_\chi^2(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2}{(1 - v_\chi^2(\hat{\mathbf{p}} \cdot \hat{\mathbf{B}}_0)^2)} \right\} \end{aligned} \quad (20)$$

and in the absence of \mathbf{B} , we recover Eq. (2).

In the following sections, I will use Eqs. (2) and (20) to study soft photon production in heavy-ion collisions.

III. ELECTRICAL CONDUCTIVITY OF QGP

The thermal photon momentum spectrum produced during the evolution of the radiating fireball can be written as

$$\omega \frac{dN_\gamma}{d^3\mathbf{p}} = \int d^4x \omega_{\text{shift}} \frac{d\Gamma_\gamma}{d^3\mathbf{p}'} \Big|_{\omega_{\text{shift}}=|\mathbf{p}'|}, \quad (21)$$

where the photon energy, which is ω in the laboratory frame, is red-shifted to ω_{shift} in the frame that fluid is at rest:

$$\begin{aligned} \omega_{\text{shift}} &= p^\mu u_\mu \\ &= \omega u^\tau [\cosh(\xi - Y) - v_x \cos \phi_p - v_y \sin \phi_p]. \end{aligned} \quad (22)$$

Here, assuming the boost-invariance, the four-velocity of the flow field is $u^\mu = (u^t, u^z, u^x, u^y) = u^\tau (\cosh \xi, \sinh \xi, v_x, v_y)$. I use Bjorken's coordinates τ, ξ, x, y , with $\tau = \sqrt{t^2 - z^2}$ the longitudinal proper time and $\xi = \tanh^{-1}(z/t)$ the space-time rapidity that $d^4x = \tau d\tau d\xi dx dy$. The photon momentum is parametrized by its rapidity Y , transverse momentum p_\perp , and azimuthal emission angle ϕ_p , i.e., $p^\mu = p_\perp (\cosh Y, \sinh Y, \cos \phi_p, \sin \phi_p)$.

In the present section, I will estimate the value of the electrical conductivity by neglecting possible modifications due to the magnetic field. I will return to the effects of magnetic field in the next section. For the soft photon production at heavy-ion collisions, I have from Eq. (2) that

$$\frac{dN_\gamma}{d^2p_\perp dY} = \frac{\alpha_{EM}}{\pi^2} \int d^4x \left(\frac{\sigma_0}{e^2 T} \right) \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1}. \quad (23)$$

I will concentrate on the photoproductions at midrapidity $Y = 0$ and expand the photon production in Fourier Harmonics:

$$\frac{dN_\gamma}{d^2p_\perp dY} = \frac{dN_\gamma}{2\pi p_\perp dp_\perp dY} [1 + 2v_2(p_\perp) \cos(2\phi_p) + \dots]. \quad (24)$$

I therefore have

$$\frac{dN_\gamma}{2\pi p_\perp dp_\perp dY} = \frac{\alpha_{EM}}{\pi^2} \int_0^{2\pi} \frac{d\phi_p}{2\pi} \int d^4x \left(\frac{\sigma}{e^2 T} \right) \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1}. \quad (25)$$

Now introducing the dimensionless quantity:

$$\begin{aligned} \left\langle \frac{\sigma}{e^2 T} \right\rangle_{\text{QGP}} &\equiv \frac{\frac{\alpha_{EM}}{\pi^2} \int_0^{2\pi} \frac{d\phi_p}{2\pi} \int d^4x \left(\frac{\sigma}{e^2 T} \right) \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1}}{\frac{\alpha_{EM}}{\pi^2} \int_0^{2\pi} \frac{d\phi_p}{2\pi} \int d^4x \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1}} \\ &= \frac{\left(\frac{dN_\gamma}{2\pi p_\perp dp_\perp dY} \right)_{\text{data}}}{\frac{\alpha_{EM}}{\pi^2} \int_0^{2\pi} \frac{d\phi_p}{2\pi} \int d^4x \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1}}. \end{aligned} \quad (26)$$

As the conductivity in hadronic phase is much smaller than that in QGP state due to the reduction of the charge carriers in the medium, $\langle \sigma/e^2 T \rangle_{\text{QGP}}$ provides us an estimation of $\sigma/e^2 T$ at typical QGP temperature.

In Fig. 1, I show $\langle \sigma/e^2 T \rangle_{\text{QGP}}$, the average $\sigma/(e^2 T)$ in QGP as defined by Eq. (26). The direct photon production data (after subtraction of hard-scattering component) are taken from results by the PHENIX Collaboration [2] at p_\perp bin $0.4 \text{ GeV} < p_\perp < 0.6 \text{ GeV}$ in $\sqrt{s_{NN}} = 200 \text{ GeV}$ Au + Au

collisions. The denominator of the last term of Eq. (26) is evaluated at $p_\perp = 0.5$ GeV using realistic hydrodynamic background. To model such background, I employ results computed with “VISH2+1”, a viscous hydrocode, developed by Song and Heinz [15], in $2 + 1$ dimensions assuming longitudinal boost invariance. Those simulations, which reproduce hadron spectrum in the experiment well, were performed by Shen [16] and the results are accessible to the public via the website <https://wiki.bnl.gov/TECHQM/index.php>. The hydrodynamic evolution starts at $\tau_i = 0.4$ fm and ends on an isothermal surface at $T_{\text{dec}} = 130$ MeV with $\eta/s = 0.20$ and the lattice-based equation of state “s95p-PCE” [16,17].

I have performed my analysis for three different centrality bins: 0–20%, 20–40%, 40–60%.⁴ As the conductivity σ_0 reflects the transport properties of QGP and the effective temperature as extracted from thermal photon spectrum for those three centrality bins are similar [2], we expect that $\langle\sigma/(e^2T)\rangle_{\text{QGP}}$ would have a weak dependence on centrality. This is indeed the case as one can see in Fig. 1: while both soft photon production and hydrodynamic backgrounds are different for those centrality bins, the resulting $\langle\sigma/(e^2T)\rangle_{\text{QGP}}$ shows little dependence on the centralities.

The error bars shown in Fig. 1 are determined from the experimental (systematic) uncertainties in the photon production as $\langle\sigma/(e^2T)\rangle_{\text{QGP}}$ given in Eq. (26) linearly depends on photon production measured in experiment. On the theory side, the major source of uncertainty is from the correction to Eq. (2) at $p_\perp = 0.5$ GeV. One may get an idea on the magnitude of such corrections from strongly coupled QCD-like theories. For example, for $\mathcal{N} = 4$ super Yang-Mills theory in the strongly coupling limit, the corrections to Eq. (2) is at most 20% for p_\perp from 0 to πT (see Fig. 1 of Ref. [19]). In my calculations, I did not include the contributions at pre-equilibrium stage. To estimate the resulting uncertainty, I have extrapolated from the initial time $\tau_i = 0.4$ fm to a 3 times smaller value assuming 1-dimensional boost-invariant expansion between these times and computed the photon production during that interval. The corrections is a few percent at most.

I now compare my results with the electrical conductivity as measured on lattice. A recent quenched study using Wilson-Clover fermions [18] in the continuum limit found that $0.33C_{\text{EM}} \leq \sigma/T \leq 1C_{\text{EM}}$ at $T = 1.45T_c$. This result is consistent with other lattice measurements [20]. Here $C_{\text{EM}} = \sum_f q_f^2$ counts number of charge carriers. For example, for $f = u, d$, $C_{\text{EM}} = (5/9)e^2$ and for $f = u, d, s$, $C_{\text{EM}} = (2/3)e^2$. For comparison, I plot the range of $\sigma/(e^2T)$ at $T = 1.45T_c$ as indicated in Ref. [18] in Fig. 1 in dashes horizontal lines with $C_{\text{EM}} = (2/3)e^2$ by assuming in QGP, u, d, s all contribute to the conductivity. It is seen there that my results

are completely comparable with lattice measurement. My results are also consistent with Ref. [21] using the off-shell parton-hadron-string dynamics transport approach.

IV. SOFT PHOTON v_2 , CHIRAL ANOMALY AND MAGNETIC FIELD

In this section, I will study the effects of magnetic field and chiral anomaly on soft photon v_2 . I will evolve soft photon production rate in the presence of magnetic field and chiral anomaly Eq. (20) as derived in Sec. II in the realistic hydrodynamic background. As at RHIC energy, the typical speed of chiral magnetic wave v_χ is around $0.1 \sim 0.4$ [22] using the susceptibility measured on lattice [23], I then neglect v_χ^2 terms in Eq. (20) and approximate the photon rate at midrapidity $Y = 0$ as

$$\begin{aligned} & \left(\frac{\omega}{e^{\omega/T} - 1} \right)^{-1} \left[\omega \frac{d\Gamma_{\gamma,B}}{d^3p} \right] \\ & \approx \frac{\alpha_{\text{EM}}}{2\pi^2 e^2} [2(\sigma_0 - \sigma_{B,T}) + (\sigma_{B,T} + \sigma_{B,L}) \cos^2 \phi_p] + \mathcal{O}(v_\chi^2) \\ & = \frac{\alpha_{\text{EM}} \sigma_0}{\pi^2 e^2} \left[\left(1 - \frac{3}{4} r_T + \frac{1}{4} r_L \right) + \frac{1}{4} (r_T + r_L) \cos(2\phi_p) \right]. \end{aligned} \quad (27)$$

Here, I have introduced dimensionless ratio r_T, r_L :

$$r_T \equiv \sigma_{B,T}/\sigma_0, \quad r_L \equiv \sigma_{B,L}/\sigma_0 \quad (28)$$

to characterize the relative change of conductivity in the presence of magnetic field.

I now estimate the contribution from the magnetic field to photon v_2 as

$$v_2(B) \approx \frac{\frac{\alpha_{\text{EM}}}{8\pi^2} \int_0^{2\pi} \frac{d\phi_p}{2\pi} \int d^4x \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1} (r_T + r_L)}{\frac{\alpha_{\text{EM}}}{\pi^2} \int_0^{2\pi} \frac{d\phi_p}{2\pi} \int d^4x \frac{\omega_{\text{shift}} T}{\exp(\frac{\omega_{\text{shift}}}{T}) - 1} (1 - \frac{3}{4} r_T + \frac{1}{4} r_L)}. \quad (29)$$

To compute $v_2(B)$, I need to determine r_T and r_L . Let us first consider $r_T = \sigma_{B,T}/\sigma_0$ under the Drude approximation (relaxation time approximate). Recall the equation of motion for a massive particle in the presence of EM field and a drag force:

$$\frac{d\mathbf{p}}{dt} = q_f \mathbf{E} + \frac{\mathbf{p}}{M} \times q_f \mathbf{B} - \frac{\mathbf{p}}{\tau_{\text{rel}}}, \quad (30)$$

where τ_{rel} denotes the relaxation time. By imposing the steady-state condition $\frac{d\mathbf{p}}{dt} = 0$ and computing the current in response to \mathbf{E} , one finds

$$r_T = \frac{\sigma_{B,T}}{\sigma_0} = \frac{(q_f B \tau_{\text{rel}}/M)^2}{1 + (q_f B \tau_{\text{rel}}/M)^2}, \quad r_L = \frac{\sigma_{B,L}}{\sigma_0} = 0. \quad (31)$$

As charge carriers moving along the direction of magnetic field \mathbf{B} do not feel the Lorentz force, the magnetic field would not affect the longitudinal components of conductivity tensor, i.e., $r_L = 0$ under the drude estimation. Chiral anomaly may introduce a nontrivial contribution to the longitudinal conductivity. However, as the purpose of this section is to

⁴Other parameters to generate background hydrodynamic flow include the initial entropy density $s_i = 86.7 \text{ fm}^{-3}$ for all impact parameters. Results shown in the current paper are using Glauber initial conditions. I have performed the calculation for both CGC initial conditions and Glauber initial conditions at $b = 3.16$ fm and found a minor difference. I take impact parameters $b = 3.16, 5.78, 7.49, 8.87, 10.1, 11.1$ fm which correspond to centrality ranges 0–10%, 10–20%, 20–30%, 30–40%, 40–50%, 50–60%, respectively.

estimate the effects of magnetic field to soft photo v_2 , I will defer the effects due to r_L to future studies.

I now ready to evaluate $v_2(B)$ as defined by Eq. (29) in the realistic hydrodynamic background as I did in the previous section. To estimate r_T using Eq. (31), I need to estimate τ_{rel}/M in QGP. In $\mathcal{N} = 4$ supersymmetric Yang-Mills (sYM) theory in strong coupling limit, this is known for heavy quarks [24],

$$\left(\frac{\tau_{\text{rel}}}{M}\right)_{\text{sYM}} = \frac{2}{\sqrt{g^2 N_c \pi T^2}}. \quad (32)$$

Following Ref. [25], I will use Eq. (32) with $g^2 N_c = 6\pi$. I will parametrize my ignorance of τ_{rel}/M in QGP by introducing a dimensionless parameter λ :

$$\left(\frac{\tau_{\text{rel}}}{M}\right)_{\text{QGP}} = \lambda \left(\frac{\tau_{\text{rel}}}{M}\right)_{\text{sYM}}. \quad (33)$$

I will treat λ as a free parameter and study the effects of the magnetic field with various λ s. As r_T depends on the q_f . Strictly speaking, in Eq. (29), one should sum the contributions from different flavors. However, as the photon rate is proportional to q_f^2 , the number of photon produced by u quarks is roughly four times that produced by d quarks. I therefore, in my actual evaluation of Eq. (29), set $q_f = q_u = (2/3)e$.

I finally specify the profile eB during the hydrodynamic evolution. I neglect the spatial gradients of magnetic field and take it in the laboratory frame along the y direction. I use the time-varying profile of the magnetic field with a parametrization

$$eB(\tau) = \frac{(eB)_{\text{max}}}{1 + (\tau/\tau_B)^2}, \quad (34)$$

where I call τ_B the lifetime of the magnetic field. This form has been widely used in previous literature (see, for example, Refs. [8,22]). I take $(eB)_{\text{max}} = 3, 5, 7, 8, 9, 10 m_\pi^2$ for $b = 3.16, 5.78, 7.49, 8.87, 10.1, 11.1 \text{ fm}$ as guided by Ref. [27]. Due to current controversy over the medium effects on τ_B [28,29], I will leave τ_B as a free parameter in the following calculations.⁵

I have computed photon $v_2(B)$, the contribution from magnetic field to photon v_2 , as a function of the life time of magnetic field τ_B , by evolving the soft photon production rate in the presence of magnetic field based on hydrodynamics, i.e., Eqs. (20) and (27). r_T appearing in Eq. (27) is taken from Eq. (31) and $\omega_B \tau_{\text{rel}}$ is given by Eq. (33). I have presented my results for three different centrality bins (0–20%, 20–40%, 40–60%) with four different $\lambda = 2, 4, 6, 8$. The dependence of $v_2(B)$ on τ_B and λ are similar for the centrality bins under study. Perhaps not surprisingly, the contribution of magnetic field to photon v_2 increases with growing λ .⁶ Dashed curves in Fig. 3 correspond to the upper and lower bound of the direct photon v_2 at $p_\perp = 0.5 \text{ GeV}$ in the




FIG. 3. (Color online) The contribution of magnetic field to photon $v_2, v_2(B)$ [cf. Eq. (29)], at $p_\perp = 0.5 \text{ GeV}$ vs the lifetime of magnetic field τ_B . Photon production is computed based on Eqs. (20) and (27) for four different $\lambda = 2, 4, 6, 8$ (red solid, blue dashed, green dot-dashed, orange dotted curves, respectively). Here λ is a parameter appearing in Eq. (33) which parametrizes the $\omega_B \tau_{\text{rel}}$ in the plasma. Two dashed horizontal lines correspond to the upper and lower bound of direct photon v_2 at $p_\perp = 0.5 \text{ GeV}$ from the data (c.f. Fig. 2). Panels (a), (b), (c) correspond to centrality bins 0–20%, 20–40%, 40–60%, respectively.

data [3] (cf. Fig. 2). As one can see in Fig. 3, depending on the value of $\omega_B \tau_{\text{rel}}$, the magnetic field would give contribution, which is comparable to the data, to the soft photon v_2 for $\tau_B > 2 \sim 3 \text{ fm}$. I have also checked that for those τ_B, λ which reproduce the photon v_2 in the experiment, my estimation on σ_0 based the photon production rate in the absence of magnetic field will only be affected by 10–20%, within the error bar shown in the Fig. 1.

V. SUMMARY AND DISCUSSION

In this paper, I have estimated the electrical conductivity σ in the quark gluon plasma (QGP) based on the soft photon production from the data and realistic hydrodynamic evolution. I find that $\sigma/(e^2 T)$ is in the range $0.4 < \sigma/(e^2 T) < 1.1$. Previously, the electrical conductivity of QGP was mostly extracted from Euclidean correlator measured on lattice [18,20]. Those analyses always involve a nontrivial analytical continuation.

⁵It should also be pointed out that if $\tau_B \ll \tau_{\text{rel}}$, the hydrodynamic expression Eq. (27) does not apply.

⁶As λ here is the ratio of the actual ϕ_B in plasma to the characteristic ϕ_B^0 [cf. Eq. (33)], one does not expect λ to be $\mathcal{O}(10)$ and choose $\lambda = 8$ to be the largest value of λ used in the current computations.

The present work offered an alternative estimation of the electrical conductivity.

Photon production in heavy-ion collisions have been studied extensively (see for example Ref. [30]) based on the thermal photon emission rate computed from perturbative QCD(pQCD) [31]. While one may apply pQCD at high photon energy, its applicability for photon energy below a few GeV is not warranted. Indeed, experiment results [32] indicate that the hydrodynamic simulations with pQCD rate typically underestimate the thermal photon production. In this work, instead of taking the soft photon production rate of QGP as an input from certain microscopic calculations, I have extracted such rate based on hydrodynamics and the data. It would be interesting to extend the method used in this paper to obtain information on photon production rate at other p_\perp window.

The effects of magnetic field on the photon production and photon azimuthal anisotropy, v_2 , have attracted much attention recently [8,33–35]. I hope the present study based on hydrodynamics would shed light on how sizable the effects of magnetic field would be. In particular, by computing the contributions from magnetic field to photon v_2 for various τ_B , the life time of magnetic field in realistic hydrodynamic background, I found that if the life-time of $\tau_B > 2 \sim 3$ fm, the resulting soft photon v_2 is comparable to that measured in experiment. On the other hand, if the lifetime of magnetic field is as short as estimated in Ref. [28], the contribution from magnetic field to photon v_2 in low momentum region might be negligible.

It should be noticed that magnetic field even in the absence of anomaly would contribute to the photon v_2 via conventional synchrotron radiation [34]. Distinguishing the effects of chiral anomaly is not that straightforward. In hydrodynamic regime, however, a model-independent conclusion can be drawn in the light of the Eq. (20). According to Eq. (20) and the discussion in Sec. IV, the contributions due to chiral anomaly are fully parametrized by the speed of chiral magnetic wave v_χ while the effects due to the conventional cyclotron motion are parametrized by $\omega_B \tau_{\text{rel}}$. Moreover, the azimuthal angle dependence of the photon production is drastically affected by the additional pole structure of retarded Green's function due to the chiral magnetic wave. For example, by Fourier transforming Eq. (20), one can see explicitly that the Fourier

component of $\cos(4\phi_p)$ is proportional to v_χ^2 , suggesting that photon v_4 might be used to study the effects of chiral magnetic wave [35].

I will conclude this paper by pointing out that chiral anomaly may play *different* roles in soft photon production at RHIC and LHC. At RHIC energy where v_χ is not very close to 1, one may apply the approximate expression Eq. (27). At RHIC, I found that suppression of the transverse conductivity due to Lorentz force may play a dominant role to contribute to the photon v_2 . However, at LHC energy where v_χ approaches 1 due to much larger magnetic field, the pole of $G_R^{ij}(\omega, \mathbf{p})$ [cf. Eq. (15)] corresponding to chiral magnetic wave will be very close to the light cone. The photon production is largely enhanced along the direction \hat{B}_0 . Physically, this is due to the decay of the chiral magnetic wave into photon when v_χ is close to 1.⁷ This implies that chiral magnetic wave will give negative contribution to soft photon v_2 (see Ref. [35] for a holographic example). It is interesting to see if this will happen for soft photon production at LHC.

ACKNOWLEDGMENTS

I am indebted to Misha Stephanov for various discussion and suggestions during this project and Gocke Basar, Dimitri Kharzeev, Ho-Ung Yee for stimulating discussion on photon production in the presence of magnetic field and chiral anomaly. Special thanks are devoted to Karl Landsteiner who pointed out a mistake on the estimation of the longitudinal conductivity in the previous version of this paper. I would like to thank Chun Shen for email correspondence on hydrodynamic simulations and Benjamin Bannier and Richard Petti for conversations on photon production measured in experiment. I would like to express my gratitude to Jinfeng Liao, Lin Shu, Derek Teaney for helpful comments. I am in gratitude to the nuclear theory group of Stony Brook university for hospitality where this work was initiated and the funds from the Provost award of UIC to support my visit to Stony Brook University. I would also like to acknowledge the stimulating environment of the “Quantum anomalies and hydrodynamics” at Simons Center for Geometry and Physics and of the “Thermal photons and dileptons in heavy-ion collisions” workshop at RIKEN-BNL Research Center. This work is supported by the DOE Grant Nos. DE-FG0201ER41195 and DE-AC02-98CH10886.

⁷I thank D. Kharzeev for pointing this out to me.

-
- [1] J. Kapusta, *Finite-temperature Field Theory*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, 1993).
 - [2] A. Adare *et al.* (PHENIX Collaboration), [arXiv:1405.3940](#) [nucl-ex] (2014).
 - [3] B. Bannier, [arXiv:1408.0466](#) [nucl-ex] (2014).
 - [4] G. D. Moore and D. Teaney, *Phys. Rev. C* **71**, 064904 (2005).
 - [5] B. Ling, T. Springer, and M. Stephanov, *Phys. Rev. C* **89**, 064901 (2014).
 - [6] C. Shen, U. W. Heinz, J.-F. Paquet, I. Kozlov, and C. Gale, [arXiv:1308.2111](#) [nucl-th] (2013).
 - [7] D. E. Kharzeev, L. D. McLerran, and H. J. Warringa, *Nucl. Phys. A* **803**, 227 (2008).
 - [8] G. Basar, D. Kharzeev, D. Kharzeev, and V. Skokov, *Phys. Rev. Lett.* **109**, 202303 (2012).
 - [9] A. Adare *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **109**, 122302 (2012).
 - [10] D. T. Son and P. Surowka, *Phys. Rev. Lett.* **103**, 191601 (2009).

- [11] K. Fukushima, D. E. Kharzeev, and H. J. Warringa, *Phys. Rev. D* **78**, 074033 (2008).
- [12] D. E. Kharzeev, [arXiv:1312.3348](#) [hep-ph] (2013).
- [13] D. E. Kharzeev and H.-U. Yee, *Phys. Rev. D* **83**, 085007 (2011).
- [14] E. Lifshitz, L. Pitaevskii, and L. Landau, *Physical Kinetics* (Pergamon Press, Oxford, 1981), Vol. 60.
- [15] H. Song and U. W. Heinz, *Phys. Lett. B* **658**, 279 (2008); *Phys. Rev. C* **77**, 064901 (2008); **78**, 024902 (2008).
- [16] C. Shen, U. Heinz, P. Huovinen, and H. Song, *Phys. Rev. C* **82**, 054904 (2010); T. Renk, H. Holopainen, U. Heinz, and C. Shen, *ibid.* **83**, 014910 (2011).
- [17] P. Huovinen and P. Petreczky, *Nucl. Phys. A* **837**, 26 (2010).
- [18] H.-T. Ding, A. Francis, O. Kaczmarek, F. Karsch, E. Laermann, and W. Soeldner, *Phys. Rev. D* **83**, 034504 (2011).
- [19] S. Caron-Huot, P. Kovtun, G. D. Moore, A. Starinets, and L. G. Yaffe, *J. High Energy Phys.* **12** (2006) 015.
- [20] B. B. Brandt, A. Francis, H. B. Meyer, and H. Wittig, *J. High Energy Phys.* **03** (2013) 100; A. Amato, G. Aarts, C. Allton, P. Giudice, S. Hands, and J.-I. Skullerud, *Phys. Rev. Lett.* **111**, 172001 (2013); G. Aarts, C. Allton, J. Foley, S. Hands, and S. Kim, *ibid.* **99**, 022002 (2007).
- [21] W. Cassing, O. Linnyk, T. Steinert, and V. Ozvenchuk, *Phys. Rev. Lett.* **110**, 182301 (2013).
- [22] H.-U. Yee and Y. Yin, *Phys. Rev. C* **89**, 044909 (2014).
- [23] S. Borsanyi, Z. Fodor, S. D. Katz, S. Krieg, C. Ratti *et al.*, *J. High Energy Phys.* **01** (2012) 138.
- [24] C. Herzog, A. Karch, P. Kovtun, C. Kozcaz, and L. Yaffe, *J. High Energy Phys.* **07** (2006) 013; J. Casalderrey-Solana and D. Teaney, *Phys. Rev. D* **74**, 085012 (2006); S. S. Gubser, *ibid.* **74**, 126005 (2006).
- [25] U. Gursoy, D. Kharzeev, and K. Rajagopal, *Phys. Rev. C* **89**, 054905 (2014).
- [26] D. Axel (PHENIX Collaboration), talk presented at “Thermal Photons and Dileptons in Heavy-Ion Collisions” workshop at RIKEN/BNL, 2014.
- [27] A. Bzdak and V. Skokov, *Phys. Lett. B* **710**, 171 (2012).
- [28] L. McLerran and V. Skokov, [arXiv:1305.0774](#) [hep-ph] (2013).
- [29] K. Tuchin, *Phys. Rev. C* **88**, 024911 (2013).
- [30] R. Chatterjee, E. S. Frodermann, U. W. Heinz, and D. K. Srivastava, *Phys. Rev. Lett.* **96**, 202302 (2006); M. Dion, J.-F. Paquet, B. Schenke, C. Young, S. Jeon, and C. Gale, *Phys. Rev. C* **84**, 064901 (2011); C. Shen, U. W. Heinz, J.-F. Paquet, and C. Gale, *ibid.* **89**, 044910 (2014).
- [31] P. B. Arnold, G. D. Moore, and L. G. Yaffe, *J. High Energy Phys.* **12** (2001) 009.
- [32] A. Adare *et al.* (PHENIX Collaboration), *Phys. Rev. Lett.* **104**, 132301 (2010).
- [33] K. Tuchin, *Phys. Rev. C* **83**, 017901 (2011).
- [34] K. Tuchin, *Phys. Rev. C* **87**, 024912 (2013).
- [35] H.-U. Yee, *Phys. Rev. D* **88**, 026001 (2013).