

Low-energy R -matrix fits for the ${}^6\text{Li}(d,\alpha){}^4\text{He}$ S factor

J. Grineviciute and L. Lamia

Dipartimento di Fisica e Astronomia, Università di Catania, I-95125 Catania, Italy

A. M. Mukhamedzhanov

Cyclotron Institute, Texas A&M University, College Station, Texas 77843, USA

C. Spitaleri

*Dipartimento di Fisica e Astronomia, Università di Catania, I-95125 Catania, Italy
and Laboratori Nazionali del Sud-INFN, I-95125 Catania, Italy*

M. La Cognata

Laboratori Nazionali del Sud-INFN, I-95125 Catania, Italy

(Received 21 April 2013; revised manuscript received 14 October 2014; published 5 January 2015)

Background: Information about the ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction rates of astrophysical interest can be obtained by extrapolating direct data to lower energies, or by indirect methods. The indirect Trojan horse method, as well as various R -matrix and polynomial fits to direct data, estimate electron screening energies much larger than the adiabatic limit. Calculations that include the subthreshold resonance estimate smaller screening energies.

Purpose: Obtain the ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction R -matrix parameters and the bare astrophysical S factor for energies relevant to the stellar plasmas by fitting R -matrix formulas for the subthreshold resonances to the S -factor data above 60 keV.

Methods: The bare S factor is calculated using the single- and the two-level R -matrix formulas for the closest to the threshold 0^+ and 2^+ subthreshold states at 22.2, 20.2, and 20.1 MeV. The electron screening potential U_e is then obtained by fitting it as a single parameter to the low-energy data. The calculations are also done by fitting U_e simultaneously with other parameters.

Results: The low-energy S factor is dominated by the 2^+ subthreshold resonance at 22.2 MeV. The influence of the other two subthreshold states is small. The resultant electron screening is smaller than the adiabatic value. The fits that neglect the electron screening above 60 keV produce a significantly smaller electron screening potential. The calculations show a large ambiguity associated with a choice of the initial channel radius.

Conclusions: The R -matrix fits do not show a significantly larger U_e than predicted by the atomic physics models. The R -matrix best fit provides $U_e = 149.5$ eV and $S_b(0) = 21.7$ MeV b.

DOI: [10.1103/PhysRevC.91.014601](https://doi.org/10.1103/PhysRevC.91.014601)

PACS number(s): 24.30.-v, 21.10.Tg, 21.10.Jx, 29.85.-c

I. INTRODUCTION

To obtain the reaction rates relevant for nuclear astrophysics, experimental data should be extrapolated to the very-low-energy region (the Gamow window). The cross section depends strongly on the energy and therefore is expressed in terms of the astrophysical $S(E)$ factor

$$\sigma(E) = S(E)E^{-1} \exp(-2\pi\eta). \quad (1)$$

In the system of units, in which $\hbar = c = 1$, the Coulomb (Sommerfeld) parameter $\eta = Z_1 Z_2 e^2 \mu / k$. Z_i is the charge of the nucleus i , $k = \sqrt{2\mu E}$ and μ are the relative momentum and the reduced mass of the interacting nuclei, and E is their relative kinetic energy in the c.m. frame.

The extrapolation of the cross section down to low energies assumes that the Coulomb potential of the target nucleus and a projectile results from bare nuclei. In experimental conditions, however, the Coulomb potential is screened by electrons surrounding the target nucleus, thus reducing the height of the Coulomb barrier and leading to a higher cross section. As the energy approaches zero, the electron screening

potential U_e enhances the bare nucleus astrophysical factor $S(E) = S_b(E) \exp(\pi\eta U_e/E)$.

The estimated electron screening potential for the ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction in the adiabatic limit is a difference in atomic binding energies between Li and Be^+ , that is, 186 eV [1,2]. The Trojan horse (TH) experiment has indirectly measured the bare nucleus astrophysical factor [3]. A comparison of the TH results and a direct measurement led to $U_e = 340 \pm 51$ eV [3]. Large U_e values were obtained by Engstler *et al.* [4] using the polynomial fits, as well as by Barker [5] using the R -matrix fits. Barker also noted [5] that fixing U_e at 175 eV results in a reasonable fit with only slightly higher χ^2 . Ruprecht *et al.* [6] reported $U_e = 190 \pm 50$ eV and concluded that the lower screening energy is due to the influence of the 2^+ subthreshold resonance.¹

The current work presents a new R -matrix analysis for the low-energy ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction that considers three subthreshold resonances. The largest contribution to the

¹The subthreshold resonance is defined as a state which is bound in the entry channel and is a resonance in the exit channel.

low-energy $S(E)$ factor comes from the resonances and the subthreshold resonances closest to the threshold of the compound nucleus. The three subthreshold resonances closest to the threshold of ${}^8\text{Be}$ are the $2^+;0$ subthreshold state at -80 keV, followed by the $0^+;0$ subthreshold resonance at -2.08 MeV and a $2^+;0$ state at -2.18 MeV. The $2^+;0$ resonance at 2.92 MeV is not included in the calculation. The astrophysical S factor is first calculated including only the 2^+ subthreshold resonance at -80 keV and then by adding two more subthreshold resonances.

II. CONDITIONS

A. Experimental data

The fitting to the direct low energy experimental data [4,7–10] is done using the nonlinear least-squares procedure. Golovkov *et al.* [11] data are the outliers and hence are not included in the calculation.

The errors of the experimental data of Engstler *et al.* [4] do not include the reported uncertainties arising from the number of counts, angular distributions, the effective energy, and the target stoichiometry and hence are reduced by 5–10%. Including the latter errors would result in a normalized $\chi_n^2 \ll 1$. Here, the normalized χ_n^2 is defined as $\chi_n^2 = \chi^2/(N - n_p)$, where N is the number of experimental points used in the fit calculation and n_p is the number of parameters.

The error bars of Elwyn *et al.* experimental data [7] are enlarged to get $\chi_n^2 \approx 1$, as the underestimated errors may lead to a bias in the derived slope. The chosen error bars for Elwyn *et al.* data [7] are set to the 10% of the measured S -factor value.

B. Parameters

The R -matrix fits are calculated using the modified R -matrix formulas [12], which use the “observed” rather than the “formal” parameters. The alternate parametrization allows us to set the resonances’ energies at the experimental values, instead of conventionally [5,13] choosing the random formal energies in the vicinity of the resonance, and calculating the resulting observed energies after the R -matrix fit.

In the three-level fit, the two 2^+ resonances interfere, and the two-level R -matrix equations are used. The contribution of the 0^+ state is added incoherently. Considering only two channels for each state, the low-energy astrophysical $S(E)$ factor is defined as [13]

$$S(E) = \frac{\pi}{2\mu} \exp(2\pi\eta) (g_{0+} |U_{cc'}^{0+}|^2 + g_{2+} |U_{cc'}^{2+}|^2). \quad (2)$$

Here, the statistical spin factor $g_{J\pi}$ is

$$g_{J\pi} = \frac{2J + 1}{(2J_c + 1)(2J_{c'} + 1)}. \quad (3)$$

The collision matrix $U_{cc'}$ in terms of the observed parameters is identical to the one that is expressed in terms of the formal parameters [12]

$$U_{cc'} = \Omega_c \Omega_{c'} (\delta_{cc'} + 2i(P_{c'} P_c)^{1/2} \gamma_{c'}^T A \gamma_c), \quad (4)$$

and the A^{-1} matrix in terms of the observed parameters is defined as

$$(A^{-1})_{ij} = (E_i - E) \delta_{ij} - \sum_c \gamma_{ic} \gamma_{jc} (S_c + i P_c) + \sum_c \begin{cases} \gamma_{ic}^2 S_{ic} & i = j \\ \gamma_{ic} \gamma_{jc} \frac{S_{ic}(E - E_j) - S_{jc}(E - E_i)}{E_i - E_j} & i \neq j \end{cases}. \quad (5)$$

Here, S_{ic} is the shift function S_c , evaluated at energy E_i , P_c is the penetrability, and $\Omega_c = (I_c/O_c)^{1/2}$.

The R -matrix formula of Lane and Thomas [13] requires inclusion of all partial waves. For the initial 2^+ state one would have $(l = 0, s = 2)$, $(l = 2, s = 0)$, $(l = 2, s = 1)$, $(l = 2, s = 2)$ partial waves and $(l = 0, s = 0)$, $(l = 2, s = 2)$ partial waves for the initial 0^+ state. The exit channel for the 2^+ state is $(l = 2, s = 0)$, and for the 0^+ state it is $(l = 0, s = 0)$. The asymptotic normalization coefficients (ANC’s) for the bound states as well as the reduced width amplitudes γ_α ’s for the final channels are unknown and therefore are treated as parameters in the R -matrix calculation. Assuming that the s -wave approximation for the incoming deuteron is reasonable at very low deuteron energies, the number of free parameters reduces significantly.

C. Constraints

Unconstrained fits to direct data do not allow reliable determination of all parameters; therefore present calculations use constrained fits, which are subject to assumptions of the physical parameters. First, the R -matrix fits depend on the restrictions placed on the experimentally unknown $\alpha + \alpha$ channels’ partial widths.

The subthreshold resonances under consideration are broad. The experimental resonance widths of ${}^8\text{Be}$ are 0.8 and 0.88 MeV for the 2^+ states at 22.2 and 20.1 MeV, respectively, and 0.72 MeV for the 0^+ state at 20.2 MeV [14]. The total width for each level is a sum of all partial widths. The threshold of the ${}^6\text{Li}(d, \alpha){}^4\text{He}$ reaction entrance channel corresponds to a high excitation energy in a compound nucleus and hence many reaction channels are open. Considering only the ${}^6\text{Li}(d, \alpha){}^4\text{He}$ reaction, the major contribution to the total width at the subthreshold energy comes from the $\alpha + \alpha$ channel due to the large Q value of the reaction ($Q = 22.37$ MeV) and varies slowly with energy. The relative widths $\frac{\Gamma(\alpha)}{\Gamma(p)}$ of the 2^+ subthreshold resonance at 20.1 MeV have been determined experimentally to be 4.5 ± 0.6 [15]. $\Gamma_\alpha/\Gamma < 0.5$ has been reported [16] for the 0^+ state.

Page [17] performed a many-level multichannel simultaneous R -matrix fit to known $\alpha + \alpha$ elastic scattering data, as well as (d, α) and other channels’ data. However, a single-level fit to data, as well as the fit that includes all three aforementioned subthreshold resonances, fails for the bare ${}^6\text{Li}(d, \alpha){}^4\text{He}$ astrophysical S factor using the suggested formal $\alpha + \alpha$ channels’ partial widths $\Gamma_\alpha = 0.11$ MeV for the 2^+ subthreshold state at 22.2 MeV (experimental total width 0.8 MeV), $\Gamma_\alpha = 0.55$ MeV for the 0^+ subthreshold state at 20.2 MeV (experimental total width 0.72 MeV), and $\Gamma_\alpha = 0.17$ MeV for the 2^+ subthreshold state at 20.1 MeV

TABLE I. The 2^+ subthreshold resonance energy, the partial width of the $\alpha + \alpha$ channel, and the bare S factor resulting from a single-level R -matrix best fit to the low-energy Engstler data [4] above 60 keV (32 data points) for various channel radii. The reduced width amplitudes for both channels and the energy are treated as free parameters. The electron screening U_e is then fit as a single parameter to the Engstler data (64 data points).

R_i (fm)	R_f (fm)	E_{2^+} (MeV)	Γ_α (MeV)	$S_b(0)$ (MeV b)	χ_n^2	U_e (eV)
4.5	4.0	22.2610	0.6952	22.5222	0.7678	76.9987
4.5	4.5	22.2608	0.6947	22.5279	0.7681	76.5234
4.5	5.0	22.2606	0.6943	22.5278	0.7682	76.6777
4.5	5.5	22.2605	0.6940	22.5309	0.7684	76.4070
5.0	4.0	22.1911	0.7928	22.4021	0.7494	83.2557
5.0	4.5	22.1907	0.7923	22.4066	0.7496	82.8810
5.0	5.0	22.1904	0.7919	22.4087	0.7497	82.7473
5.0	5.5	22.1902	0.7917	22.4115	0.7498	82.4966
5.5	4.0	22.1368	0.9330	22.2113	0.7303	92.8859
5.5	4.5	22.1362	0.9327	22.2158	0.7305	92.4693
5.5	5.0	22.1357	0.9325	22.2155	0.7306	92.6005
5.5	5.5	22.1354	0.9323	22.2184	0.7307	92.3145

(experimental total width 0.88 MeV) in Ref. [17]. The fit also fails if one includes an arbitrary background.

Reference [18] notes that because the resonances are broad, the resonance contribution cannot be separated from the background contribution and therefore the elastic scattering data do not provide accurate resonance parameters. Also, the best fit of Ref. [17] includes an additional unknown level at 580 keV (excitation energy 22.78 MeV) with the formal $\Gamma_\alpha = 0.04$ MeV and $\Gamma_d = 0.23$ MeV for the s -wave deuteron.

The relative width $\Gamma(\alpha)/\Gamma(p)$ for the 2^+ subthreshold resonance at 20.1 MeV, found in Ref. [17], is significantly smaller than that given in literature [15] and obtained from the $p + {}^7\text{Li} \rightarrow \alpha + \alpha$ and $n + {}^7\text{Be} \rightarrow \alpha + \alpha$ reaction data. Also, the Γ_α of the 2^+ subthreshold resonance at 22.2 MeV obtained by Ref. [17] is much smaller than that obtained by other R -matrix or polynomial fits to data [4–6]. Hence, the present study considers only those constraints that are imposed by the experimental measurements.

For the bound states the ANC's are related to the reduced width amplitudes [19]

$$\gamma_c^2 = \frac{1}{2\mu} \frac{W_{-\eta_{kc}, l+1/2}^2(2\kappa_c r_0)}{r_0} |C|^2, \quad (6)$$

where $W_{-\eta_{kc}, l+1/2}(2\kappa_c r_0)$ is a Whittaker function. Without the experimentally imposed constraints, it is not possible to say which range of the ANC's is more appropriate. Therefore the ANC's are treated as free parameters.

D. Channel radii

The channel radii associated with the range of the nuclear force are calculated by a conventional formula of Lane and Thomas [13] $r = r_0(A_1^{1/3} + A_2^{1/3})$, where A_1 and A_2 are the mass numbers and r_0 is a numerical value between 1.4 and 1.5 fm. In principle, the collision matrix is independent of the choice of the channel radii, provided a large enough number of levels is included into the analysis, and consequentially is the astrophysical S factor. Partial widths and energies of the

resonances resulting from an R -matrix fit calculation should also be channel radii independent [20]. The sensitivity of the resonance parameters to the adopted channel radii in the initial and final channels is illustrated in Table I for a single-level calculation that uses the s -wave approximation for the deuteron. The parameters depend strongly on the ${}^6\text{Li} + d$ channel radius, which may indicate a need for the inclusion of the additional initial channel partial waves in the R -matrix fit. Also, as the analysis deals with broad resonances, the strong dependence on channel radii may support a need for an inclusion of additional levels into the R -matrix fit.

When allowing the channel radius to vary as one of the parameters, the single-level best fit sets the initial channel radius $R_i = 6.6$ fm. The best fit places the subthreshold resonance at 22.252 MeV with $\Gamma_\alpha = 0.2351$ MeV. The corresponding $U_e = 95.097$ eV. There is no obvious reason, however, to set the initial channel radius to this value, as one looks for a range of radii in which the conclusions of the calculation are reasonably stable, rather than for a single value which produces the lowest χ^2 . In this analysis, the chosen values for the initial and final channel radii are 5.0 and 4.5 fm, respectively. The sensitivity to the initial channel radius is evaluated in the error bars.

As deviations of the conclusions could be attributed to the effects of other levels, the single-level R -matrix fit for the 2^+ subthreshold resonance and the R -matrix fit that includes the three aforementioned subthreshold states are considered. The calculations use the s -wave approximation for the deuteron, and as with the existing data including more partial waves would only introduce more unknown fitting parameters.

III. RESULTS

A. Single-level fit

An unrestricted single-level R -matrix best fit to the low-energy experimental data [4,7–10] that uses the energy E_R of a 2^+ subthreshold resonance near the 22.2-MeV excitation energy, the electron screening potential U_e , as well as the




FIG. 1. (Color online) The astrophysical factor $S(E)$ for the ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction, resulting from a single-level R -matrix best fit to the experimental data [4,7–10], as a function of the relative kinetic energy E in the entrance channel. The solid line and the dotted line correspond to the observed and the bare astrophysical S factor, respectively. The top panel shows the best fit to data below 1 MeV that uses four free parameters: the reduced width amplitudes γ_α and γ_d , the 2^+ subthreshold resonance energy E_R near the 22.2-MeV excitation energy, and the electron screening potential U_e . The bottom panel shows the best fit to data in a 60 keV to 1 MeV range that uses γ_α and γ_d as the only parameters. The electron screening potential U_e is then fit to data [4,7–10] below 1 MeV as a single parameter. Engstler *et al.*'s data [4] are shown as pluses for atomic target and crosses for molecular target. The circles are the bare $S_b(E)$ factor from the Trojan horse experiment [3]. Other data [7–11,21] are taken from Ref. [22]. The errors of Engstler *et al.*'s data [4] do not include reported uncertainties arising from number of counts, angular distributions, effective energy, and target stoichiometry and hence are reduced by 5–10%. The error bars of experimental data of Elwyn *et al.* [7] are shown as in Ref. [22] but are enlarged in the calculation.

reduced width amplitudes γ_α and γ_d as free parameters, is shown in the top panel of Fig. 1. The best fit places the subthreshold resonance at $22.1692^{+0.0775}_{-0.0625}$ MeV with $\Gamma_\alpha = 0.8378^{+0.1564}_{-0.1044}$ MeV. The resulting bare astrophysical S factor at zero energy $S_b(0) = 21.6144^{+0.0725}_{-0.0573}$ MeV b and the electron screening $U_e = 161.953^{+7.648}_{-4.557}$ eV. The χ^2 of the fit, which uses four parameters and 82 data points, is 118.61. The normalized χ_n^2 is then 1.5207.

The subthreshold energy of the 2^+ state is then fixed at the experimental value of -80 keV (excitation energy 22.2 MeV) and the partial width of the $\alpha + \alpha$ channel is restricted not to exceed the total width of the state. The best fit that treats U_e , γ_α , and γ_d as parameters provides $S_b(0) = 22.1571^{+0.0322}_{-0.5271}$

MeV b, $\Gamma_\alpha = 0.5132^{+0.1952}_{-0.1141}$ MeV, and the electron screening $U_e = 126.665^{+43.231}_{-2.807}$ eV. The $\chi^2 = 119.69$ is only slightly higher than that resulting from the former fit, in which the subthreshold resonance energy is allowed to vary, resulting in a slightly smaller $\chi_n^2 = 1.5147$.

The ambiguity of the fit due to the sensitivity to the choice of the channel radii is larger for the restricted fit. The larger errors correspond to the fits for which the initial channel radius was reduced, resulting in larger values of the electron screening, the larger α partial width, and the smaller $S_b(0)$.

To neglect the electron screening, the single-level R matrix is fit to data above 60 keV [6]. The calculation for the bare S factor is done treating γ_α and γ_d as the only parameters. The observed S factor is then obtained by varying U_e as a single parameter to fit data below 1 MeV [4,7–10]. The best fit is shown in the bottom panel of Fig. 1, resulting in $S_b(0) = 22.7446^{+0.0104}_{-1.0660}$ MeV b and $\Gamma_\alpha = 0.5244^{+0.1864}_{-0.1212}$ MeV. χ^2 of the best fit to 50 data points is 66.6727, $\chi_n^2 = 1.3890$. The reduced width amplitudes are $\gamma_\alpha = -0.2105$ MeV $^{-1/2}$ and $\gamma_d = 0.4341$ MeV $^{-1/2}$. The corresponding ANC is $C = 3.1449$ fm $^{-1/2}$. Resultant $U_e = 69.288^{+98.537}_{-0.678}$ eV.

The sensitivity for the latter fit tracks the same way as the previous fit: Reducing the channel radius strongly increases the electron screening. Also, the single-level R -matrix best fit at a fixed energy of the 2^+ subthreshold resonance at -80 keV results in different sets of parameters, when fitting the electron screening simultaneously with other parameters and when the electron screening is neglected below 60 keV. Fitting all data below 1 MeV with the screening potential U_e included in the set of the fitting parameters results in $U_e = 126.7$ eV, while fitting data at energies above 60 keV, where supposedly the electron screening can be neglected, and then using the U_e as a single fitting parameter to fit all data below 1 MeV results in the screening potential nearly twice as low, $U_e = 69.3$ eV.

B. Two-level fit

The second closest subthreshold resonance to the threshold of ${}^8\text{Be}$ (excitation energy 22.28 MeV) is the $0^+;0$ state at -2.08 MeV (excitation energy 20.2 MeV). The S factors for the two states are added incoherently.

The R -matrix best fit that treats the γ_α 's and the ANC's as free parameters results in a much too large width of the $0^+;0$ state. Restricting the partial widths of the $\alpha + \alpha$ channels not to exceed the total widths of each state leads to $S_b(0) = 22.9535^{+0.0054}_{-1.1700}$ MeV b, $\Gamma_{2^+} = 0.5070^{+0.1883}_{-0.1186}$ MeV, and $\Gamma_{0^+} = 0.7115^{+0.0816}_{-0.5849}$ MeV. The best fit parameters are $\text{ANC} = 3.0533$ fm $^{-1/2}$ and $\gamma_\alpha = -0.2069$ MeV $^{-1/2}$ for the 2^+ state and $\text{ANC} = 12.6073$ fm $^{-1/2}$ and $\gamma_\alpha = 0.242$ MeV $^{-1/2}$ for the 0^+ state. χ^2 of the fit that uses four parameters and 50 data points is 65.8849, $\chi_n^2 = 1.4323$. The electron screening is then fit to data below 1 MeV as a single parameter, resulting in $U_e = 56.477^{+103.391}_{-0.200}$ eV.

By restricting the Γ_α of the $0^+;0$ subthreshold resonance not to exceed the half of the total width, as reported in Ref. [16], the best fit parameters become $\text{ANC} = 3.1437$ fm $^{-1/2}$ and $\gamma_\alpha = -0.2104$ MeV $^{-1/2}$ for the 2^+ state and $\text{ANC} = 1.1062$ fm $^{-1/2}$ and $\gamma_\alpha = 0.0823$ MeV $^{-1/2}$ for the 0^+ state. χ^2 of the best fit is

TABLE II. The R -matrix best fit observed parameters for three subthreshold resonances to experimental data [4,7–10]. The first parameter set results from the best fit to data between 60 keV and 1 MeV (50 data points), treating the reduced width amplitudes γ_α 's and the asymptotic normalization coefficients ANC's as parameters. U_e is then fit as a single parameter to all data. The second set results from the fit to data below 1 MeV (82 data points), simultaneously fitting the reduced width amplitudes γ_α 's, ANC's and the electron screening potential U_e .

$J^\pi T$	E_x (MeV)	Parameter set 1		U_e (eV)	Γ_α (MeV)	Parameter set 2			
		C (fm $^{-1/2}$)	γ_α (MeV $^{-1/2}$)			C (fm $^{-1/2}$)	γ_α (MeV $^{-1/2}$)	U_e (eV)	Γ_α (MeV)
2^+0	22.2	2.2607	0.2420		0.6932	2.2378	0.2558		0.7746
0^+0	20.2	1.7430	0.2009		0.4892	1.2790	0.0366		0.0163
2^+0	20.1	2.4390	0.1146	71.393	0.1462	2.1159	0.0925	149.521	0.0952
		$\chi_n^2 = 1.4889$				$\chi_n^2 = 1.5671$			

66.6689 and $\chi_n^2 = 1.4493$. Then $S_b(0) = 22.7512^{+0.1075}_{-1.0184}$ MeV b, $\Gamma_{2^+} = 0.5243^{+0.1786}_{-0.1213}$ MeV, and $\Gamma_{0^+} = 0.0823^{+0.3084}_{-0.0325}$ MeV, and the electron screening U_e is $68.670^{+95.031}_{-1.134}$ eV.

Thus, adding the second subthreshold resonance hardly affects the extracted screening potential from that obtained for a single subthreshold state. It also does not reduce the sensitivity to the channel radii. One notes that whether a single 2^+ subthreshold state is used in the R -matrix fit or the two subthreshold states 2^+ and 0^+ are considered, the determined α partial width of the 2^+ state is significantly larger than the one obtained in Ref. [17].

C. Three-level fit

In this section, the R -matrix fit considers three subthreshold resonances: two 2^+ and one 0^+ . The 2^+ states are interfering and the 0^+ state is added incoherently.

The partial widths of the $\alpha + \alpha$ channels for each state are constrained not to exceed the total experimental widths. In the absence of further constraints, the R -matrix best fit to the low-energy experimental data [4,7–10] for the bare astrophysical factor between 60 keV and 1 MeV results in $S_b(0) = 22.6507^{+0.3972}_{-0.8690}$ MeV b. The best fit parameters are shown in Table II as set 1. The partial widths of the $\alpha + \alpha$ channels are $\Gamma_{2^+} = 0.6932^{+0.0020}_{-0.1867}$ MeV, $\Gamma_{0^+} = 0.4892^{+0.2204}_{-0.4790}$ MeV, and $\Gamma_{2^+} = 0.1462^{+0.0128}_{-0.1451}$ MeV for the subthreshold levels at 22.2, 20.2, and 20.1 MeV respectively. χ^2 of the fit that uses six parameters and 50 data points is 65.5116, $\chi_n^2 = 1.4889$. The electron screening is then fit to all data below 1 MeV as a single parameter, resulting in $U_e = 71.393^{+88.7313}_{-19.603}$ eV. The best fit looks identical to Fig. 1(b).

The sensitivity of the fit to the choice of channel radii for the three-level fit tracks the same way as the sensitivity for a single-level fit, possibly indicating that this problem is not due to the lack of the background levels. $\chi^2 = 65.5116$ for the three-level fit is only slightly smaller than the χ^2 of a single-level fit, producing a slightly higher χ_n^2 . Due to a strong sensitivity to the channel radii, it is difficult to precisely determine the partial widths of the subthreshold resonances; however, it is clear that the fit prefers a partial width of the 2^+ state at 20.1 MeV significantly lower than the experimental total width 0.88 MeV [22] and its effect on the S factor is small.

The R -matrix best fit to the low-energy experimental data [4,7–10] that fits the ANC's, the reduced width amplitudes

γ_α 's, and the electron screening simultaneously looks very similar to Fig. 1(a). $S_b(0)$ of the best fit is $21.7186^{+0.5402}_{-0.0580}$ MeV b and $U_e = 149.521^{+19.307}_{-30.461}$ eV. The best fit parameters are shown in Table II as set 2. Resulting partial widths of $\alpha + \alpha$ channels are $\Gamma_{2^+} = 0.7746^{+0.0086}_{-0.3690}$ MeV, $\Gamma_{0^+} = 0.0163^{+0.4928}_{-0.0}$ MeV, and $\Gamma_{2^+} = 0.0952^{+0.0013}_{-0.0941}$ MeV for the subthreshold levels at 22.2, 20.2, and 20.1 MeV respectively. χ^2 of the fit that uses 7 parameters and 82 data points is 117.5280, $\chi_n^2 = 1.5671$.

The electron screening potential $U_e = 149.521$ eV resulting from the best fit that varies all parameters simultaneously is halved when the R matrix is fit to the experimental data above 60 keV to neglect the electron screening. A comparison of the best fit parameters, when U_e is fit simultaneously and separately, listed in Table II, possibly suggests that the electron screening is not negligible above 60 keV, contrary to Ref. [6], which states that for the deuteron energies larger than 50 keV the enhancement of the cross section due to the screening effect can be neglected. Therefore, the parameters we recommend are those resulting from a fit that varies U_e simultaneously with other parameters.

The bare S factor resulting from different R -matrix fits is larger than many previously reported calculations [5] and agrees with $S_b(0) = 23 \pm 2.5$ MeV b obtained by Ruprecht *et al.* [6]. While the value of the electron screening potential U_e is sensitive to the choice of channel radii, it is smaller than the adiabatic limit. We did not observe a significantly larger U_e , as was reported in Ref. [3–5].

IV. CONCLUSION

The astrophysical S factor for the low-energy ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction dominated by broad subthreshold resonances has been analyzed using the single-, two-, and three-level R -matrix fits. For the low-energy R matrix we use the s -wave approximation for the deuteron. The resulting ambiguity due to the choice of channel radii is large in a single-level fit, Table I, as well as in the fit that considers three subthreshold resonances.

Our goal is to check possibility of determining the electron screening potential from the low-energy astrophysical S factors and to determine the ANC's of the subthreshold states and α partial widths. We find that parameters depend on the number of the subthreshold states involved in the fitting. We consider the fit with three subthreshold states as the most reliable. We find that the extracted screening potential depends

on the used procedure. If we first fit the S factor varying all the parameters at energies above 60 keV, at which according to Ref. [6] the electron screening potential can be neglected, and then fix all the parameters and vary only U_e to fit the astrophysical factor at energies below 1 MeV, we obtain $U_e = 71.4$ eV. However, if we fit the S factor at energies below 1 MeV varying all the parameters simultaneously including U_e , we get $U_e = 149.5$ eV. Thus, the result strongly depends on the fitting procedure. We may conclude that the assumption of Ref. [6] that electron screening effects are negligible at energies above 50 keV is not valid and our recommended value is $U_e = 149.5$ eV. Note that other obtained fitting parameters are also sensitive to the fitting procedure. Our recommended values are the parameters from set 2 in Table II.

Our obtained α width for the 2^+ subthreshold resonance at 22.2 MeV, $\Gamma_\alpha = 0.77$ MeV, is higher than the 0.56-MeV value obtained in Ref. [6] and is closer to the 0.76-MeV value obtained in Ref. [5]. The partial width is lower than the total experimental value of 0.8 MeV [22], which was confirmed by corresponding theoretical calculations. The α partial width for the 0^+ subthreshold state at 20.2 MeV is small and agrees

with $\Gamma_\alpha/\Gamma < 0.5$ as reported in Ref. [16]. The partial width of 2^+ state at 20.1 MeV is much lower than the experimental value of the total width [22]. Our attempt to fit the S factor for the ${}^6\text{Li}(d,\alpha){}^4\text{He}$ reaction using α widths from Ref. [17] failed, and we believe that the main reason for that is that our α partial width for the dominant 2^+ state at 22.2 MeV is significantly higher than 0.11 MeV obtained in Ref. [17]. We conclude that while it is difficult to pinpoint accurately the electron screening potential by fitting existing direct measurements, the obtained electron screening potential is definitely smaller than the adiabatic limit $U_e = 186$ eV [1,2].

ACKNOWLEDGMENTS

This work has been partially supported by the Italian Ministry of the University under Grant No. RFBR082838 (FIRB2008). A.M.M. acknowledges support by the U.S. Department of Energy under Grants No. DE-FG02-93ER40773 and No. DE-FG52-09NA29467 and NSF under Grant No. PHY-1415656.

-
- [1] H. J. Assenbaum, K. Langake, and C. Rolfs, *Z. Phys. A* **327**, 461 (1987).
 - [2] T. D. Shoppa, S. E. Koonin, K. Langanke, and R. Seki, *Phys. Rev. C* **48**, 837 (1993).
 - [3] C. Spitaleri *et al.*, *Phys. Rev. C* **63**, 055801 (2001).
 - [4] S. Engstler, G. Raimann, C. Angulo, U. Greife, C. Rolfs, U. Schröder, E. Somorjai, B. Kirch, and K. Langanke, *Z. Phys. A* **342**, 471 (1992).
 - [5] F. C. Barker, *Nucl. Phys. A* **707**, 277 (2002).
 - [6] G. Ruprecht, K. Czerski, D. Bemmerer, M. Hoefft, and P. Heide, *Phys. Rev. C* **70**, 025803 (2004).
 - [7] A. J. Elwyn, R. E. Holland, C. N. Davids, L. Meyer-Schutzmeister, J. E. Monahan, F. P. Mooring, and W. Ray, Jr., *Phys. Rev. C* **16**, 1744 (1977).
 - [8] C. Dunjiu, Z. Enchen, and J. Chenglie (unpublished).
 - [9] R. Risler, W. Grüebler, A. A. Debenham, V. König, P. A. Schmelzbach, and D. O. Boerma, *Nucl. Phys. A* **286**, 115 (1977).
 - [10] J. M. F. Jeronimo, G. S. Mani, F. Picard, and A. Sadeghi, *Nucl. Phys.* **38**, 11 (1962).
 - [11] M. S. Golovkov *et al.*, *Sov. J. Nucl. Phys.* **34**, 480 (1981).
 - [12] C. R. Brune, *Phys. Rev. C* **66**, 044611 (2002).
 - [13] A. M. Lane and R. G. Thomas, *Rev. Mod. Phys.* **30**, 257 (1958).
 - [14] D. R. Tilley, *Nucl. Phys. A* **745**, 155 (2004).
 - [15] V. M. Pugach *et al.*, *Bull. Rus. Acad. Sci. Phys.* **56**, 1768 (1992).
 - [16] J. Benn, E. B. Dally, H. H. Müller, R. E. Pixley, H. H. Staub, and H. Winkler, *Nucl. Phys. A* **106**, 296 (1967).
 - [17] P. R. Page, *Phys. Rev. C* **72**, 054312 (2005).
 - [18] A. M. Mukhamedzhanov, B. F. Irgaziev, V. Z. Goldberg, Y. V. Orlov, and I. Qazi, *Phys. Rev. C* **81**, 054314 (2010).
 - [19] A. M. Mukhamedzhanov and R. E. Tribble, *Phys. Rev. C* **59**, 3418 (1999).
 - [20] P. Descouvemont and D. Baye, *Rep. Prog. Phys.* **73**, 036301 (2010).
 - [21] C. R. Gould, J. M. Joyce, and J. R. Boyce, *Proceedings of Conference on Nuclear Cross Sections and Technology*, Publication 421 (NBS, Washington, DC, 1975), p. 697.
 - [22] D. R. Tilley *et al.* (unpublished); this is part of the updated version of F. Ajzenberg-Selove, *Nucl. Phys. A* **490**, 1 (1988).