

Pairing phase transition: A finite-temperature relativistic Hartree-Fock-Bogoliubov studyJia Jie Li,^{1,2,3} Jérôme Margueron,² Wen Hui Long,³ and Nguyen Van Giai¹¹*Institut de Physique Nucléaire, IN2P3-CNRS, Université Paris-Sud, F-91406 Orsay, France*²*Institut de Physique Nucléaire de Lyon, IN2P3-CNRS, Université de Lyon, F-69622 Villeurbanne, France*³*School of Nuclear Science and Technology, Lanzhou University, Lanzhou 730000, China*

(Received 8 April 2015; revised manuscript received 27 May 2015; published 2 July 2015)

Background: The relativistic Hartree-Fock-Bogoliubov (RHFB) theory has recently been developed and it provides a unified and highly predictive description of both nuclear mean-field and pairing correlations. Ground-state properties of finite nuclei can accurately be reproduced without neglecting exchange (Fock) contributions.

Purpose: Finite-temperature RHFB (FT-RHFB) theory has not yet been developed, leaving yet unknown its predictions for phase transitions and thermal excitations in both stable and weakly bound nuclei.

Method: FT-RHFB equations are solved in a Dirac Woods-Saxon (DWS) basis considering two kinds of pairing interactions: finite or zero range. Such a model is appropriate for describing stable as well as loosely bound nuclei since the basis states have correct asymptotic behavior for large spatial distributions.

Results: Systematic FT-RH(F)B calculations are performed for several semimagic isotopic/isotonic chains comparing the predictions of a large number of Lagrangians, among which are PKA1, PKO1, and DD-ME2. It is found that the critical temperature for a pairing transition generally follows the rule $T_c = 0.60\Delta(0)$ for a finite-range pairing force and $T_c = 0.57\Delta(0)$ for a contact pairing force, where $\Delta(0)$ is the pairing gap at zero temperature. Two types of pairing persistence are analyzed: type I pairing persistence occurs in closed subshell nuclei while type II pairing persistence can occur in loosely bound nuclei strongly coupled to the continuum states.

Conclusions: This FT-RHFB calculation shows very interesting features of the pairing correlations at finite temperature and in finite systems such as pairing re-entrance and pairing persistence.

DOI: [10.1103/PhysRevC.92.014302](https://doi.org/10.1103/PhysRevC.92.014302)

PACS number(s): 05.70.Ce, 21.60.Jz, 24.10.Pa, 64.60.Ht

I. INTRODUCTION

Over the past several decades, the thermodynamical properties of excited nuclei have drawn renewed attention due to the advanced accurate measurements of level densities at low excitation energies [1–5]. Pairing correlations play an essential role in many fermion systems and have thus a strong influence on nuclear structure at low excitation energies [6–10]. Pairing correlations in finite systems such as nuclei or Wigner-Seitz cells, and in infinite ones, such as in neutron star matter, may exhibit different behaviors reflected in the specific heat and the level density [11–15]. Moreover, the phase transition is a complex and rich phenomenon, where pairing re-entrance in asymmetric matter, in odd-nuclei, rotating nuclei, and even in doubly magic nuclei close to the drip line may occur [6,14,16,17]. The interplay between temperature and shell effects in superfluid systems, giving rise to re-entrance or its opposite phenomenon—suppression—still remains to be studied.

The competition between temperature and pairing correlations in nuclei at low excitation energies has been studied for several decades, with the pioneering works based on finite-temperature Bardeen-Cooper-Schrieffer (FT-BCS) theory [18] and finite-temperature Hartree-Fock-Bogoliubov (FT-HFB) theory [19]. It was predicted that the critical temperature T_c for pair correlation quenching could be expressed, as in uniform matter, as a function of the average pairing gap at zero temperature $\Delta(0)$ following the relations $T_c = 0.57\Delta(0)$ for a constant pairing force [18]. The more evolved Bogoliubov approach later has been applied to finite nuclei confirming the existence of such relations between T_c and $\Delta(0)$. For the simplified degenerate model, the relation was found to

be $T_c = 0.50\Delta(0)$ [19], while in rare-earth transition nuclei the interplay with deformation induces shape transitions in the superfluid phase, leading to the ratio $T_c/\Delta(0) = 0.57$ for protons and 0.63 for neutrons [20]. In addition, pairing correlations are expected to play an important role in the decay of compound nuclei formed in heavy-ion collisions, as illustrated in the seminal work presented in Ref. [21]. More recently, the BCS and HFB approaches have been extended to self-consistent mean-field models in order to improve the description of the pairing transition in spherical nuclei [22–24], as well as in deformed nuclei where shape transitions have been predicted [25–29]. In summary, the ratio $T_c/\Delta(0)$ lies in the interval 0.50–0.60, where the uncertainty originates mainly from the detailed level structure of spherical and deformed nuclei which depends itself on models.

It is worth noticing that in most of the quoted studies, the calculations were performed either on the harmonic oscillator basis or within the nonrelativistic framework. Due to the limitation of the harmonic oscillator basis in giving an appropriate asymptotic behavior of the single particle (s.p.) wave functions, the nucleon densities at large distance converge very slowly with respect to the size of the basis. The situation becomes even more serious in the weakly bound nuclei close to the drip line [30,31]. At present a realistic framework is to perform the calculations in an appropriate basis that can provide a reasonable description of both the overall and asymptotic behaviors of the density profiles, for instance, the Woods-Saxon (WS) basis [32,33]. In some applications, the small component of the Dirac spinors were usually neglected in determining the relativistic Hartree-Bogoliubov (RHB) and

finite-temperature RHB (FT-RHB) pairing tensor [24,34]. We therefore present, in this paper, the first, as far as we know, fully relativistic Hartree-Fock-Bogoliubov (RHFB) calculations at finite temperature (FT-RHFB): both the large and small components of the Dirac spinors contribute to the pairing channel, and the contributions of the Fock terms are naturally included. In addition, the Dirac Woods-Saxon (DWS) basis [32] is employed to better describe weakly bound nuclei.

This paper is organized as follows. The general formalism of the FT-RHFB theory is outlined in Sec. II. In Sec. III, we compare the results obtained with different covariant density functional (CDF) models [35–40] using the finite- and zero-range pairing interactions [41,42]. Systematic FT-RHFB calculations are performed for several isotopes/isotones. Moreover, the questions of pairing persistence and re-entrance phenomena with increasing temperature are addressed. Finally, the main conclusions are drawn in Sec. IV.

II. GENERAL FORMALISM AND NUMERICAL DETAILS

We briefly recall here the general features of RHFB theory and thermodynamics and then their generalization to the finite-temperature case.

A. RHFB framework

The relativistic Hartree-Fock (RHF) theory [35,40,43] is designed to describe bulk and s.p. nuclear properties. The relevant degrees of freedom are the isospin doublet nucleon field $\psi_{n(p)}$, the meson fields which mediate the nuclear interactions (two isoscalar fields σ and ω , two isovector fields ρ and π), and the photon field (A) that accounts for the electromagnetic interaction. A general effective Lagrangian is constructed in terms of various currents or densities: $(\bar{\psi}\tau\Gamma\psi)$, where $\tau \in \{1, \boldsymbol{\tau}\}$ (1 stands for the isoscalar quantities and the isospin Pauli matrix $\boldsymbol{\tau}$ corresponds to the isovector ones) and Γ denotes the Dirac matrices $\{1, \gamma_\mu, \gamma_5\gamma_\mu, \sigma_{\mu\nu}\}$. Applying the standard variations of the Lagrangian, one may obtain the field equations for nucleon, meson, and photon fields (respectively, the Dirac, Klein-Gordon, and Proca equations) and the continuity equations leading to the Hamiltonian. With the creation and annihilation operators (c^\dagger, c) defined from stationary solutions of the Dirac equation, the effective Hamiltonian \mathcal{H} is formally expressed in the second quantized form as

$$\mathcal{H} = \sum_{ij} c_i^\dagger c_j T_{ij} + \frac{1}{2} \sum_{ijkl;\phi} c_i^\dagger c_j^\dagger c_l c_k V_{ijkl}^\phi, \quad (1)$$

where T_{ij} represents the Dirac kinetic energy, and the two-body terms V_{ijkl} correspond to different types of meson- (or photon-) nucleon couplings denoted by ϕ . In the mean-field approximation, the energy functional \mathcal{E} is obtained by taking the expectation value of the Hamiltonian \mathcal{H} on a Slater determinant, where both the Hartree and Fock terms are considered. The RHFB theory [44–46] is deduced by incorporating the Bogoliubov transformation

$$\begin{pmatrix} \alpha \\ \alpha^\dagger \end{pmatrix} = \begin{pmatrix} \psi_U^* & \psi_V^* \\ \psi_V & \psi_U \end{pmatrix} \begin{pmatrix} c \\ c^\dagger \end{pmatrix} \quad (2)$$

with the RHF model. It provides a unified and self-consistent description of both particle-hole (ph) and particle-particle (pp) channels at the mean-field level.

B. Thermodynamics and statistical mechanics

The thermodynamical properties of a system are calculated here in the canonical ensemble. For a statistical N -body system at finite temperature T , the equilibrium state is obtained from the variational principle applied to the grand-canonical potential Ω [19,21,47],

$$\Omega(T, \lambda) = F - \lambda N = E - TS - \lambda N, \quad (3)$$

where F is the free energy, S the entropy, E the total energy, and λ the associated Lagrange multiplier. Namely the variation

$$\delta\Omega = 0 \quad (4)$$

defines the density operator \mathcal{D} with trace equal to 1, and the grand partition function Z , respectively, read as:

$$\mathcal{D} = Z^{-1} \exp\{-\beta(\mathcal{H} - \lambda\mathcal{N})\}, \quad (5a)$$

$$Z = \text{Tr}[\exp\{-\beta(\mathcal{H} - \lambda\mathcal{N})\}], \quad (5b)$$

where \mathcal{N} is the particle number operator and $\beta = T^{-1}$. As is conventional, the temperature T is given in energy units. For arbitrary operator \mathcal{O} , the thermal average over the excited states populated at finite temperature is defined as:

$$\langle \mathcal{O} \rangle = \text{Tr}(\mathcal{D}\mathcal{O}), \quad (6)$$

where the trace is taken over all possible excited states. At finite temperature, the total energy, entropy, and number of particles are therefore expressed as

$$E = \langle \mathcal{H} \rangle = \text{Tr}(\mathcal{D}\mathcal{H}), \quad (7a)$$

$$S = -\langle \ln \mathcal{D} \rangle = -\text{Tr}(\mathcal{D} \ln \mathcal{D}), \quad (7b)$$

$$N = \langle \mathcal{N} \rangle = \text{Tr}(\mathcal{D}\mathcal{N}). \quad (7c)$$

C. Extension to finite temperature

The FT-RHFB theory is a straightforward generalization of the RHFB theory that readily incorporates a statistical ensemble of excited states. At the finite-temperature mean-field level, the density operator is approximated by [19]:

$$\mathcal{D} = \prod_{\alpha} [f_{\alpha} \mathcal{N}_{\alpha} + (1 - f_{\alpha})(1 - \mathcal{N}_{\alpha})], \quad (8)$$

where f_{α} is the Fermi-Dirac distribution

$$f_{\alpha} = \langle \mathcal{N}_{\alpha} \rangle = \frac{1}{1 + e^{\beta E_{\alpha}}}. \quad (9)$$

Notice that, for $T = 0$, we have $f_{\alpha} = 0$ for all states α . The quasiparticle energy E_{α} is obtained as the solution of the FT-RHFB equations, see Eq. (11).

In spherically symmetric systems the Dirac-Bogoliubov spinors in Eq. (2) can be written as

$$\psi_{U_{\alpha}}(\mathbf{r}) = \frac{1}{r} \begin{pmatrix} iG_{U_{\alpha}}(r) \\ F_{U_{\alpha}}(r)\boldsymbol{\sigma} \cdot \hat{\mathbf{r}} \end{pmatrix} \mathcal{Y}_{jm}^l(\hat{\mathbf{r}}) \chi_{\frac{1}{2}}^l(\tau), \quad (10a)$$

$$\psi_{V_{\alpha}}(\mathbf{r}) = \frac{1}{r} \begin{pmatrix} iG_{V_{\alpha}}(r) \\ F_{V_{\alpha}}(r)\boldsymbol{\sigma} \cdot \hat{\mathbf{r}} \end{pmatrix} \mathcal{Y}_{jm}^l(\hat{\mathbf{r}}) \chi_{\frac{1}{2}}^l(\tau), \quad (10b)$$

where $G_{U(V)}$ and $F_{U(V)}$ correspond to the radial parts of the upper and lower components, respectively, $\mathcal{Y}_{jm}^l(\hat{\mathbf{r}})$ is the

spinor spherical harmonics, and $\chi_{\frac{1}{2}}(\tau)$ is the isospinor. Here the subindex $\alpha = (a, m) = (n, l, j, m)$ contains the quantum numbers n (number of nodes of the upper component $G_{U(V)}$), l (orbital angular momentum), and j, m (total angular momentum and its projection to the z axis).

The minimization of Ω with respect to \mathcal{D} leads to the FT-RHFB equations,

$$\int d\mathbf{r}' \begin{pmatrix} h(\mathbf{r}, \mathbf{r}') & \Delta(\mathbf{r}, \mathbf{r}') \\ -\Delta(\mathbf{r}, \mathbf{r}') & h(\mathbf{r}, \mathbf{r}') \end{pmatrix} \begin{pmatrix} \psi_U(\mathbf{r}') \\ \psi_V(\mathbf{r}') \end{pmatrix} = \begin{pmatrix} \lambda + E & 0 \\ 0 & \lambda - E \end{pmatrix} \begin{pmatrix} \psi_U(\mathbf{r}) \\ \psi_V(\mathbf{r}) \end{pmatrix}, \quad (11)$$

which are formally identical to the RHFB equations [44–46]. The Dirac Hamiltonian $h(\mathbf{r}, \mathbf{r}')$ contains the kinetic energy h^K , the direct local potential h^D , and exchange nonlocal potential h^E ,

$$h^K(\mathbf{r}, \mathbf{r}') = [\boldsymbol{\alpha} \cdot \mathbf{p} + \beta M] \delta(\mathbf{r} - \mathbf{r}'), \quad (12a)$$

$$h^D(\mathbf{r}, \mathbf{r}') = [\Sigma_T(\mathbf{r})\gamma_5 + \Sigma_0(\mathbf{r}) + \beta \Sigma_S(\mathbf{r})] \delta(\mathbf{r} - \mathbf{r}'), \quad (12b)$$

$$h^E(\mathbf{r}, \mathbf{r}') = \begin{bmatrix} Y_G(\mathbf{r}, \mathbf{r}') & Y_F(\mathbf{r}, \mathbf{r}') \\ X_G(\mathbf{r}, \mathbf{r}') & X_F(\mathbf{r}, \mathbf{r}') \end{bmatrix}. \quad (12c)$$

$$X_{G_a}^{(\phi)}(r, r') = \sum_b \mathcal{T}_{ab}^\phi \hat{j}_b^2 [(g_\phi F_{V_b})_r \mathcal{R}_{ab}^{X_G}(\mathbf{m}_\phi; r, r') (g_\phi G_{V_b})_{r'} (1 - f_b) + (g_\phi F_{U_b})_{r'} \mathcal{R}_{ab}^{X_G}(\mathbf{m}_\phi; r, r') (g_\phi G_{U_b})_{r'} f_b], \quad (14a)$$

$$X_{F_a}^{(\phi)}(r, r') = \sum_b \mathcal{T}_{ab}^\phi \hat{j}_b^2 [(g_\phi F_{V_b})_r \mathcal{R}_{ab}^{X_F}(\mathbf{m}_\phi; r, r') (g_\phi F_{V_b})_{r'} (1 - f_b) + (g_\phi F_{U_b})_{r'} \mathcal{R}_{ab}^{X_F}(\mathbf{m}_\phi; r, r') (g_\phi F_{U_b})_{r'} f_b], \quad (14b)$$

$$Y_{G_a}^{(\phi)}(r, r') = \sum_b \mathcal{T}_{ab}^\phi \hat{j}_b^2 [(g_\phi G_{V_b})_r \mathcal{R}_{ab}^{Y_G}(\mathbf{m}_\phi; r, r') (g_\phi G_{V_b})_{r'} (1 - f_b) + (g_\phi G_{U_b})_{r'} \mathcal{R}_{ab}^{Y_G}(\mathbf{m}_\phi; r, r') (g_\phi G_{U_b})_{r'} f_b], \quad (14c)$$

$$Y_{F_a}^{(\phi)}(r, r') = \sum_b \mathcal{T}_{ab}^\phi \hat{j}_b^2 [(g_\phi G_{V_b})_r \mathcal{R}_{ab}^{Y_F}(\mathbf{m}_\phi; r, r') (g_\phi F_{V_b})_{r'} (1 - f_b) + (g_\phi G_{U_b})_{r'} \mathcal{R}_{ab}^{Y_F}(\mathbf{m}_\phi; r, r') (g_\phi F_{U_b})_{r'} f_b]. \quad (14d)$$

In these expressions, \mathcal{T}_{ab}^ϕ denotes the isospin factors: δ_{ab} for isoscalar channels and $2 - \delta_{ab}$ for isovector channels, $\hat{j}_b^2 = 2j_b + 1$ is the degeneracy number of the corresponding energy level, g_ϕ represents the coupling constants, and \mathcal{R}_{ab} denotes the multipole expansions of meson propagators. Equation (14) is a generalization at finite temperature of the expressions given in Ref. [45].

Next we consider the pairing field $\Delta(r, r')$ of Eq. (11),

$$\Delta_a(r, r') = -\frac{1}{2} \sum_b V_{ab}^{\text{pp}}(r, r') \kappa_b(r, r'), \quad (15)$$

with the pairing interaction V^{pp} and the pairing tensor κ . If we take a finite-range pairing force, the pairing tensor κ will read as

$$\kappa_a(r, r') = \hat{j}_a^2 \{ [G_{V_a}(r)G_{U_a}(r') + F_{V_a}(r)F_{U_a}(r')] + [G_{U_a}(r)G_{V_a}(r') + F_{U_a}(r)F_{V_a}(r')] \} (1 - 2f_a). \quad (16)$$

Notice that the temperature dependence of the solution ($E_\alpha; \psi_{U_\alpha}, \psi_{V_\alpha}$) of the FT-RHFB Eq. (11) comes implicitly

In the above expressions, the local self-energies Σ_S, Σ_0 , and Σ_T contain the contributions from the Hartree (direct) terms and the rearrangement terms, which depend directly on various local quasiparticle densities. They are the vector, scalar, and tensor densities, respectively,

$$\rho_v(r) = \frac{1}{4\pi r^2} \sum_a \hat{j}_a^2 \{ [G_{V_a}^2(r) + F_{V_a}^2(r)](1 - f_a) + [G_{U_a}^2(r) + F_{U_a}^2(r)]f_a \}, \quad (13a)$$

$$\rho_s(r) = \frac{1}{4\pi r^2} \sum_a \hat{j}_a^2 \{ [G_{V_a}^2(r) - F_{V_a}^2(r)](1 - f_a) + [G_{U_a}^2(r) - F_{U_a}^2(r)]f_a \}, \quad (13b)$$

$$\rho_t(r) = \frac{1}{4\pi r^2} \sum_a \hat{j}_a^2 \{ 2G_{V_a}(r)F_{V_a}(r)(1 - f_a) + 2G_{U_a}(r)F_{U_a}(r)f_a \}. \quad (13c)$$

Notice that, at the limit $T = 0$ where $f_a = 0$, Eqs. (13a)–(13c) reduce to the usual RH(F)B expressions at zero temperature, e.g., given in Refs. [39,45]. The nonlocal self-energies $X_{G(F)}$ and $Y_{G(F)}$ come from the Fock (exchange) terms,

through the quasiparticle densities $\rho_{v,s,t}$, nonlocal potentials $X(Y)$, and pairing tensor κ .

For practical evaluation of the pairing correlations, an average pairing gap is introduced and defined as the ratio of the pairing energy over the pairing tensor,

$$\Delta = \frac{\text{Tr}(\Delta\kappa)}{\text{Tr}\kappa}. \quad (17)$$

This quantity is calculated for neutrons (Δ_n) and protons (Δ_p) separately and is discussed in detail in the next section.

The total FT-RHFB energy E of the system is calculated with the microscopic two-body center-of-mass correction [48]. The entropy S of the system can be evaluated from

$$S(T) = - \sum_\alpha [f_\alpha \ln f_\alpha + (1 - f_\alpha) \ln(1 - f_\alpha)], \quad (18)$$

and the specific heat is defined by

$$C_v(T) = T \left. \frac{\partial S(T)}{\partial T} \right|_N. \quad (19)$$

They correspond to the first and second derivatives of the free energy F , respectively.

The integrodifferential FT-RHFB Eq. (11) is solved by using a DWS basis [32] with a radial cutoff $R = 26$ fm. The numbers of positive and negative energy states in the basis expansion for each s.p. angular momentum (l, j) are chosen to be 36 and 12, respectively. We have verified that this truncation scheme provides sufficient numerical accuracy for the description of weakly bound nuclei (Pb isotopes).

III. RESULTS AND DISCUSSION

In this section, we compare the predictions based on several models which are determined by the model Lagrangian as well as the pairing interaction. The latter interaction is either a finite-range Gogny D1S interaction [41] or a density-dependent contact interaction (DDCI) of the form [42]:

$$V(r, r') = V_0 \frac{1}{2} (1 - P_\sigma) \left(1 - \frac{\rho(r)}{\rho_0} \right) \delta(r - r'). \quad (20)$$

Notice that the DDCI requires a regularization scheme. We have considered in this study a simple cutoff scheme, defined to be 100 MeV in quasiparticle space, and the strength V_0 is adjusted to reproduce the same average pairing gap as that obtained with the Gogny D1S interaction. These values of V_0 will be given with the results hereafter. The D1S pairing interaction depends slightly on the mass: A general factor g is therefore introduced for its strength, as in Refs. [49,50]. We consider six different model Lagrangians, which are given in Table I. Some of the bulk properties of nuclear matter determined by these Lagrangians are also shown. They are generally compatible with the expected gross properties of finite nuclei, e.g., the binding energy $E/A \simeq 16$ MeV, the saturation density $\rho_0 \simeq 0.15 \text{ fm}^{-3}$, the incompressibility modulus $K \in [220, 280]$ MeV, and the symmetry energy $J \in [32, 39]$ MeV.

In the following discussion, the persistence and re-entrance of the pairing phenomenon will be discussed and analyzed. Let us briefly recall the unifying mechanism which is at play in these various phenomena, recently discussed in Ref. [14]. Due to thermal excitations, s.p. states above the Fermi energy can be slightly populated while states below the Fermi energy can be partially depleted. This occurs if the involved new states are not too far in energy from the last occupied state, but it should also be not too close, otherwise these states would already participate to pairing correlations at zero temperature. The typical shell gap should be around 2 MeV. The participation of these states at finite temperature gives rise either to the persistence of pairing correlations slightly above the usual critical temperature for nuclei which are already superfluid at zero temperature or to pairing re-entrance at finite temperature for nuclei which have weak or no pairing at zero temperature. The best nuclei, in which such a phenomenon is expected, are those close to the drip line, as well as those located at a subshell closure as shown in this section.

In the following subsections, we evaluate the influence of the model on the pairing properties in hot finite nuclei, taking ^{124}Sn as an example for testing pairing correlations. The ratio of the critical temperature over the average pairing gap at zero temperature $T_c/\Delta_n(0)$ is also studied as a function of the model, varying either the Lagrangian or the type of pairing

interaction. Finally, a more systematic study of the critical temperature is performed on a set of semimagic nuclei.

A. Study of the ratio $T_c/\Delta_n(0)$

Since pairing correlations are active only around the Fermi level, the ratio $T_c/\Delta_n(0)$ is expected to be modified by the effective mass which influences the s.p. level spacing to a large extent. Notice that here the effective mass corresponds to the nonrelativistic one M_{NR}^* instead of the quantity $M_S^* = M + \Sigma_S$ that is named as the Dirac mass [40,55]. It can indeed be shown that, in the weak-coupling limit of the BCS approximation, the average pairing gap at the Fermi surface Δ_F can be expressed as [56]

$$\Delta_F \approx 2\epsilon_F \exp[2/(N_F v_{\text{pair}})], \quad (21)$$

where ϵ_F is the Fermi energy, $N_F = m_F^* k_F / \pi^2$ is the average density of state in uniform matter at the Fermi energy, k_F denotes the Fermi momentum, and v_{pair} is a constant pairing interaction. It is clear from Eq. (21) that the pairing gap Δ_F is quite sensitive to the effective mass at the Fermi energy ϵ_F . Equation (21) is obtained in infinite matter and it provides only a qualitative understanding of the relation between the pairing force strength and the effective mass. In the following, we present a quantitatively precise analysis of the correlation between the critical temperature and the nonrelativistic effective mass in finite nuclei.

Figure 1(a) displays the evolution of the neutron pairing gap as a function of temperature for ^{124}Sn , a good candidate for studying pairing correlations. We compare two Lagrangians, PKO2 and DD-ME2 (see Table I), both with two kinds of pairing interactions: the finite-range Gogny D1S force [41] and the contact force DDCI [42]. The two Lagrangians PKO2 and DD-ME2 mostly differ by their nonrelativistic effective mass M_{NR}^* (see Table I), and it is observed that the average pairing gap at zero temperature Δ_n scales with M_{NR}^* , as expected from




FIG. 1. (Color online) The neutron pairing gaps in ^{124}Sn as a function of temperature, calculated with PKO2 (black) and DD-ME2 (blue) corresponding to different M_{NR}^* . In the pairing channel, we compare finite-range D1S (filled circles) and contact DDCI (empty circles) forces. The analytical results (dashed lines) are also shown. Notice that here we took the factor $g = 1$ for the Gogny D1S force, and for the DDCI pairing, we have taken the following values for V_0 (in MeVfm^{-3}): 335 (PKO2) and 342 (DD-ME2) for the RH(F)B and BCS calculations. For the DDCIx force used in BCS, we took for V_0 (in MeVfm^{-3}): 526 (PKO2) and 539 (DD-ME2).

TABLE I. Bulk properties of symmetric nuclear matter at the saturation point: density ρ_0 (fm^{-3}), binding energy E_B/A (MeV), compression modulus K (MeV), symmetry energy J (MeV), and nonrelativistic effective masses M_{NR}^* (M) predicted by selected RHF and RH models. The nonrelativistic effective masses in neutron matter are also listed.

Model	Interaction	Ref.	Symmetric matter					Neutron matter	
			ρ_0	E_B/A	K	J	M_{NR}^*	$M_{\text{NR}}^*(\nu)$	$M_{\text{NR}}^*(\pi)$
RHF	PKA1	[43]	0.160	-15.83	229.96	36.02	0.68	0.68	0.70
	PKO1	[40]	0.152	-16.00	250.28	34.37	0.75	0.73	0.76
	PKO2	[51]	0.151	-16.03	249.53	32.49	0.76	0.75	0.77
RH	DD-ME2	[52]	0.152	-16.14	250.97	32.31	0.65	0.64	0.70
	PK1r	[53]	0.148	-16.27	283.68	37.83	0.68	0.64	0.72
	NL3*	[54]	0.150	-16.30	258.56	38.70	0.67	0.63	0.72

the weak-coupling expression (21). Comparing the different types of pairing interaction (finite or zero range) for the same Lagrangian, it is observed in Fig. 1 that the vanishing of pairing correlations at finite temperature slightly depends on the type of pairing force, namely with the zero-range pairing interaction, the critical temperature T_c is slightly lower than with the finite-range interaction. From Fig. 1, the ratio $T_c/\Delta_n(0)$ is obtained as 0.60 for D1S and 0.57 for DDCI. In Ref. [57], it was shown that the dependence of the pairing gap on the state around the Fermi energy qualitatively differs for contact and finite-range interactions. In addition, with a finite-range interaction, the pairing gap and the pairing tensor have nonlocal components which cannot be simply absorbed in the DDCI. The slight increase of the critical temperature with the D1S interaction is therefore an effect of the finite-range nature of the interaction, dispersing the pairing effects among more s.p. states. For comparison, and taking $T_c = 0.60\Delta_n(0)$ (finite-range pairing force) and $T_c = 0.57\Delta_n(0)$ (zero-range pairing forces), the analytical relation [58],

$$\Delta_n(T) = \Delta_n(0) \left[1 - \left(\frac{T}{T_c} \right)^m \right]^{1/2} \Theta(T - T_c), \quad (22)$$

where $m = 3.32$, is plotted for the DD-ME2 model [dashed lines in Fig. 1(a)]. There is almost no difference between the analytical model, i.e., Eq. (22), and the numerical calculations for stable nuclei like ^{124}Sn .

Figure 1(b) shows the neutron pairing gap calculated with the RHF theory plus BCS pairing at finite temperature (FT-RHF-BCS), using the same DDCI interaction as in the FT-RHFB calculations shown in Fig. 1(a) and the modified one (DDCIx) with enhanced pairing strength parameter V_0 (see caption for more details). It is found that, with the same DDCI pairing interaction, the neutron pairing gaps determined by the BCS method are reduced to about half of the Bogoliubov results. Such a distinct difference is due to the fact that the off-diagonal couplings, which account for about half of the pairing correlations, are absent in the BCS pairing. As a result, the strength parameter V_0 in BCS calculations is usually larger than in HFB, see, for instance, Ref. [59]. We have therefore readjusted V_0 in the RHF-BCS calculation at zero temperature to obtain the same pairing gap as the RHFB prediction, leading to the DDCIx interaction in Fig. 1(b). Applying such a simple modification of the parameter V_0 in ^{124}Sn , the temperature dependence of the pairing gap predicted by the FT-RHFB

and FT-RHF-BCS frameworks are almost undistinguishable. However, this is not always true and as we will see that the above simple renormalization of the pairing strength will not work towards the drip line where the coupling to continuum states becomes more and more important.

The critical temperatures is also expected to depend on the nonrelativistic effective mass, see, for instance, Refs. [22,25,60]. The dependence of the ratio $T_c/\Delta_n(0)$ on the effective mass is, however, not very well known. In Fig. 2, we plot the ratio $T_c/\Delta_n(0)$ as a function of M_{NR}^* in ^{124}Sn , and we compare the predictions of two pairing interactions (finite versus zero range). The effective mass M_{NR}^* is obtained with 14 CDFs (the 6 CDFs given in Table 1 completed with 8 other CDFs: PKO3 [51], PKDD [53], DD-ME1 [61], DD - ME δ [62], TW99 [63], PK1 [53], NL3 [64], and TM1 [65]). It is found that the ratio $T_c/\Delta_n(0)$ does not depend much on M_{NR}^* . Similar analyses were also carried out to check the relations with the incompressibility modulus K and the symmetry energy J , but no evidence of correlation is found. The small fluctuations observed in Fig. 2 are therefore mostly shell effects. In conclusion, the critical temperature in stable nuclei scales very well with the average pairing gap at zero temperature, the shell effects contributing to a dispersion of less than 4%.

In order to check whether the ratio $T_c/\Delta_n(0)$ is still constant for larger pairing strength, we have artificially varied the pairing interaction strength and correlated the critical




FIG. 2. (Color online) The ratios $T_c/\Delta_n(0)$ in ^{124}Sn as a function of the nonrelativistic effective mass M_{NR}^* , calculated using the FT-RH(F)B theory with 14 parameter sets. In the pairing channel the finite-range D1S and the contact force DDCI are employed.




FIG. 3. (Color online) The critical temperature T_c (left axes, blue circles) and the occupation number of continuum states N_{con} (right axes, black squares) at zero temperature in ^{124}Sn as a function of the neutron pairing gap calculated from PKO1 FT-RHFB with Gogny D1S (a) and DDCI (b) pairing forces. The dashed lines correspond to the relation $T_c = 0.60(0.57)\Delta_n(0)$. The strength parameter of the pairing interaction is varied to obtain different values of the pairing gap at zero temperature.

temperature T_c with the neutron pairing gap $\Delta_n(0)$ at zero temperature. Figures 3(a) and 3(b), respectively, show the results calculated with the pairing interactions D1S and DDCI using the PKO1 Lagrangian, where the average pairing gap $\Delta_n(0)$ goes from 100 keV to about 3.5 MeV. At the low pairing gap, the ratios $T_c/\Delta_n(0)$ are consistent with the analytical ones (dashed lines), as expected, until small deviations emerge beyond $\Delta_n(0) \sim 2.5$ MeV, where the continuum contributions become sizable (filled squares). These results are consistent with previous findings [24] based on a separable version of the Gogny D1S pairing force [66] and RH Lagrangian with PC-PK1 point coupling [67]. A simple calculation in infinite matter with a contact pairing interaction gives a correction to the ratio [68]

$$\frac{T_c}{\Delta(0)} \approx 0.57 \left[1 - \frac{1}{4\omega_D^2} \Delta(0)^2 \right], \quad (23)$$

where ω_D represents the pairing window. According to Eq. (23), the next-to-leading order correction has a negative sign, at variance with our results in ^{124}Sn , see Fig. 3. Notice that for the DDCI pairing interaction, the pairing window is about 100 MeV. The correction to the linear approximation of Eq. (23) is therefore very small: For the maximal pairing gap considered in this work [$\Delta(0) \approx 4.0$ MeV], the correction represents no more than a few percentages of the linear leading term.

In Fig. 3, the increase of the critical temperature for $\Delta_n(0) > 2.0$ MeV reveals an enhancement of the thermal pairing correlations, as well as the important role played by the continuum states. We remind that the next-to-the-leading order correction appearing in the simple expression in uniform matter (23) is negative. However, Fig. 3 shows a continuous enhanced continuum effect with respect to the pairing gap $\Delta_n(0)$. To better stress the increasing contribution of the continuum states and their role in the pairing correlations, we have introduced the following two quantities: the cumulative




FIG. 4. Contributions, in ^{124}Sn , of the continuum states to the pairing number \tilde{N}_{con} (a) and to the neutron number N_{con} (b), normalized to their value at zero temperature, as a function of the temperature $T/0.6\Delta(0)$. The results correspond to PKO1 Lagrangian and D1S pairing interaction.

occupation number of the neutron continuum states N_{con} ,

$$N_{\text{con}} = \sum_{a, \epsilon_a \geq 0} \int 4\pi r^2 \rho_a(r) dr, \quad (24)$$

and the cumulative pairing-occupation number of the neutron continuum states \tilde{N}_{con} ,

$$\tilde{N}_{\text{con}} = \sum_{a, \epsilon_a \geq 0} \int 4\pi r^2 \kappa_a(r) dr. \quad (25)$$

Here the continuum states a are determined in the $T = 0$ canonical basis, for simplicity [25,47], with s.p. energy ϵ_a above the continuum threshold. The increasing role of the continuum states at finite temperature is illustrated in Fig. 4 where the contributions to the pairing number \tilde{N}_{con} [Fig. 4(a)] and to the neutron number N_{con} [Fig. 4(b)] from the continuum states in ^{124}Sn , normalized to their values at zero temperature, are shown. Results with a weak pairing [$\Delta_n(0) = 1.3$ MeV] and with a stronger pairing [$\Delta_n(0) = 3.0$ MeV] are compared. Even if in the latter case the pairing is slightly larger than the expected value in finite nuclei, its inclusion in our analysis helps to understand the role of the continuum states. For the weak-pairing case, the continuum effects are very small, see Fig. 3, and both N_{con} and \tilde{N}_{con} drop to zero at the expected value $T_c = 0.6\Delta(0)$, see Fig. 4. For the strong-pairing case, a clear correlation is observed in Fig. 4 between the increase of the occupation of the continuum states at finite temperature N_{con} [see Fig. 4(b)] and the persistence of the pairing numbers \tilde{N}_{con} [see Fig. 4(a)]. The persistence of pairing correlations below the critical temperature modifies also the critical temperature, and Fig. 4(a) shows a larger value of the quantity $T/0.6\Delta(0)$ where pairing correlations in the continuum space drops to zero in the case of strong pairing compared to the weak one. Coming back to Fig. 3, we now understand better the correlation between the slight deviations of T_c from the analytical behaviors and enhanced

continuum effects for $\Delta(0) \gtrsim 2.5$ MeV. Since the presence of resonance states in the continuum is a typical feature of finite systems, the increase observed in Fig. 3, which differs from the prediction of Eq. (23), is then expected only in finite systems. Anticipating the results shown in the next figures, a similar enhancement of pairing correlations in other nuclei will also be observed, revealing here also the role of the resonant states.

B. Evolution of the critical temperature in isotopic and isotonic chains

In this subsection, we perform a systematic analysis of the evolution of the pairing gaps and critical temperature along isotopic and isotonic chains of semimagic nuclei. Through this extensive analysis, we have access to various s.p. configurations going from stable nuclei towards weakly bound drip-line nuclei, and we probe the pairing correlations inside various major shells. We consider three models, PKA1, PKO1 (RHF), and DD-ME2 (RH), which have different symmetry energies and nonrelativistic effective masses, see Table I. Anticipating our results, we will see that these models lead to rather different predictions for the pairing gap Δ .

Our results are shown for PKA1 (Fig. 5), PKO1 (Fig. 6), and DD-ME2 (Fig. 7) models, with the pairing channel described by the D1S interaction. In Figs. 5–7, we have represented isotopic (Ni, Sn, and Pb) and isotonic ($N = 50, 82,$ and 126) average pairing gap evolution as a function of N (for isotopes) and Z (for isotones). The isotopic and isotonic chains are bounded by the drip lines predicted by each of the considered models and determined by the two-nucleon separation energy. These drip lines are consistent, within a few units of uncertainty, with predictions given by other models obtained with Skyrme forces [69,70], Gogny forces [71,72], and RH Lagrangians [50,73]. We have calculated the average pairing gap at zero temperature (filled black circles) defined from Eq. (17) and compared the calculated critical temperature (filled blue circles) with the approximate relation, i.e., $0.6\Delta(0)$ (green curves). The arch structure of the results shown in Figs. 5–7 reflect the presence of magic numbers where pairing correlations completely vanish.

The PKA1 model (Fig. 5) is the most complete RHF version of the CDF theory. It contains the ρ - N Lorentz tensor coupling which is known to enhance the spin-orbit splitting [43,74,75]: in many cases the subshell structure is found to be closer to the experimental data than those predicted by other models without the ρ - N Lorentz tensor coupling, such as the RH approaches shown in Fig. 7. These subshell structures are clearly visible in Figs. 5–7 since they induce a partial quenching of the pairing gap for the associated submagic numbers. Going towards the drip lines, a reduction of the pairing gaps is often observed, revealing the presence of closed-shell nuclei at or near the drip lines. For the neutron drip line, it is the case of Sn and Pb isotopes, and for the proton drip line, it is observed for $N = 50$.

We first discuss the pairing properties of finite nuclei at zero temperature, which are influenced by the underlying s.p. structure around the Fermi energy. For the Ni isotopes, a subshell closure at $N = 40$ is predicted with PKA1 and PKO1 Lagrangians, as expected from experiments [76], while




FIG. 5. (Color online) Comparison of $\Delta(0)$ (black circles), $0.60\Delta(0)$ (green curves), and T_c (blue circles) in the even-even Ni, Sn, Pb isotopes (left panels) and $N = 50, 82, 126$ isotones (right panels), calculated in FT-RHFB with PKA1 and the Gogny pairing interaction D1S.

DD-ME2 shows a more pronounced shell closure. For neutron-rich Ni isotopes, PKA1 indicates another subshell closure at $N = 62$ which is not seen with PKO1 or DD-ME2. It is, however, beyond the present experimental limits. For the Sn isotopes, a decrease of the pairing gap by subshell closure is observed at $N = 64$ with PKO1 and DD-ME2 Lagrangians but not with PKA1. For Pb, a small decrease of the pairing gap is observed at $N = 138$ with PKA1 and at $N = 146$ with PKO1 and DD-ME2. On the isotonic side, we observe a subshell closure at $Z = 40$ for $N = 50$ with PKA1 Lagrangian but not with PKO1 or DD-ME2. For $N = 82$ isotones, PKA1 predicts a well-marked shell closure at $Z = 40$ and a subshell closure at $Z = 34$ and $Z = 64$, while at $Z = 58$ PKO1 predicts a shell closure and DD-ME2 only a subshell closure. Finally, for $N = 126$, PKO1 and DD-ME2 indicate a reduction of the pairing gap at $Z = 92$, which is not confirmed by experimental data [77] and is not present with PKA1 Lagrangian. We will see below that these structures can have an impact on the thermal properties.




FIG. 6. (Color online) The same as in Fig. 5 but calculated in the FT-RHFB theory with the effective interaction PKO1.

Turning now to the thermal properties of these isotopes and isotones, the comparison of the calculated critical temperature T_c and the approximate relation $0.60\Delta(0)$ shown in Figs. 5–7 exhibits some interesting features. The critical temperature T_c and the approximate relation $0.60\Delta(0)$ are identical in most cases with some exceptions. In heavy nuclei (Pb and $N = 126$), there are no strong deviations between these two quantities, but they are, however, more marked in lighter nuclei. Moreover, the cases where the exact and the approximate values of T_c differ are correlated with either the presence of a subshell closure or with the proximity of the drip lines. In the case of subshell closure, the effect of the temperature is to “wash out” the decrease of the pairing correlations. This can be understood as the consequence of the thermal occupation probabilities which overcome small shell gaps. Close to the neutron drip lines, the more pronounced effects are observed in Ni and Sn isotopes. This is also due to the thermal occupation of close-by resonant states as discussed in Refs. [14,78]. In the nonrelativistic Skyrme Hartree-Fock plus BCS (SHF-BCS) approach, an enhancement of the critical temperature was found in ^{140}Sn using SKT6 [22]. We do not confirm this enhancement in ^{140}Sn with the models used in




FIG. 7. (Color online) The same as in Fig. 5 but calculated in the FT-RHFB theory with the effective interaction DD-ME2.

this work. However, it is interesting to notice that the origin of such an enhancement found in Ref. [22] is also related to the existence of a subshell closure.

Let us finish this subsection with some general remarks concerning the nuclei which do not manifest any enhancement of the critical temperature. For the Pb isotopes, as shown in Figs. 5–7, we have not observed any marked enhancement of the critical temperature near the drip line as in the case of Sn isotopes. Comparing Pb and Sn, since the pairing gap for these isotopes is decreasing near the drip line, one could have expected to observe an enhancement of the critical temperature in Pb as it is observed in Sn. For instance, the last occupied state in the drip-line nucleus ^{266}Pb is indeed found to be well bound ($\epsilon_{3d_{3/2}} < -2.0$ MeV), and the lowest s.p. resonance $\epsilon_{2h_{1/2}}$ is found to be above 1.5 MeV. There is therefore a rather strong gap in the neutron-rich Pb isotopes ($N = 184$) which prevents the large coupling to the continuum. For the isotonic chains, we do not find any pairing persistence phenomenon around the drip line. This can be related to the well-developed shell closures at $Z = 50, 82,$ and 92 for proton-rich nuclei, which quench the coupling to continuum states, as it would have been expected in such exotic nuclei. In addition, the coupling to the

continuum is weaker for protons since the Coulomb barrier tends to localize the proton density in the nuclear interior [79]. For these reasons, the persistence phenomenon is strongly quenched in the proton channel.

C. Pairing persistence in ^{68}Ni and ^{174}Sn

In this subsection, we analyze in more detail the temperature dependence of the pairing gap for two representative nuclei, ^{68}Ni and ^{174}Sn . On the one hand, ^{68}Ni is an isotope which is slightly more neutron rich than the five stable isotopes $^{58-64}\text{Ni}$. As shown in the previous subsection, ^{68}Ni is considered a subshell isotope [50,80,81], and, as a consequence, the pairing gap at zero temperature is either reduced or strongly quenched depending on the model, see Figs. 5–7. On the other hand, ^{174}Sn is a very neutron-rich isotope at or close to the neutron drip line, where the continuum effects are expected to be remarkable [14,15,57,82]. However, since ^{174}Sn is close to the potentially doubly magic ^{176}Sn ($Z = 50$, $N = 126$), a gap is expected to be present in the s.p. structure between bound and resonant states. These two nuclei are therefore representative of quantum systems for which pairing at zero temperature is weakened by the presence of a gap above the Fermi energy and where a finite amount of temperature allows us to overcome the gap and provokes an enhancement of pairing correlations, giving rise to pairing persistence.

We first show the temperature dependence of the pairing gap for ^{68}Ni in Fig. 8(a), calculated with the FT-RHF model where we consider the PKO1 Lagrangian in the mean-field channel and either Gogny D1S or DDCI interaction in the pairing channel. The analytical model is also shown for reference. It is found that the predictions for $\Delta_n(T)$ do not practically depend on the pairing force. The critical temperature predicted by the FT-RHF approach is increased with respect to the reference analytical model: the FT-RHF pairing gaps vanish around $T = 0.90$ MeV, which is 0.25 MeV higher than the expected value (0.65 MeV). The pairing gap predicted by FT-RHF-BCS is shown in Fig. 8(b). Surprisingly, the pairing gap is zero if the same DDCI pairing interaction

is used. An increase of the pairing strength V_0 is therefore necessary. It is also interesting to observe that the DDCI pairing interaction, where V_0 is increased to match with the zero temperature pairing gap obtained with FT-RHF case, reproduces almost exactly the temperature dependence of the FT-RHF case and predicts as well an increase of the critical temperature with respect to the analytical model. The nucleus ^{68}Ni is a typical example of pairing persistence at finite temperature in closed subshell ($N = 40$) nuclei. We hereafter name this phenomenon type I pairing persistence. Other examples of similar behavior are the following: ^{90}Ni (PKA1), ^{114}Sn (PKO1, DD-ME2), ^{220}Pb (PKA1), and ^{230}Pb (DD-ME2) for the neutron pairing gap and ^{90}Zr (PKA1, PKO1 and DD-ME2), ^{140}Ce (DD-ME2), and $^{146,190}\text{Gd}$ (PKA1) for the proton pairing gap.

We turn now to the analysis of the results in ^{174}Sn . As stated above, this is a nucleus where pairing correlations are slightly weakened due to the proximity of shell closure. A small amount of temperature is expected to reorganize the level occupancy around the Fermi energy, opening more space below the Fermi energy and producing a nonzero occupancy of the first levels above the Fermi energy which are in the continuum. Most of the occupied states in the continuum are resonance states, but it is interesting to notice that a small number of them are also nonresonant states [83]. Without the participation of these nonresonant states, the asymptotic behavior of the density would be ill defined and present an unexpected gas component, as is also observed in the BCS theory [57]. The Bogoliubov transformation couples all states in a sub- (l, j) space and a truncation among these states breaks the unitarity of this transformation. To avoid the presence of nonphysical gas component in the density profile, it is therefore important to couple all states in the continuum within the Bogoliubov transformation.

Figure 9(a) displays the evolution of the neutron pairing gap as a function of temperature in ^{174}Sn with different Lagrangians and pairing forces. The results are very weakly affected by the choice of the pairing interaction. The effect induced by the choice of Lagrangian is also very small.




FIG. 8. (Color online) The neutron pairing gaps in ^{68}Ni as a function of temperature, calculated by FT-RHF with Gogny D1S and DDCI pairing forces, and by FT-RHF-BCS with DDCI pairing force. To evaluate the persistence provoked by the subshell, the analytical results are also shown. The pairing strength V_0 (in MeV fm^{-3}) is fixed to be 326 (DDCI) and 537 (DDCIx).




FIG. 9. (Color online) The neutron pairing gaps in ^{174}Sn as a function of temperature, calculated with PKO1 and NL3*, using Gogny D1S and DDCI pairing forces. The results of the analytical model are also shown. The pairing strength V_0 (in MeV fm^{-3}) is 333 (DDCI with PKO1), 317 (DDCI with NL3*), and 596 (DDCIx with the PKO1 and BCS frameworks).

We find a systematic increase of the critical temperature ($T_c \approx 0.65\text{--}0.70$ MeV) with respect to that expected from the analytical relation ($T_c \approx 0.47$ MeV), independently of the considered model. In addition, for temperature $T > 0.2$ MeV, an increase of the pairing gap is observed, which can also be related to the thermally induced contribution of the continuum states. We compare these results to the ones obtained with the FT-RHF-BCS framework shown in Fig. 9(b). As already discussed, the DDCI interaction predicts a reduced pairing gap in BCS compared to RHFB. On the other side, the DDCI χ interaction where the pairing strength is increased to match the $T = 0$ predictions of RHFB leads to an overestimation of the critical temperature compared to the FT-RHFB case. It shows that, in the case of the drip-line nucleus, the RHFB calculation cannot be simply reproduced by a BCS calculation where the pairing strength is just increased. Since the coupling to the continuum plays a dominant role in the persistence phenomenon in ^{174}Sn , we hereafter call it a type II phenomenon. From our results, it is also expected to occur in Ni and Sn neutron-rich nuclei, Ni ($N > 54 \sim 60$) and Sn ($N > 100$).

D. Entropy and specific heat

We now focus on the entropy S and specific heat C_v , which are the first and second derivatives of the free energy F with respect to the temperature, and thus sensitive to thermal changes of the ground state, see, for instance, Ref. [84] and Refs. therein. To test the sensitivity of these quantities to the choice of different models, we select two Sn isotopes, ^{120}Sn and ^{160}Sn . The former is a good example of a stable well-paired nucleus, while the latter is yet far from actual nuclear experiments but represents an extreme case with large isospin asymmetry.

In Fig. 10 are shown the entropy S and the specific heat C_v as functions of the temperature calculated by the RHF functionals PKO1 and PKO2, the RH ones with the nonlinear self-couplings PK1r and NL3*, and the RH one with density-dependent meson-nucleon couplings DD-ME2. In the pairing channel, the value of the scaling factor g (see the first

paragraph of Sec. III), is slightly modified to give identical pairing gaps at zero temperature for the different models. In Fig. 10(a), the entropy calculated with the pairing correlations [labeled FT-RH(F)B] is compared to that neglecting the pairing correlations [labeled FT-RH(F)]. At low temperature, if the pairing effects are ignored, the entropy is found to be largely model dependent, i.e., the model with smaller nonrelativistic effective mass (see Table I), which leads to larger s.p. level spacing on the average, presents smaller entropy. As the temperature increases, and also as the pairing correlations are switched on, the entropy becomes less model dependent.

In fact, at low temperature or without pairing correlations, the entropy is largely determined by the few states around the Fermi energy, and the number of the involved states is essentially determined by the detailed s.p. spectrum which depends on the models, therefore leading to model-dependent entropy. Both temperature and pairing correlations can disperse the particle over the states beyond the Fermi level. As the temperature increases, and/or as the pairing correlations are enhanced, more s.p. states will get involved to contribute to the entropy, and the average properties such as the density of states will become dominant, instead of a few states as in the FT-RH(F) cases. Compared to distinctly different s.p. spectra around the Fermi surface, the dispersions of the nonrelativistic effective masses (see Table I) between the models are less remarkable. However, in Fig. 10(a) it is clearly shown that the FT-RH(F)B results are grouped by the values of the effective masses when $T \gtrsim 1$ MeV, which correspond to different average densities of states. As expected, the effect of the pairing correlations is clearly visible below the critical temperature ($T_c \approx 0.8$ MeV), inducing a strong reduction of the entropy [see Fig. 10(a)] and singular behaviors of the specific heat around the critical temperature as shown in Fig. 10(b). Just above the critical temperature, we can notice that the specific heat is not linear in T , as expected from the Fermi gas model [85,86], and the linear dependence seems to be found at slightly larger temperature ($T > 1.5$ MeV). The nonlinearity of the specific heat around T_c might be related to shell effects.

The results thus clearly show that the pairing correlations contribute to the S-shaped behavior of the specific heat, as has already been inferred from the analysis of thermal excited nuclei in laboratory experiments [8,10,86]. A realistic description of the smooth S-shaped behavior in finite nuclei requires a more elaborated modeling, including, for instance, particle number projection [7,87–89]. It is, however, shown in Ref. [29] that the smooth S-shaped behavior may be even washed out in some rare-earth nuclei. The results presented in Fig. 10 should not be compared directly to the semiexperimental data.

The situation for the neutron-rich nucleus ^{160}Sn is more complex, as shown in Fig. 11. The model dependence of entropy at low temperature is reduced compared to the case of ^{120}Sn . Only PKO2 predictions differ from the other models. The predictions for the critical temperature, associated to the discontinuity in the specific heat of Fig. 11, vary among the models to a much larger extent than what was found in ^{120}Sn . ^{160}Sn is located in the region where pairing persistence is expected to appear, see Figs. 5–7. Since this




FIG. 10. (Color online) Entropy and specific heat in ^{120}Sn as a function of temperature, calculated using the FT-RH(F)B and FT-RH(F) theories.




FIG. 11. (Color online) The same as in Fig. 10 but for ^{160}Sn .

phenomenon is strongly related to the position of resonance states in the continuum, we expect to observe deviations among models predicting different positions of these states. This model dependence therefore reveals our lack of knowledge in extrapolating Lagrangians which have been adjusted for less exotic nuclei.

IV. SUMMARY and CONCLUSION

In this work, we have developed, as far as we know, the first FT-RHFB theory for spherical nuclei. The self-consistent FT-RHFB equations are solved by using a DWS basis which provides an appropriate asymptotic behavior for the continuum states. We have performed systematic FT-RHFB calculations for both stable and weakly bound nuclei and discussed their thermal properties. The influence of the pairing interaction on the pairing phase transition is evaluated. It is found that the critical temperature for a pairing transition generally follows the rule $T_c = 0.60\Delta(0)$ with a finite-range pairing force and $T_c = 0.57\Delta(0)$ with a contact pairing force. The finite- or zero-range nature of the pairing force, while generating different state-dependence pairing gaps, causes only small differences in our results. We have described the pairing persistence in two kinds of situations: nuclei at subshell closure (type I) and nuclei strongly coupled to continuum states which are close to the drip line (type II). We have observed that, while a refitting of the pairing strength could match the FT-RHF-BCS with the FT-RHFB predictions for the pairing gap in case I, it is no longer true in case II. This is due to the participation of the

continuum states in the second case which involve coupling of a different nature.

We have also analyzed the influence of the interaction on the thermal response. The results show clearly that the pairing correlations contribute to the S-shaped behavior of the specific heat curve and help to wash out the model dependence. For stable nuclei the model deviations, to some extent, can be traced back to the effective mass, since the level structure only weakly depends on the choice of the CDF. The situation for exotic nuclei is more complex since it is related to our lack of knowledge in very exotic nuclei, and the pairing persistence would have large effects on their thermal property.

In conclusion, we have illustrated the richness and complexity of pairing correlations at finite temperature and in finite systems within the FT-RHFB calculation. The discussion of correlations beyond mean field, induced, for instance, by particle number projection, is not addressed in this work. It is, however, expected that the particle number projection will contribute to an increase of the pairing correlations in the case where they are weak [29], like in the pairing persistence phenomenon discussed in this paper. In future work, a more quantitative calculation will be necessary to estimate the strength these additional correlations and how they modify the results presented in this work. Another interesting perspective which is suggested by this work is the possibility that similar phenomena can be observed in other domains of physics. For instance, it was studied whether cold atoms in a double potential could demonstrate pairing persistence as well [90]. Finally, the application of this formalism for the prediction of temperature evolution of pairing properties in the crust of neutron stars [11,91,92] will be performed in the near future. There, the thermal modification of pairing correlations could have a large impact on the thermal relaxation of the crust [12] and could be observed during the quiescent period of low mass x-ray transients.

ACKNOWLEDGMENTS

J.L. acknowledges support by China Scholarship Council (CSC) under Grant No. 201206180063. This work was partially supported by the NewCompStar COST action and the ANR project SN2NS No. ANR-10-BLAN-0503. Research at LZU was supported by the National Natural Science Foundation of China under Grants No. 11375076 and No. 11405223 and by the Specialized Research Fund for the Doctoral Program of Higher Education under Grant No. 20130211110005.

[1] E. Melby, L. Bergholt, M. Guttormsen, M. Hjorth-Jensen, F. Ingebretsen, S. Messelt, J. Rekstad, A. Schiller, S. Siem, and S. W. Ødegård, *Phys. Rev. Lett.* **83**, 3150 (1999).
 [2] U. Agvaanluvsan, A. Schiller, J. A. Becker, L. A. Bernstein, P. E. Garrett, M. Guttormsen, G. E. Mitchell, J. Rekstad, S. Siem, A. Voinov *et al.*, *Phys. Rev. C* **70**, 054611 (2004).
 [3] Y. Kalmykov, C. Özen, K. Langanke, G. Martínez-Pinedo, P. von Neumann-Cosel, and A. Richter, *Phys. Rev. Lett.* **99**, 202502 (2007).

[4] H. K. Toft, A. C. Larsen, U. Agvaanluvsan, A. Bürger, M. Guttormsen, G. E. Mitchell, H. T. Nyhus, A. Schiller, S. Siem, N. U. H. Syed *et al.*, *Phys. Rev. C* **81**, 064311 (2010).
 [5] M. Guttormsen, A. C. Larsen, F. L. B. Garrote, Y. Byun, T. K. Eriksen, F. Giacompo, A. Görgen, T. W. Hagen, M. Klintefjord, H. T. Nyhus *et al.*, *Phys. Rev. C* **90**, 044309 (2014).
 [6] R. Balian, H. Flocard, and M. Vénéroni, *Phys. Rep.* **317**, 251 (1999).
 [7] S. Liu and Y. Alhassid, *Phys. Rev. Lett.* **87**, 022501 (2001).

- [8] M. Guttormsen, M. Hjorth-Jensen, E. Melby, J. Rekstad, A. Schiller, and S. Siem, *Phys. Rev. C* **64**, 034319 (2001).
- [9] D. J. Dean and M. Hjorth-Jensen, *Rev. Mod. Phys.* **75**, 607 (2003).
- [10] U. Agvaanluvsan, A. C. Larsen, M. Guttormsen, R. Chankova, G. E. Mitchell, A. Schiller, S. Siem, and A. Voinov, *Phys. Rev. C* **79**, 014320 (2009).
- [11] N. Sandulescu, *Phys. Rev. C* **70**, 025801 (2004).
- [12] M. Fortin, F. Grill, J. Margueron, D. Page, and N. Sandulescu, *Phys. Rev. C* **82**, 065804 (2010).
- [13] A. Pastore, S. Baroni, and C. Losa, *Phys. Rev. C* **84**, 065807 (2011).
- [14] J. Margueron and E. Khan, *Phys. Rev. C* **86**, 065801 (2012).
- [15] A. Pastore, J. Margueron, P. Schuck, and X. Viñas, *Phys. Rev. C* **88**, 034314 (2013).
- [16] A. Sedrakian, T. Alm, and U. Lombardo, *Phys. Rev. C* **55**, R582 (1997).
- [17] N. Q. Hung and N. D. Dang, *Phys. Rev. C* **84**, 054324 (2011).
- [18] M. Sano and S. Yamasaki, *Prog. Theor. Phys.* **29**, 397 (1963).
- [19] A. L. Goodman, *Nucl. Phys. A* **352**, 30 (1981).
- [20] A. L. Goodman, *Phys. Rev. C* **34**, 1942 (1986).
- [21] J. L. Egido and P. Ring, *J. Phys. G: Nucl. Part. Phys.* **19**, 1 (1993).
- [22] C. Reiß, M. Bender, and P.-G. Reinhard, *Eur. Phys. J. A* **6**, 157 (1999).
- [23] E. Khan, N. Van Giai, and N. Sandulescu, *Nucl. Phys. A* **789**, 94 (2007).
- [24] Y. F. Niu, Z. M. Niu, N. Paar, D. Vretenar, G. H. Wang, J. S. Bai, and J. Meng, *Phys. Rev. C* **88**, 034308 (2013).
- [25] Y. K. Gambhir, J. P. Maharana, G. A. Lalazissis, C. P. Panos, and P. Ring, *Phys. Rev. C* **62**, 054610 (2000).
- [26] J. L. Egido, L. M. Robledo, and V. Martin, *Phys. Rev. Lett.* **85**, 26 (2000).
- [27] B. K. Agrawal, T. Sil, S. K. Samaddar, and J. N. De, *Phys. Rev. C* **63**, 024002 (2001).
- [28] V. Martin, J. L. Egido, and L. M. Robledo, *Phys. Rev. C* **68**, 034327 (2003).
- [29] D. Gambacurta, D. Lacroix, and N. Sandulescu, *Phys. Rev. C* **88**, 034324 (2013).
- [30] J. Dobaczewski, H. Flocard, and J. Treiner, *Nucl. Phys. A* **422**, 103 (1984).
- [31] M. Stoitsov, P. Ring, D. Vretenar, and G. A. Lalazissis, *Phys. Rev. C* **58**, 2086 (1998).
- [32] S.-G. Zhou, J. Meng, and P. Ring, *Phys. Rev. C* **68**, 034323 (2003).
- [33] N. Schunck and J. L. Egido, *Phys. Rev. C* **77**, 011301 (2008).
- [34] T. Nikšić, N. Paar, D. Vretenar, and P. Ring, *Comput. Phys. Commun.* **185**, 1808 (2014).
- [35] A. Bouyssy, J.-F. Mathiot, N. Van Giai, and S. Marcos, *Phys. Rev. C* **36**, 380 (1987).
- [36] P. Ring, *Prog. Part. Nucl. Phys.* **37**, 193 (1996).
- [37] M. Bender, P.-H. Heenen, and P.-G. Reinhard, *Rev. Mod. Phys.* **75**, 121 (2003).
- [38] D. Vretenar, A. V. Afanasjev, G. A. Lalazissis, and P. Ring, *Phys. Rep.* **409**, 101 (2005).
- [39] J. Meng, H. Toki, S. G. Zhou, S. Q. Zhang, W. H. Long, and L. S. Geng, *Prog. Part. Nucl. Phys.* **57**, 470 (2006).
- [40] W. H. Long, N. Van Giai, and J. Meng, *Phys. Lett. B* **640**, 150 (2006).
- [41] J. F. Berger, M. Girod, and D. Gogny, *Nucl. Phys. A* **428**, 23 (1984).
- [42] N. Tajima, P. Bonche, H. Flocard, P.-H. Heenen, and M. S. Weiss, *Nucl. Phys. A* **551**, 434 (1993).
- [43] W. H. Long, H. Sagawa, N. V. Giai, and J. Meng, *Phys. Rev. C* **76**, 034314 (2007).
- [44] H. Kucharek and P. Ring, *Z. Phys. A* **339**, 23 (1991).
- [45] W. H. Long, P. Ring, N. Van Giai, and J. Meng, *Phys. Rev. C* **81**, 024308 (2010).
- [46] J.-P. Ebran, E. Khan, D. Peña Arteaga, and D. Vretenar, *Phys. Rev. C* **83**, 064323 (2011).
- [47] P. Bonche, S. Levit, and D. Vautherin, *Nucl. Phys. A* **427**, 278 (1984).
- [48] M. Bender, K. Rutz, P.-G. Reinhard, and J. A. Maruhn, *Eur. Phys. J. A* **7**, 467 (2000).
- [49] L. J. Wang, B. Y. Sun, J. M. Dong, and W. H. Long, *Phys. Rev. C* **87**, 054331 (2013).
- [50] S. E. Agbemava, A. V. Afanasjev, D. Ray, and P. Ring, *Phys. Rev. C* **89**, 054320 (2014).
- [51] W. Long, H. Sagawa, J. Meng, and N. Van Giai, *Europhys. Lett.* **82**, 12001 (2008).
- [52] G. A. Lalazissis, T. Nikšić, D. Vretenar, and P. Ring, *Phys. Rev. C* **71**, 024312 (2005).
- [53] W. Long, J. Meng, N. Van Giai, and S.-G. Zhou, *Phys. Rev. C* **69**, 034319 (2004).
- [54] G. A. Lalazissis, S. Karatzikos, R. Fossion, D. Pena Arteaga, A. Afanasjev, and P. Ring, *Phys. Lett. B* **671**, 36 (2009).
- [55] M. Jaminon and C. Mahaux, *Phys. Rev. C* **40**, 354 (1989).
- [56] U. Lombardo, in *Nuclear Methods and the Nuclear Equation of State*, edited by M. Baldo (World Scientific, Singapore, 1999), Chap. 9.
- [57] J. Dobaczewski, W. Nazarewicz, T. R. Werner, J. F. Berger, C. R. Chinn, and J. Dechargé, *Phys. Rev. C* **53**, 2809 (1996).
- [58] S. Goriely, *Nucl. Phys. A* **605**, 28 (1996).
- [59] G. F. Bertsch, C. A. Bertulani, W. Nazarewicz, N. Schunck, and M. V. Stoitsov, *Phys. Rev. C* **79**, 034306 (2009).
- [60] E. Yüksel, E. Khan, K. Bozkurt, and G. Coló, *Eur. Phys. J. A* **50**, 160 (2014).
- [61] T. Nikšić, D. Vretenar, P. Finelli, and P. Ring, *Phys. Rev. C* **66**, 024306 (2002).
- [62] X. Roca-Maza, X. Viñas, M. Centelles, P. Ring, and P. Schuck, *Phys. Rev. C* **84**, 054309 (2011).
- [63] S. Typel and H. H. Wolter, *Nucl. Phys. A* **656**, 331 (1999).
- [64] G. A. Lalazissis, J. König, and P. Ring, *Phys. Rev. C* **55**, 540 (1997).
- [65] Y. Sugahara and H. Toki, *Nucl. Phys. A* **579**, 557 (1994).
- [66] Y. Tian, Z. Y. Ma, and P. Ring, *Phys. Lett. B* **676**, 44 (2009).
- [67] P. W. Zhao, Z. P. Li, J. M. Yao, and J. Meng, *Phys. Rev. C* **82**, 054319 (2010).
- [68] G. D. Mahan, *Many-Particle Physics* (Plenum Press, New York, 2000).
- [69] J. Erler, N. Birge, M. Kortelainen, W. Nazarewicz, E. Olsen, A. M. Perhac, and M. Stoitsov, *Nature* **486**, 509 (2012).
- [70] J. Erler, C. J. Horowitz, W. Nazarewicz, M. Rafalski, and P.-G. Reinhard, *Phys. Rev. C* **87**, 044320 (2013).
- [71] S. Hilaire and M. Girod, *Eur. Phys. J. A* **33**, 237 (2007).
- [72] J. P. Delaroche, M. Girod, J. Libert, H. Goutte, S. Hilaire, S. Péru, N. Pillet, and G. F. Bertsch, *Phys. Rev. C* **81**, 014303 (2010).

- [73] A. V. Afanasjev, S. E. Agbemava, D. Ray, and P. Ring, *Phys. Lett. B* **726**, 680 (2013).
- [74] W. Long, T. Nakatsukasa, H. Sagawa, J. Meng, H. Nakada, and Y. Zhang, *Phys. Lett. B* **680**, 428 (2009).
- [75] J. J. Li, W. H. Long, J. Margueron, and N. Van Giai, *Phys. Lett. B* **732**, 169 (2014).
- [76] O. Sorlin, S. Leenhardt, C. Donzaud, J. Duprat, F. Azaiez, F. Nowacki, H. Grawe, Z. Dombrádi, F. Amorini, A. Astier *et al.*, *Phys. Rev. Lett.* **88**, 092501 (2002).
- [77] L.-S. Geng, J. Meng, H. Toki, W.-H. Long, and G. Shen, *Chin. Phys. Lett.* **23**, 1139 (2006).
- [78] A. Pastore, *Phys. Rev. C* **86**, 065802 (2012).
- [79] J. Dobaczewski, I. Hamamoto, W. Nazarewicz, and J. A. Sheikh, *Phys. Rev. Lett.* **72**, 981 (1994).
- [80] M. Kleban, B. Nerlo-Pomorska, J. F. Berger, J. Dechargé, M. Girod, and S. Hilaire, *Phys. Rev. C* **65**, 024309 (2002).
- [81] M. Yamagami, J. Margueron, H. Sagawa, and K. Hagino, *Phys. Rev. C* **86**, 034333 (2012).
- [82] J. Meng, H. Toki, J. Y. Zeng, S. Q. Zhang, and S.-G. Zhou, *Phys. Rev. C* **65**, 041302 (2002).
- [83] K. Bennaceur, J. Dobaczewski, and M. Płoszajczak, *Phys. Rev. C* **60**, 034308 (1999).
- [84] A. Pastore, N. Chamel, and J. Margueron, *Mon. Not. R. Astron. Soc.* **448**, 1887 (2015).
- [85] A. Gilbert and A. G. W. Cameron, *Can. J. Phys.* **43**, 1446 (1965).
- [86] A. Schiller, A. Bjerve, M. Guttormsen, M. Hjorth-Jensen, F. Ingebretsen, E. Melby, S. Messelt, J. Rekstad, S. Siem, and S. W. Ødegård, *Phys. Rev. C* **63**, 021306 (2001).
- [87] K. Tanabe, K. Sugawara-Tanabe, and H. J. Mang, *Nucl. Phys. A* **357**, 20 (1981).
- [88] C. Esebbag and J. L. Egidio, *Nucl. Phys. A* **552**, 205 (1993).
- [89] N. D. Dang and A. Arima, *Phys. Rev. C* **68**, 014318 (2003).
- [90] A. Pastore, P. Schuck, M. Urban, X. Viñas, and J. Margueron, *Phys. Rev. A* **90**, 043634 (2014).
- [91] F. Grill, J. Margueron, and N. Sandulescu, *Phys. Rev. C* **84**, 065801 (2011).
- [92] J. Margueron and N. Sandulescu, in *Neutron Star Crust*, edited by C. A. Bertulani and J. Piekarewicz (World Scientific, Singapore, 2012), Chap. 4.