

^7Be solar neutrino measurement with KamLAND

A. Gando,¹ Y. Gando,¹ H. Hanakago,¹ H. Ikeda,¹ K. Inoue,^{1,2} K. Ishidoshiro,¹ H. Ishikawa,¹ Y. Kishimoto,^{1,2,*} M. Koga,^{1,2} R. Matsuda,¹ S. Matsuda,¹ T. Mitsui,¹ D. Motoki,¹ K. Nakajima,^{1,†} K. Nakamura,^{1,2} A. Obata,¹ A. Oki,¹ Y. Oki,¹ M. Otani,^{1,‡} I. Shimizu,¹ J. Shirai,¹ A. Suzuki,¹ K. Tamae,¹ K. Ueshima,¹ H. Watanabe,¹ B. D. Xu,¹ S. Yamada,^{1,‡} Y. Yamauchi,¹ H. Yoshida,^{1,§} A. Kozlov,² Y. Takemoto,² S. Yoshida,³ C. Grant,^{4,¶} G. Keefer,^{4,¶} D. W. McKee,^{4,**} A. Piepke,^{4,2} T. I. Banks,⁵ T. Bloxham,⁵ S. J. Freedman,^{5,††} B. K. Fujikawa,^{5,2} K. Han,^{5,‡‡} L. Hsu,^{5,§§} K. Ichimura,^{5,*} H. Murayama,^{5,2} T. O'Donnell,⁵ H. M. Steiner,⁵ L. A. Winslow,^{5,¶¶} D. Dwyer,^{6,¶¶} C. Mauger,^{6,***} R. D. McKeown,^{6,†††} C. Zhang,^{6,‡‡‡} B. E. Berger,^{7,2} C. E. Lane,⁸ J. Maricic,^{8,9} T. Miletic,^{8,§§§} J. G. Learned,⁹ M. Sakai,⁹ G. A. Horton-Smith,¹⁰ A. Tang,¹⁰ K. E. Downum,¹¹ K. Tolich,^{11,12,¶¶¶} Y. Efremenko,^{13,2} Y. Kamyshkov,¹³ O. Perevozchikov,^{13,¶¶¶} H. J. Karwowski,¹⁴ D. M. Markoff,¹⁴ W. Tornow,^{14,2} J. A. Detwiler,¹² S. Enomoto,^{12,2} K. Heeger,^{15,‡‡} and M. P. Decowski^{16,2}

(KamLAND Collaboration)

¹Research Center for Neutrino Science, Tohoku University, Sendai, Miyagi 980-8578, Japan²Kavli Institute for the Physics and Mathematics of the Universe (WPI), University of Tokyo, Kashiwa, Chiba 277-8583, Japan³Graduate School of Science, Osaka University, Toyonaka, Osaka 560-0043, Japan⁴Department of Physics and Astronomy, University of Alabama, Tuscaloosa, Alabama 35487, USA⁵Physics Department, University of California at Berkeley, Berkeley, California 94720, USA, and Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA⁶Kellogg Radiation Laboratory, California Institute of Technology, Pasadena, California 91125, USA⁷Department of Physics, Colorado State University, Fort Collins, Colorado 80523, USA⁸Department of Physics, Drexel University, Philadelphia, Pennsylvania 19104, USA⁹Department of Physics and Astronomy, University of Hawaii at Manoa, Honolulu, Hawaii 96822, USA¹⁰Department of Physics, Kansas State University, Manhattan, Kansas 66506, USA¹¹Department of Physics, Stanford University, Stanford, California 94305, USA¹²Center for Experimental Nuclear Physics and Astrophysics, University of Washington, Seattle, Washington 98195, USA¹³Department of Physics and Astronomy, University of Tennessee, Knoxville, Tennessee 37996, USA¹⁴Triangle Universities Nuclear Laboratory, Durham, North Carolina 27708, USA, and Departments of Physics at Duke University, Durham, North Carolina 27708, USA, North Carolina Central University, Durham, North Carolina 27701, USA, and the University of North Carolina at Chapel Hill, Chapel Hill, North Carolina 27599, USA¹⁵Department of Physics, University of Wisconsin, Madison, Wisconsin 53706, USA¹⁶Nikhef and the University of Amsterdam, Science Park, Amsterdam, the Netherlands

(Received 23 May 2014; revised manuscript received 22 July 2015; published 30 November 2015)

We report a measurement of the neutrino-electron elastic scattering rate of 862 keV ^7Be solar neutrinos based on a 165.4 kt d exposure of KamLAND. The observed rate is 582 ± 94 (kt d)⁻¹, which corresponds to an 862-keV ^7Be solar neutrino flux of $(3.26 \pm 0.52) \times 10^9$ cm⁻² s⁻¹, assuming a pure electron-flavor flux. Comparing this flux with the standard solar model prediction and further assuming three-flavor mixing, a ν_e survival probability of 0.66 ± 0.15 is determined from the KamLAND data. Utilizing a global three-flavor oscillation analysis, we obtain a total ^7Be solar neutrino flux of $(5.82 \pm 1.02) \times 10^9$ cm⁻² s⁻¹, which is consistent with the standard solar model predictions.

DOI: [10.1103/PhysRevC.92.055808](https://doi.org/10.1103/PhysRevC.92.055808)

PACS number(s): 26.65.+t, 14.60.Pq, 25.40.Sc

*Present address (The same shall apply hereinafter.): Kamioka Observatory, Institute for Cosmic Ray Research, the University of Tokyo, Hida, Gifu 506-1205, Japan.

[†]Graduate School of Engineering, University of Fukui, Fukui 910-8507, Japan.

[‡]Institute of Particle and Nuclear Studies, High Energy Accelerator Research Organization, Tsukuba, Ibaraki 305-0801, Japan.

[§]Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan.

[¶]Physics Department, University of California, Davis, Davis, California 95616, USA.

^{¶¶}Lawrence Livermore National Laboratory, Livermore, California 94550, USA.

**Chemical and Physical Sciences Department, Missouri Southern State University, Joplin, Missouri 64801, USA.

^{††}Deceased.

^{‡‡}Department of Physics, Yale University, New Haven, Connecticut 06520, USA.

^{§§}Fermi National Accelerator Laboratory, Batavia, Illinois 60510, USA.

^{¶¶}Department of Physics, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA.

^{¶¶¶}Lawrence Berkeley National Laboratory, Berkeley, California 94720, USA.

^{***}Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA.

I. INTRODUCTION

During the past 40 years, solar neutrino flux measurements have been obtained through the use of a wide range of detection techniques and neutrino interactions [1–8]. This research tested the validity of the solar fusion model and provided compelling evidence for matter-induced neutrino flavor conversion within the solar interior, called the MSW effect [9,10]. The observation of antineutrino flavor oscillations by the KamLAND experiment [11], having mixing parameters consistent with those needed to explain the solar data, showed neutrino mixing to be responsible for what had been known as the *Solar Neutrino Problem*.

The most recent solar neutrino experiments are liquid scintillator based [12,13]. The low-energy threshold possible with such detectors allows for real-time detection of the low-energy ${}^7\text{Be}$ solar neutrinos. Neutrinos from ${}^7\text{Be}$ electron capture (EC) inside the Sun are monoenergetic (862 keV) and dominate the solar neutrino spectrum above the energies of pp neutrinos (<420 keV). The flux of ${}^7\text{Be}$ solar neutrinos has been previously measured by the Borexino experiment [12]. In this article, we report a measurement of the ${}^7\text{Be}$ solar neutrino flux with KamLAND, thereby providing the first independent cross check of this important quantity.

As in Borexino, solar neutrinos are detected via neutrino-electron elastic scattering, $\nu + e \rightarrow \nu + e$ (ES), which has a well understood cross section. In the standard three-flavor mixing scheme, electron neutrinos (ν_e) produced inside the Sun can transform into muon or tau neutrinos (ν_μ or ν_τ) during flight at a rate determined by the neutrino oscillation parameters and the electron density of the solar interior; the total active neutrino flux ($\nu_e + \nu_\mu + \nu_\tau$) is conserved. At the energy of ${}^7\text{Be}$ solar neutrinos, the ES cross section of ν_e is about five times larger than that of ν_μ or ν_τ .

The expected ${}^7\text{Be}$ neutrino flux at the Earth's surface, given by the GS98 solar model [14], is $5.00 \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$ [15]. The necessity of a large detector is evident considering the small ES cross section, which gives, for the KamLAND scintillator, an expected interaction rate of 500 (kt d)^{-1} including neutrino flavor conversion. The ES-induced recoil electrons have to be measured without the benefit of a convenient event tag. Here lies the main difficulty of these experiments: The low-energy backgrounds of a kiloton-size detector have to be sufficiently suppressed to allow the observation of a signal composed of only few events, a nontrivial task. In the current work this is achieved by comparing detector background models with and without a solar recoil signal to the data.

†††Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA.

††††Brookhaven National Laboratory, Yaphank, New York 11980, USA.

§§§Department of Chemistry and Physics, Arcadia University, Glenside, Pennsylvania 19038, USA.

§§§§Department of Physics, University of Washington, Seattle, Washington 98195, USA.

¶¶¶¶Department of Physics and Astronomy, Louisiana State University, Baton Rouge, Louisiana 70803, USA.




FIG. 1. Schematic diagram of the KamLAND detector.

The presence of such a signal is then inferred from the data by means of a χ^2 statistical analysis.

II. DETECTOR AND CALIBRATION

The KamLAND detector (Fig. 1) consists of 1 kton of liquid scintillator (LS) contained in a thin plastic-film balloon of 13 m diameter. The scintillation light is viewed by an array of 1879 photomultiplier tubes (PMTs) mounted on the inner surface of an 18-m-diameter stainless-steel sphere (SSS). The space between the SSS and the balloon is filled with purified mineral oil, which shields the LS from external radiation. The SSS and its content, denoted the inner detector (ID), is contained within a cylindrical, 3.2-kton water-Cherenkov outer detector (OD). All detector materials and components were selected to have low radioactivity content to maintain the option of a low background phase.

The KamLAND detector started collecting data for the reactor antineutrino phase in March 2002. Owing to the delayed coincidence structure (a prompt positron followed by a delayed neutron capture) with which antineutrinos can be tagged, the background was low enough for their detection. However, the analysis of low-energy singles data (composed of events not benefitting from a delayed coincidence) showed that the LS contained $883 \pm 20 \mu\text{Bq/kg}$ of ${}^{85}\text{Kr}$ and $58.4 \pm 1.1 \mu\text{Bq/kg}$ of ${}^{210}\text{Pb}$, the latter inferred from the decay rates of its unstable daughters ${}^{210}\text{Bi}$ and ${}^{210}\text{Po}$. The resulting total decay rate of $8.1 \times 10^7 \text{ (kt d)}^{-1}$ made the detection of about 500 (kt d)^{-1} ${}^7\text{Be}$ solar neutrino-induced recoil electrons impossible.

To enable a low-energy physics program with KamLAND, the collaboration developed methods for the efficient removal of Kr and Pb from the LS [16]. Based on extensive small-scale laboratory studies, large scintillator distillation and nitrogen purge systems were constructed underground. Two purification campaigns were performed in 2007 and 2008–2009. During the purification campaigns the old LS was drained from the detector and simultaneously filled with recycled, purified LS such that the LS mass supported by the balloon remained constant. To maximize the efficiency of purification, the temperature and density of the purified LS was carefully controlled

to maintain a boundary between the old and the purified LS. By the end of both campaigns more than five detector volume exchanges were performed, resulting in a substantial reduction of the background-creating impurities. The overall reduction factors for rates of ^{85}Kr , ^{210}Bi , and ^{210}Po were about 6×10^{-6} , 8×10^{-4} , and 5×10^{-2} , respectively. This dramatic reduction allowed the primary trigger threshold to be lowered from 180 PMT hits to 70 PMT hits (the latter value corresponds to a threshold of ~ 0.4 MeV) and thus extended KamLAND's scientific reach into the detection of low-energy solar neutrinos. To allow more detailed study of low-energy backgrounds, the threshold is lowered once per second to ~ 0.2 MeV for a duration of 1 ms. The data presented were collected in 616 days between April 7, 2009, and June 21, 2011.

The event position and energy are reconstructed based on the time and charge of photon hits recorded by the PMTs. The KamLAND coordinate system utilizes the horizontal equatorial plane as its xy plane; the z axis points up. The reconstruction is calibrated using γ sources deployed periodically in the detector, namely ^7Be (0.478 MeV), ^{60}Co (2.506 MeV), ^{68}Ge (1.022 MeV), ^{85}Sr (0.514 MeV), ^{137}Cs (0.662 MeV), and ^{203}Hg (0.279 MeV). The effects of scintillation quenching, Cherenkov light production, and PMT dark hits on the energy scale nonlinearity are determined from this calibration data. They are corrected for in the spectral analysis discussed later. The observed vertex resolution is $\sim 13 \text{ cm}/\sqrt{E(\text{MeV})}$, and the energy resolution (σ_E/E) is $(6.9 \pm 0.1)\%/\sqrt{E(\text{MeV})}$. The deviation of the position-dependent energy from the center of the detector is evaluated as $+0.3\%$ to -1.5% inside of the $-4.5 \text{ m} < z < 4.5 \text{ m}$ region shown in Fig. 2. The




FIG. 2. (Color online) Deviation of the reconstructed energy with respect to $z = 0 \text{ m}$ and deviation of the reconstructed position with respect to the true position for γ calibration sources inside the detector.

majority of the calibrations were performed by moving the sources to specific points along the central vertical detector axis using the deployment system described in Ref. [17]. To constrain deviations from rotational symmetry after scintillator purification, a full three-dimensional (3D) calibration was performed using an off-axis system described in Ref. [18].

III. EVENT SELECTION

Candidate events are selected according to the following requirements.

- (i) The radial position of event vertices must be less than 4.5 m. An additional volume selection, defined in cylindrical coordinates and motivated by the data classification described in the next section, is also applied. The combined volume selection defines the fiducial volume (FV), which corresponds to 268.6 tons of LS.
- (ii) Cosmic-ray muons [identified by a total PMT charge of larger than 10 000 photoelectrons (p.e.) or more than 5 PMT hits in the OD] and all events within 2 ms after muons are rejected to reduce background events due to muon spallation products and electronics noise. In addition, noise events within $100 \mu\text{s}$ after high-energy events (a total PMT charge of larger than 1000 p.e.) are rejected.
- (iii) Two successive events within $1 \mu\text{s}$ are rejected to avoid the possible cross-talk effect between two events owing to the finite time spread of scintillation photons.
- (iv) Coincidence events occurring within 1.2 ms of each other are eliminated to remove ^{214}Bi - ^{214}Po and ^{212}Bi - ^{212}Po sequential decays.
- (v) Candidates must pass a vertex-time-charge (VTQ) fit quality test to eliminate noise events mainly produced by two-event pileup in a one-event time window ($\sim 200 \text{ ns}$). The VTQ cut is tuned using calibration data. The reduction of the neutrino event selection efficiency is found to be negligible in the analysis energy range.

IV. DATA CLASSIFICATION

After the introduction of purified LS into KamLAND, there were time periods of thermal instability owing to slight variations in the temperature gradient of the detector. It was further found that the containment balloon acts as a ^{210}Pb reservoir, slowly releasing ^{210}Bi into the scintillator. The result of these thermal gradients was convection in the LS FV and a nonuniformly distributed ^{210}Bi concentration; thus some regions of the FV were much cleaner than others. Choosing to only analyze regions of the FV that contain low concentrations of ^{210}Bi could introduce a selection bias. Thus, a procedure for analyzing all the data regardless of the local ^{210}Bi concentration was developed. The procedure used in this analysis is as follows.

- (1) From the perspective of a ρ^2 vs z distribution in cylindrical coordinates, where $\rho^2 = x^2 + y^2$, the FV

TABLE I. Listing of the ranks defined according to the estimated ^{210}Bi rate in ($0.5 \text{ MeV} < E < 0.8 \text{ MeV}$) and the total exposure for each rank.

Rank	^{210}Bi rate $10^{-6} (\text{m}^3 \text{ s})^{-1}$	Exposure (kt d)
1	<5	26.52
2	5–10	34.42
3	10–15	27.06
4	15–20	17.81
5	20–25	11.35
6	25–30	7.63
7	>30	40.63
Total		165.43

is divided into equal-volume partitions having dimensions $d\rho^2 = 2.0 \text{ m}^2$ and $dz = 0.2 \text{ m}$.

- (2) For events with an energy of $0.5\text{--}0.8 \text{ MeV}$ (mainly ^{210}Bi) an effective event rate is calculated for each partition. Each partition is assigned a rank based on the average rate of its spatial and temporal neighbors. An average rate is calculated for up to eight partitions bordering the cell to be ranked (a partition near the FV boundary has fewer neighbors). The time dimension is included in this by adding the rate of the ranked cell, as determined in the previous and following data-taking runs (usually 1 day long), to the average rate of its neighbors.
- (3) The effective event rate is then used to classify each partition into one of seven different *ranks*, which are defined in Table I. As an example, the rank classification for one data-taking run ($\sim 24 \text{ h}$ long) is shown in Fig. 3. We find that the rank assignment of each volume generally varies slowly with time except




FIG. 3. (Color online) Rank classification of partitions in the FV during a short data-taking interval.

after periods of thermal instability when convection occurred in the LS.

The procedure just outlined allows identification of low- ^{210}Bi regions within the detector without complex and arbitrary FV cuts. However, each partition will have some volume bias owing to vertex resolution and position-dependent vertex reconstruction. This bias is corrected in the spectral fit using the fact that the following event rates, produced from cosmogenic or astrophysical sources, must be distributed uniformly in the LS: ^{11}C , ^{10}C , ^7Be , and ^7Be solar neutrinos. Using the prescribed rank classification, a simultaneous spectral fit is performed over the data from every rank to obtain one common ^7Be solar neutrino rate.

V. BACKGROUND ESTIMATION

Accurate modeling of the energy distributions of background sources inside and outside the detector is necessary to determine the ^7Be solar neutrino flux with KamLAND. These energy distributions are based on a phenomenological detector response function. They derive their validity from the fact that they describe the data well (a) before purification when radioimpurity concentrations were high and (b) in areas of high rate after purification. For some subdominant components, that cannot be verified in this way, Monte Carlo-generated spectra are used instead. The contributions of the various background components are summed with freely varying normalizations, although some normalizations are constrained by other independent KamLAND data. A fit to the candidate event spectrum then determines the partial contributions.

This section describes how the background model is constructed and what is known about its constituents. Background sources are classified into three categories: radioactive impurities in the LS, spallation products, and radioactivity in the surrounding materials.

A. Radioactive impurities in the liquid scintillator

The abundance of ^{232}Th and ^{238}U and their daughters in the LS is constrained by KamLAND data. The decay rates of the $^{222}\text{Rn}\text{--}^{210}\text{Pb}$ subchain of the ^{238}U series and the $^{228}\text{Th}\text{--}^{208}\text{Pb}$ subchain of the ^{232}Th series can be measured, almost background free, using $^{214}\text{Bi}\text{--}^{214}\text{Po}$ and $^{212}\text{Bi}\text{--}^{212}\text{Po}$ $\beta\text{--}\alpha$ delayed coincidences, with 94.8% and 31.9% selection efficiencies, respectively. The inferred ^{238}U and ^{232}Th concentrations assuming secular equilibrium are $(5.0 \pm 0.2) \times 10^{-18} \text{ g/g}$ ($93 \pm 4 \text{ nBq/m}^3$) and $(1.3 \pm 0.1) \times 10^{-17} \text{ g/g}$ ($59 \pm 4 \text{ nBq/m}^3$), respectively. Some radioactive decay chains are found to be out of secular equilibrium. A fit to the prominent ^{210}Po α peak, quenched to about 0.3 MeV visible energy in Fig. 4, yields spatially varying activities ranging from 2.4 to 4.8 mBq/m^3 , depending on the rank of the analyzed volume. However, as discussed in the following section, this background component is not considered in the fit because it falls below the analysis threshold. The higher-mass members of the decay chains above ^{228}Th and ^{222}Rn are determined by a spectral fit and are found to be unimportant.

Below 1 MeV , the background is dominated by daughters of ^{210}Pb , namely ^{210}Bi and ^{210}Po . ^{210}Bi , the most important




FIG. 4. (Color online) Energy spectra of candidate events for each data-set rank, exhibiting different ^{210}Bi rates. All data sets are used for the simultaneous fit of the ^7Be solar neutrino spectrum, with a common rate normalization. The location of the major background sources are labeled and the theoretical electron recoil spectrum of 862-keV ^7Be solar neutrinos, whose rate is based on the standard solar model by Serenelli *et al.* [15], is shown for comparison.

background contributor, is found to be spatially nonuniform; in addition, its decay rate in the FV fluctuates in time. This is attributed to ^{210}Bi supplied from the balloon surface by irregular LS convection currents, followed by its decay with $T_{1/2} = 5.01$ days. This interpretation is supported by the observation that around the balloon surface, the spectrum is dominated by electrons from ^{210}Bi decays, and α particles from ^{210}Po decays, consistent with the hypothesis that the balloon film is contaminated with ~ 200 Bq of ^{210}Pb , a ^{222}Rn daughter introduced during detector construction. The scintillator subvolume ranking technique, described before, was devised to cope with this variability in an unbiased way. Depending on the rank, the specific ^{210}Bi activity varies between 35.7 and 681.1 $\mu\text{Bq}/\text{m}^3$ within the FV. ^{210}Bi undergoes a first forbidden β decay. The parametrization of the resulting beta spectrum will be discussed in the analysis section.

The concentration of ^{40}K is measured to be $(7.3 \pm 1.2) \times 10^{-17}$ g/g from the energy spectrum fit. Owing to the fluid circulation during LS purification, ^{238}U , ^{232}Th , and ^{40}K may have nonuniform spatial distributions. The long-lived radioactive noble gas, ^{85}Kr , was the major source of low-energy background prior to scintillator purification. The present background model includes this component to deal with any leftover activity. ^{39}Ar is included as a potential source in the background model, its abundance constrained by the ratio of ^{85}Kr and ^{39}Ar found in the atmosphere. From the spectral fit we find that its contribution to the background is negligible. This is consistent with the prepurification spectral analysis results. For volume ranks 1 through 6 the ^{85}Kr specific activity varies between 4.1 and 19.2 $\mu\text{Bq}/\text{m}^3$. For

the highest background partitions—those with rank 7—the fit only yields an upper limit for the activity. After LS purification, we collected samples from a location within the FV at $z = +1.5$ m. These samples were analyzed for their Kr content using a helium purge, a cold trap to retain the Kr, and a residual gas analyzer to measure its partial pressure. Based on these measurements, the ^{85}Kr decay rate was estimated at 8.3 ± 4.2 $\mu\text{Bq}/\text{m}^3$, assuming a recent isotopic ratio of ^{85}Kr in air [19]. As a third cross check for the ^{85}Kr content of the LS, a ^{85}Kr - $^{85\text{m}}\text{Rb}$ delayed coincidence analysis was performed, utilizing the β decay into the metastable, 514-keV excited state of ^{85}Rb . This analysis yielded 17.3 ± 5.9 $\mu\text{Bq}/\text{m}^3$, averaged over the entire FV. ^{85}Kr undergoes a unique first forbidden β decay. The β spectrum contained in the background model was calculated using a relativistic Fermi function plus a shape correction accounting for the forbiddenness following the procedure outlined in Ref. [20]. This spectral parametrization was found to fit well the high-statistics β spectrum collected before LS purification.

B. Spallation products

The 2700 mwe of rock overburden of the Kamioka Underground Laboratory suppresses the rate of cosmic ray muons traversing the KamLAND LS to 0.198 ± 0.014 s^{-1} [21]. The surviving muons can produce unstable light nuclei by spallation of carbon, whose decays result in background. The dominant cosmogenic background between 1 and 2 MeV is due to decays of ^{11}C (β^+ , $\tau = 29.4$ min, $Q = 1.98$ MeV). Owing to its relatively long half-life, these decays cannot be tagged without incurring large dead time. Its decay rate has earlier been estimated to be 1106 ± 178 $(\text{kt d})^{-1}$ [21]. The background contribution of ^{11}C is constrained by this value.

Another spallation source of interest is ^7Be [EC decay, $\tau = 76.8$ days, $Q = 0.862$ MeV]. While its production rate by muon spallation in the LS is estimated to be small [21,22], there is the possibility of a higher-than-steady-state production yield owing to the introduction of fresh, surface-exposed LS during the scintillator purification. Therefore, to be conservative, the rate of ^7Be decays is unconstrained.

C. Radioactivity in the surrounding material

The background from external γ rays is mainly caused by ^{40}K , ^{232}Th , and ^{238}U contained in the surrounding rock, stainless steel, PMT glass, balloon film, and Kevlar suspension ropes. The energy distributions resulting from these radiation sources were modeled by means of a Monte Carlo simulation. The simulation was tuned with source calibration data to reproduce the vertex distribution as well. The γ -ray attenuation in the radial direction in the simulation is consistent with that in the real data. External backgrounds dominate the energy distribution for radial positions larger than 4.5 m. These data were used to fit the relative background contributions. The Monte Carlo simulation was then used to extrapolate the background for $R < 4.5$ m. Based on this fit, it was concluded that external γ rays do not significantly impact the FV background below 1 MeV.

As discussed before, the balloon surface is a source of electrons from ^{210}Bi decays and α particles from ^{210}Po decays. While the FV cut effectively suppresses these backgrounds, accidental pileup of two external events can lead to a vertex and energy displacement, moving external events into the analysis volume. Owing to the high ^{210}Po decay rate, resulting in a pronounced peak at ~ 0.3 MeV in the energy spectrum (see Fig. 4), most of the pileup events are concentrated in the high-energy tail of this peak. Although the absolute rate of tail events is not small, the fraction of pileup events to observed events above 0.5 MeV is $< 1\%$. To eliminate this background and the systematic bias it would introduce into the measurement of the ^7Be solar neutrinos, the energy threshold was set to 0.5 MeV in the present analysis.

VI. SYSTEMATIC UNCERTAINTY

The leading contributions to the systematic uncertainty of the ^7Be solar neutrino flux measurement are listed in Table II. The measured neutrino-electron scattering rate is converted into a solar neutrino flux, the accuracy of this conversion is given by the uncertainty of the interaction cross section. Based on the evaluation of Ref. [23], a value of 1% is assigned to this error. The determination of the flux further requires knowledge of the number of electrons contained in the FV. The LS density is measured to be 0.780 g/cm^3 with an uncertainty of 0.025% at 11.5°C . The uncertainty of the dependence of the number of electrons on the temperature within the FV is estimated at 0.1%. We estimate that 9.21×10^{31} electrons are contained in the 344.3 m^3 FV. Data collected with the full volume calibration system showed vertex reconstruction deviations of less than 5 cm. This corresponds to a FV uncertainty of 3.4%. The FV event selection inefficiency owing to vertex misreconstruction was established with source calibrations and is less than 0.5% in the analysis energy region.

Other systematic uncertainties, related to the modeling of the detector response, are determined through the spectral fit procedure, which is presented in the following section. The correction for the nonlinearity of the energy scale for each

TABLE II. Uncertainties on the measurement of the ^7Be solar neutrino flux.

Source	Uncertainty (%)
Cross section	1.0
No. of target electrons	0.10
Fiducial volume	3.4
Vertex misreconstruction	0.5
Energy scale	7.9
Rank-dependent energy scale	2.9
Energy resolution	3.4
Background from ^{238}U series (^{222}Rn - ^{210}Pb)	1.7
Background from ^{232}Th series (^{228}Th - ^{208}Pb)	1.8
Background from other solar neutrinos	1.9
Systematic total	10.2
Statistics	12.4
Total	16.1

particle type (γ , e^- , and e^+) is performed by varying the energy response parameters in the χ^2 fit. The fit model, on which the solar neutrino analysis is based, uses free-floating energy scales with constraints from calibration data in the form of penalty terms. As such, the fit uncertainty already includes the energy scale uncertainty. To quantify the contribution of the energy scale uncertainty as a separate item in the error budget, we repeat the fit with the energy scale parameters fixed. The energy scale error is stated as the quadratic difference between the errors obtained from free-floating and fixed parameter fits. This method implies a 7.9% uncertainty on the best-fit ^7Be rate owing to the uncertainty of the energy scale. To check the possibility of the energy scale varying with the rank, we compared the fit result for two extreme cases, namely fully correlated and fully independent energy scale between ranks. We adopt the deviation of the best-fit ^7Be as an estimate of the associated systematic uncertainty. To propagate the $0.1\%/\sqrt{E(\text{MeV})}$ uncertainty of the energy resolution, we repeat the analysis varying the detector resolution within this uncertainty. We find that this causes the best-fit ^7Be rate to vary by 3.4%.

The definition of rank boundaries (Table I) and their effect on the best-fit ^7Be rate was also studied. It is important to note that the fit sensitivity to the solar signal comes from ranks 1 and 2, where the signal-to-background is largest; hence, changing the boundary definitions of the other ranks has negligible effect. The fitting procedure was repeated while varying the boundary between ranks 1 and 2 from 3×10^{-6} to 7×10^{-6} events/($\text{m}^3\text{ s}$). The choice of 5×10^{-6} events/($\text{m}^3\text{ s}$) as the boundary was found to have the highest fit probability, quantified by $\chi^2/\text{n.d.f.}$ (where n.d.f. stands for number of degrees of freedom), and is used in the final analysis presented here. Furthermore, we find the best-fit rate for all rank boundaries considered were consistent with each other to within 5% and conclude that the choice of boundary does not bias the result.

The largest background contribution, ^{210}Bi first forbidden β decay, has an additional uncertainty related to its shape correction. Using our data, mainly rank 7 where the background rate is highest, we derived a phenomenological fourth-order polynomial correction to the shape factor published in Ref. [24] to better model the ^{210}Bi shape. The parameters of this correction polynomial (a_n) are $a_0 = 1$, $a_1 = 41.4 \pm 0.8$, $a_2 = -101.2 \pm 1.4$, $a_3 = 102.9 \pm 3.5$, and $a_4 = -37.9 \pm 2.0$ in units of MeV^{-n} . Within the analysis energy window, deviations of the spectral shape from that of Ref. [24] were small.

Position-dependent biases in vertex reconstruction could introduce an artificial nonuniformity in the distribution of ^7Be solar neutrinos within the FV, resulting in a systematic error on the simultaneous fit to different rank data. The effects of the nonuniformity are parameterized and constrained for each rank spectrum to obtain uniformly distributed ^{11}C events.

The systematic uncertainties owing to the ^{238}U and ^{232}Th series and other solar neutrinos are evaluated by the deviation of the ^7Be solar neutrino rate on varying their rates within $\pm 1\sigma$.

TABLE III. Summary of signal and background in the FV. The best-fit signal and background rates in the whole energy range (w/o E cut) and the ^7Be solar neutrino energy range ($0.5 \text{ MeV} < E < 0.8 \text{ MeV}$) are shown.

Isotope	Event rate w/o E cut		Event rate in 0.5–0.8 MeV	
	All ranks	Rank 1	All ranks	Rank 1
	(kt d) $^{-1}$		(kt d) $^{-1}$	
Solar neutrinos				
$^7\text{Be } \nu$	582 ± 94		117 ± 19	
Other ν	1443		14.9	
Radioactive impurities in the LS				
^{210}Bi	23974 ± 883	3955 ± 238	4557 ± 168	578 ± 35
^{85}Kr	858 ± 59	453 ± 102	57 ± 4	30 ± 7
^{39}Ar	3 ± 3	2 ± 3	0.04 ± 0.04	0.02 ± 0.04
^{40}K	181 ± 29	3 ± 38	48 ± 8	1 ± 10
Spallation products				
^7Be	167 ± 173		2 ± 2	
^{11}C	973 ± 10		3.55 ± 0.04	
Other	173		8.0	
Others				
External γ	—	—	2.4 ± 0.4	2 ± 1
Pileup	—	—	10 ± 1	8 ± 2

VII. RESULT

The ^7Be solar neutrino rate is estimated by a likelihood fit to the binned energy spectra of candidate events with visible energy between 0.5 and 1.4 MeV. The background contributions from ^{210}Bi , ^{85}Kr , ^{40}K , and ^7Be EC are free parameters and their normalizations are left unconstrained in the fit. The contributions from the ^{222}Rn - ^{210}Pb and ^{228}Th - ^{208}Pb chains and ^{11}C are allowed to vary, but their normalizations are constrained by independent KamLAND measurements, as outlined in Sec. V. Distributions with different background rank are fitted simultaneously. The background rates, derived from the fit normalizations, are summarized in Table III. The backgrounds from external γ rays, and pileup events are constrained by the MC study. The model parameters of the detector response are also constrained, as discussed in Sec. VI.

Figure 5 shows the result of this procedure: the best-fit spectrum for the rank 1 data set which is lowest in background and, therefore, most sensitive to the solar neutrino signal. The inset shows the background-subtracted energy distribution, which exhibits the shape of the best-fit ^7Be neutrino signal extracted from the analysis. The backscattering edge is visible and located at the correct energy. The $\chi^2/\text{n.d.f.}$ comparing the binned data and the best-fit model is found to be 635.3/589 using data of all ranks. The $\chi^2/\text{n.d.f.}$ of the lowest background rank 1 data is 80.1/90. As can be seen from the data in Table III the signal-to-background ratio is estimated to be 1:5.5 for rank 1 data ($0.5 < E < 0.8 \text{ MeV}$). This relatively low signal-to-background ratio naturally invites the question of whether a statistically significant ^7Be solar neutrino signal is really present in the data. The fit to the data was also performed assuming the absence of a solar neutrino signal, but minimizing all other background components within their constraints as before. The observed increase in the fit χ^2 leads




FIG. 5. (Color online) Energy spectrum of selected ^7Be solar neutrino candidates for the rank 1 data set together with the best-fit curves from the simultaneous fit to all rank's spectra. The fit range is $0.5 \text{ MeV} < E < 1.4 \text{ MeV}$. The background-subtracted spectrum is shown in the inset, where a Compton-like shoulder characteristic for the ^7Be solar neutrino contribution is evident. The error bars are statistical only and do not include correlated systematic uncertainties in the background rate estimates.

us to reject the no-solar signal hypothesis at 8.2σ CL. To understand whether the solar recoil signal, preferred by the data and shown in the inset of Fig. 5, is a unique solution or whether other continuous distributions yield equally good fits, the neutrino energy (assuming a monoenergetic source) and interaction rates were floated and the fit χ^2 profile determined under their variation. The result of this procedure is shown in Fig. 6. The KamLAND data clearly prefer the presence of an edge at an energy that coincides with that expected for solar ^7Be neutrino-induced electron recoils. Assuming that the edge is indeed solar neutrino induced, we determine the solar neutrino energy to be $E_\nu = 862 \pm 16 \text{ keV}$. This is the first direct spectroscopic determination of the solar ^7Be neutrino energy.

The fit to the KamLAND data gives a solar neutrino interaction rate of $R_{\text{KL}} = 582 \pm 94 \text{ (kt d)}^{-1}$. The quoted error corresponds to the quadratic sum of the statistical and




FIG. 6. (Color online) $\Delta\chi^2$ profile from the fit to the energy of monoenergetic neutrinos discussed in the text. The vertical line indicates the neutrino energy predicted from the ^7Be electron-capture reaction at 862 keV.

systematic errors, listed in Table II. This result is in agreement with the latest interaction rate reported by the Borexino experiment: $R_B = 460^{+21}_{-22}$ (kt d) $^{-1}$ [12]. The rate difference ($R_{KL} - R_B$) deviates by 1.3σ from zero. Differences in the chemical composition of the LSs used in both experiments result in a 3.6% difference of the rates. Assuming that the ES interactions detected by KamLAND are due to a pure electron flavor flux, KamLAND's interaction rate corresponds to a 862-keV ${}^7\text{Be}$ solar neutrino flux of $(3.26 \pm 0.52) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$. The standard solar model (SSM) by Serenelli *et al.* [15] gives two ${}^7\text{Be}$ solar neutrino flux values, depending whether the older Grevesse and Sauval (GS98) [14] or the Asplund (AGSS09) [25] solar abundances are utilized. When the GS98 solar abundances are assumed, the SSM flux value is $(5.00 \pm 0.35) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$. The flux reduces to $(4.56 \pm 0.32) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$ under the assumption of the AGSS09 solar abundances. Assuming that the ν_e mix with ν_μ and ν_τ the KamLAND flux measurement corresponds to a survival probability, $P_{ee} = 0.66 \pm 0.15$. This value was obtained taking into account the cross section difference for these neutrino flavors.

Using KamLAND's 2013 best-fit neutrino oscillation parameters [11], based on a global oscillation analysis under the assumption of CPT invariance, the survival probability is better constrained. In this case, we obtain a total ${}^7\text{Be}$ neutrino flux of $(5.82 \pm 1.02) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$. This result is consistent with the flux determination provided by Borexino: $(4.75^{+0.26}_{-0.22}) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$ [12], and while it somewhat favors the GS98 model flux, the KamLAND data are not precise enough to discriminate between the two solar model fluxes.

VIII. CONCLUSION

The KamLAND collaboration reports a new measurement of the ${}^7\text{Be}$ solar neutrino interaction rate in a liquid scintillator. Performing this difficult measurement required an extensive purification campaign, reducing the scintillator's radioimpurity content by several orders of magnitude. The measured ${}^7\text{Be}$ solar neutrino rate in KamLAND is $R_{KL} = 582 \pm 94$ (kt d) $^{-1}$, which corresponds to a flux of $(5.82 \pm 1.02) \times 10^9 \text{ cm}^{-2} \text{ s}^{-1}$ under the assumption of KamLAND's 2013 best-fit oscillation parameters.

The statistical significance of this signal is estimated to be 8.2σ and provides the first independent verification of the only prior measurement of this quantity, performed by Borexino. The solar neutrino flux derived from the KamLAND data agrees with the solar model values, but is not accurate enough to shed light on the question of solar metallicity.

ACKNOWLEDGMENTS

The KamLAND experiment is supported by the Grant-in-Aid for Specially Promoted Research under Grants No. 16002002 and No. 21000001 of the Japanese Ministry of Education, Culture, Sports, Science and Technology; the World Premier International Research Center Initiative (WPI Initiative), MEXT, Japan; Stichting FOM in the Netherlands; and under the US Department of Energy (DOE) Grants No. DE-AC02-05CH11231 and No. DE-FG02-01ER41166, as well as other DOE grants to individual institutions. The Kamioka Mining and Smelting Company has provided service for activities in the mine.

-
- [1] B. T. Cleveland *et al.*, *Astrophys. J.* **496**, 505 (1998).
 [2] W. Hampel *et al.*, *Phys. Lett. B* **447**, 127 (1999).
 [3] M. Altmann *et al.*, *Phys. Lett. B* **616**, 174 (2005).
 [4] J. N. Abdurashitov *et al.*, *Phys. Rev. C* **80**, 015807 (2009).
 [5] J. Hosaka *et al.*, *Phys. Rev. D* **73**, 112001 (2006).
 [6] B. Aharmim *et al.*, *Phys. Rev. C* **81**, 055504 (2010).
 [7] B. Aharmim *et al.*, *Phys. Rev. Lett.* **101**, 111301 (2008).
 [8] C. Arpesella *et al.*, *Phys. Rev. Lett.* **101**, 091302 (2008).
 [9] L. Wolfenstein, *Phys. Rev. D* **20**, 2634 (1979).
 [10] S. P. Mikheyev and A. Yu. Smirnov, *Sov. J. Nucl. Phys.* **42**, 913 (1985).
 [11] A. Gando *et al.*, *Phys. Rev. D* **88**, 033001 (2013).
 [12] G. Bellini *et al.*, *Phys. Rev. D* **89**, 112007 (2014).
 [13] S. Abe *et al.*, *Phys. Rev. C* **84**, 035804 (2011).
 [14] N. Grevesse and A. J. Sauval, *Space Sci. Rev.* **85**, 161 (1998).
 [15] A. M. Serenelli, W. C. Haxton, and C. Peña-Garay, *Astrophys. J.* **743**, 24 (2011).
 [16] G. Keefer *et al.*, *Nucl. Instrum. Methods A* **769**, 79 (2015).
 [17] T. I. Banks *et al.*, *Nucl. Instrum. Methods A* **769**, 88 (2015).
 [18] B. E. Berger *et al.*, *JINST* **4**, P04017 (2009).
 [19] M. Hirota *et al.*, *J. Radiat. Res.* **45**, 405 (2004).
 [20] E. J. Konopinski, *The Theory of Beta Radioactivity* (Oxford University Press, New York, 1966).
 [21] S. Abe *et al.*, *Phys. Rev. C* **81**, 025807 (2010).
 [22] T. Hagner *et al.*, *Astropart. Phys.* **14**, 33 (2000).
 [23] J. N. Bahcall, M. Kamionkowski, and A. Sirlin, *Phys. Rev. D* **51**, 6146 (1995).
 [24] H. Daniel, *Nucl. Phys.* **31**, 293 (1962).
 [25] M. Asplund, N. Grevesse, A. J. Sauval, and P. Scott, *Annu. Rev. Astron. Astrophys.* **47**, 481 (2009).