

Electromagnetic transition strengths for light nuclei in the Skyrme model

M. Haberichter,^{1,*} P. H. C. Lau,^{2,3,†} and N. S. Manton^{3,‡}

¹*School of Mathematics, Statistics and Actuarial Science, University of Kent, Canterbury CT2 7NF, United Kingdom*

²*Yukawa Institute for Theoretical Physics, Kyoto University, Kitashirakawa Oiwakecho, Sakyo-ku, Kyoto City, Kyoto 606-8502, Japan*

³*Department of Applied Mathematics and Theoretical Physics, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, United Kingdom*

(Received 9 November 2015; published 7 March 2016)

We calculate reduced $B(E2)$ electromagnetic transition strengths for light nuclei of mass numbers $B = 8, 12, 16, 20, 24$, and 32 within the Skyrme model. We find that the predicted transition strengths are of the correct order of magnitude and the computed intrinsic quadrupole moments match the experimentally observed effective nuclear shapes. For the Hoyle state we predict a large $B(E2) \uparrow$ value of $0.0521 e^2 b^2$. For oxygen-16, we can obtain a quantitative understanding of the ground state rotational band and the rotational excitations of the second spin-0 state, 0_2^+ .

DOI: 10.1103/PhysRevC.93.034304

I. INTRODUCTION

Radiative electromagnetic transitions between nuclear states are an excellent way to probe nuclear structure and to test nuclear structure models [1–3]. In even-even nuclei, the reduced transition probability $B(E2 : 0_1^+ \rightarrow 2_1^+)$ from the 0_1^+ ground state to the first excited 2_1^+ state is particularly important [4,5]. $B(E2)$ transitions play a crucial role [5,6] in determining mean lifetimes of nuclear states, the nuclear potential deformation parameter β , the magnitude of intrinsic electric quadrupole moments, and the energy of low-lying levels of nuclei. Large quadrupole moments and transition strengths indicate collective effects in which many nucleons participate.

The Skyrme model [7,8] and its topological soliton solutions (known as Skyrmions) have been found to capture important features of light nuclei of even baryon number. As in the α -particle model of nuclei [9,10], Skyrmions with topological charge B a multiple of four are composed of charge four subunits [11]. Here the role of the α particle is taken by the cubic $B = 4$ Skyrmion. The arrangements of $B = 4$ cubes often resemble [11,12] those discussed in the α -particle model. In addition, the allowed quantum states for each Skyrmion of topological charge B often match [13–17] the ground and excited states of nuclei with mass number B a multiple of four. Among other successes of the Skyrme model is the prediction of the excitation energy of states in the rotational bands of carbon-12, including the excitations of the Hoyle state [18]. However, isoscalar quadrupole $E2$ transitions within the Skyrme model, which can provide us with valuable information about the internal structure of nuclei, and a nontrivial test of the model, have not yet been studied in detail. Note that isovector magnetic dipole $M1$ transitions have been discussed for Skyrmion states in the literature. The $M1$ transition from a delta to a nucleon has been considered [19], and also the transition from a deuteron to its isovector state [20]. In appendix B of Ref. [21] a connection between the isovector

magnetic moment operator and the Skyrmion's mixed inertia tensor was established for arbitrary SU(2) Skyrmions.

In the following, we briefly review the Skyrme model and its soliton solutions. For further details, we refer the interested reader to the literature [22–24]. The Skyrme model is a modified nonlinear sigma model, in which the sigma field σ and isotriplet of pion fields $\boldsymbol{\pi}$ are combined into an SU(2)-valued scalar field

$$U(\mathbf{x}, t) = \sigma(\mathbf{x}, t)1_2 + i\boldsymbol{\pi}(\mathbf{x}, t) \cdot \boldsymbol{\tau}, \quad (1)$$

where $\boldsymbol{\tau}$ denotes the triplet of Pauli matrices and the normalization constraint $\sigma^2 + \boldsymbol{\pi} \cdot \boldsymbol{\pi} = 1$ is imposed.

For a static Skyrme field $U(\mathbf{x})$, the energy in Skyrme units is

$$E = \int \left\{ -\frac{1}{2} \text{Tr}(R_i R_i) - \frac{1}{16} \text{Tr}([R_i, R_j][R_i, R_j]) + m^2 \text{Tr}(1_2 - U) \right\} d^3x. \quad (2)$$

Here, R_i are the spatial components of the SU(2)-valued current $R_\mu = (\partial_\mu U)U^\dagger$, and m is a dimensionless pion mass parameter. Skyrme units are converted to physical energies and lengths (in MeV and fm) by the factors $F_\pi/4e_{\text{sky}}$ and $2/e_{\text{sky}}F_\pi$, respectively. e_{sky} is a dimensionless constant and F_π can be interpreted as the pion decay constant. m is related to the pion tree level mass m_π via $m = 2m_\pi/e_{\text{sky}}F_\pi$. The energy and length conversion factors are fixed by comparison with experimental nuclear physics data.

Skyrmions are critical points of the potential energy (2) and are characterized by a conserved, integer-valued topological charge

$$B = -\frac{1}{24\pi^2} \int \epsilon_{ijk} \text{Tr}(R_i R_j R_k) d^3x. \quad (3)$$

B is the topological degree of the map $U : \mathbb{R}^3 \rightarrow \text{SU}(2)$ at any given time, which is well defined for fields satisfying the boundary conditions $\sigma \rightarrow 1$ and $\boldsymbol{\pi} \rightarrow \mathbf{0}$ as $|\mathbf{x}| \rightarrow \infty$. Physically, when semiclassically quantized [19,25], a Skyrmion of charge B is interpreted as a nucleus of mass number (or baryon

*m.haberichter@kent.ac.uk

†pakhang.lau@yukawa.kyoto-u.ac.jp

‡N.S.Manton@damtp.cam.ac.uk

TABLE I. Skyrmions of baryon numbers $B = 8, 12, 16, 20, 24, 32$, for $m = 1$. We list the symmetry group G of each Skyrmion, its energy relative to the Skyrme-Faddeev bound $\frac{E}{12\pi^2 B}$, and the diagonal elements of the inertia tensors U_{ij}, V_{ij}, W_{ij} (in Skyrme units). We also list the isospin-zero nuclei (in their ground states) that can be modelled by the Skyrmions. These are recognized from the symmetry group rather than the energy E .

B	G	Nucleus	$\frac{E}{12\pi^2 B}$	U_{11}	U_{22}	U_{33}	V_{11}	V_{22}	V_{33}	W_{11}	W_{22}	W_{33}
8	D_{4h} (twist)	${}^8_4\text{Be}_4$	1.279	298	292	326	4093	4094	1381	0	0	0
	D_{4h} (no twist)		1.283	287	291	350	4615	4615	1296	0	0	0
12	D_{4h}	Hoyle	1.274	440	449	456	12137	12137	2139	0	0	0
	D_{3h}	${}^{12}_6\text{C}_6$	1.278	442	442	497	5009	5006	7627	41	41	38
16	D_{2d} (bent square)		1.271	572	571	674	9123	9119	14602	0	0	0
	D_{4h} (flat square)		1.272	563	567	689	9143	9174	15682	0	0	0
20	T_d	${}^{16}_8\text{O}_8$	1.276	586	586	674	9100	9101	9128	0	0	0
	T_d		1.273	757	757	819	12820	12820	12821	0	0	0
24	D_{3h}	${}^{20}_{10}\text{Ne}_{10}$	1.276	857	735	735	18542	18591	9762	15	-15	-11
	D_{2h}		1.267	877	862	956	26980	14189	36783	0	0	0
24	D_{3d}		1.269	879	890	959	19600	19600	29863	0	0	0
	D_{3h}	${}^{24}_{12}\text{Mg}_{12}$	1.269	869	869	1006	20554	20454	16226	-99	99	99
32	O_h	${}^{32}_{16}\text{S}_{16}$	1.264	1115	1116	1367	31625	31628	31704	0	0	0

number) B . In nuclear physics, the notation for mass number is A but we will keep our notation B in this paper.

Skyrmion solutions with rescaled pion mass $m = 1$ and with baryon number B a multiple of four have been previously found [11,12,17,18]. For baryon numbers $B = 8, 12, 16, 20, 24, 32$, we recalculate the classical Skyrmion solutions using two different numerical relaxation techniques: nonlinear conjugate gradient [12,26,27] and damped full field evolution [28]. Skyrmions are the solutions of minimal energy, or sometimes local minima or saddle points with energies close to minimal. Our calculations have led us to two new solutions with $B = 24$. These are obtained by gluing together two copies of $B = 12$ solutions.

To find solutions, Skyrme fields of positive topological charge B and with a given symmetry group G are created by multilayer rational map ansätze [12], or product ansätze [11]. These initial Skyrme field configurations are relaxed on grids with $(201)^3$ grid points and a spatial grid spacing $\Delta x = 0.1$ to find precise solutions. We list in Table I the symmetry group, the energy and the diagonal elements of the isospin (U_{ij}), spin (V_{ij}), and mixed (W_{ij}) inertia tensors for Skyrmions with baryon numbers $B = 8, 12, 16, 20, 24, 32$. The Skyrmions are orientated such that all off-diagonal elements of the inertia tensors vanish. The formulas for the inertia tensors U_{ij}, V_{ij}, W_{ij} are rather complicated and have been given first in general form for arbitrary SU(2) Skyrmions in Refs. [20,29]. However, for numerical calculations it is much more convenient to express the inertia tensors in terms of the sigma field and pion field isotriplet, see formulas given in Refs. [17,30]. The baryon density isosurfaces we obtain are shown in Fig. 1. On these surfaces, the π -field values are visualized using Manton and Sutcliffe's field colouring scheme described in detail in Ref. [31].

In this paper, we calibrate the Skyrme model with properties of the carbon-12 nucleus. The root-mean-square matter radius of a nucleus can be calculated within the Skyrme model as

$$\langle r^2 \rangle^{\frac{1}{2}} = \left(\frac{\int r^2 \mathcal{E}(\mathbf{x}) d^3x}{\int \mathcal{E}(\mathbf{x}) d^3x} \right)^{\frac{1}{2}}, \quad (4)$$

where $\mathcal{E}(\mathbf{x})$ is the static energy density and $r = |\mathbf{x}|$. We list the matter radii $\langle r^2 \rangle_{\text{Sky}}^{\frac{1}{2}}$ (in Skyrme length units) of all the Skyrmions considered here in Table II. The energy and length conversion factors are tuned to match the experimental nuclear mass 11178 MeV and matter radius 2.43 fermi [32] for the carbon-12 ground state. This fixes the conversion factors to be

$$\left[\frac{F_\pi}{4e_{\text{Sky}}} \right] = 6.154 \text{ MeV}, \quad \left[\frac{2}{e_{\text{Sky}} F_\pi} \right] = 1.061 \text{ fm}, \quad (5)$$

and gives the parameter values

$$e_{\text{Sky}} = 3.889, \quad F_\pi = 95.6 \text{ MeV}, \quad \hbar = 30.2, \quad \text{and} \\ m_\pi = 185.9 \text{ MeV}, \quad (6)$$

where $\hbar = 2e_{\text{Sky}}^2$. (More precisely, the energy unit is $F_\pi/4e_{\text{Sky}} = 6.154 \text{ MeV}$ and the length unit is $2\hbar/e_{\text{Sky}}F_\pi = 1.061 \text{ fm}$, so the energy-length unit is $\hbar/2e_{\text{Sky}}^2 = 6.529 \text{ MeV fm}$. As $\hbar = 197.3 \text{ MeV fm}$ experimentally, $2e_{\text{Sky}}^2 = 30.2$ or equivalently $\hbar = 30.2$ in Skyrme units.) Note that we use in this article a value of F_π that is substantially lower than its experimental value and a value of m_π that is substantially larger than the physical pion mass. The experimental values for the pion mass and pion decay constant are given by $m_\pi = 138 \text{ MeV}$ and $F_\pi = 186 \text{ MeV}$, respectively. One can argue that m_π and F_π should be taken from experiment and that the Skyrme constant e_{Sky} should be used as the fitting parameter. Fitting Skyrmions with m_π, F_π and F_K (with the kaon mass term added) at its physical values and $e_{\text{Sky}} \simeq 4.1$ allows to reasonably describe the mass splittings [33,34] within the SU(3) multiplets of baryons. For these parameter values the absolute values of nuclear masses are not reproduced and one has to appeal to large Casimir energies from field fluctuations [35,36] to make up the difference. Even for the $B = 1$ Skyrmion, these quantum corrections to the soliton mass are difficult to calculate accurately and estimates have been given by various authors [35–40]. Except for the work on Casimir energies of




FIG. 1. Surfaces of constant baryon density (not to scale) of Skyrmons with pion mass parameter $m = 1$. The Skyrmons have baryon number and symmetry group: $B = 8$ (a) D_{4h} with 90° twist, (b) D_{4h} with no twist; $B = 12$ (c) D_{4h} , (d) D_{3h} ; $B = 16$ (e) D_{2d} (bent square), (f) D_{4h} (flat square), (g) T_d ; $B = 20$ (h) T_d , (i) D_{3h} ; $B = 24$ (j) D_{2h} , (k) D_{3d} , (l) D_{3h} ; $B = 32$ (m) O_h .

strongly bound $B = 2$ configurations reported in Ref. [40], there do not exist estimates of Casimir effects for $B > 1$. It is very difficult to pin down the exact magnitude of the Casimir contributions to Skyrmon masses because first of all this requires a full knowledge of the vibrational space of Skyrmons. New insights into the structure of vibrational spaces and their quantization have been reported recently in Ref. [41]. In this article, we adopt a different point of view which goes back to Adkins, Nappi, and Witten [19]. It has been found effective when modeling nuclei by Skyrmons to adjust the Skyrme parameters to fit nuclear masses and to

interpret m_π and F_π as renormalized quantities [42–44]. For example, calibrating the Skyrme model with properties of the carbon-12 nucleus has previously proven successful [18] in describing the spectrum of rotational excitations of carbon-12, including the excitations of the Hoyle state, and in modeling nucleon-nucleon scattering within the Skyrme model [45,46]. In the following, we will refer to Eq. (5) as the Lau-Manton (LM) calibration. We will find that the LM calibration is well suited for calculating electromagnetic transition strengths within the Skyrme model. In addition, for this calibration the predicted nuclear masses and matter radii are in reasonable

TABLE II. Nuclear masses E and root mean square matter radii $\langle r^2 \rangle^{\frac{1}{2}}$ for nuclei of baryon numbers $B = 8, 12, 16, 20, 24, 32$ and isospin zero. Here, the subscripts ‘‘Sky’’ and ‘‘LM’’ refer to Skyrme units and the Lau-Manton calibration (5), respectively. The experimental matter radii $\langle r^2 \rangle_{\text{Exp}}^{\frac{1}{2}}$ are taken from Ref. [47] and are given in fermi.

B	G	Nucleus	E_{LM} [MeV]	E_{Exp} [MeV]	$\langle r^2 \rangle_{\text{Sky}}^{\frac{1}{2}}$	$\langle r^2 \rangle_{\text{LM}}^{\frac{1}{2}}$ [fm]	$\langle r^2 \rangle_{\text{Exp}}^{\frac{1}{2}}$ [fm]
8	D_{4h} (twist)	${}^8_4\text{Be}_4$	7457.5	7451.9	2.05	2.18	2.34 ^a
	D_{4h} (no twist)		7480.9		2.15	2.28	
12	D_{4h}	Hoyle	11142.6		2.76	2.93	2.89 ^b
	D_{3h}	${}^{12}_6\text{C}_6$	11178	11178	2.29	2.43	2.43 ^c
16	D_{2d} (bent square)		14821.9		2.66	2.82	
	D_{4h} (flat square)		14833.5		2.70	2.87	
20	T_d	${}^{16}_8\text{O}_8$	14903.5	14903.9	2.35	2.50	2.70
	D_{3h}	${}^{20}_{10}\text{Ne}_{10}$	18556.5	18629.8	2.60	2.76	3.01
24	D_{2h}		18600.2		2.86	3.03	
	D_{3d}		22170.1		3.32	3.53	
32	D_{3h}	${}^{24}_{12}\text{Mg}_{12}$	22197.8	22355.8	3.13	3.33	3.06
	O_h	${}^{32}_{16}\text{S}_{16}$	22212.1	29480.5	2.87	3.04	3.26

^aNote that due to its instability there are no data available for beryllium-8, so here we give the charge radius for its isobar lithium-8.

^bThe Hoyle state’s nuclear radius has been measured in Ref. [48].

^cThe experimental matter radius for the carbon-12 ground state is taken from Ref. [32].

agreement with experimental data for a range of baryon numbers, see Table II.

In the following, we will calculate and discuss the electric quadrupole transitions within the Skyrme model. Our discussion will be mainly focused on the quadrupole transition strength, $B(E2) \uparrow$, between the 0^+ ground state and the first 2^+ state in even-even nuclei of zero isospin such as carbon-12. The large quadrupole strength in the 0^+ to 2^+ transition in beryllium-12 will serve as an example how transition strengths can be calculated in the presence of nonzero isospin.

II. ELECTROMAGNETIC TRANSITION STRENGTHS IN THE SKYRME MODEL

In the Skyrme model, the electric charge density $\rho(\mathbf{x})$ [49,50] is given by

$$\rho(\mathbf{x}) = \frac{1}{2}\mathcal{B}(\mathbf{x}) + \mathcal{I}_3(\mathbf{x}), \quad (7)$$

where $\mathcal{B}(\mathbf{x})$ denotes the baryon density, the integrand of Eq. (3), and $\mathcal{I}_3(\mathbf{x})$ is the third component of the isospin density. For quantum states with zero isospin, the charge density ρ is half the baryon density [51], and the total electric charge is $\frac{1}{2}B$ (in units of the proton charge e). Nuclei with zero isospin have equal numbers of protons and neutrons. For nonzero isospin, the isospin density $\mathcal{I}_3(\mathbf{x}) = \omega \mathcal{U}_{33}(\mathbf{x})$ contributes to the total electric charge. Here, ω is the isorotational angular frequency and the isospin inertia density is [17,30]

$$\begin{aligned} \mathcal{U}_{ij}(\mathbf{x}) = & 2\{(\boldsymbol{\pi} \cdot \boldsymbol{\pi} \delta_{ij} - \pi_i \pi_j)(1 + \partial_k \sigma \partial_k \sigma + \partial_k \boldsymbol{\pi} \cdot \partial_k \boldsymbol{\pi}) \\ & - \epsilon_{ide} \epsilon_{jfg} (\pi^d \partial_k \pi^e)(\pi^f \partial_k \pi^g)\}. \end{aligned} \quad (8)$$

The classical electric quadrupole tensor for a Skyrmion is defined as

$$Q_{ij} = \int d^3x (3x_i x_j - |\mathbf{x}|^2 \delta_{ij}) \rho(\mathbf{x}). \quad (9)$$

For a Skyrmion in its standard orientation, the tensor is diagonal and the diagonal entries Q_{11}, Q_{22}, Q_{33} are the quadrupole moments. The quadrupole tensor is traceless (up to numerical inaccuracies) so $Q_{11} + Q_{22} + Q_{33} = 0$. Almost all the Skyrmions we consider here can be orientated so that they have a cyclic symmetry greater than C_2 along the three-axis. Then $Q_{11} = Q_{22}$ and Q_{33} is the quadrupole moment of largest magnitude.

A. Zero isospin

We list in Table III our numerical results for quadrupole moments of Skyrmions and the corresponding nuclei with mass numbers $B = 8, 12, 16, 20, 24, 32$ and zero isospin. Here, the charge density ρ is half the baryon density. The Skyrme model’s predictions are given in Skyrme units and can be converted to physical units by multiplying by the square of the length scale. We orientate the classical Skyrmion such that Q_{33} is the quadrupole moment of maximal magnitude, as discussed above. Then, the nucleus’ intrinsic electric quadrupole moment Q_0 can be identified with $Q_{33} \times [\frac{2}{e_{\text{Sky}} F_\pi}]^2$. With the calibration (5), we obtain the intrinsic quadrupole moments Q_0 (in units of electron barn, eb) which are given in the penultimate column of Table III. For comparison, we also list experimental data, where available. Recall that a factor of $\frac{1}{100}$ is required to convert [fm]² to barn.

For a nuclear state, let J be the total angular momentum and k its projection on the body-fixed three-axis. The reduced electric quadrupole transition strength $B(E2)$ from an initial state $|J_i, k\rangle$ to a final state $|J_f, k\rangle$ can be obtained [1,2] from the intrinsic moment Q_0 via

$$B(E2 : J_i, k \rightarrow J_f, k) = \frac{5}{16\pi} Q_0^2 \langle J_i, k; 20 | J_f, k \rangle^2, \quad (10)$$

where the Clebsch-Gordan coefficient $\langle J_i, k; 20 | J_f, k \rangle$ governs the coupling of the angular momenta.

TABLE III. Intrinsic quadrupole moments for Skyrmions of mass numbers $B = 8, 12, 16, 20, 24, 32$ and of zero isospin. “Sky” and “LM” refer to Skyrme units and the Lau-Manton calibration (5), respectively. We use “—” to denote Skyrmions of zero quadrupole moment. Unless otherwise stated, the experimental results for intrinsic quadrupole moments (in electron barn) are taken from Ref. [5] and have been derived from experimental $B(E2)$ transition strengths via Eq. (12). Note that we state two different intrinsic quadrupole moments for the beryllium-8 nucleus. The experimental quadrupole moment given for oxygen-16 [5] is bracketed since this value has been derived from experimental $B(E2)$ transition strengths from the 0_1^+ ground state to the first-excited 2^+ state. Within the Skyrme model description, this $E2$ transition corresponds to an interband transition and hence cannot be modelled using the techniques described in this paper. See discussion and Fig. 2 in subsequent section on rotational states and transitions in oxygen-16 for more details.

B	G	Nucleus	Q_{11}^{Sky}	Q_{22}^{Sky}	Q_{33}^{Sky}	Q_0^{LM} [eb]	Q_0^{Exp} [eb]
8	D_{4h} (twist)	${}^8_4\text{Be}_4$	-8.54	-8.55	17.10	+0.192	+0.266 ^a , +0.320 ^b
	D_{4h} (no twist)		-10.5	-10.5	21.1	+0.238	
12	D_{4h}	Hoyle ${}^{12}_6\text{C}_6$	-32.1	-32.1	64.3	+0.724	-0.200
	D_{3h}		8.99	9.10	-18.0	-0.203	
16	D_{2d} (bent square)		18.2	18.4	-36.6	-0.412	(0.202)
	D_{4h} (flat square)		21.2	21.3	-42.5	-0.478	
20	T_d	${}^{16}_8\text{O}_8$	—	—	—	—	—
	T_d		—	—	—	—	
24	D_{3h}	${}^{20}_{10}\text{Ne}_{10}$	-18.8	-18.6	37.4	+0.421	+0.584
	D_{2h}		-107	116	-9.16	-0.103	
32	D_{3d}	${}^{24}_{12}\text{Mg}_{12}$	34.4	34.4	-68.9	-0.776	+0.659
	D_{3h}		-13.6	-12.5	26.1	+0.294	
32	O_h	${}^{32}_{16}\text{S}_{16}$	—	—	—	—	+0.549

^aVariational Monte Carlo (VMC) calculation [52].

^bGreens function Monte Carlo (GFMC) method [53].

For electromagnetic transitions between states $J_i = J$ and $J_f = J + 2$, with $k = 0$, the Clebsch-Gordan coefficient in Eq. (10) simplifies to

$$\langle J 0; 2 0 | (J + 2) 0 \rangle^2 = \frac{3(J + 1)(J + 2)}{2(2J + 1)(2J + 3)}. \quad (11)$$

Hence, the reduced electric quadrupole transition probability, $B(E2) \uparrow$, from the spin 0^+ ground state to the first excited spin 2^+ state is given by

$$B(E2 : 0^+ \rightarrow 2^+) = \frac{5}{16\pi} Q_0^2. \quad (12)$$

Note that electromagnetic excitation $B(E2) \uparrow$ and decay $B(E2) \downarrow$ of a nuclear state are related [1,2] by

$$B(E2 : J_f \rightarrow J_i) = \frac{2J_i + 1}{2J_f + 1} B(E2 : J_i \rightarrow J_f). \quad (13)$$

By substituting the intrinsic quadrupole moments Q_0^{LM} listed in Table III in Eq. (12), we obtain the Skyrme model's predictions for the $B(E2) \uparrow$ values (in units of $e^2\text{b}^2$) for nuclei of mass numbers $B = 8, 12, 16, 20, 24, 32$; they are presented in the fifth column of Table IV, with experimental data in the sixth column. In the fourth column of Table IV, we list the corresponding $B(E2) \uparrow$ values in Skyrme units. These $B(E2)^{\text{Sky}}$ values are obtained by substituting Q_{33}^{Sky} given in Table III into Eq. (12). They are related to physical units by the factor $[\frac{2}{e_{\text{sky}} F_\pi}]^4$, the fourth power of the Skyrme length unit. We also include in Table IV the calculated transition strength $B(E2 : 0^+ \rightarrow 2^+)$ for the short-lived ${}^{12}_4\text{Be}_8$ nucleus, to be discussed below, and for the Hoyle state of ${}^{12}_6\text{C}_6$.

To further simplify comparison with experimental data we convert to Weisskopf units W . This compares the transition

strength with the single-particle strength

$$B(E2) \uparrow_{\text{sp}} = 2.97 \times 10^{-5} B^{\frac{4}{3}} e^2\text{b}^2. \quad (14)$$

The strength in Weisskopf units is $W = B(E2) \uparrow / B(E2) \uparrow_{\text{sp}}$, and is a measure of collective quadrupole effects in nuclei. A value higher than 5 indicates substantial collectivity.

In the following subsections, we discuss each nucleus separately. The structure and excitation spectrum of oxygen-16 are particular difficult to understand within a shell-model description [55]. Recent progress has been made via *ab initio* calculations using α cluster initial states with tetrahedral and square configurations [56]. Within the Skyrme model, tetrahedral and square-like configurations of charge-4 subunits arise as $B = 16$ Skyrme solutions [11]. For this reason, we devote a separate, longer section of this paper to rotational states and transitions in oxygen-16.

1. Beryllium-8

For beryllium-8, we calculate $B(E2 : 0^+ \rightarrow 2^+)$ transition strengths using the two known D_{4h} -symmetric Skyrmions. For the twisted $B = 8$ Skyrme we find $B(E2) = 0.00366 e^2\text{b}^2$ and for the untwisted Skyrme $B(E2) = 0.00563 e^2\text{b}^2$. Due to the instability of beryllium-8 to α decay, we are unable to compare our results with actual experimental data. Instead, we include in Table IV $B(E2)$ values based on Hartree-Fock calculations [5] and on Monte Carlo methods [52,53]. Note that the available theoretical values vary significantly depending on which model is used. This makes it impossible to test the accuracy of our Skyrme model $B(E2)$ predictions. Our predicted intrinsic quadrupole moments are consistent with

TABLE IV. Quadrupole transition strengths $B(E2) \uparrow$ for nuclei of baryon numbers $B = 8, 12, 16, 20, 24, 32$. “Sky” and “LM” refer to Skyrme units and the Lau-Manton calibration (5), respectively. “—” denotes zero transition strength. Unless otherwise stated, the experimental $B(E2) \uparrow$ values are taken from Ref. [5]. Note that we state three different estimated transition strengths for the beryllium-8 nucleus.

B	G	Nucleus	$B(E2)^{\text{Sky}}$	$B(E2)^{\text{LM}} [e^2b^2]$	$B(E2)^{\text{Exp}} [e^2b^2]$	Dev. [%]
8	D_{4h} (twist)	${}^8_4\text{Be}_4$	29.1	0.00366 (7.7 W)	0.003 ^a 0.0100 ^b 0.00740 ^c	22% 63.3% 50.5%
8	D_{4h} (no twist)		44.6	0.00563 (11.8 W)		
12	D_{4h}	Hoyle	411	0.0521 (63.9 W)		
	D_{3h}	${}^{12}_6\text{C}_6$	32.5	0.00409 (5.0 W)	0.00397 (4.9 W)	3.02%
	D_{3h}	${}^{12}_4\text{Be}_8$	14.2	0.00181 (2.2 W)	0.0040 (4.9 W) ^d	54.7%
16	D_{2d} (bent square)	${}^{16}_8\text{O}_8$	133	0.0168 (14.1 W)		
	D_{4h} (flat square)		179	0.0227 (18.9 W)		
20	D_{3h}	${}^{20}_{10}\text{Ne}_{10}$	139	0.0176 (10.9 W)	0.0340 (21 W)	48.1%
24	D_{3h}	${}^{24}_{12}\text{Mg}_{12}$	68.1	0.00864 (4.2 W)	0.0432 (21 W)	80.0%
32	O_h	${}^{32}_{16}\text{S}_{16}$	—	—	0.0300 (9.8 W)	

^aHartree-Fock+BCS calculations with the Skyrme SIII force [5].

^bVariational Monte Carlo (VMC) calculation [52].

^cGreen’s function Monte Carlo (GFMC) method [53].

^dFor the short-lived beryllium-12 isotope we obtain the estimated experimental $B(E2) \uparrow$ value by multiplying by 5 the $B(E2) \downarrow$ value measured in Ref. [54].

the prolate shape found in Hartree-Fock and Monte Carlo calculations.

2. Carbon-12 and Hoyle state

For $B = 12$, rotational excitations of the D_{3h} triangular Skyrmion solution match the carbon-12 ground state band, and excitations of the D_{4h} chain solution reproduce the Hoyle band [18]. The D_{3h} -symmetric Skyrmion has the oblate shape assumed for the carbon-12 nucleus [57]. Our calculated intrinsic quadrupole moment $Q_0 = -0.203 \text{ eb}$ agrees well with the experimental value $Q_0 = -0.200 \text{ eb}$ [5] extracted from the measured strength of the $0_1^+ \rightarrow 2_1^+$ transition. The associated transition strength $B(E2 : 0_1^+ \rightarrow 2_1^+) = 0.00409 \text{ e}^2\text{b}^2$ deviates by 3% from the experimental value.

Measuring the $E2$ transition strength from the 2_2^+ Hoyle state to the 0_2^+ Hoyle state is experimentally challenging [58] and would require a highly efficient particle- γ experimental setup [3]. Interpreting the Hoyle state as a linear chain formed out of three $B = 4$ Skyrmons, we predict the transition strength in the opposite direction, $B(E2 : 0_2^+ \rightarrow 2_2^+) = 0.0521 \text{ e}^2\text{b}^2$. This corresponds to 63.9 W, arising from a strongly prolate intrinsic shape with an intrinsic quadrupole moment $Q_0 = 0.724 \text{ eb}$.

3. Neon-20

In the α -particle model, neon-20 is described in terms of five α particles arranged in a triangular bipyramid [59]. Four of the five low-lying rotational bands in neon-20 can be understood [60] using this bipyramidal α -particle arrangement. In the Skyrme model, there exists an analogous bipyramidal cluster arrangement [11] of five $B = 4$ cubes [see Fig. 1(i)]. This D_{3h} -symmetric configuration is not the global minimal energy Skyrmion with $B = 20$, but a nearby saddle point solution. In agreement with experimental data, this bipyramidal Skyrmion structure gives a prolate deformed

neon-20 ground state. The associated intrinsic quadrupole moment $Q_0 = 0.421 \text{ eb}$ is less than the experimental value $Q_0 = 0.584 \text{ eb}$ [5] deduced from the measured $B(E2) \uparrow$ value. For the electric quadrupole transition from the 0^+ ground state to the first excited 2^+ state, the corresponding $B(E2) = 0.0176 \text{ e}^2\text{b}^2$ is approximately 50% less than the experimental value. We have not yet identified any states of neon-20 with the quantized states of the T_d -symmetric $B = 20$ Skyrmion.

4. Magnesium-24

For $B = 24$, we consider three very different Skyrmion solutions: a nonplanar D_{3d} -symmetric ring formed of six $B = 4$ Skyrmion cubes with each neighboring pair being rotated through 90° around the line joining the cubes (see baryon density isosurface in Fig. 1(k) and Ref. [12]), a triaxial configuration constructed by gluing together two linear $B = 12$ Skyrmons [see Fig. 1(j)], and two triangular $B = 12$ Skyrmons bound together into a $B = 24$ solution [see Fig. 1(i)]. The ring was previously believed to be the Skyrmion of minimal energy, but at least one of the other, newly found solutions appears to have lower energy.

Among these Skyrmons, we find that magnesium-24 is probably best described by the D_{3h} -symmetric solution made of two triangular $B = 12$ Skyrmons. The quadrupole moment is found to be $Q_0 = 0.294 \text{ eb}$ which is still significantly less than the experimental value $Q_0 = 0.659 \text{ eb}$. The corresponding quadrupole transition strength $B(E2) = 0.00864 \text{ e}^2\text{b}^2$ is much lower than the experimental value $B(E2) = 0.0432 \text{ e}^2\text{b}^2$. The quadrupole moment of the Skyrmion ring solution has the wrong sign (compare Table III) and does not reproduce the prolate ground state of magnesium-24. The new $B = 24$ solution in Fig. 1(j) is triaxial and is badly approximated as an axially symmetric solution. Hence our analysis cannot be applied to this Skyrmion solution. However, this might give a better quadrupole moment and $B(E2)$ value.

5. Sulphur-32

Calculating nuclear properties of sulphur-32 has proven to be difficult in the past [61] and earlier work using Hartree-Fock calculations on the rotational spectra in sulphur-32 yielded contradictory, model-dependent results. The experimental excitation energies of the 0^+ , 2^+ , and 4^+ states of sulphur-32 agree very well with the vibrational excitations of a spherically shaped nucleus [62]. However, experimentally sulphur-32 possesses a relatively large positive quadrupole moment [5,63] which suggests a significant prolate nuclear deformation. This can be understood within the nuclear coexistence model [64] for sulphur-32, in which spherical and prolate rotational bands coexist.

In the Skyrme model, sulphur-32 is modelled by the cubically symmetric $B = 32$ Skyrmion (see baryon density isosurface in Fig. 1(m) and Ref. [11]) and hence its intrinsic quadrupole moment vanishes. However, this Skyrmion is still of interest because it is a candidate to model the vibrational excitations of sulphur-32. As the Skyrmion spins, the Skyrmion deforms [30,42–44] and a nonzero quadrupole moment will be induced. A calculation of $E2$ transitions for nonrigidly spinning Skyrmion solutions requires different techniques and is beyond the scope of this paper.

III. ROTATIONAL STATES AND TRANSITIONS IN OXYGEN-16

Oxygen-16 has previously been investigated within the α cluster model [65] and by performing lattice effective field theory calculations [56]. The results suggest that there exist two rotational bands, one based on a tetrahedral arrangement of α clusters, and another on a square-like arrangement. This section discusses whether such an interpretation is possible in the Skyrme model.

Here, we follow a similar analysis to the previous description for the rotational bands of carbon-12 and its Hoyle state [18]. For oxygen-16, the ground state is 0^+ and the first 3^- state is lower in energy than the first 2^+ state. This is the signature of the rotational spectrum of a tetrahedral object. In the Skyrme model, several $B = 16$ solutions are known. They are constructed from four $B = 4$ cubic Skyrmions arranged in a bent square (D_{2d}), flat square (D_{4h}) and tetrahedral (T_d) configuration, respectively [see Figs. 1(e)–1(g)]. Hence, we interpret the ground state band in terms of the tetrahedral Skyrmion. The quantized T_d -symmetric Skyrmion models the ground state of oxygen-16 and its 3^- and 4^+ rotational excitations.

The bent and flat square Skyrmions have similar energy and can be seen as energy degenerate within the limits of our numerical accuracy. Note that we cannot confirm the result of the article [11] that the flat square is of noticeable higher energy than the bent square.

For the $B = 16$ D_{4h} -symmetric, flat square Skyrmion, the Finkelstein-Rubinstein (F-R) constraints on a wave function ψ with zero isospin are

$$e^{i\frac{\pi}{2}\hat{L}_3}|\psi\rangle = |\psi\rangle \quad \text{and} \quad e^{i\pi\hat{L}_1}|\psi\rangle = |\psi\rangle, \quad (15)$$

where \hat{L}_i is the spin operator projected on the body-fixed i th axis. k is the eigenvalue of \hat{L}_3 . There is a $k = 0$ rotational band, but the first constraint excludes states with $k = 2$. For $J = 2$,

the only state allowed by the D_{4h} symmetry is $|2, 0\rangle$. The parity operator of this Skyrmion quantized with zero isospin is the identity operator. Hence, all the states have positive parity. This misses important states in the oxygen-16 spectrum, so we turn to the bent square.

For the $B = 16$ D_{2d} -symmetric, bent square Skyrmion, the F-R constraints are

$$e^{i\pi\hat{L}_3}|\psi\rangle = |\psi\rangle \quad \text{and} \quad e^{i\pi\hat{L}_1}|\psi\rangle = |\psi\rangle, \quad (16)$$

and the parity operator is

$$\hat{P} = e^{i\frac{\pi}{2}\hat{L}_3}. \quad (17)$$

In this case, there is a rotational band with $k = 0$ and a band with $k = 2$, and there are two $J = 2$ states, $|2, 0\rangle$ and $\frac{1}{\sqrt{2}}(|2, 2\rangle + |2, -2\rangle)$, with $k = 0$ and $k = 2$, respectively. The states in the $k = 0$ band have positive parity while the states in the $k = 2$ band have negative parity. These bands can be identified with $0^+, 2^+, 4^+$ and $2^-, 3^-, 4^-$ states in the experimentally measured spectrum. Hence, the quantized $B = 16$ bent square is preferable for modeling the second excited spin-0 state 0_2^+ of oxygen-16 and its rotational excitations.

To test further this identification we can use similar techniques as applied in Ref. [18] to the rotational excitations of carbon-12 and its Hoyle state. All $B = 16$ Skyrmion solutions have inertia tensors of symmetric-top type with $V_{11} = V_{22}$ and V_{33} the same or distinct. For the tetrahedral solution we find $V_{11} = V_{22} = 9100$ and $V_{33} = 9128$, where the difference must be a numerical artifact. For the bent square, $V_{11} = V_{22} = 9123$ and $V_{33} = 14602$.

The energy eigenvalues of the quantum Hamiltonian for purely rotational motion of a symmetric top are given by

$$E(J, k) = C \left\{ \frac{1}{2V_{11}} J(J+1) + \left(\frac{1}{2V_{33}} - \frac{1}{2V_{11}} \right) k^2 \right\}, \quad (18)$$

where J denotes the total spin quantum number, k is the eigenvalue of \hat{L}_3 , and C is a dimensional conversion factor from Skyrme to physical units [18]. Here, C is a purely phenomenological parameter as has previously been used in the discussion of carbon-12 and its rotational states in Ref. [18]. In Fig. 2, we plot against $J(J+1)$ the energies of experimentally observed oxygen-16 states up to spin 4 in the ground-state band and in the rotational bands formed by the rotational excitations of the 0_2^+ state and the 2^- state. Taking C as our fitting parameter, we fit Eq. (18) to the $0^+, 3^-, 4^+$ states of the ground-state band, whose energies are 0.0, 6.13, and 10.36 MeV. The linear fit gives $C = 9418$ MeV, and therefore a best fit slope of 0.517 MeV, using the V_{11} value 9100 for the tetrahedral Skyrmion.

We find that the experimental slope of the $k = 0$ band based on the 0_2^+ state then agrees very well with the Skyrme model prediction. Eq. (18) gives a theoretical slope of 0.516 MeV, where we used the bent square's moment of inertia $V_{11} = 9123$ and $C = 9418$ MeV as derived above. The experimental slope is estimated from the best linear fit to the $0_2^+, 2^+, 4^+$ states with energies 6.05, 9.84, and 16.84 MeV to be 0.545 MeV. The Skyrme model prediction for the ratio of the slopes is just the ratio of the V_{11} values for the D_{2d} - and T_d -symmetric Skyrmions, which is $9123/9100 = 1.00$. Note




FIG. 2. Experimental states of oxygen-16. The symbol triangle denotes the states of the ground state band, and circle and square denote the $k = 0, 2$ states of the “bent square” bands. The $0_1^+, 3^-, 4^+$ states in the ground state band have energies 0.0, 6.13, and 10.36 MeV. For the “bent square” band, the $0_2^+, 2^+, 4^+$ states with $k = 0$ have energies 6.05, 9.84, and 16.84 MeV and the $2^-, 3^-, 4^-$ states with $k = 2$ have energies 8.87, 11.60, and 14.30 MeV. The symbol cross represents the 2_1^+ state which has been interpreted as a rotational excitation of the 0_2^+ state by Epelbaum *et al.* [56]. See the text for more details.

that the dimensional conversion factor C cancels. For comparison, the experimental ratio of the slopes is 1.05.

We also include in Fig. 2 the experimental $2^-, 3^-, 4^-$ states of energies 8.87, 11.60, and 14.30 MeV which we interpret as the $k = 2$ band formed by the rotational excitations of the bent square. The $k = 2$ band lies below the $k = 0$ band, agreeing with the oblateness of the bent square Skyrmion. For an oblate configuration, $V_{11} < V_{33}$, and according to Eq. (18), for a fixed spin J , the energy of a state with nonzero k has lower energy than a $k = 0$ state. The predicted energy difference between states with the same spin J is 0.77 MeV between the $k = 0$ and $k = 2$ bands. This agrees marginally with the experimentally measured differences of 0.97 MeV for the spin-2 states and 2.54 MeV for the spin-4 states.

We calculate the $E2$ transition strength from the 0_2^+ state of energy 6.05 MeV to the 2^+ state of energy 9.84 MeV by taking the bent square Skyrmion as the underlying structure. We obtain $B(E2 : 0_2^+ \rightarrow 2^+) = 0.0168 e^2 b^2$. The associated intrinsic, oblate quadrupole moment is $Q_0 = -0.412 eb$. For this transition, we are unable to find experimental $E2$ values in the literature. For completeness, we also include in Tables III and IV the quadrupole moment and $B(E2)$ values when modeling the 0_2^+ state and its spin-2 excitation 2^+ by the flat square Skyrmion.

Epelbaum *et al.* [56] have considered the transition between the 0_2^+ state and the lowest spin-2 state 2_1^+ of energy 6.91 MeV. These states are represented in Fig. 2 by the cross symbols. They interpret the 2_1^+ state as a rotational excitation of a square configuration of α clusters. However, it is badly described as a rotational excitation of the bent or flat square $B = 16$

Skyrmion. The $B(E2 : 2_1^+ \rightarrow 0_2^+)$ transition strength predicted in Ref. [56] is based on nuclear lattice effective field theory simulations. The up transition strength is $B(E2 : 0_2^+ \rightarrow 2_1^+) = 0.0110 e^2 b^2$. The corresponding empirical value is found to be $B(E2 : 0_2^+ \rightarrow 2_1^+) = 0.0325 e^2 b^2$ [56,66].

IV. NONZERO ISOSPIN

In the previous sections, we restricted the discussion to $E2$ transitions in the absence of isospin. In this section, we show how quadrupole transition strengths can be calculated within the Skyrme model in the presence of nonzero nuclear isospin.

A. Beryllium-12

Beryllium-12 is a nucleus in an $I = 2$ isospin multiplet, as it has four protons and eight neutrons. Nuclei of mass number 12, with isospin 0, 1, and 2 are especially well described within the Skyrme model as quantum states of the D_{3h} -symmetric $B = 12$ Skyrmion [16] [see baryon density isosurface displayed in Fig. 1(d)]. In particular, the low-lying energy levels of beryllium-12, with various spins, appear as states with $I = 2$ and $I_3 = -2$.

The quantum states $|\psi_{J^\pi, I, |L_3|, |K_3|}\rangle$ of the $B = 12$ Skyrmion allowed by the Finkelstein-Rubinstein constraints [13,14] are listed in Table VI of Ref. [16]. J and I are the total spin and isospin labels, and $|L_3|$ and $|K_3|$ are the projections on to “body-fixed” three-axes. Both for $J = 0$ and $J = 2$ there is a unique allowed $I = 2$ state, with $K_3 = 0$. The “space-fixed” isospin I_3 can take any integer value from -2 to 2 , and for beryllium-12 it is $I_3 = -2$. Suppressing the spin state, we denote the isospin state of beryllium-12 as $|2, 0; -2\rangle$.

Beryllium-12 exhibits a large quadrupole strength in the transition between the 0^+ ground state and the first 2^+ state at 2.1 MeV [54]. In Ref. [54], a $B(E2 : 2^+ \rightarrow 0^+)$ value of $0.0008 e^2 b^2$ has been determined through the lifetime measurement of the 2^+ state. Using Eq. (13) this results in a $B(E2 : 0^+ \rightarrow 2^+)$ value of $0.0040 e^2 b^2$.

Here, we consider $E2$ transitions between these spin states in the Skyrme model, using the Skyrmion’s isospin state $|2, 0; -2\rangle$. The new aspect is to take account of the contribution of the isospin to the electric charge density, and hence to the quadrupole moments.

It would be best to do a proper quantum calculation of the expectation value of the quadrupole moments, but we have not been able to do this. Instead we treat the isospin state using a classical approximation. This is analogous to the approach to nucleons adopted in Refs. [31,46]. There a nucleon is treated as a classically spinning $B = 1$ Skyrmion, which gives it both spin and isospin. To achieve the desired “space-fixed” isospin projection, the $B = 12$ Skyrmion’s red/green/blue colors spin in isospace while the black/white colours do not. (This produces a rotation among the π_1 and π_2 fields, while the π_3 field, associated with the black/white axis, remains constant.) The role of the isospin state $|2, 0; -2\rangle$ is to inform us of the most likely colouring of the Skyrmion, before the colors spin. If the projection in this state was $K_3 = \pm 2$ then the standard orientation and coloring of the Skyrmion would be the correct one, but as the projection is $K_3 = 0$ we must reorientate the colorings first.

We reorientate the classical $B = 12$ Skyrmion colouring by the angles that maximize the wave function $|2, 0; -2\rangle$. The associated Wigner \mathcal{D} function for this state takes the form $\mathcal{D}_{0,-2}^2(\alpha, \beta, \gamma) = e^{-2i\gamma} \sin^2 \beta$, where α, β, γ are the isorotational Euler angles. The factor $e^{-2i\gamma}$ is the quantum representation of the colours spinning with $I_3 = -2$. $e^{-2i\gamma} \sin^2 \beta$ has its maximum magnitude at $\beta = \pi/2$ and $\gamma = 0$ (or any other value of γ). Hence, we perform an isospin rotation of our initial classical solution with $\beta = \pi/2$ and $\gamma = 0$. This rotates the black/white points of the Skyrmion to be on the faces or edges of the $B = 4$ cube constituents, rather than at the vertices.

In detail, under an isospin rotation with $\beta = \pi/2$ and $\gamma = 0$ (and α undetermined) the pion field $\boldsymbol{\pi} = (\pi_1, \pi_2, \pi_3)$ transforms to

$$\begin{aligned} \pi'_1 &= -\sin \alpha \pi_2 + \cos \alpha \pi_3, & \pi'_2 &= \cos \alpha \pi_2 + \sin \alpha \pi_3, \\ \pi'_3 &= -\pi_1, \end{aligned} \quad (19)$$

and hence the new moment of inertia that we need is $U'_{33} = U_{11}$. The colors spin about the new three-axis in isospace, but dynamically this is equivalent to spinning about the old one-axis. For the D_{3h} -symmetric $B = 12$ Skyrmion we find $U'_{33} = U_{11} = 442$, see Table I. To classically model a beryllium-12 nucleus whose projected isospin has magnitude -2 we require $U'_{33}\omega = -2\hbar$. (Physical isospin, like spin, is a half-integer or integer multiple of \hbar .)

Hence, the isorotational angular velocity ω is given by

$$\omega = -\frac{2\hbar}{U'_{33}} = -0.14, \quad (20)$$

where $\hbar = 30.2$. For the beryllium-12 nucleus, the classical isospin density is

$$\mathcal{I}_3(\mathbf{x}) = \frac{\omega}{\hbar} \mathcal{U}'_{33}(\mathbf{x}). \quad (21)$$

This isospin contribution to the electric charge density (7) has to be taken into account when calculating quadrupole moments (9). Note that Eq. (21) is correctly normalized as its integral gives $I_3 = -2$ and decreases the total electric charge $\frac{1}{2}B + I_3$ of the $B = 12$ Skyrmion from 6 to 4.

We compute numerically the electric quadrupole moments $Q_{11}^{\text{Sky}} = 5.65$, $Q_{22}^{\text{Sky}} = 6.32$, and $Q_{33}^{\text{Sky}} = -11.9$ for the reorientated $B = 12$ Skyrmion. Thus, expressed in physical units using the calibration (5) the intrinsic quadrupole is $Q_0 = -0.135$ eb. The corresponding $B(E2 : 0 \rightarrow 2)$ value is $0.00181 e^2 b^2$ which differs by approximately 50% from the experimental value.

V. CONCLUSIONS

We have calculated for the first time within the Skyrme model electromagnetic transition strengths between the 0^+ ground state and the first-excited 2^+ state for a range of light nuclei: ${}^8_4\text{Be}$, ${}^{12}_6\text{C}$ and its Hoyle state, ${}^{12}_4\text{Be}$, ${}^{20}_{10}\text{Ne}$, and ${}^{24}_{12}\text{Mg}$. We find that the calculated $E2$ transition strengths have the correct order of magnitude and the computed intrinsic quadrupole moments match the experimentally observed effective nuclear shapes. For the Hoyle state we predict a large $B(E2) \uparrow$ value of $0.0521 e^2 b^2$. Measurements of the

electromagnetic transition strengths between the states of the Hoyle band are technically difficult and have yet to be performed [3,58].

For oxygen-16, we can obtain a quantitative understanding of the ground state band and the rotational band formed by the second excited spin-0 state 0_2^+ and its rotational excitations. Similarly to the ground state band of carbon-12 and the rotational band of the Hoyle state [18], we interpret the oxygen-16 rotational bands as rotational excitations of two Skyrmons with very different shapes, one tetrahedral and the other a bent square. The quantized tetrahedral Skyrme models the 0_1^+ ground state and its 3^- and 4^+ excitations. The quantized bent square Skyrme configuration is identified with the 0_2^+ state and its rotational excitations. We find that the $0_2^+, 2^+, 4^+$ states of energies 6.05, 9.84 and 16.84 MeV are very well modelled as $k = 0$ states of the bent square band. The almost equal values of the spin moment of inertia V_{11} for the tetrahedron and bent square are a success of the Skyrme model. The Skyrme model predicts that the ratio of the slopes of the $k = 0$ bent square band and the ground state band is the ratio of these V_{11} values, and is very close to 1. The ratio of the experimental slopes agrees with this. The $k = 2$ band of the bent square matches experimental $2^-, 3^-, 4^-$ states. Furthermore, we used Eq. (18) to predict the energy difference between $k = 0$ and $k = 2$ states of the same spin (but opposite parity). The predicted energy difference has the right sign and marginally agrees with experiment.

There remain some challenges for the Skyrme model. For baryon number 20, the triangular bipyramidal arrangement of five $B = 4$ cubes which we used to describe $E2$ transitions in neon-20 is not a minimal energy Skyrme but a saddle point. For $B = 32$, the minimal energy Skyrme is cubically symmetric and hence cannot explain the large prolate quadrupole moment of sulphur-32. However, Skyrmons deform under rotations and hence a non-zero quadrupole moment may be induced.

The approach used in this paper is limited to Skyrmons with axially symmetric inertia tensors, and with quadrupole moments satisfying $Q_{11} = Q_{22}$. The calculation of $B(E2)$ strengths for transitions between rotational levels in triaxial nuclei [67,68] using the Skyrme model requires a different approach. Further lines of investigation to consider are higher-order electric multipole transitions and magnetic dipole transitions within the Skyrme model. In particular, the 0^+ to 3^- transition strength in the oxygen-16 ground state band should be calculated. Finally, we neglected deformations in our calculations; that is, we assumed that the low-lying rotational states are well approximated by the rigid rotor states of the Skyrmons. Recently, $E2$ transitions in deformed nuclei have been studied within an effective theory for axially symmetric systems [69], and a similar study of deformed Skyrmons is desirable.

ACKNOWLEDGMENTS

This work was partly undertaken on the COSMOS Shared Memory system at DAMTP, University of Cambridge operated on behalf of the STFC DiRAC HPC Facility. This equipment is funded by BIS National E-infrastructure capital grant no. ST/J005673/1 and STFC grants no. ST/J001341/1,

ST/H008586/1, ST/K00333X/1. M.H. has been partially funded by the UK Science and Technology Facilities Council under grant no. ST/J000434/1. M.H. thanks Andrzej Wereszczynski and the Jagiellonian University, Krakow for

hospitality. P.H.C.L. thanks Ling-Yan Hung and Fudan University in Shanghai for hospitality. P.H.C.L. acknowledges support as an International Research Fellow of the Japan Society for the Promotion of Science (JSPS).

-
- [1] A. Bohr and B. R. Mottelson, *Nuclear Structure*, Vol. I: Single Particle Motion (World Scientific, Singapore, 1998).
- [2] A. Bohr and B. R. Mottelson, *Nuclear Structure*, Vol. II: Nuclear Deformations (World Scientific, Singapore, 1998).
- [3] D. G. Jenkins, in *Lecture Notes in Physics*, edited by C. Beck (Springer Verlag, Berlin, 2014), Vol. 875, p. 25.
- [4] K. H. Bhatt, C. W. Nestor, and S. Raman, *Phys. Rev. C* **46**, 164 (1992).
- [5] S. Raman, C. W. Nestor, and P. Tikkanen, *At. Data Nucl. Data Tables* **78**, 1 (2001).
- [6] B. Pritychenko, M. Birch, M. Horoi, and B. Singh, *Nucl. Data Sheets* **120**, 111 (2014).
- [7] T. H. R. Skyrme, *Proc. R. Soc. London A* **260**, 127 (1961).
- [8] T. H. R. Skyrme, *Nucl. Phys.* **31**, 556 (1962).
- [9] J. Blatt and V. Weisskopf, *Theoretical Nuclear Physics* (Wiley, New York, 1952).
- [10] D. Brink, H. Friedrich, A. Weiguny, and C. Wong, *Phys. Lett. B* **33**, 143 (1970).
- [11] R. A. Battye, N. S. Manton, and P. M. Sutcliffe, *Proc. R. Soc. London A* **463**, 261 (2007).
- [12] D. T. J. Feist, P. H. C. Lau, and N. S. Manton, *Phys. Rev. D* **87**, 085034 (2013).
- [13] S. Krusch, *Ann. Phys. (NY)* **304**, 103 (2003).
- [14] S. Krusch, *Proc. R. Soc. London A* **462**, 2001 (2006).
- [15] O. V. Manko, N. S. Manton, and S. W. Wood, *Phys. Rev. C* **76**, 055203 (2007).
- [16] R. A. Battye, N. S. Manton, P. M. Sutcliffe, and S. W. Wood, *Phys. Rev. C* **80**, 034323 (2009).
- [17] P. H. C. Lau and N. S. Manton, *Phys. Rev. D* **89**, 125012 (2014).
- [18] P. H. C. Lau and N. S. Manton, *Phys. Rev. Lett.* **113**, 232503 (2014).
- [19] G. Adkins, C. Nappi, and E. Witten, *Nucl. Phys. B* **228**, 552 (1983).
- [20] E. Braaten and L. Carson, *Phys. Rev. D* **38**, 3525 (1988).
- [21] V. B. Kopeliovich, *JETP* **93**, 435 (2001).
- [22] N. S. Manton and P. M. Sutcliffe, *Topological Solitons* (Cambridge University Press, Cambridge, 2004).
- [23] *The Multifaceted Skyrmion*, edited by G. E. Brown and M. Rho, (World Scientific, Singapore, 2010).
- [24] T. Gisiger and M. B. Paranjape, *Phys. Rep.* **306**, 109 (1998).
- [25] G. Adkins and C. Nappi, *Nucl. Phys. B* **233**, 109 (1984).
- [26] J. R. Shewchuk, An introduction to the conjugate gradient method without the agonizing pain (1994), <http://www.cs.cmu.edu/~quake-papers/painless-conjugate-gradient.pdf>.
- [27] D. T. J. Feist, *JHEP* **02** (2012) 100.
- [28] R. A. Battye and P. M. Sutcliffe, *Rev. Math. Phys.* **14**, 29 (2002).
- [29] V. B. Kopeliovich, *Yad. Fiz.* **47**, 1495 (1988) [*Sov. J. Nucl. Phys.* **47**, 949 (1988)].
- [30] R. A. Battye, M. Haberichter, and S. Krusch, *Phys. Rev. D* **90**, 125035 (2014).
- [31] N. S. Manton, *Math. Method. Appl. Sci.* **35**, 1188 (2012).
- [32] I. Tanihata, H. Hamagaki, O. Hashimoto, Y. Shida, N. Yoshikawa, K. Sugimoto, O. Yamakawa, T. Kobayashi, and N. Takahashi, *Phys. Rev. Lett.* **55**, 2676 (1985).
- [33] B. Schwesinger and H. Weigel, *Phys. Lett. B* **267**, 438 (1991).
- [34] H. Weigel, *Int. J. Mod. Phys. A* **11**, 2419 (1996).
- [35] B. Moussallam, *Ann. Phys. (NY)* **225**, 264 (1993).
- [36] F. Meier and H. Walliser, *Phys. Rep.* **289**, 383 (1997).
- [37] B. Moussallam and D. Kalafatis, *Phys. Lett. B* **272**, 196 (1991).
- [38] G. Holzwarth and H. Walliser, *Nucl. Phys. A* **587**, 721 (1995).
- [39] H. Weigel, R. Alkofer, and H. Reinhardt, *Nucl. Phys. A* **582**, 484 (1995).
- [40] F. G. Weigel, B. Schwesinger, and H. B. Geyer, *Nucl. Phys. A* **561**, 542 (1993).
- [41] C. J. Halcrow, *Nucl. Phys. B* **904**, 106 (2016).
- [42] R. A. Battye, S. Krusch, and P. M. Sutcliffe, *Phys. Lett. B* **626**, 120 (2005).
- [43] C. Houghton and S. Magee, *Phys. Lett. B* **632**, 593 (2006).
- [44] J. Fortier and L. Marleau, *Phys. Rev. D* **77**, 054017 (2008).
- [45] D. Foster and S. Krusch, *Nucl. Phys. B* **897**, 697 (2015).
- [46] D. Foster and N. S. Manton, *Nucl. Phys. B* **899**, 513 (2015).
- [47] I. Angeli and K. P. Marinova, *At. Data Nucl. Data Tables* **99**, 69 (2013).
- [48] A. N. Danilov, T. L. Belyaeva, A. S. Demyanova, S. A. Goncharov, and A. A. Ogloblin, *Phys. Rev. C* **80**, 054603 (2009).
- [49] E. Witten, *Nucl. Phys. B* **223**, 422 (1983).
- [50] E. Witten, *Nucl. Phys. B* **223**, 433 (1983).
- [51] N. S. Manton and S. W. Wood, *Phys. Rev. D* **74**, 125017 (2006).
- [52] R. B. Wiringa, S. C. Pieper, J. Carlson, and V. R. Pandharipande, *Phys. Rev. C* **62**, 014001 (2000).
- [53] V. M. Datar, D. R. Chakrabarty, Suresh Kumar, V. Nanal, S. Pastore, R. B. Wiringa, S. P. Behera, A. Chatterjee, D. Jenkins, C. J. Lister, E. T. Mirgule, A. Mitra, R. G. Pillay, K. Ramachandran, O. J. Roberts, P. C. Rout, A. Shrivastava, and P. Sugathan, *Phys. Rev. Lett.* **111**, 062502 (2013).
- [54] N. Imai *et al.*, *Phys. Lett. B* **673**, 179 (2009).
- [55] H. Nagai, *Prog. Theor. Phys.* **27**, 619 (1962).
- [56] E. Epelbaum, H. Krebs, T. A. Lähde, D. Lee, Ulf-G. Meißner, and G. Rupak, *Phys. Rev. Lett.* **112**, 102501 (2014).
- [57] H. Ebsensen, X. Tang, and C. L. Jiang, *Phys. Rev. C* **84**, 064613 (2011).
- [58] M. Freer and H. Fynbo, *Prog. Part. Nucl. Phys.* **78**, 1 (2014).
- [59] W. Wefelmeier, *Z. Phys.* **107**, 332 (1937).
- [60] M. Bouten, *Nuovo Cimento* **26**, 63 (1962).
- [61] W. Bauhoff, H. Schultheis, and R. Schultheis, *Phys. Rev. C* **22**, 861 (1980).
- [62] A. Kangasmäki, P. Tikkanen, J. Keinonen, W. E. Ormand, S. Raman, Zs. Fülöp, Á. Z. Kiss, and E. Somorjai, *Phys. Rev. C* **58**, 699 (1998).
- [63] N. Stone, *At. Data Nucl. Data Tables* **90**, 75 (2005).
- [64] Y. T. Cheng, A. Goswami, M. J. Throop, and D. K. McDaniels, *Phys. Rev. C* **9**, 1192 (1974).
- [65] B. Buck, C. Dover, and J. Vary, *Phys. Rev. C* **11**, 1803 (1975).
- [66] F. Ajzenberg-Selove, *Nucl. Phys. A* **166**, 1 (1971).
- [67] V. A. Krutov and N. V. Zackrevsky, *J. Phys. A* **2**, 448 (1969).
- [68] J. M. Allmond, R. Zaballa, A. M. Oros-Peusquens, W. D. Kulp, and J. L. Wood, *Phys. Rev. C* **78**, 014302 (2008).
- [69] E. A. Coello Pérez and T. Papenbrock, *Phys. Rev. C* **92**, 014323 (2015).