

Low-energy hypernuclear spectra within a microscopic particle-rotor model with a relativistic point-coupling hyperon-nucleon interaction

H. Mei,^{1,2} K. Hagino,^{1,3,4} J. M. Yao,^{5,2} and T. Motoba^{6,7}

¹*Department of Physics, Tohoku University, Sendai 980-8578, Japan*

²*School of Physical Science and Technology, Southwest University, Chongqing 400715, China*

³*Research Center for Electron Photon Science, Tohoku University, 1-2-1 Mikamine, Sendai 982-0826, Japan*

⁴*National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan*

⁵*Department of Physics and Astronomy, North Carolina University, Chape Hill 27599-3255, USA*

⁶*Laboratory of Physics, Osaka Electro-Communications University, Neyagawa 572-8530, Japan*

⁷*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606-8502, Japan*

(Received 4 February 2016; published 12 April 2016)

We extend the microscopic particle-rotor model for hypernuclear low-lying states by including the derivative and tensor coupling terms in the point-coupling nucleon Λ -particle ($N\Lambda$) interaction. Taking ${}^{13}_{\Lambda}\text{C}$ as an example, we show that a good overall description for excitation spectra is achieved with four sets of effective $N\Lambda$ interactions. We find that the Λ hyperon binding energy decreases monotonically with increasing strengths of the high-order interaction terms. In particular, the tensor coupling term decreases the energy splitting between the first $1/2^-$ and $3/2^-$ states and increases the energy splitting between the first $3/2^+$ and $5/2^+$ states in ${}^{13}_{\Lambda}\text{C}$.

DOI: [10.1103/PhysRevC.93.044307](https://doi.org/10.1103/PhysRevC.93.044307)

I. INTRODUCTION

Spectroscopic data on low-lying states of light Λ hypernuclei have been accumulated [1] and more data on those of medium and heavy hypernuclei are expected to be obtained with next-generation facilities such as J-PARC [2]. Rich information on the hyperon-nucleon interaction in the nuclear medium and on the impurity effect of a Λ particle on nuclear structure is contained in these data. Because hyperon-nucleon and hyperon-hyperon scattering experiments are difficult to perform, the structure of hypernuclei plays a vital role in shedding light on baryon-baryon interactions. Such information is crucial in order to understand also neutron stars, in which hyperons may emerge in the inner part [3]. However, extracting information on baryon-baryon interactions from spectroscopic data relies heavily on nuclear models.

In recent decades, several different types of theoretical models have been developed to study the structure of hypernuclei, including an *ab initio* method [4], a cluster model [5–11], a shell model [12–14], antisymmetrized molecular dynamics (AMD) [15–18], the self-consistent mean-field approach [19–27], and the generator coordinator method (GCM) based on energy density functionals [28]. In recent years, we have also developed a microscopic particle rotor model (MPRM) for hypernuclear low-lying states based on a beyond-mean-field approach [29,30]. In contrast to the GCM for whole hypernuclei [28], where the wave function of the hypernuclear states is given as a superposition of hypernuclear mean-field states, the hypernuclear states in the MPRM are constructed by coupling a hyperon to low-lying states of the core nucleus. The MPRM provides a convenient way to analyze the components of the hypernuclear wave function and has been applied to study the low-lying states of ${}^9_{\Lambda}\text{Be}$ [29], ${}^{13}_{\Lambda}\text{C}$, ${}^{21}_{\Lambda}\text{Ne}$, and ${}^{155}_{\Lambda}\text{Sm}$ hypernuclei [30]. For the sake of simplicity, only the leading-order four-fermion coupling terms of scalar and vector types were adopted for the $N\Lambda$ effective interaction in these studies.

The aim of this paper is to extend the previous calculations by implementing the higher-order derivative and tensor $N\Lambda$ interaction terms in the point-coupling interaction [31]. The derivative terms simulate to some extent the finite-range character of $N\Lambda$ interaction, and these terms are expected to be more pronounced in light hypernuclei [32]. On the other hand, the tensor $N\Lambda$ interaction is important to reproduce a small hyperon spin-orbit splitting in Λ hypernuclei [33]. It is therefore important to assess the effect of these terms on hypernuclear low-lying states.

The paper is organized as follows. In Sec. II, we present the main formulas of the microscopic MPRM for Λ hypernuclei with the full point-coupling effective $N\Lambda$ interaction. In Sec. III, we show the results for hypernuclear low-lying states in ${}^{13}_{\Lambda}\text{C}$ and discuss the influence of the higher-order terms on the energy spectra. We then summarize the paper in Sec. IV.

II. MICROSCOPIC PARTICLE-ROTOR MODEL FOR A HYPERNUCLEI

In this paper, we consider a single- Λ hypernucleus and describe the hypernuclear low-lying states using the microscopic particle-rotor model (MPRM). Since the detailed formulas for the MPRM have been given in Refs. [29,30], we give here only the main formulas of this approach. To this end, we put a particular emphasis on the implementation of the higher-order $N\Lambda$ interaction terms.

The basic idea of the MPRM is to construct a hypernuclear wave function by coupling the valence Λ hyperon to the low-lying states of the nuclear core in the laboratory frame, that is,

$$\Psi_{JM}(\mathbf{r}, \{\mathbf{r}_i\}) = \sum_{n,j,\ell,l} \mathcal{R}_{j\ell n l}(r) \mathcal{F}_{j\ell n l}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}), \quad (1)$$

with

$$\mathcal{F}_{j\ell n l}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) = [\mathcal{Y}_{j\ell}(\hat{\mathbf{r}}) \otimes \Phi_{nl}(\{\mathbf{r}_i\})]^{(JM)}, \quad (2)$$

where \mathbf{r} and \mathbf{r}_i are the coordinates of the Λ hyperon and the nucleons, respectively. Here, J is the angular momentum for the whole system, while M is its projection onto the z axis in the laboratory frame. $\mathcal{Y}_{j\ell}(\hat{\mathbf{r}})$ is the spin-angular wave function for the Λ hyperon. $|\Phi_{nI}\rangle$ are the wave functions for the low-lying states of the core nucleus, where I represents the angular momentum of the core state and $n = 1, 2, \dots$ distinguishes different core states with the same angular momentum I . In the MPRM, the core states $|\Phi_{nI}\rangle$ are constructed with the quantum-number projected GCM approach [29,30]. For convenience, hereafter we introduce the shorthand notation $k = \{j\ell n I\}$ to represent different channels.

In Eq. (1), $\mathcal{R}_k(r)$ is the radial wave function for the Λ particle. In the relativistic approach, it is given as a four-component Dirac spinor

$$\mathcal{R}_k(r) = \begin{pmatrix} f_k(r) \\ i g_k(r) \boldsymbol{\sigma} \cdot \hat{\mathbf{r}} \end{pmatrix}. \quad (3)$$

We assume that the Hamiltonian \hat{H} for the whole Λ hypernucleus is given as

$$\hat{H} = \hat{T}_\Lambda + \hat{H}_c + \sum_{i=1}^{A_c} \hat{V}^{N\Lambda}(\mathbf{r}, \mathbf{r}_i). \quad (4)$$

Here $\hat{T}_\Lambda = -i\boldsymbol{\alpha} \cdot \nabla_\Lambda + \gamma^0 m_\Lambda$ is the relativistic kinetic energy of the Λ hyperon, where m_Λ is the mass of the Λ particle and $\boldsymbol{\alpha}$ and γ^0 are the Dirac matrices. \hat{H}_c is the many-body Hamiltonian for the core nucleus [34], with which the core state $|\Phi_{nI}\rangle$ satisfies $\hat{H}_c |\Phi_{nI}\rangle = E_{nI} |\Phi_{nI}\rangle$. The last term in Eq. (4) represents the $N\Lambda$ interaction term between the valence Λ particle and the nucleons in the core nucleus, where A_c is the mass number of the core nucleus.

We construct the $N\Lambda$ interaction $\hat{V}^{N\Lambda}$ based on the relativistic point-coupling model [31], in which the energy functional for the $N\Lambda$ interaction reads

$$\begin{aligned} E_{\text{int}}^{(N\Lambda)} = & \int d\mathbf{r} [\alpha_S^{N\Lambda} \rho_S(\mathbf{r}) \rho_S^\Lambda(\mathbf{r}) + \alpha_V^{N\Lambda} \rho_V(\mathbf{r}) \rho_V^\Lambda(\mathbf{r}) \\ & + \delta_S^{N\Lambda} \rho_S(\mathbf{r}) \Delta \rho_S^\Lambda(\mathbf{r}) + \delta_V^{N\Lambda} \rho_V(\mathbf{r}) \Delta \rho_V^\Lambda(\mathbf{r}) \\ & + \alpha_T^{N\Lambda} \rho_T^\Lambda(\mathbf{r}) \rho_V(\mathbf{r})]. \end{aligned} \quad (5)$$

Here ρ_S , ρ_V , and ρ_T are the scalar, the vector, and the tensor densities defined in Ref. [31], respectively. Taking the second functional derivative of Eq. (5) with respect to the densities [35],

$$\begin{aligned} \hat{V}^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) = & \frac{\delta^2 E[\rho]}{\delta \rho_S^\Lambda(\mathbf{r}) \delta \rho_S(\mathbf{r}_i)} + \frac{\delta^2 E[\rho]}{\delta \rho_V^\Lambda(\mathbf{r}) \delta \rho_V(\mathbf{r}_i)} \\ & + \frac{\delta^2 E[\rho]}{\delta \rho_T^\Lambda(\mathbf{r}) \delta \rho_V(\mathbf{r}_i)}, \end{aligned} \quad (6)$$

we obtain the following form for the $N\Lambda$ effective interaction:

$$\hat{V}^{N\Lambda} = \hat{V}_S^{N\Lambda} + \hat{V}_V^{N\Lambda} + \hat{V}_{\text{Ten}}^{N\Lambda}, \quad (7)$$

where the scalar, vector, and tensor types of coupling terms read

$$\begin{aligned} \hat{V}_S^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) = & \alpha_S^{N\Lambda} \gamma_\Lambda^0 \delta(\mathbf{r} - \mathbf{r}_i) \gamma_N^0 + \delta_S^{N\Lambda} \gamma_\Lambda^0 [\overleftarrow{\nabla}^2 \delta(\mathbf{r} - \mathbf{r}_i) \\ & + \delta(\mathbf{r} - \mathbf{r}_i) \overleftarrow{\nabla}^2 + 2 \overleftarrow{\nabla} \cdot \delta(\mathbf{r} - \mathbf{r}_i) \overleftarrow{\nabla}] \gamma_N^0, \end{aligned} \quad (8)$$

$$\begin{aligned} \hat{V}_V^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) = & \alpha_V^{N\Lambda} \delta(\mathbf{r} - \mathbf{r}_i) + \delta_V^{N\Lambda} [\overleftarrow{\nabla}^2 \delta(\mathbf{r} - \mathbf{r}_i) \\ & + \delta(\mathbf{r} - \mathbf{r}_i) \overleftarrow{\nabla}^2 + 2 \overleftarrow{\nabla} \cdot \delta(\mathbf{r} - \mathbf{r}_i) \overleftarrow{\nabla}], \end{aligned} \quad (9)$$

$$\hat{V}_{\text{Ten}}^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) = i\alpha_T^{N\Lambda} \gamma_\Lambda^0 [\overleftarrow{\nabla} \delta(\mathbf{r} - \mathbf{r}_i) + \delta(\mathbf{r} - \mathbf{r}_i) \overleftarrow{\nabla}] \cdot \boldsymbol{\alpha}. \quad (10)$$

Here, $\overleftarrow{\nabla}$ and $\overleftarrow{\nabla}$ are understood to act on the right- and left-hand sides of the Λ hyperon coordinates, respectively. Vice versa, Eq. (5) can be obtained from the above effective $N\Lambda$ interaction (see Appendix A). We note that similar terms appear in the chiral hyperon-nucleon interaction [36], in which the nonderivative four-fermion coupling corresponds to the contact leading-order (LO) term.

With Eqs. (1) and (4), the radial wave function $\mathcal{R}_k(r)$ in Eq. (3) and the energy E_J for each hypernuclear state are obtained by solving the following coupled-channels equations:

$$\begin{aligned} \left(\frac{d}{dr} - \frac{\kappa - 1}{r} \right) g_k(r) + (E_{nI} - E_J) f_k(r) \\ + \sum_{k'} U_T^{kk'}(r) g_{k'}(r) + \sum_{k'} [U_V^{kk'}(r) + U_S^{kk'}(r)] f_{k'}(r) = 0, \end{aligned} \quad (11a)$$

$$\begin{aligned} \left(\frac{d}{dr} + \frac{\kappa + 1}{r} \right) f_k(r) - (E_{nI} - 2m_\Lambda - E_J) g_k(r) \\ - \sum_{k'} U_T^{kk'}(r) f_{k'}(r) - \sum_{k'} [U_V^{kk'}(r) - U_S^{kk'}(r)] g_{k'}(r) = 0, \end{aligned} \quad (11b)$$

where κ is defined as $\kappa = (-1)^{j+\ell+1/2}(j+1/2)$. The coupling potentials between different channels are given by

$$U_S^{kk'}(r) \equiv \langle \mathcal{F}_{jlnI}^{JM} | \sum_{i=1}^{A_c} \hat{V}_S^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) | \mathcal{F}_{j'l'n'I'}^{JM} \rangle, \quad (12a)$$

$$U_V^{kk'}(r) \equiv \langle \mathcal{F}_{jlnI}^{JM} | \sum_{i=1}^{A_c} \hat{V}_V^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) | \mathcal{F}_{j'l'n'I'}^{JM} \rangle, \quad (12b)$$

$$U_T^{kk'}(r) \equiv \langle \mathcal{F}_{jlnI}^{JM} | \sum_{i=1}^{A_c} \hat{V}_T^{N\Lambda}(\mathbf{r}, \mathbf{r}_i) \cdot \boldsymbol{\sigma} | \mathcal{F}_{j'l'n'I'}^{JM} \rangle, \quad (12c)$$

with

$$\hat{V}_T^{N\Lambda} \equiv \alpha_T^{N\Lambda} [\overleftarrow{\nabla} \delta(\mathbf{r} - \mathbf{r}_i) + \delta(\mathbf{r} - \mathbf{r}_i) \overleftarrow{\nabla}]. \quad (13)$$

By expanding each of the large $f_k(r)$ and small $g_k(r)$ components of the Dirac spinors, Eq. (3), in terms of the radial function $R_{\alpha l}(r)$ of a spherical harmonic oscillator, that is,

$$f_k(r) = \sum_{\alpha=1}^{f_{\text{max}}^{(k)}} F_{k\alpha} R_{\alpha l}^k(r), \quad (14a)$$

$$g_k(r) = \sum_{\alpha=1}^{g_{\text{max}}^{(k)}} G_{k\alpha} R_{\alpha l}^k(r), \quad (14b)$$

with $l = j \pm 1/2$ and $\tilde{l} = j \mp 1/2$, the coupled-channels equations (11a) and (11b) are transformed into a real

symmetric matrix equation,

$$\sum_{\alpha', k'} \begin{pmatrix} A_{\alpha\alpha'}^{kk'} + V_{\alpha\alpha'}^{kk'} + S_{\alpha\alpha'}^{kk'} & B_{\alpha\alpha'}^{kk'} + T_{\alpha\alpha'}^{kk'} \\ B_{\alpha\alpha'}^{kk'} + T_{\alpha\alpha'}^{kk'} & C_{\alpha\alpha'}^{kk'} + V_{\alpha\alpha'}^{kk'} - S_{\alpha\alpha'}^{kk'} \end{pmatrix} \begin{pmatrix} F_{\alpha'}^{k'} \\ G_{\alpha'}^{k'} \end{pmatrix} = E_J \begin{pmatrix} F_{\alpha}^k \\ G_{\alpha}^k \end{pmatrix}. \quad (15)$$

The dimension of the matrix is $\sum_k f_{\max}^{(k)} + g_{\max}^{(k)}$, where k represents different channels. In Eq. (15), the matrix elements are given by

$$A_{\alpha\alpha'}^{kk'} = \langle R_{\alpha l}^k(r) | E_{n l} | R_{\alpha' l'}^{k'}(r) \rangle \delta_{k, k'}, \quad (16a)$$

$$B_{\alpha\alpha'}^{kk'} = \langle R_{\alpha l}^k(r) | \frac{d}{dr} - \frac{\kappa - 1}{r} | R_{\alpha' l'}^{k'}(r) \rangle \delta_{k, k'}, \quad (16b)$$

$$C_{\alpha\alpha'}^{kk'} = \langle R_{\alpha l}^k(r) | (E_{n l} - 2m_{\Lambda}) | R_{\alpha' l'}^{k'}(r) \rangle \delta_{k, k'}, \quad (16c)$$

$$\begin{aligned} V_{\alpha\alpha'}^{kk'} &= \langle R_{\alpha l}^k(r) | U_V^{kk'}(r) | R_{\alpha' l'}^{k'}(r) \rangle \\ &= (-1)^{j'+l'+J} \sum_{\lambda} \begin{Bmatrix} J & I & j \\ \lambda & j' & I' \end{Bmatrix} \langle j \ell || Y_{\lambda} || j' \ell' \rangle \\ &\quad \times \int r^2 dr \rho_{\lambda, V}^{n l n' l'}(r) \left\{ \alpha_V^{N \Lambda} R_{\alpha l}^k(r) R_{\alpha' l'}^{k'}(r) + \delta_V^{N \Lambda} \left[\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \right) - \frac{\lambda(\lambda+1)}{r^2} \right] [R_{\alpha l}^k(r) R_{\alpha' l'}^{k'}(r)] \right\}, \end{aligned} \quad (16d)$$

$$\begin{aligned} S_{\alpha\alpha'}^{kk'} &= \langle R_{\alpha l}^k(r) | U_S^{kk'}(r) | R_{\alpha' l'}^{k'}(r) \rangle \\ &= (-1)^{j'+l'+J} \sum_{\lambda} \begin{Bmatrix} J & I & j \\ \lambda & j' & I' \end{Bmatrix} \langle j \ell || Y_{\lambda} || j' \ell' \rangle \\ &\quad \times \int r^2 dr \rho_{\lambda, S}^{n l n' l'}(r) \left\{ \alpha_S^{N \Lambda} R_{\alpha l}^k(r) R_{\alpha' l'}^{k'}(r) + \delta_S^{N \Lambda} \left[\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \right) - \frac{\lambda(\lambda+1)}{r^2} \right] [R_{\alpha l}^k(r) R_{\alpha' l'}^{k'}(r)] \right\}, \end{aligned} \quad (16e)$$

$$\begin{aligned} T_{\alpha\alpha'}^{kk'} &= \langle R_{\alpha l}^k(r) | U_T^{kk'}(r) | R_{\alpha' l'}^{k'}(r) \rangle \\ &= -\alpha_T^{N \Lambda} (-1)^{j'+l'+J} \sum_{\lambda} \begin{Bmatrix} J & I & j \\ \lambda & j' & I' \end{Bmatrix} \int r^2 dr \rho_{\lambda, V}^{n l n' l'}(r) \\ &\quad \times \left\{ \left[\frac{d R_{\alpha l}^k(r)}{dr} + \frac{\kappa+1}{r} R_{\alpha l}^k(r) \right] R_{\alpha' l'}^{k'}(r) \langle j \tilde{\ell} || Y_{\lambda} || j' \tilde{\ell}' \rangle + \left[\frac{d R_{\alpha' l'}^{k'}(r)}{dr} - \frac{\kappa'-1}{r} R_{\alpha' l'}^{k'}(r) \right] R_{\alpha l}^k(r) \langle j \ell || Y_{\lambda} || j' \ell' \rangle \right\}. \end{aligned} \quad (16f)$$

See Appendices B and C for the derivation of Eqs. (16d) and (16f), respectively. In Eq. (16), $\rho_{\lambda, V}^{n l n' l'}(r)$ and $\rho_{\lambda, S}^{n l n' l'}(r)$ are the reduced vector and scalar transition densities, respectively, between the nuclear initial state $|\Phi_{n l'}\rangle$ and the final state $|\Phi_{n l}\rangle$ defined as

$$\rho_{\lambda, V}^{n l n' l'}(r) = \langle \Phi_{n l} || \sum_{i=1}^{A_c} \frac{\delta(r-r_i)}{r_i r} Y_{\lambda}(\hat{r}_i) || \Phi_{n' l'} \rangle, \quad (17a)$$

$$\rho_{\lambda, S}^{n l n' l'}(r) = \langle \Phi_{n l} || \sum_{i=1}^{A_c} \gamma_i^0 \frac{\delta(r-r_i)}{r_i r} Y_{\lambda}(\hat{r}_i) || \Phi_{n' l'} \rangle. \quad (17b)$$

The detailed expressions for the transition densities can be found in Ref. [37].

III. APPLICATION TO ${}_{\Lambda}^{13}\text{C}$

Let us now apply the MPRM with the higher-order $N\Lambda$ interaction to ${}_{\Lambda}^{13}\text{C}$, for which several low-lying states have been observed experimentally [38,39]. To this end, we first generate several low-lying states of the core nucleus ${}^{12}\text{C}$

with a quantum-number projected GCM calculation, where the mean-field configurations are obtained from a deformation constrained relativistic mean-field plus BCS calculation using the point coupling PC-F1 for the effective nucleon-nucleon interaction [34]. A zero-range pairing force supplemented with a smooth cutoff is adopted to treat the pairing correlation among the nucleons. Axial symmetry and time-reversal invariance are imposed in the mean-field calculations. The Dirac spinor for each nucleon state is expanded on a harmonic oscillator basis with 10 shells. More numerical details have been presented in Refs. [29,30]. The wave functions and the energies of the low-lying states of ${}^{12}\text{C}$ are then used to calculate the scalar and vector transition densities given by Eq. (17) as well as the matrix elements in Eq. (15). The radial wave function for the spherical harmonic oscillator basis with 18 shells is used to expand the radial part of the hypernuclear wave function, $\mathcal{R}_k(r)$. We use the effective $N\Lambda$ interaction with the PCY-S1, PCY-S2, PCY-S3, and PCY-S4 parameter sets, which were determined by fitting to the experimental data of Λ binding energies from light to heavy mass hypernuclei [31]. We list these parameters in Table I. Notice that PCY-S2 and PCY-S4 do not include the tensor and the derivative terms, respectively.

TABLE I. The four parameter sets of the relativistic point-coupling $N\Lambda$ interaction proposed in Ref. [31].

	PCY-S1	PCY-S2	PCY-S3	PCY-S4
$\alpha_S^{N\Lambda}$ (MeV $^{-2}$)	-2.0305×10^{-4}	-4.2377×10^{-5}	-2.0197×10^{-4}	-1.8594×10^{-4}
$\alpha_V^{N\Lambda}$ (MeV $^{-2}$)	1.6548×10^{-4}	1.4268×10^{-5}	1.6449×10^{-4}	1.4981×10^{-4}
$\delta_S^{N\Lambda}$ (MeV $^{-4}$)	2.2929×10^{-9}	1.2986×10^{-9}	2.3514×10^{-9}	-1.9958×10^{-10}
$\delta_V^{N\Lambda}$ (MeV $^{-4}$)	-2.3872×10^{-9}	-1.3850×10^{-9}	-2.4993×10^{-9}	0
$\alpha_T^{N\Lambda}$ (MeV $^{-3}$)	-1.0603×10^{-7}	0	-4.082×10^{-9}	-5.5322×10^{-8}

Notice also that PCY-S3 was obtained by excluding the spin-orbit splitting of the $1p$ state of Λ in ${}^{16}_\Lambda\text{O}$ from the fitting, and the strength of the tensor coupling is considerably smaller than that in PCY-S1.

A. Low-energy spectra

Figures 1(b)–1(e) show the calculated low-energy spectra of ${}^{13}_\Lambda\text{C}$ with the higher order $N\Lambda$ interaction, in comparison with the experimental data as well as with the results of Ref. [30] obtained only with the leading-order $N\Lambda$ interaction. The first and the second states for each spin-parity (J^π) are displayed by the solid and the dashed lines, respectively. One can see that the calculated energy splitting between the $1/2^-$ and $3/2^-$ states, as well as that between the $5/2^+$ and $3/2^+$ states, are clearly different among the four different parameter sets, although the main structures of the low-lying states are the same. That is, the splitting between the $1/2^-$ and $3/2^-$ with PCY-S1 and PCY-S4 forces is smaller than that with PCY-S2 and PCY-S3 forces and much close to the experiment data. The splitting between the $5/2^+$ and $3/2^+$ states with PCY-S1 is much larger than that by the other parameter sets. In other words, the fine structure of the hypernuclear low-lying states reflects the impact of the

$N\Lambda$ interaction beyond the leading order. We have performed similar calculations for ${}^9_\Lambda\text{Be}$, and have found that the effects of the derivative and the tensor terms are similar to those in ${}^{13}_\Lambda\text{C}$. Notice that even though the tensor term is absent in the PCY-S2 force, a good description is still achieved by largely deviating from the expected relations of a naive quark model, that is, $\alpha^{N\Lambda} = \frac{2}{3}\alpha^{NN}$, etc. [40]. We will further discuss the role of the higher order terms in $N\Lambda$ interaction in the next subsections. In particular, we will demonstrate that the tensor term plays an important role if the expected relations of the naive quark model are maintained.

In Fig. 1, the $E2$ transition strengths between the low-lying states of ${}^{13}_\Lambda\text{C}$ are also presented. One can see that the $E2$ transition strengths do not much vary with the four $N\Lambda$ effective interactions and are close to those with the LO interaction.

Given the fact that all the four parameter sets of the effective $N\Lambda$ interaction were adjusted to Λ binding energy of hypernuclei at the mean-field level [31], the use of these forces in the present MPRM calculation overestimates the Λ binding energy of ${}^{13}_\Lambda\text{C}$. That is, the Λ binding energy of ${}^{13}_\Lambda\text{C}$ defined as the energy difference between the 0^+_1 state of ${}^{12}\text{C}$ and the $1/2^+_1$ state of ${}^{13}_\Lambda\text{C}$ is calculated to be 15.72, 13.63, 15.42 and




FIG. 1. The low-energy excitation spectra of ${}^{13}_\Lambda\text{C}$ obtained with the microscopic particle-rotor model with (b) PCY-S1, (c) PCY-S2, (d) PCY-S3, and (e) PCY-S4. Part (f) shows the spectrum taken from Ref. [30], which was obtained by including only the leading-order (LO) $N\Lambda$ interaction. The solid and the dashed lines represent the first and the second states for each spin-parity (J^π). The experimental data shown in (a) are taken from Refs. [1,38,39]. The numbers with the arrows indicate the $B(E2)$ values for the $3/2^+_1 \rightarrow 1/2^+_1$ and the $9/2^+_1 \rightarrow 5/2^+_1$ transitions, given in units of $e^2\text{fm}^4$. The dominant component of several hypernuclear states, together with its weight (in percent), is also given.




FIG. 2. The same as in Fig. 1, but with the scaled $N\Lambda$ interaction, in which the scaling factor is determined for each parameter set to reproduce the empirical Λ binding energy of $^{13}_{\Lambda}\text{C}$.

13.22 MeV using the PCY-S1, PCY-S2, PCY-S3, and PCY-S4 sets of $N\Lambda$ interaction, respectively, while the empirical value is $B_{\Lambda}^{\text{exp.}} = 11.38 \pm 0.05$ MeV [1]. If we want to reproduce the Λ binding energy within this approach, we need to scale all the coupling strengths in the parameters of the $N\Lambda$ interaction by 18%, 9%, 16%, and 8% for PCY-S1, PCY-S2, PCY-S3, and PCY-S4, respectively.

Figure 2 shows the calculated low-lying spectra of $^{13}_{\Lambda}\text{C}$ with those scaled effective $N\Lambda$ interactions. It is shown that the predicted low-lying excitation spectrum of $^{13}_{\Lambda}\text{C}$ is slightly compressed and the $E2$ transition strengths are somewhat increased. On the other hand, the energy splitting between the $1/2_{1}^{-}$ and $3/2_{1}^{-}$ states remains large by the PCY-S2 and PCY-S3 forces, while it is reduced from 303.7 keV (253.7 keV) to 161.5 keV (206.3 keV) after scaling the coupling strengths for the PCY-S1 (PCY-S4) interaction. Due to the slightly weaker $N\Lambda$ interaction, the configuration mixing for the $1/2_{1}^{+}$, $3/2_{1}^{+}$, $5/2_{1}^{+}$, $1/2_{1}^{-}$, and $3/2_{1}^{-}$ states becomes slightly reduced for all four parameter sets.

B. Effects of the derivative coupling terms

We now examine the effect of the derivative coupling terms on the Λ binding energy. To this end, we fix the coupling strengths for the leading order terms ($\alpha_{V}^{N\Lambda}, \alpha_{S}^{N\Lambda}$) to be the same values as those in the PCY-S2 force and study the Λ binding energy as a function of the coupling strengths ($\delta_{V}^{N\Lambda}, \delta_{S}^{N\Lambda}$) of the derivative terms. Notice that the tensor coupling is absent in PCY-S2, so that we can isolate the effect of the derivative terms. The results are shown in Fig. 3(a). A clear linear correlation is observed between $\delta_{V}^{N\Lambda}$ and $\delta_{S}^{N\Lambda}$. By selecting three sets of ($\delta_{V}^{N\Lambda}, \delta_{S}^{N\Lambda}$) along the valley in Fig. 3(a), we calculate the low-lying states of $^{13}_{\Lambda}\text{C}$ and show them in Fig. 3(b). One can see that the low-lying states are similar to each other. This implies that the coupling strengths ($\delta_{V}^{N\Lambda}, \delta_{S}^{N\Lambda}$) may not be uniquely determined by the energies of hypernuclear low-lying states.

Since the vector coupling strengths $\alpha_{V}^{N\Lambda}$ and $\delta_{V}^{N\Lambda}$ are linearly correlated with the corresponding scalar coupling strengths $\alpha_{S}^{N\Lambda}$ and $\delta_{S}^{N\Lambda}$, respectively, we next keep the ratios of $\alpha_{V}^{N\Lambda}/\alpha_{S}^{N\Lambda}$ and $\delta_{V}^{N\Lambda}/\delta_{S}^{N\Lambda}$ the same as those in PCY-S2 force and calculate the Λ binding energy as well as the low-lying spectrum as a function of $\alpha_{S}^{N\Lambda}$ and $\delta_{S}^{N\Lambda}$, as shown in Figs. 3(c) and 3(d), respectively. It is shown that the parameters $\delta_{S}^{N\Lambda}$ and $\alpha_{S}^{N\Lambda}$ are also linearly correlated when these are fitted to the Λ binding energy in $^{13}_{\Lambda}\text{C}$ [see Fig. 3(c)].

Notice that the difference between the vector transition density $\rho_{\lambda,V}^{nIn'I'}(r)$ and the scalar transition density $\rho_{\lambda,S}^{nIn'I'}(r)$ in the low-lying states of ^{12}C is small (see Fig. 4 in Ref. [30]). In the nonrelativistic approximation, with the same scalar and vector densities, the sum of LO coupling strengths $\alpha_{S}^{N\Lambda} + \alpha_{V}^{N\Lambda}$ and the sum of the derivative coupling strengths $\delta_{S}^{N\Lambda} + \delta_{V}^{N\Lambda}$ can be regarded as the depth of the central potential and the surface coupling strength, respectively. Therefore, these are also linearly correlated, as has been found in Ref. [32]. Taking three sets of the parameters along the valley with $B_{\Lambda}^{\text{th}} = B_{\Lambda}^{\text{exp}}$ in Fig. 3(c), we find that those three sets yield almost the same excitation energies (within around 0.13 MeV) for the $3/2_{1}^{+}$, $5/2_{1}^{+}$, $1/2_{2}^{+}$, and $1/2_{1}^{-}$ states, while the difference is much larger (around 0.45 MeV) for the $3/2_{1}^{-}$ and $5/2_{1}^{-}$ states. A comparison between Figs. 3(b) and 3(d) suggests that the excitation energies of the low-lying states are more sensitive to $\alpha_{S}^{N\Lambda}$ and $\alpha_{V}^{N\Lambda}$ as compared to $\delta_{S}^{N\Lambda}$ and $\delta_{V}^{N\Lambda}$.

We next examine the influence of the derivative interaction terms for the other parameter sets as well. To this end, we vary $\delta_{S}^{N\Lambda}$ and $\delta_{V}^{N\Lambda}$ by keeping the values of $\alpha_{S}^{N\Lambda}$, $\alpha_{V}^{N\Lambda}$, $\alpha_{T}^{N\Lambda}$ and the ratio $\delta_{S}^{N\Lambda}/\delta_{V}^{N\Lambda}$ the same as the original values for each parameter set. Fig. 4(a) shows the Λ binding energy so obtained as a function of $|\delta_{S}^{N\Lambda} + \delta_{V}^{N\Lambda}| = -(\delta_{S}^{N\Lambda} + \delta_{V}^{N\Lambda})$. The calculated B_{Λ} with the original value of $\delta_{S}^{N\Lambda}$ and $\delta_{V}^{N\Lambda}$ is denoted by the open circle for each parameter set. B_{Λ} decreases with increasing $|\delta_{S}^{N\Lambda} + \delta_{V}^{N\Lambda}|$ and approaches to the experimental value denoted by the thin solid line. The




FIG. 3. (a) and (c) Contour plots for the absolute value of the difference between the theoretical and the experimental hyperon binding energies of the ^{13}C hypernucleus as a function of the coupling strength parameters $(\delta_S^{N\Lambda}, \delta_V^{N\Lambda})$ and $(\alpha_S^{N\Lambda}, \alpha_V^{N\Lambda})$, respectively. In the former, $\alpha_V^{N\Lambda}$ and $\alpha_S^{N\Lambda}$ are fixed to the same values as in PCY-S2, while, in the latter, the value of $\alpha_V^{N\Lambda}$ and $\delta_V^{N\Lambda}$ is determined for each $(\alpha_S^{N\Lambda}, \delta_S^{N\Lambda})$ so as to keep the ratios $\alpha_V^{N\Lambda}/\alpha_S^{N\Lambda}$ and $\delta_V^{N\Lambda}/\delta_S^{N\Lambda}$ the same as those for PCY-S2. (b) and (d) Low-lying states in ^{13}C calculated with the strength parameters denoted by the dots in panels (a) and (c), respectively.

Λ binding energy decreases from 21.28 to 15.72 MeV by adding the derivative coupling terms to the PCY-S1 interaction (that is, by changing $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$ from 0 to the original value denoted by the open circle). For PCY-S2, PCY-S3, and PCY-S4 interactions, the shift is from 18.01, 23.29, and 21.27 MeV to 13.63, 15.42, and 13.22 MeV, respectively.

The excitation energies of the low-lying states as a function of the derivative coupling strength $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$ are shown in Fig. 4(b), where $\alpha_S^{N\Lambda}$, $\alpha_V^{N\Lambda}$, $\alpha_T^{N\Lambda}$ and $\delta_S^{N\Lambda}/\delta_V^{N\Lambda}$ are kept the same as those for PCY-S1. As one can see, the excitation energies decreases with the increase of $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$. Notice that the changes of the $3/2^+$ and $5/2^+$ states are much smaller compared to the change in the other states. Similar behaviors are found also for the PCY-S2, PCY-S3, and PCY-S4 forces (not shown). The energy splittings of $(3/2_1^+, 5/2_1^+)$ and $(1/2_1^-, 3/2_1^-)$ states as a function of the strength of the




FIG. 4. (a) The Λ binding energy in ^{13}C as a function of $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$, while keeping the same values of $\alpha_S^{N\Lambda}$, $\alpha_V^{N\Lambda}$, $\alpha_T^{N\Lambda}$, and $\delta_S^{N\Lambda}/\delta_V^{N\Lambda}$ as the original values for the PCY-S1, PCY-S2, PCY-S3, and PCY-S4 parameter sets. B_Λ with the original value of $\delta_S^{N\Lambda}$ and $\delta_V^{N\Lambda}$ is denoted by the open circles for each parameter set. The experimental value is denoted by the thin solid line. (b) The energy levels of the low-lying states as a function of $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$ for the PCY-S1 parameter set. (c) and (d) The energy splitting between the $5/2_1^+$ and $3/2_1^+$ states and that between the $1/2_1^-$ and $3/2_1^-$ states, respectively, as a function of $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$.

derivative coupling terms are shown in Figs. 4(c) and 4(d), respectively. It is found that the $5/2_1^+$ state is always slightly higher than the $3/2_1^+$ state—by less than 0.15 MeV except for PCY-S1 in the range of $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}|$ shown in the figure—although this may not be conclusive as the spin-spin $N\Lambda$ interaction [41] is not included in these calculations.

For the doublet $(1/2^-, 3/2^-)$, the $1/2^-$ state is predicted to be higher than the $3/2^-$ state for all the forces except for PCY-S1, with which the $1/2^-$ state is lower than the $3/2^-$ state for $|\delta_S^{N\Lambda} + \delta_V^{N\Lambda}| < 17.56$ MeV. As will be discussed in the next subsection, this splitting, which reflects the spin-orbit splitting of the p_Λ hyperon [30], is mainly governed by the tensor coupling term.

C. Effects of the tensor coupling term

Let us next examine the effects of the tensor coupling term on hypernuclear low-lying states. For this purpose, we adopt the PCY-S1 and PCY-S4 sets of the $N\Lambda$ interaction and vary the strength $\alpha_T^{N\Lambda}$ for the tensor coupling term. Figure 5(a) shows the Λ binding energy of ^{13}C as a function




FIG. 5. Same as Fig. 4, but as a function of the tensor coupling strength $|\alpha_T^{N\Lambda}|$ ($=-\alpha_T^{N\Lambda}$) for the PCY-S1 and PCY-S4 forces.

of $|\alpha_T^{N\Lambda}| = -\alpha_T^{N\Lambda}$. The Λ binding energy gradually decreases from 17.71 MeV (14.12 MeV) for $\alpha_T^{N\Lambda} = 0$ to 15.72 MeV (13.22 MeV) for the original value of $\alpha_T^{N\Lambda}$ for the PCY-S1 (PCY-S4) force, which is indicated by the open circle in Fig. 5(a).

Figure 5(b) shows the excitation energies for the low-lying states of $^{13}_\Lambda\text{C}$ as a function of the tensor coupling strength $|\alpha_T^{N\Lambda}|$ for PCY-S1. As already shown in the previous mean-field studies [24,40,42], the tensor coupling term makes the s_Λ hyperon less bound by increasing the energy of the $s_{1/2}$ level. Moreover, it decreases (increases) the energy of the hyperon $p_{3/2}$ ($p_{1/2}$) state. This is consistent with Fig. 5(a) for the ground state ($1/2^+$) of the $^{13}_\Lambda\text{C}$, the energy of which decreases by the tensor coupling. As a result, the Λ binding energy is reduced by 0.9 MeV for PCY-S1 and 1.99 MeV for PCY-S4 after turning on the tensor coupling term. At the same time, the tensor coupling term decreases (increases) the energy of the $3/2^-$ ($1/2^-$) state, which mainly consists of the $p_{3/2}$ ($p_{1/2}$) hyperon coupled to the ground state (0^+) of ^{12}C . Since the $1/2^-$ changes more significantly than the $3/2^-$ state, the higher lying $1/2^-$ state approaches the $3/2^-$ state and even becomes lower than the $3/2^-$ state for large values of the tensor coupling strength, indicating that the energy splitting of the $1/2^-$ and $3/2^-$ states is sensitive to the tensor coupling strength. For the PCY-S1 and PCY-S4 forces, the energy difference between $1/2_1^-$ and $3/2_1^-$ states decreases from 2.28 to 0.31 MeV, and from 1.25 to 0.25 MeV, respectively, while turning on the tensor coupling term. For the energy gap between the $3/2_1^+$ and $5/2_1^+$ states,

it increases with increasing the $|\alpha_T^{N\Lambda}|$, as shown in Fig. 5(c). Again, the tensor coupling term does not invert the energy ordering of the $3/2_1^+$ and $5/2_1^+$ states.

IV. SUMMARY

We have implemented the higher-order derivative and the tensor terms in the point coupling $N\Lambda$ interaction in the microscopic particle-rotor model for hypernuclear low-lying states. By taking $^{13}_\Lambda\text{C}$ as an example, we have adopted the four sets of effective $N\Lambda$ interaction, which were adjusted at the mean-field level to the Λ binding energy. We have shown that the four parameter sets yield qualitatively similar low-lying spectra, even though these parameter sets were obtained using only the ground state energy.

We have discussed in detail the impact of each $N\Lambda$ interaction term on hypernuclear low-lying states for $^{13}_\Lambda\text{C}$. We have shown that both the second-order derivative and the tensor coupling terms raise the energy of hypernuclear states and thus reduce the Λ binding energy. With the increase of the tensor coupling strength, the excitation energy of the $1/2^-$ state has been found to decrease faster than the $3/2^-$ states. As a result, the energy difference $E(1/2^-) - E(3/2^-)$ decreases to a small value and even changes its sign for large values of the tensor coupling term. We have also found that the $5/2_1^+$ state is higher than the $3/2_1^+$ states with the present effective $N\Lambda$ interaction. We note that the four-fermion coupling terms $(\bar{\psi}_N \Gamma_i \psi_N)(\bar{\psi}_\Lambda \Gamma_i \psi_\Lambda)$ with $\Gamma_i = \sigma_{\mu\nu}$ and $\gamma_\mu \gamma^5$, which provides the spin-spin $N\Lambda$ interaction [36], are not taken into account in the present study. This interaction term may have an important influence on the energy ordering of the $3/2_1^+$ and $5/2_1^+$ states. It will be interesting to study in near future the role of these terms in hypernuclear spectroscopy with the present microscopic particle-rotor model.

Another interesting work is to compare directly between the microscopic particle-rotor model and the generator coordinate method for the whole Λ hypernuclei [28] using the same point-coupling $N\Lambda$ interaction. A work is now in progress, and we will report on it in a separate paper.

ACKNOWLEDGMENTS

This work was supported in part by the Tohoku University Focused Research Project ‘‘Understanding the origins for matters in universe,’’ JSPS KAKENHI Grant No. 2640263, the National Natural Science Foundation of China under Grants No. 11575148, No. 11475140, and No. 11305134, and the Fundamental Research Funds for the Central University (XDJK2013C028).

APPENDIX A: THE $N\Lambda$ EFFECTIVE INTERACTION AND THE CORRESPONDING ENERGY FUNCTIONAL

In this appendix, we show that the $N\Lambda$ interaction given by Eqs. (8), (9), and (10) lead to the energy functional given by Eq. (5). The energy functional for $N\Lambda$ interaction is given by the expectation value of the effective interaction $\hat{V}^{N\Lambda}$ at the

Hartree level,

$$E_{\text{int}}^{N\Lambda} = \sum_{i=1}^{A_c} \int d\mathbf{r} d\mathbf{r}' \psi_{\Lambda}^{\dagger}(\mathbf{r}) \psi_i^{\dagger}(\mathbf{r}') \hat{V}^{N\Lambda}(\mathbf{r}, \mathbf{r}') \psi_{\Lambda}(\mathbf{r}) \psi_i(\mathbf{r}'). \quad (\text{A1})$$

Substituting the LO scalar effective interaction term,

$$\hat{V}_S^{N\Lambda}(\mathbf{r}, \mathbf{r}') = \alpha_S^{N\Lambda} \gamma_{\Lambda}^0 \delta(\mathbf{r} - \mathbf{r}') \gamma_N^0 \quad (\text{A2})$$

to Eq. (A1), one finds

$$\begin{aligned} E_S^{N\Lambda} &= \sum_{i=1}^{A_c} \int d\mathbf{r} d\mathbf{r}' \psi_{\Lambda}^{\dagger}(\mathbf{r}) \psi_i^{\dagger}(\mathbf{r}') \alpha_S^{N\Lambda} \gamma_{\Lambda}^0 \delta(\mathbf{r} - \mathbf{r}') \gamma_N^0 \psi_{\Lambda}(\mathbf{r}) \psi_i(\mathbf{r}') \\ &= \alpha_S^{N\Lambda} \sum_{i=1}^{A_c} \int d\mathbf{r} \psi_{\Lambda}^{\dagger}(\mathbf{r}) \gamma_{\Lambda}^0 \psi_{\Lambda}(\mathbf{r}) \psi_i^{\dagger}(\mathbf{r}) \gamma_N^0 \psi_i(\mathbf{r}) = \int d\mathbf{r} \alpha_S^{N\Lambda} \rho_S^{\Lambda}(\mathbf{r}) \rho_S(\mathbf{r}), \end{aligned} \quad (\text{A3})$$

where ρ_S and ρ_S^{Λ} are the scalar densities defined as

$$\rho_S(\mathbf{r}) = \sum_{i=1}^{A_c} \bar{\psi}_i(\mathbf{r}) \psi_i(\mathbf{r}), \quad \rho_S^{\Lambda}(\mathbf{r}) = \bar{\psi}_{\Lambda}(\mathbf{r}) \psi_{\Lambda}(\mathbf{r}). \quad (\text{A4})$$

The effective interaction with the scalar derivative term,

$$\hat{V}_{\text{Der}}^{N\Lambda}(\mathbf{r}, \mathbf{r}') = \delta_S^{N\Lambda} \gamma_{\Lambda}^0 [\overleftarrow{\nabla}^2 \delta(\mathbf{r} - \mathbf{r}') + \delta(\mathbf{r} - \mathbf{r}') \overrightarrow{\nabla}^2 + 2 \overleftarrow{\nabla} \cdot \delta(\mathbf{r} - \mathbf{r}') \overrightarrow{\nabla}] \gamma_N^0, \quad (\text{A5})$$

leads to

$$\begin{aligned} E_S^{N\Lambda} &= \sum_{i=1}^{A_c} \int d\mathbf{r} d\mathbf{r}' \psi_{\Lambda}^{\dagger}(\mathbf{r}) \psi_i^{\dagger}(\mathbf{r}') \delta_S^{N\Lambda} \gamma_{\Lambda}^0 [\overleftarrow{\nabla}^2 \delta(\mathbf{r} - \mathbf{r}') + \delta(\mathbf{r} - \mathbf{r}') \overrightarrow{\nabla}^2 + 2 \overleftarrow{\nabla} \cdot \delta(\mathbf{r} - \mathbf{r}') \overrightarrow{\nabla}] \gamma_N^0 \psi_{\Lambda}(\mathbf{r}) \psi_i(\mathbf{r}') \\ &= \delta_S^{N\Lambda} \sum_{i=1}^{A_c} \int d\mathbf{r} \{ [\nabla^2 \psi_{\Lambda}^{\dagger}(\mathbf{r}) \gamma_{\Lambda}^0] \psi_{\Lambda}(\mathbf{r}) + [\psi_{\Lambda}^{\dagger}(\mathbf{r}) \gamma_{\Lambda}^0] [\nabla^2 \psi_{\Lambda}(\mathbf{r})] + 2 [\nabla \psi_{\Lambda}^{\dagger}(\mathbf{r}) \gamma_{\Lambda}^0] \cdot [\nabla \psi_{\Lambda}(\mathbf{r})] \} [\psi_i^{\dagger}(\mathbf{r}) \gamma_N^0 \psi_i(\mathbf{r})] \\ &= \int d\mathbf{r} \delta_S^{N\Lambda} \rho_S(\mathbf{r}) \nabla^2 \rho_S^{\Lambda}(\mathbf{r}). \end{aligned} \quad (\text{A6})$$

A similar derivation holds also for the vector part of the $N\Lambda$ interaction.

On the other hand, the tensor effective interaction,

$$\hat{V}_T^{N\Lambda}(\mathbf{r}, \mathbf{r}') = i \alpha_T^{N\Lambda} [\overleftarrow{\nabla} \cdot \boldsymbol{\gamma} \delta(\mathbf{r} - \mathbf{r}') + \delta(\mathbf{r} - \mathbf{r}') \overrightarrow{\nabla} \cdot \boldsymbol{\gamma}] \quad (\text{A7})$$

leads to

$$\begin{aligned} E_T^{N\Lambda} &= \sum_{i=1}^{A_c} \int d\mathbf{r} d\mathbf{r}' \psi_{\Lambda}^{\dagger}(\mathbf{r}) \psi_i^{\dagger}(\mathbf{r}') i \alpha_T^{N\Lambda} [\overleftarrow{\nabla} \cdot \boldsymbol{\gamma} \delta(\mathbf{r} - \mathbf{r}') + \delta(\mathbf{r} - \mathbf{r}') \overrightarrow{\nabla} \cdot \boldsymbol{\gamma}] \psi_{\Lambda}(\mathbf{r}) \psi_i(\mathbf{r}') \\ &= \alpha_T^{N\Lambda} \sum_{i=1}^{A_c} \int d\mathbf{r} \{ [\nabla \psi_{\Lambda}^{\dagger}(\mathbf{r}) \gamma_{\Lambda}^0] \cdot i \boldsymbol{\alpha} \psi_{\Lambda}(\mathbf{r}) + [\psi_{\Lambda}^{\dagger}(\mathbf{r}) \gamma_{\Lambda}^0] [\nabla \cdot i \boldsymbol{\alpha} \psi_{\Lambda}(\mathbf{r})] \} [\psi_i^{\dagger}(\mathbf{r}) \psi_i(\mathbf{r})] \\ &= \int d\mathbf{r} \alpha_T^{N\Lambda} \rho_V(\mathbf{r}) [\nabla \cdot (\bar{\psi}_{\Lambda}(\mathbf{r}) i \boldsymbol{\alpha} \psi_{\Lambda}(\mathbf{r}))] \\ &= \int d\mathbf{r} \alpha_T^{N\Lambda} \rho_V(\mathbf{r}) \rho_T^{\Lambda}(\mathbf{r}), \end{aligned} \quad (\text{A8})$$

where ρ_V and ρ_T^{Λ} are the vector and the tensor densities defined as

$$\rho_V(\mathbf{r}) = \sum_{i=1}^{A_c} \psi_i^{\dagger}(\mathbf{r}) \boldsymbol{\gamma} \psi_i(\mathbf{r}), \quad (\text{A9a})$$

$$\rho_T^{\Lambda}(\mathbf{r}) = \nabla \cdot (\bar{\psi}_{\Lambda}(\mathbf{r}) i \boldsymbol{\alpha} \psi_{\Lambda}(\mathbf{r})). \quad (\text{A9b})$$

Putting all these together, we finally obtain Eq.(5).

**APPENDIX B: A DERIVATION OF Eq. (16d) FOR THE MATRIX ELEMENTS OF THE VECTOR
DERIVATIVE COUPLING TERM**

With the $N\Lambda$ vector derivative effective interaction $\hat{V}_D = \delta_V^{N\Lambda} [\overleftarrow{\nabla}^2 \delta(\mathbf{r} - \mathbf{r}_i) + \delta(\mathbf{r} - \mathbf{r}_i) \overrightarrow{\nabla}^2 + 2\overleftarrow{\nabla} \cdot \delta(\mathbf{r} - \mathbf{r}_i) \overrightarrow{\nabla}]$, and the definition of

$$\mathcal{F}_{jIl}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) = \sum_{m_l m} C_{Im_l j m}^{JM} \mathcal{Y}_{jlm}(\hat{\mathbf{r}}) \Phi_{Im_l}(\{\mathbf{r}_i\}), \quad (\text{B1})$$

where $\mathcal{Y}_{jlm}(\hat{\mathbf{r}})$ is the spinor spherical harmonics,

$$\mathcal{Y}_{j\ell m}(\hat{\mathbf{r}}) = \sum_{m_l m_s} C_{lm_l \frac{1}{2} m_s}^{jm} Y_{lm_l}(\vartheta, \varphi) \chi_{m_s}, \quad (\text{B2})$$

and $C_{l_1 m_1 l_2 m_2}^{jm}$ is the Clebsch-Gordan coefficient, the coupling matrix element of the vector derivative term reads

$$\begin{aligned} & \langle R_{\alpha l}^k(r) \mathcal{F}_{jIl}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) | \hat{V}_D | \mathcal{F}_{j'I'l'}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) R_{\alpha'l'}^k(r) \rangle \\ &= \delta_V^{N\Lambda} \sum_{m_l m_l'} \sum_{m_l' m_l''} C_{Im_l j m}^{JM} C_{I'm_l' j' m'}^{JM} \sum_{\lambda \mu} \int r^2 dr \int d\hat{\mathbf{r}} \langle \Phi_{Im_l} | \sum_{i=1}^{A_c} \frac{\delta(\mathbf{r} - \mathbf{r}_i)}{r r_i} Y_{\lambda \mu}(\hat{\mathbf{r}}_i) | \Phi_{I'm_l'} \rangle Y_{\lambda \mu}^*(\hat{\mathbf{r}}) \Delta[\mathcal{Y}_{j\ell m}^*(\hat{\mathbf{r}}) \mathcal{Y}_{j'\ell' m'}(\hat{\mathbf{r}}) R_{\alpha l}^k(r) R_{\alpha'l'}^k(r)]. \end{aligned} \quad (\text{B3})$$

Here, we notice

$$\begin{aligned} \langle \Phi_{Im_l} | \sum_{i=1}^{A_c} \frac{\delta(\mathbf{r} - \mathbf{r}_i)}{r r_i} Y_{\lambda \mu}(\hat{\mathbf{r}}_i) | \Phi_{I'm_l'} \rangle &= (-1)^{I-m_l} \begin{pmatrix} I & \lambda & I' \\ -m_l & \mu & m_l' \end{pmatrix} \left\langle \Phi_I \left\| \sum_{i=1}^{A_c} \frac{\delta(\mathbf{r} - \mathbf{r}_i)}{r r_i} Y_{\lambda}(\hat{\mathbf{r}}_i) \right\| \Phi_{I'} \right\rangle \\ &= (-1)^{I-m_l} \begin{pmatrix} I & \lambda & I' \\ -m_l & \mu & m_l' \end{pmatrix} \rho_{\lambda, V}^{II'}(r). \end{aligned} \quad (\text{B4})$$

With the relation of

$$\mathcal{Y}_{j\ell m}^*(\hat{\mathbf{r}}) \mathcal{Y}_{j'\ell' m'}(\hat{\mathbf{r}}) = \sum_{m_l m_s} \sum_{m_l' m_s'} C_{lm_l \frac{1}{2} m_s}^{jm} C_{l'm_l' \frac{1}{2} m_s'}^{j'm'} \delta_{m_s m_s'} (-1)^{m_l} \sum_{LM} \frac{\hat{l}'}{\sqrt{4\pi \hat{L}}} C_{l0 l'0}^{L0} C_{l-m_l l'-m_l'}^{LM} Y_{LM}(\hat{\mathbf{r}}), \quad (\text{B5})$$

we have

$$\Delta[\mathcal{Y}_{j\ell m}^*(\hat{\mathbf{r}}) \mathcal{Y}_{j'\ell' m'}(\hat{\mathbf{r}})] = \sum_{m_l m_s} \sum_{m_l' m_s'} \sum_{LM} \frac{\hat{l}'}{\sqrt{4\pi \hat{L}}} C_{lm_l \frac{1}{2} m_s}^{jm} C_{l'm_l' \frac{1}{2} m_s'}^{j'm'} \delta_{m_s m_s'} (-1)^{m_l} C_{l0 l'0}^{L0} C_{l-m_l l'-m_l'}^{LM} \left[\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \right) - \frac{L(L+1)}{r^2} \right] Y_{LM}(\hat{\mathbf{r}}). \quad (\text{B6})$$

According to the orthogonalization of the spherical harmonics,

$$\int Y_{\lambda \mu}^*(\hat{\mathbf{r}}) Y_{L m_L}(\hat{\mathbf{r}}) d\hat{\mathbf{r}} = \delta_{\lambda, L} \delta_{\mu, m_L}, \quad (\text{B7})$$

the matrix element is then given by

$$\begin{aligned} & \langle R_{\alpha l}^k(r) \mathcal{F}_{jIl}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) | \hat{V}_D | \mathcal{F}_{j'I'l'}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) R_{\alpha'l'}^k(r) \rangle = \delta_V^{N\Lambda} (-1)^{j'+I+J} \sum_{\lambda} \begin{Bmatrix} J & I & l \\ \lambda & j' & I' \end{Bmatrix} \langle j\ell || Y_{\lambda} || j'\ell' \rangle \\ & \quad \times \int r^2 dr \rho_{\lambda, V}^{II'}(r) \left[\frac{1}{r^2} \frac{d}{dr} \left(r^2 \frac{d}{dr} \right) - \frac{\lambda(\lambda+1)}{r^2} \right] [R_{\alpha l}^k(r) R_{\alpha'l'}^k(r)]. \end{aligned} \quad (\text{B8})$$

APPENDIX C: A DERIVATION OF Eq. (16f) FOR THE MATRIX ELEMENTS OF THE TENSOR COUPLING TERM

The matrix elements of the tensor coupling term are given by

$$\begin{aligned} T_{\alpha\alpha'}^{kk'} &\equiv \langle R_{\alpha l}^k(r) \mathcal{F}_{jIl}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) | \alpha_T^{N\Lambda} \sum_{i=1}^{A_c} [\overleftarrow{\nabla} \delta(\mathbf{r} - \mathbf{r}_i) + \delta(\mathbf{r} - \mathbf{r}_i) \overrightarrow{\nabla}] \cdot \boldsymbol{\sigma} | \mathcal{F}_{j'I'l'}^{JM}(\hat{\mathbf{r}}, \{\mathbf{r}_i\}) R_{\alpha'l'}^k(r) \rangle \\ &= \alpha_T^{N\Lambda} \sum_{m_l' m_l m} \sum_{m_l' m_l} C_{I'm_l' j' m'}^{JM} C_{Im_l j m}^{JM} \sum_{\lambda \mu} \int r^2 dr d\hat{\mathbf{r}} \langle \Phi_{Im_l} | \sum_{i=1}^{A_c} \frac{\delta(\mathbf{r} - \mathbf{r}_i)}{r r_i} Y_{\lambda \mu}(\hat{\mathbf{r}}_i) | \Phi_{I'm_l'} \rangle \\ & \quad \times Y_{\lambda \mu}^*(\hat{\mathbf{r}}) \nabla \cdot [R_{\alpha l}^{k*}(r) \mathcal{Y}_{j\ell m}^*(\hat{\mathbf{r}}) \boldsymbol{\sigma} R_{\alpha'l'}^k(r) \mathcal{Y}_{j'\ell' m'}(\hat{\mathbf{r}})]. \end{aligned} \quad (\text{C1})$$

Notice

$$\begin{aligned} \nabla \cdot [R_{\alpha l}^{k*}(r) \mathcal{Y}_{jlm}^*(\hat{\mathbf{r}}) \sigma R_{\alpha' \tilde{l}}^{k'}(r) \mathcal{Y}_{j' \tilde{l} m'}(\hat{\mathbf{r}})] = & \left[-\frac{dR_{\alpha l}^k(r)}{dr} - \frac{\kappa+1}{r} R_{\alpha l}^k(r) \right] [R_{\alpha' \tilde{l}}^{k'}(r) \mathcal{Y}_{jlm}^*(\hat{\mathbf{r}}) \mathcal{Y}_{j' \tilde{l} m'}(\hat{\mathbf{r}})] \\ & - \left[\frac{dR_{\alpha' \tilde{l}}^{k'}(r)}{dr} - \frac{\kappa'-1}{r} R_{\alpha' \tilde{l}}^{k'}(r) \right] [R_{\alpha l}^{k*}(r) \mathcal{Y}_{jlm}^*(\hat{\mathbf{r}}) \mathcal{Y}_{j' \tilde{l} m'}(\hat{\mathbf{r}})]. \end{aligned} \quad (\text{C2})$$

With the Wigner-Eckart theorem, one obtains

$$\int d\hat{\mathbf{r}} \mathcal{Y}_{jlm}^*(\hat{\mathbf{r}}) Y_{\lambda\mu}^*(\hat{\mathbf{r}}) \mathcal{Y}_{j'l'm'}(\hat{\mathbf{r}}) = (-1)^{\mu+j-m} \begin{pmatrix} j & \lambda & j' \\ -m & -\mu & m' \end{pmatrix} \langle j l || Y_{\lambda} || j' l' \rangle. \quad (\text{C3})$$

From the relation

$$\sum_{m'_l m_l} \sum_{m'_l m_l} \sum_{\mu} C_{l'm'_l j'm'}^{JM} C_{l m_l j m}^{JM} (-1)^{l-m_l} \begin{pmatrix} l & \lambda & l' \\ -m_l & \mu & m'_l \end{pmatrix} (-1)^{\mu+j-m} \begin{pmatrix} j & \lambda & j' \\ -m & -\mu & m' \end{pmatrix} = (-1)^{l'+j+j} \begin{Bmatrix} J & l & j \\ \lambda & j' & l' \end{Bmatrix}, \quad (\text{C4})$$

one finally obtains

$$\begin{aligned} T_{\alpha\alpha'}^{kk'} = & -\alpha_T^{N\Lambda} (-1)^{j+l'+j} \sum_{\lambda} \begin{Bmatrix} J & l & j \\ \lambda & j' & l' \end{Bmatrix} \int r^2 dr \rho_{\lambda, V}^{l'l}(r) \left\{ \left[\frac{dR_{\alpha l}^k(r)}{dr} + \frac{\kappa+1}{r} R_{\alpha l}^k(r) \right] R_{\alpha' \tilde{l}}^{k'}(r) \langle j \tilde{l} || Y_{\lambda} || j' \tilde{l}' \rangle \right. \\ & \left. + \left[\frac{dR_{\alpha' \tilde{l}}^{k'}(r)}{dr} - \frac{\kappa'-1}{r} R_{\alpha' \tilde{l}}^{k'}(r) \right] R_{\alpha l}^k(r) \langle j l || Y_{\lambda} || j' l' \rangle \right\}. \end{aligned} \quad (\text{C5})$$

-
- [1] O. Hashimoto and H. Tamura, *Prog. Part. Nucl. Phys.* **57**, 564 (2006).
- [2] H. Tamura, *Int. J. Mod. Phys. A* **24**, 2101 (2009).
- [3] N. Glendenning, *Compact Stars* (Springer-Verlag, New York, 2000).
- [4] R. Wirth, D. Gazda, P. Navratil, A. Calci, J. Langhammer, and R. Roth, *Phys. Rev. Lett.* **113**, 192502 (2014).
- [5] T. Motoba, H. Bandō, and K. Ikeda, *Prog. Theor. Phys.* **70**, 189 (1983).
- [6] E. Hiyama, M. Kamimura, K. Miyazaki, and T. Motoba, *Phys. Rev. C* **59**, 2351 (1999).
- [7] H. Bando, T. Motoba, and J. Žofka, *Int. J. Mod. Phys. A* **5**, 4021 (1990).
- [8] E. Hiyama, Y. Kino, and M. Kamimura, *Prog. Part. Nucl. Phys.* **51**, 223 (2003).
- [9] E. Cravo, A. C. Fonseca, and Y. Koike, *Phys. Rev. C* **66**, 014001 (2002).
- [10] V. M. Suslov, I. Filikhin, and B. Vlahovic, *J. Phys. G: Nucl. Part. Phys.* **30**, 513 (2004).
- [11] M. Shoeb and Sonika, *Phys. Rev. C* **79**, 054321 (2009).
- [12] R. H. Dalitz and A. Gal, *Ann. Phys. (NY)* **116**, 167 (1978).
- [13] A. Gal, J. M. Soper, and R. H. Dalitz, *Ann. Phys. (NY)* **63**, 53 (1971).
- [14] D. J. Millener, *Nucl. Phys. A* **804**, 84 (2008); **914**, 109 (2013).
- [15] M. Isaka, M. Kimura, A. Doté, and A. Ohnishi, *Phys. Rev. C* **83**, 044323 (2011).
- [16] M. Isaka, M. Kimura, A. Doté, and A. Ohnishi, *Phys. Rev. C* **83**, 054304 (2011).
- [17] M. Isaka, H. Homma, M. Kimura, A. Doté, and A. Ohnishi, *Phys. Rev. C* **85**, 034303 (2012).
- [18] M. Isaka, M. Kimura, A. Doté, and A. Ohnishi, *Phys. Rev. C* **87**, 021304(R) (2013).
- [19] X. R. Zhou, H.-J. Schulze, H. Sagawa, C. X. Wu, and E.-G. Zhao, *Phys. Rev. C* **76**, 034312 (2007).
- [20] M. T. Win and K. Hagino, *Phys. Rev. C* **78**, 054311 (2008).
- [21] H.-J. Schulze, M. T. Win, K. Hagino, and H. S. Sagawa, *Prog. Theo. Phys.* **123**, 569 (2010).
- [22] M. T. Win, K. Hagino, and T. Koike, *Phys. Rev. C* **83**, 014301 (2011).
- [23] B.-N. Lu, E.-G. Zhao, and S.-G. Zhou, *Phys. Rev. C* **84**, 014328 (2011).
- [24] W. X. Xue, J. M. Yao, K. Hagino, Z. P. Li, H. Mei, and Y. Tanimura, *Phys. Rev. C* **91**, 024327 (2015).
- [25] A. Li, E. Hiyama, X.-R. Zhou, and H. Sagawa, *Phys. Rev. C* **87**, 014333 (2013).
- [26] B.-N. Lu, E. Hiyama, H. Sagawa, and S.-G. Zhou, *Phys. Rev. C* **89**, 044307 (2014).
- [27] K. Hagino and J. M. Yao, in *Relativistic Density Functional for Nuclear Structure*, edited by J. Meng, International Review of Nuclear Physics Vol. 10 (World Scientific, Singapore, 2016), p. 263.
- [28] H. Mei, K. Hagino, and J. M. Yao, *Phys. Rev. C* **93**, 011301(R) (2016).
- [29] H. Mei, K. Hagino, J. M. Yao, and T. Motoba, *Phys. Rev. C* **90**, 064302 (2014).
- [30] H. Mei, K. Hagino, J. M. Yao, and T. Motoba, *Phys. Rev. C* **91**, 064305 (2015).
- [31] Y. Tanimura and K. Hagino, *Phys. Rev. C* **85**, 014306 (2012).
- [32] E. Hiyama, Y. Funaki, N. Kaiser, and W. Weise, *Prog. Theor. Exp. Phys.* (2014) 013D01.
- [33] J. V. Noble, *Phys. Lett. B* **89**, 325 (1980).
- [34] T. Burvenich, D. G. Madland, J. A. Maruhn, and P.-G. Reinhard, *Phys. Rev. C* **65**, 044308 (2002).
- [35] P. Ring and P. Schuck, *The Nuclear Many-Body Problem* (Springer-Verlag, Berlin, 1980).

- [36] H. Polinder, J. Haidenbauer, and Ulf-G. Meissner, *Nucl. Phys. A* **779**, 244 (2006).
- [37] J. M. Yao, M. Bender, and P.-H. Heenen, *Phys. Rev. C* **91**, 024301 (2015).
- [38] S. Ajimura, H. Hayakawa, T. Kishimoto, H. Kohri, K. Matsuoka, S. Minami, T. Mori, K. Morikubo, E. Saji, A. Sakaguchi, Y. Shimizu, M. Sumihama, R. E. Chrien, M. May, P. Pile, A. Rusek, R. Sutter, P. Eugenio, G. Franklin, P. Khaustov, K. Paschke, B. P. Quinn, R. A. Schumacher, J. Franz, T. Fukuda, H. Noumi, H. Outa, L. Gan, L. Tang, L. Yuan, H. Tamura, J. Nakano, T. Tamagawa, K. Tanida, and R. Sawafta, *Phys. Rev. Lett.* **86**, 4255 (2001).
- [39] H. Kohri, S. Ajimura, H. Hayakawa, T. Kishimoto, K. Matsuoka, S. Minami, Y. S. Miyake, T. Mori, K. Morikubo, E. Saji, A. Sakaguchi, Y. Shimizu, M. Sumihama, R. E. Chrien, M. May, P. Pile, A. Rusek, R. Sutter, P. M. Eugenio, G. Franklin, P. Khaustov, K. Paschke, B. P. Quinn, R. A. Schumacher, J. Franz, T. Fukuda, H. Noumi, H. Outa, L. Gan, L. Tang, L. Yuan, J. Nakano, T. Tamagawa, K. Tanida, R. Sawafta, H. Tamura, and H. Akikawa, *Phys. Rev. C* **65**, 034607 (2002).
- [40] Y. Sugahara and H. Toki, *Prog. Theor. Phys.* **92**, 803 (1994).
- [41] L. Canton, K. Amos, S. Karataglidis, and J. P. Svenne, *Int. J. Mod. Phys. E* **19**, 1435 (2010).
- [42] J. Mareš and B. K. Jennings, *Phys. Rev. C* **49**, 2472 (1994).