

Centrality and transverse momentum dependence of D^0 -meson production at mid-rapidity in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV

J. Adam,¹² L. Adamczyk,² J. R. Adams,³⁴ J. K. Adkins,²⁵ G. Agakishiev,²³ M. M. Aggarwal,³⁶ Z. Ahammed,⁵⁶ I. Alekseev,^{3,30} D. M. Anderson,⁵⁰ R. Aoyama,⁵³ A. Aparin,²³ D. Arkhipkin,⁵ E. C. Aschenauer,⁵ M. U. Ashraf,⁵² F. Atetalla,²⁴ A. Attri,³⁶ G. S. Averichev,²³ V. Bairathi,³¹ K. Barish,⁹ A. J. Bassill,⁹ A. Behera,⁴⁸ R. Bellwied,¹⁹ A. Bhasin,²² A. K. Bhati,³⁶ J. Bielcik,¹³ J. Bielcikova,³³ L. C. Bland,⁵ I. G. Bordyuzhin,³ J. D. Brandenburg,⁵ A. V. Brandin,³⁰ D. Brown,²⁷ J. Bryslawskyj,⁹ I. Bunzarov,²³ J. Butterworth,⁴¹ H. Caines,⁵⁹ M. Calderón de la Barca Sánchez,⁷ D. Cebra,⁷ I. Chakaberia,^{24,45} P. Chaloupka,¹³ B. K. Chan,⁸ F.-H. Chang,³² Z. Chang,⁵ N. Chankova-Bunzarova,²³ A. Chatterjee,⁵⁶ S. Chattopadhyay,⁵⁶ J. H. Chen,⁴⁶ X. Chen,⁴⁴ J. Cheng,⁵² M. Cherney,¹² W. Christie,⁵ G. Contin,²⁶ H. J. Crawford,⁶ M. Csanad,¹⁵ S. Das,¹⁰ T. G. Dedovich,²³ I. M. Deppner,¹⁸ A. A. Derevschikov,³⁸ L. Didenko,⁵ C. Dilks,³⁷ X. Dong,²⁶ J. L. Drachenberg,¹ J. C. Dunlop,⁵ T. Edmonds,³⁹ L. G. Efimov,²³ N. Elsev,⁵⁸ J. Engelage,⁶ G. Eppley,⁴¹ R. Esha,⁸ S. Esumi,⁵³ O. Evdokimov,¹¹ J. Ewigleben,²⁷ O. Eyser,⁵ R. Fatemi,²⁵ S. Fazio,⁵ P. Federic,³³ J. Fedorisin,²³ P. Filip,²³ E. Finch,⁴⁷ Y. Fisyak,⁵ C. E. Flores,⁷ L. Fulek,² C. A. Gagliardi,⁵ T. Galatyuk,¹⁴ F. Geurts,⁴¹ A. Gibson,⁵⁵ L. Greiner,²⁶ D. Grosnick,⁵⁵ D. S. Gunarathne,⁴⁹ Y. Guo,²⁴ A. Gupta,²² W. Guryn,⁵ A. I. Hamad,²⁴ A. Hamed,⁵⁰ A. Harlanderova,¹³ J. W. Harris,⁵⁹ L. He,³⁹ S. Heppelmann,⁷ S. Heppelmann,⁷ N. Herrmann,¹⁸ A. Hirsch,³⁹ L. Holub,¹³ Y. Hong,²⁶ S. Horvat,⁵⁹ B. Huang,¹¹ H. Z. Huang,⁸ S. L. Huang,⁴⁸ T. Huang,³² X. Huang,⁵² T. J. Humanic,³⁴ P. Huo,⁴⁸ G. Igo,⁸ W. W. Jacobs,²⁰ A. Jentsch,⁵¹ J. Jia,^{5,48} K. Jiang,⁴⁴ S. Jowzaee,⁵⁸ X. Ju,⁴⁴ E. G. Judd,⁶ S. Kabana,²⁴ S. Kagamaster,²⁷ D. Kalinkin,²⁰ K. Kang,⁵² D. Kapukchyan,⁹ K. Kauder,⁵ H. W. Ke,⁵ D. Keane,²⁴ A. Kechechyan,²³ M. Kelsey,²⁶ D. P. Kikola,⁵⁷ C. Kim,⁹ T. A. Kinghorn,⁷ I. Kisel,¹⁶ A. Kisiel,⁵⁷ M. Kocan,¹³ L. Kochenda,³⁰ L. K. Kosarzewski,¹³ A. F. Kraishan,⁴⁹ L. Kramarik,¹³ L. Krauth,⁹ P. Kravtsov,³⁰ K. Krueger,⁴ N. Kulathunga Mudiyanseelage,¹⁹ L. Kumar,³⁶ R. Kunnawalkam Elayavalli,⁵⁸ J. Kvapil,¹³ J. H. Kwasizur,²⁰ R. Lacey,⁴⁸ J. M. Landgraf,⁵ J. Lauret,⁵ A. Lebedev,⁵ R. Lednicky,²³ J. H. Lee,⁵ C. Li,⁴⁴ W. Li,⁴⁶ W. Li,⁴¹ X. Li,⁴⁴ Y. Li,⁵² Y. Liang,²⁴ R. Licenik,¹³ J. Lidrych,¹³ T. Lin,⁵⁰ A. Lipiec,⁵⁷ M. A. Lisa,³⁴ F. Liu,¹⁰ H. Liu,²⁰ P. Liu,⁴⁸ P. Liu,⁴⁶ X. Liu,³⁴ Y. Liu,⁵⁰ Z. Liu,⁴⁴ T. Ljubicic,⁵ W. J. Llope,⁵⁸ M. Lomnitz,²⁶ R. S. Longacre,⁵ S. Luo,¹¹ X. Luo,¹⁰ G. L. Ma,⁴⁶ L. Ma,¹⁷ R. Ma,⁵ Y. G. Ma,⁴⁶ N. Magdy,¹¹ R. Majka,⁵⁹ D. Mallick,³¹ S. Margetis,²⁴ C. Markert,⁵¹ H. S. Matis,²⁶ O. Matonoha,¹³ J. A. Mazer,⁴² K. Meehan,⁷ J. C. Mei,⁴⁵ N. G. Minaev,³⁸ S. Mioduszewski,⁵⁰ D. Mishra,³¹ B. Mohanty,³¹ M. M. Mondal,²¹ I. Mooney,⁵⁸ Z. Moravcova,¹³ D. A. Morozov,³⁸ M. K. Mustafa,²⁶ Md. Nasim,⁸ K. Nayak,¹⁰ J. M. Nelson,⁶ D. B. Nemes,⁵⁹ M. Nie,⁴⁶ G. Nigmatkulov,³⁰ T. Niida,⁵⁸ L. V. Nogach,³⁸ T. Nonaka,¹⁰ G. Odyniec,²⁶ A. Ogawa,⁵ K. Oh,⁴⁰ S. Oh,⁵⁹ V. A. Okorokov,³⁰ D. Olvitt Jr.,⁴⁹ B. S. Page,⁵ R. Pak,⁵ Y. Panebratsev,²³ B. Pawlik,³⁵ H. Pei,¹⁰ C. Perkins,⁶ R. L. Pinter,¹⁵ J. Pluta,⁵⁷ J. Porter,²⁶ M. Posik,⁴⁹ N. K. Pruthi,³⁶ M. Przybycien,² J. Putschke,⁵⁸ A. Quintero,⁴⁹ H. Qiu,²⁶ S. K. Radhakrishnan,²⁶ S. Ramachandran,²⁵ R. L. Ray,⁵¹ R. Reed,²⁷ H. G. Ritter,²⁶ J. B. Roberts,⁴¹ O. V. Rogachevskiy,²³ J. L. Romero,⁷ L. Ruan,⁵ J. Rusnak,³³ O. Rusnakova,¹³ N. R. Sahoo,⁵⁰ P. K. Sahu,²¹ S. Salur,⁴² J. Sandweiss,⁵⁹ J. Schambach,⁵¹ A. M. Schmah,²⁶ W. B. Schmidke,⁵ N. Schmitz,²⁸ B. R. Schweid,⁴⁸ F. Seck,¹⁴ J. Seger,¹² M. Sergeeva,⁸ R. Seto,⁹ P. Seyboth,²⁸ N. Shah,⁴⁶ E. Shahaiev,²³ P. V. Shanmuganathan,²⁷ M. Shao,⁴⁴ F. Shen,⁴⁵ W. Q. Shen,⁴⁶ S. S. Shi,¹⁰ Q. Y. Shou,⁴⁶ E. P. Sichtermann,²⁶ S. Siejka,⁵⁷ R. Sikora,² M. Simko,³³ J. Singh,³⁶ S. Singha,²⁴ D. Smirnov,⁵ N. Smirnov,⁵⁹ W. Solyst,²⁰ P. Sorensen,⁵ H. M. Spinka,⁴ B. Srivastava,³⁹ T. D. S. Stanislaus,⁵⁵ D. J. Stewart,⁵⁹ M. Strikhanov,³⁰ B. Stringfellow,³⁹ A. A. P. Suaide,⁴³ T. Sugiura,⁵³ M. Sumera,³³ B. Summa,³⁷ X. M. Sun,¹⁰ Y. Sun,⁴⁴ Y. Sun,⁶⁰ B. Surrow,⁴⁹ D. N. Svirida,³ M. Szelezniak,²⁶ P. Szymanski,⁵⁷ A. H. Tang,⁵ Z. Tang,⁴⁴ A. Taranenko,³⁰ T. Tarnowsky,²⁹ J. H. Thomas,²⁶ A. R. Timmins,¹⁹ T. Todoroki,⁵ M. Tokarev,²³ C. A. Tomkiel,²⁷ S. Trentalange,⁸ R. E. Tribble,⁵⁰ P. Tribedy,⁵ S. K. Tripathy,²¹ O. D. Tsai,⁸ B. Tu,¹⁰ T. Ullrich,⁵ D. G. Underwood,⁴ I. Upsal,^{5,45} G. Van Buren,⁵ J. Vanek,³³ A. N. Vasiliev,³⁸ I. Vassiliev,¹⁶ F. Videbæk,⁵ S. Vokal,²³ S. A. Voloshin,⁵⁸ A. Vossen,²⁰ F. Wang,³⁹ G. Wang,⁸ P. Wang,⁴⁴ Y. Wang,¹⁰ Y. Wang,⁵² J. C. Webb,⁵ L. Wen,⁸ G. D. Westfall,²⁹ H. Wieman,²⁶ S. W. Wissink,²⁰ R. Witt,⁵⁴ Y. Wu,²⁴ Z. G. Xiao,⁵² G. Xie,¹¹ W. Xie,³⁹ H. Xu,⁶⁰ N. Xu,²⁶ Q. H. Xu,⁴⁵ Y. F. Xu,⁴⁶ Z. Xu,⁵ C. Yang,⁴⁵ Q. Yang,⁴⁵ S. Yang,⁵ Y. Yang,³² Z. Ye,⁴¹ Z. Ye,¹¹ L. Yi,⁴⁵ K. Yip,⁵ I.-K. Yoo,⁴⁰ H. Zbroszczyk,⁵⁷ W. Zha,⁴⁴ D. Zhang,¹⁰ J. Zhang,⁴⁸ L. Zhang,¹⁰ S. Zhang,⁴⁴ S. Zhang,⁴⁶ X. P. Zhang,⁵² Y. Zhang,⁴⁴ Z. Zhang,⁴⁶ J. Zhao,³⁹ C. Zhong,⁴⁶ C. Zhou,⁴⁶ X. Zhu,⁵² Z. Zhu,⁴⁵ M. K. Zurek,²⁶ and M. Zyzak¹⁶

(STAR Collaboration)

¹Abilene Christian University, Abilene, Texas 79699²AGH University of Science and Technology, FPACS, Cracow 30-059, Poland³Alikhanov Institute for Theoretical and Experimental Physics, Moscow 117218, Russia⁴Argonne National Laboratory, Argonne, Illinois 60439⁵Brookhaven National Laboratory, Upton, New York 11973⁶University of California, Berkeley, California 94720⁷University of California, Davis, California 95616⁸University of California, Los Angeles, California 90095⁹University of California, Riverside, California 92521¹⁰Central China Normal University, Wuhan, Hubei 430079

- ¹¹University of Illinois at Chicago, Chicago, Illinois 60607
¹²Creighton University, Omaha, Nebraska 68178
¹³Czech Technical University in Prague, FNSPE, Prague 115 19, Czech Republic
¹⁴Technische Universität Darmstadt, Darmstadt 64289, Germany
¹⁵Eötvös Loránd University, Budapest, Hungary H-1117
¹⁶Frankfurt Institute for Advanced Studies FIAS, Frankfurt 60438, Germany
¹⁷Fudan University, Shanghai, 200433
¹⁸University of Heidelberg, Heidelberg 69120, Germany
¹⁹University of Houston, Houston, Texas 77204
²⁰Indiana University, Bloomington, Indiana 47408
²¹Institute of Physics, Bhubaneswar 751005, India
²²University of Jammu, Jammu 180001, India
²³Joint Institute for Nuclear Research, Dubna 141 980, Russia
²⁴Kent State University, Kent, Ohio 44242
²⁵University of Kentucky, Lexington, Kentucky 40506-0055
²⁶Lawrence Berkeley National Laboratory, Berkeley, California 94720
²⁷Lehigh University, Bethlehem, Pennsylvania 18015
²⁸Max-Planck-Institut für Physik, Munich 80805, Germany
²⁹Michigan State University, East Lansing, Michigan 48824
³⁰National Research Nuclear University MEPhI, Moscow 115409, Russia
³¹National Institute of Science Education and Research, HBNI, Jatni 752050, India
³²National Cheng Kung University, Tainan 70101
³³Nuclear Physics Institute of the CAS, Rez 250 68, Czech Republic
³⁴Ohio State University, Columbus, Ohio 43210
³⁵Institute of Nuclear Physics PAN, Cracow 31-342, Poland
³⁶Panjab University, Chandigarh 160014, India
³⁷Pennsylvania State University, University Park, Pennsylvania 16802
³⁸Institute of High Energy Physics, Protvino 142281, Russia
³⁹Purdue University, West Lafayette, Indiana 47907
⁴⁰Pusan National University, Pusan 46241, Korea
⁴¹Rice University, Houston, Texas 77251
⁴²Rutgers University, Piscataway, New Jersey 08854
⁴³Universidade de São Paulo, São Paulo, Brazil 05314-970
⁴⁴University of Science and Technology of China, Hefei, Anhui 230026
⁴⁵Shandong University, Qingdao, Shandong 266237
⁴⁶Shanghai Institute of Applied Physics, Chinese Academy of Sciences, Shanghai 201800
⁴⁷Southern Connecticut State University, New Haven, Connecticut 06515
⁴⁸State University of New York, Stony Brook, New York 11794
⁴⁹Temple University, Philadelphia, Pennsylvania 19122
⁵⁰Texas A&M University, College Station, Texas 77843
⁵¹University of Texas, Austin, Texas 78712
⁵²Tsinghua University, Beijing 100084
⁵³University of Tsukuba, Tsukuba, Ibaraki 305-8571, Japan
⁵⁴United States Naval Academy, Annapolis, Maryland 21402
⁵⁵Valparaiso University, Valparaiso, Indiana 46383
⁵⁶Variable Energy Cyclotron Centre, Kolkata 700064, India
⁵⁷Warsaw University of Technology, Warsaw 00-661, Poland
⁵⁸Wayne State University, Detroit, Michigan 48201
⁵⁹Yale University, New Haven, Connecticut 06520
⁶⁰Huzhou University, Huzhou, Zhejiang 313000



(Received 26 December 2018; published 20 March 2019)

We report a new measurement of D^0 -meson production at mid-rapidity ($|y| < 1$) in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV utilizing the heavy flavor tracker, a high resolution silicon detector at the STAR experiment. Invariant yields of D^0 mesons with transverse momentum $p_T \lesssim 9$ GeV/c are reported in various centrality bins (0–10%, 10–20%, 20–40%, 40–60%, and 60–80%). Blast-wave thermal models are used to fit the D^0 -meson p_T spectra to study D^0 hadron kinetic freeze-out properties. The average radial flow velocity extracted from the fit

is considerably smaller than that of light hadrons (π , K , and p), but comparable to that of hadrons containing multiple strange quarks (ϕ , Ξ^-), indicating that D^0 mesons kinetically decouple from the system earlier than light hadrons. The calculated D^0 nuclear modification factors reaffirm that charm quarks suffer a large amount of energy loss in the medium, similar to those of light quarks for $p_T > 4 \text{ GeV}/c$ in central 0–10% Au + Au collisions. At low p_T , the nuclear modification factors show a characteristic structure qualitatively consistent with the expectation from model predictions that charm quarks gain sizable collective motion during the medium evolution. The improved measurements are expected to offer new constraints to model calculations and help gain further insights into the hot and dense medium created in these collisions.

DOI: [10.1103/PhysRevC.99.034908](https://doi.org/10.1103/PhysRevC.99.034908)

I. INTRODUCTION

The heavy ion program at the Relativistic Heavy Ion Collider (RHIC) and Large Hadron Collider (LHC) focuses on the study of strong interactions and quantum chromodynamics (QCD) at high temperature and density. Over the last few decades, experimental results from RHIC and LHC using light flavor probes have demonstrated that a strongly coupled quark-gluon plasma (sQGP) is created in these heavy-ion collisions. The most significant evidence comes from the strong collective flow and the large high transverse momentum (p_T) suppression in central collisions for various observed hadrons including multistrange-quark hadrons ϕ and Ω [1–5].

Heavy quarks (c , b) are created predominantly through initial hard scatterings due to their large masses [6,7]. The modification to their production in transverse momentum due to energy loss and radial flow and in azimuth due to anisotropic flows is sensitive to heavy quark dynamics in the partonic sQGP phase [8]. Recent measurements of high- p_T D -meson production at RHIC and LHC show a strong suppression in the central heavy-ion collisions [9–12]. The suppression is often characterized by the nuclear modification factor R_{AA} , defined as

$$R_{AA}(p_T) = \frac{1}{\langle T_{AA} \rangle} \frac{dN_{AA}/dp_T}{d\sigma_{pp}/dp_T}, \quad (1)$$

where dN_{AA}/dp_T and $d\sigma_{pp}/dp_T$ are particle production yield and cross section in $A + A$ and $p + p$ collisions, respectively. The nuclear thickness function $T_{AA} = \langle N_{\text{bin}} \rangle / \sigma_{pp}^{\text{inel}}$ is often calculated using a Monte Carlo Glauber model, where $\langle N_{\text{bin}} \rangle$ is the average number of binary collisions and $\sigma_{pp}^{\text{inel}}$ is the total inelastic $p + p$ cross section. The D -meson R_{AA} is similar to that of light hadrons for $p_T > 4 \text{ GeV}/c$, suggesting significant energy loss for charm quarks inside the sQGP medium. The measured D -meson anisotropic flow shows that D mesons also exhibit significant elliptic and triangular flow at RHIC and LHC [13–16]. The flow magnitude, when scaled with the transverse kinetic energy, is similar to that of light and strange flavor hadrons indicating that charm quarks may have reached thermal equilibrium in these collisions at RHIC and LHC.

In this article, we report measurements of the centrality dependence of D^0 -meson transverse momentum spectra at mid-rapidity ($|\eta| < 1$) in Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$. The measurements are conducted at the solenoidal tracker at RHIC (STAR) experiment utilizing the high resolution silicon detector (the heavy flavor tracker, HFT) [17]. The paper is organized in the following order: In Sec. II, we describe the detector setup and dataset. In Sec. III, we present the topological reconstruction of D^0 mesons in the Au + Au

collision data, followed by Secs. IV and V for details on efficiency corrections and systematic uncertainties, respectively. We present our measurement results and physics discussions in Sec. VI. Finally, we end the paper with a summary in Sec. VII.

II. EXPERIMENTAL SETUP AND DATASET

The dataset used in this analysis consists of Au + Au collision events at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$ collected in the 2014 RHIC run. The main detectors used are the time projection chamber (TPC), the HFT, the time of flight (TOF) detector, and the vertex position detector (VPD).

A. Tracking and particle identification subsystems

Particle tracking for this analysis is achieved with the TPC and HFT detectors. The HFT detector provides measured space points with high precision that are used to extend track trajectories from the TPC and offer high-pointing resolution to the vicinity of the event vertex. Particle identification is achieved with a combination of the ionization energy loss (dE/dx) measurement with the TPC and the time-of-flight (tof) measurement with the TOF detector. The event start time is provided by the VPD. Both the TPC and TOF detectors have full azimuthal coverage with a pseudorapidity range of $|\eta| < 1$ [18,19]. The TPC and TOF subsystems have been extensively used in many prior STAR analyses, including D -meson measurements [4,12,20].

B. Trigger and dataset

The minimum bias trigger is defined as a coincidence between the east and west VPD detectors located at $4.4 < |\eta| < 4.9$ [21]. Each VPD detector is an assembly of 19 small detectors, each consisting of a Pb converter followed by a fast plastic scintillator read out by a photomultiplier tube. To efficiently sample the collision events in the center of the HFT acceptance, an online cut on the collision vertex position along the beam line (calculated via the time difference between the east and west VPD detectors) $|V_z^{\text{VPD}}| < 6 \text{ cm}$ is applied. The decrease in the coincidence probability in the VPD degrades the online VPD vertex resolution in peripheral low multiplicity events. These inefficiencies are corrected in the offline analysis with a method discussed in the next section.

Events are selected with the offline reconstructed collision vertex [22] within 6 cm of the TPC and HFT centers along the beam direction to ensure uniform and large acceptance. Event pileup due to the long TPC readout time ($\approx 40 \mu\text{s}$) with respect to the Au-Au collision rate (40 KHz) is rejected by




FIG. 1. (a) Uncorrected charged particle multiplicity $N_{\text{ch}}^{\text{raw}}$ distribution measured within $|\eta| < 0.5$ and $|V_z^{\text{TPC}}| < 6$ cm. The solid curve depicts the multiplicity distribution from a MC Glauber simulation fit to the experimental data. (b) Ratio between MC simulation and real data.

removing events that have an online and offline vertex position difference along the z axis ($|V_z^{\text{VPD}} - V_z^{\text{TPC}}|$) greater than 3 cm. Approximately 9×10^8 minimum bias triggered events with 0–80% centrality pass the selection criteria and are used in this analysis.

C. Centrality selection and trigger inefficiency

The centrality is selected using the measured charged global track multiplicity $N_{\text{ch}}^{\text{raw}}$ at mid-rapidity within $|\eta| < 0.5$. The events with low track multiplicity are corrected for the online VPD triggering inefficiency using a Monte Carlo (MC) Glauber simulation. 0- $X\%$ ($X = 5, 10$, etc.) centrality is defined as the 0- $X\%$ most central in terms of the total hadronic cross section determined by the impact parameter between two colliding nuclei. In this analysis, the dependence of $N_{\text{ch}}^{\text{raw}}$ on the collision vertex position and the beam luminosity has been corrected for through scaling, which takes into account the dependence of the detector acceptance and inefficiencies on these quantities [23]. The measured track multiplicity distribution from Au+Au 200 GeV from RHIC run 2014, corrected for the vertex and luminosity dependence, is shown in Fig. 1. The measured distribution is fit by the MC Glauber calculation in the high multiplicity region. One can observe that the fitted MC Glauber calculation matches the real data well for $N_{\text{ch}}^{\text{raw}} > 100$, while the discrepancy in the low multiplicity region shows the VPD trigger inefficiency. Figure 1(b) shows the ratio between MC and data. Centrality is defined according to the MC Glauber model distribution shown in Fig. 1(a). Events in the low-multiplicity region are weighted with the ratio shown in Fig. 1(b) in all the following analysis as a correction for the inefficiency in trigger.

Table I lists the extracted values of average number of binary collisions (N_{bin}), average number of participants (N_{part}),

TABLE I. Estimated values of average number of binary collisions (N_{bin}), average number of participants (N_{part}), and trigger correction factors (ϵ_{trg} , uncertainties negligible) for defined centrality bins obtained from the MC Glauber model fit to the measured multiplicity distributions.

Centrality	$\langle N_{\text{bin}} \rangle$	$\langle N_{\text{part}} \rangle$	ϵ_{trg}
0–10%	938.8 ± 26.3	319.4 ± 3.4	1.0
10–20%	579.9 ± 28.8	227.6 ± 7.9	1.0
20–40%	288.3 ± 30.4	137.6 ± 10.4	1.0
40–60%	91.3 ± 21.0	60.5 ± 10.1	0.92
60–80%	21.3 ± 8.9	20.4 ± 6.6	0.65

and trigger inefficiency correction factors (ϵ_{trg}) in the defined centrality bins. The ϵ_{trg} correction factor is applied event by event in the analysis when combining centrality bins.

D. Heavy flavor tracker

The HFT [17] is a high resolution silicon detector system that aims for topological reconstruction of secondary vertices from heavy flavor hadron decays. It consists of three silicon subsystems: the silicon strip detector (SSD), the intermediate silicon tracker (IST), and two layers of PiXeL (PXL) detectors. Table II lists the key characteristic parameters of each subsystem. The SSD detector was still under commission when the dataset was recorded, and therefore is not used in the offline data production and this analysis. The PXL detector uses the new monolithic active pixel sensors (MAPS) technology [17]. This is the first application of this technology in a collider experiment.

In the offline reconstruction, tracks are reconstructed in the TPC first and then extended to the HFT detector to find the best match to the measured high resolution spatial points. A Kalman filter algorithm that considers various detector material effects is used in the track extension [24]. Considering the level of background hits in the PXL detector due to pileup collisions, tracks are required to have at least one hit in each layer of the PXL and IST subdetectors. Figure 2 shows the track pointing resolution to the primary vertex in the transverse plane (σ_{XY}) in Fig. 2(a) and along the longitudinal direction (σ_Z) in Fig. 2(b) as a function of total momentum (p) for identified particles in 0–80% centrality Au + Au collisions. The design goal for the HFT detector was to have a pointing resolution better than $55 \mu\text{m}$ for 750 MeV charged kaon. Figure 2 demonstrates that the HFT detector system meets the design requirements. This performance enables precision measurement of D -meson production in high multiplicity heavy-ion collisions.

III. D^0 -MESON RECONSTRUCTION

D^0 and \bar{D}^0 mesons are reconstructed via the hadronic decay channel $D^0 \rightarrow K^-\pi^+$ and its charge conjugate channel with a branching ratio (B.R.) of 3.89% [25]. In what follows, we imply $(D^0 + \bar{D}^0)/2$ when using the term D^0 unless otherwise specified. D^0 mesons decay with a proper decay length of $c\tau \approx 123 \mu\text{m}$ after they are produced in Au + Au collisions. We utilize the high-pointing resolution capability enabled by

TABLE II. Several key characteristic parameters for each subsystem of the STAR HFT detector.

Subsystem	Radius (cm)	Length (cm)	Thickness at $\eta = 0(X_0)$	Pitch size (μm^2)
PXL inner layer	2.8	20	0.52% (0.39% ^a)	20.7×20.7
PXL outer layer	8.0	20	0.52%	20.7×20.7
IST	14.0	50	1.0%	600×6000
SSD ^b	22.0	106	1.0%	$95 \times 40\,000$

^aPXL inner detector material is reduced to 0.39% X_0 in 2015/2016 runs.

^bSSD is not included in this analysis.

the HFT detector to topologically reconstruct the D^0 decay vertices that are separated from the collision vertices, which drastically reduces the combinatorial background (around five orders of magnitude) and improves the measurement precision.

Charged pion and kaon tracks are reconstructed with the TPC and HFT. Tracks are required to have at least 20 measured TPC points out of a maximum of 45 to ensure a good momentum resolution. To enable high pointing precision, both daughter tracks are required to have at least one measured hit in each layer of the PXL and IST as described above. Particle identification is achieved via a combination of the ionization energy loss measurement in the TPC and the time-of-flight measurement in the TOF. The resolution-normalized dE/dx deviation from the expected values is defined as

$$n\sigma_X = \frac{1}{R} \ln \frac{\langle dE/dx \rangle_{\text{mea.}}}{\langle dE/dx \rangle_X}, \quad (2)$$

where $\langle dE/dx \rangle_{\text{mea.}}$ and $\langle dE/dx \rangle_X$ represent measured and expected values with a hypothesis of particle X , and R is the dE/dx resolution (typically $\approx 8\%$ [18]). The $n\sigma_X$ distribution should be close to a standard Gaussian for each corresponding particle species (mean = 0, $\sigma = 1$) with good dE/dx calibration. Pion (kaon) candidates are selected by a requirement of the measured dE/dx to be within three (two) standard deviations ($|n\sigma_X|$) from the expected value. When tracks have matched hits in the TOF detector, an additional requirement on the measured inverse particle velocity ($1/\beta$) to be within three standard deviations from the expected value ($|\Delta 1/\beta|$) is applied for either daughter track. Figures 3 and 4




FIG. 2. Identified particle (π^\pm , K^\pm , and $p + \bar{p}$) pointing resolution in the transverse (a) and longitudinal (b) planes as a function of particle momentum in Au + Au 0–80% collisions.

show examples of the particle identification capability from the TPC and TOF. Tracks within the kinematic acceptance $p_T > 0.6 \text{ GeV}/c$ and $|\eta| < 1$ are used to combine and make pairs. The choice of the $p_T > 0.6 \text{ GeV}/c$ cut is an optimized consideration to balance the loss of signal acceptance when tightening the cut, and the increase in background due to the HFT fake matches when loosening this cut (see Sec. IV B). The threshold has been varied for systematic uncertainty evaluation. See Sec. V for details. Table III lists the TPC and TOF selection cuts for daughter kaon and pion tracks used for D^0 reconstruction.

The D^0 decay vertex is reconstructed as the middle point on the distance of the closest approach between the candidate pion and kaon trajectories. One of the dominant background sources is the random combination of $K\pi$ pairs directly from the collision point. The following topological variables have been found to greatly reduce the combinatorial background:

- (i) decay length: the distance between the reconstructed decay vertex and the primary vertex (PV);
- (ii) distance of closest approach (DCA) between the two daughter tracks (DCA_{12});
- (iii) DCA between the reconstructed D^0 momentum vector and the PV (DCA_{D^0});
- (iv) DCA between the pion track and the PV (DCA_π);
- (v) DCA between the kaon track and the PV (DCA_K);





FIG. 3. TPC dE/dx vs particle momentum.

FIG. 4. TOF $1/\beta$ vs particle momentum.


- (vi) angle between the D^0 momentum and the direction of the decay vertex with respect to the PV (θ).

The schematic in Fig. 5 shows the topological variables used in the analysis, where \vec{P} represents the D^0 momentum. The decay length and angle θ follow the formula $\text{DCA}_{D^0} = \text{decay length} \times \sin(\theta)$. The cuts on the topological variables for this analysis are optimized using a toolkit for multivariate data analysis (TMVA) package integrated in the ROOT framework in order to obtain the greatest signal significance [26]. The rectangular cut optimization method from the TMVA package is chosen, similarly as in our previous publication [16]. The optimization is conducted for different D^0 p_T bins and different centrality bins. Table IV lists a set of topological cuts for 0–10% central Au + Au collisions.

Figure 6 shows the invariant mass distributions of $K\pi$ pairs in the p_T region of 0–10 GeV/c for 0–80% minimum bias and the 0–10% most central collisions, and 0–8 GeV/c for 60–80% peripheral collisions, respectively. The reason for choosing a different p_T range for the 60–80% centrality bin is because no signal is observed beyond the current statistics. The combinatorial background is estimated with the same-event (SE) like-sign (LS) pairs (blue histograms) and the mixed-event (ME) unlike-sign (US) (grey histograms) technique in which K and π from different events of similar characteristics (V_Z , centrality, event plane angle) are paired. The mixed-event spectra are normalized to the like-sign distributions in the

TABLE III. TPC and TOF selection cuts for K and π tracks.

Variable		K^\mp	π^\pm
p_T (GeV/c)	>	0.6	0.6
$ \eta $	<	1.0	1.0
nHitsFit (TPC)	>	20	20
$ n\sigma_X $	<	2.0	3.0
$ \Delta 1/\beta $ (if TOF matched)	<	0.03	0.03

FIG. 5. A cartoon picture for $D^0 \rightarrow K^- + \pi^+$ decay and definition of topological variables used in the reconstruction.

mass range of 1.7–2.1 GeV/ c^2 . After the subtraction of the mixed-event unlike-sign combinatorial background from the same-event unlike-sign pairs (black open circles), the remainder distributions are shown as red solid circles in each panel. Compared to the previous D^0 measurement [12], the D^0 signal significance is largely improved by a factor of ≈ 15 using the same amount of event statistics.

Figures 7 and 8 show the invariant mass distributions in the same centrality bins as Fig. 6 but for different p_T ranges: $0 < p_T < 0.5$ GeV/c in Fig. 7 and $6 < p_T < 8$ GeV/c in Fig. 8.

After the combinatorial background is subtracted, the residual $K\pi$ invariant mass distributions are fit to a Gaussian plus linear function. The linear function is used to represent remaining correlated background from either partial reconstruction of charm mesons or jet fragments. The D^0 raw yields are extracted from the Gaussian function fit results and the systematic uncertainty on the extracted raw yield is evaluated using several methods described in Sec. V.

IV. EFFICIENCIES AND CORRECTIONS

The reconstructed D^0 raw yields are calculated in each centrality and p_T bin within the rapidity window $|y| < 1$. The

TABLE IV. Topological cuts used for D^0 reconstruction in 0–10% most central collisions for separate p_T intervals.

0–10% p_T (GeV/c)	(0,0.5)	(0.5,1)	(1,2)	(2,3)	(3,5)	(5,8)	(8,10)
Decay length (μm)	> 100	199	227	232	236	255	255
DCA_{12} (μm)	< 71	64	70	63	82	80	80
DCA_{D^0} (μm)	< 62	55	40	40	40	44	44
DCA_π (μm)	> 133	105	93	97	67	55	55
DCA_K (μm)	> 138	109	82	94	76	54	54
$\cos(\theta)$	> 0.95	0.95	0.95	0.95	0.95	0.95	0.95




FIG. 6. Invariant mass $M_{K\pi}$ distributions in $0 < p_T < 10$ GeV/c from centrality bins 0–80% (a), 0–10% (b), and in $0 < p_T < 8$ GeV/c for 60–80% (c), respectively. Black open circles represent the same-event (SE) unlike-sign (US) distributions. Blue and grey shaded histograms represent the SE like-sign (LS) and mixed-event (ME) US distributions that are used to estimate the combinatorial background. The red solid circles depict the US (SE) distributions with the combinatorial background subtracted using the US (ME) distributions.

fully corrected D^0 production invariant yields are calculated using the following formula:

$$\frac{d^2N}{2\pi p_T dp_T dy} = \frac{1}{\text{B.R.}} \times \frac{N^{\text{raw}}}{N_{\text{evt}} 2\pi p_T \Delta p_T \Delta y} \times \frac{1}{\varepsilon_{\text{trg}} \times \varepsilon_{\text{TPC}} \times \varepsilon_{\text{HFT}} \times \varepsilon_{\text{PID}} \times \varepsilon_{\text{vtx}}}, \quad (3)$$

where B.R. is the $D^0 \rightarrow K^-\pi^+$ decay branching ratio, $(3.89 \pm 0.04)\%$ [25], N^{raw} is the reconstructed D^0 raw counts, N_{evt} is the total numbers of events used in this analysis, and ε_{trg} is the centrality bias correction factor described in Sec. II B. The raw yields are corrected for the TPC acceptance and tracking efficiency ε_{TPC} , the HFT acceptance and tracking

plus topological cut efficiency ε_{HFT} , the particle identification efficiency ε_{PID} , and the finite vertex resolution correction ε_{vtx} .

A. TPC acceptance and tracking efficiency ε_{TPC}

The TPC acceptance and tracking efficiency is obtained using the standard STAR TPC embedding technique, in which a small amount of MC tracks (typically 5% of the total multiplicity of the real event) are processed through the full GEANT simulation [27], then mixed with the raw data acquisition (DAQ) data in real events and reconstructed through the same reconstruction chain as the real data production. The TPC efficiency is then calculated as the ratio of the number of reconstructed MC tracks with the same offline analysis cuts




FIG. 7. Invariant mass $M_{K\pi}$ distributions in $0 < p_T < 0.5$ GeV/c from centrality bins 0–80% (a), 0–10% (b), and 60–80% (c), respectively. All histograms and markers use the same notation as in Fig. 6.




FIG. 8. Invariant mass $M_{K\pi}$ distributions in $6 < p_T < 8$ GeV/c from centrality bins 0–80% (a), 0–10% (b), and 60–80% (c), respectively. All histograms and markers use the same notation as in Fig. 6.

for geometric acceptance and other TPC requirements to that of the input MC tracks.

Figure 9 shows the TPC acceptance and tracking efficiency ε_{TPC} for D^0 mesons within $|y| < 1$ in various centrality classes. The efficiencies include the TPC and analysis acceptance cuts $p_T > 0.6$ GeV/c and $|\eta| < 1$ as well as the TPC tracking efficiency for both pion and kaon daughters. The lower efficiency observed in central collisions is due to the increased multiplicity resulting higher detector occupancy which leads to reduced tracking efficiency in these collisions.

B. HFT acceptance, tracking and topological cut efficiency ε_{HFT}

1. Data-driven simulation

In order to fully capture the real-time detector performance, the HFT-related efficiency is obtained using a data-driven simulation method, i.e., the HFT related efficiencies




FIG. 9. D^0 TPC acceptance and tracking efficiencies from different centrality classes in Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200$ GeV.

are extracted from data as a function of kinematic variables and then a Monte Carlo is used to model the D^0 decay and data-driven efficiencies are folded into the decay productions according to their kinematics. The performance of inclusive HFT tracks is characterized by a HFT-to-TPC matching efficiency ($\varepsilon_{\text{HFT}}^{\text{match}}$) and the DCA distributions with respect to the primary vertex. The HFT matching efficiency $\varepsilon_{\text{HFT}}^{\text{match}}$ is defined as the fraction of reconstructed TPC tracks that satisfy the requirement on the number of HFT hits. In this analysis, the requirement is to have at least one hit in each PXL and IST layer. The $\varepsilon_{\text{HFT}}^{\text{match}}$ includes the HFT geometric acceptance and the tracking efficiency that associate HFT hits to the extended TPC tracks. It contains the true matches for which the reconstructed tracks pick up real hits induced by these charged tracks when passing through the HFT, as well as some random fake matches. The latter has a decreasing trend as a function of p_T as the track pointing resolution gets better at high p_T , resulting in a smaller search window when associating HFT hits in the tracking algorithm. The DCA distributions are obtained for those tracks that satisfy the HFT hit requirement. Figure 10 shows an example of the HFT matching efficiency and the one-dimensional projection of the DCA_{XY} distribution for single pions at $1.0 < p_T < 1.2$ GeV/c and 0–10% central collisions. Such distributions obtained from real data are fed into a MC decay generator for $D^0 \rightarrow K^- \pi^+$, followed by the same reconstruction of D^0 secondary vertex as in the real data analysis. The same topological cuts are then applied and the HFT related efficiency for the D^0 reconstruction is calculated.

To best represent the real detector performance, we obtain the following distributions from real data in this Monte Carlo approach:

- (i) centrality-dependent V_z distributions;
- (ii) HFT matching efficiency $\varepsilon_{\text{HFT}}^{\text{match}}$, including the dependence on particle species, centrality, p_T , η , ϕ , and V_z ;
- (iii) DCA_{XY} - DCA_Z two-dimensional (2D) distributions including the dependence on particle species, centrality, p_T , η , and V_z .




FIG. 10. HFT matching efficiency $\varepsilon_{\text{HFT}}^{\text{match}}$ (a) and DCA_{XY} (c) distributions of inclusive charged pions from real data and MC simulation in 0–10% Au + Au collisions. The ratios between real data and GEANT simulation are shown in the bottom panels. The blue histogram depicts the true matches for which the reconstructed tracks pick up the correct MC hits in the HFT detector induced by the associated MC tracks in the GEANT simulation.

The DCA_{XY} - DCA_Z 2D distributions are the key to represent not only the true matches, but also the fake matches when connecting the TPC tracks with HFT hits. The distributions are obtained in two dimensions to consider the correlation between the two quantities and this is necessary and essential to reproduce the 3D DCA position distributions observed in real data. The ϕ dependence of these distributions are integrated over due to computing resource limits. We have checked the ϕ dependence (by reducing other dependencies for the same reason) and it gives a consistent result compared to the ϕ -integrated one.

In total, there are $11 (\phi) \times 10 (\eta) \times 6 (V_z) \times 9$ (centrality) $\times 2$ (particles) 1D histograms (36 p_T bins) used for the HFT matching efficiency distributions and $5 (\eta) \times 4 (V_z) \times 9$ (centrality) $\times 2$ (particles) $\times 19$ (p_T) 2D histograms (144 $\text{DCA}_{XY} \times 144 \text{DCA}_Z$ bins) for 2D DCA distributions. The number of bins chosen is optimized to balance the need of computing resources as well as the stability of the final efficiency. All dimensions have been checked so that further increase in the number of bins (in balance we need to reduce the number of bins in other dimensions) will not change the final obtained efficiency.

The procedure for this data-driven simulation package for efficiency calculation is as follows:

- (i) Sample V_z distribution according to the distribution obtained from the real data.
- (ii) Generate D^0 at the event vertex position with desired p_T (Levy function shape fitted to D^0 spectra [12]) and rapidity (flat) distributions.

- (iii) Propagate D^0 and simulate its decay to $K^-\pi^+$ daughters following the decay probability.
- (iv) Smear daughter track momentum according to the values obtained from embedding.
- (v) Smear daughter track starting position according to the DCA_{XY} - DCA_Z 2D distributions from the reconstructed data.
- (vi) Apply HFT matching efficiency according to that extracted from the reconstructed data.
- (vii) Perform the topological reconstruction of D^0 decay vertices with the same cuts as applied in the data analysis and calculate the reconstruction efficiency.

The DCA and HFT matching efficiency distributions used as the input in this simulation tool are obtained from the real data or can be the reconstructed data in MC simulation. The MC GEANT simulation is used as a systematic check to the data-driven inputs (see Sec. IV B 2).

This approach assumes these distributions obtained from real data are good representations for tracks produced at or close to the primary vertices. The impact of the secondary particle contribution will be discussed in Sec. IV B 4. The approach also neglects the finite event vertex resolution contribution which will be discussed in Sec. IV C.

Last in this MC approach, we also fold in the TPC efficiency obtained from the MC embedding so the following presented efficiency will be the total efficiency of $\varepsilon_{\text{TPC}} \times \varepsilon_{\text{HFT}}$.

2. Validation with GEANT simulation

The data-driven simulation method is validated with a full GEANT simulation with the STAR detector with HIJING




FIG. 11. Comparisons in topological variable distributions between MC GEANT simulation (black) and data-driven fast simulation with reconstructed MC data as the input (red) in 0–10% Au + Au collisions for D^0 mesons at $2 < p_T < 3$ GeV/c.

events enriched with D^0 mesons. The D^0 here is reconstructed in the same manner as real data, and therefore, a true efficiency is calculated as the ratio of reconstructed to generated D^0 candidates. A second estimate of efficiency is calculated with the data-driven simulation method, but instead using the input distributions from the HIJING event. Both methods should return the same efficiency if the data-driven simulation method works perfectly.

The GEANT simulation uses the HIJING [28] generator as its input with D^0 particles embedded to enrich the signal statistics. The full HFT detector materials (both active and inactive) have been included in the GEANT simulation as well as the offline track reconstruction. The pileup hits in the PXL detector due to finite electronic readout time have been added to realistically represent the HFT matching efficiency and DCA distributions. The overall agreement between the GEANT simulation and real data is fairly good, as can be seen in Fig. 10. The small deviations between real data and MC simulation are not considered in the systematic uncertainty estimation since the latter is not used to calculate the absolute efficiency directly, but to validate the data-driven simulation procedure as described below.

The increase in the HFT matching efficiency at low p_T range is due to the increased fake matches (in contrast to true HFT matches) and the efficiency stays flat in the high p_T range. The matching efficiency includes the tracking efficiency when associating the HFT hits as well as the HFT geometric acceptance. Therefore, the ratio has a strong dependence on the event V_z and the track η . The DCA distributions used in the package are two-dimensional distributions, as DCA_{XY} and DCA_z are strongly correlated.

With the tuned simulation setup, we use this sample to validate our data-driven simulation approach for D^0 efficiency calculation. We follow the same procedure as described in Sec. IV B 1 to obtain the HFT matching efficiency as well as the 2D DCA_{XY} - DCA_z distributions for primary particles from the reconstructed data in this simulation sample. These distributions are then fed into the data-driven simulation framework to calculate the D^0 reconstruction efficiency. The calculated D^0 efficiency from the data-driven simulation framework will be compared to the real D^0 reconstruction efficiency directly obtained from the GEANT simulation sample.

To validate the data-driven simulation tool, Fig. 11 shows the comparisons of several topological variables used in the




FIG. 12. (a) D^0 reconstruction efficiency comparison between MC GEANT simulation (black) and data-driven fast simulation with reconstructed MC data as the input (red) in central 0–10% Au + Au collisions. (b) The ratio between the two methods. The grey band around unity represents the 5% systematic uncertainties.

D^0 reconstruction obtained from the GEANT simulation directly and from the data-driven simulation with reconstructed GEANT simulation data as the input in the most central (0–10%) centrality and in $2 < p_T < 3$ GeV/c. The topological variables shown here are D^0 decay length, DCA between two D^0 decay daughters, D^0 DCA with respect to the collision vertex, pion DCA and kaon DCA with respect to the collision vertex. As seen in this figure, the data-driven simulation tool reproduces all of these topological distributions quite well. The agreements for the other p_T ranges are also decent.

Figure 12 shows the D^0 reconstruction efficiency $\epsilon_{\text{TPC}} \times \epsilon_{\text{HFT}}$ calculated with the following two methods in this GEANT simulation. The first method is the standard calculation by applying the tracking and topological cuts for reconstructed D^0 mesons in the simulation sample. In the second method, we employ the data-driven simulation method and take the reconstructed distributions from the simulation sample as the input and then calculate the D^0 reconstruction efficiency in the data-driven simulation framework. In Fig. 12(a), efficiencies from two calculation methods agree well in the whole p_T region in central 0–10% Au + Au collisions, and the ratio between the two is shown in panel (b). This demonstrates that the data-driven simulation framework can accurately reproduce the real D^0 reconstruction efficiency in central Au + Au collisions.

3. Efficiency for real data

We employ the validated data-driven simulation method for the real data analysis. Figure 13 shows comparisons of the same five topological variables between D^0 signals in real data and data-driven simulated distributions with real data as

the input in central 0–10% collisions for D^0 mesons at $2 < p_T < 3$ GeV/c. The real data distributions are extracted by reconstructing D^0 signals with the same reconstruction cuts as in Sec. III except for the interested topological variable to be compared. The distributions for D^0 candidates are generated by statistically subtracting the background using the like-sign method from the same-event unlike-sign distributions within the D^0 mass window (1.82–1.91 GeV/c²) [16,20]. The cut on the interested topological variable is loosened, but one needs to place some precuts to ensure reasonable D^0 signal reconstruction for the extraction of these topological variable distributions. These precuts effectively reduce the low-end reach for several topological variables, e.g., the D^0 decay length. In the data-driven simulation method, charged pion and kaon HFT matching efficiencies and 2D DCA distributions are used as the input to calculate these topological variables for D^0 signals. Figure 13 shows that in the selected ranges, the data-driven simulation method reproduces topological variables distributions of D^0 signals, which supports that this method can be reliably used to calculate the topological cut efficiency.

Figure 14 shows the HFT tracking and topological cut efficiency ϵ_{HFT} as a function of D^0 p_T for different centrality bins obtained using the data-driven simulation method described in this section. The smaller efficiency for central collisions is due to the lower HFT matching efficiency in higher occupancy collisions and tighter topological cuts in central collisions used to reduce the larger combinatorial background.

4. Secondary particle contribution

The impact on the data-driven efficiency from the secondary particle contribution (e.g., weak decays from K_S^0 and Λ) to the inclusive pion distributions is studied using the HIJING events processed through the GEANT simulation and the same offline reconstruction. The fraction of secondary pions from weak decay of strange hadrons (K_S^0 and Λ) to the total inclusive charged pions within DCA < 1.5 cm cut is estimated to be around 5% at pion $p_T = 0.3$ GeV/c and decreases to be <2% above 2 GeV/c. This is consistent with what was observed before in measuring the prompt charged pion spectra [29]. There is another finite contribution of low momentum antiprotons and antineutrons annihilated in the detector material and producing secondary pions. The transverse momenta of these pions are mostly around 2–3 GeV/c and the fraction of total inclusive pions is ≈ 10 –12% at $p_T = 2$ –3 GeV/c based on this simulation and contribute ≈ 5 –8% to the HFT matching efficiency. This is obtained using the GEANT simulation with GHEISHA hadronic package. With a different hadronic package, GEANT-FLUKA [30], the secondary pion fraction in 2–3 GeV/c region is significantly reduced to be negligible. The difference between the primary pions and the inclusive pions in the HFT matching efficiency has been considered as one additional correction factor in our data-driven simulation method when calculating the final efficiency. The maximum difference with respect to the result obtained using the GHEISHA hadronic package is used as the systematic uncertainty for this source. Figure 15 shows the secondary pion contribution in Au + Au collisions with the FLUKA hadronic package. Panel (a) shows the




FIG. 13. Comparison of topological variable distributions between D^0 signals in real data (black) and in data-driven simulation with real data distributions as the input (red) in 0–10% Au + Au collisions for D^0 mesons at $2 < p_T < 3 \text{ GeV}/c$. The dashed lines indicate the final topological cuts chosen for each individual topological variable.

fraction of different sources for secondary tracks including the weak decays, the scattering, and the \bar{p}/\bar{n} annihilation in the detector material. Panel (b) shows the HFT matching efficiencies for inclusive, prompt, and secondary pions. Panel (c) is the ratio of the HFT matching efficiencies between the inclusive and the primary pions from panel (b). The effect of such secondary contribution to charged kaons is found to be negligible [29].




FIG. 14. D^0 HFT tracking and topological cut efficiencies ϵ_{HFT} from different centrality classes in Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$.

C. Vertex resolution correction ϵ_{vtx}

In the data-driven approach, D^0 mesons are injected at the event vertex. In the real data, the reconstructed vertex has a finite resolution with respect to the real collision vertex. This may have some effect on the reconstructed D^0 signal counts after applying the topological cuts in small multiplicity events where the event vertex resolution decreases. We carry out similar simulation studies as described in Sec. IV B 1 for other centrality bins. Figure 16 shows the full width at half maximum (FWHM) of the difference in the vertex x position of two randomly divided subevents in various centrality bins between data and MC simulation. We choose the FWHM variable here as the distributions are not particularly Gaussian.




FIG. 15. Secondary pion contribution estimated from HIJING + GEANT simulation with FLUKA hadronic package. Panel (a) shows the fraction of different sources for secondary pion tracks. Panel (b) shows the HFT matching efficiency $\varepsilon_{\text{HFT}}^{\text{match}}$ for inclusive, primary, and secondary pions. Panel (c) shows the ratio of HFT matching efficiencies between inclusive and primary pions.

The MC simulation reproduces the vertex difference distributions seen in the real data reasonably well. This gives us confidence for using this MC simulation setup to evaluate the vertex resolution correction ε_{vtx} .




FIG. 16. Full-width at half maximum (FWHM) of vertex position difference in the X dimension between two randomly divided subevents in various centrality bins. Black solid circles present the FWHM values from real data while blue empty circles are from HIJING + GEANT simulation. Statistical uncertainties are smaller than the marker size.

To estimate the vertex resolution effect, we embed a single PYTHIA $c\bar{c}$ event into a HIJING Au + Au event, and the whole event is passed through the STAR GEANT simulation followed by the same offline reconstruction as in the real data production. The PYTHIA $c\bar{c}$ events are preselected to have at least one $D^0 \rightarrow K^-\pi^+$ decay or its charge conjugate to enhance the statistics. Figure 17 shows a comparison of the obtained D^0 reconstruction efficiency between MC simulation (black) and data-driven simulation using reconstructed MC data as the input (red) for 20–30% (left), 50–60% (middle), and 70–80% (right) centrality bins, respectively. The bottom panels show the ratios of the efficiencies obtained from the two calculation methods. In central and mid-central collisions, the data-driven simulation method can properly reproduce the D^0 real reconstruction efficiency. This is expected since the vertex resolution is small enough so that it has negligible impact on the obtained efficiency using the data-driven simulation method. However, in more peripheral collisions, the data-driven simulation method overestimates the D^0 reconstruction efficiency as shown in the middle and right panels. The vertex resolution correction factor ε_{vtx} , denoted in Eq. (3), has a mild p_T dependence but strong centrality dependence as shown in Fig. 18 for $p_T \approx 2$ and 4 GeV/c. The brackets denote the systematic uncertainties in the obtained correction factor ε_{vtx} . They are estimated by changing the multiplicity range in the HIJING + GEANT simulation so that the variation in the subevent vertex difference distributions from the real data can be covered by distributions obtained from different simulation samples. The vertex resolution corrections are applied as a function of p_T in each individual centrality class.

D. PID efficiency ε_{PID} and doubly mis-PID correction

The D^0 daughter particle identification (PID) cut efficiency includes contributions from the dE/dx selection cut efficiency




FIG. 17. D^0 reconstruction efficiency comparison between MC GEANT simulation (GEANT, black) and data-driven simulation with the reconstructed MC data as the input (D-D S., red) for 20–30% (a), 50–60% (c), and 70–80% (e) Au + Au collisions. Bottom panels (b), (d), and (f) show the ratios between the two distributions above.

as well as the TOF matching and $1/\beta$ cut efficiency. To best estimate the selection cut efficiency, we select the enriched kaon and pion samples from ϕ , K_S^0 decays following the same procedure as in Refs. [31,32] and obtain the mean and width in the dE/dx $n\sigma_X$ distributions. The dE/dx cut efficiencies for pion and kaon daughter tracks are calculated correspondingly. The TOF $1/\beta$ cut efficiency is determined by studying the $1/\beta$ distributions for kaons and pions in the clean separation region, namely $p_T < 1.5$ GeV/c. There is a mild dependence for the offset and width of $\Delta 1/\beta$ distributions vs particle

momentum and our selection cuts are generally wide enough to capture nearly all tracks once they have valid β measurements. The total PID efficiency of D^0 mesons is calculated by folding the individual track TPC and TOF PID efficiencies following the same hybrid PID algorithm as implemented in the data analysis. Figure 19 shows the total PID efficiencies for D^0 reconstruction in various centrality bins. The total PID efficiency is generally high and has nearly no centrality or p_T dependence.

When the D^0 daughter kaon track is misidentified as a pion track and the other daughter pion track is misidentified as a kaon track, the pair invariant mass distribution will have




FIG. 18. ε_{vtx} , D^0 reconstruction efficiency ratios between MC GEANT simulation and data-driven simulation with the reconstructed MC data as the input vs collision centrality for p_T at 2 and 4 GeV/c. The brackets depict the estimated systematic uncertainties.




FIG. 19. Particle identification efficiency (ε_{PID}) of D^0 mesons in different centrality classes.




FIG. 20. D^0 yield double counting fraction due to doubly mis-PID in different centrality classes. The black markers depict an estimation taking the total double counting yield in the D^0 mass window while the blue markers depict an estimation with an additional sideband (SB) subtraction. Note that most data points from different centrality bins overlap with each other.

a bump structure around the real D^0 signal peak, but the distribution is much broader in a wide mass region due to the misassigned daughter particle masses. Based on the PID performance study described above, we estimate the single kaon and pion candidate track purities. After folding the realistic particle momentum resolution, we calculate the reconstructed D^0 yield from doubly misidentified pairs (double counting) underneath the real D^0 signal and the double counting fraction is shown in Fig. 20. The black markers show the fraction by taking all doubly misidentified pairs in the D^0 mass window while the blue markers depict it with an additional sideband (SB) subtraction. The latter is used as a correction factor to the central values of reported D^0 yields while the difference between the black and blue symbols is considered as the systematic uncertainty in this source. The double counting fraction is below 10% in all p_T bins, and there is little centrality dependence.

Figure 21 shows the total D^0 reconstruction efficiency from different centrality classes in Au + Au collisions including all of the individual components discussed above.

V. SYSTEMATIC UNCERTAINTIES

The systematic uncertainty on the final measured D^0 p_T spectra can be categorized as the uncertainty of the raw D^0 yield extraction, the BR uncertainty, and the uncertainty of efficiencies and corrections.

The uncertainty of the raw yield extraction is estimated by (a) changing the D^0 raw yield counting method from the Gaussian fit to histogram bin counting, (b) varying invariant mass ranges for fit and for side bands, and (c) varying background estimation between the mixed-event and like-sign methods. For the sideband method, the D^0 raw yield is obtained by




FIG. 21. The total D^0 reconstruction efficiency from different centrality classes.

subtracting the average counts in two invariant mass ranges around the signal (1.71–1.80 and 1.93–2.02 GeV/c^2) from the counts in the signal region (1.82–1.91 GeV/c^2) [16,20]. The maximum difference between these scenarios is then converted to the standard deviation and added to the systematic uncertainties. It is the smallest in the mid- p_T bins due to the best signal significance and grows at both low and high p_T . The double counting contribution in the D^0 raw yield due to mis-PID is included as another contribution to the systematic uncertainty for the D^0 raw yield extraction as described in Sec. IV D.

The uncertainty of the TPC acceptance and efficiency correction ϵ_{TPC} is estimated via the standard procedure in STAR by comparing the TPC track distributions between real data and the embedding data. It is estimated to be $\approx 5\text{--}7\%$ for 0–10% collisions and $\approx 5\text{--}8\%$ for 60–80% collisions, and is correlated for different centralities and p_T regions.

The uncertainty of the PID efficiency correction is estimated by varying the PID selection cuts and then convoluting to the final corrected D^0 yield.

To estimate the uncertainty of the HFT tracking and topological cut efficiency correction ϵ_{HFT} , we employ the following procedures: (a) We vary the topological variable cuts such that the D^0 ϵ_{HFT} is changed to 50% and 150% from the nominal (default) efficiency and compare the efficiency-corrected final D^0 yields. The maximum difference between the two scenarios is then added to the systematic uncertainties. (b) We also vary the lower threshold cut on the daughter p_T between 0.3 to 0.6 GeV/c and the maximum difference in the final corrected D^0 yield is also included in the systematic uncertainties. (c) We add the systematic uncertainty due to limitation of the data-driven simulation approach, $\approx 5\%$, and the impact of the secondary particles, $\approx 2\%$, to the total ϵ_{HFT} systematic uncertainty.

With the corrected D^0 transverse momentum spectra, the nuclear modification factor R_{CP} is calculated as the ratio of

TABLE V. Summary of systematic uncertainties, in percentage, on the D^0 invariant yield in 0–10% and 60–80% collisions and on R_{CP} (0–10%/60–80%).

Source	Systematic uncertainty (%)			Correlation in p_T
	0–10%	60–80%	R_{CP} (0–10%/60–80%)	
Signal extra.	1–6	1–12	2–13	uncorr.
Double mis-PID	1–7	1–7	negligible	uncorr.
ε_{TPC}	5–7	5–8	3–7	largely corr.
ε_{HFT}	3–15	3–20	3–20	largely corr.
ε_{PID}	3	3	negligible	largely corr.
ε_{vtx}	5	9–18	10–18	largely corr.
B.R.		0.5	0	global
N_{bin}	2.8	42	42	global

N_{bin} -normalized yields between central and peripheral collisions, as shown in the following formula:

$$R_{CP} = \frac{d^2N/dp_T dy}{N_{bin}} \Big|_{cen} \times \frac{N_{bin}}{d^2N/dp_T dy} \Big|_{peri}. \quad (4)$$

The systematic uncertainties in the raw signal extraction in central and peripheral collisions are propagated as they are uncorrelated, while the systematic uncertainties from the other sources are correlated or partially correlated in contributing to the measured D^0 yields. To study these correlations, we vary selection cuts simultaneously in central and peripheral collisions, and the difference in the final extracted R_{CP} value is then directly counted as systematic uncertainties in the measured R_{CP} .

The nuclear modification factor R_{AA} is calculated as the ratio of N_{bin} -normalized yields between Au + Au and $p + p$ collisions. The baseline for $p + p$ collisions is the same as in Ref. [12]. The uncertainties from the $p + p$ reference dominates the systematic uncertainty for R_{AA} . They include the 1σ uncertainty from the Levy function fit to the measured spectrum and the difference between Levy and power-law function fits for extrapolation to low and high p_T , expressed as one standard deviation.

With the corrected D^0 and \bar{D}^0 transverse momentum spectra, the \bar{D}^0/D^0 ratio is calculated as a function of the transverse momentum. The systematic uncertainties in the raw signal extraction for \bar{D}^0 and D^0 are propagated as they are uncorrelated, while the systematic uncertainties from the other sources are correlated or partially correlated in contributing to the measured \bar{D}^0/D^0 ratio. As in the R_{CP} systematic uncertainty estimation, we vary selection cuts simultaneously for D^0 and \bar{D}^0 , and the difference in the final extracted \bar{D}^0/D^0 value is then directly counted as systematic uncertainties for the measured \bar{D}^0/D^0 ratio.

Table V summarizes the systematic uncertainties and their contributions, in percentage, on the D^0 invariant yield in 0–10% and 60–80% collisions and R_{CP} (0–10%/60–80%). In the last column we also comment on the correlation in p_T for each individual source. Later when reporting p_T -integrated yields or R_{CP} , systematic uncertainties are calculated under the following considerations: (a) for p_T uncorrelated sources, we take the quadratic sum of various p_T bins; (b) for sources

that are largely correlated in p_T , we take the arithmetic sum as a conservative estimate.

VI. RESULTS AND DISCUSSION

A. p_T spectra and integrated yields

Figure 22 shows the efficiency-corrected D^0 invariant yield at mid-rapidity ($|y| < 1$) as a function of p_T in 0–10%, 10–20%, 20–40%, 40–60%, and 60–80% Au + Au collisions. D^0 spectra in some centrality bins are scaled with arbitrary factors indicated in the figure for clarity. Dashed and solid lines depict fits to the spectra with the Levy function:

$$\frac{d^2N}{2\pi p_T dp_T dy} = \frac{1}{2\pi} \frac{dN}{dy} \frac{(n-1)(n-2)}{nT[nT + m_0(n-2)]} \times \left(1 + \frac{\sqrt{p_T^2 + m_0^2} - m_0}{nT}\right)^{-n}, \quad (5)$$




FIG. 22. D^0 invariant yield at mid-rapidity ($|y| < 1$) vs transverse momentum for different centrality classes. Error bars (not visible for many data points) indicate statistical uncertainties and brackets depict systematic uncertainties. Global systematic uncertainties in B.R. are not plotted. Solid and dashed lines depict Levy function fits.




FIG. 23. (a) Measured D^0 spectra from this analysis compared with the previous 2010/11 measurements for different centrality classes. Dashed lines depict Levy function fits to 2014 data. (b)–(d) Ratio of measured spectra to the fitted Levy functions in 0–10%, 10–40%, and 40–80% centrality bins, respectively.

where m_0 is the D^0 mass ($1.864 \text{ GeV}/c^2$) and dN/dy , T , and n are free parameters. The Levy function fit describes the D^0 spectra nicely in all centrality bins in our measured p_T region.

We compare our new measurements with previous measurements using the STAR TPC only. The previous measurements have been recently corrected after fixing errors in the TOF PID efficiency calculation [12]. Figure 23 shows the p_T spectra comparison in 0–10%, 10–40%, and 40–80% centrality bins in panel (a) and the ratios to the Levy fit functions in panels (b)–(d), respectively. The new measurement with the HFT detector shows a nice agreement with the measurement without the HFT, but with significantly improved precision.

The measured D^0 spectra cover a wide p_T region which allows us to extract the p_T -integrated total D^0 yield at mid-rapidity with good precision. Figure 24 shows the




FIG. 24. Integrated D^0 cross section per nucleon-nucleon collision at mid-rapidity for $p_T > 0$ (a) and $p_T > 4 \text{ GeV}/c$ (b) as a function of centrality N_{part} . The statistical and systematic uncertainties are shown as error bars and brackets on the data points. The green boxes on the data points depict the overall normalization uncertainties in $p + p$ and Au + Au data respectively.

p_T -integrated cross section for D^0 production per nucleon-nucleon collision $d\sigma^{NN}/dy|_{y=0}$ from different centrality bins for the full p_T range shown in the top panel and for $p_T > 4 \text{ GeV}/c$ shown in the bottom panel. The result from the previous $p + p$ measurement is also shown in both panels [20].

While $d\sigma^{NN}/dy|_{y=0}$ for $p_T > 4 \text{ GeV}/c$ shows a clear decreasing trend from peripheral to mid-central and central collisions, that for the full p_T range shows an approximately flat distribution as a function of N_{part} , though the systematic uncertainty in the 60–80% centrality bin is a bit large. The values for the full p_T range in mid-central to central Au + Au collisions are smaller than that in $p + p$ collisions with $\sim 1.5\sigma$ effect considering the large uncertainties from the $p + p$ measurements. The total charm quark yield in heavy-ion collisions is expected to follow the number-of-binary-collision scaling since charm quarks are believed to be predominately created at the initial hard scattering before the formation of the QGP




FIG. 25. D^0 invariant yield at mid-rapidity ($|y| < 1$) as a function of transverse kinetic energy ($m_T - m_0$) for different centrality classes. Error bars (not visible for many data points) indicate statistical uncertainties and brackets depict systematic uncertainties. Global systematic uncertainties in B.R. are not plotted. Solid and dashed black lines depict exponential function fits and the dot-dashed line depicts a power-law function fit to the spectrum in the 60–80% centrality bin.

at RHIC energies. However, the cold nuclear matter (CNM) effect including shadowing could also play an important role. In addition, hadronization through coalescence has been suggested to potentially modify the charm quark distribution in various charm hadron states which may lead to the reduction in the observed D^0 yields in Au + Au collisions [33] (as seen in Fig. 24). For instance, hadronization through coalescence can lead to an enhancement of the charmed baryon Λ_c^+ yield relative to D^0 yield [34–36] and, together with the strangeness enhancement in the hot QCD medium, can also lead to an enhancement in the charmed strange meson D_s^+ yield relative to D^0 [35–37]. Therefore, determination of the total charm quark yield in heavy-ion collisions will require measurements of other charm hadron states over a broad momentum range.

B. Collectivity

1. m_T spectra

Transverse mass spectra can be used to study the collectivity of produced hadrons in heavy-ion collisions. Figure 25 shows the D^0 invariant yield at mid-rapidity ($|y| < 1$) as a function of transverse kinetic energy ($m_T - m_0$) for different centrality classes, where $m_T = \sqrt{p_T^2 + m_0^2}$ and m_0 is the D^0 meson mass at rest. Solid and dashed black lines depict exponential function fits inspired by thermal models to data in various centrality bins up to $m_T - m_0 < 3 \text{ GeV}/c^2$ using the fit function shown below:

$$\frac{d^2N}{2\pi m_T dm_T dy} = \frac{dN/dy}{2\pi T_{\text{eff}}(m_0 + T_{\text{eff}})} e^{-(m_T - m_0)/T_{\text{eff}}}. \quad (6)$$




FIG. 26. Slope parameter T_{eff} for different particles in central Au + Au collisions [29,39–41]. The dashed lines depict linear function fits to π , K , p and ϕ , Λ , Ξ^- , Ω^- , D^0 respectively.

Such a method has been often used to analyze the particle spectra and to understand kinetic freeze-out properties from the data in heavy-ion collisions [1,38].

A power-law function [Eq. (7)] is also used to fit the spectrum in the 60–80% centrality bin:

$$\frac{d^2N}{2\pi p_T dp_T dy} = \frac{dN}{dy} \frac{4(n-1)(n-2)}{2\pi(n-3)^2 \langle p_T \rangle^2} \left(1 + \frac{2p_T}{\langle p_T \rangle (n-3)} \right)^{-n}, \quad (7)$$

where dN/dy , $\langle p_T \rangle$, and n are three free parameters.

The power-law function fit describes the 60–80% centrality data well indicating that the D^0 meson production in this peripheral bin is close to the expected feature of perturbative QCD. The D^0 meson spectra in more central collisions can be well described by the exponential function fit for $m_T - m_0 < 3 \text{ GeV}/c^2$ suggesting the D^0 mesons have gained collective motion in the medium evolution in these collisions.

The obtained slope parameter T_{eff} for D^0 mesons is compared to other light and strange hadrons measured at RHIC. Figure 26 summarizes the slope parameter T_{eff} for various identified hadrons (π^\pm , K^\pm , p/\bar{p} , ϕ , Λ , Ξ^- , Ω , D^0 , and J/ψ) in central Au + Au collisions at $\sqrt{s_{\text{NN}}} = 200 \text{ GeV}$ [29,39–41]. Point-by-point statistical and systematic uncertainties are added as a quadratic sum when performing these fits. All fits are performed up to $m_T - m_0 < 1 \text{ GeV}/c^2$ for π , K , p , $< 2 \text{ GeV}/c^2$ for ϕ , Λ , Ξ , and $< 3 \text{ GeV}/c^2$ for Ω , D^0 , J/ψ , respectively.

The slope parameter T_{eff} in a thermalized medium can be characterized by the random (generally interpreted as a kinetic freeze-out temperature T_{fo}) and collective (radial flow velocity $\langle \beta_T \rangle$) components with a simple relation [1,42,43]:

$$T_{\text{eff}} = T_{\text{fo}} + m_0 \langle \beta_T \rangle^2. \quad (8)$$

Therefore, T_{eff} will show a linear dependence as a function of particle mass m_0 with a slope that can be used to characterize the radial flow collective velocity.

The data points clearly show two different systematic trends. π , K , p data points follow one linear dependence while ϕ , Λ , Ξ^- , Ω^- , D^0 data points follow another linear dependence, as represented by the dashed lines shown in Fig. 26. Particles such as π , K , p gain radial collectivity throughout the whole system evolution, therefore the linear dependence exhibits a larger slope. On the other hand the linear dependence of ϕ , Λ , Ξ^- , Ω^- , D^0 data points shows a smaller slope indicating these particles may freeze-out from the system earlier, and therefore receive less radial collectivity.

2. Blast-wave fit

The blast-wave (BW) model is extensively used to study the particle kinetic freeze-out properties [29,44]. Assuming a hard-sphere uniform particle source with a kinetic freeze-out temperature T_{kin} and a transverse radial flow velocity β , the particle transverse momentum spectral shape is given by [45]

$$\frac{dN}{p_T dp_T} = \frac{dN}{m_T dm_T} \propto \int_0^R r dr m_T I_0 \left(\frac{p_T \sinh \rho}{T_{\text{kin}}} \right) K_1 \left(\frac{m_T \cosh \rho}{T_{\text{kin}}} \right), \quad (9)$$

where $\rho = \tanh^{-1} \beta$, and I_0 and K_1 are the modified Bessel functions. The flow velocity profile is taken as

$$\beta = \beta_s \left(\frac{r}{R} \right)^n, \quad (10)$$

where β_s is the maximum velocity at the surface and r/R is the relative radial position in the thermal source. The choice of R only affects the overall spectrum magnitude while the spectrum shape constrains the three free parameters T_{kin} , $\langle \beta \rangle = 2/(2+n)\beta_s$, and n .

In the modified Tsallis blast-wave (TBW) model, an additional parameter q is introduced to account for the nonequilibrium feature in a system [46]. The particle transverse momentum spectral shape is then described by

$$\frac{dN}{m_T dm_T} \propto m_T \int_{-Y}^{+Y} \cosh(y) dy \int_{-\pi}^{+\pi} d\phi \int_0^R r dr \times \left(1 + \frac{q-1}{T_{\text{kin}}} [m_T \cosh(y) \cosh(\rho) - p_T \sinh(\rho) \cos(\phi)] \right)^{-1/(q-1)}. \quad (11)$$

When $q - 1$ approaches zero, the TBW function returns to the regular BW function shown in Eq. (9).

Figure 27 shows the blast-wave and Tsallis blast-wave fits to the data in different centrality bins. The parameter n in these fits is fixed to be 1 due to the limited number of data points and is inspired by the fit result for light-flavor hadrons (π , K , and p) [46]. The p_T range in the BW fits is restricted to be less than $3m_0$ where m_0 is the rest mass of D^0 mesons.




FIG. 27. D^0 invariant yield at mid-rapidity ($|y| < 1$) vs transverse momentum for different centrality classes. Black and red lines depict blast-wave (BW) and Tsallis blast-wave (TBW) fits for each centrality bin respectively.

Figure 28 summarizes the fit parameters T_{kin} vs $\langle \beta \rangle$ from the blast-wave model fits to different groups of particles: black markers for the simultaneous fit to π , K , p , red markers for the simultaneous fit to ϕ , Ξ^- , and blue markers for the fit to D^0 . The data points for each group of particles represent the fit results from different centrality bins with the most central data point at the largest $\langle \beta \rangle$ value. Similar as in the fit to the m_T spectra, point-by-point statistical and systematic uncertainties are added in quadrature when performing the fit. The fit results for π , K , p are consistent with previously




FIG. 28. Results of T_{kin} vs $\langle \beta \rangle$ from the blast-wave model fits to different groups of particles. The data points for each group of particles present the results from different centrality bins with the most central data point at the largest $\langle \beta \rangle$.

TABLE VI. $\langle\beta\rangle$ and $(q-1)$ from the Tsallis blast-wave fits to the D^0 data in different centralities.

Centrality	$\langle\beta\rangle(c)$	$q-1$
0–10%	0.263 ± 0.018	0.066 ± 0.008
10–20%	0.255 ± 0.022	0.068 ± 0.010
20–40%	0.264 ± 0.015	0.070 ± 0.007
40–60%	0.251 ± 0.023	0.074 ± 0.011
60–80%	0.217 ± 0.037	0.075 ± 0.010

published results [46]. The fit results for multi-strangeness particles ϕ , Ξ^- , and for D^0 show much smaller mean transverse velocity $\langle\beta\rangle$ and larger kinetic freeze-out temperature, suggesting these particles decouple from the system earlier and gain less radial collectivity compared to light hadrons. The resulting T_{kin} parameters for ϕ , Ξ^- and for D^0 are close to the pseudocritical temperature T_c calculated from a lattice QCD calculation at zero baryon chemical potential [47], indicating negligible contribution from the hadronic stage to the observed radial flow of these particles. Therefore, the collectivity that D^0 mesons obtain is mostly through the partonic stage rescatterings in the QGP phase.

Table VI lists the fitting parameters $\langle\beta\rangle$ and $(q-1)$ for the D^0 data in different centralities. Results show a similar trend as the regular BW fit, i.e., the most central data point is located at the largest $\langle\beta\rangle$ value. The $(q-1)$ parameter in TBW, which characterizes the degree of nonequilibrium in a system, is found to be close to zero, indicating that the system is approaching thermalization in these collisions.

C. Nuclear modification factors R_{CP} and R_{AA}

Figure 29 shows the calculated $D^0 R_{\text{CP}}$ [see Eq. (4)] with the 60–80% peripheral bin as the reference for different centrality bins 0–10%, 10–20%, 20–40%, and 40–60%; the results are compared to other light and strange flavor mesons. The grey bands around unity depict the vertex resolution correction uncertainty on the measured D^0 data points, mostly originating from the 60–80% reference spectrum. The dark and light green boxes around unity on the right side indicate the global N_{bin} systematic uncertainties for the 60–80% centrality bin and for the corresponding centrality bin in each panel. The global N_{bin} systematic uncertainties should be applied to the data points of all particles in each panel.

The measured $D^0 R_{\text{CP}}$ in central 0–10% collisions shows a significant suppression at $p_T > 5$ GeV/c. The suppression level is similar to that of light and strange flavor mesons and the R_{CP} suppression gradually decreases when moving from central collisions to mid-central and peripheral collisions. The $D^0 R_{\text{CP}}$ for $p_T < 4$ GeV/c is consistent with no suppression, in contrast to light-flavor hadrons. Comparisons to dynamic model calculations for the $D^0 R_{\text{CP}}$ will be discussed in Sec. VI E.

The precision of the 60–80% centrality spectrum is limited due to the large systematic uncertainty in determining the N_{bin} based on the MC Glauber model. Figure 30 shows the $D^0 R_{\text{CP}}$ for different centralities as a function of p_T with the 40–60% centrality spectrum as the reference. The grey bands around




FIG. 29. $D^0 R_{\text{CP}}$ with the 60–80% spectrum as the reference for different centrality classes in Au + Au collisions compared to that of other light and strange mesons (π^\pm , K_s^0 , and ϕ) [48–50]. The statistical and systematic uncertainties are shown as error bars and brackets on the data points. The grey bands around unity depict the systematic uncertainty due to vertex resolution correction, mostly from the 60–80% reference spectrum. The light and dark green boxes on the right depict the normalization uncertainties in determining the N_{bin} for each centrality (light green) and the 60–80% centrality bin (dark green), respectively.

unity in each panel represent the systematic uncertainties due to the vertex resolution contribution from the 40–60% centrality. The green boxes around unity depict the global N_{bin} systematic uncertainties for the 40–60% centrality bin and for each corresponding centrality bin. As a comparison, R_{CP} of charged pions K_s^0 and ϕ in the corresponding centralities are also plotted in each panel. With much smaller systematic uncertainties, the observations seen before using the 60–80% centrality spectrum as the reference still hold.

Figure 31 shows the calculated $D^0 R_{\text{AA}}$ [see Eq. (1)] with the $p + p$ measurement [20] as the reference for different centrality bins 0–10% (a), 10–40% (b), and 40–80% (c), respectively. The new R_{AA} measurements are also compared to the previous Au + Au measurements using the STAR TPC after the recent correction [12]. The $p + p D^0$ reference




FIG. 30. $D^0 R_{CP}$ with the 40–60% spectrum as the reference for different centrality classes in Au + Au collisions compared to that of other light and strange mesons (π^\pm , K_s^0 , and ϕ) [48–50]. The statistical and systematic uncertainties are shown as error bars and brackets on the data points. The grey bands around unity depict the systematic uncertainty due to vertex resolution correction, mostly from the 40–60% reference spectrum. The light and dark green boxes on the right depict the normalization uncertainties in determining the N_{bin} for each centrality (light green) and the 40–60% centrality bin (dark green), respectively.

spectrum is updated using the latest global analysis of charm fragmentation ratios from [51] and also by taking into account the p_T dependence of the fragmentation ratio between D^0 and $D^{*\pm}$ from PYTHIA. The new measurement with the HFT detector shows a nice agreement with the measurement without the HFT. The brackets on the data points depict the total systematic uncertainty dominated by the uncertainty in the $p + p$ reference spectrum. The first two and last two data points are empty circles indicating those are calculated with an extrapolated $p + p$ reference. The light and dark green boxes around unity on the right side indicate the global N_{bin} systematic uncertainties for the corresponding centrality bin in each panel and the total inelastic cross section uncertainty in $p + p$ collisions.

The measured $D^0 R_{AA}$ in central (0–10%) and mid-central (10–40%) collisions show a significant suppression at the high p_T range which reaffirms the strong interactions between charm quarks and the medium, while the new Au + Au data points from this analysis contain much improved precision. Figure 32 shows the $D^0 R_{AA}$ in the 0–10% most central




FIG. 31. $D^0 R_{AA}$ in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV for 0–10% (a), 10–40% (b), and 40–80% (c) centrality bins, respectively. The first two and last two data points are presented as empty circles, indicating that the $p + p$ reference is extrapolated into these p_T ranges. The statistical and systematic uncertainties are shown as error bars and brackets on the data points. The light and dark green boxes on the right depict the normalization uncertainties in determining the N_{bin} in Au + Au collisions and the total inelastic cross section in $p + p$ collisions, respectively.

collisions compared to that of (a) average D meson from ALICE and (b) charged hadrons from ALICE and π^\pm from STAR [10,55,56]. The $D^0 R_{AA}$ from this measurement is comparable to that from the LHC measurements in Pb + Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV despite the large energy difference between these measurements. The comparison to that of light hadrons shows a similar suppression at high p_T , while in the intermediate range, D^0 mesons seem to be less suppressed. From the low to intermediate p_T region, the $D^0 R_{AA}$ in the central 0–10% collisions shows a characteristic bump structure that is consistent with the expectation from model predictions that charm quarks gain sizable collective motion during the medium evolution. The large uncertainty in the $p + p$ baseline needs to be further reduced before making more quantitative conclusions.

D. \bar{D}^0 and D^0 spectra and double ratio

Figure 33 shows the p_T spectra of \bar{D}^0 and D^0 mesons separately in 0–10%, 10–20%, 20–40%, 40–60%, and 60–80% centrality bins. Figure 34 shows the \bar{D}^0/D^0 ratio in various




FIG. 32. D^0 R_{AA} in 0–10% Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV compared to the ALICE D -meson result in 0–10% Pb + Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV (a) and charged hadrons from ALICE and π^\pm from STAR (b). Also shown in panel (a) are the model calculations from the LBT and Duke groups [52–54]. Notations for statistical and systematic uncertainties are the same as in previous figures.

centrality bins. Dashed lines represent constant function fits to the \bar{D}^0/D^0 ratio in each centrality bin by combining the point-by-point statistical and systematic uncertainties. The




FIG. 33. D^0 and \bar{D}^0 invariant yields at mid-rapidity ($|y| < 1$) vs transverse momentum for different centrality classes. Error bars (not visible for many data points) indicate statistical uncertainties and brackets depict systematic uncertainties. Global systematic uncertainties in B.R. and N_{bin} are not plotted. Solid lines depict Levy function fits.




FIG. 34. \bar{D}^0/D^0 invariant yield ratio at mid-rapidity ($|y| < 1$) vs transverse momentum for different centrality classes. Error bars indicate statistical uncertainties and brackets depict systematic uncertainties. Dashed lines depict constant function fits to the \bar{D}^0/D^0 ratios.

\bar{D}^0/D^0 ratio has a small but significant deviation from unity in central and mid-central collisions. Table VII lists the fitted results for the \bar{D}^0/D^0 ratio from various centralities. In the most central collisions, the \bar{D}^0 yield is higher than the D^0 yield by $\approx 4.9\sigma$. The total charm quark and anticharm quark should be conserved since they are created in pairs. A thermal model calculation predicts that the Λ_c^-/Λ_c^+ ratio will be smaller than unity at RHIC due to the finite baryon density [57]. This will then yield more \bar{D}^0 mesons formed than D^0 mesons in Au + Au collisions at RHIC. To verify the total charm quark conservation, one would need precise measure-

TABLE VII. \bar{D}^0/D^0 ratio for various centrality bins obtained from the fit to data in Fig. 34.

Centrality	\bar{D}^0/D^0
0–10%	1.104 ± 0.021
10–20%	1.071 ± 0.019
20–40%	1.060 ± 0.015
40–60%	1.073 ± 0.022
60–80%	0.943 ± 0.039




FIG. 35. $D^0 R_{CP}$ with the 60–80% spectrum as the reference for different centrality classes compared to the LBT model calculations shown by dashed lines [52,53]. Data points shown here are the same as in Fig. 29.

ments of D^+/D^- , D_s^+/D_s^- , as well as Λ_c^+/Λ_c^- ratios in the future.

E. Comparison to models

Over the past several years, there have been rapid developments in the theoretical calculations on the charm hadron production [58,59]. Here we compare our measurements to several recent calculations based on the Duke model and the linearized Boltzmann transport (LBT) model [52–54].

The Duke model [54,60] uses a Langevin stochastic simulation to trace the charm quark propagation inside the QGP medium. Both collisional and radiative energy losses are included in the calculation and charm quarks are hadronized via a hybrid approach combining both coalescence and fragmentation mechanisms. The bulk medium is simulated using a viscous hydrodynamic evolution followed by a hadronic cascade evolution using the UrQMD model [61]. The charm quark interaction with the medium is characterized using a temperature- and momentum-dependent diffusion coefficient. The medium parameters have been constrained via a statistical




FIG. 36. $D^0 R_{CP}$ with the 40–60% spectrum as the reference for different centrality classes compared to model calculations from LBT (black dotted lines) and the Duke (blue dashed lines) groups [52–54]. Data points shown here are the same as in Fig. 30.

Bayesian analysis by fitting the previous experimental data of R_{AA} and v_2 of light, strange, and charm hadrons [54]. The extracted charm quark spatial diffusion coefficient at zero momentum $2\pi T D_s|_{p=0}$ is about 1–3 near T_c and exhibits a positive slope for its temperature dependence above T_c .

The linearized Boltzmann transport (LBT) calculation [52] extends the LBT approach developed before to include both light and heavy flavor parton evolution in the QGP medium. The transport calculation includes all $2 \rightarrow 2$ elastic scattering processes for collisional energy loss and the higher-twist formalism for medium induced radiative energy loss. It uses the same hybrid approach as in the Duke model for charm quark hadronization. The heavy quark transport is coupled with a 3D viscous hydrodynamic evolution which is tuned for light flavor hadron data. The charm quark spatial diffusion coefficient is estimated via the equation $2\pi T D_s = 8\pi/\hat{q}$ (\hat{q} is the quark transport coefficient due to elastic scatterings) at parton momentum $p = 10$ GeV/c. The $2\pi T D_s$ is ≈ 3 at T_c and increases to ≈ 6 at $T = 500$ MeV [53].

Figures 35 and 36 show the measured $D^0 R_{CP}$ compared to the Duke and LBT model calculations with the 60–80% and 40–60% reference spectra respectively. The R_{CP} curves from these models are calculated based on the D^0 spectra provided by each group [52–54]. The Duke model did not calculate the spectra in the 60–80% centrality bin due to a concern about the viscous hydrodynamic implementation. In Fig. 32

for the most central collisions, there are also calculations for the $D^0 R_{AA}$ from the Duke and LBT groups, respectively. These two models also have the predictions for the $D^0 v_2$ measurements for Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV [16]. Both model calculations match our new measured R_{CP} data well. The much improved precision of these new measurements are expected to further constrain the theoretical model uncertainties in these calculations.

VII. SUMMARY

In summary, we report the improved measurement of D^0 production invariant yield at mid-rapidity ($|y| < 1$) in Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV with the STAR HFT detector. D^0 invariant yields are presented as a function of p_T in various centrality classes. The p_T -integrated D^0 production cross section per nucleon-nucleon collisions in mid-central and central Au + Au collisions seem to be smaller than that measured in $p + p$ collisions by 1.5σ , indicating that CNM effects and/or hadronization through quark coalescence may play an important role in Au + Au collisions. This calls for precise measurements of D^0 production in $p/d + A$ collisions to understand the CNM effects as well as other charm hadron states in heavy-ion collisions to better constrain the total charm quark yield.

The \bar{D}^0 yield is observed to be higher than the D^0 yield in the most central collisions, by $\approx 4.9\sigma$ on average. This is potentially consistent with the expectation, due to the finite baryon density of the system at RHIC, that the Λ_c^-/Λ_c^+ ratio should be smaller than unity, which would result in larger \bar{D}^0 yield than D^0 .

The D^0 spectra at low p_T and m_T regions are fit by the exponential function and the (Tsallis) blast-wave model to study the D^0 meson radial collectivity. The slope parameter extracted from the exponential function fit for D^0 mesons follows the same linearly increasing trend vs particle mass as ϕ , Λ , Ξ^- , Ω^- particles, but is different from the trend of π , K , and p particles. The extracted kinetic freeze-out temperature and transverse velocity from the blast-wave model fit are comparable to the fit results of ϕ , Ξ^- multi-strange-quark hadrons, but different from those of π , K , and p . These observations suggest that D^0 hadrons show a radial collective behavior with the medium, but freeze-out from the system earlier and gain less radial collectivity compared to π , K , and p particles. This observation is consistent with collective behavior observed in v_2 measurements. The fit results also suggest that D^0 mesons

have similar kinetic freeze-out properties as multi-strange-quark hadrons ϕ , Ξ^- .

The nuclear modification factors R_{CP} of D^0 mesons are presented with both 60–80% and 40–60% centrality spectra as the reference, respectively. The $D^0 R_{CP}$ is significantly suppressed at high p_T and the suppression level is comparable to that of light hadrons at $p_T > 5$ GeV/ c , reaffirming our previous observation [12]. This indicates that charm quarks lose significant energy when traversing through the hot QCD medium. The $D^0 R_{CP}$ is above the light hadron R_{CP} at low p_T . We compare our $D^0 R_{CP}$ measurements to two recent theoretical model calculations from LBT and the Duke groups. These two models have the $2\pi TD_s$ value around 1–3 near T_c and agree with our new R_{CP} measurements. The nuclear modification factor R_{AA} of D^0 mesons in 0–10% central Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV is comparable to that from the ALICE measurement in Pb + Pb at $\sqrt{s_{NN}} = 2.76$ TeV. At $p_T < 5$ GeV/ c , the $D^0 R_{AA}$ shows a characteristic bump structure. Model calculations that predict sizable collective motion for charm quarks during the medium evolution can qualitatively describe our data. We expect that the new data points with much improved precision can be used in the future to further constrain our understanding of the charm-medium interactions as well as to better determine the medium transport parameter.

ACKNOWLEDGMENTS

We thank the RHIC Operations Group and RCF at BNL, the NERSC Center at LBNL, and the Open Science Grid consortium for providing resources and support. This work was supported in part by the Office of Nuclear Physics within the US DOE Office of Science, the US National Science Foundation, the Ministry of Education and Science of the Russian Federation, National Natural Science Foundation of China, Chinese Academy of Science, the Ministry of Science and Technology of China and the Chinese Ministry of Education, the National Research Foundation of Korea, GA and MSMT of the Czech Republic, Department of Atomic Energy and Department of Science and Technology of the Government of India; the National Science Centre of Poland, National Research Foundation, the Ministry of Science, Education and Sports of the Republic of Croatia, RosAtom of Russia and German Bundesministerium für Bildung, Wissenschaft, Forschung und Technologie (BMBF), and the Helmholtz Association.

[1] J. Adams *et al.* (STAR Collaboration), *Nucl. Phys. A* **757**, 102 (2005).
 [2] K. Adcox *et al.* (PHENIX Collaboration), *Nucl. Phys. A* **757**, 184 (2005).
 [3] B. Muller, J. Schukraft, and B. Wyslouch, *Annu. Rev. Nucl. Part. Sci.* **62**, 361 (2012).
 [4] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **116**, 062301 (2016).

[5] B. B. Abelev *et al.* (ALICE Collaboration), *J. High Energy Phys.* **06** (2015) 190.
 [6] Z. Lin and M. Gyulassy, *Phys. Rev. C* **51**, 2177 (1995).
 [7] M. Cacciari, P. Nason, and R. Vogt, *Phys. Rev. Lett.* **95**, 122001 (2005).
 [8] G. D. Moore and D. Teaney, *Phys. Rev. C* **71**, 064904 (2005).
 [9] B. Abelev *et al.* (ALICE Collaboration), *J. High Energy Phys.* **09** (2012) 112.

- [10] J. Adam *et al.* (ALICE Collaboration), *J. High Energy Phys.* **03** (2016) 081.
- [11] A. M. Sirunyan *et al.* (CMS Collaboration), *Phys. Lett. B* **782**, 474 (2018).
- [12] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **113**, 142301 (2014); **121**, 229901(E) (2018).
- [13] B. Abelev *et al.* (ALICE Collaboration), *Phys. Rev. Lett.* **111**, 102301 (2013).
- [14] B. Abelev *et al.* (ALICE Collaboration), *Phys. Rev. C* **90**, 034904 (2014).
- [15] A. M. Sirunyan *et al.* (CMS Collaboration), *Phys. Rev. Lett.* **120**, 202301 (2018).
- [16] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **118**, 212301 (2017).
- [17] G. Contin *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **907**, 60 (2018).
- [18] M. Anderson *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **499**, 659 (2003).
- [19] W. J. Llope *et al.* (STAR Collaboration), *Nucl. Instrum. Methods Phys. Res., Sect. A* **661**, S110 (2012).
- [20] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. D* **86**, 072013 (2012).
- [21] W. J. Llope *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **522**, 252 (2004).
- [22] D. Smirnov, J. Lauret, V. Perevoztchikov, G. Van Buren, and J. Webb, *J. Phys.: Conf. Ser.* **898**, 042058 (2017).
- [23] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. C* **88**, 014902 (2013).
- [24] R. E. Kalman, *J. Basic Eng.* **82**, 35 (1960).
- [25] M. Tanabashi *et al.* (Particle Data Group), *Phys. Rev. D* **98**, 030001 (2018).
- [26] H. Voss *et al.*, PoS **ACAT**, 040 (2007).
- [27] R. Brun *et al.*, Report No. CERN-W5013, CERN-W-5013, W5013, W-5013 (1994).
- [28] M. Gyulassy and X.-N. Wang, *Comput. Phys. Commun.* **83**, 307 (1994).
- [29] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **92**, 112301 (2004).
- [30] A. Fasso *et al.*, in *Proceedings of the IVth International Conference on Calorimetry in High Energy Physics, La Biodola (Italy)*, (1993), Vol. C9309194, pp. 493–502.
- [31] M. Shao, O. Barannikova, X. Dong, Y. Fisyak, L. Ruan, P. Sorensen, and Z. Xu, *Nucl. Instrum. Methods Phys. Res., Sect. A* **558**, 419 (2006).
- [32] Y. Xu *et al.*, *Nucl. Instrum. Methods Phys. Res., Sect. A* **614**, 28 (2010).
- [33] V. Greco, C. Ko, and R. Rapp, *Phys. Lett. B* **595**, 202 (2004).
- [34] Y. Oh, C. M. Ko, S. H. Lee, and S. Yasui, *Phys. Rev. C* **79**, 044905 (2009).
- [35] J. Zhao, S. Shi, N. Xu, and P. Zhuang, [arXiv:1805.10858](https://arxiv.org/abs/1805.10858).
- [36] S. Plumari, V. Minissale, S. K. Das, G. Coci, and V. Greco, *Eur. Phys. J. C* **78**, 348 (2018).
- [37] M. He, R. J. Fries, and R. Rapp, *Phys. Rev. Lett.* **110**, 112301 (2013).
- [38] M. Kaneta, Ph.D. thesis, Hiroshima University, 1999.
- [39] B. I. Abelev *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **99**, 112301 (2007).
- [40] J. Adams *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **98**, 062301 (2007).
- [41] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. C* **90**, 024906 (2014).
- [42] T. Csorgo and B. Lorstad, *Phys. Rev. C* **54**, 1390 (1996).
- [43] P. F. Kolb and U. W. Heinz, *Quark Gluon Plasma* **3**, 634 (2003).
- [44] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Rev. C* **96**, 044904 (2017).
- [45] E. Schnedermann, J. Sollfrank, and U. Heinz, *Phys. Rev. C* **48**, 2462 (1993).
- [46] Z. Tang, Y. Xu, L. Ruan, G. van Buren, F. Wang, and Z. Xu, *Phys. Rev. C* **79**, 051901(R) (2009).
- [47] A. Bazavov *et al.*, *Phys. Rev. D* **85**, 054503 (2012).
- [48] B. I. Abelev *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **97**, 152301 (2006).
- [49] B. I. Abelev *et al.* (STAR Collaboration), *Phys. Rev. C* **79**, 064903 (2009).
- [50] G. Agakishiev *et al.* (STAR Collaboration), *Phys. Rev. Lett.* **108**, 072301 (2012).
- [51] M. Lisovsky, A. Verbytskyi, and O. Zenaiev, *Eur. Phys. J. C* **76**, 397 (2016).
- [52] S. Cao, T. Luo, G. Y. Qin, and X. N. Wang, *Phys. Rev. C* **94**, 014909 (2016).
- [53] S. Cao (private communication).
- [54] Y. Xu, J. E. Bernhard, S. A. Bass, M. Nahrgang, and S. Cao, *Phys. Rev. C* **97**, 014907 (2018).
- [55] B. Abelev *et al.* (ALICE Collaboration), *Phys. Lett. B* **720**, 52 (2013).
- [56] B. Abelev *et al.* (STAR Collaboration), *Phys. Lett. B* **655**, 104 (2007).
- [57] A. Andronic, P. Braun-Munzinger, K. Redlich, and J. Stachel, *Phys. Lett. B* **571**, 36 (2003).
- [58] R. Rapp *et al.*, *Nucl. Phys. A* **979**, 21 (2018).
- [59] S. Cao *et al.*, [arXiv:1809.07894](https://arxiv.org/abs/1809.07894).
- [60] S. Cao, G. Y. Qin, and S. A. Bass, *Phys. Rev. C* **92**, 024907 (2015).
- [61] M. Bleicher *et al.*, *J. Phys. G* **25**, 1859 (1999).