

Prospects for vectorlike leptons at future proton-proton colliders

Prudhvi N. Bhattiprolu and Stephen P. Martin

Department of Physics, Northern Illinois University, DeKalb, Illinois 60115, USA



(Received 8 May 2019; published 19 July 2019)

Vectorlike leptons are an intriguing possibility for physics beyond the Standard Model. We study the reach for discovering (at 5σ significance) or excluding (at 95% confidence) models of charged vectorlike leptons that mix predominantly with the tau, using multilepton signatures at various future proton-proton collider options: a high-luminosity LHC with $\sqrt{s} = 14$ TeV, a high-energy LHC with $\sqrt{s} = 27$ TeV, and possible new longer-tunnel colliders with $\sqrt{s} = 70$ or 100 TeV. For weak-isodoublet vectorlike leptons, we estimate that a 27 TeV high-energy LHC with 15 ab^{-1} could exclude masses up to about 2300 GeV, or discover them if the mass is less than about 1700 GeV, while a 100 TeV collider with 30 ab^{-1} could exclude masses up to about 5750 GeV or make a discovery if the mass is less than about 4000 GeV. However, we find that weak-isosinglet vectorlike leptons present a much more difficult challenge, with some reach for exclusion, but not for discovery at any of the collider options considered.

DOI: [10.1103/PhysRevD.100.015033](https://doi.org/10.1103/PhysRevD.100.015033)

I. INTRODUCTION

The Large Hadron Collider (LHC) has conducted searches for new physics beyond the Standard Model, at proton-proton center of mass energies up to $\sqrt{s} = 13$ TeV. Following the discovery [1] of the Higgs scalar boson with mass near $M_h = 125$ GeV, the present evidence from the LHC is consistent with the minimal version of the Standard Model, with direct and indirect limits on new particles extending well into the TeV range. However, there are many candidate new physics models that obey decoupling as the mass scale of new physics is raised. These include supersymmetric theories, where agreement with the Standard Model gets better as the scale of supersymmetry breaking is raised. An example of a nondecoupling theory is a new chiral fourth family of fermions. Because of the necessity of generating the large chiral fermion masses through Yukawa couplings and the Higgs mechanism, decoupling is violated, which causes significant contributions to precision electroweak observables as well as Higgs boson production and decay rates, in conflict [2] with the observations.

New fermions are still allowed, because they will obey decoupling, if they obtain their mass entirely or mostly as bare electroweak-singlet terms in the Lagrangian, rather than from the Yukawa couplings to the Higgs field. This implies that the new fermions are vectorlike (the antonym

of chiral). It is a notable common feature of many new physics theories that are motivated for other reasons (such as the need for a cosmological and astrophysical dark matter candidate, or to address the hierarchy problem) that they often contain vectorlike fermions. For example, in supersymmetry [3], the Higgsinos have the same gauge quantum numbers as would a vectorlike pair of weak-isodoublet leptons. It is plausible that additional vectorlike fermion masses are at the TeV scale, because whatever mechanism is responsible for the Higgsino masses can also be applied to the vectorlike fermion masses, which have the same structure. Vectorlike fermions can therefore be added on to the minimal version of supersymmetry, with one possible benefit that they can contribute to the lightest Higgs boson mass [4–18], which allows a lower supersymmetry breaking scale consistent with $M_h = 125$ GeV. For example, adding vectorlike quarks to the simplest models of gauge-mediated supersymmetry breaking allows [15–18] them to be still discoverable at the LHC. However, even ignoring such motivations, it is also worthwhile to consider vectorlike fermions merely on the merits of being a simple and consistent candidate extension of the Standard Model.

For vectorlike quarks, the LHC pair production cross section is determined from QCD and is therefore large and essentially model independent.¹ The searches for vectorlike quarks are consequently quite advanced, with current LHC limits (depending on the specific decay modes) on

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

¹Single production of vectorlike quarks is also subject to LHC limits, but in a much more model-dependent way.

pair-produced vectorlike quarks that can be found in Refs. [19,20].

In contrast, vectorlike leptons (VLLs) [21–71] are pair-produced by s -channel electroweak vector boson diagrams, leading to cross sections that are much smaller and dependent on the choices of weak isospin and weak hypercharge of the new states.² For this reason, until recently the LHC detector collaborations had not put limits on VLLs beyond those that could be inferred from the CERN large electron-positron (LEP) e^-e^+ collider, which excluded [74] masses up to 101.2 GeV. There have been several phenomenological studies that looked into the LHC discovery and exclusion capabilities. In Ref. [50], search strategies and capabilities were discussed for the optimistic case that the VLLs decayed mostly to muons, and an ATLAS search was conducted for that case in [52]. In Ref. [57], the more pessimistic case of decays to tau-rich final states was considered, and it was argued that even with the (then current, but now old) data at $\sqrt{s} = 8$ TeV it should be possible to exclude such $SU(2)_L$ -doublet VLLs up to about $M_{\ell'} = 275$ GeV, but not $SU(2)_L$ -singlet charged VLLs, which have a much smaller production cross section and unfavorable branching ratios. Since then, the CMS collaboration has published [70,71] the results of dedicated searches for doublet VLLs, based on 41.4 and 77.4 fb^{-1} data samples at $\sqrt{s} = 13$ TeV. Aided in part by a deficit of events in the signal regions compared to the background expectation, CMS was able to obtain an exclusion for $120 < M_{\ell'} < 790$ GeV for a pure $SU(2)_L$ -doublet VLL pair that mixes with and decays to tau leptons.

In this paper, we will consider the prospects for exclusion or discovery of VLLs at planned future proton-proton colliders. While there are a wide variety of possible VLL models, we will consider as benchmark models two simple minimal cases as defined in Ref. [57], a “singlet VLL” and a “doublet VLL” model, with the mixing to the Standard Model leptons assumed to be entirely with the tau and small. One advantage of these models as benchmarks is that the pair-production cross sections are uniquely determined as a function of the mass, as discussed in Ref. [57] and reviewed in the next section. As was the case in Ref. [57], our analysis below shows that the minimal singlet VLL model is very difficult to probe at proton-proton colliders. We will therefore also consider a class of non-minimal models which are similar to the minimal singlet VLL model and have the same production cross section, but have different branching ratios for the lightest charged new lepton.

We will consider the following proton-proton collider options:

- (i) a high-luminosity LHC (HL-LHC), defined as 3 ab^{-1} of proton-proton collisions at $\sqrt{s} = 14$ TeV,
- (ii) a high-energy LHC (HE-LHC), defined as 15 ab^{-1} at $\sqrt{s} = 27$ TeV,
- (iii) a very high-energy pp collider, defined as up to 30 ab^{-1} at either $\sqrt{s} = 70$ or 100 TeV, which could be realized as, e.g., a CERN hadron-hadron future circular collider [75] or a super proton-proton collider [76] in China.

The physics potential for HL-LHC and HE-LHC has been studied for various other beyond-the-Standard-Model scenarios in Ref. [77] and references therein.

For reach estimates, we will use a simple cut-based counting experiment criteria, where the significance

$$Z = \sqrt{2}\text{erfc}^{-1}(2p) \quad (1.1)$$

of an experimental result is related to the probability p of observing, in an ensemble of many repeated experiments, a result of equal or greater incompatibility with the null hypothesis. Let the number of background events be Poisson distributed with mean b , with the variance in b (corresponding intuitively to an uncertainty in the expected number of background events) given by Δ_b , and suppose the number of signal events is Poisson distributed with predicted mean s . Then the median expected significance for discovery (where the null hypothesis is background only, and the signal is assumed to be present in the data) is approximately [78–81]

$$Z_{\text{disc}} = \left[2 \left((s+b) \ln \left[\frac{(s+b)(b+\Delta_b^2)}{b^2 + (s+b)\Delta_b^2} \right] - \frac{b^2}{\Delta_b^2} \ln \left[1 + \frac{s\Delta_b^2}{b(b+\Delta_b^2)} \right] \right) \right]^{1/2}. \quad (1.2)$$

The median expected significance for exclusion (where the null hypothesis is now the background + signal model, but the signal is assumed to be actually absent in the data) can be approximated by [57]

$$Z_{\text{exc}} = \left\{ 2 \left[s - b \ln \left(\frac{b+s+x}{2b} \right) - \frac{b^2}{\Delta_b^2} \ln \left(\frac{b-s+x}{2b} \right) \right] - (b+s-x)(1+b/\Delta_b^2) \right\}^{1/2}, \quad (1.3)$$

where

$$x = \sqrt{(s+b)^2 - 4s\Delta_b^2/(1+\Delta_b^2/b)}. \quad (1.4)$$

In the idealized limit of a perfectly known background prediction, $\Delta_b = 0$, these would reduce to

$$Z_{\text{disc}} = \sqrt{2[(s+b) \ln(1+s/b) - s]}, \quad (1.5)$$

²Here we concentrate on models with new charged vectorlike leptons. Signatures from models with neutrino masses from electroweak-singlet states at the TeV scale are reviewed in [72,73], for example.

$$Z_{\text{exc}} = \sqrt{2[s - b \ln(1 + s/b)]}. \quad (1.6)$$

In the further limit of large b , these both approach $Z_{\text{disc}} = Z_{\text{exc}} = s/\sqrt{b}$. In the following, we will use as a criteria for expected five-sigma discovery that s is at least 10 and Eq. (1.2) should exceed $Z_{\text{disc}} > 5$, and for an expected 95% exclusion that Eq. (1.3) should exceed $Z_{\text{exc}} > 1.645$ corresponding to $p = 0.05$. We will use the somewhat arbitrary (since the capabilities of the detectors are unknown at present) choices $\Delta_b/b = 0.1, 0.2$, and 0.5 , corresponding to a 10%, 20%, and 50% uncertainty in the background. We have also assumed that b is always at least one event, to be conservative.

II. MINIMAL MODELS FOR PRODUCTION AND DECAY OF VECTORLIKE LEPTONS

Following the terminology and definitions in Ref. [57], the singlet VLL model contains the Standard Model fields and interactions plus an $SU(2)_L$ -singlet charged VLL τ'^- and its antiparticle τ'^+ , which consist of two-component left-handed fermions transforming under $SU(3)_c \times SU(2)_L \times U(1)_Y$ as

$$\tau' + \bar{\tau}' = (\mathbf{1}, \mathbf{1}, -1) + (\mathbf{1}, \mathbf{1}, +1). \quad (2.1)$$

The doublet VLL model contains a new charged lepton and its neutral heavy Dirac neutrino partner, realized as two-component left-handed fermions transforming as

$$\begin{aligned} L + \bar{L} &= \begin{pmatrix} \nu' \\ \tau' \end{pmatrix} + \begin{pmatrix} \bar{\nu}' \\ \bar{\tau}' \end{pmatrix} \\ &= \left(\mathbf{1}, \mathbf{2}, -\frac{1}{2}\right) + \left(\mathbf{1}, \mathbf{2}, +\frac{1}{2}\right). \end{aligned} \quad (2.2)$$

In both models, a single weak-isosinglet bare fermion mass parameter M is mostly responsible for the new fermion masses, with a small Yukawa couplings ϵ to the Higgs field providing the mixing mass with the Standard Model τ lepton, which also has its own Yukawa coupling y_τ to the Higgs field. The charged fermion mass matrix for the τ, τ' system in each case then can be written in the form

$$\mathcal{M} = \begin{pmatrix} y_\tau v & 0 \\ \epsilon v & M \end{pmatrix}, \quad (2.3)$$

where $v \approx 174$ GeV is the Standard Model Higgs vacuum expectation value. The tree-level mass eigenvalues, obtained from the square roots of the eigenvalues of $\mathcal{M}^\dagger \mathcal{M}$ after expanding for $y_\tau v, \epsilon v \ll M$, are

$$M_\tau = y_\tau v (1 - \epsilon^2 v^2 / 2M^2 + \dots), \quad (2.4)$$

$$M_{\tau'} = M (1 + \epsilon^2 v^2 / 2M^2 + \dots), \quad (2.5)$$

where the ellipses in both cases represent terms suppressed by $\epsilon^4 v^4 / M^4$ or $\epsilon^2 y_\tau^2 v^4 / M^4$, while the tree-level neutral VLL mass in the doublet model is simply $M_{\nu'} = M$. There is also a one-loop radiative correction to the physical mass splitting $M_{\nu'} - M_{\tau'}$, but it is also small, only of order of a few hundred MeV [23].

In the special case of no mixing with the tau lepton, $\epsilon = 0$, the lightest VLL would be absolutely stable due to a conserved global symmetry. Here, we assume instead that ϵ is small enough to be treated as a tiny perturbation in the mass matrix, but that it exceeds about 2×10^{-7} , which is large enough to allow [57] the VLLs to decay promptly³ on collider detector length scales to Standard Model states, with widths that dominate over the competing mode $\tau'^- \rightarrow \pi^- \nu'$ [23] in the doublet VLL case. The fermion mass eigenstates then consist of, besides the usual τ^+, τ^-, ν_τ and the rest of the Standard Model states, a charged Dirac pair τ'^-, τ'^+ in both models, and a neutral Dirac pair $\nu', \bar{\nu}'$ in the doublet VLL model only. Due to the small size of ϵ and the one-loop radiative splitting, we can take $M_{\nu'} \approx M_{\tau'}$ for the purposes of collider simulations.

The collider pair-production modes are

$$pp \rightarrow \gamma^*, \quad Z^* \rightarrow \tau'^- \tau'^+, \quad (2.6)$$

$$pp \rightarrow Z^* \rightarrow \nu' \bar{\nu}', \quad (2.7)$$

$$pp \rightarrow W^{*-} \rightarrow \tau'^- \bar{\nu}', \quad (2.8)$$

$$pp \rightarrow W^{*+} \rightarrow \tau'^+ \nu', \quad (2.9)$$

involving couplings to the γ, Z , and W^\pm vector bosons, which are listed in Ref. [57]. Of course, only the first of these processes occurs for the singlet VLL model.

In Fig. 1, we show the total pair-production cross sections at a proton-proton collider as a function of $M_{\tau'} = M_{\nu'}$, for the choices $\sqrt{s} = 8, 13, 14, 27, 70, 100$ TeV. Note that the doublet VLL model has a considerably larger total cross section, which is partly responsible for the much better search prospects as found in Ref. [57] and below in the present paper. For example, at $M_{\tau'} = 1000$ GeV, the total cross section is about 12.5 times larger in the doublet VLL model than for the singlet VLL model, nearly independent of the proton-proton collision energy over the range from 14 to 100 TeV. In the doublet VLL model, the total contribution is dominated by the W -boson-mediated $\tau'^- \bar{\nu}'$ and $\tau'^+ \nu'$ final states. However, even if one restricts attention to only the $\tau'^- \tau'^+$ final state common to both models, the doublet

³If the τ' in the singlet VLL model is quasistable, then search strategies based on time of flight and ionization rate should apply, as in the ATLAS search in Ref. [82]. Assuming that the τ' interacts similar to a chargino, we infer from the cross section limit in Fig. 10 of [82] that its mass should exceed about 750 GeV. This does not apply to the doublet VLL model, where the τ' can decay to ν' inside the detector.




FIG. 1. The total pair-production cross sections for $\tau^-\tau^+$ in the singlet VLL model (left) and for the sum of $\tau^-\tau^+$ and $\nu'\bar{\nu}'$ and $\tau^-\bar{\nu}'$ and $\tau^+\nu'$ in the doublet VLL model (right), for proton-proton collisions at $\sqrt{s} = 8, 13, 14, 27, 70, 100$ TeV (bottom to top).




FIG. 2. The individual pair-production cross sections in the singlet VLL model for $\tau^-\tau^+$ (solid red line) and in the doublet VLL model for $\tau^-\tau^+$ (dashed blue line) and $\nu'\bar{\nu}'$ (dotted blue line) and $\tau^-\bar{\nu}'$ plus $\tau^+\nu'$ (dot-dashed blue line) and total (solid blue line), for proton-proton collisions at $\sqrt{s} = 27$ TeV.

VLL model has a significantly larger cross section. This is illustrated in Fig. 2, which compares the individual cross sections for each of the final states at $\sqrt{s} = 27$ TeV. (Results are similar at other values of \sqrt{s} .)

The total pair-production cross sections for these models are also shown in Fig. 3 as a function of \sqrt{s} , for various choices of $M_{\tau'} = M_{\nu'}$. Clearly, the cross sections are getting larger at higher collider energies. In the doublet VLL model, one obtains roughly the same cross section for $M_{\tau'} = 4000$ GeV at $\sqrt{s} = 100$ TeV as one would have for $M_{\tau'} = 1870$ GeV at $\sqrt{s} = 27$ TeV or for $M_{\tau'} = 1220$ GeV at $\sqrt{s} = 14$ TeV.

As for the decays, the lepton-flavor conserving charged current process $\tau' \rightarrow \nu' \pi^-$ is kinematically allowed in the doublet VLL model, but has a width smaller than the direct lepton-flavor-violating decays to Standard Model fermions as long as $\epsilon \gtrsim 2 \times 10^{-7}$ [57] and is therefore neglected. In the doublet VLL model, the neutral VLLs decay 100% of the time to a W boson and an ordinary tau lepton, with branching ratios (BR)




FIG. 3. The total pair-production cross sections for $\tau^-\tau^+$ in the singlet VLL model (left) and for the sum of $\tau^-\tau^+$ and $\nu'\bar{\nu}'$ and $\tau^-\bar{\nu}'$ and $\tau^+\nu'$ in the doublet VLL model (right), as a function of \sqrt{s} , for various values of $M_{\tau'} = M_{\nu'}$ as labeled.

$$\text{BR}(\nu' \rightarrow W^+ \tau^-) = \text{BR}(\bar{\nu}' \rightarrow W^- \tau^+) = 1. \quad (2.10)$$

This reflects our assumptions that there is no ν' mixing with the left-handed neutrinos of the Standard Model and that the mixing Yukawa coupling ϵ involves the tau lepton, rather than muon or the electron; otherwise the discovery and exclusion strategies would be much easier, due to the higher detection efficiency and lower fake rates for e, μ compared to τ .

In both models, the charged VLLs can decay into the final states $Z\tau, h\tau$, and $W\nu_\tau$. The decay widths (neglecting the tau lepton mass) are [57]

$$\Gamma(\tau' \rightarrow h\tau) = \frac{\epsilon^2}{64\pi} M_{\tau'} (1 - r_h)^2, \quad (2.11)$$

$$\Gamma(\tau' \rightarrow Z\tau) = \frac{\epsilon^2}{64\pi} M_{\tau'} (1 + 2r_Z)(1 - r_Z)^2, \quad (2.12)$$

in both the singlet and doublet VLL models, while

$$\Gamma(\tau' \rightarrow W\nu) = \begin{cases} \frac{\epsilon^2}{32\pi} M_{\tau'} (1 + 2r_W)(1 - r_W)^2, & \text{(singlet VLL model),} \\ 0, & \text{(doublet VLL model),} \end{cases} \quad (2.13)$$

where $r_X \equiv M_X^2/M_{\tau'}^2$ for each of $X = h, Z, W$. In the decays to Z and W , the factors $(1 + 2r_Z)$ and $(1 + 2r_W)$ can be understood as coming from the longitudinal (1) and transverse ($2r_X$) components of the weak vector bosons. The longitudinal components can in turn be understood as essentially the Goldstone modes that are eaten by the vector bosons to obtain their masses. This illustrates the usual Goldstone equivalence principle, which implies that for the limit of large $M_{\tau'}$ the branching ratios should approach

$$[\text{BR}(\tau' \rightarrow h\tau), \text{BR}(\tau' \rightarrow Z\tau), \text{BR}(\tau' \rightarrow W\nu)] = \begin{cases} [0.25, 0.25, 0.5] & \text{(singlet VLL model),} \\ [0.5, 0.5, 0] & \text{(doublet VLL model).} \end{cases} \quad (2.14)$$

In Fig. 4, we plot the branching ratios for the τ' decays in the two models, as a function of the mass $M_{\tau'}$, showing the asymptotic approach to the Goldstone equivalence limit.

The doublet VLL model therefore has the following final states:

$$ZZ\tau^-\tau^+, \quad Zh\tau^-\tau^+, \quad hh\tau^-\tau^+ \quad (2.15)$$

from τ' pair-production Eq. (2.6),

$$W^+W^-\tau^-\tau^+ \quad (2.16)$$

from ν' pair-production Eq. (2.7), and

$$ZW^\pm\tau^-\tau^+, \quad hW^\pm\tau^-\tau^+ \quad (2.17)$$

from $\tau'\nu'$ production Eqs. (2.8) and (2.9). Besides the three signals in Eq. (2.15), the singlet VLL model also has final states

$$ZW^\pm\tau^\mp + E_T^{\text{miss}}, \quad hW^\pm\tau^\mp + E_T^{\text{miss}}, \quad W^+W^- + E_T^{\text{miss}}, \quad (2.18)$$

with the last being the largest in the minimal singlet VLL model.

With the above branching ratios for the minimal singlet VLL model, the analysis below does not find any reach for any of the signal regions at any of the collider options considered below, due to the low cross section and the unfortunate large branching ratio for $\tau' \rightarrow W\nu$. Therefore, we instead consider a class of nonminimal models which have the feature that the lightest new fermion is still a charged, mostly isosinglet τ' , but with other new fermions




FIG. 4. The branching ratios for $\tau' \rightarrow W\nu$ and $Z\tau$ and $h\tau$, as a function of $M_{\tau'}$, for the singlet VLL model (left) and the doublet VLL model (right), showing the rapid approach to the Goldstone boson equivalence limit for larger masses.

much heavier and therefore not contributing to the production. Thus, the new particle content first accessible to colliders is the same as in the minimal singlet VLL model, but the mixing with the much heavier vectorlike fermions allows the branching ratios of τ' to be essentially arbitrary among the three final states listed above. We therefore present results for modified singlet VLL models, with the same $\tau'^+\tau'^-$ production cross section as the minimal singlet VLL model, but with branching ratios set to the following three subcases:

$$\text{BR}(\tau' \rightarrow h\tau) = 1, \quad \text{“Higgs-philic singlet VLL,”} \quad (2.19)$$

$$\text{BR}(\tau' \rightarrow Z\tau) = 1, \quad \text{“Z-philic singlet VLL,”} \quad (2.20)$$

$$\text{BR}(\tau' \rightarrow h\tau) = \text{BR}(\tau' \rightarrow Z\tau) = 0.5, \quad \text{“W-phobic singlet VLL.”} \quad (2.21)$$

Realizations of these simplified models, from appropriate limits of mixing a vectorlike isosinglet fermion with a much heavier vectorlike isodoublet fermion, are discussed in the Appendix.

For signal simulation, we have input a Lagrangian which governs the production and decay of the doublet VLL and the singlet VLL models (as discussed above) into FEYNRULES v2.3 [83], a MATHEMATICA package, to obtain the Feynman rules as a Universal FEYNRULES Output (UFO) file, which is then imported into MADGRAPH5_AMC@NLO v2.6.5 [84].⁴ We considered Standard Model production of the final states WZ , ZZ , $t\bar{t}Z$, $t\bar{t}W$, hW , hZ , $t\bar{t}h$, WWW , WWZ , ZZW , and ZZZ as the main backgrounds contributing to multilepton final states. We did not include reducible backgrounds such as $W + \text{jets}$, $Z + \text{jets}$, and $WZ + \text{jets}$, where one or more jets fake a tau. While they could be important, we do not have a way of reliably estimating them for unknown future detectors. Therefore it must be kept in mind that if they are large, our projections may be too optimistic.

Both signal and background events were generated using MADGRAPH at leading order (LO). The decay couplings in the (VLL) Lagrangian are left as free parameters in the model file of FEYNRULES, so that one has the flexibility to adjust them during the run time of MADGRAPH. The numerical values of the decay couplings that are used are actually not important (as long as they are not too small or too large), because in each separate MADGRAPH run the decays of VLL were forced into one of the individual channels mentioned above, and later the event samples were normalized using the branching ratios shown in Fig. 4. We calculated the cross sections for signal at LO and

normalized the background cross sections as calculated at next-to-leading order, using MADGRAPH. While using MADGRAPH, we used their default set of parton distribution functions based on NNPDF2.3 set [85]. We used PYTHIA 8.2 [86,87] for showering and hadronization. Then, DELPHES 3.4 [88] was used for detector simulation. We have used the default DELPHES tagging efficiencies and misidentification rates for taus and b-jets, where the tau tagging efficiency is 0.6, and the tau misidentification rate for QCD jets is 0.01. To increase the yield for background events to pass cuts described below, we forced every particle, except for the Standard Model (SM) Higgs boson (h), to decay into leptons (including taus) and/or jets, such that they contribute to multilepton final states with at least three leptons. For generation of both signal and background events, we only considered the decay of h into W^+W^- , ZZ , and $\tau^+\tau^-$, ignoring all other decays, and have normalized using the branching ratios of h from HDECAY [89]. Thus each signal or background containing one or more h is split into three signal or background components per h . We generated at least 10^5 events for each signal component at each mass point, and at least 10^6 events for each background component, at each of $\sqrt{s} = (14, 27, 70, 100)$ TeV. These numbers of generated events are usually sufficient, with a few exceptions noted below. In the real world, better background determinations may come from data.

In our analysis, we first impose object cuts on leptons $\ell = e, \mu, \tau$, where, from now on, τ refers to a tau lepton that undergoes a hadronic decay. These include minimum p_T cuts that depend on the collider option and are therefore listed in the subsections below. In all cases, the lepton candidates are required to pass the following further cuts on pseudorapidity η and isolation from other lepton candidates or jets,

$$|\eta| < 2.5, \quad (2.22)$$

$$\Delta R_{\ell,\ell'} > 0.1 \quad (\text{for each } \ell, \ell' = e, \mu, \tau), \quad (2.23)$$

$$\Delta R_{\ell,j} > 0.3 \quad (\text{for each jet and } \ell = e, \mu, \tau). \quad (2.24)$$

Here, $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$ as is usual.

Events are then selected with at least three leptons, of which at least two must be e or μ . The leading e or μ lepton is also required to satisfy a minimum p_T trigger requirement that depends on the collider option and is therefore listed separately in each of the subsections below. We also veto b -jets to reduce large backgrounds from $t\bar{t}$ production processes. Thus our event preselection common to all signal regions is

$$N(e, \mu, \tau) \geq 3, \quad (2.25)$$

$$N(e, \mu) \geq 2, \quad (2.26)$$

⁴Electronic input files relevant for VLL models are available from the authors on request.

$$p_T^{e_1} \text{ or } p_T^{\mu_1} > p_T^{\text{trigger}}, \quad (2.27)$$

$$N(b\text{-jets}) = 0. \quad (2.28)$$

Events with no e^+e^- or $\mu^+\mu^-$ pair with invariant mass within 15 GeV of M_Z are referred to below as “no Z,” and events with exactly two e/μ that have same-sign charges will be labeled as “SS.” We then considered the following six distinct types of signal regions:

$$2 \text{ SS } e/\mu + \geq 1\tau, \quad (2.29)$$

$$2 \text{ SS } e/\mu + \geq 1\tau \quad \text{with} \quad E_T^{\text{miss}} > 150 \text{ GeV}, \quad (2.30)$$

$$\geq 3e/\mu + 1\tau, \quad (2.31)$$

$$\geq 3e/\mu + 1\tau, \text{ no } Z, \quad (2.32)$$

$$\geq 2e/\mu + 2\tau, \quad (2.33)$$

$$\geq 2e/\mu + 2\tau, \text{ no } Z. \quad (2.34)$$

Finally, in each of these signal regions, we imposed a minimum lower bound cut on L_T , which is defined to be the sum of the transverse momentum of all leptons

$$L_T = \sum_{\ell=e,\mu,\tau} p_T(\ell). \quad (2.35)$$

We varied the choice of this cut to obtain good exclusion and discovery reach simultaneously for each signal region and collider option and assumed fractional background uncertainty $\Delta_b/b = (0.1, 0.2, 0.5)$. However, we have only done a very rough optimization for the L_T cut, in part because the optimization is different for exclusion and for discovery, and also because the Monte Carlo simulations are only an approximation of the actual experimental capabilities, which will rely on detector designs yet to be determined. In general, the results found below reflect that the choice of cut on L_T increases with the mass M_ν at the edge of the reach. Also, for convenience, we always choose the same L_T cut for both exclusion and discovery, even though an optimized cut would likely be somewhat different for the two cases.




FIG. 5. L_T event distributions for total background (shaded) and doublet VLL models (lines) for pp collisions at $\sqrt{s} = 14$ TeV with an integrated luminosity $\mathcal{L} = 3 \text{ ab}^{-1}$. Five different masses $M_\nu = M_\nu = 500, 700, 900, 1100,$ and 1300 GeV are shown in each panel. The four panels show results for the four best signal regions, as labeled.

Note that the signal regions considered in Eqs. (2.29)–(2.34) are far from exclusive of each other. Therefore, to be conservative we have not attempted to combine them, although doing so could lead to some extension of the reach prospects.

III. RESULTS FOR THE HL-LHC COLLIDER

In this section, we discuss the possibility of exclusion or discovery of both doublet and singlet VLL (including the minimal and nonminimal versions) at $\sqrt{s} = 14$ TeV with 3 ab^{-1} of pp collisions. In addition to the pseudorapidity, isolation, and other requirements of Eqs. (2.22)–(2.28), we require all leptons including hadronic tau candidates to satisfy

$$p_T^{\ell} > 15 \text{ GeV}. \quad (3.1)$$

Additionally, the leading e or μ in each event is required to satisfy a trigger requirement

$$p_T^{e_1} \text{ or } p_T^{\mu_1} > 30 \text{ GeV}. \quad (3.2)$$

We then considered the six signal regions mentioned in Eqs. (2.29)–(2.34).

A. Doublet VLL model

In Fig. 5, we show the L_T distributions for the best four of these signal regions, for five different choices of M_{τ} as labeled, and for the total of all backgrounds shown as the shaded histogram. For pp collisions with $\sqrt{s} = 14$ TeV with 3 ab^{-1} , we found that the expected reach is approximately maximized if we then choose a cut $L_T > 800$ GeV.

Figure 6 shows the L_T distributions for all background components, for the four best signal regions as labeled. The L_T cut is shown in the figure as a vertical dashed line. After imposing the L_T cut, the dominant SM backgrounds are WZ , $t\bar{t}V$, and VVV (where $V = W, Z$) in the two signal regions with $2 \text{ SS } e/\mu + \geq 1\tau$, while the dominant SM backgrounds are $t\bar{t}V$ and ZZ in the signal region with $\geq 2e/\mu + 2\tau$, and $t\bar{t}V$ and $t\bar{t}h$ in the signal region with $\geq 2e/\mu + 2\tau$ (no Z).

Figure 7 shows the resulting median expected significances for exclusion (left panels) and discovery (right panels), for $\Delta_b/b = 0.1$ (top row), 0.2 (middle row), and 0.5 (bottom row), with the cut requirement $L_T > 800$ GeV




FIG. 6. L_T event distributions for all processes contributing to total SM background, for pp collisions at $\sqrt{s} = 14$ TeV with an integrated luminosity $\mathcal{L} = 3 \text{ ab}^{-1}$. The four panels show results for the four best signal regions, as labeled. The vertical dashed line in all four panels shows our choice of L_T cut.




FIG. 7. The median expected significances for exclusion Z_{excl} (left) and discovery Z_{disc} (right) as a function of M_{τ} in the doublet VLL model, for pp collisions at $\sqrt{s} = 14$ TeV with integrated luminosity $\mathcal{L} = 3 \text{ ab}^{-1}$, for six different signal regions as described in the text, each including a cut $L_T > 800$ GeV. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$ (top), 0.2 (middle), and 0.5 (bottom).

imposed. In all cases, the best signal regions for both exclusion and discovery scenarios are the ones with 2 SS $e/\mu + \geq 1\tau$, with the additional requirement $E_T^{\text{miss}} > 150$ GeV providing slightly more reach. The two signal regions with $\geq 2e/\mu + 2\tau$ give slightly less reach at higher masses, but could actually provide comparable exclusion significances for lower masses.

From Fig. 7 we conclude that a 14 TeV high-luminosity LHC with 3 ab^{-1} should be able to exclude doublet VLLs with M_{τ} up to about 1250 GeV if they are indeed absent or discover them if the mass is less than about 900 GeV, assuming that the future background determination from data is subject to uncertainties of order 10% or less. The figure also shows that prospects for exclusion are much less




FIG. 8. The event distributions for three-body invariant mass of $\tau^\pm e^+ e^-$ or $\tau^\pm \mu^+ \mu^-$, for total background (shaded) and doublet VLLs (lines), such that the $e^+ e^-$ or $\mu^+ \mu^-$ pair have an invariant mass within 10 GeV of M_Z in a signal region with $\geq 2e/\mu + 2\tau$, with the cut $L_T > 800$ GeV imposed. Two different masses $M_\nu = M_\nu = 700$ and 800 GeV are shown.




FIG. 9. The median expected significances for exclusion Z_{excl} as a function of M_ν in the singlet VLL models, for pp collisions at $\sqrt{s} = 14$ TeV with integrated luminosity $\mathcal{L} = 3 \text{ ab}^{-1}$, for the best signal region for each of the singlet VLL models, including a cut on L_T as shown in the plot. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$.







FIG. 10. L_T event distributions for total background (shaded) and doublet VLL models (lines), for pp collisions at $\sqrt{s} = 27$ TeV with an integrated luminosity $\mathcal{L} = 15 \text{ ab}^{-1}$. Four different masses $M_\nu = M_\nu = 750, 1000, 1500,$ and 2000 GeV are shown in each panel. The four panels show results for the four best signal regions, as labeled.

sensitive to uncertainty in the background than the prospects for discovery. For $\Delta_b/b = 0.5$, one can still expect to exclude doublet VLLs up to about $M_{\tau'} = 1190$ GeV or discover them if the mass is less than about 690 GeV, but the latter has already been excluded at 95% confidence level by CMS [70,71].

In the case that there are enough events for a clear discovery, one can also hope to measure the mass of the τ' . In Fig. 8, we show the event distributions for the three-body invariant mass of $\tau^\pm e^+ e^-$ or $\tau^\pm \mu^+ \mu^-$, for the signal region with $\geq 2e/\mu + 2\tau$, for two different choices of input $M_{\tau'}$, and for the total of all backgrounds shown as the shaded histogram. (Combinatorial backgrounds from wrong associations of the lepton pair and tau in the signal sample are of course also present and included.) Here, we require the two-body invariant mass of the $e^+ e^-$ or $\mu^+ \mu^-$ pair to be within 10 GeV of M_Z . Additionally, we also impose the cut $L_T > 800$ GeV. From Fig. 8, we observe that the distributions for doublet VLLs are peaked just below their respective masses, which gives a possibility to measure the masses of doublet VLLs, if they are indeed discovered. At $\sqrt{s} = 14$ TeV, the range of masses that one can hope to measure in this way is limited by the

present exclusion up to $M_{\tau'} = 790$ GeV by CMS [71] and by the fact that the cross section decreases rapidly for higher masses.

B. Singlet VLL models

Singlet VLLs are much more challenging than doublet VLL, due to their much smaller production cross sections. For the minimal singlet model, we find no possible exclusion or discovery at $\sqrt{s} = 14$ TeV with integrated luminosity of 3 ab^{-1} . And, for the nonminimal singlet models, we find some exclusion possibility, but with no prospects for discovery. In view of the difficulties involved, below we consider only the case where the uncertainty in the background is $\Delta_b/b = 0.1$.

To maximize the exclusion reach in the nonminimal cases, we chose a cut $L_T > 700$ GeV for the Z-philic singlet VLL, $L_T > 400$ GeV for the W-phobic singlet VLL, and $L_T > 200$ GeV for the Higgs-philic singlet VLL. Figure 9 shows the resulting median expected significances for exclusion, for $\Delta_b/b = 0.1$, for the best signal region for each of the singlet VLL models, with the cuts on L_T imposed. In all the models, the best signal region




FIG. 11. L_T event distributions for all processes contributing to total SM background, for pp collisions at $\sqrt{s} = 27$ TeV with an integrated luminosity $\mathcal{L} = 15 \text{ ab}^{-1}$. The four panels show results for the four best signal regions, as labeled. The vertical dashed line in all four panels shows our choice of L_T cut of 1500 GeV.

for exclusion is the one which requires $\geq 2e/\mu + 2\tau$. However, for the Higgs-philic singlet VLL model, the additional requirement of no Z proved to be beneficial.

From Fig. 9, we conclude that assuming $\Delta_b/b = 0.1$, a 14 TeV pp collider with 3 ab^{-1} should be able to exclude singlet VLLs with masses up to about 600 GeV if they are

Z -philic, or exclude masses up to about 360 GeV if they are W -phobic, or exclude masses up to about 300 GeV if they are Higgs-philic. We find that there is no possible exclusion of singlet VLLs in the minimal model. Furthermore, there are no discovery prospects in minimal or nonminimal singlet VLL models.




FIG. 12. The median expected significances for exclusion Z_{excl} (left) and discovery Z_{disc} (right) as a function of $M_{\chi'}$ in the doublet VLL model, for pp collisions at $\sqrt{s} = 27 \text{ TeV}$ with integrated luminosity $\mathcal{L} = 15 \text{ ab}^{-1}$, for six different signal regions as described in the text, each including a cut $L_T > 1500 \text{ GeV}$. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$ (top), 0.2 (middle), and 0.5 (bottom).

IV. RESULTS FOR THE HE-LHC COLLIDER

In this section, we discuss the prospects for exclusion and discovery of VLLs at $\sqrt{s} = 27$ TeV with 15 ab^{-1} of pp collisions in the six signal regions mentioned in Eqs. (2.29)–(2.34). All leptons including hadronic tau candidates are required to satisfy

$$p_T^\ell > 25 \text{ GeV}, \quad (4.1)$$

along with the same pseudorapidity, isolation, and other requirements of Eqs. (2.22)–(2.28), with a trigger requirement of

$$p_T^{e_1} \text{ or } p_T^{\mu_1} > 50 \text{ GeV}. \quad (4.2)$$

A. Doublet VLL model

In Fig. 10, we show the L_T distributions for the best four of these signal regions, for four different choices of M_ν as labeled, and for the total of all backgrounds shown as the shaded histogram. The cut chosen to enhance the reach for exclusion and discovery in this case was $L_T > 1500$ GeV.

Figure 11 shows the L_T distributions for all background components, for the four best signal regions as labeled. The L_T cut is shown in the figure as a vertical dashed line. After imposing the L_T cut, the most important SM backgrounds are $t\bar{t}V$ and VVV in all four of these signal regions.

Figure 12 shows the Z_{excl} (left panels) and Z_{disc} (right panels) as a function of M_ν , for $\Delta_b/b = 0.1$ (top row), 0.2 (middle row), and 0.5 (bottom row), with the cut $L_T > 1500$ GeV imposed. The signal region with 2 SS $e/\mu + \geq 1\tau$ with $E_T^{\text{miss}} > 150$ GeV has the farthest mass reach. However, at lower masses the two signal regions with $\geq 2e/\mu + 2\tau$ give slightly better exclusion and discovery significances.

By looking at Fig. 12, we conclude that a 27 TeV high-energy LHC with 15 ab^{-1} could exclude doublet VLLs with masses up to about 2300 GeV or discover them if the mass is less than about 1700 GeV, assuming the fractional uncertainty in the background to be 0.1. On the other hand, if we assume $\Delta_b/b = 0.5$, we would still be able to exclude doublet VLLs up to $M_\nu = 2050$ GeV or discover them if the mass is less than about 1250 GeV. Just as in the case of the HL-LHC, from Fig. 12, a larger uncertainty in the background has a moderate effect on the prospects for exclusion, but a much larger impact on the prospects for discovery.

We again consider the prospects for observing a mass peak in the case where there are enough events for a clear discovery. In Fig. 13, we show the event distributions for the three-body invariant mass of $\tau^\pm e^+ e^-$ or $\tau^\pm \mu^+ \mu^-$, for the signal region with $\geq 2e/\mu + 2\tau$, for three different choices of M_ν . The total of all backgrounds is shown as the shaded histogram. We require the two-body invariant mass of the




FIG. 13. The event distributions for three-body invariant mass of $\tau^\pm e^+ e^-$ or $\tau^\pm \mu^+ \mu^-$, for total background (shaded) and doublet VLLs (lines), such that the $e^+ e^-$ or $\mu^+ \mu^-$ pair have an invariant mass within 10 GeV of M_Z in a signal region with $\geq 2e/\mu + 2\tau$, with the cut $L_T > 1500$ GeV imposed. Three different masses $M_\nu = M_\nu = 1000, 1250, \text{ and } 1500$ GeV are shown in the plot.

$e^+ e^-$ or $\mu^+ \mu^-$ pair to be within 10 GeV of M_Z . Additionally, we also impose the cut $L_T > 1500$ GeV. From Fig. 13, we observe that the distributions for doublet VLLs are peaked just below their respective masses, which gives a possibility to measure the masses of doublet VLLs, if they are indeed discovered. For higher masses, statistical limitations and the combinatorial background mean that the mass determinations will be problematic.

B. Singlet VLL models

At $\sqrt{s} = 27$ TeV with integrated luminosity of 15 ab^{-1} , we find that there is no exclusion or discovery reach for the minimal and the Higgs-philic singlet VLL models. For the other two nonminimal singlet models, we find some exclusion possibility, but again with no prospects for discovery.

We chose a cut $L_T > 1400$ GeV for the Z -philic singlet VLL and $L_T > 800$ GeV for the W -phobic singlet VLL, to maximize the exclusion reach. A cut $L_T > 600$ GeV for the Higgs-philic singlet VLL and $L_T > 800$ GeV for the minimal singlet VLL model were chosen for the best possible exclusion significance. Figure 14 shows the resulting median expected significances for exclusion, for $\Delta_b/b = 0.1$, for the best signal region for each of the singlet VLL models, with the cuts on L_T imposed, as mentioned above. The best signal region for exclusion, for all the singlet VLL models is the one which requires $\geq 2e/\mu + 2\tau$, except that an additional requirement of no Z gave better results for the Higgs-philic model.

From Fig. 14, we can conclude that a 27 TeV pp collider with 15 ab^{-1} could possibly exclude singlet VLLs with masses up to about 1200 GeV in the Z -philic model, or exclude masses up to about 670 GeV in the W -phobic model, but with no prospects for discovery. These results




FIG. 14. The median expected significances for exclusion Z_{excl} as a function of M_τ in the singlet VLL models, for pp collisions at $\sqrt{s} = 27$ TeV with integrated luminosity $\mathcal{L} = 15 \text{ ab}^{-1}$, for the best signal region for each of the singlet VLL models, including a cut on L_τ as shown in the plot. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$.

assume a fractional uncertainty in the background of $\Delta_b/b = 0.1$. In both minimal and the Higgs-philic singlet VLL models, there is no possibility for exclusion or

discovery. From the results for singlet VLL analyses for 14 and 27 TeV colliders, we can note that the exclusion reach scaled approximately linearly with \sqrt{s} , for both Z-philic and the W-phobic singlet VLL models.

V. RESULTS FOR A pp COLLIDER WITH $\sqrt{s} = 70$ TeV

In this section, we turn our attention to prospects for exclusion and discovery of VLLs at a possible future pp collider at $\sqrt{s} = 70$ TeV, with integrated luminosity of 30 ab^{-1} in the six signal regions mentioned in Eqs. (2.29)–(2.34). All the leptons including hadronic tau candidates are required to satisfy

$$p_T^\ell > 75 \text{ GeV}, \quad (5.1)$$

along with the same pseudorapidity, isolation, and other requirements of Eqs. (2.22)–(2.28), with at least one e or μ satisfying a trigger requirement

$$p_T^{e_1} \text{ or } p_T^{\mu_1} > 150 \text{ GeV}. \quad (5.2)$$




FIG. 15. L_τ event distributions for total background (shaded) and doublet VLL models (lines), for pp collisions at $\sqrt{s} = 70$ TeV with an integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$. Four different masses $M_\tau = M_\nu = 1000, 2000, 3000,$ and 4000 GeV are shown in each panel. The four panels show results for the four best signal regions, as labeled.

A. Doublet VLL model

In Fig. 15, we show the L_T distributions for the best four of the signal regions, for four different choices of $M_{\tau'}$ as labeled, and for the total of all backgrounds shown as the shaded histogram. We found that choosing the cut $L_T > 2800$ GeV provides a good reach for both exclusion and discovery in this case.

Figure 16 shows the L_T distributions for all background components, for the four best signal regions as labeled. The L_T cut is shown in the figure as a vertical dashed line. After imposing the L_T cut, the most important SM backgrounds are $t\bar{t}V$ and VVV in the two signal regions with $2e/\mu \geq 1\tau$, while the most important SM backgrounds are WZ and $t\bar{t}V$ in the signal region with $\geq 2e/\mu + 2\tau$ and $t\bar{t}h$ and $t\bar{t}V$ in the signal region with $\geq 2e/\mu + 2\tau$ (no Z). We note that the fluctuation of about 66 normalized events in the bin from 4000 to 4500 GeV in the lower left panel (i.e., signal region with $\geq 2e/\mu + 2\tau$) of Fig. 15 is due to a single simulated event of WZ background. This is because of the large cross section but extremely low yield for this background component in this signal region even after forcing decays to leptons. This is an unavoidable source of uncertainty for our analysis; given that our sample size was already 5.5×10^6 simulated events for this component,

no practically feasible increase in sample size would yield significantly better statistics. However, in the real world the background can perhaps be determined more accurately from data.

Figure 17 shows the median expected significances for exclusion Z_{excl} (left panels) and discovery Z_{disc} (right panels) as a function of $M_{\tau'}$, for $\Delta_b/b = 0.1$ (top row), 0.2 (middle row), and 0.5 (bottom row), with the cut $L_T > 2800$ GeV imposed. The two signal regions with $2e/\mu \geq 1\tau$ and the signal region with $\geq 2e/\mu + 2\tau$ (no Z) have comparable exclusion significances and reaches. However, the latter has higher discovery significances at lower masses, as well as at higher fractional uncertainties in the background, e.g., $\Delta_b/b = 0.5$.

Figure 17 shows a possibility of excluding doublet VLLs of masses up to about 4700 GeV or discovering them if mass is less than about 3400 GeV with a 70 TeV pp collider with 30 ab^{-1} , assuming the fractional uncertainty in the background to be $\Delta_b/b = 0.1$. If $\Delta_b/b = 0.5$, we can still expect to exclude doublet VLLs up to $M_{\tau'} = 4150$ GeV or discover them if the mass is less than about 2400 GeV. Again, a larger uncertainty in the background has a moderate effect on the prospects for exclusion, but a much larger impact on the prospects for discovery.




FIG. 16. L_T event distributions for all processes contributing to total SM background, for pp collisions at $\sqrt{s} = 70$ TeV with an integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$. The four panels show results for the four best signal regions, as labeled. The vertical dashed line in all four panels shows our choice of L_T cut of 2800 GeV.




FIG. 17. The median expected significances for exclusion Z_{excl} (left) and discovery Z_{disc} (right) as a function of M_{τ} in the doublet VLL model, for pp collisions at $\sqrt{s} = 70$ TeV with integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$, for six different signal regions as described in the text, each including a cut $L_T > 2800$ GeV. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$ (top), 0.2 (middle), and 0.5 (bottom).

B. Singlet VLL models

For pp collisions with $\sqrt{s} = 70$ TeV with 30 ab^{-1} , we again find that there is no possible exclusion or discovery reach for the minimal and the Higgs-philic singlet VLL models. While we find some exclusion possibility for the Z -philic and the W -phobic singlet VLL models, there are again no prospects for discovery. Figure 18 shows the resulting median expected significances for exclusion, for $\Delta_b/b = 0.1$,

for the best signal region for each of the singlet VLL models, with appropriate cuts on L_T imposed. To approximately maximize the exclusion reaches for the Z -philic and W -phobic singlet VLL models, we chose a cut $L_T > 2100$ GeV for the former and $L_T > 950$ GeV for the latter. The best signal region for exclusion for the Z -philic singlet VLL model is the one which requires $\geq 3e/\mu + 1\tau$, while it was $\geq 2e/\mu + 2\tau$, no Z for all other singlet VLL models.




FIG. 18. The median expected significances for exclusion Z_{excl} as a function of M_χ in the singlet VLL models, for pp collisions at $\sqrt{s} = 70$ TeV with integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$, for the best signal region for each of the singlet VLL models, including a cut on L_T as shown in the plot. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$.

From Fig. 18, we conclude that a 70 TeV pp collider with 30 ab^{-1} could possibly exclude singlet VLLs with masses up to about 1700 GeV in the Z-philic model or exclude masses up to 850 GeV in the W-phobic model, but with no prospects for 5σ discovery. In both minimal and the Higgs-philic singlet VLL models, there is unfortunately no possibility for exclusion or discovery, at least with the signal regions we considered.

VI. RESULTS FOR A pp COLLIDER WITH $\sqrt{s} = 100$ TeV

Finally, we consider the possibility of excluding or discovering VLLs at a future very high-energy pp collider at $\sqrt{s} = 100$ TeV, with integrated luminosity of 30 ab^{-1} , in the six signal regions mentioned in Eqs. (2.29)–(2.34). We require all leptons including hadronic tau candidates to satisfy

$$p_T^\ell > 100 \text{ GeV}, \quad (6.1)$$




FIG. 19. L_T event distributions for total background (shaded) and doublet VLL models (lines), for pp collisions at $\sqrt{s} = 100$ TeV with an integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$. Four different masses $M_\chi = M_{\chi'} = 2000, 3000, 4000,$ and 5000 GeV are shown in each panel. The four panels show results for the four best signal regions, as labeled.

along with the same pseudorapidity, isolation, and other requirements of Eqs. (2.22)–(2.28). We then require the leading e or μ in each event to satisfy a trigger requirement

$$p_T^{e_1} \text{ or } p_T^{\mu_1} > 200 \text{ GeV}. \quad (6.2)$$

A. Doublet VLL model

In Fig. 19, we show the L_T distributions for the best four of these signal regions, for four different choices of $M_{\tau'}$ as labeled, and for the total of all backgrounds shown as the shaded histogram. To obtain enhanced expected reaches for both exclusion and discovery, we then chose a cut $L_T > 3500$ GeV.

Figure 20 shows the L_T distributions for all background components, for the four best signal regions as labeled. The L_T cut is shown in the figure as a vertical dashed line. After imposing the L_T cut, the most important SM backgrounds are $t\bar{t}V$ and VVV in the two signal regions with $2 \text{ SS } e/\mu + \geq 1\tau$ and in the signal region with $\geq 2e/\mu + 2\tau$, while the most important SM background in the signal region with $\geq 2e/\mu + 2\tau$ (no Z) is VVV .




FIG. 20. L_T event distributions for all processes contributing to total SM background, for pp collisions at $\sqrt{s} = 100$ TeV with an integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$. The four panels show results for the four best signal regions, as labeled. The vertical dashed line in all four panels shows our choice of L_T cut of 3500 GeV.

Figure 21 shows the median expected significances for exclusion Z_{excl} (left panels) and discovery Z_{disc} (right panels) as a function of $M_{\tau'}$, for $\Delta_b/b = 0.1$ (top row), 0.2 (middle row), and 0.5 (bottom row), with the cut $L_T > 3500$ GeV imposed. The signal regions which require $2 \text{ SS } e/\mu + \geq 1\tau$ usually provide the best exclusion and discovery reaches. An exception is that, with the fractional uncertainty in the background $\Delta_b/b = 0.5$, the signal regions with $\geq 3e/\mu + 1\tau$ have the farthest discovery reach.

From Fig. 21, we conclude that a 100 TeV pp collider with 30 ab^{-1} could exclude doublet VLLs with $M_{\tau'}$ up to about 5750 GeV or discover them if the mass is less than about 4000 GeV, assuming the fractional uncertainty in the background to be 0.1. For $\Delta_b/b = 0.5$, one can still expect to exclude doublet VLLs if the mass is up to about 5100 GeV or discover them if the mass is less than about 3100 GeV. Again, as a recurring theme at all the collider options considered, a larger uncertainty in the background produces a moderate reduction of the exclusion reach, but has a much greater impact on the prospects for discovery.

We again consider the possibility of observing a mass peak for the τ' when a clear discovery can be made.




FIG. 21. The median expected significances for exclusion Z_{excl} (left) and discovery Z_{disc} (right) as a function of $M_{\chi'}$ in the doublet VLL model, for pp collisions at $\sqrt{s} = 100$ TeV with integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$, for six different signal regions as described in the text, each including a cut $L_T > 3500$ GeV. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$ (top), 0.2 (middle), and 0.5 (bottom).

Figure 22 shows the event distributions for three-body invariant mass of $\tau^\pm e^+ e^-$ or $\tau^\pm \mu^+ \mu^-$, for the signal region with $\geq 2e/\mu + 2\tau$, for three different choices of $M_{\chi'}$, and for the total of all backgrounds shown as the shaded histogram. We require the two-body invariant mass of the $e^+ e^-$ or $\mu^+ \mu^-$ pair to be within 10 GeV of M_Z . We also impose the cut $L_T > 3500$ GeV. From Fig. 22, we note that there are peaks in the distributions for doublet VLLs corresponding to, and slightly below, their respective

masses, giving a possibility to measure the masses of doublet VLLs, if they are indeed discovered.

B. Singlet VLL models

We find no possible exclusion of singlet VLLs in the minimal and the Higgs-philic singlet VLL models, for pp collisions with $\sqrt{s} = 100$ TeV with integrated luminosity of 30 ab^{-1} . We find some exclusion prospects for the




FIG. 22. The event distributions for three-body invariant masses of $\tau^\pm e^+ e^-$ or $\tau^\pm \mu^+ \mu^-$, for total background (shaded) and doublet VLLs (lines), such that the $e^+ e^-$ or $\mu^+ \mu^-$ pair have an invariant mass within 10 GeV of M_Z in a signal region with $\geq 2e/\mu + 2\tau$, with the cut $L_T > 3500$ GeV imposed. Three different masses $M_\nu = M_\nu = 1500, 2000,$ and 2500 GeV are shown in the plot.

Z-philic and W-phobic singlet VLL models, assuming that the fractional uncertainty in the background is $\Delta_b/b = 0.1$. Again, there are no discovery prospects in any of the singlet VLL models.

We chose cuts of $L_T > 3000$ GeV for the Z-philic singlet VLL, and $L_T > 1200$ GeV for the W-phobic singlet VLL, to approximately maximize the exclusion reach in each case. Figure 23 shows the resulting median expected significances for exclusion, for $\Delta_b/b = 0.1$, for the best signal region for each of the singlet VLL models, with the cuts on L_T imposed, as mentioned above. The best signal region for exclusion for the Z-philic singlet VLL model is the one which requires $\geq 3e/\mu + 1\tau$, while it was $\geq 2e/\mu + 2\tau$, no Z for all other singlet VLL models.




FIG. 23. The median expected significances for exclusion Z_{excl} as a function of M_τ in the singlet VLL models, for pp collisions at $\sqrt{s} = 100$ TeV with integrated luminosity $\mathcal{L} = 30 \text{ ab}^{-1}$, for the best signal region for each of the singlet VLL models, including a cut on L_T as shown in the plot. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$.

TABLE I. Summary of mass reaches for the 95% exclusion and 5σ discovery of doublet VLL at future pp colliders. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$.

Collider	Exclusion reach (GeV)	Discovery reach (GeV)
14 TeV HL-LHC, 3 ab^{-1}	1250	900
27 TeV HE-LHC, 15 ab^{-1}	2300	1700
70 TeV pp collider, 30 ab^{-1}	4700	3400
100 TeV pp collider, 30 ab^{-1}	5750	4000

From Fig. 23, assuming $\Delta_b/b = 0.1$, we conclude that a 100 TeV pp collider with 30 ab^{-1} should be able to exclude singlet VLLs with masses up to about 2850 GeV, if they are Z-philic, or exclude masses up to 1200 GeV, if they are W-phobic. We find that there are no prospects for exclusion of singlet VLLs in both minimal and the Higgs-philic singlet VLL models.

VII. OUTLOOK

Vectorlike leptons are a common feature of motivated theories of physics beyond the Standard Model. In this paper, we have studied the prospects for discovering simple VLL models at future pp colliders, assuming that the new particles decay promptly by mixing with taus, using multilepton signatures at high transverse momentum. The results of this paper should be considered as first estimates, which certainly should be reassessed as the plans and parameters for future colliders and their detectors become clearer. We have not attempted to make use of $h \rightarrow b\bar{b}$ decays, due in part to a relative lack of confidence in what should be assumed about b -jet tagging capabilities compared to e, μ identification. We have also based our analysis entirely on a cut-and-count strategy, while a more sophisticated method such as a likelihood analysis would certainly be able to do better.

Summaries of our results are shown in Table I for the doublet VLL model and in Table II for the nonminimal singlet VLL models. In the weak-isodoublet case, we have found excellent reach prospects for both 95% exclusion (if the VLLs are absent) or 5σ discovery (if the VLLs are present). In the latter case, we also showed that there is a possibility to observe an invariant mass peak, although we did not pursue this in detail and the mass peak observation is often quite limited by statistics. In the weak-isosinglet case, the situation is much more difficult. We found no discovery or exclusion reach at all,⁵ at any pp collider, for the simple and well-motivated (e.g., [8]) case of a promptly

⁵The only limits on such a particle at present are $M_\tau > 101.2$ GeV from LEP [74], and our estimate above in a footnote in Sec. II, inferred from the ATLAS long-lived chargino search of Ref. [82], of $M_\tau > 750$ GeV if it is quasistable on detector scales.

TABLE II. Summary of mass reaches for the exclusion of singlet VLL in the nonminimal models at future pp colliders. The fractional uncertainty in the background is assumed to be $\Delta_b/b = 0.1$.

Collider	Z-philic (GeV)	W-phobic (GeV)	Higgs-philic (GeV)
14 TeV HL-LHC, 3 ab ⁻¹	600	360	300
27 TeV HE-LHC, 15 ab ⁻¹	1200	670	...
70 TeV pp collider, 30 ab ⁻¹	1700	850	...
100 TeV pp collider, 30 ab ⁻¹	2850	1200	...

decaying pure isosinglet τ' that mixes with the Standard Model tau. This should stand as a challenge to future work.

ACKNOWLEDGMENTS

P. N. B. thanks Olivier Mattelaer for all the helpful suggestions regarding the usage of Madgraph. P. N. B. also thanks Sergey A. Uzunyan for his kind assistance in using the NICADD compute cluster at Northern Illinois University, and Ramanpreet Singh and Elliot Parrish for technical advice. P. N. B. would also like to express special thanks to Nicholas T. Karonis, John Winans, and the entire support team of Gaea cluster at Northern Illinois University for their help and support in using their high-performance computing facility. This work was supported in part by the National Science Foundation Grant No. PHY-1719273.

APPENDIX: MIXED SINGLET AND DOUBLET VLL MODELS

Consider a generalized model framework that contains, besides the Standard Model chiral third-family leptons ℓ_3 and $\bar{\ell}_3$, a vectorlike weak-isosinglet pair E, \bar{E} and a weak-isodoublet pair L, \bar{L} . (Here we use two-component fermion notation [90,91], so that barred fields have electric charge +1 and unbarred fields have electric charge -1. The bar is part of the name of the two-component fermion field and does not denote any kind of conjugation.) After electro-weak symmetry breaking, the Lagrangian mass terms for the charged leptons can be written in the form

$$-\mathcal{L} = (\bar{\ell}_3 \quad \bar{E} \quad \bar{L}) \mathcal{M} \begin{pmatrix} \ell_3 \\ E \\ L \end{pmatrix} + \text{c.c.}, \quad (\text{A1})$$

where the charged lepton mixing mass matrix is

$$\mathcal{M} = \begin{pmatrix} y_\tau v & 0 & \epsilon_2 v \\ \epsilon_1 v & M_1 & x_2 v \\ 0 & x_1 v & M_2 \end{pmatrix}, \quad (\text{A2})$$

where $v = 174$ GeV is the Standard Model Higgs expectation value, $y_\tau, \epsilon_1, \epsilon_2, x_1$, and x_2 are Yukawa couplings, and M_1 and M_2 are electroweak-singlet bare mass terms for the isosinglet and isodoublet vectorlike lepton pairs, respectively. This can be diagonalized to obtain mass eigenstates according to

$$R^* \mathcal{M} L^\dagger = \text{diag}(M_\tau, M_{\tau'}, M_{\tau''}), \quad (\text{A3})$$

where R and L are unitary 3×3 matrices and, by convention, $M_\tau < M_{\tau'} < M_{\tau''}$, where τ is the usual tau lepton with $M_\tau = 1.777$ GeV, and in the special cases considered here τ'' will be taken to be heavy enough to decouple from direct experimental observation. The neutral VLL ν' has mass M_2 .

For the lighter new charged VLL τ' , we have partial decay widths

$$\Gamma(\tau' \rightarrow W \nu_\tau) = \frac{M_{\tau'}}{32\pi} (1 - r_W)^2 (1 + 2r_W) |g_{\tau'\nu_\tau}^W|^2 / r_W, \quad (\text{A4})$$

$$\Gamma(\tau' \rightarrow Z \tau) = \frac{M_{\tau'}}{32\pi} (1 - r_Z)^2 (1 + 2r_Z) (|g_{\tau'\tau}^Z|^2 + |g_{\tau'\tau}^Z|^2) / r_Z, \quad (\text{A5})$$

$$\Gamma(\tau' \rightarrow h \tau) = \frac{M_{\tau'}}{32\pi} (1 - r_h)^2 (|y_{\tau'\tau}^h|^2 + |y_{\tau'\tau}^h|^2), \quad (\text{A6})$$

where $r_X = M_X^2 / M_{\tau'}^2$ for each of $X = h, Z, W$, and

$$g_{\tau_j \nu_\tau}^W = \frac{g}{\sqrt{2}} L_{j1}^*, \quad (\text{A7})$$

$$g_{\tau_j \tau_k}^Z = \frac{g}{\cos \theta_W} \left[\frac{1}{2} L_{j2}^* L_{k2} + \left(-\frac{1}{2} + \sin^2 \theta_W \right) \delta_{jk} \right], \quad (\text{A8})$$

$$g_{\tau_j \tau_k}^Z = \frac{g}{\cos \theta_W} \left[\frac{1}{2} R_{j3}^* R_{k3} - \sin^2 \theta_W \delta_{jk} \right], \quad (\text{A9})$$

$$y_{\tau_j \tau_k}^h = \frac{1}{\sqrt{2}} [y_\tau L_{j1}^* R_{k1}^* + \epsilon_2 L_{j3}^* R_{k1}^* + \epsilon_1 L_{j1}^* R_{k2}^* + x_2 L_{j3}^* R_{k2}^* + x_1 L_{j2}^* R_{k3}^*], \quad (\text{A10})$$

for (τ_1, τ_2, τ_3) corresponding to mass eigenstates (τ, τ', τ'') , respectively.

In general, the decay widths to Standard Model states are quadratic in ϵ_1, ϵ_2 for small values of those parameters. For example, expanding to obtain the terms proportional to $\epsilon_1^2, \epsilon_1 \epsilon_2$, and ϵ_2^2 , and then keeping only the leading order in a further expansion in y_τ, x_1 , and x_2 in each of these terms, we obtain for $M_2^2 \ll M_1^2$,

$$|g_{\tau\nu\tau}^W|^2/r_W \approx \left[\frac{\epsilon_1 v(x_1 M_1 + x_2 M_2)}{M_1^2} - \frac{\epsilon_2 y_\tau v}{M_2} \right]^2, \quad (\text{A11})$$

$$\begin{aligned} (|g_{\tau\tau}^Z|^2 + |g_{\tau\tau'}^Z|^2)/r_Z \approx & \epsilon_1^2 v^2 \frac{M_2^2(x_1 M_2 + x_2 M_1)^2}{2M_1^6} \\ & - \epsilon_1 \epsilon_2 y_\tau v^2 \frac{(x_1 M_1 + x_2 M_2)}{M_1^2 M_2} + \frac{\epsilon_2^2}{2}, \end{aligned} \quad (\text{A12})$$

$$\begin{aligned} |y_{\tau\tau}^h|^2 + |y_{\tau\tau'}^h|^2 \approx & \epsilon_1^2 v^2 \frac{(2x_1 M_1 + x_2 M_2)^2}{2M_1^4} \\ & - 3\epsilon_1 \epsilon_2 y_\tau v^2 \frac{(x_1 M_1 + x_2 M_2)}{M_1^2 M_2} + \frac{\epsilon_2^2}{2}, \end{aligned} \quad (\text{A13})$$

and for $M_1^2 \ll M_2^2$,

$$|g_{\tau\nu\tau}^W|^2/r_W \approx \left[\epsilon_1 - \epsilon_2 y_\tau v^2 \frac{(x_1 M_2 + x_2 M_1)}{M_1 M_2^2} \right]^2, \quad (\text{A14})$$

$$\begin{aligned} (|g_{\tau\tau}^Z|^2 + |g_{\tau\tau'}^Z|^2)/r_Z \approx & \frac{\epsilon_1^2}{2} - \epsilon_1 \epsilon_2 y_\tau v^2 \frac{(x_1 M_2 + x_2 M_1)}{M_1 M_2^2} \\ & + \epsilon_2^2 v^2 \frac{M_1^2(x_1 M_1 + x_2 M_2)^2}{2M_2^6}, \end{aligned} \quad (\text{A15})$$

$$\begin{aligned} |y_{\tau\tau}^h|^2 + |y_{\tau\tau'}^h|^2 \approx & \frac{\epsilon_1^2}{2} - 3\epsilon_1 \epsilon_2 y_\tau v^2 \frac{(x_1 M_2 + x_2 M_1)}{M_1 M_2^2} \\ & + \epsilon_2^2 v^2 \frac{(2x_1 M_2 + x_2 M_1)^2}{2M_2^4}. \end{aligned} \quad (\text{A16})$$

Some special limits of interest follow:

- (1) If $M_2^2 \ll M_1^2$ and $\epsilon_1 = 0$, then the isosinglet heavier fermion mass eigenstate τ'' decouples from experiment, and the lighter states τ' and ν' form the minimal doublet VLL model as discussed above in Sec. II with $\epsilon = \epsilon_2$ and $M_{\nu'} = M_{\tau'} = M_2$, and branching ratios that asymptotically approach $\text{BR}(\tau' \rightarrow h\tau) = \text{BR}(\tau' \rightarrow Z\tau) = 0.5$ and $\text{BR}(\nu' \rightarrow W\tau) = 1$.
- (2) If $M_1^2 \ll M_2^2$ and $\epsilon_2 = 0$, then the heavy isodoublet fermions τ'' and ν' decouple, and the result is the minimal singlet VLL model as discussed above in Sec. II with $\epsilon = \epsilon_1$ and $M_{\nu'} = M_1$, and branching ratios that asymptotically approach $\text{BR}(\tau' \rightarrow h\tau) = \text{BR}(\tau' \rightarrow Z\tau) = 0.25$ and $\text{BR}(\tau' \rightarrow W\nu_\tau) = 0.5$.
- (3) If $M_1^2 \ll M_2^2$ and $\epsilon_1 = 0$ and $\epsilon_2 \neq 0$, then the lightest new fermion will again be a mostly isosinglet τ' vectorlike lepton with mass approximately $M_{\nu'} = M_1$. Since the τ' is mostly isosinglet, its production cross section is nearly the same as in the minimal singlet VLL model. However, due to its mixing with the heavier isodoublets rather than direct mixing with the Standard Model tau lepton, the decay $\tau' \rightarrow W\nu_\tau$ is highly suppressed. The possibilities include the following subcases:
 - (a) *W-phobic singlet VLL*: if $x_1 = 0$, then $\text{BR}(\tau' \rightarrow h\tau) \approx \text{BR}(\tau' \rightarrow Z\tau) \approx 0.5$.
 - (b) *Higgs-philic singlet VLL*: if $x_2 = 0$, then $\text{BR}(\tau' \rightarrow h\tau) \approx 1$.
 - (c) *Z-philic singlet VLL*: if $x_1 \approx -x_2 M_1/2M_2$, then $\text{BR}(\tau' \rightarrow Z\tau) \approx 1$.

These of course do not exhaust the possibilities, and in a more general search it would be sensible to simply take $\Gamma(\tau' \rightarrow Z\tau)$ and $\Gamma(\tau' \rightarrow h\tau)$ and $\Gamma(\tau' \rightarrow W\nu_\tau)$ to be free parameters.

[1] G. Aad *et al.* (ATLAS Collaboration), Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC, *Phys. Lett. B* **716**, 1 (2012); S. Chatrchyan *et al.* (CMS Collaboration), Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC, *Phys. Lett. B* **716**, 30 (2012).
[2] See, for example, A. Djouadi and A. Lenz, Sealing the fate of a fourth generation of fermions, *Phys. Lett. B* **715**, 310 (2012); A. Lenz, Constraints on a fourth generation of fermions from Higgs Boson searches, *Adv. High Energy Phys.* **2013**, 910275 (2013); O. Eberhardt, G. Herbert, H. Lacker, A. Lenz, A. Menzel, U. Nierste, and M. Wiebusch, Impact of a Higgs Boson at a Mass of 126 GeV on the

Standard Model with Three and Four Fermion Generations, *Phys. Rev. Lett.* **109**, 241802 (2012), and references therein.
[3] For a review, see S. P. Martin, A supersymmetry primer, *Adv. Ser. Dir. High Energy Phys.* **21**, 1 (2010); **18**, 1 (1998).
[4] T. Moroi and Y. Okada, Radiative corrections to Higgs masses in the supersymmetric model with an extra family and antifamily, *Mod. Phys. Lett. A* **07**, 187 (1992).
[5] T. Moroi and Y. Okada, Upper bound of the lightest neutral Higgs mass in extended supersymmetric Standard Models, *Phys. Lett. B* **295**, 73 (1992).
[6] K. S. Babu, I. Gogoladze, and C. Kolda, Perturbative unification and Higgs boson mass bounds, [arXiv:hep-ph/0410085](https://arxiv.org/abs/hep-ph/0410085).

- [7] K. S. Babu, I. Gogoladze, M. U. Rehman, and Q. Shafi, Higgs Boson mass, sparticle spectrum and little hierarchy problem in extended MSSM, *Phys. Rev. D* **78**, 055017 (2008).
- [8] S. P. Martin, Extra vector-like matter and the lightest Higgs scalar boson mass in low-energy supersymmetry, *Phys. Rev. D* **81**, 035004 (2010).
- [9] P. W. Graham, A. Ismail, S. Rajendran, and P. Saraswat, A little solution to the little hierarchy problem: A vector-like generation, *Phys. Rev. D* **81**, 055016 (2010).
- [10] S. P. Martin, Raising the Higgs mass with Yukawa couplings for isotriplets in vector-like extensions of minimal supersymmetry, *Phys. Rev. D* **82**, 055019 (2010).
- [11] M. Endo, K. Hamaguchi, S. Iwamoto, and N. Yokozaki, Higgs mass and muon anomalous magnetic moment in supersymmetric models with vector-like matters, *Phys. Rev. D* **84**, 075017 (2011).
- [12] J. L. Evans, M. Ibe, and T. T. Yanagida, Probing extra matter in gauge mediation through the lightest Higgs Boson mass, [arXiv:1108.3437](https://arxiv.org/abs/1108.3437).
- [13] T. Li, J. A. Maxin, D. V. Nanopoulos, and J. W. Walker, A Higgs mass shift to 125 GeV and a multi-jet supersymmetry signal: Miracle of the Flippons at the $\sqrt{s} = 7$ TeV LHC, *Phys. Lett. B* **710**, 207 (2012).
- [14] T. Moroi, R. Sato, and T. T. Yanagida, Extra matters decree the relatively heavy Higgs of mass about 125 GeV in the supersymmetric model, *Phys. Lett. B* **709**, 218 (2012).
- [15] M. Endo, K. Hamaguchi, S. Iwamoto, and N. Yokozaki, Higgs mass, muon $g-2$, and LHC prospects in gauge mediation models with vector-like matters, *Phys. Rev. D* **85**, 095012 (2012).
- [16] M. Endo, K. Hamaguchi, S. Iwamoto, and N. Yokozaki, Vacuum stability bound on extended GMSB models, *J. High Energy Phys.* **06** (2012) 060.
- [17] K. Nakayama and N. Yokozaki, Peccei-Quinn extended gauge-mediation model with vector-like matter, *J. High Energy Phys.* **11** (2012) 158.
- [18] S. P. Martin and J. D. Wells, Implications of gauge-mediated supersymmetry breaking with vector-like quarks and a 125 GeV Higgs boson, *Phys. Rev. D* **86**, 035017 (2012).
- [19] A. M. Sirunyan *et al.* (CMS Collaboration), Search for vector-like T and B quark pairs in final states with leptons at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **08** (2018) 177; Search for top quark partners with charge 5/3 in the same-sign dilepton and single-lepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, *J. High Energy Phys.* **03** (2019) 082.
- [20] M. Aaboud *et al.* (ATLAS Collaboration), Search for pair production of vector-like top quarks in events with one lepton, jets, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector, *J. High Energy Phys.* **08** (2017) 052; Search for pair production of heavy vector-like quarks decaying to high- p_T W bosons and b quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **10** (2017) 141; Search for pair production of up-type vector-like quarks and for four-top-quark events in final states with multiple b -jets with the ATLAS detector, *J. High Energy Phys.* **07** (2018) 089; Search for pair production of heavy vector-like quarks decaying into high- p_T W bosons and top quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *J. High Energy Phys.* **08** (2018) 048; Search for pair- and single-production of vector-like quarks in final states with at least one Z boson decaying into a pair of electrons or muons in pp collision data collected with the ATLAS detector at $\sqrt{s} = 13$ TeV, [arXiv:1806.10555](https://arxiv.org/abs/1806.10555); Search for pair production of heavy vector-like quarks decaying into hadronic final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector, *Phys. Rev. D* **98**, 092005 (2018); Combination of the Searches for Pair-Produced Vector-Like Partners of the Third-Generation Quarks at $\sqrt{s} = 13$ TeV with the ATLAS Detector, *Phys. Rev. Lett.* **121**, 211801 (2018).
- [21] S. Dimopoulos, N. Tetradis, R. Esmailzadeh, and L. J. Hall, TeV dark matter, *Nucl. Phys.* **B349**, 714 (1991).
- [22] M. Sher, Charged leptons with nanosecond lifetimes, *Phys. Rev. D* **52**, 3136 (1995).
- [23] S. D. Thomas and J. D. Wells, Phenomenology of Massive Vectorlike Doublet Leptons, *Phys. Rev. Lett.* **81**, 34 (1998).
- [24] P. H. Frampton, P. Q. Hung, and M. Sher, Quarks and leptons beyond the third generation, *Phys. Rep.* **330**, 263 (2000).
- [25] F. del Aguila, J. de Blas, and M. Perez-Victoria, Effects of new leptons in electroweak precision data, *Phys. Rev. D* **78**, 013010 (2008).
- [26] F. del Aguila, A. Carmona, and J. Santiago, Neutrino masses from an A4 symmetry in holographic composite Higgs models, *J. High Energy Phys.* **08** (2010) 127; Tau Custodian searches at the LHC, *Phys. Lett. B* **695**, 449 (2011); A. Carmona and F. Goertz, Custodial leptons and Higgs decays, *J. High Energy Phys.* **04** (2013) 163.
- [27] K. Ishiwata and M. B. Wise, Higgs Properties and fourth generation leptons, *Phys. Rev. D* **84**, 055025 (2011).
- [28] R. Dermisek, Insensitive unification of gauge couplings, *Phys. Lett. B* **713**, 469 (2012).
- [29] A. Joglekar, P. Schwaller, and C. E. M. Wagner, Dark matter and enhanced Higgs to Di-photon rate from vector-like leptons, *J. High Energy Phys.* **12** (2012) 064.
- [30] J. Kearney, A. Pierce, and N. Weiner, Vectorlike Fermions and Higgs couplings, *Phys. Rev. D* **86**, 113005 (2012).
- [31] C. Arina, R. N. Mohapatra, and N. Sahu, Co-genesis of matter and dark matter with vector-like fourth generation leptons, *Phys. Lett. B* **720**, 130 (2013).
- [32] B. Batell, S. Jung, and H. M. Lee, Singlet assisted vacuum stability and the Higgs to diphoton rate, *J. High Energy Phys.* **01** (2013) 135.
- [33] R. Dermisek, Unification of gauge couplings in the standard model with extra vectorlike families, *Phys. Rev. D* **87**, 055008 (2013).
- [34] W. Z. Feng and P. Nath, Higgs diphoton rate and mass enhancement with vectorlike leptons and the scale of supersymmetry, *Phys. Rev. D* **87**, 075018 (2013).
- [35] A. Joglekar, P. Schwaller, and C. E. M. Wagner, A supersymmetric theory of vector-like leptons, *J. High Energy Phys.* **07** (2013) 046.
- [36] P. Schwaller, T. M. P. Tait, and R. Vega-Morales, Dark matter and vectorlike leptons from gauged lepton number, *Phys. Rev. D* **88**, 035001 (2013).
- [37] R. Dermisek and A. Raval, Explanation of the Muon $g-2$ anomaly with vectorlike leptons and its implications for Higgs decays, *Phys. Rev. D* **88**, 013017 (2013).

- [38] K. Ishiwata and M. B. Wise, Phenomenology of heavy vectorlike leptons, *Phys. Rev. D* **88**, 055009 (2013).
- [39] M. Fairbairn and P. Grothaus, Baryogenesis and dark matter with vector-like Fermions, *J. High Energy Phys.* **10** (2013) 176.
- [40] W. Altmannshofer, M. Bauer, and M. Carena, Exotic leptons: Higgs, flavor and collider phenomenology, *J. High Energy Phys.* **01** (2014) 060.
- [41] S. Fichtel, G. von Gersdorff, O. Kepka, B. Lenzi, C. Royon, and M. Saimpert, Probing new physics in diphoton production with proton tagging at the Large Hadron Collider, *Phys. Rev. D* **89**, 114004 (2014); S. Fichtel, G. von Gersdorff, B. Lenzi, C. Royon, and M. Saimpert, Light-by-light scattering with intact protons at the LHC: From Standard Model to new physics, *J. High Energy Phys.* **02** (2015) 165.
- [42] A. Falkowski, D. M. Straub, and A. Vicente, Vector-like leptons: Higgs decays and collider phenomenology, *J. High Energy Phys.* **05** (2014) 092.
- [43] J. Halverson, N. Orlofsky, and A. Pierce, Vectorlike leptons as the tip of the dark matter Iceberg, *Phys. Rev. D* **90**, 015002 (2014).
- [44] B. A. Dobrescu and C. Frugiuele, Hidden GeV-Scale Interactions of Quarks, *Phys. Rev. Lett.* **113**, 061801 (2014).
- [45] S. A. R. Ellis, R. M. Godbole, S. Gopalakrishna, and J. D. Wells, Survey of vector-like fermion extensions of the Standard Model and their phenomenological implications, *J. High Energy Phys.* **09** (2014) 130.
- [46] A. Falkowski and R. Vega-Morales, Exotic Higgs decays in the golden channel, *J. High Energy Phys.* **12** (2014) 037.
- [47] B. Holdom and M. Ratzlaff, Neglected heavy leptons at the LHC, *Phys. Rev. D* **90**, 013015 (2014).
- [48] R. Dermisek, A. Raval, and S. Shin, Effects of vectorlike leptons on $h \rightarrow 4\ell$ and the connection to the muon $g-2$ anomaly, *Phys. Rev. D* **90**, 034023 (2014).
- [49] B. A. Dobrescu and A. Martin, Interpretations of anomalous LHC events with electrons and jets, *Phys. Rev. D* **91**, 035019 (2015).
- [50] R. Dermisek, J. P. Hall, E. Lunghi, and S. Shin, Limits on vectorlike leptons from searches for anomalous production of multi-lepton events, *J. High Energy Phys.* **12** (2014) 013.
- [51] B. Holdom and M. Ratzlaff, Distinctive heavy Higgs boson decays, *Phys. Rev. D* **91**, 035031 (2015).
- [52] G. Aad *et al.* (ATLAS Collaboration), Search for heavy lepton resonances decaying to a Z boson and a lepton in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector, *J. High Energy Phys.* **09** (2015) 108.
- [53] K. Ishiwata, Z. Ligeti, and M. B. Wise, New vector-like fermions and flavor physics, *J. High Energy Phys.* **10** (2015) 027.
- [54] N. Bizot and M. Frigerio, Fermionic extensions of the Standard Model in light of the Higgs couplings, *J. High Energy Phys.* **01** (2016) 036.
- [55] R. Dermisek, E. Lunghi, and S. Shin, Two Higgs doublet model with vectorlike leptons and contributions to $pp \rightarrow WW$ and $H \rightarrow WW$, *J. High Energy Phys.* **02** (2016) 119.
- [56] S. Bhattacharya, N. Sahoo, and N. Sahu, Minimal vectorlike leptonic dark matter and signatures at the LHC, *Phys. Rev. D* **93**, 115040 (2016).
- [57] N. Kumar and S. P. Martin, Vectorlike leptons at the large hadron collider, *Phys. Rev. D* **92**, 115018 (2015).
- [58] R. Dermisek, E. Lunghi, and S. Shin, New decay modes of heavy Higgs bosons in a two Higgs doublet model with vectorlike leptons, *J. High Energy Phys.* **05** (2016) 148.
- [59] M. Abdullah, J. L. Feng, S. Iwamoto, and B. Lillard, Heavy bino dark matter and collider signals in the MSSM with vectorlike fourth-generation particles, *Phys. Rev. D* **94**, 095018 (2016).
- [60] R. Dermisek, E. Lunghi, and S. Shin, New constraints and discovery potential for Higgs to Higgs cascade decays through vectorlike leptons, *J. High Energy Phys.* **10** (2016) 081.
- [61] J. L. Feng, B. Fornal, I. Galon, S. Gardner, J. Smolinsky, T. M. P. Tait, and P. Tanedo, Particle physics models for the 17 MeV anomaly in beryllium nuclear decays, *Phys. Rev. D* **95**, 035017 (2017).
- [62] A. Angelescu and G. Arcadi, Dark matter phenomenology of SM and enlarged Higgs sectors extended with vector like leptons, *Eur. Phys. J. C* **77**, 456 (2017).
- [63] S. Bahrami, M. Frank, D. K. Ghosh, N. Ghosh, and I. Saha, Dark matter and collider studies in the left-right symmetric model with vectorlike leptons, *Phys. Rev. D* **95**, 095024 (2017).
- [64] A. Choudhury, L. Darme, L. Roszkowski, E. M. Sessolo, and S. Trojanowski, Muon $g-2$ and related phenomenology in constrained vector-like extensions of the MSSM, *J. High Energy Phys.* **05** (2017) 072.
- [65] Z. Poh and S. Raby, Vectorlike leptons: Muon $g-2$ anomaly, lepton flavor violation, Higgs boson decays, and lepton nonuniversality, *Phys. Rev. D* **96**, 015032 (2017).
- [66] Q. Lu, D. E. Morrissey, and A. M. Wijangco, Higgs boson decays to dark photons through the vectorized lepton portal, *J. High Energy Phys.* **06** (2017) 138.
- [67] J. Y. Araz, S. Banerjee, M. Frank, B. Fuks, and A. Goudelis, Dark matter and collider signals in an MSSM extension with vector-like multiplets, *Phys. Rev. D* **98**, 115009 (2018).
- [68] N. F. Bell, M. J. Dolan, L. S. Friedrich, M. J. Ramsey-Musolf, and R. R. Volkas, Electroweak baryogenesis with vector-like leptons and scalar singlets, [arXiv:1903.11255](https://arxiv.org/abs/1903.11255).
- [69] S. Zheng, Minimal vectorlike model in supersymmetric unification, [arXiv:1904.10145](https://arxiv.org/abs/1904.10145).
- [70] CMS Collaboration, Search for vector-like leptons in multi-lepton final states in pp collisions at $\sqrt{s} = 13$ TeV, CERN Report No. CMS-PAS-EXO-18-005, 2018.
- [71] A. M. Sirunyan *et al.* (CMS Collaboration), Search for vector-like leptons in multilepton final states in proton-proton collisions at $\sqrt{s} = 13$ TeV, [arXiv:1905.10853](https://arxiv.org/abs/1905.10853).
- [72] Y. Cai, J. Herrero-Garcia, M. A. Schmidt, A. Vicente, and R. R. Volkas, From the trees to the forest: A review of radiative neutrino mass models, *Front. Phys.* **5**, 63 (2017).
- [73] S. Pascoli, R. Ruiz, and C. Weiland, Heavy neutrinos with dynamic jet vetoes: Multilepton searches at $\sqrt{s} = 14, 27$, and 100 TeV, *J. High Energy Phys.* **06** (2019) 049.
- [74] P. Achard *et al.* (L3 Collaboration), Search for heavy neutral and charged leptons in e^+e^- annihilation at LEP, *Phys. Lett. B* **517**, 75 (2001).
- [75] M. L. Mangano *et al.*, Physics at a 100 TeV pp collider: Standard Model processes, CERN Yellow Report No. 3, 1, 2017; T. Golling *et al.*, Physics at a 100 TeV pp collider:

- Beyond the Standard Model phenomena, CERN Yellow Report No. 3, 441, 2017; R. Contino *et al.*, Physics at a 100 TeV pp collider: Higgs and EW symmetry breaking studies, CERN Yellow Report No. 3, 255, 2017; M. Mangano, Physics at the FCC-hh, a 100 TeV pp collider, CERN Yellow Report No. CERN 2017-003-M, 2007.
- [76] CEPC-SPPC Study Group, CEPC-SPPC preliminary conceptual design report. 1. Physics and detector, Reports No. IHEP-CEPC-DR-2015-01, No. IHEP-TH-2015-01, and No. IHEP-EP-2015-01; CEPC-SPPC preliminary conceptual design report. 2. Accelerator, Reports No. IHEP-CEPC-DR-2015-01 and No. IHEP-AC-2015-01.
- [77] X. Cid Vidal *et al.*, Beyond the Standard model physics at the HL-LHC and HE-LHC, [arXiv:1812.07831](https://arxiv.org/abs/1812.07831).
- [78] T.-p. Li and Y.-q. Ma, Analysis methods for results in gamma-ray astronomy, *Astrophys. J.* **272**, 317 (1983).
- [79] R. D. Cousins, J. T. Linnemann, and J. Tucker, Evaluation of three methods for calculating statistical significance when incorporating a systematic uncertainty into a test of the background-only hypothesis for a Poisson process, *Nucl. Instrum. Methods Phys. Res., Sect. A* **595**, 480 (2008).
- [80] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, Asymptotic formulae for likelihood-based tests of new physics, *Eur. Phys. J. C* **71**, 1554 (2011).
- [81] G. Cowan, *Two Developments in Tests for Discovery: Use of Weighted Monte Carlo Events and an Improved Measure, Progress on Statistical Issues in Searches*, SLAC (2012).
- [82] M. Aaboud *et al.* (ATLAS Collaboration), Search for heavy charged long-lived particles in the ATLAS detector in 31.6 fb⁻¹ of proton-proton collision data at $\sqrt{s}=13$ TeV, *Phys. Rev. D* **99**, 092007 (2019).
- [83] A. Alloul, N. D. Christensen, C. Degrande, C. Duhr, and B. Fuks, FEYNRULES 2.0—A complete toolbox for tree-level phenomenology, *Comput. Phys. Commun.* **185**, 2250 (2014).
- [84] J. Alwall, M. Herquet, F. Maltoni, O. Mattelaer, and T. Stelzer, MADGRAPH5: Going beyond, *J. High Energy Phys.* **06** (2011) 128.
- [85] R. D. Ball, V. Bertone, S. Carrazza, L. Del Debbio, S. Forte, A. Guffanti, N. P. Hartland, and J. Rojo (NNPDF Collaboration), Parton distributions with QED corrections, *Nucl. Phys.* **B877**, 290 (2013).
- [86] T. Sjostrand, S. Mrenna, and P. Z. Skands, PYTHIA 6.4 Physics and manual, *J. High Energy Phys.* **05** (2006) 026.
- [87] T. Sjostrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, An introduction to PYTHIA 8.2, *Comput. Phys. Commun.* **191**, 159 (2015).
- [88] J. de Favereau, C. Delaere, P. Demin, A. Giammanco, V. Lemaître, A. Mertens, and M. Selvaggi (DELPHES 3 Collaboration), DELPHES 3, A modular framework for fast simulation of a generic collider experiment, *J. High Energy Phys.* **02** (2014) 057.
- [89] A. Djouadi, J. Kalinowski, and M. Spira, HDECAY: A program for Higgs boson decays in the standard model and its supersymmetric extension, *Comput. Phys. Commun.* **108**, 56 (1998).
- [90] H. K. Dreiner, H. E. Haber, and S. P. Martin, Two-component spinor techniques and Feynman rules for quantum field theory and supersymmetry, *Phys. Rep.* **494**, 1 (2010).
- [91] S. P. Martin, TASI 2011 lectures notes: Two-component fermion notation and supersymmetry, [arXiv:1205.4076](https://arxiv.org/abs/1205.4076).