

Cosmic censorship and the evolution of d -dimensional charged evaporating black holes

Hao Xu^{1,2,*}, Yen Chin Ong^{3,4,†} and Man-Hong Yung^{1,5}

¹*Institute for Quantum Science and Engineering, Department of Physics, Southern University of Science and Technology, Shenzhen 518055, China*

²*Department of Physics, University of Science and Technology of China, Hefei 230026, China*

³*Center for Gravitation and Cosmology, College of Physical Science and Technology, Yangzhou University, 180 Siwangting Road, Yangzhou City, Jiangsu Province 225002, China*

⁴*School of Aeronautics and Astronautics, Shanghai Jiao Tong University, Shanghai 200240, China*

⁵*Shenzhen Key Laboratory of Quantum Science and Engineering, Southern University of Science and Technology Shenzhen, 518055, China*



(Received 4 December 2019; accepted 19 February 2020; published 6 March 2020)

The cosmic censorship conjecture essentially states that naked singularities should not form from *generic* initial conditions. Since black hole parameters can change their values under Hawking evaporation, one has to ask whether it is possible to reach extremality by simply waiting for the black hole to evaporate. If so a slight perturbation would likely render the singularity naked. Fortunately, at least for the case of an asymptotically flat 4-dimensional Reissner-Nordström black hole, Hiscock and Weems showed that it can never reach extremality despite the fact that for a sufficiently massive black hole, its charge-to-mass ratio can increase during Hawking evaporation. Hence, cosmic censorship is never violated by Hawking emission. However, we know that under some processes, it is easier to violate cosmic censorship in higher dimensions; therefore, it is crucial to generalize the Hiscock and Weems model to dimensions above four to check cosmic censorship. We found that Hawking evaporation cannot lead to violation of cosmic censorship even in higher dimensional Reissner-Nordström spacetimes. Moreover, it seems to be more difficult to reach extremality as the number of dimensions increases.

DOI: [10.1103/PhysRevD.101.064015](https://doi.org/10.1103/PhysRevD.101.064015)

I. INTRODUCTION: HAWKING EVAPORATION AND COSMIC CENSORSHIP

Black holes hide singularities behind their horizons. Singularities themselves are problematic in general relativity, and if it ever becomes naked would render general relativity indeterministic. This is because one has to prescribe boundary conditions on the singularities, which we cannot do without understanding what they are. Nowadays we know that naked singularities can form under various circumstances [1–9], notwithstanding the (weak) cosmic censorship conjecture, which states that naked singularities should not form from generic initial conditions [10,11]. The strong cosmic censorship goes a step further, demanding that determinism should hold even inside black holes (in practice it requires the inner horizon of a black hole to be unstable).

One obvious problem is to properly understand what “generic” means in this context, and that involves the study of different mechanisms that could lead to naked singularities. It would be helpful for future research if the community can classify under what physical processes

the censorship can be violated. Since Hawking radiation can change the values of the black hole parameters (i.e., the black hole hairs; in general relativity this means mass, charge, and angular momentum), one such mechanism that should be considered is the Hawking evaporation.

Let us consider an isolated¹ asymptotically flat Reissner-Nordström black hole. The Hawking evaporation process of this type of black hole has been investigated by many previous works, e.g., [15–18], just to name a few.

¹This assumption ensures that the black hole charge is not neutralized by accreting oppositely charged particles from the surrounding environment. In an astrophysical context it has always been assumed that charge plays a minimal, if any, role. However, this might be an oversimplification, since there exists some mechanism that could “charge-up” a black hole in astrophysics (e.g., the Wald mechanism [12]), and although the amount of charges does not significantly affect the near-Schwarzschild spacetime geometry, it does affect the motion of charged particles enough to cause degeneracy with effect from rotation [13]. In addition, there exist dark matter models (e.g., minicharged dark matter) in which astrophysical black holes might be charged under a hidden U(1) symmetry, and the metric is formally the same as Kerr-Newman solution [14].

*xuh5@sustech.edu.cn
†ycong@yzu.edu.cn

Surprisingly, the charge-to-mass ratio Q/M of Reissner-Nordström black holes is not a monotonically decreasing function of time as the black holes undergo Hawking evaporation. As shown in the work of Hiscock and Weems [19], depending on the initial conditions, Q/M can actually *increase* during the course of its evaporation. Sufficiently large black holes can get very close to extremality (for Reissner-Nordström black holes, extremality is characterized by zero temperature), *even if* it initially started out with a very small charge-to-mass ratio. Suppose that these black holes can indeed become extremal under Hawking evaporation, then they could very well, under slight perturbation, become naked singularities. Of course, reaching extremality in finite time is itself a violation of the third law of black hole thermodynamics. Indeed, Davies realized a long time ago that “unattainability of absolute zero (temperature) is equivalent to the cosmic censorship hypothesis” [20]. This would spell disaster for the cosmic censorship conjecture—for any practical meaning of genericity—since a large class of black holes can evolve towards a naked singularity end state just by waiting long enough for Hawking radiation to take effect.

Fortunately, this does not occur, at least in the case of Reissner-Nordström black holes. As shown by Hiscock and Weems, although the charge-to-mass ratio Q/M might come very close to achieve extremality, it never does. Instead, the evolution would eventually turn around towards the Schwarzschild state. More specifically, on the Reissner-Nordström configuration space of $(Q/M)^2$ plotted against M , the black hole evolution curves split it into two parts separated by a narrow region which is a dissipative attractor. The attractor region approaches the Schwarzschild limit $(Q/M)^2 = 0$ when $M \rightarrow 0$, and approaches the extreme Reissner-Nordström limit $(Q/M)^2 = 1$ when $M \rightarrow \infty$. If the initial black hole is on the right of the attractor (mass dissipation zone), it will lose mass while the charge is almost unchanged—thereby increasing Q/M —until it “hits” the attractor. Then the mass and charge continue to decrease and evolve towards the Schwarzschild limit. If the initial black hole is on the left of the attractor (charge dissipation zone), it will lose charge faster than mass, and Q/M will monotonically decrease until it joins the attractor and evolves towards the Schwarzschild limit. That is to say, the attractor is approximately characterized by $dM/dt = dQ/dt$ [19,21].

Recently, the Hiscock-Weems model has been applied to asymptotically flat dilatonic charged black holes [22]. Remarkably, if we *impose* the validity of the weak cosmic censorship, then this would force us to modify the charge loss rate, the result of which agrees with direct calculation of charge loss rate via the WKB (Wentze-Kramers-Brillouin) method which considers the scattering of the charged scalar on the curved black hole background [23]. This strongly suggests that cosmic censorship does indeed hold under Hawking evaporation. (It also serves as an example that cosmic censorship can sometimes be used as a

useful principle to deduce some other physics.) In addition, a similar technique was recently applied to study charged black holes in a de Sitter universe [24].

Surprisingly, the study of charged black holes evaporation in higher dimensions using the method of Hiscock and Weems is still lacking. In view of its connection to cosmic censorship conjecture, this is a well-motivated problem. After all, we know from the literature that whether naked singularities can form from various processes does depend on the number of spacetime dimensions. For example, some of the earliest examples of naked singularity formation involves pinching-off of black strings (Gregory-Laflamme instability [5–8]) which takes place in higher dimensions. More recently, it was discovered that collisions of black holes can violate cosmic censorships in higher dimensions as energy that gets dissipated in gravitational waves decreases when dimensionality increases [9]; see also the earlier result of [25], and its recent follow-up [26]. (For a short readable account of the maximal amount of gravitational radiation emitted by black hole collision in 4-dimensions, see Hawking’s paper [27].)

Nevertheless, it is not always the case that higher dimension leads to easier violation of cosmic censorship. Recently, strong cosmic censorship of charged black holes in asymptotically de Sitter spacetime has attracted a lot of interest [28–30]. The situation in higher dimensions is not so straightforward [31]: it is indeed easier to violate strong cosmic censorship in higher dimensions, provided that the black holes are “large” (in the sense of large Λ/Λ_{\max} , where Λ is the cosmological constant, and Λ_{\max} is the value for which if $\Lambda > \Lambda_{\max}$, then the spacetime admits at most one horizon). If the black holes are small, then it becomes harder to violate the censorship. Actually the situation is even more complicated: there exists a range of Λ/Λ_{\max} for which the difficulty to violate strong cosmic censorship is not monotonic as one increases the dimensionality. For example, it could be that it is easiest to achieve this in 6-dimensions, but hardest to do so in 4-dimensions, and *then* followed by 5-dimensions (see Table I of [31]). In short, it is not easy to tell under a given process whether going to higher dimensions makes it easier to violate cosmic censorship (of either version). Therefore, although Reissner-Nordström black holes cannot attain extremality in 4-dimensions, one has to ask whether this is still true in higher dimensions, and, if so, whether the tendency to achieve extremality, in some suitable measure, is higher or lower.

Our paper is organized as follows. In the next section we will derive the differential equations describing the evolution of charged black hole as functions of time in arbitrary dimensions above four. In Sec. III we present some numerical examples of the evaporative evolution of charged black holes. In the final section we give some concluding remarks. We shall adopt the same units of Hiscock and Weems (hereinafter, HW), setting the speed of light in

vacuum c , the gravitational constant G , and the Boltzmann constant k equal to one. However the reduced Planck constant \hbar is retained explicitly.

II. HISCOCK AND WEEMS MODEL IN d -DIMENSIONS

For a d -dimensional asymptotically flat spacetime with an electric field, the action can be written as

$$S = \frac{1}{16\pi G} \int_M d^d x \sqrt{-g} [R - F^2], \quad (1)$$

where R is the Ricci scalar and F^2 the contraction of the Maxwell tensor $F^2 := F^{\mu\nu} F_{\mu\nu}$. The asymptotically flat d -dimensional ($d \geq 4$) charged black hole (Reissner-Nordström solution) can be described by the metric tensor

$$ds^2 = -f(r)dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega_{d-2}^2, \quad (2)$$

where $d\Omega_{d-2}^2$ is the canonical round metric of a unit $(d-2)$ -sphere, and the function $f(r)$ takes the form

$$f(r) = 1 - \frac{2M}{r^{d-3}} + \frac{Q^2}{r^{2d-6}}. \quad (3)$$

Here M and Q are related to the Arnowitt-Deser-Misner (ADM) mass and electric charge of the black hole (they differ from the ADM counterparts by some constant factors, the exact forms of which are not important in this work). Note that M and Q have different physical dimensions for different d . Without loss of generality we assume that $Q > 0$. The black hole radius r_+ satisfies $f(r_+) = 0$ so we have

$$r_+ = (M + \sqrt{M^2 - Q^2})^{\frac{1}{d-3}}. \quad (4)$$

The temperature is proportional to the surface gravity at the black hole horizon and is explicitly given by

$$T = \frac{(d-3)(M^2 - Q^2 + M\sqrt{M^2 - Q^2})\hbar}{2\pi(M + \sqrt{M^2 - Q^2})^{\frac{2d-5}{d-3}}}. \quad (5)$$

Now we give a brief discussion on the Stefan-Boltzmann law in $(d-1)$ -dimensional space (spatial dimension only). See also [32,33]. Considering a $(d-1)$ -dimensional cavity, the number of modes N is given by

$$dN = \frac{dx_1 dx_2 \dots dx_{d-1} dp_1 dp_2 \dots dp_{d-1}}{h^{d-1}}, \quad (6)$$

where h is the Planck constant. Integrating the above formula and multiplying the result by a factor of $(d-2)$ because of the $(d-2)$ independent polarizations of radiation (see e.g., [33–35] for more discussion in this field), we obtain

$$V \frac{(d-2)A p^{d-2} dp}{h^{d-1}} = V \frac{(d-2)A \omega^{d-2} d\omega}{(2\pi)^{d-1}}, \quad (7)$$

where

$$A = \frac{(d-1)\pi^{\frac{d-1}{2}}}{\Gamma(\frac{d+1}{2})} \quad (8)$$

is the area of an $(d-1)$ -dimensional sphere with unit radius, and V is the volume of the $(d-1)$ -dimensional cavity. For example, in 4-dimensions, we have $d = 4$ and $A = 4\pi$. After multiplying by the photon energy $\hbar\omega$ and by the Bose-Einstein factor $1/(e^{\frac{\hbar\omega}{T}} - 1)$, we obtain the $(d-1)$ -dimensional photon gas energy

$$V \frac{(d-2)A}{(2\pi)^{d-1}} \int \hbar\omega \frac{\omega^{d-2} d\omega}{e^{\frac{\hbar\omega}{T}} - 1} = V \frac{(d-2)AT^d}{(2\pi)^{d-1} \hbar^{d-1}} \int \frac{x^{d-1} dx}{e^x - 1}, \quad (9)$$

where the integration domain is from 0 to ∞ . We can find the energy density is proportional to T^d and the Stefan-Boltzmann constant reads

$$a_d = \frac{(d-2)A}{(2\pi)^{d-1} \hbar^{d-1}} \int \frac{x^{d-1} dx}{e^x - 1}. \quad (10)$$

For $d = 4$, we have the well-known result for the total energy density

$$U = \frac{\pi^2}{15\hbar^3} VT^4. \quad (11)$$

Next we need to calculate the cross section of the effective emitting surface of the black hole. We shall assume that the emitted massless particle moves along null geodesics and, following [19,36,37], we shall apply the geometrical optics approximation. Orienting the extra $(d-3)$ angular coordinates in $d\Omega_{d-2}^2$ and normalizing the affine parameter λ , the geodesic equation of the massless quanta reads

$$\left(\frac{dr}{d\lambda}\right)^2 = E^2 - J^2 \frac{f(r)}{r^2}, \quad (12)$$

where $E = f(r) \frac{dt}{d\lambda}$ and $J = r^2 \frac{d\theta}{d\lambda}$ are the energy and angular momentum, respectively. For the emitted massless particle to reach infinity rather than falling back to the black hole horizon, defining $b \equiv J/E$, one must have

$$\frac{1}{b^2} \geq \frac{f(r)}{r^2} \quad (13)$$

for any $r \geq r_+$. Therefore we need to find the maximal value of $f(r)/r^2$. In our system it corresponds to the unstable photon orbit r_p , and the impact factor b_c can be defined as $b_c \equiv r_p / \sqrt{f(r_p)}$. The exact formula for r_p and b_c are quite

complicated and we omit them here. The cross section of the black hole is just the volume of the $(d-2)$ -dimensional sphere with radius b_c , that is,

$$\sigma = \frac{\pi^{\frac{d}{2}-1}}{\Gamma(\frac{d}{2})} b_c^{d-2}. \quad (14)$$

Finally we obtain the mass loss rate of the black hole due to emission of massless particles via the simple differential equation

$$\frac{dM}{dt} = -a_d \sigma T^d. \quad (15)$$

This is the generalization of the four-dimensional Stefan-Boltzmann law in d -dimensional spacetime. Note that the Stefan-Boltzmann constant depends on spacetime dimension. With this understanding, we shall hereinafter write $a \equiv a_d$ for simplicity. The greybody factor is ignored because we are not interested in the exact lifetime of the evolution but rather its qualitative behaviors.

The charge-loss rate of the black hole, on the other hand, is described in the framework of quantum electrodynamics (QED). We can study particle creation from the vacuum by external electric fields in arbitrary spacetime dimensions. If the radius of black hole is much larger than the Compton wavelength of the electron, the production of electron-positron pairs is well described by ordinary flat-space QED. The rate of pair creation per unit d -volume reads [38]

$$\Gamma = \frac{2^{\lfloor d/2 \rfloor} - 1}{(2\pi)^{d-1}} \left(\frac{m}{\hbar}\right)^d \left(\frac{E}{E_c}\right)^{d/2} \exp\left\{\frac{-\pi E_c}{E}\right\}, \quad (16)$$

where $E_c \equiv m^2/\hbar e$ represents the Schwinger critical charge, and $\lfloor \cdot \rfloor$ denotes the floor function. For the charged black hole in d -dimensional spacetime ($d \geq 4$), the electric field² is given by

$$E = (d-3)Q/r^{d-2}. \quad (17)$$

The charge loss rate of the black hole is obtained by integrating the above formula over the entire space outside the horizon. We have

²The constant prefactor of the electric field depends on the definition of the charge and mass parameters, which differ from the ADM quantities by constant factors. The electric field is computed from the first law of black hole thermodynamics, since $dM = \Phi dQ$ gives the electric potential, from which we can compute the electric field. The fact that E seems to vanish in $(2+1)$ -dimensions is not alarming since the Reissner-Nordström metric is not a solution in $d=3$, so we do not expect the prefactor computed this way to correctly match the result obtained directly via the Gauss law of point charge for the $d=3$ case.

$$\begin{aligned} \frac{dQ}{dt} &= eA \frac{2^{\lfloor \frac{d}{2} \rfloor - 1}}{(2\pi)^{d-1}} \int_{r_+}^{\infty} \left(\frac{(d-3)Qe}{\hbar r^{d-2}}\right)^{\frac{d}{2}} r^{d-2} \\ &\quad \times \exp\left\{\frac{-\pi m^2 r^{d-2}}{(d-3)Q\hbar e}\right\} dr. \end{aligned} \quad (18)$$

The integral can be found explicitly, although this requires somewhat tedious calculations. Unlike the 4-dimensional case, we need to make use of the properties of a special function called the ‘‘WhittakerM function’’,

$$\text{WhittakerM}\left(a - \frac{1}{2}, a, z\right) = 2ae^{z/2} z^{\frac{1}{2}-a} [\Gamma(2a) - \Gamma(2a, z)], \quad (19)$$

and expand the incomplete Gamma function $\Gamma(2a, z)$ in large z ,

$$\Gamma(2a, z) = z^{2a-1} e^{-z} \sum_{k=0}^{n-1} \frac{(2a-1)(2a-2)\cdots(2a-k)}{z^k}. \quad (20)$$

Finally, we obtain the leading order term of the differential equation that governs charge loss, which reads

$$\begin{aligned} \frac{dQ}{dt} &\approx -\frac{eA 2^{\lfloor \frac{d}{2} \rfloor - 1} ((d-3)Qe)^{\frac{d}{2}+1} r_+^{-\frac{d^2}{2}+d+1}}{(2\pi)^{d-1} \pi \hbar^{\frac{d}{2}-1} m^2} \frac{1}{d-2} \\ &\quad \times \exp\left\{\frac{-\pi m^2 r_+^{d-2}}{(d-3)Q\hbar e}\right\}. \end{aligned} \quad (21)$$

This is a *universal result* in arbitrary dimensions $d \geq 4$. For $d=4$, the WhittakerM function can be described by the error function $\text{erf}(r)$, and the final result reads

$$\frac{dQ}{dt} \approx -\frac{e^4}{2\pi^3 \hbar m^2} \frac{Q^3}{r_+^3} \exp\left(\frac{-\pi m^2 r_+^2}{Q\hbar e}\right). \quad (22)$$

This is exactly the same result obtained by HW and it is also consistent with the result using the semiclassical method to investigate the tunneling of particles through the effective Dirac gap outside of the black hole [39].

The created charged particle will be repelled towards infinity. Note that since we have assumed $Q > 0$, it is actually the *positrons* that are emitted in this manner (because like charges repel). Similar discussion holds also for electrons, and so for simplicity, we shall use the term ‘‘electron’’ in the following discussions. (Since we are more used to referring to a particle rather than its antiparticle.) Since the rest mass of the electron or positron is much smaller than its electrical charge, the potential energy gained by the repelled particle, Qe/r_+^{d-3} , is also much larger than its rest mass. Consequently, the black hole mass loss rate due to the electromagnetic pair creation is

$$\frac{dM}{dt} = \frac{Q}{r_+^{d-3}} \frac{dQ}{dt}. \quad (23)$$

This is also consistent with the first law of black hole thermodynamics.

Combining Eq. (15) and Eq. (23), we finally obtain the total rate of mass loss for the charged evaporating black hole in d -dimensional spacetime:

$$\frac{dM}{dt} = -a\sigma T^d + \frac{Q}{r_+^{d-3}} \frac{dQ}{dt}. \quad (24)$$

This of course recovers the model of Hiscock and Weems when $d = 4$. Note that Eq. (24) forms a set of coupled ordinary differential equations with the charge loss rate given by Eq. (21).

A few remarks are in order before we proceed. First of all, let us explain that in this model, the black holes are required to be sufficiently large. This is because the Hawking temperature is inversely proportional to the black hole size. For small black holes, the temperature is large enough to effectively create charged particles as well, in which case it would not make sense to separate the mass loss due to massless chargeless particle and massive charged particles in the way it is done in Eq. (24). Only in the large mass regime can one model Hawking evaporation via the HW model. See also [16]. The series expansion of the charge loss formula is obtained by inserting the Eqs. (19) and (20) into (18). The leading order term is Eq. (21). This is amount to ignoring other charged particles which are heavier than the electrons (since the heavier a particle is, its production rate will be suppressed even more).

III. EVAPORATION OF HIGHER DIMENSIONAL REISSNER-NORDSTRÖM BLACK HOLES

Since we have obtained the coupled set of differential equations, Eq. (24), which can be used to describe the evaporative evolution of the charged black hole in d -dimensional spacetime, we can investigate the evolution process by numerically integrating the differential equations. Before we do that, we give a brief discussion on the validity of this method. As we emphasized earlier, the production of electron-positron pairs is well described by ordinary flat-space QED if the radius of black hole is much larger than the Compton wavelength of the electron. This requires $r_+ \gg \hbar/m$. Furthermore, the electric field of the black hole must be much smaller than the Schwinger limit E_c ; otherwise the electric field is expected to become nonlinear and the black hole will discharge rapidly. The electric field of the black hole reads

$$E = \frac{(d-3)Q}{r_+^{d-2}} = \frac{(d-3)Q}{(M + \sqrt{M^2 - Q^2})^{\frac{d-2}{d-3}}}. \quad (25)$$




FIG. 1. Evaporative evolution of the 4-dimensional asymptotically flat Reissner-Nordström black holes in the $\frac{Q^2}{M^2} - M$ diagram.

For a given black hole mass M , the electric field is a monotonic function of Q and it attains maximal value at the extremal limit $Q = M$, which reads

$$E_{\text{ext}} = \frac{d-3}{M^{\frac{1}{d-3}}}. \quad (26)$$

Thus in order to let the electric field of the black hole be much smaller than the Schwinger limit E_c , we must have

$$M \gg \left(\frac{d-3}{E_c}\right)^{d-3}. \quad (27)$$

For $d = 4$, the condition is simply $M \gg 1/E_c$. However, for the black holes in higher dimensions, their mass must be some powers of the inverse of Schwinger critical field so that they are large enough to retain a significant charge for more than a microscopic duration.

Now we are ready to investigate the mass and charge loss of these black holes. As we have discussed, in the Hiscock and Weems model, the mass loss of the black hole comes from two terms: the term $-a\sigma T^d$, which corresponds to the thermal radiation (Stefan-Boltzmann law), which depends on powers of M and Q , and the term $(Q/r_+^{d-3})dQ/dt$, which corresponds to the charge loss rate multiplied by the potential energy of the electric field, which depends exponentially on M and Q .

For a given initial mass satisfying $M_0 \gg 1/E_c^{d-3}$, if the initial charge Q_0 is small, the charge loss rate is suppressed due to the exponent, thus it is in the “mass dissipation zone”. The black hole loses its mass primarily by the




FIG. 2. Evaporative evolution of the 5-dimensional asymptotically flat Reissner-Nordström black holes in $\frac{Q^2}{M^2} - M$ diagram.

thermal radiation, until the electric field becomes larger and the charged loss is significant (though still suppressed by the Boltzmann factor). Then the black hole evolution curve joins the “attractor” in the $\frac{Q^2}{M^2} - M$ diagram, which is a very narrow region containing many different paths. Once on the attractor, the black hole will continue to lose charge and mass, and eventually evolves towards a Schwarzschild state. If the initial charge Q_0 is large, the temperature T becomes small and the black hole loses its mass primarily via charge loss, until it joins the attractor. In Fig. 1, Fig. 2,




FIG. 3. Evaporative evolution of the 6-dimensional asymptotically flat Reissner-Nordström black holes in $\frac{Q^2}{M^2} - M$ diagram.

and Fig. 3 we present some examples of the evaporative evolution of charged black holes for 4, 5, and 6 dimensional spacetimes, respectively.

Therefore, from these plots we can conclude that the qualitative features for evaporative evolution of the charged black holes are the same in higher dimensions. Although we have only plotted up to $d = 6$ dimensions, similar results probably also hold in higher dimensions since, from our analysis, it is clear that the Stefan-Boltzmann law and the Schwinger effect in different dimensions are qualitatively similar.

Note that for a fixed value of the charge-to-mass ratio Q^*/M^* close to unity, it requires that a black hole that starts from the mass dissipation zone to have higher initial mass in order to reach it. The caveat is that, in making our comparisons, we have assumed that the values of the electrical charge and mass of the electron are the same in different dimensions (the physical dimensions of charge and mass are, however, not the same in different dimensions). We are concerned with how the results in different dimensions compare, *if everything else is equal*.

IV. CONCLUSION: HAWKING EVAPORATION OBEYS COSMIC CENSORSHIP

In this work we have generalized the Hiscock and Weems model of evaporating charged black holes to higher dimensions. At the technical level, we see that it involves the WhittakerM function, whereas in $d = 4$ it only requires the error function. By solving the coupled differential equations of the mass loss and charge loss rate numerically, we have demonstrated that Reissner-Nordström black holes in higher dimensions (with explicit calculations in $d = 5, 6$) behave in qualitatively the same way as the 4-dimensional case. This means that these black holes cannot achieve extremality via Hawking evaporation, which in turn implies that they would not evolve into naked singularities.

In the Introduction, we mentioned that the role of the spacetime dimension in determining whether weak cosmic censorship is easier or harder to be violated is nontrivial. If one considers black hole collisions, then it is easier to violate cosmic censorship in higher dimensions [9], and, in view of a variety of higher dimensional cases such as black strings that form naked singularities under Gregory-Laflamme instability, it would seem to suggest that it is harder to maintain cosmic censorship in higher dimensions. Nevertheless, at this stage there is no general proof for this claim, and one needs to check it for a different physical mechanism, especially in view of the rather complicated situation in the case of the violation of the strong cosmic censorship we discussed in the Introduction.

Our work suggests that as the number of spacetime dimensions grows, the initial mass of the black hole has to be larger in order for it to evolve towards a near extremal Q/M . In this sense then, one could say that as far as Hawking radiation is concerned, cosmic censorship in

Reissner-Nordström spacetime is safest in $d = 4$. Whether this remains to be the case in other black hole spacetimes would require more studies in the future.

ACKNOWLEDGMENTS

H. X. and M.-H. Y. acknowledge the support by the Key-Area Research and Development Program of Guangdong Province (Grant No. 2018B030326001), the National Natural Science Foundation of China (Grants No. 11875160 and No. U1801661), the Natural Science Foundation of Guangdong Province (Grant

No. 2017B030308003), the Guangdong Innovative and Entrepreneurial Research Team Program (Grant No. 2016ZT06D348), the Science, Technology and Innovation Commission of Shenzhen Municipality (Grants No. JCYJ20170412152620376, No. JCYJ20170817105046702, and No. KYTDPT20181011104202253), the Economy, Trade and Information Commission of Shenzhen Municipality (Grant No. 201901161512). Y. C. O. thanks the National Natural Science Foundation of China (No. 11705162) and the Natural Science Foundation of Jiangsu Province (No. BK20170479) for funding support.

-
- [1] L. Lehner and F. Pretorius, Black Strings, Low Viscosity Fluids, and Violation of Cosmic Censorship, *Phys. Rev. Lett.* **105**, 101102 (2010).
- [2] P. Figueras, M. Kunesch, and S. Tunyasuvunakool, End Point of Black Ring Instabilities and the Weak Cosmic Censorship Conjecture, *Phys. Rev. Lett.* **116**, 071102 (2016).
- [3] O. J. C. Dias, H. S. Reall, and J. E. Santos, The BTZ black hole violates strong cosmic censorship, *J. High Energy Phys.* **12** (2019) 097.
- [4] P. Figueras, M. Kunesch, L. Lehner, and S. Tunyasuvunakool, End Point of the Ultraspinning Instability and Violation of Cosmic Censorship, *Phys. Rev. Lett.* **118**, 151103 (2017).
- [5] H. Bantilan, P. Figueras, M. Kunesch, and R. P. Macedo, The end point of nonaxisymmetric black hole instabilities in higher dimensions, *Phys. Rev. D* **100**, 086014 (2019).
- [6] R. Gregory and R. Laflamme, Black Strings and p -Branes are Unstable, *Phys. Rev. Lett.* **70**, 2837 (1993).
- [7] R. Gregory and R. Laflamme, The instability of charged black strings and p -branes, *Nucl. Phys.* **B428**, 39 (1994).
- [8] R. Gregory, The Gregory-Laflamme instability, [arXiv:1107.5821](https://arxiv.org/abs/1107.5821).
- [9] T. Andrade, R. Emparan, D. Licht, and R. Luna, Cosmic censorship violation in black hole collisions in higher dimensions, *J. High Energy Phys.* **04** (2019) 121.
- [10] R. Penrose, Gravitational Collapse and Space-Time Singularities, *Phys. Rev. Lett.* **14**, 57 (1965).
- [11] R. Penrose, Gravitational collapse: The role of general relativity, *Riv. Nuovo Cimento* **1**, 252 (1969); *Gen. Relativ. Gravit.* **34**, 1141 (2002).
- [12] R. M. Wald, Black hole in a uniform magnetic field, *Phys. Rev. D* **10**, 1680 (1974).
- [13] M. Zajaček and A. Tursunov, Electric charge of black holes: Is it really always negligible?, [arXiv:1904.04654](https://arxiv.org/abs/1904.04654).
- [14] V. Cardoso, C. F. B. Macedo, P. Pani, and V. Ferrari, Black holes and gravitational waves in models of minicharged dark matter, *J. Cosmol. Astropart. Phys.* **05** (2016) 054.
- [15] W. T. Zaumen, Upper bound on the electric charge of a black hole, *Nature (London)* **247**, 530 (1974).
- [16] G. W. Gibbons, Vacuum polarization and the spontaneous loss of charge by black holes, *Commun. Math. Phys.* **44**, 245 (1975).
- [17] D. N. Page, Particle emission rates from a black hole: Massless particles from an uncharged, nonrotating hole, *Phys. Rev. D* **13**, 198 (1976).
- [18] E. Sorkin and T. Piran, Formation and evaporation of charged black holes, *Phys. Rev. D* **63**, 124024 (2001).
- [19] W. A. Hiscock and L. D. Weems, Evolution of charged evaporating black holes, *Phys. Rev. D* **41**, 1142 (1990).
- [20] P. C. W. Davies, The thermodynamic theory of black holes, *Proc. R. Soc. A* **353**, 499 (1977).
- [21] Y. C. Ong, The attractor of evaporating reissner-nordström black holes, [arXiv:1909.09981](https://arxiv.org/abs/1909.09981).
- [22] Y. C. Ong and Y. Yao, Charged particle production rate from cosmic censorship in dilaton black hole spacetimes, *J. High Energy Phys.* **10** (2019) 129.
- [23] K. Shiraiishi, Superradiance from a charged dilaton black hole, *Mod. Phys. Lett. A* **07**, 3449 (1992).
- [24] M. Montero, T. Van Riet, and G. Venken, Festina lente: EFT constraints from charged black hole evaporation in de Sitter, *J. High Energy Phys.* **01** (2020) 039.
- [25] H. Okawa, K.-I. Nakao, and M. Shibata, Is super-planckian physics visible?—Scattering of black holes in 5 dimensions, *Phys. Rev. D* **83**, 121501 (2011).
- [26] U. Sperhake, W. Cook, and D. Wang, The high-energy collision of black holes in higher dimensions, *Phys. Rev. D* **100**, 104046 (2019).
- [27] S. Hawking, Gravitational Radiation from Colliding Black Holes, *Phys. Rev. Lett.* **26**, 1344 (1971).
- [28] V. Cardoso, J. L. Costa, K. Destounis, P. Hintz, and A. Jansen, Quasinormal Modes and Strong Cosmic Censorship, *Phys. Rev. Lett.* **120**, 031103 (2018).
- [29] V. Cardoso, J. L. Costa, K. Destounis, P. Hintz, and A. Jansen, Strong cosmic censorship in charged black-hole spacetimes: Still subtle, *Phys. Rev. D* **98**, 104007 (2018).
- [30] M. Rahman, S. Chakraborty, S. SenGupta, and A. A. Sen, Fate of strong cosmic censorship conjecture in presence of higher spacetime dimensions, *J. High Energy Phys.* **03** (2019) 178.

- [31] H. Liu, Z. Tang, K. Destounis, B. Wang, E. Papantonopoulos, and H. Zhang, Strong cosmic censorship in higher-dimensional Reissner-Nordström-de Sitter spacetime, *J. High Energy Phys.* **03** (2019) 187.
- [32] P. T. Landsberg and A. D. Vos, The Stefan-Boltzmann constant in n -dimensional space, *J. Phys. A* **22**, 1073 (1989).
- [33] T. R. Cardoso and A. S. de Castro, The blackbody radiation in D -dimensional universes, *Rev. Bras. Ens. Fis.* **27**, 559 (2005)
- [34] I. R. Lapidus, Classical electrodynamics in a universe with two space dimensions, *Am. J. Phys.* **50**, 155 (1982).
- [35] L. C. B. Crispino, A. Higuchi, and G. E. A. Matsas, Quantization of the electromagnetic field outside static black holes and its application to low-energy phenomena, *Phys. Rev. D* **63**, 124008 (2001); Erratum, *Phys. Rev. D* **80**, 029906 (2009).
- [36] H. Xu and M.-H. Yung, Black hole evaporation in love-lock gravity with diverse dimensions, *Phys. Lett. B* **794**, 77 (2019).
- [37] V. P. Frolov and A. Zelnikov, *Introduction to Black Hole Physics* (Oxford University Press, New York, 2011).
- [38] S. P. Gavrilo and D. M. Gitman, Vacuum instability in external fields, *Phys. Rev. D* **53**, 7162 (1996).
- [39] I. B. Khriplovich, Nonthermal radiation from black holes, *Yad. Fiz.* **65**, 1292 (2002) [*Phys. At. Nucl.* **65**, 1259 (2002)].