

Spontaneous vectorization of electrically charged black holes

João M. S. Oliveira¹ and Alexandre M. Pombo²

¹*Centro de Astrofísica e Gravitação—CENTRA, Departamento de Física, Instituto Superior Técnico—IST, Universidade de Lisboa—UL, Avenida Rovisco Pais 1, 1049-001 Lisboa, Portugal*

²*Departamento de Matemática da Universidade de Aveiro and Centre for Research and Development in Mathematics and Applications (CIDMA), Campus de Santiago, 3810-183 Aveiro, Portugal*



(Received 24 December 2020; accepted 14 January 2021; published 1 February 2021)

In this work, we generalize the spontaneous scalarization phenomena in Einstein-Maxwell-scalar models to a higher spin field. The result is an Einstein-Maxwell-vector model wherein a vector field is nonminimally coupled to the Maxwell invariant by an exponential coupling function. We show that the latter guarantees the circumvention of an associated no-hair theorem when the vector field has the form of an electric field. Different than its scalar counterpart, the new *spontaneously vectorized* Reissner-Nordström (RN) black holes are, always, undercharged while being entropically preferable. The solution profile and domain of existence are presented and analyzed.

DOI: [10.1103/PhysRevD.103.044004](https://doi.org/10.1103/PhysRevD.103.044004)

I. INTRODUCTION

The recent developments in black hole (BH) detection through gravitational wave emission and imaging led to one of the most important observational advances in BH history. Never before was it possible to study highly compact objects with such precision and certainty, allowing the possibility to discern between alternative theories of gravity through BH observations. One of the key differences between these theories is the phenomena of spontaneous scalarization/vectorization, which can be contained in the more generic spontaneous tensorization phenomena.

Spontaneous scalarization has been thoroughly discussed in the literature, with the most recent development in extended scalar-tensor-Gauss-Bonnet (eSTGB) gravity [1–3]. In the latter, the model contains a scalar field nonminimally coupled to the Gauss-Bonnet gravity correction term. While this scalarization phenomenon is induced by gravity, the original spontaneous scalarization mechanism [4] was induced through the presence of matter.

Alternatively, one can nonminimally couple the scalar field to the Maxwell invariant introducing the Einstein-Maxwell-Scalar (EMS) model [5]. The mechanism was also considered for Kerr BH where scalarization can be spin-induced [6–9]. In all cases, the phenomena occur due to a tachyonic instability that emerges for certain BH configurations. The tachyonic instability induces an exponential growth of the scalar field, and the resulting BH solution is now immersed in a scalar cloud.

While the eSTGB and Kerr model seems to be more astrophysically relevant in comparison to the EMS models, the latter is much simpler to deal with, while still presenting the same phenomenological behaviors. The properties of the

various types of scalarized EMS BHs have been extensively studied [10–20].

When considering further generalizations of this mechanism, it is natural to wonder if the spontaneously growing matter can be a vector or even a tensor. In this paper, we consider the phenomenon of *spontaneous vectorization* (SV).

Vector fields and their role in extended theories of gravity have been discussed before [21,22], and examples of BHs with vector hair have also been found [23,24]. The phenomenon of vectorization was later considered in the extended vector-tensor-Gauss-Bonnet (eVTGB) theory [25–27] (the vector analogue to the eSTGB theory); in theories non-minimally coupled to matter [28,29]; and other theories of gravity [25,30]. The main idea of this paper is to consider the EMS mechanism used in [5] and extend it to an *Einstein-Maxwell-vector* (EMV) model. For that let us consider the action:

$$\mathcal{S} = \frac{1}{4\pi} \int d^4x \sqrt{-g} \left[\frac{R}{4} - \frac{f(|B|^2)}{4} F^{\mu\nu} F_{\mu\nu} - \frac{1}{4} G^{\mu\nu} G_{\mu\nu}^* \right], \quad (1)$$

where R is the Ricci tensor, the Maxwell field strength $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$, and $G_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$ which represents the field strength of a (possibly complex) vector field B_μ . While A_μ and B_μ are both vector fields, for nomenclature simplicity, from now on, we will refer to A_μ as Maxwell field and B_μ as vector field.

The vector field is nonminimally coupled to the Maxwell term through the coupling function $f(|B|^2)$, where $|B|^2 = B^\mu B_\mu^*$, and

$$f(0) = 1, \quad (2)$$

so that we may recover Einstein-Maxwell when we have a trivial field B^μ . Note that this is a straightforward generalization of the massless scalar case, where a coupling $f(\phi)$ is considered.

This paper is organized as follows: Sec. II is dedicated to an analytical study of the model (1). A no-vector-hair theorem for a spherically symmetric ansatz is presented and followed by a small discussion of the flat spacetime case in Sec. III. The existence line is presented in Sec. IV as well as the full non-linear model and quantities of interest. The numerical solutions are presented and studied in Sec. V and we conclude in Sec. VI.

II. THE MODEL

The stress-energy tensor for the model described by the action (1) is

$$\begin{aligned} T_{\mu\nu} = & f(|B|^2) \left(F_\mu^\alpha F_{\nu\alpha} - \frac{1}{4} g_{\mu\nu} F^{\alpha\beta} F_{\alpha\beta} \right) \\ & + \frac{1}{2} \left(G_\mu^\alpha G_{\nu\alpha}^* + G_\mu^{*\alpha} G_{\nu\alpha} - \frac{1}{2} g_{\mu\nu} G^{\mu\nu} G_{\mu\nu}^* \right) \\ & + \frac{1}{4} \frac{df}{d|B|^2} F^{\alpha\beta} F_{\alpha\beta} (B_\mu B_\nu^* + B_\mu^* B_\nu). \end{aligned} \quad (3)$$

Note that, compared to the scalar case, the need to consider the scalar $g^{\mu\nu} B_\mu B_\nu^*$ in the coupling $f(|B|^2)$ introduces the last term in (3). The fact that it can be negative allows the possibility of a violation of the weak energy condition.

The massless vector field B_μ equations, which is described by a (massless) Proca equation, and the Einstein equations come as:

$$\nabla_\mu G^{\mu\nu} = \frac{1}{2} \frac{df}{d|B|^2} F^2 B^\nu, \quad (4)$$

$$R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R = 2T_{\mu\nu}. \quad (5)$$

The vector field B^μ , while being massless, due to the interaction with the electromagnetic field, gains an effective mass (μ_{eff})

$$\mu_{\text{eff}}^2 = \frac{1}{2} \frac{df}{d|B|^2} F^2, \quad (6)$$

which, for certain forms of the coupling function and the electromagnetic field, can be negative. This translates into a *tachyonic instability*, i.e., for an initial trivial configuration of B_μ , corresponding to a Reissner-Nordström (RN) spacetime, a small vector field perturbation, δB_μ , grows exponentially and drives the system away from the RN solution. The result is a *spontaneously vectorized RN BH* (VRN).

For a purely electric configuration $F^2 < 0$: $\mu_{\text{eff}}^2 < 0$ requires

$$\frac{df}{d|B|^2} > 0, \quad (7)$$

and the opposite sign for a purely magnetic configuration. For a deeper study on the several possible coupling function solutions in the scalarized case see [13] (the same line of thought can be applied here). Both conditions are satisfied by a quadratic exponential coupling

$$f(|B|^2) = e^{\alpha|B|^2}. \quad (8)$$

For this coupling, spontaneous vectorization of a purely electric RN BH occurs for $\alpha > 0$.

Another important property of the model is that the Lorenz condition is not implied by the equations of motion. If we take the divergence of (4)

$$\nabla^\mu B_\mu = -\frac{\nabla^\mu (\mu_{\text{eff}}^2)}{\mu_{\text{eff}}^2} B_\mu, \quad (9)$$

which, as we can see, does not correspond to the Lorenz condition, since μ_{eff}^2 is now a function.

The metric ansatz of a static, spherically symmetric spacetime can be described by

$$ds^2 = -\sigma(r)^2 N(r) dt^2 + \frac{dr^2}{N(r)} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2),$$

$$\text{with } N(r) = 1 - \frac{2m(r)}{r}, \quad (10)$$

where $m(r)$ is the Misner-Sharp mass function [31], r is the areal radius and $\sigma(r)$ is a real metric function.

We consider the following spherically symmetric ansatz for the vector field¹

$$B(r, t) = [B_t(r) dt + i B_r(r) dr] e^{-i\omega t}, \quad (11)$$

while the Maxwell field will only have an electrical component

$$A(r) = A_t(r) dt. \quad (12)$$

III. VECTOR THEOREMS

A. No-vector-hair theorem

For this section, let us follow the work done by Herdeiro *et al.* [23] and generalize their results for our current model.

Theorem: A spherically symmetric, static, asymptotically flat and electrically charged BH spacetime, regular on

¹These are not general ansatz. The A field can, for example, have magnetic components while the B field can still have a phase term of the form $e^{i\omega_1 \theta + i\omega_2 \varphi}$ (as the coupling term in (3) restricts any other angular dependence for B).

and outside the event horizon, which solves the Einstein-Maxwell complex-Proca field equations, and for which the massive Proca field inherits the spacetime spatial symmetries but can have an harmonic time dependence of the type $e^{-i\omega t}$ with $\omega \neq 0$, cannot support a nontrivial, finite on and outside the horizon, Proca field.

To prove this argument, we will follow [32]. If the vector field is given by (11), the Proca equations. in the metric (10) are

$$\frac{d}{dr} \left[\frac{r^2(B'_t(r) - \omega B_r(r))}{\sigma(r)} \right] = \frac{\mu_{\text{eff}}^2 r^2 B_t(r)}{\sigma(r) N(r)}, \quad (13)$$

$$B'_t(r) = \omega B_r(r) \left(1 - \frac{\mu_{\text{eff}}^2 \sigma^2(r) N(r)}{\omega^2} \right). \quad (14)$$

For a BH solution, we assume the existence of an outermost horizon at $r = r_H > 0$, which requires $N(r_H) = 0$. Every $r > r_H$ surface will then be a timelike surface and $N'(r_H) > 0$. As the sign of σ is irrelevant to the equations of motion, we can, without loss of generality, consider $\sigma(r_H) > 0$.

The proof comes as follows: consider that there is a small enough region close to the horizon, $r_H < r < r_1$, for which $\mu_{\text{eff}}^2 < 0$ (which is always verified for a massless field). This means that

$$1 - \frac{\mu_{\text{eff}}^2 \sigma^2(r) N(r)}{\omega^2} > 0, \quad (15)$$

is guaranteed in this region. Equation (14) then implies that the sign of B'_t is equal to the sign of B_r . If we integrate Eq. (13) in an interval $[r, r_c] \subset [r_H, r_1]$ and replace $B'_t(r)$ by Eq. (14), we get

$$B_r(r) = -\frac{\omega}{r^2 \mu_{\text{eff}}^2 \sigma(r)} \int_r^{r_c} dr \frac{\mu_{\text{eff}}^2 r^2 B_t(r)}{\sigma(r) N(r)}, \quad (16)$$

which imposes that the sign of B_r must be opposite to the sign of $B_t(r)$.

The theorem is now proven by contradiction. As we will see ahead, B_t must be zero at the horizon. So, if $B'_t > 0$ close to the horizon, then $B_t > 0$ in this region. However, as we know from the considerations above, B'_t has the same sign as B_r implying that $B_r > 0$ which is the same sign as B_t , contradicting the equation above. The exact same reasoning applies if we consider $B'_t < 0$, meaning that the only BH solution compatible with the conditions above is when $B_t = 0 = B_r$: the Reissner-Nordström family of solutions.

This same theorem can be generalized for the case where $\mu_{\text{eff}}^2 > 0$ (for example, if we have a massive field²) in the

²If the B vector was massive, μ_{eff}^2 would instead take the form $\mu_{\text{eff}}^2 = \mu_B^2 + \frac{1}{2} \frac{df}{d|B|^2} F^2$, and α_{min} would be μ_B dependent.

region $r_h < r < r_1$. As long as this region is small enough, we can always satisfy condition (15). The fact that $N(r_H) = 0$, implies that the left-hand side (lhs) of condition (15) is very close to unity in this region. Since Eq. (16) is independent of the sign of μ_{eff}^2 , the rest of the theorem follows.

Note that, this theorem is not valid for $\omega = 0$. The latter imposes a solely r dependent vector field. In that case, we can obtain the equation for B_r from the Proca Eq. (4)

$$\nabla_t G^{tr} = 0 = \mu_{\text{eff}}^2 B^r. \quad (17)$$

Since μ_{eff}^2 is assumed to be nonzero, we have that the radial component B^r must vanish. Then, the only viable vector field ansatz is

$$B(r) = B_t(r) dt. \quad (18)$$

B. Flat spacetime electric no go theorem

Let us now consider the real ansatz (18) for the vector field. By assuming a purely electric field, given by (12), the electromagnetic equation of motion is

$$\nabla_\mu (f F^{\mu\nu}) = 0 \Rightarrow A'_t = \frac{Q}{r^2 f}. \quad (19)$$

The virial identity on flat spacetime is

$$\int_0^\infty dr \frac{1}{r^2} \left(r^4 B_t'^2 + \frac{Q^2}{f} \right) = 0. \quad (20)$$

Since both terms are always positive (for $f > 0$), we find that the virial identity can only be respected for the trivial configuration $B'_t = 0$ and $Q = 0$. When $Q = 0$, the effective mass term of B vanishes, so B gains gauge freedom and becomes a typical Maxwell field, allowing us to set $B_t = 0$.

Alternatively, we can see this through a map to a scalar field, Φ . If we consider the effective action for this configuration

$$\mathcal{S} = \frac{1}{4\pi} \int d^4x \left[-\frac{1}{2} \frac{Q^2}{f r^4} + \frac{1}{2} (\partial_r B_t)^2 \right], \quad (21)$$

and the mapping $B_t(r) \rightarrow i\Phi(r)$, one recovers the effective action for the static, spherically symmetric EMS model in flat spacetime

$$\mathcal{S} = \frac{1}{4\pi} \int d^4x \left[-\frac{1}{2} \frac{Q^2}{f r^4} - \frac{1}{2} (\partial_r \Phi)^2 \right]. \quad (22)$$

This means that the B field with the ansatz (18) acts as a *ghost* scalar field. Flat, spherically symmetric spacetime solutions with a real scalar field $\Phi(r)$ have been found for this model in [33] for an arbitrary coupling $f(\Phi)$.

These solutions are then mapped to purely imaginary $B_t(r)$ solutions which do respect the virial identity (20).

IV. SPONTANEOUS VECTORIZATION

A. Bifurcation points

In the absence of backreaction, the EMV model can be seen as a Reissner-Nordström BH that suffers a perturbation from a vector field B_μ . In this case, the line element is the same as the RN BH

$$ds^2 = -N(r)dt^2 + \frac{dr^2}{N(r)} + r^2(d\theta^2 + \sin^2\theta d\phi^2), \quad \text{with}$$

$$N(r) = 1 - \frac{2M}{r} + \frac{Q^2}{r^2}, \quad (23)$$

where $M(Q)$ is the ADM mass (electric charge) of a RN BH. In this study we will consider the full model (1), however the coupling function needs to be linearly approximated in $|B|^2$ as $f(|B|^2) = e^{\alpha|B|^2} \approx 1 + \alpha|B|^2$.

The Proca Eq. (4) that describes a nodeless, massless vector field coupled to the Maxwell invariant and has the form (18), comes as:

$$\frac{g^{rr}}{\sqrt{-g}} \partial_r [\sqrt{-g} \partial^r B_t(r)] + \alpha \frac{Q^2}{r^4} B_t(r) = 0, \quad \text{with}$$

$$\mu_{\text{eff}}^2 = -\alpha \frac{Q^2}{r^4}. \quad (24)$$

A RN solution that supports SV requires an effective mass $\mu_{\text{eff}}^2 < 0$, and field equation reduces to an eigenvalue problem in M

$$r^2 B_t'' + 2r B_t' + \frac{\alpha Q^2}{r(r-2M) + Q^2} B_t = 0. \quad (25)$$

While the equations for B^t are easier to deal with, due to the divergence at the horizon of g^{tt} the value of $B^t(r_H)$ is not well defined. At the horizon, the physical vector field obeys $B_t = g_{tt} B^t(r_H) = 0$ as well as at infinity $B_t(r \rightarrow \infty) = 0$. Close to the horizon, the vector field can be approximated as

$$B_t(r) \approx b_1(r - r_H) - b_1 \frac{r_H(\alpha - 2)Q^2}{2(r_H^2 - Q^2)} (r - r_H)^2 + \dots,$$

$$\text{with } M = \frac{Q^2 + r_H^2}{2r_H}, \quad (26)$$

The field Eq. (25) has an analytical solution that obeys the proper boundary conditions Eq. (26)

$$B_t = z_2 F_1 \left[\frac{1}{4}(3-y); \frac{1}{4}(3+y); 2; -z \right], \quad (27)$$

with

$$z = \frac{4Q^2 r_H}{(Q^2 - r_H^2)^2} \left(\frac{Q^2 r_H}{r^2} - \frac{Q^2}{r} - \frac{r_H^2}{r} + r_H \right), \quad \text{and}$$

$$y = \sqrt{4\alpha - 1}, \quad (28)$$

and ${}_2F_1$ is an hypergeometric function. Observe that $\alpha_{\min} \geq \frac{1}{4}$, which occurs for an extremal RN configuration (first bifurcation point), while $Q \rightarrow 0$ implies $\alpha \rightarrow \infty$. Observe that for each value of α and Q , Eq. (26) yields a value of M at which the VRN solution bifurcates from the RN BH. The computation of all bifurcation points for a range of α gives then the *existence line*.

B. The full nonlinear model

The set of full nonlinear field equations that result from the model (1) with the ansatz (10) and (18) are

$$m' = \frac{Nr^4 B_t'^2 - Q^2 e^{\frac{\alpha B_t^2}{N\sigma^2}} (2\alpha B_t'^2 - N\sigma^2)}{2Nr^2\sigma^2}, \quad \sigma' = -\frac{B_t'^2 \alpha Q^2 e^{\frac{\alpha B_t^2}{N\sigma^2}}}{r^3 N^2 \sigma},$$

$$A_t' = -\frac{Q\sigma e^{\frac{\alpha B_t^2}{N\sigma^2}}}{r^2}, \quad B_t'' = B_t' \left(\frac{\sigma'}{\sigma} - \frac{2}{r} \right) - \frac{\alpha Q^2 e^{\frac{\alpha B_t^2}{N\sigma^2}}}{r^4 N} B_t. \quad (29)$$

where the Maxwell potential $A_t'(r)$ is under a first integral that was used to simplify the other field equations. For notation simplicity, we let the radial dependence fall. Close to the horizon, the metric functions and vector field can be approximated by a power series as

$$m \approx \frac{r_H}{2} + \frac{b_1^2 r_H^4 + Q^2}{2r_H^2} (r - r_H) + \dots,$$

$$\sigma \approx \sigma_0 - \frac{b_1^2 r_H^3 \sigma_0^3 \alpha Q^2}{(b_1^2 r_H^4 + \sigma_0^2 (Q^2 - r_H^2))^2} (r - r_H) + \dots,$$

$$A_t \approx -\frac{Q\sigma_0}{r_H^2} (r - r_H) + \dots,$$

$$B_t \approx b_1 (r - r_H) + b_2 (r - r_H)^2 + \dots,$$

$$b_2 = b_1 \left[\frac{\alpha Q^2 \sigma_0^4 (Q^2 - r_H^2)}{2r_H (b_1^2 r_H^4 + \sigma_0^2 (Q^2 - r_H^2))^2} - \frac{1}{r_H} \right], \quad (30)$$

with b_1 the value of the vector field derivative and σ_0 the value of the σ function, at the horizon. At infinity, we impose asymptotical flatness and the metric/field functions can be approximated by

$$m \approx M + \frac{Q^2 + P^2}{2r} + \dots, \quad \sigma \approx 1 - \frac{Q^2 \alpha}{2r^2} \dots,$$

$$A_t \approx \Phi_e - \frac{Q}{r} + \dots, \quad B_t \approx \frac{P}{r}, \quad (31)$$

with Φ_e the electrostatic potential difference at infinity and P the “vector charge” obtained from the asymptotic decay.

C. Quantities of interest and Smarr law

Two horizon quantities of interest are the Hawking temperature and the horizon area

$$T_H = \frac{1}{4\pi} N'(r_H) \sigma_0, \quad A_H = 4\pi r_H^2, \quad (32)$$

these, together with the horizon vector field derivative, b_1 , and the horizon σ value, σ_0 , describe the relevant horizon data.

The variation of the ADM mass is described by the first law: $dM = T_H dS + \Phi_e dQ$. The vectorized solutions obey the Smarr law

$$M = \frac{1}{2} T_H S_H + \Phi_e Q + M_P, \quad (33)$$

where M_P is the energy stored in the surrounding vector field, which can be computed through a Komar integral

$$\begin{aligned} M_P &= - \int_{\Omega} dr d\theta d\phi \sqrt{-g} (2T'_t - T) \\ &= -4\pi \int_{r_H}^{\infty} dr \frac{Q^2(\alpha B_t^2 - N\sigma^2) - r^4 N B_t'^2}{f N r^2 \sigma}, \end{aligned} \quad (34)$$

with T the trace of the stress-energy tensor.

In addition, the solutions satisfy the virial identity, which is obtained by a Derrick-type [34] scaling argument,

$$\begin{aligned} \int_{r_H}^{\infty} dr \left[(r - r_H) \frac{\alpha Q^2 B_t^2}{2r^3 N^2 f \sigma} (N - 1) \right. \\ \left. + \frac{(2r_H - r)}{2r^3 \sigma} \left(r^4 B_t'^2 + \frac{Q^2 \sigma^2}{f} \right) \right] = 0. \end{aligned} \quad (35)$$

The generic vectorized solution unknown in closed form and a numerical approach is necessary. To solve the latter, we use an adaptative step 6(5)th order Runge-Kutta method—we compare the result of a 5th and 6th order integration and decide if the integration step must be increased or decreased—(local error of 10^{-20}), the boundary conditions are imposed through a secant shooting strategy with a tolerance of 10^{-12} in terms of the unknown parameters b_1 and σ_0 . In all the presented solutions, the virial identity gave an error of $\sim 10^{-8}$, while the Smarr law gave $\sim 10^{-4}$. Both values are limited by machine precision and converge extremely fast with the decrease of the local error.

At last, observe that the model possesses the scaling symmetry $r \rightarrow \lambda r$, $Q \rightarrow \lambda Q$ where $\lambda > 0$ is a constant. Under this scaling symmetry, all other quantities change accordingly, e.g., $M \rightarrow \lambda M$, while the coupling function

$f(|B|^2)$ remains unchanged. For the physical discussion let us introduce the reduced quantities

$$q \equiv \frac{Q}{M}, \quad a_H \equiv \frac{A_H}{16\pi M^2}, \quad t_H \equiv 8\pi T_H M. \quad (36)$$

1. Light rings

One of the most important astrophysical properties of BHs is the presence of a *light ring* (LR)—since we are dealing with spherical symmetry, the LR is, in fact, a sphere: a photon sphere. To find the LR radius, r_{LR} , of a spherical spacetime, one must consider the null geodesics ($ds^2 = 0$) of the metric ansatz (10) (the dot represents a derivative with respect to an affine parameter):

$$\dot{r}^2 = \frac{E^2}{\sigma^2} - \frac{l^2 N}{r^2}, \quad (37)$$

where E and l represent the energy and angular momentum of a photon along the geodesic. The LR is circular, implying $\dot{r} = 0$ and $\ddot{r} = 0$. The first condition relates the energy with the angular momentum of the photon $E = l\sqrt{N}\sigma/r$ while the second gives us the condition necessary to find r_{LR} :

$$\sigma \left(-2m' + \frac{2m}{r} \right) + 2 \left(1 - \frac{2m}{r} \right) (r\sigma' - \sigma) = 0. \quad (38)$$

For the RN metric we have $\sigma = 1$ and $m(r) = M - Q^2/2r$, giving us

$$r_{LR}^{RN} = \frac{3M \pm \sqrt{9M^2 - 8Q^2}}{2}. \quad (39)$$

As demonstrated in [35,36], LR always come in pairs. For a BH, one of the LR is inside and the other outside the external horizon.

V. NUMERICAL RESULTS

A. Solutions profile

Let us start by studying the generic behavior of the metric functions and the vector fields of a fundamental (nodeless) state VRN BH. In Fig. 1 is represented the radial dependence of the various field functions for an illustrative solution with $\alpha = 25$, $Q = 0.25$, $r_H = 1.0$, and a charge to mass ratio $q = 0.4271$.

A universal feature of the fundamental solutions is the existence of a bulge of B_t around the event horizon. Since regularity imposes a null vector field at r_H and infinity, the only nontrivial, nodeless vector field solutions possess a sharp increase very close to the horizon ($b_1 > 0$), reaches a maximum and then “slowly” decays as r^{-1} .

Numerical analysis shows that, for a fixed charge and mass, increasing α corresponds to an increase of the




FIG. 1. Left: graphical representation of the field functions profile and Right: density plot of the electric field strength along the equatorial plane for $\alpha = 25$, $r_H = 1.0$ and $Q = 0.25$.

magnitude of the field B_r . The distance of the maximum of B_r relative to the horizon is also observed to increase slightly along with an increase in α .

Besides, due to the coupling with the Maxwell field, there will be a modulation of the electric potential that ultimately creates a nonmonotonic electric field that can have important implications in the accretion disk formation (see Fig. 1 right). As a contrast, the electro-vacuum RN solution, as well as in the scalarized case, is monotonically crescent and the modulation associated with the latter is closer to a damping.

One other interesting characteristic of this model is the presence of a region with negative energy density. Observe the $m(r)$ profile in Fig. 1 right. In the latter, there is a valley, which corresponds to the region where the B_r reaches its peak. This can be easily understood by observing that the Komar mass (mass associated with the external vector field), due to the negative energy density term in the EM tensor (3), gives a negative contribution to the ADM mass, violating the weak energy condition.

Regarding the light ring radii, we show some values in Table I. We can see that there is an increase of the LR radius when we increase the coupling constant α and that the LR radii of vectorized BHs are smaller than the corresponding RN black holes.

B. Domain of existence

Generating several solutions allows us to obtain a region of the domain of existence for the VRN BH solutions.

TABLE I. Light ring radii for four α values with $Q = 0.25$ and $r_H = 0.52$.

| α | 6 | 8 | 10 | 12 |
|----------------------|------|------|------|------|
| r_{LR} | 0.73 | 0.77 | 0.79 | 0.80 |
| r_{LR}/r_{LR}^{RN} | 0.77 | 0.83 | 0.84 | 0.88 |

The latter is delimited by the existence line—at which $b_1 \rightarrow 0$ —and a critical line—with $b_1 \rightarrow \infty$.

Different than the scalar case, all the possible solutions are undercharged, and in fact, the critical line always has a smaller q than the existence line (see Fig. 2). For a fixed α value, one can go from the existence line to the critical line through an increase in r_H , meaning that solutions never become singular. Meanwhile, b_1 and σ_0 have a growing increase at the horizon and diverge at the critical line. We have also computed the Kretschmann scalar and observed that the solution is everywhere regular along the domain of existence, including the critical line.

Concerning the vector charge, P , one observes a monotonic increase along the domain of existence for a fixed α . While it starts at zero in the existence line, it grows to almost the double of Q at the critical line.

Through the study of the domain of existence, one also observes that an increase in α implies a smaller value of the normalized electric charge for both the existence line and the critical line. However, the latter has a faster decrease in




FIG. 2. Graphical representation of the domain of existence.




FIG. 3. Left: reduced area a_H vs reduced charge q and Right: normalized horizon temperature t_H vs q for an EMV model. (Blue line) nonvectorized RN BH, (red lines) vectorized solutions and (dotted lines) constant q lines.

q and hence the domain of existence broadens, tending to the Schwarzschild case for $\alpha \rightarrow \infty$ ($q = 0.0134$ for $\alpha = 100$).

In addition, to study the thermodynamical preference of vectorized solutions over an equivalent RN BH (see Fig. 3), we have computed the entropy (left) of both solutions and the event horizon temperature (right).

From the thermodynamical study, we observe an entropic preference of the VRN in relation to an equivalent RN BH, which can be clearly understood by the fact that the ADM Mass of the vectorized BH is smaller than the mass contained in the central BH.

Concerning the horizon temperature (see Fig. 3 right), one observes a smaller horizon temperature for the VRN solution for an equivalent RN solution. In addition, the temperature decreases as one goes further from the existence line, however never reaching extremality ($t_H = 0$). In comparison with the scalarized solutions, both solutions are entropically preferable over an equivalent RN BH. However, while in the scalarized case a solution with a higher coupling constant has a higher entropy, here the opposite occurs. For a SV configuration that allows two coupling constant values, the lower one will be entropically preferable.

The numerical results show that the vector charge, P , increases along a fixed α line—starting at zero (RN solution) and reaching a maximum P_{\max} at the critical solution. From the numerical results, P_{\max} is not very α sensitive. The black dotted lines in Fig. 3 represent lines of constant q . We see that, as we lower α , we get further away from RN and closer to the critical solution (see black marker). The latter corresponds to a minimum α value for each q , since not all α solutions achieve all possible q . This behavior also implies that, for the same q , P increases as α decreases.³

³As there is no clear change in qualitative behavior of the solutions, it is reasonable to assume, that, P increases monotonically as we get further away from the RN solution by decreasing α .

VI. CONCLUSION

In this paper, we constructed electrically charged hairy black holes through a spontaneous vectorization process of Reissner-Nordström black holes. To do so, we considered an extra vector field B_μ that is nonminimally coupled to the Maxwell field.

Our main objective was to search for stable spherically symmetric configurations. With that in mind, we first showed a no-hair theorem for a generic complex vector field ansatz, along with a flat spacetime no go theorem that was an extension of the first. Notably, this no-hair theorem does not work for a vanishing field frequency $\omega = 0$, in which case we have a real vector field with only one component.

With the last ansatz, we then attempted to construct the vectorized BH solutions. An analytical study of the model showed that, for a nonminimal coupling of the form $f(B^2) = e^{\alpha B^2}$, we have a lower bound for the coupling constant $\alpha \geq 1/4$.

The numerical results show a family of vectorized solutions bifurcating from the RN existence line, reaching a critical solution. Compared to the scalarized RN solutions, the $B_r(r)$ field has a maximum away from the horizon and then slowly decays to zero at infinity. The difference occurs due to the imposition that both at the horizon and infinity, B_t must be zero.

A caveat to the vectorized solutions is that they violate the weak energy condition in a small region where the mass function $m(r)$ decreases. Since there is an additional contribution of the interaction term to the energy in the vectorized case, a negative Komar mass outside the horizon emerges, creating a region with negative energy densities.

Another peculiar property of the vectorized solutions is that they are undercharged, $Q/M < 1$. The latter behavior is amplified by an increasing coupling constant, α , tending toward Schwarzschild when $\alpha \rightarrow \infty$.

A thermodynamical study tells us that, despite this property, the vectorized solutions are entropically preferred,

having a larger horizon area and smaller temperature than the corresponding RN black holes.

At last, we would like to point out that this is a new generalization of the EMS model [5] and, while most of the behavior can be extrapolated to the VMS model, there is no guarantee that these solutions are either perturbatively stable or dynamical preferable. Such questions are a research topic for a future and more exhaustive work.

ACKNOWLEDGMENTS

We thank comments from Carlos A. R. Herdeiro and Eugen Radu. This work has been supported by Fundação para a Ciência e a Tecnologia (FCT), within Project No. UID/MAT/04106/2019 (CIDMA), by CENTRA (FCT) strategic Project No. UID/FIS/00099/2020, by national funds (OE),

through FCT, I. P., in the scope of the framework contract foreseen in the numbers 4, 5 and 6 of the article 23, of the Decree-Law 57/2016, of August 29, changed by Law 57/2017, of July 19. J. M. S. Oliveira is supported by an FCT post-doctoral grant through the Project No. PTDC/FIS-OUT/28407/2017 and A. M. Pombo is supported by the FCT grant No. PD/BD/142842/2018. This work has further been supported by the European Union's Horizon 2020 research and innovation (RISE) programmes H2020-MSCA-RISE-2015 Grant No. StronGrHEP-690904 and H2020-MSCA-RISE-2017 Grant No. FunFiCO-777740. The authors would like to acknowledge networking support by the COST Action CA16104. At last, we acknowledge support from the Projects No. CERN/FIS-PAR/0027/2019 and No. PTDC/FIS-AST/3041/2020.

-
- [1] D. D. Doneva and S. S. Yazadjiev, New Gauss-Bonnet Black Holes with Curvature-Induced Scalarization in Extended Scalar-Tensor Theories, *Phys. Rev. Lett.* **120**, 131103 (2018).
- [2] H. O. Silva, J. Sakstein, L. Gualtieri, T. P. Sotiriou, and E. Berti, Spontaneous Scalarization of Black Holes and Compact Stars from a Gauss-Bonnet Coupling, *Phys. Rev. Lett.* **120**, 131104 (2018).
- [3] G. Antoniou, A. Bakopoulos, and P. Kanti, Evasion of No-Hair Theorems and Novel Black-Hole Solutions in Gauss-Bonnet Theories, *Phys. Rev. Lett.* **120**, 131102 (2018).
- [4] T. Damour and G. Esposito-Farese, Nonperturbative Strong Field Effects in Tensor–Scalar Theories of Gravitation, *Phys. Rev. Lett.* **70**, 2220 (1993).
- [5] C. A. Herdeiro, E. Radu, N. Sanchis-Gual, and J. A. Font, Spontaneous Scalarization of Charged Black Holes, *Phys. Rev. Lett.* **121**, 101102 (2018).
- [6] P. V. Cunha, C. A. Herdeiro, and E. Radu, Spontaneously Scalarized Kerr Black Holes in Extended Scalar-Tensor–Gauss-Bonnet Gravity, *Phys. Rev. Lett.* **123**, 011101 (2019).
- [7] C. A. R. Herdeiro, E. Radu, H. O. Silva, T. P. Sotiriou, and N. Yunes, Spin-induced scalarized black holes, *Phys. Rev. Lett.* **126**, 011103 (2021).
- [8] E. Berti, L. G. Collodel, B. Kleihaus, and J. Kunz, Spin-induced black-hole scalarization in Einstein-scalar-Gauss-Bonnet theory, *Phys. Rev. Lett.* **126**, 011104 (2021).
- [9] A. Dima, E. Barausse, N. Franchini, and T. P. Sotiriou, Spin-Induced Black Hole Spontaneous Scalarization, *Phys. Rev. Lett.* **125**, 231101 (2020).
- [10] Y. S. Myung and D.-C. Zou, Instability of Reissner–Nordström black hole in Einstein–Maxwell–scalar theory, *Eur. Phys. J. C* **79**, 1 (2019).
- [11] Y. S. Myung and D.-C. Zou, Quasinormal modes of scalarized black holes in the Einstein–Maxwell–Scalar theory, *Phys. Lett. B* **790**, 400 (2019).
- [12] P. G. Fernandes, C. A. Herdeiro, A. M. Pombo, E. Radu, and N. Sanchis-Gual, Spontaneous scalarisation of charged black holes: Coupling dependence and dynamical features, *Classical Quant. Grav.* **36**, 134002 (2019).
- [13] D. Astefanesei, C. Herdeiro, A. Pombo, and E. Radu, Einstein–Maxwell–scalar black holes: Classes of solutions, dyons and extremality, *J. High Energy Phys.* **10** (2019) 078.
- [14] P. G. Fernandes, C. A. Herdeiro, A. M. Pombo, E. Radu, and N. Sanchis-Gual, Charged black holes with axionic-type couplings: Classes of solutions and dynamical scalarization, *Phys. Rev. D* **100**, 084045 (2019).
- [15] J. L. Blázquez-Salcedo, C. A. Herdeiro, J. Kunz, A. M. Pombo, and E. Radu, Einstein–Maxwell–scalar black holes: The hot, the cold and the bald, *Phys. Lett. B* **806**, 135493 (2020).
- [16] J. L. Blázquez-Salcedo, C. A. Herdeiro, S. Kahlen, J. Kunz, A. M. Pombo, and E. Radu, Quasinormal modes of hot, cold and bald Einstein–Maxwell–scalar black holes, *arXiv:2008.11744*.
- [17] D. Astefanesei, C. Herdeiro, J. M. S. Oliveira, and E. Radu, Higher dimensional black hole scalarization, *J. High Energy Phys.* **09** (2020) 186.
- [18] S. Hod, Spontaneous scalarization of charged Reissner–Nordström black holes: Analytic treatment along the existence line, *Phys. Lett. B* **798**, 135025 (2019).
- [19] S. Hod, Reissner–Nordström black holes supporting non-minimally coupled massive scalar field configurations, *Phys. Rev. D* **101**, 104025 (2020).
- [20] M. Khodadi, A. Allahyari, S. Vagnozzi, and D. F. Mota, Black holes with scalar hair in light of the Event Horizon Telescope, *J. Cosmol. Astropart. Phys.* **09** (2020) 026.
- [21] P. F. Baum, Vector fields and Gauss-Bonnet, *Bull. Am. Math. Soc.* **76**, 1202 (1970).
- [22] R. J. Gleiser and G. Dotti, Linear stability of Einstein–Gauss-Bonnet static spacetimes: Vector and scalar perturbations, *Phys. Rev. D* **72**, 124002 (2005).
- [23] C. Herdeiro, E. Radu, and H. Runarsson, Kerr black holes with proca hair, *Classical Quant. Grav.* **33**, 154001 (2016).

- [24] Z.-Y. Fan, Black holes with vector hair, *J. High Energy Phys.* **09** (2016) 039.
- [25] L. Annulli, V. Cardoso, and L. Gualtieri, Electromagnetism and hidden vector fields in modified gravity theories: Spontaneous and induced vectorization, *Phys. Rev. D* **99**, 044038 (2019).
- [26] F. M. Ramazanoğlu, Spontaneous growth of vector fields in gravity, *Phys. Rev. D* **96**, 064009 (2017).
- [27] F. M. Ramazanoğlu, Spontaneous tensorization from curvature coupling and beyond, *Phys. Rev. D* **99**, 084015 (2019).
- [28] M. Minamitsuji, Spontaneous vectorization in the presence of vector-field coupling to matter, *Phys. Rev. D* **101**, 104044 (2020).
- [29] R. Kase, M. Minamitsuji, and S. Tsujikawa, Neutron stars with a generalized Proca hair and spontaneous vectorization, *Phys. Rev. D* **102**, 024067 (2020).
- [30] F. M. Ramazanoğlu and K. I. Ünlütürk, Generalized disformal coupling leads to spontaneous tensorization, *Phys. Rev. D* **100**, 084026 (2019).
- [31] C. W. Misner and D. H. Sharp, Relativistic equations for adiabatic, spherically symmetric gravitational collapse, *Phys. Rev.* **136**, B571 (1964).
- [32] I. Pena and D. Sudarsky, Do collapsed boson stars result in new types of black holes?, *Classical Quant. Grav.* **14**, 3131 (1997).
- [33] C. A. Herdeiro, J. a. M. Oliveira, and E. Radu, A class of solitons in Maxwell-scalar and Einstein–Maxwell-scalar models, *Eur. Phys. J. C* **80**, 23 (2020).
- [34] G. Derrick, Comments on nonlinear wave equations as models for elementary particles, *J. Math. Phys. (N.Y.)* **5**, 1252 (1964).
- [35] P. V. Cunha, E. Berti, and C. A. Herdeiro, Light-Ring Stability for Ultracompact Objects, *Phys. Rev. Lett.* **119**, 251102 (2017).
- [36] P. V. Cunha and C. A. Herdeiro, Stationary Black Holes and Light Rings, *Phys. Rev. Lett.* **124**, 181101 (2020).