

Determination of \bar{m}_b/\bar{m}_c and \bar{m}_b from $n_f = 4$ lattice QCD + QEDD. Hatton and C. T. H. Davies^{*}*SUPA, School of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, United Kingdom*

J. Koponen

*PRISMA+ Cluster of Excellence and Institute for Nuclear Physics,
Johannes Gutenberg University of Mainz, D-55128 Mainz, Germany*G. P. Lepage[†]*Laboratory for Elementary-Particle Physics, Cornell University, Ithaca, New York 14853, USA*A. T. Lytle[‡]*Department of Physics, University of Illinois, Urbana, Illinois 61801, USA*(HPQCD Collaboration)[‡]

(Received 23 February 2021; accepted 17 May 2021; published 14 June 2021)

We extend HPQCD's earlier $n_f = 2 + 1 + 1$ lattice-QCD analysis of the ratio of $\overline{\text{MS}}$ masses of the b and c quark to include results from finer lattices (down to 0.03 fm) and a new calculation of QED contributions to the mass ratio. We find that $\bar{m}_b(\mu)/\bar{m}_c(\mu) = 4.586(12)$ at renormalization scale $\mu = 3$ GeV. This result is nonperturbative. Combining it with HPQCD's recent lattice QCD + QED determination of $\bar{m}_c(3 \text{ GeV})$ gives a new value for the b -quark mass: $\bar{m}_b(3 \text{ GeV}) = 4.513(26)$ GeV. The b -mass corresponds to $\bar{m}_b(\bar{m}_b, n_f = 5) = 4.202(21)$ GeV. These results are the first based on simulations that include QED.

DOI: [10.1103/PhysRevD.103.114508](https://doi.org/10.1103/PhysRevD.103.114508)**I. INTRODUCTION**

Accurate masses for heavy quarks are important for QCD-phenomenology generally, but they will be particularly important for high-precision searches for new physics in Higgs decays [1]. In this paper we present a new result for the ratio \bar{m}_b/\bar{m}_c of the $\overline{\text{MS}}$ masses of the b and c quarks. Our analysis of the mass ratio is completely nonperturbative. This is in contrast to lattice-QCD determinations of the separate quark masses, which need QCD perturbation theory to relate $\overline{\text{MS}}$ masses to lattice quantities. Thus the (nonperturbative) mass ratio provides a nontrivial check on (perturbative) determinations of the separate masses. The ratio can also be combined with recent accurate determinations of the c -quark mass to obtain new results for the b -quark mass.

Lattice simulations of b quarks are complicated by the quark's large mass, which leads to large lattice-spacing errors when the b quarks are described by the Dirac equation (as opposed to, say, NRQCD [2]). We address this problem by using a highly improved staggered-quark discretization of the Dirac equation (HISQ) [3] that is also highly efficient, making simulations at very small lattice spacings feasible. Our previous analysis of the mass ratio [4] used lattices with spacings down to 0.06 fm but still required an extrapolation in the quark mass to reach m_b . Here we reduce the lattice spacing to 0.03 fm, where $am_b \approx 0.6$, which allows us to simulate at the b mass. Lattice spacing errors at m_b are less than 1% on our finest lattice, and we are able to remove most of that error by extrapolating from results covering a range of lattice spacings and heavy-quark masses m_h .

Our new result is accurate to about 0.25%, so it becomes important to include QED effects. We recently analyzed the QED contributions to the c quark's mass [5]. Here we adapt the methods from our earlier paper to provide the first results for QED contributions to \bar{m}_b/\bar{m}_c and \bar{m}_b . Here and in our earlier paper we use the *quenched* QED approximation, which omits contributions from photons coupling to sea quarks. The quenched approximation should capture the bulk of the QED correction in mesons whose valence

^{*}christine.davies@glasgow.ac.uk[†]g.p.lepage@cornell.edu[‡]<http://www.physics.gla.ac.uk/HPQCD>.

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

quarks are both heavy; contributions from sea quarks are expected to be an order of magnitude smaller [5].

In Sec. II we describe our general strategy and the lattice QCD simulations we employed. In Sec. III, we extract a value for \bar{m}_b/\bar{m}_c using results from simulations without QED. We then add QED effects in Sec. IV. We summarize our results for the mass ratio in Sec. V and combine them with HPQCD's recent c -quark mass to obtain a new result for the b -quark mass.

II. LATTICE QCD SIMULATIONS

We use gluon configuration sets generated on a variety of lattices (by the MILC Collaboration [6]), with $n_f = 4$ flavors of HISQ sea quark and lattice spacings ranging from 0.09 fm to 0.03 fm. These sets are described in Table I. The u and d quark masses are set equal to $m_\ell \equiv (m_u + m_d)/2$; corrections to this approximation are quadratic in the light-quark masses for our analysis, and so are negligible. We include results where the light-quark masses in the sea are tuned close to their physical value, but we also include results with much larger light-quark masses in the sea. Results from these last simulations are unphysical but are easily corrected [4]. Increasing the light-quark mass in the sea significantly reduces the cost of our analysis at small lattice spacings (because smaller lattice volumes are used).

Ignoring QED for the moment, the ratio of the b and c $\overline{\text{MS}}$ masses equals the ratio of the corresponding bare quark masses used in the lattice Lagrangian, up to corrections that vanish in the continuum limit [9],

$$\frac{\bar{m}_b(\mu)}{\bar{m}_c(\mu)} = \frac{m_b^{\text{tuned}}}{m_c^{\text{tuned}}}\Big|_{\text{latt}} + \mathcal{O}(\alpha_s(\pi/a)a^2), \quad (1)$$

where the bare masses are tuned so that the QCD simulations reproduce the experimental results for meson masses. This relationship between the $\overline{\text{MS}}$ and lattice quark

masses is nonperturbative and independent of the $\overline{\text{MS}}$ renormalization scale μ .

Pseudoscalar and vector meson masses from our simulations are listed in Table II for a variety of (valence) heavy-quark masses, ranging on the finest lattices (sets 5 and 6) approximately from the c mass to the b mass. The analysis methods for extracting these masses (and most of the results) come from [5,10]. We use multiexponential fits to calculate the masses. Fig. 1 compares the result from our fit with the effective mass values at various times for the correlator closest to the η_b mass on our finest lattice.

The quark masses am_h in Table II are what is used in the HISQ Lagrangian. The $a\tilde{m}_h$ masses are corrected to remove tree-level $(am_h)^{2n}$ errors (in the pole mass) through order $2n = 10$ [3,11],

$$a\tilde{m}_h \equiv am_h \left(1 - \frac{48}{80} \left(\frac{am_h}{2} \right)^4 + \frac{1472}{2240} \left(\frac{am_h}{2} \right)^6 + \frac{456448}{537600} \left(\frac{am_h}{2} \right)^8 - \frac{78789632}{23654400} \left(\frac{am_h}{2} \right)^{10} \right). \quad (2)$$

We write the expansion as powers of $am_h/2$ because this makes the leading coefficients roughly the same size (about 1/2). The correction is -2% at $am_h = 0.9$, which is the largest value we use.

We give results in Table I for the tuned bare c mass for each of the configurations. In each case we adjust the c mass so as to reproduce the continuum value for either the η_c mass or the J/ψ mass,

$$\begin{aligned} m_{\eta_c}^{\text{cont}} &= 2.9766(13) \text{ GeV} \\ m_{J/\psi}^{\text{cont}} &= 3.09620(20) \text{ GeV}. \end{aligned} \quad (3)$$

Here we have subtracted 7.3(1.2) MeV from the experimental value for m_{η_c} [12] to account for the fact that we are not including contributions from $c\bar{c}$ annihilation in our

TABLE I. Gluon configuration sets used in this paper. Sets are grouped by approximate lattice spacing, with lattice spacings of 0.09 fm (sets 1 and 2), 0.06 fm (sets 3 and 4), 0.045 fm (set 5), and 0.03 fm (set 6). Lattice spacings are determined from the values shown for the Wilson flow parameter w_0/a [7] where $w_0 = 0.1715(9)$ fm [8]. The sea quark masses are given in lattice units for u/d quarks (am_ℓ), s quarks (am_s), and c quarks (am_c). Tuned values for the lattice c masses are also given (in GeV). These masses are adjusted to give correct masses for either the η_c or J/ψ mesons [Eq. (3)]. The c masses are tuned using slope $d\tilde{m}_c/dm_{cc}$, which is the same (within errors) for the η_c and J/ψ . The spatial and temporal sizes of the lattices, L and T , are listed, as are the number of configurations used in our analysis (the two numbers for set 1 are for quark masses am_h below and above 0.5; the three numbers for set 2 are for the pseudoscalar correlators, the vectors with mass with mass below 0.5 and the vectors with mass above 0.5). The three polarizations were averaged for vectors. Eight time sources were used on each configuration except for set 6 where four were used.

Set	w_0/a	am_ℓ^{sea}	am_s^{sea}	am_c^{sea}	$\tilde{m}_c^{\text{tuned}}(\eta_c)$	$\tilde{m}_c^{\text{tuned}}(J/\psi)$	$d\tilde{m}_c/dm_{cc}$	L/a	T/a	N_{cfg}
1	1.9006(20)	0.0074	0.037	0.44	0.9767 (25)	0.9828 (27)	0.478 (10)	32	96	300, 504
2	1.95175(70)	0.0012	0.0363	0.432	0.9671 (25)	0.9717 (27)	0.478 (10)	64	96	311, 565, 792
3	2.8960(60)	0.0048	0.024	0.286	0.9078 (24)	0.9118 (26)	0.444 (10)	48	144	333
4	3.0170(23)	0.0008	0.022	0.26	0.8944 (23)	0.8966 (25)	0.444 (10)	96	192	100
5	3.892(12)	0.00316	0.0158	0.188	0.8646 (26)	0.8675 (29)	0.433 (10)	64	192	200
6	5.243(16)	0.00223	0.01115	0.1316	0.8234 (27)	0.8251 (29)	0.423 (10)	96	288	100

TABLE II. Lattice QCD results for the ground-state pseudoscalar and vector $h\bar{h}$ mesons in lattice units: am_{hh}^P and am_{hh}^V , respectively. Results are given for each configuration sets (Table I) and a variety of bare quark masses am_h and corrected masses $a\tilde{m}_h$ (in lattice units). The uncertainties in the meson masses are negligible compared with other errors in our analysis and so have no impact on the final results. Most of these results are from [5,10].

Set	am_h	$a\tilde{m}_h$	am_{hh}^P	am_{hh}^V
1	0.45	0.44935	1.366803 (89)	1.41567 (21)
	0.6	0.59739	1.675554 (47)	1.717437 (70)
	0.8	0.79003	2.064088 (40)	2.101542 (57)
2	0.433	0.43246	1.329290 (31)	1.378280 (54)
	0.6	0.59739	1.674264 (13)	1.715453 (32)
	0.8	0.79003	2.063015 (11)	2.099940 (26)
3	0.269	0.26895	0.885242 (56)	...
	0.274	0.27394	0.896664 (33)	0.929876 (86)
	0.4	0.39963	1.175559 (29)	1.202336 (85)
	0.5	0.49891	1.387459 (27)	1.411113 (72)
	0.6	0.59739	1.593089 (25)	1.614626 (63)
	0.7	0.69461	1.793118 (23)	1.813249 (57)
	0.8	0.79003	1.987504 (22)	2.006783 (52)
4	0.26	0.25996	0.862671 (27)	0.895702 (52)
	0.4	0.39963	1.173904 (23)	1.199806 (36)
	0.6	0.59739	1.591669 (19)	1.612586 (27)
	0.8	0.79003	1.986246 (17)	2.005047 (24)
5	0.194	0.19399	0.666821 (41)	0.692026 (59)
	0.4	0.39963	1.130722 (31)	1.147617 (40)
	0.6	0.59739	1.549098 (26)	1.562884 (32)
	0.8	0.79003	1.945787 (23)	1.958252 (27)
	0.9	0.88303	2.135642 (21)	2.147903 (25)
6	0.138	0.13800	0.496969 (42)	0.516149 (61)
	0.45	0.44935	1.201328 (29)	1.211601 (28)
	0.55	0.54828	1.410659 (27)	1.420048 (24)
	0.65	0.64619	1.614877 (24)	1.623684 (21)

simulations; this correction is determined in [5]. The analogous correction to the J/ψ mass is negligible, but we have subtracted 0.7(2) MeV from the mass to account for $c\bar{c}$ annihilation to a photon; this correction is estimated perturbatively in [5]. We extrapolate the c masses to their correct values using

$$a\tilde{m}_c^{\text{tuned}} = a\tilde{m}_c - (am_{cc}^{\text{lat}} - am_{cc}^{\text{cont}}) \frac{d\tilde{m}_c}{dm_{cc}}, \quad (4)$$

where m_{cc} is either the η_c or the J/ψ mass and the slopes (see Table I) are estimated from splines fit to the entries in Table II.

III. \bar{m}_b/\bar{m}_c WITHOUT QED

We could tune the lattice b mass the same way we tuned m_c , but we have only a few simulation results near the b , and these have significant $(am_h)^{2n}$ errors. Instead we will




FIG. 1. The effective mass plotted versus time for the $am_h = 0.65$ pseudoscalar correlator from configuration set 6. For clarity, the plot includes only every third point. The orange band and dotted line show the corresponding mass (Table II) obtained from a multiexponential fit [5,10]. The error in the fit result (orange band) is almost entirely statistical in origin. In particular, possible biases due to excited states are completely negligible (50× smaller), as is typical in fits for heavy-quark ground-state masses.

use the data in Table II to define functions that relate the ratio of quark masses to the pseudoscalar (P) or vector (V) masses,

$$\begin{aligned} m_{hh}^P &= f_{hh}^P(\bar{m}_h/\bar{m}_c) \\ m_{hh}^V &= f_{hh}^V(\bar{m}_h/\bar{m}_c), \end{aligned} \quad (5)$$

where

$$\begin{aligned} f_{hh}^P(1) &\equiv m_{\eta_c}^{\text{cont}} \\ f_{hh}^V(1) &\equiv m_{J/\psi}^{\text{cont}}. \end{aligned} \quad (6)$$

Given these functions, we then obtain two estimates for \bar{m}_b/\bar{m}_c by solving each of the equations,

$$\begin{aligned} f_{hh}^P(\bar{m}_b/\bar{m}_c) &= m_{\eta_b}^{\text{cont}} \\ f_{hh}^V(\bar{m}_b/\bar{m}_c) &= m_{\Upsilon}^{\text{cont}}, \end{aligned} \quad (7)$$

for \bar{m}_b/\bar{m}_c , where

$$\begin{aligned} m_{\eta_b}^{\text{cont}} &= 9.3987(22) \text{ GeV} \\ m_{\Upsilon}^{\text{cont}} &= 9.46030(26) \text{ GeV}. \end{aligned} \quad (8)$$

The two mass ratios should agree. Here we account for the $b\bar{b}$ annihilation contribution to the η_b mass by adding an extra error of ± 1 MeV to the experimental result [12]; this estimate is based on NRQCD perturbation theory and the meson's width [3]. The analogous contribution to the Υ mass is negligible, as is Υ annihilation via a photon.

In what follows, we first describe our lattice-QCD analysis of f_{hh}^P and f_{hh}^V , and then discuss the results.

A. Analysis

We determine the f_{hh} functions [Eq. (5)] by fitting the meson masses am_{hh} from Table II to functions of the following form:

$$am_{hh} \times (1 \pm \sigma_u) = af_{hh}(r) + am_{hh}(\delta_{a^2} + \delta_{uds}^{\text{sea}} + \delta_c^{\text{sea}}), \quad (9)$$

where

$$r \equiv \frac{a\tilde{m}_h}{a\tilde{m}_c^{\text{tuned}}} \frac{\xi_m(\tilde{m}_c^{\text{tuned}}, \delta m_{uds}^{\text{sea}})}{\xi_m(\tilde{m}_h, \delta m_{uds}^{\text{sea}})} \quad (10)$$

is the effective ratio of quark masses \tilde{m}_h/\tilde{m}_c . Pseudoscalar and vector mesons are fit separately, to determine each of f_{hh}^P and f_{hh}^V . We describe each element of the fit function in turn,

- (i) We increase the fractional error on each value of am_{hh} from Table II to $\pm\sigma_u$. The am_{hh} errors listed in the table are very small. It is impossible to fit the almost six significant digits in these data with a model as simple as we use here. So we increase the fractional error on each value to σ_u , which is then a measure of the part of the variation in the data that is *unexplained* by our model. The σ_u errors are uncorrelated from one am_{hh} to another. We use the same value for σ_u for every data point and adjust its size to maximize the Bayes Factor from the fit [13]. For the parameters and model used here, we find that

$$\sigma_u = 0.00025, \quad (11)$$

which means that our model explains the individual data points to within $\pm 0.025\%$. A simpler model would have a larger σ_u : for example, σ_u more than doubles if the ξ_m factors in ratio r are dropped (but gives consistent results within the larger errors). Note that the statistical errors listed in Table II can be neglected when σ_u is included.

- (ii) We parametrize the f_{hh} functions as splines [14] with six knots evenly spaced from $r = 1$ to 4.6 ($\approx m_b/m_c$), inclusive. The fit parameters are the function values at the knots. These functions are linear up to corrections of order $v^2/c^2 \sim 0.1\text{--}0.3$, where v is the typical velocity of the heavy quarks in the meson. Therefore we use the following priors for the values at the knots with $r > 1$:

$$f_{hh}(r_{\text{knot}}) = 1.0(2) \times \left(m_{cc} + \frac{r_{\text{knot}} - 1}{3.6} (m_{bb} - m_{cc}) \right), \quad (12)$$

where m_{cc} and m_{bb} are the continuum masses of the pseudoscalar/vector mesons composed of c and b

quarks, respectively [Eqs. (3) and (8)]. At $r = 1$, we require

$$f_{hh}(r = 1) = m_{cc}. \quad (13)$$

We choose six knots to maximize the Bayes Factor from the fit. Results obtained using 5 or 7 knots agree well with those from 6 knots, with similar or smaller errors. Doubling the width of the priors has no effect on our results.

- (iii) The ξ_m factors in the mass ratio r rescale the quark masses to correct for detuned values of the light sea quarks. From [4],

$$\xi_m(m_h, \delta m_{uds}^{\text{sea}}) = 1 + \frac{g_m}{(m_h/m_c)^\zeta} \frac{\delta m_{uds}^{\text{sea}}}{m_s}, \quad (14)$$

where

$$\delta m_{uds}^{\text{sea}} \equiv \sum_{q=u,d,s} (m_q^{\text{sea}} - m_q^{\text{tuned}}) \quad (15)$$

is the difference between the masses used in the simulation and their tuned values. The tuned masses are determined from the tuned c -quark mass using results for m_c/m_s and m_s/m_ℓ from [15]. The priors for the fit parameters are

$$g_m = 0.035(5) \quad \zeta = 0.3(1), \quad (16)$$

which come from fits described in [4].

- (iv) The largest simulation errors are from the discretization. These are suppressed by $\alpha_s(\pi/a)$ in order a^2 because we are using the HISQ formalism [3]. Beyond this order they are suppressed either by $\alpha_s(\pi/a)$ or by v^2/c^2 , since we have removed the tree-level a^{2n} errors in the quark masses using Eq. (2). The fit can not distinguish easily between α_s and v^2/c^2 since both are around 0.2 for our data, so we include only an α_s correction, modeled after Eq. (2),

$$\delta_{a^2} \equiv \alpha_s(\pi/a) \sum_{n=1}^3 f_{a^2}^n(r) \left(\frac{a\tilde{m}_h}{2} \right)^{2n}. \quad (17)$$

Here functions $f_{a^2}^n(r)$ are six-knot splines with priors at the knots (same locations as above) of

$$f_{a^2}^n(r_{\text{knot}}) = 0.0(5). \quad (18)$$

Terms beyond $n = 3$ have no effect on the fit results; keeping just the $n = 1$ term gives the same final results but with errors that are 25% smaller. The splines allow for m_h dependence in the a^{2n} corrections.

- (v) We include a^2 corrections to ξ_m since $\delta m_{uds}^{\text{sea}}$ is large for some of our configuration sets,

$$\delta_{uds}^{\text{sea}} = \alpha_s (\pi/a) f_{uds}^{\text{sea}}(r) \frac{\delta m_{uds}^{\text{sea}}}{10m_s} \left(\frac{a\bar{m}_h}{2} \right)^2, \quad (19)$$

where function $f_{uds}^{\text{sea}}(r)$ is again a six-knot spline, now with priors at the knots of

$$f_{uds}^{\text{sea}}(r_{\text{knot}}) = 0.0(1), \quad (20)$$

where the width is chosen to be somewhat larger than suggested by g_m above. Omitting this correction has negligible effect on our final results.

- (vi) We also include a correction to ξ_m from detuned c -quark masses in the sea. This correction should be small because of heavy-quark decoupling [4] — the momentum transfers in the heavy-quark mesons are too small to produce $c\bar{c}$ pairs efficiently. We include the correction,

$$\delta_c^{\text{sea}} = f_c^{\text{sea}}(r) \frac{\delta m_c^{\text{sea}}}{m_c}, \quad (21)$$

where $\delta m_c \equiv m_c - m_c^{\text{tuned}}$, and $f_c^{\text{sea}}(r)$ is a six-knot spline with

$$f_c^{\text{sea}}(r_{\text{knot}}) = 0.00(1). \quad (22)$$

We choose the width to maximize the Bayes factor from the fit. Omitting this correction has negligible effect on our final results.

The fit parameters are the values of the coefficient functions (splines) at the knots, together with g_m and ζ from the ξ_m factors [Eq. (14)]. We use the `lsqfit` PYTHON module to do the fits [16,17].

B. Results

The functions $f_{hh}^{P/V}$ obtained from the fits described in the previous section (and detailed in the Appendix) are plotted in Fig. 2, together with the data from Table II. Figure 3 shows that the model [Eq. (9)], with best-fit values for the fit parameters in the corrections (on the right-hand side), reproduces the data within errors.¹ The difference between the lattice results with and without corrections is $-0.69(23)\%$ for the highest quark mass on the finest lattice (set 6).

We can use functions $f_{hh}^{P/V}$ to extract values for the ratio of $\overline{\text{MS}}$ masses by solving Eqs. (7). We obtain

¹ χ^2 is less useful as a measure of goodness-of-fit here because we adjust σ_u to give a good fit. χ^2 per degree of freedom was 0.9 for the pseudoscalar data (26 points) and 0.8 for the vector data (25 points).




FIG. 2. Lattice QCD (without QED) results for \bar{m}_h/\bar{m}_c plotted versus the $h\bar{h}$ meson masses. Values for \bar{m}_h/\bar{m}_c are corrected as in Eq. (10). The lines, which vary in thickness, show results from the best-fit values for the functions $f_{hh}^{P/V}$; the line thickness shows the 1σ uncertainty in these functions. These functions can be reconstructed from information in the Appendix. Separate results are shown using the pseudoscalar masses m_{hh}^P (top line, squares) and the vector masses m_{hh}^V (bottom line, circles). Different colors indicate different configuration sets, with sets 6 (brown) and 5 (purple) having the largest masses, followed by sets 4 (red), 3 (green), 2 (orange) and 1 (blue), in that order. Error bars are smaller than the plot symbols.

$$\bar{m}_b/\bar{m}_c = \begin{cases} 4.578(12) & \text{from the } am_{hh}^P \\ 4.578(15) & \text{from the } am_{hh}^V, \end{cases} \quad (23)$$

independent of renormalization scale. The two estimates agree to within 0.01(23)%. The weighted average, taking account of correlations, is

$$\bar{m}_b/\bar{m}_c = 4.578(12). \quad (24)$$

We tabulate the leading uncertainties in our two results in Table III. The error budgets are similar for the two mesons




FIG. 3. Relative difference between the data for am_{hh} from Table II and the model in Eq. (9) with best-fit values for the fit parameters. Results are shown for both pseudoscalar (squares, offset right) and vector (circles, offset left) mesons, where data points for the two mesons are offset slightly in opposite directions to improve visibility.

TABLE III. Contributions to the total (1σ) error in \bar{m}_b/\bar{m}_c from QCD simulations (without QED), as a percentage of the mean value. Results are given for determinations using pseudoscalar mesons (m_{hh}^p) and vector mesons (m_{hh}^v), and for the weighted average of these results. The dominant errors come from the extrapolation to zero lattice spacing and from uncertainties in the lattice spacing. Additional errors are from residual uncertainties taken in the fit data (σ_u), uncertainties in ξ_m used to correct for unphysical sea-quark masses, uncertainties in the η_c and η_b masses, and tuning uncertainties in the sea-quark masses. The error budgets are the same when QED is included aside from an additional uncertainty of 0.03% associated with the QED corrections.

	$\bar{m}_b/\bar{m}_c[m_{hh}^p]$	$\bar{m}_b/\bar{m}_c[m_{hh}^v]$	$\bar{m}_b/\bar{m}_c[\text{avg}]$
$(am_h)^2 \rightarrow 0$	0.20	0.21	0.20
$w_0, w_0/a$	0.10	0.18	0.12
σ_u	0.12	0.12	0.09
g_m, ζ	0.05	0.05	0.05
m_{cc}	0.06	0.01	0.04
m_{bb}	0.03	0.00	0.02
$(am_h)^2 \delta m_{uds}^{\text{sea}} \rightarrow 0$	0.06	0.07	0.06
$\delta m_c^{\text{sea}} \rightarrow 0$	0.03	0.03	0.03
$d\bar{m}_c/dm_{cc}$	0.03	0.02	0.02
Total:	0.27	0.32	0.27

and are dominated by uncertainties associated with discretization errors and the lattice spacings. Doubling the widths of any of the priors associated with these uncertainties has negligible effect on the central values from our fits ($<\sigma/3$), and only doubling the discretization priors [Eq. (18)] has an appreciable impact on the final uncertainties, as expected from Table III. Omitting results from the coarsest lattices (sets 1 and 2) has negligible effect on our results ($<\sigma/10$). Omitting results from the finest lattice (set 6) increases the final uncertainties significantly (by factors of 5–6) because there is then insufficient data at large masses to constrain the six-knot splines used in the fit function; reducing the number of knots decreases the errors by a third. In either case the results agree with our final results within errors.

Finally, as discussed in [5], we expect errors from the finite lattice volume and strong-isospin breaking ($m_u \neq m_d$) in the sea to be less than 0.01% and so negligible here. We have verified this for the meson masses (using configuration sets 3A–3B and 5–7 from [5]). The Wilson flow parameter w_0 should be similarly insensitive, and we have verified this to the level of our statistical errors (0.03%) for w_0/a . See [5] for further details.

IV. ADDING QED

Adding QED complicates the analysis of \bar{m}_b/\bar{m}_c because the quarks have different QED charges and therefore different mass anomalous dimensions. Thus the nonperturbative relation in Eq. (1) is only true up to $\mathcal{O}(\alpha_{\text{QED}})$

corrections. We deal with this complication by introducing QED through two ratios R ,

$$\frac{\bar{m}_b(\mu)}{\bar{m}_c(\mu)} \Big|_{\text{QCD}}^{\text{QED}} = \frac{R(\bar{m}_b/\bar{m}_c, Q_{c,b} = 0 \rightarrow \frac{1}{3})}{R(\bar{m}_c(\mu), Q_c = \frac{1}{3} \rightarrow \frac{2}{3})} \times \frac{\bar{m}_b}{\bar{m}_c} \Big|_{\text{QCD}}. \quad (25)$$

Here,

$$R\left(\bar{m}_c(\mu), Q_c = \frac{1}{3} \rightarrow \frac{2}{3}\right) \equiv \frac{\bar{m}_c(\mu) \text{ with } Q_c = \frac{2}{3}}{\bar{m}_c(\mu) \text{ with } Q_c = \frac{1}{3}} \quad (26)$$

is the ratio of the c mass in a theory with c -quark charge $\frac{2}{3}$ to the mass in a theory with c -quark charge $\frac{1}{3}$. Similarly,

$$R\left(\bar{m}_b/\bar{m}_c, Q_{c,b} = 0 \rightarrow \frac{1}{3}\right) \equiv \frac{\bar{m}_b/\bar{m}_c \text{ with } Q_{c,b} = \frac{1}{3}}{\bar{m}_b/\bar{m}_c \text{ with } Q_{c,b} = 0}, \quad (27)$$

where the c and b charges are equal ($Q_c = Q_b$) in each case (and so the ratio is μ independent). In every case, the quark masses are tuned to reproduce the continuum meson masses in Eqs. (3) and (8). Either the pseudoscalar or vector mesons can be used; they give the same results to within the precision needed here. Only the first of the R factors [Eq. (26)] depends on the $\overline{\text{MS}}$ renormalization scale μ ; we take $\mu = 3$ GeV, following [5]. We approximate full QED by *quenched* QED, where only the valence quarks carry electric charge. This is expected to be the dominant contribution in $\mathcal{O}(\alpha_{\text{QED}})$ and is much less costly to analyze. The techniques we use for introducing QED into simulations are standard and are described in [5,10].

The R factor for the c mass [Eq. (26)] is the most important and can be inferred from our earlier result [5],

$$R\left(\bar{m}_c(3 \text{ GeV}), Q_c = 0 \rightarrow \frac{2}{3}\right) = 0.99823(17). \quad (28)$$

R is quadratic in Q_c to better than 0.01% so the QED correction ($R - 1$) required to go from charge $\frac{1}{3}$ to charge $\frac{2}{3}$ is three quarters that required to go from 0 to $\frac{2}{3}$,

$$R\left(\bar{m}_c(3 \text{ GeV}), Q_c = \frac{1}{3} \rightarrow \frac{2}{3}\right) = 0.99867(13). \quad (29)$$

The other R factor, for \bar{m}_b/\bar{m}_c , is expected to be much closer to one for two reasons: the QED corrections for the b and c masses are similar and tend to cancel in the ratio, and the charges $Q_{c,b} = \frac{1}{3}$ are smaller (and the QED effect is quadratic in the charge). To estimate the effect, we calculated the ratio R_0 of meson masses m_{hh} with and without $Q_{c,b} = \frac{1}{3}$ QED, holding the quark masses constant, for two of our configuration sets; our results are in Table IV. This quantity can be related to the R factor for \bar{m}_b/\bar{m}_c by reexpressing the R -factor in terms of lattice masses, using Eq. (1) (since $Q_c = Q_b$), and writing it as

TABLE IV. Ratio R_0 of m_{hh} with QED corrections to m_{hh} without QED corrections, evaluated at the same quark mass m_h . Results are shown for ground-state pseudoscalar and vector mesons analyzed on two configuration sets. The quark's QED charge is Q times the proton's charge; results for $Q = 2/3$ can be converted to $Q = 1/3$ by replacing R_0 with $1 + (R_0 - 1)/4$.

Set	$a\bar{m}_h$	Q	$R_0(m_{hh}^P, Q)$	$R_0(m_{hh}^V, Q)$
1	0.44935	1/3	1.0002907 (26)	1.0003409 (75)
	0.59739	1/3	1.0002612 (17)	1.0003106 (30)
	0.79003	1/3	1.0002211 (11)	1.0002669 (25)
3	0.27394	2/3	1.0015755 (48)	1.001787 (11)
	0.39963	1/3	1.0003639 (20)	1.0004081 (55)
	0.49891	1/3	1.0003404 (15)	1.0003821 (43)
	0.59739	1/3	1.0003182 (14)	1.0003703 (37)
	0.69461	1/3	1.0002978 (11)	1.0003543 (37)
	0.79003	1/3	1.00027860 (97)	1.0003412 (38)

$$R(\bar{m}_b/\bar{m}_c, Q) = 1 + \frac{\delta\bar{m}_b^Q}{\bar{m}_b} - \frac{\delta\bar{m}_c^Q}{\bar{m}_c} + \mathcal{O}(\delta\bar{m}^2), \quad (30)$$

where $\delta\bar{m}_{c,b}^Q$ are the quark mass shifts needed to hold the meson masses constant when QED is added to the simulation. The mass shifts can be calculated for different heavy-quark masses m_h from the R_0 factors in Table IV,

$$\delta\bar{m}_h^Q = (1 - R_0(m_{hh}, Q))m_{hh} \frac{d\bar{m}_h}{dm_{hh}}. \quad (31)$$

Here the derivative is estimated for each configuration by fitting a cubic spline to the $a\bar{m}_h$ values in Table II as a function of the corresponding am_{hh} values.

Values for $R(\bar{m}_h/\bar{m}_c, Q = \frac{1}{3})$ are plotted versus \bar{m}_h/\bar{m}_c in Fig. 4 for both pseudoscalar (below) and vector (above) mesons from the two configuration sets. We fit these data to a simple function suggested by QED perturbation theory,

$$R = 1 + \sum_{i=1}^3 c_i \log^i(\bar{m}_h/\bar{m}_c) + \sum_{j=1}^5 d_j (a\bar{m}_h/2)^j, \quad (32)$$

with priors $c_i = 0.000(5)$ and $d_j = 0.0(5)$. Extrapolating to the b mass gives

$$R\left(\bar{m}_b/\bar{m}_c, Q = \frac{1}{3}\right) = \begin{cases} 1.000372(90) & \text{from } m_{hh}^P \\ 1.00036(19) & \text{from } m_{hh}^V. \end{cases} \quad (33)$$

The two results agree with each other, but the corrections are too small to affect our final results significantly.²

² $R(\bar{m}_b/\bar{m}_c, Q = \frac{1}{3}) = 1.00059$ to leading order in QED perturbation theory. Our results are close to this value but also include nonperturbative corrections from QCD.




FIG. 4. Ratio $R(\bar{m}_h/\bar{m}_c, Q = \frac{1}{3})$ is plotted versus \bar{m}_h/\bar{m}_c . It is the ratio of \bar{m}_h/\bar{m}_c computed with QED charge $Q = \frac{1}{3}$ to the result without QED ($Q = 0$), where the quark masses are tuned to give the same results for m_{hh}^P (bottom, red) or m_{hh}^V (top, blue). Results are shown from configuration sets 1 (squares) and 3 (circles). Errors are smaller than the plot symbols. The blue and red shaded areas show the $\pm 1\sigma$ fits to the data [Eq. (32)].

Doubling the fit priors leaves the results unchanged. We use the larger error in the error budgets for our final result.

Including both R factors, we arrive at new results for the quark mass ratio at $\mu = 3$ GeV that include (quenched) QED,

$$\left. \frac{\bar{m}_b(3 \text{ GeV})}{\bar{m}_c(3 \text{ GeV})} \right|_{\text{QCD}_{\text{QED}}} = \begin{cases} 4.586(13) & \text{from } m_{hh}^P \\ 4.586(15) & \text{from } m_{hh}^V. \end{cases} \quad (34)$$

These again agree with each other. The weighted average, which is our final result, is

$$\left. \frac{\bar{m}_b(3 \text{ GeV})}{\bar{m}_c(3 \text{ GeV})} \right|_{\text{QCD}_{\text{QED}}} = 4.586(12). \quad (35)$$

The error budgets for these ratios are the same as those in Table III, but with an additional uncertainty of 0.03% associated with the QED correction.³ Mass ratios for other values of the renormalization scale are readily calculated using QED perturbation theory,

$$\left. \frac{\bar{m}_b(\mu)}{\bar{m}_c(\mu)} \right|_{\text{QCD}_{\text{QED}}} = \left(\frac{\mu}{3 \text{ GeV}} \right)^{\alpha_{\text{QED}}/2\pi} \left. \frac{\bar{m}_b(3 \text{ GeV})}{\bar{m}_c(3 \text{ GeV})} \right|_{\text{QCD}_{\text{QED}}}, \quad (36)$$

where the additional QED correction is negligible compared to our errors for typical values of μ . Here and elsewhere we ignore the running of α_{QED} and $\mathcal{O}(\alpha_{\text{QED}}\alpha_s)$ corrections since they are also negligible compared with our errors.

³The QED uncertainty is obtained by adding (in quadrature) the 0.013% uncertainty in Eq. (29), the 0.019% uncertainty in Eq. (33), and 0.017% for possible corrections due to quenching QED (10% of the QED correction).

V. CONCLUSIONS

In this paper we described a new calculation of the ratio of the $\overline{\text{MS}}$ masses of the b and c quarks,

$$\left. \frac{\bar{m}_b(3 \text{ GeV}, n_f = 4)}{\bar{m}_c(3 \text{ GeV}, n_f = 4)} \right|_{\text{QCD}} = 4.586(12), \quad (37)$$

where n_f is the number of quark flavors in the sea. This is the first calculation of the mass ratio based on simulations that include QED, and it makes no use of weak-coupling perturbation theory. Earlier analyses used phenomenological models to estimate QED corrections to the mass ratio, but the precision of the most recent results requires a more accurate treatment, like the one described in this paper. QED increased the mass ratio by 0.17(3)% relative to our ratio without QED [Eq. (24)], which is almost equal to the standard deviation of our final result.⁴

Our new result is consistent at the 1σ level with earlier results from $n_f = 4$ simulations that did not include QED (and so are μ independent),

$$\frac{\bar{m}_b}{\bar{m}_c} = \begin{cases} 4.578(8)(10) & \text{Fermilab/MILC/TUMQCD [15]} \\ 4.528(54) & \text{HPQCD [4].} \end{cases} \quad (38)$$

In both cases the listed uncertainties include estimates of the QED effects.⁵ Our new result and HPQCD's previous results are nonperturbative; the Fermilab/MILC/TUMQCD result relies upon perturbation theory (and heavy quark effective theory), although sensitivity to the perturbative contributions mostly cancels in the ratio. The Fermilab/MILC/TUMQCD result comes from simulations of heavy-light mesons (D_s and B_s) rather than the heavy-heavy mesons used here.

In a recent paper, HPQCD presented a new value for the c mass that includes (quenched) QED effects as we do here,

$$\bar{m}_c(3 \text{ GeV}, n_f = 4)|_{\text{QED}} = 0.9841(51) \text{ GeV}. \quad (39)$$

Combining this result with our mass ratio gives a new result for the b -quark's $\overline{\text{MS}}$ mass,

$$\bar{m}_b(3 \text{ GeV}, n_f = 4)|_{\text{QCD}} = 4.513(26) \text{ GeV}, \quad (40)$$

⁴Note that the ‘‘QED correction’’ to a QCD-only analysis depends in detail on how parameters are set in the QCD-only simulation. Since QCD without QED is not the real world, it makes a difference, for example, which hadron mass is used to tune a quark mass; and the QED correction will differ for different choices. Our QED correction is relative to the specific QCD-only theories defined in Sec. III.

⁵The second error in the Fermilab/MILC/TUMQCD result is their estimate of residual QED uncertainties not included in their main analysis (and therefore not included in the errors stated in their abstract) [15].




FIG. 5. Values for the $\overline{\text{MS}}$ mass of the b quark from lattice QCD simulations with $n_f = 2 + 1 + 1$ flavors of sea quark. Results are shown from: HPQCD '21 (this paper), Fermilab/MILC/TUMQCD [15], Gambino *et al.* [28], ETM [29], HPQCD '14 (NRQCD) [30], and HPQCD '14 (HISQ) [4]. The gray band corresponds to the top result (HPQCD '21), the only one from simulations that include QED.

which is the first based on simulations that include QED. Using perturbation theory to run to the b mass gives⁶

$$\bar{m}_b(\bar{m}_b)|_{\text{QED}} = \begin{cases} 4.209(21) \text{ GeV} & n_f = 4 \\ 4.202(21) \text{ GeV} & n_f = 5, \end{cases} \quad (41)$$

where we now include an evolution factor from QED,

$$Z_m^{\text{QED}}(\mu) = (\mu/3 \text{ GeV})^{-\alpha_{\text{QED}}/6\pi}, \quad (42)$$

with $\mu = \bar{m}_b$ (which shifts the result by less than 0.02% and so is negligible). This new result for the b quark is compared with earlier results in Fig. 5. All of these results agree to within errors.

ACKNOWLEDGMENTS

We are grateful to the MILC collaboration for the use of their configurations. We are also grateful for the use of MILC's QCD code. We have modified it to generate quenched $U(1)$ gauge fields and incorporate those into the quark propagator calculation as described here. This work used the DiRAC Data Analytic system at the University of Cambridge, operated by the University of Cambridge High Performance Computing Service on behalf of the STFC DiRAC HPC Facility ([31]). This equipment was funded by BIS National E-infrastructure capital grant (Grant No. ST/K001590/1), STFC capital Grants No. ST/H008861/1 and No. ST/H00887X/1, and STFC DiRAC Operations Grant No. ST/K00333X/1. DiRAC is part of the National E-Infrastructure. We are grateful to the Cambridge HPC support staff for assistance. Funding for

⁶We use $\alpha_{\overline{\text{MS}}}(5 \text{ GeV}, n_f = 4) = 0.2128(25)$ from [4], together with five-loop results for the beta function and mass anomalous dimension, and four-loop results for adding a flavor [18–27].

this work came from the Science and Technology Facilities Council and the National Science Foundation.

APPENDIX: BEST-FIT SPLINES

The $f_{hh}^P(r)$ function plotted in Fig. 2 can be recreated from the its values at the knot locations $r = \bar{m}_h/\bar{m}_c$,

$$f_{hh}^P(r) = \begin{cases} 2.9766(13) \\ 4.3951(34) \\ 5.7058(61) \\ 6.9835(93) \\ 8.216(14) \\ 9.435(22) \end{cases} \text{ at } r = \begin{cases} 1.0 \\ 1.72 \\ 2.44 \\ 3.16 \\ 3.88 \\ 4.6 \end{cases}, \quad (\text{A1})$$

together with the correlation matrix for these values,

$$\begin{pmatrix} 1.0000 & 0.5857 & 0.4282 & 0.3546 & 0.2838 & 0.2113 \\ 0.5857 & 1.0000 & 0.5531 & 0.5205 & 0.4244 & 0.3565 \\ 0.4282 & 0.5531 & 1.0000 & 0.6617 & 0.4088 & 0.4113 \\ 0.3546 & 0.5205 & 0.6617 & 1.0000 & 0.5206 & 0.6074 \\ 0.2838 & 0.4244 & 0.4088 & 0.5206 & 1.0000 & 0.4216 \\ 0.2113 & 0.3565 & 0.4113 & 0.6074 & 0.4216 & 1.0000 \end{pmatrix} \cdot \begin{pmatrix} 1.0000 & 0.0999 & 0.0678 & 0.0480 & 0.0403 & 0.0302 \\ 0.0999 & 1.0000 & 0.8547 & 0.6582 & 0.5779 & 0.4747 \\ 0.0678 & 0.8547 & 1.0000 & 0.7891 & 0.6215 & 0.5017 \\ 0.0480 & 0.6582 & 0.7891 & 1.0000 & 0.7001 & 0.5338 \\ 0.0403 & 0.5779 & 0.6215 & 0.7001 & 1.0000 & 0.6628 \\ 0.0302 & 0.4747 & 0.5017 & 0.5338 & 0.6628 & 1.0000 \end{pmatrix}.$$

The analogous results from the vector mesons are

$$f_{hh}^V(r) = \begin{cases} 3.09620(20) \\ 4.4836(32) \\ 5.7951(63) \\ 7.054(11) \\ 8.282(16) \\ 9.497(25) \end{cases} \text{ at } r = \begin{cases} 1.0 \\ 1.72 \\ 2.44 \\ 3.16 \\ 3.88 \\ 4.6 \end{cases}, \quad (\text{A2})$$

with correlation matrix,

-
- [1] G. P. Lepage, P. B. Mackenzie, and M. E. Peskin, [arXiv:1404.0319](https://arxiv.org/abs/1404.0319).
- [2] A. Lee, C. Monahan, R. Horgan, C. Davies, R. Dowdall, and J. Koponen (HPQCD Collaboration), *Phys. Rev. D* **87**, 074018 (2013).
- [3] E. Follana, Q. Mason, C. Davies, K. Hornbostel, G. P. Lepage, J. Shigemitsu, H. Trotter, and K. Wong (HPQCD and UKQCD Collaborations), *Phys. Rev. D* **75**, 054502 (2007).
- [4] B. Chakraborty, C. Davies, B. Galloway, P. Knecht, J. Koponen, G. C. Donald, R. J. Dowdall, G. P. Lepage, and C. McNeile, *Phys. Rev. D* **91**, 054508 (2015).
- [5] D. Hatton, C. Davies, B. Galloway, J. Koponen, G. P. Lepage, and A. Lytle (HPQCD Collaboration), *Phys. Rev. D* **102**, 054511 (2020).
- [6] A. Bazavov *et al.* (MILC Collaboration), *Phys. Rev. D* **87**, 054505 (2013).
- [7] S. Borsanyi *et al.*, *J. High Energy Phys.* **09** (2012) 010.
- [8] R. J. Dowdall, C. T. H. Davies, G. P. Lepage, and C. McNeile, *Phys. Rev. D* **88**, 074504 (2013).
- [9] Ratios of bare masses are scheme independent in pure QCD. For more discussion see: C. Davies, C. McNeile, K. Wong, E. Follana, R. Horgan, K. Hornbostel, G. P. Lepage, J. Shigemitsu, and H. Trotter, *Phys. Rev. Lett.* **104**, 132003 (2010).
- [10] D. Hatton, C. T. H. Davies, J. Koponen, G. P. Lepage, and A. T. Lytle, *Phys. Rev. D* **103**, 054512 (2021).
- [11] E. McLean, C. T. H. Davies, A. T. Lytle, and J. Koponen, *Phys. Rev. D* **99**, 114512 (2019).
- [12] P. Zyla *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **(2020)**, 083C01.
- [13] See Section V.2 on the Empirical Bayes criterion in G. P. Lepage, B. Clark, C. Davies, K. Hornbostel, P. Mackenzie, C. Morningstar, and H. Trotter, *Nucl. Phys. B, Proc. Suppl.* **106**, 12 (2002).
- [14] We use the monotonic spline described in M. Steffen, *Astron. Astrophys.* **239**, 443 (1990).
- [15] A. Bazavov *et al.* (Fermilab Lattice, MILC, and TUMQCD Collaborations), *Phys. Rev. D* **98**, 054517 (2018).
- [16] G. P. Lepage, lsqfit v. 11.7, <https://github.com/gplepage/lsqfit> (2020); <https://doi.org/10.5281/zenodo.4037174>.
- [17] The splines are implemented using the gvar PYTHON module: G. P. Lepage, gvar v. 11.9.1, <https://github.com/gplepage/gvar> (2020); <https://doi.org/10.5281/zenodo.4290884>.
- [18] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *Phys. Rev. Lett.* **118**, 082002 (2017).
- [19] F. Herzog, B. Ruijl, T. Ueda, J. A. M. Vermaseren, and A. Vogt, *J. High Energy Phys.* **02** (2017) 090.
- [20] T. Luthe, A. Maier, P. Marquard, and Y. Schroder, *J. High Energy Phys.* **10** (2017) 166.
- [21] K. G. Chetyrkin, G. Falcioni, F. Herzog, and J. A. M. Vermaseren, *J. High Energy Phys.* **10** (2017) 179; **12** (2017) 006.

- [22] P. A. Baikov, K. G. Chetyrkin, and J. H. Kühn, *J. High Energy Phys.* **10** (2014) 076.
- [23] T. Luthe, A. Maier, P. Marquard, and Y. Schröder, *J. High Energy Phys.* **01** (2017) 081.
- [24] Y. Schroder and M. Steinhauser, *J. High Energy Phys.* **01** (2006) 051.
- [25] B. A. Kniehl, A. V. Kotikov, A. I. Onishchenko, and O. L. Veretin, *Phys. Rev. Lett.* **97**, 042001 (2006).
- [26] K. G. Chetyrkin, J. H. Kuhn, and C. Sturm, *Nucl. Phys.* **B744**, 121 (2006).
- [27] T. Liu and M. Steinhauser, *Phys. Lett. B* **746**, 330 (2015).
- [28] P. Gambino, A. Melis, and S. Simula, *Phys. Rev. D* **96**, 014511 (2017).
- [29] A. Bussone *et al.* (ETM Collaboration), *Phys. Rev. D* **93**, 114505 (2016).
- [30] B. Colquhoun, R. J. Dowdall, C. T. H. Davies, K. Hornbostel, and G. P. Lepage, *Phys. Rev. D* **91**, 074514 (2015).
- [31] www.dirac.ac.uk.