

## Macdonald indices for four-dimensional $\mathcal{N} = 3$ theories

Prarit Agarwal<sup>⊗,\*</sup>, Enrico Andriolo<sup>⊗,†</sup>, Gergely Kántor<sup>⊗,‡</sup>, and Constantinos Papageorgakis<sup>⊗,§</sup>  
*CRST and School of Physics and Astronomy, Queen Mary University of London,  
 Mile End Road, London E1 4NS, United Kingdom*



(Received 15 March 2021; accepted 22 April 2021; published 4 June 2021)

We brute-force evaluate the vacuum character for  $\mathcal{N} = 2$  vertex operator algebras labeled by crystallographic complex reflection groups  $G(k, 1, 1) = \mathbb{Z}_k$ ,  $k = 3, 4, 6$ , and  $G(3, 1, 2)$ . For  $\mathbb{Z}_{3,4}$  and  $G(3, 1, 2)$  these vacuum characters have been conjectured to respectively reproduce the Macdonald limit of the superconformal index for rank one and rank two S-fold  $\mathcal{N} = 3$  theories in four dimensions. For the  $\mathbb{Z}_3$  case, and in the limit where the Macdonald index reduces to the Schur index, we find agreement with predictions from the literature.

DOI: [10.1103/PhysRevD.103.L121701](https://doi.org/10.1103/PhysRevD.103.L121701)

### I. INTRODUCTION

In recent years there has been intense activity pertaining to the study of superconformal theories (SCFTs) that do not admit a Lagrangian description. Theories with  $\mathcal{N} \geq 2$  superconformal symmetry are ideal for such explorations. Despite the lack of perturbative control, one can still extract nontrivial data by exploiting the large amount of symmetry, e.g., by employing the power of dualities [1,2], implementing the bootstrap program [3,4], or evaluating superconformal indices [5].

In this context, “pure”  $\mathcal{N} = 3$  SCFTs— $\mathcal{N} = 3$  theories which do not automatically enhance to  $\mathcal{N} = 4$ —were envisioned in [6,7] and engineered in string theory through the  $S_k$ -fold constructions of [8,9]. These are isolated, holographic SCFTs ( $a_{4D} = c_{4D}$ ) with an F-theory dual on an  $\text{AdS}_5 \times (S^5 \times T^2)/\mathbb{Z}_k$  background. The gravity description was used in [10,11] to evaluate the superconformal index in the large-rank limit. Candidates for additional rank-one and rank-two  $\mathcal{N} = 3$  examples were presented in [12,13], by constructing corresponding Coulomb-branch geometries via gaugings of  $\mathcal{N} = 4$  theories by a discrete subgroup of the R-symmetry and electromagnetic duality groups. The “Coulomb” limit of the superconformal index [14] and the Higgs-branch

Hilbert series for these models were evaluated in [15]; see also [16].

As  $\mathcal{N} = 3$  theories are automatically  $\mathcal{N} = 2$ , a concrete computational handle can be established through the description of a “Schur” Bogomol’nyi–Prasad–Sommerfield (BPS) subsector of any  $\mathcal{N} = 2$  4D theory [14] by a (nonunitary) vertex operator algebra (VOA) [17]. VOAs for  $\mathcal{N} = 3$  theories were initially constructed in [18,19] culminating in the work of [20]. In that reference, it was conjectured that certain VOAs labelled by non-Coxeter crystallographic complex reflection groups encode the Schur subsector of the known  $\mathcal{N} = 3$  S-fold theories. In particular, [20] gave a prescription for an elegant free-field realization of such VOAs, along the lines of [21]. By constructing the latter, one is able to recover the “Macdonald” limit of the superconformal index [14] for  $\mathcal{N} = 3$  S-fold theories, from the VOA vacuum character. See also [22] for an alternate prescription on implementing a Macdonald grading of the chiral algebra.

Albeit concrete, implementing the findings of [20] in practice quickly becomes computationally intensive. It is difficult to write down the explicit free-field realization of the relevant VOAs in all but the simplest of cases, and also to evaluate the corresponding vacuum characters in a fugacity expansion for increasing conformal weights. The goal of this short paper is to show how far one can get by implementing a brute-force approach using mathematical software, for the VOAs labeled by the complex reflection groups  $G(k, 1, 1) = \mathbb{Z}_k$ ,  $k = 3, 4, 6$  ( $\mathbb{Z}_{3,4}$  label rank-one S-fold models) and  $G(3, 1, 2)$  (labels a rank-two S-fold model). We employ the  $G(k, p, N)$  notation of [23], where  $p$  is a divisor of  $k$ ; general complex reflection groups are denoted as  $\mathbb{G}$ .

Towards that end, we reconstruct and explicitly exhibit the free-field realizations of [20] for the theories of interest. We then provide algorithms for automating the process of

\* agarwalprarit@gmail.com

† e.andriolo@qmul.ac.uk

‡ g.kantor@qmul.ac.uk

§ c.papageorgakis@qmul.ac.uk

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP<sup>3</sup>.

finding null states and for evaluating the VOA vacuum characters. Our code, appended to this paper in the Supplemental Material [24], can in principle be executed to obtain the corresponding Macdonald index at arbitrary orders in a fugacity expansion. Note however that the vacuum character computation time increases exponentially as a function of the conformal weight. Our code is also customizable—and we have clearly signposted how to do so—for the reader interested in extending it to the evaluation of vacuum characters for VOAs labeled by other complex reflection groups, once the complete free-field realization of the VOA has been found.

Our results can be used to check the conjecture of [20] against independent calculations of the Macdonald index of 4D  $\mathcal{N} = 3$  S-fold theories and vice versa. For example, a proposal for the Schur limit of the superconformal index—a special case of the Macdonald index—for the rank-one  $\mathbb{Z}_3$  S-fold theory was put forward in [25]. In that limit, their and our findings are in complete agreement.

## II. FROM VACUUM CHARACTERS TO INDICES

We begin with a minimal introduction to the results of [20], to which we refer the reader for a full account. 4D  $\mathcal{N} = 2$  SCFTs,  $\mathcal{T}$ , contain a BPS subsector that is isomorphic to a VOA,  $\chi[\mathcal{T}]$  [4]. In this correspondence, the VOA central charge  $c$  is related to the type-B Weyl-anomaly coefficient in four dimensions as  $c = -12c_{4D}$ , while the VOA vacuum character reproduces the Schur limit of the 4D superconformal index. The Schur subsector of pure  $\mathcal{N} = 3$  4D S-fold SCFTs was conjectured to be isomorphic to “ $\mathcal{N} = 2$  VOAs”—i.e., VOAs containing the 2D  $\mathcal{N} = 2$  superconformal algebra (SCA) as a subalgebra— $\mathcal{W}_{\mathbb{G}}$  labelled by non-Coxeter crystallographic complex reflection groups. In fact, [20] proposed a free-field realization of  $\mathcal{W}_{\mathbb{G}}$  in terms of a subalgebra of  $\text{rank}(\mathbb{G})$  copies of the  $\beta\gamma bc$  ghost system. This subalgebra was identified with the kernel of a screening operator,  $\mathbb{S} = \int \mathbb{J}$ , acting on the  $\beta\gamma bc$  systems. The action of  $\mathbb{S}$  is defined as  $\mathbb{S} \cdot X = \{\mathbb{J}X\}_1$ , where  $\{\mathbb{J}X\}_1$  denotes the coefficient of the order-one pole in the holomorphic OPE of the screening current  $\mathbb{J}$  with some operator  $X$ .

A nice feature of the free-field description, which we will use extensively in the evaluation of the vacuum characters, is that null states built out of strong generators are identically zero. The free-field realization also facilitates the introduction of a corresponding R-filtration, inherited from the R-symmetry of the 4D  $\mathcal{N} = 2$  SCFT.

The vacuum character of the R-filtered VOA is

$$\chi_{\mathcal{W}_{\mathbb{G}}}(q, \xi, z) := \text{Tr}(-1)^F q^h \xi^{R+r} z^m, \quad (1)$$

where  $F$  is the fermion number,  $h$  is the conformal dimension, and  $r, m$  are associated with the  $\mathfrak{gl}(1)$  outer automorphism and  $\mathfrak{gl}(1)$  subalgebra of the 2D  $\mathcal{N} = 2$  SCA  $\mathfrak{osp}(2|2)$  respectively.  $R$  is the weight under the R-filtration

TABLE I. Quantum numbers for the  $\beta_{\ell}\gamma_{\ell}b_{\ell}c_{\ell}$  ghost systems used in VOA free-field realizations. The  $\ell = 1, \dots, \text{rank}(\mathbb{G})$  labels the ghost-system species and  $p_{\ell}$  are the degrees of the fundamental invariants of  $\mathbb{G}$ . These are given for  $\mathbb{Z}_k$  by  $p_1 = k$  and for  $G(3, 1, 2)$  by  $(p_1, p_2) = (3, 6)$ .

	$h$	$m$	$r$	$R$
$\beta_{\ell}$	$\frac{1}{2}p_{\ell}$	$\frac{1}{2}p_{\ell}$	0	$\frac{1}{2}p_{\ell}$
$b_{\ell}$	$\frac{1}{2}(p_{\ell} + 1)$	$\frac{1}{2}(p_{\ell} - 1)$	$+\frac{1}{2}$	$1 - \frac{1}{2}p_{\ell}$
$c_{\ell}$	$-\frac{1}{2}(p_{\ell} - 1)$	$-\frac{1}{2}(p_{\ell} - 1)$	$-\frac{1}{2}$	$\frac{1}{2}p_{\ell}$
$\gamma_{\ell}$	$1 - \frac{1}{2}p_{\ell}$	$-\frac{1}{2}p_{\ell}$	0	$1 - \frac{1}{2}p_{\ell}$
$\partial$	1	0	0	0

and the vacuum character is normalized so as to start with a “1” in its  $q$  expansion. This vacuum character can be further refined by taking  $\xi^{R+r} \rightarrow y^R v^r$ . In the free-field realization the  $\beta\gamma bc$  fields carry the quantum numbers presented in Table I.

Equation (1) was conjectured to correspond to the Macdonald limit of the 4D superconformal index of a theory for which  $\mathcal{W}_{\mathbb{G}}$  is the associated VOA,  $\mathcal{W}_{\mathbb{G}} = \chi[\mathcal{T}]$ . This is defined through [14]

$$\mathcal{I}_{\text{Macdonald}}^{\mathcal{T}}(q, t, z) := \text{Tr}(-1)^F q^{E-2R-r} t^{R+r} z^m, \quad (2)$$

where the trace is taken over the set of Schur operators of the 4D SCFT. Here  $E$  is the 4D conformal dimension, while  $R$  and  $r$  are charges for the Cartan generators of the  $SU(2)_R$  and  $U(1)_r$  R-symmetry groups respectively. Note that while from the point of view of this 4D  $\mathcal{N} = 2$  Macdonald index  $m$  is a quantum number for a global  $U(1)_F$ , in the full  $\mathcal{N} = 3$  description it is part of the  $U(3) \supset SU(2)_R \times U(1)_r \times U(1)_F$  R-symmetry group. We should emphasize that in order to connect with (1) one needs to redefine  $t \rightarrow \xi q$  so that

$$\mathcal{I}_{\text{Macdonald}}^{\mathcal{T}}(q, \xi, z) = \text{Tr}(-1)^F q^{E-R} \xi^{R+r} z^m. \quad (3)$$

## III. IMPLEMENTATION

We now describe the strategy behind our code, while detailed results for each case are presented in subsequent subsections. We are interested in the evaluation of the vacuum character for VOAs labeled by crystallographic, non-Coxeter complex reflection groups  $\mathbb{Z}_{3,4,6}$  and  $G(3, 1, 2)$ , and interpreting them as Macdonald indices for 4D  $\mathcal{N} = 3$  S-fold theories. To do so one needs to consider (1) and trace over all the states created by acting only with normal-ordered products and derivatives of the strong generators of the VOA on the  $\mathfrak{sl}(2)$ -invariant vacuum, up to a given conformal weight, while removing the contributions from null states. To speed up the calculation, we have taken advantage of the symmetry of the spectrum under conjugation, by only constructing

positively-charged states under the two  $\mathfrak{gl}(1)$  symmetries of the VOA.

The identification of null states is usually laborious and this is where the free-field realization becomes helpful. The construction algorithm of [20] starts with a simple free-field prescription for the generators of the 2D  $\mathcal{N} = 2$  SCA and the chiral strong generators of the  $\mathcal{W}_{\mathbf{G}}$ -algebra. One then introduces an ansatz for the remaining bosonic strong generators constructed out of all possible super-Virasoro primary, free-field combinations with the requisite quantum numbers and undetermined coefficients, and fixes the latter by closing the VOA under the OPE.

We have diverged slightly from this recipe in the following way. For the VOAs  $\chi[\mathcal{T}]$  that have appeared thus far in the literature, the OPE coefficients can be completely fixed by writing down the most general expressions for the expected set of generators and then imposing associativity by solving the Jacobi identities [19]. The W-algebra presentations of the  $\mathcal{W}_{\mathbf{G}}$  VOAs of interest to us were already given in [20] and inserting the free-field ansatz into the existing OPEs straightforwardly fixes the undetermined coefficients for all remaining strong generators (using the OPEdefs [26] and/or SOPEN2defs [27] *Mathematica* packages); we will expand upon this point on a case-by-case basis when discussing our results.

Equipped with this information we proceed to our main algorithm. In summary, all null states at a given conformal weight can be identified by looking for all possible combinations of states with the same quantum numbers that are identically zero upon using the free-field realization. This last step requires manipulating normal-ordered products of  $\beta\gamma bc$  ghosts, making heavy use of the OPEdefs [26] and ope.math [28] *Mathematica* packages. The null states that we find contain all those predicted in [20]. It is then straightforward to write down the vacuum character for given values of quantum numbers. Note that by unrefining in the  $R$  fugacity (e.g., if one were interested in the Schur index) the algorithm becomes faster; we have a dedicated section in our code for this special case.

In addition to the above method, we have cross-checked our refined  $\mathbb{Z}_{3,4}$  results using a second algorithm that makes no connection to the W-algebra presentation. This procedure constructs the VOA spectrum using all states in  $\text{rank}(\mathbf{G})$  copies of the  $\beta\gamma bc$  ghost system that lie in the kernel of a screening operator,  $\mathbb{S}$ . For  $\mathbf{G} = \mathbb{Z}_k$  this is given by [20]

$$\mathbf{J} = be^{(k^{-1}-1)(\chi+\phi)}, \quad (4)$$

where  $\chi, \phi$  are chiral bosons

$$\beta = e^{\chi+\phi}, \quad \gamma = \partial\chi e^{-\chi-\phi}, \quad (5)$$

and all expressions should be considered as normal-ordered. Such an approach is conceptually more




FIG. 1. Computation times for carrying out the calculation of the vacuum character for VOAs at different conformal weights. Several different VOAs are shown on the graph for comparison, along with the  $\mathbb{Z}_3$  theory computed using the kernel of the screening operator. We used a desktop PC with an Intel Core i7-6700K CPU clocked at 4 GHz, and 32 GB RAM.

straightforward—construct all states using free fields and then keep those in the kernel of  $\mathbb{S}$ —but is computationally more expensive as can be seen from Fig. 1. E.g., at  $h = 9/2$  one already needs to check 941 and 881 terms for  $\mathbb{Z}_3$  and  $\mathbb{Z}_4$  respectively.

#### IV. RESULTS: $\mathbf{G} = \mathbb{Z}_3$

This is a rank-one VOA with central charge  $c = -15$ . Its W-algebra presentation involves the following strong generators:  $\mathcal{T}$ ,  $\mathcal{J}$ ,  $\mathcal{G}$ , and  $\tilde{\mathcal{G}}$  from the 2D  $\mathcal{N} = 2$  SCA, as well as the chiral and antichiral generators  $\mathcal{W}_3$ ,  $\overline{\mathcal{W}}_3$ , and their superpartners  $\mathcal{G}_{\mathcal{W}_3}$  and  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_3}$ . Here  $\mathcal{G}_{\mathcal{W}_3} := \{\mathcal{G}\mathcal{W}_3\}_1$  and so on. For the explicit free-field realization in terms of a single  $\beta\gamma bc$  ghost system one starts with a prescription for the  $\mathcal{N} = 2$  SCA generators as well as for  $\mathcal{W}_3$ ,  $\mathcal{G}_{\mathcal{W}_3}$ . The ansatz for  $\overline{\mathcal{W}}_3$  contains 8 undetermined coefficients. We always count these before imposing the super-Virasoro primary constraint. Through the OPEs from the W-algebra presentation one can use it to also determine the free-field realization of  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_3}$ . We have calculated the fully refined vacuum character (1) up to  $O(q^8)$  in the accompanying “null states. nb” and have cross-checked this result using the screening-operator approach in “screening. nb” up to  $O(q^4)$ .

This VOA is expected to encode the Schur sector of a rank one, 4D S-fold  $\mathcal{N} = 3$  SCFT, with a Coulomb-branch operator of dimension  $\Delta = 3$  and trace-anomaly coefficient

$c_{4D} = \frac{5}{4}$ . Through (3) the Macdonald index of this S-fold theory—including the global  $U(1)_F$  fugacity—can be identified with the refined vacuum character. Below we only present the simpler, Schur limit of these expressions for brevity, where  $\xi \rightarrow 1$ ,  $z \rightarrow 1$ . Then:

$$\begin{aligned} \mathcal{I}_{\text{Schur}}^{\mathbb{Z}_3} = & 1 + q + q^2 + 2q^3 - 2q^{7/2} + 3q^4 - 2q^{9/2} + 4q^5 - 4q^{11/2} \\ & + 6q^6 - 6q^{13/2} + 8q^7 - 8q^{15/2} + 11q^8 + O(q^{17/2}). \end{aligned} \quad (6)$$

It is interesting to observe that the Schur index for this theory matches the expansion of the following closed-form expression up to  $O(q^{10})$ , although we currently have neither a derivation for it nor a justification for why it should hold to all orders:

$$\frac{1}{3} \sum_{\epsilon \in \mathbb{Z}_3} \frac{\epsilon}{\sqrt{q}} \text{P.E.} \left[ \frac{1}{2} i_{\mathcal{N}=4}(q) \left( \epsilon + \frac{1}{\epsilon} \right) \right]. \quad (7)$$

Here  $\text{P.E.}[f(z)] := \exp[\sum_{n=1}^{+\infty} \frac{1}{n} f(z^n)]$  is the plethystic exponential, while  $i_{\mathcal{N}=4}(q) = \frac{2q^{\frac{1}{2}}(1-q^{\frac{1}{2}})}{1-q}$  coincides with the single-letter Schur index of  $\mathcal{N} = 4$  super-Yang–Mills.

In [25], an independent argument for determining the Schur index of the  $\mathbf{G} = \mathbb{Z}_3$  S-fold theory was presented. This entailed starting from an  $\mathcal{N} = 1$  4D UV Lagrangian theory, and flowing to an interacting  $\mathcal{N} = 1$  SCFT in the IR, which can also be reached from the  $\mathbb{Z}_3$  S-fold theory via an  $\mathcal{N} = 1$  preserving marginal deformation. Upon relabeling  $q \rightarrow p^2$ , our result (6) agrees with that of [25]—listed to  $O(q^7)$ —providing a strong consistency check of both calculations.

## V. RESULTS: $\mathbf{G} = \mathbb{Z}_4$

This is a rank-one VOA with central charge  $c = -21$ . Its W-algebra presentation involves the following strong generators:  $\mathcal{T}$ ,  $\mathcal{J}$ ,  $\mathcal{G}$  and  $\tilde{\mathcal{G}}$  from the 2D  $\mathcal{N} = 2$  SCA, as well as the chiral and anti-chiral generators  $\mathcal{W}_4$ ,  $\overline{\mathcal{W}}_4$  and their superpartners  $\mathcal{G}_{\mathcal{W}_4}$  and  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_4}$ . For the explicit free-field realization in terms of a single  $\beta\gamma bc$  ghost system one starts with a prescription for the  $\mathcal{N} = 2$  SCA generators as well as for  $\mathcal{W}_4$ ,  $\mathcal{G}_{\mathcal{W}_4}$ . The ansatz for  $\overline{\mathcal{W}}_4$  contains 19 undetermined coefficients. Through the OPEs in the W-algebra presentation one can use it to also determine the free-field realization of  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_4}$ . We have calculated the fully-refined vacuum character (1) up to  $O(q^8)$  in the accompanying “null states.nb” and have cross-checked this result using the screening-operator approach in “screening.nb” up to  $O(q^4)$ .

This VOA is expected to encode the Schur sector of a rank one 4D S-fold  $\mathcal{N} = 3$  SCFT, with a Coulomb-branch operator of dimension  $\Delta = 4$  and trace-anomaly coefficient

$c_{4D} = \frac{7}{4}$ . Through (3) the Macdonald index of this S-fold theory can be identified with the vacuum character. The Schur limit of these expressions yields:

$$\begin{aligned} \mathcal{I}_{\text{Schur}}^{\mathbb{Z}_4} = & 1 + q - 2q^{3/2} + 5q^2 - 6q^{5/2} + 10q^3 - 16q^{7/2} + 27q^4 \\ & - 38q^{9/2} + 56q^5 - 86q^{11/2} + 129q^6 - 178q^{13/2} \\ & + 251q^7 - 362q^{15/2} + 511q^8 + O(q^{17/2}). \end{aligned} \quad (8)$$

In this case the W-algebra construction is such that the bosonic states always appear with integer while the fermionic ones with half-integer conformal weights. Therefore there are no cancellations between bosonic and fermionic states at each level and the chiral algebra partition function reproduces the partition function of Schur operators in the corresponding 4D  $\mathcal{N} = 3$  theory.

## VI. RESULTS: $\mathbf{G} = \mathbb{Z}_6$

This is a rank-one VOA with central charge  $c = -33$ . Its W-algebra presentation involves the following strong generators:  $\mathcal{T}$ ,  $\mathcal{J}$ ,  $\mathcal{G}$  and  $\tilde{\mathcal{G}}$  from the  $\mathcal{N} = 2$  SCA, as well as the chiral and anti-chiral generators  $\mathcal{W}_6$ ,  $\overline{\mathcal{W}}_6$  and their superpartners  $\mathcal{G}_{\mathcal{W}_6}$  and  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_6}$ . For the explicit free-field realization in terms of a single  $\beta\gamma bc$  ghost system one starts with a prescription for the 2D  $\mathcal{N} = 2$  SCA generators as well as for  $\mathcal{W}_6$ ,  $\mathcal{G}_{\mathcal{W}_6}$ . The ansatz for  $\overline{\mathcal{W}}_6$  contains 87 undetermined coefficients. Through the OPEs in the W-algebra presentation one can use it to also determine the free-field realization of  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_6}$ . We have calculated the fully-refined vacuum character (1) up to  $O(q^8)$  in the accompanying “nullstates.nb”. In the limit  $\xi \rightarrow 1$ ,  $z \rightarrow 1$  this reads:

$$\begin{aligned} \chi_{\mathcal{W}_6} = & 1 + q - 2q^{3/2} + 3q^2 - 4q^{5/2} + 8q^3 - 12q^{7/2} + 19q^4 \\ & - 26q^{9/2} + 38q^5 - 58q^{11/2} + 85q^6 - 116q^{13/2} \\ & + 165q^7 - 236q^{15/2} + 326q^8 + O(q^{17/2}). \end{aligned} \quad (9)$$

No known S-fold theory is associated with this VOA [9].

## VII. RESULTS: $\mathbf{G} = \mathbf{G}(3,1,2)$

This is our only rank 2 example, with central charge  $c = -48$ . Its W-algebra presentation involves the following strong generators:  $\mathcal{T}$ ,  $\mathcal{J}$ ,  $\mathcal{G}$  and  $\tilde{\mathcal{G}}$  from the 2D  $\mathcal{N} = 2$  SCA, the chiral and anti-chiral generators  $\mathcal{W}_3$ ,  $\mathcal{W}_6$ ,  $\overline{\mathcal{W}}_3$ ,  $\overline{\mathcal{W}}_6$ , plus their superpartners  $\mathcal{G}_{\mathcal{W}_3}$ ,  $\mathcal{G}_{\mathcal{W}_6}$  and  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_3}$ ,  $\tilde{\mathcal{G}}_{\overline{\mathcal{W}}_6}$ , as well as  $\mathcal{U}$ —which is self conjugate—and its superpartners  $\mathcal{G}_{\mathcal{U}}$ ,  $\tilde{\mathcal{G}}_{\mathcal{U}}$  and  $\mathcal{G}_{\tilde{\mathcal{U}}}$ . One also needs non-chiral  $\mathcal{O}$ ,  $\tilde{\mathcal{O}}$  and their superpartners  $\mathcal{G}_{\mathcal{O}}$ ,  $\tilde{\mathcal{G}}_{\mathcal{O}}$ ,  $\mathcal{G}_{\tilde{\mathcal{O}}}$ ,  $\tilde{\mathcal{G}}_{\tilde{\mathcal{O}}}$ ,  $\mathcal{G}_{\tilde{\mathcal{O}}}$ ,  $\mathcal{G}_{\tilde{\mathcal{O}}}$ .

The free-field realization requires two ghost systems,  $\beta_{\ell} \gamma_{\ell} b_{\ell} c_{\ell}$  with  $\ell = 1, 2$ . One starts with a prescription for the 2D  $\mathcal{N} = 2$  SCA generators as well as for  $\mathcal{W}_3$ ,  $\mathcal{W}_6$ ,  $\mathcal{G}_{\mathcal{W}_3}$



and  $\mathcal{G}_{\mathcal{W}_6}$ . The ansatz for  $\overline{\mathcal{W}}_3$  contains 84 undetermined coefficients. It turns out that through the OPEs in the W-algebra presentation, one can fix the coefficients of  $\overline{\mathcal{W}}_3$  and by doing so also determine the free-field realization of all remaining generators. We have calculated the fully refined vacuum character (1) up to  $O(q^{9/2})$  in the accompanying “null states.nb”.

This VOA is expected to encode the Schur sector of a rank-two 4D S-fold  $\mathcal{N} = 3$  SCFT, with Coulomb-branch operators of dimension  $\Delta = 3, 6$  and trace anomaly coefficient  $c_{4D} = 4$ . Through (3) the Macdonald index of this S-fold theory can be identified with the vacuum character. If for simplicity one considers the limit  $z \rightarrow 1$ :

$$\begin{aligned} \mathcal{I}_{\text{Macdonald}}^{G(3,1,2)} = & 1 + q\xi + q^{3/2}(-\sqrt{\xi} + \xi^{3/2}) + q^2(\xi + \xi^2) \\ & + q^{5/2}(-\sqrt{\xi} - \xi^{3/2} + 2\xi^{5/2}) + q^3(\xi - \xi^2 + 2\xi^3) \\ & + q^{7/2}(-\sqrt{\xi} - 2\xi^{3/2} + 2\xi^{5/2} + \xi^{7/2}) \\ & + q^4(2\xi - \xi^3 + 3\xi^4) \\ & + q^{9/2}(-\sqrt{\xi} - 3\xi^{3/2} + \xi^{5/2} + \xi^{7/2} + 2\xi^{9/2}) \\ & + O(q^5). \end{aligned} \quad (10)$$

### VIII. CONCLUSIONS

In this letter we have calculated vacuum characters of rank-one and rank-two VOAs labeled by non-Coxeter, crystallographic complex reflection groups. This involved a brute-force implementation of the algorithms presented in [20] and leads to the Macdonald index of certain 4D  $\mathcal{N} = 3$  S-fold SCFTs. Our results were given as an expansion in the fugacity that keeps track of the conformal weight, and were truncated to orders that require short computation times when using a desktop computer; they can be pushed to arbitrary higher orders by allocating appropriate

resources. As they stand, they can already be used as new data for  $\mathcal{N} = 3$  SCFTs. E.g., the  $\mathbf{G} = \mathbb{Z}_3$  result agrees in the Schur limit with [25].

Our code is customizable. We have clearly signposted where changes would need to be made to return vacuum characters of VOAs labeled by different complex reflection groups, for which the free-field realization is known. In particular, it would be very interesting to extend this approach to the  $\mathcal{N} = 3$  S-fold SCFT of rank two associated with  $G(4, 1, 2)$  and the rank-three example  $G(3, 3, 3)$ ; a proposal for the Schur index of the latter was also given in [25]. Unfortunately, finding the free-field realization for both these VOAs—already needed before identifying the null states—is a challenging task: the simplest antichiral strong generator ansätze involve 425 and 2265 undetermined coefficients respectively. It would perhaps be more promising to use the screening-operator approach, upon determining  $\mathcal{S}$ . Although our screening-operator code is currently more expensive to run, it could benefit from optimizations that parallelize the computations, hence making it significantly faster on multicore clusters. It will also be interesting to check these results by directly studying the BPS-states of  $\mathcal{N} = 3$  theories. One way to do so would be to study three-string junctions in S-fold backgrounds as in [29]. We hope to return to some of these questions in the near future.

### ACKNOWLEDGMENTS

We would like to thank B. Ergun, T. França, M. Martone, P. Longhi, C. Meneghelli, A. Pini, E. Pomoni, F. Schaposnik Massolo and J. Song for useful discussions and comments. We acknowledge financial support by the Royal Society Grants No. URF\180009, No. RGF\EA\181049, No. RGF\EA\180073, and the STFC Grant No. ST/P000754/1.

- 
- [1] P. C. Argyres and N. Seiberg, *J. High Energy Phys.* **12** (2007) 088.
  - [2] D. Gaiotto, *J. High Energy Phys.* **08** (2012) 034.
  - [3] R. Rattazzi, V. S. Rychkov, E. Tonni, and A. Vichi, *J. High Energy Phys.* **12** (2008) 031.
  - [4] C. Beem, M. Lemos, P. Liendo, L. Rastelli, and B. C. van Rees, *J. High Energy Phys.* **03** (2016) 183.
  - [5] J. Kinney, J. M. Maldacena, S. Minwalla, and S. Raju, *Commun. Math. Phys.* **275**, 209 (2007).
  - [6] S. Ferrara, M. Porrati, and A. Zaffaroni, *Lett. Math. Phys.* **47**, 255 (1999).
  - [7] O. Aharony and M. Evtikhiev, *J. High Energy Phys.* **04** (2016) 040.
  - [8] I. García-Etxebarria and D. Regalado, *J. High Energy Phys.* **03** (2016) 083.
  - [9] O. Aharony and Y. Tachikawa, *J. High Energy Phys.* **06** (2016) 044.
  - [10] Y. Imamura and S. Yokoyama, *J. Phys. A* **49**, 435401 (2016).
  - [11] R. Arai and Y. Imamura, *Prog. Theor. Exp. Phys.* **(2019)**, 083B04.
  - [12] P. C. Argyres, M. Lotito, Y. Lü, and M. Martone, *J. High Energy Phys.* **05** (2016) 088.
  - [13] P. C. Argyres, A. Bourget, and M. Martone, *SciPost Phys.* **9**, 083 (2020).
  - [14] A. Gadde, L. Rastelli, S. S. Razamat, and W. Yan, *Commun. Math. Phys.* **319**, 147 (2013).

- [15] T. Bourton, A. Pini, and E. Pomoni, *J. High Energy Phys.* **10** (2018) 131.
- [16] M. Evtikhiev, *J. High Energy Phys.* **06** (2020) 125.
- [17] C. Beem, M. Lemos, P. Liendo, W. Peelaers, L. Rastelli, and B. C. van Rees, *Commun. Math. Phys.* **336**, 1359 (2015).
- [18] T. Nishinaka and Y. Tachikawa, *J. High Energy Phys.* **09** (2016) 116.
- [19] M. Lemos, P. Liendo, C. Meneghelli, and V. Mitev, *J. High Energy Phys.* **04** (2017) 032.
- [20] F. Bonetti, C. Meneghelli, and L. Rastelli, *J. High Energy Phys.* **05** (2019) 155.
- [21] D. Adamović, *Transformation Groups* **21**, 299 (2016).
- [22] J. Song, *J. High Energy Phys.* **08** (2017) 044.
- [23] G. C. Shephard and J. A. Todd, *Can. J. Math.* **6**, 274 (1954).
- [24] Please see Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevD.103.L121701> for two accompanying *Mathematica* notebooks.
- [25] G. Zafirir, *J. High Energy Phys.* **01** (2021) 062.
- [26] K. Thielemans, Ph.D. thesis, Leuven University, 1994.
- [27] S. Krivonos and K. Thielemans, *Classical Quantum Gravity* **13**, 2899 (1996).
- [28] A. Fujitsu, *Comput. Phys. Commun.* **79**, 78 (1994).
- [29] P. Agarwal and A. Amariti, *J. High Energy Phys.* **09** (2016) 032.