

Classical Love number for quantum black holes

Ram Brustein^{*} and Yotam Sherf[†]

Department of Physics, Ben-Gurion University, Beer-Sheva 84105, Israel

 (Received 26 April 2021; accepted 16 December 2021; published 19 January 2022)

We present a method for comparing the classical and quantum calculations of the electric quadrupolar Love number k_2 and show that our previous derivation of the quantum Love number of a quantum black hole matches exactly the classical calculation of k_2 when quantum expectation values are replaced by the corresponding classical quantities, as dictated by the Bohr correspondence principle. The standard derivation of k_2 for classical relativistic stars relies on fixing boundary conditions on the surface of the star for the Einstein equations in the presence of an external perturbing field. An alternative method for calculating k_2 uses properties of the spectrum of the nonrelativistic fluid modes of the star. We adopt this alternative method and use it to derive an effective description of the interior modes in terms of a collection of driven harmonic oscillators characterized by different frequencies and amplitudes. We compare these two classical methods and find that most of the interior information can be integrated out, reducing the problem of calculating k_2 to fixing a single boundary condition for the perturbed Einstein equations on the surface of the deformed star. We then determine this single boundary condition in terms of the spectrum of the object and proceed to identify the relationship between classical quantities and quantum expectation values for the case of a quantum black hole and to verify the agreement between the results of the effective classical calculation and the quantum calculation.

DOI: [10.1103/PhysRevD.105.024044](https://doi.org/10.1103/PhysRevD.105.024044)

I. INTRODUCTION

The inspiral evolution of a binary system is accompanied by a mutual tidal interaction, leading to deformation on the spherical mass distribution. The tidal response of each of the companions is quantified in terms of the tidal Love numbers, which has a specific imprint on the emitted gravitational wave (GW) waveform [1–7]. In general relativity (GR), Love numbers of black holes (BHs) vanish universally as a consequence of the BH no-hair property [6,8,9] (for recent discussions, see [10,11] and for a discussion of higher dimensional BHs, see [12]).

We argued in [13] that future observations by the planned Laser Interferometer Space Antenna (LISA) [14] of GWs emitted during the inspiral phase of binary BH coalescence events will provide an opportunity for probing the quantum state of macroscopic astrophysical BHs via the imprint of this state on the emitted GW. Our argument was based on showing that the electric quadrupolar Love number, k_2 , is nonvanishing for a quantum black hole (QBH) and could be, under favourable circumstances, large enough for LISA to measure (similar arguments are also given in [15]). The vanishing of k_2 for GR BHs, and it being the largest of the dimensionless Love numbers, make k_2 a key diagnostic for deviations from classical GR.

Modifications to classical GR could perhaps be relevant in the strong-field regime, for which quantum effects might be significant. In this regime, additional quantum scales, such as the string scale, could appear [16,17]. Moreover, it was argued that modifications to the nature of BHs of quantum origin are needed to reconcile some fundamental problems such as the information loss paradox [18–20]. Among the suggested quantum ultracompact objects that could be relevant in this context are, for example, firewalls [19,21], fuzzballs [22], polymer BHs [23], and many others [13,24–30].

We are interested in calculating the Love numbers of large astrophysical BHs. As for any macroscopic object, the Bohr correspondence principle implies that some quantum state corresponds to the classical BH, no matter how large it is. We use the term “quantum black hole” to mean the quantum state that corresponds to a classical BH. The QBH is therefore an ultracompact object that possesses a horizon and, in addition, has a discrete spectrum of quantum mechanical energy levels. These energy levels can be viewed as coherent states that correspond to macroscopic, semiclassical excitations of the QBH. For example, in the polymer BH model of a QBH [23,28,31–33] (see also [24,34–36] for related models of QBHs), the interior matter can be effectively viewed as a fluid which can support pulsating modes in essentially the same way that a relativistic star does. These fluid modes would exist in addition to the standard spacetime modes of the exterior,

^{*}ramyb@bgu.ac.il

[†]sherfyo@post.bgu.ac.il

and so their spectrum should then be added onto that of the ringdown or quasinormal modes of a perturbed BH. Each of these fluid modes, when excited, represents a large-amplitude, high occupation number, coherent state of the interior matter rather than a single quantum excitation. In the ground state of the QBH, the exterior geometry is exactly the Schwarzschild geometry. But, when a QBH is in an excited state, it displays deviations from its GR description [33], and therefore it can be, in principle, distinguished from its classical counterpart.¹ This picture is consistent as long as the QBH has, to some degree, deviated from its equilibrium state, unlike a GR BH. The degree of deviation depends on the amount of energy that is injected into each of the specific modes.

One can probe the differences between a QBH and a classical BH when they are weakly out of equilibrium in response to the external field of a binary companion. The classical BH is bald, while the quantum BH has some quantum hair [33]. Here we focus on k_2 , a single number that is part of the quantum hair. The prevalent expectation is that quantum effects for large astrophysical BHs are controlled by the extremely small ratio of the Planck length squared to typical curvatures l_p^2/R_S^2 , and therefore are negligibly small. However, we argued that for QBHs, quantum effects are governed by the magnitude of the quantum hair which can be much larger. For example, in string theory the dimensionless magnitude of the hair scales with the string coupling squared, $g_s^2 \sim l_s^2/l_p^2$, where l_s is the string length. In general g_s^2 is expected to be small, but of the order of other typical gauge couplings, $g_s^2 \sim 0.1$.

A classical GR BH geometry does deform when subject to an external perturbing field, however, the deformation is not sourced by a varying matter distribution. Rather, the deformation is expressed in terms of a change in the Gaussian curvature at the Schwarzschild radius (or, equivalently, of the scalar curvature). Then, by embedding the BH in a fictitious two-dimensional sphere, one can interpret the deformation as the relative radial deviation of the position of the Schwarzschild radius. However, this geometric deformation should not be confused with a true physical effect. For an observer in the vicinity of the BH, the horizon is still fixed at the Schwarzschild radius. If this picture were not true, it would lead to a nonvanishing Love number. This argument is described clearly in [8], who demonstrated that geometrical deformations of classical BHs cannot be reflected in their asymptotic multipole moments and therefore their Love number vanishes identically. The conclusion is that a nonvanishing imprint on the object's asymptotic moment requires a physical matter deformation, or equivalently, a physical response of the state of the QBH to the external perturbation.

¹In [28,37] it is shown that due to vacuum fluctuations, GR BHs are tidally deformed. However, the deformation leads to a nonvanishing Love number that is Planckian suppressed [37].

In discussing quantum hair and its relevance for the calculation of k_2 , we would like to highlight the fact that for an external observer, the interior of the ultracompact object is affecting the result only in terms of the one boundary condition (BC) that is imposed on the external Einstein equations at the surface of the object. The second boundary condition can be expressed solely in terms of the perturbing classical field far away from the object. In effect, the interior is integrated out of the equations and any detailed information about the object's internal composition, energy density, or pressure can appear only through this one BC. The Love number is determined in terms of the ratio of the two BCs. Therefore, the only accessible information to an outside observer about the interior can be expressed in terms of the Love number(s). In this paper, we will develop the theoretical framework for applying these ideas and demonstrate explicitly how these ideas are manifested when solving the perturbed Einstein equations.

The rest of the paper is organized as follows. First, we review the calculation of the quantum Love number in [13]. Then, we review the standard calculation of k_2 and focus on the required BCs for this calculation. In Sec. III B, we review an alternative method of calculating the Love number of an ultracompact object that is driven by an external periodic force using the spectrum of its interior nonrelativistic (NR) fluid modes. We adapt this method so that the results match those for an asymptotic observer far away from the QBH. Then, we can compare the two methods and from the consistency of the two methods and find the boundary conditions that need to be imposed on the perturbed Einstein equations. Finally, we compare our previous quantum calculation of k_2 to the classical calculation of k_2 for a QBH and show that the two calculations agree for macroscopic QBHs. This agreement, as dictated by the Bohr correspondence principle, allows us to establish the correspondence between the classical and quantum methods of calculating k_2 and provides us with an explicit dictionary for comparing them and showing their agreement. We end the paper with a summary and conclusion.

II. QUANTUM LOVE NUMBER

We first briefly recall our previous calculation of the quantum Love number [13], where additional details can be found.

In GR, the interior of a BH is empty, nearly a vacuum, except for a possibly singular core. Reconciling GR with quantum mechanics led to the realization that a substantial revision to the classical GR is required. The firewall argument marked the beginning of a new era in the theory of quantum BHs [21,38], and forerunners of the argument are seen in [39–42].

We will discuss a class of solutions to the issues raised by the firewall argument. The main idea is that strong quantum effects lead to horizon scale deviations rather than restriction to the Planck length scale and are characterized by an

intrinsic excitations spectrum. In these cases, the emergence of new physics introduces a new scale, and the ratio of the new scale to the Planck scale can be viewed as a coupling constant, as in string theory. The complete picture is accompanied by a self-consistent interior description that requires a significant departure from the semiclassical gravity, as well as some exotic matter which is outside the realm of the standard model [43]: fuzzballs, [44,45] and the polymer model [46].

Since the interior is inaccessible to an external observer, and because the gravity in the interior is assumed to be strongly coupled, one cannot use the semiclassical geometric description in terms of a curved spacetime. However, the only relevant aspect of the QBH interior is that excitations are macroscopic so applying Bohr correspondence principle is justified, implying that the excited spectrum of a QBH can be described by a set of coherent states. Then, from the Bohr correspondence principle, we conclude that these states correspond to semiclassical states that can be effectively described as an oscillating classical system. These assumptions were previously discussed also in [34,47] and later in [48,49]. In these works, the macroscopic excitations and the energy spectrum of a quantized BH were described by a spectrum of a harmonic oscillator.

Then, to implement the idea, we view the exotic matter in the interior of the QBH effectively as a fluid that supports pulsating modes as for a relativistic star. These fluid modes would exist in addition to the standard spacetime modes of the exterior. The perturbations are divided into two sectors, the fluid modes and spacetime modes. Due to their low speed of sound and the compactness of the QBH, fluid modes are decoupled from the spacetime perturbations as in the Cowling approximation [50–52]. The fluid mode description is discussed extensively in Sec. III B.

As a motivation for the calculation of the Love number of the QBH, let us recall the analogous calculation of the polarizability of an atom. Consider an atom in its ground state, $|\Psi_0\rangle$, which is labeled by the quantum numbers $|n, l, m\rangle = |1, 0, 0\rangle$, and assume that the expectation value of the quantum dipole operator \hat{D}_i vanishes in the ground state. Recall that, classically, the dipole moment is given by $\vec{D} = \int \rho(\vec{x}') \vec{x}' dV' = 0$, where ρ is the charge density of the atom. The atom is placed in a region of an approximately uniform electric field \mathcal{E}_i that is induced by a weak external potential U_{ext} , $\mathcal{E}_i = \frac{\partial U_{\text{ext}}}{\partial x^i}$. The interaction of the atom with the external electric field \hat{V}_{int} is expressed in terms of \hat{D}_i , $\hat{V}_{\text{int}} = -\mathcal{E}_i \hat{D}_i$. The induced dipole moment of the perturbed atom in second-order time-independent perturbation theory is given by the standard textbook expression,

$$\langle \Psi_0 | \hat{D}_j | \Psi \rangle = -\mathcal{E}_i \sum_{n \neq 1, l, m} \frac{\langle 1, 0, 0 | \hat{D}_i | n, l, m \rangle \langle n, l, m | \hat{D}_j | 1, 0, 0 \rangle}{\Delta E_{1,n}}, \quad (2.1)$$

where $\Delta E_{1,n} = E_1 - E_n$. In this case, symmetry implies that $l = 1$, $m = -1, 0, 1$, and $i = j$. The atom's linear response to the external electric field is then $\langle \Psi_0 | \hat{D}_i | \Psi_0 \rangle = \alpha \mathcal{E}_i$, where α is the electric polarizability,

$$\alpha = \sum_{n \neq 1, m = -1, 0, 1} \frac{|\langle 1, 0, 0 | \hat{D}_i | n, 1, m \rangle|^2}{\Delta E_{1,n}}. \quad (2.2)$$

Following similar considerations, we derived in [13] an expression for the gravitational polarizability, the Love numbers. The idea is to replace the electric field and the dipole moment by the tidal fields and the mass moment.

For the quantum calculation, we considered, again, the inspiral phase of a binary system, as we did for the classical calculation. One of the companions is an object of mass M_{ext} on a circular orbit of radius b and the other is a QBH of mass M_{BH} and radius $R_S = 2M_{\text{BH}}$. In the early stages of the inspiral, the QBH responds to the external slowly varying tidal field that is generated by its companion. For $b \gg R_S$ one can expand the Newtonian potential, $U_{\text{ext}} = -M_{\text{ext}}/|\vec{b} - \vec{x}|$, of the external body in the vicinity of the QBH in its local inertial frame, $U(t, x)_{\text{ext}} = U_{\text{ext}}(0) + \frac{1}{2} \frac{\partial^2 U_{\text{ext}}}{\partial x^i \partial x^j} |_0 x^i x^j + \dots$.

As in the case of the electric polarizability, the interaction of the QBH with the external field is expressed in terms of the mass moment expectation value, $\hat{Q}^{(l)}$. These operators are the quantum counterparts of the classical symmetric trace-free mass multipoles [53]. We further assume that the expectation value of the BH mass moment vanishes in the BH ground state, as dictated by the spherical symmetry and the classical no-hair properties. Denoting the ground state of the QBH by $|\Psi_0\rangle$, we have $\langle \Psi_0 | \hat{Q}^{(l)} | \Psi_0 \rangle = 0$. Owing to the slowly varying weak external potential, time-independent perturbation theory is a good approximation.

In [13] we evaluated the correction to the ground state energy due to the induced quadrupole, \hat{Q}_{ij} , of a nonrotating quantum BH whose Schwarzschild radius is R_S . Recall that in the classical case, $Q_{ij} = \int \rho(t, x') (x'_i x'_j - \frac{1}{3} r'^2 \delta_{ij}) dV'$, where ρ is the energy density. In an analogy to the electric polarizability calculation, the interaction energy is given by $\hat{V}_{\text{int}} = -\frac{1}{2} \mathcal{E}_{ij} \hat{Q}_{ij}$, where $\mathcal{E}_{ij} = \frac{\partial^2 U_{\text{ext}}}{\partial x^i \partial x^j}$.

The leading order corrections to the QBH ground state quadrupole in second-order time-independent perturbation theory is given by

$$\langle \Psi_0 | \hat{Q}_{kl} | \Psi_0 \rangle = \mathcal{E}_{ij} \sum_{n > 1, l, m} \frac{\langle \Psi_0 | \hat{Q}_{ij} | n, l, m \rangle \langle n, l, m | \hat{Q}_{kl} | \Psi_0 \rangle}{|\Delta E_{1,n}|}, \quad (2.3)$$

where $|\Delta E_{1,n}| = E_n - E_1$. Here the radial number of the ground state Ψ_0 is denoted by $n = 1$, so the energy of the ground state is $E_1 = M_{\text{BH}}$. This choice is made for consistency with the standard treatments of second-order

perturbation theory. The electric quadrupolar Love number is defined as the proportionality coefficient between the induced electric quadrupole moment to the external tidal field,

$$\langle \Psi_0 | \hat{Q}_{ij} | \Psi_0 \rangle = -\lambda_2 \mathcal{E}_{ij}. \quad (2.4)$$

Here λ_2 is the dimensional quadrupolar Love number, which in its dimensionless form is commonly defined as $k_2 = \frac{3}{2} R^{-5} \lambda_2$. From Eq. (2.3) it follows that

$$k_2 = -\frac{3}{4R^5} \sum_{n, -2 < m < 2} \frac{|\langle \Psi_0 | \hat{Q}_{ij} | n, 2, m \rangle|^2}{|\Delta E_{1,n}|}. \quad (2.5)$$

We would like to point out that the additional QBH excitations decay at a parametrically slow rate in comparison to the Schwarzschild time due to the large redshift factor in the vicinity of the deformed QBH. This was explained in detail in [33] and we recall the essence of relevant arguments below. The excited modes therefore have a parametrically small width compared to the standard GR BH excitations, which justifies neglecting the width of the excitations in Eq. (2.5).

For an exterior observer, the wavelength of the excited modes near the source must be $\lambda_S \sim R_S$, which then asymptotically redshifts to some larger value, $\lambda_A \sim R_S z$ with $z \gg 1$. This observer then assigns a transmission cross section for such long wavelength modes through a proportionally smaller surface of area A which is determined by the ratio $A/\lambda^2 \sim 1/z^2$. The coupling or efficiency of emission then scales as $1/z^2$. The damping time for that mode τ is related to the inverse of the efficiency of emission and therefore scales as $z^2 \gg 1$.

III. CLASSICAL LOVE NUMBER

A. Method 1: Explicit internal solution for the metric

We review here the standard calculation of the Love number. This will also be used to set up notations and conventions.

The response of a body to a weak external tidal field is reflected in its induced mass (electric) and current (magnetic) moments. We will focus here on the quadrupolar electric Love number k_2 . At large distances, in the star's local asymptotic rest frame, the temporal component of the metric is given by

$$\begin{aligned} g_{tt} &= -1 + \frac{2M}{r} - \mathcal{E}_{ij} x^i x^j + 3 \frac{1}{r^5} Q_{ij} x^i x^j \\ &= -1 + \frac{2M}{r} - \mathcal{E}_{ij} x^i x^j - 2k_2 (R/r)^5 \mathcal{E}_{ij} x^i x^j, \end{aligned} \quad (3.1)$$

where M , R are the mass and the radius of the star, respectively, and k_2 is the dimensionless tidal Love number that measures the linear response to the applied field.

The first term in the deviation of g_{tt} from the Schwarzschild metric in Eq. (3.1) describes the applied tidal field, while the term proportional to k_2 describes the induced trace-free quadrupole moment $Q_{ij} = \int d^3x \rho(x) (x_i x_j - \frac{1}{3} \delta_{ij} r^2)$,

$$Q_{ij} = -\frac{2}{3} k_2 R^5 \mathcal{E}_{ij}. \quad (3.2)$$

From a Newtonian perspective, at large distances $g_{tt} = -(1 + 2U_N)$. The expansion of the Newtonian potential U_N , to second order in the body's local inertial frame reads $U_N = -\frac{M}{r} - \frac{3}{2r^5} Q_{ij} x^i x^j + \frac{1}{2} \mathcal{E}_{ij} x^i x^j$, with \mathcal{E}_{ij} being the quadrupole moment of the external potential $\mathcal{E}_{ij} = \frac{\partial^2 U_{\text{ext}}}{\partial x_i \partial x_j}$. Here we discuss the axisymmetric external potential, so the tidal field is given by $\mathcal{E}_{ij} x^i x^j = \mathcal{E} r^2 Y_{20}$ and consequently also the induced moment has the same angular dependence $Q_{ij} = Q Y_{20}$. It follows that

$$U_N = -\frac{M}{r} - \frac{3}{2r^3} Q Y_{20} + \frac{1}{2} \mathcal{E} r^2 Y_{20}, \quad (3.3)$$

and

$$\begin{aligned} g_{tt} &= -1 + \frac{2M}{r} + 3Q \frac{1}{r^3} Y_{20} - \mathcal{E} r^2 Y_{20}, \\ &= -1 + \frac{2M}{r} - 2k_2 R^5 \mathcal{E} \frac{1}{r^3} Y_{20} - \mathcal{E} r^2 Y_{20}, \end{aligned} \quad (3.4)$$

with

$$k_2 R^5 = -\frac{3Q}{2\mathcal{E}}. \quad (3.5)$$

We now turn to the review of the calculation of the Love number k_2 [5–7]. One considers a perturbation $h_{\mu\nu}$ about the Schwarzschild background $g_{\mu\nu}^{(0)}$, $g_{\mu\nu}^{(0)} = \text{diag}(-e^{\nu(r)}, e^{\lambda(r)}, r^2, r^2 \sin^2 \theta)$, with $e^{\nu(r)} = e^{-\lambda(r)} = 1 - 2M/r$. Then, $h_{\mu\nu}$ is decomposed into even-parity and odd-parity parts as in the Regge-Wheeler gauge. Focusing on the $l = 2$, $m = 0$ term, to linear order, in the limit that the external field is static, $\mathcal{E} = \text{const.}$, the even-parity perturbations can be expressed as [54]

$$h_{\mu\nu} = \text{diag}(e^{\nu(r)} H_0(r), e^{\lambda(r)} H_2(r), r^2 K(r), r^2 \sin^2 \theta K(r)) Y_{20}. \quad (3.6)$$

Then, solving the perturbed Einstein equations outside the body, one finds that $H_0 = H_2 \equiv H(r)$ and arrives at the following perturbation equations:

$$H'' + \frac{2x}{x^2 - 1} H' - \frac{6x^2 - 2}{(x^2 - 1)^2} H = 0, \quad (3.7)$$

$$K' - H' - \frac{2}{(x^2 - 1)}H = 0, \quad (3.8)$$

$$K - \frac{1}{2}H' - \frac{1}{4}\left(\frac{x+1}{x-1} + \frac{3x+5}{x+1}\right)H = 0, \quad (3.9)$$

where $x = r/M - 1$ and the prime denotes a derivative with respect to x .

The exterior solution of the perturbation equations is given by

$$H_{\text{ext}}(x) = c_1 \left[\frac{x(5 - 3x^2)}{x^2 - 1} + \frac{3}{2}(x^2 - 1) \ln\left(\frac{x+1}{x-1}\right) \right] + 3c_2(x^2 - 1), \quad (3.10)$$

$$K_{\text{ext}}(x) = -c_1 \frac{4 + 3x(x+3)}{x+1} + \frac{3}{2}c_1(x^2 + 2x - 1) \ln\left(\frac{x+1}{x-1}\right) + 3c_2(x^2 + 2x - 1), \quad (3.11)$$

and the perturbed background metric by

$$g_{tt} = -\left(\frac{x-1}{x+1}\right)(1 + H_{\text{ext}}Y_{20}). \quad (3.12)$$

We will be interested in cases for which $x - 1 = r/M - 2 \ll 1$. For later use, we expand H_{ext} and K_{ext} in this limit,

$$H_{\text{ext}}(x) = \frac{c_1}{x-1} - 3c_1 \ln\left(\frac{x-1}{2}\right) + 6c_2(x-1) + \mathcal{O}(x-1), \quad (3.13)$$

$$K_{\text{ext}}(x) = -8c_1 - 3c_1 \ln\left(\frac{x-1}{2}\right) + 6c_2 + \mathcal{O}(x-1). \quad (3.14)$$

It is convenient to relate the coefficients of the expanded metric in Eq. (3.4), \mathcal{E} and k_2 , to the coefficients of the expanded metric in Eq. (3.10), c_1 and c_2 :

$$c_1 = 40M^2 k_2 \mathcal{E} \quad (3.15)$$

and

$$c_2 = \frac{1}{3}M^2 \mathcal{E}, \quad (3.16)$$

so

$$k_2 = \frac{1}{120} \frac{c_1}{c_2}. \quad (3.17)$$

Before proceeding to discuss explicit solutions for k_2 , let us emphasize some features about its dependence on the

deformed body. First, we note that the functional form of the exterior solution, Eq. (3.10), does not depend on the interior, since it corresponds to a vacuum solution of the Einstein equations. Because c_2 is determined entirely by the external field, the only implicit dependence of H_{ext} on the interior is encoded in the ratio c_1/c_2 or, equivalently, the magnitude of k_2 . It follows that one needs to specify one more BC on H_{ext} to determine completely the form of the external metric perturbations.

The standard approach for calculating k_2 for objects whose interiors are known and well-defined is reviewed in [5,6]. One starts the calculation by specifying the interior configuration of the perturbed body in terms of its stress-energy-momentum tensor and an equation of state. Then, the perturbed Einstein equations in the interior, which have the same form as the exterior equations (3.7)–(3.9), are solved and their solution H_{int} is found. Next, demanding regularity of H_{int} at the star's center $r = 0$ and continuity of H and K at the star's surface $r = R$, one finds the coefficients c_1 and c_2 , or equivalently k_2 .

It is clear that this method of calculating k_2 , in which one needs the full detailed solution of the interior, contains a lot of redundant information if one is just interested in finding one number— k_2 . We just need one ratio of two numbers for that, so any other method for specifying this ratio would work just as well as the standard one. We will describe such an alternative method in the next section.

B. Method 2: Spectrum of nonrelativistic fluid modes

In this section we discuss a binary system, in which the object of interest—the “primary”—is driven by a weak periodic force which is exerted by the companion. We rely on the ideas presented in [55–58] to establish an effective description for the interior fluid modes of ultracompact objects as a collection of driven harmonic oscillators characterized by their frequencies. In this effective description, the interior modes of the object are described from the point of view of an asymptotic observer as NR fluid modes and are analyzed in a similar manner to the analysis of classical Newtonian NR fluid modes. As discussed in Sec. II, due to their low speed of sound and the compactness of the QBH, fluid modes are decoupled from the spacetime perturbations as in the Cowling approximation [50–52]. This splits the interior perturbation into two sectors, fluid modes, which we will discuss, and spacetime modes, which we will neglect.

We consider the oscillating modes of the object that are labeled by the radial and spherical indices n , l , m . Again, we focus on the tidal axisymmetric perturbations, so $l = 2$, $m = 0$. The analysis proceeds by writing and solving the equations for the Lagrangian displacement vector of the fluid ξ^i which is proportional to Y_{20} . The total displacement vector is given by the complete sum over the contributions from each of the radial modes,

$$\xi^i = \sum_n a_n \xi_n^i. \quad (3.18)$$

The modes ξ_n have units of length, so the coefficients a_n are dimensionless. Different radial modes are orthogonal and their precise normalization will not be important for us.

In the absence of the driving force that arises from the binary companion, the fluid modes satisfy the following Harmonic-oscillator equation:

$$\ddot{a}_n + \omega_n^2 a_n = 0, \quad (3.19)$$

with ω_n being the frequency of the n th mode. When an external tidal potential is present, the equation of motion (EOM) for the internal fluid modes becomes that of a driven harmonic oscillator (see, for example, [55]),

$$(-\omega^2 + \omega_n^2)a_n = \frac{\mathcal{E}Q_n}{MR^2}. \quad (3.20)$$

Here ω is the frequency of the external tidal field, to be discussed shortly. The quadrupole associated with the n th mode Q_n , is defined by the overlap integral

$$Q_n = - \int d^3r \delta\rho_n r^2. \quad (3.21)$$

The quadrupolar energy density perturbation, $\delta\rho_n$, is associated with the n th fluid mode and ΔE_n is the corresponding total mass quadrupole moment. $Q_n = -\gamma \Delta E_n R^2$, with γ being a dimensionless number of order unity.

We are interested in the $m = 0$ modes, so the driving is essentially at zero frequency, with $\omega = m\Omega$, $\Omega = \sqrt{M/b^3}$ being the orbital frequency. So, Eq. (3.20) simplifies

$$\omega_n^2 a_n = \frac{\mathcal{E}Q_n}{MR^2}. \quad (3.22)$$

In general, the driving frequency can be neglected also for the case $m \neq 0$ as it is small compared to the oscillator natural frequencies $\Omega^2 \ll \omega_n^2$, $\omega_n^2 \sim 1/R^2$, while $\Omega^2 = R/b^3$ and $R^3/b^3 \ll 1$. The solution of Eq. (3.22) is the following:

$$a_n = \mathcal{E} \frac{Q_n}{M\omega_n^2 R^2}. \quad (3.23)$$

Next, we identify the Love number using the asymptotic moments for a static observer at infinity which can be read off from the external metric perturbation equation (3.5), with $Q = \sum_n a_n Q_n$ [57], in contrast to [58] where the Love number is identified at the star surface. Substituting \mathcal{E} from Eq. (3.23) into Eq. (3.5), one finds

$$k_{2n} R^5 = -\frac{3}{2} \frac{Q_n}{\mathcal{E}} = -\frac{3}{2} \frac{1}{a_n} \frac{Q_n^2}{M\omega_n^2 R^2}. \quad (3.24)$$

Now, from the asymptotic moment decomposition $k_2 = \sum_n a_n k_{2n}$, we obtain

$$k_2 = -\sum_n \frac{3}{2R^5} \frac{Q_n^2}{M\omega_n^2 R^2}. \quad (3.25)$$

We can re-express the results in terms of the intrinsic energy spectrum of the driven system. We may define the intrinsic energy difference $\Delta E_n^{\text{int}} = E_n^{\text{int}} - M$ as

$$\Delta E_n^{\text{int}} = \frac{1}{2} M \omega_n^2 R^2. \quad (3.26)$$

The intrinsic energy difference ΔE_n^{int} depends only on the intrinsic properties of the object and does not depend on the external driving field. It should not be confused with the energy that is pumped into the mode n by the external field,

$$\Delta E_n^{\text{induced}} = \frac{1}{2} MR^2 \omega_n^2 a_n^2. \quad (3.27)$$

Substituting a_n from Eq. (3.23) into Eq. (3.27), we find that $\Delta E_n^{\text{induced}}$ is related to the total work done by the external tidal force,

$$\Delta E^{\text{induced}} = \sum_n \Delta E_n^{\text{induced}} = \frac{1}{2} \sum_n \mathcal{E} a_n Q_n = \frac{1}{2} \mathcal{E} Q. \quad (3.28)$$

Substituting Eq. (3.26) into Eq. (3.25), we obtain our final expression for the Love number,

$$k_2 = -\frac{3}{4R^5} \sum_n \frac{Q_n^2}{\Delta E_n^{\text{int}}}. \quad (3.29)$$

In the case that the sum is dominated by the lowest energy level $n = 1$, then

$$k_2 \simeq -\frac{3}{4R^5} \frac{Q_1^2}{\Delta E_1^{\text{int}}}. \quad (3.30)$$

We can further parametrize Q_n on dimensional grounds, as was previously done, by

$$Q_n = \gamma_n \Delta E_n^{\text{int}} R^2, \quad (3.31)$$

where γ_n is a dimensionless number that depends on the detailed functional form of the energy density profile of the object. Then,

$$k_2 = -\frac{3}{4R} \sum_n \gamma_n^2 \Delta E_n^{\text{int}}. \quad (3.32)$$

On general grounds, we expect γ_n to rapidly decrease as n increases, as we expect the higher- n modes to possess more

nodes and thus induce a smaller quadrupole moment.² If the decrease in γ_n offsets the increase in E_n as expected,

$$k_2 \simeq -\frac{3}{4R}\gamma_1^2\Delta E_1^{\text{int}}. \quad (3.33)$$

This relation can be further simplified since $\Delta E_1^{\text{int}} \sim M\omega_1^2 R^2$ and typically $\omega_1 \sim 1/R$, so $k_2 \sim \gamma_1^2$.

In summary, the calculation of the Love number amounts to a classical linear response calculation of a collection of driven harmonic oscillators by an external force in the limit that the intrinsic frequency of the oscillator is much higher than the frequency of the driving.

In the quantum case, the calculation is similar, except that the relevant quantity is the quadrupole moment of the interior modes (evaluated by a static observer at infinity) without an explicit reference to the Lagrangian displacement vector or to the Newtonian potential at the surface of the star. We emphasize that these should be viewed as means to an end: fixing one additional BC for the exterior perturbation equations. This will be our bridge to the quantum Love calculations and results in the semiclassical approximation. We need to know the energy of the lowest lying level and the quadrupole associated with this level.

C. Relating the two methods for quantum black holes

The two methods of calculating the Love number calculate the same quantity, the response of an ultra compact object to an external perturbation, and so, in principle, must agree. The only issue is to what extent the various approximations that are made along the way affect the resulting values of k_2 that are obtained by the two methods.

However, we now argue that a quantitative relation between the fluid and geometric methods is not necessary for our purposes, as we are interested in a general relationship that can then be applied to QBHs. We emphasize that for the QBH it is also impossible to find directly such a quantitative relation without a more detailed description of the interior. We bypass the fact that the interior of the QBH is not prescribed by observing that for QBHs all that is required to determine the Love number is a single BC at the surface of the QBH.

The physical assumptions that we make to identify this single BC are the following:

- (1) Both GR BHs and QBHs possess a horizon.
- (2) In the absence of the external perturbation, QBHs cannot be distinguished from GR BHs.
- (3) For QBHs, geometric deformations induced by external perturbations are reflected in their asymptotic moments, in contrast to GR BHs. This implies

that, due to changes in their state, the QBHs possess hair that leads to a nonvanishing k_2 .

- (4) For QBHs, as for their classical counterparts, the external metric perturbation must vanish on the deformed horizon.
- (5) The interior excited modes of the QBH can be described effectively as a collection of driven harmonic oscillators, which gives rise to the Love number as in Eq. (3.25). An exact solution of Q_n as in Eq (3.21) is model dependent and can be parametrized by a dimensionless number of order unity γ as in Eq. (3.31).

To apply our ideas in a concrete context, let us consider the temporal component of the metric perturbation δg_{tt} near the boundary of the QBH. For simplicity, we suppress the angular dependence of the external metric perturbation. Denoting by x_B the value of x at the deformed surface, we get

$$-\delta g_{tt}(x_B) \sim c_1 + c_2(x_B - 1)^2. \quad (3.34)$$

We assume that $x_B - 1 \ll 1$, so denoting by ΔR the difference $R - 2M$, $\Delta R/2M \ll 1$, it follows that

$$x_B - 1 = \frac{\Delta R}{M}. \quad (3.35)$$

We need to choose a single additional BC that the classical metric external to the QBHs needs to satisfy, that is, we need to choose one more BC that H_{ext} needs to satisfy at the deformed boundary of the QBH. Applying our five assumptions above, we find that the following conditions need to be imposed:

- (1) Assumption 1 implies that for both classical and quantum BHs, for $\mathcal{E} = 0$ (or equivalently, $c_2 = 0$), $x_B^{\text{BH}} = x_B^{\text{QBH}} = 1$ and $g_{tt}(x_B) = 0$, in agreement with assumption 4.
- (2) Assumption 2 implies that both GR BHs and QBHs possess a horizon and cannot be distinguished in the absence of perturbations. Furthermore, according to assumption 3, for $\mathcal{E} \neq 0$ a physical deformation of the QBH is induced, such that the surface of the QBH is shifted to $x_B = 1 + \delta x_B$. The specific value of δx_B depends on the spectrum of the QBH.
- (3) Assumption 1 implies that for classical BHs, $c_1 = 0$ and $x_B^{\text{BH}} = 1$ so $\delta g_{tt}^{\text{BH}} \sim c_2(x_B - 1)^2 = 0$, and from Eq. (3.12), $g_{tt}^{\text{BH}}(x_B^{\text{BH}}) = 0$. Assumption 4 implies that the QBH case is similar, $g_{tt}(1 + \delta x_B) = 0$ where $x_B = 1 + \delta x_B$ is the position of the deformed horizon.

These three conditions are summarized in Table I.

We wish to emphasize the origin for the difference between GR BHs to QBHs, where according to [6,7] it is the boundary condition on the BH horizon that kills the response terms [the terms proportional to c_1 in Eq. (3.13)]

²This property is typical for driven harmonic systems, as suggested by the equation (3.20).

TABLE I. Comparison of the response of classical and quantum BHs. We evaluate the Euclidean deformation $\Delta R/R$ from the Gaussian curvature Eq. (3.11) at the surface, $K(x_B) \sim \Delta R/R$, [6].

	x_B	δg_{tt}	$(\Delta R/R)$	k_2
QBH	$1 + \delta x_B$	$g_{tt}(1 + \delta x_B) = 0$	$c_2 - c_1$	$\neq 0$
BH	1	$\delta g_{tt}(x_B) = 0$	c_2	0

and leads to the vanishing of the Love number. An observer in the vicinity of a classical BH sees no deviation in the horizon position; the horizon is frozen at $R = 2M$, and the external perturbation is *singular* on it. On the other hand, according to assumptions 3 and 4, the horizon of a QBH does deform (see discussion in Sec. III B). Thus, the boundary conditions on the deformed surface are *regular* and lead to a nonvanishing Love number.

Rather than imposing $g_{tt}(1 + \delta x_B) = 0$, we impose an equivalent condition, $\delta g_{tt}(1) = -\frac{1}{2}c_1 Y_{20}$, as we now explain. The fact that some points with $x_B = 1$ are formally within the original horizon is not relevant for our discussion, as we are only interested in the perturbed metric far away from the horizon of the QBH. The classical metric is, of course, only valid outside the QBH horizon.

Let us recall the expansion of H_{ext} in Eq. (3.13):

$$\begin{aligned} H_{\text{ext}}(x_B) &= \frac{c_1}{(x_B - 1)} + \mathcal{O}(x_B - 1) \\ &= 120k_2 \frac{c_2}{x_B - 1} + \mathcal{O}(x_B - 1), \end{aligned} \quad (3.36)$$

where we used Eq. (3.17) to relate c_1 to c_2 . It follows that

$$\delta g_{tt}(x_B) = -\frac{1}{2}c_1 + \mathcal{O}(x_B - 1) = 5k_2 R^2 \mathcal{E} + \mathcal{O}(x_B - 1), \quad (3.37)$$

where we used the relationship between c_2 and \mathcal{E} in Eq. (3.16).

The choice of BC for fixing the exterior solution is clear. Since \mathcal{E} (or equivalently, c_2) is already fixed by the BC at infinity, we need to fix $\delta g_{tt}(x_B)$ to fix the value of c_1 . Here we use our assumptions 3 and 5 to choose the value of c_1 such that the value of k_2 agrees with the value obtained using the fluid calculation in Eq. (3.29),

$$c_1 = -2\delta g_{tt}(2M) = \frac{15}{2} \frac{\mathcal{E}}{(2M)^3} \sum_n \frac{Q_n^2}{\Delta E_n^{\text{int}}}. \quad (3.38)$$

Equation (3.38) completes our comparison between the two classical methods of calculating k_2 as it determines the relationship between the calculation of k_2 in terms of the spectrum of the fluid modes and the choice of BC on the perturbed relativistic Einstein equations for the case of a QBH.

If the lowest level dominates the sum, as expected, we can use the estimate in Eq. (3.30),

$$c_1 = -2\delta g_{tt}(R_B) \sim -\frac{15}{2} \frac{\mathcal{E}}{(2M)^3} \frac{Q_1^2}{\Delta E_1^{\text{int}}}. \quad (3.39)$$

Furthermore, from Eqs. (3.12) and (3.36) we find

$$-g_{tt}(x_B) = \left(\frac{x_B - 1}{x_B + 1} \right) \left(1 + \frac{c_1}{x_B - 1} \right) + \mathcal{O}(x_B - 1)^2, \quad (3.40)$$

which from the BC on the deformed surface $g_{tt}(1 + \delta x_B) = 0$, we obtain $c_1 = 1 - x_B$, as noted in Table I.

IV. COMPARISON WITH THE QUANTUM LOVE NUMBER

We wish to compare the classical and quantum calculations of the Love number of the QBH.

First, let us emphasize the striking similarity between the expression for the classical Love number, Eq. (3.29), and the expression in Eq. (2.5) for the quantum Love number. If one identifies the expectation values with the corresponding classical observables $\langle \Psi_0 | \hat{Q}_{ij} | n, 2, 0 \rangle = Q_n$ and the internal excited energy spectrum $|\Delta E_{1,n}| = \Delta E_n^{\text{int}}$, Eqs. (3.29) and (2.5) become identical. As discussed in [13], the quadrupole matrix element is given by the following integral:

$$|\langle \Psi_0 | \hat{Q} | n, 2, 0 \rangle| = \int d^3r \delta \tilde{\rho}_{n,2}(r) r^2 Y_{20} \Psi_{n,2}, \quad (4.1)$$

where $\delta \tilde{\rho}_{n,2}(r)$ and $\Psi_{n,2}$ are the effective energy density and the mode function of the n 'th excited level, respectively. In comparison to the overlap integral, Eq. (3.21), we find that the two expressions coincide when the density profile is decomposed by $\delta \rho = \sum_n \delta \tilde{\rho}_{n,2} \Psi_{n,2}$ [57].

As anticipated in [13], this correspondence is an explicit manifestation of the Bohr correspondence principle, which states that for macroscopic states associated with large quantum occupation numbers, expectation values correspond to classical quantities. The states that we consider are indeed states with large occupation numbers since for a given quantum state whose energy scales as $M\omega_n^2 R^2$, the occupation number N scales as $N\hbar\omega_n \sim M\omega_n^2 R^2$, so $N \sim (\omega_n R)MR/\hbar \sim (\omega_n R)S_{\text{BH}} \gg 1$. Here S_{BH} is the Bekenstein-Hawking entropy of the QBH.

Equation (3.38) completes the comparison of the quantum calculation to the classical calculation by specifying the required additional BC for the perturbation equations that determine the external metric.

V. SUMMARY AND CONCLUSIONS

We showed how to calculate k_2 in an explicit way from partial knowledge of the internal spectrum of an ultra-compact object, be it the spectrum of the nonrelativistic fluid modes of a classical ultracompact star or the spectrum of excited states of a QBH. The single additional BC that encodes the relevant information about the interior of the QBH is determined in terms of its spectrum. In both cases, k_2 depends most strongly on the first excited level, or on the lowest lying fluid mode, and is proportional to the relative excitation energy of this level $k_2 \sim \Delta E/M$. The proportionality coefficient depends on additional information: the order unity ratio of the quadrupole moment of the excited level to its excitation energy ΔE .

Furthermore, since finding k_2 is equivalent to finding the ratio of two numbers, one does not need the detailed solution in the interior. The interior information is accessible to an external observer only through deformations of the surface of the QBH and can be integrated out, such that the only relevant quantity is a single boundary condition.

That k_2 does not vanish for a QBH is a violation of the no-hair property and reflects the main difference between BHs to QBHs. For classical BHs, geometric deformations of the BH do not affect the asymptotic moments while for a QBH they do, and require a physical matter deformation, or equivalently, a physical response of the state of the QBH to the external perturbation and can therefore be detected by an external observer. Our results further highlight the importance of k_2 as a key diagnostic observable for probing the quantum nature of BHs.

The agreement between the classical and quantum calculations of k_2 indicates that they are equivalent ways of deriving the same answer and strengthens the validity of each of the calculations. The consistency of the classical and quantum calculations provides us with an explicit dictionary, translating quantum observables into classical GR quantities. Moreover, it demonstrates that the relation between quantum observables and the analogous GR quantities is in complete agreement with the Bohr correspondence principle.

For completeness, we wish to emphasize the differences between the Love number of QBHs to that of the semi-classical objects (like gravastars and wormholes) reviewed in [59]. First, these objects are not quantum in nature, they do not possess an event horizon and their unperturbed surface lies a finite distance away from the would-be horizon rather than at $R = 2M$. Second, the method for calculating their Love number is rather different; one must assume that a fictitious infinitely thin rigid shell surrounds these objects. The shell is made of a fictional matter violating both the weak and dominant energy conditions. The presence of a thin shell is necessary to guarantee the continuity of the interior to the exterior solutions. As a result, in order to compensate for the metric discontinuity, the boundary condition imposed on the object's surface is the so-called Israel junctions condition, which is different than the regularity condition of the external metric (see Table I). Therefore, the origin of the Love number is in the discontinuity of the metric solution, which is a purely geometric property. On the other hand, the QBH Love number originates in the deformation of the interior matter distribution or equivalently in the coupling of its ground state to higher states by the external tidal perturbation.

The current work can be extended to the spinning case, for which, as pointed out in [13], spin effects are subleading and induce small corrections to k_2 . Additionally, the spectrum of internal modes is expected to determine also the spectrum of ringdown modes and the nature of the merger. Thus, the internal structure of the QBH could potentially induce significant modifications to the merger and the ringdown phases of binary BH coalescence events; in particular, if resonance excitations of the QBH occur during the inspiral phase. We hope to report on these interesting possibilities and their imprint on the emitted GW waveform in a future publication [60].

ACKNOWLEDGMENTS

The research of R.B. and Y.S. was supported by the Israel Science Foundation Grant No. 1294/16. The research of Y.S. was supported by the Negev scholarship.

-
- [1] E. E. Flanagan and T. Hinderer, Constraining neutron star tidal Love numbers with gravitational wave detectors, *Phys. Rev. D* **77**, 021502 (2008).
 - [2] K. Yagi and N. Yunes, Love can be tough to measure, *Phys. Rev. D* **89**, 021303 (2014).
 - [3] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.* **119**, 161101 (2017).
 - [4] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW190425: Observation of a compact binary coalescence with total mass $\sim 3.4 M_\odot$, *Astrophys. J. Lett.* **892**, L3 (2020).
 - [5] T. Hinderer, Tidal Love numbers of neutron stars, *Astrophys. J.* **677**, 1216 (2008).
 - [6] T. Damour and A. Nagar, Relativistic tidal properties of neutron stars, *Phys. Rev. D* **80**, 084035 (2009).
 - [7] T. Binnington and E. Poisson, Relativistic theory of tidal Love numbers, *Phys. Rev. D* **80**, 084018 (2009).

- [8] N. Grlebeck, No-Hair Theorem for Black Holes in Astrophysical Environments, *Phys. Rev. Lett.* **114**, 151102 (2015).
- [9] K. Yagi and N. Yunes, Approximate universal relations for neutron stars and quark stars, *Phys. Rep.* **681**, 1 (2017).
- [10] P. Charalambous, S. Dubovsky, and M. M. Ivanov, On the vanishing of Love numbers for Kerr black holes, *J. High Energy Phys.* **05** (2021) 038.
- [11] P. Charalambous, S. Dubovsky, and M. M. Ivanov, Hidden Symmetry of Vanishing Love, *Phys. Rev. Lett.* **127**, 101101 (2021).
- [12] B. Kol and M. Smolkin, Black hole stereotyping: Induced gravito-static polarization, *J. High Energy Phys.* **02** (2012) 010.
- [13] R. Brustein and Y. Sherf, Preceding paper, Quantum Love, *Phys. Rev. D* **105**, 024043 (2022).
- [14] P. Amaro-Seoan, H. Audley, S. Babak, J. Baker, E. Barausse, P. Bender, E. Berti, P. Binetruy, M. Born, D. Bortoluzzi, J. Camp, C. Caprini, V. Cardoso, M. Colpi, J. Conklin, N. Cornish, C. Cutler *et al.*, Laser interferometer space antenna, [arXiv:1702.00786](https://arxiv.org/abs/1702.00786).
- [15] A. Maselli, P. Pani, V. Cardoso, T. Abdelsalhin, L. Gualtieri, and V. Ferrari, Probing Planckian Corrections at the Horizon Scale with LISA Binaries, *Phys. Rev. Lett.* **120**, 081101 (2018).
- [16] D. G. Boulware and S. Deser, String Generated Gravity Models, *Phys. Rev. Lett.* **55**, 2656 (1985).
- [17] B. Zwiebach, Curvature squared terms and string theories, *Phys. Lett.* **156B**, 315 (1985).
- [18] S. L. Braunstein, S. Pirandola, and K. Życzkowski, Better Late than Never: Information Retrieval from Black Holes, *Phys. Rev. Lett.* **110**, 101301 (2013).
- [19] S. D. Mathur, The information paradox: A pedagogical introduction, *Classical Quant. Grav.* **26**, 224001 (2009).
- [20] R. Brustein and A. J. M. Medved, Restoring predictability in semiclassical gravitational collapse, *J. High Energy Phys.* **09** (2013) 015.
- [21] A. Almheiri, D. Marolf, J. Polchinski, and J. Sully, Black holes: Complementarity or firewalls?, *J. High Energy Phys.* **02** (2013) 062.
- [22] K. Skenderis and M. Taylor, The fuzzball proposal for black holes, *Phys. Rep.* **467**, 117 (2008).
- [23] R. Brustein and A. J. M. Medved, Black holes as collapsed polymers, *Fortschr. Phys.* **65**, 1600114 (2017).
- [24] J. D. Bekenstein and V. F. Mukhanov, Spectroscopy of the quantum black hole, *Phys. Lett. B* **360**, 7 (1995).
- [25] S. B. Giddings, Nonviolent unitarization: Basic postulates to soft quantum structure of black holes, *J. High Energy Phys.* **12** (2017) 047.
- [26] V. Cardoso, V. F. Foit, and M. Kleban, Gravitational wave echoes from black hole area quantization, *J. Cosmol. Astropart. Phys.* **08** (2019) 006.
- [27] V. Cardoso and P. Pani, Testing the nature of dark compact objects: A status report, *Living Rev. Relativity* **22**, 4 (2019).
- [28] R. Brustein and Y. Sherf, Emission channels from perturbed quantum black holes, *Phys. Rev. D* **100**, 124005 (2019).
- [29] Q. Wang, N. Oshita, and N. Afshordi, Echoes from quantum black holes, *Phys. Rev. D* **101**, 024031 (2020).
- [30] I. Agullo, V. Cardoso, A. del Rio, M. Maggiore, and J. Pullin, Gravitational-Wave Signatures of Quantum Gravity, *Phys. Rev. Lett.* **127**, 101101 (2021).
- [31] R. Brustein and A. J. M. Medved, Emergent horizon, Hawking radiation and chaos in the collapsed polymer model of a black hole, *Fortschr. Phys.* **65**, 1600116 (2017).
- [32] R. Brustein, A. J. M. Medved, and K. Yagi, When black holes collide: Probing the interior composition by the spectrum of ringdown modes and emitted gravitational waves, *Phys. Rev. D* **96**, 064033 (2017).
- [33] R. Brustein and A. J. M. Medved, Quantum hair of black holes out of equilibrium, *Phys. Rev. D* **97**, 044035 (2018).
- [34] J. D. Bekenstein, Quantum black holes as atoms, [arXiv:gr-qc/9710076](https://arxiv.org/abs/gr-qc/9710076).
- [35] S. D. Mathur, The fuzzball proposal for black holes: An elementary review, *Fortschr. Phys.* **53**, 793 (2005).
- [36] B. Guo, S. Hampton, and S. D. Mathur, Can we observe fuzzballs or firewalls?, *J. High Energy Phys.* **07** (2018) 162.
- [37] J. W. Kim and M. Shim, Quantum corrections to tidal Love number for Schwarzschild black holes, *Phys. Rev. D* **104**, 046022 (2021).
- [38] D. Marolf and J. Polchinski, Gauge/Gravity Duality and the Black Hole Interior, *Phys. Rev. Lett.* **111**, 171301 (2013).
- [39] N. Itzhaki, Is the black hole complementarity principle really necessary?, [arXiv:hep-th/9607028](https://arxiv.org/abs/hep-th/9607028).
- [40] S. D. Mathur, What exactly is the information paradox?, *Lect. Notes Phys.* **769**, 3 (2009).
- [41] S. L. Braunstein, S. Pirandola, and K. Życzkowski, Entangled Black Holes as Ciphers of Hidden Information, *Phys. Rev. Lett.* **110**, 101301 (2013).
- [42] S. D. Mathur, Resolving the black hole causality paradox, *Gen. Relativ. Gravit.* **51**, 24 (2019).
- [43] R. Brustein and A. J. M. Medved, Non-singular black holes interiors need physics beyond the standard model, *Fortschr. Phys.* **67**, 1900058 (2019).
- [44] S. D. Mathur, The quantum structure of black holes, *Classical Quant. Grav.* **23**, R115 (2006).
- [45] K. Skenderis and M. Taylor, The fuzzball proposal for black holes, *Phys. Rep.* **467**, 117 (2008).
- [46] R. Brustein and A. J. M. Medved, Black holes as collapsed polymers, *Fortschr. Phys.* **65**, 0114 (2017).
- [47] J. D. Bekenstein, The quantum mass spectrum of the Kerr black hole, *Lett. Nuovo Cimento* **11**, 467 (1974).
- [48] S. Hod, Bohr's Correspondence Principle and the Area Spectrum of Quantum Black Holes, *Phys. Rev. Lett.* **81**, 4293 (1998).
- [49] M. Maggiore, The Physical Interpretation of the Spectrum of Black Hole Quasinormal Modes, *Phys. Rev. Lett.* **100**, 141301 (2008).
- [50] K. D. Kokkotas and B. G. Schmidt, Quasinormal modes of stars and black holes, *Living Rev. Relativity* **2**, 2 (1999).
- [51] G. Allen, N. Andersson, K. D. Kokkotas, and B. F. Schutz, Gravitational waves from pulsating stars: Evolving the perturbation equations for a relativistic star, *Phys. Rev. D* **58**, 124012 (1998).
- [52] N. Andersson, K. D. Kokkotas, and B. F. Schutz, Space-time modes of relativistic stars, *Mon. Not. R. Astron. Soc.* **280**, 1230 (1996).
- [53] K. S. Thorne, Multipole expansions of gravitational radiation, *Rev. Mod. Phys.* **52**, 299 (1980).
- [54] T. Regge and J. A. Wheeler, Stability of a Schwarzschild singularity, *Phys. Rev.* **108**, 1063 (1957).

- [55] D. Lai, Resonant oscillations and tidal heating in coalescing binary neutron stars, *Mon. Not. R. Astron. Soc.* **270**, 611 (1994).
- [56] W.C.G. Ho and D. Lai, Resonant tidal excitations of rotating neutron stars in coalescing binaries, *Mon. Not. R. Astron. Soc.* **308**, 153 (1999).
- [57] S. Chakrabarti, T. Delsate, and J. Steinhoff, Effective action and linear response of compact objects in Newtonian gravity, *Phys. Rev. D* **88**, 084038 (2013).
- [58] N. Andersson and P. Pnigouras, Exploring the effective tidal deformability of neutron stars, *Phys. Rev. D* **101**, 083001 (2020).
- [59] V. Cardoso, E. Franzin, A. Maselli, P. Pani, and G. Raposo, Testing strong-field gravity with tidal Love numbers, *Phys. Rev. D* **95**, 084014 (2017).
- [60] R. Brustein and Y. Sherf, Resonance excitations of quantum black holes (to be published).