


New nonsingular cosmological solution of nonlocal gravity

Ivan Kolář^{1,*}, Francisco José Maldonado Torralba^{2,†} and Anupam Mazumdar^{1,‡}

¹*Van Swinderen Institute, University of Groningen, 9747 AG, Groningen, Netherlands*

²*Laboratory of Theoretical Physics, Institute of Physics, University of Tartu,
W. Ostwaldi 1, 50411 Tartu, Estonia*

 (Received 16 September 2021; accepted 31 January 2022; published 22 February 2022)

We present a new bouncing cosmological solution of the nonlocal theory known as infinite derivative gravity, which goes beyond the recursive ansatz, $\square R = r_1 R + r_2$. The nonlocal field equations are evaluated using the spectral decomposition with respect to the eigenfunctions of the wave operator. The energy-momentum tensor computed for this geometry turns out to be much more sensitive to the choice of the nonlocal form factor, since it depends on the value of the function on a continuous infinite interval. We show that this stronger dependence on the form factor allows us to source the geometry by the perfect fluid with the non-negative energy density satisfying the strong energy condition. We show that this bouncing behavior is not possible in the local theories of gravity such as, in general, relativity or $R + R^2$ gravity sourced by a fluid which meets the non-negative energy and strong energy conditions.

DOI: [10.1103/PhysRevD.105.044045](https://doi.org/10.1103/PhysRevD.105.044045)

I. INTRODUCTION

There is no doubt that the current cosmological model, Λ cold dark matter, gives so far the most accurate description of the history of the Universe [1]. Nevertheless, it suffers from some important shortcomings, such as the tensions between early and late time measures [2], the nature of dark energy [3,4], and the initial singularity problem [5,6]. The latter is the most worrisome from a conceptual perspective, since the existence of singularities signals the limited range of validity of the theory.

By the time the Friedmann–Lemaître–Robertson–Walker (FLRW) cosmological solution was introduced, “ $t = 0$ singularity” was thought to be a consequence of spacetime symmetries. It was not until the development of the singularity theorems in the 1960s by Penrose and Hawking and co-workers [7–10] that the sufficient conditions for the appearance of singularities were elucidated. In particular, in a cosmological context, one can recall the Hawking singularity theorem, which states [8] the following:

Theorem.—Let M be a 4-dimensional manifold equipped with the metric g satisfying the following conditions:

- (1) Global hyperbolicity.
- (2) Temporal convergence condition, i.e., $\mathbf{v} \cdot \mathbf{Ric} \cdot \mathbf{v} \geq 0$ for every timelike vector \mathbf{v} , where \mathbf{Ric} is the Ricci tensor, and \cdot denotes a contraction between adjacent tensor indices.

- (3) There exists a Cauchy hypersurface Σ with expansion of the future-directed geodesic congruence greater or equal than a constant $A > 0$.

Then, no past-directed timelike curve from Σ can have a length greater than $3/|A|$. In particular, all past-directed timelike geodesics are incomplete.

It is clear that this theorem predicts the initial singularity for quite general requirements, and hence, this behavior is not due to the specific symmetry of the considered solution. When applied to a FLRW metric that describes the expansion of the Universe, the first and third conditions hold. Therefore, in order to predict singularities in such spacetimes, we just need to worry about the second condition in the theorem, the *temporal convergence condition*. Such a condition is completely theory independent, but it can be related to the energy-momentum tensor of a specific theory via the field equations, obtaining what is known as the *energy conditions*.

In the *general relativity* (GR), the temporal convergence condition written in terms of the energy-momentum tensor T can be expressed as

$$\mathbf{v} \cdot \left(T - \frac{1}{2} Tg \right) \cdot \mathbf{v} \geq 0, \quad (1.1)$$

for every timelike vector \mathbf{v} . Such a condition is usually referred to as the *strong energy condition* (SEC). In a particular case of the perfect fluid, which is a source of a FLRW spacetime, SEC is equivalent to

$$\rho + p \geq 0, \quad \rho + 3p \geq 0, \quad (1.2)$$

*i.kolar@rug.nl

†fmaldo01@ucm.es

‡anupam.mazumdar@rug.nl

where ρ and p are the *energy density* and the *pressure* of the perfect fluid, respectively. SEC has an important physical interpretation; it guarantees the attractive character of gravity in GR [10,11].

Because of issues linked to the initial singularity problem, the nonsingular cosmological solutions have become a widely studied topic in gravitation (for a comprehensive review, see [5]). The nonsingular cosmological solutions which contract and then expand are called the *bouncing cosmologies*. The first explicit solutions were obtained in the late 1970s by Novello and Salim [12] and Melnikov and Orlov [13]. Nevertheless, at that time, they did not attract too much attention of the community, since they require the violation of SEC. It was not until two decades later, with the measurement of the accelerated expansion of the Universe, that these kinds of metrics started to be regarded as possible physical solutions of the problems present in the standard cosmological models.

Resorting to Hawking's theorem, it is clear that in order to obtain nonsingular bouncing solutions in GR, SEC must be necessarily violated [14,15], implying that gravity is not attractive at the bounce. This fact has served as motivation to explore bouncing solutions in modified theories of gravity, where the unattractive character can be a consequence of the modification, not of the energy-momentum tensor. For a review of bouncing universes in various theories of gravity, we refer the reader to [16].

In this paper, we are interested in nonlocal theory of gravity known as the *infinite derivative gravity* (IDG) [17–20], which contains operators with an infinite number of derivatives in the action. The most general action (quadratic in curvature) with no ghosts or extra degrees of freedom has been constructed around Minkowski [20], de Sitter, and anti-de Sitter spacetimes [21]. The nonlocal gravitational interaction has been argued to improve ultraviolet aspects gravity at the quantum level [22–26]. At the classical level, it has been proved that the linearized IDG can yield nonsingular static solutions for point sources (including electromagnetic charges), p-branes (and cosmic strings), spinning ring distributions, sources describing Newman-Unti-Tamburino charges, null sources, and accelerated particles [20,27–33]. It was also shown that the linearized IDG prevents mini-black-hole production for small masses [28,34,35]. Also, hints of this nonsingular behavior for astrophysical objects have been studied in [36,37]. Exact gravitational waves generated by null sources in full IDG were found in [38–41].

Several cosmological studies were performed within IDG, in particular, the possibility of describing inflation [42–45] and the existence of nonsingular bouncing solutions [19,46–53]. All the previous examples made an extensive use of the recursive ansatz on the Ricci scalar R , i.e., $\square R = r_1 R + r_2$, which reduces the field equations to local second order nonlinear differential equations. As a matter of fact, in [42], it was proved that such a recursive ansatz for flat FRLW already incorporates all solutions of

the IDG field equations with a traceless energy-momentum tensor. In this paper, we go beyond the recursive ansatz and find a bouncing solution whose energy-momentum tensor is not traceless and shows much stronger dependence on the nonlocal form factor.

The paper is organized as follows: In Sec. II, we derive the field equations of IDG and explain the treatment of the nonlocal equations with the spectral analysis of the wave operator. In Sec. III, we present a bouncing solution for which the eigenvalue problem can be solved and derive an explicit form of the energy-momentum tensor. In Sec. IV, we find sufficient conditions for the form factor that ensure that the energy density of the perfect fluid is non-negative, and SEC is met. We also show that this bouncing behavior is not possible neither in GR nor in “ $R + R^2$ gravity” sourced by a fluid meeting such conditions. Moreover, we find that the nonlocality only affects the bounce region, while recovering GR at late times. Finally, we provide a brief summary of the results in Sec. V.

II. EVALUATION OF NONLOCAL FIELD EQUATIONS

In this paper, we consider a nonlocal theory described by the action of IDG,

$$\mathbf{S}[\mathbf{g}] = \frac{1}{2} \int_M \mathbf{g}^{\frac{1}{2}} (\kappa^{-1} (R - 2\Lambda) + RF(\square)R) + \mathbf{S}_m, \quad (2.1)$$

where R is the Ricci scalar, $\mathbf{g}^{\frac{1}{2}} = \sqrt{-g} dx^4$ the volume element, $\kappa = 8\pi G_N$ the Einstein gravitational constant, \square the wave operator, Λ the cosmological constant, and \mathbf{S}_m the matter sector of the action. The nonlocal operator $F(\square)$ is often referred to as the *form factor*. It is given by an analytic nonpolynomial function,¹

$$F(\square) = \sum_{n=0}^{\infty} f_n \square^n. \quad (2.2)$$

Performing the variation with respect to the metric \mathbf{g} , we find the field equations of IDG [19,54],

$$\kappa^{-1} \left(\mathbf{Ric} - \frac{1}{2} R \mathbf{g} + \Lambda \mathbf{g} \right) + \Upsilon + \frac{1}{2} \mathbf{g} (\Omega + \Theta) - \Omega = \mathbf{T}, \quad (2.3)$$

where the rank-2 tensors Υ , Ω , and the scalar Θ are given by

¹Here, the f_n are dimensionfull constant coefficients.

$$\begin{aligned} \Upsilon &\equiv \left[2\mathbf{Ric} - 2\nabla\nabla - \frac{1}{2}\mathbf{g}R + 2\mathbf{g}\square \right] F(\square)R, & \square\psi_\lambda &= \lambda\psi_\lambda, & (2.6) \\ \Omega &\equiv \sum_{n=1}^{\infty} f_n \sum_{k=0}^{n-1} \nabla\square^k R \nabla\square^{n-k-1} R \equiv R\vec{\nabla}G(\vec{\square}, \vec{\square})\vec{\nabla}R, \\ \Theta &\equiv \sum_{n=1}^{\infty} f_n \sum_{k=0}^{n-1} \square^k R \square^{n-k} R \equiv R \frac{\vec{\square} + \overleftarrow{\square}}{2} G(\vec{\square}, \overleftarrow{\square})R. & (2.4) \end{aligned}$$

The arrows over the covariant derivatives denote left/right actions of the derivative. The operator $G(\vec{\square}, \overleftarrow{\square})$ is a nonlocal bilinear operator which is defined as

$$G(\vec{\square}, \overleftarrow{\square}) \equiv \frac{F(\vec{\square}) - F(\overleftarrow{\square})}{\vec{\square} - \overleftarrow{\square}}. \quad (2.5)$$

The presented field equations are very complicated due to their nonlocal and nonlinear structure. There exist three main approaches to solving these equations in the literature:

- (i) **Linearized regime [20]:** Performing the first order perturbations with respect to a given background, one can approximate the field equations by *linear* nonlocal equations.
- (ii) **TN/TIII ansatz [40]:** In certain class of geometries, such as the almost universal spacetimes TN/TIII, the nonlinear terms may not contribute even without making any approximations. The field equation may reduce to the *linear* nonlocal differential equations.
- (iii) **Recursive ansatz [19]:** Assuming the recursive ansatz for the curvature, such as $\square R = r_1 R + r_2$, the field equations can be recast to the form of *local* (second order) nonlinear differential equations.

Only the second and third methods produce exact solutions of the full theory; however, such geometries are very special. In the TN/TIII ansatz, the nonlinear terms in the curvature do not contribute. In the recursive ansatz, the dependence on the form factor is very weak in the sense that only the values $F(r_1)$ and $F'(r_1)$ are actually important for the energy-momentum tensor. In this paper, we study a geometry that does not fall in any of these categories. Its energy-momentum tensor has a strong dependence on the nonlocal form factor, and the nonlinear terms do not automatically vanish.

When solving the linear nonlocal differential equations, for example, those arising in the linearized regime or from the TN/TIII ansatz, one often employs the spectral representation of the nonlocal operators using the eigenvalues of the wave operator \square . The linearity then allows one to invert the nonlocal operators and write the solutions using the associated integral transforms (if the spectrum is continuous). Of course, such an inversion is not possible if the equations are nonlinear; however, one can still use the method of spectral decomposition to evaluate the field equation. Namely, we can try to find a certain geometry for which the eigenvalue problem,

possesses simple (analytic) solutions $\psi_\lambda = \psi_\lambda(\mathbf{x})$ for discrete or continuous spectrum λ . If the eigenfunctions ψ_λ form a complete orthonormal set, $\langle \psi_\lambda, \psi_{\lambda'} \rangle = \delta_{\lambda\lambda'}$, then we can decompose a function f in terms of the eigenfunctions ψ_λ ,

$$f(\mathbf{x}) = \sum_{\lambda} f_{\lambda} \psi_{\lambda}(\mathbf{x}), \quad f_{\lambda} = \langle \psi_{\lambda}, f \rangle \quad (2.7)$$

where \sum_{λ} denotes summation \sum_{λ} if the spectrum is discrete and integration \int_{λ} if the spectrum is continuous. In the latter case, these expressions define direct and inverse integral transforms. (The simplest example is the Fourier transform in flat spacetime in Lorentzian coordinates.) Equipped with this, we can represent the action of the nonlocal operator $F(\square)$ by

$$F(\square)f = \sum_{\lambda} f_{\lambda} F(\square)\psi_{\lambda} = \sum_{\lambda} f_{\lambda} F(\lambda)\psi_{\lambda} \quad (2.8)$$

In the next section, we present a spacetime, where this process of evaluation of the nonlocal field equations is tractable, yet still nontrivial in the sense that the energy-momentum tensor depends on all values of F .

III. NEW NONLOCAL BOUNCING GEOMETRIES

The search for a geometry for which the field equations can be evaluated using the method described above is quite nontrivial for several reasons: (i) There are very few physically interesting geometries that actually admit exact closed-form solution of the eigenvalue problem. (ii) In many of these cases, the eigenfunctions are either too complicated or do not form a complete set. (iii) Even if this is all satisfied, the Ricci scalar may still fail to be decomposable. To deal with these issues, we focus on a flat FLRW ansatz,

$$\mathbf{g} = -\mathbf{d}t\mathbf{d}t + a^2\mathbf{q}, \quad (3.1)$$

where $a = a(t)$ denotes the *scale factor*, and \mathbf{q} is the metric of the 3-dimensional flat space. When acting on scalar functions, the wave operator \square reads

$$\square = -\partial_t^2 - 3H\partial_t, \quad (3.2)$$

where $H = \dot{a}/a$ is the Hubble parameter. The Ricci tensor \mathbf{Ric} and the Ricci scalar R of the metric (3.1) are given by the expressions

$$\begin{aligned} \mathbf{Ric} &= -\frac{3\ddot{a}}{a}\mathbf{d}t\mathbf{d}t + (a\ddot{a} + 2\dot{a}^2)\mathbf{q} \\ &= -3(\dot{H} + H^2)\mathbf{d}t\mathbf{d}t + (\dot{H} + 3H^2)a^2\mathbf{q}, \\ R &= 6\left(\frac{\ddot{a}}{a} + \frac{\dot{a}^2}{a^2}\right) = 6(\dot{H} + 2H^2). \end{aligned} \quad (3.3)$$

The individual terms in the field equations (2.3) can be expressed as

$$\begin{aligned} \kappa^{-1} \left(\mathbf{Ric} - \frac{1}{2} R \mathbf{g} + \Lambda \mathbf{g} \right) &= \kappa^{-1} (3H^2 - \Lambda) \mathbf{d}t \mathbf{d}t - \kappa^{-1} (2\dot{H} + 3H^2 - \Lambda) a^2 \mathbf{q}, \\ \Upsilon &= [(6H\partial_t - 3\dot{H})F(\square)R] \mathbf{d}t \mathbf{d}t + [-(2\partial_t^2 + 4H\partial_t + \dot{H})F(\square)R] a^2 \mathbf{q}, \\ \frac{1}{2} \mathbf{g}(\Omega + \Theta) - \Omega &= -\frac{1}{2} (\Theta + \Omega_{tt}) \mathbf{d}t \mathbf{d}t + \frac{1}{2} (\Theta - \Omega_{tt}) a^2 \mathbf{q}. \end{aligned} \quad (3.4)$$

Thus, we can find the formula for the energy-momentum tensor of the flat FLRW spacetime,

$$\begin{aligned} T &= \left[\kappa^{-1} (3H^2 - \Lambda) + (6H\partial_t - 3\dot{H})F(\square)R - \frac{1}{2} (\Theta + \Omega_{tt}) \right] \mathbf{d}t \mathbf{d}t \\ &+ \left[-\kappa^{-1} (2\dot{H} + 3H^2 - \Lambda) - (2\partial_t^2 + 4H\partial_t + \dot{H})F(\square)R + \frac{1}{2} (\Theta - \Omega_{tt}) \right] a^2 \mathbf{q}. \end{aligned} \quad (3.5)$$

As mentioned above, the spectral decomposition plays a central role in the evaluation of (3.5). The differential equation for the eigenvalue problem,

$$\square \psi_\omega = \lambda_\omega \psi_\omega, \quad (3.6)$$

is remarkably simple in the FLRW geometry (3.1), especially if we focus on spatially homogeneous eigenfunctions $\psi_\omega = \psi_\omega(t)$,

$$\ddot{\psi}_\omega + 3H\dot{\psi}_\omega + \lambda_\omega \psi_\omega = 0, \quad (3.7)$$

where the dot stands for the derivative with respect to t . Here, λ_ω denotes the eigenvalues, labeled by a parameter ω .

In order to find explicit solutions of this equation, one has to specify the geometry even further. Let us focus on the FLRW spacetimes with the following scale factor:

$$a = a_0 \cosh^\gamma(\alpha t), \quad (3.8)$$

with $\alpha > 0$ and $\gamma > 0$. This scale factor describes bouncing cosmologies with the bounce at $t = 0$, which approach de Sitter spacetime with the Ricci scalar $R = 12\alpha^2\gamma^2$ for $t \rightarrow \pm\infty$. Note that by explicitly choosing nonsingular geometries we avoid problems with introducing nonlocal operators that can act on functions living on manifolds with boundaries. The Hubble parameter H for this scale factor is given by

$$H = \gamma\alpha \tanh(\alpha t). \quad (3.9)$$

The special value $\gamma = 1$, has been studied in the literature [19]. An important property of such a spacetime is that the Ricci scalar satisfy recursive ansatz. In other words, the Ricci scalar shifted by a constant is itself an eigenfunction of \square [i.e., satisfying (3.7)] because

$$\square R = 2\alpha^2 R - 24\alpha^4 \iff \square \overset{\circ}{R} = 2\alpha^2 \overset{\circ}{R}, \quad \overset{\circ}{R} \equiv R - 12\alpha^2, \quad (3.10)$$

meaning that $\psi_0 = \overset{\circ}{R}$ and $\lambda_0 = 2\alpha^2$. Also, the particular case $\gamma = 1/2$ has been considered, which gives a constant curvature spacetime, which clearly satisfies the ansatz [48]. As mentioned above, the recursive ansatz has proved useful in generating solutions of nonlocal gravity because it effectively reduces the field equations to the local equations with only a weak dependence on the nonlocal form factor.

We study the case $\gamma = 2/3$, which goes beyond the recursive ansatz, since

$$\square R = -\frac{3}{2} \left(R - \frac{16}{3} \alpha^2 \right)^2. \quad (3.11)$$

The choice $\gamma = 2/3$ is rather unique, since the eigenfunctions take a surprisingly simple form,

$$\psi_\omega = \frac{1}{\sqrt{2\pi}} \operatorname{sech}(\alpha t) e^{-i\omega t}, \quad (3.12)$$

with the corresponding eigenvalues $\lambda_\omega = \omega^2 + \alpha^2$; therefore,

$$\square \psi_\omega = (\omega^2 + \alpha^2) \psi_\omega. \quad (3.13)$$

The eigenfunctions (3.12) are very convenient because they depend on ω only through the multiplicative factor $e^{-i\omega t}$. Thus, they give rise to the integral transform \mathbf{G} that is closely related to the Fourier transform \mathbf{F} ,

$$\begin{aligned} \mathbf{F}[h](\omega) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dt h(t) e^{i\omega t}, \\ \mathbf{F}^{-1}[h](t) &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} d\omega h(\omega) e^{-i\omega t}. \end{aligned} \quad (3.14)$$

In particular, for the inverse transform of \mathbf{G} , we find

$$\begin{aligned}
 f(t) &= \mathbf{G}^{-1}[f_\omega](t) \\
 &= \int_{\mathbb{R}} d\omega f_\omega \psi_\omega(t) \\
 &= \operatorname{sech}(\alpha t) \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} d\omega f_\omega e^{-i\omega t} \\
 &= \operatorname{sech}(\alpha t) \mathbf{F}^{-1}[f_\omega](t). \tag{3.15}
 \end{aligned}$$

Thanks to this relation, we can find also the direct integral transform,

$$\begin{aligned}
 f_\omega &= \mathbf{G}[f(t)](\omega) \\
 &= \mathbf{F}[\cosh(\alpha t)f(t)](\omega) \\
 &= \frac{1}{\sqrt{2\pi}} \int_{\mathbb{R}} dt \cosh(\alpha t) f(t) e^{i\omega t} \\
 &= \int_{\mathbb{R}} dt \cosh^2(\alpha t) \psi_\omega^*(t) f(t) \\
 &= \langle \psi_\omega, f \rangle, \tag{3.16}
 \end{aligned}$$

and identify the inner product in which ψ_ω are orthonormal. Hence, we are ensured that the integral transform \mathbf{G} that is given by eigenfunctions ψ_ω is well defined and has a corresponding inverse \mathbf{G}^{-1} .

To evaluate the nonlocal terms (2.4) in the field equations, we have to express the Ricci scalar R with the help of

the integral transform \mathbf{G} . The Ricci scalar of the flat FLRW with the scale factor (3.8) and $\gamma = 2/3$ is

$$R = \frac{4}{3} \alpha^2 (4 - \operatorname{sech}^2 \alpha t). \tag{3.17}$$

Apparently, a problem arises here, since the relevant integral diverges due to the constant part of the expression. Instead, we can decompose the Ricci scalar shifted by this constant,

$$\hat{R} \equiv R - \frac{16}{3} \alpha^2 = \mathbf{G}^{-1}[\hat{R}_\omega](t) = \int_{\mathbb{R}} d\omega \hat{R}_\omega \psi_\omega, \tag{3.18}$$

where

$$\begin{aligned}
 \hat{R}_\omega &= \mathbf{G}[\hat{R}(t)](\omega) \\
 &= \mathbf{F}[\cosh(\alpha t)\hat{R}(t)](\omega) \\
 &= -\frac{2\sqrt{2\pi}}{3} \alpha \operatorname{sech}\left(\frac{\pi\omega}{2\alpha}\right). \tag{3.19}
 \end{aligned}$$

As we see, this decomposition is sufficient to rewrite the field equations to the integral form. Indeed, if we apply the integral representation of the Ricci scalar (3.18), we can write

$$\begin{aligned}
 F(\square)R &= \frac{16}{3} \alpha^2 f_0 + \int_{\mathbb{R}} d\omega F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega, \\
 \Omega_{tt} &= \iint_{\mathbb{R}^2} d\omega d\omega' G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'}, \\
 \Theta &= -\frac{16}{3} \alpha^2 f_0 R + \left(\frac{16}{3} \alpha^2\right)^2 f_0 + \frac{16}{3} \alpha^2 \int_{\mathbb{R}} d\omega F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega \\
 &\quad + \iint_{\mathbb{R}^2} d\omega d\omega' \frac{\omega^2 + \omega'^2 + 2\alpha^2}{2} G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'}. \tag{3.20}
 \end{aligned}$$

Consequently, the individual terms in (3.5) can be expressed as follows:

$$\begin{aligned}
 \varkappa^{-1}(3H^2 - \Lambda) &= \varkappa^{-1} \left(\frac{4}{3} \alpha^2 \tanh^2(\alpha t) - \Lambda \right), \\
 -\varkappa^{-1}(2\dot{H} + 3H^2 - \Lambda) &= \varkappa^{-1} \left(-\frac{4}{3} \alpha^2 + \Lambda \right), \\
 (6H\partial_t - 3\dot{H})F(\square)R &= -16\alpha^2 f_0 \dot{H} + \int_{\mathbb{R}} d\omega [-4i\alpha \tanh(\alpha t)\omega - 2\alpha^2(1 + \tanh^2(\alpha t))] F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega, \\
 -(2\partial_t^2 + 4H\partial_t + \dot{H})F(\square)R &= -\frac{16}{3} \alpha^2 f_0 \dot{H} + \int_{\mathbb{R}} d\omega \left[2\omega^2 - \frac{4}{3} i\alpha \tanh(\alpha t)\omega + \frac{2}{3} \alpha^2 (1 + \operatorname{sech}^2(\alpha t)) \right] F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega, \\
 \mp \frac{1}{2}(\Theta \pm \Omega_{tt}) &= \mp \frac{1}{2} \left[\left(\frac{16}{3} \alpha^2\right)^2 f_0 - \frac{16}{3} \alpha^2 f_0 R + \frac{16}{3} \alpha^2 \int_{\mathbb{R}} d\omega F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega \right] \\
 &\quad \mp \iint_{\mathbb{R}^2} d\omega d\omega' \left[\frac{1}{4} (\omega \mp \omega')^2 \pm \frac{1}{2} i\alpha \tanh(\alpha t) (\omega + \omega') + \frac{1}{2} \alpha^2 (1 \pm \tanh^2(\alpha t)) \right] \\
 &\quad \times G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'}. \tag{3.21}
 \end{aligned}$$

When put together, we finally arrive at the explicit formula for the energy-momentum tensor,

$$\begin{aligned}
T = & \left[\chi^{-1} \left(\frac{4}{3} \alpha^2 \tanh^2(\alpha t) - \Lambda \right) - \frac{128}{9} \alpha^4 f_0 \operatorname{sech}^2(\alpha t) \right. \\
& + \int_{\mathbb{R}} d\omega \left[-4i\alpha \tanh(\alpha t) \omega - 2\alpha^2 \left(\frac{7}{3} + \tanh^2(\alpha t) \right) \right] F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega \\
& - \iint_{\mathbb{R}^2} d\omega d\omega' \left[\frac{1}{4} (\omega - \omega')^2 + \frac{1}{2} i\alpha \tanh(\alpha t) (\omega + \omega') + \frac{1}{2} \alpha^2 (1 + \tanh^2(\alpha t)) \right] G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'} \mathbf{d}t \mathbf{d}t \\
& + \left[\chi^{-1} \left(-\frac{4}{3} \alpha^2 + \Lambda \right) + \int_{\mathbb{R}} d\omega \left[2\omega^2 - \frac{4}{3} i\alpha \tanh(\alpha t) \omega + \frac{2}{3} \alpha^2 (5 + \operatorname{sech}^2(\alpha t)) \right] F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega \right. \\
& \left. + \iint_{\mathbb{R}^2} d\omega d\omega' \left[\frac{1}{4} (\omega + \omega')^2 - \frac{1}{2} i\alpha \tanh(\alpha t) (\omega + \omega') + \frac{1}{2} \alpha^2 (1 - \tanh^2(\alpha t)) \right] G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'} \right] a^2 \mathbf{q}.
\end{aligned} \tag{3.22}$$

In the next section, we study its properties and show that the specific nonsingular geometry we chose, i.e., (3.8) with $\gamma = 2/3$, can be actually sourced by a perfect fluid with non-negative energy density that meets SEC.

IV. PROPERTIES OF THE PERFECT FLUID

The energy-momentum tensor (3.22) strongly depends on the form factor F through expressions involving $F(\omega^2 + \alpha^2)$ and $G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2)$ integrated over all values of ω and ω' . In other words, we have a greater freedom to choose a form factor F such that the energy-momentum tensor for a given geometry meets the desired properties. We first require that the energy density is non-negative in order for the perfect fluid to be physically reasonable. Secondly, we impose that the perfect fluid satisfies SEC. Note that this implies that a geometry that would be sourced by the same perfect fluid in GR would be singular.

This section is divided in two parts: In the first part, we prove that neither in GR nor in “ $R + R^2$ gravity” is it possible to meet the requirements on the energy-momentum tensor of the perfect fluid for this geometry. In the second part, we study these requirements in the nonlocal theory. In particular, we find some sufficient conditions on the form factors and also provide a specific numerical example of a form factor that fulfills these conditions.

A. Local gravity: GR & $R + R^2$

The energy-momentum tensor of the perfect fluid in the flat FLRW spacetime is given by

$$T = \rho \mathbf{d}t \mathbf{d}t + p a^2 \mathbf{q}, \tag{4.1}$$

where ρ is the energy density, and p is the pressure. Let us see whether the bouncing cosmology (3.8) with $\gamma = 2/3$ can be sourced by a physical matter. First, we consider the general relativity, i.e., $F(\square) = 0$. The condition on non-negative energy density reads

$$\rho = \chi^{-1} \left(\frac{4}{3} \alpha^2 \tanh^2(\alpha t) - \Lambda \right) \geq 0, \tag{4.2}$$

which is satisfied for all t only if $\Lambda \leq 0$. Next, we demand fulfilment of SEC,

$$\begin{aligned}
\rho + p &= -\frac{4}{3} \chi^{-1} \alpha^2 \operatorname{sech}^2(\alpha t) \geq 0, \\
\rho + 3p &= \chi^{-1} \left(\frac{4}{3} \alpha^2 \tanh^2(\alpha t) - 4\alpha^2 + 2\Lambda \right) \geq 0.
\end{aligned} \tag{4.3}$$

As we can see, the first condition can never hold, and the second one is met for all t only if $\Lambda \geq 2\alpha^2$. Therefore, the perfect fluid generating this geometry in GR does not meet SEC. (A special example is pure GR without the cosmological term, $\Lambda = 0$, where the violation of SEC can be also argued based on the Hawking theorem applied to this nonsingular bouncing FLRW spacetime.) Before we proceed to another theory, let us investigate what happens at late/early times $t \rightarrow \pm\infty$ (i.e., far from the bounce $t = 0$). The energy density (4.2) and SEC (4.3) turn into $\Lambda \leq 4\alpha^2/3$ and $\Lambda \geq 4\alpha^2/3$, respectively. This means that both conditions are asymptotically satisfied only if $\Lambda = 4\alpha^2/3$, which corresponds to the asymptotic vacuum, $T \rightarrow 0$ for $t \rightarrow \pm\infty$. For this specific choice $\Lambda = 4\alpha^2/3$, the cosmological term accounts for the accelerated expansion, where the vacuum corresponds to de Sitter spacetime with the Ricci scalar $R = 16\alpha^2/3$.

Let us investigate if this behavior can be improved in “ $R + R^2$ gravity”, which is given by a constant form factor $F(\square) = f_0$. The energy density is non-negative if

$$\rho = \chi^{-1} \left(\frac{4}{3} \alpha^2 \tanh^2(\alpha t) - \Lambda \right) - 8f_0 \alpha^4 \operatorname{sech}^4(\alpha t) \geq 0, \tag{4.4}$$

which holds for all t if $\Lambda \leq -8\chi f_0 \alpha^4$. SEC reads

$$\rho + p = -\frac{4}{3}\kappa^{-1}\alpha^2\text{sech}^2(\alpha t) - 16f_0\alpha^4\text{sech}^4(\alpha t) \geq 0, \quad |\omega|^k \text{sech}\left(\frac{\pi\omega}{2\alpha}\right) |F(\omega^2 + \alpha^2)|, \quad k = 0, 1, 2, \quad (4.6)$$

$$\rho + 3p = \kappa^{-1}\left(\frac{4}{3}\alpha^2\text{tanh}^2(\alpha t) - 4\alpha^2 + 2\Lambda\right) - 32f_0\alpha^4\text{sech}^4(\alpha t) \geq 0. \quad (4.5)$$

These two conditions are met for all t only if $f_0 \leq -\kappa^{-1}/12\alpha^2$ and $\Lambda \geq 2\alpha^2 + 16\kappa f_0\alpha^4$, respectively. Unfortunately, f_0 should be positive to avoid Dolgov–Kawasaki instability [55,56], which obviously violates SEC. Taking the limits $t \rightarrow \pm\infty$ of (4.4) and (4.5), we again arrive at $\Lambda \leq 4\alpha^2/3$ and $\Lambda \geq 4\alpha^2/3$, respectively. As in GR, both conditions are asymptotically satisfied only for $\Lambda = 4\alpha^2/3$ corresponding to the asymptotic vacuum. This is because the contributions to the energy-momentum tensor arising from the R^2 term in the action decay faster than the contributions from the Einstein–Hilbert term.

B. Nonlocal gravity: IDG

Let us see if the nonlocal theory described by the action of IDG (2.1) can improve the properties of the perfect fluid and give rise to a physically reasonable matter. Our aim is to find sufficient conditions for the form factor under which the perfect fluid meets the required properties. First, recall that the bouncing geometry tends to de Sitter spacetime for $t \rightarrow \pm\infty$. Its Hubble parameter and Ricci scalar approach constants $H = 2\alpha/3$ and $R = 16\alpha^2/3$ with exponential rate, see (3.9) (with $\gamma = 2/3$) and (3.17). Moreover, the derivatives $\partial_t^k H$ and $\partial_t^k R$ for $k > 0$ decay exponentially. Inspecting (3.5), we see that the contributions to the energy-momentum tensor from all terms except for the Einstein–Hilbert term become negligible for $t \rightarrow \pm\infty$, and thus, IDG effectively reduces to GR. This has an important implication for the requirements on the perfect fluid because, as we have seen, they can be satisfied asymptotically only if we set $\Lambda = 4\alpha^2/3$; all other choices of Λ would necessarily lead to negative energy density or violation of SEC.

At this point, we should analyze possible form factors that allow for non-negative energy density and meet SEC. The first restriction, however, comes already from the integral transform we used to evaluate the energy-momentum tensor. All (1- and 2- dimensional) integrals in (3.22) must converge, which already constraints the form factors that we can choose. We have to demand that the absolute values of integrands are integrable functions. Therefore, a sufficient condition for convergence of the 1-dimensional integrals over \mathbb{R} is that functions

belong to $L_1(\mathbb{R})$. To find the conditions of convergence, we realize that the integrands are analytic functions (F is analytic), so it is sufficient to investigate the convergence of the integrals over the intervals $(-\infty, c_-)$ and (c_+, ∞) for some constants c_- and c_+ . We rely on the limit comparison test for improper integral, which says that $\int_{c_\pm}^{\pm\infty} d\omega f(\omega)$ converges if $\int_{c_\pm}^{\pm\infty} d\omega g(\omega)$ is a known convergent integral and

$$\lim_{\omega \rightarrow \pm\infty} f(\omega)/g(\omega) = L_\pm, \quad 0 \leq L_\pm < \infty. \quad (4.7)$$

We use the slowly decreasing function (with a convergent improper integral), $g(\omega) = |\omega|^{-1-\varepsilon}$, $\varepsilon > 0$. Then, it is clear that the most restrictive condition that we can write among the three mentioned conditions ($k = 0, 1, 2$) is

$$\lim_{\omega \rightarrow \pm\infty} |\omega|^{3+\varepsilon} e^{\mp\frac{\pi\omega}{2\alpha}} |F(\omega^2 + \alpha^2)| = L_\pm, \quad 0 \leq L_\pm < \infty, \quad \varepsilon > 0. \quad (4.8)$$

Before we examine the convergence of the 2-dimensional integrals, it is important to convince ourselves that the integrands are also analytic, which may not be apparent because of the presence of the function G . Indeed, the function $G(x, y)$ is analytic everywhere including the points $x = y$, where it can be analytically extended by its limit $G(x, x) = F'(x)$. The conditions for the convergence of the 2-dimensional integrals over \mathbb{R}^2 can be obtained from the 2-dimensional Fourier transform conditions of existence [57], namely, the absolute integrability of the integrand. Hence, sufficient conditions for convergence are that functions

$$|\omega|^k |\omega'|^l \text{sech}\left(\frac{\pi\omega}{2\alpha}\right) \text{sech}\left(\frac{\pi\omega'}{2\alpha}\right) |G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2)|, \quad k + l = 0, 1, 2, \quad (4.9)$$

belong to $L_1(\mathbb{R}^2)$. As we did for the 1-dimensional integrals, we can use the limit test to study the convergence [58]. We can choose the slowly decreasing function with a convergent improper integral over $A_R = \{(x, y) \in \mathbb{R}^2 \text{ s.t. } x^2 + y^2 \geq R\}$, which is $g(\omega, \omega') = (\omega^2 + \omega'^2)^{-1-\varepsilon}$, $\varepsilon > 0$. Applying the same reasoning as in the 1-dimensional case, the most restrictive conditions coming from the limit convergence test are now

$$\lim_{(\omega, \omega') \rightarrow \infty} (\omega^2 + \omega'^2)^{1+\varepsilon} |\omega|^k |\omega'|^l e^{-\frac{\pi|\omega|}{2\alpha}} e^{-\frac{\pi|\omega'|}{2\alpha}} |G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2)| = L, \quad k + l = 2, \quad 0 \leq L < \infty, \quad \varepsilon > 0. \quad (4.10)$$

Having established the convergence of the integrals, we can now move on to the condition for non-negative energy density $\rho \geq 0$ and SEC $\rho + p \geq 0$, $\rho + 3p \geq 0$. Since we want to satisfy all these inequalities at the same time, we can easily see that $\rho + p \geq 0$ is actually redundant for our

needs: Let us assume that $\rho \geq 0$. If $p \geq 0$ then SEC holds. On the other hand, if $p < 0$ than $\rho + 3p \geq 0$ automatically implies $\rho + p \geq 0$. Therefore, we only need to deal with the quantities ρ and $\rho + 3p$, which can be read out from (3.22) (where we set $\Lambda = 4\alpha^2/3$),

$$\begin{aligned} \rho &= -\left(\frac{4}{3}\chi^{-1} + \frac{128}{9}\alpha^2 f_0\right)\alpha^2 \text{sech}^2(\alpha t) + \int_{\mathbb{R}} d\omega \left[-4i\alpha \tanh(\alpha t)\omega - 2\alpha^2 \left(\frac{7}{3} + \tanh^2(\alpha t)\right) \right] F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega \\ &\quad - \iint_{\mathbb{R}^2} d\omega d\omega' \left[\frac{1}{4}(\omega - \omega')^2 + \frac{1}{2}i\alpha \tanh(\alpha t)(\omega + \omega') + \frac{1}{2}\alpha^2(1 + \tanh^2(\alpha t)) \right] \\ &\quad \times G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'}, \\ \rho + 3p &= -\left(\frac{4}{3}\chi^{-1} + \frac{128}{9}\alpha^2 f_0\right)\alpha^2 \text{sech}^2(\alpha t) + \int_{\mathbb{R}} d\omega \left(\frac{10}{3}\alpha^2 + 6\omega^2 + 4\alpha^2 \text{sech}^2(\alpha t) - 8i\alpha\omega \tanh(\alpha t) \right) F(\omega^2 + \alpha^2) \hat{R}_\omega \psi_\omega \\ &\quad + \frac{1}{2} \iint_{\mathbb{R}^2} d\omega d\omega' [2\alpha^2 + \omega^2 + \omega'^2 + 4\omega\omega' + 4\alpha \tanh(\alpha t)(-\alpha \tanh(\alpha t) - i(\omega + \omega'))] \\ &\quad \times G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2) \hat{R}_\omega \hat{R}_{\omega'} \psi_\omega \psi_{\omega'}. \end{aligned} \quad (4.11)$$

Let us take a look at these expression term by term. If we want to have a positive contribution from the first term of ρ , we need

$$f_0 < -\frac{3\chi^{-1}}{32\alpha^2}. \quad (4.12)$$

To understand the second term (the 1-dimensional integral), we evaluate an auxiliary integral with the constant positive form factor $F(\omega^2 + \alpha^2) = C > 0$,

$$\begin{aligned} C \int_{\mathbb{R}} d\omega \left[-4i\alpha \tanh(\alpha t)\omega - 2\alpha^2 \left(\frac{7}{3} + \tanh^2(\alpha t)\right) \right] \hat{R}_\omega \psi_\omega \\ = \frac{8}{9} C \alpha^4 (16 \cosh^2(\alpha t) - 9) \text{sech}^4(\alpha t). \end{aligned} \quad (4.13)$$

Since this integral is always positive, we might be tempted to assume that $F(x)$ is positive and approximately constant for all x in order to make the second term of ρ also positive. Nevertheless, this obviously contradicts the condition (4.12). To overcome this issue, we propose that $F(x)$ should be almost constant and positive ($\approx C > 0$) only for $x > \alpha^2$ but negative for $x \in (0, \zeta^2)$, with $\alpha > \zeta > 0$, so that f_0 meets the requirement (4.12). Although the third term (the 2-dimensional integral) might be negative, it should not contribute too much because $G(\omega^2 + \alpha^2, \omega'^2 + \alpha^2)$ is almost zero for all $(\omega, \omega') \in \mathbb{R}^2$.

Moving on to $\rho + 3p$, we realize that its first term is the equal to the first term of ρ , so it is automatically positive thanks to (4.12). The second term of $\rho + 3p$ (the single integral) can be analyzed again by calculating the approximate integral with $F(\omega^2 + \alpha^2) = C > 0$,

$$\begin{aligned} C \int_{\mathbb{R}} d\omega \left(\frac{10}{3}\alpha^2 + 6\omega^2 + 4\alpha^2 \text{sech}^2(\alpha t) - 8i\alpha\omega \tanh(\alpha t) \right) \hat{R}_\omega \psi_\omega \\ = \frac{32}{9} C \alpha^4 (4 \sinh^2(\alpha t) - 5) \text{sech}^4(\alpha t), \end{aligned} \quad (4.14)$$

which is positive only for $|t| > \text{arcsinh}(\sqrt{5}/2)/\alpha$. Let us see if we can counter the negativity of this term with the positivity of the first term of $\rho + 3p$. By summing the first term of $\rho + 3p$ with the negative contribution of (4.14) only, we obtain

$$-\left(\frac{4}{3}\chi^{-1} + \frac{128}{9}\alpha^2 f_0\right)\alpha^2 \text{sech}^2(\alpha t) - \frac{160}{9} C \alpha^4 \text{sech}^4(\alpha t). \quad (4.15)$$

Since $\text{sech}^2(x) \geq \text{sech}^4(x)$ for all x , this expression is positive if

$$f_0 < -\frac{3\chi^{-1}}{32\alpha^2} - \frac{5}{4} C. \quad (4.16)$$

The third term of $\rho + 3p$ (the 2-dimensional integral) is negligible for the same reason as the analogous term in the analysis of ρ .

To summarize, we have arrived at the following sufficient conditions for the form factor leading to a physically reasonable perfect fluid:

- (i) The form factor must ensure conditions for convergence of integrals appearing in the energy-momentum tensor, i.e., (4.8) and (4.10).
- (ii) The form factor must be almost constant and positive, $F(x) \approx C > 0$, for $x > \alpha^2$.




FIG. 1. Graphs of the dimensionless quantities $x\rho/\alpha^2$ (left) and the quantity $x(\rho + 3p)/\alpha^2$ (right) of the perfect fluid that sources the bouncing cosmology for the form factor given in (4.17), for three different values of $\alpha\ell$. We have chosen $C\alpha^2 = 1.3225$ and $\zeta/\alpha = 0.3887$, which satisfy (4.18) and (4.19) with $\epsilon = 0.034$.

(iii) The value of $f_0 \equiv F(0)$ has to be sufficiently negative to meet condition (4.16).

In order to verify these findings, we provide an explicit example of a form factor that gives rise to non-negative energy density and the fulfillment of SEC. Let us consider the following form factor:

$$F(\square) = C - Ce^{-\ell^2(\square - \zeta^2)}, \quad (4.17)$$

for which we have $f_0 = C - Ce^{\ell^2\zeta^2}$. This form factor has been constructed to recover the local limit, i.e., $F(\square) = 0$, for $\ell \rightarrow 0$. First, let us note that the function (4.17) obeys the convergence conditions (4.8) and (4.10) with $L_{\pm} = 0$ and $L = 0$, respectively. Next, we notice that it quickly approaches a constant value. To ensure that $F(x)$ is almost constant for $x > \alpha^2$, we propose

$$e^{\ell^2(\zeta^2 - \alpha^2)} < \epsilon, \quad (4.18)$$

where $\epsilon > 0$ is a sufficiently small number. Finally, we make f_0 negative enough according to (4.16),

$$e^{\ell^2\zeta^2} > \frac{3}{4} \left(\frac{x^{-1}}{8\alpha^2 C} + 3 \right). \quad (4.19)$$

For a given constant α , we can always find the values of C , ℓ , and ζ so that the constraints (4.18) and (4.19) are satisfied to a prescribed value of ϵ . The graph of the functions ρ and $\rho + 3p$ computed numerically from (4.11) are plotted in Fig. 1 in dimensionless quantities. We can see that the energy density is positive, and SEC is satisfied everywhere. Hence, in this nonlocal modification, one can obtain a bouncing cosmology sourced by a physically relevant perfect fluid. Finally, it is interesting to note that for this particular form factor (4.17), we recover the $R - \beta R^2$ gravity for large $\alpha\ell$, which we have rendered as an unphysical theory due to its instabilities. In order to have a nonlocal behavior that fulfills SEC, we need to choose ℓ to be of the same order as α , as

can be seen in Fig. 1. This means that the nonlocality scale should be of the same order as the de Sitter radius r_{ds} , because $r_{\text{ds}} = \sqrt{3/\Lambda} = \frac{3}{2}\alpha^{-1}$. Finally, it is interesting to note that the effects of nonlocality are only appreciable around the bounce, and that GR is recovered as one distances from it.

V. CONCLUSIONS

In this paper, we found a new cosmological bouncing geometry for which the field equations can be evaluated using the spectral decomposition with respect to the eigenfunctions of the wave operator. Furthermore, dependence of the resulting energy-momentum tensor (3.22) on the form factor is much stronger than for the previous solutions available in the literature. Since the new solution does not satisfy the recursive ansatz, $\square R = r_1 R + r_2$, the Ricci scalar (after a constant shift) is no longer an eigenfunction on its own, but it still admits eigenfunction decompositions. The geometry we choose is rather special in the sense that the eigenvalue problem of the wave operator in this background is exactly solvable, and the corresponding eigenfunctions are rather simple.

We showed that due to stronger dependence of the energy momentum on the form factor $F(\square)$ one can tune it in such a way that the following occur: (i) The integrals appearing in the formula for the energy-momentum tensor converge. (ii) The solution is sourced by a perfect fluid with non-negative energy density. (iii) The perfect fluid satisfies SEC. One of the simplest form factors satisfying (i)–(iii) can be constructed rather straightforwardly by taking $F(x)$ to be sufficiently negative at $x = 0$ and almost constant and positive in the domain of integration. We have also shown that the effects of nonlocality are only relevant around the bounce, and that GR is recovered at late times.

A natural extension of this work is to find other types of form factors satisfying i)–iii) [especially those for which $F(0) \geq 0$]. Another interesting but rather involved

problem is the perturbative stability of the solution, which may put further restrictions on the form factor. It will also be worth exploring more general solutions for which the eigenvalue problem of the d'Alembertian operator can be solved.

ACKNOWLEDGMENTS

We would like to thank to Sravan Kumar (Tokyo, Japan) for useful discussions. I. K. and A. M. were supported by the Netherlands Organization for Scientific Research (NWO) Grant No. 680-91-119.

F. J. M. T. acknowledges financial support from NRF Grants No. 120390, reference BSFP190416431035; No. 120396, reference CSRP190405427545; and No. 101775, reference SFH150727131568. F. J. M. T. is supported by the “Fundación Ramón Areces” and the European Regional Development Fund through the Center of Excellence TK133 “The Dark Side of the Universe”. F. J. M. T. would also like to thank the hospitality of the Institute of Theoretical Astrophysics of the University of Oslo, and the Research Council of Norway for the economical support.

-
- [1] P. A. R. Ade *et al.* (Planck Collaboration), Planck 2015 results. XIII. Cosmological parameters, *Astron. Astrophys.* **594**, A13 (2016).
- [2] A. G. Riess, The expansion of the universe is faster than expected, *Nat. Rev. Phys.* **2**, 10 (2020).
- [3] S. Weinberg, The cosmological constant problem, *Rev. Mod. Phys.* **61**, 1 (1989).
- [4] J. Martin, Everything you always wanted to know about the cosmological constant problem (but were afraid to ask), *C.R. Phys.* **13**, 566 (2012).
- [5] M. Novello and S. E. P. Bergliaffa, Bouncing cosmologies, *Phys. Rep.* **463**, 127 (2008).
- [6] J. M. M. Senovilla, Singularity theorems and their consequences, *Gen. Relativ. Gravit.* **30**, 701 (1998).
- [7] R. Penrose, Gravitational Collapse and Space-Time Singularities, *Phys. Rev. Lett.* **14**, 57 (1965).
- [8] S. Hawking, The occurrence of singularities in cosmology. III. Causality and singularities, *Proc. R. Soc. A* **300**, 187 (1967).
- [9] S. W. Hawking and R. Penrose, The singularities of gravitational collapse and cosmology, *Proc. R. Soc. A* **314**, 529 (1970).
- [10] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*, Cambridge Monographs on Mathematical Physics (Cambridge University Press, Cambridge, England, 2011).
- [11] R. M. Wald, *General Relativity* (Chicago University Press, Chicago, USA, 1984).
- [12] M. Novello and J. M. Salim, Nonlinear photons in the Universe, *Phys. Rev. D* **20**, 377 (1979).
- [13] V. N. Melnikov and S. V. Orlov, Nonsingular cosmology as a quantum vacuum effect, *Phys. Lett. A* **70**, 263 (1979).
- [14] C. Molina-Paris and M. Visser, Minimal conditions for the creation of a Friedman-Robertson-Walker universe from a ‘bounce’, *Phys. Lett. B* **455**, 90 (1999).
- [15] C. Cattoen and M. Visser, Necessary and sufficient conditions for big bangs, bounces, crunches, rips, sudden singularities, and extremality events, *Classical Quantum Gravity* **22**, 4913 (2005).
- [16] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, Modified gravity theories on a nutshell: Inflation, bounce and late-time evolution, *Phys. Rep.* **692**, 1 (2017).
- [17] N. V. Krasnikov, Nonlocal gauge theories, *Theor. Math. Phys.* **73**, 1184 (1987).
- [18] Y. V. Kuz'min, Finite nonlocal gravity, *Yad. Fiz.* **50**, 1630 (1989) [*Sov. J. Nucl. Phys.* **50**, 1011 (1989)].
- [19] T. Biswas, A. Mazumdar, and W. Siegel, Bouncing universes in string-inspired gravity, *J. Cosmol. Astropart. Phys.* **03** (2006) 009.
- [20] T. Biswas, E. Gerwick, T. Koivisto, and A. Mazumdar, Towards Singularity and Ghost Free Theories of Gravity, *Phys. Rev. Lett.* **108**, 031101 (2012).
- [21] T. Biswas, A. S. Koshelev, and A. Mazumdar, Gravitational theories with stable (anti-)de Sitter backgrounds, *Fundam. Theor. Phys.* **183**, 97 (2016).
- [22] E. T. Tomboulis, Superrenormalizable gauge and gravitational theories, [arXiv:hep-th/9702146](https://arxiv.org/abs/hep-th/9702146).
- [23] L. Modesto, Super-renormalizable quantum gravity, *Phys. Rev. D* **86**, 044005 (2012).
- [24] S. Talaganis, T. Biswas, and A. Mazumdar, Towards understanding the ultraviolet behavior of quantum loops in infinite-derivative theories of gravity, *Classical Quantum Gravity* **32**, 215017 (2015).
- [25] S. Abel and N. A. Dondi, UV completion on the worldline, *J. High Energy Phys.* **07** (2019) 090.
- [26] S. Abel, L. Buoninfante, and A. Mazumdar, Nonlocal gravity with worldline inversion symmetry, *J. High Energy Phys.* **01** (2020) 003.
- [27] L. Buoninfante, G. Harmsen, S. Maheshwari, and A. Mazumdar, Nonsingular metric for an electrically charged point-source in ghost-free infinite derivative gravity, *Phys. Rev. D* **98**, 084009 (2018).
- [28] V. P. Frolov and A. Zelnikov, Head-on collision of ultra-relativistic particles in ghost-free theories of gravity, *Phys. Rev. D* **93**, 064048 (2016).
- [29] J. Boos, V. P. Frolov, and A. Zelnikov, Gravitational field of static p-branes in linearized ghost-free gravity, *Phys. Rev. D* **97**, 084021 (2018).
- [30] L. Buoninfante, A. S. Cornell, G. Harmsen, A. S. Koshelev, G. Lambiase, J. Marto, and A. Mazumdar, Towards nonsingular rotating compact object in ghost-free infinite derivative gravity, *Phys. Rev. D* **98**, 084041 (2018).

- [31] I. Kolář and A. Mazumdar, NUT charge in linearized infinite derivative gravity, *Phys. Rev. D* **101**, 124005 (2020).
- [32] J. Boos, J. Pinedo Soto, and V. P. Frolov, Ultrarelativistic spinning objects in nonlocal ghost-free gravity, *Phys. Rev. D* **101**, 124065 (2020).
- [33] I. Kolář and J. Boos, Retarded field of a uniformly accelerated source in nonlocal scalar field theory, *Phys. Rev. D* **103**, 105004 (2021).
- [34] V. P. Frolov, A. Zelnikov, and T. de Paula Netto, Spherical collapse of small masses in the ghost-free gravity, *J. High Energy Phys.* **06** (2015) 107.
- [35] V. P. Frolov, Mass-Gap for Black Hole Formation in Higher Derivative and Ghost Free Gravity, *Phys. Rev. Lett.* **115**, 051102 (2015).
- [36] A. S. Koshelev and A. Mazumdar, Do massive compact objects without event horizon exist in infinite derivative gravity?, *Phys. Rev. D* **96**, 084069 (2017).
- [37] L. Buoninfante and A. Mazumdar, Nonlocal star as a blackhole mimicker, *Phys. Rev. D* **100**, 024031 (2019).
- [38] E. Kilicarslan, pp -waves as exact solutions to ghost-free infinite derivative gravity, *Phys. Rev. D* **99**, 124048 (2019).
- [39] S. Dengiz, E. Kilicarslan, I. Kolář, and A. Mazumdar, Impulsive waves in ghost free infinite derivative gravity in anti-de Sitter spacetime, *Phys. Rev. D* **102**, 044016 (2020).
- [40] I. Kolář, T. Málek, and A. Mazumdar, Exact solutions of nonlocal gravity in a class of almost universal spacetimes, *Phys. Rev. D* **103**, 124067 (2021).
- [41] I. Kolář, T. Málek, S. Dengiz, and E. Kilicarslan, Exact gyratons in higher and infinite derivative gravity, *Phys. Rev. D* **105**, 044018 (2022).
- [42] A. S. Koshelev, K. S. Kumar, and A. A. Starobinsky, R^2 inflation to probe non-perturbative quantum gravity, *J. High Energy Phys.* **03** (2018) 071.
- [43] K. S. Kumar and L. Modesto, Non-local Starobinsky inflation in the light of future CMB, [arXiv:1810.02345](https://arxiv.org/abs/1810.02345).
- [44] A. S. Koshelev, K. S. Kumar, A. Mazumdar, and A. A. Starobinsky, Non-Gaussianities and tensor-to-scalar ratio in non-local R^2 -like inflation, *J. High Energy Phys.* **06** (2020) 152.
- [45] A. S. Koshelev, K. S. Kumar, and A. A. Starobinsky, Analytic infinite derivative gravity, R^2 -like inflation, quantum gravity and CMB, *Int. J. Mod. Phys. D* **29**, 2043018 (2020).
- [46] T. Biswas, T. Koivisto, and A. Mazumdar, Towards a resolution of the cosmological singularity in non-local higher derivative theories of gravity, *J. Cosmol. Astropart. Phys.* **11** (2010) 008.
- [47] A. S. Koshelev and S. Y. Vernov, On bouncing solutions in non-local gravity, *Phys. Part. Nucl.* **43**, 666 (2012).
- [48] A. S. Koshelev, Stable analytic bounce in non-local Einstein-Gauss-Bonnet cosmology, *Classical Quantum Gravity* **30**, 155001 (2013).
- [49] T. Biswas, A. S. Koshelev, A. Mazumdar, and S. Y. Vernov, Stable bounce and inflation in non-local higher derivative cosmology, *J. Cosmol. Astropart. Phys.* **08** (2012) 024.
- [50] K. S. Kumar, S. Maheshwari, A. Mazumdar, and J. Peng, Stable, nonsingular bouncing universe with only a scalar mode, *Phys. Rev. D* **102**, 024080 (2020).
- [51] K. S. Kumar, S. Maheshwari, A. Mazumdar, and J. Peng, An anisotropic bouncing universe in non-local gravity, *J. Cosmol. Astropart. Phys.* **07** (2021) 025.
- [52] K. S. Kumar, S. Maheshwari, A. Mazumdar, and J. Peng, An anisotropic bouncing universe in non-local gravity, *J. Cosmol. Astropart. Phys.* **07** (2021) 025.
- [53] V. P. Frolov and A. Zelnikov, Bouncing cosmology in the limiting curvature theory of gravity, *Phys. Rev. D* **104**, 104060 (2021).
- [54] T. Biswas, A. Conroy, A. S. Koshelev, and A. Mazumdar, Generalized ghost-free quadratic curvature gravity, *Classical Quantum Gravity* **31**, 015022 (2014); **31**, 159501(E) (2014).
- [55] A. D. Dolgov and M. Kawasaki, Can modified gravity explain accelerated cosmic expansion?, *Phys. Lett. B* **573**, 1 (2003).
- [56] V. Faraoni, Matter instability in modified gravity, *Phys. Rev. D* **74**, 104017 (2006).
- [57] J. W. Goodman, *Introduction to Fourier Optics*. 2005 (Roberts and Company, Englewood, Colorado, 2008).
- [58] S. R. Ghorpade and B. V. Limaye, *A Course in Multivariable Calculus and Analysis* (Springer, New York, 2010).