

## Infinitely many 4D $\mathcal{N} = 1$ SCFTs with $a = c$

Monica Jinwoo Kang<sup>1,2,\*</sup> Craig Lawrie,<sup>3,†</sup> Ki-Hong Lee,<sup>2,‡</sup> and Jaewon Song<sup>2,§</sup>

<sup>1</sup>*Walter Burke Institute for Theoretical Physics, California Institute of Technology  
Pasadena, California 91125, USA*

<sup>2</sup>*Department of Physics, Korea Advanced Institute of Science and Technology  
Daejeon 34141, Republic of Korea*

<sup>3</sup>*Deutsches Elektronen-Synchrotron DESY Notkestr. 85, 22607 Hamburg, Germany*



(Received 22 December 2021; accepted 3 May 2022; published 2 June 2022)

We study a rich set of four-dimensional  $\mathcal{N} = 1$  superconformal field theories (SCFTs) with both central charges identical:  $a = c$ . We construct them via the diagonal  $\mathcal{N} = 1$  gauging of the flavor symmetry  $G$  of a collection of  $\mathcal{N} = 2$  Argyres–Douglas theories of type  $\mathcal{D}_p(G)$ , with or without additional adjoint chiral multiplets. In this way, we construct infinitely many theories that flow to interacting SCFTs with  $a = c$  in the infrared. Finally, we briefly highlight the features of the SCFTs without  $a = c$  that arise from generalizing this construction.

DOI: [10.1103/PhysRevD.105.126006](https://doi.org/10.1103/PhysRevD.105.126006)

### I. INTRODUCTION

When a conformal field theory in four dimensions is put on a curved manifold, the conformal symmetry becomes anomalous and characterized by two quantities  $a$  and  $c$ , commonly referred to as the central charges. The trace of the energy-momentum tensor is nonvanishing and is given by

$$T_{\mu}{}^{\mu} = cW^2 - aE_4, \quad (1.1)$$

where  $W^2$  is the square of the Weyl tensor, and  $E_4$  is the Euler density. It has been shown that the central charge  $a$  is a monotonically decreasing function along renormalization group (RG) flow [1],

$$a_{\text{IR}} < a_{\text{UV}}, \quad (1.2)$$

which justifies the intuitive notion of treating the central charge  $a$  as a measure for counting the degrees of freedom in a field theory. On the other hand, the central charge  $c$  is *not* a monotonically decreasing function along the RG flow. In many examples (especially with higher degrees of supersymmetry), the  $c$  function mostly decreases along the flow, but there are examples in which we get  $c_{\text{IR}} > c_{\text{UV}}$ .<sup>1</sup>

Unitarity puts constraints on the ratio of the central charges  $a/c$  to be [3]

$$\frac{1}{3} \leq \frac{a}{c} \leq \frac{31}{18}, \quad (1.3)$$

where the lower bound is saturated by the theory of a free scalar boson, and the upper bound is saturated by the free vector. For supersymmetric theories, the bound becomes narrower:

$$\mathcal{N} = 1 \text{ SCFTs: } \frac{1}{2} \leq \frac{a}{c} \leq \frac{3}{2}, \quad (1.4)$$

$$\mathcal{N} = 2 \text{ SCFTs: } \frac{1}{2} \leq \frac{a}{c} \leq \frac{5}{4}. \quad (1.5)$$

The lower bound is saturated by a free chiral multiplet or a free hypermultiplet, and the upper bound is saturated by an  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  vector multiplet, respectively. With higher supersymmetry, it is known that the central charges are equal [4]:

$$\mathcal{N} = 3, 4 \text{ SCFTs: } a = c. \quad (1.6)$$

Apart from  $\mathcal{N} = 3, 4$  superconformal theories, the central charges  $a$  and  $c$  are not generally related to each other. However, for the theories that are holographically dual to  $\text{AdS}_5$  gravity, the two central charges are equal (i.e.,  $a = c$ ) in the large  $N$  limit [5]. For a finite  $N$ , the difference of the central charges ( $a - c$ ) gives rise to the  $R^{\mu\nu\rho\sigma}R_{\mu\nu\rho\sigma}$  correction in the effective gravity action [5,6]. This results in corrections to the celebrated entropy-viscosity ratio bound [7] to give [8,9]

\*monica@caltech.edu

†craig.lawrie1729@gmail.com

‡khlee11812@gmail.com

§jaewon.song@kaist.ac.kr

<sup>1</sup>A simple example in the SUSY setup where  $c$  does not decrease along the RG flow can be constructed by considering  $\mathcal{N} = 1$   $SU(2)$  gauge theory with one adjoint and two fundamental chiral multiplets. See the flow from  $\mathcal{T}_0$  theory to  $H_0$  theory in [2].

$$\frac{\eta}{s} \geq \frac{1}{4\pi} \left( 1 - \frac{c-a}{c} + \dots \right). \quad (1.7)$$

The difference between the central charges ( $a - c$ ) also controls universal behavior of certain quantities of SCFTs, such as the Cardy-like limit of the superconformal index [10] and the entanglement entropy [11]. This difference ( $a - c$ ) also controls phenomena in CFT and holography such as the mixed current-gravitational anomaly [12] and the size of the single-trace higher spin gap for large  $N$  [13].

Then, a question to ponder would be whether one can have theories with  $a = c$  even in finite  $N$  without a high degree of supersymmetry and which, in turn, would comprise a rather special set of CFTs. In the context of 4D  $\mathcal{N} = 2$  SCFTs, families of such theories with  $a = c$  have been found to exist in [14]. What we find in this paper is that 4D SCFTs with  $a = c$  exist with the minimal amount of supersymmetry (namely,  $\mathcal{N} = 1$ ) as well, and the set of such theories is quite broad.

An interesting set of 4D  $\mathcal{N} = 2$  SCFTs is studied in [14] called  $\hat{\Gamma}(G)$ , associated to a choice of an affine ADE Dynkin diagram  $\hat{\Gamma}$  and an ADE gauge group  $G$ . These theories are obtained via gauging all of the  $G$  flavor symmetries of a collection of Argyres–Douglas and conformal matter theories. For a choice of  $\Gamma = D_4, E_6, E_7, E_8$  and particular choices of  $G$ , there is a fascinating connection to  $\mathcal{N} = 4$  super Yang–Mills (SYM) theory, whereby their central charges are identical,

$$a(\hat{\Gamma}(G)) = c(\hat{\Gamma}(G)), \quad (1.8)$$

and their Schur indices can be rewritten in terms of the Schur index of  $\mathcal{N} = 4$  SYM,

$$I_{\hat{\Gamma}(G)}(q) = I_G^{\mathcal{N}=4}(q^{\alpha_\Gamma}, q^{\frac{\alpha_\Gamma}{2}-1}), \quad (1.9)$$

where  $\alpha_\Gamma$  is the maximal coroot of  $\hat{\Gamma}$  [14]. This relationship occurs when<sup>2</sup>

$$\gcd(\alpha_\Gamma, h_G^\vee) = 1. \quad (1.10)$$

A natural question is then whether there exist  $\mathcal{N} = 1$  theories that have a similar connection to  $\mathcal{N} = 4$  SYM, as we just witnessed for  $\mathcal{N} = 2$  theories. We can easily generate a set of  $\mathcal{N} = 1$  SCFTs by deforming  $\hat{\Gamma}(G)$  theories via a mass term for the  $\mathcal{N} = 1$  adjoint chiral multiplet  $\phi$  in the  $\mathcal{N} = 2$  vector multiplet of  $G$ :

$$W = \frac{1}{2} m \text{Tr} \phi^2. \quad (1.11)$$

<sup>2</sup>In fact, this relationship between the Schur indices holds whenever the SCFT  $\hat{\Gamma}(G)$  has no flavor symmetry. Equation (1.10) is a sufficient, but not necessary, condition for the absence of flavor.

Upon renormalization group flow (below the mass scale set by  $m$ ), the mass-deformed theory flows to an  $\mathcal{N} = 1$  superconformal theory with the central charges given by [15]

$$a_{\mathcal{N}=1} = \frac{27}{32} a_{\mathcal{N}=2}, \quad c_{\mathcal{N}=1} = \frac{27}{32} c_{\mathcal{N}=2}, \quad (1.12)$$

which means that the mass-deformed  $\hat{\Gamma}(G)$  theories also have equal central charges  $a = c$ . The mass term generates a marginal coupling of the form,

$$W = \text{Tr} \left( \sum_i \mu_i \right)^2, \quad (1.13)$$

where we omit the coupling constant. The mother  $\mathcal{N} = 2$  SCFT has a marginal gauge coupling, and the mass deformed  $\mathcal{N} = 1$  theory also has an exactly marginal coupling given by this term [16,17]. This is a direct analog of the quartic coupling of  $SU(N)$  superconformal quantum chromodynamics (SQCD) with  $2N$  flavors that appears in many contexts.

We also obtain an  $\mathcal{N} = 1$  theory with  $a = c$  by simply replacing the  $\mathcal{N} = 2$  gauge multiplet in the  $\hat{\Gamma}(G)$  construction by an  $\mathcal{N} = 1$  gauge multiplet. We may expect that this theory is identical to the mass deformation of the mother  $\mathcal{N} = 2$  theory with the famous 27/32 ratio of central charges up on marginal deformation. However, we see that this is not the case. In fact, what we find in general is that there exists a  $W = 0$  fixed point (without any superpotential), at which the central charge  $a_{\mathcal{N}=1}$  is bigger than  $(27/32)a_{\mathcal{N}=2}$ , and where some operators of the form  $\text{Tr} \mu_i \mu_j$  are *relevant*. Upon deforming by these relevant operators, the theory flows to the mass-deformed theory with the central charge ratio given by 27/32. This is reminiscent of the phenomenon that appears in  $\mathcal{N} = 1$  class  $\mathcal{S}$  theory [18–20], where the mass-deformed theory corresponds to the case with equal degrees of normal bundles of more general compactifications. We depict such RG flows in Fig. 1. Besides these theories obtained via simple (universal) relevant deformation of  $\hat{\Gamma}(G)$  theories, we find there exist a wider class of theories with  $a = c$  that have no direct  $\mathcal{N} = 2$  origin. Such an  $\mathcal{N} = 1$  theory with  $a = c$  can be constructed via gauging several  $\mathcal{D}_{p_i}(G)$  theories with an  $\mathcal{N} = 1$  vector multiplet and possibly additional chiral multiplets in the adjoint representation of  $G$ . For some special choices, we can reproduce the mass-deformed  $\hat{\Gamma}(G)$  theory.

To verify that the 4D  $\mathcal{N} = 1$  SCFTs that we obtain in this manner truly are interacting SCFTs with  $a = c$ , it is necessary to check that they do not have a decoupled free sector, after flowing into the infrared. In this paper, we check that the Coulomb branch operators and products of the moment maps in the gauged theory satisfy the necessary




FIG. 1. RG flows triggered by the mass deformation of the  $\mathcal{N} = 2$  theory versus gauging an  $\mathcal{N} = 1$  vector multiplet. The red cross is to emphasize that there is no direct renormalization group flow between the  $\mathcal{N} = 2$  gauging and the  $\mathcal{N} = 1$  gauging.

unitarity conditions. A more refined test of unitarity involves the computation of the superconformal index; in [21], we perform this computation and confirm that the  $a = c$  theories constructed herein are indeed interacting SCFTs. The index also aids in the identification of the relevant operators of the theory; superpotential deformations via these operators may then trigger a flow to a new infrared SCFT. Intriguingly, we find that many of these deformations preserve the  $a = c$  property, and we will explore the landscape of such deformations in [22].

The structure of this paper is as follows. In Sec. II, we explain how and when the  $\mathcal{N} = 1$  gaugings of a collection of  $\mathcal{D}_{p_i}(G)$  theories lead to infrared SCFTs with  $a = c$ . In Sec. II A, we show precisely under which conditions any such gauging will flow to a theory with identical central charges, and in Sec. II B, we determine which particular combinations of  $p_i$  can be gauged in an asymptotically free or conformal manner. In Secs. II C, II D, II E, II F, and II G, we construct the infrared theories obtained from all the asymptotically free gaugings, with one to five  $\mathcal{D}_p(G)$ , respectively; and in Sec. II H, we analyze one instance of a conformal gauging that involves six  $\mathcal{D}_2(G)$ . We study if the constructed infrared theories are superconformal field theories and whether they satisfy the unitarity requirements to have  $a = c$ . In Sec. III, we consider gauging  $\mathcal{D}_{p_i}(G)$  together with additional matter charged under  $G$ . We explore gaugings with an additional adjoint chiral multiplet in Sec. III A, and we relate those to the mass-deformed  $\hat{\Gamma}(G)$  theories in Sec. III B. In Sec. III C, we consider a simple Lagrangian SCFT with  $a = c$ , which has two adjoint chirals and no  $\mathcal{D}_p(G)$ , and then in Sec. III D, we enumerate all possible gaugings of  $\mathcal{D}_p(G)$ s with two adjoint chirals such that the infrared SCFT has  $a = c$ . Finally, in Sec. IV, we consider a generalization to gaugings of collections of  $\mathcal{D}_p(G)$  theories and  $(G, G)$  conformal matter theories. We summarize and suggest some future directions in Sec. V.

## II. $\mathcal{N} = 1$ GLUING OF $\mathcal{D}_p(G)$ THEORIES

We are interested in constructing 4D  $\mathcal{N} = 1$  superconformal field theories starting from the non-

Lagrangian 4D  $\mathcal{N} = 2$  SCFTs known as the  $\mathcal{D}_p(G)$  Argyres–Douglas theories [23–26].<sup>3</sup> An interesting set of 4D  $\mathcal{N} = 2$  SCFTs, constructed out of  $\mathcal{D}_p(G)$  theories and  $(G \times G)$  conformal matter, called  $\hat{\Gamma}(G)$ , are studied in [14], where  $\Gamma$  and  $G$  are algebras of type ADE.<sup>4</sup> Specifically, the authors considered a diagonal gauging of the  $G$  flavor symmetry of a collection of  $\mathcal{D}_{p_i}(G)$  and  $(G \times G)$  conformal matter theories in such a way that, for each introduced gauge node, the introduced gauge coupling has a vanishing one-loop  $\beta$  function. In this paper, we consider an analogous construction for 4D  $\mathcal{N} = 1$  SCFTs via the diagonal gauging of the flavor symmetry  $G$  of a collection of  $\mathcal{D}_{p_i}(G)$ .

There are several subtleties that arise in the  $\mathcal{N} = 1$  case that are absent in the  $\mathcal{N} = 2$  gauging. In the simplest setup, we consider gauging by including in the theory a single  $\mathcal{N} = 1$  vector multiplet (in the adjoint representation of  $G$ ) and coupling the vector multiplet and the  $G$ -flavor currents. Post-gauging, we are interested in obtaining a superconformal field theory after performing the renormalization group flow into the infrared. In order to obtain a nontrivial theory in the IR, the gauge coupling for  $G$  must be asymptotically free. We could also consider the case where the one-loop  $\beta$  function for the gauge coupling is exactly zero; however, we then require the existence of exactly marginal operators for the theory to be an interacting SCFT. Such operators are not necessarily guaranteed to exist; however, we find that in some examples they do exist. In this paper, we will not study the conformal gaugings in detail; however, the properties of the infrared SCFTs, when they exist, can be determined via a straightforward application of the methods discussed herein. Furthermore, asymptotic freedom is not sufficient to guarantee the existence of an interacting SCFT in the infrared; we must

<sup>3</sup>Some of the  $\mathcal{D}_p(G)$  theories are actually  $\mathcal{N} = 2$  Lagrangian quivers, and sometimes they admit an  $\mathcal{N} = 1$  Lagrangian description [27–31]. We remind the reader that the  $\mathcal{D}_p(SU(2))$  theory is identical to the  $(A_1, D_p)$  theory.

<sup>4</sup>When  $\Gamma = D_4, E_6, E_7, E_8$ , these theories are known as the elliptic  $G$  models of [32]; some aspects of some of those particular theories have also been explored in [24,33–36].

check that there exists a nonanomalous superconformal  $R$ -symmetry. The superconformal  $R$ -symmetry in the infrared can be determined via  $a$ -maximization [37] and involves mixing between the  $R$ -symmetry in the ultraviolet and (possibly emergent) Abelian flavor symmetries of the theory.

### A. $a = c$ for $\mathcal{N} = 1$ gauging

One of the fascinating features of the  $\hat{\Gamma}(G)$  theories, which are the 4D  $\mathcal{N} = 2$  SCFTs studied in [14], is that a number of cases have their central charges being identical:  $a = c$ . In this section, we prove that many of the  $\mathcal{N} = 1$  gaugings share this feature.

The  $\mathcal{N} = 1$  gauging breaks the  $\mathcal{N} = 2$   $R$ -symmetry of the  $\mathcal{D}_p(G)$  theory to the  $U(1)$   $R$ -symmetry and a flavor  $U(1)$  symmetry, with generators,

$$R_0 = \frac{1}{3}R_{\mathcal{N}=2} + \frac{4}{3}I_3, \quad \mathcal{F} = -R_{\mathcal{N}=2} + 2I_3, \quad (2.1)$$

where  $R_{\mathcal{N}=2}$  and  $I_3$ , respectively, denote the  $U(1)_R$  charge and the  $SU(2)_R$  Cartan of the  $\mathcal{N} = 2$   $R$ -symmetry. Upon gauging, only the anomaly-free combinations are preserved. At the IR fixed point, the flavor  $\mathcal{F}_i$  of each  $\mathcal{D}_{p_i}(G)$  can mix with the  $R$ -symmetry before gauging,

$$R = R_0 + \sum_i \epsilon_i \mathcal{F}_i, \quad (2.2)$$

and thus, the  $R$ -charge gets modified in the infrared. Now, we have to require that the superconformal  $R$ -symmetry at the fixed point satisfies the anomaly cancellation condition,<sup>5</sup>

$$0 = \text{Tr}RGG = h_G^\vee + \sum_i \left( \left( \frac{1}{3} - \epsilon_i \right) \text{Tr}_i R_{\mathcal{N}=2} GG + \left( \frac{4}{3} + 2\epsilon_i \right) \text{Tr}_i I_3 GG \right), \quad (2.3)$$

where  $i$  runs through all the  $p_i$  of  $\mathcal{D}_{p_i}(G)$ , which is equivalent to the vanishing condition of the Novikov–Shifman–Vainshtein–Zakharov (NSVZ) exact  $\beta$  function for the gauge coupling. The first  $h_G^\vee$  term in Eq. (2.3) is the contribution from the gauginos, and for the next terms in Eq. (2.3), we utilize the flavor central charge  $k_G$  of the  $\mathcal{D}_p(G)$  theories [24]:

<sup>5</sup>When considering anomalies in 4D theories, each trace term of  $GGG$ ,  $FGG$ ,  $FFG$ , and  $FFF$ , respectively, corresponds to the gauge anomaly, the ABJ anomaly of the flavor  $F$ , a mixed gauge-global anomaly that naturally cancels if  $G$  is non-Abelian, and the 't Hooft anomaly. The gauge anomaly trivially vanishes since the theory has chiral symmetry. Here, we are considering the ABJ anomaly of the  $R$ -symmetry after gauging, which must vanish if it flows in the IR limit to an interacting SCFT. The mixed anomaly for an Abelian gauge theory can be nonvanishing and is responsible for a two-group global symmetry [38].

$$\text{Tr}R_{\mathcal{N}=2}GG = -\frac{1}{2}k_G = -\frac{p-1}{p}h_G^\vee. \quad (2.4)$$

Then we are left with the relation involving the mixing coefficients  $\epsilon_i$ ,

$$h_G^\vee + \sum_i \left( \frac{1}{3} - \epsilon_i \right) \left( -\frac{p_i-1}{p_i} h_G^\vee \right) = 0, \quad (2.5)$$

where the exact values of the  $\epsilon_i$  are fixed by  $a$ -maximization [37]. There are situations where there are more  $U(1)$  flavor symmetries than the ones we considered above (namely, the unbroken  $U(1)$  symmetries inside the  $\mathcal{N} = 2$   $R$ -symmetry). For example, when  $G = SU(N)$  and  $\mathcal{D}_p(G)$  has extra flavor symmetry beyond  $G$ , the  $\mathcal{D}_p(G)$  theory is a Lagrangian quiver gauge theory, and the additional flavor symmetries are the  $U(1)$ s rotating the bifundamental hypermultiplets. As these  $U(1)$ s are baryonic, they do not mix with the  $R$ -symmetry [37].

A novel feature of the glued  $\mathcal{N} = 1$  theories is that they all have their two central charges to be equal,  $a = c$ , for particular combinations of  $G$  and  $p_i$ . It follows from the evaluation of the difference of the central charges, and it does not rely on the knowledge of the exact superconformal  $R$ -symmetry,

$$\begin{aligned} \text{Tr}R &= 16(a - c) \\ &= \dim(G) + \sum_i \left( \left( \frac{1}{3} - \epsilon_i \right) \text{Tr}_i R_{\mathcal{N}=2} + \left( \frac{4}{3} + 2\epsilon_i \right) \text{Tr}_i I_3 \right) \\ &= \dim(G) + \sum_i \left( \frac{1}{3} - \epsilon_i \right) 48(a_i - c_i), \end{aligned} \quad (2.6)$$

where  $a_i$  and  $c_i$  are the central charges of each  $\mathcal{D}_{p_i}(G)$  theory. Here, we used the relation [12],

$$a = \frac{3}{32}(3\text{Tr}R^3 - \text{Tr}R), \quad c = \frac{1}{32}(9\text{Tr}R^3 - 5\text{Tr}R). \quad (2.7)$$

The central charges of the  $\mathcal{D}_{p_i}(G)$  theories with  $p_i$  and  $h_G^\vee$  coprime (which comprises a subset of the  $\mathcal{D}_p(G)$  theories without any extra flavor symmetry beyond  $G$ ) simplify to

$$\begin{aligned} a_i &= \frac{1}{48} \frac{(4p_i - 1)(p_i - 1)}{p_i} \dim(G), \\ c_i &= \frac{1}{12} (p_i - 1) \dim(G). \end{aligned} \quad (2.8)$$

Thus, if we assume that each  $p_i$  is coprime to  $h_G^\vee$ , Eq. (2.6) further implies the difference in the two central charges to be zero:

$$\begin{aligned}
 a - c &= \frac{1}{16} \left( \dim(G) + \sum_i \left( \frac{1}{3} - \epsilon_i \right) \left( -\frac{p_i - 1}{p_i} \dim(G) \right) \right) \\
 &= \frac{\dim(G)}{16h_G^\vee} \text{Tr} RGG = 0. \quad (2.9)
 \end{aligned}$$

This means that for these theories, we have the two central charges  $a$  and  $c$  to be equal and the  $\text{Tr}R$  term, presented in Eq. (2.6), vanishes. We note that this argument is unchanged under a superpotential deformation of the 4D  $\mathcal{N} = 1$  theory and thus, that the  $a = c$  property is preserved under such deformations. This is because the superpotential deformation only puts further constraints on the  $\epsilon_i$  and does not alter the structure of Eq. (2.9), as long as there is no accidental symmetry that mixes with  $R$ . Superpotential deformations of these gaugings will be explored in [22].

Let us make a remark that the condition that  $p$  and  $h_G^\vee$  are coprime is stronger than the condition for the  $\mathcal{D}_p(G)$  theory to have no additional flavor symmetry beyond  $G$ . For example,  $\mathcal{D}_p(SO(2N))$  theory with odd  $p$  has no extra flavor symmetry, but it does not give identical central charges after gauging as it does not necessarily follow that  $\gcd(p, 2N - 2) = 1$ .

### B. Constraints on $p_i$

We demonstrated in Sec. II A that if we consider a quiver gauge theory obtained by the diagonal gauging of the flavor symmetry  $G$  of the collection of Argyres–Douglas theories  $\mathcal{D}_{p_i}(G)$  such that  $\prod_i \gcd(p_i, h_G^\vee) = 1$ , then the  $\mathcal{N} = 1$  superconformal field theories that the quiver may flow to in the infrared have  $a = c$ .

As we are interested in the gauged  $\mathcal{N} = 1$  theories when they flow to superconformal field theories in the infrared, this requires that the gauge coupling for  $G$  is asymptotically free. This puts strong constraints on the possible  $p_i$  for which we can consider the gauging of  $\mathcal{D}_{p_i}(G)$ .

Before we dive into the asymptotic-free condition for the  $\mathcal{N} = 1$  gauging, we revisit the  $\mathcal{N} = 2$  gauging for the theories explored in [14]. For those theories, it was crucial to consider the conformal gauging, as opposed to the asymptotically free gauging. Constructing a 4D  $\mathcal{N} = 2$  SCFT via conformal gauging of  $N$   $\mathcal{D}_{p_i}(G)$  requires that

$$\sum_{i=1}^N \frac{2(p_i - 1)}{p_i} h_G^\vee = 4h_G^\vee \Rightarrow \sum_{i=1}^N \frac{1}{p_i} = N - 2, \quad (2.10)$$

where we have used that the central charge of the flavor symmetry of  $\mathcal{D}_p(G)$  is

$$k_G = \frac{2(p - 1)}{p} h_G^\vee. \quad (2.11)$$

It is straightforward to see that there are only four solutions (for finite values of  $p_i$ ) to the Diophantine Eq. (2.10), as shown in [24]:

$$\begin{aligned}
 (p_1, p_2, p_3, p_4) \\
 = (2, 2, 2, 2), (1, 3, 3, 3), (1, 2, 4, 4), (1, 2, 3, 6), \quad (2.12)
 \end{aligned}$$

where  $\mathcal{D}_1(G)$  is the empty theory.<sup>6</sup> These four solutions are associated to the affine ADE Dynkin diagrams  $\hat{D}_4$ ,  $\hat{E}_6$ ,  $\hat{E}_7$ , and  $\hat{E}_8$ , respectively [14].

A similar analysis can be performed to determine which  $\mathcal{N} = 1$  gaugings of  $\mathcal{D}_{p_i}(G)$  are asymptotically free and thus, which have a chance to flow to an interacting SCFT in the IR. Recalling the expressions for the flavor central charges in Eq. (2.11), we find that the asymptotic-free condition in this case is

$$\sum_{i=1}^N \frac{2(p_i - 1)}{p_i} h_G^\vee < 6h_G^\vee. \quad (2.13)$$

Since  $h_G^\vee$  is nonzero, this inequality (2.13) can be rewritten as a bound for a polynomial in  $p_i$ , as in the following Diophantine-like form:

$$\sum_{i=1}^N \frac{1}{p_i} > N - 3. \quad (2.14)$$

To find all solutions, we need to analyze all possible values of  $p_i \geq 2$  satisfying Eq. (2.14). It is clear that the maximum value of  $N$  for which there exists such solutions is  $N = 5$ , and thus, we consider each solution of Eq. (2.14) as an unordered five-tuple,

$$(p_1, p_2, p_3, p_4, p_5), \quad (2.15)$$

where some of the  $p_i$  may be one.<sup>7</sup> All tuples of  $p_i$  that satisfy the asymptotic freedom condition in Eq. (2.14) are listed in Table I. We write these  $\mathcal{N} = 1$  gaugings via the quivers,

$$\begin{array}{c}
 \mathcal{D}_{p_4}(G) \\
 | \\
 \mathcal{D}_{p_3}(G) - \textcircled{G} - \mathcal{D}_{p_5}(G) \\
 / \quad \backslash \\
 \mathcal{D}_{p_2}(G) \quad \mathcal{D}_{p_1}(G)
 \end{array}, \quad (2.16)$$

where a solid-bordered node represents an  $\mathcal{N} = 1$  vector multiplet, and a solid line between a  $\mathcal{D}_p(G)$  theory and an  $\mathcal{N} = 1$  vector node corresponds to the inclusion of a term,

$$\mathcal{L} \sim \int d^4\theta \mathcal{J}^a \mathcal{V}^a + \dots, \quad (2.17)$$

<sup>6</sup>For convenience, we expressed the solutions with  $i = 1, 2, 3, 4$ .

<sup>7</sup>The  $\mathcal{D}_1(G)$  refers to the empty theory, which can be simply ignored.

TABLE I. All possible tuples of  $p_i$  such that  $\mathcal{N} = 1$  gauging of the common flavor symmetry of the associated  $\mathcal{D}_{p_i}(G)$  leads to an asymptotically free theory. An entry that is left as  $p_i$  indicates that the theory will be asymptotically free for any positive integer  $p_i$ .

$p_1$	$p_2$	$p_3$	$p_4$	$p_5$	$p_1$	$p_2$	$p_3$	$p_4$	$p_5$	$p_1$	$p_2$	$p_3$	$p_4$	$p_5$
1	1	1	1	$p_5$	1	2	3	10	$\leq 14$	1	3	3	3	$p_4$
1	1	1	$p_4$	$p_5$	1	2	3	11	$\leq 13$	1	3	3	4	$\leq 11$
1	1	$p_3$	$p_4$	$p_5$	1	2	4	4	$p_5$	1	3	3	5	$\leq 7$
1	2	2	$p_4$	$p_5$	1	2	4	5	$\leq 19$	1	3	4	4	$\leq 5$
1	2	3	$\leq 6$	$p_5$	1	2	4	6	$\leq 11$	2	2	2	2	$p_5$
1	2	3	7	$\leq 41$	1	2	4	7	$\leq 9$	2	2	2	3	3
1	2	3	8	$\leq 23$	1	2	5	5	$\leq 9$	2	2	2	3	4
1	2	3	9	$\leq 17$	1	2	5	6	$\leq 7$	2	2	2	3	5

in the Lagrangian with  $\mathcal{J}^a$  the flavor supercurrent from the  $\mathcal{D}_p(G)$  SCFT and  $\mathcal{V}^a$  the vector superfield.

We may be further interested in determining the possible tuples of  $p_i$  such that the inequality in Eq. (2.14) is saturated. When this occurs, the one-loop  $\beta$  function of the gauge coupling for  $G$  vanishes directly. This does not generally lead to an interacting conformal field theory unless there is an exactly marginal operator. When we consider saturating the inequality, there is one solution where one has six  $p_i \geq 2$ , and all other solutions have either four or five  $p_i \geq 2$ . All solutions are listed in Table II. It is notable that solutions with six copies of  $\mathcal{D}_2(G)$  possess 16 exactly marginal operators, formed by products of the Coulomb branch operators, and thus, they are interacting SCFTs with  $a = c$ .

Asymptotic freedom of the gauge coupling is a necessary condition for the existence of an interacting superconformal field theory at the infrared fixed point. However, we know that not all theories satisfying the asymptotic freedom condition, given by Eq. (2.13), flow to an interacting SCFT. Henceforth, we have to further check that the theory is consistent. To this end, we perform  $a$ -maximization [37] to determine the superconformal  $R$ -symmetry at the end of the RG flow. We then need to verify that the  $R$ -charges of the operators of the gauged theory under the superconformal  $R$ -symmetry satisfy the unitarity bounds.

If an operator appears to be nonunitary after the  $a$ -maximization, then somewhere along the flow into the infrared, this operator becomes free and is decoupled from the theory [39]. One useful way to take care of the

decoupled operator  $\mathcal{O}$  is to introduce a flipper field  $M$  with a superpotential coupling [2,40,41]:

$$W = M\mathcal{O}. \quad (2.18)$$

In this way, one obtains the superconformal  $R$ -symmetry associated to the interacting sector of the infrared SCFT. This decoupling will violate the assumption made in Sec. II A that we do not have any accidental  $U(1)$  symmetries that can mix with the superconformal  $R$ -symmetry. For this case, the infrared SCFT will not have equal central charges.

We compute the superconformal  $R$ -symmetry via the procedure of  $a$ -maximization. Each  $\mathcal{D}_{p_i}(G)$  theory has a moment map operator  $\mu_i$  and a particular Coulomb branch operator with the smallest conformal dimension  $u_i^0$ . The operator  $u_i^0$  is the bottom component of an  $\mathcal{N} = 2$  chiral multiplet, which splits into two  $\mathcal{N} = 1$  multiplets with bottom components  $u_i^0$  and  $Q^2 u_i^0$ , where the  $Q$  is an  $\mathcal{N} = 2$  supercharge with  $(R_{\mathcal{N}=2}, I_3) = (-1, 1/2)$ . The  $R$ -charges of these operators under the newfound infrared superconformal  $R$ -symmetry are

$$R(\mu_i) = \frac{4}{3} + 2\epsilon_i, \quad R(u_i^0) = \left(\frac{1}{3} - \epsilon_i\right) \frac{2(p_i + 1)}{p_i},$$

$$R(Q^2 u_i^0) = \frac{4p_i + 2}{3p_i} + \frac{2(p_i - 1)}{p_i} \epsilon_i. \quad (2.19)$$

Since we require  $\text{Tr} \mu_i \mu_{j \neq i}$ ,  $u_i^0$ , and  $Q^2 u_i^0$  to be unitary operators of the SCFT at the IR fixed point, we get lower bounds on these  $R$ -charges as

TABLE II. The collections of  $p_i$  that saturate the inequality in Eq. (2.14). For these choices of  $p_i$ , one obtains a theory where the one-loop  $\beta$  function of the gauge coupling vanishes.

$p_1$	$p_2$	$p_3$	$p_4$	$p_5$	$p_6$	$p_1$	$p_2$	$p_3$	$p_4$	$p_5$	$p_6$	$p_1$	$p_2$	$p_3$	$p_4$	$p_5$	$p_6$
1	1	2	3	7	42	1	1	2	4	6	12	1	1	3	4	4	6
1	1	2	3	8	24	1	1	2	4	8	8	1	1	4	4	4	4
1	1	2	3	9	18	1	1	2	5	5	10	1	2	2	3	3	3
1	1	2	3	10	15	1	1	2	6	6	6	1	2	2	2	4	4
1	1	2	3	12	12	1	1	3	3	4	12	1	2	2	2	3	6
1	1	2	4	5	20	1	1	3	3	6	6	2	2	2	2	2	2

$$R(\mu_i) + R(\mu_{j \neq i}) > \frac{2}{3}, \quad R(u_i^0) > \frac{2}{3}, \quad R(Q^2 u_i^0) > \frac{2}{3}. \quad (2.20)$$

When these inequalities are saturated, the theory is still unitary, but the associated operators become free and decouple from the theory. When this occurs, the interacting sector of the resulting SCFT does not have  $a = c$  because decoupled free chirals do not have identical central charges. The second and the third conditions are obvious from the unitarity bound for the scalar ( $\Delta \geq 1$ ), and the first term stems from the fact that the gauge-invariant operators of lowest dimension arise from  $\text{Tr} \mu_i \mu_j$ . In fact,  $\text{Tr} \mu_i^k$  is zero in the chiral ring for all the  $\mathcal{D}_p(G)$  theories with  $(p, h^\vee) = 1$ . This can be understood from the fact that the Higgs branch for the  $\mathcal{D}_p(G)$  theory is given by a nilpotent orbit of  $G$  [42,43]. Any nilpotent element  $X$  in  $G$  should have

$$\text{Tr} X^k = 0. \quad (2.21)$$

Notice that  $\langle \mu \rangle$  parametrize the Higgs branch. However, the operators of the form  $\text{Tr} \mu_i \mu_j$  with  $i \neq j$  are always present. The unitarity constraints in Eq. (2.20) provide the upper and the lower bounds on the mixing parameters  $\epsilon_i$  as

$$-\frac{p_i + 1}{3(p_i - 1)} < \epsilon_i < \frac{1}{3(p_i + 1)}, \quad \epsilon_i + \epsilon_{j \neq i} > -1. \quad (2.22)$$

We will explore how a ‘‘conformal window’’ of possible sets of  $p_i$  can be determined for gaugings of between one and five  $\mathcal{D}_p(G)$  from these bounds on the mixing parameters  $\epsilon_i$ . It turns out that for most of the theories we consider, every  $\epsilon_i$  satisfies

$$-\frac{1}{3} < \epsilon_i < 0, \quad (2.23)$$

which is a sufficient condition for Eq. (2.22) to be satisfied.

Upon diagonally gauging a number of  $\mathcal{D}_p(G)$  theories and flowing into the infrared, we find in the end that an infinite number of 4D  $\mathcal{N} = 1$  SCFTs with identical central charges,  $a = c$ , can be obtained. From Table I, we pick any set of  $(p_1, p_2, p_3, p_4, p_5)$  with either at most two of the  $p_i$  being 1 or else pick  $(1, 1, 1, p_4, p_5)$  subject to the constraint that

$$p_4 \geq 3 \quad \text{and} \quad p_5 \geq 3. \quad (2.24)$$

We then consider the asymptotically free theory obtained by coupling the flavor currents for the  $G$  flavor symmetry of  $\mathcal{D}_{p_i}(G)$ , for each  $p_i$ , to an  $\mathcal{N} = 1$  vector multiplet. When the flavor symmetry  $G$  is such that  $\text{gcd}(p_i, h_G^\vee) = 1$ , for all of the  $p_i$ , then the quiver that we have formed by the diagonal  $\mathcal{N} = 1$  gauging flows to an SCFT in the infrared with  $a = c$ .

### C. Gauging one $\mathcal{D}_p(G)$ theory: No SCFT

The simplest scenario of gauging is when we gauge  $G$  of a single  $\mathcal{D}_p(G)$  theory. The resulting  $\mathcal{N} = 1$  quiver is

$$\mathcal{D}_p(G) \text{ --- } \textcircled{G}. \quad (2.25)$$

There is a unique anomaly-free  $R$ -charge, which is the generator of the superconformal  $R$ -symmetry in the IR:

$$R = R_0 - \left( \frac{2}{3} + \frac{1}{p-1} \right) \mathcal{F} \Rightarrow \epsilon = - \left( \frac{2}{3} + \frac{1}{p-1} \right). \quad (2.26)$$

It is clear that this  $\epsilon$  does not satisfy the unitarity condition, as it is not within the bounds in Eq. (2.22). The  $R$ -charges of the putative operators become

$$R(\mu) = -\frac{2}{p-1}, \quad R(u^0) = 2 + \frac{4}{p-1}, \quad R(Q^2 u_0) = 0. \quad (2.27)$$

Hence, this theory does not flow to an interacting SCFT in the IR. When we have a negative  $R$ -charge operator, it is possible to generate a dynamic superpotential as in the celebrated case of Affleck–Dine–Seiberg (ADS) [44], where it was shown in particular for  $SU(N_c)$  SQCD with  $N_f < N_c$  flavors. Even though the  $R$ -charge for  $\mu$  is negative, all the gauge-invariant operators of the form  $\text{Tr} \mu^k$  are not in the chiral ring; therefore, we do not expect such a runaway dynamical superpotential to be generated. Instead, we find a zero  $R$ -charge operator,  $Q^2 u^0$ , which is often responsible for quantum deformation of the vacuum moduli space as in the case of  $SU(N_c)$  SQCD with  $N_f = N_c$  flavors [45]. See also [46] for a similar phenomenon involving the non-Lagrangian  $T_N$  theory.

### D. Gauging two $\mathcal{D}_p(G)$ theories

In contrast to considering a single  $\mathcal{D}_p(G)$  theory, we can also form theories by gluing more than one  $\mathcal{D}_p(G)$ . We first consider gauging two  $\mathcal{D}_p(G)$  theories together, which leads to a quiver of the following form:

$$\mathcal{D}_{p_1}(G) \text{ --- } \textcircled{G} \text{ --- } \mathcal{D}_{p_2}(G). \quad (2.28)$$

As we determined in Sec. II B, the resulting theory is asymptotically free for arbitrary  $(p_1, p_2)$ , and we will assume without loss of generality that  $p_1 \leq p_2$ . Applying  $a$ -maximization, we discover that the superconformal  $R$ -symmetry is

$$R = R_0 + \epsilon_1 \mathcal{F}_1 + \epsilon_2 \mathcal{F}_2, \quad (2.29)$$

where the two mixing parameters  $\epsilon_1$  and  $\epsilon_2$  are fixed by the values of  $p_i$ :

$$\epsilon_1 = \frac{-p_1^2 + p_2 + p_1\sqrt{1 - p_1 - p_2 + p_1^2 - p_1p_2 + p_2^2}}{3(p_1 - 1)(p_1 - p_2)}, \quad p_1 \geq 3 \quad \text{and} \quad p_2 \geq 3. \quad (2.32)$$

(2.30a)

$$\epsilon_2 = \epsilon_1|_{p_1 \leftrightarrow p_2}. \quad (2.30b)$$

We note that while  $\epsilon_1$  and  $\epsilon_2$  appear to have poles when  $p_1 = p_2 = p$ , we find that it is actually canceled, and

$$\epsilon_1 = \epsilon_2 = \frac{2 + p}{6(1 - p)}. \quad (2.31)$$

The unitarity constraints on  $\epsilon_1$  and  $\epsilon_2$ , as in Eq. (2.22), give rise to constraints on  $p_1$  and  $p_2$  as

$$\frac{3}{2}N_c \leq N_f \leq 3N_c. \quad (2.33)$$

This should not be surprising since there exist abundant gauge theories that flow to interacting SCFTs even though the one-loop  $\beta$  function coefficients lie outside of the region set by the conformal windows for SQCD [47,48].

When the  $p_i$  satisfy the condition in Eq. (2.32), the central charges of the interacting SCFT in the infrared are

$$a = c = -\frac{(p_1p_2 - p_1 - p_2)(A - Q)(B - Q)(A + B + Q)}{48(p_1 - 1)(p_2 - 1)(p_1 - p_2)^2} \dim(G), \quad (2.34)$$

where we defined following quantities for convenience:

$$A = 2p_1 - p_2, \quad B = -p_1 + 2p_2, \quad Q = \sqrt{1 - p_1 - p_2 + p_1^2 - p_1p_2 + p_2^2}. \quad (2.35)$$

They are simplified when  $p_1 = p_2 = p$  to

$$a = c = \frac{9p(p - 2)}{64(p - 1)} \dim(G). \quad (2.36)$$

### E. Gauging three $\mathcal{D}_p(G)$ theories

Similarly to the gauging of two  $\mathcal{D}_p(G)$  theories discussed in Sec. II D, superconformal field theories formed via  $\mathcal{N} = 1$  gauging of three  $\mathcal{D}_p(G)$  are also asymptotically free for arbitrary  $(p_1, p_2, p_3)$ . These quivers have the following form:

$$\begin{array}{c} \mathcal{D}_{p_3}(G) \\ | \\ \mathcal{D}_{p_1}(G) \text{ --- } \textcircled{G} \text{ --- } \mathcal{D}_{p_2}(G) \end{array} \quad (2.37)$$

To check whether such a theory flows to an infrared SCFT with  $a = c$ , we must determine the  $R$ -charge generator of that putative SCFT. It is technically challenging to determine the exact  $R$ -charge analytically via  $a$ -maximization; however, we can study the solutions in particular limits or otherwise numerically.

We first consider the behavior when  $p_2 = p_1$ . Then we find that the mixing coefficients  $\epsilon_1$ ,  $\epsilon_2$ , and  $\epsilon_3$  appearing in the infrared  $R$ -charge are

$$\epsilon_1 = \epsilon_2 = \frac{-4p_3^2 + Q_0(p_1, p_3) + p_1(Q_1(p_1, p_3) + Q_2(p_1, p_3)^{1/2})}{3(p_1 - 1)Q_3(p_1, p_3)}, \quad (2.38a)$$

$$\epsilon_3 = -\frac{p_1(-4p_3^3 + p_1p_3^2 - p_1) + Q_0(p_1, p_3) + p_3(Q_1(p_1, p_3) + 2Q_2(p_1, p_3)^{1/2})}{3(p_3 - 1)Q_3(p_1, p_3)}, \quad (2.38b)$$



where we have defined the polynomials  $Q_j(p_1, p_3)$  as

$$Q_0(p_1, p_3) = -2p_1p_3(p_1 - 1)(p_3 - 1), \quad (2.39a)$$

$$Q_1(p_1, p_3) = p_1^2 - p_1^2p_3 + 4p_3^2, \quad (2.39b)$$

$$\begin{aligned} Q_2(p_1, p_3) &= 9p_1^2(p_1 - 1)^4 + 3(p_3 - p_1)p_1(p_1 - 1)(10p_1^3 - 24p_1^2 + 17p_1 - 4) \\ &\quad + (p_3 - p_1)^2(37p_1^4 - 106p_1^3 + 98p_1^2 - 32p_1 + 4) \\ &\quad + 4(p_3 - p_1)^3(p_1 - 1)(5p_1^2 - 7p_1 + 1) + 4(p_3 - p_1)^4(p_1 - 1)^2, \end{aligned} \quad (2.39c)$$

$$Q_3(p_1, p_3) = -p_1^2 + p_1^2p_3 - 4p_1p_3^2 + 4p_3^2. \quad (2.39d)$$

We see that the mixing coefficients satisfy the unitarity conditions in Eq. (2.22) in all cases. The central charges of the infrared SCFT are then given as

$$a = c = \left( \frac{4(p_1p_3 - p_1 - p_3)Q_4(p_1, p_3)}{3(p_3 - 1)Q_3(p_1, p_3)} + \frac{8Q_2(p_1, p_2)(Q_2(p_1, p_2)^{1/2} - Q_4(p_1, p_3))}{9(p_1 - 1)(p_3 - 1)Q_3(p_1, p_3)^2} \right) \dim(G), \quad (2.40)$$

where we additionally defined the polynomial  $Q_4(p_1, p_3)$  as

$$Q_4(p_1, p_3) = p_1 - 2p_1^2 + 2p_3 - 3p_1p_3 + 2p_1^2p_3 + 2p_3^2 - 2p_1p_3^2. \quad (2.41)$$

We further consider when all  $p_i$  are identical. We find that for such a case where  $p_1 = p_2 = p_3 = p$ , we get a very simple answer for the mixing coefficients:

$$\epsilon_1 = \epsilon_2 = \epsilon_3 = -\frac{1}{3(p-1)}. \quad (2.42)$$

This also simplifies the central charges to

$$a = c = \frac{4p(2p-3)}{3(p-1)} \dim(G). \quad (2.43)$$

We can also study the solutions to the  $a$ -maximization problem analytically in the asymptotic regime where  $p_3 \gg p_1, p_2$ . In this case, the mixing coefficients  $\epsilon_1, \epsilon_2$ , and  $\epsilon_3$  appearing in the infrared  $R$ -charge are

$$\epsilon_1 = \frac{-p_1^2(p_2 - 1)(p_1 + p_2) + p_2Q_1(p_1, p_2) + p_1Q_2(p_1, p_2)^{1/2}}{3(p_1 - 1)(p_1 - p_2)(p_1p_2 - p_1 - p_2)} + \mathcal{O}(p_3^{-1}), \quad (2.44a)$$

$$\epsilon_2 = \epsilon_1|_{1 \leftrightarrow 2}, \quad (2.44b)$$

$$\epsilon_3 = \frac{(p_1^2p_2 + p_1p_2^2 - p_1^2 - p_2^2)Q_1(p_1, p_2) - 2p_1p_2Q_2(p_1, p_2)^{1/2}}{6(p_1 - p_2)^2(p_1p_2 - p_1 - p_2)^2} \cdot \frac{Q_1(p_1, p_2)}{p_3} + \mathcal{O}(p_3^{-2}), \quad (2.44c)$$

where we have defined the polynomials  $Q_1(p_1, p_2)$  and  $Q_2(p_1, p_2)$  as

$$Q_1(p_1, p_2) = 2p_1p_2 - p_1 - p_2, \quad (2.45a)$$

$$\begin{aligned} Q_2(p_1, p_2) &= 4p_2^2(p_2 - 1)^4 + 4(p_1 - p_2)p_2(p_2 - 1)^3(3p_2 - 1) \\ &\quad + (p_1 - p_2)^2(13p_2^4 - 36p_2^3 + 33p_2^2 - 10p_2 + 1) \\ &\quad + (p_1 - p_2)^3(p_2 - 1)(6p_2^2 - 8p_2 + 1) + (p_1 - p_2)^4(p_2 - 1)^2. \end{aligned} \quad (2.45b)$$

These functions exhibit a nice simplification when  $p_1 = p_2 = p$  such that




FIG. 2. Contour plots of  $-\epsilon_1$  and  $-\epsilon_3 \cdot p_3$ . We see that  $\epsilon_1$  lies in the range  $(-\frac{1}{3}, 0)$ , and thus,  $\epsilon_2$  does also. In (b), we show that  $-\epsilon_3 \cdot p_3$  is always positive, and since we are in a large  $p_3$  limit, then we can see that  $\epsilon_3$  must also lie in the range  $(-\frac{1}{3}, 0)$ .

$$Q_1(p, p) = 2p(p-1), \quad Q_2(p_1, p_2) = 4p^2(p-1)^4. \quad (2.46)$$

In this special case, we find a simple expression for two of the mixing coefficients:

$$\epsilon_1 = \epsilon_2 = -\frac{1}{3(p-1)} + \mathcal{O}(p^{-1}), \quad (2.47)$$

where we observe that the first order term is identical to the exact solution for the case of  $p_1 = p_2 = p_3 = p$  given in Eq. (2.42). We can see that the unitarity constraints in Eq. (2.22) are satisfied for all  $p$ .

For general cases where  $p_1 \neq p_2$ , we do a numerical check of the values of the mixing parameters  $\epsilon_i$ . We can see from the contour plots on the  $(p_1, p_2)$ -plane in Fig. 2 that the  $\epsilon_i$  are in the range  $-\frac{1}{3} < \epsilon_i < 0$  when  $p_3$  is large, which is sufficient for them to flow to an SCFT with  $a = c$ . In addition, we checked numerically that the unitarity condition in Eq. (2.22) is satisfied for small values of  $p_3$ , specifically for  $p_3 = 2, 3, 4$ , as can be seen in Fig. 3. In both the small and the large  $p_3$  regime, the unitarity condition is always satisfied in the infrared limit of three  $\mathcal{D}_{p_i}(G)$  theories gauged together. Considering the monotonicity of the  $\epsilon_i$  along  $p_1$  and  $p_2$  in both regimes, we expect they will monotonically increase along with the increasing  $p_3$  and thus, that the unitarity condition is always satisfied. In short, every gluing  $(p_1, p_2, p_3)$  leads to an infrared SCFT with  $a = c$ .

### F. Gauging four $\mathcal{D}_p(G)$ theories

We turn now to the case where we gauge together four  $\mathcal{D}_p(G)$  theories. These  $4D \mathcal{N} = 1$  SCFTs can be written as

the infrared fixed points of the diagonal gauging, which can be depicted as

$$\begin{array}{c} \mathcal{D}_{p_4}(G) \\ | \\ \mathcal{D}_{p_1}(G) - \textcircled{G} - \mathcal{D}_{p_3}(G) \\ | \\ \mathcal{D}_{p_2}(G) \end{array} \quad (2.48)$$

All gaugings that satisfy the asymptotic freedom condition are listed in Table I, and each of these  $(p_1, p_2, p_3, p_4)$  satisfy the unitarity conditions in Eq. (2.22). Each case has been checked numerically. In Fig. 4, we show the contour plots for the  $\epsilon_i$  on the  $(p_3, p_4)$ -plane for the gauging  $(2, 2, p_3, p_4)$ . In Fig. 5, we write the value of  $\epsilon_i$  against the value of  $p_4$  for each of the other gaugings in Table I. We can observe that the unitarity condition is always satisfied for all possible gaugings either explicitly or via the asymptotic behavior when  $p_3$  or  $p_4$  is unbounded. Similar to the case where three  $\mathcal{D}_p(G)$  theories are gauged together, all gaugings of four  $\mathcal{D}_p(G)$  theories lead to interacting superconformal field theories with  $a = c$  in the infrared.

### G. Gauging five $\mathcal{D}_p(G)$ theories

Finally, we turn our attention to the maximal number of  $\mathcal{D}_p(G)$  theories that can be gauged together with an asymptotically free coupling: five  $\mathcal{D}_p(G)$ . There are only four classes of theories with five  $\mathcal{D}_p(G)$  theories glued together in such a way, as can be seen from Table I. These give rise to the quivers,




FIG. 3. Contours plot of  $\epsilon_1$  and  $\epsilon_3$  in the  $(p_1, p_2)$  plane for  $p_3 = 2, 3, 4$ . They all satisfy the unitarity condition in Eq. (2.22).




FIG. 4. Contour plot of  $-\epsilon_1$  and  $-\epsilon_3$  for  $(p_1, p_2, p_3, p_4) = (2, 2, p_3, p_4)$ . Every  $R$ -symmetry mixing coefficient  $\epsilon_i$  satisfies the unitarity condition in Eq. (2.22).

$$\begin{array}{c}
 \mathcal{D}_{p_4}(G) \\
 | \\
 \mathcal{D}_{p_3}(G) \text{ --- } \textcircled{G} \text{ --- } \mathcal{D}_{p_5}(G) \\
 / \quad \backslash \\
 \mathcal{D}_{p_2}(G) \quad \mathcal{D}_{p_1}(G)
 \end{array} \quad (2.49)$$

We can check analytically that the unitarity bound in Eq. (2.22) holds for each case. The  $R$ -symmetry mixing parameters after  $a$ -maximization are found to be

$$\begin{aligned}
 (2, 2, 2, 2, p): \epsilon_{1,2,3,4} &= \frac{2p^2 - 2 - Q(p)^{1/2}}{3(4p^2 - p + 1)}, & \epsilon_5 &= \frac{-4p^3 - 4p^2 + 5p - 1 + 2pQ(p)^{1/2}}{3(p-1)(4p^2 - p + 1)}, \\
 Q(p) &= 4p^4 + 8p^3 - 14p^2 + 3p + 3, & & (2.50a)
 \end{aligned}$$

$$\begin{aligned}
 (2, 2, 2, 3, 3): \epsilon_{1,2,3} &= \frac{1}{147}(163 - 4\sqrt{1731}) \sim -0.0232731, \\
 \epsilon_{4,5} &= \frac{1}{294}(-379 + 9\sqrt{1731}) \sim -0.0154844, & & (2.50b)
 \end{aligned}$$

$$(2, 2, 2, 3, 4): \epsilon_{1,2,3} \sim -0.0119828, \quad \epsilon_4 \sim -0.00798042, \quad \epsilon_5 \sim -0.00597775, \quad (2.50c)$$

$$(2, 2, 2, 3, 5): \epsilon_{1,2,3} \sim -0.00490809, \quad \epsilon_4 \sim -0.00327071, \quad \epsilon_5 \sim -0.00196062. \quad (2.50d)$$

In the first case, it is easy to check that the  $\epsilon_i$  satisfy the unitarity condition in Eq. (2.22) for any value of  $p$ . Thus, all the cases of Eq. (2.50) flow in the infrared to interacting SCFTs with identical central charges. We list the values of those central charges, in each case, in Table III.

### H. Gauging six $\mathcal{D}_2(G)$ theories: conformal gauging

Throughout Sec. II thus far, we mainly focused on asymptotically free ( $\mathcal{N} = 1$ )-gaugings of  $\mathcal{D}_{p_i}(G)$  theories,

and the  $a$ -maximization procedure that must be carried out to determine the superconformal  $R$ -symmetry. In addition to the asymptotically free gaugings, as shown in Table I, there are collections of  $p_i$  that can be combined via conformal gaugings; hence, the theory does not flow. We list the conformal gaugings in Table II. We do not study in detail the conformal gaugings in this paper, as they are not *a priori* guaranteed to have the exactly marginal operators that are required for the theories to be interacting SCFTs with  $a = c$ . However, in this subsection, we




FIG. 5. Plot of  $\epsilon_{1,2,3,4}$  in gluing four  $\mathcal{D}_{p_i}(G)$  theories with all  $(p_1, p_2, p_3, p_4)$  listed in Table I, except first case,  $(2, 2, p_3, p_4)$ , which is numerically studied in Fig. 4. All the coefficients are in the range prescribed in Eq. (2.22).

TABLE III. Central charges  $a = c$  of the infrared SCFTs obtained by gauging together five  $\mathcal{D}_{p_i}(G)$  theories with an  $\mathcal{N} = 1$  vector multiplet.

$(p_1, p_2, p_3, p_4, p_5)$	$a = c$
$(2, 2, 2, 2, p)$	$\left( \frac{p(2p^2 - p + 3)(8p^3 + 28p^2 - 55p + 15) + 2Q(p)^{3/2}}{24(p-1)(4p^2 - p + 1)^2} \right) \dim(G)$
$(2, 2, 2, 3, 3)$	$\left( \frac{3(-19458 + 577\sqrt{1731})}{19208} \right) \dim(G) \sim 0.710368 \dim(G)$
$(2, 2, 2, 3, 4)$	$\sim 0.793031 \dim(G)$
$(2, 2, 2, 3, 5)$	$\sim 0.875633 \dim(G)$

highlight a particular example. We consider the theory formed by  $\mathcal{N} = 1$  gauging of six copies of  $\mathcal{D}_2(G)$ , as depicted in the quiver:

$$\begin{array}{ccccc}
 & \mathcal{D}_2(G) & & \mathcal{D}_2(G) & \\
 & \diagdown & & \diagup & \\
 \mathcal{D}_2(G) & \text{---} & \text{---} & \text{---} & \mathcal{D}_2(G) \\
 & \diagup & & \diagdown & \\
 & \mathcal{D}_2(G) & & \mathcal{D}_2(G) & 
 \end{array} \quad (2.51)$$

Each  $\mathcal{D}_2(G)$  theory has a Coulomb branch operator of lowest dimension  $u_i^0$ , whose dimension is given by

$$\Delta(u_i^0) = \frac{p_i + 1}{p_i} = \frac{3}{2}. \quad (2.52)$$

Henceforth, we see that the gauged theory has 21 marginal operators formed from the product of these Coulomb branch operators:

$$u_i^0 u_j^0. \quad (2.53)$$

Furthermore, there are five  $U(1)$  flavor currents formed from nonanomalous combinations of the six  $U(1)$

symmetries, which is from the  $\mathcal{N} = 2$   $R$ -symmetry of each  $\mathcal{D}_2(G)$  theory. These symmetries are generically broken by the aforementioned marginal operators so that the broken currents combine with them to become marginally irrelevant [16,17]. Therefore, we are left with 16 exactly marginal operators that span a conformal manifold. All other combinations of more than one Coulomb branch operator are irrelevant. Due to the presence of the exactly marginal operators, we can see that this gauging provides nontrivial SCFTs with  $a = c$ . Indeed, we find

$$a = c = \frac{5}{8} \dim(G). \quad (2.54)$$

### III. $\mathcal{N} = 1$ GLUING WITH ADJOINT CHIRALS

In Sec. II, we studied how to glue together several  $\mathcal{D}_p(G)$  SCFTs via  $(\mathcal{N} = 1)$ -gauging of the diagonal flavor symmetry  $G$ . We introduce a new  $G$ -vector multiplet and couple it to the flavor current, as in Eq. (2.17). In this section, we consider a more general gluing, where, in addition to the  $G$ -vector multiplet, we include some number of chiral multiplets charged under  $G$ . Let us introduce  $n$  chiral multiplets in the representation  $\mathbf{R}_{\ell=1, \dots, n}$  of  $G$  and with  $R$ -charge  $R_\ell$ . Then, the condition for the  $R$ -symmetry to be nonanomalous is that

$$\begin{aligned}
 0 &= \text{Tr} RGG = h_G^\vee + \sum_i \left( \left( \frac{1}{3} - \epsilon_i \right) \text{Tr}_i R_{\mathcal{N}=2} GG + \left( \frac{4}{3} + 2\epsilon_i \right) \text{Tr}_i I_3 GG \right) + \sum_{\ell=1}^n (R_\ell - 1) I(\mathbf{R}_\ell) \\
 &= h_G^\vee + \sum_i \left( \frac{1}{3} - \epsilon_i \right) \left( -\frac{p_i - 1}{p_i} h_G^\vee \right) + \sum_{\ell} (R_\ell - 1) I(\mathbf{R}_\ell),
 \end{aligned} \quad (3.1)$$

where  $I(\mathbf{R}_\ell)$  is the Dynkin index of the representation  $\mathbf{R}_\ell$  in  $G$ , and  $\text{Tr}_i$  refers to the anomaly coefficients for the  $\mathcal{D}_{p_i}$  theories that we gauge.

On the other hand, the difference between the two central charges  $(a - c)$  is given by

$$\begin{aligned}
 \text{Tr} R &= 16(a - c) \\
 &= \dim(G) + \sum_i \left( \left( \frac{1}{3} - \epsilon_i \right) \text{Tr}_i R_{\mathcal{N}=2} + \left( \frac{4}{3} + 2\epsilon_i \right) \text{Tr}_i I_3 \right) + \sum_{\ell} (R_\ell - 1) \dim(\mathbf{R}_\ell) \\
 &= \dim(G) + \sum_i \left( \frac{1}{3} - \epsilon_i \right) 48(a_i - c_i) + \sum_{\ell} (R_\ell - 1) \dim(\mathbf{R}_\ell).
 \end{aligned} \quad (3.2)$$

We find that the RGG-anomaly cancellation guarantees  $a = c$ , as before, when

$$\frac{\dim(G)}{h_G^\vee} = \frac{48(a_i - c_i)}{-\frac{p_i - 1}{p_i} h_G^\vee} = \frac{\dim(\mathbf{R}_\ell)}{I(\mathbf{R}_\ell)}, \quad (3.3)$$

for each  $\mathcal{D}_{p_i}(G)$  theory and each chiral matter representation  $\mathbf{R}_\ell$ . Then the RGG-anomaly cancellation directly leads to  $a = c$  via

$$a - c = \frac{\dim(G)}{16h_G^\vee} \text{Tr} RGG = 0, \quad (3.4)$$

as was the case in the previous section. The first equality in Eq. (3.3) holds when  $p_i$  and  $h_G^\vee$  are coprime. The second equality holds if the matter is in the adjoint representation. Thus,  $\mathcal{D}_p(G)$  theories glued together with additional adjoint chiral multiplets may flow to interacting infrared SCFTs with  $a = c$ .

Asymptotic freedom or vanishing of the one-loop  $\beta$  function is required to have an interacting SCFT. It restricts the possible set of gaugings to those satisfying

$$\sum_i \frac{p_i - 1}{p_i} \leq 3 - n_a, \quad (3.5)$$

where  $n_a$  is the number of adjoint chiral multiplets. Clearly, a theory satisfying Eq. (3.5) can have only

$$n_a = 0, 1, 2. \quad (3.6)$$

Let us investigate these cases.<sup>8</sup> We already discussed the cases with  $n_a = 0$  in Sec. II. When  $n_a = 1$ , we have theories which may be connected to the  $\mathcal{N} = 2$  gaugings through superpotential deformations, and we study these further in Secs. III A and III B. Including two adjoint chiral multiplets (i.e.,  $n_a = 2$ ) leads to an asymptotically free theory only when one has a single  $\mathcal{D}_p(G)$  theory being gauged for any value of  $p$ . A theory with vanishing one-loop  $\beta$  function is obtained only when the diagonal of the flavor symmetry of two  $\mathcal{D}_2(G)$  theories is gauged. We explore the  $n_a = 2$  cases in Secs. III C and III D.

### A. $\mathcal{N} = 1$ gluing with one adjoint chiral

In this subsection, we consider several  $\mathcal{D}_p(G)$  theories glued by  $\mathcal{N} = 1$  gauging, together with one chiral multiplet  $\phi$  in the adjoint representation of  $G$ . There are two possibilities: The introduced gauge coupling can either be asymptotically free or it can have vanishing one-loop  $\beta$  function. In the latter case, the matter content is the same as with the theories formed via  $\mathcal{N} = 2$  conformal gauging, and the supersymmetry enhances to  $\mathcal{N} = 2$  if a superpotential term,

$$W = \sum_i \text{Tr} \mu_i \phi, \quad (3.7)$$

is turned on. The explicit sets of  $p_i$  that satisfy the condition written in Eq. (3.5), and are thus asymptotically free or have vanishing one-loop  $\beta$  function, are given by

$$p_i = (p_1), (p_1, p_2), (2, 2, p_3), (2, 3, \leq 6), (2, 4, 4), \\ (3, 3, 3), (2, 2, 2, 2), \quad (3.8)$$

where the  $p_i$  are given in ascending order as before, and when we write a  $p_i$  in the tuple, then any positive integer  $\geq 2$  satisfies Eq. (3.5).

The first case to consider is when we gauge a single  $\mathcal{D}_p(G)$  together with one additional chiral multiplet. In this case, the  $R$ -charge of the adjoint chiral multiplet  $\phi$  is fixed by the cancellation of the  $R$ -gauge-gauge anomaly:

$$R_\phi = \left(\frac{1}{3} - \epsilon\right) \left(\frac{p-1}{p}\right). \quad (3.9)$$

Maximizing  $a$  with respect to  $\epsilon$  leads to

$$\epsilon = \frac{1 - 2p^2 + p\sqrt{4p^2 - 6p + 3}}{6p^2 - 9p + 3}. \quad (3.10)$$

Thus, at the infrared fixed point, we have the following  $R$ -charges of the fields:

$$\frac{1}{9}(5 - \sqrt{7}) \leq R_\phi = \frac{4p - 3 - \sqrt{4p^2 - 6p + 3}}{6p - 3} < \frac{1}{3}, \\ \frac{2}{9}(2\sqrt{7} - 1) \leq R_\mu = \frac{4p^2 - 12p + 6 + 2p\sqrt{4p^2 - 6p + 3}}{6p^2 - 9p + 3} < \frac{4}{3}. \quad (3.11)$$

We can see that  $\text{Tr} \phi^2$  violates the unitarity bound, which signals existence of an accidental  $U(1)$  flavor symmetry that acts on the decoupled free operator  $\text{Tr} \phi^2$  [39]. This invalidates our  $a$ -maximization analysis and the previous argument in Sec. II A that the interacting SCFT has  $a = c$  does not apply.

One convenient way to remove such a decoupled operator is by adding a flipper-field  $M$  and the corresponding superpotential term [2,40,41],

$$\Delta W = M \text{Tr} \phi^2. \quad (3.12)$$

This introduces a  $U(1)$  flavor symmetry, under which, only the operator that gets decoupled is charged, which subsequently mixes with the superconformal  $R$ -symmetry during the flow into the infrared. After removing the  $\text{Tr} \phi^2$  operator via flipping and flowing into the infrared, one finds that the superconformal  $R$ -charges in the interacting sector are such that

<sup>8</sup>There is also a theory with  $n_a = 3$ , where all  $p_i = 1$ . This is simply the Lagrangian gauge theory with three adjoint chiral multiplets, which sits on the conformal manifold containing  $\mathcal{N} = 4$  super Yang–Mills. For obvious reasons, we will not discuss this case further.

$$\begin{aligned}
\frac{1}{48}(27 - \sqrt{217}) \leq R_\phi &= \frac{8p^2 - p - 3 - \sqrt{16p^4 - 15p^2 + 6p + 9}}{12p^2} < \frac{1}{3}, \\
\frac{1}{12}(\sqrt{217} - 3) \leq R_\mu &= \frac{4p^2 - 11p + 3 + \sqrt{16p^4 - 15p^2 + 6p + 9}}{6p(p-1)} < \frac{4}{3}, \\
R_M &= 2 - 2R_\phi.
\end{aligned} \tag{3.13}$$

Thus, it is straightforward to determine that the two central charges are given by

$$a = a_0 \dim(G) - a_1, \quad c = c_0 \dim(G) - c_1, \tag{3.14}$$

where the coefficients of the  $\dim(G)$  terms in both central charges are equal:  $a_0 = c_0$ . More specifically, the  $p$ -dependent constants appearing in Eq. (3.14) are

$$\begin{aligned}
a_0 = c_0 &= \frac{(p+1)(2p-3)(32p^6 + 8p^5 + 6p^4 - 19p^3 + 11p^2 + 3p - 9)}{1536(p-1)p^6} \\
&+ \frac{(p+1)Q(p)(16p^5 - 20p^4 + 12p^3 + 3p^2 - 12p + 9)}{1536(p-1)p^6},
\end{aligned} \tag{3.15a}$$

$$a_1 = \frac{(p+1)^2(2p-3)(10p^3 - 11p^2 - 3p + 9)}{192p^6} - \frac{(4p^4 - 3p^3 - 12p^2 + 6p + 9)Q(p)}{192p^6}, \tag{3.15b}$$

$$c_1 = \frac{(p+1)(2p-3)(8p^4 - p^3 - 14p^2 + 6p + 9)}{192p^6} - \frac{(p-3)(2p^3 + 3p^2 - 3p - 3)Q(p)}{192p^6}, \tag{3.15c}$$

where we have again, defined a useful function,

$$Q(p) = \sqrt{16p^4 - 15p^2 + 6p + 9}. \tag{3.15d}$$

Thus, (the interacting sector of) the infrared SCFT obtained from gauging a single  $\mathcal{D}_p(G)$  theory, together with an additional chiral multiplet in the adjoint representation of  $G$ , does not have  $a = c$ .

We now study the remaining combinations of  $\{p_i\}$  in Eq. (3.8) that can be gauged together, with an extra adjoint chiral, in such a way that the gauge coupling is either asymptotically free or conformal. We find that, in all cases in Eq. (3.8) where we gauge together more than one  $\mathcal{D}_p(G)$ , if each  $p_i$  satisfies

$$\gcd(p_i, h_G^\vee) = 1, \tag{3.16}$$

then we obtain an infrared SCFT with  $a = c$ .

Similarly to the discussion in Sec. II, analytically solving the coupled quadratic equations to obtain the mixing coefficients  $\epsilon_i$  that maximize  $a$  is technically challenging. Instead, we numerically explore the  $p_i$  dependence of the solutions to the  $a$ -maximization condition. In Fig. 7, we numerically study the values of  $\epsilon_1, \epsilon_2$ , and  $R_\phi$  of the gauged theories with  $(p_1, p_2)$ , for generic  $p_1$  and  $p_2$ . We find that

for all values of  $p_1$  and  $p_2$ , the unitarity conditions on the operators are satisfied, and hence, the theory flows to an interacting SCFT with  $a = c$  in the infrared. Similarly, Fig. 6 shows the numerical plots for  $\epsilon_1 = \epsilon_2, \epsilon_3$  and  $R_\phi$  of the theories corresponding to  $(2, 2, p_3)$ , with a generic  $p_3$ . Here, the asymptotic behavior in the large  $p_3$  limit demonstrates that the unitarity bounds will be satisfied for all values of  $p_3$ . Finally, in Table IV, we write the  $\epsilon_i$  and  $R_\phi$  for the gaugings for the remaining cases in Eq. (3.8). From this numerical analysis, we successfully observe that all  $\mathcal{N} = 1$  gaugings with more than one  $\mathcal{D}_p(G)$ , such that  $\gcd(p_i, h_G^\vee) = 1$ , and with an additional adjoint chiral multiplet, flow to superconformal field theories with  $a = c$  in the infrared.

We note that any set of  $p_i$  that saturates the inequality in Eq. (3.5) has

$$\epsilon_i = 0, \quad R_\phi = \frac{2}{3}. \tag{3.17}$$

This reflects the fact that these theories have vanishing one-loop  $\beta$  function, and thus, the theory does not flow.






FIG. 6. Plots of  $\epsilon_1, \epsilon_3$  and  $R_\phi$  of  $(2, 2, p_3)$  theories glued by  $\mathcal{N} = 1$  gauging with one adjoint chiral multiplet. We can see that  $\epsilon_1 = \epsilon_2$  and  $\epsilon_3$  lie in the range  $(-\frac{1}{3}, 0)$ . Furthermore,  $R_\phi$  is always larger than  $1/3$ . Thus, we conclude that for all values of  $p_3$ , the gauged theory flows to an SCFT in the infrared with  $a = c$ .

The  $\mathcal{N} = 2$  superpotential in Eq. (3.7) is marginal in these cases, and so the  $\hat{\Gamma}(G)$  theories, which are studied in [14], and the theories with the same gauged  $\mathcal{D}_p(G)$  and the same matter content but without the superpotential term in Eq. (3.7), are connected through a conformal manifold. This conformal manifold is parametrized by exactly

marginal operators, which can be determined by enumerating marginal operators and symmetries [16,17]. The marginal operators include

$$\text{Tr}\phi^3, \quad \text{Tr}\phi\mu_i. \tag{3.18}$$




FIG. 7. Contour plots of  $-\epsilon_1$  and  $R_\phi$  of  $\mathcal{D}_{p_1}(G)$  and  $\mathcal{D}_{p_2}(G)$  theories glued by  $\mathcal{N} = 1$  gauging with one adjoint chiral multiplet.  $\epsilon_1$  lies in the range  $(-\frac{1}{3}, 0)$ , as does  $\epsilon_2$ , and  $R_\phi$  is always larger than  $1/3$ ; thus, any such gauging satisfies the unitarity bound.

TABLE IV. In this table, we write the values for the  $\epsilon_i$  and the  $R$ -charge of the adjoint chiral multiplet,  $R_\phi$ , after gauging together the  $\mathcal{D}_{p_i}(G)$  with one additional adjoint chiral multiplet. We can see that the values of these parameters, after doing the  $a$ -maximization to find the superconformal  $R$ -symmetry, are such that we obtain an interacting SCFT with  $a = c$  in the infrared.

$\{p_i\}$	$\epsilon_1$	$\epsilon_2$	$\epsilon_3$	$\epsilon_4$	$R_\phi$
(1,2,3,3)	...	-0.0108332	-0.00721553	-0.00721553	0.626148
(1,2,3,4)	...	-0.00535965	-0.00357149	-0.00267712	0.645958
(1,2,3,5)	...	-0.00213347	-0.00142206	-0.00085290	0.658253
(1,2,3,6), (1,2,4,4), (1,3,3,3)	...	0	0	0	2/3
(2,2,2,2)	0	0	0	0	2/3

A number of these marginal operators become marginally irrelevant after combining with the generators of the Abelian flavor symmetry; however, we find that there is always at least one exactly marginal operator that in fact corresponds to the  $\mathcal{N} = 2$  gauge coupling upon suitable normalization. For some choices of  $\{p_i\}$ , one can have additional marginal operators formed out of the products of Coulomb branch operators. We will revisit the study of marginal operators in [21,22].

### B. Mass deformations of $\mathcal{N} = 2$ $\hat{\Gamma}(G)$ theory

In this section, we explore the connection to the mass deformations of the 4D  $\mathcal{N} = 2$  SCFTs  $\hat{\Gamma}(G)$  considered in [14]. These  $\hat{\Gamma}(G)$  theories are associated to a pair of ADE groups,  $\Gamma$  and  $G$ . When  $\Gamma = D_4, E_6, E_7, E_8$ , these theories correspond to an  $\mathcal{N} = 2$  diagonal gauging of the flavor symmetry of a collection of  $\mathcal{D}_{p_i}(G)$  with, respectively,

$$(p_1, p_2, p_3, p_4) = (2, 2, 2, 2), (1, 3, 3, 3), (1, 2, 4, 4), (1, 2, 3, 6). \quad (3.19)$$

These are precisely the same collections of  $p_i$  as in Eq. (3.8) that have vanishing one-loop  $\beta$  function. When the gauging further satisfies  $\gcd(p_i, h_G^\vee) = 1$  for all  $p_i$ , then the resulting  $\mathcal{N} = 2$  SCFTs have  $a = c$ .

It is clear that the  $\mathcal{N} = 2$  gaugings and the  $\mathcal{N} = 1$  gaugings together with one additional adjoint chiral multiplet are distinct theories; the former has the additional superpotential term,

$$W = \sum_i \text{Tr} \mu_i \phi, \quad (3.20)$$

from the  $\mathcal{N} = 2$  vector multiplet, and where  $\phi$  is the adjoint-valued scalar field inside that vector multiplet. From the perspective of the  $(\mathcal{N} = 1)$ -gauged theory, the terms in this superpotential are marginal operators that contribute to an  $\mathcal{N} = 1$  conformal manifold on which the  $\hat{\Gamma}(G)$  theory sits at a special point. Both theories have a relevant operator that gives a mass to either  $\phi$  in the  $\mathcal{N} = 2$  vector multiplet or the additional adjoint chiral  $\phi$ , respectively. It is natural to ask whether the two theories obtained

via these two mass deformations, which integrate out the adjoint chirals, lead to the same infrared SCFTs.

First, we introduce a mass term for the adjoint  $\mathcal{N} = 1$  chiral multiplet  $\phi$  inside of the  $\mathcal{N} = 2$  vector multiplet,

$$W = \text{Tr} \phi^2. \quad (3.21)$$

After adding such a mass term and flowing into the infrared, we obtain an  $\mathcal{N} = 1$  theory, where  $\phi$  is integrated out. The  $\mathcal{N} = 1$  deformations of  $\mathcal{N} = 2$  theories through a mass term were studied in general in [15]. The infrared  $U(1)_R$  charge after the mass deformation is fixed to be

$$R_{\text{IR}} = \frac{1}{2} R_{\mathcal{N}=2} + I_3, \quad (3.22)$$

solely by the mass term in the superpotential. The central charges of the infrared  $\mathcal{N} = 1$  theory thus differ by a factor of 27/32 from the central charges of the original  $\mathcal{N} = 2$  theory.

On the other hand, the infrared  $R$ -symmetry, shown in Eq. (2.2), of the  $(\mathcal{N} = 1)$ -gauged theory satisfies the  $R$ -gauge-gauge anomaly-free condition,

$$1 + \sum_i \left( \frac{1}{3} - \epsilon_i \right) \left( -\frac{p_i - 1}{p_i} \right) = 0. \quad (3.23)$$

First, let us consider the cases where all of the  $p_i$  are identical: where they are either (2,2,2,2) or (3,3,3). Each of the  $\epsilon_i$  are then the same and the anomaly-free condition fixes that

$$\epsilon_i = \frac{1}{3} - \frac{p}{4(p-1)} \Big|_{p=2} = -\frac{1}{6} \quad \text{or} \quad \epsilon_i = \frac{1}{3} - \frac{p}{3(p-1)} \Big|_{p=3} = -\frac{1}{6}, \quad (3.24)$$

for the  $\hat{D}_4$  and  $\hat{E}_6$  cases, respectively. The operators  $\text{Tr} \mu_i \mu_j$  are exactly marginal and span a nontrivial conformal manifold for each of these theories.<sup>9</sup> Going to the point

<sup>9</sup>There always exists at least one marginal operator  $\text{Tr} \mu_i \mu_j$ ; chiral ring relations may reduce the dimension of the conformal manifold from the naive expectation.

on the conformal manifold where the superpotential deformation is

$$\Delta W = \text{Tr} \left( \sum_i \mu_i \right)^2, \quad (3.25)$$

then we observe the mass-deformed  $\hat{D}_4(G)$  or  $\hat{E}_6(G)$  theories.

In general, when not all of the  $p_i$  are identical, the  $\mathcal{N} = 1$  gauging and the mass-deformation after  $\mathcal{N} = 2$  gauging yield different SCFTs with different central charges. For the ( $\mathcal{N} = 1$ )-gauging when  $(p_1, p_2, p_3) = (2, 4, 4)$ , we find that

$$\epsilon_1 = \frac{1}{3}(\sqrt{10} - 4), \quad \epsilon_2 = \epsilon_3 = \frac{1}{9}(2 - \sqrt{10}), \quad (3.26)$$

and thus,

$$a = \frac{1}{144}(29 + 20\sqrt{10}) \dim(G). \quad (3.27)$$

We can see that the operators,

$$\text{Tr}\mu_1\mu_2, \quad \text{Tr}\mu_1\mu_3, \quad (3.28)$$

are relevant operators of this SCFT, and further, there are no exactly marginal operators. The other quadratic trace of the moment maps,  $\text{Tr}\mu_2\mu_3$ , is an irrelevant operator.<sup>10</sup> Let us now consider further deformations by these two relevant operators, sequentially. The resulting infrared SCFT has

$$\epsilon_1 = \epsilon_2 = \epsilon_3 = -\frac{1}{6}, \quad (3.29)$$

and the central charge  $a$  is

$$a = \frac{81}{128} \dim(G) = \frac{27}{32} a(\hat{E}_7(G)). \quad (3.30)$$

We can see that this theory has the same central charge as the mass deformation of the  $\hat{E}_7(G)$  theory. This is as we expect because there are three marginal operators  $\text{Tr}\mu_i\mu_{j\neq i}$ , and only one of them combines with the flavor current multiplet to become marginally irrelevant; in turn, we are left with two remaining [16,17]. These leftover marginal operators will span an at least two-dimensional conformal manifold that connects the two theories.

A similar analysis can be done for the ( $\mathcal{N} = 1$ )-gauging of the  $\mathcal{D}_{p_i}(G)$  theories with  $(p_1, p_2, p_3) = (2, 3, 6)$ , and we obtain irrational  $R$ -charge mixing coefficients,

<sup>10</sup>The operators of the form  $\text{Tr}\mu_i^2$  are projected out by the chiral ring relations of each individual  $\mathcal{D}_p(G)$ . Only the mixed operators  $\text{Tr}\mu_i\mu_{j\neq i}$  survive, as we have discussed near Eq. (2.21).

$$\epsilon_1 \sim -0.284524, \quad \epsilon_2 \sim -0.182519, \quad \epsilon_3 \sim -0.0832703, \quad (3.31)$$

and corresponding central charge,

$$a \sim 0.722376 \dim(G). \quad (3.32)$$

At the same time, the infrared limit of the mass-deformed  $\hat{E}_8(G)$  has its  $a$  central charge by

$$a = \frac{27}{32} a(\hat{E}_8(G)) = \frac{45}{64} \dim(G) \sim 0.703125 \dim(G), \quad (3.33)$$

which is smaller than that of the ( $\mathcal{N} = 1$ )-gauged theory without superpotential. It agrees with the fact that each moment map operator  $\mu_i$  has its  $R$ -charge given by

$$R(\mu_i) = \frac{4}{3} + 2\epsilon_i. \quad (3.34)$$

We can see from the mixing coefficients in Eq. (3.31) that  $R(\mu_1)$  and  $R(\mu_2)$  are both strictly less than one. We find that

$$\text{Tr}\mu_1\mu_2, \quad \text{Tr}\mu_1\mu_3 \quad (3.35)$$

are relevant operators. We can turn on a superpotential deformation by either of these operators to trigger a renormalization group flow to an infrared fixed point. When we choose  $\text{Tr}\mu_1\mu_2$ , the fixed point is the mass-deformed  $\hat{E}_8(G)$  theory; however, if we turns on  $\text{Tr}\mu_1\mu_3$ , then the IR SCFT after the flow still has  $\text{Tr}\mu_1\mu_2$  as a relevant operator. Subsequently triggering a flow by this operator leads, again, to the mass-deformed  $\hat{E}_8(G)$  theory.

This analysis explains why the ( $\mathcal{N} = 1$ )-gauged theory has a central charge  $a$  that is at least as large as the central charge of the mass-deformed  $\hat{\Gamma}(G)$  theory; it is required by the  $a$  theorem [1].

### C. $\mathcal{N} = 1$ theory with two adjoint chirals: A Lagrangian model

Thus far, all the  $\mathcal{N} = 1$  SCFTs with  $a = c$  that we obtain are rather exotic and non-Lagrangian; all of them involve the Argyres–Douglas theories  $\mathcal{D}_p(G)$ . In fact, there is a simple Lagrangian gauge theory with  $a = c$ : consider a gauge theory with gauge group  $G$  and two adjoint chiral multiplets. We can consider this theory as a special case where we gauge *zero*  $\mathcal{D}_p(G)$  theories, together with two adjoint chiral multiplets  $\phi_1$  and  $\phi_2$ . We can see that this configuration is asymptotically free from Eq. (3.5). From the anomaly-free condition for the  $R$ -symmetry, we get

$$\text{Tr}RGG = 0 \Leftrightarrow h_G^\vee(1 + 2(R_\phi - 1)) = 0, \quad (3.36)$$

where we use the symmetry to write  $R_{\phi_1} = R_{\phi_2} = R_\phi$ . The anomaly cancellation enforces that the  $R$ -charges for the two adjoint chiral multiplets are

$$R_{\phi_1} = R_{\phi_2} = \frac{1}{2}. \quad (3.37)$$

Therefore, we find

$$16(a-c) = \text{Tr}R = \left(1 + 2\left(\frac{1}{2} - 1\right)\right) \dim(G) = 0, \quad (3.38)$$

and thus, the theory realizes  $a = c$ . The theory has an  $SU(2)$  flavor symmetry rotating the two adjoint chiral multiplets, and the central charges are given by

$$a = c = \frac{9}{32} \left(1^3 + 2\left(\frac{1}{2} - 1\right)^3\right) \dim(G) = \frac{27}{128} \dim(G). \quad (3.39)$$

This theory also belongs to the conformal manifold of the theory obtained starting from  $\mathcal{N} = 4$  super Yang–Mills with gauge group  $G$  and triggering an RG-flow by adding a mass term for one of the three adjoint chiral multiplets inside the  $\mathcal{N} = 4$  vector multiplet. As expected from Eq. (1.12), the central charges of the infrared  $\mathcal{N} = 1$  theory are 27/32 times the central charges of the  $\mathcal{N} = 4$  theory [15].

#### D. $\mathcal{N} = 1$ gluing with two adjoint chirals

The sets of  $\mathcal{D}_p(G)$  theories that can be gauged together when we include two adjoint chiral multiplets on the gauge node are highly restricted, as we can see from Eq. (3.5). In the simplest case, we consider a single  $\mathcal{D}_p(G)$  theory for any choice of  $p$ . Then we obtain the asymptotically free theory given by

$$\mathcal{D}_p(G) \text{ --- } \left( \text{---} \bigcirc \text{---} \right) \text{---} \quad (3.40)$$

A single  $\mathcal{D}_p(G)$  theory gauged with two adjoint chiral multiplets attached has its infrared  $R$ -charge given by

$$R = R_0 + \epsilon \mathcal{F}, \quad (3.41)$$

where  $\epsilon$  and the  $R$ -charge of adjoint chiral multiplets  $\phi_1$  and  $\phi_2$  are

$$\epsilon = \frac{-8p^3 - 2p^2 + p + 1 + 2p\sqrt{16p^4 + 8p^3 - 11p^2 + 3}}{3(8p^3 - 7p^2 - 2p + 1)},$$

$$R_{\phi_1} = R_{\phi_2} = \frac{20p^2 - p - 3 - \sqrt{16p^4 + 8p^3 - 11p^2 + 3}}{3(8p^2 + p - 1)}. \quad (3.42)$$

It is straightforward to check that the operators satisfy the unitarity conditions for any value of  $p$ , and thus, each theory flows in the infrared to an interacting SCFT with  $a = c$ , if  $\text{gcd}(p, h_G^\vee) = 1$ . The resulting SCFTs have a host of relevant operators,

$$\begin{aligned} & \text{Tr}\phi_1^2, \quad \text{Tr}\phi_1\phi_2, \quad \text{Tr}\phi_2^2, \quad \text{Tr}\phi_1^3, \quad \text{Tr}\phi_1^2\phi_2, \\ & \text{Tr}\phi_1\phi_2^2, \quad \text{Tr}\phi_2^3, \quad \text{Tr}\mu\phi_1, \quad \text{Tr}\mu\phi_2. \end{aligned} \quad (3.43)$$

Each of these operators provides a superpotential deformation that triggers a renormalization group flow to a new infrared SCFT. Let us note that when  $G = SU(2)$ , the cubic operators in Eq. (3.43) are absent. The landscape charted by superpotential deformations involving these relevant operators is one of the subjects of study in [22].

The only other possibility for gauging together  $\mathcal{D}_p(G)$  with two adjoint chiral multiplets is a theory with vanishing one-loop  $\beta$  function that is obtained by gauging two  $\mathcal{D}_2(G)$  theories. The resulting theory is of the form,

$$\mathcal{D}_2(G) \text{ --- } \left( \text{---} \bigcirc \text{---} \right) \text{---} \mathcal{D}_2(G) \quad (3.44)$$

As it is discussed in Sec. II, the gaugings that saturate the inequality in Eq. (2.13) do not necessarily lead to an interacting superconformal field theory, as they may not have any exactly marginal operators.<sup>11</sup> However, when two  $\mathcal{D}_2(G)$  theories are glued together with two adjoint chiral multiplets, we expect a nontrivial SCFT as there are now marginal operators built out of the adjoint chiral multiplets. There are eight marginal operators:

$$\text{Tr}\mu_i\phi_1, \quad \text{Tr}\mu_i\phi_2, \quad \text{Tr}\phi_1^3, \quad \text{Tr}\phi_1^2\phi_2, \quad \text{Tr}\phi_1\phi_2^2, \quad \text{Tr}\phi_2^3. \quad (3.45)$$

For group-theoretic reasons, a number of these operators may not exist; for example, for  $G = SU(2)$ , the four operators that are cubic in  $\phi_i$  are not present due to the absence of a cubic Casimir. Since  $\text{gcd}(2, h_G^\vee) = 1$  is required to obtain a theory with identical central charges  $a = c$ , it is necessary to have  $G = SU(2n+1)$  to ensure  $a = c$ . Thus, the cubic marginal operators are present in the theories with  $a = c$ . Among the eight marginal operators, at most, five of them may become marginally irrelevant as they combine with the generators of the  $SU(2) \times U(1)^2$  flavor symmetry. The remaining operators are exactly marginal and contribute to the conformal manifold with dimension at least three.<sup>12</sup>

<sup>11</sup>While such conformal gaugings may not necessarily admit any exactly marginal operators, we see in [21] that each gauging appearing in Table II does.

<sup>12</sup>The marginal operators belonging to the  $\mathcal{D}_p(G)$  theory before gauging will also contribute to the conformal manifold.

**IV. BEYOND  $a = c$ : INCLUDING CONFORMAL MATTER**

In the current paper, we mainly focused on  $\mathcal{N} = 1$  SCFTs with  $a = c$ , but let us discuss a natural generalization of our setup that may have  $a \neq c$ . In [14], the authors discussed 4D  $\mathcal{N} = 2$  SCFTs called  $\hat{\Gamma}(G)$ , for all choices of ADE Lie algebra  $\Gamma$ . These theories were constructed by taking a collection of Argyres–Douglas  $\mathcal{D}_{p_i}(G)$  theories and a collection of  $(G, G)$  conformal matter theories<sup>13</sup> and performing  $\mathcal{N} = 2$  conformal gaugings of all of the  $G$  flavor symmetries.<sup>14</sup> In [14], it was found that the classification of all possible such gaugings is equivalent to the ADE classification problem, and thus, the solutions are labeled by an ADE algebra  $\Gamma$ . For  $\Gamma = D_4, E_6, E_7, E_8$ , this gauging involved the introduction of a single gauge node, and no copies of the conformal matter theory; however, for  $\Gamma = A_{n-1 \geq 0}$  and  $\Gamma = D_{n \geq 5}$ , the gauging involved adding either of ring of  $n$  gauge nodes or a linear chain of  $(n - 3)$  gauge nodes, respectively.

In this paper, we have focused on gauging together a collection of  $\mathcal{D}_{p_i}(G)$ , possibly with additional adjoint chiral multiplets, via a *single*  $\mathcal{N} = 1$  vector multiplet. We study this particular setup because we desire to explore 4D  $\mathcal{N} = 1$  SCFTs with identical central charges,  $a = c$ . In the setup of  $\hat{\Gamma}(G)$  theories, it is only in the cases where  $\Gamma = D_4, E_6, E_7, E_8$ , and with specific values of  $G$ , that we obtain 4D  $\mathcal{N} = 2$  SCFTs with  $a = c$ . In a similar vein, we would expect that any  $\mathcal{N} = 1$  gauging involving multiple gauge nodes would create a deviation from  $a = c$  due to the included conformal matter theories. However, if we are willing to chart the landscape beyond  $a = c$ , there is no *a priori* problem with considering such a class of  $\mathcal{N} = 1$  SCFTs, obtained from  $\mathcal{N} = 1$  gaugings of a collection of  $\mathcal{D}_{p_i}(G)$  theories and a collection of  $(G \times G)$  conformal matter theories.

A natural starting point for studying  $\mathcal{N} = 1$  gaugings involving both Argyres–Douglas and conformal matter theories could be to consider mass deformations associated to the adjoint chiral multiplets in the  $\mathcal{N} = 2$  theories  $\hat{A}_{n-1 \geq 1}(G)$  and  $\hat{D}_{n \geq 5}(G)$ . This would give a interesting collection of theories with IR central charge being 27/32 of the UV central charges [15]. Here, we also would like to remind the reader that a mass deformation of chiral adjoint in a vector multiplet of  $\mathcal{N} = 2$  theory is not equivalent to the  $\mathcal{N} = 1$  gauging due to the superpotential generated by the mass term. We can consider various configurations

<sup>13</sup>These theories are sometimes known as the generalized  $(G \times G)$  bifundamental. They are obtained by compactifying the relevant six-dimensional minimal conformal matter theory [49–51] on a torus. Some  $\mathcal{N} = 1$  gaugings of  $\mathcal{N} = 2$  conformal matter have been explored in [52].

<sup>14</sup>The resulting theories can have residual flavor symmetries if the  $\mathcal{D}_{p_i}(G)$  or conformal matter theories have a larger flavor symmetry than  $G$  or  $(G \times G)$ , respectively.

TABLE V. We define  $\mathcal{C}_n$  as the set of  $\{p_i\}$  satisfying the inequality in Eq. (4.1). Here, we list all the elements inside  $\mathcal{C}_1$  and  $\mathcal{C}_2$ . These are identical to the collections of  $p_i$  found for gauging together with, respectively, one and two adjoint chiral multiplets in Sec. III. The conformal gaugings saturate the inequality while the asymptotically free gaugings do not.

Name	$\{p_i\}$	
	Asymptotically free gauging	Conformal gauging
$\mathcal{C}_1$	$\{p \geq 2\}$ $\{2, 2, p \geq 2\}$	$\{p_1 \geq 2, p_2 \geq 2\}$ $\{2, 3, 4\}$ $\{2, 3, 5\}$ $\{2, 4, 4\}$ $\{3, 3, 3\}$ $\{2, 3, 6\}$
$\mathcal{C}_2$	$\{p \geq 2\}$	$\{2, 2\}$

corresponding to different choices of  $\mathcal{N} = 1$  and  $\mathcal{N} = 2$  gaugings for the multiple gauge nodes as in the case of  $\mathcal{N} = 1$  class  $\mathcal{S}$  theories [18,20].

The number of options for  $\mathcal{N} = 1$  gaugings grows rapidly with the number of conformal matter theories involved. We introduce some simplifying notation: Define  $\mathcal{C}_n$  as all possible sets of  $p_i$  that solve the inequality

$$\mathcal{C}_n : \sum_i \frac{p_i - 1}{p_i} \leq 3 - n. \tag{4.1}$$

This is just the condition in Eq. (3.5) that dictates which sets of  $\mathcal{D}_{p_i}(G)$  can be  $(\mathcal{N} = 1)$ -gauged together with  $n$  additional adjoint chiral multiplets. The sets of  $p_i$  that belong to  $\mathcal{C}_0$  are listed in Tables I and II. For  $\mathcal{C}_1$  and  $\mathcal{C}_2$ , which will be of relevance in this section, we summarize which sets of  $p_i$  belong to them in Table V. We write  $\mathcal{C}_n$  in a quiver, connected to a gauge node, as a shorthand for all of the quivers involving all sets  $\{p_i\}$  inside  $\mathcal{C}_n$ . For example, the quiver,

$$\mathcal{C}_2 \text{ --- } \textcircled{G}, \tag{4.2}$$

is used to succinctly refer to the two quivers,

$$\mathcal{D}_p(G) \text{ --- } \textcircled{G}, \quad \mathcal{D}_2(G) \text{ --- } \textcircled{G} \text{ --- } \mathcal{D}_2(G). \tag{4.3}$$

Suppose that we wish to take the diagonal gauging of  $n$  flavor symmetries  $G$  coming from  $(G \times G)$  conformal matter, together with the  $G$  of a collection of  $\mathcal{D}_{p_i}(G)$ . The constraint for a gauging which is either asymptotically free or conformal is<sup>15</sup>

<sup>15</sup>It is actually possible to obtain an interacting theory in the IR even when the gauge coupling is not asymptotically free when the gauge group is given as a product form [53]. This is due to the mixing of a nonanomalous  $U(1)$  symmetry with the  $R$ -symmetry in the infrared. This can result in some of the gauge couplings being “dangerously irrelevant.”




FIG. 8. All possible asymptotically free or conformal gaugings of an arbitrary collection of  $\mathcal{D}_{p_i}(G)$  and  $(G \times G)$  theories with two gauge nodes. A node with a solid border denotes an  $\mathcal{N} = 1$  gauge node, and a solid line between gauge nodes refers to the presence of  $(G \times G)$  conformal matter. A line connecting a gauge node and  $\mathcal{C}_n$  is a shorthand notation for all options of gaugings with the collections of  $\mathcal{D}_{p_i}(G)$  theories as described in Table V.




FIG. 9. All possible gaugings of conformal matter and  $\mathcal{D}_p(G)$  with three gauge nodes. See the caption of Fig. 8 for an explanation of the notation.

$$\sum_i \frac{2(p_i - 1)}{p} h_G^\vee + 2nh_G^\vee \leq 6h_G^\vee, \quad (4.4)$$

where we have used that the flavor central charge of each  $G$  flavor symmetry of conformal matter is  $2h_G^\vee$ . We can see that this is equivalent to the inequality in Eq. (4.1), and thus, a gauge node obtained by the  $\mathcal{N} = 1$  diagonal gauging involving  $n$  conformal matter factors can also involve any collection of  $\mathcal{D}_{p_i}(G)$  from  $\mathcal{C}_n$ .

The solution of the classification problem for the  $\mathcal{N} = 1$  gaugings of  $\mathcal{D}_{p_i}(G)$  and  $(G \times G)$  conformal matter, which do not in general have identical central charges  $a = c$ , is left for future work. In fact, in addition to solving the classification problem for the gaugings, it is necessary to determine whether each quiver formed by the gauging flows in the infrared to an interacting SCFT. Here, let us discuss some preliminary examples including  $\mathcal{D}_p(G)$  and  $(G \times G)$  conformal matter theories as in Figs. 8 and 9. We depict all such solutions involving precisely two gauge nodes in Fig. 8. Similarly, we show all those configurations involving three gauge nodes in Fig. 9. As depicted in these

figures, we can see that there are in total five types of diagrams involving two gauge nodes and seven types involving three gauge nodes, where each type may contain infinitely many quivers once all the options of  $\mathcal{C}_n$  are taken into account. In fact, the number of types grows rapidly with the number of gauge nodes; for example, for the case of four gauge nodes, there are already 22 types.

The gaugings that we have briefly introduced in this section form a broad class of  $\mathcal{N} = 1$  theories, typically without any non-Abelian flavor symmetry, although we do not expect them to have identical central charges. We leave a detailed analysis of this intriguing class of 4D  $\mathcal{N} = 1$  theories for future work.

### V. DISCUSSION

Throughout the paper, we explored under what circumstances a collection of 4D  $\mathcal{N} = 2$  Argyres–Douglas  $\mathcal{D}_{p_i}(G)$  theories can be diagonally gauged together via an  $\mathcal{N} = 1$   $G$ -vector multiplet in such a way that the  $\mathcal{N} = 1$  theory in the infrared is an interacting SCFT with identical central charges,  $a = c$ . We also determined that the addition of one




FIG. 10. Deforming the theories with one  $\mathcal{D}_p(G)$  and two adjoint chiral multiplets  $(\phi_1, \phi_2)$ , which we refer to as  $\hat{O}$ , via ADE-type superpotential. When the  $\hat{O}$  is deformed by  $\text{Tr} \phi_2^2$ , the  $\text{Tr} \phi_1^2$  becomes free during the RG flow and must be flipped by the flip-operator  $M$ . The question marks represent non-ADE deformations.

or two chiral multiplets transforming in the adjoint representation of  $G$  is compatible with the  $a = c$  property. Surprisingly, we discover that almost all asymptotically free gaugings (and also almost all conformal gaugings when additional adjoint chirals are included) flow to IR SCFTs with  $a = c$ , when  $G$  satisfies  $\text{gcd}(p_i, h_G^\vee) = 1$  for all  $p_i$ . This opens up a vast but well-curated landscape of minimally supersymmetric CFTs with exactly equal central charges. In this section, we discuss possible future directions.

Beyond the constructions worked out in the current paper, we noted in Sec. II A that the addition of a superpotential deformation to the  $(\mathcal{N} = 1)$ -gauging does not spoil the argument that the infrared SCFTs have  $a = c$ , as long as the assumption that there are no emergent Abelian flavor symmetries is not violated. This opens up an even vaster landscape of 4D  $\mathcal{N} = 1$  SCFTs with  $a = c$ . Particularly interesting examples in this class of theories are those obtained by deformation of a single  $\mathcal{D}_p(G)$  theory gauged together with two adjoint chiral multiplets. In [54], the authors showed that the superpotential deformations of SQCD with adjoint chirals have a classification that coincides with the ADE classification. Similar ADE-type deformations exist for  $\mathcal{N} = 1$  theories constructed out of gauging  $\mathcal{D}_p(G)$  with two adjoint chiral multiplets, as can be seen in Fig. 10, and we explore these and other superpotential deformations of the  $(\mathcal{N} = 1)$ -gaugings in [22].

We have considered the 4D  $\mathcal{N} = 1$  SCFTs that live as the infrared fixed points of the diagonal  $(\mathcal{N} = 1)$ -gauging of  $G$  of a collection of  $\mathcal{N} = 2$   $\mathcal{D}_p(G)$  theories, when  $G$  is an ADE Lie group. Two natural extensions of the current work present themselves immediately. First, the definition of the  $\mathcal{D}_p(G)$  theories can be extended to include  $G$  being of BCFG-type [55,56]. Second, the  $\mathcal{D}_p(G)$  are the special case of the  $\mathcal{D}_p^b(G)$  theories, where  $b = h_G^\vee$  [25]. In this

broader context, we can ask which collections of  $\mathcal{D}_p^b(G \in \text{ABCDEFG})$  can be gauged together by an  $\mathcal{N} = 1$  vector multiplet in such a way that one obtains an interacting infrared SCFT with  $a = c$  as was partially investigated in the  $\mathcal{N} = 2$  setup in [35]. We leave this generalization for future work.

In another direction, some of the  $\mathcal{N} = 2$  SCFTs  $\hat{\Gamma}(G)$  have a fascinating connection to  $\mathcal{N} = 4$  super Yang–Mills. It is discovered in [14,34] that the Schur indices of particular  $\hat{\Gamma}(G)$  theories with  $a = c$  can be written as the Schur index of  $\mathcal{N} = 4$  super Yang–Mills with rescaled fugacities.<sup>16</sup> Is there a sense in which the  $\mathcal{N} = 1$  SCFTs with  $a = c$  that we obtain in this paper are related to  $\mathcal{N} = 4$  super Yang–Mills? Such a connection may be visible from a holographic perspective and would be particularly interesting to determine.

In the holographic setup of AdS/CFT, it has long been known that the leading order contributions, in a large- $N$  expansion, to both  $a$  and  $c$  are identical [5]; however, it is rather rare to find holographic duals with no corrections that force the central charges to differ at a subleading order, at least, not without enhanced supersymmetry. At one loop, corrections to the holographic central charges can come from both open string loops and closed string loops, and all these contributions must either be absent or conspire to cancel when a dual SCFT has  $a = c$  exactly. Open string loops can arise from the inclusion of D-branes in the system, and these typically introduce  $1/N$  corrections. Closed string loops can arise either from the  $R^4$  correction to the type IIB string effective action or from the Kaluza–Klein tower on the compact space. It would be fascinating

<sup>16</sup>It is found that the  $\hat{\Gamma}(G)$  theories without flavor symmetries, which is a superset of  $\hat{\Gamma}(G)$  theories with  $a = c$ , will all have their Schur indices to be the same as that of  $\mathcal{N} = 4$  SYM up to re-scaling fugacities.





comet, and the  $n_a$  adjoint chiral multiplets compose the head. The star-shaped and comet-shaped quivers appear as the 3D ( $\mathcal{N} = 4$ ) mirror theory [59] to a 4D  $\mathcal{N} = 2$  theory of class  $\mathcal{S}$  [60,61].<sup>17</sup> More precisely, a star- or comet-shaped quiver with  $n$  tails and  $g$  loops corresponds to the 3D mirror theory for the class  $\mathcal{S}$  theory obtained from 6d  $\mathcal{N} = (2, 0)$  theory compactified on a Riemann surface with genus  $g$  and  $n$  punctures of specific types. In our setup, the  $\mathcal{D}_{p_i}(SU(pN))$  theory corresponds to the quiver tail labeled by the partition  $[\ell_1 N, \ell_2 N, \ell_3 N, \dots]$ . The Higgs branch of this quiver (as a 3D  $\mathcal{N} = 4$  theory) gives the Hitchin moduli space [26,59,61], whereas the Coulomb branch gives the Higgs branch of the 4D theory. Although what we have in Fig. 11 looks similar to the star-shaped or comet-shaped quiver in 3D mirror theory, it is rather different. In particular, the gauge nodes in the quiver tail of a star-shaped quiver are given by  $U(n)$  gauge groups, whereas in 4D, we can only have  $SU(n)$  type. Moreover, in 3D, the central gauge (core) node has to be gauged with an  $\mathcal{N} = 4$  (8 supercharges) vector multiplet with possible adjoint hypermultiplets. In contrast, in 4D, we consider an  $\mathcal{N} = 1$  (4 supercharges) vector multiplet with  $n_a$  adjoint chiral multiplets at the core node.

As a generalization of the  $\mathcal{N} = 2$   $\mathcal{D}_p(SU(pN))$  quiver tail theory, we can consider an  $\mathcal{N} = 1$  quiver tail instead. Since the quiver tail can be obtained via nilpotent Higgsing of a linear quiver gauge theory, one can consider similar deformations to obtain  $\mathcal{N} = 1$  quiver tails [64]. The  $\mathcal{N} = 1$  version of the quiver tail is substantially different from the  $\mathcal{N} = 2$  counterpart, allowing much wider set of SCFTs. It would be interesting to find an  $\mathcal{N} = 1$  analog of  $\mathcal{D}_p(G)$  Argyres–Douglas type theories that realize  $a = c$  upon gauging.

## B. Toward a geometric construction

Given that we construct a vast collection of 4D minimally supersymmetric SCFTs with  $a = c$ , it may be possible for these theories to have both a holographic description in terms of  $AdS_5$  and a top-down string theory construction. Four-dimensional  $\mathcal{N} = 1$  field theories can be obtained by compactifying  $F$  theory on a noncompact elliptically fibered Calabi–Yau fourfold; however, due to the presence of fluxes and instantons, it is not generally known when these 4D theories are conformal. Nonetheless, it is natural to ask, if we suppose that there exists an  $F$ -theory construction, whether the property of  $a = c$  is encoded in the compactification data via geometry in some natural way. In this subsection, we explore some of the ways in which the  $F$ -theory geometry must be constrained if it corresponds to one of the  $a = c$  theories that we consider.

<sup>17</sup>See [62,63] for a small sample of papers discussing comet-shaped quivers.

Superconformal theories with  $a = c$  that are formed by gauging together  $N$  different  $\mathcal{D}_{p_i}(G)$  theories typically have a  $U(1)^{N-1}$  flavor symmetry, which is formed by the anomaly-free combinations of the  $U(1)$  flavor symmetries inside the  $\mathcal{N} = 2$   $R$ -symmetry from each of the  $\mathcal{D}_{p_i}(G)$  theories. In  $F$ -theory compactifications, Abelian symmetries, whether gauge or flavor, arise when the elliptic fibration has a nontrivial Mordell–Weil group. The Mordell–Weil group is generated by the rational sections of the fibration.  $F$ -theory compactifications with the nontrivial Mordell–Weil group have been explored in many works, for which we give a small sample here [65–70]. The form of the geometry is then heavily constrained by the presence of these  $U(1)$  symmetries, as the fibration is thus forced to have many independent rational sections. Furthermore, it is important to note that the absence of non-Abelian gauge or flavor symmetry imposes that the elliptic fibration has no reducible singular fibers supported on divisors in the base.

The subtlety with a full Calabi–Yau fourfold compactification of  $F$  theory is that the compactification can contain nongeometric structures, for example, arising from D3-branes wrapping compact complex surfaces inside of the base of the elliptic fibration.<sup>18</sup> Due to these nongeometric effects, it is particularly difficult to find the precise condition to ensure conformality in the resulting 4D theory via noncompact fourfold compactifications that have to be imposed. To highlight the difficulty, we note that such conditions have been determined for 6d SCFTs due to the particular rigidity of threefold compactifications. Some work has been done in [52] to explore whether certain elliptically fibered Calabi–Yau fourfolds lead to 4D  $\mathcal{N} = 1$  SCFTs.

We expect that the geometric construction would shed light upon the putative holographic understanding, or vice versa. In particular, if there exists an  $AdS_5 \times X_5$  solution of Type IIB supergravity, which is holographically dual to some of our gauged theories, then the  $X_5$  is highly constrained by the  $a = c$  condition: all string-loop contributions must cancel. An exploration of these cancellations will appear in [78]. Similarly, we expect that  $X_5$  is related to the elliptically fibered Calabi–Yau fourfold through which the same  $\mathcal{N} = 1$  theory is engineered in  $F$  theory; in this way, the  $a = c$  field-theoretic condition on the  $X_5$  should lead to geometric restrictions on the  $F$ -theory compactification manifold.

Instead of considering compactifications of  $F$  theory on Calabi–Yau fourfolds, with all of the attendant complexities, we can start from the 6d (1,0) SCFTs and consider a further compactification on a Riemann surface. A vast collection of six-dimensional SCFTs can be engineered in  $F$  theory via elliptically fibered Calabi–Yau threefolds

<sup>18</sup>Geometric and topological data relevant for elliptically fibered Calabi–Yau compactifications of  $F$  theory are systematically studied, for example, in [71–77].

[79,80].<sup>19</sup> Four-dimensional  $\mathcal{N} = 1$  theories can be constructed starting from these 6d (1,0) theories and compactifying either on a genus  $g > 1$  punctured Riemann surface, or else on a torus with flux inside of the 6d flavor symmetry. Some related aspects of the 4D theories obtained in this way have been explored in, for example, [91–94] and [95,96], respectively. In such cases, we understand all the UV Abelian symmetries that may mix with the superconformal flavor symmetry under  $a$ -maximization, and determining the superconformal  $R$ -symmetry is straightforward. The mixing coefficients, however, are not simply related to geometric quantities, and thus, it obscures the origin of  $a = c$  in the geometry.

Alternatively, 4D  $\mathcal{N} = 1$  theories can arise from compactification of the 6d (2,0) SCFTs [18,97–102]. Before approaching the  $\mathcal{N} = 1$  theories, we can consider the  $\mathcal{N} = 2$  theories that were obtained from gaugings of  $\mathcal{D}_p(G)$  and conformal matter, and called  $\hat{\Gamma}(G)$ , in [14]. Each of the  $\mathcal{D}_p(G)$  and conformal matter factors considered in the gauging individually has a description in terms of class  $\mathcal{S}$  [60,61]. The former are associated to spheres with one regular and one irregular puncture [26] and the latter to spheres with two full punctures and one simple puncture [36,51]. However, except for  $\hat{\Gamma} = \hat{A}$ , the gluing always

<sup>19</sup>Via elliptic fibrations, the geometric engineering of  $F$  theory provides different geometric perspectives for 6d (1,0) theories (see, e.g., [81–90] for some explicit geometric constructions), and we get a superconformal field theory when the Calabi–Yau threefold is noncompact and satisfies particular additional conditions.

involves the diagonal gauging of the flavor symmetries associated to at least three punctures, and this is not an operation that is geometric. Indeed, for most of the  $\hat{D}_n(G)$  and  $\hat{E}_n(G)$  theories, whether Lagrangian or not, there is no known class  $\mathcal{S}$  construction.

There exists alternative geometric construction for a large class of 4D SCFTs from Calabi–Yau 3-fold singularities [103,104], some of which includes a subset of  $\hat{\Gamma}(G)$  theories [35]. In this way, the construction of  $\mathcal{N} = 1$  SCFTs either directly from  $F$  theory on Calabi–Yau fourfolds, or indirectly from compactifications of the 6d (1,0) SCFTs, provides a promising avenue for a top-down realization of the  $a = c$  theories studied in this paper.

## ACKNOWLEDGMENTS

We thank Richard Derryberry for a helpful discussion. M. J. K. is partially supported by a Sherman Fairchild Postdoctoral Fellowship, the U.S. Department of Energy, Office of Science, Office of High Energy Physics, under Award No. DE-SC0011632, and the National Research Foundation of Korea (NRF) Grant No. NRF-2020R1A4A3079707. M. J. K., K. H. L., and J. S. are all partly supported by the NRF Grant No. NRF-2020R1C1C1007591. C. L. acknowledges support from DESY (Hamburg, Germany), a member of the Helmholtz Association HGF. K. H. L. and J. S. are partly supported by the Breaking the Scientific Wall (BTS-W) program provided by Korea Advanced Institute of Science and Technology (KAIST). J. S. is also supported by the Start-up Research Grant for new faculty provided by KAIST.

- 
- [1] Z. Komargodski and A. Schwimmer, On renormalization group flows in four dimensions, *J. High Energy Phys.* **12** (2011) 099.
  - [2] K. Maruyoshi, E. Nardoni, and J. Song, Landscape of Simple Superconformal Field Theories in 4d, *Phys. Rev. Lett.* **122**, 121601 (2019).
  - [3] D. M. Hofman and J. Maldacena, Conformal collider physics: Energy and charge correlations, *J. High Energy Phys.* **05** (2008) 012.
  - [4] O. Aharony and M. Evtikhiev, On four dimensional  $\mathcal{N} = 3$  superconformal theories, *J. High Energy Phys.* **04** (2016) 040.
  - [5] M. Henningson and K. Skenderis, The holographic Weyl anomaly, *J. High Energy Phys.* **07** (1998) 023.
  - [6] D. Anselmi and A. Kehagias, Subleading corrections and central charges in the AdS/CFT correspondence, *Phys. Lett. B* **455**, 155 (1999).
  - [7] P. Kovtun, D. T. Son, and A. O. Starinets, Viscosity in Strongly Interacting Quantum Field Theories from Black Hole Physics, *Phys. Rev. Lett.* **94**, 111601 (2005).
  - [8] A. Buchel, R. C. Myers, and A. Sinha, Beyond eta/s = 1/4 pi, *J. High Energy Phys.* **03** (2009) 084.
  - [9] Y. Kats and P. Petrov, Effect of curvature squared corrections in AdS on the viscosity of the dual gauge theory, *J. High Energy Phys.* **01** (2009) 044.
  - [10] L. Di Pietro and Z. Komargodski, Cardy formulae for SUSY theories in  $d = 4$  and  $d = 6$ , *J. High Energy Phys.* **12** (2014) 031.
  - [11] E. Perlmutter, M. Rangamani, and M. Rota, Central Charges and the Sign of Entanglement in 4D Conformal Field Theories, *Phys. Rev. Lett.* **115**, 171601 (2015).
  - [12] D. Anselmi, D. Z. Freedman, M. T. Grisaru, and A. A. Johansen, Nonperturbative formulas for central functions of supersymmetric gauge theories, *Nucl. Phys.* **B526**, 543 (1998).
  - [13] X. O. Camanho, J. D. Edelstein, J. Maldacena, and A. Zhiboedov, Causality constraints on corrections to the graviton three-point coupling, *J. High Energy Phys.* **02** (2016) 020.

- [14] M. J. Kang, C. Lawrie, and J. Song, Infinitely many 4d  $\mathcal{N} = 2$  SCFTs with  $a = c$  and beyond, *Phys. Rev. D* **104**, 105005 (2021).
- [15] Y. Tachikawa and B. Wecht, Explanation of the Central Charge Ratio 27/32 in Four-Dimensional Renormalization Group Flows between Superconformal Theories, *Phys. Rev. Lett.* **103**, 061601 (2009).
- [16] D. Green, Z. Komargodski, N. Seiberg, Y. Tachikawa, and B. Wecht, Exactly marginal deformations and global symmetries, *J. High Energy Phys.* **06** (2010) 106.
- [17] R. G. Leigh and M. J. Strassler, Exactly marginal operators and duality in four-dimensional  $\mathcal{N} = 1$  supersymmetric gauge theory, *Nucl. Phys.* **B447**, 95 (1995).
- [18] I. Bah, C. Beem, N. Bobev, and B. Wecht, Four-dimensional SCFTs from M5-branes, *J. High Energy Phys.* **06** (2012) 005.
- [19] I. Bah and N. Bobev, Linear quivers and  $\mathcal{N} = 1$  SCFTs from M5-branes, *J. High Energy Phys.* **08** (2014) 121.
- [20] F. Benini, Y. Tachikawa, and B. Wecht, Sicilian gauge theories and  $\mathcal{N} = 1$  dualities, *J. High Energy Phys.* **01** (2010) 088.
- [21] M. J. Kang, C. Lawrie, K.-H. Lee, and J. Song, Operator spectroscopy for 4d SCFTs with  $a = c$  (to be published).
- [22] M. J. Kang, C. Lawrie, K.-H. Lee, and J. Song, Landscape of 4d  $\mathcal{N} = 1$  SCFTs with  $a = c$  (to be published).
- [23] S. Cecotti and M. Del Zotto, Infinitely many  $\mathcal{N} = 2$  SCFT with ADE flavor symmetry, *J. High Energy Phys.* **01** (2013) 191.
- [24] S. Cecotti, M. Del Zotto, and S. Giacomelli, More on the  $\mathcal{N} = 2$  superconformal systems of type  $D_p(G)$ , *J. High Energy Phys.* **04** (2013) 153.
- [25] Y. Wang and D. Xie, Classification of Argyres–Douglas theories from M5 branes, *Phys. Rev. D* **94**, 065012 (2016).
- [26] D. Xie, General Argyres–Douglas theory, *J. High Energy Phys.* **01** (2013) 100.
- [27] P. Agarwal, K. Maruyoshi, and J. Song,  $\mathcal{N} = 1$  deformations and RG flows of  $\mathcal{N} = 2$  SCFTs, part II: Non-principal deformations, *J. High Energy Phys.* **12** (2016) 103; **04** (2017) 113(A).
- [28] P. Agarwal, A. Sciarappa, and J. Song,  $\mathcal{N} = 1$  Lagrangians for generalized Argyres–Douglas theories, *J. High Energy Phys.* **10** (2017) 211.
- [29] S. Benvenuti and S. Giacomelli, Lagrangians for generalized Argyres–Douglas theories, *J. High Energy Phys.* **10** (2017) 106.
- [30] K. Maruyoshi and J. Song,  $\mathcal{N} = 1$  deformations and RG flows of  $\mathcal{N} = 2$  SCFTs, *J. High Energy Phys.* **02** (2017) 075.
- [31] K. Maruyoshi and J. Song, Enhancement of Supersymmetry via Renormalization Group Flow and the Superconformal Index, *Phys. Rev. Lett.* **118**, 151602 (2017).
- [32] S. Cecotti and C. Vafa, Classification of complete  $\mathcal{N} = 2$  supersymmetric theories in 4 dimensions, [arXiv:1103.5832](https://arxiv.org/abs/1103.5832).
- [33] M. Buican and T. Nishinaka, Conformal manifolds in four dimensions and chiral algebras, *J. Phys. A* **49**, 465401 (2016).
- [34] M. Buican and T. Nishinaka,  $\mathcal{N} = 4$  SYM, Argyres–Douglas theories, and an exact graded vector space isomorphism, *J. High Energy Phys.* **04** (2022) 028.
- [35] C. Closset, S. Giacomelli, S. Schafer-Nameki, and Y.-N. Wang, 5d and 4d SCFTs: Canonical singularities, trinions and S-dualities, *J. High Energy Phys.* **05** (2021) 274.
- [36] M. Del Zotto, C. Vafa, and D. Xie, Geometric engineering, mirror symmetry and  $6d_{(1,0)} \rightarrow 4d_{(\mathcal{N}=2)}$ , *J. High Energy Phys.* **11** (2015) 123.
- [37] K. A. Intriligator and B. Wecht, The exact superconformal R symmetry maximizes a, *Nucl. Phys.* **B667**, 183 (2003).
- [38] C. Córdova, T. T. Dumitrescu, and K. Intriligator, Exploring 2-group global symmetries, *J. High Energy Phys.* **02** (2019) 184.
- [39] D. Kutasov, A. Parnachev, and D. A. Sahakyan, Central charges and  $U(1)(R)$  symmetries in  $\mathcal{N} = 1$  superYang–Mills, *J. High Energy Phys.* **11** (2003) 013.
- [40] E. Barnes, K. A. Intriligator, B. Wecht, and J. Wright, Evidence for the strongest version of the 4d a-theorem, via a-maximization along RG flows, *Nucl. Phys.* **B702**, 131 (2004).
- [41] S. Benvenuti and S. Giacomelli, Supersymmetric Gauge Theories with Decoupled Operators and Chiral Ring Stability, *Phys. Rev. Lett.* **119**, 251601 (2017).
- [42] T. Arakawa, Associated varieties of modules over Kac–Moody algebras and  $C_2$ -cofiniteness of W-algebras, *Int. Math. Res. Not.* **2015**, 11605 (2015).
- [43] J. Song, D. Xie, and W. Yan, Vertex operator algebras of Argyres–Douglas theories from M5-branes, *J. High Energy Phys.* **12** (2017) 123.
- [44] I. Affleck, M. Dine, and N. Seiberg, Dynamical supersymmetry breaking in supersymmetric QCD, *Nucl. Phys.* **B241**, 493 (1984).
- [45] N. Seiberg, Exact results on the space of vacua of four-dimensional SUSY gauge theories, *Phys. Rev. D* **49**, 6857 (1994).
- [46] K. Maruyoshi, Y. Tachikawa, W. Yan, and K. Yonekura,  $\mathcal{N} = 1$  dynamics with  $T_N$  theory, *J. High Energy Phys.* **10** (2013) 010.
- [47] P. Agarwal, K.-H. Lee, and J. Song, Classification of large  $N$  superconformal gauge theories with a dense spectrum, *J. High Energy Phys.* **10** (2021) 049.
- [48] P. Agarwal and J. Song, Large  $N$  gauge theories with a dense spectrum and the weak gravity conjecture, *J. High Energy Phys.* **05** (2021) 124.
- [49] F. Baume, M. J. Kang, and C. Lawrie, Two 6d origins of 4d SCFTs: Class S and 6d (1,0) on a torus, [arXiv:2106.11990](https://arxiv.org/abs/2106.11990).
- [50] M. Del Zotto, J. J. Heckman, A. Tomasiello, and C. Vafa, 6d conformal matter, *J. High Energy Phys.* **02** (2015) 054.
- [51] K. Ohmori, H. Shimizu, Y. Tachikawa, and K. Yonekura, 6d  $\mathcal{N} = (1, 0)$  theories on  $T^2$  and class S theories: Part I, *J. High Energy Phys.* **07** (2015) 014.
- [52] F. Apruzzi, J. J. Heckman, D. R. Morrison, and L. Tizzano, 4D gauge theories with conformal matter, *J. High Energy Phys.* **09** (2018) 088.
- [53] E. Barnes, K. A. Intriligator, B. Wecht, and J. Wright,  $\mathcal{N} = 1$  RG flows, product groups, and a-maximization, *Nucl. Phys.* **B716**, 33 (2005).
- [54] K. A. Intriligator and B. Wecht, RG fixed points and flows in SQCD with adjoints, *Nucl. Phys.* **B677**, 223 (2004).
- [55] F. Carta, S. Giacomelli, N. Mekareeya, and A. Mininno, Conformal manifolds and 3d mirrors of Argyres–Douglas theories, *J. High Energy Phys.* **08** (2021) 015.

- [56] Y. Wang and D. Xie, Codimension-two defects and Argyres–Douglas theories from outer-automorphism twist in 6d (2,0) theories, *Phys. Rev. D* **100**, 025001 (2019).
- [57] S. Kachru and E. Silverstein, 4-D Conformal Theories and Strings on Orbifolds, *Phys. Rev. Lett.* **80**, 4855 (1998).
- [58] J. McKay, Graphs, singularities, and finite groups, in *The Santa Cruz Conference on Finite Groups (University California, Santa Cruz, California, 1979)*, Proceedings of Symposia in Pure Mathematics Vol. 37 (American Mathematical Society, Providence, R.I., 1980), pp. 183–186.
- [59] F. Benini, Y. Tachikawa, and D. Xie, Mirrors of 3d Sicilian theories, *J. High Energy Phys.* **09** (2010) 063.
- [60] D. Gaiotto,  $\mathcal{N} = 2$  dualities, *J. High Energy Phys.* **08** (2012) 034.
- [61] D. Gaiotto, G. W. Moore, and A. Neitzke, Wall-crossing, hitchin systems, and the WKB approximation, [arXiv:0907.3987](https://arxiv.org/abs/0907.3987).
- [62] G. Bonelli, N. Fasola, and A. Tanzini, Defects, nested instantons and comet shaped quivers, *Lett. Math. Phys.* **111**, 34 (2021).
- [63] S. Rayan and L. P. Schaposnik, Moduli spaces of generalized hyperpolygons, *Quart. J. Math. Oxford Ser.* **72**, 137 (2021).
- [64] P. Agarwal, I. Bah, K. Maruyoshi, and J. Song, Quiver tails and  $\mathcal{N} = 1$  SCFTs from M5-branes, *J. High Energy Phys.* **03** (2015) 049.
- [65] J. Borchmann, C. Mayrhofer, E. Palti, and T. Weigand, Elliptic fibrations for  $SU(5) \times U(1) \times U(1)$  F-theory vacua, *Phys. Rev. D* **88**, 046005 (2013).
- [66] M. Cvetič, D. Klemm, and H. Piragua, F-theory compactifications with multiple  $U(1)$ -factors: Constructing elliptic fibrations with rational sections, *J. High Energy Phys.* **06** (2013) 067.
- [67] M. Esole, M. J. Kang, and S.-T. Yau, A new model for elliptic fibrations with a rank one Mordell-Weil group: I. Singular fibers and semi-stable degenerations, [arXiv:1410.0003](https://arxiv.org/abs/1410.0003).
- [68] C. Lawrie, S. Schafer-Nameki, and J.-M. Wong, F-theory and all things rational: Surveying  $U(1)$  symmetries with rational sections, *J. High Energy Phys.* **09** (2015) 144.
- [69] D. R. Morrison and D. S. Park, F-theory and the Mordell-Weil group of elliptically-fibered Calabi-Yau threefolds, *J. High Energy Phys.* **10** (2012) 128.
- [70] D. R. Morrison and W. Taylor, Sections, multisections, and  $U(1)$  fields in F-theory, [arXiv:1404.1527](https://arxiv.org/abs/1404.1527).
- [71] M. Esole, P. Jefferson, and M. J. Kang, Euler characteristics of crepant resolutions of Weierstrass models, *Commun. Math. Phys.* **371**, 99 (2019).
- [72] M. Esole and M. J. Kang, Characteristic numbers of crepant resolutions of Weierstrass models, [arXiv:1807.08755](https://arxiv.org/abs/1807.08755).
- [73] M. Esole and M. J. Kang, Characteristic numbers of elliptic fibrations with nontrivial Mordell-Weil groups, [arXiv:1808.07054](https://arxiv.org/abs/1808.07054).
- [74] J. Gray, A. S. Haupt, and A. Lukas, All complete intersection Calabi-Yau four-folds, *J. High Energy Phys.* **07** (2013) 070.
- [75] T. W. Grimm, T.-W. Ha, A. Klemm, and D. Klemm, Computing brane and flux superpotentials in F-theory compactifications, *J. High Energy Phys.* **04** (2010) 015.
- [76] S. Katz, D. R. Morrison, S. Schafer-Nameki, and J. Sully, Tate’s algorithm and F-theory, *J. High Energy Phys.* **08** (2011) 094.
- [77] W. Taylor and Y.-N. Wang, The F-theory geometry with most flux vacua, *J. High Energy Phys.* **12** (2015) 164.
- [78] C. Couzens, M. J. Kang, C. Lawrie, K.-H. Lee, and J. Song (to be published).
- [79] J. J. Heckman, D. R. Morrison, T. Rudelius, and C. Vafa, Atomic classification of 6D SCFTs, *Fortschr. Phys.* **63**, 468 (2015).
- [80] J. J. Heckman, D. R. Morrison, and C. Vafa, On the classification of 6D SCFTs and generalized ADE orbifolds, *J. High Energy Phys.* **05** (2014) 028; **06** (2015) 017(E).
- [81] M. Esole, R. Jagadeesan, and M. J. Kang, The geometry of  $G_2$ , spin(7), and spin(8)-models, [arXiv:1709.04913](https://arxiv.org/abs/1709.04913).
- [82] M. Esole, R. Jagadeesan, and M. J. Kang, 48 crepant paths to  $SU(2) \times SU(3)$ , [arXiv:1905.05174](https://arxiv.org/abs/1905.05174).
- [83] M. Esole and P. Jefferson, The geometry of  $SO(3)$ ,  $SO(5)$ , and  $SO(6)$  models, [arXiv:1905.12620](https://arxiv.org/abs/1905.12620).
- [84] M. Esole, P. Jefferson, and M. J. Kang, The geometry of  $F_4$ -models, [arXiv:1704.08251](https://arxiv.org/abs/1704.08251).
- [85] M. Esole and M. J. Kang, Flopping and slicing:  $SO(4)$  and Spin(4)-models, *Adv. Theor. Math. Phys.* **23**, 1003 (2019).
- [86] M. Esole and M. J. Kang, The geometry of the  $SU(2) \times G_2$ -model, *J. High Energy Phys.* **02** (2019) 091.
- [87] M. Esole, M. J. Kang, and S.-T. Yau, Mordell-Weil torsion, anomalies, and phase transitions, [arXiv:1712.02337](https://arxiv.org/abs/1712.02337).
- [88] M. Esole and S.-H. Shao, M-theory on elliptic Calabi–Yau threefolds and 6d anomalies, [arXiv:1504.01387](https://arxiv.org/abs/1504.01387).
- [89] M. Esole, S.-H. Shao, and S.-T. Yau, Singularities and gauge theory phases, *Adv. Theor. Math. Phys.* **19**, 1183 (2015).
- [90] M. Esole, S.-H. Shao, and S.-T. Yau, Singularities and gauge theory phases II, *Adv. Theor. Math. Phys.* **20**, 683 (2016).
- [91] D. Gaiotto and S. S. Razamat,  $\mathcal{N} = 1$  theories of class  $\mathcal{S}_k$ , *J. High Energy Phys.* **07** (2015) 073.
- [92] J. J. Heckman, P. Jefferson, T. Rudelius, and C. Vafa, Punctures for theories of class  $\mathcal{S}_r$ , *J. High Energy Phys.* **03** (2017) 171.
- [93] D. R. Morrison and C. Vafa, F-theory and  $\mathcal{N} = 1$  SCFTs in four dimensions, *J. High Energy Phys.* **08** (2016) 070.
- [94] S. S. Razamat, C. Vafa, and G. Zafrir, 4d  $\mathcal{N} = 1$  from 6d (1,0), *J. High Energy Phys.* **04** (2017) 064.
- [95] I. Bah, A. Hanany, K. Maruyoshi, S. S. Razamat, Y. Tachikawa, and G. Zafrir, 4d  $\mathcal{N} = 1$  from 6d  $\mathcal{N} = (1,0)$  on a torus with fluxes, *J. High Energy Phys.* **06** (2017) 022.
- [96] H.-C. Kim, S. S. Razamat, C. Vafa, and G. Zafrir, Compactifications of ADE conformal matter on a torus, *J. High Energy Phys.* **09** (2018) 110.
- [97] P. Agarwal, K. Intriligator, and J. Song, Infinitely many  $\mathcal{N} = 1$  dualities from  $m + 1 - m = 1$ , *J. High Energy Phys.* **10** (2015) 035.
- [98] C. Beem and A. Gadde, The  $\mathcal{N} = 1$  superconformal index for class  $\mathcal{S}$  fixed points, *J. High Energy Phys.* **04** (2014) 036.

- [99] G. Bonelli, S. Giacomelli, K. Maruyoshi, and A. Tanzini,  $N = 1$  geometries via M-theory, *J. High Energy Phys.* **10** (2013) 227.
- [100] A. Gadde, K. Maruyoshi, Y. Tachikawa, and W. Yan, New  $N = 1$  dualities, *J. High Energy Phys.* **06** (2013) 056.
- [101] D. Xie, M5 brane and four dimensional  $\mathcal{N} = 1$  theories I, *J. High Energy Phys.* **04** (2014) 154.
- [102] D. Xie and K. Yonekura, Generalized Hitchin system, spectral curve and  $\mathcal{N} = 1$  dynamics, *J. High Energy Phys.* **01** (2014) 001.
- [103] D. Xie and S.-T. Yau, 4d  $\mathcal{N} = 2$  SCFT and singularity theory Part I: Classification, [arXiv:1510.01324](https://arxiv.org/abs/1510.01324).
- [104] D. Xie and S.-T. Yau, Singularity, Sasaki-Einstein manifold, Log del Pezzo surface and  $\mathcal{N} = 1$  AdS/CFT correspondence: Part I, [arXiv:1903.00150](https://arxiv.org/abs/1903.00150).