

Photon emissivity of the quark-gluon plasma: A lattice QCD analysis of the transverse channel

Marco Cè^{1,2}, Tim Harris,³ Ardit Krasniqi,⁴ Harvey B. Meyer^{4,5,6} and Csaba Török⁴

¹*Albert Einstein Center for Fundamental Physics (AEC) and Institut für Theoretische Physik, Universität Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland*

²*Theoretical Physics Department, CERN, CH-1211 Geneva 23, Switzerland*

³*School of Physics and Astronomy, University of Edinburgh, EH9 3JZ, United Kingdom*

⁴*PRISMA+ Cluster of Excellence and Institut für Kernphysik, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany*

⁵*Helmholtz Institut Mainz, Johannes Gutenberg-Universität Mainz, D-55099 Mainz, Germany*

⁶*GSI Helmholtzzentrum für Schwerionenforschung, 64291 Darmstadt, Germany*



(Received 18 May 2022; accepted 15 August 2022; published 6 September 2022)

We present results for the thermal photon emissivity of the quark-gluon plasma derived from spatially transverse vector correlators computed in lattice QCD at a temperature of 250 MeV. The analysis of the spectral functions, performed at fixed spatial momentum, is based on continuum-extrapolated correlators obtained with two flavors of dynamical Wilson fermions. We compare the next-to-leading-order perturbative QCD correlators, as well as the $\mathcal{N} = 4$ supersymmetric Yang-Mills correlators at infinite coupling, to the correlators from lattice QCD and find them to lie within $\sim 10\%$ of each other. We then refine the comparison, performing it at the level of filtered spectral functions obtained model-independently via the Backus-Gilbert method. Motivated by these studies, for frequencies $\omega \lesssim 2.5$ GeV we use fit *Ansätze* to the spectral functions that perform well when applied to mock data generated from the NLO QCD or from the strongly coupled SYM spectral functions, while the high-frequency part, $\omega \gtrsim 2.5$ GeV, is matched to NLO QCD. We compare our results for the photon emissivity to our previous analysis of a different vector channel at the same temperature. We obtain the most stringent constraint at photon momenta around $k \simeq 0.8$ GeV, for which we find a differential photon emission rate per unit volume of $d\Gamma_\gamma/d^3k = (\alpha_{\text{em}}/(\exp(k/T) - 1)) \times (2.2 \pm 0.8) \times 10^{-3}$ GeV.

DOI: [10.1103/PhysRevD.106.054501](https://doi.org/10.1103/PhysRevD.106.054501)

I. INTRODUCTION

Photons and lepton pairs have long been considered to provide direct information on the quark-gluon plasma (QGP), since they are penetrating probes of the QGP due to their colorless nature [1–5]. During heavy-ion collisions, photons can escape the plasma without scattering via the strong interaction, but discriminating between different sources is quite challenging because of the continuous emission of photons during the spacetime evolution of the fireball. The detected photons are divided into two main categories: decay and direct photons. The former refers to photons coming from the electromagnetic decays of final-state hadrons, while the latter includes all photons created in the collision before the final hadrons completely

decouple. The decay photons give a much larger contribution to the signal than the direct photons, and they provide valuable information for particle reconstruction. However, when studying direct photons, that contribution amounts to a large background and has to be subtracted from the total photon yield.

Direct photons are produced via several mechanisms during the evolution of a heavy-ion collision (cf. Ref. [5]), but the two major sources are initial hard parton-parton scattering (prompt photons), and photons originating from the QGP (thermal photons). The latest direct photon yield results have been published by the PHENIX [6] and STAR [7] Collaborations at RHIC, and by ALICE [8] at the LHC. The direct photon yield measured by the PHENIX and ALICE Collaborations shows an excess at low transverse momentum, $p_T \lesssim 3$ GeV, with respect to the theoretical predictions [9,10]. The results of the STAR Collaboration [7], however, are in better agreement with the model results. The excess observed in the PHENIX and ALICE results is in the momentum range where the dominant contribution comes from thermal photons [10].

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

An important ingredient in the theoretical prediction is the emission rate of thermal photons per unit volume, which has to be integrated over the expanding spacetime volume of the medium to obtain the total thermal photon yield [10]. This rate has been calculated at leading order in the strong coupling constant [11], and the calculation has been extended to include corrections which arise from interactions with soft gluons [12]. The thermal photon emission rate from this extended calculation has a similar functional form to the leading-order one, and represents a 20% increase at $\alpha_s \simeq 0.3$. This modest increase reduces the tensions, though it is still insufficient to explain the excess observed by the PHENIX and ALICE Collaborations and highlights that the leading-order prediction may receive large corrections at relevant temperatures.

Therefore, a precise nonperturbative calculation of the thermal photon production rate using lattice QCD is highly desirable and would help to resolve the tensions or confirm the already existing weak coupling results. Although lattice QCD has been very successful in calculating observables in vacuum as well as in thermal equilibrium, reliably accessing near-equilibrium quantities such as the thermal photon rate is very challenging, because the calculation of real-time observables requires analytic continuation using a finite, limited number of noisy Euclidean correlator data. Moreover, the Euclidean correlators that need to be inverted to obtain information on the spectral function are rather insensitive to the infrared features of these spectral functions [13–16]. In spite of the difficulties, several methods have been devised to constrain the ill-posed numerical inversion problem using both model-dependent and independent approaches.

To determine the thermal photon production rate via analytic continuation, the starting Euclidean observable is the vector current correlation function at finite temperature. Although this correlator has been investigated extensively also on thermal ensembles, earlier analyses focused mainly on vanishing spatial momentum, which is relevant for the determination of the electrical conductivity of the plasma [17–22]. Continuum-extrapolated vector current correlators at finite spatial momenta obtained in quenched lattice QCD simulations were analyzed for the photon rate in Ref. [23]. In Ref. [24], the same goal was pursued with continuum-extrapolated vector correlators based on two-flavor dynamical simulations, focusing on an infrared-dominated channel, the difference between the transverse and longitudinal channels (T – L channel). In both Refs. [23,24], information on the spectral function was extracted from the Euclidean data using fit *Ansätze*.

In this work, we apply two different strategies to analyze the correlation functions: the Backus-Gilbert method [25,26] and a fit method where we apply various simple fit *Ansätze* matched to perturbation theory at high frequencies. In contrast to our earlier work, we investigate the spatially transverse channel of the vector current correlator

at finite spatial momentum. In Sec. II, we collect the relevant basic formulas and discuss the advantages of investigating this particular channel. We also present in that section an overview of the spectral function in this channel in different regimes. Details on the lattice configurations and the observables relevant in this study, including the continuum extrapolation, are presented in Sec. III. The analysis of the continuum-extrapolated correlator in the transverse channel is presented in Sec. IV. There, we also present our estimate on the thermal photon rate based on this channel and compare it to available results from the literature. We finally give our conclusions and an outlook in Sec. V.

II. THEORETICAL BACKGROUND

In this section, we collect the main definitions of the correlation functions to be analyzed and describe the existing theory predictions with which we will confront our lattice QCD calculations.

A. Basic definitions

The spectral function of the vector current $J_\mu = \bar{\Psi}\gamma_\mu T^a \Psi$, for a generic matrix T^a acting in quark-flavor space, is defined as

$$\rho_{\mu\nu}(\omega, \mathbf{k}) = \int dt d^3x e^{i(\omega t - \mathbf{k}\cdot\mathbf{x})} \langle [J_\mu(t, \mathbf{x}), J_\nu(0)^\dagger] \rangle. \quad (1)$$

The Minkowskian time evolution of the current is given by $J_\mu(t, \mathbf{x}) = e^{iHt} J_\mu(0, \mathbf{x}) e^{-iHt}$. The expectation value of the commutator is taken with respect to the thermal density matrix, $e^{-\beta H}/Z$, with $\beta = 1/T$ being the inverse temperature. To leading order in the fine-structure constant, $\alpha_{\text{em}} = e^2/(4\pi)$, letting $T^a = \text{diag}(\frac{2}{3}, \frac{-1}{3}, \frac{-1}{3}, \dots)$ contain the quark electric charges, the thermal photon production rate per unit volume of quark-gluon plasma can be expressed as [27]

$$\frac{d\Gamma_\gamma(k)}{d^3k} = \frac{\alpha_{\text{em}} \rho_V(\omega = k, k)}{\pi^2} \frac{1}{4k} \frac{1}{e^{k/T} - 1} + \mathcal{O}(\alpha_{\text{em}}^2), \quad (2)$$

in terms of the vector channel spectral function $\rho_V(\omega, k) = -\rho^\mu_\mu(\omega, \mathbf{k})$, where $k = |\mathbf{k}|$. It corresponds to the choice $\lambda = 1$ in the following linear combination, introduced in Refs. [24,28]:

$$\rho(\omega, k, \lambda) = 2\rho_T(\omega, k) + \lambda\rho_L(\omega, k), \quad (3)$$

where

$$\begin{aligned} \rho_T(\omega, k) &= \frac{1}{2} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) \rho_{ij} \quad \text{and} \\ \rho_L(\omega, k) &= \frac{k_i k_j}{k^2} \rho_{ij} - \rho_{00} \end{aligned} \quad (4)$$

denote the spatially transverse and longitudinal spectral functions, respectively. As a consequence of current conservation, expression (3) is independent of λ for light-cone kinematics $\omega = k$. Due to this fact, $\rho_V(k, k)$ can be replaced by $\rho(k, k, \lambda)$ with arbitrary λ in the evaluation of the photon rate [24,28] of Eq. (2). In Refs. [24,29], the $\lambda = -2$ case was investigated, which corresponds to the difference of the transverse and longitudinal channels,

$$\rho(\omega, k, -2) = 2[\rho_T(\omega, k) - \rho_L(\omega, k)]. \quad (5)$$

This channel, $\rho(\omega, k, -2)$, is particularly interesting because it is non-negative for $0 \leq \omega \leq k$ and highly suppressed when $\omega > k$. Therefore, it is very sensitive to infrared physics of interest. It vanishes in the vacuum and also satisfies a superconvergent sum rule, demonstrated in Refs. [24,28] and utilized as a constraint in the spectral reconstruction from Euclidean correlators in Ref. [24].

In this work, we investigate the transverse channel, corresponding to using $\lambda = 0$ in Eq. (3). As we shall see below, it has complementary properties to the previously studied $\lambda = -2$ channel; in particular, its spectral function is non-negative for all $\omega \geq 0$. The transverse-channel Euclidean two-point functions of the vector current carrying a definite spatial momentum are related to their corresponding spectral function via (see, e.g., Ref. [15])

$$G_T(\tau, k) = \int_0^\infty \frac{d\omega}{2\pi} \rho_T(\omega, k) K(\omega, \tau), \quad (6)$$

where the kernel is given as

$$K(\omega, \tau) = \frac{\cosh[\omega(\beta/2 - \tau)]}{\sinh(\omega\beta/2)}, \quad (7)$$

and $\tau = it$. This spectral representation of the Euclidean correlators will first be used to confront them with theory predictions for the spectral function, and, in a second stage, to fit an *Ansatz* for the spectral function to the Euclidean correlators computed in lattice QCD.

We remark that in Secs. III and IV, we use the flavor matrix $T^a = \text{diag}(\frac{1}{\sqrt{2}}, \frac{-1}{\sqrt{2}})$ in two-flavor QCD—i.e., we calculate the isovector vector current correlator. This specifies in particular the normalization of our results for G_T , G_{00} , or ρ_T . If we are willing to use the approximate SU(3) flavor symmetry in the high-temperature phase of QCD and neglect dynamical strange-quark effects as well as the charm contribution, the computed correlators simply need to be multiplied by the charge factor, $C_{\text{em}} = (2/3)^2 + (-1/3)^2 + (-1/3)^2 = 2/3$, to obtain the electromagnetic current correlator in the physical QGP.

B. Hydrodynamics

The long-wavelength behavior of the spectral function can be studied with the help of hydrodynamics. Let D be

the diffusion coefficient for the conserved electric charges. In the hydrodynamic regime $\omega, k \ll D^{-1}$, the first-order hydrodynamic prediction for the transverse channel spectral function is [30]

$$\frac{\rho_T(\omega, k)}{\omega} \approx 2D\chi_s, \quad (8)$$

where χ_s denotes the static quark susceptibility:

$$\chi_s = \int_0^\beta d\tau \int d^3x \langle J_0(\tau, \mathbf{x}) J_0(0, \mathbf{0}) \rangle. \quad (9)$$

The diffusion coefficient can be expressed using the electrical conductivity, σ , as $D = \sigma/\chi_s$.

The functional form of ρ_T in Eq. (8) reveals the advantage of investigating this channel—namely, that it does not couple to the diffusion pole, in contrast to ρ_V or $\rho(\omega, k, -2)$ [23,24]. Therefore, $\rho_T(\omega, k)/\omega$ around $\omega = 0$ does not contain any peaklike structure of arbitrarily small width¹ as $k \rightarrow 0$. At extremely high temperatures in QCD, which implies a small value of the strong coupling constant g , a kinetic theory treatment eventually becomes applicable and a narrow peak of width $\sim g^4 T$ appears [30].

Following Ref. [23], we define the effective diffusion coefficient

$$D_{\text{eff}}(k) \equiv \frac{\rho_T(\omega = k, k)}{2\chi_s k}, \quad (10)$$

which tends to D as $k \rightarrow 0$ and is the key dynamical ingredient to evaluate the thermal photon production rate of Eq. (2).

C. Resummed spectral functions at NLO in thermal QCD

Analytic predictions are also available for the spectral functions in the weak coupling limit. The spectral function in the timelike regime—which is relevant for dilepton production—has been investigated using perturbative calculations since the seminal works of Refs. [27,31]. In most of these perturbative calculations, the studied channel was the vector channel spectral function, ρ_V . Focusing on the transverse channel, leading-order (LO) results (corresponding to noninteracting quarks) are available for massless quarks in, e.g., Refs. [32–34]. Recently, this has been extended to NLO at $\mathcal{O}(g^2)$, calculating the perturbative contributions up to two loops. The details of this impressive two-loop calculation—valid both for spacelike and timelike virtualities—can be found in Ref. [35], while a comparison to lattice correlators has been presented in Ref. [36].

¹The only exception to this statement is for a plasma of strictly noninteracting particles, for which hydrodynamic predictions do not apply.

As we discussed in Sec. II A, the relevant information for the thermal photon production is coming from the spectral function determined at the light cone. The thermal photon emission rate vanishes for noninteracting quarks [33]. The NLO perturbative thermal photon emission rate can be determined either by evaluating the NLO spectral functions at the light cone or by making use of the computation in Ref. [11]. The $O(g^2)$ calculation of Ref. [11] has been extended to $O(g^3)$ by taking into account contributions from soft gluons [12].

The strict two-loop perturbative spectral function develops a logarithmic singularity at $\omega = k$ [35,36]. It originates from multiple rescatterings of a quark taking part off-shell in the inelastic annihilation process that produces a photon, or in bremsstrahlung [11,34,37–39]. This infrared (IR) singularity is also present in the NLO calculation of the real photon rate [11,37]. The effect is called the Landau-Pomeranchuk-Migdal (LPM) effect [11,37] and can be handled by implementing a proper resummation of ladder diagrams, called the LPM resummation [11,34,36,38].

In order to compare lattice and perturbation theory results, we used the publicly available implementation of the two-loop calculation presented in Ref. [35] and computed the LPM resummation based on Refs. [34,36]. In our implementation, we used a window function to restrict the LPM contribution to frequencies around the light cone. The NLO perturbative spectral function complemented by the LPM contribution is called NLO + LPM in the following. For the spatial momentum of $k = \pi T$, we display this spectral function in Fig. 5 using dashed lines.

D. $\mathcal{N} = 4$ supersymmetric Yang-Mills theory

In the $\mathcal{N} = 4$ supersymmetric Yang-Mills (SYM) theory, the thermal spectral functions can be obtained analytically not only in the weak coupling limit, but in the strong coupling (and large- N_c) regime as well [40]. The field content of the theory is an $SU(N_c)$ gauge field together with massless scalar and fermionic fields in the adjoint representation of the gauge group. In the limit of infinite 't Hooft coupling and an infinite number of colors, one can determine the spectral function by making use of the AdS/CFT correspondence and then numerically solving an ordinary differential equation. Our primary interest, the spectral function in the transverse channel, is a smooth function having a similar asymptotic behavior—proportional to ω^2 —at high frequencies to the transverse spectral function in thermal QCD. Although the spectral functions in SYM in the strong coupling limit cannot quantitatively describe those of thermal QCD, their qualitative features are nevertheless instructive and may well be of relevance at $T \simeq 250$ MeV. An example of such a spectral function at strong coupling is shown in Fig. 5 for a spatial momentum $k = \pi T$ with a solid line.

TABLE I. Overview of the $N_f = 2$ ensembles used in this study. Simulations were carried out at a fixed temperature of $T \approx 254$ MeV and a fixed aspect ratio $TL = 4$, where L is the spatial linear size of the lattice. The parameters given are the bare gauge coupling g_0 , the Wilson hopping parameter κ , the temporal size in units of the lattice spacing a , the number of configurations used N_{conf} , and the number of molecular dynamics time units (MDUs) separating these configurations. The number of point sources per configuration is 16 in all cases.

| Label | $6/g_0^2$ | κ | $1/(aT)$ | N_{conf} | $\frac{\text{MDUs}}{\text{conf}}$ |
|-------|-----------|----------|----------|-------------------|-----------------------------------|
| F7 | 5.3 | 0.13638 | 12 | 482 | 20 |
| O7 | 5.5 | 0.13671 | 16 | 305 | 20 |
| W7 | 5.685727 | 0.136684 | 20 | 1566 | 8 |
| X7 | 5.827160 | 0.136544 | 24 | 511 | 10 |

III. LATTICE COMPUTATION OF THE TRANSVERSE CORRELATORS

A. Ensembles and statistics

We employ dynamical $O(a)$ -improved Wilson fermions to simulate two degenerate flavors of quarks with an *in vacuo* pion mass around 270 MeV. The details of the lattice action can be found in Ref. [41] and references therein. The temperature is $T \simeq 250$ MeV, well above the transition temperature, estimated to be about $T_c \simeq 211$ MeV in $N_f = 2$ QCD [42]. We use the four ensembles listed in Table I, which are the same ones underlying the analysis of the $\lambda = -2$ channel in Ref. [24]. Both the number of configurations and the number of point sources are the same in this work as they were in Ref. [24]. Although the pion mass is larger than its physical value on these ensembles, we emphasize that quark-mass effects on the correlator are suppressed by $(m/T)^2$ in the chirally symmetric phase. For our two coarsest ensembles, labeled as F7 and O7, the bare quark mass and gauge coupling have been set identical to the vacuum F7 and O7 ensembles used in Refs. [41,43], while for the finer ensembles, labeled as W7 and X7, the tuning to the line of constant physics was performed in Refs. [44]. The lattice spacings for these ensembles are around $a \simeq 0.066, 0.049, 0.039,$ and 0.033 fm with an error of about 1% [43]. The physical volume for all ensembles is around $L \simeq 3.1$ fm.

B. Lattice observables

In this section, we introduce the various discretized lattice correlators that we used to perform the continuum extrapolation in Sec. III C. We consider the two-point function of the isovector vector current in QCD with exact isospin symmetry. This allows for precise comparisons with weak coupling predictions in $N_f = 2$ QCD. As discussed at the end of Sec. II A, the correlator of the electromagnetic current in the physical QGP can be obtained approximately by multiplying our correlator by the charge factor $C_{\text{em}} = 2/3$.

The bare local and the conserved vector current are defined as

$$V_\mu^L(x) = \bar{\Psi}(x) \frac{\tau_3}{\sqrt{2}} \gamma_\mu \Psi(x) \quad (11)$$

and

$$V_\mu^C(x) = \frac{1}{2} \left[\bar{\Psi}(x + a\hat{\mu})(1 + \gamma_\mu) U_\mu^\dagger(x) \frac{\tau_3}{\sqrt{2}} \Psi(x) - \bar{\Psi}(x)(1 - \gamma_\mu) U_\mu(x) \frac{\tau_3}{\sqrt{2}} \Psi(x + a\hat{\mu}) \right], \quad (12)$$

respectively, where $\Psi = (u, d)^\top$ represents the isospin doublet of mass-degenerate quark fields and τ_3 is the diagonal Pauli matrix. As in Ref. [24], we have not implemented the additive $O(a)$ improvement of vector currents, as the contribution of the improvement terms to the two-point functions would be suppressed by the quark mass in the chirally restored phase. Using the currents above, we define the following bare unimproved correlators:

$$G_{\mu\nu}^{LL}(\tau, \mathbf{k}) = a^3 \sum_{\mathbf{x}} e^{i\mathbf{k}\cdot\mathbf{x}} \langle V_\mu^L(\tau, \mathbf{x}) V_\nu^L(0, \mathbf{0}) \rangle, \quad (13)$$

$$G_{\mu\nu}^{CC}(\tau + a\delta_{\mu 0}/2 - a\delta_{\nu 0}/2, \mathbf{k}) = a^3 \sum_{\mathbf{x}} e^{i\mathbf{k}\cdot(\mathbf{x} + a\hat{\mu}/2 - a\hat{\nu}/2)} \langle V_\mu^C(\tau, \mathbf{x}) V_\nu^C(0, \mathbf{0}) \rangle, \quad (14)$$

$$G_{\mu\nu}^{LC}(\tau - a\delta_{\nu 0}/2, \mathbf{k}) = a^3 \sum_{\mathbf{x}} e^{i\mathbf{k}\cdot(\mathbf{x} - a\hat{\nu}/2)} \langle V_\mu^L(\tau, \mathbf{x}) V_\nu^C(0, \mathbf{0}) \rangle, \quad (15)$$

$$G_{\mu\nu}^{CL}(\tau + a\delta_{\mu 0}/2, \mathbf{k}) = a^3 \sum_{\mathbf{x}} e^{i\mathbf{k}\cdot(\mathbf{x} + a\hat{\mu}/2)} \langle V_\mu^C(\tau, \mathbf{x}) V_\nu^L(0, \mathbf{0}) \rangle. \quad (16)$$

Under time reflections, the local-conserved correlator is transformed into the conserved-local one, $G_{\mu\nu}^{CL}(\tau, \mathbf{k}) \xrightarrow{T} G_{\nu\mu}^{LC}(-\tau, \mathbf{k})$, and vice versa. Using this fact, we average the two appropriately, and we refer to it with the superscript LC in the following. For the transverse correlator we need only the spatial components of the correlators, which are defined on the lattice sites according to Eqs. (13)–(16). In the case of local-conserved or conserved-local correlators, however, we note that the charge-charge correlator can be evaluated on site τ by averaging $G_{00}^{LC}(\tau + a/2, \mathbf{k})$ and $G_{00}^{LC}(\tau - a/2, \mathbf{k})$.

To obtain the correlator in the transverse channel, we first evaluate

$$\hat{G}_T^\alpha(\tau, \mathbf{k}) = \frac{1}{2} \sum_{i,j=1}^3 \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) G_{ij}^\alpha(\tau, \mathbf{k}), \quad (17)$$

$$\alpha = LL, LC, CC.$$

Imposing time-reversal symmetry and translation invariance, we then symmetrize the correlators in the time direction and average over the momentum orientations:

$$G_T^\alpha(\tau, k) = \frac{1}{2N_k} \sum_{|\mathbf{k}'|=k} (\hat{G}_T^\alpha(\tau, \mathbf{k}') + \hat{G}_T^\alpha(\beta - \tau, \mathbf{k}')), \quad (18)$$

where N_k is the number of momenta of norm k .

C. Continuum extrapolation

For the continuum extrapolation, we first interpolate the lattice correlators to the time separations which correspond to our finest lattice, X7. We apply two interpolation methods, Akima and monotonic cubic spline [45–47]. For the interpolation, we normalize G_T/G_{00} by the LO continuum transverse correlator—i.e., we multiply by T^3/G_T^{LO} in order to have a flattened interpolant. After the interpolation, we remove this factor.² In order to avoid the renormalization of the local currents, we divide the bare correlator by the bare static susceptibility computed using the same discretization:

$$\bar{G}_T^\alpha \equiv \frac{G_T^\alpha(\tau, k)}{\chi_s^\alpha(\tau) T}, \quad \chi_s^\alpha(\tau) = \beta G_{00}^\alpha(\tau, 0). \quad (19)$$

In the following, we omit the label T denoting the transverse channel.

When extrapolating to the continuum, our complete dataset consists of four ensembles, and on each ensemble three different discretizations of the correlator—i.e., 12 data points in total. Using these data points, we carry out fits by using only a single discretization, or simultaneously multiple discretizations of the transverse current correlators for each τ and k . When using multiple discretizations, we perform constrained correlated fits using the following fit *Ansatz*:

$$d(a/\beta, \mathbf{c}^\alpha) = c_0 + c_1^\alpha (a/\beta)^2 + c_2^\alpha (a/\beta)^4, \quad (20)$$

where c_0 is the estimate of the continuum limit from a particular fit, and the c_1^α and c_2^α parameters are characterizing the approach to the continuum of the different discretized correlators. We take into account the correlations between the different discretizations, and minimize the chi-squared statistic

²Alternatively, one can carry out the tree-level improvement of the data at this step—i.e., before the interpolation.

$$\chi^2 = \sum_{e=1}^{N_e} \sum_{\alpha, \alpha'} [\bar{G}^\alpha(a_e/\beta) - d(a_e/\beta, \mathbf{c}^\alpha)] \times \text{Cov}_{e, \alpha \alpha'}^{-1} [\bar{G}^{\alpha'}(a_e/\beta) - d(a_e/\beta, \mathbf{c}^{\alpha'})], \quad (21)$$

where the index e runs over the ensembles and $\alpha, \alpha' = \text{LL, LC, CC}$. A regularized covariance matrix Cov is obtained by multiplying the off-diagonal elements by $\xi = 0.95$. This one-parameter regularization method, which allows one to interpolate between fully correlated ($\xi = 1$) and uncorrelated fits ($\xi = 0$), is investigated in Refs. [48,49]. The regularization has a non-negligible effect only in the case of linear fits (i.e., when setting the c_2^2 coefficients to zero), for which it leads to an increased number of acceptable fits.

To increase the robustness of the continuum limit, we implement a multiplicative tree-level improvement of the lattice data

$$\bar{G}_{\text{TLI}}^\alpha(\tau, k) \equiv \bar{G}^\alpha(\tau, k) \left[\frac{\bar{G}(\tau, k)}{\bar{G}^\alpha(\tau, k)} \right]_{\text{LO}}, \quad (22)$$

where $\bar{G}_{\text{TLI}}^\alpha$ is the tree-level-improved correlator, and \bar{G}_{LO} and $\bar{G}_{\text{LO}}^\alpha$ are the continuum and the lattice leading-order perturbation theory results, respectively. The tree-level improvement reduces the difference between the various discretizations at finite lattice spacing, which results in more fits having acceptable p values. It reduces the continuum-extrapolated value at smaller Euclidean time separations, $\tau T < 0.25$. The effect of the improvement turns out to be much milder—almost negligible on the final continuum result—for larger distances, $\tau T \geq 0.25$.

In order to estimate the systematic errors, we carry out extrapolations using the tree-level-improved as well as the

unimproved data. Further systematic variations include using two interpolation methods to interpolate to the same τT points on all ensembles and varying the number of data points used in the fits. For the linear fits, only the three finest ensembles have been used; therefore, in this case we have nine data points and four fit parameters. We investigate the robustness of the results by omitting one data point from the coarsest ensemble or two data points from the coarsest two out of three ensembles. When we leave out more than one point, we always discard them from the same discretization. Furthermore, we carry out extrapolations by using only one or two discretizations. In the case of quadratic fits, we proceed similarly, but we omit one, two, or three data points from the coarsest three out of four ensembles.

These various changes in the analysis result in a total number of 104 fits. The χ^2 values, the number of fit parameters, and the number of data points have been used to calculate the Akaike information criterion (AIC) weight of each fit [50,51]. Using these weights, we build a histogram and quote the median as our final continuum result. The histogram is used in the later stages of the analysis.

We also perform the continuum limit of the static susceptibility, which requires including the Z_V renormalization constant when the local discretization of the current is used. To obtain the values of Z_V at the bare couplings of our ensembles, we utilize the parametrization of $Z_V(g_0^2)$ given in Ref. [52]. Then, adopting a similar procedure to determine the continuum limit of the static susceptibility as we used for the transverse correlator, we obtain $\chi_s/T^2 = \beta^3 G_{00}(\beta/2, 0) = 0.882(11)_{\text{stat}}(19)_{\text{sys}}$; see the right panel of Fig. 1. Compared with our result of Ref. [24], we quote a more conservative systematic error due to the continuum limit, primarily due to the inclusion of quadratic fits in a^2 in this work. We have also investigated the use of other




FIG. 1. *Left*: Simultaneous continuum extrapolation at $\tau T = 0.25$ of the local-local (LL), local-conserved (LC), and conserved-conserved (CC) tree-level improved correlators plotted with red circles, blue triangles, and green squares, respectively. The unimproved correlator values are shown with fainter symbols. The colored bands show the linear fit using the three finest lattice spacings and all discretizations, while the dashed lines illustrate the quadratic fit using all data points. The results of the linear (quadratic) extrapolations are shown as empty squares (circles). The fainter, shifted, gray symbols show the continuum values obtained using the unimproved correlators. The leftmost gray band represents our final continuum estimate, including its systematic error. *Right*: Simultaneous continuum extrapolation of the isovector charge-charge correlator required to compute the static susceptibility. The color code is the same as in the left panel.




FIG. 2. *Top panels:* Comparison of the G_T/G_{00} correlators for $k/T = \pi$ and $k/T \approx 4.97$. The bottom insets show the various correlators normalized by the continuum-extrapolated lattice-QCD correlator. The NLO + LPM result is obtained by using $\alpha_s = 0.25$. *Bottom panels:* Comparison of G_T/T^3 . We set $N_c = 3$ to obtain the SYM theory result.

determinations of the renormalization factor [53] and the inclusion of the mass-dependent improvement; these variations lead to subleading differences compared to the systematic error from the continuum extrapolation.

In order to obtain G_T/T^3 in the continuum, we multiply the continuum limit of G_T/G_{00} by the continuum estimate of G_{00}/T^3 . The transverse correlator divided by the charge-charge correlator is shown in the top panel of Fig. 2 for $k/T \approx 4.97$, and in units of temperature in the bottom panel. The statistical error on G_T/T^3 is typically around 0.25%–0.6%, while it is in the range 1.1%–1.4% for G_T/G_{00} .

D. Continuum-extrapolated correlators vs theory predictions

Having obtained the transverse-channel correlators in the continuum, we compare them to various theory predictions. In this comparison, we neglect the $O((m/T)^2)$ quark mass effects present in the lattice results, since $m/T \approx 0.05$ in our simulations.

We find that the LO correlator is about 5%–20% larger at Euclidean time separations $\tau T > 0.4$ for momenta $k/T \gtrsim 3.85$; see Fig. 2. The deviation is smaller, around $\sim 5\%$, for smaller momenta. It reduces towards smaller time separations, and below a certain (k -dependent) τT value,

the LO correlator is smaller by about 5% than our continuum result. The NLO + LPM correlator, however, is only about 3%–4% larger than the lattice result for all momenta and for all time separations $0.17 \lesssim \tau T \leq 0.5$ for which we could reliably determine the continuum limit. We conclude that the perturbative corrections to the LO correlator noticeably improve its agreement with our lattice correlator. The strongly coupled $\mathcal{N} = 4$ SYM theory result also provides a relatively good description, even though that result concerns a different non-Abelian gauge theory and applies in the large- N_c limit. The bottom insets in each of the four panels of Fig. 2 show the ratio of the perturbative and the SYM results to the continuum-extrapolated lattice results.

A word on the multiplicative normalization of the theory predictions is in order. For the perturbative predictions, while no prescription is needed for $G(\tau, k)$, a choice must be made to obtain $\tilde{G}(\tau, k)$. Here we have normalized both the LO and the NLO + LPM correlators by the $O(g^6 \ln g)$ susceptibility [54]. In Ref. [54], the coefficient of one of the undetermined $O(g^6)$ terms was estimated to be $C(N_f = 2) \approx -45$, which results in $\chi_{\text{pert}}/T^2 \approx 0.83$, the value which we used for Fig. 2. This value, however, is 6% lower than the continuum-extrapolated value we obtained. Based on our lattice continuum result, $C(N_f = 2) \approx 33$

would result in a better agreement for χ_s/T^2 . However, we note that the contributions from successive higher orders in the perturbative calculation of χ_{pert}/T^2 are similar in size, so the estimation for $C(N_f = 2)$ is to be treated with caution. As for the strongly coupled, large- N_c $\mathcal{N} = 4$ SYM correlator, we note that $\tilde{G}(\tau, k)$ is a natural quantity to compare across different thermal systems; in particular, the dependence on the number of colors drops out. In order to compare the correlator $G(\tau, k)$ itself, a certain choice must be made for the susceptibility. We have chosen to set $\chi_{\text{SYM}}/T^2 = N_c^2/8 = 9/8$.

IV. ANALYSIS OF THE TRANSVERSE-CHANNEL SPECTRAL FUNCTION

Given the transverse-channel Euclidean correlation function in the continuum limit, we proceed to analyze the corresponding spectral function via Eq. (6). Here, we present an analysis based directly on G_T/T^3 in Secs. IV A and IV B, since we found the correlator to be statistically particularly precise. In addition, we have seen that the NLO + LPM correlator is only a few percent off the lattice correlator, which encourages us to use that prediction as a baseline in Sec. IV B. In contrast, since the primary continuum-extrapolated quantity is $\tilde{G} = G_T/G_{00}$, Ref. [55] contains an analysis based on that ratio.

We start in Sec. IV A by using the Backus-Gilbert method in order to perform the comparison between lattice data and theory predictions in frequency space without introducing any model dependence. Section IV B then presents fits to the lattice data in order to determine the spectral function, with a particular focus on lightlike kinematics, $\omega = k$. For simplicity of notation, we omit the spatial momentum from the arguments in the following sections.

A. Backus-Gilbert method: Smeared spectral functions vs theory predictions

The Backus-Gilbert method is a model-independent approach to overcome the spectral reconstruction problem [25]. By using this method, one can determine a local average of the spectral function around a given value of the frequency. In the present case, it is also favorable to introduce a rescaling function, $f(\omega)$, which in particular removes the singularity at vanishing frequency of the kernel $K(\omega, \tau)$. Thus, the new kernel is defined as

$$K_f(\omega, \tau) \equiv K(\omega, \tau)f(\omega), \quad (23)$$

with the rescaling function $f(\omega)$ being specified later, but satisfying $f(\omega) \propto \omega$ as $\omega \rightarrow 0$. The smeared, rescaled spectral function, $\hat{\rho}(\omega)/f(\omega)$, is defined in this case as

$$\frac{\hat{\rho}(\omega)}{f(\omega)} \equiv \int_0^\infty d\omega' \Delta(\omega, \omega') \frac{\rho(\omega')}{f(\omega')}, \quad (24)$$

where $\Delta(\omega, \omega')$ is the so-called resolution function or averaging kernel, which is completely specified in terms of some coefficients, $g_i(\omega)$, and is given as

$$\Delta(\omega, \omega') = \sum_i g_i(\omega) K_f(\omega', \tau_i). \quad (25)$$

Inserting back $\Delta(\omega, \omega')$ of Eq. (25) into Eq. (24), we find that the filtered spectral function is the linear combination of the Euclidean correlator data,

$$\hat{\rho}(\omega) = f(\omega) \sum_i g_i(\omega) G_T(\tau_i). \quad (26)$$

The coefficients, $g_i(\omega)$, are determined in the Backus-Gilbert method by minimizing the second moment of the squared resolution function

$$\begin{aligned} A[g] &\equiv \int_0^\infty d\omega' (\omega - \omega')^2 [\Delta(\omega, \omega')]^2 \\ &= \sum_{i,j} g_i(\omega) A_{ij}(\omega) g_j(\omega) \end{aligned} \quad (27)$$

subject to the constraint

$$\int_0^\infty d\omega' \Delta(\omega, \omega') = 1. \quad (28)$$

This minimization ensures that the width of the resolution function is as small as possible and that, at fixed ω , $\Delta(\omega, \omega')$ has unit area. The minimizing solution is given as

$$g_i(\omega) = \frac{A_{ij}^{-1}(\omega) R_j}{R_k A_{kl}^{-1}(\omega) R_l}, \quad (29)$$

where

$$\begin{aligned} A_{ij}(\omega) &= \int_0^\infty d\omega' K_f(\omega', \tau_i) K_f(\omega', \tau_j) (\omega - \omega')^2 \quad \text{and} \\ R_i &= \int_0^\infty d\omega K_f(\omega, \tau_i). \end{aligned} \quad (30)$$

The matrix \mathbf{A} is ill-conditioned in practice; therefore, an error functional is added to $A[g]$, which serves as a regulator. As a consequence, \mathbf{A} has to be replaced by \mathbf{A}^{reg} in Eq. (29), where

$$A_{ij}^{\text{reg}}(\omega) = \lambda A_{ij}(\omega) + (1 - \lambda) \text{Cov}_{ij}[G_T], \quad (31)$$

and $0 \leq \lambda \leq 1$ is the regularization parameter compromising between stability and resolution. The smaller the value of λ , the larger the regularization. In Eq. (31), $\text{Cov}[G_T]$ stands for the covariance matrix of the Euclidean correlator. We note that in our numerical implementation, we work in the units of temperature.




FIG. 3. Comparison of filtered, rescaled, transverse channel spectral functions obtained using the continuum lattice data (with colors) and the perturbative spectral function (continuous, black), as well as the $\mathcal{N} = 4$ SYM theory (dashed, black) for $k/T \approx 3.14$ and $k/T \approx 4.97$ (left and right panels, respectively). The renormalization scale is set to $\mu = 2\pi T$ or $\mu = 3\pi T$ when using the NLO + LPM spectral function, which results in two curves. The renormalization scales $2\pi T$ and $3\pi T$ correspond to $\alpha_s \approx 0.31$ and $\alpha_s \approx 0.25$, respectively. Errors are only statistical.

We emphasize again that from the filtered spectral function one cannot model-independently determine the value of $\rho(\omega)$ itself, but it could be useful to compare it to a similarly smoothed spectral function coming from other approaches. Once the $g_i(\omega)$ coefficients are determined via Eq. (29) by utilizing the covariance matrix of the data, the same resolution function can be used to build, for instance, the perturbative filtered spectral function. In Fig. 3, we compare the filtered spectral function obtained from the continuum G_T/T^3 data to the filtered spectral function of the weak coupling QCD regime as well as that of the SYM theory, using the same resolution functions. The chosen rescaling function is

$$f(\omega) = \frac{\omega^2}{\tanh(\omega/(2T))}. \quad (32)$$

As Fig. 3 shows, the filtered, rescaled spectral function obtained using the lattice data is somewhat below the perturbative one, especially for small momenta ($k/T \leq 3.14$). At small frequencies, below $\omega/T \sim 5$, the filtered spectral function is between the weak coupling and the $\mathcal{N} = 4$ SYM theory filtered spectral function. Using all nine data points for the correlator, we find that the condition number of \mathbf{A}^{reg} gets more or less tractable (smaller than $\sim 10^7$) when λ is around 10^{-4} or smaller. The elements of \mathbf{A}^{reg} are still dominated in this case by the first term of Eq. (31), though they are still several orders of magnitude larger than the elements of the covariance matrix. When $\lambda \gtrsim 10^{-2}$, the condition number is above 5×10^8 and $\hat{\rho}(\omega)$ tends to have unphysical wiggles. By omitting, e.g., the first data point from the correlator, one can use larger values of λ , but one fewer coefficient. The results, nevertheless, do not change significantly.

The resolution functions, $\Delta(\omega, \omega')$, are quite wide, and they blur the fine details of the spectral function. They are almost identical for different momenta when fixing the value of λ . Their dependence on λ is also very mild, but it increases when going to a higher target ω . Some examples can be seen in Fig. 4. We obtain similar results to those discussed above using the covariance matrix of G_T/G_{00} instead.

In summary, we have found that the filtered spectral functions derived from the continuum-extrapolated lattice data are close to the NLO + LPM spectral function filtered with the same resolution function, and that the agreement increases both with increasing k and increasing ω , up to $\omega \approx 10T$. Since beyond that point the resolution function extends to very high frequencies, residual cutoff effects in the lattice results could explain the differences observed with the NLO + LPM curves in that regime. At $k = \pi T$, the nonperturbative filtered spectral function undershoots the




FIG. 4. Resolution functions for $k/T \approx 3.14$, $\lambda = 10^{-4}$ for different values of ω_*/T .




FIG. 5. *Left*: Representative fit results using the piecewise polynomial *Ansatz* with two ($N_p = 2$) or three fit parameters ($N_p = 3$) at $k/T = \pi$. For comparison, the transverse channel spectral function in the weak coupling NLO + LPM theory, as well as in the strongly coupled SYM theory are also included. *Right*: Representative fit results using the polynomial *Ansatz*.

perturbative prediction, a point already noted in Ref. [23] for the $2\rho_T + \rho_L$ channel relevant for the dilepton rate. Finally, we remark that the value of $[\hat{\rho}(\omega, k)/f(\omega)]_{\omega=k}$ found here is very much in the ballpark of the values obtained for $[\rho(\omega, k)/f(\omega)]_{\omega=k}$ with various fit *Ansätze* for the spectral function (see Fig. 5 below), even though the resolution function is rather broad compared to k .

B. The transverse-channel spectral function from fits to the Euclidean correlators

We now turn to our direct attempt at determining the spectral function by fitting various *Ansätze* to the transverse correlator. When specifying possible *Ansätze* describing the transverse spectral function, we have taken into account the facts that it has to be odd in ω , i.e., $\rho(-\omega) = -\rho(\omega)$, as well as positive for $\omega > 0$, as shown in Ref. [24]. While the *Ansätze* were chosen to be odd by construction, models were only excluded *a posteriori* if they violated positivity with a 68% confidence level.

The high-frequency behavior of the spectral function is dictated by perturbation theory; therefore, we assume the following form for the transverse spectral function:

$$\rho(\omega) = \rho_{\text{fit}}(\omega)(1 - \Theta(\omega, \omega_0, \Delta)) + \rho_{\text{pert}}(\omega)\Theta(\omega, \omega_0, \Delta), \quad (33)$$

where ρ_{pert} is the NLO prediction for the transverse spectral function complemented with the LPM contribution near the light cone,

$$\rho_{\text{pert}}(\omega) = \rho_{\text{NLO}}(\omega) + \rho_{\text{LPM}}(\omega), \quad (34)$$

and

$$\Theta(\omega, \omega_0, \Delta) = (1 + \tanh[(\omega - \omega_0)/\Delta])/2 \quad (35)$$

is a smooth step function which controls how fast the perturbative contribution falls off around ω_0 as ω is lowered. Using the decomposition of Eq. (33), we ensure that the perturbative part gives the dominant contribution above the chosen value of ω_0 , which we call the matching frequency. Moreover, the transition from the perturbative regime—assumed to be valid in the ultraviolet—can be realized in a smooth way without constraining any of the coefficients of the fit function.

For the fit function, $\rho_{\text{fit}}(\omega)$, we consider the following two possible *Ansätze*:

$$\frac{\rho_{\text{fit},1}(\omega)}{T^2} = \sum_{n=0}^{N_p-1} A_n \left(\frac{\omega}{\omega_0}\right)^{1+2n}, \quad (36)$$

where N_p denotes the number of fit parameters, and

$$\frac{\rho_{\text{fit},2}(\omega)}{T^2} = \begin{cases} A_0 \frac{\omega}{\omega_0} + A_1 \left(\frac{\omega}{\omega_0}\right)^3, & \text{if } \omega \leq k, \\ B_0 \frac{\omega}{\omega_0} + B_1 \left(\frac{\omega}{\omega_0}\right)^3, & \text{if } \omega > k, \end{cases} \quad (37)$$

where the free parameters have been chosen to be A_0 , B_0 , B_1 , and A_1 has been fixed, imposing continuity: $A_1 = B_1 + (B_0 - A_0)\omega_0^2/k^2$. In the latter case, we also carry out fits by setting B_0 to zero—i.e., having only two fit parameters. The spectral functions of Eq. (33) including $\rho_{\text{fit},1}(\omega)$ or $\rho_{\text{fit},2}(\omega)$ are referred to in the following as the polynomial or the piecewise polynomial *Ansätze*, respectively.

These *Ansätze* are motivated by their ability to describe the spectral function of the strongly coupled $\mathcal{N} = 4$ SYM theory, as well as that of the NLO + LPM resummed perturbation theory, to a satisfactory level. The functional form of Eq. (36) is more suitable for the spectral function obtained with the AdS/CFT approach in the $\mathcal{N} = 4$ SYM theory, whereas the built-in nondifferentiability at the light cone in the piecewise polynomial *Ansatz* of Eq. (37) is




FIG. 6. Results for the effective diffusion coefficient, TD_{eff} , defined in Eq. (10). Left: Separate analyses of fit results obtained with $N_p = 2$ and $N_p = 3$ using the piecewise polynomial Ansatz [Eq. (37)]. Right: Imposing the constraint of Eq. (38) at 1 standard deviation to the fit results obtained with the piecewise polynomial Ansatz. The results obtained by using the polynomial Ansatz [Eq. (36)] are displayed in red. Analytical results from perturbative QCD [11] using $\alpha_s \simeq 0.25$ or 0.31 (dashed lines) and from the strongly coupled $\mathcal{N} = 4$ SYM theory (gray line) [40] are also included, as well as an upper bound obtained from the analysis of the T – L channel (dotted line) [24]. We use the value $\chi_s/T^2 = 0.88(2)$ to obtain TD_{eff} from the lattice data, but the free susceptibility to obtain TD_{eff} from the $N_f = 2$ weak coupling photon rate.

more apt at describing the cusp present in the NLO + LPM result. More details can be found about the expressivity of these Ansätze in Appendix B, in which we present the outcome of mock data analyses and to which we return in the next subsection.

After inserting $\rho(\omega)$ from Eq. (33) into Eq. (6), we solve the correlated χ^2 minimization problem to determine the unknown coefficients. We show some representative fit results with good χ^2 values in Fig. 5 for the momentum $k = \pi T$. We have explored many variations in the fit procedure and take into account the statistical and systematic error of the continuum-extrapolated Euclidean correlators. More details on this and the method of estimating systematic errors on the effective diffusion coefficient, $D_{\text{eff}}(k)$, can be found in Appendix A. Our result for $D_{\text{eff}}(k)$ extracted using the polynomial Ansatz is displayed in the right panel of Fig. 6. The line within the band represents the median of the distribution of results obtained, while the width of the band indicates the positions of the 16th and 84th percentiles.

The piecewise polynomial fit Ansatz turns out to yield a sizeable spread of results for the effective diffusion coefficient. This spread comes from the results for D_{eff} actually falling into two well-separated intervals. Fit results with $N_p = 2$ tend to lead to results in the lower interval, while the results obtained with $N_p = 3$ populate both intervals. To illustrate the point, in the left panel of Fig. 6 we display the results for the effective diffusion coefficient $D_{\text{eff}}(k)$ obtained with $N_p = 2$ and $N_p = 3$ separately as two colored bands. This doubly peaked distribution of results for D_{eff} is associated with the behavior of the spectral function around lightlike kinematics. When using this Ansatz to fit the correlator, we obtain spectral functions

possessing either a minimum or a spikelike maximum at the light-cone frequency. Representative fit results are shown in the left panel of Fig. 5. Since neither the NLO + LPM weak coupling nor the SYM spectral functions has a maximum at $\omega = k$, we also investigate the fit results obtained by excluding the fits satisfying

$$\left. \frac{d\rho}{d\omega} \right|_{\omega=k-\epsilon} - \left. \frac{d\rho}{d\omega} \right|_{\omega=k+\epsilon} > 0 \quad (38)$$

to at least 1 standard deviation. With this qualitative theoretical prejudice in place, the fit results are significantly more constraining. The D_{eff} values obtained this way are displayed in the right panel of Fig. 6, where they are denoted as “no-peak” solutions. Excluding fits possessing the feature Eq. (38) is also affirmed by the upper limit of the results obtained by analyzing the spectral function in the T – L channel [24,29]. As can be seen by comparing the two panels of Fig. 6, a large portion of the solutions with a peak at lightlike kinematics can be excluded by our previous analysis of the T – L channel. We note that the latter analysis has been carried out using the same ensembles that we employ in the present study. For comparison, Fig. 6 also displays the weak coupling results (dashed lines) obtained directly for the photon rate in Ref. [11], as well as the strongly coupled SYM theory results (solid gray line).

C. Final result for the photon emissivity extracted from the fits

In order to arrive at our final estimate for the photon emissivity, we need to judge the reliability of our fit Ansätze in extracting the quantity D_{eff} . For this, we return to the analysis of mock data, presented in detail in Appendix B.

There, we find that the fit *Ansatz* functions tend to overestimate the photon rate in the investigated models, irrespective of whether G_T/G_{00} or G_T/T^3 is used. We therefore first derive an upper bound for D_{eff} , and hence for the photon emissivity. By fitting a linear *Ansatz* $a \times k/T + b$ to TD_{eff} obtained from the polynomial *Ansatz* in the k/T range $[\pi/2, 2\pi]$ and then plugging the effective diffusion coefficient, TD_{eff} , into the formula giving the thermal photon emission rate per unit volume, Eq. (2), we arrive at the following upper bound:

$$\frac{1}{T} \frac{d\Gamma_\gamma(k)}{d^3k} \leq \frac{\alpha_{\text{em}} \chi_s}{\pi^2 T^2} \frac{C_{\text{em}}}{e^{k/T} - 1} \times \left(0.22(1) - 0.0062(25) \frac{k}{T} \right). \quad (39)$$

In Eq. (39), C_{em} denotes the charge factor equal to $2/3$ in the $N_f = 3$ theory. We recall that the static susceptibility has been determined in Sec. III C: $\chi_s/T^2 = 0.88(2)$ at $T \simeq 250$ MeV.

As far as a lower bound on D_{eff} is concerned, we distinguish two regimes, $k > \pi T$ and $k \leq \pi T$. In the former case, using the NLO + LPM correlator as input, the central value of the output for the effective diffusion constant is usually at least 50%–80% larger than the true value, and our error estimate typically does not cover the true value in the range $3.5 \leq k/T \leq 5.5$ (see the right panel of Fig. 9). In the case of the $\mathcal{N} = 4$ SYM theory mock data, the results from the two *Ansätze* bracket the true result for intermediate and smaller momenta $k/T \lesssim 4.5$, but our *Ansätze* overestimate TD_{eff} by 5%–25% at larger momenta. Therefore, for $k > \pi T$ we extend the lower bound down to the weak coupling prediction in Fig. 6 (right panel), which we parametrize in Eq. (40).

For the mock data tests performed in the momentum range $k \leq \pi T$, on the other hand, the resulting error band always covers the true value for at least one of the two *Ansätze* that we use. As the right panel of Fig. 6 shows, the lower bound is driven by the results of the piecewise polynomial *Ansatz*, which decreases until reaching vanishing D_{eff} values at our smallest momentum. A parametrization of the lower bound can be given by a linear function in k/T that connects zero at $k = \pi T/2$ and the NLO weak coupling result at $k = \pi T$. Thus, we parametrize our lower bound according to

$$\frac{1}{T} \frac{d\Gamma_\gamma(k)}{d^3k} \geq \frac{\alpha_{\text{em}} \chi_s}{\pi^2 T^2} \frac{C_{\text{em}}}{e^{k/T} - 1} \times \begin{cases} (-9.43 + 6.00 \frac{k}{T}) \times 10^{-2}, & \pi T/2 \leq k \leq \pi T; \\ (4.86 + 14.33 \frac{T}{k}) \times 10^{-2}, & \pi T < k \leq 2\pi T. \end{cases} \quad (40)$$

From the observations made above, it is clear that this lower bound is influenced by the theory prejudice that the spectral




FIG. 7. The photon rate per unit volume of the QGP at $T \simeq 250$ MeV (hatched band). The Bose factor $n_B = 1/(e^{k/T} - 1)$ has been divided out. The upper bound parametrization derived from the D_{eff} results of the polynomial *Ansatz* is shown as a solid line. The full leading-order results [11] obtained by setting $\alpha_s \simeq 0.25$ or 0.31 are shown for comparison. Including corrections from interactions with soft gluons, one can get a 20% increase at $\alpha_s \simeq 0.3$ for the perturbative result [12].

function at lightlike kinematics does not “dive” down to even smaller values than the weak coupling results [11] predict for realistic values of α_s ; see Fig. 5.

We remark that the curves and bands displayed in Fig. 6 have a precise statistical meaning based on percentiles (16th, 50th, 84th) of the distribution of results obtained by applying a set of procedural variations. The band defined by the k -dependent bounds of Eqs. (39) and (40) summarizes the results, giving equal weight to both fit *Ansätze* that we have used; it is displayed in Fig. 7.

V. CONCLUSION

In this study, we have presented the first investigation of the transverse channel Euclidean correlator at finite momenta, using $N_f = 2$ $O(a)$ -improved dynamical Wilson fermions at around $T \simeq 250$ MeV corresponding to $1.2T_c$. We carried out a simultaneous continuum extrapolation using three discretizations of the correlators of the isovector vector currents. We used four ensembles with lattice spacings in the range $a \simeq 0.033$ – 0.066 fm. We compared the filtered spectral functions obtained from the continuum-extrapolated transverse correlator via the Backus-Gilbert method to the spectral function of perturbation theory at next-to-leading order and the one obtained in the strongly coupled $\mathcal{N} = 4$ supersymmetric Yang-Mills theory. The small-frequency lattice results lie between the filtered spectral functions obtained in these two theories. In order to determine the thermal photon emission rate, we fitted the correlators using polynomial and piecewise polynomial fit *Ansätze* for the underlying spectral function. We validated the expressivity of these *Ansätze* by performing mock tests using the spectral functions obtained in perturbation theory

as well as in the $\mathcal{N} = 4$ super-Yang-Mills theory. We compared our results for the effective diffusion coefficient to the results obtained in these theories as shown in Fig. 6, and also to the results obtained by analyzing the differences of the transverse and the longitudinal channels, presented earlier in Ref. [24]. The ranges obtained are compatible with the weak coupling, as well as with the AdS/CFT results. Our final result for the photon emissivity is displayed in Fig. 7 as a hatched band, for which we provide a parametrization in Eqs. (39) and (40). For momenta below 1 GeV, the upper edge of the band is more constraining than the estimate derived from the analysis of the difference of the transverse and longitudinal channels. We obtain our strongest constraint on the photon emissivity around $k = \pi T \simeq 0.8$ GeV:

$$\frac{d\Gamma_\gamma}{d^3k} = \frac{\alpha_{\text{em}}}{e^{k/T} - 1} (2.2 \pm 0.8) \times 10^{-3} \text{ GeV}. \quad (41)$$

Overall, our lattice-QCD based results, while consistent within uncertainties with the full leading-order weak coupling result [11], can also accommodate a photon emissivity about 2.5 times higher. Such an enhanced emission rate would likely help explain the large photon yield measured by the PHENIX [6] and ALICE [8] Collaborations in heavy-ion collisions, but may lead to a further suppression of the predictions [10] for the momentum anisotropy of the photons, which already undershoot the experimental data.

In the future, we plan to (linearly) combine the present analysis with our previous one [24] to provide estimates of the dilepton rate at invariant masses, $M_{e^+e^-}^2 = \omega^2 - k^2 > 0$. We are also investigating a qualitatively different approach to the photon rate by computing zero-virtuality correlators on the lattice [56,57]. While numerically challenging, this approach allows one to avoid confronting the inverse problem.

ACKNOWLEDGMENTS

We thank Arianna Toniato for discussions and valuable input in the early stages of this project. This work was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program through Grant Agreement No. 771971-SIMDAMA, as well as by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through the Cluster of Excellence "Precision Physics, Fundamental Interactions and Structure of Matter" (PRISMA+ EXC 2118/1) funded by the DFG within the German Excellence strategy (Project ID No. 39083149). M. C. was supported by the European Union's Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 843134-multiQCD. T. H. is supported by UK STFC CG No. ST/P000630/1. The generation of gauge configurations, as well as the computation of

correlators, was performed on the Clover and Himster2 platforms at Helmholtz-Institut Mainz and on Mogon II at Johannes Gutenberg University Mainz. We have also benefitted from computing resources at Forschungszentrum Jülich allocated under NIC Project No. HMZ21. We made use of the following libraries: GNU Scientific Library [47], GNU MPFR [58], and GNU MP [59].

APPENDIX A: TECHNICAL ASPECTS OF THE FITS FOR THE SPECTRAL FUNCTION

In this appendix, we provide some details on the procedure we followed in Sec. IV B to obtain spectral functions via fits to the Euclidean correlators. After inserting $\rho(\omega)$ from Eq. (33) into Eq. (6), we solved the correlated χ^2 minimization problem to determine the unknown coefficients. When doing so, we regularized the covariance matrix by multiplying the off-diagonal elements by 0.95. To estimate the statistical error, the minimization problem has been solved for every jackknife sample of the continuum correlator by using the covariance matrix obtained from the full data. To quantify the systematic uncertainty of the reconstruction coming from the systematic uncertainty of the continuum limit correlator values, we used the continuum extrapolation fit results for the 16th, 50th, and 84th percentiles of the AIC-weighted histogram obtained for each τT value. However, we did not take into account all possible combinations of these histogram representatives, but chose random subsets containing 20–100 combinations for each momentum.

In order to estimate the systematic error from making parameter choices for the perturbative results or changing various parameters when fitting, we performed several fits using a set of plausible variants. Regarding the perturbative input for our analysis, the NLO + LPM result depends on the coupling constant. Using the four-loop formulas of Ref. [60], we determined the coupling constant by setting the renormalization scale either to $\mu = 2\pi T$ or to $\mu = 3\pi T$. The coupling constants corresponding to these choices are $\alpha_s \simeq 0.31$ and $\alpha_s \simeq 0.25$, respectively. We took $\Lambda_{\overline{\text{MS}}}^{(N_f=2)}$ from the FLAG report [61].

We applied two values for the matching frequency, $\omega_0 = 10T$ and $\omega_0 = 12T$, which correspond to 2.5 GeV and 3 GeV, respectively. For the parameter Δ in $\Theta(\omega, \omega_0, \Delta)$, which governs the falloff of the NLO + LPM contribution towards the infrared, we set $\Delta = 2T$, allowing also for a 20% variation. As a further systematic variation, we adjusted the fit ranges in τT to include more or fewer data points in the fits. The variations we applied for the systematic error estimation are summarized in Table II. Due to these variations, we collected 750–2000 fits with p values larger than 0.05 for each momentum when using the polynomial *Ansatz*, and around 15%–30% of these fits had p values larger than 0.5. When using the piecewise polynomial *Ansatz*, we also obtained a lot of fits with good χ^2 (and p values). The fraction

TABLE II. Summary of the systematic variations we employed in the fit approach. Using nine (six) data points corresponds to fitting from $\tau T \approx 0.167$ (0.292).

| Source of systematic error | Variations |
|-------------------------------|-----------------------|
| Number of correlator data pts | 9, 8, 7, 6 |
| Number of fit parameters | 2, 3 |
| Soft: Δ/T | 1.6, 2.0, 2.4 |
| ω_0/T | 10, 12 |
| Renormalization scale | $(2, 3) \times \pi T$ |

of the number of fits with p values greater than 0.5 was around 15%–30% in that case as well. We built an AIC-weighted histogram from the fit results that we used to estimate the systematic errors [50,51].

APPENDIX B: MOCK ANALYSES

We performed mock analyses to see whether the fit *Ansätze* are expressive enough to reproduce the transverse channel spectral function as well as to investigate the reliability of the Backus-Gilbert method. For these tests, we used two models:

- (i) The NLO weak coupling spectral function complemented with the LPM contribution near the light cone.
- (ii) The spectral function of the strongly coupled $\mathcal{N} = 4$ super-Yang-Mills (SYM) theory.

We generated mock correlators using an appropriately rescaled covariance matrix of the lattice covariance matrix, which was obtained with the help of the continuum-extrapolated correlator and the jackknife samples.

The procedure we followed consists of the following steps:

- (1) Calculation of the ratio, $r(\tau)$, of the model and the continuum lattice correlator, $r(\tau) := G_{\text{model}}(\tau)/G(\tau)$.
- (2) Rescaling the covariance matrix of G in the continuum: $\text{Cov}_{\text{resc}, \tau\tau'} := \text{Cov}_{\tau\tau'} r(\tau)r(\tau')$.
- (3) Generating multivariate Gaussian variables, $G_{\text{mock}}(\tau)$, using the rescaled covariance matrix, Cov_{resc} , as well as $G_{\text{model}}(\tau)$.
- (4) Determination of the ratio $\bar{r}(\tau)$ of the mock correlator, $G_{\text{mock}}(\tau)$, and the lattice continuum correlator.
- (5) Rescaling all jackknife samples using $\bar{r}(\tau)$ and calculating the covariance matrix of the mock correlator (Cov_{mock}) using these rescaled values.

We then used $G_{\text{mock}}(\tau)$ and Cov_{mock} in the fit analysis. The mock jackknife samples have been used to estimate the statistical error in the mock analysis. The relative errors of the mock correlator are roughly the same as the relative errors of the continuum-extrapolated correlator due to the rescaling. We found that applying only step 1 and step 2 of the procedure above, complemented with a simple rescaling of the jackknife samples with $r(\tau)$, results in the overestimation of errors.

In addition to using the same relative errors as the continuum lattice data, we also investigated the effect of

reducing the errors on the mock correlator and therefore included slight modifications to the above procedure. Introducing the error reduction factor, $s > 1$, we used $r(\tau)/s$ in step 2, and instead of a rescaling, we determined the j th jackknife mock correlator value as $\bar{G}_{\text{mock}}^{(j)}(\tau) = \bar{G}_{\text{mock}}(\tau) + (r(\tau)\bar{G}^{(j)}(\tau) - \bar{G}_{\text{mock}}(\tau))/s$ in step 5.

Since we extrapolated $\bar{G} = G_T/G_{00}$ to the continuum (see Sec. III C), we also carried out the mock analysis using this observable. When doing so, G , G_{mock} , and G_{model} should be replaced by \bar{G} , \bar{G}_{mock} , and \bar{G}_{model} , respectively, in the above procedure, and we refer to this ratio of the chosen model when we have, e.g., \bar{G}_{model} . Since the covariance matrices are quite different for G_T/T^3 and for G_T/G_{00} , the mock analyses starting from the former or the latter result in different outcomes.

When using the NLO weak coupling mock correlator as an input for the reconstruction via fitting, we applied some variations in the fit setup, similar to the case of the analysis of the continuum-extrapolated correlator (Sec. IV B). These include using either two or three fit parameters in the fit *Ansätze*, changing the number of the correlator data points utilized in the fit, and changing the features (ω_0/T , Δ/T) of the matching to the UV behavior. Since when producing the mock correlator, we generated multivariate Gaussian variables randomly, to eliminate a possible effect coming from the random input, we used six different mock correlators generated with the procedure discussed above. The systematic error has been estimated using the AIC-weighted histogram of the various fit results, and we used the mock jackknife samples to estimate the statistical error of the mock analyses. The same fit *Ansatz* functions—Eqs. (36) and (37)—have been applied as in the main analysis.

We found that both fit *Ansätze* perform reasonably well in the weak coupling case when using the G_T/G_{00} data; see Fig. 8. Namely, above $k/T \approx \pi$, both *Ansätze* reproduce the input spectral functions within errors in a wide range of ω/T values also at small and large frequencies. The errors, however, are typically larger in this case for the polynomial *Ansatz* ($\sim 10\%$ – 50% , but could even be 100% at certain frequencies) than when using the G_T/T^3 data (errors $< 2\%$ – 8%). At $\omega = k$, however, the polynomial *Ansatz* gives a much larger value of ρ —i.e., a larger photon rate, using either G_T/G_{00} or G_T/T^3 . The piecewise polynomial *Ansatz* could reproduce the features of the NLO + LPM mock data better, because it is capable of producing a sharper dip at the light cone. With the help of this *Ansatz*, we get smaller photon rates at small momenta (below $k/T \approx \pi$), and larger photon rates at larger momenta, but in this latter case with errors spreading towards smaller values, typically covering the true spectral function at the light cone. Using the G_T/T^3 mock data, this *Ansatz*—similarly to the polynomial *Ansatz*—also returns a larger photon rate at all momenta—see, e.g., Fig. 9. In this




FIG. 8. Reconstructed spectral functions using the NLO + LPM weak coupling G_T/G_{00} mock correlator as an input.




FIG. 9. Reconstructed spectral functions using the NLO + LPM weak coupling G_T/T^3 mock correlator as an input.

case, the error covers the bottom of the dip only at and below $k/T = \pi$.

The reproduction of the mock spectral function can be improved by a certain amount by reducing the errors on the input mock correlators. As Fig. 10 shows, accommodating an error reduction factor of $s = 10$, the piecewise polynomial *Ansatz* could mimic the dip at $\omega = k$ at a more satisfactory level. For comparison, see the right panel of

Fig. 8 and also that of Fig. 9. This observation shows that this particular *Ansatz*, having only a few parameters, has satisfactory expressiveness for reproducing a model spectral function relevant to the physics discussed in this paper. Conversely, it also indicates that the fact that we obtained less faithful spectral function outcomes when not reducing the error is mainly due to the covariance matrix and not due to a wrong choice of *Ansätze*.




FIG. 10. Reconstructed spectral functions using the NLO + LPM correlator G_T/G_{00} (left), or G_T/T^3 (right) as an input. The covariance matrix elements have been decreased by a factor of $s^2 = 100$. Having such small errors, the piecewise polynomial *Ansatz* could better approximate the dip.




FIG. 11. Reconstructed spectral functions using the strongly coupled $\mathcal{N} = 4$ SYM theory mock correlator, G_T/G_{00} , as an input.




FIG. 12. Reconstructed spectral functions using the strongly coupled $\mathcal{N} = 4$ SYM theory mock correlator, G_T/T^3 , as an input.

Performing mock tests using the strongly coupled $\mathcal{N} = 4$ SYM theory transverse correlators, the polynomial *Ansatz* adequately reproduces the spectral function even for small momenta with errors less than around 5%–10% for frequencies around and above $\omega = k$ —see Figs. 11 and 12. As one goes for higher momenta, the reproduction gets a bit worse, but usually with errors included, one reaches the true values of the input spectral function. The photon rates given by this *Ansatz* are always higher than the true photon rate, either using the mock G_T/G_{00} or the G_T/T^3 data—see Figs. 11 and 12, respectively. The deviation increases from about 2%, corresponding to smaller momenta, up to 25%, for high momenta.

For this model, the piecewise polynomial *Ansatz* could also give reasonably good results for high momenta, where it almost coincides with the polynomial *Ansatz* results. For momenta, $k/T < 4.44$, it always gives a smaller photon rate, not depending on using G_T/G_{00} or G_T/T^3 , but in the latter case the central value is much closer to the true value.

The mock data also served as a numerical crosscheck for the Backus-Gilbert method, both using G_T/G_{00} as well as

G_T/T^3 . As Fig. 13 shows, the Backus-Gilbert method is capable of reproducing the smeared, rescaled spectral function up to around $\omega/T \sim 16$.




FIG. 13. Reconstructed, filtered spectral functions using the $\mathcal{N} = 4$ SYM mock correlator, $\bar{G}_T^{(\text{mock})} = G_T^{(\text{mock})}/G_{00}^{(\text{mock})}$, as an input. The results obtained with the help of the six randomized mock correlators are shown in orange without including the statistical errors. These errors are comparable to the errors of the reconstructed $\hat{\rho}$ using the exact model data.

- [1] E. L. Feinberg, *Nuovo Cimento A* **34**, 391 (1976).
- [2] E. V. Shuryak, *Phys. Lett.* **78B**, 150 (1978).
- [3] T. Ferbel and W. R. Molzon, *Rev. Mod. Phys.* **56**, 181 (1984).
- [4] W. Cassing and E. L. Bratkovskaya, *Phys. Rep.* **308**, 65 (1999).
- [5] G. David, *Rep. Prog. Phys.* **83**, 046301 (2020).
- [6] A. Adare *et al.* (PHENIX Collaboration), *Phys. Rev. C* **91**, 064904 (2015).
- [7] L. Adamczyk *et al.* (STAR Collaboration), *Phys. Lett. B* **770**, 451 (2017).
- [8] J. Adam *et al.* (ALICE Collaboration), *Phys. Lett. B* **754**, 235 (2016).
- [9] J.-F. Paquet, C. Shen, G. S. Denicol, M. Luzum, B. Schenke, S. Jeon, and C. Gale, *Phys. Rev. C* **93**, 044906 (2016).
- [10] C. Gale, J.-F. Paquet, B. Schenke, and C. Shen, *Phys. Rev. C* **105**, 014909 (2022).
- [11] P. B. Arnold, G. D. Moore, and L. G. Yaffe, *J. High Energy Phys.* **12** (2001) 009.
- [12] J. Ghiglieri, J. Hong, A. Kurkela, E. Lu, G. D. Moore, and Derek Teaney, *J. High Energy Phys.* **05** (2013) 010.
- [13] G. Aarts and J. M. Martinez Resco, *J. High Energy Phys.* **04** (2002) 053.
- [14] D. Teaney, *Phys. Rev. D* **74**, 045025 (2006).
- [15] H. B. Meyer, *Eur. Phys. J. A* **47**, 86 (2011).
- [16] S. Borsányi, Z. Fodor, M. Giordano, S. D. Katz, A. Pasztor, C. Ratti, A. Schäfer, K. K. Szabo, and B. C. Tóth, *Phys. Rev. D* **98**, 014512 (2018).
- [17] H.-T. Ding, A. Francis, O. Kaczmarek, F. Karsch, E. Laermann, and W. Soeldner, *Phys. Rev. D* **83**, 034504 (2011).
- [18] B. B. Brandt, A. Francis, H. B. Meyer, and H. Wittig, *J. High Energy Phys.* **03** (2013) 100.
- [19] A. Amato, G. Aarts, C. Allton, P. Giudice, S. Hands, and J.-I. Skullerud, *Phys. Rev. Lett.* **111**, 172001 (2013).
- [20] G. Aarts, C. Allton, A. Amato, P. Giudice, S. Hands, and J.-I. Skullerud, *J. High Energy Phys.* **02** (2015) 186.
- [21] B. B. Brandt, A. Francis, B. Jaeger, and H. B. Meyer, *Phys. Rev. D* **93**, 054510 (2016).
- [22] H.-T. Ding, O. Kaczmarek, and F. Meyer, *Phys. Rev. D* **94**, 034504 (2016).
- [23] J. Ghiglieri, O. Kaczmarek, M. Laine, and F. Meyer, *Phys. Rev. D* **94**, 016005 (2016).
- [24] M. Cè, T. Harris, H. B. Meyer, A. Steinberg, and A. Toniato, *Phys. Rev. D* **102**, 091501 (2020).
- [25] G. Backus and F. Gilbert, *Geophys. J. Int.* **16**, 169 (1968).
- [26] G. Backus and F. Gilbert, *Phil. Trans. R. Soc. A* **266**, 123 (1970).
- [27] L. D. McLerran and T. Toimela, *Phys. Rev. D* **31**, 545 (1985).
- [28] B. B. Brandt, A. Francis, T. Harris, H. B. Meyer, and A. Steinberg, *EPJ Web Conf.* **175**, 07044 (2018).
- [29] B. B. Brandt, M. Cè, A. Francis, T. Harris, H. B. Meyer, A. Steinberg, and A. Toniato, *Proc. Sci., LATTICE2019* (2019) 225.
- [30] J. Hong and D. Teaney, *Phys. Rev. C* **82**, 044908 (2010).
- [31] R. Baier, B. Pire, and D. Schiff, *Phys. Rev. D* **38**, 2814 (1988).
- [32] G. Aarts and J. M. Martinez Resco, *Nucl. Phys.* **B726**, 93 (2005).
- [33] M. Laine, *J. High Energy Phys.* **11** (2013) 120.
- [34] I. Ghisoiu and M. Laine, *J. High Energy Phys.* **10** (2014) 083.
- [35] G. Jackson, *Phys. Rev. D* **100**, 116019 (2019).
- [36] G. Jackson and M. Laine, *J. High Energy Phys.* **11** (2019) 144.
- [37] P. Aurenche, F. Gelis, and H. Zaraket, *Phys. Rev. D* **62**, 096012 (2000).
- [38] P. Aurenche, F. Gelis, G. D. Moore, and H. Zaraket, *J. High Energy Phys.* **12** (2002) 006.
- [39] M. E. Carrington, A. Gynther, and P. Aurenche, *Phys. Rev. D* **77**, 045035 (2008).
- [40] S. Caron-Huot, P. Kovtun, G. D. Moore, A. Starinets, and L. G. Yaffe, *J. High Energy Phys.* **12** (2006) 015.
- [41] P. Fritzsche, F. Knechtli, B. Leder, M. Marinkovic, S. Schaefer, R. Sommer, and F. Virota, *Nucl. Phys.* **B865**, 397 (2012).
- [42] B. B. Brandt, A. Francis, H. B. Meyer, O. Philipsen, D. Robaina, and H. Wittig, *J. High Energy Phys.* **12** (2016) 158.
- [43] G. P. Engel, L. Giusti, S. Lottini, and R. Sommer, *Phys. Rev. Lett.* **114**, 112001 (2015).
- [44] A. Steinberg, Ph.D. thesis, Mainz U., 2021.
- [45] H. Akima, *J. ACM* **17**, 589 (1970).
- [46] M. Steffen, *Astron. Astrophys.* **239**, 443 (1990).
- [47] M. Galassi *et al.* (2019), <https://www.gnu.org/software/gsl/>.
- [48] E. Rinaldi, S. Syritsyn, M. L. Wagman, M. I. Buchoff, C. Schroeder, and J. Wasem, *Phys. Rev. D* **99**, 074510 (2019).
- [49] A. Touloumis, *Computational Statistics and Data Analysis* **83**, 251 (2015).
- [50] H. Akaike, in *Proceedings of the 2nd International Symposium on Information Theory, Tsahkadsor, Armenia, USSR*, edited by B. N. Petrov and F. Csáki (Akadémiai Kiadó, Budapest, Hungary, 1971), pp. 267–281.
- [51] S. Borsanyi *et al.*, *Nature (London)* **593**, 51 (2021).
- [52] M. Della Morte, R. Hoffmann, F. Knechtli, R. Sommer, and U. Wolff, *J. High Energy Phys.* **07** (2005) 007.
- [53] M. Dalla Brida, T. Korzec, S. Sint, and P. Vilaseca, *Eur. Phys. J. C* **79**, 23 (2019).
- [54] A. Vuorinen, *Phys. Rev. D* **67**, 074032 (2003).
- [55] C. Török, M. Cè, T. Harris, A. Krasniqi, H. B. Meyer, and A. Toniato, *Proc. Sci., LATTICE2021* (2021) 172 [arXiv:2112.00563].
- [56] H. B. Meyer, *Eur. Phys. J. A* **54**, 192 (2018).
- [57] M. Cè, T. Harris, H. B. Meyer, A. Toniato, and C. Török, *Proc. Sci., LATTICE2021* (2022) 269 [arXiv:2112.00450].
- [58] L. Fousse, G. Hanrot, V. Lefèvre, P. Pélissier, and P. Zimmermann, *ACM Trans. Math. Softw.* **33**, 13-es (2007).
- [59] T. Granlund and the GMP development team (2020), <https://gmplib.org/>.
- [60] A. Deur, S. J. Brodsky, and G. F. de Teramond, *Prog. Part. Nucl. Phys.* **90**, 1 (2016).
- [61] S. Aoki *et al.* (Flavour Lattice Averaging Group), *Eur. Phys. J. C* **80**, 113 (2020).