

Effective field theory of Stückelberg vector bosons

Graham D. Kribs¹, Gabriel Lee^{2,3,4}, and Adam Martin⁵

¹*Institute for Fundamental Science and Department of Physics,
University of Oregon, Eugene, Oregon 97403 USA*

²*Department of Physics, LEPP, Cornell University, Ithaca, New York 14853, USA*

³*Department of Physics, Korea University, Seoul 136-713, Korea*

⁴*Department of Physics, University of Toronto, Toronto, Ontario, M5S 1A1 Canada*

⁵*Department of Physics, University of Notre Dame, South Bend, Indiana 46556, USA*

 (Received 17 May 2022; accepted 2 September 2022; published 16 September 2022)

We explore the effective field theory of a vector field X^μ that has a Stückelberg mass. The absence of a gauge symmetry for X^μ implies Lorentz-invariant operators are constructed directly from X^μ . Beyond the kinetic and mass terms, allowed interactions at the renormalizable level include $X_\mu X^\mu H^\dagger H$, $(X_\mu X^\mu)^2$, and $X_\mu j^\mu$, where j^μ is a global current of the SM or of a hidden sector. We show that all of these interactions lead to scattering amplitudes that grow with powers of \sqrt{s}/m_X , except for the case of $X_\mu j^\mu$ where j^μ is a *nonanomalous* global current. The latter is well known when X is identified as a dark photon coupled to the electromagnetic current, often written equivalently as kinetic mixing between X and the photon. The power counting for the energy growth of the scattering amplitudes is facilitated by isolating the longitudinal enhancement. We examine in detail the interaction with an *anomalous* global vector current $X_\mu j_{\text{anom}}^\mu$, carefully isolating the finite contribution to the fermion triangle diagram. We calculate the longitudinally-enhanced observables $Z \rightarrow X\gamma$ (when $m_X < m_Z$), $f\bar{f} \rightarrow X\gamma$, and $Z\gamma \rightarrow Z\gamma$ when X couples to the baryon number current. Introducing a “fake” gauge-invariance by writing $X^\mu = A^\mu - \partial^\mu \pi/m_X$, the would-be gauge anomaly associated with $A_\mu j_{\text{anom}}^\mu$ is canceled by $j_{\text{anom}}^\mu \partial_\mu \pi/m_X$; this is the four-dimensional Green-Schwarz anomaly-cancellation mechanism at work. Our analysis demonstrates there is a much larger set of possible interactions that an EFT with a Stückelberg vector field can have, revealing scattering amplitudes that grow with energy. The growth of these amplitudes can be tamed by a dark Higgs sector, but this requires dark Higgs boson interactions (and reintroduces fine-tuning in the dark Higgs sector) that can be separated from X interactions only in the limit $g \ll 1$.

DOI: [10.1103/PhysRevD.106.055020](https://doi.org/10.1103/PhysRevD.106.055020)

I. INTRODUCTION

New massive vector bosons are ubiquitous in beyond the Standard Model (SM) physics. At masses large compared with collider energies, they provide UV completions of higher dimensional operators [1]. At intermediate masses, of order collider energies, they yield resonances that are targeted by many searches [2]. At somewhat smaller masses, they can be produced, decay, and be observed in high intensity experiments [3–5], typically when coupled to charged leptons (for reviews, see [6,7]). Also at smaller masses, they can act as mediators to permit light dark matter to interact with the SM [8–10], underpinning the viability of a large class of light dark matter detection

experiments [7]. At exceptionally small masses, vector bosons can even serve as dark matter itself [11–18].

One of the attractions of a single new massive vector boson is that a simple model [19] exists: the massive $U(1)$ dark photon A^μ (see [20] for a review),

$$\mathcal{L}_{\text{dark } \gamma} = -\frac{1}{4} F_{A,\mu\nu} F_A^{\mu\nu} + \frac{1}{2} m_X^2 A_\mu A^\mu - \epsilon F_{A,\mu\nu} F_Y^{\mu\nu}, \quad (1)$$

that involves just two parameters m_X and ϵ , respectively the mass of the $U(1)$ dark photon and its kinetic mixing to hypercharge. The simplicity of this extension hinges on the existence of a Stückelberg mass (see [21] for a review) for the dark photon. In particular, by not specifying a Higgs mechanism for the dark photon, one is able to avoid the consideration of additional interactions of the dark Higgs field ϕ_X . In particular, one does not need to address the new fine tunings from the “dark hierarchy problem” that are inevitable with a dark Higgs field or how to avoid the respective destabilization of the dark and/or SM

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article’s title, journal citation, and DOI. Funded by SCOAP³.

Higgs sectors through renormalizable interactions such as $\phi_X^\dagger \phi_X H^\dagger H$.

One of the reasons the dark photon Lagrangian *seems* simple is how the longitudinal mode is packaged in A^μ . We can introduce the longitudinal mode π such that, under a gauge transformation $A^\mu \rightarrow A^\mu + \partial^\mu \alpha(x)$, the longitudinal mode shifts $\pi \rightarrow \pi + m_X \alpha(x)$. Using the equation of motion (EOM) for the hypercharge gauge boson, $\partial_\mu F_Y^{\mu\nu} = g_Y j_Y^\nu$ in terms of the SM hypercharge current j_Y^ν , the dark photon Lagrangian can be rewritten as:

$$\mathcal{L}_{\text{dark } \gamma} = -\frac{1}{4} F_{X,\mu\nu} F_X^{\mu\nu} + \frac{1}{2} m_X^2 X_\mu X^\mu - \epsilon g_Y X_\mu j_Y^\mu \quad (2)$$

in terms of $X^\mu \equiv A_X^\mu - \partial^\mu \pi / m_X$ —the Stückelberg vector field—a vector boson without a corresponding $U(1)$ gauge invariance. The lack of gauge invariance is obvious because X^μ remains invariant under the simultaneous gauge transformations of A^μ and π . This form of the dark photon Lagrangian makes it clear that a Lagrangian with a Stückelberg mass for a vector field is best expressed in terms of X^μ ; the use of the field strength $F_X^{\mu\nu}$ for the kinetic term (or kinetic mixing with the SM) has nothing to do with gauge invariance, and instead simply ensures there are only three propagating degrees of freedom (DOF) in X^μ .¹

This naturally leads to the question of the effective field theory involving a Stückelberg vector field X^μ —what are all possible interactions of X^μ , and what are their consequences? The goal of this paper is to show that the Lagrangian Eq. (2) is a special case of a more general set of interactions for X^μ . For instance, already at the renormalizable level we can write $(X_\mu X^\mu)^2$, $X_\mu X^\mu H^\dagger H$, and $X_\mu j^\mu$ where j^μ is a global vector or axial current that may or may not be (globally) anomaly-free.² As we will see, most of these interactions have couplings of the longitudinal mode with itself or the SM fields, and thus lead to scattering amplitudes that grow with powers \sqrt{s}/m_X . This is analogous to the energy growth that arise in a Higgsless SM [23]. The range of validity of the effective theory including X in the spectrum relies on taking the coefficients of longitudinally enhanced interactions to be (sometimes exceptionally) small. Only if there are *exactly zero* couplings of the longitudinal mode with itself or with the SM can the cutoff scale of the EFT be taken arbitrarily large relative to the mass of the Stückelberg vector field.

There is a host of related literature that we will only briefly mention. Numerous papers have studied theories with a Stückelberg vector field in the context of field theory or string theory [21,22,24–34]. There is also a huge

literature on anomalous $U(1)$ symmetries and their implications for theory or phenomenology [29,35–53]. The connections between anomalous $U(1)$ symmetries and the Green-Schwarz anomaly cancellation mechanism have also been elucidated [29,36,38,39,46,54]. While we have certainly benefited from this literature and we do not claim to be the first or last word on this subject, our focus on a theory with a Stückelberg mass for X^μ , a vector field without a corresponding gauge symmetry, lays a foundation for a systematic approach to analyze the effective field theory of X^μ in terms of its leading self-interactions as well as its interactions with the SM.

The organization of this paper is as follows. First, in Sec. II, we review the Stückelberg Lagrangian, (fake) gauge fixing, BRST, the external physical states, the propagator, and the BRST current. In Sec. III we consider tree-level interactions of the Stückelberg vector field X^μ . We demonstrate that self-couplings as well as tree-level couplings of the longitudinal mode with the SM lead to amplitudes that grow with energy above the mass of the Stückelberg vector field. While these interactions are not radiatively generated by a dark photon Lagrangian that consists solely of a mass term and a coupling to a conserved vector current, there are no symmetries that forbid these terms. Consequently, the dark photon Lagrangian appears rather peculiar. In particular, we show that these interactions can be generated by a dark Higgs mechanism for a dark $U(1)$ gauge theory, and like the Higgs mechanism of the SM, the dark Higgs boson renders the amplitudes finite above the dark Higgs mass. In Sec. IV, we consider the coupling of a Stückelberg vector field to an *anomalous* vector current. This is motivated by Dror *et al.* [45], who showed that should an anomalous symmetry of the SM (e.g., baryon number) be gauged, the couplings of the longitudinal mode lead to longitudinal enhancements of the amplitudes involving the anomalous fermion triangle diagram. These longitudinal enhancements are critical in determining the viable range of parameter space in the model [43]. The Stückelberg vector field theory would appear to be special, since there is no gauge symmetry, and thus, no gauge anomalies. Nevertheless, we carefully consider the one-loop triangle diagrams that arise because of an anomalous *global* symmetry of the SM. We find that the Stückelberg vector field has couplings of its longitudinal mode to the divergence of the anomalous global current. The observable predictions of a Stückelberg vector field coupled to, say, global baryon number of the SM are identical to the case in which baryon number is gauged, so long as the “anomalons” needed to cancel the gauge anomaly are taken to be heavy. In Sec. V, we demonstrate the importance of the one-loop couplings of the longitudinal part of X^μ to an anomalous global current for several physical processes, including $Z \rightarrow X\gamma$ and $f\bar{f} \rightarrow X\gamma$, and $Z\gamma \rightarrow Z\gamma$, when X couples to baryon number. Finally, in Sec. VI, we discuss the implications

¹Contrast this with a spin-one gauge field, such as hypercharge B^μ , which only appears in the field strength $F_B^{\mu\nu}$ and covariant derivatives.

²The phenomenological implications of the quartic interaction for the electromagnetic field was explored in [22].

of our results. The appendices contain technical details of calculations relevant for results in Secs. IV and V.

II. REVIEW OF QUANTIZATION OF MASSIVE VECTOR FIELDS

A. The Lagrangian and propagator for a massive spin-one field

A massive spin-one field X^μ has three propagating degrees of freedom (DOF). We see this by decomposing the four components of the four-vector X^μ into the $\mathbf{1} \oplus \mathbf{3}$, or spin-zero and spin-one, representations of the Lorentz group. The spin-zero component leads to a negative energy density, and can be removed as a propagating DOF in the theory by imposing the Lorenz condition

$$\partial_\nu X^\nu = 0, \quad (3)$$

together with writing the kinetic term for the four-vector as a function of the field-strength tensor $F_X^{\mu\nu} = \partial^\mu X^\nu - \partial^\nu X^\mu$ [55]. The above two requirements are achieved by the Proca Lagrangian

$$\mathcal{L}_P = -\frac{1}{4} F_{X,\mu\nu} F_X^{\mu\nu} + \frac{1}{2} m_X^2 X_\mu X^\mu, \quad (4)$$

which yields the EOM and its derivative

$$\begin{aligned} \partial_\mu F_X^{\mu\nu} + m_X^2 X^\nu &= 0, \\ m_X^2 \partial_\nu X^\nu &= 0. \end{aligned} \quad (5)$$

For $m_X \neq 0$, the Lorenz condition follows from the second line and therefore is not an independent constraint. The Proca Lagrangian for X^μ is not gauge invariant: there is no $U(1)$ symmetry associated with X^μ since there is no redundancy in its description—all three of its propagating DOF are physical.

The propagator for X^μ can be derived directly from inverting the Proca Lagrangian, which is textbook material [55,56]

$$\langle X^\mu(p) X^\nu(-p) \rangle = \frac{-i}{p^2 - m_X^2} \left(g^{\mu\nu} - \frac{p^\mu p^\nu}{m_X^2} \right). \quad (6)$$

The propagator for X^μ is equivalent to the propagator of a Higgsed, massive $U(1)$ theory in unitary gauge; however, we emphasize that the result above is *not* in unitary gauge—there is no gauge invariance. This also implies that the sum of the polarization states for an on-shell X^μ coincides with that of a massive $U(1)$ theory, i.e.,

$$\sum_\lambda \epsilon_\lambda^\mu(p) \epsilon_\lambda^{\nu*}(p) = -\left(g^{\mu\nu} - \frac{p^\mu p^\nu}{m_X^2} \right). \quad (7)$$

This explicitly demonstrates the counting of the on-shell physical DOF: X^μ has three physical polarizations.

B. Stückelberg formalism: Introducing a fake gauge symmetry

The Stückelberg formalism expresses

$$X^\mu \equiv A^\mu - \frac{\partial^\mu \pi}{m_X}, \quad (8)$$

where A^μ is a “fake” $U(1)$ gauge field and π is a scalar field that also transforms under this “fake” $U(1)$ gauge invariance:

$$\begin{aligned} A^\mu &\rightarrow A^\mu + \partial^\mu \alpha(x), \\ \pi &\rightarrow \pi + m_X \alpha(x), \end{aligned} \quad (9)$$

where $\alpha(x)$ is the gauge parameter. The Proca Lagrangian becomes

$$\mathcal{L}_g = -\frac{1}{4} F_{A,\mu\nu} F_A^{\mu\nu} + \frac{1}{2} m_X^2 \left(A_\mu - \frac{\partial_\mu \pi}{m_X} \right)^2, \quad (10)$$

purely in terms of the “fake” gauge field with its field strength given by $F_A^{\mu\nu}$. While this construction introduces one additional DOF π , the “fake” $U(1)$ gauge invariance removes one DOF, leaving the same three of the massive vector field in the original Proca Lagrangian [57,58].

We use the term “fake” to describe the gauge invariance of A^μ since the physical consequences of X^μ and its interactions can be determined entirely in terms of the vector-field X^μ directly. The identification $X^\mu \equiv A^\mu - \partial^\mu \pi / m_X$ is exact, in the sense that renormalization does not disrupt the size of the coefficient of $\partial^\mu \pi / m_X$ relative to A^μ . This follows from ensuring that the gauge transformations of A^μ and π leave the combination $A^\mu - \partial^\mu \pi / m_X$ invariant.

The purpose of introducing the “fake” gauge invariance is to more easily uncover the role of the longitudinal polarization of X^μ , namely X_L^μ , which for a suitable choice of gauge, can be fully captured by the interactions of the scalar field π . Hence, we will refer to π as the “longitudinal component” synonymously with X_L^μ , though we emphasize that this identification is only strictly true in Landau gauge, as we discuss below.

C. BRST and R_ξ gauge fixing

Before we discuss the gauge fixing of Eq. (10) and applying the BRST to the Stückelberg formalism, we briefly review the general gauge-fixing and quantization procedure using BRST [59,60]. The BRST transformations of the fields are equivalent to gauge transformations like those in Eq. (9) with infinitesimal gauge parameter

$$\alpha(x) = \theta\omega(x), \quad (11)$$

where θ is an infinitesimal Grassmann constant and ω is a real, Grassmann scalar field (ghost). For the Stückelberg theory, we have the following BRST transformations of the fields:

$$\begin{aligned} \delta_\theta A &= \theta\partial\omega, \\ \delta_\theta \pi &= m_X\theta\omega, \\ \delta_\theta b &= 0, \\ \delta_\theta \omega &= 0, \\ \delta_\theta \omega^* &= \theta b, \end{aligned} \quad (12)$$

where ω^* is a real, Grassmann scalar field (antighost) and b is a Nakanishi-Lautrup auxiliary field [61,62]. The action of a BRST operator \mathbf{s} on a field φ is defined in terms of the infinitesimal BRST transformation of a field φ by

$$\delta_\theta \varphi = \theta \mathbf{s} \varphi. \quad (13)$$

For a product of fields,

$$\delta_\theta(\varphi_1 \varphi_2) = (\delta_\theta \varphi_1) \varphi_2 + \varphi_1 (\delta_\theta \varphi_2) = \theta[(\mathbf{s}\varphi_1)\varphi_2 \pm \varphi_1(\mathbf{s}\varphi_2)], \quad (14)$$

where \pm for whether φ_1 is bosonic or fermionic; i.e., \mathbf{s} can be viewed as a fermionic operator. Using the transformations in Eq. (12), the gauge-fixing part of the Lagrangian can be written as [24]

$$\mathcal{L}_{\text{gf}} = \mathbf{s} \left[\omega^* \left(\mathcal{G} + \frac{\xi}{2} b \right) \right] = -\omega^*(\mathbf{s}\mathcal{G}) + b\mathcal{G} + \frac{\xi}{2} b^2, \quad (15)$$

where $\mathcal{G}[A, \pi]$ is a gauge-fixing functional. Since b is an auxiliary field and does not propagate, we can eliminate it using its EOM, yielding an alternate form for Eq. (15),

$$\mathcal{L}_{\text{gf}} = -\omega^*(\mathbf{s}\mathcal{G}) - \frac{1}{2\xi} \mathcal{G}^2. \quad (16)$$

The R_ξ -like class of gauge-fixing choices is obtained by setting

$$\mathcal{G}_\xi = \partial_\mu A^\mu + \xi m_X \pi. \quad (17)$$

The general R_ξ -gauge Lagrangian is the sum of Eq. (10) and the gauge-fixing terms,

$$\begin{aligned} \mathcal{L}_\xi &= \mathcal{L}_g + \mathcal{L}_{\text{gf}}|_{\mathcal{G}_\xi} \\ &= -\frac{1}{4} F_{A,\mu\nu} F_A^{\mu\nu} + \frac{1}{2} m_X^2 \left(A_\mu - \frac{\partial_\mu \pi}{m_X} \right)^2 - \frac{1}{2\xi} (\partial_\mu A^\mu + \xi m_X \pi)^2 - \omega^* (\partial^2 + \xi m_X^2) \omega \\ &= -\frac{1}{4} F_{A,\mu\nu} F_A^{\mu\nu} - \frac{1}{2\xi} (\partial_\mu A^\mu)^2 + \frac{1}{2} m_X A_\mu A^\mu + \frac{1}{2} \partial_\mu \pi \partial^\mu \pi - \frac{1}{2} \xi m_X^2 \pi^2 - \omega^* (\partial^2 + \xi m_X^2) \omega, \end{aligned} \quad (18)$$

which explicitly exhibits the decoupling of $A^\mu, \partial^\mu \pi$.³

From this, we see that the Proca Lagrangian corresponds to the choice $\xi \rightarrow 0$, where the second term in the last line of Eq. (18) decouples and π becomes a free, massless scalar field. The Stückelberg Lagrangian is obtained from the choice of Stückelberg-Feynman gauge $\xi = 1$,

$$\mathcal{L}_{\text{St}} = -\frac{1}{4} F_{A,\mu\nu} F_A^{\mu\nu} + \frac{1}{2} m_X^2 \left(A_\mu - \frac{\partial_\mu \pi}{m_X} \right)^2 - \frac{1}{2} (\partial^\mu A_\mu + m_X \pi)^2. \quad (19)$$

Note that the first two terms in Eq. (19) are unchanged under the gauge transformation Eq. (9); however,

³Using R_ξ gauge fixing, the ghosts decouple in Abelian gauge theories because the ghost kinetic term involves only partial derivatives (in Yang-Mills theories, these become covariant derivatives in the adjoint representation). Hence, we omit them from the Lagrangian for the remainder of the paper.

invariance of the last term requires π to obey the EOM for a massive scalar field,

$$(\square + m_X^2)\pi = 0. \quad (20)$$

D. Propagator in R_ξ gauge

The R_ξ gauge fixing removes the mixing terms of the form $A^\mu \partial_\mu \pi$ in the original Stückelberg Lagrangian of Eq. (19), leaving just the gauge-dependent two-point functions for A^μ and π . These have the standard R_ξ -gauge forms:

$$\begin{aligned} \langle A^\mu(p) A^\nu(-p) \rangle &= \frac{-i}{p^2 - m_X^2} \left(g^{\mu\nu} - \frac{p^\mu p^\nu}{p^2 - \xi m_X^2} (1 - \xi) \right), \\ \langle \pi(p) \pi(-p) \rangle &= \frac{i}{p^2 - \xi m_X^2}. \end{aligned} \quad (21)$$

Using Eq. (12) and the decomposition in Eq. (8), the BRST transformation of X^μ is

$$\delta_\theta X^\mu = \delta_\theta A^\mu - \frac{1}{m_X} \partial^\mu \delta_\theta \pi = \theta \partial^\mu \omega - \partial^\mu (\theta \omega) = 0. \quad (22)$$

X^μ is annihilated by the BRST operator and corresponds to a physical external state. From Eq. (21), the X^μ two-point function can be reconstructed as

$$\begin{aligned} \langle X^\mu(p) X^\nu(-p) \rangle &= \langle A^\mu(p) A^\nu(-p) \rangle + \frac{1}{m_X^2} (i p^\mu)(-i p^\nu) \langle \pi(p) \pi(-p) \rangle \\ &= \frac{-i}{p^2 - m_X^2} \left(g^{\mu\nu} - (1 - \xi) \frac{p^\mu p^\nu}{p^2 - \xi m_X^2} \right) + \frac{i}{m_X^2} \frac{p^\mu p^\nu}{p^2 - \xi m_X^2} \\ &= \frac{-i}{p^2 - m_X^2} \left(g^{\mu\nu} - \frac{p^\mu p^\nu}{m_X^2} \right), \end{aligned} \quad (23)$$

which agrees with Eq. (6). The absence of ξ -dependence demonstrates that the propagator for the physical state X^μ is, unsurprisingly, itself independent of the fake gauge symmetry.

E. Current conservation

The decomposition $X^\mu \equiv A^\mu - \partial^\mu \pi / m_X$ allows us to study Stückelberg theories using techniques familiar from gauge theories. In fact, the fake gauge field A^μ has the same form as that of a massive gauge field arising from a Higgsed $U(1)$ symmetry that is spontaneously broken with mass $m_X = gv/2$. However, for a Stückelberg vector field, we know that only the combination $A^\mu - \partial^\mu \pi / m_X$ is physical and can represent an external state, while for a gauge theory, the external state is of course just A^μ . How do we reconcile this difference?

To understand when there is a distinction between the Stückelberg vector field and a spontaneously broken massive gauge field A^μ , we examine the BRST current,

$$J_{\text{BRST}}^\mu = \sum_{\text{field } \varphi} \frac{\delta L}{\delta \partial_\mu \varphi} \delta_\theta \varphi. \quad (24)$$

To keep things simple, consider a scenario in which the spin-one fields have interactions with a fermion current, i.e., $g(A_\mu - \partial_\mu \pi / m_X) j_{\text{ferm}}^\mu \equiv g X_\mu j_{\text{ferm}}^\mu$ for a Stückelberg vector field and $g A_\mu j_{\text{ferm}}^\mu$ for a spontaneously broken, gauged $U(1)$ vector field.

In the case where A^μ is a gauge field that is spontaneously broken, it is straightforward to show that the divergence of the BRST current is

$$\partial_\mu J_{\text{BRST}}^\mu = -\omega \partial_\mu j_{\text{ferm}}^\mu \quad (\text{for a massive gauge field } A^\mu). \quad (25)$$

Therefore, a conserved BRST charge requires the divergence of the fermion current to vanish.

In contrast, when the same BRST transformations are applied to the Stückelberg vector field, we obtain

$$\begin{aligned} \partial_\mu J_{\text{BRST}}^\mu &= 0 \\ &(\text{for a Stückelberg vector field } X^\mu = A^\mu - \partial^\mu \pi / m_X). \end{aligned} \quad (26)$$

A conserved BRST charge can always be formed since the divergence of the BRST current vanishes independently of the conservation of the fermion current.

Once we enforce a conserved current (in what follows, a fermionic current), under the decomposition $X^\mu = A^\mu - \partial^\mu \pi / m_X$ the scalar field π decouples from this interaction leaving X^μ and A^μ indistinguishable.

III. TREE-LEVEL COUPLINGS OF A STÜCKELBERG VECTOR FIELD

We now turn to considering the tree-level interactions of a Stückelberg vector field X^μ . As we have emphasized, X^μ does not transform under a gauge symmetry. Hence, interactions in the effective theory will be built from powers of X^μ . The goal in this section is to enumerate the possible renormalizable tree-level interactions of X^μ and identify those that lead to scattering amplitudes that grow with powers of \sqrt{s}/m_X . These amplitudes arise from couplings of the longitudinal mode X_L^μ . The absence of a (gauge) symmetry under which X^μ transforms implies that its mass does not signal spontaneous symmetry breaking (SSB) nor the existence of Goldstone bosons. Nevertheless, the longitudinal mode, X_L^μ , is physical. We now state the *longitudinal equivalence theorem*: the leading interactions of the longitudinal mode can be characterized either by working directly with X_L^μ , or by using the fake gauge invariance of Eq. (9), choosing Landau gauge, and then associating X_L^μ with the interactions of the derivatively coupled longitudinal scalar field π . This is the Stückelberg analog of the Goldstone boson equivalence theorem.

A. The generalized Ward identity and the longitudinal equivalence theorem

In a theory with an exact $U(1)$ gauge symmetry, current conservation leads to the Ward identity

$$k^\mu \mathcal{M}_\mu = 0 \quad (27)$$

for an arbitrary amplitude \mathcal{M} in momentum space. This implies that the longitudinal polarization of an external on-shell gauge boson decouples. For a spontaneously broken $U(1)$ gauge theory, in which a gauge field A^μ acquires a mass m_X , the longitudinal polarization of an external on-shell gauge boson has, of course, physical couplings. Again using current conservation, a generalized Ward identity

$$\frac{k^\mu}{m_X} \mathcal{M}_\mu(A) = i\mathcal{M}(G^0; \xi = 0) \quad (28)$$

can be constructed that relates the momentum-contracted amplitude for an on-shell external gauge boson A^μ with momentum k^μ with the same amplitude, $\mathcal{M}(G^0; \xi = 0)$, for the Goldstone boson in Landau gauge. In the limit of large momentum $|\vec{k}| \gg m_X$, $\epsilon_L^\mu(A) \simeq k^\mu/m_X$, giving the Goldstone boson equivalence theorem

$$\epsilon_L^\mu(A) \mathcal{M}_\mu(A) \xrightarrow{|\vec{k}| \gg m_X} i\mathcal{M}(G^0; \xi = 0) \quad (29)$$

for a single on-shell, longitudinally polarized gauge boson.

For the massive Stückelberg vector boson, there is no (generalized or other) Ward identity since there is no gauge symmetry and thus no conserved local current associated with X^μ . This means

$$\frac{k^\mu}{m_X} \mathcal{M}_\mu(X) \neq 0. \quad (30)$$

At large momentum $|\vec{k}| \gg m_X$, $\epsilon_L^\mu(X) \simeq k^\mu/m_X$, and so this is simply a statement that the longitudinal mode of a Stückelberg vector field couples with a strength of k^μ/m_X .

What if we follow Secs. II B and II C and decompose the Stückelberg vector field into a fake gauge boson A^μ and scalar field π and use the fake gauge invariance and R_ξ gauge fixing to remove the $A^\mu \partial_\mu \pi$ mixing terms? Here, the gauge redundancy of A^μ and π implies that there is no gauge-independent identification of X_L^μ with A_L^μ and/or π . Consider the two-point functions Eq. (21) and Eq. (23). As we have discussed, the sum of the polarizations of X^μ is gauge independent. We can match the sum of the polarizations of X^μ to that of a massive gauge field A^μ by going to unitary gauge, $\xi \rightarrow \infty$. In unitary gauge, π does not play a dynamical role because $m_\pi^2 = \xi m_X^2 \rightarrow \infty$, and so $\epsilon_L^\mu(X) = \epsilon_L^\mu(A; \xi \rightarrow \infty)$. By contrast, in Landau gauge ($\xi = 0$) the sum of the polarizations of the two-point function of A^μ is purely transverse, matching that of a

massless gauge theory that has only two propagating DOF. Hence, in Landau gauge, the longitudinal polarization X_L^μ is fully captured by $\partial^\mu \pi/m_X$. This is the same result found in a spontaneously broken gauge theory in Landau gauge, where the longitudinal polarization of a massive gauge field is fully captured by $\partial^\mu G^0/m_X$ for the eaten Goldstone scalar field.

Therefore, analogously to Eq. (29) for a spontaneously broken theory, in Landau gauge at large momentum $|\vec{k}| \gg m_X$, we can identify

$$0 \neq \epsilon_L^\mu(k) \mathcal{M}_\mu(X) \xrightarrow{|\vec{k}| \gg m_X} \frac{k^\mu}{m_X} \mathcal{M}_\mu(X) = i\mathcal{M}(\pi; \xi = 0). \quad (31)$$

This is the *longitudinal equivalence theorem*: the leading behavior for on-shell, external X_L^μ interactions can be found by replacing X_L^μ with $\partial^\mu \pi/m_X$. For Stückelberg theories, longitudinal equivalence arises as a consequence of the invariance of Green's functions under BRST transformations (Slavnov-Taylor identities) carried out on the $A_\mu - \partial_\mu \pi/m_X$ formulation. Following Eq. (26), BRST invariance holds for Stückelberg theories regardless of whether $A_\mu - \partial_\mu \pi/m_X$ couples to conserved currents. Goldstone equivalence in a Higgsed $U(1)$ theory can also be formulated from BRST invariance (assuming $\partial_\mu j_{\text{ferm}}^\mu = 0$); however, it is more commonly derived using the generalized Ward identities from $U(1)$ gauge invariance (gauge fields coupling to conserved currents).⁴ Moreover, we can also identify the leading behavior of the off-shell two-point function [65],

$$\langle X^\mu(k) X^\nu(-k) \rangle \xrightarrow{k^2 \gg m_X^2} \frac{k^\mu k^\nu}{m_X^2} \langle \pi(k) \pi(-k); \xi = 0 \rangle. \quad (32)$$

The Stückelberg formalism makes clear that the large-momentum behavior found by using Eqs. (31) and (32) yields nonrenormalizable interactions of the longitudinal mode π suppressed by powers of m_X . Below, we will utilize these results in our discussions of the leading behavior of interactions and scattering amplitudes at large momentum. We note that Lagrangians involving X^μ do not necessarily contain interactions of the longitudinal mode π . For example, one special case is the Proca Lagrangian Eq. (4)

$$-\frac{1}{4} F_{X,\mu\nu} F_X^{\mu\nu} + \frac{1}{2} m_X^2 X_\mu X^\mu \xrightarrow{k^2 \gg m_X^2} \frac{1}{2} \partial_\mu \pi \partial^\mu \pi, \quad (33)$$

i.e., by the equivalence above, the Proca Lagrangian for a free massive Stückelberg vector field becomes the Lagrangian for a free massless scalar field π .

⁴See Refs. [63,64] for more details on the relation between the BRST Slavnov-Taylor identities and the generalized Ward identity in this regard.




FIG. 1. Diagrams for 2–2 scattering amplitudes with two fermions and two gauge bosons: (a) $XX \rightarrow f\bar{f}$; (b) $f\bar{f} \rightarrow XX$; (c) $fX \rightarrow fX$. [We have omitted the u -channel diagrams for (a), (b).] When $\partial_\mu j_A^\mu \neq 0$ due to the explicit violation of the global axial current by the fermion mass, the amplitudes for the longitudinally-polarized X field grow with energy proportional to $m_f \sqrt{s}/m_X^2$.

We now turn to considering interactions of X^μ with itself or with the SM, identifying those interactions that couple to the longitudinal mode, and discussing the consequences for the effective field theory.

B. Conserved vector current

Consider the interaction

$$g_X X_\mu j_V^\mu, \quad (34)$$

in which the Stückelberg vector field couples to a conserved vector current j_V^μ with strength g_X . For the purposes of this section, the current is assumed to be exactly conserved, $\partial_\mu j_V^\mu = 0$. (The anomalous case that leads to one-loop couplings will be discussed in detail in Sec. IV.) Using the equivalence $X^\mu \equiv A^\mu - \partial^\mu \pi/m_X$, it is clear that the longitudinal component π decouples from the conserved vector current, since under integration by parts (IBP)

$$\frac{1}{m_X} (\partial_\mu \pi) j_V^\mu \rightarrow -\frac{\pi}{m_X} \partial_\mu j_V^\mu \rightarrow 0. \quad (35)$$

This is the famous example of a dark photon kinetically mixed with electromagnetism, namely $j_V^\mu = j_{\text{em}}^\mu$, with coupling strength $g_X = \epsilon e$ [19]. This coupling is equivalent to a kinetically mixed Stückelberg field with the electromagnetic field strength using the EOM $j_{\text{em}}^\mu = \partial_\rho F^{\rho\mu}$ and IBP in the electroweak theory, $j_V^\mu = j_Y^\mu$, with coupling strength $g_X = \epsilon g/c_W$, where c_W is the cosine of the Weinberg angle. While j_Y^μ is no longer a pure vector current, it of course remains anomaly-free. (The couplings of X to the axial vector part of hypercharge will be discussed in the next section.)

While kinetic mixing $\epsilon F_{X,\mu\nu} F_V^{\mu\nu}$ is equivalent to $\epsilon g_V X_\mu j_V^\mu$, it is worth emphasizing that the inverse need not be true. The Stückelberg vector field X^μ can be coupled to a conserved current that is purely global and not gauged. For example, in the SM the global current j_{B-L}^μ is exactly conserved,⁵ and so the interaction

⁵The global $U(1)_{B-L}^3$ and $U(1)_{B-L}(\text{grav})^2$ anomalies vanish in the presence of three right-handed neutrinos, though this is not critical to our argument.

$$g_X X_\mu j_{B-L}^\mu \quad (36)$$

can be written without explicitly gauging $B-L$. This has fascinating consequences when one imagines X^μ coupling to a linear combination of both j_{em}^μ and j_{B-L}^μ [66].

These statements also hold for Stückelberg vector fields coupled to currents of hidden (dark) fermions, which are commonly found in the literature. In this scenario, it is often assumed that X^μ is the gauge boson of a new $U(1)$ and that the hidden fermions are charged under this symmetry. However, provided the hidden current coupling to X^μ is vectorlike and conserved, this need not be the case—the interaction is indistinguishable from a Stückelberg vector field coupled to a global (hidden fermion) current.

C. Axial-vector current

Next, consider an interaction of X^μ with an axial current,

$$g_X X_\mu j_A^\mu. \quad (37)$$

Unlike the case of the global vector current, the global axial-vector current is not, in general, conserved already at tree-level. This is simply because an axial-vector current is explicitly violated by fermion masses (within the SM or beyond).

The consequences of the axial-vector current violation by fermion mass is most easily seen by focusing on the longitudinal component of the Stückelberg vector field, X_L^μ , or equivalently $-\partial^\mu \pi/m_X$ following Eq. (8). For an axial current of fermions $j_A^\mu = \bar{f} \gamma^\mu \gamma_5 f$, the π field is derivatively coupled, so the longitudinal part of Eq. (37) becomes

$$\begin{aligned} g_X X_{L,\mu} j_A^\mu &\rightarrow -\frac{g_X}{m_X} \partial_\mu \pi (\bar{f} \gamma^\mu \gamma_5 f) \rightarrow \frac{g_X}{m_X} \pi \partial_\mu (\bar{f} \gamma^\mu \gamma_5 f) \\ &= \frac{2i g_X m_f}{m_X} \pi (\bar{f} \gamma_5 f), \end{aligned} \quad (38)$$

proportional to the fermion mass.

We can use this result to illustrate the high energy behavior of X^μ in several scattering processes that have axial-vector couplings including $f\bar{f} \rightarrow XX$, $XX \rightarrow f\bar{f}$, and $fX \rightarrow fX$ as shown in Fig. 1.

The full expression for the scattering amplitude follows by using the vector-boson polarization tensor for the external boson X^μ . Since we are interested in the high-energy behavior of the amplitude, we can focus on just the longitudinal part using Eq. (38). Using this effective interaction, and taking the limit of $s \gg m_X^2, m_f^2$ with t fixed in the amplitude squared, we find that $X_L X_L \rightarrow \bar{f}f$ is

$$|\mathcal{M}|^2 = \frac{32g_X^4 m_f^4 s}{m_X^4 (m_f^2 - t)} + \dots \quad (39)$$

where \dots stands for terms with subdominant energy growth. We see that the amplitude grows with energy proportional to $m_f \sqrt{s}/m_X^2$.

We can obtain a crude estimate of the scale at which perturbative unitarity is violated by setting $t \rightarrow 0$ (forward scattering) and $|\mathcal{M}|^2 = 1$,

$$\sqrt{s_{\max}} \sim \frac{1}{4\sqrt{2}} \frac{m_X^2}{g_X^2 m_f}. \quad (40)$$

The effective theory for a Stückelberg vector field has a cutoff scale that is parametrically above m_X only when $m_f \ll m_X$. This is fully equivalent to the Appelquist-Chanowitz bound on scattering amplitudes involving longitudinal electroweak gauge bosons and SM fermions when the Higgs is decoupled from the SM [67].

If instead the fermions are much heavier than the scattering energy, $m_f^2 \gg s \gg m_X^2$, the fermions can be integrated out, generating an effective $(X_\mu X^\mu)^2$ quartic interaction at one-loop order that will also lead to amplitudes that grow with energy. This is investigated below in Sec. III E.

The coupling of X^μ to an axial-vector current is equivalent to a dimension-4 Higgs-derivative interaction with X^μ

$$iH^\dagger \overleftrightarrow{D}_\mu H X^\mu, \quad (41)$$

where we remind the reader that $H^\dagger \overleftrightarrow{D}_\mu H = H^\dagger (D_\mu H) - (D_\mu H^\dagger)H$ is a SM gauge singlet with fully contracted $SU(2)_L \times U(1)_Y$ indices. Focusing on the longitudinal part,

$$iH^\dagger \overleftrightarrow{D}_\mu H X_L^\mu \rightarrow -iH^\dagger \overleftrightarrow{D}_\mu H \frac{\partial^\mu \pi}{m_X}. \quad (42)$$

Using IBP, the longitudinal coupling becomes

$$i \frac{\pi}{m_X} \partial^\mu (H^\dagger \overleftrightarrow{D}_\mu H) = i \frac{\pi}{m_X} [H^\dagger D^2 H - (D^2 H^\dagger)H]. \quad (43)$$

In the last line, we are free to promote the partial derivative to a covariant derivative since the additional SM vector

boson terms needed to covariantize the left-hand side of Eq. (43) vanish under \overleftrightarrow{D} . Applying the EOM of the Higgs field, the Higgs mass and quartic will cancel, leaving just the π coupling to a pseudoscalar current proportional to Yukawa couplings,

$$\rightarrow -i \frac{\pi}{m_X} \pi (\bar{f}_L y_f f_R - \bar{f}_R y_f^\dagger f_L) \frac{(v+h)}{\sqrt{2}}. \quad (44)$$

For the leptons and one type of quark, we can diagonalize the Yukawas so their entries are real and positive. In this case,

$$\rightarrow -iy_f \frac{(v+h)}{\sqrt{2}} \frac{\pi}{m_X} (\bar{f} \gamma_5 f). \quad (45)$$

We can convert this into an axial current by using the EOM for the fermions and IBP once more. Starting with Eq. (43),

$$\begin{aligned} & i \frac{\pi}{m_X} (-\bar{e}_R y_e^\dagger (H^\dagger L) + (\bar{L} H) y_e e_R + \dots) \\ &= i \frac{\pi}{m_X} (-\bar{e}_R i \not{D} e_R + \bar{L} i \not{D} L + \dots) \\ &= -\frac{\partial_\mu \pi}{m_X} (\bar{e}_R \gamma^\mu e_R - \bar{L} \gamma^\mu L + \dots) \\ &= \frac{\partial_\mu \pi}{m_X} (\bar{f} \gamma^\mu \gamma_5 f). \end{aligned} \quad (46)$$

Hence, the Higgs-derivative interaction can be rewritten as axial-vector couplings of the SM fermions with X^μ , and thus have the same energy growth in the amplitudes.

If we do not immediately focus on the longitudinal piece of X_μ , the operator $iH^\dagger \overleftrightarrow{D}_\mu H X^\mu$ contains a mass mixing between the X and Z and appears to lead to longitudinally enhanced $h \rightarrow XZ$ decays (assuming light X). However, as shown in Ref. [68], once the mass mixing is removed all purely bosonic, longitudinally enhanced interactions are eliminated. The only longitudinally enhanced amplitudes come from SM axial current couplings to X that arise as a consequence of the $X-Z$ mixing, in agreement with our result in Eq. (46).

While $X_\mu \bar{f} \gamma^\mu \gamma_5 f$ and $iH^\dagger \overleftrightarrow{D}_\mu H X^\mu$ separately lead to amplitudes that grow with energy, a carefully chosen combination of the two terms will not. This is precisely what occurs for X^μ coupling to the axial part of the hypercharge current [68]. Explicitly,

$$X_\mu j_{A,Y}^\mu \rightarrow i \sum_f \frac{y_f (v+h)}{\sqrt{2} m_X} ((Y_{f_R} - Y_{f_L}) \pm Y_H) \pi (\bar{f} \gamma_5 f) \quad (47)$$

after carrying out the manipulations in Eq. (38) to Eq. (46) and focusing on the longitudinal piece of X^μ . Here Y_{f_L}, Y_{f_R} are the hypercharges for f_L and f_R , respectively, Y_H is the Higgs hypercharge, and the $+(-)$ sign holds for leptons

and down-type quarks (up-type quarks). Inserting the hypercharges for SM matter, Eq. (47) vanishes. Thus, $X_\mu J_{A,Y}^\mu$ does not induce any amplitudes that grow with energy.

D. Higgs portal

At the renormalizable level, there is one independent Higgs interaction with X^μ ,

$$\frac{1}{2}\lambda_2|H|^2X_\mu X^\mu. \quad (48)$$

Inserting the Higgs vacuum expectation value (vev), this leads to an additional contribution to the mass of Stückelberg vector field. The shifted mass is

$$\tilde{m}_X^2 = m_X^2 + \frac{\lambda_2 v^2}{2}. \quad (49)$$

The interactions of the longitudinal component are identified as

$$\frac{1}{2}\lambda_2|H|^2X_\mu X^\mu \rightarrow \frac{1}{2\tilde{m}_X^2}\lambda_2|H|^2(\partial_\mu\pi\partial^\mu\pi). \quad (50)$$

This yields dimension-5 and dimension-6 interactions of the longitudinal mode π with the Higgs field

$$\frac{\lambda_2(2vh+h^2)}{2\tilde{m}_X^2}(\partial_\mu\pi\partial^\mu\pi) \quad (51)$$

that lead to scattering amplitudes that grow with powers of \sqrt{s}/m_X . Explicitly, examining the process $XX \rightarrow hh$ and using $|\mathcal{M}|^2 = 1$ as the criterion for the perturbative unitarity limit, we find $\sqrt{s_{\max}} \sim \sqrt{\frac{2}{\lambda_2}}\tilde{m}_X$.

If X^μ were to acquire its mass mostly through this interaction (i.e., $\tilde{m}_X^2 \simeq \lambda_2 v^2/2$), the strength of the coupling λ_2 cancels out in Eq. (51). In this case, the Stückelberg vector boson amplitudes grow with energy above the electroweak-breaking scale independently of the mass of the Stückelberg vector boson.

Finally, we note that this operator is familiar from the scenario of a $U(1)$ gauge field spontaneously broken by a complex scalar, in which case λ_2 would be identified with g^2 , the square of the $U(1)$ gauge coupling. This suggests that $\lambda_2 < 0$ is highly suspect: in particular, the positivity of λ_2 is mandatory in the case where the mass of the Stückelberg field is obtained from this operator.

E. Quartic self-interaction

At the renormalizable level, there is one operator that leads to a self-interaction of the Stückelberg vector field:

$$\frac{1}{4!}\lambda_4(X_\mu X^\mu)^2. \quad (52)$$

For the longitudinal component this becomes

$$\frac{\lambda_4}{4!m_X^4}(\partial_\mu\pi\partial^\mu\pi)^2. \quad (53)$$

In the presence of this quartic self-interaction, the 2–2 scattering amplitude with Stückelberg vector bosons grows with energy as

$$\mathcal{A}(X_L X_L \rightarrow X_L X_L) \sim \lambda_4 \frac{s^2}{m_X^4} \quad (54)$$

due to the couplings of the longitudinal mode. The s^2/m_X^4 growth of the four-point amplitude is the same as that encountered in the SM arising from (just) the four-point interaction of longitudinal W gauge bosons. Of course, this energy growth is famously canceled in the SM by Z and h exchange diagrams.

The breakdown of the effective theory from this operator can be obtained by performing a rough estimate of the maximum allowed energy as in the previous subsection,

$$\sqrt{s_{\max}} \lesssim \frac{m_X}{\lambda_4^{1/4}}. \quad (55)$$

Separating $\sqrt{s_{\max}}$ and m_X requires $\lambda_4 \ll 1$.⁶

However, restricting to just the interactions of the normal dark photon model, $(X_\mu X^\mu)^2$ is not generated radiatively. The coupling λ_4 is multiplicatively renormalized and thus technically natural if set to an exceptionally small number (including zero). It is well known that the sign of λ_4 must be positive to ensure UV analyticity [69]. Moreover, $(X_\mu X^\mu)^2$ is generated—even in the dark photon model—if we appeal to the usual lore that quantum gravity generates all possible higher dimensional operators (suppressed by powers of M_{Pl}). Specifically, we expect the operator

$$\frac{(H^\dagger H)(X_\mu X^\mu)^2}{M_{\text{Pl}}^2}, \quad (56)$$

which, below the scale of EWSB, leads to an effective quartic $\lambda_{4,\text{eff}} \sim v^2/M_{\text{Pl}}^2$. The growth of $\mathcal{A}(X_L X_L \rightarrow X_L X_L)$ is so rapid that even such a minuscule $\lambda_{4,\text{eff}} \sim \mathcal{O}(10^{-32})$ can lead to perturbative unitarity violation at low scales (meaning well within the energy range we have probed experimentally) when m_X is small. Explicitly,

⁶For vector-boson dark matter with $m_X \sim 10^{-5}$ eV [13] and requiring the cutoff scale to be $\Lambda = M_{\text{Pl}}$, we find an exceptionally small bound on the coupling $\lambda_4 \lesssim 10^{-129}$.

plugging this “gravity-generated” $\lambda_{4,\text{eff}}$ into Eq. (55) we find $\sqrt{s_{\text{max}}} \sim 10^8 m_X$.⁷

In the case of a Higgsed $U(1)$ theory in which the vector-boson mass is acquired through SSB, the energy growth of $XX \rightarrow XX$ scattering is tamed by the Higgs exchange diagram. In the low-energy effective theory below the mass of the Higgs (but above m_X), this interaction is generated with a coefficient $\lambda_4 = 6g^4 v^2 / m_h^2$ where $m_X = gv/2$, $m_h^2 = 2\lambda_h v^2$, giving $\mathcal{A}(XX \rightarrow XX) \sim \frac{1}{\lambda_h} \frac{s^2}{v^4}$. In other words, the scattering of vector bosons in a spontaneously broken $U(1)$ theory has an amplitude that grows with energy until the vev v , where the EFT must be supplemented by the Higgs boson.

IV. COUPLING A STÜCKELBERG VECTOR FIELD TO AN ANOMALOUS VECTOR CURRENT

Perhaps the most intriguing interaction that a Stückelberg vector field could have is $X_\mu j_{\text{anom}}^\mu$, a coupling to an anomalous current. In this section we will mainly focus on coupling to an anomalous *vector* current, since we already showed in Sec. III C that a tree-level coupling to an axial current generically leads to amplitudes that grow with energy.

For a gauge field, $A_\mu j_{\text{anom}}^\mu$ gauges what is a globally anomalous $U(1)$ current associated with j_{anom}^μ . In the presence of just one $U(1)$ gauge interaction, this leads to the usual $U(1)^3$ anomaly. When the fermions contributing to the current j_{anom}^μ also transform under other gauge symmetries, such as the SM, this leads to the mixed anomalies $(\text{SM})^2 U(1)$. The presence of the gauge anomalies leads to radiative corrections to the mass of the $U(1)$ gauge boson and to certain scattering amplitudes growing with energy [35,47].

In [45], a detailed analysis of a light $U(1)$ gauge boson coupled to an anomalous current was carried out. Their focus was on baryon number, which has the mixed anomalies $[U(1)_Y]^2 U(1)_B$ and $[SU(2)_L]^2 U(1)_B$. The interaction $A_\mu j_B^\mu$ leads to couplings of the longitudinal mode of A_μ with the (anomalous) baryon current. The consequences of this nonzero coupling emphasized in [45] are longitudinally enhanced interactions, including $Z \rightarrow A\gamma$ and other anomaly-induced decays. A careful analysis of the loop functions leading to this decay was carried out in [50].

But now there is a puzzle. The Stückelberg vector field interaction $X_\mu j_{\text{anom}}^\mu$ appears to lead to an anomalous fermion triangle loop, and yet, X^μ is not a gauge field. There cannot be $U(1)^3$ or $(\text{SM})^2 U(1)$ mixed gauge anomalies because there is no $U(1)$ gauge symmetry associated with X^μ .

⁷Higher-dimensional operators formed from X_μ alone will also be generated by the same argument. Forming amplitudes from these, i.e., $\mathcal{A}(X_L X_L \rightarrow 4X_L)$ from $(X_\mu X^\mu)^3 / M_{\text{Pl}}^2$, we find a similar, but weaker bound on $\sqrt{s_{\text{max}}}$.

In this section, we resolve this puzzle and, in the course of our analysis, find several consequences for theories with a Stückelberg vector boson. When we first introduce the fake gauge symmetry of Eq. (9), the mystery seems to deepen further because now A_μ would, in fact, appear to gauge an anomalous current. We will see that the term $(\partial_\mu \pi / m_X) j_{\text{anom}}^\mu$ precisely cancels the gauge anomaly that arises from $A_\mu j_{\text{anom}}^\mu$. The mechanism responsible for canceling the anomaly can be understood essentially by IBP,

$$-\frac{\partial_\mu \pi}{m_X} j_{\text{anom}}^\mu \rightarrow \frac{\pi}{m_X} \partial_\mu j_{\text{anom}}^\mu \propto \frac{\pi}{m_X} F_{\mu\nu} \tilde{F}^{\mu\nu}, \quad (57)$$

where $\tilde{F}_{\mu\nu} \equiv \frac{1}{2} \epsilon_{\alpha\beta\mu\nu} F^{\alpha\beta}$, and we recognize that the partial derivative of the anomalous current $\partial_\mu j_{\text{anom}}^\mu$ is proportional to $F_{\mu\nu} \tilde{F}^{\mu\nu}$, the Chern-Pontryagin density. The resulting dimension-5 interaction on the right-hand side of Eq. (57) is referred to as the Peccei-Quinn term⁸ (for any of the gauge symmetries of the SM, not just QCD). When this term is combined with suitable Wess-Zumino terms⁹ (coupling a gauge/vector field to a Chern-Simons class¹⁰ [55,70]) with appropriate choices of coefficients to restore gauge invariance, we will see that the Ward identities can be satisfied for all symmetries, verifying that A_μ does not have a gauge anomaly.¹¹

We now turn to considering the coupling of a vector field to an anomalous symmetry current, $j_{\text{anom}}^\mu = \sum_\psi q^\psi \psi^\dagger \bar{\sigma}^\mu \psi$, where q^ψ are the fermion charges under the symmetry. We wish to explicitly calculate the fermion loop attaching an external A^μ to two gauged vector bosons B^ν and C^ρ . Our discussion will apply to both a gauged vector field coupled

⁸This is also referred to as a “Green-Schwarz term” in some of the literature, e.g., [40].

⁹These are also referred to as “generalized Chern-Simons terms” in the literature, e.g., [40,46].

¹⁰The Chern-Simons class for a non-Abelian gauge field is (the second term is zero for the Abelian case)

$$\Omega^\mu = \epsilon^{\mu\nu\lambda\rho} \left(A_\nu^\alpha F_{\lambda\rho}^\alpha - \frac{1}{3} f^{abc} A_\nu^a A_\lambda^b A_\rho^c \right) \Rightarrow \partial_\mu \Omega^\mu = \frac{1}{2} \epsilon^{\mu\nu\lambda\rho} F_{\mu\nu}^\alpha F_{\lambda\rho}^\alpha. \quad (58)$$

¹¹Coupling A_μ and $\partial_\mu \pi$, the two “components” of X_μ , separately to the Chern-Simons class for an unbroken gauge symmetry Ω_B^μ yields

$$A_\mu \Omega_B^\mu = A^\mu \epsilon_{\mu\nu\lambda\rho} B^\nu F_B^{\lambda\rho}, \quad (59)$$

the dimension-4 Wess-Zumino term used to cancel the mixed anomaly, and

$$\frac{\partial_\mu \pi}{m_X} \Omega_B^\mu = -\frac{\pi}{m_X} \partial_\mu \Omega_B^\mu = -\frac{\pi}{m_X} F_{B\mu\nu} \tilde{F}_B^{\mu\nu}, \quad (60)$$

the dimension-5 Peccei-Quinn term. As we will see in Sec. IV C, the four-dimensional Green-Schwarz mechanism combines these two types of terms to cancel anomalies.




FIG. 2. Triangle diagrams responsible for the coupling of A^μ (decomposed from $X^\mu = A^\mu - \partial^\mu \pi / m_X$) to two gauge bosons B^ν and C^ρ . We have labeled Lorentz indices and directions of four-momenta according to their use in the main text.

to an anomalous local symmetry current, as well as a Stückelberg vector field X^μ coupled to an anomalous global symmetry current. For the Stückelberg vector field, however, we will do this by first decomposing $X^\mu = A^\mu - \partial^\mu \pi / m_X$, carrying out the calculation of the contribution to the gauge anomaly from A^μ , and then add back in the contribution from $\partial^\mu \pi / m_X$.

A. Triple-gauge vertex from a single fermion loop

Consider the triangle diagrams that contribute to the anomaly with general vector bosons A, B, C as shown in Fig. 2. By power counting, their amplitudes are linearly divergent and thus not uniquely defined. This can be encoded by including arbitrary four-momentum shifts a and b in the fermion loops in the left- and right-hand side diagrams, respectively. We will see that these arbitrary shifts are restricted by physical requirements, e.g., gauge invariance of either B or C .

Our convention for the amplitude of the sum of the triangle diagrams in Fig. 2 is

$$\tilde{\Delta}_{\{\mathbf{r}\}}^{\rho\mu\nu}(p, q; m_\psi; a, b) = g_C^{\mathbf{r}_1} g_A^{\mathbf{r}_2} g_B^{\mathbf{r}_3} \tilde{\Gamma}_{\{\mathbf{r}\}}^{\rho\mu\nu}(p, q; m_\psi; a, b), \quad (61)$$

where the indices $\mathbf{r}_i \in \{A, V\}$ indicate axial or vector couplings, respectively, of the boson with corresponding Lorentz superscript index in the same order, and m_ψ is the mass of the fermion ψ circulating in the loop. For now, all fermion charges have been subsumed into the couplings, so one should view $g_{A,B,C}^{\mathbf{r}_{1,2,3}}$ as specific to the particular fermion in the loop, i.e., the interaction term in the Lagrangian for this fermion is

$$\mathcal{L}_{\text{int}} = \bar{\psi} \gamma^\mu (g_C^V - g_C^A \gamma_5) \psi C^\mu + (C \rightarrow A, B). \quad (62)$$

Focusing for example on the case $\mathbf{r}_1 = A, \mathbf{r}_2 = V, \mathbf{r}_3 = V$, the amplitudes for the (coupling-stripped) triangle diagrams are:

$$\begin{aligned} \tilde{\Gamma}_{\text{AVV}}^{\rho\mu\nu}(p, q; m_\psi; a, b) = & \int_\ell \text{Tr} \left\{ \gamma_5 \gamma^\rho \frac{1}{\not{\ell} + \not{a} - \not{p} - m_\psi} \gamma^\mu \frac{1}{\not{\ell} + \not{a} - m_\psi} \gamma^\nu \frac{1}{\not{\ell} + \not{a} + \not{q} - m_\psi} \right. \\ & \left. + \gamma_5 \gamma^\rho \frac{1}{\not{\ell} + \not{b} - \not{q} - m_\psi} \gamma^\nu \frac{1}{\not{\ell} + \not{b} - m_\psi} \gamma^\mu \frac{1}{\not{\ell} + \not{b} + \not{p} - m_\psi} \right\}, \end{aligned} \quad (63)$$

where

$$\int_\ell \equiv \int \frac{d^d \ell}{(2\pi)^d}. \quad (64)$$

For the VAV and VVA amplitudes, we move the γ_5 matrix in front of γ^μ or γ^ν , respectively. To avoid nonchiral anomalies, we set $b = -a$ [71–73]. In terms of the external momenta p, q , we can then express the arbitrary shift $a = zp + wq$ using two real parameters z, w . The amplitude can be written in the Rosenberg parametrization as [50,71]

$$\begin{aligned} \tilde{\Gamma}_{\{\mathbf{r}\}}^{\rho\mu\nu}(p, q; z, w) = & \frac{1}{\pi^2} \{ G_{\{\mathbf{r}\}}^1(p, q; w) \epsilon^{\rho\mu\nu;p} + G_{\{\mathbf{r}\}}^2(p, q; z) \epsilon^{\rho\mu\nu;q} \\ & + (F_3(p, q) p^\mu + F_4(p, q) q^\mu) \epsilon^{\rho\nu;pq} \\ & + (F_5(p, q) p^\nu + F_6(p, q) q^\nu) \epsilon^{\rho\mu;pq} \}, \end{aligned} \quad (65)$$

where $\epsilon^{\rho\mu\nu;q} \equiv \epsilon^{\rho\mu\nu\alpha} q_\alpha$ and we have made implicit the fermion mass dependence. The form factors F_i are finite and independent of $\{\mathbf{r}\}$, whereas G^1, G^2 are dependent on the momentum shift a (or the parameters w, z) as a consequence of the linear divergences of the triangle diagrams.

Full details of computing the form factors is given in Appendix A. We quote here the final expressions for the AVV and VAV cases that we will use in the following

sections. Employing Eq. (A8) to eliminate F_5 , we obtain for the AVV and VAV cases:

$$\begin{aligned} G_{\text{AVV}}^1 &= \frac{1}{4}(z+1) + p^2 F_3 - p \cdot q F_4, \\ G_{\text{AVV}}^2 &= \frac{1}{4}(w-1) + q^2 F_6 + p \cdot q F_4, \end{aligned} \quad (66)$$

$$\begin{aligned} G_{\text{VAV}}^1 &= \frac{1}{4}(z+1) + p^2 F_3 - p \cdot q F_4, \\ G_{\text{VAV}}^2 &= \frac{1}{4}(w-1) + q^2 F_6 + p \cdot q F_4 - m_\psi^2 C_0(m_\psi^2). \end{aligned} \quad (67)$$

B. Momentum-contracted vertex functions

Now that we have established how the triple-gauge vertex can be manipulated into purely finite terms—form factors $F_{3\dots 6}$ plus the momentum-shift parameters w, z —we turn to its phenomenological consequences. The most interesting quantity is not the triple-gauge vertex itself, but what happens when the triple-gauge vertex is contracted with a longitudinally polarized A, B , or C : as explained in Sec. III A, the longitudinal polarizations are proportional to momenta in the large-momentum limit, and these can lead to scattering or decay amplitudes that grow with energy. The relevant quantities are the momentum-contracted vertex functions (MCVF)

$$(p+q)_\rho \tilde{\Delta}^{\rho\mu\nu}, p_\mu \tilde{\Delta}^{\rho\mu\nu}, q_\nu \tilde{\Delta}^{\rho\mu\nu}, \quad (68)$$

which are exactly the quantities we calculated in Appendix A to eliminate $G_{\{\mathbf{r}\}}^{1,2}$.

In fact, the MCVF are the starting points for the calculation of the Ward identities for this vertex, e.g., for A this is $p_\mu \mathcal{M}^\mu(A) = p_\mu \tilde{\Delta}^{\rho\mu\nu}$. For a vertex that respects all of the symmetries, $(p+q)_\rho \tilde{\Delta}^{\rho\mu\nu} = p_\mu \tilde{\Delta}^{\rho\mu\nu} = q_\nu \tilde{\Delta}^{\rho\mu\nu} = 0$, while for anomalous fermion content, one or more of these Ward identities is nonvanishing. Contracting the momentum of a massive gauge boson with the vertex function also yields a nonzero result, hence the Ward identity is also not satisfied. However, as we discussed in Sec. III A, one can construct a generalized Ward identity for a massive gauge boson that relates the MCVF of the massive gauge boson with that having the massive gauge bosons swapped with the Goldstone boson (for a spontaneously broken gauge symmetry) or the longitudinal mode (for a Stückelberg vector field).

Employing the procedure described in the previous subsection, we can compute $(p+q)_\rho \tilde{\Delta}^{\rho\mu\nu}$, $p_\mu \tilde{\Delta}^{\rho\mu\nu}$, $q_\nu \tilde{\Delta}^{\rho\mu\nu}$ for C, A, B , respectively, with arbitrary combination of V, A couplings to the fermions in the loop. For the remainder of the paper, however, we will make the simplification that one of the vector fields, which we take (without loss of generality) to be B , has purely vectorial couplings. This is because the phenomenological examples we will examine in Sec. V all share this property. The MCVF simplify to [50]:

$$\begin{aligned} (p+q)_\rho \tilde{\Delta}^{\rho\mu\nu} &= \frac{g_B^V}{4\pi^2} \epsilon^{\mu\nu;pq} \{ (w-z)(g_C^V g_A^A + g_C^A g_A^V) + 4m_\psi^2 C_0(m_\psi^2) \cdot g_C^A g_A^V \}, \\ -p_\mu \tilde{\Delta}^{\rho\mu\nu} &= \frac{g_B^V}{4\pi^2} \epsilon^{\rho\nu;pq} \{ (w-1)(g_C^V g_A^A + g_C^A g_A^V) - 4m_\psi^2 C_0(m_\psi^2) \cdot g_C^V g_A^A \}, \\ -q_\nu \tilde{\Delta}^{\rho\mu\nu} &= \frac{g_B^V}{4\pi^2} \epsilon^{\rho\mu;pq} \{ (z+1)(g_C^V g_A^A + g_C^A g_A^V) \}, \end{aligned} \quad (69)$$

where

$$\tilde{\Delta}^{\rho\mu\nu} = \sum_{\mathbf{r}_1, \mathbf{r}_2 \in \{A, V\}} \tilde{\Delta}_{\mathbf{r}_1 \mathbf{r}_2}^{\rho\mu\nu}, \quad (70)$$

and C_0 is a special case of the three-point Passarino-Veltman scalar function

$$\begin{aligned} C_0(m_\psi^2) &= C_0(q^2, (p+q)^2, p^2; m_\psi, m_\psi, m_\psi) \\ &= - \int_0^1 dx \int_0^{1-x} dy \Delta^{-1}, \end{aligned} \quad (71)$$

with Δ from Eq. (A3). Two relevant limits are

$$\begin{aligned} \lim_{m_\psi^2 \rightarrow \infty} m_\psi^2 C_0(m_\psi^2) &\rightarrow -\frac{1}{2}, \\ \lim_{m_\psi^2 \rightarrow 0} m_\psi^2 C_0(m_\psi^2) &\rightarrow 0. \end{aligned} \quad (72)$$

More precisely, these are limits of m_ψ^2 with respect to the other scales $p^2, (p+q)^2, q^2$ that appear in Eq. (A3).

In a theory with a fermion content that is nonanomalous, obviously all of the MCVF in Eq. (69) vanish independently of the presence or absence of masses for the vector bosons. When the fermion content is anomalous, i.e., with respect to A and/or C (recall that we take B to couple vectorially), the MCVF are not uniquely determined due to the freedom to choose the coefficients of the most general

momentum shift $a = zp + wq$ in the vertex function. This allows for several possibilities. One possible choice of coefficients results in all three MCVF being equal,

$$(p + q)_\rho \tilde{\Delta}^{\mu\nu\rho} = p_\mu \tilde{\Delta}^{\mu\nu\rho} = q_\nu \tilde{\Delta}^{\mu\nu\rho} \neq 0, \quad (73)$$

a configuration referred to as the ‘‘consistent anomaly’’ [74–76]. This choice is convenient from an EFT perspective: we view the contributions to the gauge anomaly as arising from the SM plus a contact term that, for instance, arises from some heavy fermions that maintain anomaly cancellation. In the consistent picture, all gauge symmetries are violated, so integrating out UV physics can generate gauge-violating operators. Combining these gauge-violating operators with the SM loop (also gauge-violating in the consistent picture) and choosing its coefficient appropriately, we can cancel all anomalies.¹²

A second possibility is to utilize momentum shifts such that the anomaly resides in only a single gauge interaction, the so-called ‘‘covariant anomaly’’ [75,76]. Gauge-variant Wess-Zumino effective operators of the form $\epsilon^{\mu\nu\rho\sigma} A_\mu C_\nu F_{B,\rho\sigma}$ can be added to the Lagrangian to shift from the consistent to covariant picture. This approach is often employed for calculations with two gauge bosons, B, C , with anomaly-free couplings and one (massive) gauge boson, A , that has anomalous couplings. By taking $w = z = -1$ in Eq. (69), the terms that are independent of fermion mass appear only in the MCVF for A

$$\begin{aligned} (p + q)_\rho \tilde{\Delta}^{\rho\mu\nu} &= \frac{g_B^V}{\pi^2} \epsilon^{\mu\nu;\rho q} m_\psi^2 C_0(m_\psi^2) \cdot g_C^A g_A^V, \\ -p_\mu \tilde{\Delta}^{\rho\mu\nu} &= -\frac{g_B^V}{2\pi^2} \epsilon^{\rho\nu;\rho q} \{ (g_C^V g_A^A + g_C^A g_A^V) \\ &\quad + 2m_\psi^2 C_0(m_\psi^2) \cdot g_C^V g_A^A \}, \\ -q_\nu \tilde{\Delta}^{\rho\mu\nu} &= 0. \end{aligned} \quad (74)$$

Notice that the fermion mass-dependent terms in the first two expressions above come with different coupling structures: if the fermions have purely axial couplings to A (VAV structure), the mass-dependent term vanishes from the first line, while if the couplings to C are purely axial (AVV structure), the mass-dependent term vanishes from the second line.

¹²In the covariant picture, discussed below, the issue in the EFT is that integrating out UV physics can only change the coefficients of SM terms, or generate new, higher-dimensional terms that respect the UV symmetries. As there is no B - or C -invariant, A -violating term, there is no coefficient to change, and the power counting for higher-dimensional terms will not work out correctly to cancel the anomaly. Therefore, working in the covariant picture requires doing calculations in the full UV theory, keeping both SM and UV physics and not taking the low-energy limit of SM + effective operators.

C. Anomaly cancellation, Ward identities, and π

We are now in a position to clarify the role that the longitudinal mode π plays in anomaly cancellation. Consider a theory with massless fermions in which the vector field A^μ has anomalous couplings. For a single massless fermion ψ , the MCVF become

$$\begin{aligned} (p + q)_\rho \tilde{\Delta}^{\rho\mu\nu} &= 0, \\ -p_\mu \tilde{\Delta}^{\rho\mu\nu} &= -\frac{g_C g_X g_B q_B^\psi}{2\pi^2} \epsilon^{\rho\nu;\rho q} (q_C^{V,\psi} q_X^{A,\psi} + q_C^{A,\psi} q_X^{V,\psi}), \\ -q_\nu \tilde{\Delta}^{\rho\mu\nu} &= 0, \end{aligned} \quad (75)$$

where from Eq. (72), $m_\psi^2 C_0(m_\psi^2) \rightarrow 0$ in the massless fermion limit. Here, we have also separated the coupling constants $g_{X,B,C}$ from the individual fermion charges $q_{X,B,C}^\psi$ by writing

$$\begin{aligned} g_A^V &= g_X q_X^{V,\psi}, & g_A^A &= g_X q_X^{A,\psi} \\ g_B^V &= g_B q_B^\psi, & g_B^A &= 0, \\ g_C^V &= g_C q_C^{V,\psi}, & g_C^A &= g_C q_C^{A,\psi}. \end{aligned} \quad (76)$$

In this limit, the only nonvanishing MCVF is the one involving the A^μ . If we sum over several massless fermions, this becomes

$$-p_\mu \tilde{\Delta}^{\rho\mu\nu} = -\mathcal{A}_X \frac{g_C g_X g_B}{2\pi^2} \epsilon^{\rho\nu;\rho q} \quad (77)$$

in terms of the A^μ anomaly coefficient

$$\mathcal{A}_X \equiv \sum_\psi q_B^{V,\psi} (q_C^{V,\psi} q_X^{A,\psi} + q_C^{A,\psi} q_X^{V,\psi}). \quad (78)$$

From the start of Sec. IV A until now, we have focused solely on the contribution to the MCVF from a vector field A^μ . Aside from forming the MCVF in Eq. (77) by contracting the momentum of A^μ onto the vertex, we have not specified whether A^μ is massive or massless. In addition, there is no distinction between whether A^μ is a gauge field that gauges the fermion current to which it couples with strength g_X or is in fact a Stückelberg vector field X^μ that couples to a global fermion current with strength g_X .

Below, we identify three distinct cases.

- (1) *A^μ is a massless gauge field:* In this case, Eq. (77) manifestly violates the Ward identity, and so either A^μ must acquire a mass or the theory contains multiple massless fermions with charges chosen such that, while the contribution from any single (Weyl) fermion is nonzero, the sum in Eq. (78) vanishes.
- (2) *A^μ represents X^μ , the massive Stückelberg vector field:* In this case, Eq. (77) is the final result for the

MCVF that connects X^μ with the gauge fields C^ρ and B^ν through a loop of massless fermions. There is no (generalized or other) Ward identity since there is no symmetry or conserved current associated with X^μ .

- (3) A^μ represents a massive gauge field arising from a spontaneously broken $U(1)$ gauge symmetry: This is the conventional case, which requires additional massive fermions, “anomalous,” to cancel the anomaly. The presence of anomalous is the key distinction from case 2.

We now want to compare and contrast cases 2 and 3, but we first need to resolve the puzzle of decomposing $X^\mu = A^\mu - \partial^\mu \pi / m_X$. In this decomposition, A^μ is a gauge field, and so $A_\mu j_{\text{anom}}^\mu$ necessarily gauges the anomalous current j_{anom}^μ ; however, $X_\mu j_{\text{anom}}^\mu$ is simply an interaction of a vector field with a globally anomalous current j_{anom}^μ . How can A^μ be anomalous under its gauge symmetry while X^μ has nothing to do with a gauge symmetry or a gauge anomaly?

The resolution is found by considering the additional contribution from the scalar field π . The Lagrangian contains

$$-g_X \frac{\partial_\mu \pi}{m_X} j_{\text{anom}}^\mu = g_X \frac{\pi}{m_X} \partial_\mu j_{\text{anom}}^\mu, \quad (79)$$

where we have used IBP to get the right-hand side. The divergence of the anomalous current is given by

$$g_X \partial_\mu j_{\text{anom}}^\mu = \mathcal{A}_X \frac{g_C g_X g_B}{4\pi^2} F_{C,\mu\nu} \tilde{F}_B^{\mu\nu}, \quad (80)$$

and so the scalar field contributes a dimension-5 Peccei-Quinn term in the Lagrangian,

$$\mathcal{A}_X \frac{g_C g_X g_B}{4\pi^2} \frac{\pi}{m_X} F_{C,\mu\nu} \tilde{F}_B^{\mu\nu}. \quad (81)$$

In momentum space, this interaction becomes

$$im_X \tilde{\Delta}^{\rho\nu}(\pi) = \mathcal{A}_X \frac{g_C g_X g_B}{2\pi^2} \epsilon^{\rho\nu;pq}, \quad (82)$$

namely a dimension-5 three-point vertex among π , C_ρ , and B_ν in the effective theory. We can combine Eq. (77) with Eq. (82) as

$$p_\mu \tilde{\Delta}^{\rho\mu\nu}(A) - im_X \tilde{\Delta}^{\rho\nu}(\pi) = 0. \quad (83)$$

This is the generalized Ward identity from Sec. III A for A^μ applied to the fermion triangle diagram. That is, so long as the dimension-5 Peccei-Quinn term has the specific coefficient given in Eq. (81), A^μ satisfies the generalized Ward identity. The specific coefficient that is required is precisely the one that permits the combination of the renormalizable $A_\mu j_{\text{anom}}^\mu$ and the dimension-5 interaction

$-\partial_\mu \pi j_{\text{anom}}^\mu / m_X$ to be written as $X_\mu j_{\text{anom}}^\mu$; in other words, the combination of A_μ and $(\partial_\mu \pi) / m_X$ must maintain the fake gauge invariance. This is otherwise known as the four-dimensional Green-Schwarz anomaly cancellation mechanism [29,35,36,54,77].

Since A^μ as part of X^μ is not an external state, we remark that Eq. (83), the generalized Ward identity, is not a statement about longitudinal equivalence. Contracting an on-shell external X^μ with $\tilde{\Delta}^{\rho\mu\nu}$ in the high-momentum limit $|\vec{k}| \gg m_X$ gives Eq. (77), which we can equivalently calculate using an external on-shell π and the longitudinal equivalence theorem in Eq. (31). That is, A^μ and π “conspire” to satisfy the generalized Ward identity while there is no analogue of this for X^μ .

Finally, it is interesting to compare and contrast what happens in a theory with a massive Abelian gauge boson in which the anomalous contribution is canceled by anomalous. The general case, with arbitrary vector and axial couplings for A^μ and C^ρ , can be worked out straightforwardly from Eq. (69). For the purposes of this discussion, however, we simply illustrate the similarities and differences in the case where the massless fermions contributing to the anomaly have purely vector interactions to A^μ and B^ν and purely axial interactions to C^ρ , in which case Eq. (78) simplifies to

$$\mathcal{A}_X = \sum_\psi q_B^{V,\psi} q_C^{A,\psi} q_X^{V,\psi}. \quad (84)$$

The massive anomalous have purely axial interactions to A^μ and purely vector interactions to B^ν and C^ρ ,

$$\mathcal{A}_X^{\text{anom}} = \sum_\psi q_B^{V,\psi} q_C^{V,\psi} q_X^{A,\psi}. \quad (85)$$

Anomaly cancellation requires that the sum of the charges of the anomalous under the gauge symmetries satisfy

$$\mathcal{A}_X^{\text{anom}} = -\mathcal{A}_X, \quad (86)$$

such that Eq. (78) vanishes.

However, for massive anomalous, there are additional contributions to the momentum-contracted vertex function from the C_0 functions in Eq. (69). We further simplify this discussion by taking all of the anomalous to have the same mass m_ψ . For the specific choices in Eqs. (84) and (85), the only nonzero MCVF is

$$\begin{aligned} -p_\mu \tilde{\Delta}^{\rho\mu\nu} &= -\frac{g_C g_X g_B}{2\pi^2} \epsilon^{\rho\nu;pq} [\mathcal{A}_X - \mathcal{A}_X^{\text{anom}} (1 + 2m_\psi^2 C_0(m_\psi^2))] \\ &= \mathcal{A}_X^{\text{anom}} \frac{g_C g_X g_B}{\pi^2} \epsilon^{\rho\nu;pq} m_\psi^2 C_0(m_\psi^2), \end{aligned} \quad (87)$$

where we used the anomaly cancellation condition Eq. (86) to get the second line. If the anomalous were massless, the right-hand side above would vanish using Eq. (72); this is as expected since by definition the theory would then be

anomaly-free and the Ward identities satisfied. If the anomalous are infinitely massive, the term in parentheses on the first line multiplying $\mathcal{A}_A^{\text{anom}}$ vanishes using Eq. (72), leaving the right-hand side nonzero and equal to Eq. (77), i.e., back to where we started.

With nonzero anomalon masses, Eq. (87) does not vanish. Following our discussion in Sec. III A, we can again construct a generalized Ward identity such that

$$p_\mu \mathcal{M}^\mu(A) - im_X \mathcal{M}(G^0) = 0. \quad (88)$$

where, for this discussion, G^0 is the Goldstone boson absorbed to make A^μ massive. Applying this to the MCVF for the fermion triangle diagram,

$$p_\mu \tilde{\Delta}^{\rho\mu\nu} - im_X \tilde{\Delta}^{\rho\nu}(G^0) = 0. \quad (89)$$

From this we can deduce the required interaction that the Goldstone boson must have with the MCVF,

$$i\tilde{\Delta}^{\rho\nu}(G^0) = \mathcal{A}_X^{\text{anom}} \frac{g_C g_X g_B}{\pi^2} \epsilon^{\rho\nu;pq} \frac{m_\psi^2}{m_X} C_0(m_\psi^2). \quad (90)$$

Here, we finally see the key difference between the case of a Stückelberg vector field and a spontaneously broken massive Abelian gauge field. In the specific example above, the anomalous have axial interactions with A^μ , implying the anomalous are *chiral* with respect to the gauge symmetry associated with A^μ . The only way to give mass to these chiral fermions without explicitly breaking the symmetry is to write Yukawa interactions with the Higgs field whose vev spontaneously breaks the gauge symmetry associated with A^μ . This means that, with conventional normalizations $m_\psi = y_\psi v/\sqrt{2}$ and $m_X = gv/2$, one power of the vev drops out in Eq. (90). Hence we see that the generalized Ward identity can be satisfied with *renormalizable* Yukawa interactions of the Goldstone mode with the fermions. This key difference is what permits a spontaneously broken gauge symmetry with anomalous fermion content (and a separate set of anomalous with heavier masses) to be at least possibly viable without a divergence in the UV leading to a cutoff scale. The caveat is that this requires Yukawa couplings to be perturbative (i.e., less than order one) in order to avoid Landau poles.

V. APPLICATIONS TO BARYON NUMBER

We now consider specific cases where the Stückelberg vector field X^μ couples to a globally anomalous current in order to investigate the phenomenological consequences. One of the most interesting possibilities is X^μ coupling to baryon number. Baryon number is anomalous in the SM, but anomaly-free with respect to $SU(3)_c \times U(1)_{\text{em}}$ below the electroweak scale. Here, our focus is to investigate the observable consequences of the longitudinal enhancements

that occur in the presence of $X_\mu j_B^\mu$, specifically three observables: $Z \rightarrow X\gamma$, $f\bar{f} \rightarrow X\gamma$, and $Z\gamma \rightarrow Z\gamma$. These depend on the electroweak scale and disappear in the limit $v \rightarrow \infty$. We compare and contrast our results with those when baryon number is gauged [45,50], identifying the similarities and differences for the case of a Stückelberg vector field. In the discussion below, we take all SM fermions to be massless; however, it is straightforward to reintroduce SM fermion mass dependence (e.g., [50]). In reality, only the top quark significantly invalidates this assumption, causing the baryon anomaly coefficient to be slightly smaller than what we have assumed below.

A. Prelude: $Z \rightarrow A\gamma$ with gauged baryon number

As a prelude to the results in subsequent sections, we want to review the calculation of $Z \rightarrow A\gamma$, where A^μ is the gauge field associated with gauged baryon number [78–80]; we reserve X^μ to refer to the Stückelberg vector field. However, we will use m_X , g_X , and q_X to refer to the mass, coupling, and charges of the (gauged or ungauged) vector field coupled to the baryon current.

In the SM, the baryon current is anomalous with respect to the mixed anomalies $U(1)_Y^2 U(1)_B$ and $SU(2)_L^2 U(1)_B$ in the specific combination [45]

$$\partial_\mu j_B^\mu = \frac{\mathcal{A}_B}{8\pi^2} (g^2 B_\mu \tilde{B}^{\mu\nu} - g^2 W_{\mu\nu} \tilde{W}^{\mu\nu}). \quad (91)$$

Here \mathcal{A}_B is the anomaly coefficient

$$\mathcal{A}_B = \sum_{f \in \text{SM}} Q^f q_X^{V,f} q_Z^{A,f}, \quad (92)$$

where the sum is over all of the fermions f in the SM with electric charge Q^f , baryon number q_B^f , and axial coupling $q_Z^{A,f} = T_3^f/2 = \pm 1/4$ to the Z . This is equivalent to the anomaly coefficient for just $SU(2)_L^2 U(1)_B$ or (the negative of) $U(1)_Y^2 U(1)_B$ since $U(1)_{\text{em}}^2 U(1)_B$ vanishes. Three generations of massless SM fermions give $\mathcal{A}_B = 3/4$.

As we have learned from Sec. IV B, we are free to choose a set of Wess-Zumino terms such that the only nonzero MCVF is

$$-p_\mu \sum_f \tilde{\Delta}_{\text{SM}}^{\rho\mu\nu} = -\mathcal{A}_B \frac{e g g_X}{2\pi^2 c_W} \epsilon^{\rho\nu;pq}, \quad (93)$$

following Eq. (69) with the specific choices $w = z = -1$.¹³

Baryon number can be made anomaly-free by extending the SM with anomalous ψ with charges such that, when they are included in the sum Eq. (92), the net result is zero.

¹³Had we included nonzero SM fermion masses, the first line of Eq. (69) would also be nonzero. Adding m_Z times the Z_ρ Goldstone contribution, a $\phi_Z - A - \gamma$ vertex, to the first line would yield zero.

For certain choices of their $SU(2)_L \times U(1)_Y$ charges, these anomalous can obtain masses independently of the electro-weak vev and therefore can be much heavier than the SM fermions. The full result for the decay rate $Z \rightarrow A\gamma$ including both SM fermions and a set of massive anomalous was given in [50]. However, for our purposes, it is more convenient to separate the contributions to the triangle loop from the SM, Eq. (93), and the massive anomalous. Defining $\tilde{\Delta}_{\text{anom}}^{\rho\mu\nu}$ as the contribution to the vertex function from the anomalous, and making the same choice $w = z = -1$, the additional contribution to the MCVF can again be easily obtained from Eq. (69),

$$-p_\mu \sum_\psi \tilde{\Delta}_{\text{anom}}^{\rho\mu\nu} = \mathcal{A}_B \frac{egg_X}{2\pi^2 c_W} e^{\rho\nu;pq} (1 + 2m_\psi^2 C_0(m_\psi^2)). \quad (94)$$

To cancel the anomaly and obtain mass without electro-weak symmetry breaking, these anomalous have pure vector couplings to Z and pure axial couplings to A such that

$$\sum_\psi Q^\psi q_X^{A,\psi} q_Z^{V,\psi} = -\mathcal{A}_B. \quad (95)$$

This is the same situation we encountered in Sec. IV C—the anomalon mass only appears in the MCVF in Eq. (94).

If we were instead to take $m_\psi \rightarrow 0$ (and therefore degenerate with the SM), the two sectors would cancel exactly, as required of an anomaly-free theory. For nonzero anomalon masses, the cancellation between the two sectors is inexact, leaving

$$-p_\mu \sum_{f,\psi} \tilde{\Delta}_{\text{tot}}^{\rho\mu\nu} = \mathcal{A}_B \frac{egg_X}{\pi^2 c_W} e^{\rho\nu;pq} m_\psi^2 C_0(m_\psi^2), \quad (96)$$

as in Eq. (87). It is interesting to consider anomalous that are much heavier than the Z boson.¹⁴ Then the right-hand side simplifies to

$$-p_\mu \sum_{f,\psi} \tilde{\Delta}_{\text{tot}}^{\rho\mu\nu} = -\mathcal{A}_B \frac{egg_X}{2\pi^2 c_W} e^{\rho\nu;pq}, \quad (97)$$

which up to corrections of $\mathcal{O}(m_Z^2/m_\psi^2)$ reduces to just the original SM-only contribution in Eq. (93). Dividing both sides by m_X , the above equation becomes the amplitude for $Z \rightarrow A_L\gamma$, where A_L is the longitudinal polarization. Squaring, we can convert this to a decay rate (again, in

the limit that the SM fields are massless and the anomalous are infinitely heavy)

$$\Gamma(Z \rightarrow A\gamma) \stackrel{m_X \ll m_Z}{\simeq} \Gamma(Z \rightarrow A_L\gamma) \simeq \frac{3}{32\pi^2} \frac{\alpha_{\text{em}}^2 \alpha_X m_Z^3}{c_W^2 s_W^2 m_X^3}, \quad (98)$$

where s_W is the sine of the Weinberg angle and we have used $\mathcal{A}_B = 3/4$.

As emphasized in [45], the m_Z^2/m_X^2 longitudinal enhancement implies the decay width is unbounded in the limit $m_X \ll m_Z$. For the effective theory to be valid, $\Gamma(Z \rightarrow A\gamma) < m_Z$, which implies a lower bound on m_X of

$$m_X > \sqrt{6\pi} \times \frac{egg_X}{64\pi^3 c_W} \times m_Z. \quad (99)$$

Up to an irrelevant numerical prefactor, this is the same bound obtained by Preskill [35] for an anomalous gauge theory by requiring the divergent three-loop contribution to the (anomalous) gauge boson mass not exceed its bare mass. More precisely, Preskill derived an expression $\Lambda = \frac{64\pi^3 c_W}{egg_X} m_X$ for the cutoff scale Λ of the effective theory that has the same scaling as Eq. (99) when we reinterpret the cutoff scale Λ to be m_Z .

What happens when m_X is lowered below the bound given in Eq. (99)? In a theory with anomalous, it is no longer possible to take their mass m_ψ to be much larger than m_Z . Approximating the results in [50] in the limit $m_X \ll m_\psi \ll m_Z$ (with massless SM fermions), we find

$$\Gamma(Z \rightarrow A\gamma) \simeq \frac{3}{32\pi^2} \frac{\alpha_{\text{em}}^2 \alpha_X}{c_W^2 s_W^2} \times m_Z \times \frac{m_\psi^4}{m_X^2 m_Z^2} \log^4 \frac{m_\psi^2}{m_Z^2}, \quad (100)$$

where now the EFT requirement $\Gamma(Z \rightarrow A\gamma) < m_Z$ implies the lower bound on m_X is modified to

$$m_X > \sqrt{6\pi} \times \frac{egg_X}{64\pi^3 c_W} \times m_\psi \times \frac{m_\psi}{m_Z} \log^2 \frac{m_\psi^2}{m_Z^2}. \quad (101)$$

This implies that we can lower the mass for m_X at the price of reducing the anomalon masses below m_Z . However, the additional suppression factor $m_\psi/m_Z \log^2 m_\psi^2/m_Z^2$ on the right-hand side in Eq. (101) relative to the result in Eq. (99) implies that the separation between m_X and m_ψ can become increasingly large as m_ψ is lowered below m_Z .

B. $Z \rightarrow X\gamma$ with global baryon number

Now we are in a position to evaluate $Z \rightarrow X\gamma$ when X is a Stückelberg vector field with coupling $g_X X_\mu j_B^\mu$ to the global, anomalous baryon current of the SM. The contribution to the $Z_\rho - X_\mu - \gamma_\nu$ vertex coming from loops of SM fermions is identical to the gauged case in the last section. Therefore, the nonzero MCVF is

¹⁴To play a role in the anomaly, the anomalous must receive some of their mass from the same SSB that gives mass to A and therefore $m_\psi \sim y v_X$, where $m_X \sim g_X v_X$ and y is some Yukawa coupling. A large hierarchy between the anomalous and X requires taking $g_X \ll y$, with the validity of perturbation theory limiting $y_{\text{max}} \sim 4\pi$. More discussion on the phenomenological implications of this “maximum hierarchy” between ψ and X can be found in [50].

$$-P_\mu \sum_f \tilde{\Delta}_{\text{SM}}^{\rho\mu\nu} = -\frac{\mathcal{A}_B}{2\pi^2} \frac{egg_X}{c_W} \epsilon^{\rho\nu;pq}. \quad (102)$$

This is the total contribution since there are no anomalous present.

Using this vertex to calculate $Z \rightarrow X\gamma$, we find

$$\Gamma(Z \rightarrow X\gamma) \stackrel{m_X \ll m_Z}{\simeq} \Gamma(Z \rightarrow X_L\gamma) \simeq \frac{3}{32\pi^2} \frac{\alpha_{\text{em}}^2 \alpha_X m_Z^3}{c_W^2 s_W^2 m_X^2}, \quad (103)$$

exactly the same result as Eq. (98), the case where $U(1)_B$ is gauged and made anomaly-free via infinitely heavy anomalous. We remark that the same result could also have been obtained using the longitudinal equivalence theorem to relate $\epsilon_L^\mu \tilde{\Delta}^{\rho\mu\nu}(X)$ to $\tilde{\Delta}^{\rho\nu}(\pi)$ in Landau gauge at large momentum $|\vec{k}| \gg m_X$.

Thus we see that the gauging of the would-be anomalous baryon number symmetry is irrelevant to the presence of the physically observable decay process $Z \rightarrow X\gamma$. It is the presence of the *global* baryon number anomaly that is essential for this decay to proceed. Said differently, our results show that the decay rate alone cannot differentiate between the scenarios of a gauge boson accompanied by heavy anomalous and a Stückelberg field coupled to a global current—a perspective emphasized in [73].

The presence of the Peccei-Quinn term, a dimension-5 operator in the Stückelberg EFT, implies a UV cutoff that cannot be taken arbitrarily large. Applying Eq. (81) to the specific case of the anomalous baryon current, the dimension-5 operator is

$$\mathcal{A}_B \frac{egg_X}{4\pi^2 c_W} \frac{\pi}{m_X} F_{Z,\mu\nu} \tilde{F}_{\text{em}}^{\mu\nu}, \quad (104)$$

and requiring the coefficient of this operator be less than 4π , we obtain a cutoff scale of order

$$\sqrt{s_{\text{max}}} \sim \frac{16\pi^3 c_W m_X}{\mathcal{A}_B e g g_X}. \quad (105)$$

The existence of a cutoff scale is not surprising because we previously discovered in Eq. (99) that we could not arbitrarily separate m_X from m_Z while allowing the decay rate $\Gamma(Z \rightarrow X\gamma)$ to remain perturbative. Both bounds scale similarly (up to numerical coefficients) with couplings and mass. What we see is that a Stückelberg vector field coupled to a globally anomalous current has a nonrenormalizable interaction signaling the existence of amplitudes that can grow with energy. This is explicitly seen in the decay rate $Z \rightarrow X\gamma$, and as we will see below, also occurs for processes that have one or more factors of $\tilde{\Delta}^{\rho\mu\nu}$ with an odd number of axial couplings embedded in the amplitude.

For finite $m_\psi \gg m_Z$, there will be corrections in Eq. (98) of $\mathcal{O}(m_Z^2/m_\psi^2)$ that are absent in Eq. (103). It is tempting to think that these corrections would be observable given

sufficiently accurate measurements of m_X and $\Gamma(Z \rightarrow X\gamma)$. However, this is premature, since in the case of a Stückelberg vector field, there are additional higher-dimensional operators suppressed by Λ that can contribute to the decay process. Hence, in the absence of direct observations (on-shell production) of anomalous and/or a Higgs boson, there is no way to unambiguously determine whether the decay process signals the existence of gauged baryon number, or instead, a Stückelberg vector field coupled to global baryon number.

C. $f\bar{f} \rightarrow X\gamma$

Attaching the Z^ρ leg of the $Z^\rho - X^\mu - \gamma^\nu$ vertex to a fermion current, we can explore how the longitudinal enhancement of the vertex manifests in $f\bar{f} \rightarrow X\gamma$, where f is a SM fermion. This calculation is interesting because it allows us to probe the triple-gauge vertex and its longitudinal enhancement at a wider range of energies than in Z decay. In particular, we can consider limits such as $m_X^2 \ll s \ll m_Z^2$, where the Z has been integrated out.

The diagrams for $f\bar{f} \rightarrow X\gamma$ are shown above in Fig. 3; an s -channel diagram proceeding through the triple-gauge vertex $\tilde{\Delta}^{\rho\mu\nu}$, plus t - and u -channel diagrams. The t - and u -channel diagrams involve only vectorial couplings and lead to the usual collinear divergences in the cross section. However, at least in the limit that the SM fermions are massless, they do not couple to the longitudinal part of X and thus do not grow with s (for a fixed scattering angle).¹⁵ Therefore, we will ignore these diagrams and focus on the s -channel piece, deferring a more general calculation to Appendix B. Furthermore, we will focus on the X_L piece of the amplitude, as this contains the leading dependence on s :

$$i\mathcal{M}(\bar{f}f \rightarrow X_L\gamma) = \frac{ig}{c_W} \frac{1}{s - m_Z^2} \bar{v}(k_2) \gamma^\rho \times (q_Z^{\nu,f} - q_Z^{\text{A},f} \gamma_5) u(k_1) \tilde{\Delta}_{\text{tot}}^{\rho\mu\nu} \frac{P_\mu}{m_X} \epsilon_\nu^*(q), \quad (106)$$

where $q_Z^{\nu,f}$ ($q_Z^{\text{A},f}$) are the vectorial (axial) couplings of fermion f to the Z and $\tilde{\Delta}_{\text{tot}}^{\rho\mu\nu}$ is the triple-gauge vertex after summing over all fermions—SM and beyond—in the loop. Here we have used $\epsilon_L^\mu(X) \rightarrow p^\mu/m_X$ for large $|\vec{p}| \gg m_X$,




FIG. 3. Diagrams for $f\bar{f} \rightarrow X\gamma$, with f an SM fermion: if only anomalous ψ couple to X , only the left diagram is relevant; otherwise, if f also couples to X , there are the t - and u -channel diagrams as well (cross diagrams not shown).

and so we are implicitly imagining a scenario where \sqrt{s} of the process is large compared to m_X . Note that only the transverse part of the Z propagator enters, since $(p+q)_\rho \tilde{\Delta}_{\text{tot}}^{\rho\mu\nu} = 0$.

As shown in previous sections, $p_\mu \tilde{\Delta}_{\text{tot}}^{\rho\mu\nu}$ has the same value whether we consider a Stückelberg vector field coupled to global baryon number or a gauged baryon number with anomalous much heavier than all of the other physical scales in the process. Thus we can evaluate $p_\mu \tilde{\Delta}_{\text{tot}}^{\rho\mu\nu}$ via Eq. (102) or Eq. (97), yielding

$$i\mathcal{M}(f\bar{f} \rightarrow X_L\gamma) = \frac{ig}{c_W} \frac{\mathcal{A}_B e g g_X}{2\pi^2 c_W} \frac{1}{m_X s - m_Z^2} \frac{1}{m_X s - m_Z^2} \bar{v}(k_2) \gamma^\rho \times (q_Z^{V,f} - q_Z^{A,f} \gamma_5) u(k_1) \epsilon^{\rho\nu; pq} \epsilon_\nu^*(q). \quad (107)$$

The details of the calculation of the leading behavior of the squared, polarization-summed and initial-state spin-averaged amplitude are given in Appendix B. We employ Eq. (B14) with only the AVV terms in the second line and the massless fermion limit of Eq. (B15) to obtain

$$|\bar{\mathcal{M}}|^2 \sim \frac{1}{4(N_c)^2 \pi^4} \left(\left(\frac{g}{c_W} \right)^2 g_X e \right)^2 \frac{(q_Z^{V,f})^2 + (q_Z^{A,f})^2}{(s - m_Z^2)^2} \cdot \frac{s^2}{4r_X} \left(1 - \frac{2tu}{s^2} \right) \cdot \left(\sum_q \kappa^q Q^q q_Z^{A,q} \right)^2, \quad (108)$$

where N_c is the number of colors of the initial-state fermions, $\kappa^{V,q} = \kappa^q = 1/3$ is the baryon number of the quarks, and Q^q is the electromagnetic charge of the quark q . For up- and down-type quarks, $q_Z^{A,q} = T_3/2$, so $Q^u q_Z^{A,u} = 1/6$ and $Q^d q_Z^{A,d} = 1/12$. The entire squared-sum on the right-hand side evaluates to $1/16$ for one generation (including the color factor); there is an additional factor of 9 for three generations. This agrees with $\mathcal{A}_B^2 = 9/16$. The cross section resulting from this amplitude is

$$\sigma(f\bar{f} \rightarrow X_L\gamma) = \frac{3}{8\pi} \frac{1}{N_c^2} \frac{\alpha_{\text{em}}^3 \alpha_X}{c_W^4 s_W^4} ((q_Z^{V,f})^2 + (q_Z^{A,f})^2) \frac{(s - m_X^2)^2}{m_X^2 (s - m_Z^2)^2}. \quad (109)$$

This expression already assumes $s \gg m_X^2$ (and in the case of gauged baryon number, the masses of any anomalous are much greater than m_Z and \sqrt{s}); however, there are a couple of further limits that are interesting to explore. First, consider $s \gg m_Z^2$, with the hierarchy of scales $m_X^2 \ll m_Z^2 \ll s$. In this case, the cross section becomes a constant

$$\sigma(f\bar{f} \rightarrow X_L\gamma)_{s \gg m_Z^2} = \frac{3}{8\pi} \frac{1}{N_c^2} \frac{\alpha_{\text{em}}^3 \alpha_X}{c_W^4 s_W^4} ((q_Z^{V,f})^2 + (q_Z^{A,f})^2) \frac{1}{m_X^2}. \quad (110)$$

A 2–2 scattering cross section constant in energy implies an amplitude squared that grows as s , so an amplitude that grows linearly with energy.

A more interesting limit is $s \ll m_Z^2$, with the hierarchy of scales $m_X^2 \ll s \ll m_Z^2$. In this limit,

$$\sigma(f\bar{f} \rightarrow X_L\gamma)_{m_X^2 \ll s \ll m_Z^2} = \frac{3}{8\pi} \frac{1}{N_c^2} \frac{\alpha_{\text{em}}^3 \alpha_X}{c_W^4 s_W^4} ((q_Z^{V,f})^2 + (q_Z^{A,f})^2) \frac{s^2}{m_Z^4 m_X^2}. \quad (111)$$

This cross section implies an amplitude squared $\propto s^3$, so an amplitude $\propto s^{3/2}$. To see why this limit is intriguing, let us write the amplitude squared as $\propto \frac{s^2}{m_Z^4} \frac{s}{m_X^2}$. If we use the condition $|\mathcal{M}|^2 = 1$ to set a limit on the cutoff of the theory, we find

$$\sqrt{s_{\text{max}}} \sim \frac{1}{\alpha_{\text{em}}^{1/2} \alpha_X^{1/6}} \left(\frac{m_X}{m_Z} \right)^{1/3} m_Z. \quad (112)$$

We contrast the above with the result from a four-fermion interaction in the Fermi theory. There, the amplitude $\mathcal{M}(f\bar{f} \rightarrow f\bar{f}) \sim \frac{s}{m_Z^2}$ (using m_Z instead of v to make the comparison easier and neglecting couplings and numerical factors), implying $\sqrt{s_{\text{max}}} \sim m_Z$ —a cutoff at the scale of particles we have integrated out. Compared to this, the limit from $f\bar{f} \rightarrow X_L\gamma$ is *smaller* by a factor of $(m_X/m_Z)^{1/3}$. We remark in passing that in Eq. (112), it is curious that the cutoff scale of the theory scales as $m_X^{1/3}$ in the same way as the weak gravity conjecture suggests when m_Z is replaced with M_{Pl} [47].

The situation becomes even more intriguing once we recall that the SM below the weak scale is purely vectorial. The triple-gauge vertices formed from loops of fermions with vectorial couplings (VVV in the language introduced in Sec. IV) are zero—stated in the language of gauge anomalies, the theory is anomaly-free. As such, $f\bar{f} \rightarrow X\gamma$ cannot exhibit any pathological scaling with respect to s in the limit that we take the weak scale to be infinitely heavy. The $1/m_Z^4$ scaling on the right-hand side of Eq. (111) satisfies this requirement; however, unusually, it predicts the scale where perturbative unitarity is violated (using $|\mathcal{M}|^2 \leq 1$) to be parametrically lower than the weak scale. If we require that $f\bar{f} \rightarrow X\gamma$ remain valid at least until m_Z , this sets a lower limit on the mass of X ,

$$m_{X,\text{min}} \sim \frac{4[3\alpha_{\text{em}}^3 \alpha_X ((q_Z^{V,f})^2 + (q_Z^{A,f})^2)]^{1/2}}{c_W^2 s_W^2} m_Z. \quad (113)$$

Taking f to be a charged lepton, $m_{X,\text{min}} \sim 6 \times 10^{-3} \sqrt{\alpha_X} m_Z$.

¹⁵Additionally, the interference between the t - and u -channel diagrams and the s -channel diagram is zero.




FIG. 4. $Z\gamma \rightarrow Z\gamma$ scattering through an s -channel π (left) or A (right, with 3 cross diagrams not pictured).

The above bound is a function of α_X , so it can be made arbitrarily small by sending $\alpha_X \ll 1$; in other words, for $\alpha_X \ll 1$, the EFT cutoff implied by Eq. (112) can be pushed above the scale of current experiments. It is an interesting and open question as to whether processes such as $f\bar{f} \rightarrow X\gamma$ could place bounds on (α_X, m_X) that are competitive with bounds from $Z \rightarrow X\gamma$ and other electroweak scale processes.

D. $Z\gamma \rightarrow Z\gamma$ via X exchange

The final amplitude we calculate using the triple-gauge vertex involves a so-called ‘‘BIM’’ process [81], the scattering of gauge fields off each other through the exchange of an off-shell Stückelberg vector field. The original BIM calculation considered the scattering of massless bosons through a (massive) Stückelberg vector field. For consistency with previous sections, here we specialize the calculation to the case of a Stückelberg vector field coupling to global baryon number, and so we consider $Z\gamma \rightarrow Z\gamma$ through an s -channel X . The diagrams involving X are shown in Fig. 4.

There are, of course, additional diagrams from boxes of fermions or W bosons, but these are independent of g_X . If all loop fermions ψ are heavy relative to \sqrt{s} , we recover the Euler-Heisenberg Lagrangian from the box diagrams, with $\mathcal{M}(Z\gamma \rightarrow Z\gamma) \sim \frac{s^2}{m_\psi^4}$. Here we focus on the same scenario considered in the previous subsections, with only massless SM fermions in the loop. (We would obtain the same result with gauged baryon number so long as the anomalon masses are taken to be much heavier than all other scales). Together with the limit $s \gg m_W^2$, the box diagrams involving W bosons have no bad s behavior¹⁶ [82,83], so we will neglect them and focus on the contributions from X exchange.

The X exchange occurs through a single diagram stitching together two $Z - \gamma - X$ vertices. However, if we write X^μ as $A^\mu - \partial^\mu \pi / m_X$ and employ gauge fixing as described in Sec. II C, there appear to be two diagrams as in Fig. 4—one from π exchange and one from A exchange, each with gauge dependence. Feynman rules for these diagrams can be derived from the Lagrangian in Appendix C.

¹⁶Here we are referring to $(s/M)^n$ behavior at fixed scattering angle, where M is some other mass scale in the problem, and not to divergences in the limit of forward or backward scattering. The latter manifest as ratios of Mandelstam invariants.

The A exchange piece for $Z_\rho(p_1)\gamma_\nu(q_1) \rightarrow Z_{\rho'}(p_2)\gamma_{\nu'}(q_2)$, coming from loops of SM fermions alone, is

$$\frac{-i}{s - m_X^2} \tilde{\Delta}_{\text{SM}}^{\mu\rho\nu} \left(g^{\mu\mu'} - (1 - \xi) \frac{(p_1 + q_1)^\mu (p_2 + q_2)^{\mu'}}{s - \xi m_X^2} \right) \tilde{\Delta}_{\text{SM}}^{\rho'\mu'\nu'}, \quad (114)$$

while the π piece is

$$\frac{i}{s - \xi m_X^2} \left(\frac{\mathcal{A}_B e g g_X}{2\pi^2 c_W m_X} \right)^2 e^{\rho\nu; p_1 q_1} e^{\rho'\nu'; p_2 q_2}. \quad (115)$$

Evaluating the gauge-dependent piece of Eq. (114) using Eq. (93) with appropriate modifications, then its sum with the π exchange term,

$$\begin{aligned} & \left[\frac{i}{s - m_X^2} \frac{(1 - \xi)}{s - \xi m_X^2} + \frac{i}{m_X^2 (s - \xi m_X^2)} \right] \left(\frac{\mathcal{A}_B e g g_X}{2\pi^2 c_W} \right)^2 e^{\rho\nu; p_1 q_1} e^{\rho'\nu'; p_2 q_2} \\ &= \left(\frac{\mathcal{A}_B e g g_X}{2\pi^2 c_W} \right)^2 e^{\rho\nu; p_1 q_1} e^{\rho'\nu'; p_2 q_2} \frac{i}{s - \xi m_X^2} \left(\frac{(1 - \xi)}{s - m_X^2} + \frac{1}{m_X^2} \right) \\ &= \left(\frac{\mathcal{A}_B e g g_X}{2\pi^2 c_W} \right)^2 e^{\rho\nu; p_1 q_1} e^{\rho'\nu'; p_2 q_2} \frac{i}{m_X^2 (s - m_X^2)}, \end{aligned} \quad (116)$$

we see that the ξ dependence cancels. Notice that the final result of Eq. (116) is the same as just π exchange given by Eq. (115) in the limit $s \gg m_X^2$ in Landau gauge ($\xi = 0$), as required by the longitudinal equivalence theorem in Eq. (32).

Assuming $s \gg m_Z^2, m_X^2$, we can neglect the other terms and use Eq. (116) as an approximation to the full amplitude, deferring a more complete and general calculation to Appendix C. Forming a cross section from Eq. (116) and taking the large- s limit, we find:

$$\sigma(Z\gamma \rightarrow Z\gamma) \simeq \frac{27}{128\pi^3} \frac{\alpha_{\text{em}}^4 \alpha_X^2}{c_W^4 s_W^4 m_X^4} s + \dots \quad (117)$$

where the \dots indicates terms subleading in s .

While the diagrams in Fig. 4 are reminiscent of longitudinal W scattering in the SM, we emphasize that the external γ, Z fields in the BIM process are purely transverse. In the large- s limit, contracting the vertices above with longitudinal Z polarizations yields zero (for massless SM fermions) via the MCVF.

VI. DISCUSSION

We have investigated theories with a Stückelberg vector field, emphasizing the systematic approach to constructing an effective field theory involving X^μ . We considered several possible interactions of the Stückelberg vector field with the SM or with itself, identifying the couplings of the longitudinal mode that lead to scattering amplitudes that grow with energy. At tree-level these involve the operators $(X_\mu X^\mu)^2$, $H^\dagger H X_\mu X^\mu$ and $H^\dagger D_\mu H X^\mu$, while the interaction $X_\mu j_{\text{anom}}^\mu$ (with j_{anom}^μ an anomalous global current) induces one-loop amplitudes that grow with energy. The energy growth implies an EFT with one of these interactions requires a UV cutoff scale that appears above m_X by an amount that is parametrically $1/(\text{coupling})$ of the interaction. In the specific case of X^μ coupled to the global baryon current, we demonstrated that the finite contribution to the fermion triangle diagram leads to a variety of processes that have longitudinal enhancements in the small m_X limit, including $Z \rightarrow X\gamma$, $f\bar{f} \rightarrow X\gamma$ and $Z\gamma \rightarrow Z\gamma$.¹⁷

We performed a detailed analysis of the operator $X_\mu j_{\text{anom}}^\mu$. This interaction is, at first, somewhat puzzling since X^μ is not a gauge boson and yet it suggests X^μ is gauging an anomalous current. Preskill [35] demonstrated that anomalous gauge theories are simply effective theories with a narrow range of scales where the EFT is valid. His analysis emphasized the UV divergent contributions to the two-point function, leading to maximum separation between the mass of the gauge boson of an anomalous theory and the cutoff scale of the theory. As we have seen, this result holds for theories with a Stückelberg vector field that has no gauge symmetry. In particular, we demonstrated that the generalized Ward identity is satisfied if and only if the contributions from both A^μ , the (fake) gauge boson associated with a (fake) gauge symmetry, and $\partial^\mu \pi/m_X$ appear in the specific gauge-invariant combination $A^\mu - \partial^\mu \pi/m_X$. Our analysis demonstrates that it is the existence of the global anomaly, not the gauging of it, that leads to the physical consequence of scattering amplitudes that grow with energy in the UV. This is reminiscent of [84] and may lead to a different interpretation of anomalies when expressed directly in terms of on-shell scattering amplitudes. For example, [85] recasts the constraints from anomaly cancellation in terms of on-shell amplitudes that satisfy unitarity and locality.

As we have seen throughout the paper, the interactions of a Stückelberg vector field that grow with energy can and do arise from a spontaneously broken $U(1)$ gauge theory with a dark Higgs sector. In each case, the coefficient of the corresponding operator depends explicitly on powers of g_X ,

the $U(1)$ gauge coupling. The cutoff scale of the EFT with the vector boson is resolved by dark Higgs exchange, in analogy with the growth of the scattering of longitudinally-polarized electroweak gauge bosons in the SM. UV completing an EFT with a Stückelberg vector boson using a dark Higgs sector, in which the vector boson mass is much smaller than the dark Higgs mass, requires $g_X \ll \sqrt{\lambda_h}$. In a $U(1)$ gauge theory in which all $U(1)$ field charges are order one, this implies that all interactions are suppressed by powers of the small gauge coupling g_X , and in particular, kinetic mixing arising from integrating out matter that is charged under the dark $U(1)$ and $U(1)_Y$ is also bounded by $\epsilon \lesssim g_X e$.

A Stückelberg vector field can be obtained by ungauging a Higgsed theory while holding m_X fixed, that is, by sending $g_X \rightarrow 0$ and thus taking $v_X \rightarrow \infty$. This is distinct from spontaneously-broken gauge theories: the limit $m_X \rightarrow 0$, $g_X \rightarrow 0$ with the ratio $v_X = m_X/g_X$ held constant does not exist, demonstrating that a strict interpretation of a theory with a Stückelberg vector boson does not have anything to do with SSB. There is no Higgs mechanism, no Higgs boson, and so the interactions that lead to longitudinally enhanced scattering amplitudes that grow with energy have arbitrary coefficients. Consequently, a UV cutoff scale of the EFT is inevitable. Reece [22] has suggested that weak gravity conjecture arguments [86] prevent an arbitrarily small Stückelberg mass since the limit $m_X \rightarrow 0$ lies at infinite distance in field space. It would be interesting to further investigate the constraints on other parameters of the effective theory of Stückelberg vector bosons using arguments based on embedding the theory into quantum gravity [87,88].

In the $SU(3)_c \times U(1)_{\text{em}}$ effective theory below the electroweak scale, all fermion currents are vectorial with no (gauge or global) anomalies. Naively, there are no restrictions on coupling an arbitrarily light Stückelberg vector field to any linear combination of these currents. Of course, the weak interaction explicitly violates some global symmetries, such as baryon number, so the interactions of X with SM fermion currents are not purely vectorial. Hence, X will have scattering amplitudes that grow with powers of \sqrt{s}/m_X .¹⁸ One might think this growth is the same as four-fermion interactions that also scale with s/m_W^2 , such that the cutoff scale of the theory is the electroweak breaking scale. This is not true. Consider $f\bar{f} \rightarrow X\gamma$ with X coupling to baryon number. While there is s/m_Z^2 suppression in the amplitude from Z exchange, there is also \sqrt{s}/m_X enhancement from producing a longitudinally polarized X . By observing this energy growth in the cross section (at energies well below the electroweak scale), one could determine whether or not a vector boson has longitudinally enhanced couplings.

¹⁷The importance of $Z \rightarrow X\gamma$ for gauged baryon number was emphasized in [43,50] along with other FCNC processes involving $K \rightarrow \pi X$ and $B \rightarrow KX$ meson decays [43]. Constraints on other $U(1)$ s were discussed in [48].

¹⁸An alternative approach in which a vector field interacts only through higher-dimensional operators was discussed in [89].

Finally, we should discuss the status of dark photons that partly motivated our study of Stückelberg vector fields. In theories where the dark photon Lagrangian arises from a spontaneously broken $U(1)$ gauge symmetry by a dark Higgs field, some discussion of the dark Higgs scalar has appeared (e.g., [5,7,90–95]). Instead, we proclaim that the time is ripe to consider a general set of interactions that a Stückelberg vector field can have with coefficients that are not dictated by a dark Higgs sector. Longitudinally enhanced interactions imply the theory will have a cutoff scale: within the validity of the effective theory (i.e., \sqrt{s} less than the cutoff scale as determined by the longitudinally enhanced scattering processes), what phenomenological consequences can arise in the presence of these interactions? This is an interesting question to explore for more general vector boson dark matter as well as for dark photon models.

Ultimately our discussion of a Stückelberg vector field reiterates the lesson of the precarious nature of vector fields in quantum field theory whose mass is not associated with SSB. The longitudinal component generically couples to itself or to the SM, and the presence of these couplings leads to amplitudes that grow with energy and thus require a cutoff scale for the EFT. There are only two resolutions: craft the effective theory to have no couplings of the longitudinal mode, i.e., X coupled only to an anomaly-free global current, or introduce a Higgs mechanism with a Higgs boson to restore unitarity of longitudinal vector boson interactions. If evidence of a new vector boson were uncovered in data, we hope our analysis provides a framework to characterize the effective field theory comprising the leading interactions of the vector boson independent of its ultimate UV origin.

ACKNOWLEDGMENTS

The authors wish to acknowledge that this research was initiated at the Munich Institute for Astro- and Particle Physics (MIAPP) conference “The Weak Scale at a Crossroads: Lessons from the LHC and Beyond” in May–June 2019; MIAPP is funded by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Germany’s Excellence Strategy Grant No. EXC-2094–390783311. The work of G. D. K. was supported in part by the U.S. Department of Energy under Grant No. DE-SC0011640. G. L. acknowledges support by the Samsung Science & Technology Foundation under Project No. SSTF-BA1601-07, a Korea University Grant, and the support of the U.S. National Science Foundation through Grant No. PHY1719877. G. L. is grateful to the University of Toronto for partial support during completion of this work. The work of A. M. is partially supported by the National Science Foundation under Grants No. PHY-1820860 and No. PHY-2112540.

APPENDIX A: FORM FACTORS IN THE ROSENBERG PARAMETRIZATION OF THE TRIANGLE DIAGRAMS

In this appendix, we detail the computation of the amplitude of the triple-gauge boson triangle diagrams of Fig. 2. Factoring out couplings, the relevant expressions are of the type in Eq. (63). To compute the finite form factors $F_{3,\dots,6}$, we follow the procedure of [40]. The denominators on the first and second lines of Eq. (63) can be combined as

$$\begin{aligned} & [((\ell \pm q)^2 - m_\psi^2)(\ell^2 - m_\psi^2)((\ell \mp p)^2 - m_\psi^2)]^{-1} \\ &= \Gamma(3) \int_0^1 dx \int_0^{1-x} dy [\ell^2 \pm 2\ell \cdot k + xq^2 \\ & \quad + yp^2 - m_\psi^2 + i\epsilon]^{-3}, \end{aligned} \quad (\text{A1})$$

where $k = xq - yp$; since we are only interested in the finite form factors, we can make the change of loop momentum $\ell \rightarrow \ell \mp k$. The numerators have terms with up to three powers of ℓ : the terms proportional to ℓ^3, ℓ^2 will contribute only to $G^{1,2}$, and those linear in ℓ vanish because they are odd under integration. We use the AVV case as a prototype, finding

$$\begin{aligned} \Gamma_{\text{AVV}}^{\rho\mu\nu}|_{\text{finite}} &= \int_0^1 dx \int_0^{1-x} dy \Gamma(3) \int_\ell \frac{4i}{(\ell^2 - \Delta)^3} \\ & \quad (\{(1-x-3y)k^\mu - 2yp^\mu\} \epsilon^{\rho\nu;pq} \\ & \quad + \{(1-3x-y)k^\nu - 2xq^\nu\} \epsilon^{\rho\mu;pq} \\ & \quad - \{(x-y)k^\rho + yp^\rho + xq^\rho\} \epsilon^{\mu\nu;pq}), \end{aligned} \quad (\text{A2})$$

where

$$\Delta = m_\psi^2 - x(1-x)q^2 - y(1-y)p^2 - 2xyp \cdot q - i\epsilon. \quad (\text{A3})$$

The loop integral evaluates to

$$\int_\ell \frac{1}{(\ell^2 - \Delta)^3} = -\frac{i}{(4\pi)^2 \Gamma(3)} \Delta^{-1}. \quad (\text{A4})$$

To match Eq. (A2) to the Rosenberg parameterization in Eq. (65), we apply the Schouten identity

$$k^\rho \epsilon^{\mu\nu\alpha\beta} + k^\mu \epsilon^{\nu\alpha\beta\rho} + k^\nu \epsilon^{\alpha\beta\rho\mu} + k^\alpha \epsilon^{\beta\rho\mu\nu} + k^\beta \epsilon^{\rho\mu\nu\alpha} = 0 \quad (\text{A5})$$

to the last line, which becomes

$$\begin{aligned} & \{(x-y)k^\mu - yp^\mu - xq^\mu\} \epsilon^{\rho\nu;pq} \\ & \quad - \{(x-y)k^\nu - yp^\nu - xq^\nu\} \epsilon^{\rho\mu;pq} + (\text{terms in } G_{\text{AVV}}^{1,2}). \end{aligned} \quad (\text{A6})$$

The above lead to

$$\begin{aligned}
F_3 &= - \int_0^1 dx \int_0^{1-x} dy y(1-y) \Delta^{-1}, \\
F_4 &= - \int_0^1 dx \int_0^{1-x} dy xy \Delta^{-1}, \\
F_5 &= \int_0^1 dx \int_0^{1-x} dy xy \Delta^{-1}, \\
F_6 &= \int_0^1 dx \int_0^{1-x} dy x(1-x) \Delta^{-1}, \quad (\text{A7})
\end{aligned}$$

from which we see that

$$\begin{aligned}
F_3(p, q) &= -F_6(q, p), \\
F_4(p, q) &= -F_5(p, q). \quad (\text{A8})
\end{aligned}$$

With the $F_{3,\dots,6}$ set,¹⁹ the next step is to express G^1, G^2 in terms of $F_{3,\dots,6}$ so that the vertex function can be written in terms of the finite form factors.²⁰ To relate G^1, G^2 to $F_{3,\dots,6}$, we contract $\tilde{\Gamma}$ with the momenta of A, B , or C —respectively, p_μ, q_ν , or $(p+q)_\rho$. From the Rosenberg parametrization of Eq. (65), we obtain the following expressions for the momentum-contracted coupling-stripped vertex functions:

$$\begin{aligned}
(p+q)_\rho \tilde{\Gamma}_{\{\mathbf{r}\}}^{\rho\mu\nu} &= \frac{1}{\pi^2} (G_{\{\mathbf{r}\}}^2 - G_{\{\mathbf{r}\}}^1) \epsilon^{\mu\nu; pq}, \\
-p_\mu \tilde{\Gamma}_{\{\mathbf{r}\}}^{\rho\mu\nu} &= \frac{1}{\pi^2} (G_{\{\mathbf{r}\}}^2 - p^2 F_3 - p \cdot q F_4) \epsilon^{\rho\nu; pq}, \\
-q_\nu \tilde{\Gamma}_{\{\mathbf{r}\}}^{\rho\mu\nu} &= \frac{1}{\pi^2} (G_{\{\mathbf{r}\}}^1 - p \cdot q F_5 - q^2 F_6) \epsilon^{\rho\mu; pq}. \quad (\text{A9})
\end{aligned}$$

However, we know that G^1, G^2 are not uniquely defined. To isolate their ambiguities, we first define the triangle vertex function with *unshifted* loop momentum (i.e., when $a = b = 0$ in Fig. 2)

$$\Gamma_{\{\mathbf{r}\}}^{\rho\mu\nu}(p, q) \equiv \tilde{\Gamma}_{\{\mathbf{r}\}}^{\rho\mu\nu}(p, q; z = 0, w = 0). \quad (\text{A10})$$

The difference $\tilde{\Gamma} - \Gamma$ encapsulates the ambiguity from shifting the momentum, and for any $\{\mathbf{r}\}$ with an odd number of axial couplings, evaluates to [40]

$$\begin{aligned}
[\tilde{\Gamma} - \Gamma]_{\{\mathbf{r}\}}^{\rho\mu\nu} &= \int_\ell a^\tau \partial_{\ell^\tau} \mathcal{F}_{\{\mathbf{r}\}}^{\rho\mu\nu}(\ell) = \frac{2i\pi^2}{(2\pi)^4} a^\tau \lim_{\ell \rightarrow \infty} \ell^2 \ell_\tau \mathcal{F}_{\{\mathbf{r}\}}^{\rho\mu\nu}(\ell) \\
&= \frac{1}{4\pi^2} \epsilon^{\rho\mu\nu\delta} a^\delta = \frac{1}{4\pi^2} \epsilon^{\rho\mu\nu\delta} (z p^\delta + w q^\delta), \quad (\text{A11})
\end{aligned}$$

where $\mathcal{F}_{\{\mathbf{r}\}}^{\rho\mu\nu}$ is the integrand in, e.g., Eq. (63) for the AVV case.

We proceed to directly calculate the left-hand sides of Eq. (A9) using the explicit form in Eq. (63). The integrands in each of these contractions can be massaged into terms differing only by a shift in loop momentum, which can then be evaluated using the analog of Eq. (A11). For example:

$$\begin{aligned}
q_\nu \Gamma_{\text{AVV}}^{\rho\mu\nu} &= \int_\ell \text{Tr} \left\{ \gamma_5 \gamma_\rho \frac{1}{\not{\ell} - \not{p} - m_\psi} \gamma_\mu \frac{1}{\not{\ell} - m_\psi} - \gamma_5 \gamma_\rho \frac{1}{\not{\ell} - m_\psi} \gamma_\mu \frac{1}{\not{\ell} + \not{p} - m_\psi} \right. \\
&\quad \left. + \gamma_5 \gamma_\rho \frac{1}{\not{\ell} - \not{q} - m_\psi} \gamma_\mu \frac{1}{\not{\ell} + \not{p} - m_\psi} - \gamma_5 \gamma_\rho \frac{1}{\not{\ell} - \not{p} - m_\psi} \gamma_\mu \frac{1}{\not{\ell} + \not{q} - m_\psi} \right\} \\
&= \frac{2i\pi^2}{(2\pi)^4} \lim_{\ell \rightarrow \infty} \left[p^\tau \ell^2 \ell_\tau \text{Tr} \left\{ \gamma_5 \gamma_\rho \frac{1}{\not{\ell}} \gamma_\mu \frac{1}{\not{\ell} + \not{p}} \right\} + (p-q)^\tau \ell^2 \ell_\tau \text{Tr} \left\{ \gamma_5 \gamma_\rho \frac{1}{\not{\ell} - \not{p}} \gamma_\mu \frac{1}{\not{\ell} + \not{q}} \right\} \right] \\
&= \frac{1}{2\pi^2} \lim_{\ell \rightarrow \infty} \frac{1}{\ell^2} [p^\tau \ell_\tau \epsilon^{\mu\rho; lp} + (p-q)^\tau \ell_\tau (\epsilon^{\mu\rho; lp} + \epsilon^{\mu\rho; lq} - \epsilon^{\mu\rho; pq}) + \mathcal{O}(m_\psi^2)] \\
&= -\frac{\epsilon^{\rho\mu; pq}}{4\pi^2}, \quad (\text{A12})
\end{aligned}$$

¹⁹Recall, $F_{3,\dots,6}$ are independent of the $\mathbf{r}_i \in \{A, V\}$, so the results of Eq. (A7) hold in general and are not specific to the AVV example.

²⁰In the case of massless loop fermions, we note that F_3 [F_6] suffers infrared divergences if $p^2 = 0$ [$q^2 = 0$]. This can be seen from Eqs. (A3) and (A7).

where we have used $\ell^\alpha \ell^\beta \rightarrow \ell^2 \eta^{\alpha\beta} / 4$ to simplify the penultimate line, and the $\epsilon^{\mu\rho; pq}$ term has only one power of ℓ in the numerator and therefore vanishes when we take $\ell \rightarrow \infty$.

Below, we list the complete sets of expressions for the AVV case,

$$\begin{aligned}
(p+q)_\rho \Gamma_{AVV}^{\rho\mu\nu} &= \frac{\epsilon^{\mu\nu;pq}}{4\pi^2} 4m_\psi^2 C_0(m_\psi^2), \\
-p_\mu \Gamma_{AVV}^{\rho\mu\nu} &= -\frac{\epsilon^{\rho\nu;pq}}{4\pi^2}, \\
-q_\nu \Gamma_{AVV}^{\rho\mu\nu} &= \frac{\epsilon^{\rho\mu;pq}}{4\pi^2},
\end{aligned} \tag{A13}$$

and the VAV case,

$$\begin{aligned}
(p+q)_\rho \Gamma_{VAV}^{\rho\mu\nu} &= 0, \\
-p_\mu \Gamma_{VAV}^{\rho\mu\nu} &= -\frac{\epsilon^{\rho\nu;pq}}{4\pi^2} (1 + 4m_\psi^2 C_0(m_\psi^2)), \\
-q_\nu \Gamma_{VAV}^{\rho\mu\nu} &= \frac{\epsilon^{\rho\mu;pq}}{4\pi^2}.
\end{aligned} \tag{A14}$$

Finally, we can fix $G_{\{r\}}^{1,2}$ by combining Eq. (A11) and the above contractions of $(p+q)_\rho$, p_μ , q_ν with unshifted $\Gamma_{\{r\}}^{\rho\mu\nu}$, then equating the sum with Eq. (A9).

APPENDIX B: GENERALIZED $f\bar{f} \rightarrow X\gamma$

We examine the amplitude for the s -channel (left-hand side) diagram of Fig. 3:

$$\begin{aligned}
i\mathcal{M}_s^{\mu\nu} &= \bar{v}(k_2) \left\{ \frac{ig}{c_W} \gamma^\sigma (q_Z^{V,f} - q_Z^{A,f} \gamma_5) \frac{-i}{s - m_Z^2} (\Pi_Z^\infty)_{\sigma\rho} \right. \\
&\quad \left. + ieQ^f \gamma^\sigma \frac{-i}{s} g_{\sigma\rho} \right\} u(k_1) \cdot \tilde{\Delta}^{\rho\mu\nu},
\end{aligned} \tag{B1}$$

where Q^f is the electromagnetic charge of f , $(q_Z^{V,f}, q_Z^{A,f})$ are the (vector, axial) charges of f to Z , and the Z propagator in unitary gauge is

$$(\Pi_Z^\infty)_{\sigma\rho} = g_{\sigma\rho} - \frac{(k_1 + k_2)_\sigma (p + q)_\rho}{m_Z^2}. \tag{B2}$$

The coupling ψ must be vectorlike.

Let us isolate the contribution from the intermediate Z :

$$\begin{aligned}
\mathcal{M}_{s,Z}^{\mu\nu} &= -i \frac{g}{c_W s - m_Z^2} \tilde{\Delta}^{\rho\mu\nu} \cdot \bar{v}(k_2) \left\{ \gamma_\rho (q_Z^{V,f} - q_Z^{A,f} \gamma_5) \right. \\
&\quad \left. + \frac{2m_f(p+q)_\rho}{m_Z^2} q_Z^{A,f} \gamma_5 \right\} u(k_1).
\end{aligned} \tag{B3}$$

As expected, in the case of the $f\bar{f}Z$ vector coupling, only the transverse part of the Z propagator contributes. Moreover, we focus on the two cases for which we expect a diverging amplitude: the AVV and VAV parts of the triangle vertex functions, i.e., an axial Z coupling and a vector X coupling or vice versa. Then

$$\tilde{\Delta}^{\rho\mu\nu} \rightarrow \frac{g}{c_W} g_X e Q^\psi \{ q_Z^{V,\psi} \kappa^{A,\psi} \tilde{\Gamma}_{VAV}^{\rho\mu\nu} + q_Z^{A,\psi} \kappa^{V,\psi} \tilde{\Gamma}_{AVV}^{\rho\mu\nu} \}, \tag{B4}$$

with q_Z^ψ, κ^ψ the charge of the fermion ψ to Z , X respectively.

Squaring the amplitude, summing over final polarizations, and averaging over initial spins and fermion colors N_c , we find²¹

$$\begin{aligned}
|\overline{\mathcal{M}_{s,Z}}|^2 &= \frac{1}{4(N_c)^2} \left(\left(\frac{g}{c_W} \right)^2 g_X e \right)^2 \frac{1}{(s - m_Z^2)^2} \left(g_{\mu_1\mu_2} - \frac{p_{\mu_1} p_{\mu_2}}{m_X^2} \right) g_{\nu_1\nu_2} (Q^\psi)^2 \{ q_Z^{V,\psi} \kappa^{A,\psi} \tilde{\Gamma}_{VAV}^{\rho\mu_1\nu_1} + q_Z^{A,\psi} \kappa^{V,\psi} \tilde{\Gamma}_{AVV}^{\rho\mu_1\nu_1} \} \\
&\quad \{ q_Z^{V,\psi} \kappa^{A,\psi} \tilde{\Gamma}_{VAV}^{\sigma\mu_2\nu_2} + q_Z^{A,\psi} \kappa^{V,\psi} \tilde{\Gamma}_{AVV}^{\sigma\mu_2\nu_2} \} T_{\rho\sigma},
\end{aligned} \tag{B5}$$

where $T_{\rho\sigma}$ is the trace over the external fermion part of the squared-amplitude,

$$\begin{aligned}
T_{\rho\sigma} &= ((q_Z^{V,f})^2 + (q_Z^{A,f})^2) \cdot 4 \left(\{ k_{1\rho} k_{2\sigma} + k_{1\sigma} k_{2\rho} \} - \frac{s}{2} g_{\rho\sigma} \right) + q_Z^{V,f} q_Z^{A,f} \cdot 8i\epsilon_{\rho\sigma; k_1 k_2} \\
&\quad + (q_Z^{A,f})^2 \cdot 16r_f \left(\frac{s}{2} g_{\rho\sigma} + \frac{s}{m_Z^2} \left(1 + \frac{1}{2} \frac{s}{m_Z^2} \right) (p+q)_\rho (p+q)_\sigma \right),
\end{aligned} \tag{B6}$$

with $r_i = m_i^2/s$. We can also simplify the contraction of the external polarization tensors and the triangle vertex functions:

²¹Below, we have assumed that the coupling-stripped vertex functions are real, i.e., that $F_{3,\dots,6}$ and $C_0(m_\psi^2)$ are real. For our purposes, we ignore the imaginary parts of these functions, which originate from the possibility of pair production of fermions appearing in the loop. They can be calculated using the Sokhotski–Plemelj formula applied to the integrands of Eqs. (A7) and (71) with Δ from Eq. (A3).

$$\begin{aligned} \pi^4 g_{\mu_1 \mu_2} g_{\nu_1 \nu_2} \Gamma_{\{\mathbf{r}\}}^{\rho \mu_1 \nu_1} \Gamma_{\{\mathbf{r}'\}}^{\sigma \mu_2 \nu_2} &= \mathcal{F}_T \left\{ \frac{1}{4r_X} (1-r_X)^2 s g^{\rho\sigma} - \frac{1}{2r_X} (1-r_X) \{p^\rho q^\sigma + q^\rho p^\sigma\} + q^\rho q^\sigma \right\} \\ &\quad - (2r_X G_{\{\mathbf{r}\}}^1 G_{\{\mathbf{r}'\}}^1 + (1-r_X) \{G_{\{\mathbf{r}\}}^1 G_{\{\mathbf{r}'\}}^2 + G_{\{\mathbf{r}\}}^2 G_{\{\mathbf{r}'\}}^1\}) s g^{\rho\sigma} \\ &\quad + 2G_{\{\mathbf{r}\}}^1 G_{\{\mathbf{r}'\}}^2 q^\rho p^\sigma + 2G_{\{\mathbf{r}\}}^2 G_{\{\mathbf{r}'\}}^1 p^\rho q^\sigma + 2G_{\{\mathbf{r}\}}^1 G_{\{\mathbf{r}'\}}^1 p^\rho p^\sigma + 2G_{\{\mathbf{r}\}}^2 G_{\{\mathbf{r}'\}}^2 q^\rho q^\sigma, \end{aligned} \quad (\text{B7})$$

$$-\pi^4 \frac{P_{\mu_1} P_{\mu_2}}{m_X^2} g_{\nu_1 \nu_2} \Gamma_{\{\mathbf{r}\}}^{\rho \mu_1 \nu_1} \Gamma_{\{\mathbf{r}'\}}^{\sigma \mu_2 \nu_2} = (\mathcal{F}_L - G_{\{\mathbf{r}\}}^2 G_{\{\mathbf{r}'\}}^2) \left\{ \frac{1}{4r_X} (1-r_X)^2 s g^{\rho\sigma} - \frac{1}{2r_X} (1-r_X) \{p^\rho q^\sigma + q^\rho p^\sigma\} + q^\rho q^\sigma \right\}, \quad (\text{B8})$$

where $\mathcal{F}_{T,L}$ contain products of F_i and $G_{\{\mathbf{r}\},\{\mathbf{r}'\}}^{1,2}$, and $\{\mathbf{r}\}, \{\mathbf{r}'\} \in \{\text{AVV}, \text{VAV}\}$.

In addition to Eqs. (A8), (66), and (67), we have an additional relation between the form factors in the Rosenberg parametrization by using Eq. (A9) and either Eq. (A13) or Eq. (A14):

$$r_X \cdot sF_3 = \frac{1}{2} + m_\psi^2 C_0(m_\psi^2) - (1-r_X) sF_4, \quad (\text{B9})$$

We can use this relation to simplify the expressions for $G^{1,2}$ in this case:

$$\begin{aligned} G_{\text{AVV}}^1 &= \frac{1}{4}(z+1) - \frac{1-r_X}{2} sF_4, \\ G_{\text{AVV}}^2 &= \frac{1}{4}(w+1) - \frac{1-r_X}{2} sF_4 + m_\psi^2 C_0(m_\psi^2), \end{aligned} \quad (\text{B10})$$

$$\begin{aligned} G_{\text{VAV}}^1 &= \frac{1}{4}(z+1) - \frac{1-r_X}{2} sF_4, \\ G_{\text{VAV}}^2 &= \frac{1}{4}(w+1) - \frac{1-r_X}{2} sF_4. \end{aligned} \quad (\text{B11})$$

Then only $F_{4,6}$ and $m_\psi^2 C_0(m_\psi^2)$ are independent. In terms of these functions, the two quantities $\mathcal{F}_{T,L}$ in Eqs. (B7) and (B8) can then be written as

$$\begin{aligned} \mathcal{F}_T &= \left(\frac{1}{2} + m_\psi^2 C_0(m_\psi^2) \right) \left[\frac{1}{2} + m_\psi^2 C_0(m_\psi^2) - sF_4 \right] \\ &\quad + r_X sF_4 \left[\frac{1}{2} + m_\psi^2 C_0(m_\psi^2) - sF_6 \right] \\ &\quad + r_X^2 sF_4 (sF_4 + sF_6) + r_X (sF_4 - sF_6) (G_{\{\mathbf{r}\}}^1 + G_{\{\mathbf{r}'\}}^1) \\ &\quad - \left[\frac{1}{2} + m_\psi^2 C_0(m_\psi^2) - (1-2r_X) sF_4 \right] (G_{\{\mathbf{r}\}}^2 + G_{\{\mathbf{r}'\}}^2), \end{aligned} \quad (\text{B12})$$

$$\begin{aligned} \mathcal{F}_L &= - \left[\frac{1}{2} + m_\psi^2 C_0(m_\psi^2) - \left(G_{\{\mathbf{r}\}}^2 + \frac{1-r_X}{2} sF_4 \right) \right] \\ &\quad \times \left[\frac{1}{2} + m_\psi^2 C_0(m_\psi^2) - \left(G_{\{\mathbf{r}'\}}^2 + \frac{1-r_X}{2} sF_4 \right) \right]. \end{aligned} \quad (\text{B13})$$

We contract Eq. (B6) with Eqs. (B7) and (B8) and take the limit of massless initial-state fermions, $r_f \rightarrow 0$. Inserting these results back into Eq. (B5), we obtain the expression at leading-order in $r_X \ll 1$:

$$\begin{aligned} \overline{|M_{s,Z}|^2} &\sim \frac{1}{4N_c^2 \pi^4} \left(\left(\frac{g}{c_W} \right)^2 g_X e \right)^2 \frac{1}{(s-m_Z^2)^2} \cdot ((q_Z^{\text{V},f})^2 + (q_Z^{\text{A},f})^2) \frac{s^2}{4r_X} \left(1 - \frac{2tu}{s^2} \right) \\ &\quad \cdot (Q^\psi)^2 \left[\sum_{\mathbf{r} \in \{\text{AVV}, \text{VAV}\}} q_Z^{\mathbf{r}_1, \psi} \kappa^{\mathbf{r}_2, \psi} (2G_{\{\mathbf{r}\}}^2 - sF_4) \right]^2. \end{aligned} \quad (\text{B14})$$

Of the two loop momentum shift parameters, we see from Eqs. (B10) and (B11) that only w appears in the second line. In order that the Ward identities for the photon and Z boson be satisfied, we must have $w = z = -1$. Examining the form factor combinations on the second line in the two limits $m_\psi^2 \rightarrow 0$ and $m_\psi^2 \rightarrow \infty$, we find:

$$\begin{aligned} 2G_{\text{AVV}}^2 - sF_4 &\rightarrow \begin{cases} -\frac{1}{2} & m_\psi^2 \rightarrow \infty \\ -1 & m_\psi^2 \rightarrow 0 \end{cases}, \\ 2G_{\text{VAV}}^2 - sF_4 &\rightarrow \begin{cases} 0 & m_\psi^2 \rightarrow \infty \\ -1 & m_\psi^2 \rightarrow 0 \end{cases}. \end{aligned} \quad (\text{B15})$$

For completeness, we provide expressions for the form factors in the following two limits. If the loop fermions are infinitely heavy, then at leading order we can discard the $p^2 = m_X^2$ term appearing in Eq. (A3), such that

$$r_\psi \rightarrow \infty: \begin{cases} m_\psi^2 C_0(m_\psi^2) \rightarrow -r_\psi \left(\text{Li}_2\left(\frac{2}{1+i\sqrt{4r_\psi-1}}\right) + \text{Li}_2\left(\frac{2}{1-i\sqrt{4r_\psi-1}}\right) \right) \rightarrow -\frac{1}{2} \\ sF_4 \rightarrow \frac{1}{2} + m_\psi^2 C_0(m_\psi^2) \rightarrow 0 \\ sF_6 \rightarrow 1 - \sqrt{4r_\psi-1} \operatorname{arccot}(\sqrt{4r_\psi-1}) \rightarrow 0 \end{cases}, \quad (\text{B16})$$

where Li_2 is the dilogarithm function. If the loop fermions are massless, then F_6 suffers an infrared divergence²²:

$$r_\psi \rightarrow 0: \begin{cases} m_\psi^2 C_0(m_\psi^2) \rightarrow 0 \\ sF_4 \rightarrow \frac{1-r_X+r_X \log(r_X)}{2(1-r_X)^2} \\ sF_6 \rightarrow \frac{1}{4}(1+2 \log \varepsilon) \quad (r_X \rightarrow 0) \end{cases}. \quad (\text{B17})$$

APPENDIX C: OFF-SHELL X -EXCHANGE AMPLITUDES

In this appendix, we compute the amplitude for $BB \rightarrow BB$ scattering (the BIM amplitude after [81]), for which the diagrams are shown in Fig. 4. This calculation illustrates the impact of longitudinal enhancement from the triple-gauge vertex when the Stückelberg field is off-shell, and it is analogous to WW scattering in the SM.

A simple setup that accommodates this process is the “ A - B ” model from [38]: this consists of a single Dirac fermion ψ with an axial-vector interaction to A and a vectorlike interaction to B . The vector field A has a Stückelberg-like mass term. In order to cancel anomalies, the model includes dimension-5 Peccei–Quinn local counterterms coupling the Stückelberg scalar field π to Chern–Pontryagin densities. The Lagrangian after performing the R_ξ gauge fixing procedure as in Eq. (18) is

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4} B_{\mu\nu} B^{\mu\nu} - \frac{1}{4} F_{\mu\nu} F^{\mu\nu} - \frac{1}{2\xi} (\partial_\mu A^\mu)^2 + \frac{1}{2} m_X^2 A_\mu A^\mu \\ & + \bar{\psi} i (\not{\partial} + i e \not{B} + i g_A \not{\gamma}_5) \psi - m_\psi \bar{\psi} \psi \\ & + \frac{C_A}{2m_X} \pi F_A^{\mu\nu} \tilde{F}_{A\mu\nu} + \frac{C_B}{2m_X} \pi B^{\mu\nu} \tilde{B}_{\mu\nu}. \end{aligned} \quad (\text{C1})$$

An example Feynman diagram for the Peccei–Quinn terms in the last line of Eq. (C1) is displayed in Fig. 5.²³

The diagram on the left-hand side of Fig. 4 with s -channel π exchange evaluates to

²²As previously mentioned, sF_6 also has imaginary part $\pi/4$.
²³We can assume that $e \ll g$ such that these diagrams dominate over the standard contribution from one-loop box diagrams of fermions in, e.g., light-by-light scattering.

$$\begin{aligned} i\mathcal{M}_1^{\mu_1\mu_2\nu_1\nu_2} = & i \frac{C_B}{m_X} \epsilon^{\mu_1\alpha_1\mu_2\alpha_2} (-ip_{1,\alpha_1}) (-ip_{2,\alpha_2}) \\ & \cdot \frac{i}{s - \xi m_X^2} \cdot i \frac{C_B}{m_X} \epsilon^{\nu_1\beta_1\nu_2\beta_2} (iq_{1,\beta_1}) (iq_{2,\beta_2}) \\ \mathcal{M}_1^{\mu_1\mu_2\nu_1\nu_2} = & -\frac{1}{s - \xi m_X^2} \frac{C_B^2}{m_X^2} \epsilon^{\mu_1\mu_2;p_1p_2} \epsilon^{\nu_1\nu_2;q_1q_2}. \end{aligned} \quad (\text{C2})$$

For the diagram on the right-hand side, we are interested in the part of the amplitude that involves axial couplings of the loop fermions to A ,

$$\begin{aligned} i\mathcal{M}_2^{\mu_1\mu_2\nu_1\nu_2} = & \tilde{\Delta}_{\text{AVV}}^{\lambda\mu_1\mu_2}(-p_1, -p_2) \frac{-i}{s - m_X^2} (\Pi_X^\xi)_{\lambda\rho} \cdot \tilde{\Delta}_{\text{AVV}}^{\rho\nu_1\nu_2}(q_1, q_2), \\ (\Pi_X^\xi)_{\lambda\rho} = & g_{\lambda\rho} - (1 - \xi) \frac{(p_1 + p_2)_\lambda (q_1 + q_2)_\rho}{s - \xi m_X^2}. \end{aligned} \quad (\text{C3})$$

Rewriting the R_ξ gauge propagator as in the second line of Eq. (23), we can evaluate the ξ -dependent longitudinal terms using the MCVF of Eq. (69) with $C \rightarrow A, A \rightarrow B$,

$$\begin{aligned} \mathcal{M}_2^{\mu_1\mu_2\nu_1\nu_2}|_\xi = & \frac{1}{s - \xi m_X^2} \frac{1}{m_X^2} \epsilon^{\mu_1\mu_2;p_1p_2} \epsilon^{\nu_1\nu_2;q_1q_2} \\ & \cdot \left(\frac{e^2 g}{4\pi^2} \right)^2 \{ (w - z) + 4m_\psi^2 C_0(m_\psi^2) \}^2. \end{aligned} \quad (\text{C4})$$

For the gauge dependence to cancel, we must have

$$C_B = \pm \frac{e^2 g}{4\pi^2} (w - z). \quad (\text{C5})$$

To satisfy the anomaly-free Ward identities in Eq. (69) for the B vector bosons, we must choose $w = 1, z = -1$. The remaining ξ -independent amplitude is \mathcal{M}_2 with the intermediate A propagator in “unitary” gauge:




FIG. 5. Example diagram from dimension-5 counterterms in last line of Eq. (C1) model.

$$\begin{aligned} \mathcal{M}^{\mu_1 \mu_2 \nu_1 \nu_2} &= \Delta_{\text{AVV}}^{\lambda \mu_1 \mu_2}(-p_1, -p_2) \\ &\cdot \frac{-1}{s - m_X^2} (\Pi_X^\infty)_{\lambda \rho} \cdot \tilde{\Delta}_{\text{AVV}}^{\rho \nu_1 \nu_2}(q_1, q_2), \end{aligned} \quad (\text{C6})$$

which can be broken up into its transverse and longitudinal parts as

$$\begin{aligned} \mathcal{M}_T^{\mu_1 \mu_2 \nu_1 \nu_2} &= -\frac{(e^2 g)^2}{s - m_X^2} \tilde{\Gamma}^{\lambda \mu_1 \mu_2}(-p_1, -p_2) \tilde{\Gamma}_\lambda^{\nu_1 \nu_2}(q_1, q_2), \\ \mathcal{M}_L^{\mu_1 \mu_2 \nu_1 \nu_2} &= -\frac{1}{s - m_X^2} \frac{C_B^2}{m_X^2} e^{\mu_1 \mu_2; p_1 p_2} e^{\nu_1 \nu_2; q_1 q_2}. \end{aligned} \quad (\text{C7})$$

where the subscript AVV is implicit. The squared amplitude, averaged and summed over initial and final states, is

$$\begin{aligned} |\overline{\mathcal{M}}_T|^2 &= \frac{1}{4} \left[\frac{(e^2 g)^2}{s - m_X^2} \right]^2 \tilde{\Gamma}^{\lambda \mu_1 \mu_2}(-p_1, -p_2) \tilde{\Gamma}_{\mu_1 \mu_2}^\rho(-p_1, -p_2) \tilde{\Gamma}_\lambda^{\nu_1 \nu_2}(q_1, q_2) \tilde{\Gamma}_{\rho \nu_1 \nu_2}(q_1, q_2), \\ |\overline{\mathcal{M}}_L|^2 &= \frac{1}{4} \left[\frac{1}{s - m_X^2} \frac{C_B^2}{m_X^2} \right]^2 e^{\mu_1 \mu_2; p_1 p_2} e^{\nu_1 \nu_2; q_1 q_2} e_{\mu_1 \mu_2; p_1 p_2} e_{\nu_1 \nu_2; q_1 q_2}, \\ 2\text{Re}(\overline{\mathcal{M}}_L^* \mathcal{M}_T) &= \frac{1}{2} \left[\frac{1}{s - m_X^2} \right]^2 (e^2 g)^2 \frac{C_B^2}{m_X^2} e^{\mu_1 \mu_2; p_1 p_2} \tilde{\Gamma}^{\lambda \mu_1 \mu_2}(-p_1, -p_2) e_{\nu_1 \nu_2; q_1 q_2} \tilde{\Gamma}_\lambda^{\nu_1 \nu_2}(q_1, q_2). \end{aligned} \quad (\text{C8})$$

Since $p^2 = q^2 = 0$, the evaluation of the vertex functions is simple in the BIM case. From Eq. (A8), we have both $F_5 = -F_4$, $F_3 = -F_6$. Then from Eq. (B10),

$$G_{\text{AVV}}^2|_{p^2=q^2=0} = -G_{\text{AVV}}^1|_{p^2=q^2=0} = \frac{1}{2} s F_4. \quad (\text{C9})$$

Finally, if we have massless loop fermions, then Eq. (B17) implies

$$s F_4|_{m_\psi=0} = \frac{1 - r_X + r_X \log(r_X)}{2(1 - r_X)^2}. \quad (\text{C10})$$

Then

$$\begin{aligned} |\overline{\mathcal{M}}_T|^2 &= \frac{(e^2 g)^4}{16} \frac{1}{(1 - r_X)^2} (s F_4)^4 \rightarrow \left(\frac{e^2 g}{4} \right)^4, \\ |\overline{\mathcal{M}}_L|^2 &= \frac{C_B^4}{16} \frac{1}{r_X^2 (1 - r_X)^2} \rightarrow \left(\frac{e^2 g}{4\pi^2} \right)^4 \frac{1}{r_X^2}, \\ \text{Re}(2\overline{\mathcal{M}}_L^* \mathcal{M}_T) &= \frac{(e^2 g)^2 C_B^2}{4} \frac{1}{r_X (1 - r_X)^2} \\ &\times \left(2G^1 G^2 + \frac{t}{s} (G^1 + G^2)^2 \right) \\ &\rightarrow -\left(\frac{e^2 g}{2\pi} \right)^4 \frac{1}{8r_X}. \end{aligned} \quad (\text{C11})$$

where arrows indicate the limit $r_X \rightarrow 0$ and we omitted the AVV subscript in the interference term for clarity.

Let us consider adding Wess-Zumino terms to the Lagrangian of Eq. (C1). These are

$$\begin{aligned} &\frac{1}{2} \epsilon_{\mu\nu\lambda\rho} A^\mu B^\nu (C'_A F_A^{\lambda\rho} + C'_B F_B^{\lambda\rho}) \\ &= -\epsilon_{\mu\nu\lambda\rho} A^\mu B^\nu (C'_A \partial^\rho A^\lambda + C'_B \partial^\rho B^\lambda). \end{aligned} \quad (\text{C12})$$

For the $BB \rightarrow BB$ process, only the C'_B coefficient is relevant; the amplitude is shown in Fig. 6. Along with Fig. 4, we have three additional diagrams where one or both fermion triangle loops in the diagram on the right-hand side of Fig. 4 is replaced with a three-boson vertex from Fig. 6.




FIG. 6. Feynman amplitude for Wess-Zumino term in Eq. (C12).

We then have

$$i\mathcal{M}_3^{\mu_1\mu_2\nu_1\nu_2} = C'_B e^2 g \{ e^{\lambda\mu_1\mu_2\alpha} (p_1 - p_2)_\alpha \tilde{\Gamma}^{\rho\nu_1\nu_2}(q_1, q_2) + \tilde{\Gamma}^{\lambda\mu_1\mu_2}(-p_1, -p_2) e^{\rho\nu_1\nu_2\beta}(q_1 - q_2)_\beta \} \cdot \frac{-i}{s - m_X^2} (\Pi_X^\xi)_{\lambda\rho},$$

$$i\mathcal{M}_4^{\mu_1\mu_2\nu_1\nu_2} = -(C'_B)^2 e^{\lambda\mu_1\mu_2\alpha} (p_1 - p_2)_\alpha e^{\rho\nu_1\nu_2\beta}(q_1 - q_2)_\beta \cdot \frac{-i}{s - m_X^2} (\Pi_X^\xi)_{\lambda\rho}. \quad (\text{C13})$$

Again decomposing the R_ξ propagator as in Eq. (23), we find a modified cancellation condition for gauge independence

$$C_B^2 = \left(\frac{e^2 g}{4\pi^2} \{ (w - z) + 4m_\psi^2 C_0(m_\psi^2) \} - 2C'_B \right)^2. \quad (\text{C14})$$

We are left to calculate the squared amplitude that is the sum of Eqs. (C7) and (C13), the latter with the replacement $\Pi_X^\xi \rightarrow \Pi_X^\infty$. Examining the longitudinal pieces as in the second line of Eq. (C7), we find after using the gauge independence condition above

$$\mathcal{M}_L^{\mu_1\mu_2\nu_1\nu_2} = -\frac{1}{s - m_X^2} \frac{C_B^2 - 4(C'_B)^2 + 4C'_B(\pm C_B + 2C'_B)}{m_X^2} e^{\mu_1\mu_2;\rho_1\rho_2} e^{\nu_1\nu_2;q_1q_2}, \quad (\text{C15})$$

which yields the same result as in Eq. (C11) in the relevant limit.

-
- [1] I. Brivio and M. Trott, The standard model as an effective field theory, *Phys. Rep.* **793**, 1 (2019).
- [2] P. Langacker, The physics of heavy Z' gauge bosons, *Rev. Mod. Phys.* **81**, 1199 (2009).
- [3] B. Batell, M. Pospelov, and A. Ritz, Exploring portals to a hidden sector through fixed targets, *Phys. Rev. D* **80**, 095024 (2009).
- [4] J. D. Bjorken, R. Essig, P. Schuster, and N. Toro, New fixed-target experiments to search for dark gauge forces, *Phys. Rev. D* **80**, 075018 (2009).
- [5] J. Jaeckel and A. Ringwald, The low-energy frontier of particle physics, *Annu. Rev. Nucl. Part. Sci.* **60**, 405 (2010).
- [6] J. Alexander *et al.*, Dark Sectors 2016 Workshop: Community Report, [arXiv:1608.08632](https://arxiv.org/abs/1608.08632).
- [7] M. Battaglieri *et al.*, US cosmic visions: New ideas in dark matter 2017: Community Report, in U.S. Cosmic Visions: New Ideas in Dark Matter, [arXiv:1707.04591](https://arxiv.org/abs/1707.04591).
- [8] N. Arkani-Hamed, D. P. Finkbeiner, T. R. Slatyer, and N. Weiner, A theory of dark matter, *Phys. Rev. D* **79**, 015014 (2009).
- [9] M. Pospelov and A. Ritz, Astrophysical signatures of secluded dark matter, *Phys. Lett. B* **671**, 391 (2009).
- [10] M. Pospelov, Secluded U(1) below the weak scale, *Phys. Rev. D* **80**, 095002 (2009).
- [11] A. E. Nelson and J. Scholtz, Dark light, dark matter and the misalignment mechanism, *Phys. Rev. D* **84**, 103501 (2011).
- [12] P. Arias, D. Cadamuro, M. Goodsell, J. Jaeckel, J. Redondo, and A. Ringwald, WISPy cold dark matter, *J. Cosmol. Astropart. Phys.* **06** (2012) 013.
- [13] P. W. Graham, J. Mardon, and S. Rajendran, Vector dark matter from inflationary fluctuations, *Phys. Rev. D* **93**, 103520 (2016).
- [14] P. Agrawal, N. Kitajima, M. Reece, T. Sekiguchi, and F. Takahashi, Relic abundance of dark photon dark matter, *Phys. Lett. B* **801**, 135136 (2020).
- [15] R. T. Co, A. Pierce, Z. Zhang, and Y. Zhao, Dark photon dark matter produced by axion oscillations, *Phys. Rev. D* **99**, 075002 (2019).
- [16] M. Bastero-Gil, J. Santiago, L. Ubaldi, and R. Vega-Morales, Vector dark matter production at the end of inflation, *J. Cosmol. Astropart. Phys.* **04** (2019) 015.
- [17] J. A. Dror, K. Harigaya, and V. Narayan, Parametric resonance production of ultralight vector dark matter, *Phys. Rev. D* **99**, 035036 (2019).
- [18] R. T. Co, K. Harigaya, and A. Pierce, Gravitational waves and dark photon dark matter from axion rotations, *J. High Energy Phys.* **12** (2021) 099.
- [19] B. Holdom, Two U(1)'s and epsilon charge shifts, *Phys. Lett.* **166B**, 196 (1986).
- [20] M. Fabbrichesi, E. Gabrielli, and G. Lanfranchi, *The Physics of the Dark Photon* (Springer, New York, 2021).
- [21] H. Ruegg and M. Ruiz-Altaba, The Stueckelberg field, *Int. J. Mod. Phys. A* **19**, 3265 (2004).
- [22] M. Reece, Photon masses in the landscape and the swampland, *J. High Energy Phys.* **07** (2019) 181.
- [23] B. W. Lee, C. Quigg, and H. B. Thacker, Weak interactions at very high-energies: The role of the Higgs boson mass, *Phys. Rev. D* **16**, 1519 (1977).

- [24] R. Delbourgo, S. Twisk, and G. Thompson, Massive Yang-Mills theory: Renormalizability versus unitarity, *Int. J. Mod. Phys. A* **03**, 435 (1988).
- [25] G. T. Bodwin, E. V. Kovacs, and J. H. Sloan, Renormalization counterterms in linearized lattice QED, *Phys. Rev. D* **44**, 3924 (1991).
- [26] C. Grosse-Knetter and R. Kogerler, Unitary gauge, Stueckelberg formalism and gauge invariant models for effective Lagrangians, *Phys. Rev. D* **48**, 2865 (1993).
- [27] B. Kors and P. Nath, A Stueckelberg extension of the standard model, *Phys. Lett. B* **586**, 366 (2004).
- [28] B. Kors and P. Nath, Aspects of the Stueckelberg extension, *J. High Energy Phys.* **07** (2005) 069.
- [29] C. Coriano, N. Irges, and S. Morelli, Stueckelberg axions and the effective action of anomalous Abelian models. I. A Unitarity analysis of the Higgs-axion mixing, *J. High Energy Phys.* **07** (2007) 008.
- [30] C. P. Burgess, J. P. Conlon, L.-Y. Hung, C. H. Kom, A. Maharana, and F. Quevedo, Continuous global symmetries and hyperweak interactions in string compactifications, *J. High Energy Phys.* **07** (2008) 073.
- [31] M. Goodsell, J. Jaeckel, J. Redondo, and A. Ringwald, Naturally light hidden photons in LARGE volume string compactifications, *J. High Energy Phys.* **11** (2009) 027.
- [32] L. Heisenberg, Generalization of the Proca action, *J. Cosmol. Astropart. Phys.* **05** (2014) 015.
- [33] J. Beltran Jimenez and L. Heisenberg, Derivative self-interactions for a massive vector field, *Phys. Lett. B* **757**, 405 (2016).
- [34] A. Kachanovich, S. Kovalenko, S. Kuleshov, V. E. Lyubovitskij, and A. S. Zhevlakov, Lepton phenomenology of Stueckelberg portal to dark sector, *Phys. Rev. D* **105**, 075004 (2022).
- [35] J. Preskill, Gauge anomalies in an effective field theory, *Ann. Phys. (N.Y.)* **210**, 323 (1991).
- [36] P. Anastasopoulos, M. Bianchi, E. Dudas, and E. Kiritsis, Anomalies, anomalous U(1)'s and generalized Chern-Simons terms, *J. High Energy Phys.* **11** (2006) 057.
- [37] J. A. Harvey, C. T. Hill, and R. J. Hill, Standard model gauging of the Wess-Zumino-Witten term: Anomalies, global currents and pseudo-Chern-Simons interactions, *Phys. Rev. D* **77**, 085017 (2008).
- [38] C. Coriano, M. Guzzi, and S. Morelli, Unitarity bounds for gauged axionic interactions and the Green-Schwarz mechanism, *Eur. Phys. J. C* **55**, 629 (2008).
- [39] R. Armillis, C. Coriano, M. Guzzi, and S. Morelli, Axions and anomaly-mediated interactions: The Green-Schwarz and Wess-Zumino vertices at higher orders and g-2 of the muon, *J. High Energy Phys.* **10** (2008) 034.
- [40] A. Racioppi, Anomalies, U(1)' and the MSSM, Ph. D thesis, University of Rome Tor Vergata, 2009.
- [41] I. Antoniadis, A. Boyarsky, S. Espahbodi, O. Ruchayskiy, and J. D. Wells, Anomaly driven signatures of new invisible physics at the Large Hadron Collider, *Nucl. Phys.* **B824**, 296 (2010).
- [42] B. A. Dobrescu and C. Frugiuele, Hidden GeV-Scale Interactions of Quarks, *Phys. Rev. Lett.* **113**, 061801 (2014).
- [43] J. A. Dror, R. Lasenby, and M. Pospelov, New Constraints on Light Vectors Coupled to Anomalous Currents, *Phys. Rev. Lett.* **119**, 141803 (2017).
- [44] A. Ismail, A. Katz, and D. Racco, On dark matter interactions with the Standard Model through an anomalous Z?, *J. High Energy Phys.* **10** (2017) 165.
- [45] J. A. Dror, R. Lasenby, and M. Pospelov, Dark forces coupled to nonconserved currents, *Phys. Rev. D* **96**, 075036 (2017).
- [46] A. Ekstedt, R. Enberg, G. Ingelman, J. Löfgren, and T. Mandal, Minimal anomalous U(1) theories and collider phenomenology, *J. High Energy Phys.* **02** (2018) 152.
- [47] N. Craig, I. Garcia Garcia, and G. D. Kribs, The UV fate of anomalous U(1)s and the Swampland, *J. High Energy Phys.* **11** (2020) 063.
- [48] J. A. Dror, Discovering leptonic forces using nonconserved currents, *Phys. Rev. D* **101**, 095013 (2020).
- [49] B. C. Allanach, B. Gripaios, and J. Tooby-Smith, Anomaly Cancellation with an Extra Gauge Boson, *Phys. Rev. Lett.* **125**, 161601 (2020).
- [50] L. Michaels and F. Yu, Probing new U(1) gauge symmetries via exotic $Z \rightarrow Z'\gamma$ decays, *J. High Energy Phys.* **03** (2021) 120.
- [51] Q. Bonnefoy, L. Di Luzio, C. Grojean, A. Paul, and A. N. Rossia, The anomalous case of axion EFTs and massive chiral gauge fields, *J. High Energy Phys.* **07** (2021) 189.
- [52] M. Ekhterachian, A. Hook, S. Kumar, and Y. Tsai, Bounds on gauge bosons coupled to nonconserved currents, *Phys. Rev. D* **104**, 035034 (2021).
- [53] C.-H. Chen, C.-W. Chiang, and C.-W. Su, Ultra-high-energy neutrino scattering in an anomalous U(1) effective field theory, *Phys. Lett. B* **827**, 136988 (2022).
- [54] J. Kumar, A. Rajaraman, and J. D. Wells, Probing the Green-Schwarz mechanism at the large hadron collider, *Phys. Rev. D* **77**, 066011 (2008).
- [55] S. Weinberg, *The Quantum Theory of Fields. Vol. 1: Foundations* (Cambridge University Press, Cambridge, England, 2005), p. 6.
- [56] T. Banks, *Modern Quantum Field Theory: A Concise Introduction* (Cambridge University Press, Cambridge, England, 2014), p. 12.
- [57] W. Pauli, Relativistic field theories of elementary particles, *Rev. Mod. Phys.* **13**, 203 (1941).
- [58] R. J. Glauber, On the gauge invariance of the neutral vector meson theory*, *Prog. Theor. Phys.* **9**, 295 (1953).
- [59] C. Becchi, A. Rouet, and R. Stora, Renormalization of gauge theories, *Ann. Phys. (N.Y.)* **98**, 287 (1976).
- [60] I. Tyutin, Gauge invariance in field theory and statistical physics in operator formalism, [arXiv:0812.0580](https://arxiv.org/abs/0812.0580).
- [61] N. Nakanishi, Covariant quantization of the electromagnetic field in the Landau gauge, *Prog. Theor. Phys.* **35**, 1111 (1966).
- [62] B. Lautrup, Canonical quantum electrodynamics in covariant gauges, *Kong. Dan. Vid. Sel. Mat. Fys. Med.* **35**, 11 (1967), <https://inspirehep.net/literature/1100145>.
- [63] A. Wulzer, An equivalent gauge and the equivalence theorem, *Nucl. Phys.* **B885**, 97 (2014).
- [64] G. Cuomo, L. Vecchi, and A. Wulzer, Goldstone equivalence and high energy electroweak physics, *SciPost Phys.* **8**, 078 (2020).
- [65] J. Bagger and C. Schmidt, Equivalence theorem redux, *Phys. Rev. D* **41**, 264 (1990).

- [66] H. Esseili and G. D. Kribs, Constraints on a flavor-independent generalized vector boson (to be published).
- [67] T. Appelquist and M. S. Chanowitz, Unitarity Bound on the Scale of Fermion Mass Generation, *Phys. Rev. Lett.* **59**, 2405 (1987); Erratum, *Phys. Rev. Lett.* **60**, 1589 (1988).
- [68] J. A. Dror, R. Lasenby, and M. Pospelov, Light vectors coupled to bosonic currents, *Phys. Rev. D* **99**, 055016 (2019).
- [69] A. Adams, N. Arkani-Hamed, S. Dubovsky, A. Nicolis, and R. Rattazzi, Causality, analyticity and an IR obstruction to UV completion, *J. High Energy Phys.* **10** (2006) 014.
- [70] C.-S. Chu, P.-M. Ho, and B. Zumino, NonAbelian anomalies and effective actions for a homogeneous space G/H , *Nucl. Phys.* **B475**, 484 (1996).
- [71] L. Rosenberg, Electromagnetic interactions of neutrinos, *Phys. Rev.* **129**, 2786 (1963).
- [72] S. Weinberg, *The Quantum Theory of Fields. Vol. 2: Modern Applications* (Cambridge University Press, Cambridge, England, 2013), p. 8.
- [73] A. Dedes and K. Suxho, Heavy fermion non-decoupling effects in triple gauge boson vertices, *Phys. Rev. D* **85**, 095024 (2012).
- [74] W. A. Bardeen and B. Zumino, Consistent and covariant anomalies in gauge and gravitational theories, *Nucl. Phys.* **B244**, 421 (1984).
- [75] C. T. Hill, Anomalies, Chern-Simons terms and chiral delocalization in extra dimensions, *Phys. Rev. D* **73**, 085001 (2006).
- [76] C. T. Hill, Lecture notes for massless spinor and massive spinor triangle diagrams, [arXiv:hep-th/0601155](https://arxiv.org/abs/hep-th/0601155).
- [77] M. B. Green and J. H. Schwarz, Anomaly cancellation in supersymmetric $D = 10$ gauge theory and superstring theory, *Phys. Lett.* **149B**, 117 (1984).
- [78] A. E. Nelson and N. Tetradis, Constraints on a new vector boson coupled to baryons, *Phys. Lett. B* **221**, 80 (1989).
- [79] C. D. Carone and H. Murayama, Possible Light $U(1)$ Gauge Boson Coupled to Baryon Number, *Phys. Rev. Lett.* **74**, 3122 (1995).
- [80] P. Fileviez Perez and M. B. Wise, Baryon and lepton number as local gauge symmetries, *Phys. Rev. D* **82**, 011901 (2010); Erratum, *Phys. Rev. D* **82**, 079901 (2010).
- [81] C. Bouchiat, J. Iliopoulos, and P. Meyer, An anomaly free version of Weinberg's model, *Phys. Lett.* **38B**, 519 (1972).
- [82] G. Jikia and A. Tkabladze, Photon-photon scattering at the photon linear collider, *Phys. Lett. B* **323**, 453 (1994).
- [83] G. J. Gounaris, P. I. Porfyriadis, and F. M. Renard, The gamma gamma \rightarrow gamma gamma process in the standard and SUSY models at high-energies, *Eur. Phys. J. C* **9**, 673 (1999).
- [84] S. R. Coleman and B. Grossman, 't Hooft's consistency condition as a consequence of analyticity and unitarity, *Nucl. Phys.* **B203**, 205 (1982).
- [85] W.-M. Chen, Y.-t. Huang, and D. A. McGady, Anomalies without an action, [arXiv:1402.7062](https://arxiv.org/abs/1402.7062).
- [86] N. Arkani-Hamed, L. Motl, A. Nicolis, and C. Vafa, The string landscape, black holes and gravity as the weakest force, *J. High Energy Phys.* **06** (2007) 060.
- [87] D. Harlow, B. Heidenreich, M. Reece, and T. Rudelius, The weak gravity conjecture: A review, [arXiv:2201.08380](https://arxiv.org/abs/2201.08380).
- [88] P. Draper, I. G. Garcia, and M. Reece, Snowmass White Paper: Implications of Quantum Gravity for Particle Physics, in 2022 Snowmass Summer Study (2022), [arXiv:2203.07624](https://arxiv.org/abs/2203.07624).
- [89] P. J. Fox, J. Liu, D. Tucker-Smith, and N. Weiner, An effective Z' , *Phys. Rev. D* **84**, 115006 (2011).
- [90] M. Ahlers, J. Jaeckel, J. Redondo, and A. Ringwald, Probing hidden sector photons through the Higgs window, *Phys. Rev. D* **78**, 075005 (2008).
- [91] B. Batell, M. Pospelov, and A. Ritz, Probing a secluded $U(1)$ at B-factories, *Phys. Rev. D* **79**, 115008 (2009).
- [92] K. Harigaya and Y. Nomura, Light chiral dark sector, *Phys. Rev. D* **94**, 035013 (2016).
- [93] J. Berger, K. Jedamzik, and D. G. E. Walker, Cosmological constraints on decoupled dark photons and dark Higgs, *J. Cosmol. Astropart. Phys.* **11** (2016) 032.
- [94] L. Darmé, S. Rao, and L. Roszkowski, Light dark Higgs boson in minimal sub-GeV dark matter scenarios, *J. High Energy Phys.* **03** (2018) 084.
- [95] C. Mondino, M. Pospelov, J. T. Ruderman, and O. Slone, Dark Higgs dark matter, *Phys. Rev. D* **103**, 035027 (2021).