

Linear response, Hamiltonian, and radiative spinning two-body dynamicsGustav Uhre Jakobsen^{✉*} and Gustav Mogull^{✉†}

*Institut für Physik und IRIS Adlershof, Humboldt Universität zu Berlin,
Zum Großen Windkanal 2, 12489 Berlin, Germany;
Max Planck Institute for Gravitational Physics (Albert Einstein Institute),
Am Mühlenberg 1, 14476 Potsdam, Germany;
and Kavli Institute for Theoretical Physics, University of California,
Santa Barbara, California 93106, USA*



(Received 27 October 2022; accepted 5 January 2023; published 16 February 2023)

Using the spinning, supersymmetric worldline quantum field theory formalism we compute the momentum impulse and spin kick from a scattering of two spinning black holes or neutron stars up to quadratic order in spin at third post-Minkowskian (PM) order, including radiation-reaction effects and with arbitrarily misaligned spin directions. Parts of these observables, both conservative and radiative, are also inferred from lower-PM scattering data by extending Bini and Damour's linear response formula to include misaligned spins. By solving Hamilton's equations of motion we also use a conservative scattering angle to infer a complete 3PM two-body Hamiltonian including finite-size corrections and misaligned spin-spin interactions. Finally, we describe mappings to the bound two-body dynamics for aligned spin vectors: including a numerical plot of the binding energy for circular orbits compared with numerical relativity, analytic confirmation of the NNLO PN binding energy, and the energy loss over successive orbits.

DOI: [10.1103/PhysRevD.107.044033](https://doi.org/10.1103/PhysRevD.107.044033)**I. INTRODUCTION**

The need for accurate waveform templates for comparison with gravitational wave signals coming from the LIGO, Virgo, and KAGRA detectors of binary merger events [1–6]—and in the future LISA, the Einstein Telescope, and Cosmic Explorer [7]—has provoked enormous interest in the gravitational two-body problem. One of the most important physical properties influencing the paths of massive objects following inspiral trajectories, which as they accelerate produce gravitational waves, is their spins. Accurately determining the spins of black holes and neutron stars in binary orbits yields crucial information about their origins: if the spins are approximately aligned with the orbital plane, then this suggests formation of the binary system by slow accretion of matter; if they are misaligned (precessing), then this indicates formation of the binary by a random capture event.

A fruitful path has been effective field theory (EFT)-based methods, which tackle the inspiral stage of the gravitational two-body problem using its natural separation

of length scales [8–12]: the size of the massive bodies is far less than their separation, which in turn is far less than their distance from us, the observer. Partial results for the nonspinning two-body Hamiltonian are up to sixth post-Newtonian (PN) order [13–18]; in the spinning case a body-fixed frame on the worldline is often used [19–22], and results are up to $N^3\text{LO}$ in the spin-orbit sector [23–25] and in the spin-spin sector [26–33].

However, an excellent alternative approach to the bound two-body problem comes by way of studying two-body scattering: here it is natural to define gauge-invariant scattering observables in terms of the states at past/future infinity, where the gravitational field is weak. It is also natural here to adopt the post-Minkowskian (PM) expansion in Newton's constant G , which resums terms from infinitely high velocities in the post-Newtonian series. One may use analytic continuation to directly produce PM observables for bound orbits [34–37]; alternatively, conservative scattering observables may be used to infer a Hamiltonian for the two-body system [38–44]. A more sophisticated version of this strategy is to infer an effective-one-body (EOB) Hamiltonian [45–49], which may be extended to include spin [50–54] and resums information from the test-body limit.

The worldline quantum field theory (WQFT) is a new formalism for producing gravitational scattering observables [55–63]. It builds on the highly successful PM-based worldline EFT approach [64], which has been used to produce scattering observables at 3PM [65–67] and 4PM

*gustav.uhre.jakobsen@physik.hu-berlin.de

†gustav.mogull@aei.mpg.de

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

orders [68–71]; the worldline EFT has also produced gravitational bremsstrahlung and radiative observables including tidal effects and spin [72–75]. The WQFT goes a step further by quantizing worldline degrees of freedom, which bypasses the need for intermediate off-shell objects such as the effective action. A supersymmetric extension to the worldline accounts for quadratic-in-spin effects [57,58], conveniently avoiding the typical use of a body-fixed frame. In Ref. [61] we used the WQFT to produce conservative scattering observables—the momentum impulse Δp_i^μ and spin kick ΔS_i^μ —at 3PM order.

In this paper, we upgrade these observables to include radiation-reaction (dissipative) effects, using the Schwinger-Keldysh in-in formalism [76–80] that has recently been incorporated into both the WQFT and PM-based worldline EFT frameworks [63,67]. Our results confirm the radiated four-momentum P_{rad}^μ recently predicted with the worldline EFT approach [75]. Given these new observables, we postulate and confirm an extension to Bini and Damour’s linear response relation [81–83] which allows us to predict terms in the conservative and radiative parts of the full scattering observables, depending on their behavior under the time-reversal operation $v_i^\mu \rightarrow -v_i^\mu$. The extension holds for arbitrary spin orientations, and goes beyond linear response.

The WQFT is inspired by QFT amplitudes-based methods for tackling the classical two-body problem [84–87]. These build on well-honed strategies for deriving scattering amplitudes [88–92] and performing the associated loop integrals [93,94]. In the nonspinning case a slew of two-body results have been produced at 3PM order (two loops) [95–102] and at 4PM order (three loops) [71,103,104]. There has also recently been work on N -body scattering and potentials [105]. Radiation-reaction effects have been incorporated [106–112], and an in-in style formalism for directly producing observables has been introduced [113–116]. To handle spin, higher-spin fields are used [117–126] and results have been produced at 2PM order at quadratic [52,127,128], quartic [129], and higher orders in spin [130–132]. Similar results have also been achieved with the closely related heavy-particle EFT [133–138].

Most notably, a 3PM quadratic-in-spin Hamiltonian has now been derived using amplitudes-based methods [139], involving spin on one of the two massive bodies only and without finite-size corrections. In this paper, using the conservative scattering observables derived in Ref. [61] we both confirm this result and extend it to include spin-spin effects and finite-size corrections relevant for neutron stars. Quite remarkably, we find that knowledge of a single scattering angle suffices to completely determine the Hamiltonian, also when arbitrarily misaligned spin vectors are involved.

Our paper is structured as follows. In Sec. II we review the dynamics of spinning massive bodies, including their description up to quadratic order in spin in terms of an

$\mathcal{N} = 2$ supersymmetric worldline action. We demonstrate how, with a suitable SUSY shift, we can switch between the canonical and covariant spin-supplementary conditions (SSCs). In Sec. III we review the Schwinger-Keldysh in-in formalism in the context of WQFT, and in Sec. IV we put it to use deriving the complete 3PM quadratic-in-spin momentum impulse Δp_1^μ and spin kick ΔS_1^μ including radiation-reaction effects. We present the results schematically, demonstrate how one may introduce scattering angles for misaligned spins, and perform various consistency checks.

Next, in Sec. V we upgrade the linear response relation to misaligned spin directions, generating both conservative and radiative terms from the full 3PM scattering observables Δp_1^μ and ΔS_1^μ . In Sec. VI we use the conservative scattering observables, and in particular the scattering angle, to build a complete 3PM quadratic-in-spin Hamiltonian. Finally, in Sec. VII we discuss unbound-to-bound mappings for the specific case of aligned spins: we generate the binding energy for circular orbits, both numerically and analytically and up to 4PN order, and produce plots of the binding energy as a function of the orbital frequency close to merger—comparing our results with numerical relativity. We also determine the energy radiated per orbit using an appropriate analytic continuation [37]. In Sec. VIII we conclude.

II. SPINNING MASSIVE BODIES

A pair of black holes or neutron stars interacting through a gravitational field in D -dimensional Einstein gravity is described by

$$S = S_{\text{EH}}[g_{\mu\nu}] + S_{\text{gf}}[g_{\mu\nu}] + \sum_{i=1}^2 S^{(i)}[g_{\mu\nu}, x_i^\mu, \psi_i^a], \quad (1)$$

where S_{EH} is the Einstein-Hilbert action ($\kappa = \sqrt{32\pi G}$),

$$S_{\text{EH}} = -\frac{2}{\kappa^2} \int d^D x \sqrt{-g} R, \quad (2)$$

S_{gf} is a gauge-fixing term, and $S^{(i)}$ are the two worldline actions. Up to quadratic order in spin [57,58]

$$\begin{aligned} \frac{S^{(i)}}{m_i} = & - \int d\tau_i \left[\frac{1}{2} g_{\mu\nu} \dot{x}_i^\mu \dot{x}_i^\nu + i \bar{\psi}_{i,a} \frac{D\psi_i^a}{D\tau_i} + \frac{1}{2} R_{abcd} \bar{\psi}_i^a \psi_i^b \bar{\psi}_i^c \psi_i^d \right. \\ & \left. + C_{E,i} E_{i,ab} \bar{\psi}_i^a \psi_i^b P_{i,cd} \bar{\psi}_i^c \psi_i^d \right], \end{aligned} \quad (3)$$

where the projector is $P_{i,ab} := \eta_{ab} - e_{a\mu} e_{b\nu} \dot{x}_i^\mu \dot{x}_i^\nu / \dot{x}_i^2$, η_{ab} is the (mostly minus) Minkowski metric, and $E_{i,ab} := R_{a\mu b\nu} \dot{x}_i^\mu \dot{x}_i^\nu / \dot{x}_i^2$. The finite-size multipole moment coefficients $C_{E,i}$ are defined such that $C_{E,i} = 0$ for black holes, and

$$\frac{D\psi_i^a}{D\tau_i} := \dot{x}_i^\mu \nabla_\mu \psi_i^a = \dot{\psi}_i^a + \dot{x}_i^\mu \omega_\mu^{ab} \psi_{i,b}. \quad (4)$$

The two bodies with masses m_i have positions $x_i^\mu(\tau_i)$; the complex anticommuting fields $\psi_i^a(\tau_i)$, defined in a local frame $e_\mu^a(x)$ with $g_{\mu\nu} = e_\mu^a e_\nu^b \eta_{ab}$, encode spin degrees of freedom.

The worldline action (3) enjoys a global $\mathcal{N} = 2$ supersymmetry:

$$\delta x_i^\mu = i\bar{\epsilon}_i \psi_i^\mu + i\epsilon_i \bar{\psi}_i^\mu, \quad \delta \psi_i^a = -\epsilon_i e_\mu^a \dot{x}_i^\mu - \delta x_i^\mu \omega_\mu^a{}_b \psi_i^b, \quad (5)$$

with constant SUSY parameters ϵ_i and $\bar{\epsilon}_i = \epsilon_i^\dagger$. As shown in Ref. [58], these shifts are generated by the conserved supercharges $\dot{x}_i \cdot \psi_i$ and $\dot{x}_i \cdot \bar{\psi}_i$. There is also a U(1) symmetry:

$$\delta \psi_i^a = i\alpha_i \psi_i^a, \quad \delta \bar{\psi}_i^a = -i\alpha_i \bar{\psi}_i^a, \quad \delta x_i^\mu = 0, \quad (6)$$

generated by the conserved charge $\psi_i \cdot \bar{\psi}_i$. Lastly, reparametrization invariance of the worldlines in τ_i implies

$$\dot{x}_i^2 = 1 + R_{abcd} \bar{\psi}_i^a \psi_i^b \bar{\psi}_i^c \psi_i^d + 2C_{E,i} E_{i,ab} \bar{\psi}_i^a \psi_i^b P_{i,cd} \bar{\psi}_i^c \psi_i^d \quad (7)$$

is also preserved. As $\dot{x}_i^2 \neq 1$ generically along the worldlines this implies that τ_i are not the proper times; however, as we are generally only interested in the asymptotic behavior this subtlety will not be important.

A. Background symmetries

Fields are perturbatively expanded around their background values at past infinity:

$$g_{\mu\nu}(x) = \eta_{\mu\nu} + \kappa h_{\mu\nu}(x), \quad (8a)$$

$$x_i^\mu(\tau_i) = b_i^\mu + \tau_i v_i^\mu + z_i^\mu(\tau_i), \quad (8b)$$

$$\psi_i^a(\tau_i) = \Psi_i^a + \psi_i^a(\tau_i), \quad (8c)$$

where $p_i^\mu = m_i v_i^\mu$ is the initial momentum; the initial value of the spin tensor is given by

$$S_i^{ab} = -2im_i \bar{\Psi}_i^{[a} \Psi_i^{b]}. \quad (9)$$

The antisymmetrization $[ab]$ includes a factor $1/2$ —note that this normalization of the spin tensor differs from that used in Refs. [57,58,61]. The vierbein is similarly expanded as

$$e_\mu^a = \eta^{a\nu} \left(\eta_{\mu\nu} + \frac{\kappa}{2} h_{\mu\nu} - \frac{\kappa^2}{8} h_{\mu\rho} h^\rho{}_\nu + \mathcal{O}(\kappa^3) \right), \quad (10)$$

which allows us to drop the distinction between spacetime μ, ν, \dots and local frame a, b, \dots indices. The global $\mathcal{N} = 2$ SUSY in the far past is

$$\begin{aligned} \delta b_i^\mu &= i\bar{\epsilon}_i \Psi_i^\mu + i\epsilon_i \bar{\Psi}_i^\mu, & \delta v_i^\mu &= 0, & \delta \Psi_i^\mu &= -\epsilon_i v_i^\mu, \\ &\Rightarrow \delta S_i^{\mu\nu} &= 2p_i^{[\mu} \delta b_i^{\nu]}. \end{aligned} \quad (11)$$

To fix these symmetries we find it convenient to enforce the covariant SSC:

$$p_i \cdot \Psi_i = 0 \Rightarrow p_{i,\mu} S_i^{\mu\nu} = 0. \quad (12)$$

Using the reparametrization symmetry we also enforce $v_i^2 = 1$ and $b \cdot v_i = 0$, where $b^\mu = b_2^\mu - b_1^\mu$ is the impact parameter pointing from the first to the second massive body. Finally, $\gamma = v_1 \cdot v_2$; we will also make use of unit-normalized ‘‘hatted’’ variables, e.g. $\hat{b}^\mu = b^\mu/|b|$.

The total initial angular momentum of the system is

$$\begin{aligned} J^{\mu\nu} &= L^{\mu\nu} + S_1^{\mu\nu} + S_2^{\mu\nu}, \\ L^{\mu\nu} &= 2b_1^{[\mu} p_1^{\nu]} + 2b_2^{[\mu} p_2^{\nu]}, \end{aligned} \quad (13)$$

where $L^{\mu\nu}$ is the orbital component. In this context, we see that the background symmetries (11) correspond simply to invariance of the system’s total angular momentum under shifts in the origins of the two bodies b_i^μ . We can also shift the center of our coordinate system $x^\mu \rightarrow x^\mu + a^\mu$, in which case $L^{\mu\nu} \rightarrow L^{\mu\nu} + 2a^{[\mu} p^{\nu]}$ as discussed in e.g. Ref. [140]. The orbital and spin angular momentum vectors, defined specifically in $D = 4$ dimensions, are invariant under these shifts:

$$L^\mu := \frac{1}{2} \epsilon^\mu{}_{\nu\rho\sigma} L^{\nu\rho} \hat{P}^\sigma = -\frac{1}{E} \epsilon^\mu{}_{\nu\rho\sigma} b^\nu p_1^\rho p_2^\sigma, \quad (14a)$$

$$S_i^\mu := m_i a_i^\mu = \frac{1}{2} \epsilon^\mu{}_{\nu\rho\sigma} S_i^{\nu\rho} v_i^\sigma, \quad (14b)$$

where

$$P^\mu = p_1^\mu + p_2^\mu, \quad (15a)$$

$$p^\mu = \frac{m_1 m_2}{E^2} [(\gamma m_1 + m_2) v_1^\mu - (\gamma m_2 + m_1) v_2^\mu] \quad (15b)$$

are respectively the total and center-of-mass (CoM) momentum, $p^\mu = (0, \mathbf{p}_\infty)$. Here $E = |P| = M\Gamma = M\sqrt{1+2\nu(\gamma-1)}$ is the energy in the CoM frame, $M = m_1 + m_2$, $\nu = \mu/M = m_1 m_2 / M^2$ are the total mass and symmetric mass ratio; $p_\infty = |\mathbf{p}_\infty| = \mu\sqrt{\gamma^2 - 1}/\Gamma$ is the center-of-mass momentum. With the covariant SSC choice (12) the total angular momentum J^μ is given by

$$\begin{aligned} J^\mu &:= \frac{1}{2} \epsilon^\mu{}_{\nu\rho\sigma} J^{\nu\rho} \hat{P}^\sigma \\ &= L^\mu + \sum_i (v_i \cdot \hat{P} S_i^\mu - S_i \cdot \hat{P} v_i^\mu). \end{aligned} \quad (16)$$

$$\begin{aligned}
 h_{\mu\nu}(k) &= -i \frac{m_i k}{2} e^{ik \cdot b_i} \delta(k \cdot v_i) \left(v_i^\mu v_i^\nu + ik_\rho S_i^{\rho(\mu} v_i^{\nu)} \right. \\
 &\quad \left. + \frac{1}{2} k_\rho k_\sigma S_i^{\rho\mu} S_i^{\nu\sigma} + \frac{C_{E,i}}{2} v_i^\mu v_i^\nu k_\rho S_i^{\rho\sigma} S_i^{\sigma\lambda} k^\lambda \right), \quad (29)
 \end{aligned}$$

where $\delta(\omega) := 2\pi\delta(\omega)$. At tree level the WQFT simply provides a diagrammatic mechanism for solving the classical equations of motion in momentum space, and so the use of retarded propagators ensures that boundary conditions are fixed in the far past.

IV. RADIATIVE OBSERVABLES

Building on Ref. [61], we compute the momentum impulse and change in ψ_i^μ :

$$\Delta p_i^\mu := [m_i \dot{x}_i^\mu]_{\tau_i=-\infty}^{\tau_i=+\infty} = -m_i \omega^2 \langle z_i^\mu(\omega) \rangle_{\text{in-in}}|_{\omega=0}, \quad (30a)$$

$$\Delta \psi_i^\mu := [\psi_i^\mu]_{\tau_i=-\infty}^{\tau_i=+\infty} = -i\omega \langle \psi_i^\mu(\omega) \rangle_{\text{in-in}}|_{\omega=0}, \quad (30b)$$

but now also including radiation-reaction effects. Using the definitions of the spin tensor $S_i^{\mu\nu}$ (9) and spin vector S_i^μ (14b) we can then also derive the spin kick ΔS_i^μ :

$$\begin{aligned}
 \Delta S_i^{\mu\nu} &= -2im_i (\bar{\Psi}_i^{\mu} \Delta \psi_i^{\nu}] + \Delta \bar{\psi}_i^{\mu} \Psi_i^{\nu}] + \Delta \bar{\psi}_i^{\mu} \Delta \psi_i^{\nu}], \\
 \Delta S_i^\mu &= \frac{1}{2m_i} \epsilon^\mu{}_{\nu\rho\sigma} (S_i^{\nu\rho} \Delta p_i^\sigma + \Delta S_i^{\nu\rho} p_i^\sigma + \Delta S_i^{\nu\rho} \Delta p_i^\sigma). \quad (31)
 \end{aligned}$$

We seek the 3PM components in a PM expansion:




FIG. 1. The 32 types of diagrams contributing to the $m_1^2 m_2^2$ components of $\Delta p_1^{(3)\mu}$ and the $m_1 m_2^2$ components of $\Delta \psi_1^{(3)\mu}$. Diagrams (a)–(v) were already present in the conservative calculation [61], though their expressions are modified by the inclusion of radiation; the mushrooms (w)–(ff) are purely radiative, and did not appear in the strictly conservative calculation [61]. Diagram (y) includes the same worldline propagator with opposite $i0$ prescriptions, and so belongs to the K integral family (35). For brevity we use solid lines to represent both propagating deflection z_i^μ and spin modes ψ_i^μ .

$$\Delta X = \sum_n G^n \Delta X^{(n)}, \quad (32)$$

where ΔX could be any of these observables: Δp_i^μ , ΔS_i^μ , $\Delta S_i^{\mu\nu}$, or $\Delta \psi_i^\mu$.

The relevant Feynman diagrams for both calculations are drawn in Fig. 1. These diagrams make no distinction between conservative and radiative effects. As only the $m_1^2 m_2^2$ component of $\Delta p_1^{(3)\mu}$ and the $m_1 m_2^2$ component of $\Delta \psi_1^{(3)\mu}$ are specifically affected by the inclusion of radiation-reaction effects we recompute these components; for the rest, we simply bring forward our previous results from Ref. [61]. Integrands are assembled using the WQFT Feynman rules [58] in $D = 4 - 2\epsilon$ dimensions, which involves integration on the momenta or energies of all internal lines; vertices contain either energy- or momentum-conserving δ functions, whichever is appropriate. The energy integrals, corresponding to internal propagation of z_i^μ or ψ_i^μ modes, are trivial: conservation of energy at the worldline vertices resolves them immediately.

Each graph has three unresolved four-momenta to integrate over. The first of these integrals is a Fourier transform:

$$\begin{aligned}
 \Delta X(b^\mu, v_i^\mu, S_i^{\mu\nu}) \\
 = \int_q e^{iq \cdot b} \delta(q \cdot v_1) \delta(q \cdot v_2) \Delta X(q^\mu, v_i^\mu, S_i^{\mu\nu}), \quad (33)
 \end{aligned}$$

where q^μ is the total momentum exchanged from the second to the first worldline, and $\int_q := \int d^D q / (2\pi)^D$.

Here we have implicitly defined the momentum-space observables $\Delta X(q^\mu, v_i^\mu, S_i^{\mu\nu})$, which are given as linear combinations of two-loop Feynman integrals:

$$\begin{aligned}
& I_{n_1, n_2, \dots, n_7}^{(\sigma_1; \sigma_2; \sigma_3)} [\ell_1^{\mu_1} \dots \ell_1^{\mu_n} \ell_2^{\nu_1} \dots \ell_2^{\nu_n}] \\
& := \int_{\ell_1, \ell_2} \frac{\delta(\ell_1 \cdot v_2) \delta(\ell_2 \cdot v_1) \ell_1^{\mu_1} \dots \ell_1^{\mu_n} \ell_2^{\nu_1} \dots \ell_2^{\nu_n}}{D_1^{n_1} D_2^{n_2} \dots D_7^{n_7}}, \\
& D_1 = \ell_1 \cdot v_1 + \sigma_1 i0, \quad D_2 = \ell_2 \cdot v_2 + \sigma_2 i0, \\
& D_3 = (\ell_1 + \ell_2 - q)^2 + \sigma_3 \text{sgn}(\ell_1^0 + \ell_2^0 - q^0) i0, \\
& D_4 = \ell_1^2, \quad D_5 = \ell_2^2, \quad D_6 = (\ell_1 - q)^2, \\
& D_7 = (\ell_2 - q)^2. \tag{34}
\end{aligned}$$

These integrals with retarded propagators were discussed at length in Ref. [63]: propagators D_4 – D_7 are prevented from going on-shell by the requirement that $\ell_1 \cdot v_2 = \ell_2 \cdot v_1 = 0$, so we can safely ignore their $i0$ prescriptions. We also require

$$\begin{aligned}
& K_{n_1, n_2, n_3, n_4, n_5}^{(\sigma)} [\ell^{\mu_1} \dots \ell^{\mu_n} k^{\nu_1} \dots k^{\nu_n}] \\
& := \int_{\ell, k} \frac{\delta((k - \ell) \cdot v_1) \delta(\ell \cdot v_2) \ell^{\mu_1} \dots \ell^{\mu_n} k^{\nu_1} \dots k^{\nu_n}}{D_1^{n_1} D_2^{n_2} D_3^{n_3} D_4^{n_4} D_5^{n_5}}, \\
& D_1 = \ell \cdot v_1 + i0, \quad D_2 = \ell \cdot v_1 - i0, \\
& D_3 = k^2 + \sigma \text{sgn}(k^0) i0, \quad D_4 = \ell^2, \quad D_5 = (\ell - q)^2, \tag{35}
\end{aligned}$$

which accounts for the possibility of a worldline propagator appearing twice, but with different $i0$ prescriptions—diagram (y) in Fig. 1.

The subsequent integration steps were discussed in Ref. [61], and are not substantially different with the inclusion of radiation-reaction effects in the observables. Tensorial two-loop integrals are reduced to scalar type by expanding on a suitable basis, and then reduced to master integrals using integration-by-parts identities. Expressions for these master integrals were provided in Ref. [63], and once the Fourier transform (33) has been performed on the exchanged momentum q^μ we are left with the observables in D dimensions. The scalar integrals themselves have simple reality properties:

$$\begin{aligned}
& I_{n_1, n_2, \dots, n_7}^{(\sigma_1; \sigma_2; \sigma_3)*} = (-1)^{n_1 + n_2} I_{n_1, n_2, \dots, n_7}^{(\sigma_1; \sigma_2; \sigma_3)}, \\
& K_{n_1, n_2, \dots, n_5}^{(\sigma)*} = (-1)^{n_1 + n_2} K_{n_1, n_2, \dots, n_5}^{(\sigma)}, \tag{36}
\end{aligned}$$

i.e. they are either purely real or imaginary, depending on whether they have an even or odd number of worldline propagators respectively. While this implies that the momentum-space observables $\Delta X(q^\mu, v_i^\mu, S_i^{\mu\nu})$ are complex functions, the Fourier transform (33) introduces additional factors of i , giving rise to purely real observables $\Delta X(b^\mu, v_i^\mu, S_i^{\mu\nu})$.

The final observables $\Delta p_i^{(3)\mu}$ and $\Delta \psi_i^{(3)\mu}$ in four dimensions are found by taking the limit $D \rightarrow 4$, checking to ensure the cancellation of all poles in the dimensional regularization parameter $\epsilon = 2 - \frac{D}{2}$. We generated our integrand in $D = 4 - 2\epsilon$ dimensions in order to account for possible cancellations of ϵ poles with the two-loop integrals. Like in Ref. [61] we have verified that the supercharges p_i^2 , $\psi_i \cdot \bar{\psi}_i$, $p_i \cdot \psi_i$, and $p_i \cdot \bar{\psi}_i$ are conserved. This means that the following identities,

$$\begin{aligned}
0 &= p_1 \cdot \Delta p_1^{(3)} + \Delta p_1^{(1)} \cdot \Delta p_1^{(2)}, \\
0 &= \bar{\Psi}_1 \cdot \Delta \psi_1^{(3)} + \Delta \bar{\psi}_1^{(3)} \cdot \Psi_1 + \Delta \bar{\psi}_1^{(1)} \cdot \Delta \psi_1^{(2)} + \Delta \bar{\psi}_1^{(2)} \cdot \Delta \psi_1^{(1)}, \\
0 &= p_1 \cdot \Delta \psi_1^{(3)} + \Delta p_1^{(3)} \cdot \Psi_1 + \Delta p_1^{(1)} \cdot \Delta \psi_1^{(2)} + \Delta p_1^{(2)} \cdot \Delta \psi_1^{(1)}. \tag{37}
\end{aligned}$$

are satisfied.

A. Results

We find it convenient to decompose the observables into four gauge-invariant parts each:

$$\Delta p_i^\mu = \Delta p_{i, \text{cons}}^{(+)\mu} + \Delta p_{i, \text{cons}}^{(-)\mu} + \Delta p_{i, \text{rad}}^{(+)\mu} + \Delta p_{i, \text{rad}}^{(-)\mu}, \tag{38a}$$

$$\Delta S_i^{\mu\nu} = \Delta S_{i, \text{cons}}^{(+)\mu\nu} + \Delta S_{i, \text{cons}}^{(-)\mu\nu} + \Delta S_{i, \text{rad}}^{(+)\mu\nu} + \Delta S_{i, \text{rad}}^{(-)\mu\nu}. \tag{38b}$$

The split into conservative ‘‘cons’’ and radiative ‘‘rad’’ pieces is done with respect to the integrals (34), (35): the potential and radiative regions [63]. Meanwhile the split into (\pm) sectors is defined with respect to behavior under a time-reversal operation:

$$\Delta X^{(\pm)}|_{v_i^\mu \rightarrow -v_i^\mu} = \pm \Delta X^{(\pm)}, \tag{39}$$

which flips the signs on the timelike vectors v_i^μ (and the momenta $p_i^\mu = m_i v_i^\mu$) but not the spacelike vectors b^μ and a_i^μ , leaving $\gamma = v_1 \cdot v_2$ invariant. Under this operation, $S_i^{\mu\nu} = \epsilon^{\mu\nu\rho\sigma} p_i^\rho a_i^\sigma$ changes sign.

For the impulse, we have

$$\begin{aligned}
\Delta p_{1, \text{cons}}^{(3; +)\mu} &= \frac{m_1^2 m_2^2}{|b|^3} \left[c_1^{(+)\mu} \frac{\text{arccosh} \gamma}{\sqrt{\gamma^2 - 1}} + \sum_{n=1}^3 \left(\frac{m_1}{m_2} \right)^{n-2} c_{n+1}^{(+)\mu} \right], \\
\Delta p_{1, \text{rad}}^{(3; +)\mu} &= \frac{m_1^2 m_2^2}{|b|^3} \mathcal{I}(v) c_5^{(+)\mu} \\
\Delta p_{1, \text{cons}}^{(3; -)\mu} &= \sum_{n=1}^3 \frac{\pi m_1^2 m_2^2}{|b|^3} \left(\frac{m_1}{m_2} \right)^{n-2} c_n^{(-)\mu}, \\
\Delta p_{1, \text{rad}}^{(3; -)\mu} &= \frac{\pi m_1^2 m_2^2}{|b|^3} \\
&\quad \times \left[c_4^{(-)\mu} + c_5^{(-)\mu} \frac{\text{arccosh} \gamma}{\sqrt{\gamma^2 - 1}} + c_6^{(-)\mu} \log \left(\frac{1 + \gamma}{2} \right) \right], \tag{40}
\end{aligned}$$

where

$$\mathcal{I}(v) = -\frac{8}{3} + \frac{1}{v^2} + \frac{(3v^2 - 1)}{v^3} \operatorname{arccosh}(\gamma) \quad (41)$$

is a universal prefactor, and $v = \sqrt{\gamma^2 - 1}/\gamma$. The vectors are given by

$$c_n^{(\pm)\mu} = \vec{f}_n^{(\pm)}(\gamma, C_{E,i}) \cdot \vec{\rho}_\pm^\mu, \quad (42)$$

with basis elements even/odd under time reversal:

$$\vec{\rho}_+^\mu = \left\{ \hat{b}^\mu, \frac{a_i \cdot \hat{L}}{|b|} \hat{b}^\mu, \frac{a_i \cdot \hat{b}}{|b|} \hat{L}^\mu, \frac{a_i \cdot a_j}{|b|^2} \hat{b}^\mu, \frac{a_i \cdot \hat{b} a_j \cdot \hat{b}}{|b|^2} \hat{b}^\mu, \right. \\ \left. \frac{a_i \cdot v_i a_j \cdot v_j}{|b|^2} \hat{b}^\mu, \frac{a_i \cdot \hat{b} a_j \cdot \hat{L}}{|b|^2} \hat{L}^\mu, \frac{a_i \cdot \hat{b} a_j \cdot v_j}{|b|^2} v_k^\mu \right\}, \quad (43a)$$

$$\vec{\rho}_-^\mu = \left\{ v_i^\mu, \frac{a_i \cdot \hat{L}}{|b|} v_j^\mu, \frac{a_i \cdot v_i}{|b|} \hat{L}^\mu, \frac{a_i \cdot a_j}{|b|^2} v_k^\mu, \frac{a_i \cdot v_i a_j \cdot v_j}{|b|^2} v_k^\mu, \right. \\ \left. \frac{a_i \cdot \hat{b} a_j \cdot \hat{b}}{|b|^2} v_k^\mu, \frac{a_i \cdot v_i a_j \cdot \hat{L}}{|b|^2} \hat{L}^\mu, \frac{a_i \cdot v_i a_j \cdot \hat{b}}{|b|^2} \hat{b}^\mu \right\}, \quad (43b)$$

where $i, j, k = 1, 2$ and $\bar{1} = 2, \bar{2} = 1$, $\hat{L}^\mu = L^\mu/|L|$ and $\hat{b}^\mu = b^\mu/|b|$ being unit-normalized vectors. Except for denominators of $(\gamma^2 - 1)^{n/2}$ the remaining scalar components $\vec{f}_n^{(\pm)}$ are polynomials in γ , $C_{E,i}$, so we refrain from providing explicit expressions in the text; instead, we refer the reader to the Supplemental Material [145] for full expressions.

Let us remark on certain properties of this result. $\Delta p_i^{(3;+)\mu}$ and $\Delta p_i^{(3;-)\mu}$ are respectively associated with the real (imaginary) integrals (36), i.e. those with an even (odd) number of worldline propagators. The factors of π in $\Delta p_i^{(3;-)\mu}$ thus arise from the overall factors of $i\pi$ in the purely imaginary master integrals—see Ref. [63]. We also note the behavior under $v \rightarrow -v$: in each case, the radiative components pick up the opposite sign from the conservative components. Finally, the function $\mathcal{I}(v)$ is familiar: it appears in the 2PM radiated angular momentum (56). As we shall see in Sec. V, $\Delta p_{i,\text{rad}}^{(3;+)\mu}$ and $\Delta p_{i,\text{cons}}^{(3;-)\mu}$ can be inferred directly from lower-PM observables, using a generalization to the linear response relation [81–83].

From the impulse Δp_i^μ we straightforwardly recover the four-momentum radiated from the scattering event:

$$P_{\text{rad}}^\mu = -\Delta P^\mu = -\Delta p_1^\mu - \Delta p_2^\mu, \quad (44)$$

which vanishes if we consider only conservative scattering. Here we agree with a recent 3PM worldline EFT result for P_{rad}^μ obtained by Riva, Vernizzi, and Wong [75]. We also agree with our own previous result for the leading-order radiated energy in the CoM frame $E_{\text{rad}} = \hat{P} \cdot P_{\text{rad}}$,

produced in collaboration with Plefka and Steinhoff [57], in which performing the required integrals necessitated a PN expansion—now we no longer need to do so. While knowledge of both θ and P_{rad}^μ allows one to reconstruct Δp_i^μ for aligned spins, this is not true in general—thus, in this work we fill in the missing pieces from Ref. [75]. However, we note that a corresponding expression for J_{rad}^μ at 3PM order is still lacking [the leading-order 2PM is known, see Eq. (56) below].

The 3PM spin kick takes a similar form as the impulse:

$$\Delta S_{1,\text{cons}}^{(3;-)\mu} = \frac{m_1^2 m_2^2}{|b|^3} \left[d_1^{(-)\mu} \frac{\operatorname{arccosh} \gamma}{\sqrt{\gamma^2 - 1}} + \sum_{n=1}^3 \left(\frac{m_1}{m_2} \right)^{n-2} d_{n+1}^{(-)\mu} \right], \\ \Delta S_{1,\text{rad}}^{(3;-)\mu} = \frac{m_1^2 m_2^2}{|b|^3} \mathcal{I}(v) d_5^{(-)\mu} \\ \Delta S_{1,\text{cons}}^{(3;+)\mu} = \sum_{n=1}^3 \frac{\pi m_1^2 m_2^2}{|b|^3} \left(\frac{m_1}{m_2} \right)^{n-2} d_n^{(+)\mu}, \\ \Delta S_{1,\text{rad}}^{(3;+)\mu} = \frac{\pi m_1^2 m_2^2}{|b|^3} \\ \times \left[d_4^{(+)\mu} + d_5^{(+)\mu} \frac{\operatorname{arccosh} \gamma}{\sqrt{\gamma^2 - 1}} + d_6^{(+)\mu} \log \left(\frac{1 + \gamma}{2} \right) \right], \quad (45)$$

but in this case $\Delta S_1^{(3;-)\mu}$ is associated with real integrals and $\Delta S_1^{(3;+)\mu}$ with imaginary integrals (36). The vectors d_n^μ, \tilde{d}_n^μ are given by

$$d_n^{(\pm)\mu} = \vec{g}_n^{(\pm)}(\gamma, C_{E,i}) \cdot \vec{\rho}_\pm^\mu, \quad (46)$$

and involve the same basis of vectors (43). Again, we refer the interested reader to the Supplemental Material [145] for fully explicit results.

B. Scattering angles

Conservative dynamics with spin vectors aligned to the scattering plane are described by a single angle:

$$\Delta p_{1,\text{cons}}^\mu = p^\mu (\cos \theta_{\text{cons}} - 1) + p_\infty \hat{b}^\mu \sin \theta_{\text{cons}}, \quad (47)$$

where p^μ is the CoM momentum (15b). However, generic spins and radiative effects both require a generalization of this simple parametrization. Generic spins result in non-planar motion and a nonzero spin kick; radiative effects in loss of total four-momentum P_{rad}^μ —we will present a more generic parametrization below (57).

For generic misaligned spin directions the impulse and spin kick are parametrized in spherical coordinates in terms of several angles. We focus on the following two, including radiation-reaction effects:

$$\sin\left(\frac{\theta_1}{2}\right) = \frac{|\Delta p_1|}{2p_\infty}, \quad \sin\left(\frac{\theta_2}{2}\right) = \frac{|\Delta p_2|}{2p_\infty}, \quad (48a)$$

$$\sin(\phi_1) = \frac{\hat{b} \cdot \Delta p_1}{p_\infty}, \quad \sin(\phi_2) = -\frac{\hat{b} \cdot \Delta p_2}{p_\infty}. \quad (48b)$$

The conservative counterparts of these angles, θ_{cons} and ϕ_{cons} , are defined by instead inserting $\Delta p_{i,\text{cons}}^\mu$ on the right-hand side. In this case the particle label on the angles is superficial as $\Delta p_{1,\text{cons}}^\mu = -\Delta p_{2,\text{cons}}^\mu$. For aligned spins at 3PM order these two definitions are equivalent to each other: $\theta_i = \phi_i$, which using Eq. (47) holds to all PM orders for strictly conservative scattering. Up to linear order in spin the angles equate even for misaligned spin vectors: $\theta_i = \phi_i + \mathcal{O}(S^2)$, as dependence on the spin vectors S_i^μ only enters through the spin-orbit terms $L \cdot S_i$. Surprisingly though, and in contrast to the use of spherical coordinates, we will find that only one of these angles suffices to fully describe the conservative impulse and spin kick.

In the conservative case the interpretation of θ_{cons} and ϕ_{cons} is simple. First, ϕ_{cons} measures the total scattering angle in the CoM frame in the plane spanned by \hat{b}^μ and \hat{p}^μ . Second, θ_{cons} measures the total angle between the initial and final momentum (which may point out of the initial plane). Including radiation-reaction we may still compute θ_i and ϕ_i using Eqs. (48) although their physical interpretation as scattering angles is less clear. We also note that, while θ_i is independent of the choice of SSC, ϕ_i is not due to the manifest dependence on b^μ , which transforms under SUSY shifts. To put it another way: the notion of initial plane of scattering depends on the SSC. For this reason we will mostly focus on θ_{cons} and later use it to parametrize the Hamiltonian in Sec. VI.

We expand the angle θ_i in G and spins:

$$\begin{aligned} \frac{\theta_k}{\Gamma} = & \sum_{n=1}^3 \left(\frac{GM}{|b|}\right)^n \left[\theta^{(n;0)} - \sum_i \theta^{(n;1,i)} \frac{\hat{L} \cdot a_i}{|b|} \right. \\ & + \sum_{i,j} \frac{a_i^\mu a_j^\nu}{|b|^2} (-\theta^{(n;2,1,i,j)} \eta^{\mu\nu} + \theta^{(n;2,2,i,j)} \hat{b}^\mu \hat{b}^\nu \\ & \left. + \theta^{(n;2,3,i,j)} v_i^\mu v_j^\nu + \theta_k^{(n;2,4,i,j)} \hat{b}^\mu v_j^\nu \right] + \mathcal{O}(S^3, G^4). \quad (49) \end{aligned}$$

Here i and j take the values 1,2. The coefficients $\theta^{(n;A)}$ are functions only of γ , ν , and $C_{E,i}$. Note that only the final coefficients $\theta_k^{(n;2,4,i,j)}$ depend on the particle label. The coefficients $\theta^{(3;A)}$ are provided in Appendix B and full expressions for the angles in the Supplemental Material [145]. Expansion coefficients for θ_{cons} , i.e. $\theta_{\text{cons}}^{(n;A)}$, are defined in an equivalent manner.

For aligned spins we verify our results for $\theta_1 = \theta_2$ against several results from the literature. First, we reproduce the result of our earlier work [61] where the radiative

part of θ_i was computed using linear response (see Sec. V) and has subsequently been extended to all spin orders by Alessio and Di Vecchia [146]. Second, we match our results for the probe limit and comparable-mass PN results [52,147,148]. Finally, for misaligned spin vectors in the high-energy limit where we let $\gamma \rightarrow \infty$ while keeping E and a_i^μ constant we recover a finite result:

$$\begin{aligned} \theta_i = & 4 \frac{GE}{|b|} \left[1 + \frac{\hat{L} \cdot a_+}{|b|} - \frac{2a_+^2 + 3(\hat{b} \cdot a_+)^2}{2|b|^2} \right. \\ & + \sum_j C_{E,j} \frac{a_j^2 + 2(\hat{b} \cdot a_j)^2}{|b|^2} \left. \right] + \frac{32}{3} \left(\frac{GE}{|b|}\right)^3 \left[1 + 3 \frac{\hat{L} \cdot a_+}{|b|} \right. \\ & - \frac{3}{20} \frac{41a_+^2 + a_-^2 + 50(\hat{b} \cdot a_+)^2}{|b|^2} - \frac{945\pi}{8192} C_{E,i} \frac{\hat{b} \cdot a_i v_i \cdot a_i}{|b|^2} \\ & \left. + \frac{6}{5} \sum_j C_{E,j} \frac{2a_j^2 + 5(\hat{b} \cdot a_j)^2}{|b|^2} \right] + \mathcal{O}(\gamma^{-1/2}, G^4). \quad (50) \end{aligned}$$

Here we use the notation $a_\pm^\mu = a_1^\mu \pm a_2^\mu$. Cancellations between the conservative and radiative pieces are essential to ensure the finiteness of this result. Note the dependence on the particle label in the second term of the second line, which disappears for Kerr black holes.

Finally, let us discuss parametrizations of the full radiative observables. One may introduce Lorentz transformations $\Lambda_i^\mu{}_\nu$ that transform the initial momenta and spin vectors to the final ones [51,53,149]:

$$\Delta p_i^\mu = (\Lambda_i^\mu{}_\nu - \delta_\nu^\mu) p_i^\nu, \quad (51a)$$

$$\Delta S_i^\mu = (\Lambda_i^\mu{}_\nu - \delta_\nu^\mu) S_i^\nu. \quad (51b)$$

The same transformation acts on both p_i^μ and S_i^μ : one sees this naturally given the requirement that p_i^2 , S_i^2 , and $p_i \cdot S_i$ must all be explicitly conserved. Conservation of p_i^2 (S_i^2) implies that the final momentum (spin vector) is given by a boost (rotation) of the initial one. Conservation of $p_i \cdot S_i$ implies that the boost and rotation may be combined into a single Lorentz transformation.

V. BEYOND LINEAR RESPONSE

The Bini-Damour linear response relation is used to infer linearly radiative contributions to the scattering angle θ from (angular) momentum loss at lower-PM orders. For aligned spins [81–83]

$$\theta_{\text{rad}} = -\frac{1}{2} \left(\frac{\partial \theta}{\partial J} J_{\text{rad}} + \frac{\partial \theta}{\partial E} E_{\text{rad}} \right), \quad (52)$$

where J_{rad} and E_{rad} are respectively the angular momentum and energy losses. In Ref. [61] this was used to deduce the radiative part of the quadratic-in-spin 3PM scattering angle

$\theta_{\text{rad}}^{(3)}$ for aligned spins, which we have now reconfirmed with our full calculation of $\Delta p_1^{(3)\mu}$. At 3PM order, given that $E_{\text{rad}} \sim \mathcal{O}(G^3)$ the second term plays no role: the linear response is entirely accounted for by the radiated angular momentum $J_{\text{rad}} \sim \mathcal{O}(G^2)$.

Using our newly derived observables we have checked and can confirm that at 3PM order the linear response relation (52) generalizes for misaligned spin vectors to

$$\Delta p_{1,\text{rad}}^{(+)\mu} = \frac{1}{2} \left(\frac{\partial \Delta p_1^\mu}{\partial J^\nu} J_{\text{rad}}^\nu + \frac{\partial \Delta p_1^\mu}{\partial P^\nu} P_{\text{rad}}^\nu \right). \quad (53)$$

Again, the part of this formula carrying P_{rad}^μ vanishes at 3PM order, but we include it to maintain the analogy with Eq. (52); in Eq. (38a) we defined $\Delta p_i^{(+)\mu}$ as the part of the impulse even under a time-reversal operation, where $v_i^\mu \rightarrow -v_i^\mu$. As a special case of Eq. (53) we derive a linear response relation for θ_i (48a) at 3PM order:

$$\theta_{i,\text{rad}}^{(+)} = -\frac{1}{2} \left(\frac{\partial \theta_i}{\partial J^\mu} J_{\text{rad}}^\mu + \frac{\partial \theta_i}{\partial P^\mu} P_{\text{rad}}^\mu \right), \quad (54)$$

where $\theta_{i,\text{rad}} = \theta_i - \theta_{\text{cons}}$ and the (+) superscript was defined in Eq. (39).

For the spin kick we learn about the odd part $\Delta S_i^{(-)\mu}$:

$$\Delta S_{1,\text{rad}}^{(-)\mu} = \frac{1}{2} \left(\frac{\partial \Delta S_1^\mu}{\partial J^\nu} J_{\text{rad}}^\nu + \frac{\partial \Delta S_1^\mu}{\partial P^\nu} P_{\text{rad}}^\nu \right). \quad (55)$$

The J^ν derivative is equivalent to an L^ν derivative (16): when taking these vectorial derivatives, we ignore all constraints (e.g. $L \cdot p_i = 0$) and treat the vectors involved (L^μ, p_i^μ, a_i^μ) as independent. All dependence on b^μ should be reexpressed in terms of L^μ by inverting $L^\mu = -E^{-1} \epsilon^\mu{}_{\nu\rho\sigma} b^\nu p_1^\rho p_2^\sigma$.

Both of these linear response relations involve the full vectorial radiated angular momentum J_{rad}^μ . We require it only up to 2PM order:

$$J_{\text{rad}}^{(2)\mu} = \Re \left[-\frac{4M^3 v^2 (2\gamma^2 - 1)}{|b|\Gamma} \mathcal{I}(v) \zeta^\mu \right. \\ \left. \times \left(1 + \frac{2va_3 \cdot \zeta}{|b|(1+v^2)} + \frac{(a_3 \cdot \zeta)^2}{|b|^2} - \sum_{i=1}^2 \frac{C_{E,i}}{|b|^2} (a_i \cdot \zeta)^2 \right) \right], \quad (56)$$

where $v = \sqrt{\gamma^2 - 1}/\gamma$ is the relative velocity, $a_3^\mu = a_1^\mu + a_2^\mu$, and the complex vector is $\zeta^\mu = \hat{L}^\mu + i\hat{b}^\mu$; the universal prefactor $\mathcal{I}(v)$ was given in Eq. (41).

For zero or aligned spins, one may straightforwardly show that the new linear response relation (53) reduces to the Bini-Damour formula (52) by inserting [36]

$$\Delta p_1^\mu = p_\infty \hat{b}^\mu \sin \theta - v_2 \cdot P_{\text{rad}} \frac{\gamma v_1^\mu - v_2^\mu}{\gamma^2 - 1} \\ + (\cos \theta - 1) p^\mu, \quad (57)$$

which holds up to the desired 3PM order. This schematic form of the impulse shows that Δp_1^μ is fully characterized by θ and P_{rad}^μ : as $\Delta p^{(+)\mu} = p_\infty \sin \theta \hat{b}^\mu$ knows only about the scattering angle, the linear response relationship yields no information concerning P_{rad}^μ . We use the fact that, for aligned spins, $J_{\text{rad}}^\mu = -J_{\text{rad}} \hat{L}^\mu$.

However, in this section we generalize beyond simple linear response. The linear response relation (53) forms part of a more general pair of relationships that allow us to reconstruct conservative and radiative parts of the scattering observables at higher-PM orders:

$$\Delta p_{i,\text{cons}}^\mu = \frac{1}{2} (\Delta p_i^\mu(L^\mu, p_i^\mu, S_i^\mu) \\ + \Delta p_i^\mu(L^\mu + \Delta L^\mu, -p_i^\mu - \Delta p_i^\mu, S_i^\mu + \Delta S_i^\mu)), \quad (58a)$$

$$\Delta p_{i,\text{rad}}^\mu = \frac{1}{2} (\Delta p_i^\mu(L^\mu, p_i^\mu, S_i^\mu) \\ - \Delta p_i^\mu(L^\mu + \Delta L^\mu, -p_i^\mu - \Delta p_i^\mu, S_i^\mu + \Delta S_i^\mu)) \quad (58b)$$

for the impulse, and

$$\Delta S_{i,\text{cons}}^\mu = \frac{1}{2} (\Delta S_i^\mu(L^\mu, p_i^\mu, S_i^\mu) \\ - \Delta S_i^\mu(L^\mu + \Delta L^\mu, -p_i^\mu - \Delta p_i^\mu, S_i^\mu + \Delta S_i^\mu)), \quad (59a)$$

$$\Delta S_{i,\text{rad}}^\mu = \frac{1}{2} (\Delta S_i^\mu(L^\mu, p_i^\mu, S_i^\mu) \\ + \Delta S_i^\mu(L^\mu + \Delta L^\mu, -p_i^\mu - \Delta p_i^\mu, S_i^\mu + \Delta S_i^\mu)) \quad (59b)$$

for the spin kick; we will define the split $\Delta X = \Delta X_{\text{cons}} + \Delta X_{\text{rad}}$ of the full observables into conservative and radiative parts below. We interpret all observables as real functions of the initial kinematic vectors: the orbital angular momentum vector L^μ (14a), the spin vectors $S_i^\mu = m_i a_i^\mu$, and the momenta $p_i^\mu = m_i v_i^\mu$.

A. Derivation

We define the conservative part of a single-operator expectation value as the average of its value evaluated in the in-in and out-out prescriptions:

$$\langle \mathcal{O} \rangle_{\text{cons}} := \frac{1}{2} (\langle \mathcal{O} \rangle_{\text{in-in}} + \langle \mathcal{O} \rangle_{\text{out-out}}). \quad (60)$$

The expectation $\langle \mathcal{O} \rangle_{\text{out-out}}$ is computed using precisely the same Feynman rules as $\langle \mathcal{O} \rangle_{\text{in-in}}$, but with advanced propagators pointing towards the outgoing line instead of retarded, both on the worldlines and in the bulk. At 3PM

order, one may verify by explicit calculation that this definition of the conservative dynamics coincides precisely with evaluating integrals only in the potential region—the approach previously taken for the conservative 3PM spinning dynamics in Ref. [61].

Specializing to the impulse Δp_i^μ , we therefore have

$$\begin{aligned} \Delta p_{i,\text{cons}}^\mu &= \frac{1}{2}(\Delta p_{i,\text{in-in}}^\mu(L_-^\mu, p_{i-}^\mu, S_{i-}^\mu) + \Delta p_{i,\text{out-out}}^\mu(L_+^\mu, p_{i+}^\mu, S_{i+}^\mu)). \end{aligned} \quad (61)$$

While $\Delta p_{i,\text{in-in}}^\mu$ is given in terms of background parameters defined at past infinity (− subscript), $\Delta p_{i,\text{out-out}}^\mu$ is evaluated in terms of parameters defined at future infinity (+ subscript). As we prefer to express $\Delta p_{i,\text{cons}}^\mu$ in terms of initial variables we insert

$$L_-^\mu = L^\mu, \quad L_+^\mu = L^\mu + \Delta L^\mu, \quad (62a)$$

$$p_{i-}^\mu = p_i^\mu, \quad p_{i+}^\mu = p_i^\mu + \Delta p_i^\mu, \quad (62b)$$

$$S_{i-}^\mu = S_i^\mu, \quad S_{i+}^\mu = S_i^\mu + \Delta S_i^\mu. \quad (62c)$$

Thus p_{i+}^μ and S_{i+}^μ are given using our preexisting knowledge of the impulse Δp_i^μ and spin kick ΔS_i^μ . We can infer ΔL^μ —and therefore L_+^μ —up to 2PM order from the known 2PM angular momentum loss J_{rad}^μ (56). Using Eq. (16)

$$\begin{aligned} \Delta J^\mu &= -J_{\text{rad}}^\mu \\ &= \Delta L^\mu + \sum_i \frac{1}{m_i} (\hat{P} \cdot \Delta p_i S_i^\mu + \hat{P} \cdot (p_i + \Delta p_i) \Delta S_i^\mu \\ &\quad - \hat{P} \cdot \Delta S_i p_i^\mu - \hat{P} \cdot (S_i + \Delta S_i) \Delta p_i^\mu), \end{aligned} \quad (63)$$

which we can rearrange to find ΔL^μ —ignoring the (linear) momentum loss $P_{\text{rad}}^\mu \sim \mathcal{O}(G^3)$. In the nonspinning case $\Delta L^\mu = -J_{\text{rad}}^\mu$, i.e. the change in the orbital angular momentum vector is given precisely by the total loss of angular momentum.

Finally, to obtain Eq. (58) we use the fact that

$$\Delta p_{i,\text{out-out}}^\mu(L^\mu, p_i^\mu, S_i^\mu) = \Delta p_{i,\text{in-in}}^\mu(L^\mu, -p_i^\mu, S_i^\mu), \quad (64)$$

which simply tells us that, having computed $\Delta p_{i,\text{in-in}}^\mu$, we may easily derive $\Delta p_{i,\text{out-out}}^\mu$ by continuing $p_i^\mu \rightarrow -p_i^\mu$. This works because the time-reversal operation induces a change of sign on the $i0$ prescription of the propagators (27) and (28). For the graviton propagator (27) $\text{sgn}(k^0)i0 = (k \cdot v_i)i0$: the sign on the energy component of k^μ is defined by the direction of either velocity vector v_i^μ . For the worldline propagators (28), $\omega \rightarrow -\omega$: the z_i propagator (28a) remains the same, but with $i0 \rightarrow -i0$,

while the ψ_i^μ propagator (28b) also picks up an overall sign. This overall sign is compensated for in the WQFT Feynman rules for by the vertices, which are themselves invariant under time reversal except for $S_i^{\mu\nu} = \epsilon^{\mu\nu\rho\sigma} v_i^\rho S_i^\sigma$, which flips as $S_i^{\mu\nu} \rightarrow -S_i^{\mu\nu}$. Each time we propagate an internal spin mode we pick up a factor of $S_i^{\mu\nu}$, which compensates for the additional sign.

The derivation of Eq. (59) for the spin kick proceeds similarly, although in this case as ΔS_i^μ is defined indirectly via Δp_i^μ and $\Delta S_i^{\mu\nu}$ (31) we obtain the different relative signs. One can see why this is necessary by examining the spin kick at 1PM order:

$$\Delta S_1^{(1)\mu} = \frac{4m_1 m_2 a_{1\nu} b^\nu (v_1 - 2\gamma v_2)^\mu}{|b|^2 \sqrt{\gamma^2 - 1}} + \mathcal{O}(S^2), \quad (65)$$

which is of course purely conservative. It is odd under the time-reversal $v_i^\mu \rightarrow -v_i^\mu$, which agrees with Eq. (59b) we have $\Delta S_i^{(1)\mu}(L^\mu, p_i^\mu, S_i^\mu) = -\Delta S_i^{(1)\mu}(L^\mu, -p_i^\mu, S_i^\mu)$.

B. Interpretation

We can now derive $\Delta p_{i,\text{cons}}^{(m;-)\mu}$ at any PM order m . To do so, we insert the PM decomposition (32) into Eq. (58a) and perform a Taylor-series expansion of the right-hand side, picking out the desired PM order m :

$$\begin{aligned} \Delta p_{i,\text{cons}}^{(m)\mu} &= \frac{1}{2} \left(\Delta p_i^{(m)\mu}(L^\mu, p_i^\mu, S_i^\mu) + \Delta p_i^{(m)\mu}(L^\mu, -p_i^\mu, S_i^\mu) \right. \\ &\quad + \frac{\partial \Delta p_i^{(m-1)\mu}(L^\mu, -p_i^\mu, S_i^\mu)}{\partial L^\nu} \Delta L^{(1)\nu} \\ &\quad + \sum_{j=1}^2 \left(-\frac{\partial \Delta p_i^{(m-1)\mu}(L^\mu, -p_i^\mu, S_i^\mu)}{\partial p_j^\nu} \Delta p_j^{(1)\nu} \right. \\ &\quad + \left. \frac{\partial \Delta p_i^{(m-1)\mu}(L^\mu, -p_i^\mu, S_i^\mu)}{\partial S_j^\nu} \Delta S_j^{(1)\nu} \right) \\ &\quad \left. + \dots \right). \end{aligned} \quad (66)$$

Taking the difference between this formula and its counterpart with $p_i^\mu \rightarrow -p_i^\mu$ gives $\Delta p_{i,\text{cons}}^{(m;-)\mu}$ on the left-hand side, and the first two terms on the right-hand side cancel out. Thus, $\Delta p_{i,\text{cons}}^{(m;-)\mu}$ is given entirely by lower-PM observables.

Similarly, using Eq. (59a) we may predict $\Delta S_{i,\text{cons}}^{(m;+)\mu}$.

Using Eqs. (58b) and (59b) by the same procedure we may determine $\Delta p_{i,\text{rad}}^{(+)\mu}$ and $\Delta S_{i,\text{rad}}^{(-)\mu}$. However, at 3PM order an additional simplification is possible: using the fact that the conservative and radiative observables have opposite behaviors under $v \rightarrow -v$. As J_{rad}^μ is the only nonzero radiative observable at 2PM order, it follows that all other contributions to the linear response relation cancel out at 3PM order, leaving us with Eqs. (53) and (55) as proposed

earlier. This we have checked carefully by direct calculation.

We anticipate that Eq. (59) will be useful for future 4PM computations, as we will not need to calculate the complete radiative observables: for the impulse we need only calculate $\Delta p_{1,\text{cons}}^{(4;+)\mu}$ and $\Delta p_{1,\text{rad}}^{(4;-)\mu}$ directly. This cuts down on the regions within which we need to evaluate the master integrals: only the conservative sector for the real integrals, and the radiative sector for the pseudoreal integrals. However, in the radiative sector this is predicated on our knowing the 3PM angular momentum loss $J_{\text{rad}}^{(3)\mu}$, which currently we have only up to the leading 2PM order (56) in the spinning case. For nonspinning bodies, the 3PM angular momentum loss has been determined and used to infer contributions to 4PM scattering observables [140]; a similar concept will certainly apply in the presence of spin.

VI. HAMILTONIAN

Let us now focus on the strictly conservative part of the dynamics, encoded by $\Delta p_{i,\text{cons}}^\mu$ and $\Delta S_{i,\text{cons}}^\mu$. Computing a 3PM quadratic-in-spin Hamiltonian maps these unbound observables into bound dynamics, which in the spinning context is especially useful given the current lack of direct analytic continuations between bound and unbound observables with generic misaligned spin directions. In line with recent literature on post-Minkowskian dynamics [127,130,139] we work in the CoM frame with canonical variables $\mathbf{p}(t)$ and $\mathbf{x}(t)$ describing the relative momentum and position of the two bodies respectively. These dynamical variables satisfy canonical Poisson brackets:

$$\{\mathbf{x}^m(t), \mathbf{p}^n(t)\}_{\text{P.B.}} = \delta^{mn}, \quad (67a)$$

$$\{\mathbf{S}_i^m(t), \mathbf{S}_i^n(t)\}_{\text{P.B.}} = \epsilon^{mnk} \mathbf{S}_i^k(t). \quad (67b)$$

The spin of each body is described by the spin vectors $\mathbf{S}_i(t)$, with the canonical SSC described in Sec. II B.

The Hamiltonian H is fixed in isotropic gauge, meaning that it does not depend on $\mathbf{x}(t) \cdot \mathbf{p}(t)$. It takes the general form

$$H(\mathbf{x}, \mathbf{p}, \mathbf{S}_i) = \sqrt{\mathbf{p}^2 + m_1^2} + \sqrt{\mathbf{p}^2 + m_2^2} + V(\mathbf{x}, \mathbf{p}, \mathbf{S}_i) \quad (68)$$

with gravitational potential

$$\begin{aligned} V(\mathbf{x}, \mathbf{p}, \mathbf{S}_i) &= \sum_A \mathcal{O}^A V^A(\mathbf{x}, \mathbf{p}) + \mathcal{O}(S^3) \\ &= V^{(0)} + \sum_i V^{(1,i)} \mathcal{O}^{(1,i)} + \sum_{i,j,a} V^{(2,a,i,j)} \mathcal{O}^{(2,a,i,j)} \\ &\quad + \mathcal{O}(S^3). \end{aligned} \quad (69)$$

The potential is expanded in spin structures:

$$\mathcal{O}^{(0)} = 1, \quad (70a)$$

$$\mathcal{O}^{(1,i)} = \frac{(\mathbf{x} \times \mathbf{p}) \cdot \mathbf{a}_i}{|\mathbf{x}|^2}, \quad (70b)$$

$$\mathcal{O}^{(2,1,i,j)} = \frac{\mathbf{a}_i \cdot \mathbf{a}_j}{|\mathbf{x}|^2}, \quad (70c)$$

$$\mathcal{O}^{(2,2,i,j)} = \frac{\mathbf{x} \cdot \mathbf{a}_i \mathbf{x} \cdot \mathbf{a}_j}{|\mathbf{x}|^4}, \quad (70d)$$

$$\mathcal{O}^{(2,3,i,j)} = \frac{\mathbf{p} \cdot \mathbf{a}_i \mathbf{p} \cdot \mathbf{a}_j}{|\mathbf{x}|^2}, \quad (70e)$$

where $\mathbf{a}_i = \mathbf{S}_i/m_i$. In each case the first index counts the spin order; subsequent indices count the specific structures involved. Note that the symmetric spin structures $\mathcal{O}^{(2,a,1,2)} = \mathcal{O}^{(2,a,2,1)}$ are counted twice and their coefficients are equal. Finally, we PM-expand each component:

$$V^A(\mathbf{x}, \mathbf{p}) = \sum_n \left(\frac{GM}{|\mathbf{x}|} \right)^n c^{(n;A)}(\mathbf{p}^2). \quad (71)$$

These coefficients $c^{(n;A)}$ fully encode the Hamiltonian.

We fix the coefficients $c^{(n;A)}(\mathbf{p}^2)$ by matching observables computed from the Hamiltonian H with scattering observables from the WQFT. Hamilton's equations for the dynamical variables are

$$\dot{\mathbf{x}} = \frac{\partial H}{\partial \mathbf{p}}, \quad \dot{\mathbf{p}} = -\frac{\partial H}{\partial \mathbf{x}}, \quad \dot{\mathbf{S}}_i = -\mathbf{S}_i \times \frac{\partial H}{\partial \mathbf{S}_i}, \quad (72)$$

and we solve them perturbatively up to third order in G :

$$\mathbf{x}(t) = \mathbf{x}^{(0)} + \sum_{n=1}^3 G^n \mathbf{x}^{(n)}(t) + \mathcal{O}(G^4), \quad (73a)$$

$$\mathbf{p}(t) = \mathbf{p}^{(0)} + \sum_{n=1}^3 G^n \mathbf{p}^{(n)}(t) + \mathcal{O}(G^4), \quad (73b)$$

$$\mathbf{S}_i(t) = \mathbf{S}_i^{(0)} + \sum_{n=1}^3 G^n \mathbf{S}_i^{(n)}(t) + \mathcal{O}(G^4). \quad (73c)$$

The zeroth-order scattering trajectories are

$$\mathbf{x}^{(0)} = t \frac{\mathbf{p}_\infty}{\xi E} - \mathbf{b}_{\text{can}}, \quad \mathbf{p}^{(0)} = \mathbf{p}_\infty, \quad \mathbf{S}_i^{(0)} = \mathbf{S}_{i,\infty}, \quad (74)$$

where $p^\mu = (0, \mathbf{p}_\infty)$, $S_{i,\text{can}}^\mu = (0, \mathbf{S}_{i,\infty})$ and $b_{\text{can}}^\mu = (0, \mathbf{b}_{\text{can}})$. The dimensionless parameter ξ is defined as $\xi = E_1 E_2 / E^2$, where $E = E_1 + E_2$ and $E_i = \sqrt{\mathbf{p}_\infty^2 + m_i^2}$. Inserting the expansions of the dynamical variables (73) into Hamilton's equations (72) we get perturbative equations of motion at

each PM order. The spatial components of the impulse and spin kick in the CoM frame are then given by

$$\Delta \mathbf{p}^{(n)} = \int_{-\infty}^{\infty} dt \dot{\mathbf{p}}^{(n)}(t), \quad (75a)$$

$$\Delta \mathbf{S}_i^{(n)} = \int_{-\infty}^{\infty} dt \dot{\mathbf{S}}_i^{(n)}(t), \quad (75b)$$

where these are the conservative 3-vector components of Δp_1^μ and $\Delta S_{i,\text{can}}^\mu$. The change in the canonical spin vector is given in terms of the covariant spin kick by

$$\begin{aligned} \Delta S_{i,\text{can}}^\mu - \Delta S_i^\mu \\ = - \frac{m_i \hat{\mathbf{P}} \cdot \Delta S_i (\hat{\mathbf{P}}^\mu + v_i^\mu) + \hat{\mathbf{P}} \cdot (S_i + \Delta S_i) \Delta p_i^\mu}{E_i + m_i}, \end{aligned} \quad (76)$$

having used the definition of $S_{i,\text{can}}^\mu$ (18), and assuming conservative scattering.

Quite remarkably though, we find that knowledge of θ_{cons} (48) suffices in order to fully fix all coefficients in the Hamiltonian. The impulse and spin kick may then be expressed in terms of the coefficients $c^{(n;A)}(\mathbf{p}^2)$ and derivatives thereof. The derivatives come about when we evaluate $\dot{\mathbf{x}}$ by differentiating H with respect to \mathbf{p} :

$$\frac{\partial c^{(n;A)}(\mathbf{p}^2)}{\partial \mathbf{p}} = 2\mathbf{p} \frac{\partial c^{(n;A)}(\mathbf{p}^2)}{\partial p^2} = 2\mathbf{p} c^{(n;A;1)}(\mathbf{p}^2), \quad (77)$$

where $c^{(n;A;m)} := \partial^m c^{(n;A)} / (\partial \mathbf{p}^2)^m$ —the observables are written as functions of $c^{(n;A;m)}$. We then match those expressions to the explicit WQFT-derived results and solve for the coefficients. We print the results until linear in spins here, and the results quadratic in spins in Appendix C:

$$\begin{aligned} c^{(3;0)}(\mathbf{p}^2) &= -\frac{p_\infty^2}{4E\xi} \theta_{\text{can}}^{(3;0)} + \frac{1}{2\pi p_\infty^2} \mathcal{D} \left[\frac{p_\infty^4}{E\xi} \theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(2;0)} \right] \\ &\quad - \frac{1}{48 p_\infty^2} \mathcal{D}^2 \left[\frac{p_\infty^4}{E\xi} (\theta_{\text{can}}^{(1;0)})^3 \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \\ c^{(3;1,i)}(\mathbf{p}^2) &= -\frac{p_\infty}{4E\xi} \theta_{\text{can}}^{(3;1,i)} + \frac{1}{2\pi p_\infty^4} \mathcal{D} \left[\frac{p_\infty^5}{E\xi} (\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(2;1,i)} \right. \\ &\quad \left. + \theta_{\text{can}}^{(2;0)} \theta_{\text{can}}^{(1;1,i)}) \right] \\ &\quad - \frac{1}{16 p_\infty^4} \mathcal{D}^2 \left[\frac{p_\infty^5}{E\xi} (\theta_{\text{can}}^{(1;0)})^2 \theta_{\text{can}}^{(1;1,i)} \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}. \end{aligned} \quad (78)$$

Here $\phi_{\text{can}}^{(n;A)}$ are canonical expansion coefficients defined in Appendix A and related to the covariant expansion coefficients of Eq. (49) by

$$\theta_{\text{can}}^{(n;0)} = \Gamma \theta_{\text{cons}}^{(n;0)}, \quad (79a)$$

$$\theta_{\text{can}}^{(n;1,i)} = \Gamma \left(\theta_{\text{cons}}^{(n;1,i)} + n \frac{p_\infty}{E_i + m_i} \theta_{\text{cons}}^{(n;0)} \right). \quad (79b)$$

A particular subtlety here is that $\theta_{\text{can}}^{(n;A)}$ is given in terms of the previously defined background variables p_∞ , γ , $E_i = \sqrt{\mathbf{p}_\infty^2 + m_i^2}$ evaluated at past infinity, e.g.

$$\gamma = \frac{p_1 \cdot p_2}{m_1 m_2} = \frac{E_1 E_2 + \mathbf{p}_\infty^2}{m_1 m_2}, \quad (80)$$

instead of the dynamical momentum $\mathbf{p}(t)$: we interpolate to the full dynamical coefficients $c^{(n;A;m)}$ simply by replacing one with the other. We have also introduced the differential operator

$$\mathcal{D}[X] := \frac{\partial(p_\infty X)}{\partial p_\infty}, \quad (81)$$

and $\mathcal{D}^2[X] = \mathcal{D}[\mathcal{D}[X]]$. The angle and its coefficients are given in Appendix B, together with full expressions for the Hamiltonian coefficients given in the Supplemental Material [145].

We have checked this Hamiltonian numerically against the recent results obtained in Ref. [139], which also included 3PM quadratic-in-spin terms. Our results complement those by adding $S_1 S_2$ contributions to the Hamiltonian together with finite-size effects ($C_{E,i}$ terms) in the S_i^2 sector. We have also verified that the PN expansion of this Hamiltonian correctly reproduces 4PN results in the isotropic gauge [29]. We did so by PN-expanding the $c^{(n;A)}(\mathbf{p}^2)$ coefficients in powers of \mathbf{p}^2 .

Finally, let us observe that having expressed the coefficients of the Hamiltonian in terms only of the scattering angle ϕ_{can} , this implies that the full conservative scattering observables $\Delta p_{i,\text{cons}}^{(3)\mu}$ and $\Delta S_{i,\text{cons}}^{(3)\mu}$ may themselves be expressed in terms of this angle. We would obtain the precise relationship by solving Hamilton's equations again for the impulse and spin kick, but this time plugging in expressions in terms of ϕ_{cons} . In contrast to the Hamiltonian, the relations thus obtained are gauge invariant and will be an intriguing topic of future studies.

VII. UNBOUND-TO-BOUND MAPPINGS

Let us now discuss how our results may be applied to describe bound orbits, which the now-complete 3PM quadratic-in-spin Hamiltonian (68) gives us partial access to. This will allow us to determine the binding energy, which together with the radiative fluxes may be used to inform complete gravitational waveform models. In this section we specialize to spin vectors aligned with the orbital angular momentum vector:

$$S_i^\mu = m_i a_i^\mu = G m_i^2 \chi_i \hat{L}^\mu, \quad (82)$$

where χ_i are the directed spin lengths and $S_{i,\text{can}}^\mu = S_i^\mu$. For Kerr black holes $m_i|\chi_i|$ are the radii of the ring singularities, and $-1 < \chi_i < 1$. Using Eq. (19) we see that for aligned spins $\hat{L}^\mu = \hat{L}_{\text{can}}^\mu$; however, their magnitudes differ and so using Eq. (19) we introduce

$$\lambda = \frac{|L_{\text{can}}|}{GM\mu} = \frac{|b|p_\infty}{GM\mu} + \frac{\mathcal{E}}{2} \left(\chi_+ + \frac{\delta}{\Gamma} \chi_- \right), \quad (83)$$

where $\mathcal{E} = (E - M)/\mu$ is the reduced binding energy, $\delta = (m_2 - m_1)/M$ and

$$\chi_\pm = \frac{m_1\chi_1 \pm m_2\chi_2}{M}, \quad (84)$$

$$\chi_{E,\pm}^2 = \frac{C_{E,1}m_1^2\chi_1^2 \pm C_{E,2}m_2^2\chi_2^2}{M^2}. \quad (85)$$

Using axial symmetry the Hamiltonian $E = H(r, p_r, \lambda, \chi_i)$ depends—besides the masses m_i and finite-size coefficients $C_{E,i}$ —on the radial coordinate r and momentum p_r , where in axial coordinates

$$\mathbf{p}^2 = p_r^2 + \frac{p_\phi^2}{r^2}. \quad (86)$$

The axial momentum $p_\phi = |L_{\text{can}}|$ is a constant of motion. In all of these expressions we leave the dependence on masses m_i and finite-size coefficients $C_{E,i}$ implicit.

A. Numerical PM binding energy

In Fig. 2 we plot the reduced binding energy $\mathcal{E} = (H - M)/\mu$ for circular orbits as a function of the orbital frequency $GM\Omega$ leading up to merger. It is compared with a numerical relativity (NR) simulation provided by the Simulating eXtreme Spacetimes (SXS) Collaboration [150], extracted in Ref. [151]. Our plots are also determined numerically: within the Hamiltonian we set $p_r = 0$ and, using $\dot{p}_r = 0 = -\partial H/\partial r$, we solve for $\lambda(r)$ for different orbital separations r and with specific values of χ_1, χ_2 . The reduced binding energy \mathcal{E} is plotted against the orbital frequency:

$$x^{3/2} = GM\Omega = \frac{d\mathcal{E}}{d\lambda}, \quad (87)$$

with the number of orbits leading up to merger provided by NR—see Refs. [48,71] for more details.

The conclusion of these plots is somewhat disappointing: the quadratic-in-spin part of the Hamiltonian yields little improvement over the spin-orbit contribution. However, there is a far more noticeable improvement when going from 2PM to 3PM order, which suggests that producing a 4PM spinning Hamiltonian will be a worthwhile endeavor. A similar improvement of the 4PM (hyperbolic)




FIG. 2. The reduced binding energy $\mathcal{E} = (H - M)/\mu$ for circular orbits determined numerically and plotted as a function of orbital frequency $GM\Omega$ up to the innermost stable circular orbit. It is compared for different PM and spin orders with a NR simulation provided by the SXS Collaboration [150].

Hamiltonian over the 3PM seen in the nonspinning case [71] also encourages us in this direction; however, it will also be important to resum in the test-body limit by feeding these results into a suitable EOB model [45]. It is also worth noting that, as these plots are generated for circular orbits, they do not showcase PM results in the best possible light: it is anticipated that PM-based results will perform better for highly elliptical orbits, where the velocity at closest approach between the massive bodies is large [71].

B. Analytic PN binding energy

Working in the PN expansion we may derive precise analytic formulas for the binding energy \mathcal{E} and periastron advance $\Delta\phi$. Following closely the discussion in Ref. [148] (see also Refs. [35,149]) our starting point is the radial action for unbound orbits:

$$w_r(\mathcal{E}, \lambda, \chi_i) = \frac{1}{GM\mu\pi} \text{Pf} \int_{r_{\min}}^{\infty} dr p_r(r, \mathcal{E}, \lambda, \chi_i), \quad (88)$$

where Pf denotes the *partie finie* of the radial action; the energy constraint $E = H(r, p_r, \lambda, \chi_i)$ can be solved for the radial momentum p_r —but we refrain from doing so explicitly. The innermost point r_{\min} is given by the root of $p_r(r_{\min}, \mathcal{E}, \lambda, \chi_i) = 0$. This unbound radial action is related to the bound radial action i_r by analytic continuation:

$$i_r(\mathcal{E}, \lambda, \chi_i) = w_r(\mathcal{E}, \lambda, \chi_i) - w_r(\mathcal{E}, -\lambda, -\chi_i), \quad (89)$$

which sends $\lambda \rightarrow -\lambda$ and $\chi_i \rightarrow -\chi_i$. For unbound dynamics the reduced binding energy $\mathcal{E} > 0$ ($\gamma > 1$), but now we

consider values $\mathcal{E} < 0$ ($0 < \gamma < 1$) as we see in Fig. 2. The reduced binding energy \mathcal{E} for circular orbits is derived by setting the bound radial action to zero:

$$i_r(\mathcal{E}, \lambda, \chi_i) = 0. \quad (90)$$

Using Eqs. (90) and (87) we may write \mathcal{E} as a function of x , the spins χ_{\pm} , $\chi_{E,\pm}$, and ν .

Rather than the Hamiltonian, we prefer to derive the radial action—and thus the binding energy \mathcal{E} —from the gauge-invariant conservative scattering angle θ_{cons} . It is given by a λ derivative of the unbound radial action w_r :

$$2\pi \frac{\partial}{\partial \lambda} w_r(\mathcal{E}, \lambda, \chi_i) = -(\theta_{\text{cons}}(\mathcal{E}, \lambda, \chi_i) + \pi), \quad (91)$$

and by analytic continuation it is related to the periastron advance for bound orbits [34–37]:

$$\Delta\phi(\mathcal{E}, \lambda, \chi_i) = \theta_{\text{cons}}(\mathcal{E}, \lambda, \chi_i) + \theta_{\text{cons}}(\mathcal{E}, -\lambda, -\chi_i). \quad (92)$$

We PM-expand the angle in λ as

$$\theta_{\text{cons}} = \sum_n \frac{\tilde{\theta}^{(n)}}{\lambda^n}, \quad (93)$$

where $\tilde{\theta}^{(n)}$ depends on χ_i through the ratios χ_i/λ . The analytic continuation in λ (89) is trivial for all terms in w_r except the one coming from the nonspinning part of $\tilde{\theta}^{(1)}$, wherein the dependence on λ is $\log(\lambda)$. With this exception the odd-in- G terms in the PM expansion disappear in the difference (89), and the analytic continuation of $\log(-\lambda) - \log(\lambda)$ leaves behind a finite piece. We therefore have

$$i_r(\mathcal{E}, \lambda, \chi_i) = -\lambda + \frac{2\gamma^2 - 1}{\sqrt{1 - \gamma^2}} - \frac{1}{\pi} \sum_n \int d\lambda \frac{\tilde{\theta}^{(2n)}}{\lambda^{2n}}. \quad (94)$$

Here γ should be reexpressed in terms of \mathcal{E} . The integral on λ is elementary but left unresolved because of the remaining λ dependence inside $\tilde{\theta}^{(2n)}$.

Naively this result indicates that our 3PM results have no relevance for the mapping to bound results. However, this obstacle is conveniently circumvented in the PN expansion by use of the so-called impetus formula with PM-coefficients f_k :

$$\mathbf{p}^2 = p_{\infty}^2 + \sum_{k=1}^{\infty} \frac{G^k}{r^k} f_k. \quad (95)$$

This is fed into the definition of the scattering angle by way of the radial action (88):

$$\begin{aligned} \pi + \theta_{\text{cons}} &= -\frac{2}{GM\mu} \int_{r_{\text{min}}}^{\infty} dr \frac{\partial}{\partial \lambda} p_r(r, \mathcal{E}, \lambda) \\ &= -2 \int_{r_{\text{min}}}^{\infty} dr \frac{\partial}{\partial \lambda} \sqrt{\frac{1}{G^2 M^2 \mu^2} \left(p_{\infty}^2 + \sum_{k=1}^{\infty} \frac{G^k}{r^k} f_k \right) - \frac{\lambda^2}{r^2}}. \end{aligned} \quad (96)$$

This integral has been performed up to high orders in G in Ref. [43] and the scattering angle is then expressed in terms of f_k . Knowledge of the 1PM, 2PM, and 3PM scattering angles suffices in order to fully determine $f_{k \leq 3}$, which may in turn be used to reconstruct the leading-PN terms of the angle at higher PM orders.

In order to reconstruct the quadratic-in-spin binding energy up to 4PN order, we find it necessary to reconstruct the leading-PN and subleading-PN parts of $\tilde{\theta}^{(6)}$ and $\tilde{\theta}^{(4)}$ respectively, at both linear and quadratic order in spin. Following this procedure, we discover that the binding energy is

$$\begin{aligned} -2\mathcal{E} &= x \left[1 - x \frac{9 + \nu}{12} - x^2 \frac{81 - 57\nu + \nu^2}{24} + \dots \right] \\ &+ x^{5/2} \left[\frac{7\chi_+ - \delta\chi_-}{3} + x \frac{(99 - 61\nu)\chi_+ - (45 - \nu)\delta\chi_-}{18} + x^2 \frac{(405 - 1101\nu + 29\nu^2)\chi_+ - (243 - 165\nu - \nu^2)\delta\chi_-}{24} + \dots \right] \\ &- x^3 \left[\chi_+^2 + \frac{5x}{36} ((5 - 6\nu)\chi_+^2 - 44\chi_+\chi_- - (1 + 8\nu)\chi_-^2) \right. \\ &+ \frac{7x^2}{216} ((198 - 680\nu + 3\nu^2)\chi_+^2 - 2(171 - 137\nu)\delta\chi_-\chi_+ + (63 - 251\nu + 56\nu^2)\chi_-^2) \\ &\left. + \chi_{E,+}^2 + \frac{5x}{6} ((5 - \nu)\chi_{E,+}^2 - 2\delta\chi_{E,-}^2) + \frac{x^2}{72} ((1125 - 1025\nu + 7\nu^2)\chi_{E,+}^2 - 2\delta(279 - 70\nu)\chi_{E,-}^2) + \dots \right]. \end{aligned} \quad (97)$$

At each order in spin, we give the terms up to and including next-to-next-to-leading order in the PN expansion. With the nonspinning terms appearing up to 2PN order, the spin-orbit and spin-spin terms appear up to 3.5PN and 4PN order respectively.

C. Radiated energy

Finally, we may also determine the energy radiated per orbit in the CoM frame [37]:

$$E_{\text{rad}}^{\text{bound}}(\mathcal{E}, \lambda, \chi_i) = E_{\text{rad}}(\mathcal{E}, \lambda, \chi_i) - E_{\text{rad}}(\mathcal{E}, -\lambda, -\chi_i), \quad (98)$$

where $E_{\text{rad}} = \hat{P} \cdot P_{\text{rad}}$ derives from the full radiative momentum impulse Δp_1^μ (44). The same analysis was also done successfully by Riva, Vernizzi, and Wong [75], who confirmed that this result for the radiated energy for bound orbits agrees with the corresponding 4PN terms from Ref. [32]. To leading-PN at each spin order, we find that

$$E_{\text{rad}}^{\text{bound}} = \frac{2\pi\mu^2}{M} \left(\frac{148\mathcal{E}^2}{15\lambda^3} - \frac{4(67\chi_+ - 2\delta\chi_-)\mathcal{E}^3}{5\lambda^4} + \frac{(236(\chi_+^2 - \chi_{E,+}^2) + 9\chi_-^2)\mathcal{E}^3}{5\lambda^5} \right) + \dots \quad (99)$$

Like in the binding energy (97), we see that the linear-in-spin terms appear at 1.5PN above the nonspinning; the quadratic-in-spin terms at 2PN above.

However, we emphasize that, unlike in the case of the binding energy, this result for the radiated energy does not encode the full leading-PN result. To see why, we recall that in the nonspinning case the leading-PN radiated energy may be derived from Einstein's quadrupole formula:

$$E_{\text{rad}}^{\text{bound}} = \frac{2\pi\mu^2}{M} \left(\frac{148\mathcal{E}^2}{15\lambda^3} + \frac{244\mathcal{E}}{5\lambda^5} + \frac{85}{3\lambda^7} \right) + \dots \quad (100)$$

While we do reproduce the λ^{-3} contribution, gaining access to the λ^{-5} and λ^{-7} contributions via PM-based scattering calculations would necessitate 5PM and 7PM calculations respectively: a seemingly impossible task. However, a more encouraging conclusion was reached in Ref. [36]: that by instead using the PM-scattering data to fix the form of the (gauge-dependent) instantaneous fluxes of energy and angular momentum, the leading-PN information could instead be extracted from a future 4PM derivation. Such an approach might be especially beneficial in the spinning case, given the unbound-to-bound mapping's current limitation to aligned spin vectors. This would be similar to our present use of the conservative scattering observables to reconstruct a Hamiltonian, a prospect that we leave for future work.

VIII. CONCLUSIONS

In this paper we have for the first time provided complete expressions for the impulse Δp_1^μ and spin kick ΔS_1^μ at third post-Minkowskian (3PM) order to quadratic order in the spins of two scattering bodies, including finite-size corrections. The computation relied on our use of the spinning, supersymmetric WQFT formalism [57,58] and its extension to utilize the Schwinger-Keldysh in-in formalism

[63,76–80]. These results upgrade the previously obtained conservative observables provided by the present authors [61], and include knowledge of the total radiated four-momentum P_{rad}^μ which has also recently been computed using worldline EFT methods [75]. We also wrote down a scattering angle that encapsulates the motion for arbitrarily misaligned spin directions.

Next, we demonstrated how both conservative and radiative parts of these observables may be reconstructed using an extension of Bini and Damour's linear response relation [81–83], incorporating the full 2PM radiated angular momentum J_{rad}^μ . These relations build on a split into conservative and radiative parts as an average of the full in-in and out-out observables. It will be exciting to see how these relations may help produce results at 4PM order. For spin effects this will require a 3PM computation of J_{rad}^μ with spin, but one may already explore nonspinning applications of the formula. Similar studies have already been initiated in the nonspinning case in Ref. [140].

Using the conservative parts of our results—already known from Ref. [61]—we constructed a two-body Hamiltonian mapping our unbound results to bound motion. This Hamiltonian describes the conservative two-body dynamics up to 3PM order and to quadratic order in their spins—it complements the Hamiltonian of Ref. [139] by adding the terms $\mathcal{O}(S_1 S_2)$ and finite-size effects. We note that the coefficients of the Hamiltonian are more complicated than the scattering observables. While the observables can easily be reduced to a number of polynomials in γ , the Hamiltonian coefficients depend on the center-of-mass variables in a complicated manner. This is partly due to its gauge dependence, but also the necessity of using a canonical spin-supplementary condition. This canonical Pryce-Newton-Wigner SSC [142–144] we showed is related to the covariant SSC by a supersymmetry transformation.

It is an interesting study to explore whether gauge choices other than the isotropic gauge could lead to simpler coefficients in the Hamiltonian. Quite intriguingly, we found it possible to fix all coefficients of the Hamiltonian by matching to a single scattering angle defined for generic spins. This in turn leads to the exciting result that the conservative dynamics for generic spins can be described by a single scalar. The expressions for the impulse and spin kick in terms of the scattering angle thus obtained are gauge invariant. They deserve further study and preferably a direct relation highlighting the gauge invariance. Such relations are similar to the eikonal relations explored in Ref. [127].

We also studied mappings to bound orbits. This began with using the complete 3PM Hamiltonian to produce numerical plots of the binding energy for circular orbits leading up to merger, in comparison with numerical relativity simulations [150]. We successfully reproduced the known 4PN quadratic-in-spin binding energy, and the

leading-PM radiated energy for bound orbits. Unfortunately, these plots did not show a significant improvement of the quadratic-in-spin Hamiltonian over its spin-orbit counterpart; however, the effect of going from 2PM to 3PM order was more significant. This calls for the future determination of the 4PM spinning Hamiltonian, which—similar to the nonspinning case, and due to the presence of tails—encounters nonlocalities that distinguish between bound and unbound dynamics [68–71,103,104]. To inform realistic waveform models, it will also be important to incorporate knowledge of the test-body limit by way of the EOB formalism [45,46].

Finally, we determined the energy radiated per orbit in the CoM frame from an appropriate unbound-to-bound mapping [37]. Current limitations in this mapping restrict us to considering only aligned spins; furthermore, this approach does not reproduce the full leading-PN result—which in the nonspinning case may be derived from Einstein’s quadrupole formula. To overcome both of these limitations, and following the suggestion of Ref. [36], we believe that in the future it will be more profitable to focus on reconstructing the (gauge-dependent) instantaneous momentum and angular momentum fluxes. In this case, a complete 4PM result would suffice to reconstruct the leading-PN form of the radiated energy. Alternatively, we hope that improved unbound-to-bound mappings for spinning bodies will further alleviate these issues.

A natural continuation of this work will therefore be to progress upwards in the perturbative series—both to higher PM orders and higher spin orders. While higher PM orders will present a challenge for the integration step, there has recently been a promising development in this area: the first complete analytic result for the 4PM momentum impulse including radiation-reaction effects [70], wherein loop integrals with retarded propagators were also used. With a similar basis of master integrals, it will be possible to tackle spin effects at 4PM order. While higher PM orders present a challenge regarding the integration steps, higher spin orders rather challenge the construction of the integrand. In this case, it will be necessary to upgrade the spinning $\mathcal{N} = 2$ supersymmetric worldline action to include more supersymmetry—a tantalizing prospect that we leave for future work.

ACKNOWLEDGMENTS

We thank Alessandra Buonanno, Gregor Kälin, Jan Plefka, Muddu Saketh, Benjamin Sauer, Chia-Hsien Shen, and Justin Vines for enlightening discussions. We are especially grateful to Jan Steinhoff for his assistance in comparing with the 4PN quadratic-in-spin Hamiltonian, and to Mohammed Khalil for helping us produce the numerical binding energy plot. We also thank the organizers of the “High-Precision Gravitational Waves” program at the Kavli Institute for Theoretical Physics (KITP) for their hospitality. This work is funded by the Deutsche

Forschungsgemeinschaft (DFG, German Research Foundation) Projekt-nummer 417533893/GRK2575 “Rethinking Quantum Field Theory.” It was also supported in part by the National Science Foundation under Grant No. NSF PHY-1748958.

APPENDIX A: CANONICAL EXPANSION OF SCATTERING ANGLE

Here we discuss the relationship between the canonical and covariant expansions of the conservative scattering angle θ_{cons} . As the angle is independent of the choice of SSC, only its expansion coefficients change when we expand in canonical rather than covariant variables. The canonical expansion is defined analogously to the covariant expansion in Eq. (49):

$$\begin{aligned} \theta_{\text{cons}} = & \sum_{n=1}^3 \left(\frac{GM}{|b_{\text{can}}|} \right)^n \left[\theta_{\text{can}}^{(n;0)} - \sum_i \theta_{\text{can}}^{(n;1,i)} \frac{\hat{L}_{\text{can}} \cdot a_{i,\text{can}}}{|b_{\text{can}}|} \right. \\ & + \sum_{i,j} \frac{a_{i,\text{can}}^\mu a_{j,\text{can}}^\nu}{|b_{\text{can}}|^2} \left(-\theta_{\text{can}}^{(n;2,1,i,j)} \eta^{\mu\nu} + \theta_{\text{can}}^{(n;2,2,i,j)} \hat{b}_{\text{can}}^\mu \hat{b}_{\text{can}}^\nu \right. \\ & \left. \left. + \theta_{\text{can}}^{(n;2,3,i,j)} \hat{p}^\mu \hat{p}^\nu + \theta_{\text{can}}^{(n;2,4,i,j)} \hat{b}_{\text{can}}^\mu \hat{p}^\nu \right) \right] \\ & + \mathcal{O}(S^3, G^4). \end{aligned} \quad (\text{A1})$$

Using the definitions of the canonical variables from Sec. II B we relate the canonical expansion coefficients, $\theta_{\text{can}}^{(n;A)}$, to the covariant ones, $\theta_{\text{cons}}^{(n;A)}$:

$$\theta_{\text{can}}^{(n;0)} = \Gamma \theta_{\text{cons}}^{(n;0)}, \quad (\text{A2a})$$

$$\theta_{\text{can}}^{(n;1,i)} = \Gamma \left(\theta_{\text{cons}}^{(n;1,i)} + n \frac{p_\infty}{E_i + m_i} \theta_{\text{cons}}^{(n;0)} \right), \quad (\text{A2b})$$

$$\begin{aligned} \theta_{\text{can}}^{(n;2,1,i,j)} = & \Gamma \left(\theta_{\text{cons}}^{(n;2,1,i,j)} + \frac{(n+1)p_\infty}{2(E_j + m_j)} \theta_{\text{cons}}^{(n;1,i)} \right. \\ & + \frac{(n+1)p_\infty}{2(E_i + m_i)} \theta_{\text{cons}}^{(n;1,j)} \\ & \left. + \frac{n(n+1)p_\infty^2}{2(E_i + m_i)(E_j + m_j)} \theta_{\text{cons}}^{(n;0)} \right), \end{aligned} \quad (\text{A2c})$$

$$\begin{aligned} \theta_{\text{can}}^{(n;2,2,i,j)} = & \Gamma \left(\theta_{\text{cons}}^{(n;2,2,i,j)} - \frac{(n+2)p_\infty}{2(E_j + m_j)} \theta_{\text{cons}}^{(n;1,i)} \right. \\ & - \frac{(n+2)p_\infty}{2(E_i + m_i)} \theta_{\text{cons}}^{(n;1,j)} \\ & \left. - \frac{n(n+2)p_\infty^2}{2(E_i + m_i)(E_j + m_j)} \theta_{\text{cons}}^{(n;0)} \right), \end{aligned} \quad (\text{A2d})$$

$$\begin{aligned}
 \theta_{\text{can}}^{(n;2,3,i,j)} &= \Gamma \left((-1)^{i+j} \frac{p_\infty^2 \Gamma^2}{\mu^2} \theta_{\text{cons}}^{(n;2,3,i,j)} \right. \\
 &+ \frac{1 - (-1)^{i+j} E_1 E_2 - m_1 m_2 + p_\infty^2}{2 m_1 m_2} \theta_{\text{cons}}^{(n;2,1,1,2)} \\
 &- \frac{(n+1)p_\infty}{2(E_j + m_j)} \theta_{\text{cons}}^{(n;1,i)} - \frac{(n+1)p_\infty}{2(E_i + m_i)} \theta_{\text{cons}}^{(n;1,j)} \\
 &\left. - \frac{n(n+1)p_\infty^2}{2(E_i + m_i)(E_j + m_j)} \theta_{\text{cons}}^{(n;0)} \right). \quad (\text{A2e})
 \end{aligned}$$

The covariant coefficients $\theta_{\text{cons}}^{(n;A)}$ are given in Appendix B.

APPENDIX B: SCATTERING ANGLE

In this section we print the 3PM contributions to the covariant expansion coefficients of the scattering angle θ defined in Eqs. (48) and (49). First, we print the conservative contributions which are used for the Hamiltonian. Then, we print the radiative contributions. The coefficients that we have not printed here may be obtained by exchanging the two particles. The conservative nonspinning and spin-orbit coefficients are

$$\theta_{\text{cons}}^{(3;0)} = \frac{2(64\gamma^6 - 120\gamma^4 + 60\gamma^2 - 5)\Gamma^2}{3(\gamma^2 - 1)^3} - \frac{8\gamma(14\gamma^2 + 25)\nu}{3(\gamma^2 - 1)} - \frac{8(4\gamma^4 - 12\gamma^2 - 3)\nu \text{arccosh}\gamma}{(\gamma^2 - 1)^{3/2}}, \quad (\text{B1})$$

$$\theta_{\text{cons}}^{(3;1,1)} = -\frac{2\gamma(16\gamma^4 - 20\gamma^2 + 5)(5\Gamma^2 - \delta)}{(\gamma^2 - 1)^{5/2}} + \frac{4(44\gamma^4 + 100\gamma^2 + 41)\nu}{(\gamma^2 - 1)^{3/2}} + \frac{48\gamma(\gamma^2 - 6)(2\gamma^2 + 1)\nu \text{arccosh}\gamma}{(\gamma^2 - 1)^2}. \quad (\text{B2})$$

These agree with the ones in Ref. [61] as expected from the discussion beneath Eq. (48). The coefficients $\theta_{\text{cons}}^{(3;2,1,i,j)}$ are

$$\begin{aligned}
 \theta_{\text{cons}}^{(3;2,1,1,1)} &= \Gamma^2 \left(\frac{4(96\gamma^6 - 160\gamma^4 + 70\gamma^2 - 5)}{(\gamma^2 - 1)^3} - \frac{4(1772\gamma^6 - 2946\gamma^4 + 1346\gamma^2 - 137)C_{E,1}}{35(\gamma^2 - 1)^3} \right) \\
 &+ \delta \left(-\frac{8(4\gamma^2 - 2\gamma - 1)(4\gamma^2 + 2\gamma - 1)}{(\gamma^2 - 1)^2} + \frac{8(214\gamma^4 - 223\gamma^2 + 44)C_{E,1}}{35(\gamma^2 - 1)^2} \right) \\
 &- \frac{16\gamma(148\gamma^4 + 374\gamma^2 + 383)\nu}{5(\gamma^2 - 1)^2} + \frac{8\gamma(3244\gamma^4 + 7972\gamma^2 + 4639)C_{E,1}\nu}{105(\gamma^2 - 1)^2} \\
 &+ \text{arccosh}\gamma \left(-\frac{192(\gamma^6 - 8\gamma^4 - 7\gamma^2 - 1)\nu}{(\gamma^2 - 1)^{5/2}} + \frac{16(8\gamma^6 - 56\gamma^4 - 24\gamma^2 - 3)C_{E,1}\nu}{(\gamma^2 - 1)^{5/2}} \right), \quad (\text{B3})
 \end{aligned}$$

$$\theta_{\text{cons}}^{(3;2,1,1,2)} = \frac{4(96\gamma^6 - 160\gamma^4 + 70\gamma^2 - 5)\Gamma^2}{(\gamma^2 - 1)^3} - \frac{32\gamma(15\gamma^4 + 46\gamma^2 + 47)\nu}{(\gamma^2 - 1)^2} - \frac{48(4\gamma^6 - 36\gamma^4 - 35\gamma^2 - 5)\nu \text{arccosh}\gamma}{(\gamma^2 - 1)^{5/2}}. \quad (\text{B4})$$

The coefficients $\theta_{\text{cons}}^{(3;2,2,i,j)}$ are

$$\begin{aligned}
 \theta_{\text{cons}}^{(3;2,2,1,1)} &= \frac{4\gamma(9000\gamma^{10} + 4404\gamma^8 - 2152\gamma^6 - 12152\gamma^4 + 8379\gamma^2 - 1479)\nu}{15(\gamma^2 - 1)^3(2\gamma^2 - 1)^2} \\
 &+ \pi^2 \left(-\frac{(5\gamma^2 - 3)(10\gamma^4 - 5\gamma^2 + 9)\gamma^2 \delta}{128(\gamma^2 - 1)^2(2\gamma^2 - 1)^2} + \frac{(100\gamma^8 - 160\gamma^6 + 193\gamma^4 - 78\gamma^2 + 45)\gamma^2 \Gamma^2}{128(\gamma^2 - 1)^2(2\gamma^2 - 1)^3} \right. \\
 &- \frac{(100\gamma^9 - 160\gamma^7 - 60\gamma^6 + 193\gamma^5 - 12\gamma^4 - 78\gamma^3 + 12\gamma^2 + 45\gamma - 36)\gamma^2 \nu}{64(\gamma^2 - 1)^2(2\gamma^2 - 1)^3} \left. + \delta \left(\frac{10(4\gamma^2 - 2\gamma - 1)(4\gamma^2 + 2\gamma - 1)}{(\gamma^2 - 1)^2} \right) \right. \\
 &- \frac{2(1192\gamma^4 - 1382\gamma^2 + 295)C_{E,1}}{35(\gamma^2 - 1)^2} \left. - \frac{4\gamma(10744\gamma^6 + 13474\gamma^4 + 2665\gamma^2 + 9237)C_{E,1}\nu}{105(\gamma^2 - 1)^3} \right) \\
 &+ \Gamma^2 \left(\frac{2(6568\gamma^6 - 11114\gamma^4 + 5079\gamma^2 - 463)C_{E,1}}{35(\gamma^2 - 1)^3} - \frac{2(960\gamma^{10} - 2560\gamma^8 + 2540\gamma^6 - 1150\gamma^4 + 225\gamma^2 - 13)}{(\gamma^2 - 1)^3(2\gamma^2 - 1)^2} \right) \\
 &+ \text{arccosh}\gamma \left(\frac{16\gamma^2(60\gamma^{10} - 600\gamma^8 + 551\gamma^6 - 63\gamma^4 - 63\gamma^2 + 15)\nu}{(\gamma^2 - 1)^{7/2}(2\gamma^2 - 1)^2} - \frac{32(8\gamma^8 - 56\gamma^6 + 26\gamma^4 - 18\gamma^2 - 3)C_{E,1}\nu}{(\gamma^2 - 1)^{7/2}} \right), \quad (\text{B5})
 \end{aligned}$$

$$\begin{aligned}
\theta_{\text{cons}}^{(3;2,2,1,2)} = & -\frac{2(960\gamma^{10} - 2560\gamma^8 + 2540\gamma^6 - 1150\gamma^4 + 225\gamma^2 - 13)\Gamma^2}{(\gamma^2 - 1)^3(2\gamma^2 - 1)^2} \\
& + \frac{4\gamma(1800\gamma^{10} + 2020\gamma^8 - 424\gamma^6 - 3032\gamma^4 + 1703\gamma^2 - 231)\nu}{3(\gamma^2 - 1)^3(2\gamma^2 - 1)^2} \\
& + \frac{16(60\gamma^{12} - 664\gamma^{10} + 519\gamma^8 - 31\gamma^6 - 43\gamma^4 + 7\gamma^2 - 1)\nu \text{arccosh}\gamma}{(\gamma^2 - 1)^{7/2}(2\gamma^2 - 1)^2} \\
& + \pi^2 \left(\frac{3(\gamma^2 + 1)(5\gamma^4 - 4\gamma^2 + 3)\gamma^2\Gamma^2}{32(\gamma^2 - 1)^2(2\gamma^2 - 1)^3} + \frac{(100\gamma^8 - 60\gamma^7 - 160\gamma^6 - 12\gamma^5 + 193\gamma^4 + 12\gamma^3 - 78\gamma^2 - 36\gamma + 45)\gamma^2\nu}{64(\gamma^2 - 1)^2(2\gamma^2 - 1)^3} \right). \tag{B6}
\end{aligned}$$

The coefficients $\theta_{\text{cons}}^{(3;2,3,i,j)}$ are

$$\begin{aligned}
\theta_{\text{cons}}^{(3;2,3,1,1)} = & \frac{\gamma(18624\gamma^8 + 24848\gamma^6 - 45192\gamma^4 - 58631\gamma^2 + 36351)\nu}{15(\gamma^2 - 1)^4(2\gamma^2 - 1)} + \delta \left(\frac{704\gamma^6 - 880\gamma^4 + 312\gamma^2 - 23}{2(\gamma^2 - 1)^3(2\gamma^2 - 1)} \right. \\
& \left. - \frac{2(1376\gamma^4 - 1294\gamma^2 + 233)C_{E,1}}{35(\gamma^2 - 1)^3} \right) - \frac{4\gamma(8720\gamma^6 + 14894\gamma^4 - 22663\gamma^2 - 37071)C_{E,1}\nu}{105(\gamma^2 - 1)^4} \\
& + \Gamma^2 \left(\frac{2(4064\gamma^6 - 6562\gamma^4 + 2997\gamma^2 - 359)C_{E,1}}{35(\gamma^2 - 1)^4} - \frac{1728\gamma^8 - 3664\gamma^6 + 2584\gamma^4 - 673\gamma^2 + 41}{2(\gamma^2 - 1)^4(2\gamma^2 - 1)} \right) \\
& + \text{arccosh}\gamma \left(\frac{64(3\gamma^8 - 35\gamma^6 + 9\gamma^4 + 42\gamma^2 + 6)\nu}{(\gamma^2 - 1)^{9/2}} - \frac{16(8\gamma^8 - 80\gamma^6 + 44\gamma^4 + 99\gamma^2 + 15)C_{E,1}\nu}{(\gamma^2 - 1)^{9/2}} \right), \tag{B7}
\end{aligned}$$

$$\begin{aligned}
\theta_{\text{cons}}^{(3;2,3,1,2)} = & \frac{4\gamma(96\gamma^6 - 148\gamma^4 + 55\gamma^2 - 1)\Gamma^2}{(\gamma^2 - 1)^4} - \frac{16\gamma(12\gamma^8 - 136\gamma^6 - 21\gamma^4 + 210\gamma^2 + 88)\nu \text{arccosh}\gamma}{(\gamma^2 - 1)^{9/2}} \\
& - \frac{(2880\gamma^{10} + 7712\gamma^8 - 5664\gamma^6 - 22688\gamma^4 + 9219\gamma^2 + 1197)\nu}{3(\gamma^2 - 1)^4(2\gamma^2 - 1)}. \tag{B8}
\end{aligned}$$

The coefficients $\theta_{\text{cons}}^{(3;2,4,i,j)}$ are

$$\begin{aligned}
\theta_{\text{cons}}^{(3;2,4,1,1)} = & \pi \left(-\frac{(80\gamma^9 - 144\gamma^7 - 12\gamma^6 + 42\gamma^5 + 33\gamma^4 + 32\gamma^3 - 21\gamma^2 - 18\gamma + 6)\nu}{4(\gamma^2 - 1)^{5/2}(2\gamma^2 - 1)^2} - \frac{3(30\gamma^3 - 15\gamma^2 - 6\gamma + 1)C_{E,1}\nu}{4(\gamma^2 - 1)^{3/2}} \right. \\
& + \Gamma^2 \left(\frac{80\gamma^8 - 104\gamma^6 + 12\gamma^5 + 20\gamma^4 - 15\gamma^3 + 18\gamma^2 + 9\gamma - 6}{8(\gamma^2 - 1)^{5/2}(2\gamma^2 - 1)^2} + \frac{3(30\gamma^4 + 15\gamma^3 - 21\gamma^2 - \gamma + 3)C_{E,1}}{8(\gamma^2 - 1)^{5/2}} \right) \\
& \left. + \delta \left(-\frac{80\gamma^8 - 104\gamma^6 - 12\gamma^5 + 20\gamma^4 + 15\gamma^3 + 18\gamma^2 - 9\gamma - 6}{8(\gamma^2 - 1)^{5/2}(2\gamma^2 - 1)^2} - \frac{3(30\gamma^4 - 15\gamma^3 - 21\gamma^2 + \gamma + 3)C_{E,1}}{8(\gamma^2 - 1)^{5/2}} \right) \right), \tag{B9}
\end{aligned}$$

$$\begin{aligned}
\theta_{\text{cons}}^{(3;2,4,1,2)} = & \pi \left(-\frac{(20\gamma^7 + 12\gamma^6 - 21\gamma^5 - 24\gamma^4 + 4\gamma^3 + 15\gamma^2 + 3\gamma - 3)\Gamma^2}{4(\gamma^2 - 1)^{5/2}(2\gamma^2 - 1)^2} \right. \\
& + \frac{(20\gamma^7 - 12\gamma^6 - 21\gamma^5 + 24\gamma^4 + 4\gamma^3 - 15\gamma^2 + 3\gamma + 3)\delta}{4(\gamma^2 - 1)^{5/2}(2\gamma^2 - 1)^2} \\
& \left. + \frac{(120\gamma^8 - 56\gamma^7 - 194\gamma^6 + 48\gamma^5 + 124\gamma^4 - \gamma^3 - 36\gamma^2 - 9\gamma + 6)\nu}{4(\gamma^2 - 1)^{5/2}(2\gamma^2 - 1)^2} \right). \tag{B10}
\end{aligned}$$

Let us then print the radiative contributions. All coefficients are proportional to the universal function $\mathcal{I}(v)$ except $\theta_{k,\text{rad}}^{3;2,4,i,j}$. These coefficients depend on the particle label k :

$$\theta_{\text{rad}}^{(3;0)} = \frac{4(1-2\gamma^2)^2\nu}{(\gamma^2-1)^{3/2}}\mathcal{I}(v), \quad (\text{B11})$$

$$\theta_{\text{rad}}^{(3;1,1)} = -\frac{24\gamma(2\gamma^2-1)\nu}{\gamma^2-1}\mathcal{I}(v), \quad (\text{B12})$$

$$\theta_{\text{rad}}^{(3;2,1,1,1)} = -\frac{16\nu(\gamma^4(4C_{E,1}-6) + \gamma^2(6-4C_{E,1}) + C_{E,1}-1)}{(\gamma^2-1)^{3/2}}\mathcal{I}(v), \quad (\text{B13})$$

$$\theta_{\text{rad}}^{(3;2,1,1,2)} = \frac{16(6\gamma^4-6\gamma^2+1)\nu}{(\gamma^2-1)^{3/2}}\mathcal{I}(v), \quad (\text{B14})$$

$$\theta_{\text{rad}}^{(3;2,2,1,1)} = \frac{8\nu(\gamma^4(16C_{E,1}-15) + \gamma^2(15-16C_{E,1}) + 4(C_{E,1}-1))}{(\gamma^2-1)^{3/2}}\mathcal{I}(v), \quad (\text{B15})$$

$$\theta_{\text{rad}}^{(3;2,2,1,2)} = -\frac{8(15\gamma^4-15\gamma^2+4)\nu}{(\gamma^2-1)^{3/2}}\mathcal{I}(v), \quad (\text{B16})$$

$$\theta_{\text{rad}}^{(3;2,3,1,1)} = \frac{16\nu(\gamma^4(4C_{E,1}-6) + \gamma^2(6-4C_{E,1}) + C_{E,1}-1)}{(\gamma^2-1)^{5/2}}\mathcal{I}(v), \quad (\text{B17})$$

$$\theta_{\text{rad}}^{(3;2,3,1,2)} = \frac{16\gamma(6\gamma^4-6\gamma^2+1)\nu}{(\gamma^2-1)^{5/2}}\mathcal{I}(v). \quad (\text{B18})$$

The coefficients $\theta_{k,\text{rad}}^{3;2,4,i,j}$ are

$$\begin{aligned} & \theta_{1,\text{rad}}^{(3;2,4,1,1)} \\ &= \frac{3\pi\gamma(28\gamma^8-80\gamma^6+41\gamma^4+34\gamma^2-15)\nu\text{arccosh}\gamma}{32(\gamma^2-1)^{7/2}(2\gamma^2-1)} \\ & \quad - \frac{3\pi(14\gamma^7+14\gamma^6+101\gamma^5-323\gamma^4+236\gamma^3-60\gamma^2-23\gamma+25)\nu\log(\frac{\gamma+1}{2})}{16(\gamma-1)^2(\gamma+1)^3(2\gamma^2-1)} \\ & \quad + \frac{\pi(280\gamma^{10}+470\gamma^9+1930\gamma^8+5397\gamma^7-88373\gamma^6+244865\gamma^5-273845\gamma^4+73199\gamma^3+112249\gamma^2-102411\gamma+25759)\nu}{160(\gamma-1)^3(\gamma+1)^5(2\gamma^2-1)} \\ & \quad + C_{E,1} \left(\frac{3\pi\gamma(98\gamma^6-239\gamma^4+164\gamma^2-39)\nu\text{arccosh}\gamma}{64(\gamma^2-1)^{7/2}} - \frac{3\pi(49\gamma^5+169\gamma^4-306\gamma^3+150\gamma^2-63\gamma+33)\nu\log(\frac{\gamma+1}{2})}{32(\gamma-1)^2(\gamma+1)^3} \right. \\ & \quad \left. - \frac{\pi(1575\gamma^9-810\gamma^8+1020\gamma^7+4140\gamma^6+26442\gamma^5-196568\gamma^4+442476\gamma^3-521244\gamma^2+321447\gamma-80398)\nu}{640(\gamma-1)^3(\gamma+1)^5} \right), \quad (\text{B19}) \end{aligned}$$

$$\begin{aligned} & \theta_{1,\text{rad}}^{(3;2,4,2,2)} \\ &= -\frac{3\pi(2\gamma^4-13\gamma^2+15)\gamma^2\nu\text{arccosh}\gamma}{16(\gamma^2-1)^{7/2}(2\gamma^2-1)} + \frac{3\pi(-110\gamma^5+259\gamma^4-186\gamma^3+8\gamma^2+40\gamma-19)\nu\log(\frac{\gamma+1}{2})}{8(\gamma-1)^2(\gamma+1)^3(2\gamma^2-1)} \\ & \quad + \frac{\pi(-2070\gamma^9+2740\gamma^8+20777\gamma^7-153646\gamma^6+408765\gamma^5-463000\gamma^4+119039\gamma^3+196158\gamma^2-168991\gamma+40708)\nu}{160(\gamma-1)^3(\gamma+1)^5(2\gamma^2-1)} \\ & \quad + C_{E,2} \left(-\frac{3\pi(30\gamma^4-59\gamma^2+21)\gamma^2\nu\text{arccosh}\gamma}{32(\gamma^2-1)^{7/2}} - \frac{3\pi(65\gamma^4-190\gamma^3+156\gamma^2-74\gamma+27)\nu\log(\frac{\gamma+1}{2})}{16(\gamma-1)^2(\gamma+1)^3} \right. \\ & \quad \left. + \frac{\pi(-1075\gamma^7+980\gamma^6-6738\gamma^5+73384\gamma^4-198749\gamma^3+236162\gamma^2-135438\gamma+30994)\nu}{160(\gamma-1)^3(\gamma+1)^5} \right), \quad (\text{B20}) \end{aligned}$$

$$\begin{aligned}
& \theta_{1,\text{rad}}^{(3;2,4,1,2)} \\
&= -\frac{3\pi(8\gamma^6 - 22\gamma^4 + 9\gamma^2 + 9)\gamma^2 \nu \operatorname{arccosh} \gamma}{16(\gamma^2 - 1)^{7/2}(2\gamma^2 - 1)} + \frac{3\pi(17\gamma^6 - 63\gamma^5 + 107\gamma^4 - 81\gamma^3 + 8\gamma^2 + 16\gamma - 8)\nu \log(\frac{\gamma+1}{2})}{4(\gamma-1)^2(\gamma+1)^3(2\gamma^2-1)} \\
&+ \frac{\pi(-630\gamma^{10} - 480\gamma^9 + 4481\gamma^8 + 11476\gamma^7 - 88357\gamma^6 + 195970\gamma^5 - 191553\gamma^4 + 34776\gamma^3 + 87891\gamma^2 - 69710\gamma + 16328)\nu}{64(\gamma-1)^3(\gamma+1)^5(2\gamma^2-1)},
\end{aligned} \tag{B21}$$

$$\begin{aligned}
& \theta_{1,\text{rad}}^{(3;2,4,2,1)} = \frac{3\pi\gamma(4\gamma^6 - 24\gamma^4 + 33\gamma^2 - 9)\nu \operatorname{arccosh} \gamma}{16(\gamma^2 - 1)^{7/2}(2\gamma^2 - 1)} + \frac{3\pi(20\gamma^6 - 76\gamma^5 + 120\gamma^4 - 73\gamma^3 - \gamma^2 + 25\gamma - 11)\nu \log(\frac{\gamma+1}{2})}{4(\gamma-1)^2(\gamma+1)^3(2\gamma^2-1)} \\
&+ \frac{\pi(-98\gamma^9 + 1488\gamma^8 + 9431\gamma^7 - 75972\gamma^6 + 188037\gamma^5 - 200914\gamma^4 + 48417\gamma^3 + 85464\gamma^2 - 73947\gamma + 17902)\nu}{64(\gamma-1)^3(\gamma+1)^5(2\gamma^2-1)}.
\end{aligned} \tag{B22}$$

APPENDIX C: HAMILTONIAN COEFFICIENTS

Here we give the Hamiltonian coefficients in terms of those of the scattering angle. We do not print the expressions for the spin-spin coefficients at 3PM order which are found in the Supplemental Material [145]. We start with the simpler spinless and spin-orbit coefficients. At 1PM order we find

$$\begin{aligned}
c^{(1;0)}(\mathbf{p}^2) &= -\frac{P_\infty^2}{2E\xi} \theta_{\text{can}}^{(1;0)} \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \\
c^{(1;1,i)}(\mathbf{p}^2) &= -\frac{P_\infty}{2E\xi} \theta_{\text{can}}^{(1;1,i)} \Big|_{p_\infty \rightarrow |\mathbf{p}|}.
\end{aligned} \tag{C1}$$

At 2PM order we find

$$\begin{aligned}
c^{(2;0)}(\mathbf{p}^2) &= -\frac{P_\infty^2}{\pi E\xi} \theta_{\text{can}}^{(2;0)} + \frac{1}{8p_\infty} \mathcal{D} \left[\frac{P_\infty^3}{E\xi} (\theta_{\text{can}}^{(1;0)})^2 \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \\
c^{(2;1,i)}(\mathbf{p}^2) &= -\frac{P_\infty}{\pi E\xi} \theta_{\text{can}}^{(2;1,i)} + \frac{1}{4p_\infty^3} \mathcal{D} \left[\frac{P_\infty^4}{E\xi} \theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;1,i)} \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}.
\end{aligned} \tag{C2}$$

Finally we reprint the nonspinning and spin-orbit coefficients at 3PM order from Eq. (78):

$$\begin{aligned}
c^{(3;0)}(\mathbf{p}^2) &= -\frac{P_\infty^2}{4E\xi} \theta_{\text{can}}^{(3;0)} + \frac{1}{2\pi p_\infty^2} \mathcal{D} \left[\frac{P_\infty^4}{E\xi} \theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(2;0)} \right] - \frac{1}{48p_\infty^2} \mathcal{D}^2 \left[\frac{P_\infty^4}{E\xi} (\theta_{\text{can}}^{(1;0)})^3 \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \\
c^{(3;1,i)}(\mathbf{p}^2) &= -\frac{P_\infty}{4E\xi} \theta_{\text{can}}^{(3;1,i)} + \frac{1}{2\pi p_\infty^4} \mathcal{D} \left[\frac{P_\infty^5}{E\xi} (\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(2;1,i)} + \theta_{\text{can}}^{(2;0)} \theta_{\text{can}}^{(1;1,i)}) \right] - \frac{1}{16p_\infty^4} \mathcal{D}^2 \left[\frac{P_\infty^5}{E\xi} (\theta_{\text{can}}^{(1;0)})^2 \theta_{\text{can}}^{(1;1,i)} \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}.
\end{aligned} \tag{C3}$$

The spin-spin coefficients do not seem to obey the same simplicity as the above spinless and spin-orbit coefficients. At the first post-Minkowskian order we find

$$c^{(1;2,1,i,j)}(\mathbf{p}^2) = -\frac{P_\infty^2}{4E\xi} \theta_{\text{can}}^{(1;2,1,i,j)} \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \tag{C4a}$$

$$c^{(1;2,2,i,j)}(\mathbf{p}^2) = -\frac{3P_\infty^2}{8E\xi} \theta_{\text{can}}^{(1;2,2,i,j)} + \frac{3P_\infty^2}{16E\xi} \frac{\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}}{\theta_{\text{can}}^{(1;0)}} \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \tag{C4b}$$

$$c^{(1;2,3,i,j)}(\mathbf{p}^2) = -\frac{1}{4E\xi} \left(\theta_{\text{can}}^{(1;2,3,i,j)} - \frac{1}{2} \theta_{\text{can}}^{(1;2,2,i,j)} \right) - \frac{1}{16E\xi} \frac{\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}}{\theta_{\text{can}}^{(1;0)}} \Big|_{p_\infty \rightarrow |\mathbf{p}|}. \quad (\text{C4c})$$

At second post-Minkowskian order we find

$$c^{(2;2,1,i,j)}(\mathbf{p}^2) = -\frac{2p_\infty^2}{3\pi E\xi} \theta_{\text{can}}^{(2;2,1,i,j)} + \frac{1}{32p_\infty^3} \mathcal{D} \left[\frac{p_\infty^5}{E\xi} (3\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)} + 4\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;2,1,i,j)}) \right] \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \quad (\text{C5a})$$

$$\begin{aligned} c^{(2;2,2,i,j)}(\mathbf{p}^2) = & -\frac{p_\infty^2}{E\xi} \left(\frac{8\theta_{\text{can}}^{(2;2,2,i,j)}}{9\pi} + \frac{\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;2,2,i,j)}}{8} - \frac{\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}}{16} + \frac{p_\infty}{16} \left(\frac{\theta_{\text{can}}^{(1;1,i)}}{m_i} + \frac{\theta_{\text{can}}^{(1;2,j)}}{m_j} \right) \theta_{\text{can}}^{(1;2,2,i,j)} \right) \\ & + \frac{1}{32p_\infty^3} \mathcal{D} \left[\frac{p_\infty^5}{E\xi} (6\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;2,2,i,j)} - 7\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}) \right] - \frac{4p_\infty^2}{9\pi E\xi} \frac{\theta_{\text{can}}^{(2;0)} \theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}}{(\theta_{\text{can}}^{(1;0)})^2} \\ & + \frac{p_\infty^2}{E\xi \theta_{\text{can}}^{(1;0)}} \left(\frac{p_\infty}{32} \left(\frac{\theta_{\text{can}}^{(1;1,i)}}{m_i} + \frac{\theta_{\text{can}}^{(1;1,j)}}{m_j} \right) \theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)} + \frac{2}{9\pi} (\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(2;1,j)} + \theta_{\text{can}}^{(2;1,i)} \theta_{\text{can}}^{(1;1,j)}) \right) \Big|_{p_\infty \rightarrow |\mathbf{p}|}, \quad (\text{C5b}) \end{aligned}$$

$$\begin{aligned} c^{(2;2,3,i,j)}(\mathbf{p}^2) = & -\frac{1}{E\xi} \left(\frac{2\theta_{\text{can}}^{(2;2,3,i,j)}}{3\pi} - \frac{2\theta_{\text{can}}^{(2;2,2,i,j)}}{9\pi} - \frac{\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;2,2,i,j)}}{8} - \frac{p_\infty}{16} \left(\frac{\theta_{\text{can}}^{(1;1,i)}}{m_i} + \frac{\theta_{\text{can}}^{(1;2,j)}}{m_j} \right) \theta_{\text{can}}^{(1;2,2,i,j)} \right) \\ & + \frac{\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}}{16} - \frac{1}{32p_\infty^5} \mathcal{D} \left[\frac{p_\infty^5}{E\xi} (2\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;2,2,i,j)} + \theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)} - 4\theta_{\text{can}}^{(1;0)} \theta_{\text{can}}^{(1;2,3,i,j)}) \right] \\ & - \frac{1}{E\xi \theta_{\text{can}}^{(1;0)}} \left(\frac{p_\infty}{32} \left(\frac{\theta_{\text{can}}^{(1;1,i)}}{m_i} + \frac{\theta_{\text{can}}^{(1;1,j)}}{m_j} \right) \theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)} + \frac{1}{18\pi} (\theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(2;1,j)} + \theta_{\text{can}}^{(2;1,i)} \theta_{\text{can}}^{(1;1,j)}) \right) \\ & + \frac{1}{9\pi E\xi} \frac{\theta_{\text{can}}^{(2;0)} \theta_{\text{can}}^{(1;1,i)} \theta_{\text{can}}^{(1;1,j)}}{(\theta_{\text{can}}^{(1;0)})^2} \Big|_{p_\infty \rightarrow |\mathbf{p}|}. \quad (\text{C5c}) \end{aligned}$$

The 3PM spin-spin coefficients are found in the Supplemental Material [145].

-
- [1] B. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), Observation of Gravitational Waves from a Binary Black Hole Merger, *Phys. Rev. Lett.* **116**, 061102 (2016).
- [2] B. P. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral, *Phys. Rev. Lett.* **119**, 161101 (2017).
- [3] B. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs, *Phys. Rev. X* **9**, 031040 (2019).
- [4] R. Abbott *et al.* (LIGO Scientific and Virgo Collaborations), GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo during the First Half of the Third Observing Run, *Phys. Rev. X* **11**, 021053 (2021).
- [5] R. Abbott *et al.* (LIGO Scientific and VIRGO Collaborations), GWTC-2.1: Deep extended catalog of compact binary coalescences observed by LIGO and Virgo during the first half of the third observing run, [arXiv:2108.01045](https://arxiv.org/abs/2108.01045).
- [6] R. Abbott *et al.* (LIGO Scientific, VIRGO, and KAGRA Collaborations), GWTC-3: Compact binary coalescences observed by LIGO and Virgo during the second part of the third observing run, [arXiv:2111.03606](https://arxiv.org/abs/2111.03606).
- [7] S. W. Ballmer *et al.*, Snowmass2021 cosmic frontier white paper: Future gravitational-wave detector facilities, [arXiv:2203.08228](https://arxiv.org/abs/2203.08228).
- [8] W. D. Goldberger and I. Z. Rothstein, An effective field theory of gravity for extended objects, *Phys. Rev. D* **73**, 104029 (2006).
- [9] W. D. Goldberger and I. Z. Rothstein, Towers of gravitational theories, *Gen. Relativ. Gravit.* **38**, 1537 (2006).
- [10] W. D. Goldberger and A. Ross, Gravitational radiative corrections from effective field theory, *Phys. Rev. D* **81**, 124015 (2010).
- [11] R. A. Porto, The effective field theorist's approach to gravitational dynamics, *Phys. Rep.* **633**, 1 (2016).

- [12] M. Levi, Effective field theories of post-Newtonian gravity: A comprehensive review, *Rep. Prog. Phys.* **83**, 075901 (2020).
- [13] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, Testing binary dynamics in gravity at the sixth post-Newtonian level, *Phys. Lett. B* **807**, 135496 (2020).
- [14] D. Bini, T. Damour, and A. Geralico, Sixth post-Newtonian local-in-time dynamics of binary systems, *Phys. Rev. D* **102**, 024061 (2020).
- [15] D. Bini, T. Damour, and A. Geralico, Sixth post-Newtonian nonlocal-in-time dynamics of binary systems, *Phys. Rev. D* **102**, 084047 (2020).
- [16] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, The fifth-order post-Newtonian Hamiltonian dynamics of two-body systems from an effective field theory approach: Potential contributions, *Nucl. Phys.* **B965**, 115352 (2021).
- [17] S. Foffa, R. Sturani, and W. J. Torres Bobadilla, Efficient resummation of high post-Newtonian contributions to the binding energy, *J. High Energy Phys.* **02** (2021) 165.
- [18] J. Blümlein, A. Maier, P. Marquard, and G. Schäfer, The 6th post-Newtonian potential terms at $O(G_N^4)$, *Phys. Lett. B* **816**, 136260 (2021).
- [19] R. A. Porto, Post-Newtonian corrections to the motion of spinning bodies in NRGR, *Phys. Rev. D* **73**, 104031 (2006).
- [20] M. Levi and J. Steinhoff, Spinning gravitating objects in the effective field theory in the post-Newtonian scheme, *J. High Energy Phys.* **09** (2015) 219.
- [21] W. D. Goldberger, J. Li, and I. Z. Rothstein, Non-conservative effects on spinning black holes from world-line effective field theory, *J. High Energy Phys.* **06** (2021) 053.
- [22] M. V. S. Saketh and J. Vines, Scattering of gravitational waves off spinning compact objects with an effective worldline theory, *Phys. Rev. D* **106**, 124026 (2022).
- [23] M. Levi, A. J. McLeod, and M. Von Hippel, N^3 LO gravitational spin-orbit coupling at order G^4 , *J. High Energy Phys.* **07** (2021) 115.
- [24] J.-W. Kim, M. Levi, and Z. Yin, N^3 LO spin-orbit interaction via the EFT of spinning gravitating objects, [arXiv:2208.14949](https://arxiv.org/abs/2208.14949).
- [25] M. K. Mandal, P. Mastrolia, R. Patil, and J. Steinhoff, Gravitational spin-orbit Hamiltonian at NNNLO in the post-Newtonian framework, [arXiv:2209.00611](https://arxiv.org/abs/2209.00611).
- [26] M. Levi, Binary dynamics from spin1-spin2 coupling at fourth post-Newtonian order, *Phys. Rev. D* **85**, 064043 (2012).
- [27] M. Levi and J. Steinhoff, Equivalence of ADM Hamiltonian and effective field theory approaches at next-to-next-to-leading order spin1-spin2 coupling of binary inspirals, *J. Cosmol. Astropart. Phys.* **12** (2014) 003.
- [28] M. Levi and J. Steinhoff, Next-to-next-to-leading order gravitational spin-squared potential via the effective field theory for spinning objects in the post-Newtonian scheme, *J. Cosmol. Astropart. Phys.* **01** (2016) 008.
- [29] M. Levi and J. Steinhoff, Complete conservative dynamics for inspiralling compact binaries with spins at the fourth post-Newtonian order, *J. Cosmol. Astropart. Phys.* **09** (2021) 029.
- [30] J.-W. Kim, M. Levi, and Z. Yin, Quadratic-in-spin interactions at fifth post-Newtonian order probe new physics, *Phys. Lett. B* **834**, 137410 (2022).
- [31] M. Levi, A. J. McLeod, and M. Von Hippel, N^3 LO gravitational quadratic-in-spin interactions at G^4 , *J. High Energy Phys.* **07** (2021) 116.
- [32] G. Cho, R. A. Porto, and Z. Yang, Gravitational radiation from inspiralling compact objects: Spin effects to the fourth post-Newtonian order, *Phys. Rev. D* **106**, L101501 (2022).
- [33] J.-W. Kim, M. Levi, and Z. Yin, N^3 LO quadratic-in-spin interactions for generic compact binaries, [arXiv:2209.09235](https://arxiv.org/abs/2209.09235).
- [34] G. Kälin and R. A. Porto, From boundary data to bound states, *J. High Energy Phys.* **01** (2020) 072.
- [35] G. Kälin and R. A. Porto, From boundary data to bound states. Part II. Scattering angle to dynamical invariants (with twist), *J. High Energy Phys.* **02** (2020) 120.
- [36] M. V. S. Saketh, J. Vines, J. Steinhoff, and A. Buonanno, Conservative and radiative dynamics in classical relativistic scattering and bound systems, *Phys. Rev. Res.* **4**, 013127 (2022).
- [37] G. Cho, G. Kälin, and R. A. Porto, From boundary data to bound states. Part III. Radiative effects, *J. High Energy Phys.* **04** (2022) 154.
- [38] N. Bjerrum-Bohr, J. F. Donoghue, and P. Vanhove, On-shell techniques and universal results in quantum gravity, *J. High Energy Phys.* **02** (2014) 111.
- [39] C. Cheung, I. Z. Rothstein, and M. P. Solon, From Scattering Amplitudes to Classical Potentials in the Post-Minkowskian Expansion, *Phys. Rev. Lett.* **121**, 251101 (2018).x
- [40] D. Neill and I. Z. Rothstein, Classical space-times from the S matrix, *Nucl. Phys.* **B877**, 177 (2013).
- [41] V. Vaidya, Gravitational spin Hamiltonians from the S matrix, *Phys. Rev. D* **91**, 024017 (2015).
- [42] T. Damour, High-energy gravitational scattering and the general relativistic two-body problem, *Phys. Rev. D* **97**, 044038 (2018).
- [43] N. E. J. Bjerrum-Bohr, A. Cristofoli, and P. H. Damgaard, Post-Minkowskian scattering angle in Einstein gravity, *J. High Energy Phys.* **08** (2020) 038.
- [44] A. Cristofoli, P. H. Damgaard, P. Di Vecchia, and C. Heissenberg, Second-order post-Minkowskian scattering in arbitrary dimensions, *J. High Energy Phys.* **07** (2020) 122.
- [45] A. Buonanno and T. Damour, Effective one-body approach to general relativistic two-body dynamics, *Phys. Rev. D* **59**, 084006 (1999).
- [46] T. Damour, Introductory lectures on the effective one body formalism, *Int. J. Mod. Phys. A* **23**, 1130 (2008).
- [47] T. Damour, Classical and quantum scattering in post-Minkowskian gravity, *Phys. Rev. D* **102**, 024060 (2020).
- [48] A. Antonelli, A. Buonanno, J. Steinhoff, M. van de Meent, and J. Vines, Energetics of two-body Hamiltonians in post-Minkowskian gravity, *Phys. Rev. D* **99**, 104004 (2019).
- [49] P. H. Damgaard and P. Vanhove, Remodeling the effective one-body formalism in post-Minkowskian gravity, *Phys. Rev. D* **104**, 104029 (2021).

- [50] J. Vines, D. Kunst, J. Steinhoff, and T. Hinderer, Canonical Hamiltonian for an extended test body in curved space-time: To quadratic order in spin, *Phys. Rev. D* **93**, 103008 (2016).
- [51] J. Vines, Scattering of two spinning black holes in post-Minkowskian gravity, to all orders in spin, and effective-one-body mappings, *Classical Quantum Gravity* **35**, 084002 (2018).
- [52] J. Vines, J. Steinhoff, and A. Buonanno, Spinning-black-hole scattering and the test-black-hole limit at second post-Minkowskian order, *Phys. Rev. D* **99**, 064054 (2019).
- [53] D. Bini and T. Damour, Gravitational spin-orbit coupling in binary systems, post-Minkowskian approximation and effective one-body theory, *Phys. Rev. D* **96**, 104038 (2017).
- [54] D. Bini and T. Damour, Gravitational spin-orbit coupling in binary systems at the second post-Minkowskian approximation, *Phys. Rev. D* **98**, 044036 (2018).
- [55] G. Mogull, J. Plefka, and J. Steinhoff, Classical black hole scattering from a worldline quantum field theory, *J. High Energy Phys.* **02** (2021) 048.
- [56] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, Classical Gravitational Bremsstrahlung from a Worldline Quantum Field Theory, *Phys. Rev. Lett.* **126**, 201103 (2021).
- [57] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, Gravitational Bremsstrahlung and Hidden Supersymmetry of Spinning Bodies, *Phys. Rev. Lett.* **128**, 011101 (2022).
- [58] G. U. Jakobsen, G. Mogull, J. Plefka, and J. Steinhoff, SUSY in the sky with gravitons, *J. High Energy Phys.* **01** (2022) 027.
- [59] C. Shi and J. Plefka, Classical double copy of worldline quantum field theory, *Phys. Rev. D* **105**, 026007 (2022).
- [60] F. Bastianelli, F. Comberiati, and L. de la Cruz, Light bending from eikonal in worldline quantum field theory, *J. High Energy Phys.* **02** (2022) 209.
- [61] G. U. Jakobsen and G. Mogull, Conservative and Radiative Dynamics of Spinning Bodies at Third Post-Minkowskian Order Using Worldline Quantum Field Theory, *Phys. Rev. Lett.* **128**, 141102 (2022).
- [62] T. Wang, Binary dynamics from worldline QFT for scalar-QED, *arXiv:2205.15753*.
- [63] G. U. Jakobsen, G. Mogull, J. Plefka, and B. Sauer, All things retarded: Radiation-reaction in worldline quantum field theory, *J. High Energy Phys.* **10** (2022) 128.
- [64] G. Kälin and R. A. Porto, Post-Minkowskian effective field theory for conservative binary dynamics, *J. High Energy Phys.* **11** (2020) 106.
- [65] G. Kälin, Z. Liu, and R. A. Porto, Conservative Dynamics of Binary Systems to Third Post-Minkowskian Order from the Effective Field Theory Approach, *Phys. Rev. Lett.* **125**, 261103 (2020).
- [66] G. Kälin, Z. Liu, and R. A. Porto, Conservative tidal effects in compact binary systems to next-to-leading post-Minkowskian order, *Phys. Rev. D* **102**, 124025 (2020).
- [67] G. Kälin, J. Neef, and R. A. Porto, Radiation-reaction in the effective field theory approach to post-Minkowskian dynamics, *arXiv:2207.00580*.
- [68] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, Dynamics of binary systems to fourth Post-Minkowskian order from the effective field theory approach, *Phys. Lett. B* **831**, 137203 (2022).
- [69] C. Dlapa, G. Kälin, Z. Liu, and R. A. Porto, Conservative Dynamics of Binary Systems at Fourth Post-Minkowskian Order in the Large-Eccentricity Expansion, *Phys. Rev. Lett.* **128**, 161104 (2022).
- [70] C. Dlapa, G. Kälin, Z. Liu, J. Neef, and R. A. Porto, Radiation reaction and gravitational waves at fourth post-Minkowskian order, *arXiv:2210.05541*.
- [71] M. Khalil, A. Buonanno, J. Steinhoff, and J. Vines, Energetics and scattering of gravitational two-body systems at fourth post-Minkowskian order, *Phys. Rev. D* **106**, 024042 (2022).
- [72] S. Mougiakakos, M. M. Riva, and F. Vernizzi, Gravitational bremsstrahlung in the post-Minkowskian effective field theory, *Phys. Rev. D* **104**, 024041 (2021).
- [73] M. M. Riva and F. Vernizzi, Radiated momentum in the post-Minkowskian worldline approach via reverse unitarity, *J. High Energy Phys.* **11** (2021) 228.
- [74] S. Mougiakakos, M. M. Riva, and F. Vernizzi, Gravitational Bremsstrahlung with Tidal Effects in the Post-Minkowskian Expansion, *Phys. Rev. Lett.* **129**, 121101 (2022).
- [75] M. M. Riva, F. Vernizzi, and L. K. Wong, Gravitational bremsstrahlung from spinning binaries in the post-Minkowskian expansion, *Phys. Rev. D* **106**, 044013 (2022).
- [76] J. S. Schwinger, Brownian motion of a quantum oscillator, *J. Math. Phys. (N.Y.)* **2**, 407 (1961).
- [77] L. V. Keldysh, Diagram technique for nonequilibrium processes, *Zh. Eksp. Teor. Fiz.* **47**, 1515 (1964).
- [78] R. D. Jordan, Effective field equations for expectation values, *Phys. Rev. D* **33**, 444 (1986).
- [79] S. Weinberg, Quantum contributions to cosmological correlations, *Phys. Rev. D* **72**, 043514 (2005).
- [80] C. R. Galley and M. Tiglio, Radiation reaction and gravitational waves in the effective field theory approach, *Phys. Rev. D* **79**, 124027 (2009).
- [81] D. Bini and T. Damour, Gravitational radiation reaction along general orbits in the effective one-body formalism, *Phys. Rev. D* **86**, 124012 (2012).
- [82] T. Damour, Radiative contribution to classical gravitational scattering at the third order in G , *Phys. Rev. D* **102**, 124008 (2020).
- [83] D. Bini, T. Damour, and A. Geralico, Radiative contributions to gravitational scattering, *Phys. Rev. D* **104**, 084031 (2021).
- [84] N. J. Bjerrum-Bohr, P. H. Damgaard, G. Festuccia, L. Planté, and P. Vanhove, General Relativity from Scattering Amplitudes, *Phys. Rev. Lett.* **121**, 171601 (2018).
- [85] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Planté, and P. Vanhove, The SAGEX review on scattering amplitudes, Chapter 13: Post-Minkowskian expansion from scattering amplitudes, *J. Phys. A* **55**, 443014 (2022).
- [86] D. A. Kosower, R. Monteiro, and D. O'Connell, The SAGEX review on scattering amplitudes, Chapter 14: Classical gravity from scattering amplitudes, *J. Phys. A* **55**, 443015 (2022).
- [87] A. Buonanno, M. Khalil, D. O'Connell, R. Roiban, M. P. Solon, and M. Zeng, Snowmass white paper: Gravitational waves and scattering amplitudes, *arXiv:2204.05194*.

- [88] L. J. Dixon, Calculating scattering amplitudes efficiently, [arXiv:hep-ph/9601359](#).
- [89] H. Elvang and Y.-t. Huang, Scattering amplitudes, [arXiv:1308.1697](#).
- [90] J. M. Henn and J. C. Plefka, *Scattering Amplitudes in Gauge Theories* (Springer, Berlin, 2014), Vol. 883, [10.1007/978-3-642-54022-6](#).
- [91] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson, and R. Roiban, The duality between color and kinematics and its applications, [arXiv:1909.01358](#).
- [92] G. Travaglini *et al.*, The SAGEX review on scattering amplitudes, *J. Phys. A* **55**, 443001 (2022).
- [93] S. Weinzierl, Feynman integrals, [arXiv:2201.03593](#).
- [94] J. Blümlein and C. Schneider, The SAGEX review on scattering amplitudes, Chapter 4: Multi-loop Feynman integrals, *J. Phys. A* **55**, 443005 (2022).
- [95] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes and the Conservative Hamiltonian for Binary Systems at Third Post-Minkowskian Order, *Phys. Rev. Lett.* **122**, 201603 (2019).
- [96] Z. Bern, C. Cheung, R. Roiban, C.-H. Shen, M. P. Solon, and M. Zeng, Black hole binary dynamics from the double copy and effective theory, *J. High Energy Phys.* **10** (2019) 206.
- [97] Z. Bern, H. Ita, J. Parra-Martinez, and M. S. Ruf, Universality in the Classical Limit of Massless Gravitational Scattering, *Phys. Rev. Lett.* **125**, 031601 (2020).
- [98] C. Cheung and M. P. Solon, Classical gravitational scattering at $\mathcal{O}(G^3)$ from Feynman diagrams, *J. High Energy Phys.* **06** (2020) 144.
- [99] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, Universality of ultra-relativistic gravitational scattering, *Phys. Lett. B* **811**, 135924 (2020).
- [100] C. Cheung and M. P. Solon, Tidal Effects in the Post-Minkowskian Expansion, *Phys. Rev. Lett.* **125**, 191601 (2020).
- [101] Z. Bern, J. Parra-Martinez, R. Roiban, E. Sawyer, and C.-H. Shen, Leading nonlinear tidal effects and scattering amplitudes, *J. High Energy Phys.* **05** (2021) 188.
- [102] M. Accattulli Huber, A. Brandhuber, S. De Angelis, and G. Travaglini, From amplitudes to gravitational radiation with cubic interactions and tidal effects, *Phys. Rev. D* **103**, 045015 (2021).
- [103] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes and Conservative Binary Dynamics at $\mathcal{O}(G^4)$, *Phys. Rev. Lett.* **126**, 171601 (2021).
- [104] Z. Bern, J. Parra-Martinez, R. Roiban, M. S. Ruf, C.-H. Shen, M. P. Solon, and M. Zeng, Scattering Amplitudes, the Tail Effect, and Conservative Binary Dynamics at $\mathcal{O}(G^4)$, *Phys. Rev. Lett.* **128**, 161103 (2022).
- [105] C. R. T. Jones and M. Solon, Scattering amplitudes and N-body post-Minkowskian Hamiltonians in general relativity and beyond, [arXiv:2208.02281](#).
- [106] E. Herrmann, J. Parra-Martinez, M. S. Ruf, and M. Zeng, Gravitational Bremsstrahlung from Reverse Unitarity, *Phys. Rev. Lett.* **126**, 201602 (2021).
- [107] E. Herrmann, J. Parra-Martinez, M. S. Ruf, and M. Zeng, Radiative classical gravitational observables at $\mathcal{O}(G^3)$ from scattering amplitudes, *J. High Energy Phys.* **10** (2021) 148.
- [108] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, Radiation reaction from soft theorems, *Phys. Lett. B* **818**, 136379 (2021).
- [109] P. Di Vecchia, C. Heissenberg, R. Russo, and G. Veneziano, The eikonal approach to gravitational scattering and radiation at $\mathcal{O}(G^3)$, *J. High Energy Phys.* **07** (2021) 169.
- [110] C. Heissenberg, Infrared divergences and the eikonal exponentiation, *Phys. Rev. D* **104**, 046016 (2021).
- [111] N. E. J. Bjerrum-Bohr, P. H. Damgaard, L. Planté, and P. Vanhove, The amplitude for classical gravitational scattering at third post-Minkowskian order, *J. High Energy Phys.* **08** (2021) 172.
- [112] P. H. Damgaard, L. Plante, and P. Vanhove, On an exponential representation of the gravitational S-matrix, *J. High Energy Phys.* **11** (2021) 213.
- [113] D. A. Kosower, B. Maybee, and D. O’Connell, Amplitudes, observables, and classical scattering, *J. High Energy Phys.* **02** (2019) 137.
- [114] B. Maybee, D. O’Connell, and J. Vines, Observables and amplitudes for spinning particles and black holes, *J. High Energy Phys.* **12** (2019) 156.
- [115] A. Cristofoli, R. Gonzo, D. A. Kosower, and D. O’Connell, Waveforms from amplitudes, *Phys. Rev. D* **106**, 056007 (2022).
- [116] A. Cristofoli, R. Gonzo, N. Moynihan, D. O’Connell, A. Ross, M. Sergola *et al.*, The uncertainty principle and classical amplitudes, [arXiv:2112.07556](#).
- [117] A. Guevara, A. Ochirov, and J. Vines, Scattering of spinning black holes from exponentiated soft factors, *J. High Energy Phys.* **09** (2019) 056.
- [118] Y. F. Bautista and A. Guevara, From scattering amplitudes to classical physics: Universality, double copy and soft theorems.
- [119] A. Guevara, A. Ochirov, and J. Vines, Black-hole scattering with general spin directions from minimal-coupling amplitudes, *Phys. Rev. D* **100**, 104024 (2019).
- [120] N. Arkani-Hamed, Y.-t. Huang, and D. O’Connell, Kerr black holes as elementary particles, *J. High Energy Phys.* **01** (2020) 046.
- [121] A. Guevara, B. Maybee, A. Ochirov, D. O’Connell, and J. Vines, A worldsheet for Kerr, *J. High Energy Phys.* **03** (2021) 201.
- [122] Y. F. Bautista, A. Guevara, C. Kavanagh, and J. Vines, From scattering in black hole backgrounds to higher-spin amplitudes: Part I, [arXiv:2107.10179](#).
- [123] M. Chiodaroli, H. Johansson, and P. Pichini, Compton black-hole scattering for $s \leq 5/2$, *J. High Energy Phys.* **02** (2022) 156.
- [124] R. Aoude and A. Ochirov, Classical observables from coherent-spin amplitudes, *J. High Energy Phys.* **10** (2021) 008.
- [125] Y. F. Bautista and N. Siemonsen, Post-Newtonian waveforms from spinning scattering amplitudes, *J. High Energy Phys.* **01** (2022) 006.
- [126] L. Cangemi and P. Pichini, Classical limit of higher-spin string amplitudes, [arXiv:2207.03947](#).

- [127] Z. Bern, A. Luna, R. Roiban, C.-H. Shen, and M. Zeng, Spinning black hole binary dynamics, scattering amplitudes, and effective field theory, *Phys. Rev. D* **104**, 065014 (2021).
- [128] D. Kosmopoulos and A. Luna, Quadratic-in-spin Hamiltonian at $\mathcal{O}(G^2)$ from scattering amplitudes, *J. High Energy Phys.* **07** (2021) 037.
- [129] W.-M. Chen, M.-Z. Chung, Y.-t. Huang, and J.-W. Kim, The 2PM Hamiltonian for binary Kerr to quartic in spin, *J. High Energy Phys.* **08** (2022) 148.
- [130] Z. Bern, D. Kosmopoulos, A. Luna, R. Roiban, and F. Teng, Binary dynamics through the fifth power of spin at $\mathcal{O}(G^2)$, [arXiv:2203.06202](https://arxiv.org/abs/2203.06202).
- [131] R. Aoude, K. Haddad, and A. Helset, Searching for Kerr in the 2PM amplitude, *J. High Energy Phys.* **07** (2022) 072.
- [132] R. Aoude, K. Haddad, and A. Helset, Classical Gravitational Spinning-Spinless Scattering at $\mathcal{O}(G^2 S^\infty)$, *Phys. Rev. Lett.* **129**, 141102 (2022).
- [133] P. H. Damgaard, K. Haddad, and A. Helset, Heavy black hole effective theory, *J. High Energy Phys.* **11** (2019) 070.
- [134] R. Aoude, K. Haddad, and A. Helset, On-shell heavy particle effective theories, *J. High Energy Phys.* **05** (2020) 051.
- [135] A. Brandhuber, G. Chen, G. Travaglini, and C. Wen, A new gauge-invariant double copy for heavy-mass effective theory, *J. High Energy Phys.* **07** (2021) 047.
- [136] A. Brandhuber, G. Chen, G. Travaglini, and C. Wen, Classical gravitational scattering from a gauge-invariant double copy, *J. High Energy Phys.* **10** (2021) 118.
- [137] R. Aoude, K. Haddad, and A. Helset, Tidal effects for spinning particles, *J. High Energy Phys.* **03** (2021) 097.
- [138] K. Haddad, Exponentiation of the leading eikonal phase with spin, *Phys. Rev. D* **105**, 026004 (2022).
- [139] F. Febres Cordero, M. Kraus, G. Lin, M. S. Ruf, and M. Zeng, Conservative binary dynamics with a spinning black hole at $\mathcal{O}(G^3)$ from scattering amplitudes, [arXiv:2205.07357](https://arxiv.org/abs/2205.07357).
- [140] A. V. Manohar, A. K. Ridgway, and C.-H. Shen, Radiated Angular Momentum and Dissipative Effects in Classical Scattering, *Phys. Rev. Lett.* **129**, 121601 (2022).
- [141] E. Barausse, E. Racine, and A. Buonanno, Hamiltonian of a spinning test-particle in curved spacetime, *Phys. Rev. D* **80**, 104025 (2009).
- [142] M. H. L. Pryce, Commuting co-ordinates in the new field theory, *Proc. R. Soc. A* **150**, 166 (1935).
- [143] M. H. L. Pryce, The mass center in the restricted theory of relativity and its connection with the quantum theory of elementary particles, *Proc. R. Soc. A* **195**, 62 (1948).
- [144] T. D. Newton and E. P. Wigner, Localized states for elementary systems, *Rev. Mod. Phys.* **21**, 400 (1949).
- [145] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevD.107.044033> for full expressions of the results presented in this paper.
- [146] F. Alessio and P. Di Vecchia, Radiation reaction for spinning black-hole scattering, *Phys. Lett. B* **832**, 137258 (2022).
- [147] P. H. Damgaard, J. Hoogeveen, A. Luna, and J. Vines, Scattering angles in Kerr metrics, *Phys. Rev. D* **106**, 124030 (2022).
- [148] A. Antonelli, C. Kavanagh, M. Khalil, J. Steinhoff, and J. Vines, Gravitational spin-orbit and aligned spin₁-spin₂ couplings through third-subleading post-Newtonian orders, *Phys. Rev. D* **102**, 124024 (2020).
- [149] Z. Liu, R. A. Porto, and Z. Yang, Spin effects in the effective field theory approach to post-Minkowskian conservative dynamics, *J. High Energy Phys.* **06** (2021) 012.
- [150] M. Boyle *et al.*, The SXS Collaboration catalog of binary black hole simulations, *Classical Quantum Gravity* **36**, 195006 (2019).
- [151] S. Ossokine, T. Dietrich, E. Foley, R. Katebi, and G. Lovelace, Assessing the energetics of spinning binary black hole systems, *Phys. Rev. D* **98**, 104057 (2018).