

**Entanglement wedges for gravitating regions**

Raphael Bousso\* and Geoff Penington†

*Center for Theoretical Physics and Department of Physics,  
University of California, Berkeley, California 94720, USA* (Received 21 December 2022; accepted 7 March 2023; published 4 April 2023)

Motivated by properties of tensor networks, we conjecture that an arbitrary gravitating region  $a$  can be assigned a generalized entanglement wedge  $E \supset a$ , such that quasilocal operators in  $E$  have a holographic representation in the full algebra generated by quasilocal operators in  $a$ . The Universe need not be asymptotically flat or anti-de Sitter (AdS), and  $a$  need not be asymptotic or weakly gravitating. On a static Cauchy surface  $\Sigma$ , we propose that  $E$  is the superset of  $a$  that minimizes the generalized entropy. We prove that  $E$  satisfies a no-cloning theorem and appropriate forms of strong subadditivity and nesting. If  $a$  lies near a portion  $A$  of the conformal boundary of AdS space, our proposal reduces to the quantum minimal surface prescription applied to  $A$ . We also discuss possible covariant extensions of this proposal, although none prove completely satisfactory. Our results are consistent with the conjecture that information in  $E$  that is spacelike to  $a$  in the semiclassical description can nevertheless be recovered from  $a$ , by microscopic operators that break that description. We thus propose that  $E$  quantifies the range of holographic encoding, an important nonlocal feature of quantum gravity, in general spacetimes.

DOI: [10.1103/PhysRevD.107.086002](https://doi.org/10.1103/PhysRevD.107.086002)**I. INTRODUCTION**

We seek a theory of quantum gravity that can describe our own Universe. One possible approach to this challenge is to analyze the well-known theories [1] that instead describe asymptotically anti-de Sitter (AdS) space. In particular, one would like to identify elements of the AdS/CFT correspondence that have broader validity. The entanglement wedge prescription is a crucial part of the AdS/CFT dictionary: it identifies the spacetime region dual to a given conformal field theory (CFT) subregion. We will propose a more general prescription that associates an entanglement wedge to arbitrary regions in arbitrary spacetimes.

**A. Entanglement wedge of a boundary region**

The Ryu-Takayanagi (RT) proposal [2] was first introduced in the context of the AdS/CFT correspondence, as a method for computing the von Neumann entropy of a stationary CFT state reduced to any spatial region  $A$  of the boundary. The proposal was refined to allow for time-dependent states by Hubeny *et al.* [3]. Quantum corrections

were first included by Faulkner *et al.* [4]. Engelhardt and Wall [5] introduced the present, most powerful formulation of RT in terms of the quantum-extremal surface  $\text{QES}_{\min}(A)$  anchored on  $A$  with smallest generalized entropy among all such surfaces. This prescription can be applied in the semiclassical regime to any order in  $G\hbar$ .

The entanglement wedge  $\text{EW}(A)$  is defined as the homology wedge, i.e., the causal development of any spatial region bounded by  $\text{QES}_{\min}(A)$  and  $A$ . There is strong evidence that, in a large class of states,<sup>1</sup>  $\text{EW}(A)$  is precisely the bulk dual to the boundary region  $A$  [8,9].<sup>2</sup> That is, any local bulk operator in  $\text{EW}(A)$  can be implemented, or “reconstructed,” by a CFT operator in the algebra associated with the region  $A$ , and no bulk operator outside  $\text{EW}(A)$  can be so implemented. This result is known as entanglement wedge reconstruction.

Its definition does not require the entanglement wedge to be connected. Any component of  $\text{EW}(A)$  that is not connected to  $A$  is called an entanglement island [13,14]. An island arises naturally in the course of black hole evaporation, if the radiation is stored in a distant, localized reservoir  $a$  [15]. Consider a small boundary region  $A$  whose entanglement wedge contains  $a$ , as shown in Fig. 1. After the Page time, i.e., when the entropy of the radiation

\*bousso@berkeley.edu

†geoffp@berkeley.edu

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP<sup>3</sup>.

<sup>1</sup>See Refs. [6,7] for subtleties and appropriate refinements of the entanglement wedge. Our proposal admits analogous refinements, which may be important for its generalization to time-dependent settings; see Sec. IV C.

<sup>2</sup>See Refs. [10–12] for early work on this problem.




FIG. 1. Entanglement island. Hawking radiation emitted from a black hole (gray circle) is collected in the reservoir  $a$  (red) located near the boundary region  $A$ . Before the Page time, the (ordinary) entanglement wedge  $EW(A)$  consists only of the blue region adjacent to  $A$ . After the Page time,  $EW(A)$  also contains an island  $b$  in the black hole interior.

exceeds the Bekenstein-Hawking entropy of the remaining black hole,  $EW(A)$  will also include an island  $b$  just inside of the black hole horizon. The inclusion of  $b$  in  $EW(A)$  decreases the generalized entropy since the radiation entropy in  $a$  is purified by the Hawking partners in  $b$ , and the price paid is only the Bekenstein-Hawking entropy of the island boundary (roughly that of the black hole).

## B. Going beyond AdS space

With certain assumptions, the entanglement wedge prescription can be derived from the gravitational path integral [16]. This implies that the entanglement wedge prescription can be used to compute the entropy not only of subregions of the conformal boundary of AdS space, but of more general systems that are coupled to a gravitating spacetime, regardless of whether an exact description is known.<sup>3</sup>

Crucially, this suggests that the entanglement wedge prescription is not confined to the context of AdS/CFT. Here we propose that an entanglement wedge prescription should be applicable in general spacetimes. Like the generalized second law [18,19] and the covariant entropy bound [20,21], the entanglement wedge reveals aspects of the (usually unknown) full quantum gravity theory, extracted from the gravitational path integral.<sup>4</sup>

<sup>3</sup>The earliest example of this phenomenon was the observation that, if a nongravitating external system  $R$  purifies matter near the QES that bounds  $EW(A)$ , then access to  $R$  can increase the size of the reconstructible bulk region [6,17]. In other words,  $EW(A \cup R)$  can differ from  $EW(A)$ .

<sup>4</sup>When a full theory is available, it can be used to verify the predictions of the entanglement wedge prescription. For example, by the AdS/CFT duality, the Page curve for an evaporating AdS black hole follows independently from the scrambling properties and the unitarity of the CFT.




FIG. 2. Examples that motivate and constrain our definition of a generalized entanglement wedge, a map  $E: a \rightarrow E(a)$ , that takes bulk regions as its input. Left: Hawking radiation arrives on a portion  $A$  of future null infinity. After the Page time, the (original) entanglement wedge of  $A$  includes an island:  $EW(A) = A \cup b$ . When the radiation is still in the distant spacetime region  $a$  (green shaded), it should already possess a generalized entanglement wedge, such that  $E(a) \supset a \cup b$ . Right: the (original) entanglement wedge  $EW(A)$  of an AdS boundary region  $A$  can be regarded as the bulk algebra of operators encoded in the CFT algebra of operators in  $A$ . The latter is generated by local operators in  $A$ , which are dual to quasilocally bulk operators in the near-boundary region  $a$ . Hence  $a$  should possess a generalized entanglement wedge such that  $E(a) = EW(A)$ .

The full generality of the entanglement wedge prescription—its applicability beyond the AdS/CFT correspondence—was long obscured by the lack of nontrivial boundary-anchored extremal surfaces in non-AdS spacetimes or auxiliary systems. Entanglement islands, however, are detached from the conformal boundary by definition. Hence, islands furnish nontrivial examples of entanglement wedges of systems that have nothing to do with the conformal boundary of AdS space.

For example, if the Hawking radiation is transferred to an external system  $R$ , then after the Page time,  $R$  will possess a nontrivial entanglement wedge:  $EW(R) = R \cup b$ . (Islands were first discovered in this setting.) This should also be the case for Hawking radiation far from the black hole in asymptotically flat spacetimes (see Fig. 2, left panel). In both cases, the presence of an entanglement island  $b$  is crucial to deriving the Page curve and thus the unitarity of the Hawking process. And in neither case is the entanglement wedge computed for a portion of the conformal boundary of AdS space.

What should be regarded as the proper input for computing an entanglement wedge: a portion of the conformal boundary? Nongravitating systems outside the spacetime? In this paper, we will propose that the answer is neither: rather, the entanglement wedge prescription should be formulated as a map that takes any gravitating spacetime region as input and outputs another (equal or larger) bulk spacetime region. The entanglement wedges of nongravitating systems and of boundary regions should be interpreted as limits of this more general prescription that arise if

the input region is weakly gravitating or approaches the asymptotic boundary.

Because the real world has gravity, such a proposal might be testable at least in principle, unlike a boundary-to-bulk map. Importantly, it should extend the notion of an entanglement wedge to arbitrary spacetimes, including cosmology.

For static spacetimes, we propose that the generalized entanglement wedge  $E(a)$  is simply the superset of  $a$  with smallest generalized entropy.<sup>5</sup> We motivate this prescription using tensor network toy models of quantum gravity and show that it satisfies a number of desirable properties that suggest it correctly quantifies the range of holographic encoding.

For general regions in time-dependent spacetimes, the correct definition of the generalized entanglement wedge seems less clear. We discuss one possible definition in some detail and show that it satisfies some but not all of the properties satisfied by  $E$ .

### C. Outline

In Sec. II, we motivate our prescription. We argue in Sec. II A that it must be possible to assign a larger entanglement wedge at least to some gravitating regions. Universality then suggests that all gravitating regions have an entanglement wedge. In Sec. II B, we analyze the example of tensor network toy models and argue that a sensible entanglement wedge may be assigned to bulk regions.

In Sec. III, we define the entanglement wedge  $E(a)$  of a bulk region  $a$ . We focus on the “static” case, when  $a$  and  $E(a)$  are part of time-reversal invariant initial data. That is, we propose a generalization of the Ryu-Takayanagi prescription [2] including quantum corrections [4]. We establish some definitions and notation in Sec. III A. In Sec. III B, we define  $E(a)$  as the spatial region with smallest generalized entropy, among all regions that contain  $a$ . In Sec. III C, we show that  $E$  obeys nontrivial properties consistent with its interpretation as an entanglement wedge: no cloning, nesting, and strong subadditivity.

In Sec. IV, we discuss possible generalizations of our prescription to the time-dependent case, analogous to the Hubeny-Rangamani-Takayanagi prescription [3] and its quantum extension by Engelhardt and Wall [5]. After introducing relevant concepts in Sec. IV A, we consider  $E_n(a)$ , the wedge with the smallest generalized entropy among all quantum-normal wedges that contain  $a$  and share the conformal boundary (if any) of  $a$ . We show that  $E_n$  reduces to the entanglement wedge prescription for boundary regions and static regions. Moreover,  $E_n$  satisfies an appropriate no-cloning theorem. However,  $E_n$  fails to satisfy strong subadditivity and nesting. We briefly discuss some alternative proposals.

<sup>5</sup>Throughout, the symbols  $\subset$  and  $\supset$  are understood to allow set equality.



FIG. 3. Left: two tensors  $T_1$  and  $T_2$  in a tensor network. The in-plane legs (horizontal) connect different tensors together, while the bulk legs (vertical) input matter degrees of freedom. Right: by bending the in-plane leg connecting  $T_1$  and  $T_2$  around, we can reinterpret it as two maximally entangled bulk legs, one on each tensor.

### D. Discussion

By analogy with the interpretation of the usual entanglement wedge, our results suggest that information in  $E(a)$  can be manipulated or summoned by a bulk observer in  $a$ . This is a striking implication.

We expect that such operations would not admit an interpretation in terms of a continuous, classical spacetime. Indeed, even simple operations in AdS space, such as the instantaneous, local addition of a particle deep in the bulk, would violate the Bianchi identity and hence cannot be represented as a continuous geometry without introducing time folds. Yet, this operation can be implemented instantaneously on a global boundary slice  $\sigma$ . Continuity of the boundary manifold across  $\sigma$  then suggests that the bulk exists as a semiclassical spacetime in the past and future of  $\sigma$ , but not spacelike to  $\sigma$ . We expect that the reconstruction, from a bulk region  $a$ , of operators in  $E(a) \cap a'$  involves comparably violent breakdowns of the spacetime geometry, at least outside of  $a$ .

We stress that the full physical significance of our proposal is not yet clear. In particular, we are not proposing a specific reconstruction map for operators in  $E(a) \cap a'$  or any other generalization of the known entries of the holographic dictionary beyond the entanglement wedge prescription. But given its nontrivial properties, we expect  $E$  to play a role in formulating quantum gravity as a holographic duality beyond AdS/CFT, for arbitrary spacetimes.

### E. Relation to other work

In Ref. [22], a definition was given for entanglement islands of *low-energy* bulk fields in a weakly gravitating region. The fields have an explicit momentum cutoff, and higher-energy operators, such as those that would create black holes, are explicitly excluded. Hence, when applied to an asymptotic bulk region, the prescription [22] would not give the entanglement wedge of the corresponding boundary region. Thus, while we believe that Ref. [22] correctly treats the problem formulated there, its construction cannot be the right answer to the question we pose.

In Ref. [23], a restricted maximin prescription was proposed for the entanglement wedge of a boundary region in a *cutoff* spacetime. Again, this prescription is intended to answer a different question than the one we are interested in here. As we describe in Sec. IV C, a closely related

prescription—with the edge of  $a$  playing the role of the cutoff surface—yields a time-dependent generalization of our proposal that obeys strong subadditivity, but which violates the nesting property required for the entanglement wedges of bulk regions.

Finally, the present work does not support claims that there is no Page curve for an evaporating black hole [24]. The Page curve is expected to describe the entropy of low-energy Hawking radiation modes, not all the information, in principle, accessible far from the black hole. For example, let  $a$  be the exterior of a large sphere in asymptotically flat space. Then the generalized entanglement wedge we define,  $E(a)$ , can include the entire interior enclosed by  $a$ . But this does not mean that the information in  $E(a)\setminus a$  can be accessed via low-energy operators in  $a$ .

## II. MOTIVATION

### A. Why a bulk region should have an entanglement wedge

We begin by arguing that at least *some* gravitating bulk regions constitute legitimate input to the entanglement wedge prescription. We will discuss the two examples shown in Fig. 2.

First, consider the Hawking radiation emitted by a black hole in asymptotically flat space, after the Page time. The radiation must contain the same information, whether it has been extracted into an auxiliary non-gravitating system, or arrived at a portion  $A$  of future null infinity, or is still traversing a distant, weakly gravitating region  $a$ . Operationally, the presence of an island  $b$  in the former cases indicates unitarity: sufficiently careful experiments would show that the ultimate state of the radiation is pure. But if such experiments could only succeed in the complete absence of gravity, they would fail in the real world, so the question of unitarity would be operationally meaningless.

These considerations make it implausible that the radiation has an entanglement island only if its self-gravity is completely turned off, but not if its self-gravity is arbitrarily small. It follows that the weakly gravitating region  $a$  must be assigned an entanglement wedge that includes the island  $b$ ,

$$E(\text{distant Hawking radiation } a) \supset a \cup b. \quad (2.1)$$

Let us pause for a moment to examine what is happening here. Careful measurements on one part of the bulk spacetime, the Hawking radiation, are allowing us to extract information that resides (from the semiclassical viewpoint) in a distant spacelike-separated region: the black hole interior. This is a striking example of a fundamental nonlocality that appears to be present in quantum gravity; sufficiently complex operators do not have to respect the semiclassical structure of spacetime.

Importantly, this fundamental nonlocality is not solely a feature of black hole physics; instead it is an essential aspect of holography. This brings us to our second example (Fig. 2, right panel).

In the AdS/CFT correspondence, local CFT operators in a conformal boundary subregion  $A$  are dual to (quasi)local bulk operators near  $A$  [25,26]. The notion of “near” can be made precise by defining a bulk region  $a$  as the union of the entanglement wedges of tiny boundary regions containing slightly smeared local boundary operators.<sup>6</sup> But local CFT operators generate the entire algebra of the CFT. It follows that the algebra generated by bulk operators in the near-boundary bulk region  $a$  encodes the *entire* entanglement wedge of  $A$ .<sup>7</sup> Information apparently located deep in the bulk must be secretly, and highly nonlocally, encoded in quasilocal bulk operators near the asymptotic boundary.<sup>8</sup>

Thus, the traditional entanglement wedge prescription for boundary subregions,  $EW(A)$ , can be reinterpreted as a statement about the algebra generated by quasilocal bulk operators in the asymptotic bulk region  $a$ , without any reference to a CFT dual. It follows that an asymptotic bulk region  $a$  in asymptotically AdS spacetime must possess an entanglement wedge—indeed, the same entanglement wedge that would have been assigned to its conformal boundary  $A$ ,

$$E(\text{asymptotic AdS bulk region } a) = EW(A). \quad (2.2)$$

If two such apparently dissimilar bulk regions—Hawking radiation and asymptotic regions in AdS—encode a larger entanglement wedge, then Occam’s razor suggests that

- (i) *any* bulk region should have an associated entanglement wedge, and
- (ii) *only* bulk regions possess an entanglement wedge. (Boundary regions and auxiliary systems can be viewed as idealized limits of bulk regions that are near asymptotic infinity or very weakly gravitating, respectively.)

<sup>6</sup>High-dimensional local operators in the CFT will not correspond to low-energy fields in the bulk effective field theory. Instead, they will generically create large black holes. However, these black holes will still be localized near the asymptotic boundary. See, e.g., the discussion in [27].

<sup>7</sup>In  $2+1$  boundary dimensions and higher, Wilson loops play an important role; these are nonlocal operators that (in non-Abelian gauge theories) cannot be written as a product of gauge-invariant local operators. However, any Wilson loop will be contained in a thin boundary strip with arbitrarily shallow entanglement wedge. Therefore, it should also be accessible as a bulk operator in the corresponding near-boundary region and should still be contained in the associated bulk algebra.

<sup>8</sup>In fact, this encoding is exponentially simpler than the encoding of the black hole interior in Hawking radiation [28,29].

## B. Motivation from tensor networks

Additional motivation for such a proposal comes from tensor networks, a useful toy model of static states in quantum gravity [30–33]. In these models, each tensor represents a patch of the gravitational spacetime, with the logarithm of the dimension of an edge capturing the area (in Planck units) of the surface connecting two neighboring patches. Bulk degrees of freedom within the patch are described by additional “out-of-plane” legs that feed into the network.

Importantly, the distinction between out-of-plane and in-plane tensor network legs is not precise. Mathematically, highly entangled Rindler modes on either side of a cut play an identical role in the tensor network as in-plane legs representing area, as shown in Fig. 3. The same effect is seen in gravity where the Rindler modes and geometrical area are equivalent under renormalization group flow. As we increase the UV cutoff on the bulk quantum field theory, the amount of Rindler entanglement across a surface increases; this increase is canceled in a computation of generalized entropy by the decrease in  $A/4G$  created by the renormalization of Newton’s constant.

If we extrapolate the UV cutoff all the way to the Planck scale, it is natural to conjecture that the area term, or analogously the in-plane tensor network legs, go away entirely, with the entanglement coming entirely from out-of-plane bulk legs. In such a limit, the tensor network consists purely of a set of projection operators that map the bulk state to the boundary state.

Why does this suggest that a region  $a$  encodes the larger region  $E(a)$ , as suggested in Fig. 4. If in-plane legs are really bulk Planck scale bulk legs, then an operator  $O$  can be reconstructed by a (potentially very-high-energy) operator within the region  $a$  if the action of  $O$  is the same as an operator  $\tilde{O}$  that acts on the out-of-plane legs within  $a$ , along with the in-plane legs at the boundary of  $a$ . After all, both sets of degrees of freedom should be, in principle, accessible from  $a$ , even if the Planckian degrees of freedom may not be described by an effective field theory.




FIG. 4. Left: in this example, the edge of  $a$  is a nonconvex surface on a static slice of Minkowski space, and the edge of  $E(a)$  is its convex hull. Right: a tensor network analog of the region  $a$ . For random tensor networks, the map from the boundary of  $E(a)$  (blue) to the boundary of  $a$  (green) will be an approximate isometry.

In a tensor network,  $E(a)$  corresponds to the smallest-entropy cut through the network such that  $a$  is contained entirely within the cut (see Fig. 4, right panel). Let us assume for simplicity that the dominant contribution to the entropy of any given cut comes from the maximally entangled in-plane legs. (In gravity, this corresponds to taking the semiclassical limit where the area term gives the dominant contribution to generalized entropy.) Now consider the map  $V$  induced by the tensor network from the boundary of  $E(a)$ , together with bulk legs in  $E(a)\setminus a$ , to the boundary of  $a$ . By definition, any intermediate cut through this subnetwork describing  $V$  has larger area (i.e., much larger bond dimension) than the input to  $V$ . For networks with sufficiently generic tensors, this ensures that the map  $V$  is an (approximate) isometry,<sup>9</sup> and hence that the operator  $\tilde{O} = V^\dagger O V$ , which acts on the in-plane legs at the boundary of  $a$ , successfully reconstructs any operator  $O$  acting in  $E(a)\setminus a$ .

## III. GENERALIZED ENTANGLEMENT WEDGES IN TIME-REFLECTION SYMMETRIC SLICES

### A. Preliminaries

*Definition 3.1.*—A spacetime  $M$  is time-reflection symmetric if there exists a  $\mathbb{Z}_2$  action on  $M$  that exchanges past and future timelike directions and preserves all points in a Cauchy surface  $\Sigma$ .

*Convention 3.2.*—In this section, we will only be concerned with the time-reflection symmetric Cauchy surface  $\Sigma$ , regarded as a manifold with Riemannian metric  $h$ . We assume that  $\Sigma$  is inextendible.

*Definition 3.3.*—We use  $\partial s$  to denote the boundary of a set  $s \subset \Sigma$  in the induced topology of  $\Sigma$ . Also  $\text{cl } s \equiv s \cup \partial s$  and  $\text{int } s = s \cap \text{cl } s^C$ , where the superscript  $C$  denotes the set complement in  $\Sigma$ .

*Definition 3.4.*—A wedge  $a$  is any open subset of  $\Sigma$  that is the interior of its closure:  $a = \text{int cl } a$ . (The name is slightly more natural in the more general time-dependent context; it is adopted for compatibility with the existing phrase entanglement wedge).

*Definition 3.5.*—The intersection of two wedges is a wedge; but the union, complement, and relative complement need not be. Given two wedges  $a$  and  $b$ , the wedge union, wedge complement, and wedge relative complement are wedges defined by

$$a \uplus b \equiv \text{int cl}(a \cup b), \quad (3.1)$$

$$a' \equiv \text{int } a^C, \quad (3.2)$$

$$a \setminus b \equiv a \cap b'. \quad (3.3)$$

<sup>9</sup>Concretely, this will be the case with very high probability if the tensors are drawn from a Haar random ensemble [32].

*Definition 3.6.*—The area of a wedge  $a$  is the geometric area of  $\partial a$  and is denoted  $\text{Area}(a)$ .

*Definition 3.7.*—The generalized entropy of a wedge  $a$  is

$$S_{\text{gen}}(a) \equiv \frac{\text{Area}(\partial a)}{4G\hbar} + S(a) + \dots \quad (3.4)$$

Here

$$S(a) = -\text{tr} \rho_a \log \rho_a \quad (3.5)$$

is the von Neumann entropy of

$$\rho_a = \text{tr}_{a'} \rho, \quad (3.6)$$

the density operator of the quantum fields restricted to  $a$ .

*Remark 3.8.*—The area in Eq. (3.4) can be thought of as a counterterm that cancels the leading divergence in the von Neumann entropy  $S$ . Additional counterterms for subleading divergences have been omitted. See the appendix of Ref. [21] for details.

*Definition 3.9.*—Let  $\tilde{\Sigma} \equiv (M, \tilde{h})$  denote the conformal completion of  $\Sigma$ , obtained by conformally mapping  $\Sigma$  to a subset of a compact set (for example, a sphere) and taking the closure of the image.  $\tilde{\Sigma}$  may be a manifold with boundary [34]. The boundary of the image  $\partial\tilde{\Sigma}$  is called the conformal boundary of  $\Sigma$ . We will not distinguish notationally between a set  $a \subset \Sigma$  and its image in  $\tilde{\Sigma}$ . However, such a distinction is crucial for boundaries of sets.

*Definition 3.10.*—The boundary of a wedge  $a \subset \tilde{\Sigma}$  will be denoted  $\delta a$ . As shown in Fig. 5, we define the conformal boundary of  $a$  as the set

$$\tilde{\delta} a \equiv \delta a \cap \partial\tilde{\Sigma}; \quad (3.7)$$




FIG. 5. The boundary  $\delta a \in \tilde{\Sigma}$  of a wedge  $a$  in the conformal completion  $\tilde{\Sigma}$  decomposes into the original boundary  $\delta a \in \Sigma$  and a conformal boundary  $\tilde{\delta} a \in \partial\tilde{\Sigma}$ .

thus,

$$\delta a = \partial a \sqcup \tilde{\delta} a. \quad (3.8)$$

## B. Definition and basic properties

*Definition 3.11.*—Given a wedge  $a \subset \Sigma$ , we define its generalized entanglement wedge  $E(a)$  as the wedge that satisfies

$$a \subset E(a) \subset \Sigma \quad \text{and} \quad \tilde{\delta} a = \tilde{\delta} E(a) \quad (3.9)$$

and which has the smallest generalized entropy among all such wedges. We assume without proof that  $E(a)$  exists, and for convenience we shall assume that it is unique.

*Lemma 3.12: Monotonicity.*—Let  $a$  and  $b$  be wedges with the same conformal boundary,  $\tilde{\delta} a = \tilde{\delta} b$ . Then

$$a \subset b \Rightarrow S_{\text{gen}}[E(a)] \leq S_{\text{gen}}[E(b)]. \quad (3.10)$$

That is, the generalized entropy of the entanglement wedge increases monotonically with the input wedge under inclusion, if the conformal boundary is held fixed.

*Proof.*— $E(b) \supset b \supset a \Rightarrow S_{\text{gen}}[E(b)] \geq S_{\text{gen}}[E(a)]$  by Definition 3.11. ■

*Definition 3.13: Entanglement wedge of a boundary region.*—Let  $B \subset \partial\tilde{\Sigma}$  be a subregion of the conformal boundary  $\partial\tilde{\Sigma}$ . The (ordinary) entanglement wedge  $\text{EW}(B) \subset \Sigma$  [2,4] is the wedge with conformal boundary  $B$  and smallest generalized entropy among all such sets.

*Lemma 3.14: EW as a special case of E.*—If the wedge  $a$  lies in the (ordinary) entanglement wedge of its conformal boundary,  $a \subset \text{EW}(\tilde{\delta} a)$ , then its generalized entanglement wedge is

$$E(a) = \text{EW}(\tilde{\delta} a). \quad (3.11)$$

*Proof.*—The result follows immediately from Definitions 3.11 and 3.13. ■

*Remark 3.15: Asymptotic bulk regions.*—Let  $a_n$  be an infinite sequence of wedges with  $a_{n+1} \subset a_n$ ,  $\tilde{\delta} a_{n+1} = \tilde{\delta} a_n$ , and  $\bigcap_n a_n = \emptyset$ . Then  $E(a_n) = \text{EW}(\tilde{\delta} a_n)$  for all sufficiently large  $n$ .

## C. No cloning, strong subadditivity, and nesting

*Theorem 3.16: No cloning.*—

$$a \cap E(b) = \emptyset, \quad b \cap E(a) = \emptyset. \quad (3.12)$$

Then

$$E(a) \cap E(b) = \emptyset. \quad (3.13)$$




FIG. 6. A configuration excluded by Theorem 3.16 (no cloning): if  $a \cap E(b) = \emptyset$  and  $b \cap E(a) = \emptyset$  then  $E(a) \cap E(b) = \emptyset$ , because otherwise removing the overlap from each entanglement wedge would fail to increase their combined generalized entropy.

*Proof.*—As shown in Fig. 6,

$$\begin{aligned} & \text{Area}[E(a) \setminus E(b)] + \text{Area}[E(b) \setminus E(a)] \\ & \leq \text{Area}[E(a)] + \text{Area}[E(b)]. \end{aligned} \quad (3.14)$$

[This need not be an equality; for example, there may be a disconnected shared boundary in  $E(a) \cap E(b)$ .] Strong subadditivity implies the same inequality for the von Neumann entropies; hence

$$\begin{aligned} & S_{\text{gen}}[E(a) \setminus E(b)] + S_{\text{gen}}[E(b) \setminus E(a)] \\ & \leq S_{\text{gen}}[E(a)] + S_{\text{gen}}[E(b)]. \end{aligned} \quad (3.15)$$

That is, the generalized entropy of  $E(a)$  or  $E(b)$  decreases when  $E(a) \cap E(b)$  is removed. By the assumption of the Lemma,  $a \subset E(a) \setminus E(b)$  and  $b \subset E(b) \setminus E(a)$ , so this contradicts Definition 3.11 unless  $E(a) \cap E(b) = \emptyset$ . ■

*Theorem 3.17: Strong subadditivity.*—

$$\begin{aligned} & S_{\text{gen}}[E(a \uplus b)] + S_{\text{gen}}[E(b \uplus c)] \\ & \geq S_{\text{gen}}[E(b)] + S_{\text{gen}}[E(a \uplus b \uplus c)]. \end{aligned} \quad (3.16)$$

*Proof.*—The setup is illustrated in Fig. 7. Rearranging components of surfaces, we have

$$\begin{aligned} & \text{Area}[E(a \uplus b)] + \text{Area}[E(b \uplus c)] \\ & \geq \text{Area}[E(a \uplus b) \cap E(b \uplus c)] + \text{Area}[E(a \uplus b) \uplus E(b \uplus c)]. \end{aligned} \quad (3.17)$$

(This need not be an equality since the wedge union can erase boundary portions.) Strong subadditivity implies the same inequality for the von Neumann entropies, and hence for the generalized entropy

$$\begin{aligned} & S_{\text{gen}}[E(a \uplus b)] + S_{\text{gen}}[E(b \uplus c)] \\ & \geq S_{\text{gen}}[E(a \uplus b) \cap E(b \uplus c)] + S_{\text{gen}}[E(a \uplus b) \uplus E(b \uplus c)]. \end{aligned} \quad (3.18)$$




FIG. 7. An illustration of the setup from Theorem 3.17 (strong subadditivity): The combined  $S_{\text{gen}}$  of  $E(a \uplus b)$  and  $E(b \uplus c)$  is at least big as the combined  $S_{\text{gen}}$  of their intersection and union. In turn, these upper bound the combined  $S_{\text{gen}}$  of  $E(b)$  and  $E(a \uplus b \uplus c)$ , respectively.

The first set on the right contains  $b$  and the second contains  $a \uplus b \uplus c$ , so Definition 3.11 implies Eq. (3.16). ■

*Theorem 3.18: Nesting.*—

$$a \subset b \Rightarrow E(a) \subset E(b). \quad (3.19)$$

*Proof.*—The setup is illustrated in Fig. 8. Since  $a \subset b$ , we have  $a \subset E(a) \cap E(b)$ , so by Definition 3.11,

$$S_{\text{gen}}[E(a)] \leq S_{\text{gen}}[E(a) \cap E(b)], \quad (3.20)$$

$$S_{\text{gen}}[E(b)] \leq S_{\text{gen}}[E(a) \uplus E(b)]. \quad (3.21)$$

But

$$\begin{aligned} & \text{Area}[E(b)] + \text{Area}[E(a)] \\ & \geq \text{Area}[E(a) \cap E(b)] + \text{Area}[E(a) \uplus E(b)]. \end{aligned} \quad (3.22)$$

(This need not be an inequality since the wedge union can erase boundary portions.) Strong subadditivity implies the same inequality for the von Neumann entropies, and hence for the generalized entropy




FIG. 8. A hypothetical counterexample to Theorem (3.19) (nesting): if  $a \subset b$  but  $E(a) \not\subset E(b)$ , we can replace the two entanglement wedges by  $E(a) \cap E(b)$  and  $E(a) \uplus E(b)$ , respectively, decreasing their combined area.

$$\begin{aligned}
 & S_{\text{gen}}[E(b)] + S_{\text{gen}}[E(a)] \\
 & \geq S_{\text{gen}}[E(a) \cap E(b)] + S_{\text{gen}}[E(a) \uplus E(b)]. \quad (3.23)
 \end{aligned}$$

To avoid a contradiction, all three of the above inequalities must be saturated. The assumed uniqueness of  $E(a)$  implies  $E(a) \cap E(b) = E(a)$ , and hence  $E(a) \subset E(b)$ . ■

#### IV. TIME DEPENDENCE

In this section, we discuss possible generalizations of our proposal to the case where  $a$  does not lie on a time-reflection symmetric Cauchy surface. We will sketch approaches, obtain some partial results, and outline specific challenges. We focus primarily on one possible generalization—the smallest-generalized-entropy quantum-normal wedge  $E_n(a)$ —which turns out to obey a no-cloning theorem, but not strong subadditivity or nesting. We briefly comment on other possible generalizations that preserve strong subadditivity and nesting, respectively, but do not find any single definition that preserves all three.

##### A. Preliminaries

We begin by introducing natural generalizations of the key objects introduced in Sec. III A. Let  $M$  be a globally hyperbolic Lorentzian spacetime with metric  $g$ . The chronological and causal future and past  $I^\pm$  and  $J^\pm$ , and the future and past domains of dependence and Cauchy horizons  $D^\pm$  and  $H^\pm$ , are defined as in Wald [34]. Given  $s \subset M$ , we use  $\partial s$  to denote the boundary of  $s$  in  $M$  and  $s'$  to denote the interior of the set of points that are spacelike related to all points in  $s$ , that is, points outside the causal future and past of  $s$ .

*Definition 4.1.*—A wedge is a set  $a \subset M$  that satisfies  $a = a''$  (see Fig. 9, left).

*Remark 4.2.*—It can be shown that the intersection of two wedges  $a, b$  is a wedge (see Fig. 9, right); and the complement wedge  $a'$  is a wedge,

$$(a \cap b)'' = a \cap b; \quad a''' = a'. \quad (4.1)$$




FIG. 9. Left: a spherical wedge  $a$ , its complement wedge  $a'$ , and their shared edge  $\delta a$  in Minkowski space. The wedge  $a$  is diamond shaped in this spacetime diagram; a Cauchy slice of  $a$  is shown in dark green. This wedge is “normal,” since outgoing orthogonal light rays expand. Right: the intersection of two wedges is itself a wedge, with an edge that decomposes as  $\delta(a \cap b) = cI[(\delta a \cap b) \sqcup (H^+(a) \cap H^-(b)) \sqcup \{a \leftrightarrow b\}]$ .




FIG. 10. The wedge union (thick purple diamond) of two wedges (blue diamonds) is defined as the wedge complement of the intersection of their wedge complements. Two examples are shown.

*Definition 4.3.*—Given two wedges  $a$  and  $b$ , we define the relative complement of  $b$  in  $a$  as  $a \cap b'$  and the wedge union as  $a \uplus b \equiv (a' \cap b')'$  (see Fig. 10); these objects are wedges by the above remark.

*Remark 4.4.*—The wedge union satisfies  $a \uplus b \supset a \cup b$ . It is minimal in the sense that any wedge that contains  $a \cup b$  must contain  $a \uplus b$ .

*Definition 4.5.*—The edge  $\delta a$  of a wedge  $a$  is defined by  $\delta a \equiv \partial a \cap \partial a'$ . Note that  $a$  is fully characterized by specifying  $\delta a$  and one spatial side of  $\delta a$ .

*Definition 4.6.*—The area and generalized entropy of a wedge  $a$  are defined as in the static case, Definitions 3.6 and 3.7, with  $\partial a$  replaced by  $\delta a$ . Note that neither depend on a choice of Cauchy slice of  $a$ .

*Definition 4.7.*—Given a wedge  $a$ , we distinguish between its edge  $\delta a$  in  $M$  and its edge  $\tilde{\delta a}$  as a subset of the conformal completion  $\tilde{M}$ . The latter can contain an additional piece, the conformal edge

$$\tilde{\delta a} \equiv \delta a \cap \partial \tilde{M}. \quad (4.2)$$

*Definition 4.8.*—Given a wedge  $a$  and a point  $p \in \delta a$ , the past (future) quantum expansion,  $\Theta^-(a, p)$  [ $\Theta^+(a, p)$ ] is the shape derivative of the generalized entropy under outward deformations of  $a$  along the past (future) null vector field orthogonal to  $\delta a$  at  $p$ . A precise definition can be given in terms of a functional derivative,

$$\Theta^\pm(a, p) \equiv 4G\hbar \frac{\delta S_{\text{gen}}[a(X^\pm(p))]}{\delta X^\pm(p)}. \quad (4.3)$$

Here  $X^\pm$  are null coordinates orthogonal to  $\delta a$ , and  $a(X^\pm(p))$  are wedges obtained by deforming  $\delta a$  along them. See Ref. [21] for further details.

*Remark 4.9.*—By Eq. (3.4), the quantum expansion can be decomposed into a classical and a quantum piece,

$$\Theta^\pm(a, p) = \theta(p) + 4G\hbar \frac{\delta S[a(X^\pm(p))]}{\delta X^\pm(p)}. \quad (4.4)$$

The first term is the classical expansion [34], which depends only on the shape of  $\delta a$  near  $p$ . The second term in Eq. (4.4) is nonlocal.

*Lemma 4.10.*—Let  $a \subset b$  be wedges whose edges coincide in an open neighborhood  $O$ . At any point  $p \in \delta a \cap O$ ,

$$\Theta^\pm(a, p) \geq \Theta^\pm(b, p). \quad (4.5)$$

*Proof.*—Since  $\delta a$  and  $\delta b$  coincide near  $p$ , their shape, and hence their classical expansion  $\theta$ , will agree at  $p$ . The Lemma then follows from strong subadditivity of the von Neumann entropy, applied to an infinitesimal deformation of  $a$ , and of  $b$ , at  $p$ ; see Fig. 3(a) in Ref. [21]. ■

*Definition 4.11.*—The wedge  $a$  is called quantum normal at  $p \in \delta a$  if  $\Theta^+(a, p) \geq 0$  and  $\Theta^-(a, p) \geq 0$ . Other combinations of signs correspond to quantum-antinormal ( $\leq, \leq$ ), quantum trapped ( $\leq, \geq$ ), quantum antitrapped ( $\geq, \leq$ ), and quantum extremal ( $=, =$ ). Marginal cases arise if one expansion vanishes at  $p$ . In relations that hold for all  $p \in \delta a$ , we drop the argument  $p$ .

*Conjecture 4.12: Quantum focusing conjecture.*—At all orders in  $G\hbar$  [21],

$$\frac{\delta\Theta(a, p)}{\delta X^\pm(\bar{p})} \leq 0. \quad (4.6)$$

*Remark 4.13.*—In the ‘‘off-diagonal’’ case  $p \neq \bar{p}$ , the quantum focusing conjecture (QFC) follows from Lemma 4.10. A quantum field theory limit of the QFC, the quantum null energy condition [21], was proven in Refs. [35,36]; see also Refs. [37–40]. The general diagonal case  $p = \bar{p}$  remains a conjecture.

*Lemma 4.14.*—The intersection of two quantum-normal wedges is quantum normal.

*Proof.*—Let  $a$  and  $b$  be quantum normal. We use the decomposition of  $\delta(a \cap b)$  described in Fig. 9. For  $p \in \delta a \cap b$  (or  $p \in a \cap \delta b$ ), the condition  $\Theta^\pm(a \cap b, p) \geq 0$  follows from the normalcy of  $a$  (or  $b$ ) at  $p$  and Lemma 4.10. For  $p \in H^\pm(a) \cap H^\mp(b)$ , let  $a_p \subset a$  and  $b_p \subset b$  be deformations of  $a$  and  $b$  along  $H^\pm(a)$  and  $H^\mp(b)$ , respectively, such that  $\delta a_p, \delta b_p$  coincide with  $H^\mp(a) \cap H^\pm(b)$  in a neighborhood of  $p$ . Since  $\Theta^\pm(a) \geq 0$  and  $\Theta^\mp(b) \geq 0$  in a neighborhood of the points lightlike separated from  $p$ , Conjecture 4.12 implies that  $\Theta^\pm(a_p, p) \geq 0$  and  $\Theta^\mp(b_p, p) \geq 0$ . Hence  $\Theta^\pm(a \cap b, p) \geq 0$  by Lemma 4.10. ■

## B. Smallest-generalized-entropy normal wedge

*Definition 4.15.*—Given a wedge  $a$ , let  $E_n(a)$  denote the quantum-normal wedge that contains  $a$ , has the same conformal boundary as  $a$ , and has the smallest generalized entropy among all such wedges. As in the static case (Definition 3.11), we assume without proof that this wedge exists, and we assume for convenience that it is unique.

*Lemma 4.16: Monotonicity.*—Let  $a$  and  $b$  be wedges with the same conformal boundary,  $\tilde{\delta}a = \tilde{\delta}b$ . Then

$$a \subset b \Rightarrow S_{\text{gen}}[E_n(a)] \leq S_{\text{gen}}[E_n(b)]. \quad (4.7)$$

*Proof.*—The proof is analogous to the static case, Lemma 3.12. ■

*Lemma 4.17: Quantum expansion.*—By definition,  $E_n$  is quantum normal. Marginal cases arise at points  $p \in \delta$  that do not lie on  $\delta a$ :

- (i)  $E_n(a)$  is marginally quantum antitrapped,  $\Theta^-[E_n(a), p] = 0$ , at  $p \in H^+(a')$ .
- (ii)  $E_n(a)$  is marginally quantum trapped,  $\Theta^+[E_n(a), p] = 0$ , at points  $p \in H^-(a')$ .
- (iii)  $E_n(a)$  is quantum extremal,  $\Theta^+[E_n(a), p] = \Theta^-[E_n(a); p] = 0$ , at points  $p \in a'$ .

*Proof.*—By quantum normalcy of  $E(a)$ ,  $S_{\text{gen}}$  decreases when  $\delta E(a)$  is deformed toward  $\delta a$  at  $p$ . Lemma 4.10 ensures quantum normalcy of  $\delta E(a)$  away from  $p$  under this inward deformation. This conflicts with the  $S_{\text{gen}}$ -minimization requirement in Definition 4.15, unless the quantum expansion at  $p$  in the other null direction immediately becomes negative under the deformation. ■

*Definition 4.18: Entanglement wedge of a boundary region.*—Let  $B \subset \partial\tilde{M}$  be a subregion of a Cauchy surface of the conformal boundary  $\partial\tilde{M}$ . The entanglement wedge of  $B$ ,  $\text{EW}(B) \subset M$ , is the smallest-generalized-entropy quantum-extremal wedge with conformal edge  $B$  [5].

*Lemma 4.19: Reduction of  $E_n$  to EW.*—Let  $a$  be contained in the ordinary quantum-entanglement wedge of its conformal edge  $\tilde{\delta}a$ , i.e.,  $a \subset \text{EW}(\tilde{\delta}a)$ . Then

$$E_n(a) = \text{EW}(\tilde{\delta}a). \quad (4.8)$$

*Proof.*—By the arguments in the proof of Lemma 4.17, the smallest- $S_{\text{gen}}$  quantum-normal wedge with conformal edge  $\tilde{\delta}a$  is necessarily extremal, and hence is given by  $\text{EW}(\tilde{\delta}a)$ . By assumption  $a \subset \text{EW}(\tilde{\delta}a)$ , so this continues to be true when we restrict to normal wedges containing  $a$ . ■

*Lemma 4.20: Reduction of  $E_n$  to  $E$ .*—Let  $\Sigma_0$  be a time-reflection symmetric Cauchy surface, and let  $a$  be a wedge with  $\delta a \subset \Sigma_0$ . Then  $E_n(a)$  reduces to the static entanglement wedge  $E(a)$ ,<sup>10</sup>

$$\delta E_n(a) = \partial E(a). \quad (4.9)$$

*Proof.*—Let  $T$  be the time-reflection operator. The set  $E(a) \cap T[E(a)]$  is a wedge by Remark 4.2, normal by Lemma 4.14, and it contains  $a$ . It is also time-reflection symmetric, so its edge lies on  $\Sigma_0$ . This conflicts with

<sup>10</sup>In Sec. III,  $\partial$  denotes the boundary in the topology of  $\Sigma_0$ , and we revert to this usage in this Lemma and its proof. Elsewhere in the present section,  $\partial$  denotes the boundary in  $M$ .

Conjecture 4.12 unless  $E(a) = T[E(a)]$ . Hence  $\delta E_n(a) \subset \Sigma_0$ . Moreover,  $\partial E(a \cap \Sigma_0)$  must be pointwise quantum normal or antinormal by time-reflection symmetry. If it were antinormal at some point  $p$ , we could decrease  $S_{\text{gen}}$  by deforming  $\partial E(a \cap \Sigma_0)$  outward on  $\Sigma_0$  at  $p$ , contradicting its definition. ■

*Theorem 4.21: No cloning for  $E_n$ .*—Let  $a, b$  be wedges that satisfy

$$a \subset E_n(b)', \quad b \subset E_n(a)'. \quad (4.10)$$

Then

$$E_n(a) \subset E_n(b)', \quad (4.11)$$

or, equivalently,  $E_n(b) \subset E_n(a)'$ .

*Proof.*—For arbitrary wedges  $c$  and  $d$ , let

$$c/d \equiv (c \cap d) \cup (c \cap d'). \quad (4.12)$$

Lemma 4.17 and Conjecture 4.12 can be shown to imply

$$\begin{aligned} S_{\text{gen}}[E_n(a)'/_{E_n(b)}] &\leq S_{\text{gen}}[E_n(a)] \quad \text{and} \\ S_{\text{gen}}[E_n(b)'/_{E_n(a)}] &\leq S_{\text{gen}}[E_n(b)]. \end{aligned} \quad (4.13)$$

We add these inequalities and use strong subadditivity of the von Neumann entropy to obtain

$$\begin{aligned} S_{\text{gen}}[E_n(a) \cap E_n(b)'] + S_{\text{gen}}[E_n(a)' \cap E_n(b)] \\ \leq S_{\text{gen}}[E_n(a)] + S_{\text{gen}}[E_n(b)]. \end{aligned} \quad (4.14)$$

We will now show that  $E_n(a) \cap E_n(b)'$  is quantum normal. Let us begin by substituting  $a \rightarrow E_n(a)$  and  $b \rightarrow E_n(b)'$  in the proof of Lemma 4.14. The Lemma assumed that  $a$  and  $b$  are normal, whereas now  $E_n(b)'$  is quantum antinormal. However, our assumption, Eq. (4.10), implies that  $\Theta^+[E_n(b)', p] = 0$  or  $\Theta^-[E_n(b)', p] = 0$  at any points  $p$  where these expansions actually appear in the proof. Indeed, since  $E_n(a)$  is spacelike to  $b$ ,  $\Theta^\pm[E_n(b)', p] = 0$  for  $p \in E_n(a) \cap \delta E_n(b)$  by Lemma 4.17. If  $p$  is the end point of a geodesic generator of  $H^-[E_n(b)']$  that intersects  $H^+[E_n(a)]$ , then  $\Theta^-[E_n(b)', p] = 0$  by Lemma 4.17, or else there would be a causal curve from  $\delta E_n(a)$  to  $b$ , in conflict with Eq. (4.10).

By the same reasoning,  $E_n(a)' \cap E_n(b)$  is also quantum normal. Moreover,  $a \subset E_n(a) \cap E_n(b)'$  and  $b \subset E_n(a)' \cap E_n(b)$  by Eq. (4.10). By Definition 4.15,  $E_n(a)$  and  $E_n(b)$  are the unique smallest- $S_{\text{gen}}$  quantum-normal wedges containing, respectively,  $a$  and  $b$ . Equation (4.14) thus implies  $E_n(a) \cap E_n(b)' = E_n(a)$ , which is equivalent to Eq. (4.11). ■




FIG. 11. Collapse of a null shell purified by a distant reference system  $r$ . The wedge  $a$  is marginally quantum trapped, so  $a = E_n(a)$ . One can arrange that  $S_{\text{gen}}[E_n(a)] + S_{\text{gen}}[E_n(r)] \gg S_{\text{gen}}[E_n(a \cup r)]$ . If  $r$  is the Python's lunch [29] (dashed red) of an asymptotic region  $b$ , one can arrange that  $E_n(a \cup b) \not\subset E_n(b)$ . Hence  $E_n$  satisfies neither subadditivity nor nesting.

### C. Discussion

$E_n$  does not satisfy subadditivity or nesting. This suggests that  $E_n$  is not quite the correct generalization of  $E$ . A counterexample to both properties is shown in Fig. 11. A purely classical counterexample to strong subadditivity is furnished by taking  $b$  to be a rectangle of length and width  $\Delta y \gg \tau \gg \Delta x \gg \ell_p$  at  $t = 0$  in two-dimensional Minkowski space and  $a$  and  $c$  to be identical rectangles obtained by moving  $b$  by  $1 + 2\Delta x$  in the  $\pm x$  direction and by  $\tau$  in the  $t$  direction.

Ideally, we would like to find a prescription  $e(a)$  that satisfies all of the following properties:

- (1)  $e$  reduces to the time-reflection symmetric prescription given in Sec. III:  $e(a) = E(a)$ .
- (2)  $e$  reduces to the QES prescription for the entanglement wedge of boundary regions:  $a \subset \text{EW}(\delta a) \Rightarrow e(a) = \text{EW}(\delta a)$ .
- (3)  $e$  is well defined.
- (4)  $e$  satisfies no cloning, strong subadditivity, and nesting, in the sense shown for  $E$  in Sec. III.

We have seen that  $E_n$  satisfies the first three requirements as well as no cloning. We are aware of alternate candidate prescriptions satisfying other subsets of these requirements, but of no *single* prescription that satisfies all. For example, one can ensure nesting in addition to the requirements satisfied by  $E_n$ , by taking the wedge union  $\cup_{b \subset a} E_n(b)$  over all subwedges  $b \subset a$ . However, this definition is somewhat artificial, and it still does not obey strong subadditivity.

Reference [23] proposed a restricted maximin prescription for the entanglement wedge of a cutoff boundary region. This prescription can be converted into a proposal for the generalized entanglement wedge of a bulk region, which (like  $E_n$ ) satisfies the first three requirements above. In addition, it obeys strong subadditivity; however, it

violates nesting.<sup>11</sup> Explicitly, let  $\delta a$  be the cutoff surface denoted  $\gamma$  in Ref. [23], and let  $a$  be its exterior, i.e., the portion discarded by the cutoff. Choosing  $A = \gamma$  in their notation, the prescription of Ref. [23] defines a wedge  $\mathcal{E}(a) \subset a'$  bounded by  $\delta a$  and its restricted maximin surface. The wedge union  $e(a) = a \cup \mathcal{E}(a)$  is then a candidate for the generalized entanglement wedge. The proof of strong subadditivity from Ref. [23] directly implies strong subadditivity for the generalized entanglement wedges, so long as  $e(a \cup b) \cap c = e(b \cup c) \cap a = \emptyset$ . (This ensures that the maximin surfaces lie in  $a' \cap b' \cap c'$ , the spacetime region that survives all three cutoffs. From an information-theoretic perspective, it is roughly analogous to the conditions in Theorem 3.16 that ensure that  $a$ ,  $b$ , and  $c$  are genuinely independent degrees of freedom.) However, the statement of nesting proved in Ref. [23], while appropriate in that context, is very different from the general statement  $e(a) \subset e(b)$  that must hold for  $e$  to be an entanglement wedge of a bulk region. In fact, it is easy to see that the latter statement fails, since  $e(a) = a$  for any region  $a$  with edge  $\delta a$  piecewise lightlike.

However, it may not be appropriate to seek a single definition of the entanglement wedge of bulk regions. In recent work, it was shown that the QES prescription for the entanglement wedge EW of boundary regions must be refined, by replacing the von Neumann entropy with certain

<sup>11</sup>We also expect no cloning to hold but we have not verified this.

one-shot entropies [7]. For static spacetimes, instead of a single entanglement wedge, the refined prescription defines *two* wedges, called the max-EW and the min-EW. They describe the bulk regions that are, respectively, fully encoded in, and partially influenced by, the boundary region. This refinement has been generalized to the full time-dependent setting [41]. For a boundary input region, the difference between its max- and min-EW is always a quantum effect. However, if one tries to generalize this prescription further to allow for bulk input regions, it appears that, in time-dependent settings, the max- and min-EW may differ even at the classical level. We are hopeful that this may resolve the difficulties that we encountered in this section when trying to define a single generalized entanglement wedge with all our desired properties. We leave the details of any such prescription to future work.

### ACKNOWLEDGMENTS

We thank Chris Akers, Ben Freivogel, Adam Levine, and Juan Maldacena for valuable discussions. This work was supported in part by the Berkeley Center for Theoretical Physics, by the Department of Energy, Office of Science, Office of High Energy Physics under QuantISED Award No. DE-SC0019380, and under Contract No. DE-AC02-05CH11231. R. B. was supported by the National Science Foundation under Grant No. 2112880. G. P. was supported by the Simons Foundation through the ‘‘It from Qubit’’ program; by AFOSR Grant No. FA9550-22-1-0098, and by an IBM Einstein Fellowship at the Institute for Advanced Study.

- 
- [1] J. M. Maldacena, The large  $N$  limit of superconformal field theories and supergravity, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
  - [2] S. Ryu and T. Takayanagi, Holographic Derivation of Entanglement Entropy from AdS/CFT, *Phys. Rev. Lett.* **96**, 181602 (2006).
  - [3] V. E. Hubeny, M. Rangamani, and T. Takayanagi, A covariant holographic entanglement entropy proposal, *J. High Energy Phys.* **07** (2007) 062.
  - [4] T. Faulkner, A. Lewkowycz, and J. Maldacena, Quantum corrections to holographic entanglement entropy, *J. High Energy Phys.* **11** (2013) 074.
  - [5] N. Engelhardt and A. C. Wall, Quantum extremal surfaces: Holographic entanglement entropy beyond the classical regime, *J. High Energy Phys.* **01** (2015) 073.
  - [6] P. Hayden and G. Penington, Learning the Alpha-bits of black holes, *J. High Energy Phys.* **12** (2019) 007.
  - [7] C. Akers and G. Penington, Leading order corrections to the quantum extremal surface prescription, *J. High Energy Phys.* **04** (2021) 062.
  - [8] A. C. Wall, Maximin surfaces, and the strong subadditivity of the covariant holographic entanglement entropy, *Classical Quantum Gravity* **31**, 225007 (2014).
  - [9] D. L. Jafferis, A. Lewkowycz, J. Maldacena, and S. J. Suh, Relative entropy equals bulk relative entropy, *J. High Energy Phys.* **06** (2016) 004.
  - [10] R. Bousso, S. Leichenauer, and V. Rosenhaus, Light-sheets and AdS/CFT, *Phys. Rev. D* **86**, 046009 (2012).
  - [11] V. E. Hubeny and M. Rangamani, Causal holographic information, *J. High Energy Phys.* **06** (2012) 114.
  - [12] B. Czech, J. L. Karczmarek, F. Nogueira, and M. Van Raamsdonk, The gravity dual of a density matrix, *Classical Quantum Gravity* **29**, 155009 (2012).
  - [13] G. Penington, Entanglement wedge reconstruction and the information paradox, *J. High Energy Phys.* **09** (2020) 002.
  - [14] A. Almheiri, N. Engelhardt, D. Marolf, and H. Maxfield, The entropy of bulk quantum fields and the entanglement wedge of an evaporating black hole, *J. High Energy Phys.* **12** (2019) 063.

- [15] R. Bousso and M. Tomašević, Unitarity from a smooth horizon?, *Phys. Rev. D* **102**, 106019 (2020).
- [16] A. Lewkowycz and J. Maldacena, Generalized gravitational entropy, *J. High Energy Phys.* **08** (2013) 090.
- [17] C. Akers, S. Leichenauer, and A. Levine, Large breakdowns of entanglement wedge reconstruction, *Phys. Rev. D* **100**, 126006 (2019).
- [18] J.D. Bekenstein, Black holes and the second law, *Lett. Nuovo Cimento* **4**, 737 (1972).
- [19] A.C. Wall, A proof of the generalized second law for rapidly changing fields and arbitrary horizon slices, *Phys. Rev. D* **85**, 104049 (2012).
- [20] R. Bousso, A covariant entropy conjecture, *J. High Energy Phys.* **07** (1999) 004.
- [21] R. Bousso, Z. Fisher, S. Leichenauer, and A.C. Wall, Quantum focusing conjecture, *Phys. Rev. D* **93**, 064044 (2016).
- [22] X. Dong, X.-L. Qi, Z. Shangnan, and Z. Yang, Effective entropy of quantum fields coupled with gravity, *J. High Energy Phys.* **10** (2020) 052.
- [23] B. Grado-White, D. Marolf, and S.J. Weinberg, Radial cutoffs and holographic entanglement, *J. High Energy Phys.* **01** (2021) 009.
- [24] S. Raju, Failure of the split property in gravity and the information paradox, *Classical Quantum Gravity* **39**, 064002 (2022).
- [25] T. Banks, M.R. Douglas, G.T. Horowitz, and E.J. Martinec, AdS dynamics from conformal field theory, [arXiv:hep-th/9808016](https://arxiv.org/abs/hep-th/9808016).
- [26] A. Hamilton, D.N. Kabat, G. Lifschytz, and D.A. Lowe, Holographic representation of local bulk operators, *Phys. Rev. D* **74**, 066009 (2006).
- [27] D. Harlow and H. Ooguri, Symmetries in quantum field theory and quantum gravity, *Commun. Math. Phys.* **383**, 1669 (2021).
- [28] D. Harlow and P. Hayden, Quantum computation vs. firewalls, *J. High Energy Phys.* **06** (2013) 085.
- [29] A. R. Brown, H. Gharibyan, G. Penington, and L. Susskind, The Python's Lunch: Geometric obstructions to decoding Hawking radiation, *J. High Energy Phys.* **08** (2020) 121.
- [30] B. Swingle, Entanglement renormalization and holography, *Phys. Rev. D* **86**, 065007 (2012).
- [31] F. Pastawski, B. Yoshida, D. Harlow, and J. Preskill, Holographic quantum error-correcting codes: Toy models for the bulk/boundary correspondence, *J. High Energy Phys.* **06** (2015) 149.
- [32] P. Hayden, S. Nezami, X.-L. Qi, N. Thomas, M. Walter, and Z. Yang, Holographic duality from random tensor networks, *J. High Energy Phys.* **11** (2016) 009.
- [33] N. Bao, G. Penington, J. Sorce, and A.C. Wall, Beyond toy models: Distilling tensor networks in full AdS/CFT, *J. High Energy Phys.* **11** (2019) 069.
- [34] R. M. Wald, *General Relativity* (Chicago University Press, Chicago, 1984).
- [35] R. Bousso, Z. Fisher, J. Koeller, S. Leichenauer, and A.C. Wall, Proof of the quantum null energy condition, *Phys. Rev. D* **93**, 024017 (2016).
- [36] S. Balakrishnan, T. Faulkner, Z. U. Khandker, and H. Wang, A general proof of the quantum null energy condition, *J. High Energy Phys.* **09** (2019) 020.
- [37] J. Koeller and S. Leichenauer, Holographic proof of the quantum null energy condition, *Phys. Rev. D* **94**, 024026 (2016).
- [38] A.C. Wall, Lower Bound on the Energy Density in Classical and Quantum Field Theories, *Phys. Rev. Lett.* **118**, 151601 (2017).
- [39] F. Ceyhan and T. Faulkner, Recovering the QNEC from the ANEC, *Commun. Math. Phys.* **377**, 999 (2020).
- [40] S. Balakrishnan, V. Chandrasekaran, T. Faulkner, A. Levine, and A. Shahbazi-Moghaddam, Entropy variations and light ray operators from replica defects, *J. High Energy Phys.* **09** (2022) 217.
- [41] C. Akers, A. Levine, G. Penington, and E. Wildenhain (to be published).