

Collider probe of heavy additional Higgs bosons solving the muon $g - 2$ and dark matter problems

Monika Blanke^{*} and Syuhei Iguro[†]

*Institute for Theoretical Particle Physics (TTP), Karlsruhe Institute of Technology (KIT),
Engesserstraße 7, 76131 Karlsruhe, Germany*

*and Institute for Astroparticle Physics (IAP), Karlsruhe Institute of Technology (KIT),
Hermann-von-Helmholtz-Platz 1, 76344 Eggenstein-Leopoldshafen, Germany*



(Received 2 November 2022; accepted 4 May 2023; published 16 May 2023)

We study the Large Hadron Collider (LHC) search potential of a $\mathbb{Z}_4$ -based two Higgs doublet model which can simultaneously explain the muon $g - 2$ anomaly and the observed dark matter. The neutral scalars in the second Higgs doublet couple to μ and τ and largely contribute to the muon anomalous magnetic moment through the one-loop diagram involving τ and scalars. An additional singlet scalar which is charged under the discrete symmetry can be a dark matter candidate. An upper limit on the scalar mass originates from the unitarity constraint, and the $\mu\tau$ flavor-violating nature of the scalars predicts nonstandard signatures at the LHC. However, the previously proposed $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ signal via the electroweak heavy neutral scalar pair production at the LHC loses sensitivity for increasing scalar mass. We revisit this model and investigate the LHC prospects for the single production of the $\mu\tau$ flavor-violating neutral scalar. It is shown that the single scalar process helps to extend the LHC reach to the 1 TeV mass regime of the scenario. The search potential at the high energy LHC is also discussed.

DOI: [10.1103/PhysRevD.107.095024](https://doi.org/10.1103/PhysRevD.107.095024)

I. INTRODUCTION

Most experimental results so far support the standard model (SM) of particle physics. However, the SM falls short of explaining dark matter, the baryon asymmetry of the universe, neutrino masses and so on. Each of these problems has many possible solutions, and thus more experimental hints are required to specify the correct new physics (NP) scenario. One of the most notorious and long-lived discrepancies between the SM prediction and the measurement exists in the muon anomalous magnetic moment (a_μ) [1–3]. The comparison of the SM prediction with the experimental value is given as

$$\Delta a_\mu = a_\mu^{\text{exp}} - a_\mu^{\text{SM}} = (2.51 \pm 0.59) \times 10^{-9}. \quad (1)$$

The SM prediction is taken from the theory white paper [1]¹ which is mainly based on the data-driven determination

of the hadronic vacuum-polarization contribution.² It is known that the discrepancy is of the same order as the electroweak contribution, i.e., a new $\mathcal{O}(100)$ GeV weakly coupled particle can explain the discrepancy. However, no signal of NP at this scale has been found at the Large Hadron Collider (LHC) so far. This fact implies that in order to explain the discrepancy in terms of NP, some enhancement mechanism in the NP contribution to $g - 2$ is necessary.³

A popular method to enhance the $g - 2$ contribution is the introduction of a new flavor-violating particle. The dipole operator underlying $g - 2$ requires a chirality flip, which corresponds to the muon mass within flavor-conserving scenarios. A one-loop contribution involving a $\mu\tau$ flavor-violating particle is instead enhanced by a factor of $m_\tau/m_\mu \simeq 17$ [32–60].⁴ This mechanism can lift the mass scale of the new particle by more than a factor of four.

^{*}monika.blanke@kit.edu

[†]igurosyuhei@gmail.com

¹See Refs. [4–23] for relevant original work.

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

²We note that the estimate based on the recent lattice simulation differs and is more consistent with the measured muon $g - 2$ [24–26]. Recent results from other lattice groups are converging toward the BMW result [24,27]. However, the lattice results are in tension with the low energy $\sigma(e^+e^- \rightarrow \text{hadrons})$ data [28–30], so that further clarification is needed. In this paper we consider the discrepancy as quoted in Eq. (1).

³See Ref. [31] for a recent review.

⁴Due to the loop function, scalar mediators receive a further enhancement.




FIG. 1. Representative Feynman diagrams that contribute to the $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ signal at the LHC. The left diagram displays the electroweak pair production channel. The middle and right diagrams correspond to the single production process where ϕ denotes A or H . In addition, there are also contributions obtained by exchanging μ and τ which are included in our numerical calculation.

However, lepton flavor-violating (LFV) interactions are stringently constrained and easily spoil the model if the particle also has lepton flavor-conserving couplings. Therefore one needs to ensure the absence of flavor-diagonal couplings for the τ mass enhanced muon $g-2$ solution to be viable.

This specific coupling alignment can be realized by a discrete $\mathbb{Z}_4$ flavor symmetry within the two Higgs doublet model (2HDM) [45].⁵ In this model the $g-2$ contribution is proportional to the $\mu\tau$ LFV coupling and the mass difference of the additional neutral scalars. Recently it was proposed that the singlet scalar extension of the model can explain the relic density of the dark matter (DM) through the thermal freeze-out mechanism [60]. The $\mathbb{Z}_4$ symmetry is then used both to stabilize the DM candidate and also to realize the flavor alignment.

Since the new scalars are quark-phobic within the $\mathbb{Z}_4$ -based model, their production cross section at the LHC is not large. However, the unique coupling structure predicts that the neutral scalars $\phi = A, H$ decay into $\mu^\pm\tau^\mp$. Previously we pointed out the smoking-gun signature of a $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ final state via electroweak scalar pair production (left of Fig. 1) with a special focus on the case where all Yukawa and scalar potential couplings are smaller than one [47]. We argued that the full Run 2 dataset can test the model up to 500 GeV scalar mass thanks to the very unique double $\mu\tau$ LFV resonance nature of the signal events. However, if we accept relatively large coupling of $\mathcal{O}(1)$, the model can still explain the discrepancy with 1 TeV scalars.

In this paper we revisit the model's collider prospects in the presence of larger couplings. The pair production cross section is governed only by the electroweak coupling and decreases rapidly when the scalars get heavier. We thus propose the single scalar production process (middle and right of Fig. 1) to assist to cover the heavier scalar scenario. To search for the heavy leptophilic bosons, it is known that the inclusion of photon-initiated processes is important [51]. We combine those processes and evaluate the search potential at the future LHC.

⁵Note that a $\mathbb{Z}_4$ -symmetric 2HDM always carries an accidental $U(1)$ symmetry [61], however further extensions of the scalar sector can break the latter symmetry [62].

TABLE I. Relevant field content and charge assignment of the model. The notation of SM gauge quantum numbers is given as $(SU(3)_C, SU(2)_L)_{U(1)_Y}$.

Field	H_1	H_2	(L_e, L_μ, L_τ)	(e_R, μ_R, τ_R)	S
SM gauge	$(1, 2)_{1/2}$	$(1, 2)_{1/2}$	$(1, 2)_{-1/2}$	$(1, 1)_{-1}$	$(1, 1)_0$
$\mathbb{Z}_4$	1	-1	$(1, i, -i)$	$(1, i, -i)$	i

The layout of the paper is given as follows. In Sec. II, we briefly introduce our setup of the 2HDM and review the muon $g-2$ explanation. There we determine how heavy the scalar can be and discuss relevant constraints. In Sec. III, we focus on the collider phenomenology and show the impact of the single scalar production process to evaluate the future LHC reach. Section IV is devoted to the summary and discussion.

II. MODEL, MUON $g-2$ AND DARK MATTER

We consider a two Higgs doublet model with an additional scalar singlet (S) and a discrete $\mathbb{Z}_4$ symmetry under which the Higgs and lepton fields transform as given in Table I. The gauge charge assignments of other SM fields, e.g., quarks, are the same as in the SM, and they trivially transform under $\mathbb{Z}_4$.⁶

We assume the $\mathbb{Z}_4$ symmetry to be unaffected by electroweak symmetry breaking, so that the two Higgs doublets $H_{1,2}$ are in the Higgs basis [63,64] in which only one Higgs doublet has a nonvanishing vacuum expectation value (vev) of $v \simeq 246$ GeV. In this basis, the two Higgs doublets can be decomposed as

$$H_1 = \begin{pmatrix} G^+ \\ \frac{v+h+iG}{\sqrt{2}} \end{pmatrix}, \quad H_2 = \begin{pmatrix} H^+ \\ \frac{H+iA}{\sqrt{2}} \end{pmatrix}, \quad (2)$$

where G^+ and G are the SM Nambu-Goldstone bosons, and H^+ and h are a charged Higgs boson and the discovered

⁶In order to obtain realistic neutrino masses and mixings the model needs to be extended. See Ref. [60] for details.

CP -even Higgs boson, respectively. H and A correspond to additional neutral scalars. Note that a nonzero vev of the singlet S would spontaneously break the $\mathbb{Z}_4$ symmetry. The presence of $\langle S \rangle \neq 0$ would not alter the phenomenology discussed in the present paper, hence we do not discuss this possibility further.

The scalar potential of our model is given by [60]

$$\begin{aligned} V = & M_1^2 H_1^\dagger H_1 + M_2^2 H_2^\dagger H_2 + \lambda_1 (H_1^\dagger H_1)^2 + \lambda_2 (H_2^\dagger H_2)^2 \\ & + \lambda_3 (H_1^\dagger H_1)(H_2^\dagger H_2) + \lambda_4 (H_1^\dagger H_2)(H_2^\dagger H_1) \\ & + \frac{\lambda_5}{2} [(H_1^\dagger H_2)^2 + \text{H.c.}] \\ & + M_S^2 |S|^2 + \lambda_S |S|^4 + [\lambda'_S S^4 + \text{H.c.}] + \lambda_{S1} (H_1^\dagger H_1) |S|^2 \\ & + \lambda_{S2} (H_2^\dagger H_2) |S|^2 + \kappa [(H_1^\dagger H_2) S^2 + \text{H.c.}]. \end{aligned} \quad (3)$$

Since the mass spectrum of the scalars is of crucial importance for the muon $g-2$ as well as the collider phenomenology, we explicitly show the mass relations:

$$\begin{aligned} m_h^2 &= \lambda_1 v^2, & m_A^2 &= M_2^2 + \frac{\lambda_3 + \lambda_4 - \lambda_5}{2} v^2, \\ m_{H^\pm}^2 &= m_A^2 + \lambda_5 v^2, & m_{H^\pm}^2 &= m_A^2 - \frac{\lambda_4 - \lambda_5}{2} v^2, \\ m_S^2 &= M_S^2 + \frac{\lambda_{S1}}{2} v^2. \end{aligned} \quad (4)$$

Note that due to the $\mathbb{Z}_4$ charge assignments, neither h nor S mix with the neutral components H , A of H_2 . For later convenience we define the mass difference of the heavy neutral scalars H , A as

$$\begin{aligned} \Delta_{H-A} &= m_H - m_A \\ &\simeq 50 \text{ GeV} \left(\frac{\lambda_5}{1.5} \right) \left(\frac{1800 \text{ GeV}}{m_H + m_A} \right). \end{aligned} \quad (5)$$

Note that $\lambda_5 \geq 0$ corresponds to $m_H \geq m_A$. The mass difference Δ_{H-A} decreases for heavier scalars, since it is proportional to the $SU(2)_L$ breaking.

Thanks to the $\mathbb{Z}_4$ symmetry the Yukawa sector of the model contains only two new parameters, significantly less than in the general 2HDM. Following the notation in Ref. [38], the Yukawa sector of the model based on the $\mathbb{Z}_4$ charge assignment, in addition to the SM part, is given as

$$-\mathcal{L}_Y = \rho_e^{\mu\tau} \bar{L}_\mu H_2 \tau_R + \rho_e^{\tau\mu} \bar{L}_\tau H_2 \mu_R + \text{H.c.}, \quad (6)$$

where $\rho_e^{\mu\tau}$ and $\rho_e^{\tau\mu}$ are free parameters. The scalar singlet S does not couple to the SM fermions.

In this model, a sizable contribution to Δa_μ is generated via the one-loop diagram mediated by the extra neutral




FIG. 2. The black contours show the value of $\rho_e^{\mu\tau} \rho_e^{\tau\mu}$ to explain the central value of α_μ in the m_A vs. λ_5 plane. The blue region is excluded by the lepton flavor universality of τ decays. The orange region corresponds to $\Gamma_\phi/m_\phi \geq 30\%$. The purple contours depict the cutoff scale of the model.

Higgs bosons H and A . The τ mass enhanced contribution⁷ is given as [38,39]

$$\begin{aligned} \Delta a_\mu &\simeq \frac{m_\mu m_\tau \rho_e^{\mu\tau} \rho_e^{\tau\mu}}{16\pi^2} \left(\frac{\ln \frac{m_H^2}{m_\tau^2} - \frac{3}{2}}{m_H^2} - \frac{\ln \frac{m_A^2}{m_\tau^2} - \frac{3}{2}}{m_A^2} \right) \\ &\simeq -2.5 \times 10^{-9} \left(\frac{\rho_e^{\mu\tau} \rho_e^{\tau\mu}}{1.0} \right) \left(\frac{\lambda_5}{1.0} \right) \left(\frac{700 \text{ GeV}}{m_A} \right)^4, \end{aligned} \quad (7)$$

where Eq. (5) is used to derive the second relation. Notice the dependence on the product of the two new Yukawa couplings, $\Delta a_\mu \propto \rho_e^{\mu\tau} \rho_e^{\tau\mu}$. For given masses $m_{H,A}$, the NP effect in a_μ is thus largest if both couplings $\rho_e^{\mu\tau}$ and $\rho_e^{\tau\mu}$ are large. As we are interested in the heaviest possible scenario, we thus set $|\rho_e^{\mu\tau}| = |\rho_e^{\tau\mu}|$ in the following.

Figure 2 shows the value of $\rho_e^{\mu\tau} \rho_e^{\tau\mu}$ required to explain the central value of the discrepancy with black contours. If we allow for large Yukawa couplings, heavy scalars of $\mathcal{O}(1)$ TeV can explain the muon $g-2$ discrepancy. Furthermore the product $\rho_e^{\mu\tau} \rho_e^{\tau\mu} \lambda_5$ must be negative to obtain a positive contribution to Δa_μ . In summary, we find that the parameters relevant for Δa_μ are $\rho_e^{\mu\tau} \rho_e^{\tau\mu}$, m_A and λ_5 . It is noted that the τ mass enhanced $g-2$ contribution picks up the $SU(2)_L$ -breaking effect and is proportional to m_A^{-4} .

The charged Higgs mass is set to $m_{H^\pm} = m_A$ ($m_{H^\pm} = m_H$) for $\lambda_5 \geq 0$ ($\lambda_5 \leq 0$) to respect the constraints from electroweak oblique parameters [65] and vacuum

⁷We also include non- m_τ enhanced terms of the neutral scalar loop diagram numerically, however their impact is small in our case. The $H^\pm$ -loop contribution does not have an m_τ enhancement and thus its numerical impact is also small.




FIG. 3. Left: fiducial cross section of $\mu^\pm \mu^\pm \tau_h^\mp \tau_h^\mp$ via electroweak HA production at a 14 TeV pp collider, as a function of m_A in GeV. The horizontal gray lines correspond to the sensitivities with different luminosities. Right: the collider sensitivities are overlaid on Fig. 2. The cyan region can be tested with 3 ab^{-1} of the electroweak pair production, while the magenta region can additionally be probed by including the single production channel. The blue and orange regions and purple contours are the same as in Fig. 2. $\Delta a_\mu = 2.51 \times 10^{-9}$ is fixed and $|\rho_e^{\mu\tau}| = |\rho_e^{\tau\mu}|$ is assumed in the figure.

stability [66,67]. The size of the couplings is bounded from above by the requirement that the theory remains perturbative, we hence choose $|\lambda_i| \leq 4\pi$ and $|\rho_e^{\mu\tau}| \leq \sqrt{4\pi}$ [45]. Furthermore, the perturbative unitarity condition for which we require tree-level unitarity of $2 \rightarrow 2$ processes is imposed [68–70]. Even if the couplings satisfy those constraints at the mass scale of the additional scalars, renormalization group (RG) running effects alter them and the theory would become nonperturbative at a high energy scale with $\mathcal{O}(1)$ couplings.⁸ To quantify this, we solve the coupled RG equations (RGEs) for the β functions of the SM third generation fermion Yukawa couplings, λ_i , $\rho_e^{\mu\tau}$ and $\rho_e^{\tau\mu}$, and then determine the cutoff scale, Λ_{LP} , where LP stands for Landau pole.⁹ These conditions imply the existence of an upper mass limit for the heavy scalars. In Fig. 2 dash-dotted, dashed and dotted lines in purple depict $\Lambda_{\text{LP}} = 30 \text{ TeV}$, 10 TeV and 5 TeV . It is observed that if we require the theory to be perturbative up to 30 (5) TeV, the upper limit on the scalar mass is given as

$$m_A \leq 1250(1650) \text{ GeV}. \quad (8)$$

⁸In order to keep the model under perturbative control, additional particles and their interactions at the scale Λ_{LP} would be necessary. Such $\mathcal{O}(10 \text{ TeV})$ new particles can in principle contribute to the muon $g-2$, however the effect is small since it scales as $1/m_{\Lambda_{\text{LP}}}^2$. Similarly, they are out of reach of direct LHC searches. Finally, the contribution of the new particles to the beta functions and their impact on the RGE evolution highly depends on the nature of the new particles, e.g. their coupling structure and spin.

⁹See Refs. [45,71] for the β functions. At the initial scale, we set $\lambda_2, \lambda_3 \ll 1$ to maximize the cutoff scale. It is noted that there are typos in the β functions of Ref. [45].

The large λ_5 case is constrained by the $2 \rightarrow 2$ unitarity bound while the small λ_5 region is disfavored by the unitarity constraint on the RGE-evolved Yukawa couplings.

The presence of a charged Higgs with a sizable product of the relevant Yukawa couplings can modify the decay rate of $\tau \rightarrow \mu\nu\bar{\nu}$. Following Ref. [44], lepton flavor universality in τ decays puts an upper limit on the interaction:

$$\left| \frac{\rho_e^{\mu\tau} \rho_e^{\tau\mu}}{1.9} \right| \left(\frac{700 \text{ GeV}}{m_{H^\pm}} \right)^2 \leq 1. \quad (9)$$

The corresponding exclusion is shown in blue in Fig. 2.¹⁰ Furthermore the one-loop corrections to Z and Higgs boson couplings to $\tau\bar{\tau}$ and $\mu\bar{\mu}$ are known to be less constraining.

Since we are interested in the collider sensitivity utilizing resonant production of the new scalars $\phi = A, H$, their width-to-mass ratio is important. This ratio is approximately given as $\Gamma_\phi/m_\phi \sim |\rho_e^{\mu\tau}|^2 \times 4\%$. We assume that the narrow width approximation is valid up to 30%. The parameter region which predicts $\Gamma_\phi/m_\phi \geq 30\%$ is shown in orange in Fig. 2. Depending on the desired accuracy for LHC cross-section predictions, a more conservative limit on the validity of the narrow width approximation may be in order. However, since the cross section is a steeply falling function of m_A , as seen in the left panels of Figs. 3 and 4, and we are interested in the collider reach in terms of the mass m_A , even a somewhat imprecise prediction of the cross section using the narrow width approximation will yield sufficiently precise results for the (HL/HE-)LHC reach.

¹⁰The Belle II experiment will improve the sensitivity, however, a quantitative evaluation is not available [72].




FIG. 4. Left: fiducial cross section of $\mu^\pm\mu^\mp\tau_h^\pm\tau_h^\mp$ via electroweak HA production as a function of m_A in GeV. The colored lines show the model prediction at a 27 TeV pp collider. The horizontal gray lines correspond to the sensitivities with different luminosities. Right: the cyan region shows the sensitivity of the HE-LHC based on HA production. The magenta region can additionally be probed by including single production. The gray region displays the sensitivity of the HL-LHC. See also the caption in Figs. 2 and 3.

As shown in Ref. [60] and mentioned in the introduction, the singlet scalar S is a viable dark matter candidate. In the early universe S is in thermal equilibrium with the SM fields through its interaction with the scalar doublets, see Eq. (3). Note that in Ref. [60] the coupling of S to the SM Higgs doublet, λ_{S1} , has been set to zero and S has no vacuum expectation value. S then couples only to the new heavy scalars $H^\pm$, H , A which serve as $\mu\tau$ -flavored mediators. It is then possible to explain the relic abundance with the thermal freeze-out mechanism. Thanks to the $\mathbb{Z}_4$ charge assignment, the singlet scalar does not couple to the nucleon at the tree level, evading the stringent constraints from direct detection experiments. Similarly, due to the absence of couplings to SM particles, S is not produced directly at colliders. However, one-loop Z penguin induced DM-nucleon scattering can constrain the model. The viable mass range for S is broad and the next generation experiments are important to probe the interesting parameter space. More details on the phenomenology of S and its dark matter interpretation can be found in [60].

We stress that while the explanation of the $g-2$ anomaly in terms of A , H contributions is independent of the introduction of S and its dark matter interpretation, the reverse statement is not true. On the one hand, dark matter stability is guaranteed by the same $\mathbb{Z}_4$ symmetry that is responsible for the lepton flavor-violating coupling structure of the heavy scalars. On the other hand, since the heavy scalars serve as mediators between the visible and dark sectors of the model, their mass and interaction strength with the SM leptons directly influences the dark matter phenomenology of the model.

In passing we note that charged scalar pair production is probed by the left-handed slepton search [73,74]. However, it is difficult to constrain $O(1)$ TeV $H^\pm$ with $\text{BR}(H^\pm \rightarrow \mu\nu) \simeq 0.5$ even at the HL-LHC.

III. LHC PROBE OF TeV SCALAR SCENARIO

In this section we investigate the LHC sensitivity to our model. Reference [47] pointed out the smoking-gun signature $\mu^\pm\mu^\mp\tau^\pm\tau^\mp$ via pair production of the heavy neutral scalars HA , each decaying into $\mu^\pm\tau^\mp$. This signature violates lepton flavor and is therefore free from irreducible SM backgrounds. Note that while the final state $\mu^\pm\mu^\mp\tau^\pm\tau^\mp$ is produced at comparable rates, the latter is lepton flavor-conserving and therefore subject to sizeable SM background from, e.g., ZZ production. The signature $\mu^\pm\mu^\mp\tau^\pm\tau^\mp$ is thus less suitable as a discovery channel—however in case of a $\mu^\pm\mu^\mp\tau^\pm\tau^\mp$ discovery it could potentially be used to validate the underlying model. In the present paper we focus on the discovery prospects of the lepton flavor-violating signature $\mu^\pm\mu^\mp\tau^\pm\tau^\mp$.

As a first step we extend the previous study [47] and evaluate the high luminosity (HL)-LHC reach of the model based on the electroweak pair production of the scalars. Reference [47] focused on the weakly-coupled scenario with $|\lambda_5|, |\rho_e^{\mu\tau}|, |\rho_e^{\tau\mu}| \leq 1$, and thus $m_A \leq 700$ GeV was considered. It was argued that the very distinctive $\mu^\pm\mu^\mp\tau^\pm\tau^\mp$ final state via decays of an electroweakly produced pair of neutral scalars is useful to test the model and it was shown that 139 fb^{-1} of the data can probe the scenario with $m_A \lesssim 500$ GeV. The electroweak production cross section depends only on the heavy scalar masses.

Throughout our analysis, we use MADGRAPH5_aMC@NLO [75] with NNPDF3.1luxQED [76] to calculate the signal cross sections. To account for the minimal kinematic cuts, $|p_T^\mu|, |p_T^\tau| \geq 20$ GeV, $|\eta_\mu|, |\eta_\tau| \leq 2.7$, and $\Delta R \geq 0.1$ are imposed for all pairs of charged leptons. We assume a hadronic τ -tagging efficiency of 70% [77] and the hadronic τ decay branching ratio of about 65% [65]. It is noted that an excellent τ charge reconstruction is reported in Ref. [78].

Our signal of same sign $\mu^\pm\mu^\pm$ and $\tau^\mp\tau^\mp$ pairs of which $\mu^\pm\tau^\mp$ forms a resonance is very distinctive. Specifically, due to the resonance structure, the final-state leptons are very energetic. Therefore we can safely assume the SM background (SMBG) to be negligible.¹¹ In this situation Poisson statistics is applicable and the sensitivity at 95% confidence level is given when $\simeq 3$ signal events are predicted, if no events are observed in the data. The additional decay channels $H \rightarrow W^\pm H^\mp$ and $A \rightarrow HZ$ open in the case of large mass differences.¹² However, such large mass splittings are difficult for $\mathcal{O}(1)$ TeV scalars. We numerically included this dilution effect.

Interpreting the DM candidate S as the relic of a thermal freeze-out [60] requires a nonvanishing portal interaction between the Higgs doublets and S in Eq. (3),

$$V \supset \kappa(H_1^\dagger H_2)S^2 + \text{H.c.} \quad (10)$$

This coupling induces the invisible scalar decay $\phi \rightarrow SS$, where $\phi = H, A$. In turn this would suppress $\text{BR}(\phi \rightarrow \mu\tau)$ and hence reduce the $\mu^\pm\mu^\pm\tau_h^\mp\tau_h^\mp$ signal number. However, $\text{BR}(\phi \rightarrow \mu\tau)$ is proportional to m_ϕ while $\text{BR}(\phi \rightarrow SS)$ is proportional to m_ϕ^{-1} because of the dimensionful coupling κv originating from the portal interaction. As a result $\text{BR}(\phi \rightarrow SS)$ is suppressed in the heavy scalar regime and $\text{BR}(\phi \rightarrow \mu\tau) \simeq 1$ holds well. Therefore the $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ mode remains a viable and important probe of the model also in the presence of the DM candidate S , underlining the broad applicability of this search channel. It is noted that the relic abundance requires κ of $\mathcal{O}(0.1)$ and hence its presence does not have a significant impact on the phenomenology discussed in the present paper.

The colored contours in Fig. 3 (left) show the fiducial $\mu^\pm\mu^\pm\tau_h^\mp\tau_h^\mp$ cross sections based on the electroweak HA production with $\sqrt{s} = 14$ TeV. The horizontal lines correspond to the sensitivity for integrated luminosities of 139 fb^{-1} , 1 ab^{-1} , and 3 ab^{-1} , and correspond to cross sections of 0.02 fb , 0.003 fb , and 0.001 fb , respectively. For these cross sections $\simeq 3$ signal events are expected at the respective integrated luminosities so that an exclusion at 95% confidence level can be obtained in the absence of signal.

Therefore, the HL-LHC data of 1 ab^{-1} and 3 ab^{-1} would be sensitive to mass scales of $m_A \simeq 800 \text{ GeV}$ and $m_A \simeq 960 \text{ GeV}$. It is worth mentioning that the pair production channel is powerful since once the neutral scalars are produced they dominantly decay into $\mu\tau$, as long as a sizable τ -mass enhanced contribution to a_μ is

¹¹Background events come from $pp \rightarrow ZZ \rightarrow 4\tau$. We confirmed that the contribution is smaller than $\mathcal{O}(10^{-5}) \text{ fb}$ and can safely be neglected in the resonant regime.

¹²Note that the decay $A \rightarrow hZ$ is forbidden by the $\mathbb{Z}_4$ symmetry.

postulated. Nevertheless there is a mass gap between the sensitivity and the theoretical upper limit of Eq. (8). The loss of sensitivity for larger m_A mainly comes from two factors: the contributing coupling constant is a weak gauge coupling which is independent of Δa_μ and the production cross section is suppressed by the heaviness of the pair-produced scalars.

One possible way to extend the LHC reach to our model is to include the single heavy scalar production channels corresponding to the middle and right diagrams of Fig. 1. Especially for $\mathcal{O}(1)$ TeV leptophilic particles the inclusion of the photon initiated process is important [51]. Again, following the procedure in Ref. [41] we assume the SMBG to be negligible. There are two terms in the single scalar production amplitude to generate $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ events. One is proportional to $(|\rho_e^{\mu\tau}|^2 + |\rho_e^{\tau\mu}|^2)$ which vanishes in the $m_A = m_H$ limit. On the other hand the term proportional to $\rho_e^{\mu\tau}\rho_e^{\tau\mu}$ does not disappear in this limit. Since the m_A^{-4} scaling in Eq. (7) requires a large product of the LFV Yukawa couplings and the first term is suppressed by the mass difference, the second contribution will be important for $\mathcal{O}(1)$ TeV scalars.

In Fig. 3 (right) we overlay the collider sensitivity in the m_A vs. λ_5 plane. The cyan region shows the sensitivity of the pair production channel. The sensitivity is asymmetric in λ_5 , since $\lambda_5 \geq 0$ corresponds to $m_H \geq m_A$ and thus the production cross section will be smaller compared to $m_H \leq m_A$. The magenta regions can additionally be covered by including also the single scalar production. We note that there is also a nonresonant signal contribution which comes from t -channel A/H exchange. Since the lepton p_T in this case is generally small and we are interested in the high p_T region where the BG is negligible, this contribution is separated and subtracted to evaluate the sensitivity. We find that the inclusion of the single production process can improve the experimental reach by 130 and 60 GeV for $|\lambda_5| \simeq 1$ and 2, respectively, when the Yukawa couplings are fixed to explain the central value of the muon $g-2$. This 130 GeV gain approximately reduces to 90 GeV (50 GeV) when the size of the NP contribution in $g-2$ is reduced by -1σ (-2σ). This still leaves a gap between the experimental reach and theoretical upper limit.

In order to further boost the sensitivity to the large-mass scenario it is important to increase the center of mass energy from $\sqrt{s} = 14$ TeV to, for instance, 27 TeV [79]. The colored lines in Fig. 4 (left) show the fiducial pair production cross section with $\sqrt{s} = 27$ TeV. The horizontal lines correspond to the sensitivity for integrated luminosities of 300 fb^{-1} , 1 ab^{-1} , and 3 ab^{-1} . Thanks to the larger center of mass energy we see that 1250 (1550) GeV can be covered with 1 (3) ab^{-1} of data. It is noted that the same kinematic cut introduced above has been applied for simplicity. The sensitivity is shown in cyan in the right panel.

Again the reach of the $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ channel can be extended by including the single production process. The magenta regions in Fig. 4 (right) in the right corners can also be probed. Compared to the sensitivity with $\sqrt{s} = 14$ TeV shown in gray on the left, the high energy (HE)-LHC is significantly more sensitive to the heavy scalar scenario. As a result, we observe that all the theoretically viable parameter region in Fig. 4 (right) can be covered.

Finally, a comment is in order concerning the impact of the assumptions on the Yukawa couplings $\rho_e^{\mu\tau}, \rho_e^{\tau\mu}$ entering our analysis. As mentioned in Sec. II for given scalar masses m_ϕ their effect in a_μ is maximized for equal couplings $|\rho_e^{\mu\tau}| = |\rho_e^{\tau\mu}|$ and we hence restricted our attention to this limit. While moving away from the equal coupling scenario decreases Δa_μ , the HA pair-production cross section and the branching ratio into the $\mu^\pm\mu^\pm\tau_h^\mp\tau_h^\mp$ final state remain unaffected. As a consequence, for $|\rho_e^{\mu\tau}| \neq |\rho_e^{\tau\mu}|$ the HA pair-production channel becomes more sensitive to the scalar solution to the $g-2$ anomaly in this model. The single-production cross section, as discussed above, decomposes into two contributions, the dominant of which is proportional to the same product $\rho_e^{\mu\tau}\rho_e^{\tau\mu}$ as Δa_μ . The sensitivity of this channel is therefore independent on the relative size of $\rho_e^{\mu\tau}$ and $\rho_e^{\tau\mu}$, as long as Δa_μ is fixed. It will on the other hand decrease (increase) if eventually a smaller (larger) NP contribution to $g-2$ is required.

IV. SUMMARY AND DISCUSSION

The new FNAL experimental data for the muon $g-2$ is consistent with the previously measured value at the Brookhaven experiment, and the significance of the long-standing discrepancy now amounts to 4.2σ . In this article we revisited the collider prospects of the $\mathbb{Z}_4$ -based 2HDM which can explain the discrepancy using new one-loop contributions involving τ and neutral scalars. A distinctive model prediction is the $\mu^\pm\mu^\pm\tau^\mp\tau^\mp$ signature at the LHC. Since the viable parameter space of the model cannot be fully probed at the LHC with the previously proposed electroweak scalar pair production, we investigated the impact of the single scalar production. We have shown that the latter mode helps to extend the reach for the $\mathcal{O}(1)$ TeV scalar solution of the muon $g-2$ discrepancy. For instance, the HL-LHC with 3 ab^{-1} can test up to $m_A = 1100$ GeV. We also examined the search potential at the HE-LHC and showed that increasing the center of the

mass energy is crucial to fully probe our scenario. While the model can also explain the DM relic abundance, the reach of the proposed search does not depend on the DM interpretation.

In this article we did not discuss $H^\pm\phi$ production. Due to a combinatorial factor the cross section is four times larger than the one of HA pair production [47]. The final state contains three charged leptons $2\tau + \mu$ or $2\mu + \tau$ and a neutrino at parton level, with their relative rates depending on the ratio of $|\rho_e^{\mu\tau}|^2$ and $|\rho_e^{\tau\mu}|^2$ [46]. Hence $H^\pm\phi$ production is expected to have better sensitivity to $|\rho_e^{\tau\mu}|^2$. Allowing for an imbalance in the couplings, $\rho_e^{\tau\mu} < \rho_e^{\mu\tau}$, while keeping the product $\rho_e^{\tau\mu}\rho_e^{\mu\tau}$ fixed, will thus decrease the LHC sensitivity to the $g-2$ solution in $H^\pm\phi$ production. At the same time the required larger value of $\rho_e^{\mu\tau}$ also lowers the cut-off scale Λ_{LP} of the theory. Note that the single scalar production cross section, being proportional to $(|\rho_e^{\mu\tau}|^2 + |\rho_e^{\tau\mu}|^2)$, is enhanced in this case. Thus, the signal discussed here is more universal. Combining those various search channels to further enhance the sensitivity would be an interesting future direction.

Motivated by the muon $g-2$ discrepancy, we focused on the 2HDM with $\mu\tau$ LFV couplings. While the $e\mu$ - and $e\tau$ -philic scenarios lack such motivation, LFV particles can be a viable DM mediator and they predict a similar collider phenomenology. Especially the $e\mu$ case is attractive in this respect, since the particle reconstruction is easier and the fiducial cross section via electroweak pair production is larger by a factor of five since the hadronic τ decay branching ratio and tagging efficiency no longer reduce the signal rate. This means that the relevant cross sections can be obtained from Fig. 3 and Fig. 4 by lifting the predictions by a factor of five.

ACKNOWLEDGMENTS

We thank Ulrich Nierste, Shohei Okawa, and Yuji Omura for valuable comments and discussion. We enjoy the support from the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) under Grant No. 396021762-TRR257. We also appreciate KIT for providing the necessary computational resources. M. B. thanks the Mainz Institute for Theoretical Physics (MITP) of the Cluster of Excellence PRISMA⁺ (Project ID 39083149) for its hospitality and support during the completion of this project.

- [1] T. Aoyama *et al.*, *Phys. Rep.* **887**, 1 (2020).
- [2] G. W. Bennett *et al.* (Muon g-2 Collaboration), *Phys. Rev. D* **73**, 072003 (2006).
- [3] B. Abi *et al.* (Muon g-2 Collaboration), *Phys. Rev. Lett.* **126**, 141801 (2021).
- [4] T. Aoyama, M. Hayakawa, T. Kinoshita, and M. Nio, *Phys. Rev. Lett.* **109**, 111808 (2012).
- [5] T. Aoyama, T. Kinoshita, and M. Nio, *Atoms* **7**, 28 (2019).
- [6] A. Czarnecki, W. J. Marciano, and A. Vainshtein, *Phys. Rev. D* **67**, 073006 (2003); **73**, 119901(E) (2006).
- [7] C. Gnendiger, D. Stöckinger, and H. Stöckinger-Kim, *Phys. Rev. D* **88**, 053005 (2013).
- [8] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, *Eur. Phys. J. C* **77**, 827 (2017).
- [9] A. Keshavarzi, D. Nomura, and T. Teubner, *Phys. Rev. D* **97**, 114025 (2018).
- [10] G. Colangelo, M. Hoferichter, and P. Stoffer, *J. High Energy Phys.* **02** (2019) 006.
- [11] M. Hoferichter, B.-L. Hoid, and B. Kubis, *J. High Energy Phys.* **08** (2019) 137.
- [12] M. Davier, A. Hoecker, B. Malaescu, and Z. Zhang, *Eur. Phys. J. C* **80**, 241 (2020); **80**, 410(E) (2020).
- [13] A. Keshavarzi, D. Nomura, and T. Teubner, *Phys. Rev. D* **101**, 014029 (2020).
- [14] A. Kurz, T. Liu, P. Marquard, and M. Steinhauser, *Phys. Lett. B* **734**, 144 (2014).
- [15] K. Melnikov and A. Vainshtein, *Phys. Rev. D* **70**, 113006 (2004).
- [16] P. Masjuan and P. Sanchez-Puertas, *Phys. Rev. D* **95**, 054026 (2017).
- [17] G. Colangelo, M. Hoferichter, M. Procura, and P. Stoffer, *J. High Energy Phys.* **04** (2017) 161.
- [18] M. Hoferichter, B.-L. Hoid, B. Kubis, S. Leupold, and S. P. Schneider, *J. High Energy Phys.* **10** (2018) 141.
- [19] A. Gérardin, H. B. Meyer, and A. Nyffeler, *Phys. Rev. D* **100**, 034520 (2019).
- [20] J. Bijnens, N. Hermansson-Truedsson, and A. Rodríguez-Sánchez, *Phys. Lett. B* **798**, 134994 (2019).
- [21] G. Colangelo, F. Hagelstein, M. Hoferichter, L. Laub, and P. Stoffer, *J. High Energy Phys.* **03** (2020) 101.
- [22] T. Blum, N. Christ, M. Hayakawa, T. Izubuchi, L. Jin, C. Jung, and C. Lehner, *Phys. Rev. Lett.* **124**, 132002 (2020).
- [23] G. Colangelo, M. Hoferichter, A. Nyffeler, M. Passera, and P. Stoffer, *Phys. Lett. B* **735**, 90 (2014).
- [24] M. Cè *et al.*, *Phys. Rev. D* **106**, 114502 (2022).
- [25] S. Borsanyi *et al.*, *Nature (London)* **593**, 51 (2021).
- [26] G. Colangelo, A. X. El-Khadra, M. Hoferichter, A. Keshavarzi, C. Lehner, P. Stoffer, and T. Teubner, *Phys. Lett. B* **833**, 137313 (2022).
- [27] C. Alexandrou *et al.*, *Phys. Rev. D* **107**, 074506 (2023).
- [28] A. Crivellin, M. Hoferichter, C. A. Manzari, and M. Montull, *Phys. Rev. Lett.* **125**, 091801 (2020).
- [29] A. Keshavarzi, W. J. Marciano, M. Passera, and A. Sirlin, *Phys. Rev. D* **102**, 033002 (2020).
- [30] G. Colangelo, M. Hoferichter, and P. Stoffer, *Phys. Lett. B* **814**, 136073 (2021).
- [31] A. Crivellin and M. Hoferichter, *J. High Energy Phys.* **07** (2021) 135; **10** (2022) 30.
- [32] S. Nie and M. Sher, *Phys. Rev. D* **58**, 097701 (1998).
- [33] R. A. Diaz, R. Martinez, and J. A. Rodriguez, *Phys. Rev. D* **64**, 033004 (2001).
- [34] S. Baek, N. G. Deshpande, X. G. He, and P. Ko, *Phys. Rev. D* **64**, 055006 (2001).
- [35] E. O. Iltan and H. Sundu, *Acta Phys. Slovaca* **53**, 17 (2003).
- [36] Y.-L. Wu and Y.-F. Zhou, *Phys. Rev. D* **64**, 115018 (2001).
- [37] K. A. Assamagan, A. Deandrea, and P.-A. Delsart, *Phys. Rev. D* **67**, 035001 (2003).
- [38] Y. Omura, E. Senaha, and K. Tobe, *J. High Energy Phys.* **05** (2015) 028.
- [39] Y. Omura, E. Senaha, and K. Tobe, *Phys. Rev. D* **94**, 055019 (2016).
- [40] J. Heeck, *Phys. Lett. B* **758**, 101 (2016).
- [41] W. Altmannshofer, C.-Y. Chen, P. S. Bhupal Dev, and A. Soni, *Phys. Lett. B* **762**, 389 (2016).
- [42] C.-W. Chiang and K. Tsumura, *J. High Energy Phys.* **05** (2018) 069.
- [43] P. S. B. Dev, R. N. Mohapatra, and Y. Zhang, *Phys. Rev. Lett.* **120**, 221804 (2018).
- [44] S. Iguro and Y. Omura, *J. High Energy Phys.* **05** (2018) 173.
- [45] Y. Abe, T. Toma, and K. Tsumura, *J. High Energy Phys.* **06** (2019) 142.
- [46] L. Wang and Y. Zhang, *Phys. Rev. D* **100**, 095005 (2019).
- [47] S. Iguro, Y. Omura, and M. Takeuchi, *J. High Energy Phys.* **11** (2019) 130.
- [48] A. Crivellin, D. Müller, and C. Wiegand, *J. High Energy Phys.* **06** (2019) 119.
- [49] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, *Phys. Rev. Lett.* **124**, 211803 (2020).
- [50] C. Cornella, P. Paradisi, and O. Sumensari, *J. High Energy Phys.* **01** (2020) 158.
- [51] S. Iguro, K. A. Mohan, and C. P. Yuan, *Phys. Rev. D* **101**, 075011 (2020).
- [52] S. Iguro, Y. Omura, and M. Takeuchi, *J. High Energy Phys.* **09** (2020) 144.
- [53] S. Jana, P. K. Vishnu, and S. Saad, *Phys. Rev. D* **101**, 115037 (2020).
- [54] H.-X. Wang, L. Wang, and Y. Zhang, *Eur. Phys. J. C* **81**, 1007 (2021).
- [55] A. J. Buras, A. Crivellin, F. Kirk, C. A. Manzari, and M. Montull, *J. High Energy Phys.* **06** (2021) 068.
- [56] N. Ghosh and J. Lahiri, *Eur. Phys. J. C* **81**, 1074 (2021).
- [57] W.-S. Hou, R. Jain, C. Kao, G. Kumar, and T. Modak, *Phys. Rev. D* **104**, 075036 (2021).
- [58] X.-F. Han, F. Wang, L. Wang, J. M. Yang, and Y. Zhang, *Chin. Phys. C* **46**, 103105 (2022).
- [59] J. Kriewald, J. Orloff, E. Pinsard, and A. M. Teixeira, *Eur. Phys. J. C* **82**, 844 (2022).
- [60] K. Asai, C. Miyao, S. Okawa, and K. Tsumura, *Phys. Rev. D* **106**, 035017 (2022).
- [61] P. M. Ferreira and J. P. Silva, *Phys. Rev. D* **83**, 065026 (2011).
- [62] I. P. Ivanov and C. C. Nishi, *J. High Energy Phys.* **11** (2013) 069.
- [63] H. Georgi and D. V. Nanopoulos, *Phys. Lett.* **82B**, 95 (1979).
- [64] J. F. Donoghue and L. F. Li, *Phys. Rev. D* **19**, 945 (1979).
- [65] P. A. Zyla *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **2020**, 083C01 (2020).

- [66] M. Sher, *Phys. Rep.* **179**, 273 (1989).
- [67] M. Muhlleitner, M. O. P. Sampaio, R. Santos, and J. Wittbrodt, *J. High Energy Phys.* **03** (2017) 094.
- [68] B. W. Lee, C. Quigg, and H. B. Thacker, *Phys. Rev. D* **16**, 1519 (1977).
- [69] S. Kanemura, T. Kubota, and E. Takasugi, *Phys. Lett. B* **313**, 155 (1993).
- [70] A. Arhrib, in *Workshop on Noncommutative Geometry, Superstrings and Particle Physics* (2000), [arXiv:hep-ph/0012353](#).
- [71] S. Iguro, S. Okawa, and Y. Omura, *J. High Energy Phys.* **03** (2023) 010.
- [72] W. Altmannshofer *et al.* (Belle-II Collaboration), *Prog. Theor. Exp. Phys.* **2019**, 123C01 (2019); **2020**, 029201(E) (2020).
- [73] G. Aad *et al.* (ATLAS Collaboration), *Phys. Rev. D* **101**, 052005 (2020).
- [74] G. Aad *et al.* (ATLAS Collaboration), *Eur. Phys. J. C* **80**, 123 (2020).
- [75] J. Alwall, R. Frederix, S. Frixione, V. Hirschi, F. Maltoni, O. Mattelaer, H. S. Shao, T. Stelzer, P. Torrielli, and M. Zaro, *J. High Energy Phys.* **07** (2014) 079.
- [76] V. Bertone, S. Carrazza, N. P. Hartland, and J. Rojo (NNPDF Collaboration), *SciPost Phys.* **5**, 008 (2018).
- [77] ATLAS Collaboration, Report No. ATL-PHYS-PUB-2019-033.
- [78] ATLAS Collaboration, Report No. ATL-PHYS-PUB-2015-025.
- [79] A. Abada *et al.* (FCC Collaboration), *Eur. Phys. J. Special Topics* **228**, 1109 (2019).