

# Dimensional regularization and two-loop vacuum polarization operator: Master integrals, analytic results, and energy shifts

Stefano Laporta<sup>1</sup> and Ulrich D. Jentschura<sup>2</sup>

<sup>1</sup>*Dipartimento di Fisica e Astronomia, Università di Padova e Istituto Nazionale di Fisica Nucleare, Sezione di Padova, Via Marzolo 8, I-35131 Padova, Italy*

<sup>2</sup>*Department of Physics and LAMOR, Missouri University of Science and Technology, Rolla, Missouri 65409, USA*



(Received 1 January 2024; accepted 9 March 2024; published 13 May 2024)

We present a complete reevaluation of the irreducible two-loop vacuum-polarization correction to the photon propagator in quantum electrodynamics, i.e. with an electron-positron pair in the fermion propagators. The integration is carried out by reducing the integrations to a limited set of master integrals, which are calculated using integration-by-parts identities. Dimensional regularization is used in  $D = 4 - 2\epsilon$  dimensions, and on-mass shell renormalization is employed. The one-loop effect is given to order  $\epsilon$ , to be combined with the  $1/\epsilon$  divergence of the two-loop amplitude. Master integrals are given. Final evaluations of two-loop energy shifts for  $1S$ ,  $2S$ , and  $2P$  states are done analytically, and results are presented, with an emphasis on muonic hydrogen. For relativistic Dirac-Coulomb reference states, higher-order coefficients are obtained for the  $Z\alpha$ -expansion. We compare the results obtained to the existing literature.

DOI: [10.1103/PhysRevD.109.096020](https://doi.org/10.1103/PhysRevD.109.096020)

## I. INTRODUCTION

The two-loop vacuum-polarization correction to bound-state energy levels is an important contribution to the Lamb shift in muonic bound systems. Specifically, the irreducible two-loop effect lowers the  $2S$  level energy in muonic hydrogen by roughly 1.25 meV in comparison to the  $2P$  level [1,2]. This energy shift needs to be compared to the proton-size discrepancy commonly referred to as the proton radius puzzle (0.88 fm versus 0.84 fm, for a recent brief summary see Ref. [3]). The proton radius puzzle corresponds to a 0.3 meV shift of the  $2S$  states, when expressed in energy units [1,2]. The two-loop effect is thus phenomenologically extremely relevant; any conceivable inaccuracy in its evaluation could lead to at least a partial theoretical explanation of the proton size discrepancy. Vacuum-polarization effects are drastically enhanced in muonic as compared to electronic bound systems [4–7]. Here, we present a reevaluation of the irreducible two-loop diagrams, on the basis of modern integration-by-parts techniques, employing dimensional regularization. We note that dimensional regularization was not yet sufficiently developed in 1973 (despite Refs. [8,9] which appeared in the literature at the time) to make an evaluation of the

two-loop effect using dimensional regularization techniques feasible [10]. For vacuum-polarization effects in particular, the modern techniques lead to drastic simplifications of the calculations. We find both the imaginary as well as the real part of the two-loop vacuum tensor in closed analytic form.

We also derive analytic expressions for the expectation value of the two-loop potential, evaluated on (nonrelativistic) Schrödinger-Coulomb eigenstates, generalizing the treatment originally outlined in Ref. [4] to two-loop order. In general, for predominantly nonrelativistic bound systems, energy shifts can be represented in terms of a semianalytic expansion (in powers of  $Z\alpha$  and  $\ln[(Z\alpha)^{-2}]$ , where  $\alpha$  is the fine-structure constant and  $Z$  is the nuclear charge number). As a byproduct, we derive a few higher-order coefficients from the expectation value of two-loop potential, evaluated on the relativistic Dirac-Coulomb eigenstates.

This paper is organized as follows. We start by briefly discussing the evaluation of the one-loop vacuum-polarization effect in Sec. II. Throughout this paper, we use dimensional regularization with the number  $D$  of dimensions expressed as

$$D = 4 - 2\epsilon. \quad (1)$$

The one-loop effect needs to be evaluated to order  $\epsilon$ , because finite contributions are generated at two-loop order when the one-loop terms of order  $\epsilon$  are multiplied by the  $1/\epsilon$ -terms from the two-loop amplitudes. The irreducible two-loop diagrams are discussed in Sec. III,

---

*Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP<sup>3</sup>.*

including master integrals and their series expansions, as well as complete results. A comparison to the available literature is also performed. In Sec. IV, we demonstrate that energy shifts due to the two-loop effect can be evaluated analytically, for muonic bound systems. Conclusions are reserved for Sec. V.

## II. ONE-LOOP VACUUM POLARIZATION

### A. Orientation

Our aim is to evaluate the one-loop vacuum-polarization effect to order  $\epsilon$ , i.e., in a form suitable for the later two-loop calculation. For simplicity we set the lepton mass (electron mass) in the vacuum-polarization loops to be  $m = 1$ . We use for a metric tensor  $g_{\mu\nu} = \text{diag}(1, -1, -1, -1)$  (see, for example, Refs. [11,12]), where

$$p^\mu = (p_0, \vec{p}), \quad p_\mu = (p_0, -\vec{p}), \quad (2)$$

$$\{\gamma^\mu, \gamma^\nu\} = 2g^{\mu\nu}, \quad S(p) = \frac{1}{\not{p} - m + i\epsilon}, \quad m = 1. \quad (3)$$

Here,  $m = 1$  is the electron mass.

Due to current conservation, the vacuum polarization function  $\Pi_{\mu\nu}$  has this tensorial structure

$$\Pi_{\mu\nu}(q^2) = (q_\mu q_\nu - q^2 g_{\mu\nu})\Pi(q^2). \quad (4)$$

We follow the convention of Ref. [13] so that

$$\left. \frac{d\Pi(q^2)}{dq^2} \right|_{q^2=0} > 0 \quad (5)$$

at the one-loop and two-loop level. For completeness, we note that the opposite conventions are used elsewhere in the literature, e.g., in Refs. [14,15]. The conventions employed here are consistent with those employed in Ref. [13]. Summation over the Lorentz indices in  $D$  dimensions leads to the result

$$\Pi^\mu{}_\mu(q^2) = (-q^2 g^\mu{}_\mu + q^\mu q_\mu)\Pi(q^2) = q^2(1-D)\Pi(q^2). \quad (6)$$

So, we can extract the scalar vacuum polarization function  $\Pi(q^2)$  as

$$\Pi(q^2) = \frac{\Pi^\mu{}_\mu(q^2)}{q^2(1-D)}. \quad (7)$$

The threshold for pair production of  $\Pi(q^2 = s)$  is  $q^2 = s = 4$ . The argument of the polarization tensor is

$$q^2 = (q^0)^2 - \vec{q}^2, \quad (8)$$

which is the four-momentum square of the photon entering the vacuum-polarization loop.

### B. Calculation

For the renormalization of the two-loop diagram with a self-mass insertion we need the one-loop vacuum polarization function  $\Pi^{(1)}$ , and the function  $\Pi^{(1a)}$  from the same diagram with iterated electron propagators on one side of the loop. The unrenormalized tensors read as follows:

$$\Pi_{\mu\nu}^{(1)}(q^2) = -ie^2 \int \frac{d^D k}{(2\pi)^D} [-\text{Tr}(\gamma_\mu S(k)\gamma_\nu S(k-q))], \quad (9)$$

$$\Pi_{\mu\nu}^{(1a)}(q^2) = ie^2 \int \frac{d^D k}{(2\pi)^D} [-\text{Tr}(\gamma_\mu [S(k)]^2 \gamma_\nu S(k-q))]. \quad (10)$$

The needed quantities are expressed in terms of two one-loop master integrals,

$$\Pi^{(1)}(q^2) = \left( \frac{e^2 C(\epsilon)}{4\pi\alpha} \right) \left( \frac{\alpha}{\pi} \right) P^{(1)}(q^2), \quad (11a)$$

$$\Pi^{(1a)}(q^2) = \left( \frac{e^2 C(\epsilon)}{4\pi\alpha} \right) \left( \frac{\alpha}{\pi} \right) P^{(1a)}(q^2). \quad (11b)$$

The prefactor in these results simplifies to unity provided we set

$$e^2 = \frac{4\pi\alpha}{C(\epsilon)}, \quad C(\epsilon) = \Gamma(1+\epsilon)(4\pi)^\epsilon \mu^{-2\epsilon}. \quad (12)$$

One observes that the identification (12) is not completely canonical in dimensional regularization. Namely [see, e.g., Eq. (10.173) of Ref. [15]], in the  $\overline{\text{MS}}$  scheme, one normally has the relation  $e^2 = (4\pi)^{1-\epsilon} \alpha \mu^{2\epsilon} e^{\gamma_E \epsilon}$ , where  $\gamma_E = 0.5772156649\dots$  is the Euler-Mascheroni constant, and  $\mu$  is the renormalization scale induced by dimensional regularization. This is equivalent to Eq. (12) up to terms linear in  $\epsilon$ . The prefactor  $1/C(\epsilon)$  helps to simplify results after integration.

The one-loop integrals are as follows:

$$P^{(1)}(q^2) = \frac{(D-2)q^2 + 4}{2(D-1)q^2} M_{12}(q^2) + \frac{(D-2)M_2(q^2)}{(D-1)q^2}, \quad (13)$$

$$P^{(1a)}(q^2) = \frac{(D-4)q^2 + 4}{q^2(q^2-4)} M_{12}(q^2) + \frac{2(D-2)M_2(q^2)}{q^2(q^2-4)}, \quad (14)$$

where

$$M_{12}(q^2) = N(\epsilon) \int \frac{[d^D k_1]}{D_1 D_2}, \quad (15a)$$

$$M_2(q^2) = N(\epsilon) \int \frac{[d^D k_1]}{D_2}, \quad (15b)$$

$$D_j = -p_j^2 + 1 - i\epsilon, \quad j = 1, \dots, 2. \quad (15c)$$

$$p_1 = k_1, \quad p_2 = k_1 - q, \quad (15d)$$

$$[d^D k] = \mu^{-2\epsilon} d^D k. \quad (15e)$$

The above definition of  $[d^D k]$  contains the factor  $\mu^{-2\epsilon}$ . It simplifies with the factor  $1/\mu^{-2\epsilon}$  of  $N(\epsilon)$  so that the master integrals are independent of  $\mu$ . The normalization factor is

$$N(\epsilon) = \frac{\mu^{2\epsilon}}{\pi^{2-\epsilon}\Gamma(1+\epsilon)} = \frac{4^\epsilon}{\pi^{2-2\epsilon}C(\epsilon)} = \frac{4(4\pi^2)}{(2\pi)^{4-2\epsilon}C(\epsilon)}. \quad (16)$$

The latter form exhibits the dimension  $D = 4 - 2\epsilon$ . With the normalization factor  $N(\epsilon)$ , all master integrals  $M$  occurring in our calculations are dimensionless and not only uncluttered from logarithms of  $\mu$ , but also, from logarithms of  $4\pi$ .

One can show that the one-loop master integrals satisfy the following differential equations:

$$M_2'(q^2) = 0, \quad (17a)$$

$$\begin{aligned} [(D-4)q^2 + 4]M_{12}(q^2) + 2(D-2)M_2(q^2) \\ - 2(q^2 - 4)q^2 M_2'(q^2) = 0, \end{aligned} \quad (17b)$$

where the differentiation is with respect to  $q^2$ . One finds that  $M_2$  is a momentum-independent vacuum integral,

$$M_2(q^2) = -\frac{1}{\epsilon(1-\epsilon)}. \quad (18)$$

For  $M_{12}(q^2)$ , one writes the result as a function of the variable  $v = v(q^2)$ ,

$$v = \sqrt{1 - \frac{4}{q^2}}. \quad (19)$$

The result is

$$\begin{aligned} (1-2\epsilon)M_{12}(q^2) \\ = \frac{1}{\epsilon} + v \ln\left(\frac{v-1}{v+1}\right) - \frac{\epsilon}{6}v \left[ \pi^2 + 12\text{Li}_2\left(\frac{1-v}{v+1}\right) \right. \\ \left. + 12 \ln\left(\frac{2v}{v+1}\right) \ln\left(\frac{v-1}{v+1}\right) - 3\ln^2\left(\frac{v-1}{v+1}\right) \right] \\ + \epsilon^2 \frac{v}{6} N_{12}(v) + \mathcal{O}(\epsilon^3), \end{aligned} \quad (20)$$

where we include the terms of order  $\epsilon$  and  $\epsilon^2$  which will be useful in the context of the two-loop calculation. The explicit form of  $N_{12}(q^2)$  is given in Appendix A.

### C. Renormalization

Substituting Eqs. (18) and (20) into (13), we can find the expansions of  $P^{(1)}(q^2)$  and  $P^{(1a)}(q^2)$ . But for the renormalization we need to subtract the values at  $q^2 = 0$ ,

$$P^{(1)}(0) = \frac{1}{3\epsilon}, \quad P^{(1a)}(0) = \frac{1}{3}. \quad (21)$$

In our formalism, the scaled functions  $P^{(1)}$  and  $P^{(2)}$  do not have powers of  $\mu$ , but the scalar vacuum polarization functions  $\Pi^{(1)}$ ,  $\Pi^{(2)}$  depend on  $\mu$ . Yet, after renormalization the dependence disappears in the limit  $\epsilon \rightarrow 0$ . The renormalized expression for the one-loop function  $P_R^{(1)}(q^2)$  is as follows:

$$\begin{aligned} P_R^{(1)}(q^2) &= P^{(1)}(q^2) - P^{(1)}(0) \\ &= \frac{1}{9}(8-3v^2) - \frac{v}{6}(v^2-3) \ln\left(\frac{v-1}{v+1}\right) \\ &\quad + \epsilon Q_1^{(1)}(v) + \epsilon^2 Q_2^{(1)}(v) + \mathcal{O}(\epsilon^3). \end{aligned} \quad (22)$$

The coefficient of the term linear in  $\epsilon$  is

$$\begin{aligned} Q_1^{(1)}(v) &= \frac{v}{36}(v^2-3) \left[ 12\text{Li}_2\left(\frac{1-v}{v+1}\right) - 3\ln^2\left(\frac{v-1}{v+1}\right) \right. \\ &\quad \left. + 12 \ln\left(\frac{2v}{v+1}\right) \ln\left(\frac{v-1}{v+1}\right) + \pi^2 \right] \\ &\quad + \left( v - \frac{4v^3}{9} \right) \ln\left(\frac{v-1}{v+1}\right) - \frac{8v^2}{9} + \frac{52}{27}. \end{aligned} \quad (23)$$

For the explicit form of the term of order  $\epsilon^2$ , we refer to Appendix A.

The renormalized value for the function  $P_R^{(1a)}(q^2)$  with iterated electron propagators is as follows:

$$\begin{aligned} P_R^{(1a)}(q^2) &= P^{(1a)}(q^2) - P^{(1a)}(0) \\ &= \frac{v^2-1}{4v} \left\{ \frac{2v(3v^2-5)}{3(v^2-1)} + (v^2-1) \ln\left(\frac{v-1}{v+1}\right) \right\} \\ &\quad + \epsilon Q_1^{(1a)}(v) + \epsilon^2 Q_2^{(1a)}(v) + \mathcal{O}(\epsilon^3). \end{aligned} \quad (24)$$

The term linear in  $\epsilon$  is given as

$$\begin{aligned} Q_1^{(1a)}(v) &= \frac{v^2-1}{4v} \left[ (v^2-1) \left( -2\text{Li}_2\left(\frac{1-v}{v+1}\right) \right. \right. \\ &\quad \left. \left. + \frac{1}{2}\ln^2\left(\frac{v-1}{v+1}\right) - \frac{\pi^2}{6} \right) + 2\ln\left(\frac{v-1}{v+1}\right) \right. \\ &\quad \left. \times \left( v^2 - (v^2-1) \ln\left(\frac{2v}{v+1}\right) \right) + 4v \right]. \end{aligned} \quad (25)$$

The quadratic term in  $\epsilon$ , denoted as  $Q_2^{(1a)}(v)$ , is given in Appendix A. Actually, only the terms proportional to  $\epsilon$  are

necessary for the renormalization of the two-loop vacuum polarization, but higher-order terms are useful for higher-loop calculations, and are thus relegated to Appendix A.

In the limit  $\epsilon \rightarrow 0^+$ , one recovers the known result for the one-loop vacuum polarization,

$$\begin{aligned} \Pi_{\text{R}}^{(1)}(q^2) &= \left(\frac{\alpha}{\pi}\right) P_{\text{R}}^{(1)}(q^2) \\ &= \left(\frac{\alpha}{\pi}\right) \left[ \frac{1}{9} (8 - 3v^2) - \frac{1}{6} v(v^2 - 3) \ln\left(\frac{v-1}{v+1}\right) \right]. \end{aligned} \quad (26)$$

The dispersion relation is

$$\Pi_{\text{R}}^{(1)}(q^2) = \frac{q^2}{\pi} \int_4^\infty dq'^2 \frac{\text{Im}\Pi_{\text{R}}^{(1)}(q'^2 + i\epsilon)}{q'^2(q'^2 - q^2)}. \quad (27)$$

The imaginary part is positive infinitesimally above the cut, and negative infinitesimally below the cut. One has the simple representation

$$\text{Im}\Pi_{\text{R}}^{(1)}(q^2 + i\epsilon) = \frac{\alpha}{6} v(3 - v^2) \Theta(q^2 - 4). \quad (28)$$

On the cut, the  $v$  variable runs from  $v = 0$  to  $v = 1$ . Here,  $\Theta$  denotes the Heaviside step function. The above result given in Eq. (26) in the  $v$  representation fulfills [see also Eq. (5)]

$$\Pi_{\text{R}}^{(1)}(q^2) = \frac{\alpha q^2}{\pi 15} + \mathcal{O}(q^4), \quad (29)$$

for  $q^2 = (q^0)^2 - \vec{q}^2$ .

### III. TWO-LOOP VACUUM POLARIZATION

#### A. Orientation

We focus on the diagrams in Fig. 1. The first diagram 1 is the proper two-loop vacuum polarization diagram, while diagram 2 is a one-loop vacuum polarization diagram with a self-energy insertion. The expression  $\Pi_{\mu\nu}^{(2;1)}(q^2)$  corresponds to diagram 1,

$$\begin{aligned} \Pi_{\mu\nu}^{(2;1)}(q^2) &= e^4 \int \frac{d^D k_1}{(2\pi)^D} \frac{d^D k_2}{(2\pi)^D} \\ &\times \left[ -\text{Tr}(\gamma_\mu S(p_4) \gamma^\rho S(p_2) \gamma_\nu S(-p_1) \gamma^\sigma \right. \\ &\times \left. S(-p_3) D_{\rho\sigma}(p_5) \right], \end{aligned} \quad (30a)$$

while  $\Pi_{\mu\nu}^{(2;2)}(q^2)$  corresponds to the self-energy insertion into a fermionic line of the vacuum-polarization loop,




FIG. 1. The diagrams concern the electronic two-loop vacuum-polarization diagrams in muonic hydrogen (the negatively charged muon line is denoted by  $\mu^-$ ). These are naturally divided into diagram (1), which is a proper two-loop diagram, diagram (2), which is a one-loop diagram with a self-energy insertion, and the loop-after-loop (reducible) diagram (3).

$$\begin{aligned} \Pi_{\mu\nu}^{(2;2)}(q^2) &= e^4 \int \frac{d^D k_1}{(2\pi)^D} \frac{d^D k_2}{(2\pi)^D} \\ &\times \left[ -\text{Tr}(\gamma_\mu S(p_4) \gamma^\rho S(p_2) \gamma^\sigma S(p_4) \gamma_\nu \right. \\ &\times \left. S(-p_3) D_{\rho\sigma}(p_5) \right]. \end{aligned} \quad (30b)$$

Here, the photon propagator is given by [see Eq. (9.133) of Ref. [15]]

$$D_{\rho\sigma}(k) = -\frac{1}{k^2 + i\epsilon} \left( g_{\rho\sigma} - \lambda \frac{k_\rho k_\sigma}{k^2} \right), \quad (31)$$

and the momenta and denominators are

$$p_1 = k_1, \quad p_2 = q - k_1, \quad p_3 = k_2, \quad (32a)$$

$$p_4 = q - k_2, \quad p_5 = k_1 - k_2, \quad (32b)$$

$$D_j = -p_j^2 + 1 - i\epsilon, \quad j = 1, \dots, 4, \quad (32c)$$

$$D_5 = -p_5^2 - i\epsilon. \quad (32d)$$

These conventions are adapted to FORM [16,17]; in an East Coast metric with  $g_{\mu\nu} = \text{diag}(-1, -1, -1, 1)$ , one would replace  $p_j^2 \rightarrow -p_j^2$ .

#### B. Reduction to master integrals

After the calculation of traces, performed with computer algebra (using FORM, see Refs. [16,17]), and a Wick rotation, the scalar contribution  $\Pi^{(1)}(q^2)$  and  $\Pi^{(2)}(q^2)$  of the two diagrams are reduced to a combination of 24 and 41 Feynman integrals, respectively. In order to further reduce the expressions, we use the integration by parts (IBP) identities [18,19]. A system of IBP identities is built and solved with the program SYS, which is based on an algorithm described in Refs. [20,21]. These identities allow one to reduce  $\Pi^{(1)}(q^2)$  and  $\Pi^{(2)}(q^2)$  to a linear combination of five irreducible master integrals.

The choice of the master integrals is somewhat arbitrary; we chose the following ones:

$$M_{1234} = [N(\epsilon)]^2 \int \frac{[d^D k_1][d^D k_2]}{D_1 D_2 D_3 D_4}, \quad (33a)$$

$$M_{234} = [N(\epsilon)]^2 \int \frac{[d^D k_1][d^D k_2]}{D_2 D_3 D_4}, \quad (33b)$$

$$M_{235} = [N(\epsilon)]^2 \int \frac{[d^D k_1][d^D k_2]}{D_2 D_3 D_5}, \quad (33c)$$

$$M_{235k} = [N(\epsilon)]^2 \int \frac{[d^D k_1][d^D k_2] q \cdot k_2}{D_2 D_3 D_5}, \quad (33d)$$

$$M_{24} = [N(\epsilon)]^2 \int \frac{[d^D k_1][d^D k_2]}{D_2 D_4}. \quad (33e)$$

The following master integrals are reducible and factorize into a product of two one-loop master integrals,

$$M_{24}(q^2) = [M_2(q^2)]^2, \quad (34a)$$

$$M_{234}(q^2) = M_2(q^2) M_{12}(q^2), \quad (34b)$$

$$M_{1234}(q^2) = [M_{12}(q^2)]^2. \quad (34c)$$

The expressions for the contribution of diagrams 1 and 2 are given as

$$\Pi^{(2;1)}(q^2) = [C(\epsilon)]^2 \left(\frac{\alpha}{\pi}\right)^2 (P^{(2;1)}(q^2) + \lambda G^{(2;1)}(q^2)), \quad (35a)$$

$$\Pi^{(2;2)}(q^2) = [C(\epsilon)]^2 \left(\frac{\alpha}{\pi}\right)^2 (P^{(2;2)}(q^2) + \lambda G^{(2;2)}(q^2)), \quad (35b)$$

where  $\lambda$  is the gauge parameter [see Eq. (31)], and

$$\begin{aligned} P^{(2;1)}(q^2) &= \mathcal{P}_{24}^{(2;1)} M_{24}(q^2) + \mathcal{P}_{234}^{(2;1)} M_{234}(q^2) \\ &+ \mathcal{P}_{1234}^{(2;1)} M_{1234}(q^2) + \mathcal{P}_{235}^{(2;1)} M_{235}(q^2) \\ &+ \mathcal{P}_{235k}^{(2;1)} M_{235k}(q^2). \end{aligned} \quad (36)$$

The coefficient functions  $\mathcal{P}_{24}^{(2;1)}$ ,  $\mathcal{P}_{234}^{(2;1)}$ ,  $\mathcal{P}_{1234}^{(2;1)}$ ,  $\mathcal{P}_{235}^{(2;1)}$  and  $\mathcal{P}_{235k}^{(2;1)}$  are given in Appendix B. The gauge-dependent part of diagram 1 is given as

$$\begin{aligned} G^{(2;1)}(q^2) &= \mathcal{G}_{24}^{(2;1)} M_{24}(q^2) + \mathcal{G}_{235}^{(2;1)} M_{235}(q^2) \\ &+ \mathcal{G}_{235k}^{(2;1)} M_{235k}(q^2), \end{aligned} \quad (37)$$

with the coefficient functions  $\mathcal{G}_{24}^{(2;1)}$ ,  $\mathcal{G}_{235}^{(2;1)}$  and  $\mathcal{G}_{235k}^{(2;1)}$  also being indicated in Appendix B. For diagram 2, one has the following reduction:

$$\begin{aligned} P^{(2;2)}(q^2) &= \mathcal{P}_{234}^{(2;2)} M_{234}(q^2) + \mathcal{P}_{235k}^{(2;2)} M_{235k}(q^2) \\ &+ \mathcal{P}_{235}^{(2;2)} M_{235}(q^2) + \mathcal{P}_{24}^{(2;2)} M_{24}(q^2). \end{aligned} \quad (38)$$

Again, the coefficient functions  $\mathcal{P}_{24}^{(2;2)}$ ,  $\mathcal{P}_{234}^{(2;2)}$ ,  $\mathcal{P}_{1234}^{(2;2)}$ ,  $\mathcal{P}_{235}^{(2;2)}$ , and  $\mathcal{P}_{235k}^{(2;2)}$  are given in Appendix B. The gauge-dependent part of the second diagram cancels the gauge-dependent part of the first diagram, in view of the relation  $G^{(2;2)}(q^2) = -\frac{1}{2}G^{(2;1)}(q^2)$ .

### C. Differential equations and master integrals

A system of differential equations in  $q^2$  is being created; we need to compute the five master integrals. Three of them are reducible into products of one-loop integrals, as outlined in Eq. (34). We obtain a system of differential equations in  $q^2$  satisfied by the irreducible master integrals [22–24]. The first of these equations is of second order,

$$\begin{aligned} q^2[-3((D-4)q^2+4)M'_{235}(q^2) \\ + 2q^2(q^2-4)M''_{235}(q^2)] + (D-2)M_{24}(q^2) \\ + (D-3)[D(q^2+2)-4(q^2+1)]M_{235}(q^2) = 0, \end{aligned} \quad (39a)$$

while the remaining one is a first-order differential equation,

$$\begin{aligned} [D(q^2+2)-q^2-6]M_{235}(q^2) + 6(D-2)M_{235k}(q^2) \\ + (D-2)M_{24}(q^2) + q^2(q^2-4)M'_{235}(q^2) = 0, \end{aligned} \quad (39b)$$

where the differentiation is with respect to  $q^2$ . The initial conditions at  $q^2 = 0$  for these diagrams are simple vacuum integrals.

### D. Results for the master integrals

Due to the appearance of the factor  $D-4$  in the denominators of Eqs. (36)–(38) (for the coefficients, see Appendix B), the master integrals have to be evaluated up to order  $\mathcal{O}(\epsilon)$ . It is not sufficient to keep only the finite part. The results are as follows. For  $M_{235}$ , one finds the following relation, where we note the prefactor  $(1-v^2)$ :

$$\begin{aligned} (1-v^2)M_{235}(q^2) &= \frac{v^2-1}{\epsilon^2} + \frac{3v^2-2}{\epsilon} + \left[ 6v^2 + \frac{1}{2} \right. \\ &+ \frac{1}{4}(-v^4-2v^2+3)\ln^2\left(\frac{v-1}{v+1}\right) \\ &- (v^2-3)v\ln\left(\frac{v-1}{v+1}\right) \left. \right] \\ &+ \epsilon N_{235}(q^2) + \mathcal{O}(\epsilon^2). \end{aligned} \quad (40)$$

The order- $\epsilon$  coefficient  $N_{235}(q^2)$  is found in Appendix C. For  $M_{235k}$ , one finds the relation

$$\begin{aligned}
 & (1-v^2)^2 M_{235k}(q^2) \\
 &= \frac{1-v^2}{\epsilon^2} + \frac{1}{\epsilon} \left[ \frac{13}{6} - \frac{7v^2}{2} \right] \\
 &+ \frac{1}{8} (-v^6 + 3v^4 + 5v^2 - 7) \ln^2 \left( \frac{v-1}{v+1} \right) \\
 &- \frac{1}{6} v(3v^4 - 8v^2 + 21) \ln \left( \frac{v-1}{v+1} \right) \\
 &+ \frac{1}{12} (-6v^4 - 91v^2 - 7) + \epsilon N_{235k}(q^2) + \mathcal{O}(\epsilon^2). \quad (41)
 \end{aligned}$$

Again,  $N_{235k}(q^2)$  can be found in Appendix C. Similar master integrals have recently been considered in Refs. [25,26].

### E. Renormalization

We now substitute the master integrals (34a)–(34c), (40), and (41) into the reduction formulas given in Eqs. (36) and (38). We can thus find the series expansions of the vacuum-polarization functions  $\Pi^{(2;1)}(q^2)$  and  $\Pi^{(2;2)}(q^2)$ . For the renormalization, we need the values at  $q^2 = 0$ . The behavior, for small  $q^2$ , is the following:

$$\begin{aligned}
 P^{(2;1)}(q^2) &= -\frac{12\epsilon^5 + 4\epsilon^4 - 23\epsilon^3 - 14\epsilon^2 - 21\epsilon - 6}{12(\epsilon-2)(\epsilon-1)\epsilon^2(2\epsilon+1)(2\epsilon+3)} \\
 &- \frac{1}{q^2} \frac{1}{(1-\epsilon)(1+2\epsilon)(1-2\epsilon)} + \mathcal{O}(q^2), \\
 P^{(2;2)}(0) &= -\frac{(\epsilon+1)(4\epsilon^4 - 8\epsilon^3 - 8\epsilon^2 + 21\epsilon + 3)}{12(\epsilon-2)(\epsilon-1)\epsilon^2(2\epsilon+1)(2\epsilon+3)} \\
 &+ \frac{1}{2q^2} \frac{1}{(1-\epsilon)(1+2\epsilon)(1-2\epsilon)} + \mathcal{O}(q^2). \quad (42)
 \end{aligned}$$

Terms containing  $1/q^2$  appear, due to the fact that the single diagrams are not gauge invariant. Taking the gauge invariant sum  $P^{(2;1)}(q^2) + 2P^{(2;2)}(q^2)$ , the divergent terms cancel out, so that, taking the limit  $q^2 \rightarrow 0$  one gets

$$P^{(2;1)}(0) + 2P^{(2;2)}(0) = -\frac{10\epsilon^2 - 7\epsilon - 9}{12(\epsilon-2)\epsilon(2\epsilon+1)}. \quad (43)$$

In order to carry out the renormalization, we need the renormalization constants

$$Z_J = 1 + \sum_L Z_J^{(L)} \left( \frac{\alpha}{\pi} \right)^L \left( \frac{e^2 C(\epsilon)}{4\pi\alpha} \right)^L, \quad (44)$$

where  $J$  stands for the vertex renormalization ( $J = 1$ ), the wave function renormalization ( $J = 2$ ), and the mass

renormalization ( $J = m$ ), and  $L$  enumerates the loop order. We define the one-loop quantities

$$F = -Z_1^{(1)}, \quad S_0 = \delta m = -Z_m^{(1)}, \quad S_1 = -Z_2^{(1)}, \quad (45)$$

and obtain the results

$$F = \frac{3-2\epsilon}{4\epsilon-8\epsilon^2}, \quad S_0 = \frac{3-2\epsilon}{4\epsilon-8\epsilon^2}, \quad S_1 = S_0. \quad (46)$$

We note that  $Z_1^{(1)} = Z_2^{(1)}$  because of the Ward-Takahashi identity. The identity  $Z_2^{(1)} = Z_m^{(1)}$  is a simple coincidence, which occurs only at one-loop order in dimensional regularization, but to all orders in  $\epsilon$ . The constants  $F$ ,  $S_0$  and  $S_1$  in Eq. (45) are defined with the prefactor  $(-1)$  so that the expansion of the self-energy insertion into the fermion line about the mass shell acquires positive terms, for  $\not{q} \approx m = 1$ :

$$\begin{aligned}
 \frac{1}{\not{q}-m} \Sigma(q) \frac{1}{\not{q}-m} &= \frac{1}{\not{q}-m} [S_0 + (\not{q}-m)S_1] \frac{1}{\not{q}-m} \\
 &= \frac{S_1}{\not{q}-m} + \frac{S_0}{(\not{q}-m)^2}. \quad (47)
 \end{aligned}$$

Finally, the renormalized contributions of the diagrams are found as

$$P_R^{(2;1)}(q^2) = P^{(2;1)}(q^2) - P^{(2;1)}(0) - 2FP_R^{(1)}(q^2), \quad (48)$$

$$\begin{aligned}
 P_R^{(2;2)}(q^2) &= P^{(2;2)}(q^2) - P^{(2;2)}(0) \\
 &- (S_1 P_R^{(1)}(q^2) + S_0 P_R^{(1a)}(q^2))(-1), \quad (49)
 \end{aligned}$$

and the renormalized two-loop function  $P_R^{(2)}(q^2)$  is obtained as

$$P_R^{(2)}(q^2) = P_R^{(2;1)}(q^2) + 2P_R^{(2;2)}(q^2). \quad (50)$$

### F. Results

We recall the definition of  $v$  in Eq. (19). In the limit  $\epsilon \rightarrow 0^+$ , one obtains the two-loop vacuum-polarization function

$$\Pi_R^{(2)}(q^2) = \left( \frac{\alpha}{\pi} \right)^2 P_R^{(2)}(q^2). \quad (51)$$

We find a compact representation (see also Ref. [27])

$$\begin{aligned} \Pi_{\text{R}}^{(2)}(q^2) = & \left(\frac{\alpha}{\pi}\right)^2 \left\{ \frac{v^4 - 2v^2 - 3}{12} \left[ \Phi_1\left(\frac{v-1}{v+1}\right) \ln^2\left(\frac{v-1}{v+1}\right) - 4\Phi_2\left(\frac{v-1}{v+1}\right) \ln\left(\frac{v-1}{v+1}\right) + 6\Phi_3\left(\frac{v-1}{v+1}\right) \right. \right. \\ & + 3\zeta(3) \left. \right] + \frac{3v - v^3}{12} \left[ 4\Phi_1\left(\frac{v-1}{v+1}\right) \ln\left(\frac{v-1}{v+1}\right) - 4\Phi_2\left(\frac{v-1}{v+1}\right) + 3\ln^2\left(\frac{v-1}{v+1}\right) \right] + \frac{5v - 3v^3}{8} \ln\left(\frac{v-1}{v+1}\right) \\ & + \frac{1}{96} (7v^4 - 22v^2 - 33) \ln^2\left(\frac{v-1}{v+1}\right) - \frac{13(v^2 - 1)}{24} + \frac{5}{24} \left. \right\}, \end{aligned} \quad (52)$$

where

$$\Phi_n(x) = \text{Li}_n(x) + 2\text{Li}_n(-x), \quad \text{Li}_1(x) = -\ln(1-x). \quad (53)$$

For small positive  $q^2$ , one obtains the expression

$$\Pi_{\text{R}}^{(2)}(q^2) = \left(\frac{\alpha}{\pi}\right)^2 \frac{41q^2}{162} + \dots \quad (54)$$

The representation for the imaginary part just above the cut is even more compact,

$$\begin{aligned} \text{Im}\Pi_{\text{R}}^{(2)}(q^2 + i\epsilon) &= \frac{\alpha^2}{\pi} \Theta(q^2 - 4) \left\{ \frac{v^4 - 2v^2 - 3}{6} \left[ \Phi_1\left(\frac{v-1}{v+1}\right) \ln\left(\frac{1-v}{1+v}\right) \right. \right. \\ & - 2\Phi_2\left(\frac{v-1}{v+1}\right) \left. \right] + \frac{3v - v^3}{6} \left[ 2\Phi_1\left(\frac{v-1}{v+1}\right) \right. \\ & + 3\ln\left(\frac{1-v}{1+v}\right) \left. \right] + \frac{1}{48} (7v^4 - 22v^2 - 33) \ln\left(\frac{1-v}{1+v}\right) \\ & + \frac{5v - 3v^3}{8} \left. \right\}. \end{aligned} \quad (55)$$

### G. Comparison with the literature

An essential ingredient of our calculations is the master integrals, which should be compared to results communicated in Refs. [28–30]. In Refs. [28,29], the authors define a master integral  $J_{011}$  which is equivalent to our  $M_{12}$  up a factor

$$J_{011} = i\pi^{2-\epsilon} \Gamma(1 + \epsilon) M_{12}. \quad (56)$$

In Eq. (1) of Ref. [28], the general term of the  $\epsilon$ -expansion of  $J_{011}$  is given in terms of log-sine integrals which can be written in terms of Nielsen polylogarithms; the analytic continuation is shown in Eqs. (2.9)–(2.23) of Ref. [29]. Moreover, an closed analytical expression for  $J_{011}$  is given in Eqs. (2.10)–(2.14) of Ref. [29], containing hypergeometric  ${}_2F_1$  functions.

Changing the variable from  $t = q^2$  to  $v$  and using the transformations for Nielsen polylogarithm  $S_{n,p}(-1/z) \rightarrow S_{n,p}(-z)$ , we found that the results in our Eqs. (20) and (A1) agree perfectly with the corresponding terms of

Eq. (1) of Ref. [28] and the expansion given in Eqs. (2.10)–(2.14) of Ref. [29].

In Eqs. (4.9), (4.10), and (E.4) of Ref. [30], analytical results for master integrals equivalent to our  $M_{235}$  and  $M_{235k}$  were presented. These master integrals are related to ours by the following relations: one finds for the ultraviolet-convergent integral (4.9) of Ref. [30],

$$\begin{aligned} & 4q^2(q^2 - 4)^2 J_{011}(1, 2, 2) \\ &= -24(D-3)(D-2)(q^2 + 2) M_{235k} \\ & - (D-2)((D-2)q^4 - 2Dq^2 \\ & + 16D - 40) M_{24} - 4(D-3)(2(D-2)q^4 \\ & - (D-6)q^2 + 8D - 20) M_{235}, \end{aligned} \quad (57)$$

while, for the ultraviolet-convergent linear combination of integrals given in Eq. (4.10) of Ref. [30], one has

$$\begin{aligned} & 4(D-4)q^2(q^2 - 4)^3 (J_{011}(1, 2, 2) + 2J_{011}(2, 1, 2)) \\ &= -(D-2)[(D-4)(D-2)q^6 + 2(D-6)(D-4)q^4 \\ & + 16(D(5D-31) + 47)q^2 - 32(2D-7)(2D-5)] M_{24} \\ & - 4(D-3)[4(D-4)(D-2)q^6 + 3D(3D-10)q^4 \\ & + 24(D-4)(D-3)q^2 - 16(2D-7)(2D-5)] M_{235} \\ & - 48(D-3)(D-2)(q^2 - 1) \\ & \times (D(q^2 + 8) - 4(q^2 + 7)) M_{235k}, \end{aligned} \quad (58)$$

and for the ultraviolet divergent integral (E.4) of Ref. [30],

$$J_{011}(1, 1, 1) = M_{235}. \quad (59)$$

The  $\epsilon$ -expansion obtained from those of  $M_{235}$  and  $M_{235k}$  agree perfectly with those presented in Ref. [30].

Our results for the complete two-loop function given in Eq. (52) and for its imaginary part, as given in Eq. (55), agree with the calculation of Ref. [31]. One notes that Eq. (49) of Ref. [31] contains the imaginary part, while Eq. (57) of Ref. [31] also contains the real part of the two-loop vacuum polarization. One should note, though, that some integrals are left unevaluated in Ref. [31]. Furthermore, it needs to be pointed out that the expressions in Ref. [31] also contain the reducible diagram (see diagram 3 in Fig. 1) with two iterated one-loop vacuum

polarizations on the same line. Our result for the imaginary part also agrees with the result given in Eq. (5-4.200) of Ref. [32]. Our results are also in agreement with the paper [33], specifically, with the results given in Eqs. (71), (78)–(81) of Ref. [33]. Furthermore, we can refer to related investigations on the vector part of the quantum chromodynamic vacuum-polarization tensor [34–36].

In Ref. [10], the two-loop vacuum polarization was also calculated. The result given in Eq. (49) of Ref. [10] is expressed in terms of the variable  $\theta$ , which in our notation reads as  $\theta = -\frac{1-v}{1+v}$ . The last-but-one line of Eq. (49) of Ref. [10] contains two terms, which, in the notation of Ref. [10], read as  $2\ln(\theta)\text{Li}_2(\theta)$  and  $2\ln(\theta)\text{Li}_2(-\theta)$ . The term  $2\ln(\theta)\text{Li}_2(-\theta)$  contains a misprint; it should have read as  $4\ln(\theta)\text{Li}_2(-\theta)$ . Formula (49) of Ref. [10] was referred to in Ref. [37], and it was stated in Ref. [37] that Eq. (49) of Ref. [10] has a misprint. Indeed, Eq. (49) of Ref. [10] is rewritten as Eq. (6) of Ref. [37], with a change of notation: The variable  $\delta$  used in Ref. [37] is related to the variable  $\theta$  used in Eq. (49) of Ref. [10] by the relation  $\delta = (1+\theta)/(1-\theta) = v$ . In Eq. (6) of Ref. [37], the wrong term  $2\ln(\theta)\text{Li}_2(-\theta)$  is corrected to the right  $4\ln(\theta)\text{Li}_2(-\theta)$ , but unfortunately new misprints are inserted. The terms (in the notation adapted in Ref. [10])

$$T_1 = -(\delta(5 - 3\delta^2)/8) \ln \theta, \quad (60a)$$

$$T_2 = +\frac{1}{4} \ln^2 \theta \left[ \frac{33 + 22\delta^2 - 7\delta^4}{24} - \delta(3 - \delta^2) \right], \quad (60b)$$

have the wrong signs and should have read

$$T_1 \rightarrow \tilde{T}_1 = +(\delta(5 - 3\delta^2)/8) \ln \theta, \quad (61a)$$

$$T_2 \rightarrow \tilde{T}_2 = -\frac{1}{4} \ln^2 \theta \left[ \frac{33 + 22\delta^2 - 7\delta^4}{24} - \delta(3 - \delta^2) \right]. \quad (61b)$$

In Ref. [38], Eq. (6) of Ref. [37] is rewritten as Eq. (140) of Ref. [38], but the two terms involving  $\ln(\theta)$ 's keep the wrong sign.

In Ref. [39], the work from Ref. [31] is being cited for the imaginary part of the proper two-loop vacuum polarization, and it is expressed in terms of the variable  $d$ , which equals our  $v$ . However, Eq. (16) of Ref. [39] contains a misprint. Namely, the term  $+3/2\ln((1+d)/(1-d))$  (in Ref. [39]) has a logarithm missing and should read  $+3/2\ln((1+d)/(1-d))\ln((1+d)/2)$ . We should clarify that, despite the typographical error in Eq. (16) of Ref. [39], final results for the energy shifts due to the two-loop vacuum polarization obtained in Ref. [39] are in agreement with those reported here.

## IV. CONTRIBUTIONS TO ENERGY SHIFTS

### A. Muonic hydrogen

Armed with the compact expressions given in Eqs. (52) and (55) for the imaginary part of the two-loop vacuum-polarization function, we can evaluate energy shifts in hydrogenlike ions. In Chap. 10 of Ref. [15], the calculation of the vacuum-polarization energy shift is outlined in detail; specifically, we refer to Eqs. (10.244) and (10.245) of Ref. [15]. (We note the different sign conventions for the sign of the imaginary part of the renormalized scalar function  $\Pi_R$ , as compared to Ref. [15].) From the one-loop and two-loop scalar functions, one infers the one-loop and two-loop irreducible vacuum-polarization potentials  $V_R^{(1)}(r)$  and  $V_R^{(2)}(r)$ , respectively,

$$V_R^{(1)}(r) = -\frac{Z\alpha}{\pi} \int_4^\infty \frac{d(q^2)}{q^2} \frac{e^{-qr}}{r} \text{Im}[\Pi_R^{(1)}(q^2 + i\epsilon)], \quad (62)$$

$$V_R^{(2)}(r) = -\frac{Z\alpha}{\pi} \int_4^\infty \frac{d(q^2)}{q^2} \frac{e^{-qr}}{r} \text{Im}[\Pi_R^{(2)}(q^2 + i\epsilon)]. \quad (63)$$

Of particular phenomenological importance for the proton radius puzzle [3] is the contribution of the irreducible two-loop vacuum-polarization effect on the  $2P$ - $2S$  energy interval in muonic hydrogen. We use the following results:

$$\langle 2S | \frac{\exp(-qr)}{r} | 2S \rangle = \frac{1 + 2(\beta q)^2}{4\beta(1 + \beta q)^4}, \quad (64a)$$

$$\langle 2P | \frac{\exp(-qr)}{r} | 2P \rangle = \frac{1}{4\beta(1 + \beta q)^4}. \quad (64b)$$

Here, the  $\beta$  parameter is

$$\beta = \frac{1}{Z\alpha\mu}, \quad (65)$$

where  $Z$  is the nuclear charge number,  $\alpha$  is the fine-structure constant, and  $\mu$  is the reduced mass of the system. For muonic hydrogen, one has  $Z = 1$ . In units where the electron mass is  $m = 1$  and the muon mass is  $m_\mu$ , one has  $\mu = m_p/(1 + m_p)$  for ordinary hydrogen and  $\mu = m_\mu m_p/(m_\mu + m_p)$  for muonic hydrogen.

The contribution of the irreducible two-loop vacuum polarization diagrams to the  $2P$ - $2S$  splitting in muonic hydrogen is obtained as

$$E^{(2)}(2P - 2S) = \left(\frac{\alpha}{\pi}\right)^2 (Z\alpha)^2 \mu f^{(2)}(\beta),$$

$$f^{(2)}(\beta) = \int_4^\infty \frac{d(q^2)}{q^2} \frac{\text{Im}\Pi_R^{(2)}(q^2 + i\epsilon)}{\alpha^2/\pi} \frac{(\beta q)^2}{2(1 + \beta q)^4}. \quad (66)$$



Quite surprisingly, the two-loop energy shift can be expressed analytically. We define

$$\beta = \frac{\sqrt{1-w^2}}{2}, \quad w = \sqrt{1-4\beta^2}, \quad (67)$$

and find the result

$$\begin{aligned} f^{(2)}(w) = & \frac{1}{288w^6} \left\{ -480(w^2-1)^2 w^6 I_1 \left( \sqrt{\frac{1-w}{1+w}} \right) \right. \\ & + 128(15w^6 - 25w^4 + 8w^2 + 3) w^3 I_2 \left( \sqrt{\frac{1-w}{1+w}} \right) \\ & + 404w^8 - 312w^6 - 176w^4 + 96w^2 \\ & - 32\pi\sqrt{1-w^2}(15w^4 - 19w^2 + 3)w^4 + 4(w^2-1) \\ & \times (41w^6 + 58w^4 - 10w^2 - 60)w \ln\left(\frac{1-w}{1+w}\right) \\ & - 16\pi^2(1-w^2)^{3/2}(15w^6 - 5w^4 - 2w^2 + 6) \\ & - (199w^{10} - 302w^8 + 51w^6 - 4w^4 + 176w^2 - 96) \\ & \left. \times \left[ \ln^2\left(\frac{1-w}{1+w}\right) + \pi^2 \right] \right\}, \quad (68) \end{aligned}$$

where the expressions  $I_j(a)$  with  $j = 1, 2$  read as follows:

$$\begin{aligned} I_1(a) = & \frac{1}{4} [6\Phi_3(-a^2) - 4\Phi_2(-a^2) \ln(a^2) \\ & + \Phi_1(-a^2) \ln^2(a^2) - 4i\pi(\text{Li}_2(ia) - \text{Li}_2(-ia)) \\ & - \pi^2 \ln(1+a^2) - 4\pi^2 \ln(1+a) + 3\zeta(3)], \quad (69) \end{aligned}$$

$$\begin{aligned} I_2(a) = & \frac{1}{16} [-4\Phi_2(-a^2) + 4\Phi_1(-a^2) \ln(a^2) \\ & + 3(\ln^2(a^2) + \pi^2) - 8\pi \arctan(a)]. \quad (70) \end{aligned}$$

Some of the terms could also be expressed in terms of the Legendre  $\chi$  function,  $\chi_\nu(z) = \frac{1}{2}[\text{Li}_\nu(z) - \text{Li}_\nu(-z)]$ .

For muonic hydrogen ( $Z = 1$ ), one obtains for the nonrelativistic energy shift due to the irreducible two-loop diagrams, using CODATA values for the relevant physical constants [40,41] (see also Ref. [27]),

$$E^{(2)}(2P - 2S)|_{\mu\text{H}} = 1.25298 \text{ meV}. \quad (71)$$

This is consistent with the literature (e.g., Ref. [2]). The result (71) confirms the numerical value of a contribution to the proton radius puzzle in muonic hydrogen whose numerical magnitude could have potentially contributed to an explanation of the discrepancy. The confirmation is obtained based on modern quantum-field theoretical methods.

The reducible diagram, according to Eqs. (D8) and (D9), leads to an energy shift of

$$E^{(r)}(2P - 2S)|_{\mu\text{H}} = 0.25495 \text{ meV}. \quad (72)$$

For selected individual low-lying atomic reference states, an analytic integration of the one-loop and two-loop energy shifts, for arbitrary  $\beta$ , still in the nonrelativistic approximation, is presented in Appendix D.

## B. Relativistic ordinary hydrogen

Given our analytic approach, we may also study the energy shifts to low-lying  $S$  and  $P$  states for ordinary (electronic) hydrogen, with a relativistic reference state. We focus on higher-order terms in the semianalytic expansion in powers and logarithms of  $Z\alpha$ , for relativistic Dirac-Coulomb reference states (see Chap. 8 of Ref. [15]).

The potential  $V_R^{(r)}$  due to the reducible diagram is

$$V_R^{(r)}(r) = \frac{Z\alpha}{\pi} \int_4^\infty \frac{d(q^2)}{q^2} \frac{e^{-qr}}{r} \text{Im} \left\{ [\Pi_R^{(1)}(q^2 + i\epsilon)]^2 \right\}. \quad (73)$$

We study the energy shifts

$$\Delta E^{(r)}(nS) = \langle \psi | V_R^{(r)} | \psi \rangle, \quad (74)$$

where  $|\psi\rangle$  is the relativistic reference-state wave function, and  $\langle \psi$  stands for the Hermitian adjoint (not for the Dirac adjoint  $\tilde{\psi} = \psi^\dagger \gamma^0$ ). The relevant semianalytic expansion is as follows:

$$\begin{aligned} \Delta E^{(r)} = & \left( \frac{\alpha}{\pi} \right)^2 \frac{(Z\alpha)^4}{n^3} \left\{ (Z\alpha) B_{50}^{(r)} + (Z\alpha)^2 B_{60}^{(r)} \right. \\ & \left. + (Z\alpha)^3 [B_{71}^{(r)} \ln[(Z\alpha)^{-2}] + B_{70}^{(r)}] \right\}. \quad (75) \end{aligned}$$

For the  $nS$  levels, the dependence on  $n$  is given, for the reducible diagram, by the following formulas (see also Table I):

$$B_{50}^{(r)}(nS) = -\frac{23\pi}{567}, \quad (76a)$$

$$B_{60}^{(r)}(nS) = \frac{29}{810} + \frac{8}{225n^2}, \quad (76b)$$

$$B_{71}^{(r)}(nS) = \frac{1}{2} B_{50}^{(r)}(nS), \quad (76c)$$

$$\begin{aligned} B_{70}^{(r)}(nS) = & -\frac{23\pi}{567} W_n - \frac{570029\pi}{7858620} - \frac{46}{567} \pi \ln(2) \\ & + \frac{1294\pi}{18711n^2}, \quad (76d) \end{aligned}$$

$$W_n = \ln\left(\frac{n}{2}\right) - \Psi(n) - \gamma_E + \frac{1}{n}. \quad (76e)$$

Here,  $\Psi(x) = \Gamma'(x)/\Gamma(x)$  is the logarithmic derivative of the Gamma function. For the  $nP_{1/2}$  levels, one has

TABLE I. Coefficients from the reducible diagram are indicated for  $S$  and  $P$  states.

	1S	2S	$2P_{1/2}$	$2P_{3/2}$
$B_{50}^{(r)}$	$-\frac{23\pi}{567}$	$-\frac{23\pi}{567}$	0	0
$B_{60}^{(r)}$	$\frac{289}{4050}$	$\frac{181}{4050}$	$\frac{2}{225}$	$\frac{2}{225}$

$B_{50}^{(r)}(nP) = B_{71}^{(r)}(nP) = 0$  and the following results for the other coefficients for the reducible diagram,

$$B_{60}^{(r)}(nP_{1/2}) = \frac{8}{675} \frac{n^2 - 1}{n^2}, \quad (77)$$

$$B_{70}^{(r)}(nP_{1/2}) = -\frac{983\pi}{93555} \frac{n^2 - 1}{n^2}. \quad (78)$$

For the  $nP_{3/2}$  levels, one has the results

$$B_{60}^{(r)}(nP_{3/2}) = B_{60}^{(r)}(nP_{1/2}), \quad (79a)$$

$$B_{70}^{(r)}(nP_{3/2}) = -\frac{127\pi}{17820} \frac{n^2 - 1}{n^2}. \quad (79b)$$

The two-loop energy shift from the irreducible diagrams, evaluated on the relativistic wave function, is

$$\Delta E^{(2)} = \langle \psi | V_R^{(2)} | \psi \rangle, \quad (80)$$

where  $V_R^{(2)}$  is given in Eq. (63). It gives rise to the following semianalytic expansion,

$$\begin{aligned} \Delta E^{(2)} = & \left(\frac{\alpha}{\pi}\right)^2 \frac{(Z\alpha)^4}{n^3} \left\{ B_{40}^{(2)} + (Z\alpha) B_{50}^{(2)} \right. \\ & + (Z\alpha)^2 \left[ B_{61}^{(2)} \ln[(Z\alpha)^{-2}] + B_{60}^{(2)} \right] \\ & \left. + (Z\alpha)^3 \left[ B_{71}^{(2)} \ln[(Z\alpha)^{-2}] + B_{70}^{(2)} \right] \right\}. \quad (81) \end{aligned}$$

For the irreducible diagrams, the dependence on  $n$  is the following (see also Table II):

$$B_{40}^{(2)}(nS) = -\frac{82}{81}, \quad (82a)$$

$$B_{50}^{(2)}(nS) = \frac{15647\pi}{13230} - \frac{25\pi^2}{63} + \frac{52}{63} \pi \ln(2), \quad (82b)$$

$$B_{60}^{(2)}(nS) = B_{40}^{(2)}(nS) W_n + \frac{77\zeta(3)}{72} - \frac{2311}{405} + \frac{1313}{675n^2}, \quad (82c)$$

$$B_{61}^{(2)}(nS) = \frac{1}{2} B_{40}^{(2)}(nS), \quad B_{71}^{(2)}(nS) = \frac{1}{2} B_{50}^{(2)}(nS). \quad (82d)$$

 TABLE II. Coefficients from the irreducible diagrams are indicated for  $S$  and  $P$  states.

	1S	2S	$2P_{1/2}$	$2P_{3/2}$
$B_{40}^{(2)}$	$-\frac{82}{81}$	$-\frac{82}{81}$	0	0
$B_{50}^{(2)}$	$\frac{15647\pi}{13230} - \frac{25\pi^2}{63} + \frac{52}{63} \pi \ln(2)$	$\frac{15647\pi}{13230} - \frac{25\pi^2}{63} + \frac{52}{63} \pi \ln(2)$	0	0
$B_{60}^{(2)}$	$-\frac{82}{81} \ln(\frac{1}{2}) + \frac{77\zeta(3)}{72} - \frac{358}{75}$	$\frac{77\zeta(3)}{72} - \frac{12727}{2700}$	$-\frac{737}{2700}$	$-\frac{449}{5400}$
$B_{61}^{(2)}$	$-\frac{82}{162}$	$-\frac{82}{162}$	0	0

The Catalan constant  $G = 0.915965\dots$  enters the  $B_{70}^{(2)}$  coefficient,

$$B_{70}^{(2)}(nS) = B_{50}^{(2)} W_n + B_{70}^{(a)} + \frac{B_{70}^{(b)}}{n^2},$$

$$\begin{aligned} B_{70}^{(a)} = & \frac{104}{63} \pi \ln^2(2) - \frac{25}{63} \pi^2 \ln(2) - \frac{200\pi}{63} G + \frac{13\pi^3}{189} \\ & + \frac{79318\pi \ln(2)}{19845} - \frac{82361\pi^2}{79380} + \frac{89515961\pi}{20003760}, \quad (82e) \end{aligned}$$

$$B_{70}^{(b)} = -\frac{3304831\pi}{1428840} + \frac{1891\pi^2}{2268} - \frac{1076}{567} \pi \ln(2). \quad (82f)$$

For the  $nP_{1/2}$  levels, the coefficients are

$$B_{60}^{(2)}(nP_{1/2}) = -\frac{737}{2025} \frac{n^2 - 1}{n^2}, \quad (83a)$$

$$B_{70}^{(2)}(nP_{1/2}) = \pi \left[ \frac{1764797}{7144200} - \frac{809\pi}{11340} + \frac{328 \ln(2)}{2835} \right] \frac{n^2 - 1}{n^2}. \quad (83b)$$

For the  $nP_{3/2}$  levels, the coefficients are

$$B_{60}^{(2)}(nP_{3/2}) = -\frac{449}{4050} \frac{n^2 - 1}{n^2}, \quad (84a)$$

$$B_{70}^{(2)}(nP_{3/2}) = \pi \left[ \frac{75763}{510300} - \frac{31\pi}{810} + \frac{19 \ln(2)}{405} \right] \frac{n^2 - 1}{n^2}. \quad (84b)$$

The fine-structure difference

$$B_{60}^{(2)}(nP_{3/2}) - B_{60}^{(2)}(nP_{1/2}) = \frac{41}{162} \frac{n^2 - 1}{n^2} \quad (85)$$

is consistent with the result communicated in Eq. (7.7) of Ref. [42].

## V. CONCLUSIONS

We have completed a complete reevaluation of the two-loop vacuum-polarization tensor in dimensional regularization, on the basis of integration-by-parts identities. The two-loop vacuum-polarization tensor constitutes a numerically significant contribution to the Lamb shift of muonic hydrogen which influences the determination of the proton radius from muonic hydrogen spectroscopy [2].

In Sec. II, we have discussed the evaluation of the one-loop vacuum polarization insertion into the photon propagator, evaluated to order  $\epsilon$ , and  $\epsilon^2$ , and thus, in a form suitable for inclusion into higher-order loop calculation where knowledge of the terms of higher orders in  $\epsilon$  is indispensable. We note that we use somewhat nonstandard conventions for the  $\overline{MS}$  charge, as given in Eq. (12).

In Sec. III, the irreducible two-loop vacuum polarization insertion has been evaluated, by expressing it in terms of master integrals. The renormalization has been carried out, and final results have been presented for the real and imaginary parts, in Eqs. (52) and (55). A comparison to the existing literature is being performed as well (Sec. III G).

In Sec. IV and Appendix D, we demonstrate that, for arbitrary reduced mass of a two-body bound system, the two-loop vacuum-polarization corrections to the energy can be evaluated analytically (for nonrelativistic reference states) and expressed in terms of dilogarithmic, and trilogarithmic, functions. This applies both to the  $2P$ - $2S$  difference (see Sec. IVA) as well as to individual hydrogenic levels (see Appendix D). Higher-order coefficients for the semianalytic expansion of the two-loop vacuum-polarization energy shift could be evaluated with the help of the fully relativistic hydrogen wave function. Results have been presented in Sec. IV B.

## ACKNOWLEDGMENTS

The authors acknowledge insightful conversations with Professor Ettore Remiddi. Support from the National Science Foundation (Grant No. PHY-2110294) is gratefully acknowledged. S. L. also acknowledges support from Italian Ministry of University and Research (MUR) via the PRIN 2022 Project No. 20225X52RA-MUS4GM2 funded by the European Union via the Next Generation EU package.

## APPENDIX A: TERMS OF ORDER $\epsilon^2$

We first give the terms quadratic in  $\epsilon$  for Eq. (20), for the master integral  $M_{12}$ ,

$$N_{12}(v) = -12\text{Li}_3\left(\frac{1-v}{v+1}\right) - 24\text{Li}_3\left(\frac{2v}{v+1}\right) + \ln^3\left(\frac{v-1}{v+1}\right) - 6\ln\left(\frac{2v}{v+1}\right)\ln^2\left(\frac{v-1}{v+1}\right) + 12\ln^2\left(\frac{2v}{v+1}\right)\left(\ln\left(\frac{v-1}{v+1}\right) - \ln\left(-\frac{v-1}{v+1}\right)\right) - \pi^2\ln\left(\frac{v-1}{v+1}\right) + 6\pi^2\ln\left(\frac{2v}{v+1}\right) + 12\zeta(3). \quad (\text{A1})$$

We now give the expressions for the terms of order  $\epsilon^2$  in the one-loop effect, as discussed in Eq. (22). For the quadratic term  $Q_2^{(1)}(v)$  in Eq. (22), one obtains the result

$$Q_2^{(1)}(v) = \left(v - \frac{4v^3}{9}\right) \left[ -2\text{Li}_2\left(\frac{1-v}{v+1}\right) + \frac{1}{2}\ln^2\left(\frac{v-1}{v+1}\right) - 2\ln\left(\frac{2v}{v+1}\right)\ln\left(\frac{v-1}{v+1}\right) - \frac{\pi^2}{6} \right] + \frac{1}{36}v(v^2-3) \left[ 12\text{Li}_3\left(\frac{1-v}{v+1}\right) + 24\text{Li}_3\left(\frac{2v}{v+1}\right) - \ln^3\left(\frac{v-1}{v+1}\right) + 6\ln\left(\frac{2v}{v+1}\right)\ln^2\left(\frac{v-1}{v+1}\right) + 12\left(\ln\left(\frac{1-v}{v+1}\right) - \ln\left(\frac{v-1}{v+1}\right)\right)\ln^2\left(\frac{2v}{v+1}\right) + \pi^2\ln\left(\frac{v-1}{v+1}\right) - 6\pi^2\ln\left(\frac{2v}{v+1}\right) - 12\zeta(3) \right] - \frac{2}{81} \left( (39v^3 - 81v)\ln\left(\frac{v-1}{v+1}\right) + 78v^2 - 160 \right). \quad (\text{A2})$$

For the term  $Q_2^{(1a)}(v)$  from the integral with two fermion propagators, one finds for the term quadratic in  $\epsilon$  the expression

$$Q_2^{(1a)}(v) = \frac{v^2-1}{4v} \left\{ -\frac{1}{6}(v^2-1) \left[ 12\text{Li}_3\left(\frac{1-v}{v+1}\right) + 24\text{Li}_3\left(\frac{2v}{v+1}\right) - \ln^3\left(\frac{v-1}{v+1}\right) + 6\ln\left(\frac{2v}{v+1}\right)\left(\ln^2\left(\frac{v-1}{v+1}\right) - \pi^2\right) - 24\ln^2\left(\frac{2v}{v+1}\right)\ln\left(\frac{v-1}{v+1}\right) + 12\left(\ln\left(\frac{1-v}{v+1}\right) + \ln\left(\frac{v-1}{v+1}\right)\right)\ln^2\left(\frac{2v}{v+1}\right) + \pi^2\ln\left(\frac{v-1}{v+1}\right) - 12\zeta(3) \right] + v^2 \left[ -4\text{Li}_2\left(\frac{1-v}{v+1}\right) + \ln^2\left(\frac{v-1}{v+1}\right) - 4\left(\ln\left(\frac{2v}{v+1}\right) - 1\right)\ln\left(\frac{v-1}{v+1}\right) - \frac{\pi^2}{3} + 8v \right] \right\}. \quad (\text{A3})$$

## APPENDIX B: COEFFICIENT FUNCTIONS

The coefficients in Eq. (36) are as follows:

$$\mathcal{P}_{234}^{(2;1)} = -\frac{(2-D)(D^2(-q^2-4) - D(q^4 - 5q^2 - 28) + 2(q^4 - 2q^2 - 20))}{2(D-4)(D-3)(D-1)q^2(4-q^2)}, \quad (\text{B1})$$

$$\mathcal{P}_{24}^{(2;1)} = -\frac{(D-2)(3D^2q^4 - 4D(3q^4 - 2q^2 + 8) + 4(3q^4 - 8q^2 + 20))}{2(D-4)(D-3)(D-1)q^4(4-q^2)^2}, \quad (\text{B2})$$

$$\mathcal{P}_{1234}^{(2;1)} = \frac{-(D^3 - 9D^2 + 30D - 32)q^4 + 4(D^3 - 11D^2 + 38D - 40)q^2 + 32}{8(D-4)(D-1)q^2(4-q^2)}, \quad (\text{B3})$$

$$\mathcal{P}_{235k}^{(2;1)} = -\frac{1}{(D-4)(D-3)(D-1)q^4(4-q^2)^2} [3(-4D^3q^2(1-q^2) - D^2(-q^6 + 30q^4 - 36q^2 + 16) + 4D(-q^6 + 18q^4 - 28q^2 + 20) - 4(-q^6 + 14q^4 - 28q^2 + 24))], \quad (\text{B4})$$

$$\mathcal{P}_{235}^{(2;1)} = -\frac{1}{4(D-4)(D-3)(D-1)q^4(4-q^2)^2} [-D^4q^2(4-q^2)^2 + D^3q^2(23q^4 - 100q^2 + 176) + 4D^2(q^8 - 35q^6 + 119q^4 - 176q^2 - 32) - 8D(2q^8 - 40q^6 + 127q^4 - 148q^2 - 88) + 8(2q^8 - 31q^6 + 98q^4 - 84q^2 - 120)]. \quad (\text{B5})$$

The coefficients in Eq. (37) are

$$\mathcal{G}_{235k}^{(2;1)} = -\frac{3(D-2)(-D^2(-q^2-2)q^2 - D(4q^4 + 6q^2 + 8) + 4(q^4 - q^2 + 6))}{(D-4)(D-1)q^4(4-q^2)^2}, \quad (\text{B6})$$

$$\mathcal{G}_{24}^{(2;1)} = \frac{(2-D)(3D^2q^4 - 4D(3q^4 - 2q^2 + 8) + 4(3q^4 - 8q^2 + 20))}{4(D-4)(D-1)q^4(4-q^2)^2}, \quad (\text{B7})$$

$$\mathcal{G}_{235}^{(2;1)} = \frac{1}{4(D-4)(D-1)q^4(4-q^2)^2} [-3D^3(q^2+2)q^4 + D^2(18q^6 + 26q^4 + 64) + 4D(-9q^6 - 4q^4 + 8q^2 - 88) + 8(3q^6 - 2q^4 - 10q^2 + 60)]. \quad (\text{B8})$$

The coefficients in Eq. (38) are

$$\mathcal{P}_{234}^{(2;2)} = \frac{(D-2)(-2D^2q^2 - D(q^4 - 14q^2 + 8) + 2(q^4 - 10q^2 + 12))}{16(D-3)q^2(4-q^2)}, \quad (\text{B9})$$

$$\mathcal{P}_{235k}^{(2;2)} = \frac{3(D-2)}{8(D-4)(D-3)(D-1)q^4(4-q^2)^2} [-D^3q^2(4-q^2)^2 + D^2q^2(7q^4 - 48q^2 + 128) + 2D(-7q^6 + 40q^4 - 136q^2 - 32) - 8(-q^6 + 4q^4 - 12q^2 - 24)], \quad (\text{B10})$$

$$\mathcal{P}_{235}^{(2;2)} = -\frac{1}{8(D-4)(D-3)(D-1)q^4(q^2-4)^2} [D^4q^2(q^2-4)^2(q^2+1) - 3D^3q^2(3q^6 - 19q^4 + 28q^2 + 48) + 4D^2(7q^8 - 40q^6 + 69q^4 + 112q^2 + 32) - 4D(9q^8 - 45q^6 + 80q^4 + 128q^2 + 176) + 16(q^8 - 4q^6 + 6q^4 + 6q^2 + 60)], \quad (\text{B11})$$

$$\mathcal{P}_{24}^{(2;2)} = \frac{D-2}{4(D-4)(D^2-4D+3)q^4(q^2-4)^2} [D^4q^2(q^2-4) - 8D^3q^2(q^2-4) + 12D^2q^2(2q^2-7) + D(-34q^4 + 96q^2 - 32) + 20q^4 - 64q^2 + 80]. \quad (\text{B12})$$

**APPENDIX C: HIGHER-ORDER TERMS IN THE MASTER INTEGRALS**

We have the following results for the higher-order terms from the master integrals, with reference to Eqs. (40) and (41). For  $N_{235}(q^2)$ , we obtain

$$\begin{aligned}
 N_{235}(q^2) = & \frac{1}{6}v(v^2 - 3) \left[ 18\Phi_2\left(\frac{v-1}{v+1}\right) - 18\ln\left(\frac{v-1}{v+1}\right)\Phi_1\left(\frac{v-1}{v+1}\right) - 6\text{Li}_2\left(\frac{v-1}{v+1}\right) - 6\ln\left(\frac{2}{v+1}\right)\ln\left(\frac{v-1}{v+1}\right) \right. \\
 & - 39\ln\left(\frac{v-1}{v+1}\right) + \pi^2 \left. \right] + \frac{1}{12}(v^4 + 2v^2 - 3) \left[ 36\Phi_3\left(\frac{v-1}{v+1}\right) - 18\ln\left(\frac{v-1}{v+1}\right)\Phi_2\left(\frac{v-1}{v+1}\right) \right. \\
 & - 6\text{Li}_2\left(\frac{v-1}{v+1}\right)\ln\left(\frac{v-1}{v+1}\right) - 3\ln^3\left(\frac{v-1}{v+1}\right) - 6\ln\left(\frac{2}{v+1}\right)\ln^2\left(\frac{v-1}{v+1}\right) + \pi^2\ln\left(\frac{v-1}{v+1}\right) + 18\zeta(3) \left. \right] \\
 & + \frac{1}{8} \left( 60v^2 + (-5v^4 - 16v^3 - 14v^2 + 48v + 3)\ln^2\left(\frac{v-1}{v+1}\right) + 170 \right). \tag{C1}
 \end{aligned}$$

For  $N_{235k}(q^2)$ , we obtain

$$\begin{aligned}
 N_{235k}(v) = & \frac{1}{36}v(3v^4 - 8v^2 + 21) \left[ 18\Phi_2\left(\frac{v-1}{v+1}\right) - 18\ln\left(\frac{v-1}{v+1}\right)\Phi_1\left(\frac{v-1}{v+1}\right) - 6\text{Li}_2\left(\frac{v-1}{v+1}\right) \right. \\
 & - 6\ln\left(\frac{2}{v+1}\right)\ln\left(\frac{v-1}{v+1}\right) + \pi^2 \left. \right] + \frac{1}{24}(v^6 - 3v^4 - 5v^2 + 7) \left[ 36\Phi_3\left(\frac{v-1}{v+1}\right) - 18\ln\left(\frac{v-1}{v+1}\right)\Phi_2\left(\frac{v-1}{v+1}\right) \right. \\
 & - 6\text{Li}_2\left(\frac{v-1}{v+1}\right)\ln\left(\frac{v-1}{v+1}\right) - 3\ln^3\left(\frac{v-1}{v+1}\right) - 6\ln\left(\frac{2}{v+1}\right)\ln^2\left(\frac{v-1}{v+1}\right) + \pi^2\ln\left(\frac{v-1}{v+1}\right) + 18\zeta(3) \left. \right] \\
 & + \frac{1}{48} \left[ -2(86v^4 + 233v^2 + 601) - 4v(37v^4 - 108v^2 + 279)\ln\left(\frac{v-1}{v+1}\right) \right. \\
 & \left. + (-13v^6 - 48v^5 + 63v^4 + 128v^3 + 105v^2 - 336v - 27)\ln^2\left(\frac{v-1}{v+1}\right) \right]. \tag{C2}
 \end{aligned}$$

**APPENDIX D: ANALYTIC INTEGRATIONS**
**1. One-loop diagram**

We recall the imaginary part of the one-loop effect from Eq. (28),

$$\text{Im}[\Pi_{\text{R}}^{(1)}(q^2 + i\epsilon)] = \frac{\alpha}{3} \sqrt{1 - \frac{4}{q^2}} \left( 1 + \frac{2}{q^2} \right), \tag{D1}$$

for  $q^2 \geq 4$ . We use Eq. (10.245) of Ref. [15] and write

$$V_{\text{R}}^{(1)}(r) = -\frac{Z\alpha}{\pi} \int_4^\infty \frac{d(q^2)}{q^2} \frac{e^{-qr}}{r} \text{Im}[\Pi_{\text{R}}^{(1)}(q^2 + i\epsilon)]. \tag{D2}$$

One uses the well-known formula for the generalized Bohr radius in a hydrogenlike ion with nuclear charge number  $Z$ , namely,  $\beta \equiv a_0 = 1/(Z\alpha\mu)$ , where  $\mu$  is the reduced mass of the two-body system, measured in units of the electron

mass. For the  $2S$  and  $2P$  states of atomic hydrogen, one finds, respectively,

$$\langle 2S | \frac{\exp(-qr)}{r} | 2S \rangle = \frac{1 + 2(\beta q)^2}{4\beta(1 + \beta q)^4}, \tag{D3a}$$

$$\langle 2P | \frac{\exp(-qr)}{r} | 2P \rangle = \frac{1}{4\beta(1 + \beta q)^4}. \tag{D3b}$$

Hence,

$$\begin{aligned}
 & \langle 2P | \frac{\exp(-qr)}{r} | 2P \rangle - \langle 2S | \frac{\exp(-qr)}{r} | 2S \rangle \\
 & = -\frac{(\beta q)^2}{2\beta(1 + \beta q)^4}. \tag{D4}
 \end{aligned}$$

The contribution to the  $2P - 2S$  energy shift is

$$\begin{aligned}
 E_1(2P-2S) &= \langle 2P | V_R^{(1)} | 2P \rangle - \langle 2S | V_R^{(1)} | 2S \rangle \\
 &= \frac{(Z\alpha)^2 \mu}{2\pi} \int_4^\infty \frac{d(q^2)}{q^2} \text{Im}[\Pi_R^{(1)}(q^2 + i\epsilon)] \frac{(\beta q)^2}{(1 + \beta q)^4} \\
 &= \left(\frac{\alpha}{\pi}\right) (Z\alpha)^2 \mu f_1(\beta), \\
 f_1(\beta) &= \int_4^\infty \frac{d(q^2)}{q^2} \frac{\text{Im}\Pi_R^{(1)}(q^2 + i\epsilon)}{\alpha} \frac{(\beta q)^2}{2(1 + \beta q)^4}.
 \end{aligned} \tag{D5}$$

We choose the  $2P-2S$  interval in order to demonstrate that energy shifts due to the one-loop and two-loop vacuum-polarization effects can be evaluated analytically. For muonic hydrogen, one has the result  $\mu = m_\mu m_p / (m_\mu + m_p)$ , where  $m_\mu$  and  $m_p$  are the muon and proton masses, respectively.

It turns out to be advantageous to express the result in terms of the variable  $w$  defined in Eq. (67), with the result

$$\begin{aligned}
 f_1(w) &= \frac{(w^2 - 1)^2}{72w^5} \left[ 3(8w^4 + 4w^2 + 3) \ln\left(\frac{1-w}{1+w}\right) \right. \\
 &\quad \left. + \frac{24\pi w^5}{\sqrt{1-w^2}} + \frac{2(24w^6 - 28w^4 - 3w^2 + 9)w}{(w^2 - 1)^2} \right].
 \end{aligned} \tag{D6}$$

We can get the expansions for small and large  $\beta$ ,

$$\begin{aligned}
 f_1(\beta) &= \frac{1}{18} - \beta^2 + \frac{8\pi\beta^3}{3} + \beta^4 \left( 10 \ln(\beta^2) + \frac{32}{3} \right) \\
 &\quad + \beta^6 \left( \frac{140 \ln(\beta^2)}{3} + \frac{236}{3} \right) + \mathcal{O}(\beta^8),
 \end{aligned} \tag{D7a}$$

$$\begin{aligned}
 f_1(\beta) &= \frac{1}{30\beta^2} - \frac{5\pi}{384\beta^3} + \frac{1}{28\beta^4} - \frac{35\pi}{4096\beta^5} + \frac{1}{54\beta^6} \\
 &\quad - \frac{63\pi}{16384\beta^7} + \mathcal{O}\left(\frac{1}{\beta^8}\right).
 \end{aligned} \tag{D7b}$$

For muonic hydrogen, one has  $\beta = 0.737384\dots$ , while, for ordinary hydrogen, one has  $\beta = 137.110\dots$ . Hence, the second of the above equations is relevant for ordinary hydrogen, while none of the expansions can be used with good accuracy for muonic hydrogen. It should be possible, though, to generalize the results reported in this appendix to reference states other than  $2S$  and  $2P$  if desired. Relativistic corrections to the one-loop vacuum-polarization shift have been analyzed in Ref. [43], with an emphasis on muonic hydrogen.

## 2. Two-loop reducible diagram

Next, we consider the contribution of the reducible diagram 3 in Fig. 1,

$$\begin{aligned}
 E^{(r)}(2P-2S) &= \left(\frac{\alpha}{\pi}\right)^2 (Z\alpha)^2 \mu f^{(r)}(\beta), \\
 f^{(r)}(\beta) &= - \int_4^\infty \frac{d(q^2)}{q^2} \frac{\text{Im}[(\Pi_R^{(1)}(q^2 + i\epsilon))^2]}{\alpha^2/\pi} \frac{(\beta q)^2}{2(1 + \beta q)^4}.
 \end{aligned} \tag{D8}$$

We use the conventions outlined in Eq. (67). An analytic evaluation leads to the formula

$$\begin{aligned}
 f^{(r)}(w) &= -\frac{1}{648w^5} \left\{ 4w(-315w^8 + 1005w^6 - 836w^4 \right. \\
 &\quad \left. + 84w^2 + 72) + 12(-105w^{10} + 370w^8 - 381w^6 \right. \\
 &\quad \left. + 96w^4 + 10w^2 + 12) \ln\left(\frac{1-w}{1+w}\right) \right. \\
 &\quad \left. - 45(w^2 - 1)^2(7w^2 - 13)w^5 \left[ \ln^2\left(\frac{1-w}{1+w}\right) + \pi^2 \right] \right\}.
 \end{aligned} \tag{D9}$$

The expansion for small  $\beta$  reads as

$$\begin{aligned}
 f^{(r)}(\beta) &= -\frac{\ln(\beta^2)}{27} - \frac{5}{81} + \beta^2 \left( \frac{2}{3} \ln(\beta^2) + \frac{32}{9} \right) \\
 &\quad - \frac{\beta^4}{27} (180 \ln^2(\beta^2) + 774 \ln(\beta^2) + 180\pi^2 \\
 &\quad - 73) + \mathcal{O}(\beta^6).
 \end{aligned} \tag{D10}$$

For large  $\beta$ , one obtains the result

$$\begin{aligned}
 f^{(r)}(\beta) &= \frac{23\pi}{4536\beta^3} - \frac{1}{45\beta^4} + \frac{127\pi}{19008\beta^5} - \frac{1}{60\beta^6} \\
 &\quad + \frac{1675\pi}{439296\beta^7} + \mathcal{O}\left(\frac{1}{\beta^8}\right).
 \end{aligned} \tag{D11}$$

## 3. Two-loop irreducible diagram

We continue with the contribution of the irreducible two-loop vacuum polarization diagrams,

$$\begin{aligned}
 E_2(2P-2S) &= \left(\frac{\alpha}{\pi}\right)^2 (Z\alpha)^2 \mu f_2(\beta), \\
 f_2(\beta) &= \int_4^\infty \frac{d(q^2)}{q^2} \frac{\text{Im}\Pi_R^{(2)}(q^2 + i\epsilon)}{\alpha^2/\pi} \frac{(\beta q)^2}{2(1 + \beta q)^4}.
 \end{aligned} \tag{D12}$$

The result is a bit more complex and given in Eq. (68). The expansion for small  $\beta$  is

$$\begin{aligned}
 f_2(\beta) &= \frac{1}{24} - \beta^2 \left( 3 \ln(\beta) + \frac{11}{2} \right) + \beta^3 \left( \frac{320\pi}{27} - \frac{56\pi^2}{9} \right) \\
 &+ \beta^4 \left( 60 \ln^2(\beta) + 69 \ln(\beta) - 20\zeta(3) + 15\pi^2 + \frac{43}{12} \right) \\
 &+ \mathcal{O}(\beta^5). \tag{D13}
 \end{aligned}$$

For large  $\beta$ , one finds the result

$$\begin{aligned}
 f_2(\beta) &= \frac{41}{324\beta^2} + \frac{1}{\beta^3} \left( -\frac{15647\pi}{105840} + \frac{25\pi^2}{504} - \frac{13}{126}\pi \ln(2) \right) \\
 &+ \frac{449}{2160\beta^4} + \frac{1}{\beta^5} \left( -\frac{75763\pi}{544320} + \frac{31\pi^2}{864} - \frac{19}{432}\pi \ln(2) \right) \\
 &+ \frac{62479}{453600\beta^6} + \frac{1}{\beta^7} \left[ -\frac{71317517\pi}{936714240} + \frac{37\pi^2}{2112} \right. \\
 &\left. - \frac{125}{8448}\pi \ln(2) \right] + \mathcal{O}\left(\frac{1}{\beta^8}\right). \tag{D14}
 \end{aligned}$$

#### 4. Individual energy shifts

In Sec. IV A, we had concentrated on the  $2P$ - $2S$  energy difference, for muonic hydrogen. It is instructive to separately give the nonrelativistic contributions to selected energy levels with principal quantum number  $n$  and orbital angular momentum quantum number  $\ell$  (in the nonrelativistic approximation) from the one-loop diagram, the two-loop irreducible and the two-loop reducible diagrams,

$$E^{(1)}(n\ell) = \left(\frac{\alpha}{\pi}\right) (Z\alpha)^2 \mu f_{n\ell}^{(1)}(\beta), \tag{D15a}$$

$$E^{(r)}(n\ell) = \left(\frac{\alpha}{\pi}\right)^2 (Z\alpha)^2 \mu f_{n\ell}^{(r)}(\beta), \tag{D15b}$$

$$E^{(2)}(n\ell) = \left(\frac{\alpha}{\pi}\right)^2 (Z\alpha)^2 \mu f_{n\ell}^{(2)}(\beta). \tag{D15c}$$

We consider the cases  $|n\ell\rangle = |1S\rangle, |2S\rangle, |2P\rangle$ . The variable  $w$  has been defined in Eq. (67). We define the variable  $u$  as follows:

$$u = \sqrt{1 - \beta^2}, \quad \beta = \sqrt{1 - u^2}. \tag{D16}$$

For the ground state, we obtain

$$\begin{aligned}
 f_{1S}^{(1)}(\beta) &= \frac{1}{18u} \left[ 3\pi\sqrt{1-u^2}(4u^2-7)u - 24u^3 + 46u \right. \\
 &\left. - 3(4u^4 - 9u^2 + 3) \ln\left(\frac{1-u}{1+u}\right) \right]. \tag{D17}
 \end{aligned}$$

For the  $2S$  state, one has the result

$$\begin{aligned}
 f_{2S}^{(1)}(\beta) &= \frac{1}{144w^5} \left[ -336w^7 + 464w^5 - 18w^3 - 54w \right. \\
 &+ 12\pi w^5(14w^2 - 17)\sqrt{1-w^2} \\
 &\left. - 3(56w^8 - 96w^6 + 27w^4 + 9) \ln\left(\frac{1-w}{1+w}\right) \right]. \tag{D18}
 \end{aligned}$$

For the  $2P$  level, we obtain

$$\begin{aligned}
 f_{2P}^{(1)}(\beta) &= \frac{1}{144w^5} \left[ -240w^7 + 352w^5 - 30w^3 - 18w \right. \\
 &+ 12\pi w^5(10w^2 - 13)\sqrt{1-w^2} \\
 &- 3(40w^8 - 72w^6 + 21w^4 + 4w^2 + 3) \\
 &\left. \times \ln\left(\frac{1-w}{1+w}\right) \right]. \tag{D19}
 \end{aligned}$$

For the reducible diagram, the results are as follows. Let us first indicate the result for the ground state,

$$\begin{aligned}
 f_{1S}^{(r)}(\beta) &= -\frac{1}{1620u} \left[ -288\pi u\sqrt{1-u^2} \right. \\
 &+ 20u(63u^4 - 282u^2 + 304) \\
 &+ 45u(7u^6 - 36u^4 + 51u^2 - 18) \left[ \ln^2\left(\frac{1-u}{1+u}\right) + \pi^2 \right] \\
 &\left. + 60(21u^6 - 101u^4 + 126u^2 - 24) \ln\left(\frac{1-u}{1+u}\right) \right]. \tag{D20}
 \end{aligned}$$

For the  $2S$  state, one has the result

$$\begin{aligned}
 f_{2S}^{(r)}(\beta) &= -\frac{1}{6480w^5} \left[ -576\pi w^5\sqrt{1-w^2} \right. \\
 &+ 20(1386w^8 - 4719w^6 + 4132w^4 - 396w^2 - 216)w \\
 &+ 45w^5(154w^6 - 627w^4 + 792w^2 - 315) \\
 &\times \left[ \ln^2\left(\frac{1-w}{1+w}\right) + \pi^2 \right] + 60(462w^{10} - 1727w^8 \\
 &\left. + 1851w^6 - 468w^4 - 54w^2 - 36) \ln\left(\frac{1-w}{1+w}\right) \right], \tag{D21}
 \end{aligned}$$

while the  $2P$  result is

$$\begin{aligned}
f_{2P}^{(r)}(\beta) = & -\frac{1}{6480w^5} \left[ -576\pi w^5 \sqrt{1-w^2} + 60(252w^8 - 903w^6 + 820w^4 - 76w^2 - 24)w \right. \\
& + 45w^5(84w^6 - 357w^4 + 462w^2 - 185) \left[ \ln^2\left(\frac{1-w}{1+w}\right) + \pi^2 \right] + 60(252w^{10} - 987w^8 + 1089w^6 \\
& \left. - 276w^4 - 34w^2 - 12) \ln\left(\frac{1-w}{1+w}\right) \right]. \tag{D22}
\end{aligned}$$

The difference of the results given in Eqs. (D21) and (D22) confirms Eq. (D9).

The results for the irreducible two-loop diagram are more complicated and read as follows [we refer to the definitions of  $I_1$  and  $I_2$  in Eqs. (69) and (70)]. For the ground state, one obtains

$$\begin{aligned}
f_{1S}^{(2)}(\beta) = & \frac{1}{96u^2} \left[ 128(11 - 5u^2)u^3 I_2 \left( \sqrt{\frac{1-u}{1+u}} \right) + (43u^6 - 94u^4 - 45u^2 - 48) \left( \ln^2\left(\frac{1-u}{1+u}\right) + \pi^2 \right) \right. \\
& + 4(43 - 37u^2)u^3 \ln\left(\frac{1-u}{1+u}\right) + 4u^2 \left( 8(5u^4 - 10u^2 + 1) I_1 \left( \sqrt{\frac{1-u}{1+u}} \right) - 57u^2 + 74 \right) \\
& \left. + \sqrt{1-u^2} \left[ \pi \left( \frac{32}{9}(45u^2 - 113)u^2 + \frac{1024}{3}u^2 \ln(2) \right) + \pi^2 \left( -80u^4 + \frac{320u^2}{3} + 48 \right) \right] \right]. \tag{D23}
\end{aligned}$$

For the metastable  $2S$  level, the analytic integration leads to the result

$$\begin{aligned}
f_{2S}^{(2)}(\beta) = & \frac{1}{1152w^6} \left[ 3 \left( (653w^{10} - 1058w^8 + 93w^6 + 56w^4 + 256w^2 - 192) \left( \ln^2\left(\frac{1-w}{1+w}\right) + \pi^2 \right) \right. \right. \\
& \left. \left. - 4w^2(447w^6 - 430w^4 - 88w^2 + 48) \right) - 384(55w^6 - 97w^4 + 28w^2 + 6)w^3 I_2 \left( \sqrt{\frac{1-w}{1+w}} \right) \right. \\
& + 96(55w^4 - 110w^2 + 51)w^6 I_1 \left( \sqrt{\frac{1-w}{1+w}} \right) + 12(-227w^8 + 77w^6 + 224w^4 + 52w^2 - 120)w \\
& \times \ln\left(\frac{1-w}{1+w}\right) + \sqrt{1-w^2} \left( \pi(2048w^6 \ln(2) + \frac{32}{3}(495w^4 - 757w^2 + 54)w^4) - 16\pi^2 \right. \\
& \left. \left. \times (165w^8 - 220w^6 + 21w^4 + 30w^2 - 36) \right) \right]. \tag{D24}
\end{aligned}$$

Finally, the irreducible two-loop diagram, for the  $2P$  level, leads to the energy shift

$$\begin{aligned}
f_{2P}^{(2)}(\beta) = & \frac{1}{1152w^6} \left[ (1163w^{10} - 1966w^8 + 75w^6 + 184w^4 + 64w^2 - 192) \left( \ln^2\left(\frac{1-w}{1+w}\right) + \pi^2 \right) \right. \\
& - 128(105w^6 - 191w^4 + 52w^2 + 6)w^3 I_2 \left( \sqrt{\frac{1-w}{1+w}} \right) + 4w^2(24(35w^4 - 70w^2 + 31)w^4 I_1 \left( \sqrt{\frac{1-w}{1+w}} \right) \\
& - 937w^6 + 978w^4 + 88w^2 - 48) - 4(517w^8 - 299w^6 - 400w^4 + 44w^2 + 120)w \\
& \times \ln\left(\frac{1-w}{1+w}\right) + \sqrt{1-w^2} \left( \pi(2048w^6 \ln(2) + \frac{32}{3}(315w^4 - 529w^2 + 18)w^4) \right. \\
& \left. \left. - 16\pi^2(105w^8 - 140w^6 + 9w^4 - 2w^2 - 12) \right) \right]. \tag{D25}
\end{aligned}$$

The difference of the  $2P$  and  $2S$  energy shifts confirms Eq. (71) for muonic hydrogen.



- [1] K. Pachucki, Theory of the Lamb shift in muonic hydrogen, *Phys. Rev. A* **53**, 2092 (1996).
- [2] U. D. Jentschura, Lamb shift in muonic hydrogen. I. Verification and update of theoretical predictions, *Ann. Phys. (N.Y.)* **326**, 500 (2011).
- [3] U. D. Jentschura, Proton radius: A puzzle or a solution!?, *J. Phys. Conf. Ser.* **2391**, 012017 (2022).
- [4] G. Pustovalov, Vacuum polarization in mesonic atoms, *Zh. Éksp. Teor. Fiz.* **32**, 1519 (1957) [*Sov. Phys. JETP* **5**, 1234 (1957)].
- [5] U. D. Jentschura, G. Soff, V. G. Ivanov, and S. G. Karshenboim, Bound  $\mu^+\mu^-$  system, *Phys. Rev. A* **56**, 4483 (1997).
- [6] S. G. Karshenboim, V. G. Ivanov, U. D. Jentschura, and G. Soff, Bound states of the muon-antimuon system: Lifetimes and hyperfine splitting, *Zh. Eksp. Teor. Fiz.* **113**, 409 (1998) [*Sov. Phys. JETP* **86**, 226 (1998)].
- [7] S. G. Karshenboim, U. D. Jentschura, V. G. Ivanov, and G. Soff, Next-to-leading and higher-order corrections to the decay rate of dimuonium, *Phys. Lett. B* **424**, 397 (1998).
- [8] G. 't Hooft and M. Veltman, Regularization and renormalization of gauge fields, *Nucl. Phys.* **B44**, 189 (1972).
- [9] C. G. Bollini and J. J. Giambiagi, Dimensional renormalization: The number of dimensions is a regularizing parameter, *Nuovo Cimento* **12B**, 20 (1972).
- [10] R. Barbieri and E. Remiddi, Infra-red divergences and adiabatic switching. Fourth-order vacuum polarization, *Nuovo Cimento A* **13**, 99 (1973).
- [11] J. D. Bjorken and S. D. Drell, *Relativistic Quantum Fields* (McGraw-Hill, New York, 1965).
- [12] M. E. Peskin and D. V. Schroeder, *An Introduction to Quantum Field Theory* (Perseus, Cambridge, MA, 1995).
- [13] C. Itzykson and J.-B. Zuber, *Quantum Field Theory* (McGraw-Hill, New York, 1980).
- [14] V. B. Berestetskii, E. M. Lifshitz, and L. P. Pitaevskii, *Quantum Electrodynamics, Volume 4 of the Course on Theoretical Physics*, 2nd ed. (Pergamon Press, Oxford, UK, 1982).
- [15] U. D. Jentschura and G. S. Adkins, *Quantum Electrodynamics: Atoms, Lasers and Gravity* (World Scientific, Singapore, 2022).
- [16] J. A. M. Vermaseren, New features of FORM, [arXiv:math-ph/0010025](https://arxiv.org/abs/math-ph/0010025).
- [17] J. Kuipers, T. Ueda, J. A. M. Vermaseren, and J. Vollinga, FORM version 4.0, *Comput. Phys. Commun.* **184**, 1453 (2013).
- [18] K. G. Chetyrkin and F. V. Tkachov, Integration by parts: The algorithm to calculate beta functions in 4 loops, *Nucl. Phys.* **B192**, 159 (1981).
- [19] F. V. Tkachov, A theorem on analytical calculability of four loop renormalization group functions, *Phys. Lett.* **100B**, 65 (1981).
- [20] S. Laporta, High precision calculation of multiloop Feynman integrals by difference equations, *Int. J. Mod. Phys. A* **15**, 5087 (2000).
- [21] S. Laporta, Calculation of master integrals by difference equations, *Phys. Lett. B* **504**, 188 (2001).
- [22] A. V. Kotikov, Differential equations method: New technique for massive Feynman diagrams calculation, *Phys. Lett. B* **254**, 158 (1991).
- [23] E. Remiddi, Differential equations for Feynman graph amplitudes, *Nuovo Cimento A* **254**, 158 (1991).
- [24] T. Gehrmann and E. Remiddi, Differential equations for two loop four point functions, *Nucl. Phys.* **B580**, 458 (2000).
- [25] R. Scharf and J. B. Tausk, Scalar two loop integrals for gauge boson self-energy diagrams with a massless fermion loop, *Nucl. Phys.* **B412**, 523 (1994).
- [26] C. Anastasiou, S. Beerli, S. Bucherer, A. Daleo, and Z. Kunszt, Two-loop amplitudes and master integrals for the production of a Higgs boson via a massive quark and a scalar-quark loop, *J. High Energy Phys.* **01** (2007) 082.
- [27] See Supplemental Material at <http://link.aps.org/supplemental/10.1103/PhysRevD.109.096020> for numerical examples and a verification of the dispersion relations.
- [28] A. I. Davydychev, Explicit results for all orders of the epsilon expansion of certain massive and massless diagrams, *Phys. Rev. D* **61**, 087701 (2000).
- [29] A. I. Davydychev and M. Y. Kalmykov, New results for the epsilon expansion of certain one, two and three loop Feynman diagrams, *Nucl. Phys.* **B605**, 266 (2001).
- [30] A. I. Davydychev and M. Y. Kalmykov, Massive Feynman diagrams and inverse binomial sums, *Nucl. Phys.* **B699**, 3 (2004).
- [31] G. Källén and A. Sabry, Two-loop vacuum polarization, *Kong. Dan. Vid. Sel. Mat. Fys. Med.* **29**, 1 (1955).
- [32] J. Schwinger, *Particles, Sources and Fields (Volume III)* (Addison-Wesley, Reading, MA, 1970).
- [33] D. J. Broadhurst, J. Fleischer, and O. V. Tarasov, Two-loop two-point functions with masses: Asymptotic expansions and Taylor series, in any dimension, *Z. Phys. C* **60**, 287 (1993).
- [34] B. A. Kniehl, Two-loop corrections to the vacuum polarizations in perturbative QCD, *Nucl. Phys.* **B347**, 86 (1990).
- [35] A. Djouadi and P. Gambino, Electroweak gauge boson self-energies: Complete QCD corrections, *Phys. Rev. D* **49**, 3499 (1994).
- [36] K. G. Chetyrkin, J. H. Kühn, and M. Steinhauser, Three-loop polarization function and  $O(\alpha_s^2)$  corrections to the production of heavy quarks, *Nucl. Phys.* **B482**, 213 (1996).
- [37] E. Borie, Lamb shift in light muonic atoms, *Z. Phys. A* **275**, 347 (1975).
- [38] E. Borie and G. A. Rinker, The energy levels of muonic atoms, *Rev. Mod. Phys.* **54**, 67 (1982).
- [39] K. Pachucki, Contributions to the binding, two-loop correction to the Lamb shift, *Phys. Rev. A* **48**, 2609 (1993).
- [40] P. J. Mohr, B. N. Taylor, and D. B. Newell, CODATA recommended values of the fundamental physical constants: 2006, *Rev. Mod. Phys.* **80**, 633 (2008).
- [41] E. Tiesinga, P. J. Mohr, D. B. Newell, and B. N. Taylor, CODATA recommended values of the fundamental physical constants: 2018, *Rev. Mod. Phys.* **93**, 025010 (2021).
- [42] U. D. Jentschura, A. Czarnecki, and K. Pachucki, Non-relativistic QED approach to the Lamb shift, *Phys. Rev. A* **72**, 062102 (2005).
- [43] U. D. Jentschura, Relativistic reduced-mass and recoil corrections to vacuum polarization in muonic hydrogen, muonic deuterium, and muonic helium Ions, *Phys. Rev. A* **84**, 012505 (2011).