

Novel approaches to determine B^\pm and B^0 meson production fractions

Florian Bernlochner,¹ Martin Jung², Munira Khan¹, Greg Landsberg³, and Zoltan Ligeti⁴

¹*Physikalisches Institut der Rheinischen Friedrich-Wilhelms-Universität Bonn, 53115 Bonn, Germany*

²*Dipartimento di Fisica, Università di Torino & INFN, Sezione di Torino, I-10125 Torino, Italy*

³*Department of Physics, Brown University, Providence, Rhode Island 02912, USA*

⁴*Ernest Orlando Lawrence Berkeley National Laboratory, University of California, Berkeley, California 94720, USA*



(Received 31 July 2023; accepted 21 May 2024; published 9 July 2024)

We propose novel methods to determine the $\Upsilon(4S) \rightarrow B^+B^-$ and $\Upsilon(4S) \rightarrow B^0\bar{B}^0$ decay rates. The precision to which they and their ratio are known yields at present a limiting uncertainty around 2% in measurements of absolute B decay rates, and thus in a variety of applications, such as precision determinations of elements of the Cabibbo-Kobayashi-Maskawa matrix and flavor-symmetry relations. The new methods we propose are based in one case on exploiting the $\Upsilon(5S)$ datasets, in the other case on the different average number of charged tracks in B^\pm and B^0 decays. We estimate future sensitivities using these methods and discuss possible measurements of f_d/f_u at the (HL-)LHC.

DOI: [10.1103/PhysRevD.110.014007](https://doi.org/10.1103/PhysRevD.110.014007)

I. INTRODUCTION

Precise knowledge of the absolute branching fractions of charged and neutral B meson decays is crucial for a large part of the flavor physics program, spanning the range from understanding hadronic physics to new-physics searches. They enter precision determinations of fundamental Standard Model (SM) parameters, such as elements of the Cabibbo-Kobayashi-Maskawa (CKM) matrix, and flavor-symmetry relations, which in turn impact the sensitivity of CP -violation measurements to physics beyond the SM. As Belle and BABAR recorded hundreds of millions of B meson decays, projected to increase by nearly two orders of magnitude at Belle II, a significant uncertainty in the otherwise ever more precise measurements has become the ratio of $\Upsilon(4S)$ decay to charged vs neutral B mesons,

$$R^{\pm 0} = \frac{\Gamma(\Upsilon(4S) \rightarrow B^+B^-)}{\Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0)}. \quad (1)$$

Although the mass difference $m_{B^0} - m_{B^+} = (0.32 \pm 0.05)$ MeV [1] is small, the restricted phase space in the $\Upsilon(4S) \rightarrow B\bar{B}$ decay of merely 20 MeV and the resulting small velocity of the B mesons give rise to enhanced electromagnetic effects and isospin violation [2]. The range

of theoretical predictions for these effects is substantial, spanning $R^{\pm 0}$ values beyond 1–1.2 [2–9], much above the desired precision. Therefore, direct experimental determinations are crucial. The world average obtained by Heavy Flavor Averaging Group (HFLAV) [10] is

$$R^{\pm 0} = 1.057^{+0.024}_{-0.025}, \quad (2)$$

indicating a notable deviation from unity, albeit smaller than predicted by some theoretical estimates.

Given the above range of applications, reducing this uncertainty to the (sub-)percent level would be very important (as discussed, e.g., in Refs. [11–15].) An intrinsic challenge of such a determination is the difficulty of separating the production fractions from B meson decay rates, since the most often measured quantities determine only their product. Moreover, the measurements usually assume that $\Upsilon(4S)$ decays exclusively to B meson pairs. What is meant by this is that the enhancement of the total e^+e^- cross section near the $\Upsilon(4S)$ resonance equals (within a few permille) the B meson production rate, with B meson production kinematically forbidden for $\sqrt{s} \lesssim m_{\Upsilon(4S)} - \Gamma_{\Upsilon(4S)}$. We denote $f_{\pm} = \Gamma(\Upsilon(4S) \rightarrow B^+B^-)/\Gamma_{\Upsilon(4S)}$, $f_{00} = \Gamma(\Upsilon(4S) \rightarrow B^0\bar{B}^0)/\Gamma_{\Upsilon(4S)}$, and $f_{\cancel{B}} = 1 - f_{\pm} - f_{00}$. Clearly, $R^{\pm 0} = f_{\pm}/f_{00}$, so to relate $R^{\pm 0}$ to absolute branching fractions, knowledge of $f_{\cancel{B}}$ is required. A lower bound on $f_{\cancel{B}}$ is obtained from the sum of measured $\Upsilon(4S)$ decays to lighter bottomonia and pions [10],

$$f_{\cancel{B}} > (0.264 \pm 0.021)\%. \quad (3)$$

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

The strongest constraint, not yet included by HFLAV, is from the CLEO experiment [16],

$$f_{\psi} = (-0.11 \pm 1.43 \pm 1.07)\%, \quad (4)$$

with a larger uncertainty than the desired precision, but limiting f_{ψ} from above.

Another implicit assumption when averaging the available measurements is that the center-of-mass energy and the beam-energy spread at which the $\Upsilon(4S)$ are produced are similar at the relevant colliders and data-taking periods. This will be further discussed in Sec. VI.

In this paper we propose new methods to address these challenges, both in the short term and in the long term. In Sec. II, we reappraise the theoretical assumptions in various $R^{\pm 0}$ measurements, and update its world average. In Sec. III, we propose a new method to determine $R^{\pm 0}$ with small theoretical uncertainties, using $\Upsilon(5S)$ data anticipated at Belle II in the next decade. In Sec. IV, we propose a new method based on the different average number of charged-particle tracks in charged and neutral B decays. In Sec. V we discuss possible measurements of the B^0 to B^+ meson production ratio f_d/f_u at the (HL-)LHC. Finally, Sec. VI discusses additional issues related to collider running conditions and concludes.

II. PRESENT STATUS OF $R^{\pm 0}$

In this section we update the analysis of Ref. [15], with the main difference that we allow for $f_{\psi} \neq 0$. In order to separately determine the production fractions and decay rates, three categories of measurements have been commonly used so far (which we label I, II, and III below):

- (I) Cancellation of final-state dependence. This technique relies on the observation that for double-tagged events in $\Upsilon(4S)$ decays, the B^+B^- and $B^0\bar{B}^0$ production fractions enter linearly, while the decay rate enters quadratically (a technique developed for $\psi(3770)$ [17]). This allows for a cancellation of the dependence on the decay rates in the ratio of the number of single-tag events squared and the number of double-tag events, while retaining that on the production fractions, thus making a theoretically clean measurement of isospin violation in production possible.
- (II) Known ratio of decay rates. Considering any ratio of a charged to a neutral B meson decay, the experimentally determined quantity is proportional to $R^{\pm 0}$ times the ratio of the corresponding decay rates. If the ratio of decay rates is known, it is possible to extract the ratio $R^{\pm 0}$. Taking the ratio of decay rates from an external measurement relies on the determination of $R^{\pm 0}$ at the corresponding experiment, while for an extraction without

external inputs, the knowledge of the ratio of decay rates has to stem from theory. Given the required level of precision, presently the only method available relies on isospin symmetry. While generally, a precise theoretical determination of isospin violation is extremely difficult, there are a few cases in which two decays are not only related by isospin symmetry, but isospin breaking is additionally suppressed. This is the case, e.g., for inclusive semileptonic B meson decays, where the operator product expansion and heavy-quark symmetry provide an additional $\Lambda_{\text{QCD}}^2/m_{c,b}^2$ suppression [18] of the isospin breaking from both the strong interaction (as discussed, e.g., in Ref. [19]) and from electromagnetic effects. In this case, isospin breaking can be safely assumed to be below 1%.

- (III) (Pseudo-)Isospin symmetry. Given the potential enhancement of isospin breaking in production, it is possible to extract it assuming that the breaking for (pseudo-)isospin-related decays is small compared to the one in production. We call *pseudoisospin relations* those in which the amplitudes are *not* equal by isospin symmetry alone, but the unrelated contributions are expected to be of similar size as generic isospin breaking. This is the case, e.g., for $B \rightarrow J/\psi K$ decays, in which the annihilation amplitude contributing only to the charged mode is often argued to be negligible. The remaining isospin breaking is expected to be at the percent level. Clearly, making this assumption precludes the extraction of isospin violation in decay at the same order (and especially in the same decays for which this assumption has been made). This holds also for the values quoted in Eq. (2), since some of the measurements in the average use this assumption. Furthermore, this strategy relies on the assumption that the isospin breaking in production is much larger than that in decay. While reasonable, this assumption is not firmly established experimentally yet, given that the result in Eq. (2) is only about two standard deviations from unity.

The available measurements of $R^{\pm 0}$ are collected in Table I. The only measurement from category I is the BABAR result (using about 82 fb^{-1} of data) for the production fraction of B^0 mesons [20],

$$f_{00} = 0.487 \pm 0.010 \pm 0.008, \quad (5)$$

where the dominant systematic uncertainty stems from the number of $B\bar{B}$ pairs. To turn this into the determination of $R^{\pm 0}$ in Table I, information regarding the non- $B\bar{B}$ fraction in $\Upsilon(4S)$ is required.

From category II, Belle [22] used inclusive semileptonic decays to measure $R^{\pm 0}$. This result needs to be updated to the common ratio of B meson lifetimes, the dominant

TABLE I. Available measurements for $R^{\pm 0}$ from the three categories, as explained in detail in the main text.

$R^{\pm 0}$	Method	Comment	Reference
1.047(44)(36)	Single vs double-tag	Uses f_{ψ} , see text	[10,16,20]
1.039(31)(50)	$B \rightarrow X_c \ell \nu$	Assumes negligible isospin violation	[21,22]
1.068(32)(20)(21)	$B \rightarrow X_s \gamma$	Third uncertainty due to resolved photon contributions	[23]
1.055(30)		Average categories I and II	
1.065(12)(19)(32)	$B \rightarrow J/\psi K$	Third uncertainty due to isospin violation in $B \rightarrow J/\psi K$	[24,25]
1.013(36)(27)(30)	$B \rightarrow J/\psi K$	Third uncertainty due to isospin violation in $B \rightarrow J/\psi K$	[26]
1.100(35)(35)(33)	$B \rightarrow J/\psi(ee)K$	Third uncertainty due to isospin violation in $B \rightarrow J/\psi K$	[27]
1.066(32)(34)(32)	$B \rightarrow J/\psi(\mu\mu)K$	Systematic uncertainties $\sim 100\%$ correlated with ee mode	[27]
1.060(18)(32)		Average for $B \rightarrow J/\psi K$	
1.057(23)		Average of all categories I–III	

systematic uncertainty in this case.¹ It would also be interesting to revisit this analysis technique, where $R^{\pm 0}$ was determined simultaneously with Δm_d in a self-consistent way. The data set used, 30 fb^{-1} , was only a small fraction of the full Belle or current Belle II data.

Another measurement belonging to category II is from the isospin asymmetry A_I between CP -averaged rates in $B \rightarrow X_s \gamma$ decays,

$$A_I(B \rightarrow X_s \gamma) \equiv \frac{\Gamma(\bar{B}^0 \rightarrow X_s \gamma) - \Gamma(B^- \rightarrow X_s \gamma)}{\Gamma(\bar{B}^0 \rightarrow X_s \gamma) + \Gamma(B^- \rightarrow X_s \gamma)}. \quad (6)$$

While the so-called resolved photon contributions affect the isospin asymmetry, this effect is probably subdominant. Therefore, this mode might not be suitable to achieve percent-level precision, but it is still useful, given the current measurements and uncertainties. Assuming isospin symmetry, except for a 2% uncertainty due to the isospin-violating part of the resolved photon contributions [28–30], the Belle measurement $A_I(B \rightarrow X_s \gamma) = (-0.48 \pm 1.49 \pm 0.97 \pm 1.15)\%$ [23] (using all Belle data) translates to the value listed in Table I.

Measurements in category III are presently dominated by $B \rightarrow J/\psi K$ decays. The experimentally determined quantity in these measurements (for a pair of pseudoisospin-related final states, F) is the ratio

$$q_F \equiv R^{\pm 0} \frac{\mathcal{B}(B^- \rightarrow F^-)}{\mathcal{B}(\bar{B}^0 \rightarrow F^0)} = R^{\pm 0} \frac{\tau_{B^-} \Gamma(B^- \rightarrow F^-)}{\tau_{B^0} \Gamma(\bar{B}^0 \rightarrow F^0)}. \quad (7)$$

The B^0 and B^\pm lifetimes, τ_{B^0} and τ_{B^\pm} , respectively, are typically determined separately (but need to be used consistently when combining measurements), and *either* the ratio of rates *or* the ratio of production fractions can be determined, making an assumption about the other. The

¹We follow the HFLAV procedure to account for the change in the central value and uncertainty of the lifetime ratio. Notably, $R^{\pm 0}$ and τ_{B^\pm}/τ_{B^0} are anticorrelated [21].

values in Table I correspond to the assumption that the ratio of rates is equal to unity, and assigning a 3% uncertainty to that assumption (as discussed below). Turning this around, using our averaged value for $R^{\pm 0}$ based on the measurements from the first two categories, we obtain

$$\frac{\Gamma(B^- \rightarrow J/\psi K^-)}{\Gamma(\bar{B}^0 \rightarrow J/\psi \bar{K}^0)} = 1.005 \pm 0.033, \quad (8)$$

or, equivalently,

$$A_I(B \rightarrow J/\psi K) = -0.002 \pm 0.017, \quad (9)$$

where now the uncertainty due to the production fractions is taken into account consistently. This shows no indication of a sizable violation of the pseudoisospin relation.

A few comments regarding the values in Table I are in order:

- (i) The measurements show excellent consistency, even among the different categories. While the uncertainties are sizable, this indicates that the isospin asymmetry is not anomalously large in the modes used for this determination, namely in $B \rightarrow J/\psi K$ decays. This is quantified in Eq. (8), which also motivates the uncertainty assigned to it above: isospin conservation was not assumed in obtaining Eq. (8), but is experimentally seen to hold at this level.
- (ii) The average of all the values in Table I, including our estimates for the uncertainty due to isospin violation, results in

$$R_{\text{I+II+III}}^{\pm 0} = 1.057 \pm 0.023, \quad (10)$$

which is numerically close to the HFLAV average, but more robust, since it includes additional uncertainties for the assumptions made. The reason is that additional measurements are included here [16,23,24,27].




FIG. 1. Impact of the treatment of $f_{\not{B}}$ on the determination of the $B\bar{B}$ production fractions. The black line corresponds to setting $f_{\not{B}} = 0$, i.e., $f_{00} + f_{\pm} = 1$. The red line corresponds to setting $f_{\not{B}}$ to the lower bound in Eq. (3). The yellow-shaded areas use our results in Table I and treat Eq. (3) as a lower limit, while the blue-shaded constraints include the CLEO measurement in Eq. (4). The lighter and darker regions show $\Delta\chi^2 \leq 5.99$ and 2.28, respectively, while the dashed lines correspond to $\Delta\chi^2 = 1, 4$ (illustrating one-dimensional limits).

- (iii) In principle, $R^{\pm 0}$ and f_{00} together determine $f_{\not{B}}$ via $f_{\not{B}} = 1 - f_{00}(1 + R^{\pm 0}) = -0.003 \pm 0.029$; however, this uncertainty is still larger than that in Eq. (4).
- (iv) Since we include $f_{\not{B}}$ in our calculations, the average for $R^{\pm 0}$ does not represent the full information from our analysis. Specifically, while with $f_{\not{B}} = 0$ the value for $R^{\pm 0}$ is in a one-to-one correspondence with f_{00} and f_{\pm} , this no longer holds for $f_{\not{B}} \neq 0$. For instance, determining f_{\pm} now requires two of the measured quantities:

$$\begin{aligned} f_{\pm} &= f_{00}R^{\pm 0} = 1 - f_{00} - f_{\not{B}} \\ &= \frac{R^{\pm 0}(1 - f_{\not{B}})}{1 + R^{\pm 0}}. \end{aligned} \quad (11)$$

However, it is f_{00} and f_{\pm} that determine the precision of absolute branching fractions, not $R^{\pm 0}$. While our result for $R^{\pm 0}$ is numerically close to the one from HFLAV [10], the results for the production fractions are quite different, since we include the CLEO measurement of $f_{\not{B}}$ [16]. This is illustrated in Fig. 1, where we compare the impact of different treatments of $f_{\not{B}}$ on f_{00} and f_{\pm} . This highlights the importance of determining $f_{\not{B}}$ with better precision.

The resulting uncertainties are asymmetric and highly correlated, and the central values and $\Delta\chi^2 = 1$ ($\Delta\chi^2 = 4$) ranges of the production fractions from the fit including the CLEO measurement are

$$\begin{aligned} f_{\pm} &= 0.512[0.504, 0.518]([0.493, 0.523]), \\ f_{00} &= 0.485[0.478, 0.491]([0.470, 0.496]), \\ f_{\not{B}} &= 0.003[0.002, 0.014]([0.002, 0.029]). \end{aligned} \quad (12)$$

The fit results without the CLEO measurement are

$$\begin{aligned} f_{\pm} &= 0.512[0.497, 0.518]([0.479, 0.523]), \\ f_{00} &= 0.485[0.474, 0.491]([0.461, 0.496]), \\ f_{\not{B}} &= 0.003[0.002, 0.027]([0.002, 0.056]), \end{aligned} \quad (13)$$

so our analysis reduces the uncertainties in f_{\pm} and f_{00} from about 2 to 1.5%. The difference would be even larger without the measurement of f_{00} ; this shows again the necessity to determine individual production fractions for either B mesons or non- $B\bar{B}$ states.

Interestingly, the values in Table I are not only consistent with one another, but also with the value obtained considering only the phase-space difference between $\Upsilon(4S) \rightarrow B^0\bar{B}^0$ and $\Upsilon(4S) \rightarrow B^+B^-$,

$$R_{\text{PS}}^{\pm 0} = \frac{p_{\pm}^3}{p_0^3} \approx 1.048. \quad (14)$$

This value is larger than may be naively expected, due to the small phase space, which amplifies the impact of the small mass difference between the charged and neutral B mesons. On the other hand, the naive Coulomb enhancement of the charged mode, in the nonrelativistic limit and assuming pointlike mesons, is [2]

$$R_{\text{CE}}^{\pm 0} = \frac{2\pi\lambda(1 + \lambda^2)}{1 - \exp(-2\pi\lambda)}, \quad (15)$$

where $\lambda = \alpha/(2v_{\pm})$ denotes the Coulomb parameter (and $v_{\pm} = (1 - 4m_{B^{(*)\pm}}^2/m_{\Upsilon}^2)^{1/2}$, as appropriate for the B or B^* states in the $\Upsilon(4S)$ or $\Upsilon(5S)$ decays), which yields the values in the second to last column of Table II.² Evidently, the large enhancement expected from this estimate is reduced, given that

$$R_{\text{I+II+III}}^{\pm 0}/R_{\text{PS}}^{\pm 0} = 1.008 \pm 0.022, \quad (16)$$

²Since $m_{B^0} > m_{B^+}$, in $\Upsilon(nS) \rightarrow B\bar{B}$, the phase-space difference and the Coulomb enhancement of the charged mode go in the same direction. (This also holds for $\Upsilon(5S) \rightarrow B^*\bar{B}^*$ discussed below, though in that case the phase-space effect is very small.)

TABLE II. Relative phase-space factors $R_{\text{PS}}^{\pm 0}$ for $\Upsilon(4S)$ and $\Upsilon(5S)$ decays, together with the naive Coulomb enhancement for pointlike particles $R_{\text{CE}}^{\pm 0}$ and their product, corresponding to the naive prediction for $R^{\pm 0}$.

Decay mode	$R_{\text{PS}}^{\pm 0}$	$R_{\text{CE}}^{\pm 0}$	$R_{\text{PS}}^{\pm 0} R_{\text{CE}}^{\pm 0}$
$\Upsilon(4S) \rightarrow B\bar{B}$	1.048	1.20	1.26
$\Upsilon(5S) \rightarrow B\bar{B}$	1.003	1.05	1.05
$\Upsilon(5S) \rightarrow B^* \bar{B}^*$	1.004	1.06	1.06

consistent with small isospin violation beyond the phase-space factor. Nevertheless, the (additional) production asymmetry from isospin violation in the $\Upsilon(4S)$ decay may still be larger than without any enhancement.

While the determinations of $R^{\pm 0}$ in Table I can be considered robust, as they explicitly include uncertainty estimates for the assumptions made, they are still unsatisfactory in several ways:

- (1) The overall precision is not at the level necessary for high-precision measurements at current and future colliders.
- (2) There is no clear path to reduce the uncertainties related to isospin breaking, so additional measurements from category III (or $B \rightarrow X_s \gamma$, from category II) would not reduce this uncertainty further.
- (3) The average uses decay modes, whose isospin asymmetries are themselves of interest. This concerns for instance the resolved photon contributions in $B \rightarrow X_s \gamma$ or the annihilation contributions in $B \rightarrow J/\psi K$ decays. Overall, it would be desirable for applications in B physics to have a determination of $R^{\pm 0}$ that does not rely on specific B meson decays, but rather only on properties of the Υ system. Of the methods employed so far, only the double-tag technique fulfills this criterion.
- (4) The difficulty in using only the double-tag technique is that it requires very large datasets, due to the low efficiency for double-tag events, even if using a semi-inclusive tagging.

For these reasons, having independent methods that do not rely on assumptions about specific B decays would be important. Below, we propose two such methods.

III. DETERMINING $R^{\pm 0}$ USING $\Upsilon(5S)$ DECAYS

As discussed in the previous sections, the main reason for the sizable isospin violation causing $R^{\pm 0}$ to deviate from unity is the small phase space in $\Upsilon(4S)$ decays, $m_{\Upsilon(4S)} - 2m_B \simeq 20$ MeV, while the mass difference near the $\Upsilon(5S)$ resonance is more substantial, $m_{\Upsilon(5S)} - 2m_B \simeq 326$ MeV. However, an e^+e^- collider running near this resonance produces many different final states.

Experimentally, $\Gamma(\Upsilon(5S) \rightarrow BBX) = (76.2_{-4.0}^{+2.7})\%$ [1,31], of which only about 5.5% is direct $B\bar{B}$ production,

complemented by BB^* (13.7%) and B^*B^* (38.1%) production.³ Additionally multibody final states, such as $B^{(*)}B^{(*)}\pi$ and $BB\pi\pi$ contribute. For the (quasi-)two-body final states, we expect

$$R_{5S}^{\pm 0} = \frac{\Gamma(\Upsilon(5S) \rightarrow B^{(*)+}B^{(*)-})}{\Gamma(\Upsilon(5S) \rightarrow B^{(*)0}\bar{B}^{(*)0})} \simeq 1 \quad (17)$$

to be similarly reduced, in light of $R_{\text{PS}}^{\pm 0}|_{\Upsilon(5S)} - 1 < 0.5\%$, and additional effects seen above to be $\lesssim 2\%$ for the $\Upsilon(4S)$ system. This allows for a novel determination of $R^{\pm 0}$, by studying the double ratio of pairs of decays at the $\Upsilon(4S)$ and $\Upsilon(5S)$ resonances,

$$r(f, f') = \frac{[N(B^+ \rightarrow f)]}{[N(B^0 \rightarrow f')]_{\Upsilon(4S)}} \bigg/ \frac{[N(B^+ \rightarrow f)]}{[N(B^0 \rightarrow f')]_{\Upsilon(5S)}}. \quad (18)$$

Here N denotes the acceptance- and efficiency-corrected yields in $\Upsilon(4S)$ and $\Upsilon(5S)$ decays, in the latter case including B mesons from all (quasi-)two-body decays $\Upsilon(5S) \rightarrow \bar{B}^{(*)}B^{(*)}$. Crucially, in this ratio the $\mathcal{B}(B^+ \rightarrow f)$ and $\mathcal{B}(B^0 \rightarrow f')$ branching fractions cancel, so no information on the size of isospin breaking in the decay rates is needed. In fact, f and f' do not have to be (pseudo-)isospin related, and any pair of states can be chosen to minimize the experimental uncertainties. Thus, the double ratio in Eq. (18) directly probes the ratio of production rates, $R^{\pm 0}$, assuming Eq. (17) holds.

One aspect that could spoil Eq. (17) is the contamination from final states other than $B^{(*)}\bar{B}^{(*)}$, where the reduced phase space may enhance isospin violation. However, if the $B \rightarrow f$ decay is reconstructed in a fully hadronic final state, its kinematic properties can be used to separate many-body from the (quasi-)two-body production, using the beam-constrained mass $M_{\text{bc}} = \sqrt{s/4 - |\vec{p}_B|^2}$ [33], where \vec{p}_B is the three-momentum of the reconstructed B meson. The M_{bc} method can also be used for semileptonic decays [34].

In Table III we present estimates of projected sensitivities to $r(f, f')$ using this method, with the existing Belle, as well as anticipated Belle II data, the latter split into partial (10%) and full datasets. We studied a few promising modes, corresponding to different parton-level transitions and different experimental signatures and uncertainties. We base our uncertainty estimates on Refs. [24,35–37] and

³Regarding phase-space differences caused by the B^* masses, the mass difference $m_{B^{*0}} - m_{B^{*+}} = (0.91 \pm 0.26)$ MeV [32] has been measured by the CMS Collaboration. Curiously, this value is approximately $-m_c/m_b \simeq -\frac{1}{3}$ times $m_{D^{*0}} - m_{D^{*+}} \simeq -3.4$ MeV [1], as expected from heavy-quark symmetry. The isospin splittings of the ground-state mesons, $m_{B^0} - m_{B^+} \approx 0.3$ MeV and $m_{D^0} - m_{D^+} \approx -4.8$ MeV [1], are far from this relation, probably due to electromagnetic effects.

TABLE III. Estimated sensitivity to $r(f, f')$ in Eq. (18), with available Belle data and anticipated partial and full Belle II data.

	Belle	Belle II partial	Belle II full
$\mathcal{L}_{\Upsilon(5S)}/\mathcal{L}_{\Upsilon(4S)}$ [ab ⁻¹ /ab ⁻¹]	0.12/0.71	0.5/5	5/50
$N_{B^{(*)}B^{(*)}}^{\Upsilon(5S)}/N_{BB}^{\Upsilon(4S)}$	$2.74 \times 10^7/7.72 \times 10^8$	$1.13 \times 10^8/5.55 \times 10^9$	$1.13 \times 10^9/5.55 \times 10^{10}$
f, f'		$\Delta r(f, f')/r(f, f')$	
$J/\psi K^+, J/\psi K^0$	7.1%	3.5%	1.1%
$\bar{D}^0 \pi^+, D^- \pi^+$	2.4%	1.2%	0.4%
$\bar{D}^{*0} \ell^+ \nu, D^{*-} \ell^+ \nu$	4.5%	2.2%	0.7%
$\bar{D}^0 \pi^+, D^{*-} \ell^+ \nu$	1.8%	0.9%	0.3%

assume for Belle II an improvement on the systematic uncertainties by a factor of 2. We scale the statistical uncertainties with the integrated luminosity ratios and base our estimates for the $\Upsilon(5S)$ analyses on the precision of the $\Upsilon(4S)$ measurements, assuming the same systematic uncertainties, but correspondingly larger statistical uncertainties. We further assume that common systematic uncertainties cancel between the $\Upsilon(4S)$ and $\Upsilon(5S)$ measurements.

For $B \rightarrow J/\psi K$ decays and the currently available Belle $\Upsilon(4S)$ and $\Upsilon(5S)$ datasets, a precision of 7.1% on $R^{\pm 0}$ can be reached, limited by the statistical uncertainty of the $\Upsilon(5S)$ measurement. A determination focusing on $B \rightarrow D\pi^{\pm}$ decays could already reach a precision similar to the current world average. Semileptonic $B \rightarrow D^* \ell \bar{\nu}$ also offer a clean avenue, but are limited by the $B^+ \rightarrow \bar{D}^{*0} \ell^+ \nu$ precision at $\Upsilon(5S)$. An additional improvement could be obtained by focusing on $B \rightarrow DX \ell \bar{\nu}$ decays. These three decays look promising to reach 1% or even sub-1% uncertainties with a Belle II dataset of 5 ab⁻¹ of $\Upsilon(5S)$ data. We illustrate the fact that f and f' can be chosen independently to minimize the experimental uncertainties by studying mixed $D\pi$ and semileptonic channels, further improving the precision of the $r(f, f')$ determination.

We conclude that the double-ratio method using either $B \rightarrow D\pi^{\pm}$ decays or mixed $D\pi$ and semileptonic decays is a promising way to study the feasibility of this method with the existing Belle data.

IV. DECAY-CHANNEL-INDEPENDENT DETERMINATION OF $R^{\pm 0}$

An alternative to using specific decay channels for determinations of $R^{\pm 0}$ would be the use of the full range of B meson decays. This would constitute another way to remove assumptions about isospin violation. To our knowledge, such a determination has not been attempted; in the following we discuss a possible strategy and estimate the corresponding sensitivities. This idea utilizes the fact that B mesons leave a fairly easy signature to trigger on, and that

e^+e^- B -factory experiments operate often with nearly 100% efficiency to record events. Most triggers rely on properties that are nearly identical for $B^0 \bar{B}^0$ and $B^+ B^-$ events: a typical selection requires at least three tracks and more than 1-GeV energy deposition in the calorimeter with four isolated clusters. Such inclusive samples are in fact regularly analyzed to count the number of B meson pairs and to subtract backgrounds from continuum processes [38].

To separate $B^0 \bar{B}^0$ and $B^+ B^-$ events, another event property can be combined with this approach: the total number of detector-stable charged daughter particles. This is a difficult quantity to reconstruct, but has a reliable proxy with the total number of charged-particle tracks. Figure 2 (left; top and bottom) shows the number of charged daughter particles for B^+ and B^0 meson decays as simulated by EvtGen [39], without any selection. A distinctive feature is that for B^+ (B^0) the number of charged daughters must be odd (even). This separation is reduced if one looks at the number of charged daughters of a pair of B mesons produced in $\Upsilon(4S)$ decay (as shown in Fig. 2, right). A key problem is that these distributions are sensitive to the modeling of B meson decays. For instance, in EvtGen thousands of exclusive decays are mixed with final states from Pythia8 [40] to simulate inclusive B meson decays. One way to control this is to measure this distribution or rather its proxy (the number of charged-particle tracks) in data using decays of B mesons, which identify their charge. For instance, one can consider $B^0 \rightarrow D^- \pi^+$ decays, which, despite their small branching fraction of $\approx 2.5 \times 10^{-3}$, can be reconstructed with excellent experimental precision.

With the final-state particles of one B meson decay precisely assigned to this signal, the rest of the collision event can be assessed and the multiplicity distribution of the number of charged-particle tracks can be precisely measured. Similar measurements can be carried out with B^{\pm} decays and with other exclusive channels. This way one can obtain the key ingredients for the prediction of the B meson pair distributions from data, as their decays progress fully independently from each other.




FIG. 2. Number of charged daughters from B^+ and B^0 decay (left) and from a pair of B decays (right), from EvtGen.

We construct an Asimov fit [41] to assess the separation power using the number of charged particles. We assume that a calibration of the charged and neutral B meson multiplicities can be carried out with $B^0 \rightarrow D^- \pi^+$ and $B^+ \rightarrow D^0 \pi^+$ decays. We scale the statistical uncertainty of Ref. [35] by the expected increase in the integrated luminosity to evaluate the future Belle II sensitivities for 5 ab^{-1} and 50 ab^{-1} . As we do not need to measure a branching fraction, but rather the distribution of reconstructed tracks, many of the leading systematic uncertainties in Ref. [35] do not dilute the sensitivity. Using the number of events and the expected distributions, we determine templates and correlated uncertainties for the $B^0 \bar{B}^0$ and $B^+ B^-$ multiplicity distributions. Notably, the predictions in the bins of the multiplicity for a pair of B mesons are correlated, as they are predicted from sampling twice the distribution of a single B meson decay. We fit the resulting distributions with different assumptions for the three luminosity scenarios and report the achievable relative uncertainties in Table IV, taking into account the uncertainties from the expected calibration precision.

In practice, additional reconstruction effects will cause differences between the number of charged particles and the number of tracks, such as the finite detector acceptance

TABLE IV. The estimated $R^{\pm 0}$ sensitivity achievable using the number of charged-particle tracks. This includes the calibration uncertainty in the number of charged-particle tracks from $B^0 \rightarrow D^- \pi^+$ decays and assumes a similar sensitivity can be achieved in $B^\pm \rightarrow D^0 \pi^+$ decays. Without the calibration uncertainty, the statistical component would be subpercent even with the data available now.

	Belle	Belle II partial	Belle II full
$\mathcal{L}_{\Upsilon(4S)}$ [ab^{-1}]	0.71	5	50
$\Delta(R^{\pm 0})/R^{\pm 0}$	2.2%	0.9%	0.3%

or the occurrence of misidentified or duplicate tracks. Such effects shift and broaden the $B^0 \bar{B}^0$ and $B^+ B^-$ distributions, and a more robust study on the feasibility of this method can only be done within the experiments.

V. PRODUCTION FRACTION RATIOS AT HADRON COLLIDERS

At hadron colliders, the production fractions of charged and neutral B mesons, denoted f_u and f_d , respectively, play an analogous role to that of the Υ decay rates at $e^+ e^- B$ factories. However, symmetry considerations are not as easily applicable: *a priori*, we *cannot* expect the relation $f_u = f_d$ to hold, since both the initial and final states are more complicated than at a B factory. At the Tevatron, the initial $p\bar{p}$ state is a superposition of an isosinglet and an isotriplet, while at the LHC the pp initial state is a pure isotriplet. Furthermore, the presence of additional particles in the final state does not allow for a determination of the isospin state of the b hadron pair. However, the dominant $b\bar{b}$ production mechanisms at the LHC (gluon splitting and t -channel flavor creation) are isospin invariant. At the same time, the fragmentation into B mesons is a complicated process. Regarding fragmentation to B_s mesons, corrections to the $SU(3)$ flavor symmetry are large, as measured by the ratio $f_s/f_d \approx 0.25$, with a dependence on the center-of-mass energy and kinematics [42]. Therefore, the size of the ratio f_u/f_d is ultimately an experimental question and $f_u/f_d = 1$ cannot be assumed, but should be determined experimentally, including a possible kinematic dependence, as observed for f_s and f_{Λ_b} [42,43].

The experimental determination of this quantity is again complicated by the difficulty of decoupling the production fractions from the decay rates. An additional complication arises due to the uncorrelated hadronization of the b and \bar{b} quarks produced, such that category I measurements discussed above are not possible. This leaves us with categories II and III.

A measurement falling into category II with external inputs of the ratio of decay rates (and thereby $R^{\pm 0}$) for $B^0 \rightarrow J/\psi K^{*0}$ and $B^+ \rightarrow J/\psi K^+$ has been carried out by the CMS Collaboration [44], yielding

$$\frac{f_d}{f_u} = 1.015 \pm 0.051. \quad (19)$$

The precision of this measurement is presently limited by the uncertainty in the CMS analysis and to lesser extent by the uncertainty in $R^{\pm 0}$.

On the other hand, it would be desirable to obtain a measurement of f_d/f_u that does not rely on the external measurement of $R^{\pm 0}$, using the large samples of B mesons that already exist at the LHC and will be significantly enlarged in the HL-LHC era. To that aim, we propose to use the approximate equality of rates of the semi-inclusive decays,

$$\Gamma(B^0 \rightarrow \bar{D}^{(*)} X_{\mu\nu}) \approx \Gamma(B^+ \rightarrow \bar{D}^{(*)} X_{\mu\nu}). \quad (20)$$

This relation follows from the equality of inclusive rates discussed above, given the small fraction of decays that do not result in a $D^{(*)}$ meson in the final state, specifically final states including $D_s^{(*)} \bar{K}^{(*)}$ or baryons. An analogous method has been employed in the determination of the ratio of production fractions f_s/f_d from semileptonic decays by the LHCb collaboration [43]. These final states also include decays of B_s and Λ_b . While most of these decays have not been observed explicitly, the ones that have been seen sum to a branching fraction of $\sim 1\%$. Their contributions are additionally suppressed by the smaller production fractions, f_s and f_{Λ_b} , respectively, so accounting for them should not be too difficult [45]. In order to separate the neutral and charged B mesons decaying into these final states, one possibility is to employ the oscillations in B^0 meson mixing, which are absent for B^{\pm} mesons. This has been used by the LHCb collaboration in a time-dependent semi-inclusive measurement of Δm_d [45] to remove the background from B^{\pm} mesons; here, this background is considered instead part of the signal.

Whether the desired $\mathcal{O}(1\%)$ precision can be reached via this method is an experimental question; we leave the detailed studies to dedicated experimental analyses, and simply point out their potential use for measuring B meson production fractions.

Finally, large samples of $t\bar{t}$ events accumulated at the LHC by the ATLAS and CMS Collaborations could also be used as a way to test isospin invariance in production and/or decay of B mesons. Unlike the pp initial state, the $t\bar{t}$ system is an isospin singlet. If the interaction with the rest of the event, often referred to as ‘‘color reconnection,’’ is small, and we consider the case in which the W bosons from the subsequent $t \rightarrow bW^+$ process decay leptonically, we can

expect $f_u = f_d$ for B mesons produced in this process, based on the isospin symmetry.

In this sense, the $t\bar{t}$ system at the LHC can play a similar role as the $\Upsilon(4S)$ or Z at e^+e^- colliders, as an isosinglet source of B mesons. It is therefore interesting to experimentally test the $f_d = f_u$ relationship in top-quark decays, and also to test the equality of B and \bar{B} production, which could be affected, e.g., by the valence quarks in the protons. This can be achieved by tagging the top quark (or antiquark) in the event by measuring the charge of the lepton in a leptonic W boson decay from the $t \rightarrow bW^+$ or $\bar{t} \rightarrow \bar{b}W^-$ process and then compare the yield of charged and neutral B mesons produced in the fragmentation of the b jet accompanying the W boson. If this ratio is different from unity, it could have a profound impact on our understanding of color reconnection [46]. In any case, if $f_{u,d}$ can be determined with good precision in this process, the $t\bar{t}$ system can be used to probe isospin invariance in B meson decays.

Again, we leave the detailed studies of the feasibility of this approach to experiments, and simply mention them as a complementary approach to test the isospin invariance and determine production fractions of b hadrons using decays of top quarks, which has not been done before.

VI. DISCUSSION AND CONCLUSIONS

Before concluding, we would like to mention a few aspects regarding the experimental environment that also affect the picture discussed so far. Throughout this paper we have assumed that $R^{\pm 0}$ is a constant. This would be correct if the $\Upsilon(4S)$ would itself be produced in a decay process, but at a e^+e^- collider its mass is constrained by the center-of-mass-energy \sqrt{s} of the colliding beams. The typical beam-energy spread at B factories, such as PEP-II, KEKB, or SuperKEKB, is about 4–6 MeV [47,48], which is several times smaller than the width of the $\Upsilon(4S)$, 20.5 MeV [1]. Assuming a Gaussian distribution for the beam-energy spread, the functional dependence of $R^{\pm 0}$ on \sqrt{s} results in a small bias. There is again a substantial range of predictions for this energy dependence [4–9]. Using as examples the phase-space estimate in Eq. (14) or the simplified Coulomb factor in Eq. (15) (which appears to be an overestimate), we find that the impact of the beam energy spread is 0.3 or 0.4%, respectively, which is currently an order of magnitude smaller than the experimental accuracy. We expect that this effect is much smaller at the $\Upsilon(5S)$ resonance.

Another interesting question is what happens to $R^{\pm 0}$ if different experiments run at different center-of-mass energies, near, but not exactly on, the peak cross section of $e^+e^- \rightarrow b\bar{b}$ in the vicinity of the $\Upsilon(4S)$ resonance. Such shifts can also occur during different runs of a single experiment. If the total data of a given experiment are used to extract branching fractions of the same experiment using




FIG. 3. The ratio $R^{\pm 0}$ as a function of \sqrt{s} using Eqs. (14) (solid) and (15) (dashed), normalized to the respective values at $\sqrt{s} = m_{\Upsilon(4S)} = 10.5794$ GeV [1].

an identical dataset, there is of course no problem: the recovered $R^{\pm 0}$ values correspond to the recorded data. However, if several experiments are combined, or within an experiment $R^{\pm 0}$ determinations use different data-taking periods and conditions, biases may emerge. We can estimate possible shifts by studying Fig. 3 qualitatively: if the phase-space dependence is the leading contribution that changes $R^{\pm 0}$ as a function of \sqrt{s} , shifts of the order 1% can occur for order 5 MeV shifts away from the peak cross section. If Eq. (15) is used instead, these shifts can be as large as 3%. Since, as discussed above, we cannot rely on any of these estimates, the energy dependence of $R^{\pm 0}$ should be experimentally determined. This can be done, in a limited range, by exploiting the varying running conditions that provide samplings around the peak $e^+e^- \rightarrow b\bar{b}$ cross section. For a more complete exploration, a dedicated energy scan and measurements using modes that can be reliably identified, such as $B \rightarrow J/\psi K$, are required.

Another key question resides in the experimental determination of the number of B meson pairs and its robustness against $f_{\neq} \neq 0$. In order for Belle II to achieve percent-level precision goals in the study of branching fractions and other observables, a consistent treatment is needed that takes into account the correlated aspects of $R^{\pm 0}$ determinations and B meson counting.

In summary, we investigated the determinations of the $\Upsilon(4S) \rightarrow B^+B^-$ and $B^0\bar{B}^0$ decay rates and proposed new methods to improve them. Presently the limited precision of

these decay rates constitutes a lower limit of $\sim 2\%$ on the uncertainties in absolute branching fraction measurements, and thereby in applications, such as precision determinations of CKM matrix elements or flavor-symmetry relations. We revisited the theoretical assumptions in $R^{\pm 0}$ measurements, and updated its world average in Sec. II, emphasizing underestimated uncertainties in prior evaluations, in particular due to isospin violation and nonzero f_{\neq} value (as shown in Fig. 1). Due to the inclusion of additional measurements, we obtained nevertheless an improved precision for $R^{\pm 0}$ and the individual production fractions of about 2 and 1.5%, given in Eqs. (10) and (12), respectively. When using both f_{\pm} and f_{00} , care must be taken to include their correlations shown in Fig. 1. We proposed two new methods in Secs. III and IV to determine $R^{\pm 0}$ precisely, using $\Upsilon(5S)$ data anticipated at Belle II over the next decade, or using the different average number of charged-particle tracks between charged and neutral B meson decays. Tables III and IV summarize our estimates of future sensitivities. Section V proposed possible measurements of f_d/f_u at the (HL-)LHC.

The issues raised and methods developed in this article will remain important at future colliders. In the meantime, progress could already be made by revisiting the measurement of Ref. [22] with the full dataset, performing a double-tag analysis at Belle or Belle II, and by using our methods with the existing data.

ACKNOWLEDGMENTS

We thank Paolo Gambino, Dean Robinson, Frank Tackmann, and Kerstin Tackmann for helpful conversations. We thank the organizers of the workshop on ‘‘Challenges in Semileptonic B Decays’’ for the stimulating Barolo ambiance when this work started. We also thank the CERN theory group for hospitality. F. B. and M. K. are supported by DFG Emmy-Noether Grant No. BE 6075/1-1 and BMBF Grant No. 05H21PDKBA. F. B. also thanks the LBNL theory group for its hospitality. The work of M. J. is supported by the Italian Ministry of Research (MIUR) under Grant No. PRIN 20172LNEEZ. The work of G. L. and Z. L. was supported in part by the Office of High Energy Physics of the U.S. Department of Energy under Contracts No. DE-SC0010010 and No. DE-AC02-05CH11231, respectively.

[1] R. L. Workman *et al.* (Particle Data Group), *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).

[2] D. Atwood and W. J. Marciano, *Phys. Rev. D* **41**, 1736 (1990).

[3] G. P. Lepage, *Phys. Rev. D* **42**, 3251 (1990).

[4] N. Byers and E. Eichten, *Phys. Rev. D* **42**, 3885 (1990).

[5] R. Kaiser, A. V. Manohar, and T. Mehen, *Phys. Rev. Lett.* **90**, 142001 (2003).

[6] M. B. Voloshin, *Mod. Phys. Lett. A* **18**, 1783 (2003).

- [7] M. B. Voloshin, *Phys. At. Nucl.* **68**, 771 (2005).
- [8] S. Dubynskiy, A. Le Yaouanc, L. Oliver, J. C. Raynal, and M. B. Voloshin, *Phys. Rev. D* **75**, 113001 (2007).
- [9] A. I. Milstein and S. G. Salnikov, *Phys. Rev. D* **104**, 014007 (2021).
- [10] Y. S. Amhis *et al.* (Heavy Flavor Averaging Group, HFLAV Collaborations), *Phys. Rev. D* **107**, 052008 (2023).
- [11] M. Jung, *Phys. Rev. D* **86**, 053008 (2012).
- [12] Y. Grossman and D. J. Robinson, *J. High Energy Phys.* **04** (2013) 067.
- [13] M. Jung and S. Schacht, *Phys. Rev. D* **91**, 034027 (2015).
- [14] Z. Ligeti and D. J. Robinson, *Phys. Rev. Lett.* **115**, 251801 (2015).
- [15] M. Jung, *Phys. Lett. B* **753**, 187 (2016).
- [16] B. Barish *et al.* (CLEO Collaboration), *Phys. Rev. Lett.* **76**, 1570 (1996).
- [17] R. M. Baltrusaitis *et al.* (MARK-III Collaboration), *Phys. Rev. Lett.* **56**, 2140 (1986).
- [18] J. Chay, H. Georgi, and B. Grinstein, *Phys. Lett. B* **247**, 399 (1990).
- [19] M. Gronau, Y. Grossman, and J. L. Rosner, *Phys. Rev. D* **73**, 057501 (2006).
- [20] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **95**, 042001 (2005).
- [21] N. C. Hastings, Neutral B meson mixing and CPT violation with the Belle experiment, Ph.D. thesis, Charles University, 2002.
- [22] N. C. Hastings *et al.* (Belle Collaboration), *Phys. Rev. D* **67**, 052004 (2003).
- [23] S. Watanuki *et al.* (Belle Collaboration), *Phys. Rev. D* **99**, 032012 (2019).
- [24] S. Choudhury *et al.* (Belle Collaboration), *Phys. Rev. D* **107**, L031102 (2023).
- [25] S. Choudhury *et al.* (Belle Collaboration), *J. High Energy Phys.* **03** (2021) 105.
- [26] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. D* **69**, 071101 (2004).
- [27] F. Abudinén *et al.* (Belle II Collaboration), [arXiv:2207.11275](https://arxiv.org/abs/2207.11275).
- [28] F. U. Bernlochner, H. Lacker, Z. Ligeti, I. W. Stewart, F. J. Tackmann, and K. Tackmann, *Phys. Rev. Lett.* **127**, 102001 (2021).
- [29] A. Gunawardana and G. Paz, *J. High Energy Phys.* **11** (2019) 141.
- [30] M. Benzke, S. J. Lee, M. Neubert, and G. Paz, *J. High Energy Phys.* **08** (2010) 099.
- [31] R. Mizuk *et al.* (Belle Collaboration), *J. High Energy Phys.* **06** (2021) 137.
- [32] A. M. Sirunyan *et al.* (CMS Collaboration), *Eur. Phys. J. C* **78**, 939 (2018).
- [33] A. Drutskoy *et al.* (Belle Collaboration), *Phys. Rev. D* **81**, 112003 (2010).
- [34] P. del Amo Sanchez *et al.* (BABAR Collaboration), *Phys. Rev. D* **83**, 032007 (2011).
- [35] E. Waheed *et al.* (Belle Collaboration), *Phys. Rev. D* **105**, 012003 (2022).
- [36] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. D* **77**, 032002 (2008).
- [37] B. Aubert *et al.* (BABAR Collaboration), *Phys. Rev. Lett.* **100**, 231803 (2008).
- [38] A. J. Bevan *et al.* (BABAR, Belle Collaborations), *Eur. Phys. J. C* **74**, 3026 (2014).
- [39] D. J. Lange, *Nucl. Instrum. Methods Phys. Res., Sect. A* **462**, 152 (2001).
- [40] T. Sjöstrand, S. Ask, J. R. Christiansen, R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C. O. Rasmussen, and P. Z. Skands, *Comput. Phys. Commun.* **191**, 159 (2015).
- [41] G. Cowan, K. Cranmer, E. Gross, and O. Vitells, *Eur. Phys. J. C* **71**, 1554 (2011); **73**, 2501(E) (2013).
- [42] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. D* **104**, 032005 (2021).
- [43] R. Aaij *et al.* (LHCb Collaboration), *Phys. Rev. D* **100**, 031102 (2019).
- [44] A. Tumasyan *et al.* (CMS Collaboration), *Phys. Rev. Lett.* **131**, 121901 (2023).
- [45] R. Aaij *et al.* (LHCb Collaboration), *Eur. Phys. J. C* **76**, 412 (2016).
- [46] S. Argyropoulos and T. Sjöstrand, *J. High Energy Phys.* **11** (2014) 043.
- [47] B. Aubert *et al.* (BABAR Collaboration, PEP-II Machine Group), in *21st International Symposium on Lepton and Photon Interactions at High Energies (LP 03)* (2003); *Int. J. Mod. Phys. A* **19**, 785 (2004).
- [48] K. Akai, K. Furukawa, and H. Koiso (on behalf of the SuperKEKB Accelerator Team), *Nucl. Instrum. Methods Phys. Res., Sect. A* **907**, 188 (2018).