

Exact evaluation of large-charge correlation functions in nonrelativistic conformal field theory

Silas R. Beane^{1,3}, Domenico Orlando^{2,1,4} and Susanne Reffert^{2,1}

¹*Albert Einstein Center for Fundamental Physics, Institut für Theoretische Physik, Universität Bern, Sidlerstrasse 5, CH-3012 Bern, Switzerland*

²*Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 650-0047, Japan*

³*Department of Physics, University of Washington, Seattle, Washington 98195, USA*

⁴*INFN sezione di Torino, via Pietro Giuria 1, 10125 Torino, Italy*



(Received 10 April 2024; accepted 10 June 2024; published 15 July 2024)

The large-charge master field which generates all n -point correlation functions with an insertion of large charge Q in nonrelativistic conformal field theory is obtained. This field is used to compute Schrödinger-invariant n -point correlation functions of large-charge operators via a direct evaluation of the path integral. Conformal dimensions are found to agree with calculations based on the state-operator correspondence. The master field solution exhibits an emergent harmonic trap whose frequency is a function of the Euclidean time. The large-charge effective action with operator insertions describes a droplet of superfluid matter whose spatial size scales with the time separation of sources. The solution is used to compute Schrödinger symmetry breaking corrections in the large-charge effective field theory (EFT) due to a finite scattering length in the fundamental theory of fermions near unitarity. The scaling of these effects in the large-charge power counting scheme is established, and the size of the effects is quantified using input from quantum Monte Carlo simulations of the near-unitary gas, as well as from the large- N expansion at large charge.

DOI: [10.1103/PhysRevD.110.025011](https://doi.org/10.1103/PhysRevD.110.025011)

I. INTRODUCTION

In physical systems with many-body complexity, theoretical progress relies on finding small parameters which enable a perturbative expansion of physical observables. These small parameters may arise from ratios of characteristic physical scales as in EFT constructions, as well as from the large dimensionality of group representations or large particle number. For momentum transfers that are small compared to the pion mass, nuclear physics can be described efficiently by an EFT of contact operators [1–3]. As s -wave scattering lengths in the nucleon-nucleon system are very large compared to the range of the interaction, this EFT is nonperturbative in the scattering length with higher-order terms in the effective range expansion treated as perturbative insertions in the EFT. If one considers a system of neutrons only, the neutron-neutron scattering length is near infinite as compared to the pion Compton wavelength, and the leading order in the nuclear EFT is very near unitarity, where the theory is at a fixed point of the renormalization group (RG) and is therefore described

by a nonrelativistic conformal field theory, constrained by Schrödinger symmetry. While this EFT is significantly constrained by its proximity to unitarity [4,5], the machinery of conformal field theory has not found many useful applications in this EFT of nucleon contact operators.

This situation has recently changed with work of Hammer and Son [6] (See also Refs. [7–9]) which has shown that in special nuclear reactions with some number of low-energy neutrons in the final state, nonrelativistic conformal field theory technology can be used to compute the part of the nuclear reaction cross sections which account for neutron final-state interactions. This EFT for nuclear reactions has been dubbed “unnuclear” physics (in analogy with the related “unparticle” physics of Ref. [10]), as the object that describes the propagating nucleus does not have a particle interpretation. In the EFT description, the leading order is a nonrelativistic conformal field theory, whose correlation functions are constrained by Schrödinger symmetry [5,11–14]. A powerful tool for such theories is the state-operator correspondence [15], which allows the simple computation of the conformal dimensions of operators that appear in the correlation functions by considering the energy eigenspectrum of the system in a harmonic trap. Schrödinger symmetry breaking effects due to a finite scattering length and effective range can be included in conformal perturbation theory [9]. This method has now been applied to nuclear reactions with up to three neutrons

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

as one of the reaction products. However, as pointed out in Ref. [9], the analysis of final states with more than three neutrons is constrained by the complexity of the few-body wave functions, and, for systems with many neutrons, it may be profitable to consider the large-charge expansion [16,17] for nonrelativistic systems.

Based on earlier work with relativistic superfluids (for a review, see Refs. [18,19]), the large-charge expansion is formulated in the superfluid phase where the fundamental degree of freedom is the Goldstone boson of spontaneously broken particle number [20,21]. Working in sectors of large global charge has been shown to lead to important simplifications and allows the computation of the conformal data in strongly coupled relativistic CFTs [18,19]. Also in the nonrelativistic case, the conformal dimensions of charged operators have been computed to several non-trivial orders, making use of the state-operator correspondence [16,22–26]. A primary goal of this paper is to verify results based on the state-operator correspondence via a direct evaluation of the path integral, with no input from the state-operator correspondence. It turns out to be straightforward to find the large-charge “master field” solution of the Goldstone field which generates all n -point functions with large-charge Q operators. A second goal of this paper is to use the large-charge solution to compute Schrödinger-symmetry breaking corrections within the large-charge expansion. In particular, Schrödinger symmetry breaking corrections to correlation functions due to a finite scattering length in the fundamental theory of fermions are obtained. These corrections are found to take a universal form in the large-charge limit, and their size is estimated using both quantum Monte Carlo simulations, and the large- N expansion, where N is the number of fundamental fermion flavors [26]. Inclusion of effective range and other shape parameter effects in the fundamental theory, which are essential for quantifying Schrödinger symmetry breaking effects in neutron matter, require hadronic modeling, and will be left to a separate publication.

This paper is organized as follows. Section II reviews an EFT of fermions which exhibits two RG fixed points which are in correspondence with the ideal Fermi gas and with the unitary Fermi gas. This EFT near the unitary fixed point describes neutrons interacting at small momentum transfers. Since the unitary Fermi gas is superfluid, it is also described by an EFT whose degree of freedom is the Goldstone boson of spontaneously broken particle number. Aspects of this EFT that are relevant for what follows are also reviewed in this section. Section III focuses on the evaluation of nonrelativistic conformal field theory correlation functions. Schrödinger symmetry constraints on two- and three-point functions are reviewed, and the technique of evaluating correlation functions via operator insertions is introduced using the example of a two-point correlation function of noninteracting bosons. As a necessary prelude to what follows, the Euclidean formulation of the superfluid

EFT, which contains various subtleties, is developed. Then, the large-charge master field solution is found and the two-, three- and general n -point correlation functions with large-charge insertions are evaluated in the large charge limit, finding results consistent with the state-operator correspondence. In Sec. IV Schrödinger symmetry breaking effects are considered. First, these effects are formulated in the fundamental fermion theory, and then through a simple spurion analysis are matched to operators in the superfluid EFT. The quantitative size of these effects is shown to follow from quantum Monte Carlo simulations. Then, the basic structure of the symmetry-breaking corrections is shown to be constrained by Schrödinger symmetry, and, the symmetry breaking terms are derived explicitly with the large-charge solution. Finally, in Sec. V, a second estimate of the size of the symmetry-breaking corrections is provided by considering the large- N expansion at large charge. Section VI summarizes and concludes.

II. NONRELATIVISTIC CONFORMAL FIELD THEORIES

A. Fermions at unitarity

Consider a system of spin-1/2 fermions which interact via two-body contact forces. At very low energies, where derivative interactions can be ignored, the Lagrange density takes the Galilean invariant form

$$\mathcal{L} = \psi_\sigma^\dagger \left[i\partial_t + \frac{\nabla^2}{2M} \right] \psi_\sigma - \frac{1}{2} C_0 (\psi_\sigma^\dagger \psi_\sigma)^2, \quad (2.1)$$

where the field ψ_σ^\dagger creates a fermion of spin $\sigma = \uparrow, \downarrow$ and C_0 is a bare low-energy constant. The position-space interaction is a delta function at the origin, and therefore the fermion interactions are highly singular at short distances.

Now consider fermion-fermion scattering. Below inelastic thresholds, the s-wave phase shift is given by the effective range expansion

$$k \cot \delta(k) = -\frac{1}{a} + \frac{1}{2} r k^2 + \mathcal{O}(k^4), \quad (2.2)$$

where k is the center-of-mass momentum, a is the scattering length and r is the effective range. The EFT described by Eq. (2.1) requires regularization and renormalization. In dimensional regularization with the PDS scheme [2,3] and renormalized at the scale μ , the relation between the low-energy constant C_0 and the scattering length is given by

$$C_0(\mu) = \frac{4\pi}{M} \frac{1}{1/a - \mu}. \quad (2.3)$$

There is a fixed point at $C_0 = 0$, corresponding to free particles ($a = 0$), and a fixed point at $C_0 = C_\star$ corresponding

to a divergent scattering length (unitarity). It is convenient to rescale the couplings to $\hat{C}_0 \equiv C_0/C_\star$. The beta-function for the rescaled coupling is then

$$\hat{\beta}(\hat{C}_0) = \mu \frac{d}{d\mu} \hat{C}_0(\mu) = -\hat{C}_0(\mu)(\hat{C}_0(\mu) - 1), \quad (2.4)$$

which has fixed points at $\hat{C}_0 = 0$ (free fermions) and 1 (fermions at unitarity).

It is convenient to rewrite the Lagrangian with an auxiliary field s as [9]

$$\mathcal{L} = \psi_\sigma^\dagger \left[i\partial_t + \frac{\vec{\nabla}^2}{2M} \right] \psi_\sigma + \frac{1}{C_0} s^\dagger s + \psi_\downarrow^\dagger \psi_\uparrow^\dagger s + s^\dagger \psi_\uparrow \psi_\downarrow. \quad (2.5)$$

Then, at unitarity, the system of low-energy fermions is described by a nonrelativistic conformal field theory, defined by the Lagrange density

$$\begin{aligned} \mathcal{L}_{\text{CFT}} = & \psi_\sigma^\dagger \left[i\partial_t + \frac{\vec{\nabla}^2}{2M} \right] \psi_\sigma + \frac{1}{C_\star} s^\dagger s + \psi_\downarrow^\dagger \psi_\uparrow^\dagger s \\ & + s^\dagger \psi_\uparrow \psi_\downarrow. \end{aligned} \quad (2.6)$$

Now consider a gas of fermions. At the noninteracting fixed point ($C_0 = 0$), the energy per particle is

$$E/N = \frac{3}{5} \frac{k_F^2}{2M}, \quad (2.7)$$

where k_F is the Fermi momentum. At the interacting fixed point (unitarity) there is no new scale and therefore

$$E/N = \frac{3}{5} \frac{k_F^2}{2M} \xi, \quad (2.8)$$

where ξ is a dimensionless parameter, known as the Bertsch parameter, that must be determined by numerical simulation or via experiment (see section IV A for details).

B. Superfluid EFT at unitarity

As the unitary limit can be approached from both negative and positive values of a , it is natural to wonder: what phase does the gas of unitary fermions find itself in the far infrared? In nuclear physics the neutron-neutron interaction is attractive, the scattering length is negative, and the neutron gas is superfluid. More generally, in atomic systems the s-wave scattering length a can be tuned at will via external magnetic fields near a Feshbach resonance, and one can smoothly crossover from a fermionic superfluid BCS state ($a < 0$) of long-range Cooper pairs to a bosonic superfluid BEC state ($a > 0$) of tightly bound, repulsive dimers at the unitary fixed point. Remarkably these seemingly very different physical systems are all described by the same superfluid EFT in the infrared [20,21].

The order parameter $\langle \psi\psi \rangle$ breaks the U(1) particle-number symmetry, giving rise to a Goldstone boson excitation, $\theta(x)$, which can be defined as (one half of) the phase of the condensate,

$$\langle \psi\psi \rangle = |\langle \psi\psi \rangle| e^{-2i\theta}. \quad (2.9)$$

A fundamental Galilean-invariant building block in the superfluid EFT is the field

$$X = D_t\theta - \frac{(\partial_i\theta)^2}{2M}, \quad (2.10)$$

where $D_t\theta = \dot{\theta} - A_0$, with $\dot{\theta} = \partial_t\theta$ and A_0 is an external field. The effect of an external trapping potential can be included by setting $A_0 = M\omega^2 r^2/2$. It is straightforward to find the leading-order (LO) Lagrange density corresponding to fermions at unitarity [21,27],

$$\mathcal{L}_{\text{LO}} = c_0 M^{3/2} X^{5/2}, \quad (2.11)$$

where c_0 is a low-energy constant. Note that assigning time, t , scaling dimension -2 , and space, x_i , scaling dimension -1 , i.e.,

$$(t, x_i) \rightarrow (e^{2\alpha}t, e^\alpha x_i), \quad (2.12)$$

a scale-invariant action implies a Lagrange density with scaling dimension 5. As X has scaling dimension 2, the Lagrange density of Eq. (2.11) is scale invariant, and can be further shown to respect the full Schrödinger symmetry [21].

The equation of motion (EOM) for θ is

$$\partial_t(X^{3/2}) = \frac{1}{M} \partial_i(\partial_i\theta X^{3/2}). \quad (2.13)$$

In spite of the apparently simple form, this is a highly nonlinear equation.

The simplest nontrivial solution $\theta = \mu t$ gives the ground state of a system with chemical potential μ . If one expands as

$$\theta(x) = \mu t - \phi(x), \quad (2.14)$$

the fluctuations around ϕ correspond to phonon excitations. The density of the system is

$$\rho = \frac{\delta\mathcal{L}}{\delta X} = \frac{5}{2} c_0 M^{3/2} X^{3/2}, \quad (2.15)$$

and one sees that the EOM takes the form of the continuity equation for the particle number symmetry:

$$\partial_t\rho - \frac{1}{M} \partial_i(\partial_i\theta\rho) = 0. \quad (2.16)$$

Identifying the Lagrange density at the ground-state solution with the grand potential, $\Omega(\mu)$, and matching with Eq. (2.8), one finds

$$c_0 = \frac{2^{5/2}}{15\pi^2 \xi^{3/2}}. \quad (2.17)$$

The Hamiltonian density is related to the Lagrange density in the usual way,

$$\mathcal{H} = \dot{\theta} \frac{\partial \mathcal{L}}{\partial \dot{\theta}} - \mathcal{L} = \left(X - A_0 - \frac{(\partial_i \theta)^2}{2M} \right) \rho - \mathcal{L}. \quad (2.18)$$

The classical solution $\phi = 0$, $X = \mu$ with $A_0 = 0$ then recovers the standard thermodynamic relation between the energy density and the grand potential.

III. CORRELATION FUNCTIONS AT LARGE CHARGE

A. Schrödinger symmetry constraints

Schrödinger symmetry places strong constraints on the form of the correlation functions that can be computed in the EFT [5,11–14]. Consider the N -point correlation function of scalar primary operators, \mathcal{O}_i ,

$$G(x_1, x_2, \dots, x_N) = -i \langle 0 | T(\mathcal{O}_1(x_1) \mathcal{O}_2(x_2) \dots \mathcal{O}_N(x_N)) | 0 \rangle. \quad (3.1)$$

In Euclidean space, Schrödinger symmetry constrains the form of the two-point function as [11–14]

$$G(x_1, x_2) = \delta_{Q_1, Q_2} \delta_{\Delta_1, \Delta_2} \theta(\tau_{12}) \tau_{12}^{-\Delta_1} \exp\left(-\frac{Q_1 M \mathbf{x}_{12}^2}{2\tau_{12}}\right) \Psi_2 + \dots, \quad (3.2)$$

where Ψ_2 is a constant normalization factor, and the dots corresponds to the other time ordering. The notation is given by $\tau_{ij} \equiv \tau_i - \tau_j$, $\mathbf{x}_{ij} \equiv \mathbf{x}_i - \mathbf{x}_j$. Similarly, the three-point correlation function of scalar primary operators takes the form

$$G(x_1, x_2, x_3) = \delta_{Q_1+Q_2, -Q_3} \theta(\tau_{13}) \theta(\tau_{23}) \tau_{13}^{-\Delta_{13,2}/2} \tau_{23}^{-\Delta_{23,1}/2} \tau_{12}^{-\Delta_{12,3}/2} \times \exp\left(-\frac{Q_1 M \mathbf{x}_{13}^2}{2\tau_{13}} - \frac{Q_2 M \mathbf{x}_{23}^2}{2\tau_{23}}\right) \Psi_3(v_{123}) + \dots, \quad (3.3)$$

where Ψ_3 is a nonuniversal function of the Schrödinger-invariant variable v_{123} , which is given by

$$v_{123} \equiv \frac{1}{2} \frac{(\mathbf{x}_{13} \tau_{23} - \mathbf{x}_{23} \tau_{13})^2}{\tau_{12} \tau_{13} \tau_{23}} = \frac{1}{2} \left(\frac{\mathbf{x}_{23}^2}{\tau_{23}} + \frac{\mathbf{x}_{12}^2}{\tau_{12}} - \frac{\mathbf{x}_{13}^2}{\tau_{13}} \right), \quad (3.4)$$

and $\Delta_{ij,k} \equiv \Delta_i - \Delta_j - \Delta_k$. The Ward identity constraints become less and less strong for higher n -point correlators, which include nonuniversal functions of the $(n-2)(n-1)/2$ Schrödinger-invariant variables that generalize the v_{123} [28].

If the correlation functions describe free-particle propagation, then the conformal dimensions are given by their naive scaling values. However, for interacting nonrelativistic conformal field theories, the conformal dimensions are generally strong-interaction physics and must be obtained from either the state-operator correspondence or an exact evaluation of the correlator.

B. Free theory with operator insertions

A useful tool for evaluating correlation functions in the semi-classical approximation involves exponentiating the operator insertions and defining a new effective action, which in turn is evaluated at its saddle point to generate the correlation functions. This method has been used in the context of the large-charge expansion in Refs. [19,29–32]. Here, as an example for what follows, it will be applied to the two-point function of free, nonrelativistic, boson operators. The correlation function of interest is

$$G(x_1, x_2) = -i \langle 0 | T(\mathcal{O}_Q(x_1) \mathcal{O}_Q^\dagger(x_2)) | 0 \rangle, \quad (3.5)$$

where \mathcal{O}_Q^\dagger is a primary operator of number charge $Q \equiv Q_{\mathcal{O}^\dagger} = -Q_{\mathcal{O}} \geq 0$, and scaling dimension Δ_Q . Assuming that $\mathcal{O}_Q | 0 \rangle = 0$, and $x_1 = (\tau_1, \mathbf{x}_1)$, $x_2 = (\tau_2, \mathbf{x}_2)$, one can write

$$G(x_1, x_2) = -i \theta(\tau_{12}) \langle 0 | \mathcal{O}_Q(\tau_1, \mathbf{x}_1) \mathcal{O}_Q^\dagger(\tau_2, \mathbf{x}_2) | 0 \rangle. \quad (3.6)$$

Consider the path integral representation of the two-point function,

$$G(x_1, x_2) = \int \mathcal{D}\phi \mathcal{D}\phi^\dagger \mathcal{O}_Q(x_1) \mathcal{O}_Q^\dagger(x_2) e^{-\int d^4x \mathcal{L}(x)}, \quad (3.7)$$

where now the Euclidean space formulation¹ of the free theory will be used, with a chemical potential μ ,

$$\mathcal{L} = \phi^\dagger \left[\partial_\tau - \frac{\vec{\nabla}^2}{2M} - \mu \right] \phi, \quad (3.8)$$

where the field ϕ^\dagger creates a boson with unit charge. An operator of charge Q can be written as

$$\mathcal{O}_Q = \mathcal{N}(\phi)^Q, \quad \mathcal{O}_Q^\dagger = \mathcal{N}(\phi^\dagger)^Q, \quad (3.9)$$

where \mathcal{N} is a normalization factor. With $\Delta_\phi = 3/2$, one has $\Delta_Q = 3Q/2$. Note that since \mathcal{O}_Q is a primary operator with

¹The Euclidean space formulation is obtained via transforming $t \rightarrow -i\tau$, accompanied by $\mathcal{L} \rightarrow -\mathcal{L}$.

$\mathcal{O}_Q|0\rangle = 0$, there is always an overall $\theta(\tau_{12})$ multiplying the two-point function which is ignored in what follows by assuming that $\tau_1 > \tau_2$.

Exponentiating the insertions gives the new action,

$$S = \int d^4x [\mathcal{L} - Q \log(\phi) \delta^4(x - x_1) - Q \log(\phi^\dagger) \delta^4(x - x_2)]. \quad (3.10)$$

Varying this action then gives the two EOM about the free fixed point,

$$\begin{aligned} \phi^\dagger \left[\partial_\tau - \frac{\vec{\nabla}^2}{2M} - \mu \right] \phi &= +Q \delta^4(x - x_2), \\ \phi \left[\partial_\tau + \frac{\vec{\nabla}^2}{2M} + \mu \right] \phi^\dagger &= -Q \delta^4(x - x_1). \end{aligned} \quad (3.11)$$

In order to solve these equations, consider the diffusion equation

$$\left[\partial_\tau - \frac{\vec{\nabla}^2}{2M} - \mu \right] \mathcal{G}(x; x_2) = \delta^4(x - x_2), \quad (3.12)$$

which is satisfied by the Green's function

$$\begin{aligned} \mathcal{G}(x; x_2) &= \theta(\tau - \tau_2) \left(\frac{M}{2\pi(\tau - \tau_2)} \right)^{3/2} \exp \left(-\frac{M(\mathbf{x} - \mathbf{x}_2)^2}{2(\tau - \tau_2)} \right) \\ &\times \exp(\mu(\tau - \tau_2)), \end{aligned} \quad (3.13)$$

and, likewise,

$$\left[\partial_\tau + \frac{\vec{\nabla}^2}{2M} + \mu \right] \mathcal{G}(x_2; x) = -\delta^4(x - x_2). \quad (3.14)$$

The solution of the EOM in the presence of the sources is thus found to be

$$\begin{aligned} \phi &= i(Q)^{1/2} \frac{\mathcal{G}(x; x_2)}{\mathcal{G}(x_1; x_2)^{1/2}}, \\ \phi^\dagger &= -i(Q)^{1/2} \frac{\mathcal{G}(x_1; x)}{\mathcal{G}(x_1; x_2)^{1/2}}. \end{aligned} \quad (3.15)$$

Note that the density is given by

$$\rho = -\frac{\delta \mathcal{L}}{\delta \mu} = \phi^\dagger \phi = Q \frac{\mathcal{G}(x_1; x) \mathcal{G}(x; x_2)}{\mathcal{G}(x_1; x_2)}, \quad (3.16)$$

which has support in the range $\tau_2 \leq \tau \leq \tau_1$. Using the identity

$$\mathcal{G}(x_1; x_2) = \int d^3\mathbf{x} \mathcal{G}(x_1; x) \mathcal{G}(x; x_2), \quad (3.17)$$

one finds

$$Q = \int d^3\mathbf{x} \rho, \quad (3.18)$$

as expected.

The action at the saddle point is

$$S = Q - Q \log \left(-\frac{1}{2} Q \mathcal{G}(x_1; x_2) \right), \quad (3.19)$$

and, finally,

$$G(x_1, x_2) = \theta(\tau_{12}) \tau_{12}^{-3Q/2} \exp \left(-\frac{MQ\mathbf{x}_{12}^2}{2\tau_{12}} \right) \Psi_2 e^{\mu Q \tau_{12}}, \quad (3.20)$$

where

$$\Psi_2 = \mathcal{N}^2 \left[\left(\frac{M}{2\pi} \right)^{3/2} e^{1-i\pi} 2Q \right]^Q. \quad (3.21)$$

This result is in the Schrödinger form of Eq. (3.2) only if $\mu = 0$, as the chemical potential is a Schrödinger-breaking effect.

Before moving on to the more interesting interacting case, consider several general remarks regarding this result. An operator of charge Q has been inserted at $\tau = \tau_2$ and then removed at $\tau = \tau_1 > \tau_2$. By construction this is the Euclidean time-ordered correlator. In this semiclassical calculation, the ordering [represented here by the Heaviside function $\theta(\tau_{12})$] comes from the fact that the EOM is a diffusion equation, for which the initial-value problem is well-posed for $\tau > 0$ (as opposed to the backward equation for which the problem is generically not well posed).

C. Two-point function at large charge

Before proceeding to computing the two-point function in the large-charge EFT, consider the Euclidean formulation of the superfluid EFT. The partition function is

$$Z = \int \mathcal{D}\theta \exp(-S) = \int \mathcal{D}\theta \exp \left(-\int d\tau d^3\mathbf{x} \mathcal{L} \right), \quad (3.22)$$

where, at LO,

$$\mathcal{L}_{\text{LO}} = -c_0 M^{3/2} X^{5/2}, \quad (3.23)$$

with

$$X = i\partial_\tau \theta - A_0 - \frac{(\partial_i \theta)^2}{2M}. \quad (3.24)$$

Hermiticity requires that θ be pure imaginary. Note that now

$$\rho = -\frac{\delta \mathcal{L}}{\delta X}, \quad \mathcal{H} = \mathcal{L} - \theta \frac{\partial \mathcal{L}}{\partial \theta}, \quad (3.25)$$

where here $\dot{\theta} \equiv \partial_\tau \theta$. The Euler-Lagrange equations take the form

$$\partial_\tau \left(i \frac{\delta \mathcal{L}}{\delta X} \right) = \frac{1}{M} \partial_i \left(\partial_i \theta \frac{\delta \mathcal{L}}{\delta X} \right) + \frac{\delta \mathcal{L}}{\delta \theta}. \quad (3.26)$$

At LO, one finds the ground-state solution

$$\theta = -i\mu\tau, \quad X = \mu - A_0. \quad (3.27)$$

For what follows, it is interesting to note that there is also a space-time dependent solution to the EOM (now taking $A_0 = 0$),

$$\theta_f(\tau, \mathbf{x}) = -i \frac{M \mathbf{x}^2}{4\tau}, \quad X_f(\tau, \mathbf{x}) = -\frac{M \mathbf{x}^2}{8\tau^2}. \quad (3.28)$$

As X_f is negative definite, this solution corresponds to an unstable configuration. As will be seen below, there is a sense in which this is the “free” solution, which is unsurprisingly not stable in the superfluid EFT.

The saddle configuration. Now, take $A_0 = 0$. In the EFT every operator can be written as a function of the Goldstone field by simply identifying its quantum numbers. It follows that a generic primary operator with charge Q and dimension Δ must have the form

$$\mathcal{O}_{\Delta, Q} = \mathcal{N} \left(\frac{2}{\gamma} \right)^{\Delta/2} X^{\Delta/2} \exp(iQ\theta), \quad (3.29)$$

where \mathcal{N} is a normalization constant, and the other constant factor has been chosen for future convenience and will be defined below. Define the two-point function

$$G_Q(x_1, x_2) = \int \mathcal{D}\theta \mathcal{O}_{\Delta, Q}(x_2) \mathcal{O}_{\Delta, -Q}(x_1) e^{-\int d^4x \mathcal{L}_I}, \quad (3.30)$$

where \mathcal{L}_I is the interaction Lagrange density. In this semiclassical computation, only one time ordering will appear as a consequence of the fact that the EOM are not time-reversal invariant, as was already observed in the free case above. In what follows, it will be assumed that $\tau_1 > \tau_2$.

The operator insertions can be exponentiated to give

$$G_Q(x_1, x_2) = \mathcal{N}^2 \left(\frac{2}{\gamma} \right)^\Delta \int \mathcal{D}\theta e^{-\int d^4x \mathcal{L}}, \quad (3.31)$$

where

$$\begin{aligned} \mathcal{L} = & \mathcal{L}_I - \frac{\Delta}{2} \log X [\delta^4(x - x_2) + \delta^4(x - x_1)] \\ & - iQ\theta [\delta^4(x - x_2) - \delta^4(x - x_1)]. \end{aligned} \quad (3.32)$$

Now the saddle point solution must be found in the presence of the sources. The Euler-Lagrange equation is

$$\partial_\tau \rho + \frac{1}{M} \partial_i (i \partial_i \theta \rho) = Q [\delta^4(x - x_2) - \delta^4(x - x_1)]. \quad (3.33)$$

This is the continuity equation in the presence of a source. It may appear confusing that there are now two distinct expressions for the density. First, there is the density as an observable quantity related to the specific Lagrange density which describes the system, given by Eq. (3.25), and which satisfies Eq. (3.33) in the presence and absence of sources. On the other hand, there is the density—proportional to Q —which solves Eq. (3.33) in the presence of sources and vanishes in the absence of sources. This second density is a distribution and is not directly equal to the observable density. However, the densities integrated over space must be equal, and indeed that equality fixes the charge Q of the superfluid system.

In order to solve Eq. (3.33) for $\theta = \theta(\tau, \mathbf{x})$, consider the ansatz

$$i \partial_i \theta \epsilon \rho_\epsilon = \epsilon \frac{1}{2} \partial_i \rho_\epsilon, \quad (3.34)$$

where $\epsilon = \pm$. This equation is readily integrated to give

$$\rho_\epsilon = f_\epsilon(\tau) \exp(2i\epsilon\theta), \quad (3.35)$$

where $f_\epsilon(\tau)$ is an arbitrary real function of τ .

Now plugging the ansatz, Eq. (3.34), into the EOM gives

$$\left[\partial_\tau + \epsilon \frac{\vec{\nabla}^2}{2M} \right] \rho_\epsilon = Q [\delta^4(x - x_2) - \delta^4(x - x_1)], \quad (3.36)$$

and it is clear that the ansatz restricts the initial highly nonlinear EOM to a diffusion equation, whose solution has been shown above to be Schrödinger invariant. Noting that the diffusion equation

$$\left[\partial_\tau - \frac{\vec{\nabla}^2}{2M} \right] \mathbb{G}(x; x') = \delta^4(x - x') \quad (3.37)$$

is solved by the Green's function

$$\mathbb{G}(x; x') = \theta(\tau - \tau') \left(\frac{M}{2\pi(\tau - \tau')} \right)^{3/2} \exp \left(-\frac{M(\mathbf{x} - \mathbf{x}')^2}{2(\tau - \tau')} \right), \quad (3.38)$$

and

$$\left[\partial_\tau + \frac{\vec{\nabla}^2}{2M} \right] \mathbb{G}(x'; x) = -\delta^4(x - x'), \quad (3.39)$$

one finds the solutions

$$\begin{aligned} \rho_- &= +Q[\mathbb{G}(x; x_2) - \mathbb{G}(x; x_1)], \\ \rho_+ &= -Q[\mathbb{G}(x_2; x) - \mathbb{G}(x_1; x)]. \end{aligned} \quad (3.40)$$




FIG. 1. Euclidean timeline with source charge placements (above line) and charge regions (below line).

Integrating both sides over space gives

$$\begin{aligned} \int d^3\mathbf{x} \rho_- &= +Q[\theta(\tau - \tau_2) - \theta(\tau - \tau_1)], \\ \int d^3\mathbf{x} \rho_+ &= -Q[\theta(\tau_2 - \tau) - \theta(\tau_1 - \tau)]. \end{aligned} \quad (3.41)$$

And, as expected,

$$Q = \frac{1}{(\tau_1 - \tau_2)} \int_{\tau_2+\varepsilon}^{\tau_1-\varepsilon} d^3\mathbf{x} \rho_{\pm}. \quad (3.42)$$

The temporal structure of the solution is illustrated in Fig. 1, which makes clear that the solution has nonvanishing charge only in the interval between τ_2 and τ_1 .

Consider the solution ρ_- in the range $\tau_2 + \varepsilon < \tau < \tau_1 - \varepsilon$. One has

$$\begin{aligned} \rho_- &= Q \left(\frac{M}{2\pi(\tau - \tau_2)} \right)^{3/2} \exp \left(-\frac{M(\mathbf{x} - \mathbf{x}_2)^2}{2(\tau - \tau_2)} \right) \\ &= f_-(\tau) \exp(-2i\theta_-). \end{aligned} \quad (3.43)$$

Now scale invariance suggests the power law form

$$f_-(\tau) = d \left(\frac{M}{2\pi(\tau - \tau_2)} \right)^{3/2-\gamma}, \quad (3.44)$$

where d is an irrelevant dimensionless constant that contributes to the overall normalization of the Green's function, and is here chosen for convenience to be $d = Q$, and γ is an anomalous dimension.² Now if $\gamma = 0$, then one finds from Eq. (3.43) the unstable solution $\theta_- = \theta_f$, given in Eq. (3.28). It is in this sense that θ_f may be thought of as the “free” solution. Therefore, one should expect superfluidity to be triggered by $\gamma \neq 0$. In this case, one finds from Eq. (3.43) the Goldstone field

$$\theta_- = \frac{i}{2}\gamma \log \left(\frac{M}{2\pi(\tau - \tau_2)} \right) - i \frac{M(\mathbf{x} - \mathbf{x}_2)^2}{4(\tau - \tau_2)}. \quad (3.45)$$

Next consider the solution ρ_+ in the range $\tau_2 + \varepsilon < \tau < \tau_1 - \varepsilon$. One has

² γ should not be confused with Δ , the conformal dimension that is computed below.

$$\begin{aligned} \rho_+ &= Q \left(\frac{M}{2\pi(\tau_1 - \tau)} \right)^{3/2} \exp \left(-\frac{M(\mathbf{x} - \mathbf{x}_1)^2}{2(\tau_1 - \tau)} \right) \\ &= f_+(\tau) \exp(2i\theta_+), \end{aligned} \quad (3.46)$$

where now

$$f_+(\tau) = d \left(\frac{M}{2\pi(\tau_1 - \tau)} \right)^{3/2-\gamma}. \quad (3.47)$$

And the second solution for the Goldstone field is

$$\theta_+ = -\frac{i}{2}\gamma \log \left(\frac{M}{2\pi(\tau_1 - \tau)} \right) + i \frac{M(\mathbf{x} - \mathbf{x}_1)^2}{4(\tau_1 - \tau)}. \quad (3.48)$$

Both solutions, Eq. (3.45) and Eq. (3.48), satisfy the homogeneous EOM. However, the inhomogeneous solution, and therefore the general solution, must depend on both source locations. Indeed, one expects that the solution will be invariant with respect to $Q \rightarrow -Q$ and $x_2 \leftrightarrow x_1$. The only linear combination of θ_- and θ_+ which is invariant with respect to $x_2 \leftrightarrow x_1$ is the sum³ $\theta_- + \theta_+$, and indeed the sum is evidently the only linear combination that solves the homogeneous EOM.

Finally, the general solution to the EOM and the saddle point location in the presence of sources is given by

$$\theta_s(\tau, \mathbf{x}) = \frac{i}{2}\gamma \log \left(\frac{\tau_1 - \tau}{\tau - \tau_2} \right) - \frac{i}{4}M \left[\frac{(\mathbf{x} - \mathbf{x}_2)^2}{(\tau - \tau_2)} - \frac{(\mathbf{x} - \mathbf{x}_1)^2}{(\tau_1 - \tau)} \right]. \quad (3.49)$$

As will be seen below, this is the large-charge master field. The solution is well-defined and physical for $\tau_2 < \tau < \tau_1$. Just as in the free case, the structure of the EOM imposes a time ordering. To arrive at this solution, the assumption of Schrödinger invariance has entered via the diffusion-equation ansatz of Eq. (3.34). However, it is straightforward to see that the master field solution is present in the Schrödinger algebra, and this will be shown in detail below.⁴ From the saddle solution, one finds

$$X[\theta_s] = \frac{1}{2}\gamma \frac{(\tau_1 - \tau_2)}{(\tau - \tau_2)(\tau_1 - \tau)} - \frac{1}{8}M \left[\frac{(\mathbf{x} - \mathbf{x}_2)^2}{(\tau - \tau_2)} - \frac{(\mathbf{x} - \mathbf{x}_1)^2}{(\tau_1 - \tau)} \right]^2. \quad (3.50)$$

³Invariance follows if one also takes $\gamma \rightarrow -\gamma$, however note that the EOM is solved independently of the value of γ , so the sign of this term is irrelevant until one fixes the charge.

⁴This solution is general in the following sense. Consider a Schrödinger-invariant system and let θ be the field that transforms with a shift under the action of the particle number operator. Then there always exists a solution to the EOM in which θ has the form in Eq. (3.49). For example, for the free theory, a special case of this solution has been found in [33] as an instanton which describes the decay of a metastable bosonic droplet.

It is convenient to shift the spatial variable to

$$\mathbf{x} = \mathbf{y} + \mathbf{x}_2 \left(\frac{\tau_1 - \tau}{\tau_1 - \tau_2} \right) + \mathbf{x}_1 \left(\frac{\tau - \tau_2}{\tau_1 - \tau_2} \right). \quad (3.51)$$

It then follows that

$$\begin{aligned} \mathbf{x} - \mathbf{x}_2 &= \mathbf{y} + \mathbf{x}_{12} \left(\frac{\tau - \tau_2}{\tau_1 - \tau_2} \right), \\ \mathbf{x} - \mathbf{x}_1 &= \mathbf{y} - \mathbf{x}_{12} \left(\frac{\tau_1 - \tau}{\tau_1 - \tau_2} \right), \end{aligned} \quad (3.52)$$

and the spatial delta functions become

$$\delta^4(x - x_i) = \delta^3(\mathbf{y}) \delta(\tau - \tau_i). \quad (3.53)$$

In the new coordinate,

$$\begin{aligned} \theta_s &= \frac{i}{2} \gamma \log \left(\frac{\tau_1 - \tau}{\tau - \tau_2} \right) - \frac{i}{4} M \left[\frac{1}{\tau - \tau_2} \left(\mathbf{y} + \mathbf{x}_{12} \left(\frac{\tau - \tau_2}{\tau_1 - \tau_2} \right) \right)^2 \right. \\ &\quad \left. - \frac{1}{\tau_1 - \tau} \left(\mathbf{y} - \mathbf{x}_{12} \left(\frac{\tau_1 - \tau}{\tau_1 - \tau_2} \right) \right)^2 \right] \end{aligned} \quad (3.54)$$

and

$$X = \frac{1}{2} \gamma \frac{(\tau_1 - \tau_2)}{(\tau - \tau_2)(\tau_1 - \tau)} - \frac{1}{8} M \left[\frac{(\tau_1 - \tau_2)}{(\tau - \tau_2)(\tau_1 - \tau)} \right]^2 \mathbf{y}^2. \quad (3.55)$$

Note that now all dependence of the correlation function on the spatial separation is contained in the Q insertion in the action. The emergent harmonic trap is now seen by writing

$$X = \bar{\mu}(\tau) - \frac{1}{2} M \bar{\omega}(\tau)^2 r^2, \quad (3.56)$$

where $r = |\mathbf{y}|$, and

$$\begin{aligned} \bar{\omega}(\tau) &= \frac{1}{2} \frac{(\tau_1 - \tau_2)}{(\tau - \tau_2)(\tau_1 - \tau)}, \\ \bar{\mu}(\tau) &= \frac{1}{2} \gamma \frac{(\tau_1 - \tau_2)}{(\tau - \tau_2)(\tau_1 - \tau)}. \end{aligned} \quad (3.57)$$

Here, it is clear that the time-dependent function $\bar{\mu}(\tau)$, via the anomalous dimension $\gamma > 0$, drives spontaneous symmetry breaking.⁵

The saddle from the state-operator correspondence. An alternative derivation of the saddle solution, this time based on Schrödinger invariance is obtained by comparing two equivalent frames:

- (i) The Galileian frame with coordinates (τ, \mathbf{x}) that was used above, in which there is no background

potential and the sources are inserted at finite Euclidean time, and

- (ii) the oscillator frame, with coordinates $(\tilde{\tau}, \tilde{\mathbf{x}})$ in which the sources are placed at $\tilde{\tau} = \pm\infty$ and a background field $A_0 = M\omega^2 \mathbf{y}^2/2$ is present.

These are the two sides of the nonrelativistic state-operator correspondence [34], in which the conformal dimension Δ of a primary in the Galileian frame is mapped to the energy in units of ω , \tilde{E}/ω of a state in the oscillator frame.

The two frames are completely equivalent for Schrödinger-invariant systems and are related by

$$\begin{cases} \omega\tau = \tanh(\omega\tilde{\tau}), \\ \mathbf{x} = \tilde{\mathbf{x}} \frac{1}{\cosh(\omega\tilde{\tau})} \end{cases}, \quad \begin{cases} \omega\tilde{\tau} = \operatorname{arctanh}(\omega\tau), \\ \tilde{\mathbf{x}} = \frac{\mathbf{x}}{\sqrt{1-\omega^2\tau^2}}. \end{cases} \quad (3.58)$$

Insertions at $\tilde{\tau} = \pm\infty$ are then mapped to $\tau = \pm 1/\omega$. However, since the map is singular at these points, such insertions in the oscillator frame are always mapped to $\mathbf{x} = 0$ in the Galileian frame. Then the source points are always at $\mathbf{x}_1 = \mathbf{x}_2 = 0$, and therefore $\mathbf{x} = \mathbf{y}$. In addition, $M = 1$ in this section.

Primary operators of dimension Δ and charge Q are related by

$$\begin{aligned} \mathcal{O}_{\Delta,Q}(\tau, \mathbf{x}) &= (\cosh(\omega\tilde{\tau}))^\Delta \exp \left[-\frac{Q\omega}{2} \tanh(\omega\tilde{\tau}) \tilde{\mathbf{x}}^2 \right] \\ &\quad \times \tilde{\mathcal{O}}_{\Delta,Q}(\tilde{\tau}, \tilde{\mathbf{x}}), \end{aligned} \quad (3.59)$$

$$\tilde{\mathcal{O}}_{\Delta,Q}(\tilde{\tau}, \tilde{\mathbf{x}}) = (1 - \omega^2 \tau^2)^{\Delta/2} \exp \left[\frac{Q\omega^2}{2} \frac{\mathbf{x}^2 \tau}{1 - \omega^2 \tau^2} \right] \mathcal{O}_{\Delta,Q}(\tau, \mathbf{x}). \quad (3.60)$$

The Goldstone field transforms nonlinearly under Schrödinger symmetry, so to see how the ground state solution $\tilde{\theta}(\tilde{\tau}, \tilde{\mathbf{x}}) = -i\mu\tilde{\tau}$ is mapped to $\theta(\tau, \mathbf{x})$, take a generic primary in terms of the Goldstone field and use the map above:

$$\begin{aligned} \tilde{\mathcal{O}}_{\Delta,Q}(\tilde{\tau}, \tilde{\mathbf{x}}) &= \tilde{X}(\tilde{\tau}, \tilde{\mathbf{x}})^{\Delta/2} e^{iQ\tilde{\theta}(\tilde{\tau}, \tilde{\mathbf{x}})} \mapsto \mathcal{O}_{\Delta,Q}(\tau, \mathbf{x}) \\ &= X(\tau, \mathbf{x})^{\Delta/2} e^{iQ\theta(\tau, \mathbf{x})}. \end{aligned} \quad (3.61)$$

Then one finds

$$\begin{aligned} \tilde{\theta} = -i\mu\tilde{\tau} &\mapsto \theta \\ &= i\frac{\mu}{2\omega} \log \left(\frac{1/\omega - \tau}{\tau + 1/\omega} \right) - \frac{i}{4} \left(\frac{\mathbf{x}^2}{\tau + 1/\omega} - \frac{\mathbf{x}^2}{1/\omega - \tau} \right), \end{aligned} \quad (3.62)$$

$$\begin{aligned} \tilde{X} &= \mu - \frac{\omega^2}{2} \tilde{\mathbf{x}}^2 \mapsto X \\ &= \frac{\mu}{2\omega} \left(\frac{1}{1/\omega - \tau} + \frac{1}{\tau + 1/\omega} \right) \\ &\quad - \frac{1}{8} \left(\frac{\mathbf{x}}{\tau + 1/\omega} + \frac{\mathbf{x}}{1/\omega - \tau} \right)^2, \end{aligned} \quad (3.63)$$

⁵Despite the suggestive notation, this function should not be viewed as a time-dependent chemical potential.

which is the field configuration generated by two insertions at $(\tau = \pm 1/\omega, \mathbf{x} = 0)$. These expressions are consistent with those found working directly in the Galileian frame with the identification

$$\gamma = \frac{\mu}{\omega} = \frac{\mu}{2} \tau_{12}. \quad (3.64)$$

Evaluation of the 2-point function. The density is given by

$$\rho = -\frac{\delta \mathcal{L}}{\delta X} = -\frac{\delta \mathcal{L}_I}{\delta X} + \frac{\Delta}{2} X^{-1} [\delta^4(x - x_2) + \delta^4(x - x_1)]. \quad (3.65)$$

Now, using Eq. (3.53) and Eq. (3.55), it is easy to see that the second term has no support and therefore gives a vanishing contribution to the density. Therefore, taking

$$\mathcal{L}_I = \mathcal{L}_{LO} = -c_0 M^{3/2} X^{5/2}, \quad \rho = \frac{5}{2} c_0 M^{3/2} X^{3/2}, \quad (3.66)$$

it is straightforward to check that ρ satisfies the homogeneous EOM at the saddle-point solution, Eq. (3.49).

To satisfy the EOM in the presence of sources, the density at the saddle-point solution must be integrated over all space to fix the charge. Now note that the density vanishes when $r \rightarrow \bar{R}(\tau)$ with

$$\bar{R}(\tau) = \sqrt{\frac{2\bar{\mu}}{M} \frac{1}{\bar{\omega}}} = \frac{2}{\sqrt{M}} \left(\gamma \frac{(\tau - \tau_2)(\tau_1 - \tau)}{(\tau_1 - \tau_2)} \right)^{1/2}. \quad (3.67)$$

Hence, in the presence of the sources the superfluid system behaves like a droplet whose size is governed by the Euclidean-time separation between sources. Figure 2

illustrates the droplet radius forming at the source τ_2 , rising to its maximum at radius $\tau_{12}/2$, and returning to zero at the sink, τ_1 .

One now easily finds

$$\int d^3 \mathbf{x} \rho = 4\pi \int_0^{\bar{R}(\tau)} dr r^2 \rho = \frac{\bar{\mu}^3}{3\bar{\omega}^3 \xi^{3/2}} = \frac{1}{3\xi^{3/2}} \gamma^3 = Q. \quad (3.68)$$

Therefore, the anomalous dimension is related to the charge by

$$\gamma = 3^{1/3} \xi^{1/2} Q^{1/3}, \quad (3.69)$$

and

$$\bar{\mu}(\tau) = 3^{1/3} \xi^{1/2} Q^{1/3} \bar{\omega}(\tau). \quad (3.70)$$

Having found the solution to the EOM, the two-point function at the saddle is straightforwardly evaluated:

$$G_Q(x_1, x_2) = \mathcal{O}_{\Delta, Q}(x_2) \mathcal{O}_{\Delta, -Q}(x_1) e^{-S[\theta]}|_{\theta=\theta_s} + \dots, \quad (3.71)$$

where the dots are subleading effects in the large-charge EFT (see below). Because of the insertions at $\tau = \tau_2$ and $\tau = \tau_1$, it is convenient to view this expression as

$$G_Q(x_1, x_2) = \lim_{\varepsilon \rightarrow 0} \mathcal{O}_{\Delta, Q}(\tau_2 + \varepsilon, \mathbf{x}_2) \mathcal{O}_{\Delta, -Q}(\tau_1 - \varepsilon, \mathbf{x}_1) \times \exp \left[- \int_{\tau_2 + \varepsilon}^{\tau_1 - \varepsilon} d\tau \int d^3 \mathbf{x} \mathcal{L}_I[\theta] \right] \Big|_{\theta=\theta_s}. \quad (3.72)$$

The existence of this limit will then give a constraint that will be used to compute the conformal dimension Δ .




FIG. 2. Left: plot of the radius of the superfluid droplet vs Euclidean time as described in the text. Right: equal-time slices representing the evolution of the droplet size for insertions at $(0, -1, -1)$ and $(1, 1, 1)$. Darker colors correspond to denser regions.

The only thing that remains to compute is the value of the action at the saddle:

$$\begin{aligned}
 S_I &= \int_{\tau_2+\varepsilon}^{\tau_1-\varepsilon} d\tau \int d^3\mathbf{x} \mathcal{L}_I \\
 &= -4\pi c_0 \int_{\tau_2+\varepsilon}^{\tau_1-\varepsilon} d\tau \int_0^{\bar{R}(\tau)} r^2 dr X(\tau, r)^{5/2} \\
 &= -\frac{5c_0\pi^2\gamma^4}{16\sqrt{2}} \log\left(\frac{\tau_{12}}{\varepsilon}\right) + \mathcal{O}(\varepsilon) \\
 &= -\frac{3^{1/3}\xi^{1/2}}{4} Q^{4/3} \log\left(\frac{\tau_{12}}{\varepsilon}\right) + \mathcal{O}(\varepsilon). \tag{3.73}
 \end{aligned}$$

Simply substituting the saddle solution into the expression for the operators one then finds

$$\begin{aligned}
 G_Q(x_1, x_2) &= \mathcal{N}^2 \varepsilon^{Q\gamma - \Delta - \frac{3^{1/3}\xi^{1/2}}{4} Q^{4/3}} \\
 &\times \left[\tau_{12}^{-Q\gamma + \frac{3^{1/3}\xi^{1/2}}{4} Q^{4/3}} \exp\left(-\frac{QM\mathbf{x}_{12}^2}{2\tau_{12}}\right) + \mathcal{O}(\varepsilon) \right]. \tag{3.74}
 \end{aligned}$$

Cancellation of the divergence for $\varepsilon \rightarrow 0$ requires

$$Q\gamma - \Delta - \frac{3^{1/3}\xi^{1/2}}{4} Q^{4/3} = 0 \tag{3.75}$$

and leads to the solution

$$G_Q(x_1, x_2) = \mathcal{N}^2 \tau_{12}^{-\Delta} \exp\left(-\frac{QM\mathbf{x}_{12}^2}{2\tau_{12}}\right), \tag{3.76}$$

where

$$\Delta = Q\gamma - \frac{3^{1/3}\xi^{1/2}}{4} Q^{4/3} = \frac{3^{4/3}}{4} \xi^{1/2} Q^{4/3} \tag{3.77}$$

in agreement with the state-operator correspondence large-charge result [17].

Note also that γ measures how much the conformal dimension changes when the charge changes by one unit:

$$\frac{d\Delta}{dQ} = \gamma. \tag{3.78}$$

This shows that equation (3.75) is simply a Legendre transform relating Δ to the value of the action at the saddle. In the oscillator frame, this is precisely the thermodynamic relation between the free energy $F(Q)$ and the grand potential $\Omega(\mu)$.

EFT power counting. Note that up to this point, there has been no explicit use of the large-charge expansion. However, the semi-classical expansion is used implicitly in a claim that the effective action, evaluated at the saddle, provides the large-charge solution. This is, of course, true

when Q is large, as expected. However, it is worth considering the various scales in the problem in some detail. This parallels consideration of the large-charge EFT in the presence of a harmonic trap [25]. However, in the state-operator correspondence, the trap frequency, ω , is an artificial parameter which is taken to zero as $Q \rightarrow \infty$ in a manner that leaves the density fixed. In the exact solution given here, the emergent trap frequency, $\bar{\omega}$, vanishes as the temporal separation of the sources increases. Say $\tau_1 = \Delta\tau$ and $\tau_2 = -\Delta\tau$. Then, $\Delta\tau = (\tau_1 - \tau_2)/2$, and for fixed τ and $\Delta\tau \rightarrow \infty$, one has

$$\bar{\omega} \rightarrow \Delta\tau^{-1} \rightarrow 0. \tag{3.79}$$

One also has

$$\begin{aligned}
 \bar{R} &\sim \frac{Q^{1/6}}{M^{1/2}\bar{\omega}^{1/2}}, & V &\sim \bar{R}^3 \sim \frac{Q^{1/2}}{M^{3/2}\bar{\omega}^{3/2}}, \\
 \rho &\sim \frac{Q}{V} \sim M^{3/2}\bar{\omega}^{3/2} Q^{1/2}, \tag{3.80}
 \end{aligned}$$

where V is a measure of the spatial volume occupied by the superfluid droplet. Therefore, one can take $Q \rightarrow \infty$, $\bar{\omega} \rightarrow 0$ while keeping ρ fixed if in the power counting $Q \sim M^3 \Delta\tau^3$. Then, taking the infrared (IR) scale to be $p_{\text{IR}} \equiv \bar{R}^{-1}$ and the ultraviolet (UV) scale to be $\Lambda_{\text{UV}} \equiv \rho^{1/3}$, the large-charge EFT can be organized as an expansion in the ratio $p_{\text{IR}}/\Lambda_{\text{UV}} \sim Q^{-1/3}$ which is small when Q is large. This power counting argument provides a basis for organizing operators and higher-order corrections in the large-charge EFT.

NLO corrections. The easiest way to identify the NLO corrections for the two-point function result is to write the next terms in the EFT in the oscillator frame. The reason is that there is a natural IR scale $p_{\text{IR}}^2 \sim \omega$ given by the oscillator frequency and a natural UV scale $\Lambda_{\text{UV}}^2 \sim \mu$. The leading scale-invariant terms, up to field redefinitions, can be chosen to be (with $M = 1$)

$$\begin{aligned}
 \mathcal{L}_{\text{osc}} &= c_0 X^{5/2} + c_1 X^{-1/2} (\partial_i X)^2 \\
 &+ X^{1/2} (c_2 (\partial_i \partial_j \theta)^2 + c_3 (\partial_i \partial_j \theta)^2 + c_4 \partial_i \partial_j A_0) \tag{3.81}
 \end{aligned}$$

with $A_0 = \omega^2 \mathbf{x}^2/2$. The c_i are low-energy constants that are determined by the short-distance physics. On the basis of power counting, each derivative counts as $Q^{-1/3}$, and therefore one expects NLO contributions to the scaling dimensions suppressed by $Q^{-2/3}$ as compared to the LO. This is indeed correct [17].

Fixing the charge breaks the Schrödinger symmetry and in general the EFT will not be invariant order-by-order in ω/μ . When passing to the flat frame, the c_0 and c_1 terms are separately invariant under the transformations in Eq. (3.58) and enter the flat-frame EFT as they are. As for the other terms, one can verify that the computation of the conformal

dimension in the oscillator frame and of the two-point function in the flat frame agree if the following condition is met:

$$c_4 = -3c_2 - c_3. \quad (3.82)$$

This relation agrees with the condition of conformal invariance when the LO EOM is used, as found in Ref. [35]. This is also consistent with Ref. [26], which considers the superfluid EFT at large-charge and large- N and finds that the explicit trap-dependent contribution to the c_4 operator in Eq. (3.81) is present in the Schrödinger-invariant EFT.

As is generally the case, there is large freedom in defining the fields and transformation properties in the EFT. In the literature different choices have appeared [17,21,35], but they must all ultimately agree on the resulting values for the physical observables.

D. Three-point function and higher at large charge

Again consider the primary operator with charge Q and dimension Δ given in Eq. (3.29), and now define the three-point function

$$G_Q(x_1, x_2, x_3) = \int \mathcal{D}\theta \mathcal{O}_{\Delta_3, Q_3}(x_3) \mathcal{O}_{\Delta_2, Q_2}(x_2) \times \mathcal{O}_{\Delta_1, Q_1}(x_1) e^{-\int d^4x \mathcal{L}_I}, \quad (3.83)$$

where \mathcal{L}_I is the interaction Lagrange density. The operator charges and dimensions are chosen as follows:

$$Q_3 = Q, \quad Q_2 = q, \quad Q_1 = -Q - q, \quad (3.84)$$

$$\Delta_3 = \Delta_Q, \quad \Delta_2 = \Delta_q, \quad \Delta_1 = \Delta_{Q+q}, \quad (3.85)$$

and the large-charge approximation is taken in which $q \ll Q$. In this limit, the small-charge insertion is seen as a probe in the field configuration generated by the two large charge insertions [36–38]. Semiclassically, this requires the insertions to be ordered as $\tau_3 < \tau_2 < \tau_1$. Concretely, one can use the same solution that was found above, corresponding to an insertion of charge Q and evaluate the three-point function at the saddle,⁶

$$G_Q(x_1, x_2, x_3) = \langle \mathcal{O}_{\Delta_Q, Q}(x_3) \mathcal{O}_{\Delta_q, q}(x_2) \mathcal{O}_{\Delta_{Q+q}, -Q-q}(x_1) \rangle \\ = \lim_{\varepsilon \rightarrow 0} \mathcal{O}_{\Delta_Q, Q}(\tau_3 + \varepsilon, \mathbf{x}_3) \mathcal{O}_{\Delta_q, q}(\tau_2, \mathbf{x}_2) \\ \times \mathcal{O}_{\Delta_{Q+q}, -Q-q}(\tau_1 - \varepsilon, \mathbf{x}_1) e^{-S_I[\theta]}|_{\theta=\theta_S} + \dots, \quad (3.86)$$

⁶One could equivalently use the solution corresponding to charge $Q + q$ as a background.




FIG. 3. Euclidean timeline with source charge placements (above line) and charge regions (below line).

where, for simplicity, $\delta \equiv \Delta_q$. The temporal structure of the solution is illustrated in Fig. 3, which makes clear that in the large-charge limit, $q \ll Q$, there is charge Q in the interval between τ_3 and τ_1 , and vanishing charge elsewhere. Note that at $\tau = \tau_2$ the solution is regular and there is no need to move the insertion point. The value of S_I is the same as was found above (with $\tau_{12} \rightarrow \tau_{13}$),

$$S_I = -\frac{3^{1/3}}{4} \xi^{1/2} Q^{4/3} \log\left(\frac{\tau_{13}}{\varepsilon}\right) + \mathcal{O}(\varepsilon), \quad (3.87)$$

and all that remains to do is to insert the values at the saddle, observing that the master field in $x = x_2$ can be written in terms of the Schrödinger-invariant variable v_{123} :

$$e^{iq\theta(x_2)}|_{\theta=\theta_s} = \left(\frac{\tau_{23}}{\tau_{12}}\right)^{q\gamma/2} \exp\left[-\frac{1}{2}qMv_{123} + \frac{1}{2}qM\frac{\mathbf{x}_{23}^2}{\tau_{23}} - \frac{1}{4}qM\frac{\mathbf{x}_{13}^2}{\tau_{13}}\right], \quad (3.88)$$

$$X(x_2)|_{\theta=\theta_s} = \frac{\gamma}{2} \left(\frac{1}{\tau_{12}} + \frac{1}{\tau_{23}}\right) \left(1 - \frac{1}{2\gamma}Mv_{123}\right). \quad (3.89)$$

Then one has

$$G_Q(x_1, x_2, x_3) = \mathcal{N}^3 \lim_{\varepsilon \rightarrow 0} e^{-\frac{3^{1/3}\xi^{1/2}}{4}Q^{4/3} + q\gamma/2 + Q\gamma - (\Delta_{Q+q} + \Delta_Q)/2} \\ \times \tau_{12}^{-(q\gamma+\delta)/2} \tau_{23}^{(q\gamma-\delta)/2} \tau_{13}^{\delta/2 - 1/2(q+2Q) + 3^{1/3}/4\xi^{1/2}Q^{4/3}} \\ \times \exp[Q_1 M \mathbf{x}_{13}^2 / (2\tau_{13}) + Q_2 M \mathbf{x}_{23}^2 / (2\tau_{23})] \\ \times \left(1 - \frac{1}{2\gamma}Mv_{123}\right)^{\delta/2} e^{-qMv_{123}/2}. \quad (3.90)$$

In this case, the cancellation of the divergence reads

$$-\frac{3^{1/3}\xi^{1/2}}{4}Q^{4/3} + q\frac{\gamma}{2} + Q\gamma - \frac{\Delta_{Q+q} + \Delta_Q}{2} = 0. \quad (3.91)$$

In the discussion of the two-point function, it was already found that γ measures the variation of Δ corresponding to a (small) variation of the charge [Eq. (3.78)],

$$\Delta_{Q+q} - \Delta_Q = \gamma q. \quad (3.92)$$

Putting it all together, the three-point function takes the expected form of Eq. (3.3), fixed by Schrödinger invariance, with

$$\Psi_3(v_{123}) = \left(1 - M \frac{v_{123}}{2 \times 3^{1/3} \xi^{1/2} Q^{1/3}}\right)^{\delta/2} e^{-qMv_{123}/2}. \quad (3.93)$$

This result is consistent with what was found in Ref. [17].

The computation of the three-point function can be generalized in a straightforward way to any n -point function with two heavy insertions and $n - 2$ light ones:

$$G_Q(x_1, \dots, x_n) = \langle \mathcal{O}_{\Delta_Q, Q}(x_n) \mathcal{O}_{\delta_{n-1}, q_{n-1}}(x_{n-1}) \dots \mathcal{O}_{\delta_2, q_2}(x_2) \mathcal{O}_{\Delta_{Q'}, -Q'}(x_1) \rangle, \quad (3.94)$$

where, by charge conservation

$$Q' = Q + \sum_{i=2}^{n-1} q_i = Q + q. \quad (3.95)$$

The temporal structure of the solution is illustrated in Fig. 4. The goal is to compute this correlator semiclassically in the EFT, treating the $n - 2$ insertions at x_2, \dots, x_{n-2} as probes for the field profile generated by the heavy insertions at x_1 and x_n (which implies that the small insertions must occur between τ_1 and τ_n). Two conditions have to be met:

- (i) the inner insertions have to be light, $|q_i| \ll Q$, and $|q| \ll Q$,
- (ii) the distance between any two insertions has to be much larger than the UV scale

$$\frac{1}{\tau_{ij}} \ll \Lambda_{\text{UV}} = \rho^{1/3} \approx \frac{Q^{1/3}}{\tau_{1n}}. \quad (3.96)$$

If the insertion of charge Q at x_n is taken as a reference, one recovers precisely the same profile discussed in the previous sections and the n -point function is simply the product of n contributions evaluated at the saddle:

$$G_Q = \lim_{\varepsilon \rightarrow 0} \mathcal{O}_{\Delta_Q, Q}(\tau_n - \varepsilon, \mathbf{x}_n) \mathcal{O}_{\delta_{n-1}, q_{n-1}}(x_{n-1}) \dots \times \mathcal{O}_{\delta_2, q_2}(x_2) \mathcal{O}_{\Delta_{Q'}, -Q'}(\tau_1 - \varepsilon, \mathbf{x}_1) e^{-S_I[\theta]}|_{\theta=\theta_s} + \dots \quad (3.97)$$

The regulator ε is only needed at x_1 and x_n , so that the ε -dependence inside the limit is given by




FIG. 4. Euclidean timeline for n -point function with source charge placements (above line) and charge regions (below line).

$$\varepsilon^{-\frac{3^{1/3} \xi^{1/2}}{4}} Q^{4/3 + \gamma(Q' + Q)/2 - (\Delta_{Q'} + \Delta_Q)/2}, \quad (3.98)$$

where the fact that the value of the action at the saddle is still given by Eq. (3.73) has been used. Then the divergence is canceled if

$$-\frac{3^{1/3} \xi^{1/2}}{4} Q^{4/3} + \frac{\gamma}{2} (Q' + Q) - \frac{1}{2} (\Delta_{Q'} + \Delta_Q) = 0, \quad (3.99)$$

which is identically true since, as seen in the computation of the two-point function,

$$\Delta_Q = Q\gamma - \frac{3^{1/3} \xi^{1/2}}{4} Q^{4/3} \quad \text{and} \quad \Delta_{Q'} - \Delta_Q = \gamma(Q' - Q). \quad (3.100)$$

Consistently with the appropriate Ward identities, the result can be parametrized in terms of $(n - 1)(n - 2)/2$ independent conformally invariant variables

$$v_{ijn} \equiv \frac{1}{2} \left(\frac{\mathbf{x}_{jn}^2}{\tau_{jn}} + \frac{\mathbf{x}_{ij}^2}{\tau_{ij}} - \frac{\mathbf{x}_{in}^2}{\tau_{in}} \right) \quad i < j < n. \quad (3.101)$$

Now one can write the general expression of the master field in terms of v_{1in} :

$$e^{iq_i \theta(x_i)}|_{\theta=\theta_s} = \left(\frac{\tau_{in}}{\tau_{1i}} \right)^{q_i \gamma/2} \exp \left[-\frac{1}{2} q_i M v_{1in} + \frac{1}{2} q_i M \frac{\mathbf{x}_{in}^2}{\tau_{in}} - \frac{1}{4} q_i M \frac{\mathbf{x}_{1n}^2}{\tau_{1n}} \right], \quad (3.102)$$

$$X(x_i)|_{\theta=\theta_s} = \frac{\gamma}{2} \left(\frac{1}{\tau_{1i}} + \frac{1}{\tau_{in}} \right) \left(1 - \frac{1}{2\gamma} M v_{1in} \right), \quad (3.103)$$

to find that the n -point function takes a factorized form

$$G_Q = \mathcal{N}^2 \tau_{1n}^{-\Delta_Q - q\gamma/2 + \delta/2} \exp \left[-(Q + q) M \frac{\mathbf{x}_{1n}^2}{2\tau_{1n}} \right] \times \prod_{i=2}^{n-1} \mathcal{N}_i \tau_{1i}^{-q_i \gamma/2 - \delta_i/2} \tau_{in}^{q_i \gamma/2 - \delta_i/2} \exp \left[q_i M \frac{\mathbf{x}_{in}^2}{2\tau_{in}} \right] \times \left(1 - M \frac{v_{1in}}{2 \times 3^{1/3} \xi^{1/2} Q^{1/3}} \right)^{\delta_i/2} e^{-q_i M v_{1in}/2}, \quad (3.104)$$

where $q = q_2 + \dots + q_{n-2}$ and $\delta = \delta_2 + \dots + \delta_{n-2}$.

IV. SYMMETRY BREAKING EFFECTS

A. Fermions near unitarity

Consider now the inclusion of small Schrödinger-breaking effects in the fundamental theory. Using Eq. (2.5) one can write

$$\mathcal{L} = \mathcal{L}_{\text{CFT}} + \frac{M}{4\pi a} s^\dagger s - \frac{M^2 r}{16\pi} s^\dagger \left(i \overleftrightarrow{\partial}_t + \frac{\overleftrightarrow{\nabla}^2 + \overleftarrow{\nabla}^2}{4M} \right) s, \quad (4.1)$$

where, in addition, effective range corrections have been included [9,39]. The scattering length corrections therefore enter via a relevant dimension-4 operator (the field s at unitarity has dimension 2), and the effective range corrections enter via an irrelevant dimension-6 operator.

The energy per particle E/N of the interacting Fermi gas in the near-Schrödinger limit can be written at very-low densities⁷ as [40,41]

$$E/N = \frac{3}{5} \frac{k_F^2}{2M} \left(\xi - \frac{\zeta}{k_F a} - \frac{\zeta_2}{k_F^2 a^2} + \cdots + \eta k_F r + \cdots \right). \quad (4.2)$$

Here the various dimensionless universal parameters have been determined using quantum Monte Carlo simulations. From Ref. [42], $\xi = 0.372(5)$ (Bertsch parameter) and $\eta = 0.12(3)$. From the simulation data in Ref. [43], it is straightforward to extract $\zeta = 0.8(3)$ and $\zeta_2 = 1.0(6)$.

Note that the effective range contributions to the energy density grow with Fermi momentum with respect to the energy at unitarity. This implies that in the large-charge limit, the superfluid EFT is sensitive to arbitrarily short distance scales in the fundamental fermion theory. Hence, the effect of the effective range and all higher shape parameter corrections that arise from operators with arbitrary numbers of derivative interactions are of the same size, leading to a breakdown of the fundamental-fermion EFT expansion. This renders it difficult to assess the effect of Schrödinger-breaking effects due to finite effective-range corrections in the large-charge EFT, as evidently knowledge of the equation-of-state at high densities is required.⁸ This will be treated in a separate publication, and therefore effective range and higher shape parameter corrections in the fundamental theory are assumed to vanish in what follows.

B. Symmetry breaking in EFT

The leading conformal-breaking effects are straightforward to include using spurion formalism. In the fundamental theory, recalling that the field s has scaling dimension 2, the conformal-breaking effects can be made scale invariant by assigning $1/a$ scaling dimension 1. Correspondingly, in the EFT, the leading conformal-breaking effects due to a finite scattering length are encoded as⁹

⁷Note that the range of validity of the EFT of neutron contact operators is, strictly speaking, not set by the pion mass, M_π , but rather by $M_\pi/2$, which is the position of the t-channel branch point in the neutron-neutron scattering amplitude.

⁸Recent state-of-the-art simulations of the energy density in neutron matter [44] indicate a remarkable closeness to the unitary curve up to unexpectedly high densities.

⁹Note that this is the Euclidean Lagrange density, and therefore the relationship with the energy density is given by Eq. (3.25).

$$\mathcal{L}_{\text{SB}} = g_1 a^{-1} M X^2 + g_2 a^{-2} M^{1/2} X^{3/2}, \quad (4.3)$$

where the g_i are dimensionless constants.¹⁰ Matching to Eq. (4.2) one finds

$$g_1 = -\frac{2\zeta}{5\pi^2 \xi^2} = -0.23(10),$$

$$g_2 = -\frac{\sqrt{2}}{25\pi^2 \xi^{5/2}} (4\zeta^2 + 5\zeta_2 \xi) = -0.34(17), \quad (4.4)$$

where the Monte Carlo results quoted above have been used in the numerical determination.

Note that following this procedure for the effective range corrections, one would assign r scaling dimension -1 . Correspondingly, the range corrections would enter perturbative matching via an X^3 operator which scales as μ^3 . As the LO Schrödinger-invariant operator scales as $\mu^{5/2}$, formally there can be no such operator in the large-charge limit. This suggests that the utility of the large-charge EFT in the presence of significant effective range, and higher shape parameter effects, will be system specific.

C. Symmetry breaking at large charge

Remarkably, Schrödinger symmetry places strong constraints on the general form of the symmetry-breaking corrections due to scattering length and effective range correction [9]. In the fundamental fermion theory with a two-point function constructed from an operator of arbitrary charge Q , it can be shown in conformal perturbation theory that

$$G_Q(x_1, x_2) = G_Q^{\text{CFT}}(x_1, x_2) [1 + c' a^{-1} \tau_{12}^{1/2}], \quad (4.5)$$

where c' is an arbitrary constant that is not fixed by symmetry. In conformal perturbation theory this symmetry-breaking effect is computed by obtaining the symmetry-breaking action at the Schrödinger-symmetric master field solution.

At leading order in the symmetry breaking, the saddle solution remains unchanged. This is reflected in the fact that in computing n -point functions in the theory with symmetry breaking, in the limit in which two insertions are much heavier than the others and the symmetry-breaking is small, the only new effect is a variation in the value of the action at the saddle:

¹⁰Note that the contribution of the leading scattering length correction to the phonon dispersion relation has been considered in Ref. [45].

$$\begin{aligned}
& \left\langle \mathcal{O}_{Q,\Delta} \prod_{i=2}^{n-2} \mathcal{O}_{\delta_i, q_i} \mathcal{O}_{\Delta', -Q'} \right\rangle_{\text{SB}} \\
& \sim \mathcal{O}_{Q,\Delta} \prod_{i=2}^{n-2} \mathcal{O}_{\delta_i, q_i} \mathcal{O}_{\Delta', -Q'} e^{-S[\theta] - S_{\text{SB}}[\theta]} \Big|_{\theta=\theta_S} \\
& = \left\langle \mathcal{O}_{Q,\Delta} \prod_{i=2}^{n-2} \mathcal{O}_{\delta_i, q_i} \mathcal{O}_{\Delta', -Q'} \right\rangle_{\text{CFT}} e^{-S_{\text{SB}}[\theta_S]}, \quad (4.6)
\end{aligned}$$

where $\Delta = \Delta(Q)$, $\Delta' = \Delta(Q')$, $Q' = Q + q_2 + \dots + q_{n-2}$, $Q \gg q_i$, and $\langle \dots \rangle_{\text{CFT}}$ is the correlator in the conformal theory. In this limit the breaking of Schrödinger symmetry is measured by a universal term that is easily evaluated.

For the leading scattering length correction one finds

$$\int d^3\mathbf{x} \mathcal{H}_{\text{SB}}^a = C^a M^{-1/2} Q^{7/6} a^{-1} \tilde{\omega}(\tau)^{1/2}, \quad (4.7)$$

where

$$C^a \equiv \frac{64\sqrt{2}3^{1/6}\pi\xi^{7/4}g_1}{35}, \quad (4.8)$$

and

$$\begin{aligned}
S_{\text{SB}}^a &= \int d^4x \mathcal{H}_{\text{SB}}^a \\
&= C^a M^{-1/2} Q^{7/6} a^{-1} \frac{1}{\sqrt{2}} \int_{\tau_2+\epsilon}^{\tau_1-\epsilon} d\tau \left(\frac{(\tau_1 - \tau_2)}{(\tau - \tau_2)(\tau_1 - \tau)} \right)^{1/2} \\
&= \frac{1}{\sqrt{2}} \pi C^a M^{-1/2} Q^{7/6} a^{-1} \tau_{12}^{1/2}. \quad (4.9)
\end{aligned}$$

This effect is small as compared to LO for $a^{-1} \ll M^{1/2} Q^{1/6} \tau_{12}^{-1/2} \sim \Lambda_{\text{UV}}$. Therefore, with a large, one can take $a^{-1} \sim p_{\text{IR}}$ and scattering length effects are small at large charge. For the two-point function one finds

$$G_Q(x_1, x_2) = G_Q^{\text{CFT}}(x_1, x_2) \left[1 - \frac{1}{\sqrt{2}} \pi C^a M^{-1/2} Q^{7/6} a^{-1} \tau_{12}^{1/2} \right],$$

which is indeed of the expected form, Eq. (4.5). Now, continuing back to Minkowski space, choosing the source points $x_1 = (t, \mathbf{x})$, $x_2 = (0, \mathbf{0})$, and using the Fourier transform [6,8]

$$\begin{aligned}
& \int dt \int d^3\mathbf{x} \theta(t) t^{-\Delta} \exp\left(i \frac{QM\mathbf{x}^2}{2t}\right) \exp(iEt - i\mathbf{p} \cdot \mathbf{x}) \\
& = i^{\Delta-1} \left(\frac{2\pi}{QM} \right)^{3/2} \left(\frac{p^2}{2QM} - E \right)^{\Delta-5/2} \Gamma\left(\frac{5}{2} - \Delta\right), \quad (4.10)
\end{aligned}$$

one finds

$$\begin{aligned}
G_Q(E, \mathbf{p}) &= -i\mathcal{N}^2 \left(\frac{2\pi}{QM} \right)^{3/2} \Gamma\left(\frac{5}{2} - \Delta_Q\right) \left(\frac{p^2}{2QM} - E \right)^{\Delta_Q-5/2} \\
&\times \left[1 - \frac{1}{\sqrt{2}} \pi C^a M^{-1/2} Q^{7/6} a^{-1} \frac{\Gamma(\frac{6}{2} - \Delta_Q)}{\Gamma(\frac{5}{2} - \Delta_Q)} \right. \\
&\times \left. \left(\frac{p^2}{2QM} - E \right)^{-1/2} \right]. \quad (4.11)
\end{aligned}$$

Finally, one has

$$\text{Im } G_Q(E, \mathbf{0}) = C_0 E^{\Delta_Q-5/2} \left[1 + \frac{C_Q}{a\sqrt{ME}} \right], \quad (4.12)$$

where C_0 is a normalization constant that can be absorbed into the definition of the X field, and

$$\begin{aligned}
C_Q(Q) &= -\frac{1}{\sqrt{2}} \pi C^a Q^{7/6} \frac{\Gamma(\frac{6}{2} - \Delta_Q)}{\Gamma(\frac{5}{2} - \Delta_Q)} \tan \pi \Delta_Q \\
&\xrightarrow{Q \rightarrow \infty} 3^{5/6} \frac{64}{175} \zeta Q^{11/6} + \mathcal{O}(Q^{1/2}), \quad (4.13)
\end{aligned}$$

where on the right side the result at asymptotic Q is given. This function is plotted in Fig. 5. The $\mathcal{O}(a^{-2})$ corrections are similarly evaluated. It is noteworthy and promising that, for $Q \sim 3$, these $\mathcal{O}(a^{-1})$ corrections in the large-charge EFT are consistent with the range of values found in Ref. [9] working directly with the three-body wave functions. Of course the utility of the large-charge EFT for neutron matter requires consideration of effective-range effects, which are sizeable in the neutron-neutron system.




FIG. 5. Plot of the function $C_Q(Q)$ vs Q . The band represents the uncertainty in the parameter g_1 , determined by Monte Carlo simulations, as discussed in the text.

V. LARGE N AT LARGE CHARGE

The action in Eq. (4.1) can be studied perturbatively without the need for an EFT if one introduces N species of fermions and takes the $N \rightarrow \infty$ limit. To make contact with the superfluid EFT, a sector of fixed charge Q is also considered, and the double-scaling limit Q/N with fixed is studied. Here the effect of a finite scattering length a on the gap and on the value of the grand potential is considered. The calculation of other observables is left for future work.

The starting point is the action

$$S = \int dt d^3\mathbf{x} \left[\sum_{i=1}^N \psi_{\sigma i}^\dagger \left(i\partial_t + \frac{1}{2M} \nabla^2 + \mu \right) \psi_{\sigma i} + \psi_{\downarrow i}^\dagger \psi_{\uparrow i}^\dagger s + s^\dagger \psi_{\uparrow i} \psi_{\downarrow i} + \frac{NM}{4\pi a} s^\dagger s - \frac{M^2 r}{16\pi} s^\dagger \left(i\overleftrightarrow{\partial}_t + \frac{\overleftrightarrow{\nabla}^2}{4M} \right) s \right], \quad (5.1)$$

which is the nonrelativistic version of the Gross-Neveu-Yukawa (GNY) model. Note that the field s remains a singlet of the $U(1) \times Sp(2N)$ symmetry of the action.

One way to view this model is as the UV completion of a fermionic theory with four-Fermi interaction. At high energy, both the fermionic and the bosonic fields are dynamical. Going down in energy, the first scale that one meets is $1/(Mra)$ where the scalar field is frozen, and one is not sensitive to a finite effective range of the interactions. Below this energy, s is nondynamical, appears quadratically and can be integrated out, leading to a four-Fermi interaction with coupling proportional to a/N . In this way, the strongly coupled fixed point of the fermionic model then appears as an IR fixed point for the nonrelativistic GNY model. From this point of view, the theory under discussion lives below the first scale (the bosonic field is not dynamical) and is close enough to the IR fixed point.

The fermions appear quadratically and can be integrated out,

$$S = -N \text{Tr} \log G^{-1}[s] - \int d\tau d\mathbf{x} \frac{NM}{4\pi a} s^\dagger s, \quad (5.2)$$

where the inverse propagator is given by

$$G^{-1}[s] = \begin{pmatrix} -\partial_\tau + \frac{1}{2M} \nabla^2 + \mu & s(x) \\ s(x)^\dagger & -\partial_\tau - \frac{1}{2M} \nabla^2 - \mu \end{pmatrix}. \quad (5.3)$$

At leading order in N one can look for a saddle at constant (real) values of s , the effect of the fluctuations being controlled by $1/N$. A standard calculation leads to the following expression for the action evaluated at $s = \Sigma_0$ [26]:

$$\frac{S(\mu, \Sigma_0)}{NVT} = -4\pi M^{3/2} I_{0,0} \left(\frac{\Sigma_0}{\mu} \right) \mu^{5/2} - \frac{\Sigma_0^2}{4\pi a}, \quad (5.4)$$

where $I_{m,n}$ is written in terms of Gaussian hypergeometric functions ${}_2F_1$:

$$I_{m,n}(y) = -\frac{y^{5/2+m/2-2n}}{2(2\pi)^{7/2}} \left[\Gamma\left(\frac{3+m}{4}\right) \Gamma\left(n - \frac{5+m}{4}\right) \times {}_2F_1\left(-\frac{m+1}{4}, n - \frac{m+5}{4}, \frac{1}{2}; -\frac{1}{y^2}\right) + \frac{2}{y} \Gamma\left(\frac{5+m}{4}\right) \Gamma\left(n - \frac{3+m}{4}\right) \times {}_2F_1\left(\frac{1-m}{4}, n - \frac{m+3}{4}, \frac{3}{2}; -\frac{1}{y^2}\right) \right]. \quad (5.5)$$

Here, a heat kernel regularization has been used in which the conformal point is at $1/a^* = 0$.

The value of the action at the saddle, that is identified with the grand potential $\Omega(\mu)$, is obtained by solving the gap equation

$$\frac{d}{d\Sigma} S(\mu, \Sigma) = 0. \quad (5.6)$$

This equation can be solved perturbatively in $1/a$ using the identity

$$I'_{m,n}(y) = -2y I_{m,n+1}(y). \quad (5.7)$$

At leading order,

$$I_{0,1}(y_0) = 0, \quad (5.8)$$

which can be solved numerically to give $y_0 = \Sigma_0/\mu = 1.162$, and a Bertsch parameter of $\xi = 0.5906$, consistent with well-known results in the literature [46]. Expanding around this point in inverse powers of a , one then finds the gap

$$\Sigma_0 = \mu y_0 \left(1 - \frac{1}{32\pi^2 y_0^2 a \mu^{1/2} I_{0,2}(y_0)} + \frac{1}{2(32\pi^2 y_0^2 a \mu^{1/2})^2 I_{0,2}(y_0)^2} \dots \right), \quad (5.9)$$

and the grand potential

$$\begin{aligned} \Omega(\mu) &= 4\pi N \mu^{5/2} I_{0,0}(y_0) \left(1 - \frac{y_0^2}{16\pi^2 a \mu^{1/2} I_{0,0}(y_0)} \right. \\ &\quad \left. - \frac{1}{2(16\pi^2 a \mu^{1/2})^2 I_{0,0}(y_0) I_{0,2}(y_0)} \right. \\ &\quad \left. - \frac{1}{6(16\pi^2 a \mu^{1/2})^3 y_0^2 I_{0,0}(y_0) I_{0,2}(y_0)^2} + \dots \right) \\ &= -0.08418 \mu^{5/2} \left(1 + \frac{1.277}{a \mu^{1/2}} + \frac{1.513}{a^2 \mu} - \frac{1.195}{a^3 \mu^{3/2}} + \dots \right). \end{aligned} \quad (5.10)$$

In the limit of large chemical potential, the expected corrections to the standard result for the grand potential are found at the fixed point, which are controlled by $1/(a\mu^{1/2})$. This is in perfect analogy with the result for the n -point functions in the previous section which, in the grand canonical ensemble, are controlled by $\tau_{12}^{1/2}/(a\gamma^{1/2})$, which is consistent with the identification of $2\gamma/\tau_{12}$ with the chemical potential. The grand potential is easily read off from the Euclidean Lagrange density, including scattering length corrections, giving

$$\Omega(\mu) = -c_0\mu^{5/2} \left(1 + \frac{3\zeta}{2^{3/2}\xi^{1/2}a\mu^{1/2}} + \frac{3(4\zeta^2 + 5\zeta_2\xi)}{20\xi a^2\mu} + \dots \right). \quad (5.11)$$

Matching, one finds that in the large- N limit, $\zeta = 0.925$ and $\zeta_2 = 0.858$. These values are consistent with the Monte Carlo simulation values, if one makes a conservative estimate of uncertainties due to omitted higher orders in the large- N expansion.

VI. CONCLUSION

Schrödinger symmetry places strong constraints on the form of correlation functions in nonrelativistic conformal field theory, and the state-operator correspondence provides a powerful tool for computing the conformal dimensions of the operators that appear in the correlation functions. This paper has shown that in the large-charge limit, there exists a Goldstone boson master field that allows for the exact computation of large-charge correlation functions directly from their path integral expressions. A direct derivation of the master field was given which relies on the assumption of Schrödinger invariance of the Goldstone-boson equation of motion. However, it has also been shown that the master-field solution is present in the Schrödinger algebra, and can be obtained directly from the ground-state large-charge solution in the absence of sources via a transformation from the Galilean frame to the oscillator frame. The master field has been used to reproduce the known 2- and 3-point correlation functions, as well as the n -point correlation function with an insertion of large charge.

Additionally, the grand potential has been computed including the effect of finite scattering length in the limit of large number of fermion species without using the EFT. The result is consistent with the EFT results. In this limit, moreover, the numerical constants that are not accessible in the EFT can be computed and are consistent with Monte Carlo results.

While Schrödinger symmetry constrains the form of Schrödinger symmetry breaking corrections, the overall coefficients of the corrections are not fixed by symmetry. However, in conformal perturbation theory, these coefficients are determined by the master field solution. In particular, symmetry-breaking corrections due to finite scattering length were computed, and shown to appear in a universal manner for any large-charge n -point function. While scattering length corrections are naturally accounted for in the large-charge EFT, the effective-range corrections have to be unnaturally small to remain a perturbative effect. This suggests that the large-charge EFT may not provide an efficient description of systems of many neutrons. Here the issue is subtle as in the large-charge limit, the superfluid EFT probes all distance scales in the fundamental-fermion EFT. The issue of the effective range corrections and their quantitative contribution to symmetry breaking in neutron matter will be treated elsewhere.

It would be interesting to consider quantum corrections, which have been studied using the state-operator correspondence, in the context of the master field solution and the time-dependent superfluid droplet. In addition, given the simplicity of the master-field equation, one interesting avenue to pursue is the possibility of a holographic dual [47,48] based on a Schrödinger background [49].

ACKNOWLEDGMENTS

We would like to thank Dam Thanh Son for illuminating discussions. D.O. and S.R. thank Jan Herrmann for discussions and collaboration related to the large- N computation. This work was supported by the Swiss National Science Foundation (SNSF) under Grant No. 200021_192137 and by the U.S. Department of Energy Grant No. DE-FG02-97ER-41014 (UW Nuclear Theory).

-
- [1] U. van Kolck, Effective field theory of short range forces, *Nucl. Phys.* **A645**, 273 (1999).
 - [2] D. B. Kaplan, M. J. Savage, and M. B. Wise, A new expansion for nucleon-nucleon interactions, *Phys. Lett. B* **424**, 390 (1998).

- [3] D. B. Kaplan, M. J. Savage, and M. B. Wise, Two nucleon systems from effective field theory, *Nucl. Phys.* **B534**, 329 (1998).
- [4] T. Mehen, I. W. Stewart, and M. B. Wise, Wigner symmetry in the limit of large scattering lengths, *Phys. Rev. Lett.* **83**, 931 (1999).

- [5] T. Mehen, I.W. Stewart, and M.B. Wise, Conformal invariance for nonrelativistic field theory, *Phys. Lett. B* **474**, 145 (2000).
- [6] H.-W. Hammer and D.T. Son, Unnuclear physics, *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2108716118 (2021).
- [7] T. Schaefer and G. Baym, From nuclear to unnuclear physics, *Proc. Natl. Acad. Sci. U.S.A.* **118**, e2113775118 (2021).
- [8] E. Braaten and H.-W. Hammer, Point production of a nonrelativistic unparticle recoiling against a particle, *Phys. Rev. D* **107**, 034017 (2023).
- [9] S.D. Chowdhury, R. Mishra, and D.T. Son, Applied nonrelativistic conformal field theory: Scattering-length and effective-range corrections to unnuclear physics, *Phys. Rev. D* **109**, 016001 (2024).
- [10] H. Georgi, Unparticle physics, *Phys. Rev. Lett.* **98**, 221601 (2007).
- [11] C.R. Hagen, Scale and conformal transformations in galilean-covariant field theory, *Phys. Rev. D* **5**, 377 (1972).
- [12] U. Niederer, The maximal kinematical invariance group of the free Schrodinger equation, *Helv. Phys. Acta* **45**, 802 (1972).
- [13] M. Henkel, Schrodinger invariance in strongly anisotropic critical systems, *J. Stat. Phys.* **75**, 1023 (1994).
- [14] M. Henkel and J. Unterberger, Schrodinger invariance and space-time symmetries, *Nucl. Phys. B* **660**, 407 (2003).
- [15] Y. Nishida and D.T. Son, Nonrelativistic conformal field theories, *Phys. Rev. D* **76**, 086004 (2007).
- [16] S. Favrod, D. Orlando, and S. Reffert, The large-charge expansion for Schrödinger systems, *J. High Energy Phys.* **12** (2018) 052.
- [17] S.M. Kravec and S. Pal, Nonrelativistic conformal field theories in the large charge sector, *J. High Energy Phys.* **02** (2019) 008.
- [18] S. Hellerman, D. Orlando, S. Reffert, and M. Watanabe, On the CFT operator spectrum at large global charge, *J. High Energy Phys.* **12** (2015) 071.
- [19] L. A. Gaumé, D. Orlando, and S. Reffert, Selected topics in the large quantum number expansion, *Phys. Rep.* **933**, 1 (2021).
- [20] M. Greiter, F. Wilczek, and E. Witten, Hydrodynamic relations in superconductivity, *Mod. Phys. Lett. B* **03**, 903 (1989).
- [21] D.T. Son and M. Wingate, General coordinate invariance and conformal invariance in nonrelativistic physics: Unitary Fermi gas, *Ann. Phys. (Amsterdam)* **321**, 197 (2006).
- [22] S.M. Kravec and S. Pal, The spinful large charge sector of nonrelativistic CFTs: From phonons to vortex crystals, *J. High Energy Phys.* **05** (2019) 194.
- [23] S. Hellerman and I. Swanson, Droplet-edge operators in nonrelativistic conformal field theories, *arXiv:2010.07967*.
- [24] V. Pellizzani, Operator spectrum of nonrelativistic CFTs at large charge, *Phys. Rev. D* **105**, 125018 (2022).
- [25] S. Hellerman, D. Orlando, V. Pellizzani, S. Reffert, and I. Swanson, Nonrelativistic CFTs at large charge: Casimir energy and logarithmic enhancements, *J. High Energy Phys.* **05** (2022) 135.
- [26] S. Hellerman, D. Krichevskiy, D. Orlando, V. Pellizzani, S. Reffert, and I. Swanson, The unitary Fermi gas at large charge and large N , *J. High Energy Phys.* **05** (2024) 323.
- [27] D.T. Son, Low-energy quantum effective action for relativistic superfluids, *arXiv:hep-ph/0204199*.
- [28] A. Volovich and C. Wen, Correlation functions in non-relativistic holography, *J. High Energy Phys.* **05** (2009) 087.
- [29] S. Hellerman and S. Maeda, On the large R -charge expansion in $\mathcal{N} = 2$ superconformal field theories, *J. High Energy Phys.* **12** (2017) 135.
- [30] G. Arias-Tamargo, D. Rodriguez-Gomez, and J.G. Russo, The large charge limit of scalar field theories and the Wilson-Fisher fixed point at $\epsilon = 0$, *J. High Energy Phys.* **10** (2019) 201.
- [31] D. Orlando, S. Reffert, and F. Sannino, Near-conformal dynamics at large charge, *Phys. Rev. D* **101**, 065018 (2020).
- [32] G. Badel, G. Cuomo, A. Monin, and R. Rattazzi, Feynman diagrams and the large charge expansion in $3 - \epsilon$ dimensions, *Phys. Lett. B* **802**, 135202 (2020).
- [33] D.T. Son, M. Stephanov, and H.-U. Yee, Fate of multi-particle resonances: From Q-balls to ^3He droplets, *Phys. Rev. A* **106**, L050801 (2022).
- [34] W.D. Goldberger, Z.U. Khandker, and S. Prabhu, OPE convergence in non-relativistic conformal field theories, *J. High Energy Phys.* **12** (2015) 048.
- [35] J.L. Mañes and M.A. Valle, Effective theory for the Goldstone field in the BCS-BEC crossover at $T = 0$, *Ann. Phys. (Amsterdam)* **324**, 1136 (2009).
- [36] A. Monin, D. Pirtskhalava, R. Rattazzi, and F.K. Seibold, Semiclassics, Goldstone bosons and CFT data, *J. High Energy Phys.* **06** (2017) 011.
- [37] G. Cuomo, A note on the large charge expansion in 4d CFT, *Phys. Lett. B* **812**, 136014 (2021).
- [38] N. Dondi, I. Kalogerakis, R. Moser, D. Orlando, and S. Reffert, Spinning correlators in large-charge CFTs, *Nucl. Phys. B* **983**, 115928 (2022).
- [39] S.R. Beane and M.J. Savage, Rearranging pionless effective field theory, *Nucl. Phys. A* **694**, 511 (2001).
- [40] A. Bulgac and G.F. Bertsch, Collective oscillations of a trapped fermi gas near the unitary limit, *Phys. Rev. Lett.* **94**, 070401 (2005).
- [41] U. van Kolck, Unitarity and discrete scale invariance, *Few Body Sys.* **58**, 112 (2017).
- [42] J. Carlson, S. Gandolfi, K.E. Schmidt, and S. Zhang, Auxiliary-field quantum monte carlo method for strongly paired fermions, *Phys. Rev. A* **84**, 061602 (2011).
- [43] S.Y. Chang, V.R. Pandharipande, J. Carlson, and K.E. Schmidt, Quantum Monte Carlo studies of superfluid Fermi gases, *Phys. Rev. A* **70**, 043602 (2004).
- [44] Y. Lim and A. Schwenk, Symmetry energy and neutron star properties constrained by chiral effective field theory calculations, *Phys. Rev. C* **109**, 035801 (2024).
- [45] M.A. Escobedo, M. Mannarelli, and C. Manuel, Bulk viscosities for cold Fermi superfluids close to the unitary limit, *Phys. Rev. A* **79**, 063623 (2009).

- [46] M. Y. Veillette, D. E. Sheehy, and L. Radzihovsky, Large- N expansion for unitary superfluid Fermi gases, *Phys. Rev. A* **75**, 043614 (2007).
- [47] D. T. Son, Toward an AdS/cold atoms correspondence: A geometric realization of the Schrodinger symmetry, *Phys. Rev. D* **78**, 046003 (2008).
- [48] K. Balasubramanian and J. McGreevy, Gravity duals for nonrelativistic conformal field theories, *Phys. Rev. Lett.* **101**, 061601 (2008).
- [49] D. Israel, C. Kounnas, D. Orlando, and P. M. Petropoulos, Electric/magnetic deformations of S^3 and AdS_3 , and geometric cosets, *Fortschr. Phys.* **53**, 73 (2005).