Search for charmed baryons in the $\Lambda_c^+ \eta$ system and measurement of the branching fractions of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ decaying to $\Lambda_c^+ \eta$ and pD^0 relative to $\Sigma_c(2455)\pi$

S. X. Li[®], C. P. Shen[®], I. Adachi[®], J. K. Ahn[®], H. Aihara[®], D. M. Asner[®], H. Atmacan[®], T. Aushev[®], R. Ayad[®],
Sw. Banerjee[®], K. Belous[®], J. Bennett[®], M. Bessner[®], T. Bilka[®], D. Biswas[®], D. Bodrov[®], A. Bozek[®], M. Bračko[®],
P. Branchini[®], T. E. Browder[®], A. Budano[®], M. Campajola[®], M.-C. Chang[®], B. G. Cheon[®], K. Chilikin[®], H. E. Cho[®],
K. Cho[®], S.-K. Choi[®], Y. Choi[®], S. Choudhury[®], N. Dash[®], G. De Nardo[®], G. De Pietro[®], R. Dhamija[®],
J. Dingfelder[®], Z. Doležal[®], T. V. Dong[®], S. Dubey[®], P. Ecker[®], T. Ferber[®], B. G. Fulsom[®], V. Gaur[®], A. Garmash[®],
P. Goldenzweig[®], E. Graziani[®], B. Grube[®], Y. Guan[®], K. Gudkova[®], C. Hadjivasiliou[®], C.-L. Hsu[®], N. Ipsita[®],
R. Itoh[®], M. Iwasaki[®], W. W. Jacobs[®], Q. P. Ji[®], S. Jia[®], Y. Jin[®], K. K. Joo[®], C. Kiesling[®], D. Y. Kim[®], Y. J. Kim[®],
K. Kinoshita[®], P. Kodyš[®], A. Korobov[®], S. Korpar[®], E. Kovalenko[®], P. Križan[®], P. Krokovny[®], T. Kuhr[®], R. Kumar[®],
K. Kumara[®], Y.-J. Kwon[®], L. K. Li[®], Y. Li[®], Y. B. Li[®], D. Liventsev[®], M. Masuda[®], S. K. Maurya[®], F. Meier[®],
M. Merola[®], K. Miyabayashi[®], R. Mizuk[®], R. Mussa[®], T. Nakano[®], M. Nakao[®], A. Natochii[®], M. Nayak[®],
S. Nishida[®], P. Pakhlov[®], G. Pakhlova[®], S. Pardi[®], J. Park[®], S.-H. Park[®], S. Patra[®], S. Paul[®], T. K. Pedlar[®],
R. Pestotnik[®], L. E. Piilonen[®], T. Podobnik[®], E. Prencipe[®], M. T. Prim[®], G. Russo[®], S. Sandilya[®], V. Savinov[®],
G. Schnell⁹, C. Schwanda[®], Y. Seino[®], K. Senyo⁹, J.-G. Shiu[®], E. Solovieva[®], M. Starič[®], M. Sumihama[®],
M. Takizawa[®], U. Tamponi[®], K. Tanida[®], F. Tenchini[®], M. Uchida[®], T. Uglov[®], S. Uno[®], E. Wan[®], E. Won[®],

(The Belle Collaboration)

(Received 25 June 2024; accepted 26 July 2024; published 27 August 2024)

We search for excited charmed baryons in the $\Lambda_c^+\eta$ system using a data sample corresponding to an integrated luminosity of 980 fb⁻¹. The data were collected by the Belle detector at the KEKB e^+e^- asymmetric-energy collider. No significant signals are found in the $\Lambda_c^+\eta$ mass spectrum, including the known $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$. Clear $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signals are observed in the pD^0 mass spectrum. We set upper limits at 90% credibility level on ratios of branching fractions of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ decaying to $\Lambda_c^+\eta$ relative to $\Sigma_c(2455)\pi$ of < 0.13 for the $\Lambda_c(2880)^+$ and < 1.11 for the $\Lambda_c(2940)^+$. We measure ratios of branching fractions of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ decaying to pD^0 relative to $\Sigma_c(2455)\pi$ of 0.75 ± 0.03(stat) ± 0.07(syst) for the $\Lambda_c(2880)^+$ and 3.59 ± 0.21(stat) ± 0.56(syst) for the $\Lambda_c(2940)^+$.

DOI: 10.1103/PhysRevD.110.032021

I. INTRODUCTION

Charmed baryons provide a good opportunity to study the dynamics of quark confinement [1–5]. Usually, singlycharmed baryons are considered to be bound states of a charm quark and a light diquark. In this model, the Jacobi coordinate ρ is used to label the degree of freedom between the light quarks and λ to label the degree of freedom between the charm quark and the center of mass of the diquark. The excitation of the diquark is called ρ -mode excitation, while the excitation between the charm quark and the diquark is called λ -mode excitation. The excitation energy for the ρ -mode is expected to be higher than the one for λ -mode by a factor of $\sqrt{3}$ in the heavy-quark limit [6].

In the Λ_c^+ sector, both the $\Lambda_c(2880)^+$ and the $\Lambda_c(2940)^+$ have been observed in several decay modes. The $\Lambda_c(2880)^+$ was first observed by CLEO in the $\Lambda_c^+\pi^+\pi^$ decay mode, where some of the events resonated through $\Sigma_c(2455)\pi$ [7], and later reported by *BABAR* in the *pD*⁰ mass spectrum [8]. The $\Lambda_c(2940)^+$ was first seen by *BABAR* in the *pD*⁰ decay mode [8] and confirmed by Belle in the $\Sigma_c(2455)\pi$ decay [9]. The spin-parities of the two states were also studied. Belle analyzed the $\Sigma_c(2455/2520)\pi$ decay modes and measured the spin of the $\Lambda_c(2880)^+$ to be 5/2 [9]. LHCb studied the $\Lambda_b^0 \rightarrow$ $D^0p\pi^-$ decay and measured the favored J^P of $\Lambda_c(2940)^+$

Published by the American Physical Society under the terms of the Creative Commons Attribution 4.0 International license. Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

to be $3/2^-$, but they could not exclude spins from 1/2 to 7/2 [10]. The inner structure of the $\Lambda_c(2940)^+$ remains a puzzle. The $\Lambda_c(2940)^+$ may be a conventional excited charmed baryon, but has also been explained as an ND^* molecular state with potential spin-parity assignments including $1/2^+$, $1/2^-$, and $3/2^-$ [11–13].

 $\Lambda_c^+\eta$ is a good channel to search for excited Λ_c^+ baryons. Any signal in $\Lambda_c^+\eta$ is likely to be an excited Λ_c^+ rather than a Σ_c^+ , as for the latter decays to $\Lambda_c^+\pi$ are allowed by isospin and are likely to dominate. Furthermore, studies of the $\Lambda\eta$ channel in the strange-baryon sector suggest $\Lambda_c^+\eta$ to be an interesting channel. In addition to excited Λ states decaying into $\Lambda\eta$, such as the $\Lambda(2000)$ [14], a narrow enhancement was observed recently in the pK^- channel near the $\Lambda\eta$ threshold, which was identified as a threshold cusp [15].

Much attention has been paid to the pD^0 system as an analog to the *NK* system, where there exists the $\Lambda(1405)$ resonance as a *NK* quasibound state near threshold [16–18]. Thus the pD^0 system can be studied from the viewpoint of the description of the Λ_c charmed baryons. *BABAR* has reported $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ in the pD^0 mass spectrum [8], while Belle has not previously investigated the pD^0 spectrum in direct e^+e^- annihilation.

In this study, which is based on the full Belle dataset, we study for the first time the $\Lambda_c^+\eta$ system and search for singly-charmed baryons, including $\Lambda_c(2880)^+$, $\Lambda_c(2940)^+$ and other yet unknown states, in a mass region from 2.83 to 3.15 GeV/ c^2 . In addition, we measure ratios of branching fractions of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ decaying to $\Lambda_c^+\eta$ and pD^0 relative to the reference mode, $\Sigma_c(2455)\pi$, where $\Sigma_c(2455)\pi$ is the combination of $\Sigma_c(2455)^0\pi^+$ and $\Sigma_c(2455)^{++}\pi^-$.

II. DATA SAMPLE AND BELLE DETECTOR

We perform our analysis using a data sample corresponding to an integrated luminosity of 980 fb⁻¹, collected with the Belle detector at the KEKB asymmetricenergy e^+e^- collider [19,20]. Most of the data were recorded at the $\Upsilon(4S)$ resonance, the rest was collected at other $\Upsilon(nS)$ states, with n = 1, 2, 3, or 5, or at 60 MeV below the $\Upsilon(4S)$.

The Belle detector is a large-solid-angle magnetic spectrometer that consists of a silicon vertex detector (SVD), a 50-layer central drift chamber (CDC), an array of aerogel threshold Cherenkov counters (ACC), a barrellike arrangement of TOF scintillation counters, and an electromagnetic calorimeter (ECL) comprised of CsI(Tl) crystals. All these detector components are located inside a superconducting solenoid coil that provides a 1.5 T magnetic field. An iron flux-return located outside of the coil is instrumented to detect K_L^0 mesons and to identify muons. The detector is described in detail in Refs. [21,22].

We use samples of simulated $e^+e^- \rightarrow c\bar{c}$ Monte Carlo (MC) events for the optimization of selection criteria, the

estimation of background contributions, and the determination of signal detection efficiencies. MC samples are generated by EvtGen [23] and then propagated through a detector simulation based on Geant3 [24]. The package PYTHIA [25] is used to simulate $e^+e^- \rightarrow c\bar{c} \rightarrow \Lambda_c^{*+}X$ signal events, where Λ_c^{*+} stands for $\Lambda_c(2880)^+$, $\Lambda_c(2940)^+$, or one of possible new excited Λ_c states, and X denotes anything. The masses and widths of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ are fixed to the world-average values [26]. The masses of the unknown Λ_c s are set to values from 2.850 to 3.125 GeV/ c^2 in steps of 25 MeV/ c^2 , and for each mass value the widths are set to 0, 10, 20, 30, and 40 MeV. We generate the $\Lambda_c^{*+} \rightarrow \Lambda_c^+ \eta$ and $\Lambda_c(2880)^+ / \Lambda_c(2940)^+ \rightarrow$ $pD^0/\Sigma_c(2455)\pi$ samples with a phase-space model [27] and also the secondary decays $\Lambda_c^+ \to p K^- \pi^+ / p K_s^0, \eta \to \gamma \gamma$, $D^0 \rightarrow K^- \pi^+ / K^- \pi^+ \pi^0 / K^- \pi^+ \pi^+ \pi^-$, and $\Sigma_c(2455) \rightarrow \Lambda_c^+ \pi^$ are all generated with a phase-space model. The $\eta \rightarrow$ $\pi^0 \pi^+ \pi^-$ decay is generated with its known substructures [26]. We take the effect of final-state radiation from charged particles into account with the PHOTOS package [28]. We use the MC samples with zero width for Λ_c^{*+} in order to estimate the intrinsic mass resolution. Generic MC samples of $\Upsilon(1S, 2S, 3S)$ decays, $\Upsilon(4S) \rightarrow$ $B^+B^-/B^0\bar{B}^0$, $\Upsilon(5S) \to B_s^{(*)}\bar{B}_s^{(*)}/B^{(*)}\bar{B}^{(*)}(\pi)/\Upsilon(4S)\gamma$, and $e^+e^- \to q\bar{q} \ (q=u, d, s, c)$ at $\sqrt{s} = 10.52, 10.58,$ and 10.867 GeV, corresponding to two times the integrated luminosity of the data, are used to optimize selection criteria and perform background studies.

III. EVENT SELECTION

We study the Λ_c^{*+} in three decay modes; $\Lambda_c^+\eta$, pD^0 and $\Sigma_c(2455)\pi$ with $\Lambda_c^+ \to pK^-\pi^+/pK_S^0$, $\eta \to \gamma\gamma/\pi^0\pi^+\pi^-$, $D^0 \to K^-\pi^+/K^-\pi^+\pi^0/K^-\pi^+\pi^-$, and $\Sigma_c(2455) \to \Lambda_c^+\pi$. Event selections are optimized by maximizing a figure-ofmerit defined as ϵ/\sqrt{B} , where ϵ denotes the detection efficiency of the signal process and *B* is the number of background events obtained from generic MC samples in a mass region from 2.870 to 2.895 GeV/ c^2 , which is defined as the $\Lambda_c(2880)^+$ signal region.

Final-state charged particles, π^{\pm} , K^{\pm} , and $p(\bar{p})$, are identified using information from the tracking systems (SVD, CDC) and the particle identification detectors (CDC, ACC, TOF), which is combined into the likelihood ratio $\mathcal{L}(A|B) = \mathcal{L}(A)/[\mathcal{L}(A) + \mathcal{L}(B)]$, where *A* and *B* are π , *K*, or *p* as appropriate [29]. We require protons to have $\mathcal{L}(p|K) > 0.6$ and $\mathcal{L}(p|\pi) > 0.6$, kaons to have $\mathcal{L}(K|p) >$ 0.6 and $\mathcal{L}(K|\pi) > 0.6$, and pions to have $\mathcal{L}(\pi|p) > 0.6$ and $\mathcal{L}(\pi|K) > 0.6$. A requirement of $\mathcal{L}(e|\text{hadrons}) =$ $\mathcal{L}(e)/[\mathcal{L}(e) + \mathcal{L}(\text{hadrons})] < 0.95$ is applied for all charged particles to suppress electrons, where $\mathcal{L}(e)$ and $\mathcal{L}(\text{hadrons})$ are obtained from the ECL in addition to tracking and particle identification systems [30]. The momentum-averaged identification efficiencies of protons, kaons, and pions are 90%, 90%, and 93%, respectively. We also require the distance of the closest approach of all charged-particle tracks with respect to the interaction point to be less than 2.0 cm along the direction opposite to the e^+ beam and to be less than 0.2 cm in the transverse plane. In addition, invariant masses of pion, kaon, and proton pairs with the same sign are required to be larger than $0.28 \text{ GeV}/c^2$, $0.989 \text{ GeV}/c^2$, and $1.878 \text{ GeV}/c^2$, respectively, in order to reject duplicated tracks.

A cluster detected in the ECL not matching any tracks is identified as a photon candidate. Candidates for $\pi^0 \rightarrow \gamma \gamma$ are selected from photon pairs, where each photon has an energy greater than 0.05 GeV, and are discarded once the invariant mass of the two photons lies outside the range from 0.12 GeV/ c^2 to 0.15 GeV/ c^2 . We reconstruct the η candidates using two modes; $\gamma\gamma$ and $\pi^0\pi^+\pi^-$. Candidates for $\eta \rightarrow \gamma \gamma$ are reconstructed from photon pairs whose invariant mass satisfies $0.52 \text{ GeV}/c^2 < M(\gamma\gamma) <$ 0.57 GeV/ c^2 . The energies of both photons are required to be greater than 0.3 GeV. We select $\eta \to \pi^0 \pi^+ \pi^-$ candidates with $\pi^0 \rightarrow \gamma \gamma$ by requiring 0.540 GeV/ c^2 < $M(\pi^0 \pi^+ \pi^-) < 0.555 \text{ GeV}/c^2$, where $M(\pi^0 \pi^+ \pi^-)$ is the invariant mass of $\pi^0 \pi^+ \pi^-$. To suppress background events we require in addition that the photons from the $\eta \rightarrow$ $\pi^+\pi^-\pi^0$, $\pi^0 \to \gamma\gamma$ decay have an energy greater than 0.1 GeV. We adjust the four-momenta of the daughter particles within their uncertainties by constraining the invariant masses of the systems of final-state particles to the nominal π^0 and η masses [26], respectively.

Candidates for $K_S^0 \to \pi^+ \pi^-$ are reconstructed from pairs of oppositely charged particles taken as pions, using an artificial neural network [31,32]. We require $|M(\pi^+\pi^-) - m(K_S^0)| < 10 \text{ MeV}/c^2$, where $m(K_S^0)$ is the nominal $K_{\rm S}^0$ mass [26]. The two pions are refitted to have a common vertex and the $\pi^+\pi^-$ invariant mass is constrained to $m(K_S^0)$. The Λ_c^+ candidates are reconstructed from $pK^{-}\pi^{+}$ and pK_{S}^{0} modes. The invariant masses of $pK^{-}\pi^{+}$ and pK_s^0 are required to be within $\pm 10 \text{ MeV}/c^2$ of the nominal Λ_c^+ mass $m(\Lambda_c^+)$ [26]. For the selected events, we constrain $M(pK^{-}\pi^{+})$ and $M(pK^{0}_{S})$ to $m(\Lambda^{+}_{C})$ to improve momentum resolution. We reconstruct $\Sigma_c(2455)^{0,++}$ candidates by combining a Λ_c^+ candidate and a charged pion. The requirements of $|M(\Lambda_c^+\pi^-) - m(\Sigma_c(2455)^0)| <$ $|M(\Lambda_c^+\pi^+) - m(\Sigma_c(2455)^{++})| <$ $5 \text{ MeV}/c^2$ and 5 MeV/ c^2 , defined as $\Sigma_c(2455)$ signal region (SR), are adopted to select $\Sigma_c(2455)^{0,++}$ candidates, where $m(\Sigma_c(2455)^{0,++})$ is the nominal $\Sigma_c(2455)^{0,++}$ mass [26]. The $\Sigma_c(2455)$ sideband regions (SB) are defined as $|M(\Lambda_c^+\pi^-) - m(\Sigma_c(2455)^0)| \in (15, 20) \text{ MeV}/c^2$ and $|M(\Lambda_c^+\pi^+) - m(\Sigma_c(2455)^{++})| \in (15,20) \text{ MeV}/c^2$. Figure 1 shows the $M(\Lambda_c^+\pi^-)$ and $M(\Lambda_c^+\pi^+)$ spectra.

The D^0 candidates are reconstructed from $K^-\pi^+$, $K^-\pi^+\pi^-\pi^-$, and $K^-\pi^+\pi^0$ modes, using almost the same selection criteria as those used in the study of Ξ_c states



FIG. 1. Invariant-mass distributions of (a) $\Lambda_c^+ \pi^-$ and (b) $\Lambda_c^+ \pi^+$. Regions between the two red dotted lines are the $\Sigma_c(2455)$ signal regions, and regions between the two blue long-dashed lines are the $\Sigma_c(2455)$ sideband regions.

in the ΛD final state [33]. Candidates satisfying $|M(K^-\pi^+) - m(D^0)| < 14 \text{ MeV}/c^2$, $|M(K^-\pi^+\pi^+\pi^-) - m(D^0)| < 11 \text{ MeV}/c^2$, and $|M(K^-\pi^+\pi^0) - m(D^0)| < 27 \text{ MeV}/c^2$, where $m(D^0)$ is the nominal D^0 mass [26], are selected as D^0 candidates. For the $K^-\pi^+\pi^0$ mode, the energy of the π^0 candidate is required to be greater than 0.5 GeV to further suppress combinatorial background events. We also require the daughter particles of D^0 to originate from a common vertex and we constrain the invariant mass of the system of final-state particles to the nominal D^0 mass [26]. The sideband regions of D^0 are selected as $|M(K^-\pi^+) - m(D^0)| \in (23, 30) \text{ MeV}/c^2$, $|M(K^-\pi^+\pi^-) - m(D^0)| \in (24, 30) \text{ MeV}/c^2$, and $|M(K^-\pi^+\pi^0) - m(D^0)| \in (38, 50) \text{ MeV}/c^2$.

All the above required mass ranges around the signal peaks approximately correspond to $\pm 2.5\sigma$ intervals, where σ is the standard deviation of the peak distributions. Finally, we reconstruct Λ_c^{*+} candidates from $\Lambda_c^+\eta$, pD^0 , or $\Sigma_c(2455)\pi$. We define the scaled momentum x_p of Λ_c^{*+} as $x_p = p^*/\sqrt{s/4 - M^2}$ [34] with p^* being the momentum of Λ_c^{*+} in the e^+e^- center-of-mass frame, *s* the square of the center-of-mass energy, and *M* the mass of Λ_c^{*+} . We require

 $x_p > 0.7$ to suppress the combinatorial background, especially from *B*-meson decays. There are 5.8% of the $\Lambda_c^{*+} \rightarrow \Lambda_c^+ \eta$ events that have multiple candidates and only the candidate with the smallest $\chi^2(\Lambda_c^+) + \chi^2(\eta)$ is preserved, where χ^2 stands for the quality of the mass-constrained fit.

IV. SEARCH FOR CHARMED BARYONS IN THE $\Lambda_c^+ \eta$ SYSTEM

Figure 2 shows the invariant-mass distributions of $\Lambda_c^+ \eta$ with $\eta \to \gamma \gamma$ and $\eta \to \pi^0 \pi^+ \pi^-$, obtained after applying the selection criteria discussed in Sec. III.

A series of simultaneous binned maximum-likelihood fits to the measured mass spectra of $\Lambda_c^+ \eta$ with $\eta \to \gamma \gamma$ and $\eta \to \pi^0 \pi^+ \pi^-$ are performed. The probability density function (PDF) for the signal component is represented by a constant-width relativistic Breit-Wigner (BW) function convolved with a Gaussian distribution to take the intrinsic mass resolution into account, where the intrinsic mass resolution is determined from signal MC samples. The background component is parametrized by a threshold function of the form



FIG. 2. An example of the simultaneous fit to the $M(\Lambda_c^+ \eta)$ spectra with (a) $\eta \to \gamma \gamma$ and (b) $\eta \to \pi^0 \pi^+ \pi^-$ (points with statistical uncertainties). The blue solid curves represent the result of the fit. The red dashed curves represent a hypothetical excited Λ_c state with $M = 2.95 \text{ GeV}/c^2$ and $\Gamma = 20 \text{ MeV}$, while the magenta dash-dotted curves represent the background.



FIG. 3. Statistical significances for the various excited Λ_c^+ hypotheses considered.

$$\left(1-e^{-\frac{M-M_{\rm th}}{\mu}}\right)\left(\frac{M}{M_{\rm th}}\right)^a+b\left(\frac{M}{M_{\rm th}}-1.0\right)+c\left(\frac{M}{M_{\rm th}}-1.0\right)^2.$$
(1)

Here, *M* represents the $\Lambda_c^+\eta$ mass, $M_{\rm th}$ stands for the threshold mass, which is fixed to the sum of the nominal Λ_c^+ and η masses [26], and *a*, *b*, *c*, and μ are free parameters. The fit is performed with a 0.5 MeV/ c^2 bin width, while the histograms in Fig. 2 are shown with merged bins of 5 MeV/ c^2 .

We constrain the signal mass and width to be identical for the two η decay modes, and fix the signal mass to be a value from 2.850 GeV/ c^2 to 3.125 GeV/ c^2 in steps of 25 MeV/ c^2 . The signal width is fixed to 0, 10, 20, 30, and 40 MeV for each fixed mass. The ratio of signal yields for the $\eta \rightarrow \gamma\gamma$ mode relative to those for the $\eta \rightarrow \pi^0 \pi^+ \pi^$ mode is calculated by the product of detection efficiencies and the known η branching fractions [26], and is fixed in the fit. This ratio decreases from 5.4 to 2.7 as the signal mass increases from 2.850 GeV/ c^2 to 3.125 GeV/ c^2 . Figure 2 shows an example of the fit with the signal mass and width fixed to 2.95 GeV/ c^2 and 20 MeV, respectively.

The statistical significance of the signal is calculated using the log-likelihood difference $-2\ln(\mathcal{L}_0/\mathcal{L}_{max})$, where \mathcal{L}_{max} and \mathcal{L}_0 are the likelihood values of the simultaneous fit to the $\Lambda_c^+\eta$ mass spectra with and without including the signal PDF, respectively. The significance is negative when the number of fitted signal events is itself negative. The set of statistical significances obtained from the fits is shown in Fig. 3. Significances are below 3σ for all assumed excited Λ_c^+ hypotheses. We hence find no significant Λ_c^{*+} signals in the $\Lambda_c^+\eta$ invariant-mass spectra.

V. RELATIVE BRANCHING FRACTION

Figure 4 shows the invariant-mass distributions of $\Lambda_c^+ \eta$ with $\eta \to \gamma\gamma$, $\Lambda_c^+ \eta$ with $\eta \to \pi^0 \pi^+ \pi^-$, pD^0 , $\Sigma_c(2455)\pi$ in the $\Sigma_c(2455)$ signal region, and $\Sigma_c(2455)\pi$ in the



FIG. 4. Measured invariant-mass distributions of (a) $\Lambda_c^+\eta$ with $\eta \to \gamma\gamma$, (b) $\Lambda_c^+\eta$ with $\eta \to \pi^+\pi^-\pi^0$, (c) pD^0 , (d) $\Sigma_c(2455)\pi$ in the $\Sigma_c(2455)$ signal region, and (e) $\Sigma_c(2455)\pi$ in the normalized $\Sigma_c(2455)$ sideband regions (points with statistical uncertainties). The blue solid curve represents the result of a simultaneous fit. The red and dark green dashed curves represent the contributions of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$, respectively, while the magenta dash-dotted curves represent the background.

normalized $\Sigma_c(2455)$ sideband regions obtained after applying the selection criteria discussed in Sec. III. The number of normalized $\Sigma_c(2455)$ sideband events equals to the number of background events in the $\Sigma_c(2455)$ signal region. No clear $\Lambda_c(2880)^+$ or $\Lambda_c(2940)^+$ signals are seen in the $\Lambda_c^+\eta$ invariant-mass spectra [see Figs. 4(a) and 4(b)]. By contrast, we observe significant $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signals in the $M(pD^0)$ distribution [see Fig. 4(c)]. We do not observe any such peaks in the pD^0 mass spectra from generic MC events [35], nor in the data from the D^0 mass sideband regions. Clear $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signals are also seen in the $M(\Sigma_c(2455)\pi)$

[SR] distribution [see Fig. 4(d)]. A $\Lambda_c(2880)^+$ signal is seen in $\Sigma_c(2455)$ sideband events [see Fig. 4(e)]. This is expected and is due to the $\Lambda_c(2880)^+ \rightarrow \Lambda_c^+ \pi^+ \pi^-$ nonresonant decay, which in this analysis we consider to be a peaking background to our $\Sigma_c(2455)\pi$ mode. The contribution of this background is estimated from $\Sigma_c(2455)$ sideband events.

The yields of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ are obtained from a simultaneous binned maximum-likelihood fit to the measured mass spectra of $\Lambda_c^+\eta$ with $\eta \to \gamma\gamma$, $\Lambda_c^+\eta$ with $\eta \to \pi^0 \pi^+ \pi^-$, pD^0 , $\Sigma_c(2455)\pi$ in the $\Sigma_c(2455)$ signal region, and $\Sigma_c(2455)\pi$ in the normalized $\Sigma_c(2455)$ sideband regions. In this fit, the masses of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ are free parameters, constrained to be identical for all decay modes, and the widths are fixed to worldaverage values [26]. The fit is performed with a 0.5 MeV/ c^2 bin width, while histograms in Fig. 4 are shown with merged bins of 5 MeV/ c^2 . The fitted mass range starts at 2.83 GeV/ c^2 because of the $\Lambda_c^+\eta$ threshold. Lower $\Sigma_c(2455)\pi$ and $\Lambda_c^+\pi^+\pi^-$ masses are populated by the tail of the $\Lambda_c(2765)^+$ [7] and have been studied already in Ref. [9].

The PDF for the $\Lambda_c(2880)^+$ or $\Lambda_c(2940)^+$ signal component is represented by a relativistic BW function with mass-dependent width that is convolved with a Gaussian distribution to take the intrinsic mass resolution into account. We determine the intrinsic mass resolution based on signal MC samples. As the most plausible J^P assignment for $\Lambda_c(2880)^+$ is $5/2^+$ and that for $\Lambda_c(2940)^+$ is $3/2^{-}$, the two signals are parametrized by an F-wave BW function and a D-wave BW function, respectively, using Blatt-Weisskopf form factors [36]. The background component in the $M(\Lambda_c^+\eta)$ spectra for $\eta \to \gamma\gamma$ and $\eta \to \pi^0\pi^+\pi^$ is parametrized by the function in Eq. (1). A sixth-order Chebychev polynomial is used to model the combinatorial background component in the $M(pD^0)$ spectrum. The background components in the $M(\Sigma_c(2455)\pi)$ [SR] and $M(\Sigma_c(2455)\pi)$ [SB] spectra are parametrized by secondorder Chebychev polynomials.

Since no $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signals are observed in the $\Lambda_c^+\eta$ mass spectra, we constrain the ratios of yields of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ in the $\Lambda_c^+\eta$ decay channel with $\eta \to \gamma\gamma$ relative to those with $\eta \to \pi^0\pi^+\pi^-$ by the products of the detection efficiencies and the known η branching fractions [26], resulting in ratio values of 5.1 for $\Lambda_c(2880)^+$ and 4.4 for $\Lambda_c(2940)^+$. The result of the simultaneous fit is shown in Fig. 4, and the signal yields are summarized in Table I. The reduced χ^2 values of the fit are $\chi^2/d.o.f = 1.2$ (746.6/635) for $\Lambda_c^+\eta(\eta \to \gamma\gamma)$, 1.5 (937.3/633) for $\Lambda_c^+\eta(\eta \to \pi^0\pi^+\pi^-)$, 1.1 (695.4/629) for pD^0 , 1.6 (1049.2/634) for $\Sigma_c(2455)\pi$ in the $\Sigma_c(2455)$ signal region, and 0.5 (297.0/633) for $\Sigma_c(2455)\pi$ in the normalized $\Sigma_c(2455)$ sideband regions, where d.o.f. is the number of degrees of freedom.

TABLE I. Signal yields from the simultaneous fit to the mass spectra in Fig. 4. The uncertainties shown are statistical only.

	$\Lambda_c(2880)^+$	$\Lambda_c(2940)^+$
$\overline{N_{\Lambda_c^+\eta;\eta\to\gamma\gamma}}$	$(-0.2 \pm 0.7) \times 10^2$	$(0.7 \pm 1.4) \times 10^2$
$N_{\Lambda_c^+\eta;\eta\to\pi^0\pi^+\pi^-}$	$(-0.04 \pm 0.13) \times 10^{2}$	$(0.16 \pm 0.31) \times 10^2$
$N_{\Sigma_c(2455)\pi}$	$(4.25 \pm 0.15) \times 10^3$	$(1.19 \pm 0.19) \times 10^3$
N _{nonres}	$(1.6\pm0.4) imes10^2$	$(-0.01 \pm 0.03) \times 10^{2}$
N_{pD^0}	$(1.17 \pm 0.04) \times 10^4$	$(1.64 \pm 0.09) \times 10^4$

The possible structures around 2.93 GeV/ c^2 , 2.97 GeV/ c^2 , and 3.09 GeV/ c^2 in Fig. 4(b), which make large contributions to the χ^2 , are regarded as statistical fluctuations, since no corresponding structures are found in Fig. 4(a). The fit to the $M(\Sigma_c(2455)\pi)$ spectrum in SR, as shown in Fig. 4(d), shows deviations from the data, and implies that there might be contributions from other states not included in the fit, such as the $\Lambda_c(2910)^+$ [37]. The impact from the $\Lambda_c(2910)^+$ is considered in the systematic uncertainty, as discussed in Sec. VI.

We measure ratios of branching fractions,

$$R_{\rm X}(2880/2940) \equiv \frac{\mathcal{B}(\Lambda_c(2880/2940)^+ \to {\rm X})}{\mathcal{B}(\Lambda_c(2880/2940)^+ \to \Sigma_c(2455)\pi)},$$
(2)

where X denotes the $\Lambda_c^+\eta$ or pD^0 mode and $\mathcal{B}(\Lambda_c(2880/2940)^+ \rightarrow \Sigma_c(2455)\pi)$ is the sum of branching fractions for $\Sigma_c(2455)^0\pi^+$ and $\Sigma_c(2455)^{++}\pi^-$. The two branching fractions are taken as being the same as predicted by isospin symmetry. These ratios are obtained from the measured signal yields using

$$R_{\rm X} = \frac{N_{\rm X}}{\sum_i \epsilon_{\rm X}^i \mathcal{B}_{\rm X}^i} \left/ \frac{N_{\Sigma_c(2455)\pi} - N_{\rm nonres}}{\sum_j \epsilon_{\Sigma_c(2455)\pi}^j \mathcal{B}_{\Sigma_c(2455)\pi}^j}, \qquad (3)$$

where the index *i* represents subdecay channels for the pD^0 or $\Lambda_c^+\eta$ modes, the index *j* represents subdecay channels for the $\Sigma_c(2455)\pi$ mode, the ϵ are the detection efficiencies for the decay defined by X and *i*, the \mathcal{B} are the products of branching fractions of intermediate states for the respective decay channels, and the N stand for the signal yields given in Table I. The N_{nonres} are the fitted $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signal yields in the $M(\Sigma_c(2455)\pi)$ [SB] spectrum. We estimate the detection efficiencies from the MC samples (see Sec. II), as summarized in Table II, and we calculate the products of known branching fractions using the world-average values from Ref. [26].

We obtain branching-fraction ratios for the pD^0 mode of $R_{pD^0}(2880) = 0.75 \pm 0.03 \pm 0.07$ and $R_{pD^0}(2940) =$ $3.59 \pm 0.21 \pm 0.56$, where the uncertainties are statistical and systematic (see Sec. VI below for details on the latter ones).

TABLE II. Detection efficiencies from the MC samples for each decay mode.

Decays	$\Lambda_c(2880)^+$	$\Lambda_c(2940)^+$
$\frac{1}{\Lambda_c^+ \eta(pK^-\pi^+,\gamma\gamma)}$	0.0518	0.0567
$\Lambda_c^+ \eta(pK_S^0, \gamma\gamma)$	0.0578	0.0662
$\Lambda_{c}^{+}\eta(pK^{-}\pi^{+},\pi^{0}\pi^{+}\pi^{-})$	0.0176	0.0224
$\Lambda_c^+ \eta (pK_s^0, \pi^0 \pi^+ \pi^-)$	0.0198	0.0241
$pD^0(K^-\pi^+)$	0.273	0.289
$pD^{0}(K^{-}\pi^{+}\pi^{+}\pi^{-})$	0.091	0.096
$pD^{0}(K^{-}\pi^{+}\pi^{0})$	0.176	0.185
$\Sigma_c(2455)\pi(pK^-\pi^+)$	0.137	0.144
$\Sigma_c(2455)\pi(pK_S^0)$	0.162	0.170

Since no significant signals of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ decaying to $\Lambda_c^+\eta$ are seen, we set upper limits R^{UL} on the branching-fraction ratios at 90% credibility level (CL) by solving the equation

$$\frac{\int_0^{R^{\text{UL}}} \mathcal{L}(R) dR}{\int_0^{+\infty} \mathcal{L}(R) dR} = 0.9.$$
(4)

Here, *R* is the assumed branching-fraction ratio and $\mathcal{L}(R)$ is the corresponding likelihood value obtained from fitting the data. Before integrating, we include the systematic uncertainty σ_{sys} described in Sec. VI by convolving the likelihood function with a Gaussian function whose width is equal to σ_{sys} . We obtain upper limits on the branching-fraction ratios for the $\Lambda_c^+\eta$ mode at 90% CL of $R_{\Lambda_c^+\eta}(2880) < 0.13$ and $R_{\Lambda_c^+\eta}(2940) < 1.11$.

VI. SYSTEMATIC UNCERTAINTIES

The systematic uncertainties of the branching-fraction ratios are listed in Table III. We include uncertainties from the particle identification (PID) efficiency, the tracking efficiency, the K_s^0 , π^0 , η , and $\Sigma_c(2455)$ selection efficiency,

the branching fractions of intermediate states, the finite MC sample size, the mass resolution, the signal yield extraction, the fit bias, and the impact of $\Lambda_c(2910)^+$ in the $M(\Sigma_c(2455)\pi)$ spectra.

The uncertainty from the charged kaon and pion identification efficiency is estimated using $D^{*+} \rightarrow D^0 \pi^+$ with $D^0 \rightarrow K^- \pi^+$, and that of the proton identification efficiency is estimated using $\Lambda^0 \rightarrow p\pi^-$. The signal selection efficiency is corrected by a factor derived from the ratio of identification efficiencies of data and MC. We treat the statistical uncertainty of these correction factor as the systematic uncertainty. As the number of kaons, pions, and protons depends on the analyzed subdecay channels, the uncertainty is calculated by weighting with the product of known branching fractions and detection efficiencies. The uncertainty associated with the PID efficiencies for the Λ_c^+ daughters cancels for $R_{\Lambda_c^+}n$.

The uncertainty of the tracking efficiency is assigned to be 0.35% per track, which is estimated using partially reconstructed $D^{*+} \rightarrow D^0 \pi^+$ with $D^0 \rightarrow \pi^+ \pi^- K_S^0$. Considering that the number of tracks differs in each subdecay channel, a weighted uncertainty is calculated.

We evaluate the uncertainty coming from the K_S^0 selection efficiency using $D^* \to \pi D^0$ with $D^0 \to K_S^0 \pi^0$. The ratio of efficiencies for data and MC is $(98.57 \pm 0.40)\%$, from which we quote the systematic uncertainty of 1.8%. Since the K_S^0 only appears in $\Lambda_c^+ \to p K_S^0$, a weighted uncertainty is calculated.

Systematic uncertainties due to π^0 and η selection efficiencies are both estimated to be 3% [38]. For R_{pD^0} , a weighted uncertainty is calculated since the π^0 only appears in $D^0 \to K^- \pi^+ \pi^0$.

The ratio of efficiencies of the $M(\Lambda_c^+\pi^\pm)$ requirement for data and MC simulation is 0.968 ± 0.016 for $\Lambda_c(2880)^+$, and 0.965 ± 0.022 for $\Lambda_c(2940)^+$. We correct detection efficiencies for the $\Sigma_c(2455)\pi$ mode with these ratios and take relative uncertainties of ratios as systematic

TABLE III. Relative systematic uncertainties (in %) for the branching-fraction ratios.

Sources	$R_{pD^0}(2880)$	$R_{pD^0}(2940)$	$R_{\Lambda_c^+\eta}(2880)$	$R_{\Lambda_c^+\eta}(2940)$
PID efficiency	4.5	4.5	3.5	3.5
Tracking efficiency	0.5	0.5	0.6	0.6
$K_{\rm S}^0$ selection	0.5	0.5		
π^0 selection	0.9	0.9		
η selection			3.0	3.0
$\dot{\Sigma}_{c}(2455)$ selection	1.6	2.3	1.6	2.3
Branching fraction	5.1	5.1	0.6	0.6
MC sample size	1.3	1.2	2.2	2.2
Signal yield extraction	6.2	12.5	11.2	13.5
Fit bias	1.0	2.3	4.1	9.6
$\Lambda_c(2910)^+$	1.5	5.7	3.9	8.2
Total	9.7	15.7	13.7	19.3

uncertainties, which are 1.6% for $\Lambda_c(2880)^+$ and 2.3% for $\Lambda_c(2940)^+$.

The systematic uncertainty from the branching fractions of intermediate states are taken from world-average values in Ref. [26]. The uncertainty related to the branching fractions of the Λ_c^+ and K_s^0 decays cancels for $R_{\Lambda_r^+n}$.

The relative uncertainty due to the finite MC sample size is estimated as $\sqrt{(1-\epsilon)/(\epsilon N_{\text{gen}})}$, where ϵ is the detection efficiency and N_{gen} is the number of generated signal MC events.

The systematic uncertainty due to the fixed mass resolution is estimated by fitting the $M(pD^0)$ spectrum with the mass resolution floated. The relative difference on the ratio of the branching fractions between the one from the floated mass resolution and the nominal value is taken as the uncertainty, which is found to be negligible.

The systematic uncertainty from the signal yield extraction consists of uncertainties from the widths of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$, the background PDF, the fit interval, and the bin width. To estimate the uncertainty introduced by fixing the widths of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$, we vary the signal width by $\pm 1\sigma$ [26] and take the maximum difference from the nominal result as the systematic uncertainty. The uncertainty arising from the background PDF is determined by varying the order of the Chebychev polynomial. The uncertainty from the fit range is estimated by enlarging the fitted interval from $(2.83, 3.15) \text{ GeV}/c^2$ to (2.81, 3.17) GeV/ c^2 . We evaluate the uncertainty due to the bin width by varying it from $0.5 \text{ MeV}/c^2$ to $0.6 \text{ MeV}/c^2$ or $0.4 \text{ MeV}/c^2$, and taking the maximum difference as the systematic uncertainty. Of the above uncertainties, the ones due to the signal width and fit interval are dominant and those from the background PDF and the bin width are small. The various contributions to the yield uncertainty are added in quadrature.

The systematic uncertainty introduced by the fit bias is assessed by constructing an ensemble of MC pseudoexperiments. For each pseudoexperiment, we generate simulated MC samples, by randomly drawing signal and background MC events from the signal and background PDFs, respectively, based on the extracted numbers of corresponding events in data. The sizes of these MC samples are hence identical to the extracted yields for each decay mode. We repeat the above procedure 1000 times and then perform simultaneous binned maximum-likelihood fits on these 1000 independent pseudoexperiments. This yields Gaussian-shaped distributions of the R_X values. The relative differences between the mean values of these distributions and the nominal R_X values are assigned as the systematic uncertainties.

The systematic uncertainty from the possible $\Lambda_c(2910)^+$ signal in the $M(\Sigma_c(2455)\pi)$ spectrum around 2.90–2.95 GeV/ c^2 is estimated by adding an additional state, the $\Lambda_c(2910)^+$, in the fit to the $M(\Sigma_c(2455)\pi)$ spectra. The relative difference of the branching-fraction ratio is taken as the systematic uncertainty.

Summing all the uncertainty terms discussed above in quadrature gives the total systematic uncertainty quoted in Table III.

VII. SUMMARY

In this paper, we report on a search for excited singlycharmed baryons performed for the first time in the $\Lambda_c^+ \eta$ mass spectra in a range from 2.83 GeV/ c^2 to 3.15 GeV/ c^2 based on the full Belle dataset corresponding to an integrated luminosity of 980 fb⁻¹. No significant excess is found in the $M(\Lambda_c^+ \eta)$ spectrum. This is in contrast to excited hyperons, where resonances decaying into $\Lambda \eta$ have been observed.

Clear $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ signals are observed in the pD^0 mass spectrum. The first measurements of branching-fraction ratios of $\Lambda_c(2880)^+$ and $\Lambda_c(2940)^+$ decaying to $\Lambda_c^+\eta$ and pD^0 relative to $\Sigma_c(2455)\pi$ are performed. We measure

$$R_{pD^0}(2880) = 0.75 \pm 0.03 \pm 0.07,$$

$$R_{pD^0}(2940) = 3.59 \pm 0.21 \pm 0.56,$$

where the first uncertainty is statistical and the second one systematic. We also determine upper limits for $R_{\Lambda_c^+\eta}$ at 90% CL of

$$R_{\Lambda_c^+\eta}(2880) < 0.13,$$

 $R_{\Lambda_c^+\eta}(2940) < 1.11.$

It is predicted in Ref. [6] that the ρ -mode states decay primarily into a light meson and a heavy baryon, whereas the λ -mode states decay predominantly into a light baryon and a heavy meson. Therefore, when interpreted using this model, our branching-ratio measurements suggest that the $\Lambda_c(2880)^+$ is a ρ -mode excited state while the $\Lambda_c(2940)^+$ is a λ -mode excited state.

ACKNOWLEDGMENTS

This work, based on data collected using the Belle detector, which was operated until June 2010, was supported by the Ministry of Education, Culture, Sports, Science, and Technology (MEXT) of Japan, the Japan Society for the Promotion of Science (JSPS), and the Tau-Lepton Physics Research Center of Nagoya University; the Australian Research Council including DP210101900, DP210102831, Grants No. No. No. DE220100462, No. LE210100098, and LE230100085; Austrian Federal Ministry No. of Education, Science and Research and Austrian Science Fund (FWF) [grant DOI: 10.55776/P35891] No. P 31361-N36; National Key R&D Program of China under Contract No. 2022YFA1601903, National Natural Science Foundation of China and research Grants No. 11575017, No. 11761141009, No. 11705209, No. 11975076, No. 12135005, No. 12150004, No. 12161141008, and No. 12175041, and Shandong Provincial Natural Science Foundation Project ZR2022JQ02; the Czech Science Foundation Grant No. 22-18469S; Horizon 2020 ERC Advanced Grant No. 884719 and ERC Starting Grant No. 947006 "InterLeptons" (European Union); the Carl Zeiss Foundation, the Deutsche Forschungsgemeinschaft, the Excellence Cluster Universe, and the VolkswagenStiftung; the Department of Atomic Energy (Project Identification No. RTI 4002), the Department of Science and Technology of India, and the UPES (India) SEED finding Programs No. UPES/R&D-SEED-INFRA/ 17052023/01 and No. UPES/R&D-SOE/20062022/06; the Istituto Nazionale di Fisica Nucleare of Italy; National Research Foundation (NRF) of Korea Grants No. 2016R1D1A1B02012900, No. 2018R1A2B3003643, No. 2018R1A6A1A06024970, No. RS202200197659, No. 2019R1I1A3A01058933, No. 2021R1A6A-1A03043957, No. 2021R1F1A1060423, No. 2021R1, F1A1064008, and No. 2022R1A2C1003993; No. Radiation Science Research Institute, Foreign Large-size Research Facility Application Supporting project, the Global Science Experimental Data Hub Center of the Korea Institute of Science and Technology Information and KREONET/GLORIAD; the Polish Ministry of Science and Higher Education and the National Science Center; the Ministry of Science and Higher Education of the Russian Federation and the HSE University Basic Research Program, Moscow; University of Tabuk research Grants No. S-1440-0321, No. S-0256-1438, and No. S-0280-1439 (Saudi Arabia); the Slovenian Research Agency Grants No. J1-9124 and No. P1-0135; Ikerbasque, Basque Foundation for Science, and the State Agency for Research of the Spanish Ministry of Science and Innovation through Grant No. PID2022-136510NB-C33 (Spain); the Swiss National Science Foundation; the Ministry of Education and the National Science and Technology Council of Taiwan; and the United States Department of Energy and the National Science Foundation. These acknowledgements are not to be interpreted as an endorsement of any statement made by any of our institutes, funding agencies, governments, or their representatives. We thank the KEKB group for the excellent operation of the accelerator; the KEK cryogenics group for the efficient operation of the solenoid; and the KEK computer group and the Pacific Northwest National Laboratory (PNNL) Environmental Molecular Sciences Laboratory (EMSL) computing group for strong computing support; and the National Institute of Informatics, and Science Information NETwork 6 (SINET6) for valuable network support.

- [1] J. M. Richard, Phys. Rep. 212, 1 (1992).
- [2] E. Klempt and J. M. Richard, Rev. Mod. Phys. 82, 1095 (2010).
- [3] V. Crede and W. Roberts, Rep. Prog. Phys. **76**, 076301 (2013).
- [4] H. X. Chen, W. Chen, X. Liu, Y. R. Liu, and S. L. Zhu, Rep. Prog. Phys. 80, 076201 (2017).
- [5] H. Y. Cheng, Chin. J. Phys. (Taipei) 78, 324 (2022).
- [6] T. Yoshida, E. Hiyama, A. Hosaka, M. Oka, and K. Sadato, Phys. Rev. D 92, 114029 (2015).
- [7] M. Artuso *et al.* (CLEO Collaboration), Phys. Rev. Lett. 86, 4479 (2001).
- [8] B. Aubert *et al.* (BABAR Collaboration), Phys. Rev. Lett. 98, 012001 (2007).
- [9] R. Mizuk *et al.* (Belle Collaboration), Phys. Rev. Lett. 98, 262001 (2007).
- [10] R. Aaij *et al.* (LHCb Collaboration), J. High Energy Phys. 05 (2017) 030.
- [11] J.R. Zhang, Phys. Rev. D 89, 096006 (2014).
- [12] Y. Dong, Phys. Rev. D 81, 014006 (2010).
- [13] P. G. Ortega, D. R. Entem, and F. Fernandez, Phys. Lett. B 718, 1381 (2013).
- [14] H. Zhang, J. Tulpan, M. Shrestha, and D. M. Manley, Phys. Rev. C 88, 035205 (2013).

- [15] S. B. Yang *et al.* (Belle Collaboration), Phys. Rev. D 108, L031104 (2023).
- [16] N. Kaiser, P.B. Siegel, and W. Weise, Nucl. Phys. A594, 325 (1995).
- [17] E. Oset and A. Ramos, Nucl. Phys. A635, 99 (1998).
- [18] J. A. Oller and U. G. Meißner, Phys. Lett. B 500, 263 (2001).
- [19] S. Kurokawa and E. Kikutani, Nucl. Instrum. Methods Phys. Res., Sect. A 499, 1 (2003), and other papers included in this volume.
- [20] T. Abe *et al.*, Prog. Theor. Exp. Phys. **2013**, 03A001 (2013), and references therein.
- [21] J. Brodzicka *et al.*, Prog. Theor. Exp. Phys. **2012**, 04D001 (2012).
- [22] A. Abashian *et al.* (Belle Collaboration), Nucl. Instrum. Methods Phys. Res., Sect. A 479, 117 (2002).
- [23] D. J. Lange, Nucl. Instrum. Methods Phys. Res., Sect. A 462, 152 (2001).
- [24] R. Brun *et al.*, Geant 3: User's guide Geant 3.10, Geant 3.11, CERN Report No. DD/EE/84-1, 1984.
- [25] T. Sjöstrand, S. Mrenna, and P. Skands, Comput. Phys. Commun. 178, 852 (2008).
- [26] R. L. Workman *et al.* (Particle Data Group), Prog. Theor. Exp. Phys. **2022**, 083C01 (2022) and 2023 update.

- [27] Charge-conjugate modes are implied throughout this paper unless otherwise stated.
- [28] E. Barberio and Z. Was, Comput. Phys. Commun. **79**, 291 (1994).
- [29] E. Nakano, Nucl. Instrum. Methods Phys. Res., Sect. A 494, 402 (2002).
- [30] K. Hanagaki, H. Kakuno, H. Ikeda, T. Iijima, and T. Tsukamoto, Nucl. Instrum. Methods Phys. Res., Sect. A 485, 490 (2002).
- [31] M. Feindt and U. Kerzel, Nucl. Instrum. Methods Phys. Res., Sect. A 559, 190 (2006).
- [32] H. Nakano, Ph.D. thesis, Tohoku University, 2014, Chap. 4, http://hdl.handle.net/10097/58814.
- [33] Y. Kato *et al.* (Belle Collaboration), Phys. Rev. D **94**, 032002 (2016).
- [34] We used units in which the speed of light is c = 1.
- [35] X. Y. Zhou, S. X. Du, G. Li, and C. P. Shen, Comput. Phys. Commun. 258, 107540 (2021).
- [36] S. Godfey and N. Isgur, Phys. Rev. D 32, 189. (1985).
- [37] Y. B. Li *et al.* (Belle Collaboration), Phys. Rev. Lett. **130**, 031901 (2023).
- [38] M.C. Chang et al., Phys. Rev. D 85, 091102 (2012).