

Large-scale microwave background fluctuations: Gauge-invariant formalism

Mirosław Panek*

NASA/Fermilab Astrophysics Group, P.O. Box 500, Batavia, Illinois 60510

(Received 16 January 1986)

The explicitly gauge-invariant formula for the large-scale fluctuations of the temperature of the microwave background radiation is obtained. The formula is applicable to the wide class of cosmological models (i.e., multicomponent and nonflat) based on Robertson-Walker metrics. Some specific cases are briefly discussed and in the case of general flat models the amplitudes of the multipole moments of the temperature pattern are obtained as functions of the baryonic perturbation quantities.

I. INTRODUCTION

The observed isotropy of the microwave background radiation (MBR) puts stringent limits on parameters of cosmological models. The requirement of reconciliation of low-level MBR fluctuations with the amplitude of the density perturbations necessary to produce galaxies today is a powerful tool for eliminating cosmological scenarios.

In order to calculate the small-scale MBR fluctuations one has to take the dynamics of the decoupling into account and the theory of this process is now relatively well known.¹ When dealing with fluctuations on scales larger than the horizon at the decoupling we can assume that decoupling occurs instantaneously. The distance that photons can travel is smaller than the characteristic length of the perturbation and the details of the decoupling process cannot influence the MBR pattern. However, there is another problem with the large scales. Because of the freedom of making gauge transformations (i.e., changing the correspondence between the points in the physical spacetime and the points in the undisturbed background) the perturbations of physical quantities are different in different gauges and can contain the spurious gauge modes.

The most elegant way of treating cosmological perturbations was proposed by Bardeen² and is based on using the gauge-invariant quantities that within the horizon are perturbations of well-known physical quantities. Since the paper by Bardeen some developments of his formalism were done, e.g., on the case of uncoupled fluids in the flat universe³ and the system of fluid plus collisionless gas⁴ or on the case of matter described by the massive field.⁵ The elements of the formalism were used to find the large-scale anisotropy of MBR by Abbott and Wise^{6,7} Efstathiou and Bond.⁸

In this paper we present the gauge-invariant calculation leading to formulas for the large-scale anisotropy of the MBR in the general (nonflat, multicomponent) cosmological model based on the Robertson-Walker metric. The formulas for scalar, vector, and tensor perturbations are given and those for scalar ones are analyzed in some specific models of the Universe. For the general flat models we decompose the large-scale MBR fluctuations into multipoles and calculate the amplitude of the *l*th

multipole as a function of the baryonic density and velocity perturbation fields.

In Sec. II we define perturbation quantities used later and give the equations of evolution of the background and the equations of motion for the perturbations. In Sec. III the MBR anisotropies are found. The results for some specific cases are discussed in Sec. IV. Section V presents the formalism of the multipole decomposition of the MBR pattern. Finally, Sec. VI contains concluding remarks.

II. DEFINITIONS AND EQUATIONS OF MOTION

We try to follow the notation of Bardeen's original paper.² The background Robertson-Walker metric is

$$ds^2 = g_{ij} dx^i dx^j = S^2(\tau) (-d\tau^2 + {}^3g_{\alpha\beta} dx^\alpha dx^\beta), \quad (1)$$

where $i, j, \dots = 0, 1, 2, 3$, $\alpha, \beta, \dots = 1, 2, 3$. The derivative with respect to the conformal time will be denoted by a dot, the covariant derivative with respect to g_{ij} by a semicolon, and with respect to ${}^3g_{\alpha\beta}$ by a vertical bar.

We assume that the Universe contains N ideal fluids with the unperturbed energy-momentum tensors:

$$T_{a0}^0 = -E_{a0}, \quad T_{a\beta}^\alpha = P_{a0} \delta_\beta^\alpha, \quad (2)$$

where $a = 1, \dots, N$ and E_{a0} and P_{a0} are the background energy density and pressure of the ideal fluid a . The more careful approach would be to describe the collisionless components by the means of a distribution function.⁴

We denote (c_{Sa} ² is the speed of sound)

$$w_a = \frac{P_{a0}}{E_{a0}}, \quad c_{Sa}^2 = \frac{dP_{a0}}{dE_{a0}}$$

and assume that the fluids are coupled only by gravity.

The equations governing the evolution of the background are

$$\left(\frac{\dot{S}}{S} \right)^2 = \frac{1}{3} S^2 \sum_{a=1}^N E_{a0} - K, \quad (3a)$$

$$\left[\frac{\dot{S}}{S} \right] = -\frac{1}{6} S^2 \sum_{a=1}^N (E_{a0} + 3P_{a0}), \quad (3b)$$

$$\frac{\dot{E}_{a0}}{E_{a0} + P_{a0}} = -3 \frac{\dot{S}}{S}, \quad (3c)$$

where $K = -1, 0, 1$ is the scalar of curvature in open, flat, and closed universes, respectively (units $c = 8\pi G = 1$). We can also incorporate the nonzero cosmological constant in the model: $E_{a0} = -P_{a0} = \Lambda$.

Perturbations can be classified according to their transformation properties under spatial coordinate transformations of the background spacetime as scalar, vector, and tensor perturbations. The time- and spatial-dependent parts of perturbation quantities can be separated thanks to the homogeneity and isotropy of the background. The spatial parts may be decomposed into the solutions of the generalized Helmholtz equations. For now, we will restrict our analysis to the case of the perturbations described by the single mode.

A. Scalar perturbations

Scalar harmonics $Q(x^\mu)$ are solutions of the equation

$$Q|^\alpha|_\alpha + k^2 Q = 0. \quad (4)$$

For a flat Robertson-Walker universe the Q 's are conveniently taken to be plane waves.

The vector and traceless tensor quantities are constructed by

$$Q_\alpha = -\frac{1}{k} Q|_\alpha, \quad (5a)$$

$$Q_{\alpha\beta} = \frac{1}{k^2} Q|_{\alpha\beta} + \frac{1}{3} g_{\alpha\beta} Q. \quad (5b)$$

The metric perturbations are written as

$$g_{00} = -S^2(1 + 2AQ), \quad (6a)$$

$$g_{0\alpha} = -S^2 B Q_\alpha, \quad (6b)$$

$$g_{\alpha\beta} = S^2[(1 + 2H_L Q)^3 g_{\alpha\beta} + 2H_T Q_{\alpha\beta}]. \quad (6c)$$

Let u_a^i be the four-velocity of the rest frame of fluid a relative to the coordinate frame (the rest frame is the frame in which the energy flux of fluid a vanishes). We assume that to zeroth order all u_a^i , $a = 1, \dots, N$ are the same. The first-order perturbations of the velocity of the fluid a are

$$u_a^0 = \frac{1}{S}(1 - AQ), \quad (7a)$$

$$u_a^\alpha = \frac{1}{S} v_a Q^\alpha. \quad (7b)$$

The perturbations in the energy-momentum tensor are

$$T_{a0}^0 = -E_{a0}(1 + \delta_a Q), \quad (8a)$$

$$T_{a0}^\alpha = -(E_{a0} + P_{a0})v_a Q^\alpha, \quad (8b)$$

$$T_{a\alpha}^0 = (E_{a0} + P_{a0})(v_a - B)Q_\alpha, \quad (8c)$$

$$T_{a\beta}^\alpha = P_{a0}[(1 + \pi_{La} Q)\delta_\beta^\alpha + \pi_{Ta} Q_\beta^\alpha], \quad (8d)$$

and the entropy perturbation is

$$\eta_a = \pi_{La} - \frac{c_{Sa}^2}{w_a} \delta_a. \quad (9)$$

The general gauge transformation of the wave number k is

$$\tilde{\tau} = \tau + T(\tau)Q(x^\mu), \quad (10a)$$

$$\tilde{x}^\alpha = x^\alpha + L(\tau)Q^\alpha(x^\mu). \quad (10b)$$

The gauge-invariant perturbation quantities are π_{Ta} , η_a , and

$$\Phi_A = A + \frac{1}{k} \dot{B} + \frac{1}{k} \frac{\dot{S}}{S} B - \frac{1}{k^2} \left[\ddot{H}_T + \frac{\dot{S}}{S} \dot{H}_T \right], \quad (11a)$$

$$\Phi_H = H_L + \frac{1}{3} H_T + \frac{1}{k} \frac{\dot{S}}{S} B - \frac{1}{k^2} \frac{\dot{S}}{S} \dot{H}_T, \quad (11b)$$

$$\epsilon_a = \delta_a + 3 \frac{1 + w_a}{k} \frac{\dot{S}}{S} (v_a - B), \quad (11c)$$

$$v_{Sa} = v_a - \frac{1}{k} \dot{H}_T. \quad (11d)$$

The equations of the evolution of perturbations derived from the Einstein equations are

$$\frac{2(k^2 - 3K)}{S^2} \Phi_H = \sum_{a=1}^N E_{a0} \epsilon_a, \quad (12a)$$

$$- \frac{k^2}{S^2} (\Phi_A + \Phi_H) = \sum_{a=1}^N P_{a0} \pi_{Ta}. \quad (12b)$$

And from the conservation equations $T_{aj;i}^i = 0$ we obtain ($a = 1, \dots, N$)

$$(E_{a0} \epsilon_a S^3)' + \frac{3(E_{a0} + P_{a0}) S^3}{k} \left\{ \left[\left[\frac{\dot{S}}{S} \right]^2 - \left[\frac{\dot{S}}{S} \right] \right] v_{Sa} + k \dot{\Phi}_H + \frac{1}{3} k^2 v_{Sa} \right\} - 3S^3 \frac{\dot{S}}{S} \left[(E_{a0} + P_{a0}) \Phi_A - \frac{2}{3} \left[1 - \frac{3K}{k^2} \right] P_{a0} \pi_{Ta} \right] = 0, \quad (13a)$$

$$\dot{v}_{Sa} + \frac{\dot{S}}{S} v_{Sa} = k \Phi_A + \frac{k}{1 + w_a} (c_{Sa}^2 \epsilon_a + w_a \eta_a) - \frac{2}{3} k \left[1 - \frac{3K}{k^2} \right] \frac{w_a}{1 + w_a} \pi_{Ta}. \quad (13b)$$

Equation (13a) for the case $K=0$ was first derived by Abbott and Wise.³

B. Vector perturbations

In this mode the quantities being scalars under spatial coordinate transformations remain unperturbed. The vector harmonics $Q^{(1)\alpha}$ are solutions of

$$Q^{(1)\alpha}{}_{|\beta} + k^2 Q^{(1)\alpha} = 0. \quad (14)$$

The tensor quantity is obtained by

$$Q^{(1)\alpha\beta} = -\frac{1}{2k} (Q^{(1)\alpha|\beta} + Q^{(1)\beta|\alpha}). \quad (15)$$

In analogy to the scalar perturbations we have

$$\begin{aligned} g_{0\alpha} &= -S^2 B^{(1)} Q_\alpha^{(1)}, \\ g_{\alpha\beta} &= S^2 ({}^3g_{\alpha\beta} + 2H_T^{(1)} Q_{\alpha\beta}^{(1)}), \\ u_a^\alpha &= \frac{1}{S} v_a^{(1)} Q^{(1)\alpha}, \\ T_{\alpha\alpha}^0 &= (E_{a0} + P_{a0})(v_a^{(1)} - B^{(1)}) Q_\alpha^{(1)}, \\ T_{a0}^\alpha &= -(E_{a0} + P_{a0}) v_a^{(1)} Q^{(1)\alpha}, \\ T_{\alpha\beta}^\alpha &= P_{a0} (\delta_\beta^\alpha + \pi_{Ta}^{(1)} Q_\beta^{(1)\alpha}). \end{aligned} \quad (16)$$

The allowed gauge transformation is $\bar{x}^\alpha = x^\alpha + L^{(1)}(\tau) Q^\alpha(x^\mu)$ and the gauge-invariant quantities are $\pi_{Ta}^{(1)}$ and

$$\Psi = B^{(1)} - \frac{1}{k} \dot{H}_T^{(1)}, \quad (17a)$$

$$v_{Sa}^{(1)} = v_a^{(1)} - \frac{1}{k} \dot{H}_T^{(1)} \quad \text{or} \quad v_{Ca} = v_{Sa}^{(1)} - \Psi. \quad (17b)$$

The equations of motion are

$$\frac{k^2 - 2K}{2S^2} \Psi = \sum_{a=1}^N (E_{a0} + P_{a0}) v_{Ca}, \quad (18a)$$

$$\dot{v}_{Ca} + \frac{\dot{S}}{S} (1 - 3c_{Sa}^2) v_{Ca} = -\frac{k}{2} \frac{w_a}{1 + w_a} \pi_{Ta}^{(1)}. \quad (18b)$$

C. Tensor perturbations

Now only tensor quantities are perturbed and the tensor harmonics are solutions of

$$Q^{(2)\alpha\beta}{}_{|\gamma} + k^2 Q^{(2)\alpha\beta} = 0. \quad (19)$$

We have

$$\begin{aligned} g_{\alpha\beta} &= S^2 ({}^3g_{\alpha\beta} + 2H_T^{(2)} Q_{\alpha\beta}^{(2)}), \\ T_{\alpha\beta}^\alpha &= P_{a0} (\delta_\beta^\alpha + \pi_{Ta}^{(2)} Q_\beta^{(2)\alpha}). \end{aligned} \quad (20)$$

The quantities $H_T^{(2)}$ and $\pi_{Ta}^{(2)}$ are automatically gauge invariant. The equation of motion is

$$\ddot{H}_T^{(2)} + 2\frac{\dot{S}}{S} \dot{H}_T^{(2)} + (k^2 + 2K) H_T^{(2)} = S^2 \sum_{a=1}^N P_{a0} \pi_{Ta}^{(2)}. \quad (21)$$

III. THE TEMPERATURE OF MBR IN THE PERTURBED UNIVERSE

Our aim here is to find the MBR pattern $T_R(\theta, \phi)$ (subscript R denotes reception and subscript E denotes emission) in a cosmological model based on a Robertson-Walker metric if the solutions of the perturbation equations for density and velocity fields are known. We assume that the decoupling is instantaneous and that we are interested in large angular scales only.

The temperature of the MBR coming from a given direction is determined by the red-shift that photons acquired from emission until reception. The density and velocity perturbations of the fluids filling the Universe cause the perturbations in the metric that in turn influence the motion of light, resulting in red-shift differences of photons coming from different directions. (The interaction of light and matter can also have the direct form of scattering, if the intergalactic medium was reionized in the epoch of galaxy or star formation.)

In order to obtain the pattern of the MBR in a perturbed universe we should in principle integrate the Boltzmann equation for a generalized, gauge-invariant distribution function of photons through the decoupling phase and further, until today. However, details of the decoupling are important only for small angular scales, and in this case the use of gauge-invariant quantities is not necessary. For large scales the opposite is true. By large angular scales we mean scales larger than those subtended today by the light rays emitted at decoupling from two points at separation equal to the size of the observable Universe at decoupling (i.e., twice the distance to the horizon). For the matter-dominated universe this angular scale is⁹

$$\theta_H \simeq 3^\circ \left(\frac{1300}{z} \Omega_R \right)^{1/2},$$

where z is the red-shift of the decoupling and Ω_R is the density parameter today.

Therefore a suitable approximation is to follow the movement of the single photon emitted from the last scattering hypersurface using the gauge-invariant quantities.

The light moves along the null-like geodesics $x^i(\lambda)$ where λ is the affine parameter. The null vector tangent to the geodesic is

$$k^i = \frac{dx^i}{d\lambda}, \quad k^i = (v, P^\alpha), \quad k_i k^i = 0. \quad (22)$$

We write

$$v = {}^0v(1 - M), \quad P^\alpha = {}^0P^\alpha + {}^1P^\alpha, \quad (23)$$

where 0vM and ${}^1P^\alpha$ are the first-order corrections to components of vector k^i . The equation of motion is

$$\frac{Dk^i}{d\lambda} = \frac{dk^i}{d\lambda} + \Gamma_{kl}^i k^k k^l = 0. \quad (24)$$

The temperature of the MBR now observed is

$$\frac{T_R}{T_E} = \frac{1}{1+z} = \frac{(k^i u_{bi})_R}{(k^i u_{bi})_E}, \quad (25)$$

where u_{bi} is the four-vector of velocity of observer at rest relative to baryonic fluid (observers are made of baryons). We take no account of local, gravity-induced motions, e.g., of the Galaxy, that result in additional dipole term.

It is convenient to introduce a new parameter $s(\lambda)$ (the derivative with respect to s will be denoted by a prime) such that

$$\frac{d\lambda}{ds} = \frac{S^2}{S_R^2}, \quad \frac{d}{ds} = \frac{S^2}{S_R^2} \frac{d}{d\lambda} = \frac{S^2}{S_R^2} \left[v \frac{\partial}{\partial \tau} + P^\alpha \frac{\partial}{\partial x^\alpha} \right]. \quad (26)$$

From (22) and (24) we obtain, in the zeroth order

$${}^0v = \frac{S_R^2}{S^2}, \quad {}^0v^2 = {}^0P^\alpha {}^0P_\alpha. \quad (27)$$

Defining ${}^0P^\alpha = -{}^0vR^\alpha$ where R^α is the spatial unit vector ($R^\alpha R_\alpha = 1$) in the direction of observation and using the normalization ${}^0v_R = 1$ we obtain the solutions for the lightlike geodesics to zeroth order:

$$\tau = \tau_E + s, \quad (28a)$$

$$x^\alpha = R^\alpha (\tau_R - \tau_E - s), \quad (28b)$$

$$\frac{dR^\alpha}{ds} = {}^3\Gamma_{\beta\gamma}^\alpha R^\beta R^\gamma, \quad (28c)$$

where

$$\frac{d}{ds} = \frac{\partial}{\partial \tau} - R^\alpha \frac{\partial}{\partial x^\alpha}, \quad R^\alpha = -\frac{dx^\alpha}{ds}.$$

The coordinates of the emission event are $s=0$, $\tau=\tau_E$, $x_E^\alpha = R_E^\alpha (\tau_R - \tau_E)$ and of the reception event are $s=\tau_R - \tau_E$, $\tau=\tau_R$, $x_R^\alpha = 0$.

The equations to first order look differently for scalar, vector, and tensor perturbations.

A. Scalar perturbations

Using (22) and (24) to first order we obtain

$$M' = \dot{A}Q + 2kAQ_{\alpha}R^\alpha + \frac{k}{3}BQ + \dot{H}_L Q + (\dot{H}_T - kB)Q_{\alpha\beta}R^\alpha R^\beta \quad (29)$$

and from (25)

$$\frac{T_E}{T_R} = \frac{S_R}{S_E} \left\{ 1 + \int_E^R \left[\left[\dot{H}_L + \frac{k}{3}v_b \right] Q - (\dot{v}_b - kA - \dot{B})Q_{\alpha}R^\alpha - (kv_b - \dot{H}_T)Q_{\alpha\beta}R^\alpha R^\beta \right] ds \right\}, \quad (30)$$

where the integral is along the zeroth-order lightlike geodesic.

The above formula is gauge invariant because S transforms under (10a) as $S(\bar{\tau}) = S(\tau)[1 + (\dot{S}/S)TQ]$ and then we should be able to express it using only the gauge-invariant variables.

To show that the details of the recombination are not important in most of the models we derive a simple approximation for them. In our model of decoupling the emission of radiation occurs on the hypersurface of the constant density of free electrons that couple to photons by Thomson scattering. This density is a function of the local temperature and density of baryons and thus for general perturbations the hypersurface of emission is neither the hypersurface of constant temperature nor that of constant baryon density. In the presence of perturbations the emission in a given point in space occurs in the moment $\tau_{E_0} + \Delta\tau$ where $\Delta\tau$ is a function of perturbations. Denoting the density of free electrons at the emission by n_{eE} and the moment of emission in the zeroth order by the subscript E_0 we have on the hypersurface of last scattering:

$$n_{eE} = \text{const} = n_e(\tau_{E_0} + \Delta\tau) = n_{e0}(\tau_{E_0} + \Delta\tau)(1 + \delta_e Q), \quad (31)$$

so

$$\Delta\tau = -\frac{n_{e0}}{\dot{n}_{e0}} \delta_e Q, \quad (32)$$

where δ_e is a perturbation of density of electrons and in general

$$n_e = f(E_b)g(T). \quad (33)$$

Using this functional form we can express $\Delta\tau$ as a function of the perturbations of baryons and photons ($E_\gamma = \sigma T^4$). The result is

$$\frac{\dot{S}}{S} \Delta\tau = \frac{1}{3+D} \delta_b Q + \frac{D}{4(3+D)} \delta_\gamma Q, \quad (34)$$

where

$$D = \left[\frac{f(dg/dT)T}{(df/dE_b)gE_b} \right]_E. \quad (35)$$

In the simplest model of decoupling we can use the Saha formula for the fractional ionization¹⁰ and we obtain $D \simeq \frac{3}{2} + B/k_B T_E$ where $B = 13.6$ eV and for $T_E = 3500$ K, $D \simeq 47$.

At the moment of emission

$$S_E = S_{E_0} \left[1 + \frac{\dot{S}}{S} \Delta\tau \right]_E. \quad (36)$$

Now we can rewrite (30) in gauge-invariant quantities. We define

$$\left[\frac{\delta T}{T} \right]_R = \frac{T_R - T_{R_0}}{T_{R_0}}, \quad T_{R_0} = \frac{S_{E_0} T_E}{S_{R_0}} \quad (37)$$

to obtain

$$\left[\frac{\delta T}{T} \right]_R = \left[\frac{\delta T}{T} \right]_E - \left[\left[\Phi_H - \frac{1}{k} \frac{\dot{S}}{S} v_{Sb} \right] Q \right]_E^R - \int_E^R \left[\left[\frac{\dot{v}_{Sb}}{k} + \Phi_H - \Phi_A \right] Q_{|\alpha} R^\alpha - \frac{v_{Sb}}{k} Q_{|\alpha\beta} R^\alpha R^\beta \right] ds \quad (38)$$

where

$$\left(\frac{\delta T}{T}\right)_E \equiv \left[\left[\frac{1}{3+D}\epsilon_b + \frac{D}{4(3+D)}\epsilon_\gamma \right] Q - \frac{1}{k} \frac{\dot{S}}{S} \frac{D}{3+D} (v_{S\gamma} - v_{Sb}) Q \right]_E \quad (39)$$

and we used $w_b=0$, $w_\gamma=\frac{1}{3}$ and the fact that any part of T_R independent of the direction of observation can be incorporated into the definition of T_{R_0} . This is the explicitly gauge-invariant form of the temperature fluctuations in the general case.

In the most popular (and suggested by many theories of the very early Universe) case of adiabatic perturbations we have $\delta_\gamma = \frac{4}{3}\delta_b$ and the initial fluctuations of temperature reduce to $\frac{1}{3}\epsilon_b Q$ at the emission. We will analyze further this case only, and, as we will see this term can be dropped. However, we should emphasize that in nonadiabatic models the initial fluctuations given by the formula (39) can be important—for example, the term $\sim \epsilon_\gamma$ is responsible for the increase of the large-scale angular fluctuations in the model of isocurvature axion perturbations analyzed by Efstathiou and Bond.⁸

We can put $c_{Sb}^2=0$ if we are interested in scales larger than the baryon Jeans mass. This is the case because for $\lambda \geq (ct)_E$ this Jeans mass is much less than the mass in a sphere of diameter $\sim \lambda$. The equations of motion (13) for the baryonic perturbations are then

$$\dot{\epsilon}_b + 3 \left[\left(\frac{\dot{S}}{S} \right)^2 - \left[\frac{\dot{S}}{S} \right] \right] \frac{v_{Sb}}{k} + 3\dot{\Phi}_H - 3\frac{\dot{S}}{S}\Phi_A + kv_{Sb} = 0, \quad (40a)$$

$$\dot{v}_{Sb} + \frac{\dot{S}}{S}v_{Sb} = k\Phi_A. \quad (40b)$$

We can use these equations to rewrite (38) in the form

$$\left(\frac{\delta T}{T}\right)_R = \left(\frac{1}{3}\epsilon_b Q\right)_E + \int_E^R \left[\frac{1}{3}(\dot{\epsilon}_b + kv_{Sb})Q + \frac{v_{Sb}}{k} Q_{|\alpha\beta} R^\alpha R^\beta \right] ds. \quad (41)$$

$$(E_{a0}\epsilon_a S^3)' + \frac{3(E_{a0}+P_{a0})S^3}{k} \left\{ \left[\left(\frac{\dot{S}}{S} \right)^2 - \left[\frac{\dot{S}}{S} \right] \right] v_{Sa} + k\dot{\Phi}_H + \frac{1}{3}k^2 v_{Sa} \right\} + 3S^3 \frac{\dot{S}}{S} (E_{a0}+P_{a0})\Phi_H = 0, \quad (44a)$$

$$\dot{v}_{Sa} + \frac{\dot{S}}{S}v_{Sa} = -k\Phi_H + \frac{kc_{Sa}^2}{1+w_a}\epsilon_a, \quad (44b)$$

$$\Phi_H = \frac{S^2}{2(k^2-3K)} \sum_{a=1}^N E_{a0}\epsilon_a. \quad (44c)$$

The next simplification is obtained if we are interested in the models dominated by nonrelativistic components with $P_{a0}=0$ [e.g., cold-dark-matter (CDM) models] since decoupling. If we omit the radiation and relativistic neutrinos in the equations of motion, we have ($a=1, \dots, N$)

The first term represents the influence of the density perturbations at the emission and can be dropped because in our case it is much smaller than the integral term (see Sec. IV). The integral term describes how the motion of light is influenced by the geometry perturbations generated by all density and anisotropic pressure perturbations present in the model [see (12)] and described by the baryonic quantities [$(k/S)v_{Sb}$ is the magnitude of shear of the baryonic velocity field]. Some specific cases of (41) are described in the next section.

B. Vector perturbations

An analysis similar to the one performed before gives us the result

$$\left(\frac{\delta T}{T}\right)_R^{(1)} = \int_E^R (\dot{v}_{Cb} Q_\alpha^{(1)} R^\alpha + kv_{Sb}^{(1)} Q_{\alpha\beta}^{(1)} R^\alpha R^\beta) ds. \quad (42)$$

C. Tensor perturbations

We find

$$\left(\frac{\delta T}{T}\right)_R^{(2)} = - \int_E^R \dot{H}_T^{(2)} Q_{\alpha\beta}^{(2)} R^\alpha R^\beta ds. \quad (43)$$

The vector and tensor cases will not be analyzed further.

IV. SPECIFIC CASES OF THE SCALAR PERTURBATIONS

Although the general formula describing the MBR pattern in the presence of scalar perturbations (41) is very simple the troubles arise with its use because of the complicated form of the equations of motion in the general case (12), (13). However we can simplify them in some specific models.

First of all we expect that the anisotropic stress and entropy perturbations in any component of the Universe operate only at very early stages of the evolution. Thus we can omit them in our analysis. Now the equations of motion are ($a=1, \dots, N$)

$$\dot{\epsilon}_a \left[1 + \frac{3V_a}{2(k^2 - 3K)S} \right] + kv_{Sa} \left[1 - \frac{3K}{k^2} + \frac{3}{2k^2S} \sum_{c=1}^N V_c \right] + \frac{3}{2(k^2 - 3K)S} \sum_{\substack{c=1 \\ c \neq a}}^N V_c \dot{\epsilon}_c = 0, \quad (45a)$$

$$\dot{v}_{Sa} + \frac{\dot{S}}{S} v_{Sa} = - \frac{1}{2k} \left[1 - \frac{3K}{k^2} \right] S \sum_{c=1}^N V_c \epsilon_c. \quad (45b)$$

$V_a = E_{a0} S^3$ are constants closely related to the present density parameters Ω_{Ra} (H_R is the Hubble constant measured today):

$$\Omega_{Ra} = \frac{E_{a0R}}{3H_R^2} = \frac{V_a}{\sum_{c=1}^N V_c - 3KS_R}. \quad (46)$$

In the case $K=0$ suggested by the inflationary scenarios we have $\Omega_a = \text{const} = \Omega_{Ra}$ and we denote the sum of all V_c by V :

$$\dot{\epsilon}_a \left[1 + \frac{3\Omega_a V}{2k^2 S} \right] + kv_{Sa} \left[1 + \frac{3V}{2k^2 S} \right] + \frac{3V}{2k^2 S} \sum_{\substack{c=1 \\ c \neq a}}^N \Omega_c \dot{\epsilon}_c = 0, \quad (47a)$$

$$\dot{v}_{Sa} + \frac{\dot{S}}{S} v_{Sa} = - \frac{V}{2kS} \sum_{c=1}^N \Omega_c \epsilon_c. \quad (47b)$$

Greatly simplified equations of motion are obtained from (45a) and (45b) if $N=1$ or all $\dot{\epsilon}_c$ $c=1, \dots, N$ are equal. For baryons

$$\dot{\epsilon}_b + \left[1 - \frac{3K}{k^2} \right] kv_{Sb} = 0. \quad (48a)$$

This is the case in most of the dark-matter scenarios—on scales larger than the Jeans mass, the baryons sink in the potential wells of the dark component. In a few expansion times all the ϵ_c 's become equal. Then the second equation of motion is

$$\dot{v}_{Sb} + \frac{\dot{S}}{S} v_{Sb} = - \frac{V}{2k} \left[1 - \frac{3K}{k^2} \right] S \epsilon_b. \quad (48b)$$

Now we can write formula (41) under the assumption of equal ϵ_c 's:

$$\left[\frac{\delta T}{T} \right]_R = \left(\frac{1}{3} \epsilon_b Q \right)_E - \frac{1}{k^2 - 3K} \times \int_E^R \dot{\epsilon}_b (KQ + Q_{|\alpha\beta} R^\alpha R^\beta) ds. \quad (49)$$

In the flat universe $K=0$, the covariant derivatives be-

come normal and Q is taken to be a plane wave with the wave vector k^α : $Q = \exp(ik_\alpha x^\alpha)$. From (49) we obtain

$$\left[\frac{\delta T}{T} \right]_R = \left(\frac{1}{3} \epsilon_b Q \right)_E + \frac{1}{k^2} \int_E^R \dot{\epsilon}_b (k_\alpha R^\alpha)^2 Q ds. \quad (50)$$

In this case the equation for density perturbations has the solution

$$\epsilon_b = A\tau^2 + B\tau^{-3}. \quad (51)$$

For the growing mode $\epsilon_b = \epsilon_{bE} (\tau/\tau_E)^2$ the integral in (50) can be integrated by parts:

$$\left[\frac{\delta T}{T} \right]_R = \frac{1}{3} \epsilon_{bE} Q_E + \frac{2\epsilon_{bE}}{k^2 \tau_E^2} [\tau_R (R^\alpha Q_{,\alpha})_R - \tau_E (R^\alpha Q_{,\alpha})_E + Q_R - Q_E]. \quad (52)$$

The comoving coordinate distance to the horizon at τ_E is equal τ_E . Then the criterion for the comoving scale k to be larger than the horizon at τ_E is $k\tau_E \ll 1$. In this regime we can drop the first term because it is much smaller than the integral term. (In fact when the first term is important, on scales $k\tau_E \geq 1$, it does not look so simple.) We can incorporate the term $\sim Q_R$ into the definition of T_{R_0} . This gives

$$\left[\frac{\delta T}{T} \right]_R = \frac{2\epsilon_{bE}}{k^2 \tau_E^2} [\tau_R (R^\alpha Q_{,\alpha})_R - \tau_E (R^\alpha Q_{,\alpha})_E - Q_E]. \quad (53)$$

This is the well-known result of Sachs and Wolfe.

V. MULTIPOLE DECOMPOSITION

The anisotropies of the MBR calculated in the cosmological models can be compared with the limits from observations to restrict the parameters of these models. For the large-scale anisotropies the convenient quantities are the amplitudes of the multipoles in the decomposition of the function $(\delta T/T)_R(\theta, \phi)$. In further analysis we will restrict ourselves to the case of the flat universe.

We usually assume that the field of the density perturbations of all components consists of the sum of plane waves with random wave vectors and phases. The amplitude of the wave with the wave vector k^α is assumed to be the function of the modulus of the wave vector only, usually the power law at the prescribed moment.

The power-law behavior is understandable on scales smaller than the horizon but its continuation on scales larger than the horizon is not obvious. This is because there is no unique choice of gauge-invariant quantity for density perturbations. However, we argue that we can assume the continuation of the functional behavior for the quantity ϵ used here (this is ϵ_m in Bardeen's notation), because it is uniform on scales smaller and larger than the horizon (Bardeen's ϵ_g is not) and because it directly couples to the potential Φ_H [see (12a)].

Some spectra that are power laws at the time of horizon

crossing at a given scale deviate later from a pure power law as a result of the dynamical evolution of perturbations in a radiation-dominated universe after horizon crossing (e.g., CDM spectra). Therefore we will assume the general functional dependence of the amplitude of density perturbations at a given scale at the emission. The single plane wave is then replaced by the sum:

$$\sum_{\mathbf{k} \neq 0} \epsilon_a(k) e^{i\mathbf{k} \cdot \mathbf{x}}. \quad (54)$$

Our initial conditions are the density perturbations at decoupling. We assume that the equations of motion were solved and we can write the appropriate form of $\epsilon_b(k)$ and $v_{Sb}(k)$ for all $\tau_E \leq \tau \leq \tau_R$. Now the formula (41) for the flat universe can be written in the form

$$\left[\frac{\delta T}{T} \right]_R = \sum_{\mathbf{k} \neq 0} \int_E^R \left[\frac{1}{3} \dot{\epsilon}_b(k) + kv_{Sb}(k) \left(\frac{1}{3} - \xi_{\mathbf{k}}^2 \right) \right] e^{i\mathbf{k} \cdot \mathbf{x}} ds, \quad (55)$$

where $\xi_{\mathbf{k}}$ is the cosine of the angle between the wave vector \mathbf{k} and the direction of observation \mathbf{R} .

Our aim is to find the coefficients of the decomposition

$$\left[\frac{\delta T}{T} \right]_R = \sum_{l,m} a_l^m Y_l^m(\Omega), \quad a_l^m = \int Y_l^{*m} \left[\frac{\delta T}{T} \right]_R d\Omega, \quad (56)$$

where $\Omega = (\theta, \phi)$. We use the following mathematical formulas:

$$e^{i\mathbf{k} \cdot \mathbf{x}} = \sum_{l=0}^{\infty} i^l (2l+1) j_l(kx) P_l(\xi_{\mathbf{k}}), \quad (57)$$

$$\xi_{\mathbf{k}}^2 P_l(\xi_{\mathbf{k}}) = \frac{1}{2l+1} [b_l P_{l+2}(\xi_{\mathbf{k}}) + c_l P_l(\xi_{\mathbf{k}}) + d_l P_{l-2}(\xi_{\mathbf{k}})], \quad (58)$$

$$P_l(\xi_{\mathbf{k}}) = \frac{4\pi}{2l+1} \sum_{m=-l}^l Y_l^{*m}(\Omega_{\mathbf{k}}) Y_l^m(\Omega), \quad (59)$$

where

$$b_l = \frac{(l+1)(l+2)}{2l+3}, \quad c_l = \frac{(l+1)^2}{2l+3} + \frac{l^2}{2l-1}, \quad (60)$$

$$d_l = \frac{l(l-1)}{2l-1},$$

and $\Omega_{\mathbf{k}} = (\theta_{\mathbf{k}}, \phi_{\mathbf{k}})$ are the angular coordinates of the vector \mathbf{k} .

In (56) we decompose the plane waves into a series of Legendre polynomials, which in turn are represented by the spherical harmonics. We then integrate over $d\Omega$. The result is

$$(a_l)^2 = \frac{16\pi}{(2l+1)^2 \tau_E^4} \int_0^{k_{\max}} k^{-2} |\epsilon_{bE}(k)|^2 \{ (2l+1) j_l(k s_R) + k \tau_E [l j_{l-1}(k s_R) - (l+1) j_{l+1}(k s_R)] - \delta_{l1} k \tau_R \}^2 dk \quad (67)$$

and the integral term $\sim k \tau_E$ is dominated by the others.

The results of observations on medium angular scales are represented in the form of the angular correlation function of fluctuations,

$$a_l^m = 4\pi i^l \sum_{\mathbf{k} \neq 0} A_l(k) Y_l^{*m}(\Omega_{\mathbf{k}}), \quad (61)$$

where

$$A_l(k) = \int_E^R \left\{ \frac{1}{3} [\dot{\epsilon}_b(k) + kv_{Sb}(k)] j_l(kx) + \frac{kv_{Sb}(k)}{2l+1} [d_l j_{l-2}(kx) - c_l j_l(kx) + b_l j_{l+2}(kx)] \right\} ds. \quad (62)$$

This result can be generalized on the case of nonadiabatic models by simply adding to a_l^m the contribution from the decomposition of initial fluctuations (39).

The quantity used to compare with observations is

$$(a_l)^2 = \langle |a_l^m|^2 \rangle \quad (63)$$

(in fact, for a given l , all a_l^m are statistically independent with the same expectation values). We use the assumption that the phases of $\epsilon_b(k)$ and $v_{Sb}(k)$ are random and change the sum over \mathbf{k} into the integral

$$\sum_{\mathbf{k} \neq 0} \rightarrow \int k^2 dk \frac{d\Omega_{\mathbf{k}}}{4\pi} \quad (64)$$

to obtain

$$(a_l)^2 = 4\pi \int_0^{k_{\max}} k^2 |A_l(k)|^2 dk. \quad (65)$$

The cutoff k_{\max} reflects the limits of our simplified model. They can be related to the horizon size and the scale of the Silk damping of baryonic perturbations, but another requirement for the small scales is not to be smaller than the scales that are nonlinear today. Anyway we should expect the results to be practically cutoff independent for small l (if k_{\max} is large enough) because small scale perturbations averaged over large scales give the result ~ 0 .

The formula (62) is simplified in the case of the universe dominated by the nonrelativistic matter such that all ϵ_c , $c = 1, \dots, N$ are equal (or $N = 1$) because then $\dot{\epsilon}_b + kv_{Sb} = 0$ and

$$A_l(k) = -\frac{1}{2l+1} \int_E^R \dot{\epsilon}_b(k) [d_l j_{l-2}(kx) - c_l j_l(kx) + b_l j_{l+2}(kx)] ds. \quad (66)$$

For the growing mode of the density perturbations $\epsilon_b = \epsilon_{bE}(\tau/\tau_E)^2$ the integral can be calculated⁶ (or equivalently we can integrate the Sachs-Wolfe formula). The result is ($s_R = \tau_R - \tau_E$)

$$W(\theta) = \frac{1}{8\pi^2} \int \int d\Omega_1 d\Omega_2 \left[\frac{\delta T}{T}(\Omega_1) \right]_R \left[\frac{\delta T}{T}(\Omega_2) \right]_R \times \delta_D(\cos\theta_{12} - \cos\theta), \quad (68)$$

which can be expressed using the multipole coefficients as

$$W(\theta) = \frac{1}{4\pi} \sum_{l=1}^{\infty} (2l+1) a_l^2 P_l(\cos\theta). \quad (69)$$

VI. CONCLUSIONS

The gauge-invariant approach² solves many ambiguities about the behavior of the large-scale perturbations. Its application to the large-scale fluctuations of the MBR temperature is presented in this paper. The explicitly gauge-invariant formula for these fluctuations (38) is obtained for the wide class of cosmological models based on the Robertson-Walker metrics. The only assumptions used in its derivation are that all constituents of the Universe can be described as perfect fluids and that the decoupling of matter and radiation occurs instantaneously on the hypersurface of the last scattering. However, relaxing these assumptions will not substantially change the results.

For the most popular model of the flat universe we obtain the multipole coefficients of the decomposition of the MBR pattern into spherical harmonics (65). They can be explicitly calculated if the equations of motion for baryonic perturbations were solved.

The multipole coefficients and the angular correlation function of fluctuations found in a model can be com-

pared with the results of observations to provide valuable constraints on parameters of the model. The higher multipole moments have not been observed yet and we know only the upper limits of the two first moments.

Unfortunately the dipole moment ($l=1$) is influenced by the nonlinear, gravity induced motions of our Galaxy that cannot be reliably subtracted. However, the observations^{11,12} indicate that the intrinsic dipole moment of MBR can be $a_1 \leq 10^{-4}$ (Ref. 12). The recent observational limits on the quadrupole ($l=2$) moment complied by Efstathiou and Bond⁸ give the value $a_2 \leq 10^{-4}$.

Any comparison of $W(\theta)$ calculated in the model with the observations requires additional information about the characteristic of the antenna used (see Ref. 8 for examples). The dipole momentum is usually excluded.

The reheating of intergalactic medium during star or galaxy formation could influence the pattern of the MBR and this effect will be addressed in another paper.

ACKNOWLEDGMENTS

The author would like to thank Roman Juszkiewicz and Michael Turner for encouragement and helpful comments. This work was supported in part by NASA and the Department of Energy at Fermilab.

*Permanent address: Copernicus Astronomical Center, Bartyc-ka 18, 00-716 Warsaw, Poland.

¹P. J. E. Peebles and J. T. Yu, *Astrophys. J.* **162**, 815 (1970); M. L. Wilson and J. Silk, *ibid.* **243**, 14 (1981).

²J. M. Bardeen, *Phys. Rev. D* **22**, 1882 (1980).

³L. F. Abbott and M. B. Wise, *Nucl. Phys.* **B237**, 226 (1984).

⁴M. Kasai and K. Tomita, Hiroshima University report, 1985 (unpublished).

⁵R. Brandenberger, R. Kahn, and W. Press, *Phys. Rev. D* **28**, 1809 (1983).

⁶L. F. Abbott and M. B. Wise, *Phys. Lett.* **135B**, 279 (1984).

⁷L. F. Abbott and M. B. Wise, *Astrophys. J.* **282**, L47 (1984).

⁸G. Efstathiou and J. R. Bond, *Mon. Not. R. Astron. Soc.* **218**, 103 (1986).

⁹S. Weinberg, *Gravitation and Cosmology* (Wiley, New York, 1972).

¹⁰P. J. E. Peebles, *Physical Cosmology* (Princeton University Press, Princeton, N.J., 1971).

¹¹L. Hart and R. D. Davies, *Nature (London)* **275**, 514 (1982); G. A. Tammann and A. Sandage, *Astrophys. J.* **294**, 81 (1985).

¹²G. de Vaucouleurs and W. L. Peters, *Astrophys. J.* **287**, 1 (1984).