

Transverse-flow effects in dilepton emission

K. Kajantie,* M. Kataja,† L. McLerran,† and P. V. Ruuskanen†

Research Institute for Theoretical Physics, University of Helsinki, Helsinki, Finland

(Received 3 February 1986)

Dilepton emission from expanding QCD matter formed in ultrarelativistic nuclear collisions is computed. The energy density and the velocity field of the expanding matter are computed numerically with a (1+3)-dimensional numerical code which assumes cylindrical invariance in the transverse and boost invariance in the longitudinal direction. Effects of transverse flow are studied and shown to be important in the region of pair mass $M \lesssim 1.2$ GeV. The existence of transverse flow results in the persistent appearance of the ρ peak in the M distribution at fixed transverse mass M_T for large M_T . The initial temperature T_i and the thermalization time τ_i of the hydrodynamical flow are related to the associated pion multiplicity dN/dy and the dilepton emission rate at $M > 1.5$ GeV.

I. INTRODUCTION

A large amount of work has been devoted to studying possible signals for the production of QCD matter in its different phases as it might be formed in ultrarelativistic nucleus-nucleus collisions.¹ The physics which can be studied in such collisions includes the determination of state of hadronic matter, as well as properties of exotic new forms of matter such as quark-gluon plasma.

Because of the complexity of the problem, one has been forced to simplify the space-time history of the expanding matter. At best² one has coupled the dilution due to longitudinal boost-invariant expansion³ with transverse expansion using a one-phase equation of state.

The purpose of this paper is to consider the dilepton signal⁴⁻⁹ by numerically computing¹⁰⁻¹² the matter flow assuming boost-invariant longitudinal flow and cylindrical symmetry for the flow in the transverse direction and, secondly, assuming an equation of state containing a first-order phase transition. The principal extension of the treatment² of the transverse expansion is that we use an equation of state which allows one to continue past the quark-gluon plasma phase, through a phase transition, and finally through a hadronic gas phase. This treatment involves using a hydrodynamic code which properly treats shock discontinuities and is discussed in Ref. 10. In this paper we shall exclusively consider central collisions of heavy nuclei such as uranium.

What happens in the central region (at rest in the center of mass) of a central (zero impact parameter) U + U collision is essentially as follows. Very early in the collision (at times less than $1/M$) a single hard quark-antiquark collision may produce a massive dilepton pair through the Drell-Yan process. After this, collisions between the partons start thermalizing the system. Dileptons may also be emitted during this pre-equilibrium stage.¹³⁻¹⁶ During the pre-equilibrium stage the total comoving entropy $S(\eta)$ of the system increases rapidly. From some proper time τ_i and temperature T_i on S is essentially constant and the system is regarded as thermalized. We shall, in the following, unless otherwise stated, consider only the

case when T_i is so large that the system initially is in the quark-plasma phase. It may happen that this is not the case for an average experimental event.

Because of quantum fluctuations the value of $S(\eta)$ may vary from event to event, but its value may be experimentally measured by measuring the associated pion multiplicity dN/dy and using the relationship⁹

$$S(\eta) = c \frac{dN}{dy}(\eta=y) = \pi R_A^2 4a T_i^3 \tau_i, \quad (1.1)$$

where $c=3.6$ and $a=4.6$ (for 2.5 flavors). Note that this only gives the combination $T_i^3 \tau_i$. The value of τ_i is, however, not so well determined for a variety of reasons such as fluctuations in the initial conditions. In view of these uncertainties, we will consider τ_i in the range of 0.1 to 1.5 fm. From the point of view of the hydrodynamic flow, this variation in the value of τ_i is not important: increasing τ_i from 0.1 to 1.5 fm gives, apart from small initial condition ($v_r=0$) effects, exactly the same flow, if T_i is simultaneously decreased by a factor $(1.5/0.1)^{1/3}$ (Sec. II). The change from the point of view of signals is, however, significant, because the high-temperature period between 0.1 and 1.5 fm in the history of the system would be excluded (Sec. III).

Assume T_i is so large that the system initially is in the quark-plasma phase. The system then will expand through quark, mixed, and hadron phases.^{10,11} We also assume that the nucleation of the first-order transition is fast and do not discuss scenarios with phase separation¹⁷ or deep supercooling.¹⁸ What one finds (Sec. II) is that the behavior of the quark phase is little affected by the transverse flow. The longitudinal cooling is so fast that it cools the system out of the quark-plasma phase before the transverse rarefaction wave reaches the center, i.e., before $\tau = R_A/c_s \simeq 12$ fm. Only for very large S_i does the quark phase experience transverse motion. However, one finds that when the system is in the mixed or even more so in the hadron phase, it is in very rapid transverse motion. For instance, in the numerical examples of Sec. II the temperature $T=50$ MeV (this is the lowest temperature we consider for the hadron gas) is reached at a proper

time $\tau=35\text{--}65$ fm, and the system is then essentially a disk of radius $15\text{--}50$ fm in which the radial velocity varies from 0 at center to almost 1 at the sides.

How is this flow pattern experimentally diagnosed by dilepton emission? One finds (Sec. III) that small- M ($M \lesssim 1.2$ GeV) pairs are dominantly emitted while the system is in the mixed and hadron phases. As the system is then in rapid transverse motion, the system imparts the pairs in this mass range a large transverse momentum. We find that the ρ peak will persist until largest transverse masses M_T for all reasonable values of the initial temperature, formation time, and hadron multiplicity. Since the neglect of the transverse motion predicts the disappearance of the ρ peak,¹⁹ the observation of the ρ peak will permit one to experimentally identify the transverse flow. A different method of observing the flow is by comparing transverse-momentum distributions of pions to those in pp at equivalent energy.¹⁰

Similarly, the larger- M ($M \gtrsim 2$ GeV) pairs are dominantly emitted while the system is in the quark phase. As this is the initial stage, one can now attempt to experimentally measure the values of T_i and τ_i separately. First, a fixed value of $T_i^3 \tau_i$ is chosen by considering only events with some fixed dN/dy of pions. Second, for these events one measures the rate $dN/dM dy$ for pairs in the mass range $2 < M < 3$, say. The value of this rate depends on τ_i separately and thus one can measure T_i and τ_i separately. Note that this value of τ_i (and the associated T_i) is the average value of τ_i for the events with the given dN/dy ; it is not the τ_i for each event separately. The average value of τ_i also contains an averaging over the transverse coordinates.

The upper limit $M \lesssim 3$ GeV in the determination of τ_i follows from the fact that there is a mass-emission time anticorrelation: large- M pairs are emitted first, small- M pairs last. Very-large- M pairs, say, $M \gtrsim 10$ GeV, are emitted at times $\tau \sim 1/M$ by the single-collision and totally unthermalized Drell-Yan mechanism. This mechanism is well under theoretical control, but does not involve the thermal system discussed here. While the system thermalizes, it can also emit pairs with intermediate masses. What this mass range is depends on the total multiplicity of the event. For the rapidity densities discussed below (of the order of 5000 in central U + U collisions), its lower limit could be $M \gtrsim 3$ GeV. For small particle densities, it is possible that there is no thermal and even little pre-equilibrium emission. Ultimately these questions will be settled by experiment.

With dilepton measurements one has thus a fairly detailed handle on all the stages of the collision history from the earliest to the latest times. The required multiplicities are so large that the experiments are clearly very demanding. On the other hand, if either these large multiplicities are never produced in the collisions or if one is unable to experimentally observe them, one may never see evidence of these very interesting collective phenomena on the nuclear scale.

II. THE FLOW

The general features of a hydrodynamic flow through a quark-plasma (Q), mixed (M), and hadron (H) phase

under an assumed first-order equation of state, have been discussed in several places (see, for instance, Refs. 10–11) and we do not go into detail here. Just to fix our notation, we note that our equation of state is, for the quark-plasma and hadron phases, given by

$$\begin{aligned} p_q(T) &= g_q \frac{\pi^2}{90} T^4 - B, \\ p_h(T) &= g_h \frac{\pi^2}{90} T^4, \end{aligned} \quad (2.1)$$

where

$$\begin{aligned} g_q &= 2 \times 8 + 2.5 \times 2 \times 2 \times 3 \times \frac{7}{8} = 42.25, \\ g_h &= 3 \end{aligned} \quad (2.2)$$

(we use $N_F=2.5$ to simulate the effect of the strange-quark mass). Many quantities depend only on the ratio

$$r = \frac{g_q}{g_h} \approx 14. \quad (2.3)$$

The bag constant B and the transition temperature T_c are related by

$$B = (g_q - g_h) \frac{\pi^2}{90} T_c^4, \quad (2.4)$$

and the values of the energy density in the quark and hadron phases at T_c are

$$\begin{aligned} \epsilon_Q(T_c) &\equiv \epsilon_Q = \epsilon_H + 4B, \\ \epsilon_h(T_c) &\equiv \epsilon_H = \frac{3B}{r-1}. \end{aligned} \quad (2.5)$$

We shall mostly use the value $T_c=200$ MeV for the transition temperature and $T_f=50$ MeV for the decoupling temperature. These correspond to (in units of GeV/fm³)

$$\begin{aligned} \epsilon_Q &= 3.62, \\ \epsilon_H &= 0.21, \\ \epsilon_f &= 0.001. \end{aligned} \quad (2.6)$$

In the hydrodynamic flow with no transverse motion the times between the different phases are given by

$$\begin{aligned} \tau_Q &= \left(\frac{T_i}{T_c} \right)^3 \tau_i, \\ \tau_H &= r \tau_Q, \\ \tau_f &= \left(\frac{T_c}{T_f} \right)^3 \tau_H. \end{aligned} \quad (2.7)$$

Note here that a proper treatment of decoupling from the hadron phase would require the use of kinetic theory. The use of hydrodynamics down to $T_f=50$ MeV in the dilepton-emission calculations below is to be understood as an estimate of the upper limit of the contribution from the hadron phase. A conservative lower limit is obtained by including only the mixed phase (i.e., increasing T_f to T_c) and the results from this estimate are in qualitative accord with the results obtained with $T_f=50$ MeV.

The hydrodynamic flows corresponding to the parame-

ters in Table I were computed using the methods of Ref. 10 and some of the results are exhibited in Figs. 1–5. One can observe the following (the sizes and lifetimes given below refer, for definiteness, to a decoupling temperature $T_f=50$ MeV; sizes corresponding to other values of T_f can be read from the figures).

(1) At $T_i=250$ MeV, $\tau_i=1$ fm, the quark phase lasts only very briefly, from 1 to 2 fm, and its transverse motion is totally negligible. Even the mixed phase is hardly at all affected by transverse motion, while the hadron phase moves in the transverse direction. The system lives for about 35 fm and its largest transverse extent is 17 fm.

(2) At $T_i=500$ MeV, $\tau_i=0.5$ fm, the value of τ_Q ($=7.8$ fm) is already so close to the transverse time scale ($R_A/c_s \approx 12$ fm) that there appears a moderate buildup of transverse expansion at the sides of the quark phase. As a result, large parts of the mixed phase and the entire hadron phase flow transversally. This reduces significantly the growth of the lifetime of the system (relative to $v_r=0$), which now is 50 fm, the largest transverse extent being 30 fm.

(3) The flows with $T_i=350$ MeV, $\tau_i=1.5$ fm and $T_i=833$ MeV, $\tau_i=0.11$ fm have the same $T_i^3\tau_i$, i.e., the same dN/dy . They are thus expected to be essentially the same flows, just started at different times and, correspondingly, different temperatures. This is verified by the numerical computation (results not shown). Actually small differences, caused by initial transverse velocity effects, appear: if we vary τ_i between 1.4, 0.5, and 0.11 fm, the transverse velocity field clearly cannot be assumed to vanish at all these times. Thus the differences also give a feeling of the effects of changing the initial radial velocity.

(4) At $T_i=833$ MeV and $\tau_i=0.33$ fm, $\tau_Q=24$ fm is much larger than the transverse time scale and also the quark phase is set in transverse motion. Thus also its lifetime is reduced from the value 24 fm, valid for no transverse motion, to about 12 fm. Because of the rapid transverse motion the lifetime of the mixed phase is reduced even relative to the previous $dN/A dy=26$ case, but ultimately the system lives long as a very dilute pion gas (more properly treated by kinetic theory) so that its total lifetime is 65 fm and its largest extent about 50 fm.

Note that physically this example is rather extreme. The initial energy density implied in this circumstance

corresponds to all of the kinetic energy of the two nuclei being deposited in a rapidity interval of one unit when the collision energy is $100+100$ GeV.

(5) If T_c is decreased from 200 to 160 MeV, everything else remaining unchanged, the qualitatively essential change is the increase of τ_Q by a factor 2.0. This implies that already the quark phase is set in rapid transverse motion, which further propagates to the mixed and hadron phases. The expected net result is an increase in the space-time volume of the quark phase, a decrease in that of the mixed and hadron phases, and an increase in the overall transverse motion. The expectations are borne out by the numerical computation (results in Fig. 4).

(6) All the above flows assume that T_i is independent of r . The expected r dependence is correlated with the A dependence of dN/dy : if

$$\frac{dN}{dy} \propto A^{(2/3)(1+p)},$$

then

$$s_i(r)\tau_i \propto \left[1 - \frac{r^2}{R_A^2}\right]^p.$$

We expect p to be about $\frac{1}{2}$, $p=1$ being a rather extreme value. The flow corresponding to this extreme r dependence,

$$T_i(r) = T_{i0} \left[1 - \frac{r^2}{R_A^2}\right]^{1/3}, \quad (2.8)$$

with $\tau_i=0.5$ and $T_{i0}=630$ MeV (so that the total entropy of the flow is normalized to the same value $dN/A dy=26$ as for the flow with $T_i=500$ MeV, $\tau_i=0.5$ fm) is given in Fig. 5. As expected, the transverse flow is now more rapid. Even the mixed phase now flows transversally and its lifetime near $r=0$ is considerably reduced, less so at larger r . However, now the system lives longer in the hadron phase so that the total lifetime at $r=0$ is about the same as with constant T_i .

The huge lifetimes of the mixed and hadron phases (Table I) in the approximation $v_r=0$ are thus considerably reduced. Still, this reduction in the time direction is partially compensated by the expansion of the system in the transverse direction.

TABLE I. Values of parameters used (in units of MeV or fm, $A=238$). Note the logic in the table: the primary quantity is the associated pion multiplicity dN/dy , which can be experimentally fixed and which fixes $T_i^3\tau_i$. All the flows with the same dN/dy and T_c have the same one-dimensional ($v_r=0$) time scales [Eq. (2.7)]. Changing T_c changes the relative balance of Q and M but not the total one-dimensional duration. The true three-dimensional lifetimes are discussed in Sec. II (see Figs. 1–5).

$dN/A dy$	T_i	T_c	τ_i	τ_Q	τ_H	τ_f
6.4	250	200	1.0	1.9	27	1 800
26	350	200	1.5	7.8	110	7 000
26	500	200	0.5	7.8	110	7 000
26	833	200	0.11	7.8	110	7 000
26	500	160	0.5	15	215	7 000
78	833	200	0.33	24	340	21 500




FIG. 1. Curves of (a) constant energy density (in units of GeV/fm^3) and (b) constant radial velocity for $T_i=250$ MeV, $\tau_i=1$ fm, and $T_c=200$ MeV. The velocity contours near the light cone in this and the following figures are an artifact of the numerical method and graphics and have no physical meaning.

III. EMISSION OF DILEPTONS

The calculation of dilepton emission rate per collision proceeds straightforwardly by convoluting the rest-frame emission rates from the quark-plasma and the hadron phases with the flows computed above. In the mixed phase, the system emits both from the quark and hadron phases, the relative amount of these being given by entropy conservation.

The emission rate from the quark phase is well known to be

$$\frac{dN}{d^4x dM^2 d^3p/E} = \frac{\alpha^2}{8\pi^4} \sum e_q^2 e^{-E/T}. \quad (3.1)$$

The calculation of the emission rate from the hadrons is much more complicated.⁶ Since we model the hadron




FIG. 2. Same as Fig. 1 but for $T_i=500$ MeV and $\tau_i=0.5$ fm. The curves for the cases having the same dN/dy in Table I are essentially the same apart from the starting time τ_i of the flow.

phase by a pion gas, we only include the process $\pi^+\pi^-\rightarrow l^+l^-$ and include only the ρ pole in the form factor:

$$F^2(M) = \frac{m_\rho^4}{(m_\rho^2 - M^2)^2 + m_\rho^2 \Gamma_\rho^2}, \quad (3.2)$$

with $m_\rho=0.77$ and $\Gamma_\rho=0.155$ GeV. With this form factor, the emission rate is

$$\frac{dN}{d^4x dM^2 d^3p/E} = \frac{\alpha^2}{8\pi^4} \left[1 + \frac{2m_l^2}{M^2} \right] \left[1 - \frac{4m_\pi^2}{M^2} \right] \times \frac{1}{12} F^2(M) e^{-E/T}. \quad (3.3)$$

This could be improved by including more resonances in the form factor or, even better, by using e^+e^- data for





FIG. 3. Same as Fig. 1 but for $T_i = 833$ MeV and $\tau_i = 0.33$ fm.





FIG. 4. Same as Fig. 2 but for $T_c = 160$ MeV.

pion production. Also, one might want to include a temperature-dependent mass and width^{20,21} for the ρ , to obtain a method for experimentally testing these T dependences. If the ρ significantly alters its mass in the pion phase, this shift might indeed appear in the experimentally observed dilepton spectrum. The degree of such a shift near the deconfinement temperature in the hadron phase is unknown. However, the main approximation is probably the neglect of bremsstrahlung production of dileptons: $\pi^+\pi^-\rightarrow\pi^+\pi^-\gamma^*$. This will certainly dominate in the region $M < 2m_\pi$. It can be calculated, but the calcula-

tion is technically rather complicated. We believe that the simple ρ -pole approximation used here is quite adequate for pair masses near the ρ mass and above.

We are also neglecting the sharp resonances ϕ (Ref. 22) and ψ (Ref. 23). A calculation of their production rate with the above realistic flows would be another interesting problem. Since the ψ mass is so large, it is emitted very early in the collision from the quark-plasma phase and we actually expect the no-transverse-flow approximation of Ref. 23 to be good.

The convolution over the flow is now of the form

$$\begin{aligned}
 \frac{dN}{dM^2 dy d^2 p_T} &= \int d^4 x e^{-u \cdot p/T} \\
 &= \int \tau d\tau d\eta r dr d\varphi \exp \left[-\frac{\gamma_r}{T} [M_T \cosh(\eta - y) - v_r p_T \cos\varphi] \right] \\
 &= 4\pi \int \tau d\tau r dr K_0 \left[\frac{\gamma_r M_T}{T} \right] I_0 \left[\frac{\gamma_r v_r p_T}{T} \right], \tag{3.4}
 \end{aligned}$$

where we have used

$$\tau = (t^2 - z^2)^{1/2},$$

$$\eta = \frac{1}{2} \ln \frac{t+z}{t-z},$$

$$u^\mu = \gamma_r \left[\frac{t}{\tau}, v_r \cos \varphi, v_r \sin \varphi, \frac{z}{\tau} \right],$$

$$p^\mu = (E, p_T, 0, p_L)$$

$$M_T = (M^2 + p_T^2)^{1/2}, \quad E = M_T \cosh \eta,$$

and K_0 and I_0 are modified Bessel functions. The transverse velocity v_r [$\gamma_r = 1/(1-v_r^2)^{1/2}$] and the temperature T are given as functions of τ and r by the numerical calculation of the flow.

Noting that the relative volumes of the quark-plasma and hadron phases within the mixed phase are given by

$$\frac{V_Q}{V} = \frac{\epsilon - \epsilon_H}{\epsilon_Q - \epsilon_H}, \quad \frac{V_H}{V} = \frac{\epsilon_Q - \epsilon}{\epsilon_Q - \epsilon_H},$$

we can write the dilepton rate in the final form

$$\begin{aligned} & \frac{dN}{dM^2 dy d^2 p_T} \\ &= \frac{\alpha^2}{2\pi^3} \int \tau d\tau r dr I_0 \left[\frac{\gamma_r v_r p_T}{T} \right] K_0 \left[\frac{\gamma_r M_T}{T} \right] \\ & \quad \times \left[\sum e_q^2 \theta(Q) \right. \\ & \quad \left. + \left[\frac{\epsilon - \epsilon_H}{\epsilon_Q - \epsilon_H} \sum e_q^2 + \frac{\epsilon_Q - \epsilon}{\epsilon_Q - \epsilon_H} G(M^2) \right] \right. \\ & \quad \left. \times \theta(M) + G(M^2) \theta(H) \right], \quad (3.5) \end{aligned}$$

where the θ functions test whether $\epsilon > \epsilon_Q$ [$\theta(Q)$], $\epsilon_H < \epsilon < \epsilon_Q$ [$\theta(M)$], or $\epsilon_f < \epsilon < \epsilon_H$ [$\theta(H)$] and (we take $m_l = 0$)

$$G(M^2) = \frac{1}{12} F^2(M) \left[1 - \frac{4m_\pi^2}{M^2} \right].$$

Further, the result (3.5) can be integrated analytically over the p_T of the emitted dilepton with the result

$$\frac{dN}{dM^2 dy} = \frac{\alpha^2}{\pi^2} \int \tau d\tau r dr M T K_1 \left[\frac{M}{T} \right] (\dots), \quad (3.6)$$

where the ellipsis represents the quantity within the square brackets in (3.5). Surprisingly, v_r does not appear at all in Eq. (3.6). The result looks very simple, but T (and the ϵ in the square brackets) is only given by the numerical hydrodynamic calculation as a function of τ and r .

If the radial flow is neglected ($v_r = 0$), the above results simplify to the form

$$\begin{aligned} \frac{dN}{dM^2 dy d^2 p_T} &= \frac{\alpha^2}{4\pi^4} \pi R_A^2 (T_i^3 \tau_i)^2 \frac{3}{M_T^6} \left[\int_{M_T/T_i}^{M_T/T_c} dy y^5 K_0(y) \sum e_q^2 + \frac{r-1}{18} \left[\sum e_q^2 + r G(M^2) \right] \left[\frac{M_T}{T_c} \right]^6 K_0 \left[\frac{M_T}{T_c} \right] \right. \\ & \quad \left. + r^2 \int_{M_T/T_c}^{M_T/T_f} dy y^5 K_0(y) G(M^2) \right] \quad (3.7) \end{aligned}$$




FIG. 5. Same as Fig. 2 but for the radially varying initial temperature distribution in Eq. (2.8).

and

$$\frac{dN}{dM^2 dy} = \frac{\alpha^2}{4\pi^4} \pi R_A^2 (T_i^3 \tau_i)^2 \frac{6\pi}{M^4} \left[\int_{M/T_i}^{M/T_c} dy y^4 K_1(y) \sum e_q^2 + \frac{r-1}{18} \left(\sum e_q^2 + rG(M^2) \right) \left(\frac{M}{T_c} \right)^5 K_1 \left(\frac{M}{T_c} \right) + r^2 \int_{M/T_c}^{M/T_f} dy y^4 K_1(y) G(M^2) \right]. \quad (3.8)$$

Within the domain of validity of the approximations, one could replace the Bessel functions by $K_m(y) = (\pi/2y)^{1/2} \exp(-y)$.

IV. NUMERICAL RESULTS

The formulas given for the dilepton rate in Sec. III are easily integrated over the flows in Sec. II. Results are given in Figs. 6–13. Note that we give all numbers as rates per a collision at zero impact parameter b . The experiments will attempt to trigger for central collisions and averages over b are not needed. However, when comparing with a Drell-Yan rate (which is averaged over b), the averaging has to be taken into account. The rates can be turned to cross sections by multiplying with $\sigma \simeq \pi(R_A + R_B)^2$ [and dividing by $(1 + R_A/R_B)^2$, if averaging over b is required].

Figure 6 gives the wholly differential rate $dN/dM^2 dy d^2 p_T$ for the flow with $T_i = 500$ MeV, $\tau_i = 0.5$ fm, corresponding to $dN/A dy = 26$. For comparison, the same quantity is given in Fig. 7, calculated if only longitudinal cooling with no transverse flow is included, in Fig. 8 for a flow which has the same comoving entropy but which starts later, at $\tau_i = 1.5$ fm corresponding to $T_i = 350$ MeV, and in Fig. 9 for a flow starting at $\tau_i = 0.5$ with the same comoving entropy but with an initially r -dependent temperature distribution [Eq. (2.8)].

Consider first the quark-plasma contribution, which has, in detail, been studied in Refs. 8 and 9. This is “the new phase of matter” one expects to diagnose with these dilepton measurements. It is formed very early in the process and thus, as is explicitly shown by Figs. 1–5 and a comparison of Figs. 6 and 7, the transverse flow has little effect on it. If one goes to details, one observes that the transverse flow reduces the rate somewhat and the reduction is largest at small M_T . This is natural, since small- M_T pairs are dominantly produced late in the quark-plasma phase, so that the transverse flow has had time to penetrate towards the center and further reduce the temperature.

Another consequence of the insensitivity of the quark-matter contribution to the transverse flow is that the rate from it is essentially independent of the pair mass M when plotted against M_T . For the $v_r = 0$ case this observation was made already in Ref. 8.

Another relevant remark can be made by considering two different flows having the same total entropy, i.e., the value of $T_i^3 \tau_i$. If we change the value of τ_i from 0.5 to 1.5 fm (as in Figs. 6 and 8), the flow after 1.5 fm is un-

changed, if the initial condition at 1.5 fm is taken to be the initial condition at 0.5 fm hydrodynamically evolved to 1.5 fm. In the calculations of Sec. II $v_r = 0$ was actually used as the initial condition at both times. This causes a small change in the later flows. Apart from this small change, the only change in the dilepton rate is then that, when increasing τ_i from 0.5 to 1.5 fm, we lose the contribution from this time interval. As the matter is then in a very hot quark-plasma phase (between $T = 350$ and 500 MeV), the loss is essentially in the production of large- M_T pairs. In the case of Figs. 6 and 8 this loss is an order of magnitude or more at $M_T > 3.5$ GeV. We again emphasize that this effect is an early-time large- M_T effect and thus independent of transverse flow.

Ideally, one could use the sensitivity of the large- M_T dilepton rate to very early times to experimentally determine the initial thermalization time τ_i and temperature T_i separately as follows. First separate the dilepton events according to the value of the associated pion multiplicity dN/dy . This (given R_A) fixes the value of the product $T_i^3 \tau_i$ according to Eq. (1.1). Considering now this fixed value of dN/dy one measures the rate of dilepton production at large M_T . Its value depends on T_i and τ_i separately and, when measured, can thus be used to give a numerical value for τ_i and T_i separately. This is not a value of τ_i for each event separately (as is the product $T_i^3 \tau_i$) but an average τ_i for all the events having a fixed value of $T_i^3 \tau_i$. Since the initial value of T_i may and will depend on r and will also stochastically fluctuate across the nuclear disc, the measured value of τ_i will also contain an average across the transverse coordinate.

As discussed in the Introduction, note also that M_T should neither be so large that one is in the single-collision Drell-Yan region nor even in the intermediate pre-equilibrium region. We shall give an estimate for this when discussing $dN/dM dy$ below.

Consider then the mixed- and hadron-phase contributions. A comparison between Figs. 6 and 7 shows that the result for correct transverse flow is significantly different from that obtained by taking $v_r = 0$ [Eqs. (3.7) and (3.8)]: the rate at small M_T is reduced by orders of magnitude and at large M_T increased by orders of magnitude. This is a direct effect of the transverse flow. Pairs with $M < 1$ GeV are dominantly emitted at late times from relatively cool matter, which is moving transversally with a large velocity (see Figs. 1–5). This imparts a large p_T to the particles and makes the M_T distribution much flatter than for $v_r = 0$. For $v_r = 0$ the M_T distribution of pairs from the mixed phase is [Eq. (3.7)]




FIG. 6. The quark-phase (Q), mixed-phase (M), and hadron-phase (H) contributions to the differential dilepton production rate $dN/dM^2 dy d^2 p_T$ plotted vs M_T for $M=0.3, 0.8$, and 1.5 GeV for the flow with $T_i=500$ MeV and $\tau_i=0.5$ fm. Up to negligible corrections, the Q contribution is independent of pair mass M when plotted vs M_T .

$$K_0 \left[\frac{M_T}{T_c} \right] \sim M_T^{-1/2} e^{-M_T/T_c}$$

(this also holds for H if $M_T > 5T_c$). For true flow this is found to be much flatter but not quite as flat as $\exp(-M_T/T_i)$.

The transverse-flow effect is particularly clear if one chooses a mass which maximizes the M and H contributions relative to Q : the region near $M=m_\rho$. From Figs. 6 and 8 it is clear that M and H will dominate over Q at all values of M_T . In detail the persistence of the ρ peak is shown in Fig. 10. Its relative importance thus even increases with increasing M_T . This situation is quite different from that at $v_r=0$. As shown in Fig. 7, the ρ peak there will vanish under the mass-independent quark-plasma contribution for $M_T > 3$ GeV (Ref. 19). The observation of the ρ peak at large M_T would be direct evidence for a collective flow of a hadronic system. It is hard to see how this effect could be reproduced by the




FIG. 7. Same as Fig. 6 but for zero radial flow. Q is independent of pair mass M .




FIG. 8. Same as Fig. 6 but for the flow with $T_i=350$ MeV and $\tau_i=1.5$ fm. This flow has the same comoving entropy as the one in Fig. 6.

various models for particle production in nucleus-nucleus collisions, which make no reference to collective behavior. This clearly is an issue deserving further study.

In principle, if QCD matter existed in the hadron phase up to very high temperatures, the pions in the high- T hadron gas would also produce ρ 's at large M_T and our conclusion about the persistence of the ρ peak would be weakened. However, by simple density arguments we find it physically very improbable that the hadron phase could exist up to such high temperatures that our conclusion about the ρ peak at $M_T=4$ GeV would be significantly affected. And in this improbable case other hadronic degrees of freedom, not contributing to production of pairs with $M=m_\rho$, would be increasingly important. As seen from Fig. 9, the persistence of the ρ peak is also stable against introducing r dependence in the initial temperature, although the flows in the mixed phase seem considerably different (compare Figs. 2 and 5). The rapid evolution of the system in the mixed phase in Fig. 5 means that energy is not wasted in longitudinal expansion. It goes into transverse expansion and maintains the large emission rate into large M_T near $M=m_\rho$.

Consider then the results for the p_T -integrated rate




FIG. 9. Same as Fig. 6 but now calculated for the flow in Fig. 5 with the radially varying initial temperature distribution in Eq. (2.8).




FIG. 10. Same as Fig. 6 but now plotted vs M for transverse masses $M_T=1, 2$, and 4 GeV. The dashed lines show the separate contributions of the different phases at $M_T=4$. Note the persistence and even strengthening of the ρ peak. The ρ peak persists even if the hadron-phase contribution is neglected.

$dN/dM dy$ [Eq. (3.6)], shown in Figs. 11–13. At this point it is also convenient to compare the result with the Drell-Yan background. A rough estimate for this at $M \ll \sqrt{s}$ is

$$\frac{dN_{A+A}^{DY}}{dM dy} = 4 \frac{A^{4/3} \times 3.8 \times 10^{-8}}{M^3} \left(\frac{4\pi R_A^2}{\sigma_{in}} \right) \times \left(\frac{S^4(0) + \text{valence-sea effects}}{0.25A} \right)^2 \quad (4.1)$$

which is directly obtained from the standard Drell-Yan formula (no K factor) and in which we expect the quantities in large parentheses to be of order 1. The factor 4 in front corrects the formula for central collisions and M is expressed in GeV units to give the result in $1/\text{GeV}$. If T_i is clearly above T_c , the Drell-Yan background is clearly below thermal emission for the range of masses shown ($M < 3$ GeV); for high dilepton masses the Drell-Yan rate will, of course, ultimately win. For $T_i=250$ MeV (Fig. 13) there is so little of the quark-plasma phase that the Drell-Yan mechanism—especially if supplemented by the so-far uncalculated pre-equilibrium emission—gives a relatively large contribution.

The following comments can be made concerning Figs. 11–13. The three parts of Fig. 11 illustrate, as discussed above, how one gains more quark-plasma contribution by starting the flow earlier. The gain is from high temperatures and goes, accordingly, to large dilepton masses. The quantitative observation of the high-mass tail in Fig. 11 is precisely “diagnosis of quark plasma.” Figure 12 illustrates what happens if T_c is decreased from 200 to 160 MeV. As discussed in Sec. II, the quark phase then lasts longer at temperatures close to T_c . The small-mass part

of Q is thus somewhat increased. The M and H contributions decrease above the ρ faster at $T_c=160$ MeV, roughly proportionally to $\exp(-M/T_c)$. However, at this large T_i it is not possible to observe this behavior, since it is hidden below the large- Q contribution. To measure the value of T_c one has to go to smaller values of T_i (Fig. 13). The Q term then is much smaller and the $\exp(-M/T_c)$ behavior is exposed. However, the Drell-Yan contribution is threateningly close.

We conclude this section by noting that the calculated $dN/dM dy$ for produced dilepton pairs in the ρ -mass re-




FIG. 11. The quark (Q), the mixed (M), the hadron phase (H) contributions, and the total differential dilepton production rate $dN/dM dy$ plotted vs pair mass M for the three flows (a) $T_i=350$ MeV, $\tau_i=1.5$ fm, (b) $T_i=500$ MeV, $\tau_i=0.5$ fm, (c) $T_i=833$ MeV, $\tau_i=0.11$ fm having the same associated pion multiplicity $dN/dy=26A$ (Table I). Note how M and H are essentially unchanged while the large- M behavior of Q increases when τ_i is decreased and more of the hotter initial matter is included.




FIG. 12. Same as Fig. 11(b) but for $T_c = 160$ MeV.

gion can be used to estimate the number of $\pi\pi$ collisions which took place in the mixed and hadron phases. The pions, namely, dominantly interact via the ρ resonance (apart from $\pi^0\pi^0$), and what we have calculated is how often this resonance, after a lifetime of about 1 fm, decays to a dilepton pair. To estimate the total number of interactions we simply write

$$\text{number of } \pi\pi \text{ collisions} = \frac{3\Gamma_\rho}{B(\rho \rightarrow e^+e^-)} \frac{dN}{dM dy}, \quad (4.2)$$

where the 3 accounts for $\pi^+\pi^0$, $\pi^+\pi^-$, and $\pi^-\pi^0$. For instance, for the $T_i = 500$ MeV, $\tau_i = 0.5$ fm case this gives about 3000 interactions in the mixed phase and 1500 in the hadron phase. Counting collisions per particle gives a further factor of 2. In the mixed phase there are in addition pion-quark and pion-gluon interactions. There should not be too much difficulty maintaining thermal equilibration in the mixed phase. In the hadron phase, this is however enough for kinetic theory but marginally so for hydrodynamics. Note also that these final remarks are in agreement with the standard observation that essentially all the pions produced in hadronic collisions are produced via the ρ . This observation just counts the last interaction via a ρ as the produced particle, while we have been considering the pions produced in the previous interactions as elements of the hadronic fluid.

V. CONCLUSIONS

We have in this paper considered the effects of transverse collective flow of QCD matter as it might be formed in ultrarelativistic nucleus-nucleus collisions. This is clearly an important question; we cannot really talk about “matter” unless we can prove that the system behaves collectively. The question of the existence of collective flow has also been a central issue in connection of nuclear collisions in the 1-GeV range.²⁴

Under reasonable initial conditions, the quark-plasma phase is formed, if at all, very early in the collision. It thus is little affected by transverse motion and its collective motion is essentially longitudinal. This, however, is next to impossible to separate from the longitudinal motion, which is present in the produced particles. Instead, we have attempted to discuss how the parameters T_i and τ_i of the flow could be determined.

The situation is quite different with the later stages of

the process. If the system has started high up in the quark-plasma phase, it is already in rapid transverse motion after having cooled down to the mixed phase. Whatever is emitted at this stage, will get a boost in the transverse direction from this motion. As the system is already rather cool, it will dominantly emit dileptons in the 1-GeV mass range and we have shown that this emission from matter in transverse motion may lead to the persistence of the ρ peak until very large M_T . The effect is further amplified by the emission from the very late and cool hadron phase, which should properly be treated with kinetic theory.

If the persistence of the ρ peak at large M_T is experimentally observed, one further has to prove that this cannot convincingly be reproduced by any of the mechanism of particle production not containing collective motion. The discussion will perhaps parallel that around cascade versus collective models in the 1-GeV range.²⁵

Concerning the experimental observation of the effects discussed here one should note that an average $A + A$ event might have $T_i = T_c$ and starts expansion in the mixed phase. For the effects to be fully developed, the expansion time has to be long and the system must start at high temperature in the quark-plasma phase, the higher the better. One will thus be studying the tail of the multiplicity distribution and the length of this tail will determine how well these collective effects can be studied. The average and the width of the multiplicity distribution are crucial unknown parameters, on which even the first-generation relativistic heavy-ion experiments will give important information.

ACKNOWLEDGMENTS

We have profited from discussions with P. Glässel, J. Kapusta, and A. Mekjian.




FIG. 13. Same as Fig. 11 but for the flow with $T_i = 250$ MeV, $\tau_i = 1$ fm. The dashed line gives the single-scattering Drell-Yan contribution according to Eq. (4.1).

- *Permanent address: Academy of Finland and Department of Theoretical Physics, Siltavuorenpenger 20 C, SF-00170 Helsinki, Finland.
- †Permanent address: Department of Physics, Seminaarinkatu 15, SF-40100 Jyväskylä, Finland.
- ‡Permanent address: Fermilab, P.O. Box 500, Batavia, IL 60510.
- ¹For a review of the present state of quark-matter physics, see *Quark Matter '84*, proceedings of the Fourth International Conference on Ultrarelativistic Nucleus-Nucleus Collisions, edited by K. Kajantie (Lecture Notes in Physics Vol. 221) (Springer, New York, 1985); or *Proceedings of the RHIC Workshop*, edited by P. Hausteiner and C. Woody (BNL Report No. 51921, 1985).
- ²Gordon Baym, B. Friman, J.-P. Blaizot, M. Soyeur, and W. Czyz, Nucl. Phys. **A407**, 541 (1983).
- ³J. D. Bjorken, Phys. Rev. D **27**, 140 (1983).
- ⁴E. L. Feinberg, Nuovo Cimento A **34**, 39 (1976); E. Shuryak, Phys. Lett. **78B**, 150 (1978); Yad. Fiz. **28**, 796 (1978) [Sov. J Nucl. Phys. **28**, 1548 (1978)].
- ⁵V. Cerny, P. Lichard, and J. Pisut, Phys. Lett. **70B**, 61 (1977); Phys. Rev. D **16**, 2822 (1977); **18**, 2409 (1978); **24**, 652 (1981).
- ⁶G. Domokos and J. Goldman, Phys. Rev. D **23**, 203 (1981); G. Domokos, *ibid.* **28**, 123 (1983).
- ⁷K. Kajantie and H. I. Miettinen, Z. Phys. C **9**, 341 (1981); **14**, 357 (1982).
- ⁸L. D. McLerran and T. Toimela, Phys. Rev. D **31**, 545 (1985).
- ⁹R. Hwa and K. Kajantie, Phys. Rev. D **32**, 1109 (1985).
- ¹⁰E. von Gersdorff, M. Kataja, L. McLerran, and P. V. Ruuskanen, Phys. Rev. D **34**, 794 (1986).
- ¹¹B. Friman, K. Kajantie, and P. V. Ruuskanen, Nucl. Phys. **B266**, 468 (1986).
- ¹²A. Białas, W. Czyz, and A. Kolawa, Acta Phys. Pol. **B15**, 229 (1984); S. Pratt, University of Minnesota report, 1985 (unpublished).
- ¹³A. Białas and J.-P. Blaizot, Phys. Rev. D **32**, 2954 (1985).
- ¹⁴K. Kajantie and T. Matsui, Phys. Lett. **164B**, 373 (1985).
- ¹⁵M. Gyulassy and A. Iwazaki, Phys. Lett. **165B**, 157 (1985).
- ¹⁶A. Kerman, T. Matsui, and B. Svetitsky, Phys. Rev. Lett. **56**, 219 (1986).
- ¹⁷L. Van Hove, Z. Phys. C **27**, 135 (1985).
- ¹⁸M. Gyulassy, K. Kajantie, H. Kurki-Suonio, and L. McLerran, Nucl. Phys. **B237**, 477 (1984); N. K. Glendenning and T. Matsui, Phys. Lett. **141B**, 419 (1984); B. Friman, *ibid.* **159B**, 369 (1985).
- ¹⁹P. Siemens and S. A. Chin, Phys. Rev. Lett. **55**, 1266 (1985).
- ²⁰A. I. Bochkarev and M. E. Shaposnikov, Phys. Lett. **145B**, 276 (1984).
- ²¹R. Pisarski, Phys. Lett. **110B**, 155 (1982).
- ²²A. Shor, Phys. Rev. Lett. **54**, 1122 (1985).
- ²³J. Cleymans and C. Vanderzande, Phys. Lett. **147B**, 186 (1984).
- ²⁴H. A. Gustafsson *et al.*, Phys. Rev. Lett. **52**, 1590 (1984).
- ²⁵J. Cugnon, in *Quark Matter '84* (Ref. 1).