

Quantization on the light cone: Response to a Comment by Hagen

Kent Hornbostel and Stanley J. Brodsky

Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305

Hans-Christian Pauli

Max Planck Institute für Kernphysik, D-6900 Heidelberg 1, Germany

(Received 31 August 1987)

We show that, if carefully defined, light-cone commutation relations are not inconsistent with the charge and Lorentz invariance of the vacuum. We demonstrate this for charge invariance with an explicit example.

The method of discretized light-cone quantization¹ (DLCQ) has recently been applied to a series of field theories in one space and one time dimension, including Yukawa theory¹ (charged fermions coupled to scalar bosons), quantum electrodynamics,² and quantum chromodynamics (for arbitrary N_{color}) (Ref. 3). By diagonalizing the Hamiltonian defined at a given light-cone time on a discrete momentum Fock basis, one obtains not only the charge-zero spectrum but also the bound-state wave functions of each field theory for arbitrary coupling constant and fermion mass. The numerical results agree with previous calculations of the spectrum where they are available.

The preceding Comment by Hagen⁴ raises several issues concerning the consistency of field theory quantized on the light cone. These should not be confused with gauge-fixing problems, such as the choice of light-cone gauge $A^+ = 0$. A recent discussion of such problems is given in Ref. 5. Indeed Hagen's remarks are directed to the application of DLCQ to Yukawa theory where there are no gauge fields.

Hagen's first remark⁴ [see his Eq. (1)] concerns a possible unwanted term in the Poincaré algebra. Our choice of periodic boundary conditions trivially eliminates this term. The remainder of his Comment reiterates claims made in his earlier papers that the light-cone vacuum is neither charge nor Lorentz invariant. Although these issues are not directly related to our paper, they should be evaluated on their merit, independently of their relevance to our work. In fact, we will present an explicit counterexample to the central theorem on which Hagen's main objections rely. In particular, we will investigate the claim that a charge-invariant light-cone vacuum cannot be defined;⁶ the discussion of Lorentz invariance is similar.

Hagen's theorem assumes a conserved current, the existence of a spectral decomposition for vacuum expectation values of this current, and standard light-cone equal- x^+ commutation relations for scalar fields. To avoid any possible ambiguity or controversy arising from light-cone quantization techniques we will employ noninteracting scalar fields of mass m quantized at equal time. Although quantized conventionally, these fields will be

shown to satisfy all the requirements necessary to test Hagen's theorem.

For simplicity, we work in the relevant two dimensions throughout. In equal-time quantization, commutators are fixed at $x^0 = 0$, but because in this example the fields are free, they are known for all x . In particular, $\langle [\phi(x), \phi(0)] \rangle$ is known at $x^+ \equiv x^0 + x^1 = 0$, where

$$\langle [\phi(x), \phi(0)] \rangle_{x^+ = 0} \equiv i \Delta(x; m^2)_{x^+ = 0} = \frac{-i}{4} \epsilon(x^-). \quad (1)$$

This will be demonstrated below. [$\epsilon(x)$ is the antisymmetric step function.]

Now allow for a conserved current by making $\phi(x)$ complex, with $(1/\sqrt{2})\phi_1(x)$ and $(1/\sqrt{2})\phi_2(x)$ the real and imaginary parts. The commutator of ϕ and ϕ^* is as in Eq. (1). The current-current commutator, with

$$j^\mu(x) = :\phi_2(x)\partial^\mu\phi_1(x) - \phi_1(x)\partial^\mu\phi_2(x): \quad (2)$$

may be expressed in a general spectral form

$$\langle [j^\mu(x), j^\nu(0)] \rangle = (g^{\mu\nu}\partial^2 - \partial^\mu\partial^\nu) \times \int_0^\infty dM^2 \sigma(M^2) i \Delta(x; M^2), \quad (3)$$

where, for free fields,

$$\sigma(M^2) = \Theta \left[1 - \frac{4m^2}{M^2} \right] \frac{\left[1 - \frac{4m^2}{M^2} \right]^{1/2}}{2\pi M^2}. \quad (4)$$

The assumptions of Hagen's proof⁶ are therefore satisfied.

The charge operator on the light cone may be computed, also unambiguously:

$$\begin{aligned} Q &\equiv \int \frac{dx^-}{2} j^+(x) \\ &= \int \frac{dx^-}{2} [j^0(x) + j^1(x)] \\ &= i \int dk^1 [a_1^\dagger(k^1) a_2(k^1) - a_2^\dagger(k^1) a_1(k^1)], \end{aligned} \quad (5)$$

where $a_i(k^1)$ and $a_i^\dagger(k^1)$ are the usual equal-time creation and annihilation operators. The charge defined

in Eq. (5) is thus explicitly identical to the charge Q defined conventionally at equal time,

$$Q \equiv \int dx^1 j^0(x), \quad (6)$$

as current conservation guarantees. It is obvious that

$$Q |0\rangle = 0 \quad (7)$$

and that, since $\partial_0 Q = 0$ and $\partial_1 Q = 0$,

$$\partial_+ Q = 0, \quad (8)$$

in conflict with Hagen's published claim that a charge-invariant light-cone vacuum does not exist. In this case the light-cone and ordinary vacuum are identical. We reemphasize that although quantized at equal time, free charged scalar fields satisfy all the assumptions enumerated in Hagen's theorem, and yet contradict his result.

To uncover the source of the discrepancy between this example and Hagen's theorem, it is necessary to examine his calculation in detail. From Eq. (3), for a general field theory,

$$\langle [j^+(x), j^+(0)] \rangle = -4\partial_-^2 \int_0^\infty dM^2 \sigma(M^2) i\Delta(x; M^2). \quad (9)$$

The commutator $\Delta(x; M^2)$ is singular on the light cone,

$$i\Delta(x; M^2) = \lim_{\eta \rightarrow 0^+} \int_0^\infty \frac{dk^+}{2\pi k^+} \left\{ \exp \left[\frac{-i}{2} \left[k^+(x^- - i\eta) + \frac{M^2}{k^+}(x^+ - i\eta) \right] \right] - \exp \left[\frac{i}{2} \left[k^+(x^- + i\eta) + \frac{M^2}{k^+}(x^+ + i\eta) \right] \right] \right\}. \quad (11)$$

Convergence factors are explicitly and necessarily included to ensure that the integral exists, and that manipulations such as the rearrangement of integration and differentiation make sense.

Near the light cone Eq. (11) reduces to

$$i\Delta(x; M^2)_{x^2 \sim 0} = \frac{-i}{4} [\epsilon(x^+) + \epsilon(x^-)] + O(x^2)$$

or equivalently,

$$= \frac{-i}{2} \epsilon(x^0) \Theta(x^2) + O(x^2). \quad (12)$$

so the evaluation of Eq. (9) and related commutators at $x^+ = 0$ may depend on the procedure employed. Hagen formally shows $\langle [\partial_+ Q, j^+(0)] \rangle$ to be nonzero by first differentiating Eq. (9) with respect to x^+ and performing one x^- derivative, then setting x^+ identically to zero and replacing $\Delta(x; M^2)$ with the standard equal- x^+ commutator. Finally, he performs the second x^- derivative and integrates over x^- .

Several features of Hagen's procedure imply that it produces results which are unreliable. First, it gives zero for $\langle [Q(x^+ = 0), j^+(0)] \rangle$, a possibly finite number for $\langle [\partial_+ Q(x^+ = 0), j^+(0)] \rangle$, and divergences for higher derivatives in x^+ . However, as Q is conserved, it should have no x^+ dependence, and there is no good reason to set x^+ to zero at an intermediate step. Had it been left arbitrary, Hagen would have obtained the conventional result that $\langle [Q(x^+), j^+(0)] \rangle$ is zero for *all* x^+ . Higher derivatives in x^+ computed at this stage would then trivially give zero for all x^+ . Also, it can be seen from Eq. (9) that these quantities all involve an integral over a total derivative in x^- . That Hagen does not obtain zero for this, and that his result depends on a particular sequence of integration and differentiation, suggests that he is not working with well-defined integrals. That this is the case may be seen by carefully defining the commutator in Eq. (9). The formal representation in two dimensions,

$$i\Delta(x; M^2) = \int \frac{d^2 k}{2\pi} \delta(k^2 - M^2) \epsilon(k^0) \exp(-ik \cdot x), \quad (10)$$

is, in light-cone variables,

The damping prescription in Eq. (11) uniquely defines $\epsilon(y)$ and $\Theta(y)$ in Eq. (12) to be zero and one-half, respectively, when y is identically zero. It therefore reproduces the standard equal- x^+ commutator

$$i\Delta(x; M^2)_{x^+ = 0} = \frac{-i}{4} \epsilon(x^-). \quad (13)$$

Substituting Eq. (11) into Eq. (9) yields, after appropriate differentiation and integration,

$$\langle [\partial_+ Q(x^+), j^+(0)] \rangle = \lim_{\eta \rightarrow 0^+} \frac{-i}{4} \int_0^\infty dM^2 M^2 \sigma(M^2) \int_{-\infty}^\infty dk^+ \delta(k^+) \exp \left[\frac{-i}{2} \frac{M^2}{k^+} x^+ - \frac{M^2 \eta}{2 |k^+|} \right]. \quad (14)$$

[If desired, a limiting procedure could also be introduced to explicitly define $\delta(k^+)$, with the same conclusion.] As long as all the steps leading to this were well defined, that is, as long as η was kept infinitesimal but not zero, then Eq. (14) and all higher derivatives in x^+ are unambiguously zero independently of x^+ . This is consistent with an invariant vacuum as illustrated in our initial free scalar field example. Hagen's result is obtained if both x^+ and η are set to zero identically, but then his intermediate manipulations are apparently suspect.

Finally, if one insists on employing the commutator at equal- x^+ explicitly, Eq. (11) with x^+ at zero has been

shown to provide an adequate definition of the scalar light-cone commutator, and it produces consistent results as long as η is taken to zero only at the end.⁷

Computations involving singular quantities in a field theory are always potentially ambiguous. In general, one must define such quantities by a prescription, when one exists, consistent with the desired properties of that theory. That at least one prescription does exist for the light-cone commutators in question consistent with the charge and Lorentz invariance of the vacuum has been demonstrated above.

¹H. C. Pauli and S. J. Brodsky, Phys. Rev. D **32**, 1993 (1985); **32**, 2001 (1985).

²T. Eller, H. C. Pauli, and S. J. Brodsky, Phys. Rev. D **35**, 1493 (1987).

³K. Hornbostel (in preparation).

⁴C. R. Hagen, preceding Comment, Phys. Rev. D **37**, 2362 (1988).

⁵Light-cone quantization and the various pitfalls are discussed in detail in the literature. See, for example, P. J. Steinhardt, Ann. Phys. **128**, 425 (1980), and references therein. Noncovariant gauges are reviewed by G. Leibbrandt, Rev. Mod.

Phys. **59**, 1067 (1987).

⁶C. R. Hagen, Phys. Lett. **90B**, 405 (1980); Phys. Rev. D **24**, 2605 (1981).

⁷Alternatively, the equal- x^+ commutator may be defined in analogy to the Johnson-Low (equal-time) prescription [cf. K. Johnson and F. E. Low, Prog. Theor. Phys. Suppl. **37-38**, 74 (1966)], by subtracting the x^+ -ordered product at $-\eta$ from that at $+\eta$ as $\eta \rightarrow 0^+$. This also produces the standard commutator, Eq. (1), and in this case it is the Feynman $i\epsilon$ prescription in the propagator which serves to regulate the small- k^+ region in Eq. (14).