

Signals of unconventional E_6 models at e^+e^- colliders

Enrico Nardi*

Randall Laboratory of Physics, University of Michigan, Ann Arbor, Michigan 48109-1120

(Received 10 September 1993)

Generation-dependent discrete symmetries often appear in models derived from superstring theories. In particular, in the framework of E_6 models the presence of such symmetries is required in order to allow for the radiative generation of naturally small neutrino masses. Recently it was shown that by imposing suitable generation-dependent discrete symmetries, a class of models can be consistently constructed in which the three sets of known fermions in each generation do not have the same assignments with respect to the 27 representation of E_6 . In this scenario, the different embedding in the gauge group of the three generations implies in particular that the known charged leptons couple in a nonuniversal way to the new neutral gauge bosons (Z_β) present in these models. We exploit this fact to study the signature of this class of models at present and future e^+e^- colliders. We show that some signals of deviation from lepton universality as well as some other discrepancies with the standard model predictions which have been observed at the KEK TRISTAN collider in the production rate of μ and τ , can be accounted for if the Z_β mass is not much heavier than 300 GeV. We also study the discovery limits for lepton universality violation of this type at CERN LEP 2 and at the 500 GeV e^+e^- Next Linear Collider (NLC). We show that models predicting unconventional assignments for the leptons will give an unmistakable signature, when the Z_β mass is as heavy as ~ 800 GeV (LEP 2) and ~ 2 TeV (NLC).

PACS number(s): 12.10.Dm, 12.15.Ff, 13.10.+q, 14.60.-z

I. INTRODUCTION

During the past few years, extended electroweak models based on the group E_6 have been thoroughly studied, since they were found to lead to interesting phenomenology. In particular, all the possible embeddings of the standard model (SM) gauge group in E_6 , as well as various possibilities for the assignments of the matter fields to the fundamental representation of the group, were analyzed in Ref. [1]. Recently it has been shown [2] that in the framework of superstring-derived E_6 models, it is possible to implement an unconventional scenario in which some of the known fermions of the three families are embedded in the fundamental 27 representation of the group in a generation-dependent way, meaning that their gauge quantum numbers do not replicate throughout the three generations. It was also argued [2] that if E_6 models are required to allow for small neutrino masses (as is needed for any particle physics solution of the solar neutrino problem, and/or for explaining the atmospheric ν_μ deficit via ν_μ oscillations) the unconventional assignments (UA) scenario should be considered as a natural alternative to the standard schemes. In fact, in the framework of superstring-inspired E_6 theories the most natural way for generating small neutrino masses is through radiative corrections [3,4] since in these models the Higgs representation necessary to implement a seesaw mechanism [5]

is absent.¹ In order to implement the generation of ν masses at the loop level, a suitable discrete symmetry must be imposed on the superpotential to ensure that at the lowest order $m_\nu=0$. Branco and Geng [7] have shown that no generation-blind symmetry exists that forbids nonvanishing neutrino masses at the tree level and at the same time allows for their radiative generation. As a result, in order to implement in a consistent way the generation of ν masses via loop diagrams, a symmetry that does not act in the same way on the three generations is needed.

The main motivation for investigating the UA schemes stems from the observation that once we choose to build a model based on a symmetry that distinguishes among the different generations, there is no reason in principle to expect that this symmetry will result in a set of *light* fermions (i.e., the known states) that will exactly replicate throughout the three generations [2]. This implies the possibility that the known states belonging to different generations could have different E_6 gauge interactions. Of course, experimentally we know that the $SU(2)\times U(1)$ gauge interactions of the known fermions do respect universality with a high degree of precision. However, since the SM gauge group is rank 4, while E_6 is rank 6, as many as two additional massive neutral gauge bosons (Z_β) can be present, possibly with $M_\beta\sim 1$ TeV or less,

*Electronic address: nardi@umiphys.bitnet

¹See, however, Ref. [6] for a discussion of a seesaw mechanism induced by gravitational effects.

and the possibility that the additional $U_{\beta}(1)$ interactions could violate universality is still phenomenologically viable.

Since the fundamental representation of E_6 is 27 dimensional, the fermion content of models based on this group is enlarged with respect to the SM. In fact, in addition to the standard fermions, two additional leptonic $SU(2)$ doublets, two $SU(2)$ singlet neutral states, and two color-triplet $SU(2)$ singlet d -type quarks are present. UA models are realized by identifying in a generation-dependent way some of the known doublets of lepton and/or d -type singlet quarks with the additional fermion multiplets [2]. Models based on the UA scenario can be implemented without conflicting with phenomenological or theoretical constraints. For example, the model described in Ref. [2] was shown to be consistent with a large number of experimental constraints, ranging from the direct and cosmological limits on the neutrino masses, to the stringent limits on flavor-changing neutral currents (FCNC's). In this model the left- (L -) handed lepton doublet " $(\begin{smallmatrix} \nu \\ \tau \end{smallmatrix})_L$ " and the right- (R -) handed quark singlet b_R of the third generation are assigned to $SU(2)$ multiplets having a different embedding in E_6 with respect to the corresponding states of the first two generations. The nonstandard phenomenology resulting from this model implies in particular that the " τ " neutrinos have different neutral current (NC) interactions than do " ν_e " and " ν_μ ".

In Sec. II we will briefly outline the main features of the E_6 models based on the UA scenario, and establish our conventions and notation. A more complete discussion of the theoretical framework can be found in Ref. [2]. A very clean signature for the UA models would be the detection of deviations from universality in neutral current (NC) processes. Because of the clean experimental environment of e^+e^- annihilation, such a signature could be more easily detected at e^+e^- colliders rather than at hadron colliders. In Sec. III we will investigate the phenomenology of UA models at the present and future e^+e^- machines. Since the UA for the known fermions would result in a violation of universality only in the fermion couplings to the new Z_β bosons without affecting the couplings to the standard Z_0 , the large amount of data collected at the Z_0 resonance by the CERN Large Electron Positron (LEP) Collaborations are not effective for the search of these kind of effects. In fact, the contribution of Z_β - Z_0 interference to the various cross sections and asymmetries measured at LEP 1 is

too small to be measured at the peak, and the contribution of pure Z_β exchange is also vanishingly small. Some effects could still be detected if the Z_0 had a sizable mixing with Z_β ; however, the existing bounds on the Z_0 - Z_β mixing angle are extremely tight [8], so that we will disregard this possibility throughout this paper.

Among presently operating colliders, the one best suited to reveal the kind of effects we are looking for is the TRISTAN collider at the KEK laboratories, which is collecting data at about 60 GeV c.m. energy. It is intriguing, though certainly not compelling, that a few discrepancies between the TRISTAN data on the total hadronic and leptonic cross sections and the SM predictions do exist [9], and they point toward the existence of a Z_β at rather low energies $M_\beta \lesssim 300$ GeV [10]. At the same time the data on the rate of production of τ and μ leptons do show a signal of violation of universality (at the level of 1.6 standard deviations) which could not be explained by conventional extended gauge models. However, as we will show, these data can be well accounted for in the framework of models with UA. We will complete our discussion by analyzing the discovery limits for UA models at LEP 2 and at the 500 GeV Next Linear e^+e^- Collider (NLC). Finally, in Sec. IV we will summarize our results and draw the conclusions.

II. UNCONVENTIONAL ASSIGNMENTS IN E_6 MODELS

In E_6 grand unified theories, matter fields belong to the fundamental 27 representation of the group. E_6 contains $SO(1) \times U_\psi(1)$ as a maximal subalgebra, and the 27 branches to the $1 + 10 + 16$ representation of $SO(10)$. In turn $SU(10)$ contains $SU(5) \times U_\chi(1)$. The $SO(10)$, $SU(5)$, $U_\psi(1)$, and $U_\chi(1)$ assignments for the fermions in the 27 representation are listed in Table I. Usually the known particles of the three generations are assigned to the 16 of $SO(10)$ that also contains an $SU(2)$ singlet neutrino " ν^c ":

$$[16]_i = \left[Q \equiv \begin{pmatrix} u \\ d \end{pmatrix}, u^c, e^c, d^c, L \equiv \begin{pmatrix} \nu \\ e \end{pmatrix}, \nu^c \right]_i, \quad i = 1, 2, 3. \quad (2.1)$$

The 10 and the 1 of $SO(10)$ contain the new fields

$$[10]_i = \left[H^c \equiv \begin{pmatrix} E^c \\ N^c \end{pmatrix}, h, H \equiv \begin{pmatrix} N \\ E \end{pmatrix}, h^c \right]_i, \quad (2.2)$$

$$[1]_i = [S^c]_i, \quad i = 1, 2, 3.$$

TABLE I. $SO(10)$, $U_\psi(1)$, $SU(5)$, and $U_\chi(1)$ assignments for the left-handed fermions of the 27 fundamental representation of E_6 . The $SU(2)$ doublets H^c , H , L , and Q are explicitly written in components. The Abelian charges Q_ψ and Q_χ can be obtained from the quantum numbers in the brackets by dividing by $c_\psi = 6\sqrt{\frac{2}{3}}$ and $c_\chi = 6\sqrt{\frac{2}{3}}$, respectively. The charges are normalized to the hypercharge according to $\sum_{f=1}^{27} (Q_{\psi,\chi}^f)^2 = \sum_{f=1}^{27} (\frac{1}{2} Y^f)^2 = 5$.

	S^c	$(\begin{smallmatrix} E^c \\ N^c \end{smallmatrix}) h$	$(\begin{smallmatrix} N \\ E \end{smallmatrix}) h^c$	ν^c	$(\begin{smallmatrix} \nu \\ e \end{smallmatrix}) d^c$	$e^c u^c (\begin{smallmatrix} u \\ d \end{smallmatrix})$
$SO(10) (c_\psi Q_\psi)$	1 (4)	10 (-2)			16(1)	
$SU(5) (c_\chi Q_\chi)$	1 (0)	5 (2)	$\bar{5}$ (-2)	1 (-5)	$\bar{5}$ (3)	10 (-1)

As is clear from Table I, there is an ambiguity in assigning the known states to the 27 representation, since under the SM gauge group

$$\mathcal{G}_{\text{SM}} \equiv \text{SU}(3)_c \times \text{SU}(2)_L \times \text{U}(1)_Y$$

the $\bar{5}_{(10)}$ in the 10 of SO(10) has the same field content as the $\bar{5}_{(16)}$ in the 16. The same ambiguity is also present for the two \mathcal{G}_{SM} singlets: namely, $1_{(1)}$ and $1_{(16)}$. In the present paper we will concentrate on the consequences of having different assignments for the known L -handed leptons. Since these leptons might not correspond to the entries as listed in Table I, we use quotation marks to denote the known states with their conventional labels, while labels not enclosed within quotation marks will always refer to the fields listed in the table.

In the models under investigation as many as two additional neutral gauge bosons can be present, corresponding, for example, to some linear combinations of the $U_\chi(1)$ and $U_\psi(1)$ generators. The interaction of the fermions in the $\bar{5}$ of SU(5) with these gauge bosons will depend on the specific assignments to the 16 or to the 10 of SO(10). The two additional neutral gauge bosons are usually parametrized as

$$\begin{aligned} Z'_\beta &= Z_\psi \sin\beta + Z_\chi \cos\beta, \\ Z''_\beta &= Z_\psi \cos\beta - Z_\chi \sin\beta, \end{aligned} \quad (2.3)$$

where β is a model-dependent parameter. In the following we will denote the lightest of the two new gauge bosons as Z_β . In the presence of a “light” Z_β different assignments will lead to a different phenomenology. In contrast, in the limit $M_\beta \rightarrow \infty$ the choice of the assignment is irrelevant as long as we are only concerned with the gauge interactions. However, even in this limit the requirement of $U_{\psi,\chi}(1)$ gauge invariance for the superpotential, together with the phenomenological constraints on the absence of FCNC’s in the Higgs sector, strongly constrain the structure of the viable models [2].

A model realizing a UA scenario, in which what we call “ τ_L ” corresponds to the charged component of the H_3 weak doublet belonging to $\bar{5}_{10}$, while the “ e_L ” and the “ μ_L ” leptons are as usual assigned to the $\bar{5}_{16}$, was recently proposed in Ref. [2]. This model is realized by imposing on the superpotential a particular family-nonblind $Z_2 \times Z_3$ discrete symmetry. As a result of such a symmetry, the masses of the known (light) chiral leptons are generated by vacuum expectation values (VEV’s) of Higgs doublets, through the terms $m_\tau E_{3L} e_{3R}$ (with $m_\tau \sim \langle \bar{L}_3 \rangle_0$) and $m_{\alpha\beta} e_{\alpha L} e_{\beta R}$ (with $m_{\alpha\beta} \sim \langle \bar{H}_2 \rangle_0$ and $\alpha, \beta = 1, 2$). The remaining charged leptons e_{3L} , E_{3R} , $E_{\alpha L}$, and $E_{\alpha R}$ are vectorlike, and acquire large masses from VEV’s of Higgs singlets.

As it was argued in Ref. [2], several interesting features of this model are peculiar to the UA schemes in general, independently of this particular realization. For example, in contrast with the conventional E_6 models [3], in the UA schemes rank 6 models are not disfavored with respect to rank 5, so that the general parametrization of the two additional gauge bosons given in (2.3) is well motivated. We stress that other assignments, leading to

models with a structure similar to the model proposed in Ref. [2], but implying a different phenomenology, can be easily obtained by means of some different discrete symmetries.

If some of the ν^c and/or S^c SU(2) neutral singlets are massless or are very light ($m_s \lesssim 1$ MeV), cosmological arguments suggest that the Z_β bosons should be heavier than about ~ 1 –2 TeV [11], thus excluding the possibility of detecting any signal at TRISTAN and LEP 2. In fact, though singlet under \mathcal{G}_{SM} , these states do have $U(1)_\beta$ interactions. Then, not to conflict with the limit of 3.6 relativistic neutrinos in thermal equilibrium at the time of nucleosynthesis [12] (which can be derived from the data on the light element abundances) we have to require this interaction to be weak enough to allow for the decoupling of the light \mathcal{G}_{SM} singlets at a sufficiently early time (for example, before the QCD phase transition) so that their number density can be safely diluted. Requiring the $U_\beta(1)$ interaction to be “superweak” results in the quoted lower bound on the Z_β mass [11]. We would like to mention that there are two models corresponding to the particular values of the angle β in (2.3) ($\tan 2\beta = -\sqrt{15}/7, 0$) in which the nucleosynthesis constraints on M_β can be evaded even in the presence of light SU(2) singlets [2]. In fact, for these two values of β , respectively, the ν^c and the S^c degrees of freedom decouple from the lightest Z_β , behaving as “effective singlets” with respect to all the “light” gauge bosons. Then they could play the role of the helicity partners of the standard neutrinos, allowing in particular for nonzero neutrino Dirac masses, while at the same time their gauge interactions would not be effective to keep them in thermal equilibrium in the early Universe.

However, for the sake of generality, in the present analysis we will assume that all the ν^c and S^c \mathcal{G}_{SM} singlets are heavy ($m_s \gg 1$ MeV). In this case, independently of the value of β , the nucleosynthesis constraints on M_β are evaded. Therefore the Z_β boson could be as light as allowed by the present limits from direct searches at colliders [13] and from the analysis of Z_β indirect effects [8], resulting in both cases in $M_\beta \gtrsim 200$ –300 GeV. As we will see, in the UA schemes the presence of a Z_β with a mass in this range can give rise to lepton universality-violating effects that could be detected at the colliders presently in operation.

III. SIGNALS OF UNCONVENTIONAL ASSIGNMENTS AT e^+e^- COLLIDERS

In the presence of additional neutral gauge bosons, the lowest order cross section² for the process $e^+e^- \rightarrow l^+l^-$ with $l \neq e$ is

$$\sigma(s) = \frac{4}{3} \frac{\pi\alpha^2}{s} \sum_{m,n=0}^N C(m,n) \chi_m(s) \chi_n^*(s), \quad (3.1)$$

²In the numerical computations we have taken into account the leading one-loop corrections by using an improved Born approximation [14].

$$C(m, n) = [v_m(e)v_n^*(e) + a_m(e)a_n^*(e)] \\ \times [v_m(l)v_n^*(l) + a_m(l)a_n^*(l)], \quad (3.2)$$

$$\chi_m(s) = \frac{g_m^2}{4\pi\alpha} \frac{s}{s - M_m^2 - iM_m\Gamma_m}. \quad (3.3)$$

We will henceforth assume that one of the two new bosons in (2.3) is heavy enough so that its effects on the low energy physics are negligible. Then $m, n = 0, 1, 2$ correspond respectively to the γ , Z_0 , and Z_β amplitudes. The couplings in (3.2) and (3.3) are

$$g_0 = e, \quad v_0(l) = Q_{\text{em}}^l, \quad a_0(l) = 0, \\ g_1 = (\sqrt{2}G_\mu M_Z^2)^{1/2}, \quad v_1(l) = T_{3L}^l - 2Q_{\text{em}}^l s_W^2, \\ a_1(l) = T_{3L}^l, \quad (3.4) \\ g_2 = s_W g_1, \quad v_2(l) = Q_\beta^l - Q_\beta^{l,c}, \\ a_2(l) = Q_\beta^l + Q_\beta^{l,c}, \quad l = e, \mu, \tau,$$

where $Q_{\text{em}}^l = -1$ is the electric charge of the leptons, $T_{3L}^l = -\frac{1}{2}$ is the third component of the weak isospin, $s_W \equiv \sin\theta_W$ with θ_W the weak mixing angle, and $Q_\beta^{l,c} = Q_\psi^{l,c} \sin\beta + Q_\chi^{l,c} \cos\beta$ is the lepton coupling to the Z_β boson in (2.3). The new charges $Q_\psi^{l,c}, Q_\chi^{l,c}$ are given in Table I, and are normalized to the hypercharge generator $Y/2$. In addition, in (3.4) we have assumed for the Abelian coupling g_2 a renormalization group evolution down to the electroweak scale similar to that of the hypercharge coupling $g_Y \simeq s_W g_1$.

In (3.4) the vector and axial-vector couplings $v_{0,1}(l)$ and $a_{0,1}(l)$ do not depend on the specific assignments for the leptons, and are unmodified with respect to the SM. In contrast, $v_2(l)$ and $a_2(l)$ depend on the particular assignments of the l lepton. With the notation given in (2.1) and (2.2), and referring to the 16 and to the 10 representations of SO(10), the possible assignments for the L -handed $l_1 = "e_L", l_2 = "\mu_L, l_3 = "\tau_L$ charged leptons are

$$"l_i" \in L_i \in 16,$$

or

$$"l_i" \in H_i \in 10, \quad i = 1, 2, 3. \quad (3.5)$$

With these assignments three different cross sections for the process $e^+e^- \rightarrow l^+l^-$ ($l \neq e$) are possible. They are $\sigma_{16 \rightarrow 16}, \sigma_{16 \rightarrow 10} = \sigma_{10 \rightarrow 16},$ and $\sigma_{10 \rightarrow 10}$, where the subscripts refer to the specific embedding of the L -handed components of the initial e^- and final l^- states in the 16 or in the 10 of SO(10). In the following we will give results for the quantities $R_{16 \rightarrow 16}, R_{16 \rightarrow 10},$ and $R_{10 \rightarrow 10}$ corresponding to the different cross sections normalized to the point-like QED cross section for muon pair production.

In Fig. 1 we compare the theoretical values for $R_{16 \rightarrow 16}, R_{16 \rightarrow 10},$ and $R_{10 \rightarrow 10}$ at $\sqrt{s} = 58$ GeV c.m. energy, with the SM prediction $R_{ll}^{\text{SM}} = 1.053$ (heavy solid line), and

with the TRISTAN experimental data $R_{\tau\tau} = 1.026 \pm 0.037$ and $R_{\mu\mu} = 0.982 \pm 0.036$. These figures have been obtained by combining the results of the AMY, TOPAZ, and VENUS Collaborations given in Ref. [9]. In deriving the averages we have assigned a common systematic error of ± 0.030 for the uncertainty in the luminosity. Figure 1 shows that the measured values of $R_{\tau\tau}$ and $R_{\mu\mu}$ are both lower than the SM prediction. However, while the value of $R_{\tau\tau}$ is consistent with the SM within one standard deviation, $R_{\mu\mu}$ is about two standard deviations off the expected value. The shaded areas show the predictions for the three ratios $R_{16 \rightarrow 16}, R_{16 \rightarrow 10},$ and $R_{10 \rightarrow 10}$ for a Z_β mass ranging between 200 and 300 GeV.

This range corresponds to the range of the *most conservative* lower bounds on M_β , as obtained in the framework of conventional E_6 models. Direct bounds have been derived at the Fermilab Tevatron from the combined limits on the two processes $p\bar{p} \rightarrow Z_\beta \rightarrow e^+e^-, \mu^+\mu^-$ [13]. For the models usually referred to as $\psi, \eta,$ and χ which correspond to the particular values $\sin\beta = -\sqrt{5}/8, 0,$ and 1 [15], the published direct bounds are respectively 200, 230, and 280 GeV [13].³

Other indirect limits have been obtained from high precision electroweak data by analyzing the Z_β indirect effects on NC observables. The indirect bounds also suggest $M_\beta \gtrsim 200$ GeV for all the values of the parameter β [8].

Since all these limits have been derived under the assumption that both e and μ are embedded in the same way in the group, they cannot be straightforwardly applied to the Z_β of UA models. For example, it is clear that in the present case the e and μ gauge couplings can be different. However we have no reason to expect that by assuming UA the bounds could be greatly relaxed (or strengthened), and hence we regard our choice of the conservative lower limits on M_β as a safe one.

From Fig. 1, it is apparent that in the presence of a light Z_β , the experimental data on $R_{\mu\mu}$ would be better accounted for by either assigning both " e_L " and " μ_L " to the 16 representation of SO(10) (as in conventional E_6 models), and for values of $\sin\beta$ centered around zero, or by assigning both these leptons to the 10 and for $\sin\beta$ close to unity. At the same time, for any choice of the assignments and for any value of β the various R are in good agreement with the experimental value of $R_{\tau\tau}$. Only a small region in the vicinity of $\sin\beta \sim 0.6$ is slightly

³The quoted bounds are very conservative since they correspond to the assumption that the Z_β decays to *all the fermions and supersymmetric fermions* present in the model, and that all phase space suppression factors can be neglected. In the case that *only decays to the SM fermions* are allowed, the present bounds are of order 300 GeV or slightly higher. We also stress that the quoted limits correspond to an integrated luminosity of about 4 pb^{-1} . In the near future, the limits for the already accumulated luminosity of 25 pb^{-1} will be able to rule out the UA models as a possible explanation of the TRISTAN data.

disfavored if the assignments “ e_L ” \in 16 and “ τ_L ” \in 10 are chosen.

One of the most spectacular signals of UA models would be a deviation of the ratio $\rho_{\mu/\tau} \equiv \sigma(e^+e^- \rightarrow \mu^+\mu^-)/\sigma(e^+e^- \rightarrow \tau^+\tau^-)$ from unity. Since many systematic errors, as for example the uncertainties in the luminosity measurements, cancel in this ratio, the experimental error is statistically dominated, implying a very transparent significance for such a measurement. At the Z_0 resonance, the measured value for this observable $\rho_{\mu/\tau} = 0.998 \pm 0.006$ [16] is in striking agreement with μ - τ universality. Undoubtedly it would be difficult to accommodate any large deviation from unity at $\sqrt{s} \neq M_Z$ by means of some mechanism different from the one discussed here. For this reason we believe that if a value $\rho_{\mu/\tau} \neq 1$ is measured off Z_0 resonance, this would represent a very clean and almost unmistakable signature of the UA models.

According to the assignments in (3.5), and without




FIG. 1. TRISTAN total cross sections for μ pair production ($R_{\mu\mu}$) and τ pair production ($R_{\tau\tau}$), normalized to the pointlike QED cross section, compared to the standard model prediction R_{ii}^{SM} . The dotted and dashed lines correspond to the one standard deviation experimental errors for $R_{\mu\mu}$ and $R_{\tau\tau}$, respectively. The shaded areas, enclosed between the solid lines ($M_\beta = 200$ GeV) and the dot-dashed lines ($M_\beta = 300$ GeV), depict the predictions for lepton pair production at the TRISTAN c.m. energy $\sqrt{s} = 58$ GeV in the unconventional assignments E_6 models. The results are given for a general Z_β from E_6 , as a function of $\sin\beta$. $R_{16 \rightarrow 16}$ refers to the case in which the L components of both the initial and final leptons are assigned to the 16 representation of $SO(10)$, and similarly for $R_{16 \rightarrow 10}$ and $R_{10 \rightarrow 10}$.

referring to any specific lepton flavor, we can write two expressions for this ratio which deviate from unity, $\rho_{16} \equiv \sigma_{16 \rightarrow 16}/\sigma_{16 \rightarrow 10}$ and $\rho_{10} \equiv \sigma_{10 \rightarrow 10}/\sigma_{10 \rightarrow 16}$, where the subscripts label the assignment for the L electron in the initial state. Figure 2 shows the two ratios ρ_{16} and ρ_{10} compared to the combined TRISTAN measurement $\rho_{\mu/\tau} = 0.957 \pm 0.027$. This value is about 1.6 standard deviations off the value of unity predicted by any model which assumes lepton universality. Again it is apparent that the experimental data can be better accounted for by taking

$$"e_L", " \mu_L" \in 16, " \tau_L" \in 10, \text{ and } -0.5 \lesssim \sin\beta \lesssim 0.8, \quad (3.6)$$

or

$$"e_L", " \mu_L" \in 10, " \tau_L" \in 16, \text{ and } \sin\beta \gtrsim 0.4, \quad (3.7)$$

and assuming $M_\beta \lesssim 300$ GeV. We note that the set of as-




FIG. 2. TRISTAN result for the ratio of μ to τ pair productions $(\sigma_{\mu\mu}/\sigma_{\tau\tau})_{\text{expt}}$ compared to the standard model prediction $(\sigma_{\mu\mu}/\sigma_{\tau\tau})_{\text{SM}} = 1$. The dotted lines correspond to the one standard deviation experimental error. The shaded areas, enclosed between the solid lines ($M_\beta = 200$ GeV) and the dot-dashed lines ($M_\beta = 300$ GeV) depict the predictions for the ratio of pair productions of two different lepton flavors at $\sqrt{s} = 58$ GeV, in the unconventional assignments E_6 models. The results are given for a general Z_β from E_6 , as a function of $\sin\beta$. $\sigma_{16 \rightarrow 16}$ refers to the case in which the L components of both the initial and final leptons are assigned to the 16 representation of $SO(10)$, and similarly for $\sigma_{16 \rightarrow 10} = \sigma_{10 \rightarrow 16}$ and $\sigma_{10 \rightarrow 10}$.

signments in (3.6) coincides with the assignments in the model discussed in Ref. [2].

For the particular value $\sin\beta = -\sqrt{\frac{5}{8}}$, which corresponds to the rank 5 η model [15], the cross section (3.1) is invariant with respect to the different choices of the assignments. This is apparent from Fig. 1, and in particular Fig. 2 shows that in this case lepton universality is preserved. This follows from the fact that for this value of β , the Q_η charges for the $\bar{5}_{(10)}$ and for the $\bar{5}_{(16)}$ are equal [15,17] [this is true also for $Q_\eta(1_{(1)})$ and $Q_\eta(1_{(16)})$] implying that for all the leptons the couplings to the Z_η are the same independently of the UA.

In Fig. 3 we depict the theoretical values of ρ_{16} and ρ_{10} for M_β in the range 600–800 GeV, and for $\sqrt{s} = 190$ GeV, corresponding to the c.m. energy at LEP 2. The dotted lines depict the one standard deviation statistical error achievable with 500 pb^{-1} of integrated luminosity corresponding to a one year run [18] ($\sim 3 \times 10^3$ leptonic events per flavor). It is apparent that the signature of UA models could be easily recognized for Z_β bosons corresponding to most of the $\sin\beta$ positive values.




FIG. 3. Predictions for the ratio of pair productions of two different lepton flavors in the unconventional assignments E_6 model at the LEP 2 c.m. energy $\sqrt{s} = 190$ GeV (shaded areas), compared to the standard model prediction (thick solid line). The dotted lines depict the expected one standard deviation experimental error, based on an integrated luminosity of 500 pb^{-1} . The solid lines enclosing the shaded areas correspond to $M_\beta = 600$ GeV while the dot-dashed lines correspond to $M_\beta = 800$ GeV. The results are given for a general Z_β from E_6 , as a function of $\sin\beta$.

Figure 4 depicts the ratios ρ_{16} and ρ_{10} for M_β in the range 1500–2200 GeV, and for $\sqrt{s} = 500$ GeV, corresponding to the NLC c.m. energy. The one standard deviation error corresponds to a statistics of $\sim 8.6 \times 10^3$ leptonic events per flavor, based on an integrated luminosity of 20 fb^{-1} (one year run) and taking into account the efficiency for a cut to suppress the two-photon background [19]. The violation of lepton universality, intrinsic in the UA models, would produce striking effects for a Z_β as heavy as ~ 2 TeV and for values of $\sin\beta$ not too close to the η model.

We have seen that by measuring the various quantities R_{ll} and $\rho_{l/l'}$, it will be easy to detect the effects of the UA. However, from Figs. 1–4 it is also apparent that these observables alone would not be sufficient to determine the exact pattern of lepton assignments, since different assignments can account for the same set of experimental data by means of a different choice of the β parameter. Working out a procedure for identifying unequivocally the correct pattern of assignments is beyond the scope of the present analysis; however, we believe that once signals of UA are detected in the leptonic cross sections, a measurement of the various leptonic asymmetries would be quite effective to achieve this result. Also, we would like to point out that if some mechanism resulting in generation-dependent assignments for the




FIG. 4. Same as Fig. 3, at the NLC c.m. energy $\sqrt{s} = 500$ GeV, for $M_\beta = 1500$ GeV (solid lines) and $M_\beta = 2200$ GeV (dot-dashed lines). The dotted lines depict the expected one standard deviation experimental error, based on an integrated luminosity of 20 fb^{-1} and including an efficiency cut.

lepton doublets is effective, quite naturally the same mechanism would imply UA for the d -type SU(2) singlet quarks as well. For example, in the model discussed in Ref. [2], the assignments " e_L ," " μ_L " \in 16, and " τ_L " \in 10 did imply, for the consistency of the model, the UA " d_L^c ," " s_L^c " \in 10, and " b_L^c " \in 16 in the quark sector. Clearly, due to the experimental difficulties in tagging the quark flavors, identifying UA for the d quarks would be a much harder task.

IV. CONCLUSIONS

In conclusion, we have examined the possibility of detecting with the present and future e^+e^- colliders, signals of models predicting UA for the charged leptons. We have shown that a class of models based on the gauge group E_6 , in which the known SU(2) lepton doublets are embedded in the fundamental representation of the group in a generation-dependent way, would result in a unique type of violation of lepton universality, which is induced by the exchange of new Z_β bosons. In agreement with LEP 1 data, no observable effects are predicted at the Z_0 resonance; however, some signals could be detected off Z_0 resonance. For example, we have shown that a few anomalies in the production rate of leptons, as well a hint of violation of μ - τ universality which have been observed at the TRISTAN collider, could be well accounted for in the UA scenarios. As we have discussed, though these anomalies are not statistically compelling, it will not be

easy to find an alternative particle physics mechanism that could simultaneously account for the LEP 1 and the TRISTAN observations. However, the mechanism proposed here would be effective only if the Z_β mass is not much heavier than ~ 300 GeV. Though this value is still consistent with the direct limits from the $p\bar{p}$ collider [13], in the near future the data obtainable at Tevatron will be able to probe or rule out such an explanation of the leptonic cross section anomalies [20].

We have also discussed the discovery potential for this class of models at LEP 2, operating at 190 GeV c.m. energy, and at the NLC, operating at 500 GeV c.m. energy. We have shown that at these future colliders, striking effects of lepton universality violation resulting from the various UA could be observed up to $M_\beta \sim 800$ GeV (LEP 2) and $M_\beta \sim 2200$ GeV (NLC) for most of the values of the model-dependent parameter β .

ACKNOWLEDGMENTS

It is a pleasure to thank the Theory Group at KEK and the Particle Physics Group at the Yukawa Institute in Kyoto for their kind hospitality during the initial stage of this work. I also thank K. Hagiwara, G. Kane, and T. G. Rizzo for some useful discussions, and J. F. Dodge, I. Rothstein, and D. Tommasini for a critical reading of the manuscript.

-
- [1] M. Dine *et al.*, Nucl. Phys. **B259**, 549 (1985); F. del Aguila, J. A. Gonzales, and M. Quiros, *ibid.* **B307**, 571 (1988).
- [2] E. Nardi, Phys. Rev. D **48**, 3277 (1993).
- [3] B. A. Campbell *et al.*, Int. J. Mod. Phys. A **2**, 831 (1987).
- [4] A. Masiero, D. V. Nanopoulos, and A. I. Sanda, Phys. Rev. Lett. **57**, 663 (1986).
- [5] M. Gell-Mann, P. Ramond, and R. Slansky, in *Supergravity*, proceedings of the Workshop, Stony Brook, New York, 1979, edited by F. van Nieuwenhuizen and D. Freedman (North-Holland, Amsterdam, 1979), p. 315; T. Yanagida, in *Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe*, Tsukuba, Japan, 1979, edited by A. Sawada and A. Sugamoto (KEK Report No. 79-18, Tsukuba, 1979).
- [6] S. Nandi and U. Sarkar, Phys. Rev. Lett. **56**, 564 (1986); M. Cvetič and P. Langacker, Phys. Rev. D **46**, R2759 (1992).
- [7] G. C. Branco and C. Q. Geng, Phys. Rev. Lett. **58**, 969 (1986).
- [8] E. Nardi, E. Roulet, and D. Tommasini, Phys. Rev. D **46**, 3040 (1992); P. Langacker and M. Luo, *ibid.* **45**, 278 (1992); F. del Aguila, W. Hollik, J. M. Moreno, and M. Quirós, Nucl. Phys. **B372**, 3 (1992); J. Layssac, F. M. Renard, and C. Verzegnassi, Z. Phys. C **53**, 97 (1992); M. C. Gonzalez García and J. W. F. Valle, Phys. Lett. B **259**, 365 (1991); G. Altarelli *et al.*, *ibid.* **263**, 459 (1991).
- [9] AMY Collaboration, T. Kumita *et al.*, in *Proceedings of the Workshop on TRISTAN Physics at High Luminosities*, KEK, Tsukuba, Japan, 1992, edited by M. Yamauchi (KEK Proceedings No. 93-2), Tsukuba (1993); TOPAZ Collaboration, B. L. Howell *et al.*, *ibid.*; VENUS Collaboration, T. Sumiyoshi *et al.*, *ibid.*; see also A. Maki, in *Colmar 1991, Proceedings of the 11th International Conference on Physics in Collision*, Colmar, France, 1991 (KEK Report No. 91-100, Tsukuba, 1991).
- [10] K. Hagiwara, R. Najima, M. Sakuda, and T. Terunuma, Phys. Rev. D **41**, 815 (1990).
- [11] M. C. Gonzales-Garcia and J. W. Valle, Phys. Lett. B **240**, 163 (1990); J. L. Lopez and D. V. Nanopoulos, *ibid.* **241**, 392 (1990); A. E. Faraggi and D. V. Nanopoulos, Mod. Phys. Lett. A **6**, 61 (1991).
- [12] T. Walker, in *Texas/PASCOS 92: Relativistic Astrophysics and Particle Cosmology*, proceedings of the 16th Texas Symposium on Relativistic Astrophysics and 3rd Particles, Strings, and Cosmology Symposium, Berkeley, CA, 1992, edited by C. W. Akerlof and M. A. Srednicki [N.Y. Acad. Sci. **688**, 745 (1993)].
- [13] CDF Collaboration, F. Abe *et al.*, Phys. Rev. Lett. **67**, 2609 (1991); **68**, 1463 (1992).
- [14] M. Consoli and W. Hollik, in *Z Physics at LEP 1*, Proceedings of the Workshop, Geneva, Switzerland, 1989, edited by G. Altarelli *et al.* (CERN Report No. 89-08, Geneva, 1989), Vol. 1, p. 7; G. Burgers and F. Jegerlehner, *ibid.* p. 55; G. Burgers and W. Hollik, in *Polarization at LEP*, edited by G. Alexander *et al.* (CERN Report No. 88-06, Geneva, Switzerland, 1989), Vol. 1, p. 136; D. C. Kennedy and B. W. Lynn, Nucl. Phys. B **322**, 1 (1989).
- [15] See J. L. Hewett and T. G. Rizzo, Phys. Rep. **183**, 195 (1989), and references therein.
- [16] See A. Gurtu, presented at 10th DAE Symposium on High

- Energy Physics, Bombay, India, 1992 [Report No. TIFR/EHEP 93-1 (unpublished)].
- [17] E. Nardi, *Phys. Rev. D* **48**, 1240 (1993).
- [18] D. H. Perkins, in *Proceedings of the ECFA Workshop on LEP 200, Aachen, Germany, 1986*, edited by A. Böhm and W. Hoogland (CERN Report No. 87-08, Geneva, Switzerland, 1987), p. 1.
- [19] A. Djouadi *et al.*, in *e^+e^- Collisions at 500 GeV: The Physics Potential*, Proceedings of the Workshop, Munich, Annecy, Hamburg, 1991, edited by P. M. Zerwas (DESY Report No. 92-123B, Hamburg, Germany, 1992), p. 491.
- [20] F. del Aguila, M. Cvetič, and P. Langacker, in *Physics and Experiments with Linear e^+e^- Colliders*, Proceedings of the Workshop, Waikoloa, Hawaii, 1993, edited by F. A. Harris, S. L. Olsen, S. Pakvasa, and X. Tata (World Scientific, Singapore, 1993), Vol. 2, p. 430.




FIG. 1. TRISTAN total cross sections for μ pair production ($R_{\mu\mu}$) and τ pair production ($R_{\tau\tau}$), normalized to the pointlike QED cross section, compared to the standard model prediction R_{ii}^{SM} . The dotted and dashed lines correspond to the one standard deviation experimental errors for $R_{\mu\mu}$ and $R_{\tau\tau}$, respectively. The shaded areas, enclosed between the solid lines ($M_\beta = 200$ GeV) and the dot-dashed lines ($M_\beta = 300$ GeV), depict the predictions for lepton pair production at the TRISTAN c.m. energy $\sqrt{s} = 58$ GeV in the unconventional assignments E_6 models. The results are given for a general Z_β from E_6 , as a function of $\sin \beta$. $R_{16 \rightarrow 16}$ refers to the case in which the L components of both the initial and final leptons are assigned to the 16 representation of $SO(10)$, and similarly for $R_{16 \rightarrow 10}$ and $R_{10 \rightarrow 10}$.




FIG. 2. TRISTAN result for the ratio of μ to τ pair productions $(\sigma_{\mu\mu}/\sigma_{\tau\tau})_{\text{expt}}$ compared to the standard model prediction $(\sigma_{\mu\mu}/\sigma_{\tau\tau})_{\text{SM}}=1$. The dotted lines correspond to the one standard deviation experimental error. The shaded areas, enclosed between the solid lines ($M_\beta=200$ GeV) and the dot-dashed lines ($M_\beta=300$ GeV) depict the predictions for the ratio of pair productions of two different lepton flavors at $\sqrt{s}=58$ GeV, in the unconventional assignments E_6 models. The results are given for a general Z_β from E_6 , as a function of $\sin\beta$. $\sigma_{16\rightarrow 16}$ refers to the case in which the L components of both the initial and final leptons are assigned to the 16 representation of $SO(10)$, and similarly for $\sigma_{16\rightarrow 10}=\sigma_{10\rightarrow 16}$ and $\sigma_{10\rightarrow 10}$.




FIG. 4. Same as Fig. 3, at the NLC c.m. energy $\sqrt{s} = 500$ GeV, for $M_\beta = 1500$ GeV (solid lines) and $M_\beta = 2200$ GeV (dot-dashed lines). The dotted lines depict the expected one standard deviation experimental error, based on an integrated luminosity of 20 fb^{-1} and including an efficiency cut.