

REVIEW OF PARTICLE PROPERTIES[†]

Particle Data Group

L. Montanet and Technical Associate: K. Gieselmann

CERN, European Laboratory for Particle Physics, CH-1211 Genève 23, Switzerland

R.M. Barnett, D.E. Groom, T.G. Trippe, and C.G. Wohl

Technical Associates: B. Armstrong and G.S. Wagman

Physics Division, Lawrence Berkeley Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA

H. Murayama

*Physics Division, Lawrence Berkeley Laboratory, 1 Cyclotron Road, Berkeley, CA 94720, USA;
and Department of Physics, Tohoku University, Sendai 980, Japan*

J. Stone

Department of Physics, Boston University, Boston, MA 02215, USA

J.J. Hernandez

IFIC — Instituto de Física Corpuscular, Universitat de València — C.S.I.C., E-46100 Burjassot, València, Spain; and CERN

F.C. Porter

Department of Physics, California Institute of Technology, Pasadena, CA 91125, USA

R.J. Morrison

Department of Physics, University of California at Santa Barbara, Santa Barbara, CA 93106, USA

A. Manohar

Department of Physics, University of California at San Diego, La Jolla, CA 92093, USA

M. Aguilar-Benitez

C.I.E.M.A.T., E-28040, Madrid, Spain; and CERN

C. Caso and Technical Associate: P. Lantero

Dipartimento di Fisica e INFN, Università di Genova, I-16146 Genova, Italy

R.L. Crawford

Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, Scotland

M. Roos

*Physics Department, PB 9, FIN-00014, University of Helsinki, Finland;
and Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, I-50125 Firenze, Italy*

N.A. Törnqvist

Research Institute of High Energy Physics, PB 9, FIN-00014, University of Helsinki, Finland; and CERN

K.G. Hayes

Department of Physics, Hillsdale College, Hillsdale, MI 49242, USA

G. Höhler

Institut für Theoretische Teilchenphysik, Universität Karlsruhe, Postfach 6980, 76128 Karlsruhe, Germany

S. Kawabata

KEK, National Laboratory for High Energy Physics, Oho, Tsukuba-shi, Ibaraki-ken 305, Japan

D.M. Manley

Department of Physics, Kent State University, Kent, OH 44242, USA

K. Olive

School of Physics and Astronomy, University of Minnesota, Minneapolis, MN 55455, USA

R.E. Shrock

Institute for Theoretical Physics, State University of New York, Stony Brook, NY 11794, USA

S. Eidelman

Institute of Nuclear Physics, SU-630090, Novosibirsk, Russia, USSR

R.H. Schindler

Stanford Linear Accelerator Center, Stanford, CA 94309, USA

A. Gurtu

Tata Institute of Fundamental Research, Bombay 400 005, India; and CERN

K. Hikasa

Physics Department, Tohoku University, Aoba-ku, Sendai 980-77, Japan

G. Conforto

Università degli Studi, I-61029 Urbino, Italy; Istituto Nazionale di Fisica Nucleare, Sezione di Firenze, I-50125 Firenze, Italy

R.L. Workman

Department of Physics, Virginia Polytechnic Institute and State University, Blacksburg, VA 24061, USA

C. Grab

Institute for High Energy Physics, ETH-Hönggerberg, CH-8093 Zürich, Switzerland

Abstract

This biennial review summarizes much of Particle Physics. Using data from previous editions, plus 2300 new measurements from 700 papers, we list, evaluate, and average measured properties of gauge bosons, leptons, quarks, mesons, and baryons. We also summarize searches for hypothetical particles such as Higgs bosons, heavy neutrinos, monopoles, and supersymmetric particles. All the particle properties and search limits are listed in Summary Tables. We also give numerous tables, figures, formulae, and reviews of topics such as the Standard Model, particle detectors, probability, and statistics. A booklet is available containing the Summary Tables and abbreviated versions of some of the other sections of this full *Review*.

[†]The publication of the *Review of Particle Properties* is supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098, and by the U.S. National Science Foundation under Agreement No. PHY-9320551. Partial funding to cover the cost of this *Review* is also provided by the European Laboratory for Particle Physics (CERN), the Italian National Institute of Nuclear Physics (INFN), and by an implementing arrangement between the governments of Japan (Monbusho) and the United States (DOE) on cooperative research and development.

TABLE OF CONTENTS

Introduction			
I.	Overview	1177	
II.	Authors and consultants	1177	
III.	The naming scheme for hadrons	1179	
IV.	Procedures	1179	
	A. Selection and treatment of data	1179	
	B. Averages and fits	1180	
	1. Treatment of errors	1180	
	2. Unconstrained averaging	1180	
	3. Constrained fits	1181	
	C. Discussion	1182	
	History plots	1183	
	Using High-Energy Physics Databases	1185	
Summary Tables of Particle Properties			
	Gauge and Higgs bosons	1191	
	Leptons and quarks	1193	
	Mesons	1196	
	Baryons	1218	
	Searches	1227	
	Tests of conservation laws	1228	
Reviews, Tables, and Plots			
1.	Physical constants (rev.)	1233	
2.	Astrophysical constants (rev.)	1234	
3.	Big-bang cosmology (rev.)	1236	
4.	Dark matter (rev.)	1238	
5.	International System of Units (SI)	1240	
6.	Atomic and nuclear properties of materials (rev.)	1241	
7.	Periodic table of the elements (rev.)	1243	
8.	Electronic structure of the elements	1244	
9.	High-energy collider parameters (rev.)	1246	
10.	Passage of particles through matter (rev.)	1251	
11.	Photon and electron attenuation plots (rev.)	1259	
12.	Particle detectors	1261	
13.	Radioactivity and radiation protection	1268	
14.	Cosmic ray fluxes	1269	
15.	Commonly used radioactive sources	1270	
16.	Probability (rev.)	1271	
17.	Statistics (rev.)	1275	
18.	Monte Carlo techniques (rev.)	1283	
19.	Electromagnetic relations	1285	
20.	Clebsch-Gordan coefficients, spherical harmonics, and d functions	1287	
21.	SU(3) isoscalar factors and representation matrices	1288	
22.	SU(n) multiplets and Young diagrams	1289	
23.	Kinematics	1290	
24.	Cross-section formulae for specific processes (rev.)	1294	
25.	Quantum chromodynamics (rev.)	1297	
26.	Standard model of electroweak interactions (rev.)	1304	
27.	Constraints on new physics from electroweak analyses (new)	1312	
28.	Cabibbo-Kobayashi-Maskawa mixing matrix (rev.)	1315	
29.	CP violation (new)	1318	
30.	Quark model	1319	
31.	Naming scheme for hadrons	1323	
32.	Monte Carlo particle numbering scheme (rev.)	1325	
33.	Plots of cross sections and related quantities (rev.)	1327	
Full Listings*			
	Illustrative key and abbreviations	1343	
Gauge and Higgs bosons			
	(γ , gluon, graviton, W , Z , Higgs searches, Axions)	1351	
Leptons and quarks			
	(ν , e , μ , τ , lepton searches)	1385	
	(d , u , s , c , t , b' (4^{th} Generation))	1433	
Mesons			
	Light unflavored ($S = C = B = 0$)	1443	
	(π , ρ , a , b) (η , ω , f , ϕ , h)		
	Other light unflavored ($S = C = B = 0$)	1517	
	Strange ($S = \pm 1$, $C = B = 0$) (K , K^*)	1521	
	Charmed ($C = \pm 1$, $S = B = 0$) (D , D^*)	1564	
	Charmed, strange ($C = \pm 1$, $S = \pm 1$, $B = 0$)		
	(D_s , D_s^*)	1596	
	Bottom ($B = \pm 1$, $S = C = 0$) (B , B^*)	1601	
	Bottom, strange		
	($B = \pm 1$, $S = \mp$, $C = 0$) (B_s , B_s^*)	1639	
	$c\bar{c}$ (η_c , $J/\psi(1S)$, χ_c , ψ)	1642	
	$b\bar{b}$ (Υ , χ_b)	1661	
	Non- $q\bar{q}$ candidates	1670	
Baryons			
	N ($S = 0$, $I = 1/2$)	1673	
	Δ ($S = 0$, $I = 3/2$)	1710	
	Z ($S = +1$)		
	Λ ($S = -1$, $I = 0$)	1728	
	Σ ($S = -1$, $I = 1$)	1745	
	Ξ ($S = -2$, $I = 1/2$)	1769	
	Ω ($S = -3$, $I = 0$)	1780	
	Λ_c , Σ_c , Ξ_c , Ω_c ($C = +1$)	1782	
	Λ_b ($B = -1$)	1790	
Miscellaneous searches			
	Free quarks	1791	
	Monopoles	1793	
	Supersymmetry	1795	
	Compositeness	1805	
	Other	1811	
Other Compilations of Interest			
		1815	
Index			
		1817	

*The colored divider sheets give more detailed indices for each main section of the Full Listings.

INTRODUCTION

I. Overview	1177
II. Authors and consultants	1177
III. The naming scheme for hadrons	1179
IV. Procedures	1179
A. Selection and treatment of data	1179
B. Averages and fits	1180
1. Treatment of errors	1180
2. Unconstrained averaging	1180
3. Constrained fits	1181
C. Discussion	1182
History plots	1183

USING HIGH-ENERGY PHYSICS DATABASES

High-Energy Physics Databases	1185
Accessing the High-Energy Physics Databases	1188
Menu-Driven RPP Database at LBL	1188
Databases on the INTERNET/World-Wide Web	1188
Library Databases on ALICE at CERN	1188
Databases on SPIRES at SLAC	1188
Online Access to HEP at DESY	1189
Databases on SPIRES at Yukawa Institute	1189
High-Energy Physics Bulletin Board Preprint Archives	1189
IHEP-LBL Particle Physics Data System (PPDS) at LBL	1189
Durham-RAL Databases on BDMS at Durham and CERN	1189

INTRODUCTION

1. Overview

The *Review of Particle Properties* and the abbreviated version, the *Particle Physics Booklet*, are reviews of the field of Particle Physics. This complete *Review* includes a compilation/evaluation of data on particle properties, called the "Full Listings." These Listings include 2300 new measurements from 700 papers, in addition to the 12,000 measurements from 3500 papers that first appeared in previous editions. Both books include Summary Tables with our best values and limits for particle properties such as masses, widths or lifetimes, and branching fractions, as well as reviews, tables, and figures on a variety of topics. We also give an extensive summary of searches for hypothetical particles.

The *Review* and the *Booklet* are published in even-numbered years. This edition is an updating through December 1993 (and, in some areas, well into 1994). As described in the section "Using Particle Physics Databases" following this introduction, the content of this *Review* is available on the World-Wide Web, and there is a public access database allowing user-designed searches.

The Summary Tables give our best values of the properties of the particles we consider to be well established, a summary of search limits for hypothetical particles, and a summary of experimental tests of conservation laws.

The Full Listings contain all the data used to get the values given in the Summary Tables. Other measurements considered recent enough or important enough to mention, but which for one reason or another are not used to get the best values, appear separately just beneath the data we do use for the Summary Tables. The Full Listings also give information on unconfirmed particles and on particle searches, as well as short "minireviews" on subjects of particular interest or controversy.

The Full Listings were once an archive of all published data on particle properties. This is no longer possible because of the large quantity of data. We refer interested readers to earlier editions for data now considered to be obsolete.

We organize the particles into five categories:

- Gauge and Higgs bosons
- Leptons and quarks
- Mesons
- Baryons

Searches for free quarks, monopoles, supersymmetry, compositeness, *etc.*

The last category is for searches for particles that do not belong to the previous groups; searches for heavy charged leptons and massive neutrinos, for example, are with the leptons.

In addition to the compilations of measurements and best values, we give a long section of "Reviews, Tables, and Plots," a quick reference for the practicing particle physicist.

In Sec. 2 of this Introduction, we list the main areas of responsibility of the authors, and also list our large number of consultants, without whom we would not have been able to produce this *Review*. In Sec. 3, we mention briefly the naming scheme for hadrons. In Sec. 4, we discuss our procedures for choosing among measurements of particle

properties and for obtaining best values of the properties from the measurements.

The accuracy and usefulness of this *Review* depend in large part on interaction between its users and the authors. We appreciate comments, criticisms, and suggestions for improvements of any kind. Please send them to the appropriate author, according to the list of responsibilities in Sec. 2 below, or to

Particle Data Group, MS 50-308
Lawrence Berkeley Laboratory
Berkeley, CA 94720, USA

Or send them via computer mail to

PDG@LBL.GOV on INTERNET or
LBL::PDG on HEPNET, or
PDG@LBL on BITNET.

To order a copy of the *Review* or the *Particle Physics Booklet* from North and South America, Australia, and the Far East, write to

Particle Data Group, MS 50-308
Lawrence Berkeley Laboratory
Berkeley, CA 94720, USA

To order more than one copy of the *Review* or more than five Booklets, write to

Technical Information Division, MS 50B-2265
Lawrence Berkeley Laboratory
Berkeley, CA 94720, USA

From all other areas, write to

CERN Scientific Information Service
CH-1211 Geneva 23
Switzerland

2. Authors and consultants

The authors' main areas of responsibility are as follows (an asterisk indicates the person to contact with questions or comments):

Gauge and Higgs bosons

γ	C. Grab, C.G. Wohl*
Gluons	D.E. Groom*
Graviton	D.E. Groom* R.E. Shrock
W, Z	R.M. Barnett,* C. Caso, G. Conforto, A. Gurtu
Higgs bosons	R.M. Barnett,* K. Hikasa, H. Murayama
Heavy bosons	K. Hikasa, T.G. Trippe*
Axions	R.M. Barnett,* K. Hikasa, H. Murayama, K. Olive

Leptons and quarks

Neutrinos	D.E. Groom,* K. Olive, R.E. Shrock
e, μ	C. Grab, C.G. Wohl*
ν_τ, τ	D.E. Groom,* K.G. Hayes
Quarks	R.M. Barnett,* A. Manohar
Top quark	K. Hikasa, S. Kawabata, T.G. Trippe*

Mesons

π, η	D.E. Groom, C. Grab, C.G. Wohl*
Unstable mesons	M. Aguilar-Benitez, C. Caso, S. Eidelman, J.J. Hernandez, L. Montanet, F.C. Porter, M. Roos,* N.A. Törnqvist
K (stable)	C. Grab, T.G. Trippe*
D (stable)	R.J. Morrison, R.H. Schindler, C.G. Wohl*
B (stable)	R.J. Morrison, R.H. Schindler, T.G. Trippe*

Baryons

Stable baryons	C. Grab, R.H. Schindler, C.G. Wohl*
Unstable baryons	R.L. Crawford, G. Höhler, D.M. Manley, C.G. Wohl*, R.L. Workman

Miscellaneous searches

Quark & monopole	D.E. Groom,* J. Stone
Supersymmetric	R.M. Barnett,* K. Hikasa, H. Murayama, K. Olive
Compositeness	K. Hikasa, T.G. Trippe*
Other	K. Hikasa, T.G. Trippe*

Reviews, tables, figures, and formulae

R.M. Barnett, D.E. Groom,* T.G. Trippe, C.G. Wohl

Technical support

B. Armstrong, K. Gieselmann, P. Lantero, G.S. Wagman

We have also relied on the expertise of the following people for advice on particular topics:

- L. Addis (SLAC)
- M. Artuso (Syracuse University)
- V. Balakin (Novosibirsk)
- V.I. Balbekov (Serpukhov)
- T. Barnes (University of Tennessee)
- O. Beibel (RWTH, Aachen)
- D. Besson (Cornell University)
- S. Bethke (RWTH, Aachen)
- V. Bharadwaj (Fermilab)
- S. Bilenky (Joint Inst. for Nuclear Research, Dubna)
- M. Billing (Cornell University)
- G. Brianti (CERN)
- E. Browne
- J.N. Butler (Fermilab)
- R.N. Cahn (LBL)
- M. Chanowitz (LBL)
- A. Chao (SSC)
- Z. Chuang (IHEP, Beijing)
- E.D. Commins (University of California, Berkeley)
- COMPAS Group (IHEP, Serpukhov)
- O. Dahl (LBL)
- R.H. Dalitz (Oxford University)
- J. Donoghue (University of Massachusetts, Amherst)
- J. Dorfan (SLAC)
- S. Ecklund (SLAC)
- J. Ellis (CERN)
- J. Erler (University of Pennsylvania)
- L. Evans (CERN)
- V.V. Ezhela (Serpukhov)
- W. Fetscher (ETH, Zürich)
- V. Flaminio (University of Pisa)
- R. Flores (University of Minnesota)
- W. Frazer (University of California, Berkeley)
- S. Freedman (LBL and UC, Berkeley)
- R. Frosch (ETH, Zürich)
- A. Garren (UCLA)
- J. Gasser (University of Bern)
- S. Geer (Fermilab)
- H.-J. Gerber (ETH, Zürich)
- F.J. Gilman (SSC)
- H.A. Gould (LBL)
- N.A. Greenhouse (LBL)
- H.E. Haber (University of California, Santa Cruz)
- R. Hagstrom (ANL)
- I. Hinchliffe (LBL)
- H. Hirayama (KEK)
- P. Ivanov (Novosibirsk)
- Yu.M. Ivanov (Petersburg Nuclear Physics Inst.)
- J.D. Jackson (LBL)
- F. James (CERN)
- D. Karlen (Carleton University)
- R.W. Kenney (LBL)
- M. Klein (DESY)
- K. Kleinknecht (Universität Dortmund)
- S. Kurokawa (KEK)
- R. Kutschke (Cornell University)
- P. Langacker (University of Pennsylvania)
- D. Lanske (RWTH, Aachen)
- H. Leutwyler (University of Bern)
- M. Luty (LBL)
- G.R. Lynch (LBL)
- W.C. Martin (NIST)
- G. Moneti (Syracuse University)
- D. Morgan (Rutherford Appleton Lab)
- T. Nakada (PSI)
- K. Nakamura (Inst. Cosmic Ray Research, U. of Tokyo)
- W.R. Nelson (SLAC)
- Y. Oyanagi (University of Tsukuba, Japan)
- S.I. Parker (University of Hawaii)
- M.R. Pennington (University of Durham)
- M. Perl (SLAC)
- M. Peskin (SLAC)
- J. Primack (University of California, Santa Cruz)
- I. Protopopov (Novosibirsk)
- H.S. Pruis (Zürich University)
- H. Quinn (SLAC)
- B. Renk (Universität Mainz)
- J. Richman (University of California, Santa Barbara)
- B.L. Roberts (Boston University)
- B.P. Roe (University of Michigan)
- N.A. Roe (LBL)
- M. Ronan (LBL)
- L. Rosenberg (Stanford University)
- S. Rudaz (University of Minnesota)
- D. Schaile (CERN)
- D. Schramm (University of Chicago)

- V. Serbo (Novosibirsk State University)
- Yu. Shatunov (Novosibirsk)
- J.G. Smith (University of Colorado)
- E.M. Standish, Jr. (Jet Propulsion Lab., Pasadena)
- S. Stone (Cornell University)
- M. Suzuki (LBL)
- Y. Takaiwa (KEK)
- B.N. Taylor (U.S. National Bureau of Standards)
- G.H. Trilling (LBL)
- R.D. Tripp (LBL)
- M.S. Turner (Fermilab)
- G. Vignola (Frascati)
- M. Virchaux (Saclay)
- P. von Handel (DESY)
- R. Voss (CERN)
- H. Wahl (CERN)
- S.R. Wasserbaech (University of Washington)
- S. Willenbrock (University of Illinois)
- M.S. Witherell (University of California, Santa Barbara)
- L. Wolfenstein (Carnegie-Mellon University)
- B.N. Taylor (NIST)

In addition, the Particle Data Group benefits greatly from the assistance of some 700 physicists who are asked to verify every piece of data entered into this *Review*. Of special value is the advice of the PDG Advisory Committee which meets annually and thoroughly reviews all aspects of our operation. The members of the 1993 committee were:

W. Toki (Colorado State University), Chair
 D. Schaile (CERN)
 P. Kreitz (SLAC)
 P. Langacker (University of Pennsylvania)
 M. Turner (Fermilab)

3. Naming scheme for hadrons

We introduced in the 1986 edition [2] a new naming scheme for the hadrons. Changes from older terminology affected mainly the heavier mesons made of u , d , and s quarks. Otherwise, the only important change to known hadrons was that the F^\pm became the D_s^\pm . None of the lightest pseudoscalar or vector mesons changed names, nor did the $c\bar{c}$ or $b\bar{b}$ mesons (we do, however, now use χ_c for the $c\bar{c}$ X states), nor did any of the established baryons. The Summary Tables give both the new and old names whenever a change has occurred.

The scheme is described in “Naming Scheme for Hadrons” (p. 1323) of this *Review*.

We give here our conventions on type-setting style. Particle symbols are italic (or slanted) characters: e^- , p , Λ , π^0 , K_L , D_s^+ , b . Charge is indicated by a superscript: B^- , Δ^{++} . Charge is not normally indicated for p , n , or the quarks, and is optional for neutral isosinglets: η or η^0 . Antiparticles and particles are distinguished by charge for charged leptons and mesons: τ^+ , K^- . Otherwise, distinct antiparticles are indicated by a bar (overline): $\bar{\nu}_\mu$, \bar{l} , \bar{p} , \bar{K}^0 , and $\bar{\Sigma}^+$ (the antiparticle of the Σ^-).

4. Procedures

4.1. Selection and treatment of data: The Full Listings contain all relevant data known to us that are published in journals. With very few exceptions, we do not include results from preprints or conference reports. Nor do we include data that are of historical importance only (the Listings are not an archival record). We search every volume of 20 journals through our cutoff date for relevant data. We also include later published papers that are sent to us by the authors (or others).

In the Full Listings, we clearly separate measurements that are used to calculate or estimate values given in the Summary Tables from measurements that are not used. We give explanatory comments in many such cases. Among the reasons a measurement might be excluded are the following:

- It is superseded by or included in later results.
- No error is given.
- It involves assumptions we question.
- It has a poor signal-to-noise ratio, low statistical significance, or is otherwise of poorer quality than other data available.
- It is clearly inconsistent with other results that appear to be more reliable. Usually we then state the criterion, which sometimes is quite subjective, for selecting “more reliable” data for averaging. See Sec. 4.
- It is not independent of other results.
- It is not the best limit (see below).
- It is quoted from a preprint or a conference report.

In some cases, *none* of the measurements is entirely reliable and no average is calculated. For example, the masses of many of the baryon resonances, obtained from partial-wave analyses, are quoted as estimated ranges thought to probably include the true values, rather than as averages with errors. This is discussed in the Baryon Full Listings.

For upper limits, we normally quote in the Summary Tables the strongest limit. We do not average or combine upper limits except in a very few cases where they may be re-expressed as measured numbers with Gaussian errors.

As is customary, we assume that particle and antiparticle share the same spin, mass, and mean life. The Tests of Conservation Laws table, following the Summary Tables, lists tests of CPT as well as other conservation laws.

We use the following indicators in the Full Listings to tell how we get values from the tabulated measurements:

- OUR AVERAGE—From a weighted average of selected data.
- OUR FIT—From a constrained or overdetermined multi-parameter fit of selected data.
- OUR EVALUATION—Not from a direct measurement, but evaluated from measurements of related quantities.
- OUR ESTIMATE—Based on the observed range of the data. Not from a formal statistical procedure.
- OUR LIMIT—For special cases where the limit is evaluated by us from measured ratios or other data. Not from a direct measurement.

An experimentalist who sees indications of a particle will of course want to know what has been seen in that region in the past. Hence we include in the Full Listings all reported

states that, in our opinion, have sufficient statistical merit and that have not been disproved by more reliable data. However, we promote to the Summary Tables only those states that we feel are well established. This judgment is, of course, somewhat subjective and no precise criteria can be given. For more detailed discussions, see the minireviews in the Full Listings.

4.2. Averages and fits: We divide this discussion on obtaining averages and errors into three sections: (1) treatment of errors; (2) unconstrained averaging; (3) constrained fits.

4.2.1. Treatment of errors: In what follows, the “error” δx means that the range $x \pm \delta x$ is intended to be a 68.3% confidence interval about the central value x . We treat this error as if it were Gaussian. Thus when the error is Gaussian, δx is the usual one standard deviation (1σ). Many experimenters now give statistical and systematic errors separately, in which case we usually quote both errors, with the statistical error first. For averages and fits, we then add the two errors in quadrature and use this combined error for δx .

When experimenters quote asymmetric errors $(\delta x)^+$ and $(\delta x)^-$ for a measurement x , the error that we use for that measurement in making an average or a fit with other measurements is a continuous function of these three quantities. When the resultant average or fit \bar{x} is less than $x - (\delta x)^-$, we use $(\delta x)^-$; when it is greater than $x + (\delta x)^+$, we use $(\delta x)^+$. In between, the error we use is a linear function of x . Since the errors we use are functions of the result, we iterate to get the final result. Asymmetric output errors are determined from the input errors assuming a linear relation between the input and output quantities.

In fitting or averaging, we usually do not include correlations between different measurements, but we try to select data in such a way as to reduce correlations. Correlated errors are, however, treated explicitly when there are a number of results of the form $A_i \pm \sigma_i \pm \Delta$ that have identical systematic errors Δ . In this case, one can first average the $A_i \pm \sigma_i$ and then combine the resulting statistical error with Δ . One obtains, however, the same result by averaging $A_i \pm (\sigma_i^2 + \Delta^2)^{1/2}$, where $\Delta_i = \sigma_i \Delta [\sum (1/\sigma_j^2)]^{1/2}$. This procedure has the advantage that, with the modified systematic errors Δ_i , each measurement may be treated as independent and averaged in the usual way with other data. Therefore, when appropriate, we adopt this procedure. We tabulate Δ and invoke an automated procedure that computes Δ_i before averaging and we include a note saying that there are common systematic errors.

Another common case of correlated errors occurs when experimenters measure two quantities and then quote the two and their difference, e.g., m_1 , m_2 , and $\Delta = m_2 - m_1$. We cannot enter all of m_1 , m_2 and Δ into a constrained fit because they are not independent. In some cases, it is a good approximation to ignore the quantity with the largest error and put the other two into the fit. However, in some cases correlations are such that the errors on m_1 , m_2 and Δ are comparable and none of the three values can be ignored. In this case, we put all three values into the fit and invoke an automated procedure to increase the errors prior to fitting such that the three quantities can be treated as independent

measurements in the constrained fit. We include a note saying that this has been done.

4.2.2. Unconstrained averaging: To average data, we use a standard weighted least-squares procedure and in some cases, discussed below, increase the errors with a “scale factor.” We begin by assuming that measurements of a given quantity are uncorrelated, and calculate a weighted average and error as

$$\bar{x} \pm \delta\bar{x} = \frac{\sum_i w_i x_i}{\sum_i w_i} \pm (\sum_i w_i)^{-1/2}, \quad (1)$$

where

$$w_i = 1/(\delta x_i)^2.$$

Here x_i and δx_i are the value and error reported by the i th experiment, and the sums run over the N experiments. We then calculate $\chi^2 = \sum w_i (\bar{x} - x_i)^2$ and compare it with $N - 1$, which is the expectation value of χ^2 if the measurements are from a Gaussian distribution.

If $\chi^2/(N - 1)$ is less than or equal to 1, and there are no known problems with the data, we accept the results.

If $\chi^2/(N - 1)$ is very large, we may choose not to use the average at all. Alternatively, we may quote the calculated average, but then make an educated guess of the error, a conservative estimate designed to take into account known problems with the data.

Finally, if $\chi^2/(N - 1)$ is greater than 1, but not greatly so, we still average the data, but then also do the following:

(a) We increase our quoted error, $\delta\bar{x}$ in Eq. (1), by a scale factor S defined as

$$S = [\chi^2/(N - 1)]^{1/2}. \quad (2)$$

Our reasoning is as follows. The large value of the χ^2 is likely to be due to underestimation of errors in at least one of the experiments. Not knowing which of the errors are underestimated, we assume they are all underestimated by the same factor S . If we scale up all the input errors by this factor, the χ^2 becomes $N - 1$, and of course the output error $\delta\bar{x}$ scales up by the same factor. See Ref. 3.

When combining data with widely varying errors, we modify this procedure slightly. We evaluate S using only the experiments with smaller errors. Our cutoff or ceiling on δx_i is arbitrarily chosen to be

$$\delta_0 = 3N^{1/2} \delta\bar{x},$$

where $\delta\bar{x}$ is the unscaled error of the mean of all the experiments. Our reasoning is that although the low-precision experiments have little influence on the values \bar{x} and $\delta\bar{x}$, they can make significant contributions to the χ^2 , and the contribution of the high-precision experiments thus tends to be obscured. Note that if each experiment has the same error δx_i , then $\delta\bar{x}$ is $\delta x_i/N^{1/2}$, so each δx_i is well below the cutoff. (More often, however, we simply exclude measurements with relatively large errors from averages and fits: new, precise data chase out old, imprecise data.)

Our scaling procedure has the property that if there are two values with comparable errors separated by much more than their stated errors (with or without a number of other values of lower accuracy), the scaled-up error $\delta\bar{x}$ is

approximately half the interval between the two discrepant values.

We emphasize that our scaling procedure for *errors* in no way affects central values. And if you wish to recover the unscaled error $\delta\bar{x}$, simply divide the quoted error by S .

(b) If the number M of experiments with an error smaller than δ_0 is at least three, and if $\chi^2/(M-1)$ is greater than 1.25, we show in the Full Listings an ideogram of the data. Fig. 1 is an example. Sometimes one or two data points lie apart from the main body; other times the data split into two or more groups. We extract no numbers from these ideograms; they are simply visual aids, which the reader may use as he or she sees fit.




Figure 1: A typical ideogram. The “data point” at the top shows the position of the weighted average, while the width of the error bar (and the shaded pattern beneath it) shows the error in the average after scaling by the factor S . The column on the right gives the χ^2 contribution of each of the experiments. Note that the next-to-last experiment, denoted by the incomplete error flag (\perp), is not used in the calculation of S (see the text).

Each measurement in an ideogram is represented by a Gaussian with a central value x_i , error δx_i , and area proportional to $1/\delta x_i$. The choice of $1/\delta x_i$ for the area is somewhat arbitrary. With this choice, the center of gravity of the ideogram corresponds to an average that uses weights $1/\delta x_i$ rather than the $(1/\delta x_i)^2$ actually used in the averages. This may be appropriate when some of the experiments have seriously underestimated systematic errors. However, since for this choice of area the height of the Gaussian for each measurement is proportional to $(1/\delta x_i)^2$, the peak position of the ideogram will often favor the high-precision measurements at least as much as does the least-squares average. See our 1986 edition [2] for a detailed discussion of the use of ideograms.

4.2.3. Constrained fits: Except for trivial cases, all branching ratios and rate measurements are analyzed by making a simultaneous least-squares fit to all the data and extracting the partial decay fractions P_i , the partial widths Γ_i , the full width Γ (or mean life), and the associated error matrix.

Assume, for example, that a state has m partial decay fractions P_i , where $\sum P_i = 1$. These have been measured in N_r different ratios R_r , where, e.g., $R_1 = P_1/P_2$, $R_2 = P_1/P_3$, etc. [We can handle any ratio R of the form $\sum \alpha_i P_i / \sum \beta_i P_i$, where α_i and β_i are constants, usually 1 or 0. The forms $R = P_i P_j$ and $R = (P_i P_j)^{1/2}$ are also allowed.] Further assume that *each* ratio R has been measured by N_k experiments (we designate each experiment with a subscript k , e.g., R_{1k}). We then find the best values of the fractions P_i by minimizing the χ^2 as a function of the $m-1$ independent parameters:

$$\chi^2 = \sum_{r=1}^{N_r} \sum_{k=1}^{N_k} \left(\frac{R_{rk} - R_r}{\delta R_{rk}} \right)^2, \quad (3)$$

where the R_{rk} are the measured values and R_r are the fitted values of the branching ratios.

In addition to the fitted values \bar{P}_i , we calculate an error matrix $\langle \delta \bar{P}_i \delta \bar{P}_j \rangle$. We tabulate the diagonal elements of $\delta \bar{P}_i = \langle \delta \bar{P}_i \delta \bar{P}_i \rangle^{1/2}$ (except that some errors are scaled as discussed below). In the Full Listings, we give the complete correlation matrix; we also calculate the fitted value of each ratio, for comparison with the input data, and list it above the relevant input, along with a simple unconstrained average of the same input.

Three comments on the example above:

(1) There was no connection assumed between measurements of the full width and the branching ratios. But often we also have information on partial widths Γ_i as well as the total width Γ . In this case we must introduce Γ as a parameter in the fit, along with the P_i , and we give correlation matrices for the widths in the Full Listings.

(2) We do *not* allow for correlations between input data. We *do* try to pick those ratios and widths that are as independent and as close to the original data as possible. When one experiment measures all the branching fractions and constrains their sum to be one, we leave one of them (usually the least well-determined one) out of the fit to make the set of input data more nearly independent.

(3) We calculate scale factors for both the R_r and P_i when the measurements for any R give a larger-than-expected contribution to the χ^2 . According to Eq. (3), the double sum for χ^2 is first summed over experiments $k = 1$ to N_k , leaving a single sum over ratios $\chi^2 = \sum \chi_r^2$. One is tempted to define a scale factor for the ratio r as $S_r^2 = \chi_r^2 / \langle \chi_r^2 \rangle$. However, since $\langle \chi_r^2 \rangle$ is not a fixed quantity (it is somewhere between N_k and N_{k-1}), we do not know how to evaluate this expression. Instead we define

$$S_r^2 = \frac{1}{N_k} \sum_{k=1}^{N_k} \frac{(R_{rk} - \bar{R}_r)^2}{(\delta R_{rk})^2 - (\delta \bar{R}_r)^2}, \quad (4)$$

where $\delta \bar{R}_r$ is the fitted error for ratio r . With this definition the expected value of S_r^2 is one.

The fit is redone using errors for the branching ratios that are scaled by the larger of S_r and unity, from which new

and often larger errors $\delta\bar{P}_i'$ are obtained. The scale factors we finally list in such cases are defined by $S_i = \delta\bar{P}_i'/\delta\bar{P}_i$. However, in line with our policy of not letting S affect the central values, we give the values of \bar{P}_i obtained from the original (unscaled) fit.

There is one special case in which the errors that are obtained by the preceding procedure may be changed. When a fitted branching ratio (or rate) \bar{P}_i turns out to be less than three standard deviations ($\delta\bar{P}_i'$) from zero, a new smaller error $(\delta\bar{P}_i'')^-$ is calculated on the low side by requiring the area under the Gaussian between $\bar{P}_i - (\delta\bar{P}_i'')^-$ and \bar{P}_i to be 68.3% of the area between zero and \bar{P}_i . A similar correction is made for branching fractions that are within three standard deviations of one. This keeps the quoted errors from overlapping the boundary of the physical region.

4.3. Discussion: The problem of averaging data containing discrepant values is nicely discussed by Taylor in Ref. 4. He considers a number of algorithms that attempt to incorporate inconsistent data into a meaningful average. However, it is difficult to develop a procedure that handles simultaneously in a reasonable way two basic types of situations: (a) data that lie apart from the main body of the data are incorrect (contain unreported errors); and (b) the opposite—it is the main body of data that is incorrect. Unfortunately, as Taylor shows, case (b) is not infrequent. He concludes that the choice of procedure is less significant than the initial choice of data to include or exclude.

We place much emphasis on this choice of data. Often we solicit the help of outside experts (consultants). Sometimes, however, it is simply impossible to determine which of a set of discrepant measurements are correct. Our scale-factor technique is an attempt to address this ignorance by increasing the error. In effect, we are saying that present experiments do not allow a precise determination of this quantity because of unresolvable discrepancies, and one must await further measurements. The reader is warned of this situation by the size of the scale factor, and if he or she desires can go back to the literature (via the Full Listings) and redo the average with a different choice of data.

Our situation is less severe than most of the cases Taylor considers, such as estimates of the fundamental constants like \hbar , *etc.* Most of the errors in his case are dominated by systematic effects. For our data, statistical errors are often at least as large as systematic errors, and statistical errors are usually easier to estimate. A notable exception occurs in partial-wave analyses, where different techniques applied to the same data yield different results. In this case, as stated earlier, we often do not make an average but just quote a range of values.

A brief history of early Particle Data Group averages is given in Ref. 3. Figure 0.2 shows some histories of our values of a few particle properties. Sometimes large changes occur. These usually reflect the introduction of significant new data or the discarding of older data. Older data are discarded in favor of newer data when it is felt that the newer data have smaller systematic errors, or have more checks on systematic errors, or have made corrections unknown at the time of the older experiments, or simply have much smaller errors. Sometimes, the scale factor becomes large near the time at which a large jump takes place, reflecting the uncertainty introduced by the new and inconsistent data.

By and large, however, a full scan of our history plots shows a dull progression toward greater precision at central values quite consistent with the first data points shown.

We conclude that the reliability of the combination of experimental data and our averaging procedures is usually good, but it is important to be aware that fluctuations outside of the quoted errors can and do occur.

ACKNOWLEDGMENTS

The publication of the *Review of Particle Properties* is supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract No. DE-AC03-76SF00098, by the U.S. National Science Foundation under Agreement No. PHY-9320551, and by the Italian Istituto Nazionale di Fisica Nucleare (INFN). Partial funding to cover the cost of this *Review* is also provided by the European Laboratory for Particle Physics (CERN) and by an implementing arrangement between the governments of Japan (Monbusho) and the United States (DOE) on cooperative research and development.

We thank all those who have assisted in the many phases of preparing this *Review*. We particularly thank the many who have responded to our requests for verification of data entered in the Listings, and those who have made suggestions or pointed out errors. The Berkeley members of the Particle Data Group acknowledge the administrative and baking skills of Gail Harper and the graphics expertise of Charles Groom.

REFERENCES

1. The previous edition was Particle Data Group: K. Hikasa, K. Hagiwara, S. Kawabata, R.M. Barnett, D.E. Groom, T.G. Trippe, C.G. Wohl, G.P. Yost, B. Armstrong, G.S. Wagman, J. Stone, J.J. Hernández, F.C. Porter, R.J. Morrison, R.E. Cutkosky, L. Montanet, K. Gieselmann, M. Aguilar-Benitez, C. Caso, R.L. Crawford, M. Roos, N.A. Törnqvist, K.G. Hayes, G. Höhler, D.M. Manley, K.A. Olive, R.E. Shrock, S. Eidelman, R.H. Schindler, G. Conforto, and R.A. Eichler *Phys. Rev. D* **45**, 1 June 1992, Part II.
2. Particle Data Group: M. Aguilar-Benitez, F.C. Porter, J.J. Hernandez, L. Montanet, K.R. Schubert, M. Roos, N.A. Törnqvist, G. Höhler, R.M. Barnett, I. Hinchliffe, G.R. Lynch, A. Rittenberg, T.G. Trippe, G.P. Yost, B. Armstrong, G.S. Wagman, T. Shimada, G.P. Gopal, J. Primack, K.G. Hayes, R.H. Schindler, R.E. Shrock, R.A. Eichler, R. Frosch, L.D. Roper, and W.P. Trower, *Phys. Lett.* **170B** (1986).
3. A.H. Rosenfeld, *Ann. Rev. Nucl. Sci.* **25**, 555 (1975).
4. B.N. Taylor, "Numerical Comparisons of Several Algorithms for Treating Inconsistent Data in a Least-Squares Adjustment of the Fundamental Constants," U.S. National Bureau of Standards NBSIR 81-2426 (1982).




Figure 2: An historical perspective of values of a few particle properties tabulated in this *Review* as a function of date of publication of the *Review*. A full error bar indicates the quoted error; a thick-lined portion indicates the same but without the “scale factor.”

Revised by G.S. Wagman and B. Armstrong, June 1994

The Full Listings in this *Review of Particle Properties*, as well as other particle physics databases, are computer accessible. Some of the databases help find papers of interest, while others contain actual numerical data. Here we describe some databases maintained at LBL, SLAC, CERN, Durham, IHEP, KEK, and Yukawa, and how to start using them.

1. High-Energy Physics Databases

This section describes publicly accessible databases of interest to high-energy physicists. See Table 1 for availability, contacts, and user guides. For databases that are derived from other databases, see also the description of the originating database. See Section 2 for information on accessing these databases.

RPP contains the Full Listings from this *Review of Particle Properties*. Through user-friendly menus, users may query by paper, particle, mass range, quantum numbers, or detector, and can select specific properties or classes of properties like masses or decay parameters. All other relevant information (*e.g.*, footnotes and references) is included. Complete instructions are available online. The database is completely updated in the Summer of even-numbered years before publication of the *Review*, and is less thoroughly updated in the Summer of odd-numbered years.

The Particle Data Group provides, through the World-Wide Web (see Section 2.2), full-text PostScript of the *Review of Particle Properties*, excluding the Full Listings. For example, users may access the articles, tables, figures, and formulae.

Other High-Energy Physics Databases:

- **BOOKS** contains bibliographic summaries of more than 20,000 textbooks, conference proceedings, lecture-notes, monographs, and serials covering high-energy physics and related subjects.
- **CONF (CERN)**, a subset of **LIB**, contains forthcoming conferences of interest for high-energy physics and accelerator research and contains past conferences back to 1986.
- **CONF (SLAC)** contains almost 5,000 listings of conferences, schools, and meetings of interest to high-energy physicists. Information on forthcoming conferences is entered regularly, with detailed descriptions and links to further World-Wide Web information when available.
- **CS**, regularly updated from the **REACTIONS** database, contains data from CERN-HERA, UCRL, and LBL cross-section compilations covering 1950 to the present.
- **DIR**, the Directory of Research Institutes in High Energy Physics, contains addresses, telephone, fax, and telex numbers, and e-mail nodes, as well as brief information on research programs and accelerators.

To obtain **DIR** in a Filemaker PRO format for Macintosh computers, contact the SIS Secretariat at CERN or Wolfgang Simon (ISI@CERNVM.CERN.CH).

- **DOCUMENTS** contains two groups of keys:

(1) Bibliographic: ID, references, year of preprinting or publication, authors and affiliations, document title, experiment number, collaboration name, and related references.

(2) Topical: beam particle, target particle, reactions, particles in the final states of reactions, momenta in initial states, types of data obtained, particles whose property has been measured, accelerator and/or detector, and initial state polarization.

It covers 1895 to the present, with coverage since 1950 being more complete, and is updated monthly. The report "A Guide to Experimental Elementary Particle Physics Literature" [1] is produced from it.

- **E-MAIL IDS**, derived from **HEP NAMES**, contains e-mail addresses of many people working in high-energy physics.
- **EXPERIMENTS** contains summaries of approved experiments at major laboratories. It covers approximately 1975 to the present, with coverage since 1980 being more complete. It is searchable by experiment number, author, accelerator, detector, reaction, beam momentum, journal paper, and other items. The report "Current Experiments in Elementary Particle Physics" [2] is produced from it.
- **HEP-PREPRINT** is a collection of bulletin board archives reserved for high-energy physics preprints. It includes **HEP-EX** for experimental HEP preprints (since April 1994), **HEP-LAT** for lattice/computational preprints (since December 1991), **HEP-PH** for particle phenomenology preprints (since March 1992), and **HEP-TH** for string/conformal/field theory preprints (since August 1991). Other archives are described in the HELP facility of these archives.
- **HEP (SPIRES-HEP)**, a joint project of the SLAC and DESY libraries, contains 270,000 bibliographic entries on particle physics papers (journal articles, preprints, reports, theses, *etc.*). It covers 1974 to the present and is updated daily. It is searchable by author, institution, title, topic, report number, citation, bulletin board number, and other bibliographic items. It is an indexing and access tool (via the World-Wide Web) to more than 12,000 full-text PostScript documents, including bulletin board articles processed jointly by the SLAC and DESY libraries and the CERN publication group.
- **HEP NAMES** contains 23,000 e-mail addresses of many people working in high-energy physics.
- **INSTITUTIONS** contains 3,000 addresses and, often, phone and fax numbers of high-energy physics-related institutions.
- **LIB** contains the CERN library's catalogue of books, reports, preprints, and other information.
- **PP** contains information on particle properties derived from the Summary Tables in this *Review of Particle Properties*.

Table 1. Summary of High-Energy Physics Databases

NAME	SYSTEM	AVAILABILITY	CONTACT	USER
		RESPONSIBILITY*		GUIDE
<i>BOOKS</i>	SPIRES	WWW,QSPIRES, SLAC ,Yukawa	<i>a,b</i>	<i>c</i>
<i>CONF (CERN)</i> ¹	ALICE	WWW, CERN	<i>d</i>	<i>e</i>
<i>CONF (SLAC)</i> ¹	SPIRES	WWW,QSPIRES, SLAC ,Yukawa	<i>f,b</i>	<i>c</i>
<i>CS</i> ²	PPDS	IHEP ,LBL	<i>g</i>	<i>g,h</i>
<i>DIR</i>	ALICE	WWW, CERN	<i>d</i>	<i>i</i>
<i>DOCUMENTS</i> ³	PPDS	IHEP ,LBL	<i>g</i>	<i>g,h</i>
<i>E-MAIL IDS</i> ⁴	BDMS	WWW, DURHAM ,CERN	<i>j</i>	<i>j,i</i>
<i>EXPERIMENTS</i> ⁵	BDMS	WWW,DURHAM,CERN	<i>j</i>	<i>j,i</i>
<i>EXPERIMENTS</i> ⁵	PPDS	IHEP	<i>g</i>	<i>g,h</i>
<i>EXPERIMENTS</i> ⁵	SPIRES	WWW,QSPIRES, SLAC ,Yukawa	<i>k,b</i>	<i>c</i>
<i>HEP-PREPRINT</i>	—	WWW,FTP,EMAIL	<i>l</i>	<i>m</i>
<i>HEP (SPIRES-HEP)</i> ⁶	SPIRES	WWW,QSPIRES, SLAC , DESY ,KEK,Yukawa	<i>n,o,p,b</i>	<i>c</i>
<i>HEPNAMES</i>	SPIRES	WWW,QSPIRES, SLAC ,Yukawa	<i>q,b</i>	<i>c</i>
<i>INSTITUTIONS</i> ⁷	SPIRES	WWW,QSPIRES, SLAC ,Yukawa	<i>r,b</i>	<i>c</i>
<i>LIB</i>	ALICE	WWW, CERN	<i>d</i>	<i>e</i>
<i>PP</i> ⁸	PPDS	IHEP ,LBL	<i>g</i>	<i>g,h</i>
<i>PREP</i>	ALICE	WWW, CERN	<i>d</i>	<i>e</i>
<i>REACTION DATA</i> ⁹	BDMS	WWW, DURHAM ,CERN	<i>j</i>	<i>j,i</i>
<i>REACTIONS</i> ¹⁰	PPDS	IHEP ,LBL	<i>g</i>	<i>g,h</i>
<i>RPP</i> ¹¹	MENU-DRIVEN	WWW, LBL	<i>h</i>	—
<i>SLACPPF/CITATIONS</i> ¹²	BDMS	WWW, DURHAM ,CERN	<i>j</i>	<i>j,i</i>
<i>VOCABULARY</i> ¹³	PPDS	IHEP ,LBL	<i>g</i>	<i>g,h</i>

* Institutions listed in **BOLD** are responsible for the content of the database.

(continued)

Table 1 (continuation)

-
-
- ¹ *CONF (CERN)* and *CONF (SLAC)* are similar in nature, but are not identical.
- ² Updated mainly from *REACTIONS* and maintained by the IHEP COMPAS Group.
- ³ Maintained by the IHEP COMPAS Group with input from the LBL Particle Data Group, ITEP, and JINR. Bibliographic data is extracted from *SPIRES-HEP* and *LIB*.
- ⁴ Derived from *HEPNames*.
- ⁵ Maintained by SLAC, transferred to other institutions. (Ref. 2).
- ⁶ Maintained by SLAC in collaboration with the DESY HEP Index Group, transferred to other institutions nightly.
- ⁷ Maintained by SLAC in collaboration with the LBL Particle Data Group.
- ⁸ Derived from the Summary Tables in this *Review of Particle Properties*; maintained by the IHEP COMPAS Group.
- ⁹ Maintained by the HEP Database Group at Durham; data exchanged twice yearly with IHEP.
- ¹⁰ Maintained by the IHEP COMPAS Group with input from ITEP; data exchanged twice yearly with Durham.
- ¹¹ Derived from the Full Listings in this *Review of Particle Properties*.
- ¹² A subset of *SPIRES-HEP*.
- ¹³ Maintained by the IHEP COMPAS Group with input from *RPP* and *EXPERIMENTS*.
-

- a*: LIBRARY@SLAC.STANFORD.EDU (SLAC Library, MS-82, P.O. Box 4349, Stanford, CA 94309, USA)
- b*: AOKI@HEP.S.KANAZAWA-U.AC.JP (Ken-Ichi Aoki, Yukawa Inst., Kyoto Univ., Kyoto 606, Japan)
- c*: TECHPUB@SLAC.STANFORD.EDU (Order: "Guide to QSPIRES and the Particle Physics Databases on SLACVM," SLAC-393 Report, by Hrvoje Galić)
- d*: MALICE@VXLIB.CERN.CH
- e*: LIBDESK@CERN.CH
- f*: CONF@SLAC.STANFORD.EDU (Georgia Row, SLAC Library, MS-82, P.O. Box 4349, Stanford, CA 94309, USA)
- g*: ALEKHIN@MX.IHEP.SU or EZHEL@MX.IHEP.SU (Sergey Alekhin or Vladimir Ezhela, IHEP, Protvino, Moscow Region, Russian Federation, RU-142284)
- h*: PDG@LBL.GOV (Particle Data Group, LBL, 50-308, Berkeley, CA 94720, USA)
- i*: ANTON@CERNVM.CERN.CH (Anna Anton, CERN Library, CH-1211 Geneva 23 Switzerland)
- j*: M.R.WHALLEY@DURHAM.AC.UK (Mike Whalley, Durham Univ., South Rd., Durham City, DH1 3LE, UK)
or RGR@V2.RL.AC.UK (Dick Roberts, Rutherford Appleton Laboratory, Chilton, Didcot, Oxon. OX11 0QX, UK)
- k*: EXPBASE@SLAC.STANFORD.EDU (Hrvoje Galić, SLAC, MS-82, P.O. Box 4349, Stanford, CA 94309, USA)
- l*: HEP-EX@XXX.LANL.GOV, HEP-LAT@FTP.SCRI.FSU.EDU, HEP-PH@XXX.LANL.GOV, or HEP-TH@XXX.LANL.GOV
(Send e-mail with subject COMMENT)
- m*: HEP-EX@XXX.LANL.GOV, HEP-LAT@FTP.SCRI.FSU.EDU, HEP-PH@XXX.LANL.GOV, or HEP-TH@XXX.LANL.GOV
(Send e-mail with subject HELP and no message.)
- n*: HEP@SLAC.STANFORD.EDU (SLAC Library, MS-82, P.O. Box 4349, Stanford, CA 94309, USA)
- o*: LOOHTP@DSYIBM.DESY.DE (Hartmut Preissner, DESY, Notkestrasse 85, D-22603 Hamburg, Germany)
- p*: MIURA@KEKVAX.KEK.JP (Yasuko Miura, KEK, 1-1 Oho, Tsukuba-shi, Ibaraki-ken 305, Japan)
- q*: HEPNames@SLAC.STANFORD.EDU (Hrvoje Galić, SLAC, MS-82, P.O. Box 4349, Stanford, CA 94309, USA)
- r*: LIRYG@SLAC.STANFORD.EDU (Robert Gex, SLAC, MS-82, P.O. Box 4349, Stanford, CA 94309, USA)
-
-

- **PREP**, a subset of **LIB**, contains entries for preprints, reports, conference papers, theses, *etc.* It covers all preprints and reports received in the CERN Library since 1980. It also has publication details for all papers published with CERN as an affiliation and for many conference papers published in proceedings since about 1987.
- **REACTION DATA** is a compilation of numerical experimental particle physics reaction data, including data from 2-body (and quasi-2-body) scattering, e^+e^- annihilation, and inclusive hadron, photon, and lepton physics such as total and differential cross sections, fragmentation functions, structure functions, and polarization measurements.
- **REACTIONS**, in conjunction with **REACTION DATA**, contains numerical data on reactions: differential and total cross sections, structure functions, polarization measurements, and other quantities. It covers 1952 to the present and is updated approximately quarterly.
- **SLACPPF/CITATIONS**, a subset of the **SPIRES-HEP** literature-searching guide, contains references to papers and preprints since 1980, being comprised of the SLAC PPF (preprint) records with PPA (published references to PPF) updates compiled by the SLAC library. Many journal publications compiled by the DESY library are also included.
- **VOCABULARY** controls usage of particle names, accelerator names, detector names, and data descriptors in all PPDS databases.

2. Accessing the High-Energy Physics Databases

2.1. Menu-Driven RPP Database at LBL

To access the **RPP** database on INTERNET: TELNET MUSE.LBL.GOV (131.243.48.11) or on DECNET: SET HOST MUSE (42062). Then login to the captive account PDG_PUBLIC (a password is not required).

2.2. Databases on the INTERNET/World-Wide Web

Many databases, including several of the high-energy physics databases discussed above, are accessible via the World-Wide Web (W3 or WWW) (see Table 1), which is an INTERNET-based wide-area hypermedia information retrieval system. There are several browsers available (WWW line-mode browser, NCSA's Mosaic browser for X Windows, SLAC's MidasWWW browser for X Windows, Cello browser for MSDOS, *etc.*) that may already be installed at your institution; if not, try TELNET INFO.CERN.CH, which will connect you to WWW and will explain how to acquire the browsers.

Much of this *Review of Particle Properties* can be accessed through the World-Wide Web link

<http://www-pdg.lbl.gov/>

2.3. Library Databases on ALICE at CERN

The CERN Library ALICE databases are accessible through the World-Wide Web link

<http://www.cern.ch/>

This link and the subsequent "Preprints" link also provide access to full-text preprints received since 1994 from bulletin board archives and from scanned papers.

To access ALICE on INTERNET: TELNET ALICE.CERN.CH (128.141.201.44) or on DECNET: SET HOST VXLIB (22748). Then login to account ALICE (a password is not required) and select the terminal type according to the menu. ALICE is a full-screen system using the DEC international character set, which can be displayed on suitable terminals. Simple searching can be done by using a menu system or by using the full power of the ISO Common Command Language; HELP displays are provided to guide searching. With the MAIL command, the results of searches can be sent to any e-mail address for printing.

People without login access to CERN can use QALICE.

Typical messages from OpenVMS to QALICE:

```
mshg VXLIB QALICE base prep;f black hole?;
mshg VXLIB QALICE base and 1991-->1992/yr;show
mshg VXLIB QALICE base dir;f org=cern;show full
```

Alternately, send a blank e-mail message to

QALICE@VXLIB.CERN.CH

and put the query in the subject field. For further information, send the subject HELP to QALICE.

Regular weekly or monthly searches of the CERN databases can be arranged according to a personal search 'profile', with the results sent automatically by e-mail. For details on this Selective Dissemination of Information (SDI) service on QALICE, contact David Dallman, Scientific Information Service (SIS), CERN, CH-1211 Geneva 23, Switzerland (DALLMAN@CERNVM.CERN.CH).

2.4. Databases on SPIRES at SLAC

SLAC encourages the high-energy physics community to access its databases through the World-Wide Web link <http://www-slac.slac.stanford.edu/find/spires.html> leading to **SPIRES-HEP**, **BOOKS**, **CONF**, **HEPNAMES**, **INSTITUTIONS**, **EXPERIMENTS**, and other SLAC databases.

People without login access or World-Wide Web access to SLAC can use QSPIRES. QSPIRES, as described in the 1992 edition of the *Review of Particle Properties*, is a remote server which enables e-mail access to SPIRES databases. Effective March 1994, registration or authorization is no longer needed to access QSPIRES. For further information on QSPIRES, contact QSPI@SLAC.STANFORD.EDU.

2.5. Online Access to HEP at DESY

DESY offers accounts for remote login from Europe restricted to *SPIRES-HEP*. Contact Hartmut Preissner (LOOHTP@DSYIBM.DESY.DE).

2.6. Databases on SPIRES at Yukawa Institute

The Yukawa Institute provides Gopher access (GOPHER.YUKAWA.KYOTO-U.AC.JP) and QSPIRES access (JPNYITP.YUKAWA.KYOTO-U.AC.JP) from the Far East to SPIRES databases. Contact Ken-Ichi Aoki (AOKI@HEP.S.KANAZAWA-U.AC.JP).

2.7. High-Energy Physics Bulletin Board Preprint Archives

E-mail listings of high-energy physics preprint titles and abstracts submitted to the archives can be received daily by sending a blank e-mail message to the appropriate archive (HEP-EX@XXX.LANL.GOV, HEP-LAT@FTP.SCRI.FSU.EDU, HEP-PH@XXX.LANL.GOV, or HEP-TH@XXX.LANL.GOV) with the subject:

SUBSCRIBE *your_name*

To receive detailed instructions on submitting and recovering papers, send a blank e-mail message with the subject:

HELP

The listings and papers can also be accessed through the World-Wide Web link

<http://xxx.lanl.gov/>

2.8. IHEP-LBL Particle Physics Data System (PPDS) at LBL

The databases maintained by the IHEP Protvino COMPAS Group under the Berkeley Database Management System (BDMS) with input from the world-wide Particle Data Group collaboration can be accessed interactively on INTERNET: TELNET MUSE.LBL.GOV (131.243.48.11) or on DECNET: SET HOST MUSE (42062). Then login to the captive account PPDS_PUBLIC (a password is not required).

Otherwise, remote interactive access can be achieved from other OpenVMS computers with DECNET access to MUSE. The remote software (20,000 blocks) can be obtained from LUGOVSKY@MX.IHEP.SU or PDG@LBL.GOV and can then be initialized by the system manager or by having each user type:

@disk:[*directory*.COMPAS.BDMS.COM]BDMSINI

In the following description, words in Typewriter Font must be typed as given. Only the letters in UPPER CASE are necessary and these must be entered in upper case. *Italic words* are variables for which the user substitutes an appropriate value, again in upper case.

- To enter the system and obtain general information, type:

PPDS

or, to open a particular database, type:

PPDS *database_name*

(*e.g.*, PPDS DOCUMENTS)

- For a short explanation of the database, type:

HELpbase

- For a list of BDMS commands, type:

?

- For an explanation of a particular BDMS command, type:

?*command_word*

(*e.g.*, ?FInd, ?HELpbase, ??)

- To see the record structure and names of keys for searching, type:

FDT

- To browse the index of a key, type:

INDEx,*key_name*

(*e.g.*, INDEx,AC)

- To search an index, type:

FInd *key_name=key_value*;**

Note the use of '**' to terminate each search statement and the use of ';' to separate data elements.

The following examples typify the FIND search command.

FInd AC=BNL;**

FInd AC=BNL; OR AC=BONN;**

Each successful search produces a list of all previous searches and labels them with a 'set number.' A previous search result can then be combined with a current search by use of set numbers:

FInd (1) and RE=PI+ P --> PI+ P;**

FInd (1) and (2) **

Note that ';' is not used in searches that only use 'sets'.

Enter DIR to get a list of these set numbers and search commands.

- To do a truncated search, use a slash after the key value:

FInd DE=HBC/**

This finds all detectors that begin with HBC.

- To do a string search, use /C after the key name:

FInd DE/C=BC;**

This finds all detectors that have BC anywhere in the name.

- The following examples are WRONG:

find ac=bnl;** (Error: uses lowercase)

FInd AC BNL;** (Error: no '=')

FInd AC=BNL ** (Error: no ';')

FInd AC=BNL OR BONN;** (Error: no ';' & no 'AC=')

- To see the results of a search with key names, type:

LISt

- Or to restrict data elements shown, append the desired key names. For example:

LISt,AC,RE,SC.

The leading comma and terminal period are required.

- Or for an attractive listing, type:

DOcument then

LOokfile

- Remote users may save the result of a search in a file by typing one of the following:

DOcument

DUmp

PRInt

The results are stored in the files DOC.DOC, DOC.DUM, or DOC.PRN. The first file contains a user-friendly listing, the second contains a highly compressed dump of each record (with data element and value), and the third contains a line-by-line decompressed version of the second file. Another file automatically created, DOC.AUD, contains a history of your commands.

2.9. Durham-RAL Databases on BDMS at Durham and CERN

Databases running under the Berkeley Database Management System (BDMS) that are menu driven with on-line help information are available on the CERN IBM/VM system and on the Durham OpenVMS system. To access the VM system on INTERNET: TELNET CERNVM.CERN.CH (128.141.2.4) and enter GIME UDISK followed by HEPDATA. To access the Durham OpenVMS system on INTERNET: TELNET DURPDG.DUR.AC.UK (129.234.8.100) or on DECNET: SET HOST DURPDG (19788). A guest account PDG, password HEPDATA, is available on this machine.

Retrieve data by using simple keyword-based searches; resulting data records can be listed on the terminal or transferred to the user's own host machine.

References:

1. S.I. Alekhin *et al.*, "A Guide to Experimental Elementary Particle Physics Literature," LBL-90 (revised 1993).
2. H. Galić *et al.*, "Current Experiments in Elementary Particle Physics," LBL-91 (revised 1994).

SUMMARY TABLES OF PARTICLE PROPERTIES

Gauge and Higgs Bosons	1191
Leptons and Quarks	1193
Mesons	1196
Baryons	1218
Searches*	1227
Tests of conservation laws	1228
Meson Quick Reference Table	1216
Baryon Quick Reference Table	1217

* There are also search limits in the Summary Tables for the Gauge and Higgs Bosons, the Leptons and Quarks, and the Mesons.

Gauge & Higgs Boson Summary Table

SUMMARY TABLES OF PARTICLE PROPERTIES

July 1994

Particle Data Group

M. Aguilar-Benitez, R.M. Barnett, C. Caso, G. Conforto, R.L. Crawford, S. Eidelman, C. Grab, D.E. Groom, A. Gurtu, K.G. Hayes, J.J. Hernandez, K. Hikasa, G. Höhler, S. Kawabata, D.M. Manley, A. Manohar, L. Montanet, R.J. Morrison, H. Murayama, K. Olive, F.C. Porter, M. Roos, R.H. Schindler, R.E. Shrock, J. Stone, N.A. Törnqvist, T.G. Trippe, C.G. Wohl, and R.L. Workman
 Technical Associates: B. Armstrong, K. Gieselmann, P. Lantero, G.S. Wagman

(Approximate closing date for data: January 1, 1994)

GAUGE AND HIGGS BOSONS

γ

$$I(J^{PC}) = 0,1(1^{-})$$

Mass $m < 3 \times 10^{-27}$ eV
 Charge $q < 2 \times 10^{-32}$ e
 Mean life $\tau =$ Stable

**g
or gluon**

$$I(J^P) = 0(1^{-})$$

Mass $m = 0$ [a]
 SU(3) color octet

W

$$J = 1$$

Charge = ± 1 e
 Mass $m = 80.22 \pm 0.26$ GeV
 $m_Z - m_W = 10.96 \pm 0.26$ GeV
 $m_{W^+} - m_{W^-} = -0.2 \pm 0.6$ GeV
 Full width $\Gamma = 2.08 \pm 0.07$ GeV

W^- modes are charge conjugates of the modes below.

W^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
$e^+\nu$	(10.8±0.4) %		40100
$\mu^+\nu$	(10.6±0.7) %		40100
$\tau^+\nu$	(10.8±1.0) %		40100
$\ell^+\nu$	[b] (10.7±0.5) %		40100
hadrons	(67.8±1.5) %		-
$\pi^+\gamma$	< 5 $\times 10^{-4}$	95%	40110

Z

$$J = 1$$

Charge = 0
 Mass $m = 91.187 \pm 0.007$ GeV [c]
 Full width $\Gamma = 2.490 \pm 0.007$ GeV
 $\Gamma(\ell^+\ell^-) = 83.84 \pm 0.27$ MeV [b]
 $\Gamma(\text{invisible}) = 498.2 \pm 4.2$ MeV [d]
 $\Gamma(\text{hadrons}) = 1740.7 \pm 5.9$ MeV
 $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-) = 1.000 \pm 0.005$
 $\Gamma(\tau^+\tau^-)/\Gamma(e^+e^-) = 0.998 \pm 0.005$ [e]
 $g_V^f = -0.0377 \pm 0.0016$
 $g_A^f = -0.5008 \pm 0.0008$

Asymmetry parameters

$A_e = 0.161 \pm 0.012$ [f] ($S = 1.7$)
 $A_\tau = 0.141 \pm 0.021$ [f] ($S = 1.2$)

Charge asymmetry at Z pole

$A_{FB}^{(0,\ell)} = (1.59 \pm 0.18) \times 10^{-2}$
 $A_{FB}^{(0,c)} = (5.8 \pm 2.2) \times 10^{-2}$
 $A_{FB}^{(0,b)} = (10.7 \pm 1.3) \times 10^{-2}$

Z DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	p (MeV/c)
e^+e^-	(3.366±0.008) %		45600
$\mu^+\mu^-$	(3.367±0.013) %		45600
$\tau^+\tau^-$	(3.360±0.015) %		45600
$\ell^+\ell^-$	[b] (3.367±0.006) %		45600
invisible	(20.01 ±0.16) %		45600
hadrons	(69.90 ±0.15) %		-
$(u\bar{u}+c\bar{c})/2$	(9.7 ±1.8) %		-
$(d\bar{d}+s\bar{s}+b\bar{b})/3$	(16.8 ±1.2) %		-
$c\bar{c}$	(11.9 ±1.4) %		-
$b\bar{b}$	(15.45 ±0.21) %		-
$\pi^0\gamma$	< 5.5 $\times 10^{-5}$	95%	45600
$\eta\gamma$	< 5.1 $\times 10^{-5}$	95%	45600
$\omega\gamma$	< 6.5 $\times 10^{-4}$	95%	45600
$\eta'(958)\gamma$	< 4.2 $\times 10^{-5}$	95%	45600
$\gamma\gamma$	< 5.5 $\times 10^{-5}$	95%	45600
$\gamma\gamma\gamma$	< 1.7 $\times 10^{-5}$	95%	45600
$\pi^\pm W^\mp$	[g] < 7 $\times 10^{-5}$	95%	10300
$\rho^\pm W^\mp$	[g] < 8.3 $\times 10^{-5}$	95%	10300
$K^0 X$	(61.5 ±0.6) %		-
$K^*(892)^+ X$	(51 ±5) %		-
ΛX	(20.9 ±0.6) %		-
$\Xi^- X$	(1.42 ±0.14) %		-
$\Sigma(1385)^\pm X$	(2.6 ±0.4) %		-
$\Xi(1530)^0 X$	(4.4 ±1.0) $\times 10^{-3}$		-
$\Omega^- X$	(3.5 ±1.0) $\times 10^{-3}$		-
$J/\psi(1S) X$	(3.8 ±0.5) $\times 10^{-3}$		-
$\chi_{c1}(1P) X$	(7.5 ±3.0) $\times 10^{-3}$		-
$(D^0/\bar{D}^0) X$	(28 ±4) %		-
$D^\pm X$	(13.9 ±2.1) %		-
$D^*(2010)^\pm X$	[g] (12.5 ±1.3) %		-
$B_s^0 X$	seen		-
anomalous γ + hadrons	[h] < 3.2 $\times 10^{-3}$	95%	-
$e^+e^-\gamma$	[h] < 5.2 $\times 10^{-4}$	95%	45600
$\mu^+\mu^-\gamma$	[h] < 5.6 $\times 10^{-4}$	95%	45600
$\tau^+\tau^-\gamma$	[h] < 7.3 $\times 10^{-4}$	95%	45600
$\ell^+\ell^-\gamma\gamma$	[i] < 6.8 $\times 10^{-6}$	95%	45600
$q\bar{q}\gamma\gamma$	[i] < 5.5 $\times 10^{-6}$	95%	-
$\nu\bar{\nu}\gamma\gamma$	[i] < 3.1 $\times 10^{-6}$	95%	45600
$e^\pm\mu^\mp$	LF [g] < 6 $\times 10^{-6}$	95%	45600
$e^\pm\tau^\mp$	LF [g] < 1.3 $\times 10^{-5}$	95%	45600
$\mu^\pm\tau^\mp$	LF [g] < 1.9 $\times 10^{-5}$	95%	45600

Gauge & Higgs Boson Summary Table

Searches for Higgs Bosons — H^0 and H^\pm

H^0 Mass $m > 58.4$ GeV, CL = 95%

H^\pm_1 In Supersymmetric Models ($m_{H^\pm_1} < m_{H^0}$) [1]

Mass $m > 44$ GeV, CL = 95% for $\tan\beta > 1$

A^0 Pseudoscalar Higgs Boson In Supersymmetric Models [1]

Mass $m > 22$ GeV, CL = 95% for $50 > \tan\beta > 1$

H^\pm Mass $m > 41.7$ GeV, CL = 95%

See the Full Listings for a Note giving details of Higgs Bosons.

Searches for Heavy Bosons Other Than Higgs Bosons

Additional W Bosons

W_R — right-handed W

Mass $m > 406$ GeV, CL = 90%

(assuming light right-handed neutrino)

W' with standard couplings decaying to $e\nu, \mu\nu$

Mass $m > 520$ GeV, CL = 95%

Additional Z Bosons

Z'_{SM} with standard couplings

Mass $m > 412$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 779$ GeV, CL = 95% (electroweak fit)

Z'_{LR} of $SU(2)_L \times SU(2)_R \times U(1)$

(with $g_L = g_R$)

Mass $m > 310$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 389$ GeV, CL = 95% (electroweak fit)

Z'_χ of $SO(10) \rightarrow SU(5) \times U(1)_\chi$

(coupling constant derived from G.U.T.)

Mass $m > 340$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 321$ GeV, CL = 95% (electroweak fit)

Z'_ψ of $E_6 \rightarrow SO(10) \times U(1)_\psi$

(coupling constant derived from G.U.T.)

Mass $m > 320$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 160$ GeV, CL = 95% (electroweak fit)

Z'_η of $E_6 \rightarrow SU(3) \times SU(2) \times U(1) \times U(1)_\eta$

(coupling constant derived from G.U.T.;

charges are $Q_\eta = \sqrt{3}/8Q_\chi - \sqrt{5}/8Q_\psi$)

Mass $m > 340$ GeV, CL = 95% ($p\bar{p}$ direct search)

Mass $m > 182$ GeV, CL = 95% (electroweak fit)

Scalar Leptoquarks

Mass $m > 120$ GeV, CL = 95% (1st generation, pair prod.)

Mass $m > 181$ GeV, CL = 95% (1st gener., single prod.)

Mass $m > 44.5$ GeV, CL = 95% (2nd gener., pair prod.)

Mass $m > 73$ GeV, CL = 95% (2nd gener., single prod.)

Mass $m > 45$ GeV, CL = 95% (3rd gener., pair prod.)

(last four limits are for charge $-1/3$, weak isoscalar)

Searches for Axions (A^0) and Other Very Light Bosons

The standard Peccei-Quinn axion is ruled out. Variants with reduced couplings or much smaller masses are constrained by various data. The Full Listings in the full Review contain a Note discussing axion searches.

The best limit for the half-life of neutrinoless double beta decay with Majoron emission is $> 7.2 \times 10^{24}$ years (CL = 90%).

NOTES

In this Summary Table:

When a quantity has "(S = ...)" to its right, the error on the quantity has been enlarged by the "scale factor" S, defined as $S = \sqrt{\chi^2/(N-1)}$, where N is the number of measurements used in calculating the quantity. We do this when $S > 1$, which often indicates that the measurements are inconsistent. When $S > 1.25$, we also show in the Full Listings an ideogram of the measurements. For more about S, see the Introduction.

A decay momentum p is given for each decay mode. For a 2-body decay, p is the momentum of each decay product in the rest frame of the decaying particle. For a 3-or-more-body decay, p is the largest momentum any of the products can have in this frame.

- [a] Theoretical value. A mass as large as a few MeV may not be precluded.
- [b] ℓ indicates each type of lepton ($e, \mu, \text{ and } \tau$), not sum over them.
- [c] The Z -boson mass listed here corresponds to a Breit-Wigner resonance parameter. It lies approximately 34 MeV above the real part of the position of the pole (in the energy plane) in the Z -boson propagator.
- [d] This partial width takes into account Z decays into $\nu\bar{\nu}$ and any other possible undetected modes.
- [e] This ratio has not been corrected for the τ mass.
- [f] Here $A \equiv 2g_V g_A / (g_V^2 + g_A^2)$.
- [g] The value is for the sum of the charge states indicated.
- [h] See the Z Full Listings for the γ energy range used in this measurement.
- [i] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.
- [j] The limits assume no invisible decays.

Lepton & Quark Summary Table

LEPTONS

Neutrinos

See the Full Listings for a Note giving details of neutrinos, masses, mixing, and the status of experimental searches.

 ν_e

$$J = \frac{1}{2}$$

Mass m : The formal upper limit, as obtained from the m^2 average (see the Full Listings), is 5.1 eV at the 95% CL. Caution is urged in interpreting this result, since the m^2 average is positive with only a 3.5% probability. If the weighted average m^2 were forced to zero, the limit would increase to 7.0 eV.

Mean life/mass, $\tau/m_{\nu_e} > 300$ s/eV, CL = 90%

Magnetic moment $\mu < 1.08 \times 10^{-9} \mu_B$, CL = 90%

 ν_μ

$$J = \frac{1}{2}$$

Mass $m < 0.27$ MeV, CL = 90%

Mean life/mass, $\tau/m_{\nu_\mu} > 15.4$ s/eV, CL = 90%

Magnetic moment $\mu < 7.4 \times 10^{-10} \mu_B$, CL = 90%

 ν_τ

$$J = \frac{1}{2}$$

Mass $m < 31$ MeV, CL = 95%

Magnetic moment $\mu < 5.4 \times 10^{-7} \mu_B$, CL = 90%

 e

$$J = \frac{1}{2}$$

Mass $m = 0.51099906 \pm 0.00000015$ MeV [a]
 $= (5.48579903 \pm 0.00000013) \times 10^{-4} u$

$(m_{e^+} - m_{e^-})/m < 4 \times 10^{-8}$, CL = 90%

$|q_{e^+} + q_{e^-}|/e < 4 \times 10^{-8}$

Magnetic moment $\mu = 1.001159652193 \pm 0.000000000010 \mu_B$

$(g_{e^+} - g_{e^-})/g_{\text{average}} = (-0.5 \pm 2.1) \times 10^{-12}$

Electric dipole moment $d = (-0.3 \pm 0.8) \times 10^{-26}$ e cm

Mean life $\tau > 2.7 \times 10^{23}$ yr, CL = 68% [b]

 μ

$$J = \frac{1}{2}$$

Mass $m = 105.658389 \pm 0.000034$ MeV [a]

$= 0.113428913 \pm 0.000000017 u$

Mean life $\tau = (2.19703 \pm 0.00004) \times 10^{-6}$ s

$\tau_{\mu^+}/\tau_{\mu^-} = 1.00002 \pm 0.00008$

$c\tau = 658.654$ m

Magnetic moment $\mu = 1.001165923 \pm 0.000000008 e\hbar/2m_\mu$

$(g_{\mu^+} - g_{\mu^-})/g_{\text{average}} = (-2.6 \pm 1.6) \times 10^{-8}$

Electric dipole moment $d = (3.7 \pm 3.4) \times 10^{-19}$ e cm

Decay parameters [c]

$\rho = 0.7518 \pm 0.0026$

$\eta = -0.007 \pm 0.013$

$\delta = 0.749 \pm 0.004$

$\xi P_\mu = 1.003 \pm 0.008$ [d]

$\xi P_\mu \delta / \rho > 0.99682$, CL = 90% [d]

$\xi' = 1.00 \pm 0.04$

$\xi'' = 0.7 \pm 0.4$

$\alpha/A = (0 \pm 4) \times 10^{-3}$

$\alpha'/A = (0 \pm 4) \times 10^{-3}$

$\beta/A = (4 \pm 6) \times 10^{-3}$

$\beta'/A = (2 \pm 6) \times 10^{-3}$

$\bar{\eta} = 0.02 \pm 0.08$

μ^+ modes are charge conjugates of the modes below.

μ^- DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$e^- \bar{\nu}_e \nu_\mu$	$\approx 100\%$		53
$e^- \bar{\nu}_e \nu_\mu \gamma$	[e] $(1.4 \pm 0.4)\%$		53
$e^- \bar{\nu}_e \nu_\mu e^+ e^-$	[f] $(3.4 \pm 0.4) \times 10^{-5}$		53

Lepton Family number (LF) violating modes

$e^- \nu_e \bar{\nu}_\mu$	LF	$ g < 1.2$	%	90%	53
$e^- \gamma$	LF	< 4.9	$\times 10^{-11}$	90%	53
$e^- e^+ e^-$	LF	< 1.0	$\times 10^{-12}$	90%	53
$e^- 2\gamma$	LF	< 7.2	$\times 10^{-11}$	90%	53

 τ

$$J = \frac{1}{2}$$

Mass $m = 1777.1^{+0.4}_{-0.5}$ MeV

Mean life $\tau = (295.6 \pm 3.1) \times 10^{-15}$ s

$c\tau = 88.6 \mu\text{m}$

Electric dipole moment $d < 5 \times 10^{-17}$ e cm, CL = 95%

Weak dipole moment $< 3.7 \times 10^{-17}$ e cm, CL = 95%

Decay parameters

See the τ Full Listings for a note concerning τ -decay parameters.

$\rho^\tau(e \text{ or } \mu) = 0.74 \pm 0.04$

$\rho^\tau(e) = 0.72 \pm 0.04$

$\rho^\tau(\mu) = 0.76 \pm 0.05$

$\xi^\tau(e \text{ or } \mu) = 0.90 \pm 0.18$

W - τ couplings $2g_A g_V / (g_A^2 + g_V^2) = 1.25^{+0.27}_{-0.24}$

τ^+ modes are charge conjugates of the modes below. " h^\pm " stands for π^\pm or K^\pm . " e " stands for e or μ . "Neutral" means neutral hadron whose decay products include γ 's and/or π^0 's.

τ^- DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Modes with one charged particle			
particle $^- \geq 0$ neutrals ν_τ ("1-prong")	(85.49 \pm 0.24) %	S=1.5	-
$\mu^- \bar{\nu}_\mu \nu_\tau$	(17.65 \pm 0.24) %	S=1.1	885
$\mu^- \bar{\nu}_\mu \nu_\tau \gamma$ ($E_\gamma > 37$ MeV)	(2.3 \pm 1.1) $\times 10^{-3}$		-
$e^- \bar{\nu}_e \nu_\tau$	(18.01 \pm 0.18) %	S=1.1	889
$h^- \geq 0$ neutrals ν_τ	(49.83 \pm 0.35) %	S=1.3	-
$h^- \nu_\tau$	(12.88 \pm 0.34) %	S=1.2	-
$\pi^- \nu_\tau$	(11.7 \pm 0.4) %	S=1.3	883
$K^- \geq 0$ neutrals ν_τ	(1.68 \pm 0.24) %		-
$K^- \nu_\tau$	(6.7 \pm 2.3) $\times 10^{-3}$	S=1.3	820
$K^- \geq 1$ neutrals ν_τ	(1.2 \pm 0.5) -0.6 %		-
$h^- \geq 1$ neutrals ν_τ	(36.9 \pm 0.4) %	S=1.3	-
$h^- \pi^0 \nu_\tau$	(25.7 \pm 0.4) %	S=1.7	-
$\pi^- \pi^0 \nu_\tau$	(25.2 \pm 0.4) %	S=1.7	878
$h^- \geq 2\pi^0 \nu_\tau$	(11.2 \pm 0.4) %	S=1.5	-
$h^- 2\pi^0 \nu_\tau$	(9.6 \pm 0.4) %	S=1.5	-
$h^- \geq 3\pi^0 \nu_\tau$	(1.48 \pm 0.26) %	S=1.7	-
$h^- 3\pi^0 \nu_\tau$	(1.28 \pm 0.24) %	S=1.7	-
$h^- 4\pi^0 \nu_\tau$	(1.9 \pm 1.1) -1.0 $\times 10^{-3}$	S=1.6	-
Modes with three charged particles			
$2h^- h^+ \geq 0$ neutrals ν_τ ("3-prong")	(14.38 \pm 0.24) %	S=1.5	-
$h^- h^- h^+ \nu_\tau$	(8.42 \pm 0.31) %	S=1.3	-
$h^- h^- h^+ \geq 1$ neutrals ν_τ	(5.63 \pm 0.30) %	S=1.2	-
$h^- h^- h^+ 2\pi^0 \nu_\tau$	(4.9 \pm 0.5) $\times 10^{-3}$		-
$\omega \pi^- \geq 0$ neutrals ν_τ	(1.6 \pm 0.4) %		-
$\omega \pi^- \nu_\tau$	(1.6 \pm 0.5) %		708
$h^- \omega \pi^0 \nu_\tau$	(4.0 \pm 0.6) $\times 10^{-3}$		-
$K^- h^+ h^- \geq 0$ neutrals ν_τ	$< 6 \times 10^{-3}$	CL=90%	-
$K^- \pi^+ \pi^- \geq 0$ neutrals ν_τ	(2.2 \pm 1.6) -1.3 $\times 10^{-3}$		-
$K^- K^+ \pi^- \nu_\tau$	(2.2 \pm 1.7) -1.1 $\times 10^{-3}$		685
Modes with five charged particles			
$3h^- 2h^+ \geq 0$ neutrals ν_τ ("5-prong")	(1.25 \pm 0.24) $\times 10^{-3}$		-
$3h^- 2h^+ \nu_\tau$	(5.6 \pm 1.6) $\times 10^{-4}$		-
$3h^- 2h^+ \pi^0 \nu_\tau$	(5.1 \pm 2.2) $\times 10^{-4}$		-

Lepton & Quark Summary Table

Miscellaneous other allowed modes			
$4h^-3h^+ \geq 0$ neutrals ν_τ	< 1.9	$\times 10^{-4}$	CL=90%
("7-prong")			
$K^*(892)^- \geq 0$ neutrals ν_τ	(1.43±0.17) %		-
$K^*(892)^- \nu_\tau$	(1.45±0.18) %		665
$K^*(892)^0 K^- \geq 0$ neutrals ν_τ	(3.2 ±1.4) $\times 10^{-3}$		-
$\bar{K}^*(892)^0 \pi^- \geq 0$ neutrals ν_τ	(3.8 ±1.7) $\times 10^{-3}$		-
$K^0 h^- \geq 0$ neutrals ν_τ	(1.30±0.30) %		-
$K^- K^0 \geq 0$ neutrals ν_τ	< 8	$\times 10^{-3}$	CL=90%
$K^0 K^- \nu_\tau$	< 2.6	$\times 10^{-3}$	CL=95%
$K^0 K^- \geq 1$ neutrals ν_τ	< 2.6	$\times 10^{-3}$	CL=95%
$K^0 h^+ h^- h^- \geq 0$ neutrals ν_τ	< 1.7	$\times 10^{-3}$	CL=95%
$K_2^*(1430)^- \nu_\tau$	< 3	$\times 10^{-3}$	CL=95%
$\eta \pi^- \geq 0$ neutrals ν_τ	< 1.3	%	CL=95%
$\eta \pi^- \nu_\tau$	< 3.4	$\times 10^{-4}$	CL=95%
$\eta \pi^- \pi^0 \nu_\tau$	(1.70±0.28) $\times 10^{-3}$		778
$\eta \pi^- \pi^0 \pi^0 \nu_\tau$	< 4.3	$\times 10^{-4}$	CL=95%
$\eta K^- \nu_\tau$	< 4.7	$\times 10^{-4}$	CL=95%
$\eta \pi^+ \pi^- \pi^- \geq 0$ neutrals ν_τ	< 3	$\times 10^{-3}$	CL=90%
$\eta \eta \pi^- \geq 0$ neutrals ν_τ	< 5	$\times 10^{-3}$	CL=90%
$\eta \eta \pi^- \nu_\tau$	< 1.1	$\times 10^{-4}$	CL=95%
$\eta \eta \pi^- \pi^0 \nu_\tau$	< 2.0	$\times 10^{-4}$	CL=95%

Lepton Family number (LF), Lepton number (L), or Baryon number (B) violating modes
(In the modes below, ℓ means a sum over e and μ modes)

L means lepton number violation (e.g. $\tau^- \rightarrow e^+ \pi^- \pi^-$). Following common usage, LF means lepton family violation and *not* lepton number violation (e.g. $\tau^- \rightarrow e^- \pi^+ \pi^-$).

$e^- \gamma$	LF	< 1.2	$\times 10^{-4}$	CL=90%	889
$\mu^- \gamma$	LF	< 4.2	$\times 10^{-6}$	CL=90%	885
$e^- \pi^0$	LF	< 1.4	$\times 10^{-4}$	CL=90%	883
$\mu^- \pi^0$	LF	< 4.4	$\times 10^{-5}$	CL=90%	880
$e^- K^0$	LF	< 1.3	$\times 10^{-3}$	CL=90%	819
$\mu^- K^0$	LF	< 1.0	$\times 10^{-3}$	CL=90%	815
$e^- \eta$	LF	< 6.3	$\times 10^{-5}$	CL=90%	804
$\mu^- \eta$	LF	< 7.3	$\times 10^{-5}$	CL=90%	800
$e^- \rho^0$	LF	< 1.9	$\times 10^{-5}$	CL=90%	723
$\mu^- \rho^0$	LF	< 2.9	$\times 10^{-5}$	CL=90%	718
$e^- K^*(892)^0$	LF	< 3.8	$\times 10^{-5}$	CL=90%	665
$\mu^- K^*(892)^0$	LF	< 4.5	$\times 10^{-5}$	CL=90%	660
$\pi^- \gamma$	L	< 2.8	$\times 10^{-4}$	CL=90%	883
$\pi^- \pi^0$	L	< 3.7	$\times 10^{-4}$	CL=90%	878
$\ell^- \ell^+ \ell^+$	LF	[h] < 3.4	$\times 10^{-5}$	CL=90%	-
$e^- e^+ e^-$	LF	< 1.3	$\times 10^{-5}$	CL=90%	889
$(e \mu \mu)^-$	LF	< 2.7	$\times 10^{-5}$	CL=90%	882
$e^- \mu^+ \mu^-$	LF	< 1.9	$\times 10^{-5}$	CL=90%	882
$e^+ \mu^- \mu^-$	LF	< 1.6	$\times 10^{-5}$	CL=90%	882
$(\mu e e)^-$	LF	< 2.7	$\times 10^{-5}$	CL=90%	885
$\mu^- e^+ e^-$	LF	< 1.4	$\times 10^{-5}$	CL=90%	885
$\mu^+ e^- e^-$	LF	< 1.4	$\times 10^{-5}$	CL=90%	885
$\mu^- \mu^+ \mu^-$	LF	< 1.7	$\times 10^{-5}$	CL=90%	873
$\ell^\pm \pi^\mp \pi^-$	LF,L	[h,i] < 6.3	$\times 10^{-5}$	CL=90%	-
$e^\mp \pi^\pm \pi^-$	LF,L	[i] < 6.0	$\times 10^{-5}$	CL=90%	877
$e^- \pi^+ \pi^-$	LF	< 2.7	$\times 10^{-5}$	CL=90%	877
$e^+ \pi^- \pi^-$	L	< 1.7	$\times 10^{-5}$	CL=90%	877
$\mu^\mp \pi^\pm \pi^-$	LF,L	[i] < 3.9	$\times 10^{-5}$	CL=90%	866
$\mu^- \pi^+ \pi^-$	LF	< 3.6	$\times 10^{-5}$	CL=90%	866
$\mu^+ \pi^- \pi^-$	L	< 3.9	$\times 10^{-5}$	CL=90%	866
$\ell^\pm \pi^\mp K^-$	LF,L	[h,i] < 1.2	$\times 10^{-4}$	CL=90%	-
$(e \pi K)^-$, all charged	LF,L	< 7.7	$\times 10^{-5}$	CL=90%	814
$e^- \pi^\pm K^\mp$	LF	[i] < 5.8	$\times 10^{-5}$	CL=90%	814
$e^- \pi^+ K^-$	LF	< 2.9	$\times 10^{-5}$	CL=90%	814
$e^- \pi^- K^+$	LF	< 5.8	$\times 10^{-5}$	CL=90%	814
$e^+ \pi^- K^-$	L	< 2.0	$\times 10^{-5}$	CL=90%	814
$(\mu \pi K)^-$, all charged	LF,L	< 7.7	$\times 10^{-5}$	CL=90%	800
$\mu^- \pi^\pm K^\mp$	LF	[i] < 7.7	$\times 10^{-5}$	CL=90%	800
$\mu^- \pi^+ K^-$	LF	< 7.7	$\times 10^{-5}$	CL=90%	800
$\mu^- \pi^- K^+$	LF	< 7.7	$\times 10^{-5}$	CL=90%	800
$\mu^+ \pi^- K^-$	L	< 4.0	$\times 10^{-5}$	CL=90%	800
$\bar{p} \gamma$	L,B	< 2.9	$\times 10^{-4}$	CL=90%	641
$\bar{p} \pi^0$	L,B	< 6.6	$\times 10^{-4}$	CL=90%	632
$\bar{p} \eta$	L,B	< 1.30	$\times 10^{-3}$	CL=90%	476
e^- light spinless boson	LF	< 3.2	$\times 10^{-3}$	CL=95%	-
μ^- light spinless boson	LF	< 6	$\times 10^{-3}$	CL=95%	-

Number of Light Neutrino Types

(including $\nu_e, \nu_\mu,$ and ν_τ)
 Number $N = 2.983 \pm 0.025$ (Standard Model fits to Z data)
 Number $N = 2.97 \pm 0.17$ (Direct measurement of invisible Z width)

Heavy Lepton Searches

- L^\pm – charged lepton**
Mass $m > 44.3$ GeV, CL = 95% $m_\nu \approx 0$
- L^\pm – stable charged heavy lepton**
Mass $m > 42.8$ GeV, CL = 95%
- L^0 – stable neutral heavy lepton**
Mass $m > 45.0$ GeV, CL = 95% (Dirac)
Mass $m > 39.5$ GeV, CL = 95% (Majorana)
- Neutral heavy lepton**
Mass $m > 19.6$ GeV, CL = 95% (all $|U_{\ell j}|^2$) (Dirac)
Mass $m > 45.7$ GeV or $m < 25$, CL = 95% ($|U_{\ell j}|^2 > 10^{-13}$) (Dirac)

Searches for Massive Neutrinos and Lepton Mixing

For excited leptons, see Compositeness Limits below.

See the Full Listings for a Note giving details of neutrinos, masses, mixing, and the status of experimental searches.

No direct, uncontested evidence for massive neutrinos or lepton mixing has been obtained. Sample limits are:

- ν oscillation: $\bar{\nu}_e \not\leftrightarrow \bar{\nu}_e$**
 $\Delta(m^2) < 0.0083$ eV², CL = 90% (if $\sin^2 2\theta = 1$)
 $\sin^2 2\theta < 0.14$, CL = 68% (if $\Delta(m^2)$ is large)
- ν oscillation: $\nu_\mu \rightarrow \nu_e$ ($\theta =$ mixing angle)**
 $\Delta(m^2) < 0.09$ eV², CL = 90% (if $\sin^2 2\theta = 1$)
 $\sin^2 2\theta < 2.5 \times 10^{-3}$, CL = 90% (if $\Delta(m^2)$ is large)

Lepton & Quark Summary Table

QUARKS

The u -, d -, and s -quark masses are estimates of so-called "current-quark masses," in a mass-independent subtraction scheme such as \overline{MS} at a scale $\mu \approx 1$ GeV. The c - and b -quark masses are estimated from charmonium, bottomonium, D , and B masses. They are the "running" masses in the \overline{MS} scheme. These can be different from the heavy quark masses obtained in potential models.

u	$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$	
	Mass $m = 2$ to 8 MeV [U]	Charge = $\frac{2}{3} e$ $I_z = +\frac{1}{2}$
	$m_u/m_d = 0.25$ to 0.70	
d	$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$	
	Mass $m = 5$ to 15 MeV [U]	Charge = $-\frac{1}{3} e$ $I_z = -\frac{1}{2}$
	$m_s/m_d = 17$ to 25	
s	$I(J^P) = 0(\frac{1}{2}^+)$	
	Mass $m = 100$ to 300 MeV [U]	Charge = $-\frac{1}{3} e$ Strangeness = -1
	$(m_s - (m_u + m_d)/2)/(m_d - m_u) = 34$ to 51	
c	$I(J^P) = 0(\frac{1}{2}^+)$	
	Mass $m = 1.0$ to 1.6 GeV	Charge = $\frac{2}{3} e$ Charm = $+1$
b	$I(J^P) = 0(\frac{1}{2}^+)$	
	Mass $m = 4.1$ to 4.5 GeV	Charge = $-\frac{1}{3} e$ Bottom = -1
Searches for t Quark	$I(J^P) = 0(\frac{1}{2}^+)$	
	Charge = $\frac{2}{3} e$	Top = $+1$

Mass $m > 62$ GeV, CL = 95% (all decays)
 Mass $m > 131$ GeV, CL = 95% (assumes $t \rightarrow Wb$ decay)
 Mass $m = 174 \pm 10^{+13}_{-12}$ GeV (top candidate events)
 Mass $m = 169^{+18}_{-18} +^{17}_{-20}$ GeV (Standard Model electroweak fit)
 The first result is from a CDF $\Gamma(W)$ measurement; the second is from a $D\bar{0}$ direct search; the third is from a CDF observation of top candidate events. CDF observes a 2.8σ effect which is not sufficient to firmly establish the existence of top but which, if interpreted as top, yields the third result. The fourth result is from a Standard Model electroweak fit to Z , W , and νN data not including direct m_t measurements. The central value assumes $m_H = 300$ GeV while the second upper (lower) error corresponds to $m_H = 1000$ (60) GeV.

Searches for b' (4th Generation) Quark

Mass $m > 85$ GeV, CL = 95% ($\rho\bar{\rho}$, charged current decays)
 Mass $m > 46.0$ GeV, CL = 95% (e^+e^- , all decays)

NOTES

In this Summary Table:

When a quantity has "(S = . . .)" to its right, the error on the quantity has been enlarged by the "scale factor" S , defined as $S = \sqrt{\chi^2/(N-1)}$, where N is the number of measurements used in calculating the quantity. We do this when $S > 1$, which often indicates that the measurements are inconsistent. When $S > 1.25$, we also show in the Full Listings an ideogram of the measurements. For more about S , see the Introduction.

A decay momentum p is given for each decay mode. For a 2-body decay, p is the momentum of each decay product in the rest frame of the decaying particle. For a 3-or-more-body decay, p is the largest momentum any of the products can have in this frame.

- [a] The masses of the e and μ are most precisely known in u (unified atomic mass units). The conversion factor to MeV, $1 u = 931.49432(28)$ MeV, is less well known than are the masses in u .
- [b] This is the best "electron disappearance" limit. The best limit for the mode $e^- \rightarrow \nu\gamma$ is $> 2.35 \times 10^{25}$ yr (CL=68%).
- [c] See the "Note on Muon Decay Parameters" in the μ Full Listings for definitions and details.
- [d] P_μ is the longitudinal polarization of the muon from pion decay. In standard $V-A$ theory, $P_\mu = 1$ and $\rho = \delta = 3/4$.
- [e] This only includes events with the γ energy > 10 MeV. Since the $e^- \bar{\nu}_e \nu_\mu$ and $e^- \bar{\nu}_e \nu_\mu \gamma$ modes cannot be clearly separated, we regard the latter mode as a subset of the former.
- [f] See the μ Full Listings for the energy limits used in this measurement.
- [g] A test of additive vs. multiplicative lepton family number conservation.
- [h] ℓ means a sum over e and μ modes.
- [i] The value is for the sum of the charge states indicated.
- [j] The ratios m_u/m_d and m_s/m_d are extracted from pion and kaon masses using chiral symmetry. The estimates of u and d masses are not without controversy and remain under active investigation. Within the literature there are even suggestions that the u quark could be essentially massless. The s -quark mass is estimated from SU(3) splittings in hadron masses.

Meson Summary Table

LIGHT UNFLAVORED MESONS ($S = C = B = 0$)

For $l = 1$ (π, b, ρ, a): $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$;
for $l = 0$ ($\eta, \eta', h, h', \omega, \phi, f, f'$): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

π^\pm

$$J^{PC} = 1^-(0^-)$$

Mass $m = 139.56995 \pm 0.00035$ MeV [a]
Mean life $\tau = (2.6030 \pm 0.0024) \times 10^{-8}$ s
 $c\tau = 7.804$ m

$\pi^\pm \rightarrow \ell^\pm \nu \gamma$ form factors [b]

$F_V = 0.017 \pm 0.008$
 $F_A = 0.0116 \pm 0.0016$ ($S = 1.3$)
 $R = 0.059^{+0.009}_{-0.008}$

π^\mp modes are charge conjugates of the modes below.

π^\pm DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	Scale factor/ Confidence level	ρ (MeV/c)
$\mu^+ \nu_\mu$	[c] (99.98770 \pm 0.00004) %			30
$\mu^+ \nu_\mu \gamma$	[d] (1.24 \pm 0.25) $\times 10^{-4}$			30
$e^+ \nu_e$	[c] (1.230 \pm 0.004) $\times 10^{-4}$			70
$e^+ \nu_e \gamma$	[d] (1.61 \pm 0.23) $\times 10^{-7}$			70
$e^+ \nu_e \pi^0$	(1.025 \pm 0.034) $\times 10^{-8}$			4
$e^+ \nu_e e^+ e^-$	(3.2 \pm 0.5) $\times 10^{-9}$			70
$e^+ \nu_e \nu \bar{\nu}$	< 5 $\times 10^{-6}$	90%		70
Lepton Family number (LF) or Lepton number (L) violating modes				
$\mu^+ \bar{\nu}_e$	L [e] < 1.5	$\times 10^{-3}$	90%	30
$\mu^+ \nu_e$	LF [e] < 8.0	$\times 10^{-3}$	90%	30
$\mu^- e^+ e^+ \nu$	LF < 1.6	$\times 10^{-6}$	90%	30

π^0

$$J^{PC} = 1^-(0^{++})$$

Mass $m = 134.9764 \pm 0.0006$ MeV [a]
 $m_{\pi^\pm} - m_{\pi^0} = 4.5936 \pm 0.0005$ MeV
Mean life $\tau = (8.4 \pm 0.6) \times 10^{-17}$ s ($S = 3.0$)
 $c\tau = 25.1$ nm

π^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	Scale factor/ Confidence level	ρ (MeV/c)
2γ	(98.798 \pm 0.032) %		S=1.1	67
$e^+ e^- \gamma$	(1.198 \pm 0.032) %		S=1.1	67
γ positronium	(1.82 \pm 0.29) $\times 10^{-9}$			67
$e^+ e^+ e^- e^-$	(3.14 \pm 0.30) $\times 10^{-5}$			67
$e^+ e^-$	(7.5 \pm 2.0) $\times 10^{-8}$			67
4γ	< 2 $\times 10^{-8}$	CL=90%		67
$\nu \bar{\nu}$	[f] < 8.3 $\times 10^{-7}$	CL=90%		67
$\nu_e \bar{\nu}_e$	< 1.7 $\times 10^{-6}$	CL=90%		67
$\nu_\mu \bar{\nu}_\mu$	< 3.1 $\times 10^{-6}$	CL=90%		67
$\nu_\tau \bar{\nu}_\tau$	< 2.1 $\times 10^{-6}$	CL=90%		67

Charge conjugation (C) or Lepton Family number (LF) violating modes

3γ	C < 3.1	$\times 10^{-8}$	CL=90%	67
$\mu^+ e^- + e^- \mu^+$	LF < 1.72	$\times 10^{-8}$	CL=90%	26

η

$$J^{PC} = 0^+(0^{-+})$$

Mass $m = 547.45 \pm 0.19$ MeV ($S = 1.6$)
Full width $\Gamma = 1.20 \pm 0.11$ keV [g] ($S = 1.8$)

G-nonconserving decay parameters [h]

$\pi^+ \pi^- \pi^0$ Left-right asymmetry = $(0.09 \pm 0.17) \times 10^{-2}$
 $\pi^+ \pi^- \pi^0$ Sextant asymmetry = $(0.18 \pm 0.16) \times 10^{-2}$
 $\pi^+ \pi^- \pi^0$ Quadrant asymmetry = $(-0.17 \pm 0.17) \times 10^{-2}$
 $\pi^+ \pi^- \gamma$ Left-right asymmetry = $(0.9 \pm 0.4) \times 10^{-2}$
 $\pi^+ \pi^- \gamma$ β (D-wave) = 0.05 ± 0.06 ($S = 1.5$)

η DECAY MODES

	Fraction (Γ_i/Γ)	Confidence level	Scale factor/ Confidence level	ρ (MeV/c)
neutral modes				
2γ	(70.8 \pm 0.8) %		S=1.2	-
$3\pi^0$	[g] (38.8 \pm 0.5) %		S=1.2	274
$\pi^0 2\gamma$	(31.9 \pm 0.4) %		S=1.2	180
$\pi^0 2\gamma$	(7.1 \pm 1.4) $\times 10^{-4}$			258
charged modes				
$\pi^+ \pi^- \pi^0$	(29.2 \pm 0.8) %		S=1.2	-
$\pi^+ \pi^- \gamma$	(23.6 \pm 0.6) %		S=1.2	175
$e^+ e^- \gamma$	(4.88 \pm 0.15) %		S=1.2	236
$\mu^+ \mu^- \gamma$	(5.0 \pm 1.2) $\times 10^{-3}$			274
$e^+ e^-$	(3.1 \pm 0.4) $\times 10^{-4}$			253
$\mu^+ \mu^-$	< 3 $\times 10^{-4}$		CL=90%	274
$\mu^+ \mu^-$	(5.7 \pm 0.8) $\times 10^{-6}$			253
$\pi^+ \pi^- e^+ e^-$	(1.3 \pm 0.8) $\times 10^{-3}$			236
$\pi^+ \pi^- 2\gamma$	< 2.1 $\times 10^{-3}$			236
$\pi^+ \pi^- \pi^0 \gamma$	< 6 $\times 10^{-4}$		CL=90%	175
$\pi^0 \mu^+ \mu^- \gamma$	< 3 $\times 10^{-6}$		CL=90%	211

**Charge conjugation (C), Parity (P), or
Charge conjugation \times Parity (CP) violating modes**

$\pi^+ \pi^-$	P, CP < 1.5	$\times 10^{-3}$		236
3γ	C < 5	$\times 10^{-4}$	CL=95%	274
$\pi^0 e^+ e^-$	C [i] < 4	$\times 10^{-5}$	CL=90%	258
$\pi^0 \mu^+ \mu^-$	C [i] < 5	$\times 10^{-6}$	CL=90%	211

$\rho(770)$

$$J^{PC} = 1^+(1^{--})$$

Mass $m = 769.9 \pm 0.8$ MeV ($S = 1.8$)
Full width $\Gamma = 151.2 \pm 1.2$ MeV
 $\Gamma_{ee} = 6.77 \pm 0.32$ keV

$\rho(770)$ DECAY MODES

	Fraction (Γ_i/Γ)	Confidence level	Scale factor/ Confidence level	ρ (MeV/c)
$\pi \pi$	~ 100 %			359

$\rho(770)^\pm$ decays

$\pi^\pm \gamma$	(4.5 \pm 0.5) $\times 10^{-4}$		S=2.2	372
$\pi^\pm \eta$	< 6 $\times 10^{-3}$		CL=84%	147
$\pi^\pm \pi^+ \pi^- \pi^0$	< 2.0 $\times 10^{-3}$		CL=84%	250

$\rho(770)^0$ decays

$\pi^+ \pi^- \gamma$	(9.9 \pm 1.6) $\times 10^{-3}$			359
$\pi^0 \gamma$	(7.9 \pm 2.0) $\times 10^{-4}$			373
$\eta \gamma$	(3.8 \pm 0.7) $\times 10^{-4}$			190
$\mu^+ \mu^-$	[j] (4.60 \pm 0.28) $\times 10^{-5}$			370
$e^+ e^-$	[j] (4.46 \pm 0.21) $\times 10^{-5}$			385
$\pi^+ \pi^- \pi^0$	< 1.2 $\times 10^{-4}$		CL=90%	320
$\pi^+ \pi^- \pi^+ \pi^-$	< 2 $\times 10^{-4}$		CL=90%	247
$\pi^+ \pi^- \pi^0 \pi^0$	< 4 $\times 10^{-5}$		CL=90%	253

$\omega(782)$

$$J^{PC} = 0^-(1^{--})$$

Mass $m = 781.94 \pm 0.12$ MeV ($S = 1.5$)
Full width $\Gamma = 8.43 \pm 0.10$ MeV
 $\Gamma_{ee} = 0.60 \pm 0.02$ keV

$\omega(782)$ DECAY MODES

	Fraction (Γ_i/Γ)	Confidence level	Scale factor/ Confidence level	ρ (MeV/c)
$\pi^+ \pi^- \pi^0$	(88.8 \pm 0.7) %			327
$\pi^0 \gamma$	(8.5 \pm 0.5) %			379
$\pi^+ \pi^-$	(2.21 \pm 0.30) %			365
neutrals (excluding $\pi^0 \gamma$)				
	(5.3 \pm 8.7 \pm 3.5) $\times 10^{-3}$			-
$\eta \gamma$	(8.3 \pm 2.1) $\times 10^{-4}$			199
$\pi^0 e^+ e^-$	(5.9 \pm 1.9) $\times 10^{-4}$			379
$\pi^0 \mu^+ \mu^-$	(9.6 \pm 2.3) $\times 10^{-5}$			349
$e^+ e^-$	(7.15 \pm 0.19) $\times 10^{-5}$			391
$\pi^+ \pi^- \pi^0 \pi^0$	< 2 %		90%	261
$\pi^+ \pi^- \gamma$	< 3.6 $\times 10^{-3}$		95%	365
$\pi^+ \pi^- \pi^+ \pi^-$	< 1 $\times 10^{-3}$		90%	256
$\pi^0 \pi^0 \gamma$	< 4 $\times 10^{-4}$		90%	367
$\mu^+ \mu^-$	< 1.8 $\times 10^{-4}$		90%	376

Meson Summary Table

 $\eta'(958)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Mass $m = 957.77 \pm 0.14$ MeV
 Full width $\Gamma = 0.201 \pm 0.016$ MeV ($S = 1.3$)

$\eta'(958)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$\pi^+\pi^-\eta$	(43.7 \pm 1.5) %	S=1.1	232
$\rho^0\gamma$	(30.2 \pm 1.3) %	S=1.1	169
$\pi^0\pi^0\eta$	(20.8 \pm 1.3) %	S=1.2	239
$\omega\gamma$	(3.02 \pm 0.30) %		160
$\gamma\gamma$	(2.12 \pm 0.13) %	S=1.2	479
$3\pi^0$	(1.55 \pm 0.26) $\times 10^{-3}$		430
$\mu^+\mu^-\gamma$	(1.04 \pm 0.26) $\times 10^{-4}$		467
$\pi^+\pi^-\pi^0$	< 5 %	CL=90%	427
$\pi^0\rho^0$	< 4 %	CL=90%	118
$\pi^+\pi^-$	< 2 %	CL=90%	458
$\pi^0e^+e^-$	< 1.3 %	CL=90%	469
ηe^+e^-	< 1.1 %	CL=90%	322
$\pi^+\pi^+\pi^-\pi^-$	< 1 %	CL=90%	372
$\pi^+\pi^+\pi^-\pi^-\text{ neutrals}$	< 1 %	CL=95%	-
$\pi^+\pi^+\pi^-\pi^-\pi^0$	< 1 %	CL=90%	298
6π	< 1 %	CL=90%	189
$\pi^+\pi^-e^+e^-$	< 6 $\times 10^{-3}$	CL=90%	458
$\pi^0\pi^0$	< 9 $\times 10^{-4}$	CL=90%	459
$\pi^0\gamma\gamma$	< 8 $\times 10^{-4}$	CL=90%	469
$4\pi^0$	< 5 $\times 10^{-4}$	CL=90%	379
3γ	< 1.0 $\times 10^{-4}$	CL=90%	479
$\mu^+\mu^-\pi^0$	< 6.0 $\times 10^{-5}$	CL=90%	445
$\mu^+\mu^-\eta$	< 1.5 $\times 10^{-5}$	CL=90%	274
$\pi^+\pi^-\gamma$ (including $\rho^0\gamma$)	(27.9 \pm 2.3) %		-
e^+e^-	< 2.1 $\times 10^{-7}$	CL=90%	479

 **$f_0(980)$
was $S(975)$**

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass $m = 980 \pm 10$ MeV
 Full width $\Gamma = 40$ to 400 MeV

$f_0(980)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\pi\pi$	(78.1 \pm 2.4) %		470
$K\bar{K}$	(21.9 \pm 2.4) %		-
$\gamma\gamma$	(1.19 \pm 0.33) $\times 10^{-5}$		490
e^+e^-	< 3 $\times 10^{-7}$	90%	490

 **$a_0(980)$
was $\delta(980)$**

$$I^G(J^{PC}) = 1^-(0^{++})$$

Mass $m = 982.4 \pm 1.4$ MeV
 Full width $\Gamma = 50$ to 300 MeV

$a_0(980)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\eta\pi$	dominant	319
$K\bar{K}$	seen	-
$\gamma\gamma$	seen	491

 $\phi(1020)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 1019.413 \pm 0.008$ MeV
 Full width $\Gamma = 4.43 \pm 0.06$ MeV
 $\Gamma_{ee} = 1.37 \pm 0.05$ keV

$\phi(1020)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
K^+K^-	(49.1 \pm 0.9) %	S=1.3	127
$K_S^0 K_S^0$	(34.3 \pm 0.7) %	S=1.2	110
$\rho\pi$	(12.9 \pm 0.7) %		181
$\pi^+\pi^-\pi^0$	(2.5 \pm 0.9) %	S=1.1	462
$\eta\gamma$	(1.28 \pm 0.06) %	S=1.2	363
$\pi^0\gamma$	(1.31 \pm 0.13) $\times 10^{-3}$		501
e^+e^-	(3.09 \pm 0.07) $\times 10^{-4}$		510
$\mu^+\mu^-$	(2.48 \pm 0.34) $\times 10^{-4}$		499
ηe^+e^-	(1.3 $^{+0.8}_{-0.6}$) $\times 10^{-4}$		363
$\pi^+\pi^-$	(8 $^{+5}_{-4}$) $\times 10^{-5}$	S=1.5	490

$\omega\gamma$	< 5 %	CL=84%	210
$\rho\gamma$	< 2 %	CL=84%	219
$\pi^+\pi^-\gamma$	< 7 $\times 10^{-3}$	CL=90%	490
$f_0(980)\gamma$	< 2 $\times 10^{-3}$	CL=90%	39
$\pi^0\pi^0\gamma$	< 1 $\times 10^{-3}$	CL=90%	492
$\pi^+\pi^-\pi^+\pi^-$	< 8.7 $\times 10^{-4}$	CL=90%	410
$\eta'(958)\gamma$	< 4.1 $\times 10^{-4}$	CL=90%	60
$\pi^+\pi^+\pi^-\pi^-\pi^0$	< 1.5 $\times 10^{-4}$	CL=95%	341
$\pi^0e^+e^-$	< 1.2 $\times 10^{-4}$	CL=90%	501
$\pi^0\eta\gamma$	< 2.5 $\times 10^{-3}$	CL=90%	346
$a_0(980)\gamma$	< 5 $\times 10^{-3}$	CL=90%	36

 $h_1(1170)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

Mass $m = 1170 \pm 20$ MeV
 Full width $\Gamma = 360 \pm 40$ MeV

$h_1(1170)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\rho\pi$	seen	310

 $b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

Mass $m = 1231 \pm 10$ MeV [k]
 Full width $\Gamma = 142 \pm 8$ MeV ($S = 1.1$)

$b_1(1235)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\omega\pi$	dominant		348
	[D/S amplitude ratio = 0.26 \pm 0.04]		
$\pi^\pm\gamma$	(1.6 \pm 0.4) $\times 10^{-3}$		608
$\eta\rho$	seen		-
$\pi^+\pi^+\pi^-\pi^0$	< 50 %	84%	536
$(K\bar{K})^\pm\pi^0$	< 8 %	90%	248
$K_S^0 K_S^0 \pi^\pm$	< 6 %	90%	238
$K_S^0 K_S^0 \pi^\pm$	< 2 %	90%	238
$\pi\phi$	< 1.5 %	84%	146

 $a_1(1260)$

$$I^G(J^{PC}) = 1^-(1^{++})$$

Mass $m = 1230 \pm 40$ MeV [k]
 Full width $\Gamma \sim 400$ MeV

$a_1(1260)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\rho\pi$	dominant		356
$\pi\gamma$	seen		607
$\pi(\pi\pi)S$ -wave	[k] < 0.7 %	90%	575

 $f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 1275 \pm 5$ MeV [k]
 Full width $\Gamma = 185 \pm 20$ MeV [k]

$f_2(1270)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$\pi\pi$	(84.9 $^{+2.5}_{-1.3}$) %	S=1.3	622
$\pi^+\pi^-2\pi^0$	(6.9 $^{+1.5}_{-2.7}$) %	S=1.4	562
$K\bar{K}$	(4.6 \pm 0.5) %	S=2.8	403
$2\pi^+2\pi^-$	(2.8 \pm 0.4) %	S=1.2	559
$\eta\eta$	(4.5 \pm 1.0) $\times 10^{-3}$	S=2.4	327
$4\pi^0$	(3.0 \pm 1.0) $\times 10^{-3}$		564
$\gamma\gamma$	(1.32 $^{+0.18}_{-0.16}$) $\times 10^{-5}$	S=1.1	637
$\eta\pi\pi$	< 8 $\times 10^{-3}$	CL=95%	475
$K_S^0 K^- \pi^+ + \text{c.c.}$	< 3.4 $\times 10^{-3}$	CL=95%	293
e^+e^-	< 9 $\times 10^{-9}$	CL=90%	637

Meson Summary Table

$f_1(1285)$		$I^G(J^{PC}) = 0^+(1^{++})$	
Mass $m = 1282 \pm 5$ MeV [k] Full width $\Gamma = 24 \pm 3$ MeV [k]			
$f_1(1285)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
4π	(29 ± 6) %		563
$\pi^0 \pi^+ \pi^-$	(15 ± 9) (8) %	S=1.1	–
$2\pi^+ 2\pi^-$	(15 ± 6) %		563
$\rho^0 \pi^+ \pi^-$	dominates $2\pi^+ 2\pi^-$		340
$4\pi^0$	< 7 × 10 ⁻⁴	CL=90%	568
$\eta \pi \pi$	(54 ± 15) %		479
$a_0(980)\pi$ [ignoring $a_0(980) \rightarrow K\bar{K}$]	(44 ± 7) %	S=1.1	234
$\eta \pi \pi$ [excluding $a_0(980)\pi$]	(10 ± 7) %	S=1.1	–
$K\bar{K}\pi$	(9.7 ± 1.6) %	S=1.2	308
$K\bar{K}^*(892)$	not seen		–
$\gamma \rho^0$	(6.6 ± 1.3) %	S=1.5	410
$\phi \gamma$	(8.0 ± 3.1) × 10 ⁻⁴		236

$\eta(1295)$		$I^G(J^{PC}) = 0^+(0^{-+})$	
Mass $m = 1295 \pm 4$ MeV Full width $\Gamma = 53 \pm 6$ MeV			
$\eta(1295)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$\eta \pi^+ \pi^-$	seen		488
$a_0(980)\pi$	seen		245

$f_0(1300)$ was $f_0(1400)$ was $\epsilon(1200)$		$I^G(J^{PC}) = 0^+(0^{++})$	
Mass $m = 1000$ -1500 MeV Full width $\Gamma = 150$ to 400 MeV $\Gamma_{\gamma\gamma} = 5.4 \pm 2.3$ keV $\Gamma_{ee} < 20$ eV, CL = 90%			
$f_0(1300)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$\pi \pi$	(93.6 ± 1.9) (1.5) %		–
$K\bar{K}$	(7.5 ± 0.9) %		–
$\eta \eta$	seen		–
$\gamma \gamma$	seen		–
$e^+ e^-$	not seen		–

$\pi(1300)$		$I^G(J^{PC}) = 1^-(0^{-+})$	
Mass $m = 1300 \pm 100$ MeV [k] Full width $\Gamma = 200$ to 600 MeV			
$\pi(1300)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$\rho \pi$	seen		406
$\pi(\pi\pi)$ S-wave	seen		612

$a_2(1320)$		$I^G(J^{PC}) = 1^-(2^{++})$	
Mass $m = 1318.4 \pm 0.6$ MeV (S = 1.1) (3π and $K^\pm K_S^0$ modes) Full width $\Gamma = 107 \pm 5$ MeV [k] ($K^\pm K_S^0$ and $\eta \pi$ modes)			
$a_2(1320)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$\rho \pi$	(70.1 ± 2.7) %	S=1.2	419
$\eta \pi$	(14.5 ± 1.2) %		535
$\omega \pi \pi$	(10.6 ± 3.2) %	S=1.3	362
$K\bar{K}$	(4.9 ± 0.8) %		437
$\eta'(958)\pi$	(5.7 ± 1.1) × 10 ⁻³		287
$\pi^\pm \gamma$	(2.8 ± 0.6) × 10 ⁻³		652
$\gamma \gamma$	(9.7 ± 1.0) × 10 ⁻⁶		659
$\pi^+ \pi^- \pi^-$	< 8 %	CL=90%	621
$e^+ e^-$	< 2.3 × 10 ⁻⁷	CL=90%	659

$f_1(1420)$ [l]		$I^G(J^{PC}) = 0^+(1^{++})$	
Mass $m = 1426.8 \pm 2.3$ MeV (S = 1.3) Full width $\Gamma = 52 \pm 4$ MeV			
$f_1(1420)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$K\bar{K}\pi$	dominant		439
$\eta \pi \pi$	possibly seen		571

$\omega(1420)$ [m]		$I^G(J^{PC}) = 0^-(1^{--})$	
Mass $m = 1419 \pm 31$ MeV Full width $\Gamma = 174 \pm 60$ MeV			
$\omega(1420)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$\rho \pi$	dominant		488

$\eta(1440)$ [n] was $\iota(1440)$		$I^G(J^{PC}) = 0^+(0^{-+})$	
Mass $m = 1420 \pm 20$ MeV [k] Full width $\Gamma = 60 \pm 30$ MeV [k]			
$\eta(1440)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$K\bar{K}\pi$	seen		433
$\eta \pi \pi$	seen		567
$a_0(980)\pi$	seen		350
4π	seen		640

$\rho(1450)$ [o]		$I^G(J^{PC}) = 1^+(1^{--})$	
Mass $m = 1465 \pm 25$ MeV [k] Full width $\Gamma = 310 \pm 60$ MeV [k]			
$\rho(1450)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\pi \pi$	seen		719
4π	seen		665
$e^+ e^-$	seen		732
$\eta \rho$	< 4 %		317
$\omega \pi$	< 2.0 %	95%	512
$\phi \pi$	< 1 %		358
$K\bar{K}$	< 1.6 × 10 ⁻³	95%	541

$f_1(1510)$		$I^G(J^{PC}) = 0^+(1^{++})$	
Mass $m = 1512 \pm 4$ MeV Full width $\Gamma = 35 \pm 15$ MeV			
$f_1(1510)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$K\bar{K}^*(892) + c.c.$	seen		292

$f_2'(1525)$		$I^G(J^{PC}) = 0^+(2^{++})$	
Mass $m = 1525 \pm 5$ MeV [k] Full width $\Gamma = 76 \pm 10$ MeV [k]			
$f_2'(1525)$ DECAY MODES	Fraction (Γ_i/Γ)		ρ (MeV/c)
$K\bar{K}$	(71.2 ± 2.0) (2.5) %		581
$\eta \eta$	(27.9 ± 2.5) (2.0) %		531
$\pi \pi$	(8.2 ± 1.6) × 10 ⁻³		750
$\gamma \gamma$	(1.23 ± 0.22) × 10 ⁻⁶		763

Meson Summary Table

 $f_0(1590)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

Seen by one group only.

Mass $m = 1581 \pm 10$ MeV

Full width $\Gamma = 180 \pm 17$ MeV ($S = 1.2$)

$f_0(1590)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\eta\eta'(958)$	dominant	234
$\eta\eta$	large	570
$4\pi^0$	large	732

 $\omega(1600)$ [ρ]

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 1662 \pm 13$ MeV

Full width $\Gamma = 280 \pm 24$ MeV

$\omega(1600)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\rho\pi$	seen	644
$\omega\pi\pi$	seen	610
e^+e^-	seen	831

 $\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

Mass $m = 1668 \pm 5$ MeV

Full width $\Gamma = 173 \pm 11$ MeV [k]

$\omega_3(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\rho\pi$	seen	647
$\omega\pi\pi$	seen	614
$b_1(1235)\pi$	possibly seen	359

 $\pi_2(1670)$

$$I^G(J^{PC}) = 1^-(2^{-+})$$

Mass $m = 1670 \pm 20$ MeV [k]

Full width $\Gamma = 240 \pm 15$ MeV [k] ($S = 1.1$)

$\Gamma_{ee} = 1.35 \pm 0.26$ keV

$\pi_2(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$f_2(1270)\pi$	$(56.2 \pm 3.2)\%$	325
$\pi^\pm\pi^+\pi^-$	$(53 \pm 4)\%$	-
$\rho\pi$	$(31 \pm 4)\%$	649
$f_0(1300)\pi$	$(8.7 \pm 3.4)\%$	-
$K\bar{K}^*(892) + c.c.$	$(4.2 \pm 1.4)\%$	453
$\gamma\gamma$	$(5.6 \pm 1.1) \times 10^{-6}$	835
$\eta\pi$	$< 5\%$	738
$\pi^\pm 2\pi^+ 2\pi^-$	$< 5\%$	734

 $\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

Not a well-established resonance.

Mass $m = 1680 \pm 50$ MeV [k]

Full width $\Gamma = 150 \pm 50$ MeV [k]

$\phi(1680)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\bar{K}^*(892) + c.c.$	dominant	462
$K_S^0 K\pi$	seen	619
$K\bar{K}$	seen	680
e^+e^-	seen	840
$\omega\pi\pi$	not seen	621

 $\rho_3(1690)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

 J^P from the 2π and $K\bar{K}$ modes.

Mass $m = 1691 \pm 5$ MeV [k] ($2\pi, K\bar{K}$, and $K\bar{K}\pi$ modes)

Full width $\Gamma = 215 \pm 20$ MeV [k] ($2\pi, K\bar{K}$, and $K\bar{K}\pi$ modes)

$\rho_3(1690)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	ρ (MeV/c)
4π	$(71.1 \pm 1.9)\%$		788
$\pi^\pm\pi^+\pi^-\pi^0$	$(67 \pm 22)\%$		788
$\pi\pi$	$(23.6 \pm 1.3)\%$		834
$\omega\pi$	$(16 \pm 6)\%$		656
$K\bar{K}\pi$	$(3.8 \pm 1.2)\%$		628
$K\bar{K}$	$(1.58 \pm 0.26)\%$	1.2	686
$\eta\pi^+\pi^-$	seen		728

 $\rho(1700)$ [ω]

$$I^G(J^{PC}) = 1^+(1^{--})$$

Mass $m = 1700 \pm 20$ MeV [k] ($\eta\rho^0$ and mixed modes)

Full width $\Gamma = 235 \pm 50$ MeV [k] ($\eta\rho^0, \pi^+\pi^-$, and mixed modes)

$\rho(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\rho\pi\pi$	dominant	640
$\rho^0\pi^+\pi^-$	large	640
$\rho^\pm\pi^\mp\pi^0$	[q] large	642
$2(\pi^+\pi^-)$	large	792
$\pi^+\pi^-$	seen	838
$K\bar{K}^*(892) + c.c.$	seen	479
$\eta\rho$	seen	533
$K\bar{K}$	seen	692
e^+e^-	seen	850

 $f_J(1710)$ was $\theta(1690)$

$$I^G(J^{PC}) = 0^+(\text{even}^{++})$$

Mass $m = 1709 \pm 5$ MeV

Full width $\Gamma = 140 \pm 12$ MeV

$f_J(1710)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\bar{K}$	seen	697
$\pi\pi$	seen	843

 $\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

Mass $m = 1854 \pm 7$ MeV

Full width $\Gamma = 87_{-23}^{+28}$ MeV ($S = 1.2$)

$\phi_3(1850)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\bar{K}$	seen	785
$K\bar{K}^*(892) + c.c.$	seen	602

 $f_2(2010)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Seen by one group only.

Mass $m = 2011_{-80}^{+60}$ MeV

Full width $\Gamma = 202 \pm 60$ MeV

$f_2(2010)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\phi\phi$	seen	-

Meson Summary Table

 $f_4(2050)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

Mass $m = 2044 \pm 11$ MeV ($S = 1.4$)Full width $\Gamma = 208 \pm 13$ MeV ($S = 1.2$)

$f_4(2050)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\omega\omega$	(26 \pm 6) %	658
$\pi\pi$	(17.0 \pm 1.5) %	1012
$K\bar{K}$	(6.8 $^{+3.4}_{-1.8}$) $\times 10^{-3}$	895
$\eta\eta$	(2.1 \pm 0.8) $\times 10^{-3}$	863
$4\pi^0$	< 1.2 %	977

 $f_2(2300)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2297 \pm 28$ MeVFull width $\Gamma = 149 \pm 40$ MeV

$f_2(2300)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi\phi$	seen	529

 $f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 2339 \pm 60$ MeVFull width $\Gamma = 319^{+80}_{-70}$ MeV

$f_2(2340)$ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\phi\phi$	seen	573

STRANGE MESONS ($S = \pm 1, C = B = 0$)

 $K^+ = u\bar{s}, K^0 = d\bar{s}, \bar{K}^0 = \bar{d}s, K^- = \bar{u}s$, similarly for K^{*} 's **K^\pm**

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 493.677 \pm 0.016$ MeV ($S = 2.8$)Mean life $\tau = (1.2371 \pm 0.0029) \times 10^{-8}$ s ($S = 2.2$) $c\tau = 3.709$ mSlope parameter g [1]

(See Full Listings for quadratic coefficients)

$$K^+ \rightarrow \pi^+\pi^+\pi^- = -0.2154 \pm 0.0035 \quad (S = 1.4)$$

$$K^- \rightarrow \pi^-\pi^-\pi^+ = -0.217 \pm 0.007 \quad (S = 2.5)$$

$$K^\pm \rightarrow \pi^\pm\pi^0\pi^0 = 0.594 \pm 0.019 \quad (S = 1.3)$$

 K^\pm decay form factors [b,s]

$$K_{e3}^+ \quad \lambda_+ = 0.0286 \pm 0.0022$$

$$K_{\mu 3}^+ \quad \lambda_+ = 0.033 \pm 0.008 \quad (S = 1.6)$$

$$K_{\mu 3}^+ \quad \lambda_0 = 0.004 \pm 0.007 \quad (S = 1.6)$$

$$K_{e3}^+ \quad |f_S/f_+| = 0.084 \pm 0.023 \quad (S = 1.2)$$

$$K_{e3}^+ \quad |f_T/f_+| = 0.38 \pm 0.11 \quad (S = 1.1)$$

$$K_{\mu 3}^+ \quad |f_T/f_+| = 0.02 \pm 0.12$$

$$K^+ \rightarrow e^+\nu_e\gamma \quad |F_A + F_V| = 0.148 \pm 0.010$$

$$K^+ \rightarrow \mu^+\nu_\mu\gamma \quad |F_A + F_V| < 0.23, \text{ CL} = 90\%$$

$$K^+ \rightarrow e^+\nu_e\gamma \quad |F_A - F_V| < 0.49$$

$$K^+ \rightarrow \mu^+\nu_\mu\gamma \quad |F_A - F_V| = -2.2 \text{ to } 0.3$$

 K^- modes are charge conjugates of the modes below.

K^+ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	p (MeV/c)
$\mu^+\nu_\mu$	(63.51 \pm 0.18) %	$S=1.3$	236
$e^+\nu_e$	(1.55 \pm 0.07) $\times 10^{-5}$		247
$\pi^+\pi^0$	(21.16 \pm 0.14) %	$S=1.1$	205
$\pi^+\pi^+\pi^-$	(5.59 \pm 0.05) %	$S=1.9$	125
$\pi^+\pi^0\pi^0$	(1.73 \pm 0.04) %	$S=1.2$	133
$\pi^0\mu^+\nu_\mu$	(3.18 \pm 0.08) %	$S=1.5$	215
Called $K_{\mu 3}^+$.			
$\pi^0e^+\nu_e$	(4.82 \pm 0.06) %	$S=1.3$	228
Called K_{e3}^+ .			
$\pi^0\pi^0e^+\nu_e$	(2.1 \pm 0.4) $\times 10^{-5}$		206
$\pi^+\pi^-e^+\nu_e$	(3.91 \pm 0.17) $\times 10^{-5}$		203
$\pi^+\pi^-\mu^+\nu_\mu$	(1.4 \pm 0.9) $\times 10^{-5}$		151
$\pi^0\pi^0\pi^0e^+\nu_e$	< 3.5 $\times 10^{-6}$	CL=90%	135
$\pi^+\gamma\gamma$	[t] < 1 $\times 10^{-6}$	CL=90%	227
$\pi^+3\gamma$	[t] < 1.0 $\times 10^{-4}$	CL=90%	227
$e^+\nu_e\nu_\nu$	< 6 $\times 10^{-5}$	CL=90%	247
$\mu^+\nu_\mu\nu_\nu$	< 6.0 $\times 10^{-6}$	CL=90%	236
$\mu^+\nu_\mu e^+e^-$	(1.06 \pm 0.32) $\times 10^{-6}$		236
$e^+\nu_e e^+e^-$	(2.1 $^{+2.1}_{-1.1}$) $\times 10^{-7}$		247
$\mu^+\nu_\mu\mu^+\mu^-$	< 4.1 $\times 10^{-7}$	CL=90%	185
$\mu^+\nu_\mu\gamma$	[t,u] (5.50 \pm 0.28) $\times 10^{-3}$		236
$\pi^+\pi^0\gamma$	[t,u] (2.75 \pm 0.15) $\times 10^{-4}$		205
$\pi^+\pi^0\gamma(\text{DE})$	[t,v] (1.8 \pm 0.4) $\times 10^{-5}$		205
$\pi^+\pi^+\pi^-\gamma$	[t,u] (1.04 \pm 0.31) $\times 10^{-4}$		125
$\pi^+\pi^0\pi^0\gamma$	[t,u] (7.4 $^{+5.5}_{-2.9}$) $\times 10^{-6}$		133
$\pi^0\mu^+\nu_\mu\gamma$	[t,u] < 6.1 $\times 10^{-5}$	CL=90%	215
$\pi^0e^+\nu_e\gamma$	[t,u] (2.62 \pm 0.20) $\times 10^{-4}$		228
$\pi^0e^+\nu_e\gamma(\text{SD})$	[w] < 5.3 $\times 10^{-5}$	CL=90%	228
$\pi^0\pi^0e^+\nu_e\gamma$	< 5 $\times 10^{-6}$	CL=90%	206

Lepton Family number (LF), Lepton number (L), $\Delta S = \Delta Q$ (SQ) violating modes, or $\Delta S = 1$ weak neutral current ($S1$) modes

$\pi^+\pi^+\pi^-\bar{\nu}_e$	SQ	< 1.2 $\times 10^{-8}$	CL=90%	203
$\pi^+\pi^+\mu^-\bar{\nu}_\mu$	SQ	< 3.0 $\times 10^{-6}$	CL=95%	151
$\pi^+e^+e^-$	$S1$	(2.74 \pm 0.23) $\times 10^{-7}$		227
$\pi^+\mu^+\mu^-$	$S1$	< 2.3 $\times 10^{-7}$	CL=90%	172
$\pi^+\nu_\nu$	$S1$	< 5.2 $\times 10^{-9}$	CL=90%	227
$\mu^-\nu_e e^+$	LF	< 2.0 $\times 10^{-8}$	CL=90%	236
$\mu^+\nu_e$	LF	[e] < 4 $\times 10^{-3}$	CL=90%	236
$\pi^+\mu^+e^-$	LF	< 2.1 $\times 10^{-10}$	CL=90%	214
$\pi^+\mu^-e^+$	LF	< 7 $\times 10^{-9}$	CL=90%	214
$\pi^-\mu^+e^+$	L	< 7 $\times 10^{-9}$	CL=90%	214
$\pi^-e^+e^+$	L	< 1.0 $\times 10^{-8}$	CL=90%	227
$\pi^-\mu^+\mu^+$	L	< 1.5 $\times 10^{-4}$	CL=90%	172
$\mu^+\bar{\nu}_e$	L	[e] < 3.3 $\times 10^{-3}$	CL=90%	236
$\pi^0e^+\bar{\nu}_e$	L	[e] < 3 $\times 10^{-3}$	CL=90%	228

 K^0

$$I(J^P) = \frac{1}{2}(0^-)$$

50% K_S , 50% K_L Mass $m = 497.672 \pm 0.031$ MeV $m_{K^0} - m_{K^\pm} = 3.995 \pm 0.034$ MeV ($S = 1.1$)

Meson Summary Table

K_S^0			
$I(J^P) = \frac{1}{2}(0^-)$			
Mean life $\tau = (0.8926 \pm 0.0012) \times 10^{-10}$ s			
$c\tau = 2.676$ cm			
CP-violation parameters [x]			
$\text{Im}(\eta_{+-0})^2 < 0.12$, CL = 90%			
$\text{Im}(\eta_{000})^2 < 0.1$, CL = 90%			
K_S^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$\pi^+\pi^-$	(68.61 ± 0.28) %	S=1.2	206
$\pi^0\pi^0$	(31.39 ± 0.28) %	S=1.2	209
$\pi^+\pi^-\gamma$	[u,y] (1.78 ± 0.05) × 10 ⁻³		206
$\gamma\gamma$	(2.4 ± 1.2) × 10 ⁻⁶		249
$\pi^+\pi^-\pi^0$	< 8.5 × 10 ⁻⁵	CL=90%	133
$3\pi^0$	< 3.7 × 10 ⁻⁵	CL=90%	139
$\pi^\pm e^\mp \nu$	[z] (6.68 ± 0.10) × 10 ⁻⁴	S=1.3	229
$\pi^\pm \mu^\mp \nu$	[z] (4.66 ± 0.07) × 10 ⁻⁴	S=1.2	216
$\Delta S = 1$ weak neutral current (SI) modes			
$\mu^+\mu^-$	SI < 3.2 × 10 ⁻⁷	CL=90%	225
e^+e^-	SI < 1.0 × 10 ⁻⁵	CL=90%	249
$\pi^0 e^+ e^-$	SI < 1.1 × 10 ⁻⁶	CL=90%	231

K_L^0			
$I(J^P) = \frac{1}{2}(0^-)$			
$m_{K_L} - m_{K_S} = (0.5333 \pm 0.0027) \times 10^{10} \hbar s^{-1}$ (S = 1.2)			
$= (3.510 \pm 0.018) \times 10^{-12}$ MeV			
Mean life $\tau = (5.17 \pm 0.04) \times 10^{-8}$ s			
$c\tau = 15.49$ m			
Slope parameter g [r]			
(See Full Listings for quadratic coefficients)			
$K_L^0 \rightarrow \pi^+\pi^-\pi^0 = 0.670 \pm 0.014$ (S = 1.6)			
K_L decay form factors [s]			
K_{e3}^0	$\lambda_+ = 0.0300 \pm 0.0016$	(S = 1.2)	
$K_{\mu 3}^0$	$\lambda_+ = 0.034 \pm 0.005$	(S = 2.3)	
$K_{\mu 3}^0$	$\lambda_0 = 0.025 \pm 0.006$	(S = 2.3)	
K_{e3}^0	$ f_S/f_+ < 0.04$, CL = 68%		
K_{e3}^0	$ f_T/f_+ < 0.23$, CL = 68%		
$K_{\mu 3}^0$	$ f_T/f_+ = 0.12 \pm 0.12$		
$K_L \rightarrow e^+e^-\gamma: \alpha_{K^*} = -0.28 \pm 0.08$			

CP-violation parameters [x]			
$\delta = (0.327 \pm 0.012)\%$			
$ \eta_{00} = (2.259 \pm 0.023) \times 10^{-3}$ (S = 1.1)			
$ \eta_{+-} = (2.269 \pm 0.023) \times 10^{-3}$ (S = 1.1)			
$ \eta_{00}/\eta_{+-} = 0.9955 \pm 0.0023$ [aa] (S = 1.8)			
$\epsilon'/\epsilon = (1.5 \pm 0.8) \times 10^{-3}$ [aa] (S = 1.8)			
$\phi_{+-} = (44.3 \pm 0.8)^\circ$			
$\phi_{00} = (43.3 \pm 1.3)^\circ$			

$\Delta S = -\Delta Q$ in K_S^0 decay			
$\text{Re } x = 0.006 \pm 0.018$ (S = 1.3)			
$\text{Im } x = -0.003 \pm 0.026$ (S = 1.2)			

K_L^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$3\pi^0$	(21.6 ± 0.8) %	S=1.5	139
$\pi^+\pi^-\pi^0$	(12.38 ± 0.21) %	S=1.5	133
$\pi^\pm \mu^\mp \nu$	[q] (27.0 ± 0.4) %	S=1.3	216
Called $K_{\mu 3}^0$.			
$\pi^\pm e^\mp \nu$	[q] (38.7 ± 0.5) %	S=1.4	229
Called $K_{e 3}^0$.			
2γ	(5.73 ± 0.27) × 10 ⁻⁴	S=2.0	249
$\pi^0 2\gamma$	[bb] (1.70 ± 0.28) × 10 ⁻⁶		231
$\pi^0 \pi^\pm e^\mp \nu$	[q] (5.18 ± 0.29) × 10 ⁻⁵		207
$(\pi \mu \text{atom}) \nu$	(1.05 ± 0.11) × 10 ⁻⁷		216
$\pi^\pm e^\mp \nu_e \gamma$	[q,u,bb] (1.3 ± 0.8) %		229
$\pi^+\pi^-\gamma$	[u,bb] (4.61 ± 0.14) × 10 ⁻⁵		206
$\pi^0 \pi^0 \gamma$	< 5.6 × 10 ⁻⁶		-

Charge conjugation × Parity (CP) or Lepton Family number (LF) violating modes, or $\Delta S = 1$ weak neutral current (SI) modes

$\pi^+\pi^-$	CPV	(2.03 ± 0.04) × 10 ⁻³	S=1.2	206
$\pi^0\pi^0$	CPV	(9.14 ± 0.34) × 10 ⁻⁴	S=1.8	209
$\mu^+\mu^-$	SI	(7.4 ± 0.4) × 10 ⁻⁹		225
$\mu^+\mu^-\gamma$	SI	(2.8 ± 2.8) × 10 ⁻⁷		225
e^+e^-	SI	< 4.1 × 10 ⁻¹¹	CL=90%	249
$e^+e^-\gamma$	SI	(9.1 ± 0.5) × 10 ⁻⁶		249
$e^+e^-\gamma\gamma$	SI [bb]	(6.6 ± 3.2) × 10 ⁻⁷		249
$\pi^+\pi^-e^+e^-$	SI	< 2.5 × 10 ⁻⁶	CL=90%	206
$\mu^+\mu^-e^+e^-$	SI	< 4.9 × 10 ⁻⁶	CL=90%	225
$e^+e^-e^+e^-$	SI [cc]	(3.9 ± 0.7) × 10 ⁻⁸		249
$\pi^0 \mu^+ \mu^-$	CP,SI [dd]	< 5.1 × 10 ⁻⁹	CL=90%	177
$\pi^0 e^+ e^-$	CP,SI [dd]	< 4.3 × 10 ⁻⁹	CL=90%	231
$\pi^0 \nu \bar{\nu}$	CP,SI [ee]	< 2.2 × 10 ⁻⁴	CL=90%	231
$e^\pm \mu^\mp$	LF [q]	< 3.3 × 10 ⁻¹¹	CL=90%	238

 $K^*(892)$

$$I(J^P) = \frac{1}{2}(1^-)$$

$K^*(892)^\pm$ mass $m = 891.59 \pm 0.24$ MeV	(S = 1.1)
$K^*(892)^0$ mass $m = 896.10 \pm 0.28$ MeV	(S = 1.4)
$K^*(892)^\pm$ full width $\Gamma = 49.8 \pm 0.8$ MeV	
$K^*(892)^0$ full width $\Gamma = 50.5 \pm 0.6$ MeV	(S = 1.1)

$K^*(892)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$K\pi$	~ 100 %		291
$K^0\gamma$	(2.30 ± 0.20) × 10 ⁻³		310
$K^\pm\gamma$	(1.01 ± 0.09) × 10 ⁻³		309
$K\pi\pi$	< 7 × 10 ⁻⁴	95%	224

 $K_1(1270)$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 1273 \pm 7$ MeV [k]
Full width $\Gamma = 90 \pm 20$ MeV [k]

$K_1(1270)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\rho$	(42 ± 6) %	76
$K_S^0(1430)\pi$	(28 ± 4) %	-
$K^*(892)\pi$	(16 ± 5) %	301
$K\omega$	(11.0 ± 2.0) %	-
$K f_0(1300)$	(3.0 ± 2.0) %	-

 $K_1(1400)$

$$I(J^P) = \frac{1}{2}(1^+)$$

Mass $m = 1402 \pm 7$ MeV
Full width $\Gamma = 174 \pm 13$ MeV (S = 1.6)

$K_1(1400)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K^*(892)\pi$	(94 ± 6) %	401
$K\rho$	(3.0 ± 3.0) %	298
$K f_0(1300)$	(2.0 ± 2.0) %	-
$K\omega$	(1.0 ± 1.0) %	285

Meson Summary Table

 $K^*(1410)$

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass $m = 1412 \pm 12$ MeV ($S = 1.1$)Full width $\Gamma = 227 \pm 22$ MeV ($S = 1.1$)

$K^*(1410)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$K^*(892)\pi$	> 40 %	95%	408
$K\pi$	(6.6 \pm 1.3) %		611
$K\rho$	< 7 %	95%	309

 $K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

Mass $m = 1429 \pm 6$ MeVFull width $\Gamma = 287 \pm 23$ MeV

$K_0^*(1430)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\pi$	(93 \pm 10) %	621

 $K_2^*(1430)$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $K_2^*(1430)^\pm$ mass $m = 1425.4 \pm 1.3$ MeV ($S = 1.1$) $K_2^*(1430)^0$ mass $m = 1432.4 \pm 1.3$ MeV $K_2^*(1430)^\pm$ full width $\Gamma = 98.4 \pm 2.3$ MeV $K_2^*(1430)^0$ full width $\Gamma = 109 \pm 5$ MeV ($S = 1.9$)

$K_2^*(1430)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$K\pi$	(49.7 \pm 1.2) %		622
$K^*(892)\pi$	(25.2 \pm 1.7) %		423
$K^*(892)\pi\pi$	(13.0 \pm 2.3) %		375
$K\rho$	(8.8 \pm 0.8) %	$S=1.2$	331
$K\omega$	(2.9 \pm 0.8) %		319
$K^+\gamma$	(2.4 \pm 0.5) $\times 10^{-3}$		627
$K\eta$	(1.4 $^{+2.8}_{-0.9}$) $\times 10^{-3}$	$S=1.1$	492
$K\omega\pi$	< 7.2 $\times 10^{-4}$	CL=95%	110
$K^0\gamma$	< 9 $\times 10^{-4}$	CL=90%	631

 $K^*(1680)$

$$I(J^P) = \frac{1}{2}(1^-)$$

Mass $m = 1714 \pm 20$ MeV ($S = 1.1$)Full width $\Gamma = 323 \pm 110$ MeV ($S = 4.2$)

$K^*(1680)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\pi$	(38.7 \pm 2.5) %	779
$K\rho$	(31.4 $^{+4.7}_{-2.1}$) %	571
$K^*(892)\pi$	(29.9 $^{+2.2}_{-4.7}$) %	615

 **$K_2(1770)$ [η']
was $L(1770)$**

$$I(J^P) = \frac{1}{2}(2^-)$$

Mass $m = 1773 \pm 8$ MeVFull width $\Gamma = 186 \pm 14$ MeV

$K_2(1770)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\pi\pi$		–
$K_2^*(1430)\pi$	dominant	287
$K^*(892)\pi$	seen	653
$Kf_2(1270)$	seen	–
$K\phi$	seen	441
$K\omega$	seen	608

 $K_3^*(1780)$

$$I(J^P) = \frac{1}{2}(3^-)$$

Mass $m = 1770 \pm 10$ MeV ($S = 1.7$)Full width $\Gamma = 164 \pm 17$ MeV ($S = 1.1$)

$K_3^*(1780)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$K\rho$	(45 \pm 4) %	$S=1.4$	612
$K^*(892)\pi$	(27.3 \pm 3.2) %	$S=1.5$	651
$K\pi$	(19.3 \pm 1.0) %		810
$K\eta$	(8.0 \pm 1.5) %	$S=1.4$	715
$K_2^*(1430)\pi$	< 21 %	CL=95%	284

 $K_2(1820)$

$$I(J^P) = \frac{1}{2}(2^-)$$

Mass $m = 1816 \pm 13$ MeVFull width $\Gamma = 276 \pm 35$ MeV

$K_2(1820)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\phi$	possibly seen	481
$K_2^*(1430)\pi$	seen	325
$K^*(892)\pi$	seen	680
$Kf_2(1270)$	seen	186
$K\omega$	seen	638

 $K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

Mass $m = 2045 \pm 9$ MeV ($S = 1.1$)Full width $\Gamma = 198 \pm 30$ MeV

$K_4^*(2045)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$K\pi$	(9.9 \pm 1.2) %	958
$K^*(892)\pi\pi$	(9 \pm 5) %	800
$K^*(892)\pi\pi\pi$	(7 \pm 5) %	764
$\rho K\pi$	(5.7 \pm 3.2) %	742
$\omega K\pi$	(5.0 \pm 3.0) %	736
$\phi K\pi$	(2.8 \pm 1.4) %	591
$\phi K^*(892)$	(1.4 \pm 0.7) %	363

Meson Summary Table

CHARMED MESONS (C = ±1)

$D^+ = c\bar{d}$, $D^0 = c\bar{u}$, $\bar{D}^0 = \bar{c}u$, $D^- = \bar{c}d$, similarly for D^{*s}

D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1869.4 \pm 0.4$ MeV

Mean life $\tau = (1.057 \pm 0.015) \times 10^{-12}$ s

$c\tau = 317$ μm

D^- modes are charge conjugates of the modes below.

D⁺ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Inclusive modes			
e^+ anything	(17.2 ± 1.9) %	—	—
K^- anything	(24.2 ± 2.8) %	S=1.4	—
\bar{K}^0 anything + K^0 anything	(59 ± 7) %	—	—
K^+ anything	(5.8 ± 1.4) %	—	—
η anything	[gg] < 13 %	CL=90%	—
Leptonic and semileptonic modes			
$\mu^+ \nu_\mu$	< 7.2 × 10 ⁻⁴	CL=90%	932
$\bar{K}^0 e^+ \nu_e$	[hh] (6.7 ± 0.8) %	—	868
$\bar{K}^0 e^+ \nu_e$	(6.6 ± 0.9) %	—	868
$\bar{K}^0 \mu^+ \nu_\mu$	(7.0 $\frac{+3.0}{-2.0}$) %	—	865
$\bar{K}^0 \ell^+ \nu_\ell$	(6.7 ± 3.5) %	—	—
$K^- \pi^+ e^+ \nu_e$	(4.2 $\frac{+0.9}{-0.7}$) %	—	863
$\bar{K}^*(892)^0 e^+ \nu_e$	(3.2 ± 0.33) %	—	720
$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 × 10 ⁻³	CL=90%	863
$K^- \pi^+ \mu^+ \nu_\mu$	(3.2 ± 1.7) %	—	851
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(3.0 ± 0.4) %	—	715
$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(2.7 ± 1.1) × 10 ⁻³	—	851
$(\bar{K}^*(892)^0 \pi^0 e^+ \nu_e)$	< 1.2 %	CL=90%	714
$(\bar{K}^*(892)^0 \pi^0 e^+ \nu_e \text{ non-}\bar{K}^*(892))$	< 9 × 10 ⁻³	CL=90%	846
$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.4 × 10 ⁻³	CL=90%	825
$\pi^0 \ell^+ \nu_\ell$	[ij] (5.7 ± 2.2) × 10 ⁻³	—	—
Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.			
$\bar{K}^*(892)^0 e^+ \nu_e$	[hh] (4.8 ± 0.4) %	—	720
$\bar{K}^*(892)^0 e^+ \nu_e$	(4.8 ± 0.5) %	—	720
$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(4.5 ± 0.6) %	S=1.1	715
$\rho^0 e^+ \nu_e$	< 3.7 × 10 ⁻³	CL=90%	776
$\rho^0 \mu^+ \nu_\mu$	(2.0 $\frac{+1.5}{-1.3}$) × 10 ⁻³	—	772
$\phi e^+ \nu_e$	< 2.09 %	CL=90%	657
$\phi \mu^+ \nu_\mu$	< 3.72 %	CL=90%	651
$\eta'(958) \mu^+ \nu_\mu$	< 9 × 10 ⁻³	CL=90%	684
Hadronic modes with one or three K's			
$\bar{K}^0 \pi^+$	(2.74 ± 0.29) %	—	862
$K^- \pi^+ \pi^+$	[W] (9.1 ± 0.6) %	—	845
$\bar{K}^*(892)^0 \pi^+$	(1.5 ± 0.3) %	—	712
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	—	—	—
$\bar{K}_0(1430)^0 \pi^+$	(2.3 ± 0.3) %	—	368
$\times B(\bar{K}^*(1430)^0 \rightarrow K^- \pi^+)$	—	—	—
$\bar{K}^*(1680)^0 \pi^+$	(2.6 ± 1.3) × 10 ⁻³	—	65
$\times B(\bar{K}^*(1680)^0 \rightarrow K^- \pi^+)$	—	—	—
$K^- \pi^+ \pi^+$ nonresonant	(7.3 ± 1.4) %	—	845
$\bar{K}^0 \pi^+ \pi^0$	[W] (9.7 ± 3.0) %	S=1.1	845
$\bar{K}^0 \rho^+$	(6.6 ± 2.5) %	—	680
$\bar{K}^*(892)^0 \pi^+$	(0.7 ± 0.2) %	—	712
$\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$	—	—	—
$\bar{K}^0 \pi^+ \pi^0$ nonresonant	(1.3 ± 1.1) %	—	845

$K^- \pi^+ \pi^+ \pi^0$	[W] (6.4 ± 1.1) %	816
$\bar{K}^*(892)^0 \rho^+$ total	(1.4 ± 0.9) %	423
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	—	—
$\bar{K}_1(1400)^0 \pi^+$	(2.2 ± 0.6) %	390
$\times B(\bar{K}_1(1400)^0 \rightarrow K^- \pi^+ \pi^0)$	—	—
$K^- \rho^+ \pi^+$ total	(3.1 ± 1.1) %	616
$\bar{K}^*(892)^0 \pi^+ \pi^0$ total	(4.5 ± 0.9) %	687
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	—	—
$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body	(2.8 ± 0.9) %	687
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	—	—
$K^*(892)^- \pi^+ \pi^+$ 3-body	(1.4 ± 0.6) %	688
$\times B(K^{*-} \rightarrow K^- \pi^0)$	—	—
$K^- \pi^+ \pi^+ \pi^0$ nonresonant	[kk] (1.2 ± 0.6) %	816
$\bar{K}^0 \pi^+ \pi^+ \pi^-$	[W] (7.0 ± 1.0) %	814
$\bar{K}^0 a_1(1260)^+$	(4.0 ± 0.8) %	328
$\times B(a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-)$	—	—
$\bar{K}_1(1400)^0 \pi^+$	(2.2 ± 0.6) %	390
$\times B(\bar{K}_1(1400)^0 \rightarrow \bar{K}^0 \pi^+ \pi^-)$	—	—
$K^*(892)^- \pi^+ \pi^+$ 3-body	(1.4 ± 0.6) %	688
$\times B(K^{*-} \rightarrow \bar{K}^0 \pi^-)$	—	—
$\bar{K}^0 \rho^0 \pi^+$ total	(4.2 ± 0.9) %	614
$\bar{K}^0 \pi^+ \pi^+ \pi^-$ nonresonant	(8 ± 4) × 10 ⁻³	814
$K^- \pi^+ \pi^+ \pi^+$	(8.2 ± 1.4) × 10 ⁻³	772
$\bar{K}^*(892)^0 \pi^+ \pi^+$	(6.8 ± 1.8) × 10 ⁻³	642
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	—	—
$\bar{K}^*(892)^0 \rho^0 \pi^+$	(5.1 ± 2.2) × 10 ⁻³	242
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	—	—
$K^- \pi^+ \pi^+ \pi^0 \pi^0$	(2.2 $\frac{+5.0}{-0.9}$) %	775
$\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^-$	(5.4 $\frac{+3.0}{-1.4}$) %	773
$\bar{K}^0 \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(8 ± 7) × 10 ⁻⁴	714
$K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0$	(2.0 ± 1.8) × 10 ⁻³	718
$\bar{K}^0 \bar{K}^0 K^+$	(3.1 ± 0.7) %	545

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\bar{K}^0 \rho^+$	(6.6 ± 2.5) %	680	
$\bar{K}^0 a_1(1260)^+$	(8.1 ± 1.7) %	328	
$\bar{K}^0 a_2(1320)^+$	< 3 × 10 ⁻³	CL=90% 199	
$\bar{K}^*(892)^0 \pi^+$	(2.2 ± 0.4) %	712	
$\bar{K}^*(892)^0 \rho^+$ total	(2.1 ± 1.4) %	423	
$\bar{K}^*(892)^0 \rho^+$ S-wave	[kk] (1.7 ± 1.6) %	423	
$\bar{K}^*(892)^0 \rho^+$ P-wave	< 1 × 10 ⁻³	CL=90% 423	
$\bar{K}^*(892)^0 \rho^+$ D-wave	(10 ± 7) × 10 ⁻³	423	
$\bar{K}^*(892)^0 \rho^+$ D-wave longitudinal	< 7 × 10 ⁻³	CL=90% 423	
$\bar{K}_1(1270)^0 \pi^+$	< 7 × 10 ⁻³	CL=90% 487	
$\bar{K}_1(1400)^0 \pi^+$	(5.0 ± 1.3) %	390	
$\bar{K}^*(1410)^0 \pi^+$	< 7 × 10 ⁻³	CL=90% 382	
$\bar{K}_0^*(1430)^0 \pi^+$	(3.4 ± 0.4) %	368	
$\bar{K}^*(1680)^0 \pi^+$	(1.0 ± 0.5) %	65	
$\bar{K}^*(892)^0 \pi^+ \pi^0$ total	(6.7 ± 1.4) %	687	
$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body	(4.2 ± 1.4) %	687	
$K^*(892)^- \pi^+ \pi^+$ 3-body	(2.1 ± 0.9) %	688	
$K^- \rho^+ \pi^+$ total	(3.1 ± 1.1) %	616	
$K^- \rho^+ \pi^+$ 3-body	(1.1 ± 0.4) %	616	
$\bar{K}^0 \rho^0 \pi^+$ total	(4.2 ± 0.9) %	CL=90% 614	
$\bar{K}^0 \rho^0 \pi^+$ 3-body	(5 ± 5) × 10 ⁻³	614	
$\bar{K}^0 f_0(980) \pi^+$	< 5 × 10 ⁻³	CL=90% 461	
$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$	(1.02 ± 0.27) %	642	
$\bar{K}^*(892)^0 \rho^0 \pi^+$	(7.7 ± 3.3) × 10 ⁻³	242	
Plionic modes			
$\pi^+ \pi^0$	(2.5 ± 0.7) × 10 ⁻³	925	
$\pi^+ \pi^+ \pi^-$	(3.2 ± 0.6) × 10 ⁻³	908	
$\rho^0 \pi^+$	< 1.4 × 10 ⁻³	CL=90% 769	
$\pi^+ \pi^+ \pi^-$ nonresonant	(2.5 ± 0.7) × 10 ⁻³	908	
$\pi^+ \pi^+ \pi^- \pi^0$	(1.9 $\frac{+1.5}{-1.2}$) %	883	
$\eta \pi^+ \times B(\eta \rightarrow \pi^+ \pi^- \pi^0)$	(1.8 ± 0.6) × 10 ⁻³	848	
$\omega \pi^+ \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	< 6 × 10 ⁻³	CL=90% 764	
$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(1.0 $\frac{+0.8}{-0.7}$) × 10 ⁻³	845	
$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	(2.9 $\frac{+2.9}{-2.0}$) × 10 ⁻³	799	

Meson Summary Table

Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\rho^0 \pi^+$	< 1.4	$\times 10^{-3}$	CL=90%	769
$\eta \pi^+$	(7.5 \pm 2.5)	$\times 10^{-3}$		848
$\omega \pi^+$	< 7	$\times 10^{-3}$	CL=90%	764
$\eta \rho^+$	< 1.2	%	CL=90%	658
$\eta'(958) \pi^+$	< 9	$\times 10^{-3}$	CL=90%	680
$\eta'(958) \rho^+$	< 1.5	%	CL=90%	355

Hadronic modes with two K^* 's

$\bar{K}^0 K^+$	(7.8 \pm 1.7)	$\times 10^{-3}$		792
$K^+ K^- \pi^+$	(1.13 \pm 0.13)	%		744
$\phi \pi^+ \times B(\phi \rightarrow K^+ K^-)$	(3.3 \pm 0.4)	$\times 10^{-3}$		647
$\bar{K}^*(892)^0 K^+$	(3.4 \pm 0.7)	$\times 10^{-3}$		610
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$				
$K^+ K^- \pi^+$ nonresonant	(4.6 \pm 0.9)	$\times 10^{-3}$		744
$K^+ K^- \pi^+$ nonresonant				682
$\phi \pi^+ \pi^0 \times B(\phi \rightarrow K^+ K^-)$	(1.2 \pm 0.5)	%		619
$\phi \rho^+ \times B(\phi \rightarrow K^+ K^-)$	< 7	$\times 10^{-3}$	CL=90%	268
$K^+ K^- \pi^+ \pi^0$ non- ϕ	(1.5 \pm 0.7)	%		682
$K^+ \bar{K}^0 \pi^+ \pi^-$	< 2	%	CL=90%	678
$K^0 K^- \pi^+ \pi^+$	(1.0 \pm 0.6)	%		678
$K^*(892)^+ \bar{K}^*(892)^0$	(1.2 \pm 0.5)	%		273
$\times B^2(K^* \rightarrow K \pi^+)$				
$K^0 K^- \pi^+ \pi^+ \text{ non-} K^* \bar{K}^{*0}$	< 7.9	$\times 10^{-3}$	CL=90%	678
$K^+ K^- \pi^+ \pi^+ \pi^-$				600
$\phi \pi^+ \pi^+ \pi^-$	< 1	$\times 10^{-3}$	CL=90%	566
$\times B(\phi \rightarrow K^+ K^-)$				
$K^+ K^- \pi^+ \pi^+ \pi^-$ nonresonant	< 3	%	CL=90%	600

Fractions of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

$\phi \pi^+$	(6.7 \pm 0.8)	$\times 10^{-3}$		647
$\bar{K}^*(892)^0 K^+$	(5.1 \pm 1.0)	$\times 10^{-3}$		610
$\phi \pi^+ \pi^0$	(2.3 \pm 1.0)	%		619
$\phi \rho^+$	< 1.5	%	CL=90%	268
$K^*(892)^+ \bar{K}^*(892)^0$	(2.6 \pm 1.1)	%		273
$\phi \pi^+ \pi^+ \pi^-$	< 2	$\times 10^{-3}$	CL=90%	566

Doubly Cabibbo suppressed (DC) modes, $\Delta C = 1$ weak neutral current (CI) modes, or Lepton Family number (LF) or Lepton number (L) violating modes

$K^+ \pi^+ \pi^-$	DC	< 5	$\times 10^{-3}$	CL=90%	845
$K^+ K^+ K^-$	DC	(5.2 \pm 2.0)	$\times 10^{-3}$		550
ϕK^+	DC	(3.9 \pm 2.2)	$\times 10^{-4}$		527
$\pi^+ e^+ e^-$	CI	< 2.5	$\times 10^{-3}$	CL=90%	929
$\pi^+ \mu^+ \mu^-$	CI	< 2.9	$\times 10^{-3}$	CL=90%	917
$K^+ e^+ e^-$	[II]	< 4.8	$\times 10^{-3}$	CL=90%	870
$K^+ \mu^+ \mu^-$	[II]	< 9.2	$\times 10^{-3}$	CL=90%	856
$\pi^+ e^+ \mu^-$	LF	< 3.8	$\times 10^{-3}$	CL=90%	926
$\pi^+ e^+ \mu^-$	LF	< 3.3	$\times 10^{-3}$	CL=90%	926
$\pi^+ e^+ \mu^-$	LF	< 3.3	$\times 10^{-3}$	CL=90%	926
$K^+ e^+ \mu^-$	LF	< 3.4	$\times 10^{-3}$	CL=90%	866
$K^+ e^- \mu^+$	LF	< 3.4	$\times 10^{-3}$	CL=90%	866
$\pi^- e^+ e^+$	L	< 4.8	$\times 10^{-3}$	CL=90%	929
$\pi^- \mu^+ \mu^+$	L	< 6.8	$\times 10^{-3}$	CL=90%	917
$\pi^- e^+ \mu^+$	L	< 3.7	$\times 10^{-3}$	CL=90%	926
$K^- e^+ e^+$	L	< 9.1	$\times 10^{-3}$	CL=90%	870
$K^- \mu^+ \mu^+$	L	< 4.3	$\times 10^{-3}$	CL=90%	856
$K^- e^+ \mu^+$	L	< 4.0	$\times 10^{-3}$	CL=90%	866

 D^0

$$I(J^P) = \frac{1}{2}(0^-)$$

Mass $m = 1864.6 \pm 0.5$ MeV

$$|m_{D_1^0} - m_{D_2^0}| < 20 \times 10^{10} \hbar s^{-1}, \text{ CL} = 90\% [mm]$$

$$m_{D^\pm} - m_{D^0} = 4.78 \pm 0.10 \text{ MeV}$$

$$\text{Mean life } \tau = (0.415 \pm 0.004) \times 10^{-12} \text{ s}$$

$$c\tau = 124.4 \mu\text{m}$$

$$|\tau_{D_1^0} - \tau_{D_2^0}|/\tau_{D^0} < 0.17, \text{ CL} = 90\% [mm]$$

$$\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+) < 0.0037, \text{ CL} = 90\%$$

$$\Gamma(\mu^- X \text{ (via } \bar{D}^0))/\Gamma(\mu^+ X) < 0.0056, \text{ CL} = 90\%$$

$$[\Gamma(D^0 \rightarrow K^+ K^-) - \Gamma(\bar{D}^0 \rightarrow K^+ K^-)]/\text{sum} < 0.45, \text{ CL} = 90\%$$

 \bar{D}^0 modes are charge conjugates of the modes below.

D^0 DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Inclusive modes			
e^+ anything	(7.7 \pm 1.2) %	S=1.1	-
μ^+ anything	(10.0 \pm 2.6) %		-
K^- anything	(53 \pm 4) %	S=1.3	-
\bar{K}^0 anything + K^0 anything	(42 \pm 5) %		-
K^+ anything	(3.4 \pm 0.6) %		-
η anything	[gg] < 13 %	CL=90%	-
Semileptonic modes			
$K^- e^+ \nu_e$	[hh] (3.68 \pm 0.21) %	S=1.1	867
$K^- e^+ \nu_e$	(3.80 \pm 0.22) %	S=1.1	867
$K^- \mu^+ \nu_\mu$	(3.2 \pm 0.4) %		864
$K^- \pi^0 e^+ \nu_e$	[nn] (1.6 \pm 1.3) %		861
$\bar{K}^0 \pi^- e^+ \nu_e$	[nn] (2.8 \pm 1.7) %		860
$\bar{K}^*(892)^- e^+ \nu_e$	(1.3 \pm 0.3) %		719
$\times B(K^{*-} \rightarrow \bar{K}^0 \pi^+)$			
$\bar{K}^*(892)^0 \pi^- e^+ \nu_e$	[oo] < 1.3 %	CL=90%	709
$K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.2 $\times 10^{-3}$	CL=90%	821
$(\bar{K}^*(892) \pi)^- \mu^+ \nu_\mu$	< 1.4 $\times 10^{-3}$	CL=90%	694
$\pi^- e^+ \nu_e$	(3.9 \pm 2.3) $\times 10^{-3}$		927

A fraction of the following resonance mode has already appeared above as a submode of a particular charged-particle mode.

$K^*(892)^- e^+ \nu_e$	(2.0 \pm 0.4) %		719
Hadronic modes with one or three K^*'s			
$K^- \pi^+$	(4.01 \pm 0.14) %		861
$\bar{K}^0 \pi^0$	(2.05 \pm 0.26) %	S=1.1	860
$\bar{K}^0 \pi^+ \pi^-$	[ll] (5.3 \pm 0.6) %	S=1.2	842
$\bar{K}^0 \rho^0$	(1.10 \pm 0.18) %		676
$\bar{K}^0 f_0(980)$	(2.4 \pm 1.0) $\times 10^{-3}$		549
$\times B(f_0 \rightarrow \pi^+ \pi^-)$			
$\bar{K}^0 f_2(1270)$	(2.6 \pm 1.2) $\times 10^{-3}$		263
$\times B(f_2 \rightarrow \pi^+ \pi^-)$			
$\bar{K}^0 f_0(1300)$	(4.3 \pm 1.7) $\times 10^{-3}$		223
$\times B(f_0 \rightarrow \pi^+ \pi^-)$			
$K^*(892)^- \pi^+$	(3.3 \pm 0.4) %		711
$\times B(K^{*-} \rightarrow \bar{K}^0 \pi^-)$			
$K_0^*(1430)^- \pi^+$	(7 \pm 3) $\times 10^{-3}$		364
$\times B(K_0^*(1430)^- \rightarrow \bar{K}^0 \pi^-)$			
$\bar{K}^0 \pi^+ \pi^-$ nonresonant	(1.43 \pm 0.26) %		842
$K^- \pi^+ \pi^0$	[ll] (13.8 \pm 1.0) %	S=1.1	844
$K^- \rho^+$	(10.4 \pm 1.3) %		678
$K^*(892)^- \pi^+$	(1.6 \pm 0.2) %		711
$\times B(K^{*-} \rightarrow K^- \pi^0)$			
$\bar{K}^*(892)^0 \pi^0$	(2.0 \pm 0.3) %		709
$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$			
$K^- \pi^+ \pi^0$ nonresonant	(6.0 \pm 2.7) $\times 10^{-3}$		844
$\bar{K}^0 \pi^0 \pi^0$			843
$\bar{K}^*(892)^0 \pi^0$	(1.0 \pm 0.2) %		709
$\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$			
$\bar{K}^0 \pi^0 \pi^0$ nonresonant	(7.6 \pm 2.1) $\times 10^{-3}$		843

Meson Summary Table

$K^- \pi^+ \pi^+ \pi^-$	[<i>ll</i>] (8.1 ± 0.5) %	812
$K^- \pi^+ \rho^0$ total	(6.8 ± 0.5) %	612
$K^- \pi^+ \rho^0$ 3-body	(5.1 ± 2.3) × 10 ⁻³	612
$\bar{K}^*(892)^0 \rho^0$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(1.1 ± 0.3) %	418
$K^- a_1(1260)^+$ × B($a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-$)	(3.9 ± 0.6) %	327
$\bar{K}^*(892)^0 \pi^+ \pi^-$ total	(1.6 ± 0.4) %	683
× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body	(1.01 ± 0.22) %	683
× B($\bar{K}^{*0} \rightarrow K^- \pi^+$)		
$K_1(1270)^- \pi^+$ × B($K_1(1270)^- \rightarrow K^- \pi^+ \pi^-$)	(3.5 ± 1.1) × 10 ⁻³	483
$K^- \pi^+ \pi^+ \pi^-$ nonresonant	(1.89 ± 0.28) %	812
$\bar{K}^0 \pi^+ \pi^- \pi^0$	[<i>ll</i>] (9.8 ± 1.4) %	812
$\bar{K}^0 \eta \times B(\eta \rightarrow \pi^+ \pi^- \pi^0)$	(1.61 ± 0.26) × 10 ⁻³	772
$\bar{K}^0 \omega \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	(1.8 ± 0.4) %	670
$K^*(892)^- \rho^+$ × B($K^{*-} \rightarrow \bar{K}^0 \pi^-$)	(3.9 ± 1.6) %	422
$\bar{K}^*(892)^0 \rho^0$ × B($\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0$)	(5.3 ± 1.4) × 10 ⁻³	418
$K_1(1270)^- \pi^+$ × B($K_1(1270)^- \rightarrow \bar{K}^0 \pi^- \pi^0$)	[<i>kk</i>] (5.0 ± 1.5) × 10 ⁻³	483
$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body	(5.1 ± 1.1) × 10 ⁻³	683
× B($\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0$)		
$\bar{K}^0 \pi^+ \pi^- \pi^0$ nonresonant	(2.1 ± 2.1) %	812
$K^- \pi^+ \pi^0 \pi^0$	(15 ± 5) %	815
$K^- \pi^+ \pi^+ \pi^- \pi^0$	(4.3 ± 0.4) %	771
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(1.3 ± 0.6) %	641
$\bar{K}^*(892)^0 \eta$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) × B($\eta \rightarrow \pi^+ \pi^- \pi^0$)	(3.0 ± 0.8) × 10 ⁻³	580
$K^- \pi^+ \omega \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	(2.8 ± 0.5) %	605
$\bar{K}^*(892)^0 \omega$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$) × B($\omega \rightarrow \pi^+ \pi^- \pi^0$)	(7 ± 3) × 10 ⁻³	406
$\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^-$	(5.6 ± 1.7) × 10 ⁻³	768
$\bar{K}^0 \pi^+ \pi^- \pi^0 \pi^0 (\pi^0)$	(10.6 ± 7.3) %	771
$\bar{K}^0 K^+ K^-$	(9.1 ± 1.2) × 10 ⁻³	544
$\bar{K}^0 \phi \times B(\phi \rightarrow K^+ K^-)$	(4.2 ± 0.6) × 10 ⁻³	520
$\bar{K}^0 K^+ K^-$ non- ϕ	(4.9 ± 0.9) × 10 ⁻³	544
$K_S^0 K_S^0 K_S^0$	(8.6 ± 2.5) × 10 ⁻⁴	538
$K^+ K^- \bar{K}^0 \pi^0$	(7.2 ± 4.8) × 10 ⁻³	435

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\bar{K}^*(892)\rho$ submodes only appear below.)

$\bar{K}^0 \eta$	(6.8 ± 1.1) × 10 ⁻³	772
$\bar{K}^0 \rho^0$	(1.10 ± 0.18) %	676
$K^- \rho^+$	(10.4 ± 1.3) %	679
$\bar{K}^0 \omega$	(2.0 ± 0.4) %	670
$\bar{K}^0 \eta'(958)$	(1.66 ± 0.29) %	565
$\bar{K}^0 f_0(980)$	(4.6 ± 2.0) × 10 ⁻³	549
$\bar{K}^0 \phi$	(8.3 ± 1.2) × 10 ⁻³	520
$K^- a_1(1260)^+$	(7.9 ± 1.2) %	327
$\bar{K}^0 a_1(1260)^0$	< 1.9 %	322
$\bar{K}^0 f_2(1270)$	(4.6 ± 2.1) × 10 ⁻³	263
$\bar{K}^0 f_0(1300)$	(6.9 ± 2.7) × 10 ⁻³	223
$K^- a_2(1320)^+$	< 2 × 10 ⁻³	197
$K^*(892)^- \pi^+$	(4.9 ± 0.6) %	711
$\bar{K}^*(892)^0 \pi^0$	(3.0 ± 0.4) %	709
$\bar{K}^*(892)^0 \pi^+ \pi^-$ total	(2.4 ± 0.6) %	683
$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body	(1.52 ± 0.33) %	683
$K^- \pi^+ \rho^0$ total	(6.8 ± 0.5) %	612
$K^- \pi^+ \rho^0$ 3-body	(5.1 ± 2.3) × 10 ⁻³	612
$\bar{K}^*(892)^0 \rho^0$	(1.6 ± 0.4) %	418
$\bar{K}^*(892)^0 \rho^0$ transverse	(1.6 ± 0.5) %	418
$\bar{K}^*(892)^0 \rho^0$ S-wave	(3.0 ± 0.6) %	418
$\bar{K}^*(892)^0 \rho^0$ S-wave long.	< 3 × 10 ⁻³	418
$\bar{K}^*(892)^0 \rho^0$ P-wave	< 3 × 10 ⁻³	418
$\bar{K}^*(892)^0 \rho^0$ D-wave	(2.1 ± 0.6) %	418
$K^*(892)^- \rho^+$	(5.9 ± 2.4) %	422
$K^*(892)^- \rho^+$ longitudinal	(2.8 ± 1.2) %	422
$K^*(892)^- \rho^+$ transverse	(3.1 ± 1.8) %	422
$K^*(892)^- \rho^+$ P-wave	< 1.5 %	422

$K^- \pi^+ f_0(980)$	< 1.1 %	CL=90%	459
$\bar{K}^*(892)^0 f_0(980)$	< 7 × 10 ⁻³	CL=90%	-
$K_1(1270)^- \pi^+$	[<i>kk</i>] (1.04 ± 0.31) %	483	
$K_1(1400)^- \pi^+$	< 1.2 %	CL=90%	386
$\bar{K}_1(1400)^0 \pi^0$	< 3.7 %	CL=90%	378
$K^*(1410)^- \pi^+$	< 1.2 %	CL=90%	378
$K_0^*(1430)^- \pi^+$	(1.1 ± 0.4) %	364	
$K_2^*(1430)^- \pi^+$	< 8 × 10 ⁻³	CL=90%	367
$\bar{K}_2^*(1430)^0 \pi^0$	< 4 × 10 ⁻³	CL=90%	363
$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	(1.9 ± 0.9) %	641	
$\bar{K}^*(892)^0 \eta$	(1.9 ± 0.5) %	580	
$K^- \pi^+ \omega$	(3.1 ± 0.6) %	605	
$\bar{K}^*(892)^0 \omega$	(1.1 ± 0.5) %	406	
$K^- \pi^+ \eta'(958)$	(7.5 ± 2.0) × 10 ⁻³	479	
$\bar{K}^*(892)^0 \eta'(958)$	< 1.1 × 10 ⁻³	CL=90%	100
Plonic modes			
$\pi^+ \pi^-$	(1.59 ± 0.12) × 10 ⁻³	922	
$\pi^0 \pi^0$	(8.8 ± 2.3) × 10 ⁻⁴	922	
$\pi^+ \pi^- \pi^0$	(1.6 ± 1.1) %	S=2.7	907
$\pi^+ \pi^+ \pi^- \pi^-$	(8.3 ± 0.9) × 10 ⁻³	880	
$\pi^+ \pi^+ \pi^+ \pi^- \pi^0$	(1.9 ± 0.4) %	844	
$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$	(4.0 ± 3.0) × 10 ⁻⁴	795	
Hadronic modes with two K's			
$K^+ K^-$	(4.54 ± 0.29) × 10 ⁻³	791	
$K^0 \bar{K}^0$	(1.1 ± 0.4) × 10 ⁻³	788	
$K^0 K^- \pi^+$	(6.3 ± 1.1) × 10 ⁻³	S=1.2	739
$\bar{K}^*(892)^0 K^0$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	< 1.0 × 10 ⁻³	CL=90%	605
$K^*(892)^+ K^-$ × B($K^{*+} \rightarrow K^0 \pi^+$)	(2.3 ± 0.5) × 10 ⁻³	610	
$K^0 K^- \pi^+$ nonresonant	(2.4 ± 2.4) × 10 ⁻³	739	
$\bar{K}^0 K^+ \pi^-$	(4.9 ± 1.0) × 10 ⁻³	739	
$K^*(892)^0 \bar{K}^0$ × B($K^{*0} \rightarrow K^+ \pi^-$)	< 5 × 10 ⁻⁴	CL=90%	605
$K^*(892)^- K^+$ × B($K^{*-} \rightarrow \bar{K}^0 \pi^-$)	(1.2 ± 0.7) × 10 ⁻³	610	
$\bar{K}^0 K^+ \pi^-$ nonresonant	(4.0 ± 2.4) × 10 ⁻³	739	
$K^+ K^- \pi^+ \pi^-$	(2.4 ± 0.5) × 10 ⁻³	677	
$\phi \pi^+ \pi^- \times B(\phi \rightarrow K^+ K^-)$	(1.3 ± 0.4) × 10 ⁻³	614	
$\phi \rho^0 \times B(\phi \rightarrow K^+ K^-)$	(1.0 ± 0.25) × 10 ⁻³	260	
$K^*(892)^0 K^- \pi^+ + c.c. \times$ B($K^{*0} \rightarrow K^+ \pi^-$)	(5 ± 9) × 10 ⁻⁴	528	
$K^*(892)^0 \bar{K}^*(892)^0$ × B ² ($K^{*0} \rightarrow K^+ \pi^-$)	(1.3 ± 0.7) × 10 ⁻³	257	
$K^+ K^- \pi^+ \pi^-$ non- ϕ	(1.7 ± 0.5) × 10 ⁻³	677	
$K^+ K^- \pi^+ \pi^-$ nonresonant	(8 ± 90) × 10 ⁻⁵	677	
$K^+ K^- \pi^+ \pi^- \pi^0$	(3.1 ± 2.0) × 10 ⁻³	600	
Fractions of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.			
$\bar{K}^*(892)^0 K^0$	< 1.5 × 10 ⁻³	CL=90%	605
$K^*(892)^+ K^-$	(3.4 ± 0.8) × 10 ⁻³	610	
$K^*(892)^0 \bar{K}^0$	< 8 × 10 ⁻⁴	CL=90%	605
$K^*(892)^- K^+$	(1.8 ± 1.0) × 10 ⁻³	610	
$\phi \pi^+ \pi^-$	(2.6 ± 0.7) × 10 ⁻³	614	
$\phi \rho^0$	(1.9 ± 0.5) × 10 ⁻³	260	
$K^*(892)^0 K^- \pi^+ + c.c.$	(8 ± 13) × 10 ⁻⁴	528	
$K^*(892)^0 \bar{K}^*(892)^0$	(2.9 ± 1.6) × 10 ⁻³	257	
Doubly Cabibbo suppressed (DC) modes, ΔC = 2 forbidden via mixing (C2M) modes, ΔC = 1 weak neutral current (C1) modes, or Lepton Family number (LF) violating modes			
$K^+ \pi^-$	DC (3.1 ± 1.4) × 10 ⁻⁴	861	
$K^+ \pi^-$ (via \bar{D}^0)	C2M < 1.5 × 10 ⁻⁴	CL=90%	861
$K^+ \pi^+ \pi^- \pi^-$	DC < 1.5 × 10 ⁻³	CL=90%	812
μ^- anything (via \bar{D}^0)	C2M < 6 × 10 ⁻⁴	CL=90%	-
$e^+ e^-$	C1 < 1.3 × 10 ⁻⁴	CL=90%	932
$\mu^+ \mu^-$	C1 < 1.1 × 10 ⁻⁵	CL=90%	926
$\bar{K}^0 e^+ e^-$	< 1.7 × 10 ⁻³	CL=90%	866
$\rho^0 e^+ e^-$	C1 < 4.5 × 10 ⁻⁴	CL=90%	773
$\rho^0 \mu^+ \mu^-$	C1 < 8.1 × 10 ⁻⁴	CL=90%	756
$\mu^\pm e^\mp$	LF [<i>q</i>] < 1.0 × 10 ⁻⁴	CL=90%	929

Meson Summary Table

$$D^*(2007)^0 \quad I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

Mass $m = 2006.7 \pm 0.5$ MeV

$m_{D^{*0}} - m_{D^0} = 142.12 \pm 0.07$ MeV

Full width $\Gamma < 2.1$ MeV, CL = 90%

$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.

$D^*(2007)^0$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$D^0\pi^0$	(63.6 ± 2.8) %	43
$D^0\gamma$	(36.4 ± 2.8) %	137

$$D^*(2010)^\pm \quad I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation.

Mass $m = 2010.0 \pm 0.5$ MeV

$m_{D^{*(2010)^+}} - m_{D^+} = 140.64 \pm 0.09$ MeV

$m_{D^{*(2010)^+}} - m_{D^0} = 145.42 \pm 0.05$ MeV

Full width $\Gamma < 0.131$ MeV, CL = 90%

$D^*(2010)^-$ modes are charge conjugates of the modes below.

$D^*(2010)^\pm$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$D^0\pi^+$	(68.1 ± 1.3) %	39
$D^+\pi^0$	(30.8 ± 0.8) %	38
$D^+\gamma$	(1.1 $^{+1.4}_{-0.7}$) %	136

$$D_1(2420)^0 \quad I(J^P) = \frac{1}{2}(1^+)$$

I, J, P need confirmation.

Mass $m = 2422.8 \pm 3.2$ MeV ($S = 1.6$)

Full width $\Gamma = 18^{+6}_{-4}$ MeV

$\bar{D}_1(2420)^0$ modes are charge conjugates of modes below.

$D_1(2420)^0$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$D^*(2010)^+\pi^-$	seen	355
$D^+\pi^-$	not seen	474

$$D_2^*(2460) \quad I(J^P) = \frac{1}{2}(2^+)$$

$J^P = 2^+$ assignment strongly favored (ALBRECHT 89B).

Mass $m_{D_2^*(2460)^0} = 2457.7 \pm 1.9$ MeV

Mass $m_{D_2^*(2460)^\pm} = 2456 \pm 6$ MeV ($S = 2.0$)

$m_{D_2^*(2460)^\pm} - m_{D_2^*(2460)^0} = 2 \pm 5$ MeV ($S = 1.4$)

Full width $\Gamma_{D_2^*(2460)^0} = 21 \pm 5$ MeV

Full width $\Gamma_{D_2^*(2460)^\pm} = 23 \pm 10$ MeV

$\bar{D}_2^*(2460)$ modes are charge conjugates of modes below.

$D_2^*(2460)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$D_2^*(2460)^0 \rightarrow D^+\pi^-$	seen	503
$D_2^*(2460)^0 \rightarrow D^*(2010)^+\pi^-$	seen	387
$D_2^*(2460)^\pm \rightarrow D^0\pi^\pm$	seen	505

CHARMED, STRANGE MESONS ($C = S = \pm 1$)

$$D_s^+ = c\bar{s}, D_s^- = \bar{c}s, \quad \text{similarly for } D_s^{* \pm}$$

$$D_s^\pm$$

was F^\pm

$$I(J^P) = 0(0^-)$$

Mass $m = 1968.5 \pm 0.7$ MeV ($S = 1.2$)

$m_{D_s^\pm} - m_{D^\pm} = 99.1 \pm 0.6$ MeV ($S = 1.1$)

Mean life $\tau = (0.467 \pm 0.017) \times 10^{-12}$ s

$c\tau = 140$ μm

Branching fractions for modes below with a resonance in the final state include all the decay modes of the resonance. D_s^- modes are charge conjugates of the modes below.

Nearly all other modes are measured relative to the $\phi\pi^+$ mode. However, none of the determinations of the $\phi\pi^+$ branching fraction are direct measurements: all rely on calculated relations between D^+ and D_s^+ decay widths, on estimates of D_s^+ cross sections, or on other model-dependent assumptions. Thus a better determination of the $\phi\pi^+$ branching fraction could cause the other branching fractions to slide up or down, all together.

D_s^\pm DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Inclusive modes			
K^- anything	(13 $^{+14}_{-12}$) %		–
\bar{K}^0 anything + K^0 anything	(39 ± 28) %		–
K^+ anything	(20 $^{+18}_{-14}$) %		–
non- $K\bar{K}$ anything	(64 ± 17) %		–
e^+ anything	< 20 %	CL=90%	–
Leptonic and semileptonic modes			
$\mu^+\nu_\mu$	(5.9 ± 2.2) $\times 10^{-3}$	$S=1.1$	981
$\phi\ell^+\nu_\ell$	[pp] (1.88 ± 0.29) %		–
$\eta\mu^+\nu_\mu + \eta'(958)\mu^+\nu_\mu$	(7.4 ± 3.2) %		–
$\eta\mu^+\nu_\mu$			905
$\eta'(958)\mu^+\nu_\mu$	< 3.0 %	CL=90%	747
Hadronic modes with two K's (including from ϕ's)			
$K^+\bar{K}^0$	(3.5 ± 0.7) %		850
$K^+K^-\pi^+$	[aq] (4.8 ± 0.7) %		805
$\phi\pi^+$	(3.5 ± 0.4) %		712
$K^+\bar{K}^*(892)^0$	(3.3 ± 0.5) %		682
$K^+K^-\pi^+$ nonresonant	(8.7 ± 3.2) $\times 10^{-3}$		805
$K^0\bar{K}^0\pi^+$			802
$K^*(892)^+\bar{K}^0$	(4.2 ± 1.0) %		683
$K^+K^-\pi^+\pi^0$			748
$\phi\pi^+\pi^0$	(8 ± 4) %		687
$\phi\rho^+$	(6.5 $^{+1.6}_{-1.8}$) %		407
$\phi\pi^+\pi^0$ 3-body	< 2.5 %	CL=90%	687
$K^+K^-\pi^+\pi^0$ non- ϕ	< 8 %	CL=90%	748
$K^+\bar{K}^0\pi^+\pi^-$	< 2.7 %	CL=90%	744
$K^0K^-\pi^+\pi^+$	(4.2 ± 1.1) %		744
$K^*(892)^+\bar{K}^*(892)^0$	(5.6 ± 2.1) %		412
$K^0K^-\pi^+\pi^+$ non- $K^*\bar{K}^*$	< 2.8 %	CL=90%	744
$K^+K^-\pi^+\pi^+$			673
$\phi\pi^+\pi^+\pi^-$	(1.8 ± 0.5) %		640
$K^+K^-\pi^+\pi^+\pi^-$ non- ϕ	(3.0 $^{+3.0}_{-2.0}$) $\times 10^{-3}$		673

Meson Summary Table

Other hadronic modes		
$\pi^+ \pi^+ \pi^-$	$(1.35 \pm 0.31) \%$	959
$\rho^0 \pi^+$	$< 2.8 \times 10^{-3}$	CL=90% 827
$f_0(980) \pi^+$	$(10 \pm 4) \times 10^{-3}$	732
$\pi^+ \pi^+ \pi^-$ nonresonant	$(1.01 \pm 0.35) \%$	959
$\pi^+ \pi^+ \pi^- \pi^0$	$< 12 \%$	CL=90% 935
$\eta \pi^+$	$(1.9 \pm 0.4) \%$	902
$\omega \pi^+$	$< 1.7 \%$	CL=90% 822
$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	$(3.0 \pm 4.0) \times 10^{-3}$	899
$\pi^+ \pi^+ \pi^- \pi^0 \pi^0$		902
$\eta \rho^+$	$(10.0 \pm 2.2) \%$	727
$\eta \pi^+ \pi^0$ 3-body	$< 2.9 \%$	CL=90% 787
$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	$(4.9 \pm 3.2) \%$	856
$\eta'(958) \pi^+$	$(4.7 \pm 1.4) \%$	743
$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0 \pi^0$		803
$\eta'(958) \rho^+$	$(12.0 \pm 3.0) \%$	470
$\eta'(958) \pi^+ \pi^0$ 3-body	$< 3.0 \%$	CL=90% 720
$K^0 \pi^+$	$< 7 \times 10^{-3}$	CL=90% 916
$K^+ \pi^+ \pi^-$	$(3.0 \pm 4.0) \times 10^{-3}$	900
$K^+ K^- K^+$		628
ϕK^+	$< 2.5 \times 10^{-3}$	CL=90% 607

 $D_s^{*\pm}$

$$I(J^P) = ?(??)$$

Mass $m = 2110.0 \pm 1.9$ MeV ($S = 1.2$)
 $m_{D_s^{*\pm}} - m_{D_s^\pm} = 141.6 \pm 1.8$ MeV ($S = 1.2$)
 Full width $\Gamma < 4.5$ MeV, CL = 90%

D_s^{*-} modes are charge conjugates of the modes below.

$D_s^{*\pm}$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$D_s^{*+} \gamma$	dominant	137

 $D_{s1}(2536)^\pm$

$$I(J^P) = 0(1^+)$$

I, J, P need confirmation.

Mass $m = 2535.35 \pm 0.34$ MeV
 Full width $\Gamma < 2.3$ MeV, CL = 90%

$D_{s1}(2536)^-$ modes are charge conjugates of the modes below.

$D_{s1}(2536)^+$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$D^*(2010)^+ K^0$	seen	150
$D^*(2007)^0 K^+$	seen	169
$D^+ K^0$	not seen	382
$D^0 K^+$	not seen	392
$D_s^{*+} \gamma$	possibly seen	389

BOTTOM MESONS ($B = \pm 1$)

$$B^+ = \bar{u}b, B^0 = d\bar{b}, \bar{B}^0 = \bar{d}b, B^- = \bar{b}u, \text{ similarly for } B^{*+}$$

 B^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions. Measurements which do not identify the charge state of B also appear here.

Mass $m_{B^\pm} = 5278.7 \pm 2.0$ MeV

Mean life $\tau = (1.54 \pm 0.11) \times 10^{-12}$ s

Mean life τ (avg over B hadrons) = $(1.537 \pm 0.021) \times 10^{-12}$ s [a]

$c\tau = 388 \mu\text{m}$

B^- modes are charge conjugates of the modes below.

Only data from $\Upsilon(4S)$ decays are used for branching fractions, with rare exceptions. The branching fractions listed below assume a 50:50 $B^0 \bar{B}^0 : B^+ B^-$ production ratio at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D, D_s, D^* , and ψ branching ratios to current values whenever this would effect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

B^\pm DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Semileptonic modes			
$B^+ \rightarrow \bar{D}^0 \ell^+ \nu$	[b] $(1.6 \pm 0.7) \%$		—
$B^+ \rightarrow \bar{D}^*(2007)^0 \ell^+ \nu$	[b] $(6.6 \pm 2.2) \%$		—
$B^+ \rightarrow \pi^0 e^+ \nu_e$	$< 2.2 \times 10^{-3}$	CL=90%	2638
$B^+ \rightarrow \omega \ell^+ \nu_\ell$	[b] $< 2.1 \times 10^{-4}$	CL=90%	—
$B^+ \rightarrow \omega \mu^+ \nu_\mu$	seen		2580
$B^+ \rightarrow \rho^0 \ell^+ \nu_\ell$	[b] $< 2.1 \times 10^{-4}$	CL=90%	—
D, D^*, or D_s modes			
$B^+ \rightarrow \bar{D}^0 \pi^+$	$(5.3 \pm 0.5) \times 10^{-3}$		2308
$B^+ \rightarrow \bar{D}^0 \rho^+$	$(1.34 \pm 0.18) \%$		2237
$B^+ \rightarrow \bar{D}^0 \pi^+ \pi^+ \pi^-$	$(1.1 \pm 0.4) \%$		2289
$B^+ \rightarrow \bar{D}^0 \pi^+ \pi^+ \pi^-$ nonresonant	$(5 \pm 4) \times 10^{-3}$		2289
$B^+ \rightarrow \bar{D}^0 \pi^+ \rho^0$	$(4.2 \pm 3.0) \times 10^{-3}$		2208
$B^+ \rightarrow \bar{D}^0 a_1(1260)^+$	$(5 \pm 4) \times 10^{-3}$		2123
$B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+$	$(2.1 \pm 0.6) \times 10^{-3}$		2247
$B^+ \rightarrow D^- \pi^+ \pi^+$	$< 1.4 \times 10^{-3}$	CL=90%	2299
$B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+$	$(5.2 \pm 0.8) \times 10^{-3}$		2255
$B^+ \rightarrow \bar{D}^*(2007)^0 \rho^+$	$(1.55 \pm 0.31) \%$		2182
$B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-$	$(9.4 \pm 2.6) \times 10^{-3}$		2236
$B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^0$	$(1.5 \pm 0.7) \%$		2235
$B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^-$	$< 1 \%$	CL=90%	2217
$B^+ \rightarrow \bar{D}_1^*(2420)^0 \pi^+$	$(1.1 \pm 0.5) \times 10^{-3}$		2081
$B^+ \rightarrow \bar{D}_1^*(2420)^0 \rho^+$	$< 1.4 \times 10^{-3}$	CL=90%	1996
$B^+ \rightarrow \bar{D}_2^*(2460)^0 \pi^+$	$< 1.3 \times 10^{-3}$	CL=90%	2064
$B^+ \rightarrow \bar{D}_2^*(2460)^0 \rho^+$	$< 4.7 \times 10^{-3}$	CL=90%	1979
$B^+ \rightarrow \bar{D}^0 D_s^+$	$(1.7 \pm 0.6) \%$		1814
$B^+ \rightarrow \bar{D}^0 D_s^{*+}$	$(1.2 \pm 1.0) \%$		1735
$B^+ \rightarrow \bar{D}^*(2007)^0 D_s^+$	$(1.0 \pm 0.7) \%$		1737
$B^+ \rightarrow \bar{D}^*(2007)^0 D_s^{*+}$	$(2.4 \pm 1.3) \%$		1652
$B^+ \rightarrow D_s^+ \pi^0$	$< 2.1 \times 10^{-4}$	CL=90%	2270
$B^+ \rightarrow D_s^{*+} \pi^0$	$< 3.4 \times 10^{-4}$	CL=90%	2215
$B^+ \rightarrow D_s^+ \eta$	$< 5 \times 10^{-4}$	CL=90%	2235
$B^+ \rightarrow D_s^{*+} \eta$	$< 8 \times 10^{-4}$	CL=90%	2178
$B^+ \rightarrow D_s^+ \rho^0$	$< 4 \times 10^{-4}$	CL=90%	2197
$B^+ \rightarrow D_s^{*+} \rho^0$	$< 5 \times 10^{-4}$	CL=90%	2139
$B^+ \rightarrow D_s^+ \omega$	$< 5 \times 10^{-4}$	CL=90%	2195
$B^+ \rightarrow D_s^{*+} \omega$	$< 7 \times 10^{-4}$	CL=90%	2137
$B^+ \rightarrow D_s^+ a_1(1260)^0$	$< 2.3 \times 10^{-3}$	CL=90%	2079

Meson Summary Table

$B^+ \rightarrow D_s^{*+} a_1(1260)^0$	< 1.7	$\times 10^{-3}$	CL=90%	2015
$B^+ \rightarrow D_s^+ \phi$	< 3.3	$\times 10^{-4}$	CL=90%	2140
$B^+ \rightarrow D_s^{*+} \phi$	< 4	$\times 10^{-4}$	CL=90%	2080
$B^+ \rightarrow D_s^+ \bar{K}^0$	< 1.1	$\times 10^{-3}$	CL=90%	2241
$B^+ \rightarrow D_s^{*+} \bar{K}^0$	< 1.2	$\times 10^{-3}$	CL=90%	2185
$B^+ \rightarrow D_s^+ \bar{K}^*(892)^0$	< 5	$\times 10^{-4}$	CL=90%	2171
$B^+ \rightarrow D_s^{*+} \bar{K}^*(892)^0$	< 5	$\times 10^{-4}$	CL=90%	2111
$B^+ \rightarrow D_s^- \pi^+ K^+$	< 9	$\times 10^{-4}$	CL=90%	2222
$B^+ \rightarrow D_s^{*-} \pi^+ K^+$	< 1.2	$\times 10^{-3}$	CL=90%	2165
$B^+ \rightarrow D_s^- \pi^+ K^*(892)^+$	< 7	$\times 10^{-3}$	CL=90%	2137
$B^+ \rightarrow D_s^{*-} \pi^+ K^*(892)^+$	< 9	$\times 10^{-3}$	CL=90%	2076

Charmonium modes

$B^+ \rightarrow J/\psi(1S) K^+$	$(1.02 \pm 0.14) \times 10^{-3}$			1683
$B^+ \rightarrow J/\psi(1S) K^+ \pi^+ \pi^-$	$(1.4 \pm 0.6) \times 10^{-3}$			1612
$B^+ \rightarrow J/\psi(1S) K^*(892)^+$	$(1.7 \pm 0.5) \times 10^{-3}$			1571
$B^+ \rightarrow \psi(2S) K^+$	$(6.9 \pm 3.1) \times 10^{-4}$	S=1.3		1284
$B^+ \rightarrow \psi(2S) K^*(892)^+$	< 3.0	$\times 10^{-3}$	CL=90%	1115
$B^+ \rightarrow \psi(2S) K^*(892)^+ \pi^+ \pi^-$	$(1.9 \pm 1.2) \times 10^{-3}$			909
$B^+ \rightarrow \chi_{c1}(1P) K^+$	$(1.0 \pm 0.4) \times 10^{-3}$			1411
$B^+ \rightarrow \chi_{c1}(1P) K^*(892)^+$	< 2.1	$\times 10^{-3}$	CL=90%	1265

K or K* modes

$B^+ \rightarrow K^0 \pi^+$	< 1.0	$\times 10^{-4}$	CL=90%	2614
$B^+ \rightarrow K^*(892)^0 \pi^+$	< 1.5	$\times 10^{-4}$	CL=90%	2561
$B^+ \rightarrow K^+ \pi^- \pi^+$ (no charm)	< 1.9	$\times 10^{-4}$	CL=90%	2609
$B^+ \rightarrow K_1(1400)^0 \pi^+$	< 2.6	$\times 10^{-3}$	CL=90%	2451
$B^+ \rightarrow K_2^*(1430)^0 \pi^+$	< 6.8	$\times 10^{-4}$	CL=90%	2443
$B^+ \rightarrow K^+ \rho^0$	< 8	$\times 10^{-5}$	CL=90%	2559
$B^+ \rightarrow K^*(892)^+ \pi^+ \pi^-$	< 1.1	$\times 10^{-3}$	CL=90%	2556
$B^+ \rightarrow K^*(892)^+ \rho^0$	< 9.0	$\times 10^{-4}$	CL=90%	2505
$B^+ \rightarrow K_1(1400)^+ \rho^0$	< 7.8	$\times 10^{-4}$	CL=90%	2388
$B^+ \rightarrow K_2^*(1430)^+ \rho^0$	< 1.5	$\times 10^{-3}$	CL=90%	2382
$B^+ \rightarrow K^+ K^- K^+$	< 3.5	$\times 10^{-4}$	CL=90%	2522
$B^+ \rightarrow K^+ \phi$	< 9	$\times 10^{-5}$	CL=90%	2516
$B^+ \rightarrow K^*(892)^+ K^+ K^-$	< 1.6	$\times 10^{-3}$	CL=90%	2466
$B^+ \rightarrow K^*(892)^+ \phi$	< 1.3	$\times 10^{-3}$	CL=90%	2460
$B^+ \rightarrow K_1(1400)^+ \phi$	< 1.1	$\times 10^{-3}$	CL=90%	2339
$B^+ \rightarrow K_2^*(1430)^+ \phi$	< 3.4	$\times 10^{-3}$	CL=90%	2332
$B^+ \rightarrow K^+ f_0(980)$	< 8	$\times 10^{-5}$	CL=90%	2524
$B^+ \rightarrow K^*(892)^+ \gamma$	$(5.7 \pm 3.3) \times 10^{-5}$			2564
$B^+ \rightarrow K_1(1270)^+ \gamma$	< 7.3	$\times 10^{-3}$	CL=90%	2486
$B^+ \rightarrow K_1(1400)^+ \gamma$	< 2.2	$\times 10^{-3}$	CL=90%	2453
$B^+ \rightarrow K_2^*(1430)^+ \gamma$	< 1.4	$\times 10^{-3}$	CL=90%	2447
$B^+ \rightarrow K^*(1680)^+ \gamma$	< 1.9	$\times 10^{-3}$	CL=90%	2361
$B^+ \rightarrow K_3^*(1780)^+ \gamma$	< 5.5	$\times 10^{-3}$	CL=90%	2343
$B^+ \rightarrow K_4^*(2045)^+ \gamma$	< 9.9	$\times 10^{-3}$	CL=90%	2243

Light unflavored meson modes

$B^+ \rightarrow \pi^+ \pi^0$	< 2.4	$\times 10^{-4}$	CL=90%	2636
$B^+ \rightarrow \pi^+ \pi^+ \pi^-$	< 1.9	$\times 10^{-4}$	CL=90%	2630
$B^+ \rightarrow \rho^0 \pi^+$	< 1.5	$\times 10^{-4}$	CL=90%	2581
$B^+ \rightarrow \pi^+ f_0(980)$	< 1.4	$\times 10^{-4}$	CL=90%	2546
$B^+ \rightarrow \pi^+ f_2(1270)$	< 2.4	$\times 10^{-4}$	CL=90%	2483
$B^+ \rightarrow \pi^+ \pi^0 \pi^0$	< 8.9	$\times 10^{-4}$	CL=90%	2631
$B^+ \rightarrow \rho^+ \pi^0$	< 5.5	$\times 10^{-4}$	CL=90%	2581
$B^+ \rightarrow \pi^+ \pi^- \pi^+ \pi^0$	< 4.0	$\times 10^{-3}$	CL=90%	2621
$B^+ \rightarrow \rho^+ \rho^0$	< 1.0	$\times 10^{-3}$	CL=90%	2525
$B^+ \rightarrow a_1(1260)^+ \pi^0$	< 1.7	$\times 10^{-3}$	CL=90%	2494
$B^+ \rightarrow a_1(1260)^0 \pi^+$	< 9.0	$\times 10^{-4}$	CL=90%	2494
$B^+ \rightarrow \omega \pi^+$	< 4.0	$\times 10^{-4}$	CL=90%	2580
$B^+ \rightarrow \eta \pi^+$	< 7.0	$\times 10^{-4}$	CL=90%	2609
$B^+ \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	< 8.6	$\times 10^{-4}$	CL=90%	2608
$B^+ \rightarrow \rho^0 a_1(1260)^+$	< 6.2	$\times 10^{-4}$	CL=90%	2433
$B^+ \rightarrow \rho^0 a_2(1320)^+$	< 7.2	$\times 10^{-4}$	CL=90%	2411
$B^+ \rightarrow \pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 6.3	$\times 10^{-3}$	CL=90%	2592
$B^+ \rightarrow a_1(1260)^+ a_1(1260)^0$	< 1.3	%	CL=90%	2335

Baryon modes

$B^+ \rightarrow p \bar{p} \pi^+$	< 1.6	$\times 10^{-4}$	CL=90%	2438
$B^+ \rightarrow p \bar{p} \pi^+ \pi^+ \pi^-$	< 5.2	$\times 10^{-4}$	CL=90%	2369
$B^+ \rightarrow p \bar{\Lambda}$	< 6	$\times 10^{-5}$	CL=90%	2430
$B^+ \rightarrow p \bar{\Lambda} \pi^+ \pi^-$	< 2.0	$\times 10^{-4}$	CL=90%	2367
$B^+ \rightarrow \Delta^0 p$	< 3.8	$\times 10^{-4}$	CL=90%	2402
$B^+ \rightarrow \Delta^{++} \bar{p}$	< 1.5	$\times 10^{-4}$	CL=90%	2402

Lepton Family number (LF) or Lepton number (L) violating modes, or $\Delta B = 1$ weak neutral current (BI) modes

$B^+ \rightarrow \pi^+ e^+ e^-$	B1	< 3.9	$\times 10^{-3}$	CL=90%	2638
$B^+ \rightarrow \pi^+ \mu^+ \mu^-$	B1	< 9.1	$\times 10^{-3}$	CL=90%	2633
$B^+ \rightarrow K^+ e^+ e^-$	B1	< 6	$\times 10^{-5}$	CL=90%	2616
$B^+ \rightarrow K^+ \mu^+ \mu^-$	B1	< 1.7	$\times 10^{-4}$	CL=90%	2612
$B^+ \rightarrow K^*(892)^+ e^+ e^-$	B1	< 6.9	$\times 10^{-4}$	CL=90%	2564
$B^+ \rightarrow K^*(892)^+ \mu^+ \mu^-$	B1	< 1.2	$\times 10^{-3}$	CL=90%	2560
$B^+ \rightarrow \pi^+ e^+ \mu^-$	LF	< 6.4	$\times 10^{-3}$	CL=90%	2636
$B^+ \rightarrow \pi^+ e^- \mu^+$	LF	< 6.4	$\times 10^{-3}$	CL=90%	2636
$B^+ \rightarrow K^+ e^+ \mu^-$	LF	< 6.4	$\times 10^{-3}$	CL=90%	2615
$B^+ \rightarrow K^+ e^- \mu^+$	LF	< 6.4	$\times 10^{-3}$	CL=90%	2615
$B^+ \rightarrow \pi^- e^+ e^+$	L	< 3.9	$\times 10^{-3}$	CL=90%	2638
$B^+ \rightarrow \pi^- \mu^+ \mu^+$	L	< 9.1	$\times 10^{-3}$	CL=90%	2633
$B^+ \rightarrow \pi^- e^+ \mu^+$	L	< 6.4	$\times 10^{-3}$	CL=90%	2636
$B^+ \rightarrow K^- e^+ e^+$	L	< 3.9	$\times 10^{-3}$	CL=90%	2616
$B^+ \rightarrow K^- \mu^+ \mu^+$	L	< 9.1	$\times 10^{-3}$	CL=90%	2612
$B^+ \rightarrow K^- e^+ \mu^+$	L	< 6.4	$\times 10^{-3}$	CL=90%	2615

B DECAY MODES

\bar{B} modes are charge conjugates of the modes below.

For the following modes, the charge of B was not determined. The measurements are for an admixture of B mesons at the $\Upsilon(4S)$ unless otherwise indicated by a footnote and a "B" instead of "B" in the initial state.

Semileptonic and leptonic modes

$B \rightarrow e^+ \nu_e$ anything	[c]	$(10.4 \pm 0.4) \%$	S=1.3	-
$B \rightarrow \bar{D}^*(2010) e^+ \nu_e$		$(7.0 \pm 2.3) \%$		-
$B \rightarrow \bar{p} e^+ \nu_e$ anything		< 1.6	$\times 10^{-3}$	CL=90%
$B \rightarrow \mu^+ \nu_\mu$ anything	[c]	$(10.3 \pm 0.5) \%$		-
$B \rightarrow \ell^+ \nu_\ell$ anything	[b,c]	$(10.43 \pm 0.24) \%$		-
$B \rightarrow D^- \ell^+ \nu_\ell$ anything	[b]	$(2.7 \pm 0.8) \%$		-
$B \rightarrow \bar{D}^0 \ell^+ \nu_\ell$ anything	[b]	$(7.0 \pm 1.4) \%$		-
$B \rightarrow D^{*-} \ell^+ \nu_\ell$	[b,d]	$(2.7 \pm 0.7) \%$		-
$B \rightarrow D_s^- \ell^+ \nu_\ell$ anything	[b]	< 9	$\times 10^{-3}$	CL=90%
$B \rightarrow D_s^- \ell^+ \nu_\ell K^+$ anything	[b]	< 6	$\times 10^{-3}$	CL=90%
$B \rightarrow D_s^- \ell^+ \nu_\ell K^0$ anything	[b]	< 9	$\times 10^{-3}$	CL=90%
$B \rightarrow K^+ \ell^+ \nu_\ell$ anything	[b]	$(5.6 \pm 1.0) \%$		-
$B \rightarrow K^- \ell^+ \nu_\ell$ anything	[b]	$(1.0 \pm 0.6) \%$		-
$B \rightarrow K^0 / \bar{K}^0 \ell^+ \nu_\ell$ anything	[b]	$(4.1 \pm 0.8) \%$		-
$\bar{b} \rightarrow \tau^+ \nu_\tau$ anything	[e]	$(4.1 \pm 1.0) \%$		-

D, D*, or D_s modes

$B \rightarrow D^-$ anything		$(26 \pm 4) \%$		-
$B \rightarrow \bar{D}^0$ anything		$(54 \pm 6) \%$		-
$B \rightarrow D^*(2010)^- anything$		$(23 \pm 4) \%$	S=1.4	-
$B \rightarrow D_s^+ anything$	[f]	$(8.9 \pm 1.1) \%$		-
$B \rightarrow D_s D, D_s^* D, D_s D^*, or D_s^* D^*$	[f]	$(5.0 \pm 0.9) \%$		-
$B \rightarrow D^*(2010) \gamma$		< 1.1	$\times 10^{-3}$	CL=90%
$B \rightarrow D_s^+ \pi^-, D_s^{*+} \pi^-, D_s^+ \rho^-, D_s^{*+} \rho^-, D_s^+ \pi^0, D_s^{*+} \pi^0, D_s^+ \eta, D_s^{*+} \eta, D_s^+ \rho^0, D_s^{*+} \rho^0, D_s^+ \omega, D_s^{*+} \omega$		< 5	$\times 10^{-4}$	CL=90%

Charmonium modes

$B \rightarrow J/\psi(1S) anything$		$(1.30 \pm 0.17) \%$		-
$B \rightarrow \psi(2S) anything$		$(4.6 \pm 2.0) \times 10^{-3}$		-
$B \rightarrow \chi_{c1}(1P) anything$		$(1.1 \pm 0.4) \%$		-

K or K* modes

$B \rightarrow K^\pm anything$	[f]	$(85 \pm 11) \%$		-
$B \rightarrow K^0 / \bar{K}^0 anything$		$(63 \pm 8) \%$		-
$b \rightarrow s \gamma$	[g]	< 1.2	$\times 10^{-3}$	CL=90%
$B \rightarrow K^*(892) \gamma$		< 2.4	$\times 10^{-4}$	CL=90%
$B \rightarrow K_1(1400) \gamma$		< 4.1	$\times 10^{-4}$	CL=90%
$B \rightarrow K_2^*(1430) \gamma$		< 8.3	$\times 10^{-4}$	CL=90%
$B \rightarrow K_2(1770) \gamma$		< 1.2	$\times 10^{-3}$	CL=90%
$B \rightarrow K_3^*(1780) \gamma$		< 3.0	$\times 10^{-3}$	CL=90%
$B \rightarrow K_4^*(2045) \gamma$		< 1.0	$\times 10^{-3}$	CL=90%

Light unflavored meson modes

$B \rightarrow \phi anything$		$(2.3 \pm 0.8) \%$		-
-------------------------------	--	--------------------	--	---

Meson Summary Table

Baryon modes				Charmonium modes			
$B \rightarrow$ charmed-baryon anything	(6.4 ± 1.1) %	-		$D^*(2010)^- \pi^+ \pi^- \pi^0$	(3.4 ± 1.8) %		2218
$B \rightarrow \bar{\Sigma}_c^-$ anything	(4.8 ± 2.5) × 10 ⁻³	-		$\bar{D}_s^*(2460)^- \pi^+$	< 2.2 × 10 ⁻³	90%	2065
$B \rightarrow \bar{\Sigma}_c^0$ anything	< 1.1 %	CL=90%	-	$\bar{D}_s^*(2460)^- \rho^+$	< 4.9 × 10 ⁻³	90%	1980
$B \rightarrow \bar{\Sigma}_c^{\prime 0}$ anything	(5.3 ± 2.5) × 10 ⁻³	-		$D^- D_s^+$	(8 ± 4) × 10 ⁻³		1812
$B \rightarrow \bar{\Sigma}_c^{\prime 0} N(N = p \text{ or } n)$	< 1.7 × 10 ⁻³	CL=90%	-	$D^*(2010)^- D_s^+$	(1.2 ± 0.6) %		1735
$B \rightarrow p$ anything + \bar{p} anything	(8.0 ± 0.5) %	-		$D^- D_s^{*+}$	(2.1 ± 1.5) %		1733
$B \rightarrow p(\text{direct})$ anything + $\bar{p}(\text{direct})$ anything	(5.6 ± 0.7) %	-		$D^*(2010)^- D_s^{*+}$	(2.0 ± 1.2) %		1650
$B \rightarrow \Lambda$ anything + $\bar{\Lambda}$ anything	(4.0 ± 0.5) %	-		$D_s^+ \pi^-$	< 2.9 × 10 ⁻⁴	90%	2270
$B \rightarrow \Xi^-$ anything + $\Xi^{\prime -}$ anything	(2.7 ± 0.6) × 10 ⁻³	-		$D_s^+ \pi^0$	< 5 × 10 ⁻⁴	90%	2215
$B \rightarrow$ baryons anything	(6.8 ± 0.6) %	-		$D_s^+ \rho^-$	< 7 × 10 ⁻⁴	90%	2198
$B \rightarrow p\bar{p}$ anything	(2.47 ± 0.23) %	-		$D_s^+ \rho^0$	< 8 × 10 ⁻⁴	90%	2140
$B \rightarrow \Lambda p\bar{p}$ anything + $\bar{\Lambda} p\bar{p}$ anything	(2.5 ± 0.4) %	-		$D_s^+ a_1(1260)^-$	< 2.7 × 10 ⁻³	90%	2079
$B \rightarrow \Lambda \bar{\Lambda}$ anything	< 5 × 10 ⁻³	CL=90%	-	$D_s^+ a_1(1260)^0$	< 2.2 × 10 ⁻³	90%	2015
$\Delta B = 1$ weak neutral current ($B1$) modes				$D_s^- K^+$	< 2.4 × 10 ⁻⁴	90%	2242
$\bar{b} \rightarrow e^+ e^-$ anything	$B1$ $ g < 2.4$	× 10 ⁻³	-	$D_s^- K^+$	< 1.8 × 10 ⁻⁴	90%	2186
$\bar{b} \rightarrow \mu^+ \mu^-$ anything	$B1$ $ g < 5.0$	× 10 ⁻⁵	CL=90%	$D_s^- K^*(892)^+$	< 1.0 × 10 ⁻³	90%	2172
B^0				$D_s^- K^*(892)^+$	< 1.2 × 10 ⁻³	90%	2113
$I(J^P) = \frac{1}{2}(0^-)$				$D_s^- \pi^+ K^0$	< 6 × 10 ⁻³	90%	2221
I, J, P need confirmation. Quantum numbers shown are quark-model predictions.				$D_s^- \pi^+ K^0$	< 3.2 × 10 ⁻³	90%	2164
Mass $m_{B^0} = 5279.0 \pm 2.0$ MeV				$D_s^- \pi^+ K^*(892)^0$	< 4 × 10 ⁻³	90%	2136
$m_{B^0} - m_{B^\pm} = 0.34 \pm 0.29$ MeV ($S = 1.1$)				$D_s^- \pi^+ K^*(892)^0$	< 2.1 × 10 ⁻³	90%	2075
Mean life $\tau = (1.50 \pm 0.11) \times 10^{-12}$ s				$\bar{D}^0 \pi^0$	< 4.8 × 10 ⁻⁴	90%	2308
$c\tau = 449$ μm				$\bar{D}^0 \rho^0$	< 5.5 × 10 ⁻⁴	90%	2238
$\tau_{B^+}/\tau_{B^0} = 0.98 \pm 0.09$				$\bar{D}^0 \eta$	< 6.8 × 10 ⁻⁴	90%	2274
$B^0 - \bar{B}^0$ mixing parameters				$\bar{D}^0 \eta'$	< 8.6 × 10 ⁻⁴	90%	2197
$\chi_d = 0.156 \pm 0.024$				$\bar{D}^0 \omega$	< 6.3 × 10 ⁻⁴	90%	2235
$\Delta m_{B^0} = m_{B_\mu^0} - m_{B_l^0} = (0.51 \pm 0.06) \times 10^{12} \hbar \text{ s}^{-1}$				$\bar{D}^*(2007)^0 \pi^0$	< 9.7 × 10 ⁻⁴	90%	2256
$\chi_d = \Delta m_{B^0}/\Gamma_{B^0} = 0.71 \pm 0.06$ [a]				$\bar{D}^*(2007)^0 \rho^0$	< 1.17 × 10 ⁻³	90%	2182
\bar{B}^0 modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing. Decays in which the charge of the B is not determined are in the B^\pm section.				$\bar{D}^*(2007)^0 \eta$	< 6.9 × 10 ⁻⁴	90%	2220
Only data from $\Upsilon(4S)$ decays are used for branching fractions, with rare exceptions. The branching fractions listed below assume a 50:50 $B^0 \bar{B}^0 : B^+ B^-$ production ratio at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D, D_s, D^* , and ψ branching ratios to current values whenever this would effect our averages and best limits significantly.				$\bar{D}^*(2007)^0 \eta'$	< 2.7 × 10 ⁻³	90%	2140
Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.				$\bar{D}^*(2007)^0 \omega$	< 2.1 × 10 ⁻³	90%	2180
B^0 DECAY MODES				$J/\psi(1S) K^0$	(7.5 ± 2.1) × 10 ⁻⁴		1682
Semileptonic and leptonic modes				$J/\psi(1S) K^+ \pi^-$	(1.2 ± 0.6) × 10 ⁻³		1652
$\ell^+ \nu_\ell$ anything	[b] (9.5 ± 1.6) %	-		$J/\psi(1S) K^*(892)^0$	(1.58 ± 0.28) × 10 ⁻³		1569
$D^- \ell^+ \nu_\ell$	[b] (1.9 ± 0.5) %	-		$\psi(2S) K^0$	< 8 × 10 ⁻⁴	90%	1283
$D^*(2010)^- \ell^+ \nu_\ell$	[b] (4.4 ± 0.4) %	-		$\psi(2S) K^+ \pi^-$	< 1 × 10 ⁻³	90%	1238
$\rho^- \ell^+ \nu_\ell$	[b] < 4.1 × 10 ⁻⁴	90%	-	$\psi(2S) K^*(892)^0$	(1.4 ± 0.9) × 10 ⁻³		1113
$\pi^- \mu^+ \nu_\mu$	seen		2636	$\chi_{c1}(1P) K^0$	< 2.7 × 10 ⁻³	90%	1410
D, D^*, or D_s modes				$\chi_{c1}(1P) K^*(892)^0$	< 2.1 × 10 ⁻³	90%	1263
$D^- \pi^+$	(3.0 ± 0.4) × 10 ⁻³		2306	K or K^* modes			
$D^- \rho^+$	(7.8 ± 1.4) × 10 ⁻³		2236	$K^+ \pi^-$	< 2.6 × 10 ⁻⁵	90%	2615
$\bar{D}^0 \pi^+ \pi^-$	< 1.6 × 10 ⁻³	90%	2301	$K^+ K^-$	< 7 × 10 ⁻⁶	90%	2593
$D^*(2010)^- \pi^+$	(2.6 ± 0.4) × 10 ⁻³		2254	$K^0 \pi^+ \pi^-$	< 4.4 × 10 ⁻⁴	90%	2609
$D^- \pi^+ \pi^+ \pi^-$	(8.0 ± 2.5) × 10 ⁻³		2287	$K^0 \rho^0$	< 3.2 × 10 ⁻⁴	90%	2559
($D^- \pi^+ \pi^+ \pi^-$) nonresonant	(3.9 ± 1.9) × 10 ⁻³		2287	$K^0 f_0(980)$	< 3.6 × 10 ⁻⁴	90%	2523
$D^- \pi^+ \rho^0$	(1.1 ± 1.0) × 10 ⁻³		2207	$K^*(892)^+ \pi^-$	< 3.8 × 10 ⁻⁴	90%	2562
$D^- a_1(1260)^+$	(6.0 ± 3.3) × 10 ⁻³		2121	$K_2^*(1430)^+ \pi^-$	< 2.6 × 10 ⁻³	90%	2445
$D^*(2010)^- \pi^+ \pi^0$	(1.5 ± 0.5) %		2247	$K^0 K^+ K^-$	< 1.3 × 10 ⁻³	90%	2522
$D^*(2010)^- \rho^+$	(7.3 ± 1.5) × 10 ⁻³		2181	$K^0 \phi$	< 4.2 × 10 ⁻⁴	90%	2516
$D^*(2010)^- \pi^+ \pi^+ \pi^-$	(1.19 ± 0.27) %		2235	$K^*(892)^0 \pi^+ \pi^-$	< 1.4 × 10 ⁻³	90%	2556
($D^*(2010)^- \pi^+ \pi^+ \pi^-$) non-resonant	(0.0 ± 2.5) × 10 ⁻³		2235	$K^*(892)^0 \rho^0$	< 4.6 × 10 ⁻⁴	90%	2504
$D^*(2010)^- \pi^+ \rho^0$	(5.7 ± 3.1) × 10 ⁻³		2151	$K^*(892)^0 f_0(980)$	< 1.7 × 10 ⁻⁴	90%	2467
$D^*(2010)^- a_1(1260)^+$	(1.5 ± 0.7) %		2061	$K_1(1400)^+ \pi^-$	< 1.1 × 10 ⁻³	90%	2451
				$K^*(892)^0 K^+ K^-$	< 6.1 × 10 ⁻⁴	90%	2465
				$K^*(892)^0 \phi$	< 3.2 × 10 ⁻⁴	90%	2459
				$K_1(1400)^0 \rho^0$	< 3.0 × 10 ⁻³	90%	2388
				$K_1(1400)^0 \phi$	< 5.0 × 10 ⁻³	90%	2339
				$K_2^*(1430)^0 \rho^0$	< 1.1 × 10 ⁻³	90%	2380
				$K_2^*(1430)^0 \phi$	< 1.4 × 10 ⁻³	90%	2330
				$K^*(892)^0 \gamma$	(4.0 ± 1.9) × 10 ⁻⁵		2563
				$K_1(1270)^0 \gamma$	< 7.0 × 10 ⁻³	90%	2486
				$K_1(1400)^0 \gamma$	< 4.3 × 10 ⁻³	90%	2453
				$K_2^*(1430)^0 \gamma$	< 4.0 × 10 ⁻⁴	90%	2445
				$K^*(1680)^0 \gamma$	< 2.0 × 10 ⁻³	90%	2361
				$K_2^*(1780)^0 \gamma$	< 1.0 %	90%	2343
				$K_4^*(2045)^0 \gamma$	< 4.3 × 10 ⁻³	90%	2243

Meson Summary Table

Light unflavored meson modes			
$\pi^+\pi^-$	< 2.9	$\times 10^{-5}$	90%
$\pi^+\pi^-\pi^0$	< 7.2	$\times 10^{-4}$	90%
$\rho^0\pi^0$	< 4.0	$\times 10^{-4}$	90%
$\rho^\mp\pi^\pm$	[c] < 5.2	$\times 10^{-4}$	90%
$\pi^+\pi^-\pi^+\pi^-$	< 6.7	$\times 10^{-4}$	90%
$\rho^0\rho^0$	< 2.8	$\times 10^{-4}$	90%
$a_1(1260)^\mp\pi^\pm$	[c] < 4.9	$\times 10^{-4}$	90%
$a_2(1320)^\mp\pi^\pm$	[c] < 3.0	$\times 10^{-4}$	90%
$\pi^+\pi^-\pi^0\pi^0$	< 3.1	$\times 10^{-3}$	90%
$\rho^+\rho^-$	< 2.2	$\times 10^{-3}$	90%
$a_1(1260)^0\pi^0$	< 1.1	$\times 10^{-3}$	90%
$\omega\pi^0$	< 4.6	$\times 10^{-4}$	90%
$\eta\pi^0$	< 1.8	$\times 10^{-3}$	90%
$\pi^+\pi^+\pi^-\pi^-\pi^0$	< 9.0	$\times 10^{-3}$	90%
$a_1(1260)^+\rho^-$	< 3.4	$\times 10^{-3}$	90%
$a_1(1260)^0\rho^0$	< 2.4	$\times 10^{-3}$	90%
$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^-$	< 3.0	$\times 10^{-3}$	90%
$a_1(1260)^+a_1(1260)^-$	< 2.8	$\times 10^{-3}$	90%
$\pi^+\pi^+\pi^+\pi^-\pi^-\pi^-\pi^0$	< 1.1	%	90%

Baryon modes			
$\rho\bar{\rho}$	< 3.4	$\times 10^{-5}$	90%
$\rho\bar{\rho}\pi^+\pi^-$	< 2.5	$\times 10^{-4}$	90%
$\rho\bar{\Lambda}\pi^-$	< 1.8	$\times 10^{-4}$	90%
$\Delta^0\bar{\Delta}^0$	< 1.5	$\times 10^{-3}$	90%
$\Delta^{++}\Delta^{--}$	< 1.1	$\times 10^{-4}$	90%
$\Sigma^- \bar{\Delta}^{++}$	< 1.2	$\times 10^{-3}$	90%

Lepton Family number (LF) violating modes, $\Delta B = 2$ forbidden decay via mixing (B2M) modes, or $\Delta B = 1$ weak neutral current (B1) modes			
e^+e^-	B1	< 5.9	$\times 10^{-6}$
$\mu^+\mu^-$	B1	< 5.9	$\times 10^{-6}$
$K^0e^+e^-$	B1	< 3.0	$\times 10^{-4}$
$K^0\mu^+\mu^-$	B1	< 3.6	$\times 10^{-4}$
$K^*(892)^0e^+e^-$	B1	< 2.9	$\times 10^{-4}$
$K^*(892)^0\mu^+\mu^-$	B1	< 2.3	$\times 10^{-5}$
$e^\pm\mu^\mp$	LF	[c] < 5.9	$\times 10^{-6}$
$e^\pm\tau^\mp$	LF	[c] < 5.3	$\times 10^{-4}$
$\mu^\pm\tau^\mp$	LF	[c] < 8.3	$\times 10^{-4}$

B^* $I(J^P) = \frac{1}{2}(1^-)$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

Mass $m_{B^*} = 5324.8 \pm 2.1$ MeV

$m_{B^*} - m_B = 46.0 \pm 0.6$ MeV

BOTTOM, STRANGE MESONS

$(B = \pm 1, S = \mp 1)$

$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b$, similarly for $B_s^{*\pm}$'s

B_s^0 $I(J^P) = \frac{1}{2}(0^-)$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

Mass $m_{B_s^0} = 5375 \pm 6$ MeV ($S = 1.3$)

Mean life $\tau = (1.34_{-0.27}^{+0.32}) \times 10^{-12}$ s ($S = 1.4$)

$B_s^0-\bar{B}_s^0$ mixing parameters

$\chi_s = 0.62 \pm 0.13$

$\Delta m_{B_s^0} = m_{B_{s1}^0} - m_{B_{s2}^0} > 1.8 \times 10^{12} \hbar s^{-1}$, CL = 95%

$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0} > 2.0$, CL = 95%

B_s^0 DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
D_s^- anything	seen	-
$D_s^- \ell^+ \nu_\ell$ anything	seen	-
<i>(ℓ means sum of e and μ)</i>		
$D_s^- \pi^+$	seen	2325
$J/\psi(1S)\phi$	seen	1594
$\psi(2S)\phi$	seen	1128

HEAVY QUARK SEARCHES

Searches for Top and Fourth Generation Hadrons

See the sections "Searches for *t* Quark" and "Searches for *b'* (4^{th} Generation) Quark" at the end of the QUARKS section.

$c\bar{c}$ MESONS

**$\eta_c(1S)$
or $\eta_c(2980)$** $I^G(J^{PC}) = 0^+(0^{-+})$

Mass $m = 2978.8 \pm 1.9$ MeV ($S = 1.8$)

Full width $\Gamma = 10.3_{-3.4}^{+3.8}$ MeV

$\eta_c(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
Decays involving hadronic resonances			
$\eta'(958)\pi\pi$	(4.1 \pm 1.7) %		1319
$\rho\rho$	(2.6 \pm 0.9) %		1275
$K^*(892)^0 K^- \pi^+ + c.c.$	(2.0 \pm 0.7) %		1273
$K^*(892)\bar{K}^*(892)$	(8.5 \pm 3.1) $\times 10^{-3}$		1193
$\phi\phi$	(7.1 \pm 2.8) $\times 10^{-3}$		1086
$a_0(980)\pi$	< 2 %	90%	1323
$a_2(1320)\pi$	< 2 %	90%	1193
$K^*(892)\bar{K} + c.c.$	< 1.28 %	90%	1307
$f_2(1270)\eta$	< 1.1 %	90%	1142
$\omega\omega$	< 3.1 $\times 10^{-3}$	90%	1268
Decays into stable hadrons			
$K\bar{K}\pi$	(6.6 \pm 1.8) %		1378
$\eta\pi\pi$	(4.9 \pm 1.8) %		1425
$\pi^+\pi^- K^+ K^-$	(2.0 \pm 0.7) %		1342
$2(\pi^+\pi^-)$	(1.2 \pm 0.4) %		1457
$\rho\bar{\rho}$	(1.2 \pm 0.4) $\times 10^{-3}$		1157
$K\bar{K}\eta$	< 3.1 %	90%	1262
$\pi^+\pi^-\rho\bar{\rho}$	< 1.2 %	90%	1023
$\Lambda\bar{\Lambda}$	< 2 $\times 10^{-3}$	90%	987
Radiative decays			
$\gamma\gamma$	(6 \pm 6) $\times 10^{-4}$		1489

Meson Summary Table

J/ψ(1S)
or **J/ψ(3097)**

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 3096.88 \pm 0.04$ MeV

Full width $\Gamma = 88 \pm 5$ keV

$\Gamma_{ee} = 5.26 \pm 0.37$ keV (Assuming $\Gamma_{ee} = \Gamma_{\mu\mu}$)

J/ψ(1S) DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
hadrons	(86.0 ± 2.0) %		—
virtual $\gamma \rightarrow$ hadrons	(17.0 ± 2.0) %		—
e^+e^-	(5.99 ± 0.25) %		1548
$\mu^+\mu^-$	(5.97 ± 0.25) %	S=1.1	1545
Decays involving hadronic resonances			
$\rho\pi$	(1.28 ± 0.10) %		1449
$\rho^0\pi^0$	(4.2 ± 0.5) × 10 ⁻³		1449
$a_2(1320)\rho$	(1.09 ± 0.22) %		1125
$\omega\pi^+\pi^+\pi^-\pi^-$	(8.5 ± 3.4) × 10 ⁻³		1392
$\omega\pi^+\pi^-$	(7.2 ± 1.0) × 10 ⁻³		1435
$K^*(892)^0\bar{K}_2^*(1430)^0 + c.c.$	(6.7 ± 2.6) × 10 ⁻³		1005
$\omega K^*(892)\bar{K} + c.c.$	(5.3 ± 2.0) × 10 ⁻³		1098
$\omega f_2(1270)$	(4.3 ± 0.6) × 10 ⁻³		1143
$K^+\bar{K}^*(892)^- + c.c.$	(5.0 ± 0.4) × 10 ⁻³		1373
$K^0\bar{K}^*(892)^0 + c.c.$	(4.2 ± 0.4) × 10 ⁻³		1371
$\omega\pi^0\pi^0$	(3.4 ± 0.8) × 10 ⁻³		1436
$b_1(1235)^\pm\pi^\mp$	[q] (3.0 ± 0.5) × 10 ⁻³		1299
$\omega K^\pm K_S^0\pi^\mp$	[q] (3.0 ± 0.7) × 10 ⁻³		1210
$b_1(1235)^0\pi^0$	(2.3 ± 0.6) × 10 ⁻³		1299
$\phi K^*(892)\bar{K} + c.c.$	(2.04 ± 0.28) × 10 ⁻³		969
$\omega K\bar{K}$	(1.9 ± 0.4) × 10 ⁻³		1268
$\omega f_J(1710) \rightarrow \omega K\bar{K}$	(4.8 ± 1.1) × 10 ⁻⁴		878
$\phi 2(\pi^+\pi^-)$	(1.60 ± 0.32) × 10 ⁻³		1318
$\Delta(1232)^{++}\bar{p}\pi^-$	(1.6 ± 0.5) × 10 ⁻³		1030
$\omega\eta$	(1.58 ± 0.16) × 10 ⁻³		1394
$\phi K\bar{K}$	(1.48 ± 0.22) × 10 ⁻³		1179
$\phi f_J(1710) \rightarrow \phi K\bar{K}$	(3.6 ± 0.6) × 10 ⁻⁴		875
$\rho\bar{\rho}\omega$	(1.30 ± 0.25) × 10 ⁻³	S=1.3	769
$\Delta(1232)^{++}\bar{\Delta}(1232)^{--}$	(1.10 ± 0.29) × 10 ⁻³		938
$\Sigma(1385)^-\bar{\Sigma}(1385)^+ (or c.c.)$	[q] (1.03 ± 0.13) × 10 ⁻³		692
$\rho\bar{\rho}\eta(958)$	(9 ± 4) × 10 ⁻⁴	S=1.7	596
$\phi f_2'(1525)$	(8 ± 4) × 10 ⁻⁴	S=2.7	871
$\phi\pi^+\pi^-$	(8.0 ± 1.2) × 10 ⁻⁴		1365
$\phi K^\pm K_S^0\pi^\mp$	[q] (7.2 ± 0.9) × 10 ⁻⁴		1114
$\omega f_1(1420)$	(6.8 ± 2.4) × 10 ⁻⁴		1062
$\phi\eta$	(6.5 ± 0.7) × 10 ⁻⁴		1320
$\Xi(1530)^-\bar{\Xi}^+$	(5.9 ± 1.5) × 10 ⁻⁴		597
$\rho K^-\bar{\Sigma}(1385)^0$	(5.1 ± 3.2) × 10 ⁻⁴		645
$\omega\pi^0$	(4.2 ± 0.6) × 10 ⁻⁴	S=1.4	1447
$\phi\eta'(958)$	(3.3 ± 0.4) × 10 ⁻⁴		1192
$\phi f_0(980)$	(3.2 ± 0.9) × 10 ⁻⁴	S=1.9	1182
$\Xi(1530)^0\bar{\Xi}^0$	(3.2 ± 1.4) × 10 ⁻⁴		608
$\Sigma(1385)^-\bar{\Sigma}^+ (or c.c.)$	[q] (3.1 ± 0.5) × 10 ⁻⁴		857
$\phi f_1(1285)$	(2.6 ± 0.5) × 10 ⁻⁴	S=1.1	1032
$\rho\eta$	(1.93 ± 0.23) × 10 ⁻⁴		1398
$\omega\eta'(958)$	(1.67 ± 0.25) × 10 ⁻⁴		1279
$\omega f_0(980)$	(1.4 ± 0.5) × 10 ⁻⁴		1271
$\rho\eta'(958)$	(1.05 ± 0.18) × 10 ⁻⁴		1283
$\rho\bar{\rho}\phi$	(4.5 ± 1.5) × 10 ⁻⁵		527
$a_2(1320)^\pm\pi^\mp$	[q] < 4.3 × 10 ⁻³	CL=90%	1263
$K\bar{K}_2^*(1430) + c.c.$	< 4.0 × 10 ⁻³	CL=90%	1159
$K_2^*(1430)^0\bar{K}_2^*(1430)^0$	< 2.9 × 10 ⁻³	CL=90%	588
$K^*(892)^0\bar{K}^*(892)^0$	< 5 × 10 ⁻⁴	CL=90%	1263
$\phi f_2(1270)$	< 3.7 × 10 ⁻⁴	CL=90%	1036
$\rho\bar{\rho}\rho$	< 3.1 × 10 ⁻⁴	CL=90%	779
$\phi\eta(1440) \rightarrow \phi\eta\pi\pi$	< 2.5 × 10 ⁻⁴	CL=90%	946
$\omega f_2'(1525)$	< 2.2 × 10 ⁻⁴	CL=90%	1003
$\Sigma(1385)^0\bar{\Lambda}$	< 2 × 10 ⁻⁴	CL=90%	911
$\Delta(1232)^+\bar{p}$	< 1 × 10 ⁻⁴	CL=90%	1100
$\Sigma^0\bar{\Lambda}$	< 9 × 10 ⁻⁵	CL=90%	1032
$\phi\pi^0$	< 6.8 × 10 ⁻⁶	CL=90%	1377

Decays into stable hadrons

$2(\pi^+\pi^-)\pi^0$	(3.37 ± 0.26) %		1496
$3(\pi^+\pi^-)\pi^0$	(2.9 ± 0.6) %		1433
$\pi^+\pi^-\pi^0$	(1.50 ± 0.20) %		1533
$\pi^+\pi^-\pi^0 K^+K^-$	(1.20 ± 0.30) %		1368
$4(\pi^+\pi^-)\pi^0$	(9.0 ± 3.0) × 10 ⁻³		1345
$\pi^+\pi^-K^+K^-$	(7.2 ± 2.3) × 10 ⁻³		1407
$K\bar{K}\pi$	(6.1 ± 1.0) × 10 ⁻³		1440
$\rho\bar{\rho}\pi^+\pi^-$	(6.0 ± 0.5) × 10 ⁻³	S=1.3	1107
$2(\pi^+\pi^-)$	(4.0 ± 1.0) × 10 ⁻³		1517
$3(\pi^+\pi^-)$	(4.0 ± 2.0) × 10 ⁻³		1466
$\eta\bar{\eta}\pi^+\pi^-$	(4 ± 4) × 10 ⁻³		1106
$\Sigma\bar{\Sigma}$	(3.8 ± 0.5) × 10 ⁻³		992
$2(\pi^+\pi^-)K^+K^-$	(3.1 ± 1.3) × 10 ⁻³		1320
$\rho\bar{\rho}\pi^+\pi^-\pi^0$	[xx] (2.3 ± 0.9) × 10 ⁻³	S=1.9	1033
$\rho\bar{\rho}$	(2.14 ± 0.10) × 10 ⁻³		1232
$\rho\bar{\rho}\eta$	(2.09 ± 0.18) × 10 ⁻³		948
$\rho\bar{\rho}\pi^-$	(2.00 ± 0.10) × 10 ⁻³		1174
$\eta\bar{\eta}$	(1.9 ± 0.5) × 10 ⁻³		1231
$\Xi\bar{\Xi}$	(1.8 ± 0.4) × 10 ⁻³	S=1.8	818
$\Lambda\bar{\Lambda}$	(1.35 ± 0.14) × 10 ⁻³	S=1.2	1074
$\rho\bar{\rho}\pi^0$	(1.09 ± 0.09) × 10 ⁻³		1176
$\Lambda\bar{\Sigma}^-\pi^+ (or c.c.)$	[q] (1.06 ± 0.12) × 10 ⁻³		945
$\rho K^-\bar{\Lambda}$	(8.9 ± 1.6) × 10 ⁻⁴		876
$2(K^+K^-)$	(7.0 ± 3.0) × 10 ⁻⁴		1131
$\rho K^-\bar{\Sigma}^0$	(2.9 ± 0.8) × 10 ⁻⁴		820
K^+K^-	(2.37 ± 0.31) × 10 ⁻⁴		1468
$\Lambda\bar{\Lambda}\pi^0$	(2.2 ± 0.7) × 10 ⁻⁴		998
$\pi^+\pi^-$	(1.47 ± 0.23) × 10 ⁻⁴		1542
$K_S^0 K_L^0$	(1.08 ± 0.14) × 10 ⁻⁴		1466
$\Lambda\bar{\Sigma}^+ + c.c.$	< 1.5 × 10 ⁻⁴	CL=90%	1032
$K_S^0 K_S^0$	< 5.2 × 10 ⁻⁶	CL=90%	1466
Radiative decays			
$\gamma\eta_c(1S)$	(1.3 ± 0.4) %		116
$\gamma\pi^+\pi^-\pi^0$	(8.3 ± 3.1) × 10 ⁻³		1518
$\gamma\eta\pi\pi$	(6.1 ± 1.0) × 10 ⁻³		1487
$\gamma\eta(1440) \rightarrow \gamma K\bar{K}\pi$	[n] (9.1 ± 1.8) × 10 ⁻⁴		1223
$\gamma\eta(1440) \rightarrow \gamma\gamma\rho^0$	(6.4 ± 1.4) × 10 ⁻⁵		1223
$\gamma\rho\rho$	(4.5 ± 0.8) × 10 ⁻³		1343
$\gamma\eta'(958)$	(4.31 ± 0.30) × 10 ⁻³		1400
$\gamma 2\pi^+ 2\pi^-$	(2.8 ± 0.5) × 10 ⁻³	S=1.9	1517
$\gamma f_4(2050)$	(2.7 ± 0.7) × 10 ⁻³		874
$\gamma\omega\omega$	(1.59 ± 0.33) × 10 ⁻³		1337
$\gamma\eta(1440) \rightarrow \gamma\rho^0\rho^0$	(1.4 ± 0.4) × 10 ⁻³		1223
$\gamma f_2(1270)$	(1.38 ± 0.14) × 10 ⁻³		1286
$\gamma f_J(1710) \rightarrow \gamma K\bar{K}$	(9.7 ± 1.2) × 10 ⁻⁴		1075
$\gamma\eta$	(8.6 ± 0.8) × 10 ⁻⁴		1500
$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	(8.3 ± 1.5) × 10 ⁻⁴		1220
$\gamma f_1(1285)$	(6.5 ± 1.0) × 10 ⁻⁴		1283
$\gamma f_2'(1525)$	(6.3 ± 1.0) × 10 ⁻⁴		1173
$\gamma\phi\phi$	(4.0 ± 1.2) × 10 ⁻⁴	S=2.1	1166
$\gamma\rho\bar{\rho}$	(3.8 ± 1.0) × 10 ⁻⁴		1232
$\gamma\eta(2225)$	(2.9 ± 0.6) × 10 ⁻⁴		834
$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	(1.3 ± 0.9) × 10 ⁻⁴		1048
$\gamma\pi^0$	(3.9 ± 1.3) × 10 ⁻⁵		1546
$\gamma\rho\bar{\rho}\pi^+\pi^-$	< 7.9 × 10 ⁻⁴	CL=90%	1107
$\gamma\gamma$	< 5 × 10 ⁻⁴	CL=90%	1548
$\gamma\Lambda\bar{\Lambda}$	< 1.3 × 10 ⁻⁴	CL=90%	1074
3γ	< 5.5 × 10 ⁻⁵	CL=90%	1548

Meson Summary Table

$\chi_{c0}(1P)$
or **$\chi_{c0}(3415)$**

$$I^G(J^{PC}) = 0^+(0^{++})$$

Mass $m = 3415.1 \pm 1.0$ MeV
Full width $\Gamma = 14 \pm 5$ MeV

$\chi_{c0}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
Hadronic decays			
$2(\pi^+\pi^-)$	(3.7±0.7) %		1679
$\pi^+\pi^-K^+K^-$	(3.0±0.7) %		1580
$\rho^0\pi^+\pi^-$	(1.6±0.5) %		1608
$3(\pi^+\pi^-)$	(1.5±0.5) %		1633
$K^+\bar{K}^*(892)^0\pi^- + c.c.$	(1.2±0.4) %		1522
$\pi^+\pi^-$	(7.5±2.1) × 10 ⁻³		1702
K^+K^-	(7.1±2.4) × 10 ⁻³		1635
$\pi^+\pi^-\rho\bar{p}$	(5.0±2.0) × 10 ⁻³		1320
$\pi^0\pi^0$	(3.1±0.6) × 10 ⁻³		1702
$\eta\eta$	(2.5±1.1) × 10 ⁻³		1617
$\rho\bar{p}$	< 9.0 × 10 ⁻⁴	90%	1427
Radiative decays			
$\gamma J/\psi(1S)$	(6.6±1.8) × 10 ⁻³		303
$\gamma\gamma$	(4.0±2.3) × 10 ⁻⁴		1708

$\chi_{c1}(1P)$
or **$\chi_{c1}(3510)$**

$$I^G(J^{PC}) = 0^+(1^{++})$$

Mass $m = 3510.53 \pm 0.12$ MeV
Full width $\Gamma = 0.88 \pm 0.14$ MeV

$\chi_{c1}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
Hadronic decays		
$3(\pi^+\pi^-)$	(2.2±0.8) %	1683
$2(\pi^+\pi^-)$	(1.6±0.5) %	1727
$\pi^+\pi^-K^+K^-$	(9 ± 4) × 10 ⁻³	1632
$\rho^0\pi^+\pi^-$	(3.9±3.5) × 10 ⁻³	1659
$K^+\bar{K}^*(892)^0\pi^- + c.c.$	(3.2±2.1) × 10 ⁻³	1576
$\pi^+\pi^-\rho\bar{p}$	(1.4±0.9) × 10 ⁻³	1381
$\rho\bar{p}$	(8.6±1.2) × 10 ⁻⁵	1483
$\pi^+\pi^- + K^+K^-$	< 2.1 × 10 ⁻³	-
Radiative decays		
$\gamma J/\psi(1S)$	(27.3±1.6) %	389

$\chi_{c2}(1P)$
or **$\chi_{c2}(3555)$**

$$I^G(J^{PC}) = 0^+(2^{++})$$

Mass $m = 3556.17 \pm 0.13$ MeV
Full width $\Gamma = 2.00 \pm 0.18$ MeV

$\chi_{c2}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
Hadronic decays			
$2(\pi^+\pi^-)$	(2.2 ± 0.5) %		1751
$\pi^+\pi^-K^+K^-$	(1.9 ± 0.5) %		1656
$3(\pi^+\pi^-)$	(1.2 ± 0.8) %		1707
$\rho^0\pi^+\pi^-$	(7 ± 4) × 10 ⁻³		1683
$K^+\bar{K}^*(892)^0\pi^- + c.c.$	(4.8 ± 2.8) × 10 ⁻³		1601
$\pi^+\pi^-\rho\bar{p}$	(3.3 ± 1.3) × 10 ⁻³		1410
$\pi^+\pi^-$	(1.9 ± 1.0) × 10 ⁻³		1773
K^+K^-	(1.5 ± 1.1) × 10 ⁻³		1708
$\rho\bar{p}$	(10.0 ± 1.0) × 10 ⁻⁵		1510
$\pi^0\pi^0$	(1.10±0.28) × 10 ⁻³		1773
$\eta\eta$	(8 ± 5) × 10 ⁻⁴		1692
$J/\psi(1S)\pi^+\pi^-\pi^0$	< 1.5 %	90%	185
Radiative decays			
$\gamma J/\psi(1S)$	(13.5 ± 1.1) %		430
$\gamma\gamma$	(1.6 ± 0.5) × 10 ⁻⁴		1778

$\psi(2S)$
or **$\psi(3685)$**

$$I^G(J^{PC}) = 0^-(1^{--})$$

Mass $m = 3686.00 \pm 0.09$ MeV
Full width $\Gamma = 277 \pm 31$ keV ($S = 1.1$)
 $\Gamma_{ee} = 2.14 \pm 0.21$ keV (Assuming $\Gamma_{ee} = \Gamma_{\mu\mu}$)

$\psi(2S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	Scale factor/ ρ (MeV/c)
Decays into $J/\psi(1S)$ and anything			
hadrons	(98.10±0.30) %		-
virtual $\gamma \rightarrow$ hadrons	(2.9 ± 0.4) %		-
e^+e^-	(8.8 ± 1.3) × 10 ⁻³		1843
$\mu^+\mu^-$	(7.7 ± 1.7) × 10 ⁻³		1840
Hadronic decays			
$3(\pi^+\pi^-)\pi^0$	(3.5 ± 1.6) × 10 ⁻³		1746
$2(\pi^+\pi^-)\pi^0$	(3.1 ± 0.7) × 10 ⁻³		1799
$\pi^+\pi^-K^+K^-$	(1.6 ± 0.4) × 10 ⁻³		1726
$\pi^+\pi^-\rho\bar{p}$	(8.0 ± 2.0) × 10 ⁻⁴		1491
$K^+\bar{K}^*(892)^0\pi^- + c.c.$	(6.7 ± 2.5) × 10 ⁻⁴		1673
$2(\pi^+\pi^-)$	(4.5 ± 1.0) × 10 ⁻⁴		1817
$\rho^0\pi^+\pi^-$	(4.2 ± 1.5) × 10 ⁻⁴		1751
$\bar{p}p$	(1.9 ± 0.5) × 10 ⁻⁴		1586
$3(\pi^+\pi^-)$	(1.5 ± 1.0) × 10 ⁻⁴		1774
$\bar{p}p\pi^0$	(1.4 ± 0.5) × 10 ⁻⁴		1543
K^+K^-	(1.0 ± 0.7) × 10 ⁻⁴		1776
$\pi^+\pi^-\pi^0$	(9 ± 5) × 10 ⁻⁵		1830
$\pi^+\pi^-$	(8 ± 5) × 10 ⁻⁵		1838
$\Lambda\bar{\Lambda}$	< 4 × 10 ⁻⁴	CL=90%	1467
$\Xi^-\bar{\Xi}^+$	< 2 × 10 ⁻⁴	CL=90%	1285
$\rho\pi$	< 8.3 × 10 ⁻⁵	CL=90%	1760
$K^+K^-\pi^0$	< 2.96 × 10 ⁻⁵	CL=90%	1754
$K^+\bar{K}^*(892)^-\pi^0 + c.c.$	< 1.79 × 10 ⁻⁵	CL=90%	1698
Radiative decays			
$\gamma\chi_{c0}(1P)$	(9.3 ± 0.8) %		261
$\gamma\chi_{c1}(1P)$	(8.7 ± 0.8) %		171
$\gamma\chi_{c2}(1P)$	(7.8 ± 0.8) %		127
$\gamma\eta_c(1S)$	(2.8 ± 0.6) × 10 ⁻³		639
$\gamma\pi^0$	< 5.4 × 10 ⁻³	CL=95%	1841
$\gamma\eta'(958)$	< 1.1 × 10 ⁻³	CL=90%	1719
$\gamma\gamma$	< 1.6 × 10 ⁻⁴	CL=90%	1843
$\gamma\eta(1440) \rightarrow \gamma K\bar{K}\pi$	[n] < 1.2 × 10 ⁻⁴	CL=90%	1569

$\psi(3770)$

$$I^G(J^{PC}) = ?^?(1^{--})$$

Mass $m = 3769.9 \pm 2.5$ MeV ($S = 1.8$)
Full width $\Gamma = 23.6 \pm 2.7$ MeV ($S = 1.1$)
 $\Gamma_{ee} = 0.26 \pm 0.04$ keV ($S = 1.2$)

$\psi(3770)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ ρ (MeV/c)
$D\bar{D}$	dominant	242
e^+e^-	(1.12±0.17) × 10 ⁻⁵	1.2 1885

$\psi(4040)$ [yy]

$$I^G(J^{PC}) = ?^?(1^{--})$$

Mass $m = 4040 \pm 10$ MeV
Full width $\Gamma = 52 \pm 10$ MeV
 $\Gamma_{ee} = 0.75 \pm 0.15$ keV

$\psi(4040)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
e^+e^-	(1.4±0.4) × 10 ⁻⁵	2020
$D^0\bar{D}^0$	seen	777
$D^*(2007)^0\bar{D}^0 + c.c.$	seen	578
$D^*(2007)^0\bar{D}^*(2007)^0$	seen	232

Meson Summary Table

 $\psi(4160)$ [$\psi\psi$]

$$I^G(J^{PC}) = ?^?(1^- -)$$

Mass $m = 4159 \pm 20$ MeV
 Full width $\Gamma = 78 \pm 20$ MeV
 $\Gamma_{ee} = 0.77 \pm 0.23$ keV

$\psi(4160)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$e^+ e^-$	$(10 \pm 4) \times 10^{-6}$	2079

 $\psi(4415)$ [$\psi\psi$]

$$I^G(J^{PC}) = ?^?(1^- -)$$

Mass $m = 4415 \pm 6$ MeV
 Full width $\Gamma = 43 \pm 15$ MeV ($S = 1.8$)
 $\Gamma_{ee} = 0.47 \pm 0.10$ keV

$\psi(4415)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
hadrons	dominant	-
$e^+ e^-$	$(1.1 \pm 0.4) \times 10^{-5}$	2207

 $b\bar{b}$ MESONS **$T(1S)$
or $T(9460)$**

$$I^G(J^{PC}) = ?^?(1^- -)$$

Mass $m = 9460.37 \pm 0.21$ MeV ($S = 2.7$)
 Full width $\Gamma = 52.5 \pm 1.8$ keV
 $\Gamma_{ee} = 1.32 \pm 0.03$ keV

$T(1S)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	ρ (MeV/c)
$\tau^+ \tau^-$	$(2.97 \pm 0.35) \%$		4384
$e^+ e^-$	$(2.52 \pm 0.17) \%$		4730
$\mu^+ \mu^-$	$(2.48 \pm 0.07) \%$	$S=1.1$	4729
Hadronic decays			
$J/\psi(1S)$ anything	$(1.1 \pm 0.4) \times 10^{-3}$		4223
$\rho\pi$	$< 2 \times 10^{-4}$	CL=90%	4698
$\pi^+ \pi^-$	$< 5 \times 10^{-4}$	CL=90%	4728
$K^+ K^-$	$< 5 \times 10^{-4}$	CL=90%	4704
$\rho\bar{\rho}$	$< 9 \times 10^{-4}$	CL=90%	4636
Radiative decays			
$\gamma 2h^+ 2h^-$	$(7.0 \pm 1.5) \times 10^{-4}$		4720
$\gamma 3h^+ 3h^-$	$(5.4 \pm 2.0) \times 10^{-4}$		4703
$\gamma 4h^+ 4h^-$	$(7.4 \pm 3.5) \times 10^{-4}$		4679
$\gamma \pi^+ \pi^- K^+ K^-$	$(2.9 \pm 0.9) \times 10^{-4}$		4686
$\gamma 2\pi^+ 2\pi^-$	$(2.5 \pm 0.9) \times 10^{-4}$		4720
$\gamma 3\pi^+ 3\pi^-$	$(2.5 \pm 1.2) \times 10^{-4}$		4703
$\gamma 2\pi^+ 2\pi^- K^+ K^-$	$(2.4 \pm 1.2) \times 10^{-4}$		4658
$\gamma \pi^+ \pi^- \rho\bar{\rho}$	$(1.5 \pm 0.6) \times 10^{-4}$		4604
$\gamma 2\pi^+ 2\pi^- \rho\bar{\rho}$	$(4 \pm 6) \times 10^{-5}$		4563
$\gamma 2K^+ 2K^-$	$(2.0 \pm 2.0) \times 10^{-5}$		4601
$\gamma \eta'(958)$	$< 1.3 \times 10^{-3}$	CL=90%	4682
$\gamma \eta$	$< 3.5 \times 10^{-4}$	CL=90%	4714
$\gamma f_2'(1525)$	$< 1.4 \times 10^{-4}$	CL=90%	4607
$\gamma f_2(1270)$	$< 1.3 \times 10^{-4}$	CL=90%	4644
$\gamma \eta(1440)$	$< 8.2 \times 10^{-5}$	CL=90%	4624
$\gamma f_j(1710) \rightarrow \gamma K\bar{K}$	$< 2.6 \times 10^{-4}$	CL=90%	4576
$\gamma f_4(2220) \rightarrow \gamma K^+ K^-$	$< 1.5 \times 10^{-5}$	CL=90%	4469

 **$X_{b0}(1P)$ [zz]
or $X_{b0}(9860)$**

$$I^G(J^{PC}) = ?^?(0 \text{ preferred } ++)$$

 J needs confirmation.

Mass $m = 9859.8 \pm 1.3$ MeV

$X_{b0}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\gamma T(1S)$	$< 6 \%$	90%	391

 **$X_{b1}(1P)$ [zz]
or $X_{b1}(9890)$**

$$I^G(J^{PC}) = ?^?(1^+ +)$$

 J needs confirmation.

Mass $m = 9891.9 \pm 0.7$ MeV

$X_{b1}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\gamma T(1S)$	$(35 \pm 8) \%$	422

 **$X_{b2}(1P)$ [zz]
or $X_{b2}(9915)$**

$$I^G(J^{PC}) = ?^?(2^+ +)$$

 J needs confirmation.

Mass $m = 9913.2 \pm 0.6$ MeV

$X_{b2}(1P)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\gamma T(1S)$	$(22 \pm 4) \%$	443

 **$T(2S)$
or $T(10023)$**

$$I^G(J^{PC}) = ?^?(1^- -)$$

Mass $m = 10.02330 \pm 0.00031$ GeV
 Full width $\Gamma = 44 \pm 7$ keV

$T(2S)$ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$T(1S) \pi^+ \pi^-$	$(18.5 \pm 0.8) \%$		475
$T(1S) \pi^0 \pi^0$	$(8.8 \pm 1.1) \%$		480
$\tau^+ \tau^-$	$(1.7 \pm 1.6) \%$		4686
$\mu^+ \mu^-$	$(1.31 \pm 0.21) \%$		5011
$e^+ e^-$	seen		5012
$T(1S) \pi^0$	$< 8 \times 10^{-3}$	90%	531
$T(1S) \eta$	$< 2 \times 10^{-3}$	90%	127
$J/\psi(1S)$ anything	$< 6 \times 10^{-3}$	90%	4533

Radiative decays

$\gamma X_{b1}(1P)$	$(6.7 \pm 0.9) \%$		131
$\gamma X_{b2}(1P)$	$(6.6 \pm 0.9) \%$		110
$\gamma X_{b0}(1P)$	$(4.3 \pm 1.0) \%$		162
$\gamma f_j(1710)$	$< 5.9 \times 10^{-4}$	90%	4866
$\gamma f_2'(1525)$	$< 5.3 \times 10^{-4}$	90%	4896
$\gamma f_2(1270)$	$< 2.41 \times 10^{-4}$	90%	4931

 **$X_{b0}(2P)$ [zz]
or $X_{b0}(10235)$**

$$I^G(J^{PC}) = ?^?(0 \text{ preferred } ++)$$

 J needs confirmation.

Mass $m = 10.2321 \pm 0.0006$ GeV

$X_{b0}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\gamma T(2S)$	$(4.6 \pm 2.1) \%$	210
$\gamma T(1S)$	$(9 \pm 6) \times 10^{-3}$	746

 **$X_{b1}(2P)$ [zz]
or $X_{b1}(10255)$**

$$I^G(J^{PC}) = ?^?(1 \text{ preferred } ++)$$

 J needs confirmation.

Mass $m = 10.2552 \pm 0.0005$ GeV
 $m_{X_{b1}(2P)} - m_{X_{b0}(2P)} = 23.5 \pm 1.0$ MeV

$X_{b1}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	ρ (MeV/c)
$\gamma T(2S)$	$(21 \pm 4) \%$	1.5	229
$\gamma T(1S)$	$(8.5 \pm 1.3) \%$	1.3	764

 **$X_{b2}(2P)$ [zz]
or $X_{b2}(10270)$**

$$I^G(J^{PC}) = ?^?(2 \text{ preferred } ++)$$

 J needs confirmation.

Mass $m = 10.2685 \pm 0.0004$ GeV
 $m_{X_{b2}(2P)} - m_{X_{b1}(2P)} = 13.5 \pm 0.6$ MeV

$X_{b2}(2P)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\gamma T(2S)$	$(16.2 \pm 2.4) \%$	242
$\gamma T(1S)$	$(7.1 \pm 1.0) \%$	776

Meson Summary Table

T(3S)
or **T(10355)**

$${}^1G(J^{PC}) = ?^?(1^{--})$$

Mass $m = 10.3553 \pm 0.0005$ GeV
Full width $\Gamma = 26.3 \pm 3.5$ keV

T(3S) DECAY MODES	Fraction (Γ_i/Γ)	Scale factor	ρ (MeV/c)
T(2S) anything	(10.6 \pm 0.8) %		296
T(2S) $\pi^+ \pi^-$	(2.8 \pm 0.6) %	2.2	177
T(2S) $\pi^0 \pi^0$	(2.00 \pm 0.32) %		190
T(2S) $\gamma\gamma$	(5.0 \pm 0.7) %		–
T(1S) $\pi^+ \pi^-$	(4.48 \pm 0.21) %		814
T(1S) $\pi^0 \pi^0$	(2.06 \pm 0.28) %		816
$\mu^+ \mu^-$	(1.81 \pm 0.17) %		5177
$e^+ e^-$	seen		5177
Radiative decays			
$\gamma X_{b2}(2P)$	(11.4 \pm 0.8) %	1.3	87
$\gamma X_{b1}(2P)$	(11.3 \pm 0.6) %		100
$\gamma X_{b0}(2P)$	(5.4 \pm 0.6) %	1.1	123

T(4S)
or **T(10580)**

$${}^1G(J^{PC}) = ?^?(1^{--})$$

Mass $m = 10.5800 \pm 0.0035$ GeV
Full width $\Gamma = 23.8 \pm 2.2$ MeV
 $\Gamma_{ee} = 0.24 \pm 0.05$ keV ($S = 1.7$)

T(4S) DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$e^+ e^-$	(1.01 \pm 0.21) $\times 10^{-5}$		5290
D^{*+} anything + c.c.	< 7.4 %	90%	5099
ϕ anything	< 2.3 $\times 10^{-3}$	90%	5240
T(1S) anything	< 4 $\times 10^{-3}$	90%	1053

T(10860)

$${}^1G(J^{PC}) = ?^?(1^{--})$$

Mass $m = 10.865 \pm 0.008$ GeV ($S = 1.1$)
Full width $\Gamma = 110 \pm 13$ MeV
 $\Gamma_{ee} = 0.31 \pm 0.07$ keV ($S = 1.3$)

T(10860) DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$e^+ e^-$	(2.8 \pm 0.7) $\times 10^{-6}$	5432

T(11020)

$${}^1G(J^{PC}) = ?^?(1^{--})$$

Mass $m = 11.019 \pm 0.008$ GeV
Full width $\Gamma = 79 \pm 16$ MeV
 $\Gamma_{ee} = 0.130 \pm 0.030$ keV

T(11020) DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$e^+ e^-$	(1.6 \pm 0.5) $\times 10^{-6}$	5509

Searches for Top and Fourth Generation HadronsSee the sections "Searches for t Quark" and "Searches for b' (4^{th} Generation) Quark" at the end of the QUARKS section.

NOTES

In this Summary Table:

When a quantity has "($S = \dots$)" to its right, the error on the quantity has been enlarged by the "scale factor" S , defined as $S = \sqrt{\chi^2/(N-1)}$, where N is the number of measurements used in calculating the quantity.

We do this when $S > 1$, which often indicates that the measurements are inconsistent. When $S > 1.25$, we also show in the Full Listings an ideogram of the measurements. For more about S , see the Introduction.

A decay momentum p is given for each decay mode. For a 2-body decay, p is the momentum of each decay product in the rest frame of the decaying particle. For a 3-or-more-body decay, p is the largest momentum any of the products can have in this frame.

[a] The π^\pm mass has increased by three (old) standard deviations since our 1992 edition, and the π^0 mass, which is determined using the mass difference ($m_{\pi^\pm} - m_{\pi^0}$), has increased accordingly. See the "Note on the Charged Pion Mass" in the π^\pm Full Listings for a discussion.

[b] See the "Note on $\pi^\pm \rightarrow \ell^\pm \nu \gamma$ and $K^\pm \rightarrow \ell^\pm \nu \gamma$ Form Factors" in the π^\pm Full Listings for definitions and details.

[c] Measurements of $\Gamma(e^+ \nu_e)/\Gamma(\mu^+ \nu_\mu)$ always include decays with γ 's, and measurements of $\Gamma(e^+ \nu_e \gamma)$ and $\Gamma(\mu^+ \nu_\mu \gamma)$ never include low-energy γ 's. Therefore, since no clean separation is possible, we consider the modes with γ 's to be subreactions of the modes without them, and let $[\Gamma(e^+ \nu_e) + \Gamma(\mu^+ \nu_\mu)]/\Gamma_{\text{total}} = 100\%$.

[d] See the π^\pm Full Listings for the energy limits used in this measurement; low-energy γ 's are not included.

[e] Derived from an analysis of neutrino-oscillation experiments.

[f] Astrophysical and cosmological arguments give limits of order 10^{-13} ; see the π^0 Full Listings.

[g] See the "Note on the Decay Width $\Gamma(\eta \rightarrow \gamma\gamma)$ " in the η Full Listings.

[h] See the "Note on η Decay Parameters" in the η Full Listings.

[i] C parity forbids this to occur as a single-photon process.

[j] The $e^+ e^-$ branching fraction is from $e^+ e^- \rightarrow \pi^+ \pi^-$ experiments only. The $\omega \rho$ interference is then due to $\omega \rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+ \mu^-) = \Gamma(\rho^0 \rightarrow e^+ e^-) \times 0.99785$.

[k] This is only an educated guess; the error given is larger than the error on the average of the published values. See the Full Listings for details.

[l] See the "Note on the $f_1(1420)$ " in the $f_1(1420)$ Full Listings.

[m] See also the $\omega(1600)$ Full Listings.

[n] See the "Note on the $\eta(1440)$ " in the $\eta(1440)$ Full Listings.

[o] See the "Note on the $\rho(1450)$ and the $\rho(1700)$ " in the $\rho(1700)$ Full Listings.

[p] See also the $\omega(1420)$ Full Listings.

[q] The value is for the sum of the charge states indicated.

[r] The definition of the slope parameter g of the $K \rightarrow 3\pi$ Dalitz plot is as follows (see also "Note on Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays" in the K^\pm Full Listings):

$$|M|^2 = 1 + g(s_3 - s_0)/m_{\pi^+}^2 + \dots$$

[s] For more details and definitions of parameters see the Full Listings.

[t] See the K^\pm Full Listings for the energy limits used in this measurement.

[u] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ 's.

[v] Direct-emission branching fraction.

[w] Structure-dependent part.

[x] The CP -violation parameters are defined as follows (see also "Note on CP Violation in $K_S \rightarrow 3\pi$ " and "Note on CP Violation in K_L^0 Decay" in the Full Listings):

$$\eta_{+-} = |\eta_{+-}| e^{i\phi_{+-}} = \frac{A(K_L^0 \rightarrow \pi^+ \pi^-)}{A(K_S^0 \rightarrow \pi^+ \pi^-)} = \epsilon + \epsilon'$$

$$\eta_{00} = |\eta_{00}| e^{i\phi_{00}} = \frac{A(K_L^0 \rightarrow \pi^0 \pi^0)}{A(K_S^0 \rightarrow \pi^0 \pi^0)} = \epsilon - 2\epsilon'$$

$$\delta = \frac{\Gamma(K_L^0 \rightarrow \pi^- \ell^+ \nu) - \Gamma(K_L^0 \rightarrow \pi^+ \ell^- \nu)}{\Gamma(K_L^0 \rightarrow \pi^- \ell^+ \nu) + \Gamma(K_L^0 \rightarrow \pi^+ \ell^- \nu)},$$

$$\text{Im}(\eta_{+-0})^2 = \frac{\Gamma(K_S^0 \rightarrow \pi^+ \pi^- \pi^0)_{CP \text{ viol.}}}{\Gamma(K_L^0 \rightarrow \pi^+ \pi^- \pi^0)},$$

$$\text{Im}(\eta_{000})^2 = \frac{\Gamma(K_S^0 \rightarrow \pi^0 \pi^0 \pi^0)}{\Gamma(K_L^0 \rightarrow \pi^0 \pi^0 \pi^0)}.$$

where for the last two relations CPT is assumed valid, *i.e.*, $\text{Re}(\eta_{+-0}) \simeq 0$ and $\text{Re}(\eta_{000}) \simeq 0$.

[y] See the K_S^0 Full Listings for the energy limits used in this measurement.

[z] Calculated from K_L^0 semileptonic rates and the K_S^0 lifetime assuming $\Delta S = \Delta Q$.

[aa] ϵ'/ϵ is derived from $|\eta_{00}/\eta_{+-}|$ measurements using theoretical input on phases.

[bb] See the K_L^0 Full Listings for the energy limits used in this measurement.

[cc] $m_{e^+e^-} > 470$ MeV

[dd] Allowed by higher-order electroweak interactions.

[ee] Violates CP in leading order. Test of direct CP violation since the indirect CP -violating and CP -conserving contributions are expected to be suppressed.

[ff] See the note in the $L(1770)$ Full Listings in Reviews of Modern Physics **56** No. 2 Pt. II (1984), p. S200.

[gg] This is a weighted average of D^\pm (44%) and D^0 (56%) branching fractions. See " D^+ and $D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$ " under " D^+ Branching Ratios" in the Full Listings.

[hh] This value combines the e^+ and μ^+ branching fractions, making a small phase-space adjustment to the μ^+ fraction to be able to use it as an e^+ fraction; hence the " e^+ ." In fact, some of the e^+ measurements already use μ^+ events in this way.

[ii] ℓ indicates e or μ mode, not sum over modes.

[jj] The branching fractions for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers in the Full Listings.

[kk] The two experiments determining this ratio are in serious disagreement. See the Full Listings.

[ll] This mode is not a useful test for a $\Delta C=1$ weak neutral current because both quarks must change flavor in this decay.

[mm] The $D_1^0 D_2^0$ limits are inferred from the limit on $D^0 \rightarrow \bar{D}^0 \rightarrow K^+ \pi^-$.

[nn] See the "Note on Semileptonic Decays of D and B Mesons" in the D^+ Full Listings for a comparison of inclusive and summed-inclusive branching fractions.

[oo] The limit on $(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$ just below is much stronger.

[pp] For now, we average together measurements of the $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$ branching fractions.

[qq] This branching fraction is calculated from appropriate fractions of the next three branching fractions.

[rr] For admixture of B hadrons at LEP and Tevatron energies.

[ss] These values are model dependent. See note on "Semileptonic Decays" in the B^+ Full Listings.

[tt] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.

[uu] B^0 , B^+ , B_s^0 , and B baryon states not separated.

[vv] B^0 , B^+ , and B_s^0 not separated.

[ww] Derived from measurements of χ_d and of Δm_{B^0} times B^0 mean life.

[xx] Includes $p\bar{p}\pi^+\pi^-\gamma$ and excludes $p\bar{p}\eta$, $p\bar{p}\omega$, $p\bar{p}\eta'$.

[yy] J^{PC} known by production in e^+e^- via single photon annihilation. J^G is not known; interpretation of this state as a single resonance is unclear because of the expectation of substantial threshold effects in this energy region.

[zz] Spectroscopic labeling for these states is theoretical, pending experimental information.

Baryon Summary Table

This short table gives the name, the quantum numbers (where known), and the status of baryons in the Review. Only the baryons with 3- or 4-star status are included in the main Baryon Summary Table. Due to insufficient data or uncertain interpretation, the other entries in the short table are not established as baryons. The names with masses are of baryons that decay strongly. See our 1986 edition (Physics Letters **170B**) for listings of evidence for Z baryons (KN resonances).

p	P_{11}	****	$\Delta(1232)$	P_{33}	****	Λ	P_{01}	****	Σ^+	P_{11}	****	Ξ^0	P_{11}	****
n	P_{11}	****	$\Delta(1600)$	P_{33}	***	$\Lambda(1405)$	S_{01}	****	Σ^0	P_{11}	****	Ξ^-	P_{11}	****
$N(1440)$	P_{11}	****	$\Delta(1620)$	S_{31}	****	$\Lambda(1520)$	D_{03}	****	Σ^-	P_{11}	****	$\Xi(1530)$	P_{13}	****
$N(1520)$	D_{13}	****	$\Delta(1700)$	D_{33}	****	$\Lambda(1600)$	P_{01}	***	$\Sigma(1385)$	P_{13}	****	$\Xi(1620)$		*
$N(1535)$	S_{11}	****	$\Delta(1750)$	P_{31}	*	$\Lambda(1670)$	S_{01}	****	$\Sigma(1480)$		*	$\Xi(1690)$		***
$N(1650)$	S_{11}	****	$\Delta(1900)$	S_{31}	***	$\Lambda(1690)$	D_{03}	****	$\Sigma(1560)$		**	$\Xi(1820)$	D_{13}	***
$N(1675)$	D_{15}	****	$\Delta(1905)$	F_{35}	****	$\Lambda(1800)$	S_{01}	***	$\Sigma(1580)$	D_{13}	**	$\Xi(1950)$		***
$N(1680)$	F_{15}	****	$\Delta(1910)$	P_{31}	****	$\Lambda(1810)$	P_{01}	***	$\Sigma(1620)$	S_{11}	**	$\Xi(2030)$		***
$N(1700)$	D_{13}	***	$\Delta(1920)$	P_{33}	***	$\Lambda(1820)$	F_{05}	****	$\Sigma(1660)$	P_{11}	***	$\Xi(2120)$		*
$N(1710)$	P_{11}	***	$\Delta(1930)$	D_{35}	***	$\Lambda(1830)$	D_{05}	****	$\Sigma(1670)$	D_{13}	****	$\Xi(2250)$		**
$N(1720)$	P_{13}	****	$\Delta(1940)$	D_{33}	*	$\Lambda(1890)$	P_{03}	****	$\Sigma(1690)$		**	$\Xi(2370)$		**
$N(1900)$	P_{13}	*	$\Delta(1950)$	F_{37}	****	$\Lambda(2000)$		*	$\Sigma(1750)$	S_{11}	***	$\Xi(2500)$		*
$N(1990)$	F_{17}	**	$\Delta(2000)$	F_{35}	*	$\Lambda(2020)$	F_{07}	*	$\Sigma(1770)$	P_{11}	*			
$N(2000)$	F_{15}	**	$\Delta(2150)$	S_{31}	*	$\Lambda(2100)$	G_{07}	****	$\Sigma(1775)$	D_{15}	****	Ω^-		****
$N(2080)$	D_{13}	**	$\Delta(2200)$	G_{37}	*	$\Lambda(2110)$	F_{05}	***	$\Sigma(1840)$	P_{13}	*	$\Omega(2250)^-$		***
$N(2090)$	S_{11}	*	$\Delta(2300)$	H_{39}	**	$\Lambda(2325)$	D_{03}	*	$\Sigma(1880)$	P_{11}	**	$\Omega(2380)^-$		**
$N(2100)$	P_{11}	*	$\Delta(2350)$	D_{35}	*	$\Lambda(2350)$	H_{09}	***	$\Sigma(1915)$	F_{15}	****	$\Omega(2470)^-$		**
$N(2190)$	G_{17}	****	$\Delta(2390)$	F_{37}	*	$\Lambda(2585)$		**	$\Sigma(1940)$	D_{13}	***	Λ_c^+		****
$N(2200)$	D_{15}	**	$\Delta(2400)$	G_{39}	**				$\Sigma(2000)$	S_{11}	*	$\Lambda_c(2625)^+$		***
$N(2220)$	H_{19}	****	$\Delta(2420)$	$H_{3,11}$	****				$\Sigma(2030)$	F_{17}	****	$\Sigma_c(2455)$		****
$N(2250)$	G_{19}	****	$\Delta(2750)$	$l_{3,13}$	**				$\Sigma(2070)$	F_{15}	*	$\Sigma_c(2530)$		*
$N(2600)$	$l_{1,11}$	***	$\Delta(2950)$	$K_{3,15}$	**				$\Sigma(2080)$	P_{13}	**	Ξ_c^+		***
$N(2700)$	$K_{1,13}$	**							$\Sigma(2100)$	G_{17}	*	Ξ_c^0		***
									$\Sigma(2250)$		***	Ω_c^0		**
									$\Sigma(2455)$		**	Λ_b^0		***
									$\Sigma(2620)$		**			
									$\Sigma(3000)$		*			
									$\Sigma(3170)$		*			

**** Existence is certain, and properties are at least fairly well explored.

*** Existence ranges from very likely to certain, but further confirmation is desirable and/or quantum numbers, branching fractions, etc. are not well determined.

** Evidence of existence is only fair.

* Evidence of existence is poor.

Baryon Summary Table

N BARYONS

$(S = 0, I = 1/2)$

$$p, N^+ = uud; \quad n, N^0 = udd$$

p

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 938.27231 \pm 0.00028$ MeV [a]
 $= 1.007276470 \pm 0.000000012$ u
 $m_{\bar{p}}/m_p = 0.99999998 \pm 0.00000004$
 $|q_p + q_{\bar{p}}|/e < 2 \times 10^{-5}$
 $|q_p + q_e|/e < 1.0 \times 10^{-21}$ [b]
Magnetic moment $\mu = 2.79284739 \pm 0.00000006 \mu_N$
Electric dipole moment $d = (-4 \pm 6) \times 10^{-23}$ e cm
Electric polarizability $\bar{\alpha} = (10.2 \pm 0.9) \times 10^{-4}$ fm³
Magnetic polarizability $\bar{\beta} = (4.0 \pm 0.9) \times 10^{-4}$ fm³
Mean life $\tau > 1.6 \times 10^{25}$ years (independent of mode)
 $> 10^{31} - 5 \times 10^{32}$ years [c] (mode dependent)

For N decays, p and n distinguish proton and neutron partial lifetimes. See also the "Note on Proton Mean Life Limits" in the Full Listings.

The "partial mean life" limits tabulated here are the limits on τ/B_j , where τ is the total mean life and B_j is the branching fraction for the mode in question.

p DECAY MODES	Partial mean life (10 ³⁰ years)	Confidence level	$\frac{p}{c}$ (MeV/c)
Antilepton + meson			
$N \rightarrow e^+ \pi$	> 130 (n), > 550 (p)	90%	459
$N \rightarrow \mu^+ \pi$	> 100 (n), > 270 (p)	90%	453
$N \rightarrow \nu \pi$	> 100 (n), > 25 (p)	90%	459
$p \rightarrow e^+ \eta$	> 140	90%	309
$p \rightarrow \mu^+ \eta$	> 69	90%	296
$n \rightarrow \nu \eta$	> 54	90%	310
$N \rightarrow e^+ \rho$	> 58 (n), > 75 (p)	90%	153
$N \rightarrow \mu^+ \rho$	> 23 (n), > 110 (p)	90%	119
$N \rightarrow \nu \rho$	> 19 (n), > 27 (p)	90%	153
$p \rightarrow e^+ \omega$	> 45	90%	142
$p \rightarrow \mu^+ \omega$	> 57	90%	104
$n \rightarrow \nu \omega$	> 43	90%	144
$N \rightarrow e^+ K$	> 1.3 (n), > 150 (p)	90%	337
$p \rightarrow e^+ K_S^0$	> 76	90%	337
$p \rightarrow e^+ K_L^0$	> 44	90%	337
$N \rightarrow \mu^+ K$	> 1.1 (n), > 120 (p)	90%	326
$p \rightarrow \mu^+ K_S^0$	> 64	90%	326
$p \rightarrow \mu^+ K_L^0$	> 44	90%	326
$N \rightarrow \nu K$	> 86 (n), > 100 (p)	90%	339
$p \rightarrow e^+ K^*(892)^0$	> 82	90%	45
$N \rightarrow \nu K^*(892)$	> 22 (n), > 20 (p)	90%	45
Antilepton + mesons			
$p \rightarrow e^+ \pi^+ \pi^-$	> 21	90%	448
$p \rightarrow e^+ \pi^0 \pi^0$	> 38	90%	449
$n \rightarrow e^+ \pi^- \pi^0$	> 32	90%	449
$p \rightarrow \mu^+ \pi^+ \pi^-$	> 17	90%	425
$p \rightarrow \mu^+ \pi^0 \pi^0$	> 33	90%	427
$n \rightarrow \mu^+ \pi^- \pi^0$	> 33	90%	427
$n \rightarrow e^+ K^0 \pi^-$	> 18	90%	319
Lepton + meson			
$n \rightarrow e^- \pi^+$	> 65	90%	459
$n \rightarrow \mu^- \pi^+$	> 49	90%	453
$n \rightarrow e^- \rho^+$	> 62	90%	154
$n \rightarrow \mu^- \rho^+$	> 7	90%	120
$n \rightarrow e^- K^+$	> 32	90%	340
$n \rightarrow \mu^- K^+$	> 57	90%	330
Lepton + mesons			
$p \rightarrow e^- \pi^+ \pi^+$	> 30	90%	448
$n \rightarrow e^- \pi^+ \pi^0$	> 29	90%	449
$p \rightarrow \mu^- \pi^+ \pi^+$	> 17	90%	425
$n \rightarrow \mu^- \pi^+ \pi^0$	> 34	90%	427
$p \rightarrow e^- \pi^+ K^+$	> 20	90%	320
$p \rightarrow \mu^- \pi^+ K^+$	> 5	90%	279

Antilepton + photon(s)

$p \rightarrow e^+ \gamma$	> 460	90%	469
$p \rightarrow \mu^+ \gamma$	> 380	90%	463
$n \rightarrow \nu \gamma$	> 24	90%	470
$p \rightarrow e^+ \gamma \gamma$	> 100	90%	469

Three leptons

$p \rightarrow e^+ e^+ e^-$	> 510	90%	469
$p \rightarrow e^+ \mu^+ \mu^-$	> 81	90%	457
$p \rightarrow e^+ \nu \nu$	> 11	90%	469
$n \rightarrow e^+ e^- \nu$	> 74	90%	470
$n \rightarrow \mu^+ e^- \nu$	> 47	90%	464
$n \rightarrow \mu^+ \mu^- \nu$	> 42	90%	458
$p \rightarrow \mu^+ e^+ e^-$	> 91	90%	464
$p \rightarrow \mu^+ \mu^+ \mu^-$	> 190	90%	439
$p \rightarrow \mu^+ \nu \nu$	> 21	90%	463
$p \rightarrow e^- \mu^+ \mu^+$	> 6	90%	457
$n \rightarrow 3\nu$	> 0.0005	90%	470

Inclusive modes

$N \rightarrow e^+$ anything	> 0.6 (n, p)	90%	-
$N \rightarrow \mu^+$ anything	> 1.2 (n, p)	90%	-
$N \rightarrow e^+ \pi^0$ anything	> 0.6 (n, p)	90%	-

$\Delta B = 2$ dinucleon modes

The following are lifetime limits per iron nucleus.

$p p \rightarrow \pi^+ \pi^+$	> 0.7	90%	-
$p n \rightarrow \pi^+ \pi^0$	> 2	90%	-
$n n \rightarrow \pi^+ \pi^-$	> 0.7	90%	-
$n n \rightarrow \pi^0 \pi^0$	> 3.4	90%	-
$p p \rightarrow e^+ e^+$	> 5.8	90%	-
$p p \rightarrow e^+ \mu^+$	> 3.6	90%	-
$p p \rightarrow \mu^+ \mu^+$	> 1.7	90%	-
$p n \rightarrow e^+ \bar{\nu}$	> 2.8	90%	-
$p n \rightarrow \mu^+ \bar{\nu}$	> 1.6	90%	-
$n n \rightarrow \nu_e \bar{\nu}_e$	> 0.000012	90%	-
$n n \rightarrow \nu_\mu \bar{\nu}_\mu$	> 0.000006	90%	-

\bar{p} DECAY MODES

\bar{p} DECAY MODES	Partial mean life (years)	Confidence level	$\frac{p}{c}$ (MeV/c)
$\bar{p} \rightarrow e^- \gamma$	> 1848	95%	469
$\bar{p} \rightarrow e^- \pi^0$	> 554	95%	459
$\bar{p} \rightarrow e^- \eta$	> 171	95%	309
$\bar{p} \rightarrow e^- K_S^0$	> 29	95%	337
$\bar{p} \rightarrow e^- K_L^0$	> 9	95%	337

n

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 939.56563 \pm 0.00028$ MeV [a]
 $= 1.008664904 \pm 0.000000014$ u
 $m_n - m_p = 1.293318 \pm 0.000009$ MeV
 $= 0.001388434 \pm 0.000000009$ u
Mean life $\tau = 887.0 \pm 2.0$ s ($S = 1.3$)
 $c\tau = 2.659 \times 10^8$ km
Magnetic moment $\mu = -1.9130428 \pm 0.0000005 \mu_N$
Electric dipole moment $d < 11 \times 10^{-26}$ e cm, CL = 95%
Electric polarizability $\alpha = (1.16^{+0.19}_{-0.23}) \times 10^{-3}$ fm³
Charge $q = (-0.4 \pm 1.1) \times 10^{-21}$ e
Mean time for $n\bar{n}$ oscillations $> 1.2 \times 10^8$ s, CL = 90% [d]

Decay parameters [e]

$p e^- \bar{\nu}_e$	$g_A/g_V = -1.2573 \pm 0.0028$
"	$A = -0.1127 \pm 0.0011$
"	$B = 0.997 \pm 0.028$
"	$a = -0.102 \pm 0.005$
"	$\phi_{AV} = (180.07 \pm 0.18)^\circ$ [f]
"	$D = (-0.5 \pm 1.4) \times 10^{-3}$

n DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	$\frac{p}{c}$ (MeV/c)
$p e^- \bar{\nu}_e$	100 %		1.19
Charge conservation (Q) violating mode			
$p \nu_e \bar{\nu}_e$	$Q < 9 \times 10^{-24}$	90%	1.29

Baryon Summary Table

 $N(1440) P_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 1430$ to 1470 (≈ 1440) MeV
 Full width $\Gamma = 250$ to 450 (≈ 350) MeV
 $p_{\text{beam}} = 0.61$ GeV/c $4\pi\chi^2 = 31.0$ mb

$N(1440)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	60–70 %	397
$N\pi\pi$	30–40 %	342
$\Delta\pi$	20–30 %	143
$N\rho$	<8 %	†
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	5–10 %	–
$p\gamma$	0.04–0.07 %	414
$n\gamma$	0.001–0.05 %	413

 $N(1520) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Mass $m = 1515$ to 1530 (≈ 1520) MeV
 Full width $\Gamma = 110$ to 135 (≈ 120) MeV
 $p_{\text{beam}} = 0.74$ GeV/c $4\pi\chi^2 = 23.5$ mb

$N(1520)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	50–60 %	456
$N\pi\pi$	40–50 %	410
$\Delta\pi$	15–25 %	228
$N\rho$	15–25 %	†
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	<8 %	–
$p\gamma$	0.45–0.53 %	470
$n\gamma$	0.34–0.48 %	470

 $N(1535) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Mass $m = 1520$ to 1555 (≈ 1535) MeV
 Full width $\Gamma = 100$ to 250 (≈ 150) MeV
 $p_{\text{beam}} = 0.76$ GeV/c $4\pi\chi^2 = 22.5$ mb

$N(1535)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	35–55 %	467
$N\eta$	30–55 %	182
$N\pi\pi$	1–10 %	422
$\Delta\pi$	<1 %	242
$N\rho$	<4 %	†
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	<3 %	–
$N(1440)\pi$	<7 %	†
$p\gamma$	0.45–0.53 %	481
$n\gamma$	0.34–0.48 %	480

 $N(1650) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-)$$

Mass $m = 1640$ to 1680 (≈ 1650) MeV
 Full width $\Gamma = 145$ to 190 (≈ 150) MeV
 $p_{\text{beam}} = 0.96$ GeV/c $4\pi\chi^2 = 16.4$ mb

$N(1650)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	60–80 %	547
ΛK	3–11 %	161
$N\pi\pi$	5–20 %	511
$\Delta\pi$	3–7 %	344
$N\rho$	4–14 %	†
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	<4 %	–
$N(1440)\pi$	<5 %	147
$p\gamma$	0.10–0.18 %	558
$n\gamma$	0.03–0.18 %	557

 $N(1675) D_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^-)$$

Mass $m = 1670$ to 1685 (≈ 1675) MeV
 Full width $\Gamma = 140$ to 180 (≈ 150) MeV
 $p_{\text{beam}} = 1.01$ GeV/c $4\pi\chi^2 = 15.4$ mb

$N(1675)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	40–50 %	563
ΛK	<1 %	209
$N\pi\pi$	50–60 %	529
$\Delta\pi$	50–60 %	364
$N\rho$	<1–3 %	†
$p\gamma$	0.005–0.014 %	575
$n\gamma$	0.07–0.11 %	574

 $N(1680) F_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^+)$$

Mass $m = 1675$ to 1690 (≈ 1680) MeV
 Full width $\Gamma = 120$ to 140 (≈ 130) MeV
 $p_{\text{beam}} = 1.01$ GeV/c $4\pi\chi^2 = 15.2$ mb

$N(1680)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	60–70 %	567
$N\pi\pi$	30–40 %	532
$\Delta\pi$	5–15 %	369
$N\rho$	3–15 %	†
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	5–20 %	–
$p\gamma$	0.21–0.35 %	578
$n\gamma$	0.02–0.04 %	577

 $N(1700) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

Mass $m = 1650$ to 1750 (≈ 1700) MeV
 Full width $\Gamma = 50$ to 150 (≈ 100) MeV
 $p_{\text{beam}} = 1.05$ GeV/c $4\pi\chi^2 = 14.5$ mb

$N(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	5–15 %	580
ΛK	<3 %	250
$N\pi\pi$	85–95 %	547
$N\rho$	<35 %	†
$p\gamma$	~ 0.01 %	591

 $N(1710) P_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 1680$ to 1740 (≈ 1710) MeV
 Full width $\Gamma = 50$ to 250 (≈ 100) MeV
 $p_{\text{beam}} = 1.07$ GeV/c $4\pi\chi^2 = 14.2$ mb

$N(1710)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–20 %	587
ΛK	5–25 %	264
$N\pi\pi$	40–90 %	554
$\Delta\pi$	15–40 %	393
$N\rho$	5–25 %	48
$N(\pi\pi)_{S\text{-wave}}^{I=0}$	10–40 %	–

 $N(1720) P_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

Mass $m = 1650$ to 1750 (≈ 1720) MeV
 Full width $\Gamma = 100$ to 200 (≈ 150) MeV
 $p_{\text{beam}} = 1.09$ GeV/c $4\pi\chi^2 = 13.9$ mb

$N(1720)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–20 %	594
ΛK	1–15 %	278
$N\pi\pi$	>70 %	561
$N\rho$	70–85 %	104
$p\gamma$	0.01–0.06 %	–

Baryon Summary Table

 $N(2190) G_{17}$

$$I(J^P) = \frac{1}{2}(\frac{7}{2}^-)$$

Mass $m = 2100$ to 2200 (≈ 2190) MeV
 Full width $\Gamma = 350$ to 550 (≈ 450) MeV
 $p_{\text{beam}} = 2.07$ GeV/c $4\pi\chi^2 = 6.21$ mb

$N(2190)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–20 %	888

 $N(2220) H_{19}$

$$I(J^P) = \frac{1}{2}(\frac{9}{2}^+)$$

Mass $m = 2180$ to 2310 (≈ 2220) MeV
 Full width $\Gamma = 320$ to 550 (≈ 400) MeV
 $p_{\text{beam}} = 2.14$ GeV/c $4\pi\chi^2 = 5.97$ mb

$N(2220)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–20 %	905

 $N(2250) G_{19}$

$$I(J^P) = \frac{1}{2}(\frac{9}{2}^-)$$

Mass $m = 2170$ to 2310 (≈ 2250) MeV
 Full width $\Gamma = 290$ to 470 (≈ 400) MeV
 $p_{\text{beam}} = 2.21$ GeV/c $4\pi\chi^2 = 5.74$ mb

$N(2250)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	5–15 %	923

 $N(2600) h_{1,11}$

$$I(J^P) = \frac{1}{2}(\frac{11}{2}^-)$$

Mass $m = 2550$ to 2750 (≈ 2600) MeV
 Full width $\Gamma = 500$ to 800 (≈ 650) MeV
 $p_{\text{beam}} = 3.12$ GeV/c $4\pi\chi^2 = 3.86$ mb

$N(2600)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	5–10 %	1126

Δ BARYONS ($S = 0, I = 3/2$)

$$\Delta^{++} = uuu, \quad \Delta^+ = uud, \quad \Delta^0 = udd, \quad \Delta^- = ddd$$

 $\Delta(1232) P_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$$

Mass $m = 1230$ to 1234 (≈ 1232) MeV
 Full width $\Gamma = 115$ to 125 (≈ 120) MeV
 $p_{\text{beam}} = 0.30$ GeV/c $4\pi\chi^2 = 94.8$ mb

$\Delta(1232)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	>99 %	227
$N\gamma$	0.55–0.61 %	259

 $\Delta(1600) P_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$$

Mass $m = 1550$ to 1700 (≈ 1600) MeV
 Full width $\Gamma = 250$ to 450 (≈ 350) MeV
 $p_{\text{beam}} = 0.87$ GeV/c $4\pi\chi^2 = 18.6$ mb

$\Delta(1600)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–25 %	512
$N\pi\pi$	75–90 %	473
$\Delta\pi$	40–70 %	301
$N\rho$	<25 %	†
$N(1440)\pi$	10–35 %	74
$p\gamma$	~0 %	–

 $\Delta(1620) S_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$$

Mass $m = 1615$ to 1675 (≈ 1620) MeV
 Full width $\Gamma = 120$ to 180 (≈ 150) MeV
 $p_{\text{beam}} = 0.91$ GeV/c $4\pi\chi^2 = 17.7$ mb

$\Delta(1620)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	20–30 %	526
$N\pi\pi$	70–80 %	488
$\Delta\pi$	30–60 %	318
$N\rho$	7–25 %	†
$N\gamma$	0.02–0.06 %	538

 $\Delta(1700) D_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^-)$$

Mass $m = 1670$ to 1770 (≈ 1700) MeV
 Full width $\Gamma = 200$ to 400 (≈ 300) MeV
 $p_{\text{beam}} = 1.05$ GeV/c $4\pi\chi^2 = 14.5$ mb

$\Delta(1700)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–20 %	580
$N\pi\pi$	80–90 %	547
$\Delta\pi$	30–60 %	385
$N\rho$	30–55 %	†
$N\gamma$	0.16–0.28 %	591

 $\Delta(1900) S_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$$

Mass $m = 1850$ to 1950 (≈ 1900) MeV
 Full width $\Gamma = 140$ to 240 (≈ 200) MeV
 $p_{\text{beam}} = 1.44$ GeV/c $4\pi\chi^2 = 9.71$ mb

$\Delta(1900)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–30 %	710

 $\Delta(1905) F_{35}$

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^+)$$

Mass $m = 1870$ to 1920 (≈ 1905) MeV
 Full width $\Gamma = 280$ to 440 (≈ 350) MeV
 $p_{\text{beam}} = 1.46$ GeV/c $4\pi\chi^2 = 9.62$ mb

$\Delta(1905)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	5–15 %	713
$N\pi\pi$	85–95 %	687
$\Delta\pi$	<25 %	542
$N\rho$	>60 %	421
$N\gamma$	0.01–0.04 %	721

 $\Delta(1910) P_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$$

Mass $m = 1870$ to 1920 (≈ 1910) MeV
 Full width $\Gamma = 190$ to 270 (≈ 250) MeV
 $p_{\text{beam}} = 1.46$ GeV/c $4\pi\chi^2 = 9.54$ mb

$\Delta(1910)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	15–30 %	716

 $\Delta(1920) P_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$$

Mass $m = 1900$ to 1970 (≈ 1920) MeV
 Full width $\Gamma = 150$ to 300 (≈ 200) MeV
 $p_{\text{beam}} = 1.48$ GeV/c $4\pi\chi^2 = 9.37$ mb

$\Delta(1920)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	5–20 %	722

Baryon Summary Table

$\Delta(1930) D_{35}$	$I(J^P) = \frac{3}{2}(\frac{5}{2}^-)$
Mass $m = 1920$ to 1970 (≈ 1930) MeV	
Full width $\Gamma = 250$ to 450 (≈ 350) MeV	
$\rho_{\text{beam}} = 1.50$ GeV/c	$4\pi\chi^2 = 9.21$ mb

$\Delta(1930)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	10–20 %	729

$\Delta(1950) F_{37}$	$I(J^P) = \frac{3}{2}(\frac{7}{2}^+)$
Mass $m = 1940$ to 1960 (≈ 1950) MeV	
Full width $\Gamma = 290$ to 350 (≈ 300) MeV	
$\rho_{\text{beam}} = 1.54$ GeV/c	$4\pi\chi^2 = 8.91$ mb

$\Delta(1950)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	35–40 %	741
$N\pi\pi$		716
$\Delta\pi$	20–30 %	574
$N\rho$	<10 %	469
$N\gamma$	0.10–0.15 %	749

$\Delta(2420) H_{3,11}$	$I(J^P) = \frac{3}{2}(\frac{11}{2}^+)$
Mass $m = 2300$ to 2500 (≈ 2420) MeV	
Full width $\Gamma = 300$ to 500 (≈ 400) MeV	
$\rho_{\text{beam}} = 2.64$ GeV/c	$4\pi\chi^2 = 4.68$ mb

$\Delta(2420)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\pi$	5–15 %	1023

Λ BARYONS

$(S = -1, I = 0)$

$$\Lambda^0 = uds$$

Λ	$I(J^P) = 0(\frac{1}{2}^+)$
Mass $m = 1115.684 \pm 0.006$ MeV	
Mean life $\tau = (2.632 \pm 0.020) \times 10^{-10}$ s ($S = 1.6$)	
$c\tau = 7.89$ cm	
Magnetic moment $\mu = -0.613 \pm 0.004 \mu_N$	
Electric dipole moment $d < 1.5 \times 10^{-16}$ ecm, CL = 95%	

Decay parameters

$\rho\pi^-$	$\alpha_- = 0.642 \pm 0.013$
"	$\phi_- = (-6.5 \pm 3.5)^\circ$
"	$\gamma_- = 0.76$ [g]
"	$\Delta_- = (8 \pm 4)^\circ$ [g]
$n\pi^0$	$\alpha_0 = +0.65 \pm 0.05$
$\rho e^- \bar{\nu}_e$	$g_A/g_V = -0.718 \pm 0.015$ [e]

Λ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\rho\pi^-$	(63.9 \pm 0.5) %	101
$n\pi^0$	(35.8 \pm 0.5) %	104
$n\gamma$	(1.75 \pm 0.15) $\times 10^{-3}$	162
$\rho\pi^- \gamma$	[h] (8.4 \pm 1.4) $\times 10^{-4}$	101
$\rho e^- \bar{\nu}_e$	(8.32 \pm 0.14) $\times 10^{-4}$	163
$\rho\mu^- \bar{\nu}_\mu$	(1.57 \pm 0.35) $\times 10^{-4}$	131

$\Lambda(1405) S_{01}$	$I(J^P) = 0(\frac{1}{2}^-)$
Mass $m = 1407 \pm 4$ MeV	
Full width $\Gamma = 50.0 \pm 2.0$ MeV	
Below $\bar{K}N$ threshold	

$\Lambda(1405)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\Sigma\pi$	100 %	152

$\Lambda(1520) D_{03}$	$I(J^P) = 0(\frac{3}{2}^-)$
Mass $m = 1519.5 \pm 1.0$ MeV [1]	
Full width $\Gamma = 15.6 \pm 1.0$ MeV [1]	
$\rho_{\text{beam}} = 0.39$ GeV/c	$4\pi\chi^2 = 82.8$ mb

$\Lambda(1520)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	45 \pm 1%	244
$\Sigma\pi$	42 \pm 1%	267
$\Lambda\pi\pi$	10 \pm 1%	252
$\Sigma\pi\pi$	0.9 \pm 0.1%	152
$\Lambda\gamma$	0.8 \pm 0.2%	351

$\Lambda(1600) P_{01}$	$I(J^P) = 0(\frac{1}{2}^+)$
Mass $m = 1560$ to 1700 (≈ 1600) MeV	
Full width $\Gamma = 50$ to 250 (≈ 150) MeV	
$\rho_{\text{beam}} = 0.58$ GeV/c	$4\pi\chi^2 = 41.6$ mb

$\Lambda(1600)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	15–30 %	343
$\Sigma\pi$	10–60 %	336

$\Lambda(1670) S_{01}$	$I(J^P) = 0(\frac{1}{2}^-)$
Mass $m = 1660$ to 1680 (≈ 1670) MeV	
Full width $\Gamma = 25$ to 50 (≈ 35) MeV	
$\rho_{\text{beam}} = 0.74$ GeV/c	$4\pi\chi^2 = 28.5$ mb

$\Lambda(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	15–25 %	414
$\Sigma\pi$	20–60 %	393
$\Lambda\eta$	15–35 %	64

$\Lambda(1690) D_{03}$	$I(J^P) = 0(\frac{3}{2}^-)$
Mass $m = 1685$ to 1695 (≈ 1690) MeV	
Full width $\Gamma = 50$ to 70 (≈ 60) MeV	
$\rho_{\text{beam}} = 0.78$ GeV/c	$4\pi\chi^2 = 26.1$ mb

$\Lambda(1690)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	20–30 %	433
$\Sigma\pi$	20–40 %	409
$\Lambda\pi\pi$	~ 25 %	415
$\Sigma\pi\pi$	~ 20 %	350

$\Lambda(1800) S_{01}$	$I(J^P) = 0(\frac{1}{2}^-)$
Mass $m = 1720$ to 1850 (≈ 1800) MeV	
Full width $\Gamma = 200$ to 400 (≈ 300) MeV	
$\rho_{\text{beam}} = 1.01$ GeV/c	$4\pi\chi^2 = 17.5$ mb

$\Lambda(1800)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	25–40 %	528
$\Sigma\pi$	seen	493
$\Sigma(1385)\pi$	seen	345
$N\bar{K}^*(892)$	seen	†

$\Lambda(1810) P_{01}$	$I(J^P) = 0(\frac{1}{2}^+)$
Mass $m = 1750$ to 1850 (≈ 1810) MeV	
Full width $\Gamma = 50$ to 250 (≈ 150) MeV	
$\rho_{\text{beam}} = 1.04$ GeV/c	$4\pi\chi^2 = 17.0$ mb

$\Lambda(1810)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	20–50 %	537
$\Sigma\pi$	10–40 %	501
$\Sigma(1385)\pi$	seen	356
$N\bar{K}^*(892)$	30–60 %	†

Baryon Summary Table

 $\Lambda(1820) F_{05}$

$I(J^P) = 0(\frac{5}{2}^+)$

Mass $m = 1815$ to 1825 (≈ 1820) MeV
 Full width $\Gamma = 70$ to 90 (≈ 80) MeV
 $p_{\text{beam}} = 1.06$ GeV/c $4\pi\lambda^2 = 16.5$ mb

$\Lambda(1820)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	55–65 %	545
$\Sigma\pi$	8–14 %	508
$\Sigma(1385)\pi$	5–10 %	362

 $\Lambda(1830) D_{05}$

$I(J^P) = 0(\frac{5}{2}^-)$

Mass $m = 1810$ to 1830 (≈ 1830) MeV
 Full width $\Gamma = 60$ to 110 (≈ 95) MeV
 $p_{\text{beam}} = 1.08$ GeV/c $4\pi\lambda^2 = 16.0$ mb

$\Lambda(1830)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	3–10 %	553
$\Sigma\pi$	35–75 %	515
$\Sigma(1385)\pi$	>15 %	371

 $\Lambda(1890) P_{03}$

$I(J^P) = 0(\frac{3}{2}^+)$

Mass $m = 1850$ to 1910 (≈ 1890) MeV
 Full width $\Gamma = 60$ to 200 (≈ 100) MeV
 $p_{\text{beam}} = 1.21$ GeV/c $4\pi\lambda^2 = 13.6$ mb

$\Lambda(1890)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	20–35 %	599
$\Sigma\pi$	3–10 %	559
$\Sigma(1385)\pi$	seen	420
$N\bar{K}^*(892)$	seen	233

 $\Lambda(2100) G_{07}$

$I(J^P) = 0(\frac{7}{2}^-)$

Mass $m = 2090$ to 2110 (≈ 2100) MeV
 Full width $\Gamma = 100$ to 250 (≈ 200) MeV
 $p_{\text{beam}} = 1.68$ GeV/c $4\pi\lambda^2 = 8.68$ mb

$\Lambda(2100)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	25–35 %	751
$\Sigma\pi$	~ 5 %	704
$\Lambda\eta$	<3 %	617
ΞK	<3 %	483
$\Lambda\omega$	<8 %	443
$N\bar{K}^*(892)$	10–20 %	514

 $\Lambda(2110) F_{05}$

$I(J^P) = 0(\frac{5}{2}^+)$

Mass $m = 2090$ to 2140 (≈ 2110) MeV
 Full width $\Gamma = 150$ to 250 (≈ 200) MeV
 $p_{\text{beam}} = 1.70$ GeV/c $4\pi\lambda^2 = 8.53$ mb

$\Lambda(2110)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	5–25 %	757
$\Sigma\pi$	10–40 %	711
$\Lambda\omega$	seen	455
$\Sigma(1385)\pi$	seen	589
$N\bar{K}^*(892)$	10–60 %	524

 $\Lambda(2350) H_{09}$

$I(J^P) = 0(\frac{9}{2}^+)$

Mass $m = 2340$ to 2370 (≈ 2350) MeV
 Full width $\Gamma = 100$ to 250 (≈ 150) MeV
 $p_{\text{beam}} = 2.29$ GeV/c $4\pi\lambda^2 = 5.85$ mb

$\Lambda(2350)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	~ 12 %	915
$\Sigma\pi$	~ 10 %	867

 Σ BARYONS
($S = -1, I = 1$)

$\Sigma^+ = uus, \Sigma^0 = uds, \Sigma^- = dds$

 Σ^+

$I(J^P) = 1(\frac{1}{2}^+)$

Mass $m = 1189.37 \pm 0.07$ MeV ($S = 2.2$)
 Mean life $\tau = (0.799 \pm 0.004) \times 10^{-10}$ s
 $c\tau = 2.396$ cm

Magnetic moment $\mu = 2.458 \pm 0.010 \mu_N$ ($S = 2.1$)
 $\Gamma(\Sigma^+ \rightarrow n\ell^+\nu)/\Gamma(\Sigma^- \rightarrow n\ell^-\bar{\nu}) < 0.043$

Decay parameters

$\rho\pi^0$	$\alpha_0 = -0.980^{+0.017}_{-0.015}$
"	$\phi_0 = (36 \pm 34)^\circ$
"	$\gamma_0 = 0.16$ [g]
"	$\Delta_0 = (187 \pm 6)^\circ$ [g]
$n\pi^+$	$\alpha_+ = 0.068 \pm 0.013$
"	$\phi_+ = (167 \pm 20)^\circ$ ($S = 1.1$)
"	$\gamma_+ = -0.97$ [g]
"	$\Delta_+ = (-73^{+133}_{-10})^\circ$ [g]
$\rho\gamma$	$\alpha_\gamma = -0.76 \pm 0.08$

Σ^+ DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\rho\pi^0$	(51.57 ± 0.30) %		189
$n\pi^+$	(48.30 ± 0.30) %		185
$\rho\gamma$	$(1.25 \pm 0.07) \times 10^{-3}$		225
$n\pi^+\gamma$	[h] $(4.5 \pm 0.5) \times 10^{-4}$		185
$\Lambda e^+\nu_e$	$(2.0 \pm 0.5) \times 10^{-5}$		71

 $\Delta S = \Delta Q$ (SQ) violating modes or
 $\Delta S = 1$ weak neutral current (SI) modes

$ne^+\nu_e$	SQ	< 5	$\times 10^{-6}$	90%	224
$n\mu^+\nu_\mu$	SQ	< 3.0	$\times 10^{-5}$	90%	202
ρe^+e^-	SI	< 7	$\times 10^{-6}$		225

 Σ^0

$I(J^P) = 1(\frac{1}{2}^+)$

J^P not measured; assumed to be the same as for the Σ^+ and Σ^- .

Mass $m = 1192.55 \pm 0.08$ MeV ($S = 1.2$)
 $m_{\Sigma^-} - m_{\Sigma^0} = 4.88 \pm 0.08$ MeV ($S = 1.2$)
 $m_{\Sigma^0} - m_\Lambda = 76.87 \pm 0.08$ MeV ($S = 1.2$)
 Mean life $\tau = (7.4 \pm 0.7) \times 10^{-20}$ s
 $c\tau = 2.22 \times 10^{-11}$ m
 Transition magnetic moment $|\mu_{\Sigma\Lambda}| = 1.61 \pm 0.08 \mu_N$

Σ^0 DECAY MODES	Fraction (Γ_i/Γ)	Confidence level	ρ (MeV/c)
$\Lambda\gamma$	100 %		74
$\Lambda\gamma\gamma$	< 3 %	90%	74
Λe^+e^-	[j] 5×10^{-3}		74

 Σ^-

$I(J^P) = 1(\frac{1}{2}^+)$

Mass $m = 1197.436 \pm 0.033$ MeV ($S = 1.2$)
 $m_{\Sigma^-} - m_{\Sigma^+} = 8.07 \pm 0.08$ MeV ($S = 1.9$)
 $m_{\Sigma^-} - m_\Lambda = 81.752 \pm 0.034$ MeV ($S = 1.2$)
 Mean life $\tau = (1.479 \pm 0.011) \times 10^{-10}$ s ($S = 1.3$)
 $c\tau = 4.434$ cm
 Magnetic moment $\mu = -1.160 \pm 0.025 \mu_N$ ($S = 1.7$)

Decay parameters

$n\pi^-$	$\alpha_- = -0.068 \pm 0.008$
"	$\phi_- = (10 \pm 15)^\circ$
"	$\gamma_- = 0.98$ [g]
"	$\Delta_- = (249^{+120}_{-120})^\circ$ [g]
$ne^-\bar{\nu}_e$	$g_A/g_V = 0.340 \pm 0.017$ [e]
"	$f_2(0)/f_1(0) = 0.97 \pm 0.14$
"	$D = 0.11 \pm 0.10$
$\Lambda e^-\bar{\nu}_e$	$g_V/g_A = 0.01 \pm 0.10$ [e] ($S = 1.5$)
"	$g_{WM}/g_A = 2.4 \pm 1.7$ [e]

Baryon Summary Table

Σ^- DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$n\pi^-$	$(99.848 \pm 0.005) \%$	193
$n\pi^- \gamma$	$[h] (4.6 \pm 0.6) \times 10^{-4}$	193
$ne^- \bar{\nu}_e$	$(1.017 \pm 0.034) \times 10^{-3}$	230
$n\mu^- \bar{\nu}_\mu$	$(4.5 \pm 0.4) \times 10^{-4}$	210
$\Lambda e^- \bar{\nu}_e$	$(5.73 \pm 0.27) \times 10^{-5}$	79

$\Sigma(1385) P_{13}$	$I(J^P) = 1(\frac{3}{2}^+)$
$\Sigma(1385)^+$ mass $m = 1382.8 \pm 0.4$ MeV	(S = 2.0)
$\Sigma(1385)^0$ mass $m = 1383.7 \pm 1.0$ MeV	(S = 1.4)
$\Sigma(1385)^-$ mass $m = 1387.2 \pm 0.5$ MeV	(S = 2.2)
$\Sigma(1385)^+$ full width $\Gamma = 35.8 \pm 0.8$ MeV	
$\Sigma(1385)^0$ full width $\Gamma = 36 \pm 5$ MeV	
$\Sigma(1385)^-$ full width $\Gamma = 39.4 \pm 2.1$ MeV	(S = 1.7)
Below $\bar{K}N$ threshold	

$\Sigma(1385)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$\Lambda\pi$	$88 \pm 2 \%$	208
$\Sigma\pi$	$12 \pm 2 \%$	127

$\Sigma(1660) P_{11}$	$I(J^P) = 1(\frac{1}{2}^+)$
Mass $m = 1630$ to 1690 (≈ 1660) MeV	
Full width $\Gamma = 40$ to 200 (≈ 100) MeV	
$\rho_{\text{beam}} = 0.72$ GeV/c	$4\pi\chi^2 = 29.9$ mb

$\Sigma(1660)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	10 – 30%	405
$\Lambda\pi$	seen	439
$\Sigma\pi$	seen	385

$\Sigma(1670) D_{13}$	$I(J^P) = 1(\frac{3}{2}^-)$
Mass $m = 1665$ to 1685 (≈ 1670) MeV	
Full width $\Gamma = 40$ to 80 (≈ 60) MeV	
$\rho_{\text{beam}} = 0.74$ GeV/c	$4\pi\chi^2 = 28.5$ mb

$\Sigma(1670)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	7 – 13%	414
$\Lambda\pi$	5 – 15%	447
$\Sigma\pi$	30 – 60%	393

$\Sigma(1750) S_{11}$	$I(J^P) = 1(\frac{1}{2}^-)$
Mass $m = 1730$ to 1800 (≈ 1750) MeV	
Full width $\Gamma = 60$ to 160 (≈ 90) MeV	
$\rho_{\text{beam}} = 0.91$ GeV/c	$4\pi\chi^2 = 20.7$ mb

$\Sigma(1750)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	10 – 40%	486
$\Lambda\pi$	seen	507
$\Sigma\pi$	$< 8 \%$	455
$\Sigma\eta$	15 – 55%	81

$\Sigma(1775) D_{15}$	$I(J^P) = 1(\frac{5}{2}^-)$
Mass $m = 1770$ to 1780 (≈ 1775) MeV	
Full width $\Gamma = 105$ to 135 (≈ 120) MeV	
$\rho_{\text{beam}} = 0.96$ GeV/c	$4\pi\chi^2 = 19.0$ mb

$\Sigma(1775)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	37 – 43%	508
$\Lambda\pi$	14 – 20%	525
$\Sigma\pi$	2 – 5%	474
$\Sigma(1385)\pi$	8 – 12%	324
$\Lambda(1520)\pi$	17 – 23%	198

$\Sigma(1915) F_{15}$	$I(J^P) = 1(\frac{5}{2}^+)$
Mass $m = 1900$ to 1935 (≈ 1915) MeV	
Full width $\Gamma = 80$ to 160 (≈ 120) MeV	
$\rho_{\text{beam}} = 1.26$ GeV/c	$4\pi\chi^2 = 12.8$ mb

$\Sigma(1915)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	5 – 15%	618
$\Lambda\pi$	seen	622
$\Sigma\pi$	seen	577
$\Sigma(1385)\pi$	$< 5 \%$	440

$\Sigma(1940) D_{13}$	$I(J^P) = 1(\frac{3}{2}^-)$
Mass $m = 1900$ to 1950 (≈ 1940) MeV	
Full width $\Gamma = 150$ to 300 (≈ 220) MeV	
$\rho_{\text{beam}} = 1.32$ GeV/c	$4\pi\chi^2 = 12.1$ mb

$\Sigma(1940)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	$< 20 \%$	637
$\Lambda\pi$	seen	639
$\Sigma\pi$	seen	594
$\Sigma(1385)\pi$	seen	460
$\Lambda(1520)\pi$	seen	354
$\Delta(1232)\bar{K}$	seen	410
$N\bar{K}^*(892)$	seen	320

$\Sigma(2030) F_{17}$	$I(J^P) = 1(\frac{7}{2}^+)$
Mass $m = 2025$ to 2040 (≈ 2030) MeV	
Full width $\Gamma = 150$ to 200 (≈ 180) MeV	
$\rho_{\text{beam}} = 1.52$ GeV/c	$4\pi\chi^2 = 9.93$ mb

$\Sigma(2030)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	17 – 23%	702
$\Lambda\pi$	17 – 23%	700
$\Sigma\pi$	5 – 10%	657
ΞK	$< 2 \%$	412
$\Sigma(1385)\pi$	5 – 15%	529
$\Lambda(1520)\pi$	10 – 20%	430
$\Delta(1232)\bar{K}$	10 – 20%	498
$N\bar{K}^*(892)$	$< 5 \%$	438

$\Sigma(2250)$	$I(J^P) = 1(?^?)$
Mass $m = 2210$ to 2280 (≈ 2250) MeV	
Full width $\Gamma = 60$ to 150 (≈ 100) MeV	
$\rho_{\text{beam}} = 2.04$ GeV/c	$4\pi\chi^2 = 6.76$ mb

$\Sigma(2250)$ DECAY MODES	Fraction (Γ_i/Γ)	ρ (MeV/c)
$N\bar{K}$	$< 10 \%$	851
$\Lambda\pi$	seen	842
$\Sigma\pi$	seen	803

Baryon Summary Table

Ξ BARYONS (S = -2, I = 1/2)

$$\Xi^0 = uss, \quad \Xi^- = dss$$

Ξ⁰

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

P is not yet measured; + is the quark model prediction.

$$\text{Mass } m = 1314.9 \pm 0.6 \text{ MeV}$$

$$m_{\Xi^-} - m_{\Xi^0} = 6.4 \pm 0.6 \text{ MeV}$$

$$\text{Mean life } \tau = (2.90 \pm 0.09) \times 10^{-10} \text{ s}$$

$$c\tau = 8.71 \text{ cm}$$

$$\text{Magnetic moment } \mu = -1.250 \pm 0.014 \mu_N$$

Decay parameters

$$\Lambda\pi^0 \quad \alpha = -0.411 \pm 0.022 \quad (S = 2.1)$$

$$" \quad \phi = (21 \pm 12)^\circ$$

$$" \quad \gamma = 0.85 [g]$$

$$" \quad \Delta = (218_{-19}^{+12})^\circ [g]$$

$$\Lambda\gamma \quad \alpha = 0.4 \pm 0.4$$

$$\Sigma^0\gamma \quad \alpha = 0.20 \pm 0.32$$

Ξ ⁰ DECAY MODES	Fraction (Γ _i /Γ)	Confidence level	ρ (MeV/c)
Λπ ⁰	(99.54 ± 0.05) %		135
Λγ	(1.06 ± 0.16) × 10 ⁻³		184
Σ ⁰ γ	(3.5 ± 0.4) × 10 ⁻³		117
Σ ⁺ e ⁻ ν _e	< 1.1 × 10 ⁻³	90%	120
Σ ⁺ μ ⁻ ν _μ	< 1.1 × 10 ⁻³	90%	64

ΔS = ΔQ (S_Q) violating modes or ΔS = 2 forbidden (S₂) modes

Σ ⁻ e ⁺ ν _e	S _Q < 9 × 10 ⁻⁴	90%	112
Σ ⁻ μ ⁺ ν _μ	S _Q < 9 × 10 ⁻⁴	90%	49
ρπ ⁻	S ₂ < 4 × 10 ⁻⁵	90%	299
ρe ⁻ ν _e	S ₂ < 1.3 × 10 ⁻³		323
ρμ ⁻ ν _μ	S ₂ < 1.3 × 10 ⁻³		309

Ξ⁻

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

P is not yet measured; + is the quark model prediction.

$$\text{Mass } m = 1321.32 \pm 0.13 \text{ MeV}$$

$$\text{Mean life } \tau = (1.639 \pm 0.015) \times 10^{-10} \text{ s}$$

$$c\tau = 4.91 \text{ cm}$$

$$\text{Magnetic moment } \mu = -0.6507 \pm 0.0025 \mu_N$$

Decay parameters

$$\Lambda\pi^- \quad \alpha = -0.456 \pm 0.014 \quad (S = 1.8)$$

$$" \quad \phi = (4 \pm 4)^\circ$$

$$" \quad \gamma = 0.89 [g]$$

$$" \quad \Delta = (188 \pm 8)^\circ [g]$$

$$\Lambda e^- \bar{\nu}_e \quad g_A/g_V = -0.25 \pm 0.05 [e]$$

Ξ ⁻ DECAY MODES	Fraction (Γ _i /Γ)	Confidence level	ρ (MeV/c)
Λπ ⁻	(99.887 ± 0.035) %		139
Σ ⁻ γ	(1.27 ± 0.23) × 10 ⁻⁴		118
Λe ⁻ ν _e	(5.63 ± 0.31) × 10 ⁻⁴		190
Λμ ⁻ ν _μ	(3.5 $_{-2.2}^{+3.5}$) × 10 ⁻⁴		163
Σ ⁰ e ⁻ ν _e	(8.7 ± 1.7) × 10 ⁻⁵		122
Σ ⁰ μ ⁻ ν _μ	< 8 × 10 ⁻⁴	90%	70
Ξ ⁰ e ⁻ ν _e	< 2.3 × 10 ⁻³	90%	6

ΔS = 2 forbidden (S₂) modes

nπ ⁻	S ₂ < 1.9 × 10 ⁻⁵	90%	303
ne ⁻ ν _e	S ₂ < 3.2 × 10 ⁻³	90%	327
nμ ⁻ ν _μ	S ₂ < 1.5 %	90%	314
ρπ ⁻ π ⁻	S ₂ < 4 × 10 ⁻⁴	90%	223
ρπ ⁻ e ⁻ ν _e	S ₂ < 4 × 10 ⁻⁴	90%	304
ρπ ⁻ μ ⁻ ν _μ	S ₂ < 4 × 10 ⁻⁴	90%	250
ρμ ⁻ μ ⁻	L < 4 × 10 ⁻⁴	90%	-

Ξ(1530) P₁₃

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$$

$$\Xi(1530)^0 \text{ mass } m = 1531.80 \pm 0.32 \text{ MeV} \quad (S = 1.3)$$

$$\Xi(1530)^- \text{ mass } m = 1535.0 \pm 0.6 \text{ MeV}$$

$$\Xi(1530)^0 \text{ full width } \Gamma = 9.1 \pm 0.5 \text{ MeV}$$

$$\Xi(1530)^- \text{ full width } \Gamma = 9.9_{-1.9}^{+1.7} \text{ MeV}$$

Ξ(1530) DECAY MODES	Fraction (Γ _i /Γ)	Confidence level	ρ (MeV/c)
Ξπ	100 %		152
Ξγ	< 4 %	90%	200

Ξ(1690)

$$I(J^P) = \frac{1}{2}(?^?)$$

$$\text{Mass } m = 1690 \pm 10 \text{ MeV} [l]$$

$$\text{Full width } \Gamma < 50 \text{ MeV}$$

Ξ(1690) DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
ΛK̄	seen	240
ΣK̄	seen	51
Ξ ⁻ π ⁺ π ⁻	possibly seen	214

Ξ(1820) D₁₃

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-)$$

$$\text{Mass } m = 1823 \pm 5 \text{ MeV} [l]$$

$$\text{Full width } \Gamma = 24_{-10}^{+15} \text{ MeV} [l]$$

Ξ(1820) DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
ΛK̄	large	400
ΣK̄	small	320
Ξπ	small	413
Ξ(1530)π	small	234

Ξ(1950)

$$I(J^P) = \frac{1}{2}(?^?)$$

$$\text{Mass } m = 1950 \pm 15 \text{ MeV} [l]$$

$$\text{Full width } \Gamma = 60 \pm 20 \text{ MeV} [l]$$

Ξ(1950) DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
ΛK̄	seen	522
ΣK̄	possibly seen	460
Ξπ	seen	518

Ξ(2030)

$$I(J^P) = \frac{1}{2}(\geq \frac{5}{2}^?)$$

$$\text{Mass } m = 2025 \pm 5 \text{ MeV} [l]$$

$$\text{Full width } \Gamma = 20_{-5}^{+15} \text{ MeV} [l]$$

Ξ(2030) DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
ΛK̄	~ 20 %	589
ΣK̄	~ 80 %	533
Ξπ	small	573
Ξ(1530)π	small	421
ΛK̄π	small	501
ΣK̄π	small	430

Baryon Summary Table

Ω BARYONS (S = -3, I = 0)

$$\Omega^- = sss$$

Ω⁻

$$I(J^P) = 0(\frac{3}{2}^+)$$

J^P is not yet measured; $\frac{3}{2}^+$ is the quark model prediction.

Mass $m = 1672.45 \pm 0.29$ MeV

Mean life $\tau = (0.822 \pm 0.012) \times 10^{-10}$ s

$c\tau = 2.46$ cm

Magnetic moment $\mu = -1.94 \pm 0.22 \mu_N$

Decay parameters

ΛK^- $\alpha = -0.026 \pm 0.026$

$\Xi^0 \pi^-$ $\alpha = 0.09 \pm 0.14$

$\Xi^- \pi^0$ $\alpha = 0.05 \pm 0.21$

Ω ⁻ DECAY MODES	Fraction (Γ _i /Γ)	Confidence level	ρ (MeV/c)
ΛK^-	(67.8 ± 0.7) %		211
$\Xi^0 \pi^-$	(23.6 ± 0.7) %		294
$\Xi^- \pi^0$	(8.6 ± 0.4) %		290
$\Xi^- \pi^+ \pi^-$	(4.3 ^{+3.4} _{-1.3}) × 10 ⁻⁴		190
$\Xi(1530)^0 \pi^-$	(6.4 ^{+5.1} _{-2.0}) × 10 ⁻⁴		17
$\Xi^0 e^- \bar{\nu}_e$	(5.6 ± 2.8) × 10 ⁻³		319
$\Xi^- \gamma$	< 2.2 × 10 ⁻³	90%	314
ΔS = 2 forbidden (S2) modes			
$\Lambda \pi^-$	S2 < 1.9 × 10 ⁻⁴	90%	449

Ω(2250)⁻

$$I(J^P) = 0(?^?)$$

Mass $m = 2252 \pm 9$ MeV

Full width $\Gamma = 55 \pm 18$ MeV

Ω(2250) ⁻ DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
$\Xi^- \pi^+ K^-$	seen	531
$\Xi(1530)^0 K^-$	seen	437

CHARMED BARYONS (C = +1)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc,$$

$$\Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

Λ_c⁺

$$I(J^P) = 0(\frac{1}{2}^+)$$

J not confirmed; $\frac{1}{2}$ is the quark model prediction.

Mass $m = 2285.1 \pm 0.6$ MeV

Mean life $\tau = (0.200^{+0.011}_{-0.010}) \times 10^{-12}$ s

$c\tau = 60.0 \mu\text{m}$

Decay asymmetry parameters

$\Lambda \pi^+$ $\alpha = -1.03 \pm 0.29$

$\Lambda e^+ \nu_e$ $\alpha = -0.89^{+0.19}_{-0.12}$

Λ _c ⁺ DECAY MODES	Fraction (Γ _i /Γ)	Scale factor/ Confidence level	ρ (MeV/c)
Hadronic modes with a p and one K			
$p \bar{K}^0$	(2.1 ± 0.4) %		872
$p K^- \pi^+$	(4.4 ± 0.6) %		822
$p \bar{K}^*(892)^0$	[k] (1.6 ± 0.4) %		681
$\Delta(1232)^{++} K^-$	(7 ± 4) × 10 ⁻³		709
$\Lambda(1520) \pi^+$	[k] (3.9 ^{+2.0} _{-1.7}) × 10 ⁻³		626
$p K^- \pi^+$ nonresonant	(2.4 ^{+0.5} _{-0.6}) %		822
$p \bar{K}^0 \pi^+ \pi^-$	(2.4 ± 0.8) %	S=1.3	753

$p K^- \pi^+ \pi^0$	seen	758
$p \bar{K}^*(892)^- \pi^+$	seen	579
$p(K^- \pi^+)^0_{\text{nonresonant}} \pi^0$	(3.2 ± 0.7) %	758
$\Delta(1232) \bar{K}^*(892)$	seen	417
$p K^- \pi^+ \pi^+ \pi^-$	(10 ± 7) × 10 ⁻⁴	670
$p K^- \pi^+ \pi^0 \pi^0$	(7.0 ± 3.5) × 10 ⁻³	676
$p K^- \pi^+ \pi^0 \pi^0 \pi^0$	(4.4 ± 2.8) × 10 ⁻³	573

Hadronic modes with a p and zero or two K's

$p \pi^+ \pi^-$	(3.0 ± 1.6) × 10 ⁻³	926
$p f_0(980)$	[k] (2.4 ± 1.6) × 10 ⁻³	621
$p \pi^+ \pi^+ \pi^- \pi^-$	(1.6 ± 1.0) × 10 ⁻³	851
$p K^+ K^-$	(3.0 ± 1.1) × 10 ⁻³	615
$p \phi$	[k] < 1.7 × 10 ⁻³	CL=90% 589

Hadronic modes with a hyperon

$\Lambda \pi^+$	(7.9 ± 1.8) × 10 ⁻³	863
$\Lambda \pi^+ \pi^0$	(3.2 ± 0.9) %	843
$\Lambda \rho^0$	< 4 %	CL=95% 639
$\Lambda \pi^+ \pi^+ \pi^-$	(2.7 ± 0.6) %	806
$\Sigma^0 \pi^+$	(8.7 ± 2.0) × 10 ⁻³	824
$\Sigma^0 \pi^+ \pi^0$	(1.6 ± 0.6) %	802
$\Sigma^0 \pi^+ \pi^+ \pi^-$	(9.2 ± 3.3) × 10 ⁻³	762
$\Sigma^+ \pi^0$	(8.7 ± 2.2) × 10 ⁻³	826
$\Sigma^+ \pi^+ \pi^-$	(3.0 ± 0.6) %	803
$\Sigma^+ \rho^0$	< 1.2 %	CL=95% 579
$\Sigma^- \pi^+ \pi^+$	(1.6 ± 0.6) %	798
$\Sigma^+ \pi^+ \pi^- \pi^0$		569
$\Sigma^+ \omega$	[k] (2.4 ± 0.7) %	569
$\Sigma^+ \pi^+ \pi^+ \pi^- \pi^-$	(2.6 ^{+3.5} _{-1.8}) × 10 ⁻³	707
$\Sigma^+ K^+ K^-$	(3.1 ± 0.8) × 10 ⁻³	346
$\Sigma^+ \phi$	[k] (3.0 ± 1.3) × 10 ⁻³	292
$\Sigma^+ K^+ \pi^-$	(5.7 ^{+5.3} _{-3.1}) × 10 ⁻³	668
$\Xi^0 K^+$	(3.4 ± 0.9) × 10 ⁻³	652
$\Xi^- K^+ \pi^+$	(3.8 ± 1.2) × 10 ⁻³	564
$\Xi(1530)^0 K^+$	[k] (2.3 ± 0.9) × 10 ⁻³	471

Inclusive modes

p anything	(50 ± 16) %	-
p anything (no Λ)	(12 ± 19) %	-
n anything	(50 ± 16) %	-
n anything (no Λ)	(29 ± 17) %	-
Λ anything	(35 ± 11) %	S=1.4
Σ [±] anything	[l] (10 ± 5) %	-
e ⁺ anything	(4.5 ± 1.7) %	-
ρ e ⁺ anything	(1.8 ± 0.9) %	-
Λ e ⁺ anything	(1.4 ± 0.5) %	-
Λ μ ⁺ anything	(1.5 ± 0.9) %	-

Λ_c(2625)⁺

$$I = 0$$

Mass $m = 2625.6 \pm 0.8$ MeV

$m - m_{\Lambda_c^+} = 340.6 \pm 0.6$ MeV

Full width $\Gamma < 3.2$ MeV, CL = 90%

Λ _c (2625) ⁺ DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
$\Lambda_c^+ \pi^+ \pi^-$	seen	182
$\Sigma_c(2455)^{++} \pi^- +$	seen	99
$\Sigma_c(2455)^0 \pi^+$		
$\Lambda_c^+ \pi^+ \pi^-$ nonresonant	seen	182

Σ_c(2455)

$$I(J^P) = 1(\frac{1}{2}^+)$$

J^P not confirmed; $\frac{1}{2}^+$ is the quark model prediction.

$\Sigma_c(2455)^{++}$ mass $m = 2453.1 \pm 0.6$ MeV

$\Sigma_c(2455)^+$ mass $m = 2453.8 \pm 0.9$ MeV

$\Sigma_c(2455)^0$ mass $m = 2452.4 \pm 0.7$ MeV (S = 1.1)

Σ _c (2455) DECAY MODES	Fraction (Γ _i /Γ)	ρ (MeV/c)
$\Lambda_c^+ \pi$	100 %	91

Baryon Summary Table



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$I(J^P)$ not confirmed; $\frac{1}{2}(\frac{1}{2}^+)$ is the quark model prediction.

$$\text{Mass } m = 2465.1 \pm 1.6 \text{ MeV}$$

$$\text{Mean life } \tau = (0.35^{+0.07}_{-0.04}) \times 10^{-12} \text{ s}$$

$$c\tau = 106 \text{ } \mu\text{m}$$

Ξ^+ DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Lambda K^- \pi^+ \pi^+$	seen	785
$\Sigma^+ K^- \pi^+$	seen	808
$\Sigma^0 K^- \pi^+ \pi^+$	seen	733
$\Xi^- \pi^+ \pi^+$	seen	850



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

$I(J^P)$ not confirmed; $\frac{1}{2}(\frac{1}{2}^+)$ is the quark model prediction.

$$\text{Mass } m = 2470.3 \pm 1.8 \text{ MeV } (S = 1.3)$$

$$m_{\Xi^0} - m_{\Xi^+} = 5.2 \pm 2.2 \text{ MeV } (S = 1.1)$$

$$\text{Mean life } \tau = (0.098^{+0.023}_{-0.015}) \times 10^{-12} \text{ s}$$

$$c\tau = 29 \text{ } \mu\text{m}$$

A few branching *ratios* but no absolute branching *fractions* have been measured.

Ξ^0 DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$\Xi^- \ell^+$ anything	[<i>m</i>] seen	–
$\Xi^- \pi^+$	seen	875
$\Xi^- \pi^+ \pi^+ \pi^-$	seen	816
$\rho K^- \bar{K}^*(892)^0$	seen	406
$\Omega^- K^+$	seen	522

BOTTOM (BEAUTY) BARYON ($B = -1$)

$$\Lambda_b^0 = udb$$



$$I(J^P) = 0(\frac{1}{2}^+)$$

$I(J^P)$ not yet measured; $0(\frac{1}{2}^+)$ is the quark model prediction.

$$\text{Mass } m = 5641 \pm 50 \text{ MeV}$$

$$\text{Mean life } \tau = (1.07^{+0.19}_{-0.16}) \times 10^{-12} \text{ s}$$

Λ_b^0 DECAY MODES	Fraction (Γ_i/Γ)	p (MeV/c)
$J/\psi(1S)\Lambda$	seen	1756
$\rho D^0 \pi^-$	seen	2383
$\Lambda_c^+ \pi^+ \pi^- \pi^-$	seen	2336
$\Lambda \ell^- X$	seen	–
$\Lambda^+ \ell^- X$	seen	–

NOTES

This Summary Table only includes established baryons. The Full Listings include evidence for other baryons. The masses, widths, and branching fractions for the resonances in this Table are Breit-Wigner parameters. The Full Listings also give, where available, pole parameters. See, in particular, the *Note on N and Δ Resonances*.

For most of the resonances, the parameters come from various partial-wave analyses of more or less the same sets of data, and it is not appropriate to treat the results of the analyses as independent or to average them together. Furthermore, the systematic errors on the results are not well understood. Thus, we usually only give ranges for the parameters. We then also give a best guess for the mass (as part of the name of the resonance) and for the width. The *Note on N and Δ Resonances* and the *Note on Λ and Σ Resonances* in the Full Listings review the partial-wave analyses.

When a quantity has "($S = \dots$)" to its right, the error on the quantity has been enlarged by the "scale factor" S , defined as $S = \sqrt{\chi^2/(N-1)}$, where N is the number of measurements used in calculating the quantity. We do this when $S > 1$, which often indicates that the measurements are inconsistent. When $S > 1.25$, we also show in the Full Listings an ideogram of the measurements. For more about S , see the Introduction.

A decay momentum p is given for each decay mode. For a 2-body decay, p is the momentum of each decay product in the rest frame of the decaying particle. For a 3-or-more-body decay, p is the largest momentum any of the products can have in this frame. For any resonance, the *nominal* mass is used in calculating p . A dagger (" \dagger ") in this column indicates that the mode is forbidden when the nominal masses of resonances are used, but is in fact allowed due to the nonzero widths of the resonances.

[a] The masses of the p and n are most precisely known in u (unified atomic mass units). The conversion factor to MeV, $1 u = 931.49432 \pm 0.00028$ MeV, is less well known than are the masses in u .

[b] The limit is from neutrality-of-matter experiments; it assumes $q_n = q_p + q_e$. See also the charge of the neutron.

[c] The first limit is geochemical and independent of decay mode. The second limit assumes the dominant decay modes are among those investigated. For antiprotons the best limit, inferred from the observation of cosmic ray \bar{p} 's is $\tau_{\bar{p}} > 10^7$ yr, the cosmic-ray storage time, but this limit depends on a number of assumptions. The best direct observation of stored antiprotons gives $\tau_{\bar{p}}/B(\bar{p} \rightarrow e^- \gamma) > 1848$ yr.

[d] There is some controversy about whether nuclear physics and model dependence complicate the analysis for bound neutrons (from which the best limit comes). For reactor experiments with free neutrons, the best limit is $> 10^7$ s.

[e] The parameters g_A , g_V , and g_{WM} for semileptonic modes are defined by $\bar{B}_f[\gamma_\lambda(g_V + g_A\gamma_5) + i(g_{WM}/m_{B_i})\sigma_{\lambda\nu}q^\nu]B_i$, and ϕ_{AV} is defined by $g_A/g_V = |g_A/g_V|e^{i\phi_{AV}}$. See the "Note on Baryon Decay Parameters" in the neutron Full Listings.

[f] Time-reversal invariance requires this to be 0° or 180° .

[g] The decay parameters γ and Δ are calculated from α and ϕ using

$$\gamma = \sqrt{1-\alpha^2} \cos\phi, \quad \tan\Delta = -\frac{1}{\alpha} \sqrt{1-\alpha^2} \sin\phi.$$

See the "Note on Baryon Decay Parameters" in the neutron Full Listings.

[h] See the Full Listings for the pion momentum range used in this measurement.

[i] The error given here is only an educated guess. It is larger than the error on the weighted average of the published values.

[j] A theoretical value using QED.

[k] The branching fraction includes all the decay modes of the final-state resonance.

[l] The value is for the sum of the charge states indicated.

[*m*] ℓ indicates e or μ mode, not sum over modes.

SEARCHES FOR FREE QUARKS, MONOPOLES, SUPERSYMMETRY, COMPOSITENESS, etc.

Free Quark Searches

All searches since 1977 have had negative results.

Magnetic Monopole Searches

Isolated candidate events have not been confirmed. Most experiments obtain negative results.

Supersymmetric Particle Searches

Limits are based on the Minimal Supersymmetric Standard Model.

Assumptions include: 1) $\tilde{\chi}_1^0$ (or $\tilde{\gamma}$) is lightest supersymmetric particle; 2) R -parity is conserved; 3) $m_{\tilde{t}_L} = m_{\tilde{t}_R}$, and all scalar quarks (except \tilde{t}_L and \tilde{t}_R) are degenerate in mass.

See the Full Listings for a Note giving details of supersymmetry.

$\tilde{\chi}_i^0$ — neutralinos (mixtures of $\tilde{\gamma}$, \tilde{Z}^0 , and \tilde{H}^0)

Mass $m_{\tilde{\gamma}} > 15$ GeV, CL = 90%	[if $m_{\tilde{\gamma}} = 100$ GeV (from cosmology)]
Mass $m_{\tilde{\chi}_1^0} > 18$ GeV, CL = 90%	[GUT relations assumed]
Mass $m_{\tilde{\chi}_2^0} > 45$ GeV, CL = 95%	[GUT relations assumed]
Mass $m_{\tilde{\chi}_3^0} > 70$ GeV, CL = 95%	[GUT relations assumed]
Mass $m_{\tilde{\chi}_4^0} > 108$ GeV, CL = 95%	[GUT relations assumed]

$\tilde{\chi}_i^\pm$ — charginos (mixtures of \tilde{W}^\pm and \tilde{H}^\pm)

Mass $m_{\tilde{\chi}_1^\pm} > 45$ GeV, CL = 95%	[all $m_{\tilde{\chi}_1^0}$]
Mass $m_{\tilde{\chi}_2^\pm} > 99$ GeV, CL = 95%	[GUT relations assumed]

$\tilde{\nu}$ — scalar neutrino (sneutrino)

Mass $m > 37.1$ GeV, CL = 95%	[one flavor]
Mass $m > 41.8$ GeV, CL = 95%	[three degenerate flavors]

\tilde{e} — scalar electron (selectron)

Mass $m > 65$ GeV, CL = 95%	[if $m_{\tilde{\gamma}} = 0$]
Mass $m > 50$ GeV, CL = 95%	[if $m_{\tilde{\gamma}} < 5$ GeV]
Mass $m > 45$ GeV, CL = 95%	[if $m_{\tilde{\chi}_1^0} < 41$ GeV]

$\tilde{\mu}$ — scalar muon (smuon)

Mass $m > 45$ GeV, CL = 95%	[if $m_{\tilde{\chi}_1^0} < 41$ GeV]
-----------------------------	--------------------------------------

$\tilde{\tau}$ — scalar tau (stau)

Mass $m > 45$ GeV, CL = 95%	[if $m_{\tilde{\chi}_1^0} < 38$ GeV]
-----------------------------	--------------------------------------

\tilde{q} — scalar quark (squark)

These limits include the effects of cascade decay, for a particular choice of parameters, $\mu = -250$ GeV, $\tan\beta = 2$. The limits are weakly sensitive to these parameters over much of parameter space. Limits assume GUT relations between gaugino masses and the gauge coupling; in particular that for $|\mu|$ not small, $m_{\tilde{\chi}_1^0} \approx m_{\tilde{g}}/6$.

Mass $m > 90$ GeV, CL = 90%	[any $m_{\tilde{g}} < 410$ GeV]
Mass $m > 218$ GeV, CL = 90%	[if $m_{\tilde{g}} = m_{\tilde{q}}$]

\tilde{g} — gluino

There is some controversy about a low-mass window ($1 \lesssim m_{\tilde{g}} \lesssim 4$ GeV). Several experiments cast doubt on the existence of this window.

These limits include the effects of cascade decay, for a particular choice of parameters, $\mu = -250$ GeV, $\tan\beta = 2$. The limits are weakly sensitive to these parameters over much of parameter space. Limits assume GUT relations between gaugino masses and the gauge coupling; in particular that for $|\mu|$ not small, $m_{\tilde{\chi}_1^0} \approx m_{\tilde{g}}/7$.

Mass $m > 100$ GeV, CL = 90%	[any $m_{\tilde{q}}$]
Mass $m > 218$ GeV, CL = 90%	[if $m_{\tilde{q}} \leq m_{\tilde{g}}$]

Searches for Quark and Lepton Compositeness

Scale Limits Λ for Contact Interactions (the lowest dimensional interactions with four fermions)

If the Lagrangian has the form

$$\pm \frac{g^2}{2\Lambda^2} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L$$

(with $g^2/4\pi$ set equal to 1), then we define $\Lambda \equiv \Lambda_{LL}^\pm$. For the full definitions and for other forms, see the Note in the Listings on Searches for Quark and Lepton Compositeness in the full Review and the original literature.

$\Lambda_{LL}^+(e e e e) > 1.6$ TeV, CL = 95%
$\Lambda_{LL}^-(e e e e) > 3.6$ TeV, CL = 95%
$\Lambda_{LL}^+(e e \mu \mu) > 2.6$ TeV, CL = 95%
$\Lambda_{LL}^-(e e \mu \mu) > 1.9$ TeV, CL = 95%
$\Lambda_{LL}^+(e e \tau \tau) > 1.9$ TeV, CL = 95%
$\Lambda_{LL}^-(e e \tau \tau) > 2.9$ TeV, CL = 95%
$\Lambda_{LL}^+(l l l l) > 3.5$ TeV, CL = 95%
$\Lambda_{LL}^-(l l l l) > 2.8$ TeV, CL = 95%
$\Lambda_{LL}^+(e e q q) > 1.7$ TeV, CL = 95%
$\Lambda_{LL}^-(e e q q) > 2.2$ TeV, CL = 95%
$\Lambda_{LL}^+(\mu \mu q q) > 1.4$ TeV, CL = 95%
$\Lambda_{LL}^-(\mu \mu q q) > 1.6$ TeV, CL = 95%
$\Lambda_{LR}^\pm(\nu_\mu \nu_e \mu e) > 3.1$ TeV, CL = 90%
$\Lambda_{LL}^\pm(q q q q) > 1.4$ TeV, CL = 95%

Excited Leptons

The limits from $\ell^{*+} \ell^{*-}$ do not depend on λ (where λ is the $\ell \ell^*$ transition coupling). The λ -dependent limits assume chiral coupling, except for the third limit for e^* which is for nonchiral coupling. For chiral coupling, this limit corresponds to $\lambda_\gamma = \sqrt{2}$.

$e^{*\pm}$ — excited electron

Mass $m > 46.1$ GeV, CL = 95%	(from $e^{*+} e^{*-}$)
Mass $m > 91$ GeV, CL = 95%	(if $\lambda_Z > 1$)
Mass $m > 127$ GeV, CL = 95%	(if $\lambda_\gamma = 1$)

$\mu^{*\pm}$ — excited muon

Mass $m > 46.1$ GeV, CL = 95%	(from $\mu^{*+} \mu^{*-}$)
Mass $m > 91$ GeV, CL = 95%	(if $\lambda_Z > 1$)

$\tau^{*\pm}$ — excited tau

Mass $m > 46.0$ GeV, CL = 95%	(from $\tau^{*+} \tau^{*-}$)
Mass $m > 90$ GeV, CL = 95%	(if $\lambda_Z > 0.18$)

ν^* — excited neutrino

Mass $m > 47$ GeV, CL = 95%	(from $\nu^* \bar{\nu}^*$)
Mass $m > 91$ GeV, CL = 95%	(if $\lambda_Z > 1$)

q^* — excited quark

Mass $m > 45.6$ GeV, CL = 95%	(from $q^* \bar{q}^*$)
Mass $m > 88$ GeV, CL = 95%	(if $\lambda_Z > 1$)
Mass $m > 540$ GeV, CL = 95%	($p \bar{p} \rightarrow q^* X$)

Color Sextet and Octet Particles

Color Sextet Quarks (q_6)

Mass $m > 84$ GeV, CL = 95%	(Stable q_6)
-----------------------------	-----------------

Color Octet Charged Leptons (ℓ_8)

Mass $m > 86$ GeV, CL = 95%	(Stable ℓ_8)
-----------------------------	--------------------

Color Octet Neutrinos (ν_8)

Mass $m > 110$ GeV, CL = 90%	($\nu_8 \rightarrow \nu g$)
------------------------------	-------------------------------

Tests of Conservation Laws

TESTS OF CONSERVATION LAWS

Revised by L. Wolfenstein and T.G. Trippe, June 1994.

In keeping with the current interest in tests of conservation laws, we collect together a Table of experimental limits on all weak and electromagnetic decays, mass differences, and moments, and on a few reactions, whose observation would violate conservation laws. The Table is given only in the full *Review of Particle Properties*, not in the Particle Physics Booklet. For the benefit of Booklet readers, we include the best limits from the Table in the following text. The Table is in two parts: "Discrete Space-Time Symmetries," *i.e.*, C , P , T , CP , and CPT ; and "Number Conservation Laws," *i.e.*, lepton, baryon, hadronic flavor, and charge conservation. The references for these data can be found in the Full Listings in the *Review*. A discussion of these tests follows.

CPT INVARIANCE

General principles of relativistic field theory require invariance under the combined reformation CPT . The simplest tests of CPT invariance are the equality of the masses and lifetimes of a particle and its antiparticle. The best test comes from the limit on the mass difference between K^0 and \bar{K}^0 . Any such difference contributes to the CP -violating parameter ϵ . Assuming CPT invariance, ϕ_ϵ , the phase of ϵ should be very close to 44° . (See the "Note on CP Violation in K_L^0 Decay" in the Full Listings.) In contrast, if the entire source of CP violation in K^0 decays were a $K^0 - \bar{K}^0$ mass difference, ϕ_ϵ would be $44^\circ + 90^\circ$. It is possible to deduce that [1]

$$m_{\bar{K}^0} - m_{K^0} \approx \frac{2(m_{K_L^0} - m_{K_S^0}) |\eta| (\frac{2}{3}\phi_{+-} + \frac{1}{3}\phi_{00} - \phi_\epsilon)}{\sin \phi_\epsilon}.$$

Using our best values of the CP -violation parameters, we get $|(m_{\bar{K}^0} - m_{K^0})/m_{K^0}| \leq 10^{-18}$ (CL = 90%). Limits can also be placed on specific CPT -violating decay amplitudes. Given the small value of $(1 - |\eta_{00}/\eta_{+-}|)$, the value of $\phi_{00} - \phi_{+-}$ provides a measure of CPT violation in $K_L^0 \rightarrow 2\pi$ decay. Results from CERN [1] and Fermilab [2] indicate no CPT -violating effect.

CP AND T INVARIANCE

Given CPT invariance, CP violation and T violation are equivalent. So far the only evidence for CP or T violation comes from the measurements of η_{+-} , η_{00} , and the semileptonic decay charge asymmetry for K_L , *e.g.*, $|\eta_{+-}| = |A(K_L^0 \rightarrow \pi^+\pi^-)/A(K_S^0 \rightarrow \pi^+\pi^-)| = (2.269 \pm 0.023) \times 10^{-3}$ and $[\Gamma(K_L^0 \rightarrow \pi^-e^+\nu) - \Gamma(K_L^0 \rightarrow \pi^+e^-\bar{\nu})]/[\text{sum}] = (0.333 \pm 0.014)\%$. Other searches for CP or T violation divide into (a) those that involve weak interactions or parity violation, and (b) those that involve processes allowed by the strong or electromagnetic interactions. In class (a) the most sensitive is probably the search for an electric dipole moment of the neutron, measured to be $< 1.1 \times 10^{-25}$ e cm (95% CL). A nonzero value requires both P and T violation. Class (b) includes the search for C violation in η decay, believed to be an electromagnetic process, *e.g.*, as measured by $\Gamma(\eta \rightarrow \mu^+\mu^-\pi^0)/\Gamma(\eta \rightarrow \text{all}) < 5 \times 10^{-6}$, and searches for T violation in a number of nuclear and electromagnetic reactions.

CONSERVATION OF LEPTON NUMBERS

Present experimental evidence and the standard electroweak theory are consistent with the absolute conservation of three separate lepton numbers: electron number L_e , muon number L_μ , and tau number L_τ . Searches for violations are of the following types:

a) $\Delta L = 2$ for one type of lepton. The best limit comes from the search for neutrinoless double beta decay $(Z, A) \rightarrow (Z + 2, A) + e^- + e^-$. The best laboratory limit is $t_{1/2} > 1.4 \times 10^{24}$ yr (CL=90%) for ^{76}Ge .

b) Conversion of one lepton type to another. For purely leptonic processes, the best limits are on $\mu \rightarrow e\gamma$ and $\mu \rightarrow 3e$, measured as $\Gamma(\mu \rightarrow e\gamma)/\Gamma(\mu \rightarrow \text{all}) < 5 \times 10^{-11}$ and $\Gamma(\mu \rightarrow 3e)/\Gamma(\mu \rightarrow \text{all}) < 1.0 \times 10^{-12}$. For semileptonic processes, the best limit comes from the coherent conversion process in a muonic atom, $\mu^- + (Z, A) \rightarrow e^- + (Z, A)$, measured as $\Gamma(\mu^- \text{Ti} \rightarrow e^- \text{Ti})/\Gamma(\mu^- \text{Ti} \rightarrow \text{all}) < 4 \times 10^{-12}$. Of special interest is the case in which the hadronic flavor also changes, as in $K_L \rightarrow e\mu$ and $K^+ \rightarrow \pi^+e^-\mu^+$, measured as $\Gamma(K_L \rightarrow e\mu)/\Gamma(K_L \rightarrow \text{all}) < 3.3 \times 10^{-11}$ and $\Gamma(K^+ \rightarrow \pi^+e^-\mu^+)/\Gamma(K^+ \rightarrow \text{all}) < 2.1 \times 10^{-10}$. Limits on the conversion of τ into e or μ are found in τ decay and are much less stringent than those for $\mu \rightarrow e$ conversion, *e.g.*, $\Gamma(\tau \rightarrow \mu\gamma)/\Gamma(\tau \rightarrow \text{all}) < 4.2 \times 10^{-6}$ and $\Gamma(\tau \rightarrow e\gamma)/\Gamma(\tau \rightarrow \text{all}) < 1.2 \times 10^{-4}$.

c) Conversion of one type of lepton into another type of antilepton. The case most studied is $\mu^- + (Z, A) \rightarrow e^+ + (Z - 2, A)$, the strongest limit being $\Gamma(\mu^- \text{Ti} \rightarrow e^+ \text{Ca})/\Gamma(\mu^- \text{Ti} \rightarrow \text{all}) < 9 \times 10^{-11}$.

d) Relation to neutrino mass. If neutrinos have mass, then it is expected even in the standard electroweak theory that the lepton numbers are not separately conserved, as a consequence of lepton mixing analogous to Cabibbo quark mixing. However, in this case lepton-number-violating processes such as $\mu \rightarrow e\gamma$ are expected to have extremely small probability. For small neutrino masses, the lepton-number violation would be observed first in neutrino oscillations, which have been the subject of extensive experimental searches. For example, searches for $\bar{\nu}_e$ disappearance, which we label as $\bar{\nu}_e \not\rightarrow \bar{\nu}_e$, give measured limits $\Delta(m^2) < 0.0083 \text{ eV}^2$ for $\sin^2(2\theta) = 1$, and $\sin^2(2\theta) < 0.14$ for large $\Delta(m^2)$, where θ is the neutrino mixing angle. Searches for $\nu_\mu \rightarrow \nu_e$ set limits $\Delta(m^2) < 0.09 \text{ eV}^2$ for $\sin^2(2\theta) = 1$, and $\sin^2(2\theta) < 0.0025$ for large $\Delta(m^2)$. For larger neutrino masses ($\gg 1 \text{ keV}$), lepton-number violation is searched for by looking for anomalous decays such as $\pi \rightarrow e\nu_x$, where ν_x is a massive neutrino. If the $\Delta L = 2$ type of violation occurs, it is expected that neutrinos will have a nonzero mass of the Majorana type.

CONSERVATION OF HADRONIC FLAVORS

In strong and electromagnetic interactions, hadronic flavor is conserved, *i.e.* the conversion of a quark of one flavor (d, u, s, c, b, t) into a quark of another flavor is forbidden. In the Standard Model, the weak interactions violate these conservation laws in a manner described by the Cabibbo-Kobayashi-Maskawa mixing (see the section "Cabibbo-Kobayashi-Maskawa Mixing Matrix"). The way in which these conservation laws are violated is tested as follows:

a) $\Delta S = \Delta Q$ rule. In the semileptonic decay of strange particles, the strangeness change equals the change in charge of the hadrons. Tests come from limits on decay rates such as $\Gamma(\Sigma^+ \rightarrow ne^+\nu)/\Gamma(\Sigma^+ \rightarrow \text{all}) < 5 \times 10^{-6}$, and from a detailed analysis of $K_L \rightarrow \pi e\nu$, which yields the parameter x , measured to be $(\text{Re } x, \text{Im } x) = (0.006 \pm 0.018, -0.003 \pm 0.026)$. Corresponding rules are $\Delta C = \Delta Q$ and $\Delta B = \Delta Q$.

b) Change of flavor by two units. In the Standard Model this occurs only in second-order weak interactions. The classic example is $\Delta S = 2$ via $K^0 - \bar{K}^0$ mixing, which is directly measured by $m(K_S) - m(K_L) = (3.510 \pm 0.018) \times 10^{-12} \text{ MeV}$. There

is now evidence for $B^0 - \bar{B}^0$ mixing ($\Delta B = 2$), with the corresponding mass difference between the eigenstates ($m_{B_H^0} - m_{B_L^0}$) = $(0.71 \pm 0.06) \Gamma_B = (3.1 \pm 0.4) \times 10^{-10}$ MeV. No evidence exists for $D^0 - \bar{D}^0$ mixing, which is expected to be much smaller in the Standard Model.

c) Flavor-changing neutral currents. In the Standard Model the neutral-current interactions do not change flavor. The low rate $\Gamma(K_L \rightarrow \mu^+ \mu^-)/\Gamma(K_L \rightarrow \text{all}) = (7.4 \pm 0.4) \times 10^{-9}$ puts limits on such interactions; the nonzero value for this rate is attributed to a combination of the weak and electromagnetic interactions. The best test should come from a limit on $K^+ \rightarrow \pi^+ \nu \bar{\nu}$, which occurs in the Standard Model only as a second-order weak process with a branching fraction of $(1 \text{ to } 8) \times 10^{-10}$. The current limit is $\Gamma(K^+ \rightarrow \pi^+ \nu \bar{\nu})/\Gamma(K^+ \rightarrow \text{all}) < 5.2 \times 10^{-9}$. Limits for charm-changing or bottom-changing neutral currents are much less stringent: $\Gamma(D^0 \rightarrow \mu^+ \mu^-)/\Gamma(D^0 \rightarrow \text{all}) < 1.1 \times 10^{-5}$ and $\Gamma(B^0 \rightarrow \mu^+ \mu^-)/\Gamma(B^0 \rightarrow \text{all}) < 5.9 \times 10^{-6}$. One cannot isolate flavor-changing neutral current (FCNC) effects in non leptonic decays. For example, the FCNC transition $s \rightarrow d + (\bar{u} + u)$ is equivalent to the charged-current transition $s \rightarrow u + (\bar{u} + d)$. Tests for FCNC are therefore limited to hadron decays into lepton pairs. Such decays are expected only in second-order in the electroweak coupling in the Standard Model.

References

1. R. Carosi *et al.*, Phys. Lett. **B237**, 303 (1990).
2. M. Karlsson *et al.*, Phys. Rev. Lett. **64**, 2976 (1990); L.K. Gibbons *et al.*, Phys. Rev. Lett. **70**, 1199 (1993).

TESTS OF DISCRETE SPACE-TIME SYMMETRIES

CHARGE CONJUGATION (C)

$\Gamma(\eta \rightarrow 3\gamma)/\Gamma_{\text{total}}$	$< 3.1 \times 10^{-8}$, CL = 90%
$\Gamma((e^+ e^-)_{J=0} \rightarrow 3\gamma)/\Gamma((e^+ e^-)_{J=0} \rightarrow 2\gamma)$	[a] $< 1 \times 10^{-5}$, CL = 90%
$\Gamma((e^+ e^-)_{J=1} \rightarrow 4\gamma)/\Gamma((e^+ e^-)_{J=1} \rightarrow 3\gamma)$	[a] $< 1 \times 10^{-5}$, CL = 90%
η C-nonconserving decay parameters	
$\pi^+ \pi^- \pi^0$ left-right asymmetry parameter	$(0.09 \pm 0.17) \times 10^{-2}$
$\pi^+ \pi^- \pi^0$ sextant asymmetry parameter	$(0.18 \pm 0.16) \times 10^{-2}$
$\pi^+ \pi^- \pi^0$ quadrant asymmetry parameter	$(-0.17 \pm 0.17) \times 10^{-2}$
$\pi^+ \pi^- \gamma$ left-right asymmetry parameter	$(0.9 \pm 0.4) \times 10^{-2}$
$\pi^+ \pi^- \gamma$ parameter β (D -wave)	0.05 ± 0.06 ($S = 1.5$)
$\Gamma(\eta \rightarrow 3\gamma)/\Gamma_{\text{total}}$	$< 5 \times 10^{-4}$, CL = 95%
$\Gamma(\eta \rightarrow \pi^0 e^+ e^-)/\Gamma_{\text{total}}$	[b] $< 4 \times 10^{-5}$, CL = 90%
$\Gamma(\eta \rightarrow \pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$	[b] $< 5 \times 10^{-6}$, CL = 90%

PARITY (P)

e electric dipole moment	$(-0.3 \pm 0.8) \times 10^{-26}$ ecm
μ electric dipole moment	$(3.7 \pm 3.4) \times 10^{-19}$ ecm
τ electric dipole moment	$< 5 \times 10^{-17}$ ecm, CL = 95%
$\Gamma(\eta \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$	$< 1.5 \times 10^{-3}$
p electric dipole moment	$(-4 \pm 6) \times 10^{-23}$ ecm
n electric dipole moment	$< 11 \times 10^{-26}$ ecm, CL = 95%
Λ electric dipole moment	$< 1.5 \times 10^{-16}$ ecm, CL = 95%

TIME REVERSAL (T)

e electric dipole moment	$(-0.3 \pm 0.8) \times 10^{-26}$ ecm
μ electric dipole moment	$(3.7 \pm 3.4) \times 10^{-19}$ ecm
μ decay parameters	
transverse e^+ polarization normal to plane of μ spin, e^+ momentum	0.007 ± 0.023
α'/A	$(0 \pm 4) \times 10^{-3}$
β'/A	$(2 \pm 6) \times 10^{-3}$
τ electric dipole moment	$< 5 \times 10^{-17}$ ecm, CL = 95%
$\text{Im}(\xi)$ in $K_{\mu 3}^\pm$ decay (from transverse μ pol.)	-0.017 ± 0.025
$\text{Im}(\xi)$ in $K_{\mu 3}^0$ decay (from transverse μ pol.)	-0.007 ± 0.026
p electric dipole moment	$(-4 \pm 6) \times 10^{-23}$ ecm
n electric dipole moment	$< 11 \times 10^{-26}$ ecm, CL = 95%
$n \rightarrow p e^- \nu$ decay parameters	
ϕ_{AV} , phase of g_A relative to g_V	[c] $(180.07 \pm 0.18)^\circ$
triple correlation coefficient D	$(-0.5 \pm 1.4) \times 10^{-3}$
Λ electric dipole moment	$< 1.5 \times 10^{-16}$ ecm, CL = 95%
triple correlation coefficient D for $\Sigma^- \rightarrow n e^- \bar{\nu}_e$	0.11 ± 0.10

CHARGE CONJUGATION TIMES PARITY (CP)

τ weak dipole moment	$< 3.7 \times 10^{-17}$ ecm, CL = 95%
$\Gamma(\eta \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$	$< 1.5 \times 10^{-3}$
$K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ rate difference/average	$(0.07 \pm 0.12)\%$
$K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ rate difference/average	$(0.0 \pm 0.6)\%$
$K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ rate difference/average	$(0.9 \pm 3.3)\%$
$(g_{\tau^+} - g_{\tau^-}) / (g_{\tau^+} + g_{\tau^-})$ for $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$	$(-0.7 \pm 0.5)\%$
CP-violation parameters in K_S^0 decay	
$\text{Im}(\eta_{+-0})^2 = \Gamma(K_S^0 \rightarrow \pi^+ \pi^- \pi^0, \text{CP-violating}) / \Gamma(K_L^0 \rightarrow \pi^+ \pi^- \pi^0)$	< 0.12 , CL = 90%
$\text{Im}(\eta_{000})^2 = \Gamma(K_S^0 \rightarrow 3\pi^0) / \Gamma(K_L^0 \rightarrow 3\pi^0)$	< 0.1 , CL = 90%
charge asymmetry J for $K_L^0 \rightarrow \pi^+ \pi^- \pi^0$	0.0011 ± 0.0008
$ \eta_{+-\gamma} = A(K_L^0 \rightarrow \pi^+ \pi^- \gamma) / A(K_S^0 \rightarrow \pi^+ \pi^- \gamma) $	$(2.15 \pm 0.26 \pm 0.20) \times 10^{-3}$
$\phi_{+-\gamma}$ = phase of $\eta_{+-\gamma}$	$(72 \pm 23 \pm 17)^\circ$
$ \epsilon_{+-\gamma}' /\epsilon$	< 0.3 , CL = 90%
$\Gamma(K_L^0 \rightarrow \pi^0 \mu^+ \mu^-)/\Gamma_{\text{total}}$	[d] $< 5.1 \times 10^{-9}$, CL = 90%
$\Gamma(K_L^0 \rightarrow \pi^0 e^+ e^-)/\Gamma_{\text{total}}$	[d] $< 4.3 \times 10^{-9}$, CL = 90%
$\Gamma(K_L^0 \rightarrow \pi^0 \nu \bar{\nu})/\Gamma_{\text{total}}$	[e] $< 2.2 \times 10^{-4}$, CL = 90%
$[\Gamma(D^0 \rightarrow K^+ K^-) - \Gamma(\bar{D}^0 \rightarrow K^+ K^-)]/\text{sum}$	< 0.45 , CL = 90%
$ \text{Re}(\epsilon_{B^0}) $	< 0.045
$[\alpha_-(\Lambda) + \alpha_+(\bar{\Lambda})] / [\alpha_-(\Lambda) - \alpha_+(\bar{\Lambda})]$	-0.03 ± 0.06

CHARGE CONJUGATION TIMES PARITY (CP) VIOLATION OBSERVED

charge asymmetry in K_{L}^0 decays	
$\delta(\mu) = [\Gamma(\pi^- \mu^+ \nu_\mu) - \Gamma(\pi^+ \mu^- \bar{\nu}_\mu)]/\text{sum}$	$(0.304 \pm 0.025)\%$
$\delta(e) = [\Gamma(\pi^- e^+ \nu_e) - \Gamma(\pi^+ e^- \bar{\nu}_e)]/\text{sum}$	$(0.333 \pm 0.014)\%$
parameters for $K_L^0 \rightarrow 2\pi$ decay	
$ \eta_{00} = A(K_L^0 \rightarrow 2\pi^0) / A(K_S^0 \rightarrow 2\pi^0) $	$(2.259 \pm 0.023) \times 10^{-3}$ ($S = 1.1$)
$ \eta_{+-} = A(K_L^0 \rightarrow \pi^+ \pi^-) / A(K_S^0 \rightarrow \pi^+ \pi^-) $	$(2.269 \pm 0.023) \times 10^{-3}$ ($S = 1.1$)
$e'/\epsilon \approx \text{Re}(e'/\epsilon) = (1 - \eta_{00}/\eta_{+-})/3$	[f] $(1.5 \pm 0.8) \times 10^{-3}$ ($S = 1.8$)
ϕ_{+-} , phase of η_{+-}	$(44.3 \pm 0.8)^\circ$
ϕ_{00} , phase of η_{00}	$(43.3 \pm 1.3)^\circ$
$\Gamma(K_L^0 \rightarrow \pi^+ \pi^-)/\Gamma_{\text{total}}$	$(2.03 \pm 0.04) \times 10^{-3}$ ($S = 1.2$)
$\Gamma(K_L^0 \rightarrow \pi^0 \pi^0)/\Gamma_{\text{total}}$	$(9.14 \pm 0.34) \times 10^{-4}$ ($S = 1.8$)

Tests of Conservation Laws

CPT INVARIANCE

$(m_{W^+} - m_{W^-}) / m_{\text{average}}$	-0.002 ± 0.007
$(m_{e^+} - m_{e^-}) / m_{\text{average}}$	$< 4 \times 10^{-8}$, CL = 90%
$ q_{e^+} + q_{e^-} /e$	$< 4 \times 10^{-8}$
$(g_{e^+} - g_{e^-}) / g_{\text{average}}$	$(-0.5 \pm 2.1) \times 10^{-12}$
$\tau_{\mu^+} / \tau_{\mu^-}$ mean life ratio	1.00002 ± 0.00008
$(\tau_{\mu^+} - \tau_{\mu^-}) / \tau_{\text{average}}$	$(2 \pm 8) \times 10^{-5}$
$(g_{\mu^+} - g_{\mu^-}) / g_{\text{average}}$	$(-2.6 \pm 1.6) \times 10^{-8}$
$(m_{\pi^+} - m_{\pi^-}) / m_{\text{average}}$	$(2 \pm 5) \times 10^{-4}$
$(\tau_{\pi^+} - \tau_{\pi^-}) / \tau_{\text{average}}$	$(6 \pm 7) \times 10^{-4}$
$(m_{K^+} - m_{K^-}) / m_{\text{average}}$	$(-0.6 \pm 1.8) \times 10^{-4}$
$(\tau_{K^+} - \tau_{K^-}) / \tau_{\text{average}}$	$(0.11 \pm 0.09)\%$ (S = 1.2)
$\kappa^{\pm} \rightarrow \mu^{\pm} \nu_{\mu}$ rate difference/average	$(-0.5 \pm 0.4)\%$
$\kappa^{\pm} \rightarrow \pi^{\pm} \pi^0$ rate difference/average	[g] $(0.8 \pm 1.2)\%$
$ m_{K^0} - m_{\bar{K}^0} / m_{\text{average}}$	[h] $< 9 \times 10^{-19}$
phase difference $\phi_{00} - \phi_{+-}$	$(-1.0 \pm 1.0)^{\circ}$
$(m_{\rho^-} - m_{\bar{\rho}^-}) / m_{\text{average}}$	$(2 \pm 4) \times 10^{-8}$
$ q_{\rho^-} + q_{\bar{\rho}^-} /e$	$< 2 \times 10^{-5}$
$(\mu_{\rho^-} - \mu_{\bar{\rho}^-}) / \mu_{\text{average}} $	$(-2.6 \pm 2.9) \times 10^{-3}$
$(m_{\eta} - m_{\bar{\eta}}) / m_{\text{average}}$	$(9 \pm 5) \times 10^{-5}$
$(m_{\Lambda} - m_{\bar{\Lambda}}) / m_{\Lambda}$	$(-1.0 \pm 0.9) \times 10^{-5}$
$(\tau_{\Lambda} - \tau_{\bar{\Lambda}}) / \tau_{\text{average}}$	0.04 ± 0.09
$(\mu_{\Sigma^+} - \mu_{\bar{\Sigma}^-}) / \mu_{\text{average}} $	0.014 ± 0.015
$(m_{\Xi^-} - m_{\bar{\Xi}^+}) / m_{\text{average}}$	$(1.1 \pm 2.7) \times 10^{-4}$
$(\tau_{\Xi^-} - \tau_{\bar{\Xi}^+}) / \tau_{\text{average}}$	0.02 ± 0.18
$(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\text{average}}$	$(0 \pm 5) \times 10^{-4}$

TESTS OF NUMBER CONSERVATION LAWS

LEPTON FAMILY NUMBER

Lepton family number conservation means separate conservation of each of L_e, L_{μ}, L_{τ} .

$\Gamma(Z \rightarrow e^{\pm} \mu^{\mp}) / \Gamma_{\text{total}}$	[i] $< 6 \times 10^{-6}$, CL = 95%
$\Gamma(Z \rightarrow e^{\pm} \tau^{\mp}) / \Gamma_{\text{total}}$	[i] $< 1.3 \times 10^{-5}$, CL = 95%
$\Gamma(Z \rightarrow \mu^{\pm} \tau^{\mp}) / \Gamma_{\text{total}}$	[i] $< 1.9 \times 10^{-5}$, CL = 95%
limit on $\mu^- \rightarrow e^-$ conversion $\sigma(\mu^- 32S \rightarrow e^- 32S) / \sigma(\mu^- 32S \rightarrow \nu_{\mu} 32P^*)$	$< 7 \times 10^{-11}$, CL = 90%
$\sigma(\mu^- \text{Ti} \rightarrow e^- \text{Ti}) / \sigma(\mu^- \text{Ti} \rightarrow \text{capture})$	$< 4.3 \times 10^{-12}$, CL = 90%
limit on muonium \rightarrow antimuonium conversion $R_{\bar{G}} = G_{\bar{C}} / G_F$	< 0.13 , CL = 90%
$\Gamma(\mu^- \rightarrow e^- \nu_e \bar{\nu}_{\mu}) / \Gamma_{\text{total}}$	[j] $< 1.2 \times 10^{-2}$, CL = 90%
$\Gamma(\mu^- \rightarrow e^- \gamma) / \Gamma_{\text{total}}$	$< 4.9 \times 10^{-11}$, CL = 90%
$\Gamma(\mu^- \rightarrow e^- e^+ e^-) / \Gamma_{\text{total}}$	$< 1.0 \times 10^{-12}$, CL = 90%
$\Gamma(\mu^- \rightarrow e^- 2\gamma) / \Gamma_{\text{total}}$	$< 7.2 \times 10^{-11}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \gamma) / \Gamma_{\text{total}}$	$< 1.2 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \gamma) / \Gamma_{\text{total}}$	$< 4.2 \times 10^{-6}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \pi^0) / \Gamma_{\text{total}}$	$< 1.4 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \pi^0) / \Gamma_{\text{total}}$	$< 4.4 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- K^0) / \Gamma_{\text{total}}$	$< 1.3 \times 10^{-3}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- K^0) / \Gamma_{\text{total}}$	$< 1.0 \times 10^{-3}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \eta) / \Gamma_{\text{total}}$	$< 6.3 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \eta) / \Gamma_{\text{total}}$	$< 7.3 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \rho^0) / \Gamma_{\text{total}}$	$< 1.9 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \rho^0) / \Gamma_{\text{total}}$	$< 2.9 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- K^*(892)^0) / \Gamma_{\text{total}}$	$< 3.8 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- K^*(892)^0) / \Gamma_{\text{total}}$	$< 4.5 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \ell^- \ell^+) / \Gamma_{\text{total}}$	[k] $< 3.4 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- e^+ e^-) / \Gamma_{\text{total}}$	$< 1.3 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 2.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^+ \mu^- \mu^-) / \Gamma_{\text{total}}$	$< 1.6 \times 10^{-5}$, CL = 90%

$\Gamma(\tau^- \rightarrow (\mu e e^-) / \Gamma_{\text{total}}$	$< 2.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- e^+ e^-) / \Gamma_{\text{total}}$	$< 1.4 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 1.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \ell^{\pm} \pi^{\mp} \pi^-) / \Gamma_{\text{total}}$	[l,k] $< 6.3 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^{\mp} \pi^{\pm} \pi^-) / \Gamma_{\text{total}}$	[l] $< 6.0 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \pi^+ \pi^-) / \Gamma_{\text{total}}$	$< 2.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^{\mp} \pi^{\pm} \pi^-) / \Gamma_{\text{total}}$	[l] $< 3.9 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \pi^+ \pi^-) / \Gamma_{\text{total}}$	$< 3.6 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \ell^{\pm} \pi^{\mp} K^-) / \Gamma_{\text{total}}$	[l,k] $< 1.2 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow (e \pi K)^-, \text{ all charged}) / \Gamma_{\text{total}}$	$< 7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \pi^{\pm} K^{\mp}) / \Gamma_{\text{total}}$	[l] $< 5.8 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \pi^+ K^-) / \Gamma_{\text{total}}$	$< 2.9 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \pi^- K^+) / \Gamma_{\text{total}}$	$< 5.8 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow (\mu \pi K)^-, \text{ all charged}) / \Gamma_{\text{total}}$	$< 7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \pi^{\pm} K^{\mp}) / \Gamma_{\text{total}}$	[l] $< 7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \pi^+ K^-) / \Gamma_{\text{total}}$	$< 7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^- \pi^- K^+) / \Gamma_{\text{total}}$	$< 7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^- \text{ light spinless boson}) / \Gamma_{\text{total}}$	$< 3.2 \times 10^{-3}$, CL = 95%
$\Gamma(\tau^- \rightarrow \mu^- \text{ light spinless boson}) / \Gamma_{\text{total}}$	$< 6 \times 10^{-3}$, CL = 95%
ν oscillations. (For other lepton mixing effects in particle decays, see the Full Listings.)	
$\bar{\nu}_e \neq \bar{\nu}_e$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 0.0083 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	< 0.14 , CL = 68%
$\nu_e \rightarrow \nu_{\tau}$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 9 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	< 0.12 , CL = 90%
$\bar{\nu}_e \rightarrow \bar{\nu}_{\tau}$	
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	< 0.7 , CL = 90%
$\nu_{\mu} \rightarrow \nu_e$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 0.09 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	$< 2.5 \times 10^{-3}$, CL = 90%
$\bar{\nu}_{\mu} \rightarrow \bar{\nu}_e$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 0.14 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	< 0.004 , CL = 95%
$\nu_{\mu}(\bar{\nu}_{\mu}) \rightarrow \nu_e(\bar{\nu}_e)$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 0.075 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	$< 3 \times 10^{-3}$, CL = 90%
$\nu_{\mu} \rightarrow \nu_{\tau}$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 0.9 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	< 0.004 , CL = 90%
$\bar{\nu}_{\mu} \rightarrow \bar{\nu}_{\tau}$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 2.2 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	$< 4.4 \times 10^{-2}$, CL = 90%
$\nu_{\mu}(\bar{\nu}_{\mu}) \rightarrow \nu_{\tau}(\bar{\nu}_{\tau})$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 1.5 \text{ eV}^2$, CL = 90%
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	$< 8 \times 10^{-3}$, CL = 90%
$\nu_e \neq \nu_e$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$< 2.3 \text{ eV}^2$
$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$	$< 7 \times 10^{-2}$, CL = 90%
$\nu_{\mu} \neq \nu_{\mu}$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	< 0.23 or $> 1500 \text{ eV}^2$
$\sin^2(2\theta)$ for $\Delta(m^2) = 100 \text{ eV}^2$	[l] < 0.02 , CL = 90%
$\bar{\nu}_{\mu} \neq \bar{\nu}_{\mu}$	
$\Delta(m^2)$ for $\sin^2(2\theta) = 1$	< 7 or $> 1200 \text{ eV}^2$
$\sin^2(2\theta)$ for $190 \text{ eV}^2 < \Delta(m^2) < 320 \text{ eV}^2$	[m] < 0.02 , CL = 90%
$\Gamma(\pi^+ \rightarrow \mu^+ \nu_e) / \Gamma_{\text{total}}$	[n] $< 8.0 \times 10^{-3}$, CL = 90%
$\Gamma(\pi^+ \rightarrow \mu^- e^+ e^+ \nu) / \Gamma_{\text{total}}$	$< 1.6 \times 10^{-6}$, CL = 90%
$\Gamma(\pi^0 \rightarrow \mu^+ e^- + e^- \mu^+) / \Gamma_{\text{total}}$	$< 1.72 \times 10^{-8}$, CL = 90%
$\Gamma(K^+ \rightarrow \mu^- \nu_e e^+) / \Gamma_{\text{total}}$	$< 2.0 \times 10^{-8}$, CL = 90%
$\Gamma(K^+ \rightarrow \mu^+ \nu_e) / \Gamma_{\text{total}}$	[n] $< 4 \times 10^{-3}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^+ \mu^+ e^-) / \Gamma_{\text{total}}$	$< 2.1 \times 10^{-10}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^+ \mu^- e^+) / \Gamma_{\text{total}}$	$< 7 \times 10^{-9}$, CL = 90%
$\Gamma(K_L^0 \rightarrow e^{\pm} \mu^{\mp}) / \Gamma_{\text{total}}$	[l] $< 3.3 \times 10^{-11}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^+ e^{\pm} \mu^{\mp}) / \Gamma_{\text{total}}$	[l] $< 3.8 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^+ e^+ \mu^-) / \Gamma_{\text{total}}$	$< 3.3 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^+ e^- \mu^+) / \Gamma_{\text{total}}$	$< 3.3 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow K^+ e^+ \mu^-) / \Gamma_{\text{total}}$	$< 3.4 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow K^+ e^- \mu^+) / \Gamma_{\text{total}}$	$< 3.4 \times 10^{-3}$, CL = 90%
$\Gamma(D^0 \rightarrow \mu^{\pm} e^{\mp}) / \Gamma_{\text{total}}$	[l] $< 1.0 \times 10^{-4}$, CL = 90%
$\Gamma(B^+ \rightarrow \pi^+ e^+ \mu^-) / \Gamma_{\text{total}}$	$< 6.4 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow \pi^+ e^- \mu^+) / \Gamma_{\text{total}}$	$< 6.4 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow K^+ e^+ \mu^-) / \Gamma_{\text{total}}$	$< 6.4 \times 10^{-3}$, CL = 90%

Limits are given at the 90% confidence level, while errors are given as ± 1 standard deviation.

Tests of Conservation Laws

$\Gamma(B^+ \rightarrow K^+ e^- \mu^+)/\Gamma_{\text{total}}$	$<6.4 \times 10^{-3}$, CL = 90%
$\Gamma(B^0 \rightarrow e^\pm \mu^\mp)/\Gamma_{\text{total}}$	[f] $<5.9 \times 10^{-6}$, CL = 90%
$\Gamma(B^0 \rightarrow e^\pm \tau^\mp)/\Gamma_{\text{total}}$	[f] $<5.3 \times 10^{-4}$, CL = 90%
$\Gamma(B^0 \rightarrow \mu^\pm \tau^\mp)/\Gamma_{\text{total}}$	[f] $<8.3 \times 10^{-4}$, CL = 90%

TOTAL LEPTON NUMBER

Violation of total lepton number conservation also implies violation of lepton family number conservation.

limit on $\mu^- \rightarrow e^+$ conversion	
$\sigma(\mu^- 32\text{S} \rightarrow e^+ 32\text{Si}^*) / \sigma(\mu^- 32\text{S} \rightarrow \nu_\mu 32\text{P}^*)$	$<9 \times 10^{-10}$, CL = 90%
$\sigma(\mu^- 127\text{I} \rightarrow e^+ 127\text{Sb}^*) / \sigma(\mu^- 127\text{I} \rightarrow \text{anything})$	$<3 \times 10^{-10}$, CL = 90%
$\sigma(\mu^- \text{Ti} \rightarrow e^+ \text{Ca}) / \sigma(\mu^- \text{Ti} \rightarrow \text{capture})$	$<8.9 \times 10^{-11}$, CL = 90%
$\Gamma(\tau^- \rightarrow \pi^- \gamma)/\Gamma_{\text{total}}$	$<2.8 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \pi^- \pi^0)/\Gamma_{\text{total}}$	$<3.7 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^+ e^- e^-)/\Gamma_{\text{total}}$	$<1.4 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \ell^\pm \pi^\mp \pi^-)/\Gamma_{\text{total}}$	[i,k] $<6.3 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^\mp \pi^\pm \pi^-)/\Gamma_{\text{total}}$	[f] $<6.0 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^+ \pi^- \pi^-)/\Gamma_{\text{total}}$	[f] $<1.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^\mp \pi^\pm \pi^-)/\Gamma_{\text{total}}$	[f] $<3.9 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^+ \pi^- \pi^-)/\Gamma_{\text{total}}$	[f] $<3.9 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \ell^\pm \pi^\mp K^-)/\Gamma_{\text{total}}$	[i,k] $<1.2 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow (e\pi K)^-, \text{all charged})/\Gamma_{\text{total}}$	$<7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow e^+ \pi^- K^-)/\Gamma_{\text{total}}$	$<2.0 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow (\mu\pi K)^-, \text{all charged})/\Gamma_{\text{total}}$	$<7.7 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \mu^+ \pi^- K^-)/\Gamma_{\text{total}}$	$<4.0 \times 10^{-5}$, CL = 90%
$\Gamma(\tau^- \rightarrow \bar{p}\gamma)/\Gamma_{\text{total}}$	$<2.9 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \bar{p}\pi^0)/\Gamma_{\text{total}}$	$<6.6 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \bar{p}\eta)/\Gamma_{\text{total}}$	$<1.30 \times 10^{-3}$, CL = 90%
$\nu_e \rightarrow (\bar{\nu}_e)_L$	
$\alpha\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$<0.14 \text{ eV}^2$, CL = 90%
$\alpha^2 \sin^2(2\theta)$ for "Large" $\Delta(m^2)$	<0.032 , CL = 90%
$\nu_\mu \rightarrow (\bar{\nu}_\mu)_L$	
$\alpha\Delta(m^2)$ for $\sin^2(2\theta) = 1$	$<0.16 \text{ eV}^2$, CL = 90%
$\alpha^2 \sin^2(2\theta)$ for "Large" $\Delta(m^2)$	<0.001 , CL = 90%
$\Gamma(\pi^+ \rightarrow \mu^+ \bar{\nu}_e)/\Gamma_{\text{total}}$	[n] $<1.5 \times 10^{-3}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^- \mu^+ e^+)/\Gamma_{\text{total}}$	$<7 \times 10^{-9}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^- e^+ e^+)/\Gamma_{\text{total}}$	$<1.0 \times 10^{-8}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$	$<1.5 \times 10^{-4}$, CL = 90%
$\Gamma(K^+ \rightarrow \mu^+ \bar{\nu}_e)/\Gamma_{\text{total}}$	[n] $<3.3 \times 10^{-3}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^0 e^+ \bar{\nu}_e)/\Gamma_{\text{total}}$	[n] $<3 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^- e^+ e^+)/\Gamma_{\text{total}}$	$<4.8 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$	$<6.8 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^- e^+ \mu^+)/\Gamma_{\text{total}}$	$<3.7 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow K^- e^+ e^+)/\Gamma_{\text{total}}$	$<9.1 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$	$<4.3 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow K^- e^+ \mu^+)/\Gamma_{\text{total}}$	$<4.0 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow \pi^- e^+ e^+)/\Gamma_{\text{total}}$	$<3.9 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow \pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$	$<9.1 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow \pi^- e^+ \mu^+)/\Gamma_{\text{total}}$	$<6.4 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow K^- e^+ e^+)/\Gamma_{\text{total}}$	$<3.9 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$	$<9.1 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow K^- e^+ \mu^+)/\Gamma_{\text{total}}$	$<6.4 \times 10^{-3}$, CL = 90%
$\Gamma(\Xi^- \rightarrow p\mu^- \mu^-)/\Gamma_{\text{total}}$	$<4 \times 10^{-4}$, CL = 90%

BARYON NUMBER

$\Gamma(\tau^- \rightarrow \bar{p}\gamma)/\Gamma_{\text{total}}$	$<2.9 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \bar{p}\pi^0)/\Gamma_{\text{total}}$	$<6.6 \times 10^{-4}$, CL = 90%
$\Gamma(\tau^- \rightarrow \bar{p}\eta)/\Gamma_{\text{total}}$	$<1.30 \times 10^{-3}$, CL = 90%
p mean life	$>1.6 \times 10^{25}$ years
A few examples of proton or bound neutron decay follow. For limits on many other nucleon decay channels, see the Baryon Summary Table.	
$\tau(N \rightarrow e^+ \pi)$	>130 (n), >550 (p) $\times 10^{30}$ years, CL = 90%
$\tau(N \rightarrow \mu^+ \pi)$	>100 (n), >270 (p) $\times 10^{30}$ years, CL = 90%
$\tau(N \rightarrow e^+ K)$	>1.3 (n), >150 (p) $\times 10^{30}$ years, CL = 90%
$\tau(N \rightarrow \mu^+ K)$	>1.1 (n), >120 (p) $\times 10^{30}$ years, CL = 90%
mean time for $n\bar{n}$ transition in vacuum	[o] $>1.2 \times 10^8$ s, CL = 90%

ELECTRIC CHARGE (Q)

e mean life / branching fraction	[p] $>2.7 \times 10^{23}$ yr, CL = 68%
$\Gamma(n \rightarrow p\nu_e \bar{\nu}_e)/\Gamma_{\text{total}}$	$<9 \times 10^{-24}$, CL = 90%

 $\Delta S = \Delta Q$ RULE

Allowed in second-order weak interactions.

$\Gamma(K^+ \rightarrow \pi^+ \pi^+ e^- \bar{\nu}_e)/\Gamma_{\text{total}}$	$<1.2 \times 10^{-8}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^+ \pi^+ \mu^- \bar{\nu}_\mu)/\Gamma_{\text{total}}$	$<3.0 \times 10^{-6}$, CL = 95%
$x = A(\bar{K}^0 \rightarrow \pi^- \ell^+ \nu)/A(K^0 \rightarrow \pi^- \ell^+ \nu) = A(\Delta S = -\Delta Q)/A(\Delta S = \Delta Q)$	
real part of x	0.006 ± 0.018 (S = 1.3)
imaginary part of x	-0.003 ± 0.026 (S = 1.2)
$\Gamma(\Sigma^+ \rightarrow n\ell^+ \nu)/\Gamma(\Sigma^- \rightarrow n\ell^- \bar{\nu})$	<0.043
$\Gamma(\Sigma^+ \rightarrow n e^+ \nu_e)/\Gamma_{\text{total}}$	$<5 \times 10^{-6}$, CL = 90%
$\Gamma(\Sigma^+ \rightarrow n \mu^+ \nu_\mu)/\Gamma_{\text{total}}$	$<3.0 \times 10^{-5}$, CL = 90%
$\Gamma(\Xi^0 \rightarrow \Sigma^- e^+ \nu_e)/\Gamma_{\text{total}}$	$<9 \times 10^{-4}$, CL = 90%
$\Gamma(\Xi^0 \rightarrow \Sigma^- \mu^+ \nu_\mu)/\Gamma_{\text{total}}$	$<9 \times 10^{-4}$, CL = 90%

 $\Delta S = 2$ FORBIDDEN

Allowed in second-order weak interactions.

$\Gamma(\Xi^0 \rightarrow p\pi^-)/\Gamma_{\text{total}}$	$<4 \times 10^{-5}$, CL = 90%
$\Gamma(\Xi^0 \rightarrow p e^- \bar{\nu}_e)/\Gamma_{\text{total}}$	$<1.3 \times 10^{-3}$
$\Gamma(\Xi^0 \rightarrow p \mu^- \bar{\nu}_\mu)/\Gamma_{\text{total}}$	$<1.3 \times 10^{-3}$
$\Gamma(\Xi^- \rightarrow n\pi^-)/\Gamma_{\text{total}}$	$<1.9 \times 10^{-5}$, CL = 90%
$\Gamma(\Xi^- \rightarrow n e^- \bar{\nu}_e)/\Gamma_{\text{total}}$	$<3.2 \times 10^{-3}$, CL = 90%
$\Gamma(\Xi^- \rightarrow n \mu^- \bar{\nu}_\mu)/\Gamma_{\text{total}}$	$<1.5 \times 10^{-2}$, CL = 90%
$\Gamma(\Xi^- \rightarrow p\pi^- \pi^-)/\Gamma_{\text{total}}$	$<4 \times 10^{-4}$, CL = 90%
$\Gamma(\Xi^- \rightarrow p\pi^- e^- \bar{\nu}_e)/\Gamma_{\text{total}}$	$<4 \times 10^{-4}$, CL = 90%
$\Gamma(\Xi^- \rightarrow p\pi^- \mu^- \bar{\nu}_\mu)/\Gamma_{\text{total}}$	$<4 \times 10^{-4}$, CL = 90%
$\Gamma(\Omega^- \rightarrow \Lambda\pi^-)/\Gamma_{\text{total}}$	$<1.9 \times 10^{-4}$, CL = 90%

 $\Delta S = 2$ VIA MIXING

Allowed in second-order weak interactions, e.g. mixing.

$m_{K_L^0} - m_{K_S^0}$	$(0.5333 \pm 0.0027) \times 10^{10} \hbar s^{-1}$ (S = 1.2)
$m_{K_L^0} - m_{K_S^0}$	$(3.510 \pm 0.018) \times 10^{-12} \text{ MeV}$ (S = 1.3)

 $\Delta C = 2$ VIA MIXING

Allowed in second-order weak interactions, e.g. mixing.

$ m_{D_1^0} - m_{D_2^0} $	[q] $<20 \times 10^{10} \hbar s^{-1}$, CL = 90%
$\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+)$	<0.0037 , CL = 90%
$\Gamma(\mu^- \text{ anything (via } \bar{D}^0))/\Gamma(\mu^+ \text{ anything})$	<0.0056 , CL = 90%

$\Delta B = 2$ VIA MIXING

Allowed in second-order weak interactions, e.g. mixing.

χ_d	0.156 ± 0.024
$\Delta m_{B^0} = m_{B^0_H} - m_{B^0_L}$	$(0.51 \pm 0.06) \times 10^{12} \hbar s^{-1}$
$\chi_d = \Delta m_{B^0} / \Gamma_{B^0}$	0.71 ± 0.06
χ_s	0.62 ± 0.13
$\Delta m_{B^0_s} = m_{B^0_{sH}} - m_{B^0_{sL}}$	$> 1.8 \times 10^{12} \hbar s^{-1}$, CL = 95%
$\chi_s = \Delta m_{B^0_s} / \Gamma_{B^0_s}$	> 2.0 , CL = 95%

 $\Delta S = 1$ WEAK NEUTRAL CURRENT FORBIDDEN

Allowed by higher-order electroweak interactions.

$\Gamma(K^+ \rightarrow \pi^+ e^+ e^-) / \Gamma_{\text{total}}$	$(2.74 \pm 0.23) \times 10^{-7}$
$\Gamma(K^+ \rightarrow \pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 2.3 \times 10^{-7}$, CL = 90%
$\Gamma(K^+ \rightarrow \pi^+ \nu \bar{\nu}) / \Gamma_{\text{total}}$	$< 5.2 \times 10^{-9}$, CL = 90%
$\Gamma(K^0_S \rightarrow \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 3.2 \times 10^{-7}$, CL = 90%
$\Gamma(K^0_S \rightarrow e^+ e^-) / \Gamma_{\text{total}}$	$< 1.0 \times 10^{-5}$, CL = 90%
$\Gamma(K^0_S \rightarrow \pi^0 e^+ e^-) / \Gamma_{\text{total}}$	$< 1.1 \times 10^{-6}$, CL = 90%
$\Gamma(K^0_L \rightarrow \mu^+ \mu^-) / \Gamma_{\text{total}}$	$(7.4 \pm 0.4) \times 10^{-9}$
$\Gamma(K^0_L \rightarrow \mu^+ \mu^- \gamma) / \Gamma_{\text{total}}$	$(2.8 \pm 2.8) \times 10^{-7}$
$\Gamma(K^0_L \rightarrow e^+ e^-) / \Gamma_{\text{total}}$	$< 4.1 \times 10^{-11}$, CL = 90%
$\Gamma(K^0_L \rightarrow e^+ e^- \gamma) / \Gamma_{\text{total}}$	$(9.1 \pm 0.5) \times 10^{-6}$
$\Gamma(K^0_L \rightarrow e^+ e^- \gamma \gamma) / \Gamma_{\text{total}}$	[r] $(6.6 \pm 3.2) \times 10^{-7}$
$\Gamma(K^0_L \rightarrow \pi^+ \pi^- e^+ e^-) / \Gamma_{\text{total}}$	$< 2.5 \times 10^{-6}$, CL = 90%
$\Gamma(K^0_L \rightarrow \mu^+ \mu^- e^+ e^-) / \Gamma_{\text{total}}$	$< 4.9 \times 10^{-6}$, CL = 90%
$\Gamma(K^0_L \rightarrow e^+ e^- e^+ e^-) / \Gamma_{\text{total}}$	[s] $(3.9 \pm 0.7) \times 10^{-8}$
$\Gamma(K^0_L \rightarrow \pi^0 \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 5.1 \times 10^{-9}$, CL = 90%
$\Gamma(K^0_L \rightarrow \pi^0 e^+ e^-) / \Gamma_{\text{total}}$	$< 4.3 \times 10^{-9}$, CL = 90%
$\Gamma(K^0_L \rightarrow \pi^0 \nu \bar{\nu}) / \Gamma_{\text{total}}$	$< 2.2 \times 10^{-4}$, CL = 90%
$\Gamma(\Sigma^+ \rightarrow p e^+ e^-) / \Gamma_{\text{total}}$	$< 7 \times 10^{-6}$

 $\Delta C = 1$ WEAK NEUTRAL CURRENT FORBIDDEN

Allowed by higher-order electroweak interactions.

$\Gamma(D^+ \rightarrow \pi^+ e^+ e^-) / \Gamma_{\text{total}}$	$< 2.5 \times 10^{-3}$, CL = 90%
$\Gamma(D^+ \rightarrow \pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 2.9 \times 10^{-3}$, CL = 90%
$\Gamma(D^0 \rightarrow e^+ e^-) / \Gamma_{\text{total}}$	$< 1.3 \times 10^{-4}$, CL = 90%
$\Gamma(D^0 \rightarrow \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 1.1 \times 10^{-5}$, CL = 90%
$\Gamma(D^0 \rightarrow \rho^0 e^+ e^-) / \Gamma_{\text{total}}$	$< 4.5 \times 10^{-4}$, CL = 90%
$\Gamma(D^0 \rightarrow \rho^0 \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 8.1 \times 10^{-4}$, CL = 90%

 $\Delta B = 1$ WEAK NEUTRAL CURRENT FORBIDDEN

Allowed by higher-order electroweak interactions.

$\Gamma(B^+ \rightarrow \pi^+ e^+ e^-) / \Gamma_{\text{total}}$	$< 3.9 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow \pi^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 9.1 \times 10^{-3}$, CL = 90%
$\Gamma(B^+ \rightarrow K^+ e^+ e^-) / \Gamma_{\text{total}}$	$< 6 \times 10^{-5}$, CL = 90%
$\Gamma(B^+ \rightarrow K^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 1.7 \times 10^{-4}$, CL = 90%
$\Gamma(B^+ \rightarrow K^*(892)^+ e^+ e^-) / \Gamma_{\text{total}}$	$< 6.9 \times 10^{-4}$, CL = 90%
$\Gamma(B^+ \rightarrow K^*(892)^+ \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 1.2 \times 10^{-3}$, CL = 90%
$\Gamma(\bar{B} \rightarrow e^+ e^- \text{ anything}) / \Gamma_{\text{total}}$	[t] $< 2.4 \times 10^{-3}$, CL = 90%
$\Gamma(\bar{B} \rightarrow \mu^+ \mu^- \text{ anything}) / \Gamma_{\text{total}}$	[t] $< 5.0 \times 10^{-5}$, CL = 90%
$\Gamma(B^0 \rightarrow e^+ e^-) / \Gamma_{\text{total}}$	$< 5.9 \times 10^{-6}$, CL = 90%
$\Gamma(B^0 \rightarrow \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 5.9 \times 10^{-6}$, CL = 90%
$\Gamma(B^0 \rightarrow K^0 e^+ e^-) / \Gamma_{\text{total}}$	$< 3.0 \times 10^{-4}$, CL = 90%
$\Gamma(B^0 \rightarrow K^0 \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 3.6 \times 10^{-4}$, CL = 90%
$\Gamma(B^0 \rightarrow K^*(892)^0 e^+ e^-) / \Gamma_{\text{total}}$	$< 2.9 \times 10^{-4}$, CL = 90%
$\Gamma(B^0 \rightarrow K^*(892)^0 \mu^+ \mu^-) / \Gamma_{\text{total}}$	$< 2.3 \times 10^{-5}$, CL = 90%

NOTES

In this Summary Table:

When a quantity has "(S = ...)" to its right, the error on the quantity has been enlarged by the "scale factor" S, defined as $S = \sqrt{\chi^2 / (N - 1)}$, where N is the number of measurements used in calculating the quantity. We do this when $S > 1$, which often indicates that the measurements are inconsistent. When $S > 1.25$, we also show in the Full Listings an ideogram of the measurements. For more about S, see the Introduction.

- [a] Positronium data are from A.P. Mills and S. Berko, Physical Review Letters **18** 420 (1967); and K. Marko and A. Rich, Physical Review Letters **33** 980 (1974). Values for 90% confidence level, are from A.P. Mills, private communication.
- [b] C parity forbids this to occur as a single-photon process.
- [c] Time-reversal invariance requires this to be 0° or 180° .
- [d] Allowed by higher-order electroweak interactions.
- [e] Violates CP in leading order. Test of direct CP violation since the indirect CP-violating and CP-conserving contributions are expected to be suppressed.
- [f] ϵ'/ϵ is derived from $|\eta_{00}/\eta_{+-}|$ measurements using theoretical input on phases.
- [g] Neglecting photon channels. See, e.g., A. Pais and S.B. Treiman, Phys. Rev. **D12**, 2744 (1975).
- [h] Derived from measured values of ϕ_{+-} , ϕ_{00} , $|\eta|$, $\tau_{K_S^0}$, and $|m_{K_L^0} - m_{K_S^0}|$, as described in the introduction to this Table.
- [i] The value is for the sum of the charge states indicated.
- [j] A test of additive vs. multiplicative lepton family number conservation.
- [k] ℓ means a sum over e and μ modes.
- [l] $\Delta(m^2) = 100 \text{ eV}^2$.
- [m] $190 \text{ eV}^2 < \Delta(m^2) < 320 \text{ eV}^2$.
- [n] Derived from an analysis of neutrino-oscillation experiments.
- [o] There is some controversy about whether nuclear physics and model dependence complicate the analysis for bound neutrons (from which the best limit comes). For reactor experiments with free neutrons, the best limit is $> 10^7 \text{ s}$.
- [p] This is the best "electron disappearance" limit. The best limit for the mode $e^- \rightarrow \nu \gamma$ is $> 2.35 \times 10^{25} \text{ yr}$ (CL=68%).
- [q] The $D_1^0 - D_2^0$ limits are inferred from the limit on $D^0 \rightarrow \bar{D}^0 \rightarrow K^+ \pi^-$.
- [r] See the K_L^0 Full Listings for the energy limits used in this measurement.
- [s] $m_{e^- e^-} > 470 \text{ MeV}$.
- [t] B^0 , B^+ , and B_s^0 not separated.

REVIEWS, TABLES, AND PLOTS

1. Physical constants (revised)	1233
2. Astrophysical constants (revised)	1234
3. Big-bang cosmology (revised)	1236
4. Dark matter (revised)	1238
5. International System of units (SI)	1240
6. Atomic and nuclear properties of materials (revised)	1241
7. Periodic table of the elements (revised)	1243
8. Electronic structure of the elements	1244
9. High-energy collider parameters (revised)	1246
10. Passage of particles through matter (revised)	1251
11. Photon and electron attenuation plots (revised)	1259
12. Particle detectors	1261
13. Radioactivity and radiation protection	1268
14. Cosmic ray fluxes	1269
15. Commonly used radioactive sources	1270
16. Probability (revised)	1271
17. Statistics (revised)	1275
18. Monte Carlo techniques (revised)	1283
19. Electromagnetic relations	1285
20. Clebsch-Gordan coefficients, spherical harmonics, and d functions	1287
21. $SU(3)$ isoscalar factors and representation matrices	1288
22. $SU(n)$ multiplets and Young diagrams	1289
23. Kinematics	1290
24. Cross-section formulae for specific processes (revised)	1294
25. Quantum chromodynamics (revised)	1297
26. Standard model of electroweak interactions (revised)	1304
27. Constraints on new physics from electroweak analyses (new)	1312
28. Cabibbo-Kobayashi-Maskawa mixing matrix (revised)	1315
29. CP violation (new)	1318
30. Quark model	1319
31. Naming scheme for hadrons	1323
32. Monte Carlo particle numbering scheme (revised)	1325
33. Plots of cross sections and related quantities (revised)	1327

Additional Reviews and Notes related to specific particles are located in the Full Listings.

1. PHYSICAL CONSTANTS

Table 1.1. Reviewed 1993 by B.N. Taylor, NIST. Based mainly on the “1986 Adjustment of the Fundamental Physical Constants” by E.R. Cohen and B.N. Taylor, Rev. Mod. Phys. **59**, 1121 (1987). The last group of constants (beginning with the Fermi coupling constant) comes from the Particle Data Group. The figures in parentheses after the values give the 1-standard-deviation uncertainties in the last digits; the corresponding uncertainties in parts per million (ppm) are given in the last column. This set of constants (aside from the last group) is recommended for international use by CODATA (the Committee on Data for Science and Technology).

Since the 1986 adjustment, new experiments have yielded improved values for a number of constants, including the Rydberg constant R_∞ , the Planck constant h , the fine-structure constant α , and the molar gas constant R , and hence also for constants directly derived from these, such as the Boltzmann constant k and Stefan-Boltzmann constant σ . The new results and their impact on the 1986 recommended values are discussed extensively in “Recommended Values of the Fundamental Physical Constants: A Status Report,” B.N. Taylor and E.R. Cohen, J. Res. Natl. Inst. Stand. Technol. **95**, 497 (1990); see also E.R. Cohen and B.N. Taylor, “The Fundamental Physical Constants,” Phys. Today, August 1993 Part 2, BG9. In general, the new results give uncertainties for the affected constants that are 5 to 7 times smaller than the 1986 uncertainties, but the changes in the values themselves are smaller than twice the 1986 uncertainties. Because the output values of a least-squares adjustment are correlated, the new results cannot readily be incorporated with the 1986 values. Until the next complete adjustment of the constants, the 1986 CODATA set, given (in part) below, remains the set of choice.

Quantity	Symbol, equation	Value	Uncert. (ppm)
speed of light in vacuum	c	299 792 458 m s ⁻¹	exact*
Planck constant	h	6.626 075 5(40) × 10 ⁻³⁴ J s	0.60
Planck constant, reduced	$\hbar \equiv h/2\pi$	1.054 572 66(63) × 10 ⁻³⁴ J s = 6.582 122 0(20) × 10 ⁻²² MeV s	0.60 0.30
electron charge magnitude	e	1.602 177 33(49) × 10 ⁻¹⁹ C = 4.803 206 8(15) × 10 ⁻¹⁰ esu	0.30, 0.30
conversion constant	$\hbar c$	197.327 053(59) MeV fm	0.30
conversion constant	$(\hbar c)^2$	0.389 379 66(23) GeV ² mbarn	0.59
electron mass	m_e	0.510 999 06(15) MeV/c ² = 9.109 389 7(54) × 10 ⁻³¹ kg	0.30, 0.59
proton mass	m_p	938.272 31(28) MeV/c ² = 1.672 623 1(10) × 10 ⁻²⁷ kg = 1.007 276 470(12) u = 1836.152 701(37) m_e	0.30, 0.59 0.012, 0.020
deuteron mass	m_d	1875.613 39(57) MeV/c ²	0.30
unified atomic mass unit (u)	(mass ¹² C atom)/12 = (1 g)/(N _A mol)	931.494 32(28) MeV/c ² = 1.660 540 2(10) × 10 ⁻²⁷ kg	0.30, 0.59
permittivity of free space	ϵ_0	8.854 187 817 ... × 10 ⁻¹² F m ⁻¹ 4π × 10 ⁻⁷ N A ⁻² = 12.566 370 614 ... × 10 ⁻⁷ N A ⁻²	exact
permeability of free space	μ_0		exact
fine-structure constant	$\alpha = e^2/4\pi\epsilon_0\hbar c$	1/137.035 989 5(61) [†]	0.045
classical electron radius	$r_e = e^2/4\pi\epsilon_0 m_e c^2$	2.817 940 92(38) × 10 ⁻¹⁵ m	0.13
electron Compton wavelength	$\lambda_e = \hbar/m_e c = r_e \alpha^{-1}$	3.861 593 23(35) × 10 ⁻¹³ m	0.089
Bohr radius ($m_{\text{nucleus}} = \infty$)	$a_\infty = 4\pi\epsilon_0 \hbar^2/m_e e^2 = r_e \alpha^{-2}$	0.529 177 249(24) × 10 ⁻¹⁰ m	0.045
wavelength of 1 eV/c particle	hc/e	1.239 842 44(37) × 10 ⁻⁶ m	0.30
Rydberg energy	$hcR_\infty = m_e e^4/2(4\pi\epsilon_0)^2 \hbar^2 = m_e c^2 \alpha^2/2$	13.605 698 1(40) eV	0.30
Thomson cross section	$\sigma_T = 8\pi r_e^2/3$	0.665 246 16(18) barn	0.27
Bohr magneton	$\mu_B = e\hbar/2m_e$	5.788 382 63(52) × 10 ⁻¹¹ MeV T ⁻¹	0.089
nuclear magneton	$\mu_N = e\hbar/2m_p$	3.152 451 66(28) × 10 ⁻¹⁴ MeV T ⁻¹	0.089
electron cyclotron freq./field	$\omega_{\text{cycl}}^e/B = e/m_e$	1.758 819 62(53) × 10 ¹¹ rad s ⁻¹ T ⁻¹	0.30
proton cyclotron freq./field	$\omega_{\text{cycl}}^p/B = e/m_p$	9.578 830 9(29) × 10 ⁷ rad s ⁻¹ T ⁻¹	0.30
gravitational constant	G_N	6.672 59(85) × 10 ⁻¹¹ m ³ kg ⁻¹ s ⁻² = 6.707 11(86) × 10 ⁻³⁹ $\hbar c$ (GeV/c ²) ⁻²	128 128
standard grav. accel., sea level	g	9.806 65 m s ⁻²	exact
Avogadro constant	N_A	6.022 136 7(36) × 10 ²³ mol ⁻¹	0.59
Boltzmann constant	k	1.380 658(12) × 10 ⁻²³ J K ⁻¹ = 8.617 385(73) × 10 ⁻⁵ eV K ⁻¹	8.5 8.4
molar volume, ideal gas at STP	$N_A k(273.15 \text{ K})/(101 325 \text{ Pa})$	22.414 10(19) × 10 ⁻³ m ³ mol ⁻¹	8.4
Wien displacement law constant	$b = \lambda_{\text{max}} T$	2.897 756(24) × 10 ⁻³ m K	8.4
Stefan-Boltzmann constant	$\sigma = \pi^2 k^4/60\hbar^3 c^2$	5.670 51(19) × 10 ⁻⁸ W m ⁻² K ⁻⁴	34
Fermi coupling constant [‡]	$G_F/(\hbar c)^3$	1.166 39(2) × 10 ⁻⁵ GeV ⁻²	20
weak mixing angle	$\sin^2 \hat{\theta}(M_Z) (\overline{\text{MS}})$	0.2319(5)	2200
W^\pm boson mass	m_W	80.22(26) GeV/c ²	3200
Z^0 boson mass	m_Z	91.187(7) GeV/c ²	77
strong coupling constant	$\alpha_s(m_Z)$	0.116(5)	43000
$\pi = 3.141 592 653 589 793 238$ $e = 2.718 281 828 459 045 235$ $\gamma = 0.577 215 664 901 532 861$			
1 in ≡ 0.0254 m	1 G ≡ 10 ⁻⁴ T	1 eV = 1.602 177 33(49) × 10 ⁻¹⁹ J	kT at 300 K = [38.681 49(33)] ⁻¹ eV
1 Å ≡ 10 nm	1 dyne ≡ 10 ⁻⁵ N	1 eV/c ² = 1.782 662 70(54) × 10 ⁻³⁶ kg	0° C ≡ 273.15 K
1 barn ≡ 10 ⁻²⁸ m ²	1 erg ≡ 10 ⁻⁷ J	2.997 924 58 × 10 ⁹ esu = 1 C	1 atmosphere ≡ 760 torr ≡ 101 325 Pa

* The meter is defined to be the length of path traveled by light in vacuum in 1/299 792 458 s. See B.W. Petley, Nature **303**, 373 (1983).

† At $Q^2 = 0$. At $Q^2 \approx m_W^2$ the value is approximately 1/128.

‡ See discussion in Sec. 26 “Standard Model of electroweak interactions.”

2. ASTROPHYSICAL CONSTANTS

Table 2.1. Written and revised with the help of K.R. Lang, K.A. Olive, J. Primack, S. Rudaz, E. M. Standish, Jr., and M.S. Turner. The figures in parentheses after some values give the 1-standard deviation uncertainties in the last digit(s). While every effort has been made to obtain the most accurate current values of the listed quantities, the table does not represent a critical review or adjustment of the constants, and is not intended as a primary reference.

Quantity	Symbol, equation	Value	Reference
speed of light	c	$299\,792\,458\text{ m s}^{-1}$	defined [1]
Newtonian gravitational constant	G_N	$6.672\,59(85) \times 10^{-11}\text{ m}^3\text{ kg}^{-1}\text{ s}^{-2}$	[2]
astronomical unit	AU	$1.495\,978\,706\,6(2) \times 10^{11}\text{ m}$	[3,4]
tropical year (equinox to equinox) (1994)	yr	$31\,556\,925.2\text{ s}$	[3]
sidereal year (fixed star to fixed star) (1994)		$31\,558\,149.8\text{ s}$	[3]
mean sidereal day		$23^{\text{h}}\,56^{\text{m}}\,04^{\text{s}}.90\,53$	[3]
Jansky	Jy	$10^{-26}\text{ W m}^{-2}\text{ Hz}^{-1}$	
Planck mass	$\sqrt{\hbar c/G_N}$	$1.221\,047(79) \times 10^{19}\text{ GeV}/c^2$ $= 2.176\,71(14) \times 10^{-8}\text{ kg}$	uses [2]
parsec (1 AU/1 arc sec)	pc	$3.085\,677\,580\,7(4) \times 10^{16}\text{ m} = 3.262\dots\text{ ly}$	[5]
light year (deprecated unit)	ly	$0.306\,6\dots\text{ pc} = 0.946\,1\dots \times 10^{16}\text{ m}$	
Schwarzschild radius of the Sun	$2G_N M_\odot/c^2$	$2.953\,250\,08\text{ km}$	[6]
solar mass	M_\odot	$1.988\,92(25) \times 10^{30}\text{ kg}$	[7]
solar luminosity	L_\odot	$3.846 \times 10^{26}\text{ W}$	[8]
solar equatorial radius	R_\odot	$6.96 \times 10^8\text{ m}$	[3]
Earth equatorial radius	R_\oplus	$6.378\,140 \times 10^6\text{ m}$	[3]
Earth mass	M_\oplus	$5.973\,70(76) \times 10^{24}\text{ kg}$	[9]
luminosity conversion	L	$3.02 \times 10^{28} \times 10^{-0.4 M_b}\text{ W}$ (M_b = absolute bolometric magnitude = bolometric magnitude at 10 pc)	[10]
flux conversion	\mathcal{F}	$2.52 \times 10^{-8} \times 10^{-0.4 m_b}\text{ W m}^{-2}$ (m_b = apparent bolometric magnitude)	from above
v_\odot around center of Galaxy	Θ_\odot	$220(20)\text{ km s}^{-1}$	[11]
solar distance from galactic center	R_\odot	$8.0(5)\text{ kpc}$	[12]
Hubble constant [†]	H_0	$100 h_0\text{ km s}^{-1}\text{ Mpc}^{-1}$ $= h_0 \times (9.778\,13\text{ Gyr})^{-1}$	[13]
normalized Hubble constant [†]	h_0	$0.5 < h_0 < 0.85$	[14,15]
critical density of the universe [†]	$\rho_c = 3H_0^2/8\pi G_N$	$2.775\,366\,27 \times 10^{11} h_0^2 M_\odot\text{ Mpc}^{-3}$ $= 1.878\,82(24) \times 10^{-29} h_0^2\text{ g cm}^{-3}$ $= 1.053\,94(13) \times 10^{-5} h_0^2\text{ GeV cm}^{-3}$	
local disk density	ρ_{disk}	$3\text{--}12 \times 10^{-24}\text{ g cm}^{-3} \approx 3\text{--}7\text{ GeV}/c^2\text{ cm}^{-3}$	[16]
local halo density	ρ_{halo}	$3\text{--}7 \times 10^{-25}\text{ g cm}^{-3} \approx 0.2\text{--}0.4\text{ GeV}/c^2\text{ cm}^{-3}$	[17]
density parameter of the universe [†]	$\Omega_0 \equiv \rho_0/\rho_c$	$0.1 < \Omega_0 < 2$	[18]
scaled cosmological constant [†]	$\lambda_0 = \Lambda c^2/3H_0^2$	$-1 < \lambda_0 < 2$	[19,20]
scale factor for cosmological constant [†]	$c^2/3H_0^2$	$2.853 \times 10^{51} h_0^{-2}\text{ m}^2$	
age of the universe [†]	t_0	$15(5)\text{ Gyr}$	[10]
	$\Omega_0 h_0^2$	≤ 2.4 for $t_0 \geq 10\text{ Gyr}$ ≤ 1 for $t_0 \geq 10\text{ Gyr}$, $h_0 > 0.4$	[10] [10]
cosmic background radiation (CBR) temperature [†]	T_0	$2.726 \pm 0.005\text{ K}$	[21]
solar velocity with respect to CBR		$369.5 \pm 3.0\text{ km s}^{-1}$	[22]
energy density of CBR	ρ_γ	$4.647\,7 \times 10^{-34} (T/2.726)^4\text{ g cm}^{-3}$ $= 0.260\,71 (T/2.726)^4\text{ eV cm}^{-3}$	[10]
number density of CBR photons	n_γ	$410.89 (T/2.726)^3\text{ cm}^{-3}$	[10]
entropy density/Boltzmann constant	s/k	$2\,892.4 (T/2.726)^3\text{ cm}^{-3}$	[10]

[†] Subscript 0 indicates present-day values.

References:

1. B.W. Petley, *Nature* **303**, 373 (1983).
2. E.R. Cohen and B.N. Taylor, *Rev. Mod. Phys.* **59**, 1121 (1987). The set of constants resulting from this adjustment has been recommended for international use by CODATA (Committee on Data for Science and Technology).
3. *The Astronomical Almanac for the year 1994*, U.S. Government Printing Office, Washington, and Her Majesty's Stationary Office, London (1993). Where possible, the values as adjusted for the fitting of the ephemerides to all the observational data are used.
4. JPL Planetary Ephemerides, E. Myles Standish, Jr., private communication (1989).
5. 1 AU divided by $\pi/648\,000$; quoted error is from the JPL Planetary Ephemerides value of the AU [4].
6. Heliocentric gravitational constant from Ref. 3 times $2/c^2$. The given 9-place accuracy appears to be consistent with uncertainties in actually defining the earth's orbital parameters.
7. Obtained from the heliocentric gravitational constant [3] and G_N [2]. The error is the 128 ppm standard deviation quoted for G_N .
8. It is surprisingly difficult to find a definitive value for this important constant. In all cases, the solar luminosity is calculated as $4\pi \times (1 \text{ AU})^2$ times the solar constant (or total solar irradiance, TSI). The luminosity given is reduced from $\text{TSI} = 1367.51 \pm 0.01 \text{ W m}^{-2}$, obtained from SMM/ACRIMI spacecraft measurements during the interval 2/80–6/89 [23]. While the time constant for energy production by the sun is very long, radiation from the surface might be modulated or otherwise modified by sunspots; this has apparently not been taken into account. Accordingly, we quote 4-place accuracy. We do not know the actual error, but suppose it might be 5 or 10 in the last place. Sackmann *et al.* [24] use $\text{TSI} = 1370 \pm 2 \text{ W m}^{-2}$, but conclude that the solar luminosity ($L_\odot = 3.853 \times 10^{26} \text{ J s}^{-1}$) has an uncertainty of 1.5%. Their value is based on three 1977–83 papers, and they comment that the error is based on scatter among the reported values, which is substantially in excess of that expected from the individual quoted errors. The conclusion of the 1971 review by Thekaekara and Drummond [25] ($1353 \pm 1\% \text{ W m}^{-2}$) is often quoted [26], and a luminosity based on this value was tabulated in the last two editions of this Review. The conversion to luminosity is not given in the Thekaekara and Drummond paper, and we cannot exactly reproduce the solar luminosity given in Ref. 26. Finally, a value based on the 1954 spectral curve due to Johnson [27] ($1395 \pm 1\% \text{ W m}^{-2}$, or $L_\odot = 3.92 \times 10^{26} \text{ J s}^{-1}$) has been used widely, and may be the basis for higher value of the solar luminosity and corresponding lower value of the solar absolute bolometric magnitude (4.72) still common in the literature [10].
9. Obtained from the geocentric gravitational constant [3] and G_N [2]. The error is the 128 ppm standard deviation quoted for G_N .
10. E.W. Kolb and M.S. Turner, *The Early Universe*, Addison-Wesley (1990).
11. F.J. Kerr and D. Lynden-Bell, *Mon. Not. R. Astr. Soc.* **221**, 1023–1038 (1985). “On the basis of this review these [$R_o = 8.5 \pm 1.1 \text{ kpc}$ and $\Theta_o = 220 \pm 20 \text{ km s}^{-1}$] were adopted by resolution of IAU Commission 33 on 1985 November 21 at Delhi”.
12. M.J. Reid, *Annu. Rev. Astron. Astrophys.* **31**, 345–372 (1993). Note that Θ_o from the 1985 IAU Commission 33 recommendations is adopted in this review, although the new value for R_o is smaller.
13. Conversion using length of tropical year.
14. P.J.E. Peebles, *Principles of Physical Cosmology*, Princeton University Press (1993).
15. Kolb and Turner [10] give the more conservative limits $0.4 < h_0 < 1$. For other conclusions, see the recent reviews by Jacoby *et al.* [28], who say “Using the weighted or unweighted Virgo distances to bootstrap to the Coma cluster, we find the Hubble constant to be either 80 ± 11 or $73 \pm 11 \text{ km s}^{-1} \text{ Mpc}^{-1}$, respectively,” and Huchra [29], who concludes that “Values are still clustered about two numbers, but these numbers are now 50 and 85. A preponderance of the newest local estimates favors the higher value of $85 \text{ km s}^{-1} \text{ Mpc}^{-1}$...”.
16. G. Gilmore, R.F.G. Wyse, and K. Kuijken, *Annu. Rev. Astron. Astrophys.* **27**, 555 (1989).
17. J.N. Bahcall, M. Schmidt, and R.M. Soneira, *Astrophys. J.* **265**, 730 (1983); E. Gates and M.S. Turner, *Phys. Rev. Lett.* **72**, 2520 (1994) state that “...in plausible CDM models the local CDM density is at least $10^{-25} \text{ g cm}^{-3}$. The value $0.3 \text{ GeV}/c^2$ has been taken as “standard” in several papers setting limits on WIMP mass limits, e.g. in M. Mori *et al.*, *Phys. Lett.* **B289**, 463 (1992).
18. Tonry gives $0.2 < \Omega_0 < 2$ [19]. We extend the lower limit so as not to exclude the absence of nonbaryonic dark matter.
19. J. L. Tonry, in *Proc. Texas/PASCOS 92: Relativistic Astrophysics and Particle Cosmology*, ed. C.W. Akerlof and M. Srednicki (Ann. N. Y. Acad. Sci. **688**, 113 (1993).
20. S.M. Carroll and W. H. Press, *Annu. Rev. Astron. Astrophys.* **30**, 499 (1992).
21. J. C. Mather *et al.*, *Astrophys. J.* **420**, 439 (1994). Error quoted here is 1σ .
22. A. Kogut *et al.*, *Astrophys. J.* **419**, 1 (1993).
23. R.C. Willson, pp 5–18 in *Atlas of Satellite Observations Related to Global Change*, ed. J.L. Foster and C.L. Parkinson, Cambridge University Press, (1993).
24. I.-J. Sackmann, A.I. Boothroyd, and K.E. Kraemer, *Astrophys. J.* **418**, 457 (1993).
25. M.P. Thekaekara and A.J. Drummond, *Nature Phys. Sci.* **229**, 6 (1971).
26. K.R. Lang, *Astrophysical Formulae*, Springer-Verlag (1974); K.R. Lang, *Astrophysical Data: Planets and Stars*, Springer-Verlag (1992).
27. F.S. Johnson, *J. Meteorol.* **11**, 431 (1954).
28. G.H. Jacoby *et al.*, *J. Astron. Soc. Pacific* **104**, 599–662 (1992).
29. J.P. Huchra, *Science* **256**, 321–325 (1992).

3. BIG-BANG COSMOLOGY

Revised November 1993 by K.A. Olive.

At early times, and today on a sufficiently large scale, our Universe is very nearly homogeneous and isotropic. The most general space-time metric for a homogeneous, isotropic space is the Friedmann-Robertson-Walker metric (with $c = 1$) [1,2,3]:

$$ds^2 = dt^2 - R^2(t) \left[\frac{dr^2}{1 - \kappa r^2} + r^2 (d\theta^2 + \sin^2 \theta d\phi^2) \right]. \quad (3.1)$$

$R(t)$ is a scale factor for distances in comoving coordinates. With appropriate rescaling of the coordinates, κ can be chosen to be +1, -1, or 0, corresponding to closed, open, or spatially flat geometries. Einstein's equations lead to the Friedmann equation

$$H^2 \equiv \left(\frac{\dot{R}}{R} \right)^2 = \frac{8\pi G_N \rho}{3} - \frac{\kappa}{R^2} + \frac{\Lambda}{3}, \quad (3.2)$$

as well as to

$$\frac{\ddot{R}}{R} = \frac{\Lambda}{3} - \frac{4\pi G_N}{3} (\rho + 3p), \quad (3.3)$$

where $H(t)$ is the Hubble parameter, ρ is the total mass-energy density, p is the isotropic pressure, and Λ is the cosmological constant. (For limits on Λ , see the Table of Astrophysical Constants; we will assume here $\Lambda = 0$.) The Friedmann equation serves to define the density parameter Ω_0 (subscript 0 indicates present-day values):

$$\kappa/R_0^2 = H_0^2(\Omega_0 - 1), \quad \Omega_0 = \rho_0/\rho_c; \quad (3.4)$$

and the critical density is defined as

$$\rho_c \equiv \frac{3H_0^2}{8\pi G_N} = 1.88 \times 10^{-29} h^2 \text{ g cm}^{-3}, \quad (3.5)$$

with

$$H_0 = 100 h_0 \text{ km s}^{-1} \text{ Mpc}^{-1} = h_0/(9.78 \text{ Gyr}). \quad (3.6)$$

Observational bounds give $0.4 < h_0 < 1$. The three curvature signatures $\kappa = +1, -1$, and 0 correspond to $\Omega_0 > 1$, < 1 , and $= 1$. Knowledge of Ω_0 is even poorer than that of h_0 . Luminous matter (stars and associated material) contribute $\Omega_{\text{lum}} \leq 0.01$. There is no lack of evidence for copious amounts of dark matter: rotation curves of spiral galaxies, virial estimates of cluster masses, gravitational lensing by clusters and individual galaxies, and so on. The minimum amount of dark matter required to explain the flat rotation curves of spiral galaxies only amounts to $\Omega_0 \sim 0.1$, while estimates for Ω_0 based upon cluster virial masses suggests $\Omega_0 \sim 0.2 - 0.4$. The highest estimates for the mass density come from studies of the peculiar motions of galaxies (including our own); estimates for Ω_0 obtained by relating peculiar velocity measurements to the distribution galaxies within a few hundred Mpc approach unity. A conservative range for the mass density is: $0.1 \leq \Omega_0 \leq 2$. The excess of Ω_0 over Ω_{lum} leads to the inference that most of the matter in the Universe is nonluminous dark matter.

In an expanding universe, the wavelength of light emitted from a distant source is shifted towards the red. The redshift z is defined such that $1 + z$ is the ratio of the detected wavelength (λ) to emitted (laboratory) wavelength (λ_e) of some electromagnetic spectral feature. It follows from the metric given in Eq. (3.1) that

$$1 + z = \lambda/\lambda_e = R_0/R_e \quad (3.7)$$

where R_e is the value of the scale factor at the time the light was emitted. For light emitted in the not too distant past, one can expand R_e and write $R_e \simeq R_0 + (t_e - t_0)\dot{R}_0$. For small (compared to H_0^{-1}) $\Delta t = (t_e - t_0)$, Eq. (3.7) takes the form of Hubble's law

$$z \approx \Delta t \frac{\dot{R}_0}{R_0} \approx \ell H_0, \quad (3.8)$$

where ℓ is the distance to the source.

Energy conservation implies that

$$\dot{\rho} = -3(\dot{R}/R)(\rho + p), \quad (3.9)$$

so that for a matter-dominated ($p = 0$) universe $\rho \propto R^{-3}$, while for a radiation-dominated ($p = \rho/3$) universe $\rho \propto R^{-4}$. Thus the less singular curvature term κ/R^2 in the Friedmann equation can be neglected at early times when R is small. If the Universe expands adiabatically, the entropy per comoving volume ($\equiv R^3 s$) is constant, where the entropy density is $s = (\rho + p)/T$ and T is temperature. The energy density of radiation can be expressed (with $\hbar = c = 1$) as

$$\rho_r = \frac{\pi^2}{30} N(T) (kT)^4, \quad (3.10)$$

where $N(T)$ counts the effectively massless degrees of freedom of bosons and fermions:

$$N(T) = \sum_B g_B + \frac{7}{8} \sum_F g_F. \quad (3.11)$$

For example, for $m_\mu > kT > m_e$, $N(T) = g_\gamma + 7/8(g_e + 3g_\nu) = 2 + 7/8[4 + 3(2)] = 43/4$. For $m_\pi > kT > m_\mu$, $N(T) = 57/4$. At temperatures less than about 1 MeV, neutrinos have decoupled from the thermal background, *i.e.*, the weak interaction rates are no longer fast enough compared with the expansion rate to keep neutrinos in equilibrium with the remaining thermal bath consisting of γ, e^\pm . Furthermore, at temperatures $kT < m_e$, by entropy conservation, the ratio of the neutrino temperature to the photon temperature is given by $(T_\nu/T_\gamma)^3 = g_\gamma/(g_\gamma + \frac{7}{8}g_e) = 4/11$.

In the early Universe when $\rho \approx \rho_r$, then $\dot{R} \propto 1/R$, so that $R \propto t^{1/2}$ and $Ht \rightarrow 1/2$ as $t \rightarrow 0$. The time-temperature relationship at very early times can then be found from the above equations:

$$t = \frac{2.42}{\sqrt{N(T)}} \left(\frac{1 \text{ MeV}}{kT} \right)^2 \text{ sec}. \quad (3.12)$$

At later times, since the energy density in radiation falls off as R^{-4} and the energy density in non-relativistic matter falls off as R^{-3} , the Universe eventually became matter dominated. The epoch of matter-radiation density equality is determined by equating the matter density at t_{eq} , $\rho_m = \Omega_0 \rho_c (R_0/R_{\text{eq}})^3$ to the radiation density, $\rho_r = (\pi^2/30)[2 + (21/4)(4/11)^{4/3}](kT_0)^4 (R_0/R_{\text{eq}})^4$ where T_0 is the present temperature of the microwave background (see below). Solving for $(R_0/R_{\text{eq}}) = 1 + z_{\text{eq}}$ gives

$$\begin{aligned} z_{\text{eq}} + 1 &= \Omega_0 h_0^2 / 4.2 \times 10^{-5} = 2.4 \times 10^4 \Omega_0 h_0^2; \\ kT_{\text{eq}} &= 5.6 \Omega_0 h_0^2 \text{ eV}; \\ t_{\text{eq}} &\approx 0.39 (\Omega_0 H_0^2)^{-1/2} (1 + z_{\text{eq}})^{-3/2} \\ &= 3.2 \times 10^{10} (\Omega_0 h_0^2)^{-2} \text{ sec}. \end{aligned} \quad (3.13)$$

Prior to this epoch the density was dominated by radiation (relativistic particles; see Eq. (3.10)), and at later epochs matter density dominated. Atoms formed at $z \approx 1300$, and by $z_{\text{dec}} \approx 1100$ the free electron density was low enough that space became essentially transparent to photons and matter and radiation were decoupled. These are the photons observed in the microwave background today.

The age of the Universe today, t_0 , is related to both the Hubble parameter and the value of Ω_0 (still assuming that $\Lambda = 0$). In the standard model, $t_0 \gg t_{\text{eq}}$ and we can write

$$t_0 = H_0^{-1} \int_0^1 \left(1 - \Omega_0 + \Omega_0 x^{-1} \right)^{-1/2} dx. \quad (3.14)$$

Constraints on t_0 yield constraints on the combination $\Omega_0 h_0^2$. For example, $t_0 \geq 13 \times 10^9 \text{ yr}$ implies that $\Omega_0 h_0^2 \leq 0.25$ for $h_0 \geq 0.5$.

or $\Omega_0 h_0^2 \leq 0.45$ for $h_0 \geq 0.4$, while $t_0 \geq 10 \times 10^9$ yr implies that $\Omega_0 h_0^2 \leq 0.8$ for $h_0 \geq 0.5$, or $\Omega_0 h_0^2 \leq 1.1$ for $h_0 \geq 0.4$.

The present temperature of the microwave background is $T_0 = 2.726 \pm 0.005$ K as measured by COBE [4], and the number density of photons $n_\gamma = (2\zeta(3)/\pi^2)(kT_0)^3 \approx 411 \text{ cm}^{-3}$. The energy density in photons (for which $g_\gamma = 2$) is $\rho_\gamma = (\pi^2/15)(kT_0)^4$. At the present epoch, $\rho_\gamma = 4.65 \times 10^{-34} \text{ g cm}^{-3} = 0.26 \text{ eV cm}^{-3}$. For nonrelativistic matter (such as baryons) today, the energy density is $\rho_B = m_B n_B$ with $n_B \propto R^{-3}$, so that for most of the history of the Universe n_B/s is constant. Today, the entropy density is related to the photon density by $s = (4/3)(\pi^2/30)[2 + (21/4)(4/11)](kT_0)^3 = 7.0 n_\gamma$. Big Bang nucleosynthesis calculations limit $\eta = n_B/n_\gamma$ to $2.8 \times 10^{-10} \leq \eta \leq 4.0 \times 10^{-10}$. The parameter η is also related to the portion of Ω in baryons

$$\Omega_B = 3.66 \times 10^7 \eta h_0^{-2} (T_0/2.726 \text{ K})^3, \quad (3.15)$$

so that $0.010 < \Omega_B h_0^2 < 0.015$, and hence the Universe cannot be closed by baryons.

References:

1. S. Weinberg, *Gravitation and Cosmology*, John Wiley and Sons (1972).
2. G. Börner, *The Early Universe: Facts and Fiction*, Springer-Verlag (1988).
3. E.W. Kolb and M.S. Turner, *The Early Universe*, Addison-Wesley (1990).
4. J.C. Mather *et al.*, *Astrophys. J.* **420**, 439 (1994). Error quoted here is 1σ .

4. DARK MATTER

Written September 1989 by R. Flores and K.A. Olive. Revised November 1991 and November 1993.

There is increasing evidence for the existence of large quantities of dark matter in the Universe. The most direct evidence comes from the astronomical observation of the motion of visible matter (stars and regions of neutral hydrogen gas) in galaxies. The observed velocities due to rotational motion in spiral galaxies are measured to be largely independent of the distance to the center of these galaxies [1] (they are said to have “flat rotational curves”). In the absence of any unseen component, we would expect that the velocity falls off with increasing distance, $v^2 \approx G_N M_{\text{vis}}/r$. In contrast, a flat rotation curve implies a total mass $M_{\text{tot}} \approx G_N^{-1} v_{\text{obs}}^2 r \approx 10^{11} M_{\odot} (v_{\text{obs}}/200 \text{ km s}^{-1})^2 (r/10 \text{ kpc})$ in excess of the visible mass M_{vis} . It can be inferred from these observations that there exists a dark matter component distributed in a (roughly) spherical halo about the galaxy. The dynamics of groups of galaxies and clusters, as well as the presence of very hot gas in elliptical galaxies require large quantities of unseen matter as well [2]. More recent observations of hot x-ray emitting gas in clusters of galaxies [3], and of gravitational lensing of high redshift background galaxies by foreground clusters [4] (which do not require the cluster to be relaxed), also indicate the presence of dark matter on cluster scales. In addition, theories of cosmological inflation predict that the density parameter of the Universe $\Omega_{\text{tot}} = 1$, whereas standard Big Bang nucleosynthesis requires $\Omega_{\text{baryon}} \lesssim 0.1$ [5]. This implies the existence of nonbaryonic dark matter. Further indirect evidence comes from our theoretical understanding of the growth of density perturbations as seeds for galaxy formation. Without the presence of dark matter, it is very difficult to reconcile the existence of galaxies (and quasars) at high redshifts with the new measurements by COBE of the anisotropy of the microwave background radiation [6]. Perturbations in baryon density can grow only after the time of recombination, *i.e.* when the baryons decouple from the microwave background. When $\Omega_{\text{tot}} = 1$ due to dark matter, matter domination occurs much earlier and dark matter perturbations grow for a longer period, thus avoiding a conflict with the magnitude of the microwave background anisotropy.

In our own galaxy, the distribution of the visible matter and its observed circular motion determine the local (solar neighborhood) dark matter density $\rho^{\text{DM}} \approx 0.3 \text{ GeV cm}^{-3}$ [7]. Regardless of the nature of the dark matter, it must behave as a collisionless gas, with a broad velocity distribution (typically assumed to be Maxwellian); $\langle v \rangle \approx \Delta v \approx 300 \text{ km s}^{-1}$.

We do not know the identity of the dark matter nor whether there is more than one type of dark matter. Baryons are difficult to conceal [8] and in the standard Big Bang model cannot make up all of the dark matter if $\Omega_{\text{tot}} = 1$. By the same token, by comparing the lower limit of $\Omega_{\text{baryon}} \gtrsim 0.01$ from nucleosynthesis [5], it is very likely that some of the baryons are dark. Though it is theoretically unlikely that galactic halos could be made of very dim objects, such as low-mass stars with masses $\lesssim 0.1 M_{\odot}$, recent gravitational microlensing searches for such objects may have a positive detection [9] (these objects are sometimes referred to as MACHOS, massive compact halo objects). There are several theoretical elementary particle candidates (WIMPs, weakly interacting massive particles) that could explain the existence of dark matter, of which the most commonly discussed are: a neutrino (if massive), a neutralino (from supersymmetry), and the axion (from the strong CP problem). These are summarized in Table 4.1.

Table 4.1: Dark Matter Candidates

Type	Candidate	Possible mass
Hot	neutrino	1–10 eV
Cold	neutralino; photino, Higgsino, or bino	20–350 GeV
Cold	axion	10^{-5} – 10^{-3} eV

Regardless of the exact identity of the dark matter (DM), its kinetic energy at the time when dark-matter domination begins

determines the subsequent evolution of the density perturbations that seed galactic and large structures [10]. If the dark matter is relativistic (hot dark matter, HDM) only the largest (supercluster) structures survive and they must fragment to form galactic structure, whereas if it is nonrelativistic (cold dark matter, CDM), structure on all scales is preserved. The large-scale distribution of matter in n -body simulations of a HDM-dominated universe is not compatible with observations (unless there are point-like density perturbations), whereas a flat CDM-dominated universe requires that the visible matter be predominantly concentrated in the denser regions of the DM distribution (biased galaxy formation). In point of fact, both the HDM- and CDM-dominated universes have some degree of difficulty with the size of the microwave background anisotropy measured by COBE. A mixture of cold and hot dark matter may provide a better solution to the problem of generating large scale structures. An example of such a mixture would be a ν_{τ} with a mass of order a few eV and a more massive neutralino.

For a cold dark matter particle species with equal particle (X) and antiparticle (\bar{X}) densities (except for the axions), its cosmological density at present is [11]

$$\Omega_X h^2 \approx 1.6 \times 10^{-10} N_F^{1/2} (T_X/T_\gamma)^3 \times \left(a + \frac{1}{12} b \langle v^2 \rangle_f \right)^{-1} \langle v^2 \rangle_f^{-1} \quad (4.1)$$

with a and b determined from the (velocity averaged) annihilation cross section, expanded in powers of momentum, $\langle v \sigma_{X\bar{X}} \rangle = a + \frac{1}{6} b \langle v^2 \rangle_f$, at freezeout temperature T_f ($\langle v^2 \rangle_f = 6T_f/M_X$) at which the X 's drop from thermal equilibrium (typically $T_f \approx \frac{1}{20} M_X$). In Eq. (4.1), N_F is the total number of relativistic degrees of freedom at T_f and (T_X/T_γ) is the ratio of the temperatures of X 's and photons at T_f . In the halo of our galaxy $\langle v^2 \rangle \approx 10^{-6}$, thus $\langle v \sigma_{X\bar{X}} \rangle_{\text{halo}}$ and Ω_X are closely related.

Several proposals or experiments exist to detect cold dark matter candidates. For the case of heavy ($M \gtrsim 1 \text{ GeV}$) particles, elastic scattering from nuclei would produce nuclear recoils with energies of $\gtrsim 1 \text{ keV}$, and several techniques have been proposed to detect these recoils. The expected collision rate for a target nucleus mass m_N is:

$$R = 4.3 \text{ kg}^{-1} \text{ day}^{-1} \left(\frac{1 \text{ GeV}^2}{m_N m_x} \right) \left(\frac{\sigma_{\text{el}}}{10^{-38} \text{ cm}^2} \right) \times \left(\frac{\rho^{\text{DM}}}{0.3 \text{ GeV cm}^{-3}} \right) \left(\frac{\langle v_E \rangle}{300 \text{ km s}^{-1}} \right), \quad (4.2)$$

where $\langle v_E \rangle$ is the average velocity at which they strike the detector. Since crossing symmetry relates σ_{el} to $\sigma_{X\bar{X}}$, R is closely related to Ω_X . Dirac neutrinos (and sneutrinos) with masses 0.012–4.7 TeV have already been excluded by searches done using double- β decay detectors [12]. Axions could be detected by their expected coherent conversion to microwave photons in a tuned cavity. Products of DM annihilation in the halo (*e.g.*, cosmic ray \bar{p} 's, e^+ 's, γ 's) and the core of the Sun (ν 's) would indirectly signal the existence of particle DM. The absence of a signal in high energy solar- ν searches using underground detectors rules out sneutrinos whereas cosmic ray searches do not constrain theory so far. Experimental limits concerning a number of dark matter candidates are given in the Full Listings. See the index under “Dark matter limits.”

Recent LEP results when combined with the above experimental constraints now completely eliminate massive 4th-generation neutrinos or sneutrinos as dark matter candidates. Sneutrinos and additional Dirac and Majorana neutrinos with masses $\lesssim 40 \text{ GeV}$ are excluded by LEP. This alone eliminates a Majorana neutrino, since the relic abundance for the neutrinos with masses $\gtrsim 40 \text{ GeV}$ would be $\Omega h^2 \lesssim 2 \times 10^{-3}$, making them cosmologically uninteresting. In the case of Dirac neutrinos, if there were a density asymmetry between ν and $\bar{\nu}$, it would in principle be possible to have a cosmologically interesting density even though $m_\nu > 40 \text{ GeV}$. However, as described

in the previous paragraph, Dirac neutrinos (and sneutrinos) below 4.7 TeV are ruled out by direct cold dark matter searches.

References:

1. S.M. Faber and J.J. Gallagher, *Ann. Rev. Astron. Astrophys.* **17**, 135 (1979).
2. D. Fabricant and P. Gorenstein, *Astrophys. J.* **267**, 535 (1983); G.C. Stewart, C.R. Canizares, A.C. Fabian, and P.E.J. Nilsen, *Astrophys. J.* **278**, 53 (1984) and references therein.
3. R. Mushotzky, in *Proceedings Texas/PASCOS 92: Relativistic Astrophysics and Particle Cosmology*, ed. C.W. Akerlof and M. Srednicki (*Ann. N.Y. Acad. Sci.* **688**, 184 (1993)); J.S. Mulchaey, D.S. Davis, R.F. Mushotzky, and D. Burstein, preprint (1993).
4. J.A. Tyson, F. Valdes, and R.A. Wenk, *Astrophys. J.* **349**, L1 (1990).
5. T.P. Walker, G. Steigman, D.N. Schramm, K.A. Olive, and H.-S. Kang, *Astrophys. J.* **376**, 51 (1991).
6. E.L. Wright *et al.*, *Astrophys. J.* **396**, L13 (1992).
7. R. Flores, *Phys. Lett.* **B215**, 73 (1988).
8. D.J. Hegyi and K.A. Olive, *Phys. Lett.* **126B**, 28 (1983); *Astrophys. J.* **303**, 56 (1986).
9. C. Alcock *et al.*, *Nature* **365**, 621 (1993); E. Aubourg *et al.*, *Nature* **365**, 623 (1993).
10. See: J.R. Primack in *Enrico Fermi Course 92*, ed. N. Cabibbo (North Holland, Amsterdam, 1987), p. 137; V. Trimble, *Ann. Rev. Astron. Astrophys.* **25** (1987) 425; J. Primack, D. Seckel, and B. Sadulek, *Ann. Rev. Nucl. and Part. Sci.* **38**, 751 (1988); *Dark Matter*, ed. M. Srednicki (North-Holland, Amsterdam, 1989).
11. J. Ellis, J. Hagelin, D.V. Nanopoulos, K.A. Olive and M. Srednicki, *Nucl. Phys.* **B238**, 453 (1984).
12. D. Caldwell *et al.*, *Phys. Rev. Lett.* **61**, 510 (1988).

5. INTERNATIONAL SYSTEM OF UNITS (SI)

Sec "The International System of Units (SI)," NIST special publication 330, B.N. Taylor, ed. (USGPO, Washington, DC, 1991).

Physical quantity	Name of unit	Symbol
<i>Base units</i>		
length	meter	m
mass	kilogram	kg
time	second	s
electric current	ampere	A
thermodynamic temperature	kelvin	K
amount of substance	mole	mol
luminous intensity	candela	cd
<i>Supplementary units</i>		
plane angle	radian	rad
solid angle	steradian	sr
<i>Derived units with special names</i>		
frequency	hertz	Hz
energy	joule	J
force	newton	N
pressure	pascal	Pa
power	watt	W
electric charge	coulomb	C
electric potential	volt	V
electric resistance	ohm	Ω
electric conductance	siemens	S
electric capacitance	farad	F
magnetic flux	weber	Wb
inductance	henry	H
magnetic flux density	tesla	T
luminous flux	lumen	lm
illuminance	lux	lx
celsius temperature	degree celsius	$^{\circ}\text{C}$
activity (of a radioactive source)*	becquerel	Bq
absorbed dose (of ionizing radiation)*	gray	Gy
dose equivalent*	sievert	Sv

SI prefixes

10^{24}	yotta	(Y)
10^{21}	zetta	(Z)
10^{18}	exa	(E)
10^{15}	peta	(P)
10^{12}	tera	(T)
10^9	giga	(G)
10^6	mega	(M)
10^3	kilo	(k)
10^2	hecto	(h)
10	deca	(da)
10^{-1}	deci	(d)
10^{-2}	centi	(c)
10^{-3}	milli	(m)
10^{-6}	micro	(μ)
10^{-9}	nano	(n)
10^{-12}	pico	(p)
10^{-15}	femto	(f)
10^{-18}	atto	(a)
10^{-21}	zepto	(z)
10^{-24}	yocto	(y)

*See our section 13, on "Radioactivity and radiation protection," p. 1268.

6. ATOMIC AND NUCLEAR PROPERTIES OF MATERIALS

Table 6.1. Table revised June 1994. Gases are evaluated at 20°C, 1 atm, (in parentheses) or at STP [square brackets].

Material	Z	A	Nuclear ^a total cross section σ_T [barn]	Nuclear ^b inelastic cross section σ_I [barn]	Nuclear ^c collision length λ_T [g/cm ²]	Nuclear ^c interaction length λ_I [g/cm ²]	$dE/dx _{\min}^d$ [MeV] [g/cm ²] () is for gas	Radiation length ^e X_0 [g/cm ²] [cm] () is for gas	Density ^f [g/cm ³] [g/l] () is for gas	Refractive index n^f ($n-1$) $\times 10^6$ for gas	
H ₂ gas	1	1.01	0.0387	0.033	43.3	50.8	(4.103)	61.28	865	(0.0838)[0.090]	[140]
H ₂ (B.C., 26K)	1	1.01	0.0387	0.033	43.3	50.8	4.045	61.28	865	0.0708	1.112
D ₂	1	2.01	0.073	0.061	45.7	54.7	(2.052)	122.6	757	0.162[0.177]	1.128
He	2	4.00	0.133	0.102	49.9	65.1	(1.937)	94.32	755	0.125[0.178]	1.024[35]
Li	3	6.94	0.211	0.157	54.6	73.4	1.639	82.76	155	0.534	—
Be	4	9.01	0.268	0.199	55.8	75.2	1.594	65.19	35.3	1.848	—
C	6	12.01	0.331	0.231	60.2	86.3	1.745	42.70	18.8	2.265 ^g	—
N ₂	7	14.01	0.379	0.265	61.4	87.8	(1.825)	37.99	47.0	0.808[1.25]	1.205[300]
O ₂	8	16.00	0.420	0.292	63.2	91.0	(1.801)	34.24	30.0	1.14[1.43]	1.22[266]
Ne	10	20.18	0.507	0.347	66.1	96.6	(1.724)	28.94	24.0	1.207[0.900]	1.092[67]
Al	13	26.98	0.634	0.421	70.6	106.4	1.615	24.01	8.9	2.70	—
Si	14	28.09	0.660	0.440	70.6	106.0	1.664	21.82	9.36	2.33	—
Ar	18	39.95	0.868	0.566	76.4	117.2	(1.519)	19.55	14.0	1.40[1.782]	1.233[283]
Ti	22	47.88	0.995	0.637	79.9	124.9	1.476	16.17	3.56	4.54	—
Fe	26	55.85	1.120	0.703	82.8	131.9	1.451	13.84	1.76	7.87	—
Cu	29	63.55	1.232	0.782	85.6	134.9	1.403	12.86	1.43	8.96	—
Ge	32	72.59	1.365	0.858	88.3	140.5	1.371	12.25	2.30	5.323	—
Sn	50	118.69	1.967	1.21	100.2	163	1.264	8.82	1.21	7.31	—
Xe	54	131.29	2.120	1.29	102.8	169	(1.255)	8.48	2.77	3.057[5.858]	[705]
W	74	183.85	2.767	1.65	110.3	185	1.145	6.76	0.35	19.3	—
Pt	78	195.08	2.861	1.708	113.3	189.7	1.129	6.54	0.305	21.45	—
Pb	82	207.19	2.960	1.77	116.2	194	1.123	6.37	0.56	11.35	—
U	92	238.03	3.378	1.98	117.0	199	1.082	6.00	≈0.32	≈18.95	—
Air, (20°C, 1 atm.), [STP]					62.0	90.0	(1.815)	36.66	[30420]	(1.205)[1.29]	(273)[293]
H ₂ O					60.1	84.9	1.991	36.08	36.1	1.00	1.33
CO ₂					62.4	90.5	(1.819)	36.2	[18310]	[1.977]	[410]
Shielding concrete ^h					67.4	99.9	1.711	26.7	10.7	2.5	—
Borosilicate glass (Pyrex) ^l					66.2	97.6	1.695	28.3	12.7	2.23	1.474
SiO ₂ (fused quartz) ^m					67.0	99.2	1.697	27.05	11.7	2.32 ^m	1.458
Methane (CH ₄)					54.7	74.0	(2.417)	46.5	[64850]	0.423[0.717]	[444]
Ethane (C ₂ H ₆)					55.73	75.71	(2.304)	45.66	[34035]	0.509(1.356) ⁿ	(1.038) ⁿ
Propane (C ₃ H ₈)					—	—	(2.262)	—	—	(1.879)	—
Isobutane ((CH ₃) ₂ CHCH ₃)					56.3	77.4	(2.239)	45.2	[16930]	[2.67]	[1900]
Octane, liquid (CH ₃ (CH ₂) ₆ CH ₃)					—	—	2.123	—	—	0.703	—
Paraffin wax (CH ₃ (CH ₂) _n CH ₃ , (n) ≈ 25)					—	—	2.087	—	—	0.93	—
Nylon, type 6					—	—	1.974	—	—	1.14	—
Polycarbonate (Lexan)					—	—	1.886	—	—	1.200	—
Polyethylene terephthlate (Mylar) (C ₅ H ₄ O ₂)					60.2	85.7	1.848	39.95	28.7	1.39	—
Polyethylene (monomer CH ₂ =CH ₂)					56.9	78.8	2.076	44.8	≈47.9	0.92–0.95	—
Polyimide film (Kapton)					—	—	1.820	—	—	1.420	—
Polymethylmethacralate (Lucite, Plexiglas) (monomer (CH ₂ =C(CH ₃)CO ₂ CH ₃))					59.2	83.6	1.929	40.55	≈34.4	1.16–1.20	≈1.49
Polystyrene, scintillator (monomer C ₆ H ₅ CH=CH ₂)					58.4	82.0	1.936	43.8	42.4	1.032	1.581
Polytetrafluoroethylene (Teflon) (monomer CF ₂ =CF ₂)					—	—	1.671	—	—	2.20	—
Polyvinyltolulene, scintillator (monomer 2-CH ₃ C ₆ H ₄ CH=CH ₂)					—	—	1.956	—	—	1.032	—
Barium fluoride (BaF ₂)					92.1	146	1.303	9.91	2.05	4.89	1.56
Bismuth germanate (BGO) (Bi ₄ Ge ₃ O ₁₂)					97.4	156	1.251	7.98	1.12	7.1	2.15
Cesium iodide (CsI)					—	—	1.243	—	—	4.51	—
Lithium fluoride (LiF)					62.00	88.24	1.614	39.25	14.91	2.632	1.392
Sodium fluoride (NaF)					66.78	97.57	1.69	29.87	11.68	2.558	1.336
Sodium iodide (NaI)					94.8	152	1.305	9.49	2.59	3.67	1.775
Silica Aerogel ^o					65.5	95.7	1.83	29.85	≈150	0.1–0.3	1.0+0.25ρ
NEMA G10 plate ^p					62.6	90.2	1.87	33.0	19.4	1.7	—

Material	Dielectric constant ($\kappa = \epsilon/\epsilon_0$) () is $(\kappa-1) \times 10^6$ for gas	Young's modulus [10^6 psi]	Coeff. of thermal expansion [10^{-6} cm/cm-°C]	Specific heat [cal/g-°C]	Electrical resistivity [$\mu\Omega$ cm(@°C)]	Thermal conductivity [cal/cm-°C-sec]
H ₂	(253.9)	—	—	—	—	—
He	(64)	—	—	—	—	—
Li	—	—	56	0.86	8.55(0°)	0.17
Be	—	37	12.4	0.436	5.885(0°)	0.38
C	—	0.7	0.6–4.3	0.165	1375(0°)	0.057
N ₂	(548.5)	—	—	—	—	—
O ₂	(495)	—	—	—	—	—
Ne	(127)	—	—	—	—	—
Al	—	10	23.9	0.215	2.65(20°)	0.53
Si	11.9	16	2.8–7.3	0.162	—	0.20
Ar	(517)	—	—	—	—	—
Ti	—	16.8	8.5	0.126	50(0°)	—
Fe	—	28.5	11.7	0.11	9.71(20°)	0.18
Cu	—	16	16.5	0.092	1.67(20°)	0.94
Ge	16.0	—	5.75	0.073	—	0.14
Sn	—	6	20	0.052	11.5(20°)	0.16
Xe	—	—	—	—	—	—
W	—	50	4.4	0.032	5.5(20°)	0.48
Pt	—	21	8.9	0.032	9.83(0°)	0.17
Pb	—	2.6	29.3	0.038	20.65(20°)	0.083
U	—	—	36.1	0.028	29(20°)	0.064

σ_T , σ_I , λ_T , and λ_I are energy dependent. Values quoted apply to high energy range given in footnote a or b, where energy dependence is weak.

- a. σ_{total} at 80–240 GeV for neutrons ($\approx \sigma$ for protons) from Murthy *et al.*, Nucl. Phys. **B92**, 269 (1975). This scales approximately as $A^{0.77}$.
- b. $\sigma_{\text{inelastic}} = \sigma_{\text{total}} - \sigma_{\text{elastic}} - \sigma_{\text{quasielastic}}$; for neutrons at 60–375 GeV from Roberts *et al.*, Nucl. Phys. **B159**, 56 (1979). For protons and other particles, see Carroll *et al.*, Phys. Lett. **80B**, 319 (1979); note that $\sigma_I(p) \approx \sigma_I(n)$. σ_I scales approximately as $A^{0.71}$.
- c. Mean free path between collisions (λ_T) or inelastic interactions (λ_I), calculated from $\lambda = A/(N \times \sigma)$, where N is Avogadro's number.
- d. For minimum-ionizing heavy particles (muons, pions, protons, *etc.*). Minimum dE/dx calculated in 1994, using density effect correction coefficients from R. M. Sternheimer, M. J. Berger, and S. M. Seltzer, Atomic Data and Nuclear Data Tables **30**, 261–271 (1984). For electrons and positrons see S.M. Seltzer and M.J. Berger, Int. J. Appl. Radiat. **35**, 665–676 (1984). Ionization energy loss is discussed in Sec. 10.
- e. From Y.S. Tsai, Rev. Mod. Phys. **46**, 815 (1974); X_0 data for all elements up to uranium may be found here. Corrections for molecular binding applied for H₂ and D₂. Parentheses refer to gaseous form at STP (0°C, 1 atm.).
- f. Values for solids, or the liquid phase at boiling point, except as noted. Refractive index given for sodium D line.
- g. For pure graphite; industrial graphite density may vary 2.1–2.3 g/cm³.
- h. Standard shielding blocks, typical composition O₂ 52%, Si 32.5%, Ca 6%, Na 1.5%, Fe 2%, Al 4%, plus reinforcing iron bars. The attenuation length, $\ell = 115 \pm 5$ g/cm², is also valid for earth (typical $\rho = 2.15$), from CERN-LRL-RHEL Shielding exp., UCRL-17841 (1968).
- i. Density may vary about $\pm 3\%$, depending on operating conditions.
- j. Values for typical working conditions with H₂ target: 50 mole percent, 29°K, 7 atm.
- k. Typical scintillator; *e.g.*, PILOT B and NE 102A have an atomic ratio H/C = 1.10.
- l. Main components: 80% SiO₂ + 12% B₂O₃ + 5% Na₂O.
- m. For typical fused quartz; density may vary. The specific gravity of crystalline quartz is 2.64.
- n. Solid ethane density at –60°C; gaseous refractive index at 0°C, 546 mm pressure.
- o. $n(\text{SiO}_2) + 2n(\text{H}_2\text{O})$ used in Čerenkov counters, $\rho =$ density in g/cm³. From M. Cantin *et al.*, Nucl. Instr. and Meth. **118**, 177 (1974).
- p. G10-plate, typical 60% SiO₂ and 40% epoxy.

7. PERIODIC TABLE OF THE ELEMENTS

Table 7.1. Revised 1993. The atomic number (top left) is the number of protons in the nucleus. The atomic mass (bottom) is weighted by isotopic abundances in the Earth's surface. Atomic masses are relative to the mass of the carbon-12 isotope, defined to be exactly 12 unified atomic mass units (u). Errors range from 1 to 9 in the last digit quoted. Relative isotopic abundances often vary considerably, both in natural and commercial samples. A number in parentheses is the mass of the longest-lived isotope of that element—no stable isotope exists. However, although Th, Pa, and U have no stable isotopes, they do have characteristic terrestrial compositions, and meaningful weighted masses can be given. Adapted from "Atomic Weights of the Elements 1991," Pure and Applied Chemistry **64**, 1519 (1992), and G. Audi and A.H. Wapstra, "The 1993 Mass Evaluation," Nucl. Phys. **A565**, 1 (1993). The names for elements 104–109 have been adopted by the American Chemical Society Nomenclature Committee, and recommended to IUPAC.

IA		VIIA										VIIIA					
1	2											10	11				
Hydrogen	Helium											Neon	Argon				
1.00794	4.002602											20.1797	39.948				
IIA		VIA										VIIA					
3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18		
Lithium	Beryllium	Boron	Carbon	Nitrogen	Oxygen	Fluorine	Neon	Silicon	Phosphorus	Sulfur	Chlorine	Argon	Potassium	Calcium	Scandium		
6.941	9.012182	10.811	12.011	14.00674	15.9994	18.9984032	20.1797	28.0855	30.973762	32.066	35.4527	39.948	39.0983	40.078	44.955910		
11	12	VIII										19	20				
Sodium	Magnesium <th colspan="10"></th> <th>Potassium</th> <th>Calcium</th>											Potassium	Calcium				
22.989768	24.3050											39.0983	40.078				
IIIB		IVB		VB		VIB		VIIB		VIII		IIB					
13	14	15	16	17	18	19	20	21	22	23	24	25	26				
Aluminum	Silicon	Phosphorus	Sulfur	Chlorine	Argon	Cobalt	Nickel	Copper	Zinc	Gallium	Germanium	Arsenic	Selenium				
26.981539	28.0855	30.973762	32.066	35.4527	39.948	58.93320	58.93320	63.546	65.39	69.723	72.61	74.92159	78.96				
19	20	21	22	23	24	25	26	27	28	29	30	31	32				
Potassium	Calcium	Scandium	Titanium	Vanadium	Chromium	Manganese	Iron	Cobalt	Nickel	Copper	Zinc	Gallium	Germanium				
39.0983	40.078	44.955910	47.88	50.9415	51.9961	54.93805	55.847	58.93320	58.93320	63.546	65.39	69.723	72.61				
37	38	39	40	41	42	43	44	45	46	47	48	49	50				
Rubidium	Strontium	Yttrium	Zirconium	Niobium	Molybdenum	Technetium	Ruthenium	Rhodium	Palladium	Silver	Cadmium	Indium	Tin				
85.4678	87.62	88.90585	91.224	92.90638	95.94	(97.9072)	101.07	102.90550	106.42	107.8682	112.411	114.818	118.710				
55	56	57–71	72	73	74	75	76	77	78	79	80	81	82				
Cesium	Barium	Lanthanides	Hafnium	Tantalum	Tungsten	Rhenium	Osmium	Iridium	Platinum	Gold	Mercury	Thallium	Lead				
132.90543	137.327		178.49	180.9479	183.84	186.207	192.22	192.22	195.08	196.96654	200.59	204.3833	207.2				
87	88	89–103	104 (Rf)	105 (Ha)	106 (Sg)	107 (Nh)	108 (Hs)	109 (Mt)									
Francium	Radium	Actinides	Rutherfordium	Dubnium	Seaborgium	Nielsbohrium	Hassium	Meitnerium									
(223.0197)	(226.0254)		(261.1089)	(262.1144)	(263.1186)	(262.1231)	(265.1306)	(266.1378)									
Lanthanide series		Actinide series															
57	89	90	91	92	93	94	95	96	97	98	99	100	101	102	103	104	
Lanthanum	Actinium	Thorium	Protactinium	Uranium	Neptunium	Plutonium	Americium	Curium	Berkelium	Californium	Einsteinium	Fermium	Mendelevium	Nobelium	Lawrencium		
138.9055	227.0278	232.0381	231.03588	238.0289	(237.0482)	(244.0642)	(243.0614)	(247.0703)	(247.0703)	(251.0796)	(252.0830)	(257.0951)	(258.0984)	(259.1011)	(262.1098)		

8. ELECTRONIC STRUCTURE OF THE ELEMENTS

Table 8.1. Reviewed 1993 by W.C. Martin, NIST. The electronic configurations and ionization energies here are taken from the *CRC Handbook of Chemistry and Physics*, 74th Edition, ed. D.R. Lide (CRC Press, Boca Raton, FL, 1993). The electron configuration for, say, iron indicates an argon electronic core (see argon) plus six 3d electrons and two 4s electrons. The ionization energy is the least energy necessary to remove to infinity one electron from an *atom* of the element.

Element		Electron configuration (3d ⁵ = five 3d electrons, etc.)	Ground state $2S+1L_J$	Ionization energy (eV)
1	H Hydrogen	1s	$^2S_{1/2}$	13.5984
2	He Helium	1s ²	1S_0	24.5874
3	Li Lithium	(He) 2s	$^2S_{1/2}$	5.3917
4	Be Beryllium	(He) 2s ²	1S_0	9.3226
5	B Boron	(He) 2s ² 2p	$^2P_{1/2}$	8.2980
6	C Carbon	(He) 2s ² 2p ²	3P_0	11.2603
7	N Nitrogen	(He) 2s ² 2p ³	$^4S_{3/2}$	14.5341
8	O Oxygen	(He) 2s ² 2p ⁴	3P_2	13.6181
9	F Fluorine	(He) 2s ² 2p ⁵	$^2P_{3/2}$	17.4228
10	Ne Neon	(He) 2s ² 2p ⁶	1S_0	21.5645
11	Na Sodium	(Ne) 3s	$^2S_{1/2}$	5.1391
12	Mg Magnesium	(Ne) 3s ²	1S_0	7.6462
13	Al Aluminum	(Ne) 3s ² 3p	$^2P_{1/2}$	5.9858
14	Si Silicon	(Ne) 3s ² 3p ²	3P_0	8.1517
15	P Phosphorus	(Ne) 3s ² 3p ³	$^4S_{3/2}$	10.4867
16	S Sulfur	(Ne) 3s ² 3p ⁴	3P_2	10.3600
17	Cl Chlorine	(Ne) 3s ² 3p ⁵	$^2P_{3/2}$	12.9676
18	Ar Argon	(Ne) 3s ² 3p ⁶	1S_0	15.7596
19	K Potassium	(Ar) 4s	$^2S_{1/2}$	4.3407
20	Ca Calcium	(Ar) 4s ²	1S_0	6.1132
21	Sc Scandium	(Ar) 3d 4s ²	$^2D_{3/2}$	6.5614
22	Ti Titanium	(Ar) 3d ² 4s ²	3F_2	6.8282
23	V Vanadium	(Ar) 3d ³ 4s ²	$^4F_{3/2}$	6.7463
24	Cr Chromium	(Ar) 3d ⁵ 4s	7S_3	6.7666
25	Mn Manganese	(Ar) 3d ⁵ 4s ²	$^6S_{5/2}$	7.4340
26	Fe Iron	(Ar) 3d ⁶ 4s ²	5D_4	7.9024
27	Co Cobalt	(Ar) 3d ⁷ 4s ²	$^4F_{9/2}$	7.8810
28	Ni Nickel	(Ar) 3d ⁸ 4s ²	3F_4	7.6398
29	Cu Copper	(Ar) 3d ¹⁰ 4s	$^2S_{1/2}$	7.7264
30	Zn Zinc	(Ar) 3d ¹⁰ 4s ²	1S_0	9.3941
31	Ga Gallium	(Ar) 3d ¹⁰ 4s ² 4p	$^2P_{1/2}$	5.9993
32	Ge Germanium	(Ar) 3d ¹⁰ 4s ² 4p ²	3P_0	7.900
33	As Arsenic	(Ar) 3d ¹⁰ 4s ² 4p ³	$^4S_{3/2}$	9.8152
34	Se Selenium	(Ar) 3d ¹⁰ 4s ² 4p ⁴	3P_2	9.7524
35	Br Bromine	(Ar) 3d ¹⁰ 4s ² 4p ⁵	$^2P_{3/2}$	11.8138
36	Kr Krypton	(Ar) 3d ¹⁰ 4s ² 4p ⁶	1S_0	13.9996
37	Rb Rubidium	(Kr) 5s	$^2S_{1/2}$	4.1771
38	Sr Strontium	(Kr) 5s ²	1S_0	5.6948
39	Y Yttrium	(Kr) 4d 5s ²	$^2D_{3/2}$	6.217
40	Zr Zirconium	(Kr) 4d ² 5s ²	3F_2	6.6339
41	Nb Niobium	(Kr) 4d ⁴ 5s	$^6D_{1/2}$	6.7589
42	Mo Molybdenum	(Kr) 4d ⁵ 5s	7S_3	7.0924
43	Tc Technetium	(Kr) 4d ⁵ 5s ²	$^6S_{5/2}$	7.28
44	Ru Ruthenium	(Kr) 4d ⁷ 5s	5F_5	7.3605
45	Rh Rhodium	(Kr) 4d ⁸ 5s	$^4F_{9/2}$	7.4589
46	Pd Palladium	(Kr) 4d ¹⁰	1S_0	8.3369
47	Ag Silver	(Kr) 4d ¹⁰ 5s	$^2S_{1/2}$	7.5762
48	Cd Cadmium	(Kr) 4d ¹⁰ 5s ²	1S_0	8.9937

49	In	Indium	(Kr)4d ¹⁰ 5s ² 5p			² P _{1/2}	5.7864
50	Sn	Tin	(Kr)4d ¹⁰ 5s ² 5p ²			³ P ₀	7.3438
51	Sb	Antimony	(Kr)4d ¹⁰ 5s ² 5p ³			⁴ S _{3/2}	8.64
52	Te	Tellurium	(Kr)4d ¹⁰ 5s ² 5p ⁴			³ P ₂	9.0096
53	I	Iodine	(Kr)4d ¹⁰ 5s ² 5p ⁵			² P _{3/2}	10.4513
54	Xe	Xenon	(Kr)4d ¹⁰ 5s ² 5p ⁶			¹ S ₀	12.1299
55	Cs	Cesium	(Xe) 6s			² S _{1/2}	3.8939
56	Ba	Barium	(Xe) 6s ²			¹ S ₀	5.2117
57	La	Lanthanum	(Xe) 5d 6s ²			² D _{3/2}	5.5770
58	Ce	Cerium	(Xe)4f 5d 6s ²			¹ G ₄	5.5387
59	Pr	Praseodymium	(Xe)4f ³ 6s ²	L		⁴ I _{9/2}	5.464
60	Nd	Neodymium	(Xe)4f ⁴ 6s ²	a		⁵ I ₄	5.5250
61	Pm	Promethium	(Xe)4f ⁵ 6s ²	a		⁶ H _{5/2}	5.55
62	Sm	Samarium	(Xe)4f ⁶ 6s ²	n		⁷ F ₀	5.6437
63	Eu	Europium	(Xe)4f ⁷ 6s ²	t		⁸ S _{7/2}	5.6704
64	Gd	Gadolinium	(Xe)4f ⁷ 5d 6s ²	h		⁹ D ₂	6.1500
65	Tb	Terbium	(Xe)4f ⁹ 6s ²	a		⁶ H _{15/2}	5.8639
66	Dy	Dysprosium	(Xe)4f ¹⁰ 6s ²	n		⁵ I ₈	5.9389
67	Ho	Holmium	(Xe)4f ¹¹ 6s ²	i		⁴ I _{15/2}	6.0216
68	Er	Erbium	(Xe)4f ¹² 6s ²	d		³ H ₆	6.1078
69	Tm	Thulium	(Xe)4f ¹³ 6s ²	e		² F _{7/2}	6.1843
70	Yb	Ytterbium	(Xe)4f ¹⁴ 6s ²	s		¹ S ₀	6.2542
71	Lu	Lutetium	(Xe)4f ¹⁴ 5d 6s ²			² D _{3/2}	5.4259
72	Hf	Hafnium	(Xe)4f ¹⁴ 5d ² 6s ²	T		³ F ₂	6.8251
73	Ta	Tantalum	(Xe)4f ¹⁴ 5d ³ 6s ²	r	e	⁴ F _{3/2}	7.89
74	W	Tungsten	(Xe)4f ¹⁴ 5d ⁴ 6s ²	a	l	⁵ D ₀	7.98
75	Re	Rhenium	(Xe)4f ¹⁴ 5d ⁵ 6s ²	n	e	⁶ S _{5/2}	7.88
76	Os	Osmium	(Xe)4f ¹⁴ 5d ⁶ 6s ²	s	m	⁵ D ₄	8.7
77	Ir	Iridium	(Xe)4f ¹⁴ 5d ⁷ 6s ²	i	e	⁴ F _{9/2}	9.1
78	Pt	Platinum	(Xe)4f ¹⁴ 5d ⁹ 6s	t	n	³ D ₃	9.0
79	Au	Gold	(Xe)4f ¹⁴ 5d ¹⁰ 6s	i	t	² S _{1/2}	9.2257
80	Hg	Mercury	(Xe)4f ¹⁴ 5d ¹⁰ 6s ²	o	s	¹ S ₀	10.4375
81	Tl	Thallium	(Xe)4f ¹⁴ 5d ¹⁰ 6s ² 6p			² P _{1/2}	6.1083
82	Pb	Lead	(Xe)4f ¹⁴ 5d ¹⁰ 6s ² 6p ²			³ P ₀	7.4167
83	Bi	Bismuth	(Xe)4f ¹⁴ 5d ¹⁰ 6s ² 6p ³			⁴ S _{3/2}	7.289
84	Po	Polonium	(Xe)4f ¹⁴ 5d ¹⁰ 6s ² 6p ⁴			³ P ₂	8.4167
85	At	Astatine	(Xe)4f ¹⁴ 5d ¹⁰ 6s ² 6p ⁵			² P _{3/2}	
86	Rn	Radon	(Xe)4f ¹⁴ 5d ¹⁰ 6s ² 6p ⁶			¹ S ₀	10.7485
87	Fr	Francium	(Rn) 7s			² S _{1/2}	
88	Ra	Radium	(Rn) 7s ²			¹ S ₀	5.2789
89	Ac	Actinium	(Rn) 6d 7s ²			² D _{3/2}	5.17
90	Th	Thorium	(Rn) 6d ² 7s ²			³ F ₂	6.08
91	Pa	Protactinium	(Rn)5f ² 6d 7s ²	A		⁴ K _{11/2}	5.89
92	U	Uranium	(Rn)5f ³ 6d 7s ²	c		⁵ L ₆	6.1941
93	Np	Neptunium	(Rn)5f ⁴ 6d 7s ²	t		⁶ L _{11/2}	6.2657
94	Pu	Plutonium	(Rn)5f ⁶ 7s ²	i		⁷ F ₀	6.06
95	Am	Americium	(Rn)5f ⁷ 7s ²	n		⁸ S _{7/2}	5.993
96	Cm	Curium	(Rn)5f ⁷ 6d 7s ²	i		⁹ D ₂	6.02
97	Bk	Berkelium	(Rn)5f ⁹ 7s ²	d		⁶ H _{15/2}	6.23
98	Cf	Californium	(Rn)5f ¹⁰ 7s ²	e		⁵ I ₈	6.30
99	Es	Einsteinium	(Rn)5f ¹¹ 7s ²	s		⁴ I _{15/2}	6.42
100	Fm	Fermium	(Rn)5f ¹² 7s ²			³ H ₆	6.50
101	Md	Mendelevium	(Rn)5f ¹³ 7s ²			² F _{7/2}	6.58
102	No	Nobelium	(Rn)5f ¹⁴ 7s ²			¹ S ₀	6.65
103	Lr	Lawrencium	(Rn)5f ¹⁴ 6d 7s ² ?			² D _{3/2} ?	
104	Rf	Rutherfordium	(Rn)5f ¹⁴ 6d ² 7s ² ?				

9. HIGH-ENERGY COLLIDER PARAMETERS: e^+e^- Colliders (I)

The numbers here were received from representatives of the colliders in 1993 or early 1994. Quantities are, where appropriate, r.m.s. H and V indicate horizontal and vertical directions. Many of the numbers of course change over the lifetime of a collider; only the latest values are given here.

	SPEAR (SLAC)	DORIS (DESY)	CESR [CESR+ (phase2)] (Cornell)	PETRA (DESY)	PEP (SLAC)
Physics start date	1972	1973	1979 [1995]	1978	1980
Physics end date	1990	—	—	1986	1990
Maximum beam energy (GeV)	4	5.6	6	23.4	15
Luminosity ($10^{30}\text{cm}^{-2}\text{s}^{-1}$)	10 at 3 GeV	33 at 5.3 GeV	290 at 5.3 GeV [600 in 1995]	24 at 17.5 GeV	60
Time between collisions (μs)	0.75	0.965	0.36 [0.028 in 1995]	3.8	2.44
Crossing half angle (μ rad)	0	0	0 [2500 in 1995]	0	0
Energy spread (units 10^{-3})	1	1.2 at 5 GeV	0.6 at 5.3 GeV	1.1 at 17.5 GeV	1
Bunch length (cm)	$\sigma_z \approx 4$	$\sigma \sim 2$ at 5 GeV	1.7	$\sigma \sim 1.3$ at 17.5 GeV	$\sigma_z = 2$
Beam radius (10^{-6} m)	$H: 700$ $V: 50$	$H: 740$ $V: \sim 30$ } at 5 GeV	$H: 500$ $V: 11$	$H: 430$ $V: 13$ } at 17.5 GeV	$H: 340$ $V: 14$
Free space at interaction point (m)	± 2.5	± 1.2	± 2.2 (± 0.6 to REC quads)	± 4.5	± 3.7
Luminosity lifetime (hr)	≈ 3	1.0–1.5	3–4	4 at 17.5 GeV	4
Filling time (min)	15	≈ 15	10	20	15
Acceleration period (s)	≤ 100	—	—	—	≤ 100
Injection energy (GeV)	2.5	up to 5.6	6	7	15
Transverse emittance ($10^{-9}\pi$ rad-m)	$H \approx 430$	$H: 500$ $V: 5-50$ } at 5 GeV	$H: 240$ $V: 8$	$H: 140$ $V: 2$	$H \approx 120$
β^* , amplitude function at interaction point (m)	$H: 1.2$ $V: 0.08$	$H: 0.59/12.3$ $V: 0.04/0.79$	$H: 1.0$ $V: 0.018$	$H: 1.3$ $V: 0.08$	$H: 1.0$ $V: 0.05$
Beam-beam tune shift per crossing (units 10^{-4})	300	≤ 280 (space charge limit at 5.3 GeV)	420 [300 in 1995]	$H: 160$ $V: 400$ } at 17.5 GeV	550
RF frequency (MHz)	358	500	500	500	352
Particles per bunch (units 10^{10})	15	27	24 [17 in 1995]	26	35
Bunches per ring per species	1	1	7 [27 in 1995]	2	3
Average beam current per species (mA)	30	45 at 5.3 GeV	110 [300 in 1995]	11 at 17.5 GeV	21
Circumference (km)	0.234	0.2892	0.768	2.304	2.2
Interaction regions	2	2	1	4	1
Utility insertions	18	10	2	4	5
Magnetic length of dipole (m)	2.35	3.2/1.1	1.6–6.6	5.38	5.4
Length of standard cell (m)	11.4	13.2	16	14.4	14.35
Phase advance per cell (deg)	$H: 79$ $V: 90$	$H: 140$ $V: 50$	45–90 (no standard cell)	$H: 47$ $V: 40$	$H: 56$ $V: 33$
Dipoles in ring	36	$H: 28$ $V: 6$	86	224	192
Quadrupoles in ring	46	68	106	360	248
Peak magnetic field (T)	1.1	1.5	0.3 normal 0.8 high field } at 8 GeV	0.4 at 23 GeV	0.36

HIGH-ENERGY COLLIDER PARAMETERS: e^+e^- Colliders (II)

The numbers here were received from representatives of the colliders in 1993 or early 1994. Numbers are subject to change. Quantities are, where appropriate, r.m.s. H , V , and, s.c. indicate horizontal and vertical directions, and superconducting.

	BEPC (China)	VEPP-4M (Novosibirsk)	TRISTAN (KEK)	SLC (SLAC)	LEP (CERN)
Physics start date	1989	1994	1987	1989	1989
Maximum beam energy (GeV)	2.2	6	32	50	55
Luminosity ($10^{30}\text{cm}^{-2}\text{s}^{-1}$)	10	50	37	0.35	11
Time between collisions (μs)	0.8	0.6	5	8300	22
Crossing angle (μ rad)	0	0	0	0	0
Energy spread (units 10^{-3})	0.58	1	2.3	3	1.0
Bunch length (cm)	≈ 5	5	1.5	0.1	1.8
Beam radius (10^{-6} m)	H : 926 V : 61	H : 1000 V : 30	H : 280 V : 8	H : 2.5 V : 0.8	H : 200 V : 8
Free space at interaction point (m)	± 2.5	± 2	± 2.51	± 2.8	± 3.5
Luminosity lifetime (hr)	7–12	2	2	—	20
Filling time (min)	30	15	40	—	90
Acceleration period (s)	120	150	300	—	320
Injection energy (GeV)	1.3	2	8	50	20
Transverse emittance ($10^{-9}\pi$ rad-m)	H : 660 V : 43	H : 400 V : 20	H : 25.5 at 29 GeV	H : 0.6 V : 0.6	H : 36 V : 2
β^* , amplitude function at interaction point (m)	H : 1.3 V : 0.085	H : 0.75 V : 0.05	H : 1.0 V : 0.04	H : 0.01 V : 0.006	H : 1.00 V : 0.04
Beam-beam tune shift per crossing (units 10^{-4})	420	500	340	—	420
RF frequency (MHz)	199.53	180	508.5808	—	352.2
Particles per bunch (units 10^{10})	20 at 2 GeV	15	22	3.0	41.6
Bunches per ring per species	1	2	2	1	$4e^+ + 4e^-$ $8e^+ + 8e^-$
Average beam current per species (mA)	40 at 2 GeV	40	7	0.0006	3
Beam polarization (%)	—	—	—	e^- : 62	—
Circumference or length (km)	0.2404	0.366	3.02	1.45 + 1.47	26.66
Interaction regions	2	1	4	1	4
Utility insertions	4	1	8	—	4
Magnetic length of dipole (m)	1.6	2	5.86	2.5	11.66/pair
Length of standard cell (m)	6.6	7.2	16.1	5.2	79
Phase advance per cell (deg)	≈ 60	65	60	108	60+90
Dipoles in ring	40 + 4 weak	78	264 + 8 weak	460+440	3280+24 inj. + 64 weak
Quadrupoles in ring	68	150	392	—	520+288 + 8 s.c.
Peak magnetic field (T)	0.9028	0.6	0.41 at 30 GeV	0.597	0.135

HIGH-ENERGY COLLIDER PARAMETERS: e^+e^- Colliders (III)

Proposed e^+e^- colliders. The numbers here were received from representatives of the colliders in 1993 or early 1994. Numbers are subject to change and many are only estimates; those in parentheses are for later upgrades. Quantities are, where appropriate, r.m.s. H and V indicate horizontal and vertical directions.

	KEKB (KEK)	PEP-II (SLAC)	VLEPP, INP (Serpukhov)
Physics start date	1998	1999	?
Maximum beam energy (GeV)	8×3.5	$e^- \times e^+$: 9×3.1 (6.5 GeV c.m. max)	250
Luminosity ($10^{30} \text{cm}^{-2} \text{s}^{-1}$)	2000 (\rightarrow 10000)	3000	3000
Time between collisions (μs)	0.01 (\rightarrow 0.002)	0.0042	—
Crossing angle (μ rad)	± 2800 ($\rightarrow \pm 10,000$)	0	?
Energy spread (units 10^{-3})	0.7	e^-/e^+ : 0.61/0.81	5–100
Bunch length (cm)	0.5	1.0	0.075
Beam radius (10^{-6} m)	H : 140 V : 1.4	H : 155 V : 6.2	H : 1 V : 0.007
Free space at interaction point (m), angular spread	± 0.2 , (+300/ – 500) mrad cone	± 0.2 , ± 300 mrad cone	± 1.2
Luminosity lifetime (hr)	3	2	—
Filling time (min)	6 (\rightarrow 13) topping up	3 (topping up)	—
Acceleration period (s)	—	—	0.0033
Injection energy (GeV)	8/3.5	2.8–12	3.5
Transverse emittance ($10^{-9} \pi$ rad-m)	H : 19 V : 0.19	e^- : 48 (H), 1.9 (V) e^+ : 64 (H), 2.6 (V)	H : 0.2 V : 3×10^{-4}
β^* , amplitude function at interaction point (m)	H : 1.0 V : 0.01	e^- : 0.50 (H), 0.02 (V) e^+ : 0.375 (H), 0.015 (V)	H : 5×10^{-3} V : 10^{-4}
Beam-beam tune shift per crossing (units 10^{-4})	500	300	—
RF frequency (MHz)	508	476	1.4×10^4
Particles per bunch (units 10^{10})	1.3/3.2	e^-/e^+ : 2.7/5.9	10–20
Bunches per ring per species	1024 (\rightarrow 5120)	1658	1
Average beam current per species (mA)	220/520 (\rightarrow 1100/2600)	e^-/e^+ : 990/2140	0.003
Circumference or length (km)	3.02	2.2	2×3
Interaction regions	1	1 (2 possible)	1
Utility insertions	3	5	—
Magnetic length of dipole (m)	2.56/0.42	e^-/e^+ : 5.4/0.45	—
Length of standard cell (m)	19	15.2	1.2
Phase advance per cell (deg)	90	e^-/e^+ : 60/90	20–90
Dipoles in ring	224	e^-/e^+ : 192/192	—
Quadrupoles in ring	343/341	e^-/e^+ : 282/282	20,000
Peak magnetic field (T)	0.3/0.85	e^-/e^+ : 0.18/0.75	—

HIGH-ENERGY COLLIDER PARAMETERS: e^+e^- Colliders (IV)

The numbers here were received from representatives of the colliders in 1993 or early 1994. Numbers are subject to change. Quantities are, where appropriate, r.m.s. H , V , and, s.c. indicate horizontal and vertical directions, and superconducting.

	VEPP-2M	DAΦNE	φ FACTORY (UCLA)	φ FACTORY (Novosibirsk)
Physics start date	1992	1996	?	1997
Maximum beam energy (GeV)	0.7	2×0.510 (1.5 c.m. max.)	0.18×1.5	0.55
Luminosity ($10^{30} \text{cm}^{-2} \text{s}^{-1}$)	6	135(→540)	10	1000
Time between collisions (μs)	0.03	0.0108(→0.0027)	—	0.06
Crossing angle (μ rad)	0	$(1.0 \text{ to } 1.5) \times 10^4$	—	0
Energy spread (units 10^{-3})	0.6	0.40	—	0.5
Bunch length (cm)	3	3.0	—	1
Beam radius (10^{-6} m)	H : 400 V : 10	H : 2100 V : 21	—	H : 65 V : 65
Free space at interaction point (m)	± 1	± 0.46	—	± 2
Luminosity lifetime (hr)	0.3	3.0	—	continuous
Filling time (min)	continuous	< 2 (topping up)	—	continuous
Acceleration period (s)	—	—	—	—
Injection energy (GeV)	—	0.510	—	—
Transverse emittance ($10^{-9} \pi$ rad-m)	H : 400 V : 4	H : 1000 V : 10	Lo E: H/V : 312/13 Hi E: H/V : 156/6.2	H : 400 V : 400
β^* , amplitude function at interaction point (m)	H : 0.48 V : 0.04	H : 4.5 V : 0.045	Lo E: H/V : 0.50/0.02 Hi E: H/V : 1.00/0.04	H : 0.01 V : 0.01
Beam-beam tune shift per crossing (units 10^{-4})	500	400	—	1000
RF frequency (MHz)	200	368.25	Lo E/Hi E=212/500	700
Particles per bunch (units 10^{10})	4	8.9	—	16
Bunches per ring per species	1	30(→120)	Lo E/Hi E=1/3	1
Average beam current per species (mA)	100	1313(→5250)	Lo E/Hi E=180/43	200
Circumference or length (m)	18	97.7	Lo E/Hi E=8.5/25.5	36
Interaction regions	2	2	1	1
Utility insertions	1	2×2	—	1
Magnetic length of dipole (m)	1	1.21/0.99	—	0.9
Length of standard cell (m)	4.5	—	—	9
Phase advance per cell (deg)	280	—	—	548
Dipoles in ring	8	8(+4 wigglers)	—	16
Quadrupoles in ring	20	51	—	28
Peak magnetic field (T)	1.8	1.2(→1.76) dipoles 1.8 wigglers	Lo E/Hi E=1.0/7.0	2.2

HIGH-ENERGY COLLIDER PARAMETERS: p , $\bar{p}p$, and ep Colliders

The numbers here were received from representatives of the colliders in 1993 or early 1994. Numbers are subject to change, and many are only estimates. Quantities are, where appropriate, r.m.s. H , V , and, s.c. indicate horizontal and vertical directions, and superconducting.

	SppS (CERN)	TEVATRON (Fermilab)	HERA (DESY)	UNK (Serpukhov)	LHC (CERN)		SSC (USA)
Physics start date	1981	1987	1990	?	2002		Terminated
Particles collided	$p\bar{p}$	$p\bar{p}$	ep	pp	pp	Pb Pb	pp
Maximum beam energy (TeV)	0.315 (0.45 in pulsed mode)	0.9–1.0	e : 0.030 p : 0.82	0.4 (3)	7.0	574	20
Luminosity ($10^{30}\text{cm}^{-2}\text{s}^{-1}$)	6	7.5 (1993) 10 (1994)	16	1000	1.0×10^4	0.002	1000
Time between collisions (μs)	3.8	3.5	0.096	0.165	0.025	0.135	0.016678
Crossing angle (μrad)	0	0	0	0	200	≤ 100	100 to 200 (135 nominal)
Energy spread (units 10^{-3})	0.35	0.15	e : 0.91 p : 0.2	± 1 (± 0.3)	0.1	0.1	0.055
Bunch length (cm)	20	50	e : 0.83 p : 8.5	70 (40)	7.5	7.5	6.0
Beam radius (10^{-6}m)	p : 73(H), 36(V) \bar{p} : 55(H), 27(V)	36	e : 280(H), 37(V) p : 265(H), 84(V)	70	16	15	4.8
Free space at interaction point (m)	16	± 6.5	± 5.5	± 8	32	32	± 20
Luminosity lifetime (hr)	15	10–40	10	10	10	10	~ 24
Filling time (min)	0.5	120	e : 30 p : 20	20	7	16	72
Acceleration period (s)	10	86	—	100	1200		1500
Injection energy (TeV)	0.026	0.15	e : 0.014 p : 0.040	0.065 (0.4)	0.450		2
Transverse emittance ($10^{-9}\pi\text{rad}\cdot\text{m}$)	p : 9 \bar{p} : 5	p : 2.6 \bar{p} : 2.6	e : 39(H), 2(V) p : 10(H), 10(V)	18 (2.3)	0.5	0.5	0.047
β^* , amplitude function at interaction point (m)	0.6 (H) 0.15 (V)	0.5 \rightarrow 0.25	e : 2(H), 0.7(V) p : 7(H), 0.7(V)	0.2 (1.5)	0.5	0.5	0.5
Beam-beam tune shift per crossing (units 10^{-4})	50	p : 20 \bar{p} : 70	e : 190(H), 210(V) p : 12(H), 9(V)	50	32		8 head on 13 long range
RF frequency (MHz)	100+200	53	e : 499.7 p : 208.2/52.05	200	400	200+ 400	359.75
Particles per bunch (units 10^{10})	p : 15 \bar{p} : 8	p : 15 \bar{p} : 4.5	e : 3.65 p : 10	30	10	0.009	0.8
Bunches per ring per species	6	6	210	348	2835	496	17,424
Average beam current per species (mA)	p : 6 \bar{p} : 3	p : 6.9 \bar{p} : 2.0	e : 58 p : 158	240	536	6.9	71
Circumference (km)	6.911	6.28	6.336	20.772	26.659		87.12
Interaction regions	2	2 high \mathcal{L}	3	4	2 high \mathcal{L}	1	4
Utility insertions	—	4	1	2	2		2
Magnetic length of dipole (m)	6.26	6.12	e : 9.185 p : 8.82	5.8	Mostly 13.50		Mostly 14.928
Length of standard cell (m)	64	59.5	e : 23.5 p : 47	91.8	102.04		180
Phase advance per cell (deg)	90	67.8	e : 60 p : 90	82.5	90		90
Dipoles in ring	744	774	e : 396 p : 416	2204 (2192)	1296 24 crossing dipoles		H : 8336 V : 88 } in 2 rings
Quadrupoles in ring	232	216	e : 580 p : 280	560 (474)	538		2084 } 2 rings
Magnet type	H type with bent-up coil ends	s.c. $\cos\theta$ warm iron	e : C-shaped p : s.c., collared, cold iron	H type (s.c.)	s.c. 2 in 1 cold iron		s.c. $\cos\theta$ cold iron
Peak magnetic field (T)	1.4 (2 in pulsed mode)	4.4	e : 0.274 p : 4.65	0.67 (5)	8.65		6.790
\bar{p} source accum. rate (hr^{-1})	6×10^{10}	5×10^{10}	—	—	—		—
Max. no. \bar{p} in accum. ring	1.2×10^{12}	1×10^{12}	—	—	—		—

10. PASSAGE OF PARTICLES THROUGH MATTER

Revised June 1994.

10.1. Notation

Table 10.1: The notation and values given in Table 1.1 are used. The kinematic variables β and γ have the usual meanings. Definitions of other variables used in this section are summarized below.

Symbol	Definition	Units or Value
α	Fine structure constant	$1/137.0359895(61)$
M	Incident particle mass	MeV/c^2
E	Incident particle energy $\gamma M c^2$	MeV
T	Kinetic energy	MeV
$m_e c^2$	Electron mass $\times c^2$	$0.51099906(15)$ MeV
r_e	Classical electron radius $e^2/4\pi\epsilon_0 c^2$	$2.81794092(38)$ fm
N_A	Avogadro's number	$6.0221367(36) \times 10^{23}$ mol $^{-1}$
ze	Charge of incident particle	
Z	Atomic number of medium	
A	Atomic mass of medium	g mol $^{-1}$
K/A	$4\pi N_A r_e^2 m_e c^2 / A$	0.307075 MeV g $^{-1}$ cm 2 for $A = 1$ g mol $^{-1}$
δ	Density effect correction to ionization energy loss	
$\hbar\omega_p$	Plasma energy $\sqrt{4\pi N_e r_e^3} m_e c^2 / \alpha$	$= 28.816 \sqrt{\rho(Z/A)}$ eV $^{(a)}$
w_j	Fraction by weight of the j th element in a compound or mixture	
n_j	\propto number of j th kind of atoms in a compound or mixture	
X_0	Radiation length	MeV g $^{-1}$ cm 2
E_c	Critical energy	MeV
E_s	Scale energy $\sqrt{4\pi/\alpha} m_e c^2$	21.2052 MeV
R_M	Molière radius	MeV g $^{-1}$ cm 2

(a) For ρ in g cm $^{-3}$.

10.2. Ionization energy loss by heavy particles [1-5]

Moderately relativistic charged particles other than electrons lose energy in matter primarily by ionization. If the incident particle velocity βc is larger than that of orbital electrons ($\sim Z\alpha c$) and small enough that radiative effects do not dominate (for example, pion energy smaller than 100–200 GeV in iron), then the mean rate of energy loss (or stopping power) is given by the Bethe-Bloch equation,

$$-\frac{dE}{dx} = K z^2 \frac{Z}{A} \frac{1}{\beta^2} \left[\frac{1}{2} \ln \frac{2m_e c^2 \beta^2 \gamma^2 T_{\max}}{I^2} - \beta^2 - \frac{\delta}{2} \right]. \quad (10.1)$$

Here T_{\max} is the maximum kinetic energy which can be imparted to a free electron in a single collision, and the other variables are defined in Table 10.1. The units are chosen so that dx is measured in mass per unit area, *e.g.*, in g cm $^{-2}$. The function as computed for pions on copper is shown by the solid curve in Fig. 10.1, and for pions on other materials in Fig. 10.2. A minor dependence on M at the highest energies is introduced through T_{\max} , but for all practical purposes in high-energy physics dE/dx in a given material is a function only of β . Except in hydrogen, particles of the same velocity have very similar rates of energy loss in different materials; there is a slow decrease in the rate of energy loss with increasing Z . The qualitative difference in stopping power behavior at high energies between a gas (He) and the other materials shown in Fig. 10.2 is due to the density effect correction, δ , discussed below. The stopping power functions are characterized by broad minima whose position drops from $\beta\gamma = 3.5$ to 3.0 as Z goes from 7 to 100.

In practical cases, most relativistic particles (*e.g.*, cosmic-ray muons) have energy loss rates close to the minimum, and are said to be minimum ionizing particles, or mip's.

Eq. (10.1) may be integrated to find the total range R for a particle which loses energy only through ionization. Since dE/dx depends only on β , R/M is a function of E/M or pc/M . In practice, range is a useful concept only for low-energy hadrons ($R \lesssim \lambda_I$, where λ_I is the nuclear interaction length), and for muons below a few hundred




Figure 10.1: Energy loss rate in copper. The function without the density effect correction is also shown, as is the shell correction and two low-energy approximations.




Figure 10.2: Energy loss rate in liquid (bubble chamber) hydrogen, gaseous helium, carbon, aluminum, tin, and lead.

GeV (above which radiative effects dominate). R/M as a function of $\beta\gamma = pc/M$ is shown for a variety of materials in Fig. 10.3.

For a point-like charged particle with mass M and momentum $M\beta\gamma$, T_{\max} is given by

$$T_{\max} = \frac{2m_e c^2 \beta^2 \gamma^2}{1 + 2\gamma m_e/M + (m_e/M)^2}. \quad (10.2)$$

It is usual [1,2] to make the low-energy approximation $T_{\max} = 2m_e c^2 \beta^2 \gamma^2$, valid for $2\gamma m_e/M \ll 1$; this, in fact, is done implicitly in many standard references. For pion in copper, the error thus introduced into dE/dx is greater than 6% at 100 GeV. The correct expression should be used.

At such energies, the maximum 4-momentum transfer to the electron can exceed 1 GeV/c, where structure effects significantly




Figure 10.3: Range of heavy charged particles in liquid (bubble chamber) hydrogen, helium gas, carbon, iron, and lead. For example: For a K^+ whose momentum is 700 MeV/c, $\beta\gamma = 1.42$. For lead we read $R/M \approx 396$, and so the range is 195 g cm^{-2} .

modify the cross sections. This problem has been investigated by J.D. Jackson [6], who concluded that corrections to dE/dx are negligible below energies where radiative effects dominate. While the cross section for rare hard collisions is modified, the average stopping power, dominated by many softer collisions, is almost unchanged.

The mean excitation energy I is $10 \pm 1 \text{ eV}$ for elements heavier than oxygen. The values adopted by the ICRU for the chemical elements [7] are now in wide use; these are shown in Fig. 10.4. Machine-readable versions can also be found [8]. Given the availability of these constants and their variation with atomic structure, there seems little point to depending upon approximate formulae, as was done in the past.

A shell correction is often included in the square brackets of Eq. (10.1) [3,5,7], to correct for atomic binding having been neglected in calculating some of the contributions to Eq. (10.1). We show the Barkas form [3] in Fig. 10.1. For copper it contributes about 1% at $\beta\gamma = 0.3$ (kinetic energy 6 MeV for a pion), and the correction decreases very rapidly with energy. While it is negligible for high-energy physics applications, this and other low-energy corrections must be taken into account at lower energies, such as those encountered in medical physics.

As the particle energy increases, its electric field flattens and extends, so that the distant-collision contribution to Eq. (10.1) increases as $\ln \beta\gamma$. However, real media become polarized, limiting the field extension and effectively truncating this part of the logarithmic rise [4,9–13]. At very high energies,

$$\delta/2 \rightarrow \ln(\hbar\omega_p/I) + \ln \beta\gamma - 1/2, \quad (10.3)$$

where $\delta/2$ is the density effect correction introduced in Eq. (10.1) and $\hbar\omega_p$ is the plasma energy defined in Table 10.1. A comparison with Eq. (10.1) shows that $|dE/dx|$ then grows as $\ln \beta\gamma$ rather than $\ln \beta^2\gamma^2$, and that the mean excitation energy I is replaced by the plasma energy $\hbar\omega_p$. The stopping power as calculated with and




Figure 10.4: Excitation energies (divided by Z) as adopted by the ICRU [7]. Those based on measurement are shown by points with error flags; the interpolated values are simply joined. The solid point is for liquid H_2 ; the open point at 19.2 is for H_2 gas. Also shown are curves based on two approximate formulae.

without the density effect correction is shown in Fig. 10.1. Since the plasma frequency scales as the square root of the electron density, the correction is much larger for a liquid or solid than for a gas, as is illustrated by the examples in Fig. 10.2.

The remaining relativistic rise can be attributed to large energy transfers to a few electrons. If these escape or are otherwise accounted for separately, the energy deposited in an absorbing layer (in contrast to the energy lost by the particle) approaches a constant value, the Fermi plateau (see Sec. 10.3 below). The curve in Fig. 10.1 labeled “ $T_{\text{cut}} = 0.5 \text{ MeV}$ ” illustrates this behavior. At extreme energies (e.g., 400 GeV for muons or pions in iron), radiative effects become important. These are especially relevant for high-energy muons, as discussed in Sec. 10.9.

For particles moving more slowly than atomic electrons, the above discussion is inapplicable. At velocities $\alpha z \gtrsim \beta \gtrsim 10^{-3}$ or slightly lower, the total energy-loss rate is proportional to β , and non-ionizing nuclear recoil energy loss contributes substantially to the total [14]. For protons in silicon, $|dE/dx| = 61.2 \beta \text{ GeV cm}^2 \text{ g}^{-1}$ for $\beta < 0.005$; the peak occurs at $\beta = 0.0126$, where $|dE/dx| = 522 \text{ MeV cm}^2 \text{ g}^{-1}$. In neutron-scattering experiments, light output in scintillator has been observed for recoil protons with energies as low as 30 eV [15].

It is often stated that for $\beta \gtrsim z/137$, $|dE/dx|$ falls as β^{-2} before reaching the broad minimum at $\beta\gamma \approx 3.0\text{--}3.5$. In fact, the slope is nowhere this great, and $|dE/dx| \propto \beta^{-5/3}$ provides a very good approximation to the actual function out to $\beta\gamma > 1$. This behavior is shown in Fig. 10.1, along with the traditional β^{-2} proportionality.

The quantity $(dE/dx)\delta x$ is the mean energy loss via interaction with electrons in a layer of the medium with thickness δx . For finite δx , there are fluctuations in the actual energy loss. The distribution is skewed toward high values (the Landau tail) [1,16]. Only for a thick layer $[(dE/dx)\delta x \gg 2m_e c^2 \beta^2 \gamma^2]$ is the distribution nearly Gaussian. The large fluctuations in the energy loss are due to the small number of collisions involving large energy transfers. The fluctuations are smaller for the so-called restricted energy loss rate, as discussed in Sec. 10.3 below.

A mixture or compound can be thought of as made up of thin layers of pure elements in the right proportion (Bragg additivity). In this case,

$$\frac{dE}{dx} = \sum w_j \left. \frac{dE}{dx} \right|_j, \quad (10.4)$$

where $dE/dx|_j$ is the mean rate of energy loss (in MeV g cm^{-2}) in the j th element. Eq. (10.1) can be inserted into Eq. (10.4) to find expressions for $\langle Z/A \rangle$, $\langle I \rangle$, and $\langle \delta \rangle$; for example, $\langle Z/A \rangle =$

$\sum w_j Z_j / A_j = \sum n_j Z_j / \sum n_j A_j$. However, $\langle I \rangle$ as defined this way is an underestimate, because in a compound electrons are more tightly bound than in the free elements, and $\langle \delta \rangle$ as calculated this way has little relevance, because it is the electron density which matters. If possible, one uses the tables given in Refs. 13 and 12, which include effective excitation energies and interpolation coefficients for calculating the density effect correction for the chemical elements and nearly 200 mixtures and compounds. If a compound or mixture is not found, then one uses the recipe for δ given in Ref. 10 (or Ref. 8), and calculates $\langle I \rangle$ according to the discussion in Ref. 11. (Note the "13%" rule!)

Ionization losses by electrons and positrons [12] are not discussed here. Above the critical energy, which is a few tens of MeV in most materials, bremsstrahlung is the dominant source of energy loss. This important case is discussed below. The contributions of various electron energy-loss processes in lead are shown in Fig. 11.4.

10.3. Restricted energy loss rates for relativistic ionizing particles

Fluctuations in energy loss are due mainly to the production of a few high-energy knock-on electrons. Practical detectors often measure the energy *deposited*, not the energy *lost*. When energy is carried off by energetic knock-on electrons, it is more appropriate to consider the mean energy loss excluding energy transfers greater than some cutoff T_{cut} . The restricted energy loss rate is

$$-\frac{dE}{dx} \Big|_{T < T_{\text{cut}}} = K z^2 \frac{Z}{A} \frac{1}{\beta^2} \left[\frac{1}{2} \ln \frac{2m_e c^2 \beta^2 \gamma^2 T_{\text{upper}}}{I^2} - \frac{\beta^2}{2} \left(1 + \frac{T_{\text{upper}}}{T_{\text{max}}} \right) - \frac{C}{Z} - \frac{\delta}{2} \right] \quad (10.5)$$

where $T_{\text{upper}} = \text{MIN}(T_{\text{cut}}, T_{\text{max}})$. This form agrees with the equation given in previous editions of this *Review* [17] for $T_{\text{cut}} \ll T_{\text{max}}$ but smoothly joins the normal Bethe-Bloch function (Eq. (10.1)) for $T_{\text{cut}} > T_{\text{max}}$.

10.4. Energetic knock-on electrons (δ rays)

The distribution of secondary electrons with kinetic energies $T \gg I$ is given by [1]

$$\frac{d^2 N}{dT dx} = \frac{1}{2} K z^2 \frac{Z}{A} \frac{1}{\beta^2} \frac{F(T)}{T^2} \quad (10.6)$$

for $I \ll T \leq T_{\text{max}}$, where T_{max} is given by Eq. (10.2). The factor F is spin-dependent, but is about unity for $T \ll T_{\text{max}}$. For spin-0 particles $F(T) = (1 - \beta^2 T / T_{\text{max}})$; forms for spins 1/2 and 1 are also given by Rossi [1]. When Eq. (10.6) is integrated from T_{cut} to T_{max} , one obtains the difference between Eq. (10.1) and Eq. (10.5). For incident electrons, the indistinguishability of projectile and target means that the range of T extends only to half the kinetic energy of the incident particle. Additional formulae are given in Ref. 18. Equation (10.6) is inaccurate for T close to I : for $2I \lesssim T \lesssim 10I$, the $1/T^2$ dependence above becomes approximately $T^{-\eta}$, with $3 \lesssim \eta \lesssim 5$ [19].

10.5. Ionization yields

Physicists frequently relate total energy loss to the number of ion pairs produced near the particle's track. This relation becomes complicated for relativistic particles due to the wandering of energetic knock-on electrons whose ranges exceed the dimensions of the fiducial volume. For a qualitative appraisal of the nonlocality of energy deposition in various media by such modestly energetic knock-on electrons, see Ref. 20. The mean local energy dissipation per local ion pair produced, W , while essentially constant for relativistic particles, increases at slow particle speeds [21]. For gases, W can be surprisingly sensitive to trace amounts of various contaminants [21]. Furthermore, ionization yields in practical cases may be greatly influenced by such factors as subsequent recombination [22].

10.6. Multiple scattering through small angles

A charged particle traversing a medium is deflected by many small-angle scatters. Most of this deflection is due to Coulomb scattering from nuclei, and hence the effect is called multiple Coulomb scattering. (However, for hadronic projectiles, the strong interactions also contribute to multiple scattering.) The Coulomb scattering distribution is well represented by the theory of Molière [23]. It is roughly Gaussian for small deflection angles, but at larger angles (greater than a few θ_0 , defined below) it behaves like Rutherford scattering, having larger tails than does a Gaussian distribution.

If we define

$$\theta_0 = \theta_{\text{plane}}^{\text{rms}} = \frac{1}{\sqrt{2}} \theta_{\text{space}}^{\text{rms}} \quad (10.7)$$

then it is sufficient for many applications to use a Gaussian approximation for the central 98% of the projected angular distribution, with a width given by [24,25]

$$\theta_0 = \frac{13.6 \text{ MeV}}{\beta c p} z \sqrt{x/X_0} \left[1 + 0.038 \ln(x/X_0) \right] \quad (10.8)$$

Here p , βc , and z are the momentum, velocity, and charge number of the incident particle, and x/X_0 is the thickness of the scattering medium in radiation lengths (defined below). This value of θ_0 is from a fit to Molière distribution [23] for singly charged particles with $\beta = 1$ for all Z , and is accurate to 11% or better for $10^{-3} < x/X_0 < 100$.

Eq. (10.8) describes scattering from a single material, while the usual problem involves the multiple scattering of a particle traversing many different layers and mixtures. Since it is from a fit to a Molière distribution, it is incorrect to add the individual θ_0 contributions in quadrature; the result is systematically too small. It is much more accurate to apply Eq. (10.8) once, after finding x and X_0 for the combined scatterer.

Lynch and Dahl have extended this phenomenological approach, fitting Gaussian distributions to a variable fraction of the Molière distribution for arbitrary scatterers [25], and achieve accuracies of 2% or better.




Figure 10.5: Quantities used to describe multiple Coulomb scattering. The particle is incident in the plane of the figure.

The nonprojected (space) and projected (plane) angular distributions are given approximately by [23]

$$\frac{1}{2\pi \theta_0^2} \exp \left(-\frac{\theta_{\text{space}}^2}{2\theta_0^2} \right) d\Omega, \quad (10.9)$$

$$\frac{1}{\sqrt{2\pi} \theta_0} \exp \left(-\frac{\theta_{\text{plane}}^2}{2\theta_0^2} \right) d\theta_{\text{plane}}, \quad (10.10)$$

where θ is the deflection angle. In this approximation, $\theta_{\text{space}}^2 \approx (\theta_{\text{plane},x}^2 + \theta_{\text{plane},y}^2)$, where the x and y axes are orthogonal to the direction of motion, and $d\Omega \approx d\theta_{\text{plane},x} d\theta_{\text{plane},y}$. Deflections into $\theta_{\text{plane},x}$ and $\theta_{\text{plane},y}$ are independent and identically distributed.

Figure 10.5 shows these and other quantities sometimes used to describe multiple Coulomb scattering. They are

$$\psi_{\text{plane}}^{\text{rms}} = \frac{1}{\sqrt{3}} \theta_{\text{plane}}^{\text{rms}} = \frac{1}{\sqrt{3}} \theta_0, \quad (10.11)$$

$$y_{\text{plane}}^{\text{rms}} = \frac{1}{\sqrt{3}} x \theta_{\text{plane}}^{\text{rms}} = \frac{1}{\sqrt{3}} x \theta_0, \quad (10.12)$$

$$s_{\text{plane}}^{\text{rms}} = \frac{1}{4\sqrt{3}} x \theta_{\text{plane}}^{\text{rms}} = \frac{1}{4\sqrt{3}} x \theta_0. \quad (10.13)$$

All the quantitative estimates in this section apply only in the limit of small $\theta_{\text{plane}}^{\text{rms}}$ and in the absence of large-angle scatters. The random variables s , ψ , y , and θ in a given plane are distributed in a correlated fashion (see Sec. 16.1 of this *Review* for the definition of the correlation coefficient). Obviously, $y \approx x\psi$. In addition, y and θ have the correlation coefficient $\rho_{y\theta} = \sqrt{3}/2 \approx 0.87$. For Monte Carlo generation of a joint $(y_{\text{plane}}, \theta_{\text{plane}})$ distribution, or for other calculations, it may be most convenient to work with independent Gaussian random variables (z_1, z_2) with mean zero and variance one, and then set

$$\begin{aligned} y_{\text{plane}} &= z_1 x \theta_0 (1 - \rho_{y\theta}^2)^{1/2} / \sqrt{3} + z_2 \rho_{y\theta} x \theta_0 / \sqrt{3} \\ &= z_1 x \theta_0 / \sqrt{12} + z_2 x \theta_0 / 2; \end{aligned} \quad (10.14)$$

$$\theta_{\text{plane}} = z_2 \theta_0. \quad (10.15)$$

Note that the second term for y_{plane} equals $x \theta_{\text{plane}}/2$ and represents the displacement that would have occurred had the deflection θ_{plane} all occurred at the single point $x/2$.

For heavy ions the multiple Coulomb scattering has been measured and compared with various theoretical distributions [26].

10.7. Radiation length and associated quantities

In dealing with electrons and photons at high energies, it is convenient to measure the thickness of the material in units of the radiation length X_0 . This is the mean distance over which a high-energy electron loses all but $1/e$ of its energy by bremsstrahlung, and is the appropriate scale length for describing high-energy electromagnetic cascades. X_0 has been calculated and tabulated by Y.S. Tsai [27]:

$$\frac{1}{X_0} = 4\alpha r_e^2 \frac{N_A}{A} \left\{ Z^2 [L_{\text{rad}} - f(Z)] + Z L'_{\text{rad}} \right\}, \quad (10.16)$$

where L_{rad} and L'_{rad} are given in Table 10.2. The function $f(Z)$ is an infinite sum, but for elements up to uranium can be represented to 4-place accuracy by

$$\begin{aligned} f(Z) &= a^2 [(1 + a^2)^{-1} + 0.20206 \\ &\quad - 0.0369 a^2 + 0.0083 a^4 - 0.002 a^6], \end{aligned} \quad (10.17)$$

where $a = \alpha Z$ [28].

Table 10.2: Tsai's L_{rad} and L'_{rad} , for use in calculating the radiation length in an element using Eq. (10.16).

Element	Z	L_{rad}	L'_{rad}
H	1	5.31	6.144
He	2	4.79	5.621
Li	3	4.74	5.805
Be	4	4.71	5.924
Others	> 4	$\ln(184.15 Z^{-1/3})$	$\ln(1194 Z^{-2/3})$

Although it is easy to use Eq. (10.16) to calculate X_0 , the functional dependence on Z is somewhat hidden. Dahl provides a compact fit to the data [29]:

$$X_0 = \frac{716.4 \text{ g cm}^{-2} A}{Z(Z+1) \ln(287/\sqrt{Z})} \quad (10.18)$$

Results obtained with this formula agree with Tsai's values to better than 2.5% for all elements except helium, where the result is about 5% low.




Figure 10.6: Two definitions of the critical energy E_c .




Figure 10.7: Critical energy for the chemical elements, using Rossi's definition [1]. The fits shown are for solids and liquids (solid line) and gases (dashed line). The rms deviation is 2.2% for the solids and 4.0% for the gases.

The radiation length in a mixture or compound may be approximated by

$$1/X_0 = \sum w_j / X_j, \quad (10.19)$$

where w_j and X_j are the fraction by weight and the radiation length for the i th element.

An electron loses energy by bremsstrahlung at a rate nearly proportional to its energy, while the ionization loss rate varies only logarithmically with the electron energy. The *critical energy* E_c is sometimes defined as the energy at which the two loss rates are equal [30]. Berger and Seltzer [30] also give the approximation $E_c = (800 \text{ MeV})/(Z + 1.2)$. This formula has been widely quoted, and has been given in previous editions of this *Review* [17]. Among alternate definitions is that of Rossi [1], who defines the critical energy as the energy at which the ionization loss per radiation length is equal to the electron energy. Equivalently, it is the same as the first definition with the approximation $|dE/dx|_{\text{brems}} \approx E/X_0$. These definitions are illustrated in the case of copper in Fig. 10.6 [31].

The accuracy of approximate forms for E_c has been limited by the failure to distinguish between gases and solid or liquids, where there is a substantial difference in ionization at the relevant energy because of the density effect. We distinguish these two cases in Fig. 10.7. Fits

were also made with functions of the form $a/(Z+b)^\alpha$, but α was essentially unity.

The transverse development of electromagnetic showers in different materials scales fairly accurately with the *Molière radius* R_M , given by [32,33]

$$R_M = X_0 E_s / E_c, \quad (10.20)$$

where $E_s \approx 21$ MeV (see Table 10.1), and the Rossi definition of E_c is used.

In a material containing a weight fraction w_j of the element with critical energy E_{cj} and radiation length X_j , the Molière radius is given by

$$\frac{1}{R_M} = \frac{1}{E_s} \sum \frac{w_j E_{cj}}{X_j}. \quad (10.21)$$

For very high-energy photons, the total e^+e^- pair-production cross section is approximately

$$\sigma = \frac{7}{9}(A/X_0 N_A), \quad (10.22)$$

where A is the atomic weight of the material and N_A is Avogadro's number. Equation Eq. (10.22) is accurate to within a few percent down to energies as low as 1 GeV. The cross section decreases at lower energies, as shown in Fig. 11.4 of this *Review*. As the energy decreases, a number of other processes become important, as is shown in Fig. 11.3 of this *Review*.

10.8. Electromagnetic cascades

When a high-energy electron or photon is incident on a thick absorber, it initiates an electromagnetic cascade as pair production and bremsstrahlung generate more electrons and photons with lower energy. The longitudinal development is governed by the high-energy part of the cascade, and therefore scales as the radiation length in the material. Electron energies eventually fall below the critical energy, and then dissipate their energy by ionization and excitation rather than by the generation of more shower particles. In describing shower behavior, it is therefore convenient to introduce the scale variables

$$\begin{aligned} t &= x/X_0 \\ y &= E/E_c, \end{aligned} \quad (10.23)$$

so that distance is measured in units of radiation length and energy in units of critical energy.




Figure 10.8: An EGS4 simulation of a 30 GeV electron-induced cascade in iron. The histogram shows fractional energy deposition per radiation length, and the curve is a gamma-function fit to the distribution. Circles indicate the number of electrons with total energy greater than 1.5 MeV crossing planes at $X_0/2$ intervals (scale on right) and the squares the number of photons with $E \geq 1.5$ MeV crossing the planes (scaled down to have same area as the electron distribution).

Longitudinal profiles for an EGS4 [8] simulation of a 30 GeV electron-induced cascade in iron are shown in Fig. 10.8. The number of particles crossing a plane (very close to Rossi's Π function [1]) is sensitive to the cutoff energy, here chosen as a total energy of 1.5 MeV for both electrons and photons. The electron number falls off more quickly than energy deposition. This is because, with increasing depth, a larger fraction of the cascade energy is carried by photons. Exactly what a calorimeter measures depends on the device, but it is not likely to be exactly any of the profiles shown. In gas counters it may be very close to the electron number, but in glass Čerenkov detectors and other devices with "thick" sensitive regions it is closer to the energy deposition (total track length). In such detectors the signal is proportional to the "detectable" track length T_d , which is in general less than the total track length T . Practical devices are sensitive to electrons with energy above some detection threshold E_d , and $T_d = T F(E_d/E_c)$. An analytic form for $F(E_d/E_c)$ obtained by Rossi [1] is given by Fabjan [34]; see also Amaldi [35].

The mean longitudinal profile of the energy deposition in an electromagnetic cascade is reasonably well described by a gamma distribution [36]:

$$\frac{dE}{dt} = E_0 b \frac{(bt)^{a-1} e^{-bt}}{\Gamma(a)} \quad (10.24)$$

The maximum t_{\max} occurs at $(a-1)/b$. We have made fits to shower profiles in elements ranging from carbon to uranium, at energies from 1 GeV to 100 GeV. The energy deposition profiles are well described by Eq. (10.24) with

$$t_{\max} = (a-1)/b = 1.0 \times (\ln y + C_j), \quad j = e, \gamma, \quad (10.25)$$

where $C_e = -0.5$ for electron-induced cascades and $C_\gamma = +0.5$ for photon-induced cascades. To use Eq. (10.24), one finds $(a-1)/b$ from Eq. (10.25) and Eq. (10.23), then finds a either by assuming $b \approx 0.5$ or by finding a more accurate value from Fig. 10.9. The results are very similar for the electron number profiles, but there is some dependence on the atomic number of the medium. A similar form for the electron number maximum was obtained by Rossi in the context of his "Approximation B," [1] (see Fabjan's review in Ref. 34), but with $C_e = -1.0$ and $C_\gamma = -0.5$; we regard this as superseded by the EGS4 result.




Figure 10.9: Fitted values of the scale factor b for energy deposition profiles obtained with EGS4 for a variety of elements for incident electrons with $1 \leq E_0 \leq 100$ GeV. Values obtained for incident photons are essentially the same.

The "shower length" $X_s = X_0/b$ is less conveniently parametrized, since b depends upon both Z and incident energy, as shown in Fig. 10.9. As a corollary of this Z dependence, the number of electrons crossing a plane near shower maximum is underestimated using Rossi's

approximation for carbon and seriously overestimated for uranium. Essentially the same b values are obtained for incident electrons and photons. For many purposes it is sufficient to take $b \approx 0.5$.

The gamma distribution is very flat near the origin, while the EGS4 cascade (or a real cascade) increases more rapidly. As a result Eq. (10.24) fails badly for about the first two radiation lengths; it was necessary to exclude this region in making fits.

Because fluctuations are important, Eq. (10.24) should be used only in applications where average behavior is adequate. Grindhammer *et al.* have developed fast simulation algorithms in which the variance and correlation of a and b are obtained by fitting Eq. (10.24) to individually simulated cascades, then generating profiles for cascades using a and b chosen from the correlated distributions [37].

Measurements of the lateral distribution in electromagnetic cascades are shown in Refs. 32 and 33. On the average, only 10% of the energy lies outside the cylinder with radius R_M . About 99% is contained inside of $3.5R_M$, but at this radius and beyond composition effects become important and the scaling with R_M fails. The distributions are characterized by a narrow core, and broaden as the shower develops. They are often represented as the sum of two Gaussians, and Grindhammer [37] describes them with the function

$$f(r) = \frac{2r R^2}{(r^2 + R^2)^2}, \quad (10.26)$$

where R is a phenomenological function of x/X_0 and $\ln E$.

10.9. Muon energy loss at high energy

At sufficiently high energies, radiative processes become more important than ionization for all charged particles. For muons and pions in materials such as iron, this “critical energy” occurs at several hundred GeV. Radiative effects dominate the energy loss of energetic muons found in cosmic rays or produced at the newest accelerators. These processes are characterized by small cross sections, hard spectra, large energy fluctuations, and the associated generation of electromagnetic and (in the case of photonuclear interactions) hadronic showers. As a consequence, at these energies the treatment of energy loss as a uniform and continuous process is for many purposes inadequate.

It is convenient to write the average rate of muon energy loss as [38]

$$-dE/dx = a(E) + b(E)E. \quad (10.27)$$

Here $a(E)$ is the ionization energy loss given by Eq. (10.1), and $b(E)$ is the sum of e^+e^- pair production, bremsstrahlung, and photonuclear contributions. To the approximation that these slowly-varying functions are constant, the mean range x_0 of a muon with initial energy E_0 is given by

$$x_0 \approx (1/b) \ln(a + bE_0). \quad (10.28)$$

Figure 10.10 shows contributions to $b(E)$ for iron. Since $a(E) \approx 0.002 \text{ GeV g}^{-1} \text{ cm}^2$, $b(E)E$ dominates the energy loss above several hundred GeV, where $b(E)$ is nearly constant. The rate of energy loss for muons in hydrogen, uranium, and iron is shown in Fig. 10.11 [39].

QED calculations of cross sections for bremsstrahlung and e^+e^- pair production have long been known, but were much improved around 1970 to meet the needs of cosmic ray physics [40–44]. Rozenal showed that the screened atomic electron contribution could be included by replacing Z^2 with $Z(Z+1.2)$ in the nuclear bremsstrahlung cross sections and by $Z(Z+1.3)$ in the case of e^+e^- pair production [45], and that other corrections might reduce the cross section by as much as 5%. We take this as the present uncertainty. Cross sections for both processes have been evaluated independently by Tsai [27].

A comparison of various improvements to the Bethe-Heitler formula is given by Wright [46]. For muon energies above 100 GeV, $\mu^+\mu^-$ pair production is also possible. This process is potentially troublesome because it can lead to charge misassignment, but it contributes less than 0.01% to the total energy loss [39].




Figure 10.10: Contributions to the fractional energy loss by muons in iron due to e^+e^- pair production, bremsstrahlung, and photonuclear interactions, as obtained from Lohmann *et al.* [39].




Figure 10.11: The average energy loss of a muon in hydrogen, iron, and uranium as a function of muon energy. Contributions to dE/dx in iron from ionization and the processes shown in Fig. 10.10 are also shown.

Photonuclear interactions account for about 5% of the total energy loss of high-energy muons in iron, and for about 2% in uranium [47]. The losses are concentrated in rare, relatively hard events.

These radiative cross sections are expressed as functions of the fractional energy loss ν . The bremsstrahlung cross section goes roughly as $1/\nu$ over most of the range, while for the pair production case the distribution goes as ν^{-3} to ν^{-2} (see Ref. 48). “Hard” losses are therefore more probable in bremsstrahlung, and in fact energy losses due to pair production may very nearly be treated as continuous. The momentum distribution of an incident 1 TeV/c muon beam after it crosses 3 m of iron is shown in Fig. 10.12. The most probable loss is 9 GeV, or $3.8 \text{ MeV g}^{-1} \text{ cm}^2$. The full width at half maximum is 7 GeV/c, or 0.7%. The radiative tail is almost entirely due to bremsstrahlung; this includes most of the 10% that lost more than 2.8% of their energy. Most of the 3.3% that lost more than 10% of their incident energy experienced photonuclear interactions. The latter can exceed nominal detector resolution [49], necessitating the reconstruction of lost energy. Electromagnetic and hadronic cascades in detector materials can obscure muon tracks in detector planes and reduce tracking efficiency [50].




Figure 10.12: The momentum distribution of 1 TeV/c muons after traversing 3 m of iron, as obtained with Van Ginniken's TRAMU muon transport code [48].

10.10. Čerenkov and transition radiation [4,51,52]

A charged particle radiates if its velocity is greater than the local phase velocity of light (Čerenkov radiation) or if it crosses suddenly from one medium to another with different optical properties (transition radiation). Neither process is important for energy loss, but both are used in high-energy physics detectors.

Čerenkov Radiation. The half-angle θ_c of the Čerenkov cone for a particle with velocity βc in a medium with index of refraction n is

$$\theta_c = \arccos(1/n\beta) \approx \sqrt{2(1-1/n\beta)} \quad \text{for small } \theta_c, \text{ e.g. in gases.} \quad (10.29)$$

The threshold velocity β_t is $1/n$, and $\gamma_t = 1/(1-\beta_t^2)^{1/2}$. Therefore, $\beta_t \gamma_t = 1/(2\delta + \delta^2)^{1/2}$, where $\delta = n - 1$. Values of δ for various commonly used gases are given as a function of pressure and wavelength in Ref. 53. For values at atmospheric pressure, see Table 6.1. Data for other commonly used materials are given in Ref. 54.

The number of photons produced per unit path length of a particle with charge ze and per unit energy interval of the photons is

$$\frac{d^2 N}{dE dx} = \frac{\alpha z^2}{\hbar c} \sin^2 \theta_c = \frac{\alpha^2 z^2}{r_e m_e c^2} \left(1 - \frac{1}{\beta^2 n^2(E)}\right) \approx 370 \sin^2 \theta_c(E) \text{ eV}^{-1} \text{ cm}^{-1} \quad (z=1), \quad (10.30)$$

or, equivalently,

$$\frac{d^2 N}{dx d\lambda} = \frac{2\pi \alpha z^2}{\lambda^2} \left(1 - \frac{1}{\beta^2 n^2(\lambda)}\right). \quad (10.31)$$

The index of refraction is a function of photon energy E , as is the sensitivity of the transducer used to detect the light. For practical use, Eq. (10.30) must be multiplied by the the transducer response function and integrated over the region for which $\beta n(E) > 1$. Further details are given in the discussion of Čerenkov detectors in the Detectors section (Sec. 12 of this Review).

Transition Radiation. The energy radiated when a particle with charge ze crosses the boundary between vacuum and a medium with plasma frequency ω_p is

$$I = \alpha z^2 \gamma \hbar \omega_p / 3, \quad (10.32)$$

where

$$\begin{aligned} \hbar \omega_p &= \sqrt{4\pi N_e r_e^3} m_e c^2 / \alpha \\ &= \sqrt{4\pi N_e a_\infty^3} 2 \times 13.6 \text{ eV}. \end{aligned} \quad (10.33)$$

Here N_e is the electron density in the medium, r_e is the classical electron radius, and a_∞ is the Bohr radius. For styrene and similar materials, $\sqrt{4\pi N_e a_\infty^3} \approx 0.8$, so that $\hbar \omega_p \approx 20$ eV. The typical emission angle is $1/\gamma$.

The radiation spectrum is logarithmically divergent at low energies and decreases rapidly for $\hbar \omega / \gamma \hbar \omega_p > 1$. About half the energy is emitted in the range $0.1 \leq \hbar \omega / \gamma \hbar \omega_p \leq 1$. For a particle with $\gamma = 10^3$, the radiated photons are in the soft x-ray range 2 to 20 eV. The γ dependence of the emitted energy thus comes from the hardening of the spectrum rather than from an increased quantum yield. For a typical radiated photon energy of $\gamma \hbar \omega_p / 4$, the quantum yield is

$$\begin{aligned} N_\gamma &\approx \frac{1}{2} \frac{\alpha z^2 \gamma \hbar \omega_p}{3} / \frac{\gamma \hbar \omega_p}{4} \\ &\approx \frac{2}{3} \alpha z^2 \approx 0.5\% \times z^2. \end{aligned} \quad (10.34)$$

More precisely, the number of photons with energy $\hbar \omega > \hbar \omega_0$ is given by [55]

$$N_\gamma(\hbar \omega > \hbar \omega_0) = \frac{\alpha z^2}{\pi} \left[\left(\ln \frac{\gamma \hbar \omega_p}{\hbar \omega_0} - 1 \right)^2 + \frac{\pi^2}{12} \right], \quad (10.35)$$

within corrections of order $(\hbar \omega_0 / \gamma \hbar \omega_p)^2$. The number of photons above a fixed energy $\hbar \omega_0 \ll \gamma \hbar \omega_p$ thus grows as $(\ln \gamma)^2$, but the number above a fixed fraction of $\gamma \hbar \omega_p$ (as in the example above) is constant. For example, for $\hbar \omega > \gamma \hbar \omega_p / 10$, $N_\gamma = 2.519 \alpha z^2 / \pi = 0.59\% \times z^2$.

The yield can be increased by using a stack of plastic foils with gaps between. However, interference can be important, and the soft x rays are readily absorbed in the foils. The first problem can be overcome by choosing thicknesses and spacings large compared to the "formation length" $D = \gamma c / \omega_p$, which in practical situations is tens of μm . Other practical problems are discussed in Sec. 12.

References:

1. B. Rossi, *High Energy Particles*, Prentice-Hall, Inc., Englewood Cliffs, NJ, 1952.
2. U. Fano, *Ann. Rev. Nucl. Sci.* **13**, 1 (1963).
3. W. H. Barkas and M. J. Berger, *Tables of Energy Losses and Ranges of Heavy Charged Particles*, NASA-SP-3013 (1964).
4. J.D. Jackson, *Classical Electrodynamics*, 2nd edition, (John Wiley & Sons, New York, 1975).
5. "Stopping Powers and Ranges for Protons and Alpha Particles," ICRU Report No. 41 (1993).
6. J. D. Jackson, "Effect of Form Factor on dE/dx from Close Collisions," Particle Data Group Note PDG-93-04 (19 October 1993) (unpublished).
7. ICRU Report 37, "Stopping Powers for Electrons and Positrons," International Commission on Radiation Units and Measurements, Washington, DC (1984).
8. W.R. Nelson, H. Hirayama, and D.W.O. Rogers, "The EGS4 Code System," SLAC-265, Stanford Linear Accelerator Center (Dec. 1985).
9. A. Crispin and G.N. Fowler, *Rev. Mod. Phys.* **42**, 290 (1970).
10. R. M. Sternheimer and R. F. Peierls, *Phys. Rev.* **B3**, 3681-3692 (1971).
11. S. M. Seltzer and M. J. Berger, *Int. J. Appl. Radiat.* **33**, 1189-1218 (1982).
12. S. M. Seltzer and M. J. Berger, *Int. J. Appl. Radiat.* **35**, 665-676 (1984). This paper corrects and extends the results of Ref. 11.
13. R.M. Sternheimer, S.M. Seltzer, and M.J. Berger, "The Density Effect for the Ionization Loss of Charged Particles in Various Substances," *Atomic Data & Nucl. Data Tables* **30**, 261 (1984).
14. Low-velocity energy loss has been extensively treated by J.F. Ziegler and his collaborators; see e.g. J.F. Ziegler, Jp.P. Bier-sac, and U. Littmark, *The Stopping and Range of Ions in Solids*, Pergamon Press 1985, and references therein.
15. D.J. Ficenec *et al.*, *Phys. Rev.* **D36**, 311 (1987).

16. L.D. Landau, J. Exp. Phys. (USSR) **8**, 201 (1944); See, for instance, K.A. Ispirian, A.T. Margarian, and A.M. Zverev, Nucl. Instr. and Meth. **117**, 125 (1974).
17. K. Hikasa *et al.*, *Review of Particle Properties*, Phys. Rev. **D46** (1992) S1.
18. For unit-charge projectiles, see E.A. Uehling, Ann. Rev. Nucl. Sci. **4**, 315 (1954). For highly charged projectiles, see J.A. Doggett and L.V. Spencer, Phys. Rev. **103**, 1597 (1956). A Lorentz transformation is needed to convert these center-of-mass data to knock-on energy spectra.
19. N.F. Mott and H.S.W. Massey, *The Theory of Atomic Collisions*, Oxford Press, London, 1965.
20. L.V. Spencer "Energy Dissipation by Fast Electrons," Nat'l Bureau of Standards Monograph No. 1 (1959).
21. "Average Energy Required to Produce an Ion Pair," ICRU Report No. 31 (1979).
22. N. Hadley *et al.*, "List of Poisoning Times for Materials," Lawrence Berkeley Lab Report TPC-LBL-79-8 (1981).
23. H.A. Bethe, Phys. Rev. **89**, 1256 (1953). A thorough review of multiple scattering is given by W.T. Scott, Rev. Mod. Phys. **35**, 231 (1963). However, the data of Shen *et al.*, (Phys. Rev. **D20**, 1584 (1979)) show that Bethe's simpler method of including atomic electron effects agrees better with experiment than does Scott's treatment. For a thorough discussion of simple formulae for single scatters and methods of compounding these into multiple-scattering formulae, see W.T. Scott, Rev. Mod. Phys. **35**, 231 (1963). For detailed summaries of formulae for computing single scatters, see J.W. Motz, H. Olsen, and H.W. Koch, Rev. Mod. Phys. **36**, 881 (1964).
24. V.L. Highland, Nucl. Instr. and Meth. **129**, 497 (1975), and Nucl. Instr. and Meth. **161**, 171 (1979).
25. G.R. Lynch and O.I. Dahl, Nucl. Instr. and Meth. **B58**, 6 (1991).
26. M. Wong *et al.*, Med. Phys. **17**, 163 (1990).
27. Y.S. Tsai, Rev. Mod. Phys. **46**, 815 (1974).
28. H. Davies, H.A. Bethe, and L.C. Maximon, Phys. Rev. **93**, 788 (1954).
29. O.I. Dahl, private communication.
30. M.J. Berger and S.M. Seltzer, "Tables of Energy Losses and Ranges of Electrons and Positrons," National Aeronautics and Space Administration Report NASA-SP-3012 (Washington DC 1964).
31. We are indebted to H. Hirayama for pointing out both the distinctions among definitions and the appropriateness of the Rossi definition (private communication, 1992; informal KEK note (unpublished)).
32. W.R. Nelson, T.M. Jenkins, R.C. McCall, and J.K. Cobb, Phys. Rev. **149**, 201 (1966).
33. G. Bathow *et al.*, Nucl. Phys. **B20**, 592 (1970).
34. *Experimental Techniques in High Energy Physics*, ed. by T. Ferbel (Addison-Wesley, Menlo Park CA 1987).
35. U. Amaldi, Phys. Scripta **23**, 409 (1981).
36. E. Longo and I. Sestili, Nucl. Instr. and Meth. **128**, 283 (1987).
37. G. Grindhammer *et al.*, in *Proceedings of the Workshop on Calorimetry for the Supercollider*, Tuscaloosa, AL, March 13-17, 1989, edited by R. Donaldson and M.G.D. Gilchriese (World Scientific, Teaneck, NJ, 1989), p. 151.
38. P.H. Barrett, L.M. Bollinger, G. Cocconi, Y. Eisenberg, and K. Greisen, Rev. Mod. Phys. **24**, 133 (1952).
39. W. Lohmann, R. Kopp, and R. Voss, "Energy Loss of Muons in the Energy Range 1-10000 GeV," CERN Report 85-03 (1985).
40. H.A. Bethe and W. Heitler, *Proc. Roy. Soc. A* **146**, 83 (1934); H.A. Bethe, *Proc. Cambridge Phil. Soc.* **30**, 542 (1934).
41. A.A. Petrukhin and V.V. Shestakov, Can. J. Phys. **46**, S377 (1968).
42. V.M. Galitskii and S.R. Kel'ner, Sov. Phys. JETP **25**, 948 (1967).
43. S.R. Kel'ner and Yu.D. Kotov, Sov. J. Nucl. Phys. **7**, 237 (1968). S.R. Kel'ner and Yu.D. Kotov, Can. J. Phys. **46**, S387 (1968).
44. R.P. Kokoulin and A.A. Petrukhin, in *Proceedings of the International Conference on Cosmic Rays*, Hobart, Australia, August 16-25, 1971, Vol. 4, p. 2436.
45. I.L. Rozental, Sov. Phys. Usp. **11**, 49 (1968).
46. A.G. Wright, J. Phys. **A6**, 79 (1973).
47. L.B. Bezrukov and E.V. Bugaev, Sov. J. Nucl. Phys. **33**, 635 (1981).
48. A. Van Ginneken, Nucl. Instr. and Meth. **A251**, 21 (1986).
49. U. Becker *et al.*, Nucl. Instr. and Meth. **A253**, 15 (1986).
50. J.J. Eastman and S.C. Loken, in *Proceedings of the Workshop on Experiments, Detectors, and Experimental Areas for the Supercollider*, Berkeley, CA, July 7-17, 1987, edited by R. Donaldson and M.G.D. Gilchriese (World Scientific, Singapore, 1988), p. 542.
51. *Methods of Experimental Physics*, L.C.L. Yuan and C.-S. Wu, editors, Academic Press, 1961, Vol. 5A, p. 163.
52. W.W.M. Allison and P.R.S. Wright, "The Physics of Charged Particle Identification: dE/dx , Čerenkov Radiation, and Transition Radiation," p. 371 in *Experimental Techniques in High Energy Physics*, T. Ferbel, editor, (Addison-Wesley 1987).
53. E.R. Hayes, R.A. Schluter, and A. Tamosaitis, "Index and Dispersion of Some Čerenkov Counter Gases," ANL-6916 (1964).
54. T. Ypsilantis, "Particle Identification at Hadron Colliders", CERN-EP/89-150 (1989), or ECFA 89-124. 2 661 (1989).
55. J.D. Jackson, private communication (1992).

11. PHOTON AND ELECTRON ATTENUATION

Photon Attenuation Length




Figure 11.1: The photon mass attenuation length $\lambda = 1/(\mu/\rho)$ (also known as mfp, mean free path) for various absorbers as a function of photon energy, where μ is the mass attenuation coefficient. For a homogeneous medium of density ρ , the intensity I remaining after traversal of thickness t is given by the expression $I = I_0 \exp(-t\rho/\lambda)$. The accuracy is a few percent. Interpolation to other Z should be done in the cross section $\sigma = A/\lambda N_A \text{ cm}^2/\text{atom}$, where A is the atomic weight of the absorber material in grams and N_A is the Avogadro number. For a chemical compound or mixture, use $(1/\lambda)_{\text{eff}} \approx \sum w_i(1/\lambda)_i$, accurate to a few percent, where w_i is the proportion by weight of the i^{th} constituent. The processes responsible for attenuation are given in Fig. 11.4. Not all of these processes necessarily result in detectable attenuation. For example, coherent Rayleigh scattering off an atom may occur at such low momentum transfer that the change in energy and momentum of the photon may not be significant.

(a) Low-energy region.

(b) The photon mass attenuation length, high-energy range (note that ordinate is linear scale). The attenuation length is constant beyond the range shown for at least two decades in energy.

From Hubbell, Gimm, and Øverbø, *J. Phys. Chem. Ref. Data* **9**, 1023 (80). See also J.H. Hubbell, *Int. J. of Applied Rad. and Isotopes* **33**, 1269 (82). Data courtesy J.H. Hubbell.

Photon Pair Conversion Probability




Figure 11.2: Probability P that a photon interaction will result in conversion to an e^+e^- pair. Except for a few-percent contribution from photonuclear absorption around 10 or 20 MeV, essentially all other interactions result in Compton scattering off an atomic electron. For a photon attenuation length λ (g/cm^2) (Fig. 11.1), the probability that a given photon will produce an electron pair (without first Compton scattering) in thickness t (cm) of absorber of density ρ (g/cm^3) is $P[1 - \exp(-t\rho/\lambda)]$.

Contributions to Photon Cross Section in Carbon and Lead




Figure 11.3: Photon total cross sections as a function of energy in carbon and lead, showing the contributions of different processes.

- $\sigma_{p.e.}$ = Atomic photo-effect (electron ejection, photon absorption)
- $\sigma_{coherent}$ = Coherent scattering (Rayleigh scattering—atom neither ionized nor excited)
- $\sigma_{incoherent}$ = Incoherent scattering (Compton scattering off an electron)
- κ_n = Pair production, nuclear field
- κ_e = Pair production, electron field
- σ_{nuc} = Photonuclear absorption (nuclear absorption, usually followed by emission of a neutron or other particle)

From Hubbell, Gimm, and Øverbø, J. Phys. Chem. Ref. Data 9, 1023 (80). The photon total cross section is assumed approximately flat for at least two decades beyond the energy range shown. Figures courtesy J.H. Hubbell.

Fractional Energy Loss for Electrons and Positrons in Lead




Figure 11.4: Fractional energy loss per radiation length in lead as a function of electron or positron energy. Electron (positron) scattering is considered as ionization when the energy loss per collision is below 0.255 MeV, and as Moller (Bhabha) scattering when it is above. Adapted from Fig. 3.2 from Messel and Crawford, *Electron-Photon Shower Distribution Function Tables for Lead, Copper, and Air Absorbers*, Pergamon Press, 1970. Messel and Crawford use $X_0(\text{Pb}) = 5.82 \text{ g/cm}^2$, but we have modified the figures to reflect the value given in the Table of Atomic and Nuclear Properties of Materials, namely $X_0(\text{Pb}) = 6.4 \text{ g/cm}^2$. The development of electron-photon cascades is approximately independent of absorber when the results are expressed in terms of inverse radiation lengths (*i.e.*, scale on left of plot).

12. PARTICLE DETECTORS

Updated 1992 by D.G. Coyne, R.W. Fast, R.D. Kephart, B. Mansoulie, H.F.W. Sadrozinski, H.G. Spieler, and C.L. Woody

In this section we give various parameters for common detector components. The quoted numbers are usually based on typical devices, and should be regarded only as rough approximations for new designs. A more detailed discussion of detectors can be found in Ref. 1. In Table 12.1 are given typical spatial and temporal resolutions of common detectors.

Table 12.1: Typical detector characteristics.

Detector Type	Accuracy (rms)	Resolution Time	Dead Time
Bubble chamber	10 to 150 μm	1 ms	50 ms ^a
Streamer chamber	300 μm	2 μs	100 ms
Proportional chamber	$\geq 300 \mu\text{m}^{b,c}$	50 ns	200 ns
Drift chamber	50 to 300 μm	2 ns ^d	100 ns
Scintillator	—	150 ps	10 ns
Emulsion	1 μm	—	—
Silicon strip	$\frac{\text{pitch}^e}{3 \text{ to } 7}$	f	f
Silicon pixel	2 μm^g	f	f

^a Multiple pulsing time.

^b 300 μm is for 1 mm pitch.

^c Delay line cathode readout can give $\pm 150 \mu\text{m}$ parallel to anode wire.

^d For two chambers.

^e The highest resolution (“7”) is obtained for small-pitch detectors ($\lesssim 25 \mu\text{m}$) with pulse-height-weighted center finding.

^f Limited at present by properties of the readout electronics. (Time resolution of ≤ 15 ns is planned for the SDC silicon tracker.)

^g Analog readout of 34 μm pitch, monolithic pixel detectors.

12.1. Plastic scintillators

The photon yield in the frequency range of practical photomultiplier tubes is ≈ 1 photon per 100 eV of charged particle ionization energy loss in plastic scintillator [2]. One must take into account the light collection efficiency ($\lesssim 10\%$ for typical 1-cm-thick scintillator), the attenuation length (1 to 4 m for typical scintillators [3]), and the quantum efficiency of the photomultiplier cathode ($\lesssim 25\%$ when folded with a typical scintillator emission spectrum).

12.2. Inorganic scintillators

Table 12.2 gives a partial list of commonly-used inorganic scintillators in high-energy and nuclear physics [4–11]. These scintillating crystals are generally used where high density and good energy resolution are required. In a crystal which contains nearly all of the energy deposited by an incident particle, the energy resolution is determined largely, but not totally, by the light output. The table gives the light output of the various materials relative to NaI, which has an intrinsic light output of about 40000 photons per MeV of energy deposit. The detected signal is usually quoted in terms of photoelectrons per MeV produced by a given photodetector. The relationship between photons/MeV produced and p.e.’s/MeV detected involves factors for light collection efficiency (typically 10–50%, depending on geometry) and the quantum efficiency of the detector (~ 15 –20% for photomultiplier tubes and $\sim 70\%$ for silicon photodiodes for visible wavelengths). The quantum efficiency of the detector is usually highly wavelength dependent and should be matched to the particular crystal of interest to give the highest quantum yield at the wavelength corresponding to the peak of the scintillation emission. The comparison of the light output given in Table 12.2 is for a standard photomultiplier tube with a bialkali photocathode. For scintillators which emit in the UV, a detector with a quartz window should be used.

Table 12.2: Properties of several inorganic crystal scintillators.

	NaI(Tl)	BGO	BaF ₂	CsI(Tl)	CsI(pure)
Density (g/cm ³)	3.67	7.13	4.89	4.53	4.53
Radiation length (cm)	2.59	1.12	2.05	1.85	1.85
Molière radius (cm)	4.5	2.4	3.4	3.8	3.8
dE/dx (MeV/cm)	4.8	9.2	6.6	5.6	5.6
(per mip)					
Nucl. int. length (cm)	41.4	22.0	29.9	36.5	36.5
Decay time (ns)	250	300	0.7 ^f	1000	10, 36 ^f
			620 ^s		$\sim 1000^s$
Peak emission λ (nm)	410	480	220 ^f	565	305 ^f
			310 ^s		$\sim 480^s$
Refractive index	1.85	2.20	1.56	1.80	1.80
Relative light output	1.00	0.15	0.05 ^f	0.40	0.10 ^f
			0.20 ^s		0.02 ^s
Hygroscopic	very	no	slightly	somewhat	somewhat

f = fast component, s = slow component

12.3. Čerenkov detectors

Čerenkov detectors utilize one or more of the properties of Čerenkov radiation discussed in the Passages of Particles through Matter section (Sec. 10 of this Review): the existence of a *threshold* for radiation; the dependence of the Čerenkov cone half-angle θ_c on the *velocity* of the particle; the dependence of the *number of emitted photons* on the particle’s velocity. The presence of the refractive index n in the relations allows tuning these quantities for a particular experimental application (*e.g.*, using pressurized gas and/or various liquids as radiators).

The number of photoelectrons (p.e.’s) detected in a given device or channel is

$$N_{\text{p.e.}} = L \frac{\alpha^2 z^2}{r_e m_e c^2} \int \epsilon_{\text{coll}}(E) \epsilon_{\text{det}}(E) \sin^2 \theta_c(E) dE, \quad (12.1)$$

where L is the path length in the radiator, ϵ_{coll} is the efficiency for collecting the Čerenkov light, ϵ_{det} is the quantum efficiency of the transducer (photomultiplier or equivalent), and $\alpha^2/(r_e m_e c^2) = 370 \text{ cm}^{-1} \text{ eV}^{-1}$. The quantities ϵ_{coll} , ϵ_{det} , and θ_c are all functions of the photon energy E , although in typical detectors θ_c (or, equivalently, the index of refraction) is nearly constant over the useful range of photocathode sensitivity. In this case,

$$N_{\text{p.e.}} \approx LN_0 \langle \sin^2 \theta_c \rangle \quad (12.2)$$

with

$$N_0 = \frac{\alpha^2 z^2}{r_e m_e c^2} \int \epsilon_{\text{coll}} \epsilon_{\text{det}} dE. \quad (12.3)$$

We take $z = 1$, the usual case in high-energy physics, in the following discussion.

Threshold Čerenkov detectors make a simple yes/no decision based on whether the particle is above/below the Čerenkov threshold velocity $\beta_t = 1/n$. Careful designs give $\langle \epsilon_{\text{coll}} \rangle \gtrsim 90\%$. For a photomultiplier with a typical bialkali cathode, $\int \epsilon_{\text{det}} dE \approx 0.27$, so that

$$N_{\text{p.e.}}/L \approx 90 \text{ cm}^{-1} \langle \sin^2 \theta_c \rangle \quad (\text{i.e., } N_0 = 90 \text{ cm}^{-1}). \quad (12.4)$$

Suppose, for example, that n is chosen so that the threshold for species a is p_t ; that is, at this momentum species a has velocity $\beta_a = 1/n$. A second, lighter, species b with the same momentum has velocity β_b , so $\cos \theta_c = \beta_a/\beta_b$, and

$$\frac{N_{\text{p.e.}}}{L} \approx 90 \text{ cm}^{-1} \frac{m_a^2 - m_b^2}{p_t^2 + m_a^2}. \quad (12.5)$$

For K/π separation at $p = 1 \text{ GeV}/c$, $N_{\text{p.e.}}/L \approx 16 \text{ cm}^{-1}$ for π ’s and (by design) 0 for K ’s.

For limited path lengths $N_{p.e.}$ can be small, and some minimum number is required to trigger external electronics. The overall efficiency of the device is controlled by Poisson fluctuations, which can be especially critical for separation of species where one particle type is dominant [12].

A related class of detectors uses the number of observed photoelectrons (or the calibrated pulse height) to discriminate between species or to set probabilities for each particle species [13].

Differential Čerenkov detectors exploit the dependence of θ_c on β , using optical focusing and/or geometrical masking to select particles having velocities in a specified region. With careful design, a velocity resolution of $\sigma_\beta/\beta \approx 10^{-4}$ – 10^{-5} can be obtained [12,14].

Ring-Imaging Čerenkov detectors use all three properties of Čerenkov radiation in both small-aperture and 4π geometries. They are principally used as hypothesis-testing rather than yes/no devices; that is, the probability of various identification possibilities is established from θ_c and $N_{p.e.}$ for a particle of known momentum. In most cases the optics map the Čerenkov cone onto a circle at the photodetector, often with distortions which must be understood.

The 4π devices [15,16] typically have both liquid (C_6F_{14} , $n = 1.276$) and gas (C_5F_{12} , $n = 1.0017$) radiators, the light from the latter being focused by mirrors. They achieve 3σ separation of $e/\pi/K/p$ over wide ranges, as shown in Table 12.3. Great attention to detail, especially with the minimization of UV-absorbing impurities, is required to get $\langle\epsilon_{coll}\rangle \gtrsim 50\%$.

Table 12.3: Momentum range for 3σ separation in the SLD ring-imaging Čerenkov detector.

Particle pair	Mom. range for 3σ separation
e/π	$p \lesssim 5 \text{ GeV}/c$
π/K	$0.23 \lesssim p \lesssim 20 \text{ GeV}/c$
K/p	$0.82 \lesssim p \lesssim 30 \text{ GeV}/c$

The phototransducer is typically a TPC/wire-chamber combination sensitive to single photoelectrons and having charge division or pads. This construction permits three-dimensional reconstruction of photoelectron origins, which is important for transforming the Čerenkov cone into a ring. Single photoelectrons are generated by doping the TPC gas (for instance, ethane/methane in some proportion) with $\sim 0.05\%$ TMAE [tetrakis(dimethylamino)ethylene] [17], leading to photon absorption lengths along the Čerenkov cone of ~ 30 mm. The readout wires must be equipped with special structures (blinds or wire gates) to prevent photon feedback from avalanches generating cross-talk photoelectrons in the TPC. Drift-gas purity must be maintained to assure mean drift lengths of the order of meters without recombination (*i.e.*, lifetimes of $\gtrsim 100 \mu\text{s}$ at typical drift velocities of $\gtrsim 4 \text{ cm}/\mu\text{s}$). The net $\langle\epsilon_{det}\rangle$'s reach 30%, with the limitation being the TMAE quantum efficiency.

Photon energy cutoffs are set by the TMAE ($E > 5.4 \text{ eV}$), the UV transparency of fused silica glass ($E < 7.4 \text{ eV}$), and the C_6F_{14} ($E < 7.1 \text{ eV}$). With effort one gets $50 \leq N_0 \leq 100$ for complete rings using liquid or gas. This includes losses due to electrostatic shielding wires and window/mirror reflections, but not gross losses caused by total internal reflection or inadequate coverage by the TPC's.

Such numbers allow determination of ring radii to $\sim 0.5\%$ (liquid) and $\sim 2\%$ (gas), leading to the particle species separations quoted above. Since the separation efficiencies may have "holes" as a function of p , detailed calculations are necessary.

12.4. Transition radiation detectors (TRD's)

It is evident from the discussion in the Passages of Particles Through Matter section (Sec. 10 of this *Review*) that transition radiation (TR) only becomes useful for particle detectors when the Lorentz factor $\gamma \gtrsim 10^3$. In practice, TRD's are used to provide e/π separation when $p \gtrsim 1 \text{ GeV}/c$. (The momentum is usually measured elsewhere in the detector.) Since a soft x ray is radiated with about 1% probability per boundary crossing, practical detectors use radiators with several hundred interfaces, *e.g.* foils of lithium or plastic in a gas. Absorption inside the radiator and interference effects between interfaces are important [18,19].

A practical detector is composed of several similar modules, each consisting of a radiator and an x-ray detector. The radiator is made of foils or fibers of a low- Z material (for low absorption) in a low- Z gas such as helium. The x-ray detector is usually a wire chamber operated with a xenon-rich mixture in order to obtain a high conversion efficiency. As transition radiation is emitted at small angles, the chamber usually detects the sum of the ionization of the particle and of converted TR photons. The discrimination between electrons and pions can be based on the charges measured in each set, or on more sophisticated methods using pulse-shape analysis. The TRD in the $D\theta$ experiment serves as an example [20,21].

The major factor in the performance of a TRD is its overall length. Very roughly, the pion rejection factor for a detector with 90% electron efficiency is $10(L/20 \text{ cm})$, where L is the overall length of a radiator with foils. Radiators with fibers are easier to build, but generally provide a rejection factor which is at least a factor of two lower.

12.5. Silicon photodiodes and particle detectors

Silicon detectors are p - n junction diodes operated at reverse bias. This forms a sensitive region depleted of mobile charge and sets up an electric field that sweeps charge liberated by radiation to the electrodes. The thickness of the depleted region is

$$W = \sqrt{\frac{2\epsilon(V + V_{bi})}{ne}} = \sqrt{2\rho\epsilon(V + V_{bi})}, \quad (12.6)$$

where V = external bias voltage

V_{bi} = "built-in" voltage ($\approx 0.8 \text{ V}$ for resistivities typically used in detectors)

n = doping concentration

e = electron charge

ϵ = dielectric constant = $11.9 \epsilon_0 \approx 1 \text{ pF}/\text{cm}$

ρ = resistivity (typically 1 – $10 \text{ k}\Omega \text{ cm}$)

μ = charge carrier mobility

= $1350 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ for electrons (n -type material)

= $450 \text{ cm}^2 \text{ V}^{-1} \text{ s}^{-1}$ for holes (p -type material)

or

$$W = 0.5 \mu\text{m} \times \sqrt{\rho(V + V_{bi})} \quad \text{for } n\text{-type material}, \quad (12.7)$$

and

$$W = 0.3 \mu\text{m} \times \sqrt{\rho(V + V_{bi})} \quad \text{for } p\text{-type material}, \quad (12.8)$$

where V is in volts and ρ is in $\Omega \text{ cm}$.

The corresponding capacitance per unit area is

$$C = \frac{\epsilon}{W} \approx 1 \text{ [pF/cm]} \frac{1}{W}. \quad (12.9)$$

In strip detectors the capacitance is dominated by the strip-to-strip fringing capacitance of ~ 1 – 1.5 pF cm^{-1} of strip length at a strip pitch of 25 – $50 \mu\text{m}$.

About 3.6 eV is required to create an electron-hole pair. For minimum-ionizing particles, the most probable charge deposition in a

300 μm thick silicon detector is about 4 fC (25000 electrons). Readily available photodiodes have quantum efficiencies $> 70\%$ for wavelengths between 600 nm and 1 μm . UV extended photodiodes have useful efficiency down to 200 nm. In applications in which photodiodes detect light from scintillators, care must be taken so that signal from the scintillator is larger than that produced by particles going through the photodiode.

Collection time decreases with increased depletion voltage, and can be reduced further by operating the detector with “overbias,” *i.e.*, a bias voltage exceeding the value required to fully deplete the device. The collection time is limited by velocity saturation at high fields; at an average field of 10^4 V/cm, the collection times is about 15 ps/ μm for electrons and 30 ps/ μm for holes. In typical strip detectors of 300 μm thickness, electrons are collected within about 8 ns, and holes within about 25 ns.

Position resolution is limited by transverse diffusion during charge collection (typically 5 μm for 300 μm thickness) and by knock-on electrons. Resolutions of 3–4 μm (rms) have been obtained in beam tests. In magnetic fields, the Lorentz drift can increase the spatial spread appreciably (see “Hall effect” in semiconductor textbooks).

Radiation damage occurs through two basic mechanisms:

1. Bulk damage due to displacement of atoms from their lattice sites. This leads to increased leakage current, carrier trapping, and changes in doping concentration. Displacement damage depends on the nonionizing energy loss, *i.e.*, particle type and energy. The dose should be specified as a fluence of particles of a specific type and energy.
2. Surface damage due to charge build-up in surface layers, which leads to increased surface leakage currents. In strip detectors the inter-strip isolation is affected. The effects of charge build-up are strongly dependent on the device structure and on fabrication details. Since the damage is determined directly by the absorbed energy, the dose should be specified in these units (rad or Gray).

The increase in leakage current due to bulk damage is $\Delta i = \alpha \phi$ per unit volume, where ϕ is the particle fluence and α the damage coefficient ($\alpha \approx 2 \times 10^{-17}$ A/cm for minimum ionizing protons and pions after long-term annealing; roughly the same value applies for 1 MeV neutrons). The doping concentration in *n*-type silicon changes as $n = n_0 \exp(-\delta \phi) - \beta \phi$, where n_0 is the initial donor concentration, $\delta \approx 6 \times 10^{14}$ cm² determines donor removal, and $\beta \approx 0.03$ cm⁻¹ describes acceptor creation. This leads to an initial increase in resistivity until type-inversion changes the net doping from *n* to *p*. At this point the resistivity decreases, with a corresponding increase in depletion voltage. The safe operating limit of depletion voltage ultimately limits the detector lifetime. Strip detectors have remained functional at fluences beyond 10^{14} cm⁻² for minimum ionizing protons. At this damage level, charge loss due to recombination and trapping also seems to become significant.

12.6. Proportional and drift chambers

Proportional chamber wire instability. The limit on the voltage V for a wire tension T , due to mechanical effects when the electrostatic repulsion of adjacent wires exceeds the restoring force of wire tension, is given by (SI units) [22]

$$V \leq \frac{s}{\ell C} \sqrt{4\pi\epsilon_0 T}, \quad (12.10)$$

where s , ℓ , and C are the wire spacing, length, and capacitance per unit length. An approximation to C for chamber half-gap t and wire diameter d (good for $s \lesssim t$) gives [23]

$$V \lesssim 59T^{1/2} \left[\frac{t}{\ell} + \frac{s}{\pi\ell} \ln \left(\frac{s}{\pi d} \right) \right], \quad (12.11)$$

where V is in kV, and T is in grams-weight equivalent.

Proportional and drift chamber potentials The potential distributions and fields in a proportional or drift chamber can usually be calculated with good accuracy from the exact formula for the potential

around an array of parallel line charges q (coul/m) along z and located at $y = 0$, $x = 0, \pm s, \pm 2s, \dots$,

$$V(x, y) = -\frac{q}{4\pi\epsilon_0} \ln \left\{ 4 \left[\sin^2 \left(\frac{\pi x}{s} \right) + \sinh^2 \left(\frac{\pi y}{s} \right) \right] \right\}. \quad (12.12)$$

Errors from the presence of cathodes, mechanical defects, TPC-type edge effects, *etc.*, are usually small and are beyond the scope of this review.

12.7. Calorimeters

Electromagnetic calorimeters. The development of electromagnetic showers is discussed in the “Passage of Particles Through Matter” section (Sec. 10 of this *Review*). Formulae are given for the approximate description of average showers, but since the physics of electromagnetic showers is well understood, detailed and reliable Monte Carlo simulation is possible. EGS4 has emerged as the standard [24].

The resolution of sampling calorimeters (hadronic and electromagnetic) is usually dominated by sampling fluctuations, leading to fractional resolution σ/E scaling inversely as the square root of the incident energy. Homogenous calorimeters, such as solid NaI(Tl), will in general not have resolution varying as $1/\sqrt{E}$. At high energies deviations from $1/\sqrt{E}$ occur because of noise, pedestal fluctuations, nonuniformities, calibration errors, and incomplete shower containment. Such effects are usually included by adding a constant term to σ/E , either in quadrature or (incorrectly) directly. In the case of the hadronic cascades discussed below, noncompensation also contributes to the constant term.

In Table 12.4 we give resolution as measured in detectors using typical EM calorimeter technologies. In almost all cases the installed calorimeters yield worse resolution than test beam prototypes for a variety of practical reasons. Where possible actual detector performance is given. For a fixed number of radiation lengths, the FWHM in sandwich detectors would be expected to be proportional to $\sqrt{\ell}$ for ℓ (= plate thickness) ≥ 0.2 radiation lengths [25].

Given sufficient transverse granularity early in the calorimeter, position resolution of the order of a millimeter can be obtained.

Table 12.4: Resolution of typical electromagnetic calorimeters. E is in GeV.

Detector	Resolution
NaI(Tl) (Crystal Ball [26]; 20 X_0)	2.7%/ $E^{1/4}$
Lead glass (OPAL [27])	5%/ \sqrt{E}
Lead-liquid argon (NA31 [28]; 80 cells: 27 X_0 , 1.5 mm Pb + 0.6 mm Al + 0.8 mm G10 + 4 mm LA)	7.5%/ \sqrt{E}
Lead-scintillator sandwich (ARGUS [29], LAPP-LAL [30])	9%/ \sqrt{E}
Lead-scintillator spaghetti (CERN test module) [31]	13%/ \sqrt{E}
Proportional wire chamber (MAC; 32 cells: 13 X_0 , 2.5 mm typemetal + 1.6 mm Al) [32]	23%/ \sqrt{E}

Hadronic calorimeters [33,34]. The length scale appropriate for hadronic cascades is the nuclear interaction length, given very roughly by

$$\lambda_I \approx 35 \text{ g cm}^{-2} A^{1/3}. \quad (12.13)$$

Longitudinal energy deposition profiles are characterized by a sharp peak near the first interaction point (from the fairly local deposition of EM energy resulting from π^0 's produced in the first interaction), followed by a more gradual development with a maximum at

$$x/\lambda_I \equiv t_{\text{max}} \approx 0.2 \ln(E/1 \text{ GeV}) + 0.7 \quad (12.14)$$

as measured from the front of the detector.

The depth required for containment of a fixed fraction of the energy also increases logarithmically with incident particle energy. The thickness of iron required for 95% (99%) containment of cascades induced by single hadrons is shown in Fig. 12.1 [35]. Two of the sets of data are from large neutrino experiments, while the third is from a commonly used parametrization. Depths as measured in nuclear interaction lengths presumably scale to other materials. From the same data it can be concluded that the requirement that 95% of the energy in 95% of the showers be contained requires 40 to 50 cm (2.4 to 3.0 λ_I) more material than for an average 95% containment.




Figure 12.1: Required calorimeter thickness for 95% and 99% hadronic cascade containment in iron, on the basis of data from two large neutrino detectors and the parametrization of Bock *et al.* [35].

The transverse dimensions of hadronic showers also scale as λ_I , although most of the energy is contained in a narrow core.

The energy deposit in a hadronic cascade consists of a prompt EM component due to π^0 production and a slower component mainly due to low-energy hadronic activity. In general, these energy depositions are converted to electrical signals with different efficiencies [36]. The ratio of the conversion efficiencies is usually called the intrinsic e/h ratio. If $e/h = 1.0$ the calorimeter is said to be *compensating*. If it differs from unity by more than 5% or 10%, detector performance is compromised because of fluctuations in the π^0 content of the cascades. Problems include:

- A skewed signal distribution;
- A response ratio for electrons and hadrons (the “ e/π ratio”) which is different from unity and depends upon energy;
- A nonlinear response to hadrons (the response per GeV is proportional to the reciprocal of e/π);
- A constant contribution to detector resolution, almost proportional to the degree of noncompensation. The coefficient relating the constant term to $|1 - e/h|$ is 14% according to FLUKA simulations, and 21% according to Wigman’s calculations [33].

In most cases e/h is greater than unity, particularly if little hydrogen is present or if the gate time is short. This is because much of the low-energy hadronic energy is “hidden” in nuclear binding energy release, low-energy spallation products, *etc.* Partial correction for these losses occurs in a sampling calorimeter with thick plates, because a disproportionate fraction of electromagnetic energy is deposited in the inactive region. For this reason, it is very unlikely that a fully sensitive detector such as BGO or glass can be made compensating.

Compensation has been demonstrated in calorimeters with 2.5 mm scintillator sheets sandwiched between 3 mm depleted uranium plates [38] or 10 mm lead plates [39]; resolutions σ/E of $0.34/\sqrt{E}$

and $0.44/\sqrt{E}$ were obtained for these cases (E in GeV). The former was shown to be linear to within 2% over three orders of magnitude in energy, with approximately Gaussian signal distributions.

dE/dx resolution in argon. Particle identification by dE/dx is dependent on the width of the distribution. For relativistic incident particles with charge e in a multiple-sample Ar gas counter with no lead [40],

$$\frac{dE}{dx} \Big|_{\text{FWHM}} / \frac{dE}{dx} \Big|_{\text{most probable}} = 0.96 N^{-0.46} (xp)^{-0.32}, \quad (12.15)$$

where N = number of samples, x = thickness per sample (cm), p = pressure (atm.). Most commonly used chamber gases (except Xe) give approximately the same resolution.

Free electron drift velocities in liquid ionization chambers [41–44]. Velocity as a function of electric field strength is given in Fig. 12.2.




Figure 12.2: Electron drift velocity as a function of field strength for commonly used liquids.

12.8. Measurement of particle momenta in a uniform magnetic field [45]

The trajectory of a particle with momentum p (in GeV/c) and charge ze in a constant magnetic field \vec{B} is a helix, with radius of curvature R and pitch angle λ . The radius of curvature and momentum component perpendicular to \vec{B} are related by

$$p \cos \lambda = 0.3 z B R, \quad (12.16)$$

where B is in tesla and R is in meters.

The distribution of measurements of the curvature $k \equiv 1/R$ is approximately Gaussian. The curvature error for a large number of uniformly spaced measurements on the trajectory of a charged particle in a uniform magnetic field can be approximated by

$$(\delta k)^2 = (\delta k_{\text{res}})^2 + (\delta k_{\text{ms}})^2, \quad (12.17)$$

where δk = curvature error

δk_{res} = curvature error due to finite measurement resolution

δk_{ms} = curvature error due to multiple scattering.

If many (≥ 10) uniformly spaced position measurements are made along a trajectory in a uniform medium,

$$\delta k_{\text{res}} = \frac{\epsilon}{L^2} \sqrt{\frac{720}{N+4}}, \quad (12.18)$$

where N = number of points measured along track

L' = the projected length of the track onto the bending plane
 ϵ = measurement error for each point, perpendicular to the trajectory.

If a vertex constraint is applied at the origin of the track, the coefficient under the radical becomes 320.

The contribution due to multiple Coulomb scattering is approximately

$$\delta k_{\text{ms}} \approx \frac{(0.016)(\text{GeV}/c)z}{Lp\beta \cos^2 \lambda} \sqrt{\frac{L}{X_0}}, \quad (12.19)$$

where p = momentum (GeV/ c)

z = charge of incident particle in units of e

L = the total track length

X_0 = radiation length of the scattering medium (in units of length; the X_0 defined elsewhere must be multiplied by density)

β = the kinematic variable v/c .

More accurate approximations for multiple scattering may be found in the section on Passage of Particles Through Matter (Sec. 10 of this *Review*). The contribution to the curvature error is given approximately by $\delta k_{\text{ms}} \approx 8s_{\text{plane}}^{\text{rms}}/L^2$, where $s_{\text{plane}}^{\text{rms}}$ is defined there.

12.9. Superconducting solenoids for collider detectors

12.9.1. Basic (approximate) equations: In all cases SI units are assumed, so that B is in tesla, E is in joules, dimensions are in meters, and $\mu_0 = 4\pi \times 10^{-7}$.

Magnetic field. The magnetic field at the center of a solenoid of length L and radius R , having N total turns and a current I is

$$B(0,0) = \frac{\mu_0 NI}{\sqrt{L^2 + 4R^2}}. \quad (12.20)$$

Stored energy. The energy stored in the magnetic field of any magnet is calculated by integrating B^2 over all space:

$$E = \frac{1}{2\mu_0} \int B^2 dV. \quad (12.21)$$

For a solenoid with an iron flux return in which the magnetic field is $< 2T$, the field in the aperture is approximately uniform and equal to $\mu_0 NI/L$. If the thickness of the coil is small, (which is the case if it is superconducting), then

$$E \approx (\pi/2\mu_0)B^2 R^2 L. \quad (12.22)$$

Cost of a superconducting solenoid [46]:

$$\text{Cost (in M\$)} = 0.523 [(E/(1 \text{ MJ}))^{0.662}] \quad (12.23)$$

Magnetostatic computer programs. It is too difficult to solve the Biot-Savart equation for a magnetic circuit which includes iron components and so iterative computer programs are used. These include POISSON, TOSCA [47], and ANSYS [48].

12.9.2. Scaling laws for thin solenoids: For a detector in which the calorimetry is outside the aperture of the solenoid, the coil must be thin in terms of radiation and absorption lengths. This usually means that the coil is superconducting and that the vacuum vessel encasing it is of minimum real thickness and fabricated of a material with long radiation length. There are two major contributors to the thickness of a thin solenoid:




Figure 12.3: Ratio of stored energy to cold mass for existing thin detector solenoids.

1. The conductor, consisting of the current-carrying superconducting material (usually Cu/Nb-Ti) and the quench protecting stabilizer (usually aluminum), is wound on the inside of a structural support cylinder (usually aluminum also). This package typically represents about 60% of the total thickness in radiation lengths. The thickness scales approximately as $B^2 R$.
2. Approximately another 25% of the thickness of the magnet comes from the outer cylindrical shell of the vacuum vessel. Since this shell is susceptible to buckling collapse, its thickness is determined by the diameter, length, and the modulus of the material of which it is fabricated. When designing this shell to a typical standard, the real thickness is

$$t = P_c D^{2.5} [(L/D) - 0.45(t/D)^{0.5}] / 2.6Y^{0.4}, \quad (12.24)$$

where t = shell thickness (in), D = shell diameter (in), L = shell length (in), Y = modulus of elasticity (psi), and P_c = design collapse pressure (= 30 psi). For most large-diameter detector solenoids, the thickness to within a few percent is given by [49]

$$t = P_c D^{2.5} (L/D) / 2.6Y^{0.4}. \quad (12.25)$$

12.9.3. Properties of collider detector solenoids: The physical dimensions, central field, stored energy and thickness in radiation lengths normal to the beam line of the superconducting solenoids associated with the major colliders are given in Table 12.5.

Table 12.5: Properties of superconducting collider detector solenoids.

Experiment-Lab	Field (T)	Bore Dia (m)	Length (m)	Energy (MJ)	Thickness (X_0)
CDF-Fermilab	1.5	2.86	5.07	30	0.86
Topaz-KEK	1.2	2.72	5.4	19.5	0.70
Venus-KEK	0.75	3.4	5.64	12	0.52
Cleo II-Cornell	1.5	2.9	3.8	25	2.5
Aleph-CERN	1.5	5.0	7.0	130	1.7
Delphi-CERN	1.2	5.2	7.4	109	4.0
H1-DESY	1.2	5.2	5.75	120	1.2
Zeus-DESY	1.8	1.72	2.85	10.5	0.9

The ratio of stored energy to cold mass (E/M) is a useful performance measure. One would like the cold mass to be as small as possible to minimize the thickness, but temperature rise during a quench must also be minimized. Ratios as large as 8 kJ/kg may be possible (final temperature of 80 K after a fast quench with homogenous energy dump), but some contingency is desirable. This quantity is shown as a function of total stored energy for some major collider detectors in Fig. 12.3.

12.10. Radiation levels in detectors at hadron colliders

An SSC Central Design Group task force made a study of radiation levels to be expected in SSC detectors [50]. Its model assumed

- The machine luminosity at $\sqrt{s} = 40$ TeV is $\mathcal{L} = 10^{33}$ cm⁻²s⁻¹, and the pp inelastic cross section is $\sigma_{\text{inel}} = 100$ mb. This luminosity is effectively achieved for 10^7 s yr⁻¹. The interaction rate is thus 10^8 s⁻¹, or 10^{15} yr⁻¹;
- All radiation comes from pp collisions at the interaction point;
- The charged particle distribution is (a) flat in pseudorapidity for $|\eta| < 6$ and (b) has a momentum distribution whose perpendicular component is independent of rapidity, which is taken as independent of pseudorapidity:

$$\frac{d^2 N_{\text{ch}}}{d\eta dp_{\perp}} = H f(p_{\perp}) \quad (12.26)$$

(where $p_{\perp} = p \sin \theta$). Integrals involving $f(p_{\perp})$ are simplified by replacing $f(p_{\perp})$ by $\delta(p_{\perp} - \langle p_{\perp} \rangle)$; in the worst case this approximation introduces an error of less than 10%;

- Gamma rays from π^0 decay are as abundant as charged particles. They have approximately the same η distribution, but half the mean momentum;
- At the SSC ($\sqrt{s} = 40$ TeV), $H \approx 7.5$ and $\langle p_{\perp} \rangle \approx 0.6$ GeV/c; assumed values at other energies are given in Table 12.7. Together with the model discussed above, these values are thought to describe particle production to within a factor of two or better.

It then follows that the flux of charged particles from the interaction point passing through a normal area da located a distance r_{\perp} from the beam line is given by

$$\frac{dN_{\text{ch}}}{da} = \frac{1.2 \times 10^8 \text{ s}^{-1}}{r_{\perp}^2} \quad (12.27)$$

In a typical organic material, a relativistic charged particle flux of 3×10^9 cm⁻² produces an ionizing radiation dose of 1 Gy, where 1 Gy \equiv 1 joule kg⁻¹ (= 100 rads). The above result may thus be rewritten as dose rate,

$$\dot{D} = \frac{0.4 \text{ MGy yr}^{-1}}{(r_{\perp}/1 \text{ cm})^2} \quad (12.28)$$

If a magnetic field is present, “loopers” may increase this dose rate by a factor of two.

In a medium in which cascades can develop, the ionizing dose or neutron fluence is proportional to dN_{ch}/da multiplied by $\langle E \rangle^{\alpha}$, where $\langle E \rangle$ is the mean energy of the particles going through da and the power α is slightly less than unity. Since $E \approx p = p_{\perp}/\sin \theta$ and $r_{\perp} = r \sin \theta$, the above expression for dN_{ch}/da becomes

$$\text{Dose or fluence}^* = \frac{A}{r^2} \cosh^{2+\alpha} \eta = \frac{A}{r^2 \sin^{2+\alpha} \theta} \quad (12.29)$$

The constant A contains the total number of interactions $\sigma_{\text{inel}} \int \mathcal{L} dt$, so the ionizing dose or neutron fluence at another accelerator scales as $\sigma_{\text{inel}} \int \mathcal{L} dt H \langle p_{\perp} \rangle^{\alpha}$.

The dose or fluence in a calorimeter scales as $1/r^2$, as does the neutron fluence inside a central cavity with characteristic dimension r .

Under all conditions so far studied, the neutron spectrum shows a broad log-normal distribution peaking at just under 1 MeV. In a 2 m radius central cavity of a detector with coverage down to $|\eta| = 3$, the average neutron flux is 2×10^{12} cm⁻²yr⁻¹, including secondary scattering contributions.

Values of A and α are given in Table 12.6 for several relevant situations. Examples of scaling to other accelerators are given in Table 12.7. It should be noted that the assumption that all radiation comes from the interaction point does not apply to the present generation of accelerators.

Table 12.6: Coefficients $A/(100 \text{ cm})^2$ and α for the evaluation of calorimeter radiation levels at cascade maxima under SSC nominal operating conditions. At a distance r and angle θ from the interaction point the annual fluence or dose is $A/(r^2 \sin^{2+\alpha} \theta)$.

Quantity	$A/(100 \text{ cm})^2$	Units	$\langle p_{\perp} \rangle$	α
Neutron flux	1.5×10^{12}	cm ⁻² yr ⁻¹	0.6 GeV/c	0.67
Dose rate from photons	124	Gy yr ⁻¹	0.3 GeV/c	0.93
Dose rate from hadrons	29	Gy yr ⁻¹	0.6 GeV/c	0.89

Table 12.7: A rough comparison of beam-collision induced radiation levels at the Tevatron, UNK, high-luminosity LHC, and SSC.

	Tevatron	UNK-3	LHC	SSC
\sqrt{s} (TeV)	1.8	6	16	40
\mathcal{L}_{nom} (cm ⁻² s ⁻¹)	2×10^{30}	4×10^{32}	4×10^{34a}	1×10^{33}
σ_{inel}	59 mb	80 mb	86 mb	100 mb
H	4.1	4.5	6.3	7.5
$\langle p_{\perp} \rangle$ (GeV/c)	0.46	0.52	0.55	0.60
Relative dose rate ^b	5×10^{-4}	0.2	27	1

^a High-luminosity option.

^b Proportional to $\mathcal{L}_{\text{nom}} \sigma_{\text{inel}} H \langle p_{\perp} \rangle^{0.7}$

The constant A includes factors evaluated with cascade simulation programs as well as constants describing particle production at the interaction point. It is felt that each could introduce an error as large as a factor of two in the results.

References:

- * *Dose* is the time integral of *dose rate*, and *fluence* is the time integral of *flux*.
- 1. *Experimental Techniques in High Energy Physics*, T. Ferbel (ed.) (Addison-Wesley, Menlo Park, CA, 1987).
- 2. *Methods of Experimental Physics*, L.C.L. Yuan and C.-S. Wu, editors, Academic Press, 1961, Vol. 5A, p. 127.
- 3. Nuclear Enterprises Catalogue.
- 4. R.K. Swank, *Ann. Rev. Nucl. Sci.* **4**, 137 (1954); G.T. Wright, *Proc. Phys. Soc.* **B68**, 929 (1955).
- 5. M. Laval *et al.*, *Nucl. Instr. and Meth.* **206**, 169 (1983).
- 6. M. Moszynski *et al.*, *Nucl. Instr. and Meth.* **A226**, 534 (1984).
- 7. E. Blucher *et al.*, *Nucl. Instr. and Meth.* **A249**, 201 (1986).
- 8. C. Bebek, *Nucl. Instr. and Meth.* **A265**, 258 (1988).
- 9. S. Kubota *et al.*, *Nucl. Instr. and Meth.* **A268**, 275 (1988).
- 10. B. Ádeva *et al.*, *Nucl. Instr. and Meth.* **A289**, 35 (1990).
- 11. I. Holl, E. Lorentz, G. Mageras, *IEEE Trans. Nucl. Sci.* **35**, 105 (1988).
- 12. J. Litt and R. Meunier, *Ann. Rev. Nucl. Sci.* **23**, 1 (1973).
- 13. D. Bartlett *et al.*, *Nucl. Instr. and Meth.* **A260**, 55 (1987).
- 14. P. Duteil *et al.*, *Review of Scientific Instruments* **35**, 1523 (1964).
- 15. M. Cavalli-Sforza *et al.*, *Construction and Testing of the SLC Čerenkov Ring Imaging Detector*, *IEEE* **37**, N3:1132 (1990).
- 16. E.G. Anassontzis *et al.*, *Recent Results from the DELPHI Barrel Ring Imaging Čerenkov Counter*, *IEEE* **38**, N2:417 (1991).
- 17. R.T. Rewick *et al.*, *Anal. Chem.* **60**, 2095 (1989).
- 18. X. Artru *et al.*, *Phys. Rev.* **D12**, 1289 (1975).
- 19. G.M. Garibian *et al.*, *Nucl. Instr. and Meth.* **125**, 133 (1975).
- 20. J.F. Detoeuf *et al.*, *Nucl. Instr. and Meth.* **A265**, 157 (1988).
- 21. Y. Ducros *et al.*, *Nucl. Instr. and Meth.* **A277**, 401 (1989).
- 22. T. Trippe, CERN NP Internal Report 69-18 (1969).
- 23. S. Parker and R. Jones, LBL-797 (1972);

- P. Morse and H. Feshbach, *Methods of Theoretical Physics*, McGraw-Hill, New York, 1953, p. 1236.
24. W.R. Nelson, H. Hirayama and D.W.O. Rogers, "The EGS4 Code System," SLAC-265, Stanford Linear Accelerator Center (Dec. 1985).
 25. D. Hitlin *et al.*, Nucl. Instr. and Meth. **137**, 225 (1976). See also W. J. Willis and V. Radeka, Nucl. Instr. and Meth. **120**, 221 (1974), for a more detailed discussion.
 26. E. Bloom and C. Peck, Ann. Rev. Nucl. and Part. Sci. **33**, 143 (1983).
 27. M.A. Akrawy *et al.*, Nucl. Instr. and Meth. **A290**, 76 (1990).
 28. H. Burkhardt *et al.*, Nucl. Instr. and Meth. **A268**, 116 (1988).
 29. W. Hoffman *et al.*, Nucl. Instr. and Meth. **163**, 77 (1979).
 30. M.A. Schneegans *et al.*, Nucl. Instr. and Meth. **193**, 445 (1982).
 31. C. Fabjan and R. Wigmans, Rept. Prog. Phys. **52**, 1519 (1989).
 32. J.V. Allaby *et al.*, Nucl. Instr. and Meth. **A281**, 291 (1989).
 33. R. Wigmans, Nucl. Instr. and Meth. **A259**, 389 (1987).
 34. R. Wigmans, Nucl. Instr. and Meth. **A265**, 273 (1988).
 35. D. Binting, in *Proceedings of the Workshop on Calorimetry for the Supercollider*, Tuscaloosa, AL, March 13–17, 1989, edited by R. Donaldson and M.G.D. Gilchriese (World Scientific, Teaneck, NJ, 1989), p. 91.
 36. T.A. Gabriel, D.E. Groom, P.K. Job, N.V. Mokhov, and G.R. Stevenson, Nucl. Instr. and Meth. **A338**, 336 (1994).
 37. R.K. Bock, T. Hansl-Kozanecka, and T.P. Shah, Nucl. Instr. and Meth. **186**, 533 (1981).
 38. T. Akesson *et al.*, Nucl. Instr. and Meth. **A262**, 243 (1987).
 39. E. Bernardi *et al.*, Nucl. Instr. and Meth. **A262**, 229 (1987).
 40. W.W.M. Allison and J.H. Cobb, "Relativistic Charged Particle Identification by Energy-Loss," Ann. Rev. Nucl. Sci. **30**, 253 (1980), see p. 287.
 41. E. Shibamura *et al.*, Nucl. Instr. and Meth. **131**, 249 (1975).
 42. T.G. Ryan and G.R. Freeman, J. Chem. Phys. **68**, 5144 (1978).
 43. W.F. Schmidt, "Electron Migration in Liquids and Gases," HMI B156 (1974).
 44. A.O. Allen, "Drift Mobilities and Conduction Band Energies of Excess Electrons in Dielectric Liquids," NSRDS-NBS-58 (1976).
 45. R.L. Gluckstern, Nucl. Instr. and Meth. **24**, 381 (1963).
 46. M.A. Green, R.A. Byrns, and S.J. St. Lorant, "Estimating the cost of superconducting magnets and the refrigerators needed to keep them cold," in *Advances in Cryogenic Engineering*, Vol. 37, Plenum Press, New York (1992).
 47. Vector Fields, Inc., 1700 N. Farnsworth Ave., Aurora, IL.
 48. Swanson Analysis Systems, Inc., P.O. Box 65, Johnson Rd., Houston, PA.
 49. CGA-341-1987, "Standard for insulated cargo tank specification for cryogenic liquids," Compressed Gas Association, Inc., Arlington, VA (1987).
 50. Report of the Task Force on Radiation Levels in the SSC Interaction Regions, SSC Central Design Group Report SSC-SR-1033 (June 1988). Abridged versions of this report may be found in Nucl. Instr. and Meth. **A279**, 1 (1989) and in *Proceedings of the 1988 Summer Study on High Energy Physics in the 1990's*, Snowmass, CO, June 27 – July 15, 1990, edited by F.J. Gilman and S. Jensen, (World Scientific, Teaneck, NJ, 1989).

13. RADIOACTIVITY & RADIATION PROTECTION

Revised Sept. 1991 with assistance from N.A. Greenhouse.

The International Commission on Radiation Units and Measurements (ICRU) recommends the use of SI units. Therefore we list SI units first, followed by cgs (or other common) units in parentheses, where they differ.

- **Unit of activity** = becquerel (curie):
1 Bq = 1 disintegration s^{-1} [= $1/(3.7 \times 10^{10})$ Ci]
- **Unit of absorbed dose** = gray (rad):
1 Gy = 1 joule kg^{-1} (= 10^4 erg g^{-1} = 100 rad)
= 6.24×10^{12} MeV kg^{-1} deposited energy
- **Unit of exposure**, the quantity of x - or γ - radiation at a point in space integrated over time, in terms of charge of either sign produced by showering electrons in a small volume of air about the point:
= 1 coul kg^{-1} of air (roentgen; 1 R = 2.58×10^{-4} coul kg^{-1})
= 1 esu cm^{-3} (= 87.8 erg released energy per g of air)

Implicit in the definition is the assumption that the small test volume is embedded in a sufficiently large uniformly irradiated volume that the number of secondary electrons entering the volume equals the number leaving. This unit is somewhat historical, but appears on many measuring instruments.

- **Unit of equivalent dose** (for biological damage) = sievert [= 100 rem (roentgen equivalent for man)]: Equivalent dose in Sv = absorbed dose in grays $\times w_R$, where w_R (radiation weighting factor, formerly the quality factor Q) expresses long-term risk (primarily cancer and leukemia) from low-level chronic exposure. It depends upon the type of radiation and other factors, as follows [1]:

Table 13.1: Radiation weighting factors.

Radiation	w_R
X- and γ -rays, all energies	1
Electrons and muons, all energies	1
Neutrons < 10 keV	5
10–100 keV	10
> 100 keV to 2 MeV	20
2–20 MeV	10
> 20 MeV	5
Protons (other than recoils) > 2 MeV	5
Alphas, fission fragments, & heavy nuclei	20

- **Natural annual background**, all sources: Most world areas, whole-body equivalent dose rate \approx (0.4–4) mSv (40–400 millirems). Can range up to 50 mSv (5 rems) in certain areas. U.S. average \approx 3.6 mSv, including \approx 2 mSv (\approx 200 mrem) from inhaled natural radioactivity, mostly radon and radon daughters (0.1–0.2 mSv in open areas. Average is for a typical house and varies by more than an order of magnitude. It can be more than two orders of magnitude higher in poorly ventilated mines).

- **Cosmic ray background** in counters (Earth's surface): $\sim 1 \text{ min}^{-1} \text{ cm}^{-2} \text{ sr}$. For more accurate estimates and details, see the Cosmic Rays section (Sec. 14 of this *Review*).

- **Fluxes** (per cm^2) to deposit one Gy, assuming uniform irradiation:
 \approx (**charged particles**) $6.24 \times 10^9 / (dE/dx)$, where dE/dx (MeV $g^{-1} \text{ cm}^2$), the energy loss per unit length, may be obtained from the Mean Range and Energy Loss figures.
 $\approx 3.5 \times 10^9 \text{ cm}^{-2}$ minimum-ionizing singly-charged particles in carbon.

- \approx (**photons**) $6.24 \times 10^9 / [Ef/\lambda]$, for photons of energy E (MeV), attenuation length λ ($g \text{ cm}^{-2}$) (see Photon Attenuation Length figure), and fraction $f \lesssim 1$ expressing the fraction of the photon's energy deposited in a small volume of thickness $\ll \lambda$ but large enough to contain the secondary electrons.

- $\approx 2 \times 10^{11}$ photons cm^{-2} for 1 MeV photons on carbon ($f \approx 1/2$).
(Quoted fluxes are good to about a factor of 2 for all materials.)

- **Recommended limits to exposure (whole-body dose):***

CERN: 15 mSv yr^{-1}

U.K.: 15 mSv yr^{-1}

U.S.: 50 mSv yr^{-1} (5 rem yr^{-1})[†]

- **Lethal dose:** Whole-body dose from penetrating ionizing radiation resulting in 50% mortality in 30 days (assuming no medical treatment) 2.5–3.0 Gy (250–300 rads), as measured internally on body longitudinal center line. Surface dose varies due to variable body attenuation and may be a strong function of energy.

For a recent review, see E. Pochin, *Nuclear Radiation: Risks and Benefits* (Clarendon Press, Oxford, 1983).

- * The ICRP recommendation [1] is 20 mSv yr^{-1} averaged over 5 years, with the dose in any one year \leq 50 mSv.

[†] Many laboratories in the U.S. and elsewhere set lower limits.

Reference:

1. ICRP Publication 60, *1990 Recommendation of the International Commission on Radiological Protection* Pergamon Press (1991).

14. COSMIC RAY FLUXES

The fluxes of particles of different types depend at the $\sim 10\%$ level on the latitude, their energy, and the conditions of measurement. Some typical sea-level values [1] for charged particles are given below:

I_v flux per unit solid angle per unit horizontal area about vertical direction

$$\equiv j(\theta = 0, \phi) [\theta = \text{zenith angle, } \phi = \text{azimuthal angle}] ;$$

J_1 total flux crossing unit horizontal area from above

$$\equiv \int_{\theta \leq \pi/2} j(\theta, \phi) \cos \theta \, d\Omega \quad [d\Omega = \sin \theta \, d\theta \, d\phi] ;$$

J_2 total flux from above (impinging on a sphere of unit cross-sectional area)

$$\equiv \int_{\theta \leq \pi/2} j(\theta, \phi) \, d\Omega .$$

	<u>Total Intensity</u>	<u>Hard Component</u>	<u>Soft Component</u>	
I_v	1.1×10^2	0.8×10^2	0.3×10^2	$\text{m}^{-2} \text{s}^{-1} \text{sterad}^{-1}$
J_1	1.8×10^2	1.3×10^2	0.5×10^2	$\text{m}^{-2} \text{s}^{-1}$
J_2	2.4×10^2	1.7×10^2	0.7×10^2	$\text{m}^{-2} \text{s}^{-1}$

Very approximately, about 75% of all particles at sea level are penetrating, and are muons (the dominant portion of the hard

component at sea level). The sea-level vertical flux ratio for protons to muons (both charges together) is about 3.5% at 1 GeV/c, decreasing to about 0.5% at 10 GeV/c.

The muon flux at sea level has a mean energy of 2 GeV and a differential spectrum falling as E^{-2} , steepening smoothly to $E^{-3.6}$ above a few TeV. The angular distribution is $\cos^2 \theta$, changing to $\sec \theta$ at energies above a TeV, where θ is the zenith angle at production. The \pm charge ratio is 1.25–1.30. The mean energy of muons originating in the atmosphere is roughly 300 GeV at slant depths \gtrsim a few hundred meters. Beyond slant depths of ~ 10 km water-equivalent, the muons are due primarily to in-the-earth neutrino interactions (roughly $1/8$ interaction $\text{ton}^{-1} \text{year}^{-1}$ for $E_\nu > 300$ MeV, \sim constant throughout the earth) [2]. Muons from this source arrive with a mean energy of 20 GeV, and have a flux of $2 \times 10^{-9} \text{m}^{-2} \text{s}^{-1} \text{sterad}^{-1}$ in the vertical direction and about twice that in the horizontal [3], down at least as far as the deepest mines.

Updated April 1986.

References:

1. B. Rossi, *Rev. Mod. Phys.* **20**, 537 (1948). See also C. Grupen, "News from Cosmic Rays at High Energies," Siegen University preprint SI-84-01, and Allkofer and Grieder, *Cosmic Rays on Earth*, Fachinformationszentrum, Karlsruhe (1984); flux ratio for protons at sea level from G. Brook and A.W. Wolfendale, *Proc. of the Phys. Soc. of London*, Vol. 83 (1964), p. 843.
2. J.G. Learned, F. Reines, and A. Soni, *Phys. Rev. Lett.* **43**, 907 (1979).
3. M.F. Crouch *et al.*, *Phys. Rev.* **D18**, 2239 (1978).

15. COMMONLY USED RADIOACTIVE SOURCES

Table 15.1. Updated November 1993 by E. Browne.

Nuclide	Half-life	Type of decay	Particle		Photon	
			Energy (MeV)	Emission prob.	Energy (MeV)	Emission prob.
$^{22}_{11}\text{Na}$	2.603 y	β^+ , EC	0.545	90%	0.511 Annih. 1.275 100%	
$^{54}_{25}\text{Mn}$	0.855 y	EC			0.835 100% Cr K x rays 26%	
$^{55}_{26}\text{Fe}$	2.73 y	EC			Mn K x rays: 0.00589 25% 0.00649 3.4%	
$^{57}_{27}\text{Co}$	0.744 y	EC			0.014 9% 0.122 86% 0.136 11% Fe K x rays 58%	
$^{60}_{27}\text{Co}$	5.271 y	β^-	0.316	100%	1.173 100% 1.333 100%	
$^{68}_{32}\text{Ge}$	0.742 y	EC			Ga K x rays 44%	
$\rightarrow ^{68}_{31}\text{Ga}$		β^+ , EC	1.899	90%	0.511 Annih. 1.077 3%	
$^{90}_{38}\text{Sr}$	28.5 y	β^-	0.546	100%		
$\rightarrow ^{90}_{39}\text{Y}$		β^-	2.283	100%		
$^{106}_{44}\text{Ru}$	1.020 y	β^-	0.039	100%		
$\rightarrow ^{106}_{45}\text{Rh}$		β^-	3.541	79%	0.512 21% 0.622 10%	
$^{109}_{48}\text{Cd}$	1.267 y	EC	0.063 e^- 0.084 e^- 0.087 e^-	41% 45% 9%	0.088 3.6% Ag K x rays 100%	
$^{113}_{50}\text{Sn}$	0.315 y	EC	0.364 e^- 0.388 e^-	29% 6%	0.392 65% In K x rays 97%	
$^{137}_{55}\text{Cs}$	30.2 y	β^-	0.514 e^- 1.176 e^-	94% 6%	0.662 85%	
$^{133}_{56}\text{Ba}$	10.54 y	EC	0.045 e^- 0.075 e^-	50% 6%	0.081 34% 0.356 62% Cs K x rays 121%	
$^{207}_{83}\text{Bi}$	31.8 y	EC	0.481 e^- 0.975 e^- 1.047 e^-	2% 7% 2%	0.569 98% 1.063 75% 1.770 7% Pb K x rays 78%	
$^{228}_{90}\text{Th}$	1.912 y	6α : $3\beta^-$:	5.341 to 8.785 0.334 to 2.246		0.239 44% 0.583 31% 2.614 36%	
$(\rightarrow ^{224}_{88}\text{Ra} \rightarrow ^{220}_{86}\text{Rn} \rightarrow ^{216}_{84}\text{Po} \rightarrow ^{212}_{82}\text{Pb} \rightarrow ^{212}_{83}\text{Bi} \rightarrow ^{212}_{84}\text{Po})$						
$^{241}_{95}\text{Am}$	432.7 y	α	5.443 5.486	13% 85%	0.060 36% Np L x rays 38%	
$^{241}_{95}\text{Am}/\text{Be}$	432.2 y	6×10^{-5} neutrons (4–8 MeV) and 4×10^{-5} γ 's (4.43 MeV) per Am decay				
$^{244}_{96}\text{Cm}$	18.11 y	α	5.763 5.805	24% 76%	Pu L x rays \sim 9%	
$^{252}_{98}\text{Cf}$	2.645 y	α (97%) Fission (3.1%)	6.076 6.118	15% 82%		
		≈ 20 γ 's/fission; 80% < 1 MeV ≈ 4 neutrons/fission; $\langle E_n \rangle = 2.14$ MeV				

"Emission probability" is the probability per decay of a given emission; because of cascades these may total more than 100%. Only principal emissions are listed. EC means electron capture, and e^- means monoenergetic internal conversion (Auger) electron. The intensity of 0.511 MeV e^+e^- annihilation photons depends upon the number of stopped positrons. Endpoint β^\pm energies are listed. In some cases when energies are closely spaced, the γ -ray values are approximate weighted averages. Radiation from short-lived daughter isotopes is included where relevant.

Half-lives, energies, and intensities are from E. Browne and R.B. Firestone, *Table of Radioactive Isotopes* (John Wiley & Sons, New York, 1986), recent *Nuclear Data Sheets*, and *X-ray and Gamma-ray Standards for Detector Calibration*, IAEA-TECDOC-619 (1991).

Neutron data are from *Neutron Sources for Basic Physics and Applications* (Pergamon Press, 1983).

16. PROBABILITY

Revised June 1994.

16.1. General [1–5]

Let x be a possible outcome of an observation. The probability of x is the relative frequency with which that outcome occurs out of a (possibly hypothetical) large set of similar observations. If x can take any value from a *continuous* range, we write $f(x; \theta) dx$ as the probability of observing x between x and $x + dx$. The function $f(x; \theta)$ is the *probability density function* (p.d.f.) for the *random variable* x , which may depend upon one or more parameters θ . If x can take on only *discrete* values (e.g., the non-negative integers), then $f(x; \theta)$ is itself a probability, but we shall still call it a p.d.f. The p.d.f. is always normalized to unit area (unit sum, if discrete). Both x and θ may have multiple components and are then often written as column vectors. If θ is unknown and we wish to estimate its value from a given set of data measuring x , we may use statistics (see Sec. 17).

The *cumulative distribution function* $F(a)$ is the probability that $x \leq a$:

$$F(a) = \int_{-\infty}^a f(x) dx. \quad (16.1)$$

Here and below, if x is discrete-valued, the integral is replaced by a sum. The endpoint a is expressly included in the integral or sum. Then $0 \leq F(x) \leq 1$, $F(x)$ is nondecreasing, and $\text{Prob}(a < x \leq b) = F(b) - F(a)$. If x is discrete, $F(x)$ is flat except at allowed values of x , where it has discontinuous jumps equal to $f(x)$.

Any function of random variables is itself a random variable, with (in general) a different p.d.f. The *expectation value* of any function $u(x)$ is

$$E[u(x)] = \int_{-\infty}^{\infty} u(x) f(x) dx, \quad (16.2)$$

assuming the integral is finite. For $u(x)$ and $v(x)$ any two functions of x , $E(u + v) = E(u) + E(v)$. For c and k constants, $E(cu + k) = cE(u) + k$.

The n th moment of a distribution is

$$\alpha_n \equiv E(x^n) = \int_{-\infty}^{\infty} x^n f(x) dx, \quad (16.3a)$$

and the n th moment about the mean of x , α_1 , is

$$m_n \equiv E[(x - \alpha_1)^n] = \int_{-\infty}^{\infty} (x - \alpha_1)^n f(x) dx. \quad (16.3b)$$

The most commonly used moments are the mean μ and variance σ^2 :

$$\mu \equiv \alpha_1 \quad (16.4a)$$

$$\sigma^2 \equiv \text{Var}(x) \equiv m_2 = \alpha_2 - \mu^2. \quad (16.4b)$$

The mean is the location of the “center of mass” of the probability density function, and the variance is a measure of the square of its width. Note that $\text{Var}(cx + k) = c^2 \text{Var}(x)$.

Any odd moment about the mean is a measure of the skewness of the p.d.f. The simplest of these is the dimensionless coefficient of skewness $\gamma_1 \equiv m_3/\sigma^3$.

Besides the mean, another useful indicator of the “middle” of the probability distribution is the *median* x_{med} , defined by $F(x_{\text{med}}) = 1/2$; i.e., half the probability lies above and half lies below x_{med} . For a given *sample* of events, x_{med} is the value such that half the events have larger x and half have smaller x (not counting any that have the same x as the median). If the sample median lies between two observed x values, it is set by convention halfway between them. If the p.d.f. for x has the form $f(x - \mu)$ and μ is both mean and median, then for a large number of events N , the variance of the median approaches $1/[4Nf^2(0)]$, provided $f(0) > 0$.

Let x and y be two random variables with a joint p.d.f. $f(x, y)$. The *marginal* p.d.f. of x (the distribution of x with y unobserved) is

$$f_1(x) = \int_{-\infty}^{\infty} f(x, y) dy, \quad (16.5)$$

and similarly for the marginal p.d.f. $f_2(y)$. We define the *conditional* p.d.f. of x , given fixed y , by

$$f_3(y|x) f_1(x) = f(x, y). \quad (16.6a)$$

Similarly, the conditional p.d.f. of y , given fixed x , is

$$f_4(x|y) f_2(y) = f(x, y). \quad (16.6b)$$

From these definitions we immediately obtain Bayes' theorem [2]:

$$f_4(x|y) = \frac{f_3(y|x) f_1(x)}{f_2(y)} = \frac{f_3(y|x) f_1(x)}{\int f_3(y|x) f_1(x) dx}. \quad (16.7)$$

The mean of x is

$$\mu_x = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} x f(x, y) dx dy = \int_{-\infty}^{\infty} x f_1(x) dx, \quad (16.8)$$

and similarly for y . The *correlation* between x and y is a measure of the dependence of one on the other:

$$\rho_{xy} = E[(x - \mu_x)(y - \mu_y)] / \sigma_x \sigma_y = \text{Cov}(x, y) / \sigma_x \sigma_y, \quad (16.9)$$

where σ_x and σ_y are defined in analogy with Eq. (16.4b). It can be shown that $-1 \leq \rho_{xy} \leq 1$. Here “Cov” is the covariance of x and y , a 2-dimensional analogue of the variance.

Two random variables are *independent* if and only if

$$f(x, y) = f_1(x) f_2(y). \quad (16.10)$$

If x and y are independent then $\rho_{xy} = 0$; the converse is not necessarily true except for Gaussian-distributed x and y . If x and y are independent, $E[u(x)v(y)] = E[u(x)]E[v(y)]$, and $\text{Var}(x + y) = \text{Var}(x) + \text{Var}(y)$; otherwise, $\text{Var}(x + y) = \text{Var}(x) + \text{Var}(y) + 2\text{Cov}(x, y)$, and $E(uv)$ does not factor.

In a *change of continuous random variables* from $\mathbf{x} \equiv (x_1, \dots, x_n)$, with p.d.f. $f(\mathbf{x}) = f(x_1, \dots, x_n)$, to $\mathbf{y} \equiv (y_1, \dots, y_n)$, a one-to-one function of the x_i 's, the p.d.f. $g(\mathbf{y}) = g(y_1, \dots, y_n)$ is found by substitution for (x_1, \dots, x_n) in f followed by multiplication by the absolute value of the Jacobian of the transformation; that is,

$$g(\mathbf{y}) = f[w_1(\mathbf{y}), \dots, w_n(\mathbf{y})] |J|. \quad (16.11)$$

The functions w_i express the *inverse* transformation, $x_i = w_i(\mathbf{y})$ for $i = 1, \dots, n$, and $|J|$ is the absolute value of the determinant of the square matrix $J_{ij} = \partial x_i / \partial y_j$. If the transformation from \mathbf{x} to \mathbf{y} is not one-to-one, the situation is more complex and a unique solution may not exist. For example, if the change is to $m < n$ variables, then a given \mathbf{y} may correspond to more than one \mathbf{x} , leading to multiple integrals over the contributions [1].

To change variables for discrete random variables simply substitute; no Jacobian is necessary because now f is a probability rather than a probability density.

If f depends upon a parameter set θ , a change to a different parameter set $\phi_i = \phi_i(\theta)$ is made by simple substitution; no Jacobian is used.

16.2. Characteristic functions

The characteristic function $\phi(u)$ associated with the p.d.f. $f(x)$ is essentially its (inverse) Fourier transform, or the expectation value of $\exp(iux)$:

$$\phi(u) = E(e^{iux}) = \int_{-\infty}^{\infty} e^{iux} f(x) dx . \tag{16.12}$$

It is often useful, and several of its properties follow [1].

It follows from Eqs. (16.3a) and (16.12) that the n th moment of the distribution $f(x)$ is given by

$$i^{-n} \frac{d^n \phi}{du^n} \Big|_{u=0} = \int_{-\infty}^{\infty} x^n f(x) dx = \alpha_n . \tag{16.13}$$

Thus it is often easy to calculate all the moments of a distribution defined by $\phi(u)$, even when $f(x)$ is difficult to obtain.

If $f_1(x)$ and $f_2(y)$ have characteristic functions $\phi_1(u)$ and $\phi_2(u)$, then the characteristic function of the weighted sum $ax + by$ is $\phi_1(au)\phi_2(bu)$. The addition rules for common distributions (e.g., that the sum of two numbers from Gaussian distributions also has a Gaussian distribution) easily follow from this observation.

Let the (partial) characteristic function corresponding to the conditional p.d.f. $f_2(x|z)$ be $\phi_2(u|z)$, and the p.d.f. of z be $f_1(z)$. The characteristic function after integration over the conditional value is

$$\phi(u) = \int \phi_2(u|z) f_1(z) dz . \tag{16.14}$$

Suppose we can write ϕ_2 in the form

$$\phi_2(u|z) = A(u)e^{ig(u)z} . \tag{16.15}$$

Then

$$\phi(u) = A(u)\phi_1(g(u)) . \tag{16.16}$$

The semi-invariants κ_n are defined by

$$\phi(u) = \exp \left(\sum_1^{\infty} \frac{\kappa_n}{n!} (iu)^n \right) = \exp \left(i\kappa_1 u - \frac{1}{2} \kappa_2 u^2 + \dots \right) . \tag{16.17}$$

The κ_n 's are related to the moments α_n and m_n . The first few relations are

$$\begin{aligned} \kappa_1 &= \alpha_1 (= \mu, \text{ the mean}) \\ \kappa_2 &= m_2 = \alpha_2 - \alpha_1^2 (= \sigma^2, \text{ the variance}) \\ \kappa_3 &= m_3 = \alpha_3 - 3\alpha_1\alpha_2 + 2\alpha_1^3 . \end{aligned} \tag{16.18}$$

16.3. Some probability distributions

Table 16.1 gives a number of common probability density functions and corresponding characteristic functions, means, and variances. Further information may be found in Refs. 1-6; Ref. 6 has particularly detailed tables. Monte Carlo techniques for generating each of them may be found in our Sec. 18.3. We comment below on all except the trivial uniform distribution.

16.3.1. Binomial distribution: A random process with exactly two possible outcomes is called a *Bernoulli* process. If the probability of obtaining a certain outcome (a "success") in each trial is p , then the probability of obtaining exactly r successes ($r = 0, 1, 2, \dots, n$) in n trials, without regard to the order of the successes and failures, is given by the binomial distribution $f(r; n, p)$ in Table 16.1. If r successes are observed in n_r trials with probability p of a success, and if s successes are observed in n_s similar trials, then $t = r + s$ is also binomial with $n_t = n_r + n_s$.

16.3.2. Poisson distribution: The Poisson distribution $f(r; \mu)$ gives the probability of finding exactly r events in a given interval of x (e.g., space and time) when the events occur independently of one another and of x at an average rate of μ per the given interval. The variance σ^2 equals μ . It is the limiting case $p \rightarrow 0, n \rightarrow \infty, np = \mu$ of the binomial distribution. The Poisson distribution approaches the Gaussian distribution for large μ .

Two or more Poisson processes (e.g., *signal + background*, with parameters μ_s and μ_b) that independently contribute amounts n_s and n_b to a given measurement will produce an observed number $n = n_s + n_b$, which is distributed according to a new Poisson distribution with parameter $\mu = \mu_s + \mu_b$.

16.3.3. Normal or Gaussian distribution: The normal (or Gaussian) probability density function $f(x; \mu, \sigma^2)$ given in Table 16.1 has mean $\bar{x} = \mu$ and variance σ^2 . Comparison of the characteristic function $\phi(u)$ given in Table 16.1 with Eq. (16.17) shows that all semi-invariants κ_n beyond κ_2 vanish; this is a unique property of the Gaussian distribution. Some properties of the distribution are:

- rms deviation = σ
- probability x in the range $\mu \pm \sigma = 0.6827$
- probability x in the range $\mu \pm 0.6745\sigma = 0.5$
- expectation value of $|x - \mu|, E(|x - \mu|) = (2/\pi)^{1/2}\sigma = 0.7979\sigma$
- half-width at half maximum = $(2 \ln 2)^{1/2}\sigma = 1.177\sigma$

The cumulative distribution, Eq. (16.1), for a Gaussian with $\mu = 0$ and $\sigma^2 = 1$ is related to the error function $\text{erf}(y)$ by

$$F(x; 0, 1) = \frac{1}{2} \left[1 + \text{erf}(x/\sqrt{2}) \right] . \tag{16.19}$$

The error function is tabulated in Ref. 6 and is available in computer math libraries and personal computer spreadsheets. For a mean μ and variance σ^2 , replace x by $(x - \mu)/\sigma$. The probability of x in a given range can be calculated with Eq. (17.29).

For x and y independent and normally distributed, $z = ax + by$ obeys $f(z; a\mu_x + b\mu_y, a^2\sigma_x^2 + b^2\sigma_y^2)$; that is, the weighted means and variances add.

The Gaussian gets its importance in large part from the *central limit theorem*: If a continuous random variable x is distributed according to any p.d.f. with finite mean and variance, then the sample mean, \bar{x}_n , of n observations of x will have a p.d.f. that approaches a Gaussian as n increases. Therefore the end result $\sum^n x_i \equiv n\bar{x}_n$ of a large number of small fluctuations x_i will be distributed as a Gaussian, even if the x_i themselves are not.

For \mathbf{x} a set of n (not necessarily independent) Gaussian random variables x_i arranged into a column vector, the joint p.d.f. is the *multivariate Gaussian*:

$$\begin{aligned} f(\mathbf{x}; \boldsymbol{\mu}, V) &= \frac{1}{(2\pi)^{n/2}} |V|^{-1/2} \\ &\times \exp \left[-\frac{1}{2} (\mathbf{x} - \boldsymbol{\mu})^T V^{-1} (\mathbf{x} - \boldsymbol{\mu}) \right] , \quad |V| \neq 0 . \end{aligned} \tag{16.20}$$

Here V is the *covariance matrix* of the x 's, with $V_{ii} = \text{Var}(x_i)$ and $V_{ij} = E[(x_i - \mu_i)(x_j - \mu_j)] \equiv \rho_{ij} \sigma_i \sigma_j$, and $|V|$ is the determinant of V . (If V is singular, there is a linear relation among some of the variables, and so one should eliminate dependent variables and work with an independent set.) The quantity ρ_{ij} is the correlation coefficient for x_i and x_j , and $|\rho_{ij}|^2 \leq 1$. For $n = 2, f(\mathbf{x}; \boldsymbol{\mu}, V)$ is

$$\begin{aligned} f(x_1, x_2; \mu_1, \mu_2, \sigma_1, \sigma_2, \rho) &= \frac{1}{2\pi\sigma_1\sigma_2\sqrt{1-\rho^2}} \\ &\times \exp \left\{ \frac{-1}{2(1-\rho^2)} \left[\frac{(x_1 - \mu_1)^2}{\sigma_1^2} - \frac{2\rho(x_1 - \mu_1)(x_2 - \mu_2)}{\sigma_1\sigma_2} \right. \right. \\ &\quad \left. \left. + \frac{(x_2 - \mu_2)^2}{\sigma_2^2} \right] \right\} . \end{aligned} \tag{16.21}$$

Table 16.1: Some common probability density functions, with corresponding characteristic functions and means and variances. In the Table, $\Gamma(k)$ is the gamma function, equal to $(k - 1)!$ when k is an integer.

Distribution	Probability density function f (variable; parameters)	Characteristic function $\phi(u)$	Mean	Variance σ^2
Uniform	$f(x; a, b) = \begin{cases} 1/(b - a) & a \leq x \leq b \\ 0 & \text{otherwise} \end{cases}$	$\frac{e^{ibu} - e^{iau}}{(b - a)iu}$	$\bar{x} = \frac{a + b}{2}$	$\frac{(b - a)^2}{12}$
Binomial	$f(r; n, p) = \frac{n!}{r!(n - r)!} p^r q^{n-r}$ $r = 0, 1, 2, \dots, n; \quad 0 \leq p \leq 1; \quad q = 1 - p$	$(q + pe^{iu})^n$	$\bar{r} = np$	npq
Poisson	$f(r; \mu) = \frac{\mu^r e^{-\mu}}{r!}; \quad r = 0, 1, 2, \dots; \quad \mu > 0$	$\exp[\mu(e^{iu} - 1)]$	$\bar{r} = \mu$	μ
Normal (Gaussian)	$f(x; \mu, \sigma^2) = \frac{1}{\sigma\sqrt{2\pi}} \exp(-(x - \mu)^2/2\sigma^2)$ $-\infty < x < \infty; \quad -\infty < \mu < \infty; \quad \sigma > 0$	$\exp(i\mu u - \frac{1}{2}\sigma^2 u^2)$	$\bar{x} = \mu$	σ^2
χ^2	$f(z; n) = \frac{z^{n/2-1} e^{-z/2}}{2^{n/2} \Gamma(n/2)}; \quad z \geq 0$	$(1 - 2iu)^{-n/2}$	$\bar{z} = n$	$2n$
Student's t	$f(t; n) = \frac{1}{\sqrt{n\pi}} \frac{\Gamma[(n + 1)/2]}{\Gamma(n/2)} \left(1 + \frac{t^2}{n}\right)^{-(n+1)/2}$ $-\infty < t < \infty; \quad n$ not required to be integer	—	$\bar{t} = 0$ for $n \geq 2$	$n/(n - 2)$ for $n \geq 3$
Gamma	$f(x; \lambda, k) = \frac{x^{k-1} \lambda^k e^{-\lambda x}}{\Gamma(k)}; \quad 0 < x < \infty;$ k not required to be integer	$(1 - iu/\lambda)^{-k}$	$\bar{x} = k/\lambda$	k/λ^2

The marginal distribution of any x_i is a Gaussian with mean μ_i and variance V_{ii} . V is $n \times n$, symmetric, and positive definite. Therefore for any vector \mathbf{X} , the quadratic form $\mathbf{X}^T V^{-1} \mathbf{X} = C$, where C is any positive number, traces an n -dimensional ellipsoid as \mathbf{X} varies. If $X_i = (x_i - \mu_i)/\sigma_i$, then C is a random variable obeying the $\chi^2(n)$ distribution, discussed in the following section. The probability that \mathbf{X} corresponding to a set of Gaussian random variables \mathbf{x} lies outside the ellipsoid characterized by a given value of $C (= \chi^2)$ is given by Eq. (16.22) and may be read from Fig. 16.1. For example, the “ s -standard-deviation ellipsoid” occurs at $C = s^2$. For the two-variable case ($n = 2$), the point \mathbf{X} lies outside the one-standard-deviation ellipsoid with 61% probability. (This assumes that μ_i and σ_i are correct.) For $X_i = x_i/\sigma_i$, the ellipsoids of constant χ^2 have the same size and orientation but are centered at $\boldsymbol{\mu}$. The use of these ellipsoids as indicators of probable error is described in Sec. 17.5.1.

It is a characteristic of the multivariate Gaussian that $\rho_{ij} = 0$ is necessary and sufficient for x_i and x_j to be independent. For a given covariance matrix V , there always exist nonsingular $n \times n$ matrices H such that $HH^T = V$; H is usually upper or lower triangular in the most efficient algorithms. Then $\mathbf{z} = H^{-1}(\mathbf{x} - \boldsymbol{\mu})$ is a vector of n independent Gaussian random variables with zero mean and with a covariance matrix equal to the identity matrix.

16.3.4. χ^2 distribution: If x_1, \dots, x_n are independent Gaussian distributed random variables, the sum $z = \sum_{i=1}^n (x_i - \mu_i)^2/\sigma_i^2$ is distributed as a χ^2 with n degrees of freedom, $\chi^2(n)$. Under a linear transformation to n dependent Gaussian variables x'_i , the χ^2 at each transformed point retains its value; then $z = \mathbf{X}'^T V^{-1} \mathbf{X}'$ as in the previous section. For a set of z_i , each of which is $\chi^2(n_i)$, $\sum z_i$ is a new random variable which is $\chi^2(\sum n_i)$.

Fig. 16.1 shows the confidence level (CL) obtained by integrating the tail of $f(z; n)$:

$$CL(\chi^2) = \int_{\chi^2}^{\infty} f(z; n) dz. \tag{16.22}$$

This is shown for a special case in Fig. 16.2, and is equal to 1.0 minus the cumulative distribution function $F(z = \chi^2; n)$. It is useful




Figure 16.1: The confidence level versus χ^2 for n degrees of freedom, as defined in Eq. (16.22). The curve for a given n gives the probability that a value at least as large as χ^2 will be obtained in an experiment; e.g., for $n = 10$, a value $\chi^2 \gtrsim 18$ will occur in 5% of a large number of experiments. For a fit, the CL is a measure of goodness-of-fit, in that a good fit to a correct model is expected to yield a low χ^2 (see Sec. 17.4.0). For a confidence interval, α measures the probability that the interval does not cover the true value of the quantity being estimated (see Sec. 17.5). The dashed curve for $n = 20$ is calculated using the approximation of Eq. (16.23).

in evaluating the consistency of data with a model (see Sec. 17): The CL is the probability that a random repeat of the given experiment would observe a greater χ^2 , assuming the model is correct. It is also useful for confidence intervals for statistical estimators (see Sec. 17.5), in which case one is interested in the unshaded area of Fig. 16.2.




Figure 16.2: Illustration of the confidence level integral given in Eq. (16.22). This particular example is for $n = 10$, where the area above 15.99 is 0.1.

Since the mean of the χ^2 distribution is equal to n , one expects in a “reasonable” experiment to obtain $\chi^2 \approx n$. While caution is necessary because of the width and skewness of the distribution, the “reduced χ^2 ” $\equiv \chi^2/n$ is a sometimes useful quantity. Figure 16.3 shows χ^2/n for useful CL’s as a function of n .




Figure 16.3: Confidence levels as a function of the “reduced χ^2 ” $\equiv \chi^2/n$ and the number of degrees of freedom n . Curves are labeled by the probability that a measurement will give a value of χ^2/n greater than that given on the y axis; e.g., for $n = 10$, a value $\chi^2/n \gtrsim 1.8$ can be expected 5% of the time.

For large n , the CL is approximately given by [1,7]

$$CL(\chi^2) \approx \frac{1}{\sqrt{2\pi}} \int_y^\infty e^{-x^2/2} dx, \tag{16.23}$$

where $y = \sqrt{2\chi^2} - \sqrt{2n-1}$. This approximation was used to draw the dashed curves in Fig. 16.1 (for $n = 20$) and Fig. 16.3 (for $CL = 5\%$). Since all the functions and their inverses are now readily available in standard mathematical libraries (such as IMSL, used to generate these figures, and personal computer spreadsheets, such as Microsoft® Excel [8]), the approximation (and even figures and tables) are seldom needed.

16.3.5. Student’s t distribution: Suppose that x and x_1, \dots, x_n are independent and Gaussian distributed with mean 0 and variance 1. We then define

$$z = \sum_1^n x_i^2, \text{ and } t = \frac{x}{\sqrt{z/n}}. \tag{16.24}$$

The variable z thus belongs to a $\chi^2(n)$ distribution. Then t is distributed according to a Student’s t distribution with n degrees of freedom, $f(t; n)$, given in Table 16.1.

The Student’s t distribution resembles a Gaussian distribution with wide tails. As $n \rightarrow \infty$, the distribution approaches a Gaussian. If $n = 1$, the distribution is a *Cauchy* or *Breit-Wigner* distribution. The mean is finite only for $n > 1$ and the variance is finite only for $n > 2$, so for $n = 1$ or $n = 2$, t does not obey the central limit theorem.

As an example, consider the *sample mean* $\bar{x} = \sum x_i/n$ and the *sample variance* $s^2 = \sum (x_i - \bar{x})^2/(n-1)$ for normally distributed random variables x_i with unknown mean μ and variance σ^2 . The sample mean has a Gaussian distribution with a variance σ^2/n , so the variable $(\bar{x} - \mu)/\sqrt{\sigma^2/n}$ is normal with mean 0 and variance 1. Similarly, $(n-1)s^2/\sigma^2$ is independent of this and is χ^2 distributed with $n-1$ degrees of freedom. The ratio

$$t = \frac{(\bar{x} - \mu)/\sqrt{\sigma^2/n}}{\sqrt{(n-1)s^2/\sigma^2}/\sqrt{n-1}} = \frac{\bar{x} - \mu}{\sqrt{s^2/n}} \tag{16.25}$$

is distributed as $f(t; n-1)$. The unknown true variance σ^2 cancels, and t can be used to test the probability that the true mean is some particular value μ .

In Table 16.1, n in $f(t; n)$ is not required to be an integer. A Student’s t distribution with nonintegral $n > 0$ is useful in certain applications.

16.3.6. Gamma distribution: For a process that generates events as a function of x (e.g., space or time) according to a Poisson distribution, the distance in x from an arbitrary starting point (which may be some particular event) to the k^{th} event belongs to a *gamma* distribution, $f(x; \lambda, k)$. The Poisson parameter μ is λ per unit x . The special case $k = 1$ (i.e., $f(x; \lambda, 1) = \lambda e^{-\lambda x}$) is called the *exponential* distribution. A sum of k' exponential random variables x_i is distributed as $f(\sum x_i; \lambda, k')$.

The parameter k is not required to be an integer. For $\lambda = 1/2$ and $k = n/2$, the gamma distribution reduces to the $\chi^2(n)$ distribution.

References:

1. H. Cramér, *Mathematical Methods of Statistics*, Princeton Univ. Press, New Jersey (1958).
2. F.T. Solmitz, *Ann. Rev. Nucl. Sci.* **14**, 375 (1964).
3. W.T. Eadie, D. Drijard, F.E. James, M. Roos, and B. Sadoulet, *Statistical Methods in Experimental Physics* (North Holland, Amsterdam and London, 1971).
4. L. Lyons, *Statistics for Nuclear and Particle Physicists* (Cambridge University Press, New York, 1986).
5. B.R. Roe, *Probability and Statistics in Experimental Physics*, (Springer-Verlag, New York, 208 pp., 1992).
6. M. Abramowitz and I. Stegun, eds., *Handbook of Mathematical Functions* (Dover, New York, 1972).
7. R.A. Fischer, *Statistical Methods for Research Workers*, 8th edition, Edinburgh and London (1941).
8. Microsoft® is a registered trademark of Microsoft corporation.

17. STATISTICS

Revised June 1994 with the help of R. Cousins, F. James, G. Lynch, B. P. Roe, and M. Roos

17.1. General [1–6]

A probability density function with known parameters enables us to predict the frequency with which a random variable will take on a particular value (if discrete) or lie in a given range (if continuous). In *parametric* statistics we have the opposite problem of estimating the parameters of the p.d.f. from a set of actual observations.

We refer to the true p.d.f. as the *population*; the data form a *sample* from this population. A *statistic* is any function of the data, plus known constants, which does not depend upon any of the unknown parameters. A statistic is a random variable if the data have random errors. An *estimator* is any statistic whose value is intended as a meaningful guess for the value of an unknown parameter; we denote estimators with hats, e.g., $\hat{\alpha}$.

Often it is possible to construct more than one reasonable estimator. Let α represent the true value of a parameter to be estimated; α is a vector α if there is more than one parameter. Then if $\hat{\alpha}$ is an estimator for α , desirable properties for $\hat{\alpha}$ are: (a) *Unbiased*; bias $b = E(\hat{\alpha}) - \alpha$, where the expectation value is taken over a hypothetical set of similar experiments in which $\hat{\alpha}$ is constructed the same way. The bias may be due to statistical properties of the estimator or to *systematic* errors in the experiment. If we can estimate the average bias b we usually subtract it from $\hat{\alpha}$ to obtain a new $\hat{\alpha}' \equiv \hat{\alpha} - b$. However, b may depend upon α or other unknowns, in which case we usually try to choose an estimator which minimizes its average size. (b) *Minimum variance*; the minimum possible value of $\text{Var}(\hat{\alpha})$ is given by the Rao-Cramér-Frechet bound:

$$\text{Var}_{\min} = [1 + \partial b / \partial \alpha]^2 / I(\alpha); \quad (17.1)$$

$$I(\alpha) = E \left\{ \left[\frac{\partial}{\partial \alpha} \sum_{i=1}^n \ln f(x_i; \alpha) \right]^2 \right\}.$$

(Compare with Eq. (17.6) below.) The sum is over all data and b is the bias, if any; the x_i are assumed independent and distributed as $f(x_i; \alpha)$, and the allowed range of x must not depend upon α . The ratio $\epsilon = \text{Var}_{\min} / \text{Var}(\hat{\alpha})$ is the *efficiency*. An *efficient* estimator (with $\epsilon = 1$) exists only for certain cases. The square root of the variance expresses the expected spread of $\hat{\alpha}$ about its average value, as would be observed in a large number of repeats of the same measurement. (c) *Minimum mean-squared error* (mse); $\text{mse} = E[(\hat{\alpha} - \alpha)^2] = V(\hat{\alpha}) + b^2$. The mse combines the error due to any bias quadratically with the variance, which expresses only the spread about $E(\hat{\alpha})$, as distinct from α , the true value. (d) *Robust*; a robust estimator is not sensitive to errors in our assumptions, e.g., to departures from the assumed p.d.f. due to such factors as noise.

These criteria (and others) allow us to evaluate any procedure for obtaining $\hat{\alpha}$. In many cases these criteria conflict. The bias, variance, and mse may depend on the unknown α . In this case the optimum prescription for $\hat{\alpha}$ may depend on the range in which we assume α to lie.

Following are techniques in common use for obtaining estimators and their standard errors $\sigma(\hat{\alpha}) = \sqrt{\text{Var}(\hat{\alpha})}$. When the conditions of the central limit theorem are satisfied, the interval $\hat{\alpha} \pm \sigma(\hat{\alpha})$ forms a 68.3% *confidence interval*. This is a random interval in that its endpoints depend upon the randomly sampled data; its meaning here will be taken to be that in 68.3% of all similar experiments the interval will include the true value α . One should be aware that in most practical cases the central limit theorem is only approximately satisfied and accordingly confidence intervals which depend on that are only approximate. Confidence intervals are discussed in Section 17.5 below.

17.2. Data with a common mean

Suppose we have a set of N independent measurements y_i assumed to be unbiased measurements of the same unknown quantity μ with a common, but unknown, variance σ^2 resulting from measurement error. Then

$$\hat{\mu} = \frac{1}{N} \sum_{i=1}^N y_i \quad (17.2)$$

$$\hat{\sigma}^2 = \frac{1}{N-1} \sum_{i=1}^N (y_i - \hat{\mu})^2 = \frac{N}{N-1} (E(y^2) - \hat{\mu}^2) \quad (17.3)$$

are unbiased estimators of μ and σ^2 . The variance of $\hat{\mu}$ is σ^2/N . If the common p.d.f. of the y_i is Gaussian, these statistics are independent. Then, for large N , the variance of $\hat{\sigma}^2$ is $2\sigma^4/N$. If the y_i are Gaussian or N is large enough that the central limit theorem applies, then $\hat{\mu}$ is an efficient estimator for μ . Otherwise $\hat{\mu}$ is sometimes subject to large fluctuations, e.g., if the p.d.f. for y_i has long tails. In this case the median of the y_i may be a more robust estimator for μ , provided the median and mean are expected to lie at the same point in the p.d.f. for y . For Gaussian y , the median has asymptotic (large- N) efficiency $2/\pi \approx 0.64$. Student's t -distribution provides an example in which there are large tails. In this case, for large N the efficiency of the sample median relative to the sample mean is $(\infty, \infty, 1.62, 1.12, 0.96, 0.80, 0.64)$ for $(1, 2, 3, 4, 5, 8, \infty)$ degrees of freedom.

If σ^2 is known, $\hat{\mu}$ as given in Eq. (17.2) is still the best estimator for μ ; if μ is known, substitute it for $\hat{\mu}$ in Eq. (17.3) and replace $N-1$ by N , to obtain a somewhat better estimator $\hat{\sigma}^2$.

If the y_i have different, known, variances σ_i^2 , then

$$\hat{\mu} = \frac{1}{w} \sum_{i=1}^N w_i y_i, \quad (17.4)$$

is an unbiased estimator for μ with smaller variance than Eq. (17.2), where $w_i = 1/\sigma_i^2$ and $w = \sum w_i$. The variance of $\hat{\mu}$ is $1/w$.

17.3. The method of maximum likelihood

17.3.1. General:

"From a theoretical point of view, the most important general method of estimation so far known is the *method of maximum likelihood*" [1]. We suppose that a set of independently measured quantities x_i came from a p.d.f. $f(x; \alpha)$, where α is an unknown set of parameters. The method of maximum likelihood consist of finding the set of values, $\hat{\alpha}$, which maximizes the joint probability density for all the data, given by

$$\mathcal{L}(\alpha) = \prod_i f(x_i; \alpha), \quad (17.5)$$

where \mathcal{L} is called the likelihood. It is usually easier to work with $\ln \mathcal{L}$, and since both are maximized for the same set of α , it is sufficient to solve the *likelihood equation*

$$\frac{\partial \ln \mathcal{L}}{\partial \alpha_n} = 0. \quad (17.6)$$

The solution is called the *maximum likelihood estimate* of α . The importance of the approach is shown by the following proposition, proved in Ref. 1:

If an efficient estimate $\hat{\alpha}$ of α exists, the likelihood equation will have a unique solution equal to $\hat{\alpha}$.

In evaluating \mathcal{L} , it is important that any normalization factors in the f 's which involve α be included. However, we will only be interested in the maximum of \mathcal{L} and in ratios of \mathcal{L} at different α 's; hence any multiplicative factors which do not involve the parameters

we want to estimate may be dropped; this includes factors which depend on the data but not on α .

If the solution to Eq. (17.6) is at a maximum, $\partial \ln \mathcal{L} / \partial \alpha_n$ will have negative slope in its vicinity. In many practical problems, one often uses nonlinear algorithms for finding the maximum, and must be alert to various possibilities for error: (a) Eq. (17.6) may yield a minimum, therefore one must check the second derivative; (b) there may be more than one maximum—one must try to find the global maximum; (c) the global maximum may lie at a boundary of the physical region, in which case Eq. (17.6) will not find it.

If an unbiased, efficient estimator exists, this method will find it. If $\partial \ln \mathcal{L} / \partial \alpha_n$ is linear in the vicinity of the root, an efficient estimator is guaranteed; other efficient cases are discussed in the literature. For large data samples, the central limit theorem will usually assure this condition in some significant neighborhood of zero; hence the estimator is usually efficient in that case, provided certain conditions are met (e.g., that the solution does not lie on a boundary). In this case, in the neighborhood of the maximum $\ln \mathcal{L}$ is a downward-curving paraboloid and \mathcal{L} is proportional to a multivariate Gaussian.

The results of two or more experiments may be combined by forming the product of the \mathcal{L} 's, or the sum of the $\ln \mathcal{L}$'s.

Under a one-to-one change of parameters from α to $\beta = \beta(\alpha)$, the maximum likelihood estimate is simply $\beta(\hat{\alpha})$, given the solution $\hat{\alpha}$ for α . That is, the maximum likelihood solution for β is found by simple substitution of $\hat{\alpha}$ into the transformation equation. It is possible that the new solution $\hat{\beta}$ will be a biased solution for the true value of β even if $\hat{\alpha}$ is not biased, and vice-versa. In the asymptotic limit (of large amounts of data) both $\hat{\alpha}$ and $\hat{\beta}$ will (usually) converge to unbiased solutions, but at different rates.

Except in special cases like the least-squares method, the value of the likelihood function at the solution does not necessarily tell us whether the final fit was a sensible description of the data or not. In special cases such as the one discussed in Sec. 17.3.3, one can define a quantity which approaches the χ^2 -distribution in the limit of a large number of counts in the experiment, but in general some other strategy must be used. For example, data generated by Monte Carlo simulations of the experiment can be analyzed by the same method. If the experimental likelihood is lower than that of some agreed-upon fraction of these results, one should question the appropriateness of the p.d.f. At the same time one can check for bias in the solution.

17.3.2. Error estimates:

The covariance matrix V may be estimated from

$$V_{nm} = \left(E \left[- \frac{\partial^2 \ln \mathcal{L}}{\partial \alpha_n \partial \alpha_m} \Big|_{\hat{\alpha}} \right] \right)^{-1} \tag{17.7}$$

If $\partial \ln \mathcal{L} / \partial \alpha_n$ is linear, the “expectation” operation in Eq. (17.7) has no effect because the second derivative of $\ln \mathcal{L}$ is constant. Otherwise, it may be approximated by taking the average of the quantity in square brackets over a range of α_n and α_m near the solution. For complex cases it may be more practical to evaluate s -standard-deviation errors from the contour

$$\ln \mathcal{L}(\alpha) = \ln \mathcal{L}_{\max} - s^2 / 2, \tag{17.8}$$

where $\ln \mathcal{L}_{\max}$ is the value of $\ln \mathcal{L}$ at the solution point (compare with Eq. (17.27), below). The extreme limits of this contour parallel to the α_n axis give an approximate s -standard-deviation confidence interval in α_n . These intervals may not be symmetric and they may even consist of two or more disjoint intervals. This procedure gives one-standard-deviation errors in α_n equal to $\sqrt{V_{nn}}$ (not summed) of Eq. (17.7) if the estimator is efficient. If it is not efficient, the level of confidence implied by the value of s is only approximate.

17.3.3. Application to Poisson-distributed data:

In the case of Poisson-distributed data in a counting experiment, the unbinned maximum likelihood method (where the index i in Eq. (17.5) labels events) is preferred if the total the number of events is very small. If there are enough events to justify binning them in a histogram, then one may alternatively maximize the likelihood function for the contents of the bins (so i labels bins). This is equivalent to minimizing [7]

$$\chi^2 = \sum_i \left[2(N_i^{\text{th}} - N_i^{\text{obs}}) + 2N_i^{\text{obs}} \ln(N_i^{\text{obs}}/N_i^{\text{th}}) \right]. \tag{17.9}$$

where N_i^{obs} and N_i^{th} are the observed and theoretical (from f) contents of the i th bin. In bins where $N_i^{\text{obs}} = 0$, the second term is zero. This function asymptotically behaves like a classical χ^2 for purposes of point estimation, interval estimation, and goodness-of-fit. It also guarantees that the area under the fitted function f is equal to the sum of the histogram contents (as long as the overall normalization of f is effectively left unconstrained during the fit), which is not the case for χ^2 statistics based on a least-squares procedure with traditional weights.

17.4. Method of least squares

The *method of least squares* can be derived from the maximum likelihood theorem. We suppose a set of N measurements at points x_i . The i th measurement y_i is assumed to be chosen from a Gaussian distribution with mean $F(x_i; \alpha)$ and variance σ_i^2 . Then

$$\chi^2 = -2 \ln \mathcal{L} + \text{constant} = \sum_1^N \frac{[y_i - F(x_i; \alpha)]^2}{\sigma_i^2}. \tag{17.10}$$

Finding the set of parameters α which maximizes \mathcal{L} is equivalent to finding the set which minimizes χ^2 .

In many practical cases one further restricts the problem to the situation in which $F(x_i; \alpha)$ is a linear function of the α_n 's,

$$F(x_i; \alpha) = \sum_n \alpha_n f_n(x), \tag{17.11}$$

where the f_n are k linearly independent functions (e.g., $1, x, x^2, \dots$, or Legendre polynomials) which are single-valued over the allowed range of x . We require $k \leq N$, and at least k of the x_i must be distinct. We wish to estimate the linear coefficients α_n . Later we will discuss the nonlinear case.

If the point errors $\epsilon_i = y_i - F(x_i; \alpha)$ are Gaussian, then the minimum χ^2 will be distributed as a χ^2 random variable with $n = N - k$ degrees of freedom. We can then evaluate the goodness-of-fit (confidence level) from Figs. 16.1 or 16.3, as per the earlier discussion. The confidence level expresses the probability that a *worse* fit would be obtained in a large number of similar experiments under the assumptions that: (a) the model $y = \sum \alpha_n f_n$ is correct and (b) the errors ϵ_i are Gaussian and unbiased with variance σ_i^2 . If this probability is larger than an agreed-upon value (0.001, 0.01, or 0.05 are common choices), the data are *consistent* with the assumptions; otherwise we may want to find improved assumptions. As for the converse, most people do not regard a model as being truly *inconsistent* unless the probability is as low as that corresponding to four or five standard deviations for a Gaussian (6×10^{-3} or 6×10^{-5} ; see Sec. 17.5.1). If the ϵ_i are not Gaussian, the method of least squares still gives an answer, but the goodness-of-fit test would have to be done using the correct distribution of the random variable which is still called “ χ^2 .”

Finding the minimum of χ^2 in the linear case is straightforward:

$$\begin{aligned} -\frac{1}{2} \frac{\partial \chi^2}{\partial \alpha_m} &= \sum_i f_m(x_i) \left(\frac{y_i - \sum_n \alpha_n f_n(x_i)}{\sigma_i^2} \right) \\ &= \sum_i \frac{y_i f_m(x_i)}{\sigma_i^2} - \sum_n \alpha_n \sum_i \frac{f_n(x_i) f_m(x_i)}{\sigma_i^2}. \end{aligned} \tag{17.12}$$

With the definitions

$$g_m = \sum_i y_i f_m(x_i) / \sigma_i^2 \tag{17.13}$$

and

$$(V_{\hat{\alpha}}^{-1})_{mn} = \sum_i f_n(x_i) f_m(x_i) / \sigma_i^2, \tag{17.14}$$

the k -element column vector of solutions $\hat{\alpha}$, for which $\partial\chi^2/\partial\alpha_m = 0$ for all m , is given by

$$\hat{\alpha} = V_{\hat{\alpha}} g. \tag{17.15}$$

Non-independent y_i 's

More generally, the measured y_i 's are not independent. Then the set of σ_i^2 's must be replaced by the $N \times N$ covariance matrix V_y . Then, if H is the $N \times k$ matrix with element $H_{in} = f_n(x_i)$, the solution $\hat{\alpha}$ is given by the solution to the normal equation

$$(H^T V_y^{-1} H) \hat{\alpha} = H^T V_y^{-1} y, \tag{17.16a}$$

or, formally,

$$\hat{\alpha} = (H^T V_y^{-1} H)^{-1} H^T V_y^{-1} y \equiv D y, \tag{17.16b}$$

where y is the N -element vector of measured y_i 's. The normal equations may be solved by numerical methods much more computationally efficient than brute application of Eq. (17.16b). In particular, $H^T V_y^{-1} H$ is sometimes singular or nearly singular. In such cases there is at least one f_n which may be expressed as a linear combination of others (or nearly so) when evaluated at the data points. The best procedure is usually to drop such functions from the expansion (or set $\hat{\alpha}_n = 0$). See Press [8], Maindonald [9], or Basilevsky [10] for discussions.

In terms of the $k \times N$ matrix D , the standard covariance matrix for the $\hat{\alpha}$ is estimated by

$$V_{\hat{\alpha}} = D V_y D^T. \tag{17.17}$$

If the measured y_i 's are independent, V_y is diagonal with ii^{th} element σ_i^2 and $V_{\hat{\alpha}}$ is obtained from Eq. (17.14) above.

The expected covariance [see Eq. (16.9)] of $\hat{\alpha}_n$ and $\hat{\alpha}_m$ is estimated by

$$E[(\alpha_n - \hat{\alpha}_n)(\alpha_m - \hat{\alpha}_m)] = (V_{\hat{\alpha}})_{nm}. \tag{17.18}$$

Even when the y_i 's are independent (diagonal V_y), $\hat{\alpha}_n$ and $\hat{\alpha}_m$ may not be (nondiagonal $V_{\hat{\alpha}}$). For the model function $y = \sum \alpha_n f_n(x)$, the estimated variance of an interpolated or extrapolated value of y at a point x is

$$E[(y - \hat{y})^2] = \sigma^2(y) = \sum_{n,m} (V_{\hat{\alpha}})_{nm} f_n(x) f_m(x). \tag{17.19}$$

Example: straight-line fit

For the case of a straight-line fit, $y(x) = \alpha_1 + \alpha_2 x$, one obtains, for independent measurements y_i , the following estimates of α_1 and α_2 ,

$$\hat{\alpha}_1 = (g_1 S_{22} - g_2 S_{12}) / D, \tag{17.20}$$

$$\hat{\alpha}_2 = (g_2 S_{11} - g_1 S_{12}) / D, \tag{17.21}$$

where

$$(S_{11}, S_{12}, S_{22}) = \sum (1, x_i, x_i^2) / \sigma_i^2, \tag{17.22a}$$

$$(g_1, g_2) = \sum (1, x_i) y_i / \sigma_i^2. \tag{17.22b}$$

respectively, and

$$D = S_{11} S_{22} - S_{12}^2. \tag{17.23}$$

The covariance matrix of the fitted parameters is:

$$\begin{pmatrix} V_{11} & V_{12} \\ V_{12} & V_{22} \end{pmatrix} = \frac{1}{D} \begin{pmatrix} S_{22} & -S_{12} \\ -S_{12} & S_{11} \end{pmatrix}. \tag{17.24}$$

The estimated variance of an interpolated or extrapolated value of y at point x is:

$$(\hat{y} - y_{\text{true}})_{\text{est}}^2 = \frac{1}{S_1} + \frac{S_1}{D} \left(x - \frac{S_x}{S_1}\right)^2. \tag{17.25}$$

17.4.1. General comments:

If y is not linear in the fitting parameters α_n , or if the errors σ_i depend upon y and therefore on α_n , the solution vector may have to be found by iteration of Eqs. (17.13)–(17.15) or Eq. (17.16b). The same results may be obtained by numerical techniques from the sum of squares, χ^2 , directly, if we have a reasonable first guess α_0 for the solution vector:

$$\hat{\alpha} = \alpha_0 - \left(\frac{\partial^2 \chi^2}{\partial \alpha^2}\right)_{\alpha_0}^{-1} \cdot \frac{\partial \chi^2}{\partial \alpha} \Big|_{\alpha_0} \tag{17.26a}$$

and

$$V_{\hat{\alpha}} = 2 \left(\frac{\partial^2 \chi^2}{\partial \alpha^2}\right)_{\hat{\alpha}}^{-1}, \tag{17.26b}$$

where $\partial\chi^2/\partial\alpha$ is a k -element vector whose n^{th} element is $\partial\chi^2/\partial\alpha_n$, $\partial^2\chi^2/\partial\alpha^2$ is a $k \times k$ matrix with mn^{th} element $\partial^2\chi^2/(\partial\alpha_m \cdot \partial\alpha_n)$, and all derivatives are to be evaluated at the points indicated. If “ χ^2 ” is a true χ^2 , the second-derivative matrix is independent of α ; therefore the shape of the χ^2 as a function of α is a paraboloid and Eq. (17.26a) will give the solution immediately. Otherwise one may need to iterate Eq. (17.26a) to arrive at a solution (Newton-Raphson method). The CERN program MINUIT [11] offers several iteration schemes for solving such problems.

Note that in Eq. (17.16b), one needs only a matrix proportional to V_y to find $\hat{\alpha}$. Hence, for example, if the variances σ_i^2 of the errors are unknown but assumed equal and independent, and $E(\epsilon_i) = 0$, one can still solve for $\hat{\alpha}$. One cannot, however, solve for $V_{\hat{\alpha}}$ or evaluate goodness-of-fit. These can be estimated from the residuals, $r_i = \hat{y}(x_i) - y_i$, where $\hat{y}(x_i)$ is the fitted curve at x_i , because study of the r_i enables one to estimate V_y . In addition, the residuals can be used to look for evidence of bias such as trends in the data not incorporated in the model [4].

The errors on the solution $\hat{\alpha}$ are independent of the value of χ^2 at minimum—they depend only upon the shape about the minimum. Eq. (17.26b) implies that s -standard-deviation limits on the elements of $\hat{\alpha}$ are given by the set of α' such that

$$\chi^2(\alpha') = \chi_{\text{min}}^2 + s^2; \tag{17.27}$$

(This is a special case of Eq. (17.8).) This equation, which defines a contour in α -space, is often convenient for estimating errors in applications of least-squares techniques to nonlinear cases, where the second derivative [Eq. (17.26b)] may be a rapidly varying function of α . If the problem is highly nonlinear, all such contours are, at best, only approximations to desired exact confidence regions which would have some given probability of covering the true value of α . It may be that Eq. (17.27) will define a set of disjoint regions. In addition, iteration of Eq. (17.26a) may require sophisticated techniques [8,11] to reach convergence in a practical amount of computation. For example, in cases involving many variables in α , especially if the correlations are large, simplex or other techniques which do not involve explicit calculation of derivatives are often to be preferred. Such techniques are designed to find their way through complicated nonlinear problems without diverging to infinite α (unless the minimum is actually at infinity).

The method of least squares is sometimes used in cases where the distribution is not Gaussian or not known to be Gaussian. In such cases it can still be used, but it is then not a special case of the maximum-likelihood method, and the theorems having to do with that approach no longer apply. However, if (a) the distribution of $y_i - \sum \alpha_k f_k(x_i)$ has an expectation value of zero (unbiased) and (b) has a finite, known, fixed variance σ_i^2 (does not depend on α), then estimates of α obtained by minimizing χ^2 will be unbiased and have the smallest possible variance of all linear unbiased estimates (Gauss-Markov theorem). This statement is more general than the least-squares method as a special case of the maximum likelihood method in that the distributions do not have to be Gaussian, but more restrictive in that it applies only when the fitting function is linear in the α_k 's.

For example, in counting experiments one often groups the data in bins in order to associate a Poisson error with each bin. In this case y_i is the bin height and the error depends on the expectation value of the theory in each bin, as estimated by the best fit of the model. Since the variances are functions of the fitting parameters, neither the conditions of the Gauss-Markov theorem nor the assumption of Gaussian distributions (with or without fixed variances) in the context of the maximum likelihood approach are valid without a large-number approximation, and so an unbiased or efficient least-squares fit is not guaranteed. In such cases it seems more sensible to use one of the approaches discussed in Sec. 17.3.3.

17.5. Errors and confidence intervals

We measure a mass, lifetime, or other physical quantity under the assumption that a “true answer” α exists. The conditions of the measurement introduce a random element, and our measurement (or combination of measurements) $\hat{\alpha}_{\text{exp}}$ samples a distribution with p.d.f. $f(\hat{\alpha}; \alpha)$. The unknown constant α appears as a parameter. We suppose that for every value of α we can find two values $\gamma_1(\alpha, \epsilon)$ and $\gamma_2(\alpha, \epsilon)$ such that repeated experiments would produce results in the interval $\gamma_1 < \hat{\alpha} < \gamma_2$ a fraction $1 - \epsilon$ of the time, where

$$1 - \epsilon = \int_{\gamma_1}^{\gamma_2} f(\hat{\alpha}; \alpha) d\hat{\alpha}. \tag{17.28}$$

This situation is shown in Fig. 17.1 (ignore the “unphysical region” part of the graph for now), where the region between the curves $\gamma_1(\alpha, \epsilon)$ and $\gamma_2(\alpha, \epsilon)$ is indicated by the domain $D(\epsilon)$. It can be argued that since the point $(\alpha_{\text{actual}}, \hat{\alpha}_{\text{exp}})$ belongs to D , then our statement that repeated experiments would produce values of $\hat{\alpha}$ in the interval $\gamma_1 < \hat{\alpha} < \gamma_2$ is equivalent to the statement that the confidence interval $c_1 < \alpha < c_2$ includes α_{actual} with probability $1 - \epsilon$ [1,6]. (We will call ϵ the *confidence coefficient*.) In this “confidence interval” or frequentist approach, α is a parameter, not a statistical variable. Instead, c_1 and c_2 vary from experiment to experiment and are statistical variables. It is very different to say that a lifetime τ is to be found in the interval $\tau_0 \pm \sigma_\tau$ with 68% probability than to say that the interval $\tau_0 \pm \sigma_\tau$ (which can vary from experiment to experiment) includes the actual, fixed, value of the lifetime with 68% probability.

The actual choice of γ_1 and γ_2 , such that $\int_{\gamma_1}^{\gamma_2} f(\hat{\alpha}; \alpha) d\hat{\alpha} = 1 - \epsilon$, can be made in an infinite number of ways, but in practical situations there are usually additional criteria. For a Gaussian distribution, for example, choosing the limits symmetric about the mean minimizes the length of the interval. The area of the excluded tail on either side is then $\epsilon/2$. For a Poisson distribution negative values cannot occur, so $\gamma(\hat{\alpha}, \alpha)$ (with $\hat{\alpha}$ an integer and α the Poisson mean) might be taken as the curve below which ϵ of the area under the distribution lies. (In this case the curve really consists of discrete points, since $\hat{\alpha}$ can have only discrete values.) For $\epsilon = 0.05$ the curve starts at $(\alpha, \hat{\alpha}) = (3.0, 0)$. If in a given experiment no decays to a certain final state are seen, we might then conclude that $\alpha < 3.0$ excludes the actual value of α with 95% probability. This statement can be converted to a similar statement about the branching fraction.

In Sec. 16 we discussed such confidence limits for a χ^2 distribution (where ϵ was called CL). Here we discuss confidence intervals for the Gaussian and Student’s t -distribution, and confidence limits for the Poisson case. We then discuss the much more contentious situation in which the horizontal line at ordinate $\hat{\alpha}$ in Fig. 17.1 enters $D(\epsilon)$ at a boundary for unphysical values of α , so that at least c_1 is undefined—for example, if we find $\hat{m}^2 = -30 \pm 50 \text{ eV}^2$.

Extensive tables and graphs were once used to find confidence intervals and limits, but by now their main function is to confirm that software is working. FORTRAN mathematical libraries (IMSL, NAG, CERLIB) are readily available, and a wide range of distributions are available in personal computer spreadsheet applications such as Microsoft® Excel [12]. Its built-in functions CHIDIST, NORMDIST, and TDIST (Student’s t -distribution), along with “Solver,” were used to produce or check the numbers given in this section.




Figure 17.1: Confidence intervals for a single unknown parameter α . One might think of the p.d.f. $f(\hat{\alpha}; \alpha)$ as being plotted out of the paper as a function of $\hat{\alpha}$ along each vertical line of constant α . The domain $D(\epsilon)$ contains a fraction $1 - \epsilon$ of the area under each of these functions.

17.5.1. Gaussian errors:

If the data are such that the distribution of the estimator(s) satisfies the central limit theorem discussed in Sec. 16.3.3, the Gaussian distribution is the basis of the error analysis. If there is more than one parameter being estimated, the multivariate Gaussian is used. For the univariate case with known σ ,

$$1 - \epsilon = \int_{\hat{\mu} - \delta}^{\hat{\mu} + \delta} f(x; \hat{\mu}, \sigma^2) dx = \text{erf} \left(\frac{x - \hat{\mu}}{\sqrt{2} \sigma} \right) \tag{17.29}$$




Figure 17.2: Illustration of a symmetric 90% confidence interval (unshaded) for a measurement of a single quantity with Gaussian errors. Integrated probabilities, defined by ϵ , are as shown.

is the probability that the true value of μ will fall within $\pm \delta$ ($\delta > 0$) of the measured $\hat{\mu}$. This interval will cover μ in a fraction $1 - \epsilon$ of all similar measurements. Fig. 17.2 shows a $\delta = 1.64\sigma$ confidence interval unshaded. The choice $\delta = \sqrt{\text{Var}(\hat{\mu})} \equiv \sigma$ gives an interval called the *standard error* which has $1 - \epsilon = 68.27\%$ if σ is known. Confidence coefficients ϵ for other frequently used choices of δ are given in Table 17.1.

For other δ , find ϵ as the ordinate of Fig. 16.1 on the $n = 1$ curve at $\chi^2 = (\delta/\sigma)^2$. We can set a one-sided (upper or lower) limit by excluding above $\hat{\mu} + \delta$ (or below $\hat{\mu} - \delta$); ϵ 's for such limits are 1/2 the values in the table above.

We have increased confidence that the interval covers the true value as $1 - \epsilon$ increases, or χ^2 increases. We must be careful to distinguish this case from the other major use of Fig. 16.1, evaluation of goodness-of-fit (Sec. 17.4.0). In that case we have increased confidence

Table 17.1: Area of the tails ϵ outside $\pm\delta$ from the mean of a Gaussian distribution.

ϵ (%)	δ	ϵ (%)	δ
31.73	1σ	20	1.28σ
4.55	2σ	10	1.64σ
0.27	3σ	5	1.96σ
6.3×10^{-3}	4σ	1	2.58σ
5.7×10^{-5}	5σ	0.1	3.29σ
2.0×10^{-7}	6σ	0.01	3.89σ

in the fit as χ^2 decreases. In an attempt to reduce possible confusion in this discussion, we will use the ϵ notation (which corresponds to notation used in hypothesis testing [4]) when discussing confidence intervals and CL notation when discussing goodness-of-fit. Elsewhere in this *Review*, where the confusion between fit confidence level and interval (usually an upper or lower limit) confidence level does not arise, we follow the common practice of using “CL” to refer to the confidence level of the interval. This CL is understood to represent $1 - \epsilon$.

If the variance σ^2 of the estimator is not known, but must be estimated from the data, then we need to incorporate the error in $\hat{\sigma}$ into our confidence interval using Student’s t distribution. If we have N data points with which we estimate k parameters, the Gaussian approximation is adequate for $N - k \gg 1$. Otherwise replace δ by a factor $T\hat{\sigma}$, T being defined by

$$1 - \epsilon = \int_{-T}^T f(t; N - k) dt, \tag{17.30}$$

where f for the Student’s t -distribution is defined in Table 16.1. T is tabulated in Ref. 13 and in Table 17.2.

Table 17.2: t limits containing $1 - \epsilon$ of the area of Student’s t -distribution $f(t; N - k)$.

$N - k$	ϵ (%)					
	31.67	10.00	5.00	4.55	1.00	0.27
1	1.84	6.31	12.71	13.97	63.66	235.78
2	1.32	2.92	4.30	4.53	9.92	19.21
3	1.20	2.35	3.18	3.31	5.84	9.22
4	1.14	2.13	2.78	2.87	4.60	6.62
5	1.11	2.01	2.57	2.65	4.03	5.51
10	1.05	1.81	2.23	2.28	3.17	3.96
20	1.03	1.72	2.09	2.13	2.85	3.42
∞	1.00	1.64	1.96	2.00	2.58	3.00

For multivariate α we must consider pairwise correlations. Assuming a multivariate Gaussian, Eq. (16.20), and subsequent discussion the standard error ellipse for the pair $(\hat{\alpha}_m, \hat{\alpha}_n)$ may be drawn as in Fig. 17.3.




Figure 17.3: Standard error ellipse for the estimators $\hat{\alpha}_m$ and $\hat{\alpha}_n$. In this case the correlation is negative.

The minimum χ^2 or maximum likelihood solution is at $(\hat{\alpha}_m, \hat{\alpha}_n)$. The standard errors σ_m and σ_n are defined as shown, where the ellipse is at a constant value of $\chi^2 = \chi^2_{\min} + 1$ or

$\ln \mathcal{L} = \ln \mathcal{L}_{\max} - 1/2$. The angle of the major axis of the ellipse is given by

$$\tan 2\phi = \frac{2\rho_{mn} \sigma_m \sigma_n}{\sigma_m^2 - \sigma_n^2}. \tag{17.31}$$

For non-Gaussian or nonlinear cases, one may construct an analogous contour from the same χ^2 or $\ln \mathcal{L}$ relations. Any other parameters $\hat{\alpha}_\ell, \ell \neq m, n$ must be allowed freely to find their optimum values for every trial point.

For any unbiased procedure (e.g., least squares or maximum likelihood) being used to estimate k parameters $\alpha_i, i = 1, \dots, k$, the probability $1 - \epsilon$ that the true values of all k lie within the s -standard deviation ellipsoid may be found from Fig. 16.1. Read the ordinate as ϵ ; the correct value of ϵ occurs on the $n = k$ curve at $\chi^2 = s^2$. For example, for $k = 2$, the probability that the true values of α_1 and α_2 simultaneously lie within the one-standard-deviation error ellipse ($s = 1$), centered on $\hat{\alpha}_1$ and $\hat{\alpha}_2$, is 39%. This probability only assumes Gaussian errors, unbiased estimators, and that the model describing the data in terms of the α_i is correct.

17.5.2. Poisson processes—upper limits:

Because the outcome of a Poisson process is an integral number of events, n_0 , it is usually not possible to set confidence intervals for the true Poisson parameter μ at a certain exact ϵ . For large n_0 an approximate interval can be set using the Gaussian approximation, in our section on Probability, Sec. 16.3.2, and the techniques of Sec. 17.5.1.

For small n_0 we can define an upper limit N for μ as being that value of μ such that it would be at least $1 - \epsilon$ (e.g., 90% or 95%) probable that a random observation of n would then lie above the observed n_0 . Thus

$$1 - \epsilon = \sum_{n=n_0+1}^{\infty} f(n; N); \quad \epsilon = \sum_{n=0}^{n_0} f(n; N). \tag{17.32}$$




Figure 17.4: Illustration of Eq. (17.32) Poisson probabilities for an assumed mean of N . With an observed count $n_0 = 2$, $N = 5.3$ as shown gives summed probability $1 - \epsilon = 90\%$. The dotted summed probability curve (scale on right) has been displaced by -0.5 for clarity.

Fig. 17.4 illustrates the case with $n_0 = 2$ and $1 - \epsilon = 90\%$, for which it may be shown that $N = 5.3$. For any given n_0 and desired ϵ we can obtain N from the χ^2 Confidence Level figure because of a relation between the Poisson and the χ^2 : read the ordinate as ϵ , find χ^2 on the curve for $n = 2(n_0 + 1)$; then $N = \chi^2/2$. Some useful values are given in Table 17.3.

The meaning of these upper limits is that, for a given true μ , the probability is at least $1 - \epsilon$ that one will observe n_0 which will result in N which is $\geq \mu$. The probability for that to occur may be higher than $1 - \epsilon$; for example, if $\mu \leq 2.30$ a “90%” upper limit will actually exceed μ 100% of the time. Note from Eq. (17.32) that for $n_0 = 0$, $N = -\ln \epsilon$.

Table 17.3: Poisson upper limits N for n_0 observed events.

n_0	$\epsilon =$		n_0	$\epsilon =$	
	10%	5%		10%	5%
0	2.30	3.00	6	10.53	11.84
1	3.89	4.74	7	11.77	13.15
2	5.32	6.30	8	13.00	14.44
3	6.68	7.75	9	14.21	15.71
4	7.99	9.15	10	15.41	16.96
5	9.27	10.51	11	16.60	18.21

17.5.3. Bounded physical region*:

The measurement of a physical constant α results in an estimator $\hat{\alpha}$, together with some knowledge of experimental error and therefore knowledge of $f(\hat{\alpha}; \alpha)$, the parameterized p.d.f. that allows us to state the probability with which repeated experiments would produce results in a given range. It does *not* permit us to comment about α itself, which in this language is a constant, not a statistical variable. At the beginning of this section we introduced the confidence interval, or frequentist, approach to the problem, and were able to say that with a given probability the unknown parameter could be found between (statistical) limits c_1 and c_2 . But what if a physical boundary exists? Although polarization should be less than one and mass or its square should be greater than zero, experimental results do not always fall inside such a physical boundary because of statistical fluctuations.

However one might set a limit, there is little question about how to report and combine data [14]. A given experiment finds an unbiased estimator $\hat{\alpha} = -5 \pm 10$ for a physical constant (e.g. the square of the mass of a neutrino, in eV^2). This value should be reported as the primary result. In case the true value is zero, for example, this “unphysical” result would not be unlikely. It can be combined with the results of other such experiments by forming the appropriately weighted average of unbiased results, including negative ones, to find an unbiased estimator which expresses our best knowledge of the parameter.

What if we wish to extend our concept of confidence limit to such a situation? The question of how to calculate an upper limit in the vicinity of a physical boundary is one of the most divisive in high-energy physics. We present two main approaches: The confidence interval, or frequentist, method, and the Bayesian method. “Classical method” is applied to one or the other by various writers, so we avoid the term.




Figure 17.5: The situation near a physical boundary. In Fig. 17.1 the horizontal line for a given $\hat{\alpha}_{\text{exp}}$ crossed the domain $D(\epsilon)$, bounded by $\gamma_1(\alpha, \epsilon)$ and $\gamma_2(\alpha, \epsilon)$ entirely in the physical region, entering at c_1 and leaving at c_2 . The limits γ_1 and γ_2 cannot be defined in a region where α is not defined, so the functions cannot be continued into the unphysical region. As a result c_1 (for experiment A) or c_1 and c_2 (for experiment B) cannot be defined. Options 1, 2, and 3 label the ways one might define confidence intervals, as described in the text.

1. *The method of confidence intervals* [1,15]. This is the approach described in the introduction, and requires little further explanation. It is presently the method in favor [1,6,14]. For a Gaussian distribution it gives the same result as the Bayesian approach with a flat prior distribution (see below) if the region containing α with the stated probability is far from an unphysical region, as in Fig. 17.1. Two cases in which this is untrue are shown in Fig. 17.5, where as a matter of convenience we assume that α must be positive. As before, we can define limits γ_1 and γ_2 for each value of the unknown parameter α , such that we can expect that a fraction $1 - \epsilon$ of repeated experiments to produce results between these limits. Since this can be done for each value of α , the limits are described by the functions $\gamma_1(\alpha, \epsilon)$ and $\gamma_2(\alpha, \epsilon)$. However, these *cannot* be extended into a region in which α makes no sense. Experimental result $\hat{\alpha}_{\text{exp A}}$, indicated in Fig. 17.5, is positive, but if the true value is α_{actual} a significant fraction of repetitions of the experiment would produce negative $\hat{\alpha}$. In these cases there is no horizontal intercept c_1 , so without further assumptions we cannot make a statement about the region which would cover α_{actual} in a given fraction of experiments. Experimental result $\hat{\alpha}_{\text{exp B}}$ presents a more serious problem, since it is so negative that there is no physical α for which the point $(\hat{\alpha}, \alpha)$ lies in the domain $D(\epsilon)$. The reason why the frequentist method gives no confidence interval is clear: This measured value of $\hat{\alpha}$ would be unlikely *no matter what the true value of α was*.

There are several *ad hoc* ways to set confidence limits in such cases, although many frequentists would prefer to stop with the weighted average of unbiased results—if the outcome is exceedingly unlikely, one should look to the experiment, not to the statistics. The methods we list below all involve placing c_1 on the physical boundary, which in our example is at $\alpha = 0$.

1. If $\hat{\alpha}_{\text{exp}} > \gamma_1(0, \epsilon)$, as in Experiment A, c_2 is defined. Use it for the upper limit, whether or not $\hat{\alpha}_{\text{exp}} > 0$.
2. If $\hat{\alpha}_{\text{exp}} < 0$, as in Experiment B, use the c_2 corresponding to $\hat{\alpha}_{\text{exp}} = 0$.
3. If c_1 is not defined, “lift up” $\hat{\alpha}$ to $\gamma_2(0, \epsilon)$, where $c_1 = 0$. Use the corresponding c_2 as the upper limit.

These three options are shown in in Fig. 17.5; note that there are regions where more than one of them can be used, with different results. The third option is certainly the most conservative. For Gaussian $f(\hat{\alpha}; \alpha)$ the upper limit c_2 is a one-sided Gaussian confidence limit; read the tables for a 90% two-sided limit to obtain 95% one-sided limit. Alternatively, read the intercepts of the dotted lines in Fig. 17.7. (The horizontal axis is incorrectly labeled for this application.)

2. *The Bayesian approach* [3]. This is the approach favored in the older literature, and has (unfortunately and incorrectly) been referred to as the “PDG method” in certain papers. To begin with, it is argued that while α is not a statistical variable, our *knowledge* of α is less than complete, and it is fair to describe our uncertainty by treating α as a statistical variable. The parameterized p.d.f. $f(\hat{\alpha}; \alpha)$ is replaced by the conditional p.d.f. $f(\hat{\alpha}|\alpha)$. The confidence limit question can then be rephrased: Our measurements provide $f(\hat{\alpha}|\alpha)$, that is, information about $\hat{\alpha}$ for a fixed and unknown value of α , while we really want to know $g(\alpha|\hat{\alpha})$, which tells us that, given our measurement $\hat{\alpha}$, the “true answer” α lies between α and $\alpha + d\alpha$ with probability $g(\alpha|\hat{\alpha}) d\alpha$. The connection is provided by Bayes’ theorem (Eq. (16.7):

$$g(\alpha|\hat{\alpha}) = \frac{f(\hat{\alpha}|\alpha) \pi(\alpha)}{\int f(\hat{\alpha}|\alpha) \pi(\alpha) d\alpha} \tag{17.33}$$

Here $\pi(\alpha)$ represents our “advance knowledge” of the value of α . In the usual case we claim no prior knowledge, so that before the experiment all physically reasonable values of α are equally probable: $\pi(\alpha)$ is a constant over the region of interest and zero in the unphysical region. This assumption leads to the conclusion that

$$g(\alpha|\hat{\alpha}) = \begin{cases} f(\hat{\alpha}|\alpha) / \int f(\hat{\alpha}|\alpha) d\alpha & \text{if } \alpha \text{ is in the physical region;} \\ 0 & \text{otherwise;} \end{cases} \tag{17.34}$$

where this time the integral is over the physical region. In Fig. 17.6 we assume that an ensemble of experiments would produce values for $\hat{\alpha}$ which distribute as shown, with a significant probability of obtaining

results with unphysical values. With our assumed step function $\pi(\alpha)$, the effect of Eq. (17.33) or (17.34) is to replace this distribution with the function shown by the shaded region, except that it is renormalized to unit area. By stating our confidence at the 90% level that α lies below the beginning of the dark shaded region, we mean that 90% of the area in the physical region is in the light shaded region.




Figure 17.6: An example of a bounded physical region, in which a measurement $\hat{\alpha}$ can fall in an unphysical region with significant probability. If we assume that α , the quantity we are trying to measure, cannot lie in the unphysical region (0 probability) but can lie anywhere in the physical region (“no prior knowledge”), then Bayes’ theorem says that our new knowledge of the distribution of α , given our measurement $\hat{\alpha}$, is given by the shaded function after appropriate renormalization.




Figure 17.7: Application of the Bayesian scheme shown in Fig. 17.6 to the case of Gaussian $f(\hat{\alpha}|\alpha)$. For example, if our measurement $\hat{\alpha}$ is 1.0 standard deviations negative, then we conclude that $\alpha < 1.15\sigma$ with 90% probability—however, there is only a 31% probability that an experimental result as low as this would occur. Note that these are upper limits, so that the asymptote for large $\hat{\alpha}/\sigma$ corresponds to a one-sided confidence interval, e.g., the asymptote for a 95% confidence level is $\alpha < \hat{\alpha} + 1.64\sigma$, corresponding to a 90% confidence interval for a two-sided distribution. The dashed lines show the frequentist limit; if Option 3 is used, these are extended horizontally to the right for negative $\hat{\alpha}/\sigma$.

In most cases of interest in this Review, $\hat{\alpha}$ is assumed to be a random value from a Gaussian distribution. Application of the procedure sketched in Fig. 17.6 then leads to the family of curves shown in Fig. 17.7. The confidence limit set by this method is always greater than the [one-sided] confidence interval set without the restriction of an unphysical region, and approaches it from above as the tail in the

unphysical region becomes unimportant. It is also greater than any of the limits shown in Fig. 17.5. With a small modification (exclusion of that portion of the negative tail inside the physical region in the confidence interval definition), it smoothly approaches the usual two-sided confidence interval for Gaussian distributions.

Even so, it is not a valid confidence limit. If it were, the interval would include the true value of α with *exactly* $1 - \epsilon$ probability no matter what the true value was. If the true answer is zero, our procedure, by guaranteeing a limit greater than zero for any experiment, also guarantees that the confidence interval for any ϵ includes α_{actual} with 100% probability. Only as α increases does the probability decrease toward the α -independent Gaussian result.

The error function corresponding to the right axis of Fig. 17.7 shows the probability that $\hat{\alpha}/\sigma$ at or below the given value should occur. If the experimental value is exceeding improbable, then the formal confidence limit obtained by this or any other method means very little.

What about the arbitrariness of $\pi(\alpha)$? If the square of the neutrino mass is measured ($\alpha = m_\nu^2$), then should we not take the prior knowledge distribution as proportional to $\sqrt{\alpha}$? There are other attractive options. Jeffreys points out that if $\pi(\alpha) d\alpha = d\alpha/\alpha$, then the distributions for α and α^n are proportional [16], but there are practical difficulties with this approach. Lynch has investigated prior distributions that are constant in α , α^2 , and $\sqrt{\alpha}$ in the context of Gaussian $f(\hat{\alpha}|\alpha)$, and has observed that assuming a prior distribution that is flat in α gives results that are much more satisfactory than one gets from the others [17]: All three methods have the property that the probability that the calculated limit contains the correct answer is 100% when $\alpha = 0$ and approaches the proper value when $\alpha \gg \sigma$, but the approach to the proper value as α increases is much faster when the prior distribution is taken to be flat in α . In this case the approach is also monotonic, giving it the “conservative” property that for no value of α will the method produce a limit that has a probability of being correct that is less than the stated confidence limit. Although there is nothing unique about the limit calculated with a constant $\pi(\alpha)$, it has desirable features and no obvious replacement.

Summary: If there is a significant probability of obtaining an estimator value corresponding to an unphysical value for a parameter, there is no universally accepted way way to make a statement of the sort “ α is less than c_2 with probability $1 - \epsilon$.” A variety of upper limits can be defined, but no method is entirely satisfactory. The Bayesian method with a flat prior distribution gives a reasonable upper limit which combines everything we know about the unknown quantity α into a physically reasonable value, but it does not give a complete summary of the information contained in the experiment.

17.5.4. Poisson processes with background [18] :

If we observe n_0 events in a Poisson process which has two components, signal and background, estimating a limit on the signal is more complicated. Let μ_S be the unknown mean (the Poisson parameter) for the signal and μ_B be the mean for the sum of all backgrounds. Assume μ_B is known with negligible error; however we don’t know n_B , the actual number of events resulting from the background. We do know that $n_B \leq n_0$. If $\mu_B + \mu_S$ is large, the Gaussian approximation to the Poisson distribution (see Sec. 16.3.2) is usually adequate, and one can define confidence intervals or limits as above, assuming $\hat{n}_B \approx \mu_B$ and therefore $\hat{\mu}_S = n_0 - \mu_B$ with variance equal to n_0 (larger than $\hat{\mu}_S$ to allow for the error in \hat{n}_B).

Otherwise an upper limit can be defined by extension of the argument of the preceding section. Let N be the desired upper limit on μ_S with confidence coefficient ϵ . Set N to be that value of μ_S such that any random repeat of the current experiment with $\mu_S = N$ and the same μ_B would observe *more* than n_0 events in total *and* would have $n_B \leq n_0$, all with probability $1 - \epsilon$. For any assumed N and μ_B we can calculate this probability:

$$1 - \epsilon = 1 - \frac{e^{-(\mu_B + N)} \sum_{n=0}^{n_0} \frac{(\mu_B + N)^n}{n!}}{e^{-\mu_B} \sum_{n=0}^{n_0} \frac{\mu_B^n}{n!}} \tag{17.35}$$

We adjust N to obtain a desired ε . For $\mu_B = 0$ this converges to Eq. (17.32). As in that case (see the last paragraph of Section 17.5.2) this gives a *conservative* upper limit in that for any given true μ_S we get a true probability $\geq 1 - \varepsilon$ that $N \geq \mu_S$, averaged over a large set of identically performed experiments. For $\varepsilon = 0.10$, Fig. 17.8 shows N as a function of n_0 and μ_B .




Figure 17.8: 90% confidence coefficient upper limit on the number of signal events as a function of the expected number of background events. For example, if the expected background is 8 events and 5 events are observed, then the signal is 4.0 (approximately) or less with 90% confidence. Dashed portions indicate regions where it is to be expected that the number observed would exceed the number actually observed $\geq 99\%$ of the time, even in the complete absence of signal.

Averaging of experiments and other comparisons require that n_0 and μ_B be quoted and the technique used for upper limit extraction be given.

If $\mu_B \gg n_0$ the experimenter should question the probability of observing n_B as that n_0 . If this is very small the background, μ_B , may not have been calculated properly and the upper limit for μ_S obtained under those assumptions may be too low. For example, in Fig. 17.8, the dashed portions of the curves lie in the region where n_0 is expected to exceed the observed value 99% of the time (or more), even in the complete absence of signal. In these regions one should be cautious about accepting the results of the measurement.

As in the Gaussian case (Sec. 17.5.3), whenever $n_0 < \mu_B$ some experimenters may prefer to use N calculated as if $n_0 \approx \mu_B$ rather than the smaller value obtained from the observed n_0 .

17.6. Propagation of errors

Suppose we have a set of N random variables y_i which may be direct measurements or derived estimators $\hat{\alpha}_i$, and we have a covariance matrix $V(y)$ for these. We can make a transformation to a different set of variables $f_n \equiv f_n(y)$, $j = 1, \dots, M$ ($M \leq N$) and obtain best estimates for the f_n from

$$\hat{f}_n \approx f_n(\hat{y}) + \frac{1}{2} \sum_{k,n} V_{kn}(\hat{y}) \left[\frac{\partial^2 f_n}{\partial y_k \partial y_n} \right]_{\hat{y}} \quad (17.36)$$

with covariance matrix

$$V_{ij}(\hat{f}) \approx \sum_{n,m} \frac{\partial f_i}{\partial y_n} \bigg|_{\hat{y}} \frac{\partial f_j}{\partial y_m} \bigg|_{\hat{y}} V_{nm}(\hat{y}) . \quad (17.37)$$

For a single-valued function f of a single measurement y with variance σ^2 (i.e., $M = 1, N = 1$), this becomes

$$\begin{aligned} \hat{f} &\approx f(\hat{y}) + \frac{1}{2} \sigma^2 f''(\hat{y}) \\ V(\hat{f}) &\approx \sigma^2 [f'(\hat{y})]^2 , \end{aligned} \quad (17.38)$$

where the primes denote differentiation with respect to y , evaluated at \hat{y} .

These approximations are based on a Taylor expansion of f about the true value of y . If f is approximately linear in y over a range of roughly $\hat{y}_i \pm \sigma(y_i)$, the approximation is good and the second-order terms in (17.36) and (17.38) can be neglected. This is what is usually done. However, if linearity is badly violated (e.g., $f \propto 1/y$ and \hat{y} is no more than a few σ from zero), it should be recognized that propagation of errors will give very approximate results. In such cases $\hat{f} \approx f(\hat{y})$ may be a biased estimator for f even if \hat{y} is unbiased for y , and the second-order terms in (17.36) and (17.38) will help to reduce that bias.

*In addition to the references cited, communications with R.D. Cousins, F. James, G. Lynch, and B. Roe have been invaluable in formulating this section.

†There is an additional problem: Even if we set a confidence limit on m_ν^2 by some particular recipe, it translates into a different confidence limit on $\sqrt{m_\nu^2}$ except when a Bayesian procedure with Jeffreys' $\pi(\alpha) \propto 1/\alpha$ prior distribution is used.

References:

1. H. Cramér, *Mathematical Methods of Statistics*, Princeton Univ. Press, New Jersey (1958).
2. J. Orear, "Notes on Statistics for Physicists," Univ. of Calif. Radiation Lab. Report UCRL-8417 (1958), unpublished; "Notes on Statistics for Physicists, Revised," Cornell Univ. Report CLNS 82/511 (1982), unpublished.
3. F.T. Solmitz, *Annu. Rev. Nucl. Sci.* **14**, 375–402 (1964).
4. W.T. Eadie, D. Drijard, F.E. James, M. Roos, and B. Sadoulet, *Statistical Methods in Experimental Physics* (North Holland, Amsterdam and London, 1971).
5. L. Lyons, *Statistics for Nuclear and Particle Physicists*, (Cambridge University Press, New York (1986)).
6. B.P. Roe, *Probability and Statistics in Experimental Physics*, (Springer-Verlag, New York, 208 pp., 1992).
7. S. Baker and R. Cousins, *Nucl. Instr. and Meth.* **221**, 437 (1984).
8. W.H. Press *et al.*, *Numerical Recipes* (Cambridge University Press, New York, 1986).
9. W.H. Maindonald *et al.*, *Statistical Computation* (John Wiley and Sons, Inc., New York, 1984).
10. A. Basilevsky *et al.*, *Applied Matrix Algebra in the Statistical Sciences* (North Holland, New York, 1983).
11. F. James and M. Roos, "MINUIT, Function Minimization and Error Analysis," CERN D506 (Long Writeup). Available from the CERN Program Library Office, CERN-DD Division, CERN, CH-1211, Geneva 21, Switzerland.
12. Microsoft ® is a registered trademark of Microsoft Corporation.
13. M. Abramowitz and I. Stegun, eds., *Handbook of Mathematical Functions* (Dover, New York, 1972).
14. F. James and M. Roos, *Phys. Rev.* **D44**, 299 (1991).
15. J. Neyman, *Lectures and Conferences on Mathematical Statistics*, Washington (1938).
16. H. Jeffreys, *Theory of Probability*, 3rd ed. (Oxford Univ. Press, Oxford, England, 447 pp., 1961).
17. G. Lynch, "Calculations of Upper Limits," Particle Data Group Note PDG-93-03 (Sept. 1993) (unpublished).
18. O. Helene, *Nucl. Instr. and Meth.* **212**, 319 (1983).

18. MONTE CARLO TECHNIQUES

Revised June 1994 with the help of T. Adye, O. Dahl, and H.-J. Trost.

Monte Carlo techniques are used to simulate on a computer random behavior which is too complex to be derived analytically. Most calculations are based upon pseudorandom numbers, a reproducible sequence of numbers generated on the open interval (0,1) in such a way that they satisfy various statistical tests for a uniform distribution, with independent numbers. (Caution: some commercial random number generators fill the closed interval [0,1]. The occurrence of 0 or 1 can sometimes cause problems for the algorithms below). No such numbers are truly uniform and independent. Many commercial random number generators sacrifice randomness in favor of speed. It is not rare that unforeseen correlations will introduce non-negligible errors in the results. A useful test for this is to recompute the same results with a different algorithm for the pseudorandom numbers. To improve the performance of an existing generator one may use the Bays-Durham algorithm [see Ref. 1 for discussion]: (a) Initialize by generating and storing N (e.g., $N = 97$) random numbers in an array v , using the available generator. Generate a new random number u and save it. (b) On the next call, use this u as an address $j = 1 + (\text{integer part of } Nu)$ to select v_j as the random number to be returned. Also save this v_j as u for the next call. Replace v_j in the array with a new random number using the available generator. On the next call, go to (b).

A second problem sometimes encountered in computations requiring long sequences of random numbers is that all pseudorandom number generators will eventually begin over and repeat the same sequence. One may choose algorithms which minimize the number used. One may also use two or three different generators in different parts of the program.

Monte Carlo simulations of complex processes break them down into a sequence of steps. At each step a particular outcome is chosen from a set of possibilities according to a certain p.d.f. To do this we must transform our uniform random numbers into random numbers sampled from different distributions on different ranges.

Two techniques are in wide use to do this. We will discuss only single variable cases; multiple variable cases use straightforward extensions of these techniques. We assume we are in possession of a random number u chosen from a uniform distribution on (0,1).

18.1. Inverse transform method

If the desired probability density function is $f(x)$ on the range $-\infty < x < \infty$, its cumulative distribution function (expressing the probability that $x \leq a$) is given by Eq. (16.1). If a is chosen with probability density $f(a)$, then the integrated probability up to point a , $F(a)$, is itself a random variable which will occur with uniform probability density on [0, 1]. If x can take on any value, and ignoring the endpoints, we can then find a unique x chosen from the p.d.f. $f(x)$ for a given u if we set

$$u = F(x) , \tag{18.1}$$

provided we can find an inverse of F , defined by

$$x = F^{-1}(u) . \tag{18.2}$$

This method is shown in Fig. 18.1a.

For a discrete distribution, $F(x)$ will have a discontinuous jump of size $f(x_k)$ at each allowed $x_k, k = 1, 2, \dots$. Choose u from a uniform distribution on (0,1) as before. Find x_k such that

$$F(x_{k-1}) < u \leq F(x_k) \equiv \text{Prob} (x \leq x_k) = \sum_{i=1}^k f(x_i) ; \tag{18.3}$$

then x_k is the value we seek (note: $F(x_0) \equiv 0$). This algorithm is illustrated in Fig. 18.1b.




Figure 18.1: Use of a random number u chosen from a uniform distribution (0,1) to find a random number x from a distribution with cumulative distribution function $F(x)$.

18.2. Acceptance-rejection method (Von Neumann)

Very commonly an analytic form for $F(x)$ is unknown or too complex to work with, so that obtaining an inverse as in Eq. (18.2) is impractical. We suppose that for any given value of x the probability density function $f(x)$ can be computed and further that enough is known about $f(x)$ that we can enclose it entirely inside a shape which is C times an easily generated distribution $h(x)$ as illustrated in Fig. 18.2.




Figure 18.2: Illustration of the acceptance-rejection method. Random points are chosen inside the upper bounding figure, and rejected if the ordinate exceeds $f(x)$. Lower figure illustrates importance sampling.

Frequently $h(x)$ is uniform or is a normalized sum of uniform distributions. Note that both $f(x)$ and $h(x)$ must be normalized to unit area and therefore the proportionality constant $C > 1$. To generate $f(x)$, first generate a candidate x according to $h(x)$. Calculate $f(x)$ and the height of the envelope $C h(x)$; generate u and test if $u C h(x) \leq f(x)$. If so, accept x ; if not reject x and try again. If we regard x and $u C h(x)$ as the abscissa and ordinate of a point in a two-dimensional plot, these points will populate the entire area $C h(x)$ in a smooth manner; then we accept those which fall under $f(x)$. The efficiency is the ratio of areas, which must equal $1/C$; therefore we must keep C as close as possible to 1.0. Therefore we try to choose $C h(x)$ to be as close to $f(x)$ as convenience dictates, as in the lower part of Fig. 18.2. This practice is called importance sampling, because

we generate more trial values of x in the region where $f(x)$ is most important.

18.3. Algorithms

Algorithms for generating random numbers belonging to many different distributions are given by Press [1], Ahrens and Dieter [2], Rubinstein [3], Everett and Cashwell [4], Devroye [5], and Walck [6]. For many distributions alternative algorithms exist, varying in complexity, speed, and accuracy. For time-critical applications, these algorithms may be coded in-line to remove the significant overhead often encountered in making function calls. Variables named “ u ” are assumed to be independent and uniform on $(0,1)$.

In the examples given below, we use the notation for the variables and parameters given in Table 16.1.

18.3.1. Sine and cosine of random angle:

Generate u_1 and u_2 . Then $v_1 = 2u_1 - 1$ is uniform on $(-1,1)$, and $v_2 = u_2$ is uniform on $(0,1)$. Calculate $r^2 = v_1^2 + v_2^2$. If $r^2 > 1$, start over. Otherwise, the sine (S) and cosine (C) of a random angle are given by

$$S = 2v_1v_2/r^2 \quad \text{and} \quad C = (v_1^2 - v_2^2)/r^2. \tag{18.4}$$

18.3.2. Gaussian distribution:

If u_1 and u_2 are uniform on $(0,1)$, then

$$z_1 = \sin 2\pi u_1 \sqrt{-2 \ln u_2} \quad \text{and} \quad z_2 = \cos 2\pi u_1 \sqrt{-2 \ln u_2} \tag{18.5}$$

are independent and Gaussian distributed with mean 0 and $\sigma = 1$.

There are many faster variants of this basic algorithm. For example, construct $v_1 = 2u_1 - 1$ and $v_2 = 2u_2 - 1$, which are uniform on $(-1,1)$. Calculate $r^2 = v_1^2 + v_2^2$, and if $r^2 > 1$ start over. If $r^2 < 1$, it is uniform on $(0,1)$. Then

$$z_1 = v_1 \sqrt{\frac{-2 \ln r^2}{r^2}} \quad \text{and} \quad z_2 = v_2 \sqrt{\frac{-2 \ln r^2}{r^2}} \tag{18.6}$$

are independent numbers chosen from a normal distribution with mean 0 and variance 1. $z'_i = \mu + \sigma z_i$ distributes with mean μ and variance σ^2 .

For a multivariate Gaussian, see the algorithm in Ref. 7.

18.3.3. $\chi^2(n)$ distribution:

For n even, generate $n/2$ uniform numbers u_i ; then

$$y = -2 \ln \left(\prod_{i=1}^{n/2} u_i \right) \quad \text{is} \quad \chi^2(n). \tag{18.7}$$

For n odd, generate $(n - 1)/2$ uniform numbers u_i and one Gaussian z as in Sec. 18.3.2; then

$$y = -2 \ln \left(\prod_{i=1}^{(n-1)/2} u_i \right) + z^2 \quad \text{is} \quad \chi^2(n). \tag{18.8}$$

For $n \gtrsim 30$ the much faster Gaussian approximation for the χ^2 may be preferable: generate z as in Sec. 18.3.2 and use $y = [z + \sqrt{2n - 1}]^2 / 2$; if $z < -\sqrt{2n - 1}$ reject and start over.

18.3.4. Gamma distribution:

All of the following algorithms are given for $\lambda = 1$. For $\lambda \neq 1$, divide the resulting random number x by λ .

- If $k = 1$ (the exponential distribution), accept $x = -(\ln u)$.
- If $0 < k < 1$, initialize with $v_1 = (e + k)/e$ (with $e = 2.71828\dots$ being the natural log base). Generate u_1, u_2 . Define $v_2 = v_1 u_1$.

Case 1: $v_2 \leq 1$. Define $x = v_2^{1/k}$. If $u_2 \leq e^{-x}$, accept x and stop, else restart by generating new u_1, u_2 .

Case 2: $v_2 > 1$. Define $x = -\ln([v_1 - v_2]/k)$. If $u_2 \leq x^{k-1}$, accept x and stop, else restart by generating new u_1, u_2 .

Note that, for $k < 1$, the probability density has a pole at $x = 0$, so that return values of zero due to underflow must be accepted or otherwise dealt with.

- Otherwise, if $k > 1$, initialize with $c = 3k - 0.75$. Generate u_1 and compute $v_1 = u_1(1 - u_1)$ and $v_2 = (u_1 - 0.5)\sqrt{c/v_1}$. If $x = k + v_2 - 1 \leq 0$, go back and generate new u_1 ; otherwise generate u_2 and compute $v_3 = 64v_1^3 u_2^2$. If $v_3 \leq 1 - 2v_2^2/x$ or if $\ln v_3 \leq 2\{[k - 1] \ln|x/(k - 1)| - v_2\}$, accept x and stop; otherwise go back and generate new u_1 .

18.3.5. Binomial distribution:

If $p \leq 1/2$, iterate until a successful choice is made: begin with $k = 1$; compute $P_k = q^n$ [for $k \neq 1$ use $P_k \equiv f(r_k; n, p)$], and store P_k into B ; generate u . If $u \leq B$ accept $r_k = k - 1$ and stop; otherwise increment k by 1 and compute next P_k and add to B ; generate a new u and repeat. If we arrive at $k = n + 1$, stop and accept $r_{n+1} = n$. If $p > 1/2$ it will be more efficient to generate r from $f(r; n, q)$, i.e., with p and q interchanged, and then set $r_k = n - r$.

18.3.6. Poisson distribution:

Iterate until a successful choice is made: Begin with $k = 1$ and set $A = 1$ to start. Generate u . Replace A with uA ; if now $A < \exp(-\mu)$, where μ is the Poisson parameter, accept $n_k = k - 1$ and stop. Otherwise increment k by 1, generate a new u and repeat, always starting with the value of A left from the previous try. For large $\mu (\gtrsim 10)$ it may be satisfactory (and much faster) to approximate the Poisson distribution by a Gaussian distribution (see our Probability chapter, Sec. 16.3.3) and generate z from $f(z; 0, 1)$; then accept $x = \max(0, [\mu + z\sqrt{\mu} + 0.5])$ where $[\]$ signifies the greatest integer \leq the expression.

18.3.7. Student's t distribution:

For $n > 0$ degrees of freedom (n not necessarily integer), generate x from a Gaussian with mean 0 and $\sigma^2 = 1$ according to the method of 18.3.2. Next generate y , an independent gamma random variate with $k = n/2$ degrees of freedom. Then $z = x\sqrt{2n}/\sqrt{y}$ is distributed as a t with n degrees of freedom.

For the special case $n = 1$, the Breit-Wigner distribution, generate u_1 and u_2 ; set $v_1 = 2u_1 - 1$ and $v_2 = 2u_2 - 1$. If $v_1^2 + v_2^2 \leq 1$ accept $z = v_1/v_2$ as a Breit-Wigner distribution with unit area, center at 0.0, and FWHM 2.0. Otherwise start over. For center M_0 and FWHM Γ , use $W = z\Gamma/2 + M_0$.

References:

1. W.H. Press *et al.*, *Numerical Recipes* (Cambridge University Press, New York, 1986).
2. J.H. Ahrens and U. Dieter, *Computing* **12**, 223 (1974).
3. R.Y. Rubinstein, *Simulation and the Monte Carlo Method* (John Wiley and Sons, Inc., New York, 1981).
4. C.J. Everett and E.D. Cashwell, *A Third Monte Carlo Sampler*, Los Alamos report LA-9721-MS (1983).
5. L. Devroye, *Non-Uniform Random Variate Generation* (Springer-Verlag, New York, 1986).
6. Ch. Walck, *Random Number Generation*, University of Stockholm Physics Department Report 1987-10-20 (Vers. 3.0).
7. F. James, *Rept. on Prog. in Phys.* **43**, 1145 (1980).

19. ELECTROMAGNETIC RELATIONS

Quantity	Gaussian CGS	SI
Charge:	$2.997\,924\,58 \times 10^9$ esu	$= 1\text{ C} = 1\text{ A s}$
Electron charge e :	$4.803\,206\,8 \times 10^{-10}$ esu	$= 1.602\,177\,33 \times 10^{-19}\text{ C}$
Potential:	$(1/299.792\,458)$ statvolt (ergs/esu)	$= 1\text{ V} = 1\text{ J C}^{-1}$
Magnetic field:	10^4 gauss $= 10^4$ dyne/esu	$= 1\text{ T} = 1\text{ N A}^{-1}\text{m}^{-1}$
Lorentz force:	$\mathbf{F} = q(\mathbf{E} + \frac{\mathbf{v}}{c} \times \mathbf{B})$	$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B})$
Maxwell equations:	$\nabla \cdot \mathbf{D} = 4\pi\rho$ $\nabla \times \mathbf{H} - \frac{1}{c} \frac{\partial \mathbf{D}}{\partial t} = \frac{4\pi}{c} \mathbf{J}$ $\nabla \cdot \mathbf{B} = 0$ $\nabla \times \mathbf{E} + \frac{1}{c} \frac{\partial \mathbf{B}}{\partial t} = 0$	$\nabla \cdot \mathbf{D} = \rho$ $\nabla \times \mathbf{H} - \frac{\partial \mathbf{D}}{\partial t} = \mathbf{J}$ $\nabla \cdot \mathbf{B} = 0$ $\nabla \times \mathbf{E} + \frac{\partial \mathbf{B}}{\partial t} = 0$
Materials:	$\mathbf{D} = \epsilon\mathbf{E}, \quad \mathbf{H} = \mathbf{B}/\mu$	$\mathbf{D} = \epsilon\mathbf{E}, \quad \mathbf{H} = \mathbf{B}/\mu$
Permittivity of free space:	1	$\epsilon_0 = 8.854\,187 \dots \times 10^{-12}\text{ F m}^{-1}$
Permeability of free space:	1	$\mu_0 = 4\pi \times 10^{-7}\text{ N A}^{-2}$
Fields from potentials:	$\mathbf{E} = -\nabla V - \frac{1}{c} \frac{\partial \mathbf{A}}{\partial t}$ $\mathbf{B} = \nabla \times \mathbf{A}$	$\mathbf{E} = -\nabla V - \frac{\partial \mathbf{A}}{\partial t}$ $\mathbf{B} = \nabla \times \mathbf{A}$
Static potentials: (coulomb gauge)	$V = \sum_{\text{charges}} \frac{q_i}{r_i} = \int \frac{\rho(\mathbf{r}')}{ \mathbf{r} - \mathbf{r}' } d^3x'$ $\mathbf{A} = \frac{1}{c} \sum_{\text{currents}} \frac{\mathbf{I}_i}{r_i} = \frac{1}{c} \int \frac{\mathbf{J}(\mathbf{r}')}{ \mathbf{r} - \mathbf{r}' } d^3x'$	$V = \frac{1}{4\pi\epsilon_0} \sum_{\text{charges}} \frac{q_i}{r_i} = \frac{1}{4\pi\epsilon_0} \int \frac{\rho(\mathbf{r}')}{ \mathbf{r} - \mathbf{r}' } d^3x'$ $\mathbf{A} = \frac{\mu_0}{4\pi} \sum_{\text{currents}} \frac{\mathbf{I}_i}{r_i} = \frac{\mu_0}{4\pi} \int \frac{\mathbf{J}(\mathbf{r}')}{ \mathbf{r} - \mathbf{r}' } d^3x'$
Relativistic transformations: (\mathbf{v} is the velocity of the primed frame as seen in the unprimed frame)	$\mathbf{E}'_{\parallel} = \mathbf{E}_{\parallel}$ $\mathbf{E}'_{\perp} = \gamma(\mathbf{E}_{\perp} + \frac{1}{c} \mathbf{v} \times \mathbf{B})$ $\mathbf{B}'_{\parallel} = \mathbf{B}_{\parallel}$ $\mathbf{B}'_{\perp} = \gamma(\mathbf{B}_{\perp} - \frac{1}{c} \mathbf{v} \times \mathbf{E})$	$\mathbf{E}'_{\parallel} = \mathbf{E}_{\parallel}$ $\mathbf{E}'_{\perp} = \gamma(\mathbf{E}_{\perp} + \mathbf{v} \times \mathbf{B})$ $\mathbf{B}'_{\parallel} = \mathbf{B}_{\parallel}$ $\mathbf{B}'_{\perp} = \gamma(\mathbf{B}_{\perp} - \frac{1}{c^2} \mathbf{v} \times \mathbf{E})$
	$\frac{1}{4\pi\epsilon_0} = c^2 \times 10^{-7}\text{ N A}^{-2} = 8.987\,55 \dots \times 10^9\text{ m F}^{-1}; \quad \frac{\mu_0}{4\pi} = 10^{-7}\text{ N A}^{-2};$	$c = \frac{1}{\sqrt{\mu_0\epsilon_0}} = 2.997\,924\,58 \times 10^8\text{ m s}^{-1}$

19.1. Impedances (SI units)

ρ = resistivity at room temperature in $10^{-8} \Omega \text{ m}$:
 ~ 1.7 for Cu ~ 5.5 for W
 ~ 2.4 for Au ~ 73 for SS 304
 ~ 2.8 for Al ~ 100 for Nichrome
 (Al alloys may have double the Al value.)

For alternating currents, instantaneous current I , voltage V , angular frequency ω :

$$V = V_0 e^{j\omega t} = ZI . \tag{19.1}$$

Impedance of self-inductance L : $Z = j\omega L$.

Impedance of capacitance C : $Z = 1/j\omega C$.

Impedance of free space: $Z = \sqrt{\mu_0/\epsilon_0} = 376.7 \Omega$.

Impedance per unit length of a flat conductor of width w (high frequency, ν):

$$Z = \frac{(1+j)\rho}{w\delta} , \quad \text{where } \delta = \text{effective skin depth} ; \tag{19.2}$$

$$\delta = \sqrt{\frac{\rho}{\pi\nu\mu}} \approx \frac{6.6 \text{ cm}}{\sqrt{\nu(\text{Hz})}} \quad \text{for Cu} . \tag{19.3}$$

19.2. Capacitance \hat{C} and inductance \hat{L} per unit length (SI units)

Flat rectangular plates of width w , separated by $d \ll w$:

$$\hat{C} = \epsilon \frac{w}{d} ; \quad \hat{L} = \mu \frac{d}{w} ; \tag{19.4}$$

$$\frac{\epsilon}{\epsilon_0} = 2 \text{ to } 6 \text{ for plastics; } 4 \text{ to } 8 \text{ for porcelain, glasses.} \tag{19.5}$$

Coaxial cable of inner radius r_1 , outer radius r_2 :

$$\hat{C} = \frac{2\pi\epsilon}{\ln(r_2/r_1)} ; \quad \hat{L} = \frac{\mu}{2\pi} \ln(r_2/r_1) . \tag{19.6}$$

Transmission lines (no loss):

$$\text{Impedance: } Z = \sqrt{\hat{L}/\hat{C}} . \tag{19.7}$$

$$\text{Velocity: } v = 1/\sqrt{\hat{L}\hat{C}} = 1/\sqrt{\mu\epsilon} . \tag{19.8}$$

19.3. Synchrotron radiation (CGS units)

For a particle of charge e , velocity $v = \beta c$, and energy $E = \gamma mc^2$, traveling in a circular orbit of radius R , the energy loss per revolution δE is

$$\delta E = \frac{4\pi}{3} \frac{e^2}{R} \beta^3 \gamma^4 . \tag{19.9}$$

For high-energy electrons or positrons ($\beta \approx 1$), this becomes

$$\delta E \text{ (in MeV)} \approx 0.0885 [E(\text{in GeV})]^4 / R(\text{in m}) . \tag{19.10}$$

For $\gamma \gg 1$, the energy radiated per revolution into the photon energy interval $d(\hbar\omega)$ is

$$dI = \frac{8\pi}{9} \alpha \gamma F(\omega/\omega_c) d(\hbar\omega) , \tag{19.11}$$

where $\alpha = e^2/\hbar c$ is the fine-structure constant and

$$\omega_c = \frac{3\gamma^3 c}{2R} \tag{19.12}$$

is the critical frequency. The normalized function $F(y)$ is

$$F(y) = \frac{9}{8\pi} \sqrt{3} y \int_y^\infty K_{5/3}(x) dx , \tag{19.13}$$

where $K_{5/3}(x)$ is a modified Bessel function of the third kind. For electrons or positrons,

$$\hbar\omega_c \text{ (in keV)} \approx 2.22 [E(\text{in GeV})]^3 / R(\text{in m}) . \tag{19.14}$$

Fig. 19.1 shows $F(y)$ over the important range of y .




Figure 19.1: The normalized synchrotron radiation spectrum $F(y)$.

For $\gamma \gg 1$ and $\omega \ll \omega_c$,

$$\frac{dI}{d(\hbar\omega)} \approx 3.3\alpha (\omega R/c)^{1/3} , \tag{19.15}$$

whereas for

$\gamma \gg 1$ and $\omega \gtrsim 3\omega_c$,

$$\frac{dI}{d(\hbar\omega)} \approx \sqrt{\frac{3\pi}{2}} \alpha \gamma \left(\frac{\omega}{\omega_c}\right)^{1/2} e^{-\omega/\omega_c} \left[1 + \frac{55}{72} \frac{\omega_c}{\omega} + \dots\right] . \tag{19.16}$$

The radiation is confined to angles $\lesssim 1/\gamma$ relative to the instantaneous direction of motion.

See J.D. Jackson, *Classical Electrodynamics*, 2nd edition (John Wiley & Sons, New York, 1975) for more formulae and details. In his book, Jackson uses a definition of ω_c that is twice as large as the customary one given above.

20. CLEBSCH-GORDAN COEFFICIENTS, SPHERICAL HARMONICS, AND d FUNCTIONS

Note: A $\sqrt{\quad}$ is to be understood over every coefficient, e.g., for $-8/15$ read $-\sqrt{8/15}$.

Notation:

J	J	...
M	M	...
m_1	m_2	
m_1	m_2	Coefficients
.	.	
.	.	
.	.	

$Y_1^0 = \sqrt{\frac{3}{4\pi}} \cos \theta$

$Y_1^1 = -\sqrt{\frac{3}{8\pi}} \sin \theta e^{i\phi}$

$Y_2^0 = \sqrt{\frac{5}{4\pi}} \left(\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right)$

$Y_2^1 = -\sqrt{\frac{15}{8\pi}} \sin \theta \cos \theta e^{i\phi}$

$Y_2^2 = \frac{1}{4} \sqrt{\frac{15}{2\pi}} \sin^2 \theta e^{2i\phi}$

$Y_l^{-m} = (-1)^m Y_l^{m*}$

$d_{m,0}^l = \sqrt{\frac{4\pi}{2l+1}} Y_l^m e^{-im\phi}$

$\langle j_1 j_2 m_1 m_2 | j_1 j_2 J M \rangle$
 $= (-1)^{J-j_1-j_2} \langle j_2 j_1 m_2 m_1 | j_2 j_1 J M \rangle$

$d_{m',m}^j = (-1)^{m-m'} d_{m,m'}^j = d_{-m,m'}^j$

$d_{1/2,1/2}^{1/2} = \cos \frac{\theta}{2}$

$d_{1/2,-1/2}^{1/2} = -\sin \frac{\theta}{2}$

$d_{1,1}^1 = \frac{1 + \cos \theta}{2}$

$d_{1,0}^1 = -\frac{\sin \theta}{\sqrt{2}}$

$d_{1,-1}^1 = \frac{1 - \cos \theta}{2}$

$d_{0,0}^1 = \cos \theta$

$d_{3/2,3/2}^{3/2} = \frac{1 + \cos \theta}{2} \cos \frac{\theta}{2}$

$d_{3/2,1/2}^{3/2} = -\sqrt{3} \frac{1 + \cos \theta}{2} \sin \frac{\theta}{2}$

$d_{3/2,-1/2}^{3/2} = \sqrt{3} \frac{1 - \cos \theta}{2} \cos \frac{\theta}{2}$

$d_{3/2,-3/2}^{3/2} = -\frac{1 - \cos \theta}{2} \sin \frac{\theta}{2}$

$d_{1/2,1/2}^{3/2} = \frac{3 \cos \theta + 1}{2} \cos \frac{\theta}{2}$

$d_{1/2,-1/2}^{3/2} = -\frac{3 \cos \theta + 1}{2} \sin \frac{\theta}{2}$

$d_{2,2}^2 = \left(\frac{1 + \cos \theta}{2} \right)^2$

$d_{2,1}^2 = -\frac{1 + \cos \theta}{2} \sin \theta$

$d_{2,0}^2 = \frac{\sqrt{6}}{4} \sin^2 \theta$

$d_{2,-1}^2 = -\frac{1 - \cos \theta}{2} \sin \theta$

$d_{2,-2}^2 = \left(\frac{1 - \cos \theta}{2} \right)^2$

$d_{1,1}^2 = \frac{1 + \cos \theta}{2} (2 \cos \theta - 1)$

$d_{1,0}^2 = -\sqrt{\frac{3}{2}} \sin \theta \cos \theta$

$d_{1,-1}^2 = \frac{1 - \cos \theta}{2} (2 \cos \theta + 1)$

$d_{0,0}^2 = \left(\frac{3}{2} \cos^2 \theta - \frac{1}{2} \right)$

Figure 20.1: Sign convention is that of Wigner (*Group Theory*, Academic Press, New York, 1959), also used by Condon and Shortley (*The Theory of Atomic Spectra*, Cambridge Univ. Press, New York, 1953), Rose (*Elementary Theory of Angular Momentum*, Wiley, New York, 1957), and Cohen (*Tables of the Clebsch-Gordan Coefficients*, North American Rockwell Science Center, Thousand Oaks, Calif., 1974). The signs and numbers in the current tables have been calculated by computer programs written independently by Cohen and at LBL.

21. SU(3) ISOSCALAR FACTORS AND REPRESENTATION MATRICES

The most commonly used SU(3) isoscalar factors, corresponding to the singlet, octet, and decuplet content of $8 \otimes 8$ and $10 \otimes 8$, are shown at the right. The notation uses particle names to identify the coefficients, so that the pattern of relative couplings may be seen at a glance. We illustrate the use of the coefficients below. See J.J. de Swart, Rev. Mod. Phys. 35, 916 (1963) for detailed explanations and phase conventions.

A $\sqrt{\quad}$ is to be understood over every integer in the matrices; the exponent 1/2 on each matrix is a reminder of this. For example, the $\Xi \rightarrow \Omega K$ element of the $10 \rightarrow 10 \otimes 8$ matrix is $-\sqrt{6}/\sqrt{24} = -1/2$.

Intramultiplet relative decay strengths may be read directly from the matrices. For example, in decuplet \rightarrow octet + octet decays, the ratio of $\Omega^* \rightarrow \Xi \bar{K}$ and $\Delta \rightarrow N \pi$ partial widths is, from the $10 \rightarrow 8 \times 8$ matrix,

$$\frac{\Gamma(\Omega^* \rightarrow \Xi \bar{K})}{\Gamma(\Delta \rightarrow N \pi)} = \frac{12}{6} \times (\text{phase space factors}) . \quad (21.1)$$

Including isospin Clebsch-Gordan coefficients, we obtain, e.g.,

$$\frac{\Gamma(\Omega^{*-} \rightarrow \Xi^0 K^-)}{\Gamma(\Delta^+ \rightarrow p \pi^0)} = \frac{1/2}{2/3} \times \frac{12}{6} \times p.s.f. = \frac{3}{2} \times p.s.f. \quad (21.2)$$

Partial widths for $8 \rightarrow 8 \otimes 8$ involve a linear superposition of 8_1 (symmetric) and 8_2 (antisymmetric) couplings. For example,

$$\Gamma(\Xi^* \rightarrow \Xi \pi) \sim \left(-\sqrt{\frac{9}{20}} g_1 + \sqrt{\frac{3}{12}} g_2 \right)^2 . \quad (21.3)$$

The relations between g_1 and g_2 (with de Swart's normalization) and the standard D and F couplings that appear in the interaction Lagrangian,

$$\mathcal{L} = -\sqrt{2} D \text{Tr}(\{\bar{B}, B\} M) + \sqrt{2} F \text{Tr}([\bar{B}, B] M) , \quad (21.4)$$

where $[\bar{B}, B] \equiv \bar{B}B - B\bar{B}$ and $\{\bar{B}, B\} \equiv \bar{B}B + B\bar{B}$, are

$$D = \frac{\sqrt{30}}{40} g_1 , \quad F = \frac{\sqrt{6}}{24} g_2 . \quad (21.5)$$

Thus, for example,

$$\Gamma(\Xi^* \rightarrow \Xi \pi) \sim (F - D)^2 \sim (1 - 2\alpha)^2 , \quad (21.6)$$

where $\alpha \equiv D/(D + F)$.

When acting upon a representation of dimension d , the generators of SU(3) transformations, λ_a ($a = 1, 8$), are $d \times d$ matrices that obey the following commutation and anticommutation relationships:

$$[\lambda_a, \lambda_b] \equiv \lambda_a \lambda_b - \lambda_b \lambda_a = 2i f_{abc} \lambda_c \quad (21.7)$$

$$\{\lambda_a, \lambda_b\} \equiv \lambda_a \lambda_b + \lambda_b \lambda_a = \frac{4}{3} \delta_{ab} I + 2d_{abc} \lambda_c , \quad (21.8)$$

where I is the $d \times d$ identity matrix, and δ_{ab} is the Kronecker delta symbol. The f_{abc} are odd under the permutation of any pair of indices, while the d_{abc} are even. The nonzero values are

$1 \rightarrow 8 \otimes 8$

$$(\Lambda) \rightarrow (N \bar{K} \Sigma \pi \Lambda \eta \Xi K) = \frac{1}{\sqrt{8}} (2 \ 3 \ -1 \ -2)^{1/2}$$

$8_1 \rightarrow 8 \otimes 8$

$$\begin{pmatrix} N \\ \Sigma \\ \Lambda \\ \Xi \end{pmatrix} \rightarrow \begin{pmatrix} N\pi & N\eta & \Sigma K & \Lambda K \\ N\bar{K} & \Sigma\pi & \Lambda\pi & \Sigma\eta & \Xi K \\ N\bar{K} & \Sigma\pi & \Lambda\eta & \Xi K \\ \Sigma\bar{K} & \Lambda\bar{K} & \Xi\pi & \Xi\eta \end{pmatrix} = \frac{1}{\sqrt{20}} \begin{pmatrix} 9 & -1 & -9 & -1 \\ -6 & 0 & 4 & 4 & -6 \\ 2 & -12 & -4 & -2 \\ 9 & -1 & -9 & -1 \end{pmatrix}^{1/2}$$

$8_2 \rightarrow 8 \otimes 8$

$$\begin{pmatrix} N \\ \Sigma \\ \Lambda \\ \Xi \end{pmatrix} \rightarrow \begin{pmatrix} N\pi & N\eta & \Sigma K & \Lambda K \\ N\bar{K} & \Sigma\pi & \Lambda\pi & \Sigma\eta & \Xi K \\ N\bar{K} & \Sigma\pi & \Lambda\eta & \Xi K \\ \Sigma\bar{K} & \Lambda\bar{K} & \Xi\pi & \Xi\eta \end{pmatrix} = \frac{1}{\sqrt{12}} \begin{pmatrix} 3 & 3 & 3 & -3 \\ 2 & 8 & 0 & 0 & -2 \\ 6 & 0 & 0 & 6 \\ 3 & 3 & 3 & -3 \end{pmatrix}^{1/2}$$

$10 \rightarrow 8 \otimes 8$

$$\begin{pmatrix} \Delta \\ \Sigma \\ \Xi \\ \Omega \end{pmatrix} \rightarrow \begin{pmatrix} N\pi & N\eta \\ N\bar{K} & \Sigma\pi & \Lambda\pi & \Sigma\eta & \Xi K \\ \Sigma\bar{K} & \Lambda\bar{K} & \Xi\pi & \Xi\eta \\ \Xi\bar{K} \end{pmatrix} = \frac{1}{\sqrt{12}} \begin{pmatrix} -6 & 6 \\ -2 & 2 & -3 & 3 & 2 \\ 3 & -3 & 3 & 3 \\ 12 \end{pmatrix}^{1/2}$$

$8 \rightarrow 10 \otimes 8$

$$\begin{pmatrix} N \\ \Sigma \\ \Lambda \\ \Xi \end{pmatrix} \rightarrow \begin{pmatrix} \Delta\pi & \Sigma K \\ \Delta\bar{K} & \Sigma\pi & \Sigma\eta & \Xi K \\ \Sigma\pi & \Xi K \\ \Sigma\bar{K} & \Xi\pi & \Xi\eta & \Omega K \end{pmatrix} = \frac{1}{\sqrt{15}} \begin{pmatrix} -12 & 3 \\ 8 & -2 & -3 & 2 \\ -9 & 6 \\ 3 & -3 & -3 & 6 \end{pmatrix}^{1/2}$$

$10 \rightarrow 10 \otimes 8$

$$\begin{pmatrix} \Delta \\ \Sigma \\ \Xi \\ \Omega \end{pmatrix} \rightarrow \begin{pmatrix} \Delta\pi & \Delta\eta & \Sigma K \\ \Delta\bar{K} & \Sigma\pi & \Sigma\eta & \Xi K \\ \Sigma\bar{K} & \Xi\pi & \Xi\eta & \Omega K \\ \Xi\bar{K} & \Omega\eta \end{pmatrix} = \frac{1}{\sqrt{24}} \begin{pmatrix} 15 & 3 & -6 \\ 8 & 8 & 0 & -8 \\ 12 & 3 & -3 & -6 \\ 12 & -12 \end{pmatrix}^{1/2}$$

abc	f_{abc}	abc	d_{abc}	abc	d_{abc}
123	1	118	$1/\sqrt{3}$	355	1/2
147	1/2	146	1/2	366	-1/2
156	-1/2	157	1/2	377	-1/2
246	1/2	228	$1/\sqrt{3}$	448	$-1/(2\sqrt{3})$
257	1/2	247	-1/2	558	$-1/(2\sqrt{3})$
345	1/2	256	1/2	668	$-1/(2\sqrt{3})$
367	-1/2	338	$1/\sqrt{3}$	778	$-1/(2\sqrt{3})$
458	$\sqrt{3}/2$	344	1/2	888	$-1/\sqrt{3}$
678	$\sqrt{3}/2$				

In the fundamental 3-dimensional representation, the λ_a 's are

$$\lambda_1 = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \lambda_2 = \begin{pmatrix} 0 & -i & 0 \\ i & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \quad \lambda_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 0 \end{pmatrix}$$

$$\lambda_4 = \begin{pmatrix} 0 & 0 & 1 \\ 0 & 0 & 0 \\ 1 & 0 & 0 \end{pmatrix} \quad \lambda_5 = \begin{pmatrix} 0 & 0 & -i \\ 0 & 0 & 0 \\ i & 0 & 0 \end{pmatrix} \quad \lambda_6 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{pmatrix}$$

$$\lambda_7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & -i \\ 0 & i & 0 \end{pmatrix} \quad \lambda_8 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -2 \end{pmatrix}$$


22. SU(N) MULTIPLETS AND YOUNG DIAGRAMMS

This note tells (1) how SU(n) particle multiplets are identified or labeled, (2) how to find the number of particles in a multiplet from its label, (3) how to draw the Young diagram for a multiplet, and (4) how to use Young diagrams to determine the overall multiplet structure of a composite system, such as a 3-quark or a meson-baryon system.

In much of the literature, the word “representation” is used where we use “multiplet,” and “tableau” is used where we use “diagram.”

22.1. Multiplet labels

An SU(n) multiplet is uniquely identified by a string of (n-1) nonnegative integers: (α, β, γ, ...). Any such set of integers specifies a multiplet. For an SU(2) multiplet such as an isospin multiplet, the single integer α is the number of steps from one end of the multiplet to the other (i.e., it is one fewer than the number of particles in the multiplet). In SU(3), the two integers α and β are the numbers of steps across the top and bottom levels of the multiplet diagram. Thus the labels for the SU(3) octet and decuplet



are (1,1) and (3,0). For larger n, the interpretation of the integers in terms of the geometry of the multiplets, which exist in an (n-1)-dimensional space, is not so readily apparent.

The label for the SU(n) singlet is (0, 0, ..., 0). In a flavor SU(n), the n quarks together form a (1, 0, ..., 0) multiplet, and the n antiquarks belong to a (0, ..., 0, 1) multiplet. These two multiplets are conjugate to one another, which means their labels are related by (α, β, ...) ↔ (... , β, α).

22.2. Number of particles

The number of particles in a multiplet, N = N(α, β, ...), is given as follows (note the pattern of the equations).

In SU(2), N = N(α) is

$$N = \frac{(\alpha + 1)}{1} \quad (22.1)$$

In SU(3), N = N(α, β) is

$$N = \frac{(\alpha + 1)}{1} \cdot \frac{(\beta + 1)}{1} \cdot \frac{(\alpha + \beta + 2)}{2} \quad (22.2)$$

In SU(4), N = N(α, β, γ) is

$$N = \frac{(\alpha + 1)}{1} \cdot \frac{(\beta + 1)}{1} \cdot \frac{(\gamma + 1)}{1} \cdot \frac{(\alpha + \beta + 2)}{2} \cdot \frac{(\beta + \gamma + 2)}{2} \cdot \frac{(\alpha + \beta + \gamma + 3)}{3} \quad (22.3)$$


Note that there is no factor with (α + γ + 2): only a consecutive sequence of the label integers appears in any factor. One more example should make the pattern clear for any SU(n). In SU(5), N = N(α, β, γ, δ) is

$$N = \frac{(\alpha + 1)}{1} \cdot \frac{(\beta + 1)}{1} \cdot \frac{(\gamma + 1)}{1} \cdot \frac{(\delta + 1)}{1} \cdot \frac{(\alpha + \beta + 2)}{2} \cdot \frac{(\beta + \gamma + 2)}{2} \cdot \frac{(\gamma + \delta + 2)}{2} \cdot \frac{(\alpha + \beta + \gamma + 3)}{3} \cdot \frac{(\beta + \gamma + \delta + 3)}{3} \cdot \frac{(\alpha + \beta + \gamma + \delta + 4)}{4} \quad (22.4)$$

From the symmetry of these equations, it is clear that multiplets that are conjugate to one another have the same number of particles, but so can other multiplets. For example, the SU(4) multiplets (3,0,0) and (1,1,0) each have 20 particles. Try the equations and see.

22.3. Young diagrams

A Young diagram consists of an array of boxes (or some other symbol) arranged in one or more left-justified rows, with each row being at least as long as the row beneath. The correspondence between a diagram and a multiplet label is: The top row juts out α boxes to the right past the end of the second row, the second row juts out β boxes to the right past the end of the third row, etc. A diagram in SU(n) has at most n rows. There can be any number of “completed” columns of n boxes buttressing the left of a diagram; these don’t affect the label. Thus in SU(3) the diagrams



represent the multiplets (1,0), (0,1), (0,0), (1,1), and (3,0). In any SU(n), the quark multiplet is represented by a single box, the antiquark multiplet by a column of (n-1) boxes, and a singlet by a completed column of n boxes.


22.4. Coupling multiplets together

The following recipe tells how to find the multiplets that occur in coupling two multiplets together. To couple together more than two multiplets, first couple two, then couple a third with each of the multiplets obtained from the first two, etc.

First a definition: A sequence of the letters a, b, c, ... is admissible if at any point in the sequence at least as many a’s have occurred as b’s, at least as many b’s have occurred as c’s, etc. Thus abcd and aabc are admissible sequences and abb and acb are not. Now the recipe:

(a) Draw the Young diagrams for the two multiplets, but in one of the diagrams replace the boxes in the first row with a’s, the boxes in the second row with b’s, etc. Thus, to find the multiplets that occur in the coupling of two SU(3) octets (one might be the π-meson octet, the other the baryon octet), we draw and . The unlettered diagram forms the upper left-hand corner of all the enlarged diagrams constructed below.


(b) Add the a’s from the lettered diagram to the right-hand ends of the rows of the unlettered diagram to form all possible legitimate Young diagrams that have no more than one a per column. In general, there will be several distinct diagrams, and all the a’s appear in each diagram. At this stage, the calculation of the coupling of the two SU(3) octets look as follows:



(c) Use the b’s to further enlarge the diagrams already obtained, subject to the same rules. Then throw away any diagram in which the sequence of letters formed by reading right to left in the first row, then the second row, etc., is not admissible.

(d) Proceed as in (c) with the c’s (if any), etc.

The final result of the coupling of the two octets is:



Here only the diagrams with admissible sequences of a’s and b’s and with fewer than four rows (since n = 3) have been kept. In terms of multiplet labels, the above may be written

$$(1, 1) \otimes (1, 1) = (2, 2) \oplus (3, 0) \oplus (0, 3) \oplus (1, 1) \oplus (1, 1) \oplus (0, 0) .$$

In terms of numbers of particles, it may be written

$$8 \otimes 8 = 27 \oplus 10 \oplus \bar{10} \oplus 8 \oplus 8 \oplus 1 .$$

The product of the numbers on the left here is equal to the sum on the right. (See also the section on the Quark Model.)

23. KINEMATICS

Revised June 1994.

Throughout this section units are used in which $\hbar = c = 1$. The following conversions are useful: $\hbar c = 197.3$ MeV fermi, $(\hbar c)^2 = 0.3894$ (GeV)² mb.

23.1. Lorentz transformations

The energy E and 3-momentum \mathbf{p} of a particle of mass m form a 4-vector $p = (E, \mathbf{p})$ whose square $p^2 \equiv E^2 - |\mathbf{p}|^2 = m^2$. The velocity of the particle is $\boldsymbol{\beta} = \mathbf{p}/E$. The energy and momentum (E^*, \mathbf{p}^*) viewed from a frame moving with velocity $\boldsymbol{\beta}_f$ are given by

$$\begin{pmatrix} E^* \\ p_{\parallel}^* \end{pmatrix} = \begin{pmatrix} \gamma_f & -\gamma_f \beta_f \\ -\gamma_f \beta_f & \gamma_f \end{pmatrix} \begin{pmatrix} E \\ p_{\parallel} \end{pmatrix}, \quad p_T^* = p_T, \quad (23.1)$$

where $\gamma_f = (1 - \beta_f^2)^{-1/2}$ and p_T (p_{\parallel}) are the components of \mathbf{p} perpendicular (parallel) to $\boldsymbol{\beta}_f$. The scalar product of two 4-vectors $\mathbf{p}_1 \cdot \mathbf{p}_2 = E_1 E_2 - \mathbf{p}_1 \cdot \mathbf{p}_2$ is invariant (frame independent).

In the collision of two particles of masses m_1 and m_2 the total center-of-mass energy can be expressed in the Lorentz-invariant form

$$\begin{aligned} E_{\text{cm}} &= \left[(E_1 + E_2)^2 - (\mathbf{p}_1 + \mathbf{p}_2)^2 \right]^{1/2}, \\ &= \left[m_1^2 + m_2^2 + 2E_1 E_2 (1 - \beta_1 \beta_2 \cos \theta) \right]^{1/2}, \end{aligned} \quad (23.2)$$

where θ is the angle between the particles. In the frame where one particle (of mass m_2) is at rest (lab frame),

$$E_{\text{cm}} = (m_1^2 + m_2^2 + 2E_{1\text{lab}} m_2)^{1/2}. \quad (23.3)$$

The velocity in the lab of the center-of-mass frame is

$$\boldsymbol{\beta}_{\text{cm}} = \mathbf{p}_{\text{lab}} / (E_{1\text{lab}} + m_2), \quad (23.4)$$

and

$$\gamma_{\text{cm}} = (E_{1\text{lab}} + m_2) / E_{\text{cm}}. \quad (23.5)$$

23.2. Center of mass energy and momentum

A beam of particles with mass m and momentum \mathbf{p}_{beam} is incident on a fixed target consisting of particles with mass M . The energy of the beam particles E_{beam} , the total center-of-mass energy E_{cm} , and center of mass momentum of one of the particles \mathbf{p}_{cm} are given by

$$E_{\text{beam}} = \sqrt{p_{\text{beam}}^2 + m^2} \quad (23.6)$$

$$E_{\text{cm}} = \sqrt{m^2 + 2E_{\text{beam}} M + M^2} \quad (23.7)$$

$$\mathbf{p}_{\text{cm}} = \mathbf{p}_{\text{beam}} \frac{M}{E_{\text{cm}}}. \quad (23.8)$$

For example, if a 0.80 GeV/c kaon beam is incident on a proton target, the center of mass energy is 1.699 GeV and the center of mass momentum of either particle is 0.442 GeV/c. It is also useful to note that

$$E_{\text{cm}} dE_{\text{cm}} = M dE_{\text{beam}} = M \beta_{\text{beam}} dp_{\text{beam}}. \quad (23.9)$$

23.3. Lorentz-invariant amplitudes

The invariant amplitude $-i\mathcal{M}$ for a scattering or decay process is determined in perturbation theory by a set of Feynman diagrams. The convention of Bjorken and Drell is used except that fermion spinors are normalized so that $u\bar{u} = 2m$. As an example, the S -matrix for $2 \rightarrow 2$ scattering is related to \mathcal{M} by

$$\begin{aligned} \langle p'_1 p'_2 | S | p_1 p_2 \rangle &= I - i(2\pi)^4 \delta^4(p_1 + p_2 - p'_1 - p'_2) \\ &\times \frac{\mathcal{M}(p_1, p_2; p'_1, p'_2)}{(2E_1)^{1/2} (2E_2)^{1/2} (2E'_1)^{1/2} (2E'_2)^{1/2}}. \end{aligned} \quad (23.10)$$

The state normalization is such that

$$\langle p' | p \rangle = (2\pi)^3 \delta^3(\mathbf{p} - \mathbf{p}'). \quad (23.11)$$

23.4. Particle decays

The partial decay rate of a particle of mass M into n bodies in its rest frame is given in terms of the Lorentz-invariant matrix element \mathcal{M} by

$$d\Gamma = \frac{(2\pi)^4}{2M} |\mathcal{M}|^2 d\Phi_n(P; p_1, \dots, p_n), \quad (23.12)$$

where $d\Phi_n$ is an element of n -body phase space given by

$$d\Phi_n(P; p_1, \dots, p_n) = \delta^4(P - \sum_{i=1}^n p_i) \prod_{i=1}^n \frac{d^3 p_i}{(2\pi)^3 2E_i}. \quad (23.13)$$

This phase space can be generated recursively, viz.

$$\begin{aligned} d\Phi_n(P; p_1, \dots, p_n) &= d\Phi_j(q; p_1, \dots, p_j) \\ &\times d\Phi_{n-j+1}(P; q, p_{j+1}, \dots, p_n) (2\pi)^3 dq^2, \end{aligned} \quad (23.14)$$

where $q^2 = (\sum_{i=1}^j E_i)^2 - |\sum_{i=1}^j \mathbf{p}_i|^2$. This form is particularly useful in the case where a particle decays into another particle which subsequently decays.

23.4.1. Survival probability: If a particle of mass M has mean proper lifetime τ ($= 1/\Gamma$) and has momentum (E, \mathbf{p}) , then the probability that it lives for a time t_0 or greater before decaying is given by

$$P(t_0) = e^{-t_0 \Gamma / \gamma} = e^{-M t_0 \Gamma / E}, \quad (23.15)$$

and the probability that it travels a distance x_0 or greater is

$$P(x_0) = e^{-M x_0 \Gamma / |\mathbf{p}|}. \quad (23.16)$$

23.4.2. Two-body decays:




Figure 23.1: Variable definitions for two-body decays.

In the rest frame of a particle of mass M , decaying into 2 particles labeled 1 and 2,

$$E_1 = \frac{M^2 - m_2^2 + m_1^2}{2M}, \quad (23.17)$$

$$\begin{aligned} |\mathbf{p}_1| &= |\mathbf{p}_2| \\ &= \frac{[(M^2 - (m_1 + m_2)^2)(M^2 - (m_1 - m_2)^2)]^{1/2}}{2M}, \end{aligned} \quad (23.18)$$

and

$$d\Gamma = \frac{1}{32\pi^2} |\mathcal{M}|^2 \frac{|\mathbf{p}_1|}{M^2} d\Omega, \quad (23.19)$$

where $d\Omega = d\phi_1 d(\cos \theta_1)$ is the solid angle of particle 1.




Figure 23.2: Variable definitions for three-body decays.

23.4.3. Three-body decays:

Defining $p_{ij} = p_i + p_j$, $m_{ij}^2 = p_{ij}^2$, then $m_{12}^2 + m_{23}^2 + m_{13}^2 = M^2 + m_1^2 + m_2^2 + m_3^2$ and $m_{12}^2 = (P - p_3)^2 = M^2 + m_3^2 - 2ME_3$. The relative orientation of the three final-state particles is fixed if their energies are known. Their momenta can therefore be specified by giving three Euler angles (α, β, γ) which specify the orientation of the final system relative to the initial particle. Then

$$d\Gamma = \frac{1}{(2\pi)^5} \frac{1}{16M} |\mathcal{M}|^2 dE_1 dE_2 d\alpha d\cos\beta d\gamma. \quad (23.20)$$

Alternatively

$$d\Gamma = \frac{1}{(2\pi)^5} \frac{1}{16M^2} |\mathcal{M}|^2 |\mathbf{p}_1^*| |\mathbf{p}_3| dm_{12} d\Omega_1^* d\Omega_3, \quad (23.21)$$

where $(|\mathbf{p}_1^*|, \Omega_1^*)$ is the momentum of particle 1 in the rest frame of 1 and 2, and Ω_3 is the angle of particle 3 in the rest frame of the decaying particle. $|\mathbf{p}_1^*|$ and $|\mathbf{p}_3|$ are given by

$$|\mathbf{p}_1^*| = \frac{[(m_{12}^2 - (m_1 + m_2)^2)(m_{12}^2 - (m_1 - m_2)^2)]^{1/2}}{2m_{12}}, \quad (23.22a)$$

and

$$|\mathbf{p}_3| = \frac{[(M^2 - (m_{12} + m_3)^2)(M^2 - (m_{12} - m_3)^2)]^{1/2}}{2M}. \quad (23.22b)$$

[Compare with Eq. (23.18).]

Integrating over the angles in Eq. (23.20) (this is only possible if the decaying particle is a scalar or we average over its spin states; otherwise \mathcal{M} depends on α, β , and γ) gives

$$\begin{aligned} d\Gamma &= \frac{1}{(2\pi)^3} \frac{1}{8M} |\mathcal{M}|^2 dE_1 dE_2 \\ &= \frac{1}{(2\pi)^3} \frac{1}{32M^3} |\mathcal{M}|^2 dm_{12}^2 dm_{23}^2. \end{aligned} \quad (23.23)$$

This is the standard form for the Dalitz plot.

23.4.3.1. Dalitz plot: If m_{12}^2 is fixed then the range of m_{13}^2 is determined by its values when \mathbf{p}_1 is parallel or antiparallel to \mathbf{p}_3 .

$$(m_{13}^2)_{\max} = (E_1^* + E_3^*)^2 - \left(\sqrt{E_1^{*2} - m_1^2} - \sqrt{E_3^{*2} - m_3^2} \right)^2, \quad (23.24a)$$

$$(m_{13}^2)_{\min} = (E_1^* + E_3^*)^2 - \left(\sqrt{E_1^{*2} - m_1^2} + \sqrt{E_3^{*2} - m_3^2} \right)^2, \quad (23.24b)$$

where $E_3^* = (M^2 - m_{12}^2 - m_3^2)/(2m_{12})$ and $E_1^* = (m_{12}^2 + m_1^2 - m_2^2)/(2m_{12})$. The scatter plot in m_{12}^2 and m_{13}^2 has uniform phase space density [see Eq. (23.23)] and is called a Dalitz plot.

A nonuniformity in the plot gives immediate information on $|\mathcal{M}|^2$. For example, in the case of $D \rightarrow K\pi\pi$, bands appear when $m(K\pi) = m_{K^*(892)}$, reflecting the appearance of the decay chain $D \rightarrow K^*(892)\pi \rightarrow K\pi\pi$.




Figure 23.3: Dalitz plot for a three-body final state. In this example, the state is $\bar{K}^0\pi^+p$ at 3 GeV. Four-momentum conservation restricts events to the interior of the closed curve.

23.4.4. Kinematic limits: In a three-body decay the maximum of $|\mathbf{p}_3|$, [given by Eq. (23.22)], is achieved when $m_{12} = m_1 + m_2$, i.e., particles 1 and 2 have the same vector velocity in the rest frame of the decaying particle. If, in addition, $m_3 > m_1, m_2$, then $|\mathbf{p}_3|_{\max} > |\mathbf{p}_1|_{\max}, |\mathbf{p}_2|_{\max}$.

23.4.5. Multibody decays: The above results may be generalized to final states containing any number of particles by combining some of the particles into “effective particles” and treating the final states as 2 or 3 “effective particle” states. Thus, if $p_{ijk\dots} = p_i + p_j + p_k + \dots$, then

$$m_{ijk\dots} = \sqrt{p_{ijk\dots}^2}, \quad (23.25)$$

and $m_{ijk\dots}$ may be used in place of e.g., m_{12} in the relations in Sec. 23.4.3 or 23.4.3.1 above.




Figure 23.4: Variable definitions for production of an n -body final state.

23.5. Cross sections

The differential cross section is given by

$$\begin{aligned} d\sigma &= \frac{(2\pi)^4 |\mathcal{M}|^2}{4\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2}} \\ &\times d\Phi_n(p_1 + p_2; p_3, \dots, p_{n+2}). \end{aligned} \quad (23.26)$$

[See Eq. (23.13).] In the rest frame of m_2 (lab),

$$\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2} = m_2 p_{1\text{lab}}; \quad (23.27a)$$

while in the center-of-mass frame

$$\sqrt{(p_1 \cdot p_2)^2 - m_1^2 m_2^2} = p_{1\text{cm}} \sqrt{s}. \quad (23.27b)$$




Figure 23.5: Variable definitions for a two-body final state.

23.5.1. Two-body reactions:

Two particles of momenta p_1 and p_2 and masses m_1 and m_2 scatter to particles of momenta p_3 and p_4 and masses m_3 and m_4 ; the Lorentz-invariant Mandelstam variables are defined by

$$s = (p_1 + p_2)^2 = (p_3 + p_4)^2 = m_1^2 + 2E_1 E_2 - 2\mathbf{p}_1 \cdot \mathbf{p}_2 + m_2^2, \quad (23.28)$$

$$t = (p_1 - p_3)^2 = (p_2 - p_4)^2 = m_1^2 - 2E_1 E_3 + 2\mathbf{p}_1 \cdot \mathbf{p}_3 + m_3^2, \quad (23.29)$$

$$u = (p_1 - p_4)^2 = (p_2 - p_3)^2 = m_1^2 - 2E_1 E_4 + 2\mathbf{p}_1 \cdot \mathbf{p}_4 + m_4^2, \quad (23.30)$$

and they satisfy

$$s + t + u = m_1^2 + m_2^2 + m_3^2 + m_4^2. \quad (23.31)$$

The two-body cross section may be written as

$$\frac{d\sigma}{dt} = \frac{1}{64\pi s} \frac{1}{|\mathbf{p}_{1\text{cm}}|^2} |\mathcal{M}|^2. \quad (23.32)$$

In the center-of-mass frame

$$t = (E_{1\text{cm}} - E_{3\text{cm}})^2 - (\mathbf{p}_{1\text{cm}} - \mathbf{p}_{3\text{cm}})^2 - 4p_{1\text{cm}} p_{3\text{cm}} \sin^2(\theta_{\text{cm}}/2) = t_0 - 4p_{1\text{cm}} p_{3\text{cm}} \sin^2(\theta_{\text{cm}}/2), \quad (23.33)$$

where θ_{cm} is the angle between particle 1 and 3. The limiting values t_0 ($\theta_{\text{cm}} = 0$) and t_1 ($\theta_{\text{cm}} = \pi$) for $2 \rightarrow 2$ scattering are

$$t_0(t_1) = \left[\frac{m_1^2 - m_3^2 - m_2^2 + m_4^2}{2\sqrt{s}} \right]^2 - \left\{ \left[\left(\frac{s + m_1^2 - m_2^2}{2\sqrt{s}} \right)^2 - m_1^2 \right]^{1/2} \mp \left[\left(\frac{s + m_3^2 - m_4^2}{2\sqrt{s}} \right)^2 - m_3^2 \right]^{1/2} \right\}^2. \quad (23.34)$$

Note that t_1 is always negative. In the literature the notation t_{min} (t_{max}) for t_0 (t_1) is sometimes used. This usage should be discouraged since $t_0 > t_1$. The center-of-mass energies and momenta of the incoming particles are

$$E_{\text{cm}} = \frac{s + m_1^2 - m_2^2}{2\sqrt{s}}, \quad (23.35)$$

$$p_{\text{cm}} = \frac{[(s - (m_1 + m_2)^2)(s - (m_1 - m_2)^2)]^{1/2}}{2\sqrt{s}} = \frac{p_{1\text{lab}} m_2}{\sqrt{s}}, \quad (23.36)$$

Here the subscript lab refers to the frame where particle 2 is at rest. [For other relations see Eqs. (23.2)–(23.4).]

23.5.2. Inclusive reactions: Choose some direction (usually the beam direction) for the z -axis; then the energy and momentum of a particle can be written as

$$E = m_T \cosh y, \quad p_x, p_y, p_z = m_T \sinh y, \quad (23.37)$$

where m_T is the transverse mass

$$m_T^2 = m^2 + p_x^2 + p_y^2, \quad (23.38)$$

and the rapidity y is defined by

$$y = \frac{1}{2} \ln \left(\frac{E + p_z}{E - p_z} \right) = \ln \left(\frac{E + p_z}{m_T} \right) = \tanh^{-1} \left(\frac{p_z}{E} \right). \quad (23.39)$$

Under a boost in the z -direction to a frame with velocity β , $y \rightarrow y + \tanh^{-1} \beta$. Hence the shape of the rapidity distribution dN/dy is invariant. The invariant cross section may also be rewritten

$$E \frac{d^3\sigma}{d^3p} = \frac{d^3\sigma}{d\phi dy p_T dp_T} \Rightarrow \frac{d^2\sigma}{\pi dy d(p_T^2)}. \quad (23.40)$$

The second form is obtained using the identity $dy/dp_z = 1/E$, and the third form represents the average over ϕ .

Feynman's x variable is given by

$$x = \frac{p_z}{p_{z\text{max}}} \approx \frac{E + p_z}{(E + p_z)_{\text{max}}}; \quad (23.41)$$

in the center-of-mass frame,

$$x \approx \frac{2p_{z\text{cm}}}{\sqrt{s}} \approx \frac{2m_T \sinh y_{\text{cm}}}{\sqrt{s}}. \quad (23.42)$$

For y_{cm} such that $e^{-2y_{\text{cm}}} \ll 1$,

$$x \approx \frac{m_T}{\sqrt{s}} e^{y_{\text{cm}}} \quad (23.43)$$

and

$$(y_{\text{cm}})_{\text{max}} = \ln(\sqrt{s}/m). \quad (23.44)$$

The definition of rapidity [Eq. (23.39)] may be expanded to obtain

$$y = \frac{1}{2} \ln \frac{\cos^2(\theta/2) + m^2/4p^2 + \dots}{\sin^2(\theta/2) + m^2/4p^2 + \dots} \approx -\ln \tan(\theta/2) \equiv \eta \quad (23.45)$$

if the particle has zenith angle θ . The pseudorapidity η defined by the second line is approximately equal to the rapidity y for $p \gg m$ and $\theta \gg 1/\gamma$, and in any case can be measured when the mass and momentum of the particle is unknown. From the definition one can obtain the identities

$$\sinh \eta = \cot \theta \quad (23.46)$$

$$\cosh \eta = 1/\sin \theta \quad (23.47)$$

$$\tanh \eta = \cos \theta. \quad (23.48)$$

23.5.3. Partial waves: The amplitude in the center of mass for elastic scattering of spinless particles may be expanded in Legendre polynomials

$$f(k, \theta) = \frac{1}{k} \sum_{\ell} (2\ell + 1) a_{\ell} P_{\ell}(\cos \theta), \quad (23.49)$$

where k is the c.m. momentum, θ is the c.m. scattering angle, $a_{\ell} = (\eta_{\ell} e^{2i\delta_{\ell}} - 1)/2i$, $0 \leq \eta_{\ell} \leq 1$, and δ_{ℓ} is the phase shift of the ℓ^{th} partial wave. For purely elastic scattering, $\eta_{\ell} = 1$. The differential cross section is

$$\frac{d\sigma}{d\Omega} = |f(k, \theta)|^2. \quad (23.50)$$

The optical theorem states that

$$\sigma_{\text{tot}} = \frac{4\pi}{k} \text{Im} f(k, 0), \quad (23.51)$$

and the cross section in the ℓ^{th} partial wave is therefore bounded:

$$\sigma_{\ell} = \frac{4\pi}{k^2} (2\ell + 1) |a_{\ell}|^2 \leq \frac{4\pi(2\ell + 1)}{k^2}. \quad (23.52)$$

The evolution with energy of partial-wave amplitude a_{ℓ} can be displayed as a trajectory in an Argand plot, as shown in Fig. 23.6.




Figure 23.6: Argand plot for the display of a partial-wave amplitude as a function of energy.

The usual Lorentz-invariant matrix element \mathcal{M} (see Sec. 23.3 above) for the elastic process is related to $f(k, \theta)$ by

$$\mathcal{M} = -8\pi\sqrt{s} f(k, \theta), \quad (23.53)$$

so

$$\sigma_{\text{tot}} = -\frac{1}{2k\sqrt{s}} \text{Im} \mathcal{M}(t=0), \quad (23.54)$$

where s and t are the center-of-mass energy squared and momentum transfer squared, respectively (see Sec. 23.4.1).

23.5.3.1. Resonances: The Breit-Wigner form for an elastic amplitude a_{ℓ} with a resonance at c.m. energy E_R , elastic width Γ_{el} , and total width Γ_{tot} is

$$a_{\ell} = \frac{\Gamma_{\text{el}}/2}{E_R - E - i\Gamma_{\text{tot}}/2}, \quad (23.55)$$

where E is the c.m. energy. As shown in Fig. 23.7, in the absence of background the elastic amplitude traces a counterclockwise circle with center $ix_{\text{el}}/2$ and radius $x_{\text{el}}/2$, where the elasticity $x_{\text{el}} = \Gamma_{\text{el}}/\Gamma_{\text{tot}}$. The amplitude has a pole at $E = E_R - i\Gamma_{\text{tot}}/2$.

The Breit-Wigner cross section for a spin- J resonance produced in the collision of particles of spin S_1 and S_2 is

$$\sigma_{BW}(E) = \frac{(2J+1)}{(2S_1+1)(2S_2+1)} \frac{\pi}{k^2} \frac{B_{\text{in}} B_{\text{out}} \Gamma_{\text{tot}}^2}{(E - E_R)^2 + \Gamma_{\text{tot}}^2/4}, \quad (23.56)$$

where k is the c.m. momentum, E is the c.m. energy, and B_{in} and B_{out} are the branching fractions of the resonance into the entrance and exit channels. The $2S+1$ factors are the multiplicities of the incident spin states, so they are replaced by 2 for photons, etc. This expression is valid only for a particle of narrow width. If the width




Figure 23.7: Argand plot for a resonance.

is not small, Γ_{tot} cannot be treated as a constant independent of E . There are many other forms for σ_{BW} , all of which are equivalent to the one given here in the narrow-width case. Some of these forms may be more appropriate if the resonance is broad.

The relativistic Breit-Wigner form corresponding to Eq. (23.55) is:

$$a_{\ell} = \frac{-m\Gamma_{\text{el}}}{s - m^2 + im\Gamma_{\text{tot}}}. \quad (23.57)$$

A better form incorporates the known kinematic dependences, replacing $m\Gamma_{\text{tot}}$ by $\sqrt{s}\Gamma_{\text{tot}}(s)$, where $\Gamma_{\text{tot}}(s)$ is the width the resonance particle would have if its mass were \sqrt{s} , and correspondingly $m\Gamma_{\text{el}}$ by $\sqrt{s}\Gamma_{\text{el}}(s)$ where $\Gamma_{\text{el}}(s)$ is the partial width in the incident channel for a mass \sqrt{s} :

$$a_{\ell} = \frac{-\sqrt{s}\Gamma_{\text{el}}(s)}{s - m^2 + i\sqrt{s}\Gamma_{\text{tot}}(s)}. \quad (23.58)$$

For the Z boson, all the decays are to particles whose masses are small enough to be ignored, so on dimensional grounds $\Gamma_{\text{tot}}(s) = \sqrt{s}\Gamma_0/m_Z$, where Γ_0 defines the width of the Z , and $\Gamma_{\text{el}}(s)/\Gamma_{\text{tot}}(s)$ is constant. A full treatment of the line shape requires consideration of dynamics, not just kinematics. For the Z this is done by calculating the radiative corrections in the Standard Model.

24. CROSS-SECTION FORMULAE FOR SPECIFIC PROCESSES

24.1. Leptoproduction




Figure 24.1: Kinematic quantities for description of lepton-nucleon scattering. k and k' are the four-momenta of incoming and outgoing leptons. P is the four-momentum of a nucleon with mass M . The exchanged particle is a γ , W^\pm , or Z^0 ; it transfers four-momentum $q = k - k'$ to the target.

Invariant quantities:

$\nu = \frac{q \cdot P}{M} = E - E'$ is the lepton's energy loss in the lab (in earlier literature sometimes $\nu = q \cdot P$). Here, E and E' are the initial and final lepton energies in the lab.

$Q^2 = -q^2 = 2(E E' - \vec{k} \cdot \vec{k}') - m_\ell^2 - m_{\ell'}^2$, where $m_\ell(m_{\ell'})$ is the initial (final) lepton mass. If $E E' \sin^2(\theta/2) \gg m_\ell^2, m_{\ell'}^2$, then $\approx 4E E' \sin^2(\theta/2)$, where θ is the lepton's scattering angle in the lab.

$x = \frac{Q^2}{2M\nu}$ In the parton model, x is the fraction of the target nucleon's momentum carried by the struck quark. See section on QCD.

$y = \frac{q \cdot P}{k \cdot P} = \frac{\nu}{E}$ is the fraction of the lepton's energy lost in the lab.

$W^2 = (P + q)^2 = M^2 + 2M\nu - Q^2$ is the mass squared of the system recoiling against the lepton.

$s = (k + P)^2 = \frac{Q^2}{xy} + M^2$

24.1.1. Leptoproduction cross sections:

$$\begin{aligned} \frac{d^2\sigma}{dx dy} &= \nu(s - M^2) \frac{d^2\sigma}{d\nu dQ^2} = \frac{2\pi M\nu}{E'} \frac{d^2\sigma}{d\Omega_{\text{lab}} dE'} \\ &= x(s - M^2) \frac{d^2\sigma}{dx dQ^2}. \end{aligned} \quad (24.1)$$

24.1.2. Electroproduction structure functions: The neutral-current process, $eN \rightarrow eX$, is parity conserving at low Q^2 and can be written in terms of two structure functions $F_1^{\text{NC}}(x, Q^2)$ and $F_2^{\text{NC}}(x, Q^2)$:

$$\begin{aligned} \frac{d^2\sigma}{dx dy} &= \frac{4\pi\alpha^2(s - M^2)}{Q^4} \\ &\times \left[(1 - y) F_2^{\text{NC}} + y^2 x F_1^{\text{NC}} - \frac{M^2}{(s - M^2)} xy F_2^{\text{NC}} \right]. \end{aligned} \quad (24.2)$$

The charged-current processes, $e^-N \rightarrow \nu X$, $\nu N \rightarrow e^-X$, and $\bar{\nu}N \rightarrow e^+X$, are parity violating and can be written in terms of three structure functions $F_1^{\text{CC}}(x, Q^2)$, $F_2^{\text{CC}}(x, Q^2)$, and $F_3^{\text{CC}}(x, Q^2)$:

$$\begin{aligned} \frac{d^2\sigma}{dx dy} &= \frac{G_F^2(s - M^2)}{2\pi} \frac{M_W^4}{(Q^2 + M_W^2)^2} \\ &\times \left\{ \left[1 - y - \frac{M^2 xy}{(s - M^2)} \right] F_2^{\text{CC}} + \frac{y^2}{2} 2x F_1^{\text{CC}} \pm \left(y - \frac{y^2}{2} \right) x F_3^{\text{CC}} \right\}, \end{aligned} \quad (24.3)$$

where the last term is positive for the e^- and ν reactions and negative for $\bar{\nu}N \rightarrow e^+X$.

24.1.3. The QCD parton model: In the QCD parton model, the structure functions defined above can be expressed in terms of parton distribution functions. The quantity $f_i(x, Q^2)dx$ is the probability that a parton of type i (quark, antiquark, or gluon), carries a momentum fraction between x and $x + dx$ of the nucleon's momentum in a frame where the nucleon's momentum is large. For the cross section corresponding to the *neutral-current process* $ep \rightarrow eX$, we have for $s \gg M^2$ (in the case where the incoming electron is either left- (L) or right- (R) handed):

$$\begin{aligned} \frac{d^2\sigma}{dx dy} &= \frac{\pi\alpha^2}{s^2 x^2 y^2} \left[\sum_q (x f_q(x, Q^2) + x f_{\bar{q}}(x, Q^2)) \right] \\ &\times [A_q + (1 - y)^2 B_q]. \end{aligned} \quad (24.4)$$

Here the index q refers to a quark flavor (*i.e.*, u, d, s, c, b , or t), and

$$A_q = \left(-q_q + g_{Lq} g_{Le} \frac{Q^2}{Q^2 + M_Z^2} \right)^2 + \left(-q_q + g_{Rq} g_{Re} \frac{Q^2}{Q^2 + M_Z^2} \right)^2, \quad (24.5)$$

$$B_q = \left(-q_q + g_{Rq} g_{Le} \frac{Q^2}{Q^2 + M_Z^2} \right)^2 + \left(-q_q + g_{Lq} g_{Re} \frac{Q^2}{Q^2 + M_Z^2} \right)^2. \quad (24.6)$$

Here q_q is the charge of flavor q . For a left-handed electron, $g_{Re} = 0$ and $g_{Le} = (-1/2 + \sin^2 \theta_W)/(\sin \theta_W \cos \theta_W)$, while for a right-handed one, $g_{Le} = 0$ and $g_{Re} = (\sin^2 \theta_W)/(\sin \theta_W \cos \theta_W)$. For the quarks, $g_{Lq} = (T_3 - q_q \sin^2 \theta_W)/(\sin \theta_W \cos \theta_W)$, and $g_{Rq} = (-q_q \sin^2 \theta_W)/(\sin \theta_W \cos \theta_W)$.

For neutral-current *neutrino (antineutrino) scattering*, the same formula applies with g_{Le} replaced by $g_{L\nu} = 1/(2 \sin \theta_W \cos \theta_W)$ ($g_{L\nu} = 0$) and g_{Re} replaced by $g_{R\nu} = 0$ [$g_{R\nu} = -1/(2 \sin \theta_W \cos \theta_W)$].

In the case of the *charged-current processes* $e_L^- p \rightarrow \nu X$ and $\bar{\nu} p \rightarrow e^+ X$, Eq. (24.3) applies with

$$\begin{aligned} F_2 &= 2x F_1 = 2x [f_u(x, Q^2) + f_c(x, Q^2) \\ &+ f_t(x, Q^2) + f_{\bar{d}}(x, Q^2) + f_{\bar{s}}(x, Q^2) + f_{\bar{b}}(x, Q^2)], \end{aligned} \quad (24.7)$$

$$\begin{aligned} F_3 &= 2 [f_u(x, Q^2) + f_c(x, Q^2) \\ &+ f_t(x, Q^2) - f_{\bar{d}}(x, Q^2) - f_{\bar{s}}(x, Q^2) - f_{\bar{b}}(x, Q^2)]. \end{aligned} \quad (24.8)$$

For the process $\nu p \rightarrow e^- X$:

$$\begin{aligned} F_2 &= 2x F_1 = 2x [f_d(x, Q^2) + f_s(x, Q^2) \\ &+ f_b(x, Q^2) + f_{\bar{u}}(x, Q^2) + f_{\bar{c}}(x, Q^2) + f_{\bar{t}}(x, Q^2)], \end{aligned} \quad (24.9)$$

$$\begin{aligned} F_3 &= 2 [f_d(x, Q^2) + f_s(x, Q^2) \\ &+ f_b(x, Q^2) - f_{\bar{u}}(x, Q^2) - f_{\bar{c}}(x, Q^2) - f_{\bar{t}}(x, Q^2)]. \end{aligned} \quad (24.10)$$

24.2. e^+e^- annihilation

For pointlike spin-1/2 fermions in the c.m., the differential cross section for $e^+e^- \rightarrow f\bar{f}$ via single photon annihilation is

$$\frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4s} \beta \left[1 + \cos^2 \theta + (1 - \beta^2) \sin^2 \theta \right] Q_f^2, \quad (24.11)$$

where β is the velocity of the final state fermion in the center of mass and Q_f is the charge of the fermion in units of the proton charge. For $\beta \rightarrow 1$,

$$\sigma = \frac{4\pi\alpha^2}{3s} Q_f^2 = \frac{86.8 Q_f^2 nb}{s(\text{GeV}^2)}. \quad (24.12)$$

At higher energies the Z^0 (mass M_Z and width Γ_Z) must be included, and the differential cross section for $e^+e^- \rightarrow f\bar{f}$ becomes

$$\begin{aligned} \frac{d\sigma}{d\Omega} = \frac{\alpha^2}{4s} \beta & \left[Q_f^2 [1 + \cos^2 \theta + (1 - \beta^2) \sin^2 \theta] \right. \\ & - 2Q_f \chi_1 \left\{ VV_f [1 + \cos^2 \theta + (1 - \beta^2) \sin^2 \theta] - 2a_f \beta \cos \theta \right\} \\ & + \chi_2 \left\{ V_f^2 (1 + V^2) [1 + \cos^2 \theta + (1 - \beta^2) \sin^2 \theta] \right. \\ & \left. \left. + \beta^2 a_f^2 (1 + V^2) [1 + \cos^2 \theta] - 8\beta VV_f a_f \cos \theta \right\} \right], \quad (24.13) \end{aligned}$$

$$\chi_1 = \frac{1}{16 \sin^2 \theta_W \cos^2 \theta_W} \frac{s(s - M_Z^2)}{(s - M_Z^2)^2 + \Gamma_Z^2 M_Z^2}, \quad (24.14)$$

$$\chi_2 = \frac{1}{256 \sin^4 \theta_W \cos^4 \theta_W} \frac{s^2}{(s - M_Z^2)^2 + \Gamma_Z^2 M_Z^2}, \quad (24.15)$$

$$V = -1 + 4 \sin^2 \theta_W, \quad (24.16)$$

$$a_f = 2T_{3f}, \quad (24.17)$$

$$V_f = 2T_{3f} - 4Q_f \sin^2 \theta_W, \quad (24.18)$$

where the subscript f refers to the particular fermion and

$$T_3 = +1/2 \quad \text{for } \nu_e, \nu_\mu, \nu_\tau, u, c, t, \quad (24.19a)$$

$$T_3 = -1/2 \quad \text{for } e^-, \mu^-, \tau^-, d, s, b. \quad (24.19b)$$

24.3. Two-photon process at e^+e^- colliders

When an e^+ and an e^- collide with energies E_1 and E_2 , they emit dn_1 and dn_2 virtual photons with energies ω_1 and ω_2 and 4-momenta q_1 and q_2 . In the equivalent photon approximation, the cross section for $e^+e^- \rightarrow e^+e^-X$ is related to the cross section for $\gamma\gamma \rightarrow X$ by (Ref. 1)

$$d\sigma_{e^+e^- \rightarrow e^+e^-X}(s) = dn_1 dn_2 d\sigma_{\gamma\gamma \rightarrow X}(W^2) \quad (24.20)$$

where $s = 4E_1E_2$, $W^2 = 4\omega_1\omega_2$ and

$$dn_i = \frac{\alpha}{\pi} \left[1 - \frac{\omega_i}{E_i} + \frac{\omega_i^2}{2E_i^2} - \frac{m_e^2 \omega_i^2}{(-q_i^2)E_i^2} \right] \frac{d\omega_i}{\omega_i} \frac{d(-q_i^2)}{(-q_i^2)}. \quad (24.21)$$

After integration (including that over q_i^2 in the region $m_e^2 \omega_i^2/E_i(E_i - \omega_i) \leq -q_i^2 \leq (-q^2)_{\max}$), the cross section is

$$\begin{aligned} \sigma_{e^+e^- \rightarrow e^+e^-X}(s) = \frac{\alpha^2}{\pi^2} \int_{z_{th}}^1 \frac{dz}{z} & \left[f(z) \left(\ln \frac{(-q^2)_{\max}}{m_e^2 z} - 1 \right)^2 \right. \\ & \left. - \frac{1}{3} \left(\ln \frac{1}{z} \right)^3 \right] \sigma_{\gamma\gamma \rightarrow X}(zs); \end{aligned}$$

$$f(z) = \left(1 + \frac{1}{2}z \right)^2 \ln \frac{1}{z} - \frac{1}{2}(1-z)(3+z);$$

$$z = \frac{W^2}{s}. \quad (24.22)$$

The quantity $(-q^2)_{\max}$ depends on properties of the produced system X , in particular, $(-q^2)_{\max} \sim m_\rho^2$ for hadron production ($X = h$) and $(-q^2)_{\max} \sim W^2$ for lepton pair production ($X = \ell^+\ell^-$, $\ell = e, \mu, \tau$).

For production of a resonance of mass m_R and spin $J \neq 1$

$$\begin{aligned} \sigma_{e^+e^- \rightarrow e^+e^-R}(s) = (2J+1) & \frac{8\alpha^2 \Gamma_{R \rightarrow \gamma\gamma}}{m_R^3} \\ & \times \left[f(m_R^2/s) \left(\ln \frac{sm_V^2}{m_e^2 m_R^2} - 1 \right)^2 - \frac{1}{3} \left(\ln \frac{s}{m_R^2} \right)^3 \right] \quad (24.23) \end{aligned}$$

where m_V is the mass that enters into the form factor of the $\gamma\gamma \rightarrow R$ transition: $m_V \sim m_\rho$ for $R = \pi^0, \rho^0, \omega, \phi, \dots$, $m_V \sim m_R$ for $R = c\bar{c}$ or $b\bar{b}$ resonances.

24.4. Inclusive hadronic reactions

One-particle inclusive cross sections $E d^3\sigma/d^3p$ for the production of a particle of momentum p are conveniently expressed in terms of rapidity (see above) and the momentum p_T transverse to the beam direction (defined in the center-of-mass frame)

$$E \frac{d^3\sigma}{d^3p} = \frac{d^3\sigma}{d\phi dy p_T dp_T}. \quad (24.24)$$

In the case of processes where p_T is large or the mass of the produced particle is large (here large means greater than 10 GeV), the parton model can be used to calculate the rate. Symbolically

$$\sigma_{\text{hadronic}} = \sum_{ij} \int f_i(x_1, Q^2) f_j(x_2, Q^2) dx_1 dx_2 \hat{\sigma}_{\text{partonic}}, \quad (24.25)$$

where $f_i(x, Q^2)$ is the parton distribution introduced above and Q is a typical momentum transfer in the partonic process and $\hat{\sigma}$ is the partonic cross section. Two examples will help to clarify. The production of a W^+ in pp reactions at rapidity y in the center-of-mass frame is given by

$$\begin{aligned} \frac{d\sigma}{dy} = \frac{G_F \pi \sqrt{2}}{3} & \times \tau \left[\cos^2 \theta_c \left(u(x_1, M_W^2) \bar{d}(x_2, M_W^2) \right. \right. \\ & \left. \left. + u(x_2, M_W^2) \bar{d}(x_1, M_W^2) \right) \right. \\ & \left. + \sin^2 \theta_c \left(u(x_1, M_W^2) \bar{s}(x_2, M_W^2) \right. \right. \\ & \left. \left. + s(x_2, M_W^2) \bar{u}(x_1, M_W^2) \right) \right], \quad (24.26) \end{aligned}$$

where $x_1 = \sqrt{\tau} e^y$, $x_2 = \sqrt{\tau} e^{-y}$, and $\tau = M_W^2/s$. Similarly the production of a jet in pp (or $p\bar{p}$) collisions is given by

$$\begin{aligned} \frac{d^3\sigma}{d^2p_T dy} = \sum_{ij} \int f_i(x_1, p_T^2) f_j(x_2, p_T^2) & \\ & \times \left[\hat{s} \frac{d\hat{\sigma}}{d\hat{t}} \right]_{ij} dx_1 dx_2 \delta(\hat{s} + \hat{t} + \hat{u}), \quad (24.27) \end{aligned}$$

where the summation is over quarks, gluons, and antiquarks. Here

$$s = (p_1 + p_2)^2, \quad (24.28)$$

$$t = (p_1 - p_{\text{jet}})^2, \quad (24.29)$$

$$u = (p_2 - p_{\text{jet}})^2, \quad (24.30)$$

p_1 and p_2 are the momenta of the incoming p and p (or \bar{p}) and \hat{s} , \hat{t} , and \hat{u} are s , t , and u with $p_1 \rightarrow x_1 p_1$ and $p_2 \rightarrow x_2 p_2$. The partonic cross section $\hat{s}[(d\hat{\sigma})/(d\hat{t})]$ can be found in Ref. 2. Example: for the process $gg \rightarrow q\bar{q}$,

$$\hat{s} \frac{d\sigma}{dt} = 3\alpha_s^2 \frac{(\hat{t}^2 + \hat{u}^2)}{8\hat{s}} \left[\frac{4}{9\hat{t}\hat{u}} - \frac{1}{\hat{s}^2} \right]. \quad (24.31)$$

The prediction of Eq. (24.27) is compared to data from the UA1 and UA2 collaborations in Fig. 32.7 in the Plots of Cross Sections and Related Quantities section of this *Review*.

24.5. One-particle inclusive distributions

In order to describe one-particle inclusive production in e^+e^- annihilation or deep inelastic scattering, it is convenient to introduce a fragmentation function $D_i^h(z, Q^2)/z$ which is the probability that a parton of type i and momentum p will fragment into a hadron of type h and momentum zp . The Q^2 evolution is predicted by QCD and is similar to that of the parton distribution functions [see section on Quantum Chromodynamics (Sec. 25 of this *Review*)]. The $D_i^h(z, Q^2)$ are normalized so that

$$\sum_h \int D_i^h(z, Q^2) dz = 1. \quad (24.32)$$

If the contributions of the Z boson and three-jet events are neglected, the cross section for producing a hadron h in e^+e^- annihilation is given by

$$\frac{1}{\sigma_{\text{had}}} \frac{d\sigma}{dz} = \frac{\sum_i e_i^2 D_i^h(z, Q^2)}{\sum_i e_i^2}, \quad (24.33)$$

where e_i is the charge of quark-type i , σ_{had} is the total hadronic cross section, and the momentum of the hadron is $zE_{\text{cm}}/2$.

In the case of deep inelastic muon scattering, the cross section for producing a hadron of energy E_h is given by

$$\frac{1}{\sigma_{\text{tot}}} \frac{d\sigma}{dz} = \frac{\sum_i e_i^2 q_i(x, Q^2) D_i^h(z, Q^2)}{\sum_i e_i^2 q_i(x, Q^2)}, \quad (24.34)$$

where $E_h = \nu z$. (For the kinematics of deep inelastic scattering, see Sec. 23.4.2 of the Kinematics section of this *Review*.) The fragmentation functions for light and heavy quarks have a different z dependence; the former peak near $z = 0$. They are illustrated in Fig. 32.12 in the section on Plots of Cross Sections and Related Quantities (Sec. 33 of this *Review*).

References:

1. V.M. Budnev, I.F. Ginzburg, G.V. Meledin, and V.G. Serbo, Phys. Reports **15C**, 181 (1975);
See also S. Brodsky, T. Kinoshita, and H. Terazawa, Phys. Rev. **D4**, 1532 (1971).
2. G.F. Owens, F. Reya, and M. Glück, Phys. Rev. **D18**, 1501 (1978).

25. QUANTUM CHROMODYNAMICS

25.1. The QCD Lagrangian

Prepared January 1994 by I. Hinchliffe.

Quantum Chromodynamics (QCD), the gauge field theory which describes the strong interactions of colored quarks and gluons, is one of the components of the $SU(3) \times SU(2) \times U(1)$ Standard Model. A quark of specific flavor (such as a charm quark) comes in 3 colors; gluons come in eight colors; hadrons are color-singlet combinations of quarks, anti-quarks, and gluons. The Lagrangian describing the interactions of quarks and gluons is (up to gauge-fixing terms)

$$L_{\text{QCD}} = -\frac{1}{4} F_{\mu\nu}^{(a)} F^{(a)\mu\nu} + i \sum_q \bar{\psi}_q^i \gamma^\mu (D_\mu)_{ij} \psi_q^j - \sum_q m_q \bar{\psi}_q^i \psi_{qi}, \quad (25.1)$$

$$F_{\mu\nu}^{(a)} = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g_s f_{abc} A_\mu^b A_\nu^c, \quad (25.2)$$

$$(D_\mu)_{ij} = \delta_{ij} \partial_\mu - ig_s \sum_a \frac{\lambda_{ij}^a}{2} A_\mu^a, \quad (25.3)$$

where g_s is the QCD coupling constant, and the f_{abc} are the structure constants of the $SU(3)$ algebra (the λ matrices and values for f_{abc} can be found in Sec. 21 of this *Review*). The $\psi_q^i(x)$ are the 4-component Dirac spinors associated with each quark field of (3) color i and flavor q and the $A_\mu^a(x)$ are the (8) Yang-Mills (gluon) fields. A complete list of the Feynman rules which derive from this Lagrangian, together with some useful color-algebra identities, can be found in Ref. 1.

The principle of ‘‘asymptotic freedom’’ (see below)

determines that the renormalized QCD coupling is small only at high energies, and it is only in this domain that high-precision tests—similar to those in QED—can be performed using perturbation theory. Nonetheless, there has been in recent years much progress in understanding and quantifying the predictions of QCD in the nonperturbative domain, for example in soft hadronic processes and on the lattice [2]. This short review will concentrate on QCD at short distances (large momentum transfers), where perturbation theory is the standard tool. It will discuss the processes that are used to determine the coupling constant of QCD. Other recent reviews of the coupling constant measurements may be consulted for a different perspective [3].

25.2. The QCD coupling and renormalization scheme

The renormalization scale dependence of the effective QCD coupling $\alpha_s = g_s^2/4\pi$ is controlled by the β -function:

$$\mu \frac{\partial \alpha_s}{\partial \mu} = -\frac{\beta_0}{2\pi} \alpha_s^2 - \frac{\beta_1}{4\pi^2} \alpha_s^3 - \frac{\beta_2}{64\pi^3} \alpha_s^4 - \dots, \quad (25.4a)$$

$$\beta_0 = 11 - \frac{2}{3} n_f, \quad (25.4b)$$

$$\beta_1 = 51 - \frac{19}{3} n_f, \quad (25.4c)$$

$$\beta_2 = 2857 - \frac{5033}{9} n_f - \frac{325}{27} n_f^2, \quad (25.4d)$$

here n_f is the number of quarks with mass less than the energy scale μ . In solving this differential equation for α_s , a constant of integration is introduced. This constant is the one fundamental constant of QCD that must be determined from experiment. The most sensible choice for this constant is the value of α_s at a fixed-reference scale μ_0 , but it is more conventional to introduce the dimensional parameter Λ since this provides a parametrization of the μ dependence of α_s . The definition of Λ is arbitrary. One way to define it (adopted here) is to write a solution of Eq. (25.4) as an expansion in inverse powers of $\ln(\mu^2)$:

$$\alpha_s(\mu) = \frac{4\pi}{\beta_0 \ln(\mu^2/\Lambda^2)}$$

$$\times \left[1 - \frac{2\beta_1}{\beta_0^2} \frac{\ln[\ln(\mu^2/\Lambda^2)]}{\ln(\mu^2/\Lambda^2)} + \frac{4\beta_1^2}{\beta_0^4 \ln^2(\mu^2/\Lambda^2)} \right. \\ \left. \times \left(\left(\ln[\ln(\mu^2/\Lambda^2)] - \frac{1}{2} \right)^2 + \frac{8\beta_2\beta_0}{\beta_1^2} - \frac{5}{4} \right) \right] \quad (25.5a)$$

The last term in this expansion is

$$\mathcal{O}\left(\frac{\ln^2[\ln(\mu^2/\Lambda^2)]}{\ln^3(\mu^2/\Lambda^2)}\right) \quad (25.5b)$$

and is usually neglected in the definition of Λ . We choose to include it even though its effect on $\alpha_s(\mu)$ is smaller than the experimental errors. For a fixed value of $\alpha_s(M_Z)$, the inclusion of this term shifts the value of Λ by ~ 15 MeV.

This solution illustrates the *asymptotic freedom* property: $\alpha_s \rightarrow 0$ as $\mu \rightarrow \infty$. Alternative definitions of Λ are possible. We adopt this as the standard. Values given by experiments using other definitions are adjusted as needed to meet our definition.

In the above discussion we have ignored quark-mass effects, *i.e.*, we have assumed an idealized situation where quarks of mass greater than μ are neglected completely. In this picture, the β -function coefficients change by discrete amounts as flavor thresholds are crossed when integrating the differential equation for α_s . It follows that, for a relationship such as Eq. (25.5) to remain valid for all values of μ , Λ must also change as flavor thresholds are crossed. This leads to the concept of a different Λ for each range of μ corresponding to an effective number of massless quarks: $\Lambda \rightarrow \Lambda^{(n_f)}$. There is some arbitrariness in how this relationship is set up. As an idealised case consider QCD with $n_f - 1$ massless quarks and one quark of mass M . Now imagine an experiment at energy scale μ ; for example, this could be $e^+e^- \rightarrow \text{hadrons}$ at center of mass energy μ . If $\mu \gg M$, the mass M is negligible and the process well described by QCD with n_f massless flavors and its parameter $\Lambda^{(n_f)}$ up to terms of order M^2/μ^2 . Conversely if $\mu \ll M$, the heavy quark plays no role and the process well described by QCD with $n_f - 1$ massless flavors and its parameter $\Lambda^{(n_f-1)}$ up to terms of order μ^2/M^2 . If $\mu \sim M$ the effects of the quark mass are process dependent and cannot be absorbed into the running coupling.

A mass scale μ' is chosen where the relationship between $\Lambda^{(n_f-1)}$ and $\Lambda^{(n_f)}$ will be fixed. μ' should be of order M and the relationship should not depend on it. A prescription has been given [4] which has this property. We use this procedure choosing $\mu' = M_Q$, where M_Q is the mass of the value of the running quark mass defined in the $\overline{\text{MS}}$ scheme (see the note on quark masses for more details) *i.e.* where $M_{\overline{\text{MS}}}(M_Q) = M_Q$. Then [4]

$$\beta_0^{n_f-1} \ln\left(\frac{\Lambda^{(n_f)}}{\Lambda^{(n_f-1)}}\right)^2 = (\beta_0^{n_f} - \beta_0^{n_f-1}) \ln\left(\frac{M_Q}{\Lambda^{(n_f)}}\right)^2 \\ + 2\left(\frac{\beta_1^{n_f}}{\beta_0^{n_f}} - \frac{\beta_1^{n_f-1}}{\beta_0^{n_f-1}}\right) \ln\left[\ln\left(\frac{M_Q}{\Lambda^{(n_f)}}\right)^2\right] \\ - \frac{2\beta_1^{n_f-1}}{\beta_0^{n_f-1}} \ln\left(\frac{\beta_0^{n_f}}{\beta_0^{n_f-1}}\right) \\ + \frac{4\frac{\beta_1^{n_f}}{\beta_0^{n_f}}\left(\frac{\beta_1^{n_f}}{\beta_0^{n_f}} - \frac{\beta_1^{n_f-1}}{\beta_0^{n_f-1}}\right)}{\ln\left(\frac{M_Q}{\Lambda^{(n_f)}}\right)^2} \ln\left[\ln\left(\frac{M_Q}{\Lambda^{(n_f)}}\right)^2\right] \\ + \frac{1}{\beta_0^{n_f}} \left[\left(\frac{2\beta_1^{n_f}}{\beta_0^{n_f}}\right)^2 - \left(\frac{2\beta_1^{n_f-1}}{\beta_0^{n_f-1}}\right)^2 - \frac{2\beta_2^{n_f}}{\beta_0^{n_f}} + \frac{2\beta_2^{n_f-1}}{\beta_0^{n_f-1}} - \frac{14}{9} \right] \\ \ln\left(\frac{M_Q}{\Lambda^{(n_f)}}\right)^2 \quad (25.6)$$

This result is valid to order α_s^3 (or alternatively to terms of order $1/\ln^2[(M_Q/\Lambda^{(n_f)})^2]$).

In the previous edition of this note an alternative matching was used [5]. This procedure required the equality $\alpha_s(\mu)^{(n_f)} = \alpha_s(\mu)^{(n_f-1)}$ for $\mu = M_Q$. This matching is somewhat arbitrary; a different relation between $\Lambda^{(n_f)}$ and $\Lambda^{(n_f-1)}$ would result if $\mu = M_Q/2$ were used. In practice the differences between the new and old procedures are very small. $\Lambda^{(5)} = 200$ MeV corresponds to $\Lambda^{(4)} = 289$ MeV in the old scheme and $\Lambda^{(4)} = 280$ MeV in the one now adopted. Note that these differences between $\Lambda^{(5)}$ and $\Lambda^{(4)}$ are numerically very significant.

Data from deep-inelastic scattering are in a range of energy where the bottom quark is not readily excited and hence these experiments quote $\Lambda_{\overline{MS}}^{(4)}$. Most data from PEP, PETRA, TRISTAN, and LEP quote a value of $\Lambda_{\overline{MS}}^{(5)}$ since these data are in an energy range where the bottom quark is light compared to the available energy. We have converted it to $\Lambda_{\overline{MS}}^{(4)}$ as required. A few measurements, including the lattice gauge theory values from the ψ system and from τ decay are at sufficiently low energy that $\Lambda_{\overline{MS}}^{(3)}$ is appropriate.

We turn now to a discussion of renormalization-scheme dependence in QCD. Although necessarily rather technical, this discussion is vital to understanding how α_s (or Λ) values can be measured and compared. See the review by Duke and Roberts [6] for further details.

Consider a “typical” QCD cross section which, when calculated perturbatively, starts at $\mathcal{O}(\alpha_s)$:

$$\sigma = A_1 \alpha_s + A_2 \alpha_s^2 + \dots \tag{25.7}$$

The coefficients A_1, A_2 come from calculating the appropriate Feynman diagrams. In performing such calculations various divergences arise, and these must be regulated in a consistent way. This requires a particular renormalization scheme (RS). The most commonly used one is the modified minimal subtraction (\overline{MS}) scheme [7]. This involves continuing momentum integrals from 4 to $4-2\epsilon$ dimensions and then subtracting off the resulting $1/\epsilon$ poles and also $(\ln 4\pi - \gamma_E)$, which is another artifact of continuing the dimension. (Here γ_E is the Euler-Mascheroni constant.) To preserve the dimensionless nature of the coupling, a mass scale μ must also be introduced: $g \rightarrow \mu^\epsilon g$. The finite coefficients A_i thus obtained depend implicitly on the renormalization convention used and explicitly on the scale μ .

The first two coefficients (β_0, β_1) in Eq. (25.4) are independent of the choice of RS’s. In contrast, the coefficients of terms proportional to α_s^n for $n > 3$ are RS-dependent. The form given above for β_2 is in the \overline{MS} scheme. It has become conventional to use the \overline{MS} scheme for calculating QCD cross sections beyond leading order.

The fundamental theorem of RS dependence is straightforward. Physical quantities, in particular the cross section, calculated to all orders in perturbation theory, do not depend on the RS. It follows that a truncated series *does* exhibit RS dependence. In practice, QCD cross sections are known to leading or to next-to-leading order or, in a very few cases, to next-to-next-to-leading order, and it is only the latter two cases, which have reduced RS dependence, that are useful for precision tests. At second order the RS dependence is completely given by one condition which can be taken to be the value of the renormalization scale μ . One therefore has to address the question of what is the “best” choice for μ . There is no definite answer to this question—higher-order corrections do not “fix” the scale, rather they render the theoretical predictions less sensitive to its variation.

One could imagine that choosing a scale μ characteristic of the typical energy scale (E) in the process would be most appropriate. In general a poor choice of scale generates terms of order $\ln(E/\mu)$ in the A_i ’s. More byzantine choices are the scale for which the next-to-leading-order correction vanishes (“Fastest Apparent Convergence [8]”), the scale for which the next-to-leading-order prediction is stationary [9] (*i.e.* the value of μ where $d\sigma/d\mu = 0$), or that dictated by the effective charge scheme [10].

An important corollary is that if the higher-order corrections are naturally small, then the additional uncertainties introduced by the μ

dependence are likely to be less than the experimental measurement errors. There are some processes, however, for which the choice of scheme (*i.e.* the value of μ) can influence the extracted value of $\Lambda_{\overline{MS}}$. There is no resolution to this problem other than to try to calculate even more terms in the perturbation series.

In the cases where the higher-order corrections to a process are known and are large, some caution should be exercised when quoting the value of α_s . In what follows we will attempt to indicate the size of the theoretical uncertainties on the extracted value of α_s . There are two simple ways to determine this error. First, we can estimate it by comparing the value of $\alpha_s(\mu)$ obtained by fitting data using the QCD formula to highest known order in α_s , and then comparing it with the value obtained using the next-to-highest-order formula (μ is chosen as the typical energy scale in the process). The corresponding Λ ’s are then obtained by evolving $\alpha_s(\mu)$ to $\mu = m_Z$ using Eq. (25.4) to the same order in α_s as the fit, and then converting to $\Lambda^{(4)}$ using Eq. (25.6). Alternatively, we can vary the value of μ over a reasonable range, extracting a value of Λ for each choice of μ . This method is of its nature imprecise since “reasonable” involves a subjective judgement. In either case, if the perturbation series is well behaved, the resulting error on Λ will be small.

25.3. QCD in deep-inelastic scattering

The original and still one of the most powerful quantitative tests of perturbative QCD is the breaking of Bjorken scaling in deep-inelastic lepton-hadron scattering. In the leading-logarithm approximation the measured structure functions $F_i(x, Q^2)$ are related to the quark distribution functions $q_i(x, Q^2)$ according to the naive parton model by the formulae in Sec. 24 of this *Review*. (In that section, q_i is denoted by the notation f_q .) In describing the way in which scaling is broken in QCD, it is convenient to define nonsinglet and singlet quark distributions:

$$F^{NS} = q_i - q_j \quad F^S = \sum_i (q_i + \bar{q}_i) \tag{25.8}$$

The nonsinglet structure functions have nonzero values of flavor quantum numbers such as isospin or baryon number. The variation with Q^2 of these is described by the so-called Altarelli-Parisi equations [11]:

$$Q^2 \frac{\partial F^{NS}}{\partial Q^2} = \frac{\alpha_s(|Q|)}{2\pi} P^{qq} * F^{NS} \tag{25.9a}$$

$$Q^2 \frac{\partial}{\partial Q^2} \begin{pmatrix} F^S \\ G \end{pmatrix} = \frac{\alpha_s(|Q|)}{2\pi} \begin{pmatrix} P^{qq} & 2n_f P^{qg} \\ P^{gq} & P^{gg} \end{pmatrix} * \begin{pmatrix} F^S \\ G \end{pmatrix} \tag{25.9b}$$

where $*$ denotes a convolution integral:

$$f * g = \int_x^1 \frac{dy}{y} f(y) g\left(\frac{x}{y}\right) \tag{25.10}$$

The leading-order Altarelli-Parisi splitting functions are

$$P^{qq} = \frac{4}{3} \left[\frac{1+x^2}{1-x} \right]_+ + 2\delta(1-x) \tag{25.11a}$$

$$P^{qg} = \frac{1}{2} \left[x^2 + (1-x)^2 \right] \tag{25.11b}$$

$$P^{gq} = \frac{4}{3} \left[\frac{1+(1-x)^2}{x} \right] \tag{25.11c}$$

$$P^{gg} = 6 \left[\frac{1-x}{x} + x(1-x) + \left(\frac{x}{1-x} \right)_+ + \frac{11}{12} \delta(1-x) \right] - \frac{n_f}{3} \delta(1-x) \tag{25.11d}$$

Here the gluon distribution $G(x, Q^2)$ has been introduced and $1/(1-x)_+$ means

$$\int_0^1 dx \frac{f(x)}{(1-x)_+} = \int_0^1 dx \frac{f(x) - f(1)}{(1-x)} \tag{25.12}$$

The precision of contemporary experimental data demands that higher-order corrections also be included [12]. The above results are for massless quarks. Algorithms exist for the inclusion of nonzero quark masses [13]. At low Q^2 values there are also important “higher-twist” (HT) or nonperturbative contributions of the form:

$$F_i(x, Q^2) = F_i^{(LT)}(x, Q^2) + \frac{F_i^{(HT)}(x, Q^2)}{Q^2} + \dots \quad (25.13)$$

Leading twist (LT) indicates a term whose behavior is predicted by perturbative QCD. These corrections are numerically important only for $Q^2 < \mathcal{O}(10 \text{ GeV}^2)$ except for x very close to 1.

A detailed review of the current status of the experimental data can be found, for example, in [14], [15] and only a brief summary will be presented here. Since the last version of this *Review* [16] appeared, the discrepancies between different experiments have mostly been resolved. We shall only include determinations of Λ from the recently published results; the earlier editions of this *Review* should be consulted for the earlier data. In any event, the recent results will dominate the average since their errors are smaller.

From Eq. (25.9), it is clear that a nonsinglet structure function offers in principle the most precise test of the theory, since the Q^2 evolution is independent of the unmeasured gluon distribution. The CCFRR collaboration fit to the Gross-Llewellyn Smith sum rule [17] is known to order α_s^3 [18]

$$\int_0^1 dx (F_3^{\nu p}(x, Q^2) + F_3^{\nu n}(x, Q^2)) = 3 \left[\left(1 - \frac{\alpha_s}{\pi} \left(1 + 3.58 \frac{\alpha_s}{\pi} + 19.0 \left(\frac{\alpha_s}{\pi} \right)^2 \right) - \Delta HT \right) \right]. \quad (25.14)$$

Where the higher-twist contribution $\Delta HT = 0.032 \pm 0.016$ [18] Using the CCFRR data [19] this gives $\alpha_s(1.73 \text{ GeV}) = 0.320 \pm 0.043(\text{expt.}) \pm 0.029(\text{theory})$. The error from higher-twist terms dominates the theoretical error, the higher-twist term being approximately 50% larger than the α_s^3 term. Recently a measurement of Λ has been made using F_3 in neutrino scattering [20]. The result is $\Lambda_{\overline{\text{MS}}}^{(4)} = 179 \pm 36 \pm 41 \text{ MeV}$. The errors are statistical and systematic but do not include (theoretical) errors arising from the choice of μ^2 . Measurements involving singlet-dominated structure functions such as F_2 result in correlated measurements of $\Lambda_{\overline{\text{MS}}}^{(4)}$ and the gluon distribution. By utilizing high-statistics data at large x (> 0.25) and large Q^2 , where F_2 behaves like a nonsinglet and F_3 at smaller x a nonsinglet fit can be performed with better statistical precision and hence the error on the measured value of $\Lambda_{\overline{\text{MS}}}^{(4)}$ is much reduced. CCFRR gives $\Lambda_{\overline{\text{MS}}}^{(4)} = 210 \pm 28 \pm 41 \text{ MeV}$ [20] from $F_2(\nu N)$ and $F_3(\nu N)$. There is an additional uncertainty of $\pm 59 \text{ MeV}$ from the choice of scale. The NMC collaboration [21] gives $\alpha_s(7 \text{ GeV}^2) = 0.264 \pm 0.018(\text{stat.}) \pm 0.070(\text{syst.}) \pm 0.013(\text{higher-twist})$. The systematic error is larger than the CCFRR result, partially because the data are at smaller values of x and the gluon distribution is more important. A reanalysis [22] of EMC data [23] gives $\Lambda_{\overline{\text{MS}}}^{(4)} = 211 \pm 80 \pm 80 \text{ MeV}$ from $F_2(\nu N)$. Finally a combined analysis [24] of SLAC [25] and BCDMS [26] data gives $\Lambda_{\overline{\text{MS}}}^{(4)} = 263 \pm 42 \pm 55 \text{ MeV}$. Here the systematic error is an estimate of the uncertainty due to the choice of Q^2 used in the argument of α_s and in the scale at which the structure functions (factorization scale) used in the QCD calculation are evaluated.

The results from [20–22] and [24] can be combined to give $\alpha_s(M_Z) = 0.112 \pm 0.002 \pm 0.004$ or equivalently $\Lambda_{\overline{\text{MS}}}^{(4)} = 234 \pm 26 \pm 50 \text{ MeV}$. Here the former error is a combination of statistical and systematic errors and the second error is due to the scale uncertainty. This result is an average of the results weighted by their statistical and systematic errors. The scale error which is common to all is then reapplied to the average.

Typically, Λ is extracted from the data by parametrizing the parton densities in a simple analytic way at some Q_0^2 , evolving to higher Q^2 using the next-to-leading-order evolution equations, and fitting globally to the measured structure functions to obtain $\Lambda_{\overline{\text{MS}}}^{(4)}$.

Thus an important by-product of such studies is the extraction of parton densities at a fixed-reference value of Q_0^2 . These can then be evolved in Q^2 and used as input for phenomenological studies in hadron-hadron collisions (see below). To avoid having to evolve from the starting Q_0^2 value each time, a parton density is required; it is useful to have available a simple analytic approximation to the densities valid over a range of x and Q^2 values. A package is available in the form of the CERN computer library that includes an exhaustive set of fits [27]. Some of these fits are obsolete. In using a parameterization to predict event rates, a next-to-leading order fit must be used if the process being calculated is known to next-to-leading order in QCD perturbation theory. In such a case there is an additional scheme dependence; this scheme dependence is reflected in the $\mathcal{O}(\alpha_s)$ corrections that appear in the relations between the structure functions and the quark distribution functions. There are two common schemes, a deep-inelastic scheme where there are no order α_s corrections in the formula for $F_2(x, Q^2)$ and the minimal subtraction scheme. It is important, when these next-to-leading order fits are used in other processes (see below), that the same scheme is used in the calculation of the partonic rates.




Figure 25.1: Summary of the values of $\alpha_s(M_Z)$ and $\Lambda^{(5)}$ from various processes ordered from top to bottom by increasing energy scale of the measurements. The values shown indicate the process and the measured value of α_s extrapolated up to $\mu = M_Z$. The error shown is the *total* error including theoretical uncertainties. The weighted average is obtained from the above values using the method discussed in the text.

25.4. QCD in decays of the tau lepton

The semi-leptonic branching ratio of the tau (R_τ) is an inclusive quantity. It is related to the contribution of hadrons to the imaginary part of the W self energy ($\text{Im}(\Pi(s))$), just as R (see below, Sec. 25.7) is related to the imaginary part of the photon self energy. However it is more inclusive than R since it involves an integral

$$R_\tau \sim \int_0^{m_\tau^2} \frac{ds}{m_\tau^2} \left(1 - \frac{s}{m_\tau^2} \right)^2 \text{Im}(\Pi(s))$$

Since the scale involved is low, one must take into account nonperturbative (higher-twist) contributions which are suppressed by powers of the tau mass.

$$R_\tau = 3.058 \left[1 + \frac{\alpha_s(m_\tau)}{\pi} + 5.2 \left(\frac{\alpha_s(m_\tau)}{\pi} \right)^2 + 26.4 \left(\frac{\alpha_s(m_\tau)}{\pi} \right)^3 + a \frac{m^2}{m_\tau^2} + b \frac{m\psi\bar{\psi}}{m_\tau^4} + c \frac{\psi\bar{\psi}\psi\bar{\psi}}{m_\tau^6} + \dots \right]$$

Here a, b , and c are dimensionless constants and m is a light quark mass. The term of order $1/m_\tau^2$ is a kinematical effect due to the light quark masses and is consequently very small. The nonperturbative terms are estimated using sum rules [28]. In total they are estimated to be -0.007 ± 0.004 [29]. This estimate relies on there being no term of order Λ^2/m_τ^2 (note that $\frac{\alpha_s(m_\tau)}{\pi} \sim (\frac{0.5 \text{ GeV}}{m_\tau})^2$). The a, b and c can be determined from the data [30] by fitting to moments of the $\Pi(s)$. The values so extracted [31] are consistent with the theoretical estimates. If the nonperturbative terms are omitted from the fit, the extracted value of $\alpha_s(m_\tau)$ decreases by ~ 0.02 .

For $\alpha_s(m_\tau) = 0.36$ the perturbative series for R_τ is $R_\tau \sim 3.058(1 + 0.114 + 0.073 + 0.043)$. The size (estimated error) of the nonperturbative term is 20% (7%) of the size of the order α_s^3 term. The perturbation series is not very well convergent; if the order α_s^3 term is omitted the extracted value of $\alpha_s(m_\tau)$ increases by 0.05. Using the experimental average [32] for R_τ of 3.6174 ± 0.034 gives $\alpha_s(m_\tau) = 0.360 \pm 0.031$ using the experimental error alone. We assign a theoretical error equal to 1/2 of the contribution from the order α^3 and nonperturbative contributions. This then gives $\alpha_s(m_\tau) = 0.360 \pm 0.041$ for the final result. Note that the experimental errors are dominant. The small theoretical errors have been criticised [33]. Here it is claimed that the presence of hadronic resonances limits the applicability of perturbative QCD and that the theoretical errors are underestimated by at least a factor of 2. If this argument is correct the agreement with values of α_s from other processes is accidental.

25.5. QCD in high-energy hadron collisions

There are many ways in which perturbative QCD can be tested in high-energy hadron colliders. The quantitative tests are only useful if the process in question has been calculated beyond leading order in QCD perturbation theory. The production of hadrons with large transverse momentum in hadron-hadron collisions provides a direct probe of the scattering of quarks and gluons: $qq \rightarrow qq$, $qg \rightarrow qg$, $gg \rightarrow gg$, *etc.* The present generation of $p\bar{p}$ colliders provide center-of-mass energies which are sufficiently high that these processes can be unambiguously identified in two-jet production at large transverse momentum. Recent higher-order QCD calculations of the jet rates [34] and shapes are in impressive agreement with data [35]. As an example, Fig. 32.7 in this *Review* shows the inclusive jet cross section at zero pseudorapidity as a function of the jet transverse momentum for $p\bar{p}$ collisions. Data are also available on the angular distribution of jets; these are also in agreement with QCD expectations [36,37].

QCD corrections to Drell-Yan type cross sections (*i.e.*, the production in hadron collisions by quark-antiquark annihilation of lepton pairs of invariant mass Q from virtual photons, or of real W or Z bosons) are known [38]. These $\mathcal{O}(\alpha_s)$ QCD corrections are sizable at small values of Q .

It is interesting to note that the corresponding correction to W and Z production, as measured in $p\bar{p}$ collisions at $\sqrt{s} = 0.63$ TeV and $\sqrt{s} = 1.8$ TeV has essentially the same theoretical form and is of order 30%.

The production of W and Z bosons and photons at large transverse momentum can also be used to determine α_s . The leading-order QCD subprocesses are $q\bar{q} \rightarrow \gamma g$ and $qg \rightarrow \gamma q$. If the parton distributions are taken from other processes and a value of $\Lambda_{\overline{\text{MS}}}^{(4)}$ assumed, then an absolute prediction is obtained. Conversely, the data can be used to extract information on quark and gluon distributions and on the value of $\Lambda_{\overline{\text{MS}}}^{(4)}$. The next-to-leading-order QCD corrections are known [39,40] (for photons) and for W/Z production [41], and so a precision test is possible in principle. The UA2 collaboration [42] has extracted a value of $\alpha_s(M_W) = 0.123 \pm 0.018(\text{stat.}) \pm 0.017(\text{syst.})$ from the measured ratio $R_W = \frac{\sigma(W + 1jet)}{\sigma(W + 0jet)}$. The result depends on the algorithm used to define a jet, and the dominant systematic errors due to fragmentation and corrections for underlying events are connected to the algorithm (the former causes jet energy to be lost, the latter causes it to be increased). The scale at which $\alpha_s(M)$ is to be evaluated is not clear. A change from $M = M_W$ to $M = M_W/2$

causes a shift of 0.01 in the extracted α_s . The quoted error has been increased to take this into account.

25.6. QCD in heavy quarkonium decay

Under the assumption that the hadronic and leptonic decay widths of heavy $Q\bar{Q}$ resonances can be factorized into a nonperturbative part—dependent on the confining potential—and a calculable perturbative part, the ratios of partial decay widths allow measurements of α_s at the heavy quark mass scale. The most precise data come from the decay widths of the 1^{--} $J/\psi(1S)$ and Υ resonances. The total decay width of the Υ is predicted by perturbative QCD [43]

$$R_\Upsilon = \frac{\Gamma(\Upsilon \rightarrow \text{hadrons})}{\Gamma(\Upsilon \rightarrow \mu^+\mu^-)} = \frac{10(\pi^2 - 9)\alpha_s^3(M)}{9\pi\alpha_{\text{em}}^2} \left[1 + \frac{\alpha_s}{\pi} \left(-19.4 + \frac{3\beta_0}{2} \left(1.162 + \ln\left(\frac{2M}{M_\tau}\right) \right) \right) \right]$$

Data are available for the Υ , Υ' , Υ'' and ψ . The result is very sensitive to α_s and the data are sufficiently precise ($R_\mu(\Upsilon) = 32.5 \pm 0.9$) [44] that the theoretical errors will dominate. There are theoretical corrections to this simple formula due to the relativistic nature of the $Q\bar{Q}$ system; $v^2/c^2 \sim 0.1$ for the Υ . They are more severe for the ψ . There are also nonperturbative corrections of the form Λ^2/m_τ^2 ; again these are more severe for the ψ . A fit to Υ , Υ' , and Υ'' [45] gives $\alpha_s(M_Z) = 0.108 \pm 0.001(\text{expt.})$. The results from each state separately and also from the ψ are consistent with each other. There is an uncertainty of order ± 0.005 from the choice of scale; the error from v^2/c^2 corrections is a little larger. $\alpha_s(M_Z) = 0.108 \pm 0.010$ is a fair representation of the total error including the possibility of nonperturbative corrections.

25.7. Perturbative QCD in e^+e^- collisions

The total cross section for $e^+e^- \rightarrow \text{hadrons}$ is obtained (at low values of \sqrt{s}) by multiplying the muon-pair cross section by the factor $R = 3\Sigma_q e_q^2$. The higher-order QCD corrections to this quantity have been calculated, and the results can be expressed in terms of the factor:

$$R = R^{(0)} \left[1 + \frac{\alpha_s}{\pi} + C_2 \left(\frac{\alpha_s}{\pi} \right)^2 + C_3 \left(\frac{\alpha_s}{\pi} \right)^3 + \dots \right], \quad (25.15)$$

where $C_2 = 1.411$ and $C_3 = -12.8$ [46].

$R^{(0)}$ can be obtained from the formula for $d\sigma/d\Omega$ for $e^+e^- \rightarrow f\bar{f}$ by integrating over Ω . The formula is given in Sec. 24.2 of this *Review*. This result is strictly only correct in the zero-quark-mass limit. The $\mathcal{O}(\alpha_s)$ corrections are also known for massive quarks [47].

A comparison of the theoretical prediction of Eq. (25.15) (corrected for the b -quark mass) with all the available data at values of \sqrt{s} between 20 and 65 GeV gives [48] $\alpha_s(35 \text{ GeV}) = 0.146 \pm 0.030$. The principal advantage of determining α_s from R in e^+e^- annihilation is that there is no dependence on fragmentation models, jet algorithms, *etc.* The size of the order α_s^3 term is of order 40% of that of the order α_s^2 and 3% of the order α_s . If the order α_s^3 term is not included a fit to the data yields $\alpha_s(34 \text{ GeV}) = 0.142 \pm 0.03$, indicating that the theoretical uncertainty is smaller than the experimental error.

Measurements of the ratio of hadronic to leptonic width of the Z at LEP Γ_h/Γ_μ probe the same quantity as R . Using the average of $\Gamma_h/\Gamma_\mu = 20.781 \pm 0.049$ gives $\alpha_s(M_Z) = 0.124 \pm 0.007$ [49]. There are theoretical errors arising from the values of the top quark and Higgs masses which enter due to electroweak corrections to the Z width and from the choice of scale. The dominant error is the first of these (± 0.005).

An alternative method of determining α_s in e^+e^- annihilation is from measuring quantities that are sensitive to the relative rates of two-, three-, and four-jet events. In addition to simply counting jets, there are many possible choices of such “shape variables”: thrust [50], energy-energy correlations [51], planar triple-energy correlations [52], average jet mass, *etc.* All of these are infrared safe, which means they can be reliably calculated in perturbation theory. The starting point

for all these quantities is the multijet cross section. For example, at order α_s , for the process $e^+e^- \rightarrow q\bar{q}g$:

$$\frac{1}{\sigma} \frac{d^2\sigma}{dx_1 dx_2} = \frac{2\alpha_s}{3\pi} \frac{x_1^2 + x_2^2}{(1-x_1)(1-x_2)}, \quad (25.16)$$

where

$$x_i = \frac{2E_i}{\sqrt{s}} \quad (25.17)$$

are the center-of-mass energy fractions of the final-state (massless) quarks. A distribution in a “three-jet” variable, such as those listed above, is obtained by integrating this differential cross section over an appropriate phase space region for a fixed value of the variable. The order α_s^2 corrections to this process have been computed as well as the 4-jet final states such as $e^+e^- \rightarrow q\bar{q}g\bar{g}$.

There are many methods used by the LEP groups [53–56] to determine α_s from the event topology. The jet-counting algorithm originally introduced by the JADE collaboration [57] has been used by the LEP groups. Here particles of momenta p_i and p_j are combined into a pseudo-particle of momentum $p_i + p_j$ if the invariant mass of the pair is less than $y_0\sqrt{s}$. The process is then iterated until no more pairs of particles or pseudo-particles remain. The remaining number is then defined to be the number of jets in the event and can be compared to the QCD prediction.

There are theoretical ambiguities in the way that this process is carried out; quarks and gluons are massless whereas the observed hadrons are not. So that the massive jets that result from this scheme cannot be compared directly to the massless jets of perturbative QCD. Different recombination schemes have been tried, for example combining 3-momenta and then rescaling the energy of the cluster so that it remains massless, and these result in the same data giving a slightly different value [58] of α_s . These differences can be used to determine a systematic error. In addition, since what is observed is hadrons rather than quarks and gluons, a model is needed to describe the evolution of a partonic final state into one involving hadrons so that detector corrections can be applied. The second-order matrix elements are combined with a parton-fragmentation model. This model can then be used to correct the data for a direct comparison with the parton calculation. The different hadronization models that are used [59–62] model the dynamics that are controlled by nonperturbative QCD effects which we cannot yet calculate. The fragmentation parameters of these Monte Carlos are tuned to get agreement with the observed data. The differences between these models contribute to the systematic errors. The systematic errors from recombination schemes and fragmentation effects dominate over the statistical and other errors of the LEP experiments.

The scale M at which $\alpha_s(M)$ is to be evaluated is not clear. The invariant mass of a typical jet (or $\sqrt{s}y_0$) is probably a more appropriate choice than the e^+e^- center of mass energy. If the value is allowed to float in the fit to the data, the data tend to prefer values of order $\sqrt{s}/10$ [63]; the exact value depends on the variable that is fitted. The dominant uncertainties arise from the choice of M and from the freedom in the fragmentation Monte Carlos. An average of many variables and all the LEP experiments [65] gives $\alpha_s(M_Z) = 0.119 \pm 0.006$. The dominant error is theoretical. The situation might improve if complete perturbative calculations to order α_s^3 became available.

The perturbative QCD formulae can break down in special kinematical configurations. For example, the thrust distribution contains terms of the type $\alpha_s \ln^2(1-T)$. The higher orders in the perturbation expansion contains terms of order $\alpha_s^n \ln^m(1-T)$. For $T \sim 1$ (the region populated by 2-jet events) the perturbation expansion is unreliable. The terms with $n \leq m$ can be summed to all orders in α_s [66]. The resummed results give better agreement with the data at large values of T [67]. Such resummed results are not available for all of the shape variables [69], but fits using those available yields a LEP average of $\alpha_s(M_Z) = 0.124 \pm 0.006$ [65]. Some caution should be exercised in using these resummed results because of the possibility of overcounting; the showering Monte Carlos that are used for the fragmentation corrections also generate some of these

leading log corrections. The errors in the values of $\alpha_s(M_Z)$ from these shape variables are totally dominated by the theoretical uncertainties associated with the choice of scale and the effects of hadronization Monte Carlos. It is gratifying that the shift from the result using the resummed formulae is less than the error. Since the data fitted to the resummed and not-resummed formulae are the same. We will use the average of $\alpha_s(M_Z) = 0.121 \pm 0.006$. Since the error is dominantly theoretical, it does not reduce in the average.

Similar studies on event shapes have been undertaken at TRISTAN and at PEP/PETRA. A combined result from various shape parameters by the TOPAZ collaboration gives $\alpha_s(58 \text{ GeV}) = 0.125 \pm 0.009$ using the fixed order QCD result and $\alpha_s(58 \text{ GeV}) = 0.132 \pm 0.008$ using the resummed result [70]. We average these and use $\alpha_s(58 \text{ GeV}) = 0.128 \pm 0.009$. The AMY group has fitted a next to leading order QCD Monte Carlo [68] to the single particle momentum distribution inside jets. The analysis is analogous to the fitting of moments in deep-inelastic scattering. They quote $\alpha_s(58 \text{ GeV}) = 0.134 \pm 0.006$ [71]. Jets are selected so that they are dominantly quark jets since the Monte Carlo does not well describe gluon jets. No systematic error is included for the contribution of gluon jets to the result. The errors could be understated therefore and we have chosen to use $\alpha_s(58 \text{ GeV}) = 0.134 \pm 0.008$ in order to be conservative.

The measurements of event shapes at PEP/PETRA are summarized in earlier editions of this note. The results are consistent with those from Z decay, but have larger errors. We use $\alpha_s(34 \text{ GeV}) = 0.14 \pm 0.02$ [72]. Since the errors in the event shape measurements are dominantly systematic, the results from PEP/PETRA, TRISTAN and LEP are combined to give $\alpha_s(M_Z) = 0.121 \pm 0.006$. This result is then used in forming the final average value of α_s .

There are many other ways in which QCD can be tested in electron-positron collisions. Mention should be made in particular of the interesting and important results from “two-photon” processes. See the previous edition of this *Review* [16] for more information. There are no new results and the data do not contribute significantly to the average.

25.8. Lattice QCD

Lattice gauge theory calculations can be used to calculate the energy levels of a $Q\bar{Q}$ system and then extract α_s . The FNAL group [73] uses the splitting between the $1S$ and $1P$ in the charmonium system [$m_{hc} - (3m_\psi + m_{\eta_c})/4 = 456.6 \pm 0.4 \text{ MeV}$] to determine α_s . The result quoted is $\alpha_s(M_Z) = 0.108 \pm 0.006$. The splitting is almost independent of the charm-quark mass and is therefore dependent only on Λ . The calculation does not rely on perturbation theory or on nonrelativistic approximation. The main errors are systematic associated with the finite lattice spacing (a), the matching to the perturbatively defined α_s , and quenched approximation used in the calculation. The extrapolation to zero-lattice spacing produces a shift in Λ of order 5% and is therefore quite small. The quenched approximation is more serious. No light quarks are allowed to propagate and hence the extracted value of Λ corresponds to the case of zero flavors. $\alpha_s(M)$ is evolved down from the scale ($\sim 2.3 \text{ GeV}$) of the lattice used to the scale of momentum transfers appropriate to the charmonium system ($\sim 700 \text{ MeV}$). The resulting coupling is then evolved back up with the correct number of quark flavors. This produces a shift in $\alpha_s(5 \text{ GeV})$ of order 25%, with a claimed uncertainty of 7%. This error dominates and could be an underestimate as the perturbative running of $\alpha_s(M)$ has to be used at small M . In addition a recent calculation [74] using the strength of the force between two heavy quarks computed in the quenched approximation obtains a value of $\alpha_s(5 \text{ GeV})$ that is consistent with this result. Calculations based on the \mathcal{T} spectrum using non-relativistic lattice theory give $\alpha_s(M_Z) = 0.112 \pm 0.004$ [75]. Here again the dominant error arises from the quenched approximation. We average the two results since they have a common systematic error and use $\alpha_s(M_Z) = 0.110 \pm 0.006$ in the final average.




Figure 25.2: Summary of the values of $\alpha_s(\mu)$ at the values of μ where they are measured. The lines show the central values and the $\pm 1\sigma$ limits of our average. The figure clearly shows the decrease in $\alpha_s(\mu)$ with increasing μ .

25.9. Conclusions

The need for brevity has meant that many other important topics in QCD phenomenology have had to be omitted from this review. One should mention in particular the study of exclusive processes (form factors, elastic scattering, ...), the behavior of quarks and gluons in nuclei, the spin properties of the theory and the importance of polarized scattering data, the interface of soft and hard QCD as manifest, for example, by minijet production and hard diffractive processes, and QCD effects in hadron spectroscopy.

In this short review we have focused on those high-energy processes which currently offer the most quantitative tests of perturbative QCD. Emphasis has been given to the recent data from LEP and deep-inelastic scattering. Figure 25.1 shows the values of $\alpha_s(M_Z)$ deduced from the various experiments. Figure 25.2 shows the values and the values of Q where they are measured. This figure clearly shows the experimental evidence for the variation of $\alpha_s(Q)$ with Q .

An average of the values in Fig. 25.1 gives $\alpha_s(M_Z) = 0.117$ with a total χ^2 of 5.9 for nine fitted points. The error on the average assuming that all of the errors in the contributing results are uncorrelated is ± 0.0023 and is surely an underestimate. Since, in most cases the dominant error is systematic (mainly theoretical), a more reasonable estimate is to use the smallest of the individual errors on the experimental results *i.e.* ± 0.005 . Our value is then $\alpha_s(M_Z) = 0.117 \pm 0.005$ which corresponds to $\Lambda^{(5)} = 195 + 65 - 50$ MeV.

The prospects for improvement in the error are controlled mainly by the potential improvements in the theoretical errors. Lattice calculations that do not resort to the quenched approximation offer one of the best opportunities. On the experimental side, the direct determination of the top quark mass and a smaller error on the hadronic width of the Z should result in a significantly reduced error on α_s .

References:

1. F.J. Yndurain, "Quantum Chromodynamics: an Introduction to the Theory of Quarks and Gluons", (Springer, New York, 1983).
2. For a recent review, see for example A.S. Kronfeld and P.B. Mackenzie, *Ann. Rev. Nucl. and Part. Sci.* **43**, 793 (1993).
3. For example see, S. Catani in *Proceedings of the International Europhysics Conference on High Energy Physics*, Marseille, France (July 1993), preprint DFF/194/11/93;
S. Bethke, in *Proceedings of the International Conference on High Energy Physics*, Dallas TX, (July 1992).
4. W. Bernreuther, *Ann. Phys.* **151**, 127 (1983).
5. W. Marciano, *Phys. Rev.* **D29**, 580 (1984).
6. D.W. Duke and R.G. Roberts, *Phys. Reports* **120**, 275 (1985).
7. W.A. Bardeen *et al.*, *Phys. Rev.* **D18**, 3998 (1978).
8. G. Grunberg, *Phys. Lett.* **95B**, 70 (1980), and *Phys. Rev.* **D29**, 2315 (1984).
9. P.M. Stevenson, *Phys. Rev.* **D23**, 2916 (1981), and *Nucl. Phys.* **B203**, 472 (1982).
10. S. Brodsky and H.J. Lu SLAC-PUB-6389, (Nov 1993).
11. G. Altarelli and G. Parisi, *Nucl. Phys.* **B126**, 298 (1977).
12. G. Curci, W. Furmanski, and R. Petronzio, *Nucl. Phys.* **B175**, 27 (1980);
W. Furmanski and R. Petronzio, *Phys. Lett.* **97B**, 437 (1980), and *Z. Phys.* **C11**, 293 (1982);
E.G. Floratos, C. Kounnas, and R. Lacaze, *Phys. Lett.* **98B**, 89 (1981), *Phys. Lett.* **98B**, 285 (1981), and *Nucl. Phys.* **B192**, 417 (1981);
R.T. Herrod and S. Wada, *Phys. Lett.* **96B**, 195 (1981), and *Z. Phys.* **C9**, 351 (1981).
13. M. Glück *et al.*, *Z. Phys.* **C13**, 119 (1982).
14. M. Virchaux, Saclay preprint, DAPHNIA/SPP 92-30, presented at THE "QCD 20 years later" workshop Aachen Germany (Sep. 9-13, 1992).
15. R. Voss, in *Proceedings of the 1993 International Symposium on Lepton and Photon Interactions at High Energies*, Ithaca, NY (July 1993).
16. Particle Data Group, *Phys. Rev.* **D45**, Part II (1992).
17. D. Gross and Llewellyn Smith, *Nucl. Phys.* **B14**, 337 (1969).
18. J. Chyla and A.L. Kataev, *Phys. Lett.* **B297**, 385 (1992).
19. W.C. Leung *et al.*, *Phys. Lett.* **B317**, 655 (1993).
20. P.Z. Quintas *et al.*, *Phys. Rev. Lett.* **71**, 1307 (1993);
J.P. Berge *et al.*, *Z. Phys.* **C49**, 187 (1991).
21. M. Arneodo *et al.*, *Phys. Lett.* **B309**, 222 (1993).
22. K. Bazizi and S.J. Wimpenny, UCR/DIS/91-02.
23. J.J. Aubert *et al.*, *Nucl. Phys.* **B333**, 1 (1990); *Nucl. Phys.* **293**, 740 (1987); *Nucl. Phys.* **B272**, 158 (1986); and *Nucl. Phys.* **B145**, 189 (1985).
24. M. Virchaux and A. Milsztajn, *Phys. Lett.* **B274**, 221 (1992).
25. L.W. Whitlow, Ph.D thesis, SLAC Report 357 (1990).
26. A.C. Benvenuti *et al.*, *Phys. Lett.* **B195**, 97 (1987); *Phys. Lett.* **B223**, 490 (1989); *Phys. Lett.* **B223**, 485, (1989); *Phys. Lett.* **B237**, 592 (1990); and *Phys. Lett.* **B237**, 599 (1990).
27. H. Plothow-Besch, PDFLIB package in CERNLIB.
28. M.A. Shifman, A.I. Vainshtein, and V.I. Zakharov *Nucl. Phys.* **B147**, 385 (1979).
29. S. Narison and A. Pich *Phys. Lett.* **B211**, 183 (1988);
E. Braaten, S. Narison, and A. Pich, *Nucl. Phys.* **B373**, 581 (1992).
30. F. Le Diberder and A. Pich *Phys. Lett.* **B289**, 165 (1992).
31. D. Buskulic *et al.*, *Phys. Lett.* **B307**, 209 (1993).
32. P.D. Acton *et al.*, *Phys. Lett.* **B288**, 373 (1992);
B. Adeva *et al.*, *Phys. Lett.* **B265**, 451 (1991);
P. Abreu *et al.*, *Z. Phys.* **C55**, 555 (1992);
D. Decamp *et al.*, *Z. Phys.* **C54**, 211 (1992). Average reported by E. Fernandez in *Proceedings of the Europhysics Conference on High Energy Physics* Marseilles, France (July, 1992).
33. T. Truong, *Phys. Rev.* **D47**, 3999 (1993), preprint EP-CPTH-A226-1093 (1993).
34. S.D. Ellis, Z. Kunszt, and D.E. Soper, *Phys. Rev. Lett.* **64**, 2121 (1991).
35. F. Abe *et al.*, *Phys. Rev. Lett.* **68**, 1104 (1992).
36. UA1 collaboration: G. Arnison *et al.*, *Phys. Lett.* **B177**, 244 (1986).
37. F. Abe *et al.*, *Phys. Rev.* **D45**, 1448 (1992).

38. G. Altarelli, R.K. Ellis, and G. Martinelli, Nucl. Phys. **B143**, 521 (1978).
39. P. Aurenche, R. Baier, and M. Fontannaz Phys. Rev. **D42**, 1440 (1990);
P. Aurenche *et al.*, Phys. Lett. **140B**, 87 (1984);
P. Aurenche *et al.*, Nucl. Phys. **B297**, 661 (1988).
40. H. Baer, J. Ohnemus, and J.F. Owens, Phys. Lett. **B234**, 127 (1990).
41. H. Baer and M.H. Reno, Phys. Rev. **D43**, 2892 (1991);
P.B. Arnold and M.H. Reno, Nucl. Phys. **B319**, 37 (1989).
42. J. Alitti *et al.*, Phys. Lett. **B263**, 563 (1991).
43. R. Barbieri *et al.*, Phys. Lett. **95B**, 93 (1980);
B.P. Mackenzie and G.P. Lepage Phys. Rev. Lett. **47**, 1244 (1981).
44. M. Kobel *et al.*, Z. Phys. **C53**, 193 (1992).
45. M. Kobel DESY-F31-91-03.
46. S.G. Gorishny, A. Kataev, and S.A. Larin, Phys. Lett. **B259**, 114 (1991);
L.R. Surguladze and M.A. Samuel, Phys. Rev. Lett. **66**, 560 (1991).
47. E.C. Poggio *et al.*, Phys. Rev. **D13**, 1958 (1976);
R.M. Barnett, M. Dine, and L. McLerran, Phys. Rev. **D22**, 594 (1980).
48. D. Haidt, in *Directions in High Energy Physics*, Vol. 14, ed. P. Langacker (World Scientific, 1993).
49. LEP Collaborations CERN/PPE/93-157.
50. E. Farhi, Phys. Rev. Lett. **39**, 1587 (1977).
51. C.L. Basham *et al.*, Phys. Rev. **D17**, 2298 (1978).
52. F. Csikor *et al.*, Phys. Rev. **D31**, 1025 (1985).
53. M.Z. Akrawy *et al.*, Phys. Lett. **B252**, 159 (1990), Z. Phys. **C47**, 505 (1990);
P.D. Acton *et al.*, Phys. Lett. **B276**, 547 (1992).
54. B. Adeva *et al.*, Phys. Lett. **B248**, 464 (1990), Phys. Lett. **B263**, 551 (1991).
55. D. Decamp *et al.*, Phys. Lett. **B257**, 479 (1991), Phys. Lett. **B25**, 623 (1991).
56. P. Abreu *et al.*, Phys. Lett. **B252**, 149 (1990), Phys. Lett. **B247**, 167 (1990).
57. S. Bethke *et al.*, Phys. Lett. **B213**, 235 (1988).
58. M.Z. Akrawy *et al.*, Z. Phys. **C49**, 375 (1991).
59. B. Andersson *et al.*, Phys. Reports **97**, 33 (1983).
60. A. Ali *et al.*, Nucl. Phys. **B168**, 409 (1980);
A. Ali and R. Barreiro, Phys. Lett. **118B**, 155 (1982).
61. B.R. Webber, Nucl. Phys. **B238**, 492 (1984).
62. T. Sjostrand and M. Bengtsson, Comp. Phys. Comm. **43**, 367 (1987).
63. O. Adriani *et al.*, Phys. Lett. **B284**, 471 (1992);
M. Akrawy *et al.*, Z. Phys. **C47**, 505 (1990);
B. Adeva *et al.*, Phys. Lett. **B248**, 473 (1990);
P. Abreu *et al.*, Z. Phys. **C54**, 55 (1992);
B. Decamp *et al.*, Phys. Lett. **B255**, 623 (1991).
64. P. Abreu *et al.*, CERN PPE/91-181.
65. S. Bethke, *Proceedings of the International Conference on High Energy Physics*, Dallas, TX (August 1992) and HD-PY-92/12.
66. S. Catani *et al.*, Phys. Lett. **B263**, 491 (1991).
67. B. Decamp *et al.*, Phys. Lett. **B284**, 163 (1992);
B. Adeva *et al.*, Phys. Lett. **B284**, 471 (1992);
P.D. Acton *et al.*, Z. Phys. **C55**, 1 (1992).
68. K. Kato and T. Munechisa, Comp. Phys. Comm. **64**, 67 (1991).
69. S. Catani *et al.*, Phys. Lett. **B269**, 432 (1991);
S. Catani, B.R. Webber, and G. Turnock, Phys. Lett. **B272**, 368 (1991);
N. Brown and J. Stirling, Z. Phys. **C53**, 629 (1992).
70. Y. Onishi *et al.*, Phys. Lett. **B313**, 475 (1993).
71. K.B. Lee *et al.*, Phys. Lett. **B313**, 469 (1993).
72. W.J. Stirling and M.R. Whalley, Durham-RAL Database Publication RAL/87/107 (1987);
S.L. Wu, in *Proceedings of the 1987 International Symposium on Lepton and Photon Interactions at High Energies, Hamburg, 27-31 July 1987*, ed. by W. Bartel and R. Ruckl (North-Holland, Amsterdam, 1988), p. 39;
W.J. Stirling, in *Proceedings of the 1987 International Symposium on Lepton and Photon Interactions at High Energies, Hamburg, 27-31 July 1987*, ed. by W. Bartel and R. Ruckl (North-Holland, Amsterdam, 1988), p. 715.
73. A.X. El-Khadra *et al.*, Phys. Rev. Lett. **69**, 729 (1992); A.X. El-Khadra, presented at 1993 Lattice Conference, OHSTPY-HEP-T-93-020, A.X. El-Khadra *et al.*, FNAL 94-091/T (1994).
74. M. Luscher *et al.*, Nucl. Phys. **B413**, 481 (1994).
75. C.T.H. Davies *et al.*, Phys. Rev. Lett. (to appear);
G.P. Lepage and J. Sloan presented at 1993 Lattice Conference, HEP-LAT/9312070.

26. STANDARD MODEL OF ELECTROWEAK INTERACTIONS

This section prepared May 1994 by P. Langacker and J. Erler.

The standard electroweak model is based on the gauge group [1] $SU(2) \times U(1)$, with gauge bosons W_μ^i , $i = 1, 2, 3$, and B_μ for the $SU(2)$ and $U(1)$ factors, respectively, and the corresponding gauge coupling constants g and g' . The left-handed fermion fields $\psi_i = \begin{pmatrix} \nu_i \\ \ell_i^- \end{pmatrix}$ and $\begin{pmatrix} u_i \\ d_i^- \end{pmatrix}$ of the i^{th} fermion family transform as doublets under $SU(2)$, where $d_i^- \equiv \sum_j V_{ij} d_j$, and V is the Cabibbo-Kobayashi-Maskawa mixing matrix.* The right-handed fields are $SU(2)$ singlets. In the minimal model there are three fermion families and a single complex Higgs doublet $\phi \equiv \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix}$.

After spontaneous symmetry breaking the Lagrangian is

$$\begin{aligned} \mathcal{L}_F = & \sum_i \bar{\psi}_i \left(i \not{\partial} - m_i - \frac{gm_i H}{2M_W} \right) \psi_i \\ & - \frac{g}{2\sqrt{2}} \sum_i \bar{\psi}_i \gamma^\mu (1 - \gamma^5) (T^+ W_\mu^+ + T^- W_\mu^-) \psi_i \\ & - e \sum_i q_i \bar{\psi}_i \gamma^\mu \psi_i A_\mu \\ & - \frac{g}{2 \cos \theta_W} \sum_i \bar{\psi}_i \gamma^\mu (g_V^i - g_A^i \gamma^5) \psi_i Z_\mu. \end{aligned} \quad (26.1)$$

$\theta_W \equiv \tan^{-1}(g'/g)$ is the weak angle; $e = g \sin \theta_W$ is the positron electric charge; and $A \equiv B \cos \theta_W + W^3 \sin \theta_W$ is the (massless) photon field. $W^\pm \equiv (W^1 \mp iW^2)/\sqrt{2}$ and $Z \equiv -B \sin \theta_W + W^3 \cos \theta_W$ are the massive charged and neutral weak boson fields, respectively. T^+ and T^- are the weak isospin raising and lowering operators. The vector and axial couplings are

$$g_V^i \equiv t_{3L}(i) - 2q_i \sin^2 \theta_W \quad (26.2)$$

$$g_A^i \equiv t_{3L}(i), \quad (26.3)$$

where $t_{3L}(i)$ is the weak isospin of fermion i (+1/2 for u_i and ν_i ; -1/2 for d_i and e_i) and q_i is the charge of ψ_i in units of e .

The second term in \mathcal{L}_F represents the charged-current weak interaction [2]. For example, the coupling of a W to an electron and a neutrino is

$$-\frac{e}{2\sqrt{2} \sin \theta_W} \left[W_\mu^- \bar{\nu} \gamma^\mu (1 - \gamma^5) \nu + W_\mu^+ \bar{\nu} \gamma^\mu (1 - \gamma^5) e \right]. \quad (26.4)$$

For momenta small compared to M_W , this term gives rise to the effective four-fermion interaction with the Fermi constant given (at tree level, *i.e.*, lowest order in perturbation theory) by $G_F/\sqrt{2} = g^2/8M_W^2$. CP violation is incorporated in the Standard Model by a single observable phase in V_{ij} . The third term in \mathcal{L}_F describes electromagnetic interactions (QED), and the last is the weak neutral-current interaction.

In Eq. (26.1), m_i is the mass of the i^{th} fermion ψ_i . For the quarks these are the current masses. For the light quarks, a typical estimate [3] gives $m_u \approx 5.6 \pm 1.1$ MeV, $m_d \approx 9.9 \pm 1.1$ MeV, $m_s \approx 199 \pm 33$ MeV, and $m_c \approx 1.35 \pm 0.05$ GeV (these are running masses evaluated at 1 GeV). For the heavier quarks $m_b \approx 4.7$ GeV (the ‘‘pole’’ mass), and $m_t > 131$ GeV [4]. (The CDF Collaboration has reported [5] evidence for top events with $m_t = 174 \pm 16$ GeV.) See ‘‘The Note on Quark Masses’’ in the Full Listings for more information.

H is the physical neutral Higgs scalar which is the only remaining part of ϕ after spontaneous symmetry breaking. The Yukawa coupling of H to ψ_i , which is flavor diagonal in the minimal model, is $gm_i/2M_W$. The H mass is not predicted by the model. Experimental limits are given in the Higgs section. In nonminimal models there are additional charged and neutral scalar Higgs particles [6].

26.1. Renormalization and radiative corrections

The Standard Model has three parameters (not counting M_H and the fermion masses and mixings). A particularly useful set is:

- (a) The fine structure constant $\alpha = 1/137.036$, determined from the quantum Hall effect. In most electroweak-renormalization schemes, it is convenient to define a running α dependent on the energy scale of the process, with $\alpha^{-1} \sim 137$ appropriate at low energy. At energies of order M_Z , $\alpha^{-1} \sim 128$. For example, in the modified minimal subtraction ($\overline{\text{MS}}$) scheme, one has $\hat{\alpha}(M_Z)^{-1} = 127.9 \pm 0.1$ [7], with the uncertainty due to the low-energy hadronic contribution to vacuum polarization.
- (b) The Fermi constant, $G_F = 1.16639(2) \times 10^{-5}$ GeV $^{-2}$, determined from the muon lifetime formula [8]:

$$\begin{aligned} \tau_\mu^{-1} = & \frac{G_F^2 m_\mu^5}{192\pi^3} F \left(\frac{m_e^2}{m_\mu^2} \right) \left(1 + \frac{3}{5} \frac{m_\mu^2}{M_W^2} \right) \\ & \times \left[1 + \frac{\alpha(m_\mu)}{2\pi} \left(\frac{25}{4} - \pi^2 \right) \right], \end{aligned} \quad (26.5a)$$

where

$$F(x) = 1 - 8x + 8x^3 - x^4 - 12x^2 \ln x \quad (26.5b)$$

and

$$\alpha(m_\mu)^{-1} = \alpha^{-1} - \frac{2}{3\pi} \ln \left(\frac{m_\mu}{m_e} \right) + \frac{1}{6\pi} \approx 136. \quad (26.5c)$$

The uncertainty in G_F from the input quantities is 1.1×10^{-10} GeV $^{-2}$. The quoted uncertainty of 2×10^{-10} is dominated by second order radiative corrections, estimated from the magnitude of the known $\alpha^2 \ln(m_\mu/m_e)$ term to be $\sim 1.8 \times 10^{-10}$ (alternately, one can view Eq. (26.5) as the exact definition of G_F ; then the theoretical uncertainty appears instead in the formulae for quantities derived from G_F).

- (c) $\sin^2 \theta_W$, determined from the Z mass and other Z -pole observables, the W mass, and neutral-current processes [9]. The value of $\sin^2 \theta_W$ depends on the renormalization prescription. There are a number of popular schemes [10–15] leading to $\sin^2 \theta_W$ values which differ by small factors which depend on m_t and M_H . The notation for these schemes is shown in Table 26.1. Discussion of the schemes follows the table.

Table 26.1: Notations used to indicate the various schemes discussed in the text. Each definition of $\sin \theta_W$ leads to values that differ by small factors depending on m_t and M_H .

Scheme	Notation
On-shell	$s_W = \sin \theta_W$
NOV	$s_{M_Z} = \sin \theta_W$
$\overline{\text{MS}}$	$\hat{s}_Z = \sin \theta_W$
$\overline{\text{MS}} ND$	$\hat{s}_{ND} = \sin \theta_W$
Effective angle	$\bar{s}_f = \sin \theta_W$

- (i) The on-shell scheme promotes the tree-level formula $\sin^2 \theta_W = 1 - M_W^2/M_Z^2$ to a definition of the renormalized $\sin^2 \theta_W$ to all orders in perturbation theory, *i.e.*, $\sin^2 \theta_W \rightarrow s_W^2 \equiv 1 - M_W^2/M_Z^2$. This scheme is simple conceptually. However, M_W is known much less precisely than M_Z and in practice one extracts s_W^2 from M_Z alone using

$$M_W = \frac{A_0}{s_W(1 - \Delta r)^{1/2}} \quad (26.6a)$$

$$M_Z = \frac{M_W}{c_W}, \quad (26.6b)$$

where $s_W \equiv \sin \theta_W$, $c_W \equiv \cos \theta_W$, $A_0 = (\pi\alpha/\sqrt{2}G_F)^{1/2} = 37.2802$ GeV, and Δr includes the radiative corrections relating α , $\alpha(M_Z)$, G_F , M_W , and M_Z . One finds $\Delta r \sim \Delta r_0 - \rho_t/\tan^2 \theta_W$, where $\Delta r_0 \approx 1 - \alpha/\alpha(M_Z) \approx 0.07$ is due to the running of α and $\rho_t = 3G_F m_t^2/8\sqrt{2}\pi^2 \approx 0.0031$ ($m_t/100$ GeV)² represents the dominant (quadratic) m_t dependence. There are additional contributions to Δr from bosonic loops, including those which depend logarithmically on the Higgs mass M_H . For example, $\Delta r = 0.0475 \pm 0.0009$ for $(m_t, M_H) = (150, 300)$, while $\Delta r = 0.0327 \pm 0.0009$ for $(190, 300)$, where the 0.0009 uncertainty is from $\alpha(M_Z)$. Thus the value of s_W^2 extracted from M_Z includes a large uncertainty (~ 0.002) from the currently allowed range of m_t .

- (ii) A more precisely determined quantity $s_{M_Z}^2$ can be obtained from M_Z by removing the (m_t, M_H) dependent term from Δr [11], i.e.,

$$s_{M_Z}^2 c_{M_Z}^2 \equiv \frac{\pi \bar{\alpha}}{\sqrt{2} G_F M_Z^2}, \quad (26.7)$$

where $\bar{\alpha}^{-1} = \alpha(M_Z)^{-1} = 128.87 \pm 0.12$. [This is defined using the conventional QED renormalization, and differs by finite constants from $\hat{\alpha}(M_Z)^{-1}$.] This yields $s_{M_Z}^2 = 0.2312 \pm 0.0003$, with most of the uncertainty from $\bar{\alpha}$ rather than M_Z . Scheme (ii) is equivalent to using M_Z rather than $\sin^2 \theta_W$ as the third fundamental parameter. However, it recognizes that $s_{M_Z}^2$ is still a useful derived quantity. The small uncertainty in $s_{M_Z}^2$ compared to other schemes is because the m_t dependence has been removed by definition. However, the m_t uncertainty reemerges when other quantities (e.g., M_W or other Z -pole observables) are predicted in terms of M_Z . Both s_W^2 and $s_{M_Z}^2$ depend not only on the gauge couplings but also on the spontaneous-symmetry breaking, and both definitions are awkward in the presence of any extension of the Standard Model which perturbs the value of M_Z (or M_W). Other definitions are motivated by the tree-level coupling constant definition $\theta_W = \tan^{-1}(g'/g)$.

- (iii) In particular, the modified minimal subtraction ($\overline{\text{MS}}$) scheme introduces the quantity $\sin^2 \hat{\theta}_W(\mu) \equiv \hat{g}'^2(\mu)/[\hat{g}^2(\mu) + \hat{g}'^2(\mu)]$, where the couplings \hat{g} and \hat{g}' are defined by modified minimal subtraction and the scale μ is conveniently chosen to be M_Z for electroweak processes. The value of $\hat{s}_Z^2 = \sin^2 \hat{\theta}_W(M_Z)$ extracted from M_Z is less sensitive than s_W^2 to m_t (by a factor of $\tan^2 \theta_W$), and is less sensitive to most types of new physics than s_W^2 or $s_{M_Z}^2$. It is also very useful for comparing with the predictions of grand unification. There are actually several variant definitions of $\sin^2 \hat{\theta}_W(M_Z)$, differing according to whether or how finite $\alpha \ln(m_t/M_Z)$ terms are decoupled (subtracted from the couplings). One cannot entirely decouple the $\alpha \ln(m_t/M_Z)$ terms from all electroweak quantities because $m_t \gg m_b$ breaks SU(2) symmetry. The scheme that will be adopted here decouples the $\alpha \ln(m_t/M_Z)$ terms from the $\gamma - Z$ mixing [7,12], essentially eliminating any $\ln(m_t/M_Z)$ dependence in the formulae for asymmetries at the Z pole when written in terms of \hat{s}_Z^2 . The various definitions are related by

$$\hat{s}_Z^2 = c(m_t, M_H) s_W^2 = \bar{c}(m_t, M_H) s_{M_Z}^2, \quad (26.8)$$

where $c = 1.025$ (1.041) for $m_t = 150$ (190) GeV and $M_H = 300$ GeV. Similarly $\bar{c}(150, 300) = 1.006$ and $\bar{c}(190, 300) = 1.0003$. The quadratic m_t dependence is given by $c \sim 1 + \rho_t/\tan^2 \theta_W$. The expressions for M_W and M_Z in the $\overline{\text{MS}}$ scheme are

$$M_W = \frac{A_0}{\hat{s}_Z(1 - \Delta \hat{r}_W)^{1/2}} \quad (26.9a)$$

$$M_Z = \frac{M_W}{\hat{\rho}^{1/2} \hat{c}_Z}. \quad (26.9b)$$

One predicts $\Delta \hat{r}_W = 0.0703$ (0.0708) \pm 0.0009 for $m_t = 150$ (190) GeV and $M_H = 300$ GeV. $\Delta \hat{r}_W$ has no quadratic m_t dependence, because shifts in M_W are absorbed into the observed G_F , so that $\Delta \hat{r}_W$ is dominated by $\Delta r_0 = 1 - \alpha/\alpha(M_Z)$. Similarly, $\hat{\rho} \sim 1 + \rho_t$. Including bosonic loops, $\hat{\rho} = 1.0072$ (1.0118) for $m_t = 150$ (190) GeV.

- (iv) A variant $\overline{\text{MS}}$ quantity \hat{s}_{ND}^2 (used in the previous edition of this *Review*) does not decouple the $\alpha \ln(m_t/M_Z)$ terms [13]. It is related to \hat{s}_Z^2 by

$$\hat{s}_Z^2 = \hat{s}_{\text{ND}}^2 / \left(1 + \frac{\hat{\alpha}}{\pi} d\right) \quad (26.10a)$$

$$d = \frac{1}{3} \left(\frac{1}{\hat{s}^2} - \frac{8}{3} \right) \left[\left(1 + \frac{\hat{\alpha}_s}{\pi}\right) \ln \frac{m_t}{M_Z} - \frac{15\hat{\alpha}_s}{\pi} \right], \quad (26.10b)$$

where $\hat{\alpha}_s$ is the QCD coupling at M_Z . Thus, $\hat{s}_Z^2 - \hat{s}_{\text{ND}}^2 \sim -0.0001$ (-0.0002) for $m_t = 150$ (190) GeV and $M_H = 300$ GeV.

- (v) Yet another definition, the effective angle [14,15] \hat{s}_f^2 for Z coupling to fermion f , is described below.

Experiments are now at such a level of precision that complete $\mathcal{O}(\alpha)$ radiative corrections must be applied. These corrections are conveniently divided into two classes:

1. QED diagrams involving the emission of real photons or the exchange of virtual photons in loops, but not including vacuum polarization diagrams. These graphs yield finite and gauge-invariant contributions to observable processes. However, they are dependent on energies, experimental cuts, etc., and must be calculated individually for each experiment.
2. Electroweak corrections, including $\gamma\gamma$, γZ , ZZ , and WW vacuum polarization diagrams, as well as vertex corrections, box graphs, etc., involving virtual W 's and Z 's. Many of these corrections are absorbed into the renormalized Fermi constant defined in Eq. (26.5). Others modify the tree-level expressions for Z -pole observables and neutral-current amplitudes in several ways [9]. One-loop corrections are included for all processes. In addition, certain two-loop corrections are also important. In particular, two-loop corrections involving the top-quark [16] modify ρ_t in $\hat{\rho}$, Δr , and elsewhere by

$$\rho_t \rightarrow \rho_t [1 + R(M_H/m_t) \rho_t / 3], \quad (26.11)$$

where R is strongly dependent on M_H/m_t , with $R(0) = 19 - 2\pi^2$. Similarly, mixed $(\alpha\alpha_s m_t^2)$ QCD-electroweak loops [17] multiply ρ_t by $1 - \alpha_s(m_t)(\pi^2 + 3)/9\pi \sim 0.9$, increasing the predicted value of m_t by $\sim 5\%$. Analogous mixed terms modify the $Z \rightarrow b\bar{b}$ vertex [18]. Recently, there has been discussion of threshold or higher-order corrections to this term, which have been estimated by both dispersion relation and perturbative methods [19,20]. One estimate [20] suggests that $\alpha_s(m_t)$ should be replaced by $\alpha_s(0.15 m_t)$, raising the predicted m_t by ~ 3 GeV. However, these threshold effects are still uncertain and are therefore not included here.

26.2. Cross section and asymmetry formulas

It is convenient to write the four-fermion interactions relevant to ν -hadron, νe , and parity-violating e -hadron neutral-current processes in a form that is valid in an arbitrary gauge theory (assuming massless left-handed neutrinos). One has

$$-\mathcal{L}^{\nu\text{Hadron}} = \frac{G_F}{\sqrt{2}} \bar{\nu} \gamma^\mu (1 - \gamma^5) \nu \times \sum_i \left[\epsilon_L(i) \bar{q}_i \gamma_\mu (1 - \gamma^5) q_i + \epsilon_R(i) \bar{q}_i \gamma_\mu (1 + \gamma^5) q_i \right], \quad (26.12)$$

$$-\mathcal{L}^{\nu e} = \frac{G_F}{\sqrt{2}} \bar{\nu}_\mu \gamma^\mu (1 - \gamma^5) \nu_\mu \bar{e} \gamma_\mu (g_V^{\nu e} - g_A^{\nu e} \gamma^5) e \quad (26.13)$$

(for $\nu_e e$ or $\bar{\nu}_e e$, the charged-current contribution must be included), and

$$-\mathcal{L}^{\text{eHadron}} = -\frac{G_F}{\sqrt{2}} \times \sum_i \left[C_{1i} \bar{e} \gamma_\mu \gamma^5 e \bar{q}_i \gamma^\mu q_i + C_{2i} \bar{e} \gamma_\mu e \bar{q}_i \gamma^\mu \gamma^5 q_i \right] \quad (26.14)$$

(One must add the parity-conserving QED contribution.)

The Standard Model expressions for $\epsilon_{L,R}(i)$, $g_{V,A}^{\nu e}$, and C_{ij} are given in Table 26.2. Note that $g_{V,A}^{\nu e}$ and the other quantities are coefficients of effective four-fermi operators, which differ from the quantities defined in Eq. (26.2) and Eq. (26.3) in the radiative corrections and in the presence of possible physics beyond the Standard Model.

A precise determination of the on-shell s_W^2 , which depends only very weakly on m_t and M_H , is obtained from deep inelastic neutrino scattering from (approximately) isoscalar targets. The ratio $R_\nu \equiv \sigma_{\nu N}^{NC} / \sigma_{\nu N}^{CC}$ of neutral- to charged-current cross sections has been measured to 1% accuracy by the CDHS [21] and CHARM [22] collaborations [23,24] at CERN, and the CCFR collaboration at Fermilab [25] has recently reported an even more precise result, so it is important to obtain theoretical expressions for R_ν and $R_{\bar{\nu}} \equiv \sigma_{\bar{\nu} N}^{NC} / \sigma_{\bar{\nu} N}^{CC}$ (as functions of $\sin^2 \theta_W$) to comparable accuracy. Fortunately, most of the uncertainties from the strong interactions and neutrino spectra cancel in the ratio.

Table 26.2: Standard Model expressions for the neutral-current parameters for ν -hadron, νe , and e -hadron processes. If radiative corrections are ignored, $\rho = \kappa = 1$, $\lambda = 0$. At $\mathcal{O}(\alpha)$ in the on-shell scheme, $\rho_{\nu N}^{NC} = 1.0083$, $\kappa_{\nu N} = 1.0330$, $\lambda_{uL} = -0.0032$, $\lambda_{dL} = -0.0026$, and $\lambda_{uR} = 1/2 \lambda_{dR} = 3.6 \times 10^{-5}$ for $m_t = 169$ GeV, $M_H = 300$ GeV, $M_Z = 91.187$ GeV, and $\langle Q^2 \rangle = 20$ GeV². For νe scattering, $\kappa_{\nu e} = 1.0332$ and $\rho_{\nu e} = 1.0131$ (at $\langle Q^2 \rangle = 0$). For atomic parity violation, $\rho'_{eq} = 0.9874$ and $\kappa'_{eq} = 1.033$. For the SLAC polarized electron experiment, $\rho_{eq} = 0.978$, $\kappa_{eq} = 1.031$, $\rho_{eq} = 1.001$, and $\kappa_{eq} = 1.06$ after incorporating additional QED corrections, while $\lambda_{2u} = -0.013$, $\lambda_{2d} = 0.003$. The dominant m_t dependence is given by $\rho \sim 1 + \rho_t$, while $\kappa \sim 1 + \rho_t / \tan^2 \theta_W$ (on-shell) or $\kappa \sim 1(\overline{\text{MS}})$.

Quantity	Standard Model Expression
$\epsilon_L(u)$	$\rho_{\nu N}^{NC} \left(\frac{1}{2} - \frac{2}{3} \kappa_{\nu N} \sin^2 \theta_W + \lambda_{uL} \right)$
$\epsilon_L(d)$	$\rho_{\nu N}^{NC} \left(-\frac{1}{2} + \frac{1}{3} \kappa_{\nu N} \sin^2 \theta_W + \lambda_{dL} \right)$
$\epsilon_R(u)$	$\rho_{\nu N}^{NC} \left(-\frac{2}{3} \kappa_{\nu N} \sin^2 \theta_W + \lambda_{uR} \right)$
$\epsilon_R(d)$	$\rho_{\nu N}^{NC} \left(\frac{1}{3} \kappa_{\nu N} \sin^2 \theta_W + \lambda_{dR} \right)$
$g_V^{\nu e}$	$\rho_{\nu e} \left(-\frac{1}{2} + 2\kappa_{\nu e} \sin^2 \theta_W \right)$
$g_A^{\nu e}$	$\rho_{\nu e} \left(-\frac{1}{2} \right)$
C_{1u}	$\rho'_{eq} \left(-\frac{1}{2} + \frac{4}{3} \kappa'_{eq} \sin^2 \theta_W \right)$
C_{1d}	$\rho'_{eq} \left(\frac{1}{2} - \frac{2}{3} \kappa'_{eq} \sin^2 \theta_W \right)$
C_{2u}	$\rho_{eq} \left(-\frac{1}{2} + 2\kappa_{eq} \sin^2 \theta_W \right) + \lambda_{2u}$
C_{2d}	$\rho_{eq} \left(\frac{1}{2} - 2\kappa_{eq} \sin^2 \theta_W \right) + \lambda_{2d}$

A simple zeroth-order approximation is

$$R_\nu = g_L^2 + g_R^2 \quad (26.15a)$$

$$R_{\bar{\nu}} = g_L^2 + \frac{g_R^2}{r}, \quad (26.15b)$$

where

$$g_L^2 \equiv \epsilon_L(u)^2 + \epsilon_L(d)^2 \approx \frac{1}{2} - \sin^2 \theta_W + \frac{5}{9} \sin^4 \theta_W \quad (26.16a)$$

$$g_R^2 \equiv \epsilon_R(u)^2 + \epsilon_R(d)^2 \approx \frac{5}{9} \sin^4 \theta_W, \quad (26.16b)$$

and $r \equiv \sigma_{\bar{\nu} N}^{CC} / \sigma_{\nu N}^{CC}$ is the ratio of $\bar{\nu}$ and ν charged-current cross sections, which can be measured directly. [In the simple parton model, ignoring hadron energy cuts, $r \approx (\frac{1}{3} + \epsilon) / (1 + \frac{1}{3}\epsilon)$, where $\epsilon \sim 0.125$ is the ratio of the fraction of the nucleon's momentum carried by antiquarks to that carried by quarks.] In practice, Eq. (26.15) must be corrected for quark mixing, the s and c seas, c -quark threshold effects, nonisoscalar target effects, W - Z propagator differences, and radiative corrections (which lower the extracted value of $\sin^2 \theta_W$ by ~ 0.009). Details of the neutrino spectra, experimental cuts, x and Q^2 dependence of structure functions, and longitudinal structure functions enter only at the level of these corrections and therefore lead to very small uncertainties. The largest theoretical uncertainty is associated with the c threshold, which mainly affects σ^{CC} . Using the slow rescaling prescription [9] the central value of $\sin^2 \theta_W$ varies as $0.013 [m_c(\text{GeV}) - 1.3]$, where m_c is the effective mass. For $m_c = 1.31 \pm 0.24$ GeV (determined from ν -induced dimuon production [25]) this contributes ± 0.003 to the total theoretical uncertainty $\Delta \sin^2 \theta_W \sim \pm 0.004$. This would require a high-energy neutrino beam for improvement. (The experimental uncertainty is ± 0.003). The CCFR group quotes $s_W^2 = 0.2218 \pm 0.0059$ for $(m_t, M_H) = (150, 100)$, but this result is insensitive to (m_t, M_H) . Combining all of the precise deep-inelastic measurements, one obtains $s_W^2 = 0.2260 \pm 0.0048$ for (m_t, M_H) in the allowed range.

The laboratory cross section for $\nu_\mu e \rightarrow \nu_\mu e$ or $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ elastic scattering is

$$\frac{d\sigma_{\nu_\mu, \bar{\nu}_\mu}}{dy} = \frac{G_F^2 m_e E_\nu}{2\pi} \times \left[(g_V^{\nu e} \pm g_A^{\nu e})^2 + (g_V^{\nu e} \mp g_A^{\nu e})^2 (1-y)^2 - (g_V^{\nu e 2} - g_A^{\nu e 2}) \frac{y m_e}{E_\nu} \right], \quad (26.17)$$

where the upper (lower) sign refers to $\nu_\mu(\bar{\nu}_\mu)$, and $y \equiv E_e/E_\nu$ [which runs from 0 to $(1 + m_e/2E_\nu)^{-1}$] is the ratio of the kinetic energy of the recoil electron to the incident ν or $\bar{\nu}$ energy. For $E_\nu \gg m_e$ this yields a total cross section

$$\sigma = \frac{G_F^2 m_e E_\nu}{2\pi} \left[(g_V^{\nu e} \pm g_A^{\nu e})^2 + \frac{1}{3} (g_V^{\nu e} \mp g_A^{\nu e})^2 \right]. \quad (26.18)$$

The most accurate leptonic measurements [26–28] of $\sin^2 \theta_W$ are from the ratio $R \equiv \sigma_{\nu_\mu e} / \sigma_{\bar{\nu}_\mu e}$ in which many of the systematic uncertainties cancel. Radiative corrections (other than m_t effects) are small compared to the precision of present experiments and have negligible effect on the extracted $\sin^2 \theta_W$. The most precise (CHARM II) experiment [28] determined not only $\sin^2 \theta_W$ but $g_{V,A}^{\nu e}$ as well. The cross sections for $\nu_e e$ and $\bar{\nu}_e e$ may be obtained from Eq. (26.17) by replacing $g_{V,A}^{\nu e}$ by $g_{V,A}^{\nu e} + 1$, where the 1 is due to the charged-current contribution.

The SLAC polarized-electron experiment [29] measured the parity-violating asymmetry

$$A = \frac{\sigma_R - \sigma_L}{\sigma_R + \sigma_L}, \quad (26.19)$$

where $\sigma_{R,L}$ is the cross section for the deep-inelastic scattering of a right- or left-handed electron: $e_{R,L} N \rightarrow eX$. In the quark parton model

$$\frac{A}{Q^2} = a_1 + a_2 \frac{1 - (1-y)^2}{1 + (1-y)^2}, \quad (26.20)$$

where $Q^2 > 0$ is the momentum transfer and y is the fractional energy transfer from the electron to the hadrons. For the deuteron or other isoscalar target, one has, neglecting the s quark and antiquarks,

$$a_1 = \frac{3G_F}{5\sqrt{2}\pi\alpha} \left(C_{1u} - \frac{1}{2}C_{1d} \right) \approx \frac{3G_F}{5\sqrt{2}\pi\alpha} \left(-\frac{3}{4} + \frac{5}{3}\sin^2\theta_W \right) \quad (26.21a)$$

$$a_2 = \frac{3G_F}{5\sqrt{2}\pi\alpha} \left(C_{2u} - \frac{1}{2}C_{2d} \right) \approx \frac{9G_F}{5\sqrt{2}\pi\alpha} \left(\sin^2\theta_W - \frac{1}{4} \right). \quad (26.21b)$$

Radiative corrections (other than m_t effects) lower the extracted value of $\sin^2\theta_W$ by ~ 0.005 .

Experiments measuring atomic parity violation [30] are now quite precise, and the uncertainties associated with atomic wave functions are relatively small (especially for cesium, for which the theoretical uncertainty is $\sim 1\%$ [31]). For heavy atoms one determines the ‘‘weak charge’’

$$Q_W = -2[C_{1u}(2Z+N) + C_{1d}(Z+2N)] \\ \approx Z(1 - 4\sin^2\theta_W) - N. \quad (26.22)$$

Radiative corrections increase the extracted $\sin^2\theta_W$ by ~ 0.008 .

The forward-backward asymmetry for $e^+e^- \rightarrow \ell\bar{\ell}$, $\ell = \mu$ or τ , is defined as

$$A_{FB} \equiv \frac{\sigma_F - \sigma_B}{\sigma_F + \sigma_B}, \quad (26.23)$$

where $\sigma_F(\sigma_B)$ is the cross section for ℓ^- to travel forward (backward) with respect to the e^- direction. A_{FB} and R , the total cross section relative to pure QED, are given by

$$R = F_1 \quad (26.24)$$

$$A_{FB} = 3F_2/4F_1, \quad (26.25)$$

where

$$F_1 = 1 - 2\chi_0 g_V^e g_V^f \cos\delta_R + \chi_0^2 (g_V^{e2} + g_A^{e2}) (g_V^{f2} + g_A^{f2}) \quad (26.26a)$$

$$F_2 = -2\chi_0 g_A^e g_A^f \cos\delta_R + 4\chi_0^2 g_A^e g_A^f g_V^e g_V^f, \quad (26.26b)$$

where

$$\tan\delta_R = \frac{M_Z \Gamma_Z}{M_Z^2 - s} \quad (26.27)$$

$$\chi_0 = \frac{G_F}{2\sqrt{2}\pi\alpha} \frac{sM_Z^2}{[(M_Z^2 - s)^2 + M_Z^2 \Gamma_Z^2]^{1/2}} \quad (26.28)$$

and \sqrt{s} is the CM energy. Eq. (26.26) is valid at tree level. If the data are radiatively corrected for QED effects (as described above), then the remaining electroweak corrections can be incorporated [32] (in an approximation adequate for existing PEP, PETRA, and TRISTAN data, which are well below the Z pole) by replacing χ_0 by $\chi(s) \equiv (1 + \rho_t)\chi_0(s)\alpha/\hat{\alpha}(s)$, where $\hat{\alpha}(s)$ is the running QED coupling, and evaluating g_V in the \overline{MS} scheme. Formulas for $e^+e^- \rightarrow hadrons$ may be found in Ref. 33.

At LEP and SLC, there are high-precision measurements of various Z -pole observables. These include the Z mass and total width Γ_Z , and partial widths $\Gamma(f\bar{f})$ for $Z \rightarrow f\bar{f}$ for fermion f ($f = e, \mu, \tau$, hadrons, b, c , and ν). The data is consistent with lepton-family universality $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-) = \Gamma(\tau^+\tau^-)$, so one may work with an average width $\Gamma(\ell\bar{\ell})$. It is convenient to use the variables $M_Z, \Gamma_Z, R \equiv \Gamma(had)/\Gamma(\ell\bar{\ell})$, $\sigma_{had} \equiv 12\pi\Gamma(e^+e^-)\Gamma(had)/M_Z^2\Gamma_Z^2$, $R_b \equiv \Gamma(b\bar{b})/\Gamma(had)$, and $R_c \equiv \Gamma(c\bar{c})/\Gamma(had)$, which are weakly correlated experimentally. ($\Gamma(had)$ is the partial width into hadrons.) R is insensitive to m_t (except for $Z \rightarrow b\bar{b}$ vertex corrections) and is especially useful for constraining α_s . The width for invisible decays, $\Gamma(inv) = \Gamma_Z - 3\Gamma(\ell\bar{\ell}) - \Gamma(had) = 498.2 \pm 4.2$ MeV, can be used to determine the number of neutrino flavors lighter than $M_Z/2$, $N_\nu = \Gamma_{inv}/\Gamma(\nu\bar{\nu}) \sim 2.98 \pm 0.03$.

There are also measurements of various asymmetries. These include the polarization or left-right asymmetry

$$A_{LR} \equiv \frac{\sigma_L - \sigma_R}{\sigma_L + \sigma_R}, \quad (26.29)$$

where $\sigma_L(\sigma_R)$ is the cross section for a left- (right)-handed incident electron. A_{LR} has been measured precisely by the SLD collaboration at SLC [34] and has the advantages of being extremely sensitive to $\sin^2\theta_W$ and insensitive to QED radiative corrections. Other asymmetries are the forward-backward asymmetries $A_{FB}^{(0,f)}$ for $f = e, \mu, \tau, b, c$ ($A_{FB}^{(0,e)}, A_{FB}^{(0,\mu)}, A_{FB}^{(0,\tau)}$ are consistent with lepton-family universality, allowing an average value $A_{FB}^{(0,\ell)}$), the jet-charge asymmetry, the τ polarization P_τ , and its angular distribution. Further details, including references to the data from the LEP experiments (ALEPH, DELPHI, L3, OPAL) may be found in the Full Listings in the ‘‘Note on the Z Boson.’’ At tree level and neglecting QED effects and terms of order $(\Gamma_Z/M_Z)^2$, one has

$$A_{FB}^{(0,f)} \approx \frac{3}{4} A_f \frac{A_e + P_e}{1 + P_e A_e} \quad (26.30)$$

$$A_{LR} \approx A_e P_e, \quad (26.31)$$

where P_e is the initial e^- polarization and

$$A_f \equiv \frac{2g_V^f g_A^f}{g_V^{f2} + g_A^{f2}}. \quad (26.32)$$

Similarly, A_τ is given by the negative total τ polarization, and A_e can be extracted from the angular distribution of the polarization.

It has become customary for the experimental groups to present corrected asymmetries A^0 , in which photon exchange and γ - Z interference, QED corrections, and corrections for $\sqrt{s} \neq M_Z$ are removed from the data, leaving the pure electroweak asymmetries. Ignoring negligible electroweak boxes, these corrected asymmetries are expressed using effective tree-level expression *e.g.*, $A_{FB}^{(0,f)} = \frac{3}{4} \bar{A}_f \bar{A}_e$ (for $P_e = 0$) and $A_{LR}^0 = \bar{A}_e$, where

$$\bar{A}_f = \frac{2\bar{g}_V^f \bar{g}_A^f}{\bar{g}_V^{f2} + \bar{g}_A^{f2}}, \quad (26.33a)$$

and

$$\bar{g}_V^f = \sqrt{\rho_f} (t_{3L}^{(f)} - 2q_f \kappa_f \sin^2\theta_W) \quad (26.33b)$$

$$\bar{g}_A^f = \sqrt{\rho_f} t_{3L}^{(f)}. \quad (26.33c)$$

The electroweak-radiative corrections have been absorbed into corrections $\rho_f - 1$ and $\kappa_f - 1$, which depend on the fermion f and on the renormalization scheme. In the on-shell scheme, the quadratic m_t dependence is given by $\rho_f \sim 1 + \rho_t$, $\kappa_f \sim \kappa_f^{os} \sim 1 + \rho_t/\tan^2\theta_W$, while in \overline{MS} , $\rho_f \sim \hat{\rho}$, $\kappa_f \equiv \hat{\kappa}_f \sim 1$. In practice, additional bosonic loops, vertex corrections, *etc.*, must be included. For example, in the \overline{MS} scheme one has, for $(m_t, M_H) = (169, 300)$, $\rho_t = 1.0045$ and $\hat{\kappa}_t = 1.0009$. It is convenient to define an effective angle $\bar{s}_t^2 \equiv \sin^2\bar{\theta}_W^f \equiv \hat{\kappa}_f \hat{s}_t^2 = \kappa_f^{os} s_W^2$, in terms of which \bar{g}_V^f and \bar{g}_A^f are given by $\sqrt{\rho_f}$ times their tree-level formulae. Because \bar{g}_V^f is very small, not only $A_{LR}^0, A_{FB}^{(0,\ell)}$, and P_τ^0 , but also $A_{FB}^{(0,b)}, A_{FB}^{(0,c)}$, and the jet-charge asymmetry are mainly sensitive to \bar{s}_t^2 . One finds that $\hat{\kappa}_f$ is almost independent of (m_t, M_H) , so that

$$\bar{s}_t^2 \sim \hat{s}_t^2 + 0.0002 \quad (26.34)$$

using Ref. 15, or $\bar{s}_t^2 \sim \hat{s}_t^2 + 0.00028$ from Ref. 14 (the small difference is an indication of theoretical uncertainties from higher-order terms, *etc.*). In any case, the asymmetries determine values of \bar{s}_t^2 and \hat{s}_t^2 almost independent of m_t , while the κ 's for the other schemes are m_t dependent.

26.3. W and Z decays

The partial decay width for gauge bosons to decay into massless fermions $f_1\bar{f}_2$ is

$$\Gamma(W^+ \rightarrow e^+\nu_e) = \frac{G_F M_W^3}{6\sqrt{2}\pi} \approx 227 \pm 1 \text{ MeV} \quad (26.35a)$$

$$\Gamma(W^+ \rightarrow u_i\bar{d}_i) = \frac{CG_F M_W^3}{6\sqrt{2}\pi} |V_{ij}|^2 \approx (707 \pm 3) |V_{ij}|^2 \text{ MeV} \quad (26.35b)$$

$$\Gamma(Z \rightarrow \psi_i\bar{\psi}_i) = \frac{CG_F M_Z^3}{6\sqrt{2}\pi} [g_V^2 + g_A^2] \quad (26.35c)$$

$$\approx \begin{cases} 167.1 \pm 0.3 \text{ MeV} (\nu\bar{\nu}), & 83.9 \pm 0.2 \text{ MeV} (e^+e^-), \\ 298.0 \pm 0.6 \text{ MeV} (u\bar{u}), & 384.5 \pm 0.8 \text{ MeV} (d\bar{d}), \\ 375.2 \mp 0.4 \text{ MeV} (b\bar{b}). \end{cases}$$

For leptons $C = 1$, while for quarks $C = 3 \left(1 + \alpha_s(M_V)/\pi + 1.409\alpha_s^2/\pi^2 - 12.77\alpha_s^3/\pi^3\right)$, where the 3 is due to color and the factor in parentheses is a QCD correction. Corrections to Eq. (26.35) for massive fermions are given in Refs. 10 and 35 and the mass/Yukawa effects in the QCD corrections in Refs. [18,36]. Expressing the widths in terms of $G_F M_{W,Z}^3$ incorporates the bulk of the low-energy radiative corrections [10,35]. The $Z \rightarrow f\bar{f}$ widths have an additional QED correction $1 + 3\alpha g_f^2/4\pi$. The electroweak corrections are incorporated by replacing $g_{V,A}^2$ by $\bar{g}_{V,A}^2$. Hence, the widths are proportional to $\rho_i \sim 1 + \rho_t$. There is additional (negative) quadratic m_t dependence in the $Z \rightarrow b\bar{b}$ vertex corrections [37] which causes $\Gamma(b\bar{b})$ to decrease with m_t . The dominant effect is to multiply $\Gamma(b\bar{b})$ by the vertex correction $1 + \delta\rho_{b\bar{b}}$, where $\delta\rho_{b\bar{b}} \sim 10^{-2} \left(-\frac{1}{2} \frac{m_t^2}{M_Z^2} + \frac{1}{5}\right)$. In practice, the corrections are included in ρ_b and κ_b .

For 3-fermion families the total widths are predicted to be

$$\Gamma_Z \approx 2.493 \pm 0.004 \text{ GeV} \quad (26.36)$$

$$\Gamma_W \approx 2.09 \pm 0.01 \text{ GeV} \quad (26.37)$$

The numerical values for the widths assume $M_Z = 91.187 \pm 0.007$ GeV, $M_W = 80.29 \pm 0.11$ GeV, $\alpha_s = 0.120$, and $m_t = 169_{-18}^{+16}$ GeV, where the M_W , α_s , and m_t values are predicted by the global fits for $M_H = 300$ GeV. The uncertainties for Γ_W and Γ_Z are dominated by ΔM_W and Δm_t , respectively. The uncertainty in α_s , ± 0.007 , introduces an additional uncertainty of 0.2% in the hadronic widths, corresponding to ± 4 MeV in Γ_Z .

These predictions are to be compared with the experimental results $\Gamma_Z = 2.490 \pm 0.007$ GeV and $\Gamma_W = 2.08 \pm 0.07$ GeV.

26.4. Experimental results

The values of the principal Z -pole observables are listed in Table 26.3, along with the Standard Model predictions for $M_Z = 91.187 \pm 0.007$, $m_t = 169_{-18}^{+16}$ GeV (for $M_H = 300$ GeV), 60 GeV $< M_H < 1$ TeV, and $\alpha_s = 0.120 \pm 0.007$. The values and predictions of M_W , M_W/M_Z , the Q_W for cesium [30,31], and recent results from deep inelastic and $\nu_e e$ scattering are also listed. The agreement is generally excellent, although $\Gamma_b = \Gamma(b\bar{b})/\Gamma(\text{had})$ is $\sim 1.8\sigma$ above the Standard Model prediction, and the left-right A_{LR}^0 is 2σ above the Standard Model prediction. There is also an experimental difference of $\sim 2\sigma$ between the SLD value of $A_e^0 = A_{LR}^0$ and the LEP value $A_{e\text{LEP}}^0 \sim 0.1434 \pm 0.0073$ obtained from $A_{FB}^{(0,e)}$, $A_e^0(P_\tau)$, $A_\tau^0(P_\tau)$ assuming lepton family universality. The observables in Table 26.3 (including correlations on the LEP observables), as well as all low-energy neutral-current data [9], are used in the global fits described below. The parameter $\sin^2\theta_W$ can be determined from the Z -pole observables and M_W , and from a variety of neutral-current processes spanning a very wide Q^2 range. The results [9], shown in Table 26.4, are in impressive agreement with each other, indicating the quantitative success of the Standard Model. The one discrepancy is the value $\hat{s}_Z^2 = 0.2292 \pm 0.0010$ from A_{LR}^0 which is 2.4σ below the




Figure 26.1: One-standard-deviation uncertainties in $\sin^2\hat{\theta}_W$ as a function of m_t , the direct constraint $m_t > 131$ GeV, the CDF range 174 ± 16 GeV, and the 90% CL region in $\sin^2\hat{\theta}_W - m_t$ allowed by all data, assuming $M_H = 300$ GeV.

value (0.2319 ± 0.005) from the global fit to all data and 2.8σ below the value 0.2323 ± 0.0005 obtained from all data other than A_{LR}^0 .

The data allow a simultaneous determination of $\sin^2\theta_W$, m_t , and the strong coupling $\alpha_s(M_Z)$. The latter is determined mainly from Γ_Z and R , and is almost uncorrelated from the other variables. The global fit to all indirect data yields

$$\hat{s}_Z^2 = 0.2319 \pm 0.0005 \pm 0.0002$$

$$m_t = 169_{-18-20}^{+16+17} \text{ GeV}$$

$$\alpha_s(M_Z) = 0.120 \pm 0.007 \pm 0.002, \quad (26.38)$$

where the central values are for a Higgs mass of 300 GeV, and the second error bars are for $M_H \rightarrow 1000(+)$ or $60(-)$ GeV. In all fits, the errors include full statistical, systematic, and theoretical uncertainties. The \hat{s}_Z^2 error is dominated by m_t , and \hat{s}_Z^2 and m_t have a strong negative correlation of ~ -0.85 . In the on-shell scheme one has $s_W^2 = 0.2247 \pm 0.0019$, the larger error due to the stronger sensitivity to m_t . The extracted value of α_s is based on a formula which has almost no theoretical uncertainty, and is in excellent agreement with the values 0.121 ± 0.006 from jet-event shapes, at the Z pole, and 0.116 ± 0.005 extracted from low-energy and jet data (see the section on Quantum Chromodynamics). Including the latter value as a separate constraint in the fits has negligible effect on $\sin^2\theta_W$ and m_t .

The value of m_t predicted by the precision data is in remarkable agreement with the value $m_t = 174 \pm 16$ GeV suggested by the CDF candidate top events. (The indirect prediction is for the pole mass, which should correspond approximately to the kinematic mass extracted from the CDF events.) One can carry out a combined fit of the indirect and direct (CDF) data, with the result $m_t = 172_{-12-9}^{+11+8}$ GeV, with little change in the $\sin^2\theta_W$ and α_s values. The results of fits to various combinations of the data are shown in Table 26.5 and the relation between \hat{s}_Z^2 and m_t for various observables in Fig. 26.1.

The data indicate a preference for a small Higgs mass. This is because there is a strong correlation between the quadratic ρ_t terms and logarithmic M_H effects in all of the indirect data except the $Z \rightarrow b\bar{b}$ vertex. The latter favor a smaller m_t and therefore a smaller M_H . The difference in χ^2 for the global fit is $\Delta\chi^2 = \chi^2(M_H = 1000 \text{ GeV}) - \chi^2(M_H = 60 \text{ GeV}) = 2.9$. However, this sensitivity is reduced to $\Delta\chi^2 \sim 2.1$ when the direct CDF value of m_t is included. Hence, the data favor a small value of M_H , as in supersymmetric extensions of the Standard Model, and m_t on the lower side of the allowed range; but the effect is marginal statistically:

including the direct constraint $M_H \geq 60$ GeV, the best fit is for $M_H = 60$ GeV, with the weak limit $M_H < 840$ GeV at 90% CL. The value of $\hat{s}_Z^2 = 0.2319 \pm 0.0005$ is in striking agreement with the prediction 0.233 ± 0.003 of grand unified theories based on the minimal supersymmetric extension of the Standard Model, but disagrees with the prediction 0.210 ± 0.003 of nonsupersymmetric unified theories.

One can also determine the radiative correction parameters Δr : including the CDF data, one obtains $\Delta r = 0.040 \pm 0.004$ and $\Delta \hat{r}_W = 0.068 \pm 0.003$, where the error includes m_t and M_H , in excellent agreement with the predictions 0.041 ± 0.007 and 0.0705 ± 0.0003 .

Table 26.3: Principal LEP and other recent observables, compared with the Standard Model predictions for $M_Z = 91.187 \pm 0.007$ GeV, $60 \text{ GeV} < M_H < 1 \text{ TeV}$, the global best fit value $m_t = 169_{-18}^{+16}$ GeV (for $M_H = 300$ GeV), and $\alpha_s = 0.012 \pm 0.007$. The LEP averages [38] of the ALEPH, DELPHI, L3, and OPAL results include common systematic errors and correlations [38]. $\bar{\alpha}_\ell(A_{FB}^{(0,q)})$ is the effective angle extracted from the quark-charge asymmetry. A_{LR}^0 includes both 1992 and 1993 data. The more accurate 1993 run only yields 0.1656 ± 0.0076 . In the fits, the values of $A_e^0(P_\tau)$ and A_{LR}^0 are combined to give 0.161 ± 0.012 , including a scale factor of 1.7. The values of $\Gamma(\ell\bar{\ell})$, $\Gamma(\text{had})$, and $\Gamma(\text{inv})$ are not independent of Γ_Z , R , and σ_{had} . The M_W and M_W/M_Z values are from the PDG fit. In the fits shown here the (uncorrelated) values from the individual experiments are used. The two values of s_W^2 from deep-inelastic scattering are from CCFR [25] and the global average, respectively. The $g_{V,A}^{\nu e}$ are from CHARM II [28]. The second error in Q_W (for cesium) is theoretical [31]. Older low-energy results are not listed but are included in the fits. In the Standard Model predictions, the first uncertainty is from M_Z and Δr , while the second is from m_t and M_H . The $\Delta\alpha_s = 0.007$ uncertainty leads to additional errors of 0.004 (Γ_Z), 0.05 (R), 0.04 (σ), 3.7 ($\Gamma(\text{had})$).

Quantity	Value	Standard Model
M_Z (GeV)	91.187 ± 0.007	input
Γ_Z (GeV)	2.490 ± 0.007	$2.493 \pm 0.001 \pm 0.004$
R	20.76 ± 0.05	$20.74 \pm 0.01 \pm 0.005$
$\sigma_{\text{had}}(\text{nb})$	41.55 ± 0.14	$41.48 \pm 0.01 \pm 0.01$
R_b	0.2210 ± 0.0029	$0.2157 \pm 0 \pm 0.0004$
R_c	0.171 ± 0.020	$0.171 \pm 0 \pm 0$
$A_{FB}^{(0,\ell)}$	0.0159 ± 0.0018	$0.0151 \pm 0.0005 \pm 0.0010$
$A_\tau^0(P_\tau)$	0.141 ± 0.021	$0.142 \pm 0.003 \pm 0.005$
$A_e^0(P_\tau)$	0.127 ± 0.025	$0.142 \pm 0.003 \pm 0.005$
$A_{FB}^{(0,b)}$	0.107 ± 0.013	$0.0995 \pm 0.002 \pm 0.003$
$A_{FB}^{(0,c)}$	0.058 ± 0.022	$0.071 \pm 0.002 \pm 0.003$
A_{LR}^0	0.1637 ± 0.0075	$0.142 \pm 0.003 \pm 0.005$
$\bar{\alpha}_\ell^2(A_{FB}^{(0,q)})$	0.2320 ± 0.0016	$0.2322 \pm 0.0003 \pm 0.0006$
$\Gamma(\ell\bar{\ell})$ (MeV)	83.84 ± 0.27	$83.90 \pm 0.02 \pm 0.16$
$\Gamma(\text{had})$ (MeV)	1740.7 ± 5.9	$1739.8 \pm 1 \pm 3$
$\Gamma(\text{inv})$ (MeV)	498.2 ± 4.2	$501.3 \pm 0.4 \pm 0.8$
M_W (GeV)	80.22 ± 0.26	$80.29 \pm 0.02 \pm 0.11$
M_W/M_Z	0.8798 ± 0.0028	$0.8805 \pm 0.0002 \pm 0.001$
Q_W	$-71.04 \pm 1.58 \pm 0.88$	$-72.92 \pm 0.07 \pm 0.07$
$s_W^2 = 1 - \frac{M_W^2}{M_Z^2}$	0.2218 ± 0.0059 0.2260 ± 0.0048	$0.2247 \pm 0.0003 \pm 0.0021$
$g_A^{\nu e}$	-0.503 ± 0.018	$-0.506 \pm 0 \pm 0.001$
$g_V^{\nu e}$	-0.025 ± 0.019	$-0.037 \pm 0 \pm 0.001$

Table 26.4: Values obtained for s_W^2 (on-shell) and $\hat{s}_Z^2(\overline{\text{MS}})$ from various reactions assuming the global best fit value $m_t = 169_{-18}^{+16}$ GeV (for $M_H = 300$ GeV), and $\alpha_s = 0.120 \pm 0.007$. The uncertainties include the effect of $60 \text{ GeV} < M_H < 1 \text{ TeV}$.

Reaction	s_W^2	\hat{s}_Z^2
M_Z	0.2247 ± 0.0021	0.2320 ± 0.0006
$M_W, M_W/M_Z$	0.2264 ± 0.0025	0.2338 ± 0.0022
$\Gamma_Z, R, \sigma_{\text{had}}$	0.2250 ± 0.0018	0.2322 ± 0.0006
$A_{FB}^{(0,\ell)}$	0.2243 ± 0.0018	0.2315 ± 0.0011
LEP asymmetries	0.2245 ± 0.0017	0.2317 ± 0.0008
A_{LR}^0	0.2221 ± 0.0017	0.2292 ± 0.0010
Deep inelastic (isocalar)	0.2260 ± 0.0048	0.233 ± 0.005
$\nu_\mu(\bar{\nu}_\mu)p \rightarrow \nu_\mu(\bar{\nu}_\mu)p$	0.205 ± 0.031	0.212 ± 0.032
$\nu_\mu(\bar{\nu}_\mu)e \rightarrow \nu_\mu(\bar{\nu}_\mu)e$	0.224 ± 0.009	0.231 ± 0.009
atomic parity violation	0.216 ± 0.008	0.223 ± 0.008
SLAC eD	0.216 ± 0.017	0.223 ± 0.018
All data	0.2247 ± 0.0019	0.2319 ± 0.0005

Table 26.5: Values of \hat{s}_Z^2 and s_W^2 (in parentheses), α_s , and m_t for various combinations of observables. The central values are for $M_H = 300$ GeV, and the second set of errors is for $M_H \rightarrow 1000(+), 60(-)$.

Data	\hat{s}_Z^2 (s_W^2)	α_s (M_Z)	m_t (GeV)
All indirect	$0.2319(5)(2)$ (0.2247 ± 0.0019)	$0.120(7)(2)$	169_{-18-20}^{+16+17}
Indirect + CDF direct	$0.2319(4)(3)$ (0.2244 ± 0.0013)	$0.120(6)(2)$	172_{-12-9}^{+11+8}
All LEP	$0.2322(6)(2)$ (0.2256 ± 0.0023)	$0.121(7)(2)$	163_{-22-20}^{+19+17}
SLD + M_Z	$0.2294(9)(1)$ (0.2154 ± 0.0032)	—	238_{-23-20}^{+20+17}
Z pole (LEP + SLD)	$0.2318(5)(2)$ (0.2241 ± 0.0021)	$0.120(7)(2)$	175_{-19-20}^{+18+17}

26.5. Deviations from the Standard Model

The Z pole, W mass, and neutral-current data can be used to search for and set limits on deviations from the Standard Model. For example, the relation between M_W and M_Z is modified if there are Higgs multiplets with weak isospin $> 1/2$ with significant vacuum expectation values. In order to calculate to higher orders in such theories one must define a set of four fundamental renormalized parameters. It is convenient to take these as α , G_F , M_Z , and M_W , since M_W and M_Z are directly measurable. Then \hat{s}_Z^2 and ρ_0 can be considered dependent parameters defined by

$$\hat{s}_Z^2 \equiv A_0^2/M_W^2(1 - \Delta \hat{r}_W) \quad (26.39)$$

and

$$\rho_0 \equiv M_W^2/(M_Z^2 \hat{c}_Z^2 \hat{\rho}). \quad (26.40)$$

Provided that the new physics which yields $\rho_0 \neq 1$ is a small perturbation which does not significantly affect the radiative corrections, ρ_0 can be regarded as a phenomenological parameter which multiplies G_F in Eqs. (26.12)–(26.14), (26.28), and Γ_Z in

Eq. (26.35). (Also, the expression for M_Z is divided by $\sqrt{\rho_0}$; the M_W formula is unchanged.) There is now enough data to determine ρ_0 , $\sin^2 \theta_W$, m_t , and α_s simultaneously. In particular, R_b and the CDF events yield m_t independent of ρ_0 , the asymmetries yield \hat{s}_Z^2 , R gives α_s , and M_Z and the widths constrain ρ_0 . From the global fit (including CDF),

$$\rho_0 = 1.0004 \pm 0.0022 \pm 0.002 \quad (26.41)$$

$$\hat{s}_Z^2 = 0.2318 \pm 0.0005 \quad (26.42)$$

$$\alpha_s = 0.120 \pm 0.007 \quad (26.43)$$

$$m_t = 170_{-15}^{+16}, \quad (26.44)$$

where the second error on ρ_0 is from M_H (the other parameters are insensitive to M_H). This is in remarkable agreement with the Standard Model expectation $\rho_0 = 1$, and constrains any higher-dimensional Higgs representation to have vacuum expectation values of less than a few percent of those of the doublets. The allowed regions in the $\rho_0 - \hat{s}_Z^2$ plane are shown in Fig. 26.2.




Figure 26.2: The allowed regions in $\sin^2 \hat{\theta}_W - \rho_0$ at 90% CL. m_t is a free parameter and $M_H = 300$ GeV is assumed. (Varying M_H in the range 60–1000 GeV moves the contour down or up by 0.002.)

Most of the parameters relevant to ν -hadron, νe , e -hadron, and e^+e^- processes are determined uniquely and precisely from the data in “model independent” fits (*i.e.*, fits which allow for an arbitrary electroweak gauge theory). The values for the parameters defined in Eqs. (26.12)–(26.14) are given in Table 26.6 along with the predictions of the Standard Model. The agreement is excellent. The low-energy e^+e^- results are difficult to present in a model-independent way because Z -propagator effects are non-negligible at TRISTAN, PETRA, and PEP energies. However, assuming e - μ - τ universality, the lepton asymmetries imply [33] $4(g_A^{\nu e})^2 = 0.99 \pm 0.05$, in good agreement with the Standard Model prediction $\simeq 1$. The much more precisely measured Z -pole parameters in Table 26.3 are in excellent agreement with the Standard Model.

* Constraints on V are discussed in the section on the Cabibbo-Kobayashi-Maskawa mixing matrix.

References:

1. S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967);
A. Salam in *Elementary Particle Theory*, ed. N. Svartholm (Almqvist and Wiksells, Stockholm, 1969) p. 367;

Table 26.6: Values of the model-independent neutral-current parameters, compared with the Standard Model prediction using $M_Z = 91.187$ GeV for $m_t = 169_{-18}^{+16}$ GeV and $M_H = 300$ GeV. There is a second $g_{V,A}^{\nu e}$ solution, given approximately by $g_V^{\nu e} \leftrightarrow g_A^{\nu e}$, which is eliminated by e^+e^- data under the assumption that the neutral current is dominated by the exchange of a single Z . θ_i , $i = L$ or R , is defined as $\tan^{-1}[\epsilon_i(u)/\epsilon_i(d)]$.

Quantity	Experimental Value	Standard Model Prediction	Correlation
$\epsilon_L(u)$	0.332 ± 0.016	0.345 ± 0.001	
$\epsilon_L(d)$	-0.438 ± 0.012	-0.429 ± 0.001	non-
$\epsilon_R(u)$	-0.178 ± 0.013	-0.155	Gaussian
$\epsilon_R(d)$	$-0.026 \begin{smallmatrix} +0.075 \\ -0.048 \end{smallmatrix}$	0.078	
g_L^2	0.3017 ± 0.0033	0.303 ± 0.001	
g_R^2	0.0326 ± 0.0033	0.030	small
θ_L	2.50 ± 0.035	2.46	
θ_R	$4.58 \begin{smallmatrix} +0.46 \\ -0.28 \end{smallmatrix}$	5.18	
$g_A^{\nu e}$	-0.506 ± 0.015	-0.506 ± 0.001	-0.04
$g_V^{\nu e}$	-0.039 ± 0.017	-0.037 ± 0.001	
C_{1u}	-0.214 ± 0.046	-0.189 ± 0.001	-0.995 -0.79
C_{1d}	0.359 ± 0.041	0.341 ± 0.002	0.79
$C_{2u} - \frac{1}{2}C_{2d}$	-0.04 ± 0.13	-0.051 ± 0.002	

S.L. Glashow, J. Iliopoulos, and L. Maiani, Phys. Rev. **D2**, 1285 (1970).

2. For reviews, see G. Barbiellini and C. Santoni, Riv. Nuovo Cimento **9(2)**, 1 (1986);
E.D. Commins and P.H. Bucksbaum, *Weak Interactions of Leptons and Quarks* (Cambridge Univ. Press, Cambridge, 1983).
3. C.A. Dominguez and E. de Rafael, Ann. Phys. **174**, 372 (1987);
J. Gasser and H. Leutwyler, Phys. Reports **87**, 77 (1982);
S. Narison, Phys. Lett. **B216**, 191 (1989);
J.F. Donoghue, Ann. Rev. Nucl. and Part. Sci. **39**, 1 (1989).
4. DØ: S. Abachi *et al.*, Phys. Rev. Lett. **72**, 2138 (1994).
5. CDF: F. Abe *et al.*, Fermilab-Pub-94/097-E.
6. For reviews, see J. Gunion *et al.*, “*The Higgs Hunter’s Guide*,” (Addison-Wesley, Redwood City, 1990);
M. Sher, Phys. Reports **179**, 273 (1989).
7. S. Fanchiotti, B. Kniehl, and A. Sirlin, Phys. Rev. **D48**, 307 (1993) and references therein.
8. W.J. Marciano and A. Sirlin, Phys. Rev. Lett. **61**, 1815 (1988).
9. The results given here are updated from U. Amaldi *et al.*, Phys. Rev. **D36**, 1385 (1987);
P. Langacker and M. Luo, Phys. Rev. **D44**, 817 (1991);
Very similar conclusions are reached in an analysis by G. Costa *et al.*, Nucl. Phys. **B297**, 244 (1988);
Deep inelastic scattering is considered by G.L. Fogli and D. Haidt, Z. Phys. **C40**, 379 (1988).
10. A. Sirlin, Phys. Rev. **D22**, 971 (1980); **D29**, 89 (1984);
W. Hollik, Fortsch. Phys. **38**, 165 (1990);
D.C. Kennedy *et al.*, Nucl. Phys. **B321**, 83 (1989);
D.C. Kennedy and B.W. Lynn, Nucl. Phys. **B322**, 1 (1989);
D.Yu Bardin *et al.*, Z. Phys. **C44**, 493 (1989);
For recent reviews, see the articles by W. Hollik and W. Marciano, in “Precision Tests of the Standard Electroweak Model,” ed. P. Langacker (World, Singapore, 1994) Extensive references to other papers are given in Ref. 9.
11. V.A. Novikov, L.B. Okun, and M.I. Vysotsky, Nucl. Phys. **B397**, 35 (1993).

12. W.J. Marciano and J.L. Rosner, Phys. Rev. Lett. **65**, 2963 (1990).
13. G. Degross, S. Fanchiotti, and A. Sirlin, Nucl. Phys. **B351**, 49 (1991).
14. P. Gambino and A. Sirlin, Phys. Rev. **D49**, 1160 (1994).
15. Zfitter: D. Bardin *et al.*, CERN-TH.6443/92, and references therein.
16. R. Barbieri *et al.*, Phys. Lett. **B288**, 95 (1992); Nucl. Phys. **B409**, 105 (1993).
17. A. Djouadi and C. Verzegnassi, Phys. Lett. **B195**, 265 (1987); A. Djouadi, Nuovo Cimento **100A**, 357 (1988).
18. J. Fleischer *et al.*, Phys. Lett. **B293**, 437 (1992); K.G. Chetyrkin *et al.*, Mod. Phys. Lett. **A8**, 2785 (1993).
19. S. Fanchiotti *et al.*, (see Ref. 7 above); A. Sirlin, Brookhaven preprint BNL-60161 (1994).
20. B.H. Smith and M.B. Voloshin, Minnesota preprint UMN-TH-1241/94.
21. CDHS: H. Abramowicz *et al.*, Phys. Rev. Lett. **57**, 298 (1986); A. Blondel *et al.*, Z. Phys. **C45**, 361 (1990).
22. CHARM: J.V. Allaby *et al.*, Z. Phys. **C36**, 611 (1987).
23. BEBC: D. Allasia *et al.*, Nucl. Phys. **B307**, 1 (1988).
24. Previous Fermilab results are CCFR:P.G. Reutens *et al.*, Z. Phys. **C45**, 539 (1990); FMM: T.S. Mattison *et al.*, Phys. Rev. **D42**, 1311 (1990).
25. CCFR: C.G. Arroyo *et al.*, CCFR preprint (1994).
26. CHARM I: J. Dorenbosch *et al.*, Z. Phys. **C41**, 567 (1989).
27. BNL E734: L.A. Ahrens *et al.*, Phys. Rev. **D41**, 3297 (1990).
28. CHARM II: P. Vilain *et al.*, Phys. Lett. **B281**, 159 (1992).
29. C.Y. Prescott *et al.*, Phys. Lett. **84B**, 524 (1979).
30. Boulder: M.C. Noecker *et al.*, Phys. Rev. Lett. **61**, 310 (1988). For reviews of earlier work, see M.A. Bouchiat L. Pottier, Science **234**, 1203 (1986); M.A. Bouchiat, *12th International Atomic Physics Conference*, (Ann Arbor, MI, July 1990).
31. S.A. Blundell, W.R. Johnson, and J. Sapirstein, Phys. Rev. Lett. **65**, 1411 (1990); V.A. Dzuba *et al.*, Phys. Lett. **141A**, 147 (1989).
32. B.W. Lynn and R.G. Stuart, Nucl. Phys. **B253**, 216 (1985); *Physics at LEP*, ed. J. Ellis and R. Peccei, CERN 86-02, Vol. I.
33. R. Marshall, Z. Phys. **C43**, 607 (1989); C. Kiesling, *Tests of the Standard Theory of Electroweak Interactions* (Springer-Verlag, NY, 1988); Y. Mori *et al.*, Phys. Lett. **B218**, 499 (1989).
34. SLD: K. Abe *et al.*, SLAC-PUB-6456.
35. D. Albert *et al.*, Nucl. Phys. **B166**, 460 (1980); F. Jegerlehner, Z. Phys. **C32**, 425 (1986); W. Beenakker and W. Hollik, Z. Phys. **C40**, 141 (1988); A. Djouadi *et al.*, Z. Phys. **C46**, 411 (1990); A.A. Akhundov *et al.*, Nucl. Phys. **B276**, 1 (1986); A. Borrelli *et al.*, Nucl. Phys. **B333**, 357 (1990).
36. B.A. Kniehl and J.H. Kuhn, Nucl. Phys. **B329**, 547 (1990); K.G. Chetyrkin *et al.*, Phys. Lett. **B305**, 285 (1993); K.G. Chetyrkin and O.V. Tarasov, Bielefeld TTP-93-38.
37. W. Beenakker and W. Hollik, Z. Phys. **C40**, 141 (1988); A.A. Akhundov *et al.*, Nucl. Phys. **B276**, 1 (1986); J. Bernabeu, A. Pich, and A. Santamaria, Nucl. Phys. **B363**, 326 (1991).
38. See the Z-Boson Listings, and the LEP Collaborations, CERN/PPE/93-157.

27. CONSTRAINTS ON NEW PHYSICS FROM ELECTROWEAK ANALYSES

This section prepared May 1994 by P. Langacker and J. Erler.

Precision electroweak experiments are sensitive to loop effects, allowing a prediction of the top quark mass m_t , constraints on the Higgs mass M_H , and a search for certain types of new physics that have not been directly detected. This article will mainly discuss m_t , M_H , and the effects of exotic particles with masses large compared to M_Z on the gauge boson self-energies. Brief remarks are made on new physics which is not of this type. The effects of m_t and M_H on the radiative corrections are treated exactly to one-loop order. This can in principle be done for other types of new physics, but this necessitates a case-by-case discussion. Instead, the article will discuss in detail only the constraints on particles with heavy masses $M_{\text{new}} \gg M_Z$ in an expansion in M_Z/M_{new} . In this case, most of the effects on precision measurements can be described by three gauge self-energy parameters S , T , and U , and a $Zb\bar{b}$ vertex correction parameter γ_b .

A large value of $|m_t - m_b|$ breaks vector SU(2) symmetry and significantly affects many precision electroweak observables. The major sensitivity for processes involving light external fermions is through t - and b -quark loop contributions to the W and Z self-energies [1]. Most of the shift in M_W is absorbed into the measured value of the Fermi constant G_F , while the prediction for M_Z

$$M_Z = \frac{M_W}{\hat{\rho}^{1/2} \hat{c}_Z}, \quad (27.1)$$

decreases rapidly for large m_t . In Eq. (27.1) $\hat{\rho} \simeq 1 + \rho_t$, where

$$\rho_t = \frac{3G_F m_t^2}{8\sqrt{2}\pi^2} \sim 0.0031 \left(\frac{m_t}{100 \text{ GeV}} \right)^2, \quad (27.2)$$

and $\hat{c}_Z = \cos\hat{\theta}_W(M_Z)$, the cosine of the weak angle in the $\overline{\text{MS}}$ scheme evaluated at M_Z [2]. In addition to M_Z itself, neutral current amplitudes and the coefficient of $G_F M_Z^2$ in the expression for Γ_Z are multiplied by $\hat{\rho}$. There is additional logarithmic m_t dependence in these quantities and in M_W . Vertex and box diagrams also introduce large (quadratic) m_t dependence, which is especially important in quantities involving external b quarks (in order to avoid mixing angle suppressions), such as in the $Z \rightarrow b\bar{b}$ partial width or in $B - \bar{B}$ mixing. Finally, in the on-shell renormalization scheme, significant but somewhat artificial m_t dependence is introduced into Z vertices through the definition [2] $s_W^2 \equiv 1 - M_W^2/M_Z^2$.

The CDF collaboration has recently presented [3] a number of candidate top quark events, with a mass $m_t = 174 \pm 16$ GeV. However, they have not yet claimed a definite discovery. Therefore, the indirect constraints on m_t from precision experiments will first be summarized, and then the results when the CDF value is combined with the indirect constraints.

As discussed in the section on the Standard Model of Electroweak Interactions (see especially Figure 1 of that section), the consistency of the various observables allows a prediction for m_t . A global fit to all data (see Table 26.5 of the Standard Model Section) yields

$$m_t = 169_{-18-20}^{+16+17} \text{ GeV}, \quad (27.3)$$

where the central value is for a Higgs mass $M_H = 300$ GeV and the second uncertainty is from varying M_H in the range 60 GeV (–) to 1000 GeV (+). One can also use the indirect data to set upper limits on m_t . Taking the $D0$ lower limit $m_t > 131$ GeV [4] into account, one finds $m_t < 205$ (210) GeV at 90 (95)% CL for $M_H \leq 1000$ GeV.

The upper limit on m_t is unchanged or strengthened in the presence of many types of new physics. For example, nondegenerate multiplets of heavy fermions or scalars break the vector part of weak SU(2) and lead to a decrease in the value of M_Z/M_W . A nondegenerate SU(2) doublet (f_2^I) yields a positive contribution to ρ_t of [1]

$$\frac{CG_F}{8\sqrt{2}\pi^2} \Delta m^2, \quad (27.4)$$

where

$$\Delta m^2 \equiv m_1^2 + m_2^2 - \frac{4m_1^2 m_2^2}{m_1^2 - m_2^2} \ln \frac{m_1}{m_2} \geq (m_1 - m_2)^2, \quad (27.5)$$

and $C = 1$ (3) for color singlets (triplets). Thus, to a good first approximation (*i.e.*, except for $\Gamma(Z \rightarrow b\bar{b})$ and logarithmic effects) the 90% CL upper limit $m_t < 205$ GeV in the Standard Model can be reinterpreted as

$$m_t^2 + \sum_i \frac{C_i}{3} \Delta m_i^2 < (205 \text{ GeV})^2, \quad (27.6)$$

where the sum includes fourth-family quark or lepton doublets, (f_2^I) or (E^0), and scalar doublets such as (\hat{f}) in supersymmetry (in the absence of $L - R$ mixing). Similarly, heavy Z' bosons decrease the prediction for M_Z due to mixing and generally strengthen the m_t limit [5]. Additional Higgs doublets which participate in spontaneous symmetry breaking [6], heavy lepton doublets involving Majorana neutrinos [7], and the presence of heavy degenerate chiral multiplets (the S parameter, to be discussed below) can weaken the limits on m_t , though the effect is usually small for reasonable parameter ranges. The only known way to significantly weaken the limits is to allow for the presence of Higgs triplets (or higher-dimensional representations), whose vacuum expectation values can cancel all of the quadratic m_t dependence except for the $b\bar{b}$ vertex. Even in that case one has an upper limit of 220 (237) GeV at 90 (95)% CL, mainly from $R_b \equiv \Gamma(Z \rightarrow b\bar{b})/\Gamma(Z \rightarrow \text{had})$ [2,8].

All of these constraints become much tighter when one incorporates the mass $m_t = 174 \pm 16$ GeV of the CDF candidate events [3], which is remarkably consistent with the prediction in Eq. (27.3). (The indirect prediction is for the pole mass, which should approximately coincide with the kinematic mass determined by CDF.) The joint fit to indirect and direct data yields [2] $m_t = 172_{-12-9}^{+11+8}$ GeV, again with the central value for $M_H = 300$ GeV.

As discussed in Ref. 2, the combination of indirect data with the CDF value for m_t allows stringent limits on new physics. In particular, many extensions of the Standard Model are described by the ρ_0 parameter:

$$\rho_0 \equiv M_W^2 / (M_Z^2 \hat{c}_Z^2 \hat{\rho}), \quad (27.7)$$

which describes new sources of SU(2) breaking that cannot be accounted for by Higgs doublets or m_t effects. It has previously been difficult to distinguish ρ_0 from $\hat{\rho} \simeq 1 + \rho_t$ experimentally, though some separation could be done utilizing R_b [8]. Using the CDF m_t value as an independent constraint, however, one can calculate $\hat{\rho}$ and thus obtain the precise value [2] $\rho_0 = 1.0004 \pm 0.0022 \pm 0.002$, where the second error is from M_H . In Ref. 2, this result was used to constrain the vacuum expectation values of higher-dimensional Higgs representations. It can also be used to constrain Δm^2 from nondegenerate SU(2) multiplets, as defined in Eq. (27.5). That is, in the presence of such multiplets, one has

$$\frac{3G_F}{8\sqrt{2}\pi^2} \sum_i \frac{C_i}{3} \Delta m_i^2 = \rho_0 - 1, \quad (27.8)$$

implying

$$\sum_i \frac{C_i}{3} \Delta m_i^2 < (95 \text{ GeV})^2, (111 \text{ GeV})^2, (128 \text{ GeV})^2 \quad (27.9)$$

for $M_H = 60, 300, \text{ or } 1000$ GeV at 90% CL.

As discussed in the Standard Model of Electroweak Interactions section, the indirect data exhibit a moderate preference for a smaller Higgs mass. The best fit to m_t as a function of M_H is roughly

$$m_t \sim 169_{-18}^{+16} + 13 \ln \left(\frac{M_H}{300 \text{ GeV}} \right) \quad (27.10)$$

without the direct CDF constraint or

$$m_t \sim 172_{-12}^{+11} + 6 \ln \left(\frac{M_H}{300 \text{ GeV}} \right) \quad (27.11)$$

with it. The χ^2 for $M_H = 60$ GeV is lower by 2.9(2.1) than that for $M_H = 1000$ GeV for the two cases, implying $M_H \lesssim 840$ GeV at 90% CL. While not compelling statistically, this result is consistent with the minimal supersymmetric extension of the Standard Model, which acts much like the Standard Model with a light Higgs as far as precision experiments are concerned. Of course, the conclusions for M_H could be invalidated if other new physics modifies the precision observables significantly.

A number of authors [9–14] have considered the general effects on neutral current and Z and W -pole observables of various types of heavy (*i.e.*, $M > M_Z$) physics which contribute to the W and Z self-energies but which do not have any direct coupling to the ordinary fermions. In addition to nondegenerate multiplets, which break the vector part of weak SU(2), these include heavy degenerate multiplets of chiral fermions which break the axial generators. The effects of one degenerate chiral doublet are small, but in technicolor theories there may be many chiral doublets and therefore significant effects [9].

All such effects can be described by just three parameters, S , T , and U . T is proportional to the difference between the W and Z self-energies at $Q^2 = 0$ (*i.e.*, vector SU(2)-breaking), while S ($S + U$) is associated with the difference between the Z (W) self-energy at $Q^2 = M_{Z,W}^2$ and $Q^2 = 0$ (axial SU(2)-breaking). In the $\overline{\text{MS}}$ scheme [10]

$$\begin{aligned} \alpha T &\equiv \frac{\Pi_{WW}^{\text{new}}(0)}{M_W^2} - \frac{\Pi_{ZZ}^{\text{new}}(0)}{M_Z^2} \\ \frac{\alpha}{4\hat{s}_Z^2 c_Z^2} S &\equiv \frac{\Pi_{ZZ}^{\text{new}}(M_Z^2) - \Pi_{ZZ}^{\text{new}}(0)}{M_Z^2} \\ \frac{\alpha}{4\hat{s}_Z^2} (S + U) &\equiv \frac{\Pi_{WW}^{\text{new}}(M_W^2) - \Pi_{WW}^{\text{new}}(0)}{M_W^2}, \end{aligned} \quad (27.12)$$

where Π_{WW}^{new} and Π_{ZZ}^{new} are respectively the contributions of the new physics to the W and Z self-energies, $\hat{s}_Z^2 = \sin^2 \hat{\theta}_W(M_Z)$, $c_Z^2 = 1 - \hat{s}_Z^2$, and α is the running coupling evaluated at M_Z . S , T , and U are defined with a factor of α removed, so that they are expected to be of order unity in the presence of new physics. S , T , and U are related to other parameters ($\hat{\epsilon}_i$, h_i , S_i) defined in [10–12] by

$$\begin{aligned} T &= h_V = \hat{\epsilon}_1 / \alpha \\ S &= h_{AZ} = S_Z = 4\hat{s}_Z^2 \hat{\epsilon}_3 / \alpha \\ U &= h_{AW} - h_{AZ} = S_W - S_Z = -4\hat{s}_Z^2 \hat{\epsilon}_2 / \alpha. \end{aligned} \quad (27.13)$$

A heavy nondegenerate multiplet of fermions or scalars contributes to T as

$$\rho_0 = \frac{1}{1 - \alpha T} \simeq 1 + \alpha T, \quad (27.14)$$

where ρ_0 is given in Eq. (27.8). If there are non-doublet Higgs representations, their vacuum expectation values also contribute to ρ_0 . The effects of such nonstandard Higgs representations cannot be separated from heavy nondegenerate multiplets unless the new physics has other consequences, such as vertex corrections.

A multiplet of heavy degenerate chiral fermions yields

$$S = C \sum_i \left(t_{3L}(i) - t_{3R}(i) \right)^2 / 3\pi, \quad (27.15)$$

where $t_{3L,R}(i)$ is the third component of weak isospin of the left-(right-) handed component of fermion i and C is the number of colors. For example, a heavy degenerate family would contribute $2/3\pi$ to S . In technicolor models with QCD-like dynamics, one expects [9] $S \sim 0.45$ for an isodoublet of technifermions, assuming $N_{TC} = 4$ technicolors, while $S \sim 1.62$ for a full technigeneration with $N_{TC} = 4$; T is harder to estimate because it is model dependent. In these examples one has $S \geq 0$. However, it is possible to find situations in which $S < 0$ [15]. In particular, these estimates do not apply

to models of walking technicolor, for which S can be smaller or even negative [16]. Supersymmetric extensions of the Standard Model generally give very small effects [17]. Most simple types of new physics yield $U = 0$, although there are counter-examples, such as the effects of anomalous triple-gauge vertices [12].

It is also possible to parametrize the effects of large $m_t \gg M_Z$ (except for the $b\bar{b}$ vertex) or $M_H \gg M_Z$ in terms of S , T , and U . If one takes $m_t = m_t^{\text{ref}}$, $M_H = M_H^{\text{ref}}$ as a reference point, then other values of m_t and M_H can be expressed for large m_t , M_H as [11]

$$\begin{aligned} \Delta T &= \frac{\rho_t(m_t) - \rho_t(m_t^{\text{ref}})}{\alpha} \\ &\quad - \frac{3G_F}{4\sqrt{2}\alpha\pi^2} (M_Z^2 - M_W^2) \ln(M_H/M_H^{\text{ref}}) \\ \Delta S &= c_S \ln(m_t/m_t^{\text{ref}}) + \frac{1}{6\pi} \ln(M_H/M_H^{\text{ref}}) \\ \Delta U &= c_U \ln(m_t/m_t^{\text{ref}}), \end{aligned} \quad (27.16)$$

where the coefficients c_S and c_U depend on the renormalization scheme. Various authors use different reference values for m_t and M_H when determining S , T , and U from the data. In the following $M_H^{\text{ref}} = 300$ GeV and the global best fit value $m_t^{\text{ref}} = 169$ GeV will be used when the CDF constraint is not imposed. The allowed ranges of the parameters will then represent both the effects of new physics and of values of m_t and M_H different from the reference values, *i.e.*, $S = S_{\text{new}} + \Delta S$, $T = T_{\text{new}} + \Delta T$, $U = U_{\text{new}} + \Delta U$. For later use, note that $\Delta S = \Delta S_t + \Delta S_{\text{Higgs}}$ and similarly for ΔT .

The Standard Model expressions for observables are replaced by

$$\begin{aligned} M_Z^2 &= M_{Z0}^2 \frac{1 - \alpha T}{\rho_0} \frac{1}{1 - G_F M_{Z0}^2 S / 2\sqrt{2}\pi} \\ M_W^2 &= M_{W0}^2 \frac{1}{1 - G_F M_{W0}^2 (S + U) / 2\sqrt{2}\pi}, \end{aligned} \quad (27.17)$$

where M_{Z0} and M_{W0} are the Standard Model expressions in the $\overline{\text{MS}}$ scheme. Furthermore,

$$\begin{aligned} \Gamma_Z &= \frac{\rho_0}{1 - \alpha T} M_Z^3 \beta_Z \\ \Gamma_W &= M_W^3 \beta_W \\ A_i &= \frac{\rho_0}{1 - \alpha T} A_{i0}, \end{aligned} \quad (27.18)$$

where β_Z and β_W are the Standard Model expressions for the reduced widths Γ_{Z0}/M_{Z0}^3 and Γ_{W0}/M_{W0}^3 , M_Z and M_W are the physical masses, and A_i (A_{i0}) is a neutral current amplitude (in the Standard Model).

The $Z \rightarrow b\bar{b}$ vertex is sensitive to certain types of new physics which primarily couple to heavy families [18,19]. It is useful to introduce an additional parameter γ_b by [20]

$$\Gamma(Z \rightarrow b\bar{b}) = \Gamma^0(Z \rightarrow b\bar{b})(1 + \gamma_b), \quad (27.19)$$

where Γ^0 is the Standard Model expression (or the expression modified by S , T , U , and ρ_0). Experimentally, $R_b \equiv \Gamma(Z \rightarrow b\bar{b})/\Gamma(\text{had})$ is about 1.8σ above the Standard Model expectations, favoring a positive γ_b . Extended technicolor models generally yield negative values of γ_b of a few percent [18], while supersymmetry can yield (typically small) contributions of either sign [19].

The indirect data allow a simultaneous determination of \hat{s}_Z^2 (*e.g.*, from the Z -pole asymmetries), S (from M_Z), U (from M_W), T (*e.g.*, from the Z -decay widths), α_s (from $\Gamma(Z \rightarrow \text{had})/\Gamma(\ell\bar{\ell})$), and γ_b (from R_b) with little correlation. For the reference values for m_t and M_H , one obtains

$$\begin{aligned} S &= -0.42 \pm 0.36 \\ T &= -0.37 \pm 0.39 \\ U &= -1.3 \pm 1.3 \\ \gamma_b &= 0.032 \pm 0.016, \end{aligned} \quad (27.20a)$$

and $\hat{s}_Z^2 = 0.2311 \pm 0.0007$, $\alpha_s = 0.099 \pm 0.013$. Combining with the CDF direct value of m_t , one does not need to introduce a reference value for m_t . That is, one can determine S_{new} , T_{new} , U_{new} , and γ_b directly, in a simultaneous fit with \hat{s}_Z^2 , α_s , and m_t . One obtains

$$\begin{aligned} S_{\text{new}} &= -0.42 \pm 0.36_{-0.17}^{+0.08} \\ T_{\text{new}} &= -0.35 \pm 0.44_{-0.11}^{+0.18} \\ U_{\text{new}} &= -1.3 \pm 1.3 \\ \gamma_b &= 0.032 \pm 0.017, \end{aligned} \quad (27.20b)$$

and the allowed region in $S'_{\text{new}} - T'_{\text{new}}$ is shown in Figure 1. (S'_{new} includes the M_H dependence explicitly, $S'_{\text{new}} \equiv S_{\text{new}} + \Delta S_{\text{Higgs}}$, and similarly for T'_{new} , see Eq. (27.16).) The values in Eq. (27.20b) are almost identical to those in Eq. (27.20a) except for a small increase in the errors due to the variation of m_t around the best fit value 175 ± 16 . The second error bars represent the uncertainty due to M_H ranging from the reference value of 300 GeV to 1000 (upper) or 60 (lower) GeV. From Eq. (27.20b) one obtains $S_{\text{new}} < 0.09 (+0.23)$, and $T < 0.27 (0.43)$ at 90 (95)% CL. If one requires the constraint $S_{\text{new}} \geq 0$ then (as in QCD-like technicolor models) $S_{\text{new}} < 0.38 (0.46)$. Allowing arbitrary S_{new} , only one heavy generation of ordinary fermions is allowed at 95% CL. The favored value of S_{new} is problematic for technicolor models with many techni-doublets and QCD-like dynamics, as is the value of γ_b . Although S_{new} is consistent with zero, the electroweak asymmetries, especially the SLD left-right asymmetry, favor $S_{\text{new}} < 0$. The simplest origin of $S_{\text{new}} < 0$ would probably be an additional heavy Z' boson [5], which could mimic $S_{\text{new}} < 0$.

There is no simple parametrization that is powerful enough to describe the effects of every type of new physics on every possible observable. The S , T , and U formalism describes many types of heavy physics which affect only the gauge self-energies, and it can be applied to all precision observables. However, new physics which couples directly to ordinary fermions, such as heavy Z' bosons or mixing with exotic fermions cannot be fully parametrized in the S , T , and U framework. It is convenient to treat these types of new physics by parametrizations that are specialized to that particular class of theories (e.g., extra Z' bosons), or to consider specific models (which might contain, e.g., Z' bosons and exotic fermions with correlated parameters). An alternate formalism [21] defines parameters, ϵ_1 , ϵ_2 , ϵ_3 , ϵ_b in terms of the specific observables M_W/M_Z , the leptonic Z width $\Gamma_{\ell\ell}$, the forward-backward asymmetry [2] at the Z pole, $A_{FB}^{(0,\ell)}$, and R_b . The definitions coincide with those for $\hat{\epsilon}_i$ in Eqs. (27.12) and (27.13) for physics which affects gauge self-energies only, but the ϵ 's now parametrize arbitrary types of new physics and can also incorporate all of the effects of m_t and M_H on the four basic observables. However, the ϵ 's are not related to other observables unless additional model-dependent assumptions are made. Another approach [22] parametrizes new physics in terms of gauge-invariant sets of operators. It is especially powerful in studying the effects of new physics on nonabelian gauge vertices. The most general approach introduces deviation vectors [24]. Each type of new physics defines a deviation vector, the components of which are the deviations of each observable from its Standard Model prediction, normalized to the experimental uncertainty. The length (direction) of the vector represents the strength (type) of new physics.

References:

1. M. Veltman, Nucl. Phys. **B123**, 89 (1977); M. Chanowitz, M.A. Furman, and I. Hinchliffe, Phys. Lett. **B78**, 285 (1978).
2. See the section on the Standard Model of Electroweak Interactions.
3. CDF: F. Abe *et al.*, FERMILAB-PUB-94/097-E.
4. DØ: S. Abachi *et al.*, Phys. Rev. Lett. **72**, 2138 (1994).
5. See P. Langacker and M. Luo, Phys. Rev. **D45**, 278 (1992) and references therein.
6. A. Denner *et al.*, Phys. Lett. **B240**, 438 (1990).
7. S. Bertolini and A. Sirlin, Phys. Lett. **B257**, 179 (1991).
8. P. Langacker and M. Luo, Phys. Rev. **D44**, 817 (1991).




Figure 27.1: 90% CL limits on S'_{new} and T'_{new} from various inputs. S'_{new} and T'_{new} have the m_t dependence removed (it is fit separately), but contain the small M_H dependence. In the Standard Model, one expects $S'_{\text{new}} = T'_{\text{new}} = 0$ for $M_H = 300$ GeV (indicated by a box). The expected values for $M_H = 60$ GeV (circle) and 1000 GeV (diamond) are also indicated. The fit to M_W and M_Z assumes $U = 0$, while U is arbitrary in the other fits.

9. M. Peskin and T. Takeuchi, Phys. Rev. Lett. **65**, 964 (1990) and Phys. Rev. **D46**, 381 (1992); M. Golden and L. Randall, Nucl. Phys. **B361**, 3 (1991).
10. W. Marciano and J. Rosner, Phys. Rev. Lett. **65**, 2963 (1990).
11. D. Kennedy and P. Langacker, Phys. Rev. Lett. **65**, 2967 (1990) and Phys. Rev. **D44**, 1591 (1991).
12. G. Altarelli and R. Barbieri, Phys. Lett. **B253**, 161 (1990).
13. B. Holdom and J. Terning, Phys. Lett. **B247**, 88 (1990).
14. B.W. Lynn, M.E. Peskin, and R.G. Stuart, in *Physics at LEP*, CERN 86-02, Vol. I, p. 90 (1986).
15. H. Georgi, Nucl. Phys. **B363**, 301 (1991); M.J. Dugan and L. Randall, Phys. Lett. **B264**, 154 (1991); E. Gates and J. Terning, Phys. Rev. Lett. **67**, 1840 (1991).
16. R. Sundrum and S.D.H. Hsu, Nucl. Phys. **B391**, 127 (1993); R. Sundrum, Nucl. Phys. **B395**, 60 (1993); M. Luty and R. Sundrum, Phys. Rev. Lett. **70**, 529 (1993); T. Appelquist and J. Terning, Phys. Lett. **B315**, 139 (1993).
17. R. Barbieri, M. Frigeni, F. Giuliani, and H.E. Haber, Nucl. Phys. **B341**, 309 (1990).
18. R.S. Chivukula, B. Selipsky, and E.H. Simmons, Phys. Rev. Lett. **69**, 575 (1992); R.S. Chivukula *et al.*, Phys. Lett. **B311**, 157 (1993).
19. A. Djouadi *et al.*, Nucl. Phys. **B349**, 48 (1991); M. Boulware and D. Finnell, Phys. Rev. **D44**, 2054 (1991); G. Altarelli *et al.*, Phys. Lett. **B314**, 357 (1993).
20. G. Altarelli, R. Barbieri, and F. Caravaglios, Nucl. Phys. **B405**, 3 (1993).
21. G. Altarelli, R. Barbieri, and S. Jadach, Nucl. Phys. **B369**, 3 (1992); **B376**, 444(E) (1992).
22. A. De Rújula, M.B. Gavela, P. Hernandez, and E. Massó, Nucl. Phys. **B384**, 3 (1992).
23. C.P. Burgess and D. London, Phys. Rev. **D48**, 4337 (1993).
24. P. Langacker, M. Luo, and A.K. Mann, Rev. Mod. Phys. **64**, 87 (1992).

28. THE CABIBBO-KOBAYASHI-MASKAWA MIXING MATRIX

Updated 1993 by F.J. Gilman, K. Kleinknecht, and B. Renk.

In the Standard Model with $SU(2) \times U(1)$ as the gauge group of electroweak interactions, both the quarks and leptons are assigned to be left-handed doublets and right-handed singlets. The quark mass eigenstates are not the same as the weak eigenstates, and the matrix relating these bases was defined for six quarks and given an explicit parametrization by Kobayashi and Maskawa [1] in 1973. It generalizes the four-quark case, where the matrix is parametrized by a single angle, the Cabibbo angle [2].

By convention, the three charge 2/3 quarks (u , c , and t) are unmixed, and all the mixing is expressed in terms of a 3×3 unitary matrix V operating on the charge $-1/3$ quarks (d , s , b):

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}. \quad (28.1)$$

The values of individual matrix elements can in principle all be determined from weak decays of the relevant quarks, or, in some cases, from deep inelastic neutrino scattering. Using the constraints discussed below together with unitarity, and assuming only three generations, the 90% confidence limits on the magnitude of the elements of the complete matrix are:

$$\begin{pmatrix} 0.9747 \text{ to } 0.9759 & 0.218 \text{ to } 0.224 & 0.002 \text{ to } 0.005 \\ 0.218 \text{ to } 0.224 & 0.9738 \text{ to } 0.9752 & 0.032 \text{ to } 0.048 \\ 0.004 \text{ to } 0.015 & 0.030 \text{ to } 0.048 & 0.9988 \text{ to } 0.9995 \end{pmatrix}. \quad (28.2)$$

The ranges shown are for the individual matrix elements. The constraints of unitarity connect different elements, so choosing a specific value for one element restricts the range of the others.

There are several parametrizations of the Cabibbo-Kobayashi-Maskawa matrix. In view of the need for a "standard" parametrization in the literature, we advocate:

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta_{13}} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta_{13}} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta_{13}} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta_{13}} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta_{13}} & c_{23}c_{13} \end{pmatrix} \quad (28.3)$$

proposed by Chau and Keung [3]. The choice of rotation angles follows earlier work of Maiani [4], and the placement of the phase follows that of Wolfenstein [5]. The notation used is that of Harari and Leurer [6] who, along with Fritzsch and Plankl [7], proposed this parametrization as a particular case of a form generalizable to an arbitrary number of "generations." The general form was also put forward by Botella and Chau [8]. Here $c_{ij} = \cos \theta_{ij}$ and $s_{ij} = \sin \theta_{ij}$, with i and j being "generation" labels, $\{i, j = 1, 2, 3\}$. In the limit $\theta_{23} = \theta_{13} = 0$ the third generation decouples, and the situation reduces to the usual Cabibbo mixing of the first two generations with θ_{12} identified with the Cabibbo angle [2]. The real angles θ_{12} , θ_{23} , θ_{13} can all be made to lie in the first quadrant by an appropriate redefinition of quark field phases. Then all s_{ij} and c_{ij} are positive, $|V_{us}| = s_{12}c_{13}$, $|V_{ub}| = s_{13}$, and $|V_{cb}| = s_{23}c_{13}$. As c_{13} is known to deviate from unity only in the fifth decimal place, $|V_{us}| = s_{12}$, $|V_{ub}| = s_{13}$, and $|V_{cb}| = s_{23}$ to an excellent approximation. The phase δ_{13} lies in the range $0 \leq \delta_{13} < 2\pi$, with non-zero values generally breaking CP invariance for the weak interactions. The generalization to the n generation case contains $n(n-1)/2$ angles and $(n-1)(n-2)/2$ phases [6,7,8]. The range of matrix elements in Eq. (28.2) corresponds to 90% CL limits on the angles of $s_{12} = 0.218$ to 0.224 , $s_{23} = 0.032$ to 0.048 , and $s_{13} = 0.002$ to 0.005 .

Kobayashi and Maskawa [1] originally chose a parametrization involving the four angles, θ_1 , θ_2 , θ_3 , δ :

$$\begin{pmatrix} d' \\ s' \\ b' \end{pmatrix} = \begin{pmatrix} c_1 & -s_1c_3 & -s_1s_3 \\ s_1c_2 & c_1c_2c_3 - s_2s_3e^{i\delta} & c_1c_2s_3 + s_2c_3e^{i\delta} \\ s_1s_2 & c_1s_2c_3 + c_2s_3e^{i\delta} & c_1s_2s_3 - c_2c_3e^{i\delta} \end{pmatrix} \begin{pmatrix} d \\ s \\ b \end{pmatrix}, \quad (28.4)$$

where $c_i = \cos \theta_i$ and $s_i = \sin \theta_i$ for $i = 1, 2, 3$. In the limit $\theta_2 = \theta_3 = 0$, this reduces to the usual Cabibbo mixing with θ_1

identified (up to a sign) with the Cabibbo angle [2]. Slightly different forms of the Kobayashi-Maskawa parametrization are found in the literature. The CKM matrix used in the 1982 *Review of Particle Properties* is obtained by letting $s_1 \rightarrow -s_1$ and $\delta \rightarrow \delta + \pi$ in the matrix given above. An alternative is to change Eq. (28.4) by $s_1 \rightarrow -s_1$ but leave δ unchanged. With this change in s_1 , the angle θ_1 becomes the usual Cabibbo angle, with the "correct" sign (*i.e.* $d' = d \cos \theta_1 + s \sin \theta_1$) in the limit $\theta_2 = \theta_3 = 0$. The angles θ_1 , θ_2 , θ_3 can, as before, all be taken to lie in the first quadrant by adjusting quark field phases. Since all these parametrizations are referred to as "the" Kobayashi-Maskawa form, some care about which one is being used is needed when the quadrant in which δ lies is under discussion.

Other parametrizations, mentioned above, are due to Maiani [4] and to Wolfenstein [5]. The latter emphasizes the relative sizes of the matrix elements by expressing them in powers of the Cabibbo angle. Still other parametrizations [9] have come into the literature in connection with attempts to define "maximal CP violation". No physics can depend on which of the above parametrizations (or any other) is used as long as a single one is used consistently and care is taken to be sure that no other choice of phases is in conflict.

Our present knowledge of the matrix elements comes from the following sources:

- (1) Nuclear beta decay, when compared to muon decay, gives [10–13]

$$|V_{ud}| = 0.9744 \pm 0.0010. \quad (28.5)$$

This includes refinements in the analysis of the radiative corrections, especially the order $Z\alpha^2$ effects, which have brought the ft-values from low and high Z Fermi transitions into good agreement.

- (2) Analysis of K_{e3} decays yields [14]

$$|V_{us}| = 0.2196 \pm 0.0023. \quad (28.6)$$

(The notation K_{e3}^+ refers to $K^+ \rightarrow \pi^0 e^+ \nu_e$ and K_{e3}^0 refers to $K_L^0 \rightarrow \pi^\pm e^\mp \nu_e$.) The isospin violation between K_{e3}^+ and K_{e3}^0 decays has been taken into account, bringing the values of $|V_{us}|$ extracted from these two decays into agreement at the 1% level of accuracy. The analysis of hyperon decay data has larger theoretical uncertainties because of first order $SU(3)$ symmetry breaking effects in the axial-vector couplings, but due account of symmetry breaking [15] applied to the WA2 data [16] gives a corrected value [17] of 0.222 ± 0.003 . We average these two results to obtain:

$$|V_{us}| = 0.2205 \pm 0.0018. \quad (28.7)$$

(3) The magnitude of $|V_{cd}|$ may be deduced from neutrino and antineutrino production of charm off valence d quarks. The dimuon production cross sections of the CDHS group [18] yield $\overline{B}_c |V_{cd}|^2 = 0.41 \pm 0.07 \times 10^{-2}$, where \overline{B}_c is the semileptonic branching fraction of the charmed hadrons produced. The corresponding value from a recent Tevatron experiment [19] is $\overline{B}_c |V_{cd}|^2 = 0.534_{-0.078}^{+0.052} \times 10^{-2}$. Averaging these two results gives $\overline{B}_c |V_{cd}|^2 = 0.47 \pm 0.05 \times 10^{-2}$. Supplementing this with measurements of the semileptonic branching fractions of charmed mesons [20], weighted by a production ratio of $D^0/D^+ = (60 \pm 10)/(40 \mp 10)$, to give $\overline{B}_c = 0.113 \pm 0.015$, yields

$$|V_{cd}| = 0.204 \pm 0.017 \quad (28.8)$$

(4) Values of $|V_{cs}|$ from neutrino production of charm are dependent on assumptions about the strange quark density in the parton-sea. The most conservative assumption, that the strange-quark sea does not exceed the value corresponding to an $SU(3)$ symmetric sea, leads to a lower bound [18], $|V_{cs}| > 0.59$. It is more advantageous to proceed analogously to the method used for extracting $|V_{us}|$ from K_{e3} decay; namely, we compare the experimental value for the width of D_{e3} decay with the expression [21] that follows from the standard weak interaction amplitude:

$$\Gamma(D \rightarrow \overline{K} e^+ \nu_e) = |f_+^D(0)|^2 |V_{cs}|^2 (1.54 \times 10^{11} \text{ s}^{-1}). \quad (28.9)$$

Here $f_+^D(q^2)$, with $q = p_D - p_K$, is the form factor relevant to D_{e3} decay; its variation has been taken into account with the parametrization $f_+^D(t)/f_+^D(0) = M^2/(M^2 - t)$ and $M = 2.1 \text{ GeV}/c^2$, a form and mass consistent with Mark III and E691 measurements [22,23]. Combining data on branching ratios for D_{e3} decays from Mark III, E687, E691, and CLEO experiments [22–24] with accurate values [25] for τ_{D^+} and τ_{D^0} , gives the value $(0.762 \pm 0.055) \times 10^{11} \text{ s}^{-1}$ for $\Gamma(D \rightarrow \bar{K}e^+\nu_e)$. Therefore

$$|f_+^D(0)|^2 |V_{cs}|^2 = 0.495 \pm 0.036. \quad (28.10)$$

A very conservative assumption is that $|f_+^D(0)| < 1$, from which it follows that $|V_{cs}| > 0.62$. Calculations of the form factor either performed [26,27] directly at $q^2 = 0$, or done [28] at the maximum value of $q^2 = (m_D - m_K)^2$ and interpreted at $q^2 = 0$ using the measured q^2 dependence, yield $f_+^D(0) = 0.7 \pm 0.1$. It follows that

$$|V_{cs}| = 1.01 \pm 0.18. \quad (28.11)$$

The constraint of unitarity when there are only three generations gives a much tighter bound (see below).

(5) The ratio $|V_{ub}/V_{cb}|$ can be obtained from the semileptonic decay of B mesons produced on the $\Upsilon(4S) b\bar{b}$ resonance by measuring the lepton energy spectrum above the endpoint of the $b \rightarrow c\ell\nu$ spectrum. There the $b \rightarrow u\ell\nu$ decay rate can be obtained by subtracting the background from nonresonant e^+e^- reactions. This continuum background is determined from auxiliary measurements off the $\Upsilon(4S)$. Both the CLEO [29] and ARGUS [30] collaborations have reported evidence for $b \rightarrow u$ transitions in semileptonic B decays. The interpretation of the result in terms of $|V_{ub}/V_{cb}|$ depends fairly strongly on the theoretical model used to generate the lepton energy spectrum, especially for $b \rightarrow u$ transitions [27,28,31]. Combining the experimental and theoretical uncertainties, we quote

$$|V_{ub}/V_{cb}| = 0.08 \pm 0.02. \quad (28.12)$$

(6) The magnitude of V_{cb} itself can be determined if the measured semileptonic bottom hadron partial width is assumed to be that of a quark decaying through the usual $V - A$ interaction:

$$\Gamma(b \rightarrow c\ell\bar{\nu}_\ell) = \frac{B(b \rightarrow c\ell\bar{\nu}_\ell)}{\tau_b} = \frac{G_F^2 m_b^5}{192\pi^3} F(m_c/m_b) |V_{cb}|^2, \quad (28.13)$$

where τ_b is the b lifetime and $F(m_c/m_b)$ is a phase space factor that is approximately one-half. Most of the error on $|V_{cb}|$ derived from Eq. (28.13) is not from the experimental uncertainties, but in the theoretical uncertainties in choosing a value of m_b and in the use of the quark model to represent inclusively semileptonic decays which, at least for the B meson, are dominated by a few exclusive channels. Instead we use the nearly model-independent treatment in the heavy quark effective theory [32], where, in the case of $B \rightarrow D^*$ transitions, the decay rates at zero recoil are fixed by a normalization condition, with vanishing $1/m_q$ corrections [33]. From data of the ARGUS [34] and CLEO [35] experiments, we quote a value [36] derived from the decay of $\bar{B} \rightarrow D^*\ell\bar{\nu}_\ell$ of

$$|V_{cb}| = 0.040 \pm 0.005 \quad (28.14)$$

that is deduced using a B -lifetime of $(1.49 \pm 0.04) \text{ ps}$ [37]. The central value and the error are now comparable to what is obtained from the inclusive semileptonic decays, but ultimately, with more data, exclusive semileptonic decays should provide the most accurate value of $|V_{cb}|$.

The results for three generations of quarks, from Eqs. 28.5, 28.7, 28.8, 28.11, 28.12, and 28.14 plus unitarity, are summarized in the matrix in Eq. (28.2). The ranges given there are different from those given in Eqs. (28.5)–(28.14) (because of the inclusion of unitarity), but are consistent with the one standard deviation errors on the input matrix elements.

The data do not preclude there being more than three generations. Moreover, the entries deduced from unitarity might be altered when

the CKM matrix is expanded to accommodate more generations. Conversely, the known entries restrict the possible values of additional elements if the matrix is expanded to account for additional generations. For example, unitarity and the known elements of the first row require that any additional element in the first row have a magnitude $|V_{ub'}| < 0.07$. When there are more than three generations the allowed ranges (at 90% CL) of the matrix elements connecting the first three generations are

$$\begin{pmatrix} 0.9728 \text{ to } 0.9757 & 0.218 \text{ to } 0.224 & 0.002 \text{ to } 0.005 & \dots \\ 0.180 & \text{to } 0.228 & 0.800 \text{ to } 0.975 & 0.032 \text{ to } 0.048 & \dots \\ 0 & \text{to } 0.13 & 0 & \text{to } 0.56 & 0 & \text{to } 0.9995 & \dots \\ \vdots & & \vdots & & \vdots & & \vdots \end{pmatrix}, \quad (28.15)$$

where we have used unitarity (for the expanded matrix) and Eqs. 28.5, 28.7, 28.8, 28.11, 28.12, and 28.14.

Further information on the angles requires theoretical assumptions. For example, $B_d - \bar{B}_d$ mixing, if it originates from short distance contributions to ΔM_B dominated by box diagrams involving virtual t quarks, gives information on $V_{tb}V_{td}^*$ once hadronic matrix elements and the t quark mass are known. A similar comment holds for $V_{ts}V_{td}^*$ and $B_s - \bar{B}_s$ mixing.

Direct and indirect information on the CKM matrix is neatly summarized in terms of the “unitarity triangle.” The name arises since unitarity of the 3×3 CKM matrix applied to the first and third columns yields

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0. \quad (28.16)$$

The unitarity triangle is just a geometrical presentation of this equation in the complex plane [38]. We can always choose to orient the triangle so that $V_{cd}V_{cb}^*$ is horizontal; in the parametrization we have chosen, V_{cb} is real, and V_{cd} is real to a very good approximation in any case. Setting cosines of small angles to unity, Eq. (28.16) becomes

$$V_{ub}^* + V_{td} = s_{12}V_{cb}^*, \quad (28.17)$$

which is shown as the unitarity triangle in Fig. 28.1(a). Rescaling the triangle by a factor $[1/|s_{12}V_{cb}|]$, the coordinates of the vertices become

$$A(\text{Re}(V_{ub})/|s_{12}V_{cb}|, -\text{Im}(V_{ub})/|s_{12}V_{cb}|), B(1,0), C(0,0). \quad (28.18)$$

In the approximation of the Wolfenstein parametrization [5], with matrix elements expressed in powers of the Cabibbo angle, $\lambda \sim s_{12}$:

$$\begin{aligned} V_{us} &\sim \lambda \\ V_{ub} &\sim \lambda^3 A(\rho - i\eta) \\ V_{cb} &\sim \lambda^2 A \\ V_{td} &\sim \lambda^3 A(1 - \rho - i\eta), \end{aligned} \quad (28.19)$$

the coordinates of the vertex A of the unitarity triangle are simply (ρ, η) , as shown in Fig. 28.1(b).

CP -violating processes will involve the phase in the CKM matrix, assuming that the observed CP violation is solely related to a nonzero value of this phase. This allows additional constraints to be imposed. More specifically, a necessary and sufficient condition for CP violation with three generations can be formulated in a parametrization-independent manner in terms of the non-vanishing of the determinant of the commutator of the mass matrices for the charge $2e/3$ and charge $-e/3$ quarks [39]. CP violating amplitudes or differences of rates all are proportional to the CKM factor in this quantity. This is the product of factors $s_{12}s_{13}s_{23}c_{12}c_{13}^2c_{23}s_{\delta_{13}}$ in the parametrization adopted above, and is $s_1^2s_2s_3c_1c_2c_3s_\delta$ in that of Ref. 1. With the approximation of setting cosines to unity, this is just twice the area of the unitarity triangle. While hadronic matrix elements whose values are imprecisely known generally now enter, the constraints from CP violation in the neutral kaon system are tight enough to very much restrict the range of angles and the phase of




Figure 28.1: (a) Representation in the complex plane of the triangle formed by the CKM matrix elements V_{ub}^* , V_{td} , and $s_{12} V_{cb}^*$. (b) Rescaled triangle with vertices $A(\rho, \eta)$, $B(1, 0)$, and $C(0, 0)$.

the CKM matrix. For CP -violating asymmetries of neutral B mesons decaying to CP eigenstates, there is a direct relationship between the magnitude of the asymmetry in a given decay and $\sin 2\phi$, where $\phi = \alpha, \beta, \gamma$ is an appropriate angle of the unitarity triangle [38].

The combination of all the direct and indirect information can be used to find the overall constraints on the CKM matrix and thence the implications for future measurements of CP violation in the B system [40].

References:

1. M. Kobayashi and T. Maskawa, *Prog. Theor. Phys.* **49**, 652 (1973).
2. N. Cabibbo, *Phys. Rev. Lett.* **10**, 531 (1963).
3. L.-L. Chau and W.-Y. Keung, *Phys. Rev. Lett.* **53**, 1802 (1984).
4. L. Maiani, *Phys. Lett.* **62B**, 183 (1976) and in *Proceedings of the 1977 International Symposium on Lepton and Photon Interactions at High Energies* (DESY, Hamburg, 1977), p. 867.
5. L. Wolfenstein, *Phys. Rev. Lett.* **51**, 1945 (1983).
6. H. Harari and M. Leurer, *Phys. Lett.* **B181**, 123 (1986).
7. H. Fritzsch and J. Plankl, *Phys. Rev.* **D35**, 1732 (1987).
8. F.J. Botella and L.-L. Chau, *Phys. Lett.* **B168**, 97 (1986).
9. See, for example, M. Gronau and J. Schechter, *Phys. Rev. Lett.* **54**, 385 (1985), where various parametrizations are discussed, including one equivalent to that in Eq. (28.3).
10. W.J. Marciano and A. Sirlin, *Phys. Rev. Lett.* **56**, 22 (1986).
11. A. Sirlin and R. Zucchini, *Phys. Rev. Lett.* **57**, 1994 (1986).
12. W. Jaus and G. Rasche, *Phys. Rev.* **D35**, 3420 (1987).
13. A. Sirlin, *Phys. Rev.* **D35**, 3423 (1987).
14. H. Leutwyler and M. Roos, *Z. Phys.* **C25**, 91 (1984);
See also the earlier work of R.E. Shrock and L.L. Wang, *Phys. Rev. Lett.* **41**, 1692 (1978).
15. J.F. Donoghue, B.R. Holstein, and S.W. Klimt, *Phys. Rev.* **D35**, 934 (1987).
16. M. Bourquin *et al.*, *Z. Phys.* **C21**, 27 (1983).
17. J.M. Gaillard and G. Sauvage, private communication;
Using different $SU(3)$ symmetry breaking corrections, a somewhat higher value of $|V_{us}|$ is obtained by A. Garcia, R. Huerta, and P. Kielanowski, *Phys. Rev.* **D45**, 879 (1992).
18. H. Abramowicz *et al.*, *Z. Phys.* **C15**, 19 (1982).
19. C. Foudas *et al.*, *Phys. Rev. Lett.* **64**, 1207 (1990).
20. D. Hitlin, in *Proceedings of the 1987 International Symposium on Lepton and Photon Interactions at High Energies*, Hamburg, July 27–31, 1987, edited by W. Bartel and R. Rückl (North Holland, Amsterdam, 1988), p. 179.
21. The result for $M = 2.2$ GeV is found in F. Bletzacker, H.T. Nieh, and A. Soni, *Phys. Rev.* **D16**, 732 (1977).
22. CLEO: A. Bean *et al.*, *Phys. Lett.* **B317**, 647 (1993).
23. Mark III: Z. Bai *et al.*, *Phys. Rev. Lett.* **66**, 1011 (1991).
24. Mark III: J. Adler *et al.*, *Phys. Rev. Lett.* **62**, 1821 (1989).
25. E691: J.R. Raab *et al.*, *Phys. Rev.* **D37**, 2391 (1988);
E687: P.L. Frabetti *et al.*, *Phys. Lett.* **B263**, 584 (1991).
26. T.M. Aliev *et al.*, *Yad. Fiz.* **40**, 823 (1984) [*Sov. J. Nucl. Phys.* **40**, 527 (1984)].
27. M. Bauer, B. Stech, and M. Wirbel, *Z. Phys.* **C29**, 637 (1985).
28. B. Grinstein, N. Isgur, and M.B. Wise, *Phys. Rev. Lett.* **56**, 298 (1986);
B. Grinstein, N. Isgur, D. Scora, and M.B. Wise, *Phys. Rev.* **D39**, 799 (1989).
29. F. Bartelt *et al.*, *Phys. Rev. Lett.* **71**, 4111 (1993).
30. H. Albrecht *et al.*, *Phys. Lett.* **B255**, 297 (1991);
M. Danilov, rapporteur's talk at the *1993 EPS Conference on High-Energy Physics*, (Marseille, France, July 1993).
31. G. Altarelli *et al.*, *Nucl. Phys.* **B208**, 365 (1982).
32. N. Isgur and M.B. Wise, *Phys. Lett.* **B237**, 527 (1990); **B232**, 113 (1989);
E. Eichten and B. Hill, *Phys. Lett.* **B234**, 511 (1990).
33. M.E. Luke, *Phys. Lett.* **B252**, 447 (1990).
34. H. Albrecht *et al.*, *Z. Phys.* **C57**, 533 (1993);
H. Albrecht *et al.*, *Phys. Lett.* **B275**, 195 (1992).
35. G. Crawford *et al.*, CLEO Conf 93-30, submitted to *1993 EPS Conference on High-Energy Physics*, (Marseille, France, July 1993).
36. M. Neubert, *Phys. Lett.* **B264**, 455 (1991).
37. D. Karlen, *5th International Conference on Heavy Flavor Physics* (Montreal, Canada, July 1993).
38. L.-L. Chau and W.-Y. Keung, Ref. 3;
J.D. Bjorken, private communication and *Phys. Rev.* **D39**, 1396 (1989);
C. Jarlskog and R. Stora, *Phys. Lett.* **B208**, 268 (1988);
J.L. Rosner, A.I. Sanda, and M.P. Schmidt, in *Proceedings of the Workshop on High Sensitivity Beauty Physics at Fermilab*, Fermilab, November 11–14, 1987, edited by A.J. Slaughter, N. Lockyer, and M. Schmidt (Fermilab, Batavia, IL, 1988), p. 165;
C. Hamzaoui, J.L. Rosner and A.I. Sanda, *ibid.*, p. 215.
39. C. Jarlskog, *Phys. Rev. Lett.* **55**, 1039 (1985) and *Z. Phys.* **C29**, 491 (1985).
40. C.O. Dib *et al.*, *Phys. Rev.* **D41**, 1522 (1990);
A.I. Sanda, invited talk at the KEK Topical Conference on Electron-Positron Collision Physics, Tsukuba, Japan, May 17–19, 1989 and KEK report 89-70, 1989 (unpublished);
C.S. Kim, J.L. Rosner, and C.-P. Yuan, *Phys. Rev.* **D42**, 96 (1990).

29. CP VIOLATION

This section prepared April 1994 by L. Wolfenstein.

The symmetries C (particle-antiparticle interchange) and P (space inversion) hold for strong and electromagnetic interactions. After the discovery of large C and P violation in the weak interactions, it appeared that the product CP was a good symmetry. Then CP violation was observed in K^0 decays at a level given by the parameter $\epsilon = 2.3 \times 10^{-3}$. Larger CP -violation effects are anticipated in B^0 decays.

The eigenstates of the K^0 - \bar{K}^0 system can be written

$$|K_S\rangle = p|K^0\rangle + q|\bar{K}^0\rangle, \quad |K_L\rangle = p|K^0\rangle - q|\bar{K}^0\rangle. \quad (29.1)$$

If CP invariance held, we would have $q = p$ so that K_S would be CP even and K_L CP odd. (We define $|\bar{K}^0\rangle$ as CP $|K^0\rangle$). CP violation in K^0 - \bar{K}^0 mixing gives

$$\frac{p}{q} = \frac{(1 + \tilde{\epsilon})}{(1 - \tilde{\epsilon})}. \quad (29.2)$$

CP violation can also occur in the decay amplitudes

$$A(K^0 \rightarrow \pi\pi(I)) = A_I e^{i\delta_I}, \quad A(\bar{K}^0 \rightarrow \pi\pi(I)) = A_I^* e^{i\delta_I}, \quad (29.3)$$

where I is the isospin of $\pi\pi$, δ_I is the final-state phase shift, and A_I would be real if CP invariance held. The ratios of CP -violating to CP -conserving amplitudes $\eta_{+-} = A(K_L^0 \rightarrow \pi^+\pi^-)/A(K_S^0 \rightarrow \pi^+\pi^-)$ and $\eta_{00} = A(K_L^0 \rightarrow \pi^0\pi^0)/A(K_S^0 \rightarrow \pi^0\pi^0)$ can be written as

$$\eta_{+-} = \epsilon + \epsilon', \quad \eta_{00} = \epsilon - 2\epsilon', \quad (29.4a)$$

$$\epsilon = \tilde{\epsilon} + i (\text{Im } A_0 / \text{Re } A_0), \quad (29.4b)$$

$$|\sqrt{2}\epsilon'| = (\text{Re } A_2 / \text{Re } A_0) (\text{Im } A_2 / \text{Re } A_2 - \text{Im } A_0 / \text{Re } A_0). \quad (29.4c)$$

If CP violation is confined to the mass matrix, as in a superweak theory, ϵ' is zero and $\eta_{+-} = \eta_{00} = \epsilon = \tilde{\epsilon}$. The measurement of ϵ'/ϵ has as its goal finding an effect that requires CP violation in the decay amplitude; this corresponds to a relative phase between A_2 and A_0 as seen in Eq. (29.4c).

In the Standard Model CP violation arises as a result of a single phase entering the CKM matrix (q.v.). As a result in what is now the standard phase convention, two elements have large phases, $V_{ub} \sim e^{-i\gamma}$, $V_{td} \sim e^{-i\beta}$. Because these elements have small magnitudes and involve the third generation, CP violation in the K^0 system is small. A definite nonzero value for ϵ'/ϵ is expected but hadronic uncertainties allow theoretical values between 10^{-4} and 3×10^{-3} . On the other hand, large effects are expected in the B^0 system, which is a major motivation for B factories.

The most clearcut experiments would be those that measure asymmetries between B^0 and \bar{B}^0 decays. The time-dependent rate to a CP eigenstate a is given by

$$\Gamma_a \sim e^{-\Gamma t} \left([1 + |r_a|^2] \pm [1 - |r_a|^2] \cos(\Delta M t) \mp 2\eta_a \text{Im } r_a \sin(\Delta M t) \right), \quad (29.5)$$

where the top sign is for B^0 and the bottom for \bar{B}^0 , η_a is the CP eigenvalue and

$$r_a = (q_B/p_B) \bar{A}_a/A_a. \quad (29.6)$$

The quantity (q_B/p_B) comes from the analogue for B^0 of Eq. (29.1); however, for B^0 the eigenstates have a negligible lifetime difference and are distinguished only by the mass difference ΔM ; also as a result $|q_B/p_B| \approx 1$ so that $\tilde{\epsilon}_B$ is purely imaginary. A_a (\bar{A}_a) are the decay amplitudes to a for B^0 (\bar{B}^0). If only one quark weak transition contributes to the decay $|\bar{A}_a/A_a| = 1$ so that $|r_a| = 1$ and the $\cos(\Delta M t)$ term vanishes. The basic goal of the B factories is to observe the asymmetric $\sin(\Delta M t)$ term. For B^0 (\bar{B}^0) $\rightarrow \psi K_s$ from the transition $b \rightarrow c\bar{c}s$, one finds in the Standard Model the asymmetry parameter

$$-2\text{Im } r_a = \sin 2\beta. \quad (29.7)$$

The asymmetry is given directly in terms of a CKM phase with no hadronic uncertainty and is expected to be between 0.2 and 0.8. For B^0 (\bar{B}^0) $\rightarrow \pi^+\pi^-$ from the transition $b \rightarrow u\bar{u}d$

$$-2\text{Im } r_a = \sin 2(\beta + \gamma). \quad (29.8)$$

(This result has some hadronic uncertainty due to penguin contributions, but these should be able to be estimated from other observations.) While either of these asymmetries could be ascribed to B^0 - \bar{B}^0 mixing (q_B/p_B or $\tilde{\epsilon}_B$), the difference between the two asymmetries is evidence for direct CP violation. From Eq. (29.6) (with $\bar{A}_a/A_a = 1$) it is seen this corresponds to a phase difference between $A_{\psi K_S}$ and $A_{\pi^+\pi^-}$. Thus this is analogous to ϵ' . In the standard phase convention 2β in Eq. (29.7) and (29.8) arises from B^0 - \bar{B}^0 mixing whereas the 2γ comes from V_{bu} in the transition $b \rightarrow u\bar{u}d$.

CP violation in the decay amplitude is also revealed by the $\cos(\Delta M t)$ in Eq. (29.5) or by a difference in rates of B^+ and B^- to charge-conjugate states. These effects, however, require two contributing amplitudes to the decay (such as a tree amplitude plus a penguin) and also require final-state interaction phases. Predicted effects are very uncertain and are generally small.

For further details, see the notes on CP violation in the K_L^0 , K_S^0 , and B^0 Full Data Listings of this Review.

30. QUARK MODEL

30.1. Quantum numbers of the quarks

Each quark has spin 1/2 and baryon number 1/3. Table 30.1 gives the additive quantum numbers (other than baryon number) of the three generations of quarks. Our convention is that the *flavor* of a quark (I_z , S, C, B, or T) has the same sign as its *charge*. With this convention, any flavor carried by a *charged* meson has the same sign as its charge; e.g., the strangeness of the K^+ is +1, the bottomness of the B^+ is +1, and the charm and strangeness of the D_s^- are each -1.

By convention, each quark is assigned positive parity. Then each antiquark has negative parity.

Table 30.1: Additive quantum numbers of the quarks.

Property \ Quark	d	u	s	c	b	t
Q - electric charge	$-\frac{1}{3}$	$+\frac{2}{3}$	$-\frac{1}{3}$	$+\frac{2}{3}$	$-\frac{1}{3}$	$+\frac{2}{3}$
I_z - isospin z-component	$-\frac{1}{2}$	$+\frac{1}{2}$	0	0	0	0
S - strangeness	0	0	-1	0	0	0
C - charm	0	0	0	+1	0	0
B - bottomness	0	0	0	0	-1	0
T - topness	0	0	0	0	0	+1

30.2. Mesons: $q\bar{q}$ states

Nearly all known mesons are bound states of a quark q and an antiquark \bar{q}' (the flavors of q and q' may be different). If the orbital angular momentum of the $q\bar{q}'$ state is L , then the parity P is $(-1)^{L+1}$. A state $q\bar{q}'$ of a quark and its own antiquark is also an eigenstate of charge conjugation, with $C = (-1)^{L+S}$, where the spin S is 0 or 1. The $L = 0$ states are the pseudoscalars, $J^P = 0^-$, and the vectors, $J^P = 1^-$. Assignments for many of the known mesons are given in Table 30.2. States in the "normal" spin-parity series, $P = (-1)^J$, must, according to the above, have $S = 1$ and hence $CP = +1$. Thus mesons with normal spin-parity and $CP = -1$ are forbidden in the $q\bar{q}'$ model. The $J^{PC} = 0^{- -}$ state is forbidden as well. Mesons with such J^{PC} may exist, but would lie outside the $q\bar{q}'$ model.

The nine possible $q\bar{q}'$ combinations containing u , d , and s quarks group themselves into an octet and a singlet:

$$3 \otimes \bar{3} = 8 \oplus 1 \quad (30.1)$$

States with the same IJ^P and additive quantum numbers can mix. (If they are eigenstates of charge conjugation, they must also have the same value of C .) Thus the $I = 0$ member of the ground-state pseudoscalar octet mixes with the corresponding pseudoscalar singlet to produce the η and η' . These appear as members of a nonet, which is shown as the middle plane in Fig. 30.1(a). Similarly, the ground-state vector nonet appears as the middle plane in Fig. 30.1(b).

A fourth quark such as charm can be included in this scheme by extending the symmetry to SU(4), as shown in Fig. 30.1. Bottom extends the symmetry to SU(5); to draw the multiplets would require four dimensions.

For the pseudoscalar mesons, the Gell-Mann-Okubo formula is

$$m_\eta^2 = \frac{1}{3}(4m_K^2 - m_\pi^2), \quad (30.2)$$

assuming no octet-singlet mixing. However, the octet η_8 and singlet η_1 mix because of SU(3) breaking. In general, the mixing angle is mass dependent and becomes complex for resonances of finite width.




Figure 30.1: SU(4) 16-plets for the (a) pseudoscalar and (b) vector mesons made of u , d , s , and c quarks. The nonets of light mesons occupy the central planes, to which the $c\bar{c}$ states have been added. The neutral mesons at the centers of these planes are mixtures of $u\bar{u}$, $d\bar{d}$, $s\bar{s}$, and $c\bar{c}$ states.

Neglecting this, the physical states η and η' are given in terms of a mixing angle θ_P by

$$\eta = \eta_8 \cos \theta_P - \eta_1 \sin \theta_P \quad (30.3a)$$

$$\eta' = \eta_8 \sin \theta_P + \eta_1 \cos \theta_P. \quad (30.3b)$$

These combinations diagonalize the mass-squared matrix

$$M^2 = \begin{pmatrix} M_{11}^2 & M_{18}^2 \\ M_{18}^2 & M_{88}^2 \end{pmatrix}, \quad (30.4)$$

where $M_{88}^2 = \frac{1}{3}(4m_K^2 - m_\pi^2)$. It follows that

$$\tan^2 \theta_P = \frac{M_{88}^2 - m_\eta^2}{m_\eta^2 - M_{88}^2}. \quad (30.5)$$

The sign of θ_P is meaningful in the quark model. If

$$\eta_1 = (u\bar{u} + d\bar{d} + s\bar{s})/\sqrt{3} \quad (30.6a)$$

$$\eta_8 = (u\bar{u} + d\bar{d} - 2s\bar{s})/\sqrt{6}, \quad (30.6b)$$

then the matrix element M_{18}^2 , which is due mostly to the strange quark mass, is negative. From the relation

$$\tan \theta_P = \frac{M_{88}^2 - m_\eta^2}{M_{18}^2}, \quad (30.7)$$

we find that $\theta_P < 0$. However, caution is suggested in the use of the η - η' mixing-angle formulas, as they are extremely sensitive to SU(3)

Table 30.2: Suggested $q\bar{q}$ quark-model assignments for most of the known mesons. Some assignments, especially for the 0^{++} multiplet and for some of the higher multiplets, are controversial. Mesons in bold face are included in the Meson Summary Table. Of the light mesons in the Summary Table, the $f_1(1420)$, $f_0(1590)$, $f_J(1710)$, $f_2(2300)$, $f_2(2340)$, and the two peaks in the $\eta(1440)$ entry are not in this table. Within the $q\bar{q}$ model, it is especially hard to find a place for the first three of these f mesons and for one of the $\eta(1440)$ peaks. See the “Note on Non- $q\bar{q}$ Mesons” at the end of the Meson Listings.

$N^{2S+1}L_J$	J^{PC}	$u\bar{d}, u\bar{u}, d\bar{d}$ $I = 1$	$u\bar{u}, d\bar{d}, s\bar{s}$ $I = 0$	$c\bar{c}$ $I = 0$	$b\bar{b}$ $I = 0$	$\bar{s}u, \bar{s}d$ $I = 1/2$	$c\bar{u}, c\bar{d}$ $I = 1/2$	$c\bar{s}$ $I = 0$	$\bar{b}u, \bar{b}d$ $I = 1/2$
1^1S_0	0^{-+}	π	η, η'	η_c		K	D	D_s	B
1^3S_1	1^{--}	ρ	ω, ϕ	$J/\psi(1S)$	$\Upsilon(1S)$	$K^*(892)$	$D^*(2010)$	$D_s^*(2110)$	$B^*(5330)$
1^1P_1	1^{+-}	$\mathbf{b}_1(1235)$	$\mathbf{h}_1(1170), h_1(1380)$	$h_c(1P)$		K_{1B}^\dagger	$D_1(2420)$	$D_{s1}(2536)$	
1^3P_0	0^{++}	$\mathbf{a}_0(980)$	$\mathbf{f}_0(1300), \mathbf{f}_0(980)$	$\chi_{c0}(1P)$	$\chi_{b0}(1P)$	$K_0^*(1430)$			
1^3P_1	1^{++}	$\mathbf{a}_1(1260)$	$\mathbf{f}_1(1285), \mathbf{f}_1(1510)$	$\chi_{c1}(1P)$	$\chi_{b1}(1P)$	K_{1A}^\dagger			
1^3P_2	2^{++}	$\mathbf{a}_2(1320)$	$\mathbf{f}_2(1270), \mathbf{f}_2'(1525)$	$\chi_{c2}(1P)$	$\chi_{b2}(1P)$	$K_2^*(1430)$	$D_2^*(2460)$		
1^1D_2	2^{-+}	$\pi_2(1670)$				$K_2(1770)$			
1^3D_1	1^{--}	$\rho(1700)$	$\omega(1600)$	$\psi(3770)$		$K^*(1680)^\ddagger$			
1^3D_2	2^{--}					$K_2(1820)$			
1^3D_3	3^{--}	$\rho_3(1690)$	$\omega_3(1670), \phi_3(1850)$			$K_3^*(1780)$			
1^3F_4	4^{++}	$a_4(2040)$	$\mathbf{f}_4(2050), f_4(2220)$			$K_4^*(2045)$			
2^1S_0	0^{-+}	$\pi(1300)$	$\eta(1295)$	$\eta_c(2S)$		$K(1460)$			
2^3S_1	1^{--}	$\rho(1450)$	$\omega(1420), \phi(1680)$	$\psi(2S)$	$\Upsilon(2S)$	$K^*(1410)^\ddagger$			
2^3P_2	2^{++}		$f_2(1810), \mathbf{f}_2(2010)$		$\chi_{b2}(2P)$	$K_2^*(1980)$			
3^1S_0	0^{-+}	$\pi(1770)$	$\eta(1760)$			$K(1830)$			

[†] The K_{1A} and K_{1B} are nearly equal (45°) mixes of the $K_1(1270)$ and $K_1(1400)$.

[‡]The $K^*(1410)$ could be replaced by the $K^*(1680)$ as the 2^3S_1 state.

breaking. If we allow $M_{88}^2 = \frac{1}{3}(4m_K^2 - m_\pi^2)(1 + \Delta)$, the mixing angle is determined by

$$\tan^2 \theta_P = 0.0319(1 + 17\Delta) \quad (30.8)$$

$$\theta_P = -10.1^\circ(1 + 8.5\Delta) \quad (30.9)$$

to first order in Δ . A small breaking of the Gell-Mann-Okubo relation can produce a major modification of θ_P .

For the vector mesons, $\pi \rightarrow \rho$, $K \rightarrow K^*$, $\eta \rightarrow \phi$, and $\eta' \rightarrow \omega$, so that

$$\phi = \omega_8 \cos \theta_V - \omega_1 \sin \theta_V \quad (30.10)$$

$$\omega = \omega_8 \sin \theta_V + \omega_1 \cos \theta_V. \quad (30.11)$$

For “ideal” mixing, $\phi = s\bar{s}$, so $\tan \theta_V = 1/\sqrt{2}$ and $\theta_V = 35.3^\circ$. Experimentally, θ_V is near 35° , the sign being determined by a formula like that for $\tan \theta_P$. Following this procedure we find the mixing angles given in Table 30.3.

Table 30.3: Singlet-octet mixing angles for several nonets, neglecting possible mass dependence and imaginary parts. The sign conventions are given in the text. The values of θ_{quad} are obtained from the equations in the text, while those for θ_{lin} are obtained by replacing m^2 by m throughout. Of the two isosinglets in a nonet, the mostly octet one is listed first.

J^{PC}	Nonet members	θ_{quad}	θ_{lin}
0^{-+}	π, K, η, η'	-10°	-23°
1^{--}	$\rho, K^*(892), \phi, \omega$	39°	36°
2^{++}	$a_2(1320), K_2^*(1430), f_2'(1525), f_2(1270)$	28°	26°
3^{--}	$\rho_3(1690), K_3^*(1780), \phi_3(1850), \omega_3(1670)$	29°	28°

supermultiplets, specified by (D, L_N^P) , where D is the dimensionality of the SU(6) representation, L is the total quark orbital angular momentum, and P is the total parity. Supermultiplets contained in bands up to $N = 12$ are given in Ref. 3. The $N = 0$ band, which contains the nucleon and $\Delta(1232)$, consists only of the $(56, 0_0^+)$ supermultiplet. The $N = 1$ band consists only of the $(70, 1_1^-)$ multiplet and contains the negative-parity baryons with masses below about 1.9 GeV. The $N = 2$ band contains five supermultiplets: $(56, 0_2^+)$, $(70, 0_2^+)$, $(56, 2_2^+)$, $(70, 2_2^+)$, and $(20, 1_2^+)$. Baryons belonging to the $(20, 1_2^+)$ supermultiplet are not ever likely to be observed, since a coupling from the ground-state baryons requires a two-quark excitation. Selection rules are similarly responsible for the fact that many other baryon resonances have not been observed [4].

In Table 30.4, quark-model assignments are given for many of the established baryons whose SU(6)⊗O(3) compositions are relatively unmixed. We note that the unestablished resonances $\Sigma(1480)$, $\Sigma(1560)$, $\Sigma(1580)$, $\Sigma(1770)$, and $\Xi(1620)$ in our Baryon Full Listings are too low in mass to be accommodated in most quark models [4,5].

Table 30.4: Quark-model assignments for many of the known baryons in terms of a flavor-spin SU(6) basis. Only the dominant representation is listed. Assignments for some states, especially for the $\Lambda(1810)$, $\Lambda(2350)$, $\Xi(1820)$, and $\Xi(2030)$, are merely educated guesses.

J^P	(D, L_N^P)	S	Octet members			Singlets
$1/2^+$	$(56, 0_0^+)$	$1/2$	$N(939)$	$\Lambda(1116)$	$\Sigma(1193)$	$\Xi(1318)$
$1/2^+$	$(56, 0_2^+)$	$1/2$	$N(1440)$	$\Lambda(1600)$	$\Sigma(1660)$	$\Xi(?)$
$1/2^-$	$(70, 1_1^-)$	$1/2$	$N(1535)$	$\Lambda(1670)$	$\Sigma(1620)$	$\Xi(?)$ $\Lambda(1405)$
$3/2^-$	$(70, 1_1^-)$	$1/2$	$N(1520)$	$\Lambda(1690)$	$\Sigma(1670)$	$\Xi(1820)$ $\Lambda(1520)$
$1/2^-$	$(70, 1_1^-)$	$3/2$	$N(1650)$	$\Lambda(1800)$	$\Sigma(1750)$	$\Xi(?)$
$3/2^-$	$(70, 1_1^-)$	$3/2$	$N(1700)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$
$5/2^-$	$(70, 1_1^-)$	$3/2$	$N(1675)$	$\Lambda(1830)$	$\Sigma(1775)$	$\Xi(?)$
$1/2^+$	$(70, 0_2^+)$	$1/2$	$N(1710)$	$\Lambda(1810)$	$\Sigma(1880)$	$\Xi(?)$ $\Lambda(?)$
$3/2^+$	$(56, 2_2^+)$	$1/2$	$N(1720)$	$\Lambda(1890)$	$\Sigma(?)$	$\Xi(?)$
$5/2^+$	$(56, 2_2^+)$	$1/2$	$N(1680)$	$\Lambda(1820)$	$\Sigma(1915)$	$\Xi(2030)$
$7/2^-$	$(70, 3_3^-)$	$1/2$	$N(2190)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$ $\Lambda(2100)$
$9/2^-$	$(70, 3_3^-)$	$3/2$	$N(2250)$	$\Lambda(?)$	$\Sigma(?)$	$\Xi(?)$
$9/2^+$	$(56, 4_4^+)$	$1/2$	$N(2220)$	$\Lambda(2350)$	$\Sigma(?)$	$\Xi(?)$
Decuplet members						
$3/2^+$	$(56, 0_0^+)$	$3/2$	$\Delta(1232)$	$\Sigma(1385)$	$\Xi(1530)$	$\Omega(1672)$
$1/2^-$	$(70, 1_1^-)$	$1/2$	$\Delta(1620)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$
$3/2^-$	$(70, 1_1^-)$	$1/2$	$\Delta(1700)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$
$5/2^+$	$(56, 2_2^+)$	$3/2$	$\Delta(1905)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$
$7/2^+$	$(56, 2_2^+)$	$3/2$	$\Delta(1950)$	$\Sigma(2030)$	$\Xi(?)$	$\Omega(?)$
$11/2^+$	$(56, 4_4^+)$	$3/2$	$\Delta(2420)$	$\Sigma(?)$	$\Xi(?)$	$\Omega(?)$

The quark model for baryons is extensively reviewed in Ref. 6 and 7.

30.4. Dynamics

Many specific quark models exist, but most contain the same basic set of dynamical ingredients. These include:

- i) A confining interaction, which is generally spin-independent.
- ii) A spin-dependent interaction, modeled after the effects of gluon exchange in QCD. For example, in the S -wave states, there is a spin-spin hyperfine interaction of the form

$$H_{HF} = -\alpha_S M \sum_{i>j} (\vec{\sigma}^i \lambda_a)_i (\vec{\sigma}^j \lambda_a)_j, \quad (30.19)$$

where M is a constant with units of energy, λ_a ($a = 1, \dots, 8$) is the set of SU(3) unitary spin matrices, defined in Sec. 21, on "SU(3) Isoscalar Factors and Representation Matrices," and the sum runs over constituent quarks or antiquarks. Spin-orbit interactions, although allowed, seem to be small.

- iii) A strange quark mass somewhat larger than the up and down quark masses, in order to split the SU(3) multiplets.
- iv) In the case of isoscalar mesons, an interaction for mixing $q\bar{q}$ configurations of different flavors (*e.g.*, $u\bar{u} \leftrightarrow d\bar{d} \leftrightarrow s\bar{s}$), in a manner which is generally chosen to be flavor independent.

These four ingredients provide the basic mechanisms that determine the hadron spectrum.

References:

1. F.E. Close, in *Quarks and Nuclear Forces* (Springer-Verlag, 1982), p. 56.
2. Particle Data Group, Phys. Lett. **111B** (1982).
3. R.H. Dalitz and L.J. Reinders, in *Hadron Structure as Known from Electromagnetic and Strong Interactions, Proceedings of the Hadron '77 Conference* (Veda, 1979), p. 11.
4. N. Isgur and G. Karl, Phys. Rev. **D18**, 4187 (1978); *ibid.* **D19**, 2653 (1979); *ibid.* **D20**, 1191 (1979); K.-T. Chao, N. Isgur, and G. Karl, Phys. Rev. **D23**, 155 (1981).
5. C.P. Forsyth and R.E. Cutkosky, Z. Phys. **C18**, 219 (1983).
6. A.J.G. Hey and R.L. Kelly, Phys. Reports **96**, 71 (1983). Also see S. Gasiorowicz and J.L. Rosner, Am. J. Phys. **49**, 954 (1981).
7. N. Isgur, Int. J. Mod. Phys. **E1**, 465 (1992); G. Karl, Int. J. Mod. Phys. **E1**, 491 (1992).

31. NAMING SCHEME FOR HADRONS

Table 31.1: Symbols for mesons with the strangeness and all heavy-flavor quantum numbers equal to zero.

J^{PC}	$\begin{cases} 0^{-+} & 1^{+-} & 1^{--} & 0^{++} \\ 2^{-+} & 3^{+-} & 2^{--} & 1^{++} \\ \vdots & \vdots & \vdots & \vdots \end{cases}$				
$q\bar{q}$ content	$2S+1L_J$	${}^1(L\text{ even})_J$	${}^1(L\text{ odd})_J$	${}^3(L\text{ even})_J$	${}^3(L\text{ odd})_J$
$u\bar{d}, u\bar{u} - \bar{d}\bar{d}, \bar{d}\bar{u}$ ($I=1$)	π	b	ρ	a	
$d\bar{d} + u\bar{u}$ and/or $s\bar{s}$ } ($I=0$)	η, η'	h, h'	ω, ϕ	f, f'	
$c\bar{c}$	η_c	h_c	ψ^\dagger	χ_c	
$b\bar{b}$	η_b	h_b	Υ	χ_b	
$t\bar{t}$	η_t	h_t	θ	χ_t	

[†]The J/ψ remains the J/ψ .

31.1. Introduction

We introduced in the 1986 edition [1] a new naming scheme for the hadrons. Changes from older terminology affected mainly the heavier mesons made of the light ($u, d,$ and s) quarks. Otherwise, the only important change to known hadrons was that the F^\pm became the D_s^\pm . None of the lightest pseudoscalar or vector mesons changed names, nor did the $c\bar{c}$ or $b\bar{b}$ mesons (we do, however, now use χ_c for the $c\bar{c}$ χ states), nor did any of the established baryons. The Summary Tables give both the new and old names whenever a change has occurred.

We follow custom and use spectroscopic names such as $\Upsilon(1S)$ as the primary name for most of those $\psi, \Upsilon,$ and χ states whose spectroscopic identity is known. We use the form $\Upsilon(9460)$ as an alternate, and as the primary name when the spectroscopic identity is not known.

31.2. “Neutral-flavor” mesons ($S=C=B=T=0$)

Table 31.1 shows the naming scheme for mesons having the strangeness and all heavy-flavor quantum numbers equal to zero. The scheme is designed for all mesons, whether ordinary or exotic. (This isn't quite true. We haven't proposed names for mesons whose charge $Q,$ strangeness $S,$ or other *additive* quantum numbers can't be matched by a $q\bar{q}$ state. For example, we have no name for a meson with $Q=2,$ or for one with $Q=-1$ and $S=+1.$)

First, we assign names to those states with quantum numbers compatible with being $q\bar{q}$ states. The rows of the Table give the possible $q\bar{q}$ content. The columns give the possible parity/charge-conjugation states, $PC = -+, +-, --,$ and $++;$ these combinations correspond one-to-one with the

angular-momentum state $2S+1L_J$ of the $q\bar{q}$ system being ${}^1(L\text{ even})_J, {}^1(L\text{ odd})_J, {}^3(L\text{ even})_J,$ or ${}^3(L\text{ odd})_J.$ Here $S, L,$ and J are the spin, orbital, and total angular momenta of the $q\bar{q}$ system. The relations between the quantum numbers are $P = (-1)^{L+1}, C = (-1)^{L+S},$ and G parity $= (-1)^{L+S+I},$ where of course the C quantum number is only relevant to neutral mesons.

The entries in the Table give the particle symbols. The meson spin J is added as a subscript except for pseudoscalar and vector mesons, and the mass is added in parentheses for any meson that decays strongly. However, for the lightest meson resonances, we sometimes omit the mass, as in ρ for $\rho(770), \phi$ for $\phi(1020),$ etc.

Measurements of the mass, quark content (where relevant), and quantum numbers $I, J, P,$ and C (or G) of a meson thus fix its symbol. Conversely, these properties may be inferred unambiguously from the symbol.

If the main symbol cannot be assigned because the quantum numbers are unknown, X is used. Sometimes it is not known whether a meson is mainly the isospin-0 mix of $u\bar{u}$ and $d\bar{d}$ or is mainly $s\bar{s}.$ A prime (or symbol ϕ) may be used to distinguish two such mixing states.

Names are assigned for $t\bar{t}$ mesons, even though the top quark is evidently so heavy that it is expected to decay too rapidly for bound states to form.

Gluonium states or other mesons that are not $q\bar{q}$ states are, if the quantum numbers are *not* exotic, to be named just as are the $q\bar{q}$ mesons. Such states will probably be difficult to distinguish from $q\bar{q}$ states and will likely mix with them, and we make no attempt to distinguish those “mostly gluonium” from those “mostly $q\bar{q}.$ ”

An “exotic” meson with J^{PC} quantum numbers that a $q\bar{q}$ system cannot have, namely $J^{PC} = 0^{--}, 0^{+-}, 1^{-+}, 2^{+-}, 3^{-+}, \dots,$ would use the same symbol as does an ordinary meson with all the same quantum numbers as the exotic meson except for the C parity. Then a caret or “hat” is added; for example, an isospin-1 0^{--} meson would be a $\hat{\pi},$ an isospin-0 1^{-+} meson would be an $\hat{\omega}.$

The results of all this are as follows. Established mesons whose names changed slightly in 1986 are:

Old name	New name	Old name	New name
$H(1190)$	$h_1(1170)$	$A_2(1320)$	$a_2(1320)$
$B(1235)$	$b_1(1235)$	$f'(1525)$	$f'_2(1525)$
$A_1(1270)$	$a_1(1260)$	$\omega(1670)$	$\omega_3(1670)$
$f(1270)$	$f_2(1270)$		

Established mesons whose names changed completely are:

Old name	New name	Old name	New name
$S(975)$	$f_0(980)$	$g(1690)$	$\rho_3(1690)$
$\delta(980)$	$a_0(980)$	$\theta(1690)$	$f_J(1710)$
$D(1285)$	$f_1(1285)$	$X(1850)$	$\phi_3(1850)$
$\epsilon(1200)$	$f_0(1300)$	$g_T(2010)$	$f_2(2010)$
$E(1420)$	$f_1(1420)$	$h(2030)$	$f_4(2050)$
$\iota(1440)$	$\eta(1440)$	$g'_T(2300)$	$f_2(2300)$
$D(1530)$	$f_1(1510)$	$g''_T(2340)$	$f_2(2340)$
$A_3(1680)$	$\pi_2(1670)$		

The old $S(975), D(1285), \epsilon(1200), E(1420), D(1530), \theta(1690), g_T(2010), h(2030), g'_T(2300),$ and $g''_T(2340)$ all became f mesons; the new scheme revealed that they are all $PC = ++, {}^3(L\text{ odd})_J$ states.

31.3. Mesons with nonzero $S, C, B,$ and/or T

Since the strangeness or a heavy flavor of these mesons is nonzero, none of them are eigenstates of charge conjugation, and in each of them one of the quarks is heavier than the other. The rules are:

1. The main symbol is an upper-case italic letter indicating the heavier quark as follows:

$$s \rightarrow \bar{K} \quad c \rightarrow D \quad b \rightarrow \bar{B} \quad t \rightarrow T.$$

We use the convention that *the flavor and the charge of a quark have the same sign.* Thus the strangeness of the s quark is negative, the charm of the c quark is positive, and the bottom of the b quark is negative. In addition, I_3 of the u and d quarks are positive and negative, respectively. The effect of this convention is as follows: *Any flavor carried by a charged meson has the same sign as its charge.* Thus the $K^+, D^+,$ and B^+ have positive strangeness, charm, and bottom, respectively, and all have positive $I_3.$ The D_s^+ has positive charm *and* strangeness. Furthermore, the $\Delta(\text{flavor}) = \Delta Q$ rule, best known for the kaons, applies to every flavor.

2. If the lighter quark is not a u or a d quark, its identity is given by a subscript. The D_s^+ is an example.
3. If the spin-parity is in the “normal” series, $J^P = 0^+, 1^-, 2^+, \dots,$ a superscript “*” is added.
4. The spin is added as a subscript unless the meson is a pseudoscalar or a vector particle.

Thus the pseudoscalar and vector K , K^* , D , D^* , and B mesons did not change names. Established mesons whose names did change were:

Old name	New name	Old name	New name
$Q_1(1280)$	$K_1(1270)$	$L(1770)$	$K_2(1770)$
$Q_2(1400)$	$K_1(1400)$	$K^*(1780)$	$K_3^*(1780)$
$\kappa(1350)$	$K_0^*(1430)$	$K^*(2060)$	$K_4^*(2045)$
$K^*(1430)$	$K_2^*(1430)$	F	D_s

Most notably, the F (the $c\bar{s}$ state) became the D_s .

31.4. Baryons

The symbols N , Δ , Λ , Σ , Ξ , and Ω used for 30 years for the baryons made of light quarks (u , d , and s quarks) tell the isospin and quark content, and the same information is conveyed by the symbols now used for the baryons containing one or more heavy quarks (c , b , and t quarks). The rules are:

1. Baryons with *three* u and/or d quarks are N 's (isospin 1/2) or Δ 's (isospin 3/2).
2. Baryons with *two* u and/or d quarks are Λ 's (isospin 0) or Σ 's (isospin 1). If the third quark is a c , b , or t quark, its identity is given by a subscript.
3. Baryons with *one* u or d quark are Ξ 's (isospin 1/2). One or two subscripts are used if one or both of the remaining quarks are heavy: thus Ξ_c , Ξ_{cc} , Ξ_b , *etc.*
4. Baryons with *no* u or d quarks are Ω 's (isospin 0), and subscripts indicate any heavy-quark content.

In short, the number of u plus d quarks together with the isospin determine the main symbol, and subscripts indicate any content of heavy quarks. A Σ always has isospin 1, an Ω always has isospin 0, *etc.*

Reference:

1. Particle Data Group: M. Aguilar-Benitez *et al.*, Phys. Lett. **170B** (1986).

32. MONTE CARLO PARTICLE NUMBERING SCHEME

Written April 1988 by G.R. Lynch and T.G. Trippe.

Most particle physics Monte Carlo and analysis systems use a numbering scheme to represent particles. The lack of standardization of such schemes inhibits interfacing different programs. The following table proposes a standard numbering scheme. Some of the properties of this scheme are:

1. Quarks and leptons are ordered by family, and within the family, by isospin. This puts the u and d in the opposite order than is often used in other numbering schemes. In our scheme we call the highest numbered quark the heaviest quark.
2. For multiple quark systems (mesons, baryons, and diquarks), the rightmost digit is generally $L = 2J + 1$. (The K_S^0 and K_L^0 are exceptions.) Particles with $J > 4$ have not been assigned numbers.
3. Mesons are represented by the form NML and baryons by $NMKL$, where N , M , and K are quark numbers.
4. For these systems the heaviest quark is usually on the left and the quarks are in decreasing mass order from left to right. One exception to this convention is the K_L^0 - K_S^0 pair. A second exception is for the A 's for which we invert the up and down quarks to distinguish the A from the Σ^0 .
5. The other exception to this mass order rule is for some N 's and Δ 's. For N 's, the u and d quark are reversed for spins $3/2$ and $7/2$. For Δ 's, they are reversed for spins $1/2$ and $5/2$. The quarks are in the normal decreasing order when $I + J$ is odd.
6. Mesons, and only mesons, have the third digit nonzero and the fourth digit zero. (We designate the rightmost digit as the first digit.)
7. Only baryons and diquarks have the fourth digit nonzero.
8. Only quarks and diquarks have the second digit equal to zero.
9. Particles have positive numbers; each antiparticle has the negative of its counterpart.

10. The particle-antiparticle convention is the one used by the Particle Data Group, so that the K^+ and B^+ are particles.
11. The above rules imply that for mesons (as opposed to anti-mesons), when the number of the leftmost (heaviest) quark is even, it is a quark, and when the number of the leftmost quark is odd, it is an antiquark.
12. The gluon has two numbers. Its official number is 21 to place it with the other gauge bosons. Its number is also 9 so that a glueball is specified as 99.
13. The fifth digit is used to differentiate different particles with the same quark content and spin.
14. Although isospin is not manifest in this scheme, the isospin of any hadron can be determined from the number. Mesons with $11L$ are isospin 1 and those with $22L$ are isospin 0. For nonstrange baryons, if the quarks are in the normal decreasing order, then $I + J$ is odd, otherwise $I + J$ is even. If a strange baryon does not have the normal decreasing quark order, it has $I = 0$.

More details about the motivation behind, and properties of, this scheme can be found in Ref. 1. Although this scheme has the advantage that a particle's number has considerable physics content, it has the disadvantage that it is not compact. An algorithm that translates this scheme into a more compact scheme is needed for its implementation. Contact the Berkeley Particle Data Group for further information on such an algorithm.

A list of particle numbers follows.

Reference:

1. T.G. Trippe and G.R. Lynch, "Particle I.D. Numbers, Decay Tables, and Other Possible Contributions of the Particle Data Group to Monte Carlo Standards," LBL-24287, in *Proceedings of the Workshop on Detector Simulation for the SSC* (August 1987).

QUARKS

d	1
u	2
s	3
c	4
b	5
t	6

GAUGE AND HIGGS BOSONS

γ	22
W	24
Z	23
g	21 and 9
H_1^0	25
H_2^0	35
H_3^0	36
H^+	37

LEPTONS

ν_e	12
ν_μ	14
ν_τ	16
e	11
μ	13
τ	15

DIQUARKS

$(dd)_1$	1103
$(ud)_0$	2101
$(ud)_1$	2103
$(uu)_1$	2203
$(sd)_0$	3101
$(sd)_1$	3103
$(su)_0$	3201
$(su)_1$	3203

MESONS

π^+	211
π^0	111
η	221
$\rho(770)$	113, 213
$\omega(782)$	223
$\eta'(958)$	331
$f_0(980)$	10221
$a_0(980)$	10111, 10211
$\phi(1020)$	333
$h_1(1170)$	10223
$b_1(1235)$	10113, 10213
$a_1(1260)$	20113, 20213
$f_2(1270)$	225
$f_1(1285)$	20223
$\eta(1295)$	20221
$f_0(1300)$	30221

MESONS (Cont'd)

$\pi(1300)$	20111, 20211
$a_2(1320)$	115, 215
$f_1(1420)$	30223
$\omega(1420)$	50223
$\eta(1440)$	40221
$\rho(1450)$	40113, 40213
$f_1(1510)$	40223
$f_2'(1525)$	335
$f_0(1590)$	50221
$\omega(1600)$	60223
$\omega_3(1670)$	227
$\pi_2(1670)$	10115, 10215
$\phi(1680)$	10333
$\rho_3(1690)$	117, 217
$\rho(1700)$	30113, 30213
$f_7(1710)$	30113, 30213
$\phi_3(1850)$	337
$f_2(2010)$	20225
$f_4(2050)$	229
$f_2(2300)$	30225
$f_2(2340)$	40225
K^+	321
K^0	311
K_S^0	310
K_L^0	130

MESONS (Cont'd)

$K^*(892)$	313, 323
$K_1(1270)$	10313, 10323
$K_1(1400)$	20313, 20323
$K^*(1410)$	30313, 30323
$K_0^*(1430)$	10311, 10321
$K_2^*(1430)$	315, 325
$K^*(1680)$	40313, 40323
$K_2(1770)$	10315, 10325
$K_3^*(1780)$	317, 327
$K_2(1820)$	20315, 20325
$K_4^*(2045)$	319, 329
D^+	411
D^0	421
$D^*(2007)^0$	423
$D^*(2010)^+$	413
$D_1(2420)^0$	10423
$D_2^*(2460)$	425, 415
D_s^+	431
D_s^{*+}	433
$D_{s1}(2536)^+$	10433
B^+	521
B^0	511
B^*	513, 523
B_s^0	531
$\eta_c(1S)$	441
$J/\psi(1S)$	443
$\chi_{c0}(1P)$	10441
$\chi_{c1}(1P)$	10443
$\chi_{c2}(1P)$	445
$\psi(2S)$	20443
$\psi(3770)$	30443
$\psi(4040)$	40443
$\psi(4160)$	50443
$\psi(4415)$	60443
$\Upsilon(1S)$	553
$\chi_{b0}(1P)$	551
$\chi_{b1}(1P)$	10553
$\chi_{b2}(1P)$	555
$\Upsilon(2S)$	20553
$\chi_{b0}(2P)$	10551
$\chi_{b1}(2P)$	70553
$\chi_{b2}(2P)$	10555
$\Upsilon(3S)$	30553
$\Upsilon(4S)$	40553
$\Upsilon(10860)$	50553
$\Upsilon(11020)$	60553

BARYONS

p	P_{11}	2212
n	P_{11}	2112
$N(1440)$	P_{11}	12112, 12212
$N(1520)$	D_{13}	1214, 2124
$N(1535)$	S_{11}	22112, 22212

BARYONS (Cont'd)

$N(1650)$	S_{11}	32112, 32212
$N(1675)$	D_{15}	2116, 2216
$N(1680)$	F_{15}	12116, 12216
$N(1700)$	D_{13}	21214, 22124
$N(1710)$	P_{11}	42112, 42212
$N(1720)$	P_{13}	31214, 32124
$N(2190)$	G_{17}	1218, 2128
$\Delta(1232)$	P_{33}	1114, 2114, 2214, 2224
$\Delta(1600)$	P_{33}	31114, 32114, 32214, 32224
$\Delta(1620)$	S_{31}	1112, 1212, 2122, 2222
$\Delta(1700)$	D_{33}	11114, 12114, 12214, 12224
$\Delta(1900)$	S_{31}	11112, 11212, 12122, 12222
$\Delta(1905)$	F_{35}	1116, 1216, 2126, 2226
$\Delta(1910)$	P_{31}	21112, 21212, 22122, 22222
$\Delta(1920)$	P_{33}	21114, 22114, 22214, 22224
$\Delta(1930)$	D_{35}	11116, 11216, 12126, 12226
$\Delta(1950)$	F_{37}	1118, 2118, 2218, 2228
Λ	P_{01}	3122
$\Lambda(1405)$	S_{01}	13122
$\Lambda(1520)$	D_{03}	3124
$\Lambda(1600)$	P_{01}	23122
$\Lambda(1670)$	S_{01}	33122
$\Lambda(1690)$	D_{03}	13124
$\Lambda(1800)$	S_{01}	43122
$\Lambda(1810)$	P_{01}	53122
$\Lambda(1820)$	F_{05}	3126
$\Lambda(1830)$	D_{05}	13126
$\Lambda(1890)$	P_{03}	23124
$\Lambda(2100)$	G_{07}	3128
$\Lambda(2110)$	F_{05}	23126
Σ^+	P_{11}	3222
Σ^0	P_{11}	3212
Σ^-	P_{11}	3112
$\Sigma(1385)$	P_{13}	3114, 3214, 3224
$\Sigma(1660)$	P_{11}	13112, 13212, 13222
$\Sigma(1670)$	D_{13}	13114, 13214, 13224
$\Sigma(1750)$	S_{11}	23112, 23212, 23222
$\Sigma(1775)$	D_{15}	3116, 3216, 3226
$\Sigma(1915)$	F_{15}	13116, 13216, 13226
$\Sigma(1940)$	D_{13}	23114, 23214, 23224
$\Sigma(2030)$	F_{17}	3118, 3218, 3228
Ξ^0	P_{11}	3322
Ξ^-	P_{11}	3312
$\Xi(1530)$	P_{13}	3314, 3324
$\Xi(1820)$	D_{13}	13314, 13324
Ω^-		3334
Λ_c^+		4122
$\Sigma_c(2455)$		4112, 4212, 4222
Ξ_c^+		4322
Ξ_c^0		4312
Ω_c^0		4332
Λ_b^0		5122

33. PLOTS OF CROSS SECTIONS AND RELATED QUANTITIES

NOTE: THE FIGURES IN THIS SECTION ARE INTENDED TO SHOW THE REPRESENTATIVE DATA.
THEY ARE NOT MEANT TO BE COMPLETE COMPILATIONS OF ALL THE WORLD'S RELIABLE DATA

Structure Functions




Figure 32.1: The proton structure function F_2^p measured in electromagnetic scattering of 26.7 GeV electrons on 820 GeV protons by the HERA experiments, versus x , for fixed bins of Q^2 . The closed symbols show H1 and the open symbols ZEUS data obtained with luminosities of 22.5 nb^{-1} and 24.7 nb^{-1} , respectively. A QCD prescription for $R = \sigma_L/\sigma_T$ was used to extract the structure function from the measured cross sections. The inner error bars are statistical, the outer error bars are statistical and systematic errors combined in quadrature. An overall normalization error of $\approx 8\%$ is not shown in the figure. References: H1—I. Abt *et al.*, Nucl. Phys. **B407**, 515 (1993) and G. Bernardi, *Proceedings International Europhysics Conference on High Energy Physics*, (Marseille 1993), ed. by J. Carr and M. Perrottet; ZEUS—M. Derrick *et al.*, Phys. Lett. **B316**, 412 (1993). (Courtesy of M. Virchaux and R. Voss, 1994.)




Figure 32.2: The nucleon structure function F_2 measured in deep inelastic scattering of electrons (SLAC), muons (BCDMS, NMC), and neutrinos (CCFR) on deuterium (BCDMS, NMC, SLAC), carbon (BCDMS C), and iron (CCFR Fe) targets. The data are shown versus Q^2 , for fixed bins of x , and have been scaled by the factors shown in parentheses for convenience in plotting. The error bars show statistical and systematic errors combined in quadrature. The heavy target data were corrected for nuclear effects and the neutrino data for electromagnetic quark charges and the excess of the strange quark sea. The overall normalization of the BCDMS, CCFR, and NMC data has been adjusted by typically 1%. The solid line represents a perturbative QCD fit which includes a parameterization of higher twist effects (M. Virchaux and A. Milsztajn, Phys. Lett. **B274**, 221 (1992)). References: BCDMS—A.C. Benvenuti *et al.*, Phys. Lett. **B237**, 592 (1990); BCDMS C—A.C. Benvenuti *et al.*, Phys. Lett. **B195**, 91 (1987); CCFR—S.R. Mishra *et al.*, NEVIS-1465 (1992); NMC—P. Amaudruz *et al.*, Phys. Lett. **B295**, 159 (1992); SLAC—L.W. Whitlow *et al.*, Phys. Lett. **B282**, 475 (1992). (Courtesy of M. Virchaux and R. Voss, 1994.)

Structure Functions




Figure 32.3: The deuteron structure function F_2^d measured in electromagnetic scattering of electrons (SLAC) and muons (BCDMS, NMC) on deuterium targets, versus Q^2 , for fixed bins of x . The data have been scaled by the factors shown in parentheses for convenience in plotting. The error bars show statistical and systematic errors combined in quadrature. A phenomenological parameterization of $R = \sigma_L/\sigma_T$ (L.W. Whitlow *et al.*, Phys. Lett. **B250**, 193 (1990)) was used in the analysis of the SLAC and the NMC data and a QCD prescription for R in the analysis of the BCDMS data. Where necessary, the SLAC and BCDMS data were interpolated to the x bins of the NMC data; SLAC and BCDMS data at $x > 0.5$ are not shown in this figure. The solid lines represent a QCD inspired phenomenological fit. References: **BCDMS**—A.C. Benvenuti *et al.*, Phys. Lett. **B237**, 592 (1990); **NMC**—P. Amaudruz *et al.*, Phys. Lett. **B295**, 159 (1992); **SLAC**—L.W. Whitlow *et al.*, Phys. Lett. **B282**, 475 (1992). Similar data are available for the proton structure function F_2^p (NMC and SLAC—same references; BCDMS—A.C. Benvenuti *et al.*, Phys. Lett. **B223**, 485 (1989)). (Courtesy of M. Virchaux and R. Voss, 1994.)

Structure Functions




Figure 32.4: The spin-dependent structure function g_1 measured in deep inelastic scattering of polarised electrons (b) and muons (a,c) on polarised protons (a, EMC and SMC), neutrons (b, E142), and deuterons (c, SMC). For the muon data, only statistical errors are shown with the data points and the size of the systematic errors is indicated by the shaded bands. For the electron data, statistical and systematic errors are combined in quadrature. The data are shown versus x ; for each of the four experiments, the Q^2 of the data increases with increasing x . The average Q^2 is also different for each of the four experiments; all data in this plot are at $1 \text{ GeV}^2 < Q^2 < 60 \text{ GeV}^2$. References: **E142**—P.L. Anthony *et al.*, Phys. Rev. Lett. **71**, 959 (1993); **EMC**—J. Ashman *et al.*, Nucl. Phys. **B328**, 1 (1989); **SMC**—B. Adeva *et al.*, Phys. Lett. **B302**, 533 (1993); D. Adams *et al.*, CERN-PPE/94-57. (Courtesy of M. Virchaux and R. Voss, 1994.)

Jet Production in pp and $p\bar{p}$ Interactions




Figure 32.5: Differential cross sections for observation of a single jet of pseudorapidity $\eta = 0$ as a function of the jet transverse momentum. CDF—F. Abe *et al.*, Phys. Rev. Lett. **70**, 1376 (1993); UA1—G. Arnison *et al.*, Phys. Lett. **B172**, 461 (1986); UA2—J. Alitti *et al.*, Phys. Lett. **B257**, 232 (1991); R807—T. Akesson *et al.*, Phys. Lett. **B123**, 133 (1983). (Courtesy of S. Geer, FNAL, 1994.)

Direct γ Production in $p\bar{p}$ Interactions




Figure 32.6: Differential cross sections for observation of a single photon of pseudorapidity $\eta = 0$ as a function of the photon transverse momentum UA1—C. Albajar *et al.*, Phys. Lett. **B209**, 385 (1988); UA2—J. Alitti *et al.*, Phys. Lett. **B288**, 386 (1992); CDF—Fermilab-CONF-94/148-E. (Courtesy of S. Geer, FNAL, 1994.)

Pseudorapidity Distributions in $p\bar{p}$ Interactions




Figure 32.7: Charge particle pseudorapidity distributions in $p\bar{p}$ collisions for $53 \text{ GeV} \leq \sqrt{s} \leq 900 \text{ GeV}$. The number per pseudorapidity interval is about 10% higher if the rate is normalized excluding singly diffractive events rather than to the total inelastic rate. $Sp\bar{p}S$ data are from G.J. Alner *et al.*, Z. Phys. **C33**, 1 (1986), and ISR data are from K. Alpgård *et al.*, Phys. Lett. **112B**, 193 (1982). CDF nonsingle-diffractive results at $\sqrt{s} = 630$ and 1800 GeV are given in F. Abe *et al.*, Phys. Rev. **D41**, 2330 (1990). (Courtesy of D.R. Ward, Cambridge Univ., 1991.)

Average Hadron Multiplicities in e^+e^- Annihilation Events

Table 33.1: Average hadron multiplicity per e^+e^- annihilation event at $\sqrt{s} \approx 10, 29\text{--}35,$ and 91 GeV. The rates given include decay products from resonances with $cr < 10$ cm, and include charge conjugated states. (Updated April 1994 by S. Bethke and O. Biebel.)

Particle	$\sqrt{s} \approx 10$ GeV	$\sqrt{s} = 29\text{--}35$ GeV	$\sqrt{s} = 91$ GeV
Pseudoscalar mesons:			
π^+	6.6 \pm 0.2	10.3 \pm 0.4	17.1 \pm 0.4
π^0	3.2 \pm 0.3	5.6 \pm 0.3	9.9 \pm 0.08 ^(a)
K^+	0.90 \pm 0.04	1.48 \pm 0.09	2.42 \pm 0.13
K^0	0.91 \pm 0.05	1.48 \pm 0.07	2.12 \pm 0.06
η	0.20 \pm 0.04	0.61 \pm 0.07	0.73 \pm 0.07 ^(a)
$\eta'(958)$	0.03 \pm 0.01	0.26 \pm 0.10	0.17 \pm 0.05 ^(a)
D^+	0.16 \pm 0.03	0.17 \pm 0.03	0.20 \pm 0.03
D^0	0.37 \pm 0.06	0.45 \pm 0.07	0.40 \pm 0.06
B^-, B^0	—	—	0.34 \pm 0.06 ^(b)
Scalar mesons:			
$f_0(980)$	0.024 \pm 0.006	0.11 \pm 0.04	0.14 \pm 0.06 ^(a)
Vector mesons:			
$\rho(770)^0$	0.35 \pm 0.04	0.81 \pm 0.08	1.4 \pm 0.1 ^(c)
$K^*(892)^+$	0.27 \pm 0.03	0.64 \pm 0.05	0.78 \pm 0.08
$K^*(892)^0$	0.29 \pm 0.03	0.56 \pm 0.06	0.77 \pm 0.09
$\phi(1020)$	0.044 \pm 0.006	0.085 \pm 0.011	0.086 \pm 0.018
$D^*(2010)^+$	0.22 \pm 0.04	0.43 \pm 0.07	0.17 \pm 0.02
$D^*(2007)^0$	0.23 \pm 0.06	0.27 \pm 0.11	—
$\omega(782)$	0.30 \pm 0.08	—	—
$J/\psi(1S)$	—	—	0.0036 \pm 0.0006
Pseudovector mesons:			
$\chi_{c1}(1P)$	—	—	0.008 \pm 0.003
Tensor mesons:			
$f_2(1270)$	0.09 \pm 0.02	0.14 \pm 0.04	0.31 \pm 0.12
$K_2^*(1430)^+$	—	0.09 \pm 0.03	—
$K_2^*(1430)^0$	—	0.12 \pm 0.06	—
Baryons:			
p	0.253 \pm 0.016	0.640 \pm 0.050	0.92 \pm 0.11
Λ	0.080 \pm 0.007	0.205 \pm 0.010	0.348 \pm 0.013
Σ^0	0.023 \pm 0.008	—	—
$\Delta(1232)^{++}$	0.040 \pm 0.010	—	—
Ξ^-	0.0059 \pm 0.0007	0.0176 \pm 0.0027	0.0238 \pm 0.0024
$\Xi(1530)^0$	0.0015 \pm 0.006	—	0.0063 \pm 0.0014
$\Sigma(1385)^-$	0.006 \pm 0.002	0.017 \pm 0.004	—
$\Sigma(1385)^+$	0.005 \pm 0.001	0.017 \pm 0.004	—
$\Sigma(1385)^\pm$	0.0106 \pm 0.0020	0.033 \pm 0.008	0.0380 \pm 0.0062
Ω^-	0.0007 \pm 0.0004	0.014 \pm 0.007	0.0051 \pm 0.0013
Λ_c^+	0.100 \pm 0.030 ^(d)	0.110 \pm 0.050	—
Λ_b^0	—	—	0.031 \pm 0.016
$\Sigma_c^{++}, \Sigma_c^0$	0.014 \pm 0.007	—	—
$\Lambda(1520)$	0.008 \pm 0.002	—	—

- (a) Extrapolation to the unobserved region using the shape predicted by JETSET.
(b) The Standard Model branching ratio $B(Z \rightarrow b\bar{b}) = 0.220$ was used.
(c) Mass of $\rho(770)$ determined to 757 ± 2 MeV/ c^2 . The reason for that is possibly the Bose-Einstein effect affecting the pions from the $\rho(770)$ decay.
(d) The value was taken from the cross section of the $\Lambda_c^+ \rightarrow p\pi K$, assuming the branching fraction to be $(3.2 \pm 0.7)\%$ (RPP 1992).

References:

- RPP, Phys. Rev. **D45** (1992) and references therein
R. Marshall, Rept. on Prog. in Phys. **52**, 1329 (1989)
A. De Angelis, J. Phys. **G19**, 1233 (1993) and references therein
Ch. Bieler *et al.*, (Crystal Ball), Z. Phys. **C49**, 225 (1991)
H. Albrecht *et al.*, (ARGUS), Z. Phys. **C46**, 15 (1990)
D.D. Pietzl *et al.*, (JADE), Z. Phys. **C46**, 1 (1990)
H.J. Behrend *et al.*, (CELLO), Z. Phys. **C47**, 1 (1990)
H. Aihara *et al.*, (TPC), Phys. Rev. Lett. **53**, 2378 (1984)
H. Schellman *et al.*, (MARK II), Phys. Rev. **D31**, 3013 (1985)
M. Derrick *et al.*, (HRS), Phys. Rev. **D35**, 2639 (1987)
Ch. Berger *et al.*, (PLUTO), Phys. Lett. **104B**, 79 (1981)
W. Bartel *et al.*, (JADE), Z. Phys. **C20**, 187 (1983)
H. Aihara *et al.*, (TASSO), Z. Phys. **C27**, 27 (1985)
H.J. Behrend *et al.*, (CELLO), Z. Phys. **C46**, 397 (1990)
P. Abreu *et al.*, (DELPHI), Z. Phys. **C59**, 533 (1993)
P. Abreu *et al.*, (DELPHI), Z. Phys. **C57**, 181 (1993)
H. Albrecht *et al.*, (ARGUS), Z. Phys. **C58**, 199 (1993)
S. Abachi *et al.*, (HRS), Phys. Rev. Lett. **57**, 1990 (1986)
A. Albrecht *et al.*, (ARGUS), Z. Phys. **C61**, 1 (1994)
P.D. Acton *et al.*, (OPAL), Z. Phys. **C56**, 521 (1992)
O. Adriani *et al.*, (L3), Phys. Lett. **B317**, 467 (1993)
H. Albrecht *et al.*, (ARGUS), Z. Phys. **C44**, 547 (1989)
H. Albrecht *et al.*, (ARGUS), Phys. Lett. **B230**, 169 (1989)
R. Akers *et al.*, (OPAL), CERN PPE/94-49 (1994)

Fragmentation in e^+e^- Annihilation




Figure 32.8: Fragmentation into π^\pm in e^+e^- annihilations: Inclusive cross sections $(1/\sigma\beta)(d\sigma/dx)$, with $x = 2E/\sqrt{s}$. The indicated errors are statistical and systematic errors added in quadrature. Δ : rate at $\sqrt{s} = 9.98$ GeV; an overall uncertainty of 1.8% [H. Albrecht *et al.* (ARGUS), *Z. Phys.* C44, 547 (1989)]. \square : rate at $\sqrt{s} = 29$ GeV [TPC—H. Aihara *et al.*, *Phys. Rev. Lett.* 61, 1263 (1988)]. \circ : rate for hadronic decays of the Z at $\sqrt{s} = 91.2$ GeV [OPAL—R. Akers *et al.*, CERN PPE/94-49 (1994)]. (Courtesy of O. Biebel, S. Bethke, and D. Lanske, RWTH, Aachen, 1994.)




Figure 32.9: Fragmentation into $p\bar{p}$ in e^+e^- annihilations: Inclusive cross sections $(1/\sigma\beta)(d\sigma/dx)$, with $x = 2E/\sqrt{s}$. The indicated errors are statistical and systematic errors added in quadrature. Δ : rate at $\sqrt{s} = 9.98$ GeV; an overall uncertainty of 1.8%. This rate is obtained from the measured \bar{p} rate by scaling with a factor of two [H. Albrecht *et al.* (ARGUS), *Z. Phys.* C44, 547 (1989)]. \square : rate at $\sqrt{s} = 29$ GeV [TPC—H. Aihara *et al.*, *Phys. Rev. Lett.* 61, 1263 (1988)]. \circ : rate for hadronic decays of the Z at $\sqrt{s} = 91.2$ GeV [OPAL—R. Akers *et al.*, CERN PPE/94-49 (1994)]. (Courtesy of O. Biebel, S. Bethke, and D. Lanske, RWTH, Aachen, 1994.)




Figure 32.10: Fragmentation into K^\pm in e^+e^- annihilations: Inclusive cross sections $(1/\sigma\beta)(d\sigma/dx)$, with $x = 2E/\sqrt{s}$. The indicated errors are statistical and systematic errors added in quadrature. Δ : rate at $\sqrt{s} = 9.98$ GeV; an overall uncertainty of 1.8% [ARGUS—H. Albrecht *et al.*, *Z. Phys.* C44, 547 (1989)]. \square : rate at $\sqrt{s} = 29$ GeV [TPC—H. Aihara *et al.*, *Phys. Rev. Lett.* 61, 1263 (1988)]. \circ : rate for hadronic decays of the Z at $\sqrt{s} = 91.2$ GeV [OPAL—R. Akers *et al.*, CERN PPE/94-49 (1994)]. (Courtesy of O. Biebel, S. Bethke, and D. Lanske, RWTH, Aachen, 1994.)


Heavy Quark Fragmentation in e^+e^- Annihilation

Figure 32.12: Heavy quark fragmentation: Shown are the CLEO (D. Bortoletto *et al.*, Phys. Rev. **D37**, 1719 (1988)) and ARGUS (H. Albrecht *et al.*, Z. Phys. **C52**, 353 (1991)) inclusive cross sections ($\sigma \cdot B \cdot d\sigma/dx_p$, with $x_p = p/p_{\max}$) for the production of pseudoscalar D^0 and vector D^{*+} in e^+e^- annihilations at $\sqrt{s} \approx 10$ GeV. These inclusive spectra have not been corrected for cascades from higher states, nor for radiative effects. Many functional forms have been suggested to describe these “hard” spectra, characteristic of charmed particles produced in e^+e^- annihilations. The parameterization given by Peterson *et al.*, (Phys. Rev. **D27**, 105, (1983)) in terms of just one variable ϵ_p ($dN/dx_p = [x_p(1 - 1/x_p - \epsilon_p/(1 - x_p))]^{-2}$) has found the most use. Fits to the combined CLEO and ARGUS D^0 and D^{*+} data give $\epsilon_p(D^0) = 0.135 \pm 0.010$ and $\epsilon_p(D^{*+}) = 0.078 \pm 0.008$; these are indicated by the solid curves.

Spin-dependent effects have been observed in, *e.g.*, the polarization of D^{*+} mesons as a function of x_p . Recent measurements of ϵ_p for D^{*+} and D_{sJ} mesons by CLEO—J. Alexander *et al.*, Phys. Lett. **B303**, 377 (1993) and ARGUS—H. Albrecht *et al.*, Phys. Lett. **B221**, 422 (1989) and Phys. Lett. **B232**, 398, (1989) also indicate that the fragmentation functions of such orbitally excited charmed mesons are distinctly harder than for D or D^* mesons. How much of this is a mass effect and how much is truly a spin effect has not yet been fully determined. (Courtesy of D. Besson, Univ. of Kansas, 1994.)


Average e^+e^- , pp , and $\bar{p}p$ Multiplicity

Figure 32.13: Average multiplicity as a function of \sqrt{s} for e^+e^- and $p\bar{p}$ annihilations and pp collisions. The indicated errors are statistical and systematic errors added in quadrature, except when no systematic errors are given.

e^+e^- : All measurements include contributions from K_S^0 and Λ decays. The $\gamma\gamma 2$ and MARK I measurements contain a systematic 5% error. The five points at the Z resonance have been spread horizontally for clarity [OPAL—P.D. Acton *et al.*, Z. Phys. **C53**, 539 (1992) and references therein].

$p\bar{p}$: The values measured by UA5 exclude single diffractive dissociation [J. Benecke *et al.* (bubble chamber), Nucl. Phys. **B76**, 29 (1976), W.M. Morse *et al.* (bubble chamber), Phys. Rev. **D15**, 66 (1977), ISR—A. Breakstone *et al.*, Phys. Rev. **D30**, 528 (1984), UA5—G.J. Alner *et al.*, Phys. Lett. **167B**, 476 (1986), UA5—Ansgor *et al.*, Z. Phys. **C43**, 357 (1989)]. (Courtesy of O. Biebel, S. Bethke, and D. Lanske, RWTH, Aachen, 1994.)


Annihilation Cross Section Near M_Z 

Figure 32.14: Data from the Mark II, ALEPH, DELPHI, L3, and OPAL Collaborations for the cross section in e^+e^- annihilation into hadronic final states as a function of c.m. energy near the Z . LEP detectors obtained data at the same energies; some of the points are obscured by overlap. The curves show the predictions of the Standard Model with three species (solid curve) and four species (dashed curve) of light neutrinos. The asymmetry of the curves is produced by initial-state radiation. References:

ALEPH: D. Decamp *et al.*, Z. Phys. **C53**, 1 (1992).

DELPHI: P. Abreu *et al.*, Nucl. Phys. **B367**, 511 (1992).

L3: B. Adeva *et al.*, Z. Phys. **C51**, 179 (1991).

OPAL: G. Alexander *et al.*, Z. Phys. **C52**, 175 (1991).




Figure 32.11: Selected measurements of $R \equiv \sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$, where the annihilation in the numerator proceeds via one photon or via the Z . Measurements in the vicinity of the Z mass are shown in the following figure. The denominator is the calculated QED single-photon process; see the section on Cross-Section Formulae for Specific Processes. Radiative corrections and, where important, corrections for two-photon processes and τ production have been made. Note that the ADONE data ($\gamma\gamma 2$ and MEA) is for ≥ 3 hadrons. The points in the $\psi(3770)$ region are from the MARK I—Lead Glass Wall experiment. To preserve clarity only a representative subset of the available measurements is shown—references to additional data are included below. Also for clarity, some points have been combined or shifted slightly ($< 4\%$) in E_{cm} , and some points with low statistical significance have been omitted. Systematic normalization errors are not included; they range from ~ 5 – 20% , depending on experiment. We caution that especially the older experiments tend to have large normalization uncertainties. Note the suppressed zero. The horizontal extent of the plot symbols has no significance. The positions of the $J/\psi(1S)$, $\psi(2S)$, and the four lowest Υ vector-meson resonances are indicated. Two curves are overlaid for $E_{\text{cm}} > 11$ GeV, showing the theoretical prediction for R , including higher order QCD [M. Dine and J. Sapirstein, Phys. Rev. Lett. **43**, 668 (1979)] and electroweak corrections. The Λ values are for 5 flavors in the $\overline{\text{MS}}$ scheme and are $\Lambda_{\overline{\text{MS}}}^{(5)} = 60$ MeV (lower curve) and $\Lambda_{\overline{\text{MS}}}^{(5)} = 250$ MeV (upper curve). (Courtesy of F. Porter, 1992.) References (including several references to data not appearing in the figure and some references to preliminary data):

- AMY:** T. Mori *et al.*, Phys. Lett. **B218**, 499 (1989);
CELLO: H.-J. Behrend *et al.*, Phys. Lett. **144B**, 297 (1984);
 and H.-J. Behrend *et al.*, Phys. Lett. **183B**, 400 (1987);
CLEO: R. Giles *et al.*, Phys. Rev. **D29**, 1285 (1984);
 and D. Besson *et al.*, Phys. Rev. Lett. **54**, 381 (1985);
CUSB: E. Rice *et al.*, Phys. Rev. Lett. **48**, 906 (1982);
CRYSTAL BALL: A. Osterheld *et al.*, SLAC-PUB-4160;
 and Z. Jakubowski *et al.*, Z. Phys. **C40**, 49 (1988);
DASP: R. Brandelik *et al.*, Phys. Lett. **76B**, 361 (1978);
DASP II: Phys. Lett. **116B**, 383 (1982);
DCI: G. Cosme *et al.*, Nucl. Phys. **B152**, 215 (1979);
DHHM: P. Bock *et al.* (DESY-Hamburg-Heidelberg-
 MPI München Collab.), Z. Phys. **C6**, 125 (1980);
 $\gamma\gamma 2$: C. Bacci *et al.*, Phys. Lett. **86B**, 234 (1979);
HRS: D. Bender *et al.*, Phys. Rev. **D31**, 1 (1985);
JADE: W. Bartel *et al.*, Phys. Lett. **129B**, 145 (1983);
 and W. Bartel *et al.*, Phys. Lett. **160B**, 337 (1985);
LENA: B. Niczyporuk *et al.*, Z. Phys. **C15**, 299 (1982);
MAC: E. Fernandez *et al.*, Phys. Rev. **D31**, 1537 (1985);
MARK J: B. Adeva *et al.*, Phys. Rev. Lett. **50**, 799 (1983);
 and B. Adeva *et al.*, Phys. Rev. **D34**, 681 (1986);
MARK I: J.L. Siegrist *et al.*, Phys. Rev. **D26**, 969 (1982);
MARK I + Lead Glass Wall: P.A. Rapidis *et al.*,
 Phys. Rev. Lett. **39**, 526 (1977); and P.A. Rapidis, thesis,
 SLAC-Report-220 (1979);
MARK II: J. Patrick, Ph.D. thesis, LBL-14585 (1982);
MEA: B. Esposito *et al.*, Lett. Nuovo Cimento **19**, 21 (1977);
PLUTO: A. Bäcker, thesis Gesamthochschule Siegen,
 DESY F33-77/03 (1977); C. Gerke, thesis, Hamburg Univ. (1979);
 Ch. Berger *et al.*, Phys. Lett. **81B**, 410 (1979);
 and W. Lackas, thesis, RWTH Aachen, DESY Pluto-81/11 (1981);
TASSO: R. Brandelik *et al.*, Phys. Lett. **113B**, 499 (1982);
 and M. Althoff *et al.*, Phys. Lett. **138B**, 441 (1984);
TOPAZ: I. Adachi *et al.*, Phys. Rev. Lett. **60**, 97 (1988); and
VENUS: H. Yoshida *et al.*, Phys. Lett. **198B**, 570 (1987).

Table 33.2: Regge theory provides a simple and compact description of total cross sections. It is sufficient to write $\sigma_{\text{total}} = X s^\epsilon + Y s^{-\eta}$, where the first term arises from pomeron exchange and the second from ρ , ω , f , a exchange. We list the results of A. Donnachie and P.V. Landshoff, Phys. Lett. **B296**, 227 (1992). Simultaneous fits were first made to pp and $\bar{p}p$ data for $\sqrt{s} > 10$ GeV, requiring the same values of X , ϵ , and η for both reactions. They obtained

$$\epsilon = 0.0808 \quad \eta = 0.4525 .$$


The same values of ϵ and η were used in fits to other reactions. For other pairs of reactions of the form ab and $a\bar{b}$, fits were made for a common value of X and a value of Y , using data with $\sqrt{s} > 6$ GeV. These fits are shown by the gray background curves on the following graphs.

Reaction	X (mb)	Y (mb)
$\bar{p}p$	21.70	98.39
pp	21.70	56.08
$\bar{p}n$	21.70*	92.71
pn	21.70*	54.77
π^-p	13.63	36.02
π^+p	13.63	27.56
K^-p	11.82	26.36
K^+p	11.82	8.15
γp	0.0677	0.129


*Requiring the same pomeron exchange coefficient as for $\sigma(pp)$ and $\sigma(\bar{p}p)$. Fits without the constraint result in an almost identical value, $X = 22.15$ mb.

Table 33.3: Extensively revised 1991 by A. Baldini, V. Flaminio, and O. Yushchenko. The CERN-HERA and COMPAS Groups have made least-squares fits to many high-energy cross sections. The parametrization is $\sigma(p) = A + Bp^n + C \ln^2(p) + D \ln(p)$, where σ is in mb and p is in GeV/c. The best-fit coefficients A , B , C , and D , and the exponent n are tabulated below; where indicated, not all the terms in $\sigma(p)$ are included in the fit. The errors on the parameters are highly correlated since the terms in $\sigma(p)$ are far from orthogonal. Also given is the range of momentum over which the fit was done; extrapolation outside this range is likely to give incorrect results.


Reaction	Momentum range (GeV/c)	Fitted parameters				
		A	B	n	C	D
γp total	3.0–183	0.147 ± 0.001	—	—	0.0022 ± 0.0001	-0.0170 ± 0.0007
γd total	2.0–17.8	0.300 ± 0.005	—	—	0.0095 ± 0.0020	-0.057 ± 0.007
π^+p total	4.0–340	16.4 ± 1.2	19.3 ± 0.8	-0.42 ± 0.05	0.19 ± 0.02	—
π^+p elastic	2.0–200	—	11.4 ± 0.3	-0.4 ± 0.2	0.079 ± 0.005	—
π^-p total	2.5–370	33.0 ± 1.2	14.0 ± 1.8	-1.36 ± 0.29	0.456 ± 0.049	-4.03 ± 0.48
π^-p elastic	2.0–360	1.76 ± 0.42	11.2 ± 0.3	-0.64 ± 0.07	0.043 ± 0.011	—
$\pi^\pm d$ total	2.5–370	56.8 ± 3.6	42.2 ± 8.4	-1.45 ± 0.38	0.65 ± 0.14	-5.39 ± 1.43
K^+p total	2.0–310	18.1 ± 0.1	—	—	0.26 ± 0.03	-1.0 ± 0.1
K^+p elastic	2.0–175	5.0 ± 1.2	8.1 ± 1.5	-1.8 ± 0.7	0.16 ± 0.06	-1.3 ± 0.5
K^+n total	2.0–310	18.7 ± 0.2	—	—	0.21 ± 0.02	-0.89 ± 0.14
K^+d total	2.0–310	34.2 ± 1.2	7.9 ± 3.8	-2.1 ± 1.1	0.346 ± 0.074	-0.99 ± 0.61
K^-p total	3.0–310	32.1 ± 0.2	—	—	0.66 ± 0.01	-5.6 ± 0.1
K^-p elastic	3.0–175	7.3 ± 0.1	—	—	0.29 ± 0.01	-2.40 ± 0.09
K^-n total	1.8–310	25.2 ± 0.5	—	—	0.38 ± 0.03	-2.9 ± 0.3
K^-d total	3.0–310	57.6 ± 0.4	—	—	1.17 ± 0.03	-9.5 ± 0.2
pp total	3.0–2100	48.0 ± 0.1	—	—	0.522 ± 0.005	-4.51 ± 0.05
pp elastic	2.0–2100	11.9 ± 0.8	26.9 ± 1.7	-1.21 ± 0.11	0.169 ± 0.021	-1.85 ± 0.26
pn total	3.0–370	47.30 ± 0.17	—	—	0.513 ± 0.023	-4.27 ± 0.15
pd total	3.0–370	91.3 ± 0.2	—	—	1.05 ± 0.03	-8.8 ± 0.2
pd elastic	2.0–384	16.1 ± 0.7	—	—	0.32 ± 0.04	-3.4 ± 0.4
$\bar{p}p$ total	$5.0-1.73 \times 10^6$	38.4 ± 4.4	77.6 ± 2.8	-0.64 ± 0.07	0.26 ± 0.05	-1.2 ± 0.9
$\bar{p}p$ elastic	$5.0-1.73 \times 10^6$	10.2 ± 0.7	52.7 ± 1.8	-1.16 ± 0.05	0.125 ± 0.014	-1.28 ± 0.20
$\bar{p}n$ total	1.1–280	—	133.6 ± 4.6	-0.70 ± 0.03	-1.22 ± 0.13	13.7 ± 0.7
$\bar{p}n$ elastic	1.1–5.55	36.5 ± 1.5	—	—	—	-11.9 ± 1.8
$\bar{p}d$ total	2.0–280	112 ± 13	125 ± 8	-1.08 ± 0.15	1.14 ± 0.49	-12.4 ± 4.9




Hadronic total and elastic cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; and COMPAS Group, IHEP, Serpukhov, Russia. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988). Gray curve shows Regge fit from Table 33.2 .




Hadronic total and elastic cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; and COMPAS Group, IHEP, Serpukhov, Russia. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988). Gray curve shows Regge fit from Table 33.2 .



Hadronic total and elastic cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; and COMPAS Group, IHEP, Serpukhov, Russia. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988). Gray curve shows Regge fit from Table 33.2 .



Hadronic total and elastic cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; and COMPAS Group, IHEP, Serpukhov, Russia. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988). Gray curve shows Regge fit from Table 33.2 .



Hadronic total and elastic cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; and COMPAS Group, IHEP, Serpukhov, Russia. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988). Gray curve shows Regge fit from Table 33.2 .




Figure 32.20: Λp total and elastic cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; and COMPAS Group, IHEP, Serpukhov, USSR. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988).




Figure 33.7: Photon cross sections vs. laboratory beam momentum and total center-of-mass energy. Data courtesy A. Baldini, V. Flaminio, W.G. Moorhead, and D.R.O. Morrison, CERN; COMPAS Group, IHEP, Serpukhov, USSR; and G.M. Lewis, Glasgow. See *Total Cross Sections for Reactions of High Energy Particles*, Landolt-Börnstein, New Series Vol. I/12 a and I/12 b, ed. H. Schopper (1988). Gray curve shows Regge fit from Table 33.2.

INTRODUCTION TO THE FULL LISTINGS

Illustrative key	1343
Abbreviations	1344

Illustrative Key to the Full Listings

Name of particle. "Old" name used before 1986 renaming scheme also given if different. See the section "Naming Scheme for Hadrons" for details.

$a_0(1200)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

Particle quantum numbers (where known).

OMITTED FROM SUMMARY TABLE

Evidence not compelling, may be a kinematic effect.

Indicates particle omitted from Particle Properties Summary Table, implying particle's existence is not confirmed.

Quantity tabulated below.

$a_0(1200)$ MASS

Top line gives our best value (and error) of quantity tabulated here, based on weighted average of measurements used. Could also be from fit, best limit, estimate, or other evaluation. See next page for details.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1206 ± 7 OUR AVERAGE					
$1210 \pm 8 \pm 9$	3000	FENNER 87	MMS	-	$3.5 \pi^- p$
1198 ± 10		PIERCE 83	ASPK	+	$2.1 K^- p$
$1216 \pm 11 \pm 9$	1500	MERRILL 81	HBC	0	$3.2 K^- p$
1192 ± 16	200	LYNCH 81	HBC	\pm	$2.7 \pi^- p$

General comments on particle.

"Document id" for this result; full reference given below.

Measurement technique. (See abbreviations on next page.)

Footnote number linking measurement to text of footnote.

¹ Systematic error was added quadratically by us in our 1986 edition.

$a_0(1200)$ WIDTH

Number of events above background.


Measured value used in averages, fits, limits, etc.

Error in measured value (often statistical only; followed by systematic if separately known; the two are combined in quadrature for averaging and fitting.)

Measured value *not used* in averages, fits, limits, etc. See the Introductory Text for explanations.

Top "data point" indicates average. Width of error bar (and shaded pattern below) is \pm error on average, scaled by "scale factor" S.

Value and error for each experiment.



Scale factor > 1 indicates possibly inconsistent data.

Reaction producing particle, or general comments.

"Change bar" indicates result added or changed since previous edition.

Charge(s) of particle(s) detected.

Ideogram to display possibly inconsistent data. Curve is sum of Gaussians, one for each experiment (area of Gaussian = 1/error; width of Gaussian = \pm error). See Introductory Text for discussion.

Contribution of experiment to χ^2 (if no entry present, experiment not used in calculating χ^2 or scale factor because of very large error).

$a_0(1200)$ DECAY MODES

Partial decay mode (labeled by Γ_i).

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 3π	$(65.2 \pm 1.3) \%$	S=1.7
Γ_2 KK	$(34.8 \pm 1.3) \%$	S=1.7
Γ_3 $\eta \pi^\pm$	$< 4.9 \times 10^{-4}$	CL=95%

Our best value for branching fraction as determined from data averaging, fitting, evaluating, limit selection, etc. This list is basically a compact summary of results in the Branching Ratio section below.

$a_0(1200)$ BRANCHING RATIOS

Branching ratio.

Our best value (and error) of quantity tabulated, as determined from constrained fit (using all significant measured branching ratios for this particle).

Weighted average of measurements of this ratio only.

Footnote (referring to LYNCH 81).

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_i/Γ
0.652 ± 0.013 OUR FIT				Error Includes scale factor of 1.7.	
0.643 ± 0.010 OUR AVERAGE					
0.64 ± 0.01	PIERCE 83	ASPK	+	$2.1 K^- p$	
0.74 ± 0.06	MERRILL 81	HBC	0	$3.2 K^- p$	
0.48 ± 0.15	² LYNCH 81	HBC	\pm	$2.7 \pi^- p$	
² Data has questionable background subtraction.					
$\Gamma(KK)/\Gamma_{total}$					Γ_2/Γ
0.348 ± 0.013 OUR FIT				Error Includes scale factor of 1.7.	
0.35 ± 0.05	PIERCE 83	ASPK	+	$2.1 K^- p$	
$\Gamma(KK)/\Gamma(3\pi)$					Γ_2/Γ_1
0.538 ± 0.030 OUR FIT				Error Includes scale factor of 1.7.	
0.50 ± 0.03	MERRILL 81	HBC	0	$3.2 K^- p$	
$\Gamma(\eta(\text{neutral decay})\pi^\pm)/\Gamma_{total}$					$0.71\Gamma_3/\Gamma$
< 3.5					
< 3.5	PIERCE 83	ASPK	+	$2.1 K^- p$	

Branching ratio in terms of partial decay mode(s) Γ_i above.

Confidence level for measured upper limit.

References, ordered inversely by year, then author.

"Document id" used on data entries above.

Journal, report, preprint, etc. (See abbreviations on next page.)

$a_0(1200)$ REFERENCES

FENNER 87	PRL 55 14	+Watson, Willis, Zorn	(SLAC)
PIERCE 83	PL 123B 230	+Jones+	(FNAL) [JP]
LYNCH 81	PR D24 610	+Armstrong, Harper, Rittenberg, Wagman	(CLEO Collab.)
MERRILL 81	PRL 47 143		(SACL, CERN)

Partial list of author(s) in addition to first author.

Quantum number determinations in this reference.

Institution(s) of author(s). (See abbreviations on next page.)

Abbreviations Used in the Full Listings

Indicator of Procedure Used to Obtain Our Result

OUR AVERAGE	From a weighted average of selected data.
OUR FIT	From a constrained or overdetermined multiparameter fit of selected data.
OUR EVALUATION	Not from a direct measurement, but evaluated from measurements of other quantities.
OUR ESTIMATE	Based on the observed range of the data. Not from a formal statistical procedure.
OUR LIMIT	For special cases where the limit is evaluated by us from measured ratios or other data. Not from a direct measurement.

Measurement Techniques

(i.e., Detectors and Methods of Analysis)

ACCM	ACCMOR Collaboration	FRAM	ADONE MEA group detector
AEMS	Argonne effective mass spectrometer	FREJ	FREJUS Collaboration - modular flash chamber detector (calorimeter)
ALEP	ALEPH - CERN LEP detector	GALX	GALLEX solar neutrino detector in the Gran Sasso Underground Lab.
AMY	AMY detector at KEK-TRISTAN	GAM2	IHEP hodoscope Cherenkov γ calorimeter GAMS-2000
ARG	ARGUS detector at DORIS	GAM4	CERN hodoscope Cherenkov γ calorimeter GAMS-4000
ARGD	Fit to semicircular amplitude path on Argand diagram	GOLI	CERN Goliath spectrometer
ASP	Anomalous single-photon detector	H1	H1 detector at DESY/HERA
ASPK	Automatic spark chambers	HBC	Hydrogen bubble chamber
ASTE	ASTERIX detector at LEAR	HDHC	Hydrogen and deuterium bubble chambers
ASTR	Astronomy	HEBC	Helium bubble chamber
B845	BNL experiment 845 detector	HEPT	Helium proportional tubes
BAKS	Baksan underground scintillation telescope	HLBC	Heavy-liquid bubble chamber
BC	Bubble chamber	HOME	Homestake underground scintillation detector
BDMP	Beam dump	HPW	Harvard-Pennsylvania-Wisconsin detector
BEBC	Big European bubble chamber at CERN	HRS	SLAC high-resolution spectrometer
BES	BES Beijing Spectrometer at Beijing Electron-Positron Collider	HYBR	Hybrid: bubble chamber + electronics
BIS2	BIS-2 spectrometer at Serpukhov	IMB	Irvine-Michigan-Brookhaven underground Cherenkov detector
BKEI	BENKEI spectrometer system at KEK Proton Synchrotron	IMB3	Irvine-Michigan-Brookhaven underground Cherenkov detector
BONA	Bonanza nonmagnetic detector at DORIS	INDU	Magnetic induction
BPWA	Barrelet-zero partial-wave analysis	IPWA	Energy-independent partial-wave analysis
CALO	Calorimeter	JADE	JADE detector at DESY
CBAL	Crystal Ball detector at SLAC-SPEAR or DORIS	KAM2	KAMIOKANDE-II underground Cherenkov detector
CBAR	Crystal Barrel detector at CERN-LEAR	KAMI	KAMIOKANDE underground Cherenkov detector
CBOX	Crystal Box at LAMPF	KOLR	Kolar Gold Field underground detector
CC	Cloud chamber	L3	L3 detector at LEP
CCFR	Columbia-Chicago-Fermilab-Rochester detector	LASS	Large-angle superconducting solenoid spectrometer at SLAC
CDF	Collider detector at Fermilab	LEBC	Little European bubble chamber at CERN
CDHS	CDHS neutrino detector at CERN	LENA	Nonmagnetic lead-glass NaI detector at DORIS
CELL	CELLO detector at DESY	MAC	MAC detector at PEP/SLAC
CHER	Cherenkov detector	MBR	Molecular beam resonance technique
CHM2	CHARM-II neutrino detector (glass) at CERN	MCRO	MACRO detector in Gran Sasso
CHRM	CHARM neutrino detector (marble) at CERN	MD1	Magnetic detector at VEPP-4, Novosibirsk
CIBS	CERN-IHEP boson spectrometer	MDRP	Millikan drop measurement
CLE2	CLEO II detector at CESR	MICA	Underground mica deposits
CLEO	Cornell magnetic detector at CESR	MLEV	Magnetic levitation
CMD	Cryogenic magnetic detector at VEPP-2M, Novosibirsk	MMS	Missing mass spectrometer
CNTR	Counters	MPS	Multiparticle spectrometer at BNL
COSM	Cosmology and astrophysics	MPS2	Multiparticle spectrometer upgrade at BNL
CSB2	Columbia U. - Stony Brook BGO calorimeter inserted in NaI array	MPSF	Multiparticle spectrometer at Fermilab
CUSB	Columbia U. - Stony Brook segmented NaI detector at CESR	MPWA	Model-dependent partial-wave analysis
D0	D0 detector at Fermilab Tevatron Collider	MRK1	SLAC Mark-I detector
DASP	DESY double-arm spectrometer	MRK2	SLAC Mark-II detector
DBC	Deuterium bubble chamber	MRK3	SLAC Mark-III detector
DLCO	DELCO detector at SLAC-SPEAR or SLAC-PEP	MRKJ	Mark-J detector at DESY
DLPH	DELPHI detector at LEP	MRS	Magnetic resonance spectrometer
DM1	Magnetic detector no. 1 at Orsay DCI collider	NA14	CERN
DM2	Magnetic detector no. 2 at Orsay DCI collider	NA31	CERN NA31 Spectrometer-Calorimeter
DPWA	Energy-dependent partial-wave analysis	NA32	CERN NA32 Spectrometer
E653	Fermilab E653 detector	ND	NaI detector at VEPP-2M, Novosibirsk
E687	Fermilab E687 detector	NICE	Serpukhov nonmagnetic precision spectrometer
E691	Fermilab E691 detector	NMR	Nuclear magnetic resonance
E731	Fermilab E731 Spectrometer-Calorimeter	NUSX	Mont Blanc NUSEX underground detector
E761	Fermilab E761 detector	OBLX	OBELIX detector at LEAR
E799	Fermilab E799 Spectrometer-Calorimeter	OLYA	Detector at VEPP-2M and VEPP-4, Novosibirsk
EHS	Four-pi detector at CERN	OMEG	CERN OMEGA spectrometer
ELEC	Electronic combination	OPAL	OPAL detector at LEP
EMC	European muon collaboration detector at CERN	OSPK	Optical spark chamber
EMUL	Emulsions	PLAS	Plastic detector
FBC	Freon bubble chamber	PLUT	DESY PLUTO detector
FIT	Fit to previously existing data	PWA	Partial-wave analysis
FMPS	Fermilab Multiparticle Spectrometer	REDE	Resonance depolarization
FRAB	ADONE $B\bar{B}$ group detector	RVUE	Review of previous data
FRAG	ADONE $\gamma\gamma$ group detector	SAGE	US - Russian Gallium Experiment
		SFM	CERN split-field magnet
		SHF	SLAC Hybrid Facility Photon Collaboration
		SIGM	Serpukhov CERN-IHEP magnetic spectrometer (SIGMA)
		SILI	Silicon detector
		SLD	SLC Large Detector for e^+e^- colliding beams at SLAC
		SOUJ	Soudan underground detector
		SPEC	Spectrometer
		SPED	From maximum of speed plot or resonant amplitude
		SPRK	Spark chamber
		SQID	SQUID device
		STRC	Streamer chamber
		TASS	DESY TASSO detector

Abbreviations Used in the Full Listings (*Cont'd*)

THEO	Theoretical or heavily model-dependent result
THY	Theory
TOF	Time-of-flight
TOPZ	TOPAZ detector at KEK-TRISTAN
TPC	TPC detector at PEP/SLAC
TPS	Tagged photon spectrometer at Fermilab
TRAP	Penning trap
UA1	UA1 detector at CERN
UA2	UA2 detector at CERN
UA5	UA5 detector at CERN
VES	Vertex Spectrometer Facility at 70 GeV IHEP accelerator
VNS	VENUS detector at KEK-TRISTAN
WA75	CERN WA75 experiment
WA82	CERN WA82 experiment
WIRE	Wire chamber
XEBC	Xenon bubble chamber
ZEUS	ZEUS detector at DESY/HERA

Conferences

Conferences are generally referred to by the location at which they were held (e.g., HAMBURG, TORONTO, CORNELL, BRIGHTON, etc.).

Journals

AA	Astronomy and Astrophysics
ADVP	Advances in Physics
AFIS	Anales de Fisica
AJP	American Journal of Physics
ANP	Annals of Physics
ANPL	Annals of Physics (Leipzig)
ANYAS	Annals of the New York Academy of Sciences
AP	Atomic Physics
APAH	Acta Physica Academiae Scientiarum Hungaricae
APJ	Astrophysical Journal
APP	Acta Physica Polonica
ARNPS	Annual Review of Nuclear and Particle Science
ARNS	Annual Review of Nuclear Science
BAPS	Bulletin of the American Physical Society
BASUP	Bulletin of the Academy of Science, USSR (Physics)
CJP	Canadian Journal of Physics
CNPP	Comments on Nuclear and Particle Physics
CZJP	Czechoslovak Journal of Physics
DANS	Doklady Akademii nauk SSSR
EPL	Europhysics Letters
FECAY	Fizika Elementarnykh Chastits i Atomnogo Yadra
HADJ	Hadronic Journal
IJMP	International Journal of Modern Physics
JAP	Journal of Applied Physics
JETP	English Translation of Soviet Physics ZETF
JETPL	English Translation of Soviet Physics ZETF Letters
JINR	Joint Inst. for Nuclear Research
JPA	Journal of Physics, A
JPB	Journal of Physics, B
JPCRd	Journal of Physical and Chemical Reference Data
JPG	Journal of Physics, G
JPSJ	Journal of the Physical Society of Japan
LNC	Lettere Nuovo Cimento
MNRA	Monthly Notices of the Royal Astronomical Society
MPL	Modern Physics Letters
NAT	Nature
NC	Nuovo Cimento
NIM	Nuclear Instruments and Methods
NP	Nuclear Physics
NPBPS	Nuclear Physics B Proceedings Supplement
PDAT	Physik Daten
PL	Physics Letters
PN	Particles and Nuclei
PPN	Physics of Particles and Nuclei (formerly SJPN)
PPNP	Progress in Particles and Nuclear Physics
PPSL	Proc. of the Physical Society of London
PR	Physical Review
PRAM	Pramana
PRL	Physical Review Letters
PRPL	Physics Reports (Physics Letters C)
PRSE	Proc. of the Royal Society of Edinburgh
PRSL	Proc. of the Royal Society of London, Section A
PS	Physica Scripta

PTP	Progress of Theoretical Physics
PTRSL	Phil. Trans. Royal Society of London
RA	Radiochimica Acta
RMP	Reviews of Modern Physics
RNC	La Rivista del Nuovo Cimento
RPP	Reports on Progress in Physics
RRP	Revue Roumaine de Physique
SCI	Science
SJNP	Soviet Journal of Nuclear Physics
SJPN	Soviet Journal of Particles and Nuclei
SPD	Soviet Physics Doklady (Magazine)
SPU	Soviet Physics - Uspekhi
YAF	Yadernaya Fizika
ZETF	Zhurnal Eksperimental'noi i Teoreticheskoi Fiziki
ZETFP	Zhurnal Eksperimental'noi i Teoreticheskoi Fiziki, Pis'ma v Redakts
ZNAT	Zeitschrift fur Naturforschung
ZPHY	Zeitschrift fur Physik

Institutions

AACH	Phys. Inst. der Techn. Hochschule Aachen (Historical, use for general Inst. der Techn. Hochschule)	Aachen, Germany
AACH1	I Phys. Inst. der Techn. Hochschule Aachen	Aachen, Germany
AACH3	III Phys. Inst. der Techn. Hochschule Aachen	Aachen, Germany
AACHT	Institut für Theoretische Physik	Aachen , Germany
AARH	Aarhus Univ.	Aarhus C, Denmark
ABO	Åbo Akademi University	Åbo (Turku), Finland
ADEL	Adelphi Univ.	Garden City, NY, USA
ADLD	The Univ. of Adelaide	South Australia, Australia
AERE	Atomic Energy Research Estab.	Didcot, United Kingdom
AFRR	Armed Forces Radiobiology Res. Inst.	Bethesda, MD, USA
AHMED	Physical Research Lab.	Ahmedabad , Gujarat, India
AICH	Aichi Univ. of Education	Aichi, Japan
AKIT	Akita Univ.	Akita, Japan
ALAH	Univ. of Alabama (Huntsville)	Huntsville, AL, USA
ALAT	Univ. of Alabama (Tuscaloosa)	Tuscaloosa, AL, USA
ALBA	SUNY at Albany	Albany, NY, USA
ALBE	Univ. of Alberta	Edmonton, AB, Canada
AMES	Ames Lab.	Ames, IA, USA
AMHT	Amherst College	Amherst, MA, USA
AMST	Univ. van Amsterdam	Amsterdam, The Netherlands
ANIK	NIKHEF	Amsterdam , The Netherlands
ANKA	Middle East Technical Univ.	Ankara, Turkey
ANL	Argonne National Lab.	Argonne, IL, USA
ANSM	St. Anselm Coll.	Manchester, NH, USA
ARCBO	Arecibo Observatory	Arecibo, PR, USA
ARIZ	Univ. of Arizona	Tucson, AZ, USA
ARZS	Arizona State Univ.	Tempe, AZ, USA
ASCI	Academy of Sciences	Moscow , Russian Federation
AST	Inst. of Phys.	Nankang, Taipei, The Republic of China (Taiwan)
ATEN	NCSR "Demokritos"	Aghia Paraskevi Attikis, Greece
ATHU	Univ. of Athens	Athens, Greece
AUCK	Univ. of Auckland	Auckland, New Zealand
BAKU	Inst. of Physics	Baku , Azerbaijan
BANGB	Bangabasi College	Calcutta, India
BARC	Univ. Autónoma de Barcelona	Bellaterra (Barcelona), Spain
BARI	Univ. di Bari	Bari, Italy
BART	Univ. of Delaware ; Bartol Research Inst.	Newark, DE, USA
BASL	Inst. für Physik der Univ. Basel	Basel, Switzerland
BAYR	Univ. Bayreuth	Bayreuth, Germany
BCEN	Centre d'Etudes Nucleaires de Bordeaux-Gradignan	Gradignan, France
BELG	Inter-University Inst. for High Energies (ULB-VUB)	Bruxelles , Belgium

Abbreviations Used in the Full Listings (*Cont'd*)

BELL	AT & T Bell Labs	Murray Hill, NJ, USA	CIT	California Inst. of Tech.	Pasadena, CA, USA
BERG	Univ. of Bergen	Bergen, Norway	CLER	Univ. de Clermont-Ferrand	Aubière, France
BERL	DESY - Inst. für Hochenergiephysik Zeuthen	Zeuthen , Germany	CLEV	Cleveland State Univ.	Cleveland, OH, USA
BERN	Univ. of Berne	Berne, Switzerland	CMNS	Comenius Univ.	Bratislava , Slovak Republic
BGNA	Univ. di Bologna	Bologna, Italy	CMU	Carnegie Mellon Univ.	Pittsburgh, PA, USA
BGUN	Ben-Gurion Univ.	Beer-Sheva, Israel	CNEA	Comisión Nacional de Energía Atómica	Buenos Aires, Argentina
BHAB	Bhabha Atomic Research Center	Trombay, Bombay, India	CNRC	Centre for Research in Particle Physics	Ottawa, ON, Canada
BHEP	Inst. of High Energy Physics	Beijing , The People's Republic of China	CNRS	Centre National de la Recherche Scientifique, Luminy	Marseille , France
BIEL	Univ. Bielefeld	Bielefeld, Germany	COLO	Univ. of Colorado	Boulder, CO, USA
BING	SUNY at Binghamton	Binghamton, NY, USA	COLU	Columbia Univ.	New York, NY, USA
BIRM	Univ. of Birmingham	Edgbaston, Birmingham, United Kingdom	CONC	Concordia University	Montreal, PQ, Canada
BLSU	Bloomsburg Univ.	Bloomsburg, PA, USA	CORN	Cornell Univ.	Ithaca, NY, USA
BNL	Brookhaven National Lab.	Upton, NY, USA	COSU	Colorado State Univ.	Fort Collins, CO, USA
BOCH	Ruhr Univ. Bochum	Bochum, Germany	CRAC	Kraków Inst. of Nuclear Physics	Kraków, Poland
BOHR	Niels Bohr Inst.	Copenhagen Ø, Denmark	CRNL	Chalk River Labs.	Chalk River, ON, Canada
BOIS	Boise State Univ.	Boise, ID, USA	CSOK	Oklahoma Central State Univ.	Edmond, OK, USA
BOMB	Univ. of Bombay	Bombay, India	CSULB	California State Univ.	Long Beach, CA, USA
BONN	Univ. Bonn	Bonn, Germany	CUNY	City College of New York	New York, NY, USA
BORD	Univ. de Bordeaux I	Gradignan, France	CURIN	Univ. Pierre et Marie Curie (Paris VI), LPNHE	Paris, France
BOSE	S.N. Bose National Centre for Basis Sciences	Calcutta, India	CURIT	Univ. Pierre et Marie Curie (Paris VI), LPTHE	Paris, France
BOSK	"Rudjer Bošković" Inst.	Zagreb, Croatia	DALH	Dalhousie Univ.	Halifax, NS, Canada
BOST	Boston Univ.	Boston, MA, USA	DARE	Daresbury Lab	Cheshire, United Kingdom
BRAN	Brandeis Univ.	Waltham, MA, USA	DARM	Tech. Hochschule Darmstadt	Darmstadt, Germany
BRCO	Univ. of British Columbia	Vancouver, BC, Canada	DELA	Univ. of Delaware ; Dept. of Physics & Astronomy	Newark, DE, USA
BRIS	Univ. of Bristol	Bristol, United Kingdom	DELH	Univ. of Delhi	Delhi, India
BROW	Brown Univ.	Providence, RI, USA	DESY	DESY , Deutsches Elektronen-Synchrotron	Hamburg, Germany
BRUX	Univ. Libre de Bruxelles ; Service de Physique des Particules Élémentaires	Bruxelles, Belgium	DOE	Department of Energy	Washington, DC, USA
BRUXT	Univ. Libre de Bruxelles ; Physique Théorique	Bruxelles, Belgium	DORT	Univ. Dortmund	Dortmund, Germany
BUCH	Univ. of Bucharest	Bucharest-Magurele, Romania	DUKE	Duke Univ.	Durham, NC, USA
BUDA	KFKI Research Inst. for Particle & Nuclear Physics	Budapest , Hungary	DURH	Univ. of Durham	Durham City, United Kingdom
BUFF	SUNY at Buffalo	Buffalo, NY, USA	DUUC	University College	Dublin, Ireland
BURE	Inst. des Hautes Etudes Scientifiques	Bures-sur-Yvette , France	EDIN	Univ. of Edinburgh	Edinburgh, United Kingdom
CAEN	Lab. de Physique Corpusculaire, ISMRA	CAEN , France	EFI	Enrico Fermi Inst.	Chicago , IL, USA
CAGL	Univ. di Cagliari	Cagliari, Italy	ELMT	Elmhurst College	Elmhurst, IL, USA
CAIR	Cairo University	Orman, Giza, Cairo, Egypt	ENSP	L'Ecole Normale Supérieure	Paris , France
CAIW	Carnegie Inst. of Washington	Washington, DC, USA	EOTV	Eötvös University	Budapest, Hungary
CALC	Univ. of Calcutta	Calcutta, India	EPOL	École Polytechnique	Palaiseau , France
CAMB	Univ. of Cambridge	Cambridge, United Kingdom	ERLA	Univ. Erlangen-Nurnberg	Erlangen, Germany
CAMP	Univ. de Campinas	Campinas , SP, Brasil	ETH	Inst. for High Energy Physics	Zürich , Switzerland
CANB	Australian National Univ.	Canberra, ACT, Australia	FERR	Univ. di Ferrara	Ferrara, Italy
CAPE	University of Capetown	Rondebosch, Cape, South Africa	FIRZ	Univ. di Firenze	Firenze, Italy
CARA	Univ. Central de Venezuela	Caracas, Venezuela	FISK	Fisk Univ.	Nashville, TN, USA
CARL	Carleton Univ.	Ottawa, ON, Canada	FLOR	Univ. of Florida	Gainesville, FL, USA
CARLC	Carleton College	Northfield, MN, USA	FNAL	Fermilab	Batavia, IL, USA
CASE	Case Western Reserve Univ.	Cleveland, OH, USA	FOM	FOM , Stichting voor Fundamenteel Onderzoek der Materie	JP Utrecht , The Netherlands
CAST	China Center of Advanced Science and Technology	Beijing, The People's Republic of China	FRAN	Univ. Frankfurt	Frankfurt am Main, Germany
CATA	Univ. di Catania	Catania, Italy	FRAS	Lab. Nazionali de Frascati dell'INFN	Frascati (Roma), Italy
CATH	Catholic Univ. of America	Washington, DC, USA	FREIB	Albert-Ludwigs Univ.	Freiburg , Germany
CAVE	Univ. of Cambridge	Cambridge, United Kingdom	FREIE	Freie Univ. Berlin	Berlin, Germany
CBNM	CBNM	Geel , Belgium	FRIB	Univ. de Fribourg	Fribourg, Switzerland
CCAC	Allegheny College	Meadville, PA, USA	FSU	Florida State University	Tallahassee, FL, USA
CDEF	College de France	Paris, France	FSUSC	Florida State Univ.	Tallahassee, FL, USA
CEA	Cambridge Electron Accelerator (Historical in <i>Review</i>)	Cambridge, MA , USA	FUKI	Fukui Univ.	Fukui, Japan
CENG	Centre d'Etudes Nucleaires	Grenoble , France	FUKU	Fukushima Univ.	Fukushima, Japan
CERN	CERN , European Laboratory for Particle Physics	Genève, Switzerland	GENO	Univ. di Genova	Genova, Italy
CFPA	Univ. of California, Berkeley	Berkeley, CA, USA	GEOR	Georgian Academy of Sciences	Tbilisi, Republic of Georgia
CHIC	Univ. of Chicago	Chicago, IL, USA	GESC	General Electric Co.	Schenectady, NY, USA
CINC	Univ. of Cincinnati	Cincinnati, OH, USA	GEVA	Univ. de Genève	Genève, Switzerland
CINV	CINVESTAV-IPN, Centro de Investigacion y de Estudios Avanzados del IPN	México , DF, Mexico	GIFU	Gifu Univ.	Gifu, Japan
			GLAS	Univ. of Glasgow	Glasgow, United Kingdom
			GMAS	George Mason Univ.	Fairfax, VA, USA
			GOET	Univ. Göttingen	Göttingen, Germany

Abbreviations Used in the Full Listings (*Cont'd*)

GRAN	Univ. de Granada	Granada, Spain	IUPU	Indiana Univ., Purdue Univ. Indianapolis	Indianapolis, IN, USA
GRAZ	Univ. Graz	Graz, Austria	JADA	Jadavpur Univ.	Calcutta, India
GRON	Univ. of Groningen	Groningen, The Netherlands	JAGL	Jagiellonian Univ.	Kraków, Poland
GSCO	Geological Survey of Canada	Ottawa, ON, Canada	JHU	Johns Hopkins Univ.	Baltimore, MD, USA
GS1	Darmstadt Gesellschaft fur Schwerionenforschung	Darmstadt, Germany	JINR	JINR, Joint Inst. for Nucl. Research	Dubna, Russian Federation
GUEL	Univ. of Guelph	Guelph, ON, Canada	JULI	Julich, Forschungszentrum	Julich, Germany
GWU	George Washington Univ.	Washington, DC, USA	JYV	Univ. of Jyväskylä	Jyväskylä, Finland
HAIF	Technion - Israel Inst. of Tech.	Technion, Haifa, Israel	KAGO	Univ. of Kagoshima	Kagoshima-shi, Japan
HAMB	Univ. Hamburg	Hamburg, Germany	KANS	Univ. of Kansas	Lawrence, KS, USA
HANN	Univ. Hannover	Hannover, Germany	KARL	Univ. Karlsruhe ; (unspecified division) (<i>Historical in Review</i>)	Karlsruhe, Germany
HARC	Houston Advanced Research Ctr.	The Woodlands, TX, USA	KARLE	Univ. Karlsruhe ; Inst. für Experimentelle Kernphysik	Karlsruhe, Germany
HARV	Harvard Univ.	Cambridge, MA, USA	KARLK	Univ. Karlsruhe ; Inst. für Kernphysik	Karlsruhe, Germany
HAWA	Univ. of Hawai'i	Honolulu, HI, USA	KARLT	Univ. Karlsruhe ; Inst. für Theoretische Teilchenphysik	Karlsruhe, Germany
HEBR	Hebrew Univ.	Jerusalem, Israel	KAZA	Kazakh Inst. of High Energy Physics	Alma Ata, Kazakhstan
HEID	Univ. Heidelberg ; (unspecified division) (<i>Historical in Review</i>)	Heidelberg, Germany	KEK	KEK, National Lab. for High Energy Phys.	Ibaraki-ken, Japan
HEIDH	Univ. Heidelberg ; Inst. für Hochenergiephysik	Heidelberg, Germany	KENT	Univ. of Kent	Canterbury, United Kingdom
HEIDP	Univ. Heidelberg ; Physik Inst.	Heidelberg, Germany	KEYN	Open Univ.	Milton Keynes, United Kingdom
HEIDT	Univ. Heidelberg ; Inst. für Theoretische Physik	Heidelberg, Germany	KFTI	Kharkiv Inst. of Physics and Tech. (KFTI)	Kharkiv, Ukraine
HEL5	Univ. of Helsinki	Helsinki, Finland	KIAE	Kurchatov, Inst. of Atomic Energy	Moscow, Russian Federation
HIRO	Hiroshima Univ.	Higashi-Hiroshima, Japan	KIAM	Keldysh Inst. of Applied Math., Acad. Sci., Russia	Moscow, Russian Federation
HOUS	Univ. of Houston	Houston, TX, USA	KIDR	Inst. of Nuclear Sciences, Vinča (Formerly Boris Kidrič Inst.)	Beograd, Serbia, Yugoslavia
HPC	Hewlett-Packard Corp.	Cupertino, CA, USA	KINK	Kinki Univ.	Osaka, Japan
HSCA	Harvard-Smithsonian Center for Astrophysics	Cambridge, MA, USA	KNTY	Univ. of Kentucky	Lexington, KY, USA
IAS	Inst. for Advanced Study	Princeton, NJ, USA	KOBE	Kobe Univ.	Kobe, Japan
IASD	Dublin Inst. for Advanced Studies	Dublin, Ireland	KOMAB	Univ. of Tokyo, Komaba	Tokyo, Japan
IBAR	Ibaraki Univ.	Ibaraki, Japan	KONAN	Konan Univ.	Kobe, Japan
IBM	IBM Corp.	Palo Alto, CA, USA	KOSI	Inst. of Experimental Physics	Košice, Slovak Republic
IBMY	IBM	Yorktown Heights, NY, USA	KYOT	Kyoto Univ.	Kyoto, Japan
IBS	Inst. for Boson Studies	Pasadena, CA, USA	KYOTY	Kyoto Univ.; Yukawa Inst. for Theor. Physics	Kyoto, Japan
ICEPP	Univ. of Tokyo ; Int. Center for Elementary Particle Physics (ICEPP)	Tokyo, Japan	LALO	LAL, Laboratoire de l'Accélérateur Linéaire	Orsay, France
ICRR	Univ. of Tokyo ; Inst. for Cosmic Ray Research	Tokyo, Japan	LANC	Univ. of Lancaster	Lancaster, United Kingdom
ICTP	Int'l Centre for Theoretical Physics	Trieste, Italy	LANL	Los Alamos National Lab. (LANL)	Los Alamos, NM, USA
IFIC	Univ. de Valencia - CSIC	Burjassot, Valencia, Spain	LAPP	LAPP, Lab. d'Annecy-le-Vieux de Phys. des Particules	Annecy-le-Vieux, France
IFRJ	Univ. Federal do Rio de Janeiro	Rio de Janeiro, RJ, Brasil	LASL	U.C. Los Alamos Scientific Lab. (Old name for LANL)	Los Alamos, NM, USA
IIT	Illinois Inst. of Tech.	Chicago, IL, USA	LAUS	Univ. de Lausanne	Lausanne, Switzerland
ILL	Univ. of Illinois at Urbana-Champaign	Urbana, IL, USA	LAVL	Univ. Laval	Quebec, PQ, Canada
ILLC	Univ. of Illinois at Chicago	Chicago, IL, USA	LBL	Lawrence Berkeley Lab.	Berkeley, CA, USA
ILLG	Inst. Laue-Langevin	Grenoble, France	LCGT	Univ. di Torino	Turin, Italy
IND	Indiana Univ.	Bloomington, IN, USA	LEBD	Lebedev Physical Inst.	Moscow, Russian Federation
INEL	E G and G Idaho, Inc.	Idaho Falls, ID, USA	LECE	Univ. di Lecce	Lecce, Italy
INFN	Ist. Nazionale di Fisica Nucleare (Generic INFN, unknown location)	Various places, Italy	LEED	Univ. of Leeds	Leeds, United Kingdom
INNS	Univ. Innsbruck	Innsbruck, Austria	LEHI	Lehigh Univ.	Bethlehem, PA, USA
INRM	INR, Inst. for Nucl. Research	Moscow, Russian Federation	LEHM	Lehman College of CUNY	Bronx, NY, USA
INUS	Univ. of Tokyo ; Inst. for Nuclear Study	Tokyo, Japan	LEID	Univ. of Leiden	Leiden, The Netherlands
IOAN	Univ. of Ioannina	Ioannina, Greece	LEMO	Le Moyne Coll.	Syracuse, NY, USA
IOFF	A.F. Ioffe Phys. Tech. Inst.	St. Petersburg, Russian Federation	LEUV	Katholieke Univ. Leuven	Leuven, Belgium
IOWA	Univ. of Iowa	Iowa City, IA, USA	LINZ	Univ. Linz	Linz, Austria
IPN	IPN, Inst. de Phys. Nucl.	Orsay, France	LISB	Inst. Nacional de Investigacion Cientifica	Lisboa CODEX, Portugal
IPNP	Univ. Pierre et Marie Curie (Paris VI)	Paris, France	LISBT	Univ. Técnica de Lisboa, Inst. Superior Técnico	Lisboa, Portugal
IRAD	Inst. du Radium (Historical)	Paris, France	LIVP	Univ. of Liverpool	Liverpool, United Kingdom
ISNG	Inst. des Sciences Nucleaires (ISN)	Grenoble, France	LLL	Lawrence Livermore Lab. (Old name for LLNL)	Livermore, CA, USA
ISU	Iowa State Univ.	Ames, IA, USA	LLNL	Lawrence Livermore National Lab.	Livermore, CA, USA
ITEP	ITEP, Inst. of Theor. and Exp. Physics	Moscow, Russian Federation	LOCK	Lockheed Palo Alto Res. Lab	Palo Alto, CA, USA
ITHA	Ithaca College	Ithaca, NY, USA			

Abbreviations Used in the Full Listings (*Cont'd*)

LOIC	Imperial College of Science Tech. & Medicine	London, United Kingdom	NDAM	Univ. of Notre Dame	Notre Dame, IN, USA
LOQM	Univ. of London , Queen Mary & Westfield College	London, United Kingdom	NEAS	Northeastern Univ.	Boston, MA, USA
LOUC	University College London	London, United Kingdom	NEUC	Univ. de Neuchâtel	Neuchâtel, Switzerland
LOWC	Westfield College (Historical, see LOQM (Queen Mary and Westfield joined))	London, United Kingdom	NICEA	Univ. de Nice	Nice, France
LRL	U.C. Lawrence Radiation Lab. (Old name for LBL)	Berkeley, CA, USA	NICEO	Observatoire de Nice	Nice, France
LSU	Louisiana State Univ.	Baton Rouge, LA, USA	NIHO	Nihon Univ.	Tokyo, Japan
LUND	Univ. of Lund	Lund, Sweden	NIIG	Niigata Univ.	Niigata, Japan
LVLN	Univ. Catholique de Louvain	Louvain-la-Neuve, Belgium	NIJM	Univ. of Nijmegen	Nijmegen, The Netherlands
LYON	Institute de Physique Nucléaire de Lyon (IPN)	Villeurbanne, France	NIRS	Nat. Inst. Radiological Sciences	Chiba, Japan
MADE	Inst. de Estructura de la Materia	Madrid, Spain	NIU	Northern Illinois Univ.	De Kalb, IL, USA
MADR	C.I.E.M.A.T	Madrid, Spain	NMSU	New Mexico State Univ.	Las Cruces, NM, USA
MADU	Univ. Autónoma de Madrid C-XI	Madrid, Spain	NORD	Nordita	Copenhagen Ø, Denmark
MANI	Univ. of Manitoba	Winnipeg, MB, Canada	NOTT	Univ. of Nottingham	Nottingham, United Kingdom
MANZ	Johannes-Gutenberg-Univ.	Mainz, Germany	NOVO	BINP, Budker Inst. of Nuclear Physics	Novosibirsk, Russian Federation
MARB	Univ. Marburg	Marburg, Germany	NPOL	Polytechnic of North London	London, United Kingdom
MARS	Centre de Physique des Particules de Marseille	Marseille, France	NRL	Naval Research Lab	Washington, DC, USA
MASA	Univ. of Massachusetts	Amherst, MA, USA	NSF	National Science Foundation	Washington, DC, USA
MASB	Univ. of Massachusetts at Boston	Boston, MA, USA	NTUA	National Tech. Univ. of Athens	Athens, Greece
MASD	Univ. of Massachusetts Dartmouth	N. Dartmouth, MA, USA	NWES	Northwestern Univ.	Evanston, IL, USA
MCGI	McGill Univ.	Montreal, PQ, Canada	NYU	New York Univ.	New York, NY, USA
MCHS	Univ. of Manchester	Manchester, United Kingdom	OBER	Oberlin College	Oberlin, OH, USA
MCMS	McMaster Univ.	Hamilton, ON, Canada	OHIO	Ohio Univ.	Athens, OH, USA
MEIS	Meisei Univ.	Tokyo, Japan	OKAY	Okayama Univ.	Okayama, Japan
MELB	Univ. of Melbourne	Parkville, Victoria, Australia	OKLA	Univ. of Oklahoma	Norman, OK, USA
MEUD	Observatoire de Meudon	Meudon, France	OKSU	Oklahoma State Univ.	Stillwater, OK, USA
MICH	Univ. of Michigan	Ann Arbor, MI, USA	OREG	Univ. of Oregon	Eugene, OR, USA
MILA	Univ. di Milano	Milano, Italy	ORNL	Oak Ridge National Laboratory	Oak Ridge, TN, USA
MINN	Univ. of Minnesota	Minneapolis, MN, USA	ORSAY	Univ. de Paris Sud	Orsay, France
MISS	Univ. of Mississippi	University, MS, USA	ORST	Oregon State Univ.	Corvallis, OR, USA
MIT	MIT Massachusetts Inst. of Technology	Cambridge, MA, USA	OSAK	Osaka Univ.	Osaka, Japan
MIU	Maharishi International Univ.	Fairfield, IA, USA	OSKC	Osaka City Univ.	Osaka-shi, Japan
MIYA	Miyazaki Univ.	Miyazaki-shi, Japan	OSLO	Univ. of Oslo	Oslo, Norway
MONP	Univ. de Montpellier II	Montpellier, France	OSU	Ohio State Univ.	Columbus, OH, USA
MONS	Univ. de Mons-Hainaut	Mons, Belgium	OTTA	Univ. of Ottawa	Ottawa, ON, Canada
MONT	Univ. de Montréal ; Laboratoire de physique nucléaire	Montréal, PQ, Canada	OXF	University of Oxford	Oxford, United Kingdom
MONTC	Univ. de Montréal ; Centre de recherches mathématiques	Montréal, PQ, Canada	OXFTP	Univ. of Oxford	Oxford, United Kingdom
MOSU	Moscow State Univ.	Moscow, Russian Federation	PADO	Univ. di Padova , "G. Galilei"	Padova, Italy
MPCM	Max Planck Inst. fur Chemie	Mainz, Germany	PARIN	Univ. Paris VI et Paris VII, IN2P3/CNRS	Paris, France
MPEI	Moscow Physical Engineering Inst.	Moscow, Russian Federation	PARIS	Univ. de Paris (Historical)	Paris, France
MPIH	Max-Planck-Inst. für Kernphysik	Heidelberg, Germany	PARM	Univ. di Parma	Parma, Italy
MPIM	Max-Planck-Inst. für Physik	München, Germany	PAST	Institut Pasteur	Paris, France
MSU	Michigan State Univ.	East Lansing, MI, USA	PATR	Univ. of Patras	Patras, Greece
MTHO	Mount Holyoke College	South Hadley, MA, USA	PAVI	Univ. di Pavia	Pavia, Italy
MULH	Centre Univ. du Haut-Rhin	Mulhouse, France	PENN	Univ. of Pennsylvania	Philadelphia, PA, USA
MUNI	Univ. of München	Garching, Germany	PGIA	Univ. di Perugia	Perugia, Italy
MUNT	Tech. Univ. München	Garching, Germany	PISA	Univ. di Pisa	Pisa, Italy
MURA	Midwestern Univ. Research Assoc. (Historical in <i>Review</i>)	Stroughton, WI, USA	PISAI	INFN, Sez. di Pisa	Pisa, Italy
NAAS	North Americal Aviation Science Center (Historical in <i>Review</i>)	Thousand Oaks, CA, USA	PITT	Univ. of Pittsburgh	Pittsburgh, PA, USA
NAGO	Nagoya Univ.	Nagoya, Japan	PLAT	SUNY at Plattsburgh	Plattsburgh, NY, USA
NAPL	Univ. di Napoli	Napoli, Italy	PLRM	Univ. di Palermo	Palermo, Italy
NASA	NASA	Greenbelt, MD, USA	PNL	Battelle Memorial Inst.	Richland, WA, USA
NBS	U.S National Bureau of Standards (Old name for NIST)	Gaithersburg, MD, USA	PNPI	Petersburg Nuclear Physics Inst.	Gatchina, Russian Federation
NBSB	National Inst. Standards Tech.	Boulder, CO, USA	PPA	Princeton-Penn. Proton Accelerator (Historical in <i>Review</i>)	Princeton, NJ, USA
NCAR	National Center for Atmospheric Research	Boulder, CO, USA	PRAG	Inst. of Physics, ASCR	Prague, Czech Republic
			PRIN	Princeton Univ.	Princeton, NJ, USA
			PSI	Paul Scherrer Inst. (Old name for VILL)	Villigen PSI, Switzerland
			PSLL	Physical Science Lab	Las Cruces, NM, USA
			PSU	Penn State Univ.	University Park, PA, USA
			PUCB	Pontificia Univ. Católica do Rio de Janeiro	Rio de Janeiro, RJ, Brasil
			PUEB	High Energy Physics Group, Colegio de Fisica	Puebla, Pue, Mexico
			PURD	Purdue Univ.	Lafayette, IN, USA
			QUKI	Queen's Univ.	Kingston, ON, Canada
			RAL	Rutherford Appleton Lab.	Chilton, Didcot, Oxon., United Kingdom
			REGE	Univ. Regensburg	Regensburg, Germany

Abbreviations Used in the Full Listings (*Cont'd*)

REHO	Weizmann Inst. of Science	Rehovot, Israel	STLO	St. Louis Univ.	St. Louis, MO, USA
RHBL	Royal Holloway & Bedford New College	Egham, Surrey, United Kingdom	STOH	Stockholm Univ.	Stockholm, Sweden
RHEL	Rutherford High Energy Lab (Old name for RAL)	Chilton, Didcot, Oxon., United Kingdom	STON	SUNY at Stony Brook	Stony Brook, NY, USA
RICE	Rice Univ.	Houston, TX, USA	STRB	CRN, Centre des Recherches Nucl.	Strasbourg, France
RIKEN	Riken Inst. (Physical & Chemical Research)	Saitama, Japan	STUT	Univ. Stuttgart	Stuttgart, Germany
RIKK	Rikkyo Univ.	Tokyo, Japan	STUTM	Max-Planck-Inst.	Stuttgart, Germany
RIS	Rowland Inst. for Science	Cambridge, MA, USA	SUGI	Sugiyama Jogakuen Univ.	Aichi, Japan
RISC	Rockwell International	Thousand Oaks, CA, USA	SURR	Univ. of Surrey	Guildford, Surrey, United Kingdom
RISL	Universities Research Re- actor	Risley, Warrington, United Kingdom	SUSS	Univ. of Sussex	Brighton, United Kingdom
RISO	Riso National Laboratory	Roskilde, Denmark	SYDN	Univ. of Sydney	Sydney, NSW, Australia
RL	Rutherford High Energy Lab (Old name for RAL)	Chilton, Didcot, Oxon., United Kingdom	SYRA	Syracuse Univ.	Syracuse, NY, USA
RMCS	Royal Military Coll. of Sci- ence	Swindon, Wilts., United Kingdom	TAJK	Acad. Sci., Tadjhik SSR	Dushanbe, Tadjhikstan
ROCH	Univ. of Rochester	Rochester, NY, USA	TAMU	Texas A&M Univ.	College Station, TX, USA
ROCK	Rockefeller Univ.	New York, NY, USA	TATA	Tata Inst. of Fundamental Research	Bombay, India
ROMA	Univ. di Roma (Historical)	Roma, Italy	TBIL	Tbilisi State University	Tbilisi, Republic of Georgia
ROMAI	INFN, Sez. di Roma	Roma, Italy	TELA	Tel-Aviv Univ.	Tel Aviv, Israel
ROSE	Rose-Hulman Inst. of Tech- nology	Terre Haute IN, USA	TELE	Teledyne Brown Engineer- ing	Huntsville, AL, USA
RPI	Rensselaer Polytechnic Inst.	Troy, NY, USA	TEMP	Temple Univ.	Philadelphia, PA, USA
RUTG	Rutgers Univ.	Piscataway, NJ, USA	TENN	Univ. of Tennessee	Knoxville, TN, USA
SACL	CE Saclay	Gif-sur-Yvette, France	TEXA	Univ. of Texas at Austin	Austin, TX, USA
SACLD	CE Saclay; DAPNIA	Gif-sur-Yvette, France	TGAK	Tokyo Gakugei Univ.	Tokyo, Japan
SAGA	Saga Univ.	Saga-shi, Japan	TGU	Tohoku Gakuin Univ.	Miyagi, Japan
SANG	Kyoto Sangyo Univ.	Kyoto-shi, Japan	THES	Univ. of Thessaloniki	Thessaloniki, Greece
SANI	Physics Lab., Ist. Superiore di Sanità	Roma, Italy	TINT	Tokyo Inst. of Technology	Tokyo, Japan
SASK	Univ. of Saskatchewan	Saskatoon, SK, Canada	TISA	Sagamihara Inst. of Space & Astronautical Sci.	Kanagawa, Japan
SASSO	Lab. Naz. del Gran Sasso dell'INFN	Assergi (L'Aquila), Italy	TMSK	Inst. Nuclear Physics	Tomsk, Russian Federation
SAVO	Univ. de Savoie	Chambery, France	TMTC	Tokyo Metropolitan Coll. Tech.	Tokyo, Japan
SBER	California State Univ.	San Bernardino, CA, USA	TMU	Tokyo Metropolitan Univ.	Tokyo, Japan
SCIT	Science Univ. of Tokyo	Tokyo, Japan	TNTO	Univ. of Toronto	Toronto, ON, Canada
SCOT	Scottish Univ. Research and Reactor Ctr.	Glasgow, United Kingdom	TOHO	Toho Univ.	Chiba, Japan
SCUC	Univ. of South Carolina	Columbia, SC, USA	TOHOK	Tohoku Univ.	Sendai, Japan
SEAT	Seattle Pacific Coll.	Seattle, WA, USA	TOKA	Tokai Univ.	Shimizu, Japan
SEIB	Austrian Research Center, Seibersdorf LTD.	Seibersdorf, Austria	TOKMS	Univ. of Tokyo; Meson Sci- ence Laboratory	Tokyo, Japan
SEOU	Korea Univ.	Seoul, Republic of Korea	TOKU	Univ. of Tokushima	Tokushima-shi, Japan
SEOUL	Seoul National Univ.	Seoul, Republic of Korea	TOKY	Univ. of Tokyo; Physics Dept.	Tokyo, Japan
SERP	IHEP, Inst. for High Energy Physics (Also known as Ser- pukhov)	Protvino, Russian Federation	TORI	Univ. degli Studi di Torino	Torino, Italy
SETO	Seton Hall Univ.	South Orange, NJ, USA	TPTI	Lab. of High Energy Phys.	Tashkent, Republic of Uzbek- istan
SFLA	Univ. of South Florida	Tampa, FL, USA	TRIN	Trinity College	Dublin, Ireland
SFRA	Simon Fraser University	Burnaby, BC, Canada	TRIU	TRIUMF	Vancouver, BC, Canada
SFSU	California State Univ.	San Francisco, CA, USA	TRST	Univ. degli Studi di Trieste	Trieste, Italy
SHEF	Univ. of Sheffield	Sheffield, United Kingdom	TRSTI	INFN, Sez. di Trieste	Trieste, Italy
SHMP	Univ. of Southampton	Southampton, United Kingdom	TRSTT	Univ. di Trieste	Trieste, Italy
SIEG	Univ.-Gesamthochschule- Siegen	Siegen, Germany	TSUK	Univ. of Tsukuba	Ibaraki-ken, Japan
SILES	Univ. of Silesia	Katowice, Poland	TTAM	Tamagawa Univ.	Tokyo, Japan
SIN	Swiss Inst. of Nuclear Re- search (Old name for VILL)	Villigen, Switzerland	TUAT	Tokyo Univ. of Agriculture Tech.	Tokyo, Japan
SISSA	Scuola Internazionale Superi- ore di Studi Avanzati	Trieste, Italy	TUBIN	Univ. Tübingen	Tübingen, Germany
SLAC	Stanford Linear Acceler- ator Center	Stanford, CA, USA	TUFTS	Tufts Univ.	Medford, MA, USA
SLOV	Inst. of Physics, Slovak Acad. of Sciences	Bratislava, Slovak Republic	TUW	Technische Univ. Wien	Vienna, Austria
SMU	Southern Methodist Univ.	Dallas, TX, USA	UCB	Univ. of California (Berke- ley); Dept. of Physics	Berkeley, CA, USA
SNSP	Scuola Normale Superiore	Pisa, Italy	UCD	Univ. of California (Davis)	Davis, CA, USA
SOFI	Inst. for Nuclear Research and Nuclear Energy	Sofia, Bulgaria	UCI	Univ. of California (Irvine)	Irvine, CA, USA
SOFU	Univ. of Sofia	Sofia, Bulgaria	UCLA	Univ. of California (Los Angeles)	Los Angeles, CA, USA
SPAUL	Univ. de São Paulo	São Paulo, SP, Brasil	UCND	Union Carbide Corp.	Oak Ridge, TN, USA
SPIFT	Inst. de Física Teórica (IFT)	São Paulo, SP, Brasil	UCR	Univ. of California (River- side)	Riverside, CA, USA
SSL	Univ. of California (Berke- ley); Space Sciences Lab	Berkeley, CA, USA	UCSB	Univ. of California (Santa Barbara); Physics Dept.	Santa Barbara, CA, USA
STAN	Stanford Univ.	Stanford, CA, USA	UCSBT	Univ. of California (Santa Barbara); Institute of Theo- retical Physics	Santa Barbara, CA, USA
STEV	Stevens Inst. of Tech.	Hoboken, NJ, USA	UCSC	Univ. of California (Santa Cruz)	Santa Cruz, CA, USA
			UCSD	Univ. of California (San Diego)	La Jolla, CA, USA
			UMD	Univ. of Maryland	College Park, MD, USA
			UNC	Univ. of North Carolina	Greensboro, NC, USA

Abbreviations Used in the Full Listings (*Cont'd*)

UNCCH	Univ. of North Carolina at Chapel Hill	Chapel Hill, NC, USA
UNCS	Union College	Schenectady, NY, USA
UNH	Univ. of New Hampshire	Durham, NH, USA
UNM	Univ. of New Mexico	Albuquerque, NM, USA
UOEH	Univ. of Occupational and Environmental Health	Kitakyushu , Japan
UPNJ	Upsala College	East Orange, NJ, USA
UPPS	Uppsala Univ.	Uppsala , Sweden
UPR	Univ. of Puerto Rico	Rio Piedras, PR, USA
URI	Univ. of Rhode Island	Kingston, RI, USA
USC	Univ. of Southern California	Los Angeles, CA, USA
USF	Univ. of San Francisco	San Francisco, CA, USA
UTAH	Univ. of Utah	Salt Lake City, UT, USA
UTRE	Univ. of Utrecht	Utrecht, The Netherlands
UTRO	Univ. of Trondheim	Dragvoll, Norway
UZINR	Acad. Sci., Ukrainian SSR	Uzhgorod , Ukraine
VALE	Univ. de Valencia	Burjassot, Valencia , Spain
VALP	Valparaiso Univ.	Valparaiso, IN, USA
VAND	Vanderbilt Univ.	Nashville, TN, USA
VASS	Vassar College	Poughkeepsie, NY, USA
VICT	Univ. of Victoria	Victoria, BC, Canada
VIEN	Inst. für Hochenergiephysik d. Österr. Akademie d. Wissenschaften	Vienna , Austria
VILL	Inst. for Particle Physics of ETH Zürich (Was Paul Scherrer Institute)	Villigen PSI, Switzerland
VIRG	Univ. of Virginia	Charlottesville, VA, USA
VPI	Virginia Polytechnic Inst. and State Univ.	Blacksburg, VA, USA
VRJ	Vrije Univ.	HV Amsterdam , The Netherlands
WABRNE	Eidgenössisches Amt für Messwesen	Waber , Switzerland
WARS	Warsaw Univ.	Warsaw, Poland
WASCR	Waseda Univ.; Cosmic Ray Division	Tokyo, Japan
WASH	Univ. of Washington	Seattle, WA, USA
WASU	Waseda Univ.; Dept. of Physics, High Energy Physics Group	Tokyo, Japan
WAYN	Wayne State Univ.	Detroit, MI, USA
WESL	Wesleyan Univ.	Middletown, CT, USA
WIEN	Univ. Wien	Vienna, Austria
WILL	Coll. of William and Mary	Williamsburg, VA, USA
WINR	Inst. for Nuclear Studies	Warsaw , Poland
WISC	Univ. of Wisconsin	Madison, WI, USA
WITW	Univ. of the Witwatersrand	Wits, South Africa
WMIU	Western Michigan Univ.	Kalamazoo, MI, USA
WONT	The Univ. of Western Ontario	London, ON, Canada
WOOD	Woodstock College (No longer in existence)	Woodstock, MD, USA
WUPP	Univ. of Wuppertal	Wuppertal, Germany
WURZ	Univ. Würzburg	Würzburg, Germany
WUSL	Washington Univ.	St. Louis, MO, USA
WYOM	Univ. of Wyoming	Laramie, WY, USA
YALE	Yale Univ.	New Haven, CT, USA
YCC	Yokohama Coll. of Commerce	Yokohama, Japan
YERE	Yerevan Physics Inst.	Yerevan, Armenia
YOKO	Yokohama National Univ.	Yokohama-shi, Japan
YORKC	York Univ.	North York, ON, Canada
ZAGR	Zagreb Univ.	Zagreb, Croatia
ZARA	Univ. de Zaragoza	Zaragoza, Spain
ZEEM	Univ. van Amsterdam	TV Amsterdam, The Netherlands
ZURI	Univ. Zürich	Zürich, Switzerland

GAUGE AND HIGGS BOSONS

γ	1351
g (gluon)	1351
graviton	1351
W	1351
Z	1353
Higgs Bosons — H^0 and H^\pm	1365
Heavy Bosons Other than Higgs Bosons	1372
Axions (A^0) and Other Very Light Bosons	1377

Notes in the Gauge and Higgs Boson Listings

Note on the Z Boson	1353
Note on the Higgs Boson	1365
Note on the Z'_i Searches	1372
Note on Axions	1377
Invisible A^0 (Axion) Mass Limits from Astrophysics and Cosmology	1382

See key on page 1343

Gauge & Higgs Boson Full Listings
 $\gamma, g, \text{graviton}, W$

GAUGE AND HIGGS BOSONS

γ

$I(J^{PC}) = 0,1(1^{--})$

γ MASS

For a review of the photon mass, see BYRNE 77.

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 3	$\times 10^{-27}$	CHIBISOV	76	Galactic magnetic field
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<(4.73 \pm 0.45) \times 10^{-12}$		¹ CHERNIKOV	92 SQID	Ampere-law null test
$<(9.0 \pm 8.1) \times 10^{-10}$		² RYAN	85	Coulomb-law null test
< 6	$\times 10^{-16}$	99.7 DAVIS	75	Jupiter magnetic field
< 7.3	$\times 10^{-16}$	HOLLWEG	74	Alfvén waves
< 6	$\times 10^{-17}$	³ FRANKEN	71	Low freq. res. cir.
< 1	$\times 10^{-14}$	WILLIAMS	71 CNTR	Tests Gauss law
< 2.3	$\times 10^{-15}$	GOLDHABER	68	Satellite data
< 6	$\times 10^{-15}$	³ PATEL	65	Satellite data
< 6	$\times 10^{-15}$	GINTSBURG	64	Satellite data

¹ CHERNIKOV 92 measures the photon mass at 1.24 K, following a theoretical suggestion that electromagnetic gauge invariance might break down at some low critical temperature. See the erratum for a correction, included here, to the published result.
² RYAN 85 measures the photon mass at 1.36 K (see the footnote to CHERNIKOV 92).
³ See criticism questioning the validity of these results in KROLL 71 and GOLDHABER 71.

γ CHARGE

VALUE (e)	DOCUMENT ID	TECN	COMMENT
$< 2 \times 10^{-32}$	COCCONI	88 TOF	Pulsar $f_1 - f_2$ TOF
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$< 2 \times 10^{-28}$	⁴ COCCONI	92	VLBA radio telescope resolution

⁴ See COCCONI 92 for less stringent limits in other frequency ranges.

γ REFERENCES

CHERNIKOV	92	PRL 68 3383	+Gerber, Ott, Gerber	(ETH)
Also	92B	PRL 69 2999 (erratum)	Chernikov, Gerber, Ott, Gerber	(ETH)
COCCONI	88	AJP 60 750		(CERN)
COCCONI	88	PL B206 705		(CERN)
RYAN	85	PR D32 802	+Accetta, Austin	(PRIN)
BYRNE	77	Ast.Sp.Sci. 46 115		(LOIC)
CHIBISOV	76	SPU 19 624		(LEBD)
DAVIS	75	PRL 35 1402	+Goldhaber, Nieto	(CIT, STON, LASL)
HOLLWEG	74	PRL 32 961		(NCAR)
FRANKEN	71	PRL 26 115	+Ampulski	(MICH)
GOLDHABER	71	RMP 43 277	+Nieto	(STON, BOHR, UCSB)
KROLL	71	PRL 26 1395		(SLAC)
WILLIAMS	71	PRL 26 721	+Faller, Hill	(WESL)
GOLDHABER	68	PRL 21 567	+Nieto	(STON)
PATEL	65	PL 14 105		(DUKE)
GINTSBURG	64	Sov. Astr. AJ7 536		(ASCI)

**g
or gluon**

$I(J^P) = 0(1^-)$

SU(3) color octet

Mass $m = 0$. Theoretical value. A mass as large as a few MeV may not be precluded.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	ABREU	92E DLPH	Spin 1, not 0
	ALEXANDER	91H OPAL	Spin 1, not 0
	BEHREND	82D CELL	Spin 1, not 0
	BERGER	80D PLUT	Spin 1, not 0
	BRANDELIK	80C TASS	Spin 1, not 0

ABREU	92E	PL B274 498	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ALEXANDER	91H	ZPHY C52 543	+Allison, Allport, Anderson+	(OPAL Collab.)
BEHREND	82D	PL B110 329	+Chen, Field, Guempel, Schroeder+	(CELLO Collab.)
BERGER	80D	PL B97 459	+Genzel, Grigull, Lackas+	(PLUTO Collab.)
BRANDELIK	80C	PL B97 453	+Braunschweig, Gather, Kadansky+	(TASSO Collab.)

graviton

$J = 2$

OMITTED FROM SUMMARY TABLE

graviton MASS

All of the following limits are obtained assuming Yukawa potential in weak field limit. VANDAM 70 argue that a massive field cannot approach general relativity in the zero-mass limit; however, see GOLDHABER 74 and references therein. h_0 is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$.

VALUE (eV)	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
$< 2 \times 10^{-29} h_0^{-1}$	¹ DAMOUR	91 Binary pulsar PSR 1913+16
	GOLDHABER	74 Rich clusters
$< 7 \times 10^{-28}$	HARE	73 Galaxy
$< 8 \times 10^4$	HARE	73 2γ decay

¹ DAMOUR 91 is an analysis of the orbital period change in binary pulsar PSR 1913+16, and confirms the general relativity prediction to 0.8%. "The theoretical importance of the [rate of orbital period decay] measurement has long been recognized as a direct confirmation that the gravitational interaction propagates with velocity c (which is the immediate cause of the appearance of a damping force in the binary pulsar system) and thereby as a test of the existence of gravitational radiation and of its quadrupolar nature." TAYLOR 93 adds that orbital parameter studies now agree with general relativity to 0.5%, and set limits on the level of scalar contribution in the context of a family of tensor [spin 2]-biscalar theories.

graviton REFERENCES

TAYLOR	93	Nature 355 132	+Wolsczcan, Damour+	(PRIN, ARCO, BURE, CARL) J
DAMOUR	91	APJ 366 501	+Taylor	(BURE, MEUD, PRIN)
GOLDHABER	74	PR D9 119	+Nieto	(LANL, STON)
HARE	73	CJP 51 431		(SASK)
VANDAM	70	NP B22 397	van Dam, Veltman	(UTRE)

W

$J = 1$

W MASS

The fit uses the W and Z mass, mass difference, and mass ratio measurements.

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
80.22 ± 0.26 OUR FIT				
80.1 ± 0.4 OUR AVERAGE				
$80.84 \pm 0.22 \pm 0.83$	2065	¹ ALITTI	92B UA2	$E_{cm}^{pp} = 630 \text{ GeV}$
79.91 ± 0.39	1722	² ABE	90G CDF	$E_{cm}^{pp} = 1800 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$80.79 \pm 0.31 \pm 0.84$		³ ALITTI	90B UA2	$E_{cm}^{pp} = 546,630 \text{ GeV}$
$80.0 \pm 3.3 \pm 2.4$	22	⁴ ABE	89I CDF	$E_{cm}^{pp} = 1800 \text{ GeV}$
$82.7 \pm 1.0 \pm 2.7$	149	⁵ ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630 \text{ GeV}$
$81.8 \pm 6.0 \pm 2.6$	46	⁶ ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630 \text{ GeV}$
$89 \pm 3 \pm 6$	32	⁷ ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630 \text{ GeV}$
$80.2 \pm 0.6 \pm 1.4$	251	⁸ ANSARI	87 UA2	Repl. by ALITTI 90B
$81.2 \pm 1.0 \pm 1.4$	119	⁸ APPEL	86 UA2	Repl. by ANSARI 87
$83.5 \pm 1.1 \pm 1.0$	86	⁹ ARNISON	86 UA1	Repl. by ALBAJAR 89
$81. \pm 6. \pm 7.$	14	¹⁰ ARNISON	84D UA1	Repl. by ALBAJAR 89
$83.1 \pm 1.9 \pm 1.3$	37	BAGNAIA	84 UA2	Repl. by ALITTI 90B
$81. \pm 5.$	6	ARNISON	83 UA1	Repl. by ARNISON 83D
80.9 ± 2.9	27	ARNISON	83D UA1	Repl. by ARNISON 86
81.0 ± 2.8		BAGNAIA	83 UA2	Repl. by BAGNAIA 84
$80. \pm 10. \pm 6.$	4	BANNER	83B UA2	Repl. by ALITTI 90B

¹ ALITTI 92B result has two contributions to the systematic error (± 0.83); one (± 0.81) cancels in m_W/m_Z and one (± 0.17) is noncancelling. These were added in quadrature. We choose the ALITTI 92B value without using the LEP m_Z value, because we perform our own combined fit.
² ABE 90G result from $W \rightarrow e\nu$ is $79.91 \pm 0.35 \pm 0.24 \pm 0.19(\text{scale}) \text{ GeV}$ and from $W \rightarrow \mu\nu$ is $79.90 \pm 0.53 \pm 0.32 \pm 0.08(\text{scale}) \text{ GeV}$.
³ There are two contributions to the systematic error (± 0.84): one (± 0.81) which cancels in m_W/m_Z and one (± 0.21) which is non-cancelling. These were added in quadrature.
⁴ ABE 89I systematic error dominated by the uncertainty in the absolute energy scale.
⁵ ALBAJAR 89 result is from a total sample of 299 $W \rightarrow e\nu$ events.
⁶ ALBAJAR 89 result is from a total sample of 67 $W \rightarrow \mu\nu$ events.
⁷ ALBAJAR 89 result is from $W \rightarrow \tau\nu$ events.
⁸ There are two contributions to the systematic error (± 1.4): one (± 1.3) which cancels in m_W/m_Z and one (± 0.5) which is non-cancelling. These were added in quadrature.
⁹ This is enhanced subsample of 172 total events.
¹⁰ Using $W^\pm \rightarrow \mu^\pm \nu$.

Gauge & Higgs Boson Full Listings

W

W/Z MASS RATIO

The fit uses the W and Z mass, mass difference, and mass ratio measurements.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.8798 ± 0.0028 OUR FIT				
0.8813 ± 0.0036 ± 0.0019	156	11 ALITTI	92B UA2	$E_{cm}^{pp} = 630$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.8831 ± 0.0048 ± 0.0026		11 ALITTI	90B UA2	$E_{cm}^{pp} = 546,630$ GeV
11 Scale error cancels in this ratio.				

 $m_Z - m_W$

The fit uses the W and Z mass, mass difference, and mass ratio measurements.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.96 ± 0.26 OUR FIT			
10.4 ± 1.4 ± 0.8	ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
11.3 ± 1.3 ± 0.9	ANSARI	87 UA2	$E_{cm}^{pp} = 546,630$ GeV

 $m_{W^+} - m_{W^-}$

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.19 ± 0.58	1722	ABE	90G CDF	$E_{cm}^{pp} = 1800$ GeV

W WIDTH

The widths labelled "extracted value" are obtained by measuring $R = \sigma(W \rightarrow e\nu)/\sigma(Z \rightarrow e^+e^-)$ which is equal to $[\sigma(W)/\sigma(Z)] [\Gamma(W \rightarrow e\nu)/\Gamma(Z \rightarrow e^+e^-)]$. The bracketed quantities can be calculated with plausible reliability. $\Gamma(W)$ is then extracted by using a value of $\Gamma(Z)$ measured at LEP. The Standard Model prediction is 2.067 ± 0.021 (ROSNER 94).

VALUE (GeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.08 ± 0.07 OUR AVERAGE					
2.063 ± 0.061 ± 0.060			12 ABE	94B CDF	Extracted value
2.10 $^{+0.14}_{-0.13}$ ± 0.09	3559		13 ALITTI	92 UA2	Extracted value
2.18 $^{+0.26}_{-0.24}$ ± 0.04			14 ALBAJAR	91 UA1	Extracted value
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.16 ± 0.17			15 ABE	92I CDF	Repl. by ABE 94B
2.12 ± 0.20			16 ABE	90 CDF	Repl. by ABE 92I
2.30 ± 0.19 ± 0.06			17 ALITTI	90C UA2	Extracted value
< 5.4	90	149	18 ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV
2.8 $^{+1.4}_{-1.5}$ ± 1.3	149		18 ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV
< 7	90	251	ANSARI	87 UA2	$E_{cm}^{pp} = 546,630$ GeV
< 7	90	119	APPEL	86 UA2	$E_{cm}^{pp} = 546,630$ GeV
< 6.5	90	86	19 ARNISON	86 UA1	Repl. by ALBAJAR 89
< 7	90	27	ARNISON	83D UA1	Repl. by ARNISON 86

12 ABE 94B measured $R = 10.90 \pm 0.32 \pm 0.29$. The values of $\sigma(Z)$ and $\sigma(W)$ come from $O(\alpha_s^2)$ calculations using $m_W = 80.24 \pm 0.10$ GeV, and $m_Z = 91.188 \pm 0.007$ GeV along with the corresponding value of $\sin^2\theta_W = 0.2325 \pm 0.005$. They use $\sigma(W)/\sigma(Z) = 3.33 \pm 0.03$, $\Gamma(Z) = 2.492 \pm 0.007$ GeV, and $\Gamma(W \rightarrow e\nu_e)/\Gamma(Z \rightarrow e^+e^-) = 2.710 \pm 0.018$.

13 ALITTI 92 measured $R = 10.4^{+0.7}_{-0.6} \pm 0.3$. The values of $\sigma(Z)$ and $\sigma(W)$ come from $O(\alpha_s^2)$ calculations using $m_W = 80.14 \pm 0.27$ GeV, and $m_Z = 91.175 \pm 0.021$ GeV along with the corresponding value of $\sin^2\theta_W = 0.2274$. They use $\sigma(W)/\sigma(Z) = 3.26 \pm 0.07 \pm 0.05$ and $\Gamma(Z) = 2.487 \pm 0.010$ GeV.

14 ALBAJAR 91 measured $R = 9.5^{+1.1}_{-1.0}$ (stat. + syst.). $\sigma(W)/\sigma(Z)$ is calculated in QCD at the parton level using $m_W = 80.18 \pm 0.28$ GeV and $m_Z = 91.172 \pm 0.031$ GeV along with $\sin^2\theta_W = 0.2322 \pm 0.0014$. They use $\sigma(W)/\sigma(Z) = 3.23 \pm 0.05$ and $\Gamma(Z) = 2.498 \pm 0.020$ GeV.

15 ABE 92I report $1216 \pm 38^{+27}_{-31}$ $W \rightarrow \mu\nu$ and $106 \pm 10^{+0.2}_{-0.1}$ $Z \rightarrow \mu^+\mu^-$ events which are combined with 2426 $W \rightarrow e\nu$ events of ABE 91C to derive the ratio σ_W/σ_Z . $B(W \rightarrow \ell\nu)/\sigma_B(Z \rightarrow \ell^+\ell^-) = 10.0 \pm 0.6 \pm 0.4$. Finally the value of $\Gamma(Z)$ measured by LEP 92 is used to extract $\Gamma(W)$.

16 ABE 90 extract $\Gamma(W) = 2.19 \pm 0.20$ by using the value $\Gamma(Z) = 2.57 \pm 0.07$ GeV. However, in ABE 91C they update their analysis with a new LEP value $\Gamma(Z) = 2.496 \pm 0.016$; the value $\Gamma(W) = 2.12 \pm 0.20$ above reflects this update. They measured $R = 10.2 \pm 0.8 \pm 0.4$, assumed $\sin^2\theta_W = 0.229 \pm 0.007$, and took predicted values $\sigma(W)/\sigma(Z) = 3.23 \pm 0.03$ and $\Gamma(W \rightarrow e\nu)/\Gamma(Z \rightarrow e^+e^-) = 2.70 \pm 0.02$. This yields $\Gamma(W)/\Gamma(Z) = 0.85 \pm 0.08$. The quoted error for $\Gamma(W)$ includes systematic uncertainties. $E_{cm}^{pp} = 1800$ GeV.

17 ALITTI 90C used the same technique as described for ABE 90. They measured $R = 9.38^{+0.82}_{-0.72} \pm 0.25$, obtained $\Gamma(W)/\Gamma(Z) = 0.902 \pm 0.074 \pm 0.024$. Using $\Gamma(Z) = 2.546 \pm 0.032$ GeV, they obtained the $\Gamma(W)$ value quoted above and the limits $\Gamma(W) < 2.56$ (2.64) GeV at the 90% (95%) CL. $E_{cm}^{pp} = 546,630$ GeV.

18 ALBAJAR 89 result is from a total sample of 299 $W \rightarrow e\nu$ events.

19 If systematic error is neglected, result is $2.7^{+1.4}_{-1.5}$ GeV. This is enhanced subsample of 172 total events.

W ANOMALOUS MAGNETIC MOMENT ($\Delta\kappa$)

The full magnetic moment is given by $\mu_W = e(1 + \kappa + \lambda)/2m_W$. In the Standard Model, at tree level, $\kappa = 1$ and $\lambda = 0$. Some papers have defined $\Delta\kappa = 1 - \kappa$ and assume that $\lambda = 0$. Note that the electric quadrupole moment is given by $-e(\kappa - \lambda)/m_W^2$. A description of the parameterization of these moments and additional references can be found in HAGIWARA 87 and BAUR 88. The parameter Λ appearing in the theoretical limits below is a regularization cutoff which roughly corresponds to the energy scale where the structure of the W boson becomes manifest.

VALUE ($e/2m_W$)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
	20 ALITTI	92C UA2
	21 SAMUEL	92 THEO
	22 SAMUEL	91 THEO
	23 GRIFOLS	88 THEO
	24 GROTCHE	87 THEO
	25 VANDERBIJ	87 THEO
	26 GRAU	85 THEO
	27 SUZUKI	85 THEO
	28 HERZOG	84 THEO

20 ALITTI 92C measure $\kappa = 1^{+2.6}_{-2.2}$ and $\lambda = 0^{+1.7}_{-1.8}$ in $p\bar{p} \rightarrow e\nu\gamma + X$ at $\sqrt{s} = 630$ GeV. At 95%CL they report $-3.5 < \kappa < 5.9$ and $-3.6 < \lambda < 3.5$.

21 SAMUEL 92 use preliminary CDF and UA2 data and find $-2.4 < \kappa < 3.7$ at 96%CL and $-3.1 < \lambda < 4.2$ at 95%CL respectively. They use data for $W\gamma$ production and radiative W decay.

22 SAMUEL 91 use preliminary CDF data for $p\bar{p} \rightarrow W\gamma X$ to obtain $-11.3 \leq \Delta\kappa \leq 10.9$. Note that their $\kappa = 1 - \Delta\kappa$.

23 GRIFOLS 88 uses deviation from ρ parameter to set limit $\Delta\kappa \lesssim 65 (M_W^2/\Lambda^2)$.

24 GROTCHE 87 finds the limit $-37 < \Delta\kappa < 73.5$ (90% CL) from the experimental limits on $e^+e^- \rightarrow \nu\bar{\nu}\gamma$ assuming three neutrino generations and $-19.5 < \Delta\kappa < 56$ for four generations. Note their $\Delta\kappa$ has the opposite sign as our definition.

25 VANDERBIJ 87 uses existing limits to the photon structure to obtain $|\Delta\kappa| < 33 (m_W/\Lambda)$. In addition VANDERBIJ 87 discusses problems with using the ρ parameter of the Standard Model to determine $\Delta\kappa$.

26 GRAU 85 uses the muon anomaly to derive a coupled limit on the anomalous magnetic dipole and electric quadrupole (λ) moments $1.05 > \Delta\kappa \ln(\Lambda/m_W) + \lambda/2 > -2.77$. In the Standard Model $\lambda = 0$.

27 SUZUKI 85 uses partial-wave unitarity at high energies to obtain $|\Delta\kappa| \lesssim 190 (m_W/\Lambda)^2$. From the anomalous magnetic moment of the muon, SUZUKI 85 obtains $|\Delta\kappa| \lesssim 2.2/\ln(\Lambda/m_W)$. Finally SUZUKI 85 uses deviations from the ρ parameter and obtains a very qualitative, order-of-magnitude limit $|\Delta\kappa| \lesssim 150 (m_W/\Lambda)^4$ if $|\Delta\kappa| \ll 1$.

28 HERZOG 84 consider the contribution of W -boson to muon magnetic moment including anomalous coupling of $W W\gamma$. Obtain a limit $-1 < \Delta\kappa < 3$ for $\Lambda \gtrsim 1$ TeV.

W⁺ DECAY MODES

W^+ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^+\nu$	(10.8 ± 0.4) %	
Γ_2 $\mu^+\nu$	(10.6 ± 0.7) %	
Γ_3 $\tau^+\nu$	(10.8 ± 1.0) %	
Γ_4 $\ell^+\nu$	[a] (10.7 ± 0.5) %	
Γ_5 hadrons	(67.8 ± 1.5) %	
Γ_6 $\pi^+\gamma$	< 5 × 10 ⁻⁴	95%

[a] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 7 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.1$ for 4 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	34		
x_3	43	15	
x_5	-70	-64	-82
	x_1	x_2	x_3

See key on page 1343

Gauge & Higgs Boson Full Listings

W, Z

W BRANCHING RATIOS

$\Gamma(e^+\nu)/\Gamma_{\text{total}}$				Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.108 ± 0.004 OUR FIT				
0.109 ± 0.004 OUR AVERAGE				
0.1094 ± 0.0033 ± 0.0031		29 ABE	94B CDF	$E_{\text{cm}}^{\text{PD}} = 1800 \text{ GeV}$
0.10 ± 0.014 $^{+0.02}_{-0.03}$	248	30 ANSARI	87C UA2	$E_{\text{cm}}^{\text{PD}} = 546,630 \text{ GeV}$
0.106 ± 0.0096	2426	31 ABE	91C CDF	Repl. by ABE 94B
seen	299	32 ALBAJAR	89 UA1	$E_{\text{cm}}^{\text{PD}} = 546,630 \text{ GeV}$
seen	119	APPEL	86 UA2	$E_{\text{cm}}^{\text{PD}} = 546,630 \text{ GeV}$
seen	172	ARNISON	86 UA1	Repl. by ALBAJAR 89

29 ABE 94B result is from a measurement of $\sigma_B(W \rightarrow e\nu)/\sigma_B(Z \rightarrow e^+e^-) = 10.90 \pm 0.32 \pm 0.29$, the theoretical prediction for the cross section ratio, the experimental knowledge of $\Gamma(Z \rightarrow e^+e^-) = 83.33 \pm 0.30 \text{ MeV}$, and $\Gamma(Z) = 2.492 \pm 0.007$.

30 The first error was obtained by adding the statistical and systematic experimental uncertainties in quadrature. The second error reflects the dependence on theoretical prediction of total W cross section: $\sigma(546 \text{ GeV}) = 4.7^{+1.4}_{-0.7} \text{ nb}$ and $\sigma(630 \text{ GeV}) = 5.8^{+1.8}_{-1.0} \text{ nb}$. See ALTARELLI 85b.

31 ABE 91C result is from a measurement of $\sigma_B(W \rightarrow e\nu)/\sigma_B(Z \rightarrow e^+e^-)$, the theoretical prediction for the cross section ratio, and the experimental knowledge of $\Gamma(Z \rightarrow e^+e^-)/\Gamma(Z \rightarrow \text{all})$.

32 ALBAJAR 89 experiment determines values of branching ratio times production cross section.

$\Gamma(\mu^+\nu)/\Gamma_{\text{total}}$				Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.10 ± 0.01	1216	33 ABE	92I CDF	$E_{\text{cm}}^{\text{PD}} = 1.8 \text{ TeV}$

33 ABE 92I quote the inverse quantity as 9.9 ± 1.2 which we have inverted.

$\Gamma(\tau^+\nu)/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.108 ± 0.010 OUR FIT				

$\Gamma(\ell^+\nu)/\Gamma_{\text{total}}$				$\Gamma_4/\Gamma = (\frac{1}{3}\Gamma_1 + \frac{1}{3}\Gamma_2 + \frac{1}{3}\Gamma_3)/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.107 ± 0.005 OUR FIT				

0.104 ± 0.008

34 1216 ± 38 $^{+27}_{-31}$ $W \rightarrow \mu\nu$ events from ABE 92I and 2426 $W \rightarrow e\nu$ events of ABE 91C. ABE 92I give the inverse quantity as 9.6 ± 0.7 and we have inverted.

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$				Γ_5/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.678 ± 0.016 OUR FIT				

$\Gamma(\mu^+\nu)/\Gamma(e^+\nu)$				Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.02 ± 0.08	1216	35 ABE	92I CDF	$E_{\text{cm}}^{\text{PD}} = 1.8 \text{ TeV}$

1.00 ± 0.14 ± 0.08

1.24 $^{+0.6}_{-0.4}$

35 ABE 92I obtain $\sigma_W B(W \rightarrow \mu\nu) = 2.21 \pm 0.07 \pm 0.21$ and combine with ABE 91C $W B(W \rightarrow e\nu)$ to give a ratio of the couplings from which we derive this measurement.

$\Gamma(\tau^+\nu)/\Gamma(e^+\nu)$				Γ_3/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.00 ± 0.08 OUR FIT				
1.00 ± 0.08 OUR AVERAGE				

0.94 ± 0.14

1.04 ± 0.08 ± 0.08

1.02 ± 0.20 ± 0.12

0.995 ± 0.112 ± 0.083

1.02 ± 0.20 ± 0.10

36 ABE 92E use two procedures for selecting $W \rightarrow \tau\nu$ events. The missing E_T trigger leads to $132 \pm 14 \pm 8$ events and the τ trigger to $47 \pm 9 \pm 4$ events. Proper statistical and systematic correlations are taken into account to arrive at $\sigma_B(W \rightarrow \tau\nu) = 2.05 \pm 0.27 \text{ nb}$. Combined with ABE 91C result on $\sigma_B(W \rightarrow e\nu)$, ABE 92E quote a ratio of the couplings from which we derive this measurement.

37 This measurement is derived by us from the ratio of the couplings of ALITTI 92F.

$\Gamma(\pi^+\gamma)/\Gamma(e^+\nu)$				Γ_6/Γ_1
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.5 \times 10^{-3}$	95	ABE	92K CDF	$E_{\text{cm}}^{\text{PD}} = 1.8 \text{ TeV}$
$< 4.9 \times 10^{-3}$	95	38 ALITTI	92D UA2	$E_{\text{cm}}^{\text{PD}} = 630 \text{ GeV}$
$< 58 \times 10^{-3}$	95	39 ALBAJAR	90 UA1	$E_{\text{cm}}^{\text{PD}} = 546,630 \text{ GeV}$

38 ALITTI 92D limit is 3.8×10^{-3} at 90%CL.
39 ALBAJAR 90 obtain < 0.048 at 90%CL.

W REFERENCES

ABE	94B	PRL (to be pub.)	+Albrow, Amidei, Anway-Wiese+	(CDF Collab.)
Fermilab-PUB-94-051-E				
ROSNER	94	PR D49 1363	+Worah, Takeuchi	(EFI, FNAL)
ABE	92E	PRL 68 3398	+Amidei, Apollinari, Atac+	(CDF Collab.)
ABE	92I	PRL 69 28	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	92K	PRL 69 2160	+Amidei, Anway-Weiss+	(CDF Collab.)
ALITTI	92	PL B276 365	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92B	PL B276 354	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92C	PL B277 194	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92D	PL B277 203	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALITTI	92F	PL B280 137	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
LEP	92	PL B276 247	+ALEPH, DELPHI, L3, OPAL	(LEP Collab.)
SAMUEL	92	PL B280 124	+Li, Sinha, Sinha, Sundaresan	(OKSU, CARL)
ABE	91C	PR D44 29	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ALBAJAR	91	PL B253 503	+Albrow, Altkofer, Ankoviak, Apsimon+	(UA1 Collab.)
ALITTI	91C	ZPHY C52 209	+Ambrosini, Ansari, Autiero+	(UA2 Collab.)
SAMUEL	91	PRL 67 9	+Li, Sinha, Sinha, Sundaresan	(OKSU, CARL)
Also	91C	PRL 67 2920 erratum		
ABE	90	PRL 64 152	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
Also	91C	PR D44 29	Abe, Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE	90G	PRL 65 2243	+Amidei, Apollinari, Atac+	(CDF Collab.)
Also	91B	PR D43 2070	Abe, Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ALBAJAR	90	PL B241 283	+Albrow, Altkofer+	(UA1 Collab.)
ALITTI	90B	PL B241 150	+Ansari, Ansoerg, Autiero+	(UA2 Collab.)
ALITTI	90C	ZPHY C47 11	+Ansari, Ansoerg, Bagnaia+	(UA2 Collab.)
ABE	89I	PRL 62 1005	+Amidei, Apollinari, Ascoli, Atac+	(CDF Collab.)
ALBAJAR	89	ZPHY C44 15	+Albrow, Altkofer, Arnison, Astbury+	(UA1 Collab.)
BAUR	88	NP B308 127	+Zeppenfeld	(FSU, WISC)
GRIFOLS	88	IJMP A3 225	+Peris, Sola	(BARC, DESY)
Also	87	PL B197 437	Grifols, Peris, Sola	(BARC, DESY)
ALBAJAR	87	PL B185 233	+Albrow, Altkofer, Arnison, Astbury+	(UA1 Collab.)
ANSARI	87	PL B186 440	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
ANSARI	87C	PL B194 158	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
GROTCH	87	PR D36 2153	+Robinett	(PSU)
HAGIWARA	87	NP B282 253	+Peccei, Zeppenfeld, Hikasa	(KEK, UCLA, FSU)
VANDERBIJ	87	PR D35 1088	van der Bij	(FNAL)
APPEL	86	ZPHY C30 1	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
ARNISON	86	PL B165 484	+Albrow, Altkofer, Astbury+	(UA1 Collab.)
ALTARELLI	85B	ZPHY C27 617	+Ellis, Martinelli	(CERN, FNAL, FRAS)
GRAU	85	PL 154B 283	+Grifols	(BARC)
SUZUKI	85	PL 153B 289		(LBL)
ARNISON	84D	PL 134B 469	+Astbury, Aubert, Bacchi+	(UA1 Collab.)
BAGNAIA	84	ZPHY C24 1	+Banner, Battiston, Blech+	(UA2 Collab.)
HERZOG	84	PL 148B 355		(WISC)
Also	85B	PL 155B 468 erratum	Herzog	(WISC)
ARNISON	83	PL 122B 103	+Astbury, Aubert, Bacchi+	(UA1 Collab.)
ARNISON	83D	PL 129B 273	+Astbury, Aubert, Bacchi+	(UA1 Collab.)
BAGNAIA	83	PL 129B 130	+Banner, Battiston, Bloch+	(UA2 Collab.)
BANNER	83B	PL 122B 476	+Battiston, Bloch, Bonaudi+	(UA2 Collab.)

Z

J = 1

NOTE ON THE Z BOSON

Precision measurements at the Z -boson resonance using electron-positron colliding beams began in 1989 at the SLC and at LEP. During 1989–93, the four CERN experiments have made high-statistics studies of the Z . The availability of longitudinally polarized electron beams at the SLC in 1993 has enabled a precision determination of the effective electroweak mixing angle $\sin^2\theta_W$ that is competitive with the CERN results on this parameter.

The Z -boson properties reported in this section may broadly be categorized as:

- The standard ‘lineshape’ parameters of the Z consisting of its mass, M_Z , its total width, Γ_Z , and its partial decay widths, $\Gamma(\text{hadrons})$ and $\Gamma(\ell\bar{\ell})$ where $\ell = e, \mu, \tau, \nu$;
- The b - and c -quark-related partial widths and charge asymmetries which require special techniques;
- Determination of rare Z decay modes and the search for modes that violate known conservation laws.

Gauge & Higgs Boson Full Listings

Z

Details on Z -parameter determination and the study of $Z \rightarrow b\bar{b}, c\bar{c}$ at LEP are given in this note.

The standard ‘lineshape’ parameters of the Z are determined with increasing precision from an analysis of the production cross sections of these final states in e^+e^- collisions at LEP. The $Z \rightarrow \nu\bar{\nu}(\gamma)$ state is identified directly by detecting single photon production and indirectly by subtracting the visible partial widths from the total width. Inclusion in this analysis of the forward-backward asymmetry of charged leptons, $A_{FB}^{(0,\ell)}$, of the τ polarization, $P(\tau)$, and its forward-backward asymmetry, $P(\tau)^{fb}$, enables the separate determination of the effective vector (\bar{g}_V) and axial vector (\bar{g}_A) couplings of the Z to these leptons and the ratio (\bar{g}_V/\bar{g}_A) which is related to the effective electroweak mixing angle $\sin^2\bar{\theta}_W$ (see the Standard Model review).

Determination of the b - and c -quark-related partial widths and charge asymmetries involves tagging the b and c quarks. Traditionally this was done by requiring the presence of a prompt lepton in the event with high momentum and high transverse momentum (with respect to the accompanying jet). Precision vertex measurement with silicon detectors enables one to do impact parameter and lifetime tagging. Lately, sophisticated neural-network techniques have been used to classify events as b or non- b on a statistical basis using event-shape variables.

Z-parameter determination at LEP

LEP is run at a few energy points on and around the Z mass constituting an energy ‘scan.’ The shape of the cross-section variation around the Z peak can be described by a Breit-Wigner ansatz with an energy-dependent total width [1]. The **three** main properties of this distribution, viz., the **position** of the peak, the **width** of the distribution, and the **height** of the peak, determine respectively the values of M_Z , Γ_Z , and $\Gamma(e^+e^-) \times \Gamma(f\bar{f})$, where $\Gamma(e^+e^-)$ and $\Gamma(f\bar{f})$ are the electron and fermion partial widths of the Z . The quantitative determination of these parameters is done by writing analytic expressions for these cross sections in terms of the parameters and fitting the calculated cross sections to the measured ones by varying these parameters, taking properly into account all the errors. Single-photon exchange and γ - Z interference are included, and the large ($\sim 30\%$) initial-state radiation (ISR) effects are taken into account by convoluting the analytic expressions over a ‘Radiator Function [1,2]’ $H(s, s')$. Thus for the process $e^+e^- \rightarrow f\bar{f}$:

$$\sigma_f(s) = \int H(s, s') \sigma_f^0(s') ds' \quad (1)$$

$$\sigma_f^0(s) = \sigma_Z^0 + \sigma_\gamma^0 + \sigma_{\gamma Z}^0 \quad (2)$$

$$\sigma_Z^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma(e^+e^-)\Gamma(f\bar{f})}{\Gamma_Z^2} \frac{s \Gamma_Z^2}{(s - M_Z^2)^2 + s^2 \Gamma_Z^2/M_Z^2} \quad (3)$$

$$\sigma_\gamma^0 = \frac{4\pi\alpha^2(s)}{3s} Q_f^2 N_c^f \quad (4)$$

$$\sigma_{\gamma Z}^0 = -\frac{2\sqrt{2}\alpha(s)}{3} (Q_f G_F N_c^f g_{V_e} g_{V_f}) \times \frac{(s - M_Z^2) M_Z^2}{(s - M_Z^2)^2 + s^2 \Gamma_Z^2/M_Z^2} \quad (5)$$

where Q_f is the charge of the fermion, $N_c^f = 3(1)$ for quark (lepton) and g_{V_f} is the neutral vector coupling of the Z to the fermion-antifermion pair $f\bar{f}$.

Since $\sigma_{\gamma Z}^0$ is expected to be much less than σ_Z^0 , the LEP collaborations have generally calculated the interference term in the framework of the Standard Model using the best known values of g_V . This fixing of $\sigma_{\gamma Z}^0$ leads to a tighter constraint on M_Z and consequently a smaller error on its fitted value.

Defining

$$A_f = 2 \frac{g_{V_f} \cdot g_{A_f}}{(g_{V_f}^2 + g_{A_f}^2)} \quad (6)$$

where g_{A_f} is the neutral axial-vector coupling of the Z to $f\bar{f}$, the lowest-order expressions for the various lepton-related asymmetries on the Z pole are [3] $A_{FB}^{(0,\ell)} = (3/4)A_e A_f$, $P(\tau) = -A_\tau$, $P(\tau)^{fb} = -(3/4)A_e$, $A_{LR} = A_e$. The full analysis takes into account the energy dependence of the asymmetries. Experimentally A_{LR} is defined as $(\sigma_L - \sigma_R)/(\sigma_L + \sigma_R)$ where $\sigma_{L(R)}$ are the $e^+e^- \rightarrow Z$ production cross sections with left- (right)-handed electrons.

In terms of g_A and g_V , the partial decay width of the Z to $f\bar{f}$ can be written as

$$\Gamma(f\bar{f}) = \frac{G_F M_Z^3}{6\sqrt{2}\pi} (g_{V_f}^2 + g_{A_f}^2) N_c^f (1 + \delta_{\text{QED}})(1 + \delta_{\text{QCD}}) \quad (7)$$

where $\delta_{\text{QED}} = 3\alpha Q_f^2/4\pi$ accounts for final-state photonic corrections and $\delta_{\text{QCD}} = 0$ for leptons and $\delta_{\text{QCD}} = (\alpha_s/\pi) + 1.409(\alpha_s/\pi)^2 - 12.805(\alpha_s/\pi)^3$ for quarks, α_s being the strong coupling constant at $\mu = M_Z$.

In the above framework, the QED radiative corrections have been explicitly taken into account by convoluting over the ISR and allowing the electromagnetic coupling constant to run [4]: $\alpha(s) = \alpha/(1 - \Delta\alpha)$. On the other hand, weak radiative corrections that depend upon the assumptions of the electroweak theory and on the values of the unknown M_{top} and M_{Higgs} are accounted for by **absorbing them into the couplings**, which are then called the *effective* couplings \bar{g}_V and \bar{g}_A (or alternatively the effective parameters of the \star scheme of Kennedy and Lynn [5]).

S-matrix approach to the Z

While practically all experimental analyses of LEP/SLC data have followed the ‘Breit-Wigner’ approach described above, an alternative S-matrix-based analysis is also possible. The Z , like all unstable particles, is associated with a complex pole in the S matrix. The pole position is process independent and gauge invariant. The mass, \bar{M}_Z , and width, $\bar{\Gamma}_Z$, can be defined in terms of the pole in the energy plane via [6]

$$\bar{s} = \bar{M}_Z^2 - i\bar{M}_Z\bar{\Gamma}_Z \quad (8)$$

leading to the relations

$$\overline{M}_Z = M_Z / \sqrt{1 + \Gamma_Z^2 / M_Z^2} \quad (9)$$

$$\approx M_Z - 34 \text{ MeV} \quad (10)$$

$$\overline{\Gamma}_Z = \Gamma_Z / \sqrt{1 + \Gamma_Z^2 / M_Z^2} \quad (11)$$

$$\approx \Gamma_Z - 0.9 \text{ MeV} \quad (12)$$

Some authors [7] choose to define the Z mass and width via

$$\overline{s} = (\overline{M}_Z - \frac{i}{2}\overline{\Gamma}_Z)^2 \quad (13)$$

which yields $\overline{M}_Z \approx M_Z - 26 \text{ MeV}$, $\overline{\Gamma}_Z \approx \Gamma_Z - 1.2 \text{ MeV}$.

The L3 collaboration at LEP (ADRIANI 93H, ADRIANI 93M) have analyzed their data using the S-matrix approach as defined in Eq. (8), in addition to the conventional one. As expected, they observe a downward shift of $\approx 34 \text{ MeV}$ in the Z mass.

Handling the large-angle e^+e^- final state

Unlike other $f\bar{f}$ decay final states of the Z , the e^+e^- final state has a contribution not only from the s -channel but also from the t -channel and s - t interference. The full amplitude is not amenable to fast calculation, which is essential if one has to carry out minimization fits within reasonable computer time. The usual procedure is to calculate the non- s channel part of the cross section separately using the Standard Model program ALIBABA [8] using the ‘most reasonable’ values of M_{top} , and M_{Higgs} and add it to the s -channel cross section calculated as for other channels. This leads to two additional sources of error in the analysis: firstly, the theoretical calculation in ALIBABA itself is known to be accurate to $\sim 0.5 \%$, and secondly, there is uncertainty due to unknown M_{top} and M_{Higgs} . These additional errors are propagated into the analysis by including them in the systematic error on the e^+e^- final state.

Errors due to uncertainty in LEP energy determination [9]

The systematic errors related to the LEP energy measurement can be classified as:

- The absolute energy scale error;
- Energy-point-to-energy-point errors due to the non-linear response of the magnets to the exciting currents;
- Energy-point-to-energy-point errors due to possible higher-order effects in the relationship between the dipole field and beam energy;
- Energy reproducibility errors due to various unknown uncertainties in temperatures, tidal effects, corrector settings, RF status *etc.* Since one groups together data taken at ‘nominally same’ energies in different fills, it can be assumed that these errors are uncorrelated and are reduced by $\sqrt{N_{\text{fill}}}$ where N_{fill} is the (luminosity weighted) effective number of fills at a particular energy point.

At each energy point the last two errors can be summed into one point-to-point error.

Choice of fit parameters

The LEP collaborations have chosen the following primary set of parameters for fitting: M_Z , Γ_Z , σ_{hadron}^0 , $R(\text{lepton})$, $A_{FB}^{(0,\ell)}$, where $R(\text{lepton}) = \Gamma(\text{hadrons})/\Gamma(\text{lepton})$, $\sigma_{\text{hadron}}^0 = 12\pi\Gamma(e^+e^-)\Gamma(\text{hadrons})/M_Z^2\Gamma_Z^2$. With a knowledge of these fitted parameters and their covariance matrix, any other parameter can be derived. The main advantage of these parameters is that they form the **least correlated** set of parameters, so that it becomes easy to combine results from the different LEP experiments.

Thus, the most general fit carried out to cross section and asymmetry data determines the **nine parameters**: M_Z , Γ_Z , σ_{hadron}^0 , $R(e)$, $R(\mu)$, $R(\tau)$, $A_{FB}^{(0,e)}$, $A_{FB}^{(0,\mu)}$, $A_{FB}^{(0,\tau)}$. Assumption of lepton universality leads to a **five-parameter fit** determining M_Z , Γ_Z , σ_{hadron}^0 , $R(\text{lepton})$, $A_{FB}^{(0,\ell)}$. The use of **only** cross-section data leads to six- or four-parameter fits if lepton universality is or is not assumed, *i.e.*, $A_{FB}^{(0,\ell)}$ values are not determined.

In order to determine the best values of the effective vector and axial vector couplings of the charged leptons to the Z , the above mentioned nine- and five-parameter fits are carried out with added constraints from the measured values of A_τ and A_e obtained from τ polarization studies at LEP and the determination of A_{LR} at SLC.

Combining results from the four LEP experiments [10]

Each LEP experiment provides the values of the parameters mentioned above together with the full covariance matrix. The statistical and experimental systematic errors are assumed to be uncorrelated among the four experiments. The sources of **common** systematic errors are i) the LEP energy uncertainties, and ii) the effect of theoretical uncertainty in calculating the small-angle Bhabha cross section for luminosity determination and in estimating the non- s channel contribution to the large-angle Bhabha cross section. Using this information, a full covariance matrix, V , of all the input parameters is constructed and a combined parameter set is obtained by minimizing $\chi^2 = \Delta^T V^{-1} \Delta$, where Δ is the vector of residuals of the combined parameter set to the results of individual experiments.

Study of $Z \rightarrow b\bar{b}, c\bar{c}$ at LEP

These studies lead to the experimental determination of the ratios of the partial widths $\Gamma(b\bar{b})/\Gamma(\text{hadrons})$, $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$, and the forward-backward (charge) asymmetries $A_{FB}^{(0,b)}$ and $A_{FB}^{(0,c)}$. Each LEP experiment has used more than one technique of b, c tagging, and then used certain assumptions, some model dependent, to extract the final numbers. This makes the **task of combining** these LEP results quite daunting. A first step in this direction was made recently [10] for a joint LEP presentation for the 1993 summer conferences. In principle, the task is the same as for combining results on the lineshape: identification of the **common systematic errors**, which in this case exist not only among different LEP experiments, but

Gauge & Higgs Boson Full Listings

Z

also within one collaboration if different (partially correlated) methods are employed to extract the same quantity.

Extraction of $\Gamma(b\bar{b})/\Gamma(\text{hadrons})$

Three methods have been used to tag a b quark: lepton tagging, analysis of event shape, and lifetime tagging. The latter includes impact-parameter or decay-length double tags, or in combination with the lepton or event-shape tags. The various sources of common systematic errors are:

- Semileptonic decay model;
- Semileptonic branching ratios of b and c quarks;
- $c\bar{c}$ contamination, $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$;
- Variation of Monte Carlo parameters, in particular the b -fragmentation function;
- Hemisphere correlations.

Some other possible common errors due to charmed hadron composition, b, c production in uds events, *etc.* are expected to be small.

Extraction of $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$

In addition to the methods of lepton tagging, event-shape analysis, and lifetime tags, $D^*(2010)$ tagging is also used to identify c quarks. The common systematic errors are basically the same as for the b -tagging case, where instead of $c\bar{c}$ contamination, it is $b\bar{b}$ contamination.

Extraction of $A_{FB}^{(0,b)}$ and $A_{FB}^{(0,c)}$

For determining the asymmetry, it is essential to identify the charge of the quark in addition to tagging an event as due to $b\bar{b}$ or $c\bar{c}$. Thus, asymmetry measurements have utilized a lepton tag and a lifetime tag in combination with a jet-charge measurement and a $D^*(2010)$ tag. The values of $A_{FB}^{(0,b)}$ are corrected for the effect of $B^0\bar{B}^0$ mixing. The common errors in $A_{FB}^{(0,b)}$ are due to:

- Semileptonic decay model and branching ratios;
- Fragmentation;
- $\Gamma(b\bar{b})$ and $\Gamma(c\bar{c})$;
- $B^0\bar{B}^0$ mixing;
- $A_{FB}^{(0,c)}$.

For $A_{FB}^{(0,c)}$ the first four common errors are the same as for $A_{FB}^{(0,b)}$. The fifth error is due to $A_{FB}^{(0,b)}$; the sixth is due to the uncertainty in the probability of producing a $D^*(2010)$ in $c\bar{c}$ and $b\bar{b}$ events and in the $D^*(2010)$ branching ratio.

References

- R.N. Cahn, Phys. Rev. **D36**, 2666 (1987); F.A. Berends *et al.*, "Z Physics at LEP 1", CERN Report 89-08 (1989), Vol. 1, eds. G. Altarelli, R. Kleiss, and C. Verzegnassi, p. 89; A. Borrelli *et al.*, Nucl. Phys. **B333**, 357 (1990).
- D. Bardin *et al.*, Nucl. Phys. **B351**, 1 (1991).
- M. Consoli *et al.*, "Z Physics at LEP 1", CERN Report 89-08 (1989), Vol. 1, eds. G. Altarelli, R. Kleiss, and C. Verzegnassi, p. 7; M. Bohm *et al.*, *ibid.*, p. 203; S. Jadach *et al.*, *ibid.*, p. 235.
- G. Burgers *et al.*, *ibid.*, p. 55.
- D.C. Kennedy and B.W. Lynn, SLAC-PUB 4039 (1986, revised 1988).
- R. Stuart, Phys. Lett. **B262**, 113 (1991); A. Sirlin, Phys. Rev. Lett. **67**, 2127 (1991); A. Leike, T. Riemann, and J. Rose, Phys. Lett. **B273**, 513 (1991); See also D. Bardin *et al.*, Phys. Lett. **B206**, 539 (1988).
- S. Willenbrock and G. Valencia, Phys. Lett. **B259**, 373 (1991).
- W. Beenakker, F.A. Berends, and S.C. van der Marck, Nucl. Phys. **B349**, 323 (1991).
- L. Arnaudon *et al.* (Working Group on LEP Energy and LEP Collaborations), Phys. Lett. **B307**, 187 (1993); L. Arnaudon *et al.* (Working Group on LEP Energy), CERN-PPE/92-1254 (1992); L. Arnaudon *et al.*, Phys. Lett. **B284**, 431 (1992); R. Baily *et al.*, 'LEP Energy Calibration' CERN-SL 90-95.
- The LEP Collaborations ALEPH, DELPHI, L3, OPAL, and LEP Electroweak Working Group, CERN-PPE/93-157, (26 Aug 1993).

Z MASS

The fit is performed using the Z mass and width, the Z hadronic pole cross section, the ratios of hadronic to leptonic partial widths, and the Z pole forward-backward lepton asymmetries. We believe that this set is the most free of correlations. Common systematic errors are taken into account. For more details, see the 'Note on the Z Boson.'

The Z-boson mass listed here corresponds to a Breit-Wigner resonance parameter. The value is 34 MeV greater than the real part of the position of the pole (in the energy plane) in the Z-boson propagator. Also the LEP experiments have generally assumed a fixed value of the $\gamma - Z$ interferences term based on the standard model. Keeping this term as free parameter leads to a somewhat larger error on the fitted Z mass. See ADRIANI 93H for a detailed investigation of both these issues.

VALUE(GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
91.187±0.007 OUR FIT				
91.188±0.007 OUR AVERAGE				
91.187±0.007±0.006	1.15M	¹ ABREU	94 DLPH	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.195±0.006±0.007	1.2M	¹ ACCIARRI	94 L3	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.182±0.007±0.006	1.33M	¹ AKERS	94 OPAL	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.187±0.007±0.006	1.28M	¹ BUSKULIC	94 ALEP	$E_{cm}^{e\bar{e}} = 88-94$ GeV
90.9 ± 0.3 ± 0.2	188	² ABE	89C CDF	$E_{cm}^{p\bar{p}} = 1800$ GeV
91.14 ± 0.12	480	³ ABRAMS	89B MRK2	$E_{cm}^{e\bar{e}} = 89-93$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
91.181±0.007±0.006	512k	⁴ ACTON	93D OPAL	Repl. by AKERS 94
91.195±0.009	460k	⁵ ADRIANI	93F L3	Repl. by ACCIARRI 94
91.160±0.010	463k	⁶ ADRIANI	93H L3	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.187±0.009	520k	⁷ BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
91.187±0.007	2.2M	⁸ LEP	93 RVUE	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.187±0.007	1.9M	⁹ QUAST	93 RVUE	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.74 ± 0.28 ± 0.93	156	¹⁰ ALITTI	92B UA2	$E_{cm}^{p\bar{p}} = 630$ GeV
91.177±0.006±0.02	550k	^{11,12} BANERJEE	92 RVUE	$E_{cm}^{e\bar{e}} = 88-94$ GeV
91.182±0.009±0.02	190k	¹² DECAMP	92B ALEP	Repl. by LEP 93
91.175±0.021	650k	¹³ LEP	92 RVUE	Repl. by LEP 93
91.177±0.010±0.02	150k	¹² ABREU	91F DLPH	Repl. by ABREU 94
91.181±0.010±0.02	125k	¹² ADEVA	91E L3	Repl. by LEP 93
91.161±0.009±0.02	184k	¹² ALEXANDER	91F OPAL	Repl. by LEP 93
93.1 ± 1.0 ± 3.0	24	^{14,15} ALBAJAR	89 UA1	$E_{cm}^{p\bar{p}} = 546,630$ GeV

¹ The second error of 6.3 MeV is due to a common LEP energy uncertainty.

² First error of ABE 89 is combination of statistical and systematic contributions; second is mass scale uncertainty.

³ ABRAMS 89B uncertainty includes 35 MeV due to the absolute energy measurement.

⁴ The systematic error in ACTON 93D is from the uncertainty in the LEP energy calibration.

⁵ The error in ADEVA 91E includes 6 MeV due to the uncertainty in LEP energy calibration.

⁶ ADRIANI 93H use the S-matrix approach to determine the pole position for the Z boson. Note the shift of this result with respect to the standard Breit-Wigner parameterization.

⁷ BUSKULIC 93J supersedes DECAMP 92B. The error includes 6 MeV due to the uncertainty in LEP energy calibration.

⁸ The LEP 93 error is due to the experiments is 4 MeV and the uncertainty due to the absolute LEP energy scale is 6 MeV.

⁹ QUAST 93 is a combined analysis of LEP results as of Feb. 1993. A common systematic error of 6 MeV is taken into account.

- ¹⁰ Enters fit through W/Z mass ratio below. The ALITTI 92B systematic error (± 0.93) has two contributions: one (± 0.92) cancels in m_W/m_Z and one (± 0.12) is noncancelling. These were added in quadrature.
- ¹¹ BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.
- ¹² The systematic error (0.02) is an error in common to the 4 LEP experiments.
- ¹³ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.
- ¹⁴ Enters fit through $Z \rightarrow W$ mass difference below.
- ¹⁵ ALBAJAR 89 result is from a total sample of 33 $Z \rightarrow e^+e^-$ events.

Z WIDTH

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.490 ± 0.007 OUR FIT				
2.491 ± 0.007 OUR AVERAGE				
2.483 ± 0.011 ± 0.0045	1.15M	16 ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
2.494 ± 0.009 ± 0.0045	1.2M	16 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
2.483 ± 0.011 ± 0.0045	1.33M	16 AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
2.501 ± 0.011 ± 0.0045	1.28M	16 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
3.8 ± 0.8 ± 1.0	188	ABE	89C CDF	$E_{cm}^{pp} = 1800$ GeV
2.42 +0.45 -0.35	480	17 ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV
2.7 +1.2 -1.0 ± 1.3	24	18 ALBAJAR	89 UA1	$E_{cm}^{pp} = 546,630$ GeV
2.7 ± 2.0 ± 1.0	25	19 ANSARI	87 UA2	$E_{cm}^{pp} = 546,630$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.483 ± 0.011 ± 0.004	512k	20 ACTON	93D OPAL	Repl. by AKERS 94
2.490 ± 0.011	460k	21 ADRIANI	93F L3	Repl. by ACCIARRI 94
2.492 ± 0.012	463k	22 ADRIANI	93H L3	$E_{cm}^{ee} = 88-94$ GeV
2.501 ± 0.012	520k	23 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
2.490 ± 0.007	1.9M	24 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
2.481 ± 0.010	550k	25 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
2.484 ± 0.017 ± 0.005	190k	26 DECAMP	92B ALEP	Repl. by BUSKULIC 93J
2.487 ± 0.010	650k	27 LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
2.465 ± 0.019 ± 0.005	150k	26 ABREU	91F DLPH	Repl. by ABREU 94
2.501 ± 0.017 ± 0.005	125k	26 ADEVA	91E L3	Repl. by ADRIANI 93F
2.492 ± 0.015 ± 0.005	184k	26 ALEXANDER	91F OPAL	Repl. by ACTON 93D

- ¹⁶ The second error of 4.5 MeV is due to a common LEP energy uncertainty.
- ¹⁷ ABRAMS 89B uncertainty includes 50 MeV due to the minISAM background subtraction error.
- ¹⁸ ALBAJAR 89 result is from a total sample of 33 $Z \rightarrow e^+e^-$ events.
- ¹⁹ Quoted values of ANSARI 87 are from direct fit. Ratio of Z and W production gives either $\Gamma(Z) < (1.09 \pm 0.07) \times \Gamma(W)$, CL = 90% or $\Gamma(Z) = (0.82 \pm 0.19 \pm 0.06) \times \Gamma(W)$. Assuming Standard-Model value $\Gamma(W) = 2.65$ GeV then gives $\Gamma(Z) < 2.89 \pm 0.19$ or $= 2.17 \pm 0.50 \pm 0.37$.
- ²⁰ The systematic error is from the uncertainty in the LEP energy calibration.
- ²¹ The error in ADRIANI 93F includes 4 MeV due to the uncertainty in LEP energy calibration.
- ²² ADRIANI 93H use the S-matrix approach to determine the pole position for the Z boson.
- ²³ The error in BUSKULIC 93J includes 4 MeV due to the uncertainty in LEP energy calibration.
- ²⁴ QUAST 93 is a combined analysis of LEP results as of Feb. 1993. A common systematic error of 4 MeV is taken into account.
- ²⁵ BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.
- ²⁶ The systematic error (0.005) is an error in common to the 4 LEP experiments.
- ²⁷ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.

Z DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 e^+e^-$	(3.366 ± 0.008) %	
$\Gamma_2 \mu^+\mu^-$	(3.367 ± 0.013) %	
$\Gamma_3 \tau^+\tau^-$	(3.360 ± 0.015) %	
$\Gamma_4 \ell^+\ell^-$	[a] (3.367 ± 0.006) %	
Γ_5 invisible	(20.01 ± 0.16) %	
Γ_6 hadrons	(69.90 ± 0.15) %	
$\Gamma_7 (u\bar{u} + c\bar{c})/2$	(9.7 ± 1.8) %	
$\Gamma_8 (d\bar{d} + s\bar{s} + b\bar{b})/3$	(16.8 ± 1.2) %	
$\Gamma_9 c\bar{c}$	(11.9 ± 1.4) %	
$\Gamma_{10} b\bar{b}$	(15.45 ± 0.21) %	
$\Gamma_{11} \pi^0\gamma$	< 5.5 × 10 ⁻⁵	95%
$\Gamma_{12} \eta\gamma$	< 5.1 × 10 ⁻⁵	95%
$\Gamma_{13} \omega\gamma$	< 6.5 × 10 ⁻⁴	95%
$\Gamma_{14} \eta'(958)\gamma$	< 4.2 × 10 ⁻⁵	95%
$\Gamma_{15} \gamma\gamma$	< 5.5 × 10 ⁻⁵	95%
$\Gamma_{16} \gamma\gamma\gamma$	< 1.7 × 10 ⁻⁵	95%
$\Gamma_{17} \pi^\pm W^\mp$	[b] < 7 × 10 ⁻⁵	95%
$\Gamma_{18} \rho^\pm W^\mp$	[b] < 8.3 × 10 ⁻⁵	95%
$\Gamma_{19} K^0 X$	(61.5 ± 0.6) %	
$\Gamma_{20} K^*(892)^+ X$	(51 ± 5) %	

$\Gamma_{21} AX$	(20.9 ± 0.6) %	
$\Gamma_{22} \Xi^- X$	(1.42 ± 0.14) %	
$\Gamma_{23} \Sigma(1385)^+ X$	(2.6 ± 0.4) %	
$\Gamma_{24} \Xi(1530)^0 X$	(4.4 ± 1.0) × 10 ⁻³	
$\Gamma_{25} \Omega^- X$	(3.5 ± 1.0) × 10 ⁻³	
$\Gamma_{26} J/\psi(1S) X$	(3.8 ± 0.5) × 10 ⁻³	
$\Gamma_{27} \chi_{c1}(1P) X$	(7.5 ± 3.0) × 10 ⁻³	
$\Gamma_{28} (D^0/\bar{D}^0) X$	(28 ± 4) %	
$\Gamma_{29} D^\pm X$	(13.9 ± 2.1) %	
$\Gamma_{30} D^*(2010)^\pm X$	[b] (12.5 ± 1.3) %	
$\Gamma_{31} B_s^0 X$	seen	
Γ_{32} anomalous γ + hadrons	[c] < 3.2 × 10 ⁻³	95%
$\Gamma_{33} e^+e^- \gamma$	[c] < 5.2 × 10 ⁻⁴	95%
$\Gamma_{34} \mu^+\mu^- \gamma$	[c] < 5.6 × 10 ⁻⁴	95%
$\Gamma_{35} \tau^+\tau^- \gamma$	[c] < 7.3 × 10 ⁻⁴	95%
$\Gamma_{36} \ell^+\ell^- \gamma\gamma$	[d] < 6.8 × 10 ⁻⁶	95%
$\Gamma_{37} q\bar{q}\gamma\gamma$	[d] < 5.5 × 10 ⁻⁶	95%
$\Gamma_{38} \nu\bar{\nu}\gamma\gamma$	[d] < 3.1 × 10 ⁻⁶	95%
$\Gamma_{39} e^\pm \mu^\mp$	LF [b] < 6 × 10 ⁻⁶	95%
$\Gamma_{40} e^\pm \tau^\mp$	LF [b] < 1.3 × 10 ⁻⁵	95%
$\Gamma_{41} \mu^\pm \tau^\mp$	LF [b] < 1.9 × 10 ⁻⁵	95%

[a] ℓ indicates each type of lepton (e , μ , and τ), not sum over them.

[b] The value is for the sum of the charge states indicated.

[c] See the Full Listings below for the γ energy range used in this measurement.

[d] For $m_{\gamma\gamma} = (60 \pm 5)$ GeV.

Z PARTIAL WIDTHS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83.83 ± 0.30 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
83.31 ± 0.54	31.4k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.43 ± 0.52	38k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
83.63 ± 0.53	42k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.61 ± 0.49	45.8k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
83.03 ± 0.66	17k	ACTON	93D OPAL	Repl. by AKERS 94
83.0 ± 0.6	16k	ADRIANI	93M L3	Repl. by ACCIARRI 94
84.43 ± 0.60		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
83.30 ± 0.35	70k	28 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
82.6 ± 0.7	16k	29 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
83.8 ± 0.9	6947	DECAMP	92B ALEP	Repl. by BUSKULIC 93J
83.20 ± 0.55	19k	30 LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
82.4 ± 1.1 ± 0.5	2772	ABREU	91F DLPH	Repl. by ABREU 94
83.3 ± 1.1	4175	ADEVA	91E L3	Repl. by ADRIANI 93M
82.9 ± 1.0	5507	ALEXANDER	91F OPAL	Repl. by ACTON 93D
28 QUAST 93 is a combined analysis of LEP results as of Feb. 1993.				
29 BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.				
30 LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.				

$\Gamma(\mu^+\mu^-)$
This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83.84 ± 0.39 OUR FIT				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
84.15 ± 0.77	45.6k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.20 ± 0.79	34k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
83.83 ± 0.65	57k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
83.62 ± 0.75	46.4k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
84.43 ± 0.92	23k	ACTON	93D OPAL	Repl. by AKERS 94
82.8 ± 1.0	14k	ADRIANI	93M L3	Repl. by ACCIARRI 94
83.66 ± 0.95		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
83.82 ± 0.52	70k	31 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
83.7 ± 1.1	16k	32 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
81.4 ± 1.4	6691	DECAMP	92B ALEP	Repl. by BUSKULIC 93J
83.35 ± 0.86	21k	33 LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
86.9 ± 1.9 ± 0.9	3428	ABREU	91F DLPH	Repl. by ABREU 94
84.5 ± 2.0	3245	ADEVA	91E L3	Repl. by ADRIANI 93M
83.2 ± 1.5	7240	ALEXANDER	91F OPAL	Repl. by ACTON 93D
31 QUAST 93 is a combined analysis of LEP results as of Feb. 1993.				
32 BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.				
33 LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.				

Gauge & Higgs Boson Full Listings

Z

 $\Gamma(\tau^+\tau^-)$ Γ_3

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83.68 ± 0.44 OUR FIT				
83.55 ± 0.91	25k	ABREU 94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
84.04 ± 0.94	25k	ACCIARRI 94	L3	$E_{cm}^{ee} = 88-94$ GeV
82.90 ± 0.77	47k	AKERS 94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.18 ± 0.79	45.1k	BUSKULIC 94	ALEP	$E_{cm}^{ee} = 88-94$ GeV
82.2 ± 1.1	18k	ACTON 93D	OPAL	Repl. by AKERS 94
84.6 ± 1.2	10k	ADRIANI 93M	L3	Repl. by ACCIARRI 94
84.09 ± 1.10		BUSKULIC 93J	ALEP	Repl. by BUSKULIC 94
83.54 ± 0.62	50k	³⁴ QUAST 93	RVUE	$E_{cm}^{ee} = 88-94$ GeV
83.1 ± 1.2	16k	³⁵ BANERJEE 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
82.4 ± 1.6	6260	DECAMP 92B	ALEP	Repl. by BUSKULIC 93J
82.76 ± 1.02	17k	³⁶ LEP 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
82.7 ± 2.1 ± 1.1	2345	ABREU 91F	DLPH	Repl. by ABREU 94
84.0 ± 2.7	2540	ADEVA 91E	L3	Repl. by ADRIANI 93M
82.7 ± 1.9	5559	ALEXANDER 91F	OPAL	Repl. by ACTON 93D

³⁴QUAST 93 is a combined analysis of LEP results as of Feb. 1993.

³⁵BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.

³⁶LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.

 $\Gamma(\ell^+\ell^-)$ Γ_4

In our fit $\Gamma(\ell^+\ell^-)$ is defined as the partial Z width for the decay into a pair of massless charged leptons. This parameter is not directly used in the 5-parameter fit assuming lepton universality but is derived using the fit results. See the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
83.84 ± 0.27 OUR FIT				
83.56 ± 0.45	102k	ABREU 94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
83.49 ± 0.46	97k	ACCIARRI 94	L3	$E_{cm}^{ee} = 88-94$ GeV
83.55 ± 0.44	146k	AKERS 94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
84.40 ± 0.43	137.3k	BUSKULIC 94	ALEP	$E_{cm}^{ee} = 88-94$ GeV
83.27 ± 0.50	58k	ACTON 93D	OPAL	Repl. by AKERS 94
83.1 ± 0.5	40k	ADRIANI 93F	L3	Repl. by ACCIARRI 94
84.22 ± 0.48		BUSKULIC 93J	ALEP	Repl. by BUSKULIC 94
83.40 ± 0.29	190k	³⁷ QUAST 93	RVUE	$E_{cm}^{ee} = 88-94$ GeV
83.0 ± 0.4	50k	³⁸ BANERJEE 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
83.1 ± 0.7	20k	DECAMP 92B	ALEP	Repl. by BUSKULIC 93J
83.24 ± 0.42	57k	³⁹ LEP 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
83.4 ± 0.8	10k	ABREU 91F	DLPH	Repl. by ABREU 94
83.6 ± 0.8	10k	ADEVA 91E	L3	Repl. by ADRIANI 93F
83.0 ± 0.7	18k	ALEXANDER 91F	OPAL	Repl. by ACTON 93D

³⁷QUAST 93 is a combined analysis of LEP results as of Feb. 1993.

³⁸BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.

³⁹LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.

 $\Gamma(\text{invisible})$ Γ_5

We use only direct measurements of the invisible partial width to obtain the average value quoted below. The fit value is obtained as a difference between the total and the observed partial widths assuming lepton universality.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
496.2 ± 4.2 OUR FIT				
496 ± 29 OUR AVERAGE				
450 ± 34 ± 34	258	BUSKULIC 93L	ALEP	$E_{cm}^{ee} = 88-94$ GeV
540 ± 80 ± 40	61	ADEVA 92	L3	$E_{cm}^{ee} = 88-94$ GeV
524 ± 40 ± 20	172	⁴⁰ ADRIANI 92E	L3	$E_{cm}^{ee} = 89-94$ GeV
500 ± 70 ± 30	73	AKRAWY 91D	OPAL	$E_{cm}^{ee} = 88-94$ GeV
509.4 ± 7.0		ABREU 94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
496.5 ± 7.9		ACCIARRI 94	L3	$E_{cm}^{ee} = 88-94$ GeV
490.3 ± 7.3		AKERS 94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
501 ± 6		BUSKULIC 94	ALEP	$E_{cm}^{ee} = 88-94$ GeV
495 ± 10		ACTON 93D	OPAL	Repl. by AKERS 94
494 ± 10		ADRIANI 93M	L3	Repl. by ACCIARRI 94
498 ± 9		BUSKULIC 93J	ALEP	Repl. by BUSKULIC 94
499 ± 6		⁴¹ QUAST 93	RVUE	$E_{cm}^{ee} = 88-94$ GeV
491 ± 13		DECAMP 92B	ALEP	Repl. by BUSKULIC 93J
498 ± 8		⁴² LEP 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
488 ± 17		ABREU 91F	DLPH	Repl. by ABREU 94
508 ± 17		ADEVA 91E	L3	Repl. by ADRIANI 93M
504 ± 15		ALEXANDER 91F	OPAL	Repl. by ACTON 93D

⁴⁰ADRIANI 92E improves but does not supersede ADEVA 92, obtained with 1990 data only.

⁴¹QUAST 93 is a combined analysis of LEP results as of Feb. 1993. Assumes lepton universality.

⁴²LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included. Assumes lepton universality.

 $\Gamma(\text{hadrons})$ Γ_6

This parameter is not directly used in the 5-parameter fit assuming lepton universality, but is derived using the fit results. See the 'Note on the Z Boson.'

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1740.7 ± 5.9 OUR FIT				
1723 ± 10	1.05M	ABREU 94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
1748 ± 10	1.1M	ACCIARRI 94	L3	$E_{cm}^{ee} = 88-94$ GeV
1741 ± 10	1.2M	⁴³ AKERS 94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
1746 ± 10	1.14M	BUSKULIC 94	ALEP	$E_{cm}^{ee} = 88-94$ GeV
1738 ± 12	454k	⁴⁴ ACTON 93D	OPAL	Repl. by AKERS 94
1747 ± 11	420k	ADRIANI 93F	L3	Repl. by ACCIARRI 94
1751 ± 11		BUSKULIC 93J	ALEP	Repl. by BUSKULIC 94
1741 ± 7	1.7M	⁴⁵ QUAST 93	RVUE	$E_{cm}^{ee} = 88-94$ GeV
1734 ± 10	500k	^{46,47} BANERJEE 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
1740 ± 12	570k	⁴⁸ LEP 92	RVUE	Repl. by LEP 93
1726 ± 19	124k	ABREU 91F	DLPH	Repl. by ABREU 94
⁴³ AKERS 94 assumes lepton universality. Without this assumption, it becomes 1742 ± 11 MeV.				
⁴⁴ ACTON 93D assumes lepton universality. Without this assumption it becomes 1743 ± 15 MeV.				
⁴⁵ QUAST 93 is a combined analysis of LEP results as of Feb. 1993. Assumes lepton universality.				
⁴⁶ Assuming lepton universality. Without this assumption it becomes 1741 ± 15 MeV.				
⁴⁷ BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.				
⁴⁸ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included. Assumes lepton universality.				

Z BRANCHING RATIOS

 $\Gamma(\text{hadrons})/\Gamma(e^+e^-)$ Γ_6/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
20.76 ± 0.08 OUR FIT				
20.74 ± 0.18	31.4k	ABREU 94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
20.96 ± 0.15	38k	ACCIARRI 94	L3	$E_{cm}^{ee} = 88-94$ GeV
20.83 ± 0.16	42k	AKERS 94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
20.59 ± 0.15	45.8k	BUSKULIC 94	ALEP	$E_{cm}^{ee} = 88-94$ GeV
20.7 ± 11.7 - 8.8	12	⁴⁹ ABRAMS 89D	MRK2	$E_{cm}^{ee} = 89-93$ GeV
20.99 ± 0.25	17k	ACTON 93D	OPAL	Repl. by AKERS 94
20.69 ± 0.21		BUSKULIC 93J	ALEP	Repl. by BUSKULIC 94
20.92 ± 0.12	70k	⁵⁰ QUAST 93	RVUE	$E_{cm}^{ee} = 88-94$ GeV
20.91 ± 0.22	19k	⁵¹ LEP 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
21.19 ± 0.49 ± 0.18	2772	⁵² ABREU 91F	DLPH	Repl. by ABREU 94
21.01 ± 0.40 ± 0.22	4175	⁵³ ADEVA 91E	L3	Repl. by ACCIARRI 94

⁴⁹ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

⁵⁰QUAST 93 is a combined analysis of LEP results as of Feb. 1993.

⁵¹LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.

⁵²ABREU 91F report $\Gamma(ee) = 82.4 \pm 1.1 \pm 0.5$ MeV and provided us with this branching ratio from the same data and analysis.

⁵³ADEVA 91E report $B(ee) = 3.33 \pm 0.04\%$ and $\Gamma(ee) = 83.3 \pm 1.1$ MeV and provided us with this branching ratio from the same data and analysis.

 $\Gamma(\text{hadrons})/\Gamma(\mu^+\mu^-)$ Γ_6/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
20.76 ± 0.07 OUR FIT				
20.78 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.3. See the Ideogram below.
20.54 ± 0.14	45.6k	ABREU 94	DLPH	$E_{cm}^{ee} = 88-94$ GeV
21.02 ± 0.16	34k	ACCIARRI 94	L3	$E_{cm}^{ee} = 88-94$ GeV
20.78 ± 0.11	57k	AKERS 94	OPAL	$E_{cm}^{ee} = 88-94$ GeV
20.83 ± 0.15	46.4k	BUSKULIC 94	ALEP	$E_{cm}^{ee} = 88-94$ GeV
18.9 ± 7.1 - 5.3	13	⁵⁴ ABRAMS 89D	MRK2	$E_{cm}^{ee} = 89-93$ GeV
20.65 ± 0.17	23k	ACTON 93D	OPAL	Repl. by AKERS 94
20.88 ± 0.20		BUSKULIC 93J	ALEP	Repl. by BUSKULIC 94
20.79 ± 0.10	70k	⁵⁵ QUAST 93	RVUE	$E_{cm}^{ee} = 88-94$ GeV
21.26 ± 0.29	6691	DECAMP 92B	ALEP	Repl. by BUSKULIC 93J
20.88 ± 0.18	21k	⁵⁶ LEP 92	RVUE	$E_{cm}^{ee} = 88-94$ GeV
19.89 ± 0.40 ± 0.19	2475	⁵⁷ ABREU 91F	DLPH	Repl. by ABREU 94
20.08 ± 0.36 ± 0.16	3428	⁵⁸ ABREU 91E	DLPH	Repl. by ABREU 94
20.75 ± 0.39 ± 0.17	3245	⁵⁹ ADEVA 91E	L3	Repl. by ACCIARRI 94
20.92 ± 0.31	7240	⁶⁰ ALEXANDER 91F	OPAL	Repl. by ACTON 93D

⁵⁴ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.

⁵⁵QUAST 93 is a combined analysis of LEP results as of Feb. 1993.

⁵⁶LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.


⁵⁷ABREU 91 also obtain $[\Gamma(e)\Gamma(\mu)]^{1/2} = 85.0 \pm 0.9 \pm 0.8$ MeV, assuming $m_Z = 91.181$ GeV and $\Gamma(Z) = 2.455$ GeV.

See key on page 1343

Gauge & Higgs Boson Full Listings

Z

- ⁵⁸ ABREU 91F report $\Gamma(\mu\mu) = 86.9 \pm 1.9 \pm 0.9$ MeV and provided us with this branching ratio from the same data and analysis.
- ⁵⁹ ADEVA 91E report $B(\mu\mu) = 3.38 \pm 0.08\%$ and $\Gamma(\mu\mu) = 84.5 \pm 2.0$ MeV and provided us with this branching ratio from the same data and analysis.
- ⁶⁰ ALEXANDER 91F report $\Gamma(\mu\mu) = 83.2 \pm 1.5$ MeV and provided us with this branching ratio from the same data and analysis.




VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
20.80 ± 0.08 OUR FIT					
20.81 ± 0.08 OUR AVERAGE					
20.68 ± 0.18	25k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV	
20.80 ± 0.20	25k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV	
21.01 ± 0.15	47k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV	
20.70 ± 0.16	45.1k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
15.2 $\begin{smallmatrix} +4.8 \\ -3.9 \end{smallmatrix}$	21	⁶¹ ABRAMS	89D MRK2	$E_{cm}^{ee} = 89-93$ GeV	
••• We do not use the following data for averages, fits, limits, etc. •••					
21.22 ± 0.25	18k	ACTON	93D OPAL	Repl. by AKERS 94	
20.77 ± 0.23		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94	
20.86 ± 0.13	50k	⁶² QUASt	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
21.00 ± 0.36	6260	DECAMP	92B ALEP	Repl. by BUSKULIC 93J	
21.02 ± 0.23	17k	⁶³ LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
21.14 ± 0.49 ± 0.27	2345	⁶⁴ ABREU	91F DLPH	Repl. by ABREU 94	
20.83 ± 0.43 ± 0.43	2540	⁶⁵ ADEVA	91E L3	Repl. by ACCIARRI 94	
21.05 ± 0.44	5559	⁶⁶ ALEXANDER	91F OPAL	Repl. by ACTON 93D	

- ⁶¹ ABRAMS 89D have included both statistical and systematic uncertainties in their quoted errors.
- ⁶² QUASt 93 is a combined analysis of LEP results as of Feb. 1993.
- ⁶³ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.
- ⁶⁴ ABREU 91F report $\Gamma(\tau\tau) = 82.7 \pm 2.1 \pm 1.1$ MeV and provided us with this branching ratio from the same data and analysis.
- ⁶⁵ ADEVA 91E report $B(\tau\tau) = 3.36 \pm 0.11\%$ and $\Gamma(\tau\tau) = 84.0 \pm 2.7$ MeV and provided us with this branching ratio from the same data and analysis.
- ⁶⁶ ALEXANDER 91F report $\Gamma(\tau\tau) = 82.7 \pm 1.9$ MeV and provided us with this branching ratio from the same data and analysis.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_4
20.76 ± 0.05 OUR FIT					
20.77 ± 0.07 OUR AVERAGE					
20.62 ± 0.10	102k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV	
20.93 ± 0.10	97k	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV	
20.835 ± 0.086	146k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV	
20.69 ± 0.09	137.3k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
18.9 $\begin{smallmatrix} +3.6 \\ -3.2 \end{smallmatrix}$	46	ABRAMS	89B MRK2	$E_{cm}^{ee} = 89-93$ GeV	
••• We do not use the following data for averages, fits, limits, etc. •••					
20.88 ± 0.13	58k	ACTON	93D OPAL	Repl. by AKERS 94	
21.00 ± 0.15	40k	ADRIANI	93M L3	Repl. by ACCIARRI 94	
20.78 ± 0.13		BUSKULIC	93J ALEP	Repl. by BUSKULIC 94	
20.87 ± 0.07	190k	⁶⁷ QUASt	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
20.90 ± 0.15	50k	⁶⁸ BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
21.00 ± 0.20	20k	DECAMP	92B ALEP	Repl. by BUSKULIC 93J	
20.89 ± 0.13	57k	⁶⁹ LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
20.70 ± 0.25 ± 0.14	10k	ABREU	91F DLPH	Repl. by ABREU 94	

- 20.84 ± 0.29 10k ADEVA 91E L3 Repl. by ADRIANI 93M
- 20.95 ± 0.22 18k ALEXANDER 91F OPAL Repl. by ACTON 93D
- ⁶⁷ QUASt 93 is a combined analysis of LEP results as of Feb. 1993. Assumes lepton universality.
- ⁶⁸ BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991.
- ⁶⁹ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.



VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
0.6990 ± 0.0015 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.6983 ± 0.0023	1.14M	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
0.6993 ± 0.0031	570k	⁷⁰ LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
⁷⁰ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included. Assumes lepton universality.					

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.03366 ± 0.00008 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.03383 ± 0.00013	45.8k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
0.03345 ± 0.00020	19k	⁷¹ LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
⁷¹ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.					

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.03367 ± 0.00013 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.03344 ± 0.00026	46.4k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
0.03351 ± 0.00034	21k	⁷² LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
⁷² LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.					

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.03360 ± 0.00015 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.03366 ± 0.00028	45.1k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
0.03328 ± 0.00040	17k	⁷³ LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
⁷³ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.					

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.03367 ± 0.00006 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.03375 ± 0.00009	137.3k	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV	
0.03347 ± 0.00013	57k	⁷⁴ LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV	
⁷⁴ LEP 92 is combined analysis by the four LEP experiments as of December 1991. Systematic errors are included.					

Gauge & Higgs Boson Full Listings

Z

 $\Gamma(\text{Invisible})/\Gamma_{\text{total}}$ See the data, the note, and the fit result for the partial width, Γ_5 , above. Γ_5/Γ

VALUE	DOCUMENT ID
0.2001 ± 0.0016 OUR FIT	

 $\Gamma(\mu^+ \mu^-)/\Gamma(e^+ e^-)$

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

 Γ_2/Γ_1

VALUE	DOCUMENT ID
1.000 ± 0.005 OUR FIT	

 $\Gamma(\tau^+ \tau^-)/\Gamma(e^+ e^-)$

This parameter is not directly used in the overall fit but is derived using the fit results; see the 'Note on the Z Boson.'

 Γ_3/Γ_1

VALUE	DOCUMENT ID
0.998 ± 0.005 OUR FIT	

 $\Gamma((u\bar{u} + c\bar{c})/2)/\Gamma(\text{hadrons})$ This quantity is the branching ratio of $Z \rightarrow$ "up-type" quarks to $Z \rightarrow$ hadrons. Γ_7/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.139 ± 0.026	75	ACTON	93F OPAL $E_{\text{cm}}^e = 88-94$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.191 ± 0.031 ± 0.040 76 ALEXANDER 91E OPAL Repl. by ACTON 93F

75 ACTON 93F use the LEP 92 value of $\Gamma(\text{hadrons}) = 1740 \pm 12$ MeV.

76 ALEXANDER 91E result is from analysis of final state photons.

 $\Gamma((d\bar{d} + s\bar{s} + b\bar{b})/3)/\Gamma(\text{hadrons})$ This quantity is the branching ratio of $Z \rightarrow$ "down-type" quarks to $Z \rightarrow$ hadrons. Γ_8/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.241 ± 0.017	77	ACTON	93F OPAL $E_{\text{cm}}^e = 88-94$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.206 ± 0.021 ± 0.028 78 ALEXANDER 91E OPAL Repl. by ACTON 93F

77 ACTON 93F use the LEP 92 value of $\Gamma(\text{hadrons}) = 1740 \pm 12$ MeV.

78 ALEXANDER 91E result is from analysis of final state photons.

 $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$

Our average consists of a simple weighted average assuming no common errors.

 Γ_9/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.171 ± 0.020 OUR AVERAGE				

0.187 ± 0.031 ± 0.023 79 ABREU 93I DLPH $E_{\text{cm}}^e = 88-94$ GeV0.151 ± 0.008 ± 0.041 80 ABREU 92O DLPH $E_{\text{cm}}^e = 88-94$ GeV0.186 ± 0.035 ± 0.020 115 81 ALEXANDER 91B OPAL $E_{\text{cm}}^e = 88-95$ GeV0.162 ± 0.030 ± 0.050 381 82 ABREU 90H DLPH $E_{\text{cm}}^e = 91$ GeV0.148 ± 0.044 $^{+0.045}_{-0.038}$ 1383 83 DECAMP 90L ALEP $E_{\text{cm}}^e = 88-94$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.223 ± 0.032 ± 0.059 84 AKRAWY 91E OPAL $E_{\text{cm}}^e = 88-95$ GeV79 ABREU 93I assume that the D_s and charmed baryons are equally produced at LEP and CLEO (10 GeV) energies.

80 ABREU 92O use the neural network technique to tag heavy flavour events among a sample of 123k selected hadronic events. The systematic error consists of three parts: due to Monte Carlo (MC) parametrization (0.023), choice of MC model (0.033) and detector effects (0.009) added in quadrature.

81 ALEXANDER 91B (OPAL) obtains the result from an analysis of the $D^{*\pm}$ momentum distribution (c production is tagged via $D^{*\pm}$ production). ALEXANDER 91B include all errors due to their experiment in the first quoted error and all others in the second (± 0.020).82 ABREU 90H use CLEO probability for $c\bar{c} \rightarrow D^*(2010)^\pm X$ with $D^*(2010)^\pm \rightarrow D^0 \pi^\pm$. Systematic error includes ± 0.026 due to uncertainties in branching ratios.83 DECAMP 90L find $B(c \rightarrow e)\Gamma(c\bar{c})/\Gamma(\text{hadrons}) = 0.0133 \pm 0.0040 \pm 0.0038$. Assumes $B(c \rightarrow e) = 0.090 \pm 0.013$. Systematic error includes about ± 0.025 due to uncertainties in branching ratios.84 AKRAWY 91E (OPAL) performs a fit to the p and p_T spectra of muon candidates, used to tag heavy flavor semileptonic decays. AKRAWY 91E systematic error includes the uncertainty from semileptonic branching ratios (± 0.025) plus other systematics (± 0.053). $\Gamma(b\bar{b})/\Gamma(\text{hadrons})$ Following the procedure of the joint LEP effort to extract $\Gamma(b\bar{b})/\Gamma(\text{hadrons})$ (see the 'Note on the Z Boson'), we have divided the measurements into 3 classes, based on lepton tagging, event shapes, and lifetime tagging. For each group, the systematic errors are split into those specific to an experiment (uncorrelated systematics) and those in common between the results (common systematics). For the overall average, the results from the three tagging techniques are assumed to be entirely uncorrelated. Γ_{10}/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2210 ± 0.0029 OUR EVALUATION				

0.218 ± 0.006 ± 0.010 85 AKERS 94D OPAL $E_{\text{cm}}^e = 88-94$ GeV0.220 ± 0.002 ± 0.013 11893 86 ACTON 93I OPAL $E_{\text{cm}}^e = 88-94$ GeV0.222 ± 0.007 ± 0.008 87 ACTON 93M OPAL $E_{\text{cm}}^e = 88-94$ GeV0.222 ± 0.003 ± 0.007 88 ADRIANI 93E L3 $E_{\text{cm}}^e = 88-94$ GeV0.2187 ± 0.0022 ± 0.0031 89 BUSKULIC 93M ALEP $E_{\text{cm}}^e = 91.3$ GeV0.228 ± 0.005 ± 0.005 90 BUSKULIC 93N ALEP $E_{\text{cm}}^e = 88-94$ GeV0.232 ± 0.005 ± 0.017 91 ABREU 92O DLPH $E_{\text{cm}}^e = 88-94$ GeV0.221 ± 0.004 ± 0.013 3893 92 ADEVA 91C L3 $E_{\text{cm}}^e = 88-95$ GeV0.251 ± 0.049 ± 0.030 32 93 JACOBSEN 91 MRK2 $E_{\text{cm}}^e = 91$ GeV0.215 ± 0.017 ± 0.024 1383 94 DECAMP 90L ALEP $E_{\text{cm}}^e = 88-94$ GeV0.23 $^{+0.10}_{-0.08}$ ± 0.05 $^{-0.04}$ 15 95 KRAL 90 MRK2 $E_{\text{cm}}^e = 89-93$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

0.222 ± 0.011 ± 0.007 96 AKERS 93B OPAL $E_{\text{cm}}^e = 88-94$ GeV0.222 $^{+0.033}_{-0.031}$ ± 0.017 97 ABREU 92 DLPH $E_{\text{cm}}^e = 88-94$ GeV0.219 ± 0.014 ± 0.019 98 ABREU 92K DLPH $E_{\text{cm}}^e = 88-94$ GeV

0.226 ± 0.008 ± 0.018 1180 99 ACTON 92I OPAL Repl. by ACTON 93I

0.193 ± 0.006 ± 0.024 1494 100 AKRAWY 91E OPAL Repl. by ACTON 93I

0.204 ± 0.014 ± 0.024 171 101 ADEVA 90E L3 Repl. by ADRIANI 93E

85 AKERS 94D perform an analysis based on a "mixed tag" method (impact parameter and lepton tagging). The systematic error includes a contribution (± 0.007) due to the $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$ uncertainty.86 ACTON 93I use both electrons and muons to tag B semileptonic decays. The systematic error includes components due to b and c quark fragmentation uncertainties, decay branching ratios, and $\Gamma(c\bar{c})/\Gamma(\text{hadrons})$.87 ACTON 93M tagged $Z \rightarrow b\bar{b}$ events using the impact parameter technique.

88 ADRIANI 93E use a multidimensional analysis based on a neural network approach.

89 BUSKULIC 93M use a method which tags the $Z \rightarrow b\bar{b}$ decays through the lifetime of the produced heavy hadrons. The systematic error includes a contribution of ± 0.0016 due to the uncertainty of the charm partial width (for compatibility, we quote the result obtained with the Standard Model prediction $\Gamma(c\bar{c})/\Gamma(\text{hadrons}) = 0.171$).90 BUSKULIC 93N use event shape and high p_T lepton discriminators applied to both hemispheres.

91 ABREU 92O use the neural network technique to tag heavy flavour events among a sample of 123k selected hadronic events. The systematic error consists of three parts: due to Monte Carlo (MC) parametrization (0.010), choice of MC model (0.008), and detector effects (0.011) added in quadrature.

92 ADEVA 91C report $\Gamma(b\bar{b}) = 385 \pm 7 \pm 11 \pm 19$ MeV; we use their $\Gamma(\text{hadrons}) = 1742 \pm 19$ MeV to obtain the branching ratio. The systematic error includes the semileptonic branching ratio uncertainty (± 0.011) plus other systematics (± 0.006).93 JACOBSEN 91 tagged $b\bar{b}$ events by requiring coincidence of ≥ 3 tracks with significant impact parameters using vertex detector. Systematic error includes lifetime and decay uncertainties (± 0.014).94 DECAMP 90L find $B(b \rightarrow e)\Gamma(b\bar{b})/\Gamma(\text{hadrons}) = 0.0219 \pm 0.0017 \pm 0.0010$. They assume $B(b \rightarrow e) = 0.102 \pm 0.007 \pm 0.007$. The quoted systematic error is dominated by that from the semileptonic branching ratio.95 KRAL 90 used isolated leptons and found $\Gamma(b\bar{b})/\Gamma(\text{total}) = 0.17 \pm 0.07 \pm 0.04$ 96 AKERS 93B use a simultaneous fit to single and dilepton events (electrons and muons) to tag $Z \rightarrow b\bar{b}$.97 ABREU 92 result is from an indirect technique. They measure the lifetime τ_B , but use a world average of τ_B independent of $\Gamma(b\bar{b})$ and compare to their $\Gamma(b\bar{b})$ dependent lifetime from a hadron sample.98 ABREU 92K use boosted-sphericity technique to tag and enrich the $b\bar{b}$ content with a sample of 50k hadronic events. Most of the systematic error is from hadronization uncertainty.99 ACTON 92I use high p (> 4 GeV), high p_T (> 0.8 GeV) electrons to tag $Z \rightarrow b\bar{b}$ events.100 For AKRAWY 91E, the systematic error includes the uncertainty from semileptonic branching ratios (± 0.021) plus other systematics (± 0.011).101 ADEVA 90E used isolated muons and found $B(B \rightarrow \mu)\Gamma(b\bar{b}) = 41.7 \pm 2.9 \pm 3.0$ MeV. The systematic error of ± 0.024 above includes 0.02 due to uncertainty in $B(B \rightarrow \mu)$ added in quadrature to ± 0.014 systematic. $\Gamma(\pi^0 \gamma)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 5.5 × 10⁻⁵	95	102 ABREU	94B DLPH	$E_{\text{cm}}^e = 88-94$ GeV
< 1.2 × 10 ⁻⁴	95	103 ADRIANI	92B L3	$E_{\text{cm}}^e = 88-94$ GeV
< 2.1 × 10 ⁻⁴	95	DECAMP	92 ALEP	$E_{\text{cm}}^e = 88-94$ GeV
< 1.4 × 10 ⁻⁴	95	AKRAWY	91F OPAL	$E_{\text{cm}}^e = 88-94$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.3 × 10⁻⁴ 95 BARDADIN... 92 RVUE $E_{\text{cm}}^e = 88-94$ GeV< 1.5 × 10⁻⁴ 95 ABREU 91E DLPH Repl. by ABREU 94B< 2.9 × 10⁻⁴ 95 ADEVA 90K L3 Repl. by ADRIANI 92B< 3.9 × 10⁻⁴ 95 AKRAWY 90F OPAL $E_{\text{cm}}^e = 88-95$ GeV< 4.9 × 10⁻⁴ 95 DECAMP 90J ALEP $E_{\text{cm}}^e = 88-95$ GeV

102 ABREU 94B supersedes ABREU 91E.

103 This limit is for both decay modes $Z^0 \rightarrow \pi^0 \gamma/\gamma\gamma$ which are indistinguishable in ADRIANI 92B. $\Gamma(\eta\gamma)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 8.0 × 10 ⁻⁵	95	104 ABREU	94B DLPH	$E_{\text{cm}}^e = 88-94$ GeV
< 1.8 × 10 ⁻⁴	95	ADRIANI	92B L3	$E_{\text{cm}}^e = 88-94$ GeV
< 5.1 × 10⁻⁵	95	DECAMP	92 ALEP	$E_{\text{cm}}^e = 88-94$ GeV
< 2.0 × 10 ⁻⁴	95	AKRAWY	91F OPAL	$E_{\text{cm}}^e = 88-94$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.9 × 10⁻⁴ 95 BARDADIN... 92 RVUE $E_{\text{cm}}^e = 88-94$ GeV< 2.8 × 10⁻⁴ 95 ABREU 91E DLPH Repl. by ABREU 94B< 4.1 × 10⁻⁴ 95 ADEVA 90K L3 Repl. by ADRIANI 92B< 5.8 × 10⁻⁴ 95 AKRAWY 90F OPAL $E_{\text{cm}}^e = 88-95$ GeV< 4.6 × 10⁻⁴ 95 DECAMP 90J ALEP $E_{\text{cm}}^e = 88-95$ GeV

104 ABREU 94B supersedes ABREU 91E.

 $\Gamma(\omega\gamma)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 6.5 × 10⁻⁴	95	ABREU	94B DLPH	$E_{\text{cm}}^e = 88-94$ GeV

$\Gamma(\eta(958)\gamma)/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<2.2 \times 10^{-4}$	95	DECAMP	90J ALEP	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{15}/Γ

This decay would violate the Landau-Yang theorem.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-5}$	95	ABREU	94B DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
$<1.2 \times 10^{-4}$	95	105 ADRIANI	92B L3	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
$<1.4 \times 10^{-4}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<2.9 \times 10^{-4}$	95	ADEVA	90K L3	Repl. by ADRIANI 92B
$<3.7 \times 10^{-4}$	95	AKRAWY	90F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV

105 This limit is for both decay modes $Z^0 \rightarrow \pi^0 \gamma/\gamma\gamma$ which are indistinguishable in ADRIANI 92B.

 $\Gamma(\gamma\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-5}$	95	106 ABREU	94B DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
$<3.3 \times 10^{-5}$	95	ADRIANI	92B L3	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
$<6.6 \times 10^{-5}$	95	AKRAWY	91F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<2.4 \times 10^{-5}$	95	BARADIN...	92 RVUE	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
$<1.4 \times 10^{-4}$	95	ABREU	91E DLPH	Repl. by ABREU 94B
$<1.2 \times 10^{-4}$	95	ADEVA	90K L3	Repl. by ADRIANI 92B
$<2.8 \times 10^{-4}$	95	AKRAWY	90F OPAL	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV

106 ABREU 94B supersedes ABREU 91E.

 $\Gamma(\pi^\pm W^\mp)/\Gamma_{\text{total}}$ Γ_{17}/Γ

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

 $\Gamma(\rho^\pm W^\mp)/\Gamma_{\text{total}}$ Γ_{18}/Γ

The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

 $\Gamma(K^0 X)/\Gamma(\text{hadrons})$ Γ_{19}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.879 ± 0.008 OUR AVERAGE			
0.880 ± 0.009	107 ABREU	92G DLPH	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV
0.877 ± 0.017	107 ALEXANDER	91C OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV

107 From multiplicity measurement in Z hadronic decays. The quoted value is summed over particle plus antiparticle. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(K^*(892)^\pm X)/\Gamma(\text{hadrons})$ Γ_{20}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.736 ± 0.068	108 ABREU	92G DLPH	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV

108 From multiplicity measurement in Z hadronic decays. The quoted value is summed over particle plus antiparticle. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(\Lambda X)/\Gamma(\text{hadrons})$ Γ_{21}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.298 ± 0.009 OUR AVERAGE			
0.300 ± 0.012	109 ABREU	93L DLPH	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV
0.296 ± 0.014	109 ACTON	92J OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••			
0.302 ± 0.049	109 ABREU	92G DLPH	Repl. by ABREU 93L

109 From multiplicity measurement in Z hadronic decays. The quoted value is summed over particle plus antiparticle. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(\Xi^- X)/\Gamma(\text{hadrons})$ Γ_{22}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.0203 ± 0.0019 OUR AVERAGE			
0.020 ± 0.005	110 ABREU	92G DLPH	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV
0.0204 ± 0.0021	110 ACTON	92J OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV

110 From multiplicity measurement in Z hadronic decays. The quoted value is summed over particle plus antiparticle. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(\Sigma(1385)^\pm X)/\Gamma(\text{hadrons})$ Γ_{23}/Γ_6

The quoted value is summed over two charge states and over particle plus antiparticle.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0373 ± 0.0060	111 ACTON	92J OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV

111 From multiplicity measurement in Z hadronic decays. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(\Xi(1530)^0 X)/\Gamma(\text{hadrons})$ Γ_{24}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.0063 ± 0.0014	112 ACTON	92J OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV

112 From multiplicity measurement in Z hadronic decays. The quoted value is summed over particle plus antiparticle. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(\Omega^- X)/\Gamma(\text{hadrons})$ Γ_{25}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
0.0063 ± 0.0015	113 ACTON	92J OPAL	$E_{\text{cm}}^{\text{ee}} = 91.2$ GeV

113 From multiplicity measurement in Z hadronic decays. The quoted value is summed over particle plus antiparticle. We have calculated this value as the probability of producing at least one particle of this kind in the final state, using as average the total yield per hadronic event as measured by the authors. We have assumed Poisson statistics but have not added a systematic error to account for possible particle/antiparticle production correlation and for deviations from Poisson statistics.

 $\Gamma(J/\psi(1S)X)/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
3.8 ± 0.5 OUR AVERAGE				
3.6 ± 0.5 ± 0.4	121	114 ADRIANI	93J L3	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
4.5 ± 0.8 ± 0.7	115	ALEXANDER	91G OPAL	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

114 ADRIANI 93J combine $\mu^+ \mu^-$ and $e^+ e^-$ channels and take into account the common systematic errors.

115 ALEXANDER 91G systematic error includes 0.4×10^{-3} systematic plus 0.6×10^{-3} from error on $J/\psi(1S) \rightarrow \ell^+ \ell^-$ branching fraction. The value is obtained by multiplying the value in ALEXANDER 91G by $\Gamma(\text{hadrons})/\Gamma(\text{total})$.

 $\Gamma(X_{c1}(1P)X)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
7.5 ± 2.9 ± 0.6	19	116 ADRIANI	93J L3	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

116 ADRIANI 93J measure this branching ratio via the decay channel $X_{c1} \rightarrow J/\psi + \gamma$, with $J/\psi \rightarrow \mu^+ \mu^-$.

 $\Gamma((D^0/\bar{D}^0)X)/\Gamma(\text{hadrons})$ Γ_{28}/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.403 ± 0.038 ± 0.044	369	117 ABREU	93I DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

117 The (D^0/\bar{D}^0) states in ABREU 93I are detected by the $K\pi$ decay mode.

 $\Gamma(D^\pm X)/\Gamma(\text{hadrons})$ Γ_{29}/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.199 ± 0.019 ± 0.024	539	118 ABREU	93I DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

118 The D^\pm states in ABREU 93I are detected by the $K\pi\pi$ decay mode.

 $\Gamma(D^*(2010)^\pm X)/\Gamma(\text{hadrons})$ Γ_{30}/Γ_6

The value is for the sum of the charge states indicated.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.179 ± 0.018 OUR AVERAGE				
0.171 ± 0.012 ± 0.016	358	119 ABREU	93I DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV
0.21 ± 0.04	362	120 DECAMP	91J ALEP	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

119 $D^*(2010)^\pm$ in ABREU 93I are reconstructed from $D^0 \pi^\pm$, with $D^0 \rightarrow K^- \pi^+$. The new CLEO II measurement of $B(D^{*\pm} \rightarrow D^0 \pi^\pm) = (68.1 \pm 1.6)\%$ is used.

120 DECAMP 91J report $B(D^*(2010)^+ \rightarrow D^0 \pi^+) B(D^0 \rightarrow K^- \pi^+) \Gamma(D^*(2010)^\pm X)/\Gamma(\text{hadrons}) = (5.11 \pm 0.34) \times 10^{-3}$. They obtained above number assuming $B(D^0 \rightarrow K^- \pi^+) = (3.62 \pm 0.34 \pm 0.44)\%$ and $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (55 \pm 4)\%$. We have rescaled their original result of 0.26 ± 0.05 taking into account the new CLEO II branching ratio $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (68.1 \pm 1.6)\%$.

 $\Gamma(B_s^0 X)/\Gamma(\text{hadrons})$ Γ_{31}/Γ_6

VALUE	DOCUMENT ID	TECN	COMMENT
seen	121 ABREU	92M DLPH	$E_{\text{cm}}^{\text{ee}} = 88-94$ GeV

121 ABREU 92M reported value is $\Gamma(B_s^0 X) \cdot B(B_s^0 \rightarrow D_s \mu \nu_\mu X) \cdot B(D_s \rightarrow \phi \pi)/\Gamma(\text{hadrons}) = (18 \pm 8) \times 10^{-5}$.

 $\Gamma(\text{anomalous } \gamma + \text{hadrons})/\Gamma_{\text{total}}$ Γ_{32}/Γ

Limits on additional sources of prompt photons beyond expectations for final-state bremsstrahlung.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-3}$	95	122 AKRAWY	90J OPAL	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV

122 AKRAWY 90J report $\Gamma(\gamma X) < 8.2$ MeV at 95%CL. They assume a three-body $\gamma q \bar{q}$ distribution and use $E(\gamma) > 10$ GeV.

Gauge & Higgs Boson Full Listings

Z

$\Gamma(e^+e^-\gamma)/\Gamma_{total}$ Γ₃₃/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-4}$	95	123 ACTON	91B OPAL	$E_{cm}^{ee} = 91.1$ GeV

123 ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy (> 0.9 GeV).

$\Gamma(\mu^+\mu^-\gamma)/\Gamma_{total}$ Γ₃₄/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.6 \times 10^{-4}$	95	124 ACTON	91B OPAL	$E_{cm}^{ee} = 91.1$ GeV

124 ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy (> 0.9 GeV).

$\Gamma(\tau^+\tau^-\gamma)/\Gamma_{total}$ Γ₃₅/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.3 \times 10^{-4}$	95	125 ACTON	91B OPAL	$E_{cm}^{ee} = 91.1$ GeV

125 ACTON 91B looked for isolated photons with $E > 2\%$ of beam energy (> 0.9 GeV).

$\Gamma(\ell^+\ell^-\gamma\gamma)/\Gamma_{total}$ Γ₃₆/Γ

The value is the sum over $\ell = e, \mu, \tau$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-6}$	95	126 ACTON	93E OPAL	$E_{cm}^{ee} = 88-94$ GeV

126 For $m_{\gamma\gamma} = 60 \pm 5$ GeV.

$\Gamma(q\bar{q}\gamma\gamma)/\Gamma_{total}$ Γ₃₇/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.5 \times 10^{-6}$	95	127 ACTON	93E OPAL	$E_{cm}^{ee} = 88-94$ GeV

127 For $m_{\gamma\gamma} = 60 \pm 5$ GeV.

$\Gamma(\nu\bar{\nu}\gamma\gamma)/\Gamma_{total}$ Γ₃₈/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.1 \times 10^{-6}$	95	128 ACTON	93E OPAL	$E_{cm}^{ee} = 88-94$ GeV

128 For $m_{\gamma\gamma} = 60 \pm 5$ GeV.

$\Gamma(e^{\pm}\mu^{\mp})/\Gamma(e^+e^-)$ Γ₃₉/Γ₁

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.07	90	ALBAJAR	89 UA1	$E_{cm}^{p\bar{p}} = 546,630$ GeV

$\Gamma(e^{\pm}\mu^{\mp})/\Gamma_{total}$ Γ₃₉/Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-5}$	95	ABREU	93B DLPH	$E_{cm}^{ee} = 88-94$ GeV
$<0.6 \times 10^{-5}$	95	ADRIANI	93i L3	$E_{cm}^{ee} = 88-94$ GeV
$<2.6 \times 10^{-5}$	95	DECAMP	92 ALEP	$E_{cm}^{ee} = 88-94$ GeV
$<4.6 \times 10^{-5}$	95	AKRAWY	91B OPAL	$E_{cm}^{ee} = 88-94$ GeV

$\Gamma(e^{\pm}\tau^{\mp})/\Gamma_{total}$ Γ₄₀/Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-4}$	95	ABREU	93B DLPH	$E_{cm}^{ee} = 88-94$ GeV
$<1.3 \times 10^{-5}$	95	ADRIANI	93i L3	$E_{cm}^{ee} = 88-94$ GeV
$<1.2 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{cm}^{ee} = 88-94$ GeV
$<7.2 \times 10^{-5}$	95	AKRAWY	91B OPAL	$E_{cm}^{ee} = 88-94$ GeV

$\Gamma(\mu^{\pm}\tau^{\mp})/\Gamma_{total}$ Γ₄₁/Γ

Test of lepton family number conservation. The value is for the sum of the charge states indicated.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-4}$	95	ABREU	93B DLPH	$E_{cm}^{ee} = 88-94$ GeV
$<1.9 \times 10^{-5}$	95	ADRIANI	93i L3	$E_{cm}^{ee} = 88-94$ GeV
$<1.0 \times 10^{-4}$	95	DECAMP	92 ALEP	$E_{cm}^{ee} = 88-94$ GeV
$<3.5 \times 10^{-4}$	95	AKRAWY	91B OPAL	$E_{cm}^{ee} = 88-94$ GeV

Z HADRONIC POLE CROSS SECTION

This quantity is defined as

$$\sigma_h^0 = \frac{12\pi}{M_Z^2} \frac{\Gamma(e^+e^-)\Gamma(\text{hadrons})}{\Gamma_Z^2}$$

It is one of the parameters used in the Z lineshape fit. (See the 'Note on the Z Boson'.)

VALUE (nb)	EVTs	DOCUMENT ID	TECN	COMMENT
41.55 ± 0.14 OUR FIT				
41.49 ± 0.10 OUR AVERAGE				
41.23 ± 0.20	1.15M	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
41.39 ± 0.26	1.2M	ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
41.70 ± 0.23	1.2M	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
41.60 ± 0.16	1.28M	BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
42 ± 4	450	ABRAMS	89B MRK2	$E_{cm}^{ee} = 89.2-93.0$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

41.45 ± 0.31	512k	ACTON	93D OPAL	Repl. by AKERS 94
41.34 ± 0.28	460k	ADRIANI	93M L3	Repl. by ACCIARRI 94
41.60 ± 0.27	520k	BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
41.33 ± 0.23	650k	129 LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
41.84 ± 0.45	150k	ABREU	91F DLPH	Repl. by ABREU 94

129 LEP 92 is a combined analysis by the four LEP experiments as of Dec 1991.

Z VECTOR COUPLINGS

These quantities are the effective vector couplings of the Z to charged leptons. Their magnitude is derived from a measurement of the Z lineshape and the forward-backward lepton asymmetries as a function of energy around the Z mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the Z asymmetry parameters, A_e and A_τ , or ν_e scattering. The fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and A_e and A_τ measurements. See "Note on the Z boson" for details.

Within the current data set, the reason for the smallness of g_V^μ compared to g_V^e and g_V^τ is due to the large value of A_e which is heavily weighted by the SLD result. This large value of A_e leads to a large value of g_V^e . Since g_V^μ is obtained using the relation $A_{FB}^\mu = 0.75 \times A_e \times A_\mu$, a large value of g_V^e leads to a SMALL value of g_V^μ . Concerning the τ , its g_V^μ gets mainly determined directly from A_τ which is obtained from a measurement of the τ polarization (see "Note on the Z boson").

g_V^e

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

-0.0398 +0.0024 -0.0023 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.0364 +0.0096 -0.0082	38k	130 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.036 ± 0.005	45.8k	131 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.040 +0.013 -0.011		132 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.034 +0.006 -0.005		130 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
-0.035 ± 0.005	70k	133 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.006 +0.003 -0.004	16k	134 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.045 +0.013 -0.011	6947	135 DECAMP	92B ALEP	Repl. by BUSKULIC 93J
-0.035 +0.013 -0.014	6947	136 DECAMP	92B ALEP	Repl. by BUSKULIC 93J

130 The τ polarization result has been included.

131 BUSKULIC 94 use the added constraint of τ polarization.

132 ADRIANI 93M use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

133 QUAST 93 is a combined analysis of LEP results as of Feb. 1993. QUAST 93 use also the average LEP values for τ polarization and the forward-backward τ polarisation asymmetry.

134 BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991. Only forward-backward lepton asymmetries are used.

135 DECAMP 92B use their measurement of the τ polarization in addition to the forward-backward lepton asymmetries.

136 Using only forward-backward lepton asymmetries.

g_V^μ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

-0.0273 +0.0056 -0.0058 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.0402 +0.0153 -0.0211	34k	137 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.034 ± 0.013	46.4k	138 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.048 +0.021 -0.033		139 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.019 +0.018 -0.019		137 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
-0.029 ± 0.010	70k	140 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.191 +0.088 -0.081	16k	141 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.018 +0.023 -0.026	6691	142 DECAMP	92B ALEP	Repl. by BUSKULIC 93J
-0.023 +0.029 -0.037	6691	143 DECAMP	92B ALEP	Repl. by BUSKULIC 93J

137 The τ polarization result has been included.

138 BUSKULIC 94 use the added constraint of τ polarization.

139 ADRIANI 93M use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.

140 QUAST 93 is a combined analysis of LEP results as of Feb. 1993. QUAST 93 use also the average LEP values for τ polarization and the forward-backward τ polarisation asymmetry.

141 BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991. Only forward-backward lepton asymmetries are used.

142 DECAMP 92B use their measurement of the τ polarization in addition to the forward-backward lepton asymmetries.

143 Using only forward-backward lepton asymmetries.

δ_V
VALUE EVTS DOCUMENT ID TECN COMMENT
-0.0389 ± 0.0043 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.0384 ± 0.0078	25k	144 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.038 ± 0.005	45.1k	145 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.037 ± 0.008	7441	146 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.039 ± 0.006		144 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
-0.039 ± 0.004	50k	147 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.287 +0.127 -0.077	16k	148 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.045 +0.010 -0.011	6260	149 DECAMP	92B ALEP	Repl. by BUSKULIC 93J
-0.104 +0.040 -0.066	6260	150 DECAMP	92B ALEP	Repl. by BUSKULIC 93J

- 144 The τ polarization result has been included.
 145 BUSKULIC 94 use the added constraint of τ polarization.
 146 ADRIANI 93M use their measurement of the τ polarization in addition to forward-backward lepton asymmetries.
 147 QUAST 93 is a combined analysis of LEP results as of Feb. 1993. QUAST 93 use also the average LEP values for τ polarization and the forward-backward τ polarisation asymmetry.
 148 BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991. Only forward-backward lepton asymmetries are used.
 149 DECAMP 92B use their measurement of the τ polarization in addition to the forward-backward lepton asymmetries.
 150 Using only forward-backward lepton asymmetries.

δ_V
VALUE EVTS DOCUMENT ID TECN COMMENT
-0.0377 ± 0.0016 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.039 ± 0.004	50.3k	151 ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
-0.0378 +0.0045 -0.0042	97k	152 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.034 ± 0.004	146k	151 AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
-0.038 ± 0.004	137.3k	151 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.027 ± 0.008	58k	151 ACTON	93D OPAL	Repl. by AKERS 94
-0.040 +0.006 -0.005		152 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.034 +0.004 -0.003		152 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
-0.0355 ± 0.0025	190k	153 QUAST	93 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.034 +0.009 -0.007	50k	154 BANERJEE	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.041 +0.007 -0.006	20k	152 DECAMP	92B ALEP	Repl. by BUSKULIC 93J
-0.042 +0.009 -0.007	20k	151 DECAMP	92B ALEP	Repl. by BUSKULIC 93J
-0.034 ± 0.006	57k	155 LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.017 ± 0.029	9k	151 ABREU	91F DLPH	Repl. by ABREU 94
-0.046 +0.015 -0.012	10k	151 ADEVA	91E L3	Repl. by ADRIANI 93M
-0.024 ± 0.015	18k	151 ALEXANDER	91F OPAL	Repl. by AKERS 94

- 151 Using forward-backward lepton asymmetries.
 152 The τ polarization result has been included.
 153 QUAST 93 is a combined analysis of LEP results as of Feb. 1993. QUAST 93 use also the average LEP values for τ polarization and the forward-backward τ polarisation asymmetry. Assumes lepton universality.
 154 BANERJEE 92 is a combined analysis of the four LEP experiments as of March 1991. Forward-backward lepton asymmetries are used.
 155 LEP 92 is a combined analysis of the four LEP experiments as of December 1991. Forward-backward lepton asymmetries are used.

Z AXIAL-VECTOR COUPLINGS

These quantities are the effective axial-vector couplings of the Z to charged leptons. Their magnitude is derived from a measurement of the Z lineshape and the forward-backward lepton asymmetries as a function of energy around the Z mass. The relative sign among the vector to axial-vector couplings is obtained from a measurement of the Z asymmetry parameters, A_e and A_τ , or ν_e scattering. The fit values quoted below correspond to global nine- or five-parameter fits to lineshape, lepton forward-backward asymmetry, and A_e and A_τ measurements. See "Note on the Z boson" for details.

δ_A
VALUE EVTS DOCUMENT ID TECN COMMENT
-0.5007 ± 0.0009 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.4998 ± 0.0016	38k	156 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.503 ± 0.002	45.8k	156 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.4980 ± 0.0021		156 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.5029 ± 0.0018		156 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

- 156 The τ -polarization constraint has been included.

δ_A
VALUE EVTS DOCUMENT ID TECN COMMENT
-0.5015 ± 0.0009 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.4987 +0.0030 -0.0026	34k	157 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.501 ± 0.002	46.4k	157 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.4968 +0.0050 -0.0037		157 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.5014 ± 0.0029		157 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

- 157 The τ -polarization constraint has been included.

δ_A
VALUE EVTS DOCUMENT ID TECN COMMENT
-0.5005 ± 0.0010 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.5014 ± 0.0029	25k	158 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.502 ± 0.003	45.1k	158 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.5032 ± 0.0038	7441	158 ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.5016 ± 0.0033		158 BUSKULIC	93J ALEP	Repl. by BUSKULIC 94

- 158 The τ -polarization constraint has been included.

δ_A
VALUE EVTS DOCUMENT ID TECN COMMENT
-0.5008 ± 0.0008 OUR FIT

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.4999 ± 0.0014	71k	ABREU	94 DLPH	$E_{cm}^{ee} = 88-94$ GeV
-0.4998 ± 0.0014	97k	159 ACCIARRI	94 L3	$E_{cm}^{ee} = 88-94$ GeV
-0.500 ± 0.001	146k	AKERS	94 OPAL	$E_{cm}^{ee} = 88-94$ GeV
-0.502 ± 0.001	137k	159 BUSKULIC	94 ALEP	$E_{cm}^{ee} = 88-94$ GeV
-0.4998 ± 0.0016	58k	ACTON	93D OPAL	Repl. by AKERS 94
-0.4986 ± 0.0015	159	ADRIANI	93M L3	Repl. by ACCIARRI 94
-0.5022 ± 0.0015	159	BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
-0.499 ± 0.001	57k	160 LEP	92 RVUE	$E_{cm}^{ee} = 88-94$ GeV
-0.501 ± 0.003	9k	ABREU	91F DLPH	Repl. by ABREU 94

- 159 The τ -polarization constraint has been included.

- 160 LEP 92 is a combined analysis of the four LEP experiments as of December 1991.

Z ASYMMETRY PARAMETERS

For each fermion-antifermion pair coupling to the Z these quantities are defined as

$$A_f = \frac{2g_V^f g_A^f}{(g_V^f)^2 + (g_A^f)^2}$$

where g_V^f and g_A^f are the effective vector and axial-vector couplings. For their relation to the various lepton asymmetries see the "Note on the Z Boson."

A_e
 Using polarized beams, this quantity can also be measured as $(\sigma_L - \sigma_R) / (\sigma_L + \sigma_R)$, where σ_L and σ_R are the e^+e^- production cross sections for Z bosons produced with left-handed and right-handed electrons respectively.

VALUE EVTS DOCUMENT ID TECN COMMENT
0.161 ± 0.012 OUR AVERAGE Error includes scale factor of 1.7. See the Ideogram below.


0.1656 ± 0.0071 ± 0.0028	49392	161 ABE	94C SLD	$E_{cm}^{ee} = 91.26$ GeV
0.097 ± 0.044 ± 0.004	10224	162 ABE	93 SLD	$E_{cm}^{ee} = 91.26$ GeV
0.120 ± 0.026		163 BUSKULIC	93P ALEP	$E_{cm}^{ee} = 88-94$ GeV
0.29 ± 0.13	3245	163 ALEXANDER	91D OPAL	$E_{cm}^{ee} = 88-94$ GeV

- 161 ABE 94C measured the left-right asymmetry in Z production. This value leads to $\sin^2\theta_W = 0.2292 \pm 0.0009 \pm 0.0004$.

- 162 ABE 93 measured the left-right asymmetry in Z production.

- 163 Derived from the measurement of forward-backward τ polarization asymmetry.

WEIGHTED AVERAGE
 0.161 ± 0.012 (Error scaled by 1.7)



A_e

Gauge & Higgs Boson Full Listings

Z

 A_τ

This quantity is derived from the measurement of the average τ polarization.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.141 ± 0.021 OUR AVERAGE		Error includes scale factor of 1.2.		
0.132 ± 0.033	10732	ADRIANI 93M L3		$E_{cm}^{ee} = 88-94$ GeV
0.143 ± 0.023		BUSKULIC 93P ALEP		$E_{cm}^{ee} = 88-94$ GeV
0.24 ± 0.07	2021	ABREU 92N DLPH		$E_{cm}^{ee} = 88-94$ GeV
0.01 ± 0.09	3245	ALEXANDER 91D OPAL		$E_{cm}^{ee} = 88-94$ GeV

$A_{FB}^{(0,e)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow e^+e^-$ (Including radiative corrections)

For the Z peak, we report the pole asymmetry defined by $(3/4)A_e^2$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data. For details see the "Note on the Z boson."

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.51 ± 0.40 OUR FIT				
1.5 ± 0.4 OUR AVERAGE				
2.5 ± 0.9		91.2	ABREU 94 DLPH	
1.04 ± 0.92		91.2	ACCIARRI 94 L3	
0.62 ± 0.80		91.2	AKERS 94 OPAL	
1.85 ± 0.66		91.2	BUSKULIC 94 ALEP	

$A_{FB}^{(0,\mu)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \mu^+\mu^-$ (Including radiative corrections)

For the Z peak, we report the pole asymmetry defined by $(3/4)A_e A_\mu$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data. For details see the "Note on the Z boson."

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.33 ± 0.26 OUR FIT				
1.34 ± 0.24 OUR AVERAGE				
1.4 ± 0.5		91.2	ABREU 94 DLPH	
1.79 ± 0.61		91.2	ACCIARRI 94 L3	
0.99 ± 0.42		91.2	AKERS 94 OPAL	
1.46 ± 0.48		91.2	BUSKULIC 94 ALEP	

• • • We do not use the following data for averages, fits, limits, etc. • • •

-29.0 ± 5.0 ± 0.5	-32.1	56.9	164 ABE	90I VNS
18 ± 8	+1	91.28	ADEVA	90D L3
-9.9 ± 1.5 ± 0.5	-9.2	35	HEGNER	90 JADE
0.05 ± 0.22	0.026	91.14	165 ABRAMS	89D MRK2
-43.4 ± 17.0	-24.9	52.0	166 BACALA	89 AMY
-11.0 ± 16.5	-29.4	55.0	166 BACALA	89 AMY
-30.0 ± 12.4	-31.2	56.0	166 BACALA	89 AMY
-46.2 ± 14.9	-33.0	57.0	166 BACALA	89 AMY
-29 ± 13	-25.9	53.3	ADACHI	88C TOPZ
+5.3 ± 5.0 ± 0.5	-1.2	14.0	ADEVA	88 MRKJ
-10.4 ± 1.3 ± 0.5	-8.6	34.8	ADEVA	88 MRKJ
-12.3 ± 5.3 ± 0.5	-10.7	38.3	ADEVA	88 MRKJ
-15.6 ± 3.0 ± 0.5	-14.9	43.8	ADEVA	88 MRKJ
-1.0 ± 6.0	-1.2	13.9	BRAUNSCH...	88D TASS
-9.1 ± 2.3 ± 0.5	-8.6	34.5	BRAUNSCH...	88D TASS
-10.6 ± 2.2 ± 0.5	-8.9	35.0	BRAUNSCH...	88D TASS
-17.6 ± 4.4 ± 0.5	-15.2	43.6	BRAUNSCH...	88D TASS
-4.8 ± 6.5 ± 1.0	-11.5	39	BEHREND	87C CELL
-18.8 ± 4.5 ± 1.0	-15.5	44	BEHREND	87C CELL
+2.7 ± 4.9	-1.2	13.9	BARTEL	86C JADE
-11.1 ± 1.8 ± 1.0	-8.6	34.4	BARTEL	86C JADE
-17.3 ± 4.8 ± 1.0	-13.7	41.5	BARTEL	86C JADE
-22.8 ± 5.1 ± 1.0	-16.6	44.8	BARTEL	86C JADE
-6.3 ± 0.8 ± 0.2	-6.3	29	ASH	85 MAC
-4.9 ± 1.5 ± 0.5	-5.9	29	DERRICK	85 HRS
-7.1 ± 1.7	-5.7	29	LEVI	83 MRK2
-16.1 ± 3.2	-9.2	34.2	BRANDELIK	82C TASS

164 ABE 90I measurements in the range $50 \leq \sqrt{s} \leq 60.8$ GeV.

165 ABRAMS 89D asymmetry includes both $9 \mu^+\mu^-$ and $15 \tau^+\tau^-$ events.

166 BACALA 89 systematic error is about 5%.

$A_{FB}^{(0,\tau)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \tau^+\tau^-$ (Including radiative corrections)

For the Z peak, we report the pole asymmetry defined by $(3/4)A_e A_\tau$ as determined by the nine-parameter fit to cross-section and lepton forward-backward asymmetry data. For details see the "Note on the Z boson."

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
2.12 ± 0.32 OUR FIT				
2.13 ± 0.31 OUR AVERAGE				
2.2 ± 0.7		91.2	ABREU 94 DLPH	
2.65 ± 0.88		91.2	ACCIARRI 94 L3	
2.05 ± 0.52		91.2	AKERS 94 OPAL	
1.97 ± 0.56		91.2	BUSKULIC 94 ALEP	

• • • We do not use the following data for averages, fits, limits, etc. • • •

-32.8 ± 6.4 ± 1.5	-32.1	56.9	167 ABE	90I VNS
-8.1 ± 2.0 ± 0.6	-9.2	35	HEGNER	90 JADE
-18.4 ± 19.2	-24.9	52.0	168 BACALA	89 AMY
-17.7 ± 26.1	-29.4	55.0	168 BACALA	89 AMY
-45.9 ± 16.6	-31.2	56.0	168 BACALA	89 AMY
-49.5 ± 18.0	-33.0	57.0	168 BACALA	89 AMY
-20 ± 14	-25.9	53.3	ADACHI	88C TOPZ
-10.6 ± 3.1 ± 1.5	-8.5	34.7	ADEVA	88 MRKJ
-8.5 ± 6.6 ± 1.5	-15.4	43.8	ADEVA	88 MRKJ
-6.0 ± 2.5 ± 1.0	8.8	34.6	BARTEL	85F JADE
-11.8 ± 4.6 ± 1.0	14.8	43.0	BARTEL	85F JADE
-5.5 ± 1.2 ± 0.5	-0.063	29.0	FERNANDEZ	85 MAC
-4.2 ± 2.0	0.057	29	LEVI	83 MRK2
-10.3 ± 5.2	-9.2	34.2	BEHREND	82 CELL
-0.4 ± 6.6	-9.1	34.2	BRANDELIK	82C TASS

167 ABE 90I measurements in the range $50 \leq \sqrt{s} \leq 60.8$ GeV.

168 BACALA 89 systematic error is about 5%.

$A_{FB}^{(0,l)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow \ell^+\ell^-$ (Including radiative corrections)

For the Z peak, we report the pole asymmetry defined by $(3/4)A_\ell^2$ as determined by the five-parameter fit to cross-section and lepton forward-backward asymmetry data assuming lepton universality. For details see the "Note on the Z boson."

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
1.59 ± 0.18 OUR FIT				
1.60 ± 0.18 OUR AVERAGE				
1.77 ± 0.37		91.2	ABREU 94 DLPH	
1.84 ± 0.45		91.2	ACCIARRI 94 L3	
1.28 ± 0.30		91.2	AKERS 94 OPAL	
1.71 ± 0.33		91.2	BUSKULIC 94 ALEP	

$A_{FB}^{(0,c)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow c\bar{c}$

Our estimate of the asymmetry is the value for the Z pole only. To estimate this average pole asymmetry, the various systematic errors were split into uncorrelated and correlated parts (model dependence, semileptonic branching ratios, fragmentation, $\Gamma(c\bar{c})/\Gamma(b\bar{b})$, and D^* branching ratios). QCD and QED corrections are also taken into account.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
5.8 ± 2.2 OUR ESTIMATE				
1.4 ± 3.0 ± 2.0	5.6	91.24	169 ACTON	93K OPAL
5.2 ± 2.8 ± 1.2	5.4	91.28	170 AKERS	93D OPAL
8.3 ± 3.8 ± 2.7	5.6	91.24	171 ADRIANI	92D L3
6.4 ± 3.9 ± 3.0	9.1	172	DECAMP	91E ALEP

• • • We do not use the following data for averages, fits, limits, etc. • • •

-14 ± 14 ± 3	-2	89.75	170 AKERS	93D OPAL
18 ± 12 ± 3	12	92.64	170 AKERS	93D OPAL
-12.9 ± 7.8 ± 5.5	-13.6	35	BEHREND	90D CELL
7.7 ± 13.4 ± 5.0	-22.1	43	BEHREND	90D CELL
-12.8 ± 4.4 ± 4.1	-13.6	35	ELSEN	90 JADE
-10.9 ± 12.9 ± 4.6	-23.2	44	ELSEN	90 JADE
-14.9 ± 6.7	-13.3	35	OULD-SAADA	89 JADE

169 ACTON 93K use the lepton tagging technique.

170 AKERS 93D identify the b and c decays using D^* .

171 ADRIANI 92D use both electron and muon semileptonic decays.

172 DECAMP 91E use the lepton-tagging technique.

$A_{FB}^{(0,b)}$ CHARGE ASYMMETRY IN $e^+e^- \rightarrow b\bar{b}$

Our estimate of the asymmetry is the value at the Z pole only and uses an average $LEP B^0\bar{B}^0$ mixing parameter of $(12.2 \pm 1.1)\%$. To estimate this average pole asymmetry, the various systematic errors were split into uncorrelated and correlated parts (model dependence, semileptonic branching ratios, fragmentation, $\Gamma(c\bar{c})/\Gamma(b\bar{b})$, mixing, and $c\bar{c}$ asymmetry). QCD and QED corrections are also taken into account.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
10.7 ± 1.3 OUR ESTIMATE				
9.2 ± 1.8 ± 0.8	8.5	91.24	173 ACTON	93K OPAL
13.9 ± 9.7 ± 4.9	9.4	91.28	174 AKERS	93D OPAL
16.1 ± 6.0 ± 2.1	9.1	91.2	175 ABREU	92H DLPH
9.7 ± 1.7 ± 0.7	8.2	91.24	176 ADRIANI	92D L3
12.6 ± 2.8 ± 1.2	9.1	177	DECAMP	91E ALEP

See key on page 1343

Gauge & Higgs Boson Full Listings
Z, Higgs Bosons — H^0 and H^\pm

• • • We do not use the following data for averages, fits, limits, etc. • • •

$7.1 \pm 5.4 \pm 0.7$	5.2	89.66	173	ACTON	93K	OPAL
$13.1 \pm 4.7 \pm 1.3$	10.8	92.75	173	ACTON	93K	OPAL
9.3 ± 1.1		91.2	178	QUAST	93	RVUE
$8.6 \pm 1.5 \pm 0.7$	8.2	91.24	179	ADRIANI	92D	L3
$2.5 \pm 5.1 \pm 0.7$	5.3	89.67	176	ADRIANI	92D	L3
$6.2 \pm 4.2 \pm 0.7$	10.8	92.81	176	ADRIANI	92D	L3
$9.7 \pm 5.7 \pm 1.4$	9.	91		AKRAWY	91E	OPAL
$-71 \pm 34 \pm 7$ -8	-58	58.3		SHIMONAKA	91	TOPZ
$-22.2 \pm 7.7 \pm 3.5$	-26.0	35		BEHREND	90D	CELL
$-49.1 \pm 16.0 \pm 5.0$	-39.7	43		BEHREND	90D	CELL
-28 ± 11	-23	35		BRAUNSCH...	90	TASS
$-16.6 \pm 7.7 \pm 4.8$	-24.3	35		ELSEN	90	JADE
$-33.6 \pm 22.2 \pm 5.2$	-39.9	44		ELSEN	90	JADE
$3.4 \pm 7.0 \pm 3.5$	-16.0	29.0		BAND	89	MAC
$-72 \pm 28 \pm 13$	-56	55.2		SAGAWA	89	AMY

173 ACTON 93K use the lepton tagging technique. The systematic error includes the uncertainty on the mixing parameter.

174 AKERS 93D identify the b and c decays using D^* .175 B tagging via its semimuonic decay. Experimental value corrected using average LEP B^0 - \bar{B}^0 mixing parameter $\chi = 0.143 \pm 0.023$.

176 ADRIANI 92D use both electron and muon semileptonic decays. The quoted systematic error is common to all measurements.

177 DECAMP 91E use the lepton-tagging technique.

178 QUAST 93 is a combined analysis of LEP results as of Feb. 1993.

179 ADRIANI 92D use both electron and muon semileptonic decays. For this measurement ADRIANI 92D average over all \sqrt{s} values to obtain a single result.CHARGE ASYMMETRY IN $e^+e^- \rightarrow q\bar{q}$

Summed over five lighter flavors.

Experimental and Standard Model values are somewhat event-selection dependent. Standard Model expectations contain some assumptions on B^0 - \bar{B}^0 mixing and on other electroweak parameters.

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
3.93 ± 0.65		91.2	180	QUAST 93 RVUE
$-0.76 \pm 0.12 \pm 0.15$		91.2	181	ABREU 92I DLPH
$4.0 \pm 0.4 \pm 0.63$	4.0	91.3	182	ACTON 92L OPAL
$9.1 \pm 1.4 \pm 1.6$	9.0	57.9		ADACHI 91 TOPZ
$-0.84 \pm 0.15 \pm 0.04$		91		DECAMP 91B ALEP
$8.3 \pm 2.9 \pm 1.9$	8.7	56.6		STUART 90 AMY
$11.4 \pm 2.2 \pm 2.1$	8.7	57.6		ABE 89L VNS
6.0 ± 1.3	5.0	34.8		GREENSHAW 89 JADE
8.2 ± 2.9	8.5	43.6		GREENSHAW 89 JADE

180 QUAST 93 is a combined analysis of LEP results as of Feb. 1993.

181 ABREU 92I has 0.14 systematic error due to uncertainty of quark fragmentation.

182 ACTON 92L use the weight function method on 259k selected $Z \rightarrow$ hadrons events.The systematic error includes a contribution of 0.2 due to B^0 - \bar{B}^0 mixing effect, 0.4 due to Monte Carlo (MC) fragmentation uncertainties and 0.3 due to MC statistics. ACTON 92L derive a value of $\sin^2\theta_W^{\text{eff}}$ to be $0.2321 \pm 0.0017 \pm 0.0028$.CHARGE ASYMMETRY IN $p\bar{p} \rightarrow Z \rightarrow e^+e^-$

ASYMMETRY (%)	STD. MODEL	\sqrt{s} (GeV)	DOCUMENT ID	TECN
$5.2 \pm 5.9 \pm 0.4$		91	ABE	91E CDF

• • • We do not use the following data for averages, fits, limits, etc. • • •

ABREU 92G ZPHY C53 567	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU 92G PL B275 231	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU 92H PL B276 536	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU 92I PL B277 371	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU 92M PL B281 383	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU 92M PL B289 199	+Adam, Adye, Agasi, Alekseev+	(DELPHI Collab.)
ABREU 92N ZPHY C55 555	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU 92O PL B295 383	+Adam, Adami, Adye+	(DELPHI Collab.)
ACTON 92I ZPHY C55 191	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON 92J PL B291 503	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON 92L PL B294 436	+Adriani, Aguilari-Benitez, Ahlen+	(L3 Collab.)
ADEVA 92E PL B275 209	+Aguilar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ADRIANI 92B PL B288 404	+Aguilar-Benitez, Ahlen, Akbari+	(L3 Collab.)
ADRIANI 92D PL B292 454	+Aguilar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ADRIANI 92E PL B292 463	+Aguilar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ALITTI 92B PL B276 354	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
BANERJEE 92E LMP A7 1853	+Ganguli, Guru	(TATA CLER)
BARDADIN... 92 ZPHY C55 163	+Bardadin-Otinowska	(CLER)
DECAMP 92 PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
DECAMP 92B ZPHY C53 1	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
LEP 92 PL B276 247	+ALEPH, DELPHI, L3, OPAL	(LEP Collabs.)
ABE 91E PRL 67 1502	+Amidei, Apollinari+	(CDF Collab.)
ABREU 91E PL B260 240	+Adam, Adami+	(DELPHI Collab.)
ABREU 91E PL B268 296	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ABREU 91F NP B367 511	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ACTON 91B PL B273 338	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ADACHI 91 PL B255 613	+Anazawa, Doser, Enomoto+	(TOPAZ Collab.)
ADEVA 91C PL B261 177	+Adriani, Aguilari-Benitez, Akbari+	(L3 Collab.)
ADEVA 91E ZPHY C51 179	+Adriani, Aguilari-Benitez, Akbari, Alcaraz+	(L3 Collab.)
AKRAWY 91B PL B254 293	+Alexander, Allison+	(OPAL Collab.)
AKRAWY 91D ZPHY C50 373	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
AKRAWY 91E PL B263 311	+Alexander, Allison+	(OPAL Collab.)
AKRAWY 91F NP B367 511	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ALEXANDER 91B PL B262 341	+Allison, Allport, Anderson, Arcelli+	(OPAL Collab.)
ALEXANDER 91C PL B264 467	+Allison, Allport, Anderson+	(OPAL Collab.)
ALEXANDER 91D PL B266 201	+Allison, Allport, Anderson+	(OPAL Collab.)
ALEXANDER 91E PL B264 219	+Allison, Allport+	(OPAL Collab.)
ALEXANDER 91F ZPHY C52 175	+Allison, Allport, Anderson, Arcelli+	(OPAL Collab.)
ALEXANDER 91G PL B266 485	+Allison, Allport+	(OPAL Collab.)
DECAMP 91B PL B259 377	+Deschizeaux, Goy+	(ALEPH Collab.)
DECAMP 91E PL B263 325	+Deschizeaux, Goy+	(ALEPH Collab.)
DECAMP 91J PL B266 218	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
JACOBSEN 91 PRL 67 3347	+Koetke, Adolphsen, Fujino+	(Mark II Collab.)
SHIMONAKA 91 PL B268 457	+Fujii, Miyamoto+	(TOPAZ Collab.)
ABE 90I ZPHY C48 13	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ABREU 90H PL B252 140	+Adam, Adami, Adye+	(DELPHI Collab.)
ADEVA 90D PL B238 122	+Adriani, Aguilari-Benitez, Akbari+	(L3 Collab.)
ADEVA 90E PL B241 416	+Adriani, Aguilari-Benitez, Akbari+	(L3 Collab.)
ADEVA 90K PL B250 199	+Adriani, Aguilari-Benitez, Akbari, Alcaraz+	(L3 Collab.)
AKRAWY 90F PL B241 133	+Alexander, Allison, Allport+	(OPAL Collab.)
AKRAWY 90J PL B246 285	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
BEHREND 90D ZPHY C47 333	+Criegee, Field, Franke, Jung+	(CELLO Collab.)
BRAUNSCH... 90 ZPHY C48 433	+Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
DECAMP 90J PL B241 635	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
DECAMP 90L PL B244 351	+Deschizeaux, Goy+	(ALEPH Collab.)
ELSEN 90 ZPHY C46 349	+Allison, Ambrus, Barlow, Bartel+	(JADE Collab.)
HEGNER 90 ZPHY C46 547	+Naroska, Schroth, Allison+	(JADE Collab.)
KRAL 90 PRL 64 1211	+Abrams, Adolphsen, Averill, Ballam+	(Mark II Collab.)
STUART 90 PRL 64 983	+Breedon, Kim, Ko, Lander, Maeshima+	(AMY Collab.)
ABE 89P PRL 62 613	+Amidei, Apollinari, Ascari, Atac+	(CDF Collab.)
ABE 89C PRL 63 720	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABE 89L PL B232 425	+Amako, Arai, Asano, Chiba+	(VENUS Collab.)
ABRAMS 89P PRL 63 2173	+Adolphsen, Averill, Ballam, Barish+	(Mark II Collab.)
ABRAMS 89D PR 63 2780	+Adolphsen, Averill, Ballam, Barish+	(Mark II Collab.)
ALBAJAR 89 ZPHY C44 15	+Albrow, Altkofer, Arnison, Astbury+	(UJAL Collab.)
BACALA 89 PL B218 1142	+Malchow, Sparks, Imlay, Kirk+	(AMY Collab.)
BAND 89 PL B218 369	+Camporesi, Chadwick, Delfino, Desangro+	(MAC Collab.)
GREENSHAW 89 ZPHY C42 1	+Warming, Allison, Ambrus, Barlow+	(JADE Collab.)
OULD-SAADA 89 ZPHY C44 567	+Allison, Ambrus, Barlow, Bartel+	(JADE Collab.)
SAGAWA 89 PRL 63 2341	+Lim, Abe, Fujii, Higashi+	(AMY Collab.)
ADACHI 88C PL B208 319	+Aihara, Dijkstra, Enomoto, Fujii+	(TOPAZ Collab.)
ADEVA 88 PR D38 2665	+Banderhub, Ansari, Becker+	(Mark-II Collab.)
BRAUNSCH... 88D ZPHY C40 163	+Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
ANSARI 87 PL B186 440	+Bagnaia, Banner, Battiston+	(UA2 Collab.)
BEHREND 87C PL B191 209	+Buerger, Criegee, Dainton+	(CELLO Collab.)
BARTEL 86C ZPHY C30 371	+Becker, Cords, Felst, Haidt+	(JADE Collab.)
Also 85B ZPHY C26 507	+Bartel, Becker, Bowdery, Cords+	(JADE Collab.)
Also 82 PL 108B 140	+Bartel, Cords, Dittmann, Eichler+	(JADE Collab.)
ASH 88 PRL 55 1831	+Band, Blume, Camporesi+	(MAC Collab.)
BARTEL 85F PL 161B 188	+Becker, Cords, Felst+	(JADE Collab.)
DERRICK 85 PR D31 2352	+Fernandez, Fries, Hyman+	(HRS Collab.)
FERNANDEZ 85 PRL 54 1624	+Ford, Qi, Read+	(MAC Collab.)
LEVI 83 PRL 51 1941	+Blocker, Strait+	(Mark II Collab.)
BEHREND 82 PL 114B 282	+Chen, Fenner, Field+	(CELLO Collab.)
BRANDELIC 82C PL 110B 173	+Braunschweig, Gather	(TASSO Collab.)

Z REFERENCES

ABE 94C PRL (to be pub.)	+Aht, Ash, Aston, Bacchetta, Baird+	(SLD Collab.)
ABREU 94 NP B418 403	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU 94B PL B327 386	+Adam, Adye, Agasi+	(DELPHI Collab.)
ACCIARRI 94 ZPHY C62 551	+Adam, Adriani, Aguilari-Benitez+	(L3 Collab.)
AKERS 94 ZPHY C61 19	+Alexander, Allison+	(OPAL Collab.)
AKERS 94D ZPHY C61 357	+Alexander, Allison+	(OPAL Collab.)
BUSKULIC 94 ZPHY C62 539	+Casper, De Bonis, Decamp, Ghez, Goy+	(ALEPH Collab.)
ABE 93 PRL 70 2515	+Aht, Acton+	(SLD Collab.)
ABREU 93B PL B298 247	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU 93I ZPHY C59 533	+Adam, Adye, Agasi+	(DELPHI Collab.)
ABREU 93L PL B318 249	+Adam, Adami, Adye+	(DELPHI Collab.)
ACTON 93D ZPHY C58 219	+Alexander, Allison+	(OPAL Collab.)
ACTON 93E PL B311 391	+Akers, Alexander+	(OPAL Collab.)
ACTON 93F ZPHY C58 405	+Alexander, Allison+	(OPAL Collab.)
ACTON 93I ZPHY C58 523	+Alexander, Allison+	(OPAL Collab.)
ACTON 93K ZPHY C60 19	+Akers, Alexander+	(OPAL Collab.)
ACTON 93M ZPHY C60 579	+Akers, Alexander+	(OPAL Collab.)
ADRIANI 93E PL B307 237	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI 93F PL B309 451	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI 93H PL B315 494	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI 93I PL B316 427	+Aguilar-Benitez, Ahlen+	(L3 Collab.)
ADRIANI 93J PL B317 467	+Aguilar-Benitez, Ahlen, Alcaraz+	(L3 Collab.)
ADRIANI 93M PRPL 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
Also 92H PL B294 466	+Adriani, Aguilari-Benitez, Ahlen, Alcaraz+	(L3 Collab.)
AKERS 93B ZPHY C60 199	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
AKERS 93D ZPHY C60 601	+Alexander, Allison+	(OPAL Collab.)
BUSKULIC 93J ZPHY C60 71	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
BUSKULIC 93L PL B313 520	+De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC 93M PL B313 535	+De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC 93N PL B313 549	+De Bonis, Decamp+	(ALEPH Collab.)
BUSKULIC 93P ZPHY C59 369	+Decamp, Goy+	(ALEPH Collab.)
LEP 93 PL B307 187	+LEP Energy Group, LEP Collabs	(LEP Collabs.)
QUAST 93 MPL A8 675		(DESY)

Searches for Higgs Bosons — H^0 and H^\pm

NOTE ON THE HIGGS BOSON

(by I. Hinchliffe, LBL)

The Standard Model [1] contains one neutral scalar Higgs boson, which is a remnant of the mechanism that breaks the $SU(2) \times U(1)$ symmetry and generates the W and Z boson masses. The Higgs couples to quarks and leptons of mass m_f with a strength $gm_f/2M_W$. Its coupling to W and Z bosons is of strength g , where g is the coupling constant of the $SU(2)$ gauge theory. Consequently its coupling to stable matter is very small, and its production and detection in experiments is difficult. An exception is its production in the decay of the Z boson. Since large numbers of Z 's can be produced and the coupling of the Z to the Higgs is unsuppressed, experiments

Gauge & Higgs Boson Full Listings

Higgs Bosons — H^0 and H^\pm

at LEP are now able to rule out a significant range of Higgs masses.

If the Higgs mass is very large, the couplings of the Higgs to itself and to longitudinally polarized gauge bosons become large. Requiring that these couplings remain weak enough so that perturbation theory is applicable implies that $M_H \lesssim 1$ TeV [2]. While this is not an absolute bound, it is an indication of the mass scale at which one can no longer speak of an elementary Higgs boson. This fact is made more clear if one notes that the width of the Higgs boson is proportional to the cube of its mass and that a boson of mass 1 TeV has a width of 500 GeV.

It is believed that scalar field theories of the type used to describe Higgs self-interactions can only be effective theories valid over a limited range of energies if the Higgs self-coupling and hence Higgs mass is nonzero. A theory of this type that is valid at all energy scales must have zero coupling. The range of energies over which the interacting theory is valid is a function of the Higgs self-coupling and hence its mass. An upper bound on the Higgs mass can then be determined by requiring that the theory be valid (*i.e.*, have a nonzero value of the renormalized Higgs self-coupling) at all scales up to the Higgs mass [3]. Non-perturbative calculations using lattice [4] gauge theory that can be used to compute at arbitrary values of the Higgs mass indicate that $M_H \lesssim 750$ GeV.

If the Higgs mass were small, then the vacuum (ground) state with the correct value of M_W would cease to be the true ground state of the theory [5]. A theoretical constraint can then be obtained from the requirement that this is not the case, *i.e.*, that the our universe is in the true minimum of the Higgs potential. The constraint can be parametrized approximately as [6]

$$M_H > 1.85(m_{\text{top}} - 85 \text{ GeV}) .$$

This constraint may be too restrictive. Strictly speaking we can only require that the predicted lifetime of our universe, if it is not at the true minimum of the Higgs potential, be longer than its observed age. This constraint can be approximated by [7,8]

$$M_H > 5.9(m_{\text{top}} - 170 \text{ GeV}) .$$

Experiments at LEP are able to exclude a large range of Higgs masses. They search for the decay $Z \rightarrow HZ^*$. Here Z^* refers to a virtual Z boson that can appear in the detector as e^+e^- , $\mu^+\mu^-$, $\tau^+\tau^-$, $\nu\bar{\nu}$ (*i.e.*, missing energy) or hadrons. The experimental searches have considered both $H \rightarrow$ hadrons and $H \rightarrow \tau^+\tau^-$. The best limits are shown in the listings below.

Precision measurement of electroweak parameters such as M_W and the various asymmetries at LEP and SLC are becoming sensitive enough that they can in principle constrain the Higgs mass through its effect in radiative corrections. Currently, the precision tests allow the entire range from the direct LEP limit ($M_H \gtrsim 60$ GeV) to 1 TeV [9]. However, a measurement of the top mass might enable a constraint on M_H to be extracted.

Extensions of the standard model, such as those based on supersymmetry [10], can have more complicated spectra of Higgs bosons. The simplest extension has two Higgs doublets whose neutral components have vacuum expectation values v_1 and v_2 , both of which contribute to the W and Z masses. The physical particle spectrum contains one charged Higgs boson (H^\pm), two neutral scalars (H_1^0, H_2^0), and one pseudoscalar (A^0) if CP is conserved in the scalar sector [11]. In the simplest version of the supersymmetric model, there is an upper bound on the mass of one of these scalars. The bound depends upon the top quark mass; for $m_t = 150$ GeV, the bound is $M_{H_1^0} \lesssim 110$ GeV [12]. In models where all fermions of the same electric charge receive their masses from only one of the two doublets (v_2 gives mass to the charge 2/3 quarks, while v_1 gives mass to the charged leptons and the charge 1/3 quarks), there are, as in the standard model, no flavor-changing neutral currents at lowest order in perturbation theory. The H_i^0 and A^0 couplings to fermions depend on v_2/v_1 and are either enhanced or suppressed relative to the couplings in the standard model. Experiments at LEP are able to exclude ranges of masses for neutral Higgs particles in these models. These ranges depend on the values of v_2/v_1 . See the listings below on H_1^0 , Mass Limits in Supersymmetric Models.

Searches for charged Higgs bosons depend on the assumed branching fractions to $\nu\tau$, $c\bar{s}$, and $c\bar{b}$. See the listings for H^\pm Mass Limit.

References

1. S. Weinberg, Phys. Rev. Lett. **19**, 1264 (1967); A. Salam, in "Elementary Particle Theory," W. Svartholm, ed., Almqvist and Wiksell, Stockholm (1968); S.L. Glashow, J. Iliopoulos, and L. Maiani, Phys. Rev. **D2**, 1285 (1970).
2. M. Veltman, Ann. Phys. (NY) **B8**, 475 (1977); B.W. Lee, C. Quigg, and H. Thacker, Phys. Rev. **D16**, 1519 (1977).
3. L. Maiani, G. Parisi, and R. Petronzio, Nucl. Phys. **B136**, 115 (1978); R. Dashen and H. Neuberger, Phys. Rev. Lett. **50**, 1897 (1983).
4. G. Bhanot and K. Bitar, Phys. Rev. Lett. **61**, 798 (1988); J. Kuti, L. Lin, and Y. Shen, Phys. Rev. Lett. **61**, 678 (1988); M. Gockeler, K. Jansen, and T. Neuhaus, Phys. Lett. **B273**, 450 (1991); U.M. Heller, H. Neuberger, and P. Vranas, Phys. Lett. **B283**, 355 (1992).
5. A.D. Linde, JETP Lett. **23**, 64 (1976) [Pis'ma Zh. Eksp. Teor. Fiz. **23**, 73 (1976)]; S. Weinberg, Phys. Rev. Lett. **36**, 294 (1976).
6. M. Lindner, M. Sher, and H.W. Zaglauer, Phys. Lett. **B228**, 527 (1988); M.J. Duncan, R. Phillippe, and M. Sher, Phys. Lett. **153B**, 165 (1985).
7. G. Anderson, Phys. Lett. **B243**, 265 (1990).
8. P.B. Arnold, Phys. Rev. **D40**, 613 (1989).
9. W. Hollik, presented at the XVI International Symposium on Lepton-Photon Interactions, (Cornell, Aug. 1993).

Gauge & Higgs Boson Full Listings

Higgs Bosons — H^0 and H^\pm

10. For a review of these models see, for example, I. Hinchliffe, *Ann. Rev. Nucl. and Part. Sci.* **36**, 505 (1986) and J.F. Gunion, H.E. Haber, G.L. Kane, and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Redwood City, CA, 1990).
11. J.F. Gunion and H.E. Haber, *Nucl. Phys.* **B278**, 449 (1986).
12. J. Ellis, G. Ridolfi, and F. Zwirner, *Phys. Lett.* **B257**, 83 (1991);
R. Barbieri, F. Caravaglios, and M. Frigeni, *Phys. Lett.* **B258**, 167 (1991);
P.H. Chankowski, S. Pokorski, and T. Rosiek, *Phys. Lett.* **B274**, 191 (1991);
Y. Okada, M. Yamaguchi, and T. Yanagida, *Prog. Theor. Phys.* **85**, 1 (1991);
H.E. Haber and R. Hempfling, *Phys. Rev. Lett.* **66**, 1815 (1991).

H^0 (Higgs Boson) MASS LIMITS

These limits apply to the Higgs boson of the three-generation Standard Model with the minimal Higgs sector. Limits that depend on the $Ht\bar{t}$ coupling may also apply to a Higgs boson of an extended Higgs sector whose couplings to up-type quarks are comparable to or larger than those of the standard one-doublet model H^0 couplings.

For comprehensive reviews, see Gunion, Haber, Kane, and Dawson, "The Higgs Hunter's Guide," (Addison-Wesley, Menlo Park, CA, 1990) and R.N. Cahn, *Reports on Progress in Physics* **52** 389 (1989). For a review of theoretical bounds on the Higgs mass, see M. Sher, *Physics Reports* (Physics Letters C) **179** 273 (1989).

Limits from Coupling to Z/W^\pm

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>58.4 (CL = 95%) OUR LIMIT				
>55.7	95	1 ABREU	94G DLPH	$Z \rightarrow H^0 Z^*$
>56.9	95	2 AKERS	94B OPAL	$Z \rightarrow H^0 Z^*$
>57.7	95	3 ADRIANI	93C L3	$Z \rightarrow H^0 Z^*$
>58.4	95	4 BUSKULIC	93H ALEP	$Z \rightarrow H^0 Z^*$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>60	95	5 GROSS	93 RVUE	$Z \rightarrow H^0 Z^*$
		6 ABREU	92D DLPH	$Z \rightarrow H^0 \gamma$
>38	95	7 ABREU	92I DLPH	$Z \rightarrow H^0 Z^*$
>52	95	8 ADEVA	92B L3	$Z \rightarrow H^0 Z^*$
		9 ADRIANI	92F L3	$Z \rightarrow H^0 \gamma$
>48	95	10 DECAMP	92 ALEP	$Z \rightarrow H^0 Z^*$
> 0.21	99	11 ABREU	91B DLPH	$Z \rightarrow H^0 Z^*$
>11.3	95	12 ACTON	91 OPAL	$H^0 \rightarrow \text{anything}$
>41.8	95	13 ADEVA	91 L3	$Z \rightarrow H^0 Z^*$
		14 ADEVA	91D L3	$Z \rightarrow H^0 \gamma$
none 3–44	95	15 AKRAWY	91 OPAL	$Z \rightarrow H^0 Z^*$
none 3–25.3	95	16 AKRAWY	91C OPAL	$Z \rightarrow H^0 Z^*$
none 0.21–0.818	90	17 ABE	90E CDF	$p\bar{p} \rightarrow (W^\pm, Z) + H^0 + X$
none 0.846–0.987	90	17 ABE	90E CDF	$p\bar{p} \rightarrow (W^\pm, Z) + H^0 + X$
none 0.21–14	95	18 ABREU	90C DLPH	$Z \rightarrow H^0 Z^*$
none 2–32	95	19 ADEVA	90H L3	$Z \rightarrow H^0 Z^*$
> 2	99	20 ADEVA	90N L3	$Z \rightarrow H^0 Z^*$
none 3.0–19.3	95	21 AKRAWY	90C OPAL	$Z \rightarrow H^0 Z^*$
> 0.21	95	22 AKRAWY	90P OPAL	$Z \rightarrow H^0 Z^*$
none 0.032–15	95	23 DECAMP	90 ALEP	$Z \rightarrow H^0 Z^*$
none 11–24	95	24 DECAMP	90H ALEP	$Z \rightarrow H^0 Z^*$
> 0.057	95	25 DECAMP	90M ALEP	$Z \rightarrow H^0 ee, H^0 \mu\mu$
none 11–41.6	95	26 DECAMP	90N ALEP	$Z \rightarrow H^0 Z^*$

- 1 ABREU 94G searched for $Z \rightarrow H^0 + (e^+e^-, \mu^+\mu^-, \tau^+\tau^-, \nu\bar{\nu})$ with $H^0 \rightarrow q\bar{q}$. Four $\ell^+\ell^-$ candidates were found (all yielding low mass) consistent with expected backgrounds.
- 2 AKERS 94B searched for $Z \rightarrow H^0 + (e^+e^-, \mu^+\mu^-, \nu\bar{\nu})$ with $H^0 \rightarrow q\bar{q}$. One $\nu\bar{\nu}$ and one $\mu^+\mu^-$ candidate were found consistent with expected backgrounds.
- 3 ADRIANI 93C searched for $Z \rightarrow H^0 + (\nu\bar{\nu}, e^+e^-, \mu^+\mu^-)$ with H^0 decaying hadronically or to $\tau\bar{\tau}$. Two e^+e^- and one $\mu^+\mu^-$ candidates are found consistent with expected background.
- 4 BUSKULIC 93H searched for $Z \rightarrow H^0\nu\bar{\nu}$ (acoplanar jets) and $Z \rightarrow H^0 + (e^+e^-, \mu^+\mu^-)$ (lepton pairs in hadronic events).
- 5 GROSS 93 combine data taken by four LEP experiments through 1991.
- 6 ABREU 92D give $\sigma(e^+e^- \rightarrow Z \rightarrow H^0\gamma)B(H^0 \rightarrow \text{hadrons}) < 8$ pb (95% CL) for $m_{H^0} < 75$ GeV and $E_\gamma > 8$ GeV.

- 7 ABREU 92J searched for $Z \rightarrow H^0 + (ee, \mu\mu, \tau\tau, \nu\bar{\nu})$ with $H^0 \rightarrow q\bar{q}$. Only one candidate was found, in the channel $ee + 2$ jets, with a dijet mass 35.4 ± 5 GeV/ c^2 , consistent with the expected background of 1.0 ± 0.2 events in the 3 channels $e^+e^-, \mu^+\mu^-, \tau^+\tau^-$, and of 2.8 ± 1.3 events in all 4 channels. This paper excludes 12–38 GeV. The range 0–12 GeV is eliminated by combining with the analyses of ABREU 90C and ABREU 91B.
- 8 ADEVA 92B searched for $Z \rightarrow H^0 + (\nu\bar{\nu}, ee, \mu\mu, \tau\tau)$ with $H^0 \rightarrow \text{anything}$, $Z \rightarrow H^0 + \tau\tau$ with $H^0 \rightarrow q\bar{q}$, and $Z \rightarrow H^0 + q\bar{q}$ with $H^0 \rightarrow \tau\tau$. The analysis excludes the range $30 < m_{H^0} < 52$ GeV.
- 9 ADRIANI 92F give $\sigma(e^+e^- \rightarrow Z \rightarrow H^0\gamma)B(H^0 \rightarrow \text{hadrons}) < (2-10)$ pb (95% CL) for $m_{H^0} = 25-85$ GeV. Using $\sigma(e^+e^- \rightarrow Z) = 30$ nb, we obtain $B(Z \rightarrow H^0\gamma)B(H^0 \rightarrow \text{hadrons}) < (0.7-3) \times 10^{-4}$ (95% CL).
- 10 DECAMP 92 searched for most possible final states for $Z \rightarrow H^0 Z^*$.
- 11 ABREU 91B searched for $Z \rightarrow H^0 + \ell\bar{\ell}$ with missing H^0 and $Z \rightarrow H^0 + (\nu\bar{\nu}, \ell\bar{\ell}, q\bar{q})$ with $H^0 \rightarrow ee$.
- 12 ACTON 91 searched for $e^+e^- \rightarrow Z^*H^0$ where $Z^* \rightarrow e^+e^-, \mu^+\mu^-, \text{or } \nu\bar{\nu}$ and $H^0 \rightarrow \text{anything}$. Without assuming the minimal Standard Model mass-lifetime relationship, the limit is $m_{H^0} > 9.5$ GeV.
- 13 ADEVA 91 searched for $Z \rightarrow H^0 + (\mu\mu, ee, \nu\bar{\nu})$. This paper only excludes $15 < m_{H^0} < 41.8$ GeV. The 0–15 GeV range is excluded by combining with the analyses of previous L3 papers.
- 14 ADEVA 91D obtain a limit $B(Z \rightarrow H^0\gamma)B(H^0 \rightarrow \text{hadrons}) < 4.7 \times 10^{-4}$ (95% CL) for $m_{H^0} = 30-86$ GeV. The limit is not sensitive enough to exclude a standard H^0 .
- 15 AKRAWY 91 searched for the channels $Z \rightarrow H^0 + (\nu\bar{\nu}, ee, \mu\mu, \tau\tau)$ with $H^0 \rightarrow q\bar{q}, \tau\tau$, and $Z \rightarrow H^0 q\bar{q}$ with $H^0 \rightarrow \tau\tau$.
- 16 AKRAWY 91C searched the decay channels $Z \rightarrow H^0 + (\nu\bar{\nu}, ee, \mu\mu)$ with $H^0 \rightarrow q\bar{q}$.
- 17 ABE 90E looked for associated production of H^0 with W^\pm or Z in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV. Searched for H^0 decays into $\mu^+\mu^-, \pi^+\pi^-, \text{and } K^+K^-$. Most of the excluded region is also excluded at 95% CL.
- 18 ABREU 90C searched for the channels $Z \rightarrow H^0 + (\nu\bar{\nu}, ee, \mu\mu)$ and $H^0 + q\bar{q}$ for $m_H < 1$ GeV.
- 19 ADEVA 90H searched for $Z \rightarrow H^0 + (\mu\mu, ee, \nu\bar{\nu})$.
- 20 ADEVA 90N looked for $Z \rightarrow H^0 + (ee, \mu\mu)$ with missing H^0 and with $H^0 \rightarrow ee, \mu\mu, \pi^+\pi^-, K^+K^-$.
- 21 AKRAWY 90C based on 825 nb^{-1} . The decay $Z \rightarrow H^0\nu\bar{\nu}$ with $H^0 \rightarrow \tau\bar{\tau}$ or $q\bar{q}$ provides the most powerful search means, but the quoted results sum all channels.
- 22 AKRAWY 90P looked for $Z \rightarrow H^0 + (ee, \mu\mu)$ (H^0 missing) and $Z \rightarrow H^0\nu\bar{\nu}, H^0 \rightarrow e^+e^-, \gamma\gamma$.
- 23 DECAMP 90 limits based on 11,550 Z events. They searched for $Z \rightarrow H^0 + (\nu\bar{\nu}, ee, \mu\mu, \tau\tau, q\bar{q})$. The decay $Z \rightarrow H^0\nu\bar{\nu}$ provides the most powerful search means, but the quoted results sum all channels. Different analysis methods are used for $m_{H^0} < 2m_\mu$ where Higgs would be long-lived. The 99% confidence limits exclude $m_{H^0} = 0.040-12$ GeV.
- 24 DECAMP 90H limits based on 25,000 Z \rightarrow hadron events.
- 25 DECAMP 90M looked for $Z \rightarrow H^0\ell\bar{\ell}$, where H^0 decays outside the detector.
- 26 DECAMP 90N searched for the channels $Z \rightarrow H^0 + (\nu\bar{\nu}, ee, \mu\mu, \tau\tau)$ with $H^0 \rightarrow (\text{hadrons}, \tau\tau)$.

Limits from Other Techniques

H^0 Indirect Mass Limits from Electroweak Analysis

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
35 +205 - 26		27 ELLIS	94 RVUE	
73 +178 - 13		28 BLONDEL	93 RVUE	
10 +25 - 8		29 ELLIS	93B RVUE	
10 +60 - 8		30 NOVIKOV	93B RVUE	
		31 DELAGUILA	92B RVUE	
> 1.4	68	32 ELLIS	92 RVUE	Electroweak
25 +275 - 19		33 ELLIS	92E RVUE	
50 +353 - 0		34 RENTON	92 RVUE	
		35 SCHAILE	92 RVUE	

- 27 ELLIS 94 fit to LEP, SLD, W-mass, and neutrino data available in the summer of 1993. The fit to m_H, m_t , and α_S yields $m_t = 140^{+21}_{-22}$ GeV and $\alpha_S(m_Z) = 0.116^{+0.007}_{-0.006}$.
- 28 BLONDEL 93 perform two dimensional ($m_t - m_H$) fit to LEP electroweak data available in the spring of 1993. $\alpha_S = 0.117 \pm 0.005$ is used and $m_t > 108$ GeV, $m_H > 62.5$ GeV imposed. $m_{H^0} = 1$ TeV is compatible with the data within two standard deviations.
- 29 ELLIS 93B fit to LEP and neutrino data available in the summer of 1993. m_t is adjusted to minimize χ^2 and $\alpha_S(m_Z) = 0.123 \pm 0.006$ is used. 95% CL limit for $m_H < 250$ GeV is claimed.
- 30 NOVIKOV 93B use a subset of the most accurate and "gluon-free" data available in the spring of 1993. They use $m_W, \Gamma(\ell\ell)$, and $A_{FB}^{\ell\ell}$.
- 31 DELAGUILA 92B perform two dimensional ($m_t - m_H$) fit to various LEP, neutrino, eH , and $p\bar{p}$ data available through 1991 with direct limits on m_t, m_H . The result $m_H = 65^{+245}_{-4}$ is not expected from the statistical sensitivity of the data but due to deviation of the data from the Standard Model expectation.
- 32 ELLIS 92 result is from a fit to electroweak data from LEP and elsewhere. They also find $m_H < 160$ GeV at 68%CL and $0.5 < m_H < 1500$ GeV at 90%CL with m_t unconstrained.
- 33 ELLIS 92E perform fit to electroweak data available in the spring of 1992. m_t is adjusted to minimize χ^2 and $\alpha_S(m_Z) = 0.118 \pm 0.008$ is used.
- 34 RENTON 92 use electroweak data available in 1991 and require $m_H > 50$ GeV. The constraint $\alpha_S = 0.114 \pm 0.007$ was used.

Gauge & Higgs Boson Full Listings

Higgs Bosons — H^0 and H^\pm

³⁵SCHALE 92 performs fit to LEP electroweak data (as of summer 1991) as well as m_W (UA2/CDF) and νN (CDHS/CHARM). The fit with the constraint $m_H > 50$ GeV gives $m_H = 50^{+192}_{-0}$ GeV. However, the m_H dependence of the χ^2 is not consistent from that expected from the present statistics and the sensitivity to m_H arises from the fact that the measured values of g_A and A_{FB}^b deviate from the Standard Model expectation. Therefore, the result is not considered to be significant.

From Quarkonium Decay

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>0.086	90	36 ANTREASYAN 90C CBAL	$\Upsilon(1S) \rightarrow H^0 \gamma$	
none 0.29–0.57	90	37 ALBRECHT 89 ARG	$\Upsilon(1S) \rightarrow H^0 \gamma$ ($H^0 \rightarrow \pi^+ \pi^-$)	
none 0.21–5	90	38 LEE-FRANZINI 88 CUSB	$\Upsilon(1S, 3S) \rightarrow \gamma H^0$	
		39 DRUZHININ 87 ND	$\phi \rightarrow \gamma H^0$ ($H^0 \rightarrow \pi^0 \pi^0$)	
³⁶ ANTREASYAN 90C obtain $B(\Upsilon(1S) \rightarrow H^0 \gamma) < 3.5 \times 10^{-5}$ at 90% CL for $m_{H^0} < 2m_e$ and similar limits for heavier Higgs masses. The listed limit assumes the QCD/relativistic reduction factor for the width of 0.5. The limit is reduced to 39 MeV if 0.25 is used instead.				
³⁷ ALBRECHT 89 give a limit $B(\Upsilon(1S) \rightarrow H^0 \gamma) B(H^0 \rightarrow \pi^+ \pi^-) < 3.45 \times 10^{-5}$ for $m_{H^0} = 290$ –570 MeV, which is lower than the prediction including first order QCD corrections and assuming $B(H^0 \rightarrow \pi^+ \pi^-) > 45\%$.				
³⁸ LEE-FRANZINI 88 presents updated results from the CUSB experiment (see FRANZINI 87 for more details). First order QCD correction included with $\alpha_s \approx 0.2$ ($\Lambda = 0.2$ GeV and $n(f) = 4$). The order α_s correction reduced the rate for $\Upsilon(1S) \rightarrow H^0 \gamma$ by a factor of 2 (yielding these limits). The impact of order α_s^2 and of relativistic corrections are unknown. If they amounted to another factor of 2 suppression, the above limit would be essentially eliminated.				
³⁹ DRUZHININ 87 sets limit $B(\phi \rightarrow \gamma H^0) B(H^0 \rightarrow \pi^0 \pi^0) < 8 \times 10^{-5}$ at CL=90% for $m_{H^0} = 0.6$ –1 GeV which is still far from the standard Higgs model prediction and does not exclude the existence of light Higgs bosons.				

From B Decay

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 0.21–3.57		40 DAWSON 90 RVUE	$B \rightarrow \mu^+ \mu^- X$; $B \rightarrow K(\mu^+ \mu^-)$, $\pi^+ \pi^-$, $K^+ K^-$	
none 0.21–1.0	90	41 ALAM 89B CLEO	$B \rightarrow H^0 K$, ($H^0 \rightarrow \mu^+ \mu^-$, $\pi^+ \pi^-$)	
none 1.0–3.6	90	41 ALAM 89B CLEO	$B \rightarrow H^0 X$ ($H^0 \rightarrow \mu^+ \mu^-$)	
none 3.6–4.6		42 EILAM 89 RVUE	$B \rightarrow H^0 X$ ($H^0 \rightarrow \mu^+ \mu^-$)	
none 0.211–0.700		43 RABY 89 RVUE	$B \rightarrow \mu^+ \mu^- X$ $m_{\text{top}} > 80$ GeV	
none 0.07–0.21	90	44 SNYDER 89 MRK2	$B \rightarrow H^0 X$ ($H^0 \rightarrow e^+ e^-$)	
none 0.00103–3.57		43 CHIVUKULA 88 RVUE	$B \rightarrow H^0 X$, $m(\text{top}) > 80$ GeV	
none 2–3.7		43 GRINSTEIN 88 RVUE	$B \rightarrow H^0 X$, $m(\text{top}) > 80$ GeV	

⁴⁰Based on ALTHOFF 84G, ALAM 89B, and ALBRECHT 87D. Some processes considered require the assumption $B(B \rightarrow H^0 K)/B(B \rightarrow H^0 X) > 0.01$. Other processes require theoretical assumptions regarding $B(H \rightarrow \pi^+ \pi^-)$ when considering masses in the interval 0.9–1.2 GeV.

⁴¹ALAM 89B searched for inclusive and exclusive decays of B mesons into H^0 and can exclude the mass range $2m_\mu$ – $2m_\tau$ with a wide margin provided $m_t \gtrsim m_W$, possibly except for masses near $\chi_0(3410)$, where the mixing effect can reduce $B(H^0 \rightarrow \mu^+ \mu^-)$ significantly.

⁴²EILAM 89 assume $m_{\text{top}} > 90$ GeV and vary $|V_{ub}/V_{cb}|^2$ from 0 to 0.026.

⁴³Limits assume $m(\text{top}) > 80$ GeV and $|V_{ts} V_{tb}^* / V_{cb}| \approx 1$. CHIVUKULA 88 excludes m_{H^0} between $2m_e$ and $2m_\tau$ from the limits on $B \rightarrow \mu^+ \mu^- + X$ by taking the $B(H^0 \rightarrow \mu^+ \mu^-)$ estimate of VOLOSHIN 86. GRINSTEIN 88 argues that this estimate of VOLOSHIN 86 is unreliable, and excludes m_{H^0} between 2 GeV and 3.7 GeV where perturbative QCD is used to estimate $B(H^0 \rightarrow \mu^+ \mu^-)$.

⁴⁴SNYDER 89 exclude the mass range 70–210 MeV with a wide margin provided that $m_t \gtrsim m_W$. A limit $B(B \rightarrow H^0 X) B(H^0 \rightarrow e^+ e^-) < 22\%$ (90% CL) is given for $m_{H^0} = 50$ MeV.

From K Decay

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>0.026	90	45 ATIYA 90 CNTR	$K^+ \rightarrow \pi^+ H^0$	
		46 ATIYA 90B CNTR	$K^+ \rightarrow \pi^+ H^0$, $H^0 \rightarrow \gamma \gamma$	
none 0.012–0.211	90	47 BARR 90 NA31	$K_L^0 \rightarrow \pi^0 H^0$ ($H^0 \rightarrow e^+ e^-$)	
>0.32		48 DAWSON 90 RVUE	K decays	
>0.3		49 LEUTWYLER 90 RVUE	$K^+ \rightarrow \pi^+ H^0$	

none 0.22–0.32	50 ATIYA 89 CNTR	$K^+ \rightarrow \pi^+ H^0$ ($H^0 \rightarrow \mu^+ \mu^-$)	
>0.28	51 CHENG 89 RVUE	$K^\pm \rightarrow \pi^\pm H$	
>0.36	52 CHIVUKULA 88 RVUE	$K \rightarrow \pi^+ H^0$	
	90	53 BAKER 87 CALO	$K^\pm \rightarrow \pi^\pm H^0$ ($H^0 \rightarrow e^+ e^-$)
none 0.05–0.211	54 WILLEY 86 RVUE	$K^\pm \rightarrow \pi^\pm H^0$ ($H^0 \rightarrow e^+ e^-$)	
⁴⁵ ATIYA 90 sets limits on $B(K^+ \rightarrow \pi^+ H^0)$ varying from $< 6.4 \times 10^{-9}$ for $m_{H^0} \approx 0$ MeV to $< 10^{-6}$ for $m_{H^0} = 26$ MeV.			
⁴⁶ ATIYA 90B give 90% CL limits on $B(K^+ \rightarrow \pi^+ H^0) B(H^0 \rightarrow \gamma \gamma)$ for $m_{H^0} < 100$ MeV ranging from 10^{-4} to 10^{-7} depending on the mass.			
⁴⁷ BARR 90 set m_{H^0} -dependent limits on $B(K_L^0 \rightarrow \pi^0 H^0)$ in the region where $B(H^0 \rightarrow e^+ e^-) \approx 1$. The limit varies from $B(K_L^0 \rightarrow \pi^0 H^0) < 10^{-7}$ at $m_{H^0} = 12$ MeV to $< 2 \times 10^{-8}$ for $50 \leq m_{H^0} \leq 211$ MeV. BARR 90 allow for nonzero H^0 lifetime.			
⁴⁸ Based on ASANO 81B, YAMAZAKI 84, BAKER 87, ATIYA 89, and BARR 90. DAWSON 90 use theoretical calculations and various assumptions such as $m_t > 80$ GeV and $\text{Im } V_{td}^* V_{ts} > 0.2 \sin^2 \theta_c$.			
⁴⁹ LEUTWYLER 90 give a consistent analysis of the $K \rightarrow \pi H^0$ amplitude based on chiral theory and find that all contributions except the t -quark loop are unimportant numerically provided the t -quark mass is of order or bigger than 100 GeV. Hence, a light Higgs can probably be ruled out.			
⁵⁰ ATIYA 89 give a limit $B(K^+ \rightarrow \pi^+ H^0) B(H^0 \rightarrow \mu^+ \mu^-) < 1.5 \times 10^{-7}$ (90% CL) for $m_{H^0} = 220$ –320 MeV, which is lower than the prediction unless there is an accidental cancellation in the CP -conserving part of the amplitude and the CP -violating part is unexpectedly small. See WILLEY 89 and CHENG 89.			
⁵¹ CHENG 89 concludes even if real part of $K^+ \rightarrow \pi^+ H$ amplitude is cancelled accidentally, the imaginary contribution alone rules out $m_H < 2m_\pi$.			
⁵² CHIVUKULA 88 uses chiral perturbation theory to estimate $K \rightarrow \pi^+ H^0$ amplitudes with a conservative sign assignment for the relative sign of the $\Delta I = 1/2$ term, and exclude m_{H^0} below 0.36 GeV barring cancellation among terms, by using the limits on $K \rightarrow \pi^+ X$ with $X = \mu^+ \mu^-$, $e^+ e^-$, or missing particles. For a criticism see DAWSON 90.			
⁵³ BAKER 87 sets limit $B(K^\pm \rightarrow \pi^\pm H^0) B(H^0 \rightarrow e^+ e^-) < 8 \times 10^{-7}$ at CL=90% for $m_{H^0} < 100$ MeV if H^0 travels much less than 1.4 cm in the lab frame ($p(K^\pm) \approx 5.8$ GeV). The expected lifetime of the standard H^0 is too long to be effectively detected by the experiment and their limit on the branching ratio is significantly weakened accordingly. In view of the uncertainty in the theoretical prediction for $B(K \rightarrow \pi H)$, no definite conclusion can be drawn from the result. See also DAWSON 90.			
⁵⁴ WILLEY 86 re-examined the theoretical estimate of the decay $K^\pm \rightarrow \pi^\pm H^0$ rate via the one-loop $s d H^0$ coupling. The experimental bound $B(K \rightarrow \pi \mu \mu) < 2.4 \times 10^{-6}$ is not strong enough to rule out $2m_\mu < m_{H^0} < 2m_{\pi^0}$. For a criticism see DAWSON 90.			

From Coupling with Nucleons

Some of the experiments for a light Higgs utilize its coupling with nucleons. We parameterize the Higgs-nucleon coupling (which is dominantly isoscalar) as $g_{HNN} = \eta_{HNN}(\sqrt{2}G_F)^{1/2} m_N$. The limits depend on the value of η_{HNN} used. Shifman *et al.* [Physics Letters **78B** 443 (1978)] obtained $\eta_{HNN} = 0.22$ assuming three heavy flavors. More recently, T.P. Cheng [Physical Review **D38** 2869 (1988)], H.-Y. Cheng [Physics Letters **B219** 347 (1989)], and Barbieri and Curci [Physics Letters **B219** 503 (1989)] took into account the strange-quark content of the proton as well as the heavy quark effects, and derived $\eta_{HNN} \approx 0.56$.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>0.0009	95	55 BLUEMLEIN 92 BDMP	$pN \rightarrow H^0 X$, $H^0 Z \rightarrow e^+ e^- Z$ ($\ell = e, \mu$)	
>0.0128	90	56 LEEB 92 RVUE	$nN \rightarrow nN$	
none 0.001–0.08	95	BLUEMLEIN 91 BDMP	$pN \rightarrow H^0 X$ ($H^0 \rightarrow e^+ e^-$, 2γ)	
>0.018		57 GRIFOLS 89 RVUE	$\sigma_{\text{total}}(n\text{Pb})$	
none 0.03–0.20		58 YEPES 89B RVUE	$pN \rightarrow H^0 X$ ($H^0 \rightarrow e^+ e^-$)	
>0.010	68	59 BELTRAMI 86 SPEC	Muonic atoms	
none 0.003–0.012	95	60 FREEDMAN 84 CNTR	$\text{He}^+ \rightarrow \text{He} H^0$ ($H^0 \rightarrow e^+ e^-$)	
none 0.00103–0.00584		61 MUKHOPAD... 84 RVUE	$O^* \rightarrow O H^0$ ($H^0 \rightarrow e^+ e^-$)	
		62 HOFFMAN 83 CNTR	$\pi p \rightarrow n H^0$ ($H^0 \rightarrow e^+ e^-$)	
>0.006		63 BARBIERI 75 RVUE	$nN \rightarrow nN$	
⁵⁵ BLUEMLEIN 92 exclude very light Standard Model Higgs using bremsstrahlung Higgs production from beam dump followed by Bethe-Heitler production of lepton pairs by H^0 on nuclei. If combined with BLUEMLEIN 91, the range $m_H < 0.08$ GeV is excluded.				
⁵⁶ LEEB 92 use the neutron-lead total cross section measurement by SCHMIEDMAYER 91 as well as neutron optics data to obtain bounds on new long-range neutron interactions. The limit uses $\eta_{HNN} = 0.56$.				
⁵⁷ GRIFOLS 89 use the neutron-lead total cross-section measurement at kinetic energies of 50 eV – 50 keV by SCHMIEDMAYER 88 and argue that the agreement of the measured energy dependence with the prediction of a hard-core potential is lost by light-Higgs exchange. The limit of 18 MeV is obtained for $\eta_{HNN} = 0.56$ and is reduced to 12 MeV for $\eta_{HNN} = 0.22$. LEEB 92 argue against the use of a hard-core potential and obtain a weaker limit.				
⁵⁸ YEPES 89B reanalyzed a Fermilab experiment (BECHIS 78), which looked for a long-lived neutral lepton and found none, and argues that their limit is many orders of magnitude lower than expected from low-mass Higgs bremsstrahlung production followed by the decay to $e^+ e^-$.				

Gauge & Higgs Boson Full Listings

Higgs Bosons — H^0 and H^\pm

- ⁵⁹ BELTRAMI 86 measured the wavelengths of the $3d_{5/2}-2p_{3/2}$ X-ray transitions in muonic ^{24}Mg and ^{28}Si and found the deviation from QED $\delta\lambda/\lambda = (-0.2 \pm 3.1) \times 10^{-6}$. The listed limit uses $\eta_{HNN} = 0.23$. The experiment excludes $m_{H^0} \lesssim 1$ MeV by more than 3 s.d.
- ⁶⁰ FREEDMAN 84 is ANL experiment with dynamitron proton bombarding tritium to form He^* . $\eta_{HNN} = 0.30$ is used to derive the limit. They also reanalyze KOHLER 74 He^* data to find no mass region is excluded by that data. See also footnote for MUKHOPADHYAY 84 below.
- ⁶¹ MUKHOPADHYAY 84 examine KOHLER 74 He^* and C^* data. Claim that no mass region can be excluded by 74 He^* data since He^* decay width to proton is large [$B(\text{He}^* \rightarrow H^0\text{He}) = 3.4 \times 10^{-11}$ is very small]. Above limit is from KOHLER 74 O^* decay data.
- ⁶² HOFFMAN 83 looked for e^+e^- peak from Higgs produced in $\pi^-p \rightarrow H^0n$ at 300 MeV/c. Set CL = 90% limit $d\sigma/dt B(e^+e^-) < 3.5 \times 10^{-32} \text{ cm}^2/\text{GeV}^2$ for $140 < m_{H^0} < 160$ MeV, which does not exclude H^0 with the standard one-doublet-model couplings.
- ⁶³ BARBIERI 75 studied Higgs boson exchange effect in neutron-lead scattering data of ALEKSANDROV 66 and found limit $(g_{H^0nn}^2/4\pi) (m_{H^0}/\text{MeV})^{-4} \lesssim 3.4 \times 10^{-11}$ for $m_{H^0} \gtrsim 1$ MeV. This gives the listed limit for $\eta_{HNN} = 0.2$ and 10 MeV for $\eta_{HNN} = 0.56$. Lighter mass region $m_{H^0} \lesssim 1$ MeV would be incompatible with the measured angular distribution.

From Other Techniques

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 0.001-0.072	95	64 BARABASH	92 BDMP	$\eta' \rightarrow \eta H^0$
none 0.0012-0.052	90	DAVIER	89 BDMP	$e^-Z \rightarrow eH^0Z$ ($H^0 \rightarrow e^+e^-$)
none 0.010-0.10	90	65 EGLI	89 CNTR	$\pi^+ \rightarrow e^+\nu H^0$ ($H^0 \rightarrow e^+e^-$)
none 0.015-0.04	90	66 LINDNER	89 THEO	Vacuum stability
		67 YEPES	89 RVUE	$\pi^\pm \rightarrow e^\pm\nu H^0$ ($H^0 \rightarrow e^+e^-$)
		68 DZHELADIN	81	$\eta' \rightarrow \eta H^0$ ($H^0 \rightarrow \mu^+\mu^-$)
		69 WITTEN	81 COSM	
		69 GUTH	80 COSM	
		69 SHER	80 COSM	

- ⁶⁴ BARABASH 92 is a beam dump experiment that searched for $H^0 \rightarrow e^+e^-$ and $\gamma\gamma$ produced via the decays $\pi \rightarrow e\nu_e H^0$, $K \rightarrow e\nu_e H^0$, $K \rightarrow \pi H^0$, and $\eta' \rightarrow \eta H^0$. The last process gives the best limit if the theoretical calculation by RUSKOV 87 is used.
- ⁶⁵ EGLI 89 give a limit for $B(\pi^+ \rightarrow e^+\nu H^0) \cdot B(H^0 \rightarrow e^+e^-)$ ranging from 10^{-9} to 10^{-11} for the mass range 10-110 MeV. The theoretical prediction they use is too large by a factor of 162/49 (see DAWSON 89, DAWSON 90, and CHENG 89). The lower limit given above is reevaluated by us.
- ⁶⁶ LINDNER 89 require vacuum stability and numerically solve the renormalization equations to two-loop order. If $m_{\text{top}} = 100, 110, 120$ GeV, then $m_{\text{Higgs}} > 20, 34, 50$ GeV. However, it is possible that the vacuum is not stable but is very long-lived.
- ⁶⁷ YEPES 89 reanalyzed a BNL beam-dump experiment (JACQUES 80) which looked for electron pairs in 7 foot BC downstream from the dump and found none.
- ⁶⁸ DZHELADIN 81 obtained $B(\eta' \rightarrow \eta\mu^+\mu^-) < 1.5 \times 10^{-5}$ (CL = 90%), and argued that it excludes H^0 with the standard one-doublet-model couplings in $\mu^+\mu^-$ channel for $m_{H^0} = 0.25-0.409$ GeV. However, the number 0.409 is not well-determined due to theoretical uncertainties in $B(H^0 \rightarrow \mu^+\mu^-)$.
- ⁶⁹ Limits from cosmological considerations of $\text{SU}(2) \times \text{U}(1)$ symmetry-breaking phase transition occurring only after extreme supercooling, resulting in too high a ratio of entropy to baryon number. Limits apply to the standard one-doublet model H^0 , with 'zero bare mass' whose physical mass is determined by the Coleman-Weinberg mechanism of dynamical symmetry breakdown. These limits depend on the mass of the top quark approximately according to $m_{H^0} > 10.4[1 - 4m_t^4/(2m_W^4 + m_t^4)]^{1/2}$ GeV when $m_t < 80$ GeV. So for $m_t \approx 80$ GeV, there is no limit. If $m_t > 80$ GeV, then vacuum stability arguments may give bounds on m_H ; see LINDNER 89 above.

H^0 (Higgs Boson) MASS LIMITS in Extended Higgs Models

The parameter x denotes the Higgs coupling to charge $-1/3$ quarks and charged leptons relative to the value in the standard one-Higgs-doublet model.

In order to prevent flavor-changing neutral currents in models with more than one Higgs doublet, only one of the Higgs doublets can couple to quarks of charge 2/3. The same requirement applies independently to charge $-1/3$ quarks and to leptons. Higgs couplings can be enhanced or suppressed.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		70 BRAHMACH...	93 RVUE	
>65	95	71 BUSKULIC	93I ALEP	$Z \rightarrow H^0 Z^*$
		72 BUSKULIC	93I ALEP	Invisible H^0
		73 LOPEZ-FERN.	93 RVUE	
		74 ADRIANI	92G L3	$Z \rightarrow H^0 Z^*$

- > 3.57
- 95
- > 0.21
- none 0.6-6.2
- none 0.6-7.9
- none 3.7-5.6
- none 3.7-8.2
- 75 PICH
- 76 ACTON
- 77 DECAMP
- 78 DECAMP
- 79 AKRAWY
- 80 DAVIER
- 81 SNYDER
- 82 FRANZINI
- 82 FRANZINI
- 83 ALBRECHT
- 83 ALBRECHT
- 92 RVUE Very light Higgs
- 91 OPAL $Z \rightarrow H^0 Z^*$
- 91F ALEP $Z \rightarrow H^0 \ell^+ \ell^-$
- 91I ALEP Z decay
- 90P OPAL $Z \rightarrow H^0 Z^*$
- 89 BDMP $e^-Z \rightarrow eH^0Z$
($H^0 \rightarrow e^+e^-$)
- 89 MRK2 $B \rightarrow H^0 X$
($H^0 \rightarrow e^+e^-$)
- 87 CUSB $T(1S) \rightarrow \gamma H^0, x=2$
- 87 CUSB $T(1S) \rightarrow \gamma H^0, x=4$
- 85J ARG $T(1S) \rightarrow \gamma H^0, x=2$
- 85J ARG $T(1S) \rightarrow \gamma H^0, x=4$
- ⁷⁰ BRAHMACHARI 93 consider Higgs limit from Z decay when the Higgs decays to invisible modes. If H^0 coupling to Z is at least $1/\sqrt{2}$ of the Standard Model H^0 , the DECAMP 92 limit of 48 GeV changes within ± 6 GeV for arbitrary $B(H^0 \rightarrow \text{SM-like}) + B(H^0 \rightarrow \text{invisible}) = 1$.
- ⁷¹ See Fig. 1 of BUSKULIC 93i for the limit on $ZZ H^0$ coupling for a general Higgs having a similar decay signature to Standard Model Higgs boson or decaying invisibly. If the decay rate for $Z \rightarrow H^0 Z^*$ is $>10\%$ of the minimal Standard Model rate, then $m_{H^0} > 40$ GeV. For the standard rate the limit is 58 GeV.
- ⁷² BUSKULIC 93i limit for H^0 with the standard coupling to Z but decaying to weakly interacting particles.
- ⁷³ LOPEZ-FERNANDEZ 93 consider Higgs limit from Z decay when the Higgs decays to invisible modes. See Fig. 2 for excluded region in $m_{H^0} - ZZH$ coupling plane with arbitrary $B(H^0 \rightarrow \text{SM-like}) + B(H^0 \rightarrow \text{invisible}) = 1$. $m_H > 50$ GeV is obtained if the H^0 coupling strength to the Z is greater than 0.2 times the Standard Model rate.
- ⁷⁴ See Fig. 1 of ADRIANI 92G for the limit on $ZZ H^0$ coupling for a general Higgs having a similar decay signature to Standard Model Higgs boson. For most masses below 30 GeV, the rate for $Z \rightarrow H^0 Z^*$ is less than 10% of the Standard Model rate.
- ⁷⁵ PICH 92 analyse H^0 with $m_{H^0} < 2m_\mu$ in general two-doublet models. Excluded regions in the space of mass-mixing angles from LEP, beam dump, and π^\pm, η rare decays are shown in Figs. 3,4. The considered mass region is not totally excluded.
- ⁷⁶ ACTON 91 limit is valid for any H^0 having $\Gamma(Z \rightarrow H^0 Z^*)$ more than 0.24 (0.56) times that for the standard Higgs boson for Higgs masses below $2m_\mu$ ($2m_\tau$).
- ⁷⁷ DECAMP 91F search for $Z \rightarrow H^0 \ell^+ \ell^-$ where H^0 escapes before decaying. Combining this with DECAMP 90M and DECAMP 90N, they obtain $B(Z \rightarrow H^0 \ell^+ \ell^-) / B(Z \rightarrow \ell^+ \ell^-) < 2.5 \times 10^{-3}$ (95%CL) for $m_{H^0} < 60$ GeV.
- ⁷⁸ See Figs. 1, 3, 4, 5 of DECAMP 91I for excluded regions for the masses and mixing angles in general two-doublet models.
- ⁷⁹ AKRAWY 90P limit is valid for any H^0 having $\Gamma(Z \rightarrow H^0 Z^*)$ more than 0.57 times that for the Standard Higgs boson.
- ⁸⁰ DAVIER 89 give excluded region in $m_{H^0} - x$ plane for m_{H^0} ranging from 1.2 MeV to 50 MeV.
- ⁸¹ SNYDER 89 give limits on $B(B \rightarrow H^0 X) \cdot B(H^0 \rightarrow e^+e^-)$ for $100 < m_{H^0} < 200$ MeV, $x < 24$ mm.
- ⁸² First order QCD correction included with $\alpha_s \approx 0.2$. Their figure 4 shows the limits vs. x .
- ⁸³ ALBRECHT 85J found no mono-energetic photons in both $T(1S)$ and $T(2S)$ radiative decays in the range $0.5 \text{ GeV} < E(\gamma) < 4.0 \text{ GeV}$ with typically $\text{BR} < 0.01$ for $T(1S)$ and $\text{BR} < 0.02$ for $T(2S)$ at 90% CL. These upper limits are 5-10 times the prediction of the standard Higgs-doublet model. The quoted 90% limit $B(T(1S) \rightarrow H^0 \gamma) < 1.5 \times 10^{-3}$ at $E(\gamma) = 1.07$ GeV contradicts previous Crystal Ball observation of $(4.7 \pm 1.1) \times 10^{-3}$; see their reference 3. Their figure 8a shows the upper limits of x^2 as a function of $E(\gamma)$ by assuming no QCD corrections. We used $m_{H^0} = m_\gamma (1 - 2E(\gamma)/m_\gamma)^{1/2}$.

H_1^0 (Higgs Boson) MASS LIMITS in Supersymmetric Models

The minimal supersymmetric model has two complex doublets of Higgs bosons. The resulting physical states are two scalars [H_1^0 and H_2^0 , where we define $m_{H_1^0} < m_{H_2^0}$],

a pseudoscalar (A^0), and a charged Higgs pair (H^\pm). There are two free parameters in the theory which can be chosen to be m_{A^0} and $\tan\beta = \nu_2/\nu_1$, the ratio of vacuum expectation values of the two Higgs doublets. Tree-level Higgs masses are constrained by the model to be $m_{H_1^0} \leq m_Z, m_{H_2^0} \geq m_Z, m_{A^0} \geq m_{H_1^0}$, and $m_{H^\pm} \geq m_W$. However, as describe in the "Note on Supersymmetry," recent calculations of one-loop radiative corrections show that these relations may be violated. Many experimental analyses have not taken into account these corrections; footnotes indicate when these corrections are included. The results assume no invisible H^0 or A^0 decays.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>44	95	84 BUSKULIC	93I ALEP	$\tan\beta > 1$
>29	95	85 ABREU	92J DLPH	any $\tan\beta$
>42	95	86 ADRIANI	92G L3	$1 < \tan\beta < 50$
none 3-22	95	87 AKRAWY	91C OPAL	$\tan\beta > 0.5$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>34	95	85 ABREU	92J DLPH	$\tan\beta > 0.6$
> 0.21	95	88 ABREU	91B DLPH	any $\tan\beta$
>28	95	89 ABREU	91B DLPH	any $\tan\beta$
none 3-38	95	87 AKRAWY	91C OPAL	$\tan\beta > 6$
		90 BLUEMLEIN	91 BDMP	$pN \rightarrow H_1^0 X$ ($H_1^0 \rightarrow e^+e^-, 2\gamma$)
>41	95	91 DECAMP	91I ALEP	$\tan\beta > 1$
> 9	95	92 ABREU	90E DLPH	any $\tan\beta$
>13	95	92 ABREU	90E DLPH	$\tan\beta > 1$

Gauge & Higgs Boson Full Listings

Higgs Bosons — H^0 and H^\pm

>26	95	93 ADEVA	90R L3	$\tan\beta > 1$
none 0.05–3.1	95	94 DECAMP	90E ALEP	any $\tan\beta$
none 0.05–13	95	94 DECAMP	90E ALEP	$\tan\beta > 0.6$
none 0.006–20	95	94 DECAMP	90E ALEP	$\tan\beta > 2$
>37.1	95	94 DECAMP	90E ALEP	$\tan\beta > 6$
none 0.05–20	95	95 DECAMP	90H ALEP	$\tan\beta > 0.6$
none 0.006–21.4	95	95 DECAMP	90H ALEP	$\tan\beta > 2$
> 3.1	95	96 DECAMP	90M ALEP	any $\tan\beta$

84 BUSKULIC 93i search for $Z \rightarrow H_1^0 Z^*$ and $Z \rightarrow H_1^0 A^0$. One-loop corrections are included with any $m_t, m_{\bar{t}} > m_t$. Assumes no invisible H^0 or A^0 decays.

85 ABREU 92J searched for $Z \rightarrow H_1^0 Z^*$ and $Z \rightarrow H_1^0 A^0$ with $H_1^0, A^0 \rightarrow \tau\tau$ or jet-jet. Small mass values are excluded by ABREU 91B.

86 ADRIANI 92G search for $Z \rightarrow H_1^0 Z^*, Z \rightarrow H_1^0 A^0 \rightarrow 4b, bb\tau\tau, 4\tau, 6b$ (via $H^0 \rightarrow A^0 A^0$), and include constraints from $\Gamma(Z)$. One-loop corrections to the Higgs potential are included with $90 < m_t < 250$ GeV, $m_{\bar{t}} < m_{\bar{t}} < 1$ TeV.

87 AKRAWY 91C result from $Z \rightarrow H_1^0 A^0 \rightarrow 4\text{jet or } \tau^+ \tau^- jj$ or 4τ and $Z \rightarrow H_1^0 Z^*$ ($H_1^0 \rightarrow q\bar{q}, Z^* \rightarrow \nu\bar{\nu}$ or $e^+ e^-$ or $\mu^+ \mu^-$). See paper for the excluded region for the case $\tan\beta < 1$. Although these limits do not take into account the one-loop radiative corrections, the authors have reported unpublished results including these corrections and showed that the excluded region becomes larger.

88 ABREU 91B result is based on negative search for $Z \rightarrow H_1^0 f\bar{f}$ and the limit on invisible Z width $\Gamma(Z \rightarrow H_1^0 A^0) < 39$ MeV (95%CL), assuming $m_{A^0} < m_{H_1^0}$.

89 ABREU 91B result obtained by combining with analysis of ABREU 90I.

90 BLUEMLEIN 91 excluded certain range of $\tan\beta$ for $m_{H_1^0} < 120$ MeV, $m_{A^0} < 80$ MeV.

91 DECAMP 91I searched for $Z \rightarrow H_1^0 Z^*$, and $Z \rightarrow H_1^0 A^0 \rightarrow 4\text{jets or } \tau\tau jj$ or $3A^0$. Their limits take into account the one-loop radiative corrections to the Higgs potential with varied top and squark masses.

92 ABREU 90E searched for $Z \rightarrow H_1^0 A^0$ and $Z \rightarrow H_1^0 Z^*$. $m_{H_1^0} < 210$ MeV is not excluded by this analysis.

93 ADEVA 90R result is from $Z \rightarrow H_1^0 A^0 \rightarrow 4\text{jet or } \tau\tau jj$ or 4τ and $Z \rightarrow H_1^0 Z^*$. Some region of $m_{H_1^0} < 4$ GeV is not excluded by this analysis.

94 DECAMP 90E look for $Z \rightarrow H_1^0 A^0$ as well as $Z \rightarrow H_1^0 \ell^+ \ell^-, Z \rightarrow H_1^0 \nu\bar{\nu}$ with 18610 Z decays. Their search includes signatures in which H_1^0 and A^0 decay to $\gamma\gamma, e^+ e^-, \mu^+ \mu^-, \tau^+ \tau^-,$ or $q\bar{q}$. See their figures of $m_{H_1^0}$ vs. $\tan\beta$.

95 DECAMP 90H is similar to DECAMP 90E but with 25,000 Z decays.

96 DECAMP 90M looked for $Z \rightarrow H^0 \ell\ell$, where H^0 decays outside the detector. This excludes a region in the $(m_{H_1^0}, \tan\beta)$ plane centered at $m_{H_1^0} = 50$ MeV, $\tan\beta = 0.5$. This limit together with DECAMP 90E result excludes $m_{H_1^0} < 3$ GeV for any $\tan\beta$.

 A^0 (Pseudoscalar Higgs Boson) MASS LIMITS In Supersymmetric Models

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>21	95	97 BUSKULIC	93i ALEP	$\tan\beta > 1$
>32	95	98 ABREU	92J DLPH	$\tan\beta > 3$
>24	95	99 ADRIANI	92G L3	$1 < \tan\beta < 50$
none 3–40.5	95	100 AKRAWY	91C OPAL	$\tan\beta > 1$, if 3 GeV $< m_{H_1^0} < m_{A^0}$
>12	95	101 ABREU	90E DLPH	$\tan\beta < 1$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 0.21	95	102 ELLIS	93 RVUE	Electroweak
>20	95	103 BUSKULIC	92 ALEP	$\tan\beta > 1$
>34	95	104 DECAMP	91 ALEP	$\tan\beta > 1$
>39	95	105 ADEVA	90R L3	$\tan\beta > 1$, $m_{H_1^0} < m_{A^0}$

97 BUSKULIC 93i search for $Z \rightarrow H^0 Z^*$ and $Z \rightarrow H^0 A^0$. One-loop corrections to the Higgs potential are included with any $m_t, m_{\bar{t}} > m_t$. For $m_t = 140$ GeV and $m_{\bar{t}} = 1$ TeV, the limit is $m_{A^0} > 45$ GeV. Assumes no invisible H^0 or A^0 decays.

98 ABREU 92J searched for $Z \rightarrow H_1^0 Z^*$ and $Z \rightarrow H_1^0 A^0$ with $H_1^0, A^0 \rightarrow \tau\tau$ or jet-jet. Small mass values are excluded by ABREU 91B.

99 ADRIANI 92G search for $Z \rightarrow H_1^0 Z^*, Z \rightarrow H_1^0 A^0 \rightarrow 4b, bb\tau\tau, 4\tau, 6b$ (via $H^0 \rightarrow A^0 A^0$), and include constraints from $\Gamma(Z)$. One-loop corrections are included with $90 < m_t < 250$ GeV, $m_{\bar{t}} < m_{\bar{t}} < 1$ TeV. The region $m_{A^0} < 11$ GeV is allowed if $42 < m_{H_1^0} < 62$ GeV, but is excluded by other experiments.

100 AKRAWY 91C result from $Z \rightarrow H_1^0 A^0 \rightarrow 4\text{jet or } \tau^+ \tau^- jj$ or 4τ . See paper for the excluded region for the case $\tan\beta < 1$.

101 ABREU 90E searched $Z \rightarrow H_1^0 A^0$ and $Z \rightarrow H_1^0 Z^*$. $m_{A^0} < 210$ MeV is not excluded by this analysis.

102 ELLIS 93 analyze possible constraints on the MSSM Higgs sector by electroweak precision measurements and find that m_{A^0} is not constrained by the electroweak data.

103 BUSKULIC 92 limit is from $\Gamma(Z), Z \rightarrow H^0 Z^*$, and $Z \rightarrow H^0 A^0$. The limit is valid for any $m_{H_1^0}$ below the theoretical limit $m_{H_1^0} < 64$ GeV which holds for $m_{A^0} \sim 0$ in the minimal supersymmetric model. One-loop radiative corrections are included.

104 DECAMP 91I searched for $Z \rightarrow H_1^0 Z^*$, and $Z \rightarrow H_1^0 A^0 \rightarrow 4\text{jets or } \tau\tau jj$ or $3A^0$. Their limits take into account the one-loop radiative corrections to the Higgs potential with varied top and squark masses. For $m_t = 140$ GeV and $m_{\bar{t}} = 1$ TeV, the limit is $m_{A^0} > 31$ GeV.

105 ADEVA 90R result is from $Z \rightarrow H_1^0 A^0 \rightarrow 4\text{jet or } \tau\tau jj$ or 4τ and $Z \rightarrow H_1^0 Z^*$. Some region of $m_{A^0} < 5$ GeV is not excluded by this analysis.

MASS LIMITS for Associated Higgs Production in $e^+ e^-$ Interactions

In multi-Higgs models, associated production of Higgs via virtual or real Z in $e^+ e^-$ annihilation, $e^+ e^- \rightarrow H_1^0 H_2^0$, is possible if H_1^0 and H_2^0 have opposite CP eigenvalues. Limits are for the mass of the heavier Higgs H_2^0 in two-doublet models.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>45	95	106 ADRIANI	92G L3	$m_{H_1^0} < 20$ GeV
>37.5	95	107 DECAMP	90H ALEP	$m_{H_1^0} < m_{H_2^0}$
none 5–45	95	108 KOMAMIYA	90 MRK2	$m_{H_1^0} < 0.5$ GeV,
> 8	90	109 KOMAMIYA	89 MRK2	$H_2^0 \rightarrow q\bar{q}$ or $\tau^+ \tau^-$, $H_1^0 \rightarrow \mu^+ \mu^-$,
>28	95	110 LOW	89 AMY	$H_2^0 \rightarrow q\bar{q}, \tau^+ \tau^-$, $m_{H_1^0} \lesssim 20$ MeV,
none 2–9	90	111 AKERLOF	85 HRS	$H_2^0 \rightarrow q\bar{q}$, $m_{H_1^0} = 0$,
none 4–10	90	112 ASH	85C MAC	$H_2^0 \rightarrow f\bar{f}$, $m_{H_1^0} = 0.2$ GeV,
none 1.3–24.7	95	111 BARTEL	85L JADE	$H_2^0 \rightarrow \tau^+ \tau^-, c\bar{c}$, $m_{H_1^0} = 0.2$ GeV, $H_2^0 \rightarrow f\bar{f}$ or $f\bar{f} H_1^0$
none 1.2–13.6	95	111 BEHREND	85 CELL	$m_{H_1^0} = 0$, $H_2^0 \rightarrow f\bar{f}$
none 1–11	90	111 FELDMAN	85 MRK2	$m_{H_1^0} = 0, H_2^0 \rightarrow f\bar{f}$
none 1–9	90	111 FELDMAN	85 MRK2	$m_{H_1^0} = m_{H_2^0}$, $H_2^0 \rightarrow f\bar{f}$
106 ADRIANI 92G excluded regions of the $m_{H_1^0} - m_{A^0}$ plane for various decay modes with limits $B(Z \rightarrow H_1^0 H_2^0) < (2-20) \times 10^{-4}$ are shown in Figs. 2–5.				
107 DECAMP 90H search for $Z \rightarrow H_1^0 e^+ e^-, H_1^0 \mu^+ \mu^-, H_1^0 \tau^+ \tau^-, H_1^0 q\bar{q}$, low multiplicity final states, $\tau\tau$ -jet-jet final states and 4-jet final states.				
108 KOMAMIYA 90 limits valid for $\cos^2(\alpha - \beta) \approx 1$. They also search for the cases $H_1^0 \rightarrow \mu^+ \mu^-, \tau^+ \tau^-$, and $H_2^0 \rightarrow H_1^0 H_1^0$. See their Fig. 2 for limits for these cases.				
109 KOMAMIYA 89 assume $B(H_1^0 \rightarrow \mu^+ \mu^-) = 100\%$, $2m_\mu < m_{H_1^0} < m_{\tau^-}$. The limit is for maximal mixing. A limit of $m_{H_2^0} > 18$ GeV for the case $H_2^0 \rightarrow H_1^0 H_1^0$ ($H_1^0 \rightarrow \mu^+ \mu^-$) is also given. From PEP at $E_{\text{cm}} = 29$ GeV.				
110 LOW 89 assume that H_1^0 escapes the detector. The limit is for maximal mixing. A reduced limit of 24 GeV is obtained for the case $H_2^0 \rightarrow H_1^0 f\bar{f}$. Limits for a Higgs-triplet model are also discussed. $E_{\text{cm}}^{\text{max}} = 50-60.8$ GeV.				
111 The limit assumes maximal mixing and that H_1^0 escapes the detector.				
112 ASH 85 assumes that H_1^0 escapes undetected. The bound applies up to a mixing suppression factor of 5.				

 H^\pm (Charged Higgs or Techni- ρ) MASS LIMITS

Most of the following limits assume $B(H^+ \rightarrow \tau^+ \nu) + B(H^+ \rightarrow c\bar{s}) = 1$. DECAMP 90I, BEHREND 87, and BARTEL 86 assume $B(H^+ \rightarrow \tau^+ \nu) + B(H^+ \rightarrow c\bar{s}) + B(H^+ \rightarrow c\bar{b}) = 1$. All limits from Z decays as well as ADACHI 90B assume that H^+ has weak isospin $T_3 = +1/2$. For a discussion of techni-particles, see EICHTEN 86.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>41	95	113 ADRIANI	92G L3	$B(\tau\nu) = 0-1$
>41.7	95	114,115 DECAMP	92 ALEP	$B(\tau\nu) = 0-1$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>35.4	95	116 BARGER	93 RVUE	$b \rightarrow s\gamma$
>35.4	95	117 BELANGER	93 RVUE	$b \rightarrow s\gamma$
>35.4	95	116 HEWETT	93 RVUE	$b \rightarrow s\gamma$
>35.4	95	118 ALITTI	92F UA2	$t \rightarrow bH^+$, $H^+ \rightarrow \tau\nu\tau$
none 8.0–20.2	95	119 ALBAJAR	91B UA1	$t \rightarrow bH^+$, $H^+ \rightarrow \tau\nu\tau$
>29	95	120 YUZUKI	91 VNS	$B(\ell\nu) = 0-1$
>19	95	114,121 ABREU	90B DLPH	$B(\tau\nu) = 0-1$
>36.5	95	114,122 ADACHI	90B TOPZ	$B(\tau\nu) = 0-1$
>35	95	114,123 ADEVA	90M L3	$B(\tau\nu) = 0-1$
>35.4	95	114,124 AKRAWY	90K OPAL	$B(\tau\nu) = 0-1$
>35.4	95	114,125 DECAMP	90I ALEP	$B(\tau\nu) = 0-1$
none 10–20	95	126 SMITH	90B AMY	$B(\tau\nu) > 0.7$
>19	95	125 BEHREND	87 CELL	$B(\tau\nu) = 0-1$
>18	95	127 BARTEL	86 JADE	$B(\tau\nu) = 0.1-1.0$
>17	95	127 ADEVA	85 MRKJ	$B(\tau\nu) = 0.25-1.0$

- 113 ADRIANI 92G limit improves to 44 GeV if $B(\tau\nu_\tau) > 0.4$.
- 114 Studied $H^+H^- \rightarrow (\tau\nu) + (\tau\nu)$, $H^+H^- \rightarrow (\tau\nu) + \text{hadrons}$, $H^+H^- \rightarrow \text{hadrons}$.
- 115 DECAMP 92 limit improves to 45.3 GeV for $B(\tau\nu)=1$.
- 116 HEWETT 93 and BARGER 93 analyze charged Higgs contribution to $b \rightarrow s\gamma$ in two-doublet models with the CLEO limit $B(b \rightarrow s\gamma) < 8.4 \times 10^{-4}$ (90% CL) and find lower limits on m_{H^\pm} in the type of model (model II) in which different Higgs are responsible for up-type and down-type quark masses. HEWETT 93 give $m_{H^\pm} > 110$ (70) GeV for $m_t > 150$ (120) GeV using $m_b = 5$ GeV. BARGER 93 give $m_{H^\pm} > 155$ GeV for $m_t = 150$ GeV using $m_b = 4.25$ GeV. The authors employ leading logarithmic QCD corrections and emphasize that the limits are quite sensitive to m_b .
- 117 BELANGER 93 make an analysis similar to BARGER 93 and HEWETT 93 with an improved CLEO limit $B(b \rightarrow s\gamma) < 5.4 \times 10^{-4}$ (95%CL). For the Type II model, the limit $m_{H^\pm} > 540$ (300) GeV for $m_t > 150$ (120) GeV is obtained. The authors employ leading logarithmic QCD corrections.
- 118 ALITTI 92F search for $t \rightarrow bH^+, H^+ \rightarrow \tau\nu_\tau$ with τ decaying hadronically in $p\bar{p}$ collisions at $E_{cm} = 630$ GeV. m_{H^\pm} between 40 and 65 GeV is excluded if $m_t - m_H = m_b + (\lesssim \text{a few-10 GeV})$. See Figs. 5, 6 for the excluded region for $B(H^+ \rightarrow \tau\nu_\tau) = 1, 0.5$.
- 119 ALBAJAR 91B search for $W \rightarrow t\bar{b}$ and $t\bar{t}$ production in $p\bar{p}$ collisions with the decay chain $t \rightarrow H^+b, H^+ \rightarrow \tau^+\nu$. In single muon plus jets and dimuon channels. For $m_t = 60$ GeV, $m_{H^\pm} < 47$ GeV is excluded at 95%CL if $\tan\beta > 2.3$. The search is restricted to small values of m_t , and no limit on m_{H^\pm} is obtained if $m_t > 61$ GeV. Note that existing limits on m_t are not valid if $t \rightarrow H^+b$.
- 120 YUZUKI 91 assume photon exchange. The limit is valid for any decay mode $H^+ \rightarrow e\nu, \mu\nu, \tau\nu, q\bar{q}$ with five flavors. For $B(\ell\nu) = 1$, the limit improves to 25.0 GeV.
- 121 ABREU 90B limit improves to 36 GeV for $B(\tau\nu) = 1$.
- 122 ADACHI 90B limit improves to 22 GeV for $B(\tau\nu) = 0.6$.
- 123 ADEVA 90M limit improves to 42.5 GeV for $B(\tau\nu) = 1$.
- 124 AKRAWY 90K limit improves to 43 GeV for $B(\tau\nu) = 1$.
- 125 If $B(H^+ \rightarrow \tau^+\nu) = 100\%$, the DECAMP 90I limit improves to 43 GeV.
- 126 SMITH 90B limit applies for $v_2/v_1 > 2$ in a model in which H_2 couples to u -type quarks and charged leptons.
- 127 Studied $H^+H^- \rightarrow (\tau\nu) + (\tau\nu)$, $H^+H^- \rightarrow (\tau\nu) + \text{hadrons}$. Search for muon opposite hadronic shower.

MASS LIMITS for $H^{\pm\pm}$ (doubly-charged Higgs boson)

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.6	95	128 ACTON	92M OPAL	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>30.4	95	128 ACTON	92M OPAL	$T_3(H^{++}) = +1$
>25.5	95	129 ACTON	92M OPAL	$T_3(H^{++}) = 0$
none 6.5–36.6	95	130 SWARTZ	90 MRK2	$T_3(H^{++}) = +1$
none 7.3–34.3	95	130 SWARTZ	90 MRK2	$T_3(H^{++}) = 0$
128 ACTON 92M limit assumes $H^{\pm\pm} \rightarrow \ell^\pm\ell^\pm$ or $H^{\pm\pm}$ does not decay in the detector. Thus the region $g_{\ell\ell} \approx 10^{-7}$ is not excluded.				
129 ACTON 92M from $\Delta\Gamma_Z < 40$ MeV.				
130 SWARTZ 90 assume $H^{\pm\pm} \rightarrow \ell^\pm\ell^\pm$ (any flavor). The limits are valid for the Higgs-lepton coupling $g(H\ell\ell) \gtrsim 7.4 \times 10^{-7} [m_H/\text{GeV}]^{1/2}$. The limits improve somewhat for $e\bar{e}$ and $\mu\bar{\mu}$ decay modes.				

H^0 and H^\pm REFERENCES

ABREU 94G NP B421 3
AKERS 94B PL B327 397
ELLIS 94 PL B324 173
ADRIANI 93C PL B303 391
BARGER 93 PRL 70 1368
BELANGER 93 PR D48 5419
BLONDEL 93 PL B311 346
BRAHMACH... 93 PR D48 4224
BUSKULIC 93H PL B313 299
BUSKULIC 93I PL B313 312
ELLIS 93B NP B393 3
ELLIS 93B PL B318 148
GROSS 93 JUMP A8 407
HEWETT 93 PRL 70 1045
LOPEZ-FERN... 93 PL B312 240
NOVIKOV 93B PL B308 123
ABREU 92D ZPHY C53 555
ABREU 92I NP B373 3
ACTON 92M PL B295 347
ADEVA 92B PL B283 454
ADRIANI 92F PL B292 472
ADRIANI 92G PL B294 457
Also 93B ZPHY C57 355
ALITTI 92F PL B280 137
BARABASH 92B NP B381 451
Also 92B SJNP 55 1810
Translated from YAF 55 3247.
BLUEMLEIN 92 IJMP A7 3835
BUSKULIC 92 PL B285 309
DECAMP 92 PRPL 216 253
DELAGUILA 92B NP B381 451
ELLIS 92 PL B274 456
ELLIS 92E PL B292 427
LEEB 92 PRL 68 1472
PICH 92 NP B388 31
+Adam, Adye, Agasi, Ajinenko+ (DELPHI Collab.)
+Alexander, Allison, Anderson, Arcelli+ (OPAL Collab.)
+Fogli, Lisi (CERN, BARI)
+Aguiar-Benitez, Ahlen, Alcaraz, Aloiso+ (L3 Collab.)
+Berger, Phillips (WISC, RAL)
+Geng, Turcotte (MONT, ISU, AMES)
+Verzegnassi (EPOL, TRSTT, TRSTI)
Brahmachari, Josphipura, Rindani+(AHMED, TATA, CERN)
+De Bonis, Decamp, Ghez, Goy+ (ALEPH Collab.)
+De Bonis, Decamp, Ghez, Goy, Lees+ (ALEPH Collab.)
+Fogli, Lisi (CERN, BARI)
+Fogli, Lisi (CERN, BARI)
+Yepes (CERN)
(ANL, OREG)
Lopez-Fernandez, Romao+ (CERN, LISB, VALE)
(SERP, CERN)
+Adam, Adami, Adye, Akesson, Alekseev+(DELPHI Collab.)
+Adam, Adami, Adye, Akesson+ (DELPHI Collab.)
+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
+Adriani, Aguiar-Benitez, Ahlen, Akbari+ (L3 Collab.)
+Aguiar-Benitez, Ahlen, Akbari, Alcaraz+ (L3 Collab.)
+Aguiar-Benitez, Ahlen, Akbari, Alcaraz+ (L3 Collab.)
+Adriani, Aguiar-Benitez, Ahlen, Alcaraz+ (L3 Collab.)
+Ambrósini, Ansari, Autiero, Bärety+ (UA1 Collab.)
+Barabash, Barger, CERN, SERP, SUDA, (CERN)
Barabash+ (JINR, CERN, SERP, BUDA, BERL)
+Brunner, Grabosch+ (BERL, BUDA, JINR, SERP)
+Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
del Aguila, Martinez, Quiros (CERN, BARI)
+Fogli, Lisi (CERN, BARI)
+Fogli, Lisi (CERN, BARI)
+Schmiedmayer (TUW, HARV, RIS)
+Prades, Yepes (CERN, CNRS)

RENTON 92 ZPHY C56 355
SCHLE 92 ZPHY C54 387
ABREU 91B ZPHY C51 25
ACTON 91 PL B268 122
ADEVA 91 PL B257 450
ADEVA 91D PL B262 155
AKRAWY 91 PL B253 511
AKRAWY 91C ZPHY C49 1
ALBAJAR 91B PL B257 459
BLUEMLEIN 91 ZPHY C51 341
DECAMP 91F PL B262 139
DECAMP 91I PL B265 475
SCHMIEDM... 91 PRL 66 1015
YUZUKI 91 PL B267 309
ABE 90E PR D41 1717
ABREU 90B PL B241 449
ABREU 90C NP B342 7
ABREU 90E PL B245 276
ABREU 90I HEP-90 Singapore (unpub.)
CERN-PPE/90-163
ADACHI 90B PL B240 513
ADEVA 90H PL B248 203
ADEVA 90M PL B252 511
ADEVA 90N PL B252 518
ADEVA 90P PL B251 311
AKRAWY 90C PL B236 224
AKRAWY 90K PL B242 299
AKRAWY 90P PL B251 211
ANTREASYAN 90C PL B251 204
ATIYA 90 PRL 64 21
ATIYA 90B PRL 65 1188
BARR 90 PL B235 356
DAWSON 90 PR D41 2844
DECAMP 90 PL B235 233
DECAMP 90E PL B237 291
DECAMP 90H PL B241 141
DECAMP 90I PL B241 623
DECAMP 90M PL B245 289
DECAMP 90N PL B246 306
KOMAMIYA 90 PRL 64 2881
LEUTWYLER 90 NP B343 369
SMITH 90B PR D42 949
SWARTZ 90 PRL 64 2877
ALAM 89B PR D40 712
Also 89C PR D40 3790 erratum
ALBRECHT 89 ZPHY C42 349
ATIYA 89 PRL 63 2177
BARBIERI 89 PL B219 503
CAHN 89 PR D42 949
CHENG 89 PR D40 2980
CHENG 89B PL B219 347
DAVIER 89 PL B229 150
DAWSON 89 PL B222 143
EGLI 89 PL B222 533
EILAM 89 PL B231 184
GRIFOLS 89 PRL 63 1946
KOMAMIYA 89 PR D40 721
LINDNER 89 PL B228 139
LOW 89 PL B228 548
RABY 89 PR D39 828
SHER 89 PRPL 179 273
SNYDER 89 PL B229 169
WILLEY 89 PR D39 2784
YEPES 89 PL B227 182
YEPES 89B PRL 63 1956
CHENG 88 PR D38 2869
CHIVUKULA 88 PL B207 86
Also 89 PL B217 568 erratum
GRINSTEIN 88 PL B211 363
LEE-FRANZINI 88 Munich HEP Conf. p. 1432
SCHMIEDM... 88 PRL 61 1065
Also 88 PRL 61 2509 erratum
ALBRECHT 87D PL B199 451
BAKER 87 PRL 59 2832
Also 88 PRL 60 472 erratum
BEHREND 87 PL B193 376
DRUZHININ 87 ZPHY C37 1
FRANZINI 87 PR D35 2883
RUSKOV 87 PL B187 165
BARTEL 86 ZPHY C31 359
BELTRAMI 86 NP A451 679
EICHTEEN 86 PR D34 1547
VOLOSHIN 86 SJNP 43 495
+Manohar (BOST, MIT)
Chivukula, Manohar (BOST, MIT)
+Hall, Randall (LBL, UCB)
Schmiedmayer, Rauch, Rihs (CUSB Collab.)
+Andam, Abrams, Boeckmann+ (ARGUS Collab.)
+Gordon, Lazarus+ (BNL, SIN, WASH, YALE)
Baker, Gordon+ (BNL, SIN, WASH, YALE)
+Buerger, Criegee, Dainton+ (CELLO Collab.)
+Dubrovina, Eideiman, Golubev+ (NOVO)
+Son, Tuts, Yousef, Zhao+ (CUSB Collab.)
+Becker, Feist, Haidt+ (SOFI)
+Aas, Beer, Dechambrier, Goudsmit+ (ETH, FSU)
+Hincliffe, Lane, Quigg+ (FNAL, LBL, OSU)
+Okun (ITEP)
Translated from YAF 43 779
WILLEY 86 PL B173 480
ADEVA 85 PL 152B 439
AKERLOF 85 PL 156B 271
ALBRECHT 85J ZPHY C29 167
ASH 85 PRL 55 1831
ASH 85C PRL 54 2477
BARTEL 85 PL 155B 288
BEHREND 85 PL 161B 182
FELDMAN 85 PRL 54 2289
ALTHOFF 84G ZPHY C22 219
FREEDMAN 84 PRL 52 240
MUKHOPAD... 84 PR D29 565
YAMAZAKI 84 PRL 52 1089
HOFFMAN 83 PR D28 660
ASANO 81B PL 107B 159
DZHELZYADIN 81 PL 105B 239
WITTTEN 81 NP B177 477
GUTH 80 PR 45 1131
JACQUES 80 PRL D21 1206
SHER 80 PR D22 2989
Also 83 ANP 148 95
BECHIS 78 PRL 40 602
SHIFMAN 78 PL 78B 443
BARBIERI 75 PL 57B 270
KOHLER 74 PRL 33 1628
ALEKSANDROV66 JETPL 4 134
Translated from ZETF 4 196.
(OXF)
(FRIE)
(DELPHI Collab.)
+Alexander, Allison, Allport+ (OPAL Collab.)
+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
+Albrow, Altkofer, Ankoviak, Apsimon+ (UA1 Collab.)
+Brunner, Grabosch+ (BERL, BUDA, JINR, SERP)
+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
Schmiedmayer, Rihs, Harvey, Hill (TUW, ORNL)
+Haba, Abe, Amako, Arai, Asano+ (VENUS Collab.)
+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
+Adam, Adami, Adye, Alekseev+ (DELPHI Collab.)
+Aihara, Doerer, Enomoto+ (TOPAZ Collab.)
+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
+Alexander, Allison, Allport+ (OPAL Collab.)
+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
+Bartels, Besset, Bieler, Bienlein+ (Crystal Ball Collab.)
+Chiang, Frank, Haggerty, Ito, Kycia+ (BNL 787 Collab.)
+Chiang, Frank, Haggerty, Ito, Kycia+ (BNL 787 Collab.)
+Clark+ (CERN, EDIN, MANZ, BNL, PISA, SIEG)
+Gunion, Haber (BNL, UCD, UCSB)
+Deschizeaux, Lees, Minard, Crespo+ (ALEPH Collab.)
+Deschizeaux, Lees, Minard+ (ALEPH Collab.)
+Deschizeaux, Lees, Minard+ (ALEPH Collab.)
+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
+Abrams, Adolphsen, Averill, Ballam+ (Mark II Collab.)
+Shifman (BERN)
+McNeil, Breedon, Kim, Ko+ (AMY Collab.)
+Abrams, Adolphsen, Averill, Ballam+ (Mark II Collab.)
+Katayama, Kim, Li, Lou, Sun+ (CLEO Collab.)
Alam, Katayama, Kim, Li, Lou, Sun+ (CLEO Collab.)
+Boeckmann, Glaeser, Harder+ (ARGUS Collab.)
+Chiang, Frank, Haggerty, Ito, Kycia+ (BNL 787 Collab.)
+Curci
+Yu (AST)
+Nguyen Ngoc (LALO)
(BNL)
+Engfer, Grab, Hermes, Kraus+ (SINDRUM Collab.)
+Nakada, Wlyer (PSI, ZURI)
+Masso, Paris (BARC)
+Fordham, Abrams, Adolphsen, Akerlof+ (Mark II Collab.)
+Sher, Zaglauer (FNAL, WUSL)
+Xu, Abashian, Gotow, Hu, Mattson+ (AMY Collab.)
+West, Hoffman (LANL)
+Murray, Abrams, Adolphsen, Akerlof+ (Mark II Collab.)
(PITT)
(MCG)
(MCG)
+Manohar (BOST, MIT)
Chivukula, Manohar (BOST, MIT)
+Hall, Randall (LBL, UCB)
Schmiedmayer, Rauch, Rihs (CUSB Collab.)
+Andam, Abrams, Boeckmann+ (ARGUS Collab.)
+Gordon, Lazarus+ (BNL, SIN, WASH, YALE)
Baker, Gordon+ (BNL, SIN, WASH, YALE)
+Buerger, Criegee, Dainton+ (CELLO Collab.)
+Dubrovina, Eideiman, Golubev+ (NOVO)
+Son, Tuts, Yousef, Zhao+ (CUSB Collab.)
+Becker, Feist, Haidt+ (SOFI)
+Aas, Beer, Dechambrier, Goudsmit+ (ETH, FSU)
+Hincliffe, Lane, Quigg+ (FNAL, LBL, OSU)
+Okun (ITEP)
Translated from YAF 43 779
+Becker, Becker-Zsedy+ (Mark-I Collab.)
+Bonvicini, Chapman, Errede+ (HRS Collab.)
+Binder, Harder+ (ARGUS Collab.)
+Band, Blume, Camporesi+ (MAC Collab.)
+Band, Blume, Camporesi+ (MAC Collab.)
+Becker, Costa, Feist, Hagiwara+ (JADE Collab.)
+Burger, Criegee, Fenner+ (CELLO Collab.)
+Abrams, Amidei, Baden+ (Mark II Collab.)
+Braunschweig, Kirschfink+ (TASSO Collab.)
+Napolitano, Camp, Kroupa (ANL, CHIC)
Mukhopadhyay, Goudsmit+ (RPI, SIN, LISB)
+Ishikawa, Taniguchi, Yamanaka+ (INUS, KEK)
+Frank, Mischke, Moir, Schardt (LANL, ARZ)
+Kikuitani, Kurokawa, Miyachi+(KEK, TOKY, INUS, OSAK)
+Golovkin, Konstantinov, Kubarovski+ (SERP)
(HARV)
(SLAC)
+Weinberg
+Kalekar, Miller, Piano+ (RUTG, STEV, COLU)
(UCSC, UC)
Flores, Sher (UCSC, UC)
+Chang, Dombeck, Ellsworth, Glasser, Lau+ (UMD)
+Ericson (CERN)
+Watson, Becker (LOCK)
+Samostv, Serreter, Tsoi (JINR)
Translated from ZETF 4 196.

Gauge & Higgs Boson Full Listings

Heavy Bosons Other than Higgs Bosons

Searches for Heavy Bosons Other Than Higgs Bosons

We list here various limits on charged and neutral heavy vector bosons (other than W 's and Z 's), heavy scalar bosons (other than Higgs bosons), vector or scalar leptoquarks, and axiguons.

W_R (Right-Handed W Boson) MASS LIMITS

Assuming a light right-handed neutrino, except for BEALL 82, LANGACKER 89B, and COLANGELO 91. $g_R = g_L$ assumed. [Limits in the section MASS LIMITS for W' below are also valid for W_R if $m_{\nu_R} \ll m_{W_R}$.] Some limits assume manifest left-right symmetry, i.e., the equality of left- and right Cabibbo-Kobayashi-Maskawa matrices. For a comprehensive review, see LANGACKER 89B. Limits on the W_L - W_R mixing angle ζ are found in the next section. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 406	90	1 JODIDIO	86 ELEC	Any ζ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 439	90	2 BHATTACH...	93 RVUE	Z - Z' mixing
> 225	90	3 SEVERIJNS	93 CNTR	β^+ decay
		4 IMAZATO	92 CNTR	K^+ decay
> 475	90	5 POLAK	92B RVUE	μ decay
> 240	90	6 AQUINO	91 RVUE	Neutron decay
> 496	90	6 AQUINO	91 RVUE	Neutron and muon decay
> 700		7 COLANGELO	91 THEO	$m_{K_L^0} - m_{K_S^0}$
> 477	90	8 POLAK	91 RVUE	μ decay
[none 540-23000]		9 BARBIERI	89B ASTR	SN 1987A; light ν_R
> 300	90	10 LANGACKER	89B RVUE	General
> 160	90	11 BALKE	88 CNTR	$\mu \rightarrow e \nu \bar{\nu}$
> 482	90	1 JODIDIO	86 ELEC	$\zeta = 0$
> 800		MOHAPATRA	86 RVUE	$SU(2)_L \times SU(2)_R \times U(1)$
> 400	95	12 STOKER	85 ELEC	Any ζ
> 475	95	12 STOKER	85 ELEC	$\zeta < 0.041$
		13 BERGSMA	83 CHRM	$\nu_\mu e \rightarrow \mu \nu_e$
> 380	90	14 CARR	83 ELEC	μ^+ decay
> 1600		15 BEALL	82 THEO	$m_{K_L^0} - m_{K_S^0}$
[> 4000]		STEIGMAN	79 COSM	Nucleosynthesis; light ν_R

- 1 JODIDIO 86 is the same TRIUMF experiment as STOKER 85 (and CARR 83); however, it uses a different technique. The results given here are combined results of the two techniques. The technique here involves precise measurement of the end-point e^+ spectrum in the decay of the highly polarized μ^+ .
- 2 BHATTACHARYYA 93 uses Z - Z' mixing limit from LEP '90 data, assuming a specific Higgs sector of $SU(2)_L \times SU(2)_R \times U(1)$ gauge model. The limit is for $m_t = 200$ GeV and slightly improves for smaller m_t .
- 3 SEVERIJNS 93 measured polarization-asymmetry correlation in ^{107}In β^+ decay. The value $R/R_0 = 0.926 \pm 0.041$ is 1.7 standard deviations away from the Standard Model expectation, giving $[m_1^2/m_2^2 + \zeta]^2 = 0.087 \pm 0.051$. The listed limit assumes zero L - R mixing.
- 4 IMAZATO 92 measure positron asymmetry in $K^+ \rightarrow \mu^+ \nu_\mu$ decay and obtain $\xi_{P,\mu} > 0.990$ (90%CL). If W_R couples to $u\bar{s}$ with full weak strength ($V_{us}^R = 1$), the result corresponds to $m_{W_R} > 653$ GeV. See their Fig. 4 for m_{W_R} limits for general $|V_{us}^R|^2 = 1 - |V_{ud}^R|^2$.
- 5 POLAK 92B limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming $\zeta = 0$. Supersedes POLAK 91.
- 6 AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right symmetry assumed. Stronger of the two limits also includes muon decay results.
- 7 COLANGELO 91 limit uses hadronic matrix elements evaluated by QCD sum rule and is less restrictive than BEALL 82 limit which uses vacuum saturation approximation. Manifest left-right symmetry assumed.
- 8 POLAK 91 limit is from fit to muon decay parameters and is essentially determined by JODIDIO 86 data assuming $\zeta = 0$. Superseded by POLAK 92B.
- 9 BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.
- 10 LANGACKER 89B limit is for any ν_R mass (either Dirac or Majorana) and for a general class of right-handed quark mixing matrices.
- 11 BALKE 88 limit is for $m_{\nu_{eR}} = 0$ and $m_{\nu_{\mu R}} \leq 50$ MeV. Limits come from precise measurements of the muon decay asymmetry as a function of the positron energy.
- 12 STOKER 85 is same TRIUMF experiment as CARR 83. Here they measure the decay e^+ spectrum asymmetry above 46 MeV/c using a muon-spin-rotation technique. Assumed a light right-handed neutrino. Quoted limits are from combining with CARR 83.
- 13 BERGSMA 83 set limit $m_{W_2}/m_{W_1} > 1.9$ at CL = 90%.
- 14 CARR 83 is TRIUMF experiment with a highly polarized μ^+ beam. Looked for deviation from $V-A$ at the high momentum end of the decay e^+ energy spectrum. Limit from previous world-average muon polarization parameter is $m_{W_R} > 240$ GeV. Assumes a light right-handed neutrino.
- 15 BEALL 82 limit is obtained assuming that W_R contribution to $K_L^0 - K_S^0$ mass difference is smaller than the standard one, neglecting the top quark contributions. Manifest left-right symmetry assumed.

Limit on W_L - W_R Mixing Angle ζ

Lighter mass eigenstate $W_1 = W_L \cos \zeta - W_R \sin \zeta$. Light ν_R assumed unless noted. Values in brackets are from cosmological and astrophysical considerations.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.04	90	16 MISHRA	92 CCFR	νN scattering
-0.0006 to 0.0028	90	17 AQUINO	91 RVUE	
[none 0.00001-0.02]		18 BARBIERI	89B ASTR	SN 1987A
< 0.040	90	19 JODIDIO	86 ELEC	μ decay
-0.056 to 0.040	90	19 JODIDIO	86 ELEC	μ decay

16 MISHRA 92 limit is from the absence of extra large- x , large- y $\bar{\nu}_\mu N \rightarrow \bar{\nu}_\mu X$ events at Tevatron, assuming left-handed ν and right-handed $\bar{\nu}$ in the neutrino beam. The result gives $\zeta^2(1-2m_{W_1}^2/m_{W_2}^2) < 0.0015$. The limit is independent of ν_R mass.

17 AQUINO 91 limits obtained from neutron lifetime and asymmetries together with unitarity of the CKM matrix. Manifest left-right asymmetry is assumed.

18 BARBIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.

19 First JODIDIO 86 result assumes $m_{W_R} = \infty$, second is for unconstrained m_{W_R} .

MASS LIMITS for W' (A Heavy Charged Vector Boson Other Than W) in Hadron Collider Experiments

Couplings of W' to quarks and leptons are taken to be identical with those of W . The following limits are obtained from $p\bar{p} \rightarrow W'X$ with W' decaying to the mode indicated in the comments. Experiments other than ABE 91F assume no new decay channels (esp. $t\bar{b}$) are open.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 520	95	20 ABE	91F CDF	$W' \rightarrow e\nu, \mu\nu$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 251	90	21 ALITTI	93 UA2	$W' \rightarrow q\bar{q}$
none 260-600	95	22 RIZZO	93 RVUE	$W' \rightarrow q\bar{q}$
none 101-158	90	23 ALITTI	91 UA2	$W' \rightarrow q\bar{q}$
> 220	90	24 ALBAJAR	89 UA1	$W' \rightarrow e\nu$
> 209	90	25 ANSARI	87D UA2	$W' \rightarrow e\nu$
> 210	90	26 ARNISON	86B UA1	$W' \rightarrow e\nu$
> 170	90	27 ARNISON	83D UA1	$W' \rightarrow e\nu$

20 ABE 91F assume leptonic branching ratio of 1/12 for each lepton flavor. The limit from the $e\nu$ ($\mu\nu$) mode alone is 490 (435) GeV. These limits apply to W_R if $m_{\nu_R} \lesssim 15$ GeV and ν_R does not decay in the detector. Cross section limit $\sigma \cdot B < (1-10)$ pb is given for $m_{W'} = 100-550$ GeV; see Fig. 2.

21 ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes $\Gamma(W')/m_{W'} = \Gamma(W)/m_W$ and $B(W' \rightarrow jj) = 2/3$. This corresponds to W_R with $m_{\nu_R} > m_{W_R}$ (no leptonic decay) and $W_R \rightarrow t\bar{b}$ allowed. See their Fig. 4 for limits in the $m_{W'} - B(q\bar{q})$ plane.

22 RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

23 ALITTI 91 search is based on two-jet invariant mass spectrum, assuming $B(W' \rightarrow q\bar{q}) = 67.6\%$. Limit on $\sigma \cdot B$ as a function of two-jet mass is given in Fig. 7.

24 ALBAJAR 89 cross section limit at 630 GeV is $\sigma(W') B(e\nu) < 4.1$ pb (90% CL).

25 See Fig. 5 of ANSARI 87D for the excluded region in the $m_{W'} - (g_{W'q}^2) B(W' \rightarrow e\bar{\nu})$ plane. Note that the quantity $(g_{W'q}^2) B(W' \rightarrow e\bar{\nu})$ is normalized to unity for the standard W couplings.

26 ARNISON 86B find no excess at large p_T in 148 $W \rightarrow e\nu$ events. Set limit $\sigma \cdot B(e\nu) < 10$ pb at CL = 90% at $E_{cm} = 546$ and 630 GeV.

27 ARNISON 83D find among 47 $W \rightarrow e\nu$ candidates no event with excess p_T . Also set $\sigma \cdot B(e\nu) < 30$ pb with CL = 90% at $E_{cm} = 540$ GeV.

MASS LIMITS for Z' (Heavy Neutral Vector Boson Other Than Z)

NOTE ON THE Z' SEARCHES

The mass bounds depend on the gauge group and the gauge coupling of a Z' boson. The limits listed below are not exhaustive but include only typical Z' bosons that appear frequently in the literature. The following notations are used for these Z' bosons.

Z'_{SM} : Z'_{SM} is a clone of the Z and is introduced as a convenient way to gauge the limits rather than with a theoretical motivation. It is assumed to have exactly the same couplings as the Z but a different mass.

Left-right symmetric bosons: Z_{LR} is the extra neutral boson which appears in left-right symmetric models with the gauge group $SU(2)_L \times SU(2)_R \times U(1)_{B-L}$ or $SU(2)_L \times U(1)_R \times U(1)_{B-L}$, where $U(1)_R$ is the third component of $SU(2)_R$ and the weak hypercharge $Y = T_{3R} + \frac{1}{2}(B-L)$. The Z_{LR}

See key on page 1343

Gauge & Higgs Boson Full Listings Heavy Bosons Other than Higgs Bosons

couples to $\alpha T_{3R} - (1/2\alpha)(B - L)$ with the coupling strength g' (the weak hypercharge gauge coupling). The parameter α is model dependent. For left-right symmetric coupling $g_L = g_R$, $\alpha = (1 - 2\sin^2\theta_W)^{1/2} / \sin\theta_W \approx 1.53$, which is used for the limits in the listing unless noted. Another typical case $\alpha = (2/3)^{1/2}$ is identical to Z_χ (discussed below) with the coupling $g_\chi = g'$.

E₆ bosons: Two new neutral gauge bosons appear in E₆ models. One is contained in the SO(10) subgroup and the other is not:

$$E_6 \longrightarrow \text{SO}(10) \times \text{U}(1)_\psi,$$

$$\text{SO}(10) \longrightarrow \text{SU}(5) \times \text{U}(1)_\chi.$$

One Z' is assumed to be relatively light, which in general is a linear combination of the two:

$$Z_\beta = Z_\chi \cos\beta + Z_\psi \sin\beta.$$

The gauge quantum numbers of the ordinary quarks and leptons are shown in the table:

f	T_{3R}	Y	$B-L$	$\sqrt{24}Q_\chi$	$\sqrt{\frac{72}{5}}Q_\psi$	Q_η
ν_L, e_L^-	0	$-\frac{1}{2}$	-1	+3	+1	$+\frac{1}{6}$
ν_R	$+\frac{1}{2}$	0	-1	+5	-1	$+\frac{5}{6}$
e_R^-	$-\frac{1}{2}$	-1	-1	+1	-1	$+\frac{1}{3}$
u_L, d_L	0	$+\frac{1}{6}$	$+\frac{1}{3}$	-1	+1	$-\frac{1}{3}$
u_R	$+\frac{1}{2}$	$+\frac{2}{3}$	$+\frac{1}{3}$	+1	-1	$+\frac{1}{3}$
d_R	$-\frac{1}{2}$	$-\frac{1}{3}$	$+\frac{1}{3}$	-3	-1	$-\frac{1}{6}$

In particular, the χ charge is related to others by $\sqrt{24}Q_\chi = 4Y - 5(B-L)$. Also notice that the Z_ψ coupling is pure axial for all quarks and leptons.

Another typical case Z_η is defined as

$$Z_\eta = \sqrt{\frac{3}{8}}Z_\chi - \sqrt{\frac{5}{8}}Z_\psi,$$

which appears in a superstring-motivated model.

A reference gauge coupling for these bosons is $g' = e/\cos\theta_W$, which is predicted if there is no intermediate symmetry breaking scale.

In general, these Z' models require the existence of a set of new fermions (belonging to the 27 representation of E₆) to cancel gauge anomalies, and possibly superpartners. An exception is Z_χ , for which only right-handed neutrinos are necessary. For the direct limits from hadron colliders, it is often assumed that these new fermions are heavy and are not produced in the decay of the Z' .

Limits for Z'_{SM}

Z'_{SM} is assumed to have couplings with quarks and leptons which are identical to those of Z .

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>412	95	ABE	92B CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$ $\mu^+\mu^-$
>779	95	28,29 LANGACKER	92B RVUE	

••• We do not use the following data for averages, fits, limits, etc. •••

>237	90	30 ALITTI	93 UA2	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>119	90	31 ALLEN	93 CALO	$\nu e \rightarrow \nu e$
none 490-560	95	32 RIZZO	93 RVUE	$p\bar{p}; Z'_{SM} \rightarrow q\bar{q}$
>387	95	33 ABE	91D CDF	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>307	90	34 GEIREGAT	91 CHM2	$\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$
>426	90	35 ABE	90F VNS	e^+e^-
>208	90	36 HAGIWARA	90 RVUE	e^+e^-
>173	90	37 ALBAJAR	89 UA1	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>180	90	38 ANSARI	87D UA2	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$
>160	90	39 ARNISON	86B UA1	$p\bar{p}; Z'_{SM} \rightarrow e^+e^-$

²⁸ LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

²⁹ LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0086 < \theta < 0.0005$.

³⁰ ALITTI 93 search for resonances in the two-jet invariant mass. The limit assumes $B(Z' \rightarrow q\bar{q})=0.7$. See their Fig. 5 for limits in the $m_{Z'}-B(q\bar{q})$ plane.

³¹ ALLEN 93 limit is from total cross section for $\nu e \rightarrow \nu e$, where $\nu = \nu_e, \nu_\mu, \bar{\nu}_\mu$.

³² RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

³³ ABE 91D give $\sigma(Z') \cdot B(e^+e^-) < 1.31$ pb (95%CL) for $m_{Z'} > 200$ GeV at $E_{cm} = 1.8$ TeV. Limits ranging from 2 to 30 pb are given for $m_{Z'} = 100-200$ GeV.

³⁴ GEIREGAT 91 limit is from comparison of g_V^e from $\nu_\mu e$ scattering with $\Gamma(Z \rightarrow ee)$ from LEP. Zero mixing assumed.

³⁵ ABE 90F use data for $R, R_{\ell\ell}$, and $A_{\ell\ell}$. They fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

³⁶ HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-, \tau^+\tau^-$, and hadron cross sections and asymmetries.

³⁷ ALBAJAR 89 cross section limit at 630 GeV is $\sigma(Z') B(ee) < 4.2$ pb (90% CL).

³⁸ See Fig. 5 of ANSARI 87D for the excluded region in the $m_{Z'}-(g_{Z'}^e)^2 B(Z' \rightarrow e^+e^-)$ plane. Note that the quantity $(g_{Z'}^e)^2 B(Z' \rightarrow e^+e^-)$ is normalized to unity for the standard Z couplings.

³⁹ ARNISON 86B find no excess e^+e^- pairs among 13 pairs from Z . Set limit $\sigma \times B(e^+e^-) < 13$ pb at CL = 90% at $E_{cm} = 546$ and 630 GeV.

Limits for Z'_{LR}

Z'_{LR} is the extra neutral boson in left-right symmetric models. $g_L = g_R$ is assumed unless noted. Values in parentheses assume stronger constraint on the Higgs sector, usually motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>310	95	40 ABE	92B CDF	$p\bar{p}$
>389	95	41,42 LANGACKER	92B RVUE	
>130	95	43 ADRIANI	93D L3	Z parameters
(> 1500)	90	44 ALTARELLI	93B RVUE	Z parameters
none 490-560	95	45 RIZZO	93 RVUE	$p\bar{p}; Z'_{LR} \rightarrow q\bar{q}$
>230	95	46 ABE	92B CDF	$p\bar{p}$
(> 900)	90	47 DELAGUILA	92 RVUE	
(> 1400)	90	48 LAYSSAC	92B RVUE	Z parameters
(> 564)	90	49 POLAK	92 RVUE	μ decay
>474	90	50 POLAK	92B RVUE	Electroweak
(> 1340)	90	51 RENTON	92 RVUE	
(> 800)	90	52 ALTARELLI	91B RVUE	Z parameters
(> 795)	90	53 DELAGUILA	91 RVUE	
>382	90	54 POLAK	91 RVUE	Electroweak
[> 2000]		WALKER	91 COSM	Nucleosynthesis; light ν_R
[> 500]		55 GRIFOLS	90 ASTR	SN 1987A; light ν_R
(> 460)	90	56 HE	90B RVUE	
[> 2400-6800]		57 BARBIERI	89B ASTR	SN 1987A; light ν_R
>189		58 DELAGUILA	89 RVUE	$p\bar{p}$
[> 10000]		RAFFELT	88 ASTR	SN 1987A; light ν_R
>325	90	59 AMALDI	87 RVUE	
>278	90	60 DURKIN	86 RVUE	
>150	95	61 ADEVA	85B MRKJ	$e^+e^- \rightarrow \mu^+\mu^-$

⁴⁰ These limits assume that Z' decays to known fermions only.

⁴¹ LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

⁴² LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0025 < \theta < 0.0083$.

⁴³ ADRIANI 93D give limits on the Z - Z' mixing $-0.002 < \theta < 0.015$ assuming the ABE 92B mass limit.

⁴⁴ ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV. $m_H = 100$ GeV and $\alpha_s = 0.118$ assumed. The limit improves for larger m_t (see their Fig. 5). The 90%CL limit on the Z - Z' mixing angle is in Table 4.

⁴⁵ RIZZO 93 analyses CDF limit on possible two-jet resonances. The limit is sensitive to the inclusion of the assumed K factor.

⁴⁶ These limits assume that Z' decays to all E₆ fermions and their superpartners.

⁴⁷ See Fig. 7b and 8 in DELAGUILA 92 for the allowed region in $m_{Z'}$ -mixing plane and $m_{Z'} - m_t$ plane from electroweak fit including '90 LEP data.

⁴⁸ LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.

⁴⁹ POLAK 92 limit is from $m_{W_R} > 477$ GeV, which is derived from muon decay parameters assuming light ν_R . Specific Higgs sector is assumed.

Gauge & Higgs Boson Full Listings

Heavy Bosons Other than Higgs Bosons

- 50 POLAK 92B limit is from a simultaneous fit to charged and neutral sector in $SU(2)_L \times SU(2)_R \times U(1)$ model using Z parameters, m_W , and low-energy neutral current data as of 1991. Light ν_R assumed and $m_t = m_H = 100$ GeV used. Supersedes POLAK 91.
- 51 RENTON 92 limits use LEP data taken up to '90 as well as m_W , νN , and atomic parity violation data. Specific Higgs structure is assumed.
- 52 ALTARELLI 91B is based on Z mass, widths, and A_{FB} . The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_H < 1$ TeV assumed. For large m_t , the bound improves drastically. Bounds for Z - Z' mixing angle and Z mass shift without this model assumption are also given in the paper.
- 53 DELAGUILA 91 bounds have extra assumption of superstring motivated Higgs sector. From νN neutral current data with $m_Z = 91.10 \pm 0.04$ GeV, $m_t > 77$ GeV, $m_H < 1$ TeV assumed.
- 54 POLAK 91 limit is from a simultaneous fit to charged and neutral sector in $SU(2)_L \times SU(2)_R \times U(1)$ model using m_W , m_Z , and low-energy neutral current data as of 1990. Light ν_R assumed and $m_t = m_H = 100$ GeV used. Superseded by POLAK 92B.
- 55 GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.
- 56 HE 90B model assumes a specific Higgs sector. Neutral current data of COSTA 88 as well as m_Z is used. g_R is left free in the fit.
- 57 BARRIERI 89B limit holds for $m_{\nu_R} \leq 10$ MeV.
- 58 DELAGUILA 89 limit is based on $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1.8$ pb at CERN $p\bar{p}$ collider.
- 59 A wide range of neutral current data as of 1986 are used in the fit.
- 60 A wide range of neutral current data as of 1985 are used in the fit.
- 61 ADEVA 85B measure asymmetry of μ -pair production, following formalism of RIZZO 81.

Limits for Z_χ

Z_χ is the extra neutral boson in $SO(10) \rightarrow SU(5) \times U(1)_\chi$. $g_\chi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>340	95	62 ABE	92B CDF	$p\bar{p}$
>321	95	63,64 LANGACKER	92B RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>117	95	65 BUSKULIC	94 ALEP	Z parameters
(>900)	90	66 ADRIANI	93D L3	Z parameters
>80	95	67 ALTARELLI	93B RVUE	Z parameters
(>650)	90	68 ABE	92B CDF	$p\bar{p}$
(>760)	90	69 DELAGUILA	92 RVUE	
>148	95	70 LAYSSAC	92B RVUE	Z parameters
(>700)	90	71 LEIKE	92 RVUE	Z parameters
(> 500)	90	72 RENTON	92 RVUE	
(> 570)	90	73 ALTARELLI	91B RVUE	Z parameters
(> 555)	90	74 BUCHMUELLER	91 RVUE	Z parameters
>1470]	90	75 DELAGUILA	91 RVUE	
>320	90	76 FARAGGI	91 COSM	Nucleosynthesis; light ν_R
>221	90	77 GONZALEZ-G.	91 RVUE	
>231	90	78 MAHANTHAPPA	91 RVUE	Cs
>206	90	79,80 ABE	90F VNS	e^+e^-
>335	90	80,81 ABE	90F RVUE	e^+e^- , $\nu_\mu e$
(> 650)	90	82 BARGER	90B RVUE	$p\bar{p}$
[> 1140]	90	83 GLASHOW	90 RVUE	
[> 2100]	90	84 GONZALEZ-G.	90D COSM	Nucleosynthesis; light ν_R
none <150 or > 363	90	85 GRIFOLS	90 ASTR	SN 1987A; light ν_R
>177	90	86 HAGIWARA	90 RVUE	e^+e^-
>280	95	87 DELAGUILA	89 RVUE	$p\bar{p}$
>352	90	88 DORENBOSCH	89 CHRM	$g_\chi = g_Z$
>170	90	89 COSTA	88 RVUE	
>273	90	90 ELLIS	88 RVUE	$p\bar{p}$
>266	90	91 AMALDI	87 RVUE	
>283	90	92 MARCIANO	87 RVUE	
>283	90	92 DURKIN	86 RVUE	

- 62 These limits assume that Z' decays to known fermions only.
- 63 LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.
- 64 LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0048 < \theta < 0.0097$.
- 65 BUSKULIC 94 give 95%CL limits on the Z - Z' mixing $-0.0091 < \theta < 0.0023$.
- 66 ADRIANI 93D give limits on the Z - Z' mixing $-0.004 < \theta < 0.015$ assuming the ABE 92B mass limit.
- 67 ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV, $m_H = 100$ GeV and $\alpha_S = 0.118$ assumed. The limit improves for larger m_t (see their Fig. 5). The 90%CL limit on the Z - Z' mixing angle is in their Fig. 2.
- 68 These limits assume that Z' decays to all E_6 fermions and their superpartners.
- 69 See Fig. 7a and 8 in DELAGUILA 92 for the allowed region in $m_{Z'}$ -mixing plane and $m_{Z'} - m_t$ plane from electroweak fit including '90 LEP data.
- 70 LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.
- 71 LEIKE 92 is based on '90 LEP data published in LEP 92.
- 72 RENTON 92 limits use LEP data taken up to '90 as well as m_W , νN , and atomic parity violation data. Specific Higgs structure is assumed.
- 73 ALTARELLI 91B is based on Z mass, widths, and A_{FB} . The limits are for superstring motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_H < 1$ TeV assumed. For large m_t , the bound improves drastically. Bounds for Z - Z' mixing angle and Z mass shift without this model assumption are also given in the paper.

- 74 BUCHMUELLER 91 limit is from LEP data. Specific assumption is made for the Higgs sector.
- 75 DELAGUILA 91 bounds have extra assumption of superstring motivated Higgs sector. From νN neutral current data with $m_Z = 91.10 \pm 0.04$ GeV, $m_t > 77$ GeV, $m_H < 1$ TeV assumed.
- 76 FARAGGI 91 limit assumes the nucleosynthesis bound on the effective number of neutrinos $\Delta N_\nu < 0.5$ and is valid for $m_{\nu_R} < 1$ MeV.
- 77 GONZALEZ-GARCIA 91 limit is based on low-energy neutral current data, Z mass and widths, m_W from ABE 90G. $100 < m_t < 200$ GeV, $m_H = 100$ GeV assumed. Dependence on m_t is shown in Fig. 7.
- 78 MAHANTHAPPA 91 limit is from atomic parity violation in Cs with m_W , m_Z .
- 79 ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$.
- 80 ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- 81 e^+e^- data for R , $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{C\bar{C}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.
- 82 BARGER 90B limit is based on CDF limit $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.
- 83 GLASHOW 90 model assumes a specific Higgs sector. See GLASHOW 90B.
- 84 These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).
- 85 GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.
- 86 HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries. The upper mass limit disappears at 2.7 s.d.
- 87 DELAGUILA 89 limit is based on $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1.8$ pb at CERN $p\bar{p}$ collider.
- 88 DORENBOSCH 89 obtain the limit $(g_\chi/g_Z)^2 \cdot (m_Z/m_{Z'})^2 < 0.11$ at 95% CL from the processes $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$.
- 89 A wide range of neutral current data as of 1986 are used in the fit.
- 90 Z' mass limits from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.
- 91 MARCIANO 87 limit from unitarity of Cabibbo-Kobayashi-Maskawa matrix.
- 92 A wide range of neutral current data as of 1985 are used in the fit.

Limits for Z_ψ

Z_ψ is the extra neutral boson in $E_6 \rightarrow SO(10) \times U(1)_\psi$. $g_\psi = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>320	95	93 ABE	92B CDF	$p\bar{p}$
>160	95	94,95 LANGACKER	92B RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>118	95	96 ADRIANI	93D L3	Z parameters
>180	95	97 ABE	92B CDF	$p\bar{p}$
>122	95	98 LEIKE	92 RVUE	Z parameters
>105	90	99,100 ABE	90F VNS	e^+e^-
>146	90	100,101 ABE	90F RVUE	e^+e^- , $\nu_\mu e$
>320	90	102 BARGER	90B RVUE	$p\bar{p}$
[> 160]	90	103 GONZALEZ-G.	90D COSM	Nucleosynthesis; light ν_R
[> 2000]	90	104 GRIFOLS	90D ASTR	SN 1987A; light ν_R
>136	90	105 HAGIWARA	90 RVUE	e^+e^-
>154	90	106 AMALDI	87 RVUE	
>146	90	107 DURKIN	86 RVUE	

- 93 These limits assume that Z' decays to known fermions only.
- 94 LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.
- 95 LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.0025 < \theta < 0.013$.
- 96 ADRIANI 93D give limits on the Z - Z' mixing $-0.003 < \theta < 0.020$ assuming the ABE 92B mass limit.
- 97 These limits assume that Z' decays to all E_6 fermions and their superpartners.
- 98 LEIKE 92 is based on '90 LEP data published in LEP 92.
- 99 ABE 90F use data for R , $R_{\ell\ell}$, and $A_{\ell\ell}$.
- 100 ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.
- 101 e^+e^- data for R , $R_{\ell\ell}$, $A_{\ell\ell}$, and $A_{C\bar{C}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.
- 102 BARGER 90B limit is based on CDF limit $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.
- 103 These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).
- 104 GRIFOLS 90D limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also RIZZO 91.
- 105 HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-$, $\tau^+\tau^-$, and hadron cross sections and asymmetries.
- 106 A wide range of neutral current data as of 1986 are used in the fit.
- 107 A wide range of neutral current data as of 1985 are used in the fit.

Limits for Z_η

Z_η is the extra neutral boson in E_6 models, corresponding to $Q_\eta = \sqrt{3/8} Q_\chi - \sqrt{5/8} Q_\psi$. $g_\eta = e/\cos\theta_W$ is assumed unless otherwise stated. We list limits with the assumption $\rho = 1$ but with no further constraints on the Higgs sector. Values in parentheses assume stronger constraint on the Higgs sector motivated by superstring models. Values in brackets are from cosmological and astrophysical considerations and assume a light right-handed neutrino.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>340	95	108 ABE	92B CDF	$p\bar{p}$
>182	95	109,110 LANGACKER	92B RVUE	

Gauge & Higgs Boson Full Listings

Heavy Bosons Other than Higgs Bosons

• • • We do not use the following data for averages, fits, limits, etc. • • •

>100	95	111	ADRIANI	93D L3	Z parameters
(>500)	90	112	ALTARELLI	93B RVUE	Z parameters
>230	95	113	ABE	92B CDF	$\rho\bar{\rho}$
(>450)	90	114	DELAGUILA	92 RVUE	
(>315)		115	LAYSSAC	92B RVUE	Z parameters
>118	95	116	LEIKE	92 RVUE	Z parameters
(>470)		117	RENTON	92 RVUE	
(> 300)	90	118	ALTARELLI	91B RVUE	Z parameters
>120	90	119	GONZALEZ-G.	91 RVUE	
>125	90	120,121	ABE	90F VNS	e^+e^-
>115	90	121,122	ABE	90F RVUE	$e^+e^-, \nu_\mu e$
>340		123	BARGER	90B RVUE	$p\bar{p}$
[> 820]		124	GONZALEZ-G.	90D COSM	Nucleosynthesis; light ν_R
[> 3300]		125	GRIFOLS	90 ASTR	SN 1987A; light ν_R
>100	90	126	HAGIWARA	90 RVUE	e^+e^-
[> 1040]		124	LOPEZ	90 COSM	Nucleosynthesis; light ν_R
>173		127	DELAGUILA	89 RVUE	$p\bar{p}$
>129	90	128	COSTA	88 RVUE	
>156	90	129	ELLIS	88 RVUE	
>167	90	130	ELLIS	88 RVUE	$p\bar{p}$
>111	90	128	AMALDI	87 RVUE	
>143	90	131	BARGER	86B RVUE	$p\bar{p}$
>130	90	132	DURKIN	86 RVUE	
[> 760]		124	ELLIS	86 COSM	Nucleosynthesis; light ν_R
[> 500]		124	STEIGMAN	86 COSM	Nucleosynthesis; light ν_R

108 These limits assume that Z' decays to known fermions only.

109 LANGACKER 92B fit to a wide range of electroweak data including LEP results available early '91. $m_t > 89$ GeV used.

110 LANGACKER 92B give 95%CL limits on the Z - Z' mixing $-0.038 < \theta < 0.002$.

111 ADRIANI 93D give limits on the Z - Z' mixing $-0.029 < \theta < 0.010$ assuming the ABE 92B mass limit.

112 ALTARELLI 93B limit is from LEP data available in summer '93 and is for $m_t = 110$ GeV. $m_H = 100$ GeV and $\alpha_s = 0.118$ assumed. The 90%CL limit on the Z - Z' mixing angle is in Fig. 2.

113 These limits assume that Z' decays to all E_6 fermions and their superpartners.

114 See Fig. 7d in DELAGUILA 92 for the allowed region in $m_{Z'}$ -mixing plane from electroweak fit including '90 LEP data.

115 LAYSSAC 92B limit is from LEP data available spring '92. Specific Higgs sector is assumed. See also LAYSSAC 92.

116 LEIKE 92 is based on '90 LEP data published in LEP 92.

117 RENTON 92 limits use LEP data taken up to '90 as well as m_W, ν_N , and atomic parity violation data. Specific Higgs structure is assumed.

118 ALTARELLI 91B is based on Z mass, widths, and A_{FB} . The limits are for superpartner motivated models with extra assumption on the Higgs sector. $m_t > 90$ GeV and $m_{H^0} < 1$ TeV assumed. For large m_t , the bound improves drastically. Bounds for Z - Z' mixing angle and Z mass shift without this model assumption are also given in the paper.

119 GONZALEZ-GARCIA 91 limit is based on low-energy neutral current data, LEP Z mass and widths, m_W from ABE 90G. $100 < m_t < 200$ GeV, $m_{H^0} = 100$ GeV assumed. Dependence on m_t is shown in Fig. 8.

120 ABE 90F use data for $R, R_{\ell\ell}$, and $A_{\ell\ell}$.

121 ABE 90F fix $m_W = 80.49 \pm 0.43 \pm 0.24$ GeV and $m_Z = 91.13 \pm 0.03$ GeV.

122 e^+e^- data for $R, R_{\ell\ell}, A_{\ell\ell}$, and $A_{C\bar{C}}$ below Z as well as $\nu_\mu e$ scattering data of GEIREGAT 89 is used in the fit.

123 BARGER 90B limit is based on CDF limit $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1$ pb (Nodulman, EPS Conf. '89). Assumes no new threshold is open for Z' decay.

124 These authors claim that the nucleosynthesis bound on the effective number of light neutrinos ($\delta N_\nu < 1$) constrains Z' masses if ν_R is light ($\lesssim 1$ MeV).

125 GRIFOLS 90 limit holds for $m_{\nu_R} \lesssim 1$ MeV. See also GRIFOLS 90D, RIZZO 91.

126 HAGIWARA 90 perform a fit to e^+e^- data at PEP, PETRA, and TRISTAN including $\mu^+\mu^-, \tau^+\tau^-$, and hadron cross sections and asymmetries.

127 DELAGUILA 89 limit is based on $\sigma(p\bar{p} \rightarrow Z') \cdot B(Z' \rightarrow e^+e^-) < 1.8$ pb at CERN $p\bar{p}$ collider.

128 A wide range of neutral current data as of 1986 are used in the fit.

129 Z_η mass limits obtained by combining constraints from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider and the global analysis of neutral current data by COSTA 88. Least favorable spectrum of three (E_6 27) generations of particles and their superpartners are assumed.

130 Z' mass limits from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.

131 BARGER 86B limit is based on UA1/UA2 limit on $p\bar{p} \rightarrow Z', Z' \rightarrow e^+e^-$ (Lepton Photon Symp., Kyoto, '85). Extra decay channels for Z' are assumed not to be open.

132 A wide range of neutral current data as of 1985 are used in the fit.

Limits for other Z'

$$Z_\beta = Z_X \cos\beta + Z_\psi \sin\beta$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
-------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

>360		133	DELAGUILA	92 RVUE	
		134	ALTARELLI	91 RVUE	Z_β with $\tan\beta = \sqrt{3/5}$; Cs
>190		135	MAHANTHAP.	91 RVUE	Z_β with $\tan\beta = \sqrt{3/5}$; Cs
		136	GRIFOLS	90C RVUE	
		137	DELAGUILA	89 RVUE	$p\bar{p}$
>180	90	138,139	COSTA	88 RVUE	Z_β with $\tan\beta = \sqrt{15}$
>158	90	140	ELLIS	88 RVUE	Z_β ($\tan\beta = \sqrt{15}$), $p\bar{p}$

133 Fig. 7c and 7e in DELAGUILA 92 give limits for $\tan\beta = -1/\sqrt{15}$ and $\sqrt{15}$ from electroweak fit including '90 LEP data.

134 ALTARELLI 91 limit is from atomic parity violation in Cs together with LEP, CDF data. Z - Z' mixing is assumed to be zero to set the limit.

135 MAHANTHAPPA 91 limit is from atomic parity violation in Cs with m_W, m_Z . See Table III of MAHANTHAPPA 91 (corrected in erratum) for limits on various Z' models.

136 GRIFOLS 90C obtains a limit for Z' mass as a function of mixing angle β (his $\theta = \beta - \pi/2$), which is derived from a LAMPF experiment on $\sigma(\nu_e e)$ (ALLEN 90). The result is shown in Fig. 1.

137 See Table I of DELAGUILA 89 for limits on various Z' models.

138 $g_\beta = e/\cos\theta_W$ and $\rho = 1$ assumed.

139 A wide range of neutral current data as of 1986 are used in the fit.

140 Z' mass limits from non-observation of an excess of $\ell^+\ell^-$ pairs at the CERN $p\bar{p}$ collider [based on ANSARI 87D and GEER Uppsala Conf. 87]. The limits apply when Z' decays only into light quarks and leptons.

MASS LIMITS for Leptoquarks from Pair Production

These limits rely only on the color or electromagnetic charge of the leptoquark.

VALUE (GeV)	CL%	EVS	DOCUMENT ID	TECN	COMMENT
>120	95		141 ABACHI	94B D0	First generation
> 80	95		142 ABE	93I CDF	First generation
> 45.5	95		143,144 ABREU	93J DLPH	First + second generation
> 44.4	95		145 ADRIANI	93M L3	First generation
> 44.5	95		145 ADRIANI	93M L3	Second generation
> 44.6	95		146 ADRIANI	93M L3	Third generation
> 44	95		145 DECAMP	92 ALEP	First or second generation
> 45	95		145 DECAMP	92 ALEP	Third generation
> 44.2	95		145 ALEXANDER	91 OPAL	First or second generation
> 41.4	95		145 ALEXANDER	91 OPAL	Third generation

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 42.1	95		147 ABREU	92F DLPH	Second generation
> 74	95		148 ALITTI	92E UA2	First generation
> 43.2	95		145 ADEVA	91B L3	First generation
> 43.4	95		145 ADEVA	91B L3	Second generation
none 8.9-22.6	95		149 KIM	90 AMY	First generation
none 10.2-23.2	95		149 KIM	90 AMY	Second generation
none 5-20.8	95		150 BARTEL	87B JADE	
none 7-20.5	95	2	151 BEHREND	86B CELL	

141 ABACHI 94B search for $eejj$ and $e\nu jj$ events in $p\bar{p}$ collisions at $E_{cm}=1.8$ TeV. The limit is for scalar leptoquarks decaying with $B(eq)=B(\nu q)=0.5$ and improves to >133 GeV for $B(eq)=1$. This limit does not depend on the electroweak quantum numbers of the leptoquark.

142 ABE 93I search for $\ell\ell jj$ events in $p\bar{p}$ collisions at $E_{cm}=1.8$ TeV. The limit is for $B(eq) = B(\nu q) = 0.5$ and improves to >113 GeV for $B(eq) = 1$. This limit does not depend on electroweak quantum numbers of the leptoquark.

143 Limit is for charge $-1/3$ isospin-0 leptoquark with $B(\ell q) = 2/3$.

144 First and second generation leptoquarks are assumed to be degenerate. The limit is slightly lower for each generation.

145 Limits are for charge $-1/3$, isospin-0 scalar leptoquarks decaying to $\ell^- q$ or νq with any branching ratio. See paper for limits for other charge-isospin assignments of leptoquarks.

146 ADRIANI 93M limit for charge $-1/3$, isospin-0 leptoquark decaying to τb .

147 ABREU 92F limit is for charge $-1/3$ isospin-0 leptoquark with $B(\mu q)=2/3$. If first and second generation leptoquarks are degenerate, the limit is 43.0 GeV, and for a charge 2/3 second generation leptoquark 43.4 GeV. Cross-section limit for pair production of states decaying to ℓq is given in the paper.

148 ALITTI 92E search for $\ell\ell jj$ and $\ell\nu jj$ events in $p\bar{p}$ collisions at $E_{cm}=630$ GeV. The limit is for $B(eq) = 1$ and is reduced to 67 GeV for $B(eq) = B(\nu q) = 0.5$. This limit does not depend on electroweak quantum numbers of the leptoquark.

149 KIM 90 assume pair production of charge 2/3 scalar-leptoquark via photon exchange. The decay of the first (second) generation leptoquark is assumed to be a mixture of d^+e^+ and ν^+d^+ ($s^+\mu^+$ and $c^+\nu^+$). See paper for limits for specific branching ratios.

150 BARTEL 87B limit is valid when a pair of charge 2/3 spinless leptoquarks X is produced with point coupling, and when they decay under the constraint $B(X \rightarrow c\bar{\nu}_\mu) + B(X \rightarrow s\mu^+) = 1$.

151 BEHREND 86B assumed that a charge 2/3 spinless leptoquark, χ , decays either into $s\mu^+$ or $c\nu$: $B(X \rightarrow s\mu^+) + B(X \rightarrow c\nu) = 1$.

MASS LIMITS for Leptoquarks from Single Production

These limits depend on the q - ℓ -leptoquark coupling g_{LQ} . It is often assumed that

$$g_{LQ}^2/4\pi = 1/137. \text{ Limits shown are for a scalar, weak isoscalar, charge } -1/3 \text{ leptoquark.}$$

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 73	95	152 ABREU	93J DLPH	Second generation
>181	95	153 ABT	93 H1	First generation

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 65	95	152 ABREU	93J DLPH	First generation
>168	95	154 DERRICK	93 ZEUS	First generation

152 Limit from single production in Z decay. The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(\ell q) = 2/3$. The limit is 77 GeV if first and second leptoquarks are degenerate.

153 ABT 93 search for single leptoquark production in $e p$ collisions with the decays $e q$ and μq . The limit is for a leptoquark coupling of electromagnetic strength and assumes $B(eq) = B(\nu q) = 1/2$. The limit for $B(eq) = 1$ is 178 GeV. For limits on states with different quantum numbers, see their Fig. 2.

Gauge & Higgs Boson Full Listings

Heavy Bosons Other than Higgs Bosons

154 DERRICK 93 search for single leptoquark production in ep collisions with the decay eq and νq . The limit is for leptoquark coupling of electromagnetic strength and assumes $B(eq) = B(\nu q) = 1/2$. The limit for $B(eq) = 1$ is 176 GeV. For limits on states with different quantum numbers, see their Table 3.

Indirect Limits for Leptoquarks

VALUE (TeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
>350	155 LEURER 93	RVUE FCNC	
> 1	156 DESHPANDE 83	RVUE Pati-Salam X-boson	
>125	157 SHANKER 82	RVUE PS leptoquark	
	157 SHANKER 82	RVUE Vector-leptoquark	
155 LEURER 93 gives bounds for generation-changing couplings of scalar leptoquark from $K^0, \bar{K}^0, D^0, \bar{D}^0, B^0, \bar{B}^0$ mixing. See her Fig. 1.			
156 DESHPANDE 83 used upper limit on $K_L^0 \rightarrow \mu e$ decay with renormalization-group equations to estimate coupling at the heavy boson mass. See also DIMOPOULOS 81.			
157 From $(\pi \rightarrow e\nu)/(\pi \rightarrow \mu\nu)$ ratio.			

MASS LIMITS for g_A (axigluon)

Axigluons are massive color-octet gauge bosons in chiral color models and have axial-vector coupling to quarks with the same coupling strength as gluons.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 240-640	95	158 ABE	93G CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$ ets
>50	95	159 CUYPERS 91	RVUE	$\sigma(e^+e^- \rightarrow \text{hadrons})$
none 120-210	95	160 ABE	90H CDF	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$ ets
>29		161 ROBINETT 89	THEO	Partial-wave unitarity
none 150-310	95	162 ALBAJAR 88B	UA1	$p\bar{p} \rightarrow g_A X, g_A \rightarrow 2j$ ets
>20		BERGSTROM 88	RVUE	$p\bar{p} \rightarrow \gamma X$ via $g_A g$
> 9		163 CUYPERS 88	RVUE	γ decay
>25		164 DONCHESKI 88B	RVUE	γ decay
158 ABE 93G assume $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 10$.				
159 CUYPERS 91 compare α_s measured in γ decay and that from R at PEP/PETRA energies.				
160 ABE 90H assumes $\Gamma(g_A) = N\alpha_s m_{g_A}/6$ with $N = 5$ ($\Gamma(g_A) = 0.09m_{g_A}$). For $N = 10$, the excluded region is reduced to 120-150 GeV.				
161 ROBINETT 89 result demands partial-wave unitarity of $J = 0$ $t\bar{t} \rightarrow t\bar{t}$ scattering amplitude and derives a limit $m_{g_A} > 0.5 m_{t^*}$. Assumes $m_{t^*} > 56$ GeV.				
162 ALBAJAR 88B result is from the nonobservation of a peak in two-jet invariant mass distribution. $\Gamma(g_A) < 0.4 m_{g_A}$ assumed. See also BAGGER 88.				
163 CUYPERS 88 requires $\Gamma(\gamma \rightarrow g g_A) < \Gamma(\gamma \rightarrow g g g)$. A similar result is obtained by DONCHESKI 88.				
164 DONCHESKI 88B requires $\Gamma(\gamma \rightarrow g q\bar{q})/\Gamma(\gamma \rightarrow g g g) < 0.25$, where the former decay proceeds via axigluon exchange. A more conservative estimate of < 0.5 leads to $m_{g_A} > 21$ GeV.				

X^0 (Heavy Boson) Searches in Z Decays

Searches for radiative transition of Z to a lighter spin-0 state X^0 decaying to hadrons, a lepton pair, or a photon pair as shown in the comments. The limits are for the product of branching ratios.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		165 ACTON	93E OPAL	$X^0 \rightarrow \gamma\gamma$
		166 ABREU	92D DLPH	$X^0 \rightarrow \text{hadrons}$
		167 ADRIANI	92F L3	$X^0 \rightarrow \text{hadrons}$
		168 ACTON	91 OPAL	$X^0 \rightarrow \text{anything}$
$< 1.1 \times 10^{-4}$	95	169 ACTON	91B OPAL	$X^0 \rightarrow e^+e^-$
$< 9 \times 10^{-5}$	95	169 ACTON	91B OPAL	$X^0 \rightarrow \mu^+\mu^-$
$< 1.1 \times 10^{-4}$	95	169 ACTON	91B OPAL	$X^0 \rightarrow \tau^+\tau^-$
$< 2.8 \times 10^{-4}$	95	170 ADEVA	91D L3	$X^0 \rightarrow e^+e^-$
$< 2.3 \times 10^{-4}$	95	170 ADEVA	91D L3	$X^0 \rightarrow \mu^+\mu^-$
$< 4.7 \times 10^{-4}$	95	171 ADEVA	91D L3	$X^0 \rightarrow \text{hadrons}$
$< 8 \times 10^{-4}$	95	172 AKRAWY	90J OPAL	$X^0 \rightarrow \text{hadrons}$
165 ACTON 93E give $\sigma(e^+e^- \rightarrow X^0\gamma) \cdot B(X^0 \rightarrow \gamma\gamma) < 0.4$ pb (95%CL) for $m_{X^0} = 60 \pm 2.5$ GeV. If the process occurs via s-channel γ exchange, the limit translates to $\Gamma(X^0) \cdot B(X^0 \rightarrow \gamma\gamma)^2 < 20$ MeV for $m_{X^0} = 60 \pm 1$ GeV.				
166 ABREU 92D give $\sigma_Z \cdot B(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < (3-10)$ pb for $m_{X^0} = 10-78$ GeV. A very similar limit is obtained for spin-1 X^0 .				
167 ADRIANI 92F search for isolated γ in hadronic Z decays. The limit $\sigma_Z \cdot B(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < (2-10)$ pb (95%CL) is given for $m_{X^0} = 25-85$ GeV.				
168 ACTON 91 searches for $Z \rightarrow Z^* X^0$, $Z^* \rightarrow e^+e^-, \mu^+\mu^-,$ or $\nu\bar{\nu}$. Excludes any new scalar X^0 with $m_{X^0} < 9.5$ GeV/c if it has the same coupling to $Z Z^*$ as the MSM Higgs boson.				
169 ACTON 91B limits are for $m_{X^0} = 60-85$ GeV.				
170 ADEVA 91D limits are for $m_{X^0} = 30-89$ GeV.				
171 ADEVA 91D limits are for $m_{X^0} = 30-86$ GeV.				

172 AKRAWY 90J give $\Gamma(Z \rightarrow \gamma X^0) \cdot B(X^0 \rightarrow \text{hadrons}) < 1.9$ MeV (95%CL) for $m_{X^0} = 32-80$ GeV. We divide by $\Gamma(Z) = 2.5$ GeV to get product of branching ratios. For nonresonant transitions, the limit is $B(Z \rightarrow \gamma q\bar{q}) < 8.2$ MeV assuming three-body phase space distribution.

MASS LIMITS for a Heavy Neutral Boson Coupling to e^+e^-

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 55-61		173 ODAKA	89 VNS	$\Gamma(X^0 \rightarrow e^+e^-) \cdot B(X^0 \rightarrow \text{hadrons}) > 0.2$ MeV
>45	95	174 DERRICK	86 HRS	$\Gamma(X^0 \rightarrow e^+e^-) = 6$ MeV
>46.6	95	175 ADEVA	85 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-) = 10$ keV
>48	95	175 ADEVA	85 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-) = 4$ MeV
		176 BERGER	85S PLUT	
none 39.8-45.5		177 ADEVA	84 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-) = 10$ keV
>47.8	95	177 ADEVA	84 MRKJ	$\Gamma(X^0 \rightarrow e^+e^-) = 4$ MeV
none 39.8-45.2		177 BEHREND	84C CELL	
>47	95	177 BEHREND	84C CELL	$\Gamma(X^0 \rightarrow e^+e^-) = 4$ MeV
173 ODAKA 89 looked for a narrow or wide scalar resonance in $e^+e^- \rightarrow \text{hadrons}$ at $E_{cm} = 55.0-60.8$ GeV.				
174 DERRICK 86 found no deviation from the Standard Model Bhabha scattering at $E_{cm} = 29$ GeV and set limits on the possible scalar boson e^+e^- coupling. See their figure 4 for excluded region in the $\Gamma(X^0 \rightarrow e^+e^-) \cdot m_{X^0}$ plane. Electronic chiral invariance requires a parity doublet of X^0 , in which case the limit applies for $\Gamma(X^0 \rightarrow e^+e^-) > 3$ MeV.				
175 ADEVA 85 first limit is from $2\gamma, \mu^+\mu^-,$ hadrons assuming X^0 is a scalar. Second limit is from e^+e^- channel. $E_{cm} = 40-47$ GeV. Supersedes ADEVA 84.				
176 BERGER 85B looked for effect of spin-0 boson exchange in $e^+e^- \rightarrow e^+e^-$ and $\mu^+\mu^-$ at $E_{cm} = 34.7$ GeV. See Fig. 5 for excluded region in the $m_{X^0} - \Gamma(X^0)$ plane.				
177 ADEVA 84 and BEHREND 84C have $E_{cm} = 39.8-45.5$ GeV. MARK-J searched X^0 in $e^+e^- \rightarrow \text{hadrons}, 2\gamma, \mu^+\mu^-, e^+e^-$ and CELLO in the same channels plus τ pair. No narrow or broad X^0 is found in the energy range. They also searched for the effect of X^0 with $m_X > E_{cm}$. The second limits are from Bhabha data and for spin-0 singlet. The same limits apply for $\Gamma(X^0 \rightarrow e^+e^-) = 2$ MeV if X^0 is a spin-0 doublet. The second limit of BEHREND 84C was read off from their figure 2. The original papers also list limits in other channels.				

Search for X^0 Resonance in e^+e^- Collisions

The limit is for $\Gamma(X^0 \rightarrow e^+e^-) \cdot B(X^0 \rightarrow f)$, where f is the specified final state. Spin 0 is assumed for X^0 .

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 10^3$	95	178 ABE	93C VNS	$f = (ee)$
$< (0.4-10)$	95	179 ABE	93C VNS	$f = \gamma\gamma$
$< (0.3-5)$	95	180,181 ABE	93D TOPZ	$f = \gamma\gamma$
$< (2-12)$	95	180,181 ABE	93D TOPZ	$f = \text{hadrons}$
$< (4-200)$	95	181,182 ABE	93D TOPZ	$f = ee$
$< (0.1-6)$	95	181,182 ABE	93D TOPZ	$f = \mu\mu$
$< (0.5-8)$	90	183 STERNER	93 AMY	$f = \gamma\gamma$
178 Limit is for $\Gamma(X^0 \rightarrow e^+e^-) m_{X^0} = 56-63.5$ GeV for $\Gamma(X^0) = 0.5$ GeV.				
179 Limit is for $m_{X^0} = 56-61.5$ GeV and is valid for $\Gamma(X^0) \ll 100$ MeV. See their Fig. 5 for limits for $\Gamma = 1, 2$ GeV.				
180 Limit is for $m_{X^0} = 57.2-60$ GeV.				
181 Limit is valid for $\Gamma(X^0) \ll 100$ MeV. See paper for limits for $\Gamma = 1$ GeV and those for $J = 2$ resonances.				
182 Limit is for $m_{X^0} = 56.6-60$ GeV.				
183 STERNER 93 limit is for $m_{X^0} = 57-59.6$ GeV and is valid for $\Gamma(X^0) < 100$ MeV. See their Fig. 2 for limits for $\Gamma = 1, 3$ GeV.				

Search for X^0 Resonance in Two-Photon Process

The limit is for $\Gamma(X^0) \cdot B(X^0 \rightarrow \gamma\gamma)^2$. Spin 0 is assumed for X^0 .

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 2.6	95	184 ACTON	93E OPAL	$m_{X^0} = 60 \pm 1$ GeV
< 2.9	95	BUSKULIC	93F ALEP	$m_{X^0} \sim 60$ GeV
184 ACTON 93E limit for a $J = 2$ resonance is 0.8 MeV.				

Search for X^0 Resonance in $Z \rightarrow f\bar{f}X^0$

The limit is for $B(Z \rightarrow f\bar{f}X^0) \cdot B(X^0 \rightarrow F)$ where f is a fermion and F is the specified final state. Spin 0 is assumed for X^0 .

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 6.8 \times 10^{-6}$	95	185 ACTON	93E OPAL	$f = e, \mu, \tau; F = \gamma\gamma$
$< 5.5 \times 10^{-6}$	95	185 ACTON	93E OPAL	$f = q; F = \gamma\gamma$
$< 3.1 \times 10^{-6}$	95	185 ACTON	93E OPAL	$f = \nu; F = \gamma\gamma$
$< 6.5 \times 10^{-6}$	95	185 ACTON	93E OPAL	$f = e, \mu; F = \ell\bar{\ell}, q\bar{q}, \nu\bar{\nu}$
$< 7.1 \times 10^{-6}$	95	185 BUSKULIC	93F ALEP	$f = e, \mu; F = \ell\bar{\ell}, q\bar{q}, \nu\bar{\nu}$
		186 ADRIANI	92F L3	$f = q; F = \gamma\gamma$

See key on page 1343

Gauge & Higgs Boson Full Listings

Heavy Bosons Other than Higgs Bosons, Axions (A^0) and Other Very Light Bosons

185 Limit is for m_{X^0} around 60 GeV.

186 ADRIANI 92F give $\sigma_{\gamma} \cdot B(Z \rightarrow q\bar{q}X^0) \cdot B(X^0 \rightarrow \gamma\gamma) < (0.75-1.5) \text{ pb}$ (95%CL) for $m_{X^0} = 10-70 \text{ GeV}$. The limit is 1 pb at 60 GeV.

Heavy Particle Production in Quarkonium Decays

Limits are for branching ratios to modes shown.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.6 \times 10^{-5}$	90	187 ANTREASYAN 90C CBAL		$\Upsilon(1S) \rightarrow X^0 \gamma$ $m_{X^0} < 7.2 \text{ GeV}$
		188 ALBRECHT 89 ARG		

187 ANTREASYAN 90C assume that X^0 does not decay in the detector.

188 ALBRECHT 89 give limits for $B(\Upsilon(1S), \Upsilon(2S) \rightarrow X^0 \gamma) \cdot B(X^0 \rightarrow \pi^+ \pi^-, K^+ K^-, p\bar{p})$ for $m_{X^0} < 3.5 \text{ GeV}$.

REFERENCES FOR Searches for Heavy Bosons Other Than Higgs Bosons

ABACHI 94B	PRL 72 965	+Abbott, Abolins, Acharya, Adam+	(D0 Collab.)
BUSKULIC 94	ZPHY C62 539	+Casper, De Bonis, Decamp, Ghez, Goy+	(ALEPH Collab.)
ABE 93C	PL B302 119	+Amako, Arai, Arima, Asano, Chiba+	(VENUS Collab.)
ABE 93D	PL B304 373	+Adachi, Awa, Aoki, Belusevic, Emi+	(TOPAZ Collab.)
ABE 93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+	(CDF Collab.)
ABE 93I	PR D48 R3939	+Albrow, Amidei, Anway-Wiese, Apollinari+	(CDF Collab.)
ABREU 93J	PL B316 620	+Ady, Aduy, Agasi, Aleksan, Alekseev+	(DELPHI Collab.)
ABT 93	NP B396 3	+Andreev, Andrieu, Appuhn, Arpagaus+	(H1 Collab.)
ACTON 93E	PL B311 391	+Akers, Alexander+	(OPAL Collab.)
ADRIANI 93D	PL B306 187	+Aguiar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ADRIANI 93M	PRPL 236 1	+Aguiar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ALITTI 93	NP B400 3	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
ALLEN 93	PR D47 11	+Chen, Doe, Hausammann+	(UCI, LANL, ANL, UMD)
ALTARELLI 92D	PL B318 139	+Casalbuoni+ (CERN, FIRZ, GEVA, PADO)	
BHATTACH... 93	PR D47 R3693	+Bhattacharyya+ (CALC, JADA, ICTP, AHMED, BOSE)	
BUSKULIC 93F	PL B308 425	+De Bonis, Decamp, Chez, Goy, Lees+	(ALEPH Collab.)
DERRICK 93	PL B306 173	+Krackauer, Magill, Musgrave, Repond+	(ZEUS Collab.)
LEURER 93	PRL 71 1324		(REHO)
RIZZO 93	PR D48 4470		(ANL)
SEVERIJNS 93	PRL 70 4047	+Gimeno-Nogues+ (LVLN, WISC, LEUV, ETH, MASA)	
STERNER 93	PL B303 385	+Abashian, Gotow, Halm, Mattson, Morgan+(AMY Collab.)	
ABE 92B	PRL 68 1463	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ABREU 92D	ZPHY C53 555	+Adam, Adami, Aduy, Akesson, Alekseev+(DELPHI Collab.)	
ABREU 92F	PL B275 222	+Adam, Adami, Aduy, Akesson, Alekseev+(DELPHI Collab.)	
ADRIANI 92F	PL B292 472	+Aguiar-Benitez, Ahlen, Akbari, Alcaraz+	(L3 Collab.)
ALITTI 92E	PL B274 507	+Ambrosini, Ansari, Autiero, Bareyre+	(UA2 Collab.)
DECAMP 92	PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
DELAGUILA 92	NP B372 3	+del Aguilá, Moreno, Quiros (CERN, GRAN, MPIM, BRUXT, MADE)	
Also 91C	NP B361 45	+del Aguilá, Moreno, Quiros (BARC, MADE)	
IMAZATO 92	PR D49 877	+Kawashima, Tanaka+ (KEK, INUS, TOKY, TOKMS)	
LANGACKER 92B	PR D45 278	+Luo	(FENNI)
LAYSAC 92	ZPHY C53 97	+Renard, Verzegnassi (MONP, LAPP)	
LAYSAC 92B	PL B287 267	+Renard, Verzegnassi (MONP, TRSTT)	
LEIKE 92	PL B291 187	+Riemann, Riemann (BERL, CERN)	
LEP 92	PL B276 247	+ALEPH, DELPHI, L3, OPAL (LEP Collabs.)	
MISHRA 92	PRL 68 3499	+Leung, Arroyo+ (COLU, CHIC, FNAL, ROCH, WISC)	
POLAK 92	PL B276 492	+Zralek (SILES)	
POLAK 92B	PR D46 3871	+Zralek (SILES)	
RENTON 92	ZPHY C56 355		(OXF)
ABE 91D	PRL 67 2418	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)	
ABE 91F	PRL 67 2609	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)	
ACTON 91	PL B268 122	+Alexander, Allison, Allport+ (OPAL Collab.)	
ACTON 91B	PL B273 338	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)	
ADEVA 91B	PL B261 169	+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)	
ADEVA 91D	PL B262 155	+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)	
ALEXANDER 91	PL B263 123	+Allison, Allport, Anderson, Arcelli+ (OPAL Collab.)	
ALITTI 91	ZPHY C49 17	+Ansari, Ansoorge, Autiero, Bareyre+ (UA2 Collab.)	
ALTARELLI 91	PL B261 146	+Casalbuoni, De Curtis+ (CERN, FIRZ, GEVA)	
ALTARELLI 91B	PL B263 459	+Casalbuoni, De Curtis+ (CERN, FIRZ, GEVA)	
Also 90	PL B245 669	+Altarelli, Casalbuoni, Feruglio, Gatto(CERN, LECE, GEVA)	
AQUINO 91	PL B261 280	+Fernandez, Garcia (CINV, PUEB)	
BUCHMUELL... 91	PL B267 395	+Buchmueller, Greub, Minkowski (DESY, BERN)	
COLANGEL... 91	PL B262 155	+Nardulli (BAR)	
CUYPERS 91	PL B259 173	+Falk, Frampton (DURH, HARV, UNCCH)	
DELAGUILA 91	PL B254 497	+del Aguilá, Moreno, Quiros (BARC, MADE, CERN)	
FARAGGI 91	MPL A6 61	+Nanopoulos (TAMU)	
GEIREGAT 91	PL B259 499	+Vilain, Wilquet, Binder, Burkard+ (CHARM II Collab.)	
GONZALEZ-G... 91	PL B259 365	+Gonzalez-Garcia, Valle (VALE)	
Also 90C	NP B345 312	+Gonzalez-Garcia, Valle (VALE)	
MAHANATHAP... 91	PR D43 3093	+Mahanthappa, Mohapatra (COLO)	
Also 91B	PR D44 1516 erratum	+Mahanthappa, Mohapatra (COLO)	
POLAK 91	NP B363 385	+Zralek (SILES)	
RIZZO 91	PR D44 202		(WISC, ISU)
WALKER 91	APJ 376 51	+Steigman, Schramm, Olive+ (HSCA, OSU, CHIC, MINN)	
ABE 90F	PL B246 297	+Amako, Arai, Asano, Chiba+ (VENUS Collab.)	
ABE 90G	PRL 65 2243	+Amidei, Apollinari, Atac+ (CDF Collab.)	
ABE 90H	PR D41 1722	+Amidei, Apollinari, Ascoli, Atac+ (CDF Collab.)	
AKRAWAY 90J	PL B246 285	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)	
ALLEN 90	PRL 64 1330	+Chen, Doe+ (UCI, LASL, UMD)	
ANTREASYAN 90C	PL B251 204	+Bartels, Besset, Bieler, Bienlein+ (Crystal Ball Collab.)	
BARGER 90B	PR D42 152	+Hewett, Rizzo (WISC, ISU)	
GLASHOW 90	PR D42 3224	+Sarif (HARV)	
GLASHOW 90B	PRL 64 725	+Sarif (HARV)	
GONZALEZ-G... 90D	PL B240 163	+Gonzalez-Garcia, Valle (VALE)	
GRIFOLS 90	NP B331 244	+Masse (BARC)	
GRIFOLS 90C	MPL A5 2657	+Masse (BARC)	
GRIFOLS 90D	PR D42 3293	+Masse, Rizzo (BARC, CERN, WISC, ISU)	
HAGIWARA 90	PR D41 815	+Najima, Sakuda, Terunuma (KEK, DURH, YCC, HIRO)	
HE 90B	PL B240 441	+Joshi, Volkas (MELB)	
Also 90C	PL B244 580 erratum	+He, Joshi, Volkas (MELB)	
KIM 90	PL B240 243	+Breedon, Ko, Lander, Maeshima, Malchow+(AMY Collab.)	
LOPEZ 90	PL B241 392	+Nanopoulos (TAMU)	
ALBAJAR 89	ZPHY C44 15	+Albrow, Altkofer, Arnison, Astbury+ (UA1 Collab.)	
ALBRECHT 89	ZPHY C42 349	+Boeckmann, Glaeser, Harder+ (ARGUS Collab.)	
BARBIERI 89B	PR D39 1229	+Mohapatra (PISA, UMD)	
DELAGUILA 89	PR D40 2481	+del Aguilá, Moreno, Quiros (BARC, MADE)	
Also 89B	PR D41 134	+del Aguilá, Moreno, Quiros (BARC, MADE)	
Also 90C	PR D42 262 erratum	+del Aguilá, Moreno, Quiros (BARC, MADE)	
DORENBOS... 89	ZPHY C41 567	+Dorenbosch, Udo, Allaby, Amaldi+ (CHARM Collab.)	
GEIREGAT 89	PL B232 539	+Vilain, Wilquet, Bergsma, Binder+ (CHARM II Collab.)	
LANGACKER 89B	PR D40 1569	+Uma Sankar (PENN)	

ODAKA 89	JPSJ 58 3037	+Kondo, Abe, Amako+ (VENUS Collab.)	
ROBINETT 89	PR D39 834		(PSU)
ALBAJAR 89B	PL B209 127	+Albrow, Altkofer, Astbury, Aubert+ (UA1 Collab.)	
BARGER 88	PR D37 1188	+Schmidt, King (HARV, BOST)	
BALKE 88	PR D37 587	+Gidal, Jodidio+ (LBL, UCB, COLO, NWES, TRIU)	
BERGSTROM 88	PL B212 386		(STOH)
COSTA 88	NP B297 244	+Ellis, Fogli+ (PADO, CERN, BARI, WISC, LBL)	
CUYPERS 88	PRL 60 1237	+Frampton (UNCCH)	
DONCHESKI 88	PL B206 137	+Grotch, Robinett (PSU)	
DONCHESKI 88B	PR D38 412	+Grotch, Robinett (PSU)	
ELLIS 88	PL B202 417	+Ellis, Franzini, Zwirner (CERN, UCB, LBL)	
RAFFEL 88	PRL 60 1793	+Seckel (UCB, LLL, UCSC)	
AMALDI 87	PR D36 1385	+Bohm, Durkin, Langacker+ (CERN, AACH3, OSU+)	
ANSARI 87D	PL B195 613	+Bagnaia, Banner+ (UA2 Collab.)	
BARTEL 87B	ZPHY C36 15	+Becker, Feist+ (JADE Collab.)	
MARCIANO 87	PR D35 1672	+Sirlin (BNL, NYU)	
ARNISON 86B	EPL 1 327	+Albrow, Altkofer+ (UA1 Collab.)	
BARGER 86B	PRL 56 30	+Deshpande, Whisnant (WISC, OREG, FSU)	
BEHREND 86	PL B178 452	+Buerger, Criegee, Fenner, Field+ (CELLO Collab.)	
DERRICK 86	PL 166B 463	+Gan, Kooijman, Loos+ (HRS Collab.)	
Also 86B	PR D34 3286	+Derrick, Gan, Kooijman, Loos, Musgrave+ (HRS Collab.)	
DURKIN 86	PL 166B 436	+Langacker (PENN)	
ELLIS 86	PL 167B 457	+Enqvist, Nanopoulos, Sarkar (CERN, OXFTR)	
JODIDIO 86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+ (LBL, NWES, TRIU)	
Also 88	PR D37 237 erratum	+Jodidio, Balke, Carr+ (LBL, NWES, TRIU)	
MOHAPATRA 86	PR D34 909		(UMD)
STEIGMAN 86	PL B216 33	+Olive, Schramm, Turner (BART, MIM+)	
ADEVA 85	PL 152B 439	+Becker, Becker-Szendy+ (Mark-J Collab.)	
ADEVA 85B	PRL 55 665	+Becker, Becker-Szendy+ (Mark-J Collab.)	
BERGER 85B	ZPHY C27 341	+Deuter, Genzel, Lackas, Pielorz+ (PLUTO Collab.)	
STOKER 85	PRL 54 1887	+Balke, Carr, Gidal+ (LBL, NWES, TRIU)	
ADEVA 84	PRL 53 134	+Barber, Becker, Berdugo+ (Mark-J Collab.)	
BEHREND 84C	PL 140B 130	+Burger, Criegee, Fenner+ (CELLO Collab.)	
ARMISTEAD 83D	PL 129B 273	+Astbury, Aubert, Bacchi+ (UA1 Collab.)	
BERGSM 83	PL 122B 465	+Dorenbosch, Jonker+ (CHARM Collab.)	
CARR 83	PRL 51 627	+Gidal, Gobbi, Jodidio, Oram+ (LBL, NWES, TRIU)	
DESHPANDE 83	PR D27 1193	+Johnson (OREG)	
BEALL 82	PRL 48 848	+Bander, Soni (UCI, UCLA)	
SHANKER 82	NP B204 375		(TRIU)
DIMOPOUL... 81	NP B182 77	+Dimopoulos, Raby, Kane (STAN, MICH)	
RIZZO 81	PR D24 704	+Senjanovic (BNL)	
STEIGMAN 79	PRL 43 239	+Olive, Schramm (BART, EFI)	

Searches for Axions (A^0) and Other Very Light Bosons

NOTE ON AXIONS

In this section we list limits for very light neutral (pseudo) scalar bosons that couple weakly to stable matter. Typical examples are pseudo-Goldstone bosons like axions (A^0) [1], familons [2], and Majorons [3], associated, respectively, with spontaneously broken Peccei-Quinn [4], family, and lepton-number symmetries.

Peccei-Quinn symmetry gives a natural solution to the strong CP-violation problem. Axion mass and its coupling to stable particles are inversely proportional to the scale of the Peccei-Quinn symmetry breaking f_A . The original axion model [1,4] assumes $f_A = v$, where $v = (\sqrt{2}G_F)^{-1/2} = 247 \text{ GeV}$ is the scale of the electroweak symmetry breaking, and has two Higgs doublets as minimal ingredients. By requiring tree-level flavor conservation, the axion mass and its couplings are completely fixed in terms of one parameter, the ratio of the vacuum expectation values of two Higgs fields. The result of extensive experimental searches for such an axion have been negative [5].

One way to avoid these experimental constraints is to make A^0 sufficiently massive. This is achieved by introducing a new strong interaction (QC'D) with $\Lambda_{QC'D} \gg \Lambda_{QCD}$, whose anomaly couples to the axion [7]. A^0 can receive significant mass from the QC'D sector if QC'D colored quarks are massive. However one needs an explanation why $|\theta_{QCD} - \theta_{QC'D}| < 10^{-9}$ in this scenario.

Another way to save the Peccei-Quinn idea is to introduce a new scale $f_A \gg v$. Then the A^0 mass becomes smaller and its coupling weaker, thus one can easily avoid all the existing experimental limits; such models are called invisible axion models [8,9]. Various invisible axion models can be

Gauge & Higgs Boson Full Listings

Axions (A^0) and Other Very Light Bosons

constructed by identifying f_A with other large mass scales such as the Planck mass, the GUT scale, the SUSY-breaking scale, and so on. It has been found, however, that invisible axions are not completely elusive. Cosmological considerations on the matter density of our universe suggest [10] $f_A < \mathcal{O}(10^{12})$ GeV as a possible upper bound on the scale. Lower bounds of $f_A > \mathcal{O}(10^7)$ GeV are obtained from astrophysics [11], where axion emission from the center of stellar objects can speed up their evolutionary time scales. The recent observation of the supernova SN1987A improves the lower bound to $f_A > \mathcal{O}(10^{10})$ GeV. Various terrestrial experiments to detect invisible axions by making use of their coupling to photons have been proposed [12], and the first result of such experiments appeared recently.

Observation of a narrow-peak structure in positron spectra from heavy ion collisions [13] suggested a particle of mass 1.8 MeV that decays into e^+e^- . Variants of the original axion model, which keep $f_A = v$, but drop the constraints of tree-level flavor conservation, were proposed [14]. Extensive searches for this particle, $A^0(1.8 \text{ MeV})$, ended up with another negative result [6].

There is also a Note on invisible axions later in this section.

References

- S. Weinberg, Phys. Rev. Lett. **40**, 223 (1978);
F. Wilczek, Phys. Rev. Lett. **40**, 279 (1978).
 - F. Wilczek, Phys. Rev. Lett. **49**, 1549 (1982).
 - Y. Chikashige, R.N. Mohapatra, and R.D. Peccei, Phys. Lett. **98B**, 265 (1981);
G.B. Gelmini and M. Roncadelli, Phys. Lett. **99B**, 411 (1981).
 - R.D. Peccei and H. Quinn, Phys. Rev. Lett. **38**, 1440 (1977), also Phys. Rev. **D16**, 1791 (1977).
 - T.W. Donnelly *et al.*, Phys. Rev. **D18**, 1607 (1978);
S. Barshay *et al.*, Phys. Rev. Lett. **46**, 1361 (1981);
A. Barroso N. Mukhopadhyay, Phys. Lett. **106B**, 91 (1981);
R.D. Peccei, in *Proceedings of Neutrino '81*, Honolulu, Hawaii, Vol. 1, p. 149 (1981);
L.M. Krauss and F. Wilczek, Phys. Lett. **B173**, 189 (1986).
 - W.A. Bardeen, R.D. Peccei, and T. Yanagida, Nucl. Phys. **B279**, 401 (1987).
 - S.H.H. Tye, Phys. Rev. Lett. **47**, 1035 (1981).
 - J.E. Kim, Phys. Rev. Lett. **43**, 103 (1979);
M.A. Shifman, A.I. Vainstein, and V.I. Zakharov, Nucl. Phys. **B166**, 493 (1980).
 - A.R. Zhitnitsky, Sov. J. Nucl. Phys. **31**, 260 (1980);
M. Dine, W. Fischler, and M. Srednicki, Phys. Lett. **104B**, 199 (1981).
 - J. Preskill *et al.*, Phys. Lett. **120B**, 127 (1983);
L.F. Abbott, and P. Sikivie, Phys. Lett. **120B**, 133 (1983);
M. Dine and W. Fischler, Phys. Lett. **120B**, 137 (1983).
 - D.A. Dicus *et al.*, Phys. Rev. **D18**, 1829 (1978);
M. Fukugita, S. Watamura, and M. Yoshimura, Phys. Rev. Lett. **48**, 1522 (1982), also Phys. Rev. **D26**, 1840 (1982);
D.S.P. Dearborn *et al.*, Phys. Rev. Lett. **56**, 26 (1986);
G.G. Raffelt, Phys. Rev. **D33**, 897 (1986);
- For references to SN 1987a, see reviews by M.S. Turner, Phys. Rept. 197,67(1990);
G.G. Raffelt, Phys. Rept. 198,1(1990)..
- P. Sikivie, Phys. Rev. Lett. **51**, 1415 (1983), also Phys. Rev. **D32**, 2988 (1985).
 - J. Schweppe *et al.*, Phys. Rev. Lett. **51**, 2261 (1983);
T. Cowan *et al.*, Phys. Rev. Lett. **54**, 1761 (1985).
 - R.D. Peccei, T.T. Wu, and T. Yanagida, Phys. Lett. **B172**, 435 (1986).

A^0 (Axion) MASS LIMITS from Astrophysics and Cosmology

These bounds depend on model-dependent assumptions (i.e. — on a combination of axion parameters).

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
>0.2	BARROSO 82	ASTR	Standard Axion
>0.25	¹ RAFFELT 82	ASTR	Standard Axion
>0.2	² DICUS 78C	ASTR	Standard Axion
	MIKAELIAN 78	ASTR	Stellar emission
>0.3	² SATO 78	ASTR	Standard Axion
>0.2	VVSOTSKII 78	ASTR	Standard Axion

¹ Lower bound from 5.5 MeV γ -ray line from the sun.

² Lower bound from requiring the red giants' stellar evolution not be disrupted by axion emission.

A^0 (Axion) and Other Light Boson (X^0) Searches in Stable Particle Decays

Limits are for branching ratios.

VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT
<1.7 $\times 10^{-9}$	90	³ ATIYA 93	CNTR	$K^+ \rightarrow \pi^+ A^0$
<2 $\times 10^{-7}$	90	⁴ ATIYA 93B	CNTR	$K^+ \rightarrow \pi^+ A^0$
<3 $\times 10^{-13}$		⁵ NG 93	COSM	$\pi^0 \rightarrow \gamma X^0$
<1.1 $\times 10^{-8}$	90	⁶ ALLIEGRO 92	CALO	$K^+ \rightarrow \pi^+ A^0$ ($A^0 \rightarrow e^+e^-$)
<5 $\times 10^{-4}$	90	⁷ ATIYA 92	CNTR	$\pi^0 \rightarrow \gamma X^0$
<4 $\times 10^{-6}$	90	⁸ MEIJERDREES 92	SPEC	$\pi^0 \rightarrow \gamma X^0$ $X^0 \rightarrow e^+e^-$ $m_{X^0} = 100 \text{ MeV}$
<1 $\times 10^{-7}$	90	⁹ ATIYA 90B	CNTR	$K^+ \rightarrow \pi^+ A^0$ ($A^0 \rightarrow \gamma\gamma$)
<1.3 $\times 10^{-8}$	90	¹⁰ KORENCHENKO 87	SPEC	$\pi^+ \rightarrow e^+ \nu A^0$ ($A^0 \rightarrow e^+e^-$)
<1 $\times 10^{-9}$	90	0 ¹¹ EICHLER 86	SPEC	Stopped $\pi^+ \rightarrow e^+ \nu A^0$
<2 $\times 10^{-5}$	90	¹² YAMAZAKI 84	SPEC	For $160 < m_{A^0} < 260 \text{ MeV}$
<(1.5-4) $\times 10^{-6}$	90	¹² YAMAZAKI 84	SPEC	K decay, $m_{A^0} \ll 100 \text{ MeV}$
	0	¹³ ASANO 82	CNTR	Stopped $K^+ \rightarrow \pi^+ A^0$
	0	¹⁴ ASANO 81B	CNTR	Stopped $K^+ \rightarrow \pi^+ A^0$
		¹⁵ ZHITNITSKII 79		Heavy axion

³ ATIYA 93 looked for a peak in missing mass distribution. The limit is for massless stable A^0 particles and extends to $m_{A^0} = 80 \text{ MeV}$ at the same level. See paper for dependence on finite lifetime.

⁴ ATIYA 93B looked for a peak in missing mass distribution. The bound applies for stable A^0 of $m_{A^0} = 150-250 \text{ MeV}$, and the limit becomes stronger (10^{-8}) for $m_{A^0} = 180-240 \text{ MeV}$.

⁵ NG 93 studied the production of X^0 via $\gamma\gamma \rightarrow \pi^0 \rightarrow \gamma X^0$ in the early universe at $T \sim 1 \text{ MeV}$. The bound on extra neutrinos from nucleosynthesis $\Delta N_\nu < 0.3$ (WALKER 91) is employed. It applies to $m_{X^0} \ll 1 \text{ MeV}$ in order to be relativistic down to nucleosynthesis temperature. See paper for heavier X^0 .

⁶ ALLIEGRO 92 limit applies for $m_{A^0} = 150-340 \text{ MeV}$, and $B(A^0 \rightarrow e^+e^-) = 100\%$ assumed. Limit is $< 1.5 \times 10^{-8}$ at 99%CL.

⁷ ATIYA 92 looked for a peak in missing mass distribution. The limit applies to $m_{X^0} = 0-130 \text{ MeV}$ in the narrow resonance limit. See paper for the dependence on lifetime. Covariance requires X^0 to be a vector particle.

⁸ MEIJERDREES 92 limit applies for $\tau_{X^0} = 10^{-23}-10^{-11} \text{ sec}$. Limits between 2×10^{-4} and 4×10^{-6} are obtained from $m_{X^0} = 25-120 \text{ MeV}$. Angular momentum conservation requires that X^0 has spin ≥ 1 .

⁹ ATIYA 90B limit is for $B(K^+ \rightarrow \pi^+ A^0) \cdot B(A^0 \rightarrow \gamma\gamma)$ and applies for $m_{A^0} = 50 \text{ MeV}$, $\tau_{A^0} < 10^{-10} \text{ s}$. Limits are also provided for $0 < m_{A^0} < 100 \text{ MeV}$, $\tau_{A^0} < 10^{-8} \text{ s}$.

¹⁰ KORENCHENKO 87 limit assumes $m_{A^0} = 1.7 \text{ MeV}$, $\tau_{A^0} \lesssim 10^{-12} \text{ s}$, and $B(A^0 \rightarrow e^+e^-) = 1$.

¹¹ EICHLER 86 looked for $\pi^+ \rightarrow e^+ \nu A^0$ followed by $A^0 \rightarrow e^+e^-$. Limits on the branching fraction depend on the mass and lifetime of A^0 . The quoted limits are valid when $\pi(A^0) \gtrsim 3 \times 10^{-10} \text{ s}$ if the decays are kinematically allowed.

¹² YAMAZAKI 84 looked for a discrete line in $K^+ \rightarrow \pi^+ X$. Sensitive to wide mass range (5-300 MeV), independent of whether X decays promptly or not.

- ¹³ASANO 82 at KEK set limits for $B(K^+ \rightarrow \pi^+ A^0)$ for $m_{A^0} < 100$ MeV as $BR < 4. \times 10^{-8}$ for $\tau(A^0 \rightarrow n\gamma's) > 1. \times 10^{-9}$ s, $BR < 1.4 \times 10^{-6}$ for $\tau < 1. \times 10^{-9}$ s.
¹⁴ASANO 81B is KEK experiment. Set $B(K^+ \rightarrow \pi^+ A^0) < 3.8 \times 10^{-8}$ at CL = 90%.
¹⁵ZHITNITSKII 79 argue that a heavy axion predicted by YANG 78 ($3 < m < 40$ MeV) contradicts experimental muon anomalous magnetic moments.

 A^0 (Axion) Searches in Quarkonium Decays

Decay or transition of quarkonium. Limits are for branching ratio.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$< 4.0 \times 10^{-5}$	90		ANTREASYAN 90C CBAL 16 ANTREASYAN 90C RVUE	$T(1S) \rightarrow A^0 \gamma$	
$< 5 \times 10^{-5}$	90		17 DRUZHININ 87 ND	$\phi \rightarrow \gamma A^0$ ($A^0 \rightarrow e^+ e^-$)	
$< 2 \times 10^{-3}$	90		18 DRUZHININ 87 ND	$\phi \rightarrow \gamma A^0$ ($A^0 \rightarrow \gamma \gamma$)	
$< 7 \times 10^{-6}$	90		19 DRUZHININ 87 ND	$\phi \rightarrow \gamma A^0$ ($A^0 \rightarrow$ missing)	
$< 3.1 \times 10^{-4}$	90	0	20 ALBRECHT 86D ARG	$T(1S) \rightarrow \gamma A^0$ ($A^0 \rightarrow e^+ e^-$)	
$< 4 \times 10^{-4}$	90	0	20 ALBRECHT 86D ARG	$T(1S) \rightarrow \gamma A^0$ ($A^0 \rightarrow \mu^+ \mu^-$, $\pi^+ \pi^-$, $K^+ K^-$)	
$< 8 \times 10^{-4}$	90	1	21 ALBRECHT 86D ARG	$T(1S) \rightarrow \gamma A^0$	
$< 1.3 \times 10^{-3}$	90	0	22 ALBRECHT 86D ARG	$T(1S) \rightarrow \gamma A^0$ ($A^0 \rightarrow e^+ e^-$, $\gamma \gamma$)	
$< 2. \times 10^{-3}$	90		23 BOWCOCK 86 CLEO	$T(2S) \rightarrow T(1S) \rightarrow A^0$	
$< 5. \times 10^{-3}$	90		24 MAGERAS 86 CUSB	$T(1S) \rightarrow A^0 \gamma$	
$< 3. \times 10^{-4}$	90		25 ALAM 83 CLEO	$T(1S) \rightarrow A^0 \gamma$	
$< 9.1 \times 10^{-4}$	90		26 NICZYPORUK 83 LENA	$T(1S) \rightarrow A^0 \gamma$	
$< 1.4 \times 10^{-5}$	90		27 EDWARDS 82 CBAL	$J/\psi \rightarrow A^0 \gamma$	
$< 3.5 \times 10^{-4}$	90		28 SIVERTZ 82 CUSB	$T(1S) \rightarrow A^0 \gamma$	
$< 1.2 \times 10^{-4}$	90		28 SIVERTZ 82 CUSB	$T(3S) \rightarrow A^0 \gamma$	

- ¹⁶The combined limit of ANTREASYAN 90C and EDWARDS 82 excludes standard axion with $m_{A^0} < 2m_e$ at 90% CL as long as $C_T C_{J/\psi} > 0.09$, where $C_V (V = T, J/\psi)$ is the reduction factor for $\Gamma(V \rightarrow A^0 \gamma)$ due to QCD and/or relativistic corrections. The same data excludes $0.02 < x < 260$ (90% CL) if $C_T = C_{J/\psi} = 0.5$, and further combining with ALBRECHT 86D result excludes $5 \times 10^{-5} < x < 260$. x is the ratio of the vacuum expectation values of the two Higgs fields. These limits use conventional assumption $\Gamma(A^0 \rightarrow ee) \propto x^{-2}$. The alternative assumption $\Gamma(A^0 \rightarrow ee) \propto x^2$ gives a somewhat different excluded region $0.00075 < x < 44$.
¹⁷The first DRUZHININ 87 limit is valid when $\tau_{A^0}/m_{A^0} < 3 \times 10^{-13}$ s/MeV and $m_{A^0} < 20$ MeV.
¹⁸The second DRUZHININ 87 limit is valid when $\tau_{A^0}/m_{A^0} < 5 \times 10^{-13}$ s/MeV and $m_{A^0} < 20$ MeV.
¹⁹The third DRUZHININ 87 limit is valid when $\tau_{A^0}/m_{A^0} > 7 \times 10^{-12}$ s/MeV and $m_{A^0} < 200$ MeV.
²⁰ $\tau_{A^0} < 1 \times 10^{-13}$ s and $m_{A^0} < 1.5$ GeV. Applies for $A^0 \rightarrow \gamma \gamma$ when $m_{A^0} < 100$ MeV.
²¹ $\tau_{A^0} > 1 \times 10^{-7}$ s.
²²Independent of τ_{A^0} .
²³BOWCOCK 86 looked for A^0 that decays into $e^+ e^-$ in the cascade decay $T(2S) \rightarrow T(1S) \pi^+ \pi^-$ followed by $T(1S) \rightarrow A^0 \gamma$. The limit for $B(T(1S) \rightarrow A^0 \gamma) B(A^0 \rightarrow e^+ e^-)$ depends on m_{A^0} and τ_{A^0} . The quoted limit for $m_{A^0} = 1.8$ MeV is at $\tau_{A^0} \sim 2. \times 10^{-12}$ s, where the limit is the worst. The same limit $2. \times 10^{-3}$ applies for all lifetimes for masses $2m_e < m_{A^0} < 2m_\mu$ when the results of this experiment are combined with the results of ALAM 83.
²⁴MAGERAS 86 looked for $T(1S) \rightarrow \gamma A^0$ ($A^0 \rightarrow e^+ e^-$). The quoted branching fraction limit is for $m_{A^0} = 1.7$ MeV, at $\tau(A^0) \sim 4. \times 10^{-13}$ s where the limit is the worst.
²⁵ALAM 83 is at CESR. This limit combined with limit for $B(J/\psi \rightarrow A^0 \gamma)$ (EDWARDS 82) excludes standard axion.
²⁶NICZYPORUK 83 is DESY-DORIS experiment. This limit together with lower limit 9.2×10^{-4} of $B(T \rightarrow A^0 \gamma)$ derived from $B(J/\psi(1S) \rightarrow A^0 \gamma)$ limit (EDWARDS 82) excludes standard axion.
²⁷EDWARDS 82 looked for $J/\psi \rightarrow \gamma A^0$ decays by looking for events with a single γ [of energy $\sim 1/2$ the $J/\psi(1S)$ mass], plus nothing else in the detector. The limit is inconsistent with the axion interpretation of the FAISSNER 81B result.
²⁸SIVERTZ 82 is CESR experiment. Looked for $T \rightarrow \gamma A^0$, A^0 undetected. Limit for $1S$ (3S) is valid for $m_{A^0} < 7$ GeV (4 GeV).

 A^0 (Axion) Searches in Positronium Decays

Decay or transition of positronium. Limits are for branching ratio.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-5}$	90	29 AKOPYAN 91 CNTR		$o\text{-Ps} \rightarrow A^0 \gamma$ ($A^0 \rightarrow \gamma \gamma$), $m_{A^0} < 30$ keV
$< 1.1 \times 10^{-6}$	90	30 ASAI 91 CNTR		$o\text{-Ps} \rightarrow A^0 \gamma$, $m_{A^0} < 800$ keV
$< 3.8 \times 10^{-4}$	90	GNINENKO 90 CNTR		$o\text{-Ps} \rightarrow A^0 \gamma$, $m_{A^0} < 30$ keV
$< (1-5) \times 10^{-4}$	95	31 TSUCHIAKI 90 CNTR		$o\text{-Ps} \rightarrow A^0 \gamma$, $m_{A^0} = 300-900$ keV
$< 6.4 \times 10^{-5}$	90	32 ORITO 89 CNTR		$o\text{-Ps} \rightarrow A^0 \gamma$, $m_{A^0} < 30$ keV
		33 AMALDI 85 CNTR		Ortho-positronium
		34 CARBONI 83 CNTR		Ortho-positronium

²⁹The AKOPYAN 91 limit applies for a short-lived A^0 with $\tau_{A^0} < 10^{-13}$ s. m_{A^0} [keV].
³⁰ASAI 91 limit translates to $g_{A^0 e e}^2 / 4\pi < 1.1 \times 10^{-11}$ (90%CL) for $m_{A^0} < 800$ keV.
³¹The TSUCHIAKI 90 limit is based on inclusive photon spectrum and is independent of A^0 decay modes.
³²ORITO 89 limit translates to $g_{A^0 e e}^2 / 4\pi < 6.2 \times 10^{-10}$. Somewhat more sensitive limits are obtained for larger m_{A^0} : $B < 7.6 \times 10^{-6}$ at 100 keV.
³³AMALDI 85 set limits $B(A^0 \gamma) / B(\gamma \gamma \gamma) < (1-5) \times 10^{-6}$ for $m_{A^0} = 900-100$ keV which are about 1/10 of the CARBONI 83 limits.
³⁴CARBONI 83 looked for orthopositronium $\rightarrow A^0 \gamma$. Set limit for A^0 electron coupling squared, $g(e e A^0)^2 / (4\pi) < 6. \times 10^{-10} \cdot \gamma. \times 10^{-9}$ for m_{A^0} from 150-900 keV (CL = 99.7%). This is about 1/10 of the bound from $g-2$ experiments.

 A^0 (Axion) Search in Photoproduction

VALUE	CL%	DOCUMENT ID	COMMENT
		35 BASSOMPIERRE... 93	$m_{A^0} = 1.8$ MeV

³⁵BASSOMPIERRE 93 looked for a peak in invariant mass of $e^+ e^-$ at 1.8 MeV in photoproduction on an aligned Ge crystal. They quote a bound on the portion of possible resonance contribution of 10^{-3} for $\tau_{X^0} < 4 \times 10^{-13}$ s.

 A^0 (Axion) Production in Hadron CollisionsLimits are for $\sigma(A^0) / \sigma(\pi^0)$.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
			36 BLUEMLEIN 92 BDMP		$A^0 N_Z \rightarrow \ell^+ \ell^- N_Z$
			37 MEIJERDREES 92 SPEC		$\pi^- p \rightarrow n A^0, A^0 \rightarrow e^+ e^-$
			38 BLUEMLEIN 91 BDMP		$A^0 \rightarrow e^+ e^-$, 2γ
			39 FAISSNER 89 OSPK		Beam dump,
			40 DEBOER 88 RVUE		$A^0 \rightarrow e^+ e^-$
			41 EL-NADI 88 EMUL		$A^0 \rightarrow e^+ e^-$
			42 FAISSNER 88 OSPK		Beam dump, $A^0 \rightarrow 2\gamma$
			43 BADIER 86 BDMP		$A^0 \rightarrow e^+ e^-$
$< 2. \times 10^{-11}$	90	0	44 BERGSMA 85 CHRM		CERN beam dump
$< 1. \times 10^{-13}$	90	0	44 BERGSMA 85 CHRM		CERN beam dump
		24	45 FAISSNER 83 OSPK		Beam dump, $A^0 \rightarrow 2\gamma$
			46 FAISSNER 83B RVUE		LAMPF beam dump
			47 FRANK 83B RVUE		LAMPF beam dump
			48 HOFFMAN 83 CNTR		$\pi p \rightarrow n A^0$ ($A^0 \rightarrow e^+ e^-$)
			49 FETSCHER 82 RVUE		See FAISSNER 81B
		12	50 FAISSNER 81 OSPK		CERN PS ν wideband
		15	51 FAISSNER 81B OSPK		Beam dump, $A^0 \rightarrow 2\gamma$
		8	52 KIM 81 OSPK		26 GeV $p N \rightarrow A^0 X$
		0	53 FAISSNER 80 OSPK		Beam dump, $A^0 \rightarrow e^+ e^-$
$< 1. \times 10^{-8}$	90		54 JACQUES 80 HLBC		28 GeV protons
$< 1. \times 10^{-14}$	90		54 JACQUES 80 HLBC		Beam dump
			55 SOUKAS 80 CALO		28 GeV p beam dump
			56 BECHIS 79 CNTR		
$< 1. \times 10^{-8}$	90		57 COTEUS 79 OSPK		Beam dump
$< 1. \times 10^{-3}$	95		58 DISHAW 79 CALO		400 GeV pp
$< 1. \times 10^{-8}$	90		ALIBRAN 78 HYBR		Beam dump
$< 6. \times 10^{-9}$	95		ASRATYAN 78B CALO		Beam dump
$< 1.5 \times 10^{-8}$	90		59 BELLOTTI 78 HLBC		Beam dump
$< 5.4 \times 10^{-14}$	90		59 BELLOTTI 78 HLBC		$m_{A^0} = 1.5$ MeV
$< 4.1 \times 10^{-9}$	90		59 BELLOTTI 78 HLBC		$m_{A^0} = 1$ MeV
$< 1. \times 10^{-8}$	90		60 BOSETTI 78B HYBR		Beam dump
			61 DONNELLY 78		
$< 0.5 \times 10^{-8}$	90		HANSL 78D WIRE		Beam dump
			62 MICELMAC... 78		
			63 VYSOTSKII 78		

Gauge & Higgs Boson Full Listings

Axions (A^0) and Other Very Light Bosons

- 36 BLUEMLEIN 92 is a proton beam dump experiment at Serpukhov with a secondary target to induce Bethe-Heitler production of e^+e^- or $\mu^+\mu^-$ from the produce A^0 . See Fig. 5 for the excluded region in m_{A^0} - x plane. For the standard axion, $0.3 < x < 25$ is excluded at 95% CL. If combined with BLUEMLEIN 91, $0.008 < x < 32$ is excluded.
- 37 MEIJERDREES 92 give $\Gamma(\pi^- \rightarrow n A^0) B(A^0 \rightarrow e^+e^-) / \Gamma(\pi^- \rightarrow \text{all}) < 10^{-5}$ (90% CL) for $m_{A^0} = 100$ MeV, $\tau_{A^0} = 10^{-11}$ – 10^{-23} sec. Limits ranging from 2.5×10^{-3} to 10^{-7} are given for $m_{A^0} = 25$ –136 MeV.
- 38 BLUEMLEIN 91 is a proton beam dump experiment at Serpukhov. No candidate event for $A^0 \rightarrow e^+e^-$, 2γ are found. Fig. 6 gives the excluded region in m_{A^0} - x plane ($x = \tan\theta = v_2/v_1$). Standard axion is excluded for $0.2 < m_{A^0} < 3.2$ MeV for most $x > 1$, 0.2 – 1.1 MeV for most $x < 1$.
- 39 FAISSNER 89 searched for $A^0 \rightarrow e^+e^-$ in a proton beam dump experiment at SIN. No excess of events was observed over the background. A standard axion with mass $2m_e$ – 20 MeV is excluded. Lower limit on f_{A^0} of $\approx 10^4$ GeV is given for $m_{A^0} = 2m_e$ – 20 MeV.
- 40 DEBOER 88 reanalyze EL-NADI 88 data and claim evidence for three distinct states with mass ~ 1.1 , ~ 2.1 , and ~ 9 MeV, lifetimes 10^{-16} – 10^{-15} s decaying to e^+e^- and note the similarity of the data with those of a cosmic-ray experiment by Bristol group (B.M. Anand, Proc. of the Royal Society of London, Section A **A22** 183 (1953)). For a criticism see PERKINS 89, who suggests that the events are compatible with π^0 Dalitz decay. DEBOER 89b is a reply which contests the criticism.
- 41 EL-NADI 88 claim the existence of a neutral particle decaying into e^+e^- with mass 1.60 ± 0.59 MeV, lifetime $(0.15 \pm 0.01) \times 10^{-14}$ s, which is produced in heavy ion interactions with emulsion nuclei at ~ 4 GeV/c/nucleon.
- 42 FAISSNER 88 is a proton beam dump experiment at SIN. They found no candidate event for $A^0 \rightarrow \gamma\gamma$. A standard axion decaying to 2γ is excluded except for a region $x \approx 1$. Lower limit on f_{A^0} of 10^2 – 10^3 GeV is given for $m_{A^0} = 0.1$ – 1 MeV.
- 43 BADIÉ 86 did not find long-lived A^0 in 300 GeV π^- Beam Dump Experiment that decays into e^+e^- in the mass range $m_{A^0} = (20$ – $200)$ MeV, which excludes the A^0 decay constant $f(A^0)$ in the interval (60–600) GeV. See their figure 6 for excluded region on $f(A^0)$ - m_{A^0} plane.
- 44 BERGSMA 85 look for $A^0 \rightarrow 2\gamma$, e^+e^- , $\mu^+\mu^-$. First limit above is for $m_{A^0} = 1$ MeV; second is for 200 MeV. See their figure 4 for excluded region on f_{A^0} - m_{A^0} plane, where f_{A^0} is A^0 decay constant. For Peccei-Quinn PECCEI 77 A^0 , $m_{A^0} < 180$ keV and $\tau > 0.037$ s. (CL = 90%). For the axion of FAISSNER 81B at 250 keV, BERGSMA 85 expect 15 events but observe zero.
- 45 FAISSNER 83 observed 19 1 - γ and 12 2 - γ events where a background of 4.8 and 2.3 respectively is expected. A small-angle peak is observed even if iron wall is set in front of the decay region.
- 46 FAISSNER 83B extrapolate SIN γ signal to LAMPF ν experimental condition. Resulting 370 γ 's are not at variance with LAMPF upper limit of 450 γ 's. Derived from LAMPF limit that $[d\sigma(A^0)/d\omega \text{ at } 90^\circ] m_{A^0}/\tau_{A^0} < 14 \times 10^{-35}$ cm² sr⁻¹ MeV ms⁻¹. See comment on FRANK 83B.
- 47 FRANK 83B stress the importance of LAMPF data bins with negative net signal. By statistical analysis say that LAMPF and SIN-A0 are at variance when extrapolation by phase-space model is done. They find LAMPF upper limit is 248 not 450 γ 's. See comment on FAISSNER 83B.
- 48 HOFFMAN 83 set CL = 90% limit $d\sigma/dt B(e^+e^-) < 3.5 \times 10^{-32}$ cm²/GeV² for $140 < m_{A^0} < 160$ MeV. Limit assumes $\tau(A^0) < 10^{-9}$ s.
- 49 FETSCHER 82 reanalyzes SIN beam-dump data of FAISSNER 81. Claims no evidence for axion since 2 - γ peak rate remarkably decreases if iron wall is set in front of the decay region.
- 50 FAISSNER 81 see excess μe events. Suggest axion interactions.
- 51 FAISSNER 81B is SIN 590 MeV proton beam dump. Observed 14.5 ± 5.0 events of 2γ decay of long-lived neutral penetrating particle with $m_{2\gamma} \lesssim 1$ MeV. Axion interpretation with η - A^0 mixing gives $m_{A^0} = 250 \pm 25$ keV, $\tau_{(2\gamma)} = (7.3 \pm 3.7) \times 10^{-3}$ s from above rate. See critical remarks below in comments of FETSCHER 82, FAISSNER 83, FAISSNER 83B, FRANK 83B, and BERGSMA 85. Also see in the next subsection ALEK-SEEV 82, CAVAGNAC 83, and ANANEV 85.
- 52 KIM 81 analyzed 8 candidates for $A^0 \rightarrow 2\gamma$ obtained by Aachen-Padova experiment at CERN with 26 GeV protons on Be. Estimated axion mass is about 300 keV and lifetime is $(0.86 \sim 5.6) \times 10^{-3}$ s depending on models. Faissner (private communication), says axion production underestimated and mass overestimated. Correct value around 200 keV.
- 53 FAISSNER 80 is SIN beam dump experiment with 590 MeV protons looking for $A^0 \rightarrow e^+e^-$ decay. Assuming $A^0/\pi^0 = 5.5 \times 10^{-7}$, obtained decay rate limit $20/(A^0 \text{ mass})$ MeV/s (CL = 90%), which is about 10^{-7} below theory and interpreted as upper limit to $m_{A^0} < 2m_e$.
- 54 JACQUES 80 is a BNL beam dump experiment. First limit above comes from nonobservation of excess neutral-current-type events [$\sigma(\text{production})\sigma(\text{interaction}) < 7 \times 10^{-68}$ cm⁴, CL = 90%]. Second limit is from nonobservation of axion decays into 2γ 's or e^+e^- , and for axion mass a few MeV.
- 55 SOUKAS 80 at BNL observed no excess of neutral-current-type events in beam dump.
- 56 BECHIS 79 looked for the axion production in low energy electron Bremsstrahlung and the subsequent decay into either 2γ or e^+e^- . No signal found. CL = 90% limits for model parameter(s) are given.
- 57 COTEUS 79 is a beam dump experiment at BNL.
- 58 DISHAW 79 is a calorimetric experiment and looks for low energy tail of energy distributions due to energy lost to weakly interacting particles.
- 59 BELLOTTI 78 first value comes from search for $A^0 \rightarrow e^+e^-$. Second value comes from search for $A^0 \rightarrow 2\gamma$, assuming mass $< 2m_e$. For any mass satisfying this, limit is above value $\times (\text{mass}^{-4})$. Third value uses data of PL 60B 401 and quotes $\sigma(\text{production})\sigma(\text{interaction}) < 10^{-67}$ cm⁴.
- 60 BOSETTI 78B quotes $\sigma(\text{production})\sigma(\text{interaction}) < 2 \times 10^{-67}$ cm⁴.
- 61 DONNELLY 78 examines data from reactor neutrino experiments of REINES 76 and GURR 74 as well as SLAC beam dump experiment. Evidence is negative.
- 62 MICELMACHER 78 finds no evidence of axion existence in reactor experiments of REINES 76 and GURR 74. (See reference under DONNELLY 78 below).

63 VYSOTSKII 78 derived lower limit for the axion mass 25 keV from luminosity of the sun and 200 keV from red supergiants.

A^0 (Axion) Searches in Reactor Experiments

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	64 KETOV	86 SPEC	Reactor, $A^0 \rightarrow \gamma\gamma$
	65 KOCH	86 SPEC	Reactor; $A^0 \rightarrow \gamma\gamma$
	66 DATAR	82 CNTR	Light water reactor
	67 VUILLEUMIER	81 CNTR	Reactor, $A^0 \rightarrow 2\gamma$
64 KETOV 86 searched for A^0 at the Rovno nuclear power plant. They found an upper limit on the A^0 production probability of $0.8 [100 \text{ keV}/m_{A^0}]^6 \times 10^{-6}$ per fission. In the standard axion model, this corresponds to $m_{A^0} > 150$ keV. Not valid for $m_{A^0} \gtrsim 1$ MeV.			
65 KOCH 86 searched for $A^0 \rightarrow \gamma\gamma$ at nuclear power reactor Biblis A. They found an upper limit on the A^0 production rate of $\omega(A^0)/\omega(\gamma(M1)) < 1.5 \times 10^{-10}$ (CL=95%). Standard axion with $m_{A^0} = 250$ keV gives 10^{-5} for the ratio. Not valid for $m_{A^0} > 1022$ keV.			
66 DATAR 82 looked for $A^0 \rightarrow 2\gamma$ in neutron capture ($n p \rightarrow d A^0$) at Tarapur 500 MW reactor. Sensitive to sum of $l = 0$ and $l = 1$ amplitudes. With ZEHNDER 81 [($l = 0$) - ($l = 1$)] result, assert nonexistence of standard A^0 .			
67 VUILLEUMIER 81 is at Grenoble reactor. Set limit $m_{A^0} < 280$ keV.			

A^0 (Axion) and Other Light Boson (X^0) Searches in Nuclear Transitions

Limits are for branching ratio.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$< 1.2 \times 10^{-6}$	95		68 MINOWA	93 CNTR	$^{139}\text{La}^* \rightarrow ^{139}\text{La} A^0$
$< 2 \times 10^{-4}$	90		69 HICKS	92 CNTR	^{35}S decay, $A^0 \rightarrow \gamma\gamma$
$< 1.5 \times 10^{-9}$	95		70 ASANUMA	90 CNTR	^{241}Am decay
$< (0.4$ – $10) \times 10^{-3}$	95		71 DEBOER	90 CNTR	$^8\text{Be}^* \rightarrow ^8\text{Be} A^0$, $X^0 \rightarrow e^+e^-$
$< (0.2$ – $1) \times 10^{-3}$	90		72 BINI	89 CNTR	$^{16}\text{O}^* \rightarrow ^{16}\text{O} X^0$, $X^0 \rightarrow e^+e^-$
			73 AVIGNONE	88 CNTR	$\text{Cu}^* \rightarrow \text{Cu} A^0$ ($A^0 \rightarrow 2\gamma$, $A^0 e^- \rightarrow \gamma e^-$)
$< 1.5 \times 10^{-4}$	90		74 DATAR	88 CNTR	$^{12}\text{C}^* \rightarrow ^{12}\text{C} A^0$, $A^0 Z \rightarrow \gamma Z$
$< 5 \times 10^{-3}$	90		75 DEBOER	88c CNTR	$^{16}\text{O}^* \rightarrow ^{16}\text{O} X^0$, $X^0 \rightarrow e^+e^-$
$< 3.4 \times 10^{-5}$	95		76 DOEHNER	88 SPEC	$^2\text{H}^*, A^0 \rightarrow e^+e^-$
$< 4 \times 10^{-4}$	95		77 SAVAGE	88 CNTR	Nuclear decay (isovector)
$< 3 \times 10^{-3}$	95		77 SAVAGE	88 CNTR	Nuclear decay (isoscalar)
< 0.106	90		78 HALLIN	86 SPEC	^6Li isovector decay
< 10.8	90		78 HALLIN	86 SPEC	^{10}B isoscalar decays
< 2.2	90		78 HALLIN	86 SPEC	^{14}N isoscalar decays
$< 4 \times 10^{-4}$	90	0	79 SAVAGE	86B CNTR	$^{14}\text{N}^*$
			80 ANANEV	85 CNTR	$\text{Li}^*, \text{deut}^* A^0 \rightarrow 2\gamma$
			81 CAVAGNAC	83 CNTR	$^{97}\text{Nb}^*, \text{deut}^*$ transition $A^0 \rightarrow 2\gamma$
			82 ALEKSEEV	82B CNTR	$\text{Li}^*, \text{deut}^*$ transition $A^0 \rightarrow 2\gamma$
			83 LEHMANN	82 CNTR	$\text{Cu}^* \rightarrow \text{Cu} A^0$ ($A^0 \rightarrow 2\gamma$)
		0	84 ZEHNDER	82 CNTR	Li^*, Nb^* decay, n -capt.
		0	85 ZEHNDER	81 CNTR	$\text{Ba}^* \rightarrow \text{Ba} A^0$ ($A^0 \rightarrow 2\gamma$)
			86 CALAPRICE	79	Carbon
68 MINOWA 93 studied chain process, $^{139}\text{Ce} \rightarrow ^{139}\text{La}^*$ by electron capture and M1 transition of $^{139}\text{La}^*$ to the ground state. It does not assume decay modes of A^0 . The bound applies for $m_{A^0} < 166$ keV.					
69 HICKS 92 bound is applicable for $\tau_{X^0} < 4 \times 10^{-11}$ sec.					
70 The ASANUMA 90 limit is for the branching fraction of X^0 emission per ^{241}Am α decay and valid for $\tau_{X^0} < 3 \times 10^{-11}$ s.					
71 The DEBOER 90 limit is for the branching ratio $^8\text{Be}^* (18.15 \text{ MeV}, 1^+) \rightarrow ^8\text{Be} A^0$, $A^0 \rightarrow e^+e^-$ for the mass range $m_{A^0} = 4$ – 15 MeV.					
72 The BINI 89 limit is for the branching fraction of $^{16}\text{O}^* (6.05 \text{ MeV}, 0^+) \rightarrow ^{16}\text{O} X^0$, $X^0 \rightarrow e^+e^-$ for $m_X = 1.5$ – 3.1 MeV. $\tau_{X^0} \lesssim 10^{-11}$ s is assumed. The spin-parity of X is restricted to 0^+ or 1^- .					
73 AVIGNONE 88 looked for the 1115 keV transition $\text{C}^* \rightarrow \text{Cu} A^0$, either from $A^0 \rightarrow 2\gamma$ in-flight decay or from the secondary A^0 interactions by Compton and by Primakoff processes. Limits for axion parameters are obtained for $m_{A^0} < 1.1$ MeV.					
74 DATAR 88 rule out light pseudoscalar particle emission through its decay $A^0 \rightarrow e^+e^-$ in the mass range 1.02 – 2.5 MeV and lifetime range 10^{-13} – 10^{-8} s. The above limit is for $\tau = 5 \times 10^{-13}$ s and $m = 1.7$ MeV; see the paper for the τ - m dependence of the limit.					

Gauge & Higgs Boson Full Listings

Axions (A^0) and Other Very Light Bosons

- 75 The limit is for the branching fraction of $^{16}\text{O}^*(6.05 \text{ MeV}, 0^+) \rightarrow ^{16}\text{O}X^0, X^0 \rightarrow e^+e^-$ against internal pair conversion for $m_{X^0} = 1.7 \text{ MeV}$ and $\tau_{X^0} < 10^{-11} \text{ s}$. Similar limits are obtained for $m_{X^0} = 1.3\text{--}3.2 \text{ MeV}$. The spin parity of X^0 must be either 0^+ or 1^- . The limit at 1.7 MeV is translated into a limit for the X^0 -nucleon coupling constant: $g_{X^0 NN}^2/4\pi < 2.3 \times 10^{-9}$.
- 76 The DOEHNER 88 limit is for $m_{A^0} = 1.7 \text{ MeV}$, $\tau(A^0) < 10^{-10} \text{ s}$. Limits less than 10^{-4} are obtained for $m_{A^0} = 1.2\text{--}2.2 \text{ MeV}$.
- 77 SAVAGE 88 looked for A^0 that decays into e^+e^- in the decay of the 9.17 MeV $J^P = 2^+$ state in ^{14}N , 17.64 MeV state $J^P = 1^+$ in ^8Be , and the 18.15 MeV state $J^P = 1^+$ in ^8Be . This experiment constrains the isovector coupling of A^0 to hadrons, if $m_{A^0} = (1.1 \rightarrow 2.2) \text{ MeV}$ and the isoscalar coupling of A^0 to hadrons, if $m_{A^0} = (1.1 \rightarrow 2.6) \text{ MeV}$. Both limits are valid only if $\tau(A^0) \lesssim 1 \times 10^{-11} \text{ s}$.
- 78 Limits are for $\Gamma(A^0(1.8 \text{ MeV}))/\Gamma(\pi M1)$; i.e., for 1.8 MeV axion emission normalized to the rate for internal emission of e^+e^- pairs. Valid for $\tau_{A^0} < 2 \times 10^{-11} \text{ s}$. ^6Li isovector decay data strongly disfavor PECCEI 86 model I, whereas the ^{10}B and ^{14}N isoscalar decay data strongly reject PECCEI 86 model II and III.
- 79 SAVAGE 86B looked for A^0 that decays into e^+e^- in the decay of the 9.17 MeV $J^P = 2^+$ state in ^{14}N . Limit on the branching fraction is valid if $\tau_{A^0} \lesssim 1 \times 10^{-11} \text{ s}$ for $m_{A^0} = (1.1\text{--}1.7) \text{ MeV}$. This experiment constrains the iso-vector coupling of A^0 to hadrons.
- 80 ANANEV 85 with IBR-2 pulsed reactor exclude standard A^0 at CL = 95% masses below 470 keV (Li^* decay) and below $2m_e$ for deuteron* decay.
- 81 CAVIGNAC 83 at Bugey reactor exclude axion at any $m_{97}\text{Nb}^*$ decay and axion with m_{A^0} between 275 and 288 keV (deuteron* decay).
- 82 ALEKSEEV 82 with IBR-2 pulsed reactor exclude standard A^0 at CL = 95% mass-ranges $m_{A^0} < 400 \text{ keV}$ (Li^* decay) and $330 \text{ keV} < m_{A^0} < 2.2 \text{ MeV}$. (deuteron* decay).
- 83 LEHMANN 82 obtained $A^0 \rightarrow 2\gamma$ rate $< 6.2 \times 10^{-5}/\text{s}$ (CL = 95%) excluding m_{A^0} between 100 and 1000 keV.
- 84 ZEHNDER 82 used Goesgen 2.8GW light-water reactor to check A^0 production. No 2γ peak in Li^* , Nb^* decay (both single p transition) nor in n capture (combined with previous Ba^* negative result) rules out standard A^0 . Set limit $m_{A^0} < 60 \text{ keV}$ for any A^0 .
- 85 ZEHNDER 81 looked for $\text{Ba}^* \rightarrow A^0\text{Ba}$ transition with $A^0 \rightarrow 2\gamma$. Obtained 2γ coincidence rate $< 2.2 \times 10^{-5}/\text{s}$ (CL = 95%) excluding $m_{A^0} > 160 \text{ keV}$ (or 200 keV depending on Higgs mixing). However, see BARROSO 81.
- 86 CALAPRICE 79 saw no axion emission from excited states of carbon. Sensitive to axion mass between 1 and 15 MeV.

A^0 (Axion) Limits from Its Electron Coupling

Limits are for $\tau(A^0 \rightarrow e^+e^-)$.

VALUE (s)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$none\ 4 \times 10^{-16}\text{--}4.5 \times 10^{-12}$	90	87 BROSS	91 BDMP	$eN \rightarrow eA^0N$ ($A^0 \rightarrow ee$)
		88 GUO	90 BDMP	$eN \rightarrow eA^0N$ ($A^0 \rightarrow ee$)
		89 BJORKEN	88 CALO	$A \rightarrow e^+e^-$ or 2γ
		90 BLINOV	88 MD1	$ee \rightarrow eeA^0$ ($A^0 \rightarrow ee$)
$none\ 1 \times 10^{-14}\text{--}1 \times 10^{-10}$	90	91 RIORDAN	87 BDMP	$eN \rightarrow eA^0N$ ($A^0 \rightarrow ee$)
$none\ 1 \times 10^{-14}\text{--}1 \times 10^{-11}$	90	92 BROWN	86 BDMP	$eN \rightarrow eA^0N$ ($A^0 \rightarrow ee$)
$none\ 6 \times 10^{-14}\text{--}9 \times 10^{-11}$	95	93 DAVIER	86 BDMP	$eN \rightarrow eA^0N$ ($A^0 \rightarrow ee$)
$none\ 3 \times 10^{-13}\text{--}1 \times 10^{-7}$	90	94 KONAKA	86 BDMP	$eN \rightarrow eA^0N$ ($A^0 \rightarrow ee$)
87 The listed BROSS 91 limit is for $m_{A^0} = 1.14 \text{ MeV}$. $B(A^0 \rightarrow e^+e^-) = 1$ assumed. Excluded domain in the $\tau_{A^0}\text{--}m_{A^0}$ plane extends up to $m_{A^0} \approx 7 \text{ MeV}$ (see Fig. 5). Combining with electron $g\text{--}2$ constraint, axions coupling only to e^+e^- ruled out for $m_{A^0} < 4.8 \text{ MeV}$ (90%CL).				
88 GUO 90 use the same apparatus as BROWN 86 and improve the previous limit in the shorter lifetime region. Combined with $g\text{--}2$ constraint, axions coupling only to e^+e^- are ruled out for $m_{A^0} < 2.7 \text{ MeV}$ (90%CL).				
89 BJORKEN 88 reports limits on axion parameters (f_A, m_A, τ_A) for $m_{A^0} < 200 \text{ MeV}$ from electron beam-dump experiment with production via Primakoff photoproduction, bremsstrahlung from electrons, and resonant annihilation of positrons on atomic electrons.				
90 BLINOV 88 assume zero spin, $m = 1.8 \text{ MeV}$ and lifetime $< 5 \times 10^{-12} \text{ s}$ and find $\Gamma(A^0 \rightarrow \gamma\gamma)B(A^0 \rightarrow e^+e^-) < 2 \text{ eV}$ (CL=90%).				
91 Assumes $A^0\gamma\gamma$ coupling is small and hence Primakoff production is small. Their figure 2 shows limits on axions for $m_{A^0} < 15 \text{ MeV}$.				
92 Uses electrons in hadronic showers from an incident 800 GeV proton beam. Limits for $m_{A^0} < 15 \text{ MeV}$ are shown in their figure 3.				
93 $m_{A^0} = 1.8 \text{ MeV}$ assumed. The excluded domain in the $\tau_{A^0}\text{--}m_{A^0}$ plane extends up to $m_{A^0} \approx 14 \text{ MeV}$, see their figure 4.				
94 The limits are obtained from their figure 3. Also given is the limit on the $A^0\gamma\gamma\text{--}A^0e^+e^-$ coupling plane by assuming Primakoff production.				

Search for A^0 (Axion) Resonance in Bhabha Scattering

The limit is for $\Gamma(A^0)B(A^0 \rightarrow e^+e^-)^2$.

VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.3	97	95 HALLIN	92 CNTR	$m_{A^0} = 1.75\text{--}1.88 \text{ MeV}$
$none\ 0.0016\text{--}0.47$	90	96 HENDERSON	92C CNTR	$m_{A^0} = 1.5\text{--}1.86 \text{ MeV}$
< 2.0	90	97 WU	92 CNTR	$m_{A^0} = 1.56\text{--}1.86 \text{ MeV}$
< 0.013	95	TSERTOS	91 CNTR	$m_{A^0} = 1.832 \text{ MeV}$
$none\ 0.19\text{--}3.3$	95	98 WIDMANN	91 CNTR	$m_{A^0} = 1.78\text{--}1.92 \text{ MeV}$
< 5	97	BAUER	90 CNTR	$m_{A^0} = 1.832 \text{ MeV}$
$none\ 0.09\text{--}1.5$	95	99 JUDGE	90 CNTR	$m_{A^0} = 1.832 \text{ MeV}$, elastic
< 1.9	97	100 TSERTOS	89 CNTR	$m_{A^0} = 1.82 \text{ MeV}$
$<(10\text{--}40)$	97	100 TSERTOS	89 CNTR	$m_{A^0} = 1.51\text{--}1.65 \text{ MeV}$
$<(1\text{--}2.5)$	97	100 TSERTOS	89 CNTR	$m_{A^0} = 1.80\text{--}1.86 \text{ MeV}$
< 31	95	LORENZ	88 CNTR	$m_{A^0} = 1.646 \text{ MeV}$
< 94	95	LORENZ	88 CNTR	$m_{A^0} = 1.726 \text{ MeV}$
< 23	95	LORENZ	88 CNTR	$m_{A^0} = 1.782 \text{ MeV}$
< 19	95	LORENZ	88 CNTR	$m_{A^0} = 1.837 \text{ MeV}$
< 3.8	97	101 TSERTOS	88 CNTR	$m_{A^0} = 1.832 \text{ MeV}$
		102 VANKLINKEN	88 CNTR	
		103 MAIER	87 CNTR	
< 2500	90	MILLS	87 CNTR	$m_{A^0} = 1.8 \text{ MeV}$
		104 VONWIMMER.87	CNTR	
95 HALLIN 92 quote limits on lifetime, $8 \times 10^{-14}\text{--}5 \times 10^{-13} \text{ sec}$ depending on mass, assuming $B(A^0 \rightarrow e^+e^-) = 100\%$. They say that TSERTOS 91 overstated their sensitivity by a factor of 3.				
96 HENDERSON 92C exclude axion with lifetime $\tau_{A^0} = 1.4 \times 10^{-12}\text{--}4.0 \times 10^{-10} \text{ s}$, assuming $B(A^0 \rightarrow e^+e^-) = 100\%$. HENDERSON 92C also exclude a vector boson with $\tau = 1.4 \times 10^{-12}\text{--}6.0 \times 10^{-10} \text{ s}$.				
97 WU 92 quote limits on lifetime $> 3.3 \times 10^{-13} \text{ s}$ assuming $B(A^0 \rightarrow e^+e^-) = 100\%$. They say that TSERTOS 89 overestimate the limit by a factor of $\pi/2$. WU 92 also quote a bound for vector boson, $\tau > 8.2 \times 10^{-13} \text{ s}$.				
98 WIDMANN 91 bound applies exclusively to the case $B(A^0 \rightarrow e^+e^-) = 1$, since the detection efficiency varies substantially as $\Gamma(A^0)_{\text{total}}$ changes. See their Fig. 6.				
99 JUDGE 90 excludes an elastic pseudoscalar e^+e^- resonance for $4.5 \times 10^{-13} \text{ s} < \tau(A^0) < 7.5 \times 10^{-12} \text{ s}$ (95% CL) at $m_{A^0} = 1.832 \text{ MeV}$. Comparable limits can be set for $m_{A^0} = 1.776\text{--}1.856 \text{ MeV}$.				
100 See also TSERTOS 88B in references.				
101 The upper limit listed in TSERTOS 88 is too large by a factor of 4. See TSERTOS 88B, footnote 3.				
102 VANKLINKEN 88 looked for relatively long-lived resonance ($\tau = 10^{-10}\text{--}10^{-12} \text{ s}$). The sensitivity is not sufficient to exclude such a narrow resonance.				
103 MAIER 87 obtained limits $R\Gamma \lesssim 60 \text{ eV}$ (100 eV) at $m_{A^0} \approx 1.64 \text{ MeV}$ (1.83 MeV) for energy resolution $\Delta E_{\text{cm}} \approx 3 \text{ keV}$, where R is the resonance cross section normalized to that of Bhabha scattering, and $\Gamma = \Gamma_{ee}^2/\Gamma_{\text{total}}$. For a discussion implying that $\Delta E_{\text{cm}} \approx 10 \text{ keV}$, see TSERTOS 89.				
104 VONWIMMERSPERG 87 measured Bhabha scattering for $E_{\text{cm}} = 1.37\text{--}1.86 \text{ MeV}$ and found a possible peak at 1.73 with $f\sigma dE_{\text{cm}} = 14.5 \pm 6.8 \text{ keV}\cdot\text{b}$. For a comment and a reply, see VANKLINKEN 88B and VONWIMMERSPERG 88. Also see CONNELL 88.				

Search for A^0 (Axion) Resonance in $e^+e^- \rightarrow \gamma\gamma$

The limit is for $\Gamma(A^0 \rightarrow e^+e^-)\Gamma(A^0 \rightarrow \gamma\gamma)/\Gamma_{\text{total}}$

VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.6	95	105 TRZASKA	91 CNTR	$m_{A^0} = 1.8 \text{ MeV}$
< 4.4	95	WIDMANN	91 CNTR	$m_{A^0} = 1.78\text{--}1.92 \text{ MeV}$
		106 FOX	89 CNTR	
< 0.11	95	107 MINOWA	89 CNTR	$m_{A^0} = 1.062 \text{ MeV}$
< 33	97	CONNELL	88 CNTR	$m_{A^0} = 1.580 \text{ MeV}$
< 42	97	CONNELL	88 CNTR	$m_{A^0} = 1.642 \text{ MeV}$
< 73	97	CONNELL	88 CNTR	$m_{A^0} = 1.782 \text{ MeV}$
< 79	97	CONNELL	88 CNTR	$m_{A^0} = 1.832 \text{ MeV}$
105 TRZASKA 91 also give limits in the range $(6.6\text{--}30) \times 10^{-3} \text{ eV}$ (95%CL) for $m_{A^0} = 1.6\text{--}2.0 \text{ MeV}$.				
106 FOX 89 measured positron annihilation with an electron in the source material into two photons and found no signal at 1.062 MeV ($< 9 \times 10^{-5}$ of two-photon annihilation at rest).				
107 Similar limits are obtained for $m_{A^0} = 1.045\text{--}1.085 \text{ MeV}$.				

Search for X^0 (Light Boson) Resonance in $e^+e^- \rightarrow \gamma\gamma$

The limit is for $\Gamma(X^0 \rightarrow e^+e^-)\Gamma(X^0 \rightarrow \gamma\gamma\gamma)/\Gamma_{\text{total}}$. C Invariance forbids spin-0 X^0 coupling to both e^+e^- and $\gamma\gamma\gamma$.

VALUE (10^{-3} eV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 3.8	95	108 SKALSEY	92 CNTR	$m_{X^0} = 1.5 \text{ MeV}$
108 SKALSEY 92 also give limits 4.3 for $m_{X^0} = 1.54$ and 7.5 for 1.64 MeV. The spin of X^0 is assumed to be one.				

Gauge & Higgs Boson Full Listings

Axions (A^0) and Other Very Light Bosons

Searches for Goldstone Bosons (X^0)

(Including Horizontal Bosons and Majorons.) Limits are for branching ratios.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
			109 BOBRAKOV	91	Electron quasi-magnetic interaction
$< 3.3 \times 10^{-2}$	95		110 ALBRECHT	90E ARG	$\tau \rightarrow \mu X^0$. Familon
$< 1.8 \times 10^{-2}$	95		110 ALBRECHT	90E ARG	$\tau \rightarrow e X^0$. Familon
$< 6.4 \times 10^{-9}$	90		111 ATIYA	90 CNTR	$K^+ \rightarrow \pi^+ X^0$. Familon
$< 1.1 \times 10^{-9}$	90		112 BOLTON	88 CBOX	$\mu^+ \rightarrow e^+ X^0$. Familon
			113 CHANDA	88 ASTR	Sun, Majoron
			114 CHOI	88 ASTR	Majoron, SN 1987A
$< 5 \times 10^{-6}$	90		115 PICCIOTTO	88 CNTR	$\pi \rightarrow e \nu X^0$, Majoron
$< 1.3 \times 10^{-9}$	90		116 GOLDMAN	87 CNTR	$\mu \rightarrow e \gamma X^0$. Familon
$< 3 \times 10^{-4}$	90		117 BRYMAN	86B RVUE	$\mu \rightarrow e X^0$. Familon
$< 1. \times 10^{-10}$	90	0	118 EICHLER	86 SPEC	$\mu^+ \rightarrow e^+ X^0$. Familon
$< 2.6 \times 10^{-6}$	90		119 JODIDIO	86 SPEC	$\mu^+ \rightarrow e^+ X^0$. Familon
			120 BALTRUSAIT.	85 MRK3	$\tau \rightarrow \ell X^0$. Familon
			121 DICUS	83 COSM	$\nu(\text{hvy}) \rightarrow \nu(\text{light}) X^0$

109 BOBRAKOV 91 searched for anomalous magnetic interactions between polarized electrons expected from the exchange of a massless pseudoscalar boson (arion). A limit $\chi_e^2 < 2 \times 10^{-4}$ (95%CL) is found for the effective anomalous magneton parametrized as $\chi_e(G_F/8\pi\sqrt{2})^{1/2}$.

110 ALBRECHT 90E limits are for $B(\tau \rightarrow \ell X^0)/B(\tau \rightarrow \ell \nu \bar{\nu})$. Valid for $m_{X^0} < 100$ MeV. The limits rise to 7.1% (for μ), 5.0% (for e) for $m_{X^0} = 500$ MeV.

111 ATIYA 90 limit is for $m_{X^0} = 0$. The limit $B < 1 \times 10^{-8}$ holds for $m_{X^0} < 95$ MeV. For the reduction of the limit due to finite lifetime of X^0 , see their Fig. 3.

112 BOLTON 88 limit corresponds to $F > 3.1 \times 10^9$ GeV, which does not depend on the chirality property of the coupling.

113 CHANDA 88 find $v_T < 10$ MeV for the weak-triplet Higgs vev. in Gelmini-Roncadelli model, and $v_S > 5.8 \times 10^6$ GeV in the singlet Majoron model.

114 CHOI 88 used the observed neutrino flux from the supernova SN 1987A to exclude the neutrino Majoron Yukawa coupling h in the range $2 \times 10^{-5} < h < 3 \times 10^{-4}$ for the interaction $L_{\text{int}} = \frac{1}{2} h \bar{\psi}_\nu \gamma_5 \psi_\nu \phi_X$. For several families of neutrinos, the limit applies for $(\Sigma h_i^2)^{1/4}$.

115 PICCIOTTO 88 limit applies when $m_{X^0} < 55$ MeV and $\tau_{X^0} > 2$ ns, and it decreases to 4×10^{-7} at $m_{X^0} = 125$ MeV, beyond which no limit is obtained.

116 GOLDMAN 87 limit corresponds to $F > 2.9 \times 10^9$ GeV for the family symmetry breaking scale from the Lagrangian $L_{\text{int}} = (1/F) \bar{\psi}_\mu \gamma^\mu (a + b \gamma_5) \psi_e \partial_\mu \phi_{X^0}$ with $a^2 + b^2 = 1$.

This is not as sensitive as the limit $F > 9.9 \times 10^9$ GeV derived from the search for $\mu^+ \rightarrow e^+ X^0$ by JODIDIO 86, but does not depend on the chirality property of the coupling.

117 Limits are for $\Gamma(\mu \rightarrow e X^0)/\Gamma(\mu \rightarrow e \nu \bar{\nu})$. Valid when $m_{X^0} = 0-93.4, 98.1-103.5$ MeV.

118 EICHLER 86 looked for $\mu^+ \rightarrow e^+ X^0$ followed by $X^0 \rightarrow e^+ e^-$. Limits on the branching fraction depend on the mass and lifetime of X^0 . The quoted limits are valid when $\tau_{X^0} \lesssim 3. \times 10^{-10}$ s if the decays are kinematically allowed.

119 JODIDIO 86 corresponds to $F > 9.9 \times 10^9$ GeV for the family symmetry breaking scale with the parity-conserving effective Lagrangian $L_{\text{int}} = (1/F) \bar{\psi}_\mu \gamma^\mu \psi_e \partial_\mu \phi_{X^0}$.

120 BALTRUSAITIS 85 search for light Goldstone boson (X^0) of broken U(1). CL = 95% limits are $B(\tau \rightarrow \mu^+ X^0)/B(\tau \rightarrow \mu^+ \nu \bar{\nu}) < 0.125$ and $B(\tau \rightarrow e^+ X^0)/B(\tau \rightarrow e^+ \nu \bar{\nu}) < 0.04$. Inferred limit for the symmetry breaking scale is $m > 3000$ TeV.

121 The primordial heavy neutrino must decay into ν and familon, f_A , early so that the red-shifted decay products are below critical density, see their table. In addition, $K \rightarrow \pi f_A$ and $\mu \rightarrow e f_A$ are unseen. Combining these excludes $m_{\text{heavy}\nu}$ between 5×10^{-5} and 5×10^{-4} MeV (μ decay) and $m_{\text{heavy}\nu}$ between 5×10^{-5} and 0.1 MeV (K -decay).

Majoron Searches in Neutrinoless Double β Decay

Limits are for the half-life of neutrinoless $\beta\beta$ decay with a Majoron emission.

Previous indications for neutrinoless double beta decay with majoron emission have been superseded. No experiment currently claims any such evidence. For a review, see DOI 88.

VALUE (years)	CL%	DOCUMENT ID	TECN	COMMENT
$> 7.2 \times 10^{24}$	90	122 BERNATOW...	92 CNTR	^{128}Te
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$> 1.7 \times 10^{22}$	90	BECK	93 CNTR	^{76}Ge
$< 7.9 \times 10^{20}$	68	123 TANAKA	93 SPEC	^{100}Mo
$> 1.9 \times 10^{20}$	68	BARABASH	89 CNTR	^{136}Xe
$> 1.0 \times 10^{21}$	90	FISHER	89 CNTR	^{76}Ge
$> 3.3 \times 10^{20}$	90	ALSTON...	88 CNTR	^{100}Mo
$(6 \pm 1) \times 10^{20}$		AVIGNONE	87 CNTR	^{76}Ge
$> 1.4 \times 10^{21}$	90	CALDWELL	87 CNTR	^{76}Ge
$> 4.4 \times 10^{20}$	90	ELLIOTT	87 SPEC	^{82}Se
$> 1.2 \times 10^{21}$	90	FISHER	87 CNTR	^{76}Ge
		124 VERGADOS	82 CNTR	

122 BERNATOWICZ 92 studied double- β decays of ^{128}Te and ^{130}Te , and found the ratio $r(^{130}\text{Te})/r(^{128}\text{Te}) = (3.52 \pm 0.11) \times 10^{-4}$ in agreement with relatively stable theoretical predictions. The bound is based on the requirement that Majoron-emitting decay cannot be larger than the observed double-beta rate of ^{128}Te of $(7.7 \pm 0.4) \times 10^{24}$ year. We calculated 90% CL limit as $(7.7-1.28 \times 0.4=7.2) \times 10^{24}$.

123 TANAKA 93 also quote limit 5.3×10^{19} years on two Majoron emission.

124 VERGADOS 82 sets limit $g_H < 4 \times 10^{-3}$ for (dimensionless) lepton-number violating coupling, g_H , of scalar boson (Majoron) to neutrinos, from analysis of data on double β decay of ^{48}Ca .

INVISIBLE A^0 (AXION) MASS LIMITS FROM ASTROPHYSICS AND COSMOLOGY

Limits on $m(A^0)$ are obtained from the axion coupling to electrons, nucleons, or photons. Quoted limits are often expressed in terms of the axion decay constant f_A which can be defined in terms of the mass or axion-electron coupling by $m(A^0) = 3.5 \times 10^{10} g_{Ae} \cos^{-2} \beta \text{ eV} = 7.2 \times 10^7 (\text{GeV}/f_A) (N/6) \text{ eV}$ [using the conventions detailed in Srednicki [1]; for other conventions take $f_A \rightarrow 2f_A$ (Bardeen [2]) or $f_A \rightarrow 4f_A$ (Kaplan [3])] where N is the number of quarks with Peccei-Quinn charge (usually the number of quark flavors) and $\cos^2 \beta = v_1^2/(v_1^2 + v_2^2)$ is determined by the vacuum expectation values of the two Higgs doublets coupling to up and down quarks (and charged leptons). For the coupling to photons $m(A^0) = 6.9 \times 10^9 (g_{A\gamma}/\text{GeV}^{-1}) \text{ eV}$ and for the coupling to nucleons $m(A^0) = 7.7 \times 10^7 g_{AN}/c_{AN} \text{ eV}$ where c_{AN} depends on the details of the coupling of axions to nucleons. These couplings are defined by

$$\mathcal{L}_{\text{int}} = -\frac{1}{4} g_{A\gamma} \phi_A F_{\mu\nu} \tilde{F}^{\mu\nu} = g_{A\gamma} \phi_A \mathbf{E} \cdot \mathbf{B},$$

$$\mathcal{L}_{\text{int}} = i g_{Ae} \phi_A \bar{\psi}_e \gamma_5 \psi_e, \quad \text{and}$$

$$\mathcal{L}_{\text{int}} = i g_{AN} \phi_A \bar{\psi}_N \gamma_5 \psi_N.$$

The factors in these equations are model dependent, in particular $g_{Ae} = 0$ in the KSVZ [4] models. In the comment for each limit below, D indicates that the limit is specific to DFSZ [5] axions, K to KSVZ axions (The limits quoted assume $N = 6$ and $v_1 = v_2$.)

References

1. M. Srednicki, Nucl. Phys. **B260**, 689 (1985).
2. W. Bardeen and H. Tye, Phys. Lett. **74B**, 229 (1978).
3. D. Kaplan, Nucl. Phys. **B260**, 215 (1985).
4. J.E. Kim, Phys. Rev. Lett. **43**, 103 (1979); M.A. Shifman, A.I. Vainstein, and V.I. Zakharov, Nucl. Phys. **B166**, 493 (1980).
5. A.R. Zhitnitsky, Sov. J. Nucl. Phys. **31**, 260 (1980); M. Dine, W. Fischler, and M. Srednicki, Phys. Lett. **104B**, 199 (1981).

Invisible A^0 (Axion) MASS LIMITS from Astrophysics and Cosmology

$v_1 = v_2$ is usually assumed ($v_i =$ vacuum expectation values). For a review of these limits, see RAFFELT 90C and TURNER 90.

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 0.01	WANG	92 ASTR	D, white dwarf
< 0.03	WANG	92C ASTR	D, C-O burning
none 3-8	125 BERSHADY	91 ASTR	D, K, intergalactic light
< 10	126 KIM	91C COSM	D, K, mass density of the universe, super-symmetry
$< 1 \times 10^{-3}$	127 RAFFELT	91B ASTR	D,K, SN 1987A
none $10^{-3}-3$	128 RESSELL	91 ASTR	K, intergalactic light
	BURROWS	90 ASTR	D,K, SN 1987A
	129 ENGEL	90 ASTR	D,K, SN 1987A
< 0.02	130 RAFFELT	90D ASTR	D, red giant
$< 1 \times 10^{-3}$	131 BURROWS	89 ASTR	D,K, SN 1987A

See key on page 1343

Gauge & Higgs Boson Full Listings Axions (A^0) and Other Very Light Bosons

$<(1.4-10) \times 10^{-3}$	132 ERICSON	89	ASTR	D,K, SN 1987A
$< 3.6 \times 10^{-4}$	133 MAYLE	89	ASTR	D,K, SN 1987A
< 12	CHANDA	88	ASTR	D, Sun
$< 1 \times 10^{-3}$	RAFFELT	88	ASTR	D,K, SN 1987A
< 0.07	134 RAFFELT	88B	ASTR	red giant
< 0.7	FRIEMAN	87	ASTR	D, red giant
$< 2-5$	135 RAFFELT	87	ASTR	K, red giant
< 0.01	TURNER	87	COSM	K, thermal production
< 0.06	136 DEARBORN	86	ASTR	D, red giant
< 0.7	RAFFELT	86	ASTR	D, red giant
< 0.03	137 RAFFELT	86	ASTR	K, red giant
< 1	RAFFELT	86B	ASTR	D, white dwarf
$< 0.003-0.02$	138 KAPLAN	85	ASTR	K, red giant
$> 1 \times 10^{-5}$	IWAMOTO	84	ASTR	D, K, neutron star
$> 1 \times 10^{-5}$	ABBOTT	83	COSM	D,K, mass density of the universe
> 0.04	DINE	83	COSM	D,K, mass density of the universe
$> 1 \times 10^{-5}$	ELLIS	83B	ASTR	D, red giant
> 0.1	139 FUKUGITA	82	ASTR	D, stellar cooling
< 1	FUKUGITA	82B	ASTR	D, red giant

- 125 BERSHADY 91 searched for a line at wave length from 3100-8300 Å expected from 2γ decays of relic thermal axions in intergalactic light of three rich clusters of galaxies.
- 126 KIM 91C argues that the bound from the mass density of the universe will change drastically for the supersymmetric models due to the entropy production of saxion (scalar component in the axionic chiral multiplet) decay. Note that it is an upperbound rather than a lowerbound.
- 127 RAFFELT 91B argue that previous SN 1987A bounds must be relaxed due to corrections to nucleon bremsstrahlung processes.
- 128 RESSELL 91 uses absence of any intracuster line emission to set limit.
- 129 ENGEL 90 rule out $10^{-10} < g_{A\gamma} < 10^{-3}$, which for a hadronic axion with EMC motivated axion-nucleon couplings corresponds to $2.5 \times 10^{-3} \text{ eV} < m_{A^0} < 2.5 \times 10^4 \text{ eV}$. The constraint is loose in the middle of the range, i.e. for $g_{A\gamma} \sim 10^{-6}$.
- 130 RAFFELT 90D is a re-analysis of DEARBORN 86.
- 131 The region $m_{A^0} \gtrsim 2 \text{ eV}$ is also allowed.
- 132 ERICSON 89 considered various nuclear corrections to axion emission in a supernova core, and found a reduction of the previous limit (MAYLE 88) by a large factor.
- 133 MAYLE 89 limit based on naive quark model couplings of axion to nucleons. Limit based on couplings motivated by EM measurements is 2-4 times weaker. The limit from axion-electron coupling is weak: see HATSUDA 88B.
- 134 RAFFELT 88B derives a limit for the energy generation rate by exotic processes in helium-burning stars $\epsilon < 100 \text{ erg g}^{-1} \text{ s}^{-1}$, which gives a firmer basis for the axion limits based on red giant cooling.
- 135 RAFFELT 87 also gives a limit $g_{A\gamma} < 1 \times 10^{-10} \text{ GeV}^{-1}$.
- 136 DEARBORN 86 also gives a limit $g_{A\gamma} < 1.4 \times 10^{-11} \text{ GeV}^{-1}$.
- 137 RAFFELT 86 gives a limit $g_{A\gamma} < 1.1 \times 10^{-10} \text{ GeV}^{-1}$ from red giants and $< 2.4 \times 10^{-9} \text{ GeV}^{-1}$ from the sun.
- 138 KAPLAN 85 says $m_{A^0} < 23 \text{ eV}$ is allowed for a special choice of model parameters.
- 139 FUKUGITA 82 gives a limit $g_{A\gamma} < 2.3 \times 10^{-10} \text{ GeV}^{-1}$.

Search for Relic Invisible Axions

Limits are for $[G_{A\gamma\gamma}/m_{A^0}^2] \rho_A$ where $G_{A\gamma\gamma}$ denotes the axion two-photon coupling, $L_{\text{int}} = \frac{G_{A\gamma\gamma}}{4} \phi_A F_{\mu\nu} \tilde{F}^{\mu\nu} = G_{A\gamma\gamma} \phi_A \mathbf{E} \cdot \mathbf{B}$, and ρ_A is the axion energy density near the earth.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2 \times 10^{-41}$		140 HAGMANN	90 CNTR	$m_{A^0} = (5.4-5.9)10^{-6} \text{ eV}$
$< 1.3 \times 10^{-42}$	95	141 WUENSCH	89 CNTR	$m_{A^0} = (4.5-10.2)10^{-6} \text{ eV}$
$< 2 \times 10^{-41}$	95	141 WUENSCH	89 CNTR	$m_{A^0} = (11.3-16.3)10^{-6} \text{ eV}$

140 HAGMANN 90 experiment is based on the proposal of SIKIVIE 83.

141 WUENSCH 89 looks for condensed axions near the earth that could be converted to photons in the presence of an intense electromagnetic field via the Primakoff effect, following the proposal of SIKIVIE 83. The theoretical prediction with $[G_{A\gamma\gamma}/m_{A^0}^2] = 2 \times 10^{-14} \text{ MeV}^{-4}$ (the three generation DFSZ model) and $\rho_A = 300 \text{ MeV}/\text{cm}^3$ that makes up galactic halos gives $(G_{A\gamma\gamma}/m_{A^0}^2) \rho_A = 4 \times 10^{-44}$. Note that our definition of $G_{A\gamma\gamma}$ is $(1/4\pi)$ smaller than that of WUENSCH 89.

Invisible A^0 (Axion) Limits from Photon Coupling

Limits are for the axion-two-photon coupling $G_{A\gamma\gamma}$ defined by $L = G_{A\gamma\gamma} \phi_A \mathbf{E} \cdot \mathbf{B}$. Related limits from astrophysics can be found in the "Invisible A^0 (Axion) Mass Limits from Astrophysics and Cosmology" section.

VALUE (GeV $^{-1}$)	CL%	DOCUMENT ID	COMMENT
$< 3.6 \times 10^{-7}$	95	142 CAMERON	$m_{A^0} < 10^{-3} \text{ eV}$, optical rotation
$< 6.7 \times 10^{-7}$	95	143 CAMERON	$m_{A^0} < 10^{-3} \text{ eV}$, photon regeneration

$< 3.6 \times 10^{-9}$	99.7	144 LAZARUS	92	$m_{A^0} < 0.03 \text{ eV}$
$< 7.7 \times 10^{-9}$	99.7	144 LAZARUS	92	$m_{A^0} = 0.03-0.11 \text{ eV}$
$< 7.7 \times 10^{-7}$	99	145 RUOSO	92	$m_{A^0} < 10^{-3} \text{ eV}$
$< 2.5 \times 10^{-6}$		146 SEMERTZIDIS	90	$m_{A^0} < 7 \times 10^{-4} \text{ eV}$

- 142 Experiment based on proposal by MAIANI 86.
- 143 Experiment based on proposal by VANBIBBER 87.
- 144 LAZARUS 92 experiment is based on proposal found in VANBIBBER 89.
- 145 RUOSO 92 experiment is based on the proposal by VANBIBBER 87.
- 146 SEMERTZIDIS 90 experiment is based on the proposal of MAIANI 86. The limit is obtained by taking the noise amplitude as the upper limit. Limits extend to $m_{A^0} = 4 \times 10^{-3}$ where $G_{A\gamma\gamma} < 1 \times 10^{-4} \text{ GeV}^{-1}$.

Limit on Invisible A^0 (Axion) Electron Coupling

The limit is for $G_{Aee} \theta_{\mu\phi} A^0 \bar{e} \gamma_{\mu} e$ in GeV^{-1} , or equivalently, the dipole-dipole potential $\frac{G_{Aee}^2}{4\pi} ((\sigma_1 \cdot \sigma_2) - 3(\sigma_1 \cdot n)(\sigma_2 \cdot n))/r^3$ where $n=r/r$.

The limits below apply to invisible axion of $m_A \leq 10^{-6} \text{ eV}$.

VALUE (GeV $^{-1}$)	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.9 \times 10^{-5}$	95	147 BOBRAKOV	91	Induced magnetism
$< 1.9 \times 10^{-3}$	66	148 WINELAND	91	NMR
$< 8.9 \times 10^{-4}$	66	149 RITTER	90	Torsion pendulum
$< 6.4 \times 10^{-5}$	95	147 VOROBYOV	88	Induced magnetism

- 147 These experiments measured induced magnetization of a bulk material by the spin-dependent potential generated from other bulk material with aligned electron spins, where the magnetic field is shielded with superconductor.
- 148 WINELAND 91 looked for an effect of bulk matter with aligned electron spins on atomic hyperfine splitting using nuclear magnetic resonance.
- 149 RITTER 90 used a torsion pendulum to measure the potential between two bulk matter objects where the spins are polarized but without a net magnetic field in either of them.

REFERENCES FOR Searches for Axions (A^0) and Other Very Light Bosons

ATIYA	93	PRL 70 2521	+Chiang, Frank, Haggerty, Ito+	(BNL 787 Collab.)
Also	93C	PRL 71 305 (erratum)	Atiya, Chiang, Frank, Haggerty, Igo+	(BNL 787 Collab.)
ATIYA	93B	PR D48 2512	+Chiang, Frank, Haggerty, Ito+	(BNL 787 Collab.)
BASSOMPIERRE...	93	EPL 22 239	Bassompierre, Bologna	(LAPP, TORI, LYON)
BECK	93	PRL 70 2853	+Bensch, Bockholt, Heusser, Hirsch+	(MPIH, KIAE, SASSO)
CAMERON	93	PR D47 3707	+Cantatore, Melissinos+	(ROCH, BNL, FNAL, TRST)
MINOWA	93	PRL 71 4120	+Inoue, Asanuma, Imamura	(TOKY)
NG	93	PR D48 2941		(AST)
TANAKA	93	PR D48 5462	+Ejiri	(OSAK)
LAZARUS	92	PRL 68 2778	+Campagnari+	(BNL, FNAL, PSI, WASH, YALE)
ATIYA	92	PRL 69 733	+Chiang, Frank, Haggerty, Ito+	(BNL, LANL, PRIN, TRIU)
BERNATOW...	92	PRL 69 2341	+Bernatowicz, Brannon, Brazzle, Cowis+	(WUUL, TATA)
BLUEMLEIN	92	IJMP A7 3835	+Brunner, Grabosch+	(BERL, BUDA, JINR, SERP)
HALLIN	92	PR D45 3955	+Calaprice, McPherson, Saettler	(PRIN)
HENDERSON	92C	PRL 69 1733	+Asoka-Kumar, Greenberg, Lynn+	(YALE, BNL)
HICKS	92	PL B276 423	+Alburger	(OHIO, BNL)
LAZARUS	92	PL B276 2533	+Smith, Cameron, Melissinos+	(BNL, ROCH, FNAL)
MEIJERDREES	92	PL B6 3845	+Meijer, Walsman	(SINDRIM 1 Collab.)
RUOSO	92	ZPHY C56 505	+Cameron, Cantatore+	(ROCH, BNL, FNAL, TRST)
SKALSEY	92	PRL 68 456	+Kolata	(MICH, NDAM)
WANG	92	MPL A7 1497		(ILL)
WANG	92C	PL B291 97		(ILL)
WU	92	PRL 69 1729	+Asoka-Kumar, Greenberg, Henderson+	(BNL, YALE, CUNY)
AKOPYAN	91	PL B272 443	+Atayan, Ginenko, Sukhov	(INRM)
ASAI	91	PRL 66 2440	+Orto, Yoshimura, Haga	(ICEPP)
BERSHADY	91	PL 66 1398	+Messel, Turner	(CHIC, FNAL, EFT)
BLUEMLEIN	91	ZPHY C51 341	+Brunner, Grabosch+	(BERL, BUDA, JINR, SERP)
BOBRAKOV	91	JETPL 53 294	+Borisov, Lasakov, Serebrov, Tal'daev, Trofimova	(PNPI)
		Translated from ZETFP		53 283.
BROSS	91	PRL 67 2942	+Crisler, Pordes, Volk, Errede, Wrbanek	(FNAL, ILL)
KIM	91C	PRL 67 3465		(SEOUL)
RAFFELT	91B	PRL 67 2605	+Seckel	(MPIH, BART)
RESSELL	91	PR D44 3001		(CHIC, FNAL)
TRZASKA	91	PL B269 54	+Dejbaksh, Dutta, Li, Cormier	(TAMU)
TSERTOS	91	PL B266 259	+Kienle, Judge, Schreckenbach	(ILLG, GSI)
WALKER	91	APJ 376 51	+Steigman, Schramm, Olive+	(HSCA, OSU, CHIC, MINN)
WIDMANN	91	ZPHY A340 209	+Bauer, Connell, Maier, Major+	(STUT, GSI, STUTM)
WINELAND	91	PRL 67 1735	+Bollinger, Heinzen, Itano, Raizen	(NBSB)
ALBRECHT	90C	PL B246 278	+Ehrlichmann, Harder, Krueger+	(ARGUS Collab.)
ANTREASYAN	90	PL B251 204	+Bartels, Besset, Bielek, Bielek+	(Crystal Ball Collab.)
ASANUMA	90	PL B237 588	+Minowa, Tsukamoto, Orto, Tsunoda	(TOKY)
ATIYA	90	PRL 64 21	+Chiang, Frank, Haggerty, Ito, Kycia+	(BNL 787 Collab.)
ATIYA	90B	PRL 65 1188	+Chiang, Frank, Haggerty, Ito, Kycia+	(BNL 787 Collab.)
BAUER	90	NIM B50 300	+Briggmann, Carstangjen, Connell, et al	(STUT, VILL, GSI)
BURROWS	90	PR D42 3297	+Ressell, Turner	(ARIZ, CHIC, FNAL)
DEBOER	90	JPG 16 L1	de Boer, Lehmann, Steyaert	(LVLN)
ENGEL	90	PL 64 960	+Seckel, Hayes	(BART)
GINENENKO	90	PL B237 287	+Klubov, Poblaguev, Postoev	(INRM)
GUO	90	PR D41 2924	+Kaplan, Aide+	(NIU, LANL, FNAL, CASE, TEXA)
HAGMANN	90	PR D42 1297	+Sikivie, Sullivan, Tanner	(FNAL, GSI)
JUDGE	90	PL 65 972	+Krusche, Schreckenbach, Tsertos, Kienle	(ILLG, FLOR)
RAFFELT	90C	PR D41 1324		(MPIH)
RAFFELT	90D	PR D41 1324		(MPIH)
RITTER	90	PR D42 977	+Goldblum, Ni, Gillies, Speake	(VIRG)
SEMERTZIDIS	90	PRL 64 2968	+Cameron, Cantatore+	(ROCH, BNL, FNAL, TRST)
TSUCHIYAKI	90	PL B236 81	+Orto, Yoshida, Minowa	(ICEPP)
TURNER	90	PR D41 1324		(FNAL)
BARABASH	89	PL B223 273	+Kuzminov, Lobashev, Novikov+	(ITEP, INRM)
BINI	89	PL B221 99	+Fazzini, Giannatiempo, Poggi, Sona+	(FIRZ, CERN, AARH)
BURROWS	89	PR D39 1020	+Turner, Brinkmann	(ARIZ, CHIC, FNAL, BOCH)
Also	89B	PRL 60 1797	Turner	(FNAL, EFT)
DEBOER	89B	PRL 62 2639	de Boer, van Dantzig	(CERN, IPN)
ERICSON	89	PL B219 507	+Mathiot	(CERN, IFN)
FISHSNER	89	ZPHY C44 557	+Heinrigs, Preussger, Reitz, Samm+	(AACH3, BERL, PSI)
FISHER	89	PL B218 257	+Boehm, Bovet, Eggert+	(CIT, NEUC, PSI)
FOX	89	PR C39 288	+Kemper, Cottie, Zingarelli	(FSU)
MAYLE	89	PL B219 515	+Wilson, Ellis+	(LLL, CERN, MINN, FNAL, CHIC, OSU)
Also	89	PL B203 188	Mayle, Wilson+	(LLL, CERN, MINN, FNAL, CHIC, OSU)
MINOWA	89	PRL 62 2639	+Orto, Tsuchiyaki, Tsukamoto	(ICEPP)
ORITO	89	PRL 63 597	+Yoshimura, Haga, Minowa, Tsuchiyaki	(ICEPP)
PERKINS	89	PRL 62 2638		(OXF)

Gauge & Higgs Boson Full Listings

Axions (A^0) and Other Very Light Bosons

TSERTOS	89	PR D40 1397	+Kozhuharov, Armbruster, Kienle+	(GSI, ILLG)
VANBIBBER	89	PR D39 2089	+Van Bibber, McIntyre, Morris, Raffelt	(LLL, TAMU, LBL)
WUENSCH	89	PR D40 3153	+De Panfilis-Wuenssch, Semertzidis+	(ROCH, BNL, FNAL)
Also	87	PRL 59 839	+De Panfilis, Melissinos, Moskowitiz+	(ROCH, BNL, FNAL)
ALSTON-...	88	PRL 60 1928	+Alston-Garnjost, Dougherty+	(LBL, MTHO, UNM)
AVIGNONE	88	PR D37 618	+Baktash, Barker, Calaprice+(PRIN, SCUC, ORNL, WASH)	
BJORCKEN	88	PR D38 3375	+Ecklund, Nelson, Abashian+	(FNAL, SLAC, VPI)
BLINOV	88	SJNP 47 563	+Bondar, Bukin, Vorobeyev, Groshev+	(NOVO)
Translated from YAF	47	889		
BOLTON	88	PR D38 2077	+Cooper, Frank, Hallin+	(LANL, STAN, CHIC, TEMP)
Also	86	PRL 56 2461	+Bolton, Bowman, Cooper+	(LANL, STAN, CHIC, TEMP)
Also	86	PRL 57 3241	+Grosnick, Wright, Bolton+	(CHIC, LANL, STAN, TEMP)
CHANDA	88	PR D37 2714	+Nieves, Pai	(UMD, UPR, MASA)
CHOI	88	PR D37 3225	+Kim, Kim, Lam	(JHU)
CONNELL	88	PRL 60 2242	+Fearick, Hoernle, Sideras-Haddad, Sellschop	(WITW)
DATAR	88	PR C37 250	+Fortier, Gales, Hourani+	(IPN)
DEBOER	88	PRL 61 1274	+de Boer, van Dantzig	(ANIK)
Also	89	PRL 62 2644 erratum	+de Boer, van Dantzig	(ANIK)
Also	89	PRL 62 2638	+Perkins	(OXF)
Also	89B	PRL 62 2639	+de Boer, van Dantzig	(ANIK)
DEBOER	88C	JPG 14 1131	+de Boer, Deutsch, Lehmann, Prieels, Steyaert	(LLNL)
DOEHNER	88	PR D38 2722	+Last, Arnold, Freedman, Dubbers	(HEIDP, ANL, ILLG)
DOI	88	PR D37 2575	+Kotani, Takasugi	(ANIK)
EL-NADI	88	PR C37 2271	+Saday	(CAIR)
FAISSNER	88	ZPHY C37 231	+Heinrigs, Preussger, Reitz, Samm+	(AACH3, BERL, SIN)
HATSUDA	88B	PL B203 469	+Yoshimura	(KEK)
LORENZ	88	PL B214 10	+Mageras, Stiegler, Huszar	(MPIM, PSI)
MAYLE	88	PL B203 188	+Wilson+	(LLL, CERN, MINN, FNAL, CHIC, OSU)
PICCIOTTO	88	PR D37 1131	+Ahmad, Britton, Bryman, Clifford+	(TRIU, CNRC)
RAFFELT	88	PRL 60 1793	+Seckel	(UCB, LLL, UCSC)
RAFFELT	88B	PR D37 549	+Dearborn	(UCB, LLL)
SAVAGE	88	PR C37 1134	+Filippone, Mitchell	(CIT)
TSERTOS	88	PL B207 273	+Kozhuharov, Armbruster, Kienle+	(GSI, ILLG)
TSERTOS	88B	ZPHY A331 103	+Kozhuharov, Armbruster, Kienle+	(GSI, ILLG)
VANKLINKEN	88	PL B205 223	+van Klinken, Meiring, de Boer, Schaafsma+	(GRON, GSI)
VANKLINKEN	88B	PRL 60 2442	+van Klinken	(GRON)
VONWIMMER..	88	PRL 60 2443	+von Wimmersperg	(BNL)
VOROBYOV	88	PL B208 146	+Gitaris	(NOVO)
AVIGNONE	88	PR C36 1987	+Brodzinski, Miley, Reeves	(SCUC, PNL)
AIP Conf. Proc.		Salt Lake City, UT		
CALDWELL	87	PRL 59 419	+Eisberg, Grumm, Witherell+	(UCSB, LBL)
DRUZHININ	87	ZPHY C37 1	+Dubrovina, Eidelman, Golubev+	(NOVO)
ELLIOTT	87	PRL 59 1649	+Hahn, Moe	(UCI)
FISHER	87	PL B192 460	+Boehm, Bovet, Egger+	(CIT, NEUC, SIN)
FRIEMAN	87	PR D36 2201	+Dimopoulos, Turner	(SLAC, STAN, FNAL, EPI)
GOLDMAN	87	PR D36 1543	+Hallin, Hoffman+	(LANL, CHIC, STAN, TEMP)
KORENCHEN...	87	SJNP 46 192	+Korenchenko, Kostin, Mzhaviya+	(JINR)
Translated from YAF	46	313		
MAIER	87	ZPHY A326 527	+Bauer, Briggmann, Carstjen+	(STUT, GSI)
MILLS	87	PR D36 707	+Levy	(BELL)
RAFFELT	87	PR D36 2211	+Dearborn	(LLL, UCB)
RIORDAN	87	PRL 59 755	+Krasny, Lang, Barbaro, Bodek+	(ROCH, CIT+)
TURNER	87	PRL 59 2489	+Filippone	(FNAL, EPI)
VANBIBBER	87	PRL 59 759	+Van Bibber, Dagdeviren, Koonin+(LLL, CIT, MIT, STAN)	
VONWIMMER..	87	PRL 59 266	+von Wimmersperg, Connell, Hoernle, Sideras-Haddad(WITW)	
ALBRECHT	86D	PL B179 403	+Binder, Boeckmann+	(ARGUS Collab.)
BADIER	86	ZPHY C31 21	+Bemporad, Boucrot, Callot+	(NA3 Collab.)
BOWCOCK	86	PRL 56 2676	+Giles, Hassard, Kinoshita+	(CLEO Collab.)
BROWN	86	PRL 57 2101	+ (FNAL, WASH, KYOT, KEK, COLU, STON, SACL)	
BRYMAN	86B	PRL 57 2787	+Clifford	(TRIU)
DAVIER	86	PL B180 295	+Jeanjean, Nguyen Ngoc	(LALO)
DEARBORN	86	PRL 56 26	+Schramm, Steigman	(LLL, CHIC, FNAL, BART)
EICHLER	86	PL B175 101	+Felawka, Kraus, Niebuhr+	(SINDRUM Collab.)
HALLIN	86	PRL 57 2105	+Calparic, Dunford, McDonald	(PRIN)
JODDIO	86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+	(LBL, NWES, TRIU)
Also	88	PR D37 237 erratum	+Joddio, Balke, Carr+	(LBL, NWES, TRIU)
KETOV	86	JETPL 44 146	+Klimov, Nikolaev, Mikaelian+	(KIAE)
Translated from ZETFP		44 114		
KOCH	86	NC 96A 182	+Schult	(JUL)
KONAKA	86	PRL 57 659	+Imai, Kobayashi, Masaike, Miyake+	(KYOT, KEK)
MAGERAS	86	PRL 56 2672	+Franzini, Tuts, Youssef+	(MPIM, COLU, STON)
MAIANI	86	PL B175 359	+Petronzio, Zavattini	(CERN)
PECCEI	86	PL B172 435	+Wu, Yanagida	(DESY)
RAFFELT	86B	PR D35 897	+McKeown, Filippone, Mitchell	(MPIM)
RAFFELT	86B	PL 166B 402		(CIT)
SAVAGE	86B	PRL 57 178	+Carbani, Jonson, Thun	(CERN)
AMALDI	85	PL 153B 444	+Kalinina, Lushchikov, Olshevskii+	(JINR)
ANANEV	85	SJNP 41 585		
Translated from YAF	41	912		
BALTRUSAIT...	85	PRL 55 1842		
BERGSMA	85	PL 121B 193		
KAPLAN	85	NP B260 215		
IVAMOTO	84	PRL 53 1198		
YAMAZAKI	84	PRL 52 1089		
ABBOTT	83	PL 120B 133		
ALAM	83	PR D27 1665		
CARBONI	83	PL 123B 349		
CAVAIGNAC	83	PL 121B 193		
DICUS	83	PR D28 1778		
DINE	83	PL 120B 137		
ELLIS	83B	NP B223 252		
FAISSNER	83	PR D28 1198		
FAISSNER	83B	PR D28 1787		
FRANK	83B	PR D28 1790		
HOPFMAN	83	PR D28 660		
NICZYPOURK	83	ZPHY C17 197		
PRESKILL	83	PL 120B 127		
SIKIVIE	83	PRL 51 1415		
Also	84	PRL 52 695 erratum		
ALEKSEEV	82	JETP 55 591		
Translated from ZETF	82	1007		
ALEKSEEV	82B	JETPL 36 116		
Translated from ZETFP	36	94		
ASANO	82	PL 113B 195		
BARROSO	82	PL 116B 247		
DATAR	82	PL 114B 63		
EDWARDS	82	PRL 48 903		
FETSCHER	82	JPG 8 1147		
FUKUGITA	82	PR 48 1522		
FUKUGITA	82B	PR D26 1840		
LEHMANN	82	PL 115B 270		
RAFFELT	82	PL 119B 323		
SIVERTZ	82	PR D26 717		
VERGADOS	82	PL 109B 96		
ZEHNDER	82	PL 110B 419		
ASANO	81B	PL 107B 159		
BARROSO	81B	PL 106B 91		
FAISSNER	81	ZPHY C10 95		
FAISSNER	81B	PL 103B 234		
KIM	81	PL 105B 55		
VUILLEUMIER	81	PL 101B 341		
ZEHNDER	81	PL 104B 494		
FAISSNER	80	PL 96B 201		
JACQUES	80	PR D21 1206		
SOUKAS	80	PRL 44 564		
BECHIS	79	PRL 42 1511		
CALAPRICE	79	PR D20 2708		
COTEUS	79	PRL 42 1438		
DISHAW	79	PL 85B 142		
ZHITNITSKII	79	SJNP 29 517		
Translated from YAF	29	1001		
ALIBRAN	78	PL 74B 134		
ASRATYAN	78B	PL 79B 497		
BELLOTTI	78	PL 76B 223		
BOSETTI	78B	PL 74B 143		
DICUS	78C	PR D18 1829		
DONNELLY	78	PR D18 1607		
Also	76	PRL 37 315		
Also	74	PRL 33 179		
HANSL	78D	PL 74B 139		
MICELMAC...	78	PL 74B 139		
MIKAEIAN	78	PR D18 3605		
SATO	78	PTP 60 1942		
VYSOTSII	78	JETPL 27 502		
Translated from ZETFP	27	533		
YANG	78	PRL 41 523		
PECCEI	77	PR D16 1791		
Also	77B	PRL 38 1440		
REINES	76	PRL 37 315		
GURR	74	PRL 33 179		
ANAND	53	PRSL A22 183		
Baltrusaitis, Becker, Blaylock, Brown+		(Mark III Collab.)		
+Dorenbosch, Alibay, Amaldi+		(CHARM Collab.)		
		(HARY)		
		(UCSB, WUSL)		
		(INUS, KEK)		
+Ishikawa, Taniguchi, Yamanaka+		(BRAN, FLOR)		
+Sikivie		(VAND, CORN, ITHA, HARV, OHIO, ROCH+)		
		(CERN, MUNI)		
+Dahme		(Hoummada, Koang, Ost+)		
		(TEXA, UMD)		
+Fischler		(IAS, PENN)		
+Olive		(CERN)		
+Heinrigs, Preussger, Samm		(AACH)		
+Frenzel, Heinrigs, Preussger+		(AACH3)		
+ (LANL, YALE, LBL, MIT, SACL, SIN, CNRC, BERN)		(LANL, ARZS)		
+Frank, Mischke, Moir, Schardt		(LENA Collab.)		
+Jakubowski, Zeludzewicz+		(HARV, UCSBT)		
+Wise, Wilczek		(FLOR)		
		(FLOR)		
Sikivie		(KIAE)		
+Kartamyshev, Makarin+		(MOSU, JINR)		
Translated from ZETF	82	1007		
Alekhin, Kruglov, Kulikov+		(MOSU, JINR)		
+Kikutani, Kurokawa, Miyachi+(KEK, TOKY, INUS, OSAK)		(LIBS)		
+Branco		(BHAB)		
+Baba, Betigeri, Singh		(Crystal Ball Collab.)		
+Partridge, Peck, Porter+		(ETH)		
		(ETH)		
+Watanura, Yoshimura		(KEK)		
+Kukhopadhyay		(KEK)		
+Lesauy, Muller, Zylinderajch		(SACL)		
+Stodolsky		(MPIM)		
+Lee-Franzini, Horstkotter+		(CUSB Collab.)		
		(CERN)		
+Gabathuler, Vuilleumier		(ETH, SIN, CIT)		
+Kikutani, Kurokawa, Miyachi+(KEK, TOKY, INUS, OSAK)		(AACH3)		
+Frenzel, Grimm, Hansl, Hoffman+		(SIN)		
+Frenzel, Heinrigs, Preussger+		(AACH3)		
+Stamm		(AACH3)		
+Boehm, Hahn, Kwon+		(CIT, MUNI)		
		(ETH)		
+Frenzel, Heinrigs, Preussger, Samm+		(AACH3)		
+Kalelkar, Miller, Plano+		(RUTG, STEV, PENN)		
+Wanderer, Weng+		(BNL, HARV, ORNL, PENN)		
+Dombeck+		(UMD, COLU, AFRR)		
+Dunford, Kouzes, Miller+		(PRIN)		
+Diesburg, Fine, Lee, Sokolsky+		(COLU, ILL, BNL)		
+Diamant-Berger, Faessler, Liu+		(SLAC, CIT)		
+Skovpen		(NOVO)		
Translated from YAF	29	1001		
+Armenise, Arnold, Bartley		(Gargamelle Collab.)		
+Epstein, Fakhruddinov+		(ITEP, SERP)		
+Fiorini, Zanotti		(MILA)		
+Deden, Deuschmann, Fritze+		(BEBC Collab.)		
+Kolb, Tepitz, Wagoner		(TEXA, VPI, STAN)		
+Freedman, Lytel, Pecci, Schwartz		(STAN)		
+Reines, Gurr, Sobel		(UCI)		
+Gurr, Reines, Sobel		(UCI)		
+Holder, Knobloch, May, Paar+		(CDHS Collab.)		
+Micelmacher, Pontecorvo		(JINR)		
		(FNAL, NWES)		
		(KYOT)		
+Zeldovich, Khlopov, Chechetkin		(ASC)		
		(MASA)		
+Quinn		(STAN, SLAC)		
+Pecci, Quinn		(STAN, SLAC)		
+Gurr, Sobel		(UCI)		
+Reines, Sobel		(UCI)		
SREDNICKI	85	NP B260 689		
BARDEEN	78	PL 74B 229		
		(UCSB)		
		(FNAL)		

OTHER RELATED PAPERS

SREDNICKI	85	NP B260 689		
BARDEEN	78	PL 74B 229	+Tye	(UCSB)
				(FNAL)

LEPTONS

ν_e	1389
ν_μ	1392
ν_τ	1395
e	1396
μ	1397
τ	1403
Number of Light Neutrino Types	1416
Heavy Lepton Searches	1418
Massive Neutrinos and Lepton Mixing	1421
Neutrino Bounds from Astrophysics and Cosmology	1432

QUARKS

u	1437
d	1437
s	1437
c	1439
b	1439
Searches for t Quarks	1439
b' (Fourth Generation) Quark	1442

Notes in the Lepton Listings

Note on Neutrinos	1385
Note on the Muon Neutrino Mass	1392
Note on Muon Decay Parameters	1399
Note on Problems in the τ -Decay Data	1403
Note on τ -Decay Branching Fractions	1406
Note on the Number of Light Neutrino Types from Collider Experiments	1416
Note on Heavy Lepton Searches	1418
Note on Neutrino Oscillation Experiments	1424
Note on Constraints on Particles from SN 1987A	1432

Notes in the Quark Listings

Note on Quark Mass	1433
--------------------	------

LEPTONS

NOTE ON NEUTRINOS

(by R.E. Shrock, State Univ. of New York, Stony Brook)

In addition to the ν_e , ν_μ , and ν_τ sections, the *Review of Particle Properties* includes sections on "Searches for Massive Neutrinos and Lepton Mixing," "Number of Light Neutrino Types," "Heavy Lepton Searches," and "Constraints from Cosmology and Astrophysics."

The theoretical perspective concerning neutrino masses has changed considerably over the past 20 years. Before that time, a standard view was that there was no theoretical reason for neutrinos to have masses, which was in accord with the striking fact that the upper limits on their masses were much smaller than those of the associated charged leptons. It was also noted that experimental data were consistent with the "laws" of lepton family number and total lepton number conservation. (Some early discussions of neutrino oscillations and lepton mixing are given in Refs. 1 and 2). Indeed, even in the literature of the 1970's, one will often find statements asserting that in the standard $SU(2) \times U(1)$ electroweak theory (without electroweak-singlet neutrinos) the known (electroweak-doublet) neutrinos are massless.

In contrast, in the modern theoretical view based on the standard electroweak theory (and its supersymmetric extensions which stabilize the hierarchy), small but nonzero neutrino masses are expected on general grounds. The reason for this is as follows. Given only the known left-handed neutrino fields and the usual Higgs field(s) in the Standard Model (and supersymmetric extensions thereof which stabilize the hierarchy), nonzero neutrino masses result generically from higher-dimension operators which are strongly expected to occur at a scale near to that of quantum gravity. This expectation is based on the general consensus that pointlike theories of quantum gravity are nonrenormalizable, and is borne out by explicit calculations of the low-energy, field-theory limit of string theories. Such higher-dimension operators would be suppressed by associated inverse powers of the (reduced) Planck mass, $\kappa^{-1} \equiv \overline{M}_P \equiv \sqrt{\hbar c / (8\pi G_N)} = 2.4 \times 10^{18}$ GeV. For example, there would be a gauge-invariant dimension-5 operator consisting of the weak $I = 1$, $Y = -2$ Majorana bilinear $\nu_L^T C \nu_L$ (where generation indices are suppressed) contracted with a symmetric, $I = 1$ quadratic product of the usual $I = 1/2$, $Y = 1$ Higgs, with a coefficient of the form a/\overline{M}_P , where a is a dimensionless constant. From the vacuum expectation values of the Higgs, this would give rise to (Majorana) neutrino masses of order $m_\nu \sim av^2/\overline{M}_P$, where $v = 250$ GeV is the scale of electroweak-symmetry breaking. (Here \overline{M}_P is an approximate upper bound on the mass which suppresses such operators; it is possible that new physics occurs at some intermediate mass scale $v \ll M_I < \overline{M}_P$ in such a way that dimension-5 operators of this type would give rise to neutrino masses of order av^2/M_I .)

This not only leads one to expect nonzero neutrino masses but explains why they are so small.

In retrospect, one sees that this change in theoretical perspective is associated with the change in viewpoint concerning quantum field theory. For decades after the success of quantum electrodynamics in the late 1940's, renormalizability was taken as a necessary property for an acceptable fundamental quantum field theory. It was this implicit theoretical assumption which led to the oft-repeated statements in the 1970's that in the standard electroweak theory (without electroweak-singlet neutrinos) the known neutrinos are massless. More recently, it has been appreciated that renormalizability and, in particular, the great success of the Standard Model with its exclusion of any higher-dimension nonrenormalizable operators, may well be due only to the fact that the electroweak scale v is much smaller than \overline{M}_P (or possibly even some lower scale, M_I), *i.e.*, a consequence of the hierarchy. Once one includes such higher-dimension operators in one's considerations, the realization that neutrino masses are generic follows immediately. A summary of this modern view is given, *e.g.*, in Ref. 3.

In contrast to this mechanism for neutrino mass, which only relies upon the known neutrinos, together with the Higgs field(s) of the Standard Model (or its supersymmetric extensions), there is another more speculative mechanism which we mention here for completeness. It is not known whether there exist any electroweak-singlet neutrino fields. If they do exist, then they could lead, via renormalizable, dimension-4 operators, to neutrino masses $m_\nu \sim v^2/M_R$, where the scale M_R of the electroweak-singlet neutrino mass is naturally $\gg v$, again yielding, albeit for a different reason, very small m_ν [4].

In turn, a natural concomitant of (nondegenerate) neutrino masses is lepton mixing, which is thus also a general expectation. The lepton-mixing angles are functions of ratios of elements of neutrino-matrix elements and of charged lepton mass matrix elements, and even though left-handed neutrino masses are small, some of these ratios could, in principle, be $\mathcal{O}(1)$, which raises the issue of why such effects have not been seen. This question was answered as follows: a set of conditions for natural suppression of observable lepton flavor violation were formulated, and it was shown that the Standard Model (generalized to include nonzero m_ν) satisfies these [5]. This explains why the "law" of lepton family number conservation is obeyed to such high accuracy.

After these theoretical points, let us return to a description of the quantities upon which various experiments put limits. As an aid to understanding the limits on neutrino masses and lepton mixing, we recall that, in contrast to other particles in this *Review*, the neutrinos ν_e , ν_μ , and ν_τ are defined as weak eigenstates (the weak $I_3 = 1/2$ components of the $SU(2)_L$ lepton doublets) which couple with unit strength to e , μ , and τ , respectively. These neutrino weak eigenstates are not, in general, states of definite mass. If one assumes that neutrinos are massless, and hence degenerate, then it is possible to define the

Lepton & Quark Full Listings

Neutrinos

weak eigenstates to be simultaneously mass eigenstates. However, in the general case of possibly massive (nondegenerate) neutrinos, the weak eigenstates have no well-defined masses, but instead are linear combinations of mass eigenstates. Let us denote the charged leptons as the set $\{\ell_a\}$, $a = 1, \dots, n$, where $n \geq 3$, with $\ell_1 = e$, $\ell_2 = \mu$, and $\ell_3 = \tau$. From the LEP measurement of the Z width (see section on “Number of Light Neutrinos”), one knows that there are only three neutrinos which couple to the Z in the usual way and have masses $m_\nu < m_Z/2$. Of course, this measurement does not preclude the existence of electroweak-singlet neutrinos. The latter are often called “right-handed neutrino singlets,” although, since they are singlets, it is a convention whether one chooses to write them as $(\chi_j)_R$ or $(\chi'_j)_L = (\chi_j^c)_L$. The left-handed components of the weak eigenstates of the neutrinos, $(\nu_{\ell_a})_L$ can be expressed in terms of mass eigenstates by the transformation

$$(\nu_{\ell_a})_L = \sum_j U_{aj}(\nu_j)_L \quad (1)$$

where the $\{\nu_j\}$ denote these mass eigenstates and consist of n members together with possible additional $SU(2) \times U(1)$ singlet neutral leptons, often called “sterile” neutrinos. The ordering of the mass eigenbasis can be defined so that U is as nearly diagonal as possible, *i.e.* (with no sum on j) $|U_{jj}| \geq |U_{jk}|$, $k \neq j$. Of course, this does not imply that $m_{\nu_j} > m_{\nu_k}$ for $j > k$.

Thus, as was noted in Ref. 6, decays such as ${}^3\text{H} \rightarrow {}^3\text{He} e^- \bar{\nu}_e$ and $\pi^+ \rightarrow \mu^+ \nu_\mu$, which have been used to set the best bounds on the respective neutrino masses, really consist of sums of the separate decay modes ${}^3\text{H} \rightarrow {}^3\text{He} e^- \bar{\nu}_j$ and $\pi^+ \rightarrow \mu^+ \nu_k$, where the ν_j and ν_k are mass eigenstates, and the indices j and k range over all of the values allowed by phase space in these respective decays. The coupling strengths for the j 'th mode in ${}^3\text{H}$ β decay and the k 'th mode in π^+ μ_2^+ decay are given, respectively, by $|U_{1j}|^2$ and $|U_{2k}|^2$. In general, these modes are incoherent, although in the limit in which the ν_j all become degenerate they would become coherent. There are, in addition certain kinematic factors depending on the m_{ν_j} which enter in determining the branching ratio for a given decay mode. Assuming that the off-diagonal elements of the lepton mixing matrix U are small relative to the diagonal elements, the dominantly coupled decays are the ones with coupling strength $|U_{aj}|^2$, $a = j$, *i.e.*, ${}^3\text{H} \rightarrow {}^3\text{He} e^- \bar{\nu}_1$ and $\pi^+ \rightarrow \mu^+ \nu_2$.

Hence, it follows that the neutrino mass limits quoted in the literature for “ m_{ν_e} ,” “ m_{ν_μ} ,” and “ m_{ν_τ} ” should really be interpreted as limits on the corresponding mass eigenstates [6,7]. Specifically, a bound on “ m_{ν_e} ” from a study of tritium β decay, for example, really constitutes a weighted limit on each of the mass eigenstates ν_j in the weak eigenstate ν_e which are kinematically allowed to occur in tritium decay and which are coupled with strength $|U_{1j}|^2$ sufficiently large to make a significant contribution to the observed spectrum. It is thus certainly a limit on ν_1 , since this is, by the definition, of the order of the mass eigenbasis, the dominantly coupled neutrino.

If lepton mixing is hierarchical, as quark mixing is known to be, *i.e.*, if $|U_{jj}|^2 \gg |U_{jk}|^2$, $j \neq k$, then ν_1 is the only mass eigenstate significantly constrained by a bound on “ m_{ν_e} .” Furthermore, strictly speaking, a neutrino mass limit cannot be stated in isolation; it always contains some implicit dependence on the relevant lepton-mixing angles. This dependence is fortunately relatively unimportant for the dominantly coupled decay modes, *i.e.*, $e \bar{\nu}_1$, $\mu \bar{\nu}_2$, and $\tau \bar{\nu}_3$ and hence the mass limits on “ m_{ν_e} ,” “ m_{ν_μ} ,” and “ m_{ν_τ} ” can be reinterpreted as being limits on m_{ν_j} , $j = 1, 2$, and 3, respectively.

There are three general types of (Lorentz-invariant) neutrino mass terms: Dirac masses of the form $m_D \bar{\nu}_L \chi_R + h.c.$, left-handed Majorana masses of the form $m_L \bar{\nu}_L \nu_R^c + h.c. = m_L^* \nu_L^T C \nu_L + h.c.$ and right-handed Majorana masses of the form $m_R \bar{\chi}_L^c \chi_R + h.c. = m_R \chi_R^T C \chi_R + h.c.$, where C is the Dirac charge conjugation matrix. Dirac mass terms conserve total lepton number L_{tot} , while Majorana mass terms violate L_{tot} . In the standard electroweak theory, extended to include massive neutrinos, (i) a Dirac mass term transforms as a weak $I = 1/2$ operator, and is coupled to the $I = 1/2$ Higgs to make an $SU(2) \times U(1)$ singlet operator; (ii) a Majorana mass term involving the $I = 1/2$ left-handed neutrinos transforms as $I = 1$ and must be coupled to an operator with $I = 1$ (and $Y = 2$) to make a gauge-invariant singlet; (iii) a Majorana-mass term involving the $SU(2) \times U(1)$ singlet neutral leptons, conventionally considered to be right-handed, is a singlet; it could be present as a bare mass term or couple to some other singlet operator. Note that in the minimal supersymmetric standard model (MSSM), which has two Higgs doublets, of hypercharge $Y = 1$ and $Y = -1$, the Dirac neutrino mass term arises from the cubic chiral superfield coupling $\epsilon_{ij} \hat{L}^i \hat{\chi}^c \hat{H}_u^j$ (all chiral superfields are taken as left-handed), where \hat{H}_u is the same Higgs that gives mass to the $Q = 2/3$ quarks. The Dirac neutrino mass terms are thus proportional to $\sin \beta$, where $\tan \beta = v_u/v_d$ is the ratio of the vacuum expectation values of the two Higgs in the MSSM.

In general, in the Standard Model, in addition to the three known left-handed $I = 1/2$ lepton doublets, there could be some number n_s of electroweak-singlet neutrinos. In a compact notation, one can then denote ν_L as the 3-component vector of left-handed $I = 1/2$ neutrinos and χ_R to be the n_s -dimensional vector of electroweak-singlet singlets, taken to be right-handed. In supersymmetric extensions of the Standard Model, the χ_R arise as (conjugates of) the spin 1/2 component fields in (left-handed) gauge-singlet chiral superfields. The general neutrino mass term in the Lagrangian is then given by

$$-\mathcal{L}_m = \frac{1}{2} (\bar{\nu}_L, \bar{\chi}_L^c) \begin{pmatrix} M_L & M_D \\ M_D^T & M_R \end{pmatrix} \begin{pmatrix} \nu_R^c \\ \chi_R \end{pmatrix} + h.c. \quad (2)$$

where M_L is the 3×3 left-handed Majorana-mass matrix, M_R is a $n_s \times n_s$ right-handed Majorana-mass matrix, and M_D is the 3-row by n_s -column Dirac-mass matrix. In general, all of these mass matrices are complex. The anticommutativity of fermion fields and the property that $C \gamma_\mu C^{-1} = -\gamma_\mu^T$

together imply that the Majorana mass matrices are symmetric: $M_L = M_L^T, M_R = M_R^T$. The diagonalization of the full $(3 + n_s) \times (3 + n_s)$ mass matrix in Eq. (2) yields $3 + n_s$ mass eigenstates, which are, in general, of Majorana type. Since Majorana mass terms violate total lepton number, one sees from a general viewpoint that one does not expect conservation of total lepton number. In particular, the dimension-5 operators discussed above give rise to left-handed Majorana neutrino mass terms and violate total lepton number. Dirac-neutrinos can be constructed from two Majorana-neutrino mass eigenstates whose masses are equal in magnitude [8]. For this reason, Dirac neutrino masses may be considered to be a special (degenerate) case of Majorana neutrino masses, and the latter may be regarded as the generic case. From the similarity transformation which diagonalizes the neutrino mass matrix, together with the similarity transformation which diagonalizes the charged lepton mass matrix (where, of course, only Dirac masses are allowed by electric-charge conservation), one constructs the lepton-mixing matrix U . In general, since U is not the identity, neutrino masses naturally give rise to lepton family number violation.

In supersymmetric extensions of the Standard Model, the neutrinos could, *a priori*, mix with the neutralinos (higgsinos and neutral gauginos). However, the usual R parity which is invoked to forbid unacceptably rapid proton decay also prevents such mixing between neutrinos and neutralinos.

In addition to mass and lifetime limits, this *Review* includes limits on various other possible properties, including electric charge, the CPT -violating difference $m_{\nu_1} - m_{\bar{\nu}_1}$, and a magnetic dipole moment. These are of interest because a massless purely chiral Dirac neutrino cannot have a magnetic (or electric) dipole moment. In the standard electroweak theory, extended to allow for Dirac neutrino masses, the neutrino magnetic dipole moment is nonzero and given [5,9], as

$$\mu_{\nu_j} = \frac{3eG_F m_{\nu_j}}{8\pi^2 \sqrt{2}} = 3.2 \times 10^{-19} (m_{\nu_j}/1 \text{ eV}) \mu_B \quad (3)$$

where G_F is the Fermi constant and $\mu_B = e/2m_e$ is the Bohr magneton. The neutrino electric dipole moment violates both time-reversal invariance and parity; although it is nonzero in general, it is quite small (see, *e.g.* Ref. 10). Again, however, we note that Dirac neutrinos should be regarded as a special case; the generic case is Majorana neutrinos. The operator products which define the magnetic and electric-dipole moments, *viz.*, $\bar{\nu} \sigma_{\alpha\beta} \nu F^{\alpha\beta}$ and $\bar{\nu} \sigma_{\alpha\beta} \gamma_5 \nu F^{\alpha\beta}$, respectively (where $F^{\alpha\beta}$ is the electromagnetic field strength tensor) vanish identically if ν is a Majorana neutrino because of the Majorana property that $\nu^c = \pm\nu$. Thus, a Majorana neutrino has identically zero magnetic and electric dipole moments.

Only the diagonal magnetic- and electric-dipole moments are static properties of a given neutrino-mass eigenstate. Transition magnetic and electric dipole moments exist in general for both Dirac and Majorana neutrinos but are not static properties and hence are not considered here. Occasionally, one also finds references to the “neutrino charge

radius” in the literature. This is defined via the Taylor series expansion of the generalized vector Dirac form factor multiplying γ_μ in the electromagnetic current matrix element: $F_1^V(q^2) = F_1^V(0) + q^2 dF_1^V/dq^2|_{q^2=0} + \mathcal{O}[(q^2)^2]$, where q denotes the 4-momentum of the photon [see, *e.g.* Ref. 5 Eq. (2.20)]. The electric charge is $Q = F^V(0) = 0$ for a neutrino, and the charge radius is given by $\langle r^2 \rangle = (1/6)(F_1^V)'(0)$. However, since this is multiplied by q^2 in the Taylor series expansion, it never occurs for a real photon, where $q^2 = 0$, and hence is not an S-matrix element, *i.e.*, not a physical quantity. In a gauge theory, this is manifested in the fact that the charge radius is gauge-dependent.

If one considers the possibility of nonzero masses for neutrinos, for consistency one must then also consider the leptonic mixing which would in general occur concomitantly. Accordingly, this *Review* devotes a section to correlated bounds on neutrino masses and lepton mixing angles. These can be divided into two types. First, there are those due to decays involving neutrinos in the final state, which must be recognized to have the possible multimode structure pointed out above. In the two most sensitive cases suggested as tests for neutrino masses and mixing, one obtains a limit on m_{ν_j} and $|U_{aj}|^2$ individually for each j . The peak-search test proposed in Ref. 6 was applied to existing data in that paper and a subsequent one [7]; it was applied in new experiments on 2-body leptonic decays of K^+ and π^+ by several groups at SIN (PSI), KEK, and TRIUMF. The results are catalogued in corresponding subsections on limits on $|U_{1j}|^2$ and $|U_{2j}|^2$. The kink-search test was also applied by a number of groups. The experimental situation, which was controversial for many years, has recently been clarified (see below).

Second, there are those due to processes involving the propagation and subsequent interaction of neutrinos. The latter are often called neutrino-oscillation limits, although this term is strictly correct only if the differences in neutrino masses are sufficiently small relative to their momenta that the propagation is effectively coherent in a quantum mechanical sense; otherwise, the individual ν_j from a given decay such as $\pi_{\mu 2}$ or $K_{\mu 2}$ propagate in a measurably incoherent manner, and there is no oscillation. Experimentalists usually present their results in terms of a simplifying model in which mixing is assumed to occur only between two neutrino species. The relevant transformation equation becomes

$$\begin{pmatrix} \nu_{\ell_a} \\ \nu_{\ell_b} \end{pmatrix} = \begin{pmatrix} \cos \theta & \sin \theta \\ -\sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} \nu_i \\ \nu_j \end{pmatrix} \quad (4)$$

where ν_{ℓ_a} are the weak-neutrino eigenstates, with $\nu_{\ell_1} = \nu_e$, *etc.*, and ν_i are neutrino mass eigenstates. Let the distance between the source of the neutrinos and their point of interaction be denoted L , and their energy as E . Assume furthermore that the m_{ν_i} are such that the coherence assumption is valid. Then the probability of an initial ν_{ℓ_a} , having propagated for a distance $t = L$ (with $1 - v/c \ll 1$) being equal to ν_{ℓ_b} is given by

$$P \equiv |\langle \nu_{\ell_b} | \nu_{\ell_a}(t) \rangle|^2 = \sin^2 2\theta \sin^2 \left(\frac{\Delta m^2 L}{4E} \right) \quad (5)$$

Lepton & Quark Full Listings

Neutrinos

where

$$\Delta m^2 = m_{\nu_i}^2 - m_{\nu_j}^2. \quad (6)$$

Thus, neutrino-oscillation experiments cannot measure individual neutrino masses, but only differences of masses squared, and indeed these are generally weighted in a more complicated way by lepton-mixing matrix coefficients for the general case where there is mixing among more than just two species. Experimental results are presented as allowed regions on a plot, the axes of which are $|\Delta m^2|$ and $\sin^2 2\theta$. These are often summarized in terms of the upper limit on Δm^2 (the absolute value is usually suppressed in the notation) for maximal mixing, $\sin^2 2\theta = 1$, and the upper limit on $\sin^2 2\theta$ for “large” Δm^2 , *i.e.*, sufficiently large $|\Delta m^2|$ that the detector averages over many cycles of oscillation (or there ceases to be any coherence). A more complete discussion is given in the “Note on Neutrino Oscillation Experiments” just before the tables reporting such results.

An important type of experiment is the search for neutrinoless double- β decay, which tests for total lepton number violation such as would result for Majorana-neutrino masses. This process takes place when a nucleus with Z protons and $A = ZN$ nucleons decays according to $(Z, A) \rightarrow (Z+2, A)e^-e^-$, violating total lepton number by two units. In the case of neutrinos with masses which are sufficiently light, an upper limit on neutrinoless double- β decay yields a correlated upper limit on the quantity

$$\bar{m} = \left| \sum_j U_{1j}^2 m_{\nu_j} \right|. \quad (7)$$

Cancellations may occur in the sum, since U_{1j} is, in general, complex. See Ref. 11 for some recent reviews of searches for neutrinoless double- β decay.

A brief summary of the current experimental situation follows (see previous editions for discussions of various positive claims for neutrino masses and mixing, and their refutations).

1. There is no evidence at present from direct searches for nonzero neutrino masses. These include the endpoint of the Kurie plot in nuclear- β decay for m_{ν_e} , $\pi^+ \rightarrow \mu^+ \nu_\mu$ for m_{ν_μ} , and certain τ decays for m_{ν_τ} (where, as discussed above, the limits actually apply to the respective mass eigenstates ν_1 , ν_2 , and ν_3 in these three weak eigenstates). There has been some concern over the fact that the quantity which is actually measured, *i.e.*, the square of the neutrino mass, is negative by several standard deviations for ν_e . This may indicate systematic errors in the measurements which are not understood. A similar situation was noticed recently for the muon neutrino mass squared. Although there is still no resolution of this in the refereed journal literature, it has been argued [12,13] that the reason for this is that the charged-pion mass has been determined incorrectly from pionic x-rays, that this determination is actually ambiguous(!), and that when the analysis is interpreted differently, the charged pion mass is increased so as to yield a measurement of $m_{\nu_\mu}^2$ ($m_{\nu_2}^2$) which is no longer tachyonic but is consistent

with zero. See the recent preprint [13] and the more extensive discussion before the m_{ν_μ} and m_{π^\pm} Listings.

2. There are no indications of any positive neutrino masses from any of the peak search experiments in π or K decay, or from any experiments on neutrino decays. These set good upper limits on respective lepton mixing matrix coefficients.
3. There are no indications of any positive neutrino masses from nuclear- β decay spectra. The 7-year controversy over the claim by Simpson, Hime, and others of a 17 keV neutrino is finally over, with retractions by these authors of their original claims after very strong refutations by a number of high-sensitivity experiments.
4. Certain positive claims for neutrino oscillations in reactor and accelerator neutrino experiments have either been refuted or retracted, or both (see previous editions for details).
5. There is no indication of Majorana neutrino masses from searches for neutrinoless double- β decay.
6. The situation concerning possible atmospheric neutrino oscillations remains unsettled.

It is generally acknowledged that the strongest indirect evidence for neutrino masses and mixing is the observed deficit in the solar neutrino flux. The pioneering ^{37}Cl radiochemical solar neutrino experiment of R. Davis and his group started in 1967 and, with only one 18-month gap, has been running ever since. Over the years, a deficit in the measure flux of high-energy neutrinos (mainly from the ^8B reaction) has been reported [14,15]. The measured flux is about three times less than the flux predicted by theoretical calculations (Refs. 16–18, and other reviews quoted therein; see also Ref. 10). More recently, Kamiokande, a large water Čerenkov detector, has produced independent high-quality measurements of the solar neutrino flux in roughly the same energy range [19,20,21]. Unlike a radiochemical experiment, a Čerenkov detector preserves directional information, and the Kamiokande collaboration has been able to demonstrate that the neutrinos come from the direction of the sun. They have confirmed the deficiency reported by Davis *et al.* The ratio of observed to expected fluxes seen by the Kamiokande group is somewhat higher than that observed by Davis, but is still significantly below unity.

Two experiments are now operating which are sensitive to the lower-energy neutrinos from the main p - p burning chain. These are SAGE, the Soviet-American Gallium Experiment [22], and GALLEX, a gallium experiment in the Gran Sasso Underground Laboratory [23,24]. Both of these also report a deficiency in the measured solar neutrino flux. These results are especially important since the flux of the pp neutrinos constitutes the dominant part of the solar neutrino flux and is widely considered to be reliably calculable. (This is not to imply that calculations of the high-energy neutrino flux are unreliable, only that they are harder to check. The high-energy part does not arise from the reactions producing most of the sun’s luminosity,

but from minor side-chains which contribute only a very small fraction of the entire solar neutrino flux.)

One explanation of the results of these four experiments is that neutrino oscillations do take place during the transit from the production point in the sun to the interaction point in the earth. These oscillations may involve mixing either with ν_μ and ν_τ or “sterile” neutrino components. An appealing scenario is that of resonant neutrino oscillations, the Mikheyev-Smirnov-Wolfenstein (MSW) effect [25]. Details about this can be found in the reviews cited at the end of this note, and in the references in the original papers.

For some recent reviews on neutrino physics and further references to the original literature, see Refs. [15,26–29].

References

- Z. Maki, M. Nakagawa, and S. Sakata, *Prog. Theor. Phys.* **28**, 870 (1962); M. Nakagawa, H. Okonogi, S. Sakata, and A. Toyoda, *ibid.* **30** 727 (1963).
- B. Pontecorvo, *Sov. Phys. JETP* **6**, 429 (1957); **7**, 172 (1958); **26** 984 (1968) (*Zh. Eksp. Teor. Fiz.* **53**, 1717 (1967); V. Gribov and B. Pontecorvo, *Phys. Lett.* **28B**, 493 (1969); S.M. Bilenky and B. Pontecorvo, *Phys. Rev.* **41C**, 225 (1978).
- S. Weinberg, in *Proceedings of the XXVI International Conference on High Energy Physics*, (1992), ed. J. Sanford, (A.I.P., New York, 1993), p. 346.
- M. Gell-Mann, R. Slansky, and P. Ramond, in *Supergravity*, (North Holland, 1979), p. 315; T. Yanagida, in *Proceedings of the Workshop on Unified Theory and Baryon Number in the Universe*, (KEK, Japan, 1979).
- B.W. Lee and R.E. Shrock, *Phys. Rev.* **D16**, 1444 (1977).
- R.E. Shrock, *Phys. Lett.* **96B**, 159 (1980); “Note on Neutrino Mass Limits,” *Review of Particle Properties*, 1980 Edition, *Rev. Mod. Phys.* **52**, S63 (1980).
- R.E. Shrock, *Phys. Rev.* **D24**, 1232,1275 (1981); *Phys. Lett.* **112B**, 382 (1982).
- See, e.g., T.P. Cheng and L.F. Li, *Phys. Rev.* **D22**, 2860 (1980).
- K. Fujikawa and R.E. Shrock, *Phys. Rev. Lett.* **45**, 963 (1980).
- R.E. Shrock, *Nucl. Phys.* **B206**, 359 (1982).
- M. Doi, T. Kotani, and E. Takasugi, *Prog. Theor. Phys. Supp.* **83**, 1 (1985); T. Tomoda, *Rept. on Prog. in Phys.* **54**, 53 (1991).
- M. Janousch, R. Frosch, *et al.*, in *Proceedings of the International Workshop on Low Energy Muon Science (LEMS '93, April 4–8, 1993, Santa Fe, New Mexico)*, ed. M. Leon; R. Frosch, private communication.
- B. Jeckelmann, P.F.A. Goudsmit, and H.J. Leisi, to be published.
- R. Davis *et al.*, in the *Proceedings of the Conference on the Intersections between Particle and Nuclear Physics*, Steamboat Springs (1984), ed. R.E. Mischke, published in *Am. Inst. Phys.* **123**, 1037 (1984).
- R. Davis, Jr., A.K. Mann, and L. Wolfenstein, *Ann. Rev. Nucl. and Part. Sci.* **39**, 467 (1989).
- J.N. Bahcall, *Neutrino Astrophysics* (Cambridge, 1989).
- J.N. Bahcall and M.H. Pinsonneault, *Rev. Mod. Phys.* **64**, 885 (1992).
- S. Turck-Chièze *et al.*, *Phys. Reports* **230**, 57 (1993); S. Turck-Chièze and I. Lopes, *Ap. J.* **408**, 347 (1993).
- K.S. Hirata *et al.*, *Phys. Rev. Lett.* **63**, 16 (1989).
- K.S. Hirata *et al.*, *Phys. Rev. Lett.* **65**, 1297 (1990).
- K.S. Hirata *et al.*, *Phys. Rev. Lett.* **66**, 9 (1991).
- A.I. Abazov *et al.*, *Phys. Rev. Lett.* **67**, 3332 (1991).
- P. Anselmann *et al.*, *Phys. Lett.* **B314**, 445 (1993).
- P. Anselmann *et al.*, *Phys. Lett.* **B** (to be published, 1994).
- S.P. Mikheyev and A.Yu. Smirnov, *Sov. J. Nucl. Phys.* **42**, 913 (1985); [*Yad. Fiz.* **42**, 1441 (1985)]; L. Wolfenstein, *Phys. Rev.* **D17**, 2369 (1979).
- F. Boehm and P. Vogel, *Ann. Rev. Nucl. and Part. Sci.* **34**, 125 (1984); F. Boehm and P. Vogel, *Physics of Massive Neutrinos* (Cambridge Univ. Press, Cambridge, 1987).
- J.D. Vergados, *Phys. Reports* **133**, 1 (1986).
- S.M. Bilenky and S.T. Petcov, *Rev. Mod. Phys.* **59**, 671 (1987).
- H.V. Klapdor, ed., *Neutrinos* (Springer-Verlag, Berlin, New York, 1988).

Some other sources of information on neutrinos include topical neutrino conferences, and international high-energy physics and lepton-photon conferences.

ν_e

$$J = \frac{1}{2}$$

Not in general a mass eigenstate. See note on neutrino properties above.

These limits apply to ν_1 , the primary mass eigenstate in ν_e . They would also apply to any other ν_j which mixes strongly in ν_e and has sufficiently small mass that it can occur in the respective decay. The neutrino mass may be of a Dirac or Majorana type; the former conserves total lepton number while the latter violates it. Either would violate lepton family number, since nothing forces the neutrino mass eigenstates to coincide with the neutrino interaction eigenstates. For limits on a Majorana ν_e mass, see the section on “Searches for Massive Neutrinos and Lepton Mixing,” part (C), entitled “Searches for Neutrinoless Double- β Decay.”

From the analysis of neutrino events from SN 1987A it is important to get upper bounds on neutrino masses. For two examples of such studies, see Refs. 1 and 2 and references therein.

Our mass limit for ν_e is taken from the average in the ν_e “Mass Squared” section immediately below this section.

References

- D.N. Spergel and J.N. Bahcall, *Phys. Lett.* **B200**, 366 (1988).
- L.F. Abbott *et al.*, *Nucl. Phys.* **B299**, 734 (1988).

Lepton & Quark Full Listings

 ν_e ν_e MASS

Most of the data from which these limits are derived are from β^- decay experiments in which a $\bar{\nu}_e$ is produced, so that they really apply to $m_{\bar{\nu}_1}$. Assuming *CPT* invariance, a limit on $m_{\bar{\nu}_1}$ is the same as a limit on m_{ν_1} . Results from studies of electron capture transitions, given below " $m_{\nu_1} - m_{\bar{\nu}_1}$ ", give limits on m_{ν_1} itself. See also the Listings in the Neutrino Bounds from Astrophysics and Cosmology section.

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
OUR LIMIT		1 PDG	94 RVUE	See the note below.
< 7.2	95	2 WEINHEIMER 93	SPEC	$^3\text{H}\beta$ decay
< 11.7	95	3 HOLZSCHUH 92B	SPEC	$^3\text{H}\beta$ decay
< 13.1	95	4 KAWAKAMI 91	SPEC	$^3\text{H}\beta$ decay
< 9.3	95	5 ROBERTSON 91	SPEC	$^3\text{H}\beta$ decay
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	95	6 VANDYCK 93		$m_{^3\text{H}} - m_{^3\text{He}}$
< 14	95	7 DECMAN 92	CNTR	$^3\text{H}\beta$ decay
< 23		AVIGNONE 90	ASTR	Supernova SN 1987A
		LOREDO 89	ASTR	SN 1987A
		ABBOTT 88	ASTR	Supernova SN 1987A
< 29	95	8 KAWAKAMI 88	SPEC	Repl. by KAWAKAMI 91
		9 SPERGEL 88	ASTR	Supernova SN 1987A
17 to 40		10 BORIS 87	SPEC	$\bar{\nu}_e$, $^3\text{H}\beta$ decay
< 27	95	11 WILKERSON 87	SPEC	$\bar{\nu}_e$, $^3\text{H}\beta$ decay
< 18	95	12 FRITSCHI 86	SPEC	Repl. by HOLZSCHUH 92B

- 1 PDG 94 formal upper limit, as obtained from the m^2 average in the next section, is 5.1 eV at the 95%CL. Caution is urged in interpreting this result, since the m^2 average is positive with only a 3.5% probability. If the weighted average m^2 were forced to zero, the limit would increase to 7.0 eV.
- 2 WEINHEIMER 93 is a measurement of the endpoint of the tritium β spectrum using an electrostatic spectrometer with a magnetic guiding field. The source is molecular tritium frozen onto an aluminum substrate.
- 3 HOLZSCHUH 92B result is obtained from the measurement $m_{\nu_e}^2 = -24 \pm 48 \pm 61$ (1 σ errors), in eV^2 , using the PDG prescription for conversion to a limit in m_{ν_e} .
- 4 KAWAKAMI 91 experiment uses tritium-labeled arachidic acid. This result may be obtained from the $m_{\nu_1}^2$ limit by combining the errors in quadrature and using the method described in the Probability, Statistics, and Monte Carlo section in Chapter III of this Review. This was also done in ROBERTSON 91, although the authors report a different procedure.
- 5 ROBERTSON 91 experiment uses gaseous molecular tritium. The result is in strong disagreement with the earlier claims by the ITEP group [LUBIMOV 80, BORIS 87 (+ BORIS 88 erratum)] that m_{ν_1} lies between 17 and 40 eV. However, the probability of a positive m^2 is only 3% if statistical and systematic error are combined in quadrature.
- 6 VANDYCK 93 is a new measurement of the atomic masses of ^3H and ^3He . The authors note that "The excellent agreement with recent results from beta spectrometers lends strong support for new (upper) limits on the (electron) neutrino's rest mass."
- 7 DECMAN 92 was not published in a refereed journal, so we do not use it in our compilation, but we include it for the reader's convenience. If it were included our limit would be reduced to 4.9 eV, but the average m^2 would be positive with only a 1.1% probability.
- 8 KAWAKAMI 88 multiply their statistical error by the appropriate factor for 95% CL when $m^2 > 0$ is required (1.74), add this linearly to their unmultiplied systematic error (173 eV^2) and add the m^2 value (223 eV^2) to obtain their 95% CL limit ($m < 29$ eV). To adjust for our quadratic addition of errors and our multiplication of both the statistical and systematic errors by the factor 1.645 we set the systematic error to 269 eV^2 to yield the same limit.
- 9 SPERGEL 88 rule out masses greater than 16 eV.
- 10 See also comment in BORIS 87B and erratum in BORIS 88.
- 11 WILKERSON 87 multiply both statistical and systematic errors by 1.645 (for 95% CL), add them in quadrature and add the (negative) m^2 value (-57 eV^2) to obtain their 95% CL limit ($m < 27$ eV).
- 12 FRITSCHI 86 multiply their statistical error by 1.645 (for 95% CL), add this linearly to their unmultiplied systematic error (204 eV^2) and do NOT add in the m^2 value (-11 eV^2) to obtain their 95% CL limit ($m < 18$ eV). To adjust for our quadratic addition of errors, and our multiplication of both the statistical and systematic errors by the factor 1.645, we set the systematic error to 178 eV^2 .

 ν_e MASS SQUARED

The tritium experiments actually measure mass squared. A combined limit on mass must therefore be obtained from the weighted average of the results shown here. The recent results are in strong disagreement with the earlier claims by the ITEP group [LUBIMOV 80, BORIS 87 (+ BORIS 88, erratum)] that m_{ν_1} lies between 17 and 40 eV. The BORIS 87 result is excluded because of the controversy over the possibly large unreported systematic errors; see BERGVIST 85B, BERGVIST 86, SIMPSON 84, and REDONDO 89. However, the average for the new experiments given below implies only a 3.5% probability that m^2 is positive. See HOLZSCHUH 92 for a review of the recent direct m_{ν_1} measurements.

VALUE (eV^2)	DOCUMENT ID	TECN	COMMENT
- 54 ± 30 OUR AVERAGE			
- 39 ± 34 ± 15	13 WEINHEIMER 93	SPEC	$^3\text{H}\beta$ decay
- 24 ± 48 ± 61	14 HOLZSCHUH 92B	SPEC	$^3\text{H}\beta$ decay
- 65 ± 85 ± 65	15 KAWAKAMI 91	SPEC	$\bar{\nu}_e$, tritium
- 147 ± 68 ± 41	16 ROBERTSON 91	SPEC	$\bar{\nu}_e$, tritium
• • • We do not use the following data for averages, fits, limits, etc. • • •			
- 72 ± 41 ± 30	17 DECMAN 92	SPEC	$^3\text{H}\beta$ decay
223 ± 244 ± 269	18 KAWAKAMI 88	SPEC	Repl. by KAWAKAMI 91
- 57 ± 453 ± 118	19 WILKERSON 87	SPEC	Repl. by ROBERTSON 91
- 11 ± 63 ± 178	20 FRITSCHI 86	SPEC	Repl. by HOLZSCHUH 92B

- 13 WEINHEIMER 93 is a measurement of the endpoint of the tritium β spectrum using an electrostatic spectrometer with a magnetic guiding field. The source is molecular tritium frozen onto an aluminum substrate.
- 14 HOLZSCHUH 92B source is a monolayer of tritiated hydrocarbon.
- 15 KAWAKAMI 91 experiment uses tritium-labeled arachidic acid.
- 16 ROBERTSON 91 experiment uses gaseous molecular tritium. The result is in strong disagreement with the earlier claims by the ITEP group [LUBIMOV 80, BORIS 87 (+ BORIS 88 erratum)] that m_{ν_1} lies between 17 and 40 eV. However, the probability of a positive m^2 is only 3% if statistical and systematic error are combined in quadrature.
- 17 DECMAN 92 was not published in a refereed journal, so we do not use it in our compilation, but we include it for the reader's convenience. If it were included, our average would be $-58 \pm 26 \text{ eV}^2$.
- 18 KAWAKAMI 88 multiply their statistical error by the appropriate factor for 95% CL when $m^2 > 0$ is required (1.74), add this linearly to their unmultiplied systematic error (173 eV^2) and add the m^2 value (223 eV^2) to obtain their 95% CL limit ($m < 29$ eV). To adjust for our quadratic addition of errors and our multiplication of both the statistical and systematic errors by the factor 1.645 we set the systematic error to 269 eV^2 to yield the same limit.
- 19 WILKERSON 87 multiply both statistical and systematic errors by 1.645 (for 95% CL), add them in quadrature and add the (negative) m^2 value (-57 eV^2) to obtain their 95% CL limit ($m < 27$ eV).
- 20 FRITSCHI 86 multiply their statistical error by 1.645 (for 95% CL), add this linearly to their unmultiplied systematic error (204 eV^2) and do NOT add in the m^2 value (-11 eV^2) to obtain their 95% CL limit ($m < 18$ eV). To adjust for our quadratic addition of errors, and our multiplication of both the statistical and systematic errors by the factor 1.645, we set the systematic error to 178 eV^2 .

 $m_{\nu_1} - m_{\bar{\nu}_1}$

These are measurement of m_{ν_1} (in contrast to $m_{\bar{\nu}_1}$, given above). The masses can be different for a Dirac neutrino in the absence of *CPT* invariance. The test is not very strong.

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 225	95	SPRINGER 87	CNTR	ν , ^{163}Ho
< 550	68	YASUMI 86	CNTR	ν , ^{163}Ho
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1250		21 YASUMI 83	CNTR	ν , ^{163}Ho
< 1300		ANDERSEN 82	CNTR	ν , ^{163}Ho
< 4.5×10^5	90	CLARK 74	ASPK	K_{e3} decay
< 4100	67	BECK 68	CNTR	ν , ^{22}Na

- 21 Assumes upper limit on *Q*-value reported by ANDERSEN 82. Replaced by YASUMI 86.

 ν_1 CHARGE

VALUE (units: electron charge)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 2×10^{-15}	22 BARBIELLINI 87	ASTR	Supernova SN 1987A
< 1×10^{-13}	BERNSTEIN 63	ASTR	Solar energy losses

- 22 Precise limit depends on assumptions about the intergalactic or galactic magnetic fields and about the direct distance and time through the field.

 ν_1 MEAN LIFE

VALUE (s)	CL%	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
> 278	90	23 LOSECCO 87B	IMB

- 23 LOSECCO 87B assumes observed rate of 2.1 SNU (solar neutrino units) comes from sun while 7.0 ± 3.0 is theory.

ν_1 (MEAN LIFE) / MASS

VALUE (s/eV)	CL%	DOCUMENT ID	TECN	COMMENT
>300	90	24 REINES	74 CNTR	$\bar{\nu}$
••• We do not use the following data for averages, fits, limits, etc. •••				
> 2.8 × 10 ¹⁵		25,26 BLUDMAN	92 ASTR	$m_\nu < 50$ eV
> 6.4	90	27 KRAKAUER	91 CNTR	$\bar{\nu}$ at LAMPF
> 6.3 × 10 ¹⁵		26,28 CHUPP	89 ASTR	$m_\nu < 20$ eV
		29 COWSIK	89 ASTR	
> 1.7 × 10 ¹⁵		26 KOLB	89 ASTR	$m_\nu < 20$ eV
		30 RAFFELT	89 RVUE	$\bar{\nu}$ (Dirac, Majorana)
		31 RAFFELT	89B ASTR	
		32 BOUCHEZ	88 CNTR	$\bar{\nu}$ (Dirac, Majorana)
		33 FRIEMAN	88 ASTR	
		33 VONFEILIT...	88 ASTR	
> 8.3 × 10 ¹⁴		34 OBERAUER	87	$\bar{\nu}_R$ (Dirac)
> 22	68	34 OBERAUER	87	$\bar{\nu}$ (Majorana)
> 38	68	34 OBERAUER	87	$\bar{\nu}_L$ (Dirac)
> 59	68	34 OBERAUER	87	$\bar{\nu}_L$ (Dirac)
> 30	68	KETOV	86 CNTR	$\bar{\nu}$ (Dirac)
> 20	68	KETOV	86 CNTR	$\bar{\nu}$ (Majorana)
> 7 × 10 ⁹		35 RAFFELT	85 ASTR	
		36 HENRY	81 ASTR	$m_\nu = 16-20$ eV
		37 KIMBLE	81 ASTR	$m_\nu = 10-100$ eV
> 2 × 10 ²¹		38 STECKER	80 ASTR	$m_\nu = 10-100$ eV
<p>²⁴ REINES 74 looked for ν_e of nonzero mass decaying to a neutral of lesser mass + γ. Used liquid scintillator detector near fission reactor. Finds lab lifetime $6. \times 10^7$ s or more. Above value of (mean life)/mass assumes average effective neutrino energy of 0.2 MeV. To obtain the limit $6. \times 10^7$ s REINES 74 assumed that the full $\bar{\nu}_e$ reactor flux could be responsible for yielding decays with photon energies in the interval 0.1 MeV - 0.5 MeV. This represents some overestimate so their lower limit is an over-estimate of the lab lifetime (P. Vogel, private communication, 1984).</p> <p>²⁵ BLUDMAN 92 sets additional limits by this method for higher mass ranges. Cosmological limits are also obtained.</p> <p>²⁶ Nonobservation of γ's in coincidence with ν's from SN 1987A.</p> <p>²⁷ KRAKAUER 91 quotes the limit $\tau/m_{\nu_1} > (0.3a^2 + 9.8a + 15.9) \text{ s/eV}$, where a is a parameter describing the asymmetry in the neutrino decay defined as $dN_\gamma/d\cos\theta = (1/2)(1 + a \cos\theta)$ $a = 0$ for a Majorana neutrino, but can vary from -1 to 1 for a Dirac neutrino. The bound given by the authors is the most conservative (which applies for $a = -1$).</p> <p>²⁸ CHUPP 89 should be multiplied by a branching ratio (about 1) and a detection efficiency (about 1/4), and pertains to radiative decay of any neutrino to a lighter or sterile neutrino.</p> <p>²⁹ COWSIK 89 use observations of supernova SN 1987A to set the limit for the lifetime of a neutrino with $1 < m < 50$ MeV decaying through $\nu_H \rightarrow \nu_1 ee$ to be $\tau > 4 \times 10^{15} \exp(-m/5 \text{ MeV})$ s.</p> <p>³⁰ RAFFELT 89 uses KYULDJIEV 84 to obtain $\tau m^3 > 3 \times 10^{18} \text{ s eV}^3$ (based on $\bar{\nu}_e e^-$ cross sections). The bound is not valid if electric and magnetic transition moments are equal for Dirac neutrinos.</p> <p>³¹ RAFFELT 89B analyze stellar evolution and exclude the region $3 \times 10^{12} < \tau m^3 < 3 \times 10^{21} \text{ s eV}^3$.</p> <p>³² BOUCHEZ 88 reports limits in the nearly degenerate mass case.</p> <p>³³ Model-dependent theoretical analysis of SN 1987A neutrinos.</p> <p>³⁴ OBERAUER 87 bounds are from comparison of observed and expected rate of reactor neutrinos.</p> <p>³⁵ RAFFELT 85 limit is from solar x^- and γ-ray fluxes.</p> <p>³⁶ HENRY 81 uses UV flux from clusters of galaxies to find $\tau > 1.1 \times 10^{25}$ s for radiative decay.</p> <p>³⁷ KIMBLE 81 uses extreme UV flux limits to find $\tau > 10^{22}-10^{23}$ s.</p> <p>³⁸ STECKER 80 limit based on UV background; result given is $\tau > 4 \times 10^{22}$ s at $m_\nu = 20$ eV.</p>				

$|(\mathbf{v} - \mathbf{c}) / \mathbf{c}|$ ($\mathbf{v} \equiv \nu_1$ VELOCITY)

Expected to be zero for massless neutrino, but tests also whether photons and neutrinos have the same limiting velocity in vacuum.

VALUE (units 10 ⁻⁸)	EVTS	DOCUMENT ID	TECN	COMMENT
<1	17	39 STODOLSKY	88 ASTR	SN 1987A
<p>³⁹ STODOLSKY 88 result based on <10 hr between $\bar{\nu}_e$ detection in IMB and KAMI detectors and beginning of light signal. Inclusion of the problematic 5 neutrino events from FREJ (four hours later) does not change the result.</p>				

ν_1 MAGNETIC MOMENT

Must vanish for Majorana neutrino or purely chiral massless Dirac neutrino. The value of the magnetic moment for the standard SU(2)×U(1) electroweak theory extended to include massive neutrinos (see FUJIKAWA 80) is $\mu_\nu = 3eG_F m_\nu / (8\pi^2 \sqrt{2}) = (3.20 \times 10^{-19}) m_\nu \mu_B$ where m_ν is in eV and $\mu_B = e\hbar/2m_e$ is the Bohr magneton. Given the upper bound $m_{\nu_1} < 7.3$ eV, it follows that for the extended standard electroweak theory, $\mu(\nu_1) < 2.3 \times 10^{-18} \mu_B$. Current experiments are not yet challenging this limit. There is considerable controversy over the validity of many of the claimed upper limits on the magnetic moment from the astrophysical data. For example, VOLOSHIN 90 states that "in connection with the astrophysical limits on μ_ν , ... there is by now a general consensus that contrary to the initial claims (BARBIERI 88, LATTIMER 88, GOLDMAN 88, NOTZOLD 88), essentially no better than quoted limits (from previous constraints) can be derived from detection of the neutrino flux from the supernova SN1987A." See VOLOSHIN 88 and VOLOSHIN 88C.

VALUE (μ_B)	CL%	DOCUMENT ID	TECN	COMMENT
<1.08 × 10 ⁻⁹		40 KRAKAUER	90 CNTR	LAMPF $\nu_e e \rightarrow \nu_e e$
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.9 × 10 ⁻¹⁰	95	DERBIN	93 CNTR	Reactor $\bar{\nu}_e e \rightarrow \bar{\nu}_e e$
<7.7 × 10 ⁻¹⁰	95	MOURAO	92 ASTR	HOME/KAM2 ν rates
<2.4 × 10 ⁻¹⁰		41 VIDYAKIN	92 CNTR	Reactor $\bar{\nu}_e e \rightarrow \bar{\nu}_e e$
		42 FIORENTINI	91 ASTR	
<2 × 10 ⁻¹²	95	43 RAFFELT	90 ASTR	Red giant luminosity
<1 × 10 ⁻¹¹		44 RAFFELT	89B ASTR	Cooling helium stars
<(2-8) × 10 ⁻¹²		44,45,46 BARBIERI	88 ASTR	Supernova SN 1987A
		47 FUKUGITA	88 COSM	Primordial magn. fields
<1 × 10 ⁻¹²	45,46,48	GOLDMAN	88 ASTR	Supernova SN 1987A
<5 × 10 ⁻¹³	44,46	LATTIMER	88 ASTR	Supernova SN 1987A
≤ 1.5 × 10 ⁻¹²	44,46	NOETZOLD	88 ASTR	Supernova SN 1987A
≤ 3 × 10 ⁻¹¹	44	RAFFELT	88B ASTR	He burning stars
<1.1 × 10 ⁻¹¹	44	FUKUGITA	87 ASTR	Cooling helium stars
<4 × 10 ⁻¹¹		LYNN	81 ASTR	
<1-2 × 10 ⁻¹¹		MORGAN	81 COSM	⁴ He abundance
<8.5 × 10 ⁻¹¹		BEG	78 ASTR	Stellar plasmons
<6 × 10 ⁻¹¹		49 SUTHERLAND	76 ASTR	Red giants + deg. dwarfs
<1 × 10 ⁻¹⁰		BERNSTEIN	63 ASTR	Cooling white dwarfs
<1.4 × 10 ⁻⁹		COWAN	57 CNTR	Reactor $\bar{\nu}_e$
<p>⁴⁰ KRAKAUER 90 experiment fully reported in ALLEN 93.</p> <p>⁴¹ VIDYAKIN 92 limit is from a $(enu)_e$ elastic scattering experiment. No experimental details are given except for the cross section from which this limit is derived. Signal/noise was 1/10. The limit uses $\sin^2\theta_W = 0.23$ as input.</p> <p>⁴² FIORENTINI 91 is a study of the statistical significance of possible correlation of solar neutrino flux with sunspot cycle. Data do not imply any evidence for a nonzero neutrino magnetic moment, although they are consistent with a moment of order $1 \times 10^{-10} / \mu_B$.</p> <p>⁴³ RAFFELT 90 limit applies for a diagonal magnetic moment of a Dirac neutrino, or for a transition magnetic moment of a Majorana neutrino. In the latter case, the same analysis gives $< 1.4 \times 10^{-12}$. Limit at 95%CL obtained from δM_c.</p> <p>⁴⁴ Significant dependence on details of stellar models.</p> <p>⁴⁵ A limit of 10^{-13} is obtained with even more model-dependence.</p> <p>⁴⁶ These papers have assumed that the right-handed neutrino is inert; see BARBIERI 88b.</p> <p>⁴⁷ FUKUGITA 88 find magnetic dipole moments of any two neutrino species are bounded by $\mu < 10^{-16} [10^{-9} G/B_0]$ where B_0 is the present-day intergalactic field strength.</p> <p>⁴⁸ Some dependence on details of stellar models.</p> <p>⁴⁹ We obtain above limit from SUTHERLAND 76 using their limit $f < 1/3$.</p>				

NONSTANDARD CONTRIBUTIONS TO NEUTRINO SCATTERING

We report limits on the so-called neutrino charge radius squared in this section. This quantity is not an observable, physical quantity and this is reflected in the fact that it is gauge dependent (see LEE 77C). It is not necessarily positive. A more general interpretation of the experimental results is that they are limits on certain nonstandard contributions to neutrino scattering.

VALUE (10 ⁻³² cm ²)	CL%	DOCUMENT ID	TECN	COMMENT
0.9±2.7		ALLEN	93 CNTR	LAMPF $\nu_e e \rightarrow \nu_e e$
••• We do not use the following data for averages, fits, limits, etc. •••				
<2.3 × 10 ⁻³²	95	MOURAO	92 ASTR	HOME/KAM2 ν rates
<7.3		50 VIDYAKIN	92 CNTR	Reactor $\bar{\nu}_e e \rightarrow \bar{\nu}_e e$
1.1±2.3		ALLEN	91 CNTR	Repl. by ALLEN 93
		51 GRIFOLS	89B ASTR	SN 1987A
<p>⁵⁰ VIDYAKIN 92 limit is from a $e\bar{\nu}_e$ elastic scattering experiment. No experimental details are given except for the cross section from which this limit is derived. Signal/noise was 1/10. The limit uses $\sin^2\theta_W = 0.23$ as input.</p> <p>⁵¹ GRIFOLS 89B sets a limit of $(r^2) < 0.2 \times 10^{-32} \text{ cm}^2$ for right-handed neutrinos.</p>				

Lepton & Quark Full Listings

 ν_e, ν_μ ν_e REFERENCES

PDG	94	PR D50	Montanet+	(CERN, LBL, BOST, IFIC+)
ALLEN	93	PR D47 11	+Chen, Doe, Hausammann+	(UCI, LANL, ANL, UMD)
DERBIN	93	JETPL 57 768	+Chernyi, Popeko, Muratova+	(PNPI)
		Translated from ZETFP	57 755.	
VANDYCK	93	PRL 70 2888	Van Dyck, Farnham, Schwinberg	(WASH)
WEINHEIMER	93	PL B300 210	+Przyrembel, Backe+	(MANZ)
BLUDMAN	92	PR D45 4720		(CFPA)
DECMAN	92	BAPS 37 1286	+Stoeffl	(LLNL)
HOLZSCHUH	92	RPP 55 1035		(ZURI)
HOLZSCHUH	92	PL B287 381	+Fritsch, Kuendig	(ZURI)
MOURAO	92	PL B285 364	+Pulido, Ralston	(LIBS, LISBT, CERN, KANS)
VIDYAKIN	92	JETPL 55 206	+Vyrodov, Gurevich, Koslov+	(KIAE)
		Translated from ZETFP	55 212.	
ALLEN	91	PR D43 R1	+Chen, Doe, Hausammann	(UCI, LANL, UMD)
FIorentINI	91	PL B253 181	+Mezzorani	(CAGL, INFN)
KAWAKAMI	91	PL B256 105	+Kato, Ohshima+	(INUS, TOHOK, TINT, KOBE, KEK)
KRAKAUER	91	PR D44 R6	+Talaga, Allen, Chen+	(LAMPF E225 Collab.)
ROBERTSON	91	PRL 67 957	+Bowles, Stephenson, Wark, Wilkerson, Knapp	(LASL, LLL)
AVIGNONE	90	PR D41 682	+Collar	(SCUC)
KRAKAUER	90	PL B252 177	+Talaga, Allen, Chen+	(LAMPF E225 Collab.)
RAFFELT	90	PRL 64 2856		(MPIM)
VOLOSHIN	90	NP B (Proc. Suppl) 19 433		(ITEP)
		Neutrino 90 Conference		
CHUPP	89	PRL 62 505	+Vestrand, Reppin	(UNH, MPIM)
COWSIK	89	PL B218 91	+Schramm, Hoflich	(WUSL, TATA, CHIC, MPIM)
GRIFOLS	89	PR D40 3819	+Masso	(BARC)
KOLB	89	PRL 62 509	+Turner	(CHIC, FNAL)
LOREDO	89	ANYAS 571 601	+Lamb	(CHIC)
RAFFELT	89	PR D39 2066		(PRIN, UCB)
RAFFELT	89	APJ 336 61	+Dearborn, Silk	(UCB, LLL)
REDONDO	89	PR C40 368	+Robertson	(LANL)
ASBOTT	88	NP D29 734	+De Rujula, Waiker	(BRAN, CERN, USC)
BARBIERI	88	PRL 61 27	+Mohapatra	(PISA, UMD)
BARBIERI	88	PL B213 69	+Mohapatra, Yanagida	(PISA, UMD, MICH)
BORIS	88	PRL 61 245 erratum	+Golutin, Laptin+	(ITEP, ASCI)
BOUCHEZ	88	PL B207 217	+Pichard, Soirat, Spiro, Declais	(SACL, MARS)
FRIEMAN	88	PL B200 115	+Haber, Freese	(SLAC, UCSC, UCSB)
FUKUGITA	88	PRL 60 879	+Notzold, Raffelt, Silk	(KYOTY, MPIM, UCB)
GOLDMAN	88	PL 60 1789	+Aharanov, Alexander, Nussinov	(TELA)
KAWAKAMI	88	JPL 57 2873	+Kato, Naito, Nisimura+	(INUS, TOKY, TINT, KEK)
LATTIMER	88	PRL 61 23	+Cooperstein	(STON, BNL)
		Also	Lattimer, Cooperstein	(STON, BNL)
NOETZOLD	88	PR D38 1658		(MPIM)
NOTZOLD	88	PR D38 1658		(MPIM)
RAFFELT	88	PR D37 549	+Dearborn	(UCB, LLL)
SPERGL	88	PL B200 366	+Bahcall	(IAS)
STODOLSKY	88	PL B201 353		(MPIM)
VOLOSHIN	88	PL B209 360		(ITEP)
		Also	Voloshin	(ITEP)
VOLOSHIN	88	JETPL 47 501		(ITEP)
VONFEILIT...	88	PL B200 580	Von Feilitzsch, Oberauer	(MUNT)
BARBIELLINI	87	Nature 329 21	+Cocconi	(CERN)
BORIS	87	PRL 58 2019	+Golutin, Laptin+	(ITEP, ASCI)
		Also	Boris, Golutin, Laptin+	(ITEP, ASCI)
BORIS	87	JETPL 45 333	+Golutin, Laptin+	(ITEP)
		Translated from ZETFP	45 267.	
FUKUGITA	87	PR D36 3817	+Yazaki	(KYOTY, TOKY)
LOSECCO	87	PR D35 2073	+Bionta, Blewitt, Bratton+	(IMB Collab.)
OBERAUER	87	PL B198 113	+von Feilitzsch, Mossbauer	(MUNT)
SPRINGER	87	PR A35 679	+Bennet, Balsden+	(LLNL)
WILKERSON	87	PRL 58 2023	+Bowles, Browne+	(LANL, PRIN, UCSD)
BERGKVIST	86	Moriond Conf., Vol. M48, 465		(STOH)
FRITSCHI	86	PL B173 485	+Holzschuh, Kundig+	(ZURI, SIN)
KETOV	86	JETPL 44 146	+Klimov, Nikolaev, Mikaelyan+	(KIAE)
		Translated from ZETFP	44 114.	
YASUMI	86	PL B181 169	+Ando+	(KEK, OSAK, TOHOK, TSUK, KYOT, INUS+)
BERGKVIST	85	PL 159B 408		(STOH)
RAFFELT	85	PR D31 3002		(MPIM)
KYULDJIEV	84	NP B243 387		(SOFI)
SIMPSON	84	PR D30 1110		(GUEL)
YASUMI	83	PL 122B 461	+Rajasekaran+	(KEK, OSAK, TINT, TOHOK, TSUK)
ANDERSEN	82	PL 113B 72	+Beyer, Charpak, Derujula+	(AARN, CERN, RISO)
HENRY	81	PRL 47 618	+Feldman	(JHU)
KIMBLE	81	PR 48 80	+Bowyer, Jakobsen	(UCB)
LYNN	81	PR D23 2151		(COLU)
MORGAN	81	PL 102B 247	Morgan	(SUSS)
FUJIKAWA	80	PRL 45 963	+Shrock	(STON)
LUBIMOV	80	PL 94B 266	+Novikov, Nozik, Tretyakov, Kosik	(ITEP)
		Also	Kozik, Lubimov, Novikov+	(ITEP)
		Translated from YAF 32	301.	
		Also	Lubimov, Novikov, Nozik+	(ITEP)
		Translated from ZETF 81	115B.	
STECKER	80	PRL 45 1460		(NASA)
BEG	78	PR D17 1395	+Marciano, Ruderman	(ROCK, COLU)
LEE	77	PR D16 1444	+Shrock	(STON)
SUTHERLAND	76	PR D13 2700	+Ng, Flowers+	(PENN, COLU, NYU)
CLARK	74	PR D9 533	+Elioff, Frisch, Johnson, Kerth, Shen+	(LBL)
REINES	74	PRL 32 180	+Sobel, Gurr	(UCI)
		Also	Barnes	(PURD)
BECK	68	ZPHY 216 229	+Daniel	(MPIH)
BERNSTEIN	63	PR 132 1227	+Ruderman, Feinberg	(NYU, COLU)
COWAN	57	PR 107 528	+Reines	(LANL)

 ν_μ

$$J = \frac{1}{2}$$

Not in general a mass eigenstate. See note on neutrinos in the ν_e section above.

NOTE ON THE MUON NEUTRINO MASS*

Particle-physics limits (as opposed to those from astrophysics) for the mass of the muon neutrino m_{ν_μ} come from measurements of the muon momentum p_μ in the decay $\pi^+ \rightarrow \mu^+ \nu_\mu$. For π^+ decay at rest,

$$m_{\nu_\mu}^2 = m_\pi^2 + m_\mu^2 - 2m_\pi \sqrt{m_\mu^2 + p_\mu^2}. \quad (1)$$

The DAUM 91 determination of p_μ for π^+ decay at rest,

$$p_\mu = 29.79177 \pm 0.00024 \text{ MeV}/c,$$

has been followed by an even more precise but not yet published measurement by the same group (ASSAMAGAN 94):

$$p_\mu = 29.79207 \pm 0.00012 \text{ MeV}/c$$

The 1986 CODATA value for the muon mass is $105.658389 \pm 0.000034 \text{ MeV}$. A present ambiguity concerning the pion mass is discussed in the π^\pm section of these Listings; the two values of m_π from a recent paper, JECKELMANN 94, are given in Table 1. The corresponding values of $m_{\nu_\mu}^2$ obtained using Eq. (1) are also given. Using the ASSAMAGAN 94 p_μ value, one finds for Solution A that $m_{\nu_\mu}^2$ is negative by 6.1 standard deviations, while for Solution B, $m_{\nu_\mu}^2$ is negative by 0.9 standard deviations.

Table 1: Values for $m_{\nu_\mu}^2$ obtained using the unpublished ASSAMAGAN 94 measurement of p_μ and the two JECKELMANN 94 solutions for m_π .

m_π (MeV)	$m_{\nu_\mu}^2$ (MeV ²)
Solution A 139.56782 ± 0.00037	-0.149 ± 0.025
Solution B 139.56995 ± 0.00035	-0.022 ± 0.023

Alternatively, one can calculate $m_{\nu_\mu}^2$ as a function of m_π . This function is shown in Fig. 1, along with the two π^- mass solutions of JECKELMANN 94.

Much of this analysis follows that of Robertson [1], who concluded (before Solution B was known to be viable) that we should revert to the ANDERHUB 82 limit, $m_{\nu_\mu} \leq 0.50 \text{ MeV}$ at the 90% CL. In that experiment, the momentum of forward-going muons from pion decay in flight was measured. The kinematics are such that the sensitivity of $m_{\nu_\mu}^2$ to m_π and m_μ decreases as the energy increases, so that the need for an independent precise measurement of m_{π^\pm} mass is bypassed. However, the result, $m_{\nu_\mu}^2 = -0.14 \pm 0.20 \text{ MeV}^2$, has an error large enough to include both of the solutions shown in Fig. 1. In principle, this kind of experiment might resolve the present ambiguity. In practice, experiments on pionic x-rays in lower- Z materials, which have just begun at the Paul Scherrer Institute, are likely to resolve the ambiguity [2].




Figure 1: $m_{\nu_\mu}^2$ as a function of π^\pm mass. The ASSAMAGAN 94 measurement of the momentum of the muon in π^+ decay at rest is represented by the diagonal line with the 1σ error band from uncertainties in p_μ and m_μ . The two solutions for the π^- mass from the JECKELMANN 94 reanalysis are shown by the vertical lines, along with their 1σ error bands. Also shown are the 1σ error ellipses from the combined fits.

Since $m_{\nu_\mu}^2$ must be nonnegative and to obtain the more conservative limit, we choose the new JECKELMANN 94 Solution B for the pion mass, and take the corresponding value $m_{\nu_\mu}^2 = 0.001 \pm 0.044$ MeV². Using a Bayesian procedure with a flat prior distribution, we find $m_{\nu_\mu}^2 < 0.073$ MeV² at the 90% CL, and so list $m_{\nu_\mu} < 0.27$ MeV at the 90% CL.

If, instead, we apply the same procedure using p_μ from the as yet unpublished ASSAMAGAN 94 and the Solution B pion mass from JECKELMANN 94, we obtain $m_{\nu_\mu}^2 = -0.022 \pm 0.023$ MeV², or $m_{\nu_\mu} < 0.17$ MeV at the 90% CL.

Notes and References

* This note was prepared with extensive help from F. Boehm, R. Frosch, P.F.A. Goudsmit, Y.K. Lee, H.J. Leisi, R.G.H. Robertson, and P. Vogel

1. R.G.H. Robertson, p. 140 in *Proceedings XXVI International Conference on High-Energy Physics*, ed. J.R. Sanford (Dallas, TX, 6–12 August, 1992).
2. R. Frosch, private communication (May 1994).

ν_μ MASS

Applies to ν_2 , the primary mass eigenstate in ν_μ . Would also apply to any other ν_j which mixes strongly in ν_μ and has sufficiently small mass that it can occur in the respective decays. (This would be nontrivial only for $j \geq 3$, given the ν_e mass limit above.) See also the Listings in the Neutrino Bounds from Astrophysics and Cosmology section.

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
<0.27 (CL = 90%) OUR EVALUATION DAUM 91 and JECKELMANN 94; see Note.				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.17		1 ASSAMAGAN 94	SPEC	$m^2 = -0.022 \pm 0.023$
<0.48		2 ENQVIST 93	COSM	Nucleosynthesis
<0.003		3,4 MAYLE 93	ASTR	SN 1987A cooling
< 0.025–0.030		4,5 BURROWS 92	ASTR	SN 1987A cooling
		MAMEDOV 92	RVUE	
<0.3		6 FULLER 91	COSM	Nucleosynthesis
<0.42		6 LAM 91	COSM	Nucleosynthesis
< 0.028–0.15		7 NATALE 91	ASTR	SN 1987A
<0.028		4 GANDHI 90	ASTR	SN 1987A
<0.014		4,8 GRIFOLS 90B	ASTR	SN 1987A
<0.06		4,9 GAEMERS 89		SN 1987A
<0.27	90	10 ABELA 84	SPEC	$m^2 = -0.097 \pm 0.072$
<0.50	90	ANDERHUB 82	SPEC	$m^2 = -0.14 \pm 0.20$
<0.52	90	11 LU 80	CNTR	$m^2 = 0.102 \pm 0.119$
<0.65	90	CLARK 74	ASPK	$K_{\mu 3}$ decay

¹ ASSAMAGAN 94 result is from a new measurement of p_μ from $\pi^+ \rightarrow \mu^+ \nu_\mu$ at rest combined with JECKELMANN 94 Solution B pion mass. It is not used in "OUR EVALUATION" because it is not yet published. See Note above.

² ENQVIST 93 bases limit on the fact that thermalized wrong-helicity Dirac neutrinos would speed up expansion of early universe, thus reducing the primordial abundance. FULLER 91 exploits the same mechanism but in the older calculation obtains a larger production rate for these states, and hence a lower limit. Neutrino lifetime assumed to exceed nucleosynthesis time, ~ 1 s.

³ MAYLE 93 recalculates cooling rate enhancement by escape of wrong-helicity Dirac neutrinos using the Livermore Supernova Explosion Code, obtains more restrictive result than the "very conservative" BURROWS 92 limit because of higher core temperature.

⁴ There would be an increased cooling rate if Dirac neutrino mass is included; this does not apply for Majorana neutrinos. Limit is on $\sqrt{m^2_{\nu_\mu} + m^2_{\nu_\tau}}$, and error becomes very large if ν_τ is nonrelativistic, which occurs near the lab limit of 31 MeV. RAJPOOT 93 notes that limit could be evaded with new physics.

⁵ BURROWS 92 limit for Dirac neutrinos only.

⁶ Assumes neutrino lifetime > 1 s. For Dirac neutrinos only. See also ENQVIST 93.

⁷ NATALE 91 published result multiplied by $\sqrt{8}\sqrt{4}$ at the advice of the author.

⁸ GRIFOLS 90B estimated error is a factor of 3.

⁹ GAEMERS 89 published result (< 0.03) corrected via the GANDHI 91 erratum.

¹⁰ ABELA 84 used the PDG 84 value for π^\pm mass, in conjunction with μ momentum measurement in $\pi \rightarrow \mu \nu_\mu$ decay to obtain $m < 0.25$ and $m^2 = -0.16 \pm 0.08$. The values shown here for mass and m^2 are corrected values obtained by JECKELMANN 86 from the ABELA 84 data using the more accurate π^\pm mass of JECKELMANN 86.

¹¹ LU 80 combines DAUM 79 $\pi^+ \rightarrow \mu^+ \nu_\mu$ measurement with new LU 80 π^- mass and replaces DAUM 79. LU 80 is not independent of ABELA 84.

$m_{\nu_2} - m_{\nu_3}$

Test of CPT for a Dirac neutrino. (Not a very strong test.)

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.45	90	CLARK 74	ASPK	$K_{\mu 3}$ decay

Lepton & Quark Full Listings

ν_μ

ν_2 (MEAN LIFE) / MASS

These limits often apply to ν_τ (ν_3) also.

VALUE (s/eV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
>15.4	90		12 KRAKAUER	91 CNTR	$\nu_\mu, \bar{\nu}_\mu$ at LAMPF
> 2.8 × 10 ¹⁵			13,14 BLUDMAN	92 ASTR	$m_\nu < 50$ eV
none 10 ⁻¹² - 5 × 10 ⁴			15 DODELSON	92 ASTR	$m_\nu=1-300$ keV
> 6.3 × 10 ¹⁵			14,16 CHUPP	89 ASTR	$m_\nu < 20$ eV
> 1.7 × 10 ¹⁵			14 KOLB	89 ASTR	$m_\nu < 20$ eV
> 3.3 × 10 ¹⁴			17,18 HATSUDA	88 ASTR	
> 0.11	90	0	19,20 VONFEILIT...	88 ASTR	
			21 FRANK	81 CNTR	$\nu\bar{\nu}$ LAMPF
			22 HENRY	81 ASTR	$m_\nu = 16-20$ eV
			23 KIMBLE	81 ASTR	$m_\nu = 10-100$ eV
			24 REPHAELI	81 ASTR	$m_\nu = 30-150$ eV
			25 DERUJULA	80 ASTR	$m_\nu = 10-100$ eV
			26 STECKER	80 ASTR	$m_\nu = 10-100$ eV
> 2 × 10 ²¹			21 BLIETSCHAU	78 HLBC	$\nu_\mu, \bar{\nu}_\mu$ CERN GGM
> 1.0 × 10 ⁻²	90	0	21 BLIETSCHAU	78 HLBC	$\bar{\nu}_\mu, \nu_\mu$ CERN GGM
> 1.7 × 10 ⁻²	90	0	21 BARNES	77 DBC	$\nu, \bar{\nu}$ ANL 12-ft
> 2.2 × 10 ⁻³	90	0	21 BELLOTTI	76 HLBC	$\nu, \bar{\nu}$ CERN GGM
> 3. × 10 ⁻³	90	0	21 BELLOTTI	76 HLBC	$\bar{\nu}$ CERN GGM
> 1.3 × 10 ⁻²	90	1	21 BELLOTTI	76 HLBC	$\bar{\nu}$ CERN GGM

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 12 KRAKAUER 91 quotes the limit $\tau/m_{\nu_1} > (0.75a^2 + 21.65a + 26.3)$ s/eV, where a is a parameter describing the asymmetry in the neutrino decay defined as $dN_\gamma/d\cos\theta = (1/2)(1 + a\cos\theta)$. The parameter $a=0$ for a Majorana neutrino, but can vary from -1 to 1 for a Dirac neutrino. The bound given by the authors is the most conservative (which applies for $a = -1$).
- 13 BLUDMAN 92 sets additional limits by this method for higher mass ranges. Cosmological limits are also obtained.
- 14 Nonobservation of γ 's in coincidence with ν 's from SN 1987A. Results should be divided by the $\tau_\nu \rightarrow \gamma X$ branching ratio.
- 15 DODELSON 92 range is for wrong-helicity keV mass Dirac ν 's from the core of neutron star in SN 1987A decaying to ν 's that would have interacted in KAM2 or IMB detectors.
- 16 CHUPP 89 should be multiplied by a branching ratio (about 1) and a detection efficiency (about 1/4), and pertains to radiative decay of any neutrino to a lighter or sterile neutrino.
- 17 Model-dependent theoretical analysis.
- 18 HATSUDA 88 argues that previous bounds on radiative decays of neutrinos produced in supernovae explosions may not be valid (because $\nu_H \rightarrow \gamma\nu$ might be dominated by processes such as $\nu_H e \rightarrow \nu e$ if e number density is high enough), and that, in fact, a neutrino mean life/mass of 0.2-0.6 s/eV may be consistent with the data.
- 19 Model-dependent theoretical analysis of SN 1987A neutrinos.
- 20 Limit applies to ν_τ also.
- 21 These experiments look for $\nu_\mu \rightarrow \nu_e \gamma$ or $\bar{\nu}_\mu \rightarrow \bar{\nu}_e \gamma$.
- 22 HENRY 81 uses UV flux from clusters of galaxies to find $\tau > 1.1 \times 10^{25}$ s for radiative decay.
- 23 KIMBLE 81 uses extreme UV flux limits to find $\tau > 10^{22}-10^{23}$ s.
- 24 REPHAELI 81 consider ν decay γ effect on neutral H in early universe; based on M31 HI concludes $\tau > 10^{24}$ s.
- 25 DERUJULA 80 finds $\tau > 3 \times 10^{23}$ s based on CDM neutrino decay contribution to UV background.
- 26 STECKER 80 limit based on UV background; result given is $\tau > 4 \times 10^{22}$ s at $m_\nu = 20$ eV.

$|(\mathbf{v} - \mathbf{c}) / \mathbf{c}|$ ($\mathbf{v} \equiv \nu_2$ VELOCITY)

Expected to be zero for massless neutrino, but also tests whether photons and neutrinos have the same limiting velocity in vacuum.

VALUE (units 10 ⁻⁴)	CL%	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
<0.4	95	9800	KALBFLEISCH 79	SPEC		
<2.0	99	77	ALSPECTOR 76	SPEC	0	>5 GeV ν
<4.0	99	26	ALSPECTOR 76	SPEC	0	<5 GeV ν

ν_2 MAGNETIC MOMENT

Must vanish for Majorana neutrino or purely chiral massless Dirac neutrino. The value of the magnetic moment for the standard SU(2)×U(1) electroweak theory extended to include massive neutrinos (see FUJIKAWA 80) is $\mu_\nu = 3eG_F m_\nu / (8\pi^2 \sqrt{2}) = (3.2 \times 10^{-19}) m_\nu \mu_B$ where m_ν is in eV and $\mu_B = e\hbar/2m_e$ is the Bohr magneton. Given the upper bound $m_{\nu_2} < 0.27$ MeV, it follows that for the extended standard electroweak theory, $\mu(\nu_2) < 0.86 \times 10^{-13} \mu_B$.

VALUE (μ_B)	CL%	DOCUMENT ID	TECN	COMMENT
<8.5 × 10 ⁻¹⁰	90	AHRENS	90 CNTR	$\nu_\mu e \rightarrow \nu_\mu e$
<7.4 × 10 ⁻¹⁰	90	27 KRAKAUER	90 CNTR	LAMPF ($\nu_\mu, \bar{\nu}_\mu$) e elast.
<9.5 × 10 ⁻¹⁰	90	ABE	87B CNTR	$\nu_\mu e \rightarrow \nu_\mu e$

• • • We do not use the following data for averages, fits, limits, etc. • • •

- <1 × 10⁻⁸
- <2 × 10⁻¹²
- <1 × 10⁻¹¹
- <1.1 × 10⁻¹¹
- <6 × 10⁻¹⁴
- <4 × 10⁻¹¹
- <8.5 × 10⁻¹¹
- <8.1 × 10⁻⁹
- <1 × 10⁻¹⁰
- 27 KRAKAUER 90 experiment fully reported in ALLEN 93.
- 28 DORENBOSCH 91 corrects an incorrect statement in DORENBOSCH 89 that the ν_2 magnetic moment is $< 1 \times 10^{-9}$ at the 95%CL. DORENBOSCH 89 measures both $\nu_\mu e$ and $\bar{\nu}_\mu e$ elastic scattering and assume $\mu(\nu_\mu) = \mu(\bar{\nu}_\mu)$.
- 29 RAFFELT 90 limit applies for a diagonal magnetic moment of a Dirac neutrino, or for a transition magnetic moment of a Majorana neutrino. In the latter case, the same analysis gives $< 1.4 \times 10^{-12}$. Limit at 95%CL obtained from δM_C .
- 30 Significant dependence on details of stellar properties.
- 31 If $m_{\nu_2} < 10$ keV.
- 32 For $m_{\nu_2} = 8-200$ eV. NUSSINOV 87 examines transition magnetic moments for $\nu_\mu \rightarrow \nu_e$ and obtain $< 3 \times 10^{-15}$ for $m_{\nu_2} > 16$ eV and $< 6 \times 10^{-14}$ for $m_{\nu_2} > 4$ eV.
- 33 KIM 74 is a theoretical analysis of $\bar{\nu}_\mu$ reaction data.
- 34 If $m_{\nu_2} < 1$ keV.

NONSTANDARD CONTRIBUTIONS TO NEUTRINO SCATTERING

We report limits on the so-called neutrino charge radius squared in this section. This quantity is not an observable, physical quantity and this is reflected in the fact that it is gauge dependent (see LEE 77C). It is not necessarily positive. A more general interpretation of the experimental results is that they are limits on certain nonstandard contributions to neutrino scattering.

VALUE (10 ⁻³² cm ²)	DOCUMENT ID	TECN	COMMENT
-1.1 ± 1.0	35 AHRENS	90 CNTR	$\nu_\mu e$ elas scat
-0.3 ± 1.5	35 DORENBOSCH...	89 CHRM	$\nu_\mu e$ elas scat

35 Result is obtained from reanalysis given in ALLEN 91, followed by our reduction to obtain 1 σ errors.

ν_μ REFERENCES

ASSAMAGAN 94	PL B (subm.)	+Bronnimann, Daum+	(PSI, ZURI, VILL, VIRG)
JECKELMANN 94	PL B (accepted)	+Goudsmit, Leisi	(WABRN, VILL)
ALLEN 93	PR D47 11	+Chen, Doe, Hausamann+	(UCI, LANL, ANL, UMD)
ENQVIST 93	PL B301 376	+Ulbro	(NORD)
MAYLE 93	PL B317 119	+Schramm, Turner, Wilson	(LLNL, CHIC)
RAIPOOT 93	MPL A8 1179		(CSULB)
BLUDMAN 92	PR D45 4720		(CFPA)
BURROWS 92	PRL 68 3834	+Gandhi, Turner	(ARIZ, CHIC)
DODELSON 92	PRL 68 2572	+Frieman, Turner	(FNAL, CHIC)
MAMEDOV 92	SJPN 23 339		(JINR)
Translated from FECAV 23 767.			
ALLEN 91	PR D43 R1	+Chen, Doe, Hausamann	(UCI, LANL, UMD)
DAUM 91	PL B265 425	+Frosch, Herter, Janusch, Kettle	(VILL)
DORENBOSCH...	91 ZPHY C51 142	+Dorenbosch, Udo, Allaby, Amaldi+	(CHARM Collab.)
FULLER 91	PR D43 3136	+Malaney	(UCSD)
GANDHI 91	PL B261 519E (erratum)	+Burrows	(ARIZ)
KRAKAUER 91	PR D44 86	+Talaga, Allen, Chen+	(LAMPF E225 Collab.)
LAM 91	PR D44 3345	+Ng	(AST)
NATALE 91	PL B258 227		(SPIET)
AHRENS 90	PR D41 3297	+ (BNL, BROW, HIRO, KEK, OSAK, PENN, STON)	
GANDHI 90	PL B246 149	+Burrows	(ARIZ)
Also 91	PL B261 519E (erratum)	+Gandhi, Burrows	(ARIZ)
GRIFOLS 90B	PL B242 77	+Masso	(BARC, CERN)
KRAKAUER 90	PL B252 177	+Talaga, Allen, Chen+	(LAMPF E225 Collab.)
RAFFELT 90	PR 64 2856		(MPIM)
CHUPP 89	PRL 62 505	+Vestrand, Reppin	(UNH, MPIM)
DORENBOSCH...	89 ZPHY C41 567	+Dorenbosch, Udo, Allaby, Amaldi+	(CHARM Collab.)
GAEMERS 89	PR D40 309	+Gandhi, Lattimer	(ANIK, STON)
KOLB 89	PRL 62 509	+Turner	(CHIC, FNAL)
RAFFELT 89B	APJ 336 61	+Dearborn, Silk	(UCB, LLL)
HATSUDA 88	PL B203 462	+Lim, Yoshimura	(KEK)
VONFEILIT... 88	PL B200 580	+Von Felitzsch, Oberauer	(MUNT)
ABE 87B	PRL 58 636	+ (BNL, BROW, HIRO, KEK, OSAK, PENN, STON)	
FUKUGITA 87	PR D36 3817	+Yazaki	(KYOTU, TOKY)
NUSSINOV 87	PR D36 2278	+Rephaeli	(TELA)
JECKELMANN 86	PRL 56 1444	+Nakada, Beer+	(ETH, FRIB)
ABELA 84	PL 146B 431	+Daum, Eaton, Frosch, Jost, Kettle+	(SIN)
PDG 84	RMP 56 No. 2 Pt. II	+Wohl, Cahn, Rittenberg+	(LBL, CIT, CERN)
ANDERHUB 92	PL 14B 76	+Boecklin, Hofer, Kottmann+	(ETH, SIN)
FRANK 81	PR D24 2001	+Burman+	(STON)
HENRY 81	PRL 47 618	+Feldman	(LASL, YALE, MIT, SACL, SIN+)
KIMBLE 81	PRL 46 80	+Bowyer, Jakobsen	(JHU)
LYNN 81	PR D23 2151		(UCB)
REPHAELI 81	PL 106B 73	+Szalay	(UCSB, CHIC)
DERUJULA 80	PRL 45 942	+Glashow	(MIT, HARV)
FUJIKAWA 80	PRL 45 963	+Shrock	(STON)
LU 80	PRL 45 1066	+Delker, Dugan, Wu, Caffrey+	(YALE, COLU, JHU)
STECKER 80	PRL 45 1460		(NASA)
DAUM 79	PR D20 2692	+Eaton, Frosch, Hirschmann+	(SIN)
Also 76	PL 60B 380	+Daum, Dubal, Eaton, Frosch+	(SIN, ETH)
Also 78	PL 74B 126	+Daum, Eaton, Frosch, Hirschmann+	(SIN)

See key on page 1343

Lepton & Quark Full Listings

ν_μ, ν_τ

KALBFLEISCH 79	PRL 43 1361	+Baggett, Fowler+	(FNAL, PURD, BELL)
BEG 78	PR D17 1395	+Marciano, Ruderman	(ROCK, COLU)
BLIETSCHAU 78	NP B133 205	+Deden, Hasert, Krenz+	(Gargamelle Collab.)
BARNES 77	PRL 38 1049	+Carmony, Dauwe, Fernandez+	(PURD, ANL)
LEE 77C	PR D16 1444	+Shrock	(STON)
ALSPECTOR 76	PRL 36 837	+ (BNL, PURD, CIT, FNAL, ROCK)	
BELLOTTI 76	LNC 17 553	+Cavalli, Fiorini, Rollier	(MILA)
CLARK 74	PR D9 533	+Elioff, Frisch, Johnson, Kerth, Shen+	(LBL)
KIM 74	PR D9 3050	+Mather, Okubo	(ROCH)
BERNSTEIN 63	PR 132 1227	+Ruderman, Feinberg	(NYU, COLU)

ν_τ

$$J = \frac{1}{2}$$

Existence indirectly established from τ decay data combined with ν reaction data. See for example FELDMAN 81. ALBRECHT 92Q rules out $J = 3/2$ by establishing that the ρ^- is not in a pure $H_{\rho^-} = -1$ helicity state in $\tau^- \rightarrow \rho^- \nu_\tau$.

Not in general a mass eigenstate. See note on neutrinos in the ν_e section above.

ν_τ MASS

Applies to ν_3 , the primary mass eigenstate in ν_τ . Would also apply to any other ν_j which mixes strongly in ν_τ and has sufficiently small mass that it can occur in the respective decays. (This would be nontrivial only for a hypothetical $j \geq 4$, given the ν_e and ν_μ mass limits above.) See also the Listings in the Neutrino Bounds from Astrophysics and Cosmology section.

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 31	95	19	1 ALBRECHT 92M	ARG	$E_{cm}^{ee} = 9.4\text{--}10.6$ GeV
< 75	95		2 BALEST 93	CLEO	$E_{cm}^{ee} = 10.6$ GeV
< 32.6	95	113	3 CINABRO 93	CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
< 0.3 or > 35			4 DOLGOV 93	COSM	Nucleosynthesis
< 0.74			5 ENQVIST 93	COSM	Nucleosynthesis
< 0.003			6,7 MAYLE 93	ASTR	SN 1987A cooling
< 0.025–0.030			7,8 BURROWS 92	ASTR	SN 1987A cooling
< 0.3			9 FULLER 91	COSM	Nucleosynthesis
< 0.5 or > 25			10 KOLB 91	COSM	Nucleosynthesis
< 0.42			9 LAM 91	COSM	Nucleosynthesis
< 0.028–0.15			11 NATALE 91	ASTR	SN 1987A
< 0.028			7 GANDHI 90	ASTR	SN 1987A
< 0.014 or > 34			7,12 GRIFOLS 90B	ASTR	SN 1987A
< 0.06			7,13 GAEMERS 89	SN	1987A
< 35	95	12	14 ALBRECHT 88B	ARG	Repl. by AL-BRECHT 92M
< 76	95	13	15 ABACHI 87	HRS	$E_{cm}^{ee} = 29$ GeV
< 85	95		16 CSORNA 87B	CLEO	$E_{cm}^{ee} = 10\text{--}11$ GeV
< 84	95	10	17 ABACHI 86	HRS	Repl. by ABACHI 87
< 70	95	102	18 ALBRECHT 85I	ARG	$E_{cm}^{ee} = 10$ GeV
< 125	95	3	19 BURCHAT 85	MRK2	$E_{cm}^{ee} = 29$ GeV
< 143	95	22	20 MATTEUZZI 85	MRK2	$E_{cm}^{ee} = 29$ GeV
< 157	95	4	21 MILLS 85	DLCO	$E_{cm}^{ee} = 29$ GeV
< 250	95		22 BLOCKER 82D	MRK2	$E_{cm}^{ee} = 5.2$ GeV
< 250	95	594	23,24 BACINO 79B	DLCO	$E_{cm}^{ee} = 3.5\text{--}7.4$ GeV

- ALBRECHT 92M reports measurement of a slightly lower τ mass, which has the effect of reducing the ν_τ mass reported in ALBRECHT 88B. Bound is from analysis of $\tau^- \rightarrow 3\pi^- 2\pi^+ \nu_\tau$ mode.
- BALEST 93 derive limit by comparing their m_τ measurement (which depends on m_{ν_τ}) to BAI 92 and BACINO 78B m_τ threshold measurements.
- CINABRO 93 bound comes from analysis of $\tau^- \rightarrow 3\pi^- 2\pi^+ \nu_\tau$ and $\tau^- \rightarrow 2\pi^- \pi^+ 2\pi^0 \nu_\tau$ decay modes.
- DOLGOV 93 assumes neutrino lifetime > 100 s. For Majorana neutrinos, the low mass limit is 0.5 MeV. KAWANO 92 points out that these bounds can be overcome for a Dirac neutrino if it possesses a magnetic moment.
- ENQVIST 93 bases limit on the fact that thermalized wrong-helicity Dirac neutrinos would speed up expansion of early universe, thus reducing the primordial abundance. FULLER 91 exploits the same mechanism but in the older calculation obtains a larger production rate for these states, and hence a lower limit. Neutrino lifetime assumed to exceed nucleosynthesis time, ~ 1 s.
- MAYLE 93 recalculates cooling rate enhancement by escape of wrong-helicity Dirac neutrinos using the Livermore Supernova Explosion Code, obtains more restrictive result than the "very conservative" BURROWS 92 limit because of higher core temperature.
- There would be an increased SN 1987A cooling rate if Dirac neutrino mass is included; this does not apply for Majorana neutrinos. Limit is on $\sqrt{m^2 \nu_\mu + m^2 \nu_\tau}$, and error becomes very large if ν_τ is nonrelativistic, which occurs near the lab limit of 31 MeV. RAJPOOT 93 notes that limit could be evaded with new physics.
- BURROWS 92 limit for Dirac neutrinos only.
- Assumes neutrino lifetime > 1 s. For Dirac neutrinos. See also ENQVIST 93.

- KOLB 91 exclusion region is for Dirac neutrino with lifetime > 1 s; other limits are given.
- NATALE 91 published result multiplied by $\sqrt{8/4}$ at the advice of the author.
- GRIFOLS 90B estimated error is a factor of 3.
- GAEMERS 89 published result (< 0.03) corrected via the GANDHI 91 erratum.
- ALBRECHT 88 bound comes from analysis of $\tau \rightarrow 5\pi^\pm \nu_\tau$ decay mode. Same data reanalyzed with revised τ mass in ALBRECHT 92M.
- Bound comes from analysis of $\tau \rightarrow 5\pi^\pm (\pi^0) \nu_\tau$ decay mode in 13 decay events.
- CSORNA 87B also quote result as $31 \pm 25 \pm 20$ MeV. Bound comes from analysis of $\tau \rightarrow 3\pi^\pm (\pi^0) \nu_\tau$ decay mode.
- Bound comes from analysis of $\tau \rightarrow 5\pi^\pm \pi^0 \nu_\tau$ decay mode (5 events) and to a lesser extent from $\tau \rightarrow 5\pi^\pm \nu_\tau$ mode (5 events).
- Bound comes from analysis of $\tau \rightarrow 3\pi^\pm \nu_\tau$ decay mode.
- Bound comes from analysis of $\tau \rightarrow 5\pi^\pm (\pi^0) \nu_\tau$ decay.
- Bound comes from analysis of $\tau \rightarrow 3\pi^\pm \pi^0 \nu_\tau$ decay mode.
- Bound comes from analysis of $\tau \rightarrow K^\pm K^\mp \pi^\pm \nu_\tau$ decay mode.
- Bound comes from analysis of $\tau \rightarrow \pi \nu_\tau$ decay mode.
- Bound comes from analysis of leptonic decay spectrum.
- BACINO 79B experiment rules out V+A decay, disfavors pure V or A, and is in good agreement with V-A.

ν_3 (MEAN LIFE) / MASS

These limits often apply to ν_μ (ν_2) also.

VALUE (s/eV)	DOCUMENT ID	TECN	COMMENT
$> 2.8 \times 10^{15}$	25,26 BLUDMAN 92	ASTR	$m_\nu < 50$ eV
$< 10^{-12}$ or $> 5 \times 10^4$	27 DODELSON 92	ASTR	$m_\nu = 1\text{--}300$ keV
	28 GRANEK 91	COSM	Decaying L^0
	29 WALKER 90	ASTR	$m_\nu = 0.03\text{--}2$ MeV
$> 6.3 \times 10^{15}$	26,30 CHUPP 89	ASTR	$m_\nu < 20$ eV
$> 1.7 \times 10^{15}$	26 KOLB 89	ASTR	$m_\nu < 20$ eV
	31 TERASAWA 88	COSM	$m_\mu = 30\text{--}70$ MeV
	32 KAWASAKI 86	COSM	$m_\nu > 10$ MeV
	33 LINDLEY 85	COSM	$m_\nu > 10$ MeV
	34 BINETRUY 84	COSM	$m_\nu \sim 1$ MeV
	35 SARKAR 84	COSM	$m_\nu = 10\text{--}100$ MeV
	36 HENRY 81	ASTR	$m_\nu = 16\text{--}20$ eV
	37 KIMBLE 81	ASTR	$m_\nu = 10\text{--}100$ eV
	38 REPHAELI 81	ASTR	$m_\nu = 30\text{--}150$ eV
	39 DERUJULA 80	ASTR	$m_\nu = 10\text{--}100$ eV
$> 2 \times 10^{21}$	40 STECKER 80	ASTR	$m_\nu = 10\text{--}100$ eV
	41 DICUS 78	COSM	$m_\nu = 0.5\text{--}30$ MeV
$< 3 \times 10^{-11}$	42 FALK 78	ASTR	$m_\nu < 10$ MeV
	43 COWSIK 77	ASTR	

- • • We do not use the following data for averages, fits, limits, etc. • • •
 - $> 2.8 \times 10^{15}$
 - $< 10^{-12}$ or $> 5 \times 10^4$
 - $> 6.3 \times 10^{15}$
 - $> 1.7 \times 10^{15}$
 - $> 2 \times 10^{21}$
 - $< 3 \times 10^{-11}$
- BLUDMAN 92 sets additional limits by this method for higher mass ranges. Cosmological limits are also obtained.
 - Nonobservation of γ 's in coincidence with ν 's from SN 1987A. Results should be divided by the $\tau_\nu \rightarrow \gamma X$ branching ratio.
 - DODELSON 92 range is for wrong-helicity keV mass Dirac ν 's from the core of neutron star in SN 1987A decaying to ν 's that would have interacted in KAM2 or IMB detectors.
 - GRANEK 91 considers heavy neutrino decays to $\gamma \nu_L$ and $3\nu_L$, where $m_{\nu_L} < 100$ keV. Lifetime is calculated as a function of heavy neutrino mass, branching ratio into $\gamma \nu_L$ and m_{ν_L} .
 - WALKER 90 uses SN 1987A γ flux limits after 289 days to find $m_\tau > 1.1 \times 10^{15}$ eV s.
 - CHUPP 89 should be multiplied by a branching ratio (about 1) and a detection efficiency (about 1/4), and pertains to radiative decay of any neutrino to a lighter or sterile neutrino.
 - TERASAWA 88 finds only $10^2 < \tau < 10^4$ allowed for 30–70 MeV ν 's from primordial nucleosynthesis.
 - KAWASAKI 86 concludes that light elements in primordial nucleosynthesis would be destroyed by radiative decay of neutrinos with $10 \text{ MeV} < m_\nu < 1 \text{ GeV}$ unless $\tau \lesssim 10^4$ s.
 - LINDLEY 85 considers destruction of cosmologically-produced light elements, and finds $\tau < 2 \times 10^3$ s for $10 \text{ MeV} < m_\nu < 100 \text{ MeV}$. See also LINDLEY 79.
 - BINETRUY 84 finds $\tau < 10^8$ s for neutrinos in a radiation-dominated universe.
 - SARKAR 84 finds $\tau < 20$ s at $m_\nu = 10$ MeV, with higher limits for other m_ν , and claims that all masses between 1 MeV and 50 MeV are ruled out.
 - HENRY 81 uses UV flux from clusters of galaxies to find $\tau > 1.1 \times 10^{25}$ s for radiative decay.
 - KIMBLE 81 uses extreme UV flux limits to find $\tau > 10^{22}\text{--}10^{23}$ s.
 - REPHAELI 81 consider ν decay γ effect on neutral H in early universe; based on M31 HI concludes $\tau > 10^{24}$ s.
 - DERUJULA 80 finds $\tau > 3 \times 10^{23}$ s based on CDM neutrino decay contribution to UV background.
 - STECKER 80 limit based on UV background; result given is $\tau > 4 \times 10^{22}$ s at $m_\nu = 20$ eV.
 - DICUS 78 considers effect of ν decay photons on light-element production, and finds lifetime must be less than "hours." See also DICUS 77.
 - FALK 78 finds lifetime constraints based on supernova energetics.
 - COWSIK 77 considers variety of scenarios. For neutrinos produced in the big bang, present limits on optical photon flux require $\tau > 10^{23}$ s for $m_\nu \sim 1$ eV. See also COWSIK 79 and GOLDMAN 79.

Lepton & Quark Full Listings

ν_τ, e

ν_3 MAGNETIC MOMENT

Must vanish for Majorana neutrino or purely chiral massless Dirac neutrino. The value of the magnetic moment for the standard $SU(2) \times U(1)$ electroweak theory extended to include massive neutrinos (see FUJIKAWA 80) is $\mu_\nu = 3eG_F m_\nu / (8\pi^2 \sqrt{2}) = (3.20 \times 10^{-19}) m_\nu \mu_B$ where m_ν is in eV and $\mu_B = e\hbar/2m_e$ is the Bohr magneton. Given the upper bound $m_{\nu_3} < 35$ MeV, it follows that for the extended standard electroweak theory, $\mu(\nu_3) < 1.1 \times 10^{-11} \mu_B$.

VALUE (μ_B)	CL%	DOCUMENT ID	TECN	COMMENT
$< 5.4 \times 10^{-7}$	90	44 COOPER...	92 BEBC	$\nu_\tau e^- \rightarrow \nu_\tau e^-$
$> 10^{-8}$		45 KAWANO	92 ASTR	Primordial ^4He abundance
$< 5.6 \times 10^{-6}$	90	DESHPANDE	91 RVUE	$e^+ e^- \rightarrow \nu \bar{\nu} \gamma$
$< 2 \times 10^{-12}$	95	46 RAFFELT	90 ASTR	Red giant luminosity
$< 1 \times 10^{-11}$		47 RAFFELT	89B ASTR	Cooling helium stars
$< 4 \times 10^{-6}$	90	48 GROTCHE	88 RVUE	$e^+ e^- \rightarrow \nu \bar{\nu} \gamma$
$< 1.1 \times 10^{-11}$		47,49 FUJIKAWA	87 ASTR	Cooling helium stars
$< 6 \times 10^{-14}$		50 NUSSINOV	87 ASTR	Cosmic EM backgrounds
$< 8.5 \times 10^{-11}$		49 BEG	78 ASTR	Stellar plasmons
44 COOPER-SARKAR 92 assume $f_{D_s}/f_\tau = 2$ and D_s, \bar{D}_s production cross section = $2.6 \mu\text{b}$ to calculate ν_τ flux.				
45 KAWANO 92 lower limit is that needed to circumvent ^4He production if m_{ν_τ} is between 5 and ~ 30 MeV/ c^2 .				
46 RAFFELT 90 limit valid if $m_{\nu_3} < 5$ keV. It applies for a diagonal magnetic moment of a Dirac neutrino, or for a transition magnetic moment of a Majorana neutrino. In the latter case, the same analysis gives $< 1.4 \times 10^{-12}$. Limit at 95%CL obtained from δM_C .				
47 Significant dependence on details of stellar properties.				
48 GROTCHE 88 combined data from MAC, ASP, CELLO, and Mark J.				
49 If $m_{\nu_3} < 10$ keV.				
50 For $m_{\nu_3} = 8-200$ eV. NUSSINOV 87 examines transition magnetic moments for $\nu_\tau \rightarrow \nu_e$ and obtain $< 3 \times 10^{-15}$ for $m_{\nu_3} < 16$ eV and $< 6 \times 10^{-14}$ for $m_{\nu_3} > 4$ eV.				

LIMIT ON ν_τ PRODUCTION IN BEAM DUMP EXPERIMENT

VALUE	DOCUMENT ID	TECN
$> 10^{-8}$	51 DORENBOS...	88 CHRМ
	52 BOFILL	87 CNTR
	53 TALEBZADEH	87 BEBC
	54 USHIDA	86C EMUL
	55 ASRATYAN	81 HLBC
	56 FRITZE	80 BEBC
51 DORENBOSCH 88 is CERN SPS beam dump experiment with the CHARM detector. $\nu_\tau + \bar{\nu}_\tau$ flux is $< 21\%$ of the total prompt flux at 90% CL.		
52 BOFILL 87 is a Fermilab narrow-band ν beam with a fine-grained neutrino detector.		
53 TALEBZADEH 87 is a CERN SPS beam dump experiment with the BEBC detector. Mixing probability $P(\nu_e \rightarrow \nu_\tau) < 18\%$ at 90% CL.		
54 USHIDA 86C is a Fermilab wide-band ν beam with a hybrid emulsion spectrometer. Mixing probabilities $P(\nu_e \rightarrow \nu_\tau) < 7.3\%$ and $P(\nu_\mu \rightarrow \nu_\tau) < 0.2\%$ at 90% CL.		
55 ASRATYAN 81 is a Fermilab wide-band $\bar{\nu}$ beam with a 15 foot bubble chamber. Mixing probability $P(\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau) < 2.2\%$ at 90% CL.		
56 FRITZE 80 is CERN SPS experiment with BEBC. Neutral-current/charged-current ratio corresponds to $R = (\text{prompt-}\nu\text{-induced events})/(\text{all prompt-}\nu\text{ events}) < 0.1$. Mixing probability $P(\nu_e \rightarrow \nu_\tau) < 0.35$ at CL = 90%.		

ν_τ REFERENCES

BALEST	93	PR D47 R3671	+Daoudi, Ford, Johnson+ (CLEO Collab.)
CINABRO	93	PRL 70 3700	+Henderson, Kinoshita+ (CLEO Collab.)
DOLGOV	93	PRL 71 476	+Rothstein (MICH)
ENQVIST	93	PL B301 376	+Uibo (NORD)
MAYLE	93	PL B317 119	+Schramm, Turner, Wilson (LLNL, CHIC)
RAJPOOT	93	MPL A8 1179	(CSULB)
ALBRECHT	92M	PL B292 221	+Ehrlichmann, Hamacher, Hofmann+ (ARGUS Collab.)
ALBRECHT	92Q	ZPHY C56 339	+Ehrlichmann, Hamacher+ (ARGUS Collab.)
BAI	92	PRL 69 3021	+Bardon, Becker-Szendy, Burnett+ (BES Collab.)
BLUDMAN	92	PR D45 4720	(CFPA)
BURROWS	92	PRL 68 3834	+Gandhi, Turner (ARIZ, CHIC)
COOPER...	92	PL B280 153	Cooper-Sarkar, Sarkar, Guy, Venus+ (BEBC WA66 Collab.)
DODELSON	92	PRL 68 2572	+Frieman, Turner (FNAL, CHIC)
KAWANO	92	PL B275 487	+Fuller, Malaney, Savage (CIT, UCSD, LLL, RUTG)
DESHPANDE	91	PR D43 943	+Sarma (OREG, TATA)
FULLER	91	PR D43 3136	+Malaney (UCSD)
GANDHI	91	PL B261 519E (erratum)	Burrows (ARIZ)
GRANEK	91	IJMP A6 2387	+McKellar (MELB)
KOLB	91	PRL 67 533	+Turner, Chakravorty, Schramm (FNAL, CHIC)
LAM	91	PR D44 3345	+Ng (AST)
NATALE	91	PL B258 227	(SPIFT)
GANDHI	90	PL B246 149	+Burrows (ARIZ)
Also	91	PL B261 519E (erratum)	Gandhi, Burrows (ARIZ)
GRIFOLS	90B	PL B242 77	+Masso (BARC, CERN)
RAFFELT	90	PRL 64 2856	(MPIM)
WALKER	90	PR D41 689	(HARV)
CHUPP	89	PRL 62 505	+Vestrand, Reppin (UNH, MPIM)
GAEMERS	89	PR D40 309	+Gandhi, Lattimer (ANIK, STON)
KOLB	89	PRL 62 509	+Turner (CHIC, FNAL)
RAFFELT	89B	APJ 336 61	+Dearborn, Silk (UCB, LLL)
ALBRECHT	88	PL B207 349	+Binder, Boeckmann+ (ARGUS Collab.)
ALBRECHT	88B	PL B202 149	+Binder, Boeckmann+ (ARGUS Collab.)
DORENBOS...	88	ZPHY C40 497	Dorenbosch, Allaby, Amaldi, Barbiellini+ (CHARM Collab.)
GROTCHE	88	ZPHY C39 553	+Robinett (PSU)
TERASAWA	88	NP B302 697	+Kawasaki, Sato (TOKY)
ABACHI	87	PR D35 2880	+Baringer, Bylsma, DeBonte+ (HRS Collab.)

BOFILL	87	PR D36 3309	+Busza, Eldridge+ (MIT, FNAL, MSU)
CSORNA	87B	PR D35 2747	+Mestayer, Panvini, Word+ (CLEO Collab.)
FUKUGITA	87	PR D36 3817	+Yazaki (KYOTY, TOKY)
NUSSINOV	87	PR D36 2278	+Rephael (TELA)
TALEBZADEH	87	NP B291 503	+Guy, Venus+ (BEBC WA66 Collab.)
ABACHI	86	PRL 56 1039	+Akerlof, Baringer, Beltrami+ (HRS Collab.)
KAWASAKI	86	PL B178 71	+Terasawa, Sato (TOKY)
USHIDA	86C	PRL 57 2897	+Kondo, Tasaka, Park, Song+ (FNAL E531 Collab.)
ALBRECHT	85I	PL 163B 404	+Binder, Drescher, Schubert+ (ARGUS Collab.)
BURCHAT	85	PRL 54 2489	+Schmidke, Yelton, Abrams+ (Mark II Collab.)
LINDLEY	85	APJ 294 1	(FNAL)
MATTEUZZI	85	PR D32 800	+Barklow+ (Mark II Collab.)
MILLS	85	PRL 54 624	+Pal, Atwood, Bailon+ (DELCO Collab.)
BINETRUY	84	PL 134B 174	+Girardi, Salati (LAPP)
SARKAR	84	PL 148B 347	+Cooper (OXF, CERN)
BLOCKER	82D	PL 109B 119	+Dorfan, Abrams, Alam+ (Mark II Collab.)
ASRATYAN	81	PL 105B 301	+Eftremenko, Fedotov+ (ITEP, FNAL, SERP, MICH)
FELDMAN	81	SLAC-PUB-2839	(SLAC, STAN)
Santa Cruz APS.			
HENRY	81	PRL 47 618	+Feldman (JHU)
KIMBLE	81	PRL 46 80	+Bowyer, Jakobsen (UCB)
REPHELI	81	PL 106B 73	+Szalay (UCSB, CHIC)
DERUJULA	80	PRL 45 942	+Glashow (MIT, HARV)
FRITZE	80	PL 96B 427	(AACH3, BONN, CERN, LOIC, OXF, SAFL)
FUJIKAWA	80	PRL 45 963	+Shrock (STON)
STECKER	80	PRL 45 1460	(NASA)
BACINO	79B	PRL 42 749	+Ferguson, Nodulman, Slater+ (DELCO Collab.)
COWSIK	79	PR D19 2219	(TATA)
GOLDMAN	79	PR D19 2215	+Stephenson (LASL)
LINDLEY	79	MNRAS 188 15P	(SUSS)
BACINO	78B	PRL 41 13	+Ferguson, Nodulman, Slater+ (DELCO Collab.)
BEG	78	PR D17 1395	+Mariano, Ruderman (ROCK, COLU)
DICUS	78	PR D17 1529	+Kolb, Tepitz, Wagoner (TEXA, VPI, STAN)
FALK	78	PL 79B 511	+Schramm (CHIC)
COWSIK	77	PRL 39 784	(MPIM, TATA)
DICUS	77	PRL 39 168	+Kolb, Tepitz (TEXA, VPI)

OTHER RELATED PAPERS

WEINSTEIN	93	ARNPS 43 457	+Stroynowski (CIT, SMU)
-----------	----	--------------	-------------------------



$$J = \frac{1}{2}$$

e MASS

The mass is known much more precisely in u (atomic mass units) than in MeV (see the footnote). The conversion from u to MeV, $1 u = 931.49432 \pm 0.00028$ MeV, involves the relatively poorly known electronic charge.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
$0.51099906 \pm 0.00000015$	1 COHEN	87 RVUE	1986 CODATA value
$> 10^{-4}$			• • • We do not use the following data for averages, fits, limits, etc. • • •
0.5110034 ± 0.0000014	COHEN	73 RVUE	1973 CODATA value
1 The mass is known much more precisely in u: $m = (5.48579903 \pm 0.00000013) \times 10^{-4} u$.			

$$(m_{e^+} - m_{e^-}) / m_{\text{average}}$$

A test of CPT invariance.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-8}$	90	CHU	84 CNTR	Positronium spectroscopy

$$|q_{e^+} + q_{e^-}|/e$$

A test of CPT invariance. See also similar tests involving the proton.

VALUE	DOCUMENT ID	TECN	COMMENT
$< 4 \times 10^{-8}$	2 HUGHES	92 RVUE	
$> 10^{-8}$			• • • We do not use the following data for averages, fits, limits, etc. • • •
$> 10^{-8}$	3 MUELLER	92 THEO	Vacuum polarization
2 HUGHES 92 uses recent measurements of Rydberg-energy and cyclotron-frequency ratios.			
3 MUELLER 92 argues that an inequality of the charge magnitudes would, through higher-order vacuum polarization, contribute to the net charge of atoms.			

e MAGNETIC MOMENT ANOMALY

$$\mu_e/\mu_B - 1 = (g-2)/2$$

For the most accurate theoretical calculation, see KINOSHITA 81.

Some older results have been omitted.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	CHG	COMMENT
1159.682193 ± 0.000010	4 COHEN	87 RVUE		1986 CODATA value
$1159.6521884 \pm 0.0000043$	VANDYCK	87 MRS	-	Single electron
$1159.6521879 \pm 0.0000043$	VANDYCK	87 MRS	+	Single positron
1159.652200 ± 0.000040	VANDYCK	86 MRS	-	Single electron
1159.652222 ± 0.000050	SCHWINBERG	81 MRS	+	Single positron

4 The COHEN 87 value assumes the $g/2$ values for e^+ and e^- are equal, as required by CPT.

$$(\mathcal{G}_{e^+} - \mathcal{G}_{e^-}) / \mathcal{G}_{\text{average}}$$

A test of CPT invariance.

VALUE (units 10^{-12})	CL%	DOCUMENT ID	TECN	COMMENT
-0.5 ± 2.1		⁵ VANDYCK	87 MRS	Penning trap
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 12	95	⁶ VASSERMAN	87 CNTR	Assumes $m_{e^+} = m_{e^-}$
22 ± 64		SCHWINBERG	81 MRS	Penning trap
⁵ VANDYCK 87 measured $(g_-/g_+) - 1$ and we converted it.				
⁶ VASSERMAN 87 measured $(g_+ - g_-)/(g - 2)$. We multiplied by $(g - 2)/g = 1.2 \times 10^{-3}$.				

e ELECTRIC DIPOLE MOMENT

A nonzero value is forbidden by both T invariance and P invariance.

VALUE (10^{-26} ecm)	CL%	DOCUMENT ID	TECN	COMMENT
-0.27 ± 0.83		⁷ ABDULLAH	90 MRS	205Tl beams
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
-14 ± 24		CHO	89 NMR	TI F molecules
$-1.5 \pm 5.5 \pm 1.5$		MURTHY	89	Cesium, no β field
-50 ± 110		LAMOREAUX	87 NMR	199Hg
190 ± 340	90	SANDARS	75 MRS	Thallium
70 ± 220	90	PLAYER	70 MRS	Xenon
< 300	90	WEISSKOPF	68 MRS	Cesium
⁷ ABDULLAH 90 uses the relativistic enhancement of a valence electron's electric dipole moment in a high-Z atom.				

e MEAN LIFE / BRANCHING FRACTION

A test of charge conservation. See the "Note on Testing Charge Conservation and the Pauli Exclusion Principle" following this section in our 1992 edition (Physical Review D45, 1 June, Part II (1992), p.VI.10). We use the best "disappearance" limit for the Summary Tables. The best limit for the specific channel $e^- \rightarrow \nu\gamma$ is much better.

Note that we use the mean life rather than what is often reported, the half life.

VALUE (yr)	CL%	DOCUMENT ID	TECN	COMMENT
$>2.7 \times 10^{23}$		REUSSER	91 CNTR	Ge K-shell disappearance
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
$>2.35 \times 10^{25}$	68	BALYSH	93 CNTR	$e^- \rightarrow \nu_e\gamma$, ^{76}Ge detector
$>1.5 \times 10^{25}$	68	AVIGNONE	86 CNTR	$e^- \rightarrow \nu\gamma$
$>1 \times 10^{39}$		⁸ ORITO	85 ASTR	Astrophysical argument
$>3 \times 10^{23}$	68	BELLOTTI	83B CNTR	$e^- \rightarrow \nu\gamma$
$>2 \times 10^{22}$	68	BELLOTTI	83B CNTR	Ge K-shell disappearance
$>3.5 \times 10^{23}$	68	KOVALCHUK	79 CNTR	$e^- \rightarrow \nu\gamma$
$>2 \times 10^{22}$	68	⁹ KOVALCHUK	79 CNTR	Disappearance
$>5.3 \times 10^{21}$		⁹ STEINBERG	75 CNTR	Disappearance
$>2 \times 10^{21}$	9,10	MOE	65 CNTR	Disappearance
$>4 \times 10^{22}$		MOE	65 CNTR	$e^- \rightarrow \nu\gamma$

⁸ORITO 85 assumes that electromagnetic forces extend out to large enough distances and that the age of our galaxy is 10^{10} years.

⁹These limits are for all modes in which decay particles escape from the detector without depositing energy.

¹⁰The MOE 65 limit is re-estimated by STEINBERG 75 to be 10^{20} years.

e REFERENCES

BALYSH 93	PL B298 278	+Beck, Belyaev, Bensch+	(KIAE, MPIH, SASSO)
HUGHES 92	PRL 69 578	+Deutch	(LANL, AARH)
MUELLER 92	PRL 69 3432	+Thoma	(DUKE)
PDG 92	PR D45, 1 June, Part II	Hikasa, Barnett, Stone+	(KEK, LBL, BOST+)
REUSSER 91	PL B255 143	+Treichel, Boehm, Broggini+	(NEUC, CIT, VILL)
ABDULLAH 90	PRL 65 2347	+Carberg, Commins, Gould, Ross	(LBL, UCB)
CHO 89	PRL 63 2559	+Sangster, Hinds	(YALE)
MURTHY 89	PRL 63 965	+Krause, Li, Hunter	(AMHT)
COHEN 87	RMP 59 1121	+Taylor	(RISC, NBS)
LAMOREAUX 87	PRL 59 2275	+Jacobs, Heckel, Raab, Fortson	(WASH)
VANDYCK 87	PRL 59 26	+Van Dyck, Schwinberg, Dehmelt	(WASH)
VASSERMAN 87	PL B198 302	+Vorobyov, Gluskin+	(NOVO)
Also 87B	PL B187 172	+Vasserman, Vorobyov, Gluskin+	(NOVO)
AVIGNONE 86	PR D34 97	+Brodzinski, Hensley, Miley, Reeves+	(PNL, SCUC)
VANDYCK 86	PR D34 722	+Van Dyck, Schwinberg, Dehmelt	(WASH)
ORITO 85	PRL 54 2457	+Yoshimura	(TOKY, KEK)
CHU 84	PRL 52 1689	+Mills, Hall	(BELL, NBS, COLO)
BELLOTTI 83B	PL 124B 435	+Corti, Fiorini, Liguori, Pullia+	(MILA)
KINOSHITA 81	PRL 47 1573	+Lindquist	(CORN)
SCHWINBERG 81	PRL 47 1679	+Van Dyck, Dehmelt	(WASH)
KOVALCHUK 79	JETPL 29 145	+Pomansky, Smolinikov	(INRM)
Translated from ZETFP	29 163.		
SANDARS 75	PR A11 473	+Sternerheimer	(OXF, BNL)
STEINBERG 75	PR D12 2582	+Kwiatkowski, Maenhaut+	(UMD)
COHEN 73	JPCRD 2 663	+Taylor	(RISC, NBS)
PLAYER 70	JPB 3 1620	+Sandars	(OXF)
WEISSKOPF 68	PRL 21 1645	+Carrico, Gould, Lipworth+	(BRAN)
MOE 65	PR 140B 992	+Reines	(CASE)



$$J = \frac{1}{2}$$

μ MASS

The mass is known more precisely in u (atomic mass units) than in MeV (see the footnote to COHEN 87). The conversion from u to MeV, $1 u = 931.49432 \pm 0.00028$ MeV, involves the relatively poorly known electronic charge.

Where m_{μ}/m_e was measured, we have used the 1986 CODATA value for $m_e = 0.51099906 \pm 0.0000015$ MeV.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
105.658399 ± 0.000034	¹ COHEN	87 RVUE		1986 CODATA value
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
105.65841 ± 0.00033	² BELTRAMI	86 SPEC	-	Muonic atoms
105.658432 ± 0.000064	³ KLEMPPT	82 CNTR	+	Incl. in MARIAM 82
105.658386 ± 0.000044	⁴ MARIAM	82 CNTR	+	
105.65856 ± 0.00015	⁵ CASPERSON	77 CNTR	+	
105.65836 ± 0.00026	⁶ CROWE	72 CNTR		
105.65865 ± 0.00044	⁷ CRANE	71 CNTR		

¹The mass is known more precisely in u: $m = 0.113428913 \pm 0.000000017$ u. COHEN 87 makes use of the other entries below.

²BELTRAMI 86 gives $m_{\mu}/m_e = 206.76830(64)$.

³KLEMPPT 82 gives $m_{\mu}/m_e = 206.76835(11)$.

⁴MARIAM 82 gives $m_{\mu}/m_e = 206.768259(62)$.

⁵CASPERSON 77 gives $m_{\mu}/m_e = 206.76859(29)$.

⁶CROWE 72 gives $m_{\mu}/m_e = 206.7682(5)$.

⁷CRANE 71 gives $m_{\mu}/m_e = 206.76878(85)$.

μ MEAN LIFE τ

Measurements with an error $> 0.001 \times 10^{-6}$ s have been omitted.

VALUE (10^{-6} s)	DOCUMENT ID	TECN	CHG
2.19703 ± 0.00004	OUR AVERAGE		
2.197078 ± 0.000073	BARDIN	84 CNTR	+
2.197025 ± 0.000155	BARDIN	84 CNTR	-
2.19695 ± 0.00006	GIOVANETTI	84 CNTR	+
2.19711 ± 0.00008	BALANDIN	74 CNTR	+
2.1973 ± 0.0003	DUCLOS	73 CNTR	+

$\tau_{\mu^+}/\tau_{\mu^-}$ MEAN LIFE RATIO

A test of CPT invariance.

VALUE	DOCUMENT ID	TECN	COMMENT
1.000024 ± 0.000078	BARDIN	84 CNTR	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
1.0008 ± 0.0010	BAILEY	79 CNTR	Storage ring
1.000 ± 0.001	MEYER	63 CNTR	Mean life μ^+/μ^-

$$(\tau_{\mu^+} - \tau_{\mu^-}) / \tau_{\text{average}}$$

A test of CPT invariance. Calculated from the mean-life ratio, above.

VALUE	DOCUMENT ID
$(2 \pm 8) \times 10^{-5}$	OUR EVALUATION

μ MAGNETIC MOMENT ANOMALY

$$\mu_{\mu}/(e\hbar/2m_{\mu}) - 1 = (g_{\mu} - 2)/2$$

For reviews of theory and experiments, see HUGHES 85, KINOSHITA 84, COMB-LEY 81, FARLEY 79, and CALMET 77.

VALUE (units 10^{-6})	DOCUMENT ID	TECN	CHG	COMMENT
1165.9230 ± 0.0084	COHEN	87 RVUE		1986 CODATA value
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1165.910 ± 0.011	⁸ BAILEY	79 CNTR	+	Storage ring
1165.937 ± 0.012	⁸ BAILEY	79 CNTR	-	Storage ring
1165.923 ± 0.0085	⁸ BAILEY	79 CNTR	\pm	Storage ring
1165.922 ± 0.009	⁸ BAILEY	77 CNTR	\pm	Storage ring
1166.16 ± 0.31	BAILEY	68 CNTR	\pm	Storage rings
1162.0 ± 5.0	CHARPAK	62 CNTR	+	

⁸BAILEY 79 is final result. Includes BAILEY 77 data. We use μ/p magnetic moment ratio = 3.1833452 and recalculate the BAILEY 79 values. Third BAILEY 79 result is first two combined.

Lepton & Quark Full Listings

 μ

$$(\mathcal{G}_{\mu^+} - \mathcal{G}_{\mu^-}) / \mathcal{G}_{\text{average}}$$

A test of CPT invariance.

VALUE (units 10^{-8})	DOCUMENT ID	TECN	CHG	COMMENT
-2.6 ± 1.6	BAILEY	79		

 μ ELECTRIC DIPOLE MOMENTA nonzero value is forbidden by both T invariance and P invariance.

VALUE (10^{-19} e cm)	DOCUMENT ID	TECN	CHG	COMMENT
3.7 ± 3.4	⁹ BAILEY	78	CNTR \pm	Storage ring
8.6 ± 4.5	BAILEY	78	CNTR $+$	Storage rings
0.8 ± 4.3	BAILEY	78	CNTR $-$	Storage rings

⁹This is the combination of the two BAILEY 78 results given below. μ/p MAGNETIC MOMENT RATIOThis ratio is used to obtain a precise value of the muon mass. Measurements with an error > 0.00001 have been omitted.

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
$3.18334547 \pm 0.00000047$	¹⁰ COHEN	87	RVUE	1986 CODATA value
3.1833441 ± 0.0000017	KLEMP	82	CNTR $+$	Precession strob
3.1833461 ± 0.0000011	MARIAM	82	CNTR $+$	HFS splitting
3.1833448 ± 0.0000029	CAMANI	78	CNTR $+$	See KLEMP 82
3.1833403 ± 0.0000044	CASPERSON	77	CNTR $+$	HFS splitting
3.1833402 ± 0.0000072	COHEN	73	RVUE	1973 CODATA value
3.1833467 ± 0.0000082	CROWE	72	CNTR $+$	Precession phase

¹⁰COHEN 87 (1986 CODATA) value was fitted using their own selection of the following data. Because their value is from a multiparameter fit, correlations with other quantities may be important and one cannot arrive at this result by any average of these data alone. μ^- DECAY MODES μ^+ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $e^- \bar{\nu}_e \nu_\mu$	$\approx 100\%$	
Γ_2 $e^- \bar{\nu}_e \nu_\mu \gamma$	[a] $(1.4 \pm 0.4)\%$	
Γ_3 $e^- \bar{\nu}_e \nu_\mu e^+ e^-$	[b] $(3.4 \pm 0.4) \times 10^{-5}$	
Lepton Family number (LF) violating modes		
Γ_4 $e^- \nu_e \bar{\nu}_\mu$	LF [c] < 1.2	90%
Γ_5 $e^- \gamma$	LF < 4.9	$\times 10^{-11}$ 90%
Γ_6 $e^- e^+ e^-$	LF < 1.0	$\times 10^{-12}$ 90%
Γ_7 $e^- 2\gamma$	LF < 7.2	$\times 10^{-11}$ 90%

[a] This only includes events with the γ energy > 10 MeV. Since the $e^- \bar{\nu}_e \nu_\mu$ and $e^- \bar{\nu}_e \nu_\mu \gamma$ modes cannot be clearly separated, we regard the latter mode as a subset of the former.

[b] See the Full Listings below for the energy limits used in this measurement.

[c] A test of additive vs. multiplicative lepton family number conservation.

 μ^- BRANCHING RATIOS

$\Gamma(e^- \bar{\nu}_e \nu_\mu \gamma) / \Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.014 ± 0.004		CRITTENDEN	61	CNTR γ KE > 10 MeV	

••• We do not use the following data for averages, fits, limits, etc. •••

0.0033 ± 0.0013	862	BOGART	67	CNTR γ KE > 14.5 MeV	
		CRITTENDEN	61	CNTR γ KE > 20 MeV	
	27	ASHKIN	59	CNTR	

$\Gamma(e^- \bar{\nu}_e \nu_\mu e^+ e^-) / \Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ
$3.4 \pm 0.2 \pm 0.3$	7443	¹¹ BERL	85	SPEC $+$	SINDRUM	

••• We do not use the following data for averages, fits, limits, etc. •••

2.2 ± 1.5	7	¹² CRITTENDEN	61	HLBC $+$	$E(e^+ e^-) > 10$ MeV	
2	1	¹³ GUREVICH	60	EMUL $+$		
1.5 ± 1.0	3	¹⁴ LEE	59	HBC $+$		

¹¹BERL 85 has transverse momentum cut $p_T > 17$ MeV/c. Systematic error was increased by us.¹²CRITTENDEN 61 count only those decays where total energy of either (e^+ , e^-) combination is > 10 MeV.¹³GUREVICH 60 Interpret their event as either virtual or real photon conversion. e^+ and e^- energies not measured.¹⁴In the three LEE 59 events, the sum of energies $E(e^+) + E(e^-) + E(e^+)$ was 51 MeV, 55 MeV, and 33 MeV.

$$\Gamma(e^- \nu_e \bar{\nu}_\mu) / \Gamma_{\text{total}}$$

Forbidden by the additive conservation law for lepton family number. A multiplicative law predicts this branching ratio to be 1/2. For a review see NEMETHY 81.

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ
< 0.012	90	FREEDMAN	93	CNTR $+$	ν oscillation search	
< 0.018	90	KRAKAUER	91B	CALO $+$		
< 0.05	90	¹⁵ BERGSM	83	CALO	$\bar{\nu}_\mu e^- \rightarrow \mu^- \bar{\nu}_e$	
< 0.09	90	JONKER	80	CALO	See BERGSM 83	
-0.001 ± 0.061		WILLIS	80	CNTR $+$		
0.13 ± 0.15		BLIETSCHAU	78	HLBC \pm	Avg. of 4 values	
< 0.25	90	EICHTEN	73	HLBC $+$		

¹⁵BERGSM 83 gives a limit on the inverse muon decay cross-section ratio $\sigma(\bar{\nu}_\mu e^- \rightarrow \mu^- \bar{\nu}_e) / \sigma(\nu_\mu e^- \rightarrow \mu^- \nu_e)$, which is essentially equivalent to $\Gamma(e^- \nu_e \bar{\nu}_\mu) / \Gamma_{\text{total}}$ for small values like that quoted.

$$\Gamma(e^- \gamma) / \Gamma_{\text{total}}$$

Forbidden by lepton family number conservation.

VALUE (units 10^{-11})	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ
< 4.9	90	BOLTON	88	CBOX $+$	LAMPF	
< 100	90	AZUELOS	83	CNTR $+$	TRIUMF	
< 17	90	KINNISON	82	SPEC $+$	LAMPF	
< 100	90	SCHAAF	80	ELEC $+$	SIN	

••• We do not use the following data for averages, fits, limits, etc. •••

$$\Gamma(e^- e^+ e^-) / \Gamma_{\text{total}}$$

Forbidden by lepton family number conservation.

VALUE (units 10^{-12})	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ
< 1.0	90	¹⁶ BELLGARDT	88	SPEC $+$	SINDRUM	
< 35	90	BARANOV	91	SPEC $+$	ARES	
< 36	90	BOLTON	88	CBOX $+$	LAMPF	
< 2.4	90	¹⁶ BERL	85	SPEC $+$	SINDRUM	
< 160	90	¹⁶ BERL	84	SPEC $+$	SINDRUM	
< 130	90	¹⁶ BOLTON	84	CNTR	LAMPF	

¹⁶These experiments assume a constant matrix element.

$$\Gamma(e^- 2\gamma) / \Gamma_{\text{total}}$$

Forbidden by lepton family number conservation.

VALUE (units 10^{-11})	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_7/Γ
< 7.2	90	BOLTON	88	CBOX $+$	LAMPF	
< 840	90	¹⁷ AZUELOS	83	CNTR $+$	TRIUMF	
< 5000	90	¹⁸ BOWMAN	78	CNTR	DEPOMMIER 77 data	

¹⁷AZUELOS 83 uses the phase space distribution of BOWMAN 78.¹⁸BOWMAN 78 assumes an interaction Lagrangian local on the scale of the inverse μ mass.LIMIT ON $\mu^- \rightarrow e^-$ CONVERSION

Forbidden by lepton family number conservation.

$\sigma(\mu^- 32S \rightarrow e^- 32S) / \sigma(\mu^- 32S \rightarrow \nu_\mu 32P^*)$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7 \times 10^{-11}$		90	BADERT...	80	STRC SIN
$< 4 \times 10^{-10}$		90	BADERT...	77	STRC SIN

••• We do not use the following data for averages, fits, limits, etc. •••

$\sigma(\mu^- \text{Cu} \rightarrow e^- \text{Cu}) / \sigma(\mu^- \text{Cu} \rightarrow \text{capture})$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-8}$		90	BRYMAN	72	SPEC

••• We do not use the following data for averages, fits, limits, etc. •••

$\sigma(\mu^- \text{Tl} \rightarrow e^- \text{Tl}) / \sigma(\mu^- \text{Tl} \rightarrow \text{capture})$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.3 \times 10^{-12}$		90	¹⁹ DOHMEN	93	SPEC SINDRUM II
$< 4.6 \times 10^{-12}$		90	AHMAD	88	TPC TRIUMF
$< 1.6 \times 10^{-11}$		90	BRYMAN	85	TPC TRIUMF

¹⁹DOHMEN 93 assumes $\mu^- \rightarrow e^-$ conversion leaves the nucleus in its ground state, a process enhanced by coherence and expected to dominate.

$\sigma(\mu^- \text{Pb} \rightarrow e^- \text{Pb}) / \sigma(\mu^- \text{Pb} \rightarrow \text{capture})$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.9 \times 10^{-10}$		90	AHMAD	88	TPC TRIUMF

••• We do not use the following data for averages, fits, limits, etc. •••

LIMIT ON $\mu^- \rightarrow e^+$ CONVERSION

Forbidden by total lepton number conservation.

 $\sigma(\mu^- 32\text{S} \rightarrow e^+ 32\text{S}^*) / \sigma(\mu^- 32\text{S} \rightarrow \nu_\mu 32\text{P}^*)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 9 \times 10^{-10}$	90	BADERT...	80	STRC SIN
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 1.5 \times 10^{-9}$	90	BADERT...	78	STRC SIN

 $\sigma(\mu^- 127\text{I} \rightarrow e^+ 127\text{Sb}^*) / \sigma(\mu^- 127\text{I} \rightarrow \text{anything})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3 \times 10^{-10}$	90	20 ABELA	80	CNTR Radiochemical tech.
20 ABELA 80 is upper limit for $\mu^- e^+$ conversion leading to particle-stable states of 127Sb . Limit for total conversion rate is higher by a factor less than 4 (G. Backenstoss, private communication).				

 $\sigma(\mu^- \text{Cu} \rightarrow e^+ \text{Co}) / \sigma(\mu^- \text{Cu} \rightarrow \nu_\mu \text{Ni})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 2.6 \times 10^{-8}$	90	BRYMAN	72	SPEC
$< 2.2 \times 10^{-7}$	90	CONFORTO	62	OSPK

 $\sigma(\mu^- \text{Tl} \rightarrow e^+ \text{Ca}) / \sigma(\mu^- \text{Tl} \rightarrow \text{capture})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.9 \times 10^{-11}$	90	21 DOHMEN	93	SPEC SINDRUM II
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$< 4.3 \times 10^{-12}$	90	22 DOHMEN	93	SPEC SINDRUM II
$< 1.7 \times 10^{-10}$	90	23 AHMAD	88	TPC TRIUMF

21 This DOHMEN 93 limit assumes a giant resonance excitation of the daughter Ca nucleus (mean energy and width both 20 MeV).

22 This DOHMEN 93 limit assumes the daughter Ca nucleus is left in the ground state. However, the probability of this is unknown.

23 Assuming a giant-resonance-excitation model.

LIMIT ON MUONIUM \rightarrow ANTIMUONIUM CONVERSION

Forbidden by lepton family number conservation.

 $R_g = G_C / G_F$ The effective Lagrangian for the $\mu^+ e^- \rightarrow \mu^- e^+$ conversion is assumed to be

$$\mathcal{L} = 2^{-1/2} G_C [\bar{\psi}_\mu \gamma_\lambda (1 - \gamma_5) \psi_e] [\bar{\psi}_\mu \gamma_\lambda (1 - \gamma_5) \psi_e] + \text{h.c.}$$

The experimental result is then an upper limit on G_C/G_F , where G_F is the Fermi coupling constant.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.13	90	GORDEEV	93	SPEC JINR phasotron
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 6.9	90	NI	93	CBOX LAMPF
< 0.16	90	MATTHIAS	91	SPEC LAMPF
< 0.29	90	HUBER	90B	CNTR TRIUMF
< 0.88	90	HUBER	88	CNTR See HUBER 90B
< 7.5	90	NI	87	CBOX See NI 93
< 20	95	BEER	86	CNTR TRIUMF
< 42	95	MARSHALL	82	CNTR

NOTE ON MUON DECAY PARAMETERS

(by W. Fetscher and H.-J. Gerber, ETH Zürich)

All measurements in direct muon decay, $\mu^- \rightarrow e^- + 2$ neutrals, and its inverse, $\nu_\mu + e^- \rightarrow \mu^- + \text{neutral}$, are successfully described by the “ $V-A$ interaction,” which is a particular case of a local, derivative-free, lepton-number-conserving, four-fermion interaction [1]. The matrix element is given below. The $V-A$ form and the nature of the neutrals (ν_μ and $\bar{\nu}_e$), and hence the doublet assignments $(\nu_e e^-)_L$ and $(\nu_\mu \mu^-)_L$, can be determined from experiments [2,3].

All results in direct muon decay (energy spectra, polarizations, and angular distributions) and in inverse muon decay (the reaction cross section) at energies well below $m_W c^2$ may be parametrized in terms of amplitudes $g_{e\mu}^\gamma$ and the Fermi coupling constant G_F , using the matrix element

$$\frac{4G_F}{\sqrt{2}} \sum_{\substack{\gamma=S,V,T \\ e,\mu=R,L}} g_{e\mu}^\gamma \langle \bar{e}_e | \Gamma^\gamma | (\nu_e)_n \rangle \langle (\bar{\nu}_\mu)_m | \Gamma_\gamma | \mu_\mu \rangle. \quad (1)$$

We use the notation of Fetscher *et al.* [2], who in turn use the sign conventions and definitions of Scheck [4]. Here $\gamma = S, V, T$ indicate a scalar, vector, or tensor interaction; and $\varepsilon, \mu = R, L$ indicate a right- or left-handed chirality of the electron or muon. The chiralities n and m of the ν_e and $\bar{\nu}_\mu$ are then determined by the values of γ, ε , and μ . The particles are represented by fields of definite chirality [5].

As shown by Langacker and London [6], explicit lepton-number nonconservation still leads to a matrix element equivalent to Eq. (1). They conclude that it is not possible, even in principle, to test lepton-number conservation in (leptonic) muon decay if the final neutrinos are massless and are not observed.

The ten complex amplitudes $g_{e\mu}^\gamma$ (g_{RR}^T and g_{LL}^T are identically zero) and G_F constitute 19 independent (real) parameters to be determined by experiment. The $V-A$ interaction corresponds to the single amplitude g_{LL}^V being unity and all the others being zero.

C. Jarlskog [7] has noted that certain experiments observing the decay electron are especially informative if they yield the $V-A$ values. Indeed, all (direct) muon decay experiments are compatible with an arbitrary mix of the scalar and vector amplitudes g_{LL}^S and g_{LL}^V —in the extreme, even with the purely scalar $g_{LL}^S = 2, g_{LL}^V = 0$. The decision in favor of $V-A$ comes from the quantitative observation of inverse muon decay, which would be forbidden for pure g_{LL}^S [2].

The differential decay probability to obtain an e^\pm with (reduced) energy between x and $x + dx$, emitted in the direction \hat{z} at an angle between θ and $\theta + d\theta$ with respect to the muon polarization vector \vec{P}_μ , and with its spin pointing in the arbitrary direction $\hat{\zeta}$, is given by

$$\begin{aligned} \frac{d^2\Gamma}{dx d\cos\theta} &= \frac{m_\mu}{4\pi^3} W_{e\mu}^4 G_F^2 \sqrt{x^2 - x_0^2} \\ &\times \left(F_{IS}(x) \pm P_\mu \cos\theta F_{AS}(x) \right) \\ &\times [1 + \vec{P}_e(x, \theta) \cdot \hat{\zeta}]. \end{aligned}$$

Here $W_{e\mu} = \max(E_e) = (m_\mu^2 + m_e^2)/2m_\mu$ is the maximum e^\pm energy, $x = E_e/W_{e\mu}$ is the reduced energy, and $x_0 = m_e/W_{e\mu} = 9.67 \times 10^{-3}$. The quantity $P_\mu = |\vec{P}_\mu| \cdot \hat{\zeta}$ has the significance of the direction in which a perfect polarization-sensitive electron detector would be most sensitive. The isotropic part of the spectrum, $F_{IS}(x)$, the anisotropic part, $F_{AS}(x)$, and the electron polarization, $\vec{P}_e(x, \theta)$, depend on bilinear combinations—called decay parameters—of the coupling constants $g_{e\mu}^\gamma$. Neglecting possible nonzero neutrino masses, we have, in terms of the decay parameters ρ, η, ξ, δ , etc.,

$$F_{IS}(x) = x(1-x) + \frac{2}{9}\rho(4x^2 - 3x - x_0^2) + \eta x_0(1-x)$$

$$\begin{aligned} F_{AS}(x) &= \frac{1}{3}\xi \sqrt{x^2 - x_0^2} \\ &\times \left[1 - x + \frac{2}{3}\delta(4x - 3 - (\sqrt{1 - x_0^2} - 1)) \right] \end{aligned}$$

$$\vec{P}_e(x, \theta) = P_{T_1} \hat{x} + P_{T_2} \hat{y} + P_L \hat{z}.$$

Lepton & Quark Full Listings

 μ

Here \hat{x} , \hat{y} , and \hat{z} are orthogonal unit vectors defined as follows:

\hat{z} is along the e momentum

$\hat{y} = [\hat{z} \times \vec{P}_\mu]/|[\hat{z} \times \vec{P}_\mu]|$ is transverse to the e momentum and perpendicular to the “decay plane”

$\hat{x} = \hat{y} \times \hat{z}$ is transverse to the e momentum and in the “decay plane.”

The components of \vec{P}_e then are given by

$$P_{T_1}(x, \theta) = P_\mu \sin \theta F_{T_1}(x) / \left(F_{IS}(x) \pm P_\mu \cos \theta F_{AS}(x) \right)$$

$$P_{T_2}(x, \theta) = P_\mu \sin \theta F_{T_2}(x) / \left(F_{IS}(x) \pm P_\mu \cos \theta F_{AS}(x) \right)$$

$$P_L(x, \theta) = \pm F_{IP}(x) + P_\mu \cos \theta \times F_{AP}(x) / \left(F_{IS}(x) \pm P_\mu \cos \theta F_{AS}(x) \right),$$

where

$$F_{T_1}(x) = \frac{1}{12} \left\{ -2 \left[\xi'' + 12 \left(\rho - \frac{3}{4} \right) \right] (1-x)x_0 - 3\eta(x^2 - x_0^2) + \eta''(-3x^2 + 4x - x_0^2) \right\}$$

$$F_{T_2}(x) = \frac{1}{3} \sqrt{x^2 - x_0^2} \left\{ 3 \frac{\alpha'}{A} (1-x) + 2 \frac{\beta'}{A} \sqrt{1-x_0^2} \right\}$$

$$F_{IP}(x) = \frac{1}{54} \sqrt{x^2 - x_0^2} \left\{ 9\xi'(-2x + 2 + \sqrt{1-x_0^2}) + 4\xi(\delta - \frac{3}{4})(4x - 4 + \sqrt{1-x_0^2}) \right\}$$

$$F_{AP}(x) = \frac{1}{6} \left\{ \xi''(2x^2 - x - x_0^2) + 4\left(\rho - \frac{3}{4}\right)(4x^2 - 3x - x_0^2) + 2\eta''(1-x)x_0 \right\}.$$

For the experimental values of the decay parameters ρ , ξ , ξ' , ξ'' , δ , η , η'' , α/A , β/A , α'/A , β'/A , which are not all independent, see the Data Listings below. Experiments in the past have also been analyzed using the parameters a , b , c , a' , b' , c' , α/A , β/A , α'/A , β'/A (and $\eta = (\alpha - 2\beta)/2A$), as defined by Kinoshita and Sirlin [8]. They serve as a model-independent summary of all possible measurements on the decay electron (see Listings below). The relations between the two sets of parameters are

$$\rho - \frac{3}{4} = \frac{3}{4}(-a + 2c)/A,$$

$$\eta = (\alpha - 2\beta)/A,$$

$$\eta'' = (3\alpha + 2\beta)/A,$$

$$\delta - \frac{3}{4} = \frac{9}{4} \frac{(a' - 2c')/A}{1 - [a + 3a' + 4(b + b') + 6c - 14c']/A},$$

$$1 - \xi \frac{\delta}{\rho} = 4 \frac{[(b + b') + 2(c - c')]/A}{1 - (a - 2c)/A},$$

$$1 - \xi' = [(a + a') + 4(b + b') + 6(c + c')]/A,$$

$$1 - \xi'' = (-2a + 20c)/A,$$

where

$$A = a + 4b + 6c.$$

The relations to the coupling constants are:

$$a = 16(|g_{RL}^V|^2 + |g_{LR}^V|^2) + |g_{RL}^S + 6g_{RL}^T|^2 + |g_{LR}^S + 6g_{LR}^T|^2,$$

$$a' = 16(|g_{RL}^V|^2 - |g_{LR}^V|^2) + |g_{RL}^S + 6g_{RL}^T|^2 - |g_{LR}^S + 6g_{LR}^T|^2,$$

$$\alpha = 8 \operatorname{Re} \left[g_{RL}^V (g_{LR}^S + 6g_{LR}^T)^* + g_{LR}^V (g_{RL}^S + 6g_{RL}^T)^* \right],$$

$$\alpha' = 8 \operatorname{Im} \left[-g_{LR}^V (g_{RL}^S + 6g_{RL}^T)^* - g_{RL}^V (g_{LR}^S + 6g_{LR}^T)^* \right],$$

$$b = 4(|g_{RR}^V|^2 + |g_{LL}^V|^2) + |g_{RR}^S|^2 + |g_{LL}^S|^2,$$

$$b' = 4(|g_{RR}^V|^2 - |g_{LL}^V|^2) + |g_{RR}^S|^2 - |g_{LL}^S|^2,$$

$$\beta = -4 \operatorname{Re} \left[g_{RR}^V (g_{LL}^S)^* + g_{LL}^V (g_{RR}^S)^* \right],$$

$$\beta' = 4 \operatorname{Im} \left[g_{RR}^V (g_{LL}^S)^* - g_{LL}^V (g_{RR}^S)^* \right],$$

$$c = \frac{1}{2} \left[|g_{RL}^S - 2g_{RL}^T|^2 + |g_{LR}^S - 2g_{LR}^T|^2 \right],$$

$$c' = \frac{1}{2} \left[|g_{RL}^S - 2g_{RL}^T|^2 - |g_{LR}^S - 2g_{LR}^T|^2 \right].$$

If also the electron mass is neglected, the energy and angular distribution of the electron in the rest frame of a polarized muon (μ^\mp) is given by the Michel spectrum:

$$d^2\Gamma \sim \left\{ 3(1-x) + \frac{2\rho}{3}(4x-3) \mp \xi \cos \theta \left[1-x + \frac{2\delta}{3}(4x-3) \right] \right\} x^2 dx d(\cos \theta).$$

Here θ is the angle between the electron momentum and the muon spin, and $x \equiv 2E_e/m_\mu$. For pure $V-A$ coupling, we obtain $\rho = \xi\delta = 3/4$, $\xi = 1$, and the differential decay rate is

$$d^2\Gamma = \frac{G_F^2 m_\mu^5}{192\pi^3} [3 - 2x \pm \cos \theta(1-2x)] x^2 dx d(\cos \theta).$$

Here the coefficient in front of the square bracket is the total decay rate.

In order to determine the amplitudes $g_{e\mu}^\gamma$ uniquely from experiment, Fetscher *et al.* [2] introduced four probabilities $Q_{e\mu}(\varepsilon, \mu = R, L)$ for the decay of a μ -handed muon into an ε -handed electron and showed that there exist upper bounds on Q_{RR} , Q_{LR} , and Q_{RL} , and a lower bound on Q_{LL} . These probabilities are given in terms of the $g_{e\mu}^\gamma$'s by

$$Q_{e\mu} = \frac{1}{4} |g_{e\mu}^S|^2 + |g_{e\mu}^V|^2 + 3(1 - \delta_{e\mu}) |g_{e\mu}^T|^2, \quad (2)$$

where $\delta_{e\mu} = 1$ for $\varepsilon = \mu$ and $\delta_{e\mu} = 0$ for $\varepsilon \neq \mu$. They are related to the parameters a , b , c , a' , b' , and c' by

$$Q_{RR} = 2(b + b')/A,$$

$$Q_{LR} = [(a - a') + 6(c - c')]/2A,$$

$$Q_{RL} = [(a + a') + 6(c + c')]/2A,$$

$$Q_{LL} = 2(b - b')/A,$$

with $A = 16$. In the pure $V-A$ theory, $Q_{LL} = 1$ and the others are zero.

Since the upper bounds on Q_{RR} , Q_{LR} , and Q_{RL} are found to be small, and since the helicity of the ν_μ in pion decay is known from experiment [9,10] to very high precision to be -1 [11], the cross section S of *inverse* muon decay, normalized to the $V-A$ value, yields [2]

$$|g_{LL}^S|^2 \leq 4(1 - S) \quad (3)$$

and

$$|g_{LL}^V|^2 = S. \quad (4)$$

Thus the Standard Model assumption of a pure $V-A$ leptonic charged weak interaction for e and μ is confirmed (within errors) by experiments at energies far below the mass of the W^\pm : Eq. (4) gives a lower limit for $V-A$, and Eqs. (2) and (3) give upper limits for the other four-fermion interactions. The existence of such upper limits may also be seen from $Q_{RR} + Q_{RL} = (1 - \xi')/2$ and $Q_{RR} + Q_{LR} = \frac{1}{2}(1 + \xi/3 - 16\xi\delta/9)$. Table 1 gives the current experimental limits on the magnitudes of the $g_{e\mu}^i$'s.

Table 1. Ninety-percent confidence level experimental limits for the coupling constants $g_{e\mu}^i$. The limits on $|g_{LL}^S|$ and $|g_{LL}^V|$ are from Ref. 12, and the others are from Ref. 13. The experimental uncertainty on the muon polarization in pion decay is included.

$ g_{RR}^S < 0.066$	$ g_{RR}^V < 0.033$	$ g_{RR}^T \equiv 0$
$ g_{LR}^S < 0.125$	$ g_{LR}^V < 0.060$	$ g_{LR}^T < 0.036$
$ g_{RL}^S < 0.424$	$ g_{RL}^V < 0.110$	$ g_{RL}^T < 0.122$
$ g_{LL}^S < 0.55$	$ g_{LL}^V > 0.96$	$ g_{LL}^T \equiv 0$

Limits on the "charge retention" coordinates, as used in the older literature (*e.g.*, Ref. 14), are given by Burkard *et al.* [15].

References

- L. Michel, Proc. Phys. Soc. **A63**, 514 (1950).
- W. Fetscher, H.-J. Gerber, and K.F. Johnson, Phys. Lett. **B173**, 102 (1986).
- P. Langacker, Comm. Nucl. Part. Phys. **19**, 1 (1989).
- F. Scheck, in *Leptons, Hadrons, and Nuclei* (North Holland, Amsterdam, 1983).
- K. Mursula and F. Scheck, Nucl. Phys. **B253**, 189 (1985).
- P. Langacker and D. London, Phys. Rev. **D39**, 266 (1989).
- C. Jarlskog, Nucl. Phys. **75**, 659 (1966).
- T. Kinoshita and A. Sirlin, Phys. Rev. **108**, 844 (1957).
- A. Jodidio *et al.*, Phys. Rev. **D34**, 1967 (1986);
A. Jodidio *et al.*, Phys. Rev. **D37**, 237 (1988).
- L.Ph. Roesch *et al.*, Helv. Phys. Acta **55**, 74 (1982).
- W. Fetscher, Phys. Lett. **140B**, 117 (1984).
- S.R. Mishra *et al.*, Phys. Lett. **B252**, 170 (1990);
S.R. Mishra, private communication;
See also D. Geiregat *et al.*, Phys. Lett. **B247**, 131 (1990).
- B. Balke *et al.*, Phys. Rev. **D37**, 587 (1988).
- S.E. Derenzo, Phys. Rev. **181**, 1854 (1969).
- H. Burkard *et al.*, Phys. Lett. **160B**, 343 (1985).

μ DECAY PARAMETERS

ρ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.7518 ± 0.0026		DERENZO	69	RVUE	
0.762 ± 0.008	170k	²⁴ FRYBERGER	68	ASPK +	25–53 MeV e^+
0.760 ± 0.009	280k	²⁴ SHERWOOD	67	ASPK +	25–53 MeV e^+
0.7503 ± 0.0026	800k	²⁴ PEOPLES	66	ASPK +	20–53 MeV e^+

²⁴ η constrained = 0. These values incorporated into a two parameter fit to ρ and η by DERENZO 69.

η PARAMETER

($V-A$) theory predicts $\eta = 0$.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.007 ± 0.013 OUR AVERAGE					
-0.007 ± 0.013	5.3M	²⁵ BURKARD	85B	FIT +	9–53 MeV e^+
-0.12 ± 0.21	6346	DERENZO	69	HBC +	1.6–6.8 MeV e^+
0.011 ± 0.081 ± 0.026	5.3M	BURKARD	85B	CNTR +	9–53 MeV e^+
-0.7 ± 0.5	170k	²⁷ FRYBERGER	68	ASPK +	25–53 MeV e^+
-0.7 ± 0.6	280k	²⁷ SHERWOOD	67	ASPK +	25–53 MeV e^+
0.05 ± 0.5	800k	²⁷ PEOPLES	66	ASPK +	20–53 MeV e^+
-2.0 ± 0.9	9213	²⁸ PLANO	60	HBC +	Whole spectrum

²⁵Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

²⁷ $\alpha = \alpha' = 0$ assumed.

²⁷ ρ constrained = 0.75.

²⁸Two parameter fit to ρ and η ; PLANO 60 discounts value for η .

δ PARAMETER

($V-A$) theory predicts $\delta = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.7486 ± 0.0026 ± 0.0028		²⁹ BALKE	88	SPEC +	Surface μ^+ 's
0.752 ± 0.009	490k	³⁰ VOSSLER	69		
0.782 ± 0.031		FRYBERGER	68	ASPK +	25–53 MeV e^+
0.78 ± 0.05	8354	KRUGER	61		
		PLANO	60	HBC +	Whole spectrum

²⁹BALKE 88 uses $\rho = 0.752 \pm 0.003$.

³⁰VOSSLER 69 has measured the asymmetry below 10 MeV. See comments about radiative corrections in VOSSLER 69.

$(\xi \text{ PARAMETER}) \times (\mu \text{ LONGITUDINAL POLARIZATION})$

($V-A$) theory predicts $\xi = 1$, longitudinal polarization = 1.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.0027 ± 0.0079 ± 0.0030		BELTRAMI	87	CNTR	SIN, π decay in flight
1.0013 ± 0.0030 ± 0.0053		³¹ IMAZATO	92	SPEC +	$K^+ \rightarrow \mu^+ \nu_\mu$
0.975 ± 0.015		AKHMANOV	68	EMUL	140 kG
0.975 ± 0.030	66k	GUREVICH	64	EMUL	See AKHMANOV 68
0.903 ± 0.027		³² ALI-ZADE	61	EMUL +	27 kG
0.93 ± 0.06	8354	PLANO	60	HBC +	8.8 kG
0.97 ± 0.05	9k	BARDON	59	CNTR	Bromofom target

³¹The corresponding 90% confidence limit from IMAZATO 92 is $|\xi P_\mu| > 0.990$. This measurement is of K^+ decay, not π^+ decay, so we do not include it in an average, nor do we yet set up a separate data block for K results.

³²Depolarization by medium not known sufficiently well.

$\xi \times (\mu \text{ LONGITUDINAL POLARIZATION}) \times \delta / \rho$

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
> 0.99682	90	³³ JODIDIO	86	SPEC +	TRIUMF
> 0.9966	90	³⁴ STOKER	85	SPEC +	μ -spin rotation
> 0.9959	90	CARR	83	SPEC +	11 kG

³³JODIDIO 86 includes data from CARR 83 and STOKER 85. The value here is from the erratum.

³⁴STOKER 85 find $(\xi P_\mu \delta / \rho) > 0.9955$ and > 0.9966 , where the first limit is from new μ spin-rotation data and the second is from combination with CARR 83 data. In $V-A$ theory, $(\delta / \rho) = 1.0$.

$\xi' = \text{LONGITUDINAL POLARIZATION OF } e^+$

($V-A$) theory predicts the longitudinal polarization = ± 1 for e^\pm , respectively. We have flipped the sign for e^- so our programs can average.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.00 ± 0.04 OUR AVERAGE					
0.998 ± 0.045	1M	BURKARD	85	CNTR +	Bhabha + annhil
0.89 ± 0.28	29k	SCHWARTZ	67	OSPK -	Moller scattering
0.94 ± 0.38		BLOOM	64	CNTR +	Brems. transmiss.
1.04 ± 0.18		DUCLLOS	64	CNTR +	Bhabha scattering
1.05 ± 0.30		BUHLER	63	CNTR +	Annihilation

Lepton & Quark Full Listings

 μ ξ'' PARAMETER

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.65 ± 0.36	326k	35 BURKARD	85 CNTR	+	Bhabha + annihl

35 BURKARD 85 measure $(\xi'' - \xi'')/\xi$ and ξ' and set $\xi = 1$.

TRANSVERSE e^+ POLARIZATION IN PLANE OF μ SPIN, e^+ MOMENTUM

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$0.016 \pm 0.021 \pm 0.01$	5.3M	BURKARD	85B CNTR	+	Annihil 9-53 MeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

TRANSVERSE e^+ POLARIZATION NORMAL TO PLANE OF μ SPIN, e^+ MOMENTUM

Zero if T invariance holds.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$0.007 \pm 0.022 \pm 0.007$	5.3M	BURKARD	85B CNTR	+	Annihil 9-53 MeV

 α/A

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.4 ± 4.3		36 BURKARD	85B FIT		

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$15 \pm 50 \pm 14$	5.3M	BURKARD	85B CNTR	+	9-53 MeV e^+

36 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 α'/A

Zero if T invariance holds.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.2 ± 4.3		37 BURKARD	85B FIT		

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$-47 \pm 50 \pm 14$	5.3M	38 BURKARD	85B CNTR	+	9-53 MeV e^+

37 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

38 BURKARD 85b measure e^+ polarizations P_{T1} and P_{T2} versus e^+ energy.

 β/A

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3.9 ± 6.2		39 BURKARD	85B FIT		

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$2 \pm 17 \pm 6$	5.3M	BURKARD	85B CNTR	+	9-53 MeV e^+

39 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 β'/A

Zero if T invariance holds.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.5 ± 6.3		40 BURKARD	85B FIT		

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$17 \pm 17 \pm 6$	5.3M	41 BURKARD	85B CNTR	+	9-53 MeV e^+

40 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

41 BURKARD 85b measure e^+ polarizations P_{T1} and P_{T2} versus e^+ energy.

 a/A

This comes from an alternative parameterization to that used in the Summary Table (see the "Note on Muon Decay Parameters" above).

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN
< 15.9	90	42 BURKARD	85B FIT

42 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 d/A

This comes from an alternative parameterization to that used in the Summary Table (see the "Note on Muon Decay Parameters" above).

VALUE (units 10^{-3})	DOCUMENT ID	TECN
5.3 ± 4.1	43 BURKARD	85B FIT

43 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 $(b+b)/A$

This comes from an alternative parameterization to that used in the Summary Table (see the "Note on Muon Decay Parameters" above).

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN
< 1.04	90	44 BURKARD	85B FIT

44 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 c/A

This comes from an alternative parameterization to that used in the Summary Table (see the "Note on Muon Decay Parameters" above).

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN
< 6.4	90	45 BURKARD	85B FIT

45 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 c'/A

This comes from an alternative parameterization to that used in the Summary Table (see the "Note on Muon Decay Parameters" above).

VALUE (units 10^{-3})	DOCUMENT ID	TECN
3.5 ± 2.0	46 BURKARD	85B FIT

46 Global fit to all measured parameters. Correlation coefficients are given in BURKARD 85b.

 η PARAMETER

($V-A$) theory predicts $\eta = 0$. η affects spectrum of radiative muon decay.

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.02 ± 0.08	OUR AVERAGE			
-0.014 ± 0.090	EICHENBER...	84	ELEC	+ ρ free
$+0.09 \pm 0.14$	BOGART	67	CNTR	+

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
-0.035 ± 0.098	EICHENBER...	84	ELEC	+ $\rho=0.75$ assumed

 μ REFERENCES

DOHMEN	93	PL B317 631	+Groth, Heer+	(PSI SINDRUM-II Collab.)
FREEDMAN	93	PR D47 811	+Fujikawa, Napolitano, Nelson+	(LAMPF E645 Collab.)
GORDEEV	93	JETPL 57 270	+Savchenko, Abazov+	(PNPI, JINR)
		Translated from ZETFP 57 262.		
NI	93	PR D48 1976	+Arnold, Chmely+	(LAMPF Crystal-Box Collab.)
IMAZATO	92	PRL 69 877	+Kawashima, Tanaka+	(KEK, INUS, TOKY, TOKMS)
BARANOV	91	SJNP 53 302	+Vanko, Glazov, Evtkhovich+	(JINR)
		Translated from YAF 53 1302.		
KRAKAUER	91B	PL B263 534	+Talaga, Allen, Chen, Doe+	(UMD, UCI, LANL)
MATTHIAS	91	PRL 66 2716	+Ahn+	(YALE, HEIDP, WILL, GSI, VILL, BNL)
		Also		
	91B	PRL 67 932 erratum	+Matthias, Ahn+	(YALE, HEIDP, WILL, GSI, VILL, BNL)
HUBER	90B	PR D41 2709	+ (WYOM, VICT, ARIZ, ROCH, TRIU, SFRA, BRCO)	
AHMAD	88	PR D38 2102	+Azuels+ (TRIU, VICT, VPI, BRCO, MONT, CNRC)	
	87	PR 59 870	+Ahmad+ (TRIU, VPI, VICT, BRCO, MONT, CNRC)	
BALKE	88	PR D37 587	+Gidal, Jodidio+	(LBL, UCB, COLO, NWES, TRIU)
BELLEGARDT	88	NP B299 1	+Otter, Eichler+	(SINDRUM Collab.)
BOLTON	88	PR D38 2077	+Cooper, Frank, Hallin+	(LANL, STAN, CHIC, TEMP)
	86	PRL 56 2461	+Bolton, Bowman, Cooper+	(LANL, STAN, CHIC, TEMP)
	86	PRL 57 3241	+Grosnick, Wright, Bolton+	(CHIC, LANL, STAN, TEMP)
HUBER	88	PRL 61 2189	+Beer+ (WYOM, VICT, ARIZ, ROCH, TRIU, BRCO)	
BELTRAMI	87	PL B194 326	+Burard, Von Dinkelge+	(ETH, SIM, MANZ)
COHEN	87	RMP 59 1121	+Taylor	(RISC, NBS)
NI	87	PR 59 2716	+Arnold, Chmely+ (YALE, LANL, WILL, MISS, HEIDP)	
BEER	86	PRL 57 671	+Marshall, Mason+	(VICT, TRIU, WYOM)
BELTRAMI	86	NP A451 679	+Aas, Beer, Dechambrier, Goudsmit+	(ETH, FRIB)
JODIDIO	86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+	(LBL, NWES, TRIU)
	88	PR D37 237 erratum	+Jodidio, Balke, Carr+	(LBL, NWES, TRIU)
BERTL	85	PRL 56 465	+Egli, Eichler+	(SINDRUM Collab.)
BRYMAN	85	PL 150B 242	+ (TRIU, CNRC, BRCO, LANL, CHIC, CARL+)	
BURKARD	85	PL 160B 343	+Corriveau, Egger+	(ETH, SIN, MANZ)
	85B	PL 160B 343	+Corriveau, Egger+	(ETH, SIN, MANZ)
	81B	PR D24 2004	+Corriveau, Egger, Fetscher+	(ETH, SIN, MANZ)
	83B	PL 129B 260	+Corriveau, Egger, Fetscher+	(ETH, SIN, MANZ)
HUGHES	85	CNPP 14 341	+Kinoshita	(YALE, CORN)
STOKER	85	PR 54 1887	+Balke, Carr, Gidal+	(LBL, NWES, TRIU)
BAR DIN	84	PL 123B 135	+Duclos, Magnon+	(SACL, CERN, GNA, FIRZ)
BERTL	84	PL 140B 299	+Eichler, Felawuk+	(SINDRUM Collab.)
BOLTON	84	PR 53 1415	+Bowman, Carlini+	(LANL, CHIC, STAN, TEMP)
EICHENBER...	84	NP A412 523	+Eichenberger, Engfer, Vanderschaff	(ZURI)
GIOVANETTI	84	PR D29 343	+Dey, Eckhauser, Hart+	(WILL)
KINOSHITA	84	PRL 52 717	+Nizic, Okamoto	(CORN)
AZUELOS	83	PRL 51 164	+Depommier, Leroy, Martin+	(MONT, TRIU, BRCO)
	77	PR 39 1113	+Depommier+ (MONT, BRCO, TRIU, VICT, MELB)	
BERGSMAN	83	PL 122B 465	+Dorenbosch, Jonker+	(CHARM Collab.)
CARR	83	PR 51 627	+Gidal, Gobbi, Jodidio, Oram+	(LBL, NWES, TRIU)
KINNISON	82	PR D25 2846	+Anderson, Matis, Wright+	(EFI, STAN, LANL)
	79	PR 42 556	+Bowman, Cooper, Hamm+	(LASL, EFI, STAN)
KLEMPPT	82	PR D25 652	+Schulze, Wolf, Camani, Gyga+	(MANZ, ETH)
MARIAM	82	PR 49 993	+Beer, Bolton, Egan, Gardner+	(YALE, HEIDH, BERN)
MARSHALL	82	PR D25 1174	+Warren, Oram, Kiefl	(BRCO)
COMBLEY	81	PRPL 68 93	+Farley, Picasso	(SHEF, RMCS, CERN)
NEMETHY	81	CNPP 10 147	+Hughes	(LBL, YALE)
ABELA	80	PL 95B 318	+Backenstoss, Simons, Wuest+	(BASL, KARLK, KARLE)
BADERT...	80	LNC 28 401	+Badertscher, Borer, Czapek, Flueckiger+	(BERN)
	82	NP A377 406	+Badertscher, Borer, Czapek, Flueckiger+	(BERN)
JONKER	80	PL 93B 203	+Panman, Udo, Allaby+	(CHARM Collab.)
SCHAAP	80	NP A340 249	+Engfer, Povel, Dey+	(ZURI, ETH, SIN)
	79	NP B50 1	+Povel, Dey, Walter+, Pfeiffer+	(ZURI, ETH, SIN)
WILLIS	80	PR 44 522	+Hughes+ (YALE, LBL, LASL, SACL, SIN, CNRC+)	
	80B	PR 45 1370	+Willis+ (YALE, LBL, LASL, SACL, SIN, CNRC+)	
BAILEY	79	NP B150 1	+Picasso	(CERN, DARE, MANZ)
FARLEY	79	ARNPS 29 243	+Picasso	(RMCS, CERN)
BADERT...	78	PL 79B 371	+Badertscher, Borer, Czapek, Flueckiger+	(BERN)
BAILEY	78	JPG 4 345	+ (DARE, BERN, SHEF, MANZ, RMCS, CERN, BIRM)	
	77	NP B150 1	+Bailey	(CERN, DARE, MANZ)
BLIETSCHAU	78	NP B133 205	+Dedea, Hasert, Krenz+	(CERN, gamelle Collab.)
BOWMAN	78	PRL 41 442	+Cheng, Li, Matis	(LASL, IAS, CMU, EFI)
CAMANI	78	PL 77B 326	+Gyga, Klemppt, Schenck, Schulze+	(ETH, MANZ)
BADERT...	77	PR 39 1385	+Badertscher, Borer, Czapek, Flueckiger+	(BERN)
BAILEY	77	PL 67B 225	+ (CERN Muon Storage Ring Collab.)	
	77C	PL 68B 191	+Bailey+	(CERN, DARE, BERN, SHEF, MANZ+)
	75	PL 55B 120	+Bailey+	(CERN Muon Storage Ring Collab., BIRM)
CALMET	77	RMP 49 21	+Narison, Perrottet+	(CNRS)
CASPERSON	77	PR 38 956	+Crane+	(BERN, HEIDH, LASL, WYOM, YALE)
DEPOMMIER	77	PR 39 1113	+ (MONT, BRCO, TRIU, VICT, MELB)	
BALANDIN	74	JETP 40 811	+Grebenyuk, Zinov, Konin, Ponomarev	(JINR)

Translated from ZETFP 67 1631.

COHEN	73	JPCRD 2 663	+Taylor	(RISC, NBS)
DUCLÓS	73	PL 478 491	+Magnon, Picard	(SACL)
EICHTEN	73	PL 468 281	+Deden, Hasser, Krenz+	(Gargamelie Collab.)
BRYMAN	72	PRL 28 1469	+Blecher, Gotow, Powers	(VPI)
CROWE	72	PR D5 2145	+Hague, Rothberg, Schenck+	(LBL, WASH)
CRANE	71	PRL 27 474	+Casperson, Crane, Egan, Hughes+	(YALE)
DERENZO	69	PR 181 1854		(EFI)
VOSSLER	69	NC 63A 423		(EFI)
AKHMANOV	68	SJNP 6 230	+Gurevich, Dobretsov, Makarina+	(KIAE)
BAILEY	68	PL 28B 287	+Bartl, VonBochmann, Brown, Farley+	(CERN)
Also	72	NC 9A 369	+Bailey, Bartl, VonBochmann, Brown+	(CERN)
FRYBERGER	68	PR 166 1379		(EFI)
BOGART	67	PR 156 1405	+Dicapua, Nemethy, Streizoff	(COLU)
SCHWARTZ	67	PR 162 1306		(EFI)
SHERWOOD	67	PR 156 1475		(EFI)
PEOPLES	66	Nevis 147 unpub.		(COLU)
BLOOM	64	PL 8 87	+Dick, Feuvrais, Henry, Macq, Spighel	(CERN)
DUCLÓS	64	PL 9 62	+Heintze, DeRujula, Soergel	(CERN)
GUREVICH	64	PL 11 185	+Makarina+	(KIAE)
BUHLER	63	PL 7 368	+Cabibbo, Fidecaro, Massam, Muller+	(CERN)
MEYER	63	PR 132 2693	+Anderson, Bleser, Lederman+	(COLU)
CHARPAK	62	PL 1 16	+Farley, Garwin+	(CERN)
CONFORTO	62	NC 261	+Conversi, Dilella+	(CERN)
ALI-ZADE	61	JETP 13 313	+Gurevich, Nikolski	(INFN, ROMA, CERN)
		Translated from ZETF 40 452.		
CRITTENDEN	61	PR 121 1823	+Walker, Ballam	(WISC, MSU)
KRUGER	61	UCRL 9322 unpub.		(LRL)
GUREVICH	60	JETP 10 225	+Nikolski, Surkova	(ITEP)
		Translated from ZETF 37 318.		
PLANO	60	PR 119 1400		(COLU)
ASHKIN	59	NC 14 1266	+Fazzini, Fidecaro, Lipman, Morrison+	(CERN)
BARDON	59	PRL 2 56	+Berley, Lederman	(COLU)
LEE	59	PRL 3 55	+Samios	(COLU)

T

$$J = \frac{1}{2}$$

τ discovery paper was PERL 75. $e^+e^- \rightarrow \tau^+\tau^-$ cross-section threshold behavior and magnitude are consistent with pointlike spin-1/2 Dirac particle. BRANDELIK 78 ruled out pointlike spin-0 or spin-1 particle. FELDMAN 78 ruled out $J = 3/2$. KIRKBY 79 also ruled out $J = \text{integer}$, $J = 3/2$.

NOTE ON PROBLEMS IN THE τ -DECAY DATA

(by K.G. Hayes, Hillsdale College)

Problems remain in the experimental data for τ branching ratios. The first evidence that something was wrong was noticed in 1984 in what became known as the "1-prong deficit" [1,2]: If world averages were considered and theoretical predictions were used to limit certain poorly measured modes, then the measured inclusive 1-prong topological branching ratio was significantly larger than the sum of branching ratios of exclusive 1-prong decay modes. Reviews of this problem have appeared in these Listings starting with the 1988 edition. At the time of the 1992 edition, significant discrepancies had appeared between some recent and older measurements of the branching ratios for several modes, most notably $B(h^-2\pi^0\nu_\tau)$ and $B(h^-h^-h^+\nu_\tau)$.

Since the 1992 edition, the situation has changed in two significant ways. The long-standing "tau decay puzzle" has been resolved. If one assumes unitarity, then the leptonic branching ratios ($B_e \equiv B(e^-\bar{\nu}_e\nu_\tau)$ or $B_\mu \equiv B(\mu^-\bar{\nu}_\mu\nu_\tau)$) can be predicted from masses and lifetimes of the muon and tau. Theory and prediction have differed at the 1.5 to 2 standard deviation level since 1986. Since the last edition, new precise measurements have lowered the world averages for the τ mass and lifetime by 1.9 and 1.7 standard deviations, respectively. Using our current world averages for these quantities, the predicted electron branching ratio [3] is $18.11 \pm 0.19\%$, while the world average for $B_e = 17.90 \pm 0.17\%$. This agreement provides strong evidence that the measured leptonic branching ratios of the τ are accurate, and argues against the existence of widespread systematic problems in the various strategies used to derive branching ratios from the observed numbers of $\tau^- \rightarrow e^-\bar{\nu}_e\nu_\tau$ and $\tau^- \rightarrow \mu^-\bar{\nu}_\mu\nu_\tau$ decays.

The second significant change has resulted from the publication [4,5,6] of precise new branching ratio measurements of τ decay modes of the form $\tau^- \rightarrow h^- + (\geq)n\pi^0\nu_\tau$. The CLEO collaboration [5,6] has reported values for $n = 1, 2, 3$, and 4, based upon reconstructed $\pi^0 \rightarrow \gamma\gamma$ decays for all π^0 's. These new measurements have increased the internal inconsistency of the world data for these modes. The $B(h^-3\pi^0\nu_\tau)$ and $B(h^-4\pi^0\nu_\tau)$ measurements have allowed the replacement of our fit mode $\tau^- \rightarrow h^- \geq 3\pi^0\nu_\tau$ by the two new modes $\tau^- \rightarrow h^-3\pi^0\nu_\tau$ and $\tau^- \rightarrow h^-4\pi^0\nu_\tau$ with the assumption, valid at the present level of precision, that $B(h^- \geq 5\pi^0\nu_\tau)$ is small enough to ignore. A comparison between the current and the 1992 branching ratio fit values for fit modes containing one charged hadron and one or more π^0 's is presented in Table 1. The error scale factors from the fit, which are also listed, are all just under two. It is clear that significant discrepancies exist between experimental measurements of these modes.

Table 1: Fit branching ratios (%) and scale factors for τ decay modes containing one charged hadron and one or more π^0 's.

Mode	1992 Fit	Scale factor	1994 Fit	Scale factor
$h^-\pi^0\nu_\tau$	24.4 ± 0.6	1.1	25.7 ± 0.4	1.7
$h^-2\pi^0\nu_\tau$	10.3 ± 0.9	1.7	9.6 ± 0.4	1.5
$h^-3\pi^0\nu_\tau$			1.28 ± 0.24	1.7
$h^-4\pi^0\nu_\tau$			0.19 ± 0.12 -0.10	1.7
$h^- \geq 3\pi^0\nu_\tau$	2.7 ± 0.9	1.9	1.48 ± 0.26	1.7

The new measurements have a significant impact on the constrained fit. The fit branching ratios cannot exhibit a problem such as the 1-prong deficit, since the fit 1-prong topological branching ratio is defined to be the sum of branching ratios of exclusive 1-prong modes. Since the set of fit modes sum exactly to one, the fact that the world averages exhibit a 1-prong deficit means that the fit values for some 1-prong modes must be larger than the corresponding average values; many examples are to be found in the present Listings. How much the constraint will increase the fit values above their corresponding averages depends on the errors. In the last two editions, $B(h^-2\pi^0\nu_\tau)$ and $B(h^- \geq 3\pi^0\nu_\tau)$ had relatively large errors, and the fit could account for much of the deficit by increasing these modes with only a small increase in the fit χ^2 . For example, the 1992 fit had a χ^2 of 91.8 for 91 degrees of freedom, while the current fit has a χ^2 of 134.6 for 105 degrees of freedom. The large χ^2 is not caused by one or two outlying measurements; the largest single contribution to the χ^2 is 7.0. Table 2 gives the average χ^2 contribution per datum* grouped by decay mode in various ways: all modes used in the fit, modes containing one prong and no neutrals, modes containing one prong plus $\geq 1\pi^0$, and all modes except 1-prong exclusive modes. It is clear that the 1-prong plus $\geq 1\pi^0$ modes contribute the largest amount to the χ^2 .

Lepton & Quark Full Listings

 τ **Table 2:** Average fit χ^2 contribution of τ branching ratio measurements grouped by decay mode category.

Category	Data entries	$\langle\chi^2\rangle$ per datum
All modes	116	1.17
1-prong 0 neutrals	51	0.74
1-prong ≥ 1 neutrals	28	1.87
Not 1-prong modes	37	1.23

The fact that the fit has to spread the branching fraction corresponding to the 1-prong deficit between many relatively well-measured basis modes means that the *average* values for some branching fractions (*i.e.*, B_e and B_μ) may be slightly more accurate than the *fit* values. In this edition, the method of calculating averages has been improved. In previous editions, individual branching fraction measurements were either used for both the fit and averages, or they were excluded from both the fit and averages. Many data must be excluded from the fit because they are highly correlated to other data which are used in the fit. Thus, many data were not used in averages even though there was no other reason to exclude them. With the new change, data not used in the fit may be used in averages. For branching ratios which contain some data which are to be used only in the average, a new column has been added to the Listings to indicate which data are used both for fits and averages (“f&a”) and which are flagged for use only for averages (“avg”).

Experimental measurements of the charged-prong topological branching fractions[†] and $B(h^-h^+h^+\nu_\tau)$ also exhibit some internal inconsistency. Refer to the scale factors and ideograms given in the Full Listings for further information.

Conclusions: Experimental measurements of τ branching fractions contain internal inconsistencies. Measurements have become sufficiently precise that the constrained fit to branching ratio data now has an improbably large χ^2 . Much of the inconsistency is contained in decay modes of the type $\tau^- \rightarrow h^- + (\geq)n\pi^0\nu_\tau$ for $n = 1, 2,$ and 3 , although the charged-prong topological branching fractions and $B(h^-h^+h^+\nu_\tau)$ also show signs of problems. Future precise measurements at LEP should resolve the problems in the charged-prong topological branching fractions. Additional measurements with small, well-understood systematic errors are needed to sort out the confusion in the other modes.

Notes and References

* Because of the way we have grouped the modes, the number of degrees of freedom (dof) is not available except for the “all modes” category. While not as well defined as χ^2 per dof, the number provides a useful measure of the source of the dominant χ^2 contributions.

[†] We have slightly generalized the idea of topological branching fractions to include $(\geq)n\pi^0$'s. These are *topological*, in the sense that they classify what is observed, rather than *physical*, which would classify modes at the top of the decay chain: $e, \mu, \pi, K, \rho, K^*(892), \text{etc.}$

1. T.N. Troung, Phys. Rev. **D30**, 1509 (1984).
2. F.J. Gilman and S.H. Rhie, Phys. Rev. **D31**, 1066 (1985); F.J. Gilman, Phys. Rev. **D35**, 3541 (1987).
3. See for example M.A. Samuel, Mod. Phys. Lett. **A8**, 2491 (1993).
4. R. Akers *et al.*, Phys. Lett. B (to be published).
5. M. Artuso *et al.*, Phys. Rev. Lett. (to be published).
6. M. Procario *et al.*, Phys. Rev. Lett. **70**, 1207 (1993).

 τ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1777.1^{+0.4}_{-0.5}				OUR AVERAGE
1777.8 \pm 0.7 \pm 1.7	35k	¹ BALEST	93 CLEO	$E_{CM}^{eff} = 10.6$ GeV
1776.3 \pm 2.4 \pm 1.4	11k	² ALBRECHT	92M ARG	$E_{CM}^{eff} = 9.4\text{--}10.6$ GeV
1776.9 ^{+0.4} _{-0.5} \pm 0.2	14	³ BAI	92 BES	$E_{CM}^{eff} = 3.55\text{--}3.60$ GeV
1787 \pm 10		BLOCKER	80 MRK2	$E_{CM}^{eff} = 3.5\text{--}6.7$ GeV
1783 ⁺³ ₋₄	692	⁴ BACINO	78B DLCO	$E_{CM}^{eff} = 3.1\text{--}7.4$ GeV
1787 ⁺¹⁰ ₋₁₈	299	⁵ BARTEL	78 SPEC	$E_{CM}^{eff} = 3.6\text{--}4.4$ GeV
1807 \pm 20		BRANDELIK	78 DASP	$E_{CM}^{eff} = 3.1\text{--}5.2$ GeV
•••				We do not use the following data for averages, fits, limits, etc. •••
1803 \pm 16	1138	BLOCKER	82D MRK2	Incl. in BLOCKER 80
¹ BALEST 93 fit spectra of minimum kinematically allowed τ mass in events of the type $e^+e^- \rightarrow \tau^+\tau^- \rightarrow (\pi^+n\pi^0\nu_\tau)(\pi^-m\pi^0\nu_\tau)$ $n \leq 2, m \leq 2, 1 \leq n+m \leq 3$. If $m_{\nu_\tau} \neq 0$, result increases by $(m_{\nu_\tau}^2/1100)$ MeV.				
² ALBRECHT 92M fit τ pseudomass spectrum in $\tau^- \rightarrow 2\pi^-\pi^+\nu_\tau$ decays. Result assumes $m_{\nu_\tau} = 0$.				
³ BAI 92 fit $\sigma(e^+e^- \rightarrow \tau^+\tau^-)$ near threshold using $e\mu$ events.				
⁴ BACINO 78B value comes from $e^\pm\chi^\mp$ threshold. Published mass 1782 MeV increased by 1 MeV using the high precision $\psi(2S)$ mass measurement of ZHOLENTZ 80 to eliminate the absolute SPEAR energy calibration uncertainty.				
⁵ BARTEL 78 fits energy dependence of cross section for e^\pm and μ^\pm events. Mass value not dependent on whether $V-A$ or $V+A$ decay assumed.				

 τ MEAN LIFE

The downward revision of the τ mass to 1777.1 MeV (see the τ mass above) results in a downward correction of some of the older lifetime measurements, with the net result that the average should be reduced by 0.2 fs. WASSERBAECH 93 shows that additional systematic biases are not consistently taken into account by the experimenters. When the appropriate corrections are applied, the average is decreased by an additional 0.8 fs. “OUR EVALUATION” reflects these changes.

VALUE (10^{-15} s)	EVTS	DOCUMENT ID	TECN	COMMENT
295.6\pm 3.1				OUR EVALUATION
296.6\pm 3.1				OUR AVERAGE
298 \pm 7		ABREU	93E DLPH	$E_{CM}^{eff} = 88.5\text{--}93.7$ GeV
291.9 \pm 5.1 \pm 3.1	28k	ACTON	93J OPAL	$E_{CM}^{eff} = 88\text{--}94$ GeV
293 \pm 9 \pm 12	5743	ADRIANI	93M L3	1991 LEP run
304 \pm 14 \pm 7	4100	BATTLE	92 CLEO	$E_{CM}^{eff} = 10.6$ GeV
294.7 \pm 5.4 \pm 3.0	11.8k	BUSKULIC	92H ALEP	$E_{CM}^{eff} = 88.5\text{--}93.7$ GeV
314 \pm 23 \pm 9		ABREU	91D DLPH	$E_{CM}^{eff} = 88\text{--}94$ GeV
309 \pm 23 \pm 30	2817	ADEVA	91F L3	1990 LEP run
301 \pm 29	3780	KLEINWORT	89 JADE	$E_{CM}^{eff} = 35\text{--}46$ GeV
288 \pm 16 \pm 17	807	AMIDEI	88 MRK2	$E_{CM}^{eff} = 29$ GeV
306 \pm 20 \pm 14	695	BRAUNSCH...	88C TASS	$E_{CM}^{eff} = 36$ GeV
299 \pm 15 \pm 10	1311	ABACHI	87C HRS	$E_{CM}^{eff} = 29$ GeV
295 \pm 14 \pm 11	5696	ALBRECHT	87P ARG	$E_{CM}^{eff} = 9.3\text{--}10.6$ GeV
309 \pm 17 \pm 7	3788	BAND	87B MAC	$E_{CM}^{eff} = 29$ GeV
325 \pm 14 \pm 18	8470	BEBEK	87C CLEO	$E_{CM}^{eff} = 10.5$ GeV
•••				We do not use the following data for averages, fits, limits, etc. •••
291 \pm 13 \pm 6	6621	DECAMP	92E ALEP	Repl. by BUSKULIC 92H
308 \pm 13		ACTON	91C OPAL	Repl. by ACTON 93J
315 \pm 36 \pm 40	10k	FERNANDEZ	85 MAC	$E_{CM}^{eff} = 29$ GeV
318 ⁺⁸¹ ₋₉₄	50	ALTHOFF	84D TASS	Repl. by BRAUN-SCHWEIG 88C
320 \pm 54	156	JAROS	83 MRK2	Repl. by AMIDEI 88

τ MAGNETIC MOMENT ANOMALY

$$\mu_{\tau}/(e\hbar/2m_{\tau})-1 = (g_{\tau}-2)/2$$

For a theoretical calculation $[(g_{\tau}-2)/2 = 11773(3) \times 10^{-7}]$, see SAMUEL 91b.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	6 ESCRIBANO	93 RVUE	$Z \rightarrow \tau^+ \tau^-$ at LEP
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.12	90	GRIFOLS	91 RVUE	$Z \rightarrow \tau \tau \gamma$ at LEP
<0.023	95	7 SILVERMAN	83 RVUE	$e^+ e^- \rightarrow \tau^+ \tau^-$ at PETRA

⁶ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the magnetic moment anomaly.

⁷ SILVERMAN 83 limit is derived from $e^+ e^- \rightarrow \tau^+ \tau^-$ total cross-section measurements for q^2 up to $(37 \text{ GeV})^2$.

 τ ELECTRIC DIPOLE MOMENT

A nonzero value is forbidden by both T invariance and P invariance.

VALUE (ecm)	CL%	DOCUMENT ID	TECN	COMMENT
<3.7 $\times 10^{-17}$	95	8 ESCRIBANO	93 RVUE	$Z \rightarrow \tau^+ \tau^-$ at LEP
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<7 $\times 10^{-16}$	90	GRIFOLS	91 RVUE	$Z \rightarrow \tau \tau \gamma$ at LEP
<1.6 $\times 10^{-16}$	90	DELAGUILA	90 RVUE	$e^+ e^- \rightarrow \tau^+ \tau^-$ $E_{\text{cm}}^{\text{eff}} = 35 \text{ GeV}$

⁸ ESCRIBANO 93 limit derived from $\Gamma(Z \rightarrow \tau^+ \tau^-)$, and is on the absolute value of the electric dipole moment.

 τ WEAK DIPOLE MOMENT

A nonzero value is forbidden by CP invariance.

VALUE (ecm)	CL%	DOCUMENT ID	TECN	COMMENT
<3.7 $\times 10^{-17}$	95	9 BUSKULIC	92J ALEP	$Z \rightarrow \tau^+ \tau^-$ at LEP
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<7.0 $\times 10^{-17}$	95	9 ACTON	92F OPAL	$Z \rightarrow \tau^+ \tau^-$ at LEP

⁹ Limit is on the absolute value of the weak dipole moment, and applies for $q^2 = m_{\tau}^2$.

 τ^- DECAY MODES

τ^+ modes are charge conjugates of the modes below. "h \pm " stands for π^{\pm} or K^{\pm} . "l" stands for e or μ . "Neutral" means neutral hadron whose decay products include γ 's and/or π^0 's.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Modes with one charged particle		
Γ_1 particle $^- \geq 0$ neutrals ν_{τ} ("1-prong")	(85.49 \pm 0.24) %	S=1.5
Γ_2 $\mu^- \bar{\nu}_{\mu} \nu_{\tau}$	(17.65 \pm 0.24) %	S=1.1
Γ_3 $\mu^- \bar{\nu}_{\mu} \nu_{\tau} \gamma$ ($E_{\gamma} > 37 \text{ MeV}$)	(2.3 \pm 1.1) $\times 10^{-3}$	
Γ_4 $e^- \bar{\nu}_e \nu_{\tau}$	(18.01 \pm 0.18) %	S=1.1
Γ_5 $h^- \geq 0$ neutrals ν_{τ}	(49.83 \pm 0.35) %	S=1.3
Γ_6 $h^- \nu_{\tau}$	(12.88 \pm 0.34) %	S=1.2
Γ_7 $\pi^- \nu_{\tau}$	(11.7 \pm 0.4) %	S=1.3
Γ_8 $K^- \geq 0$ neutrals ν_{τ}	(1.68 \pm 0.24) %	
Γ_9 $K^- \nu_{\tau}$	(6.7 \pm 2.3) $\times 10^{-3}$	S=1.3
Γ_{10} $K^- \geq 1$ neutrals ν_{τ}	(1.2 $^{+0.5}_{-0.6}$) %	
Γ_{11} $h^- \geq 1$ neutrals ν_{τ}	(36.9 \pm 0.4) %	S=1.3
Γ_{12} $h^- \pi^0 \nu_{\tau}$	(25.7 \pm 0.4) %	S=1.7
Γ_{13} $\pi^- \pi^0 \nu_{\tau}$	(25.2 \pm 0.4) %	S=1.7
Γ_{14} $\pi^- \pi^0 \text{non-}\rho(770) \nu_{\tau}$		
Γ_{15} $h^- \geq 2 \pi^0 \nu_{\tau}$	(11.2 \pm 0.4) %	S=1.5
Γ_{16} $h^- 2 \pi^0 \nu_{\tau}$	(9.6 \pm 0.4) %	S=1.5
Γ_{17} $h^- \geq 3 \pi^0 \nu_{\tau}$	(1.48 \pm 0.26) %	S=1.7
Γ_{18} $h^- 3 \pi^0 \nu_{\tau}$	(1.28 \pm 0.24) %	S=1.7
Γ_{19} $h^- 4 \pi^0 \nu_{\tau}$	(1.9 $^{+1.1}_{-1.0}$) $\times 10^{-3}$	S=1.6

Modes with three charged particles

Γ_{20} $2h^- h^+ \geq 0$ neutrals ν_{τ} ("3-prong")	(14.38 \pm 0.24) %	S=1.5
Γ_{21} $h^- h^- h^+ \nu_{\tau}$	(8.42 \pm 0.31) %	S=1.3
Γ_{22} $h^- h^- h^+ \geq 1$ neutrals ν_{τ}	(5.63 \pm 0.30) %	S=1.2
Γ_{23} $\pi^- \pi^- \pi^+ \pi^0 \nu_{\tau}$		
Γ_{24} ($a_1(1260) \pi^-$) $^- \nu_{\tau}$		
Γ_{25} ($\rho \pi^0$) $^0 \pi^- \nu_{\tau}$		
Γ_{26} $\rho^0 \pi^0 \pi^- \nu_{\tau}$		
Γ_{27} $\rho^+ \pi^- \pi^- \nu_{\tau}$		
Γ_{28} $\rho^- \pi^+ \pi^- \nu_{\tau}$		
Γ_{29} $h^- h^- h^+ 2 \pi^0 \nu_{\tau}$	(4.9 \pm 0.5) $\times 10^{-3}$	
Γ_{30} $\omega \pi^- \geq 0$ neutrals ν_{τ}	(1.6 \pm 0.4) %	
Γ_{31} $\omega \pi^- \nu_{\tau}$	(1.6 \pm 0.5) %	
Γ_{32} $h^- \omega \pi^0 \nu_{\tau}$	(4.0 \pm 0.6) $\times 10^{-3}$	
Γ_{33} $K^- h^+ h^- \geq 0$ neutrals ν_{τ}	< 6 $\times 10^{-3}$	CL=90%
Γ_{34} $K^- \pi^+ \pi^- \geq 0$ neutrals ν_{τ}	(2.2 $^{+1.6}_{-1.3}$) $\times 10^{-3}$	
Γ_{35} $K^- K^+ \pi^- \nu_{\tau}$	(2.2 $^{+1.7}_{-1.1}$) $\times 10^{-3}$	

Modes with five charged particles

Γ_{36} $3h^- 2h^+ \geq 0$ neutrals ν_{τ} ("5-prong")	(1.25 \pm 0.24) $\times 10^{-3}$	
Γ_{37} $3h^- 2h^+ \nu_{\tau}$	(5.6 \pm 1.6) $\times 10^{-4}$	
Γ_{38} $3h^- 2h^+ \pi^0 \nu_{\tau}$	(5.1 \pm 2.2) $\times 10^{-4}$	

Miscellaneous other allowed modes

Γ_{39} $4h^- 3h^+ \geq 0$ neutrals ν_{τ} ("7-prong")	< 1.9 $\times 10^{-4}$	CL=90%
Γ_{40} $K^*(892)^- \geq 0$ neutrals ν_{τ}	(1.43 \pm 0.17) %	
Γ_{41} $K^*(892)^- \nu_{\tau}$	(1.45 \pm 0.18) %	
Γ_{42} $K^*(892)^0 K^- \geq 0$ neutrals ν_{τ}	(3.2 \pm 1.4) $\times 10^{-3}$	
Γ_{43} $\bar{K}^*(892)^0 \pi^- \geq 0$ neutrals ν_{τ}	(3.8 \pm 1.7) $\times 10^{-3}$	
Γ_{44} $K^0 h^- \geq 0$ neutrals ν_{τ}	(1.30 \pm 0.30) %	
Γ_{45} $K^- K^0 \geq 0$ neutrals ν_{τ}	< 8 $\times 10^{-3}$	CL=90%
Γ_{46} $K^0 K^- \nu_{\tau}$	< 2.6 $\times 10^{-3}$	CL=95%
Γ_{47} $K^0 K^- \geq 1$ neutrals ν_{τ}	< 2.6 $\times 10^{-3}$	CL=95%
Γ_{48} $K^0 h^+ h^- h^- \geq 0$ neutrals ν_{τ}	< 1.7 $\times 10^{-3}$	CL=95%
Γ_{49} $K_2^*(1430)^- \nu_{\tau}$	< 3 $\times 10^{-3}$	CL=95%
Γ_{50} $a_0(980)^- \geq 0$ neutrals ν_{τ}		
Γ_{51} $\eta \pi^- \geq 0$ neutrals ν_{τ}	< 1.3 %	CL=95%
Γ_{52} $\eta \pi^- \nu_{\tau}$	< 3.4 $\times 10^{-4}$	CL=95%
Γ_{53} $\eta \pi^- \pi^0 \nu_{\tau}$	(1.70 \pm 0.28) $\times 10^{-3}$	
Γ_{54} $\eta \pi^- \pi^0 \pi^0 \nu_{\tau}$	< 4.3 $\times 10^{-4}$	CL=95%
Γ_{55} $\eta K^- \nu_{\tau}$	< 4.7 $\times 10^{-4}$	CL=95%
Γ_{56} $\eta \pi^+ \pi^- \pi^- \geq 0$ neutrals ν_{τ}	< 3 $\times 10^{-3}$	CL=90%
Γ_{57} $\eta \eta \pi^- \geq 0$ neutrals ν_{τ}	< 5 $\times 10^{-3}$	CL=90%
Γ_{58} $\eta \eta \pi^- \nu_{\tau}$	< 1.1 $\times 10^{-4}$	CL=95%
Γ_{59} $\eta \eta \pi^- \pi^0 \nu_{\tau}$	< 2.0 $\times 10^{-4}$	CL=95%

Lepton & Quark Full Listings

T

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{total}$ Γ_2/Γ
 Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
17.65 ± 0.24 OUR FIT Error includes scale factor of 1.1.				
17.44 ± 0.23 OUR AVERAGE				
17.6 ± 0.4 ± 0.4	f&a	2148	ADRIANI 93M L3	$E_{cm}^{ee} = 88-94$ GeV
17.4 ± 0.7 ± 0.6	f&a	687	ABREU 92N DLPH	$E_{cm}^{ee} = 88.2-94.2$ GeV
17.2 ± 0.4 ± 0.5	avg		15 ALBRECHT 92D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
17.35 ± 0.41 ± 0.37	f&a		DECAMP 92C ALEP	$E_{cm}^{ee} = 88-95$ GeV
16.8 ± 0.5 ± 0.4	f&a	903	ALEXANDER 91D OPAL	$E_{cm}^{ee} = 88.3-94.3$ GeV
17.7 ± 0.8 ± 0.4	f&a	568	BEHREND 90 CELL	$E_{cm}^{ee} = 35$ GeV
17.4 ± 1.0	f&a	2197	ADEVA 88 MRKJ	$E_{cm}^{ee} = 14-16$ GeV
17.7 ± 1.2 ± 0.7	f&a		AIHARA 87B TPC	$E_{cm}^{ee} = 29$ GeV
18.3 ± 0.9 ± 0.8	f&a		BURCHAT 87 MRK2	$E_{cm}^{ee} = 29$ GeV
18.6 ± 0.8 ± 0.7	avg	558	16 BARTEL 86D JADE	$E_{cm}^{ee} = 34.6$ GeV
12.9 ± 1.7 $\begin{smallmatrix} +0.7 \\ -0.5 \end{smallmatrix}$	f&a		ALTHOFF 85 TASS	$E_{cm}^{ee} = 34.5$ GeV
18.0 ± 0.9 ± 0.5	avg	473	16 ASH 85B MAC	$E_{cm}^{ee} = 29$ GeV
18.0 ± 1.0 ± 0.6	f&a		17 BALTRUSAIT..85 MRK3	$E_{cm}^{ee} = 3.77$ GeV
19.4 ± 1.6 ± 1.7	f&a	153	BERGER 85 PLUT	$E_{cm}^{ee} = 34.6$ GeV
17.6 ± 2.6 ± 2.1	f&a	47	BEHREND 83C CELL	$E_{cm}^{ee} = 34$ GeV
17.8 ± 2.0 ± 1.8	f&a		BERGER 81B PLUT	$E_{cm}^{ee} = 9-32$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •
 17.5 ± 0.8 ± 0.5 624 ADEVA 91F L3 Repl. by ADRIANI 93M
 17.4 ± 0.6 ± 0.8 1201 ADEVA 86B MRKJ Repl. by ADEVA 88
 15 Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ and $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{total}$.
 16 Modified using $B(e^- \bar{\nu}_e \nu_\tau)/B("1 \text{ prong}')$ and $B("1 \text{ prong}')$, = 0.855.
 17 Error correlated with BALTRUSAITIS 85 $e\nu\bar{\nu}$ value.

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(\text{particle}^- \geq 0 \text{ neutrals } \nu_\tau ("1\text{-prong}'))$ $\Gamma_2/\Gamma_1 = \Gamma_2/(\Gamma_2+\Gamma_4+\Gamma_7+\Gamma_9+\Gamma_{13}+\Gamma_{16}+\Gamma_{18}+\Gamma_{19}+0.771\Gamma_{41})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2065 ± 0.0028 OUR FIT Error includes scale factor of 1.2.				
0.214 ± 0.008 OUR AVERAGE				
0.217 ± 0.009 ± 0.008		BARTEL 86D JADE	$E_{cm}^{ee} = 34.6$ GeV	
0.211 ± 0.010 ± 0.006	390	ASH 85B MAC	$E_{cm}^{ee} = 29$ GeV	

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau \gamma)/\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)$ Γ_3/Γ_2
 $E_\gamma > 37$ MeV.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.013 ± 0.006				
	10	18 WU 90 MRK2	$E_{cm}^{ee} = 29$ GeV	

18 Requirements on detected γ_s correspond to a τ rest frame energy cutoff $E_\gamma > 37$ MeV.

$\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{total}$ Γ_4/Γ
 Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
18.01 ± 0.18 OUR FIT Error includes scale factor of 1.1.				
17.90 ± 0.17 OUR AVERAGE				
17.9 ± 0.4 ± 0.4	f&a	2892	ADRIANI 93M L3	$E_{cm}^{ee} = 88-94$ GeV
18.6 ± 0.8 ± 0.6	f&a	554	ABREU 92N DLPH	$E_{cm}^{ee} = 88.2-94.2$ GeV
17.97 ± 0.14 ± 0.23	f&a	3970	AKERIB 92 CLEO	$E_{cm}^{ee} = 10.6$ GeV
17.3 ± 0.4 ± 0.5	avg		19 ALBRECHT 92D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
19.1 ± 0.4 ± 0.6	avg	2960	20 AMMAR 92 CLEO	$E_{cm}^{ee} = 10.5-10.9$ GeV
18.09 ± 0.45 ± 0.45	f&a		DECAMP 92C ALEP	$E_{cm}^{ee} = 88-95$ GeV
17.4 ± 0.5 ± 0.4	f&a	964	ALEXANDER 91D OPAL	$E_{cm}^{ee} = 88.3-94.3$ GeV
17.0 ± 0.5 ± 0.6	f&a	1.7k	ABACHI 90 HRS	$E_{cm}^{ee} = 29$ GeV
18.4 ± 0.8 ± 0.4	f&a	644	BEHREND 90 CELL	$E_{cm}^{ee} = 35$ GeV
16.3 ± 1.3 ± 3.2	f&a		JANSSEN 89 CBAL	$E_{cm}^{ee} = 9.4-10.6$ GeV
18.4 ± 1.2 ± 1.0	f&a		AIHARA 87B TPC	$E_{cm}^{ee} = 29$ GeV
19.1 ± 0.8 ± 1.1	f&a		BURCHAT 87 MRK2	$E_{cm}^{ee} = 29$ GeV
16.8 ± 0.7 ± 0.9	avg	515	20 BARTEL 86D JADE	$E_{cm}^{ee} = 34.6$ GeV
20.4 ± 3.0 $\begin{smallmatrix} +1.4 \\ -0.9 \end{smallmatrix}$	f&a		ALTHOFF 85 TASS	$E_{cm}^{ee} = 34.5$ GeV
17.8 ± 0.9 ± 0.6	avg	390	20 ASH 85B MAC	$E_{cm}^{ee} = 29$ GeV
18.2 ± 0.7 ± 0.5	f&a		21 BALTRUSAIT..85 MRK3	$E_{cm}^{ee} = 3.77$ GeV
13.0 ± 1.9 ± 2.9	f&a		BERGER 85 PLUT	$E_{cm}^{ee} = 34.6$ GeV
18.3 ± 2.4 ± 1.9	f&a	60	BEHREND 83C CELL	$E_{cm}^{ee} = 34$ GeV
16.0 ± 1.3	f&a	459	22 BACINO 78B DLCO	$E_{cm}^{ee} = 3.1-7.4$ GeV


• • • We do not use the following data for averages, fits, limits, etc. • • •
 17.7 ± 0.7 ± 0.6 686 ADEVA 91F L3 Repl. by ADRIANI 93M
 18.3 ± 0.7 ± 0.5 23 AIHARA 87B TPC $E_{cm}^{ee} = 29$ GeV

19 Not independent of ALBRECHT 92D $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ and $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{total}$.
 20 Modified using $B(e^- \bar{\nu}_e \nu_\tau)/B("1 \text{ prong}')$ and $B("1 \text{ prong}')$, = 0.855.
 21 Error correlated with BALTRUSAITIS 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma_{total}$.
 22 BACINO 78B value comes from fit to events with e^\pm and one other nonelectron charged prong.
 23 Combined result of AIHARA 87B $e\nu\bar{\nu}$ and $\mu\nu\bar{\nu}$ measurements assuming $B(\mu\nu\bar{\nu})/B(e\nu\bar{\nu}) = 0.973$.

$\Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma(\text{particle}^- \geq 0 \text{ neutrals } \nu_\tau ("1\text{-prong}'))$ $\Gamma_4/\Gamma_1 = \Gamma_4/(\Gamma_2+\Gamma_4+\Gamma_7+\Gamma_9+\Gamma_{13}+\Gamma_{16}+\Gamma_{18}+\Gamma_{19}+0.771\Gamma_{41})$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.2107 ± 0.0022 OUR FIT Error includes scale factor of 1.1.				
0.213 ± 0.008 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.				
0.2231 ± 0.0044 ± 0.0073	2856	AMMAR 92 CLEO	$E_{cm}^{ee} = 10.5-10.9$ GeV	
0.196 ± 0.008 ± 0.010		BARTEL 86D JADE	$E_{cm}^{ee} = 34.6$ GeV	
0.208 ± 0.010 ± 0.007	390	ASH 85B MAC	$E_{cm}^{ee} = 29$ GeV	

WEIGHTED AVERAGE
 0.213 ± 0.008 (Error scaled by 1.3)



$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \times \Gamma(e^- \bar{\nu}_e \nu_\tau)/\Gamma_{total}^2$ $\Gamma_2\Gamma_4/\Gamma^2$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0318 ± 0.0006 OUR FIT Error includes scale factor of 1.1.				
0.0302 ± 0.0012 OUR AVERAGE				
0.0306 ± 0.0005 ± 0.0013	3230	ALBRECHT 93G ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV	
0.0288 ± 0.0017 ± 0.0019		ASH 85B MAC	$E_{cm}^{ee} = 29$ GeV	
0.030 ± 0.005	257	BLOCKER 82D MRK2	$E_{cm}^{ee} = 3.5-6.7$ GeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 0.0298 ± 0.0007 ± 0.0013 1756 ALBRECHT 92D ARG Repl. by ALBRECHT 93G

$\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau)/\Gamma(e^- \bar{\nu}_e \nu_\tau)$ Γ_2/Γ_4
 Predicted to be 1 for sequential lepton, 1/2 for para-electron, and 2 for para-muon. Para-electron also ruled out by HEILE 78.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.980 ± 0.017 OUR FIT Error includes scale factor of 1.1.				
1.00 ± 0.05 OUR AVERAGE				
0.997 ± 0.035 ± 0.040		ALBRECHT 92D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV	
1.33 ± 0.18 ± 0.36	154	BLOCKER 82D MRK2	$E_{cm}^{ee} = 3.5-6.7$ GeV	

$\Gamma(h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{total}$ $\Gamma_5/\Gamma = (\Gamma_7+\Gamma_9+\Gamma_{13}+\Gamma_{16}+\Gamma_{18}+\Gamma_{19}+0.771\Gamma_{41})/\Gamma$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
49.83 ± 0.35 OUR FIT Error includes scale factor of 1.3.			
49.0 ± 1.4 OUR AVERAGE			
48.6 ± 1.2 ± 0.9	avg	24 AIHARA 87B TPC	$E_{cm}^{ee} = 29$ GeV
51.5 ± 2.9 $\begin{smallmatrix} +1.6 \\ -2.6 \end{smallmatrix}$	f&a	ALTHOFF 85 TASS	$E_{cm}^{ee} = 34.5$ GeV

24 Not independent of AIHARA 87B $e\nu\bar{\nu}$, $\mu\nu\bar{\nu}$, and $\pi^+ 2\pi^- (\geq 0\pi^0)\nu$ values.

$\Gamma(h^- \nu_\tau)/\Gamma_{total}$ $\Gamma_6/\Gamma = (\Gamma_7 + \Gamma_9 + \frac{1}{3}\Gamma_{41})/\Gamma$
 Inclusion of the $\frac{1}{3}B(\tau^- \rightarrow K^*(892)^- \nu_\tau)$ corrects, at <0.26% level, for undetected K_L^0 's, which are predominately from $K^*(892)^-$ decay.

Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
12.88 ± 0.34 OUR FIT				Error includes scale factor of 1.2.
12.47 ± 0.33 OUR AVERAGE				
12.4 ± 0.7 ± 0.7	f&a	283	25 ABREU	92N DLPH $E_{cm}^{cc} = 88.2-94.2$ GeV
13.32 ± 0.44 ± 0.33	f&a		26 DECAMP	92C ALEP $E_{cm}^{cc} = 88-95$ GeV
12.1 ± 0.7 ± 0.5	f&a	309	ALEXANDER	91D OPAL $E_{cm}^{cc} = 88.3-94.3$ GeV
12.3 ± 0.9 ± 0.5	f&a	1338	BEHREND	90 CELL $E_{cm}^{cc} = 35$ GeV
11.1 ± 1.1 ± 1.4	f&a		27 BURCHAT	87 MRK2 $E_{cm}^{cc} = 29$ GeV
11.3 ± 0.5 ± 0.8	avg	798	28 FORD	87 MAC $E_{cm}^{cc} = 29$ GeV
12.3 ± 0.6 ± 1.1	avg	328	29 BARTEL	86D JADE $E_{cm}^{cc} = 34.6$ GeV
13.0 ± 2.0 ± 4.0	f&a		BERGER	85 PLUT $E_{cm}^{cc} = 34.6$ GeV
11.2 ± 1.7 ± 1.2	f&a	34	30 BEHREND	83C CELL $E_{cm}^{cc} = 34$ GeV

- ²⁵ ABREU 92 with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.
- ²⁶ DECAMP 92C consider $\tau^- \rightarrow h^-(K_L^0 \rightarrow \pi^+\pi^-)\nu$ to be a 1-prong mode, which affects their 1-prong topological branching ratio relative to other experiments.
- ²⁷ BURCHAT 87 with 1.1% added to remove their correction for K^- and $K^*(892)^-$ backgrounds.
- ²⁸ FORD 87 result for $B(\pi^- \nu_\tau)$ with 0.67% added to remove their K^- correction and adjusted for 1992 B("1 prong").
- ²⁹ BARTEL 86D result for $B(\pi^- \nu_\tau)$ with 0.59% added to remove their K^- correction and adjusted for 1992 B("1 prong").
- ³⁰ BEHREND 83C quote $B(\pi^- \nu_\tau) = 9.9 \pm 1.7 \pm 1.3$ after subtracting 1.3 ± 0.5 to correct for $B(K^- \nu_\tau)$.

$\Gamma(h^- \nu_\tau)/\Gamma(\text{particle}^- \geq 0 \text{ neutrals } \nu_\tau \text{ ("1-prong")})$
 $\Gamma_6/\Gamma_1 = (\Gamma_7 + \Gamma_9 + \frac{1}{3}\Gamma_{41})/(\Gamma_2 + \Gamma_4 + \Gamma_7 + \Gamma_9 + \Gamma_{13} + \Gamma_{16} + \Gamma_{18} + \Gamma_{19} + 0.771\Gamma_{41})$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.151 ± 0.004 OUR FIT				Error includes scale factor of 1.2.
0.135 ± 0.009 OUR AVERAGE				
0.131 ± 0.006 ± 0.009	798	31 FORD	87 MAC	$E_{cm}^{cc} = 29$ GeV
0.143 ± 0.007 ± 0.013	328	32 BARTEL	86D JADE	$E_{cm}^{cc} = 34.6$ GeV

- ³¹ FORD 87 result divided by 0.865, their assumed value for B("1 prong").
- ³² BARTEL 86D result with 0.6% added to remove their K^- correction and then divided by 0.865, their assumed value for B("1 prong").

$\Gamma(h^- \nu_\tau)/\Gamma(e^- \nu_e \nu_\tau)$ $\Gamma_6/\Gamma_4 = (\Gamma_7 + \Gamma_9 + \frac{1}{3}\Gamma_{41})/\Gamma_4$

VALUE (%)	DOCUMENT ID	TECN	COMMENT
0.715 ± 0.021 OUR FIT			Error includes scale factor of 1.2.
0.678 ± 0.037 ± 0.044	ALBRECHT	92D ARG	$E_{cm}^{cc} = 9.4-10.6$ GeV
0.647 ± 0.039 ± 0.061	33 BARTEL	86D JADE	$E_{cm}^{cc} = 34.6$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •
- ³³ Combined result of BARTEL 86D $e\nu\bar{\nu}$, $\mu\nu\bar{\nu}$, and $\pi^- \nu$ assuming $B(\mu\nu\bar{\nu})/B(e\nu\bar{\nu}) = 0.973$.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
11.7 ± 0.4 OUR FIT				Error includes scale factor of 1.3.
11.7 ± 0.4 ± 1.8	1138	BLOCKER	82D MRK2	$E_{cm}^{cc} = 3.5-6.7$ GeV

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.68 ± 0.24 OUR AVERAGE				
1.6 ± 0.4 ± 0.2	35	AIHARA	87B TPC	$E_{cm}^{cc} = 29$ GeV
1.71 ± 0.29	53	MILLS	84 DLCO	$E_{cm}^{cc} = 29$ GeV

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.67 ± 0.23 OUR FIT				Error includes scale factor of 1.3.
0.67 ± 0.23 OUR AVERAGE				Error includes scale factor of 1.3.
0.59 ± 0.18	16	MILLS	84 DLCO	$E_{cm}^{cc} = 29$ GeV
1.3 ± 0.5	15	BLOCKER	82B MRK2	$E_{cm}^{cc} = 3.9-6.7$ GeV


VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.2 ± 0.5 ± 0.2 - 0.4	9	AIHARA	87B TPC	$E_{cm}^{cc} = 29$ GeV

$\Gamma(h^- \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{total}$ $\Gamma_{11}/\Gamma = (\Gamma_{13} + \Gamma_{16} + \Gamma_{18} + \Gamma_{19} + 0.438\Gamma_{41})/\Gamma$

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
36.9 ± 0.4 OUR FIT				Error includes scale factor of 1.3.
36.7 ± 0.8 OUR AVERAGE				Error includes scale factor of 1.4. See the Ideogram below.
36.14 ± 0.33 ± 0.58		AKERS	94E OPAL	1991-1992 LEP runs
38.4 ± 1.2 ± 1.0		34 BURCHAT	87 MRK2	$E_{cm}^{cc} = 29$ GeV
42.7 ± 2.0 ± 2.9		BERGER	85 PLUT	$E_{cm}^{cc} = 34.6$ GeV

- ³⁴ BURCHAT 87 quote for $B(\pi^\pm \geq 1 \text{ neutrals } \nu_\tau) = 0.378 \pm 0.012 \pm 0.010$. We add 0.006 to account for contribution from $(K^* \nu_\tau)$ which they fixed at BR = 0.013.

WEIGHTED AVERAGE
 36.7 ± 0.8 (Error scaled by 1.4)




$\Gamma(h^- \pi^0 \nu_\tau)/\Gamma_{total}$ $\Gamma_{12}/\Gamma = (\Gamma_{13} + \frac{1}{3}\Gamma_{41})/\Gamma$
 Inclusion of the $\frac{1}{3}B(\tau^- \rightarrow K^*(892)^- \nu_\tau)$ corrects, at <0.26% level, for $K^- \pi^0$ from $K^*(892)$ decays.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
25.7 ± 0.4 OUR FIT				Error includes scale factor of 1.7.
25.4 ± 0.5 OUR AVERAGE				Error includes scale factor of 1.7. See the Ideogram below.
26.25 ± 0.36 ± 0.52		AKERS	94E OPAL	1991-1992 LEP runs
26.22 ± 0.12 ± 0.42	51k	35 ARTUSO	94 CLEO	$E_{cm}^{cc} = 10.6$ GeV
22.9 ± 0.8 ± 1.3	283	36 ABREU	92N DLPH	$E_{cm}^{cc} = 88.2-94.2$ GeV
23.1 ± 0.4 ± 0.9	1249	37 ALBRECHT	92Q ARG	$E_{cm}^{cc} = 10$ GeV
25.02 ± 0.64 ± 0.88	1849	DECAMP	92C ALEP	$E_{cm}^{cc} = 88-95$ GeV
22.0 ± 0.8 ± 1.9	779	ANTREASANYAN	91 CBAL	$E_{cm}^{cc} = 9.4-10.6$ GeV
22.6 ± 1.5 ± 0.7	1101	BEHREND	90 CELL	$E_{cm}^{cc} = 35$ GeV
23.1 ± 1.9 ± 1.6		BEHREND	84 CELL	$E_{cm}^{cc} = 14,22$ GeV

- ³⁵ ARTUSO 94 reports the combined result from three independent methods, one of which (23% of the $\tau^- \rightarrow h^- \pi^0 \nu_\tau$) is normalized to the inclusive one-prong branching fraction, taken as 0.854 ± 0.004 . Renormalization to the present value causes negligible change. We add 0.35% to the published result to include $\tau^- \rightarrow h^- \pi^0 K_L \nu_\tau$.

- ³⁶ ABREU 92 with 0.5% added to remove their correction for $K^*(892)^-$ backgrounds.
- ³⁷ ALBRECHT 92Q with 0.5% added to remove their correction for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ background.

WEIGHTED AVERAGE
 25.4 ± 0.5 (Error scaled by 1.7)



Lepton & Quark Full Listings

T

$\Gamma(\pi^- \pi^0 \nu_\tau)/\Gamma_{total}$		Γ_{13}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
25.2 ± 0.4 OUR FIT	Error includes scale factor of 1.7.				
22.2 ± 1.0 OUR AVERAGE					

21.5 ± 0.4 ± 1.9	4400	38,39 ALBRECHT	88L ARG	$E_{cm}^{ee} = 10$ GeV
23.0 ± 1.3 ± 1.7	582	ADLER	87B MRK3	$E_{cm}^{ee} = 3.77$ GeV
22.3 ± 0.6 ± 1.4	629	39 YELTON	86 MRK2	$E_{cm}^{ee} = 29$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
25.8 ± 1.7 ± 2.5	40	BURCHAT	87 MRK2	$E_{cm}^{ee} = 29$ GeV

³⁸The authors divide by $(\Gamma_2 + \Gamma_4 + \Gamma_7 + \Gamma_9)/\Gamma = 0.467$ to obtain this result.
³⁹Experiment had no hadron identification. Kaon corrections were made, but insufficient information is given to permit their removal.
⁴⁰BURCHAT 87 value is not independent of YELTON 86 value. Nonresonant decays included.

$\Gamma(\pi^- \pi^0 \text{non-}\rho(770)\nu_\tau)/\Gamma_{total}$		Γ_{14}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.3 ± 0.1 ± 0.3	41	BEHREND	84 CELL	$E_{cm}^{ee} = 14.22$ GeV	


⁴¹BEHREND 84 assume a flat nonresonant mass distribution down to the $\rho(770)$ mass, using events with mass above 1300 to set the level.

$\Gamma(h^- \geq 2\pi^0 \nu_\tau)/\Gamma_{total}$		$\Gamma_{15}/\Gamma = (\Gamma_{16} + \Gamma_{18} + \Gamma_{19} + 0.105\Gamma_{41})/\Gamma$			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
11.2 ± 0.4 OUR FIT	Error includes scale factor of 1.5.				
10.7 ± 1.1 OUR AVERAGE	Error includes scale factor of 2.0. See the ideogram below.				

9.89 ± 0.34 ± 0.55 avg		AKERS	94E OPAL	1991-1992 LEP runs
14.0 ± 1.2 ± 0.6 f&a	938	BEHREND	90 CELL	$E_{cm}^{ee} = 35$ GeV
12.0 ± 1.4 ± 2.5 f&a		42 BURCHAT	87 MRK2	$E_{cm}^{ee} = 29$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13.9 ± 2.0 ± 1.9	43	AIHARA	86E TPC	$E_{cm}^{ee} = 29$ GeV

⁴²Error correlated with BURCHAT 87 $\Gamma(\rho^- \nu_\tau)/\Gamma_{total}$ value.
⁴³AIHARA 86E (TPC) quote $B(2\pi^0 \pi^- \nu_\tau) + 1.6B(3\pi^0 \pi^- \nu_\tau) + 1.1B(\pi^0 \eta \pi^- \nu_\tau)$.

WEIGHTED AVERAGE
10.7 ± 1.1 (Error scaled by 2.0)



$\Gamma(h^- 2\pi^0 \nu_\tau)/\Gamma_{total}$		Γ_{16}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
9.6 ± 0.4 OUR FIT	Error includes scale factor of 1.5.				
9.0 ± 0.4 OUR AVERAGE	Error includes scale factor of 1.2.				

8.96 ± 0.16 ± 0.44 avg	44	PROCARIO	93 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV
10.38 ± 0.66 ± 0.82 f&a	809	45 DECAMP	92C ALEP	$E_{cm}^{ee} = 88-95$ GeV
5.7 ± 0.5 ± 1.7	133	46 ANTREASIAN	91 CBAL	$E_{cm}^{ee} = 9.4-10.6$ GeV
10.0 ± 1.5 ± 1.1 f&a	333	47 BEHREND	90 CELL	$E_{cm}^{ee} = 35$ GeV
8.7 ± 0.4 ± 1.1 f&a	815	48 BAND	87 MAC	$E_{cm}^{ee} = 29$ GeV
6.0 ± 3.0 ± 1.8 f&a		BEHREND	84 CELL	$E_{cm}^{ee} = 14.22$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
6.2 ± 0.6 ± 1.2	49	GAN	87 MRK2	$E_{cm}^{ee} = 29$ GeV

⁴⁴PROCARIO 93 entry is obtained from $B(h^- 2\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.
⁴⁵We subtract 0.0015 to account for $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.
⁴⁶ANTREASIAN 91 subtract 0.001 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.
⁴⁷BEHREND 90 subtract 0.002 to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution.
⁴⁸BAND 87 assume $B(\pi^- 3\pi^0 \nu_\tau) = 0.01$ and $B(\pi^- \pi^0 \eta \nu_\tau) = 0.005$.
⁴⁹GAN 87 analysis use photon multiplicity distribution.

$\Gamma(h^- 2\pi^0 \nu_\tau)/\Gamma(h^- \pi^0 \nu_\tau)$		$\Gamma_{16}/\Gamma_{12} = \Gamma_{16}/(\Gamma_{13} + \frac{1}{3}\Gamma_{41})$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.373 ± 0.019 OUR FIT	Error includes scale factor of 1.5.				
0.342 ± 0.006 ± 0.016	50	PROCARIO	93 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV	

⁵⁰PROCARIO 93 quote 0.345 ± 0.006 ± 0.016 after correction for 2 kaon backgrounds assuming $B(K^* \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We multiply by 0.990 ± 0.010 to remove these corrections to $B(h^- \pi^0 \nu_\tau)$.

$\Gamma(h^- \geq 3\pi^0 \nu_\tau)/\Gamma_{total}$		$\Gamma_{17}/\Gamma = (\Gamma_{18} + \Gamma_{19})/\Gamma$			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.48 ± 0.26 OUR FIT	Error includes scale factor of 1.7.				
1.8 ± 0.5 OUR AVERAGE					

1.53 ± 0.40 ± 0.46	186	DECAMP	92C ALEP	$E_{cm}^{ee} = 88-95$ GeV
3.2 ± 1.0 ± 1.0	51	BEHREND	90 CELL	$E_{cm}^{ee} = 35$ GeV
3.0 ± 2.2 ± 1.5		BEHREND	84 CELL	$E_{cm}^{ee} = 14.22$ GeV

⁵¹Not independent of BEHREND 90 $\Gamma(\text{hadron}^- \geq 2\pi^0 \nu_\tau)/\Gamma_{total}$ and $\Gamma(\pi^- 2\pi^0 \nu_\tau)/\Gamma_{total}$ values.

$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma_{total}$		Γ_{18}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
1.28 ± 0.24 OUR FIT	Error includes scale factor of 1.7.				
1.15 ± 0.08 ± 0.13 avg	52	PROCARIO	93 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0 ± 1.4 ± 1.1	53	GAN	87 MRK2	$E_{cm}^{ee} = 29$ GeV

⁵²PROCARIO 93 entry is obtained from $B(h^- 3\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau)$ using ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$.
⁵³Highly correlated with GAN 87 $\Gamma(\eta \pi^- \pi^0 \nu_\tau)/\Gamma_{total}$ value. Authors quote $B(\pi^\pm 3\pi^0 \nu_\tau) + 0.67B(\pi^\pm \eta \pi^0 \nu_\tau) = 0.047 \pm 0.010 \pm 0.011$.

$\Gamma(h^- 3\pi^0 \nu_\tau)/\Gamma(h^- \pi^0 \nu_\tau)$		$\Gamma_{18}/\Gamma_{12} = \Gamma_{18}/(\Gamma_{13} + \frac{1}{3}\Gamma_{41})$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.050 ± 0.009 OUR FIT	Error includes scale factor of 1.7.				
0.044 ± 0.003 ± 0.005	54	PROCARIO	93 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV	

⁵⁴PROCARIO 93 quote 0.041 ± 0.003 ± 0.005 after correction for 2 kaon backgrounds assuming $B(K^* \nu_\tau) = 1.42 \pm 0.18\%$ and $B(h^- K^0 \pi^0 \nu_\tau) = 0.48 \pm 0.48\%$. We add 0.003 ± 0.003 and multiply the sum by 0.990 ± 0.010 to remove these corrections.

$\Gamma(h^- 4\pi^0 \nu_\tau)/\Gamma_{total}$		Γ_{19}/Γ			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
0.19 ± 0.11	Error includes scale factor of 1.6.				
0.16 ± 0.05 ± 0.05	55	PROCARIO	93 CLEO	$E_{cm}^{ee} \approx 10.6$ GeV	

⁵⁵PROCARIO 93 quotes $B(h^- 4\pi^0 \nu_\tau)/B(h^- \pi^0 \nu_\tau) = 0.006 \pm 0.002 \pm 0.002$. We multiply by the ARTUSO 94 result for $B(h^- \pi^0 \nu_\tau)$ to obtain $B(h^- 4\pi^0 \nu_\tau)$.

$\Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")})/\Gamma_{total}$		$\Gamma_{20}/\Gamma = (\Gamma_{21} + \Gamma_{22} + 0.229\Gamma_{41})/\Gamma$			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT	
14.38 ± 0.24 OUR FIT	Error includes scale factor of 1.5. See the ideogram below.				
14.32 ± 0.27 OUR AVERAGE					

15.26 ± 0.26 ± 0.22		ACTON	92H OPAL	$E_{cm}^{ee} = 88.2-94.2$ GeV
13.3 ± 0.3 ± 0.8	56	ALBRECHT	92D ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
14.35 ± 0.40 ± 0.45		DECAMP	92C ALEP	$E_{cm}^{ee} = 88-95$ GeV
14.4 ± 0.6 ± 0.3		ADEVA	91F L3	$E_{cm}^{ee} = 88.3-94.3$ GeV
13.5 ± 0.3 ± 0.3		ABACHI	89B HRS	$E_{cm}^{ee} = 29$ GeV
15.0 ± 0.4 ± 0.3		BEHREND	89B CELL	$E_{cm}^{ee} = 14-47$ GeV
15.1 ± 0.8 ± 0.6		AIHARA	87B TPC	$E_{cm}^{ee} = 29$ GeV
12.1 ± 0.5 ± 1.2		RUCKSTUHL	86 DLCO	$E_{cm}^{ee} = 29$ GeV
12.8 ± 0.5 ± 0.8	1420	SCHMIDKE	86 MRK2	$E_{cm}^{ee} = 29$ GeV
15.3 ± 1.1 ± 1.3	367	ALTHOFF	85 TASS	$E_{cm}^{ee} = 34.5$ GeV
13.6 ± 0.5 ± 0.8		BARTEL	85F JADE	$E_{cm}^{ee} = 34.6$ GeV
13.3 ± 0.3 ± 0.6		FERNANDEZ	85 MAC	$E_{cm}^{ee} = 29$ GeV


Some inconsistency exists for this mode since experiments differ in how they treat $B(\tau^- \rightarrow h^- (K_S^0 \rightarrow \pi^+ \pi^-))$ decays.

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.8 ± 1.0 ± 0.7	57 BURCHAT	87 MRK2	$E_{cm}^{ee} = 29$ GeV
13.0 ± 0.2 ± 0.3	4098 AKERLOF	85B HRS	Repl. by ABACHI 89b
12.2 ± 1.3 ± 3.9	58 BERGER	85 PLUT	$E_{cm}^{ee} = 34.6$ GeV
14.8 ± 0.9 ± 1.5	660 AIHARA	84C TPC	Repl. by AIHARA 87b
14.8 ± 2.0 ± 1.3	178 BEHREND	84 CELL	Repl. by BEHREND 89b
14.5 ± 2.2 ± 1.3	182 BEHREND	84 CELL	Repl. by BEHREND 89b
15.0 ± 2.0	186 BEHREND	82 CELL	Repl. by BEHREND 89b
14 ± 2	152 BLOCKER	82C MRK2	Repl. by SCHMIDKE 86
24 ± 6	35 BRANDELIK	80 TASS	$E_{cm}^{ee} = 30$ GeV
32 ± 5	692 59 BACINO	78B DLCO	$E_{cm}^{ee} = 3.1-7.4$ GeV
35 ± 11	59 BRANDELIK	78 DASP	Assumes V-A decay
18 ± 6.5	33 59 JAROS	78 MRK1	$E_{cm}^{ee} > 6$ GeV

- 56 This ALBRECHT 92D value is not independent of their $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) \Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{total}^2$ value.
- 57 BURCHAT 87 value is not independent of SCHMIDKE 86 value.
- 58 Not independent of BERGER 85 $\Gamma(\mu^- \bar{\nu}_\mu \nu_\tau) / \Gamma_{total}$, $\Gamma(e^- \bar{\nu}_e \nu_\tau) / \Gamma_{total}$, $\Gamma(h^- \geq 1$ neutrals $\nu_\tau) / \Gamma_{total}$, and $\Gamma(h^- \nu_\tau) / \Gamma_{total}$, and therefore not used in the fit.
- 59 Low energy experiments are not in average or fit because the systematic errors in background subtraction are judged to be large.

WEIGHTED AVERAGE
8.0±0.4 (Error scaled by 1.4)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$$\Gamma(h^- h^- h^+ \nu_\tau) \times \Gamma(\text{particle}^- \geq 0 \text{ neutrals } \nu_\tau \text{ ("1-prong")}) / \Gamma_{total}^2$$

$$\Gamma_{21} / \Gamma_{20} = \Gamma_{21} / (\Gamma_{21} + \Gamma_{22} + 0.229 \Gamma_{41})$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0720 ± 0.0026 OUR FIT				Error includes scale factor of 1.3.
0.063 ± 0.001 ± 0.004	7.5k	65 ALBRECHT 93C ARG		$E_{cm}^{ee} = 9.4-10.6$ GeV

65 ALBRECHT 93C quote $B(\pi^- \pi^- \pi^+ \nu_\tau) = 6.8 \pm 0.1 \pm 0.5\%$. We add $0.5 \pm 0.3\%$ to remove their correction for charged kaon backgrounds, then multiply by 0.8613, their assumed value for B("1-prong").


$$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")})$$

$$\Gamma_{21} / \Gamma_{20} = \Gamma_{21} / (\Gamma_{21} + \Gamma_{22} + 0.229 \Gamma_{41})$$

This branching fractions is not independent of values for $\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{total}$ and $\Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")}) / \Gamma_{total}$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.585 ± 0.020 OUR FIT				Error includes scale factor of 1.2.
0.55 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
0.47 ± 0.03 ± 0.06		RUCKSTUHL 86 DLCO		$E_{cm}^{ee} = 29$ GeV
0.37 +0.35 -0.20	103	ALTHOFF 85 TASS		$E_{cm}^{ee} = 34.5$ GeV
0.61 ± 0.03 ± 0.05		FERNANDEZ 85 MAC		$E_{cm}^{ee} = 29$ GeV

WEIGHTED AVERAGE
14.32±0.27 (Error scaled by 1.5)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")}) / \Gamma_{total} (\%)$

$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma_{total}$
Some inconsistency exists for this mode since experiments differ in how they treat $B(\tau^- \rightarrow h^- (K_S^0 \rightarrow \pi^+ \pi^-) \nu_\tau)$ decays.


Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
8.42 ± 0.31 OUR FIT				Error includes scale factor of 1.3.
8.0 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
7.3 ± 0.1 ± 0.5 avg		60 ALBRECHT 93C ARG		$E_{cm}^{ee} = 9.4-10.6$ GeV
9.49 ± 0.36 ± 0.63 f&a		DECAMP 92C ALEP		$E_{cm}^{ee} = 88-95$ GeV
8.7 ± 0.7 ± 0.3 f&a	694	61 BEHREND 90 CELL		$E_{cm}^{ee} = 35$ GeV
6.4 ± 0.4 ± 0.9 avg		62 RUCKSTUHL 86 DLCO		$E_{cm}^{ee} = 29$ GeV
7.8 ± 0.5 ± 0.8 f&a	890	SCHMIDKE 86 MRK2		$E_{cm}^{ee} = 29$ GeV
8.4 ± 0.4 ± 0.7 avg	1255	62 FERNANDEZ 85 MAC		$E_{cm}^{ee} = 29$ GeV
9.7 ± 2.0 ± 1.3 f&a		BEHREND 84 CELL		$E_{cm}^{ee} = 14,22$ GeV

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 7.0 ± 0.3 ± 0.7 1566 63 BAND 87 MAC $E_{cm}^{ee} = 29$ GeV
- 6.7 ± 0.8 ± 0.9 64 BURCHAT 87 MRK2 $E_{cm}^{ee} = 29$ GeV

- 60 ALBRECHT 93C value with $0.5 \pm 0.3\%$ added to remove their corrections for charged-kaon backgrounds.
- 61 BEHREND 90 subtract 0.3% to account for the $\tau^- \rightarrow K^*(892)^- \nu_\tau$ contribution to measured events.
- 62 Value obtained by multiplying paper's $R = B(h^- h^- h^+ \nu_\tau) / B(3\text{-prong})$ by $B(3\text{-prong}) = 0.143$ and subtracting 0.3% for $K^*(892)$ background.
- 63 BAND 87 subtract for charged kaon modes; not independent of FERNANDEZ 85 value.
- 64 BURCHAT 87 value is not independent of SCHMIDKE 86 value.

WEIGHTED AVERAGE
0.55±0.07 (Error scaled by 1.6)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(h^- h^- h^+ \nu_\tau) / \Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")})$

Lepton & Quark Full Listings

T

$\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{22}/Γ
 Data marked "avg" are highly correlated with data appearing elsewhere in the Listings, and are therefore used for the average given below but not in the overall fits. "f&a" marks results used for the fit and the average.

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
5.63 ± 0.30 OUR FIT				Error includes scale factor of 1.2.
5.4 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.2.
4.95 ± 0.29 ± 0.65	f&a	570 DECAMP	92C ALEP	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV
5.6 ± 0.7 ± 0.3	avg	352 66 BEHREND	90 CELL	$E_{\text{cm}}^{\text{ee}} = 35$ GeV
4.2 ± 0.5 ± 0.9	f&a	203 67 ALBRECHT	87L ARG	$E_{\text{cm}}^{\text{ee}} = 10$ GeV
7.6 ± 0.4 ± 0.9	avg	68,69 RUCKSTUHL	86 DLCO	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
4.7 ± 0.5 ± 0.8	avg	530 70 SCHMIDKE	86 MRK2	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
5.6 ± 0.4 ± 0.7	avg	69 FERNANDEZ	85 MAC	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
6.2 ± 2.3 ± 1.7	f&a	BEHREND	84 CELL	$E_{\text{cm}}^{\text{ee}} = 14,22$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

6.1 ± 0.8 ± 0.9		71 BURCHAT	87 MRK2	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
66 BEHREND 90 value is not independent of BEHREND 90 $B(3h\nu_\tau \geq 1 \text{ neutrals}) + B(5\text{-prong})$.				
67 ALBRECHT 87L measure the product of branching ratios $B(3\pi^\pm \pi^0 \nu_\tau) B((e\bar{\nu} \text{ or } \mu\bar{\nu} \text{ or } \nu \text{ or } K \text{ or } \rho)\nu_\tau) = 0.029$ and use the PDG 86 value for the second branching ratio which sum to 0.69 ± 0.03 to get the quoted value.				
68 Contributions from kaons and from $>1\pi^0$ are subtracted. Not independent of (3-prong + π^0) and (3-prong + $\geq 0\pi^0$) values.				
69 Value obtained using paper's $R = B(h^- h^- h^+ \nu_\tau)/B(3\text{-prong})$ and current $B(3\text{-prong}) = 0.143$.				
70 Not independent of SCHMIDKE 86 $h^- h^- h^+ \nu_\tau$ and $h^- h^- h^+ (\geq 0\pi^0)\nu_\tau$ values.				
71 BURCHAT 87 value is not independent of SCHMIDKE 86 value.				

$\Gamma((\bar{s}_1(1260)\pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau))$	CL%	DOCUMENT ID	TECN	COMMENT
<0.44	95	72 ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

72 ALBRECHT 91D not independent of their $\Gamma(\omega \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, $\Gamma(\rho^0 \pi^0 \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, $\Gamma(\rho^+ \pi^- \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, and $\Gamma(\rho^- \pi^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$ values.

$\Gamma((\rho\pi)^0 \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$	CL%	DOCUMENT ID	TECN	COMMENT
0.64 ± 0.07 ± 0.03	73	ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

73 ALBRECHT 91D not independent of their $\Gamma(\rho^+ \pi^- \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, $\Gamma(\rho^- \pi^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, and $\Gamma(\rho^0 \pi^0 \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$ values.

$\Gamma(\rho^0 \pi^0 \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$	EVTs	DOCUMENT ID	TECN	COMMENT
0.30 ± 0.04 ± 0.02	393	ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

$\Gamma(\rho^+ \pi^- \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$	EVTs	DOCUMENT ID	TECN	COMMENT
0.10 ± 0.03 ± 0.04	142	ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

$\Gamma(\rho^- \pi^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$	EVTs	DOCUMENT ID	TECN	COMMENT
0.26 ± 0.06 ± 0.01	370	ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

$[\Gamma(\rho^+ \pi^- \pi^- \nu_\tau) + \Gamma(\rho^- \pi^+ \pi^- \nu_\tau)]/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$	EVTs	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.06 ± 0.01	475	74 ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

74 ALBRECHT 91D not independent of their $\Gamma(\rho^+ \pi^- \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$ and $\Gamma(\rho^- \pi^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$ values.

$\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau)/\Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")})$	EVTs	DOCUMENT ID	TECN	COMMENT
0.034 ± 0.002 ± 0.003	668	BORTOLETTO93	CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV

$\Gamma(\omega \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	COMMENT
1.65 ± 0.3 ± 0.2	1513	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ee}} \approx 10$ GeV

$\Gamma(\omega \pi^- \nu_\tau)/\Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	COMMENT
1.60 ± 0.27 ± 0.41	139	BARINGER	87 CLEO	$E_{\text{cm}}^{\text{ee}} = 10.5$ GeV

$\Gamma(\omega \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$	EVTs	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.04 ± 0.02	458	ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$[\Gamma(\rho^0 \pi^0 \pi^- \nu_\tau) + \Gamma(\rho^+ \pi^- \pi^- \nu_\tau) + \Gamma(\rho^- \pi^+ \pi^- \nu_\tau) + \Gamma(\omega \pi^- \nu_\tau)]/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$ $(\Gamma_{26} + \Gamma_{27} + \Gamma_{28} + \Gamma_{31})/\Gamma_{23}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.81	95	75 ALBRECHT	91D ARG	$E_{\text{cm}}^{\text{ee}} = 9.4-10.6$ GeV
75 ALBRECHT 91D not independent of their $\Gamma(\omega \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, $\Gamma(\rho^0 \pi^0 \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, $\Gamma(\rho^+ \pi^- \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$, and $\Gamma(\rho^- \pi^+ \pi^- \nu_\tau)/\Gamma(\pi^- \pi^- \pi^+ \pi^0 \nu_\tau)$ values.				

$\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma(2h^- h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("3-prong")})$	EVTs	DOCUMENT ID	TECN	COMMENT
0.028 ± 0.003 ± 0.003	430	76 BORTOLETTO93	CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV

76 Not independent of BORTOLETTO 93 $\Gamma(\pi^- \rightarrow h^- h^- h^+ 2\pi^0 \nu_\tau)$ value.

$\Gamma(h^- \omega \pi^0 \nu_\tau)/\Gamma(h^- h^- h^+ 2\pi^0 \nu_\tau)$	EVTs	DOCUMENT ID	TECN	COMMENT
0.81 ± 0.06 ± 0.06		BORTOLETTO93	CLEO	$E_{\text{cm}}^{\text{ee}} \approx 10.6$ GeV

$\Gamma(K^- K^+ h^+ \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT
<0.6	90	AIHARA	84C TPC	$E_{\text{cm}}^{\text{ee}} = 29$ GeV

$\Gamma(K^- \pi^+ \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	COMMENT
0.22 ± 0.16 ± 0.13	9	77 MILLS	85 DLCO	$E_{\text{cm}}^{\text{ee}} = 29$ GeV

77 Error correlated with MILLS 85 ($K K \pi \nu$) value. Excludes 23% systematic error.

$\Gamma(K^- K^+ \pi^- \nu_\tau)/\Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	COMMENT
0.22 ± 0.17 ± 0.11	9	78 MILLS	85 DLCO	$E_{\text{cm}}^{\text{ee}} = 29$ GeV

78 Error correlated with MILLS 85 ($K \pi \pi^0 \nu$) value. Excludes 23% systematic error.

$\Gamma(3h^- 2h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("5-prong")})/\Gamma_{\text{total}}$	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.125 ± 0.024 OUR FIT					
0.123 ± 0.023 OUR AVERAGE					
0.26 ± 0.06 ± 0.05			ACTON	92H OPAL	$E_{\text{cm}}^{\text{ee}} = 88.2-94.2$ GeV

0.10 ± 0.05 ± 0.04			DECAMP	92C ALEP	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV
0.16 ± 0.13 ± 0.04			BEHREND	89B CELL	$E_{\text{cm}}^{\text{ee}} = 14-47$ GeV
0.102 ± 0.029	13		BYLSMA	87 HRS	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
0.3 ± 0.1 ± 0.2			BARTEL	85F JADE	$E_{\text{cm}}^{\text{ee}} = 34.6$ GeV
0.16 ± 0.08 ± 0.04	4		BURCHAT	85 MRK2	$E_{\text{cm}}^{\text{ee}} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.34	95	ADEVA	91F L3	$E_{\text{cm}}^{\text{ee}} = 88.3-94.3$ GeV
<0.7	95	0 ALTHOFF	85 TASS	$E_{\text{cm}}^{\text{ee}} = 34.5$ GeV
0.13 ± 0.04	10	BELTRAMI	85 HRS	Repl. by BYLSMA 87
<0.17	95	2 FERNANDEZ	85 MAC	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
<0.3	90	4 AIHARA	84C TPC	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
<0.9	95	1 BEHREND	84 CELL	$E_{\text{cm}}^{\text{ee}} = 14,22$ GeV
1.0 ± 0.4	10	BEHREND	82 CELL	Repl. by BEHREND 89B
<0.5	95	2 BLOCKER	82C MRK2	$E_{\text{cm}}^{\text{ee}} = 29$ GeV
<6.0	95	BRANDELIC	80 TASS	$E_{\text{cm}}^{\text{ee}} = 30$ GeV

$[\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau) + \Gamma(3h^- 2h^+ \geq 0 \text{ neutrals } \nu_\tau \text{ ("5-prong")})]/\Gamma_{\text{total}}$ $(\Gamma_{22} + \Gamma_{36})/\Gamma$

VALUE (%)	EVTs	DOCUMENT ID	TECN	COMMENT
5.75 ± 0.30 OUR FIT				Error includes scale factor of 1.2.
5.4 ± 0.5 OUR AVERAGE				
5.05 ± 0.29 ± 0.65	570	DECAMP	92C ALEP	$E_{\text{cm}}^{\text{ee}} = 88-95$ GeV
5.8 ± 0.7 ± 0.2	352	79 BEHREND	90 CELL	$E_{\text{cm}}^{\text{ee}} = 35$ GeV

79 BEHREND 90 not independent of their $\Gamma(h^- h^- h^+ \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ measurement.

$\Gamma(3h^- 2h^+ \nu_\tau)/\Gamma_{\text{total}}$	EVTs	DOCUMENT ID	TECN	COMMENT
0.056 ± 0.016 OUR AVERAGE				
0.064 ± 0.023 ± 0.01	12	ALBRECHT	88B ARG	$E_{\text{cm}}^{\text{ee}} = 10$ GeV
0.051 ± 0.020	7	BYLSMA	87 HRS	$E_{\text{cm}}^{\text{ee}} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.067 ± 0.030	5	80 BELTRAMI	85 HRS	Repl. by BYLSMA 87
80 The error quoted is statistical only.				

$\Gamma(3h^- 2h^+ \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.051 ± 0.022	6	BYLSMA	87 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.067 ± 0.030	5	⁸¹ BELTRAMI	85 HRS	Repl. by BYLSMA 87
-------------------	---	------------------------	--------	--------------------

⁸¹ The error quoted is statistical only.

$\Gamma(4h^- 3h^+ \geq 0 \text{ neutrals } \nu_\tau (\text{"7-prong"}))/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.019	90	BYLSMA	87 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

$\Gamma(K^*(892)^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.43 \pm 0.11 \pm 0.13$	475	⁸² GOLDBERG	90 CLEO	$E_{\text{cm}}^{\text{ex}} = 9.4\text{--}10.9$ GeV

⁸² GOLDBERG 90 estimates that 10% of observed $K^*(892)$ are accompanied by a π^0 .

$\Gamma(K^*(892)^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.45 ± 0.18 OUR FIT				
1.39 ± 0.20 OUR AVERAGE				

$1.23 \pm 0.21 \pm 0.11$	54	⁸³ ALBRECHT	88L ARG	$E_{\text{cm}}^{\text{ex}} = 10$ GeV
--------------------------	----	------------------------	---------	--------------------------------------

$1.9 \pm 0.3 \pm 0.4$	44	⁸⁴ TSCHIRHART	88 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
-----------------------	----	--------------------------	--------	--------------------------------------

$1.5 \pm 0.4 \pm 0.4$	15	⁸⁵ AIHARA	87C TPC	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
-----------------------	----	----------------------	---------	--------------------------------------

$1.3 \pm 0.3 \pm 0.3$	31	YELTON	86 MRK2	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
-----------------------	----	--------	---------	--------------------------------------

1.7 ± 0.7	11	DORFAN	81 MRK2	$E_{\text{cm}}^{\text{ex}} = 4.2\text{--}6.7$ GeV
---------------	----	--------	---------	---

⁸³ The authors divide by $\Gamma_1/\Gamma = 0.865$ to obtain this result.

⁸⁴ Not independent of TSCHIRHART 88 $\Gamma(\tau^- \rightarrow K^0 h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma(\text{total})$.

⁸⁵ Decay π^- identified in this experiment, is assumed in the others.

$\Gamma(K^*(892)^0 K^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.32 \pm 0.08 \pm 0.12$	119	GOLDBERG	90 CLEO	$E_{\text{cm}}^{\text{ex}} = 9.4\text{--}10.9$ GeV

$\Gamma(K^*(892)^0 \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
$0.38 \pm 0.11 \pm 0.13$	105	GOLDBERG	90 CLEO	$E_{\text{cm}}^{\text{ex}} = 9.4\text{--}10.9$ GeV

$\Gamma(K^0 h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.3 ± 0.3	44	TSCHIRHART	88 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

$\Gamma(K^- K^0 \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{45}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.8	90	GOLDBERG	90 CLEO	$E_{\text{cm}}^{\text{ex}} = 9.4\text{--}10.9$ GeV

$\Gamma(K^0 K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{46}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.26	95	AIHARA	87C TPC	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

$\Gamma(K^0 K^- \geq 1 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.26	95	AIHARA	87C TPC	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

$\Gamma(K^0 h^+ h^- h^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.17	95	TSCHIRHART	88 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.27	90	BELTRAMI	85 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
----------	----	----------	--------	--------------------------------------

$\Gamma(K_2^*(1430)^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE (%)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.3	95		TSCHIRHART	88 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.9	95	0	DORFAN	81 MRK2	$E_{\text{cm}}^{\text{ex}} = 4.2\text{--}6.7$ GeV
---------	----	---	--------	---------	---

$\Gamma(a_0(980)^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}} \times B(a_0(980) \rightarrow K^0 K^-)$ $\Gamma_{50}/\Gamma \times B$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.8 \times 10^{-4}$	90	GOLDBERG	90 CLEO	$E_{\text{cm}}^{\text{ex}} = 9.4\text{--}10.9$ GeV

$\Gamma(\eta \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 1.3	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ex}} \approx 10$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 2.1	90	ABACHI	87B HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
---------	----	--------	---------	--------------------------------------

< 2.1	95	BARINGER	87 CLEO	$E_{\text{cm}}^{\text{ex}} = 10.5$ GeV
---------	----	----------	---------	--

$\Gamma(\eta \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.4	95		ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ex}} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 90	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ex}} \approx 10$ GeV
--------	----	----------	---------	--

< 140	90	BEHREND	88 CELL	$E_{\text{cm}}^{\text{ex}} = 14\text{--}46.8$ GeV
---------	----	---------	---------	---

< 180	95	BARINGER	87 CLEO	$E_{\text{cm}}^{\text{ex}} = 10.5$ GeV
---------	----	----------	---------	--

< 250	90	0	COFFMAN	87 MRK3	$E_{\text{cm}}^{\text{ex}} = 3.77$ GeV
---------	----	---	---------	---------	--

$510 \pm 100 \pm 120$	65	DERRICK	87 HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
-----------------------	----	---------	--------	--------------------------------------

< 100	95	GAN	87B MRK2	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
---------	----	-----	----------	--------------------------------------

$\Gamma(\eta \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$17 \pm 2 \pm 2$	125		ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ex}} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 110	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ex}} \approx 10$ GeV
---------	----	----------	---------	--

< 210	95	BARINGER	87 CLEO	$E_{\text{cm}}^{\text{ex}} = 10.5$ GeV
---------	----	----------	---------	--

$420 \pm 70 \pm 160$		⁸⁶ GAN	87 MRK2	$E_{\text{cm}}^{\text{ex}} = 29$ GeV
----------------------	--	-------------------	---------	--------------------------------------

⁸⁶ Highly correlated with GAN 87 $\Gamma(\pi^- 3\pi^0 \nu_\tau)/\Gamma(\text{total})$ value.

$\Gamma(\eta \pi^- \pi^0 \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.3	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ex}} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 120	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ex}} \approx 10$ GeV
---------	----	----------	---------	--

$\Gamma(\eta K^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 4.7	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ex}} \approx 10.6$ GeV

$\Gamma(\eta \pi^+ \pi^- \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.3	90	ABACHI	87B HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

$\Gamma(\eta \eta \pi^- \geq 0 \text{ neutrals } \nu_\tau)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE (%)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.5	90	ABACHI	87B HRS	$E_{\text{cm}}^{\text{ex}} = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1.5	95	BARINGER	87 CLEO	$E_{\text{cm}}^{\text{ex}} = 10.5$ GeV
---------	----	----------	---------	--

$\Gamma(\eta \eta \pi^- \nu_\tau)/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.1	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ex}} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 83	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ex}} \approx 10$ GeV
--------	----	----------	---------	--

$\Gamma(\eta \eta \pi^- \pi^0 \nu_\tau)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.0	95	ARTUSO	92 CLEO	$E_{\text{cm}}^{\text{ex}} \approx 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 90	95	ALBRECHT	88M ARG	$E_{\text{cm}}^{\text{ex}} \approx 10$ GeV
--------	----	----------	---------	--

$\Gamma(e^- \gamma)/\Gamma_{\text{total}}$ Γ_{60}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.2 \times 10^{-4}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{\text{ex}} = 10$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.0 \times 10^{-4}$	90	KEH	88 CBAL	$E_{\text{cm}}^{\text{ex}} = 10$ GeV
------------------------	----	-----	---------	--------------------------------------

$< 6.4 \times 10^{-4}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{\text{ex}} = 3.8\text{--}6.8$ GeV
------------------------	----	-------	---------	---

$\Gamma(\mu^- \gamma)/\Gamma_{\text{total}}$ Γ_{61}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 0.42 \times 10^{-5}$	90	BEAN	93 CLEO	$E_{\text{cm}}^{\text{ex}} = 10.6$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 3.4 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{\text{ex}} = 10$ C.V
------------------------	----	----------	---------	--------------------------------------

$< 55 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{\text{ex}} = 3.8\text{--}6.8$ GeV
-----------------------	----	-------	---------	---

$\Gamma(e^- \pi^0)/\Gamma_{\text{total}}$ Γ_{62}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 14 \times 10^{-5}$	90	KEH	88 CBAL	$E_{\text{cm}}^{\text{ex}} = 10$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 17 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{\text{cm}}^{\text{ex}} = 10$ GeV
-----------------------	----	----------	---------	--------------------------------------

$< 210 \times 10^{-5}$	90	HAYES	82 MRK2	$E_{\text{cm}}^{\text{ex}} = 3.8\text{--}6.8$ GeV
------------------------	----	-------	---------	---

Lepton & Quark Full Listings

T

$\Gamma(\mu^- \pi^0)/\Gamma_{total}$ Γ_{63}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.4 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 82 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(e^- K^0)/\Gamma_{total}$ Γ_{64}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-3}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(\mu^- K^0)/\Gamma_{total}$ Γ_{65}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.0 \times 10^{-3}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(e^- \eta)/\Gamma_{total}$ Γ_{66}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 24 \times 10^{-5}$	90	KEH 88 CBAL		$E_{cm}^{ee} = 10$ GeV

$\Gamma(\mu^- \eta)/\Gamma_{total}$ Γ_{67}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 7.3 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV

$\Gamma(e^- \rho^0)/\Gamma_{total}$ Γ_{68}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.9 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 3.9 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K
$< 37 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(\mu^- \rho^0)/\Gamma_{total}$ Γ_{69}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.9 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 3.8 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K
$< 44 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(e^- K^*(892)^0)/\Gamma_{total}$ Γ_{70}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.8 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 5.4 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K

$\Gamma(\mu^- K^*(892)^0)/\Gamma_{total}$ Γ_{71}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.5 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 5.9 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K

$\Gamma(\pi^- \gamma)/\Gamma_{total}$ Γ_{72}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 28 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV

$\Gamma(\pi^- \pi^0)/\Gamma_{total}$ Γ_{73}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 37 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV

$\Gamma(\ell^- \ell^+ \ell^+)/\Gamma_{total}$ $\Gamma_{74}/\Gamma = (\Gamma_{75} + \Gamma_{77} + \Gamma_{78} + \Gamma_{80} + \Gamma_{81} + \Gamma_{82})/\Gamma$
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 3.4 \times 10^{-5}$	90	87 BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
$< 3.8 \times 10^{-5}$	90	ALBRECHT 87M ARG		$E_{cm}^{ee} = 10$ GeV

⁸⁷Inclusion of a potentially model-dependent cut on decay track opening angles reduces BOWCOCK 90 limit to 2.6×10^{-5} .

$\Gamma(e^- e^+ e^-)/\Gamma_{total}$ Γ_{75}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.3 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 2.7 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
$< 3.8 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K
$< 40 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma((e\mu\mu^-)/\Gamma_{total}$ $\Gamma_{76}/\Gamma = (\Gamma_{77} + \Gamma_{78})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.7 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(e^- \mu^+ \mu^-)/\Gamma_{total}$ Γ_{77}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.9 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 2.7 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
$< 3.3 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K
$< 33 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(e^+ \mu^- \mu^-)/\Gamma_{total}$ Γ_{78}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 1.8 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
$< 3.8 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K

$\Gamma((\mu e e^-)/\Gamma_{total}$ $\Gamma_{79}/\Gamma = (\Gamma_{80} + \Gamma_{81})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.7 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(\mu^- e^+ e^-)/\Gamma_{total}$ Γ_{80}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 2.7 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
$< 3.3 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K
$< 44 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(\mu^+ e^- e^-)/\Gamma_{total}$ Γ_{81}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.4 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 1.6 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
$< 3.8 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K

$\Gamma(\mu^- \mu^+ \mu^-)/\Gamma_{total}$ Γ_{82}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.7 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 1.9 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
$< 2.9 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K
$< 49 \times 10^{-5}$	90	HAYES 82 MRK2		$E_{cm}^{ee} = 3.8-6.8$ GeV

$\Gamma(\ell^\pm \pi^\mp \pi^-)/\Gamma_{total}$ $\Gamma_{83}/\Gamma = (\Gamma_{85} + \Gamma_{86} + \Gamma_{88} + \Gamma_{89})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.3 \times 10^{-5}$	90	ALBRECHT 87M ARG		$E_{cm}^{ee} = 10$ GeV

$\Gamma(e^\mp \pi^\pm \pi^-)/\Gamma_{total}$ $\Gamma_{84}/\Gamma = (\Gamma_{85} + \Gamma_{86})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 6.0 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(e^- \pi^+ \pi^-)/\Gamma_{total}$ Γ_{85}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.7 \times 10^{-5}$	90	ALBRECHT 92K ARG		$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$< 6.0 \times 10^{-5}$	90	BOWCOCK 90 CLEO		$E_{cm}^{ee} = 10.4-10.9$
$< 4.2 \times 10^{-5}$	90	ALBRECHT 87M ARG		Repl. by AL-BRECHT 92K

$\Gamma(e^+\pi^-\pi^-)/\Gamma_{total}$ Γ_{86}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<1.8 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
$<6.3 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma(\mu^\mp\pi^\pm\pi^-)/\Gamma_{total}$ $\Gamma_{87}/\Gamma = (\Gamma_{88} + \Gamma_{89})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(\mu^-\pi^+\pi^-)/\Gamma_{total}$ Γ_{88}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<3.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
$<4.0 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma(\mu^+\pi^-\pi^-)/\Gamma_{total}$ Γ_{89}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<6.3 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
$<6.3 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma(e^\pm\pi^\mp K^-)/\Gamma_{total}$ $\Gamma_{90}/\Gamma = (\Gamma_{93} + \Gamma_{94} + \Gamma_{95})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-4}$	90	ALBRECHT	87M ARG	$E_{cm}^{ee} = 10$ GeV

$\Gamma((e\pi K)^-, \text{all charged})/\Gamma_{total}$ $\Gamma_{91}/\Gamma = (\Gamma_{93} + \Gamma_{94} + \Gamma_{95})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(e^-\pi^\pm K^\mp)/\Gamma_{total}$ $\Gamma_{92}/\Gamma = (\Gamma_{93} + \Gamma_{94})/\Gamma$
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.8 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(e^-\pi^+ K^-)/\Gamma_{total}$ Γ_{93}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<5.8 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
$<4.2 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma(e^-\pi^- K^+)/\Gamma_{total}$ Γ_{94}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.8 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(e^+\pi^- K^-)/\Gamma_{total}$ Γ_{95}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<4.9 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
$<12 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma((\mu\pi K)^-, \text{all charged})/\Gamma_{total}$ $\Gamma_{96}/\Gamma = (\Gamma_{98} + \Gamma_{99} + \Gamma_{100})/\Gamma$
Test of lepton number or lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(\mu^-\pi^\pm K^\mp)/\Gamma_{total}$ $\Gamma_{97}/\Gamma = (\Gamma_{98} + \Gamma_{99})/\Gamma$
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(\mu^-\pi^+ K^-)/\Gamma_{total}$ Γ_{98}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<11 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
$<12 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma(\mu^-\pi^- K^+)/\Gamma_{total}$ Γ_{99}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.7 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$

$\Gamma(\mu^+\pi^- K^-)/\Gamma_{total}$ Γ_{100}/Γ
Test of lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-5}$	90	BOWCOCK	90 CLEO	$E_{cm}^{ee} = 10.4-10.9$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<5.8 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV
$<12 \times 10^{-5}$	90	ALBRECHT	87M ARG	Repl. by ALBRECHT 92K

$\Gamma(\bar{p}\gamma)/\Gamma_{total}$ Γ_{101}/Γ
Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<29 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV

$\Gamma(\bar{p}\pi^0)/\Gamma_{total}$ Γ_{102}/Γ
Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<66 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV

$\Gamma(\bar{p}\eta)/\Gamma_{total}$ Γ_{103}/Γ
Test of lepton number and baryon number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<130 \times 10^{-5}$	90	ALBRECHT	92K ARG	$E_{cm}^{ee} = 10$ GeV

$\Gamma(e^- \text{light spinless boson})/\Gamma(e^- \nu_e \nu_\tau)$ Γ_{104}/Γ_4
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.018	95	88 ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
<0.040	95	89 BALTRUSAITIS..85	MRK3	$E_{cm}^{ee} = 3.77$ GeV
88 ALBRECHT 90E limit holds for mass < 100 MeV, and rises to 0.050 for mass = 500 MeV.				
89 BALTRUSAITIS 85 limit holds for mass < 100 MeV.				

$\Gamma(\mu^- \text{light spinless boson})/\Gamma(e^- \nu_e \nu_\tau)$ Γ_{105}/Γ_4
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.033	95	90 ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
<0.125	95	91 BALTRUSAITIS..85	MRK3	$E_{cm}^{ee} = 3.77$ GeV
90 ALBRECHT 90E limit holds for mass < 100 MeV, and rises to 0.071 for mass = 500 MeV.				
91 BALTRUSAITIS 85 limit holds for mass < 100 MeV.				

τ DECAY PARAMETERS

NOTE ON τ -DECAY PARAMETERS

Neglecting radiative corrections and terms proportional to m_ℓ^2/m_τ^2 , the energy spectrum of the charged decay lepton ℓ in the τ rest frame is given by

$$\frac{d^2\Gamma_{\tau \rightarrow \ell \nu \bar{\nu}}}{d\Omega dx} \propto x^2 \times \left\{ 12(1-x) + \rho_\tau \left(\frac{32}{3}x - 8 \right) + 24\eta_\tau \frac{m_\ell}{m_\tau} \frac{(1-x)}{x} - P_\tau \xi_\tau \cos\theta \left[4(1-x) + \delta_\tau \left(\frac{32}{3}x - 8 \right) \right] \right\}. \quad (1)$$

Here $x = 2E_\ell/m_\tau$ is the scaled lepton energy, P_τ is the τ polarization, and θ is the angle between the τ spin and the lepton momentum. With unpolarized τ 's or integrating over the full θ range, the spectrum depends only on ρ_τ and η_τ . Measurements of the other two Michel parameters, ξ_τ and δ_τ , require polarized τ 's. Where possible, we give separately the parameters for $\tau^- \rightarrow e^- \nu_\tau \bar{\nu}_e$ and $\tau^- \rightarrow \mu^- \nu_\tau \bar{\nu}_\mu$, to avoid assumptions about universality.

$\rho^\tau(e \text{ or } \mu)$ PARAMETER

($V-A$) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.74 ± 0.04 OUR AVERAGE				
0.742 ± 0.035 ± 0.020	8000	ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.79 ± 0.10 ± 0.10	3732	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.71 ± 0.09 ± 0.03	1426	BEHRENDIS	85 CLEO	e^+e^- near $\Upsilon(4S)$

Lepton & Quark Full Listings

τ , Number of Light Neutrino Types

$\rho^\tau(e)$ PARAMETER

(V-A) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.04	OUR AVERAGE			
0.79 ± 0.08 ± 0.06	3230	ALBRECHT	93G ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.747 ± 0.045 ± 0.028	5106	ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.64 ± 0.06 ± 0.07	2753	JANSSEN	89 CBAL	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.62 ± 0.17 ± 0.14	1823	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.60 ± 0.13	699	BEHREND	85 CLEO	e^+e^- near $\Upsilon(4S)$
0.72 ± 0.10 ± 0.11	594	BACINO	79B DLCO	$E_{cm}^{ee} = 3.5-7.4$ GeV

$\rho^\tau(\mu)$ PARAMETER

(V-A) theory predicts $\rho = 0.75$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.76 ± 0.05	OUR AVERAGE			
0.76 ± 0.07 ± 0.08	3230	ALBRECHT	93G ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.734 ± 0.055 ± 0.027	3041	ALBRECHT	90E ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
0.89 ± 0.14 ± 0.08	1909	FORD	87B MAC	$E_{cm}^{ee} = 29$ GeV
0.81 ± 0.13	727	BEHREND	85 CLEO	e^+e^- near $\Upsilon(4S)$

$\xi^\tau(e \text{ or } \mu)$ PARAMETER

(V-A) theory predicts $\xi = 1$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.90 ± 0.15 ± 0.10	3230	92 ALBRECHT	93G ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV

92 ALBRECHT 93G measurement determines $|\xi^\tau|$ for the case $\xi^\tau(e) = \xi^\tau(\mu)$, but the authors point out that other LEP experiments determine the sign to be positive.

CONSTRAINT ON W - τ COUPLINGS $2g_{ABV}(g_A^2 + g_V^2)$

Standard Model predicts $g_A = g_V = 1$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.25 ± 0.23 ± 0.15 -0.08	7.5k	ALBRECHT	93C ARG	$E_{cm}^{ee} = 9.4-10.6$ GeV
1.14 ± 0.34 ± 0.34 -0.17	3.9k	ALBRECHT	90I ARG	Repl. by ALBRECHT 93C

• • • We do not use the following data for averages, fits, limits, etc. • • •

τ REFERENCES

AKERS 94E PL B328 207	+Alexander, Allison, Anderson+	(OPAL Collab.)
ARTUSO 94 PRL 72 3762	+Goldberg, He, Horwitz+	(CLEO Collab.)
ABREU 93E PL B302 356	+Adam, Adye, Agasi+	(DELPHI Collab.)
ACTON 93J ZPHY C59 183	+Alexander, Allison, Allport+	(OPAL Collab.)
ADRIANI 93M PRP 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+	(L3 Collab.)
ALBRECHT 93C ZPHY C58 61	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
ALBRECHT 93G PL B316 608	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
BALEST 93 PR D47 R3671	+Daoudi, Ford, Johnson+	(CLEO Collab.)
BEAN 93 PRL 70 138	+Gronberg, Kutschke+	(CLEO Collab.)
BORTOLETTO 93 PRL 71 1791	+Brown, Fast, McIlwain+	(CLEO Collab.)
ESCRIBANO 93 PL B301 419	+Masso	(BARC)
PROCARIO 93 PRL 70 1207	+Yang, Balest, Cho+	(CLEO Collab.)
WASSERBAECH 93 PR D48 4216		(FSU/C)
ABREU 92Z ZPHY C53 567	+Adam, Adami, Adye+	(DELPHI Collab.)
ABREU 92N ZPHY C55 555	+Adam, Adye, Agasi+	(DELPHI Collab.)
ACTON 92F PL B281 405	+Alexander, Allison, Allport+	(OPAL Collab.)
ACTON 92H PL B288 373	+Allison, Allport+	(OPAL Collab.)
AKERIB 92 PRL 69 3610	+Barish, Chadha, Cowen+	(CLEO Collab.)
Also 93B PRL 71 3395 (erratum)	Akerib, Barish, Chadha, Cowen+	(CLEO Collab.)
ALBRECHT 92D ZPHY C53 367	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
ALBRECHT 92K ZPHY C55 179	+Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALBRECHT 92M PL B292 221	+Ehrlichmann, Hamacher, Hofmann+	(ARGUS Collab.)
ALBRECHT 92Q ZPHY C56 339	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
AMMAR 92 PR D45 3976	+Baringer, Coppage, Davis+	(CLEO Collab.)
ARTUSO 92 PRL 69 3278	+Goldberg, Horwitz, Kennett+	(CLEO Collab.)
BAI 92 PRL 69 3021	+Bardon, Becker-Szendy, Burnett+	(BES Collab.)
BATTLE 92 PL B291 488	+Ernst, Kroha, Roberts+	(CLEO Collab.)
BUSKULIC 92H PL B297 432	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
BUSKULIC 92I PL B299 159	+Ernst, Goy, Lees	(ALEPH Collab.)
DECAMP 92C ZPHY C54 211	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
DECAMP 92E PL B279 411	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
ABREU 91D PL B267 422	+Adam, Adami, Adye+	(DELPHI Collab.)
ACTON 91C PL B273 355	+Alexander, Allison, Allport+	(OPAL Collab.)
ADEVA 91F PL B265 451	+Adriani, Aguiar-Benitez, Akbari+	(L3 Collab.)
ALBRECHT 91D PL B260 259	+Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALEXANDER 91D PL B266 201	+Allison, Allport, Anderson+	(OPAL Collab.)
ANTREASIAN 91 PL B259 216	+Bartels, Besset, Bieler+	(Crystal Ball Collab.)
GRIFOLS 91 PL B255 611	+Mendez	(Crystal Ball Collab.)
SAMUEL 91B PRL 67 668	+Li, Mendel	(OKSU, WONT)
Also 92B PRL 69 995	Samuel, Li, Mendel	(OKSU, WONT)
Erratum.		
ABACHI 90 PR D41 1414	+Derrick, Kooijman, Musgrave+	(HRJ Collab.)
ALBRECHT 90E PL B246 278	+Ehrlichmann, Harder, Krueger+	(ARGUS Collab.)
ALBRECHT 90I PL B250 164	+Ehrlichmann, Harder, Krueger+	(ARGUS Collab.)
BEHREND 90 ZPHY C46 537	+Criegee, Field, Franke+	(CELLO Collab.)
BOWCOCK 90 PR D41 805	+Kinoshita, Pipkin, Procaro+	(CLEO Collab.)
DELAGUILA 90 PL B252 116	+Shor	(BARC, WILL)
GOLDBERG 90 PL B251 223	+Haupt, Horwitz, Jain+	(CLEO Collab.)
WU 90 PR D41 2339	+Hayes, Perli, Barklow+	(Mark II Collab.)
ABACHI 89B PR D40 902	+Derrick, Kooijman, Musgrave+	(ARGUS Collab.)
BEHREND 89 PL B227 1659	+Trilling, Abrams, Field, Franke+	(CELLO Collab.)
JANSSEN 89 PL B228 273	+Antreasian, Bartels, Besset+	(Crystal Ball Collab.)
KLEINWORT 89 ZPHY C42 7	+Allison, Ambrus, Barlow+	(JADE Collab.)
ADEVA 88 PR D38 2665	+Andershub, Ansari, Becker+	(Mark-J Collab.)
ALBRECHT 88B PL B202 149	+Binder, Boeckmann+	(ARGUS Collab.)
ALBRECHT 88M ZPHY C41 1	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
ALBRECHT 88L PL B201 921	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
ALBRECHT 88M PR D37 1350	+Trilling, Abrams, Field, Franke+	(Mark II Collab.)
BEHREND 88 PL B200 226	+Criegee, Dainton, Field+	(CELLO Collab.)
BRUNSCHE... 88C ZPHY C39 331	+Braunschweig, Kirschfink, Martyn+	(TASSO Collab.)
KEH 88 PL B212 123	+Antreasian, Bartels, Besset+	(Crystal Ball Collab.)
TSCHIRHART 88 PL B205 407	+Abachi, Akerlof, Baringer+	(HRJ Collab.)
ABACHI 87B PL B197 291	+Baringer, Bylsma, De Bonte+	(HRJ Collab.)
ABACHI 87C PRL 59 2519	+Akerlof, Baringer, Blockus+	(HRJ Collab.)
ADLER 87B PRL 59 1527	+Becker, Blaylock, Bolton+	(Mark III Collab.)

AIHARA 87B PR D35 1553	+Alston-Garnjost, Avery+	(TPC Collab.)
AIHARA 87C PRL 59 751	+Alston-Garnjost, Avery+	(TPC Collab.)
ALBRECHT 87L PL B185 223	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 87M PL B185 228	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 87P PL B199 580	+Andam, Binder, Boeckmann+	(ARGUS Collab.)
BAND 87 PL B198 297	+Camporesi, Chadwick, Delfino+	(MAC Collab.)
BAND 87B PRL 59 415	+Bosman, Camporesi, Chadwick+	(MAC Collab.)
BARINGER 87C PRL 59 1993	+McIlwain, Miller, Shibata+	(CLEO Collab.)
BEBEK 87C PR D36 690	+Berkelman, Blucher, Cassel+	(CLEO Collab.)
BURCHAT 87 PR D35 27	+Feldman, Barklow, Boyarski+	(Mark II Collab.)
BYLSMA 87 PR D35 2269	+Abachi, Baringer, DeBonte+	(HRJ Collab.)
COFFMAN 87 PR D36 2185	+Dubois, Eigen, Hauser+	(Mark III Collab.)
DERRICK 87 PL B189 260	+Kooijman, Loos, Musgrave+	(HRJ Collab.)
FORD 87 PR D35 408	+Qi, Read, Smith+	(MAC Collab.)
FORD 87B PR D36 1971	+Qi, Read, Smith+	(MAC Collab.)
GAN 87 PRL 59 411	+Abrams, Amidei, Baden+	(Mark II Collab.)
GAN 87B PL B197 561	+Abrams, Amidei, Baden+	(Mark II Collab.)
ADEVA 86B PL B179 177	+Ansari, Becker, Becker-Szendy+	(Mark-J Collab.)
AIHARA 86E PRL 57 1836	+Alston-Garnjost, Avery+	(TPC Collab.)
BARTEL 86D PL B182 216	+Becker, Felst, Haidt, Knies+	(JADE Collab.)
PDG 86 PL 170B	+Aguilar-Benitez, Porter+	(CERN, CIT+)
RUCKSTUHL 86 PRL 56 2132	+Stroynowski, Atwood, Barish+	(DELCO Collab.)
SCHMIDKE 86 PRL 57 527	+Abrams, Matteuzzi, Amidei+	(Mark II Collab.)
YELTON 86 PRL 56 812	+Dorfan, Abrams, Amidei+	(Mark II Collab.)
AKERLOF 85B PRL 55 570	+Baranko, Baringer, Beltrami+	(HRJ Collab.)
ALTHOFF 85 ZPHY C26 521	+Braunschweig, Kirschfink+	(TASSO Collab.)
ASH 85B PRL 55 2118	+Band, Blume, Camporesi+	(MAC Collab.)
BALTRUSAIT... 85 PRL 55 1842	+Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BARTEL 85F PL 161B 188	+Becker, Cords, Felst+	(JADE Collab.)
BEHREND 85 PR D32 2468	+Gentile, Guida, Guida, Morrow+	(CLEO Collab.)
BELTRAMI 85 PRL 54 1775	+Bylsma, DeBonte, Gan+	(HRJ Collab.)
BERGER 85 ZPHY C28 11	+Genzel, Lackas, Pleier+	(PLUTO Collab.)
BURCHAT 85 PRL 54 2489	+Schmidke, Yelton, Abrams+	(Mark II Collab.)
FERNANDEZ 85 PRL 54 1624	+Ford, Qi, Read+	(MAC Collab.)
MILLS 85 PRL 54 624	+Pal, Atwood, Bailion+	(DELCO Collab.)
AIHARA 84C PR D30 2436	+Alston-Garnjost, Badtke, Bakken+	(TPC Collab.)
ALTHOFF 84D PL 141B 264	+Braunschweig, Kirschfink+	(TASSO Collab.)
BEHREND 84 ZPHY C23 103	+Fenner, Schachter, Schroeder+	(CELLO Collab.)
MILLS 84 PRL 52 1944	+Ruckstuhl, Atwood, Bailion+	(DELCO Collab.)
BEHREND 83C PL 127B 370	+Chen, Fenner, Gumpel+	(CELLO Collab.)
JAROS 83 PRL 51 955	+Amidei, Trilling, Abrams+	(Mark II Collab.)
SILVERMAN 83 PR D27 1196	+Shaw	(UCI)
BEHREND 82 PL 114B 282	+Chen, Fenner, Field+	(CELLO Collab.)
BLOCKER 82B PRL 48 1586	+Abrams, Alam, Bondel+	(Mark II Collab.)
BLOCKER 82C PRL 49 1369	+Levi, Abrams, Amidei+	(Mark II Collab.)
BLOCKER 82D PL 109B 119	+Dorfan, Abrams, Alam+	(Mark II Collab.)
HAYES 82 PR D25 2869	+Peri, Alam, Boyarski+	(Mark II Collab.)
BERGER 81B PL 99B 489	+Genzel, Grigull, Lackas+	(PLUTO Collab.)
DORFAN 81 PRL 46 215	+Blocker, Abrams, Alam+	(Mark II Collab.)
BLOCKER 80 Thesis LBL-10801		(LBL)
BRANDELIK 80 PL 92B 199	+Braunschweig, Gather+	(TASSO Collab.)
ZHOLENTZ 80 PL 96B 214	+Kurdadze, Leichuk, Mishnev+	(NOVO)
Also 81 SJNP 34 814	Zholentz, Kurdadze, Leichuk+	(NOVO)
BACINO 79B PRL 42 749	+Ferguson, Nodulman, Slater+	(DELCO Collab.)
KIRKBY 79 SLAC-PUB-2419		(SLAC) J
Batavia Lepton Photon Conference.		
BACINO 78B PRL 41 13	+Ferguson, Nodulman, Slater+	(DELCO Collab.) J
Also 78B Tokyo Conf. 249	Kirz	(STON)
Also 78B PRL 37 214	Zholentz, Kurdadze, Leichuk, Mishnev+	(NOVO)
BARTEL 78 PL 77B 331	+Dittmann, Duinker, Olsson, Oneill+	(DESY, HEIDP)
BRANDELIK 78 PL 73B 109	+Braunschweig, Martyn, Sander+	(DASP Collab.) J
FELDMAN 78 Tokyo Conf. 777		(SLAC) J
HEILE 78 NP B138 189	+Peri, Abrams, Alam, Boyarski+	(SLAC, LBL)
JAROS 78 PRL 40 1120	+Abrams, Alam+	(SLAC, LBL, NWES, HAWA)
PERL 75 PRL 35 1489	+Abrams, Boyarski, Breidenbach+	(LBL, SLAC)

OTHER RELATED PAPERS

WEINSTEIN 93 ARNPS 43 457	+Stroynowski	(CIT, SMU)
PERL 92 RPP 55 653		(SLAC)
PICH 90 MPL A5 1995		(VALE)
BARSH 88 PRL 61 1571	+Stroynowski	(CIT)
GAN 88 JMP A3 531	+Perli	(SLAC)
HAYES 88 PR D38 3351	+Perli	(SLAC)
PERL 80 ARNPS 30 299		(SLAC)

Number of Light Neutrino Types

The neutrinos referred to in this section are those of the Standard $SU(2) \times U(1)$ Electroweak Model possibly extended to allow nonzero neutrino masses. Light neutrinos are those with $m_\nu \ll m_{Z_0}$. The limits are on the number of neutrino families or species, including ν_e, ν_μ, ν_τ

NOTE ON THE NUMBER OF LIGHT NEUTRINO TYPES FROM COLLIDER EXPERIMENTS

(by Dean Karlen, Carleton University)

The most precise measurements of the number of light neutrino types, N_ν , come from studies of Z production in e^+e^- collisions. At the time of this report, the most recent combined analysis of the four LEP experiments [1] included nearly 8 million visible Z decays. The invisible partial width, Γ_{inv} , is determined from these data by subtracting the measured visible partial widths, corresponding to Z decays into quarks and charged leptons, from the total Z width. The invisible width is assumed to be due to N_ν light neutrino species each contributing the neutrino partial width Γ_ν as given by the

See key on page 1343

Lepton & Quark Full Listings Number of Light Neutrino Types

Standard Model. The Standard Model value for Γ_ν , however, is uncertain by about 1% due to the unknown top quark mass. In order to reduce this uncertainty, the Standard Model value for the ratio of the neutrino to charged leptonic partial widths, $(\Gamma_\nu/\Gamma_\ell)_{SM} = 1.992 \pm 0.003$, is used instead to determine the number of light neutrino types:

$$N_\nu = \frac{\Gamma_{inv}}{\Gamma_\ell} \left(\frac{\Gamma_\ell}{\Gamma_\nu} \right)_{SM}$$

The combined LEP result is $N_\nu = 2.985 \pm 0.023 \pm 0.004$, where the first error is the combined statistical and systematic uncertainty and the second is the uncertainty from allowing the top quark mass to vary between 100 and 200 GeV.

In the past, when only small samples of Z decays had been recorded by the LEP experiments and by the Mark II at SLC, the uncertainty in N_ν was reduced by using Standard Model fits to the measured hadronic cross sections at several center-of-mass energies near the Z resonance. Since this method is much more dependent on the Standard Model and the top quark mass, the approach described above is favored.

Before the advent of the SLC and LEP, limits on the number of neutrino generations were placed by experiments at lower-energy e^+e^- colliders by measuring the cross section of the process $e^+e^- \rightarrow \nu\bar{\nu}\gamma$. The ASP, CELLO, MAC, MARK J, and VENUS experiments observed a total of 3.9 events above background [2], leading to a 95% CL limit of $N_\nu < 4.8$. This process has since been measured at LEP by the ALEPH, L3, and OPAL experiments [3], and the combined result is $N_\nu = 2.97 \pm 0.17$.

Experiments at $p\bar{p}$ colliders also placed limits on N_ν by determining the total Z width from the observed ratio of $W^\pm \rightarrow \ell^\pm\nu$ to $Z \rightarrow \ell^+\ell^-$ events [5]. This involved a calculation that assumed Standard Model values for the total W width and the ratio of W and Z leptonic partial widths, and used an estimate of the ratio of Z to W production cross sections. Now that the Z width is very precisely known from the LEP experiments, the approach is now one of those used to determine the W width.

References

1. The LEP Electroweak Working Group, LEPEWWG/94-01.
2. C. Hearty *et al.*, Phys. Rev. **D39**, 3207 (1989);
H.J. Behrend *et al.*, Phys. Lett. **B215**, 186 (1988);
W.T. Ford *et al.*, Phys. Rev. **D33**, 3472 (1986);
H. Wu, Ph.D. Thesis, Univ. Hamburg (1986);
K. Abe *et al.*, Phys. Lett. **B232**, 431 (1989).
3. D. Buskulic *et al.*, Phys. Lett. **B313**, 520 (1993);
O. Adriani *et al.*, Phys. Lett. **B292**, 463 (1992);
B. Adeva *et al.*, Phys. Lett. **B275**, 209 (1992);
M.Z. Akrawy *et al.*, Z. Phys. **C50**, 373 (1991).
4. C. Albajar *et al.*, Phys. Lett. **B198**, 271 (1987).
5. R. Ansari *et al.*, Phys. Lett. **B186**, 440 (1987).

Number from e^+e^- Colliders

Number of Light ν Types

Our evaluation uses the invisible and leptonic widths of the Z boson from our combined fit shown in the Full Listings for the Z Boson, and the Standard Model value $\Gamma_\nu/\Gamma_\ell = 1.992 \pm 0.003$.

VALUE	DOCUMENT ID	TECN	COMMENT
2.983 ± 0.025 OUR EVALUATION	Combined fit to all Z data.		
2.994 ± 0.023 OUR AVERAGE	Error Includes scale factor of 1.1.		
3.057 ± 0.040	^{1,2} ABREU	94 DLPH	1990–1992 LEP runs
2.981 ± 0.050	^{1,2} ACCIARRI	94 L3	1990–1992 LEP runs
2.946 ± 0.045	^{1,2} AKERS	94 OPAL	1989–1992 LEP runs
2.983 ± 0.034	^{1,2} BUSKULIC	94 ALEP	1989–1992 LEP runs
2.8 ± 0.6	³ ABRAMS	89B MRK2	$E_{cm}^e = 91$ GeV at SLC
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.97 ± 0.05	BUSKULIC	93J ALEP	Repl. by BUSKULIC 94
2.97 ± 0.07	¹ DECAMP	92B ALEP	Repl. by BUSKULIC 93J
3.00 ± 0.05	⁴ LEP	92 RVUE	
2.93 ± 0.04 ± 0.07	¹ ABREU	91F DLPH	Repl. by ABREU 94
3.05 ± 0.10	¹ ADEVA	91E L3	Repl. by ACCIARRI 94
3.05 ± 0.09 ± 0.005	^{1,5} ALEXANDER	91F OPAL	Repl. by AKERS 94
3.12 ± 0.24 ± 0.25	AARNIO	90 DLPH	
2.97 ± 0.26	³ ABREU	90 DLPH	
3.23 ± 0.29	⁶ ADEVA	90D L3	
3.01 ± 0.11	ADEVA	90I L3	
3.3 ± 0.7	⁶ AKRAWY	90 OPAL	
2.73 ± 0.26 +0.02 -0.04	^{6,7} AKRAWY	90E OPAL	
3.09 ± 0.19 +0.06 -0.12	^{3,7} AKRAWY	90E OPAL	
3.35 ± 0.41	⁶ DECAMP	90B ALEP	
3.01 ± 0.15 ± 0.05	^{3,7} DECAMP	90D ALEP	
2.91 ± 0.13	DECAMP	90P ALEP	
2.4 ± 0.4 ± 0.5	³ AARNIO	89 DLPH	

¹ Simultaneous fits to all measured Z data.

² Analysis based on ≥ 1.1 million visible Z decays from LEP runs through 1992.

³ These papers assume standard model couplings.

⁴ Simultaneous fits to all measured cross section data from all four LEP experiments.

⁵ Second error is from uncertainty in top and Higgs mass.

⁶ These papers measure leptonic widths and are more model independent. However, they divide the measured invisible width via the Standard Model width for neutrinos. Such results are less precise, as discussed in the minireview.

⁷ The second error is due to theoretical uncertainties.

Number of Light ν Types from Direct Measurement of Invisible Z Width

In the following, the invisible Z width is obtained from studies of single-photon events from the reaction $Z \rightarrow \nu\bar{\nu}\gamma$. All are obtained from LEP runs in the E_{cm}^e range 88–94 GeV.

VALUE	DOCUMENT ID	TECN	COMMENT
2.97 ± 0.17 OUR AVERAGE			
2.68 ± 0.20 ± 0.20	BUSKULIC	93L ALEP	1990–1991 LEP runs
3.24 ± 0.46 ± 0.22	ADEVA	92 L3	1990 LEP run
3.14 ± 0.24 ± 0.12	ADRIANI	92E L3	1991 LEP run
3.0 ± 0.4 ± 0.2	AKRAWY	91D OPAL	

Limits from Astrophysics and Cosmology

Number of Light ν Types

("light" means $<$ about 1 MeV). See also OLIVE 81. For a review of limits based on Nucleosynthesis, Supernovae, and also on terrestrial experiments, see DENEGRI 90.

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 3.3	WALKER	91 COSM	
< 3.4	OLIVE	90 COSM	
< 5.2	ELLIS	86 COSM	
< 4	STEIGMAN	86 COSM	
< 4	YANG	84 COSM	
< 4	YANG	79 COSM	
< 7	STEIGMAN	77 COSM	
< 16	PEEBLES	71 COSM	
	⁸ SHVARTSMAN	69 COSM	
	HOYLE	64 COSM	

⁸ SHVARTSMAN 69 limit inferred from his equations.

Number Coupling with Less Than Full Weak Strength

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 20	⁹ OLIVE	81C COSM	
< 20	⁹ STEIGMAN	79 COSM	

⁹ Limit varies with strength of coupling. See also WALKER 91.

See key on page 1343

Lepton & Quark Full Listings

Heavy Lepton Searches

References

1. M.L. Perl and P. Rapidis, SLAC-PUB-1496 (October 1974).
2. J.D. Bjorken and C.H. Llewellyn Smith, Phys. Rev. **D7**, 887 (1973).
3. M. Perl, SLAC-PUB-2752 (1981).

Limits apply only to heavy lepton types specified. See review above for description of types. $L, e_P, \mu_P, \tau_P, e_O, \mu_O$, and τ_O denote sequential lepton, para-electron, para-muon, para-tau, ortho-electron, ortho-muon, and ortho-tau, respectively. As noted, limits for excited leptons (e^* , μ^* , τ^*) are included in the section on "Searches for Quark and Lepton Compositeness."

Charged Heavy Lepton MASS LIMITS

Sequential Charged Heavy Lepton (L^\pm) MASS LIMITS

These experiments assumed that a fourth generation L^\pm decayed to a fourth generation ν_L (or L^0) where ν_L was stable. New data show that stable ν_L have $m_{\nu_L} > 42.7$ GeV so that the above assumption is not valid for any mass limit ≤ 42.7 GeV. One can instead assume that L^\pm decays via mixing to ν_e, ν_μ and/or ν_τ , and in that context the limits below are meaningful.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>42.8	95	ADEVA	90s L3	Dirac
>44.3	95	AKRAWY	90G OPAL	
>42.7	95	DECAMP	90F ALEP	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 12.6–29.6	95	KIM	91B AMY	Massless ν assumed
none 0.5–10	95	1 RILES	90 MRK2	For $(m_{L^-} - m_{L^0}) > 0.25-0.4$ GeV
> 8		2 STOKER	89 MRK2	For $(m_{L^+} - m_{L^0}) = 0.4$ GeV
>12		2 STOKER	89 MRK2	For $m_{L^0} = 0.9$ GeV
>27.6	95	3 ABE	88 VNS	
>25.5	95	4 ADACHI	88B TOPZ	
none 1.5–22.0	95	BEHREND	88C CELL	
>25.0	95	5 IGARASHI	88 AMY	
>27.6	95	6 KIM	88 AMY	
>41	90	7 ALBAJAR	87B UA1	
>25.0	95	YOSHIDA	87B VNS	
>22.5	95	8 ADEVA	85 MRKJ	
>18.	95	9 ADEVA	83B MRKJ	
>18.0	95	10 BARTEL	83 JADE	
>14.	95	ADEVA	82 MRKJ	
none 4–14.5	95	11 BERGER	81B PLUT	
>15.5	95	12 BRANDELIK	81 TASS	
>13.		13 AZIMOV	80	
>16.	95	14 BARBER	80B CNTR	
> 0.490		15 ROTHE	69 RVUE	

- 1 RILES 90 limits were the result of a special analysis of the data in the case where the mass difference $m_{L^-} - m_{L^0}$ was allowed to be quite small, where L^0 denotes the neutrino into which the sequential charged lepton decays. With a slightly reduced m_{L^\pm} range, the mass difference extends to about 4 GeV.
- 2 STOKER 89 (Mark II at PEP) gives bounds on charged heavy lepton (L^+) mass for the generalized case in which the corresponding neutral heavy lepton (L^0) in the SU(2) doublet is not of negligible mass.
- 3 ABE 88 search for L^+ and $L^- \rightarrow$ hadrons looking for acoplanar jets. The bound is valid for $m_\nu < 10$ GeV.
- 4 ADACHI 88B search for hadronic decays giving acoplanar events with large missing energy. $E_{cm}^{ee} = 52$ GeV.
- 5 IGARASHI 88 search for multi-hadron events with isolated leptons. $E_{cm}^{ee} = 50-52$ GeV.
- 6 KIM 88 search for $L^\pm \rightarrow$ hadrons with $L^\mp \rightarrow$ isolated lepton X and for L^\pm and $L^\mp \rightarrow$ hadrons. $E_{cm}^{ee} = 56$ GeV.
- 7 Assumes associated neutrino is approximately massless.
- 8 ADEVA 85 analyze one-isolated-muon data and sensitive to $\tau < 10$ nanosec. Assume $B(\text{lepton}) = 0.30$. $E_{cm} = 40-47$ GeV.
- 9 ADEVA 83B looked for muon opposite against a hadron jet.
- 10 BARTEL 83 limit is from PETRA e^+e^- experiment with average $E_{cm} = 34.2$ GeV.
- 11 BERGER 81B is DESY DORIS and PETRA experiment. Looking for $e^+e^- \rightarrow L^+L^-$.
- 12 BRANDELIK 81 is DESY-PETRA experiment. Looking for $e^+e^- \rightarrow L^+L^-$.
- 13 AZIMOV 80 estimated probabilities for $M + N$ type events in $e^+e^- \rightarrow L^+L^-$ deducing semi-hadronic decay multiplicities of L from e^+e^- annihilation data at $E_{cm} = (2/3)m_L$. Obtained above limit comparing these with e^+e^- data (BRANDELIK 80).
- 14 BARBER 80B looked for $e^+e^- \rightarrow L^+L^-, L^- \rightarrow \nu_L^+ X$ with MARK-J at DESY-PETRA.
- 15 ROTHE 69 examines previous data on μ pair production and π and K decays.

Stable Charged Heavy Lepton (L^\pm) MASS LIMITS

VALUE (GeV)	CL%	DOCUMENT ID	TECN
>42.8 (CL = 95%) OUR LIMIT			
>28.2	95	16 ADACHI	90C TOPZ
none 18.5–42.8	95	AKRAWY	90G OPAL
>26.5	95	DECAMP	90F ALEP
none $m_\mu - 36.3$	95	SODERSTROM90	MRK2

¹⁶ADACHI 90C put lower limits on the mass of stable charged particles with electric charge Q satisfying $2/3 < Q/e < 4/3$ and with spin 0 or 1/2. We list here the special case for a stable charged heavy lepton.

Charged Ortho-Electron (e_O^\pm) MASS LIMITS

See also the section "MASS LIMITS for Excited e " in the section on "Searches for Quark and Lepton Compositeness."

VALUE (GeV)	EVTS	DOCUMENT ID	TECN	CHG
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 0.25–2.3		17 BACCI	77B SPEC	\pm
>0.6	0	18 BACCI	73 ELEC	\pm
>2.2	0	18 BACCI	73 ELEC	\pm
none 0.263–1.32		19 LICHTENSTEIN70	SPEC	–
none 0.1–1.3		20 BOLEY	68 SPEC	–
none 0.3–0.7		21 BUDNITZ	66 SPEC	–
>1.0	0	22 BEHREND	65 SPEC	–
none 0.12–0.57		23 BÉTOURNE	65 SPEC	–

¹⁷BACCI 77B is same type as BACCI 73. Lower mass limit corresponds to λ^2 limit of 4×10^{-5} , upper value is for λ^2 limit of 1.5×10^{-3} .

¹⁸BACCI 73 is Frascati e^+e^- experiment. Looks for $e_O \rightarrow e\gamma$. Mass limit depends on coupling constant λ for this decay. First value above is for $\lambda^2 > 9 \times 10^{-5}$, second is for $\lambda^2 > 10^{-3}$.

¹⁹LICHTENSTEIN 70 is Cornell experiment measuring e Bremsstrahlung. Mass limit depends on coupling constant. First value above is for $\lambda^2 > 0.17$, second is for $\lambda^2 > 0.42$.

²⁰BOLEY 68 is CEA experiment. Looks for $e_P \rightarrow e_O p$. Mass of 0.1 corresponds to coupling constant $\lambda^2 > 3 \times 10^{-4}$, mass limit of 1.3 to $\lambda^2 > 0.01$.

²¹BUDNITZ 66 is CEA experiment. Looks for $e_P \rightarrow e_O p$.

²²BEHREND 65 is DESY experiment. Looks for $e_P \rightarrow e_O p, e_O \rightarrow e\gamma$. This mass limit corresponds to a limit on λ^2 of 6.25×10^{-4} .

²³BÉTOURNE 65 is Orsay experiment. Looks for $e_P \rightarrow e_O p$. Mass of 0.12 corresponds to coupling constant $\lambda^2 > 0.0016$, mass of 0.57 to $\lambda^2 > 0.22$.

Charged Ortho-Muon (μ_O^\pm) MASS LIMITS

See also the section "MASS LIMITS for Excited μ " in the section on "Searches for Quark and Lepton Compositeness."

VALUE (GeV)	CL%	EVTS	DOCUMENT ID	TECN	CHG
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>10.3	98	24 ASRATYAN	78	–	
> 7.5	0	25 CNOPS	78 HLBC	–	
> 1.8	90	26 ASRATYAN	74 HLBC	\pm	
none 0–2.0		27 GITTLESON	74 SPEC		
none 0.2–0.6		28 LIBERMAN	69 OSPK	–	

²⁴ASRATYAN 78 analyzes dependence of (neutral current/charged current) on energy of associated hadrons. Uses data of HOLDER 77 — ν_μ interactions at CERN-SPS.

²⁵CNOPS 78 is FNAL experiment looking for $\nu_\mu \text{Ne} \rightarrow L^\pm$, followed by $L^\pm \rightarrow e^\pm \nu_\nu$.

²⁶ASRATYAN 74 uses EICHTEN 73 data on ν nucleon $\rightarrow e^-$ hadrons and $\bar{\nu}$ nucleon $\rightarrow e^+$ hadrons to set limits on orthomuon production.

²⁷GITTLESON 74 is $\mu P \rightarrow p \mu_O$ search. Coupling constant λ^2 is < 0.01 for mass up to 0.7 GeV, limit on λ^2 rises to < 0.1 for mass of 2.0 GeV.

²⁸LIBERMAN 69 is a BNL experiment measuring muon Bremsstrahlung.

Charged Para-Muon (μ_P^\pm) MASS LIMITS

VALUE (GeV)	CL%	EVTS	DOCUMENT ID	TECN	CHG
• • • We do not use the following data for averages, fits, limits, etc. • • •					
> 9.0	0	29 CNOPS	78 HLBC	+	
>10.0		30 ERRIQUEZ	78 BEBC	+	
>12.	90	31 HOLDER	78 CNTR	+	
> 8.4	90	32 BARISH	74 SPEC	+	
> 2.0	90	33 BARISH	73B ASPK	+	
> 2.4	90	34 EICHTEN	73 HLBC	+	

²⁹CNOPS 78 is FNAL experiment looking for $\nu_\mu \text{Ne} \rightarrow L^\pm$, followed by $L^\pm \rightarrow e^\pm \nu_\nu$.

³⁰ERRIQUEZ 78 is CERN SPS experiment. Looks for ν_μ nucleon $\rightarrow \mu^- e^+ X$. Finds cross section for producing heavy lepton $\rightarrow e^+ < 0.7 \times 10^{-3} \times CC$ cross section.

³¹HOLDER 78 is a CERN ν experiment looking for ν_μ nucleon $\rightarrow \mu^+$ anything. Assumes $\mu_P^+ \rightarrow \mu^+ 2\nu_\mu$ with BR = 0.2.

³²BARISH 74 is FNAL 50,135 GeV ν experiment. Looks for $(\nu \text{ nucleon} \rightarrow \mu_P^+ X)$. Assumes $(\mu_P^+ \rightarrow \mu^+ \nu_\mu \nu_\mu)$ with BR = 0.3.

³³BARISH 73B is FNAL 50,145 GeV ν experiment. Looks for $(\nu \text{ nucleon} \rightarrow \mu_P^+ X)$. Assumes $(\mu_P^+ \rightarrow \mu^+ \nu_\mu \nu_\mu)$ with BR = 0.3.

³⁴EICHTEN 73 is CERN 1–10 GeV ν experiment. Looks for μ_P^+ produced in ν nucleon $\rightarrow \mu_P^+$ hadrons assuming 15% decay to $e^+ \nu_\mu \nu_e$.

Lepton & Quark Full Listings

Heavy Lepton Searches

Charged Long-Lived Heavy Lepton MASS LIMITS

VALUE (GeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
>0.1	0	35 ANSORGE	73B	HBC	Long-lived
none 0.55-4.5		36 BUSHNIN	73	CNTR	Long-lived
none 0.2-0.92		37 BARNA	68	CNTR	Long-lived
none 0.97-1.03		37 BARNA	68	CNTR	Long-lived

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 35 ANSORGE 73B looks for electron pair production and electron-like Bremsstrahlung.
- 36 BUSHNIN 73 is SERPUKOV 70 GeV p experiment. Masses assume mean life above 7×10^{-10} and 3×10^{-8} respectively. Calculated from cross section (see "Charged Quasi-Stable Lepton Production Differential Cross Section" below) and 30 GeV muon pair production data.
- 37 BARNA 68 is SLAC photoproduction experiment.

Doubly-Charged Heavy Lepton MASS LIMITS

VALUE (GeV)	CL%	DOCUMENT ID	TECN	CHG	COMMENT
none 1-9 GeV	90	38 CLARK	81	SPEC	++

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 38 CLARK 81 is FNAL experiment with 209 GeV muons. Bounds apply to μp which couples with full weak strength to muon. See also section on "Doubly-Charged Lepton Production Cross Section."

Stable Neutral Heavy Lepton MASS LIMITS

Note that LEP results in combination with REUSSER 91 exclude a fourth stable neutrino with $m < 2400$ GeV.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.0	95	ABREU	92B	DLPH Dirac
>39.5	95	ABREU	92B	DLPH Majorana
>44.1	95	ALEXANDER	91F	OPAL Dirac
>37.2	95	ALEXANDER	91F	OPAL Majorana
none 3-100	90	SATO	91	KAM2 Kamiokande II
>42.8	95	39 ADEVA	90S	L3 Dirac
>34.8	95	39 ADEVA	90S	L3 Majorana
>42.7	95	DECAMP	90F	ALEP Dirac

- 39 ADEVA 90S limits for the heavy neutrino apply if the mixing with the charged leptons satisfies $|U_{1j}|^2 + |U_{2j}|^2 + |U_{3j}|^2 > 6.2 \times 10^{-8}$ at $m_{L^0} = 20$ GeV and $> 5.1 \times 10^{-10}$ for $m_{L^0} = 40$ GeV.

Neutral Heavy Lepton MASS LIMITS

Limits apply only to heavy lepton type given in comment at right of data Listings. See review above for description of types. $L, e p, \mu p, e O, \mu O$ stand for sequential lepton, para-electron, para-muon, ortho-electron, ortho-muon respectively. For a review, see GAN 88.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 2.5-50	95	40 ADRIANI	92I	L3 $ U_{\tau \text{ or } \mu} ^2 < 3 \times 10^{-4}$
none 4-50	95	40 ADRIANI	92I	L3 $ U_{\tau} ^2 < 3 \times 10^{-4}$
>46.4	95	41 ADEVA	90S	L3 Dirac
>45.1	95	41 ADEVA	90S	L3 Majorana
>46.5	95	42 AKRAWY	90L	OPAL Coupling to e or μ
>45.7	95	42 AKRAWY	90L	OPAL Coupling to τ
>19.6	95	43,44 BURCHAT	90	MRK2 Dirac, all $ U_{\ell j} ^2$
none 25-45.7	95	43,45 DECAMP	90F	ALEP Dirac $ U_{\ell j} ^2 > 10^{-13}$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- >44.5 95 46 ABREU 92B DLPH Dirac
- >39.0 95 46 ABREU 92B DLPH Majorana
- >41 95 43,44 BURCHAT 90 MRK2 Dirac, $|U_{\ell j}|^2 > 10^{-10}$
- none 8.2-26.5 95 47 SHAW 89 AMY Dirac L^0 , $|U_{e j}|^2 > 10^{-6}$
- none 8.3-22.4 95 47 SHAW 89 AMY Majorana L^0 , $|U_{e j}|^2 > 10^{-6}$
- none 8.1-24.9 95 47 SHAW 89 AMY Majorana L^0 , $|U_{\mu j}|^2 > 10^{-6}$
- none 1.8-6.7 90 48 AKERLOF 88 HRS $|U_{e j}|^2 = 1$
- none 1.8-6.4 90 48 AKERLOF 88 HRS $|U_{\mu j}|^2 = 1$
- none 2.5-6.3 80 48 AKERLOF 88 HRS $|U_{\tau j}|^2 = 1$
- none 0.6-34.6 95 49 BEHREND 88C CELL $L^0 = e p, V-A$ coupling $|U_{e j}|^2 = 1$
- none 0.4-37.4 95 49 BEHREND 88C CELL $L^0 = e p, V+A$ coupling $|U_{e j}|^2 = 1$
- none 0.25-14 90 50 MISHRA 87 CNTR $|U_{\mu j}|^2 = 1$
- none 0.25-10 90 50 MISHRA 87 CNTR $|U_{\mu j}|^2 = 0.1$

none 0.25-7.7	90	50 MISHRA	87	CNTR	$ U_{\mu j} ^2 = 0.03$
none 1.-2.	90	51 WENDT	87	MRK2	$ U_e \text{ or } \mu j ^2 = 0.1$
none 2.2-4.	90	51 WENDT	87	MRK2	$ U_e \text{ or } \mu j ^2 = 0.001$
none 2.3-3.	90	51 WENDT	87	MRK2	$ U_{\tau j} ^2 = 0.1$
none 3.2-4.8	90	51 WENDT	87	MRK2	$ U_{\tau j} ^2 = 0.001$
none 0.3-0.9	90	52 BADIÉ	86	CNTR	$ U_{e j} ^2 = 0.8$
none 0.33-2.0	90	52 BADIÉ	86	CNTR	$ U_{e j} ^2 = 0.03$
none 2.3-3.	90	52 BADIÉ	86	CNTR	$ U_{\mu j} ^2 = 0.8$
none 0.6-2.0	90	52 BADIÉ	86	CNTR	$ U_{\mu j} ^2 = 0.01-0.001$
>24.5	95	53 BARTEL	83	JADE	$e p^0$ or $e O^0, V+A$
>22.5	95	53 BARTEL	83	JADE	$e p^0$ or $e O^0, V-A$
none 1-9	90	54 CLARK	81	SPEC	μp^0
> 1.2		MEYER	77	MRK1	Neutral

- 40 ADRIANI 92I is a search for isosinglet heavy lepton N_{ℓ} which might be produced from $Z \rightarrow \nu_{\ell} N_{\ell}$, then decay via a number of different channels. Limits are weaker for decay lengths longer than about 1 m.
- 41 ADEVA 90S limits for the heavy neutrino apply if the mixing with the charged leptons satisfies $|U_{1j}|^2 + |U_{2j}|^2 + |U_{3j}|^2 > 6.2 \times 10^{-8}$ at $m_{L^0} = 20$ GeV and $> 5.1 \times 10^{-10}$ for $m_{L^0} = 40$ GeV.
- 42 AKRAWY 90L limits valid if coupling strength is greater than a mass-dependent value, e.g., 4.9×10^{-7} at $m_{L^0} = 20$ GeV, 3.5×10^{-8} at 30 GeV, 4×10^{-9} at 40 GeV.
- 43 Limits apply for $\ell = e, \mu, \tau$ and for $V-A$ decays of Dirac neutrinos.
- 44 BURCHAT 90 searched for Z decay to unstable L^0 pairs at SLC. It includes the analyses reported in JUNG 90, ABRAMS 89c, and WENDT 87.
- 45 For $25 < m_{L^0} < 42.7$ GeV, DECAMP 90F exclude an L^0 for all values of $|U_{\ell j}|^2$.
- 46 ABREU 92B limit is for mixing matrix element ≈ 1 for coupling to e or μ . Reduced somewhat for coupling to τ , increased somewhat for smaller mixing matrix element. Replaces ABREU 91F.
- 47 SHAW 89 also excludes the mass region from 8.0 to 27.2 GeV for Dirac L^0 and from 8.1 to 23.6 GeV for Majorana L^0 with equal full-strength couplings to e and μ . SHAW 89 also gives correlated bounds on lepton mixing.
- 48 AKERLOF 88 is PEP $e^+ e^-$ experiment at $E_{cm} = 29$ GeV. The L^0 is assumed to decay via $V-A$ to e or μ or τ plus a virtual W .
- 49 The first bound of BEHREND 88C applies for a general L^0 . The second and third have their assumptions indicated.
- 50 MISHRA 87 is Fermilab neutrino experiment looking for either dimuon or double vertex events (hence long-lived).
- 51 WENDT 87 is MARK-II search at PEP for heavy ν with decay length 1-20 cm (hence long-lived).
- 52 BADIÉ 86 is a search for a long-lived penetrating sequential lepton produced in $\pi^- -$ nucleon collisions with lifetimes in the range from 5×10^{-7} - 5×10^{-11} s and decaying into at least two charged particles. $U_{e j}$ and $U_{\mu j}$ are mixing angles to ν_e and ν_{μ} . See also the BADIÉ 86 entry in the section "Searches for Massive Neutrinos and Lepton Mixing".
- 53 BARTEL 83 is PETRA $e^+ e^-$ experiment with average $W_{cm} = 34.2$ GeV. First (second) limit is for $V+A(V-A)$ type $W^- e p$ coupling.
- 54 CLARK 81 is FNAL experiment with 209 GeV muons. Bounds apply to para-muon which couples with full weak strength to muon. See also section on "Neutral Heavy Lepton Production Cross Section (μ Nucleon)" below.

Astrophysical Limits on Neutrino MASS for $m_{\nu} > 1$ MeV

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
none 6 - hundreds		55,56 MORI	92B	KAM2 Dirac neutrino
none 24 - hundreds		55,56 MORI	92B	KAM2 Majorana neutrino
none 10-2400	90	57 REUSSER	91	CNTR HPGe search
none 3-100	90	SATO	91	KAM2 Kamiokande II
		58 ENQVIST	89	COSM
none 12-1400		59 CALDWELL	88	COSM Dirac ν
none 4-16	90	55,59 OLIVE	88	COSM Dirac ν
none 4-35	90	OLIVE	88	COSM Majorana ν
>4.2 to 4.7		SREDNICKI	88	COSM Dirac ν
>5.3 to 7.4		SREDNICKI	88	COSM Majorana ν
none 20-1000	95	59 AHLEN	87	COSM Dirac ν
>4.1		GRIEST	87	COSM Dirac ν

- 55 Limits based on annihilations in the sun and are due to an absence of high energy neutrinos detected in underground experiments.
- 56 MORI 92B results assume that neutrinos make up dark matter in the galactic halo. Limits based on annihilations in earth are also given.
- 57 REUSSER 91 uses existing $\beta\beta$ detector (see FISHER 89) to search for CDM Dirac neutrinos.
- 58 ENQVIST 89 argue that there is no cosmological upper bound on heavy neutrinos.
- 59 These results assume that neutrinos make up dark matter in the galactic halo.

Doubly-Charged Lepton Production Cross Section (μN Scattering)

VALUE (cm ²)	EVTs	DOCUMENT ID	TECN	CHG
<6. $\times 10^{-38}$	0	60 CLARK	81	SPEC ++

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 60 CLARK 81 is FNAL experiment with 209 GeV muon. Looked for μ^+ nucleon $\rightarrow \bar{\mu}_p^0 X$, $\bar{\mu}_p^0 \rightarrow \mu^+ \mu^- \bar{\nu}_{\mu}$ and $\mu^+ n \rightarrow \mu_p^+ X, \mu_p^+ \rightarrow 2\mu^+ \nu_{\mu}$. Above limits are for $\sigma \times BR$ taken from their mass-dependence plot figure 2.

See key on page 1343

Lepton & Quark Full Listings

Heavy Lepton Searches, Massive Neutrinos and Lepton Mixing

Neutral Heavy Lepton Production Cross Section (μN)

VALUE (cm^2)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$< 4. \times 10^{-38}$	0	61 CLARK	81	SPEC	0 μ_P^0
$< 1.22 \times 10^{-34}$		62 LEBRITTON	80	SPEC	0 $M^0 \rightarrow \mu^+ \mu^- \nu$
61 CLARK 81 is FNAL experiment with 209 GeV muon. Looked for μ^+ nucleon $\rightarrow \bar{\mu}_P^0 X$, $\bar{\mu}_P^0 \rightarrow \mu^+ \mu^- \bar{\nu}_\mu$ and $\mu^+ n \rightarrow \mu_P^+ X$, $\mu_P^+ \rightarrow 2\mu^+ \nu_\mu$. Above limits are for $\sigma \times BR$ taken from their mass-dependence plot figure 2.					
62 LEBRITTON 80 is BNL experiment with 10.5 GeV muons. Trimuons are consistent with QED trident and diffractively produced ρ decay.					

Neutral Heavy Lepton Production Cross Section

$\sigma \times B(\tau \rightarrow \text{new neutral lepton}) \times B(\text{neutral lepton} \rightarrow e\pi\mu\mu)$	VALUE (10^{-5} nb)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 450	90	63 MEYER	77	MRK1	For $m_L=0.5$ GeV
< 250	90	63 MEYER	77	MRK1	For $m_L=1.5$ GeV
63 MEYER 77 experiment looks for narrow neutral resonance in $e^- \pi$ and $\mu^- \pi$ channels. See "Heavy Lepton MASS LIMITS" section above.					

$\sigma(L_L) \times [B(L \rightarrow e\nu X) + B(L \rightarrow e\nu X)]$	VALUE (10^{-5} nb)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 8	90	64 ERREDE	84	HRS	For $m_L=1$ GeV
< 18	90	64 ERREDE	84	HRS	For $m_L=2$ GeV
< 20	90	64 ERREDE	84	HRS	For $m_L=3$ GeV
< 11	90	64 ERREDE	84	HRS	For $m_L=4$ or 5 GeV
< 13	90	64 ERREDE	84	HRS	For $m_L=6$ GeV
< 17	90	64 ERREDE	84	HRS	For $m_L=7$ GeV

64 Assuming $X = \mu$. If $X = \text{meson}$, limits are 20% higher. ERREDE 84 say these limits are comparable to those expected from naive theory. $e^+ e^-$, $E_{cm} = 29$ GeV. See also GRONAU 84, RIZZO 84.

$\sigma(L_1 + L_2) \times B(L_1 \rightarrow \text{only light neutrinos})$

VALUE (10^{-5} nb)	CL%	DOCUMENT ID	TECN	COMMENT	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 4.7	90	65 AKERLOF	85	HRS	For $m_L=2$ GeV
< 18	90	65 AKERLOF	85	HRS	For $m_L=10$ GeV
65 AKERLOF 85 observe no monojets above background. They use standard couplings to Z to find $\sigma(L_1 + L_2) = 0.36$ pb. Above data then imply $B(L_1 \rightarrow \text{light neutrinos}) < 13\text{--}50\%$ for $m_L = 2\text{--}10$ GeV.					

$\sigma(L_L) \times BR_1 \times BR_2 / \sigma(\text{standard via virtual } Z)$

where BR_1 and BR_2 are branching ratios leading to events with two or four charged particles, and $\sigma(\text{standard}) = 0.35(\beta(3 + \beta^2) / 4)$ pb with $\beta = \text{velocity}/c$ of L .

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
$< 0.1\text{--}0.2$	90	0	66 PERL	85	MRK2 For $m_L < 1$ GeV
66 PERL 85 examine a variety of models and processes. They search up to $m_L = 14$ GeV but are most sensitive for $m_L < 1$ GeV. They require lepton lifetime $< m_L 10^{-11}$ s [m_L in GeV] which limits their ability to constrain the mixing of a 4th conventional generation.					

REFERENCES FOR Heavy Lepton Searches

ABREU 92B PL B274 230	+Adams, Adami, Adye+	(DELPHI Collab.)
ADRIANI 92I PL B295 371	+Aguiar-Benitez, Aihou, Akbari, Alcaraz+	(L3 Collab.)
MORI 92B PL B289 463	+Hikasa, Nijiri, Oyama+	(KAM2 Collab.)
ABREU 91F NP B367 511	+Adam, Adami, Adye, Akesson+	(DELPHI Collab.)
ALEXANDER 91F ZPHY C52 175	+Allison, Allport, Anderson, Arcelli+	(OPAL Collab.)
KIM 91B UMP A6 2583	+Smith, Breedon, Ko+	(AMY Collab.)
REUSSER 91 PL B255 143	+Treichel, Boehm, Broggin+	(NEUC, CIT, VILL)
SATO 91 PR D44 2220	+Hirata, Kajita, Kifune, Kihara+	(Kamioka Collab.)
ADACHI 90C PR D44 352	+Aihara, Doser, Enomoto+	(TOPAZ Collab.)
ADEVA 90S PL B251 321	+Adriani, Aguiar-Benitez, Akbari+	(L3 Collab.)
AKRAWY 90G PL B240 250	+Alexander, Allison, Allport+	(OPAL Collab.)
AKRAWY 90L PL B247 448	+Alexander, Allison, Allport+	(OPAL Collab.)
AKRAWY 90O PL B252 290	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
BURCHAT 90 PR D41 3542	+King, Abrams, Adolphsen+	(Mark II Collab.)
DECAMP 90F PL B236 511	+Deschizeaux, Lees, Minard+	(ALEPH Collab.)
JUNG 90 PRL 64 1091	+Van Kooten, Abrams, Adolphsen+	(Mark II Collab.)
RILES 90 PR D42 1	+Peri, Barlow+	(Mark II Collab.)
SODERSTROM 90 PRL 64 2980	+McKenna, Abrams, Adolphsen, Averill+	(Mark II Collab.)
ABRAMS 89C PRL 63 2447	+Adolphsen, Averill, Ballam+	(Mark II Collab.)
ENQUIST 89 NP B317 647	+Kainulainen, Maalampi	(HELS)
FISHER 89 PL B218 257	+Boehm, Bovet, Egger+	(CIT, NEUC, PSI)
SHAW 89 PRL 63 1342	+Blanis, Bodek, Budd+	(AMY Collab.)
STOKER 89 PR D39 1811	+Peri, Abrams+	(Mark II Collab.)
ABE 90 PRL 61 915	+Amako, Arai, Asano, Chiba	(AMY Collab.)
ADACHI 88B PR D37 1339	+Aihara, Dijkstra, Enomoto+	(TOPAZ Collab.)
AKERLOF 88 PR D37 577	+Chapman, Errede, Ken+	(HRS Collab.)
BEHREND 88C ZPHY C41 7	+Buerger, Criegee, Dainton+	(CELLO Collab.)
CALDWELL 88 PRL 61 510	+Eisberg, Grumm, Witherell+	(UCSB, UCB, LBL)
GAN 88 IJMP A3 531	+Peri	(SLAC)
IGARASHI 88 PRL 60 2359	+Myung, Chiba, Hanaoka+	(AMY Collab.)
KIM 88 PRL 61 911	+Son, Bacala, Imlay+	(AMY Collab.)
ADACHI 88B PR D37 1339	+Srednicki	(MINN, UCSB)
AKERLOF 88 PR D37 577	+Watkins, Olive	(MINN, UCSB)
BEHREND 88C ZPHY C41 7	+Avignone, Brodzinski+	(BOST, SCUC, HARV, CHIC)
CALDWELL 88 PRL 61 510	+Albrow, Allkofer, Arnison+	(UA1 Collab.)
GAN 88 IJMP A3 531	+Seckel	(UCSC, CERN)
IGARASHI 88 PRL 60 2359	+Griest, Seckel	(UCSC, CERN)
MISHRA 87 PRL 59 1397	+Auchincloss+	(COLU, CIT, FNAL, CHIC, ROCH)

WENDT 87 PRL 58 1810	+Abrams, Amidei, Baden+	(Mark II Collab.)
YOSHIDA 87B PRL 59 2915	+Chiba, Endo+	(VENUS Collab.)
BADIER 86 ZPHY C31 21	+Bemporad, Boucrot, Callot+	(NA3 Collab.)
ADEVA 85 PL 152B 439	+Becker, Becker-Szendy+	(Mark-J Collab.)
Also 84C PRPL 109 131	+Adeva, Barber, Becker+	(Mark-J Collab.)
AKERLOF 85 PL 156B 271	+Bonvicini, Chapman, Errede+	(HRS Collab.)
PERL 85 PR D32 2859	+Barlow, Boyarski+	(Mark II Collab.)
84 PL 149B 519	+Akerlof, Chapman, Harnew+	(HRS Collab.)
GRONAU 84 PR D29 2539	+Leung, Rosner	(SYRA, FNAL, CHIC)
RIZZO 84 PL 136B 251		(ISU)
ADEVA 83B PRL 51 443	+Barber, Becker, Berdugo+	(Mark-J Collab.)
BARTEL 83 PL 123B 353	+Cords, Dietrich, Eichler+	(JADE Collab.)
ADEVA 82 PRL 48 967	+Barber, Becker, Berdugo+	(Mark-J Collab.)
BERGER 81B PL 99B 489	+Genzel, Grigull, Lackas+	(PLUTO Collab.)
BRANDELIK 81 PL 99B 163	+Braunschwelg, Gather+	(TASSO Collab.)
CLARK 81 PRL 46 299	+Johnson, Kerth, Loken+	(UCB, LBL, FNAL, PRIN)
Also 82 PR D25 2762	+Smith, Clark, Johnson, Kerth+	(LBL, FNAL, PRIN)
AZIMOV 80 JETPL 32 664	+Khozze	(PNPI)
Translated from ZETFP 32 677.		
BARBER 80B PRL 45 1904	+Becker, Bei, Berghoff+	(Mark-J Collab.)
BRANDELIK 80 PL 92B 199	+Braunschwelg, Gather+	(TASSO Collab.)
LEBRITTON 78 PL 92B 271	+McCal, Melissinos+	(ROCH, BNL, NSF)
ASRATYAN 78 PL 92B 237	+Kubantsev	(ITEP)
CNOPS 78 PRL 40 144	+Connolly, Kahn, Kirk+	(BNL, COLU)
ERRIQUEZ 78 PL 77B 227	(BARI, BIRM, BRUX, EPOL, RHEL, SACL, LOUC)	
HOLDER 78 PL 74B 277	+Knobloch, May+	(CDHS Collab.)
BACCI 77B PL 71B 227	+Dezori, Penso, Stella+	(ROMA, FRAS)
HOLDER 77 PL 70B 393	+Knobloch, May+	(CDHS Collab.)
MEYER 77 PL 70B 469	+Ngyuen, Abrams+	(SLAC, LBL, NWES, HAWA)
ASRATYAN 77 PL 46B 498	+Gershtein, Kaftanov, Kubantsev, Lapin+	(SERP)
BARISH 74 PRL 32 1387	+Bartlett, Buchholz, Merritt+	(CIT, FNAL)
GITLSON 74 PR D10 1379	+Kirk+	(HARV, ROCH, COLU, FNAL)
ANSORGE 73B PR D7 26	+Baker, Krzesinski, Neale, Rushbrooke+	(CAVE)
BACCI 73 PL 44B 530	+Parisi, Penso, Salvini, Stella+	(ROMA, FRAS)
BARISH 73B PRL 31 410	+Bartlett, Buchholz, Humphrey+	(CIT, FNAL)
BUSHNIN 73 NP 85B 476	+Dunaltzev, Golovkin, Kubarovskiy+	(SERP)
Also 72 PR 15B 236	+Golovkin, Grachev, Shodyrev+	(SERP)
EICHTEN 73 PL 46B 281	+Deden, Hasert, Krenz+	(Gargamelle Collab.)
LICHTENSTEIN 70 PR D1 825	+Ash, Berkelman, Hartill+	(CORN)
LIBERMAN 69 PRL 22 663	+Hoffman, Engels+	(HARV, CASE, MCGI, SLAC)
ROTHE 69 NP B10 241	+Wolsky	(PENN)
BARNA 68 PR 173 1391	+Cox, Martin, Peri, Tan, Toner, Zipf+	(SLAC, STAN)
BOLEY 68 PR 167 1275	+Elias, Friedman, Hartmann, Kendall+	(MIT, CEAN)
BUDNITZ 66 PR 141 1313	+Dunning, Gottein, Ramsey, Walker+	(HARV)
BEHREND 65 PRL 15 900	+Brasse, Ingler, Ganssauge+	(DESY, KARL)
BETOURNE 65 PL 17 70	+Ngoc, Perez-Jorba+	(ORSAY)

OTHER RELATED PAPERS

PERL 81 SLAC-PUB-2752	(SLAC)
Physics in Collision Conference.	

Searches for Massive Neutrinos and Lepton Mixing

Searches for the effects of nonzero neutrino masses are listed here. Direct searches for masses of dominantly coupled neutrinos are listed in the appropriate section on ν_e , ν_μ , or ν_τ . The results in the present section are correlated upper bounds on mixing matrix coefficients $U_{\alpha j}$ versus neutrino mass. These results are divided into three main sections:

1. Bounds from particle and nuclear decays;
2. Bounds from neutrino reactions, including reactor and accelerator neutrino oscillation experiments, and solar neutrino measurements (see the "Note on Neutrino Oscillations" below); and
3. Searches for neutrinoless double- β decay:

Discussion of the ν_e and ν_μ mass limits, the "17 keV neutrino," and solar neutrino observations are given in the "Note on Neutrinos" by R.E. Shrock in the ν_e section near the beginning of these data listings. Several reviews are also listed there. See also the "Note on the Muon Neutrino Mass" before the ν_μ Listings.

In addition, searches for mixing of $(\mu^- e^+)$ and $(\mu^+ e^-)$ are given in the muon Listings.

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

(A) Bounds from Particle and Nuclear Decays

Limits on $|U_{1j}|^2$ as Function of m_{ν_j}

Application of Kink and Peak Search Test to Existing Data

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1 \times 10^{-4}$	68	¹ SHROCK 81	THEO	$m_{\nu_j} = 10$ MeV
$< 5 \times 10^{-6}$	68	¹ SHROCK 81	THEO	$m_{\nu_j} = 60$ MeV
< 1	95	² SIMPSON 81B		$m_{\nu_j} = 0.1$ keV
$< 4 \times 10^{-3}$	95	² SIMPSON 81B		$m_{\nu_j} = 10$ keV
< 0.1	68	³ SHROCK 80	THEO	$m_{\nu_j} = 0.1-3$ MeV
$< 1 \times 10^{-5}$	68	⁴ SHROCK 80	THEO	$m_{\nu_j} = 80$ MeV
$< 3 \times 10^{-6}$	68	⁴ SHROCK 80	THEO	$m_{\nu_j} = 160$ MeV

- ¹ Analysis of $(\pi^+ \rightarrow e^+ \nu_e)/(\pi^+ \rightarrow \mu^+ \nu_\mu)$ and $(K^+ \rightarrow e^+ \nu_e)/(K^+ \rightarrow \mu^+ \nu_\mu)$ decay ratios.
² Application of kink search test to tritium β decay Kurie plot.
³ Application of test to search for kinks in β decay Kurie plots.
⁴ Analysis of $(K^+ \rightarrow e^+ \nu_e)$ spectrum.

New Experiments to Apply Peak and Kink Search Tests

Limits on $|U_{1j}|^2$ as function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1 \times 10^{-7}$	90	⁵ BRITTON 92B	CNTR	50 MeV $< m_{\nu_j} < 130$ MeV

- • • We do not use the following data for averages, fits, limits, etc. • • •

$< 5 \times 10^{-6}$	90	DELEENER... 91		$m_{\nu_j} = 20$ MeV
$< 5 \times 10^{-7}$	90	DELEENER... 91		$m_{\nu_j} = 40$ MeV
$< 3 \times 10^{-7}$	90	DELEENER... 91		$m_{\nu_j} = 60$ MeV
$< 1 \times 10^{-6}$	90	DELEENER... 91		$m_{\nu_j} = 80$ MeV
$< 1 \times 10^{-6}$	90	DELEENER... 91		$m_{\nu_j} = 100$ MeV
$< 5 \times 10^{-7}$	90	AZUELOS 86	CNTR	$m_{\nu_j} = 60$ MeV
$< 2 \times 10^{-7}$	90	AZUELOS 86	CNTR	$m_{\nu_j} = 80$ MeV
$< 3 \times 10^{-7}$	90	AZUELOS 86	CNTR	$m_{\nu_j} = 100$ MeV
$< 1 \times 10^{-6}$	90	AZUELOS 86	CNTR	$m_{\nu_j} = 120$ MeV
$< 2 \times 10^{-7}$	90	AZUELOS 86	CNTR	$m_{\nu_j} = 130$ MeV
$< 8 \times 10^{-6}$	90	DELEENER... 86	CNTR	$m_{\nu_j} = 20$ MeV
$< 4 \times 10^{-7}$	90	DELEENER... 86	CNTR	$m_{\nu_j} = 60$ MeV
$< 2 \times 10^{-6}$	90	DELEENER... 86	CNTR	$m_{\nu_j} = 100$ MeV
$< 7 \times 10^{-6}$	90	DELEENER... 86	CNTR	$m_{\nu_j} = 120$ MeV
$< 1 \times 10^{-4}$	90	⁶ BRYMAN 83B	CNTR	$m_{\nu_j} = 5$ MeV
$< 1.5 \times 10^{-6}$	90	BRYMAN 83B	CNTR	$m_{\nu_j} = 53$ MeV
$< 1 \times 10^{-5}$	90	BRYMAN 83B	CNTR	$m_{\nu_j} = 70$ MeV
$< 1 \times 10^{-4}$	90	BRYMAN 83B	CNTR	$m_{\nu_j} = 130$ MeV

- ⁵ BRITTON 92B is from a search for additional peaks in the e^+ spectrum from $\pi^+ \rightarrow e^+ \nu_e$ decay at TRIUMF. See also BRITTON 92.

- ⁶ BRYMAN 83B obtain upper limits from both direct peak search and analysis of $B(\pi \rightarrow e\nu)/B(\pi \rightarrow \mu\nu)$. Latter limits are not listed, except for this entry (i.e. — we list the most stringent limits for given mass).

Searches for Decays of Massive ν

Limits on $|U_{1j}|^2$ as function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1 \times 10^{-5}$	90	⁷ BARANOV 93		$m_{\nu_j} = 100$ MeV
$< 1 \times 10^{-6}$	90	⁷ BARANOV 93		$m_{\nu_j} = 200$ MeV
$< 3 \times 10^{-7}$	90	⁷ BARANOV 93		$m_{\nu_j} = 300$ MeV
$< 2 \times 10^{-7}$	90	⁷ BARANOV 93		$m_{\nu_j} = 400$ MeV
$< 6.2 \times 10^{-8}$	95	ADEVA 90S L3		$m_{\nu_j} = 20$ GeV
$< 5.1 \times 10^{-10}$	95	ADEVA 90S L3		$m_{\nu_j} = 40$ GeV
all values ruled out	95	⁸ BURCHAT 90	MRK2	$m_{\nu_j} < 19.6$ GeV
$< 1 \times 10^{-10}$	95	⁸ BURCHAT 90	MRK2	$m_{\nu_j} = 22$ GeV
$< 1 \times 10^{-11}$	95	⁸ BURCHAT 90	MRK2	$m_{\nu_j} = 41$ GeV
all values ruled out	95	DECAMP 90F	ALEP	$m_{\nu_j} = 25.0-42.7$ GeV
$< 1 \times 10^{-13}$	95	DECAMP 90F	ALEP	$m_{\nu_j} = 42.7-45.7$ GeV
$< 5 \times 10^{-3}$	90	AKERLOF 88	HRS	$m_{\nu_j} = 1.8$ GeV
$< 2 \times 10^{-5}$	90	AKERLOF 88	HRS	$m_{\nu_j} = 4$ GeV
$< 3 \times 10^{-6}$	90	AKERLOF 88	HRS	$m_{\nu_j} = 6$ GeV
$< 1.2 \times 10^{-7}$	90	BERNARDI 88	CNTR	$m_{\mu_j} = 100$ MeV
$< 1 \times 10^{-8}$	90	BERNARDI 88	CNTR	$m_{\mu_j} = 200$ MeV
$< 2.4 \times 10^{-9}$	90	BERNARDI 88	CNTR	$m_{\mu_j} = 300$ MeV

$< 2.1 \times 10^{-9}$	90	BERNARDI 88	CNTR	$m_{\mu_j} = 400$ MeV
$< 2 \times 10^{-2}$	68	⁹ OBERAUER 87		$m_{\nu_j} = 1.5$ MeV
$< 8 \times 10^{-4}$	68	⁹ OBERAUER 87		$m_{\nu_j} = 4.0$ MeV
$< 8 \times 10^{-3}$	90	BADIER 86	CNTR	$m_{\nu_j} = 400$ MeV
$< 8 \times 10^{-5}$	90	BADIER 86	CNTR	$m_{\nu_j} = 1.7$ GeV
$< 8 \times 10^{-8}$	90	BERNARDI 86	CNTR	$m_{\nu_j} = 100$ MeV
$< 4 \times 10^{-8}$	90	BERNARDI 86	CNTR	$m_{\nu_j} = 200$ MeV
$< 6 \times 10^{-9}$	90	BERNARDI 86	CNTR	$m_{\nu_j} = 400$ MeV
$< 3 \times 10^{-5}$	90	DORENBOS... 86	CNTR	$m_{\nu_j} = 150$ MeV
$< 1 \times 10^{-6}$	90	DORENBOS... 86	CNTR	$m_{\nu_j} = 500$ MeV
$< 1 \times 10^{-7}$	90	DORENBOS... 86	CNTR	$m_{\nu_j} = 1.6$ GeV
$< 7 \times 10^{-7}$	90	¹⁰ COOPER... 85	HLBC	$m_{\nu_j} = 0.4$ GeV
$< 8 \times 10^{-8}$	90	¹⁰ COOPER... 85	HLBC	$m_{\nu_j} = 1.5$ GeV
$< 1 \times 10^{-2}$	90	¹¹ BERGSMA 83B	CNTR	$m_{\nu_j} = 10$ MeV
$< 1 \times 10^{-5}$	90	¹¹ BERGSMA 83B	CNTR	$m_{\nu_j} = 110$ MeV
$< 6 \times 10^{-7}$	90	¹¹ BERGSMA 83B	CNTR	$m_{\nu_j} = 410$ MeV
$< 1 \times 10^{-5}$	90	GRONAU 83		$m_{\nu_j} = 160$ MeV
$< 1 \times 10^{-6}$	90	GRONAU 83		$m_{\nu_j} = 480$ MeV

- ⁷ BARANOV 93 is a search for neutrino decays into $e^+ e^- \nu_e$ using a beam dump experiment at the 70 GeV Serpukhov proton synchrotron. The limits are not as good as those achieved earlier by BERGSMA 83 and BERNARDI 86, BERNARDI 88.

- ⁸ BURCHAT 90 includes the analyses reported in JUNG 90, ABRAMS 89C, and WENDT 87.

- ⁹ OBERAUER 87 bounds from search for $\nu \rightarrow \nu' e e$ decay mode using reactor (anti)neutrinos.

- ¹⁰ COOPER-SARKAR 85 also give limits based on model-dependent assumptions for ν_τ flux. We do not list these. Note that for this bound to be nontrivial, j is not equal to 3, i.e. ν_j cannot be the dominant mass eigenstate in ν_τ since $m_{\nu_3} < 70$ MeV (ALBRECHT 85). Also, of course, j is not equal to 1 or 2, so a fourth generation would be required for this bound to be nontrivial.

- ¹¹ BERGSMA 83B also quote limits on $|U_{13}|^2$ where the index 3 refers to the mass eigenstate dominantly coupled to the τ . Those limits were based on assumptions about the D_s mass and $D_s \rightarrow \tau \nu_\tau$ branching ratio which are no longer valid. See COOPER-SARKAR 85.

Kink Search in Nuclear β Decay

VALUE (units 10^{-3})	CL%	m_{ν_j} (keV)	ISOTOPE	METHOD	DOCUMENT ID
< 1.9	90	17	³⁵ S	Si+guide field	¹² ABFLE 93
$0.3 \pm 1.5 \pm 0.8$	17		³⁵ S	Mag spect	¹³ BERMAN 93
< 5	95	17	⁷¹ Ge	IBEC; γ det	¹⁴ DIGREGORIO 93
			³⁵ S	Si(Li) det	¹⁵ HIME 93
< 2.8	99	17	³ H	Prop chamber	¹⁶ KALBFLEISCH 93
< 1	99	14.4-15.2	³ H	Prop chamber	¹⁶ KALBFLEISCH 93
< 0.7	99	16.3-16.6	³ H	Prop chamber	¹⁶ KALBFLEISCH 93
< 2	95	13-40	³⁵ S	Si(Li)	¹⁷ MORTARA 93
0.73	95	17	⁶³ Ni	Mag spect	OHSIMA 93
1.5	95	10.5-25.0	⁶³ Ni	Mag spect	¹⁸ OHSIMA 93
< 6	95	5-25	⁵⁵ Fe	IBEC in Ge	¹⁹ WIETFELDT 93
< 2	95	17	⁵⁵ Fe	IBEC in Ge	¹⁹ WIETFELDT 93
< 4	99	17	³ H	Prop chamber	²⁰ BAHNAN 92
< 2	90	17	³⁵ S	Mag spect.	²¹ CHEN 92
< 0.95	95	17	⁶³ Ni	Mag spect	²² KAWAKAMI 92
< 1.0	95	10-24	⁶³ Ni	Mag spect	KAWAKAMI 92
$8.4 \pm 0.6 \pm 0.5$		17.0 ± 0.4	³⁵ S	Si(Li) det	HIME 91
$9.9 \pm 1.2 \pm 1.8$		$16.75 \pm 0.35 \pm 0.15$	⁶³ Ni	Solid state det	HIME 91B
$14.0 \pm 4.5 \pm 1.4$		17 ± 2	¹⁴ C	¹⁴ C in HPGe	²³ SUR 91
16 ± 7		$17.2^{+1.9}_{-1.1}$	⁷¹ Ge	THEO	²⁴ ZLIMEN 91
		17			²⁵ DRUKAREV 89
6 to 16		16.9 ± 0.4	³ H	In HPGe	HIME 89
5 to 18		17.1 ± 0.2	³ H	In Si(Li)	²⁶ HIME 89
$7.3 \pm 0.9 \pm 0.6$		16.9 ± 0.4	³⁵ S	Si(Li)	²⁷ SIMPSON 89
< 7.4	99.7	$16.4-17.4$	⁵⁵ Fe	IBEC; γ det	²⁸ ZLIMEN 88
< 3	90	17	⁶³ Ni	Mag spect	²⁹ HETHERING... 87
< 10	90	16-35	¹²⁵ I	IBEC; γ det	³⁰ BORGE 86
10 to 20		17		RVUE	³¹ SIMPSON 86
< 4	99	17	³⁵ S	Mag spect	ALTZITZOG... 85
< 7.5	99	5-50	³⁵ S	Mag spect	ALTZITZOG... 85
< 8	90	80	³⁵ S	Mag spect	³² APALIKOV 85
< 1.5	90	60	³⁵ S	Mag spect	APALIKOV 85
< 8	90	30	³⁵ S	Mag spect	APALIKOV 85
< 3	90	17	³⁵ S	Mag spect	APALIKOV 85

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

< 45	90	4	³⁵ S	Mag spect	APALIKOV	85
< 6	90	17	³⁵ S	Si(Li)	DATAR	85
< 10	90	5-30	³⁵ S	Si(Li)	DATAR	85
< 3.0	90	5-50		Mag spect	MARKEY	85
< 2.5	90	17		Mag spect	MARKEY	85
< 0.62	90	48	³⁵ S	Si(Li)	OHI	85
< 0.90	90	30	³⁵ S	Si(Li)	OHI	85
< 1.30	90	20	³⁵ S	Si(Li)	OHI	85
< 1.50	90	17	³⁵ S	Si(Li)	OHI	85
< 3.30	90	10	³⁵ S	Si(Li)	OHI	85
30 ± 10		17.1 ± 0.2	³ H	In Si(Li)	³³ SIMPSON	85
< 25	90	30	⁶⁴ Cu	Mag spect	³⁴ SCHRECK...	83
< 4	90	140	⁶⁴ Cu	Mag spect	³⁴ SCHRECK...	83
< 8	90	440	⁶⁴ Cu	Mag spect	³⁴ SCHRECK...	83
< 100	90	0.1-3000		THEO	³⁵ SHROCK	80
			³ H	Prop. cntr	³⁶ CONWAY	59

- ¹² ABELE 93: "It appears to us that we have succeeded in isolating experimentally the very same effect that was proposed in HIME 93 to resolve the "17 keV Condium."
- ¹³ BERMAN 93 uses an iron-free intermediate-image magnetic spectrometer to measure ³⁵Sβ decay over a large portion of the spectrum. Paper reports (0.01 ± 0.15)% above result revised by author on basis of analysis refinements.
- ¹⁴ DIGREGORIO 93 is an experiment on ⁷¹Ge IBEC to search for a possible admixture of a massive neutrino. The authors state that their results "exclude the presence of a massive component of 17.2 ± 1.3 ± 1.1 keV and (1.6 ± 0.6)% mixing fraction claimed by [ZLIMEN 91] for this same nucleus, at the 99.0% confidence level."
- ¹⁵ HIME 93 is a reanalysis of HIME 91 data and states that the effect, which was previously attributed to the emission of a massive neutrino, can be explained by electron energy loss without the need for any neutrino mass or mixing.
- ¹⁶ KALBFLEISCH 93 extends the 17 keV neutrino search of BAHHRAN 92, using an improved proportional chamber to which a small amount of ³H is added. Systematics are significantly reduced, allowing for an improved upper limit. The authors give a 99% confidence limit on $|U_{1j}|^2$ as a function of m_{ν_j} in the range from 13.5 keV to 17.5 keV. Typical upper limits are listed above. They report that this experiment in combination with BAHHRAN 92 gives an upper limit of 2.4×10^{-3} at the 99% CL. See also the related papers BAHHRAN 93 and BAHHRAN 93b, on theoretical aspects of beta spectra and fitting methods for heavy neutrinos.
- ¹⁷ MORTARA 93 limit is from study using a high-resolution solid-state detector with a superconducting solenoid. The authors note that "The sensitivity to neutrino mass is verified by measurement with a mixed source of ³⁵S and ¹⁴C, which artificially produces a distortion in the beta spectrum similar to that expected from the massive neutrino."
- ¹⁸ OHSHIMA 93 is the full data analysis from this experiment. The above limit on the mixing strength for a 17 keV neutrino is obtained from the measurement $|U_{1j}|^2 = (-0.11 \pm 0.33 \pm 0.30) \times 10^{-3}$ by taking zero as the best estimate and ignoring physical boundaries; see discussion in HOLZSCHUH 92b for a comparison of methods. Using the Particle Data Group recipe gives the slightly more conservative 95%CL of 0.80×10^{-3} . An earlier report of this experiment was given in KAWAKAMI 92.
- ¹⁹ WIETVELDT 93 is an extension of the NORMAN 91 experiment. However, whereas NORMAN 91 reported indications for the emission of a neutrino with mass $m_{\nu_j} = 21 \pm 2$ keV and coupling strength = 0.0085 ± 0.0045, the present experiment states that "We find no evidence for emission of a neutrino in the mass range 5-25 keV. In particular, a 17 keV neutrino with $\sin^2\theta$ ($|U_{1j}|^2$ in our notation) = 0.008 is excluded at the 7σ level." The listed limits can be obtained from the paper's Fig. 4. The authors acknowledge that this conclusion contradicts the one reported in NORMAN 91, based on a smaller data sample.
- ²⁰ BAHHRAN 92 minor errors corrected in BAHHRAN 92b.
- ²¹ CHEN 92 is a continuation and improvement of the Boehm et al. Caltech iron-free magnetic spectrometer experiment searching for emission of massive neutrinos in ³⁵S decay (MARKEY 85). The upper limit on $|U_{1j}|^2$ for $m_{\nu_j} = 17$ keV comes from the measurement $|U_{1j}|^2 = (-0.5 \pm 1.4) \times 10^{-3}$. The authors state that their results "rule out, at the 6σ level, a 17 keV neutrino admixed at 0.85% (i.e. with $|U_{1j}|^2 = 0.85 \times 10^{-2}$," the level claimed by Hime and Jelly in HIME 91. They also state that "our data show no evidence for a heavy neutrino with a mass between 12 and 22 keV" with substantial admixture in the weak admixture in the weak eigenstate ν_ρ ; see their Fig. 4 for a graphical set of measured values of $|U_{1j}|^2$ for various hypothetical values of m_{ν_j} in this range.
- ²² KAWAKAMI 92 experiment final results are given in OHSHIMA 93. The upper limit is improved to 0.73×10^{-3} , based on $|U_{1j}|^2 = (-0.11 \pm 0.33 \pm 0.30) \times 10^{-3}$. Ohshima notes that the result is 22σ away from the value $|U_{1j}|^2 = 1\%$.
- ²³ SUR 91 reports an LBL experiment using a solid state Ge crystal grown with ¹⁴C inside. In a conference report (NORMAN 91), the authors also report indications for the emission of a 17 keV neutrino in the ⁵⁵Fe inner bremsstrahlung transition: $m_{\nu_j} = 21 \pm 2$ keV with $|U_{1j}|^2 = 0.0085 \pm 0.0045$.
- ²⁴ ZLIMEN 91 used a HPGe detector to observe the inner bremsstrahlung electron capture transition of ⁷¹Ge in an external source. Reported errors on both parameters are given as 95%CL limits, which in the case of normal distributions corresponds to 1.96σ.
- ²⁵ DRUKAREV 89 claims that taking into account screening effects can explain Simpson's claims without invoking a massive neutrino or other unconventional physics. A similar criticism concerning screening corrections had been made by LINDHARD 86.
- ²⁶ HIME 89 corrects the analysis of the data of SIMPSON 85 for screening effects as suggested by LINDHARD 86, giving a smaller range for $|U_{1j}|^2$, as cited above. This value should therefore replace that given in SIMPSON 85, which has been retracted.
- ²⁷ SIMPSON 89 and HIME 89 report kinks due to the emission of a massive neutrino in ³⁵S and ³Hβ decays, respectively.
- ²⁸ ZLIMEN 88 report an experiment on ⁵⁵Fe, observing Internal bremsstrahlung in electron capture (IBEC). For a contemporary review of IBEC, see LOGAN 89.

- ²⁹ HETHERINGTON 87 reports no evidence for any massive neutrino signal for m_{ν_j} in the range from 4 to 40 keV, and, in particular, set the upper limit cited above on $|U_{1j}|^2$ for a hypothetical 17 keV neutrino.
- ³⁰ BORGE 86 results originally presented as evidence against the SIMPSON 85 claim of a 17 keV antineutrino emitted with $|U_{1j}|^2 = 0.03$ in ³H decay.
- ³¹ SIMPSON 86 is a reanalysis of the OHI 85 data and claims that these data show evidence of heavy neutrino emission with $m_{\nu_j} = 17$ keV and $|U_{1j}|^2 =$ from 0.01 to 0.02, consistent with the earlier reported observation by SIMPSON 85. This conclusion strongly disagrees with the conclusion reached by OHI 85 from their analysis of their own data. SIMPSON 86 also states that "a similar threshold effect (due to supposed heavy neutrino emission) is seen in several of the other published ³⁵S experiments as well."
- ³² This limit was taken from the figure 3 of APALIKOV 85; the text gives a more restrictive limit of 1.7×10^{-3} at CL = 90%.
- ³³ SIMPSON 85. See footnotes on SIMPSON 89 and SIMPSON 86, as well as comments by HAXTON 85, KALBFLEISCH 85, EMAN 86, LINDHARD 86, DRUKAREV 86, and further discussion by SIMPSON 89 and HIME 89.
- ³⁴ SCHRECKENBACH 83 is a combined measurement of the β⁺ and β⁻ spectrum.
- ³⁵ SHROCK 80 was a retroactive analysis of data on several superallowed β decays to search for kinks in the Kurie plot.
- ³⁶ CONWAY 59 first reported a spectral excess of about 1% at electron kinetic energy of 1 keV in ³Hβ decay, but did not interpret it as the emission of a massive neutrino. Indeed, no searches for masses admixed neutrinos were performed prior to 1980; cf. SHROCK 80. This spectral excess was again observed in SIMPSON 85, apparently without knowledge of the CONWAY 59 finding. Spectral excesses in this kinetic energy region were also reported in HAMILTON 58 and JOHNSON 58, and in other references cited therein.

Limits on $|U_{2j}|^2$ as Function of m_{ν_j}

Application of Peak Search Test to Existing Data

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
< 6 × 10 ⁻⁶	95	37 ASANO	81	$m_{\nu_j}=240$ MeV
< 5 × 10 ⁻⁷	95	37 ASANO	81	$m_{\nu_j}=280$ MeV
< 6 × 10 ⁻⁶	95	37 ASANO	81	$m_{\nu_j}=300$ MeV
< 3 × 10 ⁻²	95	38 SHROCK	81	THEO $m_{\nu_j}=7$ MeV
< 1 × 10 ⁻²	95	38 SHROCK	81	THEO $m_{\nu_j}=13$ MeV
< 1 × 10 ⁻⁴	68	38 SHROCK	81	THEO $m_{\nu_j}=13$ MeV
< 3 × 10 ⁻⁵	68	38 SHROCK	81	THEO $m_{\nu_j}=33$ MeV
< 6 × 10 ⁻³	68	39 SHROCK	81	THEO $m_{\nu_j}=80$ MeV
< 5 × 10 ⁻³	68	39 SHROCK	81	THEO $m_{\nu_j}=120$ MeV
< 5 × 10 ⁻²	95	38 SHROCK	80	THEO $m_{\nu_j}=4-6$ MeV

³⁷ Analysis of experiment on $K^+ \rightarrow \mu^+ \nu_\mu \nu_\mu \bar{\nu}_\mu$ decay.

³⁸ Analysis of magnetic spectrometer experiment, bubble chamber experiment, and emulsion experiment on $\pi^+ \rightarrow \mu^+ \nu_\mu$ decay.

³⁹ Analysis of magnetic spectrometer experiment on $K \rightarrow \mu, \nu_\mu$ decay.

Application of Peak Search Test to New Experiments

Limits on $|U_{2j}|^2$ as function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
< 2 × 10 ⁻²	90	DAUM	87 $m_{\nu_j}=1$ MeV
< 1 × 10 ⁻³	90	DAUM	87 $m_{\nu_j}=2$ MeV
< 6 × 10 ⁻⁵	90	DAUM	87 3 MeV < m_{ν_j} < 19.5 MeV
< 3 × 10 ⁻²	90	40 MINEHART	84 $m_{\nu_j}=2$ MeV
< 1 × 10 ⁻³	90	40 MINEHART	84 $m_{\nu_j}=4$ MeV
< 3 × 10 ⁻⁴	90	40 MINEHART	84 $m_{\nu_j}=10$ MeV
< 5 × 10 ⁻⁶	90	41 HAYANO	82 $m_{\nu_j}=330$ MeV
< 1 × 10 ⁻⁴	90	41 HAYANO	82 $m_{\nu_j}=70$ MeV
< 9 × 10 ⁻⁷	90	41 HAYANO	82 $m_{\nu_j}=250$ MeV
< 1 × 10 ⁻¹	90	40 ABELA	81 $m_{\nu_j}=4$ MeV
< 7 × 10 ⁻⁵	90	40 ABELA	81 $m_{\nu_j}=10.5$ MeV
< 2 × 10 ⁻⁴	90	40 ABELA	81 $m_{\nu_j}=11.5$ MeV
< 2 × 10 ⁻⁵	90	40 ABELA	81 $m_{\nu_j}=16-30$ MeV
< 2 × 10 ⁻⁵	95	41 ASANO	81 $m_{\nu_j}=170$ MeV
< 3 × 10 ⁻⁶	95	41 ASANO	81 $m_{\nu_j}=210$ MeV
< 3 × 10 ⁻⁶	95	41 ASANO	81 $m_{\nu_j}=230$ MeV
< 1 × 10 ⁻²	95	40 CALAPRICE	81 $m_{\nu_j}=7$ MeV
< 3 × 10 ⁻³	95	40 CALAPRICE	81 $m_{\nu_j}=33$ MeV

⁴⁰ π⁺ → μ⁺ ν_μ peak search experiment.

⁴¹ K⁺ → μ⁺ ν_μ peak search experiment.

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

Peak Search in Muon Capture

Limits on $|U_{2j}|^2$ as function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1 \times 10^{-1}$		DEUTSCH 83		$m_{\nu_j}=45$ MeV
$<7 \times 10^{-3}$		DEUTSCH 83		$m_{\nu_j}=70$ MeV
$<1 \times 10^{-1}$		DEUTSCH 83		$m_{\nu_j}=85$ MeV

Searches for Decays of Massive ν

Limits on $|U_{2j}|^2$ as function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.2 \times 10^{-8}$	95	ADEVA 90S L3		$m_{\nu_j} = 20$ GeV
$<5.1 \times 10^{-10}$	95	ADEVA 90S L3		$m_{\nu_j} = 40$ GeV
all values ruled out	95	42 BURCHAT 90 MRK2		$m_{\nu_j} < 19.6$ GeV
$<1 \times 10^{-10}$	95	42 BURCHAT 90 MRK2		$m_{\nu_j} = 22$ GeV
$<1 \times 10^{-11}$	95	42 BURCHAT 90 MRK2		$m_{\nu_j} = 41$ GeV
all values ruled out	95	DECAMP 90F ALEP		$m_{\nu_j} = 25.0-42.7$ GeV
$<1 \times 10^{-13}$	95	DECAMP 90F ALEP		$m_{\nu_j} = 42.7-45.7$ GeV
$<5 \times 10^{-4}$	90	43 KOPEIKIN 90 CNTR		$m_{\nu_j} = 5.2$ MeV
$<5 \times 10^{-3}$	90	AKERLOF 88 HRS		$m_{\nu_j}=1.8$ GeV
$<2 \times 10^{-5}$	90	AKERLOF 88 HRS		$m_{\nu_j}=4$ GeV
$<3 \times 10^{-6}$	90	AKERLOF 88 HRS		$m_{\nu_j}=6$ GeV
$<1 \times 10^{-7}$	90	BERNARDI 88 CNTR		$m_{\mu_j}=200$ MeV
$<3 \times 10^{-9}$	90	BERNARDI 88 CNTR		$m_{\mu_j}=300$ MeV
$<4 \times 10^{-4}$	90	44 MISHRA 87 CNTR		$m_{\nu_j}=1.5$ GeV
$<4 \times 10^{-3}$	90	44 MISHRA 87 CNTR		$m_{\nu_j}=2.5$ GeV
$<0.9 \times 10^{-2}$	90	44 MISHRA 87 CNTR		$m_{\nu_j}=5$ GeV
<0.1	90	44 MISHRA 87 CNTR		$m_{\nu_j}=10$ GeV
$<8 \times 10^{-4}$	90	BADIER 86 CNTR		$m_{\nu_j}=600$ MeV
$<1.2 \times 10^{-5}$	90	BADIER 86 CNTR		$m_{\nu_j}=1.7$ GeV
$<3 \times 10^{-8}$	90	BERNARDI 86 CNTR		$m_{\nu_j}=200$ MeV
$<6 \times 10^{-9}$	90	BERNARDI 86 CNTR		$m_{\nu_j}=350$ MeV
$<1 \times 10^{-6}$	90	DORENBOS... 86 CNTR		$m_{\nu_j}=500$ MeV
$<1 \times 10^{-7}$	90	DORENBOS... 86 CNTR		$m_{\nu_j}=1600$ MeV
$<0.8 \times 10^{-5}$	90	45 COOPER... 85 HLBC		$m_{\nu_j}=0.4$ GeV
$<1.0 \times 10^{-7}$	90	45 COOPER... 85 HLBC		$m_{\nu_j}=1.5$ GeV

42 BURCHAT 90 includes the analyses reported in JUNG 90, ABRAMS 89C, and WENDT 87.

43 KOPEIKIN 90 find no m_{ν_j} in the interval 1–6.3 MeV at 90%CL for maximal mixing.

44 See also limits on $|U_{3j}|^2$ from WENDT 87.

45 COOPER-SARKAR 85 also give limits based on model-dependent assumptions for ν_τ flux. We do not list these. Note that for this bound to be nontrivial, j is not equal to 3, i.e. ν_j cannot be the dominant mass eigenstate in ν_τ since $m_{\nu_3} < 70$ MeV (ALBRECHT 85). Also, of course, j is not equal to 1 or 2, so a fourth generation would be required for this bound to be nontrivial.

Limits on $|U_{3j}|^2$ as a Function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<6.2 \times 10^{-8}$	95	ADEVA 90S L3		$m_{\nu_j} = 20$ GeV
$<5.1 \times 10^{-10}$	95	ADEVA 90S L3		$m_{\nu_j} = 40$ GeV
all values ruled out	95	46 BURCHAT 90 MRK2		$m_{\nu_j} < 19.6$ GeV
$<1 \times 10^{-10}$	95	46 BURCHAT 90 MRK2		$m_{\nu_j} = 22$ GeV
$<1 \times 10^{-11}$	95	46 BURCHAT 90 MRK2		$m_{\nu_j} = 41$ GeV
all values ruled out	95	DECAMP 90F ALEP		$m_{\nu_j} = 25.0-42.7$ GeV
$<1 \times 10^{-13}$	95	DECAMP 90F ALEP		$m_{\nu_j} = 42.7-45.7$ GeV
$<5 \times 10^{-2}$	80	AKERLOF 88 HRS		$m_{\nu_j}=2.5$ GeV
$<9 \times 10^{-5}$	80	AKERLOF 88 HRS		$m_{\nu_j}=4.5$ GeV

46 BURCHAT 90 includes the analyses reported in JUNG 90, ABRAMS 89C, and WENDT 87.

Limits on $|U_{aj}|^2$

Where $a = 1, 2$ from ρ parameter in μ decay.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<1 \times 10^{-2}$	68	SHROCK 81B THEO		$m_{\nu_j}=10$ MeV
$<2 \times 10^{-3}$	68	SHROCK 81B THEO		$m_{\nu_j}=40$ MeV
$<4 \times 10^{-2}$	68	SHROCK 81B THEO		$m_{\nu_j}=70$ MeV

Limits on $|U_{1j} \times U_{2j}|$ as Function of m_{ν_j}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$<3 \times 10^{-5}$	90	47 BARANOV 93		$m_{\nu_j}=80$ MeV
$<3 \times 10^{-6}$	90	47 BARANOV 93		$m_{\nu_j}=160$ MeV
$<6 \times 10^{-7}$	90	47 BARANOV 93		$m_{\nu_j}=240$ MeV
$<2 \times 10^{-7}$	90	47 BARANOV 93		$m_{\nu_j}=320$ MeV
$<9 \times 10^{-5}$	90	BERNARDI 86 CNTR		$m_{\nu_j}=25$ MeV
$<3.6 \times 10^{-7}$	90	BERNARDI 86 CNTR		$m_{\nu_j}=100$ MeV
$<3 \times 10^{-8}$	90	BERNARDI 86 CNTR		$m_{\nu_j}=200$ MeV
$<6 \times 10^{-9}$	90	BERNARDI 86 CNTR		$m_{\nu_j}=350$ MeV
$<1 \times 10^{-2}$	90	BERGSMA 83B CNTR		$m_{\nu_j}=10$ MeV
$<1 \times 10^{-5}$	90	BERGSMA 83B CNTR		$m_{\nu_j}=140$ MeV
$<7 \times 10^{-7}$	90	BERGSMA 83B CNTR		$m_{\nu_j}=370$ MeV

47 BARANOV 93 is a search for neutrino decays into $e^+ e^- \nu_e$ using a beam dump experiment at the 70 GeV Serpukhov proton synchrotron.

(B) Bounds from ν Reactions

NOTE ON NEUTRINO OSCILLATION EXPERIMENTS

Experimental results on neutrino oscillations are often presented as allowed regions on a plot of $|\Delta m^2|$ as a function of $\sin^2 2\theta$, where $\Delta m^2 = m_{\nu_i}^2 - m_{\nu_j}^2$. Although there are three flavors, data are usually analyzed assuming an oscillation between just two of them, e.g., $\nu_\tau \leftrightarrow \nu_e$. The same remark applies to lepton-number violating mixing between two states, e.g., $\nu_e \leftrightarrow \bar{\nu}_\mu$ or $\nu_\mu \leftrightarrow \bar{\nu}_\mu$.

The simplest situation occurs in an “appearance” experiment, where one searches for interactions by neutrinos of a variety not expected in the beam. An example is the search for $\bar{\nu}_e$ interactions in a beam of neutrinos from the π^+ decay chain, which (among other possibilities) might be taken as evidence for $\bar{\nu}_\mu \rightarrow \bar{\nu}_e$. For oscillation between two states, the probability that the “wrong” state will appear is given by Eq. 5 in Shrock’s “Note on Neutrinos” at the beginning of the Quark and Lepton Full Listings. For our present purposes, this may be rewritten as

$$P = \sin^2 2\theta \sin^2(1.27\Delta m^2 L/E), \quad (1)$$

where L is the distance from the neutrino’s production point to its interaction point, and E is its energy. In the above, $|\Delta m^2|$ is in eV^2 and L/E is in km/GeV or m/MeV . Since in a real experiment L and E have some spread, one must average P over the appropriate distributions. As an example, let us make the somewhat unrealistic assumption that $b \equiv 1.27L/E$ has a Gaussian distribution with standard deviation σ_b about a central value b_0 . Then:

$$\langle P \rangle = \frac{1}{2} \sin^2 2\theta [1 - \cos(2b_0\Delta m^2) \exp(-2\sigma_b^2(\Delta m^2)^2)] \quad (2)$$

The value of $\langle P \rangle$ is set by the experiment. For example, if 230 interactions of the expected flavor are detected and none of the wrong flavor are seen, then $P = 0.010$ at the 90% CL. We can then solve the above expression for $\sin^2 2\theta$ as a function of $|\Delta m^2|$. This function is shown in Fig. 1 for the parameter assumptions given in the caption. Note that:

(a) since the fast oscillations are completely washed out by the resolution for large $|\Delta m^2|$, $\sin^2 2\theta = 2 \langle P \rangle$ in this region;

See key on page 1343

Lepton & Quark Full Listings Massive Neutrinos and Lepton Mixing

- (b) the maximum excursion to the left is to approximately $\langle P \rangle$, and it occurs at $|\Delta m^2| = \pi/2b_0 \text{ eV}^2$;
 (c) for large $\sin^2 2\theta$, $\Delta m^2 \propto (\sin^2 2\theta)^{-1/2}$; and
 (d) the intercept at $\sin^2 2\theta = 1$ is at $\sqrt{\langle P \rangle}/b_0$.

The intercept for large $|\Delta m^2|$ is just a measure of running time and backgrounds, while the intercept at $\sin^2 2\theta = 1$ also depends on the mean value of L/E . The wiggles depend on the experimental resolution, but aside from such details the two intercepts completely describe the exclusion region: For large $|\Delta m^2|$, $\sin^2 2\theta$ is constant, and for large $\sin^2 2\theta$ the constant slope is known. For these reasons, it is (nearly) sufficient to summarize the results of an experiment by stating the two intercepts, as is done in the following tables. The reader is referred to the original papers for the two-dimensional plots expressing the actual limits.

If a positive effect is claimed, then the excluded region becomes an included region. This is the case for the HIRATA 92 analysis of $R(\mu/e)$ for atmospheric neutrinos.

In a “disappearance” experiment, one looks for the attenuation of the beam neutrinos (for example, ν_k) by mixing with at least one other neutrino eigenstate. (We label such experiments as $\nu_k \nrightarrow \nu_k$.) These experiments fall into two general classes:

- (a) Those in which the beam neutrino flux is known, from theory or other measurements. In the high- $|\Delta m^2|$ region, where the oscillation length is small compared to the size of the apparatus, the oscillations are in both directions and the beam intensity is reduced by a factor of two (for two-component mixing). In this case, indicated qualitatively by the “Disappearance A” curve in Fig. 1, sensitivity is maintained for large $|\Delta m^2|$, but with no simple rule relating this asymptote to the maximum excursion to the left. An example is provided by the VUILLEUMIER 82 measurements at the Gösgen reactor.
- (b) Those in which the intensity must be measured in the apparatus itself (two detectors, or a “long” detector). Then above some minimum $|\Delta m^2|$ the equilibrium is established upstream, and there is no change in intensity over the length of the apparatus. As a result, sensitivity is lost at high $|\Delta m^2|$, as is qualitatively indicated by the curve “Disappearance B” in Fig. 1. See, for example, DYDAK 84.

Finally, there are more complicated cases, such as in the HIRATA 92B analysis of the Kamiokande II solar neutrino data in terms of the MSW parameters. An irregular region on the $|\Delta m^2|$ vs $\sin^2 2\theta$ is excluded for a combination of physical reasons. It is difficult to represent adequately these graphical data within the strictures of our tables.




Figure 1: Neutrino oscillation parameter ranges excluded by a toy experiment in which one searches for the appearance of neutrinos not expected in the beam. The probability of appearance, in this case 0.5% at some specified CL, is set by the number of right-flavor events observed and/or information about the flux and cross sections. Here it is assumed that $\langle L/E \rangle = 1 \text{ km GeV}^{-1}$, and that the distribution of L/E is Gaussian with a 20% standard deviation. The wiggle structure is determined by the resolution function, and the intercepts are determined by the appearance probability and $\langle L/E \rangle$. The leftmost excursion relative to the high- $|\Delta m^2|$ limit and the slope of the lower part of the curve are independent of the experiment. In a disappearance experiment, high- $|\Delta m^2|$ sensitivity is lost unless the incident flux is known. These two possibilities are shown qualitatively by the dashed lines marked “Disappearance A” and “Disappearance B.”

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

Solar ν Experiments

Theoretical calculations of the expected solar rate are shown for comparison with experiment. Note that the expected value in "solar neutrino units" (SNU; 1 SNU = 10^{-36} captures atom $^{-1}$ s $^{-1}$) depends on the detection reaction.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$79 \pm 10 \pm 6$ SNU		48 ANSELMANN 94	GALX	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
$20^{+15}_{-20} \pm 32$ SNU		49 ABAZOV 91B	SAGE	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
< 79 SNU	90	49 ABAZOV 91B	SAGE	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
$(0.46 \pm 0.05 \pm 0.06) \times \text{SSM}$		50 HIRATA 90	KAM2	Water Cerenkov
2.33 ± 0.25 SNU		51 DAVIS 89	HOME	^{37}Cl radiochemical
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$87 \pm 14 \pm 7$ SNU		52 ANSELMANN 93	GALX	Repl. by ANSELMANN 94
		53 BAHCALL 93		
		54 HAMPEL 93	RVUE	
6.4 ± 1.4 SNU		55 TURCK-CHI... 93B	THEO	^{37}Cl radiochemical
$(4.4 \pm 1.1) \times 10^{-6} \text{ cm}^{-2} \text{ s}^{-1}$		55 TURCK-CHI... 93B	THEO	Water Cerenkov, $E \geq 7.5$ MeV
123 ± 7 SNU		55 TURCK-CHI... 93B	THEO	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
$83 \pm 19 \pm 8$ SNU		56 ANSELMANN 92	GALX	$^{71}\text{Ga} \rightarrow ^{71}\text{Ge}$
		57 ANSELMANN 92B		
8.0 ± 3.0 SNU		58 BAHCALL 92		^{37}Cl prediction
132^{+21}_{-17} SNU		58 BAHCALL 92		^{71}Ga prediction
		59 GARCIA 91	CNTR	Nuclear physics
		60 HIRATA 91	KAM2	
		61 FILIPPONE 90	THY	
		62 HIRATA 90B	KAM2	
7.9 ± 2.6 SNU		63 BAHCALL 88	THEO	^{37}Cl prediction; total theor. range
132^{+20}_{-17} SNU		63 BAHCALL 88	THEO	^{71}Ga prediction; total theor. range
5.8 ± 1.3 SNU		TURCK-CHI... 88	THEO	^{37}Cl prediction
125 ± 5 SNU		TURCK-CHI... 88	THEO	^{71}Ga prediction
5.6 SNU		FILIPPONE 83	THEO	^{37}Cl prediction
7.6 ± 3.3 SNU		64 BAHCALL 82		^{37}Cl prediction
106^{+13}_{-8} SNU		64 BAHCALL 82		^{71}Ga prediction
7.0 ± 3.0 SNU		FILIPPONE 82	THEO	^{37}Cl prediction
6.9 ± 1.0 SNU		FWALLER 82	THEO	^{37}Cl prediction
7.3 SNU		BAHCALL 80	THEO	^{37}Cl prediction

See also the reviews by BAHCALL 89 and DAVIS 89.

- 48 ANSELMANN 94 result is for a total of 15 initial runs ("GALLEX I") (see footnote for ANSELMANN 93) and 15 new runs ("GALLEX II"). The new runs yield $78 \pm 13 \pm 5$ SNU, confirming to the initial result.
- 49 ABAZOV 91B uses a 30 ton gallium detector to search for the reaction $^{71}\text{Ga}(\nu_e, e)^{71}\text{Ge}$. Result is for the first 6 months of operation. The conference report GAVRIN 92 updates this result to $58^{+17}_{-24} \pm 14$ SNU. Since this reaction has a threshold neutrino energy of 0.236 MeV, it is sensitive to the low-energy neutrinos from the main pp chain (whose maximum energy is 0.420 MeV). The upper limit quoted is to be compared with the theoretical expectation of about 130 SNU; see BAHCALL 89B and references therein.
- 50 HIRATA 90 data consists of 1040 days with threshold $E_e > 9.3$ MeV (first 450 days) or $E_e > 7.5$ MeV. "The total data sample is also analyzed for short-term variations; within the statistical error, no significant variation is observed." The flux is scaled by the value relative to the standard solar model (SSM) prediction. A theoretical flux of $(5.8 \pm 2.1) \times 10^6 \text{ cm}^{-2} \text{ s}^{-1}$ is cited, with the central value corresponding to 7.9 SNU for ^{37}Cl experiment (but see TURCK-CHIEZE 93B and other theoretical calculations.) The analysis is more fully reported in HIRATA 91B.
- 51 DAVIS 89 is the average from the ^{37}Cl experiment at the Homestake Mine (HOME) from 1970–1988. Earlier averages are given in the references therein.
- 52 ANSELMANN 93 result is for 21 runs of combined GALLEX I and GALLEX II data from inverse beta decay reactions in an aqueous solution of Ga^{3+}Cl (30.3 tons of natural Ga) in the Gran Sasso underground laboratory. The 15 runs of GALLEX I yield $81 \pm 17 \pm 9$, which replaces the preliminary result of ANSELMANN 92. The first 6 runs with GALLEX II yield $97 \pm 23 \pm 7$ SNU. A typical solar model rate prediction is for 132 SNU, but see the reviews by Bahcall and others.
- 53 BAHCALL '93 is a study of 1000 solar models in which each input parameter is chosen from a normal distribution with the appropriate mean and error. It is concluded that "Even if one abuses the solar models by artificially imposing consistency with the Kamiokande experiment, the resulting predictions of all 1000 of the 'fudged' solar models are inconsistent with the result of the chlorine experiment."
- 54 HAMPEL 93, by a member of the GALLEX collaboration, is a discussion of possible scenarios to explain the combined solar neutrino experimental data.
- 55 TURCK-CHIEZE 93B proposes new results on the solar neutrino predictions and acoustic mode frequencies. See also TURCK-CHIEZE 93 for an extensive review (233 pages, 524 references) concerning the solar interior. Table 17 provides a particularly useful comparison of experiment and theory as of mid-1993.
- 56 ANSELMANN 92 reports first observation of solar neutrinos from the pp reaction, using the GALLEX solar neutrino detector in the Gran Sasso Underground Laboratory. The total rate is 2σ below the solar model predictions in the range 124–132 SNU's, but they state that "Our result is consistent with the presence of the full pp neutrino flux expected according to the 'standard solar model' together with a reduced flux of ^8B and ^7Be neutrinos as observed by the Homestake and Kamiokande experiments."
- 57 ANSELMANN 92B discusses implications of the GALLEX observations reported in ANSELMANN 92. They state that "To fit this result together with those of the chlorine and Kamiokande experiments requires severe stretching of the solar models but does not rule out such a procedure, leaving the possibility of massless neutrinos... The Mikheyev-Smirnov-Wolfenstein mechanism provides a good fit, and the GALLEX result fixes the $\Delta(m^2)$ and $\sin^2 2\theta$ parameters in two very confined ranges (around $\Delta(m^2) = 6 \times 10^{-6} \text{ eV}^2$ and $\sin^2 2\theta = 7 \times 10^{-3}$ or $\Delta(m^2) = 8 \times 10^{-6} \text{ eV}^2$ and $\sin^2 2\theta = 0.6$). Explanations of the solar neutrino problem based on the decay or magnetic interactions of neutrinos are disfavored."
- 58 BAHCALL 92 is an extensive up-to-date discussion of the solar neutrino flux calculations with predicted event rates for various different solar neutrino detectors. First published calculation that includes diffusion. "The quoted errors represent the total theoretical range and include the effects on the model predictions of 3σ errors in measured input parameters."
- 59 GARCIA 91 reports a new study of $^{37}\text{Ca}\beta$ decays, with the result that the BAHCALL 88 SSM prediction for ^{37}Cl should be increased from 7.9 to 8.1 SNU.
- 60 HIRATA 91 reports a search for day-night and semi-annual variations in the solar neutrino flux observed in the Kamiokande II Detector. The sample is the same 1040 day counting period used for HIRATA 90 and HIRATA 90B. "Within statistical error, no such short-time variations were observed." This result was used to constrain neutrino oscillation parameters, in the framework of oscillations between two mass eigenstates. "A region defined by $\sin^2 2\theta > 0.02$ and $2 \times 10^{-6} \text{ eV}^2 < \Delta(m^2) < 1 \times 10^{-5} \text{ eV}^2$ is excluded at the 90% CL without any assumptions on the absolute value of the expected solar neutrino flux."
- 61 FILIPPONE 90 is a statistical analysis of solar neutrino data to test hypotheses of time dependence. The authors state "we have shown that in our unbiased analysis, the hypothesis of a time-independent ^{37}Cl neutrino capture rate is marginally rejected, having only 2% probability. However, it is disturbing that we are not able to find a simple hypothesis of time variation that would describe the data well. A capture rate anticorrelated with sunspot number, although more probable than the constant rate hypothesis, has a probability of only 6%. One possible explanation of these results is simply the poor statistics of the ^{37}Cl experiment."
- 62 HIRATA 90B gives an analysis of the implications of these data for allowed values of $\Delta(m^2)$ and $\sin^2 2\theta$ describing neutrino mixing between two mass eigenstates, in the model of resonant (MSW) neutrino oscillations. The possibility of regeneration as the neutrinos pass through the earth is neglected. Two limits are given, the first from the measured event rate alone, and the second from the combination of the measured event rate and the recoil electron energy spectrum. The latter "disfavor the region of adiabatic solutions $\Delta(m^2) \sim 1.3 \times 10^{-4} \text{ eV}^2$ and $7.2 \times 10^{-4} < \sin^2 2\theta < 6.3 \times 10^{-3}$ at 90% CL." The allowed regions in $\sin^2 2\theta$ vs. $\Delta(m^2)$ are given graphically; see Figs. 2(a) and 2(b) in the paper.
- 63 BAHCALL 88 "total theoretical range is calculated by evaluating the 3σ uncertainties for all measured input parameters and using the full spread in calculated values for input quantities that cannot be measured; the uncertainties from different quantities are combined quadratically." (Quotation from BAHCALL 89, p. 301.)
- 64 BAHCALL 82 quotes "effective 3σ errors." First extensive discussion of formal uncertainties in the problem.

See key on page 1343

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

Deep Underground Detector Experiments

$R(\mu/e) = (\text{Measured Ratio } \mu/e) / (\text{Expected Ratio } \mu/e)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.60 ^{+0.07} _{-0.06} ± 0.05	66 HIRATA	92 KAM2	Water Cerenkov
	67 BERGER	90B FREJ	Calorimeter

65 BECKER-SZENDY 92B reports the fraction of nonshowing events (mostly muons from atmospheric neutrinos) as $0.36 \pm 0.02 \pm 0.02$, as compared with expected fraction $0.51 \pm 0.01 \pm 0.05$. After cutting the energy range to the KAM2 limits, BEIER 92 finds $R(\mu/e)$ very close to the KAM2 value.

66 HIRATA 92 uses this ratio because both experimental and theoretical uncertainties in ν_μ and ν_e flux cancel. Part of data set is same as in HIRATA 88 (below), where the equivalent ratio was $0.56^{+0.09}$ _{-0.08}.

67 BERGER 90B reports $e/\mu = 0.53-0.57 \pm 0.09$ (data) and $0.56-0.61 \pm 0.08$ (Monte Carlo), where the range is from different analysis methods. There is thus no significant departure from expectation. See also BEIER 92.

$R(\nu_\mu) = (\text{Measured Flux of } \nu_\mu) / (\text{Expected Flux of } \nu_\mu)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.59 ± 0.07	68 CASPER	91 IMB	Water Cherenkov
0.95 ± 0.22	69 AGLIETTA	89 NUSX	
0.62 ± 0.17	70 HIRATA	88 KAM2	Water Cherenkov
	71 BOLIVIEV	81 Baksan	
	72 CROUCH	78	Case Western/UCI

68 CASPER 91 correlates showering/nonshowing signature of single-ring events with parent atmospheric-neutrino flavor. They find nonshowing ($\approx \nu_\mu$ induced) fraction is $0.41 \pm 0.03 \pm 0.02$, as compared with expected 0.51 ± 0.05 (syst).

69 AGLIETTA 89 finds no evidence for any anomaly in the neutrino flux. They define $\rho = (\text{measured number of } \nu_e\text{'s}) / (\text{measured number of } \nu_\mu\text{'s})$. They report $\rho(\text{measured}) = \rho(\text{expected}) = 0.96^{+0.32}$ _{-0.28}.

70 HIRATA 88 error is statistical.

71 From this data BOLIVIEV 81 obtain the limit $\Delta(m^2) \leq 6 \times 10^{-3} \text{ eV}^2$ for maximal mixing, $\nu_\mu \leftrightarrow \nu_\mu$ type oscillation.

$\sin^2(2\theta)$ for Given $\Delta(m^2)$ ($\nu_e \leftrightarrow \nu_\mu$)

For a review see BAHCALL 89.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.33	90	72 HIRATA	92 KAM2	$\Delta(m^2) > 0.004 \text{ eV}^2$
<0.47	90	73 BERGER	90B FREJ	$\Delta(m^2) > 1 \text{ eV}^2$
<0.14	90	74 LOSECCO	87 IMB	$\Delta(m^2) = 0.00011 \text{ eV}^2$

72 HIRATA 92 states that the allowed region for $\nu_e \leftrightarrow \nu_\mu$ conflicts with the constraints from the solar neutrino data (HIRATA 90B).

73 BERGER 90B uses the Frejus detector to search for oscillations of atmospheric neutrinos. Bounds are for both neutrino and antineutrino oscillations.

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$ ($\nu_e \leftrightarrow \nu_\mu$)

VALUE (10^{-5} eV^2)	CL%	DOCUMENT ID	TECN
<150	90	74 BERGER	90B FREJ

74 BERGER 90B uses the Frejus detector to search for oscillations of atmospheric neutrinos. Bounds are for both neutrino and antineutrino oscillations.

$\sin^2(2\theta)$ for Given $\Delta(m^2)$ ($\nu_\mu \leftrightarrow \nu_\tau$)

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.5	90	75 BECKER-SZ...	92 IMB	$\Delta(m^2) = 1-2 \times 10^{-4} \text{ eV}^2$
>0.42	90	HIRATA	92 KAM2	$\Delta(m^2) > 0.001 \text{ eV}^2$
<0.6	90	76 BERGER	90B FREJ	$\Delta(m^2) > 1 \text{ eV}^2$

75 BECKER-SZENDY 92 uses upward-going muons to search for atmospheric ν_μ oscillations. The fraction of muons which stop in the detector is used to search for deviations in the expected spectrum. No evidence for oscillations is found.

76 BERGER 90B uses the Frejus detector to search for oscillations of atmospheric neutrinos. Bounds are for both neutrino and antineutrino oscillations.

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$ ($\nu_\mu \leftrightarrow \nu_\tau$)

VALUE (10^{-5} eV^2)	CL%	DOCUMENT ID	TECN
<350	90	77 BERGER	90B FREJ

77 BERGER 90B uses the Frejus detector to search for oscillations of atmospheric neutrinos. Bounds are for both neutrino and antineutrino oscillations.

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$ ($\nu_\mu \rightarrow \nu_s$)

ν_s means ν_τ or any sterile (noninteracting) ν .

VALUE (10^{-5} eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
<3000 (or <550)	90	78 OYAMA	89	Kamiokande II
<4.2 or >54.	90	BIONTA	88 IMB	Flux has $\nu_\mu, \bar{\nu}_\mu, \nu_e,$ and $\bar{\nu}_e$

78 OYAMA 89 gives a range of limits, depending on assumptions in their analysis. They argue that the region $\Delta(m^2) = (100-1000) \times 10^{-5} \text{ eV}^2$ is not ruled out by any data for large mixing.

Reactor $\bar{\nu}$ Experiments

Events (Observed/Expected) from Reactor $\bar{\nu}_e$ Experiments

VALUE	DOCUMENT ID	COMMENT
1.05 ± 0.02 ± 0.05	VUILLEUMIER 82	$\bar{\nu}_e p \rightarrow e^+ n$
0.955 ± 0.035 ± 0.110	79 KWON	81 $\bar{\nu}_e p \rightarrow e^+ n$
0.89 ± 0.15	79 BOEHM	80 $\bar{\nu}_e p \rightarrow e^+ n$
0.38 ± 0.21	80,81 REINES	80
0.40 ± 0.22	80,81 REINES	80

79 KWON 81 represents an analysis of a larger set of data from the same experiment as BOEHM 80.

80 REINES 80 involves comparison of neutral- and charged-current reactions $\bar{\nu}_e d \rightarrow n p \bar{\nu}_e$ and $\bar{\nu}_e d \rightarrow n n e^+$ respectively. Combined analysis of reactor $\bar{\nu}_e$ experiments was performed by SILVERMAN 81.

81 The two REINES 80 values correspond to the calculated $\bar{\nu}_e$ fluxes of AVIGNONE 80 and DAVIS 79 respectively.

----- $\bar{\nu}_e \rightarrow \bar{\nu}_e$ -----

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
<0.0083	90	VIDYAKIN	90	$\bar{\nu}_e p \rightarrow e^+ n$
<0.04	90	82 KETOV	92	
<0.05	90	83 AFONIN	88 CNTR	$\bar{\nu}_e p \rightarrow e^+ n$
<0.014	68	84 AFONIN	87	$\bar{\nu}_e p \rightarrow e^+ n$
<0.05	68	85 VIDYAKIN	87	$\bar{\nu}_e p \rightarrow e^+ n$
<0.019	90	86 AFONIN	86	$\bar{\nu}_e p \rightarrow e^+ n$
<0.07	90	86 ZACEK	86	$\bar{\nu}_e p \rightarrow e^+ n$
<0.02	90	87 AFONIN	85	$\bar{\nu}_e p \rightarrow e^+ n$
<0.016	90	87 ZACEK	85	$\bar{\nu}_e p \rightarrow e^+ n$
<0.1	90	88 GABATHULER	84	$\bar{\nu}_e p \rightarrow e^+ n$
<0.13	90	AFONIN	83	$\bar{\nu}_e p \rightarrow e^+ n$
		BELENKII	83	$\bar{\nu}_e p \rightarrow e^+ n$

82 KETOV 92 is a limit from search for $\bar{\nu}_e$ oscillations using the reaction $\bar{\nu}_e p \rightarrow e^+ n$ with the $\bar{\nu}_e$ flux from the Rovno power reactor. They obtain the ratio $R = 0.976 \pm 0.02 \pm 0.015$ for the corrected flux ratios at 12 m and 18 m from the reactor core, and thus report no evidence for neutrino oscillations over distances of order 10 m. Virtually no experimental details are given.

83 Several different methods of data analysis are used in AFONIN 88. We quote the most stringent limits.

84 AFONIN 86 and AFONIN 87 also give limits on $\sin^2(2\theta)$ for intermediate values of $\Delta(m^2)$.

85 VIDYAKIN 87 bound is for $L = 32.8$ and 92.3 m distance from two reactors.

86 This bound is from data for $L = 37.9$ m, 45.9 m, and 64.7 m distance from Gosgen reactor.

87 See the comment for ZACEK 85 in the section on $\sin^2(2\theta)$ below.

88 This bound comes from a combination of the VUILLEUMIER 82 data at distance 37.9 m from Gosgen reactor and new data at 45.9 m.

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.14	68	89 VIDYAKIN	87	$\bar{\nu}_e p \rightarrow e^+ n$
<0.2	90	90 AFONIN	88 CNTR	$\bar{\nu}_e p \rightarrow e^+ n$
<0.21	68	AFONIN	87	$\bar{\nu}_e p \rightarrow e^+ n$
<0.21	90	91 ZACEK	86	$\bar{\nu}_e p \rightarrow e^+ n$
<0.34	90	AFONIN	85	$\bar{\nu}_e p \rightarrow e^+ n$
<0.19	90	92 ZACEK	85	$\bar{\nu}_e p \rightarrow e^+ n$
<0.16	90	93 GABATHULER	84	$\bar{\nu}_e p \rightarrow e^+ n$
<0.4		94 BELENKII	83	$\bar{\nu}_e p \rightarrow e^+ n$

89 VIDYAKIN 87 bound is for $L = 32.8$ and 92.3 m distance from two reactors.

90 Several different methods of data analysis are used in AFONIN 88. We quote the most stringent limits. Different upper limits on $\sin^2 2\theta$ apply at intermediate values of $\Delta(m^2)$.

91 This bound is from data for $L = 37.9$ m, 45.9 m, and 64.7 m distance from Gosgen reactor.

92 ZACEK 85 (Gosgen reactor) gives two sets of bounds depending on what assumptions are used in the data analysis. The bounds in figure 3(a) of ZACEK 85 are progressively poorer for large $\Delta(m^2)$ whereas those of figure 3(b) approach a constant. We list the latter. Both sets of bounds use combination of data from 37.9, 45.9, and 64.7 m distance from reactor. ZACEK 85 states "Our experiment excludes this area (the oscillation parameter region allowed by the Bugey data, CAVIGNAC 84) almost completely, thus disproving the indications of neutrino oscillations of CAVIGNAC 84 with a high degree of confidence."

93 This bound comes from a combination of the VUILLEUMIER 82 data at distance 37.9 m from Gosgen reactor and new data at 45.9 m.

94 This bound holds for $\Delta(m^2) > 4 \text{ eV}^2$.

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

$\Delta(m^2)$ for Given $\sin^2(2\theta)$

VALUE (eV^2)	DOCUMENT ID	TECN	COMMENT
0.2 ± 0.1	95 CAVAINAC 84	EMUL	$\bar{\nu}_e p \rightarrow e^+ n$

⁹⁵ $\sin^2(2\theta) = 0.25 \pm 0.1$. These are from best fit to data; see CAVAINAC 84 for plot of allowed regions in these variables. These data from Bugey reactor.

Accelerator Experiments

(I) Appearance experiments

$\nu_e \rightarrow \nu_\tau$

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
< 9	90	USHIDA 86c	EMUL	FNAL
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 44	90	TALEBZADEH 87	HLBC	BEBC

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.12	90	USHIDA 86c	EMUL	FNAL
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.36	90	TALEBZADEH 87	HLBC	BEBC

$\bar{\nu}_e \rightarrow \bar{\nu}_\tau$

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.7	90	FRITZE 80	HYBR	BEBC CERN SPS

⁹⁶ Authors give $P(\nu_e \rightarrow \nu_\tau) < 0.35$, equivalent to above limit.

$\nu_\mu \rightarrow \nu_e$

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.09	90	ANGELINI 86	HLBC	BEBC CERN PS
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 0.1	90	ASTIER 90		
< 1.3	90	ASTIER 89	CNTR	BNL AGS
< 0.19	90	BLUMENFELD 89	CNTR	
	90	AMMOISOV 88	HLBC	SKAT at Serpukhov
	90	BERGSMA 88	CHRM	
	90	LOVERRE 88	RVUE	
< 2.4	90	AHRENS 87	CNTR	BNL AGS
< 1.8	90	BOFILL 87	CNTR	FNAL
< 2.2	90	BRUCKER 86	HLBC	15-ft FNAL
< 0.43	90	AHRENS 85	CNTR	BNL AGS E734
< 0.20	90	BERGSMA 84	CHRM	
< 1.7	90	ARMENISE 81	HLBC	GGM CERN PS
< 0.6	90	BAKER 81	HLBC	15-ft FNAL
< 1.7	90	ERRIQUEZ 81	HLBC	BEBC CERN PS
< 1.2	95	BLIETSCHAU 78	HLBC	GGM CERN PS
< 1.2	95	BELLOTTI 76	HLBC	GGM CERN PS

⁹⁷ ASTIER 90 again finds an excess of electrons, as was reported in earlier papers by this collaboration. However, the authors concede that systematic effects weaken the statistical arguments and the consequent claim (in the earlier papers) of neutrino oscillations. An interpretation of these results in terms of neutrino oscillations seems to be already excluded by the BNL E734 (AHRENS 85) and the Los Alamos E645 (DURKIN 88) experiments.

⁹⁸ ASTIER 89 reports a positive effect with $\nu_e(\text{observed})/\nu_e(\text{expected}) = 2.2 \pm 0.6$ and $\bar{\nu}_e(\text{observed})/\bar{\nu}_e(\text{expected}) = 1.6 \pm 0.9$.

⁹⁹ LOVERRE 88 reports a less stringent, indirect limit based on theoretical analysis of neutral to charged current ratios.

¹⁰⁰ 15ft bubble chamber at FNAL.

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 2.5	90	AMMOISOV 88	HLBC	SKAT at Serpukhov
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 16	90	ASTIER 89	CNTR	BNL AGS
< 8	90	BLUMENFELD 89	CNTR	
	90	BERGSMA 88	CHRM	$\Delta(m^2) \geq 30 eV^2$
	90	LOVERRE 88	RVUE	
< 10	90	AHRENS 87	CNTR	BNL AGS
< 15	90	BOFILL 87	CNTR	FNAL
< 20	90	ANGELINI 86	HLBC	BEBC CERN PS
20 to 40	90	BERNARDI 86b	CNTR	$\Delta(m^2)=5-10$
< 11	90	BRUCKER 86	HLBC	15-ft FNAL
< 3.4	90	AHRENS 85	CNTR	BNL AGS E734
< 240	90	BERGSMA 84	CHRM	
< 10	90	ARMENISE 81	HLBC	GGM CERN PS
< 6	90	BAKER 81	HLBC	15-ft FNAL
< 10	90	ERRIQUEZ 81	HLBC	BEBC CERN PS
< 4	95	BLIETSCHAU 78	HLBC	GGM CERN PS
< 10	95	BELLOTTI 76	HLBC	GGM CERN PS

¹⁰¹ ASTIER 89 reports a positive effect with $\nu_e(\text{observed})/\nu_e(\text{expected}) = 2.2 \pm 0.6$ and $\bar{\nu}_e(\text{observed})/\bar{\nu}_e(\text{expected}) = 1.6 \pm 0.9$.

¹⁰² LOVERRE 88 reports a less stringent, indirect limit based on theoretical analysis of neutral to charged current ratios.

¹⁰³ ANGELINI 86 limit reaches 13×10^{-3} at $\Delta(m^2) \approx 2 eV^2$.

¹⁰⁴ BERNARDI 86b is a typical fit to the data, assuming mixing between two species. As the authors state, this result is in conflict with earlier upper bounds on this type of neutrino oscillations.

¹⁰⁵ 15ft bubble chamber at FNAL.

$\bar{\nu}_\mu \rightarrow \bar{\nu}_e$

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.14	90	106 FREEDMAN 93	CNTR	LAMPF
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 0.11	90	DURKIN 88	CNTR	Repl. by FREEDMAN 93
< 3.1	90	BOFILL 87	CNTR	FNAL
< 2.4	90	TAYLOR 83	HLBC	15-ft FNAL
< 0.91	90	107 NEMETHY 81b	CNTR	LAMPF
< 1	95	BLIETSCHAU 78	HLBC	GGM CERN PS

¹⁰⁶ FREEDMAN 93 is a search at LAMPF for $\bar{\nu}_e$ generated from any of the three neutrino types $\nu_\mu, \bar{\nu}_\mu$, and ν_e which come from the beam stop. The $\bar{\nu}_e$'s would be detected by the reaction $\bar{\nu}_e p \rightarrow e^+ n$.

¹⁰⁷ In reaction $\bar{\nu}_e p \rightarrow e^+ n$.

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.004	95	BLIETSCHAU 78	HLBC	GGM CERN PS
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 0.024	90	108 FREEDMAN 93	CNTR	LAMPF
< 0.014	90	109 DURKIN 88	CNTR	Repl. by FREEDMAN 93
< 0.04	90	BOFILL 87	CNTR	FNAL
< 0.013	90	TAYLOR 83	HLBC	15-ft FNAL
< 0.2	90	109 NEMETHY 81b	CNTR	LAMPF

¹⁰⁸ FREEDMAN 93 is a search at LAMPF for $\bar{\nu}_e$ generated from any of the three neutrino types $\nu_\mu, \bar{\nu}_\mu$, and ν_e which come from the beam stop. The $\bar{\nu}_e$'s would be detected by the reaction $\bar{\nu}_e p \rightarrow e^+ n$.

¹⁰⁹ In reaction $\bar{\nu}_e p \rightarrow e^+ n$.

$\nu_\mu(\bar{\nu}_\mu) \rightarrow \nu_e(\bar{\nu}_e)$

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.075	90	BORODOV... 92	CNTR	BNL E776

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 3	90	BORODOV... 92	CNTR	BNL E776

$\nu_\mu \rightarrow \nu_\tau$

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
< 0.9	90	USHIDA 86c	EMUL	FNAL
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 4.5	90	BATUSOV 90b	EMUL	FNAL
< 10.2	90	BOFILL 87	CNTR	FNAL
< 6.3	90	BRUCKER 86	HLBC	15-ft FNAL
< 4.6	90	ARMENISE 81	HLBC	GGM CERN SPS
< 3	90	BAKER 81	HLBC	15-ft FNAL
< 6	90	ERRIQUEZ 81	HLBC	BEBC CERN SPS
< 3	90	USHIDA 81	EMUL	FNAL

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.004	90	USHIDA 86c	EMUL	FNAL
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 0.06	90	BATUSOV 90b	EMUL	FNAL
< 0.34	90	BOFILL 87	CNTR	FNAL
< 0.088	90	BRUCKER 86	HLBC	15-ft FNAL
< 0.11	90	BALLAGH 84	HLBC	15-ft FNAL
< 0.017	90	ARMENISE 81	HLBC	GGM CERN SPS
< 0.06	90	BAKER 81	HLBC	15-ft FNAL
< 0.05	90	ERRIQUEZ 81	HLBC	BEBC CERN SPS
< 0.013	90	USHIDA 81	EMUL	FNAL

$\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV^2)	CL%	DOCUMENT ID	TECN	COMMENT
< 2.2	90	ASRATYAN 81	HLBC	FNAL
• • • We do not use the following data for averages, fits, limits, etc. • • •				

< 6.5	90	BOFILL 87	CNTR	FNAL
< 7.4	90	TAYLOR 83	HLBC	15-ft FNAL

See key on page 1343

Lepton & Quark Full Listings Massive Neutrinos and Lepton Mixing

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.4 x 10 ⁻²	90	ASRATYAN 81	HLBC	FNAL
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.15	90	BOFILL 87	CNTR	FNAL
<8.8 x 10 ⁻²	90	TAYLOR 83	HLBC	15-ft FNAL

----- $\nu_\mu(\bar{\nu}_\mu) \rightarrow \nu_\tau(\bar{\nu}_\tau)$ -----

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV ²)	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	110 GRUWE 93	CHM2	CERN SPS

110 GRUWE 93 is a search using the CHARM II detector in the CERN SPS wide-band neutrino beam for $\nu_\mu \rightarrow \nu_\tau$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$ oscillations signalled by quasi-elastic ν_τ and $\bar{\nu}_\tau$ interactions followed by the decay $\tau \rightarrow \nu_\tau \pi$. The maximum sensitivity in $\sin^2 2\theta$ (< 6.4 x 10⁻³ at the 90% CL) is reached for $\Delta(m^2) \simeq 50$ eV².

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE (units 10 ⁻³)	CL%	DOCUMENT ID	TECN	COMMENT
<8	90	111 GRUWE 93	CHM2	CERN SPS

111 GRUWE 93 is a search using the CHARM II detector in the CERN SPS wide-band neutrino beam for $\nu_\mu \rightarrow \nu_\tau$ and $\bar{\nu}_\mu \rightarrow \bar{\nu}_\tau$ oscillations signalled by quasi-elastic ν_τ and $\bar{\nu}_\tau$ interactions followed by the decay $\tau \rightarrow \nu_\tau \pi$. The maximum sensitivity in $\sin^2 2\theta$ (< 6.4 x 10⁻³ at the 90% CL) is reached for $\Delta(m^2) \simeq 50$ eV².

----- $\nu_e \rightarrow (\bar{\nu}_e)_L$ -----

This is a limit on lepton family-number violation and total lepton-number violation. $(\bar{\nu}_e)_L$ denotes a hypothetical left-handed $\bar{\nu}_e$. The bound is quoted in terms of $\Delta(m^2)$, $\sin(2\theta)$, and α , where α denotes the fractional admixture of (V+A) charged current.

$\alpha\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV ²)	CL%	DOCUMENT ID	TECN	COMMENT
<0.14	90	112 FREEDMAN 93	CNTR	LAMPF

••• We do not use the following data for averages, fits, limits, etc. •••

<7	90	113 COOPER 82	HLBC	BECB CERN SPS
----	----	---------------	------	---------------

112 FREEDMAN 93 is a search at LAMPF for $\bar{\nu}_e$ generated from any of the three neutrino types $\nu_\mu, \bar{\nu}_\mu,$ and ν_e which come from the beam stop. The $\bar{\nu}_e$'s would be detected by the reaction $\bar{\nu}_e p \rightarrow e^+ n$.

113 COOPER 82 states that existing bounds on V+A currents require α to be small.

$\alpha^2 \sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.032	90	114 FREEDMAN 93	CNTR	LAMPF

••• We do not use the following data for averages, fits, limits, etc. •••

<0.05	90	115 COOPER 82	HLBC	BECB CERN SPS
-------	----	---------------	------	---------------

114 FREEDMAN 93 is a search at LAMPF for $\bar{\nu}_e$ generated from any of the three neutrino types $\nu_\mu, \bar{\nu}_\mu,$ and ν_e which come from the beam stop. The $\bar{\nu}_e$'s would be detected by the reaction $\bar{\nu}_e p \rightarrow e^+ n$.

115 COOPER 82 states that existing bounds on V+A currents require α to be small.

----- $\nu_\mu \rightarrow (\bar{\nu}_e)_L$ -----

See note above for $\nu_e \rightarrow (\bar{\nu}_e)_L$ limit

$\alpha\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV ²)	CL%	DOCUMENT ID	TECN	COMMENT
<0.16	90	116 FREEDMAN 93	CNTR	LAMPF

••• We do not use the following data for averages, fits, limits, etc. •••

<0.7	90	117 COOPER 82	HLBC	BECB CERN SPS
------	----	---------------	------	---------------

116 FREEDMAN 93 is a search at LAMPF for $\bar{\nu}_e$ generated from any of the three neutrino types $\nu_\mu, \bar{\nu}_\mu,$ and ν_e which come from the beam stop. The $\bar{\nu}_e$'s would be detected by the reaction $\bar{\nu}_e p \rightarrow e^+ n$. The limit on $\Delta(m^2)$ is better than the CERN BECB experiment, but the limit on $\sin^2 2\theta$ is almost a factor of 100 less sensitive.

117 COOPER 82 states that existing bounds on V+A currents require α to be small.

$\alpha^2 \sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.001	90	118 COOPER 82	HLBC	BECB CERN SPS

••• We do not use the following data for averages, fits, limits, etc. •••

<0.07	90	119 FREEDMAN 93	CNTR	LAMPF
-------	----	-----------------	------	-------

118 COOPER 82 states that existing bounds on V+A currents require α to be small.

119 FREEDMAN 93 is a search at LAMPF for $\bar{\nu}_e$ generated from any of the three neutrino types $\nu_\mu, \bar{\nu}_\mu,$ and ν_e which come from the beam stop. The $\bar{\nu}_e$'s would be detected by the reaction $\bar{\nu}_e p \rightarrow e^+ n$. The limit on $\Delta(m^2)$ is better than the CERN BECB experiment, but the limit on $\sin^2 2\theta$

----- (ii) Disappearance experiments -----

----- $\nu_e \nrightarrow \nu_e$ -----

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV ²)	CL%	DOCUMENT ID	TECN	COMMENT
<2.3 OUR LIMIT				
< 8	90	BAKER 81	HLBC	15-ft FNAL
<2.3 OR >8	90	NEMETHY 81B	CNTR	LAMPF

••• We do not use the following data for averages, fits, limits, etc. •••

<14.9	90	BRUCKER 86	HLBC	15-ft FNAL
<56	90	DEDEN 81	HLBC	BECB CERN SPS
<10	90	ERRIQUEZ 81	HLBC	BECB CERN SPS

$\sin^2(2\theta)$ for "Large" $\Delta(m^2)$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7 x 10 ⁻²	90	ERRIQUEZ 81	HLBC	BECB CERN SPS

••• We do not use the following data for averages, fits, limits, etc. •••

<0.54	90	BRUCKER 86	HLBC	15-ft FNAL
<0.6	90	BAKER 81	HLBC	15-ft FNAL
<0.3	90	DEDEN 81	HLBC	BECB CERN SPS

----- $\nu_\mu \nrightarrow \nu_\mu$ -----

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

These experiments also allow sufficiently large $\Delta(m^2)$.

VALUE (eV ²)	CL%	DOCUMENT ID	TECN	COMMENT
<0.23 OR >1500 OUR LIMIT				
<0.23 OR >100	90	DYDAK 84	CNTR	
<13 OR >1500	90	STOCKDALE 84	CNTR	

••• We do not use the following data for averages, fits, limits, etc. •••

< 0.29 OR >22	90	BERGSMA 88	CHRM	
<7	90	BELIKOV 85	CNTR	Serpukhov
<8.0 OR >1250	90	STOCKDALE 85	CNTR	
<0.29 OR >22	90	BERGSMA 84	CHRM	
<8.0	90	BELIKOV 83	CNTR	

$\sin^2(2\theta)$ for $\Delta(m^2) = 100\text{eV}^2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	120 STOCKDALE 85	CNTR	FNAL

••• We do not use the following data for averages, fits, limits, etc. •••

<0.17	90	121 BERGSMA 88	CHRM	
<0.07	90	122 BELIKOV 85	CNTR	Serpukhov
<0.27	90	123 BERGSMA 84	CHRM	CERN PS
<0.1	90	124 DYDAK 84	CNTR	CERN PS
<0.02	90	125 STOCKDALE 84	CNTR	FNAL
<0.1	90	126 BELIKOV 83	CNTR	Serpukhov

120 This bound applies for $\Delta(m^2) = 100$ eV². Less stringent bounds apply for other $\Delta(m^2)$; these are nontrivial for $8 < \Delta(m^2) < 1250$ eV².

121 This bound applies for $\Delta(m^2) = 0.7-9$ eV². Less stringent bounds apply for other $\Delta(m^2)$; these are nontrivial for $0.28 < \Delta(m^2) < 22$ eV².

122 This bound applies for a wide range of $\Delta(m^2) > 7$ eV². For some values of $\Delta(m^2)$, the value is less stringent; the least restrictive, nontrivial bound occurs approximately at $\Delta(m^2) = 300$ eV² where $\sin^2(2\theta) < 0.13$ at CL = 90%.

123 This bound applies for $\Delta(m^2) = 1-10$ eV². Less stringent bounds apply for other $\Delta(m^2)$; these are nontrivial for $0.23 < \Delta(m^2) < 90$ eV².

124 This bound applies for $\Delta(m^2) = 110$ eV². Less stringent bounds apply for other $\Delta(m^2)$; these are nontrivial for $13 < \Delta(m^2) < 1500$ eV².

125 Bound holds for $\Delta(m^2) = 20-1000$ eV².

126 Bound holds for $\Delta(m^2) = 20-1000$ eV².

----- $\bar{\nu}_\mu \nrightarrow \bar{\nu}_\mu$ -----

$\Delta(m^2)$ for $\sin^2(2\theta) = 1$

VALUE (eV ²)	CL%	DOCUMENT ID	TECN	COMMENT
<7 OR >1200 OUR LIMIT				
<7 OR >1200	90	STOCKDALE 85	CNTR	

$\sin^2(2\theta)$ for $190 \text{ eV}^2 < \Delta(m^2) < 320 \text{ eV}^2$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	126 STOCKDALE 85	CNTR	FNAL

126 This bound applies for $\Delta(m^2)$ between 190 and 320 or = 530 eV². Less stringent bounds apply for other $\Delta(m^2)$; these are nontrivial for $7 < \Delta(m^2) < 1200$ eV².

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

(C) Searches for Neutrinoless Double β Decay

Limits on an effective Majorana neutrino mass and a lepton-number violating current admixture can be obtained from lifetime limits on $0\nu\beta\beta$ nuclear decay. The derived quantities are highly model-dependent, so the half-life measurements are given first. Where possible we list the references for the matrix elements used in the subsequent analysis. Since rates for the more conventional $2\nu\beta\beta$ decay serve to calibrate the theory, results for this process are also given.

To define the limits on lepton-number violating right-handed current admixtures, we display the relevant part of a phenomenological current-current weak interaction Hamiltonian:

$$H_W = (G_F/\sqrt{2})$$

$$\times (J_L \cdot j_L^\dagger + \eta_{RL} J_R \cdot j_L^\dagger + \eta_{LR} J_L \cdot j_R^\dagger + \eta_{RR} J_R \cdot j_R^\dagger) + \text{h.c.}$$

where $j_L^\mu = \bar{e}_L \gamma^\mu \nu_{eL}$, $j_R^\mu = \bar{e}_R \gamma^\mu \nu_{eR}$, and J_L^μ and J_R^μ are left-handed and right-handed hadronic weak currents. Experiments are not sensitive to $\eta_{RL} (\equiv \kappa)$ but quote limits on η_{LR} and η_{RR} . Many authors use an alternative notation in which $\eta_{LR} \equiv \eta$ and $\eta_{RR} \equiv \lambda$. The limits on $\eta_{LR} = \eta$ are of order 10^{-8} while the limits on $\eta_{RR} = \lambda$ are of order 10^{-6} . The reader is warned that a number of earlier experiments did not distinguish between η_{LR} and η_{RR} . Because of evolving reporting conventions and matrix element calculations, we have not tabulated the admixture parameters for experiments published earlier than 1989.

See the section on Majorana searches for additional limits set by these experiments.

Half-life Measurements and Limits for Double β Decay

In all cases of double beta decay, $(Z,A) \rightarrow (Z+2,A) + 2\beta^- + (0 \text{ or } 2)\bar{\nu}_e$.

$t_{1/2}(10^{21} \text{ yr})$	CL% ISOTOPE	TRANSITION	METHOD	DOCUMENT ID
> 44	68 ^{100}Mo 0ν	$0^+ \rightarrow 0^+$	Si(Li)	ALSTON... 93
>3400	90 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	TPC	127,128 VUILLEUMIER 93
>2600	90 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	TPC	128,129 VUILLEUMIER 93
> 0.21	90 ^{136}Xe 2ν	$0^+ \rightarrow 0^+$	TPC	128 VUILLEUMIER 93
> 2.5	90 ^{130}Te 0ν	$0^+ \rightarrow 0^+$	Bolometer	ALESSAND... 92
> 0.093	90 ^{136}Xe 2ν	$0^+ \rightarrow 0^+$	Drift chamber	ARTEMJEV 92
>1400	90 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	Enriched HPGe	130 BALYSH 92
> 430	90 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	Enriched HPGe	130 BALYSH 92
2.7 \pm 0.1	130Te		Geochem	BERNATOW... 92
7200 \pm 400	128Te		Geochem	131 BERNATOW... 92
> 0.5	90 ^{100}Mo 2ν	$0^+ \rightarrow 2^+$	HPGe	132 BLUM 92
> 0.9	90 ^{100}Mo 2ν	$0^+ \rightarrow 0^+$	HPGe	132 BLUM 92
> 0.6	90 ^{100}Mo 2ν	$0^+ \rightarrow 2^+$	HPGe	132 BLUM 92
> 27	68 ^{82}Se 0ν	$0^+ \rightarrow 0^+$	TPC	ELLIOTT 92
0.108 \pm 0.026	82Se 2ν	$0^+ \rightarrow 0^+$	TPC	ELLIOTT 92
> 0.15	68 ^{100}Mo 2ν	$0^+ \rightarrow 2^+$	Spect	133 KUDOMI 92
> 1.1	68 ^{100}Mo 0ν	$0^+ \rightarrow 2^+$	Spect	133 KUDOMI 92
> 0.08	68 ^{100}Mo 2ν	$0^+ \rightarrow 0^+$	Spect	133 KUDOMI 92
> 0.56	68 ^{100}Mo 0ν	$0^+ \rightarrow 0^+$	Spect	133 KUDOMI 92
> 0.051	68 ^{100}Mo 2ν	$0^+ \rightarrow 4^+$	Spect	133 KUDOMI 92
> 0.63	68 ^{100}Mo 0ν	$0^+ \rightarrow 4^+$	Spect	133 KUDOMI 92
> 0.065	68 ^{100}Mo 2ν	$0^+ \rightarrow 2^+$	Spect	133 KUDOMI 92
> 0.12	68 ^{100}Mo 0ν	$0^+ \rightarrow 2^+$	Spect	133 KUDOMI 92
> 330	90 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	HPGe	134 REUSSER 92
> 65	90 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	HPGe	134 REUSSER 92

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.92 \pm 0.07	76Ge 2ν	$0^+ \rightarrow 0^+$	Enriched HPGe	135 AVIGNONE 91
> 12	95 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	Prop cntr	127,136 BELLOTTI 91
> 10	95 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	Prop cntr	129,136 BELLOTTI 91
> 3.3	95 ^{136}Xe 0ν	$0^+ \rightarrow 2^+$	Prop cntr	136 BELLOTTI 91
> 0.16	95 ^{136}Xe 2ν		Prop cntr	BELLOTTI 91
> 4.7	68 ^{100}Mo 0ν		Spect	EJIRI 91
0.0115 \pm 0.0030	100Mo 2ν		Spect	EJIRI 91
				0.0020
2.0 \pm 0.6	238U		Mass spect	137 HYKAWY 91
> 250	90 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	TPC	127,139 WONG 91
> 170	90 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	TPC	129,139 WONG 91
> 9.5	76 ^{48}Ca 0ν		CaF ₂ scint.	YOU 91
> 0.14	68 ^{100}Mo $0\nu+2\nu$	$0^+ \rightarrow 2^+$	γ in HPGe	BARABASH 90
> 0.042	68 ^{100}Mo $0\nu+2\nu$	$0^+ \rightarrow 0^+$	γ in HPGe	BARABASH 90
> 0.17	68 ^{116}Cd $0\nu+2\nu$	$0^+ \rightarrow 2^+$	γ in HPGe	BARABASH 90
1.12 \pm 0.48	76Ge 2ν	$0^+ \rightarrow 0^+$	HPGe	140 MILEY 90
>1300	68 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	Enriched Ge(Li)	141 VASENKO 90
0.9 \pm 0.1	76Ge 2ν		Enriched Ge(Li)	VASENKO 90
> 4.0	68 ^{100}Mo 0ν	$0^+ \rightarrow 0^+$	Si(Li)	ALSTON... 89
> 0.40	68 ^{100}Mo 0ν	$0^+ \rightarrow 2^+$	Si(Li)	ALSTON... 89
> 3.3	68 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	Ion chamber	127 BARABASH 89
> 2.9	68 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	Ion chamber	129 BARABASH 89
> 1.5	68 ^{136}Xe 0ν	$0^+ \rightarrow 2^+$	Ion chamber	BARABASH 89
> 0.084	68 ^{136}Xe 2ν	$0^+ \rightarrow 0^+$	Ion chamber	BARABASH 89
> 9	90 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	Prop chamber	127 BELLOTTI 89
> 8	90 ^{136}Xe 0ν	$0^+ \rightarrow 0^+$	Prop chamber	129 BELLOTTI 89
> 1.3	68 ^{116}Cd 0ν		116CdWO ₄ scint	DANEVICH 89
> 160	90 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	HPGe	FISHER 89
> 60	90 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	HPGe	FISHER 89
> 60	68 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	HPGe	142 MORALES 88
> 4.7	68 ^{128}Te	$0^+ \rightarrow 2^+$	Ge(Li)	143 BELLOTTI 87
> 4.5	68 ^{130}Te	$0^+ \rightarrow 2^+$	Ge(Li)	143 BELLOTTI 87
> 500	68 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	HPGe	CALDWELL 87
0.11 \pm 0.08	82Se 2ν	$0^+ \rightarrow 0^+$	TPC	ELLIOTT 87B
> 120	68 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	HPGe	BELLOTTI 86
> 12	68 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	HPGe	BELLOTTI 86
> 250	68 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	HPGe	CALDWELL 86
> 50	68 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	HPGe	CALDWELL 86
> 7.0	68 ^{82}Se 0ν	$0^+ \rightarrow 0^+$	TPC	ELLIOTT 86
> 0.10	68 ^{82}Se 2ν	$0^+ \rightarrow 0^+$	TPC	144 ELLIOTT 86
> 50	68 ^{76}Ge 0ν		HPGe	CALDWELL 85
> 2.3	68 ^{76}Ge 0ν	$0^+ \rightarrow 2^+$	Ge(Li)	145 HUBERT 85
> 120	68 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	Ge(Li)	BELLOTTI 84
> 17	68 ^{76}Ge 0ν		Ge	FORSTER 84
> 17	90 ^{76}Ge 0ν		Intrinsic Ge	AVIGNONE 83
> 20	68 ^{76}Ge 0ν	$0^+ \rightarrow 0^+$	Ge(Li)	BELLOTTI 83
> 800	95 ^{128}Te		Geochem	146 KIRSTEN 83
2.60 \pm 0.28	130Te		Geochem	146 KIRSTEN 83

127 Limit in the case of a transition induced by a Majorana mass.

128 VUILLEUMIER 93 data is from the search for neutrinoless double beta decay of ^{136}Xe in the TPC experiment in the St. Gotthard tunnel deep underground laboratory.

129 Limit for lepton-number violating right-handed current-induced (RHC) decay.

130 BALYSH 92 presents results from the Heidelberg-Moscow experiment searching for neutrinoless double beta decay of ^{76}Ge .

131 BERNATOWICZ 92 finds $^{128}\text{Te}/^{130}\text{Te}$ activity ratio from slope of $^{128}\text{Xe}/^{132}\text{Xe}$ vs $^{130}\text{Xe}/^{132}\text{Xe}$ ratios during extraction, and normalizes to lead-dated ages for the ^{130}Te lifetime. The authors state that their results imply that "(a) the double beta decay of ^{128}Te has been firmly established and its half-life has been determined ... without any ambiguity due to trapped Xe interferences. ... (b) Theoretical calculations ... underestimate the [long half-lives of ^{128}Te ^{130}Te] by 1 or 2 orders of magnitude, pointing to a real suppression in the 2ν decay rate of these isotopes. (c) Despite [this], most $\beta\beta$ -models predict a ratio of 2ν decay widths ... in fair agreement with observation." Further details of the experiment are given in BERNATOWICZ 93. Our listed half-life has been revised downward from the published value by the authors, on the basis of reevaluated cosmic-ray ^{128}Xe production corrections.

132 BLUM 92 reports lifetime limits for the decay of ^{100}Mo to several excited states of ^{100}Ru . Limits for decay to the 0_1^+ state are about 30% higher if decay to the 2^+ states are assumed negligible. Uses 99.5% enriched ^{100}Mo .

133 KUDOMI 92 reports lifetime limits for 0ν and 2ν decays to four excited states of the daughter ^{100}Ru . The limits were obtained from searches for the two individual electrons in coincidence with photons from the decays of the excited states. The experiment was performed in the Kamioka underground laboratory. See EJIRI 91 for the group's ground-state transition measurement.

134 REUSSER 92 contains the final results for the search for neutrinoless double beta decay of ^{76}Ge in the Gotthard tunnel underground laboratory.

135 AVIGNONE 91 reports confirmation of the MILEY 90 and VASENKO 90 observations of $2\nu\beta\beta$ decay of ^{76}Ge . Error is 2σ .

136 BELLOTTI 91 uses difference between natural and enriched ^{136}Xe runs to obtain $\beta\beta 0\nu$ limits, leading to "less stringent, but safer limits."

See key on page 1343

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing

- 137 HYKAWY 91 gives new mass spectrometer determination of the ^{76}Ge - ^{76}Ge mass difference, which for given input for the nuclear matrix elements gives information on limits on Majorana masses. Application to recent ^{76}Ge decay experiments produces no new evidence for $\beta\beta(0\nu)$ decay.
- 138 TURKEVICH 91 observes activity in old U sample. The authors compare their results with theoretical calculations. They state "Using the phase-space factors of Boehm and Vogel (BOEHM 87) leads to matrix element values for the ^{238}U transition in the same range as deduced for ^{130}Te and ^{76}Ge . On the other hand, the latest theoretical estimates (STAUDT 90) give an upper limit that is 10 times lower. This large discrepancy implies either a defect in the calculations or the presence of a faster path than the standard two-neutrino mode in this case." See BOEHM 87 and STAUDT 90.
- 139 WONG 91 replaced by VUILLEUMIER 93.
- 140 MILEY 90 claims only "suggestive evidence" for the decay. Error is 2σ .
- 141 VASENKO 90 limit based on background statistics. Maximum likelihood solution is >2000 .
- 142 MORALES 88 notes a 2.5 sigma coincidence rate between electrons with energy 1483.7 ± 0.5 keV in the Ge detector and photons with energy 558 ± 15 keV in the NaI detector, close to the region where neutrinoless $0^+ \rightarrow 2^+$ ^{76}Ge decay should be expected. However, a further study reported in in MORALES 91 rejects this peak at the 95% CL.
- 143 BELLOTTI 87 searches for γ rays for 2^+ state decays in corresponding Xe isotopes. Limit for ^{130}Te case argues for dominant $0^+ \rightarrow 0^+$ transition in known decay of this isotope.
- 144 ELLIOTT 86 limit agrees with the geochemical limit and strongly disagrees with nuclear theory calculations, casting doubt on their application to derive limits on Majorana neutrino masses and η parameters from limits on neutrinoless double β decay.
- 145 HUBERT 85 gives lifetime limits on neutrinoless double β decay of ^{76}Ge to excited states of ^{76}Se .
- 146 KIRSTEN 83 reports "2 σ " error. References are given to earlier determinations of the ^{130}Te lifetime.

(m_ν) , The Effective Weighted Sum of Majorana Neutrino Masses Contributing to Neutrinoless Double β Decay

$(m_\nu) = |\sum U_{ij}^2 m_{\nu_j}|$, where the sum goes from 1 to n and where n = number of neutrino generations, and ν_j is a Majorana neutrino. Note that U_{ij}^2 , not $|U_{ij}|^2$, values in the sum. The possibility of cancellations has been stressed.

VALUE (eV)	CL%	ISOTOPE	TRANSITION	METHOD	DOCUMENT ID
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
< 6.6	68	^{100}Mo	0ν	$0^+ \rightarrow 0^+$ Si(Li)	147 ALSTON... 93
< 2.8-4.3	90	^{136}Xe	0ν	$0^+ \rightarrow 0^+$ TPC	148 VUILLEUMIER 93
< 1.5	90	^{76}Ge		Enriched HPGe	149 BALTHYSH 92
< 1.1-1.5		^{128}Te		Geochem	150 BERNATOW... 92
< 5	68	^{82}Se		TPC	151 ELLIOTT 92
< 1.9-6.7	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ HPGe	152 REUSSER 92
< 11-30	95	^{136}Xe	0ν	$0^+ \rightarrow 0^+$ Prop ctnr	153 BELLOTTI 91
< 3.3-5.0		^{136}Xe	0ν	$0^+ \rightarrow 0^+$ TPC	154 WONG 91
< 8.3	76	^{48}Ca	0ν	CaF_2 scint.	YOU 91
< 1.4-8	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ Enriched Ge(Li)	155 VASENKO 90
< 4.3-28		^{136}Xe	0ν	$0^+ \rightarrow 0^+$ Prop chamber	156 BELLOTTI 89
< 12	68	^{116}Cd	0ν	$^{116}\text{CdWO}_4$ scint	157 DANEVICH 89
< 3.4-23		^{76}Ge	0ν	$0^+ \rightarrow 0^+$ HPGe	158 FISHER 89
< 1.8		^{76}Ge	0ν	$0^+ \rightarrow 0^+$ HPGe	159 CALDWELL 87
< 2.7	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ HPGe	BELLOTTI 86
< 6	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ HPGe	160 CALDWELL 86
< 6	68	^{76}Ge	0ν	HPGe	161 CALDWELL 85
< 20	68	^{76}Ge	0ν	Ge(Li)	162 HUBERT 85
< 3.8	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ Ge(Li)	163 BELLOTTI 84
< 22		^{76}Ge	0ν	$0^+ \rightarrow 0^+$ Ge	FORSTER 84
< 10	90	^{76}Ge	0ν	Intrinsic Ge	AVIGNONE 83
< 22	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ Ge(Li)	164 BELLOTTI 83
< 8.3	68	^{76}Ge	0ν	$0^+ \rightarrow 0^+$ Ge(Li)	164 BELLOTTI 83
< 5.6	95	^{128}Te		Geochem	KIRSTEN 83

- 160 CALDWELL 86 gives several limits depending on which calculation of nuclear matrix elements is used; we quote the most conservative, i.e., least stringent. Other limits are 1.0 eV and 1.9 eV. Authors note that the overall uncertainty due to the serious disagreement between nuclear calculations and both lab and geochemical measurements for regular 2-neutrino double β decay is also present in these limits.
- 161 CALDWELL 85 uses results of HAXTON 81, HAXTON 82. Authors state that limit could be "two or three times larger."
- 162 HUBERT 85 limit is obtained from analysis of data using theoretical calculations by HAXTON 81, HAXTON 82.
- 163 See Table 1 of BELLOTTI 84 for their assessment of previous bounds.
- 164 BELLOTTI 83 limits are obtained from analysis of data using theoretical calculations by DOI 83 and ROSEN 81.

Limits on Lepton-Number Violating (V+A) Current Admixture

For reasons given in the discussion at the beginning of this section, we list only results from 1989 and later.

$\eta_{RR} (10^{-6})$ CL%	$\eta_{LR} (10^{-6})$ CL%	ISOTOPE	METHOD	DOCUMENT ID		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●						
< 2.3	90	< 1.5	90	^{76}Ge	Enriched HPGe	165 BALTHYSH 92
		< 5.3		^{128}Te	Geochem	166 BERNATOW... 92
< 3.6	68	< 2.2	68	^{76}Ge	HPGe	167 REUSSER 92
< 9	68	< 8	68	^{76}Ge	Ion chamber	BELLOTTI 89

- 165 BALTHYSH 92 uses the MUTO 89 matrix elements.
- 166 BERNATOWICZ 92 takes the measured geochemical decay width as a limit on the 0ν width, and uses the SUHONEN 91 coefficients to obtain the least restrictive limit on η . Further details of the experiment are given in BERNATOWICZ 93.
- 167 REUSSER 92 uses the MUTO 89 matrix elements for this reduction.

REFERENCES FOR Searches for Massive Neutrinos and Lepton Mixing

ANSELMANN 93	PL B327 377	+Hampel, Heusser, Kiko, Kirsten+	(GALLEX Collab.)
ABELE 94	PL B316 26	+Helm, Kania, Schmidt+	(HEID, ILLG, MUNT)
ALSTON... 93	PRL 71 831	Aiston-Garnjost+	(LBL, MTHO, UNM, INEL)
ANSELMANN 93	PL B314 445	+Hampel, Heusser, Kiko, Kirsten+	(GALLEX Collab.)
BAHCALL 93	PR D47 1298	+Bethé	(IAS, CORN)
BAHRAN 93	PR D47 R754	+Kaltfleisch	(OKLA)
BAHRAN 93B	PR D47 R759	+Kaltfleisch	(OKLA)
BARANOV 93	PL B302 336	+Batusov, Bunyatov, Klimov+	(JINR, SERP, BUDA)
BERMAN 93	PR C48 R1	+Pitt, Calaprice, Lowry	(PRIN)
BERNATOW... 93	PR C47 806	Bernatowicz, Brazzle, Cowsik+	(WUSL, TATA)
DIGREGORIO 93	PR C47 2916	+Gil, Huck, Batista+	(CNEA)
FREEDMAN 93	PR D47 811	+Fujikawa, Napolitano, Nelson+	(LAMPF E645 Collab.)
GRUWE 93	PL B309 463	+Mommata, Vilain, Wilquet+	(CHARM II Collab.)
HAMPPEL 93	IPG 118 2209		(MPIH)
HIME 93	PL B299 165		(LANL)
KALBFLEISCH 93	PL B303 355	+Bahrain	(OKLA)
MORTARA 93	PRL 70 394	+Ahmad, Coulter, Freedman+	(ANL, LBL, UCB)
OHSHIMA 93	PR D47 4840	+ (KEK, TUAT, RIKEN, SCUC, ROCH, TSUK, INUS)	
TURCK-CHIL... 93	PRPL 230 57	Turck-Chieze+	(SACL, USC, NICEA, NICEO, MEUD)
TURCK-CHIL... 93B	APJ 408 347	Turck-Chieze, Lopez	(SACL)
VUILLEUMIER 93	PL B294 479	+Busto, Farine, Jogens+	(NEUC, CIT, VILL)
WIETFIELDT 93	PRL 70 1759	+Chavez de Cruz, Garcia+	(LBL, UCB, SPAUL)
ALESSAND... 92	PL B285 176	Alessandrello, Camin, Cremonesi, Fiorini+	(MILA, SASSO)
ANSELMANN 92	PL B285 376	+Hampel, Heusser+	(GALLEX Collab.)
ANSELMANN 92B	PL B285 390	+Hampel, Heusser+	(GALLEX Collab.)
ARTEMJEV 92	PL B280 159	+Brachmann, Ivanovsky, Karelin+	(ITEP)
BAHCALL 92	RMP 64 885	+Pinsonneault	(IAS, YALE)
BAHRAN 92	PL B291 336	+Kaltfleisch	(OKLA)
BAHRAN 92B	PL B294 479 (erratum)	+Beyaz, Bockholt, Demehin+	(MPIH, KIAE, SASSO)
BALTHYSH 92	PL B283 32	Becker-Szendy, Bratton, Casper, Dye+	(IMB Collab.)
BECKER-SZ... 92	PRL 69 1010	Becker-Szendy, Bratton, Casper, Dye+	(IMB Collab.)
BECKER-SZ... 92B	PR D46 3720	Becker-Szendy, Bratton, Casper, Dye+	(IMB Collab.)
BEIER 92	PL B283 446	+Frank, Frati, Kim, Mann+	(KAM2 Collab.)
Also 94	PTSL A346 63	Beier, Frank	(PENN)
BERNATOW... 92	PRL 69 2341	Bernatowicz, Brannon, Brazzle, Cowsik+	(WUSL, TATA)
BLUM 92	PL B275 506	+Busto, Campagne, Dassie, Hubert+	(NEMO Collab.)
BORODOV... 92	PL B280 146	+Borodovsky, Chi, Ho, Kondakis, Lee+	(COLU, JHU, ILL)
BRITTON 92	PRL 68 3000	+Ahmad, Bryman, Burnham+	(TRIU, CARL)
Also 94	PR D49 28	Britton, Ahmad, Bryman+	(TRIU, CARL)
BRITTON 92B	PR D46 R885	+Ahmad, Bryman+	(TRIU, CARL)
CHEN 92	PRL 69 3151	+Imel, Radcliffe, Henrickson, Boehm	(CIT)
ELLIOTT 92	PR C46 1535	+Hahn, Moe+	(PRIN)
GARVIN 92	HEP-92 Conf., p. 1101 +Ansolov+ (INRM, LANL, PENN, LSU, UIC)		
Proc. XXV Int. Proc. Conf., Dallas, TX (1992) AIP Conf. Proc. No. 272, ed. J.R. Sanford, III.			
HIRATA 92	PL B280 146	+Inoue, Ishida+	(Kamiokande II Collab.)
HOLZSCHUH 92B	PL B287 381	+Fritschl, Kuendig	(ZURI)
KAWAKAMI 92	PL B287 45	+ (INUS, KEK, SCUC, TUAT, RIKEN, ROCH, TSUK)	
KETOV 92	JETPL 55 564	+Machulin, Mikaelyan+	(KIAE)
Translated from ZETFP 55 54.			
KUDOMI 92	PR C46 R2132	+Ejiri, Nagata, Okada, Shibata+	(OSAK, INUS)
REUSSER 92	PR D45 2548	Reusser, Trichel, Boehm+	(CIT, NEUC, VILL)
ABAZOV 91B	PRL 67 3332	+Anosov, Faizov+	(SAGE Collab.)
AVIGNONE 91	PL B256 559	+Brodzinski, Guerdard+	(SCUC, PNL, ITEP, VILL)
BELLOTTI 91	PL B266 193	+Cremonesi, Fiorini, Gervasio+	(MILA, INFN)
CASPER 91	PRL 66 2561	+Becker-Szendy, Bratton, Cady+	(IMB Collab.)
DELEENER... 91	PR D43 3611	+De Leener-Rosier, Deutsch+	(LVLN, ZURI, LAUS)
EJIRI 91	PL B258 17	+Fushimi, Kamada, Kinoshita+	(OSAK)
GARCIA 91	PRL 67 3654	+Adeberger, Magnus, Swanson+	(WASH, CERN, LBL)
HIME 91	PL B257 441	+Jelley	(OXF)
HIME 91B	Thesis OUNP-91-21	+Jelley	(OXF)
HIRATA 91	PRL 66 9	+Inoue, Kajita, Kihara+	(Kamiokande II Collab.)
HIRATA 91B	PR D44 2241		
HYKAWY 91	PRL 67 1708	+Nxumalo, Unger, Lander+	(MANI)
MORALES 91	NC 104A 1581	+Morales, Nunez-Lagos, Puimedon+	(ZARA)
NORMAN 91	PL B251 321	+Suz, Lesko+	(LBL)
SUHONEN 91	NP A535 509	+Khakikar, Faessler	(JYV, AHMED, TUBIN)
SUR 91	PRL 66 2444	+Norman, Lesko+	(LBL)
TURKEVICH 91	PRL 67 3211	+Economou, Cowan	(CHIC, LANL)
WONG 91	PRL 67 1218	+Boehm, Fisher, Gabathuler+	(CIT, VILL, NEUC)
YOU 91	PL B265 53	+Zhu, Lu+	(BHEP, CAST+)
ZLIMEN 91	PRL 67 560	+Ljubicic, Kaucic, Logan	(BOSK, OTTA)
AEVIA 90S	PL B251 321	+Adrian, Aguilari-Benitez, Akbari+	(L3 Collab.)
ASTIER 90	NP B335 517	+Bernardi+	(BOST, BNL, CERN, CURIN, PARIN)
BARABASH 90	PL B249 186	+Kopylov, Cherehovsky	(ITEP, INRM)
BATUSOV 90B	ZPHY C48 209	+Bunyatov, Kuznetsov, Pozharova+	(JINR, ITEP, SERP)
BERGER 90B	PL B245 305	+Froehlich, Moench, Nisius+	(FREJUS Collab.)
BURCHART 90	PR D41 3542	+King, Abrams, Adolphsen+	(Mark II Collab.)
DECAMP 90F	PL B236 511	+Deschizeaux, Lees, Minard+	(ALEPH Collab.)
FILIPPONE 90	PL B246 546	+Vogel	(CIT)
HIRATA 90	PRL 65 1297	+Inoue, Kajita+	(Kamiokande II Collab.)
HIRATA 90B	PRL 65 1301	+Inoue, Kajita+	(Kamiokande II Collab.)
JUNG 90	PRL 64 1091	+Van Kooten, Abrams, Adolphsen+	(Mark II Collab.)
KOPEKIN 90	JETPL 51 86	+Mikaziyian, Fayans	(KIAE)
Translated from ZETFP 51 75.			

Lepton & Quark Full Listings

Massive Neutrinos and Lepton Mixing, Neutrino Bounds from Astrophysics and Cosmology

MILEY	90	PRL 65 3092	+Avignone, Brodzinski, Collar, Reeves (SCUC, PNL)
STAUDT	90	EPL 13 31	+Muto, Kiapdor-Kleingrothaus (MPIH)
VASENKO	90	MPL A5 1299	+Kirpichnikov, Kuznetsov, Starostin (ITEP, YERE)
VIDYAKIN	90	JETP 71 424	+Vyrodov, Gurevich, Koslov+ (KIAE)
		Translated from ZETP 98 764.	
ABRAMS	89C	PRL 63 2447	+Adolphsen, Averill, Ballam+ (Mark II Collab.)
AGLIETTA	89	EPL 9 611	+Battistoni, Bellotti+ (FREJUS Collab.)
ALSTON...	89	PRL 63 1671	+Alston-Garnjost, Dougherty+ (LBL, MTHO, UNM, INEL)
ASTIER	89	PL B220 646	+Bernardi, Carugno+ (CURIN, PARIN, BOST, CERN, BNL)
BAHCALL	89	Neutrino Astrophysics, Cambridge Univ. Press	(IAS)
BAHCALL	89B	PR D40 931	+Haxton (IAS, WASH)
BARABASH	89	PL B223 273	+Kuzminov, Lobashev, Novikov+ (ITEP, INRM)
BELLOTTI	89	PL B221 209	+Cremonesi, Fiorini, Gervasio+ (MILA)
BLUMENFELD	89	PRL 62 2237	+Chi, Chichura, Chien+ (COLU, ILL, JHU)
DAMEVICH	89	JETPL 49 476	+Zdesenko, Nikolaiko, Tretyak (UZJNR)
		Translated from ZETFP 49 417.	
DAVIS	89	ARNPS 39 467	+Mann, Wolfenstein (BNL, PENN, CMU)
DRUKAREV	89	SJNP 50 184	+Strikman (PNPI)
		Translated from YAF 50 294.	
FISHER	89	PL B218 257	+Boehm, Bovet, Egger+ (CIT, NEUC, PSI)
HIME	89	PR D39 1837	+Simpson (GUEL)
LOGAN	89	NIM A280 167	+Haxton (OTTA)
MUTO	89	ZPHY A334 187	+Bender, Kiapdor (TJNT, MPIH)
OYAMA	89	PR D39 1481	+Hirata, Kajita, Kifune+ (Kamiokande II Collab.)
SIMPSON	89	PR D39 1825	+Hime (GUEL)
AFONIN	88	JETP 67 213	+Ketov, Kopeikin, Mikaelyan+ (KIAE)
		Translated from ZETFP 94 294.	
AKERLOF	88	PR D37 577	+Chapman, Errede, Ken+ (HRS Collab.)
AMMOSOV	88	ZPHY C40 487	+Belikov+ (SKAT Collab.)
BAHCALL	88	RMP 60 297	+Ulrich (IAS, UCLA)
BERGSM	88	ZPHY C40 171	+Dorenbosch, Nieuwenhuis+ (CHARM Collab.)
BERNARDI	88	PL B203 332	+Carugno, Chauveau+ (PARIN, CERN, INFN, ATEN)
BIONTA	88	PR D38 768	+Blewitt, Bratton, Casper+ (IMB Collab.)
DURKIN	88	PL B1 1813	+Hager, Ling+ (OSU, ANL, CIT, LBL, LSU, LANL)
ENGEL	88	PR C37 731	+Vogel, Zimbene
HIRATA	88	PL B205 416	+Kajita, Koshiba+ (Kamiokande II Collab.)
LOVERRE	88	PL B206 711	(INFN)
MORALES	88	NC 100A 525	+Morales, Nunez-Lagos, Puimedon+ (ZARA)
TURCK-CHI...	88	APJ 335 415	Turck-Chieze, Cahen, Casse, Doom (SACL, BRUX)
ZLIMEN	88	PS 38 539	+Kaucic, Ljubicic, Logan (BOSK, CARL)
AFONIN	87	JETPL 45 257	+Bogatov, Vershinski+ (KIAE)
		Translated from ZETFP 45 201.	
AHRENS	87	PR D36 702	+ (BNL, BROW, UCI, HIRO, KEK, OSAK, PENN, STON)
BELLOTTI	87	EPL 3 889	+Cattadori, Cremonesi, Fiorini+ (MILA)
BOEHM	87	Massive Neutrinos Cambridge Univ. Press, Cambridge	+Vogel (CIT)
BOFILL	87	PR D36 3309	+Busza, Eldridge+ (MIT, FNAL, MSU)
CALDWELL	87	PR D36 419	+Eisberg, Grumm, Witherell+ (UCSB, LBL)
DALIM	87	PR D36 2624	+Kettle, Jost+ (SIN, VIRG)
ELLIOTT	87B	PRL 59 2020	+Hahn, Moe (UCI)
HETHERING...	87	PR C36 1504	+Hetherington, Graham+ (UCI)
LOSECCO	87	PL B184 305	+Bionta, Blewitt, Bratton+ (IMB Collab.)
MISHRA	87	PRL 59 1397	+Auchincloss+ (COLU, CIT, FNAL, CHIC, ROCH)
OBERAUER	87	PL B198 113	+von Feilitzsch, Mossbauer (MUNT)
TALIBZADEH	87	NP B291 503	+Guy, Jenus+ (BEBC WA66 Collab.)
TOMODA	87	PL B199 475	+Fassler, (TUBIN)
VIDYAKIN	87	JETP 66 243	+Vyrodov, Gurevich, Kozlov+ (KIAE)
		Translated from ZETFP 93 424.	
WENDT	87	PRL 58 1810	+Abrams, Amidei, Baden+ (Mark II Collab.)
AFONIN	86	JETPL 44 142	+Bogatov, Borovoi, Vershinski+ (KIAE)
		Translated from ZETFP 44 111.	
ANGELINI	86	PL B179 307	+Apostolakis, Baldini+ (PISA, ATHU, PAOO, WISC)
AZUELOS	86	PRL 56 2241	+Britton, Bryman+ (TRI, CNRC)
BADIER	86	ZPHY C31 21	+Bemporad, Boucrot, Callot+ (NA3 Collab.)
BELLOTTI	86	NC 95A 1	+Cremonesi, Fiorini, Liguori+ (MILA)
BERNARDI	86	PL 166B 479	+Carugno+ (CURIN, INFN, CDEF, ATEN, CERN)
BERNARDI	86B	PL B181 173	+Carugno+ (CURIN, INFN, CDEF, ATEN, CERN)
BORGE	86	PS 34 591	+DeRujula, Hansen, Johnson+ (ISOLDE Collab.)
BRUCKER	86	PR D34 2183	+Jacques, Kalikar, Koller+ (RUTG, BNL, COLU)
CALDWELL	86	PR D33 2737	+Eisberg, Grumm, Hale, Witherell+ (UCSB, LBL)
DELEENER...	86	PL B177 228	+DeLeener-Rosier, Deutsch+ (LVLN, ZURI, LAUS)
DORENBOS...	86	PL 166B 473	+Dorenbosch, Allaby, Amaldi+ (CHARM Collab.)
DRUKAREV	86	JETP 64 686	+Strikman (PNPI)
		Translated from ZETFP 91 1160.	
ELLIOTT	86	PRL 56 2582	+Hahn, Moe (UCI)
EMAN	86	PR C33 2128	+Tadic (BOSK)
GROTZ	86	NC C9 535	+Kiapdor (MPIH)
LINDHARD	86	PRL 57 965	+Hansen (AARH)
SIMPSON	86	PL B174 113	+Hansen (GUEL)
USHIDA	86C	PRL 57 2897	+Kondo, Tasaka, Park, Song+ (FNAL E531 Collab.)
ZACEK	86	PR D34 2621	+Feilitzsch+ (CIT-SIN-TUM Collab.)
AFONIN	85	JETPL 41 435	+Borovoi, Dobrynin+ (KIAE)
		Translated from ZETFP 41 355.	
Also	85B	JETPL 42 285	Afonin, Bogatov, Borovoi, Dobrynin+ (KIAE)
		Translated from ZETFP 42 230.	
AHRENS	85	PR D31 2732	+Aronson+ (BNL, BROW, KEK, OSAK, PENN+)
ALBRECHT	85	PL B33B 404	+Binder, Drescher, Schubert+ (ARGUS Collab.)
ALTITZOG...	85	PRL 55 799	+Altitzoglou, Calaprice, Dewey+ (TRI, CNRC)
APALIKOV	85	JETPL 42 289	+Boris, Golutvin, Laptin, Lubimov+ (ITEP)
		Translated from ZETFP 42 233.	
BELIKOV	85	SJNP 41 589	+Volkov, Kochetkov, Mukhin+ (SERP)
		Translated from YAF 41 919.	
CALDWELL	85	PRL 54 281	+Eisberg, Grumm, Hale, Witherell+ (UCSB, LBL)
COOPER...	85	PL 160B 207	+Cooper-Sarkar+ (CERN, LOIC, OXF, SACL+)
DATAR	85	Nature 318 547	+Baba, Bhattacharjee, Bhuinaya, Roy (BHAB, TATA)
HAXTON	85	PRL 55 807	+Haxton (WASH, LASL)
HUBERT	85	NC 85A 19	+Leccia, Dassie, Mennrath+ (BCEN, ZARA)
KALBFLEISCH	85	PRL 55 2225	+Milton (OKLA)
MARKY	85	PR C32 2215	+Boehm (CIT)
OH	85	PL 160B 322	+Nakajima, Tamura+ (TOKY, INUS, KEK)
SIMPSON	85	PRL 54 1891	+Hansen (GUEL)
STOCKDALE	85	ZPHY C27 53	+Bodek+ (ROCH, CHIC, COLU, FNAL)
ZACEK	85	PL 164B 193	+Zacek, Boehm+ (MUNI, CIT, SIN)
BALLAGH	84	PR D30 2271	+Bingham+ (UCB, LBL, FNAL, HAWA, WASH, WISC)
BELLOTTI	84	PL 146B 450	+Cremonesi, Fiorini, Liguori, Pullia+ (MILA)
BERGSM	84	PL 142B 103	+Dorenbosch, Allaby, Abt+ (CHARM Collab.)
CAVAIGNAC	84	PL 146B 387	+Hounmada, Koang+ (ISNG, LAPP)
DYDAK	84	PL 134B 281	+Feldman+ (CERN, DORT, HEIDH, SACL, WARS)
FORSTER	84	PL 138B 301	+Kwon, Markey, Boehm, Henrikson (CIT)
GABATHULER	84	PL 138B 449	+Boehm+ (CIT, SIN, MUNI)
HAXTON	84	PPNP 12 409	+Stevenson
MINEHART	84	PRL 52 804	+Zock, Marshall, Stephens, Daum+ (VIRG, SIN)
STOCKDALE	84	PRL 52 1384	+Bodek+ (ROCH, CHIC, COLU, FNAL)
AFONIN	83	JETPL 38 436	+Bogatov, Borovoi, Vershinski+ (KIAE)
		Translated from ZETFP 38 361.	
AVIGNONE	83	PRL 50 721	+Brodzinski, Brown, Evans, Hensley+ (SCUC, PNL)
BELENKII	83	JETPL 38 493	+Dobrynin, Zemlyakov, Mikaelyan+ (KIAE)
		Translated from ZETFP 38 406.	
BELIKOV	83	JETPL 38 661	+Volkov, Kochetkov, Mukhin, Sviridov+ (SERP)
		Translated from ZETFP 38 547.	
BELLOTTI	83	PL 121B 72	+Fiorini, Liguori, Pullia, Sarracino+ (MILA)
BERGSM	83	PL 122B 465	+Dorenbosch, Jonker+ (CHARM Collab.)
BERGSM	83B	PL 128B 361	+Dorenbosch+ (CHARM Collab.)
BRYMAN	83B	PRL 50 1546	+Dubois, Numao, Olaniya, Olin+ (TRI, CNRC)
Also	83	PRL 50 7	Bryman, Dubois, Numao, Olaniya+ (TRI, CNRC)
DEUTSCH	83	PR D27 1644	+Lebrun, Priels (LVLN)
DOI	83	PTP 69 602	+Kotani, Nishiura, Takasugi (OSAK, KYOT)
FILIPPONE	83	PRL 50 412	+Elwyn, Davids+ (ANL, CHIC, VALP)
GROUAT	83	PR D28 2762	(HAF)
KIRSTEN	83	PRL 50 474	+Richter, Jessberger (MPIH)
Also	83B	ZPHY 16 189	Kirsten, Richter, Jessberger (MPIH)
SCHRECK...	83	PL 129B 265	Schreckenbach, Colvin+ (ISNG, ILLG)
TAYLOR	83	PR D28 2705	+Cence, Harris, Jones+ (HAWA, LBL, FNAL)
BAHCALL	82	RMP 54 767	+Huebner, Lubow+ (IAS, LANL, HPC, YALE, UCLA)
COOPER	82	PL 112B 97	+Guy, Michette, Tyndel, Venus (RL)
FILIPPONE	82	APJ 253 393	+Schramm (ANL, CIT)
FOWLER	82	A.P. 95 280	(CIT)
HAXTON	82	PR D25 2360	+Stephenson, Strottman (LANL, PURD)
HAYANO	82	PRL 49 1305	+Taniguchi, Yamanka+ (TOKY, KEK, TSUK)
VUILLEUMIER	82	PL 114B 298	+Boehm, Egger+ (CIT, SIN, MUNI)
ABELA	81	PL 105B 263	+Daum, Eaton, Frosch, Jost, Kettle, Steiner (SIN)
ARMENISE	81	PL 100B 182	+Fogli-Muciaccia+ (BARI, CERN, MILA, LALO)
ASANO	81	PL 104B 81	+Hayano, Kikutani, Kurokawa+ (KEK, TOKY, INUS, OSAK)
Also	81	PL 102A 1232	Shrock (STON)
ASRATYAN	81	PL 105B 301	+Efremenko, Fedotov+ (ITEP, FNAL, SERP, MICH)
BAKER	81	PRL 47 1576	+Connolly, Kahn, Kirk, Murtugh+ (BNL, COLU)
Also	78	PRL 40 144	Cnops, Connolly, Kahn, Kirk+ (BNL, COLU)
BOLIEV	81	SJNP 34 787	+Butkevich, Zakidyshev, Makojev+ (INRM)
		Translated from YAF 34 1418.	
CALAPRICE	81	PL 106B 175	+Schreiber, Schneider+ (PRIN, IND)
DEDEN	81	PL 98B 310	+Grassler, Boeckmann, Mermikides+ (BEBC Collab.)
ERRIQUEZ	81	PL 102B 73	+Natali+ (BARI, BIRM, BRUX, EPOL, RHEL, SACL+)
HAXTON	81	PRL 47 153	+Stephenson, Strottman (PURD, LASL)
KWON	81	PR D24 1097	+Boehm, Hahn, Henrikson+ (CIT, ISNG, MUNI)
NEMETHY	81B	PR D23 262	+ (YALE, LBL, LASL, MIT, SACL, SIN, CNRC, BERN)
ROSEN	81	Nu Conf. Hawaii	(PURD)
Also	78	RMP 50 114	Bryman, Picciotto (TRI, VICT)
SHROCK	81B	PR D24 1232	(STON)
SHROCK	81B	PR D24 1275	(STON)
SILVERMAN	81	PRL 46 467	+Soni (UCI, UCLA)
SIMPSON	81B	PR D24 2971	(GUEL)
USHIDA	81	PRL 47 1694	+ (AICH, FNAL, KOBE, SEOU, MCGI, NAGO, OSU+)
AVIGNONE	80	PR C22 594	+Greenwood (SCUC)
BAHCALL	80	PRL 45 945	+Lubow, Huebner+ (IAS, LASL, YALE, LLL, UCLA)
Also	76	Science 191 264	Bahcall, Davis (IAS, BNL)
BOEHM	80	PL 92B 229	+Cavaignac, Feilitzsch+ (ILLG, CIT, ISNG, MUNI)
FRITZE	80	PL 96B 427	+Reines, Gurr, Sobel (AAACH3, BONN, CERN, LOIC, OXF, SACL)
REINES	80	PRL 45 1307	+Sobel, Pasiber (UCI)
Also	59	PL 113 273	Reines, Cowan (LASL)
Also	66	PR 142 852	Nezrick, Reines (CASE)
Also	76	PRL 37 315	Reines, Gurr, Sobel (UCI)
SHROCK	80	PL 96B 159	(STON)
DAVIS	79	PL 91B 229	+Vogel, Mann, Schenter (Gargamele Collab.)
BLIETSCHAU	78	NP B133 205	+Deden, Hahert, Krenz+ (Gargamele Collab.)
CROUCH	78	PR D18 2239	+Landecker, Lathrop, Reines+ (CASE, UCI, WITW)
BELLOTTI	76	PL 96B 153	+Cavalli, Fiorini, Rollier (MILA)
CONWAY	59	PR 116 1544	+Johnston (PURD)
HAMILTON	58	PR 112 2010	+Langer, Smith (IND)
JOHNSON	58	PR 112 2004	+Johnson, Langer (IND)

Neutrino Bounds from Astrophysics and Cosmology

OMITTED FROM SUMMARY TABLE

The limits on the number of light neutrino types now appears in a separate section (following the τ -lepton section).

See the note on neutrinos by R.E. Shrock in the ν_e section near the beginning of these Listings. For additional information see the ν_e , ν_{μ} , ν_{τ} , and heavy- ν sections above.

NOTE ON CONSTRAINTS ON PARTICLES FROM SN 1987A

(by J. Ellis, CERN and D.N. Schramm, Univ. of Chicago)

Since there have been few new developments in 1992-94, the text of this note is omitted. The reader is referred to the 1992 edition (Phys. Rev. D45, VI. 42 (Part II, June 1992)).

ν MASS

The limits on low mass ($m_{\nu} \lesssim 1$ MeV) neutrinos apply to m_{tot} given by

$$m_{\text{tot}} = \sum_{\nu} (g_{\nu}/2) m_{\nu}$$

where g_{ν} is the number of spin degrees of freedom for ν plus $\bar{\nu}$: $g_{\nu} = 4$ for neutrinos with Dirac masses; $g_{\nu} = 2$ for Majorana neutrinos. The limits on high mass ($m_{\nu} > 1$ MeV) neutrinos apply separately to each neutrino type.

Limit on Total ν MASS, m_{tot}

(Defined in the above note), of effectively stable neutrinos (i.e., those with mean lives greater than or equal to the age of the universe). These papers assumed Dirac neutrinos. When necessary, we have generalized the results reported so they apply to m_{tot} . For other limits, see SZALAY 76, VYSOTSKY 77, BERNSTEIN 81, FREESE 84, SCHRAMM 84, and COWSIK 85.

VALUE (eV)	DOCUMENT ID	TECN
------------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •		
<180	SZALAY	74 COSM
<132	COWSIK	72 COSM
<280	MARX	72 COSM
<400	GERSHTEIN	66 COSM

Astrophysical and Cosmological Limits on ν MASSES

If neutrinos are present as dark matter in galactic halos, limits on neutrino masses have been computed based on neutrino degeneracy and Fermi statistics. The results depend strongly on assumptions. See the references.

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •			
	SPERGEL	88B	COSM
	KAWASAKI	86	COSM
	KAWASAKI	86B	COSM
	TAKAHARA	86	COSM supernovae
	MADSEN	85	COSM Some anisotropy
	MADSEN	84	COSM Assume isotropy
	SARKAR	84	COSM Decaying neutrinos
	FREESE	83	COSM Degenerate ν
	LIN	83	COSM
	PRIMACK	83	COSM
	BOND	81	COSM Adiabatic
	DAVIS	81	COSM Adiabatic+decaying ν 's
	SCHRAMM	81	COSM Isothermal
	TREMAINE	79	COSM Isothermal

Limits on MASSES of Light Stable Right-Handed ν (with necessarily suppressed interaction strengths)

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •			
<100–200	¹ OLIVE	82	COSM Dirac ν
<200–2000	¹ OLIVE	82	COSM Majorana ν

¹ Depending on interaction strength g_R where $g_R < G_F$.

Limits on MASSES of Heavy Stable Right-Handed ν (with necessarily suppressed interaction strengths)

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •			
> 10	² OLIVE	82	COSM $g_R/G_F < 0.1$
>100	² OLIVE	82	COSM $g_R/G_F < 0.01$

² These results apply to heavy Majorana neutrinos and are summarized by the equation: $m_\nu > 1.2 \text{ GeV } (G_F/g_R)$.

REFERENCES FOR Neutrino Bounds from Astrophysics and Cosmology

SPERGEL	88B	PR D38 2014	+Weinberg, Gott	(PRIN)
KAWASAKI	86	PL B178 71	+Terasawa, Sato	(TOKY)
KAWASAKI	86B	PL 169B 280	+Sato	(TOKY)
TAKAHARA	86	PL B174 373	+Sato	(TOKY)
COWSIK	85	PL 151B 62		(TATA)
MADSEN	85	PRL 54 2720	+Epstein	(AARH, LANL)
FREESE	84	NP B233 167	+Schramm	(CHIC, FNAL)
MADSEN	84	APJ 282 11	+Epstein	(AARH, LANL)
SARKAR	84	PL 148B 347	+Cooper	(OXF, CERN)
SCHRAMM	84	PL 141B 357	+Steigman	(FNAL, BART)
FREESE	83	PR D27 1689	+Kolb, Turner	(CHIC, LANL)
LIN	83	APJ 266 L21	+Faber	(UCSC)
PRIMACK	83	Phil. 4th Workshop on Grand Unification		(UCSC)
Also	82	Nature 299 37	Blumenthal, Pagels, Primack	(UCSC, ROCK)
OLIVE	82	PR D25 213	+Turner	(CHIC, UCSB)
BERNSTEIN	81	PL 101B 39	+Feinberg	(STEV, COLU)
BOND	81	Nu Conf. Hawaii	+Szalay	(UCB, CHIC)
DAVIS	81	APJ 250 423	+Lecar, Pryor, Witten	(HARV, PRIN)
SCHRAMM	81	APJ 243 1	+Steigman	(CHIC, BART)
TREMAINE	79	PRL 42 407	+Gunn	(CIT, CAMB, CAIW)
VYSOTSKY	77	JETPL 26 188	+Dolgov, Zeldovich	(ITEP)
		Translated from ZETFP 26 200.		
SZALAY	76	AA 49 437	+Marx	(EOTV)
SZALAY	74	APAH 35 8	+Marx	(EOTV)
COWSIK	72	PRL 29 569	+McClelland	(UCB)
MARX	72	Nu Conf. Budapest	+Szalay	(EOTV)
GERSHTEIN	66	JETPL 4 120	+Zeldovich	(RIAM)
		Translated from ZETFP 4 189.		

QUARKS**NOTE ON QUARK MASSES****A. Introduction**

This note discusses some of the theoretical issues involved in the determination of quark masses. Unlike the leptons, quarks are confined inside hadrons and are not observed as physical particles. Quark masses cannot be measured directly, but must be determined indirectly through their influence on hadron properties. As a result, the values of the quark masses depend on precisely how they are defined; there is no one definition that is the obvious choice. Though one often speaks loosely of quark masses as one would of the electron or muon mass, any careful statement of a quark mass value must make reference to a particular computational scheme that is used to extract the mass from observations. It is important to keep this scheme dependence in mind when using the quark mass values tabulated in the data listings.

The simplest way to define the mass of a quark is by making a fit of the hadron mass spectrum to a nonrelativistic quark model. The quark masses are defined as the values obtained from the fit. The resulting masses only make sense in the limited context of a particular quark model. They depend on the phenomenological potential used, and on how relativistic effects are modelled. The quark masses used in potential models also cannot be connected with the quark mass parameters in the QCD Lagrangian. Fortunately, there exist other definitions of the quark mass that have a more general significance, though they also depend on the method of calculation. The purpose of this review is to explain the most important such definitions and their interrelations.

B. Mass parameters and the QCD Lagrangian

The QCD Lagrangian for N_F quark flavors is

$$\mathcal{L} = \sum_{k=1}^{N_F} \bar{q}_k (i \not{D} - m_k) q_k - \frac{1}{4} G_{\mu\nu} G^{\mu\nu}, \quad (1)$$

where $\not{D} = (\partial_\mu - igA_\mu) \gamma^\mu$ is the gauge covariant derivative, A_μ is the gluon field, $G_{\mu\nu}$ is the gluon field strength, m_k is the mass parameter of the k^{th} quark, and q_k is the quark Dirac field. The QCD Lagrangian Eq. (1) gives finite scattering amplitudes after renormalization, a procedure that invokes a subtraction scheme to render the amplitudes finite, and requires the introduction of a dimensionful scale parameter μ . The mass parameters in the QCD Lagrangian Eq. (1) depend on the renormalization scheme used to define the theory, and also on the scale parameter μ . The most commonly used renormalization scheme for QCD perturbation theory is the $\overline{\text{MS}}$ scheme.

The QCD Lagrangian has a chiral symmetry in the limit that the quark masses vanish. This symmetry is spontaneously broken by dynamical chiral symmetry breaking, and explicitly broken by the quark masses. The nonperturbative scale of

Lepton & Quark Full Listings

Quarks

dynamical chiral symmetry breaking, Λ_χ , is around 1 GeV. It is conventional to call quarks heavy if $m > \Lambda_\chi$, so that explicit chiral symmetry breaking dominates, and light if $m < \Lambda_\chi$, so that spontaneous chiral symmetry breaking dominates. The c , b , and t quarks are heavy, and the u , d and s quarks are light. The computations for light quarks involve an expansion in m_q/Λ_χ about the limit $m_q = 0$, whereas for heavy quarks, they involve an expansion in Λ_χ/m_q about $m_q = \infty$. The corrections are largest for the s and c quarks, which are the heaviest light quark and the lightest heavy quark, respectively.

At high energies or short distances, nonperturbative effects such as chiral symmetry breaking are unimportant, and one can in principle analyze mass-dependent effects using QCD perturbation theory to extract the quark mass values. The QCD computations are conventionally performed using the $\overline{\text{MS}}$ scheme at a scale $\mu \gg \Lambda_\chi$, and give the $\overline{\text{MS}}$ “running” mass $\overline{m}(\mu)$. The μ dependence of $\overline{m}(\mu)$ at short distances can be calculated using the renormalization group equations.

For heavy quarks, one can obtain useful information on the quark masses by studying the spectrum and decays of hadrons containing heavy quarks. One method of calculation uses the heavy quark effective theory (HQET), which defines a HQET quark mass m_Q . Other commonly used definitions of heavy quark masses such as the pole mass are discussed in Sec. C. QCD perturbation theory at the heavy quark scale $\mu = m_Q$ can be used to relate the various heavy quark masses to the $\overline{\text{MS}}$ mass $\overline{m}(\mu)$, and to each other.

For light quarks, one can obtain useful information on the quark mass ratios by studying the properties of the light pseudoscalar mesons using chiral perturbation theory, which utilizes the symmetries of the QCD Lagrangian Eq. (1). The quark mass ratios determined using chiral perturbation theory are those in a subtraction scheme that is independent of the quark masses themselves, such as the $\overline{\text{MS}}$ scheme.

A more detailed discussion of the masses for heavy and light quarks is given in the next two sections. The $\overline{\text{MS}}$ scheme applies to both heavy and light quarks. It is also commonly used for predictions of quark masses in unified theories, and for computing radiative corrections in the Standard Model. For this reason, we use the $\overline{\text{MS}}$ scheme as the standard scheme in reporting quark masses. One can easily convert the $\overline{\text{MS}}$ masses into other schemes using the formulæ given in this review.

C. Heavy quarks

The commonly used definitions of the quark mass for heavy quarks are the pole mass, the $\overline{\text{MS}}$ mass, the Georgi-Politzer mass, the potential model mass used in ψ and Υ spectroscopy, and the HQET mass.

The strong interaction coupling constant at the heavy quark scale is small, and one can compute the heavy quark propagator using QCD perturbation theory. For an observable particle such as the electron, the position of the pole in the propagator is the definition of the particle mass. In QCD this definition of the quark mass is known as the pole mass m_P , and is

independent of the renormalization scheme used. It is known that the on-shell quark propagator has no infrared divergences up to (at least) two-loop order [1], so this provides a definition of the quark mass up to order $\alpha_s(m_P)^2$. The pole mass cannot be used to arbitrarily high accuracy because of nonperturbative infrared effects in QCD. The full quark propagator has no pole because the quarks are confined, so that the pole mass cannot be defined outside of perturbation theory.

The $\overline{\text{MS}}$ running mass $\overline{m}(\mu)$ is defined by regulating the QCD theory using dimensional regularization, and subtracting the divergences using the modified minimal subtraction scheme. The $\overline{\text{MS}}$ scheme is particularly convenient for Feynman diagram computations, and is the most commonly used subtraction scheme.

The Georgi-Politzer mass $\widehat{m}(\xi)$ is defined using the momentum space subtraction scheme at the spacelike point $-p^2 = M^2$ [2], with $M^2 = (\xi + 1)m_P^2$. It is often used in computations involving QCD sum rules [3], which can be used to extract heavy quark masses. QCD sum rules are discussed in more detail in the next section on light quark masses.

Lattice gauge theory calculations can be used to obtain heavy quark masses from ψ and Υ spectroscopy. The quark masses are obtained by comparing a nonperturbative computation of the meson spectrum with the experimental data. The lattice quark mass values can then be converted into quark mass values in the continuum QCD Lagrangian Eq. (1) using lattice perturbation theory at a scale given by the inverse lattice spacing. A recent computation gives a b -quark pole mass of 4.94 ± 0.15 GeV [4].

Potential model calculations of the hadron spectrum also involve the heavy quark mass. There is no way to relate the quark mass as defined in a potential model to the quark mass parameter of the QCD Lagrangian, or to the pole mass. Even in the heavy quark limit, the two masses can differ by nonperturbative effects of order Λ_{QCD} . There is also no reason why the potential model quark mass should be independent of the particular form of the potential used.

Recent work on the heavy quark effective theory [5–9] has provided a definition of the quark mass for a heavy quark that is valid when one includes nonperturbative effects and will be called the HQET mass m_Q . The HQET mass is particularly useful in the analysis of the $1/m_Q$ corrections in HQET. The HQET mass is not the same as the pole mass, a distinction that is often overlooked in the literature. Physical quantities such as hadron masses can in principle be computed in the heavy quark effective theory in terms of the HQET mass m_Q . The computations cannot be done analytically in practice because of nonperturbative effects in QCD, which also prevent a direct extraction of the quark masses from the original QCD Lagrangian, Eq. (1). Nevertheless, for heavy quarks, it is possible to parametrize the nonperturbative effects to a given order in the $1/m_Q$ expansion in terms of a few unknown constants that can be obtained from experiment. For example,

the B and D meson masses in the heavy quark effective theory are given in terms of a single nonperturbative parameter $\bar{\Lambda}$,

$$M(B) = m_b + \bar{\Lambda} + \mathcal{O}\left(\frac{\bar{\Lambda}^2}{m_b}\right),$$

$$M(D) = m_c + \bar{\Lambda} + \mathcal{O}\left(\frac{\bar{\Lambda}^2}{m_c}\right),$$

This allows one to determine the mass difference $m_b - m_c = M(B) - M(D) = 3.4$ GeV up to corrections of order $\bar{\Lambda}^2/m_b - \bar{\Lambda}^2/m_c$. The extraction of the individual quark masses m_b and m_c requires some knowledge of $\bar{\Lambda}$. An estimate of $\bar{\Lambda}$ using QCD sum rules gives $\bar{\Lambda} = 0.57 \pm 0.07$ GeV [10]. The HQET masses with this value of $\bar{\Lambda}$ are $m_b = 4.74 \pm 0.14$ GeV and $m_c = 1.4 \pm 0.2$ GeV, where the spin averaged meson masses $(3M(B^*) + M(B))/4$ and $(3M(D^*) + M(D))/4$ have been used to eliminate the spin-dependent $\mathcal{O}(\bar{\Lambda}^2/m_Q)$ correction terms. The errors reflect the uncertainty in $\bar{\Lambda}$ and the unknown spin-averaged $\mathcal{O}(\bar{\Lambda}^2/m_Q)$ correction. The errors do not include any theoretical uncertainty in the QCD sum rules, which could be large. A quark model estimate suggests that $\bar{\Lambda}$ is the constituent quark mass (≈ 350 MeV), which differs significantly from the sum rule estimate. In HQET, the $1/m_Q$ corrections to heavy meson decay form-factors are also given in terms of $\bar{\Lambda}$. Thus an accurate enough measurement of these form-factors could be used to extract $\bar{\Lambda}$ directly from experiment, which then determines the quark masses up to corrections of order $1/m_Q$.

The quark mass m_Q of HQET can be related to other quark mass parameters using QCD perturbation theory at the scale m_Q . The relation between these masses in QCD perturbation theory at one loop is [11]

$$m_Q = \bar{m}(\mu) \left[1 + \frac{\bar{\alpha}_s(\mu)}{3\pi} (3 \log \mu^2/m^2 + 4) \right],$$

$$m_Q = \hat{m}(\xi) \left[1 + \frac{\hat{\alpha}_s(\xi)}{\pi} \frac{\xi + 2}{\xi + 1} \log(\xi + 2) \right], \quad (2)$$

$$m_Q = m_P,$$

where $\bar{\alpha}_s(\mu)$ and $\hat{\alpha}_s(\xi)$ are the strong interaction coupling constants in the $\overline{\text{MS}}$ and momentum space subtraction schemes, respectively.

D. Light quarks

For light quarks, one can use the techniques of chiral perturbation theory to extract quark mass ratios. The light quark part of the QCD Lagrangian Eq. (1) has a chiral symmetry in the limit that the light quark masses are set to zero, under which left- and right-handed quarks transform independently. The mass term explicitly breaks the chiral symmetry, since it couples the left- and right-handed quarks to each other. A systematic analysis of this explicit chiral symmetry breaking provides some information on the light quark masses.

It is convenient to think of the three light quarks u , d and s as a three component column vector Ψ , and to write the mass term for the light quarks as

$$\bar{\Psi} M \Psi = \bar{\Psi}_L M \Psi_R + \bar{\Psi}_R M \Psi_L, \quad (3)$$

where M is the quark mass matrix M ,

$$M = \begin{pmatrix} m_u & 0 & 0 \\ 0 & m_d & 0 \\ 0 & 0 & m_s \end{pmatrix}. \quad (4)$$

The mass term $\bar{\Psi} M \Psi$ is the only term in the QCD Lagrangian that mixes left- and right-handed quarks. In the limit that $M \rightarrow 0$, there is an independent $\text{SU}(3)$ flavor symmetry for the left- and right-handed quarks. This $G_\chi = \text{SU}(3)_L \times \text{SU}(3)_R$ chiral symmetry of the QCD Lagrangian is spontaneously broken, which leads to eight massless Goldstone bosons, the π 's, K 's, and η , in the limit $M \rightarrow 0$. The symmetry G_χ is only an approximate symmetry, since it is explicitly broken by the quark mass matrix M . The Goldstone bosons acquire masses which can be computed in a systematic expansion in M in terms of certain unknown nonperturbative parameters of the theory. For example, to first order in M one finds that [12,13]

$$m_{\pi^0}^2 = B(m_u + m_d),$$

$$m_{\pi^\pm}^2 = B(m_u + m_d) + \Delta_{em},$$

$$m_{K^0}^2 = m_{K^+}^2 = B(m_d + m_s), \quad (5)$$

$$m_{K^\pm}^2 = B(m_u + m_s) + \Delta_{em},$$

$$m_\eta^2 = \frac{1}{3} B(m_u + m_d + 4m_s),$$

with two unknown parameters B and Δ_{em} , the electromagnetic mass difference. From Eq. (5), one can determine the quark mass ratios [12]

$$\frac{m_u}{m_d} = \frac{2m_{\pi^0}^2 - m_{\pi^+}^2 + m_{K^+}^2 - m_{K^0}^2}{m_{K^0}^2 - m_{K^+}^2 + m_{\pi^+}^2} = 0.56,$$

$$\frac{m_s}{m_d} = \frac{m_{K^0}^2 + m_{K^+}^2 - m_{\pi^+}^2}{m_{K^0}^2 + m_{\pi^+}^2 - m_{K^+}^2} = 20.1, \quad (6)$$

to lowest order in chiral perturbation theory. The error on these numbers is the size of the second-order corrections, which are discussed at the end of this section. Chiral perturbation theory cannot determine the overall scale of the quark masses, since it uses only the symmetry properties of M , and any multiple of M has the same G_χ transformation law as M . This can be seen from Eq. (5), where all quark masses occur only in the form Bm , so that B and m cannot be determined separately.

The mass parameters in the QCD Lagrangian have a scale dependence due to radiative corrections, and are renormalization scheme dependent. Since the mass ratios extracted using

Lepton & Quark Full Listings

Quarks

chiral perturbation theory use the symmetry transformation property of M under the chiral symmetry G_χ , it is important to use a renormalization scheme for QCD that does not change this transformation law. Any quark mass independent subtraction scheme such as $\overline{\text{MS}}$ is suitable. The ratios of quark masses are scale independent in such a scheme.

The absolute normalization of the quark masses can be determined by using methods that go beyond chiral perturbation theory, such as QCD sum rules [3]. Typically, one writes a sum rule for a quantity such as B in terms of a spectral integral over all states with certain quantum numbers. This spectral integral is then evaluated by assuming it is dominated by one (or two) of the lowest resonances, and using the experimentally measured resonance parameters [14]. There are many subtleties involved, which cannot be discussed here [14].

Another method for determining the absolute normalization of the quark masses, is to assume that the strange quark mass is equal to the SU(3) mass splitting in the baryon multiplets [12,14]. There is an uncertainty in this method since in the baryon octet one can use either the Σ - N or the Λ - N mass difference, which differ by about 75 MeV, to estimate the strange quark mass. It is also not possible to relate this definition of mass to the mass parameters in the QCD Lagrangian, which are scale and scheme dependent. It is often simply assumed that the baryon mass splittings give the $\overline{\text{MS}}$ quark masses renormalized at a scale $\mu = 1$ GeV.

One can extend the chiral perturbation expansion Eq. (5) to second order in the quark masses M to get a more accurate determination of the quark mass ratios. The meson mass terms in the chiral Lagrangian to second order in the quark mass matrix M are [16]

$$\mathcal{L} = \frac{1}{4} \text{Tr} (\chi^\dagger U + \chi U^\dagger) + L_6 \left[\text{Tr} (\chi^\dagger U + \chi U^\dagger) \right]^2 + L_7 \left[\text{Tr} (\chi^\dagger U - \chi U^\dagger) \right]^2 + L_8 \text{Tr} (\chi^\dagger U \chi^\dagger U + \chi U^\dagger \chi U^\dagger) \quad (7)$$

where $\chi = 2BM$ and U is the exponential of the pion field. Eq. (7) defines the chiral Lagrangian parameters L_{6-8} . The meson masses are obtained by expanding Eq. (7) in a power series in the meson fields.

There is a subtlety that arises at this order [15], because

$$M (M^\dagger M)^{-1} \det M^\dagger \quad (8)$$

transforms in the same way under G_χ as M , so that one can make the replacement $M \rightarrow M(\lambda) = M + \lambda M (M^\dagger M)^{-1} \det M^\dagger$,

$$\begin{aligned} M(\lambda) &= \text{diag} (m_u(\lambda), m_d(\lambda), m_s(\lambda)) \\ &= \text{diag} (m_u + \lambda m_d m_s, m_d + \lambda m_u m_s, m_s + \lambda m_u m_d) \quad (9) \end{aligned}$$

One can only determine the ratios $m_i(\lambda)/m_j(\lambda)$ using second-order chiral perturbation theory, not the ratios $m_i/m_j = m_i(\lambda = 0)/m_j(\lambda = 0)$, since second-order terms in the chiral Lagrangian are indistinguishable from first-order terms with

a redefined quark mass. M and $M(\lambda)$ have the same chiral transformation properties, which implies that the meson masses depend on L_{6-8} only in the linear combinations $L_6 - L_7$ and $L_6 + 2L_8$. One linear combination is left undetermined, so one cannot determine the quark mass ratios to second order in chiral perturbation theory without an independent determination of one of the L_i 's.

Dimensional analysis can be used to estimate [17] that second-order corrections in chiral perturbation theory due to the strange quark mass are of order $\lambda m_s \sim 0.25$. The ambiguity due to the redefinition Eq. (9) (which corresponds to a second-order correction) can produce a sizeable uncertainty in the ratio m_u/m_d . The lowest-order value $m_u/m_d = 0.56$ gets corrections of order $\lambda m_s (m_d/m_u - m_u/m_d) \sim 30\%$, whereas m_s/m_d gets a smaller correction of order $\lambda m_s (m_u/m_d - m_u m_d/m_s^2) \sim 15\%$. A more quantitative discussion of second-order effects can be found in Refs. 15,16,18. Since the second-order terms have a single parameter ambiguity, the value of m_u/m_d is related to the value of m_s/m_d .

The ratio m_u/m_d is of great interest since there is no strong CP problem if $m_u = 0$. To determine m_u/m_d requires determining the parameters L_{6-8} in the meson Lagrangian Eq. (7), or equivalently, fixing λ in the mass redefinition Eq. (9). There has been considerable effort to determine the chiral Lagrangian parameters accurately enough to determine m_u/m_d , for example from the analysis of the decays $\psi' \rightarrow \psi + \pi^0, \eta$, the decay $\eta \rightarrow 3\pi$, and by using sum rules [14,19-21]. m_u/m_d can also be extracted from η' decays. This requires the knowledge of chiral lagrangian parameters for the η' , such as L_{14} which is defined in [21]. All of these methods have large uncertainties, but indicate that $m_u/m_d \neq 0$. This conclusion has been questioned by Ref. 20 which argues that instanton corrections produce effects like the redefinition Eq. (9), and can explain the experimental observations even if $m_u = 0$.

Eventually, lattice gauge theory methods will be accurate enough to be able to compute meson masses directly from the QCD Lagrangian Eq. (1), and thus determine the light quark masses. For a reliable determination of quark masses, these computations will have to be done with dynamical fermions, and with a small enough lattice spacing that one can accurately compute the relation between lattice and continuum Lagrangians.

The quark masses for light quarks discussed so far are often referred to as current quark masses. Nonrelativistic quark models use constituent quark masses, which are of order 350 MeV for the u and d quarks. Constituent quark masses model the effects of dynamical chiral symmetry breaking, and are not related to the quark mass parameters m_k of the QCD Lagrangian Eq. (1). Constituent masses are only defined in the context of a particular hadronic model.

E. Numerical values and caveats

The quark masses in the particle data listings have been obtained by using the wide variety of theoretical methods outlined

See key on page 1343

Lepton & Quark Full Listings

Quarks, u, d, s , Light Quarks — u, d, s

above. Each method involves its own set of approximations and errors. In most cases, the errors are a best guess at the size of neglected higher-order corrections. The expansion parameter for the approximations is not much smaller than unity (for example it is $m_K^2/\Lambda_\chi^2 \approx 0.25$ for the chiral expansion), so an unexpectedly large coefficient in a neglected higher-order term could significantly alter the results. It is also important to note that the quark mass values can be significantly different in the different schemes. For example, assuming that the HQET mass m_b is 4.7 GeV, and $\bar{\alpha}_s(m_b) \approx 0.2$ gives the $\overline{\text{MS}}$ b -quark mass $\bar{m}_b(\mu = m_b) = 4.3$ GeV using the one-loop formula in Eq. (2). The heavy quark masses obtained using HQET, QCD sum rules, or lattice gauge theory are consistent with each other if they are all converted into the same scheme. When using the data listings, it is important to remember that the numerical value for a quark mass is meaningless without specifying the particular scheme in which it was obtained.

References

1. R. Tarrach, Nucl. Phys. **B183**, 384 (1981).
2. H. Georgi and H.D. Politzer, Phys. Rev. **D14**, 1829 (1976).
3. M.A. Shifman, A.I. Vainshtein, and V.I. Zakharov, Nucl. Phys. **B147**, 385 (1979).
4. C.T.H. Davies, *et al.*, Report No. SCRI-94-39 [hep-lat/9404012].
5. N. Isgur and M.B. Wise, Phys. Lett. **B232**, 113 (1989), *ibid* **B237**, 527 (1990); M.B. Voloshin and M. Shifman, Sov. J. Nucl. Phys. **45**, 292 (1987), *ibid* **47**, 511 (1988); S. Nussinov and W. Wetzel, Phys. Rev. **D36**, 130 (1987).
6. H. Georgi, Phys. Lett. **B240**, 447 (1990).
7. E. Eichten and B. Hill, Phys. Lett. **B234**, 511 (1990).
8. H. Georgi, in *Perspectives of the Standard Model*, ed. R.K. Ellis, C.T. Hill, and J.D. Lykken (World Scientific, Singapore, 1992); B. Grinstein, in *High Energy Phenomenology*, ed. R. Huerta and M.A. Pérez (World Scientific, Singapore, 1992).
9. A.F. Falk, M. Neubert, and M.E. Luke, Nucl. Phys. **B388**, 363 (1992).
10. M. Neubert, SLAC Report No. SLAC-PUB-6263 (June 1993), submitted to Phys. Reports.
11. S. Narison, Phys. Lett. **B197**, 405 (1987).
12. S. Weinberg, Trans. N.Y. Acad. Sci. **38**, 185 (1977).
13. See for example, H. Georgi, *Weak Interactions and Modern Particle Theory* (Benjamin/Cummings, Menlo Park, 1984).
14. J. Gasser and H. Leutwyler, Phys. Reports **87**, 77 (1982).
15. D.B. Kaplan and A.V. Manohar, Phys. Rev. Lett. **56**, 2004 (1986).
16. J. Gasser and H. Leutwyler, Nucl. Phys. **B250**, 465 (1985).
17. A. Manohar and H. Georgi, Nucl. Phys. **B234**, 189 (1984).
18. H. Leutwyler, Nucl. Phys. **B337**, 108 (1990).
19. P. Langacker and H. Pagels, Phys. Rev. **D19**, 2070 (1979); H. Pagels and S. Stokar, Phys. Rev. **D22**, 2876 (1980); H. Leutwyler, Nucl. Phys. **B337**, 108 (1990); J. Donoghue and D. Wyler, Phys. Rev. Lett. **69**, 3444 (1992); K. Maltman, T. Goldman and G.L. Stephenson Jr., Phys. Lett. **B234**, 158 (1990).
20. K. Choi, Nucl. Phys. **B383**, 58 (1992).
21. J. Donoghue and D. Wyler, Phys. Rev. **D45**, 892 (1992).

U

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 2$ to 8 MeV Charge = $\frac{2}{3} e$ $I_z = +\frac{1}{2}$
 $m_u/m_d = 0.25$ to 0.70

d

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$

Mass $m = 5$ to 15 MeV Charge = $-\frac{1}{3} e$ $I_z = -\frac{1}{2}$
 $m_s/m_d = 17$ to 25

S

$$I(J^P) = 0(\frac{1}{2}^+)$$

Mass $m = 100$ to 300 MeV Charge = $-\frac{1}{3} e$ Strangeness = -1
 $(m_s - (m_u + m_d)/2)/(m_d - m_u) = 34$ to 51

LIGHT QUARKS — u, d, s

u -QUARK MASS

The u -, d -, and s -quark masses are estimates of so-called "current-quark masses," in a mass-independent subtraction scheme such as $\overline{\text{MS}}$ at a scale $\mu \approx 1$ GeV. The ratios m_u/m_d and m_s/m_d are extracted from pion and kaon masses using chiral symmetry. The estimates of d and u masses are not without controversy and remain under active investigation. Within the literature there are even suggestions that the u quark could be essentially massless. The s -quark mass is estimated from SU(3) splittings in hadron masses.

VALUE (MeV) DOCUMENT ID TECN

2 to 8 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.8	¹ CHOI	92b THEO
5.1 ± 1.5	² BARDUCCI	88 THEO
1.8 ± 0.7	³ GASSER	82 THEO
5.6 ± 2.9	⁴ PAGELS	80 THEO
4.2	⁵ PAGELS	80 THEO
4	⁶ WEINBERG	77 THEO
	⁷ GASSER	75 THEO

¹ CHOI 92b argues that $m_u = 0$ is okay based on instanton contributions to the chiral coefficients. Disagrees with DONOGHUE 92 and DONOGHUE 92b.

² BARDUCCI 88 renormalized quark mass at 1 GeV. Uses a calculation of the effective potential for $\bar{\psi}\psi$ in QCD, and estimates for $\Sigma(p^2)$.

³ GASSER 82 uses chiral perturbation theory for the mass ratios, and uses QCD sum rules to extract the absolute values. The renormalization scale is 1 GeV.

⁴ PAGELS 80 uses lowest-order chiral perturbation theory plus an estimate of $\langle \bar{q}q \rangle$.

⁵ PAGELS 80 uses lowest-order chiral perturbation theory plus an estimate of $\langle \bar{q}q \bar{q}q \rangle$ correlation function.

⁶ WEINBERG 77 assumes that the baryon SU(3) splittings are equal to m_s .

⁷ GASSER 75 uses inelastic electron scattering and SU(6).

d -QUARK MASS

See the comment for the u quark above.

VALUE (MeV) DOCUMENT ID TECN

5 to 15 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

8.4	⁸ ADAMI	93 THEO
	⁹ NEFKENS	92 THEO
	¹⁰ BARDUCCI	88 THEO
	¹¹ DOMINGUEZ	87 THEO
	¹² KREMER	84 THEO
8.9 ± 2.6	¹³ GASSER	82 THEO
4.3 ± 0.7	¹⁴ PAGELS	80 THEO
14.6 ± 5.7	¹⁵ PAGELS	80 THEO
7.5	¹⁶ WEINBERG	77 THEO
6	¹⁷ GASSER	75 THEO

⁸ ADAMI 93 obtain $m_d - m_u = 3 \pm 1$ MeV at $\mu = 0.5$ GeV using isospin-violating effects in QCD sum rules.

⁹ NEFKENS 92 results for $m_d - m_u$ are 3.1 ± 0.4 MeV from meson masses and 3.6 ± 0.4 MeV from baryon masses.

¹⁰ BARDUCCI 88 renormalized quark mass at 1 GeV. Uses a calculation of the effective potential for $\bar{\psi}\psi$ in QCD, and estimates for $\Sigma(p^2)$.

¹¹ DOMINGUEZ 87 uses QCD sum rules to obtain $m_u + m_d = 15.5 \pm 2.0$ MeV and $m_d - m_u = 6 \pm 1.5$ MeV.

¹² KREMER 84 obtain $m_u + m_d = 21 \pm 2$ MeV at $Q^2 = 1$ GeV² using SVZ values for quark condensates; they obtain $m_u + m_d = 35 \pm 3$ MeV at $Q^2 = 1$ GeV² using factorization values for quark condensates.

¹³ GASSER 82 uses chiral perturbation theory for the mass ratios, and uses QCD sum rules to extract the absolute values. The renormalization scale is 1 GeV.

¹⁴ PAGELS 80 uses lowest-order chiral perturbation theory plus an estimate of $\langle \bar{q}q \rangle$.

¹⁵ PAGELS 80 uses lowest-order chiral perturbation theory plus an estimate of $\langle \bar{q}q \bar{q}q \rangle$ correlation function.

¹⁶ WEINBERG 77 assumes that the baryon SU(3) splittings are equal to m_s .

¹⁷ GASSER 75 uses inelastic electron scattering and SU(6).

Lepton & Quark Full Listings

Light Quarks — u, d, s

s-QUARK MASS

See the comment for the u quark above.

VALUE (MeV) DOCUMENT ID TECN

100 to 300 OUR EVALUATION

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 194 ± 4
- 118
- 175 ± 55
- >300
- 112 ± 66
- 378 ± 220
- 150
- 135
- 18 NEFKENS 92 THEO
- 19 DOMINGUEZ 91 THEO
- 20 BARDUCCI 88 THEO
- 21 KREMER 84 THEO
- 22 GASSER 82 THEO
- 23 PENSO 82B THEO
- 24 PAGELS 80 THEO
- 25 PAGELS 80 THEO
- 26 WEINBERG 77 THEO
- 27 GASSER 75 THEO
- 18 NEFKENS 92 results for $m_s - (m_u + m_d)/2$ are 111 ± 10 MeV from meson masses and 163 ± 15 MeV from baryon masses.
- 19 DOMINGUEZ 91 uses QCD sum rules.
- 20 BARDUCCI 88 renormalized quark mass at 1 GeV. Uses a calculation of the effective potential for $\bar{\psi}\psi$ in QCD, and estimates for $\Sigma(p^2)$.
- 21 KREMER 84 obtain $m_u + m_s = 245 \pm 10$ MeV at $Q^2 = 1$ GeV² using SVZ values for quark condensates; they obtain $m_u + m_s = 270 \pm 10$ MeV at $Q^2 = 1$ GeV² using factorization values for quark condensates.
- 22 GASSER 82 uses chiral perturbation theory for the mass ratios, and uses QCD sum rules to extract the absolute values. The renormalization scale is 1 GeV.
- 23 PENSO 82 uses SVZ sum rules to put a lower bound on the strange quark mass.
- 24 PAGELS 80 uses lowest-order chiral perturbation theory plus an estimate of $\langle \bar{q}q \rangle$.
- 25 PAGELS 80 uses lowest-order chiral perturbation theory plus an estimate of $\langle \bar{q}q \bar{q}q \rangle$ correlation function.
- 26 WEINBERG 77 assumes that the baryon SU(3) splittings are equal to m_s .
- 27 GASSER 75 is based on SU(6).

LIGHT QUARK MASS RATIOS

u/d MASS RATIO

VALUE DOCUMENT ID TECN

0.25 to 0.70 OUR EVALUATION

- • • We do not use the following data for averages, fits, limits, etc. • • •
- <0.3
- 0.26
- 0.30 ± 0.07
- 0.66
- 0.4 to 0.65
- 0.05 to 0.78
- 0.0 to 0.56
- 0.0 to 0.8
- 0.57 ± 0.04
- 0.38 ± 0.13
- 0.47 ± 0.11
- 0.56
- 28 CHOI 92 THEO
- 29 DONOGHUE 92 THEO
- 30 DONOGHUE 92B THEO
- 31 GERARD 90 THEO
- 32 LEUTWYLER 90B THEO
- 33 MALTMAN 90 THEO
- 34 CHOI 89B THEO
- 35 KAPLAN 86 THEO
- 36 GASSER 82 THEO
- 37 LANGACKER 79 THEO
- 38 LANGACKER 79B THEO
- 39 WEINBERG 77 THEO
- 28 CHOI 92 result obtained from the decays $\psi(2S) \rightarrow J/\psi(1S)\pi$ and $\psi(2S) \rightarrow J/\psi(1S)\eta$, and a dilute instanton gas estimate of some unknown matrix elements.
- 29 DONOGHUE 92 result is from a combined analysis of meson masses, $\eta \rightarrow 3\pi$ using second-order chiral perturbation theory including nonanalytic terms, and $(\psi(2S) \rightarrow J/\psi(1S)\pi)/(\psi(2S) \rightarrow J/\psi(1S)\eta)$.
- 30 DONOGHUE 92B computes quark mass ratios using $(\psi(2S) \rightarrow J/\psi(1S)\pi)/(\psi(2S) \rightarrow J/\psi(1S)\eta)$, and an estimate of L_{14} using Weinberg sum rules.
- 31 GERARD 90 uses large N and η - η' mixing.
- 32 LEUTWYLER 90B determines quark mass ratios using second-order chiral perturbation theory for the meson and baryon masses, including nonanalytic corrections. Also uses Weinberg sum rules to determine L_7 .
- 33 MALTMAN 90 uses second-order chiral perturbation theory including nonanalytic terms for the meson masses. Uses a criterion of "maximum reasonableness" that certain coefficients which are expected to be of order one are ≤ 3 .
- 34 CHOI 89 uses second-order chiral perturbation theory and a dilute instanton gas estimate of second-order coefficients in the chiral lagrangian.
- 35 KAPLAN 86 uses second-order chiral perturbation theory including nonanalytic terms for the meson masses. Assumes that less than 30% of the mass squared of the pion is due to second-order corrections.
- 36 GASSER 82 uses chiral perturbation theory for the meson and baryon masses.
- 37 LANGACKER 79 result is from a fit to the meson and baryon mass spectrum, and the decay $\eta \rightarrow 3\pi$. The electromagnetic contribution is taken from Socolow rather than from Dashen's formula.
- 38 LANGACKER 79B result uses LANGACKER 79 and also ρ - ω mixing.
- 39 WEINBERG 77 uses lowest-order chiral perturbation theory for the meson and baryon masses and Dashen's formula for the electromagnetic mass differences.

s/d MASS RATIO

VALUE DOCUMENT ID TECN

17 to 25 OUR EVALUATION

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 21
- 18
- 18 to 23
- 15 to 26
- 19.6 ± 1.5
- 22 ± 5
- 24 ± 4
- 20
- 40 DONOGHUE 92 THEO
- 41 GERARD 90 THEO
- 42 LEUTWYLER 90B THEO
- 43 KAPLAN 86 THEO
- 44 GASSER 82 THEO
- 45 LANGACKER 79 THEO
- 46 LANGACKER 79B THEO
- 47 WEINBERG 77 THEO
- 40 DONOGHUE 92 result is from a combined analysis of meson masses, $\eta \rightarrow 3\pi$ using second-order chiral perturbation theory including nonanalytic terms, and $(\psi(2S) \rightarrow J/\psi(1S)\pi)/(\psi(2S) \rightarrow J/\psi(1S)\eta)$.
- 41 GERARD 90 uses large N and η - η' mixing.
- 42 LEUTWYLER 90B determines quark mass ratios using second-order chiral perturbation theory for the meson and baryon masses, including nonanalytic corrections. Also uses Weinberg sum rules to determine L_7 .
- 43 KAPLAN 86 uses second-order chiral perturbation theory including nonanalytic terms for the meson masses. Assumes that less than 30% of the mass squared of the pion is due to second-order corrections.
- 44 GASSER 82 uses chiral perturbation theory for the meson and baryon masses.
- 45 LANGACKER 79 result is from a fit to the meson and baryon mass spectrum, and the decay $\eta \rightarrow 3\pi$. The electromagnetic contribution is taken from Socolow rather than from Dashen's formula.
- 46 LANGACKER 79B result uses LANGACKER 79 and also ρ - ω mixing.
- 47 WEINBERG 77 uses lowest-order chiral perturbation theory for the meson and baryon masses and Dashen's formula for the electromagnetic mass differences.

$(m_s - m)/(m_u - m_d)$ MASS RATIO

$m \equiv (m_u + m_d)/2$

VALUE DOCUMENT ID TECN

34 to 51 OUR EVALUATION

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 36 ± 5
- 45 ± 3
- 38 ± 9
- 43.5 ± 2.2
- 34 to 51
- 48 ± 7
- 48 NEFKENS 92 THEO
- 49 NEFKENS 92 THEO
- 50 AMETLLER 84 THEO
- GASSER 82 THEO
- GASSER 81 THEO
- MINKOWSKI 80 THEO
- 48 NEFKENS 92 result is from an analysis of meson masses, mixing, and decay.
- 49 NEFKENS 92 result is from an analysis of baryon masses.
- 50 AMETLLER 84 uses $\eta \rightarrow \pi^+ \pi^- \pi^0$ and ρ dominance.

ADAMI	93	PR D48 2304	+ Drukarev, Ioffe	(CIT, ITEP, PNPI)
CHOI	92	PL B292 159		(UCSD)
CHOI	92B	NP B303 58		(UCSD)
DONOGHUE	92	PRL 69 3444	+ Holstein, Wyler	(MASA, ZURI)
DONOGHUE	92B	PR D45 892	+ Wyler	(MASA, ZURI, UCSB)
NEFKENS	92	CNPP 20 221	+ Miller, Slaus	(UCLA, WASH, ZAGR)
DOMINGUEZ	91	PL B253 241	+ van Gend, Paver	(CAPE, TRST, INFN)
GERARD	90	MPL A5 391		(MPIM)
LEUTWYLER	90B	NP B337 108		(BERN)
MALTMAN	90	PL B234 158	+ Goldman, Stephenson Jr.	(YORKC, LANL)
CHOI	89B	PRL 62 849	+ Kim	(CMU, JHU)
BARUCCI	88	PR D38 238	+ Casalbuoni, De Curtis+	(FIRZ, INFN, LECE, GEVA)
Also	87	PL B193 305	Barducci, Casalbuoni+	(FIRZ, INFN, LECE, GEVA)
DOMINGUEZ	87	ANP 174 372	+ de Rafael	(ICTP, MARS, WIEN)
KAPLAN	86	PRL 56 2004	+ Manohar	(HARV)
AMETLLER	84	PR D30 674	+ Ayala, Bramon	(BARC)
KREMER	84	PL 143B 476	+ Papadopoulos, Schilcher	(MAIZ)
GASSER	82	PRPL 87 77	+ Leutwyler	(BERN)
PENSO	82	NC 68A 213	+ Penso, Truong	(ROMA, EPOL)
PENSO	82B	NC 72A 113	+ Verzegnassi	(ROMA, INFN, TRST, SISSA)
GASSER	81	ANP 136 62		(BERN)
MINKOWSKI	80	NP B164 25	+ Zepeda	(BERN)
PAGELS	80	PR D22 2876	+ Stokar	(ROCK)
LANGACKER	79	PR D19 2070	+ Pagels	(DESY, PRIN)
LANGACKER	79B	PR D20 2983		(PERN)
WEINBERG	77	ANYAS 38 185		(HARV)
GASSER	75	NP B94 269	+ Leutwyler	(BERN)

C

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Charm} = +1$$

c-QUARK MASS

The c-quark mass is estimated from charmonium and D masses. It corresponds to the "running" mass in the $\overline{\text{MS}}$ scheme. We have converted masses in other schemes to the $\overline{\text{MS}}$ scheme using one-loop QCD perturbation theory with $\alpha_s(\mu=m_c) = 0.39$. The range 1.0–1.6 GeV for the $\overline{\text{MS}}$ mass corresponds to 1.2–1.9 GeV for the pole mass (see the "Note on Quark Masses").

VALUE (GeV)	DOCUMENT ID	TECN
-------------	-------------	------

1.0 to 1.6 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.5 ^{+0.2} _{-0.1} ± 0.2	1 ALVAREZ	93 THEO
1.27 ± 0.02	2 NARISON	89 THEO
1.25 ± 0.05	3 NARISON	87 THEO
1.27 ± 0.05	4 GASSER	82 THEO

¹ ALVAREZ 93 method is to fit the measured x_F and p_T^2 charm photoproduction distributions to the theoretical predictions of ELLIS 89c.

² NARISON 89 determines the Georgi-Politzer mass at $p^2 = -m^2$ to be 1.26 ± 0.02 GeV using QCD sum rules.

³ NARISON 87 computes pole mass of 1.46 ± 0.05 GeV using QCD sum rules, with $\Lambda(\overline{\text{MS}}) = 180 \pm 80$ MeV.

⁴ GASSER 82 uses SVZ sum rules. The renormalization point is $\mu = \text{quark mass}$.

ALVAREZ	93	ZPHY C60 53	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ELLIS	89c	NP B312 551	+Nason	(FNAL, ETH)
NARISON	89	PL B216 191		(ICTP)
NARISON	87	PL B197 405		(CERN)
GASSER	82	PRPL 87 77	+Leutwyler	(BERN)

b

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = -\frac{1}{3} e \quad \text{Bottom} = -1$$

b-QUARK MASS

The b-quark mass is estimated from bottomonium and B masses. It corresponds to the "running" mass in the $\overline{\text{MS}}$ scheme. We have converted masses in other schemes to the $\overline{\text{MS}}$ scheme using one-loop QCD perturbation theory with $\alpha_s(\mu=m_b) = 0.22$. The range 4.1–4.5 GeV for the $\overline{\text{MS}}$ mass corresponds to 4.5–4.9 GeV for the pole mass (see the "Note on Quark Masses").

VALUE (GeV)	DOCUMENT ID	TECN
-------------	-------------	------

4.1 to 4.5 OUR EVALUATION

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.32 ± 0.05	1 DOMINGUEZ	92 THEO
4.24 ± 0.05	2 NARISON	89 THEO
4.18 ± 0.02	3 REINDERS	88 THEO
4.30 ± 0.13	4 NARISON	87 THEO
4.25 ± 0.1	5 GASSER	82 THEO

¹ DOMINGUEZ 92 determines pole mass to be 4.72 ± 0.05 using next-to-leading order in $1/m$ in moment sum rule.

² NARISON 89 determines the Georgi-Politzer mass at $p^2 = -m^2$ to be 4.23 ± 0.05 GeV using QCD sum rules.

³ REINDERS 88 determines the Georgi-Politzer mass at $p^2 = -m^2$ to be 4.17 ± 0.02 using moments of $\bar{b}\gamma\mu b$. This technique leads to a value for the mass of the B meson of 5.25 ± 0.15 GeV.

⁴ NARISON 87 determines the pole mass to be 4.70 ± 0.14 using QCD sum rules, with $\Lambda(\overline{\text{MS}}) = 180 \pm 80$ MeV.

⁵ GASSER 82 uses SVZ sum rules. The renormalization point is $\mu = \text{quark mass}$.

 $m_b - m_c$ MASS DIFFERENCE

The mass difference $m_b - m_c$ in the HQET scheme is 3.4 ± 0.2 GeV (see the "Note on Quark Masses").

VALUE (GeV)	DOCUMENT ID	TECN
-------------	-------------	------

• • • We do not use the following data for averages, fits, limits, etc. • • •

≥ 3.29	6 GROSSE	78
--------	----------	----

⁶ GROSSE 78 obtain $(m_b - m_c) \geq 3.29$ GeV based on eigenvalue inequalities in potential models.

DOMINGUEZ	92	PL B293 197	+Paver	(CAPE, TRST, INFN)
NARISON	89	PL B216 191		(ICTP)
REINDERS	88	PR D38 947		(BONN)
NARISON	87	PL B197 405		(CERN)
GASSER	82	PRPL 87 77	+Leutwyler	(BERN)
GROSSE	78	PL 79B 103	+Martin	(CERN)

Searches for t Quark

$$I(J^P) = 0(\frac{1}{2}^+)$$

$$\text{Charge} = \frac{2}{3} e \quad \text{Top} = +1$$

MASS LIMITS for t Quark or Hadron Independent of t Decay Mode

These limits are derived from $\Gamma(W)$ values shown in the W width section. Independent of the top decay mode, any W decay to $t\bar{b}$ would increase the total width of the W boson.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>62	95	1 ABE	94B CDF	$E_{cm}^{p\bar{p}} = 1800$ GeV
>45	95	2 ABE	92i CDF	$E_{cm}^{p\bar{p}} = 1800$ GeV
>53	95	3 ALITTI	92 UA2	$E_{cm}^{p\bar{p}} = 630$ GeV
>55	95	4 ALITTI	92 RVUE	
>43	95	5 ABE	91c CDF	$E_{cm}^{p\bar{p}} = 1800$ GeV
>38	90	6 ALBAJAR	91 UA1	$E_{cm}^{p\bar{p}} = 630$ GeV
>51	90	7 ALBAJAR	91 RVUE	$\Gamma(W)$

¹ ABE 94B result is from $\Gamma(W) = 2.063 \pm 0.061 \pm 0.060$ GeV.

² ABE 92i data include both e and μ final states. The result is derived from $\Gamma(W) = 2.16 \pm 0.17$ GeV. At 90%CL, the limit is >49 GeV.

³ ALITTI 92 result is derived from $\Gamma(W) = 2.10 \pm 0.16$ GeV.

⁴ Limit is from combined data of ALBAJAR 91, ALITTI 92, and ABE 90: $\Gamma(W) = 2.15 \pm 0.11$ GeV.

⁵ ABE 91c result is derived from $\Gamma(W) = 2.12 \pm 0.20$ GeV. At 90%CL, the limit is > 48 GeV.

⁶ ALBAJAR 91 result is derived from $\Gamma(W) = 2.18_{-0.24}^{+0.26} \pm 0.04$ GeV.

⁷ Limit is from combined data of ALBAJAR 91, ALITTI 90c, and ABE 90.

MASS LIMITS for t Quark in $p\bar{p}$ Collisions

These experiments are based on the assumption that no nonstandard decay modes such as $t \rightarrow bH^+$ are available, except as shown in the comments. Mass limits are now sufficiently high that decay is expected to occur before hadronization.

VALUE (GeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
>131	95		8 ABACHI	94 D0	$\ell\ell + \text{jets}, \ell + \text{jets}$
$174 \pm 10_{-12}^{+13}$		7	9 ABE	94E CDF	$\ell + b\text{-jet}$
>118	95		9 ABE	94E CDF	$\ell\ell$
> 91	95		10 ABE	92 CDF	$\ell\ell, \ell + b\text{-jet}$
			11 ALITTI	92F UA2	$t \rightarrow bH^+, H^+ \rightarrow \tau\nu_\tau$
> 60	95		12 ALBAJAR	91B UA1	$t \rightarrow bH^+; H^+ \rightarrow \tau^+\nu$
			13 BAER	91B RVUE	$t \rightarrow \bar{t}_1 \bar{\chi}_1^0 \tau^+\nu$
> 72	95		14 ABE	90B CDF	$e + \mu$
> 77	95		15 ABE	90C CDF	$e + \text{jets} + \text{missing } E_T$
> 69	95		16 AKESSON	90 UA2	$e + \text{jets} + \text{missing } E_T$
> 60	95		ALBAJAR	90B UA1	e or $\mu + \text{jets}, \mu\mu + \text{jet}$
			17 BARGER	90E RVUE	$t \rightarrow bH^+$
> 41	95		18 ALBAJAR	88 UA1	e or $\mu + \text{jets}$

⁸ ABACHI 94 search for $e\mu + \text{jets}, ee + \text{jets}, e + \text{jets},$ and $\mu + \text{jets}$. Production cross section with soft-gluon resummation of LAENEN 94 is used. The limit decreases to >122 GeV if $\mathcal{O}(\alpha_s^3)$ cross section is employed for comparison with ABE 92.

⁹ ABE 94E search for $ee, e\mu,$ and $\mu\mu$ dilepton final states and single lepton + b -jet final states. They observe a total of 15 top topology tags (12 events of which three are doubly tagged) with an expected background of $5.96_{-0.44}^{+0.49}$. The mass determination is from 7 single-lepton + b -jet events which have four jets. Their $\ell\ell$ limit uses the production cross section with soft gluon resummation from LAENEN 94.

¹⁰ ABE 92 search for $ee, e\mu, \mu\mu$ dilepton final states and $(e$ or $\mu)$ plus a b -quark jet. The b jet is tagged by a soft muon. The 90%CL limit is 95 GeV. Superseded by ABE 94E $\ell\ell$ limit.

¹¹ ALITTI 92F search for $t \rightarrow bH^+, H^+ \rightarrow \tau\nu_\tau$ with τ decaying hadronically. m_t between 50 and 70 GeV is excluded if $m_t - m_{H^+} = m_b + (\lesssim \text{a few } -10 \text{ GeV})$. See their Figs. 5,6 for the excluded region for $B(H^+ \rightarrow \tau\nu_\tau) = 1, 0.5$.

¹² ALBAJAR 91B searched for the decay $t \rightarrow H^+ b$ using single muon and dimuon events and assuming $B(H^+ \rightarrow \tau^+\nu) \geq 0.95$. The limit holds for $m_{H^+} \lesssim m_t - m_b - (3-6)$ GeV.

¹³ BAER 91B argue that a top quark as light as 60 GeV (65 GeV, if the minimal SUSY framework is assumed) may have escaped detection at CDF if a supersymmetric decay mode is open.

¹⁴ ABE 90B exclude the region 28–72 GeV.

¹⁵ ABE 90C cannot exclude $m_t < 40$ GeV, but this region is ruled out by other experiments. They study events with an energetic electron, missing transverse energy and two or more jets. Only the $t\bar{t}$ contribution (not $W \rightarrow tb$) is relevant for these masses. See also ABE 91.

¹⁶ AKESSON 90 searched for events having an electron with $p_T > 12$ GeV, missing momentum > 15 GeV, and a jet with $E_T > 10$ GeV, $|\eta| < 2.2$, and excluded m_t between 30 and 69 GeV.

¹⁷ BARGER 90E claim that ABE 90C data exclude most regions of two-Higgs-doublet models with $m_t < 80$ GeV even if $t \rightarrow bH^+$ decay is allowed.

¹⁸ ALBAJAR 88 value quoted here is revised using the full $\mathcal{O}(\alpha_s^3)$ cross section of ALTARELLI 88. Superseded by ALBAJAR 90B.

Lepton & Quark Full Listings

t Quark

INDIRECT MASS LIMITS for t Quark from Standard Model Electroweak Fit

The RVUE values are based on the data described in the footnotes. Earlier RVUE's are superseded but have been left in the Listings to show the progress.

"OUR EVALUATION" below is for our fit to electroweak data described in the "Standard Model of Electroweak Interactions" section. This fit result does not include direct measurements of m_t . The second error corresponds to $m_H=300^{+700}_{-240}$ GeV.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
169^{+16+17}_{-18-20}		OUR EVALUATION		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
157^{+36+19}_{-48-20}		19 ABREU	94 DLPH	Z parameters
$158^{+32}_{-40} \pm 19$		20 ACCIARRI	94 L3	Z parameters
132^{+41+24}_{-48-18}		21 AKERS	94 OPAL	Z parameters
184^{+25+17}_{-29-18}		22 BUSKULIC	94 ALEP	Z parameters
140^{+21}_{-22}		23 ELLIS	94 RVUE	Electroweak
$91 \pm 46 \pm 9$		24 ACTON	93D OPAL	Z parameters
$152^{+36}_{-46} \pm 20$		25 ADRIANI	93M L3	Z parameters
<207	95	26 ALTARELLI	93 RVUE	Z $b\bar{b}$ vertex
143^{+19}_{-18}		27 BLONDEL	93 RVUE	Z parameters
174^{+27+17}_{-32-22}		28 BUSKULIC	93J ALEP	Z parameters
<228	95	29 BUSKULIC	93M ALEP	$\Gamma(Z \rightarrow b\bar{b})$
132^{+20}_{-22}		30 ELLIS	93B RVUE	
102^{+35+19}_{-32-18}		31 MONTAGNA	93 RVUE	Z parameters
114^{+32}_{-42}		32 MONTAGNA	93B RVUE	Z parameters
$146^{+18}_{-19} \pm 17$		33 NOVIKOV	93B RVUE	
147^{+22+17}_{-26-22}		34 PASSARINO	93 RVUE	Electroweak
160^{+50}_{-60}		35 QUAST	93 RVUE	Z parameters
$123^{+33}_{-38} \pm 19$		36 ALITTI	92B UA2	m_W, m_Z
<208	95	37 BANERJEE	92 RVUE	Electroweak
170^{+42+21}_{-55-14}		38 BLONDEL	92 RVUE	Z $b\bar{b}$ vertex
112^{+22}_{-23}		39,40 DECAMP	92B ALEP	Z parameters
122^{+25}_{-20}		41 DELAGUILA	92 RVUE	Electroweak
120^{+27}_{-28}		42 DELAGUILA	92B RVUE	
124^{+26}_{-28}		43 ELLIS	92 RVUE	Electroweak
$124^{+40}_{-56} \pm 21$		44 ELLIS	92E RVUE	
132^{+27+18}_{-31-19}		45 LANGACKER	92B RVUE	Z'
$150^{+23}_{-26} \pm 16$		46 LEP	92 RVUE	Z parameters
137^{+22+18}_{-25-22}		46 LEP	92 RVUE	Electroweak
150^{+29+20}_{-34-22}		47 PDG	92 RVUE	Electroweak
144^{+23+19}_{-26-21}		48 RENTON	92 RVUE	Electroweak
<220	95	49 SCHAILE	92 RVUE	Z parameters
<215	95	49 SCHAILE	92 RVUE	Electroweak
$193^{+52}_{-69} \pm 16$		50 ABE	91B RVUE	m_W, m_Z
100^{+70+24}_{-52-11}		51 ABREU	91F DLPH	Z parameters
119^{+39}_{-45}		51 ADEVA	91E L3	Z parameters
134^{+47}_{-48}		52,53 ALEXANDER	91F OPAL	Z parameters
124^{+28+20}_{-34-15}		54 DELAGUILA	91C RVUE	Z'
<366	90	55 GONZALEZ-G.	91 RVUE	Electroweak
<200	95	56 HIOKI	91 RVUE	Electroweak
<200	95	57 LANGACKER	91 RVUE	Electroweak
<240	90	58 ADACHI	90F RVUE	$\sigma(e^+e^- \rightarrow \text{hadrons})$
$120 \pm 40 \pm 20$		59 ADEVA	90I RVUE	Electroweak
127^{+24}_{-30}		60 BARGER	90C RVUE	Electroweak
<190	95	61 BLONDEL	90 CDHS	$\nu N \rightarrow \nu X$ or $\bar{\nu} N \rightarrow \bar{\nu} X$
140^{+43}_{-52}		62 DECAMP	90P RVUE	Electroweak
<168	90	63 ELLIS	90B RVUE	Electroweak
<153	68	64 KENNEDY	90 RVUE	Electroweak
<291	90	65 ELLIS	89B RVUE	Electroweak
<180	90	66 LANGACKER	89 RVUE	Electroweak
	90	67 COSTA	88 RVUE	Electroweak
	68	68 ELLIS	88C RVUE	Electroweak
	90	69 FOGLI	88 RVUE	$\nu N \rightarrow \nu X$
	90	70 AMALDI	87 RVUE	Electroweak

- 19 ABREU 94 value is for $\alpha_s(m_Z)$ constrained to 0.123 ± 0.005 . The second error corresponds to $m_H = 300^{+700}_{-240}$ GeV.
- 20 ACCIARRI 94 value is for $\alpha_s(m_Z)$ constrained to 0.124 ± 0.006 . The second error corresponds to $m_H = 300^{+700}_{-240}$ GeV.
- 21 AKERS 94 result is from fit with free α_s . The second error corresponds to $m_H=300^{+700}_{-240}$ GeV. The 95%CL limit is $m_t < 210$ GeV.
- 22 BUSKULIC 94 result is from fit with free α_s . The second error is from $m_H=300^{+700}_{-240}$ GeV.
- 23 ELLIS 94 is fit to electroweak data available in summer 1993. m_t , m_H , and α_s are adjusted to minimize χ^2 , yielding m_t above, $m_H = 35^{+205}_{-26}$ GeV, and $\alpha_s(m_Z) = 0.116^{+0.007}_{-0.006}$.
- 24 ACTON 93D result is from fit with free α_s . The second error corresponds to $m_H=300^{+700}_{-240}$ GeV. The 95%CL limit is $m_t < 180$ GeV. Negative statistical error is larger than -46. Superseded by AKERS 94.
- 25 ADRIANI 93M used $\alpha_s(m_Z)=0.124 \pm 0.006$. Second error corresponds to $m_H=50$ GeV - 1 TeV. Superseded by ACCIARRI 94.
- 26 ALTARELLI 93 limit is from fit to electroweak data available in summer '92 but uses only the parameter corresponding to Z $b\bar{b}$ vertex correction. $\alpha_s(m_Z) = 0.118 \pm 0.007$ is used.
- 27 BLONDEL 93 is $m_t - m_H$ fit to LEP data available in spring '93. $\alpha_s = 0.117 \pm 0.005$ is used and $m_t > 108$ GeV, $m_H > 62.5$ GeV imposed. 95%CL limit is 185 GeV.
- 28 BUSKULIC 93J second error is from $m_H=300^{+700}_{-250}$ GeV. Superseded by BUSKULIC 94.
- 29 BUSKULIC 93M limit is from $\Gamma(b\bar{b})/\Gamma(\text{had}) = 0.2193 \pm 0.0029$. The best value is $m_t = 50 \pm 70$ GeV. The CDF limit $m_t > 91$ GeV is imposed to obtain the limit.
- 30 ELLIS 93B fit to electroweak data available in spring '93. m_H is adjusted to minimize χ^2 (even below the present direct limit) and $\alpha_s(m_Z) = 0.123 \pm 0.006$ is used. 95%CL limit of $m_t < 155$ GeV is claimed.
- 31 MONTAGNA 93 perform fit to LEP cross-section and asymmetry data taken in 1989/90. The second error corresponds to $m_H = 300^{+700}_{-235}$ GeV. Direct limits on m_t , m_H are imposed. α_s is adjusted to minimize χ^2 . Updated in MONTAGNA 93B.
- 32 MONTAGNA 93B is an update of MONTAGNA 93, including fits to realistic (i.e. with experimental cuts) cross sections and asymmetries. m_Z and α_s are adjusted to minimize χ^2 . m_H is fixed at $m_H = 300$ GeV.
- 33 See Fig. 2 of NOVIKOV 93B for χ^2 contour in $m_t - m_H$ plane calculated from m_W , $\Gamma(\ell\ell)$, and $A_{FB}^{\ell\ell}$ data available in summer '92.
- 34 PASSARINO 93 fit is to LEP and m_W data available in spring '93. α_s is adjusted to minimize χ^2 . The second error is from $m_H=300^{+700}_{-240}$ GeV.
- 35 QUAST 93 is fit to LEP data taken up to 1991. α_s is fitted ($0.133 \pm 0.008 \pm 0.002$). The second error is from $m_H=300^{+700}_{-240}$ GeV.
- 36 ALITTI 92B assume $m_H = 100$ GeV. The 95%CL limit is $m_t < 250$ GeV for $m_H < 1$ TeV.
- 37 BANERJEE 92 is a fit to LEP data taken in '90 as well as m_W . The second error is from $m_H=300^{+700}_{-250}$ GeV. $\alpha_s(m_Z) = 0.118 \pm 0.008$ is used. Fit to LEP data only gives $m_t = 99 \pm 48 \pm 22$ GeV.
- 38 BLONDEL 92 limit is from $\Gamma(b\bar{b})/\Gamma(\text{had})$, $\sin^2\theta_W^{\text{eff}}$ from asymmetries, and $\alpha_s(m_Z) = 0.117 \pm 0.004$ (data available in spring '92). $m_t > 91$ GeV imposed. The limit is sensitive to the value of α_s .
- 39 Limit from Z cross sections, leptonic forward-backward asymmetries, τ polarization asymmetry, quark charge asymmetry, and $b\bar{b}$ and $c\bar{c}$ forward-backward asymmetries.
- 40 DECAMP 92B uses $\alpha_s = 0.121 \pm 0.008$. The second error is from $m_H = 200^{+800}_{-150}$ GeV. The "Electroweak" value combines ALEPH Z data and m_W/m_Z from ALITTI 90B and ABE 90G.
- 41 DELAGUILA 92 is fit to electroweak data including LEP data taken '90. The value is for $m_H = 100$ GeV, $\alpha_s(m_Z) = 0.12$ fixed. For $m_H = 1$ TeV, $m_t = 140^{+19}_{-21}$ GeV.
- 42 DELAGUILA 92B perform two-dimensional fit to various electroweak data with direct limits on m_t, m_H .
- 43 ELLIS 92 is an update of ELLIS 90B to include the latest LEP, UA2, CDF, and CHARM II results presented at the Lepton-Photon and EPS Conference, July 1991. $\alpha_s = 0.115 \pm 0.008$ assumed and m_H left free. Fit gives $1.3 \text{ GeV} < m_H < 160 \text{ GeV}$, CL = 68% for $m_t = 130$ GeV.
- 44 ELLIS 92E perform fit to electroweak data available in spring '92. m_H is adjusted to minimize χ^2 and $\alpha_s(m_Z) = 0.118 \pm 0.008$ is used.
- 45 LANGACKER 92B consider the effect of an extra Z boson ($Z'_{SM}, Z_{LR}, Z_X, Z_\psi, Z_\eta$) on the top mass determination. The fit including Z' does not change the limit on m_t .
- 46 The LEP 92 values are combined results of the four LEP collaborations: ALEPH, DELPHI, L3, and OPAL. The "Electroweak" result includes m_W and m_W/m_Z from ABE 90G and ALITTI 92B, and neutral current data from CDHS and CHARM. Uses $\alpha_s = 0.118 \pm 0.008$. Second error corresponds to $m_H = 300^{+700}_{-250}$ GeV.
- 47 PDG 92 value comes from a fit by P. Langacker to recent data as discussed in the minireview above.
- 48 RENTON 92 is a fit to LEP data taken up to '90 as well as m_W , νN , and atomic parity violation data. The second error is from $m_H=300^{+700}_{-250}$ GeV. $\alpha_s(m_Z) = 0.114 \pm 0.007$ is used. The 95%CL limit is 193 GeV. The fit to LEP data only gives $m_t=142^{+30}_{-37} \pm 20$ GeV.
- 49 SCHAILE 92 performs fit to LEP electroweak data (as of summer 1991) as well as m_W (UA2/CDF) and νN (CDHS/CHARM). The second error is from $m_H=300^{+700}_{-250}$ GeV.
- 50 The ABE 91B limit is derived from their m_W with $m_Z = 91.161$ GeV. Combining with the m_W measurement of ALITTI 90B, one obtains $m_t < 230$ GeV (95%CL).
- 51 $\alpha_s = 0.115 \pm 0.009$. The second error is from $m_H = 300^{+700}_{-250}$ GeV.
- 52 ALEXANDER 91F use $\alpha_s = 0.118 \pm 0.008$. The second error comes from $m_H = 300^{+700}_{-250}$ GeV.

- 53 The 95%CL upper limit is 218 GeV.
- 54 DELAGUILA 91c study bound on m_t in the presence of extra Z' (Z_{LR} and Z_X) from various electroweak data. The upper bound on m_t is more strict for lower Z' masses. See their Fig. 2. See also DELAGUILA 92.
- 55 GONZALEZ-GARCIA 91 result is based on low-energy neutral current data, Z mass and widths, m_W from ABE 90G. $m_{H^0} = 100$ GeV assumed.
- 56 HIOKI 91 uses m_Z , $\Gamma_{\text{total}}(Z)$, and m_W . $m_{H^0} = 100$ GeV, $\alpha_S = 0.12$ assumed. For $m_{H^0} = 1$ TeV, one finds $m_t = 162^{+43}_{-46}$ GeV.
- 57 LANGACKER 91 is a fit to various electroweak data. The second error is from $m_{H^0} = 250^{+750}_{-200}$ GeV. $\alpha_S = 0.12 \pm 0.02$ used. The 95%CL upper limit is 182 GeV [for $m_{H^0} = 1$ TeV]. For arbitrary Higgs structure, one obtains $m_t < 310$ GeV.
- 58 ADACHI 90F limit is from R at PEP, PETRA, and TOPAZ at TRISTAN. Top mass dependence enters via radiative correction. Minimal standard model with $m_Z = 91.1$ GeV, $m_{H^0} = 100$ GeV assumed. $\Lambda_{\overline{MS}}$ is varied in the fit.
- 59 ADEVA 90i analysis is based on m_Z measured by L3 and $\sin^2\theta_W = 0.2284 \pm 0.0043$ determined from m_W/m_Z and νN scattering data. $40 < m_H < 1000$ GeV assumed. The 1σ range is $m_t = 130^{+38}_{-42}$ GeV.
- 60 BARGER 90C limit is a fit using only LEP and m_W data. $m_{H^0} = 100$ GeV assumed. The most likely value is $m_t = 151$ GeV. The limit increases to 225 GeV for $m_H = 1000$ GeV.
- 61 BLONDEL 90 limit comes from $R_{\nu} = \sigma^{NC}(\nu N) / \sigma^{CC}(\nu N)$ and R_{ν} . Comparison of R_{ν} and m_W (the latter from ALBAJAR 89 and ANSARI 87) gives an independent limit $m_t < 240$ GeV (90%CL).
- 62 DECAMP 90P result is from m_Z , $\Gamma(Z \rightarrow \ell\ell)$, and m_W/m_Z from UA2 (ALITTI 90B), m_W from CDF (APS conf. '90), and νN neutral current data from CDHS and CHARM.
- 63 ELLIS 90B limit is a fit to various electroweak data. $m_{H^0} = m_Z$ assumed. $m_c = 1.45$ GeV is used for νN data.
- 64 KENNEDY 90 limit is a fit to neutral current data, W , Z masses, and Z widths. $m_H = m_Z$ assumed. For $m_H = 1$ TeV, the limit is 212 GeV. For nonminimal Higgs sector (with $\rho \neq 1$), one obtains $m_t < 350$ GeV (90%CL).
- 65 ELLIS 89B limit is a fit to various electroweak data. $m_{H^0} = m_Z$ assumed. $m_c = 1.45$ GeV is used for νN data. Superseded by ELLIS 90B.
- 66 LANGACKER 89 limit is a fit to various electroweak data. $m_{H^0} = 100$ GeV assumed. The 90%CL upper limit is 190(210) GeV for $m_{H^0} = 100(1000)$ GeV.
- 67 COSTA 88 limit is a fit to various electroweak data. $m_{H^0} = m_Z$ assumed. $m_c = 1.5$ GeV is used for νN data.
- 68 ELLIS 88C limit is a fit to neutral current data and W , Z masses. $m_{H^0} = m_Z$ assumed. $m_c = 1.45$ GeV is used for νN data. Varying m_c relaxes the limit to 185 GeV. Superseded by ELLIS 89B.
- 69 FOGLEI 88 limit is a fit to neutrino deep-inelastic scattering data.
- 70 AMALDI 87 limit is a fit to various electroweak data. $m_{H^0} < 100$ GeV assumed.

MASS LIMITS for Top Hadrons in e^+e^- Collisions

The last column specifies measured quantities: $S =$ Sphericity, $T =$ Thrust.

For limits prior to 1987, see our 1990 edition, Physics Letters **B239**, p. VII.167 (1990).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.8	95	71 DECAMP	90F ALEP	isolated charged particle and aplanarity
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
>41.8	95	72 ADRIANI	93G L3	Quarkonium
>43	95	71 ADRIANI	93M L3	$\Gamma(Z)$
>30.2	95	71 ABREU	91F DLPH	$\Gamma(Z)$
>44.5	95	71 ABE	90D VNS	Event shape
>44.0	95	71 ABREU	90D DLPH	Event shape
>44.0	95	71,73 ABREU	90D DLPH	$t \rightarrow bH^+, H^+ \rightarrow c\bar{s}$, $\tau^+\nu$
>33.5	95	74 ABREU	90D DLPH	$\Gamma(Z \rightarrow \text{hadrons})$
>44.5	95	75 AKRAWY	90B OPAL	Acoplanarity
>44.3	95	76 AKRAWY	90B OPAL	$t \rightarrow bH^+, H^+ \rightarrow c\bar{s}$, $\tau\nu$
>40.7	95	77 ABRAMS	89C MRK2	Event shape
>42.5	95	77 ABRAMS	89C MRK2	$t \rightarrow bH^+, H^+ \rightarrow c\bar{s}$
>29.9	95	78 ADACHI	89C TOPZ	μ
>29.9	95	79 ENO	89 AMY	μ, e
>25.8	95	80 ADACHI	88 TOPZ	R, T , Acoplanarity
>25.9	95	81 IGARASHI	88 AMY	$T + (\mu, e)$
>25.9	95	82 SAGAWA	88 AMY	R, T
none $E_{cm}=50$	95	83 ABE	87 VNS	R, T , Acoplanarity
>25.5	95	84 YOSHIDA	87 VNS	R, T , Acoplanarity

- 71 Search was near the Z peak at LEP.
- 72 ADRIANI 93G search for vector quarkonium states near Z and give limit on quarkonium- Z mixing parameter $\delta m^2 < (10-30)$ GeV² (95%CL) for the mass 88-94.5 GeV. Using Richardson potential, a 1S toponium state is excluded for the mass range 87.9-88.7, 89.1-94.3 GeV. This range is very sensitive to the potential choice.
- 73 Assumed $m_{H^+} < m_t - 6$ GeV.
- 74 Superseded by ABREU 91F.
- 75 AKRAWY 90B search was restricted to data near the Z peak at $E_{cm} = 91.26$ GeV at LEP. The excluded region is between 23.4 and 44.5 GeV if no H^+ decays exist.
- 76 AKRAWY 90B limit applies for any H^+ branching ratio $B(c\bar{s})$. Limit increases to 45.2 GeV if $B(c\bar{s}) = 1$. The lower end of the excluded region is $m_{H^+} + 5$ GeV.
- 77 The ABRAMS 89C limit from an isolated track search is 40.0 GeV.
- 78 ADACHI 89C search was at $E_{cm} = 56.5-60.8$ GeV at TRISTAN using multi-hadron events accompanying muons.
- 79 ENO 89 search at $E_{cm} = 50-60.8$ GeV at TRISTAN.

- 80 ADACHI 88 set limit $\sigma(\text{top}) < 8.2$ pb at CL=95% for top-flavored-hadron production from event shape analyses at $E_{cm} = 52$ GeV. By using the quark-parton model cross-section formula with first-order QCD corrections near the threshold, the above limit leads to a lower mass limit of 25.8 GeV at 95% confidence level for top quarks.
- 81 IGARASHI 88 searches for leptons in low-thrust events and gives $\Delta R(t) < 0.15$ (95% CL) at $E_{cm} = 50-52$ GeV.
- 82 SAGAWA 88 set limit $\sigma(\text{top}) < 6.1$ pb at CL=95% for top-flavored hadron production from event shape analyses at $E_{cm} = 52$ GeV. By using the quark parton model cross-section formula near threshold, the above limit leads to lower mass bounds of 25.9 GeV for charge 2/3 quarks.
- 83 ABE 87 set limit $\sigma(\text{top}) < 16$ pb at CL=95% for top-flavored hadron production, which should be compared with the full top-quark production cross section of 45.9 pb.
- 84 YOSHIDA 87 set limit $\sigma(\text{top}) < 17$ pb at CL=95% for top-flavored hadron production from event shape analyses at $E_{cm} = 52$ GeV. This limit should be compared with the full top-quark production cross section of 34 pb, which takes into account the effect of weak neutral current but neglects its axial-vector coupling contribution expected to be suppressed near threshold. After considering the radiative effects, top quarks of mass below 25.5 GeV can be excluded by the above limit.

REFERENCES FOR Searches for t Quark

ABACHI 94 PRL 72 2138	+Abbott, Abolins, Acharya, Adam+ (D0 Collab.)
ABE 94B (to be pub.)	+Albrow, Amidei, Antos, Anway-Wiese+ (CDF Collab.)
Fermilab-PUB-94-051-E	
ABE 94E PR D (to be pub.)	+Albrow, Amidei, Antos, Anway-Wiese+ (CDF Collab.)
Fermilab-PUB-94-097-E	
Also Fermilab-PUB-94-097-E (subm.)	Abe, Albrow, Amidei, Antos, Anway-Wiese+ (CDF Collab.)
ABREU 94 NP B418 403	+Adam, Adye, Agasi+ (DELPHI Collab.)
ACCARRI 94 ZPHY C62 551	+Adam, Adriani, Aguilera-Benitez+ (L3 Collab.)
AKERS 94 ZPHY C61 19	+Alexander, Allison+ (OPAL Collab.)
BUSKULIC 94 ZPHY C62 539	+Casper, De Bonis, Decamp, Ghez, Goy+ (ALEPH Collab.)
ELLIS 94 PL B324 173	+Fogli, Lisi (CERN, BARI)
LAENEN 94 PL B321 254	+Smith, van Neerven (FNAL, UTRE, LEID)
XZD ZPH C58 219	+Alexand, Allison+ (OPAL Collab.)
ADRIANI 93G PL B313 326	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+ (L3 Collab.)
ADRIANI 93M PRPL 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+ (L3 Collab.)
ALTARELLI 93 NP B405 3	+Barberis, Caravagios (CERN, PISA, PISAL, SNSP)
BLONDEL 93 PL B311 346	+Verzegnassi (EPOL, TRSTT, TRSTI)
BUSKULIC 93J ZPHY C60 71	+Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
BUSKULIC 93M PL B313 535	+De Bonis, Decamp+ (ALEPH Collab.)
ELLIS 93B PL B318 148	+Fogli, Lisi (CERN, BARI)
93 PL B303 170	+Nicosini, Passarino (PAVI, TORI)
MONTAGNA 93B NP B401 3	+Piccinini, Nicosini, Passarino, Pittau (PAVI, TORI)
NOVIKOV 93B PL B308 123	+Okun, Vysotsky, Yurov (SERP, CERN)
PASSARINO 93 PL B313 213	
QUAST 93 MPL A8 675	
ABE 92 PRL 68 447	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
Also 92G PR D46 3921	+Abe, Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE 92H PR D46 3925	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ALITTI 92 PL B276 365	+Ambrosini, Ansari, Autiero, Barezys+ (UA2 Collab.)
ALITTI 92B PL B276 354	+Ambrosini, Ansari, Autiero, Barezys+ (UA2 Collab.)
ALITTI 92F PL B280 137	+Ambrosini, Ansari, Autiero, Barezys+ (UA2 Collab.)
BANERJEE 92 IJMP A7 1853	+Ganguli, Gurtu (TATA)
BLONDEL 92 PL B293 253	+Djouadi, Verzegnassi (EPOL, DESY, TRSTI, TRSTT)
DECAMP 92B ZPHY C53 1	+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
DELAGUILA 92 NP B372 3	del Aguilua+, (CERN, GRAN, MPIM, BRUKT, MADE)
Also 91C NP B375 45	del Aguilua, Moreno, Quiros (CERN, BARI)
DELAGUILA 92B NP B381 451	del Aguilua, Martinez, Quiros (GRAN, CERN)
ELLIS 92 PL B274 456	+Fogli, Lisi (CERN, BARI)
ELLIS 92E PL B292 427	+Fogli, Lisi (CERN, BARI)
LANGACKER 92B PR D45 278	+Luo (PENN)
LEP 92 PL B276 247	+ALEPH, DELPHI, L3, OPAL (LEP Collab.)
PDG 92 PR D45, 1 June, Part II	Hikasa, Barnett, Stone+ (KEK, LBL, BOST+)
REYON 92 ZPH C56 325	
SCHAILE 92 ZPHY C54 387	(FREIE)
ABE 91 PR D43 664	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE 91B PR D43 2070	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE 91C PR D44 29	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABREU 91F NP B367 511	+Adam, Adams, Adye, Akesson+ (DELPHI Collab.)
ADEVA 91E ZPHY C51 179	+Adriani, Aguilera-Benitez, Akbari, Alcaraz+ (L3 Collab.)
ALBAJAR 91B PL B253 803	+Albrow, Allkofer, Ankovak, Apšimon+ (UA1 Collab.)
ALBAJAR 91B PR D43 2421	+Albrow, Allkofer, Ankovak, Apšimon+ (UA1 Collab.)
ALEXANDER 91F ZPHY C52 175	+Allison, Allport, Anderson, Arcelli+ (OPAL Collab.)
BAER 91B PR D44 725	+Drees, Godbole+ (FSU, DESY, BOMB, UCD, HAWA)
DELAGUILA 91C NP B361 45	del Aguilua, Moreno, Quiros (BARC, MADE)
GONZALEZ-G. 91 PL B259 365	Gonzalez-Garcia, Valle (VALE)
HIOKI 91 MPL A6 2129	(TOKU)
LANGACKER 91 PR D44 817	+Luo (PENN)
ABE 90 PR L52	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE 90C PRL 64 147	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE 90B PRL 64 142	+Amidei, Apollinari, Atac+ (CDF Collab.)
Also 91 PR D43 664	Abe, Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE 90D PL B234 382	+Amako, Arai, Asano+ (VENUS Collab.)
ABE 90G PRL 65 2243	+Amidei, Apollinari, Atac+ (CDF Collab.)
ABREU 90D PL B242 536	+Adam, Adams, Adye, Alekseev, Allaby+ (DELPHI Collab.)
ADACHI 90F PL B234 525	+Doser, Enomoto, Fujii+ (TOPAZ Collab.)
ADEVA 90I PL B249 341	+Adriani, Aguilera-Benitez, Akbari, Alcaraz+ (L3 Collab.)
AKESSON 90 ZPHY C46 179	+Alitti, Ansari, Ansoerg, Bagnaia+ (UA2 Collab.)
AKRAWY 90B PL B236 364	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
ALBAJAR 90B ZPHY C48 1	+Albrow, Allkofer, Andrieu, Ankovak+ (UA1 Collab.)
ALITTI 90C PL B241 150	+Ansari, Ansoerg, Autiero+ (UA2 Collab.)
ALITTI 90B ZPHY C47 11	+Ansari, Ansoerg, Bagnaia+ (UA2 Collab.)
BARGER 90C PRL 65 1313	+Hewett, Rizzo (WISC, ISU)
BARGER 90E PR D41 3421	+Hewett, Phillips (WISC, ILL)
BLONDEL 90 ZPHY C46 361	+Boeckmann, Burkhardt, Dydak, Grant+ (CDHSW Collab.)
DECAMP 90F PL B236 511	+Deschizeaux, Lees, Minard+ (ALEPH Collab.)
DECAMP 90B ZPHY C48 365	+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
ELLIS 90B PL B249 543	+Fogli (CERN)
KENNEDY 90 PRL 65 2967	+Langacker (PENN)
ABRAMS 89C PRL 63 2447	+Adolphsen, Averill, Ballam+ (Mark II Collab.)
ADACHI 89C PL B229 427	+Aihara, Doser, Enomoto, Fujii+ (TOPAZ Collab.)
ALBAJAR 89B PL D42 15	+Albrow, Allkofer, Arnison, Astbury+ (UA1 Collab.)
ELLIS 89B PL B232 139	+Fogli (CERN, BARI)
ENO 89 PRL 63 1910	+Auchincloss, Blanis, Bodek, Budd+ (AMY Collab.)
LANGACKER 89 PR L63 1920	(PENN)
ADACHI 88 PRL 60 97	+Aihara, Dijkstra+ (TOPAZ Collab.)
ALBAJAR 88 ZPHY C37 505	+Albrow, Allkofer+ (UA1 Collab.)
ALTARELLI 88 NP B308 727	+Diemoz, Martinelli, Nason (CERN, ROMA, ETH)
COSTA 88 NP B287 244	+Elli, Fogli+ (PADO, CERN, BARI, WISC, LBL)
ELLIS 88C PL B213 526	+Fogli (CERN, BARI)
FOGLI 88 ZPHY C40 379	+Haidt (BARI, DESY)
IGARASHI 88 PRL 60 2359	+Myung, Chiba, Hanaoka+ (AMY Collab.)
SAGAWA 88 PRL 60 93	+Mori, Abe+ (AMY Collab.)
ABE 87 JPSJ 56 3763	+Amako, Arai+ (VENUS Collab.)
AMALDI 87 PR D36 1385	+Bohm, Durkin, Langacker+ (CERN, AACHS, OSU+)
ANSARI 87 PL B186 440	+Bagnaia, Garner, Battiston+ (UA2 Collab.)
YOSHIDA 87 PL B198 570	+Chiba, Endo+ (VENUS Collab.)

Lepton & Quark Full Listings

b' (Fourth Generation) Quark

Searches for b' (4^{th} Generation) Quark

MASS LIMITS for b' (4^{th} Generation) Quark or Hadron in $p\bar{p}$ Collisions

These experiments (except for MUKHOPADHYAYA 93) assume that no two-body modes such as $b' \rightarrow b\gamma$, $b' \rightarrow bg$, or $b' \rightarrow cH^+$ are available.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>85	95	¹ ABE	92 CDF	$\ell\ell$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>75	95	² MUKHOPAD... 93	RVUE	FCNC
>72	95	³ ABE	90B CDF	$e + \mu$
>54	95	⁴ AKESSON	90 UA2	$e + \text{jets} + \text{missing } E_T$
>43	95	⁵ ALBAJAR	90B UA1	$\mu + \text{jets}$
>34	95	⁶ ALBAJAR	88 UA1	$e \text{ or } \mu + \text{jets}$

- ¹ ABE 92 dilepton analysis limit of >85 GeV at CL=95% also applies to b' quarks, as discussed in ABE 90B.
- ² MUKHOPADHYAYA 93 analyze CDF dilepton data of ABE 92G in terms of a new quark decaying via flavor-changing neutral current. The above limit assumes $B(b' \rightarrow b\ell^+\ell^-)=1\%$. For an exotic quark decaying only via virtual Z [$B(b\ell^+\ell^-)=3\%$], the limit is 85 GeV.
- ³ ABE 90B exclude the region 28–72 GeV.
- ⁴ AKESSON 90 searched for events having an electron with $p_T > 12$ GeV, missing momentum > 15 GeV, and a jet with $E_T > 10$ GeV, $|\eta| < 2.2$, and excluded $m_{b\ell}$ between 30 and 69 GeV.
- ⁵ For the reduction of the limit due to non-charged-current decay modes, see Fig. 19 of ALBAJAR 90B.
- ⁶ ALBAJAR 88 study events at $E_{cm} = 546$ and 630 GeV with a muon or isolated electron, accompanied by one or more jets and find agreement with Monte Carlo predictions for the production of charm and bottom, without the need for a new quark. The lower mass limit is obtained by using a conservative estimate for the $b'\bar{b}'$ production cross section and by assuming that it cannot be produced in W decays. The value quoted here is revised using the full $O(\alpha_s^3)$ cross section of ALTARELLI 88.

MASS LIMITS for b' (4^{th} Generation) Quark or Hadron in e^+e^- Collisions

Search for hadrons containing a fourth-generation $-1/3$ quark denoted b' .

The last column specifies the assumption for the decay mode (C denotes the conventional charged-current decay) and the event signature which is looked for.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>46.0	95	⁷ DECAMP	90F ALEP	any decay
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>44.7	95	⁸ ADRIANI	93G L3	Quarkonium
>45	95	ADRIANI	93M L3	$\Gamma(Z)$
none 19.4–28.2	95	ABREU	91F DLPH	$\Gamma(Z)$
>45.0	95	ABE	90D VNS	Any decay; event shape
>44.5	95	ABREU	90D DLPH	$B(C) = 1$; event shape
>44.5	95	⁹ ABREU	90D DLPH	$b' \rightarrow cH^-, H^- \rightarrow \bar{c}s, \tau^- \nu$
>40.5	95	¹⁰ ABREU	90D DLPH	$\Gamma(Z \rightarrow \text{hadrons})$
>28.3	95	ADACHI	90 TOPZ	$B(\text{FCNC})=100\%$; isol. γ or 4 jets
>41.4	95	¹¹ AKRAWY	90B OPAL	Any decay; acoplanarity
>45.2	95	¹¹ AKRAWY	90B OPAL	$B(C) = 1$; acoplanarity
>46	95	¹² AKRAWY	90J OPAL	$b' \rightarrow \gamma + \text{any}$
>27.5	95	¹³ ABE	89E VNS	$B(C) = 1; \mu, e$
none 11.4–27.3	95	¹⁴ ABE	89G VNS	$B(b' \rightarrow b\gamma) > 10\%$; isolated γ
>44.7	95	¹⁵ ABRAMS	89C MRK2	$B(C) = 100\%$; isol. track
>42.7	95	¹⁵ ABRAMS	89C MRK2	$B(bg) = 100\%$; event shape
>42.0	95	¹⁵ ABRAMS	89C MRK2	Any decay; event shape
>28.4	95	^{16,17} ADACHI	89C TOPZ	$B(C) = 1; \mu$
>28.8	95	¹⁸ ENO	89 AMY	$B(C) \gtrsim 90\%; \mu, e$
>27.2	95	^{18,19} ENO	89 AMY	any decay; event shape
>29.0	95	¹⁸ ENO	89 AMY	$B(b' \rightarrow bg) \gtrsim 85\%$; event shape
>24.4	95	²⁰ IGARASHI	88 AMY	μ, e
>23.8	95	²¹ SAGAWA	88 AMY	event shape
>22.7	95	²² ADEVA	86 MRKJ	μ
>21	95	²³ ALTHOFF	84C TASS	R , event shape
>19	95	²⁴ ALTHOFF	84I TASS	Aplanarity

⁷ DECAMP 90F looked for isolated charged particles, for isolated photons, and for four-jet final states. The modes $b' \rightarrow bg$ for $B(b' \rightarrow bg) > 65\%$ $b' \rightarrow b\gamma$ for $B(b' \rightarrow b\gamma) > 5\%$ are excluded. Charged Higgs decay were not discussed.

⁸ ADRIANI 93G search for vector quarkonium states near Z and give limit on quarkonium- Z mixing parameter $\delta m^2 < (10-30) \text{ GeV}^2$ (95%CL) for the mass 88–94.5 GeV. Using Richardson potential, a $1S$ ($b'\bar{b}'$) state is excluded for the mass range 87.7–94.7 GeV. This range depends on the potential choice.

⁹ ABREU 90D assumed $m_{H^-} < m_{b'} - 3 \text{ GeV}$.

¹⁰ Superseded by ABREU 91F.

¹¹ AKRAWY 90B search was restricted to data near the Z peak at $E_{cm} = 91.26 \text{ GeV}$ at LEP. The excluded region is between 23.6 and 41.4 GeV if no H^+ decays exist. For charged Higgs decays the excluded regions are between $(m_{H^+} + 1.5 \text{ GeV})$ and 45.5 GeV.

¹² AKRAWY 90J search for isolated photons in hadronic Z decay and derive $B(Z \rightarrow b'\bar{b}') \cdot B(b' \rightarrow \gamma X) / B(Z \rightarrow \text{hadrons}) < 2.2 \times 10^{-3}$. Mass limit assumes $B(b' \rightarrow \gamma X) > 10\%$.

¹³ ABE 89E search at $E_{cm} = 56-57 \text{ GeV}$ at TRISTAN for multihadron events with a spherical shape (using thrust and acoplanarity) or containing isolated leptons.

¹⁴ ABE 89G search was at $E_{cm} = 55-60.8 \text{ GeV}$ at TRISTAN.

¹⁵ If the photonic decay mode is large ($B(b' \rightarrow b\gamma) > 25\%$), the ABRAMS 89C limit is 45.4 GeV. The limit for for Higgs decay ($b' \rightarrow cH^-, H^- \rightarrow \bar{c}s$) is 45.2 GeV.

¹⁶ ADACHI 89C search was at $E_{cm} = 56.5-60.8 \text{ GeV}$ at TRISTAN using multi-hadron events accompanying muons.

¹⁷ ADACHI 89C also gives limits for any mixture of CC and bg decays.

¹⁸ ENO 89 search at $E_{cm} = 50-60.8$ at TRISTAN.

¹⁹ ENO 89 considers arbitrary mixture of the charged current, bg , and $b\gamma$ decays.

²⁰ IGARASHI 88 searches for leptons in low-thrust events and gives $\Delta R(b') < 0.26$ (95% CL) assuming charged current decay, which translates to $m_{b'} > 24.4 \text{ GeV}$.

²¹ SAGAWA 88 set limit $\sigma(\text{top}) < 6.1 \text{ pb}$ at CL=95% for top-flavored hadron production from event shape analyses at $E_{cm} = 52 \text{ GeV}$. By using the quark parton model cross-section formula near threshold, the above limit leads to lower mass bounds of 23.8 GeV for charge $-1/3$ quarks.

²² ADEVA 86 give 95%CL upper bound on an excess of the normalized cross section, ΔR , as a function of the minimum c.m. energy (see their figure 3). Production of a pair of $1/3$ charge quarks is excluded up to $E_{cm} = 45.4 \text{ GeV}$.

²³ ALTHOFF 84C narrow state search sets limit $\Gamma(e^+e^-B(\text{hadrons})) < 2.4 \text{ keV}$ CL = 95% and heavy charge $1/3$ quark pair production $m > 21 \text{ GeV}$, CL = 95%.

²⁴ ALTHOFF 84I exclude heavy quark pair production for $7 < m < 19 \text{ GeV}$ ($1/3$ charge) using aplanarity distributions (CL = 95%).

REFERENCES FOR Searches for (Fourth Generation) b' Quark

ADRIANI	93G	PL B313 326	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+ (L3 Collab.)
ADRIANI	93M	PR D 236 1	+Aguilar-Benitez, Ahlen, Alcaraz, Aloisio+ (L3 Collab.)
MUKHOPAD... 93	PR	D48 2105	Mukhopadhyaya, Roy (TATA)
ABE	92	PRL 68 447	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
Also	92G	PR D45 3921	Abe, Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE	92G	PR D45 3921	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABREU	91F	NP B367 511	+Adam, Adams, Adaye, Akesson+ (DELPHI Collab.)
ABE	90B	PRL 64 147	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE	90D	PL B234 382	+Amako, Arai, Asano+ (VENUS Collab.)
ABREU	90D	PL B242 536	+Adam, Adams, Adaye, Alekseev, Allaby+ (DELPHI Collab.)
ADACHI	90	PL B234 197	+Aihara, Doser, Enomoto+ (TOPAZ Collab.)
AKESSON	90	ZPHY C46 179	+Alitti, Ansari, Ansorge, Bagnaia+ (UA2 Collab.)
AKRAWY	90B	PL B236 364	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
AKRAWY	90J	PL B246 285	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
ALBAJAR	90B	ZPHY C48 1	+Albrow, Altkofer, Andrieu, Ankoviak+ (UA1 Collab.)
DECAMP	90F	PL B234 511	+Deschizeaux, Lees, Minard+ (ALEPH Collab.)
ABE	89E	PR D39 3524	+Amako, Arai, Asano, Chiba, Chiba+ (VENUS Collab.)
ABE	89G	PRL 63 1776	+Amako, Arai, Asano, Chiba+ (VENUS Collab.)
ABRAMS	89C	PRL 63 2447	+Adolphsen, Averil, Ballam+ (Mark II Collab.)
ADACHI	89C	PL B229 427	+Aihara, Doser, Enomoto, Fujii+ (TOPAZ Collab.)
ENO	89	PRL 63 1910	+Auchincloss, Blanis, Bodek, Budd+ (AMY Collab.)
ALBAJAR	88	ZPHY C37 505	+Albrow, Altkofer+ (UA1 Collab.)
ALTARELLI	88	NP B308 724	+Dilemoz, Martinelli, Nason (CERN, ROMA, ETH)
IGARASHI	88	PRL 60 2359	+Myung, Chiba, Hanaoka+ (AMY Collab.)
SAGAWA	88	PRL 60 93	+Mori, Abe+ (AMY Collab.)
ADEVA	86	PR D34 681	+Ansari, Becker, Becker-Szendy+ (Mark-J Collab.)
ALTHOFF	84C	PL 138B 441	+Braunschweig, Kirschfink+ (TASSO Collab.)
ALTHOFF	84I	ZPHY C22 307	+Braunschweig, Kirschfink+ (TASSO Collab.)

LIGHT UNFLAVORED MESONS ($S = C = B = 0$)

- π^\pm 1444
- π^0 1448
- η 1450
- $\rho(770)$ 1456
- $\omega(782)$ 1459
- $\eta'(958)$ 1461
- $f_0(980)$ 1464
- $a_0(980)$ 1465
- $\phi(1020)$ 1467
- $h_1(1170)$ 1469
- $b_1(1235)$ 1470
- $a_1(1260)$ 1471
- $f_2(1270)$ 1472
- $f_1(1285)$ 1475
- $\eta(1295)$ 1478
- $f_0(1300)$ 1478
- $\pi(1300)$ 1480
- $a_2(1320)$ 1480
- $f_0(1370)$ 1483
- $h_1(1380)$ 1483
- $\rho(1405)$ 1484
- $f_1(1420)$ 1484
- $\omega(1420)$ 1486
- $f_2(1430)$ 1486
- $\eta(1440)$ 1487
- $\rho(1450)$ 1489
- $f_1(1510)$ 1490
- $f_2(1520)$ 1490
- $f_2'(1525)$ 1491
- $f_0(1525)$ 1493
- $f_0(1590)$ 1493
- $\omega(1600)$ 1494
- $X(1600)$ 1494
- $f_2(1640)$ 1494
- $\omega_3(1670)$ 1495
- $\pi_2(1670)$ 1495
- $\phi(1680)$ 1497
- $\rho_3(1690)$ 1497
- $\rho(1700)$ 1501
- $X(1700)$ 1504
- $f_J(1710)$ 1504
- $X(1740)$ 1505
- $\eta(1760)$ 1506
- $\pi(1770)$ 1506
- $X(1775)$ 1506
- $f_2(1810)$ 1506
- $X(1830)$ 1507
- $\phi_3(1850)$ 1507
- $\eta_2(1870)$ 1508
- $X(1910)$ 1508
- $X(1950)$ 1509

- $f_2(2010)$ 1509
- $a_4(2040)$ 1509
- $a_3(2050)$ 1510
- $f_4(2050)$ 1510
- $\pi_2(2100)$ 1511
- $f_2(2150)$ 1511
- $\rho(2150)$ 1512
- $X(2200)$ 1512
- $\rho(2210)$ 1512
- $f_4(2220)$ 1513
- $\eta(2225)$ 1513
- $\rho_3(2250)$ 1514
- $f_2(2300)$ 1514
- $f_4(2300)$ 1514
- $f_2(2340)$ 1515
- $\rho_5(2350)$ 1515
- $a_6(2450)$ 1516
- $f_6(2510)$ 1516
- $X(3250)$ 1516

OTHER LIGHT UNFLAVORED ($S = C = B = 0$)

- $e^+e^-(1100-2200)$ 1517
- $\bar{N}N(1100-3600)$ 1517
- $X(1900-3600)$ 1519

STRANGE MESONS ($S = \pm 1, C = B = 0$)

- K^\pm 1521
- K^0 1534
- K_S^0 1534
- K_L^0 1537
- $K^*(892)$ 1549
- $K_1(1270)$ 1551
- $K_1(1400)$ 1552
- $K^*(1410)$ 1553
- $K_0^*(1430)$ 1554
- $K_2^*(1430)$ 1554
- $K(1460)$ 1556
- $K_2(1580)$ 1557
- $K_1(1650)$ 1557
- $K^*(1680)$ 1557
- $K_2(1770)$ 1558
- $K_3^*(1780)$ 1559
- $K_2(1820)$ 1560
- $K(1830)$ 1560
- $K_0^*(1950)$ 1560
- $K_2^*(1980)$ 1561
- $K_4^*(2045)$ 1561
- $K_2(2250)$ 1562
- $K_3(2320)$ 1562
- $K_5^*(2380)$ 1562
- $K_4(2500)$ 1562
- $K(3100)$ 1563

(continued on the next page)

• Indicates the particle is in the Meson Summary Table

CHARMED MESONS ($C = \pm 1$)

• D^\pm	1572
• D^0	1581
• $D^*(2007)^0$	1593
• $D^*(2010)^+$	1593
• $D_1(2420)^0$	1594
• $D_J(2440)^\pm$	1595
• $D_2^*(2460)$	1595

CHARMED, STRANGE MESONS ($C = S = \pm 1$)

• D_s^\pm	1596
• $D_s^{*\pm}$	1599
• $D_{s1}(2536)^\pm$	1600
• $D_{sJ}(2573)^\pm$	1600

BOTTOM MESONS ($B = \pm 1$)

• B^\pm	1609
• B^0	1623
• B^*	1639

BOTTOM, STRANGE MESONS ($B = \pm 1, S = \mp 1$)

• B_s^0	1639
• B_s^*	1641

 $c\bar{c}$ MESONS

• $\eta_c(1S) = \eta_c(2980)$	1642
• $J/\psi(1S) = J/\psi(3097)$	1644
• $\chi_{c0}(1P) = \chi_{c0}(3415)$	1652
• $\chi_{c1}(1P) = \chi_{c1}(3510)$	1653
• $h_c(1P)$	1654
• $\chi_{c2}(1P) = \chi_{c2}(3555)$	1654
• $\eta_c(2S) = \eta_c(3590)$	1656
• $\psi(2S) = \psi(3685)$	1656
• $\psi(3770)$	1659
• $\psi(4040)$	1659
• $\psi(4160)$	1660
• $\psi(4415)$	1660

 $b\bar{b}$ MESONS

• $\Upsilon(1S) = \Upsilon(9460)$	1662
• $\chi_{b0}(1P) = \chi_{b0}(9860)$	1664
• $\chi_{b1}(1P) = \chi_{b1}(9890)$	1664
• $\chi_{b2}(1P) = \chi_{b2}(9915)$	1665
• $\Upsilon(2S) = \Upsilon(10023)$	1665
• $\chi_{b0}(2P) = \chi_{b0}(10235)$	1666
• $\chi_{b1}(2P) = \chi_{b1}(10255)$	1667
• $\chi_{b2}(2P) = \chi_{b2}(10270)$	1667
• $\Upsilon(3S) = \Upsilon(10355)$	1668
• $\Upsilon(4S) = \Upsilon(10580)$	1669
• $\Upsilon(10860)$	1669
• $\Upsilon(11020)$	1670

NON- $q\bar{q}$ CANDIDATES

Non- $q\bar{q}$ Candidates	1670
----------------------------	------

Notes in the Meson Listings

Note on Pseudoscalar-Meson Decay Constants	1443
Note on $\pi^\pm \rightarrow \ell^\pm \nu \gamma$ and $K^\pm \rightarrow \ell^\pm \nu \gamma$ Form Factors	1447
Note on the Charged Pion Mass	1444
Note on the Decay Width $\Gamma(\eta \rightarrow \gamma\gamma)$	1451
Note on η Decay Parameters	1454
Note on the $\rho(770)$	1456
Note on the $a_0(980)$	1465
Note on the $a_1(1260)$	1471
Note on S -wave $\pi\pi$, $K\bar{K}$, and $\eta\eta$ Interactions	1478
Note on the $f_1(1420)$	1484
Note on the $\eta(1440)$	1487
Note on the $\rho(1450)$ and the $\rho(1700)$	1501
Note on the $f_J(1710)$	1504
Note on the $X(1900-3600)$ Region	1519
Note on K^\pm Mass	1521
Note on Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays	1528
Note on $K_{\ell 3}^\pm$ and $K_{\ell 3}^0$ Form Factors	1530
Note on CP Violation in $K_S \rightarrow 3\pi$	1536
Note on CP Violation in K_L^0 Decay	1543
Note on $\Delta S = \Delta Q$ in K^0 Decays	1547
Note on $K^*(892)$ Masses and Mass Differences	1550
Note on $K_2(1770)$ and the $K_2(1820)$	1558
Note on D Meson Branching Fractions	1564
Note on Semileptonic Decays of D and B Mesons, Part I	1565
Note on Experimental Highlights of B Meson Production and Decay	1601
Note on Semileptonic Decays of D and B Mesons, Part II	1602
$B^0-\bar{B}^0$ Mixing and CP Violation in B Decay	1632
Note on Width Determinations of the Υ States	1661
Note on Non- $q\bar{q}$ Mesons	1670

• Indicates the particle is in the Meson Summary Table

LIGHT UNFLAVORED MESONS ($S = C = B = 0$)

For $l = 1$ (π, b, ρ, a): $u\bar{d}, (u\bar{u}-d\bar{d})/\sqrt{2}, d\bar{u}$;
for $l = 0$ ($\eta, \eta', h, h', \omega, \phi, f, f'$): $c_1(u\bar{u} + d\bar{d}) + c_2(s\bar{s})$

NOTE ON PSEUDOSCALAR-MESON DECAY CONSTANTS

(by M. Suzuki, Lawrence Berkeley Laboratory)

Charged mesons

The decay constant f_P for pseudoscalar meson P is defined by

$$\langle 0 | A_\mu(0) | P(\mathbf{q}) \rangle = i f_P q_\mu ,$$

where A_μ is the axial-vector part of the charged weak current after a Cabibbo-Kobayashi-Maskawa mixing-matrix element $V_{qq'}$ has been removed. The state vector is normalized by $\langle P(\mathbf{q}) | P(\mathbf{q}') \rangle = (2\pi)^3 2E_q \delta(\mathbf{q} - \mathbf{q}')$, and its phase is chosen to make f_P real and positive. Note, however, that in many theoretical papers our $f_P/\sqrt{2}$ is denoted by f_P .

In determining f_P experimentally, radiative corrections must be taken into account. Since the photon-loop correction introduces an infrared divergence that is canceled by soft-photon emission, we can determine f_P only from the combined rate for $P^\pm \rightarrow \ell^\pm \nu_\ell$ and $P^\pm \rightarrow \ell^\pm \nu_\ell \gamma$. This rate is given by

$$\Gamma(P \rightarrow \ell \nu_\ell + \ell \nu_\ell \gamma) = \frac{G_F^2 |V_{qq'}|^2}{8\pi} f_P^2 m_\ell^2 m_P \left(1 - \frac{m_\ell^2}{m_P^2} \right)^2 [1 + \mathcal{O}(\alpha)] .$$

Radiative corrections contain an inner bremsstrahlung, which is independent of the structure of the meson [1-3], and a structure-dependent part [4,5]. After radiative corrections are made, there are ambiguities in extracting f_P from experimental measurements. In fact, the definition of f_P is no longer unique.

It is desirable to define f_P such that it depends only on the properties of the pseudoscalar meson, not on the final decay products. The short-distance corrections to the fundamental electroweak constants like $G_F |V_{qq'}|$ should be separated out. Following Marciano and Sirlin [6], we define f_P with the following form for the $\mathcal{O}(\alpha)$ corrections:

$$1 + \mathcal{O}(\alpha) = \left[1 + \frac{2\alpha}{\pi} \ln \left(\frac{m_\rho}{m_P} \right) \right] \left[1 + \frac{\alpha}{\pi} F(x) \right] \times \left\{ 1 - \frac{\alpha}{\pi} \left[\frac{3}{2} \ln \left(\frac{m_\rho}{m_P} \right) + C_1 + C_2 \frac{m_\ell^2}{m_\rho^2} \ln \left(\frac{m_\rho^2}{m_\ell^2} \right) + C_3 \frac{m_\ell^2}{m_\rho^2} + \dots \right] \right\} .$$

Here

$$F(x) = 3 \ln x + \frac{13 - 19x^2}{8(1-x^2)} - \frac{8 - 5x^2}{2(1-x^2)^2} x^2 \ln x - 2 \left(\frac{1+x^2}{1-x^2} \ln x + 1 \right) \ln(1-x^2) + 2 \left(\frac{1+x^2}{1-x^2} \right) L(1-x^2) ,$$

with

$$x \equiv m_\ell/m_P , \quad L(z) \equiv \int_0^z \frac{\ln(1-t)}{t} dt .$$

The first bracket in the expression for $1 + \mathcal{O}(\alpha)$ is the short-distance electroweak correction. The QCD correction reduces this factor by 0.00033. The second bracket together with the term $-(3\alpha/2\pi) \ln(m_\rho/m_P)$ in the third bracket corresponds to the radiative corrections to the point-like pion decay ($\Lambda_{\text{cutoff}} \approx m_\rho$) [2]. The rest of the corrections in the third bracket are expanded in power of m_ℓ/m_ρ . The expansion coefficients C_1 , C_2 , and C_3 depend on the hadronic structure of the pseudoscalar meson and in most cases cannot be computed accurately. In particular, C_1 absorbs the uncertainty in the matching energy scale between short- and long-distance strong interactions and thus is the main source of uncertainty in determining f_{π^+} accurately.

With the experimental value for the decay $\pi \rightarrow \mu \nu_\mu + \mu \nu_\mu \gamma$, one obtains

$$f_{\pi^+} = 130.7 \pm 0.1 \pm 0.36 \text{ MeV} ,$$

where the first error comes from the experimental uncertainty on $|V_{ud}|$ and the second comes from the uncertainty on C_1 ($= 0 \pm 0.24$) [6]. Similarly, one obtains from the decay $K \rightarrow \mu \nu_\mu + \mu \nu_\mu \gamma$ the decay constant

$$f_{K^+} = 159.8 \pm 1.4 \pm 0.44 \text{ MeV} ,$$

where the first error is due to the uncertainty on $|V_{us}|$.

For the heavy pseudoscalar mesons, uncertainties in the experimental values for the decay rates are much larger than the radiative corrections. For the D^+ , only an upper bound can be obtained from the published data:

$$f_{D^+} < 310 \text{ MeV (CL = 90\%)} .$$

Two groups have measured the $D_s^+ \rightarrow \mu^+ \nu_\mu$ branching fraction, leading to the following values of the decay constant:

$$f_{D_s^+} = 232 \pm 45 \pm 20 \pm 48 \text{ MeV [7]} ,$$

$$f_{D_s^+} = 344 \pm 37 \pm 52 \pm 42 \text{ MeV [8]} ,$$

where the first errors are statistical, the second errors are systematic, and the third errors are uncertainties involved in extracting the branching fraction $B(D_s^+ \rightarrow \mu^+ \nu_\mu)$.

There have been many attempts to extract f_P from spectroscopy and nonleptonic decays using theoretical models. Since it is difficult to estimate uncertainties for them, we have listed here only values of decay constants that are obtained directly from the observation of $P^\pm \rightarrow \ell^\pm \nu_\ell$.

Light neutral mesons

The decay constants for the light neutral pseudoscalar mesons π^0 , η , and η' are defined by

$$\langle 0 | A_\mu(0) | P^0(\mathbf{q}) \rangle = i(f_P/\sqrt{2})q_\mu ,$$

Meson Full Listings

 π^\pm

where A_μ is a neutral axial-vector current of octet or singlet. Values of f_P can be obtained from the two-photon decay $P^0 \rightarrow \gamma\gamma$, since in the $m_P = 0$ limit the decay matrix element is determined by the Adler-Bell-Jackiw anomaly [9,10]. However, large uncertainties enter values of f_P through extrapolation to the physical mass and, in the case of η and η' , through the mixing angle, too.

The CELLO Collaboration has obtained the values [11]

$$f_{\pi^0} = 119 \pm 4 \text{ MeV}$$

$$f_\eta = 133 \pm 10 \text{ MeV}$$

$$f_{\eta'} = 126 \pm 7 \text{ MeV} ,$$

while the TPC/2 γ Collaboration has obtained [12]

$$f_\eta = 129 \pm 8 \text{ MeV}$$

$$f_{\eta'} = 110 \pm 7 \text{ MeV} .$$

(We have multiplied the published values by $\sqrt{2}$ to be in accord with our definition of f_P .)

References

1. S. Berman, Phys. Rev. Lett. **1**, 468 (1958).
2. T. Kinoshita, Phys. Rev. Lett. **2**, 477 (1959).
3. A. Sirlin, Phys. Rev. **D5**, 436 (1972).
4. M.V. Terent'ev, Yad. Fiz. **18**, 870 (1973) [Sov. J. Nucl. Phys. **18**, 449 (1974)].
5. T. Goldman and W.J. Wilson, Phys. Rev. **D15**, 709 (1977).
6. W.J. Marciano and A. Sirlin, Phys. Rev. Lett. **71**, 3629 (1993).
7. S. Aoki *et al.*, Prog. Theor. Phys. **89**, 131 (1993).
8. D. Acosta *et al.*, CLNS 93/1238, CLEO-93-14 (to be published).
9. S.L. Adler, Phys. Rev. **177**, 2426 (1969).
10. J.S. Bell and R. Jackiw, Nuovo Cimento **60A**, 46 (1969).
11. H.-J. Behrend *et al.*, Z. Phys. **C49**, 401 (1991).
12. H. Aihara *et al.*, Phys. Rev. Lett. **64**, 172 (1990).

 π^\pm

$$I^G(J^P) = 1^-(0^-)$$

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1988 edition Physics Letters **B204** (1988).

NOTE ON THE CHARGED PION MASS*

The charged pion mass we give in this edition is three (old) standard deviations higher than the one we gave in our 1992 edition. This Note explains why.

Precise determinations of the charged pion mass come from two kinds of experiments: (a) measurement of the momentum of the muon in $\pi^+ \rightarrow \mu^+ \nu_\mu$ decay at rest, assuming $m_{\nu_\mu} = 0$; and (b) measurement of x-ray wavelengths of transitions in pionic atoms. Technique (b) is free of assumptions about m_{ν_μ} .

For technique (b), the pionic-atom transition is chosen to achieve maximal spectrometer performance and minimal theoretical uncertainty. Experimenters have used the wavelengths of the $4f-3d$ transition in magnesium and the $5f-4d$ transition in titanium and phosphorous. Screening by K-shell ($1s$) electrons is significant, and the observed line profile is a blend of lines from atoms with 0, 1, and 2 K electrons present during the transition.

The best pionic-atom measurement of the pion mass in our recent editions was that of JECKELMANN 86B. Their observed line profile was measurably broadened by contributions from atoms having different $1s$ populations. Fits to the line profile gave two equally good solutions: One, with probabilities of about 27%, 70%, and 3% for 0, 1, or 2 K-shell electrons; and another, in which the state with two K electrons dominated. Using the ratio of $4f-3d$ and $3d-2p$ intensities and other experimental information, JECKELMANN 86B rejected the second solution.

Recent measurements now yield a smaller value for the intensity ratio [1], so that in a reanalysis of the data JECKELMANN 94 now consider that the grounds for rejecting the second solution are marginal. The new analysis is improved in other respects as well, including use of an improved wavelength calibration standard. The situation is now considered to be ambiguous, with both of the following solutions given equal footing:

$$\text{Solution A: } m_{\pi^-} = 139.56782 \pm 0.00037 \text{ MeV}$$

$$\text{Solution B: } m_{\pi^-} = 139.56995 \pm 0.00035 \text{ MeV}$$

The two solutions differ by about six times the uncertainty in either one. The ambiguity is expected to be resolved by future pionic x-ray measurements involving lighter atoms, where it is known with fair certainty that the K shell is almost empty during the pionic transition of interest. Such measurements on pionic oxygen and nitrogen at a pressure of a few bars have begun at the Paul Scherrer Institute [2]. We now give both Solutions A and B in the Listings below.

In the previous edition, we reported five π^\pm mass measurements with uncertainties smaller than 0.005 MeV. These, together with the results from the new analysis, are shown in Fig. 1. Two of them, CARTER 76 and MARUSHENKO 77, have such large errors that there is no conflict with either A or B. JECKELMANN 86B was basically Solution A above, with a small difference (it was 0.8σ lower) because of the new analysis and revised wavelength calibration. DAUM 91 was based on technique (a), a measurement of the muon momentum from π^+ decay at rest. Since their reported mass was obtained by assuming $m_{\nu_\mu} = 0$, it is really a lower limit. It is in good agreement with Solution B but disagrees with Solution A, as we discuss further below. LU 80 is compatible with Solution A but not with Solution B or with DAUM 91. A screening correction used by LU 80 came from a theoretical calculation based on a 1961 paper by Eisenberg and Kessler [3]. It would now appear that the new intensity-ratio measurements call into doubt some of the earlier input assumptions.

The DAUM 91 measurement of the muon momentum from π^+ decay at rest has been further refined, and Assamagan *et al.* [4] report $p_\mu = 29.79207 \pm 0.00012$ MeV/c (this work has been submitted for publication). This result is close to their earlier value, but the error is much smaller. When this result is combined with Solution A, one obtains a value for $m_{\nu_\mu}^2$ that is negative by six standard deviations. By contrast, with Solution B, $m_{\nu_\mu}^2$ is negative by only about one standard deviation. (See the parallel discussion and graph in the m_{ν_μ} section of these Full Listings.)

On the grounds that $m_{\nu_\mu}^2$ ought to be nonnegative, we choose Solution B of JECKELMANN 94 as the π^\pm mass. We do not average this with any other results, given the uncertainties. Obviously, however, the DAUM 91 (or Ref. 4) result determines our choice. Thus from the 1992 edition to this one, the mass has changed from 139.5679 ± 0.0007 MeV to 139.5695 ± 0.00035 MeV, a shift upwards of 0.00205 MeV, or three of the old standard deviations.

Since the π^0 mass comes from measurements of the π^\pm - π^0 mass difference, it too jumps upward.

Notes and References

- * This note was prepared with extensive help from F. Boehm, R. Frosch, P.F.A. Goudsmit, Y.K. Lee, H.J. Leisi, R.G.H. Robertson, and P. Vogel.
1. A. Shinohara *et al.*, Nucl. Instr. and Meth. **B84**, 14 (1994).
 2. R. Frosch, private communication (May 1994).
 3. Y. Eisenberg and D. Kessler, Nuovo Cimento **19**, 1196 (1961).
 4. K. Assamagan *et al.*, Paul Scherrer Institute preprint PSI-PR-94-19 (June 1994), submitted to Phys. Lett. **B**.




Figure 1: Published π^\pm mass measurements with uncertainties less than 0.005 MeV.

Meson Full Listings

 π^\pm π^\pm MASS

Measurements with an error > 0.005 MeV have been omitted from this Listing. The fit uses m_{π^\pm} and $(m_{\pi^\pm} - m_{\pi^0})$.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
139.56996 ± 0.00035 OUR FIT				
139.56996 ± 0.00035	1	JECKELMANN 94	CNTR	π^- atom, Soln. B
• • • We do not use the following data for averages, fits, limits, etc. • • •				
139.56782 ± 0.00037	1	JECKELMANN 94	CNTR	π^- atom, Soln. A
139.56996 ± 0.00067	2	DAUM 91	SPEC	$\pi^+ \rightarrow \mu^+ \nu$
139.56752 ± 0.00037	3	JECKELMANN 86B	CNTR	Mesonic atoms
139.5704 ± 0.0011	2	ABELA 84	SPEC	See DAUM 91
139.5664 ± 0.0009	4	LU 80	CNTR	Mesonic atoms
139.5686 ± 0.0020	CARTER 76	CNTR		Mesonic atoms
139.5660 ± 0.0024	4,5	MARUSHENKO 76	CNTR	Mesonic atoms

¹ See the above "Note on the Charged Pion Mass" for a discussion of the two JECKELMANN 94 values.

² The DAUM 91 value includes the ABELA 84 result. The value is based on a measurement of the μ^+ momentum for π^+ decay at rest, $p_\mu = 29.79179 \pm 0.00053$ MeV, uses $m_\mu = 105.658389 \pm 0.000034$ MeV, and assumes that $m_{\nu_\mu} = 0$. The last assumption means that in fact the value is a lower limit.

³ JECKELMANN 86B gives $m_\pi/m_e = 273.12677(71)$. We use $m_e = 0.51099906(15)$ MeV from COHEN 87. The authors note that two solutions for the probability distribution of K-shell occupancy fit equally well, and use other data to choose the lower of the two possible π^\pm masses.

⁴ These values are scaled with a new wavelength-energy conversion factor $V\lambda = 1.23984244(37) \times 10^{-6}$ eV m from COHEN 87. The LU 80 screening correction relies upon a theoretical calculation of inner-shell refilling rates.

⁵ This MARUSHENKO 76 value used at the authors' request to use the accepted set of calibration γ energies. Error increased from 0.0017 MeV to include QED calculation error of 0.0017 MeV (12 ppm).

 $m_{\pi^+} - m_{\mu^+}$

Measurements with an error > 0.05 MeV have been omitted from this Listing.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
33.91157 ± 0.00067	6	DAUM 91	SPEC	+	$\pi^+ \rightarrow \mu^+ \nu$
33.9111 ± 0.0011		ABELA 84	SPEC	+	See DAUM 91
33.925 ± 0.025		BOOTH 70	CNTR	+	Magnetic spect.
33.881 ± 0.035	145	HYMAN 67	HEBC	+	K^- He

⁶ The DAUM 91 value assumes that $m_{\nu_\mu} = 0$ and uses our $m_\mu = 105.658389 \pm 0.000034$ MeV.

 $(m_{\pi^+} - m_{\pi^-}) / m_{\text{average}}$

A test of CPT invariance.

VALUE (units 10^{-4})	DOCUMENT ID	TECN
2 ± 5	AYRES 71	CNTR

 π^\pm MEAN LIFE

Measurements with an error $> 0.02 \times 10^{-8}$ s have been omitted.

VALUE (10^{-8} s)	DOCUMENT ID	TECN	CHG
2.6030 ± 0.0024 OUR AVERAGE			
2.609 ± 0.008	DUNAITSEV 73	CNTR	+
2.602 ± 0.004	AYRES 71	CNTR	±
2.604 ± 0.005	NORDBERG 67	CNTR	+
2.602 ± 0.004	ECKHAUSE 65	CNTR	+
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.640 ± 0.008	7 KINSEY 66	CNTR	+

⁷ Systematic errors in the calibration of this experiment are discussed by NORDBERG 67.

 $(\tau_{\pi^+} - \tau_{\pi^-}) / \tau_{\text{average}}$

A test of CPT invariance.

VALUE (units 10^{-4})	DOCUMENT ID	TECN
5.5 ± 7.1	AYRES 71	CNTR
• • • We do not use the following data for averages, fits, limits, etc. • • •		
-14 ± 29	PETRUKHIN 68	CNTR
40 ± 70	BARDON 66	CNTR
23 ± 40	8 LOBKOWICZ 66	CNTR

⁸ This is the most conservative value given by LOBKOWICZ 66.

 π^+ DECAY MODES

π^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\mu^+ \nu_\mu$	[a] (99.98770 ± 0.00004) %	
Γ_2 $\mu^+ \nu_\mu \gamma$	[b] (1.24 ± 0.25) × 10 ⁻⁴	
Γ_3 $e^+ \nu_e$	[a] (1.230 ± 0.004) × 10 ⁻⁴	
Γ_4 $e^+ \nu_e \gamma$	[b] (1.61 ± 0.23) × 10 ⁻⁷	
Γ_5 $e^+ \nu_e \pi^0$	(1.025 ± 0.034) × 10 ⁻⁸	
Γ_6 $e^+ \nu_e e^+ e^-$	(3.2 ± 0.5) × 10 ⁻⁹	
Γ_7 $e^+ \nu_e \nu \bar{\nu}$	< 5	× 10 ⁻⁶ 90%

Lepton Family number (LF) or Lepton number (L) violating modes

Γ_8 $\mu^+ \bar{\nu}_e$	L	[c] < 1.5	× 10 ⁻³ 90%
Γ_9 $\mu^+ \nu_e$	LF	[c] < 8.0	× 10 ⁻³ 90%
Γ_{10} $\mu^- e^+ e^+ \nu$	LF	< 1.6	× 10 ⁻⁶ 90%

[a] Measurements of $\Gamma(e^+ \nu_e)/\Gamma(\mu^+ \nu_\mu)$ always include decays with γ 's, and measurements of $\Gamma(e^+ \nu_e \gamma)$ and $\Gamma(\mu^+ \nu_\mu \gamma)$ never include low-energy γ 's. Therefore, since no clean separation is possible, we consider the modes with γ 's to be subreactions of the modes without them, and let $[\Gamma(e^+ \nu_e) + \Gamma(\mu^+ \nu_\mu)]/\Gamma_{\text{total}} = 100\%$.

[b] See the Full Listings below for the energy limits used in this measurement; low-energy γ 's are not included.

[c] Derived from an analysis of neutrino-oscillation experiments.

 π^+ BRANCHING RATIOS

$\Gamma(e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_3/Γ
See note [a] in the list of π^+ decay modes just above, and see also the next block of data.

VALUE (units 10^{-4})	DOCUMENT ID
1.230 ± 0.004 OUR EVALUATION	

$[\Gamma(e^+ \nu_e) + \Gamma(e^+ \nu_e \gamma)] / [\Gamma(\mu^+ \nu_\mu) + \Gamma(\mu^+ \nu_\mu \gamma)]$ $(\Gamma_3 + \Gamma_4)/(\Gamma_1 + \Gamma_2)$

See note [a] in the list of π^+ decay modes above. See NUMAO 92 for a discussion of $e-\mu$ universality.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.230 ± 0.004 OUR AVERAGE				
1.2346 ± 0.0035 ± 0.0036	120k	CZAPEK 93	CALO	Stopping π^+
1.2265 ± 0.0034 ± 0.0044	190k	BRITTON 92	CNTR	Stopping π^+
1.218 ± 0.014	32k	BRYMAN 86	CNTR	Stopping π^+
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.273 ± 0.028	11k	9 DICAPUA 64	CNTR	
1.21 ± 0.07		ANDERSON 60	SPEC	

⁹ DICAPUA 64 has been updated using the current mean life.

$\Gamma(\mu^+ \nu_\mu \gamma)/\Gamma_{\text{total}}$ Γ_2/Γ
Note that measurements here do not cover the full kinematic range.

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
1.24 ± 0.25	26	CASTAGNOLI 58	EMUL	$KE_\mu < 3.38$ MeV

$\Gamma(e^+ \nu_e \gamma) / \Gamma_{\text{total}}$ Γ_4 / Γ

Note that measurements here do not cover the full kinematic range.

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	COMMENT
16.1 ± 2.3		¹⁰ BOLOTOV	90B SPEC	17 GeV $\pi^- \rightarrow e^- \bar{\nu}_e \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.6 ± 0.7	226	¹¹ STETZ	78 SPEC	$P_e > 56$ MeV/c
3.0	143	DEPOMMIER	63B CNTR	(KE) $e^+ \gamma > 48$ MeV

¹⁰ BOLOTOV 90B is for $E_\gamma > 21$ MeV, $E_e > 70 - 0.8 E_\gamma$.¹¹ STETZ 78 is for an $e^- \gamma$ opening angle $> 132^\circ$. Obtains 3.7 when using same cutoffs as DEPOMMIER 63B. $\Gamma(e^+ \nu_e \pi^0) / \Gamma_{\text{total}}$ Γ_5 / Γ

VALUE (units 10^{-8})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.025 ± 0.034 OUR AVERAGE					
1.026 ± 0.039	1224	¹² MCFARLANE	85 CNTR	+	Decay in flight
1.00 +0.08 -0.10	332	DEPOMMIER	68 CNTR	+	
1.07 ± 0.21	38	¹³ BACASTOW	65 OSPK	+	
1.10 ± 0.26		¹³ BERTRAM	65 OSPK	+	
1.1 ± 0.2	43	¹³ DUNAITSEV	65 CNTR	+	
0.97 ± 0.20	36	¹³ BARTLETT	64 OSPK	+	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.15 ± 0.22	52	¹³ DEPOMMIER	63 CNTR	+	See DEPOMMIER 68

¹² MCFARLANE 85 combines a measured rate $(0.394 \pm 0.015)/s$ with 1982 PDG mean life.¹³ DEPOMMIER 68 says the result of DEPOMMIER 63 is at least 10% too large because of a systematic error in the π^0 detection efficiency, and that this may be true of all the previous measurements (also V. Soergel, private communication, 1972). $\Gamma(e^+ \nu_e e^+ e^-) / \Gamma(\mu^+ \nu_\mu)$ Γ_6 / Γ_1

VALUE (units 10^{-9})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.5 ± 0.2		98	EGLI	89 SPEC	Uses $R_{PCAC} = 0.068 \pm 0.004$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.46 ± 0.16 ± 0.07	7	¹⁴ BARANOV	92 SPEC		Stopped π^+
< 4.8	90	KORENCH...	76B SPEC		
< 34	90	KORENCH...	71 OSPK		

¹⁴ This measurement by BARANOV 92 is of the structure-dependent part of the decay. The value depends on values assumed for ratios of form factors. $\Gamma(e^+ \nu_e \nu \bar{\nu}) / \Gamma_{\text{total}}$ Γ_7 / Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
< 5	90	PICCIOTTO	88 SPEC	

 $\Gamma(\mu^+ \nu_e) / \Gamma_{\text{total}}$ Γ_8 / Γ

Forbidden by total lepton number conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	COOPER	82 HLBC	Wideband ν beam

 $\Gamma(\mu^+ \nu_e) / \Gamma_{\text{total}}$ Γ_9 / Γ

Forbidden by lepton family number conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
< 8.0	90	COOPER	82 HLBC	Wideband ν beam

 $\Gamma(\mu^- e^+ e^+ \nu) / \Gamma_{\text{total}}$ Γ_{10} / Γ

Forbidden by lepton family number conservation.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 1.6	90	BARANOV	91B SPEC	+	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.9959	90	¹⁵ FETSCHER	84 RVUE	+	
-0.99 ± 0.16		¹⁶ ABELA	83 SPEC	-	μ X-rays

¹⁵ FETSCHER 84 uses only the measurement of CARR 83.¹⁶ Sign of measurement reversed in ABELA 83 to compare with μ^+ measurements. π^+ — POLARIZATION OF EMITTED μ^+ $\pi^+ \rightarrow \mu^+ \nu$

Tests the Lorentz structure of leptonic charged weak interactions.

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< (-0.9959)	90	¹⁵ FETSCHER	84 RVUE	+	
-0.99 ± 0.16		¹⁶ ABELA	83 SPEC	-	μ X-rays

¹⁵ FETSCHER 84 uses only the measurement of CARR 83.¹⁶ Sign of measurement reversed in ABELA 83 to compare with μ^+ measurements.NOTE ON $\pi^\pm \rightarrow \ell^\pm \nu \gamma$ AND $K^\pm \rightarrow \ell^\pm \nu \gamma$ FORM FACTORS

(by H.S. Pruijs, Zürich University)

In the radiative decays $\pi^\pm \rightarrow \ell^\pm \nu \gamma$ and $K^\pm \rightarrow \ell^\pm \nu \gamma$, where ℓ is an e or a μ and γ is a real or virtual photon (e^+e^- pair), both the vector and the axial-vector weak hadronic currents contribute to the decay amplitude. Each current gives a

structure-dependent term (SD_V and SD_A) from virtual hadronic states, and the axial-vector current also gives a contribution from inner bremsstrahlung (IB) from the lepton and meson. The IB amplitudes are determined by the meson decay constants f_π and f_K [1]. The SD_V and SD_A amplitudes are parameterized in terms of the vector form factor F_V and the axial-vector form factors F_A and R [1–4]:

$$M(SD_V) = \frac{-eG_F V_{qq'}}{\sqrt{2} m_P} \epsilon^\mu \ell^\nu F_V \epsilon_{\mu\nu\sigma\tau} k^\sigma q^\tau,$$

$$M(SD_A) = \frac{-ieG_F V_{qq'}}{\sqrt{2} m_P} \epsilon^\mu \ell^\nu \{F_A [(s-t)g_{\mu\nu} - q_\mu k_\nu] + R t g_{\mu\nu}\}.$$

Here $V_{qq'}$ is the Cabibbo-Kobayashi-Maskawa mixing-matrix element; $\epsilon^\mu = (e/t)\bar{u}(p_-)\gamma^\mu v(p_+)$ is the polarization vector of the photon; $\ell^\nu = \bar{u}(p_\nu)\gamma^\nu(1-\gamma_5)v(p_\ell)$ is the lepton-neutrino current; q and k are the meson and photon four-momenta, with $s = q \cdot k$ and $t = k^2$; and P stands for π or K . The s and t dependence of the form factors is neglected, which is a good approximation for pions [2] but not for kaons [4]. The pion vector form factor F_V^π is related via CVC to the π^0 lifetime, $|F_V^\pi| = (1/\alpha)\sqrt{2}\Gamma_{\pi^0}/\pi m_{\pi^0}$ [1]. PCAC relates R to the electromagnetic radius of the meson [2,4], $R^P = \frac{1}{3}m_P f_P \langle r_P^2 \rangle$. The calculation of the other form factors, F_A^π , F_V^K , and F_A^K , is model dependent [1,4].

When the photon is real, the partial decay rate can be given analytically [1,5]:

$$\frac{d^2\Gamma_{P \rightarrow \ell\nu\gamma}}{dx dy} = \frac{d^2(\Gamma_{IB} + \Gamma_{SD} + \Gamma_{INT})}{dx dy},$$

where Γ_{IB} , Γ_{SD} , and Γ_{INT} are the contributions from inner bremsstrahlung, structure-dependent radiation, and their interference, and the Γ_{SD} term is given by

$$\frac{d^2\Gamma_{SD}}{dx dy} = \frac{\alpha}{8\pi} \Gamma_{P \rightarrow \ell\nu} \frac{1}{r(1-r)^2} \left(\frac{m_P}{f_P}\right)^2 \times [(F_V + F_A)^2 SD^+ + (F_V - F_A)^2 SD^-].$$

Here

$$SD^+ = (x + y - 1 - r) [(x + y - 1)(1 - x) - r],$$

$$SD^- = (1 - y + r) [(1 - x)(1 - y) + r],$$

where $x = 2E_\gamma/m_P$, $y = 2E_\ell/m_P$, and $r = (m_\ell/m_P)^2$.

In $\pi^\pm \rightarrow e^\pm \nu \gamma$ and $K^\pm \rightarrow e^\pm \nu \gamma$ decays, the interference terms are small, and thus only the absolute values $|F_A + F_V|$ and $|F_A - F_V|$ can be obtained. In $K^\pm \rightarrow \mu^\pm \nu \gamma$ decay, the interference term is important, and thus the signs of F_V and F_A can be obtained. In $\pi^\pm \rightarrow \mu^\pm \nu \gamma$ decay, bremsstrahlung completely dominates. In $\pi^\pm \rightarrow e^\pm \nu e^+ e^-$ and $K^\pm \rightarrow \ell^\pm \nu e^+ e^-$ decays, all three form factors, F_V , F_A , and R , can be determined.

We give the π^\pm form factors F_V , F_A , and R in the Listings below. In the K^\pm Listings, we give the sum $F_A + F_V$ and difference $F_A - F_V$.

Meson Full Listings

π^\pm, π^0

The electroweak decays of the pseudoscalar mesons are investigated to learn something about the unknown hadronic structure of these mesons, assuming a standard $V - A$ structure of the weak leptonic current. The experiments are quite difficult, and it is not meaningful to analyse the results using parameters for both the hadronic structure (decay constants, form factors) and the leptonic weak current (*e.g.*, to add pseudoscalar or tensor couplings to the $V - A$ coupling). Deviations from the $V - A$ interactions are much better studied in purely leptonic systems such as muon decay.

References

1. D.A. Bryman *et al.*, Phys. Reports **88**, 151 (1982). See also our "Note on Pseudoscalar-Meson Decay Constants," above.
2. A. Kersch and F. Scheck, Nucl. Phys. **B263**, 475 (1986).
3. W.T. Chu *et al.*, Phys. Rev. **166**, 1577 (1968).
4. D.Yu. Bardin and E.A. Ivanov, Sov. J. Nucl. Phys. **7**, 286 (1976).
5. S.G. Brown and S.A. Bludman, Phys. Rev. **136**, B1160 (1964).

π^\pm FORM FACTORS

F_V , VECTOR FORM FACTOR

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.017 ± 0.008 OUR AVERAGE				
0.014 ± 0.009	17	BOLOTOV	90B SPEC	17 GeV $\pi^- \rightarrow e^- \bar{\nu}_e \gamma$
0.023 ^{+0.015} _{-0.013}	98	EGLI	89 SPEC	$\pi^+ \rightarrow e^+ \nu_e e^+ e^-$


¹⁷BOLOTOV 90B only determines the absolute value.

F_A , AXIAL-VECTOR FORM FACTOR

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.0116 ± 0.0016 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.0106 ± 0.0060	18	BOLOTOV	90B SPEC	17 GeV $\pi^- \rightarrow e^- \bar{\nu}_e \gamma$
0.0135 ± 0.0016	18	BAY	86 SPEC	$\pi^+ \rightarrow e^+ \nu \gamma$
0.006 ± 0.003	18	PIILONEN	86 SPEC	$\pi^+ \rightarrow e^+ \nu \gamma$
0.011 ± 0.003	18,19	STETZ	78 SPEC	$\pi^+ \rightarrow e^+ \nu \gamma$
0.021 ^{+0.011} _{-0.013}	98	EGLI	89 SPEC	$\pi^+ \rightarrow e^+ \nu_e e^+ e^-$

¹⁸Using the vector form factor from CVC prediction $F_V = 0.0259 \pm 0.0005$. Only the absolute value of F_A is determined.
¹⁹The result of STETZ 78 has a two-fold ambiguity. We take the solution compatible with later determinations.

WEIGHTED AVERAGE
0.0116 ± 0.0016 (Error scaled by 1.3)



π^\pm axial-vector form factor

R, SECOND AXIAL-VECTOR FORM FACTOR

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.089 ± 0.009 -0.008	98	EGLI	89 SPEC	$\pi^+ \rightarrow e^+ \nu_e e^+ e^-$

π^\pm REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1988 edition Physics Letters **B204** (1988).

JECKELMANN	94	PL B (accepted)	+Goudmit, Leisi	(WABRN, VILL)
CZAPEK	93	PRL 70 17	+Federspiel, Flueckiger, Frei+	(BERN, VILL)
BARANOV	92	SJNP 55 1644	+Vanko, Glazov, Evtukhovich+	(JINR)
		Translated from YAF 55 2940.		
BRITTON	92	PRL 68 3000	+Ahmad, Bryman, Burnham+	(TRIU, CARL)
	94	PR D49 28	Britton, Ahmad, Bryman+	(TRIU, CARL)
NUMAO	92	MPL A7 3357		(TRIU)
BARANOV	91B	SJNP 54 790	+Kisei, Korenchenko, Kuchinskii+	(JINR)
		Translated from YAF 54 1298.		
DAUM	91	PL B265 425	+Frosch, Herter, Janousch, Kettle	(VILL)
BOLOTOV	90B	PL B243 308	+Gninenko, Djikibaev, Isakov+	(INRM)
EGLI	89	PL B222 533	+Engfer, Grab, Hermes, Kraus+	(SINDRUM Collab.)
	86	PL B175 97	Egli, Engfer, Grab, Hermes+	(AACH3, ETH, SIN, ZURI)
PDG	88	PL B204	Yost, Barnett+	(LBL+)
PICCIOTTO	88	PR D37 1131	+Ahmad, Britton, Bryman, Clifford+	(TRIU, CNRC)
COHEN	87	RMP 59 1121	+Taylor	(RISC, NBS)
KORENCHENKO	87	SJNP 46 192	Korenchenko, Kostin, Mzhaviya+	(JINR)
		Translated from YAF 46 313.		
BAY	86	PL B174 445	+Ruegger, Gabioud, Joseph, Loude+	(LAUS, ZURI)
BRYMAN	86	PR D33 1211	+Dubois, Macdonald, Numao+	(TRIU, CNRC)
	83	PRL 50 7	Bryman, Dubois, Numao, Olaniya+	(TRIU, CNRC)
JECKELMANN	86B	NP A457 709	+Beer, Chambrier, Elsenhans+	(ETH, FRIB)
	86	PRL 56 1444	Jeckelmann, Nakada, Beer+	(ETH, FRIB)
PIILONEN	86	PRL 57 1402	+Bolton, Cooper, Frank+	(LANL, TEMP, CHIC)
MC FARLANE	85	PR D32 547	+Auerbach, Gallie+	(TEMP, LANL)
ABELA	84	PL 146B 431	+Daum, Eaton, Frosch, Jost, Kettle+	(SIN)
	78	PL 74B 126	Daum, Eaton, Frosch, Hirschmann+	(SIN)
	79	PR D20 2692	Daum, Eaton, Frosch, Hirschmann+	(SIN)
FETSCHER	84	PL 140B 117		(ETH)
ABELA	83	NP A398 413	+Backenstoss, Kunold, Simons+	(BASL, KARLK, KARLE)
CARR	83	PRL 51 189	+Gidai, Goggi, Jodidio, Oram+	(LBL, NWES, TRIU)
COOPER	82	PL 112B 97	+Guy, Michette, Tyndel, Venus	(RI)
LU	80	PRL 45 1066	+Deiker, Dugan, Wu, Caffrey+	(YALE, COLU, JHU)
STETZ	78	NP B138 285	+Carroll, Ottendahl, Perez-Mendez+	(LBL, UCLA)
CARTER	76	PRL 37 1380	+Dixit, Sundaresan+	(CARL, CNRC, CHIC, CIT)
KORENCHENKO	76B	JETP 44 35	Korenchenko, Kostin, Micelmacher+	(JINR)
		Translated from ZETP 71 69.		
MARUSHENKO	76	JETPL 23 72	Marushenko, Mezentsev, Petrunin+	(PNPI)
		Translated from ZETFP 23 80.		
	76	Private Comm.	Shafer	(FNAL)
	78	Private Comm.	Smirnov	(PNPI)
DUNAITSEV	73	SJNP 16 292	+Prokoshkin, Razuvaev+	(SERP)
		Translated from YAF 16 524.		
AYRES	71	PR D3 1051	+Cormack, Greenberg, Kenney+	(LRL, UCSB)
	67	PR 157 1288	Ayres, Caldwell, Greenberg, Kenney, Kurz+	(LRL)
	68	PRL 21 261	Ayres, Cormack, Greenberg+	(LRL, UCSB)
	69	UCRL 18369 Thesis	Ayres	(LRL)
	69	PRL 23 1267	Greenberg, Ayres, Cormack+	(LRL, UCSB)
KORENCHENKO	71	SJNP 13 189	Korenchenko, Kostin, Micelmacher+	(JINR)
		Translated from YAF 13 339.		
BOOTH	70	PL 32B 723	+Johnson, Williams, Wormald	(LIVP)
DEPOMMIER	68	NP B4 189	+Duclos, Heintze, Kleinkecht+	(CERN)
PETRUHKHIN	68	JINR P1 3862	+Rykalin, Khazins, Cisek	(JINR)
HYMAN	67	PL 25B 376	+Loken, Pewit, McKenzie+	(ANL, CMU, NWES)
NORDBERG	67	PL 24B 594	+Lobkowicz, Burman	(ROCH)
BARDON	66	PRL 16 775	+Dore, Dorfan, Krieger+	(COLU)
KINSEY	66	PR 144 1132	+Lobkowicz, Nordberg	(ROCH)
LOBKOWICZ	66	PRL 17 548	+Melissinos, Nagashima+	(ROCH, BNL)
BACASTOW	65	PR 139B 407	+Ghesquiere, Wiegand, Larsen	(LRL, SLAC)
BERTRAM	65	PR 139B 617	+Meyer, Carrigan+	(MICH, CMU)
DUNAITSEV	65	JETP 20 58	+Petrukhnin, Prokoshkin+	(JINR)
		Translated from ZETP 47 44.		
ECKHAUSE	65	PL 19 348	+Harris, Shuler+	(VILL)
BARTLETT	64	PR 136B 1452	+Devons, Meyer, Rosen	(COLU)
DICAPUA	64	PR 133B 1333	+Garland, Pondrom, Strelzoff	(COLU)
	86	Private Comm.	Pondrom	(WISC)
DEPOMMIER	63	PL 5 61	+Heintze, Rubbia, Soergel	(CERN)
ANDERSON	60	PR 119 2050	+Fujii, Miller+	(EFI)
CASTAGNOLI	58	PR 112 1779	+Mucznik	(ROMA)

π^0

$$I^G(J^PC) = 1^-(0^-+)$$

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1988 edition Physics Letters **B204** (1988).

π^0 MASS


The value is calculated from m_{π^\pm} and $(m_{\pi^\pm} - m_{\pi^0})$. The value of m_{π^\pm} has gone up by three (old) standard deviations since our 1992 edition, and thus m_{π^0} has also jumped. See the "Note on the Charged Pion Mass" in the π^\pm Listings.

VALUE (MeV)	DOCUMENT ID		
134.9764 ± 0.0006 OUR FIT			
$m_{\pi^\pm} - m_{\pi^0}$			
Measurements with an error > 0.01 MeV have been omitted from this Listing.			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4.5936 ± 0.0005 OUR FIT			
4.5936 ± 0.0005 OUR AVERAGE			
4.59364 ± 0.00048	CRAWFORD	91 CNTR	$\pi^- p \rightarrow \pi^0 n, n$ TOF
4.5930 ± 0.0013	CRAWFORD	86 CNTR	$\pi^- p \rightarrow \pi^0 n, n$ TOF
• • • We do not use the following data for averages, fits, limits, etc. • • •			
4.59366 ± 0.00048	CRAWFORD	88B CNTR	See CRAWFORD 91
4.6034 ± 0.0052	VASILEVSKY	66 CNTR	
4.6056 ± 0.0055	CZIRR	63 CNTR	

π^0 MEAN LIFE

Measurements with an error $> 1 \times 10^{-17}$ s have been omitted.

VALUE (10^{-17} s)	EVTS	DOCUMENT ID	TECN	COMMENT
8.4 ± 0.6 OUR AVERAGE				Error includes scale factor of 3.0. See the ideogram below.
8.97 ± 0.22 ± 0.17		ATHERTON 85	CNTR	
8.2 ± 0.4		¹ BROWMAN 74	CNTR	Primakoff effect
5.6 ± 0.6		BELLETTINI 70	CNTR	Primakoff effect
9 ± 0.68		KRYSHKIN 70	CNTR	Primakoff effect
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.4 ± 0.5 ± 0.5	1182	² WILLIAMS 88	CBAL	$e^+e^- \rightarrow e^+e^-\pi^0$
¹ BROWMAN 74 gives a π^0 width $\Gamma = 8.02 \pm 0.42$ eV. The mean life is \hbar/Γ .				
² WILLIAMS 88 gives $\Gamma(\gamma\gamma) = 7.7 \pm 0.5 \pm 0.5$ eV. We give here $\tau = \hbar/\Gamma(\text{total})$.				



π^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 2γ	(98.798 ± 0.032) %	S=1.1
Γ_2 $e^+e^-\gamma$	(1.198 ± 0.032) %	S=1.1
Γ_3 γ positronium	(1.82 ± 0.29) × 10 ⁻⁹	
Γ_4 $e^+e^+e^-e^-$	(3.14 ± 0.30) × 10 ⁻⁵	
Γ_5 e^+e^-	(7.5 ± 2.0) × 10 ⁻⁸	
Γ_6 4γ	< 2 × 10 ⁻⁸	CL=90%
Γ_7 $\nu\bar{\nu}$	[a] < 8.3 × 10 ⁻⁷	CL=90%
Γ_8 $\nu_e\bar{\nu}_e$	< 1.7 × 10 ⁻⁶	CL=90%
Γ_9 $\nu_\mu\bar{\nu}_\mu$	< 3.1 × 10 ⁻⁶	CL=90%
Γ_{10} $\nu_\tau\bar{\nu}_\tau$	< 2.1 × 10 ⁻⁶	CL=90%

Charge conjugation (C) or Lepton Family number (LF) violating modes

Γ_{11} 3γ	C	< 3.1 × 10 ⁻⁸	CL=90%
Γ_{12} μ^+e^-	LF		
Γ_{13} $\mu^+e^- + e^-\mu^+$	LF	< 1.72 × 10 ⁻⁸	CL=90%

[a] Astrophysical and cosmological arguments give limits of order 10⁻¹³; see the Full Listings below.

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 1.9$ for 2 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100	
x_4	-1	0
	x_1	x_2

π^0 BRANCHING RATIOS

$\Gamma(e^+e^-\gamma)/\Gamma(2\gamma)$	Γ_2/Γ_1			
VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
1.213 ± 0.033 OUR FIT				Error includes scale factor of 1.1.
1.213 ± 0.030 OUR AVERAGE				
1.25 ± 0.04		SCHARDT 81	SPEC	$\pi^-p \rightarrow n\pi^0$
1.166 ± 0.047	3071	³ SAMIOS 61	HBC	$\pi^-p \rightarrow n\pi^0$
1.17 ± 0.15	27	BUDAGOV 60	HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.196		JOSEPH 60	THEO	QED calculation
³ SAMIOS 61 value uses a Panofsky ratio = 1.62.				

$\Gamma(\gamma\text{positronium})/\Gamma(2\gamma)$	Γ_3/Γ_1			
VALUE (units 10 ⁻⁹)	EVTS	DOCUMENT ID	TECN	COMMENT
1.84 ± 0.29				
	277	AFANASYEV 90	CNTR	pC 70 GeV

$\Gamma(e^+e^+e^-e^-)/\Gamma(2\gamma)$	Γ_4/Γ_1			
VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
3.18 ± 0.30 OUR FIT				
3.18 ± 0.30	146	⁴ SAMIOS 62B	HBC	
⁴ SAMIOS 62B value uses a Panofsky ratio = 1.62.				

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$	Γ_5/Γ			
VALUE (units 10 ⁻⁸)	EVTS	DOCUMENT ID	TECN	COMMENT
7.5 ± 2.0 OUR AVERAGE				
6.9 ± 2.3 ± 0.6	21	⁵ DESHPANDE 93	SPEC	$K^+ \rightarrow \pi^+\pi^0$
8.8 ± 4.5 ± 0.6	8	⁶ MCFARLAND 93	SPEC	$K_L^0 \rightarrow 3\pi^0$ in flight
⁵ The DESHPANDE 93 result with bremsstrahlung radiative corrections is (8.0 ± 2.6 ± 0.6) × 10 ⁻⁸ .				
⁶ The MCFARLAND 93 result with radiative corrections and excluding $[m_{e^-}/m_{\pi^0}]^2 < 0.95$ is (7.6 ± 3.9 ± 0.5) × 10 ⁻⁸ .				

$\Gamma(e^+e^-)/\Gamma(2\gamma)$	Γ_5/Γ_1			
VALUE (units 10 ⁻⁷)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 1.3	90	NIEBUHR 89	SPEC	$\pi^-p \rightarrow \pi^0n$ at rest
< 5.3	90	ZEPHAT 87	SPEC	$\pi^-p \rightarrow \pi^0n$ 0.3 GeV/c
1.7 ± 0.6 ± 0.3	59	FRANK 83	SPEC	$\pi^-p \rightarrow n\pi^0$
1.8 ± 0.6	58	MISCHKE 82	SPEC	See FRANK 83
2.23 ± 2.40 ± 1.10	90	8 FISCHER 78B	SPRK	$K^+ \rightarrow \pi^+\pi^0$

$\Gamma(4\gamma)/\Gamma_{\text{total}}$	Γ_6/Γ			
VALUE (units 10 ⁻⁸)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
< 2	90	MCDONOUGH 88	CBOX	π^-p at rest
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 160	90	BOLOTOV 86C	CALO	
< 440	90	0 AUERBACH 80	CNTR	

$\Gamma(\nu\bar{\nu})/\Gamma_{\text{total}}$
The astrophysical and cosmological limits are many orders of magnitude lower, but we use the best laboratory limit for the Summary Tables.

VALUE (units 10 ⁻⁶)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
< 0.83	90	⁷ ATIYA 91	CNTR	$K^+ \rightarrow \pi^+\nu\nu'$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 2.9 × 10 ⁻⁷		⁸ LAM 91		Cosmological limit
< 3.2 × 10 ⁻⁷		⁹ NATALE 91		SN 1987A
< 6.5	90	DORENBOS... 88	CHRM	Beam dump, prompt
< 24	90	0 ⁷ HERCZEG 81	RVUE	$K^+\nu \rightarrow \pi^+\nu\nu'$

⁷This limit applies to all possible $\nu\nu'$ states as well as to other massless, weakly interacting states.

⁸LAM 91 considers the production of right-handed neutrinos produced from the cosmic thermal background at the temperature of about the pion mass through the reaction $\gamma\gamma \rightarrow \pi^0 \rightarrow \nu\bar{\nu}$.

⁹NATALE 91 considers the excess energy-loss rate from SN 1987A if the process $\gamma\gamma \rightarrow \pi^0 \rightarrow \nu\bar{\nu}$ occurs, permitted if the neutrinos have a right-handed component. As pointed out in LAM 91 (and confirmed by Natale), there is a factor 4 error in the NATALE 91 published result (0.8 × 10⁻⁷).

$\Gamma(\nu_e\bar{\nu}_e)/\Gamma_{\text{total}}$	Γ_8/Γ			
VALUE (units 10 ⁻⁶)	CL% EVTS	DOCUMENT ID	TECN	COMMENT
< 1.7	90	DORENBOS... 88	CHRM	Beam dump, prompt ν
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 3.1	90	¹⁰ HOFFMAN 88	RVUE	Beam dump, prompt ν
¹⁰ HOFFMAN 88 analyzes data from a 400-GeV BEBC beam-dump experiment.				

Meson Full Listings

π^0, η

$\Gamma(\nu_\mu \bar{\nu}_\mu)/\Gamma_{total}$ Γ_9/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<3.1	90	¹¹ HOFFMAN 88	RVUE	Beam dump, prompt ν
••• We do not use the following data for averages, fits, limits, etc. •••				
<7.8	90	DORENBOS... 88	CHRM	Beam dump, prompt ν
¹¹ HOFFMAN 88 analyzes data from a 400-GeV BEBC beam-dump experiment.				

$\Gamma(\nu_\tau \bar{\nu}_\tau)/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	¹² HOFFMAN 88	RVUE	Beam dump, prompt ν
••• We do not use the following data for averages, fits, limits, etc. •••				
<4.1	90	DORENBOS... 88	CHRM	Beam dump, prompt ν
¹² HOFFMAN 88 analyzes data from a 400-GeV BEBC beam-dump experiment.				

$\Gamma(3\gamma)/\Gamma_{total}$ Γ_{11}/Γ

Forbidden by C invariance.

VALUE (units 10^{-8})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.1	90		MCDONOUGH 88	CBOX	$\pi^- p$ at rest
••• We do not use the following data for averages, fits, limits, etc. •••					
< 38	90	0	HIGHLAND 80	CNTR	
<150	90	0	AUERBACH 78	CNTR	
<490	90	0	¹³ DUCLOS 65	CNTR	
<490	90	13	KUTIN 65	CNTR	
¹³ These experiments give $B(3\gamma/2\gamma) < 5.0 \times 10^{-6}$.					

$\Gamma(\mu^+ e^-)/\Gamma_{total}$ Γ_{12}/Γ

Forbidden by lepton family number conservation.

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
<16	90	LEE 90	SPEC	$K^+ \rightarrow \pi^+ \mu^+ e^-$
<78	90	CAMPAGNARI 88	SPEC	See LEE 90

$[\Gamma(\mu^+ e^-) + \Gamma(e^- \mu^+)]/\Gamma_{total}$ Γ_{13}/Γ

Forbidden by lepton family number conservation.

VALUE (units 10^{-9})	CL%	DOCUMENT ID	TECN	COMMENT
< 17.2	90	KROLAK 94	E799	$\ln K_L^0 \rightarrow 3\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
<140		HERCZEG 84	RVUE	$K^+ \rightarrow \pi^+ \mu e$
< 2 $\times 10^{-6}$		HERCZEG 84	THEO	$\mu^- \rightarrow e^- \pi^0$ conversion
< 70	90	BRYMAN 82	RVUE	$K^+ \rightarrow \pi^+ \mu e$

π^0 ELECTROMAGNETIC FORM FACTOR

The amplitude for the process $\pi^0 \rightarrow e^+ e^- \gamma$ contains a form factor $F(x)$ at the $\pi^0 \gamma \gamma$ vertex, where $x = [m_{e^+ e^-}/m_{\pi^0}]^2$. The parameter a in the linear expansion $F(x) = 1 + ax$ is listed below.

All the measurements except that of BEHREND 91 are in the time-like region of momentum transfer.

LINEAR COEFFICIENT OF π^0 ELECTROMAGNETIC FORM FACTOR

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.032 ± 0.004 OUR AVERAGE				
+0.026 ± 0.024 ± 0.048	7548	FARZANPAY 92	SPEC	$\pi^- p \rightarrow \pi^0 n$ at rest
+0.025 ± 0.014 ± 0.026	54k	MEIJERDREES 92B	SPEC	$\pi^- p \rightarrow \pi^0 n$ at rest
+0.0326 ± 0.0026 ± 0.0026	127	¹⁴ BEHREND 91	CELL	$e^+ e^- \rightarrow e^+ e^- \pi^0$
-0.11 ± 0.03 ± 0.08	32k	FONVIEILLE 89	SPEC	Radiation corr.
••• We do not use the following data for averages, fits, limits, etc. •••				
0.12 ± 0.05		¹⁵ TUPPER 83	THEO	FISCHER 78 data
-0.04				
+0.10 ± 0.03	31k	¹⁶ FISCHER 78	SPEC	Radiation corr.
+0.01 ± 0.11	2200	DEVONS 69	OSPK	No radiation corr.
-0.15 ± 0.10	7676	KOBRAK 61	HBC	No radiation corr.
-0.24 ± 0.16	3071	SAMIOS 61	HBC	No radiation corr.

¹⁴BEHREND 91 estimates that their systematic error is of the same order of magnitude as their statistical error, and so we have included a systematic error of this magnitude. The value of a is obtained by extrapolation from the region of large space-like momentum transfer assuming vector dominance.

¹⁵TUPPER 83 is a theoretical analysis of FISCHER 78 including 2-photon exchange in the corrections.

¹⁶The FISCHER 78 error is statistical only. The result without radiation corrections is +0.05 ± 0.03.

π^0 REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1988 edition Physics Letters **B204** (1988).

KROLAK 94	PL B320 407	+Briere, Cheu, Harris+ (FNAL E799 Collab.)
DESHPANDE 93	PRL 71 27	+Alliegro, Chaloupka+ (BNL E851 Collab.)
MCFARLAND 93	PRL 71 31	+Briere, Cheu, Harris+ (FNAL E799 Collab.)
FARZANPAY 92	PL B278 413	+ (ORST, TRIU, BRCO, QUKI, LBL, BIRM, OXF)
MEIJERDREES 92B	PR D45 1439	+Meijer Drees, Waltham+ (PSI SINDRUM-I Collab.)
ATIYA 91	PRL 66 2189	+Chiang, Frank, Haggerty+ (BNL LANL, PRIN, TRIU)
BEHREND 91	ZPHY C49 401	+Criegee, Field, Frank+ (CELLO Collab.)
CRAWFORD 91	PR D43 46	+Daum, Frosch, Jost, Kettle+ (VILL, VIRG)
LAM 91	PR D44 3345	+Ng (AST)
NATALE 91	PL B258 227	+ (SPIFT)
AFANASYEV 90	PL B236 116	+Chvyrov, Karpukhin+ (JINR, MOSU, SERP)
Also 90B	SJNP 51 664	Afanasyev, Gorchakov, Karpukhin, Komarov+ (JINR)
LEE 90	PRL 64 165	Translated from YAF 51 1040.
FONVIEILLE 89	PL B233 65	+Alliegro, Campagnari+ (BNL, FNAL, VILL, WASH, YALE)
NIEBUHR 89	PR D40 2796	+Bensayah, Berthot, Bertin+ (CLER, LYON, SACL)
CAMPAGNARI 88	PRL 61 2062	+Eichler, Felawka, Kozlowski+ (SINDRUM Collab.)
CRAWFORD 88B	PL B213 391	+Alliegro, Chaloupka+ (BNL, FNAL, PSI, WASH, YALE)
DORENBOS... 88	ZPHY C40 497	+Daum, Frosch, Jost, Kettle, Marshall+ (PSI, VIRG)
HOFFMAN 88	PL B208 149	Dorenbosch, Allaby, Amaldi, Barbiellini+ (CHARM Collab.)
MCDONOUGH 88	PL B208 149	(LANL)
PDG 88	PL B204	+Highland, McFarlane, Bolton+ (TEMP, LANL, CHIC)
WILLIAMS 88	PR D38 1365	Yosi, Barnett+ (LBL+)
ZEPHAT 87	JPG 13 1375	+Antreasyan, Bartels, Besset+ (Crystal Ball Collab.)
BOLOTOV 86C	JETPL 43 520	+Playfer, van Doesburg, Bressani+ (OMICRON Collab.)
		+Gninenko, Dzhalikbaev, Isakov (INRM)
		Translated from ZETFP 43 405.
CRAWFORD 86	PRL 56 1043	+Daum, Frosch, Jost, Kettle+ (SIN, VIRG)
ATHERTON 85	PL 150B 81	+Bovel, Coet+ (CERN, ISU, LUND, CURIN, EPI)
HERCZEG 84	PR D29 1954	+Hoffman (LANL)
FRANK 83	PR D28 423	+Hoffman, Mischke, Moir+ (LANL, ARZS)
TUPPER 83	PR D28 2905	+Grose, Samuel (OKSU)
BRYMAN 82	PR D26 2538	(TRIU)
MISCHKE 82	PRL 48 1153	+Frank, Hoffman, Moir, Sarracino+ (LANL, ARZS)
HERCZEG 81	PL 100B 347	+Hoffman (LANL)
SCHARDT 81	PR D23 639	+Frank, Hoffmann, Mischke, Moir+ (ARZS, LANL)
AUERBACH 80	PL 90B 317	+Halik, Highland, McFarlane, Macek+ (TEMP, LASL)
HIGHLAND 80	PRL 44 628	+Auerbach, Halik, McFarlane, Macek+ (TEMP, LASL)
AUERBACH 78	PRL 41 275	+Highland, Johnson+ (TEMP, LASL)
FISCHER 78	PL 73B 359	+Extermann, Guisan, Mermoud+ (GEVA, SACL)
FISCHER 78B	PL 73B 364	+Extermann, Guisan, Mermoud+ (GEVA, SACL)
BROWMAN 74	PRL 33 1400	+Dewire, Gittelman, Hanson+ (CORN, BING)
BELLETTINI 70	NC 66A 243	+Bempador, Lubelsmeyer+ (PISA, BONN)
KRYSHKIN 70	JETP 30 1037	+Sterigov, Usov (TMSK)
DEVONS 69	PR 184 1356	+Nemethy, Nissim-Sabat, Capua+ (COLU, ROMA)
VASILEVSKY 66	PL 23 281	+Vishnyakov, Dunaitsev+ (JINR)
DUCLOS 65	PL 19 253	+Freytag, Heintze+ (CERN, HEID)
KUTIN 65	JETPL 2 243	+Petrukhin, Prokoshkin (JINR)
		Translated from unknown journal.
CZIRR 63	PR 130 341	(LRL)
SAMIOS 62B	PR 126 1844	+Piano, Prodel+ (COLU, BNL)
KOBRAK 61	NC 20 1115	(EFI)
SAMIOS 61	PR 121 275	(COLU, BNL)
BUDAGOV 60	JETP 11 755	+Viktor, Dzhelepev, Ermolov+ (JINR)
		Translated from ZETF 38 1047.
JOSEPH 60	NC 16 997	(EFI)



$$IG(J^{PC}) = 0^+(0^-+)$$


We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1988 edition Physics Letters **B204** (1988).

η MASS

Measurements with an error ≥ 2 MeV are omitted from the average.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
547.45 ± 0.19 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
547.30 ± 0.15		PLOUIN 92	SPEC	$d p \rightarrow \eta^3 \text{He}$
547.45 ± 0.25		DUANE 74	SPEC	$\pi^- p \rightarrow n$ neutrals
548.2 ± 0.65		FOSTER 65C	HBC	
549.0 ± 0.7	148	FOELSCHE 64	HBC	
548.0 ± 1.0	91	ALFF... 62	HBC	
549.0 ± 1.2	53	BASTIEN 62	HBC	
••• We do not use the following data for averages, fits, limits, etc. •••				
555.0 ± 2.0	250	JAMES 66	HBC	
552.0 ± 3.0	325	KRAEMER 64	DBC	
549.3 ± 2.9		DEL COURT 63	CNTR	
546.0 ± 4.0	35	PICKUP 62	HBC	

WEIGHTED AVERAGE
547.45±0.19 (Error scaled by 1.6)

 **η WIDTH**

This is the partial decay rate $\Gamma(\eta \rightarrow \gamma\gamma)$ divided by the fitted branching fraction for that mode. See the "Note on the Decay Width $\Gamma(\eta \rightarrow \gamma\gamma)$," below.

VALUE (keV) DOCUMENT ID
1.20±0.11 OUR FIT Error includes scale factor of 1.8.

 η DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 neutral modes	(70.8 ± 0.8) %	S=1.2
Γ_2 2γ	[a] (38.8 ± 0.5) %	S=1.2
Γ_3 $3\pi^0$	(31.9 ± 0.4) %	S=1.2
Γ_4 $\pi^0 2\gamma$	(7.1 ± 1.4) × 10 ⁻⁴	
Γ_5 charged modes	(29.2 ± 0.8) %	S=1.2
Γ_6 $\pi^+ \pi^- \pi^0$	(23.6 ± 0.6) %	S=1.2
Γ_7 $\pi^+ \pi^- \gamma$	(4.88 ± 0.15) %	S=1.2
Γ_8 $e^+ e^- \gamma$	(5.0 ± 1.2) × 10 ⁻³	
Γ_9 $\mu^+ \mu^- \gamma$	(3.1 ± 0.4) × 10 ⁻⁴	
Γ_{10} $e^+ e^-$	< 3 × 10 ⁻⁴	CL=90%
Γ_{11} $\mu^+ \mu^-$	(5.7 ± 0.8) × 10 ⁻⁶	
Γ_{12} $\pi^+ \pi^- e^+ e^-$	(1.3 ^{+1.3} _{-0.8}) × 10 ⁻³	
Γ_{13} $\pi^+ \pi^- 2\gamma$	< 2.1 × 10 ⁻³	
Γ_{14} $\pi^+ \pi^- \pi^0 \gamma$	< 6 × 10 ⁻⁴	CL=90%
Γ_{15} $\pi^0 \mu^+ \mu^- \gamma$	< 3 × 10 ⁻⁶	CL=90%

**Charge conjugation (C), Parity (P), or
Charge conjugation × Parity (CP) violating modes**

Γ_{16} $\pi^+ \pi^-$	P, CP	< 1.5 × 10 ⁻³	
Γ_{17} 3γ	C	< 5 × 10 ⁻⁴	CL=95%
Γ_{18} $\pi^0 e^+ e^-$	C	[b] < 4 × 10 ⁻⁵	CL=90%
Γ_{19} $\pi^0 \mu^+ \mu^-$	C	[b] < 5 × 10 ⁻⁶	CL=90%

[a] See the "Note on the Decay Width $\Gamma(\eta \rightarrow \gamma\gamma)$ " in these Full Listings.

[b] C parity forbids this to occur as a single-photon process.

CONSTRAINED FIT INFORMATION

An overall fit to a decay rate and 14 branching ratios uses 40 measurements and one constraint to determine 9 parameters. The overall fit has a $\chi^2 = 31.2$ for 32 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	57								
x_4	3	3							
x_6	-88	-84	-5						
x_7	-77	-74	-5	81					
x_8	-11	-10	-1	-3	-4				
x_9	0	0	0	0	0	0			
x_{12}	-3	-3	0	-13	-10	-2	0		
Γ	-13	-8	0	12	10	1	0	0	0
	x_2	x_3	x_4	x_6	x_7	x_8	x_9	x_{12}	

Mode	Rate (keV)	Scale factor
Γ_2 2γ	[a] 0.46 ± 0.04	1.8
Γ_3 $3\pi^0$	0.381 ± 0.035	1.8
Γ_4 $\pi^0 2\gamma$	(8.5 ± 1.9) × 10 ⁻⁴	1.1
Γ_6 $\pi^+ \pi^- \pi^0$	0.283 ± 0.028	1.7
Γ_7 $\pi^+ \pi^- \gamma$	0.058 ± 0.006	1.7
Γ_8 $e^+ e^- \gamma$	0.0060 ± 0.0015	1.1
Γ_9 $\mu^+ \mu^- \gamma$	(3.7 ± 0.6) × 10 ⁻⁴	1.1
Γ_{12} $\pi^+ \pi^- e^+ e^-$	0.0016 ^{+0.0015} _{-0.0010}	

NOTE ON THE DECAY WIDTH $\Gamma(\eta \rightarrow \gamma\gamma)$

(by N.A. Roe, Lawrence Berkeley Laboratory)

In the measurements of $\Gamma(\eta \rightarrow \gamma\gamma)$ listed below, the results from two-photon production disagree with those from Primakoff production. Since the 1990 edition, one new two-photon measurement has been reported by MD-1; it is consistent with previous two-photon results, though the errors are somewhat larger. The weighted average of the two-photon measurements is 0.510 ± 0.026 keV, to be compared with the Primakoff-production measurement of BROWMAN 74B, 0.324 ± 0.046 keV.

In the two-photon measurements, η 's are produced in the QED process $e^+ e^- \rightarrow e^+ e^- \gamma^* \gamma^* \rightarrow e^+ e^- \eta$. The calculation of the rate is believed to be well understood. The uncertainty due to the virtual photon form factor is small; WILLIAMS 88 quotes an uncertainty of 0.2% from this source. Backgrounds to the η signal from beam-gas interactions and other two-photon interactions with missing particles are also small.

In the Primakoff experiments, η 's are produced by the interaction of a real photon with a virtual photon in the Coulomb field of the nucleus. There is coherent background from strong production of η 's in the nuclear hadronic field, and interference between the strong and Primakoff production amplitudes. The angular dependences of the Primakoff signal and the background are different, allowing $\Gamma(\eta \rightarrow \gamma\gamma)$ to be extracted from a fit to the angular distribution. In the best fit to their data, BEMPORAD 67 found the coherent hadronic background to be consistent with zero. BROWMAN 74B had a wider range of photon energies, a higher maximum energy, better angular resolution, and higher statistics. They found a significant contribution from the hadronic background, especially at lower

Meson Full Listings

 η

energies. BROWMAN 74B also reanalyzed the data of BEMPORAD 67 and found that it was compatible with their fit, including background terms. This suggests that the background was underestimated by BEMPORAD 67, and we consider their result to be superseded by that of BROWMAN 74B.

There remains the disagreement between the two-photon results and the result of BROWMAN 74B. The errors assigned by BROWMAN 74B include a 5.3% statistical error, a 12.2% systematic error for uncertainty in the accepted photon spectrum, and a 2.5% systematic error for uncertainty in the nuclear parameters used in the calculation of the Primakoff and nuclear form factors. The Primakoff form factor F_C is a function of the momentum transfer q and the production angle θ . As $q^2 \rightarrow 0$, the uncertainty in F_C due to the q^2 dependence vanishes. The minimum q^2 in this experiment ranged from -680 MeV^2 at the lowest energy to -174 MeV^2 at the highest. In this range, the result is sensitive to details in the calculation of F_C , but it is difficult to estimate the systematic error of this dependence. Another possible source of systematic error is in the phase of the interference term, ϕ . This was a free parameter in the fit, but was not well determined by the data because the interference contribution peaks in the same angular region as the Primakoff signal and so cannot be unambiguously separated by an angular fit. A reanalysis of the data would be necessary to determine whether any of these factors was overlooked in the determination of the systematic error.

Using the same apparatus, Browman *et al.* [1] measured $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ to be $7.92 \pm 0.42 \text{ eV}$, in good agreement with our world average of $7.7 \pm 0.6 \text{ eV}$. (Our average includes the measurement of Browman *et al.*, but is dominated by a decay-length measurement by Atherton *et al.* [2] The error on the average involves a scale factor $S=3.0$ due to one outlying measurement.) However, the uncertainty due to F_C is reduced at lower momentum transfers, and q^2 was on the order of 100 times smaller in the π^0 measurement. The signal-to-background ratio is also larger, making the fit less sensitive to nuclear production.

A possible source of common systematic error in the two-photon experiments is the calculation of the two-photon luminosity function. However, WILLIAMS 88 measured the two-photon width of the π^0 as well as of the η , and their result, $7.7 \pm 0.5 \pm 0.5 \text{ eV}$, is consistent with the world average quoted above.

To summarize, the two-photon measurements seem more reliable than the best Primakoff-production measurement. However, we include the latter in our average as there is no compelling reason to exclude it. The result, $\Gamma(\eta \rightarrow \gamma\gamma) = 0.46 \pm 0.04 \text{ keV}$, is about one standard deviation from the average using only the two-photon measurements, $0.510 \pm 0.026 \text{ keV}$, and the error is larger, due to the scale factor.

References


1. A. Browman *et al.*, Phys. Rev. Lett. **33**, 1400 (1974).
2. H.W. Atherton *et al.*, Phys. Lett. **158B**, 81 (1985).

 η DECAY RATES

$\Gamma(2\gamma)$								Γ_2	
VALUE (keV)	EVTs	DOCUMENT ID	TECN	COMMENT					
0.46 ± 0.04	OUR FIT	Error includes scale factor of 1.8.							
0.46 ± 0.04	OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.							
0.51 ± 0.12 ± 0.05	36	BARU	90 MD1	$e^+e^- \rightarrow e^+e^-\eta$					
0.490 ± 0.010 ± 0.048	2287	ROE	90 ASP	$e^+e^- \rightarrow e^+e^-\eta$					
0.514 ± 0.017 ± 0.035	1295	WILLIAMS	88 CBAL	$e^+e^- \rightarrow e^+e^-\eta$					
0.53 ± 0.04 ± 0.04		BARTEL	85E JADE	$e^+e^- \rightarrow e^+e^-\eta$					
0.324 ± 0.046		BROWMAN	74B CNTR	Primakoff effect					
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●									
0.64 ± 0.14 ± 0.13		AIHARA	86 TPC	$e^+e^- \rightarrow e^+e^-\eta$					
0.56 ± 0.16	56	WEINSTEIN	83 CBAL	$e^+e^- \rightarrow e^+e^-\eta$					
1.00 ± 0.22		¹ BEMPORAD	67 CNTR	Primakoff effect					

¹BEMPORAD 67 gives $\Gamma(2\gamma) = 1.21 \pm 0.26 \text{ keV}$ assuming $\Gamma(2\gamma)/\Gamma(\text{total}) = 0.314$. Bemporad private communication gives $\Gamma(2\gamma)^2/\Gamma(\text{total}) = 0.380 \pm 0.083$. We evaluate this using $\Gamma(2\gamma)/\Gamma(\text{total}) = 0.38 \pm 0.01$. Not included in average because the uncertainty resulting from the separation of the coulomb and nuclear amplitudes has apparently been underestimated.

WEIGHTED AVERAGE
0.46±0.04 (Error scaled by 1.8)

 η BRANCHING RATIOS

$\Gamma(\text{neutral modes})/\Gamma_{\text{total}}$				$\Gamma_1/\Gamma = (\Gamma_2 + \Gamma_3 + \Gamma_4)/\Gamma$	
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
0.708 ± 0.008	OUR FIT	Error includes scale factor of 1.2.			
0.705 ± 0.008	OUR AVERAGE	Error includes scale factor of 1.2.			
0.79 ± 0.08		BUNIA TOV	67 OSPK	MM spectrometer	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					

$\Gamma(2\gamma)/\Gamma(\text{neutral modes})$				$\Gamma_2/\Gamma_1 = \Gamma_2/(\Gamma_2 + \Gamma_3 + \Gamma_4)$	
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
0.5487 ± 0.0028	OUR FIT	Error includes scale factor of 1.1.			
0.549 ± 0.004	OUR AVERAGE	Error includes scale factor of 1.1.			
0.549 ± 0.004		ALDE	84 GAM2		
0.535 ± 0.018		BUTTRAM	70 OSPK		
0.59 ± 0.033		BUNIA TOV	67 OSPK		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.52 ± 0.09	88	ABROSIMOV	80 HLBC		
0.60 ± 0.14	113	KENDALL	74 OSPK		
0.57 ± 0.09		STRUGALSKI	71 HLBC		
0.579 ± 0.052		FELDMAN	67 OSPK		
0.416 ± 0.044		DIGIUGNO	66 CNTR	Error doubled	
0.44 ± 0.07		GRUNHAUS	66 OSPK		
0.39 ± 0.06		² JONES	66 CNTR		

²This result from combining cross sections from two different experiments.

$\Gamma(3\pi^0)/\Gamma(\text{neutral modes})$				$\Gamma_3/\Gamma_1 = \Gamma_3/(\Gamma_2 + \Gamma_3 + \Gamma_4)$	
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
0.4503 ± 0.0028	OUR FIT	Error includes scale factor of 1.1.			
0.450 ± 0.004	OUR AVERAGE	Error includes scale factor of 1.1.			
0.450 ± 0.004		ALDE	84 GAM2		
0.439 ± 0.024		BUTTRAM	70 OSPK		
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
0.44 ± 0.08	75	ABROSIMOV	80 HLBC		
0.32 ± 0.09		STRUGALSKI	71 HLBC		
0.41 ± 0.033		BUNIA TOV	67 OSPK	Not indep. of $\Gamma(2\gamma)/\Gamma(\text{neutral modes})$	
0.177 ± 0.035		FELDMAN	67 OSPK		
0.209 ± 0.054		DIGIUGNO	66 CNTR	Error doubled	
0.29 ± 0.10		GRUNHAUS	66 OSPK		

$\Gamma(3\pi^0)/\Gamma(2\gamma)$ Γ_3/Γ_2

VALUE	DOCUMENT ID	TECN	COMMENT
0.821±0.009 OUR FIT			Error includes scale factor of 1.1.
0.841±0.030 OUR AVERAGE			
0.841±0.034	AMSLER	93 CBAR	LEAR $\bar{p}p$ at rest
0.91 ±0.14	COX	70b HBC	
0.75 ±0.09	DEVONS	70 OSPK	
0.88 ±0.16	BALTAY	67b DBC	
1.1 ±0.2	CENCE	67 OSPK	
••• We do not use the following data for averages, fits, limits, etc. •••			
1.25 ±0.39	BACCI	63 CNTR	Inverse BR reported

$\Gamma(\pi^0 2\gamma)/\Gamma(\text{neutral modes})$ $\Gamma_4/\Gamma_1 = \Gamma_4/(\Gamma_2+\Gamma_3+\Gamma_4)$

VALUE	DOCUMENT ID	TECN
0.0010±0.00020 OUR FIT		
0.0010 ±0.0002	ALDE	84 GAM2

$\Gamma(\pi^0 2\gamma)/\Gamma_{\text{total}}$ Γ_4/Γ

These results are summarized in the review by LANDSBERG 85.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.1±1.4 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
9.5±2.3		70	BINON	82 GAM2	See ALDE 84
<30		90	DAVYDOV	81 GAM2	$\pi^- p \rightarrow \eta n$

$\Gamma(\text{neutral modes})/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^+ \pi^- \gamma) + \Gamma(e^+ e^- \gamma)]$
 $\Gamma_1/(\Gamma_6+\Gamma_7+\Gamma_8) = (\Gamma_2+\Gamma_3+\Gamma_4)/(\Gamma_6+\Gamma_7+\Gamma_8)$

VALUE	EVTS	DOCUMENT ID	TECN
2.44±0.09 OUR FIT			Error includes scale factor of 1.2.
2.64±0.23		BALTAY	67b DBC
••• We do not use the following data for averages, fits, limits, etc. •••			
4.5 ±1.0	280	³ JAMES	66 HBC
3.20±1.26	53	³ BASTIEN	62 HBC
2.5 ±1.0	10	³ PICKUP	62 HBC

³ These experiments are not used in the averages as they do not separate clearly $\eta \rightarrow \pi^+ \pi^- \pi^0$ and $\eta \rightarrow \pi^+ \pi^- \gamma$ from each other. The reported values thus probably contain some unknown fraction of $\eta \rightarrow \pi^+ \pi^- \gamma$.

$\Gamma(2\gamma)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^+ \pi^- \gamma) + \Gamma(e^+ e^- \gamma)]$ $\Gamma_2/(\Gamma_6+\Gamma_7+\Gamma_8)$

VALUE	EVTS	DOCUMENT ID	TECN
1.34±0.05 OUR FIT			Error includes scale factor of 1.2.
1.1 ±0.4 OUR AVERAGE			
1.51±0.93	75	KENDALL	74 OSPK
0.99±0.48		CRAWFORD	63 HBC

$\Gamma(\text{neutral modes})/\Gamma(\pi^+ \pi^- \pi^0)$ $\Gamma_1/\Gamma_6 = (\Gamma_2+\Gamma_3+\Gamma_4)/\Gamma_6$

VALUE	EVTS	DOCUMENT ID	TECN
2.99±0.11 OUR FIT			Error includes scale factor of 1.2.
3.26±0.30 OUR AVERAGE			
2.54±1.89	74	KENDALL	74 OSPK
3.4 ±1.1	29	AGUILAR...	72b HBC
2.83±0.80	70	⁴ BLOODWOO...	72b HBC
3.6 ±0.6	244	FLATTE	67b HBC
2.89±0.56		ALFF...	66 HBC
3.6 ±0.8	50	KRAEMER	64 DBC
3.8 ±1.1		PAULI	64 DBC

⁴ Error increased from published value 0.5 by Bloodworth (private communication).


$\Gamma(2\gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_2/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN
1.64±0.06 OUR FIT			Error includes scale factor of 1.2.
1.69±0.21 OUR AVERAGE			
1.72±0.25	401	BAGLIN	69 HLBC
1.61±0.39		FOSTER	65 HBC

$\Gamma(3\pi^0)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_3/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN
1.35±0.05 OUR FIT			Error includes scale factor of 1.2.
1.27^{+0.15}_{-0.14} OUR AVERAGE			Error includes scale factor of 1.3. See the ideogram below.
1.50 ^{+0.15} _{-0.29}	199	BAGLIN	69 HLBC
1.47 ^{+0.20} _{-0.17}		BULLOCK	68 HLBC
1.3 ±0.4		BAGLIN	67b HLBC
0.90±0.24		FOSTER	65 HBC
2.0 ±1.0		FOELSCH	64 HBC
0.83±0.32		CRAWFORD	63 HBC

WEIGHTED AVERAGE
 1.27±0.12±0.14 (Error scaled by 1.3)



$\Gamma(\pi^+ \pi^- \gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_7/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN
0.207±0.004 OUR FIT			Error includes scale factor of 1.1.
0.207±0.004 OUR AVERAGE			Error includes scale factor of 1.1.
0.209±0.004	18k	THALER	73 ASPK
0.201±0.006	7250	GORMLEY	70 ASPK
••• We do not use the following data for averages, fits, limits, etc. •••			
0.28 ±0.04		BALTAY	67b DBC
0.25 ±0.035		LITCHFIELD	67 DBC
0.30 ±0.06		CRAWFORD	66 HBC
0.196±0.041		FOSTER	65c HBC

$\Gamma(e^+ e^- \gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_8/Γ_6

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
2.1±0.5 OUR FIT				
2.1±0.5	80	JANE	75b OSPK	See the erratum

$\Gamma(\mu^+ \mu^- \gamma)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.1±0.4 OUR FIT				
3.1±0.4	600	DZHELADIN	80 SPEC	$\pi^- p \rightarrow \eta n$
••• We do not use the following data for averages, fits, limits, etc. •••				
1.5±0.75	100	BUSHNIN	78 SPEC	See DZHELADIN 80

$\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<3	90	DAVIES	74 RVUE	Uses ESTEN 67

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
5.7±0.8 OUR AVERAGE					
5.6 ^{+0.6} _{-0.7} ±0.5		100	KESSLER	93 SPEC	$\rho d \rightarrow \eta^3 \text{He}$
6.5±2.1		27	DZHELADIN	80b SPEC	$\pi^- p \rightarrow \eta n$
••• We do not use the following data for averages, fits, limits, etc. •••					
<20		95	WEHMANN	68 OSPK	

$\Gamma(\mu^+ \mu^-)/\Gamma(2\gamma)$ Γ_{11}/Γ_2

VALUE (units 10^{-5})	DOCUMENT ID	TECN
5.9±2.2	HYAMS	69 OSPK

$\Gamma(\pi^+ \pi^- e^+ e^-)/\Gamma(\pi^+ \pi^- \gamma)$ Γ_{12}/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN
0.027^{+0.026}_{-0.017} OUR FIT			
0.026±0.026	1	GROSSMAN	66 HBC

$\Gamma(\pi^+ \pi^- e^+ e^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE (units 10^{-2})	DOCUMENT ID	TECN
0.13^{+0.13}_{-0.08} OUR FIT		
••• We do not use the following data for averages, fits, limits, etc. •••		
<0.7	RITTENBERG	65 HBC

$\Gamma(\pi^+ \pi^- 2\gamma)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_{13}/Γ_6

VALUE	CL%	DOCUMENT ID	TECN
<0.009		PRICE	67 HBC
••• We do not use the following data for averages, fits, limits, etc. •••			
<0.016	95	BALTAY	67b DBC

Meson Full Listings

 η

$\Gamma(\pi^+\pi^-\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{14}/Γ_6
VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	
<0.24	90	0	THALER	73 ASPK	
••• We do not use the following data for averages, fits, limits, etc. •••					
<1.7	90		ARNOLD	68 HLBC	
<1.6	95		BALTAY	67B DBC	
<7.0			FLATTE	67 HBC	
<0.9			PRICE	67 HBC	
$\Gamma(\pi^0\mu^+\mu^-\gamma)/\Gamma_{total}$					Γ_{15}/Γ
VALUE (units 10^{-6})	CL%		DOCUMENT ID	TECN	COMMENT
<3	90		DZHELJADIN	81 SPEC	$\pi^-p \rightarrow \eta n$
$\Gamma(\pi^+\pi^-)/\Gamma_{total}$					Γ_{16}/Γ
Forbidden by P and CP invariance.					
VALUE (units 10^{-2})	EVTS		DOCUMENT ID	TECN	
<0.15	0		THALER	73 ASPK	
$\Gamma(3\gamma)/\Gamma(\text{neutral modes})$					$\Gamma_{17}/\Gamma_1 = \Gamma_{17}/(\Gamma_2+\Gamma_3+\Gamma_4)$
Forbidden by C invariance.					
VALUE (units 10^{-4})	CL%		DOCUMENT ID	TECN	
<7	95		ALDE	84 GAM2	
$\Gamma(\pi^0e^+e^-)/\Gamma(\pi^+\pi^-\pi^0)$					Γ_{18}/Γ_6
C parity forbids this to occur as a single-photon process.					
VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	
< 1.9	90		JANE	75 OSPK	
••• We do not use the following data for averages, fits, limits, etc. •••					
< 42	90		BAGLIN	67 HLBC	
< 16	90	0	BILLING	67 HLBC	
< 77		0	FOSTER	65B HBC	
<110			PRICE	65 HBC	
$\Gamma(\pi^0e^+e^-)/\Gamma_{total}$					Γ_{18}/Γ
C parity forbids this to occur as a single-photon process.					
VALUE (units 10^{-2})	CL%	EVTS	DOCUMENT ID	TECN	
<0.016	90	0	MARTYNOV	76 HLBC	
<0.084	90		BAZIN	68 DBC	
<0.7			RITTENBERG	65 HBC	
$\Gamma(\pi^0\mu^+\mu^-)/\Gamma_{total}$					Γ_{19}/Γ
C parity forbids this to occur as a single-photon process.					
VALUE (units 10^{-4})	CL%		DOCUMENT ID	TECN	COMMENT
<0.05	90		DZHELJADIN	81 SPEC	$\pi^-p \rightarrow \eta n$
••• We do not use the following data for averages, fits, limits, etc. •••					
<5			WEHMANN	68 OSPK	

NOTE ON η DECAY PARAMETERS C violation in η decays

A number of experiments have looked for charge asymmetries in $\eta \rightarrow \pi^+\pi^-\pi^0$ and $\eta \rightarrow \pi^+\pi^-\gamma$ decays. Any difference between the π^+ and π^- spectra in either decay would indicate C violation in electromagnetic interactions. Immediately following this Note, we list measurements of the following parameters:

(a) The left-right asymmetry

$$A = (N^+ - N^-)/(N^+ + N^-),$$

where N^+ is the number of events in which the π^+ energy in the η rest frame is greater than the π^- energy, etc.

(b) For the decay $\eta \rightarrow \pi^+\pi^-\pi^0$, the sextant asymmetry

$$A_s = \frac{N_1 + N_3 + N_5 - N_2 - N_4 - N_6}{N_1 + N_2 + N_3 + N_4 + N_5 + N_6},$$

where the N_i are the numbers of events in sextants of the Dalitz plot; see, for example, Layter *et al.* [1] A_s is sensitive to an $I=0$ C -violating final state.

(c) For the decay $\eta \rightarrow \pi^+\pi^-\pi^0$, the quadrant asymmetry

$$A_q = \frac{N_1 + N_3 - N_2 - N_4}{N_1 + N_2 + N_3 + N_4},$$

where the N_i are numbers of events in quadrants of the Dalitz plot. A_q is sensitive to an $I=2$ C -violating final state.

(d) For the decay $\eta \rightarrow \pi^+\pi^-\gamma$, evidence for a D -wave contribution to the C -violating amplitude. The upper limit for this contribution is measured by the parameter β , defined by

$$dN/d|\cos\theta| \propto \sin^2\theta(1 + \beta \cos^2\theta),$$

where θ is the angle between the π^+ and the γ in the dipion center of mass. A term proportional to $\cos^2\theta$ could also come from P - and F -wave interference.

Dalitz plot for $\eta \rightarrow \pi^+\pi^-\pi^0$

The Dalitz plot for $\eta \rightarrow \pi^+\pi^-\pi^0$ decay may be fit to the distribution

$$|M(x, y)|^2 \propto (1 + ay + by^2 + cx + dx^2 + exy).$$

Here

$$x = \sqrt{3}(T_+ - T_-)/Q,$$

$$y = (3T_0/Q) - 1,$$

where T_+ , T_- , and T_0 are the kinetic energies of the π^+ , π^- , and π^0 in the η rest frame, and $Q = T_+ + T_0 + T_-$. The coefficient of the term linear in x is sensitive to C violation due to an $I=0$ or $I=2$ final state. Below, we list papers that measured a , b , c , and d , but do not tabulate values of these parameters because the assumptions made by different authors are not compatible and do not allow comparison of the numerical values.

Dalitz plot for $\eta \rightarrow \pi^0\pi^0\pi^0$

The Dalitz plot for the decay $\eta \rightarrow \pi^0\pi^0\pi^0$ may be fit to

$$|M|^2 \propto 1 + 2\alpha z,$$

where

$$z = \frac{2}{3} \sum_{i=1}^3 \left(\frac{3E_i - m_\eta}{m_\eta - 3m_{\pi^0}} \right)^2 = \frac{\rho^2}{\rho_{\max}^2}.$$

Here E_i is the energy of the i^{th} pion in the η rest frame, and ρ is the distance from the center of the Dalitz plot. We list measurements of the parameter α below.

Reference

1. J.G. Layter *et al.*, Phys. Rev. Lett. **29**, 316 (1972).

 η C-NONCONSERVING DECAY PARAMETERS $\pi^+\pi^-\pi^0$ LEFT-RIGHT ASYMMETRY PARAMETER

Measurements with an error $> 1.0 \times 10^{-2}$ have been omitted.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN
0.09 ± 0.17 OUR AVERAGE			

0.28 ± 0.26	165k	JANE	74 OSPK
-0.05 ± 0.22	220k	LAYTER	72 ASPK

••• We do not use the following data for averages, fits, limits, etc. •••

1.5 ± 0.5	37k	⁵ GORMLEY	68C ASPK
-----------	-----	----------------------	----------

⁵The GORMLEY 68C asymmetry is probably due to unmeasured ($\mathbf{E} \times \mathbf{B}$) spark chamber effects. New experiments with ($\mathbf{E} \times \mathbf{B}$) controls don't observe an asymmetry.

π⁺π⁻π⁰ SEXTANT ASYMMETRY PARAMETERMeasurements with an error > 2.0 × 10⁻² have been omitted.

VALUE (units 10 ⁻²)	EVTs	DOCUMENT ID	TECN
0.18 ± 0.16 OUR AVERAGE			
0.20 ± 0.25	165k	JANE	74 OSPK
0.10 ± 0.22	220k	LAYTER	72 ASPK
0.5 ± 0.5	37k	GORMLEY	68c WIRE

π⁺π⁻π⁰ QUADRANT ASYMMETRY PARAMETER

VALUE (units 10 ⁻²)	EVTs	DOCUMENT ID	TECN
-0.17 ± 0.17 OUR AVERAGE			
-0.30 ± 0.25	165k	JANE	74 OSPK
-0.07 ± 0.22	220k	LAYTER	72 ASPK


π⁺π⁻γ LEFT-RIGHT ASYMMETRY PARAMETERMeasurements with an error > 2.0 × 10⁻² have been omitted.

VALUE (units 10 ⁻²)	EVTs	DOCUMENT ID	TECN
0.9 ± 0.4 OUR AVERAGE			
1.2 ± 0.6	35k	JANE	74B OSPK
0.5 ± 0.6	36k	THALER	72 ASPK
1.22 ± 1.56	7257	GORMLEY	70 ASPK

π⁺π⁻γ PARAMETER β (D-wave)Sensitive to a D-wave contribution: $dN/d\cos\theta = \sin^2\theta (1 + \beta \cos^2\theta)$

VALUE	EVTs	DOCUMENT ID	TECN
0.05 ± 0.06 OUR AVERAGE			
0.11 ± 0.11	35k	JANE	74B OSPK
0.12 ± 0.06		⁶ THALER	72 ASPK
-0.060 ± 0.065	7250	GORMLEY	70 WIRE

⁶The authors don't believe this indicates D-wave because the dependence of β on the γ energy is inconsistent with theoretical prediction. A cos²θ dependence may also come from P- and F-wave interference.

η → π⁺π⁻γ parameter β (D-wave)**ENERGY DEPENDENCE OF η → π⁺π⁻π⁰ DALITZ PLOT**See the Note on η Decay Parameters above. The following experiments fit to one or more of the coefficients a, b, c, d, or e for $|matrix\ element|^2 = 1 + ay + by^2 + cx + dx^2 + exy$.

VALUE	EVTs	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •			
81k	LAYTER	73 ASPK	
220k	LAYTER	72 ASPK	
1138	CARPENTER	70 HBC	
349	DANBURG	70 DBC	
7250	GORMLEY	70 WIRE	
526	BAGLIN	69 HLBC	
7170	CNOPS	68 OSPK	
37k	GORMLEY	68c WIRE	
1300	CLPWY	66 HBC	
705	LARRIBE	66 HBC	

α PARAMETER FOR η → 3π⁰See the Note on η Decay Parameters above. The value here is of α ln |matrix element|² = 1 + 2αz.

VALUE	EVTs	DOCUMENT ID	TECN
-0.022 ± 0.023	50k	ALDE	84 GAM2
-0.32 ± 0.37	192	BAGLIN	70 HLBC

η REFERENCESWe have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1988 edition Physics Letters **B204** (1988).

AMSLER	93	ZPHY C58 175	+Armstrong, Augustin+ (Crystal Barrel Collab.)
KESSLER	93	PRL 70 892	+Abegg, Baldisseri+ (SPES-II Collab.)
PLOUIN	92	PL B276 526	+ (SACL, EPOL, IPN, SACL, GWU, UCLA, BGUN, LOUC)
BARU	90	ZPHY C48 581	+Binov, Binov+ (MD-1 Collab.)
ROE	90	PR D41 17	+Bartha, Burke, Garbincius+ (ASP Collab.)
PDG	88	PL B204	+Yost, Barnett+ (LBL+)
WILLIAMS	88	PR D38 1365	+Antreasyan, Bartels, Besset+ (Crystal Ball Collab.)
AHARA	86	PR D33 844	+Alston-Garnjost+ (TPC-2γ Collab.)
BARTEL	85E	PL 160B 421	+Becker, Cord, Felst+ (JADE Collab.)
LANDSBERG	85	PRPL 128 310	(SERP)
ALDE	84	ZPHY C25 225	+Binon, Bricman, Donskov+ (SERP, BELG, LAPP)
	84B	SJNP 40 918	+Alde, Binon, Bricman+ (SERP, BELG, LAPP)
		Translated from YAF 40 147.	
WEINSTEIN	83	PRL 29 2896	+Antreasyan, Gu, Kollman+ (Crystal Ball Collab.)
BINON	82	SJNP 36 391	+Bricman, Gouanere+ (SERP, BELG, LAPP, CERN)
		Translated from YAF 36 670.	
	82B	NC 71A 497	+Binon, Bricman+ (SERP, BELG, LAPP, CERN)
DAVYDOV	81	LNC 32 45	+Donskov, Inyakin+ (SERP, BELG, LAPP, CERN)
	81B	SJNP 33 825	+Dayvov, Binon+ (SERP, BELG, LAPP, CERN)
		Translated from YAF 33 1534.	
DZHEL'YADIN	81	PL 105B 239	+Golovkin, Konstantinov, Kubarovski+ (SERP)
	81C	SJNP 33 822	+Dzhelyadin, Viktorov, Golovkin+ (SERP)
		Translated from YAF 33 1529.	
ABROSIMOV	80	SJNP 31 195	+Ilina, Niszc, Okhrimenko+ (JINR)
		Translated from YAF 31 371.	
DZHEL'YADIN	80	PL 94B 548	+Viktorov, Golovkin+ (SERP)
	80C	SJNP 32 516	+Dzhelyadin, Golovkin, Kachanov+ (SERP)
		Translated from YAF 32 998.	
DZHEL'YADIN	80B	PL 97B 471	+Viktorov, Golovkin+ (SERP)
	80D	SJNP 32 518	+Dzhelyadin, Golovkin, Kachanov+ (SERP)
		Translated from YAF 32 1002.	
BUSHNIN	78	PL 79B 147	+Dzhelyadin, Golovkin, Gritsuk+ (SERP)
	78B	SJNP 28 775	+Bushnin, Golovkin, Gritsuk, Dzhelyadin+ (SERP)
		Translated from YAF 28 1507.	
MARTYNOV	76	SJNP 23 48	+Saltykov, Tarasov, Uzhinskii (JINR)
		Translated from YAF 23 93.	
JANE	75	PL 59B 99	+Grannis, Jones, Lipman, Owen+ (RHEL, LOWC)
JANE	75B	PL 59B 103	+Grannis, Jones, Lipman, Owen+ (RHEL, LOWC)
	78B	PL 73B 503	+Jane
		Erratum in private communication.	
BROWMAN	74B	PRL 32 1067	+Dewire, Gittelmann, Hanson, Loh+ (CORN, BING)
DAVIES	74	NC 24A 324	+Guy, Zia (BIRM, RHEL, SHMP)
DUANE	74	PRL 32 425	+Binnie, Camilleri, Carr+ (LOIC, SHMP)
JANE	74	PL 48B 260	+Jones, Lipman, Owen+ (RHEL, LOWC, SUSS)
JANE	74B	PL 48B 265	+Jones, Lipman, Owen+ (RHEL, LOWC, SUSS)
KENDALL	74	NC 21A 387	+Lanou, Massimo, Shapiro+ (BROW, BARI, MIT)
LAYTER	73	PR D7 2565	+Appel, Kotlewski, Layter, Lee, Stein (COLU)
THALER	73	PR D7 2569	+Appel, Kotlewski, Layter, Lee, Stein (COLU)
AGUILAR...	72B	PR D6 29	+Aguilar-Benitez, Chung, Eisner, Samios (BNL)
BLOODW...	72B	NP B39 525	+Bloodworth, Jackson, Prentice, Yoon (TNTO)
LAYTER	72	PR 29 316	+Appel, Kotlewski, Lee, Stein, Thaler (COLU)
THALER	72	PR 29 313	+Appel, Kotlewski, Layter, Lee, Stein (COLU)
BASILE	71D	NC 3A 796	+Bollini, Dalpiaz, Frabetti+ (CERN, BGNA, STRB)
STRUGALSKI	71	NP B27 429	+Chuvio, Gemesy, Ivanovskaya+ (JINR)
BAGLIN	70	NP B22 66	+Bezaguat, Degrang+ (EPOL, MADR, STRB)
BUTTRAM	70	PRL 25 1358	+Kreiser, Mischke (PRIN)
CARPENTER	70	PR D1 1303	+Binkley, Chapman, Cox, Dagan+ (DUKE)
COX	70B	PRL 24 534	+Fortney, Golsen (DUKE)
DANBURG	70	PR D2 2564	+Abolins, Dahl, Davies, Hoch, Kirz+ (LRL)
DEVONS	70	PR D1 1936	+Grunhaus, Kozlowski, Nemethy+ (COLU, SYRA)
GORMLEY	70	PR D2 501	+Hyman, Lee, Nash, Peoples+ (COLU, BNL)
	Also	70B Nevis 181 Thesis	+Gormley (COLU)
BAGLIN	69	PL 29B 445	+Bezaguat+ (EPOL, UCB, MADR, STRB)
	Also	70	NP B22 66 (EPOL, MADR, STRB)
HYAMS	69	PL 29B 128	+Koch, Potter, VonLindern+ (CERN, MPIM)
ARNOLD	68	PL 27B 466	+Pati, Baglin, Bingham+ (STRB, MADR, EPOL, UCB)
BAZIN	68	PRL 20 895	+Goshaw, Zacher+ (PRIN, QUKI)
BULLOCK	68	PL 27B 402	+Esten, Fleming, Govan, Henderson+ (LOUC)
CNOPS	68	PRL 21 1609	+Hough, Cohn+ (BNL, ORNL, UCND, TENN, PENN)
GORMLEY	68C	PRL 21 402	+Hyman, Lee, Nash, Peoples+ (COLU, BNL)
WEHMANN	68	PRL 20 748	+Engels+ (HARV, CASE, SLAC, CORN, MCGI)
BAGLIN	67	PL 24B 637	+Bezaguat, Degrang+ (EPOL, UCB)
BAGLIN	67B	BAPS 12 567	+Bezaguat, Degrang+ (EPOL, UCB)
BALTAY	67	PRL 19 1498	+Franzini, Kim, Newman+ (COLU, STON)
BALTAY	67D	PRL 19 1495	+Franzini, Kim, Newman+ (COLU, BRAN)
BEMPORAD	67	PL 25B 380	+Braccini, Foa, Lubelsmey+ (PISA, BONN)
	Also	67	Private Comm.
BILLING	67	PL 25B 435	+Bullock, Esten, Govan+ (LOUC, OXF)
BUNIATOV	67	PL 25B 560	+Zavattini, Deinet+ (CERN, KARL)
CENCE	67	PRL 19 1393	+Peterson, Stenger, Chiu+ (HAWA, LRL)
ESTEN	67	PL 24B 115	+Govan, Knight, Miller, Tovey+ (LOUC, OXF)
FELDMAN	67	PRL 18 968	+Frati, Gleson, Halpern+ (PENN)
FLATTE	67	PRL 18 976	(LRL)
FLATTE	67B	PR 163 1441	+Wohl (LRL)
LITCHFIELD	67	PL 24B 486	+Rangan, Segar, Smith+ (RHEL, SACL)
PRICE	67	PRL 18 1207	+Crawford (LRL)
ALFF...	66	PR 145 1072	+Alff-Steinberger, Berley+ (COLU, RUTG)
CLPWY	66	PR 149 1044	
CRAWFORD	66	PRL 16 333	+Price (SCUC, LRL, PURD, WISC, YALE)
DIGIUGNO	66	PRL 16 767	+Giorgi, Silvestri+ (LRL)
GROSSMAN	66	PR 146 993	+Price, Crawford (NAPL, TRST, STON)
GRUNHAUS	66	Thesis	(COLU)
JAMES	66	PR 142 896	+Kraybill (YALE, BNL)
JONES	66	PL 23 597	+Binnie, Duane, Horsey, Mason+ (LOIC, RHEL)
LARRIBE	66	PL 23 600	+Leveque, Muller, Pauli+ (SACL, RHEL)
FOSTER	65	PR 138B 652	+Peters, Meer, Loeffler+ (WISC, PURD)
FOSTER	65B	Athens Conf. Thesis	+Good, Meer (WISC)
FOSTER	65C	Thesis	(WISC)
PRICE	65	PRL 15 123	+Crawford (LRL)
RITTENBERG	65	PRL 15 556	+Kalbfleisch (LRL, BNL)
FOELSCH	64	PR 134B 1138	+Kraybill (YALE)
KRAEMER	64	PR 136B 496	+Madansky, Fields+ (JHU, NWES, WOOD)
PAULI	64	PL 13 351	+Muller
BACCI	63	PRL 11 37	+Penco, Salvini+ (ROMA, FRAS)
CRAWFORD	63	PRL 10 546	+Lloyd, Fowler (LRL, DUKE)
	Also	66B	Crawford, Lloyd, Fowler (LRL, DUKE)
DELCOURT	63	PL 7 215	+Lefrancois, Perez-y-Jorba+ (ORSAY)
ALFF...	62	PRL 9 322	+Alff-Steinberger, Berley, Colley+ (COLU, RUTG)
BASTIEN	62	PRL 8 114	+Berge, Dahl, Ferro-Luzzi+ (LRL)
PICKUP	62	PRL 8 329	+Robinson, Salant (CNRC, BNL)

Meson Full Listings

$\rho(770)$

$\rho(770)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

Our latest mini-review on this particle can be found in the 1984 edition.

NOTE ON THE $\rho(770)$

Because of the large width of the $\rho(770)$, determination of the resonance parameters is beset with many difficulties. It is well known that in physical-region fits the line shape does not correspond to a relativistic Breit-Wigner function with a P -wave width, but requires some additional shape parameter. This dependence on parametrization was demonstrated long ago by PISUT 68, who showed that the mass was consistent with values between 761 MeV and 783 MeV to within two standard deviations. When mass values are quoted, as below, with one-standard-deviation errors, the conflicts between them are evident.

The same model dependence afflicts any other source of the resonance parameters, such as the energy dependence of the phase shift δ_1^1 or the pole position. It is therefore not surprising that a recent study of the $\rho(770)$ dominance in the decays of η and η' reveals the need for specific dynamical effects in addition to the $\rho(770)$ pole (BENAYOUN 93).


Recently LAFFERTY 93 has demonstrated that Bose-Einstein correlations is another source of shifts in the $\rho(770)$ line shape.

$\rho(770)$ MASS

We no longer list S -wave Breit-Wigner fits, or data with high combinatorial background.

MIXED CHARGES

769.9 ± 0.8 OUR AVERAGE Includes data from the 3 datablocks that follow this one. Error includes scale factor of 1.8. See the ideogram below.



$\rho(770)$ MASS MIXED CHARGES

CHARGED ONLY

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
The data in this block is included in the average printed for a previous datablock.

766.9 ± 1.2 OUR AVERAGE

768 ± 9		AGUILAR...	91	EHS		400pp
767 ± 3	2935	1 CAPRARO	87	SPEC	-	200 $\pi^+ \pi^-$ Cu →
						$\pi^+ \pi^0$ Cu
761 ± 5	967	1 CAPRARO	87	SPEC	-	200 $\pi^+ \pi^-$ Pb →
						$\pi^+ \pi^0$ Pb
771 ± 4		HUSTON	86	SPEC	+	202 $\pi^+ \pi^-$ A →
						$\pi^+ \pi^0$ A
766 ± 7	6500	2 BYERLY	73	OSPK	-	5 $\pi^- p$
766.8 ± 1.5	9650	3 PISUT	68	RVUE	-	1.7-3.2 $\pi^- p$, t < 10
767 ± 6	900	1 EISNER	67	HBC	-	4.2 $\pi^- p$, t < 10

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
The data in this block is included in the average printed for a previous datablock.

768.1 ± 1.3 OUR AVERAGE

767.6 ± 2.7		BARTALUCCI 78	CNTR	0		$\gamma p \rightarrow e^+ e^- \rho$
775 ± 5		GLADDING 73	CNTR	0		2.9-4.7 γp
767.0 ± 4.0	1930	BALLAM 72	HBC	0		2.8 γp
770.0 ± 4.0	2430	BALLAM 72	HBC	0		4.7 γp
765.0 ± 10.0		ALVENSLEBEN 70	CNTR	0		γA , t < 0.01
767.7 ± 1.9	140k	BIGGS 70	CNTR	0		< 4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
765 ± 5.0	4000	ASBURY 67B	CNTR	0		$\gamma t Pb$

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
The data in this block is included in the average printed for a previous datablock.

770.8 ± 1.2 OUR AVERAGE

Error includes scale factor of 2.2. See the ideogram below.


773 ± 1.6		WEIDENAUER 93	ASTE			$\bar{p} p \rightarrow \pi^+ \pi^- \omega$
762.6 ± 2.6		AGUILAR... 91	EHS			400pp
775.9 ± 1.1		4 BARKOV 85	OLYA	0		$e^+ e^- \rightarrow \pi^+ \pi^-$
768.0 ± 4.0		5,6 BOHACIK 80	RVUE	0		
769.0 ± 3.0		2 WICKLUND 78	ASPK	0		3,4,6 $\pi^\pm N$
768.0 ± 1.0	76000	DEUTSCH... 76	HBC	0		16 $\pi^+ p$
767 ± 4	4100	ENGLER 74	DBC	0		6 $\pi^+ p \rightarrow \pi^+ \pi^- p$
775.0 ± 4.0	32000	5 PROTOPOP... 73	HBC	0		7.1 $\pi^+ p$, t < 0.4
764.0 ± 3.0	6800	RATCLIFF 72	ASPK	0		15 $\pi^- p$, t < 0.3
774.0 ± 3.0	1700	REYNOLDS 69	HBC	0		2.26 $\pi^- p$
775.0 ± 3.0	2250	HYAMS 68	OSPK	0		11.2 $\pi^- p$
769.2 ± 1.5	13300	7 PISUT 68	RVUE	0		1.7-3.2 $\pi^- p$, t < 10

• • • We do not use the following data for averages, fits, limits, etc. • • •

768 ± 1		8 GESHKENBEIN 89	RVUE			π^- form factor
777.4 ± 2.0		9 CHABAUD 83	ASPK	0		17 $\pi^- p$ polarized
770 ± 2		10 HEYN 81	RVUE	0		Pion form factor
769.5 ± 0.7		5,6 LANG 79	RVUE	0		
770 ± 9		6 ESTABROOKS 74	RVUE	0		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
773.5 ± 1.7	11200	1 JACOBS 72	HBC	0		2.8 $\pi^- p$

WEIGHTED AVERAGE

770.8 ± 1.2 (Error scaled by 2.2)



$\rho(770)^0$ mass (MeV)

- Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
- Phase shift analysis. Systematic errors added corresponding to spread of different fits.
- From fit of 3-parameter relativistic P -wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67B, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66B, JACOBS 66B, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.
- From the Gounaris-Sakurai parametrization of the pion form factor.
- From pole extrapolation.
- From phase shift analysis of GRAYER 74 data.

See key on page 1343

Meson Full Listings

$\rho(770)$


- ⁷ Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67b, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66b, JACOBS 66b, JAMES 66, WEST 66, GOLDBERGER 64, ABOLINS 63.
⁸ Includes BARKOV 85 data. Model-dependent width definition.
⁹ From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P-wave intensity. CHABAUD 83 includes data of GRAYER 74.
¹⁰ HEYN 81 includes all spacelike and timelike F_π values until 1978.

$m_{\rho(770)^0} - m_{\rho(770)^\pm}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.3 ± 2.2 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.			
-4.0 ± 4.0	3000	¹¹ REYNOLDS	69 HBC	-0	2.26 $\pi^- p$
-5 ± 5	3600	¹¹ FOSTER	68 HBC	±0	0.0 $\bar{p} p$
2.4 ± 2.1	22950	¹² PISUT	68 RVUE		$\pi N \rightarrow \rho N$

¹¹ From quoted masses of charged and neutral modes.

¹² Includes MALAMUD 69, ARMENISE 68, BATON 68, BACON 67, HUWE 67, MILLER 67b, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66b, JACOBS 66b, JAMES 66, WEST 66, BLIEDEN 65, CARMONY 64, GOLDBERGER 64, ABOLINS 63.



$m_{\rho(770)^0} - m_{\rho(770)^\pm}$ (MeV)

$\rho(770)$ RANGE PARAMETER

The range parameter R enters an energy-dependent correction to the width, of the form $(1 + q^2 R^2) / (1 + q^2 R^2)$, where q is the momentum of one of the pions in the $\pi\pi$ rest system. At resonance, $q = q_r$.

VALUE (GeV ⁻¹)	DOCUMENT ID	TECN	CHG	COMMENT
5.3^{+0.9}_{-0.7}	CHABAUD	83 ASPK	0	17 $\pi^- p$ polarized

$\rho(770)$ WIDTH

We no longer list S-wave Breit-Wigner fits, or data with high combinatorial background.

MIXED CHARGES

VALUE (MeV)	DOCUMENT ID
151.2 ± 1.2 OUR AVERAGE	Includes data from the 3 datablocks that follow this one.

CHARGED ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-------------	------	-------------	------	-----	---------

The data in this block is included in the average printed for a previous datablock.

149.1 ± 2.9 OUR FIT

149.1 ± 2.9 OUR AVERAGE					
155 ± 11	2935	¹³ CAPRARO	87 SPEC	-	200 $\pi^- \text{Cu} \rightarrow \pi^- \pi^0 \text{Cu}$
154 ± 20	967	¹³ CAPRARO	87 SPEC	-	200 $\pi^- \text{Pb} \rightarrow \pi^- \pi^0 \text{Pb}$
150 ± 5		HUSTON	86 SPEC	+	202 $\pi^+ \text{A} \rightarrow \pi^+ \pi^0 \text{A}$
146 ± 12	6500	¹⁴ BYERLY	73 OSPK	-	5 $\pi^- p$
148.2 ± 4.1	9650	¹⁵ PISUT	68 RVUE	-	1.7-3.2 $\pi^- p, t < 10$
146 ± 13	900	EISNER	67 HBC	-	4.2 $\pi^- p, t < 10$

NEUTRAL ONLY, PHOTOPRODUCED

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
 The data in this block is included in the average printed for a previous datablock.

150.9 ± 3.0		BARTALUCCI	78 CNTR	0	$\gamma p \rightarrow e^+ e^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
147 ± 11		GLADDING	73 CNTR	0	2.9-4.7 γp
155.0 ± 12.0	2430	BALLAM	72 HBC	0	4.7 γp
145.0 ± 13.0	1930	BALLAM	72 HBC	0	2.8 γp
140.0 ± 5.0		ALVENSLEBEN	70 CNTR	0	$\gamma A, t < 0.01$
146.1 ± 2.9	140k	BIGGS	70 CNTR	0	< 4.1 $\gamma C \rightarrow \pi^+ \pi^- C$
160.0 ± 10.0		LANZEROTTI	68 CNTR	0	γp
130 ± 5	4000	ASBURY	67b CNTR	0	$\gamma + \text{Pb}$

NEUTRAL ONLY, OTHER REACTIONS

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
 The data in this block is included in the average printed for a previous datablock.

151.9 ± 1.5 OUR FIT

151.9 ± 1.5 OUR AVERAGE		WEIDENAUER	93 ASTE		$\bar{p} p \rightarrow \pi^+ \pi^- \omega$
145.7 ± 5.3		¹⁶ BARKOV	85 OLYA	0	$e^+ e^- \rightarrow \pi^+ \pi^-$
150.5 ± 3.0		^{17,18} BOHACIK	80 RVUE	0	
148.0 ± 6.0		¹⁴ WICKLUND	78 ASPK	0	3.4, 6 $\pi^\pm p N$
152.0 ± 9.0		DEUTSCH...	76 HBC	0	16 $\pi^+ p$
154.0 ± 2.0	76000	RATCLIFF	72 ASPK	0	15 $\pi^- p, t < 0.3$
157.0 ± 8.0	6800	REYNOLDS	69 HBC	0	2.26 $\pi^- p$
143.0 ± 8.0	1700	REYNOLDS	69 HBC	0	2.26 $\pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
138 ± 1		¹⁹ GESHKENBEIN	89 RVUE		π form factor
160.0 ± 4.1		²⁰ CHABAUD	83 ASPK	0	17 $\pi^- p$ polarized
155 ± 1		²¹ HEYN	81 RVUE	0	π form factor
148.0 ± 1.3		^{17,18} LANG	79 RVUE	0	
146 ± 14	4100	ENGLER	74 DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
143 ± 13		¹⁸ ESTABROOKS	74 RVUE	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
160.0 ± 10.0	32000	¹⁷ PROTOPOP...	73 HBC	0	7.1 $\pi^+ p, t < 0.4$
145.0 ± 12.0	2250	¹³ HYAMS	68 OSPK	0	11.2 $\pi^- p$
163.0 ± 15.0	13300	²² PISUT	68 RVUE	0	1.7-3.2 $\pi^- p, t < 10$

¹³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

¹⁴ Phase shift analysis. Systematic errors added corresponding to spread of different fits.

¹⁵ From fit of 3-parameter relativistic P-wave Breit-Wigner to total mass distribution. Includes BATON 68, MILLER 67b, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66b, JACOBS 66b, JAMES 66, WEST 66, BLIEDEN 65 and CARMONY 64.

¹⁶ From the Gounaris-Sakurai parametrization of the pion form factor.

¹⁷ From pole extrapolation.

¹⁸ From phase shift analysis of GRAYER 74 data.

¹⁹ Includes BARKOV 85 data. Model-dependent width definition.

²⁰ From fit of 3-parameter relativistic Breit-Wigner to helicity-zero part of P-wave intensity. CHABAUD 83 includes data of GRAYER 74.

²¹ HEYN 81 includes all spacelike and timelike F_π values until 1978.

²² Includes MALAMUD 69, ARMENISE 68, BACON 67, HUWE 67, MILLER 67b, ALFF-STEINBERGER 66, HAGOPIAN 66, HAGOPIAN 66b, JACOBS 66b, JAMES 66, WEST 66, GOLDBERGER 64, ABOLINS 63.

$\rho(770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/Confidence level
Γ_1 $\pi\pi$	~ 100	%
$\rho(770)^\pm$ decays		
Γ_2 $\pi^\pm \pi^0$	~ 100	%
Γ_3 $\pi^\pm \gamma$	(4.5 ± 0.5) × 10 ⁻⁴	S=2.2
Γ_4 $\pi^\pm \eta$	< 6	× 10 ⁻³ CL=84%
Γ_5 $\pi^\pm \pi^+ \pi^- \pi^0$	< 2.0	× 10 ⁻³ CL=84%
$\rho(770)^0$ decays		
Γ_6 $\pi^+ \pi^-$	~ 100	%
Γ_7 $\pi^+ \pi^- \gamma$	(9.9 ± 1.6) × 10 ⁻³	
Γ_8 $\pi^0 \gamma$	(7.9 ± 2.0) × 10 ⁻⁴	
Γ_9 $\eta \gamma$	(3.8 ± 0.7) × 10 ⁻⁴	
Γ_{10} $\mu^+ \mu^-$	[a] (4.60 ± 0.28) × 10 ⁻⁵	
Γ_{11} $e^+ e^-$	[a] (4.46 ± 0.21) × 10 ⁻⁵	
Γ_{12} $\pi^+ \pi^- \pi^0$	< 1.2	× 10 ⁻⁴ CL=90%
Γ_{13} $\pi^+ \pi^- \pi^+ \pi^-$	< 2	× 10 ⁻⁴ CL=90%
Γ_{14} $\pi^+ \pi^- \pi^0 \pi^0$	< 4	× 10 ⁻⁵ CL=90%

[a] The $e^+ e^-$ branching fraction is from $e^+ e^- \rightarrow \pi^+ \pi^-$ experiments only. The $\omega\rho$ interference is then due to $\omega\rho$ mixing only, and is expected to be small. If $e\mu$ universality holds, $\Gamma(\rho^0 \rightarrow \mu^+ \mu^-) = \Gamma(\rho^0 \rightarrow e^+ e^-) \times 0.99785$.

Meson Full Listings

$\rho(770)$

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 9 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 10.2$ for 7 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{matrix} x_3 & & & \\ \Gamma & \begin{matrix} -100 & & \\ 18 & -18 & \\ & x_2 & x_3 \end{matrix} \end{matrix}$$

Mode	Rate (MeV)	Scale factor
$\Gamma_2 \quad \pi^\pm \pi^0$	149.1 ± 2.9	
$\Gamma_3 \quad \pi^\pm \gamma$	0.068 ± 0.007	2.3

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and a branching ratio uses 9 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 4.8$ for 6 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.


$$\begin{matrix} x_{10} & & & \\ x_{11} & & & \\ \Gamma & \begin{matrix} -80 & & \\ -60 & 0 & \\ 12 & 0 & -20 \end{matrix} \end{matrix}$$

Mode	Rate (MeV)	Scale factor
$\Gamma_6 \quad \pi^+ \pi^-$	151.8 ± 1.5	
$\Gamma_{10} \quad \mu^+ \mu^-$	[a] 0.0070 ± 0.0004	
$\Gamma_{11} \quad e^+ e^-$	[a] 0.00677 ± 0.00032	

$\rho(770)$ PARTIAL WIDTHS

$\Gamma(\pi^\pm \gamma)$	VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
68 ± 7 OUR FIT					Error includes scale factor of 2.3.
68 ± 7 OUR AVERAGE					Error includes scale factor of 2.2. See the ideogram below.
81.0 ± 4.0 ± 4.0		CAPRARO	87	SPEC	- 200 $\pi^- A \rightarrow \pi^+ \pi^0 A$
59.8 ± 4.0		HUSTON	86	SPEC	+ 202 $\pi^+ A \rightarrow \pi^+ \pi^0 A$
71.0 ± 7.0		JENSEN	83	SPEC	- 156-260 $\pi^- A \rightarrow \pi^- \pi^0 A$

WEIGHTED AVERAGE
68±7 (Error scaled by 2.2)



$\Gamma(e^+ e^-)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
6.77 ± 0.32 OUR FIT				
$6.77 \pm 0.10 \pm 0.30$		BARKOV	85	OLYA $e^+ e^- \rightarrow \pi^+ \pi^-$

$\Gamma(\pi^0 \gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
121 ± 31		DOLINSKY	89	ND $e^+ e^- \rightarrow \pi^0 \gamma$

$\Gamma(\eta \gamma)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
62 ± 17		23 DOLINSKY	89	ND $e^+ e^- \rightarrow \eta \gamma$
111 ± 22		24 DOLINSKY	89	ND $e^+ e^- \rightarrow \eta \gamma$

²³ Solution corresponding to constructive ω - ρ interference. The quark model predicts a relative decay phase of zero. Also much favored by the ALDE 93 model-independent measurement of $B(\omega \rightarrow \eta \gamma)$.

²⁴ Solution corresponding to destructive ρ - ω interference.

$\rho(770)$ BRANCHING RATIOS

$\Gamma(\pi^\pm \eta) / \Gamma(\pi \pi)$	VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 60		84	FERBEL	66	HBC	± $\pi^\pm p$ above 2.5

$\Gamma(\pi^\pm \pi^+ \pi^- \pi^0) / \Gamma(\pi \pi)$	VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 20		84	FERBEL	66	HBC	± $\pi^\pm p$ above 2.5
35 ± 40			JAMES	66	HBC	+ 2.1 $\pi^+ p$

$\Gamma(\mu^+ \mu^-) / \Gamma(\pi^+ \pi^-)$	VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
$4.6 \pm 0.2 \pm 0.2$		ANTIPOV	89	SIGM $\pi^- C u \rightarrow \mu^+ \mu^- \pi^- C u$
8.2 ^{+1.6} _{-3.6}		25 ROTHWELL	69	CNTR Photoproduction
5.6 ± 1.5		26 WEHMANN	69	OSPK 12 $\pi^- C, Fe$
9.7 ^{+3.1} _{-3.3}		27 HYAMS	67	OSPK 11 $\pi^- Li, H$

²⁵ Possibly large ρ - ω interference leads us to increase the minus error.
²⁶ Result contains 11 ± 11% correction using SU(3) for central value. The error on the correction takes account of possible ρ - ω interference and the upper limit agrees with the upper limit of $\omega \rightarrow \mu^+ \mu^-$ from this experiment.
²⁷ HYAMS 67's mass resolution is 20 MeV. The ω region was excluded.

$\Gamma(e^+ e^-) / \Gamma(\pi \pi)$	VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
0.41 ± 0.05		BENAKSAS	72	OSPK $e^+ e^-$

$\Gamma(\eta \gamma) / \Gamma_{\text{total}}$	VALUE (units 10^{-4})	DOCUMENT ID	TECN	CHG	COMMENT
3.8 ± 0.7 OUR AVERAGE					
4.0 ± 1.1		28 DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta \gamma$
3.6 ± 0.9		28 ANDREWS	77	CNTR 0	6.7-10 γCu
7.3 ± 1.5		29 DOLINSKY	89	ND	$e^+ e^- \rightarrow \eta \gamma$
5.4 ± 1.1		29 ANDREWS	77	CNTR 0	6.7-10 γCu

²⁸ Solution corresponding to constructive ω - ρ interference. The quark model predicts a relative phase of zero. Also much favored by the ALDE 93 model-independent measurement of $B(\omega \rightarrow \eta \gamma)$.
²⁹ Solution corresponding to destructive ω - ρ interference.

$\Gamma(\pi^+ \pi^- \pi^+ \pi^-) / \Gamma_{\text{total}}$	VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 2		90	KURDADZE	88	OLYA $e^+ e^- \rightarrow \pi^+ \pi^- \pi^+ \pi^-$

$\Gamma(\pi^+ \pi^- \pi^+ \pi^-) / \Gamma(\pi \pi)$	VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 15		90	ERBE	69	HBC	0 2.5-5.8 γp
< 20			CHUNG	68	HBC	0 3.2, 4.2 $\pi^- p$
< 20		90	HUSON	68	HLBC	0 16.0 $\pi^- p$
< 80			JAMES	66	HBC	0 2.1 $\pi^+ p$

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$	VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
< 1.2		90	VASSERMAN	88B	ND $e^+ e^- \rightarrow \pi^+ \pi^- \pi^0$

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma(\pi \pi)$	VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
~ 0.01			BRAMON	86	RVUE	0 $J/\psi \rightarrow \omega \pi^0$
< 0.01		84	30 ABRAMS	71	HBC	0 3.7 $\pi^+ p$

³⁰ Model dependent, assumes $l = 1, 2, \text{ or } 3$ for the 3π system.

See key on page 1343

Meson Full Listings

$\rho(770), \omega(782)$

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	COMMENT
<0.4	90	AULCHENKO 87C	ND	0	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
<2	90	KURDADZE 86 OLYA	0	0	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{total}$ Γ_7/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0099 ± 0.0016		³¹ DOLINSKY 91 ND		$e^+e^- \rightarrow \pi^+\pi^-\gamma$
0.0111 ± 0.0014		³² VASSERMAN 88 ND		$e^+e^- \rightarrow \pi^+\pi^-\gamma$
<0.005	90	³³ VASSERMAN 88 ND		$e^+e^- \rightarrow \pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^0\gamma)/\Gamma_{total}$ Γ_8/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
7.9 ± 2.0	DOLINSKY 89 ND	ND	$e^+e^- \rightarrow \pi^0\gamma$

$\rho(770)$ REFERENCES

ALDE 93	PAN 56 1229	+Binon+ (SERP, LAPP, LANL, BELG, BRUX, CERN)
WEIDENAUER 93	ZPHY C59 387	+Duch+ (ASTERIX Collab.)
AGUILAR... 91	ZPHY C50 405	+Le-Benitez, Allison, Batalor+ (LEBC-EHS Collab.)
DOLINSKY 91	PRPL 202 99	+Druzhinin, Dubrovin+ (NOVO)
ANTIPOV 89	ZPHY C42 185	+Batarin+ (SERP, JINR, BGNA, MILA, TBIL)
DOLINSKY 89	ZPHY C42 511	+Druzhinin, Dubrovin, Golubev+ (NOVO)
GESHKENBEIN 89	ZPHY 45 351	(ITEP)
KURDADZE 88	JETPL 47 512	+Lelchuk, Pakhtusova, Sidorov+ (NOVO)
VASSERMAN 88	SJNP 47 1035	+Golubev, Dolinsky+ (NOVO)
VASSERMAN 88B	SJNP 48 480	+Golubev, Dolinsky+ (NOVO)
AULCHENKO 87C	IYF 87-90 Preprint	+Dolinsky, Druzhinin+ (NOVO)
CAPIARO 87	NP B288 659	+Levy+ (CLER, FRAS, MILA, PISA, LCGT, TRST+)
BRAMON 86	PL B173 97	+Casulleras (BARC)
HUSTON 86	PR 33 3199	+Berg, Collick, Jonckheere+ (ROCH, FNAL, MINN)
KURDADZE 86	JETPL 43 643	+Lelchuk, Pakhtusova, Sidorov, Skrinski+ (NOVO)
BARKOV 85	NP B256 365	+Chilingarov, Eidelman, Khazin, Lelchuk+ (NOVO)
CHABAUD 83	NP B223 1	+Gorlich, Cerrada+ (CERN, CRAC, MPIM)
JENSEN 83	PR D27 26	+Berg, Biel, Collick+ (ROCH, FNAL, MINN)
HEYN 81	ZPHY C7 169	+Lang (GRAZ)
BOHACIK 80	PR D21 1342	+Kuhnelt (SLOV, WIEN)
LANG 79	PR D19 956	+Mas-Parareda (GRAZ)
BARTALUCCI 78	NC 44A 587	+Basini, Bertolucci+ (DESY, FRAS)
WICKLUND 78	PR D17 1197	+Ayres, Diebold, Greene, Kramer, Pawlicki (ANL)
ANDREWS 77	PRL 38 198	+Fukushima, Harvey, Lobkowitz, May+ (ROCH)
DEUTSCH... 76	NP B103 426	+Deutschemann+ (AACHS, BERL, BONN, CERN+)
ENGLER 74	PR D10 2070	+Kraemer, Toaff, Weisser, Diaz+ (CMU, CASE)
ESTABROOKS 74	NP B79 301	+Martin (DURH)
GRAY 74	NP B75 189	+Hyams, Blum, Dietl+ (CERN, MPIM)
BYERLY 73	PR D7 637	+Anthony, Coffin, Meanley, Meyer, Rice+ (MICH)
GLADDING 73	PR D8 3721	+Russell, Tannenbaum, Weiss, Thomson (HARV)
PROTOPOP... 73	PR D7 1279	+Protopopescu, Alston-Garnjost, Galtieri, Flatte+ (LBL)
BALLAM 72	PR D5 545	+Chadwick, Bingham, Milburn+ (SLAC, LBL, TUFTS)
BENAKSAS 72	PL 39B 289	+Cosme, Jean-Marie, Juillan, Laplanche+ (ORSAY)
JACOBS 72	PR D6 1291	(SACL)
RATCLIFF 72	PL 38B 345	+Bulos, Carnegie, Kluge, Leith, Lynch+ (SLAC)
ABRAMS 71	PR D4 653	+Barnham, Butler, Coyne, Goldhaber, Hall+ (LBL)
ALVENSELEBEN 70	PRL 24 786	+Becker, Bertram, Chen, Cohen (DESY)
BIGGS 70	PRL 24 1197	+Braben, Clifton, Gabathuler, Kitching+ (DARE)
ERBE 69	PR 188 2060	+Hilpert+ (German Bubble Chamber Collab.)
MALAMUD 69	Argonne Conf. 93	+Schlein (UCLA)
REYNOLDS 69	PR 184 1424	+Albright, Bradley, Brucker, Harms+ (FSU)
ROTHWELL 69	PL 23 1521	+Chase, Earles, Gettner, Glass, Weinstein+ (NEAS)
WEHMANN 69	PR 178 2095	+ (HARV, CASE, SLAC, CORN, MCGI)
ARMENISE 68	NC 54A 999	+Ghidini, Forino+ (BARI, BGNA, FIRZ, ORSAY)
BATON 68	PR 176 1574	+Laurens (SACL)
CHUNG 68	PR 165 1491	+Dahl, Kirz, Miller (LRL)
FOSTER 68	NP B6 107	+Gavillet, Labrosse, Montanet+ (CERN, CDEF)
HUSON 68	PL 28B 208	+Lubatti, Six, Veillet+ (ORSAY, MILA, UCLA)
HYAMS 68	NP B7 1	+Koch, Potter, Wilson, VonLindern+ (CERN, MPIM)
LANZEROTTI 68	PR 166 1365	+Blumenthal, Ehn, Faisler+ (HARV)
PISUT 68	NP B6 325	+Roos (CERN)
ASBURY 67B	PRL 19 865	+Becker, Bertram, Joos, Jordan+ (DESY, COLU)
BACON 67	PR 157 1263	+Fickinger, Hill, Hopkins, Robinson+ (BNL)
EISNER 67	PR 164 1699	+Johnson, Klein, Peters, Sahni, Yen+ (PURO)
HUWE 67	PL 24B 252	+Marqui, Oppenheimer, Schultz, Wilson (COLU)
HYAMS 67	PL 24B 534	+Koch, Pellett, Potter, VonLindern+ (CERN, MPIM)
MILLER 67B	PR 153 1423	+Freytag, Johnson, Loeffler+ (PURD)
ALFF... 66	PR 145 1072	+Alff-Steinberger, Berley+ (COLU, RUTG)
FERBEL 66	PL 21 111	(ROCH)
HAGOPIAN 66	PR 145 1128	+Selove, Ailitti, Baton+ (PENN, SACL)
HAGOPIAN 66B	PR 152 1183	+Pan (PENN, LRL)
JACOBS 66B	UCRL 16877	(LRL)
JAMES 66	PR 142 896	+Kraybill (YALE, BNL)
WEST 66	PR 149 1089	+Boyd, Erwin, Walker (WISC)
BLIEDEN 65	PL 19 444	+Freytag, Gebel+ (CERN Missing Mass Spect. Collab.)
CARMONY 64	PRL 12 254	+Lander, Rindfleisch, Xuong, Yager (UCB)
GOLDHABER 64	PRL 12 336	+Brown, Kadyk, Shen+ (LRL, UCB)
ABOLINS 63	PRL 11 381	+Lander, Mehlopp, Nguyen, Yager (UCSD)

OTHER RELATED PAPERS

BENAYOUN 93	ZPHY 58 31	+Feindt, Girono+ (CDEF, CERN, BARI)
LAFFERTY 93	ZPHY C60 659	(MCHS)
Bose-Einstein Correlations		
KAMAL 92	PL B284 421	+Xu (ALBE)
ERKAL 85	ZPHY C29 485	+Olsson (WISC)
RYBICKI 85	ZPHY C28 65	+Sakrejda (CRAC)
KURDADZE 83	JETPL 37 733	+Lelchuk, Pakhtusova+ (NOVO)
ALEKSEEV 82	Translated from ZETFP 37 613.	
	JETP 55 591	+Kartamyshev, Makarin+ (KIAE)
	Translated from ZETF 82 1007.	

KENNEY 62	PR 126 736	+Shephard, Gall (KNTY)
SAMIOS 62	PRL 9 139	+Bachman, Lea+ (BNL, CUNY, COLU, KNTY)
XUONG 62	PR 128 1849	+Lynch (LRL)
ANDERSON 61	PRL 6 365	+Bang, Burke, Carmony, Schmitz (LRL)
ERWIN 61	PRL 6 628	+March, Walker, West (WISC)

$\omega(782)$


$$I^G(J^{PC}) = 0^-(1^--)$$

$\omega(782)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
781.94 ± 0.12 OUR AVERAGE		Error includes scale factor of 1.5. See the ideogram below.		
781.96 ± 0.13 ± 0.17	15k	AMSLER 93B	CBAR	$0.0\bar{p}p \rightarrow \omega\pi^0\pi^0$
782.4 ± 0.2	270k	WEIDENAUER 93	ASTE	$\bar{p}p \rightarrow 2\pi^+2\pi^-\pi^0$
781.78 ± 0.10		BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.2 ± 0.4	1488	KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.4 ± 0.5	7000	¹ KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
783.3 ± 0.4		CORDIER 80	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
782.5 ± 0.8	33260	ROOS 80	RVUE	$0.0-3.6 \bar{p}p$
782.6 ± 0.8	3000	BENKHEIRI 79	OMEG	$9-12 \pi^\pm p$
781.8 ± 0.6	1430	COOPER 78B	HBC	$0.7-0.8 \bar{p}p \rightarrow 5\pi$
782.7 ± 0.9	535	VANAPEL... 78	HBC	$7.2 \bar{p}p \rightarrow \bar{p}\rho\omega$
783.5 ± 0.8	2100	GESSAROLI 77	HBC	$11 \pi^-p \rightarrow \omega n$
782.5 ± 0.8	418	AGUILAR... 72B	HBC	$3.9, 4.6 K^-p$
783.4 ± 1.0	248	BIZZARRI 71	HBC	$0.0 \bar{p}p \rightarrow K^+K^-\omega$
781.0 ± 0.6	510	BIZZARRI 71	HBC	$0.0 \bar{p}p \rightarrow K_1^+K_1^-\omega$
783.7 ± 1.0	3583	² COYNE 71	HBC	$3.7 \pi^+p \rightarrow \rho\pi^+\pi^+\pi^-\pi^0$
784.1 ± 1.2	750	ABRAMOVI... 70	HBC	$3.9 \pi^-p$
783.2 ± 1.6		³ BIGGS 70B	CNTR	$<4.1 \gamma C \rightarrow \pi^+\pi^-C$
782.4 ± 0.5	2400	BIZZARRI 69	HBC	$0.0 \bar{p}p$

- ¹ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.
² From best-resolution sample of COYNE 71.
³ From ω - ρ interference in the $\pi^+\pi^-$ mass spectrum assuming ω width 12.6 MeV.

WEIGHTED AVERAGE
781.94 ± 0.12 (Error scaled by 1.5)



$\omega(782)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.43 ± 0.10 OUR AVERAGE				
8.4 ± 0.1		⁴ AULCHENKO 87	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
8.30 ± 0.40		BARKOV 87	CMD	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.8 ± 0.9	1488	KURDADZE 83B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.0 ± 0.8		CORDIER 80	WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
9.1 ± 0.8		BENAKSAS 72B	OSPK	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •				
12.0 ± 2.0	1430	COOPER 78B	HBC	$0.7-0.8 \bar{p}p \rightarrow 5\pi$
9.4 ± 2.5	2100	GESSAROLI 77	HBC	$11 \pi^-p \rightarrow \omega n$
10.22 ± 0.43	20000	⁵ KEYNE 76	CNTR	$\pi^-p \rightarrow \omega n$
13.3 ± 2	418	AGUILAR... 72B	HBC	$3.9, 4.6 K^-p$
10.5 ± 1.5		BORENSTEIN 72	HBC	$2.18 K^-p$
7.70 ± 0.9 ± 1.15	940	BROWN 72	MMS	$2.5 \pi^-p \rightarrow nMM$
10.3 ± 1.4	510	BIZZARRI 71	HBC	$0.0 \bar{p}p \rightarrow K_1^+K_1^-\omega$
12.8 ± 3.0	248	BIZZARRI 71	HBC	$0.0 \bar{p}p \rightarrow K^+K^-\omega$
9.5 ± 1.0	3583	COYNE 71	HBC	$3.7 \pi^+p \rightarrow \rho\pi^+\pi^+\pi^-\pi^0$

- ⁴ Relativistic Breit-Wigner includes radiative corrections.
⁵ Observed by threshold-crossing technique. Mass resolution = 4.8 MeV FWHM.

Meson Full Listings

 $\omega(782)$ $\omega(782)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\pi^+\pi^-\pi^0$	$(88.8 \pm 0.7)\%$	
Γ_2 $\pi^0\gamma$	$(8.5 \pm 0.5)\%$	
Γ_3 $\pi^+\pi^-$	$(2.21 \pm 0.30)\%$	
Γ_4 neutrals (excluding $\pi^0\gamma$)	$(5.3^{+8.7}_{-3.5}) \times 10^{-3}$	
Γ_5 $\eta\gamma$	$(8.3 \pm 2.1) \times 10^{-4}$	
Γ_6 $\pi^0 e^+ e^-$	$(5.9 \pm 1.9) \times 10^{-4}$	
Γ_7 $\pi^0 \mu^+ \mu^-$	$(9.6 \pm 2.3) \times 10^{-5}$	
Γ_8 $e^+ e^-$	$(7.15 \pm 0.19) \times 10^{-5}$	
Γ_9 $\pi^+\pi^-\pi^0\pi^0$	$< 2\%$	90%
Γ_{10} $\pi^+\pi^-\gamma$	$< 3.6 \times 10^{-3}$	95%
Γ_{11} $\pi^+\pi^-\pi^+\pi^-$	$< 1 \times 10^{-3}$	90%
Γ_{12} $\pi^0\pi^0\gamma$	$< 4 \times 10^{-4}$	90%
Γ_{13} $\mu^+\mu^-$	$< 1.8 \times 10^{-4}$	90%
Γ_{14} $\eta\pi^0$		

CONSTRAINED FIT INFORMATION

An overall fit to 6 branching ratios uses 20 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 10.3$ for 17 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	13		
x_3	-39	-5	
x_4	-74	-68	-1
	x_1	x_2	x_3

 $\omega(782)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	DOCUMENT ID
VALUE (keV)	
0.60 ± 0.02 OUR EVALUATION	

 $\omega(782)$ BRANCHING RATIOS

$\Gamma(\text{neutrals})/\Gamma(\pi^+\pi^-\pi^0)$	EVTs	DOCUMENT ID	TECN	COMMENT	$(\Gamma_2+\Gamma_4)/\Gamma_1$
VALUE					
0.102 ± 0.008 OUR FIT					
$0.103^{+0.011}_{-0.010}$ OUR AVERAGE					
0.15 ± 0.04	46	AGUILAR-...	72B HBC	$3.9, 4.6 K^- p$	
0.10 ± 0.03	19	BARASH	67B HBC	$0.0 \bar{p} p$	
0.134 ± 0.026	850	DIGIUGNO	66B CNTR	$1.4 \pi^- p$	
0.097 ± 0.016	348	FLATTE	66 HBC	$1.4 - 1.7 K^- p \rightarrow \Lambda \text{MM}$	
$0.06^{+0.05}_{-0.02}$		JAMES	66 HBC	$2.1 \pi^+ p$	
0.08 ± 0.03	35	KRAEMER	64 DBC	$1.2 \pi^+ d$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.11 ± 0.02	20	BUSCHBECK	63 HBC	$1.5 K^- p$	

$\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
VALUE				
0.0249 ± 0.0035 OUR FIT				
0.026 ± 0.005 OUR AVERAGE				
$0.021^{+0.028}_{-0.009}$	6	RATCLIFF	72 ASPK	$15 \pi^- p \rightarrow n2\pi$
0.028 ± 0.006		BEHREND	71 ASPK	Photoproduction
$0.022^{+0.009}_{-0.01}$	7	ROOS	70 RVUE	

⁶ Significant interference effect observed. NB of $\omega \rightarrow 3\pi$ comes from an extrapolation.
⁷ ROOS 70 combines ABRAMOVICH 70 and BIZZARRI 70.

$\Gamma(\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE				
0.096 ± 0.006 OUR FIT				
0.096 ± 0.006 OUR AVERAGE				
0.099 ± 0.007		DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\gamma$
0.084 ± 0.013		KEYNE	76 CNTR	$\pi^- p \rightarrow \omega n$
0.109 ± 0.025		BENAKSAS	72C OSPK	e^+e^-
0.081 ± 0.020		BALDIN	71 HLBC	$2.9 \pi^+ p$
0.13 ± 0.04		JACQUET	69B HLBC	

$\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ_1
VALUE				
< 0.066	90	KALBFLEISCH 75	HBC	$2.18 K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$
< 0.05	90	FLATTE	66 HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \pi^+ \pi^- \gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
VALUE				
< 0.0036	95	WEIDENAUER 90	ASTE	$p\bar{p} \rightarrow \pi^+\pi^-\pi^+\pi^-\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
VALUE				
< 0.004	95	BITYUKOV	88B SPEC	$32 \pi^- p \rightarrow \pi^+\pi^-\gamma X$

$\Gamma(\pi^+\pi^-\pi^0\pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
VALUE (units 10^{-2})				
< 2	90	KURDADZE	86 OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0\pi^0$

$\Gamma(\pi^+\mu^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_1
VALUE (units 10^{-3})				
< 0.2	90	WILSON	69 OSPK	$12 \pi^- C \rightarrow Fe$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\pi^0\gamma)$	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ_2
VALUE				
< 1.7	74	FLATTE	66 HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \mu^+ \mu^-$
< 1.2		BARBARO-...	65 HBC	$2.7 K^- p$

$\Gamma(\pi^0\mu^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ_1
VALUE (units 10^{-3})				
< 0.2	90	WILSON	69 OSPK	$12 \pi^- C \rightarrow Fe$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\eta\gamma) + \Gamma(\eta\pi^0\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_5+\Gamma_{14})/\Gamma_1$
VALUE				
< 1.7	74	FLATTE	66 HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \mu^+ \mu^-$
< 1.2		BARBARO-...	65 HBC	$2.7 K^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\eta\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
VALUE				
< 0.005	90	DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$
< 0.18	95	KEYNE	76 CNTR	$\pi^- p \rightarrow \omega n$
< 0.15	90	BENAKSAS	72C OSPK	e^+e^-
< 0.14		BALDIN	71 HLBC	$2.9 \pi^+ p$
< 0.1	90	BARMIN	64 HLBC	$1.3 - 2.8 \pi^- p$

$[\Gamma(\eta\gamma) + \Gamma(\eta\pi^0\gamma)]/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_5+\Gamma_{14})/\Gamma_1$
VALUE				
< 0.016	90	8 FLATTE	66 HBC	$1.2 - 1.7 K^- p \rightarrow \Lambda \pi^+ \pi^- \text{MM}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(\text{neutrals})/\Gamma(\text{charged particles})$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_2+\Gamma_4)/(\Gamma_1+\Gamma_3)$
VALUE				
0.099 ± 0.008 OUR FIT				
0.124 ± 0.021		FELDMAN	67C OSPK	$1.2 \pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^0\mu^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ_1
VALUE				
< 0.00045	90	DOLINSKY	89 ND	$e^+e^- \rightarrow \pi^0\pi^0\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\eta\gamma)/\Gamma(\pi^0\gamma)$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
VALUE				
0.0098 ± 0.0024		ALDE	94 GAM2	$38\pi^- p \rightarrow \omega n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	DOCUMENT ID	TECN	COMMENT	
0.0082 ± 0.0033	9	DOLINSKY	89 ND	$e^+e^- \rightarrow \eta\gamma$
0.039 ± 0.007	10	DOLINSKY	89 ND	e^+e^-
0.010 ± 0.045		APEL	72B OSPK	$4-8 \pi^- p \rightarrow n3\gamma$

⁹ Solution corresponding to constructive ω - ρ interference. The quark model predicts a relative decay phase of zero.

¹⁰ Solution corresponding to destructive ρ - ω interference.

$\Gamma(\pi^0\mu^+\mu^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
VALUE (units 10^{-4})				
0.96 ± 0.23		DZHELJADIN	81B CNTR	$25-33 \pi^- p \rightarrow \omega n$

$\Gamma(\pi^0 e^+ e^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
VALUE (units 10^{-4})				
5.9 ± 1.9	43	DOLINSKY	88 ND	$e^+e^- \rightarrow \pi^0 e^+ e^-$

See key on page 1343

Meson Full Listings

$\omega(782), \eta'(958)$

$\Gamma(e^+e^-)/\Gamma_{total}$

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.715 ± 0.019 OUR AVERAGE				
0.714 ± 0.036		DOLINSKY 89 ND		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.72 ± 0.03		BARKOV 87 CMD		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.66 ± 0.05		KURDADZE 84 OLYA		$e^+e^- \rightarrow$ hadrons
0.675 ± 0.069		CORDIER 80 WIRE		$e^+e^- \rightarrow 3\pi$
0.83 ± 0.10		BENAKSAS 72B OSPK		$e^+e^- \rightarrow 3\pi$
0.77 ± 0.06		11 AUGUSTIN 69D OSPK		$e^+e^- \rightarrow 2\pi$
0.64 ± 0.04	1488	12 KURDADZE 83B OLYA		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
0.65 ± 0.13	33	13 ASTVACAT... 68 OSPK		Assume SU(3)+mixing

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹¹ Rescaled by us to correspond to ω width 8.4 MeV.
¹² Superseded by KURDADZE 84.
¹³ Not resolved from ρ decay. Error statistical only.

$\Gamma(\text{neutrals})/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.090 ± 0.006 OUR FIT				
0.081 ± 0.011 OUR AVERAGE				
0.075 ± 0.025		BIZZARRI 71 HBC		0.0 $\rho\bar{\rho}$
0.079 ± 0.019		DEINET 69B OSPK		1.5 $\pi^-\rho$
0.084 ± 0.015		BOLLINI 68C CNTR		2.1 $\pi^-\rho$
0.073 ± 0.018	42	BASILE 72B CNTR		1.67 $\pi^-\rho$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$

See also $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0221 ± 0.0030 OUR FIT			
0.021 ± 0.004 OUR AVERAGE			
0.023 ± 0.005	BARKOV 85 OLYA		e^+e^-
0.016 +0.009 -0.007	QUENZER 78 CNTR		e^+e^-
0.010 ± 0.001	14 WICKLUND 78 ASPK		3,4,6 $\pi^\pm N$
0.0122 ± 0.0030	ALVENSLEBEN ^{71C} CNTR		Photoproduction
0.013 +0.012 -0.009	MOFFEIT 71 HBC		2.8,4,7 $\gamma\rho$
0.0080 +0.0028 -0.002	15 BIGGS 70B CNTR		4.2 $\gamma C \rightarrow \pi^+\pi^-C$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹⁴ From a model-dependent analysis assuming complete coherence.
¹⁵ Re-evaluated under $\Gamma(\pi^+\pi^-\gamma)/\Gamma(\pi^+\pi^-\pi^0)$ by BEHREND 71 using more accurate $\omega \rightarrow \rho$ photoproduction cross-section ratio.

$\Gamma(\pi^0\pi^0\gamma)/\Gamma(\text{neutrals})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.22 ± 0.07		16 DAKIN 72 OSPK		1.4 $\pi^-\rho \rightarrow nMM$
<0.19	90	DEINET 69B OSPK		

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹⁶ See $\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$.

$\Gamma(\pi^0\gamma)/\Gamma(\text{neutrals})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.78 ± 0.07		17 DAKIN 72 OSPK		1.4 $\pi^-\rho \rightarrow nMM$
>0.81	90	DEINET 69B OSPK		

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹⁷ Error statistical only. Authors obtain good fit also assuming $\pi^0\gamma$ as the only neutral decay.

$\Gamma(\eta\gamma)/\Gamma_{total}$

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
8.3 ± 2.1	ALDE 94 GAM2		38 $\pi^-\rho \rightarrow \omega n$
7.3 ± 2.9	18 DOLINSKY 89 ND		$e^+e^- \rightarrow \eta\eta$
35 ± 5	19 DOLINSKY 89 ND		$e^+e^- \rightarrow \eta\eta$
3.0 +2.5 -1.8	18 ANDREWS 77 CNTR		6.7-10 γCu
29.0 ± 7.0	19 ANDREWS 77 CNTR		6.7-10 γCu

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹⁸ Solution corresponding to constructive ω - ρ interference. The quark model predicts a relative decay phase of zero.
¹⁹ Solution corresponding to destructive ω - ρ interference.

$\Gamma(\pi^0\mu^+\mu^-)/\Gamma(\mu^+\mu^-)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.2 ± 0.6	30	20 DZHELYADIN 79 CNTR		25-33 $\pi^-\rho$

• • • We do not use the following data for averages, fits, limits, etc. • • •

²⁰ Superseded by DZHELYADIN 81B result above.

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.8942 ± 0.0062	DOLINSKY 89 ND		$e^+e^- \rightarrow \pi^+\pi^-\pi^0$

$\omega(782)$ REFERENCES

ALDE 94 ZPHY C61 35	+Binon+ (SERP, LAPP, LANL, BELG, BRUX, CERN)
AMSLER 93B PL B311 362	+Armstrong, Augustin+ (Crystal Barrel Collab.)
WEIDENAUER 93 ZPHY C59 387	+Duch+ (ASTERIX Collab.)
WEIDENAUER 90 ZPHY C47 353	+Duch, Heel, Kalinowsky+ (ASTERIX Collab.)
DOLINSKY 89B ZPHY C42 511	+Druzhinin, Dubrovina, Golubev+ (NOVO)
BITYUKOV 88B SJNP 47 800	+Borison, Viktorov, Golovkin+ (SERP)
DOLINSKY 88 SJNP 48 277	+Druzhinin, Dubrovina, Golubev+ (NOVO)
KURDADZE 88 JETPL 46 164	+Lelchouk, Pakhtusova, Sidorov+ (NOVO)
AULCHENKO 87 PL B186 432	+Dolinsky, Druzhinin, Dubrovina+ (NOVO)
BARKOV 87 JETPL 46 164	+Vasserman, Vorobev, Ivanov (NOVO)
KURDADZE 86 JETPL 43 643	+Lelchouk, Pakhtusova, Sidorov, Skriniskii+ (NOVO)
BARKOV 85 NP 5256 365	+Chilingarov, Eidelman, Khazin, Lelchuk+ (NOVO)
KURDADZE 84 IYF 84-7 Preprint	+Lelchouk, Pakhtusova, Sidorov+ (NOVO)
KURDADZE 83B JETPL 36 274	+Pakhtusova, Sidorov+ (NOVO)
DZHELYADIN 81B PL 102B 296	+Golovkin, Konstantinov+ (SERP)
CORDIER 80 NP B172 13	+Delcourt, Eschstruth, Fuida+ (LALO)
ROOS 80 LNC 27 321	+Pellinen (HELS)
BENKHEIRI 79 NP B150 268	+Eisenstein+ (EPOL, CERN, CDF, LEP)
DZHELYADIN 79 PL 84B 143	+Golovkin, Gritsuk+ (SERP)
COOPER 78B NP B146 1	+Ganguli+ (TATA, CERN, CDF, MADR)
QUENZER 78 PL 76B 512	+Ribes, Rumpf, Bertrand, Bizot, Chase+ (LALO)
VANAPEL... 78 NP B133 245	+VanApeldoorn, Grundeman, Harting+ (ZHEM)
WICKLUND 78 PR D17 1197	+Ayres, Diebold, Greene, Kramer, Pawlicki (ANL)
ANDREWS 77 PRL 38 198	+Fukushima, Harvey, Lobkowicz, May+ (RACH)
CESSAROLI 77 NP B216 382	+Cosme, Jean-Marie, Julian, Laplanche+ (ORSAY)
KEYNE 76B PR D14 28	+Binnie, Carr, Debenham, Garbutt+ (LOIC, SHMP)
Also 73B PR D8 2789	+Binnie, Carr, Debenham, Duane+ (LOIC, SHMP)
KALBFLEISCH 75 PR D11 987	+Strand, Chapman (BNL, MICH)
AGUILAR... 72B PR D6 29	+Aguilar-Benitez, Chung, Eisner, Samios (BNL)
APEL 72B PL 41B 234	+Auslander, Muller, Bertolucci+ (KARLK, KARLE, PISA)
BASILE 72B Phl. Conf. 153	+Bollini, Brogini, Dalpiaz, Frabetti+ (CERN)
BENAKSAS 72B PL 42B 507	+Cosme, Jean-Marie, Julian (ORSAY)
BENAKSAS 72C PL 42B 511	+Cosme, Jean-Marie, Julian, Laplanche+ (ORSAY)
BORENSTEIN 72 PR D5 1559	+Danburg, Kalbfleisch+ (BNL, MICH)
BROWN 72 PL 42B 117	+Downing, Holloway, Huld, Bernstein+ (ILL, ILLC)
DAKIN 72 PR D6 2321	+Hauser, Kreiser, Mischke (PRIN)
RATCLIFF 72 PL 38B 345	+Bulos, Carnegie, Kluge, Leith, Lynch+ (SLAC)
ALVENSLEBEN 71C PRL 27 888	+Becker, Busza, Chen, Cohen+ (DESY)
BALDIN 71 SJNP 13 758	+Yegorov, Trebukhovskiy, Shishov (ITEP)
BEHREND 71 PR D27 61	+Lee, Nordberg, Wehmann+ (ROCH, CORN, FNAL)
BIZZARRI 71 NP B27 140	+Montanet, Nilsson, D'Andlauer+ (CERN, CDF)
COYNE 71 NP B32 333	+Butler, Fng-Landau, MacNaughton (LRL)
MOFFEIT 71 NP B29 349	+Bingham, Fretter+ (LRL, UCB, SLAC, TUFTS)
ABRAMOV... 70 NP B20 209	+Abramovich, Blumenfeld, Bruyat+ (CERN)
BIGGS 70B PRL 24 1201	+Clift, Gabathuler, Kilching, Rand (DARE)
BIZZARRI 70 PRL 25 1385	+Clapetti, Dore, Gaspari, Guidoni+ (ROMA, SYRA)
ROOS 70 DNP/77 173	
Proc. Daresbury Study Weekend No. 1.	
AUGUSTIN 69D PL 28B 513	+Benaksas, Buon, Gracco, Haisinski+ (ORSAY)
BIZZARRI 69 NP B14 169	+Foster, Gavillet, Montanet+ (CERN, CDF)
DEINET 69B PL 30B 426	+Menzione, Muller, Buniatov+ (KARL, CERN)
JACQUET 69B NC 53A 743	+Nguyen-Khac, Haatuft, Halsteinsid (EPOL, BERG)
WILSON 69 Private Comm.	
Also 69 PR 178 2095	Wehmann+ (HARV, CASE, SLAC, CORN, MCGI)
ASTVACAT... 68 PL 27B 45	Astvacaturov, Azimov, Baldin+ (JINR, MOSU)
BOLLINI 68C NC 56A 531	+Buhler, Dalpiaz, Massam+ (CERN, BGNA, STRB)
BARASH 67B PR 156 1399	+Kirsch, Miller, Tan (COLU)
FELDMAN 67C PR 159 1219	+Fratl, Gleason, Halpern, Nussbaum+ (PENN)
DIGIUGNO 66B NC 44A 1272	+Peruzzi, Troise+ (NAPL, FRAS, TRST)
FLATTE 66 PR 145 1050	+Howe, Murray, Button-Shafer, Solmitz+ (LRL)
JAMES 66 PR 142 896	+Kraybill (YALE, BNL)
BARBARO... 65 PRL 14 279	+Barbaro-Galtieri, Tripp (LRL)
BARMIN 64 JETP 18 1289	+Dolgolenko, Krestnikov+ (ITEP)
KRAEMER 64 PR 136B 496	+Madansky, Fields+ (JHU, NWES, WOOD)
BUSCHBECK 63 Siena Conf. 1 166	+Czapp+ (VIEN, CERN, ANIK)

OTHER RELATED PAPERS

DOLINSKY 86 PL B174 453	+Druzhinin, Dubrovina, Eidelman+ (NOVO)
KURDADZE 83 JETPL 37 733	+Lelchuk, Pakhtusova+ (NOVO)
ALF... 62B PRL 9 325	+Alff-Steinberger, Berley, Coiley+ (COLU, RUTG)
ARMENTEROS 62 CERN Conf. 90	+Buddle+ (CERN, CDF, EPOL)
STEVENSON 62 PR 125 687	+Alvarez, Maglich, Rosenfeld (LRL)
MAGLICH 61 PRL 7 178	+Alvarez, Rosenfeld, Stevenson (LRL)
PEVSNER 61 PRL 7 421	+Kraemer, Nussbaum, Richardson+ (JHU)
XUONG 61 PRL 7 327	+Lynch (LRL)

$\eta'(958)$

$I^G(J^{PC}) = 0^+(0^-+)$

Our latest mini-review on this particle can be found in the 1984 edition. See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)

$\eta(958)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
957.77 ± 0.14 OUR AVERAGE				
959 ± 1	630 ± 50	BELADIDZE 92C VES		36 $\pi^- Be \rightarrow \pi^- \eta' Be$
958 ± 1	340	ARMSTRONG 91B OMEG		300 $pp \rightarrow pp\eta^+\pi^-$
958.2 ± 0.4	622	AUGUSTIN 90 DM2		$J/\psi \rightarrow \gamma\eta^+\pi^-$
957.8 ± 0.2	2420	AUGUSTIN 90 DM2		$J/\psi \rightarrow \gamma\eta^+\pi^-$
956.3 ± 1.0	143 ± 12	GIDAL 87 MRK2		$e^+e^- \rightarrow e^+\pi^+\pi^-$
957.46 ± 0.33		DUANE 74 MMS		$\pi^-\rho \rightarrow nMM$
958.2 ± 0.5	1414	DANBURG 73 HBC		2.2 $K^-\rho \rightarrow \Lambda X^0$
958 ± 1	400	JACOBS 73 HBC		2.9 $K^-\rho \rightarrow \Lambda X^0$
956.1 ± 1.1	3415	BASILE 71 CNTR		1.6 $\pi^-\rho \rightarrow nX^0$
957.4 ± 1.4	535	BASILE 71 CNTR		1.6 $\pi^-\rho \rightarrow nX^0$
957 ± 1		RITTENBERG 69 HBC		1.7-2.7 $K^-\rho$

Meson Full Listings

 $\eta'(958)$ $\eta'(958)$ WIDTH

We include direct measurements of the $\eta'(958)$ total width and $\gamma\gamma$ partial width together with the measured branching ratios in the fit for the partial decay rates.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.201 ± 0.016 OUR FIT					Error includes scale factor of 1.3.
0.28 ± 0.10	1000	BINNIE	79 MMS	0	$\pi^- p \rightarrow n \text{MM}$

 $\eta'(958)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\pi^+\pi^-\eta$	(43.7 ± 1.5) %	S=1.1
Γ_2 $\rho^0\gamma$	(30.2 ± 1.3) %	S=1.1
Γ_3 $\pi^0\pi^0\eta$	(20.8 ± 1.3) %	S=1.2
Γ_4 $\omega\gamma$	(3.02 ± 0.30) %	
Γ_5 $\gamma\gamma$	(2.12 ± 0.13) %	S=1.2
Γ_6 $3\pi^0$	(1.55 ± 0.26) × 10 ⁻³	
Γ_7 $\mu^+\mu^-\gamma$	(1.04 ± 0.26) × 10 ⁻⁴	
Γ_8 $\pi^+\pi^-\pi^0$	< 5 %	CL=90%
Γ_9 $\pi^0\rho^0$	< 4 %	CL=90%
Γ_{10} $\pi^+\pi^-$	< 2 %	CL=90%
Γ_{11} $\pi^0e^+e^-$	< 1.3 %	CL=90%
Γ_{12} ηe^+e^-	< 1.1 %	CL=90%
Γ_{13} $\pi^+\pi^+\pi^-\pi^-$	< 1 %	CL=90%
Γ_{14} $\pi^+\pi^+\pi^-\pi^- \text{ neutrals}$	< 1 %	CL=95%
Γ_{15} $\pi^+\pi^+\pi^-\pi^-\pi^0$	< 1 %	CL=90%
Γ_{16} 6π	< 1 %	CL=90%
Γ_{17} $\pi^+\pi^-e^+e^-$	< 6 × 10 ⁻³	CL=90%
Γ_{18} $\pi^0\pi^0$	< 9 × 10 ⁻⁴	CL=90%
Γ_{19} $\pi^0\gamma\gamma$	< 8 × 10 ⁻⁴	CL=90%
Γ_{20} $4\pi^0$	< 5 × 10 ⁻⁴	CL=90%
Γ_{21} 3γ	< 1.0 × 10 ⁻⁴	CL=90%
Γ_{22} $\mu^+\mu^-\pi^0$	< 6.0 × 10 ⁻⁵	CL=90%
Γ_{23} $\mu^+\mu^-\eta$	< 1.5 × 10 ⁻⁵	CL=90%
Γ_{24} $\pi^+\pi^-\gamma$ (including $\rho^0\gamma$)	(27.9 ± 2.3) %	
Γ_{25} e^+e^-	< 2.1 × 10 ⁻⁷	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, 2 combinations of partial widths obtained from integrated cross section, and 16 branching ratios uses 45 measurements and one constraint to determine 7 parameters. The overall fit has a $\chi^2 = 33.4$ for 39 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-49					
x_3	-63	-35				
x_4	-27	-25	34			
x_5	-22	-13	27	8		
x_6	-23	-13	36	12	10	
Γ	35	-11	-21	-3	-83	-7
	x_1	x_2	x_3	x_4	x_5	x_6

Mode	Rate (MeV)	Scale factor
Γ_1 $\pi^+\pi^-\eta$	0.088 ± 0.009	1.2
Γ_2 $\rho^0\gamma$	0.061 ± 0.005	1.3
Γ_3 $\pi^0\pi^0\eta$	0.042 ± 0.004	1.5
Γ_4 $\omega\gamma$	0.0061 ± 0.0008	1.2
Γ_5 $\gamma\gamma$	0.00426 ± 0.00019	1.1
Γ_6 $3\pi^0$	(3.1 ± 0.6) × 10 ⁻⁴	1.1

 $\eta'(958)$ PARTIAL WIDTHS $\Gamma(\gamma\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.26 ± 0.19 OUR FIT				Error includes scale factor of 1.1.
4.34 ± 0.25 OUR AVERAGE				
4.53 ± 0.29 ± 0.51	266 ± 17	KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
3.62 ± 0.14 ± 0.48		¹ BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\eta'(958)$
4.6 ± 1.1 ± 0.6	23	BARU	90 MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
4.57 ± 0.25 ± 0.44		BUTLER	90 MRK2	$e^+e^- \rightarrow e^+e^-\eta'(958)$
4.94 ± 0.23 ± 0.72	547	² ROE	90 ASP	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$
3.8 ± 0.7 ± 0.6	34	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.8 ± 0.5 ± 0.5	136 ± 14	² WILLIAMS	88 CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
4.7 ± 0.6 ± 0.9	143 ± 12	³ GIDAL	87 MRK2	$e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
4.0 ± 0.9		⁴ BARTEL	85E JADE	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

¹ Using $B(\eta' \rightarrow \rho(770)\gamma) = (30.1 \pm 1.4)\%$.

² Using $B(\eta' \rightarrow \gamma\gamma) = (2.17 \pm 0.17)\%$.

³ Superseded by BUTLER 90.

⁴ Systematic error not evaluated.

 $\eta'(958) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

This combination of a partial width with the partial width into $\gamma\gamma$ and with the total width is obtained from the integrated cross section into channel(i) in the $\gamma\gamma$ annihilation.

 $\Gamma(\gamma\gamma) \times \Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.29 ± 0.06 OUR FIT				Error includes scale factor of 1.2.
1.26 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.2.
1.09 ± 0.04 ± 0.13		BEHREND	91 CELL	$e^+e^- \rightarrow e^+e^-\rho(770)^0\gamma$
1.35 ± 0.09 ± 0.21		AIHARA	87 TPC	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.13 ± 0.04 ± 0.13	867 ± 30	ALBRECHT	87B ARG	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.53 ± 0.09 ± 0.21		ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.14 ± 0.08 ± 0.11	243 ± 16.5	BERGER	84B PLUT	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.73 ± 0.34 ± 0.35	95	JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^-\rho\gamma$
1.49 ± 0.13 ± 0.027	213	BARTEL	82B JADE	$e^+e^- \rightarrow e^+e^-\rho\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.85 ± 0.31 ± 0.24	43	BEHREND	83B CELL	$e^+e^- \rightarrow e^+e^-\rho\gamma$

 $\Gamma(\gamma\gamma) \times \Gamma(\pi^0\pi^0\eta)/\Gamma_{\text{total}}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.88 ± 0.07 OUR FIT			Error includes scale factor of 1.1.
0.93 ± 0.06 ± 0.11	⁵ KARCH	92 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.95 ± 0.05 ± 0.08	⁶ KARCH	90 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
1.00 ± 0.08 ± 0.10	^{5,6} ANTREASYAN	87 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi^0\pi^0$
⁵ Using $BR(\eta \rightarrow 2\gamma) = (38.9 \pm 0.5)\%$.			
⁶ Superseded by KARCH 92.			

 $\eta'(958) \alpha$ PARAMETER

$$|\text{MATRIX ELEMENT}|^2 = (1 + \alpha\gamma)^2 + \alpha^2$$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.058 ± 0.013	⁷ ALDE	86 GAM2	$38 \pi^- p \rightarrow n \eta 2\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.08 ± 0.03	⁷ KALBFLEISCH	74 RVUE	$\eta' \rightarrow \eta\pi^+\pi^-$
⁷ May not necessarily be the same for $\eta' \rightarrow \eta\pi^+\pi^-$ and $\eta' \rightarrow \eta\pi^0\pi^0$.			

 $\eta'(958)$ BRANCHING RATIOS

$\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	0.310 ± 0.011 OUR FIT				Error includes scale factor of 1.1.
	0.314 ± 0.026	281	RITTENBERG	69 HBC	1.7-2.7 $K^- p$

 $\Gamma(\pi^+\pi^-\text{neutrals})/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.398 ± 0.009 OUR FIT				Error includes scale factor of 1.1.
0.36 ± 0.05 OUR AVERAGE				
0.4 ± 0.1	39	LONDON	66 HBC	2.24 $K^- p \rightarrow \Lambda\pi^+\pi^-\text{neutrals}$
0.35 ± 0.06	33	BADIER	65B HBC	3 $K^- p$

See key on page 1343

Meson Full Listings
 $\eta'(958)$ $\Gamma(\pi^+\pi^-\eta(\text{charged decay}))/\Gamma_{\text{total}}$ $0.291\Gamma_1/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.127±0.004 OUR FIT	Error includes scale factor of 1.1.			
0.116±0.013 OUR AVERAGE				
0.123±0.014	107	RITTENBERG 69	HBC	1.7-2.7 K^-p
0.10 ±0.04	10	LONDON 66	HBC	2.24 $K^-p \rightarrow \Lambda\pi^+\pi^-\pi^+\pi^-\pi^0$
0.07 ±0.04	7	BADIER 658	HBC	3 K^-p

 $[\Gamma(\pi^0\pi^0\eta(\text{charged decay})) + \Gamma(\omega(\text{charged decay})\gamma)]/\Gamma_{\text{total}}$ $(0.291\Gamma_3+0.9\Gamma_4)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.088±0.005 OUR FIT	Error includes scale factor of 1.2.			
0.045±0.029	42	RITTENBERG 69	HBC	1.7-2.7 K^-p

 $\Gamma(\text{neutrals})/\Gamma_{\text{total}}$ $(0.709\Gamma_3+0.09\Gamma_4+\Gamma_5)/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.171±0.009 OUR FIT	Error includes scale factor of 1.2.			
0.187±0.017 OUR AVERAGE				
0.185±0.022	535	BASILE 71	CNTR	1.6 $\pi^-p \rightarrow n\chi^0$
0.189±0.026	123	RITTENBERG 69	HBC	1.7-2.7 K^-p

 $\Gamma(\rho^0\gamma)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.302±0.013 OUR FIT	Error includes scale factor of 1.1.			
0.319±0.030 OUR AVERAGE				
0.329±0.033	298	RITTENBERG 69	HBC	1.7-2.7 K^-p
0.2 ±0.1	20	LONDON 66	HBC	2.24 $K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$
0.34 ±0.09	35	BADIER 658	HBC	3 K^-p

 $\Gamma(\rho^0\gamma)/\Gamma(\pi\pi\eta)$ $\Gamma_2/(\Gamma_1+\Gamma_3)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.468±0.029 OUR FIT	Error includes scale factor of 1.1.			
0.31 ±0.15	68	DAVIS 68	HBC	5.5 K^-p

 $\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.013	90	RITTENBERG 65	HBC	2.7 K^-p

 $\Gamma(\eta e^+e^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011	90	RITTENBERG 65	HBC	2.7 K^-p

 $\Gamma(\pi^0\rho^0)/\Gamma_{\text{total}}$ Γ_9/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	90	RITTENBERG 65	HBC	2.7 K^-p

 $\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.006	90	RITTENBERG 65	HBC	2.7 K^-p

 $\Gamma(6\pi)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	90	LONDON 66	HBC	Compilation

 $\Gamma(\omega\gamma)/\Gamma(\pi^+\pi^-\eta)$ Γ_4/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.069±0.008 OUR FIT	Error includes scale factor of 1.1.			
0.068±0.013	68	ZANFINO 77	ASPK	8.4 π^-p

 $\Gamma(\rho^0\gamma)/[\Gamma(\pi^+\pi^-\eta) + \Gamma(\pi^0\pi^0\eta) + \Gamma(\omega\gamma)]$ $\Gamma_2/(\Gamma_1+\Gamma_3+\Gamma_4)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.447±0.028 OUR FIT	Error includes scale factor of 1.1.			
0.25 ±0.14	64	DAUBER 64	HBC	1.95 K^-p

 $\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0212±0.0013 OUR FIT	Error includes scale factor of 1.2.			
0.0196±0.0015 OUR AVERAGE				
0.0200±0.0018		⁸ STANTON 80	SPEC	8.45 $\pi^-p \rightarrow n\pi^+\pi^-2\gamma$
0.025 ±0.007		DUANE 74	MMS	$\pi^-p \rightarrow nMM$
0.0171±0.0033	68	DALPIAZ 72	CNTR	1.6 $\pi^-p \rightarrow n\chi^0$
0.020 ±0.008	31	HARVEY 71	OSPK	3.65 $\pi^-p \rightarrow n\chi^0$
0.018 ±0.002	6000	⁹ APEL 79	NICE	15-40 $\pi^-p \rightarrow n2\gamma$

⁸ Includes APEL 79 result.
⁹ Data is included in STANTON 80 evaluation.

 $\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	COMMENT
<2.1	90	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^+\pi^-\eta$

 $\Gamma(\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{10}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.02	90	RITTENBERG 69	HBC	1.7-2.7 K^-p
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.08	95	DANBURG 73	HBC	2.2 $K^-p \rightarrow \Lambda\chi^0$

 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	RITTENBERG 69	HBC	1.7-2.7 K^-p
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.09	95	DANBURG 73	HBC	2.2 $K^-p \rightarrow \Lambda\chi^0$

 $\Gamma(\pi^+\pi^+\pi^-\text{neutrals})/\Gamma_{\text{total}}$ Γ_{14}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	95	DANBURG 73	HBC	2.2 $K^-p \rightarrow \Lambda\chi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.01	90	RITTENBERG 69	HBC	1.7-2.7 K^-p

 $\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	90	RITTENBERG 69	HBC	1.7-2.7 K^-p

 $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{13}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.01	90	RITTENBERG 69	HBC	1.7-2.7 K^-p

 $\Gamma(\rho^0\gamma)/\Gamma(\pi^+\pi^-\gamma(\text{including}\rho^0\gamma))$ Γ_2/Γ_{24}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.08±0.08 OUR AVERAGE				
1.15±0.10	473	DANBURG 73	HBC	2.2 $K^-p \rightarrow \Lambda\chi^0$
1.01±0.15	137	JACOBS 73	HBC	2.9 $K^-p \rightarrow \Lambda\chi^0$
0.94±0.20		AGUILAR-... 70D	HBC	3.9-4.6 K^-p

 $\Gamma(\pi^0\pi^0\eta(3\pi^0\text{decay}))/\Gamma_{\text{total}}$ $0.319\Gamma_3/\Gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.066±0.004 OUR FIT	Error includes scale factor of 1.2.			
0.11 ±0.06	4	BENSINGER 70	DBC	2.2 π^+d

 $\Gamma(\rho^0\gamma)/\Gamma(\pi^+\pi^-\eta(\text{neutral decay}))$ $\Gamma_2/0.709\Gamma_1$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.97±0.07 OUR FIT	Error includes scale factor of 1.1.			
1.01±0.09 OUR AVERAGE				
1.07±0.17		BELADIDZE 92C	VES	36 $\pi^-Be \rightarrow \pi^-\eta^0\eta Be$
0.92±0.14	473	DANBURG 73	HBC	2.2 $K^-p \rightarrow \Lambda\chi^0$
1.11±0.18	192	JACOBS 73	HBC	2.9 $K^-p \rightarrow \Lambda\chi^0$

 $\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta(\text{neutral decay}))$ $\Gamma_5/0.709\Gamma_3$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.144±0.010 OUR FIT	Error includes scale factor of 1.6.			
0.188±0.068	16	APEL 72	OSPK	3.8 $\pi^-p \rightarrow n\chi^0$

 $\Gamma(\mu^+\mu^-)/\Gamma(\gamma\gamma)$ Γ_7/Γ_5

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
4.9±1.2	33	VIKTOROV 80	CNTR	25,33 $\pi^-p \rightarrow 2\mu\gamma$

 $\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<1.5	90	DZHELADIN 81	CNTR	30 $\pi^-p \rightarrow \eta^0 n$

 $\Gamma(\mu^+\mu^-\pi^0)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	COMMENT
<6.0	90	DZHELADIN 81	CNTR	30 $\pi^-p \rightarrow \eta^0 n$

 $\Gamma(3\pi^0)/\Gamma(\pi^0\pi^0\eta)$ Γ_6/Γ_3

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
74±12 OUR FIT			
74±12 OUR AVERAGE			
74±15	ALDE 87B	GAM2	38 $\pi^-p \rightarrow n6\gamma$
75±18	BINON 84	GAM2	30-40 $\pi^-p \rightarrow n6\gamma$

 $\Gamma(\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_5/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.102±0.007 OUR FIT	Error includes scale factor of 1.6.		
0.105±0.010 OUR AVERAGE	Error includes scale factor of 1.9.		
0.091±0.009	AMSLER 93	CBAR	0.0 $\bar{p}p$
0.112±0.002±0.006	ALDE 87B	GAM2	38 $\pi^-p \rightarrow n2\gamma$

 $\Gamma(\omega\gamma)/\Gamma(\pi^0\pi^0\eta)$ Γ_4/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
0.145±0.014 OUR FIT			
0.147±0.016	ALDE 87B	GAM2	38 $\pi^-p \rightarrow n4\gamma$

Meson Full Listings

$\eta'(958), f_0(980)$

$\Gamma(3\gamma)/\Gamma(\pi^0\pi^0\eta)$		Γ_{21}/Γ_3	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN COMMENT
<4.6	90	ALDE	87B GAM2 38 $\pi^- \rho \rightarrow n3\gamma$

$\Gamma(\pi^0\gamma\gamma)/\Gamma(\pi^0\pi^0\eta)$		Γ_{19}/Γ_3	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN COMMENT
<37	90	ALDE	87B GAM2 38 $\pi^- \rho \rightarrow n4\gamma$

$\Gamma(\pi^0\pi^0)/\Gamma(\pi^0\pi^0\eta)$		Γ_{18}/Γ_3	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN COMMENT
<45	90	ALDE	87B GAM2 38 $\pi^- \rho \rightarrow n4\gamma$

$\Gamma(4\pi^0)/\Gamma(\pi^0\pi^0\eta)$		Γ_{20}/Γ_3	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN COMMENT
<23	90	ALDE	87B GAM2 38 $\pi^- \rho \rightarrow n8\gamma$

$\eta'(958)$ C-NONCONSERVING DECAY PARAMETER

See the note on η decay parameters in the Stable Particle Full Listings for definition of this parameter.

DECAY ASYMMETRY PARAMETER FOR $\pi^+\pi^-\gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.01 ± 0.04	OUR AVERAGE			
-0.019 ± 0.056		AIHARA 87	TPC	$2\gamma \rightarrow \pi^+\pi^-\gamma$
-0.069 ± 0.078	295	GRIGORIAN 75	STRC	$2.1 \pi^- \rho$
0.00 ± 0.10	103	KALBFLEISCH 75	HBC	$2.18 K^- \rho \rightarrow \Delta\pi^+\pi^-\gamma$
0.07 ± 0.08	152	RITTENBERG 65	HBC	$2.1-2.7 K^- \rho$

$\eta'(958)$ REFERENCES

AMSLER 93	ZPHY C58 175	+Armstrong, Augustin+	(Crystal Barrel Collab.)
BELADIDZE 92C	SJNP 55 1535	+Bityukov, Borisov	(VES Collab.)
	Translated from YAF 55 2748		
KARCH 92	ZPHY C54 33	+Antreasyan, Bartels+	(Crystal Ball Collab.)
ARMSTRONG 91B	ZPHY C52 389	+Barnes+ (ATHU, BARI, BIRM, CERN, CDEF)	
BEHREND 91	ZPHY C49 401	+Criegee, Field, Franke+	(CELLO Collab.)
AUGUSTIN 90	PR D42 10	+Cosme+	(DM2 Collab.)
BARU 90	ZPHY C48 581	+Blinov, Blinov+	(MD-1 Collab.)
BUTLER 90	PR D42 1368	+Boyer+	(Mark II Collab.)
KARCH 90	PL B249 353	+Antreasyan, Bartels+	(Crystal Ball Collab.)
ROE 90	PR D41 17	+Bartha, Burke, Garbincius+	(ASP Collab.)
AIHARA 88C	PR D38 1	+Alston-Garnjost+	(TPC-2 γ Collab.)
VOROBYEV 88	SJNP 48 273	+Golubev, Dolinsky, Druzhinin+	(NOVO)
	Translated from YAF 48 436		
WILLIAMS 88	PR D38 1365	+Antreasyan, Bartels, Besset+	(Crystal Ball Collab.)
AIHARA 87	PR D35 2650	+Alston-Garnjost+	(TPC-2 γ Collab.) JP
ALBRECHT 87B	PL B199 457	+Andam, Binder+	(ARGUS Collab.)
ALDE 87B	ZPHY C36 603	+Binon, Bricman+ (LANL, BELG, SERP, LAPP)	
ANTREASYAN 87	PR D36 2633	+Bartels, Besset+	(Crystal Ball Collab.)
GIDAL 87	PRL 59 2012	+Boyer, Butler, Cords, Abrams+ (LBL, SLAC, HARV)	
ALDE 86	PL B177 115	+Binon, Bricman+ (SERP, BELG, LANL, LAPP)	
BARTELL 85E	PL 160B 421	+Becker, Cords, Felst+	(JADE Collab.)
ALTHOFF 87	PL 147B 467	+Braunschweig, Kirschfink, Luebelsmeyer+	(TASSO Collab.)
BERGER 84B	PL 142B 125		(PLUTO Collab.)
BINON 84	PL 140B 264	+Donskov, Duteil+ (SERP, BELG, LAPP, CERN)	
BEHREND 83B	PL 125B 518	+D'Agostini+ (CELLO Collab.)	
Also 82C	PL 114B 378	+Behrend, Chen, Fenner, Field+ (CELLO Collab.)	
JENNI 83	PR D27 1031	+Burke, Telnov, Abrams, Blocker+ (SLAC, LBL)	
BARTELL 82B	PL 113B 190	+Cords+ (JADE Collab.)	
DZHEL'YADIN 81	PL 105B 239	+Golovkin, Konstantinov, Kubarovski+ (SERP)	
STANTON 80	PL 92 B 353	+Edwards, Legacey+ (OSU, CARL, MCGI, TINTO)	
VIKTOROV 80	SJNP 32 520	+Golovkin, Dzheiyadin, Zaitsev, Mukhin+ (SERP)	
	Translated from YAF 32 1005		
APEL 79	PL 83B 131	+Augenstein, Bertolucci+(KARLK, KARLE, PISA, SERP, WIEN)	
BINNIE 79	PL 83B 141	+Carr, Debenham, Jones, Karami, Keyne+ (LOIC)	
ZANFINO 77	PRL 38 930	+Brockman+ (CARL, MCGI, OHIO, TINTO)	
GRIGORIAN 75	NP B91 232	+Ladage, Mellema, Rudnick+ (UCLA)	
KALBFLEISCH 75	PR D11 987	+Strand, Chapman (BNL, MICH)	
DUANE 74	PRL 32 425	+Binnie, Camilleri, Carr+ (LOIC, SHMP)	
KALBFLEISCH 74	PR D10 916		(BNL)
DANBURG 73	PR D8 3744	+Kalbfleisch, Borenstein, Chapman+ (BNL, MICH) JP	
JACOBS 73	PR D8 18	+Chang, Gauthier+ (BRAN, UMD, SYRA, TUFTS) JP	
APEL 72	PL 40B 680	+Auslander, Muller, Bertolucci+ (KARLK, KARLE, PISA)	
DALPIAZ 72	PL 42B 377	+Frabetti, Massam, Navarria, Zichichi (CERN)	
BASILE 71	PL 34 371	+Bolini, Dalpiaz, Frabetti+ (CERN, BGNA, STRB)	
HARVEY 71	PRL 27 885	+Marquitt, Peterson, Rhoades+ (MINN, MICH)	
AGUILAR... 70D	PRL 25 1635	+Aguilar-Benitez, Bassano, Samios, Barnes+ (BNL)	
BENSINGER 70	PL 33B 505	+Erwin, Thompson, Walker (WISC)	
RITTENBERG 69	UCRL 18863 Thesis		(LRL) I
DAVIS 68	PL 27B 532	+Ammar, Mott, Dagan, Derrick+ (NWES, ANL)	
LONDON 66	PR 143 1034	+Rau, Goldberg, Lichtman+ (BNL, SYRA) JP	
BADIER 65B	PRL 17 3556	+Demoulin, Barlioutaud+ (EPOL, SACL, AMST)	
RITTENBERG 65	PRL 15 556	+Kalbfleisch (LRL, BNL)	
DAUBER 64	PRL 13 449	+Slater, Smith, Stork, Ticho (UCLA) JP	
Also 64B	Dubna Conf. 1 418	+Dauber, Slater, Smith, Ticho (UCLA)	

OTHER RELATED PAPERS

BENAYOUN 93	ZPHY 58 31	+Feindt, Girona+ (CDEF, CERN, BARI)	
KAMAL 92	PL B284 421	+Xu (ALBE)	
BICKERSTAFF 82	ZPHY C16 171	+McKellar (MELB)	
KIENZLE 65	PL 19 438	+Maglich, Levrat, Lefebvres+ (CERN)	
TRILLING 65	PL 19 427	+Brown, Goldhaber, Kadyk, Scania (LRL)	
GOLDBERG 64	PRL 12 546	+Gundzik, Lichtman, Connolly, Hart+ (SYRA, BNL)	
GOLDBERG 64B	PRL 13 249	+Gundzik, Leitner, Connolly, Hart+ (SYRA, BNL)	
KALBFLEISCH 64	PRL 12 527	+Alvarez, Barbaro-Galtieri+ (LRL) JP	
KALBFLEISCH 64B	PRL 13 349	+Dahl, Rittenberg (LRL) JP	

$f_0(980)$

was S(975)

$$I^G(J^{PC}) = 0^+(0^{++})$$

For early work using Breit-Wigner or scattering length parametrization in fits to the $K\bar{K}$ mass spectrum, see reference section and our 1972 edition.

See also the mini-review under $f_0(1300)$ and the non- $q\bar{q}$ candidates. (See the index for the page number.)

$f_0(980)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
980 ± 10 OUR ESTIMATE			
988 ± 10	1 MORGAN 93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_S \rightarrow \pi(\pi\pi)$
~ 978	2 MORGAN 93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_S \rightarrow \pi(\pi\pi)$
~ 989	1 ZOU 93	RVUE	
797-914	2 ZOU 93	RVUE	
971.1 ± 4.0	3 AGUILAR... 91	EHS	400pp
979 ± 4	4 ARMSTRONG 91	OMEG	300 pp → ppππ, ppK K
956 ± 12	BREAKSTONE 90	SFM	pp → ppπ ⁺ π ⁻
959.4 ± 6.5	3 AUGUSTIN 89	DM2	J/ψ → ωπ ⁺ π ⁻
978 ± 9	3 ABACHI 86B	HRS	e ⁺ e ⁻ → π ⁺ π ⁻ X
985.0 ± 9.0 -39.0	ETKIN 82B	MPS	23 π ⁻ ρ → n2K _S ⁰
985	5 TORNQVIST 82	RVUE	
974 ± 4	4 GIDAL 81	MRK2	J/ψ → π ⁺ π ⁻ X
975	5 ACHASOV 80	RVUE	
986 ± 10	4 AGUILAR... 78	HBC	0.7 p̄p → K _S ⁰ K _S ⁰
969 ± 5	4 LEEPER 77	ASPK	2-2.4 π ⁻ ρ → π ⁺ π ⁻ n, K ⁺ K ⁻ n
987 ± 7	4 BINNIE 73	CNTR	π ⁻ ρ → nMM
1012 ± 6	6 GRAYER 73	ASPK	17 π ⁻ ρ → π ⁺ π ⁻ n
1007 ± 20	6 HYAMS 73	ASPK	17 π ⁻ ρ → π ⁺ π ⁻ n
997 ± 6	6 PROTOPOP... 73	HBC	7 π ⁺ ρ → π ⁺ ρπ ⁺ π ⁻

- 1 On sheet II in a 2 pole solution.
- 2 On sheet III in a 2 pole solution.
- 3 From invariant mass fit.
- 4 From coupled channel analysis.
- 5 Coupled channel analysis with finite width corrections.
- 6 Included in AGUILAR-BENITEZ 78 fit.

$f_0(980)$ WIDTH

Width determination very model dependent. Peak width is about 50 MeV, but decay width can be much larger.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40 to 400 OUR ESTIMATE			
48 ± 12	7 MORGAN 93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_S \rightarrow \pi(\pi\pi)$
56 ± 12	8 MORGAN 93	RVUE	$\pi\pi(K\bar{K}) \rightarrow \pi\pi(K\bar{K}), J/\psi \rightarrow \phi\pi\pi(K\bar{K}), D_S \rightarrow \pi(\pi\pi)$
46-50	7 ZOU 93	RVUE	
370-438	8 ZOU 93	RVUE	
37.4 ± 10.6	9 AGUILAR... 91	EHS	400pp
72 ± 8	10 ARMSTRONG 91	OMEG	300 pp → ppππ, ppK K
110 ± 30	BREAKSTONE 90	SFM	pp → ppπ ⁺ π ⁻
29 ± 13	9 ABACHI 86B	HRS	e ⁺ e ⁻ → π ⁺ π ⁻ X
120 ± 281 ± 20	ETKIN 82B	MPS	23 π ⁻ ρ → n2K _S ⁰
~ 400	11 TORNQVIST 82	RVUE	
28 ± 10	10 GIDAL 81	MRK2	J/ψ → π ⁺ π ⁻ X
70 to 300	12 ACHASOV 80	RVUE	
100 ± 60	13 AGUILAR... 78	HBC	0.7 p̄p → K _S ⁰ K _S ⁰
30 ± 8	10 LEEPER 77	ASPK	2-2.4 π ⁻ ρ → π ⁺ π ⁻ n, K ⁺ K ⁻ n
48 ± 14	10 BINNIE 73	CNTR	π ⁻ ρ → nMM
32 ± 10	14 GRAYER 73	ASPK	17 π ⁻ ρ → π ⁺ π ⁻ n
30 ± 10	14 HYAMS 73	ASPK	17 π ⁻ ρ → π ⁺ π ⁻ n
54 ± 16	14 PROTOPOP... 73	HBC	7 π ⁺ ρ → π ⁺ ρπ ⁺ π ⁻

- 7 On sheet II in a 2 pole solution.
- 8 On sheet III in a 2 pole solution.
- 9 From invariant mass fit.
- 10 From coupled channel analysis.
- 11 Total Breit-Wigner width in coupled channel analysis, but peak width in ππ 50 MeV.
- 12 Coupled channel analysis with finite width corrections.
- 13 From coupled channel fit to the HYAMS 73 and PROTOPOPESCU 73 data. With a simultaneous fit to the ππ phase-shifts, inelasticity and to the K_S⁰K_S⁰ invariant mass.
- 14 included in AGUILAR-BENITEZ 78 fit.

Meson Full Listings

$f_0(980), a_0(980)$

$f_0(980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \quad \pi\pi$	$(78.1 \pm 2.4) \%$	
$\Gamma_2 \quad K\bar{K}$	$(21.9 \pm 2.4) \%$	
$\Gamma_3 \quad \gamma\gamma$	$(1.19 \pm 0.33) \times 10^{-5}$	
$\Gamma_4 \quad e^+e^-$	$< 3 \times 10^{-7}$	90%

CONSTRAINED FIT INFORMATION

An overall fit to a branching ratio uses 3 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 2.0$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_2 \begin{vmatrix} -100 \\ x_1 \end{vmatrix}$$

$f_0(980)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3
0.56 ± 0.11 OUR AVERAGE						
0.63 ± 0.14		15	MORGAN	90	RVUE $\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
0.42 ± 0.06 ± 0.18		60 ± 8	OEST	90	JADE $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29 ± 0.07 ± 0.12		17,18	BOYER	90	MRK2 $e^+e^- \rightarrow \pi^+\pi^-, \pi^0\pi^0$	
0.31 ± 0.14 ± 0.09		17,18	MARSISKE	90	CBAL $e^+e^- \rightarrow e^+e^-\pi^0\pi^0$	

¹⁵ From amplitude analysis of BOYER 90 and MARSISKE 90, data corresponds to resonance parameters $m = 989$ MeV, $\Gamma = 61$ MeV.

¹⁶ OEST 90 quote systematic errors $^{+0.08}_{-0.18}$. We use ± 0.18 .

¹⁷ From analysis allowing arbitrary background unconstrained by unitarity.

¹⁸ Data included in MORGAN 90 analysis.

$\Gamma(e^+e^-)$	VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4
< 8.4		90	VOROBYEV	88	ND $e^+e^- \rightarrow \pi^0\pi^0$	

$f_0(980)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/[\Gamma(\pi\pi) + \Gamma(K\bar{K})]$	VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/(\Gamma_1 + \Gamma_2)$	
0.781 ± 0.024 OUR FIT						
0.781 $^{+0.027}_{-0.023}$ OUR AVERAGE						
0.67 ± 0.09		19	LOVERRE	80	HBC $4\pi^-p \rightarrow n2K_S^0$	
0.81 $^{+0.09}_{-0.04}$		19	CASON	78	STRC $7\pi^-p \rightarrow n2K_S^0$	
0.78 ± 0.03		19	WETZEL	76	OSPK $8.9\pi^-p \rightarrow n2K_S^0$	

¹⁹ Measure $\pi\pi$ elasticity assuming two resonances coupled to the $\pi\pi$ and $K\bar{K}$ channels only.

$f_0(980)$ REFERENCES

MORGAN	93	PR D48 1185	+Pennington	(RAL, DURH)
ZOU	93	PR D48 R3948	+Bugg	(LOQM)
AGUILAR...	91	ZPHY C50 405	+Aguilar-Benitez, Allison, Batalor+	(LEBC-EHS Collab.)
ARMSTRONG	91	ZPHY C51 351	+Benayoun+	(ATHU, BARI, BIRM, CERN, CDEF)
BOYER	90	PR D42 1350	+Butler+	(Mark II Collab.)
BREAKSTONE	90	ZPHY C48 569	+ (ISU, BGNA, CERN, DORT, HEIDH, WARS)	
MARSISKE	90	PR D41 3324	+Antreasyan+	(Crystal Ball Collab.)
MORGAN	90	ZPHY C48 623	+Pennington	(RAL, DURH)
OEST	90	ZPHY C47 343	+Olsson+	(JADE Collab.)
AUGUSTIN	89	NP B320 1	+Cosme	(DM2 Collab.)
VOROBYEV	88	SJNP 48 273	+Golubev, Dolinsky, Druzhinin+	(NOVO)
		Translated from YAF 48 436.		
ABACHI	86B	PRL 57 1990	+Derrick, Blockus+	(PURD, ANL, IND, MICH, LBL)
ETKIN	82B	PR D25 1786	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
TORNQVIST	82	PRL 49 524	+ (HEL)	
GIDAL	81	PL 107B 153	+Goldhaber, Guy, Millikan, Abrams+	(SLAC, LBL)
ACHASOV	80	SJNP 32 566	+Devyanin, Shestakov	(NOVO)
		Translated from YAF 32 1098.		
LOVERRE	80	ZPHY C6 187	+Armenteros, Dionisi+	(CERN, CDEF, MADR, STOJ) IJP
AGUILAR...	78	NP B140 73	+Aguilar-Benitez, Cerrada+	(MADR, BOMB, CERN+)
CASON	78	PR D25 1786	+Saunbaugh, Bishop, Biswas+	(NDAM, ANL)
LEPPER	77	PR D16 2054	+Buttram, Crawley, Duke, Lamb, Peterson	(ISU)
WETZEL	76	NP B115 208	+Freudenreich, Beusch+	(ETH, CERN, LOIC)
BINNIE	73	PRL 31 1534	+Carr, Debenham, Duane, Garbutt+	(LOIC, SHMP)
GRAYVER	73	Tallahassee	+Hyams, Jones, Blum, Dietl, Koch+	(CERN, MPIM)
HYAMS	73	NP B64 134	+Jones, Weillhammer, Blum, Dietl+	(CERN, MPIM)
PROTOPOP...	73	PR D7 1279	+Protopopescu, Alston-Garnjost, Galtieri, Flatte+	(LBL)

OTHER RELATED PAPERS

AU	87	PR D35 1633	+Morgan, Pennington	(DURH, RAL)
AKESSON	86	NP B264 154	+Albrow, Aimehed+	(Axial Field Spec. Collab.)
MENNESSIER	83	ZPHY C16 241		(MONP)
BARBER	82	ZPHY C12 1	+Dainton, Brodbeck, Brookes+	(DARE, LANC, SHEF)
ETKIN	82C	PR D25 2446	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
BIGI	62	CERN Conf. 247	+Brandt, Carrara+	(CERN)
BINGHAM	62	CERN Conf. 240	+Bloch+	(EPOL, CERN)
ERWIN	62	PRL 9 34	+Hoyer, March, Walker, Wangier	(WISC, BNL)
WANG	61	JETP 13 323	+Veksier, Vrana+	(JINR)
		Translated from ZETF 40 464.		

$a_0(980)$
was $\delta(980)$

$$I^G(J^{PC}) = 1^-(0^{++})$$

NOTE ON THE $a_0(980)$

The proper $q\bar{q}$ assignment of the $a_0(980)$ scalar meson remains a problem. The observed mass and width seem to be inconsistent with those expected for a member of an $L = 1$ $q\bar{q}$ nonet. However, since the mass and width are distorted by the proximity of the $K\bar{K}$ threshold, the nature of the $a_0(980)$ can be better investigated using different experimental observations.

TORNQVIST 82 has shown that it is possible to understand the unusual experimental features of the $a_0(980)$ in a unitarized quark model. As with the $f_0(980)$, the $a_0(980)$ can be interpreted as a normal $q\bar{q}$ resonance with a large admixture of $K\bar{K}$, $\eta\pi$, and $\eta'\pi$ continuum states.

Assuming dominance of the decay $\eta'(958) \rightarrow \eta\pi\pi$ by a virtual $a_0(980)\pi$ intermediate state, BRAMON 80 concludes that the experimental value $\Gamma(\eta'(958) \rightarrow \eta\pi\pi) \approx 200$ keV is fully consistent with a $q\bar{q}$ interpretation. The same analysis also finds additional evidence in favor of this interpretation: If the $a_0(980)$ is a $q\bar{q}$ state, one expects that the decay chain $f_1 \rightarrow a_0(980)\pi \rightarrow \eta\pi\pi$ will be more important for the $f_1(1285)$ than for the $f_1(1420)$; the reverse is true if the $a_0(980)$ is a $q\bar{q}q\bar{q}$ state with a strange quark component. In fact, $f_1(1285) \rightarrow a_0(980)\pi \rightarrow \eta\pi\pi$ is observed, while $f_1(1420) \rightarrow a_0(980)\pi \rightarrow \eta\pi\pi$ is (practically) absent.

The main argument in favor of the $q\bar{q}q\bar{q}$ interpretation of the $a_0(980)$ is its degeneracy in mass with the isoscalar $f_0(980)$. A Crystal Ball measurement of $a_0(980) \rightarrow \gamma\gamma$ suppression in the reaction $\gamma\gamma \rightarrow a_0(980) \rightarrow \eta\pi$ (ANTREASYAN 86) has reinforced this four-quark interpretation. ACHASOV 88B points out that none of the calculations performed in the framework of a $q\bar{q}$ scheme can predict such a narrow $a_0(980) \rightarrow \gamma\gamma$ width as that found by the Crystal Ball. He then argues that the $a_0(980)$ is unusual and shows that a four-quark model is able to give the correct order of magnitude of suppression of 2γ production for both the scalar $a_0(980)$ and $f_0(980)$ mesons.

Another interesting non- $q\bar{q}$ interpretation is given by WEINSTEIN 83B, 89: The $q\bar{q}q\bar{q}$ system is investigated using the nonrelativistic quark model; assuming a large hyperfine interaction, the $a_0(980)$ and $f_0(980)$ are both interpreted as $K\bar{K}$ bound states, and then the P -wave $q\bar{q}$ states are all in the 1300-MeV region. With this S -wave $K\bar{K}$ mesonium assignment, many of the peculiar properties of the $a_0(980)$ and $f_0(980)$ —masses, widths, branching fractions, and two-photon widths—make sense.

Meson Full Listings

 $a_0(980)$

If the $a_0(980)$ is not the 3P_0 state, then this state should be observed near 1300 MeV, with partial decay widths close to the flavor symmetry predictions for an ideal nonet (TORNQVIST 90). The candidate $a_0(1320)$ reported by GAMS-4000 has the right mass, but the signal is weak under the dominant $a_2(1320)$, and its width is much smaller than expected.

See also the "Note on the $f_0(1300)$ ".

 $a_0(980)$ MASS

VALUE (MeV) DOCUMENT ID
982.4 ± 1.4 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

 $\eta\pi$ FINAL STATE ONLY

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
 The data in this block is included in the average printed for a previous datablock.

982.7 ± 1.4 OUR AVERAGE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
982 ± 2		¹ AMSLER 92	CBAR		0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
984 ± 4	1040	¹ ARMSTRONG 91B	OMEG ±		300 $pp \rightarrow pp\eta\pi^+\pi^-$
976 ± 6		ATKINSON 84E	OMEG ±		25-55 $\gamma p \rightarrow \eta\pi n$
986 ± 3	500	² EVANGELISTA 81	OMEG ±		12 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-p$
990.0 ± 7.0	145	² GURTU 79	HBC ±		4.2 $K^- p \rightarrow \Lambda\eta 2\pi$
977.0 ± 7.0		GRASSLER 77	HBC -		16 $\pi^+ p \rightarrow \rho\eta 3\pi$
972 ± 10	150	DEFOIX 72	HBC ±		0.7 $\bar{p}p \rightarrow 7\pi$
980 ± 11	47	CONFORTO 78	OSPK -		4.5 $\pi^- p \rightarrow \rho X^-$
978.0 ± 16.0	50	CORDEN 78	OMEG ±		12-15 $\pi^- p \rightarrow n\eta 2\pi$
989.0 ± 4.0	70	WELLS 75	HBC -		3.1-6 $K^- p \rightarrow \Lambda\eta 2\pi$
970.0 ± 15.0	20	BARNES 69C	HBC -		4-5 $K^- p \rightarrow \Lambda\eta 2\pi$
980 ± 10		CAMPBELL 69	DBC ±		2.7 $\pi^+ d$
980.0 ± 10.0	15	MILLER 69B	HBC -		4.5 $K^- N \rightarrow \eta\pi\Lambda$
980.0 ± 10.0	30	AMMAR 68	HBC ±		5.5 $K^- p \rightarrow \Lambda\eta 2\pi$

¹ From a single Breit-Wigner fit.

² From $f_1(1285)$ decay.

 $K\bar{K}$ ONLY

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
 The data in this block is included in the average printed for a previous datablock.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
976 ± 6	316	DEBILLY 80	HBC ±		1.2-2 $\bar{p}p \rightarrow f_1(1285)\omega$
1016 ± 10	100	³ ASTIER 67	HBC ±		0.0 $\bar{p}p$
1003.3 ± 7.0	143	⁴ ROSENFELD 65	RVUE ±		

³ ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.
⁴ Plus systematic errors.

 $a_0(980)$ WIDTH $\eta\pi$ FINAL STATE ONLY

Width determination very model dependent. Peak width is about 60 MeV, but decay width can be much larger.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
50 to 300 OUR ESTIMATE					
54 ± 10		⁵ AMSLER 92	CBAR		0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
95 ± 14	1040	⁵ ARMSTRONG 91B	OMEG ±		300 $pp \rightarrow pp\eta\pi^+\pi^-$
62 ± 15	500	⁶ EVANGELISTA 81	OMEG ±		12 $\pi^- p \rightarrow \eta\pi^+\pi^-\pi^-p$
60.0 ± 20.0	145	⁶ GURTU 79	HBC ±		4.2 $K^- p \rightarrow \Lambda\eta 2\pi$
60 +50 -30	47	CONFORTO 78	OSPK -		4.5 $\pi^- p \rightarrow \rho X^-$
86.0 +60.0 -50.0	50	CORDEN 78	OMEG ±		12-15 $\pi^- p \rightarrow n\eta 2\pi$
44.0 ± 22.0		GRASSLER 77	HBC -		16 $\pi^+ p \rightarrow \rho\eta 3\pi$
80 to 300		⁷ FLATTE 76	RVUE -		4.2 $K^- p \rightarrow \Lambda\eta 2\pi$
16.0 +25.0 -16.0	70	WELLS 75	HBC -		3.1-6 $K^- p \rightarrow \Lambda\eta 2\pi$
30 ± 5	150	DEFOIX 72	HBC ±		0.7 $\bar{p}p \rightarrow 7\pi$
40 ± 15		CAMPBELL 69	DBC ±		2.7 $\pi^+ d$
60.0 ± 30.0	15	MILLER 69B	HBC -		4.5 $K^- N \rightarrow \eta\pi\Lambda$
80.0 ± 30.0	30	AMMAR 68	HBC ±		5.5 $K^- p \rightarrow \Lambda\eta 2\pi$

⁵ From a single Breit-Wigner fit.

⁶ From $f_1(1285)$ decay.

⁷ Using a two-channel resonance parametrization of GAY 76B data.

 $K\bar{K}$ ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
~ 25	100	⁸ ASTIER 67	HBC ±		
57.0 ± 13.0	143	⁹ ROSENFELD 65	RVUE ±		

⁸ ASTIER 67 includes data of BARLOW 67, CONFORTO 67, ARMENTEROS 65.
⁹ Plus systematic errors.

 $a_0(980)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\pi$	dominant
Γ_2 $K\bar{K}$	seen
Γ_3 $\rho\pi$	
Γ_4 $\pi\eta'(958)$	
Γ_5 $\gamma\gamma$	seen
Γ_6 e^+e^-	

 $a_0(980)$ $\Gamma(\eta\pi)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_5/\Gamma$
0.24 +0.08 -0.07 OUR AVERAGE						
0.28 ± 0.04 ± 0.10	44 ± 7		OEST 90	JADE	$e^+e^- \rightarrow e^+e^-\pi^0\eta$	
0.19 ± 0.07 +0.10 -0.07			ANTREASYAN 86	CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\eta$	
$\Gamma(\eta\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_6/\Gamma$
< 1.5	90		VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^0\eta$	

 $a_0(980)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\eta\pi)$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
0.7 ± 0.3	10	CORDEN 78	OMEG		12-15 $\pi^- p \rightarrow n\eta 2\pi$		
0.25 ± 0.08	10	DEFOIX 72	HBC ±		0.7 $\bar{p}p \rightarrow 7\pi$		
					¹⁰ From the decay of $f_1(1285)$.		
$\Gamma(\rho\pi)/\Gamma(\eta\pi)$	VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •							
< 0.25	70		AMMAR 70	HBC ±	4.1, 5.5 $K^- p \rightarrow \Lambda\eta 2\pi$		

 $a_0(980)$ REFERENCES

AMSLER 92	PL B291 347	+Augustin, Baker+	(Crystal Barrel Collab.)
ARMSTRONG 91B	ZPHY C52 389	+Barnes+	(ATHU, BARI, BIRM, CERN, CDEF)
OEST 90	ZHPY C47 343	+Olson+	(JADE Collab.)
VOROBYEV 88	SJNP 46 273	+Golubev, Dolinsky, Druzhinin+	(NOVO)
ANTREASYAN 86	PR D33 1847	+Aschman, Besset, Bienlein+	(Crystal Ball Collab.)
ATKINSON 84E	PL 138B 459	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
EVANGELISTA 81	NP B178 197	+ (BARI, BONN, CERN, DARE, LVP+)	
DEBILLY 80	NP B176 1	+ Briand, Duboc, Levy+	(CURIN, LAUS, NEUC, GLAS)
GURTU 79	NP B151 181	+Gavillet, Blokzijl+	(CERN, ZEEM, NIJM, OXF)
CONFORTO 78	LNC 23 419	+Conforto, Key+	(RHEL, TNTO, CHIC, FNAL+)
CORDEN 78	NP B144 253	+Corbett, Alexander+	(BIRM, RHEL, TELA, LOWC)
GRASSLER 77	NP B121 189	+ (AAACH3, BERL, BONN, CERN, CRAC, HEIDH+)	(CERN)
FLATTE 76	PL 63B 224		
GAY 76B	PL 63B 220	+Chaloupka, Blokzijl, Heinen+	(CERN, AMST, NIJM, JP)
WELLS 75	NP B101 333	+Radajcic, Roscoe, Lyons	(OXF)
DEFOIX 72	NP B44 125	+Nascimento, Bizzarri+	(CDEF, CERN)
AMMAR 70	PR D2 430	+Kropac, Davis+	(KANS, NWES, ANL, WISC)
BARNES 69C	PRL 23 610	+Chung, Eisner, Bassano, Goldberg+	(BNL, SVRA)
CAMPBELL 69	PRL 22 1204	+Lichtman, Loeffler+	(PURD)
MILLER 69B	PL 29B 255	+Kramer, Carmony+	(PURD)
Also 69	PR 188 2011	+Yen, Ammann, Carmony, Eisner+	(PURD)
AMMAR 68	PRL 21 1832	+Davis, Kropac, Derrick, Fields+	(NWES, ANL)
ASTIER 67	PL 25B 294	+Montanet, Baubillier, Duboc+	(CDEF, CERN, IRAD)
Includes data of BARLOW 67, CONFORTO 67, and ARMENTEROS 65.			
BARLOW 67	NC 50A 701	+Liljestol, Montanet+	(CERN, CDEF, IRAD, LVP)
CONFORTO 67	NP B3 469	+Marechal+	(CERN, CDEF, IPNP, LVP)
ARMENTEROS 65	PL 17 344	+Edwards, Jacobsen+	(CERN, CDEF)
ROSENFELD 65	Oxford Conf. 58		(LRL)

OTHER RELATED PAPERS

TORNQVIST 90	NPBPS 21,196		(HELS)
WEINSTEIN 89	UTPT 89 03	+Isgur	(TNTO)
ACHASOV 88B	ZPHY C41 309	+Shestakov	(NOVO)
WEINSTEIN 83B	PR D27 588	+Isgur	(TNTO)
TORNQVIST 82	PRL 49 624		(HELS)
BRAMON 80	PL 93B 65	+Masso	(BARC)
TURKOT 63	Siena Conf. 1 661	+Collins, Fujii, Kemp+	(BNL, PITT)

See key on page 1343

Meson Full Listings
 $\phi(1020)$

$\phi(1020)$

$I^G(J^{PC}) = 0^-(1^{--})$

$\phi(1020)$ MASS

We average mass and width values only when the systematic errors have been evaluated.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1019.413 ± 0.008 OUR AVERAGE				
1019.7 ± 0.3	2012	DAVENPORT	86 MPSF	400 pA → 4KX
1019.411 ± 0.008	642k	¹ DIJKSTRA	86 SPEC	100-200 $\pi^\pm, \bar{p}, \rho, K^\pm$, on Be
1019.7 ± 0.1 ± 0.1	5079	ALBRECHT	85D ARG	10 $e^+e^- \rightarrow K^+K^-X$
1019.3 ± 0.1	1500	ARENTON	82 AEMS	11.8 polar. $pp \rightarrow KK$
1019.67 ± 0.17	25080	² PELLINEN	82 RVUE	
1019.54 ± 0.12	1100	BARKOV	79B EMUL	$e^+e^- \rightarrow K^+K^-$
1019.52 ± 0.13	3681	BUKIN	78C OLYA	$e^+e^- \rightarrow K^+K^-$ hadrons
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1019.8 ± 0.7		ARMSTRONG	86 OMEG	85 $\pi^+/pp \rightarrow \pi^+/p4Kp$
1020.1 ± 0.11	5526	³ ATKINSON	86 OMEG	20-70 γp
1019.7 ± 1.0		BEBEK	86 CLEO	$e^+e^- \rightarrow T(4S)$
1020.9 ± 0.2		³ FRAME	86 OMEG	13 $K^+p \rightarrow \phi K^+p$
1021.0 ± 0.2		³ ARMSTRONG	83B OMEG	18.5 $K^-p \rightarrow K^-K^+A$
1020.0 ± 0.5		³ ARMSTRONG	83B OMEG	18.5 $K^-p \rightarrow K^-K^+A$
1019.7 ± 0.3		³ BARATE	83 GOLI	190 $\pi^-Be \rightarrow 2\mu X$
1019.8 ± 0.2 ± 0.5	766	IVANOV	81 OLYA	1-1.4 $e^+e^- \rightarrow K^+K^-$
1019.4 ± 0.5	337	COOPER	78B HBC	0.7-0.8 $\bar{p}p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$
1020.0 ± 1.0	383	³ BALDI	77 CNTR	10 $\pi^-p \rightarrow \pi^- \phi p$
1018.9 ± 0.6	800	COHEN	77 ASPK	6 $\pi^\pm N \rightarrow K^+K^-N$
1019.7 ± 0.5	454	KALBFLEISCH	76 HBC	2.18 $K^-p \rightarrow \Lambda K \bar{K}$
1019.4 ± 0.8	984	BESCH	74 CNTR	2 $\gamma p \rightarrow \rho K^+K^-$
1020.3 ± 0.4	100	BALLAM	73 HBC	2.8-9.3 γp
1019.4 ± 0.7		BINNIE	73B CNTR	$\pi^-p \rightarrow \phi n$
1019.6 ± 0.5	120	⁴ AGUILAR...	72B HBC	3.9, 4.6 $K^-p \rightarrow \Lambda K^+K^-$
1019.9 ± 0.5	100	⁴ AGUILAR...	72B HBC	3.9, 4.6 $K^-p \rightarrow K^-pK^+K^-$
1020.4 ± 0.5	131	COLLEY	72 HBC	10 $K^+p \rightarrow K^+p\phi$
1019.9 ± 0.3	410	STOTTLE...	71 HBC	2.9 $K^-p \rightarrow \Sigma/\Lambda K \bar{K}$

¹Weighted and scaled average of 12 measurements of DIJKSTRA 86.
²PELLINEN 82 review includes AKERLOF 77, DAUM 81, BALDI 77, AYRES 74, DEGRON 74.
³Systematic errors not evaluated.
⁴Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

$\phi(1020)$ WIDTH

We average mass and width values only when the systematic errors have been evaluated.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
4.43 ± 0.06 OUR FIT				
4.43 ± 0.06 OUR AVERAGE				
4.45 ± 0.06	271k	DIJKSTRA	86 SPEC	100 π^-Be
4.5 ± 0.7	1500	ARENTON	82 AEMS	11.8 polar. $pp \rightarrow KK$
4.2 ± 0.6	766	⁵ IVANOV	81 OLYA	1-1.4 $e^+e^- \rightarrow K^+K^-$
4.3 ± 0.6		⁵ CORDIER	80 WIRE	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
4.58 ± 0.55	1100	⁵ BARKOV	79B EMUL	$e^+e^- \rightarrow K^+K^-$
4.36 ± 0.29	3681	⁵ BUKIN	78C OLYA	$e^+e^- \rightarrow$ hadrons
4.4 ± 0.6	984	⁵ BESCH	74 CNTR	2 $\gamma p \rightarrow \rho K^+K^-$
4.67 ± 0.72	681	⁵ BALAKIN	71 OSPK	$e^+e^- \rightarrow$ hadrons
4.09 ± 0.29		BIZOT	70 OSPK	$e^+e^- \rightarrow$ hadrons

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

4.08 ± 0.14	13714	KURDADZE	84 OLYA	$e^+e^- \rightarrow$ hadrons
3.6 ± 0.8	337	⁵ COOPER	78B HBC	0.7-0.8 $\bar{p}p \rightarrow K_S^0 K_L^0 \pi^+ \pi^-$
4.5 ± 0.50	1300	^{5,6} AKERLOF	77 SPEC	400 pA → K^+K^-X
4.5 ± 0.8	500	^{5,6} AYRES	74 ASPK	3-6 $\pi^-p \rightarrow K^+K^-n, K^-p \rightarrow K^+K^-A/\Sigma^0$
3.81 ± 0.37		COSME	74B OSPK	$e^+e^- \rightarrow K_L^0 K_S^0$
3.8 ± 0.7	454	⁵ BORENSTEIN	72 HBC	2.18 $K^-p \rightarrow K \bar{K} n$

⁵Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
⁶Systematic errors not evaluated.

$\phi(1020)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 K^+K^-$	(49.1 ± 0.9) %	S=1.3
$\Gamma_2 K_L^0 K_S^0$	(34.3 ± 0.7) %	S=1.2
$\Gamma_3 \rho\pi$	(12.9 ± 0.7) %	
$\Gamma_4 \pi^+\pi^-\pi^0$	(2.5 ± 0.9) %	S=1.1
$\Gamma_5 \eta\gamma$	(1.28 ± 0.06) %	S=1.2
$\Gamma_6 \pi^0\gamma$	(1.31 ± 0.13) × 10 ⁻³	
$\Gamma_7 e^+e^-$	(3.09 ± 0.07) × 10 ⁻⁴	
$\Gamma_8 \mu^+\mu^-$	(2.48 ± 0.34) × 10 ⁻⁴	
$\Gamma_9 \eta e^+e^-$	(1.3 $^{+0.8}_{-0.6}$) × 10 ⁻⁴	
$\Gamma_{10} \pi^+\pi^-$	(8 $^{+5}_{-4}$) × 10 ⁻⁵	S=1.5
$\Gamma_{11} \omega\gamma$	< 5 %	CL=84%
$\Gamma_{12} \rho\gamma$	< 2 %	CL=84%
$\Gamma_{13} \pi^+\pi^-\gamma$	< 7 × 10 ⁻³	CL=90%
$\Gamma_{14} f_0(980)\gamma$	< 2 × 10 ⁻³	CL=90%
$\Gamma_{15} \pi^0\pi^0\gamma$	< 1 × 10 ⁻³	CL=90%
$\Gamma_{16} \pi^+\pi^-\pi^+\pi^-$	< 8.7 × 10 ⁻⁴	CL=90%
$\Gamma_{17} \eta'(958)\gamma$	< 4.1 × 10 ⁻⁴	CL=90%
$\Gamma_{18} \pi^+\pi^+\pi^-\pi^-\pi^0$	< 1.5 × 10 ⁻⁴	CL=95%
$\Gamma_{19} \pi^0 e^+e^-$	< 1.2 × 10 ⁻⁴	CL=90%
$\Gamma_{20} \pi^0 \eta\gamma$	< 2.5 × 10 ⁻³	CL=90%
$\Gamma_{21} a_0(980)\gamma$	< 5 × 10 ⁻³	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 9 branching ratios uses 38 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 27.1$ for 33 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-70				
x_3	0	0			
x_4	-39	-10	-75		
x_5	-5	-2	0	0	
Γ	0	0	-24	18	0
	x_1	x_2	x_3	x_4	x_5

Mode	Rate (MeV)	Scale factor
$\Gamma_1 K^+K^-$	2.17 ± 0.05	1.2
$\Gamma_2 K_L^0 K_S^0$	1.52 ± 0.04	1.1
$\Gamma_3 \rho\pi$	0.570 ± 0.030	
$\Gamma_4 \pi^+\pi^-\pi^0$	0.11 ± 0.04	1.1
$\Gamma_5 \eta\gamma$	0.0569 ± 0.0029	1.2

$\phi(1020)$ PARTIAL WIDTHS

$\Gamma(\rho\pi)$				Γ_3
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
0.570 ± 0.030 OUR FIT				
0.57 ± 0.03	JULLIAN	76	OSPK	e^+e^-
$\Gamma(e^+e^-)$				Γ_7
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT	
1.37 ± 0.05 OUR EVALUATION				

Meson Full Listings

 $\phi(1020)$ $\phi(1020)$ BRANCHING RATIOS

$\Gamma(K^+K^-)/\Gamma_{total}$					Γ_1/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.491 ± 0.009 OUR FIT				Error includes scale factor of 1.3.	
0.496 ± 0.019 OUR AVERAGE					
0.44 ± 0.05	321	KALBFLEISCH 76	HBC	2.18 $K^-p \rightarrow \Lambda K^+K^-$	
0.49 ± 0.06	270	DEGROOT 74	HBC	4.2 $K^-p \rightarrow \Lambda\phi$	
0.540 ± 0.034	565	BALAKIN 71	OSPK	$e^+e^- \rightarrow K^+K^-$	
0.486 ± 0.044		CHATELUS 71	OSPK	e^+e^-	
0.48 ± 0.04	252	LINDSEY 66	HBC	2.1-2.7 $K^-p \rightarrow \Lambda K^+K^-$	

$\Gamma(K_L^0 K_S^0)/\Gamma_{total}$					Γ_2/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.343 ± 0.007 OUR FIT				Error includes scale factor of 1.2.	
0.333 ± 0.009 OUR AVERAGE					
0.326 ± 0.035		DOLINSKY 91	ND	$e^+e^- \rightarrow K_L^0 K_S^0$	
0.310 ± 0.024		DRUZHININ 84	ND	$e^+e^- \rightarrow K_L^0 K_S^0$	
0.338 ± 0.010		KURDADZE 84	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.27 ± 0.03	133	KALBFLEISCH 76	HBC	2.18 $K^-p \rightarrow \Lambda K_L^0 K_S^0$	
0.257 ± 0.030	95	BALAKIN 71	OSPK	$e^+e^- \rightarrow K_L^0 K_S^0$	
0.40 ± 0.04	167	LINDSEY 66	HBC	2.1-2.7 $K^-p \rightarrow \Lambda K_L^0 K_S^0$	

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma_{total}$					$(\Gamma_3 + \Gamma_4)/\Gamma$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.153 ± 0.006 OUR FIT				Error includes scale factor of 1.3.	
0.148 ± 0.006 OUR AVERAGE				Error includes scale factor of 1.1.	
0.143 ± 0.007		DOLINSKY 91	ND	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
0.155 ± 0.008		KURDADZE 84	OLYA	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.139 ± 0.007	7	PARROUR 76B	OSPK	e^+e^-	

⁷Using total width 4.1 MeV. The $\rho\pi$ to 3π mode is more than 80% at the 90% confidence level.

$\Gamma(K_L^0 K_S^0)/\Gamma(K\bar{K})$					$\Gamma_2/(\Gamma_1 + \Gamma_2)$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.411 ± 0.008 OUR FIT				Error includes scale factor of 1.2.	
0.45 ± 0.04 OUR AVERAGE					
0.44 ± 0.07		LONDON 66	HBC	2.24 $K^-p \rightarrow \Lambda K\bar{K}$	
0.48 ± 0.07	52	BADIER 65B	HBC	3 K^-p	
0.40 ± 0.10	34	SCHLEIN 63	HBC	1.95 $K^-p \rightarrow \Lambda K\bar{K}$	

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K\bar{K})$					$(\Gamma_3 + \Gamma_4)/(\Gamma_1 + \Gamma_2)$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.184 ± 0.009 OUR FIT				Error includes scale factor of 1.3.	
0.24 ± 0.04 OUR AVERAGE					
0.237 ± 0.039		CERRADA 77B	HBC	4.2 $K^-p \rightarrow \Lambda 3\pi$	
0.30 ± 0.15		LONDON 66	HBC	2.24 $K^-p \rightarrow \Lambda\pi^+\pi^-\pi^0$	

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K_L^0 K_S^0)$					$(\Gamma_3 + \Gamma_4)/\Gamma_2$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.447 ± 0.021 OUR FIT				Error includes scale factor of 1.3.	
0.51 ± 0.05 OUR AVERAGE					
0.56 ± 0.07	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0$	
0.47 ± 0.06	516	COSME 74	OSPK	$e^+\pi^+\pi^-\pi^0 \rightarrow \pi^+\pi^-\pi^0$	

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$					Γ_8/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.48 ± 0.34 OUR AVERAGE					
2.69 ± 0.46	8	HAYES 71	CNTR	8.3, 9.8 $\gamma C \rightarrow \mu^+\mu^- X$	
2.17 ± 0.60	8	EARLES 70	CNTR	6.0 $\gamma C \rightarrow \mu^+\mu^- X$	
2.34 ± 1.01		MOY 69	CNTR	5.0 $\gamma C \rightarrow \mu^+\mu^- X$	

⁸Neglecting interference between resonance and continuum.

$\Gamma(\eta\gamma)/\Gamma_{total}$					Γ_5/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0128 ± 0.0006 OUR FIT				Error includes scale factor of 1.2.	
0.0128 ± 0.0007 OUR AVERAGE				Error includes scale factor of 1.2.	
0.0130 ± 0.0006		9 DRUZHININ 84	ND	$e^+e^- \rightarrow 3\gamma$	
0.014 ± 0.002	10	DRUZHININ 84	ND	$e^+e^- \rightarrow 6\gamma$	
0.0088 ± 0.0020	290	KURDADZE 83C	OLYA	$e^+e^- \rightarrow 3\gamma$	
0.0135 ± 0.0029		ANDREWS 77	CNTR	6.7-10 γCu	
0.015 ± 0.004	54	9 COSME 76	OSPK	e^+e^-	

⁹From 2γ decay mode of η .

¹⁰From $3\pi^0$ decay mode of η .

$\Gamma(\pi^+\pi^-\gamma)/\Gamma_{total}$					Γ_{13}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.007	90	COSME 74	OSPK	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.06	90	KALBFLEISCH 75	HBC	2.18 $K^-p \rightarrow \Lambda\pi^+\pi^-\gamma$	
<0.04	99	LINDSEY 65	HBC	2.1-2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-\text{ neutrals}$	

$\Gamma(\omega\gamma)/\Gamma_{total}$					Γ_{11}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.05	84	LINDSEY 66	HBC	2.1-2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-\text{ neutrals}$	

$\Gamma(\rho\gamma)/\Gamma_{total}$					Γ_{12}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.02	84	LINDSEY 66	HBC	2.1-2.7 $K^-p \rightarrow \Lambda\pi^+\pi^-\text{ neutrals}$	

$\Gamma(e^+e^-)/\Gamma_{total}$					Γ_7/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.09 ± 0.07 OUR AVERAGE					
3.05 ± 0.12	13714	KURDADZE 84	OLYA	$e^+e^- \rightarrow$ hadrons	
3.00 ± 0.21	3681	BUKIN 78C	OLYA	$e^+e^- \rightarrow$ hadrons	
3.10 ± 0.14	11	PARROUR 76	OSPK	e^+e^-	
3.3 ± 0.3		COSME 74	OSPK	$e^+e^- \rightarrow$ hadrons	
2.81 ± 0.25	681	BALAKIN 71	OSPK	$e^+e^- \rightarrow$ hadrons	
3.50 ± 0.27		CHATELUS 71	OSPK	e^+e^-	

¹¹Using total width 4.2 MeV. They detect 3π mode and observe significant interference with ω tail. This is accounted for in the result quoted above.

$\Gamma(\pi^0\gamma)/\Gamma_{total}$					Γ_6/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.31 ± 0.13 OUR AVERAGE					
1.30 ± 0.13		DRUZHININ 84	ND	$e^+e^- \rightarrow 3\gamma$	
1.4 ± 0.5	32	COSME 76	OSPK	e^+e^-	

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$					Γ_{10}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
0.8 $^{+0.5}_{-0.4}$ OUR AVERAGE				Error includes scale factor of 1.5.	
0.63 $^{+0.37}_{-0.28}$	12	GOLUBEV 86	ND	$e^+e^- \rightarrow \pi^+\pi^-$	
1.94 $^{+1.03}_{-0.81}$	12	VASSERMAN 81	OLYA	e^+e^-	
<6.6	95	BUKIN 78B	OLYA	$e^+e^- \rightarrow \pi^+\pi^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<4.0	95	JULLIAN 76	OSPK	e^+e^-	
<2.7	95	ALVENSELEBEN72	CNTR	6.7 $\gamma C \rightarrow C\pi^+\pi^-$	

¹²Using $\Gamma(e^+e^-)/\Gamma_{total} = 3.1 \times 10^{-4}$.

$\Gamma(K_L^0 K_S^0)/\Gamma(K^+K^-)$					Γ_2/Γ_1
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.699 ± 0.024 OUR FIT				Error includes scale factor of 1.2.	
0.740 ± 0.031 OUR AVERAGE					
0.70 ± 0.06	2732	BUKIN 78C	OLYA	$e^+e^- \rightarrow K_L^0 K_S^0$	
0.82 ± 0.08		LOSTY 78	HBC	4.2 $K^-p \rightarrow \phi$ hyperon	
0.71 ± 0.05		LAVEN 77	HBC	10 $K^-p \rightarrow \phi^+ K^- \Lambda$	
0.71 ± 0.08		LYONS 77	HBC	3-4 $K^-p \rightarrow \Lambda\phi$	
0.89 ± 0.10	144	AGUILAR-...	72B	HBC 3.9, 4.6 K^-p	

$[\Gamma(\rho\pi) + \Gamma(\pi^+\pi^-\pi^0)]/\Gamma(K^+K^-)$					$(\Gamma_3 + \Gamma_4)/\Gamma_1$
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.313 ± 0.016 OUR FIT				Error includes scale factor of 1.3.	
0.28 ± 0.09	34	AGUILAR-...	72B	HBC 3.9, 4.6 K^-p	

$\Gamma(\eta e^+e^-)/\Gamma_{total}$					Γ_9/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.3 $^{+0.8}_{-0.6}$	7	GOLUBEV 85	ND	$e^+e^- \rightarrow \gamma\gamma e^+e^-$	

$\Gamma(\eta'(958)\gamma)/\Gamma_{total}$					Γ_{17}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<4.1	90	DRUZHININ 87	ND	$e^+e^- \rightarrow \gamma\eta\pi^+\pi^-$	

$\Gamma(\pi^0\pi^0\gamma)/\Gamma_{total}$					Γ_{15}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
<1	90	DRUZHININ 87	ND	$e^+e^- \rightarrow 5\gamma$	

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma_{total}$					Γ_{18}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
<1.5	95	BARKOV 88	CMD	$e^+e^- \rightarrow \pi^+\pi^+\pi^-\pi^0$	

See key on page 1343

Meson Full Listings
 $\phi(1020), h_1(1170)$

$\Gamma(\pi^+\pi^-\pi^+\pi^-)/\Gamma_{\text{total}}$		Γ_{16}/Γ	
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN COMMENT
<8.7	90	CORDIER 79	WIRE $e^+e^- \rightarrow 4\pi$
$\Gamma(\eta_0(980)\gamma)/\Gamma_{\text{total}}$		Γ_{14}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN COMMENT
<2	90	DRUZHININ 87	ND $e^+e^- \rightarrow \pi^0\pi^0\gamma$
$\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$		Γ_{19}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
< 1.2×10^{-4}	90	DOLINSKY 88	ND $e^+e^- \rightarrow \pi^0e^+e^-$
$\Gamma(\pi^0\eta\gamma)/\Gamma_{\text{total}}$		Γ_{20}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN COMMENT
<2.5	90	DOLINSKY 91	ND $e^+e^- \rightarrow \pi^0\eta\gamma$
$\Gamma(a_0(980)\gamma)/\Gamma_{\text{total}}$		Γ_{21}/Γ	
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN COMMENT
<5	90	DOLINSKY 91	ND $e^+e^- \rightarrow \pi^0\eta\gamma$

 $\phi(1020)$ REFERENCES

DOLINSKY 91	PRPL 202 99	+Druzhinin, Dubrovin+	(NOVO)
BARKOV 88	SJNP 47 248	+Vasserman, Vorobyev, Ivanov+	(NOVO)
	Translated from YAF 47 393.		
DOLINSKY 88	SJNP 48 277	+Druzhinin, Dubrovin, Golubev+	(NOVO)
	Translated from YAF 48 442.		
DRUZHININ 87	ZPHY C37 1	+Dubrovich, Carney+	(NOVO)
ARMSTRONG 86	PL 166B 245	(ATHU, BARI, BIRM, CERN)	
ATKINSON 86	ZPHY C30 521	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
BEBEK 86	PRL 56 1893	+ Berkelman, Blucher, Cassel+	(CLEO Collab.)
DAVENPORT 86	PR 33 2519	(TUFTS, ARIZ, FNAL, FSU, NDAM, VAND)	
DIJKSTRA 86	ZPHY C31 375	+Bailey+	(ANIK, BRIS, CERN, CRAC, MPIM, RAL)
FRAME 86	NP B276 667	+Hughes, Lynch, Minto, McFadzean+	(GLAS)
GOLUBEV 86	SJNP 44 409	+Druzhinin, Ivanchenko, Perevedentsev+	(NOVO)
	Translated from YAF 44 633.		
ALBRECHT 85D	PL 153B 343	+Drescher, Binder, Drews+	(ARGUS Collab.)
GOLUBEV 85	SJNP 41 756	+Druzhinin, Ivanchenko, Peryshkin+	(NOVO)
	Translated from YAF 41 1183.		
DRUZHININ 84	PL 144B 136	+Golubev, Ivanchenko, Peryshkin+	(NOVO)
KURDADZE 84	IVF 84-7 Preprint	+Leltchouk, Pakhtusova, Sidorov+	(NOVO)
ARMSTRONG 83B	NP B224 193	+ (BARI, BIRM, CERN, MILA, CURIN+)	
BARATE 83	PL 121B 449	+Bareyre, Bonamy+	(SACL, LOIC, SHMP, IND)
KURDADZE 83C	JETPL 38 366	+Lechuk, Root+	(NOVO)
	Translated from ZETFP 38 306.		
ARENTON 82	PR D25 2241	+Ayres, Diebold, May, Swallow+	(ANL, ILL)
PELLINEN 82	PS 25 599	+Roos	(HELS)
DALU 81	PL 100B 439	+Bardsey+ (AMST, BRIS, CERN, CRAC, MPIM+)	
IVANOV 81	PL 107B 297	+Kurdadze, Lelchuk, Sidorov, Skrinsky+	(NOVO)
	Also Private Comm.		
VASSERMAN 81	PL 99B 62	+Kurdadze, Sidorov, Skrinsky+	(NOVO)
CORDIER 80	NP B172 13	+Delcourt, Eschstruth, Fulda+	(LALO)
BARKOV 79B	IVF 79-93 Preprint	+Zolotorev, Makarina, Mishakova+	(NOVO)
CORDIER 79	PL 81B 389	+Delcourt, Eschstruth, Fulda+	(LALO)
BUKIN 78B	SJNP 27 521	+Kurdadze, Sidorov, Skrinsky+	(NOVO)
	Translated from YAF 27 985.		
BUKIN 78C	SJNP 27 516	+Kurdadze, Serednyakov, Sidorov+	(NOVO)
	Translated from YAF 27 976.		
COOPER 78B	NP B146 1	+Ganguli+ (TATA, CERN, CDEF, MADR)	
LOSTY 78	NP B133 38	+Holmgren, Blokzijl+ (CERN, AMST, NIJM, OXF)	
AKERLOF 77	PRL 39 561	+Alley, Binstinger, Ditzler+ (FNAL, MICH, PUPD)	
ANDREWS 77	PRL 39 198	+Fukushima, Harvey, Lobkowicz, May+	(NOVO)
BALDI 77	PL 68B 381	+Bohringer, Dorasz, Hungerbuhler+	(GEVA)
CERRADA 77B	NP B126 241	+Blockzijl, Heinen+ (AMST, CERN, NIJM, OXF)	
COHEN 77	PRL 38 269	+Ayres, Diebold, Kramer, Pawlicki, Wicklund (ANL)	
LAVEN 77	NP B127 43	+Otter, Klein+ (AACH3, BERL, CERN, LOIC, WIEN)	
LYONS 77	NP B125 207	+Cooper, Clark (OXF)	
COSME 76	PL 63B 352	+Courau, Dudelzak, Grelaud, Jean-Marie+	(ORSAY)
JULLIAN 76	Thesis 2 519		
KALBFLEISCH 76	PR D13 22	+Strand, Chapman (BNL, MICH)	
PARROUR 76	PL 63B 357	+Grelaud, Cosme, Courau, Dudelzak+	(ORSAY)
PARROUR 76B	PL 63B 362	+Grelaud, Cosme, Courau, Dudelzak+	(ORSAY)
KALBFLEISCH 75	PR D11 987	+Strand, Chapman (BNL, MICH)	
AYRES 74	PRL 32 1463	+Diebold, Greene, Kramer, Levine+ (ANL)	
BESCH 74	NP B70 257	+Hartmann, Kose, Krautschneider, Paul+ (BONN)	
COSME 74	PL 48B 155	+Jean-Marie, Jullian, Laplanche+ (ORSAY)	
COSME 74B	PL 48B 159	+Jean-Marie, Jullian, Laplanche+ (ORSAY)	
DEGROOT 74	NP B74 77	+Hoogland, Jongejans, Metzger+ (AMST, NIJM)	
BALLAM 73	PR D7 3150	+Chadwick, Eisenberg, Bingham+ (SLAC, LBL)	
BINNIE 73B	PR D8 2789	+Carr, Debenham, Duane+ (LOIC, SHMP)	
AGUILAR... 72B	PR D6 29	+Aguilar-Benitez, Chung, Eisner, Samios (BNL)	
ALVENSLEBEN 72	PRL 28 66	+Becker, Biggs, Binkley+ (MIT, DESY)	
BORENSTEIN 72	PR D5 1559	+Danburg, Kalbfleisch+ (BNL, MICH)	
COLLEY 72	NP B50 1	+Jobes, Riddiford, Griffiths+ (BIRM, GLAS)	
BALAKIN 71	PL 34B 328	+Budker, Pakhtusova, Sidorov, Skrinsky+ (NOVO)	
CHATELUS 71	LAL 1247 Thesis		(STRB)
	Also		
HAYES 71	PR D4 899	+Bizot, Buon, Chatelus, Jeanjean+ (ORSAY)	
STOTTLE... 71	ORO 2504 170 Thesis	+Imlay, Joseph, Keizer, Stein (CORN)	
BIZOT 70	PL 32 416	+Stottlemeyer (UMD)	
	Also		
EARLES 70	PRL 25 1312	+Buon, Chatelus, Jeanjean+ (ORSAY)	
MOY 69	Thesis 69	+Perez-y-Jorba (NEAS)	
LINDSEY 66	PR 147 913	+Falsler, Gettner, Lutz, Moy, Tang+ (NEAS)	
LONDON 66	PR 143 1034		(LRL)
BADIER 65B	PL 17 337	+Smith (BNL, SYRA) IGJPC	
LINDSEY 65	PRL 15 221	+Rau, Goldberg, Lichtman+ (BNL, SYRA) IGJPC	
	LINDSEY 65 data included in LINDSEY 66.	+Demoulin, Barlotaud+ (EPOL, SACL, AMST)	
SCHLEIN 63	PRL 10 368	+Smith (LRL)	
		+Slater, Smith, Stork, Ticho (UCLA) IGJP	

OTHER RELATED PAPERS

KAMAL 92	PL B284 421	+Xu (ALBE)
GEORGIO... 85	PL 152B 428	+Georgiopoulos+ (TUFTS, ARIZ, FNAL, FSU, NDAM+)
ARMENTEROS 63B	Siena Conf. 2 70	+Edwards, Astier+ (CERN, CDEF)
GELFAND 63B	PRL 11 438	+Miller, Nussbaum, Kirsch+ (COLU, RUTG)
BERTANZA 62	PRL 9 180	+Brisson, Connolly, Hart+ (BNL, SYRA)

 $h_1(1170)$

$$I^G(J^{PC}) = 0^-(1^{+-})$$

 $h_1(1170)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1170 ± 20 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1168 ± 4	ANDO 92	SPEC		$8\pi^-p \rightarrow \pi^+\pi^-\pi^0n$
1166 ± 5 ± 3	¹ ANDO 92	SPEC		$8\pi^-p \rightarrow \pi^+\pi^-\pi^0n$
1190 ± 60	² DANKOWY... 81	SPEC	0	$8\pi^+\pi^-\pi^0n$ $8\pi p \rightarrow 3\pi n$
				¹ Average and spread of values using 2 variants of the model of BOWLER 75.
				² Uses the model of BOWLER 75.

 $h_1(1170)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
360 ± 40 OUR ESTIMATE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
345 ± 6	ANDO 92	SPEC		$8\pi^-p \rightarrow \pi^+\pi^-\pi^0n$
375 ± 6 ± 34	³ ANDO 92	SPEC		$8\pi^-p \rightarrow \pi^+\pi^-\pi^0n$
320 ± 50	⁴ DANKOWY... 81	SPEC	0	$8\pi^+\pi^-\pi^0n$ $8\pi p \rightarrow 3\pi n$
				³ Average and spread of values using 2 variants of the model of BOWLER 75.
				⁴ Uses the model of BOWLER 75.

 $h_1(1170)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen

 $h_1(1170)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE	ATKINSON 84	OMEG	20-70 $\gamma p \rightarrow \pi^+\pi^-\pi^0p$	
seen	DANKOWY... 81	SPEC	$8\pi p \rightarrow 3\pi n$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	ANDO 92	SPEC	$8\pi^-p \rightarrow \pi^+\pi^-\pi^0n$	

 $h_1(1170)$ REFERENCES

ANDO 92	PL B291 496	+Imai+ (KEK, KYOT, NIRS, SAGA, INUS, AKIT)
ATKINSON 84	NP B231 15	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)
DANKOWY... 81	PRL 46 580	+Dankowych+ (TNTO, BNL, CARL, MCGI, OHIO)
BOWLER 75	NP B97 227	+Game, Aitchison, Dainton (OXFPT, DARE)

Meson Full Listings

$b_1(1235)$


$b_1(1235)$

$$I^G(J^{PC}) = 1^+(1^{+-})$$

$b_1(1235)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
1231 ± 10	OUR ESTIMATE	This is only an educated guess; the error given is larger than the error on the average of the published values. Error includes scale factor of 1.6. See the ideogram below.			
1230.9 ± 3.1	OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.			
1225 ± 5		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
1235 ± 15		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
1236 ± 16		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
1222 ± 6		ATKINSON 84E	OMEG ±		25-55 $\gamma p \rightarrow \omega \pi X$
1237 ± 7		ATKINSON 84E	OMEG 0		25-55 $\gamma p \rightarrow \omega \pi X$
1239 ± 5		EVANGELISTA 81	OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
1251.0 ± 8.0	450	GESSAROLI 77	HBC		11 $\pi^- p \rightarrow \pi^- \omega p$
1245.0 ± 11.0	890	FLATTE 76C	HBC		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
1222 ± 4	1400	CHALOUPKA 74	HBC		3.9 $\pi^- p$
1220 ± 7	600	KARSHON 74B	HBC		4.9 $\pi^+ p$
1243 ± 6	1163	1 OTT 72B	HBC		7.1 $\pi^+ p$
• • •		We do not use the following data for averages, fits, limits, etc. • • •			
1311 ± 10		2 TAKAMATSU 90	SPEC	0	8 $\pi^- p \rightarrow \eta \rho n$
1190 ± 10		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
1213 ± 5		ATKINSON 84C	OMEG 0		20-70 γp
1271 ± 11		COLLICK 84	SPEC		200 $\pi^+ Z \rightarrow Z \pi \omega$

¹ From fit of the mass spectrum.
² Breit-Wigner fitting of PWA of $\eta \pi \pi$ system.



$b_1(1235)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
142 ± 8	OUR AVERAGE	Error includes scale factor of 1.1.			
113 ± 12		WEIDENAUER 93	ASTE		$\bar{p}p \rightarrow 2\pi^+ 2\pi^- \pi^0$
160 ± 30		ALDE 92C	GAM2		38,100 $\pi^- p \rightarrow \omega \pi^0 n$
151 ± 31		FUKUI 91	SPEC		8.95 $\pi^- p \rightarrow \omega \pi^0 n$
170 ± 15		EVANGELISTA 81	OMEG -		12 $\pi^- p \rightarrow \omega \pi p$
170.0 ± 50.0	225	BALTAY 78B	HBC		15 $\pi^+ p \rightarrow p 4\pi$
155.0 ± 32.0	450	GESSAROLI 77	HBC		11 $\pi^- p \rightarrow \pi^- \omega p$
182.0 ± 45.0	890	FLATTE 76C	HBC		4.2 $K^- p \rightarrow \pi^- \omega \Sigma^+$
135 ± 20	1400	CHALOUPKA 74	HBC		3.9 $\pi^- p$
156 ± 22	600	KARSHON 74B	HBC		4.9 $\pi^+ p$
134 ± 23	1163	3 OTT 72B	HBC		7.1 $\pi^+ p$
• • •		We do not use the following data for averages, fits, limits, etc. • • •			
126 ± 10		4 TAKAMATSU 90	SPEC	0	8 $\pi^- p \rightarrow \eta \rho n$
210 ± 19		AUGUSTIN 89	DM2 ±		$e^+ e^- \rightarrow 5\pi$
231 ± 14		ATKINSON 84C	OMEG 0		20-70 γp
232 ± 29		COLLICK 84	SPEC		200 $\pi^+ Z \rightarrow Z \pi \omega$

³ From fit of the mass spectrum.
⁴ Breit-Wigner fitting of PWA of $\eta \pi \pi$ system.

$b_1(1235)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\omega \pi$	dominant	
Γ_2 $\pi^\pm \gamma$	(1.6 ± 0.4) × 10 ⁻³	
Γ_3 $\eta \rho$	seen	
Γ_4 $\pi^+ \pi^+ \pi^- \pi^0$	< 50 %	84%
Γ_5 $(K\bar{K})^\pm \pi^0$	< 8 %	90%
Γ_6 $K_S^0 K_L^0 \pi^\pm$	< 6 %	90%
Γ_7 $K_S^0 K_S^0 \pi^\pm$	< 2 %	90%
Γ_8 $\pi \phi$	< 1.5 %	84%

[D/S amplitude ratio = 0.26 ± 0.04]

$b_1(1235)$ PARTIAL WIDTHS

$\Gamma(\pi^\pm \gamma)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2
VALUE (keV)					
230.0 ± 60.0	COLLICK	84	SPEC	+	200 $\pi^+ Z \rightarrow Z \pi \omega$

$b_1(1235)$ D-wave/S-wave AMPLITUDE RATIO IN DECAY OF $b_1(1235) \rightarrow \omega \pi$

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
0.260 ± 0.035	OUR AVERAGE					
0.235 ± 0.047		ATKINSON	84C	OMEG	20-70 γp	
0.4 +0.1		GESSAROLI	77	HBC	- 11 $\pi^- p \rightarrow \pi^- \omega p$	
-0.1					$\pi^- \omega p$	
0.21 ± 0.08		CHUNG	75B	HBC	+	7.1 $\pi^+ p$
0.3 ± 0.1		CHALOUPKA	74	HBC	-	3.9-7.5 $\pi^- p$
0.35 ± 0.25	600	KARSHON	74B	HBC	+	4.9 $\pi^+ p$

$b_1(1235)$ BRANCHING RATIOS

$\Gamma(\eta \rho)/\Gamma(\omega \pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1	
VALUE					
• • •	We do not use the following data for averages, fits, limits, etc. • • •				
seen	TAKAMATSU 90	SPEC			
< 0.10	ATKINSON 84D	OMEG	20-70 γp		
$\Gamma(\pi^+ \pi^+ \pi^- \pi^0)/\Gamma(\omega \pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_1
VALUE					
< 0.5	ABOLINS 63	HBC	+	3.5 $\pi^+ p$	
$\Gamma((K\bar{K})^\pm \pi^0)/\Gamma(\omega \pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_1
VALUE					
< 0.08	BALTAY 67	HBC	±	0.0 $\bar{p}p$	
$\Gamma(K_S^0 K_L^0 \pi^\pm)/\Gamma(\omega \pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ_1
VALUE					
< 0.06	BALTAY 67	HBC	±	0.0 $\bar{p}p$	
$\Gamma(K_S^0 K_S^0 \pi^\pm)/\Gamma(\omega \pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_7/Γ_1
VALUE					
< 0.02	BALTAY 67	HBC	±	0.0 $\bar{p}p$	
$\Gamma(\pi \phi)/\Gamma(\omega \pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_8/Γ_1
VALUE					
< 0.015	DAHL 67	HBC		1.6-4.2 $\pi^- p$	
• • •	We do not use the following data for averages, fits, limits, etc. • • •				
< 0.04	BIZZARRI 69	HBC	±	0.0 $\bar{p}p$	

$b_1(1235)$ REFERENCES

WEIDENAUER 93	ZPHY C59 387	+Duch+	(ASTERIX Collab.)
ALDE 92C	ZPHY C54 553	+Bellazzini+	(SERP, BELG, LANL, LAPP, PISA, KEK)
FUKUI 91	PL B257 241	+Horikawa+	(SUGI, NAGO, KEK, KYOT, MIYA)
TAKAMATSU 90	Hadron 89 Conf. p 71	+Ando+	(KEK)
AUGUSTIN 89	NP B320 1	+Cosme	(DM2 Collab.)
ATKINSON 84C	NP B243 1	+	(BONN, CERN, GLAS, LANC, MCHS, CURIN+) JP
ATKINSON 84D	NP B242 269	+	(BONN, CERN, GLAS, LANC, MCHS, CURIN+)
ATKINSON 84E	PL 138B 459	+	(BONN, CERN, GLAS, LANC, MCHS, CURIN+)
COLLICK 84	PRL 53 2374	+Heppelmann, Berg+	(MINN, ROCH, FNAL)
EVANGELISTA 81	NP B178 197	+	(BARI, BONN, CERN, DARE, LIVP+)
BALTAY 78B	PR D17 62	+Cautis, Cohen, Csorna+	(COLU, BING)
GESSAROLI 77	NP B126 382	+	(BGNA, FIRZ, GENO, MILA, OXF, PAVI) JP
FLATTE 76C	PL 64B 225	+Gay, Blokzijl, Metzger+	(CERN, AMST, NIJM, OXF) JP
CHUNG 75B	PR D11 2426	+Protopopescu, Lynch, Flatte+	(BNL, LBL, UCSF) JP
CHALOUPKA 74	PL 51B 407	+Ferrando, Losty, Montanet	(CERN) JP
KARSHON 74B	PR D10 3608	+Mikeneberg, Eisenberg, Pitluck, Ronat+	(REHO) JP
OTT 72B	LBL-1547 Thesis		(LBL) JP
BIZZARRI 69	NP B14 169	+Foster, Gavillet, Montanet+	(CERN, CDEF)
BALTAY 67	PRL 18 93	+Franzini, Severiens, Yen, Zanello	(COLU)
DAHL 67	PR 163 1377	+Hardy, Hess, Kirz, Miller	(LRL)
ABOLINS 63	PRL 11 381	+Lander, Mehlopp, Nguyen, Yager	(UCSD)

OTHER RELATED PAPERS

BRAU 88	PR D37 2379	+Frank+	(SLAC Hybrid Facility Photon Collab.) JP
ATKINSON 84C	NP B243 1	+	(BONN, CERN, GLAS, LANC, MCHS, CURIN+) JP
GOLDHABER 65	PRL 15 118	+Goldhaber, Kadyk, Shen	(LRL)
CARMONY 64	PRL 12 254	+Lander, Rindfleisch, Xuong, Yager	(UCB) JP
BONDAR 63B	PL 5 209	+Dodd+	(AACH, BIRM, HAMB, LOIC, MPIM)

$a_1(1260)$

$$J^G(J^{PC}) = 1^-(1^{++})$$

NOTE ON THE $a_1(1260)$

For some time, even the existence of this broad bump as a genuine resonance in the 3π mass spectrum was questioned. Today the $a_1(1260)$ situation is still not satisfactorily understood, and its resonance parameters are not well determined. For an attempt to fit the leptonic data with two resonances, see IIZUKA 89.

The main experimental data on the $a_1(1260)$ can be grouped into two classes:

(1) Hadronic production—There are two high-statistics experiments: Diffractive production with incident π^- (DAUM 80, 81B) and charge-exchange production with low-energy π^- (DANKOWYCH 81), both on hydrogen. The extraction of the $a_1(1260)$ parameters from these experiments is troubled by the presence of coherent background, attributed to the Deck effect. Both experiments perform a partial-wave analysis. The phenomenological amplitude used to explain the $1^+S_0^+$ data consists of a rescattered Deck amplitude (calculated from one-pion exchange and not allowed to vary), plus a direct resonance-production term. Both experiments agree on a mass of about 1270 MeV, but DAUM 81B finds a somewhat smaller width than does DANKOWYCH 81 (≈ 300 MeV versus ≈ 380 MeV). Rather lower values for the $a_1(1260)$ mass and width (1121 ± 8 MeV and 239 ± 11 MeV) have been obtained from a partial-wave analysis based on the isobar model of the $\pi^+\pi^-\pi^0$ system in a high-statistics π^-p charge-exchange reaction (ANDO 92). However, in this analysis, only Breit-Wigner terms were considered.

(2) τ decay—Four experiments have reported good data on $\tau \rightarrow a_1(1260)\nu_\tau$, $\rightarrow \rho\pi\nu_\tau$ (RUCKSTUHL 86, SCHMIDKE 86, ALBRECHT 86B, and BAND 87). Here the $a_1(1260)$ from τ decay is expected to be (almost) free from background. The four sets of data show some inconsistencies in the values quoted for the mass. However, according to BOWLER 86, this can be attributed to the different assumptions and approximations made in fitting the data. Furthermore, all these τ decays seem to indicate a consistent $a_1(1260)$ width greater than 400 MeV, considerably larger than that found by DAUM 81B. A recent analysis by ALBRECHT 93C on new ARGUS τ decay data confirms that the model-dependent systematic uncertainties of the parameters are much larger than the now small statistical errors.

The discrepancies between the hadronic and the τ decay results have stimulated several reanalyses. BOWLER 86, TORNQVIST 87, ISGUR 89, and IVANOV 91 have studied

the process $\tau \rightarrow 3\pi\nu_\tau$ (BOWLER 86 has fit the data of ALBRECHT 86B and SCHMIDKE 86, while TORNQVIST 87, ISGUR 89, and IVANOV 91 have also used RUCKSTUHL 86).

BOWLER 86 assumes that the 3π state is wholly $a_1(1260)$, with no background, coherent or incoherent. His fits to the data always use the same theoretical form with a “normal” Breit-Wigner shape and various behaviors of the $a_1(1260)$ axial coupling as a function of the 3π mass.

TORNQVIST 87 fits a modified Breit-Wigner form to the data that includes, besides $\rho\pi$ and $K^*(892)\bar{K} + \bar{K}^*(892)K$ threshold effects, an energy-dependent real part of the $a_1(1260)$ mass parameter (“running mass shift function”).

ISGUR 89 deduces a full mass-dependent covariant amplitude for $\tau \rightarrow 3\pi\nu_\tau$ from theory; all the ambiguities due to the non-pointlike nature of the hadrons (such as unknown off-shell behaviors of propagators and vertices) are associated with a parameterized nonresonant background amplitude. Since this background is small anyway, the $a_1(1260)$ parameters do not depend critically on its form.

Despite these quite different approaches, all three analyses find a good overall description of all the τ decay data with an $a_1(1260)$ mass near 1230 MeV, consistent with the hadronic data; however, their widths (400 MeV for BOWLER 86, 420 MeV for ISGUR 89, and 600 MeV for TORNQVIST 87) remain significantly higher than that extracted from diffractive-hadronic data.

IVANOV 91, using a phenomenological meson Lagrangian based on a four-quark interaction, obtains $a_1(1260)$ parameters consistent with those mentioned above.

BOWLER 88 returned to the diffractive data and investigated their consistency with an $a_1(1260)$ width greater than 400 MeV, as required by the τ -decay data. He verified that a width of about 300 MeV is a direct consequence of the particular fixed shape DAUM 81B used for the Deck amplitude. Freeing this shape, good fits are obtained with a width of about 400 MeV. He then finds no contradiction between the hadronic and τ -decay data, and the $a_1(1260)$ parameters are well constrained. However, applying the same type of analysis to the ANDO 92 data, the low mass and narrow width they obtained with the Breit-Wigner PWA do not change appreciably.

No evidence for charge-exchange photoproduction of the $a_1(1260)$ is observed by CONDO 93. This lack of evidence (together with a clear signal of $a_2(1320)$ photoproduction) is shown to be consistent with either an extremely large $a_1(1260)$ hadronic width or with a small radiative width $\Gamma[a_1(1260) \rightarrow \pi\gamma]$, which this could be accommodated if the a_1 mass is somewhat below 1260 MeV.

Meson Full Listings

 $a_1(1260)$, $f_2(1270)$ $a_1(1260)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1230 ± 40	OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1211 ± 7	ALBRECHT	93C	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1121 ± 8	1 ANDO	92	SPEC	$8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
1242 ± 37	2 IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1260 ± 14	3 IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1250 ± 9	4 IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
1208 ± 15	ARMSTRONG	90	OMEG 0	$300.0 p p \rightarrow p p \pi^+ \pi^- \pi^0$
1220 ± 15	5 ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1260 ± 25	6 BOWLER	88	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1166 ± 18 ± 11	BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1164 ± 41 ± 23	BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$
1250 ± 40	5 TORNQVIST	87	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1046 ± 11	ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1056 ± 20 ± 15	RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1194 ± 14 ± 10	SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
1240.0 ± 80.0	7 DANKOWY...	81	SPEC 0	$8.45 \pi^- p \rightarrow n 3\pi$
1280.0 ± 30.0	7 DAUM	81B	CNTR	$63.94 \pi^- p \rightarrow p 3\pi$
1041.0 ± 13.0	8 GAVILLET	77	HBC +	$4.2 K^- p \rightarrow \Sigma 3\pi$

1 Average and spread of values using 2 variants of the model of BOWLER 75.

2 Reanalysis of RUCKSTUHL 86.

3 Reanalysis of SCHMIDKE 86.

4 Reanalysis of ALBRECHT 86B.

5 From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

6 From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

7 Uses the model of BOWLER 75.

8 Produced in K^- backward scattering. $a_1(1260)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
~ 400	OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
446 ± 21	ALBRECHT	93C	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
239 ± 11	ANDO	92	SPEC	$8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
266 ± 13 ± 4	9 ANDO	92	SPEC	$8 \pi^- p \rightarrow \pi^+ \pi^- \pi^0 n$
465 +228 -143	10 IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
298 +40 -34	11 IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
488 ± 32	12 IVANOV	91	RVUE	$\tau \rightarrow \pi^+ \pi^+ \pi^- \nu$
430 ± 50	ARMSTRONG	90	OMEG 0	$300.0 p p \rightarrow p p \pi^+ \pi^- \pi^0$
420 ± 40	13 ISGUR	89	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
396 ± 43	14 BOWLER	88	RVUE	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
405 ± 75 ± 25	BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
419 ± 108 ± 57	BAND	87	MAC	$\tau^+ \rightarrow \pi^+ \pi^0 \pi^0 \nu$
521 ± 27	ALBRECHT	86B	ARG	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
476 +132 -120 ± 54	RUCKSTUHL	86	DLCO	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
462 ± 56 ± 30	SCHMIDKE	86	MRK2	$\tau^+ \rightarrow \pi^+ \pi^+ \pi^- \nu$
380.0 ± 100.0	15 DANKOWY...	81	SPEC 0	$8.45 \pi^- p \rightarrow n 3\pi$
300.0 ± 50.0	15 DAUM	81B	CNTR	$63.94 \pi^- p \rightarrow p 3\pi$
230.0 ± 50.0	16 GAVILLET	77	HBC +	$4.2 K^- p \rightarrow \Sigma 3\pi$

9 Average and spread of values using 2 variants of the model of BOWLER 75.

10 Reanalysis of RUCKSTUHL 86.

11 Reanalysis of SCHMIDKE 86.

12 Reanalysis of ALBRECHT 86B.

13 From a combined reanalysis of ALBRECHT 86B, SCHMIDKE 86, and RUCKSTUHL 86.

14 From a combined reanalysis of ALBRECHT 86B and DAUM 81B.

15 Uses the model of BOWLER 75.

16 Produced in K^- backward scattering. $a_1(1260)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \rho\pi$	dominant	
$\Gamma_2 \pi\gamma$	seen	
$\Gamma_3 \pi(\pi\pi)s\text{-wave}$	[a] <0.7 %	90%

[a] This is only an educated guess; the error given is larger than the error on the average of the published values.

 $a_1(1260)$ PARTIAL WIDTHS

$\Gamma(\pi\gamma)$	DOCUMENT ID	TECN	COMMENT
VALUE (keV)			
640.0 ± 246.0	ZIELINSKI	84C	SPEC 200 $\pi^+ Z \rightarrow Z 3\pi$

 $a_1(1260)$ BRANCHING RATIOS

$\Gamma(\pi(\pi\pi)s\text{-wave})/\Gamma(\rho\pi)$	DOCUMENT ID	TECN	
VALUE			
0.003 ± 0.003	17 LONGACRE	82	RVUE

17 Uses multichannel Altchison-Bowler model (BOWLER 75). Uses data from GAVILLET 77, DAUM 80, and DANKOWYCH 81.

 $a_1(1260)$ REFERENCES

ALBRECHT	93C	ZPHY C58 61	+Ehrlichmann, Hamacher+ (ARGUS Collab.)
ANDO	92	PL B291 496	+Imai+ (KEK, KYOT, NIRS, SAGA, INUS, AKIT)
IVANOV	91	ZPHY C49 563	+Osipov, Volkov (JINR)
ARMSTRONG	90	ZPHY C48 213	+Benayoun, Beusch (WAT6 Collab.)
ISGUR	89	PR D39 1357	+Morningstar, Reader (TNT0)
BOWLER	88	PL B209 99	(OXF)
BAND	87	PL B198 297	+Camporesi, Chadwick, Delfino+ (MAC Collab.)
TORNQVIST	87	ZPHY C36 695	(HELS)
ALBRECHT	86B	ZPHY C33 7	+Donker, Gabriel, Edwards+ (ARGUS Collab.)
RUCKSTUHL	86	PRL 56 2132	+Stroynowski, Atwood, Barish+ (DELCO Collab.)
SCHMIDKE	86	PRL 57 527	+Abrams, Matteuzzi, Amidei+ (Mark II Collab.)
ZIELINSKI	84C	PRL 52 1195	+Berg, Chandler, Changir+ (ROCH, MINN, FNAL)
LONGACRE	82	PR D26 83	(BNL)
DANKOWYCH...	81	PRL 46 580	Dankowych+ (TNT0, BNL, CARL, MCGI, OHIO)
DAUM	81B	NP B182 269	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
DAUM	80	PL 89B 281	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
GAVILLET	77	PL 69B 119	+Blockzijl, Engelen+ (AMST, CERN, NIJM, OXF) JP
BOWLER	75	NP B97 227	+Game, Altchison, Dainton (OXFTP, DARE)

OTHER RELATED PAPERS

IZUKA	89	PR D39 3357	+Koibuchi, Masuda (NAGO, IBAR, TSUK)
TORNQVIST	87	ZPHY C36 695	(HELS)
ADERHOLZ	64	PL 10 226	+Brown, Kadyk, Shen+ (AACH3, BERL, BIRM, BONN, DESY, HAMB+)
GOLDWABER	64	PRL 12 336	(LRL, UCB)
LANDER	64	PRL 13 346A	+Abolins, Carmony, Hendricks, Xuong+ (UCSD) JP
BELLINI	63	NC 29 896	+Florini, Herz, Negri, Ratti (MIL)

 $f_2(1270)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See also minireview under non- $q\bar{q}$ candidates. $f_2(1270)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1275 ± 5	OUR ESTIMATE			
1274.9 ± 1.2	OUR AVERAGE			
1269.7 ± 5.2	5730	AUGUSTIN	89	DM2 $e^+e^- \rightarrow 5\pi$
1283 ± 8	400 ± 50	1 ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$
1274 ± 5		1 AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
1283 ± 6		2 LONGACRE	86	MPS $22 \pi^- p \rightarrow n 2K_S^0$
1276 ± 7		COURAU	84	DLCO $e^+e^- \rightarrow e^+e^- \pi^+ \pi^-$
1273.3 ± 2.3		3 CHABAUD	83	ASPK $17 \pi^- p$ polarized
1280 ± 4		4 CASON	82	STRC $8 \pi^+ p \rightarrow p \pi^+ 2\pi^0$
1281 ± 7	11600 ± 1000	GIDAL	81	MRK2 J/ψ decay
1282 ± 5		5 CORDEN	79	OMEG $12\text{-}15 \pi^- p \rightarrow n 2\pi$
1269 ± 4	10k	APEL	75	NICE $40 \pi^- p \rightarrow n 2\pi^0$
1272 ± 4	4600	ENGLER	74	DBC $6 \pi^+ n \rightarrow \pi^+ \pi^- p$
1277 ± 4	5300	FLATTE	71	HBC $7.0 \pi^+ p$
1273 ± 8		1 STUNTEBECK	70	HBC $8 \pi^- p, 5.4 \pi^+ d$
1265 ± 8		BOESEBECK	68	HBC $8 \pi^+ p$

See key on page 1343

Meson Full Listings

$f_2(1270)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1281 ± 6		ADAMO	91	OBLX	$\bar{n}p \rightarrow \pi^+ \pi^+ \pi^-$
1262 ± 11		AGUILAR-...	91	EHS	400pp
1275 ± 10		AKER	91	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
1220 ± 10		BREAKSTONE	90	SFM	$p\rho \rightarrow p\rho\pi^+\pi^-$
1288 ± 12		ABACHI	86B	HRS	$e^+e^- \rightarrow \pi^+\pi^-X$
1284 ± 30	3k	BINON	83	GAM2	$38\pi^-p \rightarrow n2\eta$
1280 ± 20	3k	APEL	82	CNTR	$25\pi^-p \rightarrow n2\pi^0$
1284 ± 10	16000	DEUTSCH...	76	HBC	$16\pi^+p$
1258 ± 10	600	TAKAHASHI	72	HBC	$8\pi^-p \rightarrow n2\pi$
1275 ± 13		ARMENISE	70	HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$
1261 ± 5	1960	¹ ARMENISE	68	DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$
1270 ± 10	360	¹ ARMENISE	68	DBC	$5.1\pi^+n \rightarrow p\pi^0MM$
1268 ± 6		⁶ JOHNSON	68	HBC	$3.7-4.2\pi^-p$
1276 ± 11		RABIN	67	HBC	$8.5\pi^+p$

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.
² From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
³ From an energy-independent partial-wave analysis.
⁴ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^0\pi^0$ scattering data.
⁵ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.
⁶ JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

$f_2(1270)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
185 ± 20	OUR ESTIMATE			
184.6 ± 2.8	OUR FIT			Error includes scale factor of 1.6.

183.9^{+5.4}_{-3.0} **OUR AVERAGE** Error includes scale factor of 1.8. See the ideogram below.


180 ± 24		AGUILAR-...	91	EHS	400pp
169 ± 9	5730	⁷ AUGUSTIN	89	DM2	$e^+e^- \rightarrow 5\pi$
150 ± 30	400 ± 50	⁷ ALDE	87	GAM4	$100\pi^-p \rightarrow 4\pi^0n$
186 ⁺⁹ ₋₂		⁸ LONGACRE	86	MPS	$22\pi^-p \rightarrow n2K^0_S$
179.2 ^{+6.9} _{-6.6}		⁹ CHABAUD	83	ASPK	$17\pi^-p$ polarized
160 ± 11		DENNEY	83	LASS	$10\pi^+N$
196 ± 10	3k	APEL	82	CNTR	$25\pi^-p \rightarrow n2\pi^0$
152 ± 9		¹⁰ CASON	82	STRC	$8\pi^+p \rightarrow p\pi^+2\pi^0$
186 ± 27	11600 ± 1000	GIDAL	81	MRK2	J/ψ decay
216 ± 13		¹¹ CORDEN	79	OMEG	$12-15\pi^-p \rightarrow n2\pi$
190 ± 10	10k	APEL	75	NICE	$40\pi^-p \rightarrow n2\pi^0$
192 ± 16	4600	ENGLER	74	DBC	$6\pi^+n \rightarrow \pi^+\pi^-p$
183 ± 15	5300	FLATTE	71	HBC	$7\pi^+p \rightarrow \Delta^{++}f_2$
196 ± 30		⁷ STUNTEBECK	70	HBC	$8\pi^-p, 5.4\pi^+d$
216 ± 20	1960	⁷ ARMENISE	68	DBC	$5.1\pi^+n \rightarrow p\pi^+MM^-$
128 ± 27		⁷ BOESEBECK	68	HBC	$8\pi^+p$
176 ± 21	7,12	JOHNSON	68	HBC	$3.7-4.2\pi^-p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

206 ± 19		ADAMO	91	OBLX	$\bar{n}p \rightarrow \pi^+\pi^+\pi^-$
200 ± 10		AKER	91	CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0$
240 ± 40	3k	BINON	83	GAM2	$38\pi^-p \rightarrow n2\eta$
187 ± 30	650	⁷ ANTIPOV	77	CIBS	$25\pi^-p \rightarrow p3\pi$
225 ± 38	16000	DEUTSCH...	76	HBC	$16\pi^+p$
166 ± 28	600	⁷ TAKAHASHI	72	HBC	$8\pi^-p \rightarrow n2\pi$
173 ± 53		⁷ ARMENISE	70	HBC	$9\pi^+n \rightarrow p\pi^+\pi^-$
155 ± 17		RABIN	67	HBC	$8.5\pi^+p$

⁷ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
⁸ From a partial-wave analysis of data using a K-matrix formalism with 5 poles.
⁹ From an energy-independent partial-wave analysis.
¹⁰ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^0\pi^0$ scattering data.
¹¹ From an amplitude analysis of $\pi^+\pi^- \rightarrow \pi^+\pi^-$ scattering data.
¹² JOHNSON 68 includes BONDAR 63, LEE 64, DERADO 65, EISNER 67.

WEIGHTED AVERAGE
 183.9+5.4-3.0 (Error scaled by 1.8)



$f_2(1270)$ width (MeV)

$f_2(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/Confidence level
Γ_1 $\pi\pi$	(84.9 ^{+2.5} _{-1.3}) %	S=1.3
Γ_2 $\pi^+\pi^-2\pi^0$	(6.9 ^{+1.5} _{-2.7}) %	S=1.4
Γ_3 $K\bar{K}$	(4.6 ± 0.5) %	S=2.8
Γ_4 $2\pi^+2\pi^-$	(2.8 ± 0.4) %	S=1.2
Γ_5 $\eta\eta$	(4.5 ± 1.0) × 10 ⁻³	S=2.4
Γ_6 $4\pi^0$	(3.0 ± 1.0) × 10 ⁻³	
Γ_7 $\gamma\gamma$	(1.32 ^{+0.18} _{-0.16}) × 10 ⁻⁵	S=1.1
Γ_8 $\eta\pi\pi$	< 8 × 10 ⁻³	CL=95%
Γ_9 $K^0K^-\pi^+ + c.c.$	< 3.4 × 10 ⁻³	CL=95%
Γ_{10} e^+e^-	< 9 × 10 ⁻⁹	CL=90%

Meson Full Listings

 $f_2(1270)$

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 6 branching ratios uses 37 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 67.9$ for 30 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-92						
x_3	12	-38					
x_4	11	-36	1				
x_5	2	-9	0	0			
x_6	0	-6	0	0	0		
x_7	8	-3	-15	1	0	0	
Γ	-82	76	-12	-9	-3	0	-10
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $\pi\pi$	156.8 ± 3.2 -1.3	
Γ_2 $\pi^+\pi^-2\pi^0$	12.7 ± 2.9 -5.1	1.4
Γ_3 $K\bar{K}$	8.6 ± 0.8	2.9
Γ_4 $2\pi^+2\pi^-$	5.2 ± 0.7	1.2
Γ_5 $\eta\eta$	0.83 ± 0.19	2.4
Γ_6 $4\pi^0$	0.55 ± 0.18	
Γ_7 $\gamma\gamma$	0.00244 ± 0.00032 -0.00029	

 $f_2(1270)$ PARTIAL WIDTHS

$\Gamma(\pi\pi)$	Γ_1		
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
156.8 ± 3.2 -1.3	OUR FIT		
157.0 ± 5.0 -1.0	13 LONGACRE	86 MPS	$22 \pi^- \rho \rightarrow n2K_S^0$

$\Gamma(K\bar{K})$	Γ_3		
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
8.6 ± 0.8	OUR FIT		Error includes scale factor of 2.9.
9.0 ± 0.7 -0.3	13 LONGACRE	86 MPS	$22 \pi^- \rho \rightarrow n2K_S^0$

$\Gamma(\eta\eta)$	Γ_5		
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.19	OUR FIT		Error includes scale factor of 2.4.
1.0 ± 0.1	13 LONGACRE	86 MPS	$22 \pi^- \rho \rightarrow n2K_S^0$

$\Gamma(\gamma\gamma)$	Γ_7			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.8 ± 0.4				OUR ESTIMATE
2.44 ± 0.32 -0.29				OUR FIT
2.58 ± 0.13 -0.27		14 BEHREND	92 CELL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •				
3.10 ± 0.35 ± 0.35		15 BLINOV	92 MD1	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.27 ± 0.47 ± 0.11		ADACHI	90D TOPZ	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
3.15 ± 0.04 ± 0.39		BOYER	90 MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
3.19 ± 0.16 ± 0.29 -0.28		MARSISKE	90 CBAL	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
2.35 ± 0.65		MORGAN	90 RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$
3.19 ± 0.09 ± 0.22 -0.38	2177	OEST	90 JADE	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0$
3.2 ± 0.1 ± 0.4		17 AIHARA	86B TPC	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.5 ± 0.1 ± 0.5		BEHREND	84B CELL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.85 ± 0.25 ± 0.5		18 BERGER	84 PLUT	$e^+e^- \rightarrow e^+e^-2\pi$

2.70 ± 0.05 ± 0.20		COURAU	84 DLCO	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.52 ± 0.13 ± 0.38	19	SMITH	84C MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.3 ± 0.2 ± 0.5		FRAZER	83 JADE	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.7 ± 0.2 ± 0.6		EDWARDS	82F CBAL	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-2\pi^0$
2.9 ± 0.6 -0.4	20	EDWARDS	82F CBAL	$e^+e^- \rightarrow e^+e^-2\pi^0$
3.2 ± 0.2 ± 0.6		BRANDELIK	81B TASS	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
3.6 ± 0.3 ± 0.5		ROUSSARIE	81 MRK2	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$
2.3 ± 0.8	21	BERGER	80B PLUT	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-$

 $\Gamma(e^+e^-)$ Γ_{10}

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.7	90	VOROBYEV	88 ND	$e^+e^- \rightarrow \pi^0\pi^0$
13 From a partial-wave analysis of data using a K-matrix formalism with 5 poles.				
14 Using a unitarized model with scalars.				
15 Using the unitarized model of LYTH 85.				
16 Error includes spread of different solutions. Data of MARK2 and CRYSTAL BALL used in the analysis. Authors report strong correlations with $\gamma\gamma$ width of $f_0(1300)$: $\Gamma(f_2) + 1/4 \Gamma(f_0) = 3.6 \pm 0.3$ KeV.				
17 Radiative corrections modify the partial widths; for instance the COURAU 84 value becomes 2.66 ± 0.21 in the calculation of LANDRO 86.				
18 Using the MENNESSIER 83 model.				
19 Superseded by BOYER 90.				
20 If helicity = 2 assumption is not made.				
21 Using mass, width and $B(f_2(1270) \rightarrow 2\pi)$ from PDG 78.				

 $f_2(1270) \Gamma(l)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	$\Gamma_3\Gamma_7/\Gamma$		
VALUE (keV)	DOCUMENT ID	TECN	COMMENT

0.114 ± 0.016 -0.015	OUR FIT		Error includes scale factor of 1.1.
0.091 ± 0.007 ± 0.027	22 ALBRECHT	90G ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.104 ± 0.007 ± 0.072	23 ALBRECHT	90G ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
22 Using an incoherent background.			
23 Using a coherent background.			

 $f_2(1270)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT

0.849 ± 0.025 -0.013		OUR FIT		Error includes scale factor of 1.3.
0.837 ± 0.020		OUR AVERAGE		
0.849 ± 0.025		CHABAUD	83 ASPK	$17 \pi^- \rho$ polarized
0.85 ± 0.05	250	BEAUPRE	71 HBC	$8 \pi^+ \rho \rightarrow \Delta^{++} f_2$
0.8 ± 0.04	600	OH	70 HBC	$1.26 \pi^- \rho \rightarrow \pi^+\pi^- n$

 $\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(\pi\pi)$ Γ_2/Γ_1

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.081 ± 0.019 -0.034		OUR FIT		Error includes scale factor of 1.4.
0.15 ± 0.06	600	EISENBERG	74 HBC	$4.9 \pi^+ \rho \rightarrow \Delta^{++} f_2$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.07		EMMS	75D DBC	$4 \pi^+ n \rightarrow \rho f_2$

 $\Gamma(K\bar{K})/\Gamma(\pi\pi)$ Γ_3/Γ_1

We average only experiments which either take into account $f_2(1270)$ - $a_2(1320)$ interference explicitly or demonstrate that $a_2(1320)$ production is negligible.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-------	------	-------------	------	---------

0.055 ± 0.006		OUR FIT		Error includes scale factor of 2.8.
0.040 ± 0.005 -0.006		OUR AVERAGE		
0.037 ± 0.008 -0.021		ETKIN	82B MPS	$23 \pi^- \rho \rightarrow n2K_S^0$
0.045 ± 0.009		CHABAUD	81 ASPK	$17 \pi^- \rho$ polarized
0.039 ± 0.008		LOVERRE	80 HBC	$4 \pi^- \rho \rightarrow K\bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.036 ± 0.005		24 COSTA...	80 OMEG	$1-2.2 \pi^- \rho \rightarrow K^+K^- n$
0.030 ± 0.005		25 MARTIN	79 RVUE	
0.027 ± 0.009		26 POLYCHRO...	79 STRC	$7 \pi^- \rho \rightarrow n2K_S^0$
0.025 ± 0.015		EMMS	75D DBC	$4 \pi^+ n \rightarrow \rho f_2$
0.031 ± 0.012	20	ADERHOLZ	69 HBC	$8 \pi^+ \rho \rightarrow K^+K^- \pi^+ \rho$

24 Re-evaluated by CHABAUD 83.

25 includes PAWLICKI 77 data.

26 Takes into account the $f_2(1270)$ - $f_2'(1525)$ interference.

See key on page 1343

Meson Full Listings

$f_2(1270)$, $f_1(1285)$

$\Gamma(2\pi^+2\pi^-)/\Gamma(\pi\pi)$		Γ_4/Γ_1	
VALUE	EVTS	DOCUMENT ID	TECN COMMENT
0.033 ± 0.005 OUR FIT	Error includes scale factor of 1.2.		
0.033 ± 0.004 OUR AVERAGE	Error includes scale factor of 1.1.		
0.024 ± 0.006	160	EMMS 75D DBC	$4\pi^+n \rightarrow \rho f_2$
0.051 ± 0.025	70	EISENBERG 74 HBC	$4.9\pi^+\rho \rightarrow \Delta^{++}f_2$
0.043 ± 0.007 -0.011	285	LOUIE 74 HBC	$3.9\pi^-\rho \rightarrow n f_2$
0.037 ± 0.007	154	ANDERSON 73 DBC	$6\pi^+n \rightarrow \rho f_2$
0.047 ± 0.013		OH 70 HBC	$1.26\pi^-\rho \rightarrow \pi^+\pi^-n$

$\Gamma(\eta\eta)/\Gamma_{total}$		Γ_5/Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN COMMENT
4.5 ± 1.0 OUR FIT	Error includes scale factor of 2.4.		
3.1 ± 0.8 OUR AVERAGE	Error includes scale factor of 1.3.		
2.8 ± 0.7		ALDE 86D GAM4	$100\pi^-\rho \rightarrow 2\eta n$
5.2 ± 1.7		BINON 83 GAM2	$38\pi^-\rho \rightarrow 2\eta n$

$\Gamma(\eta\eta)/\Gamma(\pi\pi)$		Γ_5/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.05	95	EDWARDS 82F CBAL	$e^+e^- \rightarrow e^+e^-2\eta$
<0.016	95	EMMS 75D DBC	$4\pi^+n \rightarrow \rho f_2$
<0.09	95	EISENBERG 74 HBC	$4.9\pi^+\rho \rightarrow \Delta^{++}f_2$

$\Gamma(4\pi^0)/\Gamma_{total}$		Γ_6/Γ	
VALUE	EVTS	DOCUMENT ID	TECN COMMENT
0.0030 ± 0.0010 OUR FIT			
0.003 ± 0.001	400 ± 50	ALDE 87 GAM4	$100\pi^-\rho \rightarrow 4\pi^0 n$

$\Gamma(\eta\pi\pi)/\Gamma(\pi\pi)$		Γ_8/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.010	95	EMMS 75D DBC	$4\pi^+n \rightarrow \rho f_2$

$\Gamma(K^0K^-\pi^+ + c.c.)/\Gamma(\pi\pi)$		Γ_9/Γ_1	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.004	95	EMMS 75D DBC	$4\pi^+n \rightarrow \rho f_2$

$f_2(1270)$ REFERENCES

BEHREND 92	ZPHY C56 381		(CELLO Collab.)
BLINOV 92	ZPHY C53 33	+Bondar, Bukin+	(NOVO)
ADAMO 91	ZPHY C96 Conf.	+Agello, Balestra+	(OBELIX Collab.)
AGUILAR... 91	ZPHY C50 405	+Aguilar-Benitez, Allison, Batalor+	(LEBC-EHS Collab.)
AKER 91	PL B260 249	+Amsler, Peters+	(Crystal Barrel Collab.)
ADACHI 90D	PL B234 185	+Doser+	(TOPAZ Collab.)
ALBRECHT 90G	ZPHY C48 183	+Ehrlichmann, Harder+	(ARGUS Collab.)
BOYER 90	PR D42 1350	+Butler+	(Mark II Collab.)
BREAKSTONE 90	ZPHY C48 569	+ (ISU, BGNA, CERN, DORT, HEIDI, WARS)	
MARSISKE 90	PR D41 3324	+Antressyan+	(Crystal Ball Collab.)
MORGAN 90	ZPHY C43 623	+Pennington	(RAL (UCSD))
OEST 90	ZPHY C47 343	+Olsson+	(JADE Collab.)
AUGUSTIN 89	NP B320 1	+Cosme	(DM2 Collab.)
VOROBYEV 88	SJNP 48 273	+Golubev, Dolinsky, Druzhinin+	(NOVO)
ALDE 87	PL B198 286	+Binon, Bricman+	(LANL, BRUX, SERP, LAPP)
AUGUSTIN 87	ZPHY C36 369	+Cosme+	(LALO, CLER, FRAS, PADO)
ABACHI 86B	PRL 57 1990	+Derrick, Blockus+	(PURD, ANL, IND, MICH, LBL)
AIHARA 86B	PRL 57 404	+Alston-Garnjost+	(TPC-2 Collab.)
ALDE 86D	NP B269 485	+Binon, Bricman+	(BELG, LAPP, SERP, CERN)
LANDRO 86	PL B172 445	+Mork, Olsen	(UTRO)
LONGACRE 86	PL B177 223	+Etkin+	(BNL, BRAN, CUNY, DUKE, NDAM)
LYTH 85	JPG 11 459		
BEHREND 84B	ZPHY C23 223	+Fenner, Schachter, Schroeder+	(CELLO Collab.)
BERGER 84	ZPHY C26 199	+Kloving, Burger+	(PLUTO Collab.)
COURAU 84	PL 147B 227	+Johnson, Sherman, Atwood, Baillon+	(CIT, SLAC)
SMITH 84C	PR D30 851	+Burke, Abrams, Blocker, Levi+	(SLAC, LBL, HARV)
BINON 83	NC 78A 313	+Donskov, Duteil+	(BELG, LAPP, SERP, CERN)
Also 83B	SJNP 38 561	+Binon, Gouanere+	(BELG, LAPP, SERP, CERN)
CHABAUD 83	NP B223 1	+Gorlich, Cerrada+	(CERN, CRAC, MPIM)
DENNEY 83	PR D28 2726	+Cranley, Firestone, Chapman+	(IOWA, MICH)
FRAZER 83	Aches Conf.		
MENNESSIER 83	ZPHY C16 241		(MONP)
APEL 82	NP B201 197	+Augenstein+(KARLK, KARLE, PISA, SERP, WIEN, CERN)	
CASON 82	PRL 48 1316	+Biswas, Baumbaugh, Bishop+	(NDAM, ANL)
EDWARDS 82F	PL 110B 82	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
ETKIN 82B	PR D25 1786	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
BRANDELIK 81B	APP B12 117	+Boerner+	(TASSO Collab.)
CHABAUD 81	APP B12 575	+Nietorius, Becker+	(CERN, CRAC, MPIM)
GIDAL 81	PL 107B 153	+Goldhaber, Guy, Millikan, Abrams+	(SLAC, LBL)
ROUSSARIE 81	PL 105B 304	+Burke, Abrams, Alam+	(SLAC, LBL)
BERGER 80B	PL 94B 254	+Genzer+	(PLUTO Collab.)
COSTA... 80	NP B175 402	+Costa De Beauregard+	(BARI, BONN, CERN+)
LOVERRE 80	ZPHY C6 187	+Armenteros, Dionisi+	(CERN, CDEF, MADR, STOII)
CORDEN 79	NP B157 250	+Dowell, Garvey+	(BIRM, RHEL, TELA, LOWC)
MARTIN 79	NP B158 520	+Ozenmullu	(DURH)
POLYCHRO... 79	PR D19 1317	+Polychronakos, Cason, Bishop+	(NDAM, ANL)
PDG 78	PL 75B		
ANTIPOV 77	NP B119 45	+Bricman+	(SERP, GEVA)
PAWLICKI 77	PR D15 3196	+Ayres, Cohen, Diebold, Kienzle+	(NDAM, ANL)
DEUTSCH... 76	NP B103 426	+Deutschmann+	(AACH3, BERL, BONN, CERN+)
APEL 75	PL 57B 398	+Augenstein+(KARLK, KARLE, PISA, SERP, WIEN, CERN)	
EMMS 75D	NP B96 155	+Kinson, Stacey, Voltruba+	(BIRM, DURH, RHEL)
EISENBERG 74	PL 52B 239	+Engler, Haber, Karshon+	(REHO)

ENGLER 74	PR D10 2070	+Kraemer, Toaff, Weisser, Diaz+	(CMU, CASE)
LOUIE 74	PL 48B 385	+Alitti, Gandois, Chaloupka+	(SACL, CERN)
ANDERSON 73	PRL 31 562	+Engler, Kraemer, Toaff, Diaz+	(CMU, CASE)
TAKAHASHI 72	PR D6 1266	+Barish+	(TOHOK, PENN, NDAM, ANL)
BEAUPRE 71	NP B28 77	+Deutschmann, Graessler+	(AACH, BERL, CERN)
FLATTE 71	PL 34B 551	+Alston-Garnjost, Barbo-Galtieri+	(LBL)
ARMENISE 70	UNC 4 199	+Ghidini, Foring, Cartacci+	(BARI, BGNA, FIRZ)
OH 70	PR D1 2494	+Garfinkel, Morse, Walker, Prentice	(WISC, TNTO) JP
STUNTEBECK 70	PL 32B 391	+Kenney, Deery, Biswas, Cason+	(NDAM)
ADERHOLZ 69	NP B11 259	+Bartsch+	(AACH3, BERL, CERN, JAGL, WARS)
ARMENISE 68	NC 54A 999	+Ghidini, Forino+	(BARI, BGNA, FIRZ, ORSAY)
ASCOLI 68D	PRL 21 1712	+Crawley, Mortara+	(ILL)
BOESEBECK 68	NP B4 501	+Deutschmann+	(AACH, BERL, CERN)
JOHNSON 68	PR 176 1651	+Poirier, Biswas, Gutay+	(NDAM, PURD, SLAC)
EISNER 67	PR 164 1699	+Johnson, Klein, Peters, Sahni, Yen+	(PURD)
RABIN 67	Thesis		(RUTG)
DERADO 65	PRL 14 872	+Kenney, Poirier, Shephard	(NDAM)
LEE 64	PRL 12 342	+Roe, Sinclair, VanderVelde	(MICH)
BONDAR 63	PL 5 153	+ (AACH, BIRM, BONN, DESY, LOIC, MPIM)	

$f_1(1285)$

$$I^G(J^{PC}) = 0^+(1^+ +)$$

See also minireview under non- $q\bar{q}$ candidates.

$f_1(1285)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1282 ± 5 OUR ESTIMATE					
1282.1 ± 0.6 OUR AVERAGE	Error includes scale factor of 1.6. See the ideogram below.				
1282 ± 4		ARMSTRONG 93C SPEC			$\bar{p}\rho \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
$1270 \pm 6 \pm 10$		ARMSTRONG 92C OMEG			$300 p\rho \rightarrow \rho\rho\pi^+\pi^-\gamma$
1279 ± 5		FUKUI 91C SPEC			$8.95\pi^-\rho \rightarrow \eta\pi^+\pi^-n$
1278 ± 2	140 ± 12	ARMSTRONG 89 OMEG			$300 p\rho \rightarrow K\bar{K}\pi\rho\rho$
1281 ± 1		ARMSTRONG 89E OMEG			$300 p\rho \rightarrow \rho\rho 2(\pi^+\pi^-)$
1278 ± 2		ARMSTRONG 89G OMEG			$85\pi^+\rho \rightarrow 4\pi\pi\rho, \rho\rho \rightarrow 4\pi\rho\rho$
1280.1 ± 2.1	60 ± 20	RATH 89 MPS			$21.4\pi^-\rho \rightarrow K_S^0 K_S^0 \pi^0 n$
1285 ± 1	4750 ± 100	¹ BIRMAN 88 MPS			$8\pi^-\rho \rightarrow K^+\bar{K}^0\pi^-n$
1280 ± 1	504 ± 84	BITYUKOV 88 SPEC			$32.5\pi^-\rho \rightarrow K^+K^-\pi^0 n$
$1279 \pm 6 \pm 10$	16 ± 6	BECKER 87 MRK3			$e^+e^- \rightarrow \phi K\bar{K}\pi$
1286 ± 9		GIDAL 87 MRK2			$e^+e^- \rightarrow e^+\pi^-\eta\pi^+\pi^-$
1280 ± 4		ANDO 86 SPEC			$8\pi^-\rho \rightarrow n\eta\pi^+\pi^-$
1277.0 ± 2.0	420	REEVES 86 SPEC			$6.6 p\bar{p} \rightarrow K\bar{K}\pi X$
1285.0 ± 2.0		CHUNG 85 SPEC			$8\pi^-\rho \rightarrow N\bar{K}\bar{K}\pi$
1279.0 ± 2.0	604	ARMSTRONG 84 OMEG			$85\pi^+\rho \rightarrow K\bar{K}\pi\pi\rho, \rho\rho \rightarrow K\bar{K}\pi\rho\rho$
1287.0 ± 5.0	353	BITUKOV 84 SPEC			$32\pi^-\rho \rightarrow K^+K^-\pi^0 n$
1286.0 ± 1.0		CHAUVAT 84 SPEC			ISR 31.5 $p\rho \rightarrow 12\pi^-\rho \rightarrow n\eta\pi^+\pi^-p$
1278 ± 4		EVANGELISTA 81 OMEG			$4\pi^-\rho \rightarrow K\bar{K}\pi X$
1275.0 ± 6.0	31	BROMBERG 80 SPEC			$4.2 K^-\rho \rightarrow n\eta 2\pi$
1283.0 ± 3.0	103	DIONISI 80 HBC			$12-15\pi^-\rho \rightarrow n5\pi$
1288.0 ± 9.0	200	GURTU 79 HBC			$0.7, 0.76 \bar{p}\rho \rightarrow K\bar{K}3\pi$
1295.0 ± 12.0	85	CORDEN 78 OMEG			$16\pi^+\rho \rightarrow 0.7 \bar{p}\rho \rightarrow 7\pi$
1282.0 ± 2.0	320	NACASCH 78 HBC			$1.2 \bar{p}\rho \rightarrow 2K4\pi$
1279.0 ± 5.0	210	GRASSLER 77 HBC			$8\pi^+\rho \rightarrow p6\pi$
1292 ± 10	150	DEFOIX 72 HBC			$16.0 \pi^+\rho \rightarrow p5\pi$
1286 ± 3	180	DUBOC 72 HBC			$2.7\pi^+\rho$
1303.0 ± 8.0		BARDADIN... 71 HBC			$0.7 \bar{p}\rho, 4,5\text{-body}$
1283.0 ± 6.0		BOESEBECK 71 HBC			$1.2 \bar{p}\rho, 5-6\text{ body}$
1270.0 ± 10.0		CAMPBELL 69 DBC			$2.7\pi^+\rho$
1285 ± 7		LORSTAD 69 HBC			
1290 ± 7		D'ANDLAU 68 HBC			
1283.0 ± 5.0		DAHL 67 HBC			


Meson Full Listings

$f_1(1285)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1264 ± 8	AUGUSTIN	90	DM2	$J/\psi \rightarrow$
				$\gamma \eta \pi^+ \pi^-$
1284 ± 4	TAKAMATSU	90	SPEC 0	$8 \pi^- \rho \rightarrow$
				$K \bar{K} \pi n$
~ 1279	2 TORNQVIST	82B	RVUE	
~ 1275.0	46 3 STANTON	79	CNTR	$8.5 \pi^- \rho \rightarrow$
				$n 2 \gamma 2 \pi$
1271.0 ± 10.0	34 CORDEN	78	OMEG	$12-15 \pi^- \rho \rightarrow$
				$K^+ K^- \pi n$
1280 ± 3	500 4 THUN	72	MMS	$13.4 \pi^- \rho$

1 From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.
 2 From a unitarized quark-model calculation.
 3 From phase shift analysis of $\eta \pi^+ \pi^-$ system.
 4 Seen in the missing mass spectrum.



$f_1(1285)$ mass (MeV)

$f_1(1285)$ WIDTH

Only experiments giving width error less than 20 MeV are kept for averaging.

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
24 ± 3	OUR ESTIMATE					
24.3 ± 1.1	OUR AVERAGE					
25 ± 4	140 ± 12		ARMSTRONG 89	OMEG		300 $pp \rightarrow$ $K \bar{K} \pi pp$
31 ± 5			ARMSTRONG 89E	OMEG		300 $pp \rightarrow$ $pp 2(\pi^+ \pi^-)$
41 ± 12			ARMSTRONG 89G	OMEG		85 $\pi^+ \rho \rightarrow$ $4 \pi \pi \rho, pp \rightarrow$ $4 \pi pp$
17.9 ± 10.9	60 ± 20		RATH 89	MPS		21.4 $\pi^- \rho \rightarrow$ $K_S^0 K_S^0 \pi^0 n$
22 ± 2	4750 ± 100	5	BIRMAN 88	MPS		8 $\pi^- \rho \rightarrow$ $K^+ \bar{K}^0 \pi^- n$
25 ± 4	504 ± 84		BITYUKOV 88	SPEC		32.5 $\pi^- \rho \rightarrow$ $K^+ K^- \pi^0 n$
14 +20 -14 ± 10	16 ± 6		BECKER 87	MRK3		$e^+ e^- \rightarrow$ $\phi K \bar{K} \pi$
19 ± 5			ANDO 86	SPEC		8 $\pi^- \rho \rightarrow$ $n \eta \pi^+ \pi^-$
32.0 ± 8.0	420		REEVES 86	SPEC		6.6 $\rho \bar{p} \rightarrow$ $K K \pi X$

22.0 ± 2.0	CHUNG 85	SPEC	$8 \pi^- \rho \rightarrow$ $N \bar{K} \bar{K} \pi$
32.0 ± 3.0	604 ARMSTRONG 84	OMEG	$85 \pi^+ \rho \rightarrow$ $K \bar{K} \pi \pi \rho,$ $pp \rightarrow$ $K \bar{K} \pi pp$
24.0 ± 3.0	CHAUVAT 84	SPEC	ISR 31.5 pp
26 ± 12	EVANGELISTA 81	OMEG	$12 \pi^- \rho \rightarrow$ $\eta \pi^+ \pi^- \pi^- \rho$
29.0 ± 10.0	103 DIONISI 80	HBC	$4 \pi^- \rho \rightarrow$ $K \bar{K} \pi n$
25.0 ± 15.0	200 GURTU 79	HBC	$4.2 K^- \rho \rightarrow$ $n \eta 2 \pi$
28.3 ± 6.7	320 NACASCH 78	HBC	$0.7, 0.76 \bar{p} p \rightarrow$ $K \bar{K} 3 \pi$
24.0 ± 18.0	210 GRASSLER 77	HBC	$16 \pi^+ \rho$
10.0 ± 10.0	BOESEBECK 71	HBC	$16.0 \pi \rho \rightarrow$
30.0 ± 15.0	CAMPBELL 69	DBC	$2.7 \pi^+ d$

• • • We do not use the following data for averages, fits, limits, etc. • • •

44 ± 20	AUGUSTIN 90	DM2	$J/\psi \rightarrow$ $\gamma \eta \pi^+ \pi^-$
22 ± 5	TAKAMATSU 90	SPEC 0	$8 \pi^- \rho \rightarrow$ $K \bar{K} \pi n$
< 20	90 TAKAMATSU 90	SPEC 0	$8.95 \pi^- \rho \rightarrow$ $\eta \pi^+ \pi^- n$
~ 10.0	6 STANTON 79	CNTR	$8.5 \pi^- \rho \rightarrow$ $n 2 \gamma 2 \pi$
28 ± 5	150 7 DEFOIX 72	HBC	$0.7 \bar{p} p \rightarrow 7 \pi$
46 ± 9	180 7 DUBOC 72	HBC	$1.2 \bar{p} p \rightarrow 2 K 4 \pi$
37 ± 5	500 8 THUN 72	MMS	$13.4 \pi^- \rho$
60 ± 15	7 LORSTAD 69	HBC	$0.7 \bar{p} p, 4,5\text{-body}$
35.0 ± 10.0	7 DAHL 67	HBC	$1.6-4.2 \pi^- \rho$

5 From partial wave analysis of $K^+ \bar{K}^0 \pi^-$ system.
 6 From phase shift analysis of $\eta \pi^+ \pi^-$ system.
 7 Resolution is not unfolded.
 8 Seen in the missing mass spectrum.

$f_1(1285)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 4π	(29 ± 6)%	
Γ_2 $\pi^0 \pi^0 \pi^+ \pi^-$	(15 ± 9)%	S=1.1
Γ_3 $2\pi^+ 2\pi^-$	(15 ± 6)%	
Γ_4 $\rho^0 \pi^+ \pi^-$	dominates $2\pi^+ 2\pi^-$	
Γ_5 $4\pi^0$	< 7 × 10 ⁻⁴	CL=90%
Γ_6 $\eta \pi \pi$	(54 ± 15)%	
Γ_7 $a_0(980) \pi$ [ignoring $a_0(980) \rightarrow K \bar{K}$]	(44 ± 7)%	S=1.1
Γ_8 $\eta \pi \pi$ [excluding $a_0(980) \pi$]	(10 ± 7)%	S=1.1
Γ_9 $K \bar{K} \pi$	(9.7 ± 1.6)%	S=1.2
Γ_{10} $K \bar{K}^*(892)$	not seen	
Γ_{11} $\gamma \rho^0$	(6.6 ± 1.3)%	S=1.5
Γ_{12} $\phi \gamma$	(8.0 ± 3.1) × 10 ⁻⁴	
Γ_{13} $\gamma \gamma^*$		
Γ_{14} $\gamma \gamma$		

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 13 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 11.4$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-87				
x_7	-33	13			
x_8	-4	-5	-78		
x_9	46	-19	-38	-12	
x_{11}	-59	45	30	-11	-41
	x_2	x_3	x_7	x_8	x_9

See key on page 1343

Meson Full Listings
 $f_1(1285)$

$f_1(1285) \Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$				
$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT
VALUE (keV)				
<0.62	95	GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma^*)/\Gamma_{\text{total}}$				
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
1.4 ± 0.4 OUR AVERAGE		Error includes scale factor of 1.4.		
$1.18 \pm 0.25 \pm 0.20$	26	^{9,10} AIHARA	88B	TPC $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$
$2.30 \pm 0.61 \pm 0.42$		^{9,11} GIDAL	87	MRK2 $e^+e^- \rightarrow e^+e^-\eta\pi^+\pi^-$

⁹ Assuming a ρ -pole form factor.¹⁰ Published value multiplied by $\eta\pi\pi$ branching ratio 0.49.¹¹ Published value divided by 2 and multiplied by the $\eta\pi\pi$ branching ratio 0.49. $f_1(1285)$ BRANCHING RATIOS

$\Gamma(K\bar{K}\pi)/\Gamma(4\pi)$					
VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma_1 = \Gamma_9/(\Gamma_2+\Gamma_3)$	
0.33 ± 0.04 OUR FIT	Error includes scale factor of 1.2.				
0.32 ± 0.04 OUR AVERAGE	Error includes scale factor of 1.2.				
0.28 ± 0.05	¹² ARMSTRONG	89E	OMEG	$300 \rho\rho \rightarrow \rho\rho f_1(1285)$	
$0.37 \pm 0.03 \pm 0.05$	¹³ ARMSTRONG	89G	OMEG	$85 \pi\rho \rightarrow 4\pi X$	
¹² Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.					
¹³ 4π consistent with being entirely $\rho\pi\pi$.					
$\Gamma(K\bar{K}\pi)/\Gamma(\eta\pi\pi)$					
VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_9/\Gamma_6 = \Gamma_9/(\Gamma_7+\Gamma_8)$	
0.18 ± 0.04 OUR FIT	Error includes scale factor of 1.1.				
0.23 ± 0.06 OUR AVERAGE	Error includes scale factor of 1.2.				
0.42 ± 0.15	GURTU	79	HBC	$4.2 K^-\rho$	
0.5 ± 0.2	CORDEN	78	OMEG	$12-15 \pi^-\rho$	
0.20 ± 0.08	¹⁴ DEFOIX	72	HBC	$0.7 \bar{p}p \rightarrow 7\pi$	
0.16 ± 0.08	CAMPBELL	69	DBC	$2.7 \pi^+d$	
¹⁴ $K\bar{K}$ system characterized by the $l = 1$ threshold enhancement. (See under $a_0(980)$).					
$\Gamma(a_0(980)\pi \text{ [ignoring } a_0(980) \rightarrow K\bar{K}])/\Gamma(\eta\pi\pi)$					
VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_7/\Gamma_6 = \Gamma_7/(\Gamma_7+\Gamma_8)$	
0.82 ± 0.12 OUR FIT	Error includes scale factor of 1.1.				
0.69 ± 0.13 OUR AVERAGE					
0.72 ± 0.15	GURTU	79	HBC	$4.2 K^-\rho$	
0.6 ± 0.3	CORDEN	78	OMEG	$12-15 \pi^-\rho$	
-0.2					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.0 ± 0.3	GRASSLER	77	HBC	$16 \pi^{\mp}\rho$	
¹⁵ Assuming $\rho\pi\pi$ and $a_0(980)\pi$ intermediate states.					
$\Gamma(4\pi)/\Gamma(\eta\pi\pi)$					
VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_1/\Gamma_6 = (\Gamma_2+\Gamma_3)/(\Gamma_7+\Gamma_8)$	
0.54 ± 0.12 OUR FIT	Error includes scale factor of 1.1.				
0.41 ± 0.14 OUR AVERAGE					
$0.37 \pm 0.11 \pm 0.11$	BOLTON	92	MRK3	$J/\psi \rightarrow \gamma f_1(1285)$	
0.64 ± 0.40	GURTU	79	HBC	$4.2 K^-\rho$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.93 ± 0.30	¹⁵ GRASSLER	77	HBC	$16 \pi^{\mp}\rho$	

$\Gamma(K\bar{K}^*(892))/\Gamma_{\text{total}}$					
VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ	
not seen	NACASCH	78	HBC	$0.7, 0.76 \bar{p}p \rightarrow K\bar{K}3\pi$	

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma(2\pi^+2\pi^-)$					
VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_3	
1.0 ± 0.4	GRASSLER	77	HBC	$16 \text{ GeV } \pi^{\pm}\rho$	

$\Gamma(4\pi^0)/\Gamma_{\text{total}}$				
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
<7	90	ALDE	87	GAM4 $100 \pi^-\rho \rightarrow 4\pi^0 n$

$\Gamma(\phi\gamma)/\Gamma(K\bar{K}\pi)$				
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
$0.82 \pm 0.21 \pm 0.20$	19	BITYUKOV	88	SPEC $32.5 \pi^-\rho \rightarrow K^+K^-\pi^0 n$

$\Gamma(\gamma\rho^0)/\Gamma(K\bar{K}\pi)$				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>0.035	90	¹⁶ COFFMAN	90	MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$
¹⁶ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma K\bar{K}\pi) < 0.72 \times 10^{-3}$.				

$\Gamma(\gamma\rho^0)/\Gamma(2\pi^+2\pi^-)$					
VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_3	
0.45 ± 0.18 OUR FIT					
0.45 ± 0.18	¹⁷ COFFMAN	90	MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	
¹⁷ Using $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma\gamma\rho^0) = 0.25 \times 10^{-4}$ and $B(J/\psi \rightarrow \gamma f_1(1285) \rightarrow \gamma 2\pi^+2\pi^-) = 0.55 \times 10^{-4}$ given by MIR 88.					

$\Gamma(\gamma\rho^0)/\Gamma(a_0(980)\pi \text{ [ignoring } a_0(980) \rightarrow K\bar{K}])$					
VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ_7	
0.15 ± 0.04 OUR FIT	Error includes scale factor of 1.6.				
$0.10 \pm 0.03 \pm 0.02$	¹⁸ BURCHELL	91	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$	
¹⁸ Uses a result from COFFMAN 90, and includes an unknown branching ratio for $a_0(980) \rightarrow \eta\pi$.					

$\Gamma(\gamma\rho^0)/\Gamma_{\text{total}}$				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	95	BITYUKOV	91B	SPEC $32 \pi^-\rho \rightarrow \pi^+\pi^-\gamma n$

$\Gamma(\eta\pi\pi)/\Gamma(\gamma\rho^0)$					
VALUE	DOCUMENT ID	TECN	COMMENT	$\Gamma_6/\Gamma_{11} = (\Gamma_7+\Gamma_8)/\Gamma_{11}$	
8.2 ± 1.6 OUR FIT	Error includes scale factor of 1.8.				
7.5 ± 1.0	¹⁹ ARMSTRONG	92C	OMEG	$300 \rho\rho \rightarrow \rho\rho\pi^+\pi^-\gamma, \rho\rho\eta\pi^+\pi^-$	
¹⁹ Published value multiplied by 1.5.					

 $f_1(1285)$ REFERENCES

ARMSTRONG	93C	PL B307 394	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
ARMSTRONG	92C	ZPHY C54 371	+Barnes, Benayoun+	(ATHU, BARI, BIRM, CERN, CDEF)
BOLTON	92	PL B278 495	+Brown, Bunnell+	(Mark III Collab.)
BITYUKOV	91B	SJNP 54 318	+Borisov, Viktorov+	(SERP)
Translated from YAF 54 529.				
BURCHELL	91	NP B21 132 (suppl)		(Mark III Collab.)
FUKUI	91C	PL B267 293	+ (SUGI, NAGO, KEK, KYOT, MIYA, AKIT)	
AUGUSTIN	90	PR D42 10	+Cosme+	(DM2 Collab.)
COFFMAN	90	PR D41 1410	+De Jongh+	(Mark III Collab.)
TAKAMATSU	90	Hadron 89 Conf. p 71	+Ando+	(KEK)
ARMSTRONG	89	PL B221 216	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)	JPC
ARMSTRONG	89E	PL B228 536	+Benayoun (ATHU, BARI, BIRM, CERN, CDEF, CURIN+)	
ARMSTRONG	89G	ZPHY C43 55	+Bloodworth+ (CERN, BIRM, BARI, ATHU, CURIN+)	
RATH	89	PR D40 693	+Cason+ (NDAM, BRAN, BNL, CUNY, DUKE)	
AIHARA	88B	PL B209 107	+Alston-Garnjost+ (TPC-2\gamma Collab.)	
BIRMAN	88	PRL 61 1557	+Chung, Peaslee+ (BNL, FSU, IND, MASN) JP	
BITYUKOV	88	PL B203 327	+Borisov, Dorofeev+ (SERP)	
MIR	88	Photon-Photon '88 Conf., 126	+Binon, Bricman+ (Mark III Collab.)	
ALDE	87	PL B198 286	+Blaylock, Bolton, Brown+ (LANL, BRUX, SERP, LAF)	
BECKER	87	PRL 59 186	+Boyer, Butler, Cords, Abrams+ (LBL, SLAC, HARV)	
GIDAL	87	PRL 59 2012	+Imai+ (KEK, KYOT, NIRS, SAGA, INUS, TSUKU+)	IJP
ANDO	86	PR L57 1296	+Chung, Crittenden+ (FLOR, BNL, IND, MASN) JP	
REEVES	86	PR L57 779	+Fenow, Boehlein+ (BNL, FLOR, IND, MASN) JP	
CHUNG	85	PRL 55 779	+Bloodworth, Burns+ (ATHU, BARI, BIRM, CERN) JP	
ARMSTRONG	84	PL 146B 273	+Dorofeev, Dzheiyadin, Golovkin, Kulik+ (SERP)	
BITYUKOV	84	PL 144B 133	+Meritet, Bonino+ (CERN, CLER, UCLA, SACL)	
CHAUVAU	84	PL 148B 382		
TORNQVIST	82B	NP B203 268		
EVANGELISTA	81	NP B178 197	+ (BARI, BONN, CERN, DARE, LVP+)	
BROMBERG	80	PR D22 1513	+Haggerty, Abrams, Dzierba (CIT, FNAL, ILLC, IND)	
DIONISI	80	NP B169 1	+Gavillet+ (CERN, MADR, CDEF, STOH)	
GURTU	79	NP B151 181	+Gavillet, Blokzijl+ (CERN, ZEEM, NIJM, OXF)	
STANTON	79	PRL 42 346	+Brockman+ (OSU, CARL, MCGI, TNTO) JP	
CORDEN	78	NP B144 253	+Corbett, Alexander+ (BIRM, RHEL, TELA, LOWC) JP	
NACASCH	78	NP B135 203	+Defoix, Dobrzynski+ (PARIS, MADR, CERN)	
GRASSLER	77	NP B121 189	+ (AACHS, BERL, BONN, CERN, CRAC, HEIDH+)	
DEFOIX	72	NP B44 125	+Nascimento, Bizzarri+ (CDEF, CERN)	
DUBOC	72	NP B46 429	+Goldberg, Makowski, Donald+ (PARIS, LVP)	
THUN	72	PRL 28 1733	+Blieden, Finocchiaro, Bowen+ (STON, NEAS)	
BARBADIN-...	71	PR D4 2711	+Bardadin-Otwinowska, Hofmoki+ (WAR5)	
BOESEBECK	71	PL 34B 659	+ (AACH, BERL, BONN, CERN, CRAC, HEID, WARS)	
CAMPBELL	69	PRL 22 1204	+Lichtman, Loeffler+ (PURD)	
LORSTAD	69	NP B14 63	+D'Andlau, Astier+ (CDEF, CERN) JP	
D'ANDLAU	68	NP B5 693	+Astier, Barlow+ (CDEF, CERN, IRAD, LVP) IJP	
DAHL	67	PR 163 1377	+Hardy, Hess, Kirz, Miller (LRL) IJP	
OTHER RELATED PAPERS				
AIHARA	88C	PR D38 1	+Alston-Garnjost+ (TPC-2\gamma Collab.) JPC	
ASTON	85	PR D32 2255	+Carnegie, Dunwoodie+ (SLAC, CARL, CNRC)	
ATKINSON	84E	PL 138B 459	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
GAVILLET	82	ZPHY C16 119	+Armenteros+ (CERN, CDEF, PADO, ROMA)	
D'ANDLAU	65	PL 17 347	+Barlow, Adamson+ (CDEF, CERN, IRAD, LVP)	
MILLER	65	PRL 14 1074	+Chung, Dahl, Hess, Hardy, Kirz+ (LRL, UCB)	

Meson Full Listings

 $\eta(1295)$, $f_0(1300)$ $\eta(1295)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)

 $\eta(1295)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1295 ± 4	FUKUI	91C SPEC	$8.95 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1275	STANTON	79 CNTR	$8.4 \pi^- p \rightarrow n \eta 2\pi$

 $\eta(1295)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
53 ± 6	FUKUI	91C SPEC	$8.95 \pi^- p \rightarrow \eta \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 70	STANTON	79 CNTR	$8.4 \pi^- p \rightarrow n \eta 2\pi$

 $\eta(1295)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta \pi^+ \pi^-$	seen
Γ_2 $a_0(980)\pi$	seen
Γ_3 $\gamma\gamma$	

 $\eta(1295)$ $\Gamma(\text{I})\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(\eta \pi^+ \pi^-) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_3/\Gamma$
< 0.6	90	AIHARA	88C TPC	$e^+e^- \rightarrow e^+e^- \eta \pi^+ \pi^-$	
< 0.3		ANTREASYAN 87	CBAL	$e^+e^- \rightarrow e^+e^- \eta \pi \pi$	

 $\eta(1295)$ BRANCHING RATIOS

$\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	BIRMAN	88 MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$	
large	ANDO	86 SPEC	$8 \pi^- p \rightarrow n \eta \pi^+ \pi^-$	
large	STANTON	79 CNTR	$8.4 \pi^- p \rightarrow n \eta 2\pi$	

 $\eta(1295)$ REFERENCES

FUKUI	91C	PL B267 293	+	(SUGI, NAGO, KEK, KYOT, MIYA, AKIT)
AIHARA	88C	PR D38 1	+Alston-Garnjost+	(TPC-2 γ Collab.)
BIRMAN	88	PRL 61 1557	+Chung, Peaslee+	(BNL, FSU, IND, MASD) JP
ANTREASYAN	87	PR D36 2633	+Bartels, Besset+	(Crystal Ball Collab.)
ANDO	86	PRL 57 1296	+Imai+	(KEK, KYOT, NIRS, SAGA, INUS, TSUK+) IJP
STANTON	79	PRL 42 346	+Brockman+	(OSU, CARL, MCGI, TNT0) JP

 $f_0(1300)$ was $f_0(1400)$ was $\epsilon(1200)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

We list here all claims for highly elastic $\pi\pi$ S -wave resonances.

NOTE ON S -WAVE $\pi\pi$, $K\bar{K}$, AND $\eta\eta$ INTERACTIONS

In this note, we discuss results on the nonstrange $I^G J^{PC} = 0^+ 0^{++}$ partial wave (S -wave). It has been observed in its coupling to the $\pi\pi$, $K\bar{K}$, $\eta\eta$, 4π , and $\eta\eta'$ systems. We list the claimed resonances under five separate entries covering the mass range from $\pi\pi$ threshold to about 1600 MeV: $f_0(980)$, $f_0(1300)$, $f_0(1370)$, $f_0(1525)$, and $f_0(1590)$. See also the Notes on the $a_0(980)$ and on Non- $q\bar{q}$ Mesons.

Below 1100 MeV, the essential contributions come from $\pi\pi$ and $K\bar{K}$ final states. The $I = 0$ S -wave $\pi\pi$ phase shift δ_0^0 (GRAYNER 74, ROSSELET 77) shows a rapid step of 180° near the $K\bar{K}$ threshold, which is the $f_0(980)$ resonance, superimposed over a large "background" phase shift which reaches

90° a little above 1000 MeV. Above this energy, it continues to grow slowly, as expected for a very broad resonance. This is the $f_0(1300)$, which was the $f_0(1400)$ of our 1992 edition and which evolved from the $\epsilon(1200)$ (our 1976 edition). The $\pi\pi$ S -wave inelasticity is not accurately known, and the $\pi\pi \rightarrow K\bar{K}$ cross sections (COHEN 80, ETKIN 82, POLYCHRONAKOS 79, WETZEL 76) may have large uncertainties.

Using the data available in 1986, including data on $\pi^+ \pi^-$ and $K^+ K^-$ central production (AKESSON 86) and on $J/\psi(1S) \rightarrow \phi \pi \pi$ and $J/\psi(1S) \rightarrow K^+ K^-$, the coupled-channel analyses of AU 87 suggested that four resonances were needed to describe the 1-GeV region. But, adding to their analysis new data on $J/\psi(1S) \rightarrow \phi \pi \pi$ and $\phi K\bar{K}$ (MALIK 86, FALVARD 88, LOCKMAN 89), on $D_s^+ \rightarrow \pi^+ \pi^+ \pi^-$ (ANJOS 89), and on $K^+ K^- \rightarrow K_S^0 K_S^0$ (ASTON 88), and relaxing the constraints on $\pi\pi \rightarrow K\bar{K}$ cross sections (LINDENBAUM 92), MORGAN 93 now favors a single Breit-Wigner resonance for the $f_0(980)$ described by two poles, $E^{II} = (988 \pm 10) - (24 \pm 6)i$ MeV and $E^{III} = 978 - 28i$ MeV, corresponding to resonance parameters $m_0 = 983$ MeV, $\Gamma_0 = 52$ MeV, and $g_{K\bar{K}}^2/g_{\pi\pi}^2 \approx 0.85$. In a similar coupled-channel analysis, but limited to fitting elastic-scattering data (GRAYNER 74, ROSSELET 77), and the centrally produced $\pi\pi$ and $K\bar{K}$ data (AKESSON 86), ZOU 93 also finds two poles in the 1 GeV region, $E^{II} = 988 - 25i$ MeV in agreement with MORGAN 93, while their $E^{III} = 914 - 219i$ MeV, has a large imaginary part.

Below 900 MeV, BECKER 79B excludes a resonance behaviour for δ_0^0 in their $\pi^- p$ (polarized) $\rightarrow \pi^+ \pi^- n$ data. In contrast, SVEC 92, using their data on $\pi^+ n$ (polarized) $\rightarrow \pi^+ \pi^- p$, which only span the range 600–900 MeV, suggests a narrow scalar state at 750 MeV ($\Gamma = 100$ –200 MeV); the associated δ_0^0 values differ substantially from the recent consensus and would reopen the old UP-DOWN ambiguity of the early 1970's (see our 1984 edition). The DOWN option was long ago selected by requiring continuity of δ_0^0 and its associated inelasticity above 900 MeV to reproduce the observed rapid variations of the $\pi\pi$ elastic cross sections through the $K\bar{K}$ threshold region (PRO-TOPOPESCU 73). Indeed, attempts to introduce an UP-type δ_0^0 tend to contradict dispersion relations (PENNINGTON 93). Thus, the interpretation of SVEC 92 must be treated with some reservation.

Both MORGAN 93 and ZOU 93 have to add a background term in the $f_0(980)$ region, which may be interpreted as a very broad ($\Gamma \approx 700$ MeV) resonance with mass in the 1000–1500 MeV range. Many broad $\pi\pi$ elastic resonances have been claimed in the 1000–1500 MeV region, and we collect them all under one entry, the $f_0(1300)$.

Above 1200 MeV, there is increasing evidence that the $0^+ 0^{++}$ sector is dominated by the 4π channel and that the $\eta\eta$ channel cannot be neglected. For the 4π channel, the information comes mainly from the analysis of $\bar{p}n \rightarrow 5\pi$ (GASPERO 93, ADAMO 93, AMSLER 94). AMSLER 94 finds a large production of a 0^{++} resonance decaying into 4π , mostly $\rho\rho$, with $M = 1374 \pm 38$ MeV and $\Gamma = 357 \pm 61$ MeV, and quote a $4\pi:2\pi$

See key on page 1343

Meson Full Listings

 $f_0(1300)$

branching ratio of order 5:1. The same resonance is observed in $\bar{p}p \rightarrow \eta\eta\pi^0$ (AMSLER 93C) and in $\bar{p}p \rightarrow 3\pi^0$ (AKER 91) leading to an $\eta\eta:\pi\pi$ branching ratio of order 1:3.

Thus we collect all the broad $\pi\pi$ inelastic S -wave resonance claims under a new entry, the $f_0(1370)$, although they could be the $f_0(1300)$, provided the inelasticity of the latter is in fact larger than is presently believed.

In a simultaneous fit to the $\bar{p}p \rightarrow 3\pi^0$ and $\bar{p}p \rightarrow \eta\eta\pi^0$ data, ANISOVITCH 94 find again the $f_0(1370)$ and, in addition, they observe another scalar resonance with $M = 1520 \pm 25$ MeV and $\Gamma = 146 \pm 25$ MeV. We list this observation under the $f_0(1525)$, although it perhaps does not belong there.

The fifth scalar entry is the $f_0(1590)$ of the GAMS collaboration (BINON 83).

 $f_0(1300)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1000-1500 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
≈ 1000	¹ MORGAN 93	RVUE	
1472 ± 12	ARMSTRONG 91	OMEG 300 $pp \rightarrow pp\pi\pi$, $ppK\bar{K}$	
1440 ± 20	CHEN 91	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-, \gamma K\bar{K}$	
1275 ± 20	BREAKSTONE 90	SFM $62 pp \rightarrow pp\pi^+\pi^-$	
1440 ± 50	BOLONKIN 88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
1420 ± 20	AKESSON 86	SPEC $63 pp \rightarrow pp\pi^+\pi^-$	
1220 ± 40	ALDE 86D	GAM4 $100 \pi^- p \rightarrow n2\eta$	
1463 ± 9	ETKIN 82B	MPS $23 \pi^- p \rightarrow n2K_S^0$	
~ 1237	TORNQVIST 82	RVUE	
1425 ± 15	WICKLUND 80	SPEC $6 \pi N \rightarrow K^+ K^- N$	
~ 1300	POLYCHRO... 79	STRC $7 \pi^- p \rightarrow n2K_S^0$	
1256	FROGGATT 77	RVUE $\pi^+\pi^-$ channel	
¹ Combining new data on $f_0(980)$ production in J/ψ and D_S decays with earlier information on central production and elastic $\pi\pi$, $K\bar{K}$ processes.			

 $f_0(1300)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 to 400 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
≈ 700	² MORGAN 93	RVUE	
195 ± 33	ARMSTRONG 91	OMEG 300 $pp \rightarrow pp\pi\pi$, $ppK\bar{K}$	
160 ± 40	CHEN 91	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-, \gamma K\bar{K}$	
285 ± 60	BREAKSTONE 90	SFM $62 pp \rightarrow pp\pi^+\pi^-$	
250 ± 80	BOLONKIN 88	SPEC $40 \pi^- p \rightarrow K_S^0 K_S^0 n$	
460 ± 50	AKESSON 86	SPEC $63 pp \rightarrow pp\pi^+\pi^-$	
320 ± 40	ALDE 86D	GAM4 $100 \pi^- p \rightarrow n2\eta$	
~ 1400	TORNQVIST 82	RVUE	
160 ± 30	WICKLUND 80	SPEC $6 \pi N \rightarrow K^+ K^- N$	
~ 150	POLYCHRO... 79	STRC $7 \pi^- p \rightarrow n2K_S^0$	
~ 400	FROGGATT 77	RVUE $\pi^+\pi^-$ channel	
² Combining new data on $f_0(980)$ production in J/ψ and D_S decays with earlier information on central production and elastic $\pi\pi$, $K\bar{K}$ processes.			
³ Width defined as distance between 45 and 135° phase shift.			

 $f_0(1300)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	$(93.6^{+1.9}_{-1.5})\%$
Γ_2 $K\bar{K}$	$(7.5 \pm 0.9)\%$
Γ_3 $\eta\eta$	seen
Γ_4 $\gamma\gamma$	seen
Γ_5 e^+e^-	not seen

 $f_0(1300)$ PARTIAL WIDTHS

$\Gamma(\gamma\gamma)$	DOCUMENT ID	TECN	COMMENT
VALUE (keV)			
5.4 ± 2.3	⁴ MORGAN 90	RVUE	$\gamma\gamma \rightarrow \pi^+\pi^-, \pi^0\pi^0$
⁴ Error includes spread of different solutions. Authors report strong correlation with $\gamma\gamma$ width of $f_2(1270)$.			

$\Gamma(e^+e^-)$	DOCUMENT ID	TECN	COMMENT
VALUE (eV)			
<20	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^0\pi^0$

 $f_0(1300)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
VALUE			
$0.936^{+0.019}_{-0.015}$	GORLICH 80	ASPK	$17.18 \pi^- p$ polarized
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.93	TORNQVIST 82	RVUE	
0.93	LOVERRE 80	HBC	$4 \pi^- p \rightarrow K\bar{K}N$
0.73	HYAMS 75	ASPK	$17.2 \pi^- p \rightarrow n\pi^+\pi^-$

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$	DOCUMENT ID	TECN	COMMENT
VALUE			
0.08 ± 0.01	COSTA... 80	OMEG	$10 \pi^- p \rightarrow K^+ K^- n$

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
VALUE			
0.02	ALDE 86D	GAM4	$100 \pi^- p \rightarrow n2\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	BINON 83	GAM2	$38 \pi^- p \rightarrow 2\eta n$

 $f_0(1300)$ REFERENCES

MORGAN 93	PR D48 1185	+Pennington	(RAL, DURH)
Also 93C	NC A Conf. Suppl.	Morgan	(RAL)
ARMSTRONG 91	ZPHY C51 351	+Benayoun+	(ATHU, BARI, BIRM, CERN, CDEF)
CHEN 91	Hadron 91 Conf.		(Mark III Collab.)
SLAC-PUB-5669			
BREAKSTONE 90	ZPHY C48 569	+	(ISU, BGNA, CERN, DORT, HEIDH, WARS)
MORGAN 90	ZPHY C48 623	+Pennington	(RAL, DURH)
BOLONKIN 88	NP B309 426	+Bioshenko, Gorin+	(ITEP, SERP)
VOROBYEV 88	SJNP 48 273	+Golubev, Dolinsky, Druzhinin+	(NOVO)
Translated from YAF 48 436.			
AKESSON 86	NP B264 154	+Albrow, Almeded+	(Axial Field Spec. Collab.)
ALDE 86D	NP B269 485	+Binon, Bricman+	(BELG, LAPP, SERP, CERN)
BINON 83	NC 78A 313	+Donskov, Dutell+	(BELG, LAPP, SERP, CERN)
BOLONKIN 88	PR D25 1786	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
ETKIN 82B	PRL 49 624		(HEL)
TORNQVIST 82	PRL 49 624		(HEL)
COSTA... 80	NP B175 402	Costa De Beauregard+	(BARI, BONN, CERN+)
GORLICH 80	NP B174 16	+Niczyporuk+	(CRAC, MPIM, CERN, ZEEM)
LOVERRE 80	ZPHY C6 187	+Armenteros, Dionisi+	(CERN, CDEF, MADR, STOH) IJP
WICKLUND 80	PRL 45 1469	+Ayres, Cohen, Diebold, Pawlicki	(ANL)
POLYCHRO... 79	NP D19 1317	Polychronakos, Cason, Bishop+	(NDAM, ANL) IJP
FROGGATT 77	NP B129 89	+Petersen	(GLAS, NORD)
HYAMS 75	NP B100 205	+Jones, Weillhammer, Blum, Dietl+	(CERN, MPIM)

OTHER RELATED PAPERS

AMSLER 94	PL B322 431	+Armstrong+	(Crystal Barrel Collab.)
ANISOVICH 94	PL B323 233	+Armstrong+	(Crystal Barrel Collab.)
ADAMO 93	NP A558 13C	+Agnello+	(OBELIX Collab.)
GASPERO 93	NP A562 407		(ROMAI)
SVEC 92	PR D45 55	+de Lesquen, van Rossum	(MCGI, SACL)
SVEC 92B	PR D45 1518	+de Lesquen, van Rossum	(MCGI, SACL)
SVEC 92C	PR D46 949	+de Lesquen, van Rossum	(MCGI, SACL)
MORGAN 91	PL B258 444	+Pennington	(RAL, DURH)
WEINSTEIN 89	UTP T 89 03	+Isgur	(TNTO)
ALDE 88	PL B201 160	+Bellazzini, Binon+	(SERP, BELG, LANL, LAPP, PISA)
AU 87	PR D35 1633	+Morgan, Pennington	(DURH, RAL)
ACHASOV 84	ZPHY C22 53	+Devyanin, Shestakov	(NOVO)
CASON 83	PR D28 1586	+Cannata, Baumbaugh, Bishop+	(NDAM, ANL)
WEINSTEIN 83B	PR D27 588	+Isgur	(TNTO)
BECKER 79	NP B151 46	+Bianar, Blum+	(MPIM, CERN, ZEEM, CRAC)
BECKER 79B	NP B150 301	+Bianar, Blum+	(MPIM, CERN, ZEEM, CRAC)
CORDEN 79	NP B157 250	+Dowell, Garvey+	(BIRM, RHEL, TEL, LOWC) JP
CASON 76	PRL 36 1485	+Polychronakos, Bishop, Biswas+	(NDAM, ANL) IJ
WETZEL 76	NP B115 208	+Freudenreich, Beusch+	(ETH, CERN, LOIC)
ESTABROOKS 74	NP B79 301	+Martin	(DURH)
GRAYR 74	NP B75 189	+Hyams, Blum, Dietl+	(CERN, MPIM)
HYAMS 73	NP B64 134	+Jones, Weillhammer, Blum, Dietl+	(CERN, MPIM)

Meson Full Listings

$\pi(1300), a_2(1320)$

$\pi(1300)$

$$I^G(J^{PC}) = 1^-(0^{-+})$$

$\pi(1300)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1300 ± 100 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1190 ± 30	ZIELINSKI 84	SPEC	200 $\pi^+ Z \rightarrow Z3\pi$
1240 ± 30	BELLINI 82	SPEC	40 $\pi^- A \rightarrow A3\pi$
1273.0 ± 50.0	¹ AARON 81	RVUE	
1342 ± 20	BONESINI 81	OMEG	12 $\pi^- p \rightarrow p3\pi$
~ 1400	DAUM 81B	SPEC	63,94 $\pi^- p$

¹ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

$\pi(1300)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200 to 600 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
440 ± 80	ZIELINSKI 84	SPEC	200 $\pi^+ Z \rightarrow Z3\pi$
360 ± 120	BELLINI 82	SPEC	40 $\pi^- A \rightarrow A3\pi$
580.0 ± 100.0	² AARON 81	RVUE	
220 ± 70	BONESINI 81	OMEG	12 $\pi^- p \rightarrow p3\pi$
~ 600	DAUM 81B	SPEC	63,94 $\pi^- p$

² Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

$\pi(1300)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\pi(\pi\pi)_{S\text{-wave}}$	seen
Γ_3 $f_0(1300)\pi$	

$\pi(1300)$ BRANCHING RATIOS

$\Gamma(\pi(\pi\pi)_{S\text{-wave}})/\Gamma(\rho\pi)$	DOCUMENT ID	TECN	Γ_2/Γ_1
2.12	³ AARON 81	RVUE	

³ Uses multichannel Aitchison-Bowler model (BOWLER 75). Uses data from DAUM 80 and DANKOWYCH 81.

$\pi(1300)$ REFERENCES

ZIELINSKI 84	PR D30 1855	+Berg, Chandee, Cihangir+ (ROCH, MINN, FNAL)
BELLINI 82	PRL 48 1697	+Frabetti, Ivanshin, Litkin+ (MILA, BGNA, JINR)
AARON 81	PR D24 1207	+Longacre (NEAS, BNL)
BONESINI 81	PL 103B 75	+Donald+ (MILA, LVP, DARE, CERN, BARI, BONN)
DANKOWYCH... 81	PRL 46 580	Dankowych+ (TNTD, BNL, CARL, MCGI, OHIO)
DAUM 81B	NP B182 269	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
DAUM 80	PL 89B 281	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
BOWLER 75	NP B97 227	+Game, Aitchison, Dainton (OXFPT, DARE)

$a_2(1320)$

$$I^G(J^{PC}) = 1^-(2^{++})$$

$a_2(1320)$ MASS

3 π AND $K^\pm K_S^0$ MODES

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1318.4 ± 0.6 OUR AVERAGE	Includes data from the 4 datablocks that follow this one. Error includes scale factor of 1.1.				
3π MODE					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT

The data in this block is included in the average printed for a previous datablock.


1318.5 ± 1.6 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
1310 ± 5	ARMSTRONG 90	OMEG 0		300,0 $\rho\rho \rightarrow \rho\rho\pi^+ \pi^- \pi^0$
1323.8 ± 2.3	4022	AUGUSTIN 89	DM2 ±	$J/\psi \rightarrow \rho^\pm a_2^\mp$
1320.6 ± 3.1	3562	AUGUSTIN 89	DM2 0	$J/\psi \rightarrow \rho^0 a_2^0$
1317.0 ± 2.0	25000	¹ DAUM 80C	SPEC -	63,94 $\pi^- p \rightarrow 3\pi\rho$
1320.0 ± 10.0	1097	¹ BALTAY 78B	HBC +0	15 $\pi^+ p \rightarrow \rho4\pi$
1306.0 ± 8.0		FERRERSORIA 78	OMEG -	9 $\pi^- p \rightarrow \rho3\pi$
1318 ± 7	1600	¹ EMMS 75	DBC 0	4 $\pi^+ n \rightarrow \rho(3\pi)^0$
1315 ± 5		¹ ANTIPOV 73C	CNTR -	25,40 $\pi^- p \rightarrow \rho\eta\pi^-$
1306 ± 9	1580	CHALOUPKA 73	HBC -	3.9 $\pi^- p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1305 ± 14	CONDO 93	SHF		$\gamma\rho \rightarrow \eta\pi^+ \pi^+ \pi^-$
1310 ± 2	¹ EVANGELISTA 81	OMEG -		12 $\pi^- p \rightarrow 3\pi\rho$
1343.0 ± 11.0	490	BALTAY 78B	HBC 0	15 $\pi^+ p \rightarrow \Delta3\pi$
1309 ± 5	5000	BINNIE 71	MMS -	$\pi^- p$ near a_2 thresh-old
1299.0 ± 6.0	28000	BOWEN 71	MMS -	5 $\pi^- p$
1300 ± 6.0	24000	BOWEN 71	MMS +	5 $\pi^+ p$
1309.0 ± 4.0	17000	BOWEN 71	MMS -	7 $\pi^- p$
1306.0 ± 4.0	941	ALSTON... 70	HBC +	7.0 $\pi^+ p \rightarrow 3\pi\rho$

¹ From a fit to $J^P = 2^+ \rho\pi$ partial wave.

WEIGHTED AVERAGE
1318.5 ± 1.6 (Error scaled by 1.3)



$a_2(1320)$ mass, 3 π mode (MeV)

$K^\pm K_S^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					
1318.1 ± 0.7 OUR AVERAGE					
1319.0 ± 5.0	4700	^{2,3} CLELAND 82B	SPEC +		50 $\pi^+ p \rightarrow K_S^0 K^+ \rho$
1324.0 ± 6.0	5200	^{2,3} CLELAND 82B	SPEC -		50 $\pi^- p \rightarrow K_S^0 K^- \rho$
1320.0 ± 2.0	4000	CHABAUD 80	SPEC -		17 $\pi^- A \rightarrow K_S^0 K^- A$
1312.0 ± 4.0	11000	CHABAUD 78	SPEC -		9.8 $\pi^- p \rightarrow K^- K_S^0 \rho$
1316.0 ± 2.0	4730	CHABAUD 78	SPEC -		18.8 $\pi^- p \rightarrow K^- K_S^0 \rho$
1318 ± 1	^{2,4} MARTIN 78D	SPEC -			10 $\pi^- p \rightarrow K_S^0 K^- \rho$
1320.0 ± 2.0	2724	MARGULIE 76	SPEC -		23 $\pi^- p \rightarrow K^- K_S^0 \rho$
1313.0 ± 4.0	730	FOLEY 72	CNTR -		20.3 $\pi^- p \rightarrow K^- K_S^0 \rho$
1319.0 ± 3.0	1500	⁴ GRAYER 71	ASPK -		17.2 $\pi^- p \rightarrow K^- K_S^0 \rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1330.0 ± 11.0	1000	^{2,3} CLELAND 82B	SPEC +		30 $\pi^+ p \rightarrow K_S^0 K^+ \rho$
1324.0 ± 5.0	350	HYAMS 78	ASPK +		12.7 $\pi^+ p \rightarrow K^+ K_S^0 \rho$

² From a fit to $J^P = 2^+$ partial wave.
³ Number of events evaluated by us.
⁴ Systematic error in mass scale subtracted.

$\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					
1320.1 ± 2.0 OUR AVERAGE					
1325.1 ± 5.1		AOYAGI 93	BKEI		$\pi^- p \rightarrow \eta\pi^- p$
1324 ± 5		ARMSTRONG 93C	SPEC 0		$\bar{p}p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$
1317.7 ± 1.4 ± 2.0		BELADIDZE 93	VES		$37\pi^- N \rightarrow \eta\pi^- N$
1323 ± 8	1000	⁵ KEY 73	OSPK -		6 $\pi^- p \rightarrow \rho\pi^- \eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1336.2 ± 1.7	2561	DELFOSSÉ 81	SPEC +		$\pi^\pm p \rightarrow \rho\pi^\pm \eta$
1330.7 ± 2.4	1653	DELFOSSÉ 81	SPEC -		$\pi^\pm p \rightarrow \rho\pi^\pm \eta$
1324 ± 8	6200	^{5,6} CONFORTO 73	OSPK -		6 $\pi^- p \rightarrow \rho\pi^- \eta$

⁵ Error includes 5 MeV systematic mass-scale error.
⁶ Missing mass with enriched MMS = $\eta\pi^-$, $\eta = 2\gamma$.

$\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1327.0 ± 10.7	BELADIDZE 93	VES	$37\pi^- N \rightarrow \eta'\pi^- N$

The data in this block is included in the average printed for a previous datablock.

See key on page 1343

Meson Full Listings
 $a_2(1320)$ $a_2(1320)$ WIDTH 3π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
103.4 ± 2.1 OUR AVERAGE						
120 ± 10		ARMSTRONG 90	OMEG	0	300.0 $p\rho \rightarrow \rho\rho\pi^+\pi^-\pi^0$	
107.0 ± 9.7	4022	AUGUSTIN 89	DM2	±	$J/\psi \rightarrow \rho^\pm a_2^\mp$	
118.5 ± 12.5	3562	AUGUSTIN 89	DM2	0	$J/\psi \rightarrow \rho^0 a_2^0$	
97 ± 5		⁷ EVANGELISTA 81	OMEG	-	12 $\pi^-\rho \rightarrow 3\pi\rho$	
96.0 ± 9.0	25000	⁷ DAUM	80C	SPEC	-	63,94 $\pi^-\rho \rightarrow 3\pi\rho$
110.0 ± 15.0	1097	⁷ BALTAY	78B	HBC	+0	15 $\pi^+\rho \rightarrow \rho 4\pi$
112 ± 18	1600	⁷ EMMS	75	DBC	0	4 $\pi^+n \rightarrow \rho(3\pi)^0$
122 ± 14	1200	^{7,8} WAGNER	75	HBC	0	7 $\pi^+\rho \rightarrow \Delta^{++}(3\pi)^0$
115 ± 15		⁷ ANTIPOV	73C	CNTR	-	25,40 $\pi^-\rho \rightarrow \rho\eta\pi^-$
99 ± 15	1580	CHALOUKKA 73	HBC	-	3,9 $\pi^-\rho$	
105.0 ± 5.0	28000	BOWEN 71	MMS	-	5 $\pi^-\rho$	
99.0 ± 5.0	24000	BOWEN 71	MMS	+	5 $\pi^+\rho$	
103.0 ± 5.0	17000	BOWEN 71	MMS	-	7 $\pi^-\rho$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
120 ± 40		CONDO 93	SHF		$\gamma\rho \rightarrow \eta\pi^+\pi^+\pi^-$	
115.0 ± 14.0	490	BALTAY 78B	HBC	0	15 $\pi^+\rho \rightarrow \Delta 3\pi$	
72 ± 26	5000	BINNIE 71	MMS	-	$\pi^-\rho$ near a_2 threshold	
79.0 ± 12.0	941	ALSTON-... 70	HBC	+	7.0 $\pi^+\rho \rightarrow 3\pi\rho$	

⁷From a fit to $J^P = 2^+ \rho\pi$ partial wave.⁸Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass. $K^\pm K_S^0$ AND $\eta\pi$ MODES

VALUE (MeV)	DOCUMENT ID
107 ± 5 OUR ESTIMATE	
110.1 ± 1.9 OUR AVERAGE	Includes data from the 2 datablocks that follow this one.

 $K^\pm K_S^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
109.8 ± 2.4 OUR AVERAGE					
112.0 ± 20.0	4700	^{9,10} CLELAND 82B	SPEC	+	50 $\pi^+\rho \rightarrow K_S^0 K^+ p$
120.0 ± 25.0	5200	^{9,10} CLELAND 82B	SPEC	-	50 $\pi^-\rho \rightarrow K_S^0 K^- p$
106.0 ± 4.0	4000	CHABAUD 80	SPEC	-	17 $\pi^-\rho \rightarrow K_S^0 K^- A$
126.0 ± 11.0	11000	CHABAUD 78	SPEC	-	9.8 $\pi^-\rho \rightarrow K^- K_S^0 p$
101.0 ± 8.0	4730	CHABAUD 78	SPEC	-	18.8 $\pi^-\rho \rightarrow K^- K_S^0 p$
113 ± 4		^{9,11} MARTIN 78D	SPEC	-	10 $\pi^-\rho \rightarrow K_S^0 K^- p$
105.0 ± 8.0	2724	¹¹ MARGULIE 76	SPEC	-	23 $\pi^-\rho \rightarrow K^- K_S^0 p$
113.0 ± 19.0	730	FOLEY 72	CNTR	-	20.3 $\pi^-\rho \rightarrow K^- K_S^0 p$
123.0 ± 13.0	1500	¹¹ GRAYNER 71	ASPK	-	17.2 $\pi^-\rho \rightarrow K^- K_S^0 p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
121.0 ± 51.0	1000	^{9,10} CLELAND 82B	SPEC	+	30 $\pi^+\rho \rightarrow K_S^0 K^+ p$
110.0 ± 18.0	350	HYAMS 78	ASPK	+	12.7 $\pi^+\rho \rightarrow K^+ K_S^0 p$

⁹From a fit to $J^P = 2^+$ partial wave.¹⁰Number of events evaluated by us.¹¹Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass. $\eta\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
110.9 ± 3.3 OUR AVERAGE					
118 ± 10		ARMSTRONG 93C	SPEC	0	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
103 ± 6 ± 3		BELADIDZE 93	VES		$37\pi^-N \rightarrow \eta\pi^-N$
112.2 ± 5.7	2561	DELFOSSÉ 81	SPEC	+	$\pi^\pm\rho \rightarrow \rho\pi^\pm\eta$
116.6 ± 7.7	1653	DELFOSSÉ 81	SPEC	-	$\pi^\pm\rho \rightarrow \rho\pi^\pm\eta$
108 ± 9	1000	KEY 73	OSPK	-	6 $\pi^-\rho \rightarrow \rho\pi^-\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
104 ± 9	6200	¹² CONFORTO 73	OSPK	-	6 $\pi^-\rho \rightarrow \rho MM^-$
¹² Model dependent.					

 $\eta'\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
106 ± 32	BELADIDZE 93	VES	$37\pi^-N \rightarrow \eta'\pi^-N$

 $a_2(1320)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\rho\pi$	(70.1 ± 2.7) %	S=1.2
Γ_2 $\eta\pi$	(14.5 ± 1.2) %	
Γ_3 $\omega\pi\pi$	(10.6 ± 3.2) %	S=1.3
Γ_4 $K\bar{K}$	(4.9 ± 0.8) %	
Γ_5 $\eta'(958)\pi$	(5.7 ± 1.1) × 10 ⁻³	
Γ_6 $\pi^\pm\gamma$	(2.8 ± 0.6) × 10 ⁻³	
Γ_7 $\gamma\gamma$	(9.7 ± 1.0) × 10 ⁻⁶	
Γ_8 $\pi^+\pi^-\pi^-$	< 8 %	CL=90%
Γ_9 e^+e^-	< 2.3 × 10 ⁻⁷	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 18 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 9.3$ for 15 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	10		
x_3	-89	-46	
x_4	-1	-2	-24
	x_1	x_2	x_3

 $a_2(1320)$ PARTIAL WIDTHS $\Gamma(\pi^\pm\gamma)$

VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT
295 ± 60	CIHANGIR 82	SPEC	+	200 π^+A
• • • We do not use the following data for averages, fits, limits, etc. • • •				
461 ± 110	¹² MAY	77	SPEC	± 9.7 γA

 $\Gamma(\gamma\gamma)$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.04 ± 0.09 OUR AVERAGE					
1.26 ± 0.26 ± 0.18	36	BARU 90	MD1		$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.00 ± 0.07 ± 0.15	415	BEHREND 90C	CELL	0	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.03 ± 0.13 ± 0.21		BUTLER 90	MRK2		$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$
1.01 ± 0.14 ± 0.22	85	OEST 90	JADE		$e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.90 ± 0.27 ± 0.15	56	¹³ ALTHOFF 86	TASS	0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
1.14 ± 0.20 ± 0.26		¹⁴ ANTREASYAN 86	CBAL	0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
1.06 ± 0.18 ± 0.19		BERGER 84C	PLUT	0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.81 ± 0.19 ± 0.42	35	¹³ BEHREND 83B	CELL	0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.84 ± 0.07 ± 0.15		¹³ FRAZER 83	JADE	0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$
0.77 ± 0.18 ± 0.27	22	¹⁴ EDWARDS 82F	CBAL	0	$e^+e^- \rightarrow e^+e^-\pi^0\eta$

¹³From $\rho\pi$ decay mode.¹⁴From $\eta\pi^0$ decay mode. $\Gamma(e^+e^-)$

VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
< 25	90	VOROBYEV 88	ND	$e^+e^- \rightarrow \pi^0\eta$

 $a_2(1320)$ $\Gamma(I)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.126 ± 0.007 ± 0.028	¹⁵ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.081 ± 0.006 ± 0.027	¹⁶ ALBRECHT 90G	ARG	$e^+e^- \rightarrow e^+e^-K^+K^-$

¹⁵Using an incoherent background.¹⁶Using a coherent background.

Meson Full Listings

$a_2(1320)$

$a_2(1320)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\rho\pi)$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_1
------------------------------------	------	-------------	------	-----	---------	---------------------

0.070 ± 0.012 OUR FIT						
0.078 ± 0.017		CHABAUD	78		RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.056 ± 0.014	50	17 CHALOUKPA	73	HBC	-	3.9 $\pi^- \rho$
0.097 ± 0.018	113	17 ALSTON...	71	HBC	+	7.0 $\pi^+ \rho$
0.06 ± 0.03		17 ABRAMOVI...	70B	HBC	-	3.93 $\pi^- \rho$
0.054 ± 0.022		17 CHUNG	68	HBC	-	3.2 $\pi^- \rho$

¹⁷Included in CHABAUD 78 review.

$\Gamma(\eta\pi)/[\Gamma(\rho\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_2/(\Gamma_1 + \Gamma_2 + \Gamma_4)$
--	------	-------------	------	-----	---------	---

0.162 ± 0.012 OUR FIT						
0.140 ± 0.028 OUR AVERAGE						
0.13 ± 0.04		ESPIGAT	72	HBC	±	0.0 $\bar{p} \rho$
0.15 ± 0.04	34	BARNHAM	71	HBC	+	3.7 $\pi^+ \rho$

$\Gamma(\eta\pi)/\Gamma(\rho\pi)$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
-----------------------------------	------	-------------	------	-----	---------	---------------------

0.207 ± 0.018 OUR FIT						
0.213 ± 0.020 OUR AVERAGE						
0.18 ± 0.05		FORINO	76	HBC	-	11 $\pi^- \rho$
0.22 ± 0.05	52	ANTIPOV	73	CNTR	-	40 $\pi^- \rho$
0.211 ± 0.044	149	CHALOUKPA	73	HBC	-	3.9 $\pi^- \rho$
0.246 ± 0.042	167	ALSTON...	71	HBC	+	7.0 $\pi^+ \rho$
0.25 ± 0.09	15	BOECKMANN	70	HBC	+	5.0 $\pi^+ \rho$
0.23 ± 0.08	22	ASCOLI	68	HBC	-	5 $\pi^- \rho$
0.12 ± 0.08		CHUNG	68	HBC	-	3.2 $\pi^- \rho$
0.22 ± 0.09		CONTE	67	HBC	-	11.0 $\pi^- \rho$

$\Gamma(\eta'(958)\pi)/\Gamma_{total}$	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ
--	-----	-------------	------	-----	---------	-------------------

0.040 ± 0.007 OUR AVERAGE						
0.047 ± 0.010 ± 0.004		21 BELADIDZE	93	VES	37 $\pi^- N \rightarrow a_2^- N$	
0.034 ± 0.008 ± 0.005		BELADIDZE	92	VES	36 $\pi^- C \rightarrow a_2^- C$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
<0.006	95	ALDE	92B	GAM2	38,100 $\pi^- \rho \rightarrow \eta' \pi^0 \rho$	
<0.02	97	BARNHAM	71	HBC	+	3.7 $\pi^+ \rho$
0.004 ± 0.004		BOESEBECK	68	HBC	+	8 $\pi^+ \rho$

$\Gamma(\eta'(958)\pi)/\Gamma(\rho\pi)$	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_1
---	-----	-------------	------	-----	---------	---------------------

0.054 ± 0.009 OUR FIT						
0.048 ± 0.012 OUR AVERAGE						
<0.011	90	EISENSTEIN	73	HBC	-	5 $\pi^- \rho$
<0.04		ALSTON...	71	HBC	+	7.0 $\pi^+ \rho$
0.04 +0.03 -0.04		BOECKMANN	70	HBC	0	5.0 $\pi^+ \rho$

$\Gamma(K\bar{K})/[\Gamma(\rho\pi) + \Gamma(\eta\pi) + \Gamma(K\bar{K})]$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_4/(\Gamma_1 + \Gamma_2 + \Gamma_4)$
---	------	-------------	------	-----	---------	---

0.054 ± 0.009 OUR FIT						
0.048 ± 0.012 OUR AVERAGE						
0.05 ± 0.02		TOET	73	HBC	+	5 $\pi^+ \rho$
0.09 ± 0.04		TOET	73	HBC	0	5 $\pi^+ \rho$
0.03 ± 0.02	8	DAMERI	72	HBC	-	11 $\pi^- \rho$
0.06 ± 0.03	17	BARNHAM	71	HBC	+	3.7 $\pi^+ \rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.020 ± 0.004	18	ESPIGAT	72	HBC	±	0.0 $\bar{p} \rho$

¹⁸Not averaged because of discrepancy between masses from $K\bar{K}$ and $\rho\pi$ modes.

$\Gamma(\pi^+ \pi^- \pi^-)/\Gamma(\rho\pi)$	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_8/Γ_1
---	-----	-------------	------	-----	---------	---------------------

<0.12	90	ABRAMOVI...	70B	HBC	-	3.93 $\pi^- \rho$
-------	----	-------------	-----	-----	---	-------------------

$\Gamma(\pi^\pm \gamma)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT
---	-------------	------	---------

0.005 ± 0.005 -0.003	19	EISENBERG	72	HBC	4.3, 5.25, 7.5 $\gamma \rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
¹⁹ Pion-exchange model used in this estimation.					

$\Gamma(\omega\pi\pi)/\Gamma(\rho\pi)$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_1
--	------	-------------	------	-----	---------	---------------------


0.15 ± 0.05 OUR FIT						
0.15 ± 0.05 OUR AVERAGE						
0.28 ± 0.09	60	DIAZ	74	DBC	0	6 $\pi^+ n$
0.18 ± 0.08		20 KARSHON	74	HBC		Avg. of above two
0.10 ± 0.05	279	CHALOUKPA	73	HBC	-	3.9 $\pi^- \rho$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.29 ± 0.08	140	20 KARSHON	74	HBC	0	4.9 $\pi^+ \rho$
0.10 ± 0.04	60	20 KARSHON	74	HBC	+	4.9 $\pi^+ \rho$
0.19 ± 0.08		DEFOIX	73	HBC	0	0.7 $\bar{p} \rho$

²⁰KARSHON 74 suggest an additional $I = 0$ state strongly coupled to $\omega\pi\pi$ which could explain discrepancies in branching ratios and masses. We use a central value and a systematic spread.

WEIGHTED AVERAGE
0.15 ± 0.05 (Error scaled by 1.3)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\eta'(958)\pi)/\Gamma(\eta\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
---	-------------	------	---------	---------------------

0.040 ± 0.007 OUR AVERAGE					
0.047 ± 0.010 ± 0.004	21	BELADIDZE	93	VES	37 $\pi^- N \rightarrow a_2^- N$
0.034 ± 0.008 ± 0.005		BELADIDZE	92	VES	36 $\pi^- C \rightarrow a_2^- C$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
²¹ Using $B(\eta' \rightarrow \pi^+ \pi^- \eta) = 0.441$, $B(\eta \rightarrow \gamma\gamma) = 0.389$ and $B(\eta \rightarrow \pi^+ \pi^- \pi^0) = 0.236$.					

$a_2(1320)$ REFERENCES

Aoyagi 93	PL B314 246	+Fukui, Hasegawa+	(BKEI Collab.)
Armstrong 93C	PL B307 394	+Bettioni+	(FNAL, FERR, GENO, UCI, NWES+)
Beladidze 93	PL 313 276	+Berdnikov, Bitukov+	(VES Collab.)
Condo 93	PR D48 3045	+Handler, Bugg+	(SLAC Hybrid Collab.)
Alde 92B	ZPHY C54 549	+Binon+	(SERP, BELG, LANL, LAPP, PISA, KEK)
Beladidze 92	ZPHY C54 235	+Bitukov, Borisov+	(VES Collab.)
Albrecht 90G	ZPHY C48 183	+Ehrlichmann, Harder+	(ARGUS Collab.)
Armstrong 90	ZPHY C48 213	+Benayoun, Beusch	(WA76 Collab.)
Baru 90	ZPHY C48 581	+Binov, Binov+	(MD-1 Collab.)
Behrend 90C	ZPHY C46 583	+Criege+	(CELLO Collab.)
Butler 90	PR D42 1368	+Boyer+	(Mark II Collab.)
Oest 90	ZPHY C47 343	+Olsson+	(JADE Collab.)
Augustin 89	NP B320 1	+Cosme	(DM2 Collab.)
Vorobyev 88	SJNP 48 273	+Golubev, Dolinsky, Druzhinin+	(NOVO)
Althoff 86	Translated from YAF 48 436		
Antreasyan 86	ZPHY C51 537	+Boch, Foster, Bernard+	(TASSO Collab.)
Berger 84C	PL D33 1847	+Aschman, Besset, Bielein+	(Crystal Ball Collab.)
Behrend 83B	PL 149B 427	+Kloving, Burger+	(PLUTO Collab.)
Frazer 83	PL 125B 518	+Kloving, Burger+	(CELLO Collab.)
Changir 82	PL 117B 123	+D'Agostini+	(UCSD)
Cleland 82B	NP B208 228	+Berg, Biel, Chandee+	(FNAL, MINN, ROCH)
Edwards 82F	PL 135B 72	+Defosse, Dorsaz, Gloor	(DURH, GEVA, LAUS, PITT)
Delosse 81	NP B183 349	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
Evangelista 81	NP B178 197	+Guisan, Martin, Muhlemann, Weill+	(GEVA, LAUS)
Chabaud 80	NP B175 189	+ (BARI, BONN, CERN, DARE, LIPP+)	
Daum 80C	PL 89B 276	+Hyams, Papadopolou+	(CERN, MPIM, AMST)
Baltay 78B	PR D17 62	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)
Chabaud 78	NP B145 349	+Cautis, Cohen, Csorna+	(COLU, BING)
Ferrersoria 78	PL 74B 287	+Hyams, Jones, Weillhammer, Blum+	(CERN, MPIM)
Hyams 78	NP B146 303	+Treille+	(ORSAY, CERN, CDF, EPIL)
Martin 78D	PL 74B 417	+Jones, Weillhammer, Blum+	(CERN, MPIM, ATEN)
May 77	PR D16 1983	+Ozmutlu, Baldi, Bohringer, Dorsaz+	(DURH, GEVA) JP
Forino 76	NC 35A 465	+Abramson, Andrews, Busnello+	(ROCH, CORN)
Margulie 76	PR D14 667	+Gessaroli+	(BGNA, FIRZ, GENO, MILA, OXF, PAVI)
EMMS 75	PL 58B 117	+Kramer, Foley, Love, Lindenbaum+	(BNL, CUNY)
Wagner 75	PL 58B 201	+Jones, Kinson, Stacey, Bell+	(BIRM, DURH, RHUL) JP
Diaz 74	PRL 32 260	+Tabak, Chew	(LBL) JP
Karshon 74	PRL 32 852	+Defosse, Fickinger, Anderson+	(CASE, CMU)
Antipov 73	NP B63 175	+Milkenberg, Pitluk, Eisenberg, Ronat+	(REHO)
Antipov 73C	NP B63 153	+Ascoli, Busnello, Focacci+	(CERN, SERP) JP
Chaloupka 73	PL 44B 211	+Ascoli, Busnello, Focacci+	(CERN, SERP) JP
Conforto 73	PL 45B 154	+Dobrzynski, Ferrando, Losty+	(CERN)
Defoix 73	PL 43B 141	+Moble, Key+	(EFI, FNAL, TINTO, WISC)
Eisenstein 73	PRL 30 503	+Dobrzynski, Espigat, Nascimento+	(CDF)
TOET 73	NP B63 248	+Schultz, Ascoli, Ioffredo+	(CDF)
Dameri 72	NC 9A 1	+Conforto, Mobley+	(TINTO, EFI, FNAL, WISC)
Eisenberg 72	PR D5 15	+Thuan, Major+	(NIJ, BONN, DURH, TORI)
Esigat 72	NP B36 93	+Borzatta, Goussu+	(GAMO, MILA, SACL)
Foley 72	PR D6 747	+Ballam, Dagan+	(REHO, SLAC, TELA)
Alston... 71	PL 34B 156	+Ghesquier, Lilestoll, Montanet	(CERN, CDF)
Barnham 71	PRL 26 1494	+Love, Ozaki, Platner, Lindenbaum+	(BNL, CUNY)
Binnie 71	PL 36B 257	+Major+, Ascoli, Barbano, Buih, Derenzo+	(LRL)
Bowen 71	PRL 26 1663	+Abrams, Butler, Coyne, Goldhaber, Hall+	(LBL)
Grayer 71	PL 34B 333	+Camilleri, Duane, Farugi, Burton+	(LOIC, SHMP)
Abramovi... 70B	NP B23 466	+Earles, Faisler, Blieden+	(NEAS, STON)
Alston... 70	PL 35B 607	+Hyams, Jones, Schlein, Blum+	(CERN, MPIM)
Boeckmann 70	NP B16 221	+Abramovich, Blumenfeld, Bruyant+	(CERN) JP
Ascoli 68	PRL 20 1321	+Alston-Garnjost, Barbano, Buih, Derenzo+	(LRL)
Boesebeck 68	NP B4 501	+Major+, Ascoli, Barbano, Buih, Derenzo+	(BONN, DURH, NIJ, EPOL, TORI)
Chung 68	PR 165 1491	+Crawley, Mortara, Shapiro, Bridges+	(ILL) JP
Conte 67	NC 51A 175	+Deutschmann+	(AACH, BERL, CERN)
		+Dahl, Kirz, Miller	(LRL)
		+Tomasini, Cords+	(GENO, HAMB, MILA, SACL)

See key on page 1343

Meson Full Listings

$a_2(1320)$, $f_0(1370)$, $h_1(1380)$

OTHER RELATED PAPERS

JENNI	83	PR D27 1031	+Burke, Telnov, Abrams, Blocker+	(SLAC, LBL)
BEHREND	82C	PL 114B 378	+Chen, Fenner, Field+	(CELLO Collab.)
ABOLINS	65	Athens Conf.	+Carmony, Lander, Xuong, Yager	(UCSD) I
ADERHOLZ	65	PR 138B 897	(AACH3, BERL, BIRM, BONN, HAMB, LOIC, MPIM)	
ALITTI	65	PL 15 69	+Baton, Deler, Crussard+	(SACL, BGNA) JP
CHUNG	65	PRL 15 325	+Dahl, Hardy, Hess, Jacobs, Kirz	(LRL)
FORINO	65B	PL 19 68	+Gessaroli+	(BGNA, BARI, FIRZ, ORSAY, SACL)
LEFEBVRES	65	PL 19 434	+Levrat+	(CERN Missing Mass Spect. Collab.)
SEIDLITZ	65	PRL 15 217	+Dahl, Miller	(LRL)
ADERHOLZ	64	PL 10 226	(AACH3, BERL, BIRM, BONN, DESY, HAMB+)	
CHUNG	64	PRL 12 621	+Dahl, Hardy, Hess, Kalbfleisch, Kirz	(LRL)
GOLDHABER	64B	Dubna Conf. 1 480	+Goldhaber, O'Halloran, Shen	(LRL)
Also	64	PRL 12 336	Goldhaber, Brown, Kadyk, Shen+	(LRL, UCB)
LANDER	64	PRL 13 346A	+Abolins, Carmony, Hendricks, Xuong+	(UCSD)

 $f_0(1370)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

We list here all claims compatible with high $\pi\pi$ S-wave inelasticity.
See also minireviews under $f_0(1300)$ and non- $q\bar{q}$ candidates.

 $f_0(1370)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1365^{+20}_{-55}	1 ANISOVICH	94 CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1374 ± 38	AMSLER	94 CBAR	$0.0 \bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
1345 ± 12	ADAMO	93 OBLX	$\bar{p}p \rightarrow 3\pi^+2\pi^-$
1430 ± 25	AMSLER	93C CBAR	$0.0 \bar{p}p \rightarrow \pi^0\eta\eta$
1386 ± 30	GASPERO	93 DBC	$0.0 \bar{p}n \rightarrow 2\pi^+3\pi^-$
1440 ± 20	CHEN	91 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-, \gamma K\bar{K}$
1440 ± 50	BOLONKIN	88 SPEC	$40 \pi^-\pi^+ \rightarrow K_S^0 K_S^0 n$
1463 ± 9	ETKIN	82B MPS	$23 \pi^-\pi^+ \rightarrow n2K_S^0$

¹ From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$. Supersedes AMSLER 93C.

 $f_0(1370)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
268 ± 70	2 ANISOVICH	94 CBAR	$0.0 \bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
375 ± 61	AMSLER	94 CBAR	$0.0 \bar{p}p \rightarrow \pi^+\pi^-3\pi^0$
398 ± 26	ADAMO	93 OBLX	$\bar{p}p \rightarrow 3\pi^+2\pi^-$
250 ± 50	AMSLER	93C CBAR	$0.0 \bar{p}p \rightarrow \pi^0\eta\eta$
310 ± 50	GASPERO	93 DBC	$0.0 \bar{p}n \rightarrow 2\pi^+3\pi^-$
160 ± 40	CHEN	91 MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-, \gamma K\bar{K}$
250 ± 80	BOLONKIN	88 SPEC	$40 \pi^-\pi^+ \rightarrow K_S^0 K_S^0 n$
118^{+138}_{-16}	ETKIN	82B MPS	$23 \pi^-\pi^+ \rightarrow n2K_S^0$

² From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$. Supersedes AMSLER 93C.

 $f_0(1370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $\pi\pi$	< 12 %	
Γ_2 4π	> 72 %	95%
Γ_3 $4\pi^0$		
Γ_4 $2\pi^+2\pi^-$		
Γ_5 $\pi^+\pi^-2\pi^0$		
Γ_6 6π		
Γ_7 $\eta\eta$	seen	
Γ_8 $\eta\eta'$		
Γ_9 2ω		
Γ_{10} $K\bar{K}X$		

 $f_0(1370)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
< 0.12	3 GASPERO	93 RVUE	$0.0 \bar{p}n \rightarrow \text{hadrons}$	

³ Based on GASPERO 93 and private communications from M. Gaspero.

$\Gamma(4\pi)/\Gamma_{\text{total}}$	CL %	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/\Gamma = (\Gamma_3 + \Gamma_4 + \Gamma_5)/\Gamma$
VALUE					
> 0.72	95	4 GASPERO	93 RVUE	$0.0 \bar{p}n \rightarrow \text{hadrons}$	

⁴ Based on model-dependent evaluation by GASPERO 93 and private communications from M. Gaspero.

$\Gamma(2\pi^+2\pi^-)/\Gamma(4\pi)$	DOCUMENT ID	TECN	COMMENT	$\Gamma_4/\Gamma_2 = \Gamma_4/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
VALUE				
0.420 ± 0.014	5 GASPERO	93 RVUE	$0.0 \bar{p}n \rightarrow 2\pi^+3\pi^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁵ GASPERO 93 RVUE $0.0 \bar{p}n \rightarrow 2\pi^+3\pi^-$

⁶ Based on model-dependent evaluation by GASPERO 93.

$\Gamma(\pi^+\pi^-2\pi^0)/\Gamma(4\pi)$	DOCUMENT ID	TECN	COMMENT	$\Gamma_5/\Gamma_2 = \Gamma_5/(\Gamma_3 + \Gamma_4 + \Gamma_5)$
VALUE				
0.512 ± 0.019	6 GASPERO	93 RVUE	$0.0 \bar{p}n \rightarrow \text{hadrons}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁶ GASPERO 93 RVUE $0.0 \bar{p}n \rightarrow \text{hadrons}$

⁶ Based on model-dependent evaluation by GASPERO 93.

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
VALUE				
seen	AMSLER	92 CBAR	$0.0 \bar{p}p \rightarrow \eta\eta\pi^0$	

 $f_0(1370)$ REFERENCES

AMSLER	94	PL B322 431	+Armstrong+	(Crystal Barrel Collab.) JPC
ANISOVICH	94	PL B323 233	+Armstrong+	(Crystal Barrel Collab.) JPC
ADAMO	93	NP A558 13C	+Agnello+	(OBELIX Collab.) JPC
AMSLER	93C	NP A558 3C	+Augustin+	(Crystal Barrel Collab.) JPC
GASPERO	93	NP A562 407		(ROMAJ) JPC
AMSLER	92	PL B291 347	+Augustin, Baker+	(Crystal Barrel Collab.)
CHEN	91	Hadron 91 Conf.		(Mark III Collab.)
SLAC-PUB-5669				
BOLONKIN	88	NP B309 426	+Bloschenko, Gorin+	(ITEP, SERP)
ETKIN	82B	PR D25 1786	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)

OTHER RELATED PAPERS

BRIDGES	86	PRL 56 211	+Brown+	(BLSU, BNL, CASE, COLU, UMD, SYRA)
BRIDGES	86B	PRL 56 215	+Daftari, Kalogeropoulos, Debbe+	(SYRA, CASE)
BRIDGES	86C	PRL 57 1534	+Daftari, Kalogeropoulos+	(SYRA)
BETTINI	66	NC 42A 695	+Cresti, Limentani, Bertanza, Bigi+	(PADO, PISA)

 $h_1(1380)$

$$I^G(J^{PC}) = ?^-(1^{+?})$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K_S^0 K^\pm \pi^\mp$ system. Evidence for $K^* \bar{K} + \bar{K}^* K$ decays (ASTON 88C). Needs confirmation.

 $h_1(1380)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1380 ± 20	ASTON	88C LASS	$11 K^-\pi^+ \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

 $h_1(1380)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 ± 30	ASTON	88C LASS	$11 K^-\pi^+ \rightarrow K_S^0 K^\pm \pi^\mp \Lambda$

 $h_1(1380)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\bar{K}^*(892) + c.c.$		

 $h_1(1380)$ REFERENCES

ASTON	88C	PL B201 573	+Awaji, Bienz+	(SLAC, NAGO, CINC, INUS)
-------	-----	-------------	----------------	--------------------------

Meson Full Listings

 $\tilde{\rho}(1405), f_1(1420)$ $\tilde{\rho}(1405)$

$$I^G(J^{PC}) = 1^-(1^-+)$$

OMITTED FROM SUMMARY TABLE

Seen by ALDE 88B in $\pi^- \rho \rightarrow \eta \pi^0 n$ amplitude analysis. Needs confirmation.See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.) $\tilde{\rho}(1405)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1406 ± 20	¹ ALDE	88B	GAM4	0 100 $\pi^- \rho \rightarrow \eta \pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

1323.1 ± 4.6 AOYAGI 93 BKEI $\pi^- \rho \rightarrow \eta \pi^- \rho$ ¹ Seen in the P_0 -wave intensity of the $\eta \pi^0$ system. $\tilde{\rho}(1405)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
180 ± 20	² ALDE	88B	GAM4	0 100 $\pi^- \rho \rightarrow \eta \pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

143.2 ± 12.5 AOYAGI 93 BKEI $\pi^- \rho \rightarrow \eta \pi^- \rho$ ² Seen in the P_0 -wave intensity of the $\eta \pi^0$ system. $\tilde{\rho}(1405)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta \pi^0$	seen
Γ_2 $\eta \pi^-$	
Γ_3 $\rho \pi$	not seen
Γ_4 $\eta' \pi$	

 $\tilde{\rho}(1405)$ BRANCHING RATIOS

$\Gamma(\eta \pi^0)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	³ ALDE	88B	GAM4	0 100 $\pi^- \rho \rightarrow \eta \pi^0 n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

not seen ⁴ APEL 81 NICE 0 40 $\pi^- \rho \rightarrow \eta \pi^0 n$ ³ Seen in the P_0 -wave intensity of the $\eta \pi^0$ system.⁴ A general fit allowing S , D , and P waves (including $m=0$) is not done because of limited statistics.

$\Gamma(\eta \pi^-)/\Gamma_{\text{total}}$	Γ_2/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
possibly seen	AOYAGI	93	BKEI $\pi^- \rho \rightarrow \eta \pi^- \rho$

$\Gamma(\rho \pi)/\Gamma_{\text{total}}$	Γ_3/Γ	
VALUE	DOCUMENT ID	COMMENT
not seen	⁵ ZIELINSKI 86	200 $\pi^+ \text{Cu,Pb} \rightarrow \pi^+ \pi^+ \pi^- X$

⁵ A general fit allowing S , D , and P waves (including $m=0$) is not done because of limited statistics.

$\Gamma(\eta' \pi)/\Gamma(\eta \pi^0)$	Γ_4/Γ_1			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.80	95	BOUTEMEUR 90	GAM4	100 $\pi^- \rho \rightarrow 4\gamma n$

 $\tilde{\rho}(1405)$ REFERENCES

AOYAGI 93	PL B314 246	+Fukui, Hasegawa+	(BKEI Collab.)
BOUTEMEUR 90	Hadron 89 Conf. p 119+Poulet	(SERP, BELG, LANL, LAPP, PISA, KEK)	
ALDE 88B	PL B205 397	+Binon, BoutemEUR+	(SERP, BELG, LANL, LAPP) IGJPC
ZIELINSKI 86	Berkeley HEP 1 736	+Berg+	(ROCH, MINN, FNAL)
APEL 81	NP B193 269	+Augenstein, Bertolucci, Donskov+	(SERP, CERN)

OTHER RELATED PAPERS

IDDIR 88	PL B205 564	+Le Yaouanc, Ono+	(ORSAY, TOKY)
TUAN 88	PL B213 537	+Ferber, Dalitz	(HAWA, ROCH, OXFPT)
ZIELINSKI 87	ZPHY C34 255		(ROCH)
ZIELINSKI 86	Berkeley HEP 1 736	+Berg+	(ROCH, MINN, FNAL)

 $f_1(1420)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See also minireview under non- $q\bar{q}$ candidates.NOTE ON THE $f_1(1420)$ This particle is the axial-vector component of the old puzzling E/ι , which has caused much trouble.

In hadron-induced reactions, the $f_1(1420)$ is observed in centrally produced $K\bar{K}\pi$ systems obtained with π and p beams (DIONISI 80, ARMSTRONG 84, 89). A Dalitz-plot analysis gives its quantum numbers and the dominant decay mode. For instance, ARMSTRONG 89 finds that the signal is totally consistent with being an 1^{++} state with a dominant quasi-two-body S -wave decay into $K^*(892)\bar{K}$; furthermore, no 0^{-+} or 1^{+-} waves are required to fit the data. A G parity of $+1$ is suggested by the positive interference between the two overlapping $K^*(892)$ (ARMSTRONG 84). No significant signals in the $\eta\pi\pi$ or 4π decay modes are found in centrally produced 4π systems (ARMSTRONG 89G). All of this is in line with the previous observations made in $\bar{p}p$ annihilations.

In $\gamma\gamma$ fusion from e^+e^- annihilations, a signal at about 1420 MeV is seen only in single-tag events (AIHARA 86C, GIDAL 87B, BEHREND 89, HILL 89), where one of the two photons is off the mass shell; by contrast, it is totally absent in the untagged events where both photons are real and hence they cannot produce a spin-1 meson, because of the Yang-Landau theorem. This clearly implies $J = 1$ and $C = +1$. As for the parity, AIHARA 88B, 88C (same analysis as AIHARA 86C, with 25% more events) and BEHREND 89 all find angular distributions with positive parity preferred, but negative parity not excluded.

Although some uncertainties still remain, the state seen in hadronic interactions and that seen in spacelike virtual photon fusion from e^+e^- annihilations are often identified with one another since there are more similarities than differences. In particular, all experiments agree that this state appears only in $K^*(892)\bar{K}$. The same conclusions are obtained from partial wave analyses of $J/\psi(1S) \rightarrow \gamma K\bar{K}\pi$ (BAI 90C, AUGUSTIN 91).

BITYUKOV 88 studied the radiative decay $1^{++} \rightarrow \phi\gamma$. Since the ϕ is (almost) a pure $s\bar{s}$ state, the $\phi\gamma$ decay seems to be a good analyser to extract the $s\bar{s}$ component in the wave function of the decaying meson. Finding the $f_1(1285)$ but not the $f_1(1420)$, BITYUKOV 88 concludes that the $f_1(1420)$ cannot be the $s\bar{s}$ isoscalar member of the $q\bar{q}$ nonet containing the $f_1(1285)$. On the other hand, AIHARA 88C argues that, assuming they both belong to the same nonet and using several hypotheses, the octet-singlet mixing angle obtained is compatible with the $f_1(1420)$ being mostly $s\bar{s}$ and the $f_1(1285)$ being mostly $(u\bar{u} + d\bar{d})/\sqrt{2}$, although both require large admixtures of other $q\bar{q}$ components.

Arguments favoring the possibility the $f_1(1420)$ is a hybrid $q\bar{q}g$ meson or a four-quark state are put forward by ISHIDA 89 and by CALDWELL 90, respectively.

See key on page 1343

Meson Full Listings
 $f_1(1420)$

LONGACRE 90 argues that this particle is inconsistent with a QCD arrangement of quarks and gluons. He then develops a final-state rescattering mechanism with successive interactions between a K , a \bar{K} , and a π . The $f_1(1420)$ would then be a molecular state formed by the π orbiting in a P wave around an S -wave $K\bar{K}$ state.

 $f_1(1420)$ MASSPRODUCED IN $p\bar{p}$ ANNIHILATION


VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1414.9 ± 3.5 OUR AVERAGE		Error includes scale factor of 1.2.		
1417.5 ± 4		NACASCH	78 HBC	0.7,0.76 $\bar{p}p$
1398 ± 10	170	DEFOIX	72 HBC	0.7 $\bar{p}p \rightarrow 7\pi$
1406 ± 7	280	DUBOC	72 HBC	1.2 $\bar{p}p \rightarrow 2K4\pi$
1420 ± 7	310	LORSTAD	69 HBC	0.7 $\bar{p}p$
1423.0 ± 10.0		FRENCH	67 HBC	3-4 $\bar{p}p$

PRODUCED IN OTHER REACTIONS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1426.8 ± 2.3 OUR AVERAGE		Error includes scale factor of 1.3. See the ideogram below.		
1430 ± 4		1 ARMSTRONG	92E OMEG	85,300 $\pi^+p, pp \rightarrow \pi^+p, pp(K\bar{K}\pi)$
1462 ± 20		2 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
1443 $\begin{smallmatrix} +7 \\ -6 \end{smallmatrix}$ $\begin{smallmatrix} +3 \\ -2 \end{smallmatrix}$	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1425 ± 10	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
1442 ± 5.0 $\begin{smallmatrix} +10.0 \\ -17.0 \end{smallmatrix}$	111 $\begin{smallmatrix} +31 \\ -26 \end{smallmatrix}$	BECKER	87 MRK3	$e^+e^- \rightarrow \omega K\bar{K}\pi$
1423 ± 4		GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
1417.0 ± 13.0	13	AIHARA	86C TPC	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
1422.0 ± 3.0		CHAUVAT	84 SPEC	ISR 31.5 pp
1440.0 ± 10.0		3 BROMBERG	80 SPEC	100 $\pi^-p \rightarrow K\bar{K}\pi X$
1426.0 ± 6.0	221	DIONISI	80 HBC	4 $\pi^-p \rightarrow K\bar{K}\pi n$
1420 ± 20		DAHL	67 HBC	1.6-4.2 π^-p
1429 ± 3	389 ± 27	ARMSTRONG	89 OMEG	300 $pp \rightarrow K\bar{K}\pi pp$
1425.0 ± 2.0	1520	ARMSTRONG	84 OMEG	85 $\pi^+p, pp \rightarrow (\pi^+, p)(K\bar{K}\pi)p$

- 1 This result supersedes ARMSTRONG 84, ARMSTRONG 89.
 2 From fit to the $K^*(892)K 1^{++}$ partial wave.
 3 Mass error increased to account for $a_0(980)$ mass cut uncertainties.

WEIGHTED AVERAGE
 1426.8 ± 2.3 (Error scaled by 1.3)

 $f_1(1420)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
52 ± 4 OUR AVERAGE				
58 ± 10		4 ARMSTRONG	92E OMEG	85,300 $\pi^+p, pp \rightarrow \pi^+p, pp(K\bar{K}\pi)$
129 ± 41		5 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
68 $\begin{smallmatrix} +29 \\ -18 \end{smallmatrix}$ $\begin{smallmatrix} +8 \\ -9 \end{smallmatrix}$	1100	BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
42 ± 22	17	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$
40 $\begin{smallmatrix} +17 \\ -13 \end{smallmatrix}$ ± 5	111 $\begin{smallmatrix} +31 \\ -26 \end{smallmatrix}$	BECKER	87 MRK3	$e^+e^- \rightarrow \omega K\bar{K}\pi$
35.0 $\begin{smallmatrix} +47.0 \\ -20.0 \end{smallmatrix}$	13	AIHARA	86C TPC	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
47.0 ± 10.0		CHAUVAT	84 SPEC	ISR 31.5 pp
62.0 ± 14.0		BROMBERG	80 SPEC	100 $\pi^-p \rightarrow K\bar{K}\pi X$

40.0 ± 15.0	221	DIONISI	80 HBC	4 $\pi^-p \rightarrow K\bar{K}\pi n$
53 ± 20.0		NACASCH	78 HBC	0.7,0.76 $\bar{p}p$
50 ± 10	170	DEFOIX	72 HBC	0.7 $\bar{p}p \rightarrow 7\pi$
50 ± 12	280	DUBOC	72 HBC	1.2 $\bar{p}p \rightarrow 2K4\pi$
60 ± 20	310	LORSTAD	69 HBC	0.7 $\bar{p}p$
60.0 ± 20.0		DAHL	67 HBC	1.6-4.2 π^-p
45 ± 20		FRENCH	67 HBC	3-4 $\bar{p}p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
58 ± 8	389 ± 27	ARMSTRONG	89 OMEG	300 $pp \rightarrow K\bar{K}\pi pp$
62.0 ± 5.0	1520	ARMSTRONG	84 OMEG	85 $\pi^+p, pp \rightarrow (\pi^+, p)(K\bar{K}\pi)p$

- 4 This result supersedes ARMSTRONG 84, ARMSTRONG 89.
 5 From fit to the $K^*(892)K 1^{++}$ partial wave.

 $f_1(1420)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}\pi$	dominant
Γ_2 $\eta\pi\pi$	possibly seen
Γ_3 $a_0(980)\pi$	
Γ_4 $\pi\pi\rho$	
Γ_5 $K\bar{K}^*(892) + c.c.$	
Γ_6 4π	
Γ_7 $\gamma\gamma^*$	
Γ_8 $\rho^0\gamma$	

 $f_1(1420)$ $\Gamma(I)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma^*)/\Gamma(\text{total})$	$\Gamma_1\Gamma_7/\Gamma$			
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.4 OUR AVERAGE				
3.0 ± 0.9 ± 0.7		6,7 BEHREND	89 CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K\pi$
2.3 $\begin{smallmatrix} +1.0 \\ -0.9 \end{smallmatrix}$ ± 0.8		HILL	89 JADE	$e^+e^- \rightarrow e^+e^- K^\pm K_S^0 \pi^\mp$
1.3 ± 0.5 ± 0.3		AIHARA	88B TPC	$e^+e^- \rightarrow e^+e^- K^\pm K_S^0 \pi^\mp$
1.6 ± 0.7 ± 0.3		6,8 GIDAL	87B MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 8.0	95	JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
6 Assume a ρ -pole form factor. 7 A ϕ -pole form factor gives considerably smaller widths. 8 Published value divided by 2.				

 $f_1(1420)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892) + c.c.)/\Gamma(K\bar{K}\pi)$	Γ_5/Γ_1			
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.76 ± 0.06	BROMBERG	80 SPEC	100 $\pi^-p \rightarrow K\bar{K}\pi X$	
0.86 ± 0.12	DIONISI	80 HBC	4 $\pi^-p \rightarrow K\bar{K}\pi n$	
$\Gamma(\pi\pi\rho)/\Gamma(K\bar{K}\pi)$	Γ_4/Γ_1			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.3	95	CORDEN	78 OMEG	12-15 π^-p
< 2.0		DAHL	67 HBC	1.6-4.2 π^-p
$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$	Γ_2/Γ_1			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.1	95	ARMSTRONG	91B OMEG	300 $pp \rightarrow pp\eta\pi^+\pi^-$
1.35 ± 0.75		KOPKE	89 MRK3	$J/\psi \rightarrow \omega\eta\pi(K\bar{K}\pi)$
< 0.6	90	GIDAL	87 MRK2	$e^+e^- \rightarrow e^+e^- \eta\pi^+\pi^-$
< 0.5	95	CORDEN	78 OMEG	12-15 π^-p
1.5 ± 0.8		DEFOIX	72 HBC	0.7 $\bar{p}p$
$\Gamma(a_0(980)\pi)/\Gamma(\eta\pi\pi)$	Γ_3/Γ_2			
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen in either mode	ANDO	86 SPEC	8 π^-p	
not seen in either mode	CORDEN	78 OMEG	12-15 π^-p	
0.4 ± 0.2	DEFOIX	72 HBC	0.7 $\bar{p}p \rightarrow 7\pi$	
$\Gamma(4\pi)/\Gamma(K\bar{K}^*(892) + c.c.)$	Γ_6/Γ_5			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.90	95	DIONISI	80 HBC	4 π^-p

Meson Full Listings

 $f_1(1420), \omega(1420), f_2(1430)$ $\Gamma(K\bar{K}\pi)/[\Gamma(a_0(980)\pi) + \Gamma(K\bar{K}^*(892) + c.c.)]$ $\Gamma_1/(\Gamma_3 + \Gamma_5)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.65 ± 0.27	⁹ DIONISI 80 HBC		4 $\pi^- \rho$

⁹ Calculated using $\Gamma(K\bar{K})/\Gamma(\eta\pi) = 0.24 \pm 0.07$ for $a_0(980)$ fractions.

 $\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}^*(892) + c.c.)$ Γ_3/Γ_5

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.04	68	ARMSTRONG 84 OMEG		85 $\pi^+ \rho$

 $\Gamma(4\pi)/\Gamma(K\bar{K}\pi)$ Γ_6/Γ_1

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.62	95	ARMSTRONG 89G OMEG		85 $\pi \rho \rightarrow 4\pi X$

 $\Gamma(\rho^0\gamma)/\Gamma_{total}$ Γ_8/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.08	95	¹⁰ ARMSTRONG 92C SPEC		300 $\rho\rho \rightarrow \rho\rho\pi^+\pi^-\gamma$

¹⁰ Using the data on the $\bar{K}K\pi$ mode from ARMSTRONG 89.

 $f_1(1420)$ REFERENCES

ARMSTRONG 92C	ZPHY C54 371	+Barnes, Benayoun+ (ATHU, BARI, BIRM, CERN, CDEF)
ARMSTRONG 92E	ZPHY 56 29	+Benayoun+ (ATHU, BARI, BIRM, CERN, CDEF) JPC
AUGUSTIN 92	PR D46 1951	+Cosme (DM2 Collab.)
ARMSTRONG 91B	ZPHY C52 389	+Barnes+ (ATHU, BARI, BIRM, CERN, CDEF)
BAI 90C	PRL 65 2507	+Blaylock+ (Mark III Collab.)
ARMSTRONG 89	PL B221 216	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+) JPC
ARMSTRONG 89G	ZPHY C43 55	+Bloodworth+ (CERN, BIRM, BARI, ATHU, CURIN+)
BEHREND 89	ZPHY C42 367	+Criegee+ (CELLO Collab.)
HILL 89	ZPHY C42 355	+Olsson+ (JADE Collab.) JP
KOPKE 89	PRPL 174 67	+Wernes+ (CERN)
AIHARA 88B	PL B209 107	+Alston-Garnjost+ (TPC-2 γ Collab.)
BECKER 87	PRL 59 186	+Blaylock, Bolton, Brown+ (Mark III Collab.) JP
GIDAL 87	PRL 59 2012	+Boyer, Butler, Cords, Abrams+ (LBL, SLAC, HARV)
GIDAL 87B	PRL 59 2016	+Boyer, Butler, Cords, Abrams+ (LBL, SLAC, HARV)
AIHARA 86C	PRL 57 2500	+Alston-Garnjost+ (TPC-2 γ Collab.) JP
ANDO 86	PRL 57 1296	+Imai+ (KEK, KYOT, NIRS, SAGA, INUS, TSUK+)
ARMSTRONG 84	PL 146B 273	+Bloodworth, Burns+ (ATHU, BARI, BIRM, CERN) JP
CHAUVAT 84	PL 148B 382	+Meritet, Bonino+ (CERN, CLER, UCLA, SACL)
JENNI 83	PR D27 1031	+Burke, Telnov, Abrams, Blocker+ (SLAC, LBL)
BROMBERG 80	PR D22 1513	+Haggerty, Abrams, Dzierba (CIT, FNAL, ILLC, IND)
DIONISI 80	NP B169 1	+Gavillet+ (CERN, MADR, CDEF, STOHI) IJP
CORDEN 78	NP B144 253	+Corbett, Alexander+ (BIRM, RHEL, TELA, LOWC)
NACASCH 78	NP B135 203	+Defoix, Dobrzynski+ (PARIS, MADR, CERN)
DEFOIX 72	NP B44 125	+Nascimento, Bizzarri+ (CDEF, CERN)
DUBOC 72	NP B46 429	+Goldberg, Makowski, Donald+ (PARIS, LIVP)
LORSTAD 69	NP B14 63	+D'Andlau, Astier+ (CDEF, CERN) JP
DAHL 67	PR 163 1377	+Hardy, Hess, Kirz, Miller (LRL) IJP
Also 65	PRL 14 1074	Miller, Chung, Dahl, Hess, Hardy, Kirz+ (LRL, UCB)
FRENCH 67	NC 52A 438	+Kinson, McDonald, Riddiford+ (CERN, BIRM)

OTHER RELATED PAPERS

CALDWELL 90	Hadron 89 Conf. p 127	(UCSB)
ISHIDA 89	PTP 82 119	+Oda, Sawazaki, Yamada (NIHO)
AIHARA 88C	PR D38 1	+Alston-Garnjost+ (TPC-2 γ) JPC
BITYUKOV 88	PL B203 327	+Borisov, Dorofeev+ (SERP)
PROTOPOP... 87B	Hadron 87 Conf.	Protopopescu, Chung (BNL)

 $\omega(1420)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

See also $\omega(1600)$.

 $\omega(1420)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1419 ± 31	315	¹ ANTONELLI 92 DM2		1.34–2.4 $e^+e^- \rightarrow \rho\pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
1440 ± 70		² DONNACHIE 91 RVUE		
1391 ± 18		DONNACHIE 89 RVUE		$e^+e^- \rightarrow \rho\pi$

¹ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

² Using data published later by ANTONELLI 92.

 $\omega(1420)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
174 ± 59	315	³ ANTONELLI 92 DM2		1.34–2.4 $e^+e^- \rightarrow \rho\pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
240 ± 70		⁴ DONNACHIE 91 RVUE		
224 ± 49		DONNACHIE 89 RVUE		$e^+e^- \rightarrow \rho\pi$

³ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

⁴ Using data published later by ANTONELLI 92.

 $\omega(1420)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	dominant
Γ_2 $\omega\pi\pi$	
Γ_3 e^+e^-	

 $\omega(1420)$ $\Gamma(I)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\rho\pi) \times \Gamma(e^+e^-)/\Gamma_{total}$	$\Gamma_1\Gamma_3/\Gamma$			
VALUE (eV)	EVTs	DOCUMENT ID	TECN	COMMENT
81 ± 31	315	⁵ ANTONELLI 92 DM2		1.34–2.4 $e^+e^- \rightarrow \rho\pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
137 ± 40		DONNACHIE 89 RVUE		$e^+e^- \rightarrow \rho\pi$

⁵ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

 $\Gamma(\omega\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{total}$ $\Gamma_2\Gamma_3/\Gamma$

$\Gamma(\omega\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{total}$	$\Gamma_2\Gamma_3/\Gamma$			
VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<41	68	DONNACHIE 89 RVUE		$e^+e^- \rightarrow \omega 2\pi$

 $\omega(1420)$ REFERENCES

ANTONELLI 92	ZPHY C56 15	+Baldini+ (DM2 Collab.)
DONNACHIE 91	ZPHY C51 689	+Ciegg (MCHS, LANC)
DONNACHIE 89	ZPHY C42 663	+Ciegg (CERN, MCHS)

OTHER RELATED PAPERS

ATKINSON 87	ZPHY C34 157	- (BONN, CERN, GLAS, LANC, MCHS, CURIN)
ATKINSON 84	NP B231 15	- (BONN, CERN, GLAS, LANC, MCHS, CURIN+)
ATKINSON 83B	PL 127B 132	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)

 $f_2(1430)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry lists nearby peaks observed in the D wave of the $K\bar{K}$ and $\pi^+\pi^-$ systems.

 $f_2(1430)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1430 OUR ESTIMATE			
••• We do not use the following data for averages, fits, limits, etc. •••			
1421 ± 5	AUGUSTIN 87 DM2		$J/\psi \rightarrow \gamma\pi^+\pi^-$
1480.0 ± 50.0	AKESSON 86 SPEC		$\rho\rho \rightarrow \rho\rho\pi^+\pi^-$
1436.0 ± 26.0	DAUM 84 CNTR		17–18 $\pi^- \rho^-$
1412.0 ± 3.0	DAUM 84 CNTR		63 $\pi^- \rho^- \rightarrow K_S^0 K_S^0 n$
1439.0 ± 5.0	¹ BEUSCH 67 OSPK		5,7,12 $\pi^- \rho^- \rightarrow K_S^0 K_S^0 n$
– 6.0			

¹ Not seen by WETZEL 76.

 $f_2(1430)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
30 ± 9	AUGUSTIN 87 DM2		$J/\psi \rightarrow \gamma\pi^+\pi^-$
150.0 ± 40.0	AKESSON 86 SPEC		$\rho\rho \rightarrow \rho\rho\pi^+\pi^-$
81.0 ± 56.0	DAUM 84 CNTR		17–18 $\pi^- \rho^-$
– 29.0			
14.0 ± 6.0	DAUM 84 CNTR		63 $\pi^- \rho^- \rightarrow K_S^0 K_S^0 n$
43.0 ± 17.0	² BEUSCH 67 OSPK		5,7,12 $\pi^- \rho^- \rightarrow K_S^0 K_S^0 n$
– 18.0			

² Not seen by WETZEL 76.

 $f_2(1430)$ DECAY MODES

Mode
Γ_1 $K\bar{K}$
Γ_2 $\pi\pi$

 $f_2(1430)$ REFERENCES

AUGUSTIN 87	ZPHY C36 369	+Cosme+ (LALO, CLER, FRAS, PADO)
AKESSON 86	NP B264 154	+Albrow, Almedeh+ (Axial Field Spec. Collab.)
DAUM 84	ZPHY C23 339	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+) JP
WETZEL 76	NP B115 208	+Freudenreich, Beusch+ (ETH, CERN, LOIC)
BEUSCH 67	PL 25B 357	+Fischer, Gobbi, Astbury+ (ETH, CERN)

See key on page 1343

Meson Full Listings
 $\eta(1440)$ $\eta(1440)$
was $\iota(1440)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)NOTE ON THE $\eta(1440)$

The first observation of a meson with $I^G J^{PC} = 0^+ 0^{-+}$ in the 1400-MeV mass region was made with $p\bar{p}$ annihilations at rest (BAILLON 67) in the channel $\eta(1440) \rightarrow K\bar{K}\pi$. It was seen to decay equally into $a_0(980)\pi$ and $\bar{K}^*(892)K$.

The $\eta(1440)$ has since also been seen in other hadronic reactions: In a partial-wave analysis of the $\eta\pi^+\pi^-$ system, confirming the decay $\eta(1440) \rightarrow a_0(980)\pi$ (FUKUI 91C); in a partial-wave analysis of the $K\bar{K}\pi$ system (CHUNG 85, BIRMAN 88); in 6-GeV $p\bar{p}$ annihilations (REEVES 86); and in nonperipherally selected $\pi^-p \rightarrow K_S^0 K_S^0 \pi^0 n$ (RATH 89). RATH 89 favors the interpretation that there are two narrow η resonances in the 1410–1480 MeV region.

Neither the $\eta(1440)$ nor the $f_1(1420)$ are observed in the $s\bar{s}$ -enriched peripheral reaction $K^-p \rightarrow K\bar{K}\pi\Lambda$ at 11 GeV/c (ASTON 87), which speaks against an $s\bar{s}$ interpretation of either state. Moreover, the $\eta(1440)$ is not seen by ARMSTRONG 84, 89 either, who studied $K\bar{K}\pi$ central production in $\pi^+p \rightarrow \pi^+(K\bar{K}\pi)p$ and $pp \rightarrow p(K\bar{K}\pi)p$ at 85 and 300 GeV/c [but the $f_1(1420)$ is seen]. This agrees with earlier results (DIONISI 80, DEFOIX 72, DUBOC 72, LORSTAD 69, etc.).

The $\eta(1440)$ is also seen as a broad enhancement in $J/\psi(1S)$ radiative decay. In the $K\bar{K}\pi$ channel, however, its mass is higher than observed in hadronic interactions, and its width is larger. It has been shown (TOKI 87, BAI 90C) that two resonances (with $M \approx 1420$ MeV and $M \approx 1490$ MeV) give a better description of the data. Moreover, the $\eta\pi^+\pi^-$ channel peaks near 1400 MeV (AUGUSTIN 90, BURCHELL 91). All these results suggest the existence of two overlapping states (favored by RATH 89 in hadronic production), one around 1400 MeV decaying into both $K\bar{K}\pi$ and $\eta\pi\pi$, the other one around 1490 MeV seen only in $K\bar{K}\pi$. Other possible decay modes, in $\pi\pi\gamma$ and 4π , are not sufficiently well established to clarify the situation.

There is considerable confusion on the partial decay modes: The $K\bar{K}\pi$ final state is usually dominated by $\bar{K}^*(892)K$ and/or $a_0(980)\pi$ contributions, but it is impossible to quote any reliable \bar{K}^*K and $a_0\pi$ branching ratios, since the analyses are highly model dependent and the experiments do not agree.

We continue to list under the $\eta(1440)$ all the results on the 0^{-+} system in the 1380–1490 MeV region, but keep in mind that it is likely that there is more than one resonance present in these observations. The masses and widths are given separately according to the various decay modes.

 $\eta(1440)$ MASS

VALUE (MeV) DOCUMENT ID
1420 ± 20 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

 $\eta\pi\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1399 ± 4	OUR AVERAGE			
1400 ± 6		¹ BURCHELL 91	MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
1398 ± 6	261 ± 24	² AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1385 ± 15		¹ BEHREND 92	DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
1388 ± 4		FUKUI 91C	SPEC	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$
1420 ± 5		ANDO 86	SPEC	$8 \pi^- p \rightarrow n\eta\pi^+\pi^-$
¹ From fit to the $a_0(980)\pi 0^{-+}$ partial wave.				
² Best fit with a single Breit Wigner.				

 $\pi\pi\gamma$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1401 ± 18		^{3,4} AUGUSTIN 90	DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma\gamma$
1440 ± 20		⁴ COFFMAN 90	MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$
³ Best fit with a single Breit Wigner.				
⁴ This peak in the $\gamma\rho$ channel may not be related to the $\eta(1440)$.				

 4π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1489 ± 12	3270	⁵ BISELLO 89B	DM2	$J/\psi \rightarrow 4\pi\gamma$
⁵ Estimated by us from various fits.				

 $K\bar{K}\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1419.8 ± 1.0	OUR AVERAGE			Error includes scale factor of 1.2.
1421 ± 14		⁶ AUGUSTIN 92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
1416 ± 8 ± $\frac{7}{5}$	700	⁷ BAI 90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1413 ± 8	500	DUCH 89	ASTE	$\bar{p}p \rightarrow \pi^+\pi^- K^\pm \pi^\mp K^0$
1419 ± 1	8800 ± 200	^{8,9} BIRMAN 88	MPS	$8 \pi^- p \rightarrow K^+ \bar{K}^0 \pi^- n$
1424 ± 3	620	^{9,10} REEVES 86	SPEC	$6.6 \bar{p}p \rightarrow K\bar{K}\pi X$
1421 ± 2		CHUNG 85	SPEC	$8 \pi^- p \rightarrow K\bar{K}\pi n$
1425 ± 7	800	^{9,11} BAILLON 67	HBC	$0.0 \bar{p}p \rightarrow K\bar{K}\pi\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1459 ± 5		⁸ AUGUSTIN 92	DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
1445 ± 8	693 ± 30	⁹ AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1433 ± 8	296 ± 20	⁹ AUGUSTIN 90	DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1490 ± $\frac{14}{-8} \pm \frac{3}{-16}$	1100	⁶ BAI 90C	MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$
1443 ± 5		⁶ TAKAMATSU 90	SPEC	$8 \pi^- p \rightarrow nK^*(892)K$
1424 ± 4		TAKAMATSU 90	SPEC	$8 \pi^- p \rightarrow nK_S^0 K^\pm \pi^\mp$
1475 ± 4		¹² RATH 89	MPS	$21.4 \pi^- p \rightarrow nK_S^0 K_S^0 \pi^0$
1452.8 ± 6.8	170 ± 15	⁹ RATH 89	MPS	$21.4 \pi^- p \rightarrow K_S^0 K_S^0 \pi^0 n$
1412.8 ± 5.4		RATH 89	MPS	$21.4 \pi^- p \rightarrow nK_S^0 K_S^0 \pi^0$
1454 ± 3		WISNIEWSKI 87	MRK3	$J/\psi \rightarrow K\bar{K}\pi\gamma$
1440 ± $\frac{20}{-15}$	174	EDWARDS 82E	CBAL	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$
1440 ± $\frac{10}{-15}$		SCHARRE 80	MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

⁶ From fit to the $K^*(892)K 0^{-+}$ partial wave.⁷ From fit to the $a_0(980)\pi 1^{++}$ partial wave. cannot rule out a $a_0(980)\pi 1^{++}$ partial wave.⁸ From fit to the $a_0(980)\pi 0^{-+}$ partial wave.⁹ Best fit with a single Breit Wigner.¹⁰ From fit of the 0^{-+} partial wave, mainly $a_0(980)\pi$.¹¹ From best fit of 0^{-+} partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi$.¹² From fit to the $a_0(980)\pi 0^{-+}$ partial wave, but $a_0(980)\pi 1^{++}$ cannot be excluded. The fit is also consistent with one resonance at 1453 MeV.

Meson Full Listings

$\eta(1440)$

$\eta(1440)$ WIDTH

VALUE (MeV) DOCUMENT ID
60±30 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

$\eta\pi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50±8 OUR AVERAGE			
46±13	13 BURCHELL	91 MRK3	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
53±11	14 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
••• We do not use the following data for averages, fits, limits, etc. •••			
~50	14 BEHREND	92 DM2	$J/\psi \rightarrow \gamma\eta\pi^+\pi^-$
59±4	FUKUI	91C SPEC	$8.95\pi^-\rho \rightarrow \eta\pi^+\pi^-n$
31±7	ANDO	86 SPEC	$8\pi^-\rho \rightarrow n\eta\pi^+\pi^-$

13 From fit to the $a_0(980)\pi^0$ partial wave.
 14 From $\eta\pi^+\pi^-$ mass distribution - mainly $a_0(980)\pi^-$ - no spin-parity determination available.

$\pi\pi\gamma$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
174±44	AUGUSTIN	90 DM2	$J/\psi \rightarrow \pi^+\pi^-\gamma$
60±30	15 COFFMAN	90 MRK3	$J/\psi \rightarrow \pi^+\pi^-2\gamma$

15 This peak in the $\gamma\rho$ channel may not be related to the $\eta(1440)$.

4π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
144±13	3270	16 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

16 Estimated by us from various fits.

$K\bar{K}\pi$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
66.0±1.9 OUR AVERAGE				
63±18		AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
91 ⁺⁶⁷ ₋₃₁ ⁺¹⁵ ₋₃₈		BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm\pi^\mp$
62±16	500	DUCH	89 ASTE	$\bar{p}p \rightarrow K\bar{K}\pi\pi$
66±2	8800±200	BIRMAN	88 MPS	$8\pi^-\rho \rightarrow K^+\bar{K}^0\pi^-n$
60±10	620	17 REEVES	86 SPEC	$6.6\rho\bar{p} \rightarrow K K\pi X$
60±10		CHUNG	85 SPEC	$8\pi^-\rho \rightarrow K\bar{K}\pi n$
80±10	800	18 BAILLON	67 HBC	$0.0\bar{p}p \rightarrow K\bar{K}\pi\pi$
••• We do not use the following data for averages, fits, limits, etc. •••				
75±9		19 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K\bar{K}\pi$
75±9	693±30	19 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm\pi^\mp$
93±14	296±20	AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K^+K^-\pi^0$
105±10	693±30	AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma K_S^0 K^\pm\pi^\mp$
54 ⁺³⁷ ₋₂₁ ⁺¹³ ₋₂₄		20 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm\pi^\mp$
59±4		20 TAKAMATSU	90 SPEC	$9\pi^-\rho \rightarrow n\eta\pi^+\pi^-$
82±8		TAKAMATSU	90 SPEC	$8\pi^-\rho \rightarrow nK_S^0 K^\pm\pi^\mp$
57±8		TAKAMATSU	90 SPEC	$8\pi^-\rho \rightarrow nK^*(892)K$
51±13		21 RATH	89 MPS	$21.4\pi^-\rho \rightarrow nK_S^0 K_S^0\pi^0$
99.9±11.4	170±15	22 RATH	89 MPS	$21.4\pi^-\rho \rightarrow K_S^0 K_S^0\pi^0 n$
19±7		RATH	89 MPS	$21.4\pi^-\rho \rightarrow nK_S^0 K_S^0\pi^0$
160±11		WISNIEWSKI	87 MRK3	$J/\psi \rightarrow K\bar{K}\pi\gamma$
55 ⁺²⁰ ₋₃₀	174	EDWARDS	82E CBAL	$J/\psi \rightarrow \gamma K^+K^-\pi^0$
50 ⁺³⁰ ₋₂₀		SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K_S^0 K^\pm\pi^\mp$

17 From best fit to 0^- partial wave, 50% $K^*(892)K$, 50% $a_0(980)\pi^-$.
 18 From fit to the 0^- partial wave, mainly $a_0(980)\pi^-$.
 19 From fit to the $a_0(980)\pi^0$ partial wave.
 20 From fit to the $K^*(892)K^0$ partial wave.
 21 From fit to the $a_0(980)\pi^0$ partial wave, but $a_0(980)\pi^1$ cannot be excluded.
 The fit is also consistent with one resonance at 1453 MeV.
 22 Best fit with a single Breit Wigner.

$\eta(1440)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}\pi$	seen
Γ_2 $\eta\pi\pi$	seen
Γ_3 $a_0(980)\pi$	seen
Γ_4 $\pi\pi\rho$	
Γ_5 $K\bar{K}^*(892) + c.c.$	
Γ_6 4π	seen
Γ_7 $\gamma\gamma$	
Γ_8 $\rho^0\gamma$	

$\eta(1440)$ $\Gamma(i)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
<1.2	95	BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm\pi^\mp$
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.6	95	AIHARA	86D TPC	$e^+e^- \rightarrow e^+e^- K_S^0 K^\pm\pi^\mp$
<2.2	95	ALTHOFF	85B TASS	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$
<8.0	95	JENNI	83 MRK2	$e^+e^- \rightarrow e^+e^- K\bar{K}\pi$

$\Gamma(\eta\pi\pi) \times \Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
<0.3	ANTREASYAN	87 CBAL	$e^+e^- \rightarrow e^+e^-\eta\pi\pi$

$\Gamma(\rho^0\gamma) \times \Gamma(\gamma\gamma)/\Gamma(\text{total})$

VALUE (keV)	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<1.5	95	ALTHOFF	84E TASS	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\gamma$

$\eta(1440)$ BRANCHING RATIOS

$\Gamma(\eta\pi\pi)/\Gamma(K\bar{K}\pi)$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••					
	<0.5	90	EDWARDS	83B CBAL	$J/\psi \rightarrow \eta\pi\pi\gamma$
	<1.1	90	SCHARRE	80 MRK2	$J/\psi \rightarrow \eta\pi\pi\gamma$
	<1.5	95	FOSTER	68B HBC	0.0 $\bar{p}p$

$\Gamma(a_0(980)\pi)/\Gamma(K\bar{K}\pi)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
~0.8	500	23 DUCH	89 ASTE	$\bar{p}p \rightarrow \pi^+\pi^-K^\pm\pi^\mp K^0$
••• We do not use the following data for averages, fits, limits, etc. •••				
~0.75		23 REEVES	86 SPEC	$6.6\rho\bar{p} \rightarrow K K\pi X$

23 Assuming that the $a_0(980)$ decays only into $K\bar{K}$.

$\Gamma(K\bar{K}^*(892) + c.c.)/\Gamma(K\bar{K}\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.50±0.10	BAILLON	67 HBC	0.0 $\bar{p}p \rightarrow K\bar{K}\pi\pi$

$\Gamma(K\bar{K}^*(892) + c.c.)/[\Gamma(a_0(980)\pi) + \Gamma(K\bar{K}^*(892) + c.c.)]$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.25	90	EDWARDS	82E CBAL	$J/\psi \rightarrow K^+K^-\pi^0\gamma$

$\Gamma(\rho^0\gamma)/\Gamma(K\bar{K}\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.0152±0.0038	24 COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$

24 Using $B(J/\psi \rightarrow \gamma\eta(1440)) \rightarrow \gamma K\bar{K}\pi = 4.2 \times 10^{-3}$ and $B(J/\psi \rightarrow \gamma\eta(1440)) \rightarrow \gamma\gamma\rho^0 = 6.4 \times 10^{-5}$ and assuming that the $\gamma\rho^0$ signal does not come from the $f_1(1420)$.

$\eta(1440)$ REFERENCES

AUGUSTIN	92	PR D46 1951	+Cosme	(DM2 Collab.)
BEHREND	92	ZPHY C36 381		(CELLO Collab.)
BURCHELL	91	NP B21 132 (suppl)		(Mark III Collab.)
FUKUI	91C	PL B267 293	+ (SUGI, NAGO, KEK, KYOT, MIYA, AKIT)	(Mark III Collab.)
AUGUSTIN	90	PR D42 10	+Cosme+	(DM2 Collab.)
BAI	90C	PR L65 2507	+Blaylock+	(Mark III Collab.)
COFFMAN	90	PR D41 1410	+De Jongh+	(Mark III Collab.)
TAKAMATSU	90	Hadron 89 Conf. p 71	+Ando+	(KEK)
BEHREND	89	ZPHY C42 367	+Criegee+	(CELLO Collab.)
BISELLO	89B	D39 701	+Busetto+	(DM2 Collab.)
DUCH	89	ZPHY 45 223	+Heel, Bailey+	(ASTERIX Collab.) JP
RATH	89	PR D40 693	+Cason+	(NDAM, BRAN, BNL, CUNY, DUKE)
BIRMAN	88	PR L61 1557	+Chung, Peaslee+	(BNL, FSU, IND, MASA) JP
ANTREASYAN	87	PR D36 2633	+Bartels, Besset+	(Crystal Ball Collab.)
WISNIEWSKI	87	Hadron 87 Conf.		(Mark III Collab.)
AIHARA	86D	PRL 57 51	+Alston-Garnjost+	(TPC-2 γ Collab.)
ANDO	86	PRL 57 1296	+Imai+	(KEK, KYOT, NIRS, SAGA, INUS, TSUKUBA) JP
REEVES	86	PR L34 1960	+Chung, Crittenden+	(FLOR, BNL, IND, MASA) JP
ALTHOFF	85B	ZPHY C29 189	+Braunschweig, Kirschfink+	(TASSO Collab.)
CHUNG	85	PRL 55 779	+Fermov, Boehnlein+	(BNL, FLOR, IND, MASA) JP
ALTHOFF	84E	PL 147B 487	+Braunschweig, Kirschfink, Luebelsmeyer+	(TASSO Collab.)
EDWARDS	83B	PL 51 859	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
JENNI	83	PR D27 1031	+Burke, Telnov, Abrams, Blocker+	(SLAC, LBL)
EDWARDS	82E	PRL 49 259	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
Also	83	PRL 50 219	+Edwards, Partridge+	(CIT, HARV, PRIN, STAN+)
SCHARRE	80	PL 97B 329	+Trilling, Abrams, Alam, Blocker+	(SLAC, LBL)
FOSTER	68B	NP B8 174	+Gavillet, Labrosse, Montanet+	(CERN, CDEF)
BAILLON	67	NC 50A 393	+Edwards, D'Andiau, Astier+	(CERN, CDEF, IRAD)

OTHER RELATED PAPERS

AHMAD 89	NP B (PROC.)8 50	+Amsler, Auld+	(ASTERIX Collab.)
ARMSTRONG 89	PL B221 216	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)	(IND)
ZIEMINSKA 88	AIP Conf.	+Bloodworth+ (CERN, BIRM, BARI, ATHU, CURIN+)	(IND)
ARMSTRONG 87	ZPHY C34 23	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
ASTON 87	NP B292 693	Protopopescu, Chung	(BNL)
PROTOPOP... 87B	Hadron 87 Conf.	+Bloodworth, Burns+	(ATHU, BARI, BIRM, CERN)
TOKI 87	Hadron 87 Conf.	+Gavillet+	(CERN, MADR, CDEF, STOH)
ARMSTRONG 84	PL 146B 273	+Nascimento, Bizzarri+	(CDEF, CERN)
DIONISI 80	NP B169 1	+Goldberg, Makowski, Donald+	(PARIS, LIVP)
DEFOIX 72	NP B44 125	+D'Andiau, Astier+	(CDEF, CERN)
DUBOC 72	NP B46 429		
LORSTAD 69	NP B14 63		

 $\rho(1450)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

See the mini-review under the $\rho(1700)$. $\rho(1450)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1465±25 OUR ESTIMATE			This is only an educated guess; the error given is larger than the error on the average of the published values.
1451± 8 OUR AVERAGE			Includes data from the 4 datablocks that follow this one.

MIXED MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1465±25	DONNACHIE 87	RVUE	

 $\eta\rho^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1470±20	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
1446±10	FUKUI 88	SPEC	$8.95\pi^-p \rightarrow \eta\pi^+\pi^-n$

 $\pi^+\pi^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1424±25	BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$

 $\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1463±25	¹ DONNACHIE 91	RVUE	
••• We do not use the following data for averages, fits, limits, etc. •••			
1250	² ASTON 80C	OMEG 20-70	$\gamma p \rightarrow \omega\pi^0 p$
1290±40	² BARBER 80C	SPEC 3-5	$\gamma p \rightarrow \omega\pi^0 p$
¹ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.			
² Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.			

 $\pi^+\pi^-\pi^+\pi^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1449±4	³ ARMSTRONG 89E	OMEG 300	$p p \rightarrow \rho p 2(\pi^+\pi^-)$
³ Not clear whether this observation has $I=1$ or 0.			

 $\phi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1480±40	⁴ BITYUKOV 87	SPEC 0		$32.5\pi^-p \rightarrow \phi\pi^0 n$
⁴ See the minireview for $\rho(1700)$ and ACHASOV 88 for a non-exotic interpretation. DONNACHIE 91 suggests this is a different particle.				

 $\rho(1450)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
310±60 OUR ESTIMATE			This is only an educated guess; the error given is larger than the error on the average of the published values.

 $\omega\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
311± 62	⁵ DONNACHIE 91	RVUE	
••• We do not use the following data for averages, fits, limits, etc. •••			
300	⁶ ASTON 80C	OMEG 20-70	$\gamma p \rightarrow \omega\pi^0 p$
320±100	⁶ BARBER 80C	SPEC 3-5	$\gamma p \rightarrow \omega\pi^0 p$
⁵ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.			
⁶ Not separated from $b_1(1235)$, not pure $J^P = 1^-$ effect.			

MIXED MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
220±25	DONNACHIE 87	RVUE	••• We do not use the following data for averages, fits, limits, etc. •••

 $\eta\rho^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
230±30	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
60±15	FUKUI 88	SPEC	$8.95\pi^-p \rightarrow \eta\pi^+\pi^-n$

 $\pi^+\pi^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
269±31	BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$

 $\pi^+\pi^-\pi^+\pi^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
78±18	⁷ ARMSTRONG 89E	OMEG 300	$p p \rightarrow \rho p 2(\pi^+\pi^-)$
⁷ Not clear whether this observation has $I=1$ or 0.			

 $\phi\pi$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
130±60	⁸ BITYUKOV 87	SPEC 0		$32.5\pi^-p \rightarrow \phi\pi^0 n$
⁸ See the minireview for $\rho(1700)$ and ACHASOV 88 for a non-exotic interpretation. DONNACHIE 91 suggests this is a different particle.				

 $\rho(1450)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \pi\pi$	seen	
$\Gamma_2 4\pi$	seen	
$\Gamma_3 e^+e^-$	seen	
$\Gamma_4 \eta\rho$	<4 %	
$\Gamma_5 \omega\pi$	<2.0 %	95%
$\Gamma_6 \phi\pi$	<1 %	
$\Gamma_7 K\bar{K}$	< 1.6×10^{-3}	95%

 $\rho(1450)$ $\Gamma(I)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\pi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_3/\Gamma$
VALUE (keV)				
0.12	⁹ DIEKMAN 88	RVUE	$e^+e^- \rightarrow \pi^+\pi^-$	
⁹ Using total width = 235 MeV.				

$\Gamma(\eta\rho) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_4\Gamma_3/\Gamma$
VALUE (eV)				
91±19	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$	

$\Gamma(\phi\pi) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_6\Gamma_3/\Gamma$
VALUE (eV)				
<70	¹⁰ AULCHENKO 87B	ND	$e^+e^- \rightarrow K_S^0 K_L^0 \pi^0$	
¹⁰ Using mass 1480 ± 40 MeV and total width 130 ± 60 MeV of BITYUKOV 87.				

 $\rho(1450)$ BRANCHING RATIOS

$\Gamma(\eta\rho)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
VALUE				
<0.04	DONNACHIE 87B	RVUE		

$\Gamma(\phi\pi)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ_5
VALUE					
>0.5	BITYUKOV 87	SPEC 0		$32.5\pi^-p \rightarrow \phi\pi^0 n$	

$\Gamma(\omega\pi)/\Gamma(4\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
VALUE				
<0.14	CLEGG 88	RVUE		

$\Gamma(\eta\rho)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_5
VALUE				
~0.24	¹¹ DONNACHIE 91	RVUE		
••• We do not use the following data for averages, fits, limits, etc. •••				
>2	FUKUI 91	SPEC	$8.95\pi^-p \rightarrow \omega\pi^0 n$	

Meson Full Listings

$\rho(1450)$, $f_1(1510)$, $f_2(1520)$

$\Gamma(\omega\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	Γ_5/Γ
VALUE			
~ 0.21	11 DONNACHIE	91 RVUE	
$\Gamma(\pi\pi)/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_1/Γ_5
VALUE			
~ 0.24	11 DONNACHIE	91 RVUE	
$\Gamma(\phi\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	Γ_6/Γ
VALUE			
< 0.01	11 DONNACHIE	91 RVUE	
$\Gamma(K\bar{K})/\Gamma(\omega\pi)$	DOCUMENT ID	TECN	Γ_7/Γ_5
VALUE			
< 0.08	11 DONNACHIE	91 RVUE	

¹¹ Using data from BISELLO 91B, DOLINSKY 86 and ALBRECHT 87L.

$\rho(1450)$ REFERENCES

BISELLO	91B	NP B21 111 (suppl)		(DM2 Collab.)
DONNACHIE	91	ZPHY C51 689	+Clegg	(MCHS, LANC)
FUKUI	91	PL B257 241	+Horikawa+	(SUGI, NAGO, KEK, KYOT, MIYA)
ARMSTRONG	89E	PL B228 536	+Benayoun (ATHU, BARI, BIRM, CERN, CDEF, CURIN+)	
BISELLO	89	PL B220 321	+Busetto+	(DM2 Collab.)
ACHASOV	88	PL B207 199	+Kozhevnikov	(NOVO)
ANTONELLI	88	PL B212 133	+Baldini+	(DM2 Collab.)
CLEGG	88	ZPHY C40 313	+Donnachie	(MCHS, LANC)
DIEKMANN	88	PRPL 159 101		(BONN)
FUKUI	88	PL B202 441	+Horikawa+	(SUGI, NAGO, KEK, KYOT, MIYA)
ALBRECHT	87L	PL B185 223	+Binder, Boeckmann, Glaser+	(ARGUS Collab.)
AULCHENKO	87B	JETPL 45 145	+Dolinsky, Druzhinin, Dubrovin+	(NOVO)
		Translated from ZETFP 45 118		
BITYUKOV	87	PL B188 383	+Dzhelyadin, Dorofeev, Golovkin+	(SERP)
DONNACHIE	87	ZPHY C33 407	+Mirzaie	(MCHS)
DONNACHIE	87B	ZPHY C34 257	+Clegg	(MCHS, LANC)
DOLINSKY	86	PL B174 453	+Druzhinin, Dubrovin, Eidelman+	(NOVO)
ASTON	80C	PL 92B 211		(BONN, CERN, EPOL, GLAS, LANC, MCHS+)
BARBER	80C	ZPHY C4 169	+Dainton, Brodbeck, Brookes+	(DARE, LANC, SHEF)

OTHER RELATED PAPERS

LANDSBERG	92	SJNP 55 1051		(SERP)
		Translated from YAF 55 1896		
BRAU	88	PR D37 2379	+Frank+	(SLAC Hybrid Facility Photon Collab.)
ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
KURDADZE	86	JETPL 43 643	+Lechuk, Pakhtusova, Sidorov, Skirinski+	(NOVO)
		Translated from ZETFP 43 497		
BARKOV	85	NP B256 365	+Chilingarov, Eidelman, Khazin, Lechuk+	(NOVO)
BISELLO	85	LAL 85-15	+Augustin, Ajaltouni+	(PADO, LALO, CLER, FRAS)
ABE	84B	PRL 53 751	+Bacon, Ballam+	(SLAC Hybrid Facility Photon Collab.)
ATKINSON	84C	NP B243 1		(BONN, CERN, GLAS, LANC, MCHS, CURIN+)
CORDIER	82	PL 109B 129	+Bisello, Bizot, Buon, Delcourt	(LALO)
KILLIAN	80	PR D21 3005	+Treadwell, Ahrens, Berkelman, Cassel+	(CORN)
COSME	76	PL 63B 352	+Courau, Dudelzak, Grelaud, Jean-Marie+	(ORSAY)
BINGHAM	72B	PL 41B 635	+Rabin, Rosenfeld, Smadja+	(LBL, UCB, SLAC)
FRENKIEL	72	NP B47 61	+Gesquiere, Lillestol, Chung+	(CDEF, CERN)
LAYSSAC	71	NC 6A 134	+Renard	(MONP)

$f_1(1510)$

$$I^G(J^{PC}) = 0^+(1^{++})$$

See also minireview under non- $q\bar{q}$ candidates.

$f_1(1510)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1512 ± 4	600 ± 200	¹ BIRMAN	88 MPS	$8\pi^-p \rightarrow K^+\bar{K}^0\pi^-n$
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 1525		² BAUER	93B	$\gamma\gamma^* \rightarrow \pi^+\pi^-\pi^0\pi^0$
1530 ± 10		ASTON	88C LASS	$11K^-p \rightarrow K_S^0 K^\pm\pi^\mp\Lambda$
1526.0 ± 6.0	271	GAVILLET	82 HBC	$4.2K^-p \rightarrow \Lambda K K\pi$

¹ From partial wave analysis of $K^+\bar{K}^0\pi^-$ state.

² Possibly a different resonance than that seen in $K\bar{K}\pi$, isospin and spin uncertain.

$f_1(1510)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
35 ± 15	600 ± 200	³ BIRMAN	88 MPS	$8\pi^-p \rightarrow K^+\bar{K}^0\pi^-n$
••• We do not use the following data for averages, fits, limits, etc. •••				
100 ± 40		ASTON	88C LASS	$11K^-p \rightarrow K_S^0 K^\pm\pi^\mp\Lambda$
107.0 ± 15.0	271	GAVILLET	82 HBC	$4.2K^-p \rightarrow \Lambda K K\pi$

³ From partial wave analysis of $K^+\bar{K}^0\pi^-$ state.

$f_1(1510)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}^*(892) + c.c.$	seen

$f_1(1510)$ REFERENCES

BAUER	93B	PR D48 3976	+Belcinski, Berg, Bingham+	(SLAC)
ASTON	88C	PL B201 573	+Awaji, Biernz+	(SLAC, NAGO, CINC, INUS) JP
BIRMAN	88	PRL 61 1557	+Chung, Peaslee+	(BNL, FSI, IND, MASD) JP
GAVILLET	82	ZPHY C16 119	+Armenteros+	(CERN, CDEF, PADO, ROMA)

$f_2(1520)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in antiproton-nucleon annihilation at rest. See also minireview under non- $q\bar{q}$ candidates. Needs confirmation.

$f_2(1520)$ MASS


VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
1510 ± 8 OUR AVERAGE					Error includes scale factor of 1.8. See the ideogram below.
1502 ± 9		¹ ADAMO	93 OBLX		$\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
1488 ± 10		ARMSTRONG	93C SPEC		$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
1508 ± 10		ARMSTRONG	93D SPEC		$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
1525 ± 10		ARMSTRONG	93D SPEC		$\bar{p}p \rightarrow \eta\pi^0\pi^0 \rightarrow 6\gamma$
1515 ± 10		AKER	91 CBAR		$0.0\bar{p}p \rightarrow 3\pi^0$
1565 ± 20		MAY	90 ASTE		$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••					
1566 ⁺⁸⁰ ₋₅₀		² ANISOVICH	94 CBAR		$0.0\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0$
1527 ± 5	435 ± 45	³ GRAY	83 DBC	0	$0.0\bar{p}N \rightarrow 3\pi$

¹ Supersedes ADAMO 92.

² From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ including AKER 91 data.

³ No fit of the Dalitz plot has been made. $J=0$ is cautiously suggested, but $J=2$ is not excluded.

WEIGHTED AVERAGE
1510±8 (Error scaled by 1.8)



$f_2(1520)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
120 ± 5 OUR AVERAGE					
130 ± 10		⁴ ADAMO	93 OBLX		$\bar{p}p \rightarrow \pi^+\pi^+\pi^-$
148 ± 27		ARMSTRONG	93C SPEC		$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
103 ± 15		ARMSTRONG	93D SPEC		$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
111 ± 10		ARMSTRONG	93D SPEC		$\bar{p}p \rightarrow \eta\pi^0\pi^0 \rightarrow 6\gamma$
120 ± 10		AKER	91 CBAR		$0.0\bar{p}p \rightarrow 3\pi^0$
170 ± 40		MAY	90 ASTE		$\bar{p}p \rightarrow \pi^+\pi^-\pi^0$
••• We do not use the following data for averages, fits, limits, etc. •••					
166 ⁺⁸⁰ ₋₂₀		⁵ ANISOVICH	94 CBAR		$0.0\bar{p}p \rightarrow 3\pi^0, \eta\eta\pi^0$
101 ± 19	435 ± 45	^{6,7} GRAY	83 DBC	0	$0.0\bar{p}N \rightarrow 3\pi$

⁴ Supersedes ADAMO 92.

⁵ From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$ including AKER 91 data.

⁶ No fit of the Dalitz plot has been made.

⁷ Width error enlarged by us to $4\Gamma/N^{1/2}$.

See key on page 1343

Meson Full Listings

 $f_2(1520), f_2'(1525)$ $f_2(1520)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \pi^+ \pi^-$	seen
$\Gamma_2 \rho^0 \rho^0$	
$\Gamma_3 \pi^0 \pi^0$	
$\Gamma_4 2\pi^+ 2\pi^-$	
$\Gamma_5 \eta \eta$	

 $f_2(1520)$ BRANCHING RATIOS

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
VALUE					
seen	MAY	89	ASTE	$\bar{p} p \rightarrow \pi^+ \pi^- \pi^0$	
seen	GRAY	83	DBC	0 0.0 $\bar{p} N \rightarrow 3\pi$	

$\Gamma(\pi^+ \pi^-)/\Gamma(\rho^0 \rho^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ_2
VALUE					
0.042 ± 0.013	BRIDGES	86B	DBC	0 $\bar{p} N \rightarrow 3\pi^- 2\pi^+$	

$\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ	
VALUE					
seen	AKER	91	CBAR	0.0 $\bar{p} p \rightarrow 3\pi^0$	

$\Gamma(\eta \eta)/\Gamma(\pi^0 \pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_3	
VALUE					
0.024 ± 0.005 ± 0.012	ARMSTRONG	93C	SPEC	$\bar{p} p \rightarrow \pi^0 \eta \eta \rightarrow 6\gamma$	

 $f_2(1520)$ REFERENCES

ANISOVICH 94 PL B323 233	+Armstrong+	(Crystal Barrel Collab.)
ADAMO 93 NP A558 13C	+Agnello+	(OBELIX Collab.)
ARMSTRONG 93C PL B307 394	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
ARMSTRONG 93D PL B307 399	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
ADAMO 92 PL B287 368	+Agnello, Balestra+	(OBELIX Collab.)
AKER 91 PL B260 249	+Amsler, Peters+	(Crystal Barrel Collab.)
MAY 90 ZPHY C46 203	+Duch, Heel+	(ASTERIX Collab.)
MAY 89 PL B225 450	+Duch, Heel+	(ASTERIX Collab.)
BRIDGES 86B PRL 56 215	+Daftari, Kalogeropoulos, Debbe+	(SYRAX, CASE) IJP
GRAY 83 PR D27 307	+Kalogeropoulos, Nandy, Roy, Zenone	(SYRAX)

 $f_2'(1525)$

$$I^G(J^{PC}) = 0^+(2^+ +)$$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)

 $f_2'(1525)$ MASS

VALUE (MeV)	DOCUMENT ID
1525 ± 5 OUR ESTIMATE	This is only an educated guess; the error given is larger than the error on the average of the published values.

PRODUCED BY PION BEAM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1547 ⁺¹⁰ ₋₂		1 LONGACRE	86 MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
1496 ⁺⁹ ₋₈		2 CHABAUD	81 ASPK	6 $\pi^- p \rightarrow K^+ K^- n$
1497 ⁺⁸ ₋₉		CHABAUD	81 ASPK	18.4 $\pi^- p \rightarrow K^+ K^- n$
1492 ± 29		GORLICH	80 ASPK	17 $\pi^- p$ polarized → $K^+ K^- n$
1502 ± 25		3 CORDEN	79 OMEG	12-15 $\pi^- p \rightarrow K^+ K^- n$
1480	14	CRENNELL	66 HBC	6.0 $\pi^- p \rightarrow K_S^0 K_S^0 n$

PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1524.5 ± 1.4 OUR AVERAGE		Includes data from the datablock that follows this one. Error includes scale factor of 1.1.		
1526.8 ± 4.3		ASTON	88D LASS	11 $K^- p \rightarrow K_S^0 K_S^0 \Lambda$
1529 ± 3		ARMSTRONG	83B OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$
1521 ± 6	650	AGUILAR...	81B HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
1521 ± 3	572	ALHARRAN	81 HBC	8.25 $K^- p \rightarrow \Lambda K \bar{K}$
1522 ± 6	123	BARREIRO	77 HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
1528 ± 7	166	EVANGELISTA	77 OMEG	10 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1527 ± 3	120	BRANDENB...	76C ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
1519 ± 7	100	AGUILAR...	72B HBC	3.9, 4.6 $K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$

PRODUCED IN $e^+ e^-$ ANNIHILATION

VALUE (MeV) DOCUMENT ID TECN COMMENT
The data in this block is included in the average printed for a previous datablock.

1519 ± 5 OUR AVERAGE	Error includes scale factor of 1.1.		
1531.6 ± 10.0	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
1515 ± 5	4 FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$
1525 ± 10 ± 10	BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1496 ± 2	5 FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$
¹	From a partial-wave analysis of data using a K-matrix formalism with 5 poles.		
²	CHABAUD 81 is a reanalysis of PAWLICKI 77 data.		
³	From an amplitude analysis where the $f_2'(1525)$ width and elasticity are in complete disagreement with the values obtained from $K \bar{K}$ channel, making the solution dubious.		
⁴	From an analysis ignoring interference with $f_J(1710)$.		
⁵	From an analysis including interference with $f_J(1710)$.		

 $f_2'(1525)$ WIDTH

VALUE (MeV)	DOCUMENT ID	COMMENT
76 ± 10 OUR ESTIMATE	This is only an educated guess; the error given is larger than the error on the average of the published values.	
85 ± 5 OUR FIT		
76 ± 10	PDG	90 For fitting

PRODUCED BY PION BEAM

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
108 ⁺⁵ ₋₂	6 LONGACRE	86 MPS	22 $\pi^- p \rightarrow K_S^0 K_S^0 n$
69 ⁺²² ₋₁₆	7 CHABAUD	81 ASPK	6 $\pi^- p \rightarrow K^+ K^- n$
137 ⁺²³ ₋₂₁	CHABAUD	81 ASPK	18.4 $\pi^- p \rightarrow K^+ K^- n$
150 ⁺⁸³ ₋₅₀	GORLICH	80 ASPK	17 $\pi^- p$ polarized → $K^+ K^- n$
165 ± 42	8 CORDEN	79 OMEG	12-15 $\pi^- p \rightarrow \pi^+ \pi^- n$
92 ⁺³⁹ ₋₂₂	9 POLYCHRO...	79 STRC	7 $\pi^- p \rightarrow n K_S^0 K_S^0$

PRODUCED BY K^\pm BEAM

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
78 ± 5 OUR AVERAGE		Includes data from the datablock that follows this one.		
90 ± 12		ASTON	88D LASS	11 $K^- p \rightarrow K_S^0 K_S^0 \Lambda$
83 ± 15		ARMSTRONG	83B OMEG	18.5 $K^- p \rightarrow K^- K^+ \Lambda$
85 ± 16	650	AGUILAR...	81B HBC	4.2 $K^- p \rightarrow \Lambda K^+ K^-$
80 ⁺¹⁴ ₋₁₁	572	ALHARRAN	81 HBC	8.25 $K^- p \rightarrow \Lambda K \bar{K}$
72 ± 25	166	EVANGELISTA	77 OMEG	10 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$
69 ± 22	100	AGUILAR...	72B HBC	3.9, 4.6 $K^- p \rightarrow K \bar{K} (\Lambda, \Sigma)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
62 ⁺¹⁹ ₋₁₄	123	BARREIRO	77 HBC	4.15 $K^- p \rightarrow \Lambda K_S^0 K_S^0$
61 ± 8	120	BRANDENB...	76C ASPK	13 $K^- p \rightarrow K^+ K^- (\Lambda, \Sigma)$

PRODUCED IN $e^+ e^-$ ANNIHILATION

VALUE (MeV) DOCUMENT ID TECN COMMENT
The data in this block is included in the average printed for a previous datablock.

67 ± 9 OUR AVERAGE			
103 ± 30	AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
62 ± 10	10 FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$
85 ± 35	BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
100 ± 3	11 FALVARD	88 DM2	$J/\psi \rightarrow \phi K^+ K^-$
⁶	From a partial-wave analysis of data using a K-matrix formalism with 5 poles.		
⁷	CHABAUD 81 is a reanalysis of PAWLICKI 77 data.		
⁸	From an amplitude analysis where the $f_2'(1525)$ width and elasticity are in complete disagreement with the values obtained from $K \bar{K}$ channel, making the solution dubious.		
⁹	From a fit to the D with $f_2(1270)$ - $f_2'(1525)$ interference. Mass fixed at 1516 MeV.		
¹⁰	From an analysis ignoring interference with $f_J(1710)$.		
¹¹	From an analysis including interference with $f_J(1710)$.		

Meson Full Listings

 $f_2'(1525)$ $f_2'(1525)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	$(71.2^{+2.0}_{-2.5})\%$
Γ_2 $\eta\eta$	$(27.9^{+2.5}_{-2.0})\%$
Γ_3 $\pi\pi$	$(8.2 \pm 1.6) \times 10^{-3}$
Γ_4 $\gamma\gamma$	$(1.23 \pm 0.22) \times 10^{-6}$
Γ_5 $K\bar{K}^*(892) + c.c.$	
Γ_6 $\pi\pi\eta$	
Γ_7 $\pi K\bar{K}$	
Γ_8 $\pi^+\pi^+\pi^-\pi^-$	

CONSTRAINED FIT INFORMATION

An overall fit to the total width, 4 partial widths, a combination of partial widths obtained from integrated cross sections, and 2 branching ratios uses 13 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 10.0$ for 9 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100			
x_3	-6	-2		
x_4	-29	29	0	
Γ	61	-61	2	-39
	x_1	x_2	x_3	x_4

Mode	Rate (MeV)
Γ_1 $K\bar{K}$	61 ± 5
Γ_2 $\eta\eta$	$23.9^{+2.2}_{-1.2}$
Γ_3 $\pi\pi$	0.70 ± 0.14
Γ_4 $\gamma\gamma$	$(1.05 \pm 0.17) \times 10^{-4}$

 $f_2'(1525)$ PARTIAL WIDTHS

$\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_1
VALUE (MeV)				
61 ± 5 OUR FIT				
63.0^{+6.0}_{-5.0}	¹² LONGACRE	86	MPS	$22 \pi^- \rho \rightarrow K_S^0 K_S^0 n$
$\Gamma(\pi\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3
VALUE (MeV)				
0.70 ± 0.14 OUR FIT				
1.4^{+1.0}_{-0.5}	¹² LONGACRE	86	MPS	$22 \pi^- \rho \rightarrow K_S^0 K_S^0 n$
$\Gamma(\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_2
VALUE (MeV)				
23.9^{+2.2}_{-1.2} OUR FIT				
24.0^{+3.0}_{-1.0}	¹² LONGACRE	86	MPS	$22 \pi^- \rho \rightarrow K_S^0 K_S^0 n$
$\Gamma(\gamma\gamma)$	DOCUMENT ID	TECN	COMMENT	Γ_4
VALUE (keV)				
0.105 ± 0.017 OUR FIT				
0.107^{+0.029}_{-0.022} OUR AVERAGE				
0.11 ^{+0.03} _{-0.02} ± 0.02	BEHREND	89C	CELL	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$
0.10 ^{+0.04} _{-0.03} ^{+0.03} _{-0.02}	BERGER	88	PLUT	$e^+e^- \rightarrow e^+e^- K_S^0 K_S^0$

¹² From a partial-wave analysis of data using a K-matrix formalism with 5 poles.

 $f_2'(1525)$ $\Gamma(I)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_4/\Gamma$
VALUE (keV)				
0.075 ± 0.012 OUR FIT				
0.074 ± 0.016 OUR AVERAGE				
0.067 ± 0.008 ± 0.015	¹³ ALBRECHT	90G	ARG	$e^+e^- \rightarrow e^+e^- K^+ K^-$
0.12 ± 0.07 ± 0.04	¹³ AIHARA	86B	TPC	$e^+e^- \rightarrow e^+e^- K^+ K^-$
0.11 ± 0.02 ± 0.04	¹³ ALTHOFF	83	TASS	$e^+e^- \rightarrow e^+e^- K^+ K^-$
0.0314 ± 0.0050 ± 0.0077	¹⁴ ALBRECHT	90G	ARG	$e^+e^- \rightarrow e^+e^- K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹³ Using an incoherent background.
¹⁴ Using a coherent background.

 $f_2'(1525)$ BRANCHING RATIOS

$\Gamma(\eta\eta)/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE				
0.39^{+0.05}_{-0.04} OUR FIT				
0.11 ± 0.04	¹⁵ PROKOSHKIN	91	GAM4	$300 \pi^- \rho \rightarrow \pi^- \rho \eta \eta$
< 0.50	BARNES	67	HBC	$4.6, 5.0 K^- \rho$

¹⁵ Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma \eta \eta$.

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ	
VALUE					
0.0082 ± 0.0016 OUR FIT					
0.0075 ± 0.0016 OUR AVERAGE					
0.007 ± 0.002	COSTA...	80	OMEG	$10 \pi^- \rho \rightarrow K^+ K^- n$	
0.027 ^{+0.071} _{-0.013}	¹⁶ GORLICH	80	ASPK	$17, 18 \pi^- \rho$	
0.0075 ± 0.0025	^{16,17} MARTIN	79	RVUE		
< 0.06	95	AGUILAR...	81B	HBC	$4.2 K^- \rho \rightarrow \Lambda K^+ K^-$
0.19 ± 0.03		CORDEN	79	OMEG	$12-15 \pi^- \rho \rightarrow \pi^+ \pi^- n$
< 0.045	95	BARREIRO	77	HBC	$4.15 K^- \rho \rightarrow \Lambda K_S^0 K_S^0$
0.012 ± 0.004	¹⁶ PAWLICKI	77	SPEC	$6 \pi N \rightarrow K^+ K^- N$	
< 0.063	90	BRANDENB...	76C	ASPK	$13 K^- \rho \rightarrow K^+ K^- (\Lambda, \Sigma)$
< 0.0086	¹⁶ BEUSCH	75B	OSPK	$8.9 \pi^- \rho \rightarrow K^0 \bar{K}^0 n$	

¹⁶ Assuming that the $f_2'(1525)$ is produced by an one-pion exchange production mechanism.
¹⁷ MARTIN 79 uses the PAWLICKI 77 data with different input value of the $f_2'(1525) \rightarrow K\bar{K}$ branching ratio.

$\Gamma(\pi\pi)/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
VALUE				
0.0115 ± 0.0022 OUR FIT				
0.075 ± 0.035	AUGUSTIN	87	DM2	$J/\psi \rightarrow \gamma \pi^+ \pi^-$

$\Gamma(\pi\pi\eta)/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_1	
VALUE					
0.0115 ± 0.0022 OUR FIT					
< 0.41	95	AGUILAR...	72B	HBC	$3.9, 4.6 K^- \rho$
< 0.3	67	AMMAR	67	HBC	

$[\Gamma(K\bar{K}^*(892) + c.c.) + \Gamma(\pi K\bar{K})]/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_5 + \Gamma_7)/\Gamma_1$	
VALUE					
0.107 ± 0.029 OUR AVERAGE					
< 0.35	95	AGUILAR...	72B	HBC	$3.9, 4.6 K^- \rho$
< 0.4	67	AMMAR	67	HBC	

$\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_1	
VALUE					
0.107 ± 0.029 OUR AVERAGE					
< 0.32	95	AGUILAR...	72B	HBC	$3.9, 4.6 K^- \rho$

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
VALUE				
0.10 ± 0.03	¹⁸ PROKOSHKIN	91	GAM4	$300 \pi^- \rho \rightarrow \pi^- \rho \eta \eta$

¹⁸ Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production and results of CBAL, MRK3 and DM2 on $J/\psi \rightarrow \gamma \eta \eta$.

See key on page 1343

Meson Full Listings

$f_2'(1525)$, $f_0(1525)$, $f_0(1590)$

 $f_2'(1525)$ REFERENCES

PROKOSHKIN 91	SPD 316 155	(GAM2 Collab.)
ALBRECHT 90G	ZPHY C48 183	+Ehrlichmann, Harder+ (ARGUS Collab.)
PDG	PL B239	+Hernandez, Stone, Porter+ (IFIC, BOST. CIT+)
BEHREND 89C	ZPHY C43 91	+Criegee, Dainton+ (CELLO Collab.)
ASTON 88D	NP B301 525	+Awaji, Bienz+ (SLAC, NAGO, CINC, INUS)
AUGUSTIN 88	PRL 60 2238	+Calcaterra+ (DM2 Collab.)
BERGER 88	ZPHY C37 329	+Genzel, Lackas+ (PLUTO Collab.)
FALVARD 88	PR D38 2706	+Ajaltouni+ (CLER, FRAS, LALO, PADO)
AUGUSTIN 87	ZPHY C36 369	+Cosme+ (LALO, CLER, FRAS, PADO)
BALTRUSAIT... 87	PR D35 2077	+Baltrusaitis, Coffman, Dubois+ (Mark III Collab.)
AIHARA 86B	PRL 57 404	+Alston-Garnjost+ (TPC-2 γ Collab.)
LONGACRE 86	PL B177 223	+Etkin+ (BNL, BRAN, CUNY, DUKE, NDAM)
ALTHOFF 83	PL 121B 216	+Brandelik, Boerner, Burkhardt+ (TASSO Collab.)
ARMSTRONG 83B	NP B224 193	+ (BARI, BIRM, CERN, MILA, CURIN+)
AGUILAR... 81B	ZPHY C8 313	+Aguilar-Benitez, Albajar+ (CERN, CDEF, MADR+)
ALHARRAN 81	NP B191 26	+Baubillier+ (BIRM, CERN, GLAS, MICH, CURIN)
CHABAUD 81	APP B12 575	+Niczyporuk, Becker+ (CERN, CRAC, MPIM)
COSTA... 80	NP B175 402	+Costa De Beauregard+ (BARI, BONN, CERN+)
GORLICH 80	NP B174 16	+Niczyporuk+ (CRAC, MPIM, CERN, ZEEM)
CORDEN 79	NP B157 250	+Dowell, Garvey+ (BIRM, RHEL, TELA, LOWC) JP
MARTIN 79	NP B158 520	+Ozmutlu (DURH)
POLYCHRO... 79	PR D19 1317	+Polychronakos, Cason, Bishop+ (NDAM, ANL)
BARREIRO 77	NP B121 237	+Diaz, Gay, Hemingway+ (CERN, AMST, NIJM, OXF)
EVANGELISTA 77	NP B127 384	+ (BARI, BONN, CERN, DARE, GLAS+)
PAWLICKI 77	PR D15 3196	+Ayres, Cohen, Diebold, Kramer, Wicklund (ANL) JJP
BRANDENB... 76C	NP B104 413	+Brandenburg, Carnegie, Cashmore+ (SLAC)
BEUSCH 75B	PL 60B 101	+Birman, Websdale, Wetzel (CERN, ETH)
AGUILAR... 72B	PR D6 29	+Aguilar-Benitez, Chung, Eisner, Samios (BNL)
AMMAR 67	PRL 19 1071	+Davis, Hwang, Dagan, Derrick+ (NWES, ANL) JP
BARNES 67	PRL 19 964	+Dornan, Goldberg, Leitner+ (BNL, SYRA) JJP
CRENNELL 66	PRL 16 1025	+Kalbfleisch, Lai, Scarr, Schumann+ (BNL) I

OTHER RELATED PAPERS

JENNI 83	PR D27 1031	+Burke, Telnov, Abrams, Blocker+ (SLAC, LBL)
ARMSTRONG 82	PL 110B 77	+Baubillier+ (BARI, BIRM, CERN, MILA, CURIN+)
ETKIN 82B	PR D25 1786	+Foley, Lai+ (BNL, CUNY, TUFTS, VAND)
LUKE 82	DESY 82/073	(DESY)
ABRAMS 67B	PRL 18 620	+Kehoe, Glasser, Sechi-Zorn, Wolsky (UMD)
BARNES 65	PRL 15 322	+Culwick, Guidoni, Kalbfleisch, Goz+ (BNL, SYRA)

 $f_0(1525)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

OMITTED FROM SUMMARY TABLE

This entry contains evidence for $K\bar{K}\pi\pi$ and $\eta\eta$ S-wave intensity peaking at the mass of the $f_2'(1525)$. Needs confirmation. **$f_0(1525)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1525 OUR ESTIMATE			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1520 \pm 35	¹ ANISOVICH 94	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
1560 \pm 25	AMSLER 93C	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
~ 1525	ASTON 88D	LASS	11 $K^-p \rightarrow K_S^0 K_S^0 \Lambda$
~ 1525	BAUBILLIER 83		8 $K^-p \rightarrow K^+ K^- \Lambda$

¹ From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$. Supersedes AMSLER 93C. **$f_0(1525)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
148 $^{+20}_{-25}$	² ANISOVICH 94	CBAR	0.0 $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$
245 \pm 50	AMSLER 93C	CBAR	0.0 $\bar{p}p \rightarrow \pi^0\eta\eta$
~ 90	BAUBILLIER 83		8 $K^-p \rightarrow K^+ K^- \Lambda$

² From a simultaneous analysis of the annihilations $\bar{p}p \rightarrow 3\pi^0, \pi^0\eta\eta$. Supersedes AMSLER 93C. **$f_0(1525)$ REFERENCES**

ANISOVICH 94	PL B323 233	+Armstrong+ (Crystal Barrel Collab.)
AMSLER 93C	NP A558 3C	+Augustin+ (Crystal Barrel Collab.)
ASTON 88D	NP B301 525	+Awaji, Bienz+ (SLAC, NAGO, CINC, INUS)
BAUBILLIER 83	ZPHY C17 309	+ (BIRM, CERN, GLAS, MSU, CURIN)

 $f_0(1590)$

$$I^G(J^{PC}) = 0^+(0^{++})$$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.) **$f_0(1590)$ MASS**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1581 \pm 10	OUR AVERAGE			
1560 \pm 25		¹ AMSLER 92	CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
1550 \pm 45 \pm 30		BELADIDZE 92C	VES	36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
1610 \pm 20		ALDE 88	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$
1570 \pm 20	600 \pm 70	ALDE 87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
1575.0 \pm 45.0		² ALDE 86D	GAM4	100 $\pi^- p \rightarrow 2\eta n$
1568.0 \pm 33.0		BINON 84C	GAM2	38 $\pi^- p \rightarrow \eta\eta' n$
1592.0 \pm 25.0		BINON 83	GAM2	38 $\pi^- p \rightarrow 2\eta n$

¹ The error is mostly systematic.² From central value and spread of two solutions. **$f_0(1590)$ WIDTH**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
180 \pm 17	OUR AVERAGE			Error includes scale factor of 1.2.
245 \pm 50		³ AMSLER 92	CBAR	0.0 $\bar{p}p \rightarrow \eta\eta\pi^0$
153 \pm 67 \pm 50		BELADIDZE 92C	VES	36 $\pi^- \text{Be} \rightarrow \pi^- \eta' \eta \text{Be}$
170 \pm 40		ALDE 88	GAM4	300 $\pi^- N \rightarrow \pi^- N 2\eta$
150 \pm 20	600 \pm 70	ALDE 87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$
265.0 \pm 65.0		⁴ ALDE 86D	GAM4	100 $\pi^- p \rightarrow 2\eta n$
260.0 \pm 60.0		BINON 84C	GAM2	38 $\pi^- p \rightarrow \eta\eta' n$
210.0 \pm 40.0		BINON 83	GAM2	38 $\pi^- p \rightarrow 2\eta n$

³ The error is mostly systematic.⁴ From central value and spread of two solutions. **$f_0(1590)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $\eta\eta'(958)$	dominant
Γ_2 $\eta\eta$	large
Γ_3 $4\pi^0$	large
Γ_4 $\pi^0\pi^0$	
Γ_5 $K\bar{K}$	

 $f_0(1590)$ BRANCHING RATIOS

$\Gamma(\eta\eta'(958))/\Gamma(\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
2.7 \pm 0.8	BINON 84C	GAM2	38 $\pi^- p \rightarrow \eta\eta' n$	

$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
large	ALDE 88	GAM4	300 $\pi^- N \rightarrow \eta\eta\pi^- N$	
large	BINON 83	GAM2	38 $\pi^- p \rightarrow 2\eta n$	

$\Gamma(4\pi^0)/\Gamma(\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
0.8 \pm 0.3	ALDE 87	GAM4	100 $\pi^- p \rightarrow 4\pi^0 n$	

$\Gamma(\pi^0\pi^0)/\Gamma(\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_2
< 0.17	90			
< 0.3	⁵ BINON 83	GAM2	38 $\pi^- p \rightarrow 2\eta n$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁵ Superseded by PROKOSHKIN 90.

$\Gamma(K\bar{K})/\Gamma(\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
< 0.6	BINON 83	GAM2	38 $\pi^- p \rightarrow 2\eta n$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁶ Combining results of GAM4 with those of WA76 on $K\bar{K}$ central production.

Meson Full Listings

$f_0(1590)$, $\omega(1600)$, $X(1600)$, $f_2(1640)$

$f_0(1590)$ REFERENCES

AMSLER	92	PL B291 347	+Augustin, Baker+	(Crystal Barrel Collab.)
BELADIDZE	92C	SJNP 55 1535	+Bityukov, Borisov	(VES Collab.) JPC
PROKOSHKIN	91	SPD 316 155	Translated from YAF 55 2748.	(GAM2 Collab.)
PROKOSHKIN	90	Hadron 89 Conf. p 27	Translated from DANS 316 900.	(SERP, BELG, LANL, LAPP, PISA, KEK)
ALDE	88	PL B201 160	+Bellazzini, Binon+	(SERP, BELG, LANL, LAPP, PISA) JP
ALDE	87	PL B198 286	+Binon, Bricman+	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	+Binon, Bricman+	(BELG, LAPP, SERP, CERN) IGJP
BINON	84C	NC 80A 363	+Bricman, Donskov+	(BELG, LAPP, SERP, CERN)
BINON	83	NC 78A 313	+Donskov, Dutell+	(BELG, LAPP, SERP, CERN) IGJP
Also	83B	SJNP 38 561	+Binon, Gouanere+	(BELG, LAPP, SERP, CERN)
			Translated from YAF 38 934.	

OTHER RELATED PAPERS

SLAUGHTER	88	MPL A3 1361	(LANL)
-----------	----	-------------	--------

$\omega(1600)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

See also $\omega(1420)$.

$\omega(1600)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1662 ± 13	435	¹ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1609 ± 20	315	² ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → $\rho\pi$
1663 ± 12	435	³ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → $\omega\pi\pi$
1594 ± 12		DONNACHIE	89	RVUE	e ⁺ e ⁻ → $\rho\pi$
1670 ± 20		ATKINSON	83B	OMEG	20-70 $\gamma\rho$ → $3\pi X$
1657 ± 13		CORDIER	81	DM1	e ⁺ e ⁻ → $\omega 2\pi$
1679 ± 34	21	ESPOSITO	80	FRAM	e ⁺ e ⁻ → 3π
1652.0 ± 17.0		COSME	79	OSPK 0	e ⁺ e ⁻ → 3π

¹ From a coupled fit of $\rho\pi$ and $\omega\pi\pi$ channels.

² From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

³ From a single Breit-Wigner fit.

$\omega(1600)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
280 ± 24	435	⁴ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •					
159 ± 43	315	⁵ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → $\rho\pi$
240 ± 25	435	⁶ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → $\omega\pi\pi$
100 ± 30		DONNACHIE	89	RVUE	e ⁺ e ⁻ → $\rho\pi$
160 ± 20		ATKINSON	83B	OMEG	20-70 $\gamma\rho$ → $3\pi X$
136 ± 46		CORDIER	81	DM1	e ⁺ e ⁻ → $\omega 2\pi$
99 ± 49	21	ESPOSITO	80	FRAM	e ⁺ e ⁻ → 3π
42.0 ± 17.0		COSME	79	OSPK 0	e ⁺ e ⁻ → 3π

⁴ From a coupled fit of $\rho\pi$ and $\omega\pi\pi$ channels.

⁵ From a fit to two Breit-Wigner functions interfering between them and with the ω, ϕ tails with fixed (+, -, +) phases.

⁶ From a single Breit-Wigner fit.

$\omega(1600)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi$	seen
Γ_2 $\omega\pi\pi$	seen
Γ_3 e^+e^-	seen

$\omega(1600)$ $\Gamma(l)\Gamma(e^+e^-)/\Gamma(\text{total})$

$\Gamma(\rho\pi) \times \Gamma(e^+e^-)/\Gamma(\text{total})$	VALUE (eV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_3/\Gamma$
134 ± 14	435	⁷ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
93 ± 27	315	ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → $\rho\pi$	
96 ± 35		DONNACHIE	89	RVUE	e ⁺ e ⁻ → $\rho\pi$	

⁷ From a coupled fit of $\rho\pi$ and $\omega\pi\pi$ channels.

$\Gamma(\omega\pi\pi) \times \Gamma(e^+e^-)/\Gamma(\text{total})$

VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_3/\Gamma$	
170 ± 17	435	⁸ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
135 ± 16	435	⁹ ANTONELLI	92	DM2	1.34-2.4e ⁺ e ⁻ → $\omega\pi\pi$	
56 ± 31		DONNACHIE	89	RVUE	e ⁺ e ⁻ → $\omega 2\pi$	

⁸ From a coupled fit of $\rho\pi$ and $\omega\pi\pi$ channels.

⁹ From a single Breit-Wigner fit.

$\omega(1600)$ REFERENCES

ANTONELLI	92	ZPHY C56 15	+Baldini+	(DM2 Collab.)
DONNACHIE	89	ZPHY C42 663	+Clegg	(CERN, MCHS)
ATKINSON	83B	PL 127B 132	+	(BONN, CERN, GLAS, LANL, MCHS, CURIN+)
CORDIER	81	PL 106B 155	+Bisello, Bizot, Buon, Delcourt, Mane	(ORSAY)
ESPOSITO	80	LNC 28 195	+Marini, Patteri+	(FRAS, NAPL, PADO, ROMA)
COSME	79	NP B152 215	+Dudeltzak, Grelaud, Jean-Marie, Julian+	(IPN)

OTHER RELATED PAPERS

ATKINSON	87	ZPHY C34 157	+	(BONN, CERN, GLAS, LANL, MCHS, CURIN)
ATKINSON	84	NP B231 15	+	(BONN, CERN, GLAS, LANL, MCHS, CURIN+)

$X(1600)$

$$I^G(J^{PC}) = 2^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Observed in the reaction $\gamma\gamma \rightarrow \rho^0\rho^0$ near threshold. The large ratio of cross-sections $\sigma(\gamma\gamma \rightarrow \rho^0\rho^0) / \sigma(\gamma\gamma \rightarrow \rho^+\rho^-) \approx 4$ and the dominance of the $J^P = 2^+$ wave in the reaction $\gamma\gamma \rightarrow \rho^0\rho^0$ is a signature consistent with the production of an exotic ($I = 2$) resonance. Needs confirmation.

$X(1600)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1600 ± 100	¹ ALBRECHT	91F	ARG	0	10.2 e ⁺ e ⁻ → e ⁺ e ⁻ 2($\pi^+\pi^-$)

¹ Our estimate.

$X(1600)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
400 ± 200	² ALBRECHT	91F	ARG	0	10.2 e ⁺ e ⁻ → e ⁺ e ⁻ 2($\pi^+\pi^-$)

² Our estimate.

$X(1600)$ REFERENCES

ALBRECHT	91F	ZPHY C50 1	+Appuan, Paulini, Funk+	(ARGUS Collab.)
----------	-----	------------	-------------------------	-----------------

OTHER RELATED PAPERS

ALBRECHT	89M	PL B217 205	+Bockmann+	(ARGUS Collab.)
BEHREND	89D	PL B218 494	+Criegee+	(CELLO Collab.)

$f_2(1640)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

Needs confirmation.

$f_2(1640)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1647 ± 7	ADAMO	92	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
1590 ± 30	BELADIDZE	92B	VES $36 \pi^- p \rightarrow \omega\omega n$
1635 ± 7	ALDE	90	GAM2 $38 \pi^- p \rightarrow n\omega$

$f_2(1640)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
58 ± 20		ADAMO	92	OBLX $\bar{n}p \rightarrow 3\pi^+ 2\pi^-$
100 ± 20		BELADIDZE	92B	VES $36 \pi^- p \rightarrow \omega\omega n$
< 70	90	ALDE	90	GAM2 $38 \pi^- p \rightarrow n\omega$

$f_2(1640)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega\omega$	seen

See key on page 1343

Meson Full Listings

$f_2(1640), \omega_3(1670), \pi_2(1670)$

$f_2(1640)$ BRANCHING RATIOS

$\Gamma(\omega\omega)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
seen	ALDE	89B GAM2	38 $\pi^- \rho \rightarrow n\omega$	

$f_2(1640)$ REFERENCES

ADAMO	92	PL B287 368	+Agnello, Balestra+	(OBELIX Collab.)
BELADIDZE	92B	ZPHY C54 367	+Bityukov, Borisov+	(VES Collab.)
ALDE	90	PL B241 600	+Binon+	(SERP, BELG, LANL, LAPP, PISA, KEK)
ALDE	89B	PL B216 451	+Binon, Bricman+	(SERP, BELG, LANL, LAPP, TBIL)IGJPC

$\omega_3(1670)$

$$I^G(J^{PC}) = 0^-(3^-)$$

$\omega_3(1670)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1668 ± 5	OUR AVERAGE			
1685.0 ± 20.0	60	BAUBILLIER	79 HBC	8.2 $K^- p$ backward
1673.0 ± 12.0	430	1,2 BALTAY	78E HBC	15 $\pi^+ \rho \rightarrow \Delta 3\pi$
1650.0 ± 12.0		CORDEN	78B OMEG	8-12 $\pi^- \rho \rightarrow N 3\pi$
1669 ± 11	600	2 WAGNER	75 HBC	7 $\pi^+ \rho \rightarrow \Delta^{++} 3\pi$
1678 ± 14	500	DIAZ	74 DBC	6 $\pi^+ n \rightarrow \rho 3\pi^0$
1660 ± 13	200	DIAZ	74 DBC	6 $\pi^+ n \rightarrow \rho \omega \pi^0 \pi^0$
1679 ± 17	200	MATTHEWS	71D DBC	7.0 $\pi^+ n \rightarrow \rho 3\pi^0$
1670 ± 20		KENYON	69 DBC	8 $\pi^+ n \rightarrow \rho 3\pi^0$
~ 1700.0	110	1 CERRADA	77B HBC	4.2 $K^- p \rightarrow \Lambda 3\pi$
1695.0 ± 20.0		BARNES	69B HBC	4.6 $K^- p \rightarrow \omega 2\pi X$
1636 ± 20		ARMENISE	68B DBC	5.1 $\pi^+ n \rightarrow \rho 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 ~ 1700.0
 1 Phase rotation seen for $J^P = 3^- \rho \pi$ wave.
 2 From a fit to $I(J^P) = 0(3^-) \rho \pi$ partial wave.

$\omega_3(1670)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
173 ± 11	OUR AVERAGE			
160.0 ± 80.0	60	3 BAUBILLIER	79 HBC	8.2 $K^- p$ backward
173.0 ± 16.0	430	4,5 BALTAY	78E HBC	15 $\pi^+ \rho \rightarrow \Delta 3\pi$
253.0 ± 39.0		CORDEN	78B OMEG	8-12 $\pi^- \rho \rightarrow N 3\pi$
173 ± 28	600	3,5 WAGNER	75 HBC	7 $\pi^+ \rho \rightarrow \Delta^{++} 3\pi$
167 ± 40	500	DIAZ	74 DBC	6 $\pi^+ n \rightarrow \rho 3\pi^0$
122 ± 39	200	DIAZ	74 DBC	6 $\pi^+ n \rightarrow \rho \omega \pi^0 \pi^0$
155 ± 40	200	3 MATTHEWS	71D DBC	7.0 $\pi^+ n \rightarrow \rho 3\pi^0$
90 ± 20		BARNES	69B HBC	4.6 $K^- p \rightarrow \omega 2\pi$
100 ± 40		KENYON	69 DBC	8 $\pi^+ n \rightarrow \rho 3\pi^0$
112 ± 60		ARMENISE	68B DBC	5.1 $\pi^+ n \rightarrow \rho 3\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 3 Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
 4 Phase rotation seen for $J^P = 3^- \rho \pi$ wave.
 5 From a fit to $I(J^P) = 0(3^-) \rho \pi$ partial wave.

$\omega_3(1670)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \rho \pi$	seen
$\Gamma_2 \omega \pi \pi$	seen
$\Gamma_3 b_1(1235) \pi$	possibly seen

$\omega_3(1670)$ BRANCHING RATIOS

$\Gamma(\omega \pi \pi)/\Gamma(\rho \pi)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE				
0.71 ± 0.27	100	DIAZ	74 DBC	6 $\pi^+ n \rightarrow \rho 5\pi^0$

$\Gamma(b_1(1235)\pi)/\Gamma(\rho \pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
VALUE				
possibly seen	DIAZ	74 DBC	6 $\pi^+ n \rightarrow \rho 5\pi^0$	

$\Gamma(b_1(1235)\pi)/\Gamma(\omega \pi \pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
VALUE				
> 0.75	68	BAUBILLIER	79 HBC	8.2 $K^- p$ backward

$\omega_3(1670)$ REFERENCES

BAUBILLIER	79	PL 89B 131	+Cautis, Kalekar	(BIRM, CERN, GLAS, MSU, ORSAY)
BALTAY	78E	PRL 40 87	+Corbett, Alexander+	(COLU) JP
CORDEN	78B	NP B138 235	+Blockzji, Heinen+	(BIRM, RHEL, TELA, LOWE)
CERRADA	77B	NP B126 241	+Tabak, Chew	(AMST, CERN, NIJM, OXF) JP
WAGNER	75	PL 58B 201		(LBL) JP
DIAZ	74	PRL 32 260	+Dibianca, Fickinger, Anderson+	(CASE, CMU)
MATTHEWS	71D	PR D3 2561	+Prentice, Yoon, Carroll+	(TNTO, WISC)
BARNES	69B	PRL 23 142	+Chung, Eisner, Flaminio+	(BNL)
KENYON	69	PRL 23 146	+Kinson, Scarr+	(BNL, UCND, ORNL)
ARMENISE	68B	PL 26B 336	+Forino, Cartacci+	(BARI, BGNA, FIRZ, ORSAY)

OTHER RELATED PAPERS

MATTHEWS	71	LCN 1 361	+Prentice, Yoon, Carroll+	(TNTO, WISC)
ARMENISE	70	LCN 4 199	+Ghidini, Foring, Cartacci+	(BARI, BGNA, FIRZ)

$\pi_2(1670)$


$$I^G(J^{PC}) = 1^-(2^-)$$

Our latest mini-review on this particle can be found in the 1984 edition.

$\pi_2(1670)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1670 ± 20	OUR ESTIMATE				This is only an educated guess; the error given is larger than the error on the average of the published values.
1673 ± 7	OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
1710 ± 20	700 ± 150	ANTIPOV	87 SIGM	-	50 $\pi^- \text{Cu} \rightarrow \mu^+ \mu^- \pi^- \text{Cu}$
1676 ± 6		1 EVANGELISTA	81 OMEG	-	12 $\pi^- \rho \rightarrow 3\pi \rho$
1657.0 ± 14.0		1,2 DAUM	80D SPEC	-	63-94 $\pi \rho \rightarrow 3\pi X$
1662.0 ± 10.0	2000	1 BALTAY	77 HBC	+	15 $\pi^+ \rho \rightarrow \rho 3\pi$
1742 ± 31 ± 49		ANTREASNYAN	90 CBAL		$e^+ e^- \rightarrow \pi^0 \pi^0 \pi^0$ $e^+ e^- \rightarrow \pi^0 \pi^0 \pi^0$
1710.0 ± 20.0		3 DAUM	81B SPEC	-	63,94 $\pi^- \rho$
1640 ± 10	575	KALELKHAR	75 HBC	+	15 $\pi^+ \rho \rightarrow \rho \pi^+ f_2$
1660 ± 10		1 ASCOLI	73 HBC	-	5-25 $\pi^- \rho \rightarrow \rho \pi_2$

1 From a fit to $J^P = 2^- S$ -wave $f_2(1270) \pi$ partial wave.
 2 Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.
 3 From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.



$\pi_2(1670)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
240 ± 15	OUR AVERAGE				Error includes scale factor of 1.1.
170 ± 80	700 ± 150	ANTIPOV	87 SIGM	-	50 $\pi^- \text{Cu} \rightarrow \mu^+ \mu^- \pi^- \text{Cu}$
260 ± 20		4 EVANGELISTA	81 OMEG	-	12 $\pi^- \rho \rightarrow 3\pi \rho$
219.0 ± 20.0		4,5 DAUM	80D SPEC	-	63-94 $\pi \rho \rightarrow 3\pi X$
285.0 ± 60.0	2000	4 BALTAY	77 HBC	+	15 $\pi^+ \rho \rightarrow \rho 3\pi$
236 ± 49 ± 36		ANTREASNYAN	90 CBAL		$e^+ e^- \rightarrow \pi^0 \pi^0 \pi^0$ $e^+ e^- \rightarrow \pi^0 \pi^0 \pi^0$
312.0 ± 50.0		6 DAUM	81B SPEC	-	63,94 $\pi^- \rho$
240 ± 30	575	KALELKHAR	75 HBC	+	15 $\pi^+ \rho \rightarrow \rho \pi^+ f_2$
270 ± 60		4 ASCOLI	73 HBC	-	5-25 $\pi^- \rho \rightarrow \rho \pi_2$

4 From a fit to $J^P = 2^- f_2(1270) \pi$ partial wave.
 5 Clear phase rotation seen in $2^- S, 2^- P, 2^- D$ waves. We quote central value and spread of single-resonance fits to three channels.
 6 From a two-resonance fit to four $2^- 0^+$ waves. This should not be averaged with all the single resonance fits.

Meson Full Listings

 $\pi_2(1670)$ $\pi_2(1670)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $f_2(1270)\pi$	$(56.2 \pm 3.2)\%$
Γ_2 $\pi^\pm \pi^+ \pi^-$	$(53 \pm 4)\%$
Γ_3 $\rho\pi$	$(31 \pm 4)\%$
Γ_4 $f_0(1300)\pi$	$(8.7 \pm 3.4)\%$
Γ_5 $K\bar{K}^*(892) + \text{c.c.}$	$(4.2 \pm 1.4)\%$
Γ_6 $\gamma\gamma$	$(5.6 \pm 1.1) \times 10^{-6}$
Γ_7 $\eta\pi$	$< 5\%$
Γ_8 $\pi^\pm 2\pi^+ 2\pi^-$	$< 5\%$

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 6 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 1.9$ for 3 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-53		
x_4	-29	-59	
x_5	-8	-21	-9
	x_1	x_3	x_4

 $\pi_2(1670)$ PARTIAL WIDTHS


$\Gamma(\gamma\gamma)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6
VALUE (keV)					
1.35 ± 0.26 OUR AVERAGE					
1.41 ± 0.23 ± 0.28	ANTREASYAN 90	CBAL	0	$e^+e^- \rightarrow e^+e^-\pi^0\pi^0\pi^0$	
1.3 ± 0.3 ± 0.2	7 BEHREND 90C	CELL	0	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$	
••• We do not use the following data for averages, fits, limits, etc. •••					
0.8 ± 0.3 ± 0.12	8 BEHREND 90C	CELL	0	$e^+e^- \rightarrow e^+e^-\pi^+\pi^-\pi^0$	
⁷ Incoherent Ansatz.					
⁸ Constructive interference between $f_2(1270), \rho\pi$ and background.					

 $\pi_2(1670)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$	DOCUMENT ID	TECN	CHG	COMMENT
VALUE				
0.29 ± 0.04 OUR FIT				
0.29 ± 0.05	⁹ DAUM	81B	SPEC	63,94 π^-p
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.3	BARTSCH 68	HBC	+	8 $\pi^+p \rightarrow 3\pi p$
<0.4	FERBEL 68	RVUE	±	
⁹ From a two-resonance fit to four 2^{-0+} waves.				

$\Gamma(f_2(1270)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$	DOCUMENT ID	TECN	CHG	COMMENT
VALUE				
0.604 ± 0.035 OUR FIT				
0.60 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.61 ± 0.04	¹⁰ DAUM	81B	SPEC	63,94 π^-p
0.76 \pm 0.24 $-$ 0.34	ARMENISE 69	DBC	+	5.1 $\pi^+d \rightarrow d3\pi$
0.35 ± 0.20	BALTAY 68	HBC	+	7-8.5 π^+p
••• We do not use the following data for averages, fits, limits, etc. •••				
0.59	BARTSCH 68	HBC	+	8 $\pi^+p \rightarrow 3\pi p$
¹⁰ From a two-resonance fit to four 2^{-0+} waves.				

WEIGHTED AVERAGE
0.60 ± 0.05 (Error scaled by 1.3)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$	DOCUMENT ID	TECN	CHG	COMMENT
VALUE				
<0.09	BALTAY 68	HBC	+	7-8.5 π^+p
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.10	CRENNELL 70	HBC	-	6 $\pi^-p \rightarrow f_2\pi^-N$

$\Gamma(\pi^\pm 2\pi^+ 2\pi^-)/\Gamma(\pi^\pm\pi^+\pi^-)$	DOCUMENT ID	TECN	CHG	COMMENT
VALUE				
<0.10	CRENNELL 70	HBC	-	6 $\pi^-p \rightarrow f_2\pi^-N$
<0.1	BALTAY 68	HBC	+	7.8.5 π^+p

$\Gamma(f_0(1300)\pi)/\Gamma(\pi^\pm\pi^+\pi^-)$	DOCUMENT ID	TECN	COMMENT	
VALUE				
0.10 ± 0.04 OUR FIT				
0.10 ± 0.05	¹¹ DAUM	81B	SPEC 63,94 π^-p	
<0.1	BALTAY 68	HBC	+	7.8.5 π^+p
¹¹ From a two-resonance fit to four 2^{-0+} waves.				

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(f_2(1270)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_1
VALUE					
0.075 ± 0.025 OUR FIT					
0.075 ± 0.025	¹² ARMSTRONG 82B	OMEG	-	16 $\pi^-p \rightarrow K^+K^-\pi^-p$	
¹² From a partial-wave analysis of $K^+K^-\pi^-p$ system.					

D-wave/S-wave RATIO FOR $\pi_2(1670) \rightarrow f_2(1270)\pi$	DOCUMENT ID	TECN	COMMENT
VALUE			
0.22 ± 0.10	¹³ DAUM	81B	SPEC 63,94 π^-p
¹³ From a two-resonance fit to four 2^{-0+} waves.			

 $\pi_2(1670)$ REFERENCES

ANTREASYAN 90	ZPHY C48 561	+Bartels, Bessel-	(Crystal Ball Collab.)
BEHREND 90C	ZPHY C46 583	+Criegee+	(CELLO Collab.)
ANTIPOV 87	EPL 4 403	+Batarin+	(SERP, JINR, INRM, TBIL, BGNA, MILA)
ARMSTRONG 82B	NP B202 1	+Baccari	(AACH3, BARI, BONN, CERN, GLAS+)
DAUM 81B	NP B182 269	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)
EVANGELISTA 81	NP B178 197	+	(BARI, BONN, CERN, DARE, LIVP+)
Also	NP B186 594	Evangelista	
DAUM 80D	PL 89B 285	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)
BALTAY 77	PRL 39 591	+Cautis, Kalelkar	(COLU) JP
KALELKAR 75	Nevis 207 Thesis		(COLU)
ASCOLI 73	PR D7 669	(ILL, TINTO, GENO, HAMB, MILA, SACL) JP	
CRENNELL 70	PRL 24 781	+Karshon, Lai, Scarr, Sims	(BNL)
ARMENISE 69	LCN 2 501	+Ghidini, Forino, Cartacci+	(BARI, BGNA, FIRZ)
BALTAY 68	PRL 20 887	+Kung, Yeh, Ferbel+	(COLU, ROCH, RUTG, YALE)
BARTSCH 68	NP B7 345	+Keppel, Kraus+	(AACH, BERL, CERN) JP
FERBEL 68	Phil. Conf. 335		(ROCH)

OTHER RELATED PAPERS

CHEN 83B	PR D38 2304	+Fenker+	(ARIZ, FNAL, FLOR, NDAM, TUFTS+)
LEEDOM 83	PR D27 1426	+DeBonte, Gaidos, Key, Wong+	(PURD, TINTO)
BELLINI 82B	NP B199 1	+	(CERN, MILA, JINR, BGNA, HELS, PAVI, WARS+)
FOCACCI 66	PRL 17 890	+Kienzle, Levrat, Maglich, Martin	(CERN)
LEVRAT 66	PL 22 714	+Tolstrup+	(CERN Missing Mass Spect. Collab.)
LUBATTI 66	Berkeley Thesis		(LRL)
VETLITSKY 66	PL 21 579	+Guszavin, Kliger, Zolgonov+	(ITEP)
FORINO 65B	PL 19 68	+Gessaroli+	(BGNA, BARI, FIRZ, ORSAY, SACL)

See key on page 1343

Meson Full Listings

 $\phi(1680), \rho_3(1690)$ $\phi(1680)$

$$I^G(J^{PC}) = 0^-(1^{--})$$

First identified using Dalitz plot analysis of $e^+e^- \rightarrow K\bar{K}^*(892)$ (BIZOT 80, DELCOURT 81). We do not list anymore ω radial excitations under this particle. See also $\omega(1420)$ and $\omega(1600)$.

 $\phi(1680)$ MASS e^+e^- PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1680 ± 50 OUR ESTIMATE		This is only an educated guess; the error given is larger than the error on the average of the published values.		

• • • We do not use the following data for averages, fits, limits, etc. • • •

1657 ± 27	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
1655 ± 17		¹ BISELLO	88B DM2	$e^+e^- \rightarrow K^+ K^-$
1680 ± 10		² BUON	82 DM1	$e^+e^- \rightarrow \text{hadrons}$
1677 ± 12		³ MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K \pi$

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1726 ± 22	BUSENITZ 89 TPS	$\gamma p \rightarrow K^+ K^- X$	
1760 ± 20	ATKINSON 85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$	
1690 ± 10	ASTON 81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1726 ± 22	BUSENITZ 89 TPS	$\gamma p \rightarrow K^+ K^- X$	
1760 ± 20	ATKINSON 85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$	
1690 ± 10	ASTON 81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$	

¹ From global fit including ρ, ω, ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.

² From global fit of ρ, ω, ϕ and their radial excitations to channels $\omega \pi^+ \pi^-, K^+ K^-, K_S^0 K_L^0, K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

³ Fit to one channel only, neglecting interference with $\omega, \rho(1700)$.

 $\phi(1680)$ WIDTH e^+e^- PRODUCTION

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
150 ± 50 OUR ESTIMATE		This is only an educated guess; the error given is larger than the error on the average of the published values.		

• • • We do not use the following data for averages, fits, limits, etc. • • •

146 ± 55	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$
207 ± 45		⁴ BISELLO	88B DM2	$e^+e^- \rightarrow K^+ K^-$
185 ± 22		⁵ BUON	82 DM1	$e^+e^- \rightarrow \text{hadrons}$
102 ± 36		⁶ MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K \pi$

PHOTOPRODUCTION

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
121 ± 47	BUSENITZ 89 TPS	$\gamma p \rightarrow K^+ K^- X$	
80 ± 40	ATKINSON 85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$	
100 ± 40	ASTON 81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

121 ± 47	BUSENITZ 89 TPS	$\gamma p \rightarrow K^+ K^- X$	
80 ± 40	ATKINSON 85C OMEG	20-70 $\gamma p \rightarrow K \bar{K} X$	
100 ± 40	ASTON 81F OMEG	25-70 $\gamma p \rightarrow K^+ K^- X$	

⁴ From global fit including ρ, ω, ϕ and $\rho(1700)$ assume mass 1570 MeV and width 510 MeV for ρ radial excitation.

⁵ From global fit of ρ, ω, ϕ and their radial excitations to channels $\omega \pi^+ \pi^-, K^+ K^-, K_S^0 K_L^0, K_S^0 K^\pm \pi^\mp$. Assume mass 1570 MeV and width 510 MeV for ρ radial excitations, mass 1570 and width 500 MeV for ω radial excitation.

⁶ Fit to one channel only, neglecting interference with $\omega, \rho(1700)$.

 $\phi(1680)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \bar{K}^*(892) + \text{c.c.}$	dominant
Γ_2 $K_S^0 K \pi$	seen
Γ_3 $K \bar{K}$	seen
Γ_4 $e^+ e^-$	seen
Γ_5 $\omega \pi \pi$	not seen
Γ_6 $K^+ K^- \pi^0$	

 $\phi(1680)$ $\Gamma(I)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel (I) in e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(I)$ or the branching ratio $\Gamma(I)/\text{total}$.

$\Gamma(K \bar{K}^*(892) + \text{c.c.}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	$\Gamma_1 \Gamma_4/\Gamma$			
VALUE (keV)	EVTS	DOCUMENT ID	TECN	COMMENT
0.48 ± 0.14	367	BISELLO	91C DM2	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\phi(1680)$ BRANCHING RATIOS

$\Gamma(K \bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K \pi)$	Γ_1/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
dominant	MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K \bar{K})/\Gamma(K \bar{K}^*(892) + \text{c.c.})$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.07 ± 0.01	BUON	82 DM1	e^+e^-

$\Gamma(\omega \pi \pi)/\Gamma(K \bar{K}^*(892) + \text{c.c.})$	Γ_5/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
<0.10	BUON	82 DM1	e^+e^-

$\Gamma(K \bar{K}^*(892) + \text{c.c.})/\Gamma(K_S^0 K \pi)$	Γ_1/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
dominant	MANE	82 DM1	$e^+e^- \rightarrow K_S^0 K^\pm \pi^\mp$

$\Gamma(K \bar{K})/\Gamma(K \bar{K}^*(892) + \text{c.c.})$	Γ_3/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
0.07 ± 0.01	BUON	82 DM1	e^+e^-

$\Gamma(\omega \pi \pi)/\Gamma(K \bar{K}^*(892) + \text{c.c.})$	Γ_5/Γ_1		
VALUE	DOCUMENT ID	TECN	COMMENT
<0.10	BUON	82 DM1	e^+e^-

 $\phi(1680)$ REFERENCES

BISELLO 91C	ZPHY C52 227	+Busetto, Castro, Nigro, Pescara+	(DM2 Collab.)
BUSENITZ 89	PR D40 1	+Olzszewski, Callahan+	(ILL, FNAL)
BISELLO 88B	ZPHY C39 13	+Busetto+	(PADO, CLER, FRAS, LALO)
ATKINSON 85C	ZPHY C27 233	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
BUON 82	PL 118B 221	+Bisello, Bizot, Cordier, Delcourt+	(LALO, MONP)
MANE 82	PL 112B 178	+Bisello, Bizot, Buon, Delcourt, Fayard+	(LALO)
ASTON 81F	PL 104B 231	+ (BONN, CERN, EPOL, GLAS, LANC, MCHS+)	
DELCOURT 81	PL 99B 257	+Bisello, Bizot, Buon, Cordier, Mane	(ORSAY)
BIZOT 80	Madison Conf. 546	+Bisello, Buon, Cordier, Delcourt+	(LALO, MONP)

OTHER RELATED PAPERS

ATKINSON 86C	ZPHY C30 541	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ATKINSON 84	NP B231 15	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ATKINSON 84B	NP B231 1	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ATKINSON 83C	NP B229 269	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
CORDIER 81	PL 106B 155	+Bisello, Bizot, Buon, Delcourt, Mane	(ORSAY)
MANE 81	PL 99B 261	+Bisello, Bizot, Buon, Cordier, Delcourt	(ORSAY)
ASTON 80F	NP B174 269	+ (BONN, CERN, EPOL, GLAS, LANC, MCHS+)	

 $\rho_3(1690)$

$$I^G(J^{PC}) = 1^+(3^{--})$$

 $\rho_3(1690)$ MASS

We include only high statistics experiments in the average for the $2\pi, K \bar{K}$, and $K \bar{K} \pi$ modes.

 $2\pi, K \bar{K}$, AND $K \bar{K} \pi$ MODES

VALUE (MeV)	DOCUMENT ID
1691 ± 5 OUR ESTIMATE	This is only an educated guess; the error given is larger than the error on the average of the published values.
1691.4 ± 2.7 OUR AVERAGE	Includes data from the 2 datablocks that follow this one.

 2π MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
The data in this block is included in the average printed for a previous datablock.					

1686 ± 4 OUR AVERAGE

1677 ± 14		EVANGELISTA 81	OMEG	-	12 $\pi^- p \rightarrow 2\pi$
1679.0 ± 11.0	476	BALTAY 78B	HBC	0	15 $\pi^- p \rightarrow \pi^+ \pi^- n$
1678.0 ± 12.0	175	¹ ANTIPOV 77	CIBS	0	25 $\pi^- p \rightarrow \rho 3\pi$
1690 ± 7	600	¹ ENGLER 74	DBC	0	6 $\pi^+ n \rightarrow \pi^+ \pi^- p$
1693 ± 8		² GRAYER 74	ASPK	0	17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1678 ± 12		MATTHEWS 71C	DBC	0	7 $\pi^+ N$
1734.0 ± 10.0		³ CORDEN 79	OMEG		12-15 $\pi^- p \rightarrow n 2\pi$
1692 ± 12		^{2,4} ESTABROOKS 75	RVUE		17 $\pi^- p \rightarrow \pi^+ \pi^- n$
1737.0 ± 23.0		ARMENISE 70	DBC	0	9 $\pi^+ N$
1650.0 ± 35.0	122	BARTSCH 70B	HBC	+	8 $\pi^+ p \rightarrow N 2\pi$
1687 ± 21		STUNTEBECK 70	HDBC	0	8 $\pi^- p, 5.4 \pi^+ d$
1683 ± 13		ARMENISE 68	DBC	0	5.1 $\pi^+ d$
1670.0 ± 30.0		GOLDBERG 65	HBC	0	6 $\pi^+ d, 8 \pi^- p$

¹ Mass errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

² Uses same data as HYAMS 75.

³ From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K \bar{K}$ result.

⁴ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.

Meson Full Listings

$\rho_3(1690)$

$K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
The data in this block is included in the average printed for a previous datablock.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1696 ± 4 OUR AVERAGE					
1699.0 ± 5.0		ALPER	80 CNTR	0	$62 \pi^- p \rightarrow K^+ K^- n$
1698 ± 12	6k 5,6	MARTIN	78D SPEC		$10 \pi p \rightarrow K_S^0 K^- p$
1692 ± 6		BLUM	75 ASPK	0	$18.4 \pi^- p \rightarrow \eta K^+ K^-$
1690.0 ± 16.0		ADERHOLZ	69 HBC	+	$8 \pi^+ p \rightarrow K\bar{K}\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1694.0 ± 8.0	7	COSTA...	80 OMEG		$10 \pi^- p \rightarrow K^+ K^- n$


⁵ From a fit to $J^P = 3^-$ partial wave.
⁶ Systematic error on mass scale subtracted.
⁷ They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

$(4\pi)^\pm$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
1675 ± 11 OUR AVERAGE Error includes scale factor of 1.9. See the Ideogram below.

1665.0 ± 15.0	177	BALTAY	78B HBC	+	$15 \pi^+ p \rightarrow p4\pi$
1670 ± 10		THOMPSON	74 HBC	+	$13 \pi^+ p$
1687 ± 20		CASON	73 HBC	-	$8,18.5 \pi^- p$
1630 ± 15		HOLMES	72 HBC	+	$10-12 K^+ p$
1680.0 ± 40.0	144	BARTSCH	70B HBC	+	$8 \pi^+ p \rightarrow N4\pi$
1705.0 ± 21.0		CASO	70 HBC	-	$11.2 \pi^- p \rightarrow n\rho 2\pi$
1720 ± 15		BALTAY	68 HBC	+	$7, 8.5 \pi^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1694 ± 6	8	EVANGELISTA	81 OMEG	-	$12 \pi^- p \rightarrow p4\pi$
1718 ± 10	9	EVANGELISTA	81 OMEG	-	$12 \pi^- p \rightarrow p4\pi$
1673 ± 9	10	EVANGELISTA	81 OMEG	-	$12 \pi^- p \rightarrow p4\pi$
1733 ± 9	66	KLIGER	74 HBC	-	$4.5 \pi^- p \rightarrow p4\pi$
1685 ± 14	11	CASON	73 HBC	-	$8,18.5 \pi^- p$
1689.0 ± 20.0	102	BARTSCH	70B HBC	+	$8 \pi^+ p \rightarrow N2\rho$

⁸ From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.
⁹ From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.
¹⁰ From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.
¹¹ From $\rho^\pm \rho^0$ mode.



$\rho_3(1690)$ mass, $(4\pi)^\pm$ mode (MeV)

$\omega\pi$ MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT

1661 ± 6 OUR AVERAGE					
1690 ± 20		ALDE	92C GAM2		$38,100 \pi^- p \rightarrow \omega\pi^0 n$
1690 ± 15		EVANGELISTA	81 OMEG	-	$12 \pi^- p \rightarrow \omega\pi p$
1666.0 ± 14.0		GESSAROLI	77 HBC		$11 \pi^- p \rightarrow \omega\pi p$
1686 ± 9		THOMPSON	74 HBC	+	$13 \pi^+ p$
1654 ± 24		BARNHAM	70 HBC	+	$10 K^+ p \rightarrow \omega\pi X$

$\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1680 ± 15		FUKUI	88 SPEC	0	$8.95 \pi^- p \rightarrow \eta\pi^+\pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1700.0 ± 47.0		12 ANDERSON	69 MMS	-	$16 \pi^- p$ backward
1632 ± 15		12,13 FOCACCI	66 MMS	-	$7-12 \pi^- p \rightarrow pMM$
1700 ± 15		12,13 FOCACCI	66 MMS	-	$7-12 \pi^- p \rightarrow pMM$
1748 ± 15		12,13 FOCACCI	66 MMS	-	$7-12 \pi^- p \rightarrow pMM$

¹² Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+2\pi^-$, with 0, 1, 2 $\pi^+\pi^-$ pairs in ρ band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1976)
¹³ Not seen by BOWEN 72.


$\rho_3(1690)$ WIDTH

We include only high statistics experiments in the average for the $2\pi, K\bar{K}, K\bar{K}\pi$ modes.

$2\pi, K\bar{K},$ AND $K\bar{K}\pi$ MODES

VALUE (MeV) EVTS DOCUMENT ID
215 ± 20 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

215 ± 6 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.8. See the Ideogram below.





$\rho_3(1690)$ width, $2\pi, K\bar{K},$ and $K\bar{K}\pi$ modes (MeV)

2π MODE

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
The data in this block is included in the average printed for a previous datablock.

186 ± 14 OUR AVERAGE					
220 ± 29		DENNEY	83 LASS		$10 \pi^+ N$
246 ± 37		EVANGELISTA	81 OMEG	-	$12 \pi^- p \rightarrow 2\pi p$
116.0 ± 30.0	476	BALTAY	78B HBC	0	$15 \pi^+ p \rightarrow \pi^+\pi^- n$
162.0 ± 50.0	175	14 ANTIPOV	77 CIBS	0	$25 \pi^- p \rightarrow \rho 3\pi$
167 ± 40	600	ENGLER	74 DBC	0	$6 \pi^+ n \rightarrow \pi^+\pi^- p$
200 ± 18		15 GRAY	74 ASPK	0	$17 \pi^- p \rightarrow \pi^+\pi^- n$
156 ± 36		MATTHEWS	71C DBC	0	$7 \pi^+ N$
171.0 ± 65.0		ARMENISE	70 DBC	0	$9 \pi^+ d$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
322.0 ± 35.0		16 CORDEN	79 OMEG		$12-15 \pi^- p \rightarrow n2\pi$
240 ± 30		15,17 ESTABROOKS	75 RVUE		$17 \pi^- p \rightarrow \pi^+\pi^- n$
180.0 ± 30.0	122	BARTSCH	70B HBC	+	$8 \pi^+ p \rightarrow N2\pi$
267 \pm $\frac{+72}{-46}$		STUNTEBECK	70 HBC	0	$8 \pi^- p, 5.4 \pi^+ d$
188 ± 49		ARMENISE	68 DBC	0	$5.1 \pi^+ d$
180.0 ± 40.0		GOLDBERG	65 HBC	0	$6 \pi^+ d, 8 \pi^- p$

¹⁴ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
¹⁵ Uses same data as HYAMS 75 and BECKER 79.
¹⁶ From a phase shift solution containing a $f_2'(1525)$ width two times larger than the $K\bar{K}$ result.
¹⁷ From phase-shift analysis. Error takes account of spread of different phase-shift solutions.



$K\bar{K}$ AND $K\bar{K}\pi$ MODES

VALUE (MeV) EVTS DOCUMENT ID TECN CHG COMMENT
 The data in this block is included in the average printed for a previous datablock.

Value (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
218 ± 4 OUR AVERAGE					
219.0 ± 4.0		ALPER 80 CNTR	0		$62 \pi^- \rho \rightarrow K^+ K^- n$
199 ± 40	6000	18 MARTIN 78D	SPEC		$10 \pi \rho \rightarrow K_S^0 K^- \rho$
205 ± 20		BLUM 75 ASPK	0		$18.4 \pi^- \rho \rightarrow n K^+ K^-$
186.0 ± 11.0		19 COSTA... 80 OMEG			$10 \pi^- \rho \rightarrow K^+ K^- n$
112.0 ± 60.0		ADERHOLZ 69 HBC	+		$8 \pi^+ \rho \rightarrow K\bar{K}\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 18 From a fit to $J^P = 3^-$ partial wave.
 19 They cannot distinguish between $\rho_3(1690)$ and $\omega_3(1670)$.

$(4\pi)^\pm$ MODE

Value (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
119 ± 13 OUR AVERAGE					
105.0 ± 30.0	177	BALTAY 78B	HBC	+	$15 \pi^+ \rho \rightarrow \rho 4\pi$
106 ± 25		THOMPSON 74	HBC	+	$13 \pi^+ \rho$
169 +70 -48		CASON 73	HBC	-	$8.18.5 \pi^- \rho$
130 ± 30		HOLMES 72	HBC	+	$10-12 K^+ \rho$
135.0 ± 30.0	144	BARTSCH 70B	HBC	+	$8 \pi^+ \rho \rightarrow N 4\pi$
100 ± 35		BALTAY 68	HBC	+	$7.8.5 \pi^+ \rho$
123 ± 13		20 EVANGELISTA 81 OMEG	-		$12 \pi^- \rho \rightarrow \rho 4\pi$
230 ± 28		21 EVANGELISTA 81 OMEG	-		$12 \pi^- \rho \rightarrow \rho 4\pi$
184 ± 33		22 EVANGELISTA 81 OMEG	-		$12 \pi^- \rho \rightarrow \rho 4\pi$
150	66	23 KLIGER 74	HBC	-	$4.5 \pi^- \rho \rightarrow \rho 4\pi$
125 +83 -35		23 CASON 73	HBC	-	$8.18.5 \pi^- \rho$
180.0 ± 30.0	90	23 BARTSCH 70B	HBC	+	$8 \pi^+ \rho \rightarrow N a_2 \pi$
160.0 ± 30.0	102	BARTSCH 70B	HBC	+	$8 \pi^+ \rho \rightarrow N 2\rho$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 20 From $\rho^- \rho^0$ mode, not independent of the other two EVANGELISTA 81 entries.
 21 From $a_2(1320)^- \pi^0$ mode, not independent of the other two EVANGELISTA 81 entries.
 22 From $a_2(1320)^0 \pi^-$ mode, not independent of the other two EVANGELISTA 81 entries.
 23 From $\rho^\pm \rho^0$ mode.

$\omega\pi$ MODE

Value (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
150 ± 31 OUR AVERAGE	Error			Includes scale factor of 1.8. See the ideogram below.
230 ± 30	ALDE 92C	GAM2		$38,100 \pi^- \rho \rightarrow \omega \pi^0 n$
190 ± 65	EVANGELISTA 81	OMEG	-	$12 \pi^- \rho \rightarrow \omega \pi \rho$
160.0 ± 56.0	GESSAROLI 77	HBC		$11 \pi^- \rho \rightarrow \omega \pi \rho$
89 ± 25	THOMPSON 74	HBC	+	$13 \pi^+ \rho$
130 +73 -43	BARNHAM 70	HBC	+	$10 K^+ \rho \rightarrow \omega \pi X$

$\eta\pi^+\pi^-$ MODE

(For difficulties with MMS experiments, see the $a_2(1320)$ mini-review in the 1973 edition.)

Value (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
106 ± 27	FUKUI 88	SPEC	0	$8.95 \pi^- \rho \rightarrow \eta \pi^+ \pi^- n$
195.0	24 ANDERSON 69	MMS	-	$16 \pi^- \rho$ backward
< 21	24,25 FOCACCI 66	MMS	-	$7-12 \pi^- \rho \rightarrow \rho MM$
< 30	24,25 FOCACCI 66	MMS	-	$7-12 \pi^- \rho \rightarrow \rho MM$
< 38	24,25 FOCACCI 66	MMS	-	$7-12 \pi^- \rho \rightarrow \rho MM$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 24 Seen in 2.5-3 GeV/c $\bar{p}p$. $2\pi^+ 2\pi^-$, with 0, 1, 2 $\pi^+ \pi^-$ pairs in ρ^0 band not seen by OREN 74 (2.3 GeV/c $\bar{p}p$) with more statistics. (Jan. 1979)
 25 Not seen by BOWEN 72.

$\rho_3(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 4π	(71.1 ± 1.9) %	
Γ_2 $\pi^\pm \pi^+ \pi^- \pi^0$	(67 ± 22) %	
Γ_3 $\pi\pi$	(23.6 ± 1.3) %	
Γ_4 $\omega\pi$	(16 ± 6) %	
Γ_5 $K\bar{K}\pi$	(3.8 ± 1.2) %	
Γ_6 $K\bar{K}$	(1.58 ± 0.26) %	1.2
Γ_7 $\eta\pi^+\pi^-$	seen	
Γ_8 $\pi\pi\rho$	Excluding 2ρ and $a_2(1320)\pi$.	
Γ_9 $a_2(1320)\pi$		
Γ_{10} $\rho\rho$		
Γ_{11} $\phi\pi$		
Γ_{12} $\eta\pi$		
Γ_{13} $\pi^\pm 2\pi^+ 2\pi^- \pi^0$		

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 10 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 14.7$ for 7 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-77		
x_5	-74	17	
x_6	-15	2	0
	x_1	x_3	x_5

Meson Full Listings

 $\rho_3(1690)$ $\rho_3(1690)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ
VALUE					
0.236±0.013 OUR FIT					
0.243±0.013 OUR AVERAGE					
0.259 ^{+0.018} _{-0.019}	BECKER	79	ASPK	0	17 $\pi^- p$ polarized
0.23 ± 0.02	CORDEN	79	OMEG		12-15 $\pi^- p \rightarrow$
0.22 ± 0.04	26 MATTHEWS	71c	HDBC	0	7 $\pi^+ n \rightarrow \pi^- p$

••• We do not use the following data for averages, fits, limits, etc. •••

0.245±0.006

27 ESTABROOKS 75 RVUE 17 $\pi^- p \rightarrow \pi^+ \pi^- n$

²⁶ One-pion-exchange model used in this estimation.

²⁷ From phase-shift analysis of HYAMS 75 data.

$\Gamma(\pi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_2
VALUE					
0.35±0.11	CASON	73	HBC	-	8,18.5 $\pi^- p$


••• We do not use the following data for averages, fits, limits, etc. •••

<0.2	HOLMES	72	HBC	+	10-12 $K^+ p$
<0.12	BALLAM	71B	HBC	-	16 $\pi^- p$

$\Gamma(\pi\pi)/\Gamma(4\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_1
VALUE					
0.332±0.026 OUR FIT	Error includes scale factor of 1.1.				
0.30 ± 0.10	BALTAY	78B	HBC	0	15 $\pi^+ p \rightarrow p4\pi$

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ_3
VALUE					
0.067±0.011 OUR FIT	Error includes scale factor of 1.2.				
0.118^{+0.039}_{-0.032} OUR AVERAGE	Error includes scale factor of 1.7. See the ideogram below.				
0.191 ^{+0.040} _{-0.037}	GORLICH	80	ASPK	0	17,18 $\pi^- p$ polarized
0.08 ± 0.03	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.08 ^{+0.08} _{-0.03}	CRENNELL	68B	HBC		6.0 $\pi^- p$

WEIGHTED AVERAGE
0.118±0.039-0.032 (Error scaled by 1.7)



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

$\Gamma(K\bar{K})/\Gamma(\pi\pi)$

$\Gamma(K\bar{K}\pi)/\Gamma(\pi\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_3
VALUE					
0.16±0.05 OUR FIT					
0.16±0.05	28 BARTSCH	70B	HBC	+	8 $\pi^+ p$

²⁸ Increased by us to correspond to $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.

$[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	$(\Gamma_8+\Gamma_9+\Gamma_{10})/\Gamma_2$
VALUE					
0.94±0.09 OUR AVERAGE					
0.96±0.21	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.88±0.15	BALLAM	71B	HBC	-	16 $\pi^- p$
1 ± 0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
consistent with 1	CASO	68	HBC	-	11 $\pi^- p$

$\Gamma(\rho\rho)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{10}/Γ_2
VALUE					
EVTS					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.12±0.11	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.56	66 KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$
0.13±0.09	29 THOMPSON	74	HBC	+	13 $\pi^+ p$
0.7 ± 0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$

²⁹ $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

$\Gamma(\rho\rho)/[\Gamma(\pi\pi\rho) + \Gamma(a_2(1320)\pi) + \Gamma(\rho\rho)]$	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_{10}/(\Gamma_8+\Gamma_9+\Gamma_{10})$
VALUE					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.48±0.16	CASO	68	HBC	-	11 $\pi^- p$

$\Gamma(a_2(1320)\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_9/Γ_2
VALUE					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.66±0.08	BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
0.36±0.14	30 THOMPSON	74	HBC	+	13 $\pi^+ p$
not seen	CASON	73	HBC	-	8,18.5 $\pi^- p$
0.6 ± 0.15	BARTSCH	70B	HBC	+	8 $\pi^+ p$
0.6	BALTAY	68	HBC	+	7,8.5 $\pi^+ p$

³⁰ $\rho\rho$ and $a_2(1320)\pi$ modes are indistinguishable.

$\Gamma(\omega\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_2
VALUE					
CL%					
0.23±0.05 OUR AVERAGE	Error includes scale factor of 1.2.				
0.33±0.07	THOMPSON	74	HBC	+	13 $\pi^+ p$
0.12±0.07	BALLAM	71B	HBC	-	16 $\pi^- p$
0.25±0.10	BALTAY	68	HBC	+	7,8.5 $\pi^+ p$
0.25±0.10	JOHNSTON	68	HBC	-	7.0 $\pi^- p$

••• We do not use the following data for averages, fits, limits, etc. •••

<0.11	95 BALTAY	78B	HBC	+	15 $\pi^+ p \rightarrow p4\pi$
<0.09	KLIGER	74	HBC	-	4.5 $\pi^- p \rightarrow p4\pi$

$\Gamma(\phi\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{11}/Γ_2
VALUE					
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.11	BALTAY	68	HBC	+	7,8.5 $\pi^+ p$

$\Gamma(\pi^\pm 2\pi^+ 2\pi^- \pi^0)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{13}/Γ_2
VALUE					
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.15	BALTAY	68	HBC	+	7,8.5 $\pi^+ p$

$\Gamma(\eta\pi)/\Gamma(\pi^\pm\pi^+\pi^-\pi^0)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{12}/Γ_2
VALUE					
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.02	THOMPSON	74	HBC	+	13 $\pi^+ p$

$\Gamma(K\bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ
VALUE					
0.0158±0.0026 OUR FIT	Error includes scale factor of 1.2.				
0.0130±0.0024 OUR AVERAGE					
0.013 ± 0.003	COSTA...	80	OMEG	0	10 $\pi^- p \rightarrow K^+ K^- n$
0.013 ± 0.004	31 MARTIN	78B	SPEC	-	10 $\pi\rho \rightarrow K_S^0 K^- p$

³¹ From $(\Gamma_3\Gamma_6)^{1/2} = 0.056 \pm 0.034$ assuming $B(\rho_3(1690) \rightarrow \pi\pi) = 0.24$.

$\Gamma(\omega\pi)/[\Gamma(\omega\pi) + \Gamma(\rho\rho)]$	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_4/(\Gamma_4+\Gamma_{10})$
VALUE					
••• We do not use the following data for averages, fits, limits, etc. •••					
0.22±0.08	CASON	73	HBC	-	8,18.5 $\pi^- p$

$\Gamma(\eta\pi^+\pi^-)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
VALUE				
seen	FUKUI	88	SPEC	8.95 $\pi^- p \rightarrow \eta\pi^+\pi^- n$

 $\rho_3(1690)$ REFERENCES

ALDE	92C	ZPHY C54 553	+Bellazzini+ (SERP, BELG, LANL, LAPP, PISA, KEK)
FUKUI	88	PL B202 441	+Horikawa+ (SUGI, NAGO, KEK, KYOT, MIYA)
DENNEY	83	PR D28 2726	+Cranley, Firestone, Chapman+ (IOWA, MICH)
EVANGELISTA	81	NP B178 197	+ (BARI, BONN, CERN, DARE, LVP+)
ALPER	80	PL 94B 422	+Becker+ (AMST, CERN, CERN, MPIM, OXF+)
COSTA...	80	NP B175 402	Costa De Beauregard+ (BARI, BONN, CERN+)
GORLICH	80	NP B174 16	+Niczyporuk+ (CRAC, MPIM, CERN, ZEEM)
BECKER	79	NP B151 46	+Blanar, Blum+ (MPIM, CERN, ZEEM, CRAC)
CORDEN	79	NP B157 250	+Dowell, Garvey+ (BIRM, RHEL, TELA, LOWC) JP
BALTAY	78B	PR D17 62	+Cautis, Cohen, Csorna+ (COLU, BING)
MARTIN	78B	NP B140 158	+Ozmutlu, Baldi, Bohringer, Dorsaz+ (DURH, GEVA)
MARTIN	78D	PL 74B 417	+Ozmutlu, Baldi, Bohringer, Dorsaz+ (DURH, GEVA)
ANTIPOV	77	NP B119 45	+Busnello, Damgaard, Kienzle+ (SERP, GEVA)
GESSAROLI	77	NP B126 382	+ (BGNA, FIRZ, GENO, MILA, OXF, PAVI)
BLUM	75	PL 57B 403	+Chaubaud, Dietl, Garelick, Grayer+ (CERN, MPIM) JP
ESTABROOKS	75	NP B95 322	+Martin (DURH)
HYAMS	75	NP B100 205	+Jones, Weillhammer, Blum, Dietl+ (CERN, MPIM)
ENGLER	74	PR D10 2070	+Kraemer, Toaff, Weisser, Diaz+ (CMU, CASE)
GRAY	74	NP B75 189	+Hyams, Blum, Dietl+ (CERN, MPIM)
KLIGER	74	SJNP 19 428	+Beketov, Grechko, Guzhavin, Dubovikov+ (ITEP)

Translated from YAF 19 839.

OREN	74	NP B71 189	+Cooper, Fields, Rhines, Allison+	(ANL, OXF)
THOMPSON	74	NP B69 220	+Gaidos, McIlwain, Miller, Mulera+	(PURD)
CASON	73	PR D7 1971	+Biswas, Kenney, Madden+	(NDAM)
BOWEN	72	PRL 29 890	+Earles, Faisler, Blieden+	(NEAS, STON)
HOLMES	72	PR D6 3336	+Ferber, Slattery, Werner	(ROCH)
BALLAM	71B	PR D3 2606	+Chadwick, Guiragossian, Johnson+	(SLAC)
MATTHEWS	71C	NP B33 1	+Prentice, Yoon, Carroll+	(TNTO, WISC) JP
ARMENISE	70	LNC 4 199	+Ghidini, Forino, Cartacci+	(BARI, BGNA, FIRZ)
BARNHAM	70	PRL 24 1083	+Colley, Jobs, Kenyon, Pathak, Riddiford	(BIRM)
BARTSCH	70B	NP B22 109	+Kraus, Tsanos, Grote+	(AACH, BERL, CERN)
CASO	70	LNC 3 707	+Conte, Tomasini+	(GENO, HAMB, MILA, SACL)
STUNTEBECK	70	PL 32B 391	+Kenney, Deery, Biswas, Cason+	(NDAM)
ADERHOLZ	69	NP B11 259	+Bartsch+	(AACH3, BERL, CERN, JAGL, WARS)
ANDERSON	69	PRL 22 1390	+Collins+	(BNL, CMU)
ARMENISE	68	NC 54A 999	+Ghidini, Forino+	(BARI, BGNA, FIRZ, ORSAY) I
BALTAY	68	PRL 20 887	+Kung, Yeh, Ferbel+	(COLU, ROCH, RUTG, YALE) I
CASO	68	NC 54A 983	+Conte, Cords, Diaz+	(GENO, HAMB, MILA, SACL)
CREWELL	68B	PL 28B 136	+Karshon, Lal, Scarr, Skillicorn	(BNL)
JOHNSTON	68	PRL 20 1414	+Prentice, Steenberg, Yoon	(TNTO, WISC) IJP
FUCCACCI	66	PRL 17 890	+Kienzle, Levrat, Maglich, Martin	(CERN)
GOLDBERG	65	PL 17 354	+ (CERN, EPOL, ORSAY, MILA, CEA, SACL)	

OTHER RELATED PAPERS

BARNETT	83B	PL 120B 455	+Blockus, Burka, Chien, Christian+	(JHU)
EHRlich	66	PR 152 1194	+Selove, Yuta	(PENN)
LEVRAT	66	PL 22 714	+Tolstrup+	(CERN Missing Mass Spect. Collab.)
SEGUINOT	66	PL 19 712	+Martin+	(CERN Missing Mass Spect. Collab.)
BELLINI	65	NC 40A 948	+DiCorato, Duimio, Fiorini	(MILA)
DEUTSCH... FORINO	65 65	PL 18 351 PL 19 65	+Deutschmann+ +Gessaroli+	(AACH3, BERL, CERN) (BGNA, ORSAY, SACL)

 $\rho(1700)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

NOTE ON THE $\rho(1450)$ AND THE $\rho(1700)$

In our 1988 edition, we replaced the $\rho(1600)$ entry with two new ones, the $\rho(1450)$ and the $\rho(1700)$, because there was emerging evidence that the 1600-MeV region actually contains two ρ -like resonances. ERKAL 86 had pointed out this possibility with a theoretical analysis on the consistency of 2π and 4π electromagnetic form factors and the $\pi\pi$ scattering length. DONNACHIE 87, with a full analysis of data in the annihilation reactions $e^+e^- \rightarrow \pi^+\pi^-$, $2\pi^+2\pi^-$, and $\pi^+\pi^-\pi^0\pi^0$, and in the photoproduction reactions $\gamma p \rightarrow \pi^+\pi^-p$, $2\pi^+2\pi^-p$, and $\pi^+\pi^-\pi^0\pi^0p$, had also argued that to obtain a consistent picture two resonances, whose masses and widths could be fixed reasonably well, were necessary. This picture was supported by the analysis of DONNACHIE 87B of $J^P = 1^- \eta\rho^0$ mass spectra obtained in photoproduction and in e^+e^- annihilations; the analysis showed the need for a contribution from a ρ meson with a mass of about 1470 MeV, but could say little about a higher-mass resonance (actually the data could be explained without it). Confirmation of the decay $\rho(1450) \rightarrow \omega\pi$, and a tight constraint on the mass due to strong interference with the $\rho(770)$ tail, was found by DONNACHIE 91 in an analysis of $e^+e^- \rightarrow \omega\pi$.

The analysis of DONNACHIE 87 was extended by CLEGG 88 to include new data on 4π systems produced in e^+e^- annihilation and in τ decay (4π τ decays and 4π annihilation reactions can be related by the Conserved Vector Current assumption). These systems were successfully analyzed using interfering contributions from two ρ -like states, and from the tail of the $\rho(770)$ decaying into two-body states. While specific conclusions on $\rho(1450) \rightarrow 4\pi$ were obtained, the quality of the data used by CLEGG 88 prevented any conclusion on $\rho(1700) \rightarrow 4\pi$ decay.

An analysis by CLEGG 90 of 6π mass spectra from e^+e^- annihilation and from diffractive photoproduction provides evidence for two ρ mesons at about 2.1 and 1.8 GeV that decay strongly into 6π states. While the former is a candidate for

a new resonance, the latter could be a manifestation of the $\rho(1700)$, distorted by threshold effects.

Independent evidence for two 1^- states is provided by KILLIAN 80 in 4π electroproduction at $\langle Q^2 \rangle = 1$ (GeV/c^2), and by FUKUI 88 in a high-statistics sample of the $\eta\pi\pi$ system in π^-p charge exchange.

This scenario with two overlapping resonances is supported by other data. BISELLO 89 measured the pion form factor in the interval 1.35–2.4 GeV with significant statistics (280 $e^+e^- \rightarrow \pi^+\pi^-$ events with very low background); a deep minimum is observed around 1.6 GeV, and the best fit to the form factor is obtained with the hypothesis of ρ -like resonances at 1420 and 1770 MeV with widths of about 250 MeV. ANTONELLI 88 found that the $e^+e^- \rightarrow \eta\pi^+\pi^-$ cross section (using three different η decay modes) is better fitted with two fully interfering Breit-Wigners, with parameters in fair agreement with those of DONNACHIE 87 and BISELLO 89.

These results (although ANTONELLI 88 is statistically less significant than BISELLO 89) have also resolved the disagreement between DONNACHIE 87 and FUKUI 88 on the $\rho(1450)$ width in favor of the DONNACHIE 87 value. From this point of view, the two experiments can be considered as solid confirmation of the $\rho(1450)$. For the possibility that its $\phi\pi$ mode actually contains two independent vector states, see LANDSBERG 92.

Several observations on the $\omega\pi$ system in the 1200-MeV region (FRENKIEL 72, COSME 76, BARBER 80C, ASTON 80C, ATKINSON 84C, BRAU 88, AMSLER 93B) may be interpreted in terms of either $J^P = 1^- \rho(770) \rightarrow \pi\omega$ production (LAYSSAC 71) or $J^P = 1^+ b_1(1235)$ production (BRAU 88, AMSLER 93B). We argue that no special entry for a $\rho(1250)$ is needed. The LASS amplitude analysis (ASTON 91B) showing evidence for $\rho(1270)$ is preliminary and needs confirmation. For completeness, the relevant observations are listed under the $\rho(1450)$.

 $\rho(1700)$ MASS **$\eta\rho^0$ AND MIXED MODES**

VALUE (MeV)

DOCUMENT ID

1700 ± 20 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.

1712 ± 13 OUR AVERAGE Includes data from the 2 datablocks that follow this one. Error includes scale factor of 1.2.

MIXED MODES

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

1700 ± 25

DONNACHIE 87 RVUE

● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●

1580 ± 20

¹ BUON 82 DM1 $e^+e^- \rightarrow$ hadrons **$\eta\rho^0$ MODE**

VALUE (MeV)

DOCUMENT ID

TECN

COMMENT

The data in this block is included in the average printed for a previous datablock.

1740 ± 20

ANTONELLI 88 DM2 $e^+e^- \rightarrow \eta\pi^+\pi^-$

1701 ± 15

FUKUI 88 SPEC 8.95 $\pi^-\rho \rightarrow \eta\pi^+\pi^-n$

Meson Full Listings

$\rho(1700)$

$\pi^+\pi^-$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1768 ± 21	BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1546 ± 26	GESHKENBEIN ⁸⁹	RVUE	
1650	2 ERKAL 85	RVUE	20-70 $\gamma\rho \rightarrow \gamma\pi$
1550 ± 70	ABE 84 ^B	HYBR	20 $\gamma\rho \rightarrow \pi^+\pi^-\rho$
1590 ± 20	3 ASTON 80	OMEG	20-70 $\gamma\rho \rightarrow \rho 2\pi$
1600.0 ± 10.0	4 ATIYA 79 ^B	SPEC	50 $\gamma C \rightarrow C 2\pi$
1598.0 ^{+24.0} _{-22.0}	BECKER 79	ASPK	17 $\pi^-\rho$ polarized
1659 ± 25	2 LANG 79	RVUE	
1575	2 MARTIN 78 ^C	RVUE	17 $\pi^-\rho \rightarrow \pi^+\pi^-\eta$
1610 ± 30	2 FROGGATT 77	RVUE	17 $\pi^-\rho \rightarrow \pi^+\pi^-\eta$
1590 ± 20	5 HYAMS ^r 73	ASPK	17 $\pi^-\rho \rightarrow \pi^+\pi^-\eta$

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1582 ± 36	1600	CLELAND 82 ^B	SPEC	±	50 $\pi\rho \rightarrow K_S^0 K^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

$2(\pi^+\pi^-)$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1520 ± 30		3 ASTON 81 ^E	OMEG	20-70 $\gamma\rho \rightarrow p 4\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1570 ± 20		6 CORDIER 82	DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1654 ± 25		7 DIBIANCA 81	DBC	$\pi^+d \rightarrow pp 2(\pi^+\pi^-)$
1666 ± 39		6 BACCI 80	FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1780	34	KILLIAN 80	SPEC	11 $e^-\rho \rightarrow 2(\pi^+\pi^-)$
1500		8 ATIYA 79 ^B	SPEC	50 $\gamma C \rightarrow C 4\pi^\pm$
1570 ± 60	65	9 ALEXANDER 75	HBC	7.5 $\gamma\rho \rightarrow p 4\pi$
1550 ± 60		3 CONVERSI 74	OSPK	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
1550 ± 50	160	SCHACHT 74	STRC	5.5-9 $\gamma\rho \rightarrow p 4\pi$
1450 ± 100	340	SCHACHT 74	STRC	9-18 $\gamma\rho \rightarrow p 4\pi$
1430 ± 50	400	BINGHAM 72 ^B	HBC	9.3 $\gamma\rho \rightarrow p 4\pi$

$\pi^+\pi^-\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1660 ± 30	ATKINSON 85 ^B	OMEG	20-70 $\gamma\rho$
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES


VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1783 ± 15	CLEGG 90	RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-) 2(\pi^+\pi^-\pi^0)$

- From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0 K_L^0$, $K_S^0 K^\pm\pi^\mp$.
- From phase shift analysis of HYAMS 73 data.
- Simple relativistic Breit-Wigner fit with constant width.
- An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.
- Included in BECKER 79 analysis.
- Simple relativistic Breit-Wigner fit with model dependent width.
- One peak fit result.
- Parameters roughly estimated, not from a fit.
- Skew mass distribution compensated by Ross-Stodolsky factor.

$\rho(1700)$ WIDTH

$\eta\rho^0, \pi^+\pi^-,$ AND MIXED MODES

235 ± 50 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.
 213 ± 21 OUR AVERAGE Includes data from the 3 datablocks that follow this one. Error includes scale factor of 1.5. See the ideogram below.



MIXED MODES

220 ± 25 DONNACHIE 87 RVUE
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 340 ± 80¹⁰ BUON 82 DM1 $e^+e^- \rightarrow$ hadrons

$\eta\rho^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 ± 30	ANTONELLI 88	DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$
282 ± 44	FUKUI 88	SPEC	8.95 $\pi^-\rho \rightarrow \eta\pi^+\pi^-\eta$

$\pi^+\pi^-\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
224 ± 22	BISELLO 89	DM2	$e^+e^- \rightarrow \pi^+\pi^-\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
620 ± 60	GESHKENBEIN ⁸⁹	RVUE	
< 315	11 ERKAL 85	RVUE	20-70 $\gamma\rho \rightarrow \gamma\pi$
280 ⁺³⁰ ₋₈₀	ABE 84 ^B	HYBR	20 $\gamma\rho \rightarrow \pi^+\pi^-\rho$
230.0 ± 80.0	12 ASTON 80	OMEG	20-70 $\gamma\rho \rightarrow \rho 2\pi$
283.0 ± 14.0	13 ATIYA 79 ^B	SPEC	50 $\gamma C \rightarrow C 2\pi$
175.0 ^{+98.0} _{-53.0}	BECKER 79	ASPK	17 $\pi^-\rho$ polarized
232 ± 34	11 LANG 79	RVUE	
340	11 MARTIN 78 ^C	RVUE	17 $\pi^-\rho \rightarrow \pi^+\pi^-\eta$
300 ± 100	11 FROGGATT 77	RVUE	17 $\pi^-\rho \rightarrow \pi^+\pi^-\eta$
180 ± 50	14 HYAMS 73	ASPK	17 $\pi^-\rho \rightarrow \pi^+\pi^-\eta$

$K\bar{K}$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
265 ± 120	1600	CLELAND 82 ^B	SPEC	±	50 $\pi\rho \rightarrow K_S^0 K^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					

$2(\pi^+\pi^-)$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
400 ± 50	12	ASTON 81 ^E	OMEG	20-70 $\gamma\rho \rightarrow p 4\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
510 ± 40	15	CORDIER 82	DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
400 ± 146	16	DIBIANCA 81	DBC	$\pi^+d \rightarrow pp 2(\pi^+\pi^-)$
700 ± 160	15	BACCI 80	FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
100	34	KILLIAN 80	SPEC	11 $e^-\rho \rightarrow 2(\pi^+\pi^-)$
600	17	ATIYA 79 ^B	SPEC	50 $\gamma C \rightarrow C 4\pi^\pm$
340 ± 160	65	18 ALEXANDER 75	HBC	7.5 $\gamma\rho \rightarrow p 4\pi$
360 ± 100	12	CONVERSI 74	OSPK	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
400 ± 120	160	19 SCHACHT 74	STRC	5.5-9 $\gamma\rho \rightarrow p 4\pi$
850 ± 200	340	19 SCHACHT 74	STRC	9-18 $\gamma\rho \rightarrow p 4\pi$
650 ± 100	400	BINGHAM 72 ^B	HBC	9.3 $\gamma\rho \rightarrow p 4\pi$

See key on page 1343

Meson Full Listings
 $\rho(1700)$ $\pi^+\pi^-\pi^0\pi^0$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 ± 50	ATKINSON	85B OMEG	20-70 γp

 $3(\pi^+\pi^-)$ AND $2(\pi^+\pi^-\pi^0)$ MODES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
285 ± 20	CLEGG	90 RVUE	$e^+e^- \rightarrow 3(\pi^+\pi^-)2(\pi^+\pi^-\pi^0)$

¹⁰ From global fit of ρ , ω , ϕ and their radial excitations to channels $\omega\pi^+\pi^-$, K^+K^- , $K_S^0K_L^0$, $K_S^0K^\pm\pi^\mp$.

¹¹ From phase shift analysis of HYAMS 73 data.

¹² Simple relativistic Breit-Wigner fit with constant width.

¹³ An additional 40 MeV uncertainty in both the mass and width is present due to the choice of the background shape.

¹⁴ Included in BECKER 79 analysis.

¹⁵ Simple relativistic Breit-Wigner fit with model-dependent width.

¹⁶ One peak fit result.

¹⁷ Parameters roughly estimated, not from a fit.

¹⁸ Skew mass distribution compensated by Ross-Stodolsky factor.

¹⁹ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

 $\rho(1700)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\rho\pi\pi$	dominant
Γ_2 $\rho^0\pi^+\pi^-$	large
Γ_3 $\rho^0\pi^0\pi^0$	
Γ_4 $\rho^\pm\pi^\mp\pi^0$	[a] large
Γ_5 $2(\pi^+\pi^-)$	large
Γ_6 $\pi^+\pi^-$	seen
Γ_7 $K\bar{K}^*(892) + c.c.$	seen
Γ_8 $\eta\rho$	seen
Γ_9 $K\bar{K}$	seen
Γ_{10} e^+e^-	seen
Γ_{11} $\rho^0\rho^0$	
Γ_{12} $\pi\omega$	

[a] The value is for the sum of the charge states indicated.

 $\rho(1700)$ $\Gamma(\rho^0\pi^+\pi^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the cross-section into channel $\rho^0\pi^+\pi^-$ annihilation.

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.83 ± 0.42	BACCI	80 FRAG	$e^+e^- \rightarrow 2(\pi^+\pi^-)$
2.6 ± 0.2	DELICOURT	81B DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.13	²⁰ DIEKMAN	88 RVUE	$e^+e^- \rightarrow \pi^+\pi^-$

²⁰ Using total width = 220 MeV.

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.305 ± 0.071	²¹ BIZOT	80 DM1	e^+e^-

VALUE (eV)	DOCUMENT ID	TECN	COMMENT
7 ± 3	ANTONELLI	88 DM2	$e^+e^- \rightarrow \eta\pi^+\pi^-$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.035 ± 0.029	²¹ BIZOT	80 DM1	e^+e^-

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
3.510 ± 0.090	²¹ BIZOT	80 DM1	e^+e^-

²¹ Model dependent.

 $\rho(1700)$ BRANCHING RATIOS

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
$0.287^{+0.043}_{-0.042}$	BECKER	79 ASPK	$17\pi^-\rho$ polarized	
0.15 to 0.30	²² MARTIN	78C RVUE	$17\pi^-\rho \rightarrow \pi^+\pi^-\pi^-$	
<0.20	²³ COSTA...	77B RVUE	$e^+e^- \rightarrow 2\pi, 4\pi$	
0.30 ± 0.05	²² FROGGATT	77 RVUE	$17\pi^-\rho \rightarrow \pi^+\pi^-\pi^-$	
<0.15	²⁴ EISENBERG	73 HBC	$5\pi^+\rho \rightarrow \Delta^++2\pi$	
0.25 ± 0.05	²⁵ HYAMS	73 ASPK	$17\pi^-\rho \rightarrow \pi^+\pi^-\pi^-$	
0.20 ± 0.05	MONTANET	73 HBC	$0.0\bar{p}\rho$	

²² From phase shift analysis of HYAMS 73 data.

²³ Estimate using unitarity, time reversal invariance, Breit-Wigner.

²⁴ Estimated using one-pion-exchange model.

²⁵ Included in BECKER 79 analysis.

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_5
0.13 ± 0.05	ASTON	80 OMEG	$20-70\gamma p \rightarrow p2\pi$	
<0.14	²⁶ DAVIER	73 STRC	$6-18\gamma p \rightarrow p4\pi$	
<0.2	²⁷ BINGHAM	72B HBC	$9.3\gamma p \rightarrow p2\pi$	

²⁶ Upper limit is estimate.
²⁷ 2σ upper limit.

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ_5
0.15 ± 0.03	²⁸ DELICOURT	81B DM1	$e^+e^- \rightarrow \bar{K}K\pi$	

²⁸ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
<0.04		DONNACHIE	87B RVUE		
<0.02	58	ATKINSON	86B OMEG	$20-70\gamma p$	

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ_5
0.123 ± 0.027	DELICOURT	82 DM1	$e^+e^- \rightarrow \pi^+\pi^-\text{MM}$	
~ 0.1	ASTON	80 OMEG	$20-70\gamma p$	

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_3+\Gamma_4+0.709\Gamma_8)/\Gamma_5$
2.6 ± 0.4	²⁹ BALLAM	74 HBC	$9.3\gamma p$	

²⁹ Upper limit. Background not subtracted.

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_9/Γ_5
0.015 ± 0.010		³⁰ DELICOURT	81B DM1		$e^+e^- \rightarrow \bar{K}K$	
<0.04	95	BINGHAM	72B HBC	0	$9.3\gamma p$	

³⁰ Assuming $\rho(1700)$ and ω radial excitations to be degenerate in mass.

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ_7
0.052 ± 0.026	BUON	82 DM1	$e^+e^- \rightarrow \text{hadrons}$	

VALUE	EVT%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_5
~ 1.0		DELICOURT	81B DM1	$e^+e^- \rightarrow 2(\pi^+\pi^-)$	
0.7 ± 0.1	500	SCHACHT	74 STRC	$5.5-18\gamma p \rightarrow p4\pi$	
0.80		³¹ BINGHAM	72B HBC	$9.3\gamma p \rightarrow p4\pi$	

³¹ The $\pi\pi$ system is in S-wave.

VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_4
<0.10	ATKINSON	85B OMEG		$20-70\gamma p$	
<0.15	ATKINSON	82 OMEG	0	$20-70\gamma p \rightarrow p4\pi$	

Meson Full Listings

$\rho(1700)$, $X(1700)$, $f_J(1710)$

$\rho(1700)$ REFERENCES

CLEGG 90	ZPHY C45 677	+Donnachie	(LANC, MCHS)
BISELLO 89	PL B220 321	+Busetto+	(DM2 Collab.)
GESHKENBEIN 89	ZPHY 45 351		(ITEP)
ANTONELLI 88	PL B212 133	+Baldini+	(DM2 Collab.)
DIEKMANN 88	PRP 159 101		(BONN)
FUKUI 88	PL B202 441	+Horikawa+	(SUGI, NAGO, KEK, KYOT, MIYA)
DONNACHIE 87	ZPHY C33 407	+Mirzaie	(MCHS)
DONNACHIE 87B	ZPHY C34 257	+Clegg	(MCHS, LANC)
ATKINSON 86B	ZPHY C30 531	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ATKINSON 85B	ZPHY C26 499	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ERKAL 85	ZPHY C29 485	+Olsson	(WISC)
ASE 84B	PRL 53 751	+Bacon, Ballam+	(SLAC Hybrid Facility Photon Collab.)
ATKINSON 82	PL 108B 55	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
BUON 82	PL 118B 221	+Bisello, Bizot, Cordier, Delcourt+	(LALO, MONP)
CLELAND 82B	NP B208 228	+Defosse, Dorsaz, Gloor	(DURH, GEVA, LAUS, PITT)
CORDIER 82	PL 109B 129	+Bisello, Bizot, Buon, Delcourt	(LALO)
DEL COURT 82	PL 113B 93	+Bisello, Bizot, Buon, Cordier, Mane	(LALO)
ASTON 81E	NP B189 15	+ (BONN, CERN, EPOL, GLAS, LANC, MCHS+)	
DEL COURT 81B	Bonn Conf. 205		(ORSAY)
		Cordier, Bisello, Bizot, Buon, Delcourt	(LALO)
DIBIANCA 81	PR D23 595	+Fickinger, Malko, Dado, Engler+	(CASE, CMU)
ASTON 80	PL 92B 215	+ (BONN, CERN, EPOL, GLAS, LANC, MCHS+)	
BACCI 80	PL 95B 139	+DeZori, Penso, Baldini-Celio+	(ROMA, FRAS)
BIZOT 80	Madison Conf. 546	+Bisello, Buon, Cordier, Delcourt+	(LALO, MONP)
KILLIAN 80	PR D21 3005	+Treadwell, Ahrens, Berkelman, Cassel+	(CORN)
ATIYA 79B	PRL 43 1691	+Holmes, Knapp, Lee, Seto+	(COLU, ILL, FNAL)
BECKER 79	NP B151 46	+Blanan, Blum+	(MPIIM, CERN, ZEEM, CERN)
LANG 79	NP D19 956	+Mas-Parareda	(GRAZ)
MARTIN 78C	ANP 114 1	+Pennington	(CERN)
COSTA... 77B	PL 71B 345	Costa De Beauregard, Pire, Truong	(EPOL)
FROGGATT 77	NP B129 89	+Peterson	(GLAS, NORD)
ALEXANDER 75	PL 57B 487	+Benary, Gandsman, Lissauer+	(TELA)
BALLAM 74	NP B76 375	+Chadwick, Bingham, Fretter+	(SLAC, LBL, MPIIM)
CONVERSI 74	PL 52B 493	+Paoluzi, Ceradini, Grilli+	(ROMA, FRAS)
SCHUCHT 74	NP B81 205	+Derado, Fries, Park, Nount	(MPIIM)
DAVIER 73	NP B58 31	+Derado, Fries, Liu, Mozley, Odian, Park+	(SLAC)
EISENBERG 73	PL 43B 149	+Karshon, Mikenberg, Pitluck+	(REHO)
HYAMS 73	NP B64 134	+Jones, Weilhammer, Blum, Dietl+	(CERN, MPIIM)
MONTANET 73B	Erice School 518		(CERN)
BINGHAM 72B	PL 41B 635	+Rabin, Rosenfeld, Smadja+	(LBL, UCB, SLAC) IGGP

OTHER RELATED PAPERS

AMSLER 93B	PL B311 362	+Armstrong, Augustin+	(Crystal Barrel Collab.)
LANDSBERG 92	SJNP 55 1051		(SERP)
	Translated from YAF 55 1896.		
ASTON 91B	NPBPS 21 105	+Awaji, Bienz+	(LASS Collab.)
ACHASOV 88C	PL B209 373	+Kozhevnikov	(NOVO)
BRAU 88	PR D37 2379	+Frane+	(SLAC Hybrid Facility Photon Collab.) JP
CLEGG 88	ZPHY C40 313	+Donnachie	(MCHS, LANC)
ASTON 87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
ERKAL 86	ZPHY C31 615	+Olsson	(WISC)
BARKOV 85	NP B256 365	+Chilingarov, Eidelman, Khazin, Lechuk+	(NOVO)
BISELLO 85	LAL 85-15	+Augustin, Ajitouni+	(PADO, LALO, CLER, FRAS)
ATKINSON 84C	NP B243 13	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ATKINSON 83B	PL 127B 132	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
ATKINSON 83C	NP B229 269	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)	
AUGUSTIN 83	LAL 83-21	+Ayach, Bisello, Baldini+	(LALO, PADO, FRAS)
SHAMBROOM 82	PR D26 1	+Wilson, Anderson, Francis+	(HARV, EFI, ILL, OXF)
BARBER 80C	ZPHY C4 169	+Dainton, Brodbeck, Brookes+	(DARE, LANC, SHEF)
KILLIAN 80	PR D21 3005	+Treadwell, Ahrens, Berkelman, Cassel+	(CORN)
COSME 76	PL 63B 352	+Courau, Dudelzalk, Grelaud, Jean-Marie+	(ORSAY)
FRENKIEL 72	NP B47 61	+Chesquiers, Lillstol, Chung+	(CDEF, CERN)
ALVENSLEBEN 71	PRL 26 273	+Becker, Bertram, Chen+	(DESY, MIT) G
BRAUN 71	NP B30 213	+Fridman, Gerber, Givernaud+	(STRB) G
BULOS 71	PRL 26 149	+Busza, Kehoe, Beniston+	(SLAC, UMD, IBM, LBL) G
LAYSSAC 71	NC 6A 134	+Renard	(MONP)

$X(1700)$

$$I^G(J^{PC}) = \text{even}^+(\text{?}^{\text{?}})$$

OMITTED FROM SUMMARY TABLE

Enhancement seen in the $\eta\pi\pi$ system produced in the radiative decay of the $J/\psi(1S)$. May contain significant substructure. Relation to other enhancements seen in radiative $J/\psi(1S)$ decay unclear (see HITLIN 83). Enhancement seen in the $J = 2, \rho\pi\pi$ wave of the $\pi^+\pi^-\pi^+\pi^-$ system produced in pomeron-pomeron collisions. Tentatively called $X(1700)$ by us. Needs confirmation.

$X(1700)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1700.0 ± 45	EDWARDS 83B	CBAL	$J/\psi \rightarrow \eta\gamma 2\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1750	BREAKSTONE 93	SFM	$\rho\rho \rightarrow \pi^+\pi^-\pi^+\pi^-$

$X(1700)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
520 ± 110	EDWARDS 83B	CBAL	$J/\psi \rightarrow \eta\gamma 2\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
200 to 300	BREAKSTONE 93	SFM	$\rho\rho \rightarrow \pi^+\pi^-\pi^+\pi^-$

$X(1700)$ REFERENCES

BREAKSTONE 93	ZPHY C58 251	+Campanini+	(IOWA, CERN, DORT, HEIDH, WARS)
EDWARDS 83B	PRL 51 859	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
HITLIN 83	Cornell Conf. 746		(CIT)

$f_J(1710)$ was $\theta(1690)$

$$I^G(J^{PC}) = 0^+(\text{even}^{++})$$

NOTE ON THE $f_J(1710)$

The $f_J(1710)$ is seen in the "gluon rich" radiative decay $J/\psi(1S) \rightarrow \gamma f_J(1710)$; therefore $C = +$. It decays into 2η and $K_S^0 K_S^0$, which implies $I^G J^{PC} = 0^+(\text{even})^{++}$. In an amplitude analysis of the $K\bar{K}$ and $\pi^+\pi^-$ systems produced in $J/\psi(1S)$ radiative decay, CHEN 91 finds a large spin-0 component for this particle, but WA 76 favors spin 2 in central production. The spin is thus uncertain. This resonance is also observed in $K\bar{K}$ systems recoiling against a ϕ or an ω in hadronic $J/\psi(1S)$ decay [however, according to FALVARD 88, $J/\psi(1S) \rightarrow \omega f_J(1710)$ is rather controversial]. The $f_J(1710)$ is not seen in $J/\psi(1S) \rightarrow \gamma\rho^0\rho^0$ (BISELLO 89B), in agreement with the indication (BALTRUSAITIS 85G) that the $\rho\rho$ enhancement in this region has $J^P = 0^-$, and hence is unrelated to the $f_J(1710)$.

Clear evidence is seen in hadroproduction (ARMSTRONG 89D, 300-GeV/c pp central production of $K\bar{K}$), both in K^+K^- and $K_S^0 K_S^0$. Mass and width determinations are complicated because the spectra are dominated by overlap with the $f_2'(1525)$. The apparent large disagreement between the widths found by ARMSTRONG 89D in the two channels (≈ 180 MeV in K^+K^- and ≈ 100 MeV in $K_S^0 K_S^0$) can be explained by the arbitrariness of the polynomial-exponential background shape, which leads to a large systematic error for the width. ARMSTRONG 93C also sees in $\eta\eta$ a broad peak at 1747 MeV, which may be the $f_J(1710)$. This resonance is not observed in the exclusive hypercharge-exchange reaction $K^-p \rightarrow K_S^0 K_S^0 \Lambda$ (ASTON 88D).

A partial-wave analysis of the $K_S^0 K_S^0$ system (BOLONKIN 88) finds a D_0 wave ($J^{PC} = 2^{++}$) behavior near 1700 MeV, but the width (≈ 30 MeV) is much narrower than that observed in $J/\psi(1S)$ decays and in hadroproduction.

Note that in our 1992 edition, this particle was named the $f_2(1710)$; see also our "Note on Non- $q\bar{q}$ Mesons."

$f_J(1710)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1709 ± 5	OUR AVERAGE		
1713 ± 10	ARMSTRONG 89D	OMEG	$300 pp \rightarrow \rho\rho K^+ K^-$
1706 ± 10	ARMSTRONG 89D	OMEG	$300 pp \rightarrow \rho\rho K_S^0 K_S^0$
1707.0 ± 10.0	AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K^+ K^-$
1698 ± 15	AUGUSTIN 87	DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$
1720 ± 10 ± 10	BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1710 ± 20	CHEN 91	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-, \gamma K\bar{K}$
1700 ± 15	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1720 ± 60	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
1638 ± 10	¹ FALVARD 88	DM2	$J/\psi \rightarrow \phi K^+ K^-$
1690 ± 4	² FALVARD 88	DM2	$J/\psi \rightarrow \phi K^+ K^-$
1730 ± 2	^{3,4} LONGACRE 86	MPS	$22 \pi^- p \rightarrow n 2K_S^0$
1742.0 ± 15.0	WILLIAMS 84	MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
1670 ± 50	BLOOM 83	CBAL	$J/\psi \rightarrow \gamma 2\eta$
1650 ± 50	BURKE 82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
1730.0 ± 10 ± 20	ETKIN 82C	MPS	$23 \pi^- p \rightarrow n 2K_S^0$
1708.0 ± 30.0	FRANKLIN 82	MRK2	$e^+e^- \rightarrow \gamma K^+ K^-$

¹ From an analysis ignoring interference with $f_2'(1525)$.
² From an analysis including interference with $f_2'(1525)$.
³ From a partial-wave analysis of data using a K -matrix formalism with 5 poles, but assuming spin 2.
⁴ Fit with constrained inelasticity.

See key on page 1343

Meson Full Listings

$f_J(1710), X(1740)$

 $f_J(1710)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
140 ± 12 OUR AVERAGE			
181 ± 30	ARMSTRONG 89D	OMEG	300 $pp \rightarrow ppK^+K^-$
104 ± 30	ARMSTRONG 89D	OMEG	300 $pp \rightarrow ppK_S^0K_S^0$
166.4 ± 33.2	AUGUSTIN 88	DM2	$J/\psi \rightarrow \gamma K^+K^-$
136 ± 28	AUGUSTIN 87	DM2	$J/\psi \rightarrow \gamma \pi^+\pi^-$
130 ± 20	BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
186 ± 30	CHEN 91	MRK3	$J/\psi \rightarrow \gamma \pi^+\pi^-, \gamma K\bar{K}$
30 ± 20	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
350 ± 150	BOLONKIN 88	SPEC	$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
148 ± 17	⁵ FALVARD 88	DM2	$J/\psi \rightarrow \phi K^+K^-$
184 ± 6	⁶ FALVARD 88	DM2	$J/\psi \rightarrow \phi K^+K^-$
122 ± 74 - 15	^{7,8} LONGACRE 86	MPS	$22 \pi^- p \rightarrow n2K_S^0$
57.0 ± 38.0	WILLIAMS 84	MPSF	$200 \pi^- N \rightarrow 2K_S^0 X$
160 ± 80	BLOOM 83	CBAL	$J/\psi \rightarrow \gamma 2\eta$
200 ± 100	BURKE 82	MRK2	$J/\psi \rightarrow \gamma 2\rho$
200.0 ± 156.0 9.0	⁹ ETKIN 82B	MPS	$23 \pi^- p \rightarrow n2K_S^0$
156.0 ± 60.0	FRANKLIN 82	MRK2	$e^+e^- \rightarrow \gamma K^+K^-$

⁵ From an analysis ignoring interference with $f_2'(1525)$.⁶ From an analysis including interference with $f_2'(1525)$.⁷ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2.⁸ Fit with constrained inelasticity.⁹ From an amplitude analysis of the $K_S^0 K_S^0$ system. **$f_J(1710)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $\eta\eta$	
Γ_3 $\pi\pi$	seen
Γ_4 $\rho\rho$	
Γ_5 $\gamma\gamma$	

 $f_J(1710)$ $\Gamma(I)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma(\text{total})$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_5/\Gamma$
<0.11	95	¹⁰ BEHREND 89C	CELL	$\gamma\gamma \rightarrow K_S^0 K_S^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.48	95	ALBRECHT 90G	ARG	$\gamma\gamma \rightarrow K^+K^-$	
<0.28	95	¹⁰ ALTHOFF 85B	TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$	
¹⁰ Assuming helicity 2.					

 $f_J(1710)$ BRANCHING RATIOS

$\Gamma(K\bar{K})/\Gamma(\text{total})$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.38 ^{+0.09} _{-0.19}	^{11,12} LONGACRE 86	MPS	$22 \pi^- p \rightarrow n2K_S^0$	
$\Gamma(\eta\eta)/\Gamma(\text{total})$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.18 ^{+0.03} _{-0.13}	^{11,12} LONGACRE 86	RVUE		
$\Gamma(\pi\pi)/\Gamma(\text{total})$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.039 ^{+0.002} _{-0.024}	^{11,12} LONGACRE 86	RVUE		
$\Gamma(\pi\pi)/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.39 ± 0.14	ARMSTRONG 91	OMEG	300 $pp \rightarrow pp\pi\pi, ppK\bar{K}$	

¹¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles, but assuming spin 2.¹² Fit with constrained inelasticity. **$f_J(1710)$ REFERENCES**

ARMSTRONG 91	ZPHY C51 351	+Benayoun+	(ATHU, BARI, BIRM, CERN, CDEF)
CHEN 91	Hadron 91 Conf.		(Mark III Collab.)
SLAC-PUB-5669			
ALBRECHT 90G	ZPHY C48 183	+Ehrlichmann, Harder+	(ARGUS Collab.)
ARMSTRONG 89D	PL B227 186	+Benayoun	(ATHU, BARI, BIRM, CERN, CDEF)
BEHREND 89C	ZPHY C43 91	+Criegee, Dainton+	(CELLO Collab.)
AUGUSTIN 88	PRL 60 2238	+Calcaterra+	(DM2 Collab.)
BOLONKIN 88	NP B309 426	+Bloschenko, Gorin+	(ITEP, SERP)
FALVARD 88	PR D38 2706	+Ajaltouni+	(CLER, FRAS, LALO, PADO)
AUGUSTIN 87	ZPHY C36 369	+Cosme+	(LALO, CLER, FRAS, PADO)
BALTRUSAIT...87	PR D35 2077	Baltrusaitis, Coffman, Dubois+	(Mark III Collab.)
LONGACRE 86	PL B177 223	+Etkin+	(BNL, BRAN, CUNY, DUKE, NDAM)
ALTHOFF 85B	ZPHY C29 189	+Braunschweig, Kirschfink+	(TASSO Collab.)
WILLIAMS 84	PR D30 877	+Diamond+	(VAND, NDAM, TUFTS, ARIZ, FNAL+)
BLOOM 83	ARNS 33 143	+Peck	(SLAC, CIT)
BURKE 82	PRL 49 632	+Trilling, Abrams, Alam, Blocker+	(LBL, SLAC)
ETKIN 82B	PR D25 1786	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
ETKIN 82C	PR D25 2446	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
FRANKLIN 82	SLAC-254		(SLAC)

OTHER RELATED PAPERS

CHEN 91	Hadron 91 Conf.		(Mark III Collab.)
SLAC-PUB-5669			
PROKOSHIN 91	SPD 316 155		(GAM2 Collab.)
Translated from DANS 316 900.			
BISELLO 89B	PR D39 701	Busetto+	(DM2 Collab.)
ASTON 88D	NP B301 525	+Awaji, Bienz+	(SLAC, NAGO, CINC, INUS)
AKESSON 86	NP B264 154	+Albrow, Almehed+	(Axial Field Spec. Collab.)
ARMSTRONG 86B	PL 167B 133	+Bloodworth, Carney+	(ATHU, BARI, BIRM, CERN)
BALTRUSAIT...86B	PR D33 1222	Baltrusaitis, Coffman, Hauser+	(Mark III Collab.)
ALTHOFF 83	PL 121B 216	+Brandelik, Boerner, Burkhardt+	(TASSO Collab.)
BARNETT 83B	PL 120B 455	+Blockus, Burke, Chien, Christian+	(JHU)
ALTHOFF 82	ZPHY C16 13	+Boerner, Burkhardt+	(TASSO Collab.)
BARNES 82	PL B116 365	+Close	(RHEL)
BARNES 82B	NP B198 360	+Close, Monaghan	(RHEL, OXFPT)
TANIMOTO 82	PL 116B 198		(BIEL)

X(1740)

$$I^G(J^{PC}) = 0^+(\text{even}^{++})$$

OMITTED FROM SUMMARY TABLE

We have collected here resonances in the $\eta\eta$ channel which may be not the same state. See also the minireview under $f_0(1710)$. $J^P = 0^+$ or 2^+ .

X(1740) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1748 ± 10	ARMSTRONG 93C	SPEC	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
1744 ± 15	¹ ALDE 92D	GAM2	$38 \pi^- p \rightarrow \eta\eta N^*$
¹ ALDE 92 combines all the GAMS-2000 data.			

X(1740) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
264 ± 25	ARMSTRONG 93C	SPEC	$\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
< 80	ALDE 92D	GAM2	$38 \pi^- p \rightarrow \eta\eta N^*$

X(1740) DECAY MODES

Mode
Γ_1 $\eta\eta$
Γ_2 $\pi^0\pi^0$
Γ_3 $\eta\eta'$

X(1740) BRANCHING RATIOS

$\Gamma(\pi^0\pi^0)/\Gamma(\eta\eta)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1	90	ALDE 92D	GAM2	$38 \pi^- p \rightarrow \eta\eta N^*$	
$\Gamma(\eta\eta')/\Gamma(\eta\eta)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1	90	ALDE 92D	GAM2	$38 \pi^- p \rightarrow \eta\eta N^*$	

X(1740) REFERENCES

ARMSTRONG 93C	PL B307 394	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
ALDE 92	PL B276 375	+ (SERP, BELG, LANC, LAPP, PISA, KEK)	
ALDE 92D	PL B284 457	+Binon, Bricman+	(GAM2 Collab.)
Translated from YAF 54 745.			

Meson Full Listings

 $\eta(1760)$, $\pi(1770)$, $X(1775)$, $f_2(1810)$ **$\eta(1760)$**

$$I^G(J^{PC}) = 0^+(0^{-+})$$

OMITTED FROM SUMMARY TABLE

Seen by DM2 in the $\rho\rho$ system BISELLO 89B. Needs confirmation. Structure in this region has been reported before in the same system BALTRUSAITIS 86B and in the $\omega\omega$ system BALTRUSAITIS 85C, BISELLO 87.

 $\eta(1760)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1760±11	320	¹ BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

¹ Estimated by us from various fits. **$\eta(1760)$ WIDTH**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
60±16	320	² BISELLO	89B DM2	$J/\psi \rightarrow 4\pi\gamma$

² Estimated by us from various fits. **$\eta(1760)$ REFERENCES**

BISELLO 89B PR D39 701	Busetto+	(DM2 Collab.)
BISELLO 87 PL B192 239	+Ajaltouni, Baldini+	(PADO, CLER, FRAS, LALO)
BALTRUSAITIS...86B PR D33 1222	Baltrusaitis, Coffman, Hauser+	(Mark III Collab.)
BALTRUSAITIS...85C PRL 55 1723	Baltrusaitis+	(CIT, UCSC, ILL, SLAC, WASH)

 $\pi(1770)$

$$I^G(J^{PC}) = 1^-(0^{-+})$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the diffractively produced 3π system. Needs confirmation.

 $\pi(1770)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1770±30	1100	BELLINI	82 SPEC	-	$40\pi^-A \rightarrow 3\pi A$

 $\pi(1770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
310±50	1100	BELLINI	82 SPEC	-	$40\pi^-A \rightarrow 3\pi A$

 $\pi(1770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $f_0(1300)\pi$	dominant
Γ_2 $\rho\pi$	not seen

 $\pi(1770)$ BRANCHING RATIOS

$\Gamma(f_0(1300)\pi)/\Gamma_{\text{total}}$	Γ_1/Γ
dominant	

$\Gamma(\rho\pi)/\Gamma_{\text{total}}$	Γ_2/Γ
not seen	

 $\pi(1770)$ REFERENCES

BELLINI 82 PRL 48 1697	+Frabetti, Ivanshin, Litkin+	(MILA, BGNA, JINR)
------------------------	------------------------------	--------------------

 $X(1775)$

$$I^G(J^{PC}) = 1^-(?^{-+})$$

OMITTED FROM SUMMARY TABLE

Seen in the charge exchange photoproduction reactions $\gamma p \rightarrow (\rho\pi^+)(\pi^+\pi^-\pi^-)$, $\gamma p \rightarrow n\pi^+\pi^+\pi^-$. Needs confirmation.

 $X(1775)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1776±13 OUR AVERAGE			
1763±20	CONDO	91 SHF	$\gamma p \rightarrow (\rho\pi^+)(\pi^+\pi^-\pi^-)$
1787±18	CONDO	91 SHF	$\gamma p \rightarrow n\pi^+\pi^+\pi^-$

 $X(1775)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
155±40 OUR AVERAGE			
192±60	CONDO	91 SHF	$\gamma p \rightarrow (\rho\pi^+)(\pi^+\pi^-\pi^-)$
118±60	CONDO	91 SHF	$\gamma p \rightarrow n\pi^+\pi^+\pi^-$

 $X(1775)$ DECAY MODES

Mode
Γ_1 $\rho\pi$
Γ_2 $f_2(1270)\pi$

 $X(1775)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(f_2(1270)\pi)$	Γ_1/Γ_2
1.43±0.26 OUR AVERAGE	
1.3 ± 0.3	CONDO 91 SHF $\gamma p \rightarrow (\rho\pi^+)(\pi^+\pi^-\pi^-)$
1.8 ± 0.5	CONDO 91 SHF $\gamma p \rightarrow n\pi^+\pi^+\pi^-$

 $X(1775)$ REFERENCES

CONDO 91 PR D43 2787	+Handler+	(SLAC Hybrid Collab.)
----------------------	-----------	-----------------------

 $f_2(1810)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

From an amplitude analysis of the K^+K^- system seen in $\pi^-p \rightarrow K^+K^-n$ at 10 GeV/c. Confirmed by LONGACRE 86. Seen also in $\pi^+\pi^- \rightarrow 2\pi^0$ amplitude analysis (CASON 82), in the partial-wave analysis of the $\eta\eta$ system (ALDE 86D) and in the $4\pi^0$ mass spectrum (ALDE 88). Needs confirmation.

 $f_2(1810)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
1858 ⁺¹⁸ ₋₇₁	¹ LONGACRE	86 RVUE	Compilation
1799±15	CASON	82 STRC	$8\pi^+p \rightarrow \rho\pi^+2\pi^0$
1857 ⁺³⁵ ₋₂₄	² COSTA...	80 OMEG	$10\pi^-p \rightarrow K^+K^-n$

¹ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.² Error increased by spread of two solutions. Included in LONGACRE 86 global analysis. **$f_2(1810)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
388 ⁺¹⁵ ₋₂₁	³ LONGACRE	86 RVUE	Compilation
280 ⁺⁴² ₋₃₅	CASON	82 STRC	$8\pi^+p \rightarrow \rho\pi^+2\pi^0$
185 ⁺¹⁰² ₋₁₃₉	⁴ COSTA...	80 OMEG	$10\pi^-p \rightarrow K^+K^-n$

³ From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.⁴ Error increased by spread of two solutions. Included in LONGACRE 86 global analysis.

See key on page 1343

Meson Full Listings

$f_2(1810)$, $X(1830)$, $\phi_3(1850)$

 $f_2(1810)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	$(21.0^{+2.0}_{-3.0})\%$
Γ_2 $\eta\eta$	$(8.0^{+28.0}_{-3.0}) \times 10^{-3}$
Γ_3 $4\pi^0$	
Γ_4 K^+K^-	$(3.0^{+19.0}_{-2.0}) \times 10^{-3}$

 $f_2(1810)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.21 $^{+0.02}_{-0.03}$	5 LONGACRE	86 RVUE	Compilation	
••• We do not use the following data for averages, fits, limits, etc. •••				
0.44 ± 0.03	6 CASON	82 STRC	$8\pi^+\rho \rightarrow \rho\pi^+\pi^0$	

$\Gamma(\eta\eta)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.008 $^{+0.028}_{-0.003}$	5 LONGACRE	86 RVUE	Compilation	

$\Gamma(K^+K^-)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.003 $^{+0.019}_{-0.002}$	5 LONGACRE	86 RVUE	Compilation	
••• We do not use the following data for averages, fits, limits, etc. •••				
seen	COSTA...	80 OMEG	$10\pi^-\rho \rightarrow K^+K^-\eta$	

⁵From a partial-wave analysis of data using a K-matrix formalism with 5 poles. Includes compilation of several other experiments.
⁶Included in LONGACRE 86 global analysis.

 $f_2(1810)$ REFERENCES

ALDE	88	PL B201 160	+Bellazzini, Binon+ (SERP, BELG, LANL, LAPP, PISA)
ALDE	86D	NP B269 485	+Binon, Bricman+ (BELG, LAPP, SERP, CERN)
LONGACRE	86	PL B177 223	+Etkin+ (BNL, BRAN, CUNY, DUKE, NDAM)
CASON	82	PRL 48 1316	+Biswas, Baumbaugh, Bishop+ (NDAM, ANL)
COSTA...	80	NP B175 402	Costa De Beauregard+ (BARI, BONN, CERN+)

OTHER RELATED PAPERS

AKER	91	PL B260 249	+Amsler, Peters+ (Crystal Barrel Collab.)
ALDE	88	PL B201 160	+Bellazzini, Binon+ (SERP, BELG, LANL, LAPP, PISA)
ALDE	86D	NP B269 485	+Binon, Bricman+ (BELG, LAPP, SERP, CERN)
CASON	83	PR D28 1586	+Cannata, Baumbaugh, Bishop+ (NDAM, ANL)
ETKIN	82B	PR D25 1786	+Foley, Lai+ (BNL, CUNY, TUFTS, VAND)

 $X(1830)$

$$I^G(J^{PC}) = 1^-(?^{?+})$$

OMITTED FROM SUMMARY TABLE

Observed in coherent production on a carbon and beryllium nucleus.
 $J^{PC} = 1^{++}$ and 2^{-+} preferred. Needs confirmation.

 $X(1830)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1832 ± 27 OUR AVERAGE	Error includes scale factor of 1.3.			
1873 $\pm 33 \pm 20$		BELADIDZE	92C VES	$36\pi^-\text{Be} \rightarrow \pi^-\eta'\eta\text{Be}$
1814 $\pm 10 \pm 23$	426 ± 57	BITYUKOV	91 VES	$36\pi^-\text{C} \rightarrow \pi^-\eta\eta\text{C}$

 $X(1830)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
214 ± 27 OUR AVERAGE				
225 $\pm 35 \pm 20$		BELADIDZE	92C VES	$36\pi^-\text{Be} \rightarrow \pi^-\eta'\eta\text{Be}$
205 $\pm 18 \pm 32$	426 ± 57	BITYUKOV	91 VES	$36\pi^-\text{C} \rightarrow \pi^-\eta\eta\text{C}$

 $X(1830)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\eta\eta$	
Γ_2 $\pi\eta\eta(958)$	
Γ_3 $\pi f_0(1590)$	seen

 $X(1830)$ BRANCHING RATIOS

$\Gamma(\pi\eta\eta(958))/\Gamma(\pi\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.29 ± 0.06 OUR AVERAGE				
0.29 ± 0.07	BELADIDZE	92C VES	$36\pi^-\text{Be} \rightarrow \pi^-\eta'\eta\text{Be}$	
0.3 ± 0.1	BITYUKOV	91 VES	$36\pi^-\text{C} \rightarrow \pi^-\eta\eta\text{C}$	

 $X(1830)$ REFERENCES

BELADIDZE	92C	SJNP 55 1535	+Bityukov, Borisov (VES Collab.)
		Translated from YAF 55 2748.	
BITYUKOV	91	PL B268 137	+Borisov+ (SERP, TBL)

OTHER RELATED PAPERS

BORISOV	92	SJNP 55 1441	+Gershtein, Zaitsev (SERP)
		Translated from YAF 55 2583.	

 $\phi_3(1850)$

$$I^G(J^{PC}) = 0^-(3^{--})$$

Seen in the $K\bar{K}$ and $K\bar{K}\pi$ mass distributions. **$\phi_3(1850)$ MASS**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1854 ± 7 OUR AVERAGE				
1855 ± 10		ASTON	88E LASS	$11K^-\rho \rightarrow K^-K^+\Lambda, K_S^0 K^\pm \pi^\mp \Lambda$
1870.0 $^{+30.0}_{-20.0}$	430	ARMSTRONG	82 OMEG	$18.5K^-\rho \rightarrow K^-K^+\Lambda$
1850.0 ± 10.0	123	ALHARRAN	81B HBC	$8.25K^-\rho \rightarrow K\bar{K}\Lambda$

 $\phi_3(1850)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
87 $^{+28}_{-23}$ OUR AVERAGE	Error includes scale factor of 1.2.			
64 ± 31		ASTON	88E LASS	$11K^-\rho \rightarrow K^-K^+\Lambda, K_S^0 K^\pm \pi^\mp \Lambda$
160.0 $^{+90.0}_{-50.0}$	430	ARMSTRONG	82 OMEG	$18.5K^-\rho \rightarrow K^-K^+\Lambda$
80.0 $^{+40.0}_{-30.0}$	123	ALHARRAN	81B HBC	$8.25K^-\rho \rightarrow K\bar{K}\Lambda$

 $\phi_3(1850)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\bar{K}$	seen
Γ_2 $K\bar{K}^*(892) + \text{c.c.}$	seen

 $\phi_3(1850)$ BRANCHING RATIOS

$\Gamma(K\bar{K}^*(892) + \text{c.c.})/\Gamma(K\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.55 $^{+0.85}_{-0.45}$	ASTON	88E LASS	$11K^-\rho \rightarrow K^-K^+\Lambda, K_S^0 K^\pm \pi^\mp \Lambda$	
0.8 ± 0.4	ALHARRAN	81B HBC	$8.25K^-\rho \rightarrow K\bar{K}\pi\Lambda$	

••• We do not use the following data for averages, fits, limits, etc. •••

 $\phi_3(1850)$ REFERENCES

ASTON	88E	PL B208 324	+Awaji, Biewz+ (SLAC, NAGO, CINC, INUS) IGJPC
ARMSTRONG	82	PL 110B 77	+Baubillier+ (BARI, BIRM, CERN, MILA, CURIN+) JP
ALHARRAN	81B	PL 101B 357	+Amirzadeh+ (BIRM, CERN, GLAS, MICH, CURIN)

OTHER RELATED PAPERS

CORDIER	82B	PL 110B 335	+Bisello, Bizot, Buon, Delcourt, Fayard+ (LALO)
ASTON	80B	PL 92B 219	(BONN, CERN, EPOL, GLAS, LANC, MCHS+)

Meson Full Listings

 $\eta_2(1870)$, $X(1910)$ $\eta_2(1870)$

$$I^G(J^{PC}) = 0^+(2^{-+})$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

 $\eta_2(1870)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1881 ± 32 ± 40	26	KARCH	92	CBAL $e^+e^- \rightarrow e^+e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1850 ± 50		FEINDT	91	CELL $\gamma\gamma \rightarrow \eta \pi^+ \pi^-$

 $\eta_2(1870)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
221 ± 92 ± 44	26	KARCH	92	CBAL $e^+e^- \rightarrow e^+e^- \eta \pi^0 \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 360		FEINDT	91	CELL $\gamma\gamma \rightarrow \eta \pi^+ \pi^-$

 $\eta_2(1870)$ DECAY MODES

Mode
Γ_1 $\eta \pi \pi$
Γ_2 $a_2(1320) \pi$
Γ_3 $f_0(980) \eta$

 $\eta_2(1870)$ REFERENCES

KARCH	92	ZPHY C54 33	+Antreasyan, Bartels+	(Crystal Ball Collab.)
FEINDT	91	Singapore Conf. 537		

OTHER RELATED PAPERS

KARCH	90	PL B249 353	+Antreasyan, Bartels+	(Crystal Ball Collab.)
-------	----	-------------	-----------------------	------------------------

 $X(1910)$

$$I^G(J^{PC}) = 0^+(?^{?+})$$

OMITTED FROM SUMMARY TABLE

We list here several bumps seen in the mass distributions of different final states.

 $X(1910)$ MASS

VALUE (MeV)	DOCUMENT ID
1810 to 1920 OUR ESTIMATE	

 $X(1910)$ $4\pi^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1806 ± 10	1600 ± 100	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$
1870 ± 40		ALDE	86D	GAM4 $100 \pi^- p \rightarrow 4\gamma n$

 $X(1910)$ $\omega\omega$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1920 ± 10	BELADIDZE	92B VES	$36 \pi^- p \rightarrow \omega\omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1924 ± 14	ALDE	90	GAM2 $38 \pi^- p \rightarrow n\omega\omega$

 $X(1910)$ $\eta\eta'$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1911 ± 10	ALDE	91B	GAM2 $38 \pi^- p \rightarrow \eta\eta' n$

 $X(1910)$ WIDTH

VALUE (MeV)	DOCUMENT ID
90 to 250 OUR ESTIMATE	

 $X(1910)$ $4\pi^0$ MODE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
190 ± 20	1600 ± 100	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$
250 ± 30		ALDE	86D	GAM4 $100 \pi^- p \rightarrow 4\gamma n$

 $X(1910)$ $\omega\omega$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
90 ± 20	BELADIDZE	92B VES	$36 \pi^- p \rightarrow \omega\omega n$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
91 ± 50	ALDE	90	GAM2 $38 \pi^- p \rightarrow n\omega\omega$

 $X(1910)$ $\eta\eta'$ MODE

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
90 ± 35	ALDE	91B	GAM2 $38 \pi^- p \rightarrow \eta\eta' n$

 $X(1910)$ DECAY MODES

Mode
Γ_1 $4\pi^0$
Γ_2 $\pi\pi$
Γ_3 $\pi^0\pi^0$
Γ_4 $K_S^0 K_S^0$
Γ_5 $\eta\eta$
Γ_6 $\omega\omega$
Γ_7 $\eta\eta'$
Γ_8 $\eta\pi\pi$
Γ_9 $\eta'\eta'$

 $X(1910)$ BRANCHING RATIOS

$\Gamma(\pi^0\pi^0)/\Gamma(4\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.25	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$	

$\Gamma(4\pi^0)/\Gamma(\eta\eta)$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_5
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.8 ± 0.3	ALDE	87	GAM4 $100 \pi^- p \rightarrow 4\pi^0 n$	

$\Gamma(\omega\omega)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	ALDE	89B	GAM2 $38 \pi^- p \rightarrow n\omega\omega$	

$\Gamma(\pi^0\pi^0)/\Gamma(\eta\eta')$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_7
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.1	ALDE	89	GAM2 $38 \pi^- p \rightarrow n\eta\eta'$	

$\Gamma(\eta\eta)/\Gamma(\eta\eta')$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_7
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.05	90	ALDE	91B	GAM2 $38 \pi^- p \rightarrow n\eta\eta'$	

$\Gamma(K_S^0 K_S^0)/\Gamma(\eta\eta')$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_7
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.066	90	BALOSHIN	86	SPEC $40 \pi^+ p \rightarrow K_S^0 K_S^0 n$

$\Gamma(\eta'\eta')/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
possibly seen	BELADIDZE	92D VES	$37 \pi^- p \rightarrow \eta'\eta' n$	

 $X(1910)$ REFERENCES

BELADIDZE	92B	ZPHY C54 367	+Bitjukov, Borisov+	(VES Collab.)
BELADIDZE	92D	ZPHY C57 13	+Berdnikov+	(VES Collab.)
ALDE	91B	SJNP 54 455	+Binon+	(SERP, BELG, LANL, LAPP, PISA, KEK)
ALDE	90	Translated from YAF 54 751.		
ALDE	89	PL B241 600	+Binon+	(SERP, BELG, LANL, LAPP, PISA, KEK)
ALDE	89B	PL B216 447	+Binon, Bricman, Donskov+	(SERP, BELG, LANL, LAPP)
ALDE	89B	PL B216 451	+Binon, Bricman+	(SERP, BELG, LANL, LAPP, TBIL)
ALDE	87	PL B198 286	+Binon, Bricman+	(LANL, BRUX, SERP, LAPP)
ALDE	86D	NP B269 485	+Binon, Bricman+	(BELG, LAPP, SERP, CERN)
BALOSHIN	86	SJNP 43 959	+Barkov, Bolonkin, Vladimirov, Grigoriev+	(ITEP)
			Translated from YAF 43 1487.	

See key on page 1343

Meson Full Listings

$X(1950)$, $f_2(2010)$, $a_4(2040)$

$X(1950)$

$$I^G(J^{PC}) = 0^+(\text{even}^{++})$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

$X(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1952 ± 14 OUR AVERAGE				
1964 ± 35	ARMSTRONG 93D SPEC			$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
1950 ± 15	ASTON 91 LASS 0			$11 K^- p \rightarrow \Lambda K \bar{K} \pi \pi$

$X(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
238 ± 35 OUR AVERAGE				
225 ± 50	ARMSTRONG 93D SPEC			$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$
250 ± 50	ASTON 91 LASS 0			$11 K^- p \rightarrow \Lambda K \bar{K} \pi \pi$

$X(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K^*(892) \bar{K}^*(892)$	seen
$\Gamma_2 \pi^0 \pi^0$	seen

$X(1950)$ BRANCHING RATIOS

$\Gamma(K^*(892) \bar{K}^*(892))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
seen	ASTON 91 LASS 0			$11 K^- p \rightarrow \Lambda K \bar{K} \pi \pi$	

$\Gamma(\pi^0 \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen	ARMSTRONG 93D SPEC		$\bar{p}p \rightarrow 3\pi^0 \rightarrow 6\gamma$	

$X(1950)$ REFERENCES

ARMSTRONG 93D	PL B307 399	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
ASTON 91	NP B21 5 (suppl)	+Awaji+	(LASS Collab.)

OTHER RELATED PAPERS

BIENZ 90	SLAC 369		(LASS Collab.)
ALBRECHT 88N	PL B212 528	+	(ARGUS Collab.)
ALBRECHT 87Q	PL B198 255	+Binder+	(ARGUS Collab.)
ARMSTRONG 87C	ZPHY C34 33	+Bloodworth+	(CERN, BIRM, BARI, ATHU, CURIN+)

$f_2(2010)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)

$f_2(2010)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2011 ± 62	1 ETKIN 88 MPS		$22 \pi^- p \rightarrow \phi \phi n$
1980 ± 20	2 BOLONKIN 88 SPEC		$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
2050.0 ± 90.0 -50.0	ETKIN 85 MPS		$22 \pi^- p \rightarrow 2\phi n$
2120.0 ± 20.0 -120.0	LINDENBAUM 84 RVUE		
2160.0 ± 50.0	ETKIN 82 MPS		$22 \pi^- p \rightarrow 2\phi n$

¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi 2^{++} S_2$, D_2 , and D_0 is 98 ± 1 , 0 ± 1 , and 2 ± 1 , respectively.
² Statistically very weak, only 1.4 s.d.

$f_2(2010)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
202 ± 67	3 ETKIN 88 MPS		$22 \pi^- p \rightarrow \phi \phi n$
145 ± 50	4 BOLONKIN 88 SPEC		$40 \pi^- p \rightarrow K_S^0 K_S^0 n$
200.0 ± 160.0 -50.0	ETKIN 85 MPS		$22 \pi^- p \rightarrow 2\phi n$
300.0 ± 150.0 -50.0	LINDENBAUM 84 RVUE		
310.0 ± 70.0	ETKIN 82 MPS		$22 \pi^- p \rightarrow 2\phi n$

³ Includes data of ETKIN 85.
⁴ Statistically very weak, only 1.4 s.d.

$f_2(2010)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \phi \phi$	seen

$f_2(2010)$ REFERENCES

BOLONKIN 88	NP B309 426	+Bloshenko, Gorin+	(ITEP, SERP)
ETKIN 88	PL B201 568	+Foley, Lindenbaum+	(BNL, CUNY)
ETKIN 85	PL 165B 217	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
LINDENBAUM 84	CNPP 13 285		(CUNY)
ETKIN 82	PRL 49 1620	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
Also 83	Brighton Conf. 351	Lindenbaum	(BNL, CUNY)

OTHER RELATED PAPERS

ARMSTRONG 89B	PL B221 221	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)
GREEN 86	PRL 56 1639	+Lai+ (FNAL, ARIZ, FSU, NDAM, TUFTS, VAND+)
BOOTH 84	NP B242 51	+Ballance, Carroll, Donald+ (LIVP, GLAS, CERN)

$a_4(2040)$

$$I^G(J^{PC}) = 1^-(4^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K\bar{K}$ and $\pi^+\pi^-\pi^0$ systems.
Needs confirmation.

$a_4(2040)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2037 ± 26 OUR AVERAGE				
2040.0 ± 30.0	1 CLELAND 82B SPEC			$\pm 50 \pi p \rightarrow K_S^0 K^\pm p$
2030.0 ± 50.0	2 CORDEN 78C OMEG 0			$15 \pi^- p \rightarrow 3\pi n$
1903.0 ± 10.0	3 BALDI 78 SPEC			$-10 \pi^- p \rightarrow p K_S^0 K^-$

¹ From an amplitude analysis.
² $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.
³ From a fit to the Y_8^0 moment. Limited by phase space.

$a_4(2040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
427 ± 120 OUR AVERAGE				
380.0 ± 150.0	4 CLELAND 82B SPEC			$\pm 50 \pi p \rightarrow K_S^0 K^\pm p$
510.0 ± 200.0	5 CORDEN 78C OMEG 0			$15 \pi^- p \rightarrow 3\pi n$
166.0 ± 43.0	6 BALDI 78 SPEC			$-10 \pi^- p \rightarrow p K_S^0 K^-$

⁴ From an amplitude analysis.
⁵ $J^P = 4^+$ is favored, though $J^P = 2^+$ cannot be excluded.
⁶ From a fit to the Y_8^0 moment. Limited by phase space.

$a_4(2040)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 K \bar{K}$	seen
$\Gamma_2 \pi^+\pi^-\pi^0$	seen

$a_4(2040)$ BRANCHING RATIOS

$\Gamma(K \bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
seen	BALDI 78 SPEC			$\pm 10 \pi^- p \rightarrow K_S^0 K^\pm p$	

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ
seen	CORDEN 78C OMEG 0			$15 \pi^- p \rightarrow 3\pi n$	

$a_4(2040)$ REFERENCES

CLELAND 82B	NP B208 228	+Delfosse, Dorsaz, Gloor	(DURH, GEVA, LAUS, PITT)
BALDI 78	PL 74B 413	+Bohringer, Dorsaz, Hungerbuhler+	(GEVA) JP
CORDEN 78C	NP B136 77	+Dowell, Garvey+	(BIRM, RHEL, TELA, LOWC) JP

OTHER RELATED PAPERS

DELFOSSO 81	NP B183 349	+Guisan, Martin, Muhlemann, Weill+	(GEVA, LAUS)
-------------	-------------	------------------------------------	--------------

Meson Full Listings

$a_3(2050)$, $f_4(2050)$

$a_3(2050)$

$$I^G(J^{PC}) = 1^-(3^{++})$$

OMITTED FROM SUMMARY TABLE
Needs confirmation.

$a_3(2050)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2080 ± 40	208	KALELKAR	75	HBC	+ 15 $\pi^+ \rho^- \rightarrow \rho \pi^+ \rho_3$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 2100		ANTIPOV	77	CIBS	- 25 $\pi^- \rho^- \rightarrow \rho \pi^- \rho_3$
2214 ± 15		BALTAY	77	HBC	0 15 $\pi^- \rho^- \rightarrow \Delta^{++} 3\pi$

$a_3(2050)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
340 ± 80	208	KALELKAR	75	HBC	+ 15 $\pi^+ \rho^- \rightarrow \rho \pi^+ \rho_3$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 500		ANTIPOV	77	CIBS	- 25 $\pi^- \rho^- \rightarrow \rho \pi^- \rho_3$
355 ± 21		BALTAY	77	HBC	0 15 $\pi^- \rho^- \rightarrow \Delta^{++} 3\pi$

$a_3(2050)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	
Γ_2 $\rho_3(1690)\pi$	dominant

$a_3(2050)$ BRANCHING RATIOS

$\Gamma(\rho_3(1690)\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
dominant	KALELKAR	75	HBC	+ 15 $\pi^+ \rho^- \rightarrow \rho_3\pi$	

$a_3(2050)$ REFERENCES

ANTIPOV	77	NP B119 45	+Busnello, Damgaard, Kienzie+	(SERP, GEVA)
BALTAY	77	PRL 39 591	+Cautis, Kalelkar	(COLU) JP
KALELKAR	75	Nevis 207 Thesis		(COLU)

OTHER RELATED PAPERS

HARRIS	81	ZPHY C9 275	+Dunn, Lubatti, Moriyasu, Podolsky+	(SEAT, UCB)
HUSON	68	PL 28B 208	+Lubatti, Six, Veillet+	(ORSAY, MILA, UCLA)
DANYSZ	67B	NC 51A 801	+French, Simak	(CERN)

$f_4(2050)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

$f_4(2050)$ MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2044 ± 11	OUR AVERAGE	Error includes scale factor of 1.4. See the ideogram below.		
1970 ± 30		BELADIDZE	92B	VES 36 $\pi^- \rho^- \rightarrow \omega \omega n$
2060 ± 20		ALDE	90	GAM2 38 $\pi^- \rho^- \rightarrow n \omega \omega$
2038 ± 30		AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
2086 ± 15		BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
2000.0 ± 60.0		ALDE	86D	GAM4 100 $\pi^- \rho^- \rightarrow n 2\eta$
2020.0 ± 20.0	40k	¹ BINON	84B	GAM2 38 $\pi^- \rho^- \rightarrow n 2\pi^0$
2015.0 ± 28.0		¹ CASON	82	STRC 8 $\pi^+ \rho^- \rightarrow \rho \pi^+ 2\pi^0$
2031.0 ⁺²⁵ ₋₃₆		ETKIN	82B	MPS 23 $\pi^- \rho^- \rightarrow n 2K^0_S$
2020 ± 30	700	APEL	75	NICE 40 $\pi^- \rho^- \rightarrow n 2\pi^0$
2050 ± 25		BLUM	75	ASPK 18.4 $\pi^- \rho^- \rightarrow n K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1978.0 ± 5.0		² ALPER	80	CNTR 62 $\pi^- \rho^- \rightarrow K^+ K^- n$
2040.0 ± 10.0		² ROZANSKA	80	SPRK 18 $\pi^- \rho^- \rightarrow p \bar{p} n$
1935.0 ± 13.0		² CORDEN	79	OMEG 12-15 $\pi^- \rho^- \rightarrow n 2\pi$
1988.0 ± 7.0		EVANGELISTA	79B	OMEG 10 $\pi^- \rho^- \rightarrow K^+ K^- n$
1922.0 ± 14.0		³ ANTIPOV	77	CIBS 25 $\pi^- \rho^- \rightarrow \rho_3\pi$

¹ From amplitude analysis of reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.

² $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.

³ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

WEIGHTED AVERAGE
2044±11 (Error scaled by 1.4)



$f_4(2050)$ mass (MeV)

$f_4(2050)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
208 ± 13	OUR AVERAGE	Error includes scale factor of 1.2.		
300 ± 50		BELADIDZE	92B	VES 36 $\pi^- \rho^- \rightarrow \omega \omega n$
170 ± 60		ALDE	90	GAM2 38 $\pi^- \rho^- \rightarrow n \omega \omega$
304 ± 60		AUGUSTIN	87	DM2 $J/\psi \rightarrow \gamma \pi^+ \pi^-$
210 ± 63		BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma \pi^+ \pi^-$
400.0 ± 100.0		ALDE	86D	GAM4 100 $\pi^- \rho^- \rightarrow n 2\eta$
140.0 ± 40.0	40k	⁴ BINON	84B	GAM2 38 $\pi^- \rho^- \rightarrow n 2\pi^0$
190.0 ± 14.0		DENNEY	83	LASS 10 $\pi^+ n/\pi^+ p$
186.0 ^{+103.0} _{-58.0}		⁴ CASON	82	STRC 8 $\pi^+ \rho^- \rightarrow \rho \pi^+ 2\pi^0$
305.0 ⁺³⁶ ₋₁₁₉		ETKIN	82B	MPS 23 $\pi^- \rho^- \rightarrow n 2K^0_S$
180 ± 60	700	APEL	75	NICE 40 $\pi^- \rho^- \rightarrow n 2\pi^0$
225 ⁺¹²⁰ ₋₇₀		BLUM	75	ASPK 18.4 $\pi^- \rho^- \rightarrow n K^+ K^-$

• • • We do not use the following data for averages, fits, limits, etc. • • •

243.0 ± 16.0		⁵ ALPER	80	CNTR 62 $\pi^- \rho^- \rightarrow K^+ K^- n$
140.0 ± 15.0		⁵ ROZANSKA	80	SPRK 18 $\pi^- \rho^- \rightarrow p \bar{p} n$
263.0 ± 57.0		⁵ CORDEN	79	OMEG 12-15 $\pi^- \rho^- \rightarrow n 2\pi$
100.0 ± 28.0		EVANGELISTA	79B	OMEG 10 $\pi^- \rho^- \rightarrow K^+ K^- n$
107.0 ± 56.0		⁶ ANTIPOV	77	CIBS 25 $\pi^- \rho^- \rightarrow \rho_3\pi$

⁴ From amplitude analysis of reaction $\pi^+ \pi^- \rightarrow 2\pi^0$.

⁵ $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.

⁶ Width errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.

$f_4(2050)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\omega \omega$	(26 ± 6) %
Γ_2 $\pi \pi$	(17.0 ± 1.5) %
Γ_3 $K \bar{K}$	(6.8 ^{+3.4} _{-1.8}) × 10 ⁻³
Γ_4 $\eta \eta$	(2.1 ± 0.8) × 10 ⁻³
Γ_5 $4\pi^0$	< 1.2 %
Γ_6 $\gamma \gamma$	

$f_4(2050)$ $\Gamma(I)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K \bar{K}) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_3 \Gamma_6/\Gamma$
< 0.29	95	ALTHOFF	85B	TASS $\gamma \gamma \rightarrow K \bar{K} \pi$	

$\Gamma(\pi \pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_2 \Gamma_6/\Gamma$
< 1.1	95	13 ± 4	OEST	90	JADE $e^+ e^- \rightarrow e^+ e^- \pi^0 \pi^0$	

$f_4(2050)$ BRANCHING RATIOS

$\Gamma(\omega \omega)/\Gamma(\pi \pi)$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
1.5 ± 0.3	ALDE	90	GAM2 38 $\pi^- \rho^- \rightarrow n \omega \omega$	

See key on page 1343

Meson Full Listings

$f_4(2050)$, $\pi_2(2100)$, $f_2(2150)$

$\Gamma(\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE				
0.170 ± 0.015 OUR AVERAGE				
0.18 ± 0.03	⁷ BINON	83C	GAM2 38 $\pi^- p \rightarrow n4\gamma$	
0.16 ± 0.03	⁷ CASON	82	STRC 8 $\pi^+ p \rightarrow \rho\pi^+ 2\pi^0$	
0.17 ± 0.02	⁷ CORDEN	79	OMEG 12-15 $\pi^- p \rightarrow n2\pi$	
⁷ Assuming one pion exchange.				
$\Gamma(K\bar{K})/\Gamma(\pi\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_2
VALUE				
0.04 ± 0.02	ETKIN	82B	MPS 23 $\pi^- p \rightarrow n2K_S^0$	
$\Gamma(\eta\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
VALUE (units 10 ⁻³)				
2.1 ± 0.8	ALDE	86D	GAM4 100 $\pi^- p \rightarrow n4\gamma$	
$\Gamma(4\pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
VALUE				
< 0.012	ALDE	87	GAM4 100 $\pi^- p \rightarrow 4\pi^0 n$	

 $f_4(2050)$ REFERENCES

BELADIDZE	92B	ZPHY C54 367	+Bityukov, Borisov+	(VES Collab.)
ALDE	90	PL B241 600	+Binon+	(SERP, BELG, LANL, LAPP, PISA, KEK)
OEST	90	ZHPY C47 343	+Olsson+	(JADE Collab.)
ALDE	87	PL B198 286	+Binon, Bricman+	(LANL, BRUX, SERP, LAPP)
AUGUSTIN	87	ZPHY C36 369	+Cosme+	(LALO, CLER, FRAS, PADO)
BALTRUSAITIS...	87	PR D35 2077	+Baltrusaitis, Coffman, Dubois+	(Mark III Collab.)
ALDE	86D	NP B269 485	+Binon, Bricman+	(BELG, LAPP, SERP, CERN)
ALTHOFF	85B	ZPHY C29 189	+Braunschweig, Kirschfink+	(TASSO Collab.)
BINON	84B	LNC 39 41	+Donskov, Duteil, Gouanere+	(SERP, BELG, LAPP)
BINON	83C	SJNP 38 723	+Gouanere, Donskov, Duteil+	(SERP, BRUX+)
DENNEY	83	PR D28 2726	Translated from YAF 38 1199.	
CASON	82	PR D22 1503	+Cranley, Firestone, Chapman+	(IOWA, MICH)
ETKIN	82B	PR D25 1786	+Biswas, Baumbaugh, Bishop+	(NDAM, ANL)
ALPER	80	PL 94B 422	+Foley, Lai+	(BNL, CUNY, TUFTS, VAND)
ROZANSKA	80	NP B162 505	+Becker+	(AMST, CERN, CRAC, MPIM, OXF+)
CORDEN	79	NP B157 250	+Blum, Dietl, Grayer, Lorenz+	(MPI, CERN)
EVANGELISTA	79B	NP B154 381	+Dowell, Garvey+	(BIRM, RHEL, TELA, LOWC) JP
ANTIPOV	77	NP B119 45	+ (BARI, BONN, CERN, DARE, GLAS, LIVP+)	
APEL	75	PL 57B 398	+Busnelo, Damgaard, Kienzle+	(SERP, GEVA)
BLUM	75	PL 57B 403	+Augenstein+(KARLK, KARLE, PISA, SERP, WIEN, CERN)JP	
			+Chabaud, Dietl, Garelick, Grayer+	(CERN, MPIM) JP

OTHER RELATED PAPERS

CASON	83	PR D28 1586	+Cannata, Baumbaugh, Bishop+	(NDAM, ANL)
GÖTTESMAN	80	PR D22 1503	+Jacobs+	(SYRA, BRAN, BNL, CINC)
WAGNER	74	London Conf. 2 27		(MPIM)

 $\pi_2(2100)$

$$I^G(J^{PC}) = 1^-(2^{-+})$$

OMITTED FROM SUMMARY TABLE

Seen in the $\rho\pi$, $f_0(1300)\pi$, and $f_2(1270)\pi$ $J^P = 2^-$ waves of the diffractively produced 3π system. Needs confirmation. **$\pi_2(2100)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2100 ± 180	¹ DAUM	81B	CNTR 63,94 $\pi^- p \rightarrow 3\pi X$

¹ From a two-resonance fit to four 2^-0^+ waves. **$\pi_2(2100)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
651 ± 50	² DAUM	81B	CNTR 63,94 $\pi^- p \rightarrow 3\pi X$

² From a two-resonance fit to four 2^-0^+ waves. **$\pi_2(2100)$ DECAY MODES**

Mode	Fraction (Γ_i/Γ)
Γ_1 3π	seen
Γ_2 $\rho\pi$	seen
Γ_3 $f_2(1270)\pi$	seen
Γ_4 $f_0(1300)\pi$	seen

 $\pi_2(2100)$ BRANCHING RATIOS

$\Gamma(\rho\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE				
0.19 ± 0.05	³ DAUM	81B	CNTR 63,94 $\pi^- p$	
$\Gamma(f_2(1270)\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
VALUE				
0.36 ± 0.09	³ DAUM	81B	CNTR 63,94 $\pi^- p$	

$\Gamma(f_0(1300)\pi)/\Gamma(3\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
VALUE				
0.45 ± 0.07	³ DAUM	81B	CNTR 63,94 $\pi^- p$	
D-wave/S-wave RATIO FOR $\pi_2(2100) \rightarrow f_2(1270)\pi$	DOCUMENT ID	TECN	COMMENT	
VALUE				
0.39 ± 0.23	³ DAUM	81B	CNTR 63,94 $\pi^- p$	

³ From a two-resonance fit to four 2^-0^+ waves. **$\pi_2(2100)$ REFERENCES**

DAUM	81B	NP B182 269	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)
------	-----	-------------	---------------	--------------------------------

 $f_2(2150)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called T_0 . **$f_2(2150)$ MASS** **$\bar{p}p \rightarrow \pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 2170.0	¹ MARTIN	80B	RVUE
~ 2150.0	¹ MARTIN	80C	RVUE
~ 2150.0	² DULUDE	78B	OSPCK 1-2 $\bar{p}p \rightarrow \pi^0\pi^0$

¹ (J^P) = 0(2⁺) from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0\pi^0$.² (J^P) = 0⁺(2⁺) from partial-wave amplitude analysis.**S-CHANNEL $\bar{p}p$ or $\bar{N}N$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 2190.0	³ CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155.0 ± 15.0	^{3,4} COUPLAND	77	CNTR	0 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{3,5} ALSPECTOR	73	CNTR	0 $\bar{p}p$ S channel

³ Isospins 0 and 1 not separated.⁴ From a fit to the total elastic cross section.⁵ Referred to as T or T region by ALSPECTOR 73.**OTHER HADRONIC MODES**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
2104 ± 20	ARMSTRONG	93C	SPEC $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
2175 ± 20	PROKOSHKIN	90	GAM4 300 $\pi^- N \rightarrow \pi^- N2\eta$, 450 $pN \rightarrow pN2\eta$

 $f_2(2150)$ WIDTH **$\bar{p}p \rightarrow \pi\pi$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 250.0	⁶ MARTIN	80B	RVUE
~ 250.0	⁶ MARTIN	80C	RVUE
~ 250.0	⁷ DULUDE	78B	OSPCK 1-2 $\bar{p}p \rightarrow \pi^0\pi^0$

⁶ (J^P) = 0(2⁺) from simultaneous analysis of $p\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0\pi^0$.⁷ (J^P) = 0⁺(2⁺) from partial-wave amplitude analysis.**S-CHANNEL $\bar{p}p$ or $\bar{N}N$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
135.0 ± 75.0	^{8,9} COUPLAND	77	CNTR	0 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
98 ± 8	⁹ ALSPECTOR	73	CNTR	0 $\bar{p}p$ S channel

⁸ From a fit to the total elastic cross section.⁹ Isospins 0 and 1 not separated.**OTHER HADRONIC MODES**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
203 ± 10	ARMSTRONG	93C	SPEC $\bar{p}p \rightarrow \pi^0\eta\eta \rightarrow 6\gamma$
150 ± 35	PROKOSHKIN	90	GAM4 300 $\pi^- N \rightarrow \pi^- N2\eta$, 450 $pN \rightarrow pN2\eta$

 $f_2(2150)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\pi\pi$	seen

Meson Full Listings

$f_2(2150)$, $\rho(2150)$, $X(2200)$, $\rho(2210)$

$f_2(2150)$ REFERENCES

ARMSTRONG 93C	PL B307 394	+Bettoni+	(FNAL, FERR, GENO, UCI, NWES+)
PROKOSHKIN 90	Hadron 89 Conf. p 27	(SERP, BELG, LANL, LAPP, PISA, KEK)	
MARTIN 80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	+Pennington	(DURH) JP
CUTTS 78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
DULUDE 78B	PL 79B 335	+Lanou, Massimo, Peaslee+	(BROW, MIT, BARI) JP
COUPLAND 77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
ALSPECTOR 73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)

OTHER RELATED PAPERS

FIELDS 71	PRL 27 1749	+Cooper, Rhines, Allison	(ANL, OXF)
YOH 71	PRL 26 922	+Barish, Caroll, Lobkowicz+	(CIT, BNL, ROCH)

$\rho(2150)$

$$I^G(J^{PC}) = 1^+(1^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $T_1(2190)$.

Our latest mini-review on this particle can be found in the 1984 edition.

$\rho(2150)$ MASS

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN
2197 ± 17	AUGUSTIN 88 DM2 0	

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2170.0	2 MARTIN 80B RVUE
~ 2100.0	2 MARTIN 80C RVUE

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2155.0 ± 15.0	3,4 COUPLAND 77 CNTR 0			$0.97-3 \bar{p}p \rightarrow \bar{N}N$
2193 ± 2	3,5 ALSPECTOR 73 CNTR			$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
2190 ± 10	6 ABRAMS 70 CNTR			$\bar{p}p$ S channel
				S channel $\bar{p}N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 2190.0	3 CUTTS 78B CNTR			$0.97-3 \bar{p}p \rightarrow \bar{N}N$
---------------	------------------	--	--	--

$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2153 ± 37	BIAGINI 91 RVUE			$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$
2110 ± 50	1 CLEGG 90 RVUE 0			$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2110 ± 50	1 CLEGG 90 RVUE 0			$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$
---------------	-------------------	--	--	---

- Includes ATKINSON 85.
- $I(J^P) = 1(1^-)$ from simultaneous analysis of $\rho\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.
- Isospins 0 and 1 not separated.
- From a fit to the total elastic cross section.
- Referred to as T or T' region by ALSPECTOR 73.
- Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

$\rho(2150)$ WIDTH

$\bar{p}p \rightarrow \pi\pi$

VALUE (MeV)	DOCUMENT ID	TECN
~ 250.0	8 MARTIN 80B RVUE	
~ 200.0	8 MARTIN 80C RVUE	

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 250.0	8 MARTIN 80B RVUE
~ 200.0	8 MARTIN 80C RVUE

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
135.0 ± 75.0	9,10 COUPLAND 77 CNTR 0			$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
98 ± 8	10 ALSPECTOR 73 CNTR			$\bar{p}p$ S channel
~ 85	11 ABRAMS 70 CNTR			S channel $\bar{p}N$

• • • We do not use the following data for averages, fits, limits, etc. • • •

135.0 ± 75.0	9,10 COUPLAND 77 CNTR 0			$0.7-2.4 \bar{p}p \rightarrow \bar{p}p$
98 ± 8	10 ALSPECTOR 73 CNTR			$\bar{p}p$ S channel
~ 85	11 ABRAMS 70 CNTR			S channel $\bar{p}N$

$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
389 ± 79	BIAGINI 91 RVUE			$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$
410 ± 100	7 CLEGG 90 RVUE 0			$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

410 ± 100	7 CLEGG 90 RVUE 0			$e^+e^- \rightarrow \pi^+\pi^-, K^+K^-, 6\pi$
---------------	-------------------	--	--	---

- Includes ATKINSON 85.
- $I(J^P) = 1(1^-)$ from simultaneous analysis of $\rho\bar{p} \rightarrow \pi^-\pi^+$ and $\pi^0\pi^0$.
- From a fit to the total elastic cross section.
- Isospins 0 and 1 not separated.
- Seen as bump in $l = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.

$\rho(2150)$ REFERENCES

BIAGINI 91	NC 104A 363	+Dubnicka+	(FRAS, PRAG)
CLEGG 90	ZPHY C45 677	+Donnachie	(LANC, MCHS)
ATKINSON 85	ZPHY C29 333	+ (BONN, CERN, GLAS, LANL, MCHS, INP+)	
MARTIN 80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN 80C	NP B169 216	+Pennington	(DURH) JP
CUTTS 78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
COUPLAND 77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
PEASLEE 75	PL 57B 189	+Demarzo, Guerriero+	(CANB, BARI, BROW, MIT)
ALSPECTOR 73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)
ABRAMS 70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)
COOPER 68	PRL 20 1059	+Hyman, Manner, Musgrave+	(ANL)

OTHER RELATED PAPERS

BRICMAN 69	PL 29B 451	+Ferro-Luzzi, Bizard+	(CERN, CAEN, SACL)
ABRAMS 67C	PRL 18 1209	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)

$X(2200)$

$$I^G(J^{PC}) = ?^?(even^{++})$$

OMITTED FROM SUMMARY TABLE

Seen at DCI in the $K_S^0 K_S^0$ system. Not seen in T radiative decays (BARU 89). Needs confirmation.

$X(2200)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2197 ± 17	AUGUSTIN 88 DM2 0			$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$X(2200)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
201 ± 51	AUGUSTIN 88 DM2 0			$J/\psi \rightarrow \gamma K_S^0 K_S^0$

$X(2200)$ REFERENCES

BARU 89	ZPHY C42 505	+Beilin, Blinov+	(NOVO)
AUGUSTIN 88	PRL 60 2238	+Calcaterra+	(DM2 Collab.)

$\rho(2210)$

$$I^G(J^{PC}) = 1^+(1^{--})$$


OMITTED FROM SUMMARY TABLE

Seen in one-pion exchange production of the $\omega\pi^0$ system and diffractive photoproduction of the $\omega\pi^+\pi^-\pi^0$ system. Needs confirmation.

$\rho(2210)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2212 ± 22 OUR AVERAGE			Error includes scale factor of 1.4. See the ideogram below.
2220 ± 20	ALDE 92C GAM2 38		$\pi^-\pi^0 \rightarrow n\omega\pi^0$
2170 ± 30	ALDE 92C GAM4 100		$\pi^-\rho \rightarrow n\omega\pi^0$
2280 ± 50	ATKINSON 85 OMEG 20-70		$\gamma p \rightarrow \rho\omega\pi^+\pi^-\pi^0$

WEIGHTED AVERAGE
2212±22 (Error scaled by 1.4)



See key on page 1343

Meson Full Listings

$\rho(2210)$, $f_4(2220)$, $\eta(2225)$

 $\rho(2210)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
206 ± 80 OUR AVERAGE	Error includes scale factor of 1.6.		
240 ± 60	ALDE	92C GAM2	38 $\pi^- p \rightarrow n\omega\pi^0$
~ 300	ALDE	92C GAM4	100 $\pi^- p \rightarrow n\omega\pi^0$
440 ± 110	ATKINSON	85 OMEG	20-70 $\gamma p \rightarrow \rho\omega\pi^+\pi^-\pi^0$

 $\rho(2210)$ REFERENCES

ALDE	92C	ZPHY C54 553	+Bellazzini+ (SERP, BELG, LANL, LAPP, PISA, KEK)
ATKINSON	85	ZPHY C29 333	+ (BONN, CERN, GLAS, LANC, MCHS, INP+)

 $f_4(2220)$ was $\xi(2220)$

$$I^G(J^{PC}) = 0^+(4^{++})$$

OMITTED FROM SUMMARY TABLE

This state has been seen at SPEAR in the $K\bar{K}$ systems (K^+K^- and $K_S^0K_S^0$) produced in the radiative decay of $J/\psi(1S)$. Seen in $\eta\eta'$ (ALDE 86B), in $K_S^0K_S^0$ (ASTON 88D), and in K^+K^- (ASTON 88F). Needs confirmation. Also J needs confirmation. Not seen in T radiative decays nor in B inclusive decay (BEHREND 84). Not seen in $\bar{p}p \rightarrow K^+K^-$ formation experiment (BARDIN 87, SCULLI 87) and $\bar{p}p \rightarrow K_S^0K_S^0$ formation experiment (BARNES 93). Not seen at DCI in either K^+K^- or $K_S^0K_S^0$ systems (AUGUSTIN 88).

 $f_4(2220)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2225 ± 6 OUR AVERAGE				
2209 ± 17 ± 10		ASTON	88F LASS	11 $K^- p \rightarrow K^+K^-A$
2230 ± 20		BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0K_S^0n$
2220 ± 10	41	ALDE	86B GAM4	38-100 $\pi p \rightarrow \pi\eta\eta'$
2230 ± 6 ± 14	93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
2232 ± 7 ± 7	23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0K_S^0$

 $f_4(2220)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
38 ± 15 ± 13 OUR AVERAGE				
60 ± 107 ± 57		ASTON	88F LASS	11 $K^- p \rightarrow K^+K^-A$
80 ± 30		BOLONKIN	88 SPEC	40 $\pi^- p \rightarrow K_S^0K_S^0n$
26 ± 20 ± 16 ± 17	93	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K^+K^-$
18 ± 23 ± 15 ± 10	23	BALTRUSAIT..86D	MRK3	$e^+e^- \rightarrow \gamma K_S^0K_S^0$

 $f_4(2220)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\bar{K}$		
Γ_2 $p\bar{p}$	$< 1.1 \times 10^{-3}$	99.7%
Γ_3 $\gamma\gamma$		
Γ_4 $\eta\eta'(958)$		

 $f_4(2220)$ $\Gamma(I)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}) \times \Gamma(\gamma\gamma)/\Gamma(\text{total})$	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_1\Gamma_3/\Gamma$
< 0.086	95	¹ ALBRECHT	90G ARG	$\gamma\gamma \rightarrow K^+K^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.0	95	² ALTHOFF	85B TASS	$\gamma\gamma, K\bar{K}\pi$	

¹ Assuming $J^P = 2^+$.² True for $J^P = 0^+$ and $J^P = 2^+$. **$f_4(2220)$ BRANCHING RATIOS**

$\Gamma(p\bar{p})/\Gamma(\text{total})$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
< 1.1	99.7	³ BARNES	93 SPEC	$1.3-1.57\bar{p}p \rightarrow K_S^0K_S^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 2.6	99.7	³ BARDIN	87 CNTR	$1.3-1.5\bar{p}p \rightarrow K^+K^-$	
< 3.6	99.7	³ SCULLI	87 CNTR	$1.29-1.55\bar{p}p \rightarrow K^+K^-$	

³ Assuming $\Gamma = 30-35$ MeV, $J^P = 2^+$ and $B(f_4(2220) \rightarrow K\bar{K}) = 10\%$. **$f_4(2220)$ REFERENCES**

BARNES	93	PL B309 469	+Brien, Breunlich (PS185 Collab.)
ALBRECHT	90G	ZPHY C48 183	+Ehrlichmann, Harder+ (ARGUS Collab.)
ASTON	88D	NP B301 525	+Awaji, Bienz+ (SLAC, NAGO, CINC, INUS)
ASTON	88F	PL B215 199	+Awaji+ (SLAC, NAGO, CINC, INUS) JP
AUGUSTIN	88	PRL 60 2238	+Calcaterra+ (DM2 Collab.)
BOLONKIN	88	NP B309 426	+Bioshenko, Gorin+ (TEP, SERP)
BARDIN	87	PL B195 292	+Burgun+ (SACL, FERR, CERN, PADO, TORI)
SCULLI	87	PRL 58 1715	+Christenson, Kreiter, Nemethy, Yamin (NYU, BNL)
ALDE	86B	PL B177 120	+Binon, Bricman+ (SERP, BELG, LANL, LAPP)
BALTRUSAIT..86D	86D	PRL 56 107	+Baltusaitis (CIT, UCSC, ILL, SLAC, WASH)
ALTHOFF	85B	ZPHY C29 189	+Braunschweig, Kirschfink+ (TASSO Collab.)
BEHREND 84	84	PL 137B 277	+Chadwick, Chauveau, Gentile+ (CLEO Collab.)

OTHER RELATED PAPERS

BARDIN	87	PL B195 292	+Burgun+ (SACL, FERR, CERN, PADO, TORI)
YAOUANC	85	ZPHY C28 309	+Oliver, Pene, Raynal, Ono (ORSAY, TOKY)
GODFREY	84	PL 141B 439	+Kokoski, Isgur (TNTO)
SHATZ	84	PL 138B 209	(CIT)
WILLEY	84	PRL 52 585	(PITT)
EINSEWILER	83	Brighton Conf. 348	(Mark III Collab.)
HITLIN	83	Cornell Conf. 746	(CIT)

 $\eta(2225)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

OMITTED FROM SUMMARY TABLESeen in $J/\psi \rightarrow \gamma\phi\phi$. Needs confirmation. **$\eta(2225)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2230 ± 25 ± 15	BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K^+K^-$
2214 ± 20 ± 13	BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K_S^0K_L^0$
~ 2220	BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+K^-K^+K^-$

 $\eta(2225)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 ± 300 ± 60	BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+K^-K^+K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 80	BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+K^-K^+K^-$

 $\eta(2225)$ REFERENCES

BAI	90B	PRL 65 1309	+Blaylock+ (Mark III Collab.)
BISELLO	86B	PL B179 294	+Busetto, Castro, Limentani+ (DM2 Collab.)

Meson Full Listings

 $\rho_3(2250)$, $f_2(2300)$, $f_4(2300)$ $\rho_3(2250)$ $I(G^{JPC}) = 1^+(3^-)$

OMITTED FROM SUMMARY TABLE

Contains results only from formation experiments. For production experiments see the $\bar{N}N(1100-3600)$ entry. See also $\rho(2150)$, $f_2(2150)$, $f_4(2300)$, $\rho_5(2350)$.

 $\rho_3(2250)$ MASS $\bar{p}p \rightarrow \pi\pi \text{ or } K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 2250.0	¹ MARTIN	80B	RVUE	
~ 2300.0	¹ MARTIN	80C	RVUE	
~ 2140.0	² CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2150.0	³ CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

- ¹ $I(J^P) = 1(3^-)$ from simultaneous analysis of $\bar{p}p \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.
² $I = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.
³ $I(J^P) = 1(3^-)$ from amplitude analysis.

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 2190.0	⁴ CUTTS	78B	CNTR	0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2155.0 ± 15.0	^{4,5} COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2193 ± 2	^{4,6} ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
2190 ± 10	⁷ ABRAMS	70	CNTR	S channel $\bar{p}N$
⁴ Isospins 0 and 1 not separated.				
⁵ From a fit to the total elastic cross section.				
⁶ Referred to as T or T' region by ALSPECTOR 73.				
⁷ Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.				

 $\rho_3(2250)$ WIDTH $\bar{p}p \rightarrow \pi\pi \text{ or } K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 250.0	⁸ MARTIN	80B	RVUE	
~ 200.0	⁸ MARTIN	80C	RVUE	
~ 150.0	⁹ CARTER	78B	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 200.0	¹⁰ CARTER	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

- ⁸ $I(J^P) = 1(3^-)$ from simultaneous analysis of $\bar{p}p \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.
⁹ $I = 0, 1$. $J^P = 3^-$ from Barrelet-zero analysis.
¹⁰ $I(J^P) = 1(3^-)$ from amplitude analysis.

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
135.0 ± 75.0	^{11,12} COUPLAND	77	CNTR 0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
98 ± 8	¹² ALSPECTOR	73	CNTR	$\bar{p}p$ S channel
~ 85	¹³ ABRAMS	70	CNTR	S channel $\bar{p}N$
¹¹ From a fit to the total elastic cross section.				
¹² Isospins 0 and 1 not separated.				
¹³ Seen as bump in $I = 1$ state. See also COOPER 68. PEASLEE 75 confirm $\bar{p}p$ results of ABRAMS 70, no narrow structure.				

 $\rho_3(2250)$ REFERENCES

MARTIN	80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	+Pennington	(DURH) JP
CARTER	78B	NP B141 467		(LOQM)
CUTTS	78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
CARTER	77	PL 67B 117	+Coupland, Eisenhandler, Astbury+	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
PEASLEE	75	PL 57B 189	+Demaro, Guerrero+	(CANB, BARI, BROW, MIT)
ALSPECTOR	73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)
COOPER	68	PRL 20 1059	+Hyman, Manner, Musgrave+	(ANL)

OTHER RELATED PAPERS

MARTIN	79B	PL 86B 93	+Pennington	(DURH)
CARTER	78	NP B132 176		(LOQM) JP
CARTER	77B	PL 67B 122		(LOQM)
CARTER	77C	NP B127 202	+Coupland, Atkinson+	(LOQM, DARE, RHEL)
MONTANET	77	Boston Conf., 260		(CERN)
ZEMANY	76	NP B103 537	+MingMa, Mounitz, Smith	(MSU)
BERTANZA	74	NC 23A 209	+Bigi, Casali, Lariccia+	(PISA, PADO, TORI)
BETTINI	73	NC 15A 563	+Alston-Garnjost, Bigi+	(PADO, LBL, PISA, TORI)
DONNACHIE	73	LNC 7 285	+Thomas	(MCHS)
NICHOLSON	73	PR D7 2572	+Delorme, Carroll+	(CIT, ROCH, BNL)
FIELDS	71	PRL 27 1749	+Cooper, Rhines, Allison	(ANL, OXF)
YOH	71	PRL 26 922	+Barish, Caroll, Lobkowicz+	(CIT, BNL, ROCH)
ABRAMS	67C	PRL 18 1209	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)

 $f_2(2300)$ $I(G^{JPC}) = 0^+(2^+)$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)

 $f_2(2300)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2297 ± 28	¹ ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi \phi n$
••• We do not use the following data for averages, fits, limits, etc. •••			
2231.0 ± 10.0	BOOTH	86	OMEG 85 $\pi^- Be \rightarrow 2\phi Be$
2220.0 ^{+90.0} _{-20.0}	LINDENBAUM	84	RVUE
2320.0 ± 40.0	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$
¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi \phi 2^+ + S_2$, D_2 , and D_0 is 6 ⁺¹⁵ ₋₅ , 25 ⁺¹⁸ ₋₁₄ , and 69 ⁺¹⁶ ₋₂₇ , respectively.			

 $f_2(2300)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
149 ± 41	² ETKIN	88	MPS 22 $\pi^- p \rightarrow \phi \phi n$
••• We do not use the following data for averages, fits, limits, etc. •••			
133.0 ± 50.0	BOOTH	86	OMEG 85 $\pi^- Be \rightarrow 2\phi Be$
200.0 ± 50.0	LINDENBAUM	84	RVUE
220.0 ± 70.0	ETKIN	82	MPS 22 $\pi^- p \rightarrow 2\phi n$
² Includes data of ETKIN 85.			

 $f_2(2300)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\phi \phi$	seen

 $f_2(2300)$ REFERENCES

ETKIN	88	PL B201 568	+Foley, Lindenbaum+	(BNL, CUNY)
BOOTH	86	NP B273 677	+Carroll, Donald, Edwards+	(LIVP, GLAS, CERN)
ETKIN	85	PL 165B 217	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
LINDENBAUM	84	CNPP 13 285		(CUNY)
ETKIN	82	PRL 49 1820	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
Also	83	Brighton Conf. 351	Lindenbaum	(BNL, CUNY)

OTHER RELATED PAPERS

ARMSTRONG	89B	PL B221 221	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)	
GREEN	86	PRL 56 1639	+Lai+ (FNAL, ARIZ, FSU, NDAM, TUFTS, VAND+)	
BOOTH	84	NP B242 51	+Ballance, Carroll, Donald+	(LIVP, GLAS, CERN)

 $f_4(2300)$ $I(G^{JPC}) = 0^+(4^+)$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_0(2350)$. Contains results only from formation experiments. For production experiments see the $\bar{N}N(1100-3600)$ entry. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $\rho_5(2350)$.

 $f_4(2300)$ MASS $\bar{p}p \rightarrow \pi\pi \text{ or } K\bar{K}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 2300	¹ MARTIN	80B	RVUE
~ 2300	¹ MARTIN	80C	RVUE
~ 2340	² CARTER	78B	CNTR 0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2330	DULUDE	78B	OSPK 1-2 $\bar{p}p \rightarrow \pi^0 \pi^0$
~ 2310	³ CARTER	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \pi\pi$

- ¹ $I(J^P) = 0(4^+)$ from simultaneous analysis of $\bar{p}p \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.
² $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.
³ $I(J^P) = 0(4^+)$ from amplitude analysis.

S-CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 2380.0	⁴ CUTTS	78B	CNTR 0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2345 ± 15.0	^{4,5} COUPLAND	77	CNTR 0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2359 ± 2	^{4,6} ALSPECTOR	73	CNTR $\bar{p}p$ S channel
2375 ± 10	ABRAMS	70	CNTR S channel $\bar{N}N$

- ⁴ Isospins 0 and 1 not separated.
⁵ From a fit to the total elastic cross section.
⁶ Referred to as U or U' region by ALSPECTOR 73.

See key on page 1343

Meson Full Listings

$f_4(2300)$, $f_2(2340)$, $\rho_5(2350)$

 $f_4(2300)$ WIDTH **$\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
~ 200	7 MARTIN	80C RVUE	
~ 150	8 CARTER	78B CNTR	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 210	9 CARTER	77 CNTR	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$
⁷ $I(J^P) = 0(4^+)$ from simultaneous analysis of $\rho\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.			
⁸ $I(J^P) = 0(4^+)$ from Barrelet-zero analysis.			
⁹ $I(J^P) = 0(4^+)$ from amplitude analysis.			

S-CHANNEL $\bar{p}p$ or $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
135.0 +150.0 - 65.0	10,11 COUPLAND	77 CNTR	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
165 + 18 - 8	11 ALSPECTOR	73 CNTR	$\bar{p}p$ S channel
~ 190	ABRAMS	70 CNTR	S channel $\bar{N}N$
¹⁰ From a fit to the total elastic cross section.			
¹¹ Isospins 0 and 1 not separated.			

 $f_4(2300)$ REFERENCES

MARTIN	80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	+Pennington	(DURH) JP
CARTER	78B	NP B141 467		(LOQM)
CUTTS	78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
DULUDE	78B	PL 79B 335	+Lanou, Massimo, Peaslee+	(BROW, MIT, BARI) JP
CARTER	77	PL 67B 117	+Coupland, Eisenhandler, Astbury+	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)
ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)

OTHER RELATED PAPERS

FIELDS	71	PRL 27 1749	+Cooper, Rhines, Allison	(ANL, OXF)
YOH	71	PRL 26 922	+Barish, Caroli, Lobkowicz+	(CIT, BNL, ROCH)
BRICMAN	69	PL 29B 451	+Ferro-Luzzi, Bizard+	(CERN, CAEN, SACL)

 $f_2(2340)$

$$I^G(J^{PC}) = 0^+(2^+ +)$$

See also the mini-review under non- $q\bar{q}$ candidates. (See the index for the page number.)

 $f_2(2340)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2339 ± 55	1 ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi\phi$
••• We do not use the following data for averages, fits, limits, etc. •••			
2392.0 ± 10.0	BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
2360.0 ± 20.0	LINDENBAUM	84 RVUE	
¹ Includes data of ETKIN 85. The percentage of the resonance going into $\phi\phi$ 2 ⁺⁺ , S ₂ , D ₂ , and D ₀ is 37 ± 19, 4 ⁺¹² / ₋₄ , and 59 ⁺²¹ / ₋₁₉ , respectively.			

 $f_2(2340)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
319 + 81 - 69	2 ETKIN	88 MPS	22 $\pi^- p \rightarrow \phi\phi$
••• We do not use the following data for averages, fits, limits, etc. •••			
198.0 ± 50.0	BOOTH	86 OMEG	85 $\pi^- Be \rightarrow 2\phi Be$
150.0 +150.0 - 50.0	LINDENBAUM	84 RVUE	
² Includes data of ETKIN 85.			

 $f_2(2340)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \phi\phi$	seen

 $f_2(2340)$ REFERENCES

ETKIN	88	PL B201 568	+Foley, Lindenbaum+	(BNL, CUNY)
BOOTH	86	NP B273 677	+Carroll, Donald, Edwards+	(LIVP, GLAS, CERN)
ETKIN	85	PL 165B 217	+Foley, Longacre, Lindenbaum+	(BNL, CUNY)
LINDENBAUM	84	CNPP 13 285		(CUNY)

OTHER RELATED PAPERS

ARMSTRONG	89B	PL B221 221	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)	
GREEN	86	PRL 56 1639	+Lai+(FNAL, ARIZ, FSU, NDAM, TUFTS, VAND+)	
BOOTH	84	NP B242 51	+Ballance, Carroll, Donald+	(LIVP, GLAS, CERN)

 $\rho_5(2350)$

$$I^G(J^{PC}) = 1^+(5^{--})$$

OMITTED FROM SUMMARY TABLE

This entry was previously called $U_1(2400)$. Contains results only from formation experiments. For production experiments see the $\bar{N}N(1100-3600)$ entry. See also $\rho(2150)$, $f_2(2150)$, $\rho_3(2250)$, $f_4(2300)$.

 $\rho_5(2350)$ MASS **$\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 2300	1 MARTIN	80B RVUE		
~ 2250	1 MARTIN	80C RVUE		
~ 2500	2 CARTER	78B CNTR	0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 2480	3 CARTER	77 CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$
¹ $I(J^P) = 1(5^-)$ from simultaneous analysis of $\rho\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.				
² $I = 0(1)$; $J^P = 5^-$ from Barrelet-zero analysis.				
³ $I(J^P) = 1(5^-)$ from amplitude analysis.				

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 2380	4 CUTTS	78B CNTR		0.97-3 $\bar{p}p \rightarrow \bar{N}N$
2345.0 ± 15.0	4,5 COUPLAND	77 CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
2359 ± 2	4,6 ALSPECTOR	73 CNTR		$\bar{p}p$ S channel
2350 ± 10	7 ABRAMS	70 CNTR		S channel $\bar{N}N$
2360.0 ± 25.0	8 OH	70B HDBC	-0	$\bar{p}(p\eta)$, $K^* K 2\pi$
⁴ Isospins 0 and 1 not separated.				
⁵ From a fit to the total elastic cross section.				
⁶ Referred to as U or U region by ALSPECTOR 73.				
⁷ For $l = 1$ $\bar{N}N$.				
⁸ No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.				

 $\rho_5(2350)$ WIDTH **$\bar{p}p \rightarrow \pi\pi$ or $\bar{K}K$**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
~ 250	9 MARTIN	80B RVUE		
~ 300	9 MARTIN	80C RVUE		
~ 150	10 CARTER	78B CNTR	0	0.7-2.4 $\bar{p}p \rightarrow K^- K^+$
~ 210	11 CARTER	77 CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \pi\pi$
⁹ $I(J^P) = 1(5^-)$ from simultaneous analysis of $\rho\bar{p} \rightarrow \pi^- \pi^+$ and $\pi^0 \pi^0$.				
¹⁰ $I = 0(1)$; $J^P = 5^-$ from Barrelet-zero analysis.				
¹¹ $I(J^P) = 1(5^-)$ from amplitude analysis.				

S-CHANNEL $\bar{N}N$

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••				
135.0 +150.0 - 65.0	12,13 COUPLAND	77 CNTR	0	0.7-2.4 $\bar{p}p \rightarrow \bar{p}p$
165 + 18 - 8	13 ALSPECTOR	73 CNTR		$\bar{p}p$ S channel
< 60.0	14 OH	70B HDBC	-0	$\bar{p}(p\eta)$, $K^* K 2\pi$
~ 140	ABRAMS	67C CNTR		S channel $\bar{N}N$
¹² From a fit to the total elastic cross section.				
¹³ Isospins 0 and 1 not separated.				
¹⁴ No evidence for this bump seen in the $\bar{p}p$ data of CHAPMAN 71B. Narrow state not confirmed by OH 73 with more data.				

 $\rho_5(2350)$ REFERENCES

MARTIN	80B	NP B176 355	+Morgan	(LOUC, RHEL) JP
MARTIN	80C	NP B169 216	+Pennington	(DURH) JP
CARTER	78B	NP B141 467		(LOQM)
CUTTS	78B	PR D17 16	+Good, Grannis, Green, Lee+	(STON, WISC)
CARTER	77	PL 67B 117	+Coupland, Eisenhandler, Astbury+	(LOQM, RHEL) JP
COUPLAND	77	PL 71B 460	+Eisenhandler, Gibson, Astbury+	(LOQM, RHEL)
ALSPECTOR	73	PRL 30 511	+Cohen, Cvijanovich+	(RUTG, UPNJ)
OH	73	NP B51 57	+Eastman, MingMa, Parker, Smith+	(MSU)
CHAPMAN	71B	PR D4 1275	+Green, Lys, Murphy, Ring+	(MICH)
ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)
OH	70B	PRL 24 1257	+Parker, Eastman, Smith, Sprafka, Ma	(MSU)
ABRAMS	67C	PRL 18 1209	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL)

OTHER RELATED PAPERS

CASO	70	LNC 3 707	+Conte, Tomasini+	(GENO, HAMB, MILA, SACL)
BRICMAN	69	PL 29B 451	+Ferro-Luzzi, Bizard+	(CERN, CAEN, SACL)

Meson Full Listings

 $a_6(2450)$, $f_6(2510)$, $X(3250)$ **$a_6(2450)$**

$$I^G(J^{PC}) = 1^-(6^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K\bar{K}$ system. Needs confirmation. **$a_6(2450)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2450 ± 130	¹ CLELAND	82B	SPEC	± 50 $\pi p \rightarrow K_S^0 K^\pm p$

¹ From an amplitude analysis. **$a_6(2450)$ WIDTH**

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
400 ± 250	² CLELAND	82B	SPEC	± 50 $\pi p \rightarrow K_S^0 K^\pm p$

² From an amplitude analysis. **$a_6(2450)$ DECAY MODES**

Mode	Γ_1
$K\bar{K}$	

 $a_6(2450)$ REFERENCES

CLELAND 82B NP B208 228 +Delfosse, Dorsaz, Gloor (DURH, GEVA, LAUS, PITT)

 $f_6(2510)$

$$I^G(J^{PC}) = 0^+(6^{++})$$

OMITTED FROM SUMMARY TABLE

Seen in $\pi^0\pi^0$. Needs confirmation. **$f_6(2510)$ MASS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2510.0 ± 30.0	BINON	84B	GAM2 38 $\pi^- p \rightarrow n 2\pi^0$

 $f_6(2510)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
240.0 ± 60.0	BINON	84B	GAM2 23 $\pi^- p \rightarrow n 2\pi^0$

 $f_6(2510)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\pi\pi$	(6.0 ± 1.0) %

 $f_6(2510)$ BRANCHING RATIOS

$\Gamma(\pi\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 ± 0.01	¹ BINON	83C	GAM2 38 $\pi^- p \rightarrow n 4\gamma$	

¹ Assuming one pion exchange. **$f_6(2510)$ REFERENCES**BINON 84B LNC 39 41 +Donskov, Duteil, Gouanere+ (SERP, BELG, LAPP) JP
BINON 83C SJNP 38 723 +Gouanere, Donskov, Duteil+ (SERP, BRUX+)
Translated from YAF 38 1199. **$X(3250)$**

$$I^G(J^{PC}) = ??(???)$$

OMITTED FROM SUMMARY TABLE

Narrow peak observed in several final states with hidden strangeness ($\Lambda\bar{p}K^+$, $\Lambda\bar{p}K^+\pi^\pm$, $K^0\rho\bar{p}K^\pm$). Needs confirmation. See also under non- $q\bar{q}$ candidates. (See the index for the page number.) **$X(3250)$ MASS****3-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3250 ± 8 ± 20	¹ ALEEV	93	BIS2 $X(3250) \rightarrow \Lambda\bar{p}K^+$
3265 ± 7 ± 20	¹ ALEEV	93	BIS2 $X(3250) \rightarrow \bar{\Lambda}pK^-$

¹ Supersedes KEKELIDZE 90.**4-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3245 ± 8 ± 20	¹ ALEEV	93	BIS2 $X(3250) \rightarrow \Lambda\bar{p}K^+\pi^\pm$
3250 ± 9 ± 20	¹ ALEEV	93	BIS2 $X(3250) \rightarrow \bar{\Lambda}pK^-\pi^\mp$
3270 ± 8 ± 20	¹ ALEEV	93	BIS2 $X(3250) \rightarrow K_S^0\rho\bar{p}K^\pm$

 $X(3250)$ WIDTH**3-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
45 ± 18	² ALEEV	93	BIS2 $X(3250) \rightarrow \Lambda\bar{p}K^+$
40 ± 18	² ALEEV	93	BIS2 $X(3250) \rightarrow \bar{\Lambda}pK^-$

² Supersedes KEKELIDZE 90.**4-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
25 ± 11	² ALEEV	93	BIS2 $X(3250) \rightarrow \Lambda\bar{p}K^+\pi^\pm$
50 ± 20	² ALEEV	93	BIS2 $X(3250) \rightarrow \bar{\Lambda}pK^-\pi^\mp$
25 ± 11	² ALEEV	93	BIS2 $X(3250) \rightarrow K_S^0\rho\bar{p}K^\pm$

 $X(3250)$ DECAY MODES

Mode	Γ_1
$\Lambda\bar{p}K^+$	
$\Lambda\bar{p}K^+\pi^\pm$	
$K^0\rho\bar{p}K^\pm$	

 $X(3250)$ REFERENCESALEEV 93 PAN 56 1358 +Baladin+ (BIS-2 Collab.)
Translated from YAF 56 100.
KEKELIDZE 90 Hadron 89 Conf. p 551+Aleev+ (BIS-2 Collab.)

See key on page 1343

Meson Full Listings

$e^+e^-(1100-2200), \bar{N}N(1100-3600)$

OTHER LIGHT UNFLAVORED MESONS ($S = C = B = 0$)

 $e^+e^-(1100-2200)$

$$I^G(J^{PC}) = ?^?(1^{--})$$

OMITTED FROM SUMMARY TABLE

This entry contains nonstrange vector mesons coupled to e^+e^- (photon) between the ϕ and $J/\psi(1S)$ mass regions. See also $\omega(1420)$, $\rho(1450)$, $\omega(1600)$, $\phi(1680)$, and $\rho(1700)$.

$e^+e^-(1100-2200)$ MASSES AND WIDTHS

We do not use the following data for averages, fits, limits, etc.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1100 to 2200 OUR LIMIT				
1097.0 ^{+16.0} _{-19.0}	BARTALUCCI 79	OSPK		$7\gamma p \rightarrow e^+e^-p$
31.0 ^{+24.0} _{-20.0}	BARTALUCCI 79	OSPK		$7\gamma p \rightarrow e^+e^-p$
1266.0 \pm 5.0	BARTALUCCI 79	DASP	0	$7\gamma p \rightarrow e^+e^-p$
110.0 \pm 35.0	BARTALUCCI 79	DASP	0	$7\gamma p \rightarrow e^+e^-p$
~ 1830.0	PETERSON 78	SPEC		$\gamma p \rightarrow K^+K^-p$
~ 120.0	PETERSON 78	SPEC		$\gamma p \rightarrow K^+K^-p$
~ 1820	¹ SPINETTI 79	RVUE		$e^+e^- \rightarrow 4\pi \pm 2\gamma$
~ 30	¹ SPINETTI 79	RVUE		$e^+e^- \rightarrow 4\pi \pm 2\gamma$
~ 2130	² ESPOSITO 78	FRAM		$e^+e^- \rightarrow K^*(892)^+ \dots$
~ 30	² ESPOSITO 78	FRAM		$e^+e^- \rightarrow K^*(892)^+ \dots$

¹ Integrated cross section of BACCI 77, BARBIELLINI 77, ESPOSITO 77.² Not seen by DELCOURT 79.

$e^+e^-(1100-2200)$ REFERENCES

BARTALUCCI 79	NC 49A 207	+Basini, Bertolucci+	(DESY, FRAS)
DELCOURT 79	PL 86B 395	+Derado, Bertrand, Bisello, Bizot, Buon+	(LALO)
SPINETTI 79	Batavia Conf. 506		(FRAS)
ESPOSITO 78	LNC 22 305	+Felicetti	(FRAS, NAPL, PADO, ROMA)
PETERSON 78	PR D18 3955	+Dixon, Ehrlich, Galik, Larson	(CORN, HARV)
BACCI 77	PL 68B 393	+DeZorzi, Penso, Stella, Baldini+	(ROMA, FRAS)
BARBIELLINI 77	PL 68B 397	+Baretta+	(FRAS, NAPL, PISA, SANI)
ESPOSITO 77	PL 68B 389	+Felicetti, Marini+	(FRAS, NAPL, PADO, ROMA)

OTHER RELATED PAPERS

BACCI 76	PL 64B 356	+Bidoli, Penso, Stella, Baldini+	(ROMA, FRAS)
BACCI 75	PL 58B 481	+Bidoli, Penso, Stella+	(ROMA, FRAS)

$\bar{N}N(1100-3600)$

OMITTED FROM SUMMARY TABLE

This entry contains various high mass, nonstrange structures coupled to the baryon-antibaryon system, as well as quasi-nuclear bound states below threshold.

$\bar{N}N(1100-3600)$ MASSES AND WIDTHS

We do not use the following data for averages, fits, limits etc.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1100 to 3600 OUR LIMIT					
1107 \pm 4	DAFTARI	87	DBC	0	$0. \bar{p}n \rightarrow \rho^- \pi^+ \pi^-$
111 \pm 8 \pm 15	DAFTARI	87	DBC	0	$0. \bar{p}n \rightarrow \rho^- \pi^+ \pi^-$
1167 \pm 7	¹ CHIBA	91	CNTR		$\bar{p}d \rightarrow \gamma X$
1191.0 \pm 9.9	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1210 \pm 5.0	1,2,3,4 RICHTER	83	CNTR	0	Stopped \bar{p}
1325 \pm 5	¹ CHIBA	91	CNTR		$\bar{p}d \rightarrow \gamma X$
1329.2 \pm 7.6	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1390.9 \pm 6.3	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1395	1,3,4,5 PAVLOPO...	78	CNTR		Stopped \bar{p}
~ 1410	BETTINI	66	DBC	0	$0. \bar{p}N \rightarrow 5\pi$
~ 100	BETTINI	66	DBC	0	$0. \bar{p}N \rightarrow 5\pi$
1468 \pm 6	⁶ BRIDGES	86B	DBC	0	$0. \bar{p}N \rightarrow 2\pi^- \pi^+ \pi^0$
88 \pm 18	⁶ BRIDGES	86B	DBC	0	$0. \bar{p}N \rightarrow 2\pi^- \pi^+ \pi^0$
1512 \pm 7	¹ CHIBA	91	CNTR		$\bar{p}d \rightarrow \gamma X$
1523.8 \pm 3.6	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1522 \pm 7	⁶ BRIDGES	86B	DBC	0	$0. \bar{p}N \rightarrow 2\pi^- \pi^+$
59 \pm 12	⁶ BRIDGES	86B	DBC	0	$0. \bar{p}N \rightarrow 2\pi^- \pi^+$
1577.8 \pm 3.4	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1594 \pm 9	⁶ BRIDGES	86B	DBC	-	$0. \bar{p}N \rightarrow 2\pi^- \pi^+ \pi^0$
81 \pm 12	⁶ BRIDGES	86B	DBC	-	$0. \bar{p}N \rightarrow 2\pi^- \pi^+ \pi^0$
1633.6 \pm 4.1	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1637.1 ^{+5.6} _{-7.3}	ADIELS	84	CNTR		$\bar{p}He$
1638 \pm 3.0	1,2,3,4 RICHTER	83	CNTR	0	Stopped \bar{p}
1644.0 ^{+5.6} _{-7.3}	ADIELS	84	CNTR		$\bar{p}He$
1646	1,3,4,5 PAVLOPO...	78	CNTR		Stopped \bar{p}
1687.1 ^{+5.0} _{-4.3}	ADIELS	84	CNTR		$\bar{p}He$
1684	1,3,4,5 PAVLOPO...	78	CNTR		Stopped \bar{p}
1693 \pm 2	¹ CHIBA	91	CNTR		$\bar{p}d \rightarrow \gamma X$
1694 \pm 2.0	1,2,3,4 RICHTER	83	CNTR	0	Stopped \bar{p}
1713.0 \pm 2.6	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$
1731.0 \pm 1.5	¹ CHIBA	87	CNTR	0	$0. \bar{p}p \rightarrow \gamma X$

- 17 From energy dependence of far backward elastic scattering. Some indication of additional structure.
 18 Not seen by ALBERI 79 with comparable statistics.
 19 Not seen by ALBERI 79 with comparable statistics.
 20 Seen as a bump in the $\bar{p}p \rightarrow K_S^0 K_L^0$ cross section with $J^{PC} = 1^{-}$.
 21 Isospin 1 favored.
 22 Not seen by BIONTA 80, CARROLL 80, HAMILTON 80, BANKS 81, CHUNG 81, BARNETT 83.
 23 Neutron spectator. See also $n\bar{p}\pi^-$ channel following.
 24 Proton spectator. See also $p\bar{p}n(n)$ channel above.
 25 $I(J^P) = 1(3^-)$ from a mass dependent partial-wave analysis taking solution A.
 26 $I(J^P) = 1(3^-)$ from amplitude analysis assuming one-pion exchange.
 27 Seen in final state $\omega\pi^+\pi^-$.
 28 $I(J^P) = 0(2^+)$ from amplitude analysis assuming one-pion exchange.
 29 ALLES-BORELLI 67b see neutral mode only $\pi^+\pi^-\pi^0$.
 30 $I(J^P) = 0(4^+)$ from a mass dependent partial-wave analysis taking solution A.
 31 $I(J^P) = 0(4^+)$ from amplitude analysis assuming one-pion exchange.
 32 $I(J^P) = 1(5^-)$ from amplitude analysis assuming one-pion exchange.
 33 $I(J^P) = 1(5^-)$ from amplitude analysis of $\bar{p}p \rightarrow \pi\pi$.
 34 $I=0,1 J^P = 5^-$ from Barrelet-zero analysis.
 35 Decays to $\bar{N}N$ and $\bar{N}N\pi$. Not seen by BARNETT 83.
 36 Decays to $4\pi^+4\pi^-$.

 $\bar{N}N(1100-3600)$ REFERENCES

CARBONELL	93	PL B306 407	+Protasov, Dalkarov (ISNG, LEBD)
CHIBA	91	PR D44 1933	+Fujitani+ (FUJI, KEK, SANG, OSAK, TMU)
GRAF	91	PR D44 1945	+Fero, Gee+(UCI, PENN, NMSU, KARLK, KARLE, ATHU)
BUSENITZ	89	PR D40 1	+Ostrowski, Callahan+ (ILL, FNAL)
CHIBA	88	PL B202 447	+Doi (FUJI, INUS, KEK, SANG, OSAK, TMU)
CHIBA	87	PR D36 3321	+Doi+ (FUJI, INUS, KEK, SANG, OSAK, TMU)
DAFTARI	87	PRL 58 859	+Gray, Kalogeropoulos, Roy (SYRA)
FRANKLIN	87	PL B184 81	
ADIELS	86	PL B182 405	+Backenstoss+ (STOH, BASL, LASL, THES, CERN)
ANGELOPO... 86	PL B178 441	Angelopoulos+(ATHU, UCI, KARLK, KARLE, NMSU, PENN)	
BRIDGES	86B	PRL 56 215	+Daftari, Kalogeropoulos, Debb+ (SYRA, CASE)
BRIDGES	86D	PL B180 313	+Brown, Daftari+ (SYRA, BNL, CASE, UMD, COLU)
ADIELS	84	PL 138B 235	+ (BASL, KARLK, KARLE, STOH, STRB, THES)
CLOUGH	84	PL 146B 299	+Beard, Bugg+ (SURR, LOQM, ANIK, TRST, GEVA)
AZOOZ	83	PL 122B 471	+Butterworth (LOIC, RHEL, SACL, SLAC, TOHOK+)
BARNETT	83	PR D27 493	+Blockus, Burka, Chien, Christian+ (JHU)
BODENKAMP	83	PL 133B 275	+Fries, Behrend, Fenner+ (KARLK, KARLE, DESY)
RICHTER	83	PL 126B 284	+Adiels (BASL, KARLK, KARLE, STOH, STRB, THES)
BANKS	81	PL 100B 191	+Booth, Campbell, Armstrong+ (LIVP, CERN)
CHUNG	81	PRL 46 395	+Bensinger+ (BNL, BRAN, CINC, FSU, MASN)
JASTRZEM... 81	PR D23 2784	Jastrzembki, Mandelkern+ (TEMP, UCI, UNM)	
ALPER	80	PL 94B 422	+Becker+ (AMST, CERN, CRAC, MPIM, OXF+)
ASTON	80D	PL 93B 517	(BONN, CERN, EPOL, GLAS, LANC, MCHS, ORSAY+)
BIONTA	80	PRL 44 909	+Carroll, Edelstein+ (BNL, CMU, FNAL, MASN)
CARROLL	80	PRL 44 1572	+Chiang, Johnson, Cester, Webb+ (BNL, PRIN)
DAUM	80E	PL 90B 475	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
DEFONK	80	PL 91B 112	+Dobrzynski, Angelini, Bigi+ (CDFE, PISA)
HAMILTON	80	PL 44 1179	+Pun, Tripp, Lazarus+ (LBL, BNL, MTHO)
HAMILTON	80B	PRL 44 1182	+Pun, Tripp, Lazarus+ (LBL, BNL, MTHO)
KREYMER	80	PR D22 36	+Baggett, Fieguth+ (IND, PURD, SLAC, VAND)
ROZANSKA	80	NP B162 505	+Blum, Dietl, Grayer, Lorenz+ (MPIM, CERN)
ALBERI	79	PL 83B 247	+Alvear, Castell, Poropat+ (TRST, CERN, IFRJ)
EVANGELISTA	79	NP B153 253	+ (BARI, BONN, CERN, DARE, GLAS, LIVP+)
EVANGELISTA	79B	NP B154 381	+ (BARI, BONN, CERN, DARE, GLAS, LIVP+)
SAKAMOTO	79	NP B158 410	+Hashimoto, Sai, Yamamoto+ (IUIS)
CARTER	78B	NP B141 467	+ (LOQM)
PAVLOPO... 78	PL 72B 415	Pavlopoulos+(KARLK, KARLE, BASL, CERN, STOH, STRB)	
BRUCKNER	77	PL 67B 222	+Granz, Ingham, Kilian+ (MPIM, HEIDP, CERN)
CARTER	77	PL 67B 117	+Coupland, Eisenhandler, Astbury+ (LOQM, RHEL, JP)
ABASHIAN	76	PR D13 5	+Watson, Gelfand, Buttram+ (ILL, ANL, CHIC, ISU)
BRAUN	76	PL 60B 481	+Brick, Fridman, Gerber, Juillot, Maurer+ (STRB)
CHALOUPIKA	76	PL 61B 487	+ (CERN, LIVP, MONS, PADO, ROMA, TRST)
ALSTON... 75	PRL 35 1885	Alston-Garnjost, Kenney, Pollard, Ross, Tripp+(LBL, MTHO)	
D'ANDLIAU	75	PL 58B 223	+Cohen-Ganouna, Laloum, Lutz, Petri (CDFE, PISA)
KALOGERO... 75	PRL 34 1047	Kalogeropoulos, Tzanakos (SYRA)	
CARROLL	74	PRL 32 247	+Chiang, Kycia, Li, Mazur, Michael+ (BNL)
DONALD	73	NP B61 333	+Edwards, Gibbins, Briand, Duboc+ (LIVP, PARIS)
ALEXANDER	72	NP B45 29	+Bar-Nir, Benary, Dagan+ (TELA)
BENVENUTI	71	PRL 27 283	+Cline, Rutz, Reeder, Scherer (WISC)
CLINE	70B	Preprint	+English, Reeder (WISC)
ALLES... 67B	NC 50A 776	Alles-Borelli, French, Frisk+ (CERN, BONN) G	
BETTINI	66	NC 42A 695	+Cresti, Limentani, Bertanza, Bigi+ (PADO, PISA)

OTHER RELATED PAPERS

TANIMORI	90	PR D41 744	+Ishimoto+ (KEK, INUS, KYOT, TOHOK, HIRO)
LIU	87	PRL 58 2288	+Kiu, Li (STON)
ARMSTRONG	86C	PL B175 383	+Chu, Clement, Elinon+ (BNL, HOUS, PENN, RICE)
BRIDGES	86	PRL 56 211	+Brown+ (BLSU, BNL, CASE, COLU, UMD, SYRA)
BRIDGES	86C	PRL 57 1534	+Daftari, Kalogeropoulos+ (SYRA) JP
DOVER	86	PRL 57 1207	+ (BNL) JP
ANGELOPO... 85	PL 159B 210	Angelopoulos+ (ATHU, UCI, UNM, PENN, TEMP)	
BODENKAMP	85	NP B255 717	+Fries, Behrend, Hesse+ (KARLK, KARLE, DESY)
AZOOZ	84	NP B244 277	+Butterworth (LOIC, RHEL, SACL, SLAC, TOHOK+)

 $X(1900-3600)$

OMITTED FROM SUMMARY TABLE

NOTE ON THE $X(1900-3600)$ REGION

The high-mass region is covered nearly continuously with evidence for peaks of various widths having various decay modes. As a satisfactory grouping into particles is not yet possible, we list all the $Y = 0$ bumps coupled neither to $\bar{N}N$ nor to e^+e^- , and having $M > 1900$ MeV, together, ordered by increasing mass.

The narrow peaks observed in a missing-mass-spectrometer experiment at 1929, 2195, and 2382 MeV, called respectively S , T , and U by the authors (CHIKOVANI 66, FOCACCI 66), were not seen by ANTIPOV 72, who performed a similar experiment at 25 and 40 GeV/c.

 $X(1900-3600)$ MASSES AND WIDTHS

We do not use the following data for averages, fits, limits, etc.

VALUE (MeV)	DOCUMENT ID				
1900 to 3600 OUR LIMIT					
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1870.0 ± 40.0	² ALDE	86D GAM4	0	100 $\pi^- p \rightarrow 2\eta X$	
250.0 ± 30.0	² ALDE	86D GAM4	0	100 $\pi^- p \rightarrow 2\eta X$	
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1898 ± 18	100	THOMPSON 74	HBC	+	13 $\pi^+ p \rightarrow 2\rho X$
108 ⁺⁴¹ ₋₂₇	100	THOMPSON 74	HBC	+	13 $\pi^+ p \rightarrow 2\rho X$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1900 ± 40	100	BOESEBECK 68	HBC	+	8 $\pi^+ p \rightarrow \pi^+ \pi^0 X$
216 ± 105	100	BOESEBECK 68	HBC	+	8 $\pi^+ p \rightarrow \pi^+ \pi^0 X$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
1901 ± 13		ARMSTRONG 89E OMEG	300 $pp \rightarrow p\rho 2(\pi^+ \pi^-)$		
312 ± 61		ARMSTRONG 89E OMEG	300 $pp \rightarrow p\rho 2(\pi^+ \pi^-)$		
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1929 ± 14	FOCACCI 66	MMS	-	3-12 $\pi^- p$	
22 ± 2	FOCACCI 66	MMS	-	3-12 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1970 ± 10	CHLIAPNIK... 80	HBC	0	32 $K^+ p \rightarrow 2K_S^0 2\pi X$	
40 ± 20	CHLIAPNIK... 80	HBC	0	32 $K^+ p \rightarrow 2K_S^0 2\pi X$	
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1973.0 ± 15.0	30	CASO 70	HBC	-	11.2 $\pi^- p \rightarrow \rho 2\pi$
80.0	30	CASO 70	HBC	-	11.2 $\pi^- p \rightarrow \rho 2\pi$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2070	50	TAKAHASHI 72	HBC	8 $\pi^- p \rightarrow N2\pi$	
160	50	TAKAHASHI 72	HBC	8 $\pi^- p \rightarrow N2\pi$	
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2103 ± 50	586	¹ BISELLO 89B	DM2		$J/\psi \rightarrow 4\pi\gamma$
187 ± 75	586	¹ BISELLO 89B	DM2		$J/\psi \rightarrow 4\pi\gamma$
2100.0 ± 40.0		³ ALDE 86D	GAM4	0	100 $\pi^- p \rightarrow 2\eta X$
250.0 ± 40.0		³ ALDE 86D	GAM4	0	100 $\pi^- p \rightarrow 2\eta X$
¹ Estimated by us from various fits.					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
2141.0 ± 12.0	389	GREEN 86	MPSF	400 $pA \rightarrow 4KX$	
49.0 ± 28.0	389	GREEN 86	MPSF	400 $pA \rightarrow 4KX$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2190.0 ± 10.0	CLAYTON 67	HBC	±	2.5 $\bar{p}p \rightarrow a_2, \omega$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

Meson Full Listings

X(1900-3600)

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2195 ± 15	FOCACCI 66	MMS	-	3-12 $\pi^- p$	
39 ± 14	FOCACCI 66	MMS	-	3-12 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2207.0 ± 22.0	⁴ CASO 70	HBC	-	11.2 $\pi^- p$	
130.0	⁴ CASO 70	HBC	-	11.2 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2300.0 ± 100.0	ATKINSON 84F	OMEG ± 0		20-70 $\gamma p \rightarrow \rho f$	
~ 250.0	ATKINSON 84F	OMEG ± 0		20-70 $\gamma p \rightarrow \rho f$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2330 ± 30	ATKINSON 88	OMEG 0		25-50 $\gamma p \rightarrow \rho^\pm \rho^0 \pi^\mp$	
435 ± 75	ATKINSON 88	OMEG 0		25-50 $\gamma p \rightarrow \rho^\pm \rho^0 \pi^\mp$	
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2340 ± 20	126	⁵ BALTAY 75	HBC	+	15 $\pi^+ p \rightarrow p 5\pi$
180 ± 60	126	⁵ BALTAY 75	HBC	+	15 $\pi^+ p \rightarrow p 5\pi$
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2382 ± 24	FOCACCI 66	MMS	-	3-12 $\pi^- p$	
62 ± 6	FOCACCI 66	MMS	-	3-12 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2500.0 ± 32.0	ANDERSON 69	MMS	-	16 $\pi^- p$ backward	
87.0	ANDERSON 69	MMS	-	16 $\pi^- p$ backward	
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2620 ± 20	550	BAUD 69	MMS	-	8-10 $\pi^- p$
85 ± 30	550	BAUD 69	MMS	-	8-10 $\pi^- p$
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2676.0 ± 27.0	⁴ CASO 70	HBC	-	11.2 $\pi^- p$	
150.0	⁴ CASO 70	HBC	-	11.2 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT		
2747 ± 32	DENNEY 83	LASS	10 $\pi^+ N$		
195 ± 75	DENNEY 83	LASS	10 $\pi^+ N$		
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2800 ± 20	640	BAUD 69	MMS	-	8-10 $\pi^- p$
46 ± 10	640	BAUD 69	MMS	-	8-10 $\pi^- p$
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2820 ± 10	15	⁶ SABAU 71	HBC	+	8 $\pi^+ p$
50 ± 10	15	⁶ SABAU 71	HBC	+	8 $\pi^+ p$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2880 ± 20	230	BAUD 69	MMS	-	8-10 $\pi^- p$
< 15	230	BAUD 69	MMS	-	8-10 $\pi^- p$
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
3025.0 ± 20.0	BAUD 70	MMS	-	10.5-13 $\pi^- p$	
~ 25.0	BAUD 70	MMS	-	10.5-13 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
3075.0 ± 20.0	BAUD 70	MMS	-	10.5-13 $\pi^- p$	
~ 25.0	BAUD 70	MMS	-	10.5-13 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
3145.0 ± 20.0	BAUD 70	MMS	-	10.5-15 $\pi^- p$	
< 10.0	BAUD 70	MMS	-	10.5-15 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
3475.0 ± 20.0	BAUD 70	MMS	-	14-15.5 $\pi^- p$	
~ 30.0	BAUD 70	MMS	-	14-15.5 $\pi^- p$	
VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
3535.0 ± 20.0	BAUD 70	MMS	-	14-15.5 $\pi^- p$	
~ 30.0	BAUD 70	MMS	-	14-15.5 $\pi^- p$	

² Seen in $J = 2$ wave in one of the two ambiguous solutions.
³ Seen in $J = 0$ wave in one of the two ambiguous solutions.
⁴ Seen in $\rho^- \pi^+ \pi^-$ (ω and η antiselected in 4π system).
⁵ Dominant decay into $\rho^0 \rho^0 \pi^+$. BALTAY 78 finds confirmation in $2\pi^+ \pi^- 2\pi^0$ events which contain $\rho^+ \rho^0 \pi^0$ and $2\rho^+ \pi^-$.
⁶ Seen in $(K\bar{K}\pi\pi)$ mass distribution.

X(1900-3600) REFERENCES

ARMSTRONG 89E	PL B228 536	+Benayoun (ATHU, BARI, BIRM, CERN, CDEF, CURIN+)
BISELLO 89B	PR D39 701	+Busetto+ (DM2 Collab.)
ATKINSON 88	ZPHY C38 535	+Axon+ (BONN, CERN, GLAS, LANC, MCHS, CURIN)
ALDE 86D	NP B269 485	+Binon, Bricman+ (BELG, LAPP, SERP, CERN)
GREEN 86	PRL 56 1639	+Lai+ (FNAL, ARIZ, FSU, NDAM, TUFTS, VAND+)
ATKINSON 84F	NP B239 1	+ (BONN, CERN, GLAS, LANC, MCHS, IPNP+)
DENNEY 83	PR D28 2726	+Cranley, Firestone, Chapman+ (IOWA, MICH) J
CHLIAPNIK... 80	ZPHY C3 285	+Chliapnikov, Gerdyukov+ (SERP, BRUX, MONS)
BALTAY 78	PR D17 52	+Cautis, Cohen, Csorna, Kalelkar+ (COLU, BING)
BALTAY 75	PRL 35 891	+Cautis, Cohen, Kalelkar, Pisello+ (COLU, BING)
THOMPSON 74	NP B69 220	+Gaidos, McIlwain, Miller, Mulera+ (PURD)
TAKAHASHI 72	PR D6 1266	+Barish+ (TOHOK, PENN, NDAM, ANL)
SABAU 71	LCN 1 514	+Uretsky (BUCH, ANL)
BAUD 70	PL 31B 549	+Benz+ (CERN Boson Spectrometer Collab.)
CASO 70	LCN 3 707	+Conte, Tomasini+ (GENO, HAMB, MILA, SACL)
ANDERSON 69	PRL 22 1390	+Collins+ (BNL, CMU)
BAUD 69	PL 30B 129	+Benz+ (CERN Boson Spectrometer Collab.)
BOESEBECK 68	NP B4 501	+Deutschmann+ (AACH, BERL, CERN)
CLAYTON 67	Heidelberg Conf. 57	+Mason, Muirhead, Filippos+ (LIVP, ATHU)
FOCACCI 66	PRL 17 890	+Kienzle, Levrat, Maglich, Martin (CERN)

OTHER RELATED PAPERS

ANTIPOV 72	PL 40 147	+Kienzle, Landsberg+ (SERP)
CHIKOVANI 66	PL 22 233	+Kienzle, Maglich+ (SERP)

STRANGE MESONS ($S = \pm 1, C = B = 0$)

$K^+ = u\bar{s}, K^0 = d\bar{s}, \bar{K}^0 = \bar{d}s, K^- = \bar{u}s$, similarly for K^{*} 's

K^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

NOTE ON THE CHARGED KAON MASS

The average of the six charged kaon mass measurements which we use in the Full Listings is

$$m_{K^\pm} = 493.677 \pm 0.013 \text{ MeV } (S = 2.4), \quad (1)$$

where the error has been increased by the scale factor S . The large scale factor indicates a serious disagreement between different input data. The average before scaling the error is

$$m_{K^\pm} = 493.677 \pm 0.005 \text{ MeV },$$

$$\chi^2 = 22.9 \text{ for } 5 \text{ D.F.}, \text{ Prob.} = 0.04\%, \quad (2)$$

where the high χ^2 and correspondingly low χ^2 probability further quantify the disagreement.

The main disagreement is between the two most recent and precise results,

$$m_{K^\pm} = 493.696 \pm 0.007 \text{ MeV} \quad \text{DENISOV 91}$$

$$m_{K^\pm} = 493.636 \pm 0.011 \text{ MeV } (S = 1.5) \text{ GALL 88}$$

$$\text{Average} = 493.679 \pm 0.006 \text{ MeV}$$

$$\chi^2 = 21.2 \text{ for } 1 \text{ D.F.}, \text{ Prob.} = 0.0004\%, \quad (3)$$

both of which are measurements of x-ray energies from kaonic atoms. Comparing the average in Eq. (3) with the overall average in Eq. (2), it is clear that DENISOV 91 and GALL 88 dominate the overall average, and that their disagreement is responsible for most of the high χ^2 .

The GALL 88 measurement was made using four different kaonic atom transitions, $K^- \text{Pb } (9 \rightarrow 8)$, $K^- \text{Pb } (11 \rightarrow 10)$, $K^- \text{W } (9 \rightarrow 8)$, and $K^- \text{W } (11 \rightarrow 10)$. The m_{K^\pm} values they obtain from each of these transitions is shown in the Full Listings and in Fig. 1. Their $K^- \text{Pb } (9 \rightarrow 8)$ m_{K^\pm} is below and somewhat inconsistent with their other three transitions. The average of their four measurements is

$$m_{K^\pm} = 493.636 \pm 0.007,$$

$$\chi^2 = 7.0 \text{ for } 3 \text{ D.F.}, \text{ Prob.} = 7.2\%. \quad (4)$$

This is a low but acceptable χ^2 probability so, to be conservative, GALL 88 scaled up the error on their average by $S=1.5$ to obtain their published error ± 0.011 shown in Eq. (3) above and used in the Full Listings average.




Figure 1: Ideogram of m_{K^\pm} mass measurements. GALL 88 and CHENG 75 measurements are shown separately for each transition they measured.

The ideogram in Fig. 1 shows that the DENISOV 91 measurement and the GALL 88 $K^- \text{Pb } (9 \rightarrow 8)$ measurement yield two well-separated peaks. One might suspect the GALL 88 $K^- \text{Pb } (9 \rightarrow 8)$ measurement since it is responsible both for the internal inconsistency in the GALL 88 measurements and the disagreement with DENISOV 91.

To see if the disagreement could result from a systematic problem with the $K^- \text{Pb } (9 \rightarrow 8)$ transition, we have separated the CHENG 75 data, which also used $K^- \text{Pb}$, into its separate transitions. Figure 1 shows that the CHENG 75 and GALL 88 $K^- \text{Pb } (9 \rightarrow 8)$ values are consistent, suggesting the possibility of a common effect such as contaminant nuclear γ rays near the $K^- \text{Pb } (9 \rightarrow 8)$ transition energy, although the CHENG 75 errors are too large to make a strong conclusion. The average of all 13 measurements has a χ^2 of 52.6 as shown in Fig. 1 and the first line of Table 1, yielding an unacceptable χ^2 probability of 0.00005%. The second line of Table 1 excludes both the GALL 88 and CHENG 75 measurements of the $K^- \text{Pb } (9 \rightarrow 8)$ transition and yields a χ^2 probability of 43%. The third [fourth] line of Table 1 excludes only the GALL 88 $K^- \text{Pb } (9 \rightarrow 8)$ [DENISOV 91] measurement and yields a χ^2 probability of 20% [8.6%]. Table 1 shows that removing both measurements of the $K^- \text{Pb } (9 \rightarrow 8)$ transition produces the most consistent set of data, but that excluding only the GALL 88 $K^- \text{Pb } (9 \rightarrow 8)$ transition or DENISOV 91 also produces acceptable probabilities.

Yu.M. Ivanov, representing DENISOV 91, has estimated corrections needed for the older experiments because of improved ^{192}Ir and ^{198}Au calibration γ -ray energies. He estimates that CHENG 75 and BACKENSTOSS 73 m_{K^\pm} values could be raised by about 15 keV and 22 keV, respectively. With these

Meson Full Listings

K^\pm

Table 1: m_{K^\pm} averages for some combinations of Fig. 1 data.

m_{K^\pm} (MeV)	χ^2	D.F.	Prob. (%)	Measurements used
493.664 ± 0.004	52.6	12	0.00005	all 13 measurements
493.690 ± 0.006	10.1	10	43	no K^- Pb(9→8)
493.687 ± 0.006	14.6	11	20	no GALL 88 K^- Pb(9→8)
493.642 ± 0.006	17.8	11	8.6	no DENISOV 91

estimated corrections, Table 1 becomes Table 2. The last line of Table 2 shows that if such corrections are assumed, then GALL 88 K^- Pb (9 → 8) is inconsistent with the rest of the data even when DENISOV 91 is excluded. Yu.M. Ivanov warns that these are rough estimates. Accordingly, we do not use Table 2 to reject the GALL 88 K^- Pb (9 → 8) transition, but we note that a future reanalysis of the CHENG 75 data could be useful because it might provide supporting evidence for such a rejection.

Table 2: m_{K^\pm} averages for some combinations of Fig. 1 data after raising CHENG 75 and BACKENSTOSS 73 values by 0.015 and 0.022 MeV respectively.

m_{K^\pm} (MeV)	χ^2	D.F.	Prob. (%)	Measurements used
493.666 ± 0.004	53.9	12	0.00003	all 13 measurements
493.693 ± 0.006	9.0	10	53	no K^- Pb(9→8)
493.690 ± 0.006	11.5	11	40	no GALL 88 K^- Pb(9→8)
493.645 ± 0.006	23.0	11	1.8	no DENISOV 91

The GALL 88 measurement uses a Ge semiconductor spectrometer which has a resolution of about 1 keV, so they run the risk of some contaminant nuclear γ rays. Studies of γ rays following stopped π^- and Σ^- absorption in nuclei (unpublished) do not show any evidence for contaminants according to GALL 88 spokesperson, B.L. Roberts. The DENISOV 91 measurement uses a crystal diffraction spectrometer with a resolution of 6.3 eV for radiation at 22.1 keV to measure the 4f-3d transition in K^- ^{12}C . The high resolution and the light nucleus reduce the probability for overlap by contaminant γ rays, compared with the measurement of GALL 88. The DENISOV 91 measurement is supported by their high-precision measurement of the 4d-2p transition energy in π^- ^{12}C , which is good agreement with the calculated energy.

While we suspect that the GALL 88 K^- Pb (9 → 8) measurements could be the problem, we are unable to find clear grounds for rejecting it. Therefore, we retain their measurement in the average and accept the large scale factor until further information can be obtained from new measurements and/or from reanalysis of GALL 88 and CHENG 75 data.

We thank B.L. Roberts (Boston Univ.) and Yu.M. Ivanov (Petersburg Nuclear Physics Inst.) for their extensive help in understanding this problem.

K^\pm MASS


VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
493.677 ± 0.016 OUR FIT	Error includes scale factor of 2.8.			
493.677 ± 0.013 OUR AVERAGE	Error includes scale factor of 2.4. See the ideogram below.			
493.696 ± 0.007	¹ DENISOV	91	CNTR	- Kaonic atoms
493.636 ± 0.011	² GALL	88	CNTR	- Kaonic atoms
493.640 ± 0.054	LUM	81	CNTR	- Kaonic atoms
493.670 ± 0.029	BARKOV	79	EMUL	± $e^+ e^- \rightarrow K^+ K^-$
493.657 ± 0.020	² CHENG	75	CNTR	- Kaonic atoms
493.691 ± 0.040	BACKENSTO...73	CNTR	-	Kaonic atoms
• • • We do not use the following data for averages, fits, limits, etc. • • •				
493.631 ± 0.007	GALL	88	CNTR	- K^- Pb (9 → 8)
493.675 ± 0.026	GALL	88	CNTR	- K^- Pb (11 → 10)
493.709 ± 0.073	GALL	88	CNTR	- K^- W (9 → 8)
493.806 ± 0.095	GALL	88	CNTR	- K^- W (11 → 10)
493.640 ± 0.022 ± 0.008	³ CHENG	75	CNTR	- K^- Pb (9 → 8)
493.658 ± 0.019 ± 0.012	³ CHENG	75	CNTR	- K^- Pb (10 → 9)
493.638 ± 0.035 ± 0.016	³ CHENG	75	CNTR	- K^- Pb (11 → 10)
493.753 ± 0.042 ± 0.021	³ CHENG	75	CNTR	- K^- Pb (12 → 11)
493.742 ± 0.081 ± 0.027	³ CHENG	75	CNTR	- K^- Pb (13 → 12)
493.662 ± 0.19	KUNSELMAN	74	CNTR	- Kaonic atoms
493.78 ± 0.17	GREINER	65	EMUL	+
493.7 ± 0.3	BARKAS	63	EMUL	-
493.9 ± 0.2	COHEN	57	RVUE	+

¹ Error increased from 0.0059 based on the error analysis in IVANOV 92.

² This value is the authors' combination of all of the separate transitions listed for this paper.

³ The CHENG 75 values for separate transitions were calculated from their Table 7 transition energies. The first error includes a 20% systematic error in the noncircular contaminant shift. The second error is due to a ±5 eV uncertainty in the theoretical transition energies.

WEIGHTED AVERAGE
493.677 ± 0.013 (Error scaled by 2.4)



$m_{K^+} - m_{K^-}$

Test of CPT.


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG
-0.032 ± 0.090	1.5M	⁴ FORD	72	ASPK ±

⁴ FORD 72 uses $m_{\pi^+} - m_{\pi^-} = +28 \pm 70$ keV.

K^\pm MEAN LIFE

VALUE (10^{-8} s)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
1.2371 ± 0.0029 OUR FIT		Error includes scale factor of 2.2.			
1.2369 ± 0.0032 OUR AVERAGE		Error includes scale factor of 2.4. See the ideogram below.			
1.2380 ± 0.0016	3M	OTT	71 CNTR	+	Stopping K
1.2272 ± 0.0036		LOBKOWICZ	69 CNTR	+	K in flight
1.221 ± 0.011		FORD	67 CNTR	±	
1.2443 ± 0.0038		FITCH	65B CNTR	+	K at rest
1.231 ± 0.011		BOYARSKI	62 CNTR	+	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
1.25 +0.22 -0.17	5	BARKAS	61 EMUL		
1.27 +0.36 -0.23	51	BHOWMIK	61 EMUL		
1.31 ± 0.08	293	NORDIN	61 HBC	-	
1.24 ± 0.07		NORDIN	61 RVUE	-	
1.38 ± 0.24	33	FREDEN	60B EMUL		
1.21 ± 0.06		BURROWES	59 CNTR		
1.60 ± 0.3	52	EISENBERG	58 EMUL		
0.95 +0.36 -0.25	5	ILOFF	56 EMUL		

⁵Old experiments with large errors excluded from averaging.



$$\frac{(\tau_{K^+} - \tau_{K^-})}{\tau_{\text{average}}}$$

This quantity is a measure of CPT invariance in weak interactions.

VALUE (%)	DOCUMENT ID	TECN
0.11 ± 0.09 OUR AVERAGE	Error includes scale factor of 1.2.	
0.090 ± 0.078	LOBKOWICZ	69 CNTR
0.47 ± 0.30	FORD	67 CNTR

K^+ DECAY MODES

K^- modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \mu^+ \nu_\mu$	(63.51 ± 0.18) %	S=1.3
$\Gamma_2 e^+ \nu_e$	(1.55 ± 0.07) × 10 ⁻⁵	
$\Gamma_3 \pi^+ \pi^0$	(21.16 ± 0.14) %	S=1.1
$\Gamma_4 \pi^+ \pi^+ \pi^-$	(5.59 ± 0.05) %	S=1.9
$\Gamma_5 \pi^+ \pi^0 \pi^0$	(1.73 ± 0.04) %	S=1.2
$\Gamma_6 \pi^0 \mu^+ \nu_\mu$	(3.18 ± 0.08) %	S=1.5
Called $K_{\mu 3}^+$.		
$\Gamma_7 \pi^0 e^+ \nu_e$	(4.82 ± 0.06) %	S=1.3
Called $K_{e 3}^+$.		
$\Gamma_8 \pi^0 \pi^0 e^+ \nu_e$	(2.1 ± 0.4) × 10 ⁻⁵	
$\Gamma_9 \pi^+ \pi^- e^+ \nu_e$	(3.91 ± 0.17) × 10 ⁻⁵	
$\Gamma_{10} \pi^+ \pi^- \mu^+ \nu_\mu$	(1.4 ± 0.9) × 10 ⁻⁵	
$\Gamma_{11} \pi^0 \pi^0 \pi^0 e^+ \nu_e$	< 3.5 × 10 ⁻⁶	CL=90%
$\Gamma_{12} \pi^+ \gamma \gamma$	[a] < 1 × 10 ⁻⁶	CL=90%
$\Gamma_{13} \pi^+ 3\gamma$	[a] < 1.0 × 10 ⁻⁴	CL=90%
$\Gamma_{14} e^+ \nu_e \nu \bar{\nu}$	< 6 × 10 ⁻⁵	CL=90%
$\Gamma_{15} \mu^+ \nu_\mu \nu \bar{\nu}$	< 6.0 × 10 ⁻⁶	CL=90%

$\Gamma_{16} \mu^+ \nu_\mu e^+ e^-$	(1.06 ± 0.32) × 10 ⁻⁶	
$\Gamma_{17} e^+ \nu_e e^+ e^-$	(2.1 +2.1 -1.1) × 10 ⁻⁷	
$\Gamma_{18} \mu^+ \nu_\mu \mu^+ \mu^-$	< 4.1 × 10 ⁻⁷	CL=90%
$\Gamma_{19} \mu^+ \nu_\mu \gamma$	[a,b] (5.50 ± 0.28) × 10 ⁻³	
$\Gamma_{20} \pi^+ \pi^0 \gamma$	[a,b] (2.75 ± 0.15) × 10 ⁻⁴	
$\Gamma_{21} \pi^+ \pi^0 \gamma$ (DE)	[a,c] (1.8 ± 0.4) × 10 ⁻⁵	
$\Gamma_{22} \pi^+ \pi^+ \pi^- \gamma$	[a,b] (1.04 ± 0.31) × 10 ⁻⁴	
$\Gamma_{23} \pi^+ \pi^0 \pi^0 \gamma$	[a,b] (7.4 +5.5 -2.9) × 10 ⁻⁶	
$\Gamma_{24} \pi^0 \mu^+ \nu_\mu \gamma$	[a,b] < 6.1 × 10 ⁻⁵	CL=90%
$\Gamma_{25} \pi^0 e^+ \nu_e \gamma$	[a,b] (2.62 ± 0.20) × 10 ⁻⁴	
$\Gamma_{26} \pi^0 e^+ \nu_e \gamma$ (SD)	[d] < 5.3 × 10 ⁻⁵	CL=90%
$\Gamma_{27} \pi^0 \pi^0 e^+ \nu_e \gamma$	< 5 × 10 ⁻⁶	CL=90%

Lepton Family number (LF), Lepton number (L), $\Delta S = \Delta Q$ (SQ) violating modes, or $\Delta S = 1$ weak neutral current (SI) modes

$\Gamma_{28} \pi^+ \pi^+ e^- \bar{\nu}_e$	SQ	< 1.2 × 10 ⁻⁸	CL=90%
$\Gamma_{29} \pi^+ \pi^+ \mu^- \bar{\nu}_\mu$	SQ	< 3.0 × 10 ⁻⁶	CL=95%
$\Gamma_{30} \pi^+ e^+ e^-$	SI	(2.74 ± 0.23) × 10 ⁻⁷	
$\Gamma_{31} \pi^+ \mu^+ \mu^-$	SI	< 2.3 × 10 ⁻⁷	CL=90%
$\Gamma_{32} \pi^+ \nu \bar{\nu}$	SI	< 5.2 × 10 ⁻⁹	CL=90%
$\Gamma_{33} \mu^- \nu e^+ e^+$	LF	< 2.0 × 10 ⁻⁸	CL=90%
$\Gamma_{34} \mu^+ \nu_e$	LF	[e] < 4 × 10 ⁻³	CL=90%
$\Gamma_{35} \pi^+ \mu^+ e^-$	LF	< 2.1 × 10 ⁻¹⁰	CL=90%
$\Gamma_{36} \pi^+ \mu^- e^+$	LF	< 7 × 10 ⁻⁹	CL=90%
$\Gamma_{37} \pi^- \mu^+ e^+$	L	< 7 × 10 ⁻⁹	CL=90%
$\Gamma_{38} \pi^- e^+ e^+$	L	< 1.0 × 10 ⁻⁸	CL=90%
$\Gamma_{39} \pi^- \mu^+ \mu^+$	L	< 1.5 × 10 ⁻⁴	CL=90%
$\Gamma_{40} \mu^+ \bar{\nu}_e$	L	[e] < 3.3 × 10 ⁻³	CL=90%
$\Gamma_{41} \pi^0 e^+ \bar{\nu}_e$	L	[e] < 3 × 10 ⁻³	CL=90%
$\Gamma_{42} \pi^+ \gamma$			

- [a] See the Full Listings below for the energy limits used in this measurement.
- [b] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ 's.
- [c] Direct-emission branching fraction.
- [d] Structure-dependent part.
- [e] Derived from an analysis of neutrino-oscillation experiments.

CONSTRAINED FIT INFORMATION

An overall fit to the mean life, 2 decay rate, and 20 branching ratios uses 60 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 74.9$ for 53 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-58						
x_4	-42	-12					
x_5	-28	-4	22				
x_6	-49	-17	14	2			
x_7	-51	-16	35	7	39		
x_8	-3	-1	2	0	2	6	
Γ	9	3	-21	-5	-3	-7	0
	x_1	x_3	x_4	x_5	x_6	x_7	x_8

Mode	Rate (10 ⁸ s ⁻¹)	Scale factor
$\Gamma_1 \mu^+ \nu_\mu$	0.5134 ± 0.0020	1.5
$\Gamma_3 \pi^+ \pi^0$	0.1711 ± 0.0012	1.2
$\Gamma_4 \pi^+ \pi^+ \pi^-$	0.0452 ± 0.0004	1.9
$\Gamma_5 \pi^+ \pi^0 \pi^0$	0.01400 ± 0.00032	1.2
$\Gamma_6 \pi^0 \mu^+ \nu_\mu$	0.0257 ± 0.0006	1.5
Called $K_{\mu 3}^+$.		
$\Gamma_7 \pi^0 e^+ \nu_e$	0.0390 ± 0.0005	1.3
Called $K_{e 3}^+$.		
$\Gamma_8 \pi^0 \pi^0 e^+ \nu_e$	(1.70 +0.34 -0.29) × 10 ⁻⁵	

Meson Full Listings

 K^\pm K^\pm DECAY RATES $\Gamma(\mu^+\nu_\mu)$ Γ_1

VALUE (units $10^6 s^{-1}$)	DOCUMENT ID	TECN	CHG
51.34 ± 0.20 OUR FIT			
51.2 ± 0.8	FORD	67	CNTR ±

 $\Gamma(\pi^+\pi^+\pi^-)$ Γ_4

VALUE (units $10^6 s^{-1}$)	EVTs	DOCUMENT ID	TECN	CHG
4.52 ± 0.04 OUR FIT				
4.511 ± 0.024		⁶ FORD	70	ASPK
4.529 ± 0.032	3.2M	⁶ FORD	70	ASPK
4.496 ± 0.030		⁶ FORD	67	CNTR ±

⁶ First FORD 70 value is second FORD 70 combined with FORD 67. $(\Gamma(K^+) - \Gamma(K^-)) / \Gamma(K)$ $K^\pm \rightarrow \mu^\pm \nu_\mu$ RATE DIFFERENCE/AVERAGE

Test of CPT conservation.

VALUE (%)	DOCUMENT ID	TECN
-0.54 ± 0.41	FORD	67

 $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ RATE DIFFERENCE/AVERAGE

Test of CP conservation.

VALUE (%)	EVTs	DOCUMENT ID	TECN	CHG
0.07 ± 0.12 OUR AVERAGE				
0.08 ± 0.12		⁷ FORD	70	ASPK
-0.50 ± 0.90		FLETCHER	67	OSPK
-0.02 ± 0.16		⁸ SMITH	73	ASPK ±
0.10 ± 0.14	3.2M	⁷ FORD	70	ASPK
-0.04 ± 0.21		⁷ FORD	67	CNTR

⁷ First FORD 70 value is second FORD 70 combined with FORD 67.⁸ SMITH 73 value of $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$ rate difference is derived from SMITH 73 value of $K^\pm \rightarrow \pi^\pm 2\pi^0$ rate difference. $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$ RATE DIFFERENCE/AVERAGE

Test of CP conservation.

VALUE (%)	EVTs	DOCUMENT ID	TECN	CHG
0.0 ± 0.6 OUR AVERAGE				
0.08 ± 0.58		SMITH	73	ASPK ±
-1.1 ± 1.8	1802	HERZO	69	OSPK

 $K^\pm \rightarrow \pi^\pm \pi^0$ RATE DIFFERENCE/AVERAGE

Test of CPT conservation.

VALUE (%)	DOCUMENT ID	TECN
0.8 ± 1.2	HERZO	69

 $K^\pm \rightarrow \pi^\pm \pi^0 \gamma$ RATE DIFFERENCE/AVERAGE

Test of CP conservation.

VALUE (%)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
0.9 ± 3.3 OUR AVERAGE					
0.8 ± 5.8	2461	SMITH	76	WIRE ±	E_π 55-90 MeV
1.0 ± 4.0	4000	ABRAMS	73B	ASPK ±	E_π 51-100 MeV
0.0 ± 24.0	24	EDWARDS	72	OSPK	E_π 58-90 MeV

 K^+ BRANCHING RATIOS $\Gamma(\mu^+\nu_\mu)/\Gamma_{total}$ Γ_1/Γ

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
63.51 ± 0.18 OUR FIT					
63.24 ± 0.44	62k	CHIANG	72	OSPK +	1.84 GeV/c K^+
56.9 ± 2.6		⁹ ALEXANDER	57	EMUL +	
58.5 ± 3.0		⁹ BIRGE	56	EMUL +	

⁹ Old experiments not included in averaging. $\Gamma(\mu^+\nu_\mu)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_1/Γ_4

VALUE	EVTs	DOCUMENT ID	TECN	CHG
11.36 ± 0.12 OUR FIT				
10.38 ± 0.82	427	¹⁰ YOUNG	65	EMUL +

¹⁰ Deleted from overall fit because YOUNG 65 constrains his results to add up to 1. Only YOUNG 65 measured $(\mu\nu)$ directly. $\Gamma(e^+\nu_e)/\Gamma_{total}$ Γ_2/Γ

VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	CHG
2.1^{+1.8}_{-1.3}		4	BOWEN	67B	OSPK +
<160.0		95	BORREANI	64	HBC +

 $\Gamma(e^+\nu_e)/\Gamma(\mu^+\nu_\mu)$ Γ_2/Γ_1

VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	CHG
2.45 ± 0.11 OUR AVERAGE				
2.51 ± 0.15	404	HEINTZE	76	SPEC +
2.37 ± 0.17	534	HEARD	75B	SPEC +
2.42 ± 0.42	112	CLARK	72	OSPK +
1.8 ^{+0.8} _{-0.6}	8	MACEK	69	ASPK +
1.9 ^{+0.7} _{-0.5}	10	BOTTERILL	67	ASPK +

 $\Gamma(\pi^+\pi^0)/\Gamma_{total}$ Γ_3/Γ

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
21.16 ± 0.14 OUR FIT					
21.18 ± 0.28	16k	CHIANG	72	OSPK +	1.84 GeV/c K^+
21.0 ± 0.6		CALLAHAN	65	HLBC	See $\Gamma(\pi^+\pi^0)/\Gamma(\pi^+\pi^+\pi^-)$
21.6 ± 0.6		TRILLING	65B	RVUE	
23.2 ± 2.2		¹¹ ALEXANDER	57	EMUL +	
27.7 ± 2.7		¹¹ BIRGE	56	EMUL +	

¹¹ Earlier experiments not averaged. $\Gamma(\pi^+\pi^0)/\Gamma(\mu^+\nu_\mu)$ Γ_3/Γ_1

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
0.3332 ± 0.0028 OUR FIT					
0.3316 ± 0.0032 OUR AVERAGE					
0.3329 ± 0.0047 ± 0.0010	45K	USHER	92	SPEC +	$p\bar{p}$ at rest
0.3355 ± 0.0057		¹² WEISSENBERG...	76	SPEC +	
0.305 ± 0.018	1600	ZELLER	69	ASPK +	
0.3277 ± 0.0065	4517	¹³ AUERBACH	67	OSPK +	
0.328 ± 0.005	25k	¹² WEISSENBERG...	74	STRC +	
0.329 ± 0.005		¹² WEISSENBERG	76	revises WEISSENBERG 74.	
0.3253 ± 0.0065		¹³ AUERBACH	67	changed from 0.3253 ± 0.0065. See comment with ratio $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\mu^+\nu_\mu)$.	


 $\Gamma(\pi^+\pi^0)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_3/Γ_4

VALUE	EVTs	DOCUMENT ID	TECN	CHG
3.79 ± 0.04 OUR FIT				
3.84 ± 0.27 OUR AVERAGE				
3.96 ± 0.15	1045	CALLAHAN	66	FBC +
3.24 ± 0.34	134	YOUNG	65	EMUL +

 $\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{total}$ Γ_4/Γ

VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
5.59 ± 0.06 OUR FIT					
5.52 ± 0.10 OUR AVERAGE					
5.34 ± 0.21	693	¹⁴ PANDOULAS	70	EMUL +	
5.71 ± 0.15		DEMARCO	65	HBC	
6.0 ± 0.4	44	YOUNG	65	EMUL +	
5.54 ± 0.12	2332	CALLAHAN	64	HLBC +	
5.1 ± 0.2	540	SHAKLEE	64	HLBC +	
5.7 ± 0.3		ROE	61	HLBC +	
5.56 ± 0.20	2330	¹⁵ CHIANG	72	OSPK +	1.84 GeV/c K^+
5.2 ± 0.3		¹⁶ TAYLOR	59	EMUL +	
6.8 ± 0.4		¹⁶ ALEXANDER	57	EMUL +	
5.6 ± 0.4		¹⁶ BIRGE	56	EMUL +	


¹⁴ Includes events of TAYLOR 59.¹⁵ Value is not independent of CHIANG 72 $\Gamma(\mu^+\nu_\mu)/\Gamma_{total}$, $\Gamma(\pi^+\pi^0)/\Gamma_{total}$, $\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{total}$, $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma_{total}$, and $\Gamma(\pi^0e^+\nu_e)/\Gamma_{total}$.¹⁶ Earlier experiments not averaged.



$\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{total}$ Γ_5/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.73±0.04 OUR FIT					Error includes scale factor of 1.2.
1.77±0.07 OUR AVERAGE					Error includes scale factor of 1.4. See the ideogram below.
1.84±0.06	1307	CHIANG 72	OSPK	+	1.84 GeV/c K^+
1.53±0.11	198	17 PANDOULAS 70	EMUL	+	
1.8 ±0.2	108	SHAKLEE 64	HLBC	+	
1.7 ±0.2		ROE 61	HLBC	+	
••• We do not use the following data for averages, fits, limits, etc. •••					
1.5 ±0.2		18 TAYLOR 59	EMUL	+	
2.2 ±0.4		18 ALEXANDER 57	EMUL	+	
2.1 ±0.5		18 BIRGE 56	EMUL	+	

¹⁷ Includes events of TAYLOR 59.
¹⁸ Earlier experiments not averaged.



$\Gamma(\pi^+\pi^0\pi^0)/\Gamma(\pi^+\pi^0\pi^-)$ Γ_5/Γ_3

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.0819±0.0020 OUR FIT					Error includes scale factor of 1.2.
0.081 ±0.005	574	¹⁹ LUCAS 73B	HBC	-	Dalitz pairs only

¹⁹ LUCAS 73B gives $N(\pi_2\pi^0) = 574 \pm 5.9\%$, $N(2\pi) = 3564 \pm 3.1\%$. We quote $0.5N(\pi_2\pi^0)/N(2\pi)$ where 0.5 is because only Dalitz pair π^0 's were used.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_5/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.310±0.007 OUR FIT					Error includes scale factor of 1.2.
0.304±0.009 OUR AVERAGE					
0.303±0.009	2027	BISI 65	BC	+	HBC+HLBC
0.393±0.099	17	YOUNG 65	EMUL	+	

$\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma_{total}$ Γ_6/Γ

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3.18±0.08 OUR FIT					Error includes scale factor of 1.5.
3.33±0.16	2345	CHIANG 72	OSPK	+	1.84 GeV/c K^+
••• We do not use the following data for averages, fits, limits, etc. •••					
2.8 ±0.4		20 TAYLOR 59	EMUL	+	
5.9 ±1.3		20 ALEXANDER 57	EMUL	+	
2.8 ±1.0		20 BIRGE 56	EMUL	+	

²⁰ Earlier experiments not averaged.

$\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\mu^+\nu_\mu)$ Γ_6/Γ_1


VALUE	EVTS	DOCUMENT ID	TECN	CHG	
0.0501±0.0013 OUR FIT					Error includes scale factor of 1.6.
0.0488±0.0026 OUR AVERAGE					
0.054 ±0.009	240	ZELLER 69	ASPK	+	
0.0480±0.0037	424	21 GARLAND 68	OSPK	+	
0.0486±0.0040	307	22 AUERBACH 67	OSPK	+	

²¹ GARLAND 68 changed from 0.055 ± 0.004 in agreement with μ -spectrum calculation of GAILLARD 70 appendix B. L.G.Pondrom, (private communication 73).
²² AUERBACH 67 changed from 0.0602 ± 0.0046 by erratum which brings the μ -spectrum calculation into agreement with GAILLARD 70 appendix B.

$\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\pi^+\pi^+\pi^-)$ Γ_6/Γ_4

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.569±0.014 OUR FIT					Error includes scale factor of 1.5.
0.517±0.032 OUR AVERAGE					Error includes scale factor of 1.8. See the ideogram below.
0.503±0.019	1505	23 HAIDT 71	HLBC	+	
0.63 ±0.07	2845	24 BISI 65B	BC	+	HBC+HLBC
0.90 ±0.16	38	YOUNG 65	EMUL	+	
••• We do not use the following data for averages, fits, limits, etc. •••					
0.510±0.017	1505	23 EICHTEN 68	HLBC	+	

²³ HAIDT 71 is a reanalysis of EICHTEN 68.
²⁴ Error enlarged for background problems. See GAILLARD 70.



$\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\pi^0e^+\nu_e)$ Γ_6/Γ_7

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.660±0.015 OUR FIT					Error includes scale factor of 1.5.
0.680±0.013 OUR AVERAGE					
0.705±0.063	554	25 LUCAS 73B	HBC	-	Dalitz pairs only
0.698±0.025	3480	26 CHIANG 72	OSPK	+	1.84 GeV/c K^+
0.667±0.017	5601	BOTTERILL 68B	ASPK	+	
0.703±0.056	1509	27 CALLAHAN 66B	HLBC		
••• We do not use the following data for averages, fits, limits, etc. •••					
0.670±0.014		28 HEINTZE 77	SPEC	+	
0.67 ±0.12		WEISSENBE... 76	SPEC	+	
0.608±0.014	1585	29 BRAUN 75	HLBC	+	
0.596±0.025		30 HAIDT 71	HLBC	+	
0.604±0.022	1398	30 EICHTEN 68	HLBC		

²⁵ LUCAS 73B gives $N(K_{\mu 3}) = 554 \pm 7.6\%$, $N(K_{e 3}) = 786 \pm 3.1\%$. We divide.
²⁶ CHIANG 72 $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\pi^0e^+\nu_e)$ is statistically independent of CHIANG 72 $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma_{total}$ and $\Gamma(\pi^0e^+\nu_e)/\Gamma_{total}$.
²⁷ From CALLAHAN 66B we use only the $K_{\mu 3}/K_{e 3}$ ratio and do not include in the fit the ratios $K_{\mu 3}/(\pi^+\pi^0)$ and $K_{e 3}/(\pi^+\pi^0)$, since they show large disagreements with the rest of the data.
²⁸ HEINTZE 77 value from fit to λ_0 . Assumes μ -e universality.
²⁹ BRAUN 75 value is from form factor fit. Assumes μ -e universality.
³⁰ HAIDT 71 is a reanalysis of EICHTEN 68. Only individual ratios included in fit (see $\Gamma(\pi^0\mu^+\nu_\mu)/\Gamma(\pi^+\pi^+\pi^-)$ and $\Gamma(\pi^0e^+\nu_e)/\Gamma(\pi^+\pi^+\pi^-)$).

$[\Gamma(\pi^+\pi^0) + \Gamma(\pi^0\mu^+\nu_\mu)]/\Gamma_{total}$ $(\Gamma_3 + \Gamma_6)/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	
24.34±0.15 OUR FIT					Error includes scale factor of 1.2.
24.6 ±1.0 OUR AVERAGE					Error includes scale factor of 1.4.
25.4 ±0.9	886	SHAKLEE 64	HLBC	+	
23.4 ±1.1		ROE 61	HLBC	+	

Meson Full Listings

 K^\pm

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{total}$						Γ_7/Γ	
VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT		
4.82 ± 0.06 OUR FIT		Error includes scale factor of 1.3.					
4.85 ± 0.09 OUR AVERAGE							
4.86 ± 0.10	3516	CHIANG	72	OSPK	+	1.84 GeV/c K^+	
4.7 ± 0.3	429	SHAKLEE	64	HLBC	+		
5.0 ± 0.5		ROE	61	HLBC	+		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
5.1 ± 1.3		³¹ ALEXANDER	57	EMUL	+		
3.2 ± 1.3		³¹ BIRGE	56	EMUL	+		
³¹ Earlier experiments not averaged.							

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma(\mu^+ \nu_\mu)$						Γ_7/Γ_1	
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT		
0.0789 ± 0.0011 OUR FIT		Error includes scale factor of 1.4.					
0.0762 ± 0.0024 OUR AVERAGE							
0.069 ± 0.006	350	ZELLER	69	ASPK	+		
0.0775 ± 0.0033	960	BOTTERILL	68C	ASPK	+		
0.069 ± 0.006	561	GARLAND	68	OSPK	+		
0.0791 ± 0.0054	295	³² AUERBACH	67	OSPK	+		

³² AUERBACH 67 changed from 0.0797 ± 0.0054. See comment with ratio $\Gamma(\pi^0 \mu^+ \nu_\mu)/\Gamma(\mu^+ \nu_\mu)$. The value 0.0785 ± 0.0025 given in AUERBACH 67 is an average of AUERBACH 67 $\Gamma(\pi^0 e^+ \nu_e)/\Gamma(\mu^+ \nu_\mu)$ and CESTER 66 $\Gamma(\pi^0 e^+ \nu_e)/[\Gamma(\mu^+ \nu_\mu) + \Gamma(\pi^+ \pi^0)]$.

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma(\pi^+ \pi^0)$						Γ_7/Γ_3	
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT		
0.228 ± 0.004 OUR FIT		Error includes scale factor of 1.3.					
0.221 ± 0.012	786	³³ LUCAS	73B	HBC	-	Dalitz pairs only	
³³ LUCAS 73B gives $N(K_{e3}) = 786 \pm 3.1\%$, $N(2\pi) = 3564 \pm 3.1\%$. We divide.							

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma(\pi^+ \pi^+ \pi^-)$						Γ_7/Γ_4	
VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT		
0.863 ± 0.011 OUR FIT		Error includes scale factor of 1.3.					
0.860 ± 0.014 OUR AVERAGE							
0.867 ± 0.027	2768	BARMIN	87	XEBC	+		
0.856 ± 0.040	2827	BRAUN	75	HLBC	+		
0.850 ± 0.019	4385	³⁴ HAIDT	71	HLBC	+		
0.94 ± 0.09	854	BELLOTTI	67B	HLBC	+		
0.90 ± 0.06	230	BORREANI	64	HBC	+		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
0.846 ± 0.021	4385	³⁴ EICHTEN	68	HLBC	+		
0.90 ± 0.16	37	YOUNG	65	EMUL	+		
³⁴ HAIDT 71 is a reanalysis of EICHTEN 68.							

$\Gamma(\pi^0 e^+ \nu_e)/[\Gamma(\mu^+ \nu_\mu) + \Gamma(\pi^+ \pi^0)]$						$\Gamma_7/(\Gamma_1 + \Gamma_3)$	
VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT		
5.69 ± 0.08 OUR FIT		Error includes scale factor of 1.4.					
6.01 ± 0.15 OUR AVERAGE							
5.92 ± 0.65		³⁵ WEISSENBE...	76	SPEC	+		
6.16 ± 0.22	5110	ESCHSTRUTH	68	OSPK	+		
5.89 ± 0.21	1679	CESTER	66	OSPK	+		

³⁵ Value calculated from WEISSENBERG 76 ($\pi^0 e \nu$), ($\mu \nu$), and ($\pi \pi^0$) values to eliminate dependence on our 1974 ($\pi 2\pi^0$) and ($\pi \pi^+ \pi^-$) fractions.

$\Gamma(\pi^0 \pi^0 e^+ \nu_e)/\Gamma(\pi^0 e^+ \nu_e)$						Γ_8/Γ_7
VALUE (units 10^{-4})	CL%	EVTs	DOCUMENT ID	TECN	CHG	
4.3 ± 0.9 OUR FIT						
4.1 ± 1.0 OUR AVERAGE						
4.2 ± 1.0		25	BOLOTOV	86B	CALO	-
3.8 ± 5.0		2	LJUNG	73	HLBC	+
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 37.0		90	ROMANO	71	HLBC	+

$\Gamma(\pi^0 \pi^0 e^+ \nu_e)/\Gamma_{total}$						Γ_8/Γ
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
2.1 ± 0.4 OUR FIT						
2.54 ± 0.89	10	BARMIN	88B	HLBC	+	

$\Gamma(\pi^+ \pi^- e^+ \nu_e)/\Gamma(\pi^+ \pi^+ \pi^-)$						Γ_9/Γ_4	
VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT		
6.99 ± 0.30 OUR AVERAGE		Error includes scale factor of 1.2.					
7.21 ± 0.32	30k	ROSSELET	77	SPEC	+		
7.36 ± 0.68	500	BOURQUIN	71	ASPK	+		
7.0 ± 0.9	106	SCHWEINB...	71	HLBC	+		
5.83 ± 0.63	269	ELY	69	HLBC	+		
• • • We do not use the following data for averages, fits, limits, etc. • • •							
6.7 ± 1.5	69	BIRGE	65	FBC	+		

$\Gamma(\pi^+ \pi^- \mu^+ \nu_\mu)/\Gamma_{total}$						Γ_{10}/Γ
VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.77 ± 0.54		1	CLINE	65	FBC	+
-0.50						

$\Gamma(\pi^+ \pi^- \mu^+ \nu_\mu)/\Gamma(\pi^+ \pi^+ \pi^-)$						Γ_{10}/Γ_4
VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
2.57 ± 1.55	7	BISI	67	DBC	-	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
~ 2.5	1	GREINER	64	EMUL	+	

$\Gamma(\pi^0 \pi^0 \pi^0 e^+ \nu_e)/\Gamma_{total}$						Γ_{11}/Γ
VALUE (units 10^{-6})	CL%	EVTs	DOCUMENT ID	TECN	CHG	
< 3.5	90	0	BOLOTOV	88	SPEC	-
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 9	90	0	BARMIN	92	XEBC	+

$\Gamma(\pi^+ \gamma \gamma)/\Gamma_{total}$						Γ_{12}/Γ
All values given here assume a phase space pion energy spectrum.						
VALUE (units 10^{-4})	CL%	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
< 0.01	90	0	ATIYA	90B	CALO	T_π 117-127 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 0.084	90	0	ASANO	82	CNTR	T_π 117-127 MeV
-0.42 ± 0.52	0	0	ABRAMS	77	SPEC	T_π < 92 MeV
< 0.35	90	0	LJUNG	73	HLBC	6-102, 114-127 MeV
< 0.5	90	0	KLEMS	71	OSPK	T_π < 117 MeV
-0.1 ± 0.6			CHEN	68	OSPK	T_π 60-90 MeV

$\Gamma(\pi^+ 3\gamma)/\Gamma_{total}$						Γ_{13}/Γ
Values given here assume a phase space pion energy spectrum.						
VALUE (units 10^{-4})	CL%	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
< 1.0	90	0	ASANO	82	CNTR	T_π 117-127 MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 3.0	90	0	KLEMS	71	OSPK	T_π > 117 MeV

$\Gamma(e^+ \nu_e \nu \bar{\nu})/\Gamma(e^+ \nu_e)$						Γ_{14}/Γ_2
VALUE	CL%	EVTs	DOCUMENT ID	TECN	CHG	
< 3.6	90	0	HEINTZE	79	SPEC	+

$\Gamma(\mu^+ \nu_\mu \nu \bar{\nu})/\Gamma_{total}$						Γ_{15}/Γ
VALUE (units 10^{-5})	CL%	EVTs	DOCUMENT ID	TECN	CHG	
< 6.0	90	0	³⁶ PANG	73	CNTR	+
³⁶ PANG 73 assumes μ spectrum from ν - ν interaction of BARDIN 70.						

$\Gamma(\mu^+ \nu_\mu e^+ e^-)/\Gamma(\pi^+ \pi^- e^+ \nu_e)$						Γ_{16}/Γ_9
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
27. ± 8.	14	³⁷ DIAMANT...	76	SPEC	+	Extrapolated BR
• • • We do not use the following data for averages, fits, limits, etc. • • •						
3.3 ± 0.9	14	³⁷ DIAMANT...	76	SPEC	+	$m_{ee} > 140$

³⁷ DIAMANT-BERGER 76 quotes this result times our 1975 $\pi^+ \pi^- e \nu$ BR ratio. The first DIAMANT-BERGER 76 value is the second value extrapolated to 0 to include low mass e pairs.

$\Gamma(e^+ \nu_e e^+ e^-)/\Gamma(\pi^+ \pi^- e^+ \nu_e)$						Γ_{17}/Γ_9
VALUE (units 10^{-2})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
0.54 ± 0.54	4	DIAMANT...	76	SPEC	+	
-0.27						

$\Gamma(\mu^+ \nu_\mu \mu^+ \mu^-)/\Gamma_{total}$						Γ_{18}/Γ
VALUE (units 10^{-7})	CL%	EVTs	DOCUMENT ID	TECN	CHG	
< 4.1	90	0	ATIYA	89	CNTR	+

$\Gamma(\mu^+ \nu_\mu \gamma)/\Gamma_{total}$						Γ_{19}/Γ
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
5.50 ± 0.28 OUR AVERAGE						
6.6 ± 1.5	38,39	DEMIDOV	90	XEBC		$P(\mu) < 231.5$ MeV/c
6.0 ± 0.9		BARMIN	88	HLBC	+	$P(\mu) < 231.5$ MeV/c
5.4 ± 0.3	40	AKIBA	85	SPEC		$P(\mu) < 231.5$ MeV/c

• • • We do not use the following data for averages, fits, limits, etc. • • •

3.5 ± 0.8 ^{39,41} DEMIDOV 90 XEBC $E(\gamma) > 20$ MeV

3.2 ± 0.5 ⁴² BARMIN 88 HLBC $E(\gamma) > 20$ MeV

5.8 ± 3.5 12 WEISSENBE... 74 STRC $E(\gamma) > 9$ MeV

³⁸ $P(\mu)$ cut given in DEMIDOV 90 paper, 235.1 MeV/c, is a misprint according to authors (private communication).

³⁹ DEMIDOV 90 quotes only inner bremsstrahlung (IB) part.

⁴⁰ Assumes μ -e universality and uses constraints from $K \rightarrow e \nu \gamma$.

⁴¹ Not independent of above DEMIDOV 90 value. Cuts differ.

⁴² Not independent of above BARMIN 88 value. Cuts differ.

$\Gamma(\pi^+\pi^0\gamma)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
2.75 ± 0.15 OUR AVERAGE							
2.71 ± 0.45		140	BOLOTOV	87	WIRE	$T\pi^-$ 55–90 MeV	
2.87 ± 0.32		2461	SMITH	76	WIRE	$T\pi^\pm$ 55–90 MeV	
2.71 ± 0.19		2100	ABRAMS	72	ASPK	$T\pi^\pm$ 55–90 MeV	
••• We do not use the following data for averages, fits, limits, etc. •••							
$1.5^{+1.1}_{-0.6}$			43 LJUNG	73	HLBC	$T\pi^+$ 55–80 MeV	
$2.6^{+1.5}_{-1.1}$			43 LJUNG	73	HLBC	$T\pi^+$ 55–90 MeV	
$6.8^{+3.7}_{-2.1}$		17	43 LJUNG	73	HLBC	$T\pi^+$ 55–102 MeV	
2.4 ± 0.8		24	EDWARDS	72	OSPK	$T\pi^+$ 58–90 MeV	
< 1.0		0	44 MALTSEV	70	HLBC	$T\pi^+$ < 55 MeV	
< 1.9		90	0	EMMERSON	69	OSPK	$T\pi^+$ 55–80 MeV
2.2 ± 0.7		18	CLINE	64	FBC	$T\pi^+$ 55–80 MeV	

43 The LJUNG 73 values are not independent.

44 MALTSEV 70 selects low π^+ energy to enhance direct emission contribution. $\Gamma(\pi^+\pi^0\gamma(\text{DE}))/\Gamma_{\text{total}}$ Γ_{21}/Γ Direct emission part of $\Gamma(\pi^+\pi^0\gamma)/\Gamma_{\text{total}}$.

VALUE (units 10^{-5})	DOCUMENT ID	TECN	CHG	COMMENT
1.8 ± 0.4 OUR AVERAGE				
$2.05 \pm 0.46^{+0.39}_{-0.23}$	BOLOTOV	87	WIRE	$T\pi^-$ 55–90 MeV
2.3 ± 3.2	SMITH	76	WIRE	$T\pi^\pm$ 55–90 MeV
$1.56 \pm 0.35 \pm 0.5$	ABRAMS	72	ASPK	$T\pi^\pm$ 55–90 MeV

 $\Gamma(\pi^+\pi^+\pi^-\gamma)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	CHG	COMMENT	
1.04 ± 0.31 OUR AVERAGE					
1.10 ± 0.48	7	BARMIN	89	XEBC	$E(\gamma) > 5$ MeV
1.0 ± 0.4		STAMER	65	EMUL	$E(\gamma) > 11$ MeV

 $\Gamma(\pi^+\pi^0\pi^0\gamma)/\Gamma(\pi^+\pi^0\pi^0)$ Γ_{23}/Γ_5

VALUE (units 10^{-4})	DOCUMENT ID	TECN	CHG	COMMENT
$4.3^{+3.2}_{-1.7}$				
	BOLOTOV	85	SPEC	$E(\gamma) > 10$ MeV

 $\Gamma(\pi^0\mu^+\nu\mu\gamma)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-5})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
< 6.1		90	0	LJUNG	73	HLBC	$E(\gamma) > 30$ MeV

 $\Gamma(\pi^0e^+\nu_e\gamma)/\Gamma(\pi^0e^+\nu_e)$ Γ_{25}/Γ_7

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.54 ± 0.04 OUR AVERAGE Error includes scale factor of 1.1.					
0.46 ± 0.08	82	45 BARMIN	91	XEBC	$E(\gamma) > 10$ MeV, $0.6 < \cos\theta_e \gamma < 0.9$
0.56 ± 0.04	192	46 BOLOTOV	86B	CALO	$E(\gamma) > 10$ MeV
0.76 ± 0.28	13	47 ROMANO	71	HLBC	$E(\gamma) > 10$ MeV
••• We do not use the following data for averages, fits, limits, etc. •••					
1.51 ± 0.25	82	45 BARMIN	91	XEBC	$E(\gamma) > 10$ MeV, $0.98 < \cos\theta_e \gamma < 0.99$
0.48 ± 0.20	16	48 LJUNG	73	HLBC	$E(\gamma) > 30$ MeV
$0.22^{+0.15}_{-0.10}$		48 LJUNG	73	HLBC	$E(\gamma) > 30$ MeV
0.53 ± 0.22		47 ROMANO	71	HLBC	$E(\gamma) > 30$ MeV
1.2 ± 0.8		BELLOTTI	67	HLBC	$E(\gamma) > 30$ MeV

45 BARMIN 91 quotes branching ratio $\Gamma(K \rightarrow e\pi^0\nu) + \Gamma(K \rightarrow \pi^+\pi^-\pi^0)/\Gamma_{\text{all}}$. The measured normalization is $[\Gamma(K \rightarrow e\pi^0\nu) + \Gamma(K \rightarrow \pi^+\pi^-\pi^0)]/\Gamma_{\text{all}}$. For comparison with other experiments we used $\Gamma(K \rightarrow e\pi^0\nu)/\Gamma_{\text{all}} = 0.0482$ to calculate the values quoted here.46 $\cos\theta(e\gamma)$ between 0.6 and 0.9.47 Both ROMANO 71 values are for $\cos\theta(e\gamma)$ between 0.6 and 0.9. Second value is for comparison with second LJUNG 73 value. We use lowest $E(\gamma)$ cut for Summary Table value. See ROMANO 71 for E_γ dependence.48 First LJUNG 73 value is for $\cos\theta(e\gamma) < 0.9$, second value is for $\cos\theta(e\gamma)$ between 0.6 and 0.9 for comparison with ROMANO 71. $\Gamma(\pi^0e^+\nu_e\gamma(\text{SD}))/\Gamma_{\text{total}}$ Γ_{26}/Γ

Structure-dependent part.

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	CHG	
< 5.3		90	BOLOTOV	86B CALO	–

 $\Gamma(\pi^0\pi^0e^+\nu_e\gamma)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
< 5		90	0	BARMIN	92	XEBC	$E_\gamma > 10$ MeV

 $\Gamma(\pi^+\pi^+e^-\nu_e)/\Gamma_{\text{total}}$ Γ_{28}/Γ Test of $\Delta S = \Delta Q$ rule.

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
••• We do not use the following data for averages, fits, limits, etc. •••							
< 9.0		95	0	SCHWEINB...	71	HLBC	+
< 6.9		95	0	ELY	69	HLBC	+
< 20		95		BIRGE	65	FBC	+

 $\Gamma(\pi^+\pi^+e^-\nu_e)/\Gamma(\pi^+\pi^-e^+\nu_e)$ Γ_{28}/Γ_9 Test of $\Delta S = \Delta Q$ rule.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
< 3		90	3	49 BLOCH	76	SPEC	
••• We do not use the following data for averages, fits, limits, etc. •••							
< 130		95	0	BOURQUIN	71	ASPK	
49 BLOCH 76 quotes 3.6×10^{-4} at CL = 95%, we convert.							

 $\Gamma(\pi^+\pi^+\mu^-\nu_\mu)/\Gamma_{\text{total}}$ Γ_{29}/Γ Test of $\Delta S = \Delta Q$ rule.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	CHG		
< 3.0		95	0	BIRGE	65	FBC	+

 $\Gamma(\pi^+e^+e^-)/\Gamma_{\text{total}}$ Γ_{30}/Γ Test for $\Delta S = 1$ weak neutral current. Allowed by combined first-order weak and electromagnetic interactions.

VALUE (units 10^{-7})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
2.74 ± 0.23 OUR AVERAGE							
$2.75 \pm 0.23 \pm 0.13$		500	50	ALLIEGRO	92	SPEC	+
2.7 ± 0.5		41	51	BLOCH	75	SPEC	+

••• We do not use the following data for averages, fits, limits, etc. •••

< 17		90		CENCE	74	ASPK	+	Three track events
< 2.7		90		CENCE	74	ASPK	+	Two track events
< 320		90		BEIER	72	OSPK	±	
< 44		90		BISI	67	DBC	±	
< 8.8		90		CLINE	67B	FBC	±	
< 24.5		90	1	CAMERINI	64	FBC	+	

50 ALLIEGRO 92 assumes a vector interaction with a form factor given by $\lambda = 0.105 \pm 0.035 \pm 0.015$ and a correlation coefficient of -0.82 .

51 BLOCH 75 assumes a vector interaction.

 $\Gamma(\pi^+\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{31}/Γ Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-7})	CL%	DOCUMENT ID	TECN	CHG		
< 2.3		90	ATIYA	89	CNTR	+
••• We do not use the following data for averages, fits, limits, etc. •••						
< 24		90	BISI	67	DBC	+
< 30		90	CAMERINI	65	FBC	+

 $\Gamma(\pi^+\nu\bar{\nu})/\Gamma_{\text{total}}$ Γ_{32}/Γ Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE (units 10^{-9})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
< 5.2		90	52	ATIYA	93	CNTR	+	
••• We do not use the following data for averages, fits, limits, etc. •••								
< 7.5		90		ATIYA	93	CNTR	+	$T(\pi)$ 115–127 MeV
< 17		90	0	ATIYA	93B	CNTR	+	$T(\pi)$ 60–100 MeV
< 34		90		ATIYA	90	CNTR	+	
< 140		90		ASANO	81B	CNTR	+	$T(\pi)$ 116–127 MeV
< 940		90		53 CABLE	73	CNTR	+	$T(\pi)$ 60–105 MeV
< 560		90		53 CABLE	73	CNTR	+	$T(\pi)$ 60–127 MeV
< 57000		90	0	54 LJUNG	73	HLBC	+	
< 1400		90		53 KLEMS	71	OSPK	+	$T(\pi)$ 117–127 MeV

52 Combining ATIYA 93 and ATIYA 93B results.

53 KLEMS 71 and CABLE 73 assume π spectrum same as K_{e3} decay. Second CABLE 73 limit combines CABLE 73 and KLEMS 71 data for vector interaction.

54 LJUNG 73 assumes vector interaction.

 $\Gamma(\mu^-\nu_e e^+)/\Gamma(\pi^+\pi^-e^+\nu_e)$ Γ_{33}/Γ_9

Test of lepton family number conservation.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	CHG		
< 0.5		90	0	55 DIAMANT-...	76	SPEC	+

55 DIAMANT-BERGER 76 quotes this result times our 1975 $\pi^+\pi^-e\nu$ BR ratio. $\Gamma(\mu^+\nu_e)/\Gamma_{\text{total}}$ Γ_{34}/Γ

Forbidden by lepton family number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
< 0.004		90	0	LYONS	81	HLBC	0	200 GeV K^+ narrow band ν beam
••• We do not use the following data for averages, fits, limits, etc. •••								
< 0.012		90		COOPER	82	HLBC		Wideband ν beam

 $\Gamma(\pi^+\mu^+e^-)/\Gamma_{\text{total}}$ Γ_{35}/Γ

Test of lepton family number conservation.

VALUE (units 10^{-10})	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT		
< 2.1		90	0	LEE	90	SPEC	+	
••• We do not use the following data for averages, fits, limits, etc. •••								
< 11		90	0	CAMPAGNARI	88	SPEC	+	In LEE 90
< 48		90	0	DIAMANT-...	76	SPEC	+	

Meson Full Listings

K^\pm

$\Gamma(\pi^+\mu^-e^+)/\Gamma_{total}$ Γ_{36}/Γ
 Test of lepton family number conservation.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	CHG
< 7	90	0	56 DIAMANT-...	76	SPEC +
••• We do not use the following data for averages, fits, limits, etc. •••					
<28	90		56 BEIER	72	OSPK ±
56 Measurement actually applies to the sum of the $\pi^+\mu^-e^+$ and $\pi^-\mu^+e^+$ modes.					

$\Gamma(\pi^-\mu^+e^+)/\Gamma_{total}$ Γ_{37}/Γ
 Test of total lepton number conservation.

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	CHG
< 7	90	0	57 DIAMANT-...	76	SPEC +
••• We do not use the following data for averages, fits, limits, etc. •••					
<28	90		57 BEIER	72	OSPK ±
57 Measurement actually applies to the sum of the $\pi^+\mu^-e^+$ and $\pi^-\mu^+e^+$ modes.					

$\Gamma(\pi^+\mu^-e^+)/\Gamma_{total}$ Γ_{36}/Γ
 Test of total lepton number conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	CHG
<1.4	90	BEIER	72	OSPK ±

$\Gamma(\pi^-e^+e^+)/\Gamma_{total}$ Γ_{38}/Γ
 Test of total lepton number conservation.

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN	CHG
<1.5		CHANG	68	HBC -

$\Gamma(\pi^-e^+e^+)/\Gamma(\pi^+\pi^-e^+\nu_e)$ Γ_{38}/Γ_9
 Test of total lepton number conservation.

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	CHG
<2.5	90	0	58 DIAMANT-...	76	SPEC +
58 DIAMANT-BERGER 76 quotes this result times our 1975 BR ratio.					

$\Gamma(\pi^-\mu^+\mu^+)/\Gamma_{total}$ Γ_{39}/Γ
 Forbidden by total lepton number conservation.

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	CHG	
<1.5	90	59 LITTENBERG	92	HBC	
59 LITTENBERG 92 is from retroactive data analysis of CHANG 68 bubble chamber data.					

$\Gamma(\mu^+\nu_e)/\Gamma_{total}$ Γ_{40}/Γ
 Forbidden by total lepton number conservation.

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
<3.3	90	COOPER	82	HLBC Wideband ν beam

$\Gamma(\pi^0e^+\nu_e)/\Gamma_{total}$ Γ_{41}/Γ
 Forbidden by total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003	90	COOPER	82	HLBC Wideband ν beam

$\Gamma(\pi^+\gamma)/\Gamma_{total}$ Γ_{42}/Γ
 Violates angular momentum conservation. Not listed in Summary Table.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN	CHG	
<1.4	90	ASANO	82	CNTR +	
<4.0	90	60 KLEMS	71	OSPK +	
60 Test of model of Selleri, Nuovo Cimento 60A 291 (1969).					

K^+ LONGITUDINAL POLARIZATION OF EMITTED μ^+

$K^+ \rightarrow \mu^+\nu$

Tests for right-handed currents in strangeness-changing decay.

VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
> 0.990	90	IMAZATO	92	SPEC +	KEK 12 GeV protons
••• We do not use the following data for averages, fits, limits, etc. •••					
-0.970 ± 0.047		YAMANAKA	86	SPEC +	
-1.0 ± 0.1		CUTTS	69	SPRK +	
-0.96 ± 0.12		COOMBES	57	CNTR +	

NOTE ON DALITZ PLOT PARAMETERS FOR $K \rightarrow 3\pi$ DECAYS

The Dalitz plot distribution for $K^\pm \rightarrow \pi^\pm\pi^\pm\pi^\mp$, $K^\pm \rightarrow \pi^0\pi^0\pi^\pm$, and $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ can be parameterized by a series expansion such as that introduced by Weinberg [1]. We use the form

$$|M|^2 \propto 1 + g \frac{(s_3 - s_0)}{am_{\pi^+}^2} + h \left[\frac{s_3 - s_0}{m_{\pi^+}^2} \right]^2 + j \frac{(s_2 - s_1)}{m_{\pi^+}^2} + ak \left[\frac{s_2 - s_1}{m_{\pi^+}^2} \right]^2 + \dots, \quad (1)$$

where $m_{\pi^+}^2$ has been introduced to make the coefficients g , h , j , and k dimensionless, and

$$s_i = (P_K - P_i)^2 = (m_K - m_i)^2 - 2m_K T_i, \quad i = 1, 2, 3,$$

$$s_0 = \frac{1}{3} \sum_i s_i = \frac{1}{3} (m_K^2 + m_1^2 + m_2^2 + m_3^2)$$

Here the P_i are four-vectors, m_i and T_i are the mass and kinetic energy of the i^{th} pion, and the index 3 is used for the odd pion.

The coefficient g is a measure of the slope in the variable s_3 (or T_3) of the Dalitz plot, while h and k measure the quadratic dependence on s_3 and $(s_2 - s_1)$, respectively. The coefficient j is related to the asymmetry of the plot and must be zero if CP invariance holds. Note also that if CP is good, g , h , and k must be the same for $K^+ \rightarrow \pi^+\pi^+\pi^-$ as for $K^- \rightarrow \pi^-\pi^-\pi^+$.

Since different experiments use different forms for $|M|^2$, in order to compare the experiments we have converted to g , h , j , and k whatever coefficients have been measured. Where such conversions have been done, the measured coefficient a_y , a_t , a_u , or a_v is given in the comment at the right. For definitions of these coefficients, details of this conversion, and discussion of the data, see the April 1982 version of this note [2].

References

- S. Weinberg, Phys. Rev. Lett. **4**, 87 (1960).
- Particle Data Group, Phys. Lett. **111B**, 69 (1982).

ENERGY DEPENDENCE OF K^\pm DALITZ PLOT

$$|\text{matrix element}|^2 = 1 + gu + hu^2 + kv^2$$

where $u = (s_3 - s_0) / m_{\pi^+}^2$ and $v = (s_1 - s_2) / m_{\pi^+}^2$


LINEAR COEFFICIENT g_{π^+} FOR $K^+ \rightarrow \pi^+\pi^+\pi^-$

Some experiments use Dalitz variables x and y . In the comments we give $a_y =$ coefficient of y term. See note above on "Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays." For discussion of the conversion of a_y to g , see the earlier version of the same note in the Review published in Physics Letters **111B** 70 (1982).

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.2154 ± 0.0035 OUR AVERAGE					Error includes scale factor of 1.4. See the Ideogram below.
-0.2221 ± 0.0065	225k	DEVAUX	77	SPEC +	$a_y = 0.2814 \pm 0.0082$
-0.2157 ± 0.0028	750k	FORD	72	ASPK +	$a_y = 0.2734 \pm 0.0035$
-0.200 ± 0.009	39819	61 HOFFMASTER	72	HLBC +	
••• We do not use the following data for averages, fits, limits, etc. •••					
-0.196 ± 0.012	17898	62 GRAUMAN	70	HLBC +	$a_y = 0.228 \pm 0.030$
-0.218 ± 0.016	9994	63 BUTLER	68	HBC +	$a_y = 0.277 \pm 0.020$
-0.22 ± 0.024	5428	63,64 ZINCHENKO	67	HBC +	$a_y = 0.28 \pm 0.03$

61 HOFFMASTER 72 includes GRAUMAN 70 data.
 62 Emulsion data added — all events included by HOFFMASTER 72.
 63 Experiments with large errors not included in average.
 64 Also includes DBC events.

WEIGHTED AVERAGE
 -0.2154 ± 0.0035 (Error scaled by 1.4)




Linear energy dependence for $K^+ \rightarrow \pi^+\pi^+\pi^-$

See key on page 1343

Meson Full Listings
K±

QUADRATIC COEFFICIENT h FOR $K^+ \rightarrow \pi^+ \pi^+ \pi^-$


VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.012 ± 0.008 OUR AVERAGE				
Error includes scale factor of 1.4. See the Ideogram below.				
-0.0006 ± 0.0143	225k	DEVAUX	77 SPEC	+
0.0187 ± 0.0062	750k	FORD	72 ASPK	+
-0.009 ± 0.014	39819	HOFFMASTER72	HLBC	+



Quadratic coefficient h for $K^+ \rightarrow \pi^+ \pi^+ \pi^-$

QUADRATIC COEFFICIENT k FOR $K^+ \rightarrow \pi^+ \pi^+ \pi^-$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
-0.0101 ± 0.0034 OUR AVERAGE				
Error includes scale factor of 2.1. See the Ideogram below.				
-0.0205 ± 0.0039	225k	DEVAUX	77 SPEC	+
-0.0075 ± 0.0019	750k	FORD	72 ASPK	+
-0.0105 ± 0.0045	39819	HOFFMASTER72	HLBC	+



Quadratic coefficient k for $K^+ \rightarrow \pi^+ \pi^+ \pi^-$

LINEAR COEFFICIENT g_y FOR $K^+ \rightarrow \pi^+ \pi^+ \pi^-$

Some experiments use Dalitz variables x and y . In the comments we give $a_y =$ coefficient of y term. See note above on "Dalitz Plot Parameters for $K \rightarrow 3\pi$ Decays." For discussion of the conversion of a_y to g_y , see the earlier version of the same note in the Review published in Physics Letters 111B 70 (1982).

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.217 ± 0.007 OUR AVERAGE					
Error includes scale factor of 2.5.					
-0.2186 ± 0.0028	750k	FORD	72 ASPK	-	$a_y = 0.2770 \pm 0.0035$
-0.193 ± 0.010	50919	MAST	69 HBC	-	$a_y = 0.244 \pm 0.013$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
-0.199 ± 0.008	81k	65 LUCAS	73 HBC	-	$a_y = 0.252 \pm 0.011$
-0.190 ± 0.023	5778	66,67 MOSCOSO	68 HBC	-	$a_y = 0.242 \pm 0.029$
-0.220 ± 0.035	1347	68 FERRO-LUZZI	61 HBC	-	$a_y = 0.28 \pm 0.045$

65 Quadratic dependence is required by K_L^0 experiments. For comparison we average only those K^\pm experiments which quote quadratic fit values.
66 Experiments with large errors not included in average.
67 Also includes DBC events.
68 No radiative corrections included.

QUADRATIC COEFFICIENT h FOR $K^- \rightarrow \pi^- \pi^- \pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
0.010 ± 0.006 OUR AVERAGE				
0.0125 ± 0.0062	750k	FORD	72 ASPK	-
-0.001 ± 0.012	50919	MAST	69 HBC	-

QUADRATIC COEFFICIENT k FOR $K^- \rightarrow \pi^- \pi^- \pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	CHG
-0.0084 ± 0.0019 OUR AVERAGE				
-0.0083 ± 0.0019	750k	FORD	72 ASPK	-
-0.014 ± 0.012	50919	MAST	69 HBC	-


$(g_{\pi^+} - g_{\pi^-}) / (g_{\pi^+} + g_{\pi^-})$ FOR $K^\pm \rightarrow \pi^\pm \pi^+ \pi^-$
A nonzero value for this quantity indicates CP violation.

VALUE (%)	EVTS	DOCUMENT ID	TECN	CHG
-0.70 ± 0.53	3.2M	FORD	70 ASPK	

LINEAR COEFFICIENT g FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$
Unless otherwise stated, all experiments include terms quadratic in $(s_3 - s_0) / m_{\pi^+}^2$. See mini-review above.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.594 ± 0.019 OUR AVERAGE					Error includes scale factor of 1.3. See the ideogram below.
0.582 ± 0.021	43k	BOLOTOV	86 CALO	-	
0.670 ± 0.054	3263	BRAUN	76B HLBC	+	
0.630 ± 0.038	5635	SHEAFF	75 HLBC	+	
0.510 ± 0.060	27k	SMITH	75 WIRE	+	
0.67 ± 0.06	1365	AUBERT	72 HLBC	+	
0.544 ± 0.048	4048	DAVISON	69 HLBC	+	Also emulsion
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.806 ± 0.220	4639	69 BERTRAND	76 EMUL	+	
0.484 ± 0.084	574	70 LUCAS	73B HBC	-	Dalitz pairs only
0.527 ± 0.102	198	69 PANDOULAS	70 EMUL	+	
0.586 ± 0.098	1874	70 BISI	65 HLBC	+	Also HBC
0.48 ± 0.04	1792	70 KALMUS	64 HLBC	+	

69 Experiments with large errors not included in average.
70 Authors give linear fit only.



Linear energy dependence for $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$

QUADRATIC COEFFICIENT h FOR $K^\pm \rightarrow \pi^\pm \pi^0 \pi^0$
See mini-review above.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.035 ± 0.015 OUR AVERAGE					
0.037 ± 0.024	43k	BOLOTOV	86 CALO	-	
0.152 ± 0.082	3263	BRAUN	76B HLBC	+	
0.041 ± 0.030	5635	SHEAFF	75 HLBC	+	
0.009 ± 0.040	27k	SMITH	75 WIRE	+	
-0.01 ± 0.08	1365	AUBERT	72 HLBC	+	
0.026 ± 0.050	4048	DAVISON	69 HLBC	+	Also emulsion
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.164 ± 0.121	4639	71 BERTRAND	76 EMUL	+	
0.018 ± 0.124	198	71 PANDOULAS	70 EMUL	+	

71 Experiments with large errors not included in average.

Meson Full Listings

K^\pm

NOTE ON K_{e3}^\pm AND K_{e3}^0 FORM FACTORS

Assuming that only the vector current contributes to $K \rightarrow \pi \ell \nu$ decays, we write the matrix element as

$$M \propto f_+(t) [(P_K + P_\pi)_\mu \bar{\ell} \gamma_\mu (1 + \gamma_5) \nu] + f_-(t) [m_\ell \bar{\ell} (1 + \gamma_5) \nu], \quad (1)$$

where P_K and P_π are the four-momenta of the K and π mesons, m_ℓ is the lepton mass, and f_+ and f_- are dimensionless form factors which can depend only on $t = (P_K - P_\pi)^2$, the square of the four-momentum transfer to the leptons. If time-reversal invariance holds, f_+ and f_- are relatively real. $K_{\mu 3}$ experiments measure f_+ and f_- , while K_{e3} experiments are sensitive only to f_+ because the small electron mass makes the f_- term negligible.

(a) $K_{\mu 3}$ experiments. Analyses of $K_{\mu 3}$ data frequently assume a linear dependence of f_+ and f_- on t , i.e.,

$$f_\pm(t) = f_\pm(0) [1 + \lambda_\pm(t/m_\pi^2)] \quad (2)$$

Most $K_{\mu 3}$ data are adequately described by Eq. (2) for f_+ and a constant f_- (i.e., $\lambda_- = 0$). There are two equivalent parametrizations commonly used in these analyses:

(1) $\lambda_+, \xi(0)$ parametrization. Analyses of $K_{\mu 3}$ data often introduce the ratio of the two form factors

$$\xi(t) = f_-(t)/f_+(t).$$

The $K_{\mu 3}$ decay distribution is then described by the two parameters λ_+ and $\xi(0)$ (assuming time reversal invariance and $\lambda_- = 0$). These parameters can be determined by three different methods:

Method A. By studying the Dalitz plot or the pion spectrum of $K_{\mu 3}$ decay. The Dalitz plot density is (see, e.g., Chounet *et al.* [1]):

$$\rho(E_\pi, E_\mu) \propto f_+^2(t) [A + B\xi(t) + C\xi(t)^2],$$

where

$$A = m_K (2E_\mu E_\nu - m_K E'_\pi) + m_\mu^2 \left(\frac{1}{4} E'_\pi - E_\nu \right),$$

$$B = m_\mu^2 \left(E_\nu - \frac{1}{2} E'_\pi \right),$$

$$C = \frac{1}{4} m_\mu^2 E'_\pi,$$

$$E'_\pi = E_\pi^{\max} - E_\pi = (m_K^2 + m_\pi^2 - m_\mu^2) / 2m_K - E_\pi.$$

Here E_π , E_μ , and E_ν are, respectively, the pion, muon, and neutrino energies in the kaon center of mass. The density ρ is fit to the data to determine the values of $\lambda_+, \xi(0)$, and their correlation.

Method B. By measuring the $K_{\mu 3}/K_{e3}$ branching ratio and comparing it with the theoretical ratio (see, e.g., Fearing *et al.* [2]) as given in terms of λ_+ and $\xi(0)$, assuming μ - e universality:

$$\Gamma(K_{\mu 3}^\pm)/\Gamma(K_{e3}^\pm) = 0.6457 + 1.4115\lambda_+ + 0.1264\xi(0) + 0.0192\xi(0)^2 + 0.0080\lambda_+\xi(0),$$

$$\Gamma(K_{\mu 3}^0)/\Gamma(K_{e3}^0) = 0.6452 + 1.3162\lambda_+ + 0.1264\xi(0) + 0.0186\xi(0)^2 + 0.0064\lambda_+\xi(0).$$

This cannot determine λ_+ and $\xi(0)$ simultaneously but simply fixes a relationship between them.

Method C. By measuring the muon polarization in $K_{\mu 3}$ decay. In the rest frame of the K , the μ is expected to be polarized in the direction \mathbf{A} with $\mathbf{P} = \mathbf{A}/|\mathbf{A}|$, where \mathbf{A} is given (Cabibbo and Maksymowicz [3]) by

$$\begin{aligned} \mathbf{A} = & a_1(\xi) \mathbf{p}_\mu \\ & - a_2(\xi) \left[\frac{\mathbf{p}_\mu}{m_\mu} \left(m_K - E_\pi + \frac{\mathbf{p}_\pi \cdot \mathbf{p}_\mu}{|\mathbf{p}_\mu|^2} (E_\mu - m_\mu) \right) + \mathbf{p}_\pi \right] \\ & + m_K \text{Im}\xi(t) (\mathbf{p}_\pi \times \mathbf{p}_\mu). \end{aligned}$$

If time-reversal invariance holds, ξ is real, and thus there is no polarization perpendicular to the K -decay plane. Polarization experiments measure the weighted average of $\xi(t)$ over the t range of the experiment, where the weighting accounts for the variation with t of the sensitivity to $\xi(t)$.

(2) λ_+, λ_0 parametrization. Most of the more recent $K_{\mu 3}$ analyses have parameterized in terms of the form factors f_+ and f_0 which are associated with vector and scalar exchange, respectively, to the lepton pair. f_0 is related to f_+ and f_- by

$$f_0(t) = f_+(t) + [t/(m_K^2 - m_\pi^2)] f_-(t).$$

Here $f_0(0)$ must equal $f_+(0)$ unless $f_-(t)$ diverges at $t = 0$. The earlier assumption that f_+ is linear in t and f_- is constant leads to f_0 linear in t :

$$f_0(t) = f_0(0) [1 + \lambda_0(t/m_\pi^2)].$$

With the assumption that $f_0(0) = f_+(0)$, the two parametrizations, $(\lambda_+, \xi(0))$ and (λ_+, λ_0) are equivalent as long as correlation information is retained. (λ_+, λ_0) correlations tend to be less strong than $(\lambda_+, \xi(0))$ correlations.

The experimental results for $\xi(0)$ and its correlation with λ_+ are listed in the K^\pm and K_L^0 sections of the Full Listings in section ξ_A , ξ_B , or ξ_C depending on whether method A, B, or C discussed above was used. The corresponding values of λ_+ are also listed.

Because recent experiments tend to use the (λ_+, λ_0) parametrization, we include a subsection for λ_0 results. Whenever possible we have converted $\xi(0)$ results into λ_0 results and vice versa.

See the 1982 version of this note [4] for additional discussion of the $K_{\mu 3}^0$ parameters, correlations, and conversion between parametrizations, and also for a comparison of the experimental results.

(b) K_{e3} experiments. Analysis of K_{e3} data is simpler than that of $K_{\mu 3}$ because the second term of the matrix element assuming a pure vector current [Eq. (1) above] can be neglected. Here f_+ is usually assumed to be linear in t , and the linear coefficient λ_+ of Eq. (2) is determined.

If we remove the assumption of a pure vector current, then the matrix element for the decay, in addition to the terms in Eq. (2), would contain

$$+2m_K f_S \bar{\ell}(1 + \gamma_5)\nu$$

$$+(2f_T/m_K)(P_K)_\lambda(P_\pi)_\mu \bar{\ell} \sigma_{\lambda\mu}(1 + \gamma_5)\nu,$$

where f_S is the scalar form factor, and f_T is the tensor form factor. In the case of the K_{e3} decays where the f_- term can be neglected, experiments have yielded limits on $|f_S/f_+|$ and $|f_T/f_+|$.

References

1. L.M. Chounet, J.M. Gaillard, and M.K. Gaillard, Phys. Reports **4C**, 199 (1972).
2. H.W. Fearing, E. Fischbach, and J. Smith, Phys. Rev. **D2**, 542 (1970).
3. N. Cabibbo and A. Maksymowicz, Phys. Lett. **9**, 352 (1964).
4. Particle Data Group, Phys. Lett. **111B**, 73 (1982).

K_{e3}^\pm FORM FACTORS

In the form factor comments, the following symbols are used.

f_+ and f_- are form factors for the vector matrix element.

f_S and f_T refer to the scalar and tensor term.

$f_0 = f_+ + f_- t/(m_K^2 - m_\pi^2)$.

λ_+ , λ_- , and λ_0 are the linear expansion coefficients of f_+ , f_- , and f_0 .

λ_+ refers to the $K_{\mu 3}^\pm$ value except in the K_{e3}^\pm sections.

$d\xi(0)/d\lambda_+$ is the correlation between $\xi(0)$ and λ_+ in $K_{\mu 3}^\pm$.

$d\lambda_0/d\lambda_+$ is the correlation between λ_0 and λ_+ in $K_{\mu 3}^\pm$.

t = momentum transfer to the π in units of m_π^2 .

DP = Dalitz plot analysis.

PI = π spectrum analysis.

MU = μ spectrum analysis.

POL = μ polarization analysis.

BR = $K_{\mu 3}^\pm/K_{e3}^\pm$ branching ratio analysis.

E = positron or electron spectrum analysis.

RC = radiative corrections.

λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN K_{e3}^\pm DECAY)

For radiative correction of K_{e3}^\pm Dalitz plot, see GINSBERG 67 and BECHERRAWY 70.

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.0286 ± 0.0022 OUR AVERAGE					
0.0284 ± 0.0027 ± 0.0020	32k	72 AKIMENKO	91 SPEC		PI, no RC
0.029 ± 0.004	62k	73 BOLOTOV	88 SPEC		PI, no RC
0.027 ± 0.008		74 BRAUN	73B HLBC	+	DP, no RC
0.029 ± 0.011	4017	CHIANG	72 OSPK	+	DP, RC negligible
0.027 ± 0.010	2707	STEINER	71 HLBC	+	DP, uses RC
0.045 ± 0.015	1458	BOTTERILL	70 OSPK		PI, uses RC
0.08 ± 0.04	960	BOTTERILL	68C ASPK	+	e^+ , uses RC
-0.02 ± 0.08	90	EISLER	68 HLBC	+	PI, uses RC
-0.12					
0.045 ± 0.017	854	BELLOTTI	67B FBC	+	DP, uses RC
-0.018					
+0.016 ± 0.016	1393	IMLAY	67 OSPK	+	DP, no RC
+0.028 ± 0.013	515	KALMUS	67 FBC	+	e^+ , PI, no RC
-0.014					
-0.04 ± 0.05	230	BORREANI	64 HBC	+	e^+ , no RC
-0.010 ± 0.029	407	JENSEN	64 XEBC	+	PI, no RC
+0.036 ± 0.045	217	BROWN	62B XEBC	+	PI, no RC

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.025 ± 0.007		75 BRAUN	74 HLBC	+	$K_{\mu 3}/K_{e3}$ vs. t
		72 AKIMENKO	91		state that radiative corrections would raise λ_+ by 0.0013.
		73 BOLOTOV	88		state radiative corrections of GINSBERG 67 would raise λ_+ by 0.002.
		74 BRAUN	73B		states that radiative corrections of GINSBERG 67 would lower λ_+^e by 0.002 but that radiative corrections of BECHERRAWY 70 disagrees and would raise λ_+^e by 0.005.
		75 BRAUN	74		is a combined $K_{\mu 3}/K_{e3}$ result. It is not independent of BRAUN 73C ($K_{\mu 3}$) and BRAUN 73B (K_{e3}) form factor results.

$\xi_A = f_-/f_+$ (determined from $K_{\mu 3}^\pm$ spectra)

The parameter ξ is redundant with λ_0 below and is not put into the Meson Summary Table.

VALUE	$d\xi(0)/d\lambda_+$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
-0.38 ± 0.15 OUR EVALUATION						From a fit discussed in note on K_{e3} form factors in 1982 edition, PL 111B (April 1982).
-0.27 ± 0.25	-17	3973	WHITMAN	80 SPEC	+	DP
-0.8 ± 0.8	-20	490	76 ARNOLD	74 HLBC	+	DP
-0.57 ± 0.24	-9	6527	77 MERLAN	74 ASPK	+	DP
-0.36 ± 0.40	-19	1897	78 BRAUN	73C HLBC	+	DP
-0.62 ± 0.28	-12	4025	79 ANKENBRA...	72 ASPK	+	PI
+0.45 ± 0.28	-15	3480	80 CHIANG	72 OSPK	+	DP
-1.1 ± 0.56	-29	3240	81 HAIDT	71 HLBC	+	DP
-0.5 ± 0.8	-26	2041	82 KIJEWski	69 OSPK	+	PI
+0.72 ± 0.93	-17	444	CALLAHAN	66B FBC	+	PI
						• • • We do not use the following data for averages, fits, limits, etc. • • •
-0.5 ± 0.9	none	78	EISLER	68 HLBC	+	PI, $\lambda_+ = 0$
0.0 ± 1.1	-0.9	2648	83 CALLAHAN	66B FBC	+	μ , $\lambda_+ = 0$
+0.7 ± 0.5		87	GIACOMELLI	64 EMUL	+	MU+BR, $\lambda_+ = 0$
-0.08 ± 0.7			84 JENSEN	64 XEBC	+	DP+BR
+1.8 ± 0.6		76	BROWN	62B XEBC	+	DP+BR, $\lambda_+ = 0$

76 ARNOLD 74 figure 4 was used to obtain ξ_A and $d\xi(0)/d\lambda_+$.

77 MERLAN 74 figure 5 was used to obtain $d\xi(0)/d\lambda_+$.

78 BRAUN 73C gives $\xi(t) = -0.34 \pm 0.20$, $d\xi(t)/d\lambda_+ = -14$ for $\lambda_+ = 0.027$, $t = 6.6$. We calculate above $\xi(0)$ and $d\xi(0)/d\lambda_+$ for their $\lambda_+ = 0.025 \pm 0.017$.

79 ANKENBRANDT 72 figure 3 was used to obtain $d\xi(0)/d\lambda_+$.

80 CHIANG 72 figure 10 was used to obtain $d\xi(0)/d\lambda_+$. Fit had $\lambda_- = \lambda_+$ but would not change for $\lambda_- = 0$. L.Pondrom, (private communication 74).

Meson Full Listings

K^\pm

⁸¹ HAIDT 71 table 8 (Dalitz plot analysis) gives $d\xi(0)/d\lambda_+ = (-1.1+0.5)/(0.050-0.029) = -29$, error raised from 0.50 to agree with $d\xi(0) = 0.20$ for fixed λ_+ .

⁸² KIJEWski 69 figure 17 was used to obtain $d\xi(0)/d\lambda_+$ and errors.

⁸³ CALLAHAN 66 table 1 (π analysis) gives $d\xi(0)/d\lambda_+ = (0.72-0.05)/(0-0.04) = -17$, error raised from 0.80 to agree with $d\xi(0) = 0.37$ for fixed λ_+ . t unknown.

⁸⁴ JENSEN 64 gives $\lambda_+^\mu = \lambda_+^e = -0.020 \pm 0.027$. $d\xi(0)/d\lambda_+$ unknown. Includes SHAKLEE 64 $\xi_B(K_{\mu 3}/K_{e 3})$.

$\xi_B = f_-/f_+$ (determined from $K_{\mu 3}^\pm/K_{e 3}^\pm$)

The $K_{\mu 3}^\pm/K_{e 3}^\pm$ branching ratio fixes a relationship between $\xi(0)$ and λ_+ . We quote the author's $\xi(0)$ and associated λ_+ but do not average because the λ_+ values differ. The fit result and scale factor given below are not obtained from these ξ_B values. Instead they are obtained directly from the fitted $K_{\mu 3}^\pm/K_{e 3}^\pm$ ratio $\Gamma(\pi^0 \mu^+ \nu_\mu)/\Gamma(\pi^0 e^+ \nu_e)$, with the exception of HEINTZE 77. The parameter ξ is redundant with λ_0 below and is not put into the Meson Summary Table.

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-0.35±0.15 OUR EVALUATION		From a fit discussed in note on $K_{f 3}$ form factors in 1982 edition, PL 111B (April 1982).			
-0.12±0.12	55k	⁸⁵ HEINTZE	77	CNTR	$\lambda_+ = 0.029$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0 ± 0.15	5825	CHIANG	72	OSPK	$\lambda_+ = 0.03$, fig.10
-0.81±0.27	1505	⁸⁶ HAIDT	71	HLBC	$\lambda_+ = 0.028$, fig.8
-0.35±0.22		⁸⁷ BOTTERILL	70	OSPK	$\lambda_+ = 0.045 \pm 0.015$
+0.91±0.82		ZELLER	69	ASPK	$\lambda_+ = 0.023$
-0.08±0.15	5601	⁸⁷ BOTTERILL	68B	ASPK	$\lambda_+ = 0.023 \pm 0.008$
-0.60±0.20	1398	⁸⁶ EICHTEN	68	HLBC	See note
+1.0 ± 0.6	986	GARLAND	68	OSPK	$\lambda_+ = 0$
+0.75±0.50	306	AUERBACH	67	OSPK	$\lambda_+ = 0$
+0.4 ± 0.4	636	CALLAHAN	66B	FBC	$\lambda_+ = 0$
+0.6 ± 0.5		BISI	65B	HBC	$\lambda_+ = 0$
+0.8 ± 0.6	500	CUTTS	65	OSPK	$\lambda_+ = 0$
-0.17±0.75		SHAKLEE	64	XEBC	$\lambda_+ = 0$

⁸⁵ Calculated by us from λ_0 and λ_+ given below.

⁸⁶ EICHTEN 68 has $\lambda_+ = 0.023 \pm 0.008$, $t = 4$, independent of λ_- . Replaced by HAIDT 71.

⁸⁷ BOTTERILL 70 is re-evaluation of BOTTERILL 68B with different λ_+ .

$\xi_C = f_-/f_+$ (determined from μ polarization in $K_{\mu 3}^\pm$)

The μ polarization is a measure of $\xi(t)$. No assumptions on λ_+ necessary, t (weighted by sensitivity to $\xi(t)$) should be specified. In λ_+ , $\xi(0)$ parametrization this is $\xi(0)$ for $\lambda_+ = 0$. $d\xi/d\lambda = \xi t$. For radiative correction to muon polarization in $K_{\mu 3}^\pm$, see GINSBERG 71. The parameter ξ is redundant with λ_0 below and is not put into the Meson Summary Table.

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-0.35±0.15 OUR EVALUATION		From a fit discussed in note on $K_{f 3}$ form factors in 1982 edition, PL 111B (April 1982).			
-0.25±1.20	1585	⁸⁸ BRAUN	75	HLBC	POL, $t=4.2$
-0.95±0.3	3133	⁸⁹ CUTTS	69	OSPK	Total pol. $t=4.0$
-1.0 ± 0.3	6000	⁹⁰ BETTELS	68	HLBC	Total pol. $t=4.9$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
-0.64±0.27	40k	⁹¹ MERLAN	74	ASPK	POL, $d\xi(0)/d\lambda_+ = +1.7$
-1.4 ± 1.8	397	⁹² CALLAHAN	66B	FBC	Total pol.
-0.7 +0.9	2950	⁹² CALLAHAN	66B	FBC	Long. pol.
-3.3					
+1.2 +2.4	2100	⁹² BORREANI	65	HLBC	Polarization
-1.8					
-4.0 to +1.7	500	⁹² CUTTS	65	OSPK	Long. pol.

⁸⁸ BRAUN 75 $d\xi(0)/d\lambda_+ = \xi t = -0.25 \times 4.2 = -1.0$.

⁸⁹ CUTTS 69 $t = 4.0$ was calculated from figure 8. $d\xi(0)/d\lambda_+ = \xi t = -0.95 \times 4 = -3.8$.

⁹⁰ BETTELS 68 $d\xi(0)/d\lambda_+ = \xi t = -1.0 \times 4.9 = -4.9$.

⁹¹ MERLAN 74 polarization result (figure 5) not possible. See discussion of polarization experiments in note on "K_{f3} Form Factors" in the 1982 edition of this Review [Physics Letters 111B (1982)].

⁹² t value not given.

Im(ξ) in $K_{\mu 3}^\pm$ DECAY (from transverse μ pol.)

Test of T reversal invariance.

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
-0.017±0.025 OUR AVERAGE					
-0.016±0.025	20M	CAMPBELL	81	CNTR	Pol.
-0.3 +0.3	3133	CUTTS	69	OSPK	Total pol. fig.7
-0.4					
-0.1 ± 0.3	6000	BETTELS	68	HLBC	Total pol.
0.0 ± 1.0	2648	CALLAHAN	66B	FBC	MU
+1.6 ± 1.3	397	CALLAHAN	66B	FBC	Total pol.
0.5 +1.4	2950	CALLAHAN	66B	FBC	Long. pol.
-0.5					

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.010±0.019 32M ⁹³ BLATT 83 CNTR Polarization

⁹³ Combined result of MORSE 80 ($K_{\mu 3}^0$) and CAMPBELL 81 ($K_{\mu 3}^+$).

λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN $K_{\mu 3}^\pm$ DECAY)

See also the corresponding entries and footnotes in sections ξ_A , ξ_C , and λ_0 . For radiative correction of $K_{\mu 3}^\pm$ Dalitz plot, see GINSBERG 70 and BECHERRAWY 70.

VALUE	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
0.033±0.008 OUR EVALUATION		From a fit discussed in note on $K_{f 3}$ form factors in 1982 edition, PL 111B (April 1982).			
+0.050±0.013	3973	WHITMAN	80	SPEC	+ DP
0.025±0.030	490	ARNOLD	74	HLBC	+ DP
0.027±0.019	6527	MERLAN	74	ASPK	+ DP
0.025±0.017	1897	BRAUN	73C	HLBC	+ DP
0.024±0.019	4025	⁹⁴ ANKENBRA...	72	ASPK	+ PI
-0.006±0.015	3480	CHIANG	72	OSPK	+ DP
0.050±0.018	3240	HAIDT	71	HLBC	+ DP
0.009±0.026	2041	KIJEWski	69	OSPK	+ PI
0.0 ± 0.05	444	CALLAHAN	66B	FBC	+ PI

⁹⁴ ANKENBRANDT 72 λ_+ from figure 3 to match $d\xi(0)/d\lambda_+$. Text gives 0.024 ± 0.022 .

λ_0 (LINEAR ENERGY DEPENDENCE OF f_0 IN $K_{\mu 3}^\pm$ DECAY)

Wherever possible, we have converted the above values of $\xi(0)$ into values of λ_0 using the associated λ_+^μ and $d\xi/d\lambda_+$.

VALUE	$d\lambda_0/d\lambda_+$	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
0.004±0.007 OUR EVALUATION			From a fit discussed in note on $K_{f 3}$ form factors in 1982 edition, PL 111B (April 1982).				
+0.029±0.011	-0.37	3973	WHITMAN	80	SPEC	+ DP	
+0.019±0.010	+0.03	55k	⁹⁵ HEINTZE	77	SPEC	+ BR	
+0.008±0.097	+0.92	1585	⁹⁶ BRAUN	75	HLBC	+ POL	
-0.040±0.040	-0.62	490	ARNOLD	74	HLBC	+ DP	
-0.019±0.015	+0.27	6527	⁹⁷ MERLAN	74	ASPK	+ DP	
-0.008±0.020	-0.53	1897	⁹⁸ BRAUN	73C	HLBC	+ DP	
-0.026±0.013	+0.03	4025	⁹⁹ ANKENBRA...	72	ASPK	+ PI	
+0.030±0.014	-0.21	3480	⁹⁹ CHIANG	72	OSPK	+ DP	
-0.039±0.029	-1.34	3240	⁹⁹ HAIDT	71	HLBC	+ DP	
-0.056±0.024	+0.69	3133	⁹⁶ CUTTS	69	OSPK	+ POL	
-0.031±0.045	-1.10	2041	⁹⁹ KIJEWski	69	OSPK	+ PI	
-0.063±0.024	+0.60	6000	⁹⁶ BETTELS	68	HLBC	+ POL	
+0.058±0.036	-0.37	444	⁹⁹ CALLAHAN	66B	FBC	+ PI	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
-0.017±0.011			100	BRAUN	74	HLBC	+ $K_{\mu 3}^\pm/K_{e 3}^\pm$ vs. t

⁹⁵ HEINTZE 77 uses $\lambda_+ = 0.029 \pm 0.003$. $d\lambda_0/d\lambda_+$ estimated by us.

⁹⁶ λ_0 value is for $\lambda_+ = 0.03$ calculated by us from $\xi(0)$ and $d\xi(0)/d\lambda_+$.

⁹⁷ MERLAN 74 λ_0 and $d\lambda_0/d\lambda_+$ were calculated by us from ξ_A , λ_+^μ , and $d\xi(0)/d\lambda_+$. Their figure 6 gives $\lambda_0 = -0.025 \pm 0.012$ and no $d\lambda_0/d\lambda_+$.

⁹⁸ This value and error are taken from BRAUN 75 but correspond to the BRAUN 73C λ_+^μ result. $d\lambda_0/d\lambda_+$ is from BRAUN 73C $d\xi(0)/d\lambda_+$ in ξ_A above.

⁹⁹ λ_0 calculated by us from $\xi(0)$, λ_+^μ , and $d\xi(0)/d\lambda_+$.

100 BRAUN 74 is a combined $K_{\mu 3}^\pm$ - $K_{e 3}^\pm$ result. It is not independent of BRAUN 73C ($K_{\mu 3}^\pm$) and BRAUN 73B ($K_{e 3}^\pm$) form factor results.

$|f_S/f_+|$ FOR $K_{e 3}^\pm$ DECAY

Ratio of scalar to f_+ couplings.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
0.084±0.023 OUR AVERAGE			Error includes scale factor of 1.2.				
0.070±0.016±0.016		32k	AKIMENKO	91	SPEC	λ_+ , f_S , f_T , ϕ fit	
0.00 ± 0.10		2827	BRAUN	75	HLBC	+	
0.14 +0.03		2707	STEINER	71	HLBC	λ_+ , f_S , f_T , ϕ fit	
-0.04							
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<0.13		90	4017	CHIANG	72	OSPK	+
<0.23		90		BOTTERILL	68C	ASPK	+
<0.18		90		BELLOTTI	67B	HLBC	+
<0.30		95		KALMUS	67	HLBC	+

$|f_T/f_+|$ FOR $K_{e 3}^\pm$ DECAY

Ratio of tensor to f_+ couplings.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	CHG	COMMENT	
0.38±0.11 OUR AVERAGE			Error includes scale factor of 1.1.				
0.53 +0.09		32k	AKIMENKO	91	SPEC	λ_+ , f_S , f_T , ϕ fit	
-0.10 ± 0.10							
0.07 ± 0.37		2827	BRAUN	75	HLBC	+	
0.24 +0.16		2707	STEINER	71	HLBC	λ_+ , f_S , f_T , ϕ fit	
-0.14							
• • • We do not use the following data for averages, fits, limits, etc. • • •							
<0.75		90	4017	CHIANG	72	OSPK	+
<0.58		90		BOTTERILL	68C	ASPK	+
<0.58		90		BELLOTTI	67B	HLBC	+
<1.1		95		KALMUS	67	HLBC	+

f_T/f_+ FOR $K_{\mu 3}^\pm$ DECAY

Ratio of tensor to f_+ couplings.

VALUE	EVTs	DOCUMENT ID	TECN
0.02±0.12	1585	BRAUN	75
			HLBC

DECAY FORM FACTORS FOR $K^{\pm} \rightarrow \pi^{\pm} \pi^{-} e^{\pm} \nu_e$

Given in ROSSELET 77, BEIER 73, and BASILE 71c.

DECAY FORM FACTOR FOR $K^{\pm} \rightarrow \pi^0 \pi^0 e^{\pm} \nu_e$

Given in BOLOTOV 86b and BARMIN 88b.

 $K^{\pm} \rightarrow \ell^{\pm} \nu \gamma$ FORM FACTORS

For definitions of the axial-vector F_A and vector F_V form factor, see the "Note on $\pi^{\pm} \rightarrow \ell^{\pm} \nu \gamma$ and $K^{\pm} \rightarrow \ell^{\pm} \nu \gamma$ Form Factors" in the π^{\pm} section. In the kaon literature, often different definitions $a_K = F_A/m_K$ and $v_K = F_V/m_K$ are used.

 $F_A + F_V$, SUM OF AXIAL-VECTOR AND VECTOR FORM FACTOR FOR $K \rightarrow e \nu_e \gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.146 ± 0.010 OUR AVERAGE				
0.147 ± 0.011	51	101 HEINTZE	79 SPEC	$K \rightarrow e \nu \gamma$
0.150 ^{+0.018} _{-0.023}	56	102 HEARD	75 SPEC	$K \rightarrow e \nu \gamma$

101 HEINTZE 79 quotes absolute value of $|F_A + F_V| \sin \theta_c$. We use $\sin \theta_c = V_{us} = 0.2205$.
102 HEARD 75 quotes absolute value of $|F_A + F_V| \sin \theta_c$. We use $\sin \theta_c = V_{us} = 0.2205$.

 $F_A + F_V$, SUM OF AXIAL-VECTOR AND VECTOR FORM FACTOR FOR $K \rightarrow \mu \nu \mu \gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.23				
	90	103 AKIBA	85 SPEC	$K \rightarrow \mu \nu \gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-1.2 to 1.1	90	DEMIDOV	90 XEBC	$K \rightarrow \mu \nu \gamma$

103 AKIBA 85 quotes absolute value.

 $F_A - F_V$, DIFFERENCE OF AXIAL-VECTOR AND VECTOR FORM FACTOR FOR $K \rightarrow e \nu_e \gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.49				
	90	104 HEINTZE	79 SPEC	$K \rightarrow e \nu \gamma$

104 HEINTZE 79 quotes $|F_A - F_V| < \sqrt{1} |F_A + F_V|$.

 $F_A - F_V$, DIFFERENCE OF AXIAL-VECTOR AND VECTOR FORM FACTOR FOR $K \rightarrow \mu \nu \mu \gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
-2.2 to 0.3 OUR EVALUATION				
-2.2 to 0.6	90	DEMIDOV	90 XEBC	$K \rightarrow \mu \nu \gamma$
-2.5 to 0.3	90	AKIBA	85 SPEC	$K \rightarrow \mu \nu \gamma$

K[±] REFERENCES

ATIYA	93	PRL 70 2521	+Chiang, Frank, Haggerty, Ito+	(BNL 787 Collab.)
Also	93C	PRL 71 305 (erratum)	+Abe, Chiang, Frank, Haggerty, Ito+	(BNL 787 Collab.)
ATIYA	93B	PR D48 R1	+Chiang, Frank, Haggerty, Ito+	(BNL 787 Collab.)
ALLIEGRO	92	PRL 68 278	+Campagnari+ (BNL, FNAL, PSI, WASH, YALE)	
BARMIN	92	SJNP 55 547	+Barylov, Cherkukha, Davidenko+	(ITEP)
		Translated from YAF 55 976.		
IMAZATO	92	PRL 69 877	+Kawahisa, Tanaka+	(KEK, INUS, TOKY, TOKMS)
IVANOV	92	PRL 68 443	+Shrock	(BNL, STON)
LITTENBERG	92	PR D45 3961	+Fero, Gee, Graf, Mandelkern, Schultz, Shultz	(UCI)
USHER	91	PL B259 225	+Belousov+ (SERP, JINR, TBIL, CMNS, SOFU, KOSI)	
AKIMENKO	91	SJNP 53 606	+Barylov, Davidenko, Demidov+	(ITEP)
BARMIN	91	SJNP 53 606	+Barylov, Davidenko, Demidov+	(ITEP)
		Translated from YAF 53 981.		
DENISOV	91	JETPL 54 558	+Zhelamkov, Ivanov, Lapina, Levchenko, Malakhov+ (PNPI)	
Also		Translated from ZETFP 54 557.		
ATIYA	90	PRL 64 211	+Chiang, Frank, Haggerty, Ito, Kycia+	(BNL 787 Collab.)
ATIYA	90B	PRL 65 1188	+Chiang, Frank, Haggerty, Ito, Kycia+	(BNL 787 Collab.)
DEMIDOV	90	SJNP 52 1006	+Dobrokhotoy, Lyublev, Nikitenko+	(ITEP)
		Translated from YAF 52 1595.		
LEE	90	PRL 64 165	+Alliegro, Campagnari+ (BNL, FNAL, VILL, WASH, YALE)	
ATIYA	89	PRL 63 2177	+Chiang, Frank, Haggerty, Ito, Kycia+	(BNL 787 Collab.)
BARMIN	89	SJNP 50 421	+Barylov, Davidenko, Demidov, Dolgolenko+	(ITEP)
		Translated from YAF 50 679.		
BARMIN	88	SJNP 47 643	+Barylov, Davidenko, Demidov, Dolgolenko+	(ITEP)
		Translated from YAF 47 1011.		
BARMIN	88B	SJNP 48 1032	+Barylov, Davidenko, Demidov, Dolgolenko+	(ITEP)
		Translated from YAF 48 1719.		
BOLOTOV	88	JETPL 47 7	+Gninenko, Dzhlkibaev, Isakov, Klubakov+	(ASCI)
		Translated from ZETFP 47 8.		
CAMPAGNARI	88	PRL 61 2062	+Alliegro, Chaloupka+ (BNL, FNAL, PSI, WASH, YALE)	
GALL	88	PRL 60 186	+Austin+ (BOST, MIT, WILL, CIT, CMU, WYOM)	
BARMIN	87	SJNP 45 62	+Barylov, Davidenko, Demidov+	(ITEP)
		Translated from YAF 45 97.		
BOLOTOV	87	SJNP 45 1023	+Gninenko, Dzhlkibaev, Isakov, Klubakov+	(INRM)
		Translated from YAF 45 1652.		
BOLOTOV	86	SJNP 44 73	+Gninenko, Dzhlkibaev, Isakov+	(INRM)
		Translated from YAF 44 117.		
BOLOTOV	86B	SJNP 44 68	+Gninenko, Dzhlkibaev, Isakov+	(INRM)
		Translated from YAF 44 108.		
YAMANAKA	86	PR D34 85	+Hayano, Taniguchi, Ishikawa+	(KEK, TOKY)
Also	84	PRL 52 329	+Hayano, Yamanaka, Taniguchi+	(TOKY, KEK)
AKIBA	85	PR D32 2911	+Ishikawa, Iwasaki+ (TOKY, TINT, TSUK, KEK)	
BOLOTOV	85	JETPL 42 481	+Gninenko, Dzhlkibaev, Isakov+	(INRM)
		Translated from ZETFP 42 390.		
BLATT	83	PR D27 1056	+Adair, Black, Campbell+	(YALE, BNL)
ASANO	82	PL 113B 195	+Kikutani, Kurokawa, Miyachi+(KEK, TOKY, INUS, OSAK)	
COOPER	82	PL 112B 97	+Guy, Michette, Tyndel, Venus	(RL)
PDG	82	PL 111B	+Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
PDG	82B	PL 111B 70	+Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
ASANO	81B	PL 107B 159	+Kikutani, Kurokawa, Miyachi+(KEK, TOKY, INUS, OSAK)	
CAMPBELL	81	PRL 47 1032	+Black, Blatt, Kasha, Schmidt+	(YALE, BNL)
Also	83	PR D27 1056	+Blatt, Adair, Black, Campbell+	(YALE, BNL)
LUM	81	PR D23 2522	+Wiegand, Kessler, Deslattes, Seki+	(LBL, NBS+)

LYONS	81	ZPHY C10 215		
MORSE	80	PR D21 1750		
WHITMAN	80	PR D21 1750		
BARKOV	79	NP B148 53		
HEINTZE	79	NP B149 365		
ABRAMS	77	PR D15 22		
DEVAUX	77	NP B126 11		
HEINTZE	77	PL 70B 482		
ROSSELET	77	PR D15 574		
BERTRAND	76	NP B114 387		
BLOCH	76	PL 60B 393		
BRAUN	76B	LNC 17 521		
DIAMANT....	76	PL 62B 485		
HEINTZE	76	PL 60B 302		
SMITH	76	NP B109 173		
WEISSENBE...	76	NP B115 55		
BLOCH	75	PL 56B 201		
BRAUN	75	NP B89 210		
CHENG	75	NP A254 391		
HEARD	75	PL 55B 324		
HEARD	75B	PL 55B 327		
SHEAFF	75	PR D12 2570		
SMITH	75	NP B91 45		
ARNOLD	74	PR D9 1221		
BRAUN	74	PL 51B 393		
CENCE	74	PR D10 776		
Also		This is unpub.		
KUNSELMAN	74	PR C9 2469		
MERLAN	74	PR D9 107		
WEISSENBE...	74	PL 48B 474		
ABRAMS	73B	PRL 30 500		
BACKENSTO...	73	PL 43B 431		
BEIER	73	PRL 30 399		
BRAUN	73B	PL 47B 185		
Also		NP B89 210		
BRAUN	73C	PL 47B 182		
Also		NP B89 210		
CABLE	73	PR D8 3807		
LJUNG	73	PR D8 1307		
Also		PRL 28 523		
Also		PRL 28 1287		
Also		PR 23 326		
LUCAS	73	PR D8 719		
LUCAS	73B	PR D8 727		
PANG	73	PR D8 1989		
Also		PL 40B 699		
SMITH	73	NP B60 411		
ABRAMS	72	PRL 29 1118		
ANKENBRAND...	72	PRL 28 1472		
AUBERT	72	NC 12A 509		
BEIER	72	PRL 29 678		
CHIANG	72	PR D6 1254		
CLARK	72	PRL 29 1274		
EDWARDS	72	PR D5 2720		
FORD	72	PL 38B 335		
HOFFMASTER	72	NP B36 1		
BASILE	71C	PL 36B 619		
BOURQUIN	71	PL 36B 615		
GINSBERG	71	PR D4 2993		
HAIDT	71	PR D3 10		
Also		NP 29B 691		
KLEMS	71	PR D4 66		
Also		PRL 24 1086		
Also		PRL 25 473		
OTT	71	PR D3 52		
ROMANO	71	PL 36B 246		
SCHWEINB...	71	PL 36B 521		
BARDIN	70	PL 32B 121		
BECHERRAWY	70	PR D1 1452		
BOTTERILL	70	PL 31B 325		
FORD	70	PRL 25 1370		
GALLARD	70	PRL 25 1472		
GINSBERG	70	PR D1 229		
GRAUMAN	70	PR D1 1277		
Also		PRL 23 737		
MALTSEV	70	SJNP 10 678		
		Translated from YAF 10 1195.		
PANDOULAS	70	PR D2 1205		
CUTTS	68	PRL 20 955		
DAVISON	69	PR 180 1333		
ELY	69	PR 180 1319		
EMMERSON	69	PR 183 2393		
HERZO	69	PR 186 1403		
KJEWSKI	69	UCRL 18433 Thesis		
LOBKOWICZ	69	PR 185 1676		
MACKEE	69	PR 17 548		
MASST	69	PR 183 1200		
SELLERI	69	NC 60A 291		
ZELLER	69	PR 182 1420		
BETTELS	68	NC 56A 1106		
Also		PR D3 10		
BOTTERILL	68B	PRL 21 766		
BOTTERILL	68C	PR 174 1661		
BUTLER	68	UCRL 18420		
CHANG	68	PRL 20 510		
CHEN	68	PRL 20 73		
EICHTEN	68	PL 27B 586		
EISLER	68	PR 169 1090		
ESCHSTRUTH	68	PR 165 1487		
GARLAND	68	PR 167 1225		
MOSCOSO	68	Thesis		
AUERBACH	67	PR 155 1505		
Also		PR D9 3216		
Erratum.				
BELLOTTI	67	Heidelberg Conf.		
BELLOTTI	67B	NC 52A 1287		
Also		PR 150 690		
BIS	67	PL 25B 572		
BOTTERILL	67	PR 159 982		
Also		PR 171 1402		
BOWEN	67B	PR 154 1314		
CLINE	67B	Herceg Novi Tbl. 4		
Proc. International School on Elementary Particle Physics.				
FLETCHER	67	PR 159 98		
FORD	67	PRL 18 1214		
GINSBERG	67	PR 162 1570		
IMLAY	67	PR 160 1203		

+Albajar, Myatt				(OXF)
+Lepuner, Larsen, Schmidt, Blatt+				(BNL, YALE)
+Abrams, Carroll, Kycia, Li+				(ILL, BNL, LBL)
+Vasserman, Zolotarev, Krupin+				(NOVO, KIAE)
+Heinzelmann, Igo-Kemenes+				(HEIDP, CERN)
+Carroll, Kycia, Li, Michael, Mockett+				(BNL)
+Bloch, Diamant-Berger, Maillard+				(SACL, GEVA)
+Heinzelmann, Igo-Kemenes+				(HEIDP, CERN)
+Extermann, Fischer, Giusan+				(GEVA, SACL)
+Sacton+				(BRUX, KIDR, DUUC, LOUC, WARS)
+Bunce, Devaux, Diamant-Berger+				(GEVA, SACL)
+Marty, Enriquez+				(AACH3, BARI, BELG, CERN)
Diamant-Berger, Bloch, Devaux+				(SACL, GEVA)
+Heinzelmann, Igo-Kemenes, Mundhenke+				(HEIDP)
+Booth, Renshall, Jones+				(GLAS, LIVP, OXF, RHEL)
Weissenberg, Egorov, Minervina+				(ITEP, LEBD)
+Brehin, Bunce, Devaux+				(SACL, GEVA)
+Cornelsen+				(AACH3, BARI, BRUX, CERN)
+Asano, Chen, Dugan, Hu, Wu+				(COLU, YALE)
+Heintze, Heinzelmann+				(CERN, HEIDH)
+Heintze, Heinzelmann+				(CERN, HEIDH)
				(WISC)
+Booth, Renshall, Jones+				(GLAS, LIVP, OXF, RHEL)
+Roe, Sinclair				(MICH)
+Cornelsen, Marty+				(AACH3, BARI, BRUX, CERN)
+Harris, Jones, Morgado+				(

Meson Full Listings

K^\pm, K^0, K_S^0

KALMUS 67	PR 159 1187	+Kernan	(LRL)
ZINCHENKO 67	Rutgers Thesis		(RUTG)
CALLAHAN 66	NC 44 90		(WISC)
CALLAHAN 66B	PR 150 1153	+Camerini+	(WISC, LRL, UCR, BARI)
CESTER 66	PL 21 343	+Eschstruth, Oneill+	(PPA)
See footnote 1 in AUERBACH 67.			
Also	67 PR 155 1505	Auerbach, Dobbs, Mann+	(PENN, PRIN)
BIRGE 65	PR 139B 1600	+Ely, Gidal, Camerini, Cline+	(LRL, WISC)
BISI 65	NC 35 768	+Borreani, Cester, Ferraro+	(TOR)
BISI 65B	PR 139B 1068	+Borreani, Marzari-Chiesa, Rinaudo+	(TOR)
BORREANI 65	PR 140B 1686	+Gidal, Rinaudo, Caforio+	(BARI, TOR)
CALLAHAN 65	PRL 15 129	+Cline	(WISC)
CAMERINI 65	NC 37 1795	+Cline, Gidal, Kalmus, Kernan	(WISC, LRL)
CLINE 65	PL 15 293	+Fry	(WISC)
CUTTS 65	PR 138B 969	+Elioff, Stiening	(LRL)
DEMARCO 65	PR 140B 1430	+Grosso, Rinaudo	(TOR, CERN)
FITCH 65B	PR 140B 1088	+Quares, Wilkins	(PRIN, MTHO)
GREINER 65	ARNS 15 67		(LRL)
STAMER 65	PR 138B 440	+Huetter, Koller, Taylor, Grauman	(STEV)
TRILLING 65B	UCRL 16473		(LRL)
Updated from 1965 Argonne Conference, page 5.			
YOUNG 65	UCRL 16362 Thesis		(LRL)
Also	67 PR 156 1464	Young, Osborne, Barkas	(TOR)
BORREANI 64	PL 12 123	+Rinaudo, Werbrouck	(TOR)
CALLAHAN 64	PR 136B 1463	+March, Stark	(WISC)
CAMERINI 64	PRL 13 318	+Cline, Fry, Powell	(WISC, LRL)
CLINE 64	PRL 13 101	+Fry	(WISC)
GIACOMELLI 64	NC 34 1134	+Monti, Quareni+	(BGNA, MUNI)
GREINER 64	PRL 13 284	+Osborne, Barkas	(LRL)
JENSEN 64	PR 136B 1431	+Shaklee, Roe, Sinclair	(MICH)
KALMUS 64	PRL 13 99	+Kernan, Pu, Powell, Dowd	(LRL, WISC)
SHAKLEE 64	PR 136B 1423	+Jensen, Roe, Sinclair	(MICH)
BARKAS 63	PRL 11 76	+Dyer, Heckman	(LRL)
BOYARSKI 62	PR 129 2398	+Loth, Niemela, Ritson	(MIT)
BROWN 62B	PRL 8 450	+Kadyk, Trilling, Roe+	(LRL, MICH)
BARKAS 61	PR 124 1209	+Dyer, Mason, Norris, Nickols, Smit	(LRL)
BHOWMIK 61	NC 20 857	+Jain, Mathur	(DELH)
FERRO-LUZZI 61	NC 22 1087	+Miller, Murray, Rosenfeld+	(LRL)
NORDIN 61	PR 123 2166		(LRL)
ROE 61	PRL 7 346	+Sinclair, Brown, Glaser+	(MICH, LRL)
FREDEN 60B	PR 118 564	+Gilbert, White	(LRL)
BURROWES 59	PRL 2 117	+Caldwell, Frisch, Hill+	(MIT)
TAYLOR 59	PR 114 359	+Harris, Orear, Lee, Baumei	(COLU)
EISENBERG 58	NC 8 663	+Koch, Lohrmann, Nikolic+	(BERN)
ALEXANDER 57	NC 6 478	+Johnston, Oceaigh	(DUUC)
COHEN 57	Fund. Cons. Phys.	+Crowe, Dumond	(NAAS, LRL, CIT)
COOMBES 57	PR 108 1348	+Cork, Galbraith, Lambertson, Wenzel	(LRL)
BIRGE 56	NC 4 834	+Perkins, Peterson, Stork, Whitehead	(LRL)
ILOFF 56	PR 102 927	+Goldhaber, Lannutti, Gilbert+	(LRL)

OTHER RELATED PAPERS

LITTENBERG 93	ARNPS 43 729	+Valencia	(BNL, FNAL)
Rare and Radiative Kaon Decays			
RITCHIE 93	RMP 65 1149	+Wojcicki	
"Rare K Decays"			
BATTISTON 92	PRPL 214 293	+Cocolicchio, Fogli, Paver	(PGIA, CERN, TRSTT)
Status and Perspectives of K Decay Physics			
BRYMAN 89	IJMP A4 79		(TRIU)
"Rare Kaon Decays"			
CHOUNET 72	PR 140B 1199	+Gailard, Gailard	(ORSAY, CERN)
FEARING 70	PR D2 542	+Fischbach, Smith	(STON, BOHR)
HAIDT 69B	PL 29B 696	+ (AACH, BARI, CERN, EPOL, NIJM, ORSAY+)	(PRIN)
CRONIN 68B	Vienna Conf. 241		
Rapporteur talk.			
WILLIS 67	Heidelberg Conf. 273		(YALE)
Rapporteur talk.			
CABIBBO 66	Berkeley Conf. 33		(CERN)
ADAIR 64	PL 12 67	+Leipuner	(YALE, BNL)
CABIBBO 64	PL 9 352	+Maksymowicz	(CERN)
Also	64B PL 11 360	Cabibbo, Maksymowicz	(CERN)
Also	65 PL 14 72	Cabibbo, Maksymowicz	(CERN)
BIRGE 63	PRL 11 35	+Ely, Gidal, Camerini+	(LRL, WISC, BARI)
BLOCK 62B	CERN Conf. 371	+Lendinara, Monari	(NWES, BGNA)
BRENE 61	NP 22 553	+Egardt, Qvist	(NORD)

K^0

$$I(J^P) = \frac{1}{2}(0^-)$$

K^0 MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
497.672 ± 0.031 OUR FIT				
497.672 ± 0.031 OUR AVERAGE				
497.661 ± 0.033	3713	BARKOV	87B CMD	$e^+e^- \rightarrow K_L^0 K_S^0$
497.742 ± 0.085	780	BARKOV	85B CMD	$e^+e^- \rightarrow K_L^0 K_S^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
497.44 ± 0.50		FITCH	67 OSPK	
498.9 ± 0.5	4500	BALTAY	66 HBC	K^0 from $\bar{p}p$
497.44 ± 0.33	2223	KIM	65B HBC	K^0 from $\bar{p}p$
498.1 ± 0.4		CHRISTENS...	64 OSPK	

$m_{K^0} - m_{K^\pm}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
3.995 ± 0.034 OUR FIT	Error includes scale factor of 1.1.				
• • • We do not use the following data for averages, fits, limits, etc. • • •					
3.95 ± 0.21	417	HILL	68B DBC	+	$K^+d \rightarrow K^0 pp$
3.90 ± 0.25	9	BURNSTEIN	65 HBC	-	
3.71 ± 0.35	7	KIM	65B HBC	-	$K^-p \rightarrow n\bar{K}^0$
5.4 ± 1.1		CRAWFORD	59 HBC	+	
3.9 ± 0.6		ROSENFELD	59 HBC	-	

$$|m_{K^0} - m_{K^\pm}| / m_{\text{average}}$$

A test of CPT invariance.

VALUE DOCUMENT ID
 $< 9 \times 10^{-19}$ OUR EVALUATION

K^0 REFERENCES

BARKOV 87B	SJNP 46 630	+Vasserman, Vorobev, Ivanov+	(NOVO)
BARKOV 85B	JETPL 42 138	+Blinov, Vasserman+	(NOVO)
Translated from YAF 46 108B.			
HILL 68B	PR 168 1534	+Robinson, Sakitt, Canter	(BNL, CMU)
FITCH 67	PR 164 1711	+Roth, Russ, Vernon	(PRIN)
BALTAY 66	PR 142 932	+Sandweiss, Stonehill+	(YALE, BNL)
BURNSTEIN 65	PR 138B 995	+Rubin	(UMD)
KIM 65B	PR 140B 1334	+Kirsch, Miller	(COLU)
CHRISTENS... 64	PRL 13 138	Christenson, Cronin, Fitch, Turlay	(PRIN)
CRAWFORD 59	PRL 2 112	+Cresti, Good, Stevenson, Ticho	(LRL)
ROSENFELD 59	PRL 2 110	+Solmitz, Tripp	(LRL)

K_S^0

$$I(J^P) = \frac{1}{2}(0^-)$$

K_S^0 MEAN LIFE

For earlier measurements, beginning with BOLDT 58B, see our our 1986 edition, Physics Letters **170B** 130 (1986).

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN
0.8926 ± 0.0012 OUR AVERAGE			
0.8929 ± 0.0016		GIBBONS 93	E731
0.8920 ± 0.0044	214k	GROSSMAN 87	SPEC
0.881 ± 0.009	26k	ARONSON 76	SPEC
0.8913 ± 0.0032		1 CARITHERS 75	SPEC
0.8937 ± 0.0048	6M	GEWENIGER 74B	ASPK
0.8958 ± 0.0045	50k	2 SKJEGGEST... 72	HBC
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.905 ± 0.007		3 ARONSON 82B	SPEC
0.867 ± 0.024	2173	4 FACKLER 73	OSPK
0.856 ± 0.008	19994	5 DONALD 68B	HBC
0.872 ± 0.009	20000	5,6 HILL 68	DBC
0.866 ± 0.016		5 ALFF... 66B	OSPK
0.843 ± 0.013	5000	5 KIRSCH 66	HBC

1 CARITHERS 75 value is for $m_{K_L^0} - m_{K_S^0}$ $\Delta(m) = 0.5348 \pm 0.0021$. The $\Delta(m)$

dependence of the total decay rate (inverse mean life) is $\Gamma(K_S^0) = [(1.122 \pm 0.004) + 0.16(\Delta(m) - 0.5348)/\Delta(m)]10^{10}/s$. Value would not change with our current $\Delta(m) = 0.5349 \pm 0.0022$.

2 HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-} . SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

3 ARONSON 82 find that K_S^0 mean life may depend on the kaon energy.

4 FACKLER 73 does not include systematic errors.

5 Pre-1971 experiments are excluded from the average because of disagreement with later more precise experiments.

6 HILL 68 has been changed by the authors from the published value (0.865 ± 0.009) because of a correction in the shift due to η_{+-} . SKJEGGESTAD 72 and HILL 68 give detailed discussions of systematics encountered in this type of experiment.

K_S^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
$\Gamma_1 \pi^+\pi^-$	(68.61 ± 0.28) %	S=1.2
$\Gamma_2 \pi^0\pi^0$	(31.39 ± 0.28) %	S=1.2
$\Gamma_3 \pi^+\pi^-\gamma$	[a,b] (1.78 ± 0.05) × 10 ⁻³	
$\Gamma_4 \gamma\gamma$	(2.4 ± 1.2) × 10 ⁻⁶	
$\Gamma_5 \pi^+\pi^-\pi^0$	< 8.5 × 10 ⁻⁵	CL=90%
$\Gamma_6 3\pi^0$	< 3.7 × 10 ⁻⁵	CL=90%
$\Gamma_7 \pi^\pm e^\mp \nu$	[c] (6.68 ± 0.10) × 10 ⁻⁴	S=1.3
$\Gamma_8 \pi^\pm \mu^\mp \nu$	[c] (4.66 ± 0.07) × 10 ⁻⁴	S=1.2
$\Delta S = 1$ weak neutral current (S_1) modes		
$\Gamma_9 \mu^+\mu^-$	S_1 < 3.2 × 10 ⁻⁷	CL=90%
$\Gamma_{10} e^+e^-$	S_1 < 1.0 × 10 ⁻⁵	CL=90%
$\Gamma_{11} \pi^0 e^+e^-$	S_1 < 1.1 × 10 ⁻⁶	CL=90%

[a] See the Full Listings below for the energy limits used in this measurement.

[b] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ 's.

[c] Calculated from K_L^0 semileptonic rates and the K_S^0 lifetime assuming $\Delta S = \Delta Q$.

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 17 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 16.5$ for 16 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100
x_1	

K_S^0 DECAY RATES

$\Gamma(\pi^\pm e^\mp \nu)$ Γ_7
 VALUE (units 10^6 s^{-1}) DOCUMENT ID TECN COMMENT

7.48 ± 0.11 OUR EVALUATION Error includes scale factor of 1.3. From K_L^0 measurements, assuming that $\Delta S = \Delta Q$ in K^0 decay so that $\Gamma(K_S^0 \rightarrow \pi^\pm e^\mp \nu) = \Gamma(K_L^0 \rightarrow \pi^\pm e^\mp \nu)$.

• • • We do not use the following data for averages, fits, limits, etc. • • •
 seen BURGUN 72 HBC $K^+ \rho \rightarrow K^0 \rho \pi^+$
 9.3 ± 2.5 AUBERT 65 HLBC $\Delta S = \Delta Q$, CP cons. not assumed

$\Gamma(\pi^\pm \mu^\mp \nu)$ Γ_8
 VALUE (units 10^6 s^{-1}) DOCUMENT ID

5.22 ± 0.08 OUR EVALUATION Error includes scale factor of 1.2. From K_L^0 measurements, assuming that $\Delta S = \Delta Q$ in K^0 decay so that $\Gamma(K_S^0 \rightarrow \pi^\pm \mu^\mp \nu) = \Gamma(K_L^0 \rightarrow \pi^\pm \mu^\mp \nu)$.

K_S^0 BRANCHING RATIOS

$\Gamma(\pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_1 / Γ
 VALUE EVTS DOCUMENT ID TECN COMMENT

0.661 ± 0.0028 OUR FIT Error includes scale factor of 1.2.

0.671 ± 0.010 OUR AVERAGE
 0.670 ± 0.010 3447 7 DOYLE 69 HBC $\pi^- \rho \rightarrow A K^0$
 0.70 ± 0.08 COLUMBIA 608 HBC
 0.68 ± 0.04 CRAWFORD 598 HBC

• • • We do not use the following data for averages, fits, limits, etc. • • •
 0.740 ± 0.024 7 ANDERSON 628 HBC
 7 Anderson result not published, events added to Doyle sample.

$\Gamma(\pi^+ \pi^-) / \Gamma(\pi^0 \pi^0)$ Γ_1 / Γ_2
 VALUE EVTS DOCUMENT ID TECN COMMENT

2.186 ± 0.028 OUR FIT Error includes scale factor of 1.2.

2.197 ± 0.026 OUR AVERAGE
 2.11 ± 0.09 1315 EVERHART 76 WIRE $\pi^- \rho \rightarrow A K^0$
 2.169 ± 0.094 16k COWELL 74 OSPK $\pi^- \rho \rightarrow A K^0$
 2.16 ± 0.08 4799 HILL 73 DBC $K^+ d \rightarrow K^0 \rho \rho$
 2.22 ± 0.10 3068 8 ALITTI 72 HBC $K^+ \rho \rightarrow \pi^+ \rho K^0$
 2.22 ± 0.08 6380 MORSE 728 DBC $K^+ n \rightarrow K^0 \rho$
 2.10 ± 0.11 701 9 NAGY 72 HLBC $K^+ n \rightarrow K^0 \rho$
 2.22 ± 0.095 6150 10 BALTAY 71 HBC $K \rho \rightarrow K^0 \text{ neutrals}$
 2.282 ± 0.043 7944 11 MOFFETT 70 OSPK $K^+ n \rightarrow K^0 \rho$
 2.10 ± 0.06 3700 MORFIN 69 HLBC $K^+ n \rightarrow K^0 \rho$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 2.12 ± 0.17 267 9 BOZOKI 69 HLBC
 2.285 ± 0.055 3016 11 GOBBI 69 OSPK $K^+ n \rightarrow K^0 \rho$


8 The directly measured quantity is $K_S^0 \rightarrow \pi^+ \pi^- / \text{all } K^0 = 0.345 \pm 0.005$.
 9 NAGY 72 is a final result which includes BOZOKI 69.
 10 The directly measured quantity is $K_S^0 \rightarrow \pi^+ \pi^- / \text{all } \bar{K}^0 = 0.345 \pm 0.005$.
 11 MOFFETT 70 is a final result which includes GOBBI 69.

$\Gamma(\pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_2 / Γ
 VALUE EVTS DOCUMENT ID TECN

0.3139 ± 0.0028 OUR FIT Error includes scale factor of 1.2.

0.316 ± 0.014 OUR AVERAGE Error includes scale factor of 1.3. See the ideogram below.
 0.335 ± 0.014 1066 BROWN 63 HLBC
 0.288 ± 0.021 198 CHRETIEN 63 HLBC
 0.30 ± 0.035 BROWN 61 HLBC
 0.26 ± 0.06 BAGLIN 60 HLBC
 0.27 ± 0.11 CRAWFORD 598 HBC

WEIGHTED AVERAGE
 0.316 ± 0.014 (Error scaled by 1.3)



$\Gamma(\pi^+ \pi^- \gamma) / \Gamma(\pi^+ \pi^-)$ Γ_3 / Γ_1

VALUE (units 10^{-3}) EVTS DOCUMENT ID TECN COMMENT

2.60 ± 0.08 OUR AVERAGE
 2.56 ± 0.09 1286 RAMBERG 93 E731 $p_\gamma > 50 \text{ MeV}/c$
 2.68 ± 0.15 12 TAUREG 76 SPEC $p_\gamma > 50 \text{ MeV}/c$
 2.8 ± 0.6 13 BURGUN 73 HBC $p_\gamma > 50 \text{ MeV}/c$
 3.3 ± 1.2 10 WEBBER 70 HBC $p_\gamma > 50 \text{ MeV}/c$
 no ratio given 27 BELLOTTI 66 HBC $p_\gamma > 50 \text{ MeV}/c$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 7.10 ± 0.22 3723 RAMBERG 93 E731 $p_\gamma > 20 \text{ MeV}/c$
 3.0 ± 0.6 29 14 BOBISUT 74 HLBC $p_\gamma > 40 \text{ MeV}/c$

12 TAUREG 76 find direct emission contribution < 0.06, CL = 90%.
 13 BURGUN 73 estimates that direct emission contribution is 0.3 ± 0.6 .
 14 BOBISUT 74 not included in average because p_γ cut differs. Estimates direct emission contribution to be 0.5 or less, CL = 95%.

$\Gamma(\gamma \gamma) / \Gamma_{\text{total}}$ Γ_4 / Γ

VALUE (units 10^{-3}) CL% EVTS DOCUMENT ID TECN COMMENT

0.0024 ± 0.0012 19 BURKHARDT 87 NA31
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 < 0.013 90 BALATS 89 SPEC
 < 0.133 90 BARMIN 868 XEBC
 < 0.2 90 VASSERMAN 86 CALO $\phi \rightarrow K_S^0 K_L^0$
 < 0.4 90 0 BARMIN 738 HLBC
 < 0.71 90 0 15 BANNER 728 OSPK
 < 2.0 90 0 MORSE 728 DBC
 < 2.2 90 0 15 REPELLIN 71 OSPK
 < 21.0 90 0 15 BANNER 69 OSPK

15 These limits are for maximum interference in $K_S^0 - K_L^0$ to 2γ 's.

$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_5 / Γ

VALUE (units 10^{-4}) CL% DOCUMENT ID TECN COMMENT

< 0.88 90 METCALF 72 ASPK
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 < 0.49 90 16 BARMIN 85 HLBC $K^+ 850 \text{ MeV}$

16 BARMIN 85 assumes that CP-allowed and CP-violating amplitudes are equally suppressed.

$\Gamma(3\pi^0) / \Gamma_{\text{total}}$ Γ_6 / Γ

VALUE (units 10^{-4}) CL% DOCUMENT ID TECN

< 0.37 90 BARMIN 83 HLBC
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 < 4.3 90 BARMIN 73 HLBC

$\Gamma(\mu^+ \mu^-) / \Gamma(\pi^+ \pi^-)$ Γ_9 / Γ_1

Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-5}) CL% DOCUMENT ID TECN

< 0.047 90 GJESDAL 73 ASPK
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 < 20.0 90 BOHM 69 OSPK
 < 1.07 90 HYAMS 698 OSPK
 < 32.6 90 17 STUTZKE 69 OSPK
 < 10.0 90 BOTT... 67 OSPK

17 Value calculated by us, using 2.3 instead of 1 event, 90% CL.

Meson Full Listings

 K_S^0

$\Gamma(\pi^+e^-)/\Gamma(\pi^+\pi^-)$ Γ_{10}/Γ_1
 Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-5})	CL%	DOCUMENT ID	TECN
< 1.5	90	BARMIN	86 XEBC
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 16.0	90	18 BITSADZE	86 CALO
< 50.0	90	BOHM	69 OSPK
18 Use $B(\pi^+\pi^-) = 0.6861$.			

$\Gamma(\pi^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN
< 1.1	90	0	BARR	93B NA31
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 45	90		GIBBONS	88 E731

NOTE ON CP VIOLATION IN $K_S \rightarrow 3\pi$

(by T. Nakada, Paul Scherrer Institute and L. Wolfenstein, Carnegie-Mellon University)

The possible final states for the decay $K^0 \rightarrow \pi^+\pi^-\pi^0$ have isospin $I = 0, 1, 2$, and 3 . The $I = 0$ and $I = 2$ states have $CP = +1$ and K_S can decay into them without violating CP symmetry, but they are expected to be strongly suppressed by centrifugal barrier effects. The $I = 1$ and $I = 3$ states, which have no centrifugal barrier, have $CP = -1$ so that the K_S decay to these requires CP violation.

In order to see CP violation in $K_S \rightarrow \pi^+\pi^-\pi^0$, it is necessary to observe the interference between K_S and K_L decay, which determines the amplitude ratio

$$\eta_{+-0} = \frac{A(K_S \rightarrow \pi^+\pi^-\pi^0)}{A(K_L \rightarrow \pi^+\pi^-\pi^0)}.$$

If η_{+-0} is obtained from an integration over the whole Dalitz plot, there is no contribution from the $I = 0$ and $I = 2$ final states and a nonzero value of η_{+-0} is entirely due to CP violation.

Only $I = 1$ and $I = 3$ states, which are $CP = -1$, are allowed for $K^0 \rightarrow \pi^0\pi^0\pi^0$ decays and the decay of K_S into $3\pi^0$ is an unambiguous sign of CP violation. Similarly to η_{+-0} , η_{000} is defined as

$$\eta_{000} = \frac{A(K_S \rightarrow \pi^0\pi^0\pi^0)}{A(K_L \rightarrow \pi^0\pi^0\pi^0)}.$$

If one assumes that CPT invariance holds and that there are no transitions to $I = 3$ (or to nonsymmetric $I = 1$ states), it can be shown that

$$\eta_{+-0} = \eta_{000} = \epsilon + i \frac{\text{Im } a_1}{\text{Re } a_1}.$$

With the Wu-Yang phase convention, a_1 is the weak decay amplitude for K^0 into $I = 1$ final states; ϵ is determined from CP violation in $K_L \rightarrow 2\pi$ decays. The real parts of η_{+-0} and η_{000} are equal to $\text{Re}(\epsilon)$. Since currently-known upper limits on $|\eta_{+-0}|$ and $|\eta_{000}|$ are much larger than $|\epsilon|$, they can be interpreted as upper limits on $\text{Im}(\eta_{+-0})$ and $\text{Im}(\eta_{000})$.

CP-VIOLATION PARAMETERS IN K_S^0 DECAY

$\text{Im}(\eta_{+-0})^2 = \Gamma(K_S^0 \rightarrow \pi^+\pi^-\pi^0, CP\text{-violating}) / \Gamma(K_L^0 \rightarrow \pi^+\pi^-\pi^0)$
 CPT assumed valid (i.e. $\text{Re}(\eta_{+-0}) \approx 0$).

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.12	90	384	METCALF	72 ASPK	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.23	90	601	19 BARMIN	85 HLBC	K^+ 850 MeV
< 1.2	90	192	BALDO...	75 HLBC	
< 0.71	90	148	MALLARY	73 OSPK	$\text{Re}(A) = -0.05 \pm 0.17$
< 0.66	90	180	JAMES	72 HBC	
< 1.2	90	99	JONES	72 OSPK	
< 1.2	90	99	CHO	71 DBC	
< 1.0	90	98	JAMES	71 HBC	Incl. in JAMES 72
< 1.2	95	50	20 MEISNER	71 HBC	$\text{CL} = 90\%$ not avail.
< 0.8	90	71	WEBBER	70 HBC	
< 0.45	90		BEHR	66 HLBC	
< 3.8	90	18	ANDERSON	65 HBC	Incl. in WEBBER 70

19 BARMIN 85 find $\text{Re}(\eta_{+-0}) = (0.05 \pm 0.17)$ and $\text{Im}(\eta_{+-0}) = (0.15 \pm 0.33)$. Includes events of BALDO-CEOLIN 75.

20 These authors find $\text{Re}(A) = 2.75 \pm 0.65$, above value at $\text{Re}(A) = 0$.

$\text{Im}(\eta_{000})^2 = \Gamma(K_S^0 \rightarrow 3\pi^0) / \Gamma(K_L^0 \rightarrow 3\pi^0)$

CPT assumed valid (i.e. $\text{Re}(\eta_{000}) \approx 0$). This limit determines branching ratio $\Gamma(3\pi^0)/\Gamma_{\text{total}}$ above.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.1	90	632	21 BARMIN	83 HLBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.28	90		22 GJESDAL	74B SPEG	Indirect meas.
< 1.2	90	22	BARMIN	73 HLBC	

21 BARMIN 83 find $\text{Re}(\eta_{000}) = (-0.08 \pm 0.18)$ and $\text{Im}(\eta_{000}) = (-0.05 \pm 0.27)$. Assuming CPT invariance they obtain the limit quoted above.

22 GJESDAL 74B uses $K_{2\pi}$, $K_{\mu 3}$, and K_{e3} decay results, unitarity, and CPT . Calculates $|\langle \eta_{000} \rangle| = 0.26 \pm 0.20$. We convert to upper limit.

 K_S^0 REFERENCES

BARR	93B	PL B304 381	+Buchholz+ (CERN, EDIN, MANZ, LAO, PISA, SIEG)
GIBBONS	93	PRL 70 1199	+Barker, Briere, Makoff+ (FNAL E731 Collab.)
RAMBERG	93	PRL 70 2525	+Bock, Coleman, Enagonio, Hsiung+ (FNAL E731 Collab.)
BALATS	89	SJNP 49 928	+Berezin, Bogdanov, Vishnevskii, Vishnyakov+ (ITEP)
		Translated from YAF 49 1332	
GIBBONS	88	PRL 61 2661	+Papadimitriou+ (FNAL E731 Collab.)
BURKHARDT	87	PL B199 139	+ (CERN, EDIN, MANZ, LAO, PISA, SIEG)
GROSSMAN	87	PRL 59 18	+Heller, James, Shupe+ (MINN, MICH, RUTG)
BARMIN	86	SJNP 44 622	+Barylov, Davidenko, Demidov+ (ITEP)
		Translated from YAF 44 965	
BARMIN	86B	NC 85A 159	+Barylov, Chistyakova, Chuvilo+ (ITEP, PADO)
BITSADZE	86	PL 167B 138	+Budagov (CMNS, SOFI, SERP, TBIL, JINR, BAKU+)
PDG	86B	PL 170B 130	+Aguilar-Benitez, Porter+ (CERN, CIT+)
VASSERMAN	86	JETPL 43 588	+Golubev, Gluskin, Druzhinin+ (NOVO)
		Translated from ZETFP 43 457	
BARMIN	85	NC 85A 67	+Barylov, Chistyakova, Chuvilo+ (ITEP, PADO)
Also	85B	SJNP 41 759	+Barmin, Barylov, Volkov+ (ITEP, PADO)
		Translated from YAF 41 1187	
BARMIN	83	PL 128B 129	+Barylov, Chistyakova, Chuvilo+ (ITEP, PADO)
Also	84	SJNP 39 269	+Barmin, Barylov, Golubchikov+ (ITEP, PADO)
		Translated from YAF 39 428	
ARONSON	82	PRL 48 1078	+Berstein+ (BNL, CHIC, STAN, WISC)
ARONSON	82B	PRL 48 1306	+Bock, Cheng, Fischbach (BNL, CHIC, PURD)
Also	82B	PL 116B 73	+Fischbach, Cheng+ (PURD, BNL, CHIC)
Also	83	PR D28 476	+Aronson, Bock, Cheng+ (BNL, CHIC, PURD)
Also	83B	PR D28 495	+Aronson, Bock, Cheng+ (BNL, CHIC, PURD)
ARONSON	76	NC 32A 236	+McIntyre, Roehrig+ (WISC, EFI, UCSD, ILLC)
EVERHART	76	PR D14 661	+Kraus, Lande, Long, Lowenstein+ (PENN)
TAUREG	76	PL 65B 92	+Zech, Dydak, Navarra+ (HEIDH, CERN, DORT)
BALDO...	75	NC 25A 698	+Baldo-Ceolin, Bobisut, Calimani+ (PADO, WISC)
CARITHERS	75	PRL 34 1244	+Modis, Nygren, Pun+ (COLU, NYU)
BOBISUT	74	LN C 11 646	+Huzita, Mattioli, Puglierin (PADO)
COWELL	74	PR D10 2083	+Lee-Franzini, Orcutt, Franzini+ (STON, COLU)
GEWENIGER	74B	PL 48B 487	+Gjesdal, Presser+ (CERN, HEIDH)
GJESDAL	74B	PL 52B 119	+Presser, Steffen+ (CERN, HEIDH)
BARMIN	73	PL 46B 465	+Barylov, Davidenko, Demidov+ (ITEP)
BARMIN	73B	PL 47B 463	+Barylov, Davidenko, Demidov+ (ITEP)
BURGUN	73	PL 46B 481	+Bertranet, Lesquoy, Muller, Pauli+ (SACL, CERN)
FACKLER	73	PRL 31 847	+Frisch, Martin, Smoot, Sompayrac (MIT)
GJESDAL	73	PL 44B 217	+Presser, Steffen, Steinberger+ (CERN, HEIDH)
HILL	73	PR D6 1290	+Sakitt, Samios, Burris, Engler+ (BNL, CMU)
MALLARY	73	PR D7 1953	+Binnie, Gallivan, Gomez, Peck, Sciuilli+ (CIT)
ALITTI	72	PL 39B 568	+Lesquoy, Muller (SACL)
BANNER	72B	PRL 29 237	+Cronin, Hoffman, Knapp, Shochet (PRIN)
BURGUN	72	NP B50 194	+Lesquoy, Muller, Pauli+ (SACL, CERN, OSLO)
JAMES	72	NP B49 119	+Montanet, Paul, Saetre+ (CERN, SACL, OSLO)
JONES	72	NC 9A 151	+Abashian, Graham, Mantsch, Orr, Smith+ (ILL)
METCALF	72	PL 40B 703	+Neuhofer, Niebergall+ (CERN, IPN, WIEN)
MORSE	72B	PRL 28 388	+Nauenberg, Bierman, Sager+ (COLO, PRIN, UMD)
NAGY	72	NP B47 94	+Telbisz, Vestergombi (BUDA)
Also	69	PL 30B 498	+Bozoki, Fenyves, Gombosi, Nagy+ (BUDA)
SKIEGGEST...	70	NP B48 343	+Skeggestad, James+ (OSLO, CERN, SACL)
BALTAY	71	PRL 27 1678	+Bridgewater, Cooper, Gershwin, Habibi+ (COLU)
Also	71	Nevis 187 Thesis	+Cooper (COLU)
CHO	71	PR D3 1557	+Dralle, Canter, Engler, Fisk+ (CMU, BNL, CASE)
JAMES	71	PL 35B 265	+Montanet, Paul, Pauli+ (CERN, SACL, OSLO)
MEISNER	71	PR D3 59	+Mann, Hertzbach, Koffer+ (MASA, BNL, YALE)
REPELLIN	71	PL 36B 603	+Wolff, Chollet, Galliard, Jane+ (ORSAY, CERN)
MOFFETT	70	RAPS 15 512	+Gobbi, Green, Hakal, Rosen+ (ROCH)
WEBBER	70	PR D1 1967	+Solmitz, Crawford, Alston-Garnjost (LRL)
Also	69	UCRL 19226 Thesis	+Webber (LRL)

BANNER 69	PR 188 2033	+Cronin, Liu, Pilcher	(PRIN)
BOHM 69	Thesis		(AACH)
BOZOKI 69	PL 308 498	+Fenyves, Gombosi, Nagy+	(BUDA)
DOYLE 69	UCRL 18139 Thesis		(LRL)
GOBBI 69	PRL 22 682	+Green, Hakel, Moffett, Rosen+	(ROCH)
HYAMS 69B	PL 29B 521	+Koch, Potter, VonLindern, Lorenz+	(CERN, MPIM)
MORFIN 69	PRL 23 660	+Sinclair	(MICH)
STUTZKE 69	PR 177 2009	+Abashian, Jones, Mantsch, Orr, Smith	(ILL)
DONALD 68B	PL 27B 58	+Edwards, Nisar+	(LIVP, CERN, IPNP, CDEF)
HILL 68	PL 171 1418	+Robinson, Sakitt+	(BNL, CMU)
BOTT... 67	PL 24B 194	Bott-Sodenhausen, DeBouard, Cassel+	(CERN)
ALFF... 66B	PL 21 595	Alff-Steinberger, Heuer, Kleinknecht+	(CERN)
BEHR 66	PL 22 540	+Brisson, Petiau+ (EPOL, MILA, PADO, ORSAY)	
BELLOTTI 66	NC 45A 737	+Pulla, Baldo-Ceolin+	(MILA, PADO)
KIRSCH 66	PR 147 939	+Schmidt	(COLU)
ANDERSON 65	PRL 14 475	+Crawford, Golden, Stern, Binford+	(LRL, WISC)
AUBERT 65	PL 17 59	+Behr, Canavan, Chounet+	(EPOL, ORSAY)
BROWN 63	PR 130 769	+Kadyk, Trilling, Roe+	(LRL, MICH)
CHRETIEN 63	PR 131 2208		(BRAN, BROW, HARV, MIT)
ANDERSON 62B	CERN Conf. 836	+Crawford+	(LRL)
BROWN 61	NC 19 1155	+Bryant, Burnstein, Glaser, Kadyk+	(MICH)
BAGLIN 60	NC 18 1043	+Bloch, Brisson, Hennessy+	(EPOL)
COLUMBIA 60B	Rochester Conf. 727	Schwartz+	(COLU)
CRAWFORD 59B	PRL 2 266	+Cresti, Douglass, Good, Ticho+	(LRL)
BOLDT 58B	PRL 1 150	+Caldwell, Pal	(MIT)

OTHER RELATED PAPERS

LITTENBERG 93	ARNPS 43 729	+Valencia	(BNL, FNAL)
Rare and Radiative Kaon Decays			
BATTISTON 92	PR 214 293	+Cocolicchio, Fogli, Paver	(PGIA, CERN, TRSTT)
Status and Perspectives of K Decay Physics			
TRILLING 65B	UCRL 16473		(LRL)
Updated from 1965 Argonne Conference, page 115.			
CRAWFORD 62	CERN Conf. 827		(LRL)
FITCH 61	NC 22 1160	+Piroue, Perkins	(PRIN, LASL)
GOOD 61	PR 124 1223	+Matsen, Muller, Piccioni+	(LRL)
BIRGE 60	Rochester Conf. 601	+Ely+	(LRL, WISC)
MULLER 60	PRL 4 418	+Birge, Fowler, Good, Piccioni+	(LRL, BNL)



$$I(J^P) = \frac{1}{2}(0^-)$$

$$m_{K_L^0} - m_{K_S^0}$$

For earlier measurements, beginning with GOOD 61 and FITCH 61, see our 1986 edition, Physics Letters **170B** 132 (1986).

VALUE (10^{10} h s^{-1})	DOCUMENT ID	TECN	COMMENT
0.5333 ± 0.0027	OUR AVERAGE		Error includes scale factor of 1.2.
0.5257 ± 0.0049	¹ GIBBONS	93C E731	20-160 GeV K beams
0.5340 ± 0.00255 ± 0.0015	² GEWENIGER	74C SPEC	Gap method
0.5334 ± 0.0040 ± 0.0015	² GJESDAL	74 SPEC	Charge asymmetry
0.542 ± 0.006	CULLEN	70 CNTR	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.5286 ± 0.0028	³ GIBBONS	93 E731	20-160 GeV K beams
0.482 ± 0.014	⁴ ARONSON	82B SPEC	E=30-110 GeV
0.534 ± 0.007	⁵ CARNEGIE	71 ASPK	Gap method
0.542 ± 0.006	⁵ ARONSON	70 ASPK	Gap method

- ¹GIBBONS 93C fits $\Delta(m)$ and ϕ_{+-} . Finds above $\Delta(m)$ and $\phi_{+-} = (42.2 \pm 1.4)^\circ$.
- ²These two experiments have a common systematic error due to the uncertainty in the momentum scale, as pointed out in WAHL 89.
- ³GIBBONS 93 value assume $\phi_{+-} = \phi_{00} = \phi_{SW} = (43.7 \pm 0.2)^\circ$.
- ⁴ARONSON 82 find that $\Delta(m)$ may depend on the kaon energy.
- ⁵ARONSON 70 and CARNEGIE 71 use K_S^0 mean life = $(0.862 \pm 0.006) \times 10^{-10}$ s. We have not attempted to adjust these values for the subsequent change in the K_S^0 mean life or in η_{+-} .

K_L^0 MEAN LIFE

VALUE (10^{-8} s)	EVTS	DOCUMENT ID	TECN
5.17 ± 0.04	OUR FIT		
5.15 ± 0.04	OUR AVERAGE		
5.154 ± 0.044	0.4M	VOSBURGH 72	CNTR
5.15 ± 0.14		DEVLIN 67	CNTR
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.0 ± 0.5		⁶ LOWYS 67	HLBC
6.1 +1.5 -1.2	1700	ASTBURY 65C	CNTR
5.3 ± 0.6		FUJII 64	OSPK
5.1 +2.4 -1.3	15	DARMON 62	FBC
8.1 +3.2 -2.4	34	BARDON 58	CNTR

⁶Sum of partial decay rates.

K_L^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $3\pi^0$	(21.6 ± 0.8) %	S=1.5
Γ_2 $\pi^+\pi^-\pi^0$	(12.38 ± 0.21) %	S=1.5
Γ_3 $\pi^\pm\mu^\mp\nu$	[a] (27.0 ± 0.4) %	S=1.3
Called $K_{\mu 3}^0$.		
Γ_4 $\pi^-\mu^+\nu_\mu$		
Γ_5 $\pi^+\mu^-\bar{\nu}_\mu$		
Γ_6 $\pi^\pm e^\mp\nu$	[a] (38.7 ± 0.5) %	S=1.4
Called $K_{e 3}^0$.		
Γ_7 $\pi^-e^+\nu_e$		
Γ_8 $\pi^+e^-\bar{\nu}_e$		
Γ_9 2γ	(5.73 ± 0.27) × 10 ⁻⁴	S=2.0
Γ_{10} $\pi^0 2\gamma$	[b] (1.70 ± 0.28) × 10 ⁻⁶	
Γ_{11} $\pi^0\pi^\pm e^\mp\nu$	[a] (5.18 ± 0.29) × 10 ⁻⁵	
Γ_{12} ($\pi\mu\text{atom}$) ν	(1.05 ± 0.11) × 10 ⁻⁷	
Γ_{13} $\pi^\pm e^\mp\nu_e\gamma$	[a,b,c] (1.3 ± 0.8) %	
Γ_{14} $\pi^+\pi^-\gamma$	[b,c] (4.61 ± 0.14) × 10 ⁻⁵	
Γ_{15} $\pi^0\pi^0\gamma$	< 5.6 × 10 ⁻⁶	

Charge conjugation × Parity (CP) or Lepton Family number (LF) violating modes, or $\Delta S = 1$ weak neutral current (SI) modes

Γ_{16} $\pi^+\pi^-$	CPV	(2.03 ± 0.04) × 10 ⁻³	S=1.2
Γ_{17} $\pi^0\pi^0$	CPV	(9.14 ± 0.34) × 10 ⁻⁴	S=1.8
Γ_{18} $\mu^+\mu^-$	SI	(7.4 ± 0.4) × 10 ⁻⁹	
Γ_{19} $\mu^+\mu^-\gamma$	SI	(2.8 ± 2.8) × 10 ⁻⁷	
Γ_{20} e^+e^-	SI	< 4.1 × 10 ⁻¹¹	CL=90%
Γ_{21} $e^+e^-\gamma$	SI	(9.1 ± 0.5) × 10 ⁻⁶	
Γ_{22} $e^+e^-\gamma\gamma$	SI	[b] (6.6 ± 3.2) × 10 ⁻⁷	
Γ_{23} $\pi^+\pi^-e^+e^-$	SI	< 2.5 × 10 ⁻⁶	CL=90%
Γ_{24} $\mu^+\mu^-e^+e^-$	SI	< 4.9 × 10 ⁻⁶	CL=90%
Γ_{25} $e^+e^-e^+e^-$	SI	[d] (3.9 ± 0.7) × 10 ⁻⁸	
Γ_{26} $\pi^0\mu^+\mu^-$	CP,SI	[e] < 5.1 × 10 ⁻⁹	CL=90%
Γ_{27} $\pi^0e^+e^-$	CP,SI	[e] < 4.3 × 10 ⁻⁹	CL=90%
Γ_{28} $\pi^0\nu\bar{\nu}$	CP,SI	[f] < 2.2 × 10 ⁻⁴	CL=90%
Γ_{29} $e^\pm\mu^\mp$	LF	[a] < 3.3 × 10 ⁻¹¹	CL=90%

- [a] The value is for the sum of the charge states indicated.
- [b] See the Full Listings below for the energy limits used in this measurement.
- [c] Most of this radiative mode, the low-momentum γ part, is also included in the parent mode listed without γ 's.
- [d] $m_{e^+e^-} > 470$ MeV
- [e] Allowed by higher-order electroweak interactions.
- [f] Violates CP in leading order. Test of direct CP violation since the indirect CP-violating and CP-conserving contributions are expected to be suppressed.

CONSTRAINED FIT INFORMATION

An overall fit to the mean life, 4 decay rate, and 12 branching ratios uses 53 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 55.3$ for 46 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-35						
x_3	-78	6					
x_6	-86	7	46				
x_9	-2	13	-2	-3			
x_{16}	-29	45	14	15	34		
x_{17}	-9	20	2	2	74	48	
Γ	2	-4	0	0	-1	-2	-1
	x_1	x_2	x_3	x_6	x_9	x_{16}	x_{17}

Meson Full Listings

K_L^0

Mode	Rate (10^8 s^{-1})	Scale factor
$\Gamma_1 \quad 3\pi^0$	0.0419 ± 0.0016	1.4
$\Gamma_2 \quad \pi^+ \pi^- \pi^0$	0.0239 ± 0.0004	1.4
$\Gamma_3 \quad \pi^\pm \mu^\mp \nu$ Called $K_{\mu 3}^0$.	[a] 0.0522 ± 0.0008	1.2
$\Gamma_6 \quad \pi^\pm e^\mp \nu$ Called $K_{e 3}^0$.	[a] 0.0748 ± 0.0011	1.3
$\Gamma_9 \quad 2\gamma$	$(1.11 \pm 0.05) \times 10^{-4}$	1.9
$\Gamma_{16} \quad \pi^+ \pi^-$	$(3.93 \pm 0.08) \times 10^{-4}$	1.2
$\Gamma_{17} \quad \pi^0 \pi^0$	$(1.77 \pm 0.07) \times 10^{-4}$	1.7

K_L^0 DECAY RATES

$\Gamma(3\pi^0)$ Γ_1

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	COMMENT
4.19 ± 0.16 OUR FIT				Error includes scale factor of 1.4.
$5.22^{+1.03}_{-0.84}$	54	BEHR	66	HLBC Assumes CP

$\Gamma(\pi^+ \pi^- \pi^0)$ Γ_2

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	COMMENT
2.39 ± 0.04 OUR FIT				Error includes scale factor of 1.4.
2.38 ± 0.09 OUR AVERAGE				
$2.32^{+0.13}_{-0.15}$	192	BALDO...	75	HLBC Assumes CP
2.35 ± 0.20	180	⁷ JAMES	72	HBC Assumes CP
2.71 ± 0.28	99	CHO	71	DBC Assumes CP
2.12 ± 0.33	50	MEISNER	71	HBC Assumes CP
2.20 ± 0.35	53	WEBBER	70	HBC Assumes CP
$2.62^{+0.28}_{-0.27}$	136	BEHR	66	HLBC Assumes CP
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.5 ± 0.3	98	⁷ JAMES	71	HBC Assumes CP
3.26 ± 0.77	18	ANDERSON	65	HBC
1.4 ± 0.4	14	FRANZINI	65	HBC

In the fit this rate is well determined by the mean life and the branching ratio $\Gamma(\pi^+ \pi^- \pi^0) / [\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$. For this reason the discrepancy between the $\Gamma(\pi^+ \pi^- \pi^0)$ measurements does not affect the scale factor of the overall fit.

⁷JAMES 72 is a final measurement and includes JAMES 71.

$\Gamma(\pi^\pm \mu^\mp \nu)$ Γ_3

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	
5.22 ± 0.08 OUR FIT			Error includes scale factor of 1.2.	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$4.54^{+1.24}_{-1.08}$	19	LOWYS	67	HLBC

$\Gamma(\pi^\pm e^\mp \nu)$ Γ_6

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	COMMENT
7.48 ± 0.11 OUR FIT				Error includes scale factor of 1.3.
7.7 ± 0.5 OUR AVERAGE				
7.81 ± 0.56	620	CHAN	71	HBC
$7.52^{+0.85}_{-0.72}$		AUBERT	65	HLBC $\Delta S = \Delta Q, CP$ assumed

$\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)$ ($\Gamma_2 + \Gamma_3 + \Gamma_6$)

K_L^0 - charged.

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	
15.10 ± 0.19 OUR FIT			Error includes scale factor of 1.3.	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
15.1 ± 1.9	98	AUERBACH	66B	OSPK

$\Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)$ ($\Gamma_3 + \Gamma_6$)

VALUE (10^6 s^{-1})	EVTS	DOCUMENT ID	TECN	COMMENT
12.70 ± 0.18 OUR FIT				Error includes scale factor of 1.2.
11.9 ± 0.6 OUR AVERAGE				Error includes scale factor of 1.2.
12.4 ± 0.7	410	⁸ BURGUN	72	HBC $K^+ p \rightarrow K^0 p \pi^+$
13.1 ± 1.3	252	⁸ WEBBER	71	HBC $K^- p \rightarrow n \bar{K}^0$
11.6 ± 0.9	393	^{8,9} CHO	70	DBC $K^+ n \rightarrow K^0 p$
$9.85^{+1.15}_{-1.05}$	109	⁸ FRANZINI	65	HBC
• • • We do not use the following data for averages, fits, limits, etc. • • •				
8.47 ± 1.69	126	⁸ MANN	72	HBC $K^- p \rightarrow n \bar{K}^0$
10.3 ± 0.8	335	⁹ HILL	67	DBC $K^+ n \rightarrow K^0 p$

⁸ Assumes $\Delta S = \Delta Q$ rule.
⁹ CHO 70 includes events of HILL 67.

K_L^0 BRANCHING RATIOS

$\Gamma(3\pi^0) / \Gamma(\pi^+ \pi^- \pi^0)$ Γ_1 / Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.75 ± 0.08 OUR FIT				Error includes scale factor of 1.4.
1.81 ± 0.13 OUR AVERAGE				
1.80 ± 0.13	1010	BUDAGOV	68	HLBC
2.0 ± 0.6	188	ALEKSANYAN	64B	FBC
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.65 ± 0.07	883	BARMIN	72B	HLBC Error statistical only

$\Gamma(3\pi^0) / [\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ $\Gamma_1 / (\Gamma_2 + \Gamma_3 + \Gamma_6)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.277 ± 0.013 OUR FIT				Error includes scale factor of 1.5.
0.260 ± 0.011 OUR AVERAGE				
0.251 ± 0.014	549	BUDAGOV	68	HLBC ORSAY measur.
0.277 ± 0.021	444	BUDAGOV	68	HLBC Ecole polytec.meas
$0.31^{+0.07}_{-0.06}$	29	KULYUKINA	68	CC
0.24 ± 0.08	24	ANIKINA	64	CC


$\Gamma(\pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_2 / Γ

VALUE	DOCUMENT ID
0.1238 ± 0.0021 OUR FIT	Error includes scale factor of 1.5.

$\Gamma(\pi^+ \pi^- \pi^0) / [\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ $\Gamma_2 / (\Gamma_2 + \Gamma_3 + \Gamma_6)$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.1586 ± 0.0026 OUR FIT				Error includes scale factor of 1.6.
0.1588 ± 0.0024 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.163 ± 0.003	6499	CHO	77	HBC
0.1605 ± 0.0038	1590	ALEXANDER	73B	HBC
0.146 ± 0.004	3200	BRANDENB...	73	HBC
0.159 ± 0.010	558	EVANS	73	HLBC
0.167 ± 0.016	1402	KULYUKINA	68	CC
0.161 ± 0.005		HOPKINS	67	HBC
0.162 ± 0.015	126	HAWKINS	66	HBC
0.159 ± 0.015	326	ASTBURY	65B	CC
0.178 ± 0.017	566	GUIDONI	65	HBC
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$0.15^{+0.03}_{-0.04}$	66	ASTBURY	65	CC
0.144 ± 0.004	1729	HOPKINS	65	HBC See HOPKINS 67
0.151 ± 0.020	79	ADAIR	64	HBC
$0.157^{+0.03}_{-0.04}$	75	LUERS	64	HBC
0.185 ± 0.038	59	ASTIER	61	CC

WEIGHTED AVERAGE
 0.1588 ± 0.0024 (Error scaled by 1.4)



K_L^0 $\Gamma(\pi^\pm \mu^\mp \nu)/\Gamma(\pi^\pm e^\mp \nu)$ Γ_3/Γ_6

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.697 ± 0.010 OUR FIT				
0.697 ± 0.010 OUR AVERAGE				
0.702 ± 0.011	33k	CHO	80	HBC
0.662 ± 0.037	10k	WILLIAMS	74	ASPK
0.741 ± 0.044	6700	BRANDENB...	73	HBC
0.662 ± 0.030	1309	EVANS	73	HLBC
0.71 ± 0.05	770	BUDAGOV	68	HLBC
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.68 ± 0.08	3548	BASILE	70	OSPK
0.71 ± 0.04	569	10 BEILLIERE	69	HLBC
0.648 ± 0.030	1309	EVANS	69	HLBC Repl. by EVANS 73
0.67 ± 0.13		11 KULYUKINA	68	CC
0.82 ± 0.10		DEBOUARD	67	OSPK
0.7 ± 0.2	273	HAWKINS	67	HBC
0.81 ± 0.08		HOPKINS	67	HBC
0.81 ± 0.19		ADAIR	64	HBC

¹⁰BEILLIERE 69 is a scanning experiment using same exposure as BUDAGOV 68.

¹¹KULYUKINA 68 $\Gamma(\pi^\pm \mu^\mp \nu)/\Gamma(\pi^\pm e^\mp \nu)$ is not measured independently from $\Gamma(\pi^+ \pi^- \pi^0)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ and $\Gamma(\pi^\pm e^\mp \nu)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$.

 $\Gamma(\pi^\pm \mu^\mp \nu)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ $\Gamma_3/(\Gamma_2 + \Gamma_3 + \Gamma_6)$

VALUE	EVTS	DOCUMENT ID	TECN
0.3456 ± 0.0030 OUR FIT			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.335 ± 0.055	330	12 KULYUKINA	68
0.39 +0.08 -0.10	172	12 ASTBURY	65
0.356 ± 0.07	251	12 LUERS	64

¹²This mode not measured independently from $\Gamma(\pi^+ \pi^- \pi^0)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ and $\Gamma(\pi^\pm e^\mp \nu)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$.

 $\Gamma(\pi^\pm e^\mp \nu)/[\Gamma(\pi^+ \pi^- \pi^0) + \Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ $\Gamma_6/(\Gamma_2 + \Gamma_3 + \Gamma_6)$

VALUE	EVTS	DOCUMENT ID	TECN
0.4950 ± 0.0032 OUR FIT			
Error includes scale factor of 1.1.			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.498 ± 0.052	500	KULYUKINA	68
0.46 +0.08 -0.10	202	ASTBURY	65
0.487 ± 0.05	153	LUERS	64
0.46 ± 0.11	24	NYAGU	61

 $\Gamma(\pi^\pm e^\mp \nu)/[\Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ $\Gamma_6/(\Gamma_3 + \Gamma_6)$

VALUE	EVTS	DOCUMENT ID	TECN
0.5093 ± 0.0033 OUR FIT			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.415 ± 0.120	320	ASTIER	61

 $[\Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]/\Gamma_{total}$ $(\Gamma_3 + \Gamma_6)/\Gamma$

VALUE	DOCUMENT ID	TECN
0.656 ± 0.007 OUR FIT		
Error includes scale factor of 1.5.		

 $\Gamma(2\gamma)/\Gamma_{total}$ Γ_9/Γ

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT
5.73 ± 0.27 OUR FIT				Error includes scale factor of 2.0.
4.9 ± 0.5 OUR AVERAGE				
4.54 ± 0.84	13	BANNER	72B	OSPK
4.5 ± 1.0	23	ENSTROM	71	OSPK K_L^0 1.5-9 GeV/c
5.5 ± 1.1	90	KUNZ	68	OSPK Norm. to $3\pi(C+N)$
6.7 ± 2.2	32	TODOROFF	67	OSPK Repl. CRIEGEE 66
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5.0 ± 1.0	14	REPELLIN	71	OSPK
7.4 ± 1.6	33	15 CRONIN	67	OSPK
1.3 ± 0.6	16	CRIEGEE	66	OSPK

¹³This value uses $(\eta_{00}/\eta_{+-})^2 = 1.05 \pm 0.14$. In general, $\Gamma(2\gamma)/\Gamma_{total} = [(4.32 \pm 0.55) \times 10^{-4}]/[(\eta_{00}/\eta_{+-})^2]$.

¹⁴Assumes regeneration amplitude in copper at 2 GeV is 22 mb. To evaluate for a given regeneration amplitude and error, multiply by (regeneration amplitude/22mb)².

¹⁵CRONIN 67 replaced by KUNZ 68.

¹⁶CRIEGEE 66 replaced by TODOROFF 67.

 $\Gamma(2\gamma)/\Gamma(3\pi^0)$ Γ_9/Γ_1

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
2.65 ± 0.16 OUR FIT				Error includes scale factor of 1.7.
2.24 ± 0.22 OUR AVERAGE				
2.13 ± 0.43	28	BARMIN	71	HLBC
2.24 ± 0.28	115	BANNER	69	OSPK
2.5 ± 0.7	16	ARNOLD	68B	HLBC Vacuum decay

 $\Gamma(2\gamma)/\Gamma(\pi^0 \pi^0)$ Γ_9/Γ_{17}

VALUE	EVTS	DOCUMENT ID	TECN
0.627 ± 0.020 OUR FIT			
Error includes scale factor of 2.3.			
0.632 ± 0.004 ± 0.008	110k	BURKHARDT 87	NA31

 $\Gamma(\pi^0 2\gamma)/\Gamma_{total}$ Γ_{10}/Γ

VALUE (units 10 ⁻⁶)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.7 ± 0.2 ± 0.2		63	BARR	92	SPEC
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.86 ± 0.60 ± 0.60		60	PAPADIMITR...91	E731	$m_{\gamma\gamma} > 280$ MeV
< 5.1		90	PAPADIMITR...91	E731	$m_{\gamma\gamma} < 264$ MeV
2.1 ± 0.6		14	17 BARR	90C	NA31 $m_{\gamma\gamma} > 280$ MeV
< 2.7		90	PAPADIMITR...89	E731	In PAPADI...91
< 230		90	0 BANNER	69	OSPK
¹⁷ BARR 90C superseded by BARR 92.					

 $\Gamma(\pi^0 \pi^\pm e^\mp \nu)/\Gamma_{total}$ Γ_{11}/Γ

VALUE (units 10 ⁻⁵)	CL%	EVTS	DOCUMENT ID	TECN
5.18 ± 0.29 OUR AVERAGE				
5.16 ± 0.20 ± 0.22		729	MAKOFF	93
6.2 ± 2.0		16	CARROLL	80C
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 220		90	18 DONALDSON	74
¹⁸ DONALDSON 74 uses $K_L^0 \rightarrow \pi^+ \pi^- \pi^0$ (all K_L^0) decays = 0.126.				

 $\Gamma((\pi \text{ atom})\nu)/\Gamma(\pi^\pm \mu^\mp \nu)$ Γ_{12}/Γ_3

VALUE (units 10 ⁻⁷)	EVTS	DOCUMENT ID	TECN
3.90 ± 0.39	155	19 ARONSON	86
• • • We do not use the following data for averages, fits, limits, etc. • • •			
seen	18	COOMBS	76
¹⁹ ARONSON 86 quote theoretical value of $(4.31 \pm 0.08) \times 10^{-7}$.			

 $\Gamma(\pi^\pm e^\mp \nu \gamma)/\Gamma(\pi^\pm e^\mp \nu)$ Γ_{13}/Γ_6

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
3.3 ± 2.0	10	PEACH	71	HLBC γ KE > 15 MeV

 $\Gamma(\pi^+ \pi^- \gamma)/\Gamma_{total}$ Γ_{14}/Γ

VALUE (units 10 ⁻⁵)	EVTS	DOCUMENT ID	TECN	COMMENT
4.61 ± 0.14 OUR AVERAGE				
4.66 ± 0.15	3136	20 RAMBERG	93	E731 $E_\gamma > 20$ MeV
4.41 ± 0.32	1062	21 CARROLL	80B	SPEC $E_\gamma > 20$ MeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.52 ± 0.16	516	22 CARROLL	80B	SPEC $E_\gamma > 20$ MeV
2.89 ± 0.28	546	23 CARROLL	80B	SPEC
6.2 ± 2.1	24	24 DONALDSON	74C	SPEC

²⁰RAMBERG 93 finds that fraction of Direct Emission (DE) decays with $E_\gamma > 20$ MeV is 0.685 ± 0.041.

²¹Both components. Uses $K_L^0 \rightarrow \pi^+ \pi^- \pi^0$ (all K_L^0) decays = 0.1239.

²²Internal Bremsstrahlung component only.

²³Direct γ emission component only.

²⁴Uses $K_L^0 \rightarrow \pi^+ \pi^- \pi^0$ (all K_L^0) decays = 0.126.

 $\Gamma(\pi^0 \pi^0 \gamma)/\Gamma_{total}$ Γ_{15}/Γ

VALUE (units 10 ⁻⁶)	DOCUMENT ID	TECN
< 5.6	BARR	94

 $\Gamma(\pi^+ \pi^-)/\Gamma_{total}$ Γ_{16}/Γ

VALUE (units 10 ⁻³)	DOCUMENT ID
2.03 ± 0.04 OUR FIT	
Error includes scale factor of 1.2.	
2.102 ± 0.063	25 ETAFIT

²⁵This ETAFIT value is computed from fitted values of $|\eta_{+-}|$, the K_L^0 and K_S^0 lifetimes, and the $K_S^0 \rightarrow \pi^+ \pi^-$ branching fraction. See the discussion in the "Note on CP violation in K_L^0 decay."

 $\Gamma(\pi^+ \pi^-)/\Gamma(\pi^+ \pi^- \pi^0)$ Γ_{16}/Γ_2

VALUE (units 10 ⁻²)	EVTS	DOCUMENT ID	TECN	COMMENT
1.639 ± 0.032 OUR FIT				Error includes scale factor of 1.1.
1.64 ± 0.04	4200	MESSNER	73	ASPK $\eta_{+-} = 2.23$

 $\Gamma(\pi^+ \pi^-)/[\Gamma(\pi^\pm \mu^\mp \nu) + \Gamma(\pi^\pm e^\mp \nu)]$ $\Gamma_{16}/(\Gamma_3 + \Gamma_6)$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
3.09 ± 0.06 OUR FIT				Error includes scale factor of 1.2.
3.08 ± 0.10 OUR AVERAGE				
3.13 ± 0.14	1687	COUPAL	85	SPEC $\eta_{+-} = 2.28 \pm 0.06$
3.04 ± 0.14	2703	DEVUE	77	SPEC $\eta_{+-} = 2.25 \pm 0.05$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.51 ± 0.23	309	26 DEBOUARD	67	OSPK $\eta_{+-} = 2.00 \pm 0.09$
2.35 ± 0.19	525	26 FITCH	67	OSPK $\eta_{+-} = 1.94 \pm 0.08$

²⁶Old experiments excluded from fit. See subsection on η_{+-} in section on "PARAMETERS FOR $K_L^0 \rightarrow 2\pi$ DECAY" below for average η_{+-} of these experiments and for note on discrepancy.

Meson Full Listings

K_L^0

$\Gamma(\pi^+\pi^-)/[\Gamma(\pi^+\pi^-\pi^0) + \Gamma(\pi^\pm\mu^\mp\nu) + \Gamma(\pi^\pm e^\mp\nu)]$ $\Gamma_{16}/(\Gamma_2+\Gamma_3+\Gamma_6)$
 Violates CP conservation.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.60 ± 0.06 OUR FIT				Error includes scale factor of 1.1.
2.60 ± 0.07	4200	27 MESSNER	73 ASPK	$\eta_{+-} = 2.23 \pm 0.05$
1.93 ± 0.26		28 BASILE	66 OSPK	$\eta_{+-} = 1.92 \pm 0.13$
1.993 ± 0.080		28 BOTT...	66 OSPK	$\eta_{+-} = 1.95 \pm 0.04$
2.08 ± 0.35	54	28 GALBRAITH	65 OSPK	$\eta_{+-} = 1.99 \pm 0.16$
2.0 ± 0.4	45	28 CHRISTENS...	64 OSPK	$\eta_{+-} = 1.95 \pm 0.20$

²⁷ From same data as $\Gamma(\pi^+\pi^-)/\Gamma(\pi^+\pi^-\pi^0)$ MESSNER 73, but with different normalization.
²⁸ Old experiments excluded from fit. See subsection on η_{+-} in section on "PARAMETERS FOR $K_L^0 \rightarrow 2\pi$ DECAY" below for average η_{+-} of these experiments and for note on discrepancy.

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$ Γ_{17}/Γ
 Violates CP conservation.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.914 ± 0.034 OUR FIT				Error includes scale factor of 1.8.
2.5 ± 0.8	189	29 GAILLARD	69 OSPK	$\eta_{00} = 3.6 \pm 0.6$
1.2 $^{+1.5}_{-1.2}$	7	30 CRIEGEE	66 OSPK	

²⁹ Latest result of this experiment given by FAISSNER 70 $\Gamma(\pi^0\pi^0)/\Gamma(3\pi^0)$.
³⁰ CRIEGEE 66 experiment not designed to measure $2\pi^0$ decay mode.


$\Gamma(\pi^0\pi^0)/\Gamma(3\pi^0)$ Γ_{17}/Γ_1
 Violates CP conservation.

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
0.422 ± 0.023 OUR FIT				Error includes scale factor of 1.6.
0.44 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
1.21 ± 0.30	150	31 REY	76 OSPK	$\eta_{00} = 3.8 \pm 0.5$
0.37 ± 0.08	29	BARMIN	70 HLBC	$\eta_{00} = 2.02 \pm 0.23$
0.32 ± 0.15	30	BUDAGOV	70 HLBC	$\eta_{00} = 1.9 \pm 0.5$
0.90 ± 0.30	172	32 FAISSNER	70 OSPK	$\eta_{00} = 3.2 \pm 0.5$
0.46 ± 0.11	57	BANNER	69 OSPK	$\eta_{00} = 2.2 \pm 0.3$

not seen BARTLETT 68 OSPK See η_{00} below
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 1.31 ± 0.31 133 31 CENCE 69 OSPK $\eta_{00} = 3.7 \pm 0.5$
 1.89 ± 0.31 109 33 CRONIN 67 OSPK $\eta_{00} = 4.9 \pm 0.5$
 1.36 ± 0.18 33 CRONIN 67B OSPK $\eta_{00} = 3.92 \pm 0.3$

³¹ CENCE 69 events are included in REY 76.
³² FAISSNER 70 contains same $2\pi^0$ events as GAILLARD 69 $\Gamma(\pi^0\pi^0)/\Gamma_{total}$.
³³ CRONIN 67B is further analysis of CRONIN 67, now both withdrawn.

WEIGHTED AVERAGE
 0.44 ± 0.09 (Error scaled by 1.6)



$\Gamma(\pi^0\pi^0)/\Gamma(\pi^+\pi^-)$ Γ_{17}/Γ_{16}
 Violates CP conservation.

VALUE	DOCUMENT ID	TECN
0.451 ± 0.015 OUR FIT		
0.4539 ± 0.0061	34 ETAFIT	94

³⁴ This ETAFIT value is computed from fitted values of $|\eta_{00} / \eta_{+-}|$ and the $\Gamma(K_S^0 \rightarrow \pi^+\pi^-) / \Gamma(K_S^0 \rightarrow \pi^0\pi^0)$ branching fraction. See the discussion in the "Note on CP violation in K_L^0 decay."

$\Gamma(\mu^+\mu^-)/[\Gamma(\pi^+\pi^-\pi^0) + \Gamma(\pi^\pm\mu^\mp\nu) + \Gamma(\pi^\pm e^\mp\nu)]$ $\Gamma_{18}/(\Gamma_2+\Gamma_3+\Gamma_6)$
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.64 ± 0.18 OUR AVERAGE					
3.9 ± 0.3 ± 0.1		178	35 AKAGI	91B SPEC	
3.45 ± 0.18 ± 0.13		368	36 HEINSON	91 SPEC	
4.0 $^{+1.4}_{-0.9}$		15	SHOCHET	79 SPEC	
4.2 $^{+5.1}_{-2.6}$		3	37 FUKUSHIMA	76 SPEC	
5.8 $^{+2.3}_{-1.5}$		9	38 CARITHERS	73 SPEC	

$\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-)$ Γ_{18}/Γ_{16}
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.64 ± 0.18 OUR AVERAGE					
4.1 ± 0.5		54	INAGAKI	89 SPEC	In AKAGI 91B
2.8 ± 0.3 ± 0.2		87	MATHIAZHA...	89B SPEC	In HEINSON 91
< 1.53	90	0	39 CLARK	71 SPEC	
< 18.	90	0	DARRIULAT	70 SPEC	
< 140.	90	0	FOETH	69 SPEC	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 4.1 ± 0.5 54 INAGAKI 89 SPEC In AKAGI 91B
 2.8 ± 0.3 ± 0.2 87 MATHIAZHA... 89B SPEC In HEINSON 91
 < 1.53 90 0 39 CLARK 71 SPEC
 < 18. 90 0 DARRIULAT 70 SPEC
 < 140. 90 0 FOETH 69 SPEC

³⁵ AKAGI 91B give this number multiplied by the 1990 PDG average for $\Gamma(K_L^0 \rightarrow \pi^+\pi^-)/\Gamma_{total}$.
³⁶ HEINSON 91 give $\Gamma(K_L^0 \rightarrow \mu\mu)/\Gamma_{total}$. We divide out the $\Gamma(K_L^0 \rightarrow \pi^+\pi^-)/\Gamma_{total}$ PDG average which they used.
³⁷ FUKUSHIMA 76 errors are at CL = 90%.
³⁸ CARITHERS 73 errors are at CL = 68%, W.Carithers, (private communication 79).
³⁹ CLARK 71 limit raised from 1.2×10^{-6} by FIELD 74 reanalysis. Not in agreement with subsequent experiments. So not averaged.

$\Gamma(\mu^+\mu^-)/\Gamma_{total}$ Γ_{19}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.28 ± 0.28		1	40 CARROLL	80D SPEC	
< 7.81	90		41 DONALDSON	74 SPEC	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 40 Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.1239.
 41 Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.126.

$\Gamma(e^+e^-)/\Gamma_{total}$ Γ_{20}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-10})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.41	90	0	42 ARISAKA	93B SPEC	
< 1.6	90	1	AKAGI	91 SPEC	
< 5.6	90		INAGAKI	89 SPEC	In AKAGI 91
< 3.2	90		MATHIAZHA...	89B SPEC	In ARISAKA 93B
< 110	90		COUSINS	88 SPEC	
< 45	90		GREENLEE	88 SPEC	Repl. by JASTRZEMSKI 88
< 12	90		JASTRZEM...	88 SPEC	
< 15.7	90		43 CLARK	71 ASPK	
< 1500	90	0	FOETH	69 ASPK	

42 ARISAKA 93B includes all events with < 6 MeV radiated energy.
 43 Possible (but unknown) systematic errors. See note on CLARK 71 $\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-)$ entry.

$\Gamma(e^+e^-)/[\Gamma(\pi^+\pi^-\pi^0) + \Gamma(\pi^\pm\mu^\mp\nu) + \Gamma(\pi^\pm e^\mp\nu)]$ $\Gamma_{20}/(\Gamma_2+\Gamma_3+\Gamma_6)$
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1 ± 0.5 OUR AVERAGE					
9.2 ± 0.5 ± 0.5		1053	BARR	90B NA31	
9.1 ± 0.4 $^{+0.6}_{-0.5}$		919	OHL	90B B845	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 17.4 ± 8.7 4 44 CARROLL 80D SPEC
 < 27 90 0 45 BARMIN 72 HLBC

$\Gamma(e^+e^-)/\Gamma_{total}$ Γ_{21}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
9.1 ± 0.5 OUR AVERAGE					
9.2 ± 0.5 ± 0.5		1053	BARR	90B NA31	
9.1 ± 0.4 $^{+0.6}_{-0.5}$		919	OHL	90B B845	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 17.4 ± 8.7 4 44 CARROLL 80D SPEC
 < 27 90 0 45 BARMIN 72 HLBC
⁴⁴ Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.1239.
⁴⁵ Uses $K_L^0 \rightarrow 3\pi^0$ /total = 0.214.

See key on page 1343

Meson Full Listings

K_L^0

$\Gamma(e^+e^-\gamma\gamma)/\Gamma_{total}$ Γ_{24}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-7})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
6.6 ± 3.2			MORSE	92 B845	$E_\gamma > 5$ MeV

$\Gamma(\pi^+\pi^-e^+e^-)/\Gamma_{total}$ Γ_{23}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	EVTs	DOCUMENT ID	TECN
< 2.5	90	0	BALATS	83 SPEC

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 8.81	90	46	DONALDSON	76 SPEC
< 30			ANIKINA	73 STRC

⁴⁶ Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.126.

$\Gamma(\mu^+\mu^-e^+e^-)/\Gamma_{total}$ Γ_{24}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-6})	CL%	DOCUMENT ID	TECN
< 4.9	90	BALATS	83 SPEC

$\Gamma(e^+e^-e^+e^-)/\Gamma_{total}$ Γ_{25}/Γ
 Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-8})	CL%	EVTs	DOCUMENT ID	TECN
3.9 ± 0.7 OUR AVERAGE				
3.96 ± 0.78 ± 0.32		27	GU	94 E799
6 ± 2 ± 1		18	AKAGI	93 CNTR
3.07 ± 1.25 ± 0.26		6	VAGINS	93 B845
4 ± 3		2	BARR	91 NA31

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 260	90		BALATS	83 SPEC
-------	----	--	--------	---------

$\Gamma(\pi^0\mu^+\mu^-)/\Gamma_{total}$ Γ_{26}/Γ
 Violates CP in leading order. Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-9})	CL%	EVTs	DOCUMENT ID	TECN
< 5.1	90	0	HARRIS	93 E799

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 1200	90	0	⁴⁷ CARRROLL	80D SPEC
< 56600	90	48	DONALDSON	74 SPEC

⁴⁷ Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.1239.
⁴⁸ Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.126.

$\Gamma(\pi^0e^+e^-)/\Gamma_{total}$ Γ_{27}/Γ
 Violates CP in leading order. Direct and indirect CP-violating contributions are expected to be comparable and to dominate the CP-conserving part. Test for $\Delta S = 1$ weak neutral current. Allowed by higher-order electroweak interaction.

VALUE (units 10^{-9})	CL%	EVTs	DOCUMENT ID	TECN
< 4.3	90	0	HARRIS	93B E799
< 7.5	90	0	BARKER	90 E731
< 5.5	90	0	OHL	90 B845

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 40	90		BARR	88 NA31
< 320	90		JASTRZEM...	88 SPEC
< 2300	90	0	⁴⁹ CARROLL	80D SPEC

⁴⁹ Uses $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ /(all K_L^0) decays = 0.1239.

$\Gamma(\pi^0\nu\bar{\nu})/\Gamma_{total}$ Γ_{28}/Γ
 Violates CP in leading order. Test of direct CP violation since the indirect CP-violating and CP-conserving contributions are expected to be suppressed. Test for $\Delta S = 1$ weak neutral current.

VALUE (units 10^{-3})	CL%	EVTs	DOCUMENT ID	TECN
< 0.22	90	0	GRAHAM	92 CNTR

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 7.6	90	50	LITTENBERG	89 RVUE
-------	----	----	------------	---------

⁵⁰ LITTENBERG 89 is from retroactive data analysis of CRONIN 67.

$\Gamma(e^\pm\mu^\mp)/\Gamma_{total}$ Γ_{29}/Γ
 Test of lepton family number conservation.

VALUE (units 10^{-11})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
< 3.3	90	0	51 ARISAKA	93 SPEC	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 3.9	90	0	ARISAKA	93 SPEC	
< 9.4	90	0	AKAGI	91 SPEC	
< 43	90		INAGAKI	89 SPEC	In AKAGI 91
< 22	90		MATHIAZHA...	89 SPEC	
< 190	90		SCHAFFNER	89 SPEC	
< 1100	90		COUSINS	88 SPEC	
< 670	90		GREENLEE	88 SPEC	Repl. by SCHAFFNER 89
< 157	90	52	CLARK	71 ASPK	

⁵¹ This is the combined result of ARISAKA 93 and MATHIAZHAGAN 89.
⁵² Possible (but unknown) systematic errors. See note on CLARK 71 $\Gamma(\mu^+\mu^-)/\Gamma(\pi^+\pi^-)$ entry.

$\Gamma(e^\pm\mu^\mp)/[\Gamma(\pi^+\pi^-\pi^0) + \Gamma(\pi^\pm\mu^\mp\nu) + \Gamma(\pi^\pm e^\mp\nu)]$ $\Gamma_{29}/(\Gamma_2+\Gamma_3+\Gamma_6)$
 Test of lepton family number conservation.

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN
< 0.1	90	BOTT...	67 OSPK
< 0.08	90	FITCH	67 OSPK
< 1.0	90	CARPENTER	66 OSPK
< 10.0		ANIKINA	65 CC

ENERGY DEPENDENCE OF K_L^0 DALITZ PLOT

For discussion, see note on Dalitz plot parameters in the K^\pm section of the Full Listings above. For definitions of a_{ν} , a_t , a_u , and a_y , see the earlier version of the same note in the 1982 edition of this Review published in Physics Letters **111B** 70 (1982).

$|\text{matrix element}|^2 = 1 + gu + hu^2 + jv + kv^2$
 where $u = (s_3 - s_0) / m_\pi^2$ and $v = (s_1 - s_2) / m_\pi^2$

LINEAR COEFFICIENT g FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$


VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.670 ± 0.014 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
0.681 ± 0.024	6499	CHO	77 HBC	
0.620 ± 0.023	4709	PEACH	77 HBC	
0.677 ± 0.010	509k	MESSNER	74 ASPK	$a_y = -0.917 \pm 0.013$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.69 ± 0.07	192	⁵³ BALDO...	75 HLBC	
0.590 ± 0.022	56k	⁵³ BUCHANAN	75 SPEC	$a_u = -0.277 \pm 0.010$
0.619 ± 0.027	20k	^{53,54} BISI	74 ASPK	$a_t = -0.282 \pm 0.011$
0.612 ± 0.032		⁵³ ALEXANDER	73B HBC	
0.73 ± 0.04	3200	⁵³ BRANDENB...	73 HBC	
0.50 ± 0.11	180	⁵³ JAMES	72 HBC	
0.608 ± 0.043	1486	⁵³ KRENZ	72 HLBC	$a_t = -0.277 \pm 0.018$
0.688 ± 0.074	384	⁵³ METCALF	72 ASPK	$a_t = -0.31 \pm 0.03$
0.650 ± 0.012	29k	⁵³ ALBROW	70 ASPK	$a_y = -0.858 \pm 0.015$
0.593 ± 0.022	36k	^{53,55} BUCHANAN	70 SPEC	$a_u = -0.278 \pm 0.010$
0.664 ± 0.056	4400	⁵³ SMITH	70 OSPK	$a_t = -0.306 \pm 0.024$
0.400 ± 0.045	2446	⁵³ BASILE	68B OSPK	$a_t = -0.188 \pm 0.020$
0.649 ± 0.044	1350	⁵³ HOPKINS	67 HBC	$a_t = -0.294 \pm 0.018$
0.428 ± 0.055	1198	⁵³ NEFKENS	67 OSPK	$a_u = -0.204 \pm 0.025$
0.64 ± 0.17	280	⁵³ ANIKINA	66 CC	$a_v = -8.2 \pm 0.9$ -1.3
0.70 ± 0.12	126	⁵³ HAWKINS	66 HBC	$a_v = -8.6 \pm 0.7$
0.32 ± 0.13	66	⁵³ ASTBURY	65 CC	$a_v = -5.5 \pm 1.5$
0.51 ± 0.09	310	⁵³ ASTBURY	65B CC	$a_v = -7.3 \pm 0.6$ -0.8
0.55 ± 0.23	79	⁵³ ADAIR	64 HBC	$a_v = -7.6 \pm 1.7$
0.51 ± 0.20	77	⁵³ LUERS	64 HBC	$a_v = -7.3 \pm 1.6$

⁵³ Quadratic dependence required by some experiments. (See sections on "QUADRATIC COEFFICIENT h " and "QUADRATIC COEFFICIENT k " below.) Correlations prevent us from averaging results of fits not including g , h , and k terms.
⁵⁴ BISI 74 value comes from quadratic fit with quad. term consistent with zero. g error is thus larger than if linear fit were used.
⁵⁵ BUCHANAN 70 result revised by BUCHANAN 75 to include radiative correlations and to use more reliable K_L^0 momentum spectrum of second experiment (had same beam).

WEIGHTED AVERAGE
 0.670 ± 0.014 (Error scaled by 1.6)



Linear coeff. g for $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ matrix element squared

Meson Full Listings

 K_L^0 QUADRATIC COEFFICIENT h FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$

VALUE	EVTS	DOCUMENT ID	TECN
0.079 ± 0.007 OUR AVERAGE			
0.095 ± 0.032	6499	CHO 77	HBC
0.048 ± 0.036	4709	PEACH 77	HBC
0.079 ± 0.007	509k	MESSNER 74	ASPK

• • • We do not use the following data for averages, fits, limits, etc. • • •

-0.011 ± 0.018	29k	56 ALBROW 70	ASPK
0.043 ± 0.052	4400	56 SMITH 70	OSPK

See notes in section "LINEAR COEFFICIENT g FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ | MATRIX ELEMENT" ²⁴ above.

⁵⁶ Quadratic coefficients h and k required by some experiments. (See section on "QUADRATIC COEFFICIENT k " below.) Correlations prevent us from averaging results of fits not including g , h , and k terms.

QUADRATIC COEFFICIENT k FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$

VALUE	EVTS	DOCUMENT ID	TECN
0.0098 ± 0.0018 OUR AVERAGE			
0.024 ± 0.010	6499	CHO 77	HBC
-0.008 ± 0.012	4709	PEACH 77	HBC
0.0097 ± 0.0018	509k	MESSNER 74	ASPK

LINEAR COEFFICIENT J FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ (CP-VIOLATING TERM)

Listed in CP-violation section below.

QUADRATIC COEFFICIENT h FOR $K_L^0 \rightarrow \pi^0\pi^0\pi^0$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN
-3.3 ± 1.1 ± 0.7	5M	57 SOMALWAR 92	E731

⁵⁷ SOMALWAR 92 chose m_{π^+} as normalization to make it compatible with the Particle Data Group $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ definitions.

 K_L^0 FORM FACTORS

For discussion, see note on form factors in the K^\pm section of the Full Listings above.

In the form factor comments, the following symbols are used.

f_+ and f_- are form factors for the vector matrix element.

f_S and f_T refer to the scalar and tensor term.

$f_0 = f_+ + f_- t / (m_K^2 - m_\pi^2)$.

λ_+ , λ_- , and λ_0 are the linear expansion coefficients of f_+ , f_- , and f_0 .

λ_+ refers to the $K_{\mu 3}^0$ value except in the $K_{e 3}^0$ sections.

$d\xi(0)/d\lambda_+$ is the correlation between $\xi(0)$ and λ_+ in $K_{\mu 3}^0$.

$d\lambda_0/d\lambda_+$ is the correlation between λ_0 and λ_+ in $K_{\mu 3}^0$.

t = momentum transfer to the π in units of m_π^2 .

DP = Dalitz plot analysis.

PI = π spectrum analysis.

MU = μ spectrum analysis.

POL = μ polarization analysis.

BR = $K_{\mu 3}^0/K_{e 3}^0$ branching ratio analysis.

E = positron or electron spectrum analysis.

RC = radiative corrections.

 λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN $K_{e 3}^0$ DECAY)

For radiative correction of $K_{e 3}^0$ DP, see GINSBERG 67 and BECHERRAWY 70.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0300 ± 0.0016 OUR AVERAGE				Error includes scale factor of 1.2.
0.0306 ± 0.0034	74k	BIRULEV 81	SPEC	DP
0.025 ± 0.005	12k	58 ENGLER 78b	HBC	DP
0.0348 ± 0.0044	18k	HILL 78	STRC	DP
0.0312 ± 0.0025	500k	GJESDAL 76	SPEC	DP
0.0270 ± 0.0028	25k	BLUMENTHAL 75	SPEC	DP
0.044 ± 0.006	24k	BUCHANAN 75	SPEC	DP
0.040 ± 0.012	2171	WANG 74	OSPK	DP
0.045 ± 0.014	5600	ALBROW 73	ASPK	DP
0.019 ± 0.013	1871	BRANDENB... 73	HBC	PI transv.
0.022 ± 0.014	1910	NEUHOFER 72	ASPK	PI
0.023 ± 0.005	42k	BISI 71	ASPK	DP
0.05 ± 0.01	16k	CHIEN 71	ASPK	DP, no RC
0.02 ± 0.013	1000	ARONSON 68	OSPK	PI
+0.023 ± 0.012	4800	BASILE 68	OSPK	DP, no RC
-0.01 ± 0.02	762	FIRESTONE 67	HBC	DP, no RC
+0.01 ± 0.015	531	KADYK 67	HBC	e, PI, no RC
+0.08 ± 0.10	240	LOWYS 67	FBC	PI
+0.15 ± 0.08	577	FISHER 65	OSPK	DP, no RC
+0.07 ± 0.06	153	LUERS 64	HBC	DP, no RC

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.029 ± 0.005	19k	58 CHO 80	HBC	DP
0.0286 ± 0.0049	26k	BIRULEV 79	SPEC	Repl. by BIRULEV 81
0.032 ± 0.0042	48k	BIRULEV 76	SPEC	Repl. by BIRULEV 81

⁵⁸ ENGLER 78b uses an unique $K_{e 3}^0$ subset of CHO 80 events and is less subject to systematic effects.

 $\xi_B = f_-/f_+$ (determined from $K_{\mu 3}^0$ spectra)

The parameter ξ is redundant with λ_0 below and is not put into the Meson Summary Table.

VALUE	$d\xi(0)/d\lambda_+$	EVTS	DOCUMENT ID	TECN	COMMENT
-0.11 ± 0.09 OUR EVALUATION					From a fit discussed in note on $K_{e 3}^0$ form factors in 1982 edition, PL 111B (April 1982).
-0.10 ± 0.09	-12	150k	59 BIRULEV 81	SPEC	DP
+0.26 ± 0.16	-13	14k	60 CHO 80	HBC	DP
+0.13 ± 0.23	-20	16k	60 HILL 79	STRC	DP
-0.25 ± 0.22	-5.9	32k	61 BUCHANAN 75	SPEC	DP
-0.11 ± 0.07	-17	1.6M	62 DONALDSON 74b	SPEC	DP
-1.00 ± 0.45	-20	1385	63 PEACH 73	HLBC	DP
-1.5 ± 0.7	-28	9086	64 ALBROW 72	ASPK	DP
+1.2 ± 0.8	-18	1341	65 CARPENTER 66	OSPK	DP
+0.50 ± 0.61	unknown	16k	66 DALLY 72	ASPK	DP
-3.9 ± 0.4	3140	67 BASILE 70	OSPK	DP, indep of λ_+	
-0.68 ^{+0.12} _{-0.20}	-26	16k	66 CHIEN 70	ASPK	DP

• • • We do not use the following data for averages, fits, limits, etc. • • •

+0.50 ± 0.61	unknown	16k	66 DALLY 72	ASPK	DP
-3.9 ± 0.4	3140	67 BASILE 70	OSPK	DP, indep of λ_+	
-0.68 ^{+0.12} _{-0.20}	-26	16k	66 CHIEN 70	ASPK	DP

⁵⁹ BIRULEV 81 error, $d\xi(0)/d\lambda_+$ calculated by us from λ_0 , λ_+ , $d\lambda_0/d\lambda_+ = 0$ used.

⁶⁰ HILL 79 and CHO 80 calculated by us from λ_0 , λ_+ , and $d\lambda_0/d\lambda_+$.

⁶¹ BUCHANAN 75 is calculated by us from λ_0 , λ_+ and $d\lambda_0/d\lambda_+$ because their appendix A value -0.20 ± 22 assumes $\xi(t)$ constant, i.e. $\lambda_- = \lambda_+$.

⁶² DONALDSON 74b gives $\xi = -0.11 ± 0.02$ not including systematics. Above error and $d\xi(0)/d\lambda_+$ were calculated by us from λ_0 and λ_+ errors (which include systematics) and $d\lambda_0/d\lambda_+$.

⁶³ PEACH 73 gives $\xi(0) = -0.95 ± 0.45$ for $\lambda_+ = \lambda_- = 0.025$. The above value is for $\lambda_- = 0$. K. Peach, private communication (1974).

⁶⁴ ALBROW 72 fit has λ_- free, gets $\lambda_- = -0.030 ± 0.060$ or $\Lambda = +0.15 ^{+0.17} _{-0.11}$.

⁶⁵ CARPENTER 66 $\xi(0)$ is for $\lambda_+ = 0$. $d\xi(0)/d\lambda_+$ is from figure 9.

⁶⁶ CHIEN 70 errors are statistical only. $d\xi(0)/d\lambda_+$ from figure 4. DALLY 72 is a reanalysis of CHIEN 70. The DALLY 72 result is not compatible with assumption $\lambda_- = 0$ so not included in our fit. The nonzero λ_- value and the relatively large λ_+ value found by DALLY 72 come mainly from a single low t bin (figures 1,2). The (f_+, ξ) correlation was ignored. We estimate from figure 2 that fixing $\lambda_- = 0$ would give $\xi(0) = -1.4 ± 0.3$ and would add 10 to χ^2 . $d\xi(0)/d\lambda_+$ is not given.

⁶⁷ BASILE 70 is incompatible with all other results. Authors suggest that efficiency estimates might be responsible.

 $\xi_B = f_-/f_+$ (determined from $K_{\mu 3}^0/K_{e 3}^0$)

The $K_{\mu 3}^0/K_{e 3}^0$ branching ratio fixes a relationship between $\xi(0)$ and λ_+ . We quote the author's $\xi(0)$ and associated λ_+ but do not average because the λ_+ values differ. The fit result and scale factor given below are not obtained from these ξ_B values. Instead they are obtained directly from the authors $K_{\mu 3}^0/K_{e 3}^0$ branching ratio via the fitted $K_{\mu 3}^0/K_{e 3}^0$ ratio ($\Gamma(\pi^\pm \mu^\mp \nu)/\Gamma(\pi^\pm e^\mp \nu)$). The parameter ξ is redundant with λ_0 below and is not put into the Meson Summary Table.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.11 ± 0.09 OUR EVALUATION				From a fit discussed in note on $K_{e 3}^0$ form factors in 1982 edition, PL 111B (April 1982).
0.5 ± 0.4	6700	BRANDENB... 73	HBC	BR, $\lambda_+ = 0.019 ± 0.013$
-0.08 ± 0.25	1309	68 EVANS 73	HLBC	BR, $\lambda_+ = 0.02$
-0.5 ± 0.5	3548	BASILE 70	OSPK	BR, $\lambda_+ = 0.02$
+0.45 ± 0.28	569	BEILLIERE 69	HLBC	BR, $\lambda_+ = 0$
-0.22 ± 0.30	1309	68 EVANS 69	HLBC	
+0.2 ^{+0.8} _{-1.2}		KULYUKINA 68	CC	BR, $\lambda_+ = 0$
+1.1 ± 1.1	389	ADAIR 64	HBC	BR, $\lambda_+ = 0$
+0.66 ^{+0.9} _{-1.3}		LUERS 64	HBC	BR, $\lambda_+ = 0$

⁶⁸ EVANS 73 replaces EVANS 69.

 $\xi_C = f_-/f_+$ (determined from μ polarization in $K_{\mu 3}^0$)

The μ polarization is a measure of $\xi(t)$. No assumptions on λ_{\pm} necessary, t (weighted by sensitivity to $\xi(t)$) should be specified. In λ_+ , $\xi(0)$ parametrization this is $\xi(0)$ for $\lambda_+ = 0$. $d\xi/d\lambda_+ = \xi t$. For radiative correction to μ polarization in $K_{\mu 3}^0$, see GINSBERG 73. The parameter ξ is redundant with λ_0 below and is not put into the Meson Summary Table.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.11 ± 0.09 OUR EVALUATION				From a fit discussed in note on $K_{e 3}^0$ form factors in 1982 edition, PL 111B (April 1982).
+0.178 ± 0.105	207k	69 CLARK 77	SPEC	POL, $d\xi(0)/d\lambda_+ = +0.68$
-0.385 ± 0.105	2.2M	70 SANDWEISS 73	CNTR	POL, $d\xi(0)/d\lambda_+ = -6.3$
-1.81 ^{+0.50} _{-0.26}		71 LONGO 69	CNTR	POL, $t = 3.3$

• • • We do not use the following data for averages, fits, limits, etc. • • •

-1.6 ± 0.5	638	72 ABRAMS 68b	OSPK	Polarization
-1.2 ± 0.5	2608	72 AUERBACH 66b	OSPK	Polarization

⁶⁹ CLARK 77 $t = +3.80$, $d\xi(0)/d\lambda_+ = \xi(t) = 0.178 \times 3.80 = +0.68$.

⁷⁰ SANDWEISS 73 is for $\lambda_+ = 0$ and $t = 0$.

⁷¹ LONGO 69 $t = 3.3$ calculated from $d\xi(0)/d\lambda_+ = -6.0$ (table 1) divided by $\xi = -1.81$.

⁷² t value not given.

Im(ξ) in $K_{\mu 3}^0$ DECAY (from transverse μ pol.)Test of T reversal invariance.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.007 ± 0.026 OUR AVERAGE				
0.009 ± 0.030	12M	MORSE	80 CNTR	Polarization
0.35 ± 0.30	207k	73 CLARK	77 SPEC	POL, $t=0$
-0.085 ± 0.064	2.2M	74 SANDWEISS	73 CNTR	POL, $t=0$
-0.02 ± 0.08		LONGO	69 CNTR	POL, $t=3.3$
-0.2 ± 0.6		ABRAMS	68B OSPK	Polarization
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.012 ± 0.026		SCHMIDT	79 CNTR	Repl. by MORSE 80
73 CLARK 77 value has additional $\xi(0)$ dependence +0.21Re $[\xi(0)]$.				
74 SANDWEISS 73 value corrected from value quoted in their paper due to new value of Re(ξ). See footnote 4 of SCHMIDT 79.				

 λ_+ (LINEAR ENERGY DEPENDENCE OF f_+ IN $K_{\mu 3}^0$ DECAY)

See also the corresponding entries and notes in section " $\xi_A = f_-/f_+$ " above and section " λ_0 (LINEAR ENERGY DEPENDENCE OF f_0 IN $K_{\mu 3}^0$ DECAY)" below. For radiative correction of $K_{\mu 3}^0$ Dalitz plot see GINSBERG 70 and BECHERAWY 70.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.034 ± 0.005 OUR EVALUATION From a fit discussed in note on $K_{\mu 3}^0$ form factors in 1982 edition, PL 111B (April 1982).				
0.0427 ± 0.0044	150k	BIRULEV	81 SPEC	DP
0.028 ± 0.010	14k	CHO	80 HBC	DP
0.028 ± 0.011	16k	HILL	79 STRC	DP
0.046 ± 0.030	32k	BUCHANAN	75 SPEC	DP
0.030 ± 0.003	1.6M	DONALDSON	74B SPEC	DP
0.085 ± 0.015	9086	ALBROW	72 ASPK	DP
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0337 ± 0.0033	129k	DZHORD...	77 SPEC	Repl. by BIRULEV 81
0.046 ± 0.008	82k	ALBRECHT	74 WIRE	Repl. by BIRULEV 81
0.11 ± 0.04	16k	DALLY	72 ASPK	DP
0.07 ± 0.02	16k	CHIEN	70 ASPK	Repl. by DALLY 72

 λ_0 (LINEAR ENERGY DEPENDENCE OF f_0 IN $K_{\mu 3}^0$ DECAY)

Wherever possible, we have converted the above values of $\xi(0)$ into values of λ_0 using the associated λ_+^{μ} and $d\xi(0)/d\lambda_+$.

VALUE	$d\lambda_0/d\lambda_+$	EVTs	DOCUMENT ID	TECN	COMMENT
0.025 ± 0.006 OUR EVALUATION From a fit discussed in note on $K_{\mu 3}^0$ form factors in 1982 edition, PL 111B (April 1982).					
0.0341 ± 0.0067	unknown	150k	75 BIRULEV	81 SPEC	DP
+0.050 ± 0.008	-0.11	14k	CHO	80 HBC	DP
+0.039 ± 0.010	-0.67	16k	HILL	79 STRC	DP
+0.047 ± 0.009	1.06	207k	76 CLARK	77 SPEC	POL
+0.025 ± 0.019	+0.5	32k	77 BUCHANAN	75 SPEC	DP
+0.019 ± 0.004	-0.47	1.6M	78 DONALDSON	74B SPEC	DP
-0.060 ± 0.038	-0.71	1385	79 PEACH	73 HLBC	DP
-0.018 ± 0.009	+0.49	2.2M	76 SANDWEISS	73 CNTR	POL
-0.043 ± 0.052	-1.39	9086	80 ALBROW	72 ASPK	DP
-0.140 ± 0.043	+0.49		76 LONGO	69 CNTR	POL
-0.08 ± 0.07	-0.54	1371	76 CARPENTER	66 OSPK	DP
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.041 ± 0.008		14k	81 CHO	80 HBC	BR, $\lambda_+ = 0.028$
+0.0485 ± 0.0076		47k	DZHORD...	77 SPEC	In BIRULEV 81
+0.024 ± 0.011		82k	ALBRECHT	74 WIRE	In BIRULEV 81
+0.06 ± 0.03		6700	82 BRANDENB...	73 HBC	BR, $\lambda_+ = 0.019 \pm 0.013$
-0.067 ± 0.227	unknown	16k	83 DALLY	72 ASPK	DP
-0.333 ± 0.034	+1.	3140	84 BASILE	70 OSPK	DP

75 BIRULEV 81 gives $d\lambda_0/d\lambda_+ = -1.5$, giving an unreasonably narrow error ellipse which dominates all other results. We use $d\lambda_0/d\lambda_+ = 0$.

76 λ_0 value is for $\lambda_+ = 0.03$ calculated by us from $\xi(0)$ and $d\xi(0)/d\lambda_+$.

77 BUCHANAN 75 value is from their appendix A and uses only $K_{\mu 3}$ data. $d\lambda_0/d\lambda_+$ was obtained by private communication, C.Buchanan, 1976.

78 DONALDSON 74B $d\lambda_0/d\lambda_+$ obtained from figure 18.

79 PEACH 73 assumes $\lambda_+ = 0.025$. Calculated by us from $\xi(0)$ and $d\xi(0)/d\lambda_+$.

80 ALBROW 72 λ_0 is calculated by us from ξ_A , λ_+ and $d\xi(0)/d\lambda_+$. They give $\lambda_0 = -0.043 \pm 0.039$ for $\lambda_- = 0$. We use our larger calculated error.

81 CHO 80 BR result not independent of their Dalitz plot result.

82 Fit for λ_0 does not include this value but instead includes the $K_{\mu 3}/K_{e 3}$ result from this experiment.

83 DALLY 72 gives $f_0 = 1.20 \pm 0.35$, $\lambda_0 = -0.080 \pm 0.272$, $\lambda_0' = -0.006 \pm 0.045$, but with a different definition of λ_0 . Our quoted λ_0 is his λ_0/f_0 . We cannot calculate true λ_0 error without his (λ_0, f_0) correlations. See also note on DALLY 72 in section ξ_A .

84 BASILE 70 λ_0 is for $\lambda_+ = 0$. Calculated by us from ξ_A with $d\xi(0)/d\lambda_+ = 0$. BASILE 70 is incompatible with all other results. Authors suggest that efficiency estimates might be responsible.

 $|f_5/f_+|$ FOR $K_{e 3}^0$ DECAYRatio of scalar to f_+ couplings.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<0.04					
<0.095	68	25k	BLUMENTHAL75	SPEC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.07	95	18k	HILL	78 STRC	
<0.19	68	48k	BIRULEV	76 SPEC	See also BIRULEV 81
<0.15	95	5600	ALBROW	73 ASPK	
<0.15	68		KULYUKINA	67 CC	

 $|f_T/f_+|$ FOR $K_{e 3}^0$ DECAYRatio of tensor to f_+ couplings.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<0.23					
<0.40	68	25k	BLUMENTHAL75	SPEC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.40	95	18k	HILL	78 STRC	
<0.34	68	48k	BIRULEV	76 SPEC	See also BIRULEV 81
<1.0	95	5600	ALBROW	73 ASPK	
<1.0	68		KULYUKINA	67 CC	

 $|f_T/f_+|$ FOR $K_{\mu 3}^0$ DECAYRatio of tensor to f_+ couplings.

VALUE	DOCUMENT ID	TECN
0.12 ± 0.12	BIRULEV	81 SPEC

 α_{K^*} DECAY FORM FACTOR FOR $K_L \rightarrow e^+ e^- \gamma$

α_{K^*} is the constant in the model of BERGSTROM 83 which measures the relative strength of the vector-vector transition $K_L \rightarrow K^* \gamma$ with $K^* \rightarrow \rho, \omega, \phi \rightarrow \gamma^*$ and the pseudoscalar-pseudoscalar transition $K_L \rightarrow \pi, \eta, \eta' \rightarrow \gamma \gamma^*$.

VALUE	DOCUMENT ID	TECN
-0.28 ± 0.08 OUR AVERAGE		
-0.28 ± 0.13	BARR	90B NA31
-0.280 ^{+0.099} _{-0.090}	OHL	90B B845

DECAY FORM FACTORS FOR $K_L^0 \rightarrow \pi^\pm \pi^0 e^\mp \nu_e$

Given in MAKOFF 93.

NOTE ON CP VIOLATION IN K_L^0 DECAY

(by L. Wolfenstein, Carnegie-Mellon University and T. Trippe, LBL)

Experimentally Measured Parameters

CP violation has been observed in the semi-leptonic decays $K_L^0 \rightarrow \pi^\mp \ell^\pm \nu$ and in the nonleptonic decay $K_L^0 \rightarrow 2\pi$. The experimental numbers that have been measured are [1]

$$\delta = \frac{\Gamma(K_L^0 \rightarrow \pi^- \ell^+ \nu) - \Gamma(K_L^0 \rightarrow \pi^+ \ell^- \nu)}{\Gamma(K_L^0 \rightarrow \pi^- \ell^+ \nu) + \Gamma(K_L^0 \rightarrow \pi^+ \ell^- \nu)} \quad (1a)$$

$$\eta_{+-} = A(K_L^0 \rightarrow \pi^+ \pi^-) / A(K_S^0 \rightarrow \pi^+ \pi^-) = |\eta_{+-}| e^{i\phi_{+-}} \quad (1b)$$

$$\eta_{00} = A(K_L^0 \rightarrow \pi^0 \pi^0) / A(K_S^0 \rightarrow \pi^0 \pi^0) = |\eta_{00}| e^{i\phi_{00}} \quad (1c)$$

Thus there are five real numbers, three magnitudes, and two phases. We list $\delta(\mu)$ for $K_L^0 \rightarrow \pi \mu \nu$ and $\delta(e)$ for $K_L^0 \rightarrow \pi e \nu$ separately and a weighted average δ . Experimentally for the $K_L^0 \rightarrow \pi^0 \pi^0$ decay the quantities directly measured (and also of greatest theoretical interest) are $|\eta_{00}/\eta_{+-}|$ and $\phi_{00} - \phi_{+-}$.

Analysis Based on CPT Invariance [2]

CP violation can occur either in the $K^0 - \bar{K}^0$ mixing or in the decay amplitudes. Assuming CPT invariance, the CP

Meson Full Listings

 K_L^0

violation in the mixing is described by a single parameter ϵ :

$$|K_L^0\rangle = \left[(1 + \epsilon) |K^0\rangle - (1 - \epsilon) |\bar{K}^0\rangle \right] / [2(1 + |\epsilon|^2)]^{1/2} \quad (2a)$$

$$|K_S^0\rangle = \left[(1 + \epsilon) |K^0\rangle + (1 - \epsilon) |\bar{K}^0\rangle \right] / [2(1 + |\epsilon|^2)]^{1/2} . \quad (2b)$$

The decay amplitudes are written

$$\langle I = 0 | T | K^0 \rangle = e^{i\delta_0} A_0 \quad (3a)$$

$$\langle I = 2 | T | K^0 \rangle = e^{i\delta_2} A_2 \quad (3b)$$

where δ_I are the $\pi\pi$ scattering phase shifts at the K^0 mass and I is the isospin of the final state. CP violation is measured by $(\text{Im } A_I / \text{Re } A_I)$. One can then write

$$\eta_{+-} = \epsilon + \epsilon' \quad (4a)$$

$$\eta_{00} = \epsilon - 2\epsilon' \quad (4b)$$

where

$$\epsilon' = \frac{i}{\sqrt{2}} e^{i(\delta_2 - \delta_0)} \frac{\text{Re } A_2}{\text{Re } A_0} \left[\frac{\text{Im } A_2}{\text{Re } A_2} - \frac{\text{Im } A_0}{\text{Re } A_0} \right] \quad (5)$$

neglecting small corrections of order ϵ' times $\text{Re}(A_2/A_0)$. Only two of the three quantities $\epsilon, (\text{Im } A_I / \text{Re } A_I)$ are meaningful because of the ambiguity in defining the phase of K^0 . The standard phase convention due to Wu and Yang [3] sets $\text{Im } A_0 = 0$. A nonzero value of ϵ' would provide definite evidence for CP violation in the decay amplitudes independent of phase convention.

By applying CPT invariance and unitarity it is possible to relate δ to ϵ and to determine the phases of ϵ . If one assumes the $\Delta S = \Delta Q$ rule (see below "Note on the $\Delta S = \Delta Q$ rule in K^0 Decay") the expression for δ becomes

$$\delta = 2\text{Re } \epsilon / (1 + |\epsilon|^2) \approx 2\text{Re } \epsilon . \quad (6)$$

This quantity is independent of phase convention and is seen from Eq. (2) to equal $\langle K_L^0 | K_S^0 \rangle$. The phase of ϵ is given by

$$\phi(\epsilon) \approx \tan^{-1} \frac{(2\Delta m \tau_3)}{\hbar} = 43.59 \pm 0.15^\circ \quad (7a)$$

while Eq. (5) gives

$$\phi(\epsilon') = \delta_2 - \delta_0 + \frac{\pi}{2} \approx 48 \pm 4^\circ . \quad (7b)$$

The approximation in Eq. (7a) depends on the neglect of CP violation in decays other than $K^0 \rightarrow 2\pi$ and is known to be good to a few tenths of a degree. Eq. (7a) is evaluated using the values of the $K_L^0 - K_S^0$ mass difference $\Delta m = (0.5333 \pm 0.0027) \times 10^{10} \hbar s^{-1}$ and the K_S^0 mean life $\tau_3 = (0.8926 \pm 0.0012) \times 10^{-10} s$ from the current edition. The value of the $\pi\pi$ phase shifts is taken from the fit given by Chell and Olsson [4]. The most important point for the analysis is that $\cos[\phi(\epsilon') - \phi(\epsilon)] \simeq 1$. The consequence of this analysis is that only two real quantities

need be measured, the magnitude of ϵ and the value of (ϵ'/ϵ) including its sign. The measured quantity $|\eta_{00}/\eta_{+-}|^2$ which is very close to unity, is given to a good approximation by

$$|\eta_{00}/\eta_{+-}|^2 \approx 1 - 6\text{Re } (\epsilon'/\epsilon) = 1 - 6(\epsilon'/\epsilon) \cos [\phi(\epsilon') - \phi(\epsilon)] . \quad (8)$$

Since the \cos in Eq. (8) is expected theoretically to be very close to unity it is customary to say that $|\eta_{00}/\eta_{+-}|^2$ determines ϵ'/ϵ .

It is possible to use the values of the ϕ_{+-} and $\phi_{00} - \phi_{+-}$ to set limits on CPT violation. [See Tests of Conservation Laws.]

Models

In the superweak model [5] CP violation is restricted to the mass mixing so that to a high degree of accuracy one expects $\epsilon' = 0$. The phase $\phi(\epsilon)$ is given in this model exactly by Eq. (7a) so that this has sometimes been referred to as the superweak phase; however, as noted above, all CPT invariant models give Eq. (7a) as a very good approximation. In the Standard Model CP violation is entirely due to the phase in the Cabibbo-Kobayashi-Maskawa mixing matrix [6](q.v.). Since CP violation occurs in first order in decay amplitudes and in second order in mass-matrix mixing, one expects a significant nonzero value of ϵ' . The calculation is uncertain partly because m_t and V_{td} are not well known and primarily because of the difficulty of estimating hadronic matrix elements [7]. The theoretical results for ϵ'/ϵ in the standard model are generally in the range 3×10^{-4} to 5×10^{-3} , but may be even lower for large values of m_t .

Fitting procedures

We list measurements of $|\eta_{+-}|$, $|\eta_{00}|$, and $|\eta_{00}/\eta_{+-}|$. Independent information on $|\eta_{+-}|$ and $|\eta_{00}|$ can be obtained from measurements of the K_L^0 and K_S^0 lifetimes (τ) and branching ratios (B) to $\pi\pi$, using the relations

$$|\eta_{+-}| = \left[\frac{B(K_L^0 \rightarrow \pi^+\pi^-)}{\tau(K_L^0)} \frac{\tau(K_S^0)}{B(K_S^0 \rightarrow \pi^+\pi^-)} \right]^{1/2} . \quad (9a)$$

$$|\eta_{00}| = \left[\frac{B(K_L^0 \rightarrow \pi^0\pi^0)}{\tau(K_L^0)} \frac{\tau(K_S^0)}{B(K_S^0 \rightarrow \pi^0\pi^0)} \right]^{1/2} . \quad (9b)$$

We approximate a global fit to these independent sources by first performing two independent fits: 1) BRFIT, a fit to the K_L^0 branching ratios, rates, and mean life, and 2) ETAFIT, a fit to the $|\eta_{+-}|$, $|\eta_{00}|$, and $|\eta_{+-}/\eta_{00}|$ measurements. The results from fit 1, along with the K_S^0 values from this edition are used to compute values of $|\eta_{+-}|$ and $|\eta_{00}|$ which are included as measurements in the $|\eta_{00}|$ and $|\eta_{+-}|$ sections with a document ID of BRFIT 94. Thus the fit values of $|\eta_{+-}|$ and $|\eta_{00}|$ given in this edition include both the direct measurements and the results from the branching ratio fit.

The process is reversed in order to include the direct $|\eta|$ measurements in the branching ratio fit. The results from

fit 2 above (before including BRFIT 94 values) are used along with the K_L^0 and K_S^0 mean lives and the $K_S^0 \rightarrow \pi\pi$ branching fractions to compute the K_L^0 branching ratios $\Gamma(K_L^0 \rightarrow \pi^+\pi^-)/\Gamma(\text{total})$ and $\Gamma(K_L^0 \rightarrow \pi^0\pi^0)/\Gamma(K_L^0 \rightarrow \pi^+\pi^-)$. These branching ratio values are included as measurements in the branching ratio section with a document ID of ETAFIT 94. Thus the K_L^0 branching ratio fit values in this edition include the results of direct measurements of $|\eta_{+-}|$, $|\eta_{00}|$, and $|\eta_{00}/\eta_{+-}|$. Details of these fits are given in the 1990 edition of this Review [8].

Since the last edition, results on ϵ'/ϵ have been published by CERN NA31 (BARR 93D) and Fermilab E731 (GIBBONS 93B). These results dominate our fit to ϵ'/ϵ , $|\eta_{+-}|$, $|\eta_{00}|$, and $B(K_L \rightarrow \pi\pi)$. The resulting value $\epsilon'/\epsilon = (1.5 \pm 0.8) \times 10^{-3}$ ($S = 1.8$) has a large scale factor S included in the error, indicating a continuing disagreement in the input data. The CERN NA31 result continues to indicate the presence of direct CP violation with a result which is more than three σ above zero while the Fermilab E731 result is consistent with zero.

The Fermilab E731 experiment also has published recent results on ϕ_{+-} , $\phi_{00}-\phi_{+-}$, and Δm . GIBBONS 93 reports $\Delta m = (0.5286 \pm 0.0028) \times 10^{10} \text{ h s}^{-1}$ assuming $\phi_{+-} = \phi_{00} = \phi_{\text{SW}}$. GIBBONS 93C reports $\Delta m = (0.5257 \pm 0.0049) \times 10^{10} \text{ h s}^{-1}$ when a joint fit with ϕ_{+-} is done. We use this value in our average rather than the GIBBONS 93 value to avoid the ϕ_{SW} constraint. This is so that when we reevaluate ϕ_{+-} experiments using our average Δm , we will avoid biasing ϕ_{+-} toward ϕ_{SW} . The resulting average $\Delta m = (0.5333 \pm 0.0027) \times 10^{10} \text{ h s}^{-1}$ ($S = 1.2$) is $0.0018 \times 10^{10} \text{ h s}^{-1}$ lower than our 1992 average as a result of the lower E731 value. GIBBONS 93 reports $\phi_{+-} = 42.2 \pm 1.4^\circ$ for their Δm and τ_s . We use their Δm and τ_s dependence quoted in NAKADA 93. We reevaluate all ϕ_{+-} measurements for the current Δm and τ_s PDG averages. The resulting value $\phi_{+-} = 44.3 \pm 0.8^\circ$ is two old standard deviations below our 1992 average and is now consistent with the "superweak" phase given in Eq. (7a).

Footnotes and References

* The S values in parentheses are scale factors by which the errors have been increased to account for discrepancies in the data.

1. K. Kleinknecht in *CP Violation* (ed. C. Jarlskog), World Scientific, (1989), p. 41.
2. V. Barmin, *et al.*, Nucl. Phys. **B247**, 293 (1984); L. Wolfenstein, Ann. Rev. Nucl. Sci. **36**, 137 (1986).
3. T.T. Wu and C.N. Yang, Phys. Rev. Lett. **13**, 380 (1964).
4. E. Chell and M.G. Olsson, Phys. Rev. **D48**, 4076 (1993).
5. L. Wolfenstein, Phys. Rev. Lett. **13**, 562 (1964).
6. M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 659 (1973).
7. J.F. Donoghue *et al.*, Phys. Reports **131**, 320 (1986).
8. J.J. Hernandez *et al.*, Phys. Lett. **B239**, 1 (1990).

CP-VIOLATION PARAMETERS IN K_L^0 DECAYS

CHARGE ASYMMETRY IN LEPTONIC DECAYS

Such asymmetry violates CP . It is related to $\text{Re}(\epsilon)$.

$$\delta(\mu) = [\Gamma(\pi^- \mu^+ \nu_\mu) - \Gamma(\pi^+ \mu^- \bar{\nu}_\mu)]/\text{SUM}$$

Only the combined value below is put into the Meson Summary Table.

VALUE (%)	EVTS	DOCUMENT ID	TECN
0.304 ± 0.025 OUR AVERAGE			
0.313 ± 0.029	15M	GEWENIGER 74	ASPK
0.278 ± 0.051	7.7M	PICCIONI 72	ASPK
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.60 ± 0.14	4.1M	MCCARTHY 73	CNTR
0.57 ± 0.17	1M	⁸⁵ PACIOTTI 69	OSPK
0.403 ± 0.134	1M	⁸⁵ DORFAN 67	OSPK

⁸⁵ PACIOTTI 69 is a reanalysis of DORFAN 67 and is corrected for $\mu^+ \mu^-$ range difference in MCCARTHY 72.

$$\delta(e) = [\Gamma(\pi^- e^+ \nu_e) - \Gamma(\pi^+ e^- \bar{\nu}_e)]/\text{SUM}$$

Only the combined value below is put into the Meson Summary Table.

VALUE (%)	EVTS	DOCUMENT ID	TECN
0.333 ± 0.014 OUR AVERAGE			
0.341 ± 0.018	34M	GEWENIGER 74	ASPK
0.318 ± 0.038	40M	FITCH 73	ASPK
0.346 ± 0.033	10M	MARX 70	CNTR
0.246 ± 0.059	10M	⁸⁶ SAAL 69	CNTR
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.36 ± 0.18	600k	ASHFORD 72	ASPK
0.224 ± 0.036	10M	⁸⁶ BENNETT 67	CNTR

⁸⁶ SAAL 69 is a reanalysis of BENNETT 67.

$$\delta = \text{weighted average of } \delta(\mu) \text{ and } \delta(e)$$

(Combination of the above two sections.)

VALUE (%)	EVTS	DOCUMENT ID	TECN	COMMENT
0.327 ± 0.012 OUR AVERAGE				
0.313 ± 0.029	15M	GEWENIGER 74	ASPK	$K_{\mu 3}$
0.341 ± 0.018	34M	GEWENIGER 74	ASPK	$K_{e 3}$
0.318 ± 0.038	40M	FITCH 73	ASPK	$K_{e 3}$
0.333 ± 0.050	33M	WILLIAMS 73	ASPK	$K_{\mu 3} + K_{e 3}$
0.278 ± 0.051	7.7M	PICCIONI 72	ASPK	$K_{\mu 3}$
0.346 ± 0.033	10M	MARX 70	CNTR	$K_{e 3}$
0.246 ± 0.059	10M	SAAL 69	CNTR	$K_{e 3}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.60 ± 0.14	4.1M	MCCARTHY 73	CNTR	$K_{\mu 3}$
0.36 ± 0.18	600k	ASHFORD 72	ASPK	$K_{e 3}$
0.57 ± 0.17	1M	PACIOTTI 69	OSPK	$K_{\mu 3}$

PARAMETERS FOR $K_L^0 \rightarrow 2\pi$ DECAY

$$\eta_{+-} = A(K_L^0 \rightarrow \pi^+\pi^-) / A(K_S^0 \rightarrow \pi^+\pi^-)$$

$$\eta_{00} = A(K_L^0 \rightarrow \pi^0\pi^0) / A(K_S^0 \rightarrow \pi^0\pi^0)$$

The fitted values of $|\eta_{+-}|$ and $|\eta_{00}|$ given below are the results of a fit to $|\eta_{+-}|$, $|\eta_{00}|$, $|\eta_{00}/\eta_{+-}|$, and $\text{Re}(\epsilon'/\epsilon)$. Independent information on $|\eta_{+-}|$ and $|\eta_{00}|$ can be obtained from the fitted values of the $K_L^0 \rightarrow \pi\pi$ and $K_S^0 \rightarrow \pi\pi$ branching ratios and the K_L^0 and K_S^0 lifetimes. This information is included as data in the $|\eta_{+-}|$ and $|\eta_{00}|$ sections with a Document ID "BRFIT." See the "Note on CP Violation in K_L^0 Decay" above for details.

$$|\eta_{00}| = |A(K_L^0 \rightarrow 2\pi^0) / A(K_S^0 \rightarrow 2\pi^0)|$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.259 ± 0.023 OUR FIT				Error includes scale factor of 1.1.
2.12 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.2.
2.084 ± 0.080		⁸⁷ BRFIT	94	
2.33 ± 0.18		CHRISTENS...	79	ASPK
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.71 ± 0.37	56	⁸⁸ WOLFF 71	OSPK	Cu reg., 4 γ 's
2.95 ± 0.63		⁸⁸ CHOLLET 70	OSPK	Cu reg., 4 γ 's

⁸⁷ This BRFIT value is computed from fitted values of the K_L^0 and K_S^0 lifetimes and branching fractions to $\pi\pi$. See the discussion in the "Note on CP violation in K_L^0 decay."

⁸⁸ CHOLLET 70 gives $|\eta_{00}| = (1.23 \pm 0.24) \times (\text{regeneration amplitude, } 2 \text{ GeV}/c \text{ Cu})/10000\text{mb}$. WOLFF 71 gives $|\eta_{00}| = (1.13 \pm 0.12) \times (\text{regeneration amplitude, } 2 \text{ GeV}/c \text{ Cu})/10000\text{mb}$. We compute both $|\eta_{00}|$ values for (regeneration amplitude, 2 GeV/c Cu) = 24 ± 2mb. This regeneration amplitude results from averaging over FAISSNER 69, extrapolated using optical-model calculations of Bohm *et al.*, Physics Letters **27B** 594 (1968) and the data of BALATS 71. (From H. Faissner, private communication).

Meson Full Listings

K_L^0

$$|\eta_{+-}| = |A(K_L^0 \rightarrow \pi^+\pi^-) / A(K_S^0 \rightarrow \pi^+\pi^-)|$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
2.269 ± 0.023 OUR FIT				Error includes scale factor of 1.1.
2.281 ± 0.022 OUR AVERAGE				

2.266 ± 0.030		89 BRFIT	94	
2.32 ± 0.14 ± 0.03	10 ⁵	ADLER	92B SPEC	K^0, \bar{K}^0 asymm.
2.27 ± 0.12		CHRISTENS...	79B ASPK	
2.30 ± 0.035		GEWENIGER	74B ASPK	

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.28 ± 0.06	1687	90 COUPAL	85 SPEC	$P(K)=70$ GeV/c
2.09 ± 0.02		91 ARONSON	82B SPEC	$E=30-110$ GeV

⁸⁹ This BRFIT value is computed from fitted values of the K_L^0 and K_S^0 lifetimes and branching fractions to $\pi\pi$. See the discussion in the "Note on CP violation in K_L^0 decay."

⁹⁰ COUPAL 85 concludes: no energy dependence of $|\eta_{+-}|$, because their value is consistent with above values which occur at lower energies. Not independent of COUPAL 85 $\Gamma(\pi^+\pi^-)/\Gamma(\pi\ell\nu)$ measurement. Enters $|\eta_{+-}|$ via BRFIT value. In editions prior to 1990, this measurement was erroneously also included in our $|\eta_{+-}|$ average and fit. We thank H. Wahl (WAHL 89) for informing us.

⁹¹ ARONSON 82B find that $|\eta_{+-}|$ may depend on the kaon energy.

$$|\eta_{00}/\eta_{+-}|$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.9955 ± 0.0023 OUR FIT				Error includes scale factor of 1.8.
0.9930 ± 0.0020 OUR AVERAGE				

0.9931 ± 0.0020		92,93 BARR	93D NA31	
0.9904 ± 0.0084 ± 0.0036		94 WOODS	88 E731	

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.9939 ± 0.0013 ± 0.0015	1M	92 BARR	93D NA31	
0.9899 ± 0.0020 ± 0.0025		92 BURKHARDT	88 NA31	
1.014 ± 0.016 ± 0.007	3152	BERNSTEIN	85B SPEC	
0.995 ± 0.025	1122	BLACK	85 SPEC	
1.00 ± 0.09		95 CHRISTENS...	79 ASPK	
1.03 ± 0.07	124	BANNER	72 OSPK	
1.00 ± 0.06	167	HOLDER	72 ASPK	

⁹² This is the square root of the ratio R given by BURKHARDT 88 and BARR 93D.

⁹³ This is the combined results from BARR 93D and BURKHARDT 88, taking into account a common systematic uncertainty of 0.0014.

⁹⁴ We calculate $|\eta_{00}/\eta_{+-}| = 1 - 3(\epsilon'/\epsilon)$ from WOODS 88 (ϵ'/ϵ) value.

⁹⁵ Not independent of $|\eta_{+-}|$ and $|\eta_{00}|$ values which are included in fit.

$$\epsilon'/\epsilon \approx \text{Re}(\epsilon'/\epsilon) = (1 - |\eta_{00}/\eta_{+-}|)/3$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.5 ± 0.8 OUR FIT				Error includes scale factor of 1.8.
1.5 ± 0.8 OUR AVERAGE				Error Includes scale factor of 1.8. See the ideogram below.

2.3 ± 0.65		96 BARR	93D NA31	
0.74 ± 0.52 ± 0.29	>5E5	GIBBONS	93B E731	
3.2 ± 2.8 ± 1.2		97 WOODS	88 E731	


• • • We do not use the following data for averages, fits, limits, etc. • • •

2.0 ± 0.7	1M	97 BARR	93D NA31	
-0.4 ± 1.4 ± 0.6		PATTERSON	90 E731	in GIBBONS 93B
3.3 ± 1.1		97 BURKHARDT	88 NA31	

⁹⁶ This is the combined results from BARR 93D and BURKHARDT 88, taking into account their common systematic uncertainty.

⁹⁷ These values are derived from $|\eta_{00}/\eta_{+-}|$ measurements. They enter the average in this section but enter the fit via the $|\eta_{00}/\eta_{+-}|$ section only.

WEIGHTED AVERAGE
1.5±0.8 (Error scaled by 1.8)



$$\epsilon'/\epsilon \approx \text{Re}(\epsilon'/\epsilon) = (1 - |\eta_{00}/\eta_{+-}|)/3$$

ϕ_{+-} , PHASE OF η_{+-}

The dependence of the phase on the $m_{K_L^0} - m_{K_S^0}$ is given for each experiment in the

comments below, where $\Delta(m)$ is (mass difference/ \hbar) in units 10^{10} s^{-1} and τ_S is the K_S^0 mean life in units 10^{-10} s . We have evaluated these mass dependences using our 1994 values, $\Delta(m) = 0.5333 \pm 0.0027$, $\tau_S = 0.8926 \pm 0.0012$ to obtain the values and average quoted below. We also give the regeneration phase ϕ_f in the comments below.

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
44.3 ± 0.8 OUR FIT				
44.1 ± 0.9 OUR AVERAGE				

42.9 ± 1.2		98 GIBBONS	93 E731	
42.3 ± 4.4 ± 1.4	10 ⁵	99 ADLER	92B SPEC	K^0, \bar{K}^0 asymm.
46.0 ± 2.2		100 CAROSI	90 NA31	
45.2 ± 2.9		101 CARITHERS	75 SPEC	C regenerator
45.6 ± 1.8		102 GEWENIGER	74B ASPK	Vacuum regen.

• • • We do not use the following data for averages, fits, limits, etc. • • •

47.7 ± 2.0 ± 0.9		103 KARLSSON	90 E731	
35.3 ± 3.9		104 ARONSON	82B SPEC	
41.7 ± 3.5		CHRISTENS...	79B ASPK	
36.2 ± 6.1		105 CARNEGIE	72 ASPK	Cu regenerator
37 ± 12		106 BALATS	71 OSPK	Cu regenerator
40 ± 4		107 JENSEN	70 ASPK	Vacuum regen.
34 ± 10		108 BENNETT	69 CNTR	Cu regenerator
44 ± 12		109 BOHM	69B OSPK	Vacuum regen.
45 ± 7		110 FAISSNER	69 ASPK	Cu regenerator
51 ± 11		111 BENNETT	68B CNTR	Cu reg. uses
70 ± 21		112 BOTT...	67B OSPK	C regenerator
25 ± 35		112 MISCHKE	67 OSPK	Cu regenerator
30 ± 45		112 FIRESTONE	66 HBC	
45 ± 50		112 FITCH	65 OSPK	Be regenerator

⁹⁸ GIBBONS 93 measures $\phi_{+-} - \phi_f$ and calculates the regeneration phase ϕ_f from the power law momentum dependence of the regeneration amplitude using analyticity. An error of 0.6° is included for possible uncertainties in the regeneration phase. They find $\phi_{+-} = 42.21 \pm 0.9 + 189 [\Delta(m) - 0.5286] - 460 [\tau_S - 0.8922]^\circ$. B. Winstein, private communication, reported in NAKADA 93. GIBBONS 93 reports $\phi_{+-} (42.2 \pm 1.4)^\circ$ for their $\Delta(m)$ and τ_S .

⁹⁹ ADLER 92B quote separately two systematic errors: ± 0.4 from their experiment and ± 1.0 degrees due to the uncertainty in the value of $\Delta(m)$.

¹⁰⁰ CAROSI 90 $\phi_{+-} = 46.9 \pm 1.4 \pm 0.7 + 579 [\Delta(m) - 0.5351] + 303 [\tau_S - 0.8922]^\circ$.

¹⁰¹ CARITHERS 75 $\phi_{+-} = (45.5 \pm 2.8) + 224 [\Delta(m) - 0.5348]^\circ$. $\phi_f = -40.9 \pm 2.6^\circ$.

¹⁰² GEWENIGER 74B $\phi_{+-} = (49.4 \pm 1.0) + 565 [\Delta(m) - 0.540]^\circ$.

¹⁰³ KARLSSON 90 systematic error does not include regeneration phase uncertainty.

¹⁰⁴ ARONSON 82 find that ϕ_{+-} may depend on the kaon energy.

¹⁰⁵ CARNEGIE 72 ϕ_{+-} is insensitive to $\Delta(m)$. $\phi_f = -56.2 \pm 5.2^\circ$.

¹⁰⁶ BALATS 71 $\phi_{+-} = (39.0 \pm 12.0) + 198 [\Delta(m) - 0.544]^\circ$. $\phi_f = -43.0 \pm 4.0^\circ$.

¹⁰⁷ JENSEN 70 $\phi_{+-} = (42.4 \pm 4.0) + 576 [\Delta(m) - 0.538]^\circ$.

¹⁰⁸ BENNETT 69 uses measurement of $(\phi_{+-}) - (\phi_f)$ of ALFF-STEINBERGER 66B. BENNETT 69 $\phi_{+-} = (34.9 \pm 10.0) + 69 [\Delta(m) - 0.544]^\circ$. $\phi_f = -49.9 \pm 5.4^\circ$.

¹⁰⁹ BOHM 69B $\phi_{+-} = (41.0 \pm 12.0) + 479 [\Delta(m) - 0.526]^\circ$.

¹¹⁰ FAISSNER 69 error enlarged to include error in regenerator phase. FAISSNER 69 $\phi_{+-} = (49.3 \pm 7.4) + 205 [\Delta(m) - 0.555]^\circ$. $\phi_f = -42.7 \pm 5.0^\circ$.

¹¹¹ BENNETT 69 is a re-evaluation of BENNETT 68B.

¹¹² Old experiments with large errors not included in average.

ϕ_{00} , PHASE OF η_{00}

See comment in ϕ_{+-} header above for treatment of $\Delta(m)$ and τ_S dependence.

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
43.3 ± 1.3 OUR FIT				
46.2 ± 2.8				

• • • We do not use the following data for averages, fits, limits, etc. • • •

47.4 ± 1.4 ± 0.9		114 KARLSSON	90 E731	
55.7 ± 5.8		CHRISTENS...	79 ASPK	
38.0 ± 25.0	56	115 WOLFF	71 OSPK	Cu reg., 4 γ 's
51.0 ± 30.0		116 CHOLLET	70 OSPK	Cu reg., 4 γ 's
first quadrant preferred		GOBBI	69B OSPK	

¹¹³ CAROSI 90 $\phi_{00} = 47.1 \pm 2.1 \pm 1.0 + 579 [\Delta(m) - 0.5351] + 252 [\tau_S - 0.8922]^\circ$.

¹¹⁴ KARLSSON 90 systematic error does not include regeneration phase uncertainty.

¹¹⁵ WOLFF 71 uses regenerator phase $\phi_f = -48.2 \pm 3.5^\circ$.

¹¹⁶ CHOLLET 70 uses regenerator phase $\phi_f = -46.5 \pm 4.4^\circ$.

PHASE DIFFERENCE $\phi_{00} - \phi_{+-}$

Test of CPT.

VALUE (°)	DOCUMENT ID	TECN
-1.0 ± 1.0 OUR FIT		
-1.2 ± 1.0 OUR AVERAGE		

-1.6 ± 1.2	117 GIBBONS	93 E731
0.2 ± 2.6 ± 1.2	118 CAROSI	90 NA31
-0.3 ± 2.4 ± 1.2	KARLSSON	90 E731

• • • We do not use the following data for averages, fits, limits, etc. • • •

12.6 ± 6.2	118 CHRISTENS...	79 ASPK
7.6 ± 18.0	119 BARBIELLINI	73 ASPK

¹¹⁷ GIBBONS 93 give detailed dependence of systematic error on lifetime (see the section on the K_S^0 mean life) and mass difference (see the section on $m_{K_L^0} - m_{K_S^0}$).

¹¹⁸ Not independent of ϕ_{+-} and ϕ_{00} values. This is taken into account in our fitting procedure.

¹¹⁹ Independent of regenerator mechanism, $\Delta(m)$, and lifetimes.

See key on page 1343

Meson Full Listings

K_L^0

CHARGE ASYMMETRY IN $\pi^+\pi^-\pi^0$ DECAYS

CHARGE ASYMMETRY J FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$

Defined at beginning of section "LINEAR COEFFICIENT g FOR $K_L^0 \rightarrow \pi^+\pi^-\pi^0$ above. Such asymmetry violates CP . See also note on Dalitz plot parameters in K^{\pm} section and note on CP violation in K_L^0 decay above.

VALUE	EVTS	DOCUMENT ID	TECN
0.0011 ± 0.0008 OUR AVERAGE			
0.001 ± 0.011	6499	CHO	77
-0.001 ± 0.003	4709	PEACH	77
0.0013 ± 0.0009	3M	SCRIBANO	70
0.0 ± 0.017	4400	SMITH	70 OSPK
0.001 ± 0.004	238k	BLANPIED	68

PARAMETERS for $K_L^0 \rightarrow \pi^+\pi^-\gamma$ DECAY

$$|\eta_{+-\gamma}| = |A(K_L^0 \rightarrow \pi^+\pi^-\gamma)/A(K_S^0 \rightarrow \pi^+\pi^-\gamma)|$$

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN
2.15 ± 0.26 ± 0.20	3671	RAMBERG	93B E731

$\phi_{+-\gamma}$ = phase of $\eta_{+-\gamma}$

VALUE (°)	EVTS	DOCUMENT ID	TECN
72 ± 23 ± 17	3671	RAMBERG	93B E731

$|\epsilon'_{+-\gamma}|/\epsilon$

VALUE	CL%	EVTS	DOCUMENT ID	TECN
< 0.3	90	3671	120 RAMBERG	93B E731

120 RAMBERG 93B limit on $|\epsilon'_{+-\gamma}|/\epsilon$ assumes that any difference between η_{+-} and $\eta_{+-\gamma}$ is due to direct CP violation.

NOTE ON $\Delta S = \Delta Q$ IN K^0 DECAYS

The relative amount of $\Delta S \neq \Delta Q$ component present is measured by the parameter x , defined as

$$x = A(\bar{K}^0 \rightarrow \pi^-\ell^+\nu)/A(K^0 \rightarrow \pi^-\ell^+\nu)$$

We list $Re\{x\}$ and $Im\{x\}$ for K_{e3} and $K_{\mu 3}$ combined.


$$x = A(\bar{K}^0 \rightarrow \pi^-\ell^+\nu)/A(K^0 \rightarrow \pi^-\ell^+\nu) = A(\Delta S = -\Delta Q)/A(\Delta S = \Delta Q)$$

REAL PART OF x

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.006 ± 0.018 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
0.10 +0.18 -0.19	79	SMITH	75B WIRE	$\pi^-p \rightarrow K^0\Lambda$
0.04 ± 0.03	4724	NIEBERGALL	74 ASPK	$K^+p \rightarrow K^0p\pi^+$
-0.008 ± 0.044	1757	FACKLER	73 OSPK	K_{e3} from K^0
-0.03 ± 0.07	1367	HART	73 OSPK	K_{e3} from $K^0\Lambda$
-0.070 ± 0.036	1079	MALLARY	73 OSPK	K_{e3} from $K^0\Lambda X$
0.03 ± 0.06	410	121 BURGUN	72 HBC	$K^+p \rightarrow K^0p\pi^+$
-0.05 ± 0.09	442	122 GRAHAM	72 OSPK	$\pi^-p \rightarrow K^0\Lambda$
0.26 +0.10 -0.14	126	MANN	72 HBC	$K^-p \rightarrow n\bar{K}^0$
0.25 +0.07 -0.09	252	WEBBER	71 HBC	$K^-p \rightarrow n\bar{K}^0$
0.12 ± 0.09	215	123 CHO	70 DBC	$K^+d \rightarrow K^0pp$
-0.020 ± 0.025	124	BENNETT	69 CNTR	Charge asym + Cu regen.
0.09 +0.14 -0.16	686	LITTENBERG	69 OSPK	$K^+n \rightarrow K^0p$
0.09 +0.07 -0.09	121	JAMES	68 HBC	$\bar{p}p$
0.17 +0.16 -0.35	116	FELDMAN	67B OSPK	$\pi^-p \rightarrow K^0\Lambda$
0.035 +0.11 -0.13	196	AUBERT	65 HLBC	K^+ charge exchange
0.06 +0.18 -0.44	152	125 BALDO...	65 HLBC	K^+ charge exchange
-0.08 +0.16 -0.28	109	126 FRANZINI	65 HBC	$\bar{p}p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.04 +0.10 -0.13	100	122 GRAHAM	72 OSPK	$K_{\mu 3}$ from $K^0\Lambda$
-0.13 ± 0.11	342	122 MANTSCH	72 OSPK	K_{e3} from $K^0\Lambda$
0.04 +0.07 -0.08	222	121 BURGUN	71 HBC	$K^+p \rightarrow K^0p\pi^+$
0.03 ± 0.03	124	BENNETT	68 CNTR	
0.17 ± 0.10	335	123 HILL	67 DBC	$K^+d \rightarrow K^0pp$

121 BURGUN 72 is a final result which includes BURGUN 71.
 122 First GRAHAM 72 value is second GRAHAM 72 value combined with MANTSCH 72.
 123 CHO 70 is analysis of unambiguous events in new data and HILL 67.
 124 BENNETT 69 is a reanalysis of BENNETT 68.
 125 BALDO-CEOLIN 65 gives x and θ converted by us to $Re(x)$ and $Im(x)$.
 126 FRANZINI 65 gives x and θ for $Re(x)$ and $Im(x)$. See SCHMIDT 67.

WEIGHTED AVERAGE
0.006 ± 0.018 (Error scaled by 1.3)



IMAGINARY PART OF x

Assumes $m_{K_L^0} - m_{K_S^0}$ positive. See Listings above.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.003 ± 0.026 OUR AVERAGE				Error includes scale factor of 1.2.
-0.10 +0.16 -0.19	79	SMITH	75B WIRE	$\pi^-p \rightarrow K^0\Lambda$
-0.06 ± 0.05	4724	NIEBERGALL	74 ASPK	$K^+p \rightarrow K^0p\pi^+$
-0.017 ± 0.060	1757	FACKLER	73 OSPK	K_{e3} from K^0
0.09 ± 0.07	1367	HART	73 OSPK	K_{e3} from $K^0\Lambda$
0.107 +0.092 -0.074	1079	MALLARY	73 OSPK	K_{e3} from $K^0\Lambda X$
0.07 +0.06 -0.07	410	127 BURGUN	72 HBC	$K^+p \rightarrow K^0p\pi^+$
0.05 ± 0.13	442	128 GRAHAM	72 OSPK	$\pi^-p \rightarrow K^0\Lambda$
0.21 +0.15 -0.12	126	MANN	72 HBC	$K^-p \rightarrow n\bar{K}^0$
0.0 ± 0.08	252	WEBBER	71 HBC	$K^-p \rightarrow n\bar{K}^0$
-0.08 ± 0.07	215	129 CHO	70 DBC	$K^+d \rightarrow K^0pp$
-0.11 +0.10 -0.11	686	LITTENBERG	69 OSPK	$K^+n \rightarrow K^0p$
+0.22 +0.37 -0.29	121	JAMES	68 HBC	$\bar{p}p$
0.0 ± 0.25	116	FELDMAN	67B OSPK	$\pi^-p \rightarrow K^0\Lambda$
-0.21 +0.11 -0.15	196	AUBERT	65 HLBC	K^+ charge exchange
-0.44 +0.32 -0.19	152	130 BALDO...	65 HLBC	K^+ charge exchange
+0.24 +0.40 -0.30	109	131 FRANZINI	65 HBC	$\bar{p}p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.12 +0.17 -0.16	100	128 GRAHAM	72 OSPK	$K_{\mu 3}$ from $K^0\Lambda$
-0.04 ± 0.16	342	128 MANTSCH	72 OSPK	K_{e3} from $K^0\Lambda$
0.12 +0.08 -0.09	222	127 BURGUN	71 HBC	$K^+p \rightarrow K^0p\pi^+$
-0.20 ± 0.10	335	129 HILL	67 DBC	$K^+d \rightarrow K^0pp$
127 BURGUN 72 is a final result which includes BURGUN 71. 128 First GRAHAM 72 value is second GRAHAM 72 value combined with MANTSCH 72. 129 Footnote 10 of HILL 67 should read +0.58, not -0.58 (private communication) CHO 70 is analysis of unambiguous events in new data and HILL 67. 130 BALDO-CEOLIN 65 gives x and θ converted by us to $Re(x)$ and $Im(x)$. 131 FRANZINI 65 gives x and θ for $Re(x)$ and $Im(x)$. See SCHMIDT 67.				

K_L^0 REFERENCES

BARR 94 PL B (to be pub.) +Buchholz+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)
 Search for the decay $K_L^0 \rightarrow \pi^0\pi^0\gamma$
 BRFIT 94 RPP
 ETAFIT 94 RPP
 GU 94 PRL 72 3000 + (RUTG, UCLA, EFI, COLO, ELMT, FNAL, ILL, OSAK)
 AKAGI 93 PR D47 R2644 +Fukuhsia, Hemmi+ (TOHOK, TOKY, KYOT, KEK)
 ARISAKA 93 PRL 70 1049 +Auerbach, Axelrod, Betz, Biery+ (BNL E791 Collab.)
 ARISAKA 93B PRL 71 3910 +Auerbach, Axelrod, Betz, Biery+ (BNL E791 Collab.)
 BARR 93D PL B317 233 +Buchholz+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)
 GIBBONS 93 PRL 70 1199 +Barker, Briere, Makoff+ (FNAL E731 Collab.)
 GIBBONS 93B PRL 70 1203 +Barker, Briere, Makoff+ (FNAL E731 Collab.)
 GIBBONS 93C Thesis +Briere, Cheu, Makoff, McFarlane+ (FNAL E799 Collab.)
 HARRIS 93 PRL 71 3914 +Briere, Cheu, Makoff, McFarlane+ (FNAL E799 Collab.)
 HARRIS 93B PRL 71 3918 +Briere, Cheu, Makoff, McFarlane+ (FNAL E799 Collab.)
 MAKOFF 93 PRL 70 1591 +Barker, Briere, Gibbons+ (FNAL E731 Collab.)
 NAKADA 93 Lepton Photon Conf. (PSI)
 AIP Conf. Proc. 302, p. 425
 RAMBERG 93 PRL 70 2525 +Bock, Coleman, Enagonio, Hsiung+ (FNAL E731 Collab.)
 RAMBERG 93B PRL 70 2529 +Bock, Coleman, Enagonio, Hsiung+ (FNAL E731 Collab.)
 VAGINS 93 PRL 71 35 +Adair, Greenlee, Kasha, Mannelli+ (BNL E845 Collab.)
 ADLER 92B PL B286 180 +Alhalel, Angelopoulos, Apostolakis+ (CPLEAR Collab.)
 Also 92 SJNP 55 840 Adler, Alhalel, Angelopoulos+ (CPLEAR Collab.)
 BARR 92 PL B284 440 +Buchholz+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)
 GRAHAM 92 PL B295 159 +Barker, Briere, Gibbons, Makoff+ (FNAL E731 Collab.)
 MORSE 92 PR D45 36 +Leipuner, Larsen, Jastrzembski+ (BNL, YALE, VASS)
 PDG 92 PR D45, 1 June, Part II Hikasa, Barnett, Stone+ (KEK, LBL, BOST+)
 SOMALWAR 92 PRL 68 2580 +Barker, Briere, Gibbons+ (FNAL E731 Collab.)

Meson Full Listings

K⁰_L

AKAGI	91	PRL 67 2614	+Fukuhisa, Hemmi+ (TOHOK, TOKY, KYOT, KEK)	MCCARTHY	73	PR D7 687	+Brewer, Budnitz, Entis, Graven, Miller+ (LBL)
AKAGI	91B	PRL 67 2618	+Fukuhisa, Hemmi+ (TOHOK, TOKY, KYOT, KEK)	Also	72	PL 42B 291	+McCarthy, Brewer, Budnitz, Entis, Graven+ (LBL)
BARR	99	PRL 859 399	+Carosi+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)	Also	71	LBL-550 Thesis	McCarthy (LBL)
HEINSON	91	PR D44 R1	+ (UCI, UCLA, LANL, PENN, STAN, TEMP, TEXA++)	MESSNER	73	PRL 30 876	+Morse, Nauenberg, Hitlin+ (COLO, SLAC, UCSC)
PAPADIMITR...	91	PR D44 R573	Padamitriou, Barker, Briere+ (FNAL E731 Collab.)	PEACH	73	PL 43B 441	+Evans, Muir, Hopkins, Krenz (EDIN, CERN, AACH)
BARKER	90	PR D41 3546	+Briere, Gibbons, Makoff+ (FNAL E731 Collab.)	SANDWEISS	73	PRL 30 1002	+Sunderland, Turner, Willis, Keller (YALE, ANL)
Also	88	PRL 61 2661	Gibbons, Papadimitriou+ (FNAL E731 Collab.)	WILLIAMS	73	PRL 31 1521	+Larsen, Leipuner, Sapp, Sessoms+ (BNL, YALE)
BARR	90B	PL B240 283	+Carosi+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)	ALBROW	72	NP B44 1	+Aston, Barber, Bird, Ellison+ (MCHS, DARE)
BARR	90C	PL B242 523	+Carosi+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)	ASHFORD	72	PL 38B 47	+Brown, Masek, Maung, Miller, Ruderman+ (UCSD)
CAROSI	90	PL B237 303	+Clarke+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)	BANNER	72	PRL 28 1597	+Cronin, Hoffman, Knapp, Shochet (PRIN)
KARLSSON	90	PRL 64 2976	+Golin, Okamoto, Tschihart, Barker+ (FNAL E731 Collab.)	BANNER	72B	PRL 29 157	+Cronin, Hoffman, Knapp, Shochet (PRIN)
OHL	90	PRL 64 2755	+Adair, Greenlee, Kasha, Mannelli+ (BNL E845 Collab.)	BARMIN	72	SJNP 15 636	+Davidenko, Demidov, Dolgolenko+ (ITEP)
OHL	90B	PRL 65 1407	+Adair, Greenlee, Kasha, Mannelli+ (BNL E845 Collab.)	BARMIN	72B	SJNP 15 638	+Barylov, Davidenko, Demidov+ (ITEP)
PATTERSON	90	PRL 64 1491	+Barker+ (FNAL E731 Collab.)	BURGUN	72	NP B50 194	+Lesquoy, Muller, Pauli+ (SACL, CERN, OSLO)
INAGAKI	89	PR D40 1712	+Kobayashi, Sato, Shinkawa+ (KEK, TOKY, KYOT)	CARNEGIE	72	PR D6 2335	+Cester, Fitch, Strovink, Sulak (PRIN)
LITTENBERG	89	PR D39 3322	Mathiazhagan+ (UCI, UCLA, LANL, PENN, STAN++)	DALLY	72	PL 41B 647	+Innocenti, Seppi+ (SLAC, JHU, UCLA)
MATHIAZHA...	89	PRL 63 2181	Mathiazhagan+ (UCI, UCLA, LANL, PENN, STAN++)	Also	70	PL 33B 627	Chien, Cox, Ettinger+ (JHU, SLAC, UCLA)
MATHIAZHA...	89B	PRL 63 2185	Padamitriou, Gibbons, Patterson+ (FNAL E731 Collab.)	Also	71	PL 35B 261	Chien, Cox, Ettinger+ (JHU, SLAC, UCLA)
PAPADIMITR...	89	PR D39 390	+Greenlee, Kasha, Mannelli, Ohi+ (YALE, BNL)	GRAHAM	72	NC 9A 166	+Abashian, Jones, Mantsch, Orr+ (ILL, NEAS)
SCHAFFNER	89	PR D39 990	+Clarke+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)	HOLDER	72	PL 40B 141	+Rademacher, Staudt+ (AACH, CERN, TORI)
WAHL	88	CERN-EP/89-86, H. Wahl	+Kingsberg+ (UCLA, LASL, PENN, STAN, TEMP, WILL)	JAMES	72	NP B49 1	+Montanet, Paul, Saelter+ (CERN, SACL, OSLO)
BARR	88	PL B214 303	+Kasha, Mannelli, Mannelli+ (YALE, BNL)	UNC 4 213	72	PL 213 1	+Hopkins, Evans, Muir, Peach (AACH, CERN, EDIN)
BURKHARDT	88	PL B206 169	+Jastrzembski, Larsen, Leipuner, Morse+ (BNL, YALE)	MANN	72	PR D6 137	+Kofler, Meisner, Hertzbach+ (MASA, BNL, YALE)
COUSINS	88	PR D38 214	+Hishikawa, Patterson, Wah, Weinstein+ (FNAL E731 Collab.)	MANTSCH	72	NC 9A 160	+Abashian, Graham, Jones, Orr+ (ILL, NEAS)
GREENLEE	88	PRL 60 893	+ (CERN, EDIN, MANZ, LALO, PISA, SIEG)	MCCARTHY	72	PL 42B 291	+Brewer, Budnitz, Entis, Graven+ (CERN, IPN, WIEN)
JASTRZEMSKI...	88	PRL 60 1695	+Berenstein, Bock+ (BNL, CHIC, STAN, WISC)	METCALF	72	PL 40B 703	+Neuhof, Budnitz, Entis, Graven+ (CERN, IPN, WIEN)
WOODS	88	PRL 60 1695	+Aronson, Bernstein+ (BNL, CHIC, STAN, WISC)	NEUHOFFER	72	PL 41B 642	+Niebergall, Regler, Seppi+ (CERN, ORSAY, VIEN)
BURKHARDT	87	PL B199 139	+Aronson, Bernstein, Porter+ (CERN, CIT++)	PICCIONI	72	PR 29 1412	+Coombes, Donaldson, Dorfan, Fryberger+ (SLAC)
ARONSON	86	PR D33 3180	+Bock, Carlsmith, Coupal+ (CHIC, SACL)	Also	74	PR D9 2939	Johnston, Donaldson+ (SLAC, UCSC, COLO)
Also	82	PRL 48 1078	+Berenstein, Bock, Carlsmith+ (CHIC, SACL)	VOSBURGH	72	PR D6 1834	+Devlin, Esterling, Goz, Bryman+ (RUTG, MASA)
PDG	86C	PL 170B 132	+Berenstein, Bogdanov, Vishnevsky+ (ITEP)	Also	71	PL 26 866	Vosburgh, Devlin, Esterling, Goz+ (RUTG, MASA)
BERNSTEIN	85B	PRL 54 1631	+Masso, Singer (CERN)	BALATS	71	SJNP 13 53	+Berezin, Vishnevsky, Galanina+ (ITEP)
BLACK	85	PRL 54 1628	+Berenstein+ (BNL, CHIC, STAN, WISC)	Also	71	PL 35B 604	+Barylov, Veselovsky, Davidenko+ (ITEP)
COUPL	85	PRL 55 66	+Bock, Cheng, Fischbach (BNL, CHIC, PURD)	BISI	71	PL 36B 533	+Darrulat, Ferrero, Rubbia+ (AACH, CERN, TORI)
COLATS	85	SJNP 38 556	+Fischbach, Cheng+ (PURD, BNL, YALE)	BURGUN	71	PL 2 1169	+Lesquoy, Muller, Pauli+ (SACL, CERN, OSLO)
Translated from YAF 38 927			+Aronson, Bock, Cheng+ (BNL, CHIC, PURD)	CARNEGIE	71	PR D4 1	+Cester, Fitch, Strovink, Sulak (PRIN)
BERGSTROM	83	PL 131B 229	+Roes, Porter, Aguilar-Benitez+ (HELSE, CIT, CERN)	CHAN	71	LBL-350 Thesis	Cox, Ettinger+ (JHU, SLAC, UCLA)
ARONSON	82B	PRL 48 1306	+Dzhordzhadze, Genchev, Grigalashvili+ (JINR)	CHIEN	71	PL 35B 261	Daily, Innocenti, Seppi+ (SLAC, JHU, UCLA)
Also	82B	PRL 116B 73	+Birelev, Vestergombi, Genchev+ (JINR)	Also	71	PL 41B 647	Dralle, Canter, Engler, Fisk+ (CMU, BNL, CASE)
Also	83	PRL D28 476	+Chiang, Kycia, Li, Littenberg, Marx+ (BNL, ROCH)	CHO	71	PR D3 1557	Elioff, Field, Frisch, Johnson, Kerth+ (LRL)
Also	83B	PR D28 495	+Chiang, Kycia, Li, Littenberg, Marx+ (BNL, ROCH)	CLARK	71	PRL 26 1667	Frisch (LRL)
PDG	82B	PL 111B 70	+Derrick, Miller, Schlereth, Engler+ (ANL, CMU)	Also	70	UCRL 19709 Thesis	Field (SLAC)
BIRULEV	81	NP B182 1	+Leipuner, Larsen, Schmidt, Blatt+ (BNL, YALE)	Also	71	UCRL 20264 Thesis	Field (SLAC)
Also	80	SJNP 31 622	+Vestergombi, Gvakhariya, Genchev+ (JINR)	Also	71	SLAC-PUB-1498 unpub.	Akavia, Coombes, Dorfan+ (SLAC, STAN)
Translated from YAF 31 1204			+Chiang, Kycia, Li, Littenberg, Marx+ (BNL, ROCH)	Also	70	SLAC-125 Thesis	Enstrom (STAN)
CARROLL	80B	PL 96B 429	+Chiang, Kycia, Li, Littenberg, Marx+ (BNL, ROCH)	JAMES	71	PL 35B 265	+Montanet, Paul, Pauli+ (CERN, SACL, OSLO)
CARROLL	80C	PL 96B 429	+Chiang, Kycia, Li, Littenberg, Marx+ (BNL, ROCH)	MEISNER	71	PR D3 59	+Mann, Hertzbach, Kofler+ (MASA, BNL, YALE)
CARROLL	80D	PL 44 525	+Derrick, Miller, Schlereth, Engler+ (ANL, CMU)	PEACH	71	PL 35B 351	+Evans, Muir, Budagov, Hopkins+ (EDIN, CERN)
CHO	80	PR D22 2688	+Leipuner, Larsen, Schmidt, Blatt+ (BNL, YALE)	REPELLIN	71	PL 35 603	+Vosif, Choulet, Gaillard, Jene+ (ORSAY, CERN)
MORSE	80	PR D21 1750	+Vestergombi, Gvakhariya, Genchev+ (JINR)	WEBBER	71	PR D3 64	+Solmitz, Crawford, Alston-Garnjost (LRL)
BIRULEV	79	SJNP 29 778	Christenson, Goldman, Hummel, Roth+ (NVU)	Also	68	PRL 21 498	Webber, Solmitz, Crawford, Alston-Garnjost (LRL)
Translated from YAF 29 1516			Christenson, Goldman, Hummel, Roth+ (NVU)	Also	69	UCRL 19226 Thesis	Webber (CERN)
CHRISTENS...	79	PRL 43 1212	+Sakitt, Snape, Stevens+ (BNL, SLAC, SBER)	WOLFF	71	PL 36B 517	+Chollet, Repellin, Gaillard+ (ORSAY, CERN)
CHRISTENS...	79B	PRL 43 1212	+Blatt, Campbell, Grannan+ (YALE, BNL)	ALBROW	70	PL 33B 516	+Aston, Barber, Bird, Ellison+ (MCHS, DARE)
HILL	79	NP B153 39	+Linsay, Glosso-Pilcher, Frisch+ (EFI, ANL)	ARONSON	70	PRL 25 1057	+Ehrlich, Hofer, Jensen+ (EFI, ILL, SLAC)
SCHMIDT	79	PRL 43 556	+Shochet, Linsay, Glosso-Pilcher+ (EFI, ANL)	BARMIN	70	PL 33B 377	+Barylov, Borison, Bysheva+ (ITEP, JINR)
SHOCHET	79	PR D19 1965	+Keves, Kraemer, Tanaka, Cho+ (CMU, ANL)	Also	70	PR D2 78	+Cronin, Thevent, Turley, Zylberajch+ (SACL)
Also	77	PRL 39 59	+Sakitt, Snape, Stevens+ (BNL, SLAC, SBER)	BECHERRAWY	70	PR D1 1452	+Drickley, Rudnick, Shepard+ (SLAC, JHU, UCLA)
ENGLER	78B	PR D18 623	+Denard, Lisauer, Miller, Engler+ (ANL, CMU)	BUCHANAN	70	PL 33B 623	Cox (Private Comm.)
HILL	78	PL 73B 483	+Field, Holley, Johnson, Kerth, Sah, Shen (LBL)	Also	71	Private Comm.	Cox (Private Comm.)
CHO	77	PR D15 587	+Shen (LBL)	BUDAGOV	70	PR D2 815	+Cundy, Myatt, Nezzick+ (CERN, ORSAY, EPOL)
CLARK	77	PR D15 553	+Cronin, Frisch, Glosso-Pilcher+ (EFI, ANL)	Also	68B	PL 28B 215	+Budagov, Cundy, Myatt+ (CERN, ORSAY, EPOL)
DEVORE	77	PR D16 565	+Dzhordzhadze, Kekelidze, Krivokhizhin+ (JINR)	CHIEN	70	PL 33B 627	+Cox, Ettinger+ (JHU, SLAC, UCLA)
DZHORDH...	77	SJNP 26 478	+Caronon+ (BGNA, EDIN, GLAS, PISA, RHEL)	Also	71	Private Comm.	Cox (Private Comm.)
Translated from YAF 26 910			+Vestergombi, Vovenko, Votruba+ (JINR)	CHOLLET	70	PR D1 2031	+Dralle, Canter, Engler, Fisk+ (CMU, BNL, CASE)
PEACH	77	NP B127 399	+Flexer, Hall, Kennelly, Kirkby+ (STAN, NYU)	CULLEN	70	PRL 19 668	Hill, Luers, Robinson, Sakitt+ (BNL, CMU)
BIRULEV	76	SJNP 24 178	+Hitlin, Kennelly, Kirkby, Liu+ (SLAC)	DARRIULAT	70	PL 31B 658	+Gaillard, Jane, Ratcliffe, Repellin+ (CERN)
Translated from YAF 24 340			Donaldson (SLAC)	FAISSNER	70	PL 32B 523	+Darrulat, Deutsch, Foeth+ (AACH, CERN, TORI)
COOMBES	76	PRL 37 249	+Jensen, Surko, Thaler+ (PRIN, MASA)	GINSBERG	70	PL 33B 249	+Ferro, Glosso, Holder+ (AACH, CERN, TORI)
DONALDSON	76	PR D14 2839	+Cence, Jones, Parker+ (NDAM, HAWA, LRL)	JENSEN	70	NC 70A 57	+Reithler, Thome, Gaillard+ (AACH3, CERN, RHEIF)
Also	74	SLAC-184 Thesis	+Cence, Jones, Peterson, Stenger+ (HAWA, LRL)	Also	69	PR D1 229	Thesis
FUKUSHIMA	76	PRL 36 348	+Baldo-Ceolin, Bobisut, Calimani+ (PADO, WISC)	MARX	70	PRL 23 615	+Jensen, Aronson, Ehrlich, Fryberger+ (EFI, ILL)
GJESDAL	76	NP B109 118	+Frickley, Nagy+ (PENN, CHIC, TEMP)	Also	70B	Nevis 179 Thesis	+Nygren, Peoples+ (COLU, HARV, CERN)
REY	76	PR D13 1161	+Drucke, Pepper, Rudnick+ (UCLA, SLAC, JHU)	SCRIBANO	70	PL 32B 224	+Mannelli, Pierazzini, Marx+ (PISA, COLU, HARV)
Also	69	PRL 22 1210	+Modis, Nygren, Pun+ (COLU, NYU)	SMITH	70	PL 32B 133	+Wang, Whately, Zorn, Hornbostel (UMD, BNL)
BALDO...	75	NC 25A 688	+Balashov, Banik+ (JINR, BERL, BUDA, PRAG, SERP, SOFI)	WEBBER	70	PR D1 1967	+Webber (LRL)
BLUMENTHAL	75	PRL 34 164	+Ferro (TORI)	Also	69	UCRL 19226 Thesis	Webber (PRIN)
BUCHANAN	75	PR D11 457	Donaldson, Hitlin, Kennelly, Kirkby, Liu+ (SLAC)	BANNER	69	PR 18B 2033	+Cronin, Liu, Pilcher (PRIN)
CARITHERS	75	PRL 34 1244	+Fryberger, Hitlin, Liu+ (SLAC, UCSC)	Also	68	PRL 21 1103	+Banner, Cronin, Liu, Pilcher (PRIN)
SMITH	75B	UCSD Thesis unpub.	+Hitlin, Kennelly, Kirkby+ (SLAC, UCSC)	BEILLIERE	69	PL 30B 202	+Boutang, Limon (EPOL)
ALBRECHT	74	PL 48B 393	Donaldson, Hitlin, Kennelly, Kirkby, Liu+ (SLAC)	BENNETT	69	PL 29B 317	+Nygren, Saal, Steinberger+ (COLU, BNL)
BISI	74	PL 50B 504	Donaldson, Hitlin, Kennelly, Kirkby, Liu+ (SLAC)	BOHM	69B	NP 89 605	+Darrulat, Glosso, Kaftanov+ (CERN)
DONALDSON	74	SLAC-184 Thesis	+Gjesdal, Kamae, Presser+ (CERN, HEIDH)	Also	68	PL 27B 321	Bohm, Darrulat, Glosso, Kaftanov (CERN)
Also	76	PR D14 2839	+Luth (CERN)	CENCE	69	PRL 22 1210	+Jones, Peterson, Stenger+ (HAWA, LRL)
DONALDSON	74B	PR D9 2960	+Gjesdal, Presser+ (CERN, HEIDH)	EVANS	69	PRL 23 427	+Golden, Muir, Peach+ (EDIN, CERN)
Also	73B	PL 31 337	+Gjesdal, Presser, Steffen+ (CERN, HEIDH)	FAISSNER	69	PL 30B 204	+Foeth, Staudt, Tittel+ (AACH3, CERN, TORI)
DONALDSON	74C	PRL 33 554	+Gjesdal, Presser, Steffen+ (CERN, HEIDH)	FOETH	69	PL 30B 282	+Holder, Rademacher+ (AACH, CERN, TORI)
Also	74	SLAC-184 Thesis	+Franklin, Morse+ (COLO, SLAC, UCSC)	GAILLARD	69	NC 59A 453	+Galbraith, Hussri, Jene+ (CERN, RHEL, AACH)
FIELD	74	SLAC-PUB-1498 unpub.	+Regler, Stier+ (CERN, ORSAY, VIEN)	GOBBI	69B	PRL 22 685	+Gaillard, Krienen, Galbraith+ (CERN, RHEL, AACH)
GEWENIGER	74	PL 48B 483	+Smith, Whately, Zorn, Hornbostel (UMD, BNL)	LITTENBERG	69	PRL 22 684	+Green, Hake, Moflett, Rosen, Goz+ (ROCH, RUTG)
Also	74	CERN Int. 74-4 Thesis	+Larsen, Leipuner, Sapp, Sessoms+ (BNL, YALE)	LONGO	69	PR 181 1808	+Field, Piccioni, Mehlopp+ (UCSD)
GEWENIGER	74B	PL 48B 487	+Aston, Barber, Bird, Ellison+ (MCHS, DARE)	PACIOTTI	69	UCRL 19446 Thesis	+Young, Helland (MICH, UCLA)
Also	74B	PL 52B 119	+Benary, Borowitz, Lande+ (TELA, CERN)	SAAL	69	Thisis	+Coly (LRL)
GEWENIGER	74C	PL 52B 108	+Balashov, Banik+ (JINR)	ABRAMS	68B	PR 176 1603	+Abashian, Mischke, Nefkens, Smith+ (ILL)
GJESDAL	74	PRL 33 1458	+Brandenburg, Johnson, Leith, Loos+ (SLAC)	ARNOLD	68	PL 28B 56	+Budagov, Cundy, Aubert+ (CERN, ORSAY)
MESSNER	74	PRL 33 1458	+Nygren, Gordon+ (COLU, BNL, CERN)	ARONSON	68	PRL 20 287	+Chen (PRIN)
NIEBERGALL	74	PL 49B 103	+Carithers, Modis, Nygren+ (COLU, CERN, NYU)	Also	69	PR 175 1706	+Chen (PRIN)
WANG	74	PR D9 540	+Muir, Peach, Budagov+ (EDIN, CERN)	BARTLETT	68	PRL 22 685	+Carnegie, Fitch+ (PRIN)
WILLIAMS	73	NP B58 22	+Evans, Golden, Muir, Peach+ (EDIN, CERN)	BASILE	68	PL 26B 542	+Cronin, Thevent, Turley+ (SACL)
ALBROW	73B	NP B65 301	+Frisch, Martin, Stovot, Sompayrac (MIT)	BASILE	68B	PL 28B 58	+Cronin, Thevent, Turley, Zylberajch+ (SACL)
ALEXANDER	73B	JINR P1 7539	+Hepp, Jensen, Smoot, Webb (PRIN)	BENNETT	68	PL 27B 248	+Nygren, Steinberger+ (COLU, CERN)
ANIKINA	73	PL 43B 529	+Webb (PRIN)	BLANPIED	68	PR 21 1650	+Levit, Engels+ (CASE, HARV, MCGI)
BARBIELLINI	73	PR D8 1978	+Smith (MIT, STON)	BOBAGOV	68B	PL 27B 54	+Burmeister, Cundy+ (CERN, ORSAY, IPNP)
BRANDENB...	73	PRL 31 1025	+Hutton, Field, Sharp, Blackmore+ (CAVE, RHEL)	Also	68B	PL 28B 215	+Budagov, Cundy, Myatt+ (CERN, ORSAY, EPOL)
Also	73B	PRL 30 1336	+Binnie, Gallivan, Binnie, Gomez+ (CIT)	JAMES	68	NP B8 365	+Briand (IPNP, CERN)
EVANS	73	PR D7 36		Also	68	JETP 26 257	Helland, Longo, Young (UCLA, MICH)
Also	69	PRL 23 427		KULYUKINA	68	JETP 26 257	+Mestvirishvili, Nyagu+ (JINR)
ACKLER	73	PRL 31 847		Translated from ZETF 53 29.			
FITZ	73	PRL 31 1524					
Also	72	COO-3072-13 Thesis					
GINSBERG	72	PR D8 3887					
HART	73	NP B66 317					
MALLARY	73	PR D7 1953					
Also	70	PRL 25 1214					

See key on page 1343

Meson Full Listings

 $K_L^0, K^*(892)$ $K^*(892)$

$$I(J^P) = \frac{1}{2}(1^-)$$

 $K^*(892)$ MASS

CHARGED ONLY

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
891.59 ± 0.24 OUR AVERAGE		Error includes scale factor of 1.1.			
890.4 ± 0.2 ± 0.5	79709 ± 801	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892.6 ± 0.5	5840	BAUBILLIER	84B	HBC	- 8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
888.0 ± 3.0		NAPIER	84	SPEC	+ 200 $\pi^- p \rightarrow 2K_S^0 X$
891.0 ± 1.0		NAPIER	84	SPEC	+ 200 $\pi^- p \rightarrow 2K_S^0 X$
891.7 ± 2.1	3700	BARTH	83	HBC	+ 70 $K^+ p \rightarrow K^0 \pi^+ X$
891.0 ± 1.0	4100	TOAFF	81	HBC	- 6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892.8 ± 1.6		AJINENK	80	HBC	+ 32 $K^+ p \rightarrow K^0 \pi^+ X$
890.7 ± 0.9	1800	AGUILAR...	78B	HBC	± 0.76 $\bar{p} p \rightarrow K^{\mp} K_S^0 \pi^{\pm}$
886.6 ± 2.4	1225	BALAND	78	HBC	± 12 $\bar{p} p \rightarrow (K\pi)^{\pm} X$
891.7 ± 0.6	6706	COOPER	78	HBC	± 0.76 $\bar{p} p \rightarrow (K\pi)^{\pm} X$
891.9 ± 0.7	9000	2 PALER	75	HBC	- 14.3 $K^- p \rightarrow (K\pi)^- X$
892.2 ± 1.5	4404	AGUILAR...	71B	HBC	- 3.9, 4.6 $K^- p \rightarrow (K\pi)^- p$
891.0 ± 2.0	1000	CRENNELL	69D	DBC	- 3.9 $K^- N \rightarrow K^0 \pi^- X$
894 ± 1.0	2886	3 FRIEDMAN	69	HBC	- 2.1 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892 ± 2	728	FRIEDMAN	69	HBC	- 2.45 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892 ± 1.0	3229	FRIEDMAN	69	HBC	- 2.6 $K^- p \rightarrow \bar{K}^0 \pi^- p$
892 ± 1.6	1027	FRIEDMAN	69	HBC	- 2.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$
890 ± 3.0	720	BARLOW	67	HBC	± 1.2 $\bar{p} p \rightarrow (K^0 \pi)^{\pm} K^{\mp}$
889 ± 3.0	600	BARLOW	67	HBC	± 1.2 $\bar{p} p \rightarrow (K^0 \pi)^{\pm} K\pi$
891 ± 2.3	620	3 DEBAERE	67B	HBC	+ 3.5 $K^+ p \rightarrow K^0 \pi^+ p$
891.0 ± 1.2	1700	4 WOJCICKI	64	HBC	- 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
890.0 ± 2.3	800	3,4 CLELAND	82	SPEC	+ 30 $K^+ p \rightarrow K_S^0 \pi^+ p$
896.0 ± 1.1	3200	3,4 CLELAND	82	SPEC	+ 50 $K^+ p \rightarrow K_S^0 \pi^+ p$
893.0 ± 1.0	3600	3,4 CLELAND	82	SPEC	- 50 $K^+ p \rightarrow K_S^0 \pi^- p$
896.0 ± 1.9	380	DELFOSSO	81	SPEC	+ 50 $K^{\pm} p \rightarrow K^{\pm} \pi^0 p$
886.0 ± 2.3	187	DELFOSSO	81	SPEC	- 50 $K^{\pm} p \rightarrow K^{\pm} \pi^0 p$
894.2 ± 2.0	765	3 CLARK	73	HBC	- 3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$
894.3 ± 1.5	1150	3,4 CLARK	73	HBC	- 3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$
888 ± 2.5	540	3 DEWIT	68	HBC	- 3 $K^- n \rightarrow \bar{K}^0 \pi^- n$
892.0 ± 2.6	341	3 SCHWEING...	68	HBC	- 5.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NEUTRAL ONLY

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
896.10 ± 0.28 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.			
895.9 ± 0.5 ± 0.2		ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
894.52 ± 0.63	25k	2 ATKINSON	86	OMEG	20-70 γp
894.63 ± 0.76	20k	2 ATKINSON	86	OMEG	20-70 γp
897 ± 1	28k	EVANGELISTA	80	OMEG	0 10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$
898.4 ± 1.4	1180	AGUILAR...	78B	HBC	0 0.76 $\bar{p} p \rightarrow K^{\mp} K_S^0 \pi^{\pm}$
894.9 ± 1.6		WICKLUND	78	ASPK	0 3.4, 6 $K^{\pm} N \rightarrow (K\pi)^0 N$
897.6 ± 0.9		BOWLER	77	DBC	0 5.4 $K^+ d \rightarrow K^+ \pi^- pp$
895.5 ± 1.0	3600	MCCUBBIN	75	HBC	0 3.6 $K^- p \rightarrow K^- \pi^+ n$
897.1 ± 0.7	22k	2 PALER	75	HBC	0 14.3 $K^- p \rightarrow (K\pi)^0 X$
896.0 ± 0.6	10k	FOX	74	RVUE	0 2 $K^- p \rightarrow K^- \pi^+ n$
896.0 ± 0.6		FOX	74	RVUE	0 2 $K^+ n \rightarrow K^+ \pi^- p$
896 ± 2		5 MATISON	74	HBC	0 12 $K^+ p \rightarrow K^+ \pi^- \Delta$
896.0 ± 1.0	3186	LEWIS	73	HBC	0 2.1-2.7 $K^+ p \rightarrow K^+ \pi^- p$
894.0 ± 1.3		5 LINGLIN	73	HBC	0 2-13 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$
898.4 ± 1.3	1700	3 BUCHNER	72	DBC	0 4.6 $K^+ n \rightarrow K^+ \pi^- p$
897.9 ± 1.1	2934	3 AGUILAR...	71B	HBC	0 3.9, 4.6 $K^- p \rightarrow K^- \pi^+ n$
898.0 ± 0.7	5362	3 AGUILAR...	71B	HBC	0 3.9, 4.6 $K^- p \rightarrow K^- \pi^+ \pi^- p$
895.0 ± 1.0	4300	4 HABER	70	DBC	0 3 $K^- N \rightarrow K^- \pi^+ X$
893.7 ± 2.0	10k	DAVIS	69	HBC	0 12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$
894.7 ± 1.4	1040	3 DAUBER	67B	HBC	0 2.0 $K^- p \rightarrow K^- \pi^+ \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
900.7 ± 1.1	5900	BARTH	83	HBC	0 70 $K^+ p \rightarrow K^+ \pi^- X$

KUNZ	68	PU 46 Thesis		(PRIN)
BENNETT	67	PR 19 993	+Nygren, Saal, Steinberger+	(COLU)
BOTT...	67	PL 24B 994	+Bott-Bodenhausen, DeBouard, Cassel+	(CERN)
BOTT...	67B	PL 24B 438	+Bott-Bodenhausen, DeBouard, Dekkers+	(CERN)
Also	66B	PL 20 212	+Bott-Bodenhausen, DeBouard, Cassel+	(CERN)
Also	66	PL 23 277	+Bott-Bodenhausen, DeBouard, Cassel+	(CERN)
CRONIN	67	PRL 18 25	+Kunz, Risk, Wheeler	(PRIN)
Also	68	Thesis (unpub.)	+Wheeler	(PRIN)
CRONIN	67B	Princeton 11/67	+Kunz, Risk, Wheeler	(PRIN)
DEBOUARD	67	NC 52A 662	+Dekkers, Jordan, Mermoud+	(CERN)
Also	65	PL 15 58	+DeBouard, Dekkers, Scharrf+	(CERN, ORSAY, MPIM)
DEVLIN	67	PRL 18 54	+Solomon, Shepard, Beall+	(PRIN, UMD)
Also	68	PR 169 1045	+Sayer, Beall, Devlin, Shephard+	(UMD, PPA, PRIN)
DORFAN	67	PRL 19 987	+Enstrom, Raymond, Schwartz+	(SLAC, LRL)
FELDMAN	67B	PR 155 1611	+Frankel, Highland, Sloan	(PENN)
FIRESTONE	67	PRL 18 176	+Kim, Lach, Sandweiss+	(YALE, BNL)
FITCH	67	PR 164 1711	+Roth, Russ, Vernon	(PRIN)
GINSBERG	67	PR 162 1570		(MARB)
HAWKINS	67	PR 156 1444		(YALE)
HILL	67	PRL 19 668	+Luers, Robinson, Sakitt+	(BNL, CMU)
HOPKINS	67	PRL 19 185	+Bacon, Eisler	(BNL)
KADYK	67	PRL 19 597	+Chan, Drijard, Oren, Sheldon	(LRL)
KULYUKINA	67	Preprint	+Mestvirishvili, Nyagu+	(JINR)
LOWYS	67	PL 24B 75	+Aubert, Chounet, Pascaud+	(EPOL, ORSAY)
MISCHKE	67	PR 18 138	+Abashian, Abrams+	(ILL)
NEFKENS	67	PR 157 233	+Abashian, Abrams, Carpenter, Fisher+	(ILL)
SCHMIDT	67	Nevis 160 Thesis		(COLU)
TODOROFF	67	Thesis		(ILL)
ALFF...	66B	PL 21 595	+Alff-Steinberger, Heuer, Kleinknecht+	(CERN)
ANIKINA	66	SJNP 2 339	+Vardenga, Zhuravleva+	(JINR)
Also	66	Translated from YAF 47		
AUERBACH	66B	PRL 17 980	+Mann, McFarlane, Sciulli	(PENN)
BASILE	66	Balaton Conf.	+Cronin, Thevenet+	(SACL)
BEHR	66	PL 22 540	+Brisson, Petiau+	(EPOL, MILA, PADO, ORSAY)
BOTT...	66	PL 23 277	+Bott-Bodenhausen, DeBouard, Cassel+	(CERN)
CARPENTER	66	PR 142 871	+Abashian, Abrams, Fisher	(ILL)
CRIGEE	66	PRL 17 150	+Fox, Frauenfelder, Hanson, Moscat+	(ILL)
FIRESTONE	66	PRL 16 556	+Kim, Lach, Sandweiss+	(YALE, BNL)
HAWKINS	66	PL 21 238		(YALE)
Also	67	PL 156 1444	+Hawkins	(YALE)
ANDERSON	65	PRL 14 475	+Crawford, Golden, Stern, Binford+	(LRL, WISC)
ANIKINA	65	JINR P 2488	+Vardenga, Zhuravleva, Kotlyas+	(JINR)
ASTBURY	65	PL 16 80	+Finocchiaro, Beusch+	(CERN, ZURI)
Also	65	HPA 39 523	+Pepin	
ASTBURY	65B	PL 18 175	+Michelini, Beusch+	(CERN, ZURI)
ASTBURY	65C	PL 18 178	+Michelini, Beusch+	(CERN, ZURI)
AUBERT	65	PL 17 599	+Behr, Canavan, Chounet+	(EPOL, ORSAY)
Also	67	PL 4B 75	+Lows, Aubert, Chounet, Pascaud+	(EPOL, ORSAY)
BALDO...	65	NC 38 684	+Baldo-Ceolin, Calimani, Ciampolillo+	(PADO)
FISHER	65	ANL 7130 83	+Abashian, Abrams, Carpenter+	(ILL)
FITCH	65	PRL 15 73	+Roth, Russ, Vernon	(PRIN)
FRANZINI	65	PR 140B 127	+Kirsch, Plano+	(COLU, RUTG)
GALBRAITH	65	PRL 14 383	+Manning, Jones+	(AERE, BRIS, RHEL)
GUIDONI	65	Argonne Conf. 49	+Barnes, Foelsche, Ferbel, Firestone+	(BNL, YALE)
HOPKINS	65	Argonne Conf. 67	+Bacon, Eisler	(VAND, RUTG)
ADAIR	64	PL 12 67	+Leipuner	(YALE, BNL)
ALEKSANYAN	64B	Dubna Conf. 2 102	+Alikhanyan, Vartazaryan+	(YERE)
Also	64	JETP 19 1019	+Aleksanyan+	(LEBD, MPEI, YERE)
ANIKINA	64	JETP 19 42	+Zhuravleva+	(GEOR, JINR)
Also	64	Translated from ZETF 46 1504		
CHRISTENS...	64	PRL 13 138	+Christenson, Cronin, Fitch, Turlay	(PRIN)
FUJII	64	Dubna Conf. 2 146	+Jovanovich, Turkot+	(BNL, UMD, MIT)
LUERS	64	PR 133B 1276	+Mittra, Willis, Yamamoto	(BNL)
DARMON	62	PL 3 57	+Rousset, Six	(EPOL)
ASTIER	61	Aix Conf. 1 227	+Blaskovic, Rivet, Siaud+	(EPOL)
FITCH	61	NC 22 1160	+Priou, Perkins	(PRIN, LASL)
GOOD	61	PR 124 1223	+Matsen, Muller, Piccioni+	(LRL)
NYAGU	61	PRL 6 552	+Okonov, Petrov, Rosanova, Rusakov	(JINR)
Also	61B	JETP 13 1138	+Nyagu, Okonov, Petrov, Rozanova+	(JINR)
Also	61B	Translated from ZETF 40 1618		
BARDON	58	ANP 5 156	+Lande, Lederman	(COLU, BNL)


OTHER RELATED PAPERS

HAYAKAWA	93	PR D48 1150	+Sanda	(NAGO)
Also	93	"Searching for $T, CP, CPT, \Delta S = \Delta Q$ Rule Violations in the Neutral K Meson System: A Guide"	+Valencia	(BNL, FNAL)
LITTEMBERG	93	ARNPS 43 729		
Also	93	Rare and Radiative Kaon Decays		
RITCHIE	93	RMP 65 1149	+Wojcicki	
WINSTEIN	93	RMP 65 1113	+Wolfenstein	
Also	93	"The Search for Direct CP Violation"		
BATTISTON	92	PRPL 214 293	+Coccolicchio, Fogli, Paver	(PGIA, CERN, TRSTT)
Also	92	Status and Perspectives of K Decay Physics		
DIB	92	PR D46 2265	+Peccei	(UCLA)
Also	92	Tests of CP conservation in the neutral kaon system.		
KLEINKNECHT	92	CNPP 20 281		(MANZ)
Also	92	New Results on CP Violation in Decays of Neutral K Mesons.		
KLEINKNECHT	90	ZPHY C46 S57		(MANZ)
PEACH	90	JPG 16 131		(EDIN)
BRYMAN	89	IJMP A4 79		(TRIUM)
Also	89	"Rare Kaon Decays"		
KLEINKNECHT	76	ARNS 26 1		(DORT)
GINSBERG	73	PR D8 3887	+Smith	(MIT, STON)
GINSBERG	70	PR D1 229		(HAIF)
HEUSSE	70	LNC 3 449	+Aubert, Pascaud, Vialle	(ORSAY)
CRONIN	68C	Vienna Conf. 281		(PRIN)
RUBBIA	67	PL 24B 531	+Steinberger	(CERN, COLU)
Also	66C	PL 23 167	+Rubbia, Steinberger	(CERN, COLU)
Also	66C	PL 20 207	+Alff-Steinberger, Heuer, Kleinknecht+	(CERN)
Also	66B	PL 21 595	+Alff-Steinberger, Heuer, Kleinknecht+	(CERN)
AUERBACH	66	PR 149 1052	+Dobbs, Lande, Mann, Sciulli+	(PENN)
Also	66	PRL 14 192	+Auerbach, Lande, Mann, Sciulli, Uto+	(PENN)
FIRESTONE	66B	PRL 17 116	+Kim, Lach, Sandweiss+	(YALE, BNL)
BEHR	65	Argonne Conf. 59	+Brisson, Bellotti+	(EPOL, MILA, PADO)
MESTVIRISH...	65	JINR P 2449	+Mestvirishvili, Nyagu, Petrov, Rusakov+	(JINR)
TRILLING	65B	UCRL 16473		(LRL)
Also	65	Updated from 1965 Argonne Conference, page 115.		
JOVANOVI...	63	BNL Conf. 42	+Jovanovich, Fischer, Burris+	(BNL, UMD)

Meson Full Listings

$K^*(892)$

WEIGHTED AVERAGE
896.10±0.28 (Error scaled by 1.4)



$K^*(892)^0$ mass (MeV)

- 1 From a partial wave amplitude analysis.
- 2 Inclusive reaction. Complicated background and phase-space effects.
- 3 Mass errors enlarged by us to Γ/\sqrt{N} . See note.
- 4 Number of events in peak reevaluated by us.
- 5 From pole extrapolation.

NOTE ON $K^*(892)$ MASSES AND MASS DIFFERENCES

Unrealistically small errors are reported by some experiments. We use simple "realistic" tests for the minimum errors on the determination of mass and width from a sample of N events:

$$\delta_{\min}(m) = \frac{\Gamma}{\sqrt{N}}, \quad \delta_{\min}(\Gamma) = 4 \frac{\Gamma}{\sqrt{N}}.$$

(For a detailed discussion, see the 1971 edition of this note.)
We consistently increase unrealistic errors before averaging.

$m_{K^*(892)^0} - m_{K^*(892)^\pm}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
6.7±1.2 OUR AVERAGE					
7.7±1.7	2980	AGUILAR-...	78B HBC	±0	0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
5.7±1.7	7338	AGUILAR-...	71B HBC	-0	3.9,4.6 $K^- p$
6.3±4.1	283	⁶ BARASH	67B HBC		0.0 $\bar{p}p$

⁶Number of events in peak reevaluated by us.

$K^*(892)$ RANGE PARAMETER

All from partial wave amplitude analyses.

VALUE (GeV^{-1})	DOCUMENT ID	TECN	CHG	COMMENT
12.1±3.2±3.0	BIRD	89 LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
3.4±0.7	ASTON	88 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

$K^*(892)$ WIDTH

CHARGED ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
49.8±0.8 OUR FIT					
49.8±0.8 OUR AVERAGE					
45.2±1 ±2	79709±801	⁷ BIRD	89 LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
49.0±2.0	5840	BAUBILLIER	84B HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
56.0±4.0		NAPIER	84 SPEC	-	200 $\pi^- p \rightarrow 2K_S^0 X$
51.0±2.0	4100	TOAFF	81 HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
50.5±5.6		AJINENKO	80 HBC	+	32 $K^+ p \rightarrow K^0 \pi^+ X$
45.8±3.6	1800	AGUILAR-...	78B HBC	±	0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
52.0±2.5	6706	⁸ COOPER	78 HBC	±	0.76 $\bar{p}p \rightarrow (K\pi)^\pm X$
52.1±2.2	9000	⁹ PALER	75 HBC	-	14.3 $K^- p \rightarrow (K\pi)^- X$
46.3±6.7	765	⁸ CLARK	73 HBC	-	3.13 $K^- p \rightarrow \bar{K}^0 \pi^- p$
48.2±5.7	1150	^{8,10} CLARK	73 HBC	-	3.3 $K^- p \rightarrow \bar{K}^0 \pi^- p$
54.3±3.3	4404	⁸ AGUILAR-...	71B HBC	-	3.9,4.6 $K^- p \rightarrow (K\pi)^- p$
53 ±4.0	2886	⁸ FRIEDMAN	69 HBC	-	2.1 $K^- p \rightarrow \bar{K}^0 \pi^- p$
49 ±7.3	728	⁸ FRIEDMAN	69 HBC	-	2.45 $K^- p \rightarrow \bar{K}^0 \pi^- p$
46 ±3.2	3229	⁸ FRIEDMAN	69 HBC	-	2.6 $K^- p \rightarrow \bar{K}^0 \pi^- p$
49 ±6.1	1027	⁸ FRIEDMAN	69 HBC	-	2.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$
46.0±5.0	1700	^{8,10} WOJCICKI	64 HBC	-	1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
42.8±7.1	3700	BARTH	83 HBC	+	70 $K^+ p \rightarrow K^0 \pi^+ X$
64.0±9.2	800	^{8,10} CLELAND	82 SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
62.0±4.4	3200	^{8,10} CLELAND	82 SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
55.0±4.0	3600	^{8,10} CLELAND	82 SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$
62.6±3.8	380	DELFOSSÉ	81 SPEC	+	50 $K^\pm p \rightarrow K^\pm \pi^0 p$
50.5±3.9	187	DELFOSSÉ	81 SPEC	-	50 $K^\pm p \rightarrow K^\pm \pi^0 p$

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
50.5±0.6 OUR FIT Error includes scale factor of 1.1.					
50.5±0.6 OUR AVERAGE Error includes scale factor of 1.1.					
50.8±0.8±0.9		ASTON	88 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
46.5±4.3	5900	BARTH	83 HBC	0	70 $K^+ p \rightarrow K^+ \pi^- X$
54 ±2	28k	EVANGELISTA	80 OMEG	0	10 $\pi^- p \rightarrow K^+ \pi^- (\Lambda, \Sigma)$
45.9±4.8	1180	AGUILAR-...	78B HBC	0	0.76 $\bar{p}p \rightarrow K^\mp K_S^0 \pi^\pm$
51.2±1.7		WICKLUND	78 ASPK	0	3,4,6 $K^\pm N \rightarrow (K\pi)^0 N$
48.9±2.5		BOWLER	77 DBC	0	5.4 $K^+ d \rightarrow K^+ \pi^- pp$
48 ⁺³ ₋₂	3600	MCCUBBIN	75 HBC	0	3.6 $K^- p \rightarrow K^- \pi^+ n$
50.6±2.5	22k	⁹ PALER	75 HBC	0	14.3 $K^- p \rightarrow (K\pi)^0 X$
47 ±2	10k	FOX	74 RVUE	0	2 $K^- p \rightarrow K^- \pi^+ n$
51 ±2		FOX	74 RVUE	0	2 $K^+ n \rightarrow K^+ \pi^- p$
46.0±3.3	3186	⁸ LEWIS	73 HBC	0	2.1-2.7 $K^+ p \rightarrow K^\pm \pi^\pm p$
51.4±5.0	1700	⁸ BUCHNER	72 DBC	0	4.6 $K^+ n \rightarrow K^+ \pi^- p$
55.8 ^{+4.2} _{-3.4}	2934	⁸ AGUILAR-...	71B HBC	0	3.9,4.6 $K^- p \rightarrow K^- \pi^+ n$
48.5±2.7	5362	AGUILAR-...	71B HBC	0	3.9,4.6 $K^- p \rightarrow K^- \pi^+ \pi^- p$
54.0±3.3	4300	^{8,10} HABER	70 DBC	0	3 $K^- N \rightarrow K^- \pi^+ X$
53.2±2.1	10k	⁸ DAVIS	69 HBC	0	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$
44 ±5.5	1040	⁸ DAUBER	67B HBC	0	2.0 $K^- p \rightarrow K^- \pi^+ \pi^- p$

⁷From a partial wave amplitude analysis.

⁸Width errors enlarged by us to $4 \times \Gamma/\sqrt{N}$; see note.

⁹Inclusive reaction. Complicated background and phase-space effects.

¹⁰Number of events in peak reevaluated by us.

$K^*(892)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K\pi$	~ 100	%
Γ_2 $(K\pi)^\pm$	(99.899±0.009)	%
Γ_3 $(K\pi)^0$	(99.770±0.020)	%
Γ_4 $K^0\gamma$	(2.30 ±0.20)	$\times 10^{-3}$
Γ_5 $K^\pm\gamma$	(1.01 ±0.09)	$\times 10^{-3}$
Γ_6 $K\pi\pi$	< 7	$\times 10^{-4}$ 95%

See key on page 1343

Meson Full Listings

$K^*(892), K_1(1270)$

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 18 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 15.2$ for 16 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_5 & -100 & \\ \Gamma & 17 & -17 \\ & x_2 & x_5 \end{array}$$

Mode	Rate (MeV)
$\Gamma_2 (K\pi)^\pm$	49.8 \pm 0.8
$\Gamma_5 K^\pm \gamma$	0.050 \pm 0.005

CONSTRAINED FIT INFORMATION

An overall fit to the total width and a partial width uses 18 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 18.4$ for 16 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$\begin{array}{c|cc} x_4 & -100 & \\ \Gamma & 14 & -14 \\ & x_3 & x_4 \end{array}$$

Mode	Rate (MeV)	Scale factor
$\Gamma_3 (K\pi)^0$	50.4 \pm 0.6	1.1
$\Gamma_4 K^0 \gamma$	0.117 \pm 0.010	

$K^*(892)$ PARTIAL WIDTHS

$\Gamma(K^0 \gamma)$	VALUE (keV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4
116 \pm 10 OUR FIT							
116.5 \pm 9.9	584	CARLSMITH	86	SPEC	0	$K_L^0 A \rightarrow K_S^0 \pi^0 A$	

$\Gamma(K^\pm \gamma)$	VALUE (keV)	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5
50 \pm 5 OUR FIT						
50 \pm 5 OUR AVERAGE						
48.0 \pm 11.0	BERG	83	SPEC	-	156 $K^- A \rightarrow \bar{K} \pi A$	
51.0 \pm 5.0	CHANDLEE	83	SPEC	+	200 $K^+ A \rightarrow K \pi A$	

$K^*(892)$ BRANCHING RATIOS

$\Gamma(K^0 \gamma) / \Gamma_{\text{total}}$	VALUE (units 10^{-3})	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4 / Γ
2.30 \pm 0.20 OUR FIT						
1.5 \pm 0.7	CARITHERS	75B	CNTR	0	8-16 $\bar{K}^0 A$	

$\Gamma(K^\pm \gamma) / \Gamma_{\text{total}}$	VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5 / Γ
1.01 \pm 0.09 OUR FIT							
<1.6	95		BEMPORAD	73	CNTR	+	10-16 $K^+ A$

$\Gamma(K\pi\pi) / \Gamma((K\pi)^\pm)$	VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6 / Γ_2
<0.0007							
<0.002	95		JONGEJANS	78	HBC	4 $K^- p \rightarrow p \bar{K}^0 2\pi$	
<0.002	64		WOJCICKI	64	HBC	- 1.7 $K^- p \rightarrow \bar{K}^0 \pi^- p$	

$K^*(892)$ REFERENCES

BIRD	89	SLAC-332	(SLAC)
ASTON	88	NP B296 493	+Awaji, Bienz, Bird+ (SLAC, NAGO, CINC, INUS)
ATKINSON	86	ZPHY C30 521	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)
CARLSMITH	86	PRL 56 18	+Bernstein, Peyaud, Turley (EPJ, SACL)
BAUBILLIER	84B	ZPHY C26 37	+ (BIRM, CERN, GLAS, MICH, CURIN)
NAPIER	84	PL 149B 514	+Chen+ (TUFTS, ARIZ, FNAL, FLOR, NDAM+)
BARTH	83	NP B223 296	+Drevermann+ (BRUX, CERN, GENO, MONS+)
BERG	83	Thesis	(ROCH)
CHANDLEE	83	PRL 51 168	+Berg, Cihangir, Collick+ (ROCH, FNAL, MINN)
CLELAND	82	NP B208 189	+Delfosse, Dorsaz, Gloor (DURH, GEVA, LAUS, PITT)
DELFOSSÉ	81	NP B183 349	+Guisan, Martin, Muhlemann, Weill+ (GEVA, LAUS)
TOAFF	81	PR D23 1500	+Musgrave, Ammar, Davis, Ecklund+ (ANL, SACL)
AJINENKO	80	ZPHY C5 177	+Barth, Dujardin+ (SERP, BRUX, MONS, SACL)
EVANGELISTA	80	NP B165 383	+ (BARI, BONN, CERN, DARE, GLAS, LIVP+)
AGUILAR-...	78B	NP B141 101	+Aguilar-Benitez+ (MADR, TATA, CERN+)
BALAND	78	NP B140 220	+Grand+ (MONS, BELG, CERN, LOIC, LALO)
COOPER	78	NP B136 365	+Gurtu+ (TATA, CERN, CDEF+)
JONGEJANS	78	NP B139 383	+Cerrada+ (ZEM, CERN, NIJM, OXF)
WICKLUND	78	PR D17 1197	+Ayres, Diebold, Greene, Kramer, Pawlicki (ANL)
BOWLER	77	NP B126 31	+Dainton, Drake, Williams (OXF)
CARITHERS	75B	PRL 35 349	+Muhlemann, Underwood+ (ROCH, MCGI)
MCCUBBIN	75	NP B86 13	+Lyons (OXF)
PALER	75	NP B96 1	+Tovey, Shah, Spiro+ (RHEL, SACL, EPOL)
FOX	74	NP B80 403	+Griss (CIT)
MATISON	74	PR D9 1872	+Galtieri, Alston-Garnjost, Flatte, Friedman+ (LBL)
BEMPORAD	73	NP B51 1	+Beusch, Freudenreich+ (CERN, ETH, LOIC)
CLARK	73	NP B54 432	+Lyons, Radjicic (OXF)
LEWIS	73	NP B60 283	+Allen, Jacobs+ (LOWC, LOIC, CDEF)
LINGLIN	73	NP B55 408	(CERN)
BUCHNER	72	NP B45 333	+Dehm, Charriere, Cornet+ (MPIM, CERN, BRUX)
AGUILAR-...	71B	PR D4 2583	+Aguilar-Benitez, Eisner, Kinson (BNL)
HABER	70	NP B17 289	+Shapira, Alexander+ (REHO, SACL, BGNA, EPOL)
CRENNELL	69D	PRL 22 4871	+Karshon, Lai, O'Neill, Scarr (BNL)
DAVIS	69	PRL 23 1071	+Derenzo, Flatte, Garnjost, Lynch, Solmitz (LRL)
FRIEDMAN	69	UCRL 18860 Thesis	(LRL)
DEWIT	68	Thesis	(ANIK)
SCHWEING...	68	PR 166 1317	+Schweingruber, Derrick, Fields+ (ANL, NWES)
BARASH	67B	PR 156 1399	+Kirsch, Miller, Tan (COLU)
BARLOW	67	NC 50A 701	+Lillestol, Montanet+ (CERN, CDEF, IRAD, LIVP)
DAUBER	67B	PR 153 1403	+Schlein, Slater, Ticho (UCLA)
DEBAERE	67B	NC 51A 401	+Goldschmidt-Clermont, Henri+ (BRUX, CERN)
WOJCICKI	64	PR 135B 484	(LRL)

OTHER RELATED PAPERS

KAMAL	92	PL B284 421	+Xu (ALBE)
NAPIER	84	PL 149B 514	+Chen+ (TUFTS, ARIZ, FNAL, FLOR, NDAM+)
CLELAND	82	NP B208 189	+Delfosse, Dorsaz, Gloor (DURH, GEVA, LAUS, PITT)
ALEXANDER	62	PRL 8 447	+Kalbfleisch, Miller, Smith (LRL)
ALSTON	62B	CERN Conf. 291	+Ticho, Wojcicki+ (LRL)
ARMENTEROS	62C	CERN Conf. 295	+Astrer, Montanet+ (CERN, CDEF)
COLLEY	62B	CERN Conf. 315	+Gelfand+ (COLU, RUTG)
ALSTON	61	PRL 6 300	+Alvarez, Eberhard, Good+ (LRL)

$K_1(1270)$

$$J(P) = \frac{1}{2}(1^+)$$

Our latest mini-review on this particle can be found in the 1984 edition.

$K_1(1270)$ MASS

VALUE (MeV)	DOCUMENT ID
1273 \pm 7 OUR AVERAGE	Includes data from the 2 datablocks that follow this one.

PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1275.0 \pm 10.0	700	GAVILLET	78	HBC	+ 4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$

PRODUCED BY K BEAMS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1270 \pm 10	DAUM	81C	CNTR	- 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1276.0	1	TORNQVIST	82B	RVUE
~ 1300.0		VERGEEST	79	HBC
1289.0 \pm 25.0	2	CARNEGIE	77	ASPK
~ 1300		BRANDENB...	76	ASPK
~ 1270.0		OTTER	76	HBC
1260		DAVIS	72	HBC
1234 \pm 12		FIRESTONE	72B	DBC

¹ From a unitarized quark-model calculation.

² From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1294 \pm 10	310	RODEBACK	81	HBC	4 $\pi^- p \rightarrow \Lambda K 2\pi$	
1300	40	CRENNELL	72	HBC	0 4.5 $\pi^- p \rightarrow \Lambda K 2\pi$	
1242.0 $^{+9.0}_{-10.0}$		3	ASTIER	69	HBC	0 $\bar{p} p$
1300	45	CRENNELL	67	HBC	0 6 $\pi^- p \rightarrow \Lambda K 2\pi$	

³ This was called the C meson.

Meson Full Listings

$K_1(1270)$, $K_1(1400)$

$K_1(1270)$ WIDTH

90 ± 20 OUR ESTIMATE This is only an educated guess; the error given is larger than the error on the average of the published values.
87 ± 7 OUR AVERAGE Includes data from the 2 datablocks that follow this one.

PRODUCED BY K^- , BACKWARD SCATTERING, HYPERON EXCHANGE

The data in this block is included in the average printed for a previous datablock.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
75.0 ± 15.0	700	GAVILLET	78	HBC	+ 4.2 $K^- p \rightarrow \Xi^- K \pi \pi$

PRODUCED BY K BEAMS

The data in this block is included in the average printed for a previous datablock.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
90 ± 8		DAUM	81C CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
150.00 ± 71.0	4	CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
~ 200		BRANDENB...	76	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
120		DAVIS	72	HBC	+ 12 $K^+ p$
188 ± 21		FIRESTONE	72B	DBC	+ 12 $K^+ d$

4 From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.

PRODUCED BY BEAMS OTHER THAN K MESONS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
66 ± 15	310	RODEBACK	81	HBC	4 $\pi^- p \rightarrow \Lambda K 2\pi$
60	40	CRENNELL	72	HBC	0 4.5 $\pi^- p \rightarrow \Lambda K 2\pi$
127.0 ± 7.0 - 25.0		ASTIER	69	HBC	0 $\bar{p} p$
60	45	CRENNELL	67	HBC	0 6 $\pi^- p \rightarrow \Lambda K 2\pi$

$K_1(1270)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\rho$	(42 ± 6)%
Γ_2 $K_0^*(1430)\pi$	(28 ± 4)%
Γ_3 $K^*(892)\pi$	(16 ± 5)%
Γ_4 $K\omega$	(11.0 ± 2.0)%
Γ_5 $K f_0(1300)$	(3.0 ± 2.0)%

$K_1(1270)$ PARTIAL WIDTHS

$\Gamma(K\rho)$ Γ_1
 VALUE (MeV) DOCUMENT ID TECN CHG COMMENT
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 57.0 ± 5.0 MAZZUCATO 79 HBC + 4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
 75.0 ± 6.0 CARNEGIE 77B ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$\Gamma(K_0^*(1430)\pi)$ Γ_2
 VALUE (MeV) DOCUMENT ID TECN CHG COMMENT
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 26.0 ± 6.0 CARNEGIE 77B ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$\Gamma(K^*(892)\pi)$ Γ_3
 VALUE (MeV) DOCUMENT ID TECN CHG COMMENT
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 14.0 ± 11.0 MAZZUCATO 79 HBC + 4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
 2.0 ± 2.0 CARNEGIE 77B ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$\Gamma(K\omega)$ Γ_4
 VALUE (MeV) DOCUMENT ID TECN CHG COMMENT
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 4.0 ± 4.00 MAZZUCATO 79 HBC + 4.2 $K^- p \rightarrow \Xi^- (K\pi\pi)^+$
 24.0 ± 3.0 CARNEGIE 77B ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$\Gamma(K f_0(1300))$ Γ_5
 VALUE (MeV) DOCUMENT ID TECN CHG COMMENT
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 22.0 ± 5.0 CARNEGIE 77B ASPK ± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

$K_1(1270)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma_{total}$ Γ_1/Γ
 VALUE DOCUMENT ID TECN COMMENT
0.42 ± 0.06 5 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 dominant RODEBACK 81 HBC 4 $\pi^- p \rightarrow \Lambda K 2\pi$

$\Gamma(K_0^*(1430)\pi)/\Gamma_{total}$ Γ_2/Γ
 VALUE DOCUMENT ID TECN COMMENT
0.28 ± 0.04 5 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K^*(892)\pi)/\Gamma_{total}$ Γ_3/Γ
 VALUE DOCUMENT ID TECN COMMENT
0.16 ± 0.05 5 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K\omega)/\Gamma_{total}$ Γ_4/Γ
 VALUE DOCUMENT ID TECN COMMENT
0.11 ± 0.02 5 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K\omega)/\Gamma(K\rho)$ Γ_4/Γ_1
 VALUE CL% DOCUMENT ID TECN COMMENT
 • • • We do not use the following data for averages, fits, limits, etc. • • •
 < 0.30 95 RODEBACK 81 HBC 4 $\pi^- p \rightarrow \Lambda K 2\pi$

$\Gamma(K f_0(1300))/\Gamma_{total}$ Γ_5/Γ
 VALUE DOCUMENT ID TECN COMMENT
0.03 ± 0.02 5 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

D-wave/S-wave RATIO FOR $K_1(1270) \rightarrow K^*(892)\pi$

VALUE DOCUMENT ID TECN COMMENT
1.0 ± 0.7 5 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$
 5 Average from low and high t data.

$K_1(1270)$ REFERENCES

TORNQVIST 82B	NP B203 268	(HLS)
DAUM 81C	NP B187 1	+ Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
RODEBACK 81	ZPHY C9 9	+ Sjogren+ (CERN, CDEF, MADR, STOH)
MAZZUCATO 79	NP B156 532	+ Pennington+ (CERN, ZEEM, NIJM, OXF)
VERGEEST 79	NP B158 265	+ Jongejans, Dionisi+ (NIJM, AMST, CERN, OXF)
GAVILLET 78	PL 76B 517	+ Diaz, Dionisi+ (AMST, CERN, NIJM, OXF) JP
CARNEGIE 77	NP B127 509	+ Cashmore, Davier, Dunwoodie, Lasinski+ (SLAC)
CARNEGIE 77B	PL 68B 287	+ Cashmore, Dunwoodie, Lasinski+ (SLAC)
BRANDENB... 76	PRL 26 703	+ Brandenburg, Carnegie, Cashmore+ (SLAC) JP
OTTER 76	NP B106 77	+ (AACH, BERL, CERN, LOIC, VIEN, EPOL+) JP
CRENNELL 72	PR D6 1220	+ Gordon, Lai, Scarr (BNL)
DAVIS 72	PR D5 2688	+ Alston-Garnjost, Barbaro, Flatte, Friedman, Lynch+ (LBL)
FIRESTONE 72B	NP D5 505	+ Goldhaber, Lissauer, Trilling (LBL)
ASTIER 69	NP B10 65	+ Marechal, Montanet+ (CDEF, CERN, IPNP, LIVP) IJP
CRENNELL 67	PRL 19 44	+ Kalbfleisch, Lai, Scarr, Schumann (BNL) I

OTHER RELATED PAPERS

SUZUKI 93	PR D47 1252	(BLB)
BAUBILLIER 82B	NP B202 21	(BIRM, CERN, GLAS, MSU, CURIN)
FERNANDEZ 82	ZPHY C16 95	+ Aguilar-Benitez+ (MADR, CERN, CDEF, STOH) JP
GAVILLET 82	ZPHY C16 119	+ Armenteros+ (CERN, CDEF, PADO, ROMA)
SHEN 66	PRL 17 726	+ Butterworth, Fu, Goldhaber, Trilling (LRL)
ALMEIDA 65	PL 16 184	+ Goldhaber (LRL)
ARMENTEROS 64	PL 9 207	+ Atherton, Byer, Dornan, Forson+ (CAVE)
ARMENTEROS 64B	PR 145 1095	+ Edwards, D'Andlau+ (CERN, CDEF)
ARMENTEROS 64B	Dubna Conf. 1 577	+ Barash, Kirsch, Miller, Tan (COLU)
Also 64C	Dubna Conf. 1 617	+ Edwards, D'Andlau+ (CERN, CDEF)

$K_1(1400)$

$$I(J^P) = \frac{1}{2}(1^+)$$

$K_1(1400)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1402 ± 7 OUR AVERAGE				
1373 ± 14 ± 18	1 ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1392 ± 18	BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
1410 ± 25	DAUM	81C CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
1415 ± 15	ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1404.0 ± 10.0	2 CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
~ 1350	3 TORNQVIST	82B	RVUE	
~ 1400.0	VERGEEST	79	HBC	- 4.2 $K^- p \rightarrow (K\pi\pi)^- p$
~ 1400	BRANDENB...	76	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$
1420	DAVIS	72	HBC	+ 12 $K^+ p$
1368 ± 18	FIRESTONE	72B	DBC	+ 12 $K^+ d$

1 From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.
 2 From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.
 3 From a unitarized quark-model calculation.

$K_1(1400)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
174 ± 13 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
188 ± 54 ± 60	4 ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
276 ± 65	BAUBILLIER	82B	HBC	0 8.25 $K^- p \rightarrow K_S^0 \pi^+ \pi^- n$
195 ± 25	DAUM	81C CNTR	-	63 $K^- p \rightarrow K^- 2\pi p$
180 ± 10	ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
142.0 ± 16.0	5 CARNEGIE	77	ASPK	± 13 $K^\pm p \rightarrow (K\pi\pi)^\pm p$

See key on page 1343

Meson Full Listings


$K_1(1400)$, $K^*(1410)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

~ 200.0	VERGEEST	79	HBC	-	4.2	$K^- p \rightarrow (\bar{K} \pi \pi)^- p$
~ 160	BRANDENB...	76	ASPK	±	13	$K^\pm p \rightarrow (K \pi \pi)^\pm p$
80	DAVIS	72	HBC	+	12	$K^+ p$
241 ± 30	FIRESTONE	72b	DBC	+	12	$K^+ d$

⁴ From partial-wave analysis of $K^0 \pi^+ \pi^-$ system.

⁵ From a model-dependent fit with Gaussian background to BRANDENBURG 76 data.



$K_1(1400)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	(94 ± 6 %) %
Γ_2 $K\rho$	(3.0±3.0) %
Γ_3 $K f_0(1300)$	(2.0±2.0) %
Γ_4 $K\omega$	(1.0±1.0) %
Γ_5 $K_0^*(1430)\pi$	

$K_1(1400)$ PARTIAL WIDTHS

$\Gamma(K^*(892)\pi)$	Γ_1
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
117.0 ± 10.0	CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$

$\Gamma(K\rho)$	Γ_2
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
2.0 ± 1.0	CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$

$\Gamma(K\omega)$	Γ_4
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
23.0 ± 12.0	CARNEGIE 77 ASPK ± 13 $K^\pm p \rightarrow (K \pi \pi)^\pm p$

$K_1(1400)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{total}$	Γ_1/Γ
VALUE	DOCUMENT ID TECN COMMENT
0.94 ± 0.06	⁶ DAUM 81c CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K\rho)/\Gamma_{total}$	Γ_2/Γ
VALUE	DOCUMENT ID TECN COMMENT
0.03 ± 0.03	⁶ DAUM 81c CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K f_0(1300))/\Gamma_{total}$	Γ_3/Γ
VALUE	DOCUMENT ID TECN COMMENT
0.02 ± 0.02	⁶ DAUM 81c CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K\omega)/\Gamma_{total}$	Γ_4/Γ
VALUE	DOCUMENT ID TECN COMMENT
0.01 ± 0.01	⁶ DAUM 81c CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K_0^*(1430)\pi)/\Gamma_{total}$	Γ_5/Γ
VALUE	DOCUMENT ID TECN COMMENT
~ 0.00	⁶ DAUM 81c CNTR 63 $K^- p \rightarrow K^- 2\pi p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

D-wave/S-wave RATIO FOR $K_1(1400) \rightarrow K^*(892)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.04 ± 0.01	⁶ DAUM	81c	CNTR 63 $K^- p \rightarrow K^- 2\pi p$

⁶ Average from low and high t data.

$K_1(1400)$ REFERENCES

ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	82b	NP B202 21	+	(BIRM, CERN, GLAS, MSU, CURIN)
TORNQVIST	82b	NP B202 268		(HELS)
DAUM	81c	NP B187 1	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP
VERGEEST	79	NP B158 265	+Jongejans, Dionisi+	(NIJM, AMST, CERN, OXF)
CARNEGIE	77	NP B127 509	+Cashmore, Davier, Dunwoodie, Lasinski+	(SLAC)
BRANDENB...	76	PRL 26 703	Brandenburg, Carnegie, Cashmore+	(SLAC) JP
DAVIS	72	PR D5 2688	+Alston-Garnjost, Barbaro, Flatte, Friedman, Lynch+	(LBL)
FIRESTONE	72b	PR D5 505	+Goldhaber, Lissauer, Trilling	(LBL)

OTHER RELATED PAPERS

SUZUKI	93	PR D47 1252		(LBL)
FERNANDEZ	82	ZPHY C16 95	+Aguilar-Benitez+	(MADR, CERN, CDEF, STOH)
SHEN	66	PRL 17 726	+Butterworth, Fu, Goldhaber, Trilling	(LRL)
Also	66	Private Comm.	Goldhaber	(LRL)
ALMEIDA	65	PL 16 184	+Atherton, Byer, Dornan, Forson+	(CAVE)
ARMENTEROS	64	PL 9 207	+Edwards, D'Andiau+	(CERN, CDEF)
Also	66	PR 145 1095	Barash, Kirsch, Miller, Tan	(COLU)
ARMENTEROS	64b	Dubna Conf. 1 577	+Edwards, D'Andiau+	(CERN, CDEF)
Also	64c	Dubna Conf. 1 617	Armenteros	

$K^*(1410)$

$$J(P) = \frac{1}{2}(1^-)$$

$K^*(1410)$ MASS

All from partial wave amplitude analyses.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1412 ± 12 OUR AVERAGE	Error includes scale factor of 1.1.			
1367 ± 54	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1380 ± 21 ± 19	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
1420 ± 7 ± 10	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1474 ± 25	BAUBILLIER	82b	HBC	0 8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
1500 ± 30	ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$K^*(1410)$ WIDTH

All from partial wave amplitude analyses.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
227 ± 22 OUR AVERAGE	Error includes scale factor of 1.1.			
114 ± 101	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
176 ± 52 ± 22	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
240 ± 18 ± 12	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
275 ± 65	BAUBILLIER	82b	HBC	0 8.25 $K^- p \rightarrow \bar{K}^0 2\pi n$
500 ± 100	ETKIN	80	MPS	0 6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

$K^*(1410)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $K^*(892)\pi$	> 40 %	95%
Γ_2 $K\pi$	(6.6±1.3) %	
Γ_3 $K\rho$	< 7 %	95%

$K^*(1410)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	Γ_3/Γ_1
VALUE	DOCUMENT ID TECN CHG COMMENT
< 0.17	95 ASTON 84 LASS 0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$	Γ_2/Γ_1
VALUE	DOCUMENT ID TECN CHG COMMENT
< 0.16	95 ASTON 84 LASS 0 11 $K^- p \rightarrow \bar{K}^0 2\pi n$

$\Gamma(K\pi)/\Gamma_{total}$	Γ_2/Γ
VALUE	DOCUMENT ID TECN CHG COMMENT
0.066 ± 0.010 ± 0.008	ASTON 88 LASS 0 11 $K^- p \rightarrow K^- \pi^+ n$

$K^*(1410)$ REFERENCES

BIRD	89	SLAC-332		(SLAC)
ASTON	88	NP B296 493	+Awaji, Blenz, Bird+	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	+Carnegie, Dunwoodie+	(SLAC, CARL, OTTA) JP
BAUBILLIER	82b	NP B202 21	+	(BIRM, CERN, GLAS, MSU, CURIN)
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP

Meson Full Listings

 $K_0^*(1430)$, $K_2^*(1430)$ $K_0^*(1430)$

$$I(J^P) = \frac{1}{2}(0^+)$$

Our latest mini-review on this particle can be found in the 1984 edition.

 $K_0^*(1430)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
1429 ± 4 ± 5	¹ ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 1420	BAUBILLIER	84B	HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 1425	^{2,3} ESTABROOKS	78	ASPK		13 $K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$
~ 1450.0	MARTIN	78	SPEC		10 $K^\pm p \rightarrow K_S^0 \pi p$

¹ Uses a model for the background, without this background they get a mass 1340 MeV, where the phase shift passes 90°.

² Mass defined by pole position.

³ From elastic $K\pi$ partial-wave analysis.

 $K_0^*(1430)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
287 ± 10 ± 21	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 200	BAUBILLIER	84B	HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
200 to 300	⁴ ESTABROOKS	78	ASPK		13 $K^\pm p \rightarrow K^\pm \pi^\pm (n, \Delta)$

⁴ From elastic $K\pi$ partial-wave analysis.

 $K_0^*(1430)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(93 ± 10) %

 $K_0^*(1430)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ	
0.93 ± 0.04 ± 0.09	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	

 $K_0^*(1430)$ REFERENCES

ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
BAUBILLIER	84B	ZPHY C26 37	+	(BIRM, CERN, GLAS, MICH, CURIN)
ESTABROOKS	78	NP B133 490	+Carnegie+	(MCGI, CARL, DURH, SLAC)
MARTIN	78	NP B134 392	+Shimada, Baldi, Bohringer+	(DURH, GEVA)

OTHER RELATED PAPERS

TORNQVIST	82	PRL 49 624		(HELS)
GOLDBERG	69	PL 30B 434	+Huffer, Laloum+	(SABRE Collab.)
SCHLEIN	69	Argonne Conf. 446		(UCLA)
TRIPPE	68	PL 28B 203	+Chien, Malamud, Mellem, Schlein+	(UCLA)

 $K_2^*(1430)$

$$I(J^P) = \frac{1}{2}(2^+)$$

We consider that phase-shift analyses provide more reliable determinations of the mass and width.

 $K_2^*(1430)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
1425.4 ± 1.3 OUR AVERAGE	Error includes scale factor of 1.1.					
1423.4 ± 2 ± 3	24809 ± 820	¹ BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1420 ± 4	1587	BAUBILLIER	84B	HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1436 ± 5.5	400	^{2,3} CLELAND	82	SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1500	^{2,3} CLELAND	82	SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
1430 ± 3.2	1200	^{2,3} CLELAND	82	SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$
1423.0 ± 5.0	935	TOAFF	81	HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1428.0 ± 4.6		⁴ MARTIN	78	SPEC	+	10 $K^\pm p \rightarrow K_S^0 \pi p$
1423.8 ± 4.6		⁴ MARTIN	78	SPEC	-	10 $K^\pm p \rightarrow K_S^0 \pi p$
1420.0 ± 3.1	1400	AGUILAR-...	71B	HBC	-	3.9,4.6 $K^- p$
1425 ± 8.0	225	^{2,3} BARNHAM	71C	HBC	+	$K^+ p \rightarrow K^0 \pi^+ p$
1416.0 ± 10.0	220	CRENNELL	69D	DBC	-	3.9 $K^- N \rightarrow \bar{K}^0 \pi^- N$
1414 ± 13.0	60	² LIND	69	HBC	+	9 $K^+ p \rightarrow K^0 \pi^+ p$
1427.0 ± 12.0	63	² SCHWEING...	68	HBC	-	5.5 $K^- p \rightarrow \bar{K} \pi N$
1423 ± 11.0	39	² BASSANO	67	HBC	-	4.6-5.0 $K^- p \rightarrow \bar{K}^0 \pi^- p$

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
1432.4 ± 1.3 OUR AVERAGE						
1431.2 ± 1.8 ± 0.7		⁵ ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
1434 ± 4 ± 6		⁵ ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1433 ± 6 ± 10		⁵ ASTON	84B	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
1471 ± 12		⁵ BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
1428 ± 3		⁵ ASTON	81C	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
1434.0 ± 2.0		⁵ ESTABROOKS	78	ASPK	0	13 $K^\pm p \rightarrow p K \pi$
1440.0 ± 10.0		⁵ BOWLER	77	DBC	0	5.5 $K^+ d \rightarrow K \pi p p$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
1420.0 ± 7.0	300	HENDRICK	76	DBC		8.25 $K^+ N \rightarrow K^+ \pi N$
1421.6 ± 4.2	800	MCCUBBIN	75	HBC	0	3.6 $K^- p \rightarrow K^- \pi^+ n$
1420.1 ± 4.3		⁶ LINGLIN	73	HBC	0	2-13 $K^+ p \rightarrow K^+ \pi^- X$
1419.1 ± 3.7	1800	AGUILAR-...	71B	HBC	0	3.9,4.6 $K^- p$
1416 ± 6	600	CORDS	71	DBC	0	9 $K^+ n \rightarrow K^+ \pi^- p$
1421.1 ± 2.6	2200	DAVIS	69	HBC	0	12 $K^+ p \rightarrow K^+ \pi^- X$

¹ From a partial wave amplitude analysis.

² Errors enlarged by us to Γ/\sqrt{N} ; see the note with the $K^*(892)$ mass.

³ Number of events in peak re-evaluated by us.

⁴ Systematic error added by us.

⁵ From phase shift or partial-wave analysis.

⁶ From pole extrapolation, using world $K^+ p$ data summary tape.

 $K_2^*(1430)$ WIDTHCHARGED ONLY, WITH FINAL STATE $K\pi$


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
98.4 ± 2.3 OUR FIT						
98.4 ± 2.4 OUR AVERAGE						
98 ± 4 ± 4	24809 ± 820	⁷ BIRD	89	LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
109 ± 22	400	^{8,9} CLELAND	82	SPEC	+	30 $K^+ p \rightarrow K_S^0 \pi^+ p$
124 ± 12.8	1500	^{8,9} CLELAND	82	SPEC	+	50 $K^+ p \rightarrow K_S^0 \pi^+ p$
113 ± 12.8	1200	^{8,9} CLELAND	82	SPEC	-	50 $K^+ p \rightarrow K_S^0 \pi^- p$
85.0 ± 16.0	935	TOAFF	81	HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
96.5 ± 3.8		MARTIN	78	SPEC	+	10 $K^\pm p \rightarrow K_S^0 \pi p$
97.7 ± 4.0		MARTIN	78	SPEC	-	10 $K^\pm p \rightarrow K_S^0 \pi p$
94.7 ± 15.1 - 12.5	1400	AGUILAR-...	71B	HBC	-	3.9,4.6 $K^- p$

NEUTRAL ONLY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	
109 ± 5 OUR AVERAGE	Error includes scale factor of 1.9. See the ideogram below.					
116.5 ± 3.6 ± 1.7		¹⁰ ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
129 ± 15 ± 15		¹⁰ ASTON	87	LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
131 ± 24 ± 20		¹⁰ ASTON	84B	LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$
143 ± 34		¹⁰ BAUBILLIER	82B	HBC	0	8.25 $K^- p \rightarrow NK_S^0 \pi \pi$
98 ± 8		¹⁰ ASTON	81C	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
140 ± 30		¹⁰ ETKIN	80	SPEC	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
98.0 ± 5.0		¹⁰ ESTABROOKS	78	ASPK	0	13 $K^\pm p \rightarrow p K \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
125.0 ± 29.0	300	⁸ HENDRICK	76	DBC		8.25 $K^+ N \rightarrow K^+ \pi N$
116 ± 18	800	MCCUBBIN	75	HBC	0	3.6 $K^- p \rightarrow K^- \pi^+ n$
61.0 ± 14.0		¹¹ LINGLIN	73	HBC	0	2-13 $K^+ p \rightarrow K^+ \pi^- X$
116.6 ± 10.3 - 15.5	1800	AGUILAR-...	71B	HBC	0	3.9,4.6 $K^- p$
144 ± 24.0	600	⁸ CORDS	71	DBC	0	9 $K^+ n \rightarrow K^+ \pi^- p$
101 ± 10	2200	DAVIS	69	HBC	0	12 $K^+ p \rightarrow K^+ \pi^- \pi^+ p$

See key on page 1343

Meson Full Listings
 $K_2^*(1430)$



⁷ From a partial wave amplitude analysis.
⁸ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
⁹ Number of events in peak re-evaluated by us.
¹⁰ From phase shift or partial-wave analysis.
¹¹ From pole extrapolation, using world $K^+\rho$ data summary tape.

$K_2^*(1430)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $K\pi$	(49.7±1.2) %	
Γ_2 $K^*(892)\pi$	(25.2±1.7) %	
Γ_3 $K^*(892)\pi\pi$	(13.0±2.3) %	
Γ_4 $K\rho$	(8.8±0.8) %	S=1.2
Γ_5 $K\omega$	(2.9±0.8) %	
Γ_6 $K^+\gamma$	(2.4±0.5) × 10 ⁻³	
Γ_7 $K\eta$	(1.4 ^{+2.8} _{-0.9}) × 10 ⁻³	S=1.1
Γ_8 $K\omega\pi$	< 7.2 × 10 ⁻⁴	CL=95%
Γ_9 $K^0\gamma$	< 9 × 10 ⁻⁴	CL=90%

CONSTRAINED FIT INFORMATION

An overall fit to the total width, a partial width, and 10 branching ratios uses 28 measurements and one constraint to determine 8 parameters. The overall fit has a $\chi^2 = 19.5$ for 21 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta p_i \delta p_j \rangle / (\delta p_i \delta p_j)$, in percent, from the fit to parameters p_i , including the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-16						
x_3	-33	-75					
x_4	-12	39	-54				
x_5	-11	-3	-25	-8			
x_6	-1	-1	-1	-1	0		
x_7	-3	-6	-4	-4	-2	0	
Γ	0	0	0	0	0	-13	0
	x_1	x_2	x_3	x_4	x_5	x_6	x_7

Mode	Rate (MeV)	Scale factor
Γ_1 $K\pi$	48.9 ± 1.7	
Γ_2 $K^*(892)\pi$	24.8 ± 1.7	
Γ_3 $K^*(892)\pi\pi$	12.8 ± 2.3	
Γ_4 $K\rho$	8.7 ± 0.8	1.2
Γ_5 $K\omega$	2.9 ± 0.8	
Γ_6 $K^+\gamma$	0.24 ± 0.04	
Γ_7 $K\eta$	0.14 ^{+0.28} _{-0.09}	1.1

$K_2^*(1430)$ PARTIAL WIDTHS

$\Gamma(K^+\gamma)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6
240 ± 40 OUR FIT					
240 ± 45	CIHANGIR	82	SPEC	+ 200 $K^+Z \rightarrow ZK^+\pi^0$, $ZK_S^0\pi^+$	

$\Gamma(K^0\gamma)$	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_9
< 84	90	CARLSMITH	87	SPEC	0 60-200 $K_L^0 A \rightarrow K_S^0\pi^0 A$	

$K_2^*(1430)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
0.497 ± 0.012 OUR FIT					
0.488 ± 0.014 OUR AVERAGE					
0.485 ± 0.006 ± 0.020	¹² ASTON	88	LASS	0	11 $K^-\rho \rightarrow K^-\pi^+\pi$
0.49 ± 0.02	¹² ESTABROOKS	78	ASPK	±	13 $K^\pm\rho \rightarrow \rho K\pi$

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_2/\Gamma_1 = \Gamma_2/(\Gamma_1+\Gamma_2+\Gamma_4)$
0.47 ± 0.10	BASSANO	67	HBC	-0	4.6, 5.0 $K^-\rho$
0.45 ± 0.13	¹³ BADIER	65C	HBC	-	3 $K^-\rho$


$\Gamma(K\rho)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	$\Gamma_4/\Gamma_1 = \Gamma_4/(\Gamma_1+\Gamma_2+\Gamma_4)$
0.14 ± 0.10	BASSANO	67	HBC	-0	4.6, 5.0 $K^-\rho$
0.14 ± 0.07	¹³ BADIER	65C	HBC	-	3 $K^-\rho$

$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
0.51 ± 0.04 OUR FIT					
0.48 ± 0.05 OUR AVERAGE					
0.44 ± 0.09	ASTON	84B	LASS	0	11 $K^-\rho \rightarrow \bar{K}^0 2\pi n$
0.62 ± 0.19	LAUSCHER	75	HBC	0	10, 16 $K^-\rho \rightarrow K^-\pi^+\pi$
0.54 ± 0.16	DEHM	74	DBC	0	4.6 K^+N
0.47 ± 0.08	AGUILAR...	71B	HBC		3.9, 4.6 $K^-\rho$

$\Gamma(K\omega)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_1
0.059 ± 0.017 OUR FIT					
0.070 ± 0.035 OUR AVERAGE					
0.05 ± 0.04	AGUILAR...	71B	HBC		3.9, 4.6 $K^-\rho$
0.13 ± 0.07	BASSOMPIE...	69	HBC	0	5 $K^+\pi$

$\Gamma(K\rho)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_1
0.178 ± 0.018 OUR FIT				Error includes scale factor of 1.2.	
0.153 ^{+0.034} _{-0.018} OUR AVERAGE					
0.18 ± 0.05	ASTON	84B	LASS	0	11 $K^-\rho \rightarrow \bar{K}^0 2\pi n$
0.02 ^{+0.10} _{-0.02}	DEHM	74	DBC	0	4.6 K^+N
0.16 ± 0.05	AGUILAR...	71B	HBC		3.9, 4.6 $K^-\rho$

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_2
0.351 ± 0.032 OUR FIT				Error includes scale factor of 1.5.	
0.354 ± 0.033 OUR AVERAGE				Error includes scale factor of 1.4. See the Ideogram below.	
0.293 ± 0.032 ± 0.020	ASTON	87	LASS	0	11 $K^-\rho \rightarrow \bar{K}^0\pi^+\pi^-\pi$
0.38 ± 0.09	BAUBILLIER	82B	HBC	0	8.25 $K^-\rho \rightarrow NK_S^0\pi\pi$
0.39 ± 0.03	DAUM	81C	CNTR		63 $K^-\rho \rightarrow K^-2\pi\rho$



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

Meson Full Listings

 $K_2^*(1430)$, $K(1460)$

$\Gamma(K\omega)/\Gamma(K^*(892)\pi)$ Γ_5/Γ_2

VALUE DOCUMENT ID TECN CHG COMMENT
0.116 ± 0.034 OUR FIT
0.10 ± 0.04 FIELD 67 HBC - 3.8 $K^- p$

$\Gamma(K\eta)/\Gamma(K^*(892)\pi)$ Γ_7/Γ_2

VALUE DOCUMENT ID TECN CHG COMMENT
0.006 ± 0.011
-0.004 OUR FIT
0.07 ± 0.04 FIELD 67 HBC - 3.8 $K^- p$

$\Gamma(K\eta)/\Gamma(K\pi)$ Γ_7/Γ_1

VALUE CL% DOCUMENT ID TECN CHG COMMENT
0.0028 ± 0.0057
-0.0019 OUR FIT Error includes scale factor of 1.1.
0 ± 0.0056 ¹⁴ ASTON 88B LASS - 11 $K^- p \rightarrow K^- \eta p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.04 95 AGUILAR... 71B HBC 3.9,4.6 $K^- p$
 <0.065 ¹³ BASSOMPIE... 69 HBC 5.0 $K^+ p$
 <0.02 BISHOP 69 HBC 3.5 $K^+ p$

$\Gamma(K^*(892)\pi\pi)/\Gamma_{total}$ Γ_3/Γ

VALUE DOCUMENT ID TECN CHG COMMENT
0.130 ± 0.023 OUR FIT
0.12 ± 0.04 ¹⁵ GOLDBERG 76 HBC - 3 $K^- p \rightarrow \rho \bar{K}^0 \pi \pi \pi$

$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$ Γ_3/Γ_1

VALUE DOCUMENT ID TECN CHG COMMENT
0.26 ± 0.05 OUR FIT
0.21 ± 0.08 ^{13,15} JONGEJANS 78 HBC - 4 $K^- p \rightarrow \rho \bar{K}^0 \pi \pi \pi$

$\Gamma(K\omega\pi)/\Gamma_{total}$ Γ_8/Γ

VALUE (units 10^{-3}) CL% EVTS DOCUMENT ID TECN COMMENT
<0.72 95 0 JONGEJANS 78 HBC 4 $K^- p \rightarrow \rho \bar{K}^0 4\pi$

¹² From phase shift analysis.

¹³ Restated by us.

¹⁴ ASTON 88B quote < 0.0092 at CL=95%. We convert this to a central value and 1 sigma error in order to be able to use it in our constrained fit.

¹⁵ Assuming $\pi\pi$ system has isospin 1, which is supported by the data.

 $K_2^*(1430)$ REFERENCES

BIRD	89	SLAC-332			(SLAC)
ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)	
ASTON	88B	PL B201 169	+Awaji, Bienz+	(SLAC, NAGO, CINC, INUS)	
ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)	
CARLSMITH	87	PR D36 3502	+Bernstein, Bock, Coupal, Peyaud, Turley+	(EFI, SACI)	
ASTON	84B	NP B247 267	+Carnegie, Dunwoodie+	(SLAC, CARL, OTTA)	
BAUBILLIER	84B	ZPHY C26 31	+ (BIRM, CERN, GLAS, MSU, CURIN)		
BAUBILLIER	82B	NP B202 21	+ (BIRM, CERN, GLAS, MSU, CURIN)		
CHANGIR	82	PL 117B 123	+Berg, Biel, Chandlee+	(FNAL, MINN, ROCH)	
CLELAND	82	NP B208 189	+Delfosse, Dorsaz, Gloor	(DURH, GEVA, LAUS, PITT)	
ASTON	81C	PL 106B 235	+Carnegie, Dunwoodie+	(SLAC, CARL, OTTA) JP	
DAUM	81C	NP B187 1	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)		
TOAFF	81	PR D23 1500	+Musgrave, Ammar, Davis, Ecklund+	(ANL, KANS)	
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP	
ESTABROOKS	78	NP B133 490	+Carnegie+ (MCGI, CARL, DURH, SLAC)		
Also	78B	PR D17 658	Estabrooks, Carnegie+	(MCGI, CARL, DURH+)	
JONGEJANS	78	NP B139 383	+Cerrada+ (ZEEB, CERN, NIJM, OXF)		
MARTIN	78	NP B134 392	+Shimada, Baldi, Bohringer+	(DURH, GEVA)	
BOWLER	77	NP B126 31	+Dainton, Drake, Williams	(OXF)	
GOLDBERG	76	LNC 17 253		(HAIF)	
HENDRICK	76	NP B112 189	+Vignaud, Burlaud+ (MONS, SACL, PARIS, BELG)		
LAUSCHER	75	NP B86 189	+Otter, Wiczorek+ (ABCLV Collab.) JP		
MCCUBBIN	75	NP B86 13	+Lyons (OXF)		
DEHM	74	NP B75 47	+Goebel, Wittek+ (MPIM, BRUX, MONS, CERN)		
LINGLIN	73	NP B55 408		(CERN)	
AGUILAR...	71B	PR D4 2583	Aguiar-Benitez, Eisner, Kinson	(BNL)	
BARNHAM	71C	NP B28 171	+Cooley, Jobs, Griffiths, Hughes+	(BIRM, GLAS)	
CORDS	71	PR D4 1974	+Carmony, Erwin, Meiere+	(PURD, UCD, IUPU)	
BASSOMPIE...	69	NP B13 189	+Bassompierre+ (CERN, BRUX) JP		
BISHOP	69	NP B9 403	+Goshaw, Erwin, Walker	(WISC)	
CRENNELL	69D	PRL 22 487	+Karshon, Lai, O'Neill, Scarr	(BNL)	
DAVIS	69	PRL 23 1071	+Derenzo, Flatte, Garnjost, Lynch, Solmitz	(LRL)	
LIND	69	NP B14 1	+Alexander, Firestone, Fu, Goldhaber	(LRL) JP	
SCHWEING...	68	PR 166 1317	+Schweingruber, Derrick, Fields+	(ANL, NWES)	
Also	67		Schweingruber	(NWES, NWES)	
BASSANO	67	PRL 19 968	+Goldberg, Goz, Barnes, Leitner+	(BNL, SYRA)	
FIELD	67	PL 24B 638	+Hendricks, Piccioni, Yager	(UCSD)	
BADIER	65C	PL 19 612	+Demoulin, Goldberg+	(EPOL, SACL, AMST)	

OTHER RELATED PAPERS

ATKINSON	86	ZPHY C30 521	+ (BONN, CERN, GLAS, LANC, MCHS, CURIN+)		
BAUBILLIER	82B	NP B202 21	+ (BIRM, CERN, GLAS, MSU, CURIN)		
CHUNG	65	PRL 15 325	+Dahl, Hardy, Hess, Jacobs, Kirz	(LRL)	
FOCARDI	65	PL 16 351	+Ranzi, Serra+	(BGNA, SACL)	
HAQUE	65	PL 14 338	+Hague+		
HARDY	65	PRL 14 401	+Chung, Dahl, Hess, Kirz, Miller	(LRL)	

 $K(1460)$

$$I(J^P) = \frac{1}{2}(0^-)$$

OMITTED FROM SUMMARY TABLE

Observed in $K\pi\pi$ partial-wave analysis. Not seen by VERGEEST 79.
Needs confirmation.

 $K(1460)$ MASS

VALUE (MeV) DOCUMENT ID TECN CHG COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •
 ~ 1460 DAUM 81C CNTR - 63 $K^- p \rightarrow K^- 2\pi p$
 ~ 1400 ¹ BRANDENB... 76B ASPK ± 13 $K^\pm p \rightarrow K\pi\pi N$

¹ Coupled mainly to $K_0^*(1300)$. Decay into $K^*(892)\pi$ seen.

 $K(1460)$ WIDTH

VALUE (MeV) DOCUMENT ID TECN CHG COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •
 ~ 260 DAUM 81C CNTR - 63 $K^- p \rightarrow K^- 2\pi p$
 ~ 250 ² BRANDENB... 76B ASPK ± 13 $K^\pm p \rightarrow K\pi\pi N$

² Coupled mainly to $K_0^*(1300)$. Decay into $K^*(892)\pi$ seen.

 $K(1460)$ DECAY MODES

Mode

Γ_1 $K^*(892)\pi$
 Γ_2 $K\rho$
 Γ_3 $K_0^*(1430)\pi$

 $K(1460)$ PARTIAL WIDTHS

$\Gamma(K^*(892)\pi)$ Γ_1

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •
 ~ 109 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K\rho)$ Γ_2

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •
 ~ 34 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

$\Gamma(K_0^*(1430)\pi)$ Γ_3

VALUE (MeV) DOCUMENT ID TECN COMMENT

• • • We do not use the following data for averages, fits, limits, etc. • • •
 ~ 117 DAUM 81C CNTR 63 $K^- p \rightarrow K^- 2\pi p$

 $K(1460)$ REFERENCES

DAUM	81C	NP B187 1	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)
VERGEEST	79	NP B158 265	+Jongeans, Dionisi+ (NIJM, AMST, CERN, OXF)
BRANDENB...	76B	PRL 36 1239	Brandenburg, Carnegie, Cashmore+ (SLAC) JP

OTHER RELATED PAPERS

BARNES	82	PL B116 365	+Close (RHEL)
TANIMOTO	82	PL 116B 198	(BIEL)
VERGEEST	79	NP B158 265	+Jongeans, Dionisi+ (NIJM, AMST, CERN, OXF)

See key on page 1343

Meson Full Listings
 $K_2(1580)$, $K_1(1650)$, $K^*(1680)$

$K_2(1580)$

$I(J^P) = \frac{1}{2}(2^-)$

OMITTED FROM SUMMARY TABLE
 Seen in partial-wave analysis of the $K^- \pi^+ \pi^-$ system. Needs confirmation.

$K_2(1580)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1580	OTTER	79	-	10,14,16 $K^- p$

$K_2(1580)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 110	OTTER	79	-	10,14,16 $K^- p$

$K_2(1580)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K^*(892)\pi$	seen
Γ_2 $K_2^*(1430)\pi$	possibly seen

$K_2(1580)$ BRANCHING RATIOS

$\Gamma(K^*(892)\pi)/\Gamma_{total}$	Γ_1/Γ
VALUE	
seen	OTTER 79 HBC - 10,14,16 $K^- p$

$\Gamma(K_2^*(1430)\pi)/\Gamma_{total}$	Γ_2/Γ
VALUE	
possibly seen	OTTER 79 HBC - 10,14,16 $K^- p$

$K_2(1580)$ REFERENCES

OTTER 79 NP B147 1 +Rudolph+ (AACH3, BERL, CERN, LOIC, WIEN) JP

$K_1(1650)$

$I(J^P) = \frac{1}{2}(1^+)$

OMITTED FROM SUMMARY TABLE
 This entry contains various peaks in strange meson systems ($K^+ \phi$, $K \pi \pi$) reported in partial-wave analysis in the 1600–1900 mass region.

$K_1(1650)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1650±50	FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1840	ARMSTRONG	83	OMEG -	18.5 $K^- p \rightarrow 3K p$
~ 1800	DAUM	81c	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

$K_1(1650)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
150±50	FRAME	86	OMEG +	13 $K^+ p \rightarrow \phi K^+ p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 250	DAUM	81c	CNTR -	63 $K^- p \rightarrow K^- 2\pi p$

$K_1(1650)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi \pi$	
Γ_2 $K \phi$	

$K_1(1650)$ REFERENCES

FRAME 86 NP B276 667 +Hughes, Lynch, Minto, McFadzean+ (GLAS)
 ARMSTRONG 83 NP B221 1 + (BARI, BIRM, CERN, MILA, CURIN+)
 DAUM 81c NP B187 1 +Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)

$K^*(1680)$

$I(J^P) = \frac{1}{2}(1^-)$

$K^*(1680)$ MASS


All from partial wave amplitude analyses.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1714±20 OUR AVERAGE				Error includes scale factor of 1.1.
1678±64	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1677±10±32	ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
1735±10±20	ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1800±70	ETKIN	80	MPS 0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
~ 1650	ESTABROOKS	78	ASPK 0	13 $K^\pm p \rightarrow K^\pm \pi^\pm n$

$K^*(1680)$ WIDTH

All from partial wave amplitude analyses.

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
323±110 OUR AVERAGE				Error includes scale factor of 4.2. See the ideogram below.
454±270	BIRD	89	LASS -	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
205±16±34	ASTON	88	LASS 0	11 $K^- p \rightarrow K^- \pi^+ n$
423±18±30	ASTON	87	LASS 0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
170±30	ETKIN	80	MPS 0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
250 to 300	ESTABROOKS	78	ASPK 0	13 $K^\pm p \rightarrow K^\pm \pi^\pm n$



$K^*(1680)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi$	(38.7±2.5) %
Γ_2 $K \rho$	(31.4+4.7-2.1) %
Γ_3 $K^*(892)\pi$	(29.9+2.2-4.7) %

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 4 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 2.9$ for 2 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-36	
x_3	-39	-72
	x_1	x_2

Meson Full Listings

 $K^*(1680)$, $K_2(1770)$ $K^*(1680)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
VALUE					
0.387 ± 0.026 OUR FIT					
$0.388 \pm 0.014 \pm 0.022$	ASTON	88	LASS	0	11 $K^-p \rightarrow K^- \pi^+ n$
$\Gamma(K\pi)/\Gamma(K^*(892)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ_3
VALUE					
$1.30^{+0.23}_{-0.14}$ OUR FIT					
2.8 ± 1.1	ASTON	84	LASS	0	11 $K^-p \rightarrow \bar{K}^0 2\pi n$
$\Gamma(K\rho)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
VALUE					
$0.81^{+0.14}_{-0.09}$ OUR FIT					
1.2 ± 0.4	ASTON	84	LASS	0	11 $K^-p \rightarrow \bar{K}^0 2\pi n$
$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_3
VALUE					
$1.05^{+0.27}_{-0.11}$ OUR FIT					
$0.97 \pm 0.09^{+0.30}_{-0.10}$	ASTON	87	LASS	0	11 $K^-p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

 $K^*(1680)$ REFERENCES

BIRD	89	SLAC-332		(SLAC)
ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
ASTON	84	PL 149B 258	+Carnegie, Dunwoodie+	(SLAC, CARL, OTTA) JP
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP
ESTABROOKS	78	NP B133 490	+Carnegie+	(MCGI, CARL, DURH, SLAC) JP

 $K_2(1770)$ was $L(1770)$

$$J^P = \frac{1}{2}(2^-)$$

NOTE ON THE $K_2(1770)$ AND THE $K_2(1820)$

A partial-wave analysis of the $K^- \omega$ system based on about 100,000 $K^-p \rightarrow K^- \omega p$ events (ASTON 93) provides evidence for two $q\bar{q}$ D -wave states near 1.8 GeV. A previous analysis based on about 200,000 diffractively produced $K^-p \rightarrow K^- \pi^+ \pi^- p$ events (DAUM 81) gave evidence for two $J^P = 2^-$ states in this region, with masses ~ 1780 MeV and ~ 1840 MeV and widths ~ 200 MeV, in good agreement with ASTON 93. In contrast, the masses obtained using a single resonance do not agree well: ASTON 93 obtains 1728 ± 7 MeV, while DAUM 81 estimates ~ 1820 MeV.

We list under the $K_2(1770)$ other measurements that do not resolve the two-resonance structure of the enhancement (previously called $L(1770)$).

 $K_2(1770)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1773 ± 8		¹ ASTON	93	LASS	11 $K^-p \rightarrow K^- \omega p$
~ 1780		² DAUM	81c	CNTR	63 $K^-p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1810 ± 20		FRAME	86	OMEG	+ 13 $K^+p \rightarrow \phi K^+ p$
~ 1730		ARMSTRONG	83	OMEG	- 18.5 $K^-p \rightarrow 3Kp$
1710 ± 15	60	CHUNG	74	HBC	- 7.3 $K^-p \rightarrow K^- \omega p$
1767 ± 6		BLIEDEN	72	MMS	- 11-16 K^-p
1730 ± 20	306	³ FIRESTONE	72B	DBC	+ 12 K^+d
1765.0 ± 40.0		⁴ COLLEY	71	HBC	+ 10 $K^+p \rightarrow K_2 \pi N$
1740.0		DENEGRI	71	DBC	- 12.6 $K^-d \rightarrow \bar{K} 2\pi d$
1745.0 ± 20.0		AGUILAR...	70c	HBC	- 4.6 K^-p
1780.0 ± 15.0		BARTSCH	70c	HBC	- 10.1 K^-p
1760.0 ± 15.0		LU DLAM	70	HBC	- 12.6 K^-p

¹ From a partial wave analysis of the $K^- \omega$ system.² From a partial wave analysis of the $K^- 2\pi$ system.³ Produced in conjunction with excited deuteron.⁴ Systematic errors added correspond to spread of different fits. $K_2(1770)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
186 ± 14		⁵ ASTON	93	LASS	11 $K^-p \rightarrow K^- \omega p$
~ 210		⁶ DAUM	81c	CNTR	63 $K^-p \rightarrow K^- 2\pi p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
140 ± 40		FRAME	86	OMEG	+ 13 $K^+p \rightarrow \phi K^+ p$
~ 220		ARMSTRONG	83	OMEG	- 18.5 $K^-p \rightarrow 3Kp$
110 ± 50	60	CHUNG	74	HBC	- 7.3 $K^-p \rightarrow K^- \omega p$
100 ± 26		BLIEDEN	72	MMS	- 11-16 K^-p
210 ± 30	306	⁷ FIRESTONE	72B	DBC	+ 12 K^+d
90 ± 70		⁸ COLLEY	71	HBC	+ 10 $K^+p \rightarrow K_2 \pi N$
130.0		DENEGRI	71	DBC	- 12.6 $K^-d \rightarrow \bar{K} 2\pi d$
100.0 ± 50.0		AGUILAR...	70c	HBC	- 4.6 K^-p
138.0 ± 40.0		BARTSCH	70c	HBC	- 10.1 K^-p
$50.0^{+40.0}_{-20.0}$		LU DLAM	70	HBC	- 12.6 K^-p

⁵ From a partial wave analysis of the $K^- \omega$ system.⁶ From a partial wave analysis of the $K^- 2\pi$ system.⁷ Produced in conjunction with excited deuteron.⁸ Systematic errors added correspond to spread of different fits. $K_2(1770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi\pi$	
Γ_2 $K_2^*(1430)\pi$	dominant
Γ_3 $K^*(892)\pi$	seen
Γ_4 $K f_2(1270)$	seen
Γ_5 $K\phi$	seen
Γ_6 $K\omega$	seen

 $K_2(1770)$ BRANCHING RATIOS

For discussion of the experimental evidence on other decay modes, see HUGHES 71, SLATTERY 71, EISNER 74.

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
VALUE					
$(K_2^*(1430) \rightarrow K\pi)$					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 0.03	DAUM	81c	CNTR	63 $K^-p \rightarrow K^- 2\pi p$	
~ 1.0	⁹ FIRESTONE	72B	DBC	+ 12 K^+d	
<1.0	COLLEY	71	HBC	10 K^+p	
0.2 ± 0.2	AGUILAR...	70c	HBC	- 4.6 K^-p	
<1.0	BARTSCH	70c	HBC	- 10.1 K^-p	
1.0	BARBARO...	69	HBC	+ 12.0 K^+p	
⁹ Produced in conjunction with excited deuteron.					
$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1	
VALUE					
~ 0.23	DAUM	81c	CNTR	63 $K^-p \rightarrow K^- 2\pi p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$\Gamma(K f_2(1270))/\Gamma(K\pi\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1	
VALUE					
$(f_2(1270) \rightarrow \pi\pi)$					
~ 0.74	DAUM	81c	CNTR	63 $K^-p \rightarrow K^- 2\pi p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$\Gamma(K\phi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ
VALUE					
seen	ARMSTRONG	83	OMEG	- 18.5 $K^-p \rightarrow K^- \phi N$	
$\Gamma(K\omega)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ
VALUE					
seen	OTTER	81	HBC	\pm 8.25, 10.16 $K^\pm p$	
seen	CHUNG	74	HBC	- 7.3 $K^-p \rightarrow K^- \omega p$	

 $K_2(1770)$ REFERENCES

ASTON	93	PL B308 186	+Bienez, Bird+	(SLAC, NAGO, CINC, INUS)
FRAME	86	NP B276 667	+Hughes, Lynch, Minto, McFadzean+	(GLAS)
ARMSTRONG	83	NP B221 1	+ (BARI, BIRM, CERN, MILA, CURIN+)	
DAUM	81c	NP B187 1	+Hertzberger+ (AMST, CERN, CRAC, MPIM, OXF+)	
OTTER	81	NP B181 1	+ (AAACH, BERL, LOIC, VIEN, BIRM, BELG, CERN+)	
CHUNG	74	PL 51B 413	+Eisner, Protopeouscu, Samios, Strand	(BNL)
EISNER	74	Boston Conf. 140		(BNL)
BLIEDEN	72	PL 39B 668	+Finocchiaro, Bowen, Earles+	(STON, NEAS)
FIRESTONE	72B	PR D5 505	+Goldhaber, Lissauer, Trilling	(LBL)
COLLEY	71	NP B26 71	+Jobes, Kenyon, Pathak, Hughes+	(BIRM, GLAS)
DENEGRI	71	NP B28 13	+Antich, Callahan, Carson, Chien, Cox+	(JHU) JP
HUGHES	71	Bologna Conf. 293		(GLAS)
SLATTERY	71	UR-875-332		(ROCH)
AGUILAR...	70c	PR L 25 54	Aguiar-Benitez, Barnes, Bassano, Chung+	(BNL)
BARTSCH	70c	PL 33B 186	+Deutschmann+ (AAACH, BERL, CERN, LOIC, VIEN)	
LU DLAM	70	PR D2 1234	+Sandweiss, Slaughter	(YALE)
BARBARO...	69	PR L 22 1207	Barbaro-Galtieri, Davis, Flatte+	(LRL)

See key on page 1343

Meson Full Listings
 $K_2(1770), K_3^*(1780)$

OTHER RELATED PAPERS

BERLINGHIERI 67	PRL 18 1087	+Farber, Ferbel, Forman	(ROCH)
CARMONY 67	PRL 18 615	+Hendricks, Lander	(UCSD)
JOBS 67	PL 26B 49	+Bassompierre, DeBaere+	(BIRM, CERN, BRUX)
BARTSCH 66	PL 22 357	+Deutschmann+	(AACH, BERL, CERN+)

$K_3^*(1780)$

$I(J^P) = \frac{1}{2}(3^-)$


Our latest mini-review on this particle can be found in the 1984 edition.

$K_3^*(1780)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1770 ± 10 OUR AVERAGE		Error includes scale factor of 1.7. See the ideogram below.			
1720 ± 10 ± 15	6111 ± 780	¹ BIRD	89 LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1781 ± 8 ± 4		² ASTON	88 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
1740 ± 14 ± 15		² ASTON	87 LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1779.0 ± 11.0		³ BALDI	76 SPEC	+	10 $K^+ p \rightarrow K^0 \pi^+ p$
1776 ± 26		⁴ BRANDENB...	76D ASPK	0	13 $K^\pm p \rightarrow K^\pm \pi^\mp n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1749 ± 10		ASTON	88B LASS	-	11 $K^- p \rightarrow K^- \eta p$
1780.0 ± 9.0	300	BAUBILLIER	84B HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1790.0 ± 15.0		BAUBILLIER	82B HBC	0	8.25 $K^- p \rightarrow K_S^0 2\pi N$
1784.0 ± 9.0	2060	CLELAND	82 SPEC	±	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
1786 ± 15		⁵ ASTON	81D LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
1762.0 ± 9.0	190	TOAFF	81 HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1850 ± 50		ETKIN	80 MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
1812.0 ± 28.0		BEUSCH	78 OMEG		10 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
1786.0 ± 8.0		CHUNG	78 MPS	0	6 $K^- p \rightarrow K^- \pi^+ n$

- ¹ From a partial wave amplitude analysis.
- ² From energy-independent partial-wave analysis.
- ³ From a fit to Y_2^0 moment. $J^P = 3^-$ found.
- ⁴ Confirmed by phase shift analysis of ESTABROOKS 78, yields $J^P = 3^-$.
- ⁵ From a fit to the Y_2^0 moment.

WEIGHTED AVERAGE
 1770 ± 10 (Error scaled by 1.7)



$K_3^*(1780)$ mass (MeV)

$K_3^*(1780)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
164 ± 17 OUR AVERAGE		Error includes scale factor of 1.1.			
187 ± 31 ± 20	6111 ± 780	⁶ BIRD	89 LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
203 ± 30 ± 8		⁷ ASTON	88 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
171 ± 42 ± 20		⁷ ASTON	87 LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
135.0 ± 22.0		⁸ BALDI	76 SPEC	+	10 $K^+ p \rightarrow K^0 \pi^+ p$

• • • We do not use the following data for averages, fits, limits, etc. • • •

193 ⁺⁵¹ ₋₃₇		ASTON	88B LASS	-	11 $K^- p \rightarrow K^- \eta p$
99.0 ± 30.0	300	BAUBILLIER	84B HBC	-	8.25 $K^- p \rightarrow \bar{K}^0 \pi^- p$
~ 130.0		BAUBILLIER	82B HBC	0	8.25 $K^- p \rightarrow K_S^0 2\pi N$
191.0 ± 24.0	2060	CLELAND	82 SPEC	±	50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
225 ± 60		⁹ ASTON	81D LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$
~ 80	190	TOAFF	81 HBC	-	6.5 $K^- p \rightarrow \bar{K}^0 \pi^- p$
240 ± 50		ETKIN	80 MPS	0	6 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^-$
181.0 ± 44.0		¹⁰ BEUSCH	78 OMEG		10 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
96.0 ± 31.0		CHUNG	78 MPS	0	6 $K^- p \rightarrow K^- \pi^+ n$
270 ± 70		¹¹ BRANDENB...	76D ASPK	0	13 $K^\pm p \rightarrow K^\pm \pi^\mp n$

- ⁶ From a partial wave amplitude analysis.
- ⁷ From energy-independent partial-wave analysis.
- ⁸ From a fit to Y_2^0 moment. $J^P = 3^-$ found.
- ⁹ From a fit to Y_2^0 moment.
- ¹⁰ Errors enlarged by us to $4\Gamma/\sqrt{N}$; see the note with the $K^*(892)$ mass.
- ¹¹ ESTABROOKS 78 find that BRANDENBURG 76D data are consistent with 175 MeV width. Not averaged.

$K_3^*(1780)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $K\rho$	(45 ± 4) %	S=1.4
Γ_2 $K^*(892)\pi$	(27.3 ± 3.2) %	S=1.5
Γ_3 $K\pi$	(19.3 ± 1.0) %	
Γ_4 $K\eta$	(8.0 ± 1.5) %	S=1.4
Γ_5 $K_2^*(1430)\pi$	< 21 %	CL=95%

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 5 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 2.2$ for 2 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{total}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-84		
x_3	-33	-4	
x_4	-35	-14	26
	x_1	x_2	x_3

$K_3^*(1780)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$		Γ_1/Γ_2			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
1.66 ± 0.31 OUR FIT	Error includes scale factor of 1.5.				
1.52 ± 0.21 ± 0.10	ASTON	87 LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	
$\Gamma(K^*(892)\pi)/\Gamma(K\pi)$		Γ_2/Γ_3			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
1.42 ± 0.19 OUR FIT	Error includes scale factor of 1.4.				
1.09 ± 0.26	ASTON	84B LASS	0	11 $K^- p \rightarrow \bar{K}^0 2\pi n$	
$\Gamma(K\pi)/\Gamma_{total}$		Γ_3/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.193 ± 0.010 OUR FIT					
0.188 ± 0.010 OUR AVERAGE					
0.187 ± 0.008 ± 0.008	ASTON	88 LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	
0.19 ± 0.02	ESTABROOKS 78	ASPK	0	13 $K^\pm p \rightarrow K\pi N$	
$\Gamma(K\eta)/\Gamma(K\pi)$		Γ_4/Γ_3			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.41 ± 0.07 OUR FIT	Error includes scale factor of 1.5.				
0.41 ± 0.050	¹² BIRD	89 LASS	-	11 $K^- p \rightarrow \bar{K}^0 \pi^- p$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.50 ± 0.18	ASTON	88B LASS	-	11 $K^- p \rightarrow K^- \eta p$	
¹² This result supersedes ASTON 88B.					
$\Gamma(K_2^*(1430)\pi)/\Gamma(K^*(892)\pi)$		Γ_5/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 0.78	95	ASTON	87 LASS	0	11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

Meson Full Listings

 $K_3^*(1780)$, $K_2(1820)$, $K(1830)$, $K_0^*(1950)$ $K_3^*(1780)$ REFERENCES

BIRD	89	SLAC-332		(SLAC)
ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
ASTON	88B	PL B201 169	+Awaji, Bienz+	(SLAC, NAGO, CINC, INUS) JP
ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
ASTON	84B	NP B247 261	+Carnegie, Dunwoodie+	(SLAC, CARL, OTTA)
BAUBILLIER	84B	ZPHY C26 37		(BIRM, CERN, GLAS, MICH, CURIN)
BAUBILLIER	82B	NP B202 21		(BIRM, CERN, GLAS, MSU, CURIN)
CLELAND	82	NP B208 189	+Delfosse, Dorsaz, Gloor	(DURH, GEVA, LAUS, PITT)
ASTON	81D	PL 99B 502	+Dunwoodie, Durkin, Fieguth+	(SLAC, CARL, OTTA) JP
TOAFF	81	PR D23 1500	+Musgrave, Ammar, Davis, Ecklund+	(ANL, KANS)
ETKIN	80	PR D22 42	+Foley, Lindenbaum, Kramer+	(BNL, CUNY) JP
BEUSCH	78	PL 74B 282	+Birman, Konigs, Otter+	(CERN, AACH3, ETH) JP
CHUNG	78	PRL 40 355	+Etkin+	(BNL, BRAN, CUNY, MASA, PENN) JP
ESTABROOKS	78	NP B133 490	+Carnegie+	(MCGI, CARL, DURH, SLAC) JP
Also	78B	PR D17 658	Estabrooks, Carnegie+	(MCGI, CARL, DURH+)
BALDI	76	PL 63B 344	+Boehringer, Dorsaz, Hungerbuhler+	(GEVA) JP
BRANDENB...	76D	PL 60B 478	+Brandenburg, Carnegie, Cashmore+	(SLAC) JP

OTHER RELATED PAPERS

AGUILAR...	73	PRL 30 672	Aguiar-Benitez, Chung, Eisner+	(BNL)
WALUCH	73	PR D8 2837	+Flatte, Friedman	(LBL)
CARMONY	71	PRL 27 1160	+Cords, Clopp, Erwin, Meiere+	(PURD, UCD, IUPU)
FIRESTONE	71	PL 36B 513	+Goldhaber, Lissauer, Trilling	(LBL)

 $K_2(1820)$

$I(J^P) = \frac{1}{2}(2^-)$

Observed by ASTON 93 from a partial wave analysis of the $K^- \omega$ system. See Minireview under $K_2(1770)$. Needs confirmation.

 $K_2(1820)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1816 ± 13	¹ ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
~ 1840	² DAUM	81c CNTR	$63K^- p \rightarrow K^- 2\pi p$

- ¹ From a partial wave analysis of the $K^- \omega$ system.
² From a partial wave analysis of the $K^- 2\pi$ system.

 $K_2(1820)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
276 ± 35	³ ASTON	93 LASS	$11K^- p \rightarrow K^- \omega p$
~ 230	⁴ DAUM	81c CNTR	$63K^- p \rightarrow K^- 2\pi p$

- ³ From a partial wave analysis of the $K^- \omega$ system.
⁴ From a partial wave analysis of the $K^- 2\pi$ system.

 $K_2(1820)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\phi$	possibly seen
Γ_2 $K\pi\pi$	
Γ_3 $K_2^*(1430)\pi$	seen
Γ_4 $K^*(892)\pi$	seen
Γ_5 $Kf_2(1270)$	seen
Γ_6 $K\omega$	seen

 $K_2(1820)$ BRANCHING RATIOS

$\Gamma(K_2^*(1430)\pi)/\Gamma(K\pi\pi)$	Γ_3/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.77	ASTON	93 CNTR	$63K^- p \rightarrow \bar{K}2\pi p$

$\Gamma(K^*(892)\pi)/\Gamma(K\pi\pi)$	Γ_4/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.05	ASTON	93 CNTR	$63K^- p \rightarrow \bar{K}2\pi p$

$\Gamma(Kf_2(1270))/\Gamma(K\pi\pi)$	Γ_5/Γ_2		
VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 0.05	ASTON	93 CNTR	$63K^- p \rightarrow \bar{K}2\pi p$

 $K_2(1820)$ REFERENCES

ASTON	93	PL B308 186	+Bienez, Bird+	(SLAC, NAGO, CINC, INUS)
DAUM	81c	NP B187 1	+Hertzberger+	(AMST, CERN, CRAC, MPIM, OXF+)

 $K(1830)$

$I(J^P) = \frac{1}{2}(0^-)$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of $K^- \phi$ system. Needs confirmation.

 $K(1830)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 1830.0	ARMSTRONG 83	OMEG -		$18.5K^- p \rightarrow 3Kp$

 $K(1830)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
~ 250.0	ARMSTRONG 83	OMEG -		$18.5K^- p \rightarrow 3Kp$

 $K(1830)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\phi$	

 $K(1830)$ REFERENCES

ARMSTRONG 83	NP B221 1			(BARI, BIRM, CERN, MILA, CURIN+) JP
--------------	-----------	--	--	-------------------------------------

 $K_0^*(1950)$

$I(J^P) = \frac{1}{2}(0^+)$

OMITTED FROM SUMMARY TABLE

Seen in partial-wave analysis of the $K^- \pi^+$ system. Needs confirmation.

 $K_0^*(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
1945 ± 10 ± 20	¹ ASTON	88 LASS	0	$11K^- p \rightarrow K^- \pi^+ n$

- ¹ We take the central value of the two solutions and the larger error given.

 $K_0^*(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
201 ± 34 ± 79	² ASTON	88 LASS	0	$11K^- p \rightarrow K^- \pi^+ n$

- ² We take the central value of the two solutions and the larger error given.

 $K_0^*(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(52 ± 14) %

 $K_0^*(1950)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.52 ± 0.08 ± 0.12	³ ASTON	88 LASS	0	$11K^- p \rightarrow K^- \pi^+ n$

- ³ We take the central value of the two solutions and the larger error given.

 $K_0^*(1950)$ REFERENCES

ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
-------	----	-------------	----------------------	--------------------------

See key on page 1343

Meson Full Listings

 $K_2^*(1980)$, $K_4^*(2045)$ $K_2^*(1980)$

$$I(J^P) = \frac{1}{2}(2^+)$$

OMITTED FROM SUMMARY TABLE

Seen in the $J^P = 2^+$ wave amplitude of the $K^0 \pi^- \pi^+$ system.
Needs confirmation. $K_2^*(1980)$ MASS

All from partial wave amplitude analyses.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1975 ± 22 OUR AVERAGE					
1978 ± 40	241 ± 47	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$
1973 ± 8 ± 25		ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$

 $K_2^*(1980)$ WIDTH

All from partial wave amplitude analyses.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
373 ± 33 ± 60					
		ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
398 ± 47	241 ± 47	BIRD	89	LASS	- 11 $K^- p \rightarrow \bar{K}^0 \pi^- p$

 $K_2^*(1980)$ DECAY MODES

Mode

Γ_1	$K^*(892)\pi$
Γ_2	$K\rho$

 $K_2^*(1980)$ BRANCHING RATIOS

$\Gamma(K\rho)/\Gamma(K^*(892)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
1.49 ± 0.24 ± 0.09	ASTON	87	LASS	0 11 $K^- p \rightarrow \bar{K}^0 \pi^+ \pi^- n$	

 $K_2^*(1980)$ REFERENCES

BIRD	89	SLAC-332			(SLAC)
ASTON	87	NP B292 693	+Awaji, D'Amore+		(SLAC, NAGO, CINC, INUS)

OTHER RELATED PAPERS

AMSLER	93C	NP A558 3C	+Augustin+		(Crystal Barrel Collab.)
--------	-----	------------	------------	--	--------------------------

 $K_4^*(2045)$

$$I(J^P) = \frac{1}{2}(4^+)$$

 $K_4^*(2045)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2045 ± 9 OUR AVERAGE					Error includes scale factor of 1.1.
2062 ± 14 ± 13		¹ ASTON	86	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
2039 ± 10	400	^{2,3} CLELAND	82	SPEC	± 50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
2070 ⁺ 100 - 40		⁴ ASTON	81C	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$

• • • We do not use the following data for averages, fits, limits, etc. • • •

2079 ± 7	431	TORRES	86	MPSF	400 $pA \rightarrow 4KX$
2088 ± 20	650	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow K_S^0 \pi^- p$
2115 ± 46	488	CARMONY	77	HBC	0 9 $K^+ d \rightarrow K^+ \pi^+ s X$

¹ From a fit to all moments.² From a fit to 8 moments.³ Number of events evaluated by us.⁴ From energy-independent partial-wave analysis. $K_4^*(2045)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
198 ± 30 OUR AVERAGE					
221 ± 48 ± 27		⁵ ASTON	86	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
189 ± 35	400	^{6,7} CLELAND	82	SPEC	± 50 $K^+ p \rightarrow K_S^0 \pi^\pm p$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
61 ± 58	431	TORRES	86	MPSF	400 $pA \rightarrow 4KX$
170 ⁺ 100 - 50	650	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow K_S^0 \pi^- p$
240 ⁺ 500 - 100		⁸ ASTON	81C	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$
300 ± 200		CARMONY	77	HBC	0 9 $K^+ d \rightarrow K^+ \pi^+ s X$

⁵ From a fit to all moments.⁶ From a fit to 8 moments.⁷ Number of events evaluated by us.⁸ From energy-independent partial-wave analysis. $K_4^*(2045)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K\pi$	(9.9 ± 1.2) %
Γ_2 $K^*(892)\pi\pi$	(9 ± 5) %
Γ_3 $K^*(892)\pi\pi\pi$	(7 ± 5) %
Γ_4 $\rho K\pi$	(5.7 ± 3.2) %
Γ_5 $\omega K\pi$	(5.0 ± 3.0) %
Γ_6 $\phi K\pi$	(2.8 ± 1.4) %
Γ_7 $\phi K^*(892)$	(1.4 ± 0.7) %

 $K_4^*(2045)$ BRANCHING RATIOS

$\Gamma(K\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
0.099 ± 0.012	ASTON	88	LASS	0 11 $K^- p \rightarrow K^- \pi^+ n$	

$\Gamma(K^*(892)\pi\pi)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
0.89 ± 0.53	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$	

$\Gamma(K^*(892)\pi\pi\pi)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_1
0.75 ± 0.49	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$	

$\Gamma(\rho K\pi)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_1
0.58 ± 0.32	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$	

$\Gamma(\omega K\pi)/\Gamma(K\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_1
0.50 ± 0.30	BAUBILLIER	82	HBC	- 8.25 $K^- p \rightarrow \rho K_S^0 3\pi$	

$\Gamma(\phi K\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
0.028 ± 0.014	⁹ TORRES	86	MPSF 400 $pA \rightarrow 4KX$	

⁹ Error determination is model dependent.

$\Gamma(\phi K^*(892))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
0.014 ± 0.007	¹⁰ TORRES	86	MPSF 400 $pA \rightarrow 4KX$	

¹⁰ Error determination is model dependent. $K_4^*(2045)$ REFERENCES

ASTON	88	NP B296 493	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
ASTON	86	PL B180 308	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
TORRES	86	PR 34 707	+Lai+	(VPI, ARIZ, FNAL, FSU, NDAM, TUFTS+)
BAUBILLIER	82	PL 1183 447	+Burns+	(BIRM, CERN, GLAS, MSU, CURIN)
CLELAND	82	NP B208 189	+Delfosse, Dorsaz, Gloor	(DURH, GEVA, LAUS, PIT)
ASTON	81C	PL 1068 235	+Carnegie, Dunwood+	(SLAC, CARL, OTTA) JP
CARMONY	77	PR D16 1251	+Clopp, Lander, Meiere, Yen+	(PURD, UCD, IUUP)

OTHER RELATED PAPERS

ASTON	87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
BROMBERG	80	PR D22 1513	+Haggerty, Abrams, Dzierba	(CIT, FNAL, ILLC, IND)
CARMONY	71	PRL 27 1160	+Cords, Clopp, Erwin, Meiere+	(PURD, UCD, IUUP)

Meson Full Listings

 $K_2(2250)$, $K_3(2320)$, $K_5^*(2380)$, $K_4(2500)$ $K_2(2250)$

$$I(J^P) = \frac{1}{2}(2^-)$$

OMITTED FROM SUMMARY TABLE

This entry contains various peaks in strange meson systems reported in the 2150–2260 MeV region, as well as enhancements seen in the antihyperon-nucleon system, either in the mass spectra or in the $J^P = 2^-$ wave.

 $K_2(2250)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2247 ± 17	OUR AVERAGE				
2200.0 ± 40.0		¹ ARMSTRONG 83C OMEG	–		18 $K^- p \rightarrow \Lambda \bar{p} X$
2235 ± 50		¹ BAUBILLIER 81 HBC	–		8 $K^- p \rightarrow \Lambda \bar{p} X$
2260 ± 20		¹ CLELAND 81 SPEC	±		50 $K^+ p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2147 ± 4	37	CHLIAPNIK... 79 HBC	+		32 $K^+ p \rightarrow \bar{\Lambda} p X$
2240 ± 20	20	LISSAUER 70 HBC			9 $K^+ p$

¹ $J^P = 2^-$ from moments analysis.

 $K_2(2250)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
180 ± 30	OUR AVERAGE				Error includes scale factor of 1.4.
150.0 ± 30.0		² ARMSTRONG 83C OMEG	–		18 $K^- p \rightarrow \Lambda \bar{p} X$
210 ± 30		² CLELAND 81 SPEC	±		50 $K^+ p \rightarrow \Lambda \bar{p} X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
~ 200		² BAUBILLIER 81 HBC	–		8 $K^- p \rightarrow \Lambda \bar{p} X$
~ 40	37	CHLIAPNIK... 79 HBC	+		32 $K^+ p \rightarrow \bar{\Lambda} p X$
80 ± 20	20	LISSAUER 70 HBC			9 $K^+ p$

² $J^P = 2^-$ from moments analysis.

 $K_2(2250)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi \pi$	(6.1 ± 1.2) %
Γ_2 $p \bar{\Lambda}$	

 $K_2(2250)$ REFERENCES

ARMSTRONG 83C	NP B227 365	+	(BARI, BIRM, CERN, MILA, CURIN+)
BAUBILLIER 81	NP B183 1	–	(BIRM, CERN, GLAS, MSU, CURIN) JP
CLELAND 81	NP B184 1	+	Nef, Martin+ (PITT, GEVA, LAUS, DURH) JP
CHLIAPNIK... 79	NP B158 253	–	Chliapnikov, Gerdyukov+ (CERN, BELG, MONS)
LISSAUER 70	NP B18 491	–	+Alexander, Firestone, Goldhaber (LBL)

OTHER RELATED PAPERS

ALEXANDER 68B	PRL 20 755	+	Firestone, Goldhaber, Shen (LRL)
---------------	------------	---	----------------------------------

 $K_3(2320)$

$$I(J^P) = \frac{1}{2}(3^+)$$

OMITTED FROM SUMMARY TABLE

Seen in the $J^P = 3^+$ wave of the antihyperon-nucleon system. Needs confirmation.

 $K_3(2320)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2324 ± 24	OUR AVERAGE			
2330.0 ± 40.0	¹ ARMSTRONG 83C OMEG	–		18 $K^- p \rightarrow \Lambda \bar{p} X$
2320.0 ± 30.0	¹ CLELAND 81 SPEC	±		50 $K^+ p \rightarrow \Lambda \bar{p} X$

¹ $J^P = 3^+$ from moments analysis.

 $K_3(2320)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
150.0 ± 30.0	OUR AVERAGE			
~ 250.0	² ARMSTRONG 83C OMEG	–		18 $K^- p \rightarrow \Lambda \bar{p} X$
~ 250.0	² CLELAND 81 SPEC	±		50 $K^+ p \rightarrow \Lambda \bar{p} X$

² $J^P = 3^+$ from moments analysis.

 $K_3(2320)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $p \bar{\Lambda}$	

 $K_3(2320)$ REFERENCES

ARMSTRONG 83C	NP B227 365	+	(BARI, BIRM, CERN, MILA, CURIN+)
CLELAND 81	NP B184 1	–	+Nef, Martin+ (PITT, GEVA, LAUS, DURH)

 $K_5^*(2380)$

$$I(J^P) = \frac{1}{2}(5^-)$$

OMITTED FROM SUMMARY TABLE

Seen in partial wave analysis of the $K^- \pi^+$ system. Needs confirmation.

 $K_5^*(2380)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
2382 ± 14 ± 19	¹ ASTON	86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

¹ From a fit to all the moments.

 $K_5^*(2380)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT	
178 ± 37 ± 32	² ASTON	86	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$

² From a fit to all the moments.

 $K_5^*(2380)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $K \pi$	(6.1 ± 1.2) %

 $K_5^*(2380)$ BRANCHING RATIOS

$\Gamma(K \pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ	
0.061 ± 0.012	ASTON	88	LASS	0	11 $K^- p \rightarrow K^- \pi^+ n$	

 $K_5^*(2380)$ REFERENCES

ASTON 88	NP B296 493	+	Awaji, Bienz, Bird+ (SLAC, NAGO, CINC, INUS)
ASTON 86	PL B180 308	–	+Awaji, D'Amore+ (SLAC, NAGO, CINC, INUS)

 $K_4(2500)$

$$I(J^P) = \frac{1}{2}(4^-)$$

OMITTED FROM SUMMARY TABLE

Seen in the $J^P = 4^-$ wave of the antihyperon-nucleon system. Needs confirmation.

 $K_4(2500)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
2490.0 ± 20.0	¹ CLELAND 81 SPEC	±		50 $K^+ p \rightarrow \Lambda \bar{p}$

¹ $J^P = 4^-$ from moments analysis.

 $K_4(2500)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
~ 250.0	² CLELAND 81 SPEC	±		50 $K^+ p \rightarrow \Lambda \bar{p}$

² $J^P = 4^-$ from moments analysis.

 $K_4(2500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $p \bar{\Lambda}$	

 $K_4(2500)$ REFERENCES

CLELAND 81	NP B184 1	–	+Nef, Martin+ (PITT, GEVA, LAUS, DURH)
------------	-----------	---	--

See key on page 1343

Meson Full Listings
K(3100)**K(3100)**

$$I^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Narrow peak observed in several ($\Lambda\bar{p} + \text{pions}$) and ($\bar{\Lambda}p + \text{pions}$) states in Σ^- Be reactions by BOURQUIN 86 and in $n\bar{p}$ and nA reactions by ALEEV 93. Not seen by BOEHNLEIN 91. If due to strong decays, this state has exotic quantum numbers ($B=0, Q=+1, S=-1$ for $\Lambda\bar{p}\pi^+\pi^+$ and $I \geq 3/2$ for $\Lambda\bar{p}\pi^-$). See also under non- $q\bar{q}$ candidates. Needs confirmation. (See the index for the page number.)

K(3100) MASS

VALUE (MeV)	DOCUMENT ID
3100 ± 11 OUR ESTIMATE	

3-BODY DECAYS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3054 ± 11 OUR AVERAGE			
$3060 \pm 7 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+$
$3056 \pm 7 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-$
$3055 \pm 8 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^-$
$3045 \pm 8 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^+$

¹ Supersedes ALEEV 90.**4-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3059 ± 11 OUR AVERAGE			
$3067 \pm 6 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
$3060 \pm 8 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$
$3055 \pm 7 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$
$3052 \pm 8 \pm 20$	¹ ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$
3105 ± 30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
3115 ± 30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

5-BODY DECAYS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3095 ± 30	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$

K(3100) WIDTH**3-BODY DECAYS**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
42 ± 16	² ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+$
36 ± 15	² ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-$
50 ± 18	² ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^-$
30 ± 15	² ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^+$

² Supersedes ALEEV 90.**4-BODY DECAYS**

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
22 ± 8		² ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
28 ± 12		² ALEEV 93	BIS2	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$
32 ± 15		² ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^-$
30 ± 15		² ALEEV 93	BIS2	$K(3100) \rightarrow \bar{\Lambda}p\pi^-\pi^+$
< 30	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+$
< 80	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^-$

5-BODY DECAYS

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 30	90	BOURQUIN 86	SPEC	$K(3100) \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$

K(3100) DECAY MODES

Mode
$\Gamma_1 K(3100)^0 \rightarrow \Lambda\bar{p}\pi^+$
$\Gamma_2 K(3100)^{-} \rightarrow \Lambda\bar{p}\pi^-$
$\Gamma_3 K(3100)^{-} \rightarrow \bar{\Lambda}p\pi^-\pi^-$
$\Gamma_4 K(3100)^+ \rightarrow \Lambda\bar{p}\pi^+\pi^+$
$\Gamma_5 K(3100)^0 \rightarrow \Lambda\bar{p}\pi^+\pi^+\pi^-$
$\Gamma_6 K(3100)^0 \rightarrow \Sigma(1385)^+\bar{p}$

$$\Gamma(\Sigma(1385)^+\bar{p})/\Gamma(\Lambda\bar{p}\pi^+)$$

$$\Gamma_6/\Gamma_1$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.04	90	ALEEV 93	BIS2	$K(3100)^0 \rightarrow \Sigma(1385)^+\bar{p}$

K(3100) REFERENCES

ALEEV 93	PAN 56 1358	+Balandin+	(BIS-2 Collab.)
BOEHNLEIN 91	NP B21 174 (suppl)	+Chung+	(FLOR, BNL, IND, RICE, MASD)
ALEEV 90	ZPHY C47 533	+Arefiev, Balandin+	(BIS-2 Collab.)
BOURQUIN 86	PL B172 113	+Brown+	(GEVA, RAL, HEIDP, LAUS, BRIS, CERN)

Meson Full Listings

D Branching Fractions

CHARMED MESONS ($C = \pm 1$)

$D^+ = c\bar{d}$, $D^0 = c\bar{u}$, $\bar{D}^0 = \bar{c}u$, $D^- = \bar{c}d$, similarly for D^{*} 's

NOTE ON D MESON BRANCHING FRACTIONS

(by R.H. Schindler, SLAC)

This edition strongly reflects the impact that CLEO-II has had on charm meson physics in the last two years. The CESR-II luminosity for data taken at the $\Upsilon(4S)$ resonance and in the nearby e^+e^- continuum since startup in 1991 has climbed to $\sim 15\text{pb}^{-1}/\text{day}$ and has totaled more than 2fb^{-1} . In addition, this edition includes the first high-statistics results of the next (post-E691) generation of fixed-target experiments at FNAL (E653, E687). We anticipate that the CLEO-II and the FNAL fixed-target programs will continue to dominate the field through the end of the decade. The BES program for charmed physics is now also underway, and its first results should appear within the next year.

In this edition, we have restructured the Full Listings to further clarify and reduce the uncertainty in the normalization of the charmed D^+ , D^0 , and D_s^+ branching fractions. We continue to enter only experimentally measured quantities and not derived quantities, except where noted. All experiments measuring ratios of branching fractions are included in our calculations. (We shall call $B_1 = \Gamma_1/\Gamma_{\text{total}}$ a branching *fraction*, and Γ_1/Γ_2 a branching *ratio*.) These are usually measurements of branching fractions relative to the most accessible channels, such as $K^-\pi^+\pi^+$ for the D^+ , $K^-\pi^+$ and $K^-\pi^+\pi^+\pi^-$ for the D^0 , and $\phi\pi^+$ for the D_s^+ . The problem is to set the absolute scales for the branching fractions.

In our 1992 edition, we combined results from several experimental methods to set the scale of branching fractions of D^+ and D^0 mesons. One source was the direct measurements of the D^+ and D^0 branching fractions as reported in BALTRUSAITUS 86 and subsequently updated in ADLER 88C, using data at the $\psi(3770)$ resonance. In the latter paper, the rates of observed production of single charm and pairs of charm particles were compared to establish the absolute branching fraction scale. A second source was the results from topologically normalized experiments (*e.g.*, AGUILAR-BENITEZ 84) and other so-called “double tagging” techniques (*e.g.*, BARLAG 90D). A third source was measurements from ALEPH of the D^0 absolute branching fraction (DECAMP 91) using the technique pioneered by HRS (ABACHI 88) applied to D^* from Z decays. This technique compares the total rate for $Z \rightarrow D^{*+}X$ followed by $D^{*+} \rightarrow D^0\pi^+$, measured by observing only the soft low Q -value π^+ , with the total rate for $Z \rightarrow D^{*+}X$ followed by $D^{*+} \rightarrow D^0\pi^+$ and then $D^0 \rightarrow K^-\pi^+$.

In this edition, we have revisited the question, in the light of new measurements of the D^+ and D^0 branching fractions from CLEO-II. In AKERIB 93, CLEO-II measures the $D^0 \rightarrow K^-\pi^+$

branching fraction as $(3.91 \pm 0.08 \pm 0.17)\%$, in excellent agreement with the previous e^+e^- results at the $\psi(3770)$ and the Z but with better precision. In BALEST 94, CLEO-II uses a similar technique applied to the D^{*0} to measure the branching fraction $D^+ \rightarrow K^-\pi^+\pi^+$ to be $(9.3 \pm 0.6 \pm 0.8)\%$, with a precision comparable to the previous e^+e^- results at the $\psi(3770)$. In this edition, we use the direct measurements of branching fractions at the $\psi(3770)$ and the recent measurements using D^* decays, but remove the topological results from the averages and fits because of the uncertainty in the topological normalization and the inherent correlations with other measurements in the fitting procedure used to obtain them. The result is that the important $D^0 \rightarrow K^-\pi^+$ branching fraction is $(4.01 \pm 0.14)\%$ and the important $D^+ \rightarrow K^-\pi^+\pi^+$ branching fraction is $(9.1 \pm 0.6)\%$.

It may help to say a word about our averages and fits. An average is of (good) measurements of a single quantity, say of $\Gamma(D^0 \rightarrow K^-\pi^+)/\Gamma_{\text{total}}$. A fit involves two or more different quantities, say several different D^0 branching ratios. Our fits only include quantities that “push” against one another. For example, if there are measurements of $\Gamma_1/\Gamma_{\text{total}}$, $\Gamma_2/\Gamma_{\text{total}}$, and Γ_1/Γ_2 , then these three quantities are included in the fit. But if Γ_3 only occurs in Γ_3/Γ_1 , then $\Gamma_3/\Gamma_{\text{total}}$ is simply calculated from the fit value of $\Gamma_1/\Gamma_{\text{total}}$. Thus for the D^0 , say, we determine all the different measured branching ratios that push against one another and include them in one big fit. (The correlation-coefficient matrix for this fit is right in front of the measurements of the branching ratios.) Since the sum of branching fractions included in the fit is less than unity, a “dummy” mode is added to account for modes absent. Once the overall fit is done, remaining branching fractions (of the sort $\Gamma_3/\Gamma_{\text{total}}$ mentioned above) are calculated from branching fractions obtained in the fit.

The absolute branching fraction scale for the D_s^+ is still quite uncertain. We anticipate in the near future a measurement at BES of the absolute branching fraction for $D_s^+ \rightarrow \phi\pi^+$, using the direct technique exploited first by MARK-III (*e.g.*, ADLER 90B). At present, we are constrained to rely on a set of experiments (ARGUS, CLEO, CLEO-II, and E687) that determine the ratio $\Gamma(D_s^+ \rightarrow \phi e^+\nu_e)/\Gamma(D_s^+ \rightarrow \phi\pi^+)$. The ratio $\Gamma(D_s^+ \rightarrow \phi e^+\nu_e)/\Gamma(D^+ \rightarrow \bar{K}^{*0}e^+\nu_e)$ is reliably calculated theoretically, and the experiments measure $B(D^+ \rightarrow \bar{K}^{*0}e^+\nu_e)$ and use the well-measured relative lifetimes (or total widths) of the D_s^+ and D^+ to then obtain $B(D_s^+ \rightarrow \phi\pi^+)$. We thus get $B(D_s^+ \rightarrow \phi\pi^\pm) = 3.5 \pm 0.4\%$ for normalizing the other decays.

A result of much theoretical interest is the measurement of the pseudoscalar weak decay constants (f_D) of the D_s^+ and the D^+ . The decay constant measures the overlap of the heavy and light quark wave functions in the meson and is measured directly through the observation of pure leptonic decays of the D^+ or D_s^+ to $\mu^+\nu_\mu$ or $\tau^+\nu_\tau$. The precise value provides an unambiguous test of lattice QCD. The best limit on the D^+ decay constant was set by MARK-III ($< 290\text{ MeV}$ at 90% CL). Two experiments, WA75 (AOKI 93) and CLEO-II

See key on page 1343

Meson Full Listings

D Branching Fractions, Semileptonic Decays of *D*'s

(ACOSTA 94) have published evidence for pure leptonic decays of the D_s^+ . BES has also presented evidence in the form of a few events recoiling from tagged D_s^+ mesons in the 4-GeV region in e^+e^- . AOKI 93 claims 8 events in an emulsion experiment and finds a decay constant value of 232 MeV. CLEO-II finds about 39 events above background, corresponding to a decay constant of about 344 MeV. Both experiments have errors of about 75 MeV. The CLEO value is surprisingly large, but the error from the background subtraction and statistics still remains large. For further discussion, see also the “Note on Pseudoscalar-Meson Decay Constants” in the π^\pm Full Listings.

See also the “Note on Semileptonic Decays of *D* and *B* Mesons” that follows for a review of the many new results in that area.

NOTE ON SEMILEPTONIC DECAYS OF *D* AND *B* MESONS, PART I

(by R.J. Morrison and J.D. Richman, University of California, Santa Barbara)

I. Introduction

Weak decays of heavy mesons are classified as hadronic, leptonic, or semileptonic, according to the type of particles present in the final state. The most complicated are hadronic decays, in which strong interactions produce nonperturbative effects that are difficult to calculate. In leptonic decays, the final state contains only a charged lepton and a neutrino; if the decaying meson is a pseudoscalar, the effects of strong interactions in the initial state can be parametrized by a single constant. (See the “Note on Pseudoscalar-Meson Decay Constants” in the π^\pm Full Listings.) The rates for leptonic decay, however, are generally small and hard to measure.

Semileptonic decays are more tractable theoretically than hadronic decays, both because the $\ell\nu_\ell$ system produced by the virtual W boson is well understood, and because this system cannot have strong interactions with the hadronic decay products. In addition, semileptonic decay rates are reasonably large and experimentally accessible. As a consequence, semileptonic processes are the primary source of information on the Cabibbo-Kobayashi-Maskawa (CKM) matrix (see “The Cabibbo-Kobayashi-Maskawa Mixing Matrix,” in Section 28 of the Reviews, Tables, Figures, and Formulae part of this *Review*).

In this Note, we discuss measurements of semileptonic decays of *D* and *B* mesons. These studies have two primary goals: first, to understand the dynamics of semileptonic decays, in particular, the effect of strong interactions on the underlying weak process; and second, to measure the magnitudes of the CKM elements V_{cb} and V_{ub} . These goals are related, because the determination of CKM elements relies on a good understanding of the decay process. Our discussion focuses on the sources of uncertainty in different types of measurements, especially the model dependence in the inclusive *B* semileptonic branching ratio and in the determination of $|V_{cb}|$ and $|V_{ub}|$.

The effects of strong interactions on the semileptonic decay amplitude can be expressed in terms of one or more Lorentz-invariant functions called form factors, which depend only on q^2 , the square of the mass of the virtual W . The number of form factors required to describe a given exclusive decay depends on the quantum numbers of the hadrons in the decay. Information on the form factors is obtained by measuring the distributions of decay angles and q^2 , as discussed in the following section.

In the charm sector, the absolute scale of theoretical predictions for decay amplitudes (expressed in terms of form factors) can be tested, because the CKM elements can be determined independently of the *D* semileptonic decay rate using the assumption of CKM unitarity and the smallness of the CKM elements for *B* decay. Thus, the study of *D*-meson semileptonic decays is concerned more with the size and q^2 dependence of the form factors than with the determination of CKM elements. However, studies of form factors in *D* semileptonic decays should provide theoretical information that will help to improve predictions for $b \rightarrow u \ell^- \bar{\nu}_\ell$ decays, and hence improve the determination of $|V_{ub}|$.

A development of major importance is the heavy quark effective theory (HQET). In the limit of infinite quark mass, the quark mass and spin decouple from the dynamics of the decay, leading to numerous symmetry relations among form factors. In this limit, the description of a process such as $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ is simplified considerably, because both the *b* and *c* quarks are heavy. The form factors can then be related to a single universal form factor, the Isgur-Wise function. HQET predictions permit a systematic approach to understanding the dynamics of *B*-meson semileptonic decays and to determining the magnitudes of CKM elements. HQET predictions for *D* semileptonic decays are less useful, because symmetry-breaking corrections in this case are expected to be large.

We begin with a brief review of the formalism of form factors for exclusive semileptonic decays and then turn to experimental results on *D* and *B* mesons. More detailed information may be found in review articles on semileptonic decays and HQET [1,2,3].

II. Form-factor formalism for exclusive semileptonic decays

The amplitude for an exclusive semileptonic process can be constructed from the available four-vectors in the decay and from form factors, which are Lorentz invariant functions of q^2 , the square of the mass of the virtual W . Because these functions describe the effect of strong interactions, nonperturbative techniques such as lattice QCD are needed to calculate them. Form factors are generally largest at the maximum value of q^2 , where the daughter meson has the smallest recoil velocity and the overlap between the parent- and daughter-meson wave functions is largest. Studies of form factors in *D* semileptonic decays have focused on the modes $D \rightarrow \bar{K} \ell^+ \nu_\ell$ and $D \rightarrow \bar{K}^* \ell^+ \nu_\ell$, which dominate the inclusive semileptonic rate. In *B* decays, the analogous modes, $B \rightarrow \bar{D} \ell^+ \nu_\ell$ and $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$, account for about two-thirds of the inclusive semileptonic rate. The decay $B \rightarrow \bar{D} \ell^+ \nu_\ell$ has a large background from

Meson Full Listings

Semileptonic Decays of D 's

$B \rightarrow \bar{D}^* \ell^+ \nu_\ell$, so in B decays, form factors have been studied only for $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$. In this section, we discuss the formalism used in form-factor measurements for the decays $P \rightarrow P' \ell \nu_\ell$, where P and P' are pseudoscalar mesons, and $P \rightarrow V \ell \nu_\ell$, where V is a vector meson.

The differential decay rate for $P(Q\bar{q}) \rightarrow P'(q'\bar{q}) \ell \nu_\ell$ is

$$\frac{d\Gamma}{dq^2} = \frac{G_F^2 |V_{q'Q}|^2 k_{P'}^3 |f_+(q^2)|^2}{24\pi^3}. \quad (1)$$

Here G_F is the Fermi decay constant, $V_{q'Q}$ is the relevant CKM matrix element, $k_{P'}$ is the momentum of P' in the rest frame of the parent meson, and $f_+(q^2)$ is a vector form factor. (Eq. (1) assumes massless charged leptons, which is almost exact for electrons and a very good approximation for muons, but it is not correct for τ 's.) The dominant q^2 dependence comes from the p -wave factor $k_{P'}^3$, which can be written in terms of q^2 and the particle masses. This factor increases the rate at low q^2 , which is opposite to the q^2 dependence of the form factor f_+ .

The exclusive decay rate for $P \rightarrow V \ell \nu_\ell$ can be expressed in terms of three q^2 -dependent helicity amplitudes, $H_\pm(q^2)$ and $H_0(q^2)$, where the subscripts indicate the helicity of either the virtual W or the vector meson. The rate is given by

$$\begin{aligned} \frac{d\Gamma}{dq^2 d\cos\theta_\ell d\cos\theta_V d\chi} &= \frac{3G_F^2 |V_{q'Q}|^2 k_V q^2}{8(4\pi)^4 M^2} \left\{ [(1 + \eta \cos\theta_\ell)^2 |H_+(q^2)|^2 \right. \\ &+ (1 - \eta \cos\theta_\ell)^2 |H_-(q^2)|^2] \sin^2\theta_V + 4 \sin^2\theta_\ell \cos^2\theta_V |H_0(q^2)|^2 - 2 \sin^2\theta_\ell \sin^2\theta_V \cos(2\chi) H_+(q^2) H_-(q^2) \\ &\left. - 4\eta \sin\theta_\ell (1 + \eta \cos\theta_\ell) \sin\theta_V \cos\theta_V \cos\chi H_+(q^2) H_0(q^2) + 4\eta \sin\theta_\ell (1 - \eta \cos\theta_\ell) \sin\theta_V \cos\theta_V \cos\chi H_-(q^2) H_0(q^2) \right\}. \quad (2) \end{aligned}$$

Here M is the mass of the parent meson, k_V is the momentum of the vector meson and is a function of q^2 , and the factor $\eta=+1$ ($\eta=-1$) applies to B (D) decays. The angles θ_ℓ , θ_V , and χ are defined in Fig. 1. The helicity amplitudes H_\pm and H_0 can be expressed in terms of two axial-vector form factors, $A_1(q^2)$ and $A_2(q^2)$, and a vector form factor $V(q^2)$:

$$\begin{aligned} H_\pm(q^2) &= (M+m)A_1(q^2) \mp \frac{2M k_V}{(M+m)} V(q^2) \\ H_0(q^2) &= \frac{1}{2m\sqrt{q^2}} \left[(M^2 - m^2 - q^2)(M+m)A_1(q^2) - \frac{4M^2 k_V^2}{(M+m)} A_2(q^2) \right], \quad (3) \end{aligned}$$

where m is the mass of the daughter meson. The form factors f_+ , A_1 , A_2 , and V are dimensionless.




Figure 1: Definition of the angles θ_ℓ , θ_V , and χ . The decay $D \rightarrow \bar{K}^* \ell^+ \nu_\ell$ is used as an example. The polar angles θ_ℓ and θ_V are defined in the rest frames of the virtual W and the \bar{K}^* , respectively, and χ is the azimuthal angle between the projections of the lepton and the kaon momentum vectors in the plane perpendicular to z .

The $V-A$ coupling results in a larger amplitude to produce a negative-helicity vector meson in $c\bar{q}$ or $b\bar{q}$ decay than one of positive helicity: $|H_-| > |H_+|$. This difference produces a forward-backward asymmetry for the charged lepton in the virtual- W rest frame, since the net angular momentum along the decay axis of the initial heavy meson must be zero. For $D(c\bar{q})$ decays, a positively charged (right-handed) lepton is produced in association with a left-handed daughter s or d quark, resulting in a softer energy spectrum for the charged lepton than for the neutrino after boosting the lepton energy into the D rest frame. (A similar argument shows that the shape of the spectrum is the same for a \bar{D} decay.) For $\bar{B}(b\bar{q})$ decays, a negatively charged (left-handed) lepton is produced in association with a left-handed daughter quark, giving a harder energy spectrum for the charged lepton than for the neutrino in the B rest frame. In $P \rightarrow P' \ell \nu_\ell$ decays, there is no asymmetry, since the P' meson can only have helicity zero. Thus, the effect of $V-A$ is to soften the inclusive lepton spectrum in D decays and to harden it in B decays. It is useful to define rates for decays into specific helicity states:

$$\Gamma_i = \frac{G_F^2 |V_{q'Q}|^2}{96\pi^3} \int dq^2 k_V \frac{q^2}{M^2} |H_i(q^2)|^2. \quad (4)$$

Experiments extract various ratios of these rates, including Γ_+/Γ_- , $\Gamma_L/\Gamma_T = \Gamma_0/(\Gamma_- + \Gamma_+)$, the lepton forward-backward asymmetry $A_{FB} = (3\eta/4)(\Gamma_- - \Gamma_+)/\Gamma$, and the polarization parameter $\alpha = 2\Gamma_0/(\Gamma_+ + \Gamma_-) - 1$. The three form factors can be obtained by fitting the measured distribution of the variables q^2 , θ_ℓ , θ_V , and χ .

III. D -meson semileptonic decays

In this Section, we review the main results on D and D_s semileptonic decays, including measurements of $D \rightarrow \bar{K}\ell^+\nu_\ell$ and $D \rightarrow \bar{K}^*\ell^+\nu_\ell$ form factors. An important question is whether the sum of the exclusive rates for these two modes, plus the corresponding Cabibbo-suppressed rates, saturate the inclusive rate. This issue can also be addressed by searching for Cabibbo-favored semileptonic decays to hadronic systems other than the K or K^* , referred to here as nonresonant states. It is also of great interest to measure the Cabibbo-suppressed semileptonic decays of the D^0 and D^+ and the semileptonic decays of the D_s^+ . These decays differ from the dominant K or K^* decays by the strange quark content of either the initial meson or the final meson, or both. Although there are differences in the q^2 ranges and the pole masses for Cabibbo-favored and Cabibbo-suppressed decays, the form-factor intercepts for these modes should be similar. We review the measurements that attempt to address these questions. In our tables, averages of experimental results labeled “*PDG value*” are not necessarily an exact average of the numbers in the tables but come from making an overall fit to the branching ratios in the Full Listings that follow this Note.

$D \rightarrow \bar{K}\ell^+\nu_\ell$ decays

With the new CLEO-II measurement [4], the ratio $\Gamma(D^0 \rightarrow K^-e^+\nu_e)/\Gamma(D^0 \rightarrow K^-\pi^+)$ is one of the most accurately determined quantities in charm physics. This measurement, together with the new, more precise value of the absolute $D^0 \rightarrow K^-\pi^+$ branching fraction, yields a much improved measurement of $B(D^0 \rightarrow K^-e^+\nu_e)$, as shown in Table 1. Note that e - μ universality is assumed, and that muon measurements have been scaled up by a factor of 1.03 to account for the reduced phase space. Therefore, all results are expressed as electron branching fractions. The transition rate, $\Gamma(D^0 \rightarrow K^-e^+\nu_e) = (8.9 \pm 0.5) \times 10^{10} \text{ s}^{-1}$, shown in Table 10, is obtained using the PDG branching fraction of Table 1 and the PDG value of the D^0 lifetime.

Table 1: Measurements of the $D^0 \rightarrow K^-e^+\nu_e$ branching fraction. The PDG branching fraction for $D^0 \rightarrow K^-\pi^+$ has been used, where needed, to normalize.

Experiment	Reference [†]	Lepton	Norm. mode	$B(D^0 \rightarrow K^-e^+\nu_e)\%$	m_p (GeV)
E691	[5] ANJOS 89F	e	$D^0 \rightarrow K^-\pi^+$	3.65 ± 0.54	$2.1_{-0.2}^{+0.4} \pm 0.2$
CLEO-I	[6] CRAWFORD 91B	e	$D^0 \rightarrow K^-\pi^+$	3.61 ± 0.37	$2.1_{-0.2}^{+0.4+0.3}$
CLEO-I*	[6] CRAWFORD 91B	μ	$D^0 \rightarrow K^-\pi^+$	3.26 ± 0.50	
E687*	[7] FRABETTI 93I	μ	$D^0 \rightarrow K^-\pi^+$	3.39 ± 0.75	
CLEO-II*	[4] BEAN 93C	e, μ	$D^0 \rightarrow K^-\pi^+$	3.92 ± 0.25	$2.00 \pm 0.12 \pm 0.18$
MARK-III	[8] ADLER 89	e	absolute	3.4 ± 0.6	
E653*	[9] KODAMA 93B	μ	μ inclusive	2.5 ± 0.6	
PDG value				3.68 ± 0.21	

* Muon measurements have been scaled up by 1.03 to be equivalent to electrons.
[†] Here and in several later Tables, the reference ID used in the Data Listings is given following the reference number.

Isospin symmetry requires equal transition rates for the Cabibbo-favored D^0 and D^+ decays. The D^+ branching fraction measurements are given in Table 2, using our branching fractions for the normalizing modes. The average $D^+ \rightarrow \bar{K}^0e^+\nu_e$ transition rate is $(6.3 \pm 0.8) \times 10^{10} \text{ s}^{-1}$, significantly lower than $\Gamma(D^0 \rightarrow K^-e^+\nu_e)$, although it is less well measured. The discrepancy may be due, in part, to poorly measured normalizing branching fractions. In any case, we use the average transition rate $\Gamma(D \rightarrow \bar{K}e^+\nu_e) = (8.2 \pm 0.4) \times 10^{10} \text{ s}^{-1}$ to compute the form factor discussed below.

Table 2: Measurements of the $D^+ \rightarrow \bar{K}^0e^+\nu_e$ branching fraction.

Experiment	Reference	Norm. mode	$B(D^+ \rightarrow \bar{K}^0e^+\nu_e)\%$
MARK-III*	[10] BAI 91	absolute	$6.8_{-1.1}^{+1.6}$
E691	[11] ANJOS 91C	$D^+ \rightarrow K^-\pi^+\pi^+$	6.0 ± 1.6
CLEO-II*	[4] BEAN 93C	$D^+ \rightarrow \bar{K}^0\pi^+$	7.1 ± 0.8
PDG value			6.7 ± 0.8

* Average of electrons and muons.

Meson Full Listings

Semileptonic Decays of D 's

The decay $D^0 \rightarrow K^- e^+ \nu_e$ is the only one so far that allows a useful study of the q^2 dependence of heavy-meson form factors. Experiments to date are precise enough to determine only an intercept and a slope parameter for the function $f_+^K(q^2)$. This form factor is commonly parametrized with a pole form,

$$f_+^K(q^2) = f_+^K(0)/(1 - q^2/m_p^2), \quad (5)$$

where m_p is an appropriate pole mass, expected to be approximately the mass of the D_s^* . Values of m_p derived from the experiments are given in Table 1 and are in agreement with $m_{D_s^*}$. The form-factor intercept is computed by integrating Eq. (1), assuming the pole form for $f_+^K(q^2)$. Using the average transition rate of the two charge states, the result is $|f_+^K(0)||V_{cs}| = 0.73 \pm 0.02 \pm 0.02$, where the first error is from the uncertainty in the decay rate and the second is from the uncertainty in the shape of $f_+^K(q^2)$. Using $|V_{cs}| = 0.974$ from the unitarity of the CKM matrix, the value $f_+^K(0) = 0.75 \pm 0.02 \pm 0.02$ is obtained. This result is in general agreement with predictions from models and lattice gauge calculations, as shown in Table 8.

$D \rightarrow \bar{K}^* \ell^+ \nu_\ell$ decays

The differential decay distribution for $D \rightarrow \bar{K}^* \ell^+ \nu_\ell$ is governed, as explained in Section II, by one vector and two axial-vector form factors. Experiments do not yet have enough data to measure the q^2 dependences of these form factors. The practice is to assume that they have a simple pole form, as for the pseudoscalar case, using for pole masses those of the lowest lying $c\bar{s}$ states with the appropriate quantum numbers: the D_s^* mass, 2.1 GeV, for the pole mass m_V , and the D_s^{**} mass, 2.5 GeV, for m_A . The ratios of form-factor intercepts, $R_2 = A_2(0)/A_1(0)$ and $R_V = V(0)/A_1(0)$, are then determined by fitting to the q^2 and angular decay distributions. These ratios have been measured in $D^+ \rightarrow \bar{K}^{*0} \ell^+ \nu_\ell$ decay and are given in Table 3. The corresponding ratios of longitudinal to transverse polarization, and of positive to negative transverse polarization, are also shown.

Table 3: $D^+ \rightarrow \bar{K}^{*0} \ell^+ \nu_\ell$ form factor ratios.

Experiment	Ref.	$R_2(0)$	$R_V(0)$	Γ_L/Γ_T	Γ_+/Γ_-
E691	[12]	$0.0 \pm 0.5 \pm 0.2$	$2.0 \pm 0.6 \pm 0.3$	$1.8^{+0.6}_{-0.4} \pm 0.3$	$0.15^{+0.07}_{-0.05} \pm 0.03$
E653	[13]	$0.82^{+0.22}_{-0.23} \pm 0.11$	$2.00^{+0.34}_{-0.32} \pm 0.16$	$1.18 \pm 0.18 \pm 0.08$	$0.16 \pm 0.05 \pm 0.02$
E687	[16]	$0.78 \pm 0.18 \pm 0.10$	$1.74 \pm 0.27 \pm 0.28$	$1.20 \pm 0.13 \pm 0.13$	
Ave.		0.73 ± 0.15	1.89 ± 0.25	1.23 ± 0.13	0.16 ± 0.04

The normalization of the form factors is given by the transition rate, which is proportional to $A_1(0)^2$ and is only weakly dependent on R_V and R_2 . Tables 4 and 5 give measurements of the $D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e$ and $D^0 \rightarrow K^{*-} e^+ \nu_e$ branching fractions with the associated average transition rates in Table 10. The D^+ measurement is the better of the two, due to the long D^+ lifetime, the high efficiency for observing the \bar{K}^{*0} , and the recent, more precise measurement of the normalizing $D^+ \rightarrow K^- \pi^+ \pi^+$ branching fraction.

Table 4: Measurements of the $D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e$ branching fraction.

Experiment	Reference	Lepton	Norm. mode	$B(D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e)\%$
E691	[12] ANJOS 90E	e	$D^+ \rightarrow K^- \pi^+ \pi^+$	4.4 ± 0.6
E653*	[18] KODAMA 92C	μ	$D^+ \rightarrow K^- \pi^+ \pi^+$	4.4 ± 1.0
			$D^0 \rightarrow K^- e^+ \nu_e$	3.9 ± 1.1
E687*	[16] FRABETTI 93E	μ	$D^+ \rightarrow K^- \pi^+ \pi^+$	5.4 ± 0.8
WA82	[14] ADAMOVICH 91	e	$D^+ \rightarrow K^- \pi^+ \pi^+$	5.6 ± 1.6
ARGUS	[19] ALBRECHT 91	e	$D^+ \rightarrow K^- \pi^+ \pi^+$	5.0 ± 1.2
CLEO-II*	[4] BEAN 93C	e, μ	$D^+ \rightarrow K^- \pi^+ \pi^+$	6.1 ± 1.1
PDG value				4.8 ± 0.4

*Muon measurements have been scaled up by 1.05 to be equivalent to electrons.

Table 5: Measurements of the $D^0 \rightarrow K^{*-} e^+ \nu_e$ branching fraction.

Experiment	Reference	Norm. mode	$B(D^0 \rightarrow K^{*-} e^+ \nu_e)\%$
CLEO-I	[6] CRAWFORD 91B	$D^0 \rightarrow K^- e^+ \nu_e$	1.9 ± 0.7
CLEO-II	[4] BEAN 93C	$D^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$	2.0 ± 0.4
PDG value			2.0 ± 0.4

The new measurements confirm the old result that the transition rates for K^* decays are significantly smaller than for K decays, contrary to expectations from models. This result is also confirmed by several direct measurements of the K^* -to- K ratio, obtained without the use of normalizing branching fractions, shown in Table 6. This ratio is about 0.6, which leads to the low values for the form-factor intercepts $A_1(0)$ and $A_2(0)$ given in Table 7. These values are compared with predictions in Table 8. Note that the form-factor intercepts in these tables have been corrected for the new average value $\Gamma(D \rightarrow \bar{K}^* e^+ \nu_e) = (4.6 \pm 0.4) \times 10^{10} \text{ s}^{-1}$.

Table 6: Direct measurements of $\Gamma(D \rightarrow \bar{K}^* \ell \nu_\ell) / \Gamma(D \rightarrow \bar{K} \ell \nu_\ell)$.

Experiment	Reference	Measured ratio	Value
CLEO-I	[6] CRAWFORD 91B	$\frac{\Gamma(D^0 \rightarrow K^{*-} e^+ \nu_e)}{\Gamma(D^0 \rightarrow K^- e^+ \nu_e)}$	$0.51 \pm 0.18 \pm 0.06$
CLEO-II	[4] BEAN 93C	$\frac{\Gamma(D^0 \rightarrow K^{*-} e^+ \nu_e)}{\Gamma(D^0 \rightarrow K^- e^+ \nu_e)}$	$0.60 \pm 0.09 \pm 0.07$
CLEO-II	[4] BEAN 93C	$\frac{\Gamma(D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e)}{\Gamma(D^+ \rightarrow \bar{K}^0 e^+ \nu_e)}$	$0.65 \pm 0.09 \pm 0.10$
E653	[18] KODAMA 92C	$\frac{\Gamma(D^+ \rightarrow \bar{K}^{*0} \mu^+ \nu_\mu)}{\Gamma(D^+ \rightarrow \bar{K}^0 \mu^+ \nu_\mu)}$	$0.43 \pm 0.09 \pm 0.09$

Table 7: Form factors for $D^+ \rightarrow \bar{K}^{*0} e^+ \nu_e$. All results are normalized to $\Gamma(D \rightarrow \bar{K}^* e^+ \nu_e) = (4.6 \pm 0.4) \times 10^{10} \text{ s}^{-1}$.

Experiment	Ref.	$A_1(0)$	$V(0)$	$A_2(0)$
E653	[18]	0.57 ± 0.08	1.2 ± 0.3	0.47 ± 0.16
E691	[12]	0.49 ± 0.07	1.0 ± 0.3	0.0 ± 0.2
E687	[15]	0.59 ± 0.05	1.0 ± 0.3	0.46 ± 0.11
Ave.		0.56 ± 0.04	1.1 ± 0.2	0.40 ± 0.08

Table 8: Form factors: comparison with theory.

Model	Exp.	$f_+(0)$	$A_1(0)$	$V(0)$	$A_2(0)$
	average	$= 0.75 \pm 0.03$	0.56 ± 0.04	1.1 ± 0.2	0.40 ± 0.08
Quark models	ISGW [30]	0.8	0.8	1.1	0.8
	WSB [31]	0.76	0.88	1.3	1.2
	KS [32]	0.7	0.82	0.8	0.8
	AW/GS [33]	0.7	0.8	1.5	0.6
Lattice gauge	BKS [34]	$0.9 \pm 0.1 \pm 0.2$	$0.8 \pm 0.1 \pm 0.3$	$1.4 \pm 0.5 \pm 0.5$	$0.6 \pm 0.1 \pm 0.2$
	LMMS [35]	0.63 ± 0.08	0.53 ± 0.03	0.9 ± 0.1	0.2 ± 0.2
Sum rules	BBD [36]	0.60	0.5	1.1	0.6

Cabibbo-suppressed semileptonic decays

The rates of Cabibbo-suppressed pseudoscalar semileptonic decays determine the product of $|V_{cd}|$ times a form factor. Assuming that $|V_{cd}|$ is known precisely from unitarity of the CKM matrix, the main goal is to determine the form factor. The form factors measured in Cabibbo-suppressed charm decays are related by HQET [38] to those for $b \rightarrow u$ semileptonic decays in the q^2 region near q_{max}^2 , approximately 18–26 GeV^2 . Assuming appropriate pole forms for the form factors, the ratios of branching fractions can be written as

$$R_\pi = \frac{\text{B}(D^0 \rightarrow \pi^- e^+ \nu_e)}{\text{B}(D^0 \rightarrow K^- e^+ \nu_e)} = 2 \times \frac{\text{B}(D^+ \rightarrow \pi^0 e^+ \nu_e)}{\text{B}(D^+ \rightarrow \bar{K}^0 e^+ \nu_e)} = 1.97 \times \frac{|f_+^\pi(0)|^2 |V_{cd}|^2}{|f_+^K(0)|^2 |V_{cs}|^2}. \quad (6)$$

The factor of 2 difference between these two modes arises from the $1/\sqrt{2}$ in the coupling of $d\bar{d}$ to a π^0 .

To date, measurements of Cabibbo-suppressed semileptonic decays suffer from poor statistics; better results should be forthcoming from the new high-statistics charm experiments. The MARK-III measurement [20] of $\text{B}(D^0 \rightarrow \pi^- e^+ \nu_e)$, together with our value

Meson Full Listings

Semileptonic Decays of D 's

for $B(D^0 \rightarrow K^- e^+ \nu_e)$ from Table 1, yields $R_\pi = 0.11_{-0.03}^{+0.06} \pm 0.1$. This can be compared with the CLEO-II measurement of $B(D^+ \rightarrow \pi^0 e^+ \nu_e)$ [21], giving $R_\pi = 0.17 \pm 0.05 \pm 0.03$. From unitarity of the CKM matrix, we have $|V_{cd}/V_{cs}|^2 = 0.051 \pm 0.002$, giving values of $|f_+^\pi(0)/f_+^K(0)| = 1.0_{-0.2}^{+0.3} \pm 0.04$ and $1.3 \pm 0.2 \pm 0.1$ for the MARK-III and CLEO-II results. Model predictions, which range from 0.7 to 1.4 [39], are in agreement with these results. The E653 collaboration [22] reports a measurement, $\Gamma(D^+ \rightarrow \rho^0 \mu^+ \nu_\mu)/\Gamma(D^+ \rightarrow \bar{K}^{*0} \mu^+ \nu_\mu) = 0.044_{-0.025}^{+0.031} \pm 0.014$, in good agreement with expectations, within the large errors.

Nonresonant semileptonic decays

There is now substantial evidence that Cabibbo-favored semileptonic decays are nearly saturated by $D \rightarrow \bar{K} e^+ \nu_e$ and $D \rightarrow \bar{K}^* e^+ \nu_e$. Table 9 gives the results of searches for other decays. Early results from the hybrid bubble chamber [37] indicating large nonresonant contributions are clearly inconsistent with much more sensitive later measurements. Limits on the sum of the nonresonant (and higher-mass resonant) decays can be inferred from the comparison of the sum of exclusive rates with the inclusive rate, as discussed below.

Table 9: Limits on specific nonresonant modes.

Experiment	Reference	Mode	$\Gamma/10^{10} \text{ s}^{-1}$ 90% CL upper limit
E687	[16] FRABETTI 93E	$D^+ \rightarrow (K^- \pi^+)_{\text{NR}} \mu^+ \nu_\mu$	< 0.4
		$D^+ \rightarrow K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 0.1
E653	[17] KODAMA 93B	$D^0 \rightarrow K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 0.3
		$D^0 \rightarrow (K^* \pi)^- \mu^+ \nu_\mu$	< 0.4

Inclusive semileptonic decays

The inclusive transition rates are $\Gamma(D^+ \rightarrow X e^+ \nu_e) = (16.3 \pm 1.8) \times 10^{10} \text{ s}^{-1}$ and $\Gamma(D^0 \rightarrow X e^+ \nu_e) = (18.6 \pm 2.9) \times 10^{10} \text{ s}^{-1}$. These rates are dominated by old MARK-III measurements [23]. Isospin symmetry requires that the Cabibbo-favored semileptonic rates for the two charged states be equal, but small differences can be expected for the Cabibbo-suppressed rates. We ignore these differences, which are expected to be less than 2% of the inclusive rate, and use the average for the two charge states, $\Gamma(D \rightarrow X e^+ \nu_e) = (16.9 \pm 1.5) \times 10^{10} \text{ s}^{-1}$. This inclusive rate can be compared with the sum of the rates for $D \rightarrow \bar{K} e^+ \nu_e$, $D \rightarrow \bar{K}^* e^+ \nu_e$, and an estimated 8% contribution for the Cabibbo-suppressed modes, as shown in Table 10. (We average the D^0 and D^+ exclusive rates here as well.) The sum of these rates implies a deficit of slightly less than two standard deviations. CLEO [4] has measured all four of the exclusive K and K^* modes, normalizing to the best measured decay, $D^0 \rightarrow K^- \pi^+$. Averaging over the charge states, CLEO obtains $\Gamma(D \rightarrow (\bar{K} + \bar{K}^*) e^+ \nu_e) = (14.8 \pm 1.3) \times 10^{10} \text{ s}^{-1}$. Adding 8% for Cabibbo-suppressed decays results in a deficit of $(0.9 \pm 2.0) \times 10^{10} \text{ s}^{-1}$, consistent with zero. The rather large errors leave some room for small contributions from other exclusive channels. As seen in Table 9, the limits on individual channels are much more stringent.

Table 10: Exclusive transition rates. For comparison, $\Gamma(D \rightarrow X e^+ \nu_e) = (16.9 \pm 1.5) \times 10^{10} \text{ s}^{-1}$.

Mode	$\Gamma(D^0)/10^{10} \text{ s}^{-1}$	$\Gamma(D^+)/10^{10} \text{ s}^{-1}$	$\Gamma_{\text{Ave}}/10^{10} \text{ s}^{-1}$
$D \rightarrow \bar{K} e \nu_e$	8.9 ± 0.5	6.3 ± 0.8	8.2 ± 0.4
$D \rightarrow \bar{K}^* e \nu_e$	4.8 ± 1.0	4.5 ± 0.4	4.6 ± 0.4
Estimated Cabibbo-suppressed			1.0 ± 0.3
Sum of exclusive rates			13.8 ± 0.6

D_s^+ semileptonic decays

Because ϕ mesons are easy to observe, the first D_s^+ semileptonic decay measured was the decay to a vector, $D_s^+ \rightarrow \phi \ell^+ \nu_\ell$. The ratio of decay rates, $R_\phi \equiv \Gamma(D_s^+ \rightarrow \phi \ell^+ \nu_\ell)/\Gamma(D_s^+ \rightarrow \phi \pi^+)$, has been measured by several experiments (Table 11). The average value given in the table includes both electron and muon results.

Table 11: Measurements of $R_\phi \equiv \Gamma(D_s^+ \rightarrow \phi \ell^+ \nu_\ell)/\Gamma(D_s^+ \rightarrow \phi \pi^+)$.

Experiment	Reference	R_ϕ
CLEO-I	[28] ALEXANDER 90B	$0.49 \pm 0.10_{-0.14}^{+0.10}$
ARGUS	[27] ALBRECHT 91	$0.57 \pm 0.15 \pm 0.15$
E687	[26] FRABETTI 93G	$0.58 \pm 0.17 \pm 0.07$
CLEO-II	[25] BUTLER 94	$0.54 \pm 0.05 \pm 0.04$
PDG value		0.54 ± 0.05

See key on page 1343

Meson Full Listings Semileptonic Decays of D^+ 's

As yet there are no direct measurements of D_s^+ branching fractions. One method for estimating the $D_s^+ \rightarrow \phi\pi^+$ branching fraction is based on the belief that $D_s^+ \rightarrow \phi\ell^+\nu_\ell$ and $D \rightarrow \bar{K}^*\ell^+\nu_\ell$ decays are quite similar. Quark-model-based estimates [40] for the ratio $\kappa = \Gamma(D_s^+ \rightarrow \phi\ell^+\nu_\ell)/\Gamma(D \rightarrow \bar{K}^*\ell^+\nu_\ell)$ are about 1.0. The branching fraction is then

$$B(D_s^+ \rightarrow \phi\pi^+) = \frac{\tau_{D_s} \kappa \Gamma(D \rightarrow \bar{K}^*\ell^+\nu_\ell)}{R_\phi}. \quad (7)$$

The average over the two $\Gamma(D \rightarrow \bar{K}^*e^+\nu_e)$ charge states is $(4.6 \pm 0.4) \times 10^{10} \text{ s}^{-1}$. Using the PDG value for the D_s^+ lifetime, the value for R_ϕ in Table 11, and $\kappa = 1.0$ gives $B(D_s^+ \rightarrow \phi\pi^+) = (4.0 \pm 0.5)\%$. This value also has a theoretical uncertainty, which we conservatively estimate to be about 25%. (The PDG value of this branching fraction, which combines several estimates, is $(3.5 \pm 0.4)\%$.)

We will have more confidence in this method for determining the absolute D_s^+ branching fraction scale, and indeed more confidence in our understanding of charmed semileptonic decays in general, if the patterns for the D_s^+ semileptonic decays are seen to be the same as those for D^0 and D^+ . It is important to measure the form factor ratios for $D_s^+ \rightarrow \phi\ell^+\nu_\ell$ decay, as has been done for $D \rightarrow \bar{K}^*\ell^+\nu_\ell$, to shed some light on the validity of the assumption of equal transition rates. With small numbers of events, Fermilab experiments E653 and E687 have made first measurements of these ratios (Table 12). These results, to be compared with the $D \rightarrow \bar{K}^*e\nu_e$ values of Table 3, are not yet precise enough to draw conclusions about the degree of similarity of the D_s^+ and D decays.

Perhaps even more important is the ratio of the rates of semileptonic decays of the D_s^+ to vector and pseudoscalar mesons, since this ratio is not as expected in D^+ and D^0 decays. The first observation of the pseudoscalar modes is from Fermilab E653 [24], who obtained $[\Gamma(D_s^+ \rightarrow \phi\mu^+\nu_\mu)]/[\Gamma(D_s^+ \rightarrow (\eta + \eta')\mu^+\nu_\mu)] = 0.26 \pm 0.11$. The result is certainly consistent with a low vector-to-pseudoscalar ratio. This experiment was unable to separate the η and η' modes.

Table 12: $D_s^+ \rightarrow \phi\ell^+\nu_\ell$ form factor ratios.

Experiment	Ref.	$R_2(0)$	$R_V(0)$	Γ_L/Γ_T
E653	[24]	$2.1^{+0.6}_{-0.5} \pm 0.2$	$2.3^{+1.1}_{-0.9} \pm 0.4$	$0.54 \pm 0.21 \pm 0.10$
E687	[29]	$1.1 \pm 0.8 \pm 0.1$	$1.8 \pm 0.9 \pm 0.2$	$1.0 \pm 0.5 \pm 0.1$
Ave.		1.8 ± 0.5	2.0 ± 0.7	0.6 ± 0.2

Conclusions on D -meson semileptonic decays

The study of semileptonic decays of D mesons has progressed enormously in the last few years. The q^2 dependence of the pseudoscalar form factor has been measured, and the intercepts of the three vector-decay form factors been determined by several experiments. New experiments confirm the old observation that the $D \rightarrow \bar{K}^*\ell^+\nu_\ell$ decay rate, and therefore the A_1 form factor, is low compared with model calculations. Presently, no model appears able to explain all of the form factors.

Problems remain in the precise quantitative comparisons between the transition rates of the different charge states in the $D \rightarrow \bar{K}^*\ell^+\nu_\ell$ exclusive decays. The difficulty may lie in the branching fractions of the normalizing modes for the D^+ decays. Strict upper limits are observed on rates of nonresonant decays. These limits are consistent with the observation that the transition rates for $D \rightarrow \bar{K}^*\ell^+\nu_\ell$ and $D \rightarrow \bar{K}^*\ell^+\nu_\ell$, plus a reasonable estimate of the Cabibbo-suppressed rate, nearly saturates the inclusive rate.

Measurements of Cabibbo-suppressed semileptonic decays have begun. Experiments are making good progress with D_s^+ semileptonic decays. First results have been obtained for the ratios of form-factor intercepts in $D_s^+ \rightarrow \phi\ell^+\nu_\ell$ and for observations of the pseudoscalar decays $D_s^+ \rightarrow \eta\ell^+\nu_\ell$ and $D_s^+ \rightarrow \eta'\ell^+\nu_\ell$.

In the near future, we can expect further significant advances. It will be very interesting to compare the form factors for Cabibbo-favored, Cabibbo-suppressed, and D_s^+ decays for the pseudoscalar- and vector-meson final states. Measurement of the q^2 dependences of the three vector-decay form factors may help in understanding the puzzle of the small semileptonic transition rate to vector-meson final states.

Continuation of this discussion can be found in the Listings for the B^\pm .

Meson Full Listings

Semileptonic Decays of D^\pm 's, D^\pm

References

1. S. Stone, *Semileptonic B Decays*, HEPHY 93-11, to appear in *B Decays*, 2nd Edition, ed. by S. Stone, World Scientific, Singapore.
2. M. Neubert, *Heavy Quark Symmetry*, SLAC-PUB-6263, June 1993, to appear in *Physics Reports*.
3. J.D. Richman and P.R. Burchat, *Leptonic and Semileptonic Decays of Charm and Bottom Hadrons*, to be published in *Rev. Mod. Phys.*
4. CLEO: A. Bean *et al.*, *Phys. Lett.* **B317**, 647 (1993).
5. E691: J.C. Anjos *et al.*, *Phys. Rev. Lett.* **62**, 1587 (1989).
6. CLEO: G. Crawford *et al.*, *Phys. Rev.* **D44**, 3394 (1991).
7. E687: P.L. Frabetti *et al.*, *Phys. Lett.* **B315**, 203 (1993).
8. MARK III: J. Adler *et al.*, *Phys. Rev. Lett.* **62**, 1821 (1989).
9. E653: K. Kodama *et al.*, *Phys. Lett.* **B313**, 260 (1993).
10. MARK III: Z. Bai *et al.*, *Phys. Rev. Lett.* **66**, 1011 (1991).
11. E691: J.C. Anjos *et al.*, *Phys. Rev. Lett.* **67**, 1507 (1991).
12. E691: J.C. Anjos *et al.*, *Phys. Rev. Lett.* **65**, 2630 (1990).
13. E653: K. Kodama *et al.*, *Phys. Lett.* **B286**, 187 (1992).
14. WA82: M. Adamovich *et al.*, *Phys. Lett.* **B268**, 142 (1991).
15. E687: R.L. Culbertson, Ph.D. thesis, Univ. of Illinois, p. 87 (1993).
16. E687: P.L. Frabetti *et al.*, *Phys. Lett.* **B307**, 262 (1993).
17. E653: K. Kodama *et al.*, *Phys. Lett.* **B313**, 260 (1993).
18. E653: K. Kodama *et al.*, *Phys. Lett.* **B286**, 187 (1992).
19. ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B255**, 634 (1991).
20. MARK III: J. Adler *et al.*, *Phys. Rev. Lett.* **62**, 1821 (1989).
21. CLEO: M.S. Alam *et al.*, *Phys. Rev. Lett.* **71**, 1311 (1993).
22. E653: K. Kodama *et al.*, *Phys. Lett.* **B316**, 455 (1993).
23. MARK III: R. Baltrusaitis *et al.*, *Phys. Rev. Lett.* **54**, 1976 (1985).
24. E653: K. Kodama *et al.*, *Phys. Lett.* **B309**, 483 (1993).
25. CLEO: F. Butler *et al.*, *Phys. Lett.* **B324**, 255 (1994).
26. E687: P.L. Frabetti *et al.*, *Phys. Lett.* **B313**, 253 (1993).
27. ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B255**, 634 (1991).
28. CLEO: J.P. Alexander *et al.*, *Phys. Rev. Lett.* **65**, 1531 (1990).
29. E687: P.L. Frabetti *et al.*, Fermilab Pub-94/049-E6 87, submitted to *Phys. Lett.*
30. N. Isgur *et al.*, *Phys. Rev.* **D39**, 799 (1989).
31. M. Wirbel *et al.*, *Z. Phys.* **C29**, 799 (1985).
32. J.G. Körner and G.A. Schuler, *Z. Phys.* **C38**, 511 (1988).
33. T. Altomari and L. Wolfenstein, *Phys. Rev.* **C37**, 681 (1988);
F.J. Gilman and R.L. Singleton, *Phys. Rev.* **D41**, 1452 (1990).
34. C.W. Bernard, A.X. El-Khadra, and A. Soni, *Phys. Rev.* **D43**, 2140 (1991);
C.W. Bernard, A.X. El-Khadra, and A. Soni, *Phys. Rev.* **D45**, 869 (1992);
C.W. Bernard, A.X. El-Khadra, and A. Soni, *Phys. Rev.* **D47**, 998 (1993).
35. V. Lubicz, G. Martinelli, M.S. McCarthy, and C.T. Sachrajda, *Phys. Lett.* **B274**, 415 (1992).
36. P. Ball, V.M. Braun, and H.G. Dosch, *Phys. Rev.* **D44**, 3567 (1991).
37. M. Aguilar-Benitez *et al.*, *Z. Phys.* **C36**, 559 (1987).
38. M.B. Wise, *Phys. Rev.* **D45**, R2188 (1992);
G. Burdman, Z. Ligeti, M. Neubert, Y. Nir, *Phys. Rev.* **D49**, 2331 (1994).
39. N. Isgur *et al.*, *Phys. Rev.* **D39**, 799 (1989);
M. Bauer, B. Stech, and M. Wirbel, *Z. Phys.* **C34**, 103 (1987);
C.A. Dominguez, *Phys. Lett.* **B207**, 499 (1988);
T.M. Aliev, A.A. Ovchinnikov, and V.A. Slobodenyuk, Trieste Report No. IC/89/382 (to be published);
M. Crisafulli *et al.*, *Phys. Lett.* **B223**, 90 (1989);
G.P. Lepage and S.J. Brodsky, *Phys. Rev.* **D22**, 2157 (1980).
40. M. Wirbel *et al.*, *Z. Phys.* **C29**, 269 (1985);
D. Scora, *Nucl. Phys.* **A743c**, 527 (1991);
D. Scora, Ph.D. thesis, Univ. of Toronto, 1993.

 D^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

 D^\pm MASS

The fit includes D^\pm , D^0 , D_s^\pm , $D^{*\pm}$, D^{*0} , and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1869.4 ± 0.4 OUR FIT				
1869.4 ± 0.5 OUR AVERAGE				
1870.0 ± 0.5 ± 1.0	317	BARLAG	90C ACCM	π^- Cu 230 GeV
1863 ± 4		DERRICK	84 HRS	e^+e^- 29 GeV
1869.4 ± 0.6		¹ TRILLING	81 RVUE	e^+e^- 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1875 ± 10	9	ADAMOVICH	87 EMUL	Photoproduction
1860 ± 16	6	ADAMOVICH	84 EMUL	Photoproduction
1868.4 ± 0.5		¹ SCHINDLER	81 MRK2	e^+e^- 3.77 GeV
1874 ± 5		GOLDHABER	77 MRK1	D^0 , D^+ recoil spectra
1868.3 ± 0.9		¹ PERUZZI	77 MRK1	e^+e^- 3.77 GeV
1874 ± 11		PICCOLO	77 MRK1	e^+e^- 4.03, 4.41 GeV
1876 ± 15	50	PERUZZI	76 MRK1	$K\bar{K}\pi^\pm\pi^\pm$

¹PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision $J/\psi(1S)$ and $\psi(2S)$ measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted.

 D^\pm MEAN LIFE

Measurements with an error $> 0.1 \times 10^{-12}$ s are omitted from the average, and those with an error $> 0.2 \times 10^{-12}$ s have been omitted from the Listings.


VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.057 ± 0.015 OUR AVERAGE				
1.048 ± 0.015 ± 0.011	9k	FRABETTI	94D E687	$D^+ \rightarrow K^- \pi^+ \pi^+$
1.075 ± 0.040 ± 0.018	2455	FRABETTI	91 E687	γ Be, $D^+ \rightarrow K^- \pi^+ \pi^+$
1.03 ± 0.08 ± 0.06	200	ALVAREZ	90 NA14	γ , $D^+ \rightarrow K^- \pi^+ \pi^+$
1.05 $\begin{smallmatrix} +0.077 \\ -0.072 \end{smallmatrix}$	317	² BARLAG	90C ACCM	π^- Cu 230 GeV
1.05 ± 0.08 ± 0.07	363	ALBRECHT	88I ARG	e^+e^- 10 GeV
1.090 ± 0.030 ± 0.025	2992	RAAB	88 E691	Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.12 $\begin{smallmatrix} +0.14 \\ -0.11 \end{smallmatrix}$	149	AGUILAR...	87D HYBR	$\pi^- p$ and pp
1.09 $\begin{smallmatrix} +0.19 \\ -0.15 \end{smallmatrix}$	59	BARLAG	87B ACCM	K^- and π^- 200 GeV
1.14 ± 0.16 ± 0.07	247	CSORNA	87 CLEO	e^+e^- 10 GeV
1.09 ± 0.14	74	³ PALKA	87B SILI	π Be 200 GeV
0.86 ± 0.13 $\begin{smallmatrix} +0.07 \\ -0.03 \end{smallmatrix}$	48	ABE	86 HYBR	γp 20 GeV

²BARLAG 90C estimates the systematic error to be negligible.

³PALKA 87B observes this in $D^+ \rightarrow \bar{K}^*(892)e^+$.

D ⁺ DECAY MODES			
D ⁻ modes are charge conjugates of the modes below.			
Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level	
Inclusive modes			
Γ_1	e^+ anything	(17.2 ± 1.9) %	
Γ_2	K^- anything	(24.2 ± 2.8) %	S=1.4
Γ_3	\bar{K}^0 anything + K^0 anything	(59 ± 7) %	
Γ_4	K^+ anything	(5.8 ± 1.4) %	
Γ_5	η anything	[a] < 13 %	CL=90%
Γ_6	μ^+ anything		
Γ_7	$\mu^+ \mu^-$ anything		
Leptonic and semileptonic modes			
Γ_8	$\mu^+ \nu_\mu$	< 7.2 × 10 ⁻⁴	CL=90%
Γ_9	$\bar{K}^0 e^+ \nu_e$	[b] (6.7 ± 0.8) %	
Γ_{10}	$\bar{K}^0 e^+ \nu_e$	(6.6 ± 0.9) %	
Γ_{11}	$\bar{K}^0 \mu^+ \nu_\mu$	(7.0 $^{+3.0}_{-2.0}$) %	
Γ_{12}	$\bar{K}^0 \ell^+ \nu_\ell$	(6.7 ± 3.5) %	
Γ_{13}	$K^- \pi^+ e^+ \nu_e$	(4.2 $^{+0.9}_{-0.7}$) %	
Γ_{14}	$\bar{K}^*(892)^0 e^+ \nu_e$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(3.2 ± 0.33) %	
Γ_{15}	$K^- \pi^+ e^+ \nu_e$ nonresonant	< 7 × 10 ⁻³	CL=90%
Γ_{16}	$K^- \pi^+ \mu^+ \nu_\mu$ In the fit as $\frac{2}{3}\Gamma_{28} + \Gamma_{18}$, where $\frac{2}{3}\Gamma_{28} = \Gamma_{17}$.	(3.2 ± 1.7) %	
Γ_{17}	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(3.0 ± 0.4) %	
Γ_{18}	$K^- \pi^+ \mu^+ \nu_\mu$ nonresonant	(2.7 ± 1.1) × 10 ⁻³	
Γ_{19}	$\bar{K}^0 \pi^+ \pi^- e^+ \nu_e$		
Γ_{20}	$K^- \pi^+ \pi^0 e^+ \nu_e$		
Γ_{21}	$(\bar{K}^*(892)\pi)^0 e^+ \nu_e$	< 1.2 %	CL=90%
Γ_{22}	$(\bar{K}\pi\pi)^0 e^+ \nu_e$ non- $\bar{K}^*(892)$	< 9 × 10 ⁻³	CL=90%
Γ_{23}	$K^- \pi^+ \pi^0 \mu^+ \nu_\mu$	< 1.4 × 10 ⁻³	CL=90%
Γ_{24}	$\pi^0 \ell^+ \nu_\ell$	[c] (5.7 ± 2.2) × 10 ⁻³	
Γ_{25}	$\pi^+ \pi^- e^+ \nu_e$ Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.		
Γ_{26}	$\bar{K}^*(892)^0 e^+ \nu_e$	[b] (4.8 ± 0.4) %	
Γ_{27}	$\bar{K}^*(892)^0 e^+ \nu_e$	(4.8 ± 0.5) %	
Γ_{28}	$\bar{K}^*(892)^0 \mu^+ \nu_\mu$	(4.5 ± 0.6) %	S=1.1
Γ_{29}	$\rho^0 e^+ \nu_e$	< 3.7 × 10 ⁻³	CL=90%
Γ_{30}	$\rho^0 \mu^+ \nu_\mu$	(2.0 $^{+1.5}_{-1.3}$) × 10 ⁻³	
Γ_{31}	$\phi e^+ \nu_e$	< 2.09 %	CL=90%
Γ_{32}	$\phi \mu^+ \nu_\mu$	< 3.72 %	CL=90%
Γ_{33}	$\eta'(958) \mu^+ \nu_\mu$	< 9 × 10 ⁻³	CL=90%
Hadronic modes with one or three K's			
Γ_{34}	$\bar{K}^0 \pi^+$	(2.74 ± 0.29) %	
Γ_{35}	$K^- \pi^+ \pi^+$	[d] (9.1 ± 0.6) %	
Γ_{36}	$\bar{K}^*(892)^0 \pi^+$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(1.5 ± 0.3) %	
Γ_{37}	$\bar{K}_0(1430)^0 \pi^+$ × B($\bar{K}^*(1430)^0 \rightarrow K^- \pi^+$)	(2.3 ± 0.3) %	
Γ_{38}	$\bar{K}^*(1680)^0 \pi^+$ × B($\bar{K}^*(1680)^0 \rightarrow K^- \pi^+$)	(2.6 ± 1.3) × 10 ⁻³	
Γ_{39}	$K^- \pi^+ \pi^+$ nonresonant	(7.3 ± 1.4) %	
Γ_{40}	$\bar{K}^0 \pi^+ \pi^0$	[d] (9.7 ± 3.0) %	S=1.1
Γ_{41}	$\bar{K}^0 \rho^+$	(6.6 ± 2.5) %	
Γ_{42}	$\bar{K}^*(892)^0 \pi^+$ × B($\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0$)	(0.7 ± 0.2) %	
Γ_{43}	$\bar{K}^0 \pi^+ \pi^0$ nonresonant	(1.3 ± 1.1) %	
Γ_{44}	$K^- \pi^+ \pi^+ \pi^0$	[d] (6.4 ± 1.1) %	
Γ_{45}	$\bar{K}^*(892)^0 \rho^+$ total × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(1.4 ± 0.9) %	
Γ_{46}	$\bar{K}_1(1400)^0 \pi^+$ × B($\bar{K}_1(1400)^0 \rightarrow K^- \pi^+ \pi^0$)	(2.2 ± 0.6) %	
Γ_{47}	$K^- \rho^+ \pi^+$ total	(3.1 ± 1.1) %	
Γ_{48}	$\bar{K}^*(892)^0 \pi^+ \pi^0$ total × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(4.5 ± 0.9) %	
Γ_{49}	$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(2.8 ± 0.9) %	
Γ_{50}	$K^*(892)^- \pi^+ \pi^+ 3$ -body × B($K^{*-} \rightarrow K^- \pi^0$)	(1.4 ± 0.6) %	
Γ_{51}	$K^- \pi^+ \pi^+ \pi^0$ nonresonant	[e] (1.2 ± 0.6) %	
Γ_{52}	$\bar{K}^0 \pi^+ \pi^+ \pi^-$	[d] (7.0 ± 1.0) %	
Γ_{53}	$\bar{K}^0 a_1(1260)^+$ × B($a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-$)	(4.0 ± 0.8) %	
Γ_{54}	$\bar{K}_1(1400)^0 \pi^+$ × B($\bar{K}_1(1400)^0 \rightarrow \bar{K}^0 \pi^+ \pi^-$)	(2.2 ± 0.6) %	
Γ_{55}	$K^*(892)^- \pi^+ \pi^+ 3$ -body × B($K^{*-} \rightarrow \bar{K}^0 \pi^-$)	(1.4 ± 0.6) %	
Γ_{56}	$\bar{K}^0 \rho^0 \pi^+$ total	(4.2 ± 0.9) %	
Γ_{57}	$\bar{K}^0 \pi^+ \pi^+ \pi^-$ nonresonant	(8 ± 4) × 10 ⁻³	
Γ_{58}	$K^- \pi^+ \pi^+ \pi^+ \pi^-$	(8.2 ± 1.4) × 10 ⁻³	
Γ_{59}	$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(6.8 ± 1.8) × 10 ⁻³	
Γ_{60}	$\bar{K}^*(892)^0 \rho^0 \pi^+$ × B($\bar{K}^{*0} \rightarrow K^- \pi^+$)	(5.1 ± 2.2) × 10 ⁻³	
Γ_{61}	$K^- \pi^+ \pi^+ \pi^0 \pi^0$	(2.2 $^{+5.0}_{-0.9}$) %	
Γ_{62}	$\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^0$	(5.4 $^{+3.0}_{-1.4}$) %	
Γ_{63}	$\bar{K}^0 \pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(8 ± 7) × 10 ⁻⁴	
Γ_{64}	$K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0$	(2.0 ± 1.8) × 10 ⁻³	
Γ_{65}	$\bar{K}^0 \bar{K}^0 K^+$	(3.1 ± 0.7) %	
Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.			
Γ_{66}	$\bar{K}^0 \rho^+$	(6.6 ± 2.5) %	
Γ_{67}	$\bar{K}^0 a_1(1260)^+$	(8.1 ± 1.7) %	
Γ_{68}	$\bar{K}^0 a_2(1320)^+$	< 3 × 10 ⁻³	CL=90%
Γ_{69}	$\bar{K}^*(892)^0 \pi^+$	(2.2 ± 0.4) %	
Γ_{70}	$\bar{K}^*(892)^0 \rho^+$ total	(2.1 ± 1.4) %	
Γ_{71}	$\bar{K}^*(892)^0 \rho^+$ S-wave	[e] (1.7 ± 1.6) %	
Γ_{72}	$\bar{K}^*(892)^0 \rho^+$ P-wave	< 1 × 10 ⁻³	CL=90%
Γ_{73}	$\bar{K}^*(892)^0 \rho^+$ D-wave	(10 ± 7) × 10 ⁻³	
Γ_{74}	$\bar{K}^*(892)^0 \rho^+$ D-wave longitudinal	< 7 × 10 ⁻³	CL=90%
Γ_{75}	$\bar{K}_1(1270)^0 \pi^+$	< 7 × 10 ⁻³	CL=90%
Γ_{76}	$\bar{K}_1(1400)^0 \pi^+$	(5.0 ± 1.3) %	
Γ_{77}	$\bar{K}^*(1410)^0 \pi^+$	< 7 × 10 ⁻³	CL=90%
Γ_{78}	$\bar{K}_0^*(1430)^0 \pi^+$	(3.4 ± 0.4) %	
Γ_{79}	$\bar{K}^*(1680)^0 \pi^+$	(1.0 ± 0.5) %	
Γ_{80}	$\bar{K}^*(892)^0 \pi^+ \pi^0$ total	(6.7 ± 1.4) %	
Γ_{81}	$\bar{K}^*(892)^0 \pi^+ \pi^0$ 3-body	(4.2 ± 1.4) %	
Γ_{82}	$K^*(892)^- \pi^+ \pi^+ 3$ -body	(2.1 ± 0.9) %	
Γ_{83}	$K^- \rho^+ \pi^+$ total	(3.1 ± 1.1) %	
Γ_{84}	$K^- \rho^+ \pi^+$ 3-body	(1.1 ± 0.4) %	
Γ_{85}	$\bar{K}^0 \rho^0 \pi^+$ total	(4.2 ± 0.9) %	CL=90%
Γ_{86}	$\bar{K}^0 \rho^0 \pi^+$ 3-body	(5 ± 5) × 10 ⁻³	
Γ_{87}	$\bar{K}^0 f_0(980) \pi^+$	< 5 × 10 ⁻³	CL=90%
Γ_{88}	$\bar{K}^*(892)^0 \pi^+ \pi^+ \pi^-$	(1.02 ± 0.27) %	
Γ_{89}	$\bar{K}^*(892)^0 \rho^0 \pi^+$	(7.7 ± 3.3) × 10 ⁻³	
Plonic modes			
Γ_{90}	$\pi^+ \pi^0$	(2.5 ± 0.7) × 10 ⁻³	
Γ_{91}	$\pi^+ \pi^+ \pi^-$	(3.2 ± 0.6) × 10 ⁻³	
Γ_{92}	$\rho^0 \pi^+$	< 1.4 × 10 ⁻³	CL=90%
Γ_{93}	$\pi^+ \pi^+ \pi^-$ nonresonant	(2.5 ± 0.7) × 10 ⁻³	
Γ_{94}	$\pi^+ \pi^+ \pi^- \pi^0$	(1.9 $^{+1.5}_{-1.2}$) %	
Γ_{95}	$\eta \pi^+ \times B(\eta \rightarrow \pi^+ \pi^- \pi^0)$	(1.8 ± 0.6) × 10 ⁻³	
Γ_{96}	$\omega \pi^+ \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	< 6 × 10 ⁻³	CL=90%
Γ_{97}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(1.0 $^{+0.8}_{-0.7}$) × 10 ⁻³	
Γ_{98}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	(2.9 $^{+2.9}_{-2.0}$) × 10 ⁻³	
Fractions of some of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.			
Γ_{99}	$\rho^0 \pi^+$	< 1.4 × 10 ⁻³	CL=90%
Γ_{100}	$\eta \pi^+$	(7.5 ± 2.5) × 10 ⁻³	
Γ_{101}	$\omega \pi^+$	< 7 × 10 ⁻³	CL=90%
Γ_{102}	$\eta \rho^+$	< 1.2 %	CL=90%
Γ_{103}	$\eta'(958) \pi^+$	< 9 × 10 ⁻³	CL=90%
Γ_{104}	$\eta'(958) \rho^+$	< 1.5 %	CL=90%

$\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$		Γ_2/Γ	
VALUE	EVTs	DOCUMENT ID	TECN COMMENT
0.242 ± 0.028 OUR AVERAGE		Error Includes scale factor of 1.4. See the Ideogram below.	
0.278 ^{+0.036} _{-0.031}		9 BARLAG	92C ACCM π^- Cu 230 GeV
0.271 ± 0.023 ± 0.024		COFFMAN	91 MRK3 e^+e^- 3.77 GeV
0.17 ± 0.07		AGUILAR-...	87E HYBR $\pi p, pp$ 360, 400 GeV
0.19 ± 0.05	26	SCHINDLER	81 MRK2 e^+e^- 3.771 GeV
0.10 ± 0.07	3	VUILLEMIN	78 MRK1 e^+e^- 3.772 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
0.16 ^{+0.08} _{-0.07}		AGUILAR-...	86B HYBR See AGUILAR-BENITEZ 87E
9 BARLAG 92c computes the branching fraction using topological normalization.			



$[\Gamma(K^0 \text{ anything}) + \Gamma(K^+ \text{ anything})]/\Gamma_{\text{total}}$		Γ_3/Γ	
VALUE	EVTs	DOCUMENT ID	TECN COMMENT
0.59 ± 0.07 OUR AVERAGE			
0.612 ± 0.065 ± 0.043		COFFMAN	91 MRK3 e^+e^- 3.77 GeV
0.52 ± 0.18	15	SCHINDLER	81 MRK2 e^+e^- 3.771 GeV
0.39 ± 0.29	3	VUILLEMIN	78 MRK1 e^+e^- 3.772 GeV

$\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$		Γ_4/Γ	
VALUE	EVTs	DOCUMENT ID	TECN COMMENT
0.058 ± 0.014 OUR AVERAGE			
0.055 ± 0.013 ± 0.009		COFFMAN	91 MRK3 e^+e^- 3.77 GeV
0.08 ^{+0.06} _{-0.05}		AGUILAR-...	87E HYBR $\pi p, pp$ 360, 400 GeV
0.06 ± 0.04	12	SCHINDLER	81 MRK2 e^+e^- 3.771 GeV
0.06 ± 0.06	2	VUILLEMIN	78 MRK1 e^+e^- 3.772 GeV

D⁺ and D⁰ → (η anything) / (total D⁺ and D⁰)
 If measured at the ψ(3770), this quantity is a weighted average of D⁺ (44%) and D⁰ (56%) branching fractions. Only the experiment at E_{cm} = 3.77 GeV is used.

VALUE	DOCUMENT ID	TECN	COMMENT
<0.13	PARTRIDGE 81	CBAL	e^+e^- 3.77 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
<0.02	10 BRANDELIK 79	DASP	e^+e^- 4.03 GeV

10 The BRANDELIK 79 result is based on the absence of an η signal at E_{cm} = 4.03 GeV. PARTRIDGE 81 observes a substantially higher η cross section at 4.03 GeV.

$\Gamma(c/\bar{c} \rightarrow \mu^+ \text{ anything})/\Gamma(c/\bar{c} \rightarrow \text{ anything})$	
VALUE	EVTs
0.081 ± 0.010 OUR AVERAGE	
0.086 ± 0.017 ^{+0.008} _{-0.007}	69
0.078 ± 0.009 ± 0.012	11 ALBRECHT 92F
0.078 ± 0.015 ± 0.02	ONG 88
0.082 ± 0.012 ^{+0.02} _{-0.01}	MRK2 87
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●	
0.089 ± 0.018 ± 0.025	JADE 85J
11 ALBRECHT 92F uses the excess of right-sign over wrong-sign leptons in a sample of events tagged by fully reconstructed D*(2010) ⁺ → D ⁰ π ⁺ decays.	

$\Gamma(c/\bar{c} \rightarrow e^+ e^- \text{ anything})/\Gamma(c/\bar{c} \rightarrow \text{ anything})$	
VALUE	CL% EVTS
<2.2 × 10 ⁻³	90
12 The normalization uses a continuum charm production estimate.	

$\Gamma(c/\bar{c} \rightarrow e^+ \mu^- \text{ anything})/\Gamma(c/\bar{c} \rightarrow \text{ anything})$	
VALUE	CL% EVTS
<3.7 × 10 ⁻³	90
13 The normalization uses a continuum charm production estimate.	

$\Gamma(c/\bar{c} \rightarrow \mu^+ \mu^- \text{ anything})/\Gamma(c/\bar{c} \rightarrow \text{ anything})$	
VALUE	CL% EVTS
<0.018	90
<0.007	95
14 The normalization uses a continuum charm production estimate.	
15 Average BR for charm → μ ⁺ μ ⁻ X. The mixture of charmed particles is unknown and may actually contain states other than D mesons.	

———— Leptonic and semileptonic modes ————

$\Gamma(\mu^+ \nu_\mu)/\Gamma_{\text{total}}$		Γ_8/Γ	
VALUE	CL% EVTS	DOCUMENT ID	TECN COMMENT
<0.00072	90	ADLER	88B MRK3 e^+e^- 3.77 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
<0.02	90	0	16 AUBERT 83 SPEC μ ⁺ Fe, 250 GeV
16 AUBERT 83 obtains an upper limit 0.014 assuming the final state contains equal amounts of (D ⁺ , D ⁻), (D ⁺ , D ⁰), (D ⁻ , D ⁰), and (D ⁰ , D ⁰). We quote the limit they get under more general assumptions.			

$\Gamma(K^0 \rightarrow e^+ \nu_e)/\Gamma_{\text{total}}$		Γ_9/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT
0.067 ± 0.008 OUR AVERAGE			
0.066 ± 0.009	PDG	94	Our $\Gamma(K^0 \rightarrow e^+ \nu_e)/\Gamma_{\text{total}}$
0.072 ^{+0.031} _{-0.021}	PDG	94	1.03 × our $\Gamma(K^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$

$\Gamma(K^0 \rightarrow e^+ \nu_e)/\Gamma_{\text{total}}$		Γ_{10}/Γ	
VALUE	EVTs	DOCUMENT ID	TECN COMMENT
0.066 ± 0.009 OUR FIT			
0.06 ^{+0.022} _{-0.013} ± 0.007	13	BAI	91 MRK3 e^+e^- ≈ 3.77 GeV

$\Gamma(K^0 \rightarrow e^+ \nu_e)/\Gamma(K^0 \pi^+)$		Γ_{10}/Γ_{34}	
VALUE	EVTs	DOCUMENT ID	TECN COMMENT
2.39 ± 0.33 OUR FIT			
2.60 ± 0.35 ± 0.26	186	17 BEAN	93C CLEO e^+e^- ≈ 7(45)
17 BEAN 93C uses $\bar{K}^0 \mu^+ \nu_\mu$ as well as $\bar{K}^0 \rightarrow e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ ⁺ events to use them as e ⁺ events.			

$\Gamma(K^0 \rightarrow e^+ \nu_e)/\Gamma(K^- \pi^+ \pi^+)$		Γ_{10}/Γ_{35}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.10 OUR FIT			
0.66 ± 0.09 ± 0.14	ANJOS	91C	E691 γ Be 80–240 GeV

$\Gamma(K^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$		Γ_{11}/Γ	
VALUE	EVTs	DOCUMENT ID	TECN COMMENT
0.07 ^{+0.028} _{-0.016} ± 0.012	14	BAI	91 MRK3 e^+e^- ≈ 3.77 GeV

$\Gamma(K^0 \mu^+ \nu_\mu)/\Gamma(\mu^+ \text{ anything})$		Γ_{11}/Γ_6	
VALUE	EVTs	DOCUMENT ID	COMMENT
0.76 ± 0.06	84	18 AOKI	88 π ⁻ emulsion
18 From topological branching ratios in emulsion with an identified muon.			

$\Gamma(K^- \pi^+ e^+ \nu_e)/\Gamma_{\text{total}}$		Γ_{13}/Γ	
VALUE	CL% EVTS	DOCUMENT ID	TECN COMMENT
0.042 ± 0.009 OUR FIT			
0.035 ^{+0.012} _{-0.007} ± 0.004	14	19 BAI	91 MRK3 e^+e^- ≈ 3.77 GeV
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
<0.057	90	20 AGUILAR-...	87F HYBR $\pi p, pp$ 360, 400 GeV
19 BAI 91 finds that a fraction 0.79 ^{+0.15+0.09} _{-0.17-0.03} of combined D ⁺ and D ⁰ decays to $\bar{K}^0 \pi^+ \nu_e$ (24 events) are $\bar{K}^*(892) \rightarrow e^+ \nu_e$.			
20 AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.			

Meson Full Listings

 D^\pm

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{26}/Γ
 We average our $\bar{K}^{*0} e^+ \nu_e$ and $\bar{K}^{*0} \mu^+ \nu_\mu$ branching fractions, multiplying the latter by a phase-space factor of 1.05 to be able to use it with the $\bar{K}^{*0} e^+ \nu_e$ fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
0.048 ± 0.004 OUR AVERAGE			
0.048 ± 0.005	PDG	94	Our $\Gamma(\bar{K}^{*0} e^+ \nu_e)/\Gamma_{\text{total}}$
0.047 ± 0.006	PDG	94	1.05 × our $\Gamma(\bar{K}^{*0} \mu^+ \nu_\mu)/\Gamma_{\text{total}}$

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)/\Gamma(K^- \pi^+ e^+ \nu_e)$ Γ_{27}/Γ_{13}
 Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.16^{+0.21}_{-0.24} OUR FIT				
1.0 ± 0.3	35	ADAMOVIČH	91	OMEG π^- 340 GeV

$\Gamma(\bar{K}^*(892)^0 e^+ \nu_e)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{27}/Γ_{35}
 Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.53 ± 0.05 OUR FIT				
0.54 ± 0.05 OUR AVERAGE				
0.67 ± 0.09 ± 0.07	710	21 BEAN	93C	CLEO $e^+ e^- \approx \gamma(4S)$
0.62 ± 0.15 ± 0.09	35	ADAMOVIČH	91	OMEG π^- 340 GeV
0.55 ± 0.08 ± 0.10	880	ALBRECHT	91	ARG $e^+ e^- \approx 10.4$ GeV
0.49 ± 0.04 ± 0.05		22 ANJOS	89B	E691 Photoproduction

²¹BEAN 93C uses $\bar{K}^{*0} \mu^+ \nu_\mu$ as well as $\bar{K}^{*0} e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

²²For measurements of the form factors for this mode, and of the ratio of longitudinal to transverse polarization of the $\bar{K}^*(892)$, see ANJOS 90E.

$\Gamma(K^- \pi^+ e^+ \nu_e \text{ nonresonant})/\Gamma_{\text{total}}$ Γ_{15}/Γ
 Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	23 ANJOS	89B	E691 Photoproduction

²³ANJOS 89B assumes a $\Gamma(D^+ \rightarrow K^- \pi^+ \pi^+)/\Gamma_{\text{total}} = 9.1 \pm 1.3 \pm 0.4\%$.

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{28}/Γ
 Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.045 ± 0.006 OUR FIT				Error includes scale factor of 1.1.
0.0325 ± 0.0071 ± 0.0075	224	24 KODAMA	92C	E653 π^- emulsion 600 GeV

²⁴KODAMA 92C measures $\Gamma(D^+ \rightarrow \bar{K}^{*0} \mu^+ \nu_\mu)/\Gamma(D^+ \rightarrow K^- \mu^+ \nu_\mu) = 0.43 \pm 0.09 \pm 0.09$ and then uses $\Gamma(D^0 \rightarrow K^- \mu^+ \nu_\mu) = (7.0 \pm 0.7) \times 10^{10} \text{ s}^{-1}$ to get the quoted branching fraction. See also the footnote to KODAMA 92C in the next data block.

$\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{28}/Γ_{35}
 Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.49 ± 0.06 OUR FIT				Error includes scale factor of 1.1.
0.53 ± 0.06 OUR AVERAGE				
0.56 ± 0.04 ± 0.06	875	25 FRABETTI	93E	E687 γ Be $\bar{E}_\gamma \approx 200$ GeV
0.46 ± 0.07 ± 0.08	224	26 KODAMA	92C	E653 π^- emulsion 600 GeV

²⁵FRABETTI 93E also gives measurements of the form factors and of the ratio of longitudinal to transverse polarizations of the \bar{K}^{*0} .

²⁶KODAMA 92C uses the same $\bar{K}^{*0} \mu^+ \nu_\mu$ events normalizing instead with $D^0 \rightarrow K^- \mu^+ \nu_\mu$ events, as reported in the preceding data block. Measurements of form factors and of the ratio of longitudinal to transverse polarizations of the \bar{K}^{*0} are given in KODAMA 92.

$\Gamma(K^- \pi^+ \mu^+ \nu_\mu \text{ nonresonant})/\Gamma(K^- \pi^+ \mu^+ \nu_\mu)$ $\Gamma_{18}/\Gamma_{16} = \Gamma_{18}/(\Gamma_{18} + \frac{2}{3}\Gamma_{28})$

VALUE	DOCUMENT ID	TECN	COMMENT
0.083 ± 0.029 OUR FIT			
0.083 ± 0.029	27 FRABETTI	93E	E687 γ Be $\bar{E}_\gamma \approx 200$ GeV

²⁷This FRABETTI 93E value is equivalent to <0.12 at the 90% confidence level.

$\Gamma(\bar{K}^0 \pi^+ \pi^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.022^{+0.047}_{-0.006} ± 0.004	1	28 AGUILAR-...	87F	HYBR π , p , p 360, 400 GeV

²⁸AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.044^{+0.052}_{-0.013} ± 0.007	2	29 AGUILAR-...	87F	HYBR π , p , p 360, 400 GeV

²⁹AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma((\bar{K}^*(892)\pi)^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.012	90	ANJOS	92	E691 Photoproduction

$\Gamma((\bar{K}\pi\pi)^0 e^+ \nu_e \text{ non-}\bar{K}^*(892))/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	ANJOS	92	E691 Photoproduction

$\Gamma(K^- \pi^+ \pi^0 \mu^+ \nu_\mu)/\Gamma(K^- \pi^+ \mu^+ \nu_\mu)$ $\Gamma_{23}/\Gamma_{16} = \Gamma_{23}/(\Gamma_{18} + \frac{2}{3}\Gamma_{28})$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.042	90	FRABETTI	93E	E687 γ Be $\bar{E}_\gamma \approx 200$ GeV

$\Gamma(\pi^0 \ell^+ \nu_\ell)/\Gamma(\bar{K}^0 \ell^+ \nu_\ell)$ Γ_{24}/Γ_{12}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.085 ± 0.027 ± 0.014	53	30 ALAM	93	CLEO $e^+ e^- \approx \gamma(4S)$

³⁰ALAM 93 thus directly measures the product of ratios squared of CKM matrix elements and form factors at $q^2=0$: $|V_{cd}/V_{cs}|^2 \cdot |f_{\pi^+}^2(0)/f_{\bar{K}^0}^2(0)|^2 = 0.085 \pm 0.027 \pm 0.014$.

$\Gamma(\pi^+ \pi^- e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.057	90	31 AGUILAR-...	87F	HYBR π , p , p 360, 400 GeV

³¹AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(\rho^0 e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0037	90	BAI	91	MRK3 $e^+ e^- \approx 3.77$ GeV

$\Gamma(\rho^0 \mu^+ \nu_\mu)/\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ Γ_{30}/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.044^{+0.031}_{-0.025} ± 0.014	4	32 KODAMA	93C	E653 π^- emulsion 600 GeV

³²This KODAMA 93C result is based on a final signal of $4.0 \pm \frac{2.8}{2.3} \pm 1.3$ events; the estimates of backgrounds that affect this number are somewhat model dependent.

$\Gamma(\phi e^+ \nu_e)/\Gamma_{\text{total}}$ Γ_{31}/Γ
 Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0209	90	BAI	91	MRK3 $e^+ e^- \approx 3.77$ GeV

$\Gamma(\phi \mu^+ \nu_\mu)/\Gamma_{\text{total}}$ Γ_{32}/Γ
 Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0372	90	BAI	91	MRK3 $e^+ e^- \approx 3.77$ GeV

$\Gamma(\eta'(958) \mu^+ \nu_\mu)/\Gamma(\bar{K}^*(892)^0 \mu^+ \nu_\mu)$ Γ_{33}/Γ_{28}
 Decay modes of the $\eta'(958)$ not included in the search are corrected for.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.20	90	KODAMA	93B	E653 π^- emulsion 600 GeV

Hadronic modes with one or three K 's

$\Gamma(\bar{K}^0 \pi^+)/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0274 ± 0.0029 OUR FIT				
0.032 ± 0.004 OUR AVERAGE				
0.032 ± 0.005 ± 0.002	161	ADLER	88C	MRK3 $e^+ e^- 3.77$ GeV
0.033 ± 0.009	36	33 SCHINDLER	81	MRK2 $e^+ e^- 3.771$ GeV
0.033 ± 0.013	17	34 PERUZZI	77	MRK1 $e^+ e^- 3.77$ GeV

³³SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.14 ± 0.03 . We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³⁴PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.14 ± 0.05 . We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

$\Gamma(\bar{K}^0 \pi^+)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{34}/Γ_{35}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.302 ± 0.031 OUR FIT				
0.274 ± 0.030 ± 0.031	264	ANJOS	90C	E691 Photoproduction

$\Gamma(K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.091 ± 0.006 OUR FIT				
0.091 ± 0.007 OUR AVERAGE				
0.093 ± 0.006 ± 0.008	1502	35 BALEST	94	CLEO $e^+ e^- \approx \gamma(4S)$
0.091 ± 0.013 ± 0.004	1164	ADLER	88C	MRK3 $e^+ e^- 3.77$ GeV
0.091 ± 0.019	239	36 SCHINDLER	81	MRK2 $e^+ e^- 3.771$ GeV
0.086 ± 0.020	85	37 PERUZZI	77	MRK1 $e^+ e^- 3.77$ GeV

³⁵BALEST 94 measures the ratio of $D^+ \rightarrow K^- \pi^+ \pi^+$ and $D^0 \rightarrow K^- \pi^+$ branching fractions to be $2.35 \pm 0.16 \pm 0.16$ and uses their absolute measurement of the $D^0 \rightarrow K^- \pi^+$ fraction (AKERIB 93).

³⁶SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.38 ± 0.05 . We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³⁷PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.36 ± 0.06 . We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

³⁸AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

$\Gamma(\bar{K}^*(892)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{69} / \Gamma_{35}$ Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.25 ± 0.04				OUR FIT
0.24 ± 0.04				OUR AVERAGE
0.255 ± 0.014 ± 0.050		ANJOS	93 E691	γ Be 90–260 GeV
0.21 ± 0.06 ± 0.06		ALVAREZ	91B NA14	Photoproduction
0.20 ± 0.02 ± 0.11		ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.053	90	SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV

 $\Gamma(\bar{K}_0^*(1430)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{78} / \Gamma_{35}$ Unseen decay modes of the $\bar{K}_0^*(1430)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.372 ± 0.029 ± 0.025	ANJOS	93 E691	γ Be 90–260 GeV

 $\Gamma(\bar{K}^*(1680)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{79} / \Gamma_{35}$ Unseen decay modes of the $\bar{K}^*(1680)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.113 ± 0.015 ± 0.050	ANJOS	93 E691	γ Be 90–260 GeV

 $\Gamma(K^- \pi^+ \pi^+ \text{ nonresonant}) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{39} / \Gamma_{35}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.80 ± 0.14			OUR AVERAGE
0.838 ± 0.088 ± 0.275	ANJOS	93 E691	γ Be 90–260 GeV
0.79 ± 0.07 ± 0.15	ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}^0 \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{40} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.097 ± 0.030				OUR FIT
0.107 ± 0.029				OUR AVERAGE
0.102 ± 0.025 ± 0.016	159	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV
0.19 ± 0.12	10	39 SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV

³⁹SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.78 ± 0.48 . We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

 $\Gamma(\bar{K}^0 \rho^+) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{41} / \Gamma_{40}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.68 ± 0.08 ± 0.12	ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}^*(892)^0 \pi^+) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{69} / \Gamma_{40}$ Includes a factor 3 to take into account $\bar{K}^*(892)^0 \rightarrow K^- \pi^+$.

VALUE	DOCUMENT ID	TECN	COMMENT
0.23 ± 0.07			OUR FIT
0.57 ± 0.18 ± 0.18	ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}^0 \pi^+ \pi^0 \text{ nonresonant}) / \Gamma(\bar{K}^0 \pi^+ \pi^0)$ $\Gamma_{43} / \Gamma_{40}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.07 ± 0.08	ADLER	87 MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(K^- \pi^+ \pi^+ \pi^0) / \Gamma_{\text{total}}$ Γ_{44} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.064 ± 0.011				OUR FIT
0.088 ± 0.012 ± 0.012	142	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.034 \pm $\frac{+0.056}{-0.070}$	40	BARLAG	92C ACCM	π^- Cu 230 GeV
0.022 \pm $\frac{+0.047}{-0.006}$ ± 0.004	1	40 AGUILAR-...	87F HYBR	$\pi p, p p$ 360, 400 GeV

0.063 \pm $\frac{+0.014}{-0.013}$ ± 0.012	175	BALTRUSAIT...86E	MRK3	See COFFMAN 92B
---	-----	------------------	------	-----------------

⁴⁰AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

 $\Gamma(K^- \pi^+ \pi^+ \pi^0) / \Gamma(K^- \pi^+ \pi^+)$ $\Gamma_{44} / \Gamma_{35}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.70 ± 0.12				OUR FIT
0.76 ± 0.11 ± 0.12	91	ANJOS	92C E691	γ Be 90–260 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.69 ± 0.10 ± 0.16		ANJOS	89E E691	See ANJOS 92C
0.57 \pm $\frac{+0.65}{-0.17}$	1	AGUILAR-...	83B HYBR	$\pi^- p$, 360 GeV

 $\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ total}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{70} / \Gamma_{44}$ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.165 ± 0.12	ANJOS	92C E691	γ Be 90–260 GeV

 $\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ S-wave}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{71} / \Gamma_{44}$ Unseen decay modes of the $\bar{K}^*(892)^0$ are included. The two experiments disagree severely here.

VALUE	DOCUMENT ID	TECN	COMMENT
0.26 ± 0.25			OUR AVERAGE
0.15 ± 0.075 ± 0.045	ANJOS	92C E691	γ Be 90–260 GeV
0.833 ± 0.116 ± 0.165	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ P-wave}) / \Gamma_{\text{total}}$ Γ_{72} / Γ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.001	90	ANJOS	92C E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.005	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ D-wave}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{73} / \Gamma_{44}$ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.15 ± 0.09 ± 0.045	ANJOS	92C E691	γ Be 90–260 GeV

 $\Gamma(\bar{K}^*(892)^0 \rho^+ \text{ D-wave longitudinal}) / \Gamma_{\text{total}}$ Γ_{74} / Γ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}_1(1400)^0 \pi^+) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{76} / \Gamma_{44}$ Unseen decay modes of the $\bar{K}_1(1400)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.77 ± 0.20			OUR FIT
0.907 ± 0.218 ± 0.180	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(K^- \rho^+ \pi^+ \text{ total}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{83} / \Gamma_{44}$ This includes $\bar{K}^*(892)^0 \rho^+$, etc. The next entry gives the specifically 3-body fraction.

VALUE	DOCUMENT ID	TECN	COMMENT
0.48 ± 0.13 ± 0.09	ANJOS	92C E691	γ Be 90–260 GeV

 $\Gamma(K^- \rho^+ \pi^+ \text{ 3-body}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{84} / \Gamma_{44}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.06			OUR AVERAGE
0.18 ± 0.08 ± 0.04	ANJOS	92C E691	γ Be 90–260 GeV
0.159 ± 0.065 ± 0.060	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ total}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{80} / \Gamma_{44}$ This includes $\bar{K}^*(892)^0 \rho^+$, etc. The next two entries gives the specifically 3-body fraction. Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
1.05 ± 0.11 ± 0.08	ANJOS	92C E691	γ Be 90–260 GeV

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ 3-body}) / \Gamma_{\text{total}}$ Γ_{81} / Γ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.008	90	41 COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
⁴¹ See, however, the next entry: ANJOS 92C sees a large signal in this channel.				

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^0 \text{ 3-body}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{81} / \Gamma_{44}$ Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.66 ± 0.09 ± 0.17	ANJOS	92C E691	γ Be 90–260 GeV

 $\Gamma(K^*(892)^- \pi^+ \pi^+ \text{ 3-body}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{82} / \Gamma_{44}$ Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.14			OUR FIT
0.24 ± 0.12 ± 0.09	ANJOS	92C E691	γ Be 90–260 GeV

 $\Gamma(K^- \pi^+ \pi^+ \pi^0 \text{ nonresonant}) / \Gamma_{\text{total}}$ Γ_{51} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.002	90	42 ANJOS	92C E691	γ Be 90–260 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
⁴² Whereas ANJOS 92C finds no signal here, COFFMAN 92B finds a fairly large one; see the next entry.				

 $\Gamma(K^- \pi^+ \pi^+ \pi^0 \text{ nonresonant}) / \Gamma(K^- \pi^+ \pi^+ \pi^0)$ $\Gamma_{51} / \Gamma_{44}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.184 ± 0.070 ± 0.050	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV

 $\Gamma(\bar{K}^0 \pi^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{52} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.070 ± 0.010				OUR FIT
0.071 ± 0.016				OUR AVERAGE

0.066 ± 0.015 ± 0.005	168	ADLER	88C MRK3	$e^+ e^-$ 3.77 GeV
0.12 ± 0.05	21	43 SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.042 \pm $\frac{+0.019}{-0.017}$	44	BARLAG	92C ACCM	π^- Cu 230 GeV
-------------------------------------	----	--------	----------	--------------------

0.243 \pm $\frac{+0.064}{-0.041}$ ± 0.041	11	44 AGUILAR-...	87F HYBR	$\pi p, p p$ 360, 400 GeV
---	----	----------------	----------	---------------------------

⁴³SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.51 ± 0.08 . We use the MARK-3 (ADLER 88C) value of $\sigma = 4.2 \pm 0.6 \pm 0.3$ nb.

⁴⁴AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.

Meson Full Listings

 D^\pm

$\Gamma(K^0 \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+)$					$\Gamma_{52} / \Gamma_{35}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.78 ± 0.10 OUR FIT					
0.77 ± 0.07 ± 0.11	229	ANJOS	92C E691	γ Be 90–260 GeV	

$\Gamma(K^0 a_1(1260)^+) / \Gamma(K^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{67} / \Gamma_{52}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $a_1(1260)^+$ are included.					
1.15 ± 0.19 OUR AVERAGE				Error includes scale factor of 1.1.	
1.66 ± 0.28 ± 0.40		ANJOS	92C E691	γ Be 90–260 GeV	
1.078 ± 0.114 ± 0.140		COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K^0 a_2(1320)^+) / \Gamma_{total}$					Γ_{68} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $a_2(1320)^+$ are included.					
<0.003	90	ANJOS	92C E691	γ Be 90–260 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.008	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K_1(1270)^0 \pi^+) / \Gamma_{total}$					Γ_{75} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $K_1(1270)^0$ are included.					
<0.007	90	ANJOS	92C E691	γ Be 90–260 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.011	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K_1(1400)^0 \pi^+) / \Gamma_{total}$					Γ_{76} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $K_1(1400)^0$ are included.					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.009	90	45 ANJOS	92C E691	γ Be 90–260 GeV	
45 ANJOS 92C sees no evidence for $K_1(1400)^0 \pi^+$ in either the $\bar{K}^0 \pi^+ \pi^+ \pi^-$ or $K^- \pi^+ \pi^+ \pi^0$ channels, whereas COFFMAN 92B finds the $K_1(1400)^0 \pi^+$ branching fraction to be large; see the next entry.					

$\Gamma(K_1(1400)^0 \pi^+) / \Gamma(K^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{76} / \Gamma_{52}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $K_1(1400)^0$ are included.					
0.70 ± 0.17 OUR FIT					
0.623 ± 0.106 ± 0.180		COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K^*(1410)^0 \pi^+) / \Gamma_{total}$					Γ_{77} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $K^*(1410)^0$ are included.					
<0.007	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K^*(892)^- \pi^+ \pi^+ 3\text{-body}) / \Gamma_{total}$					Γ_{82} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $K^*(892)^0$ are included.					
0.021 ± 0.009 OUR FIT					
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.013	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K^*(892)^- \pi^+ \pi^+ 3\text{-body}) / \Gamma(K^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{82} / \Gamma_{52}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Unseen decay modes of the $K^*(892)^-$ are included.					
0.29 ± 0.13 OUR FIT				Error includes scale factor of 1.1.	
0.50 ± 0.09 ± 0.21		ANJOS	92C E691	γ Be 90–260 GeV	

$\Gamma(K^0 \rho^0 \pi^+ \text{total}) / \Gamma(K^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{85} / \Gamma_{52}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
This includes $\bar{K}^0 a_1(1260)^+$. The next two entries gives the specifically 3-body reaction.					
0.60 ± 0.10 ± 0.17	90	ANJOS	92C E691	γ Be 90–260 GeV	

$\Gamma(K^0 \rho^0 \pi^+ 3\text{-body}) / \Gamma_{total}$					Γ_{86} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.004	90	COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K^0 \rho^0 \pi^+ 3\text{-body}) / \Gamma(K^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{86} / \Gamma_{52}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.07 ± 0.04 ± 0.06		ANJOS	92C E691	γ Be 90–260 GeV	

$\Gamma(K^0 f_0(980) \pi^+) / \Gamma_{total}$					Γ_{87} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.005	90	ANJOS	92C E691	γ Be 90–260 GeV	

$\Gamma(K^0 \pi^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{57} / \Gamma_{52}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.12 ± 0.06 OUR AVERAGE					
0.10 ± 0.04 ± 0.06		ANJOS	92C E691	γ Be 90–260 GeV	
0.17 ± 0.056 ± 0.100		COFFMAN	92B MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^-) / \Gamma_{total}$					Γ_{58} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0037 ^{+0.0012} _{-0.0010}		46 BARLAG	92C ACCM	π^- Cu 230 GeV	
46 BARLAG 92C computes the branching fraction using topological normalization.					

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+)$					$\Gamma_{58} / \Gamma_{35}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.09 ± 0.01 ± 0.01	113	ANJOS	90D E691	Photoproduction	

$\Gamma(K^*(892)^0 \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$					$\Gamma_{88} / \Gamma_{58}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0 \pi^0$.					
1.25 ± 0.12 ± 0.23		ANJOS	90D E691	Photoproduction	

$\Gamma(K^*(892)^0 \rho^0 \pi^+) / \Gamma(K^*(892)^0 \pi^+ \pi^+ \pi^-)$					$\Gamma_{89} / \Gamma_{88}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.75 ± 0.17 ± 0.19		ANJOS	90D E691	Photoproduction	

$\Gamma(K^- \pi^+ \pi^+ \pi^0 \pi^0) / \Gamma_{total}$					Γ_{61} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.022 ± 0.047 ± 0.004	1	47 AGUILAR...	87F HYBR	$\pi p, pp$ 360, 400 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.015		47 BARLAG	92C ACCM	π^- Cu 230 GeV	
47 AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.					

$\Gamma(K^0 \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{total}$					Γ_{62} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.054 ± 0.030 ± 0.014					
0.099 ^{+0.036} _{-0.070}		48 BARLAG	92C ACCM	π^- Cu 230 GeV	
0.044 ^{+0.052} _{-0.013} ± 0.007	2	48 AGUILAR...	87F HYBR	$\pi p, pp$ 360, 400 GeV	
48 AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction by topological normalization.					

$\Gamma(K^0 \pi^+ \pi^+ \pi^+ \pi^-) / \Gamma_{total}$					Γ_{63} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.0008 ± 0.0007		49 BARLAG	92C ACCM	π^- Cu 230 GeV	
49 BARLAG 92C computes the branching fraction using topological normalization.					

$\Gamma(K^- \pi^+ \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{total}$					Γ_{64} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.0020 ± 0.0018		50 BARLAG	92C ACCM	π^- Cu 230 GeV	
50 BARLAG 92C computes the branching fraction using topological normalization.					

$\Gamma(K^0 \bar{K}^0 K^+) / \Gamma(K^- \pi^+ \pi^+)$					$\Gamma_{65} / \Gamma_{35}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.34 ± 0.07	70	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV	

Pionic modes

$\Gamma(\pi^+ \pi^0) / \Gamma_{total}$					Γ_{90} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.0053	90	1	BALTRUSAIT..85E	MRK3 $e^+ e^-$ 3.77 GeV	

$\Gamma(\pi^+ \pi^0) / \Gamma(K^0 \pi^+)$					$\Gamma_{90} / \Gamma_{34}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.30	90	SCHINDLER	81 MRK2	$e^+ e^-$ 3.771 GeV	

$\Gamma(\pi^+ \pi^0) / \Gamma(K^- \pi^+ \pi^+)$					$\Gamma_{90} / \Gamma_{35}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.028 ± 0.006 ± 0.005	34	SELEN	93 CLEO	$e^+ e^- \approx \Upsilon(4S)$	

$\Gamma(\pi^+ \pi^+ \pi^-) / \Gamma_{total}$					Γ_{91} / Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.0014 ^{+0.0011} _{-0.0010}		51 BARLAG	92C ACCM	π^- Cu 230 GeV	
51 BARLAG 92C computes the branching fraction using topological normalization.					

$\Gamma(\pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+ \pi^+)$					$\Gamma_{91} / \Gamma_{35}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
0.036 ± 0.006 OUR AVERAGE					
0.032 ± 0.011 ± 0.003	20	ADAMOVICH	93 WA82	π^- 340 GeV	
0.035 ± 0.007 ± 0.003		ANJOS	89 E691	Photoproduction	
0.042 ± 0.016 ± 0.010	57	BALTRUSAIT..85E	MRK3	$e^+ e^-$ 3.77 GeV	

$\Gamma(\rho^0\pi^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{92}/Γ_{35}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.015	90	ANJOS	89 E691 Photoproduction

$\Gamma(\pi^+\pi^+\pi^-\text{ nonresonant})/\Gamma(K^-\pi^+\pi^+)$		Γ_{93}/Γ_{35}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.027±0.007±0.002		ANJOS	89 E691 Photoproduction

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$		Γ_{94}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.019±0.015 -0.012		52 BARLAG	92C ACCM π^- Cu 230 GeV
52 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^-\pi^+\pi^+)$		Γ_{94}/Γ_{35}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.4	90	ANJOS	89E E691 Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$\Gamma(\eta\pi^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{100}/Γ_{35}	
Unseen decay modes of the η are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.083±0.023±0.014		99	DAOUDI 92 CLEO $e^+e^- \approx 10.5$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.12	90	ANJOS	89E E691 Photoproduction

$\Gamma(\omega\pi^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{101}/Γ_{35}	
Unseen decay modes of the ω are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.08	90	ANJOS	89E E691 Photoproduction

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$		Γ_{97}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.0010±0.0008 -0.0007		53 BARLAG	92C ACCM π^- Cu 230 GeV
53 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{97}/Γ_{35}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.019	90	ANJOS	89 E691 Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$\Gamma(\eta\rho^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{102}/Γ_{35}	
Unseen decay modes of the η are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.13	90	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV

$\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$		Γ_{98}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.0029±0.0029 -0.0020		54 BARLAG	92C ACCM π^- Cu 230 GeV
54 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(\eta(958)\pi^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{103}/Γ_{35}	
Unseen decay modes of the $\eta(958)$ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.1	90	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV
<0.1	90	ALVAREZ	91 NA14 Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			
<0.13	90	ANJOS	91B E691 $\gamma\text{Be}, \bar{E}_\gamma \approx 145$ GeV

$\Gamma(\eta(958)\rho^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{104}/Γ_{35}	
Unseen decay modes of the $\eta(958)$ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.17	90	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV

Hadronic modes with two K's

$\Gamma(K^0K^+)/\Gamma(K^0\pi^+)$		Γ_{105}/Γ_{34}	
VALUE	EVT%	DOCUMENT ID	TECN COMMENT
0.28 ± 0.06 OUR AVERAGE			
0.271 ± 0.065 ± 0.039	69	ANJOS	90C E691 γBe
0.317 ± 0.086 ± 0.048	31	BALTRUSAIT..85E	MRK3 $e^+e^- 3.77$ GeV
0.25 ± 0.15	6	SCHINDLER	81 MRK2 $e^+e^- 3.771$ GeV

$\Gamma(K^+K^-\pi^+)/\Gamma_{\text{total}}$		$\Gamma_{106}/\Gamma = (\Gamma_{109} + \frac{1}{2}\Gamma_{121} + \frac{2}{3}\Gamma_{122})/\Gamma$	
VALUE	EVT%	DOCUMENT ID	TECN COMMENT
0.0113±0.0013 OUR FIT			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.008 ± 0.017 -0.002 ± 0.001	1	55 AGUILAR...	87F HYBR $\pi p, pp$ 360, 400 GeV

55 AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization.

$\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$		Γ_{121}/Γ	
Unseen decay modes of the ϕ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.0049±0.0021 -0.0018		56 BARLAG	92C ACCM π^- Cu 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			

56 BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(\phi\pi^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{121}/Γ_{35}	
Unseen decay modes of the ϕ are included.			
VALUE	EVT%	DOCUMENT ID	TECN COMMENT
0.074±0.007 OUR FIT			
0.074±0.007 OUR AVERAGE			
0.062±0.017±0.006	19	ADAMOVICH	93 WA82 π^- 340 GeV
0.077±0.011±0.005	128	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV
0.098±0.032±0.014	12	ALVAREZ	90C NA14 Photoproduction
0.071±0.008±0.007	84	ANJOS	88 E691 Photoproduction
0.084±0.021±0.011	21	BALTRUSAIT..85E	MRK3 $e^+e^- 3.77$ GeV

$\Gamma(\bar{K}^*(892)^0K^+)/\Gamma_{\text{total}}$		Γ_{122}/Γ	
Unseen decay modes of the $\bar{K}^*(892)^0$ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.0050±0.0022 -0.0019		57 BARLAG	92C ACCM π^- Cu 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
57 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(\bar{K}^*(892)^0K^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{122}/Γ_{35}	
Includes a factor 3/2 to take into account $\bar{K}^*(892)^0 \rightarrow \bar{K}^0\pi^0$.			
VALUE	EVT%	DOCUMENT ID	TECN COMMENT
0.056±0.010 OUR FIT			
0.056±0.010 OUR AVERAGE			
0.058±0.009±0.006	73	ANJOS	88 E691 Photoproduction
0.048±0.021±0.011	14	BALTRUSAIT..85E	MRK3 $e^+e^- 3.77$ GeV

$\Gamma(K^+K^-\pi^+ \text{ nonresonant})/\Gamma_{\text{total}}$		Γ_{109}/Γ	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.0032±0.0012		58 BARLAG	92C ACCM π^- Cu 230 GeV
58 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(K^+K^-\pi^+ \text{ nonresonant})/\Gamma(K^-\pi^+\pi^+)$		Γ_{109}/Γ_{35}	
VALUE	EVT%	DOCUMENT ID	TECN COMMENT
0.050±0.009 OUR FIT			
0.050±0.009 OUR AVERAGE			
0.049±0.008±0.006	95	ANJOS	88 E691 Photoproduction
0.059±0.026±0.009	37	BALTRUSAIT..85E	MRK3 $e^+e^- 3.77$ GeV

$\Gamma(\phi\pi^+\pi^0)/\Gamma_{\text{total}}$		Γ_{123}/Γ	
Unseen decay modes of the ϕ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.023±0.010		59 BARLAG	92C ACCM π^- Cu 230 GeV
59 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(\phi\pi^+\pi^0)/\Gamma(K^-\pi^+\pi^+)$		Γ_{123}/Γ_{35}	
Unseen decay modes of the ϕ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.58	90	ALVAREZ	90C NA14 Photoproduction
<0.28	90	ANJOS	89E E691 Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$\Gamma(\phi\rho^+)/\Gamma(K^-\pi^+\pi^+)$		Γ_{124}/Γ_{35}	
Unseen decay modes of the ϕ are included.			
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.16	90	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV

$\Gamma(K^+K^-\pi^+\pi^0 \text{ non-}\phi)/\Gamma_{\text{total}}$		Γ_{113}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.015 ± 0.007 -0.006		60 BARLAG	92C ACCM π^- Cu 230 GeV
60 BARLAG 92C computes the branching fraction using topological normalization.			

$\Gamma(K^+K^-\pi^+\pi^0 \text{ non-}\phi)/\Gamma(K^-\pi^+\pi^+)$		Γ_{113}/Γ_{35}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.25	90	ANJOS	89E E691 Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			

$\Gamma(K^+K^0\pi^+\pi^-)/\Gamma_{\text{total}}$		Γ_{114}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.02	90	ALBRECHT	92B ARG $e^+e^- \approx 10.4$ GeV

Meson Full Listings

 D^\pm $\Gamma(K^0 K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
$0.01 \pm 0.005 \pm 0.003$	ALBRECHT 92B ARG		$e^+e^- \approx 10.4$ GeV
<0.003	61 BARLAG 92C ACCM		π^- Cu 230 GeV
	61 BARLAG 92C		computes the branching fraction using topological normalization.

 $\Gamma(K^*(892)^+ \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_{125}/Γ

Unseen decay modes of the \bar{K}^* 's are included.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.026 \pm 0.008 \pm 0.007$	ALBRECHT 92B ARG		$e^+e^- \approx 10.4$ GeV

 $\Gamma(K^0 K^- \pi^+ \pi^+ \text{non-}K^* \bar{K}^{*0})/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0079	90	ALBRECHT 92B ARG		$e^+e^- \approx 10.4$ GeV

 $\Gamma(\phi \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{126}/Γ

Unseen decay modes of the ϕ are included.

VALUE	CL%	EPTS	DOCUMENT ID	TECN	COMMENT
<0.002	90	0	ANJOS 88 E691		Photoproduction

 $\Gamma(\phi \pi^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{126}/Γ_{35}

Unseen decay modes of the ϕ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.031	90	ALVAREZ 90C NA14		Photoproduction

 $\Gamma(\phi \pi^+ \pi^+ \pi^-)/\Gamma(\phi \pi^+)$ $\Gamma_{126}/\Gamma_{121}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.6	90	FRABETTI 92 E687		γ Be

 $\Gamma(K^+ K^- \pi^+ \pi^+ \pi^- \text{nonresonant})/\Gamma_{\text{total}}$ Γ_{120}/Γ

VALUE	CL%	EPTS	DOCUMENT ID	TECN	COMMENT
<0.03	90	12	ANJOS 88 E691		Photoproduction

Rare or forbidden modes

 $\Gamma(K^+ \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{127}/Γ_{35}

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.05	90	62 PICCOLO 77 MRK1		$e^+e^- 4.03$ GeV
		62 PICCOLO 77		Obtained from $\sigma \times \text{BR}$ values of Table I of PICCOLO 77.

 $\Gamma(K^+ K^+ K^-)/\Gamma(K^- \pi^+ \pi^+)$ Γ_{128}/Γ_{35}

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

VALUE	EPTS	DOCUMENT ID	TECN	COMMENT
$0.057 \pm 0.020 \pm 0.007$	13	ADAMOVICH 93 WA82		π^- 340 GeV

 $\Gamma(\phi K^+)/\Gamma(\phi \pi^+)$ $\Gamma_{129}/\Gamma_{121}$

A doubly Cabibbo-suppressed decay with no simple spectator process possible.

VALUE	EPTS	DOCUMENT ID	TECN	COMMENT
$0.058 \pm 0.032 \pm 0.007$	4	63 ANJOS 92D E691		γ Be $\bar{E}_\gamma = 145$ GeV
		63 ANJOS 92D		The evidence of ANJOS 92D is a small excess of events ($4.5^{+2.4}_{-2.0}$).

 $\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{130}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EPTS	DOCUMENT ID	TECN	COMMENT
$<2.5 \times 10^{-3}$	90		WEIR 90B MRK2		$e^+e^- 29$ GeV
$<2.6 \times 10^{-3}$	90	39	64 HAAS 88 CLEO		$e^+e^- 10$ GeV
			64 HAAS 88		The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88c.

 $\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{131}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EPTS	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-3}$	90	36	65 HAAS 88 CLEO		$e^+e^- 10$ GeV
$<5.9 \times 10^{-3}$	90		WEIR 90B MRK2		$e^+e^- 29$ GeV
			65 HAAS 88		The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88c.

 $\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{132}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.8 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.2 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(\pi^+ e^\pm \mu^\mp)/\Gamma_{\text{total}}$ Γ_{134}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	EPTS	DOCUMENT ID	TECN	COMMENT
$<3.8 \times 10^{-3}$	90	58	66 HAAS 88 CLEO		$e^+e^- 10$ GeV
			66 HAAS 88		The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88c.

 $\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{135}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{136}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.3 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{137}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{138}/Γ

A test of lepton-family-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.4 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(\pi^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{139}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.8 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{140}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.8 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(\pi^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{141}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.7 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(K^- e^+ e^+)/\Gamma_{\text{total}}$ Γ_{142}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<9.1 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(K^- \mu^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{143}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.3 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 $\Gamma(K^- e^+ \mu^+)/\Gamma_{\text{total}}$ Γ_{144}/Γ

A test of lepton-number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-3}$	90	WEIR 90B MRK2		$e^+e^- 29$ GeV

 D^\pm PRODUCTION CROSS SECTION AT $\psi(3770)$

A compilation of the cross sections for the direct production of D^\pm mesons at or near the $\psi(3770)$ peak in e^+e^- production.

VALUE (nanobarns)	DOCUMENT ID	TECN	COMMENT
$4.2 \pm 0.6 \pm 0.3$	67 ADLER 88C MRK3		$e^+e^- 3.768$ GeV
5.5 ± 1.0	68 PARTRIDGE 84 CBAL		$e^+e^- 3.771$ GeV
$6.00 \pm 0.72 \pm 1.02$	69 SCHINDLER 80 MRK2		$e^+e^- 3.771$ GeV
9.1 ± 2.0	70 PERUZZI 77 MRK1		$e^+e^- 3.774$ GeV

67 This measurement compares events with one detected D to those with two detected D mesons, to determine the absolute cross section. ADLER 88C measure the ratio of cross sections (neutral to charged) to be $1.36 \pm 0.23 \pm 0.14$. This measurement does not include the decays of the $\psi(3770)$ not associated with charmed particle production.

68 This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. PARTRIDGE 84 measures 6.4 ± 1.15 nb for the cross section. We take the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and we assume that the $\psi(3770)$ is an isosinglet to evaluate the cross sections. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

69 This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. SCHINDLER 80 assume the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and that the $\psi(3770)$ is an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

70 This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. The phase space division of neutral and charged D mesons in $\psi(3770)$ decay is taken to be 1.33, and $\psi(3770)$ is assumed to be an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction. We exclude this measurement from the average because of uncertainties in the contamination from τ lepton pairs. Also see RAPIDIS 77.

See key on page 1343

Meson Full Listings

 D^\pm, D^0 D^\pm REFERENCES

BALEST	94	PRL 72 2328	+Cho, Daoudi, Ford+	(CLEO Collab.)
FRABETTI	94D	PL B323 459	+Cheung, Cumalat+	(FNAL E687 Collab.)
PDG	94	PR D50 1173	Montanet+	(CERN, LBL, BOST, IFIC+)
ABE	93E	PL B313 288	+Amako, Arai, Arima, Asano+	(VENUS Collab.)
ADAMOVICH	93	PL B305 177	+Alexandrov, Antinori+	(CERN WAB2 Collab.)
AKERIB	93	PRL 71 3070	+Barish, Chadha, Chan+	(CLEO Collab.)
ALAM	93	PRL 71 1311	+Kim, Nemati, O'Neill+	(CLEO Collab.)
ANJOS	93	PR D48 55	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BEAN	93C	PL B317 647	+Gronberg, Kutschke, Menary+	(CLEO Collab.)
FRABETTI	93E	PL B307 262	+Grim, Paolone, Yager+	(FNAL E687 Collab.)
KODAMA	93B	PL B313 260	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
KODAMA	93C	PL B316 455	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
SELEN	93	PRL 71 1973	+Sadoff, Ammar, Ball+	(CLEO Collab.)
ALBRECHT	92B	ZPHY C53 361	+Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALBRECHT	92F	PL B278 202	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
ANJOS	92	PR D45 R2177	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	92C	PR D46 1941	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	92D	PRL 69 2892	+Appel, Bean, Bediaga+	(FNAL E691 Collab.)
BARLAG	92C	ZPHY C55 383	+Becker, Bozok, Boehringer+	(ACCMOR Collab.)
Also	90D	ZPHY C48 29	+Barlag, Becker, Boehringer, Bosman+	(ACCMOR Collab.)
COFFMAN	92B	PR D45 2196	+DeJongh, Dubois, Eigen+	(Mark III Collab.)
DAUDI	92	PR D45 3965	+Ford, Johnson, Lingel+	(CLEO Collab.)
FRABETTI	92	PL B281 167	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
KODAMA	91C	PRL 67 1507	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
KODAMA	92C	PL B286 187	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
ADAMOVICH	91	PL B268 142	+Alexandrov, Antinori, Barberis+	(WAB2 Collab.)
ALBRECHT	91	PL B255 634	+Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALVAREZ	91	PL B255 639	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ALVAREZ	91B	ZPHY C50 11	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
AMMAR	91	PR D44 3383	+Baringer, Coppage, Davis+	(CLEO Collab.)
ANJOS	91B	PR D43 R2063	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	91C	PRL 67 1507	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BAI	91	PRL 66 1011	+Bolton, Brown, Bunnell+	(Mark III Collab.)
COFFMAN	91	PL B263 135	+DeJongh, Dubois, Eigen, Hitlin+	(Mark III Collab.)
FRABETTI	91	PL B263 584	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
ALVAREZ	90	ZPHY C47 539	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ALVAREZ	90C	PL B246 261	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ANJOS	90C	PR D41 2705	+Appel, Bean+	(FNAL E691 Collab.)
ANJOS	90D	PR D42 2414	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	90E	PR 65 2630	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BARLAG	90C	ZPHY C46 563	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
WEIR	90B	PR D41 1384	+Klein, Abrams, Adolphsen, Akerlof+	(Mark II Collab.)
ANJOS	89	PRL 62 125	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	89B	PRL 62 722	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS	89E	PL B223 267	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ADLER	88B	PRL 60 1375	+Becker, Blaylock+	(Mark III Collab.)
ADLER	88C	PRL 60 89	+Becker, Blaylock+	(Mark III Collab.)
ALBRECHT	88F	PL B210 267	+Boeckmann, Glaeser+	(ARGUS Collab.)
ANJOS	88	PRL 60 897	+Appel+	(FNAL E691 Collab.)
AOKI	88	PL B209 113	+Arnold, Baroni+	(WA75 Collab.)
HAAS	88	PRL 60 1614	+Hempstead, Jensen+	(CLEO Collab.)
ONG	88	PRL 60 2587	+Weir, Abrams, Amidel+	(Mark II Collab.)
RAAB	88	PR D37 2391	+Anjos, Appel, Bracker+	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	+Alexandrov, Bolta+	(Photon Emulsion Collab.)
ADLER	87	PL B196 107	+Becker, Blaylock, Bolton+	(Mark III Collab.)
AGUILAR...	87D	PL B193 140	+Becker, Felst, Haidt+	(JADE Collab.)
Also	88B	ZPHY C40 321	+Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
AGUILAR...	87E	ZPHY C36 551	+Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also	88B	ZPHY C40 321	+Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
AGUILAR...	87F	ZPHY C36 559	+Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also	88	ZPHY C38 520 erratum		
BARLAG	87B	ZPHY C37 17	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
BARTLE	87	ZPHY C33 339	+Becker, Felst, Haidt+	(JADE Collab.)
CSORNA	87	PL B191 319	+Mestayer, Panvini, Word+	(CLEO Collab.)
PALKA	87B	ZPHY C35 151	+Bailey, Becker+	(ACCMOR Collab.)
ABE	86	PR D33 1	+ (SLAC Hybrid Facility Photon Collab.)	
AGUILAR...	86B	ZPHY C31 491	+Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
BALTRUSAIT...	86E	PRL 56 2140	+Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
PAL	86	PR D33 2708	+Atwood, Barish, Bonneaud+	(DELCO Collab.)
AIHARA	85	ZPHY C27 39	+Alston-Garnjost, Badtke, Bakken+	(TPC Collab.)
BALTRUSAIT...	85B	PRL 54 1976	+Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BALTRUSAIT...	85E	PRL 55 150	+Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BARTEL	85J	PL 163B 277	+Becker, Cords, Felst+	(JADE Collab.)
ADAMOVICH	84	PL 140B 119	+Alexandrov, Bolta, Bravo+	(CERN WAB2 Collab.)
ALTHOFF	84G	ZPHY C22 219	+Braunschweig, Kirschfink+	(TASSO Collab.)
ALTHOFF	84J	PL 146B 443	+Braunschweig, Kirschfink+	(TASSO Collab.)
DERRICK	84	PRL 53 1971	+Fernandez, Fries, Hyman+	(HRS Collab.)
KOOP	84	PRL 52 970	+Sakuda, Atwood, Bailion+	(DELCO Collab.)
PARTRIDGE	84	Thesis CALT-68-1150		
AGUILAR...	83B	PL 123B 98	+Aguilar-Benitez, Allison+	(Crystal Ball Collab.)
AUBERT	83	NP B213 31	+Bassompierre, Beck, Best+	(EMC Collab.)
PARTRIDGE	81	PRL 47 760	+Peck, Porter, Gu+	(Crystal Ball Collab.)
SCHINDLER	81	PR D24 78	+Alam, Boyarski, Breidenbach+	(Mark II Collab.)
TRILLING	81	PRPL 75 57		(LBL, UCSB) J
BACINO	80	PRL 45 329	+Ferguson+	(UCLA, SLAC, STAN, UCI, STON)
SCHINDLER	80	PR D21 2716	+Siegrist, Alam, Boyarski+	(Mark II Collab.)
ZHOLENTZ	80	PL 96B 214	+Kurdadze, Leichuk, Mishnev+	(NOVO)
Also	81	SJNP 34 814	+Zholentz, Kurdadze, Leichuk+	(NOVO)
Translated from YAF 34 1471.				
BACINO	79	PRL 43 1073	+Ferguson, Nodulman+	(DELCO Collab.)
BRANDELIK	79	PL 80B 412	+Braunschweig, Martyn, Sander+	(DASP Collab.)
FELLER	78	PRL 40 274	+Litke, Madaras, Ronan+	(LBL, SLAC, NWES, HAWA)
VUILLEMIN	78	PRL 41 1149	+Feldman, Feller+	(LBL, SLAC, NWES, HAWA)
GOLDHABER	77	PL 69B 303	+Wiss, Abrams, Alam+	(LBL, SLAC)
PERUZZI	77	PRL 39 1301	+Piccolo, Felds, HAWA	(SLAC, LBL, FELDS, HAWA)
PICCOLO	77	PL 70B 260	+Peruzzi, Luth, Nguyen, Wiss, Abrams+	(SLAC, LBL)
RAPIDIS	77	PRL 39 526	+Gobbi, Luke, Barbaro-Galtieri+	(Mark I Collab.)
PERUZZI	76	PRL 37 569	+Piccolo, Feldman, Nguyen, Wiss+	(SLAC, LBL)

OTHER RELATED PAPERS

MORRISON	89	ARNPS 39 183	+Witherell	(UCSB)
----------	----	--------------	------------	--------

 D^0

$$I(J^P) = \frac{1}{2}(0^-)$$

 D^0 MASSThe fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}$, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1864.6 ± 0.5 OUR FIT				
1864.1 ± 1.0 OUR AVERAGE				
1864.6 ± 0.3 ± 1.0	641	BARLAG	90C ACCM	π^- Cu 230 GeV
1852 ± 7	16	ADAMOVICH	87 EMUL	Photoproduction
1861 ± 4		DERRICK	84 HRS	e^+e^- 29 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1856 ± 36	22	ADAMOVICH	84B EMUL	Photoproduction
1847 ± 7	1	FIORINO	81 EMUL	$\gamma N \rightarrow \bar{D}^0 +$
1863.8 ± 0.5		¹ SCHINDLER	81 MRK2	e^+e^- 3.77 GeV
1864.7 ± 0.6		¹ TRILLING	81 RVUE	e^+e^- 3.77 GeV
1863.0 ± 2.5	238	ASTON	80E OMEG	$\gamma p \rightarrow \bar{D}^0$
1860 ± 2	143	² AVERY	80 SPEC	$\gamma N \rightarrow D^{*+}$
1869 ± 4	35	² AVERY	80 SPEC	$\gamma N \rightarrow D^{*+}$
1854 ± 6	94	² ATIYA	79 SPEC	$\gamma N \rightarrow D^0 \bar{D}^0$
1850 ± 15	64	BALTAY	78C HBC	$\nu N \rightarrow K^0 \pi \pi$
1863 ± 3		GOLDHABER	77 MRK1	D^0, D^+ recoil spectra
1863.3 ± 0.9		¹ PERUZZI	77 MRK1	e^+e^- 3.77 GeV
1868 ± 11		PICCOLO	77 MRK1	e^+e^- 4.03, 4.41 GeV
1865 ± 15	234	GOLDHABER	76 MRK1	$K \pi$ and $K^3 \pi$

¹ PERUZZI 77 and SCHINDLER 81 errors do not include the 0.13% uncertainty in the absolute SPEAR energy calibration. TRILLING 81 uses the high precision $J/\psi(1S)$ and $\psi(2S)$ measurements of ZHOLENTZ 80 to determine this uncertainty and combines the PERUZZI 77 and SCHINDLER 81 results to obtain the value quoted. TRILLING 81 enters the fit in the D^\pm mass, and PERUZZI 77 and SCHINDLER 81 enter in the $m_{D^\pm} - m_{D^0}$ below.

² Error does not include possible systematic mass scale shift, estimated to be less than 5 MeV.

$$|m_{D_1^0} - m_{D_2^0}|$$

The D_1^0 and D_2^0 are the mass eigenstates of the D^0 meson.

VALUE ($10^{10} \hbar s^{-1}$)	CL%	DOCUMENT ID	TECN	COMMENT
< 20	90	^{3,4} ANJOS	88C E691	Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 40	90	³ ALBRECHT	87K ARG	e^+e^- 10 GeV
< 24	90	⁵ LOUIS	86 SPEC	$\pi^- W$ 225 GeV
< 106	90	^{3,6} YAMAMOTO	85 DLCO	e^+e^- 29 GeV
< 99	90	⁵ BODEK	82 SPEC	$\pi^-, pFe \rightarrow D^0$

³ Limit inferred from the $D^0-\bar{D}^0$ mixing ratio $\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+)$ near the end of the D^0 Listings.

⁴ Calculated by us using $\Delta(m) = (2r/(1-r))^{1/2} \hbar/4.21 \times 10^{-13} s$, where r is the $D^0-\bar{D}^0$ mixing ratio. See the data on $r \equiv \Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+)$ near the end of the D^0 Listings.

⁵ Limit inferred from the $D^0-\bar{D}^0$ mixing ratio $\Gamma(\mu^- \text{ anything (via } \bar{D}^0))/\Gamma(\mu^+ \text{ anything})$ near the end of the D^0 Listings.

⁶ YAMAMOTO 85 gives $\Delta(m)/\Gamma < 0.44$. We use $\Gamma = \hbar/4.3 \times 10^{-13} s$.

$$m_{D^\pm} - m_{D^0}$$

The fit includes $D^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}$, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4.78 ± 0.10 OUR FIT			
4.74 ± 0.28 OUR AVERAGE			
4.7 ± 0.3	⁷ SCHINDLER	81 MRK2	e^+e^- 3.77 GeV
5.0 ± 0.8	⁷ PERUZZI	77 MRK1	e^+e^- 3.77 GeV

⁷ See the footnote on TRILLING 81 in the D^0 and D^\pm sections on the mass.

Meson Full Listings

 D^0 D^0 MEAN LIFE

Measurements with an error $> 0.05 \times 10^{-12}$ s are omitted from the average, and those with an error $> 0.1 \times 10^{-12}$ s or that have been superseded by later results have been removed from the Listings.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.415 ± 0.004 OUR AVERAGE				
0.413 ± 0.004 ± 0.003	16k	FRABETTI	94D E687	$K^- \pi^+, K^- \pi^+ \pi^+ \pi^-$
0.424 ± 0.011 ± 0.007	5118	FRABETTI	91 E687	$K^- \pi^+, K^- \pi^+ \pi^+ \pi^-$
0.417 ± 0.018 ± 0.015	890	ALVAREZ	90 NA14	$K^- \pi^+, K^- \pi^+ \pi^+ \pi^-$
0.388 ^{+0.023} _{-0.021}	641	⁸ BARLAG	90c ACCM	π^- Cu 230 GeV
0.48 ± 0.04 ± 0.03	776	ALBRECHT	88i ARG	$e^+ e^-$ 10 GeV
0.422 ± 0.008 ± 0.010	4212	RAAB	88 E691	Photoproduction
0.42 ± 0.05	90	BARLAG	87b ACCM	K^- and π^- 200 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.34 ^{+0.06} _{-0.05} ± 0.03	58	AMENDOLIA	88 SPEC	Photoproduction
0.46 ^{+0.06} _{-0.05}	145	AGUILAR...	87D HYBR	$\pi^- p$ and pp
0.50 ± 0.07 ± 0.04	317	CSORNA	87 CLEO	$e^+ e^-$ 10 GeV
0.61 ± 0.09 ± 0.03	50	ABE	86 HYBR	γp 20 GeV
0.47 ^{+0.09} _{-0.08} ± 0.05	74	GLADNEY	86 MRK2	$e^+ e^-$ 29 GeV
0.43 ^{+0.07} _{-0.05} ^{+0.01} _{-0.02}	58	USHIDA	86b EMUL	ν wideband
0.37 ^{+0.10} _{-0.07}	26	BAILEY	85 SILI	π^- Be 200 GeV

⁸ BARLAG 90c estimate systematic error to be negligible.

 $|\tau_{D_1^0} - \tau_{D_2^0}|/\tau_{D^0}$, MEAN LIFE DIFFERENCE/AVERAGE

The D_1^0 and D_2^0 are the mass eigenstates of the D^0 meson.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.17		90 9.10 ANJOS	88c E691	Photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.21	90	¹¹ LOUIS	86 SPEC	π^- W 225 GeV
<0.8	90	⁹ YAMAMOTO	85 DLCO	$e^+ e^-$ 29 GeV
<0.55	90	¹¹ BODEK	82 SPEC	π^- , p Fe $\rightarrow D^0$
⁹ This limit is inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+)$ near the end of the D^0 Listings.				
¹⁰ Calculated by us using $\Delta\Gamma = (8r/(1+r))^2/2\hbar/4.21 \times 10^{-13}$ s, where r is the D^0 - \bar{D}^0 mixing ratio. See the data on $r \equiv \Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+)$ near the end of the D^0 Listings.				
¹¹ Limit inferred from the D^0 - \bar{D}^0 mixing ratio $\Gamma(\mu^- \text{ anything (via } \bar{D}^0))/\Gamma(\mu^+ \text{ anything})$ near the end of the D^0 Listings.				

 D^0 DECAY MODES

\bar{D}^0 modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Inclusive modes		
Γ_1 e^+ anything	(7.7 ± 1.2) %	S=1.1
Γ_2 μ^+ anything	(10.0 ± 2.6) %	
Γ_3 K^- anything	(53 ± 4) %	S=1.3
Γ_4 \bar{K}^0 anything + K^0 anything	(42 ± 5) %	
Γ_5 K^+ anything	(3.4 ^{+0.6} _{-0.4}) %	
Γ_6 η anything	[a] < 13 %	CL=90%
Semileptonic modes		
Γ_7 $K^- e^+ \nu_e$	[b] (3.68 ± 0.21) %	S=1.1
Γ_8 $K^- e^+ \nu_e$	(3.80 ± 0.22) %	S=1.1
Γ_9 $K^- \mu^+ \nu_\mu$	(3.2 ± 0.4) %	
Γ_{10} $K^- \pi^0 e^+ \nu_e$	[c] (1.6 ^{+1.3} _{-0.5}) %	
Γ_{11} $\bar{K}^0 \pi^- e^+ \nu_e$	[c] (2.8 ^{+1.7} _{-0.9}) %	
Γ_{12} $\bar{K}^*(892)^- e^+ \nu_e$ $\times B(K^{*-} \rightarrow \bar{K}^0 \pi^+)$	(1.3 ± 0.3) %	
Γ_{13} $K^*(892)^- \ell^+ \nu_\ell$	[d]	
Γ_{14} $K^- \pi^0(\pi^0) e^+ \nu_e$		
Γ_{15} $\bar{K}^0 \pi^- (\pi^0) e^+ \nu_e$		
Γ_{16} $\bar{K}^*(892)^0 \pi^- e^+ \nu_e$	[e] < 1.3 %	CL=90%
Γ_{17} $K^- \pi^+ \pi^- \mu^+ \nu_\mu$	< 1.2 $\times 10^{-3}$	CL=90%
Γ_{18} $(\bar{K}^*(892) \pi)^- \mu^+ \nu_\mu$	< 1.4 $\times 10^{-3}$	CL=90%
Γ_{19} $\pi^- e^+ \nu_e$	(3.9 ^{+2.3} _{-1.2}) $\times 10^{-3}$	
A fraction of the following resonance mode has already appeared above as a submode of a particular charged-particle mode.		
Γ_{20} $K^*(892)^- e^+ \nu_e$	(2.0 ± 0.4) %	

Hadronic modes with one or three K 's

Γ_{21} $K^- \pi^+$	(4.01 ± 0.14) %	
Γ_{22} $\bar{K}^0 \pi^0$	(2.05 ± 0.26) %	S=1.1
Γ_{23} $\bar{K}^0 \pi^+ \pi^-$	[f] (5.3 ± 0.6) %	S=1.2
Γ_{24} $\bar{K}^0 \rho^0$	(1.10 ± 0.18) %	
Γ_{25} $\bar{K}^0 f_0(980)$ $\times B(f_0 \rightarrow \pi^+ \pi^-)$	(2.4 ± 1.0) $\times 10^{-3}$	
Γ_{26} $\bar{K}^0 f_2(1270)$ $\times B(f_2 \rightarrow \pi^+ \pi^-)$	(2.6 ± 1.2) $\times 10^{-3}$	
Γ_{27} $\bar{K}^0 f_0(1300)$ $\times B(f_0 \rightarrow \pi^+ \pi^-)$	(4.3 ± 1.7) $\times 10^{-3}$	
Γ_{28} $K^*(892)^- \pi^+$ $\times B(K^{*-} \rightarrow \bar{K}^0 \pi^-)$	(3.3 ± 0.4) %	
Γ_{29} $K_0^*(1430)^- \pi^+$ $\times B(K_0^*(1430)^- \rightarrow \bar{K}^0 \pi^-)$	(7 ± 3) $\times 10^{-3}$	
Γ_{30} $\bar{K}^0 \pi^+ \pi^-$ nonresonant	(1.43 ± 0.26) %	
Γ_{31} $K^- \pi^+ \pi^0$	[f] (13.8 ± 1.0) %	S=1.1
Γ_{32} $K^- \rho^+$	(10.4 ± 1.3) %	
Γ_{33} $K^*(892)^- \pi^+$ $\times B(K^{*-} \rightarrow K^- \pi^0)$	(1.6 ± 0.2) %	
Γ_{34} $\bar{K}^*(892)^0 \pi^0$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	(2.0 ± 0.3) %	
Γ_{35} $K^- \pi^+ \pi^0$ nonresonant	(6.0 ± 2.7) $\times 10^{-3}$	
Γ_{36} $\bar{K}^0 \pi^0 \pi^0$		
Γ_{37} $\bar{K}^*(892)^0 \pi^0$ $\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$	(1.0 ± 0.2) %	
Γ_{38} $\bar{K}^0 \pi^0 \pi^0$ nonresonant	(7.6 ± 2.1) $\times 10^{-3}$	
Γ_{39} $K^- \pi^+ \pi^+ \pi^-$	[f] (8.1 ± 0.5) %	
Γ_{40} $K^- \pi^+ \rho^0$ total	(6.8 ± 0.5) %	
Γ_{41} $K^- \pi^+ \rho^0$ 3-body	(5.1 ± 2.3) $\times 10^{-3}$	
Γ_{42} $\bar{K}^*(892)^0 \rho^0$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	(1.1 ± 0.3) %	
Γ_{43} $K^- a_1(1260)^+$ $\times B(a_1(1260)^+ \rightarrow \pi^+ \pi^+ \pi^-)$	(3.9 ± 0.6) %	
Γ_{44} $\bar{K}^*(892)^0 \pi^+ \pi^-$ total $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	(1.6 ± 0.4) %	
Γ_{45} $\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	(1.01 ± 0.22) %	
Γ_{46} $K_1(1270)^- \pi^+$ $\times B(K_1(1270)^- \rightarrow K^- \pi^+ \pi^-)$	(3.5 ± 1.1) $\times 10^{-3}$	
Γ_{47} $K^- \pi^+ \pi^+ \pi^-$ nonresonant	(1.89 ± 0.28) %	
Γ_{48} $\bar{K}^0 \pi^+ \pi^- \pi^0$	[f] (9.8 ± 1.4) %	S=1.1
Γ_{49} $\bar{K}^0 \eta \times B(\eta \rightarrow \pi^+ \pi^- \pi^0)$	(1.61 ± 0.26) $\times 10^{-3}$	
Γ_{50} $\bar{K}^0 \omega \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	(1.8 ± 0.4) %	
Γ_{51} $K^*(892)^- \rho^+$ $\times B(K^{*-} \rightarrow \bar{K}^0 \pi^-)$	(3.9 ± 1.6) %	
Γ_{52} $\bar{K}^*(892)^0 \rho^0$ $\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$	(5.3 ± 1.4) $\times 10^{-3}$	
Γ_{53} $K_1(1270)^- \pi^+$ $\times B(K_1(1270)^- \rightarrow \bar{K}^0 \pi^- \pi^0)$	[g] (5.0 ± 1.5) $\times 10^{-3}$	
Γ_{54} $\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body $\times B(\bar{K}^{*0} \rightarrow \bar{K}^0 \pi^0)$	(5.1 ± 1.1) $\times 10^{-3}$	
Γ_{55} $\bar{K}^0 \pi^+ \pi^- \pi^0$ nonresonant	(2.1 ± 2.1) %	
Γ_{56} $K^- \pi^+ \pi^0 \pi^0$	(15 ± 5) %	
Γ_{57} $K^- \pi^+ \pi^+ \pi^- \pi^0$	(4.3 ± 0.4) %	
Γ_{58} $\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$	(1.3 ± 0.6) %	
Γ_{59} $\bar{K}^*(892)^0 \eta$ $\times B(\bar{K}^{*0} \rightarrow \pi^+ \pi^- \pi^0)$ $\times B(\eta \rightarrow \pi^+ \pi^- \pi^0)$	(3.0 ± 0.8) $\times 10^{-3}$	
Γ_{60} $K^- \pi^+ \omega \times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	(2.8 ± 0.5) %	
Γ_{61} $\bar{K}^*(892)^0 \omega$ $\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$ $\times B(\omega \rightarrow \pi^+ \pi^- \pi^0)$	(7 ± 3) $\times 10^{-3}$	
Γ_{62} $\bar{K}^0 \pi^+ \pi^+ \pi^- \pi^-$	(5.6 ± 1.7) $\times 10^{-3}$	
Γ_{63} $\bar{K}^0 \pi^+ \pi^- \pi^0 \pi^0 (\pi^0)$	(10.6 ^{+7.3} _{-3.0}) %	
Γ_{64} $\bar{K}^0 K^+ K^-$	(9.1 ± 1.2) $\times 10^{-3}$	
In the fit as $\frac{1}{2}\Gamma_{75} + \Gamma_{66}$, where $\frac{1}{2}\Gamma_{75} = \Gamma_{65}$.		
Γ_{65} $\bar{K}^0 \phi \times B(\phi \rightarrow K^+ K^-)$	(4.2 ± 0.6) $\times 10^{-3}$	
Γ_{66} $\bar{K}^0 K^+ K^-$ non- ϕ	(4.9 ± 0.9) $\times 10^{-3}$	
Γ_{67} $K_S^0 K_S^0 K_S^0$	(8.6 ± 2.5) $\times 10^{-4}$	
Γ_{68} $K^+ K^- \bar{K}^0 \pi^0$	(7.2 ^{+4.8} _{-3.5}) $\times 10^{-3}$	

Fractions of many of the following modes with resonances have already appeared above as submodes of particular charged-particle modes. (Modes for which there are only upper limits and $\bar{K}^*(892)\rho$ submodes only appear below.)

Γ_{69}	$\bar{K}^0 \eta$	$(6.8 \pm 1.1) \times 10^{-3}$	
Γ_{70}	$\bar{K}^0 \rho^0$	$(1.10 \pm 0.18) \%$	
Γ_{71}	$K^- \rho^+$	$(10.4 \pm 1.3) \%$	S=1.2
Γ_{72}	$\bar{K}^0 \omega$	$(2.0 \pm 0.4) \%$	
Γ_{73}	$\bar{K}^0 \eta'(958)$	$(1.66 \pm 0.29) \%$	
Γ_{74}	$\bar{K}^0 f_0(980)$	$(4.6 \pm 2.0) \times 10^{-3}$	
Γ_{75}	$\bar{K}^0 \phi$	$(8.3 \pm 1.2) \times 10^{-3}$	S=1.1
Γ_{76}	$K^- a_1(1260)^+$	$(7.9 \pm 1.2) \%$	
Γ_{77}	$\bar{K}^0 a_1(1260)^0$	< 1.9	CL=90%
Γ_{78}	$\bar{K}^0 f_2(1270)$	$(4.6 \pm 2.1) \times 10^{-3}$	
Γ_{79}	$\bar{K}^0 f_0(1300)$	$(6.9 \pm 2.7) \times 10^{-3}$	
Γ_{80}	$K^- a_2(1320)^+$	< 2	CL=90%
Γ_{81}	$K^*(892)^- \pi^+$	$(4.9 \pm 0.6) \%$	S=1.3
Γ_{82}	$\bar{K}^*(892)^0 \pi^0$	$(3.0 \pm 0.4) \%$	
Γ_{83}	$\bar{K}^*(892)^0 \pi^+ \pi^-$ total	$(2.4 \pm 0.6) \%$	
Γ_{84}	$\bar{K}^*(892)^0 \pi^+ \pi^-$ 3-body	$(1.52 \pm 0.33) \%$	
Γ_{85}	$K^- \pi^+ \rho^0$ total	$(6.8 \pm 0.5) \%$	
Γ_{86}	$K^- \pi^+ \rho^0$ 3-body	$(5.1 \pm 2.3) \times 10^{-3}$	
Γ_{87}	$\bar{K}^*(892)^0 \rho^0$	$(1.6 \pm 0.4) \%$	
Γ_{88}	$\bar{K}^*(892)^0 \rho^0$ transverse	$(1.6 \pm 0.5) \%$	
Γ_{89}	$\bar{K}^*(892)^0 \rho^0$ S-wave	$(3.0 \pm 0.6) \%$	
Γ_{90}	$\bar{K}^*(892)^0 \rho^0$ S-wave long.	< 3	CL=90%
Γ_{91}	$\bar{K}^*(892)^0 \rho^0$ P-wave	< 3	CL=90%
Γ_{92}	$\bar{K}^*(892)^0 \rho^0$ D-wave	$(2.1 \pm 0.6) \%$	
Γ_{93}	$K^*(892)^- \rho^+$	$(5.9 \pm 2.4) \%$	
Γ_{94}	$K^*(892)^- \rho^+$ longitudinal	$(2.8 \pm 1.2) \%$	
Γ_{95}	$K^*(892)^- \rho^+$ transverse	$(3.1 \pm 1.8) \%$	
Γ_{96}	$K^*(892)^- \rho^+$ P-wave	< 1.5	CL=90%
Γ_{97}	$K^- \pi^+ f_0(980)$	< 1.1	CL=90%
Γ_{98}	$\bar{K}^*(892)^0 f_0(980)$	< 7	CL=90%
Γ_{99}	$K_1(1270)^- \pi^+$	$[g] (1.04 \pm 0.31) \%$	
Γ_{100}	$K_1(1400)^- \pi^+$	< 1.2	CL=90%
Γ_{101}	$\bar{K}_1(1400)^0 \pi^0$	< 3.7	CL=90%
Γ_{102}	$K^*(1410)^- \pi^+$	< 1.2	CL=90%
Γ_{103}	$K_2^*(1430)^- \pi^+$	$(1.1 \pm 0.4) \%$	
Γ_{104}	$K_2^*(1430)^- \pi^+$	< 8	CL=90%
Γ_{105}	$\bar{K}_2^*(1430)^0 \pi^0$	< 4	CL=90%
Γ_{106}	$\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0$	$(1.9 \pm 0.9) \%$	
Γ_{107}	$\bar{K}^*(892)^0 \eta$	$(1.9 \pm 0.5) \%$	
Γ_{108}	$K^- \pi^+ \omega$	$(3.1 \pm 0.6) \%$	
Γ_{109}	$\bar{K}^*(892)^0 \omega$	$(1.1 \pm 0.5) \%$	
Γ_{110}	$K^- \pi^+ \eta'(958)$	$(7.5 \pm 2.0) \times 10^{-3}$	
Γ_{111}	$\bar{K}^*(892)^0 \eta'(958)$	< 1.1	CL=90%
Plonic modes			
Γ_{112}	$\pi^+ \pi^-$	$(1.59 \pm 0.12) \times 10^{-3}$	
Γ_{113}	$\pi^0 \pi^0$	$(8.8 \pm 2.3) \times 10^{-4}$	
Γ_{114}	$\pi^+ \pi^- \pi^0$	$(1.6 \pm 1.1) \%$	S=2.7
Γ_{115}	$\pi^+ \pi^+ \pi^- \pi^-$	$(8.3 \pm 0.9) \times 10^{-3}$	
Γ_{116}	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	$(1.9 \pm 0.4) \%$	
Γ_{117}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$	$(4.0 \pm 3.0) \times 10^{-4}$	

Hadronic modes with two K 's

Γ_{118}	$K^+ K^-$	$(4.54 \pm 0.29) \times 10^{-3}$	
Γ_{119}	$K^0 \bar{K}^0$	$(1.1 \pm 0.4) \times 10^{-3}$	
Γ_{120}	$K^0 K^- \pi^+$	$(6.3 \pm 1.1) \times 10^{-3}$	S=1.2
Γ_{121}	$\bar{K}^*(892)^0 K^0$	< 1.0	CL=90%
	$\times B(\bar{K}^{*0} \rightarrow K^- \pi^+)$		
Γ_{122}	$K^*(892)^+ K^-$	$(2.3 \pm 0.5) \times 10^{-3}$	
	$\times B(K^{*+} \rightarrow K^0 \pi^+)$		
Γ_{123}	$K^0 K^- \pi^+$ nonresonant	$(2.4 \pm 2.4) \times 10^{-3}$	
Γ_{124}	$\bar{K}^0 K^+ \pi^-$	$(4.9 \pm 1.0) \times 10^{-3}$	
Γ_{125}	$K^*(892)^0 \bar{K}^0$	< 5	CL=90%
	$\times B(K^{*0} \rightarrow K^+ \pi^-)$		
Γ_{126}	$K^*(892)^- K^+$	$(1.2 \pm 0.7) \times 10^{-3}$	
	$\times B(K^{*-} \rightarrow \bar{K}^0 \pi^-)$		
Γ_{127}	$\bar{K}^0 K^+ \pi^-$ nonresonant	$(4.0 \pm 2.4) \times 10^{-3}$	
Γ_{128}	$K^+ K^- \pi^+ \pi^-$	$(2.4 \pm 0.5) \times 10^{-3}$	
Γ_{129}	$\phi \pi^+ \pi^- \times B(\phi \rightarrow K^+ K^-)$	$(1.3 \pm 0.4) \times 10^{-3}$	
Γ_{130}	$\phi \rho^0 \times B(\phi \rightarrow K^+ K^-)$	$(1.0 \pm 0.25) \times 10^{-3}$	
Γ_{131}	$K^*(892)^0 K^- \pi^+ + c.c. \times B(K^{*0} \rightarrow K^+ \pi^-)$	$(5 \pm \frac{9}{5}) \times 10^{-4}$	
Γ_{132}	$K^*(892)^0 \bar{K}^*(892)^0$	$(1.3 \pm 0.7) \times 10^{-3}$	
	$\times B^2(K^{*0} \rightarrow K^+ \pi^-)$		
Γ_{133}	$K^+ K^- \pi^+ \pi^-$ non- ϕ	$(1.7 \pm 0.5) \times 10^{-3}$	
Γ_{134}	$K^+ K^- \pi^+ \pi^-$ nonresonant	$(8 \pm \frac{90}{8}) \times 10^{-5}$	
Γ_{135}	$K^+ K^- \pi^+ \pi^- \pi^0$	$(3.1 \pm 2.0) \times 10^{-3}$	

Fractions of the following modes with resonances have already appeared above as submodes of particular charged-particle modes.

Γ_{136}	$\bar{K}^*(892)^0 K^0$	< 1.5	CL=90%
Γ_{137}	$K^*(892)^+ K^-$	$(3.4 \pm 0.8) \times 10^{-3}$	
Γ_{138}	$K^*(892)^0 \bar{K}^0$	< 8	CL=90%
Γ_{139}	$K^*(892)^- K^+$	$(1.8 \pm 1.0) \times 10^{-3}$	
Γ_{140}	$\phi \pi^+ \pi^-$	$(2.6 \pm 0.7) \times 10^{-3}$	
Γ_{141}	$\phi \rho^0$	$(1.9 \pm 0.5) \times 10^{-3}$	
Γ_{142}	$K^*(892)^0 K^- \pi^+ + c.c.$	$(8 \pm \frac{13}{8}) \times 10^{-4}$	
Γ_{143}	$K^*(892)^0 \bar{K}^*(892)^0$	$(2.9 \pm 1.6) \times 10^{-3}$	

**Doubly Cabibbo suppressed (DC) modes,
 $\Delta C = 2$ forbidden via mixing (C2M) modes,
 $\Delta C = 1$ weak neutral current (C1) modes, or
Lepton Family number (LF) violating modes**

Γ_{144}	$K^+ \pi^-$	DC	$(3.1 \pm 1.4) \times 10^{-4}$	
Γ_{145}	$K^+ \pi^-$ (via \bar{D}^0)	C2M	< 1.5	CL=90%
Γ_{146}	$K^+ \pi^+ \pi^- \pi^-$	DC	< 1.5	CL=90%
Γ_{147}	μ^- anything (via \bar{D}^0)	C2M	< 6	CL=90%
Γ_{148}	$e^+ e^-$	C1	< 1.3	CL=90%
Γ_{149}	$\mu^+ \mu^-$	C1	< 1.1	CL=90%
Γ_{150}	$\bar{K}^0 e^+ e^-$		< 1.7	CL=90%
Γ_{151}	$\rho^0 e^+ e^-$	C1	< 4.5	CL=90%
Γ_{152}	$\rho^0 \mu^+ \mu^-$	C1	< 8.1	CL=90%
Γ_{153}	$\mu^\pm e^\mp$	LF	$[h] < 1.0$	CL=90%

Γ_{154} A dummy mode used by the fit. $(24 \pm 4) \%$ S=1.1

[a] This is a weighted average of D^\pm (44%) and D^0 (56%) branching fractions. See " D^+ and $D^0 \rightarrow (\eta \text{ anything}) / (\text{total } D^+ \text{ and } D^0)$ " under " D^+ Branching Ratios" in these Full Listings.

[b] This value combines the e^+ and μ^+ branching fractions, making a small phase-space adjustment to the μ^+ fraction to be able to use it as an e^+ fraction; hence the " e^+ ." In fact, some of the e^+ measurements already use μ^+ events in this way.

[c] See the "Note on Semileptonic Decays of D and B Mesons" in the D^+ Full Listings for a comparison of inclusive and summed-inclusive branching fractions.

[d] ℓ indicates e or μ mode, not sum over modes.

[e] The limit on $(\bar{K}^*(892)\pi)^- \mu^+ \nu_\mu$ just below is much stronger.

[f] The branching fractions for this mode may differ from the sum of the submodes that contribute to it, due to interference effects. See the relevant papers.

[g] The two experiments determining this ratio are in serious disagreement. See the Full Listings.

[h] The value is for the sum of the charge states indicated.

$\Gamma(K^- e^+ \nu_e)/\Gamma(K^- \pi^+)$ Γ_8/Γ_{21}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.95 ± 0.04 OUR FIT				
0.95 ± 0.04 OUR AVERAGE				
0.978 ± 0.027 ± 0.044	2510	15 BEAN	93c CLEO	$e^+ e^- \approx \gamma(4S)$
0.90 ± 0.06 ± 0.06	584	CRAWFORD	91B CLEO	$e^+ e^- \approx 10.5$ GeV
0.91 ± 0.07 ± 0.11		ANJOS	89F E691	Photoproduction

¹⁵BEAN 93c uses $K^- \mu^+ \nu_\mu$ as well as $K^- e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

 $\Gamma(K^- \mu^+ \nu_\mu)/\Gamma(K^- \pi^+)$ Γ_9/Γ_{21}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.80 ± 0.10 OUR FIT				
0.80 ± 0.10 OUR AVERAGE				
0.82 ± 0.13 ± 0.13	99	FRABETTI	93i E687	$\gamma Be \bar{E}_\gamma = 221$ GeV
0.79 ± 0.08 ± 0.09	231	CRAWFORD	91B CLEO	$e^+ e^- \approx 10.5$ GeV

 $\Gamma(K^- \mu^+ \nu_\mu)/\Gamma(\mu^+ \text{anything})$ Γ_9/Γ_2

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.05 ± 0.05	124	KODAMA	91	EMUL pA 800 GeV

 $\Gamma(K^- \pi^0 e^+ \nu_e)/\Gamma_{total}$ Γ_{10}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.016 ± 0.013 ± 0.002	4	16 BAI	91	MRK3 $e^+ e^- \approx 3.77$ GeV

¹⁶BAI 91 finds that a fraction $0.79^{+0.15+0.09}_{-0.17-0.03}$ of combined D^+ and D^0 decays to $\bar{K} \pi e^+ \nu_e$ (24 events) are $\bar{K}^*(892)e^+ \nu_e$.

 $\Gamma(\bar{K}^0 \pi^- e^+ \nu_e)/\Gamma_{total}$ Γ_{11}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.028 ± 0.017 ± 0.008 ± 0.003	6	17 BAI	91	MRK3 $e^+ e^- \approx 3.77$ GeV

¹⁷BAI 91 finds that a fraction $0.79^{+0.15+0.09}_{-0.17-0.03}$ of combined D^+ and D^0 decays to $\bar{K} \pi e^+ \nu_e$ (24 events) are $\bar{K}^*(892)e^+ \nu_e$.

 $\Gamma(K^*(892)^- e^+ \nu_e)/\Gamma(K^- e^+ \nu_e)$ Γ_{20}/Γ_8

VALUE	DOCUMENT ID	TECN	COMMENT
0.52 ± 0.09 OUR FIT			
0.51 ± 0.18 ± 0.06	CRAWFORD	91B CLEO	$e^+ e^- \approx 10.5$ GeV

Unseen decay modes of the $K^*(892)^-$ are included.

 $\Gamma(K^*(892)^- e^+ \nu_e)/\Gamma(\bar{K}^0 \pi^+ \pi^-)$ Γ_{20}/Γ_{23}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.38 ± 0.06 OUR FIT				
0.38 ± 0.06 ± 0.03	152	18 BEAN	93c CLEO	$e^+ e^- \approx \gamma(4S)$

¹⁸BEAN 93c uses $K^* \mu^+ \nu_\mu$ as well as $K^* e^+ \nu_e$ events and makes a small phase-space adjustment to the number of the μ^+ events to use them as e^+ events.

 $\Gamma(K^*(892)^- \ell^+ \nu_\ell)/\Gamma(\bar{K}^0 \pi^+ \pi^-)$ Γ_{13}/Γ_{23}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.24 ± 0.07 ± 0.06	137	19 ALEXANDER	90B CLEO	$e^+ e^- 10.5-11$ GeV

This an average of the $K^*(892)^- e^+ \nu_e$ and $K^*(892)^- \mu^+ \nu_\mu$ ratios. Unseen decay modes of the $K^*(892)^-$ are included.

¹⁹ALEXANDER 90B cannot exclude extra π^0 's in the final state. See nearby data blocks for more detailed results.

 $\Gamma(K^- \pi^0 (\pi^0) e^+ \nu_e)/\Gamma_{total}$ Γ_{14}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.023 ± 0.050 ± 0.006 ± 0.001	1	20 AGUILAR...	87F HYBR	$\pi p, pp$ 360, 400 GeV

²⁰AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization. Does not distinguish presence of a second π^0 .

 $\Gamma(\bar{K}^0 \pi^- (\pi^0) e^+ \nu_e)/\Gamma_{total}$ Γ_{15}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.079 ± 0.069 ± 0.005	3	21 AGUILAR...	87F HYBR	$\pi p, pp$ 360, 400 GeV

²¹AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization. Does not distinguish presence of a second π^0 .

 $\Gamma(\bar{K}^*(892)^0 \pi^- e^+ \nu_e)/\Gamma(K^*(892)^- e^+ \nu_e)$ Γ_{16}/Γ_{20}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.64	90	CRAWFORD	91B CLEO	$e^+ e^- \approx 10.5$ GeV

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

 $\Gamma(K^- \pi^+ \pi^- \mu^+ \nu_\mu)/\Gamma(K^- \mu^+ \nu_\mu)$ Γ_{17}/Γ_9

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.037	90	KODAMA	93B E653	π^- emulsion 600 GeV

 $\Gamma(\bar{K}^*(892)\pi^- \mu^+ \nu_\mu)/\Gamma(K^- \mu^+ \nu_\mu)$ Γ_{18}/Γ_9

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.043	90	22 KODAMA	93B E653	π^- emulsion 600 GeV

²²KODAMA 93B searched in $K^- \pi^+ \pi^- \mu^+ \nu_\mu$, but the limit includes other $(\bar{K}^*(892)\pi^-)$ charge states.

 $\Gamma(\pi^- e^+ \nu_e)/\Gamma_{total}$ Γ_{19}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.0039 ± 0.0023 ± 0.0004	7	23 ADLER	89	MRK3 $e^+ e^- 3.77$ GeV

²³Experiment gives $|V_{cd}/V_{cs}|^2 = 0.057^{+0.038}_{-0.015} \pm 0.005$.

Hadronic modes with one or three K 's $\Gamma(K^- \pi^+)/\Gamma_{total}$ Γ_{21}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.0401 ± 0.0014 OUR FIT				
0.0397 ± 0.0016 OUR AVERAGE				
0.045 ± 0.006 ± 0.004		ALBRECHT	94	ARG $e^+ e^- \approx \gamma(4S)$
0.0391 ± 0.0008 ± 0.0017	4208	24,25 AKERIB	93	CLEO $e^+ e^- \approx \gamma(4S)$
0.0362 ± 0.0034 ± 0.0044		25 DECAMP	91J	ALEP From Z decays
0.045 ± 0.008 ± 0.005	56	25 ABACHI	88	HRS $e^+ e^- 29$ GeV
0.042 ± 0.004 ± 0.004	930	ADLER	88C	MRK3 $e^+ e^- 3.77$ GeV
0.041 ± 0.006	263	26 SCHINDLER	81	MRK2 $e^+ e^- 3.771$ GeV
0.043 ± 0.010	130	27 PERUZZI	77	MRK1 $e^+ e^- 3.77$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.0404 ± 0.0040 ± 0.0035		28 BARLAG	92C ACCM	$\pi^- Cu$ 230 GeV
0.040 ± 0.021 ± 0.010	7	28 AGUILAR...	87F HYBR	$\pi p, pp$ 360, 400 GeV

²⁴Radiative corrections increase this AKERIB 93 value to $0.0395 \pm 0.0008 \pm 0.0017$.

²⁵ABACHI 88, DECAMP 91J, and AKERIB 93 use $D^*(2010)^+ \rightarrow D^0 \pi^+$ decays. The π^+ is both slow and of low p_T with respect to the event thrust axis ($\approx D^{*+}$ direction). The excess number of such π^+ 's over background gives the number of $D^*(2010)^+ \rightarrow D^0 \pi^+$ events, and the fraction with $D^0 \rightarrow K^- \pi^+$ gives the $D^0 \rightarrow K^- \pi^+$ branching fraction.

²⁶SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.24 ± 0.02 . We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

²⁷PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.25 ± 0.05 . We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

²⁸AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization.

 $\Gamma(\bar{K}^0 \pi^0)/\Gamma(K^- \pi^+)$ Γ_{22}/Γ_{21}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.06 OUR FIT				Error includes scale factor of 1.1.
1.36 ± 0.23 ± 0.22	119	ANJOS	92B E691	γBe 80-240 GeV

 $\Gamma(\bar{K}^0 \pi^0)/\Gamma(\bar{K}^0 \pi^+ \pi^-)$ Γ_{22}/Γ_{23}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.389 ± 0.031 OUR FIT				
0.378 ± 0.033 OUR AVERAGE				
0.44 ± 0.02 ± 0.05	1942	PROCARIO	93B CLEO	$e^+ e^- 10.36-10.7$ GeV
0.34 ± 0.04 ± 0.02	92	29 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
0.36 ± 0.04 ± 0.08	104	KINOSHITA	91	CLEO $e^+ e^- \sim 10.7$ GeV

²⁹This value is calculated from numbers in Table 1 of ALBRECHT 92P.

 $\Gamma(\bar{K}^0 \pi^+ \pi^-)/\Gamma_{total}$ Γ_{23}/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.053 ± 0.006 OUR FIT				Error includes scale factor of 1.2.
0.063 ± 0.009 OUR AVERAGE				
0.064 ± 0.005 ± 0.010		ADLER	87	MRK3 $e^+ e^- 3.77$ GeV
0.052 ± 0.016	32	30 SCHINDLER	81	MRK2 $e^+ e^- 3.771$ GeV
0.079 ± 0.023	28	31 PERUZZI	77	MRK1 $e^+ e^- 3.77$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.0427 ± 0.0102 ± 0.0096		32 BARLAG	92C ACCM	$\pi^- Cu$ 230 GeV
0.045 ± 0.059 ± 0.014 ± 0.003	2	32 AGUILAR...	87F HYBR	$\pi p, pp$ 360, 400 GeV

³⁰SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.30 ± 0.08 . We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

³¹PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.46 ± 0.12 . We use the MARK-3 (ADLER 88C) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.

³²AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization.

 $\Gamma(\bar{K}^0 \pi^+ \pi^-)/\Gamma(K^- \pi^+)$ Γ_{23}/Γ_{21}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.31 ± 0.14 OUR FIT				Error includes scale factor of 1.2.
2.1 ± 0.6 OUR AVERAGE				
1.7 ± 0.8	35	AVERY	80	SPEC $\gamma N \rightarrow D^{*+}$
2.8 ± 1.0	116	PICCOLO	77	MRK1 $e^+ e^- 4.03, 4.41$ GeV

Meson Full Listings

 D^0

$\Gamma(K^0 \rho^0)/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{24}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.208 ± 0.025 OUR AVERAGE			
0.227 ± 0.032 ± 0.009	ALBRECHT	93D ARG	$e^+ e^- \approx 10$ GeV
0.215 ± 0.051 ± 0.037	ANJOS	93 E691	γ Be 90–260 GeV
0.20 ± 0.06 ± 0.03	FRABETTI	92B E687	γ Be $\bar{E}_\gamma = 221$ GeV
0.12 ± 0.01 ± 0.07	ADLER	87 MRK3	$e^+ e^- 3.77$ GeV

$\Gamma(K^0 f_0(980))/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{74}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.088 ± 0.035 ± 0.012	ALBRECHT	93D ARG	$e^+ e^- \approx 10$ GeV

$\Gamma(K^0 f_2(1270))/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{78}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.088 ± 0.037 ± 0.014	ALBRECHT	93D ARG	$e^+ e^- \approx 10$ GeV

$\Gamma(K^0 f_0(1300))/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{79}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.131 ± 0.045 ± 0.021	ALBRECHT	93D ARG	$e^+ e^- \approx 10$ GeV

$\Gamma(K^*(892)^- \pi^+)/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{81}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.07 OUR FIT			Error includes scale factor of 1.3.
0.97 ± 0.06 OUR AVERAGE			Error includes scale factor of 1.1.
1.08 ± 0.063 ± 0.045	ALBRECHT	93D ARG	$e^+ e^- \approx 10$ GeV
0.720 ± 0.145 ± 0.185	ANJOS	93 E691	γ Be 90–260 GeV
0.96 ± 0.12 ± 0.075	FRABETTI	92B E687	γ Be $\bar{E}_\gamma = 221$ GeV
0.84 ± 0.06 ± 0.08	ADLER	87 MRK3	$e^+ e^- 3.77$ GeV
1.05 +0.23 +0.07 -0.26 -0.09	25 SCHINDLER	81 MRK2	$e^+ e^- 3.771$ GeV

$\Gamma(K^0(1430)^- \pi^+)/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{103}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.208 ± 0.055 ± 0.034	ALBRECHT	93D ARG	$e^+ e^- \approx 10$ GeV

$\Gamma(K^*_2(1430)^- \pi^+)/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{104}/Γ_{23}	
VALUE	CL%	DOCUMENT ID	TECN
< 0.15	90	ALBRECHT	93D ARG

$\Gamma(K^0 \pi^+ \pi^- \text{ nonresonant})/\Gamma(K^0 \pi^+ \pi^-)$		Γ_{30}/Γ_{23}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.27 ± 0.04 OUR AVERAGE			
0.263 ± 0.024 ± 0.041	ANJOS	93 E691	γ Be 90–260 GeV
0.26 ± 0.08 ± 0.05	FRABETTI	92B E687	γ Be $\bar{E}_\gamma = 221$ GeV
0.33 ± 0.05 ± 0.10	ADLER	87 MRK3	$e^+ e^- 3.77$ GeV

$\Gamma(K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$		Γ_{31}/Γ	
VALUE	EVTS	DOCUMENT ID	TECN
0.138 ± 0.010 OUR FIT			Error includes scale factor of 1.1.
0.131 ± 0.016 OUR AVERAGE			
0.133 ± 0.012 ± 0.013	931	ADLER	88c MRK3 $e^+ e^- 3.77$ GeV
0.117 ± 0.043	37	33 SCHINDLER	81 MRK2 $e^+ e^- 3.771$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0912 +0.0173 -0.0165		34 BARLAG	92c ACCM π^- Cu 230 GeV
0.106 +0.061 ± 0.006 -0.028	5	34 AGUILAR...	87F HYBR $\pi p, p p$ 360, 400 GeV

³³SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.68 ± 0.23 . We use the MARK-3 (ADLER 88c) value of $\sigma = 5.8 \pm 0.5 \pm 0.6$ nb.


³⁴AGUILAR-BENITEZ 87F and BARLAG 92c compute the branching fraction using topological normalization.

$\Gamma(K^- \pi^+ \pi^0)/\Gamma(K^- \pi^+)$		Γ_{31}/Γ_{21}	
VALUE	EVTS	DOCUMENT ID	TECN
3.43 ± 0.24 OUR FIT			Error includes scale factor of 1.1.
3.07 ± 0.29 OUR AVERAGE			
3.04 ± 0.16 ± 0.34	931	35 ALBRECHT	92P ARG $e^+ e^- \approx 10$ GeV
4.0 ± 0.9 ± 1.0	69	ALVAREZ	91B NA14 Photoproduction
2.8 ± 0.14 ± 0.52	1050	KINOSHITA	91 CLEO $e^+ e^- \sim 10.7$ GeV
4.2 ± 1.4	41	SUMMERS	84 E691 Photoproduction

³⁵This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^- \rho^+)/\Gamma(K^- \pi^+ \pi^0)$		Γ_{32}/Γ_{31}	
VALUE	EVTS	DOCUMENT ID	TECN
0.75 ± 0.08 OUR AVERAGE			Error includes scale factor of 1.5. See the ideogram below.
0.647 ± 0.039 ± 0.150		ANJOS	93 E691 γ Be 90–260 GeV
0.81 ± 0.03 ± 0.06		ADLER	87 MRK3 $e^+ e^- 3.77$ GeV
0.31 +0.20 -0.14	13	SUMMERS	84 E691 Photoproduction
0.85 +0.11 +0.09 -0.15 -0.10	31	SCHINDLER	81 MRK2 $e^+ e^- 3.771$ GeV

WEIGHTED AVERAGE
0.75 ± 0.08 (Error scaled by 1.5)



$\Gamma(K^*(892)^- \pi^+)/\Gamma(K^- \pi^+ \pi^0)$		Γ_{81}/Γ_{31}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.35 ± 0.05 OUR FIT			Error includes scale factor of 1.4.
0.27 ± 0.04 OUR AVERAGE			
0.252 ± 0.033 ± 0.035	ANJOS	93 E691	γ Be 90–260 GeV
0.36 ± 0.06 ± 0.09	ADLER	87 MRK3	$e^+ e^- 3.77$ GeV

$\Gamma(\bar{K}^*(892)^0 \pi^0)/\Gamma(K^- \pi^+ \pi^0)$		Γ_{82}/Γ_{31}	
VALUE	DOCUMENT ID	TECN	COMMENT
0.219 ± 0.031 OUR FIT			Error includes scale factor of 1.4.
0.208 ± 0.035 OUR AVERAGE			
0.213 ± 0.027 ± 0.035	ANJOS	93 E691	γ Be 90–260 GeV
0.20 ± 0.03 ± 0.05	ADLER	87 MRK3	$e^+ e^- 3.77$ GeV

$\Gamma(K^- \pi^+ \pi^0 \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^0)$		Γ_{35}/Γ_{31}	
VALUE	EVTS	DOCUMENT ID	TECN
0.044 ± 0.019 OUR AVERAGE			Error includes scale factor of 1.1.
0.036 ± 0.004 ± 0.018		ANJOS	93 E691 γ Be 90–260 GeV
0.09 ± 0.02 ± 0.04		ADLER	87 MRK3 $e^+ e^- 3.77$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.51 ± 0.22	21	SUMMERS	84 E691 Photoproduction

$\Gamma(\bar{K}^*(892)^0 \pi^0)/\Gamma(K^0 \pi^0)$		Γ_{82}/Γ_{22}	
VALUE	EVTS	DOCUMENT ID	TECN
1.47 ± 0.26 OUR FIT			Error includes scale factor of 1.1.
1.65 +0.39 ± 0.20	122	PROCARIO	93B CLEO $\bar{K}^0 \pi^0 \pi^0$ Dalitz plot

$\Gamma(\bar{K}^*_2(1430)^0 \pi^0)/\Gamma(\bar{K}^*(892)^0 \pi^0)$		Γ_{105}/Γ_{82}	
VALUE	CL%	DOCUMENT ID	TECN
< 0.12	90	PROCARIO	93B CLEO $\bar{K}^0 \pi^0 \pi^0$ Dalitz plot

$\Gamma(K^0 \pi^0 \pi^0 \text{ nonresonant})/\Gamma(K^0 \pi^0)$		Γ_{38}/Γ_{22}	
VALUE	EVTS	DOCUMENT ID	TECN
0.37 ± 0.08 ± 0.04	76	PROCARIO	93B CLEO $\bar{K}^0 \pi^0 \pi^0$ Dalitz plot

$\Gamma(K^- \pi^+ \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{39} / Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.081 ± 0.005 OUR FIT				
0.086 ± 0.009 OUR AVERAGE				Error includes scale factor of 1.1.
0.079 ± 0.015 ± 0.009		ALBRECHT 94 ARG		$e^+ e^- \approx \gamma(4S)$
0.091 ± 0.008 ± 0.008	992	ADLER 88c MRK3		$e^+ e^- 3.77 \text{ GeV}$
0.117 ± 0.025	185	³⁶ SCHINDLER 81 MRK2		$e^+ e^- 3.771 \text{ GeV}$
0.062 ± 0.019	44	³⁷ PERUZZI 77 MRK1		$e^+ e^- 3.77 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0844 ^{+0.0064} _{-0.0053}		³⁸ BARLAG 92c ACCM		$\pi^- \text{ Cu } 230 \text{ GeV}$
0.065 ^{+0.017} _{-0.011} ± 0.019	13	³⁸ AGUILAR-... 87F HYBR		$\pi p, pp 360, 400 \text{ GeV}$

³⁶SCHINDLER 81 (MARK-2) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.68 ± 0.11 . We use the MARK-3 (ADLER 88c) value of $\sigma = 5.8 \pm 0.5 \pm 0.6 \text{ nb}$.
³⁷PERUZZI 77 (MARK-1) measures $\sigma(e^+ e^- \rightarrow \psi(3770)) \times$ branching fraction to be 0.36 ± 0.10 . We use the MARK-3 (ADLER 88c) value of $\sigma = 5.8 \pm 0.5 \pm 0.6 \text{ nb}$.
³⁸AGUILAR-BENITEZ 87F and BARLAG 92c compute the branching fraction using topological normalization.

 $\Gamma(K^- \pi^+ \pi^+ \pi^-) / \Gamma(K^- \pi^+)$ $\Gamma_{39} / \Gamma_{21}$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
2.02 ± 0.11 OUR FIT				
2.01 ± 0.13 OUR AVERAGE				
1.7 ± 0.2 ± 0.2	1745	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
1.90 ± 0.25 ± 0.20	337	ALVAREZ 91B NA14		Photoproduction
2.12 ± 0.16 ± 0.09		BORTOLETTO88 CLEO		$e^+ e^- 10.55 \text{ GeV}$
2.0 ± 0.9	48	BAILEY 86 ACCM		$\pi^- \text{ Be fixed target}$
2.17 ± 0.28 ± 0.23		ALBRECHT 85F ARG		$e^+ e^- 10 \text{ GeV}$
2.0 ± 1.0	10	BAILEY 83B SPEC		$\pi^- \text{ Be} \rightarrow D^0$
2.2 ± 0.8	214	PICCOLO 77 MRK1		$e^+ e^- 4.03, 4.41 \text{ GeV}$

 $\Gamma(K^- \pi^+ \rho^0 \text{ total}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{40} / \Gamma_{39}$

This includes $K^- a_1(1260)^+$, $\bar{K}^*(892)^0 \rho^0$, etc. The next entry gives the specifically 3-body fraction. We rely on the MARK III and E691 full amplitude analyses of the $K^- \pi^+ \pi^+ \pi^-$ channel for values of the resonant substructure.

VALUE	DOCUMENT ID	TECN	COMMENT
0.835 ± 0.035 OUR AVERAGE			
0.80 ± 0.03 ± 0.05	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
0.855 ± 0.032 ± 0.030	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.98 ± 0.12 ± 0.10	ALVAREZ 91B NA14		Photoproduction

 $\Gamma(K^- \pi^+ \rho^0 \text{ 3-body}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{41} / \Gamma_{39}$

We rely on the MARK III and E691 full amplitude analyses of the $K^- \pi^+ \pi^+ \pi^-$ channel for values of the resonant substructure.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.063 ± 0.028 OUR AVERAGE				
0.05 ± 0.03 ± 0.02		ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
0.084 ± 0.022 ± 0.04		COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.77 ± 0.06 ± 0.06	³⁹	ALVAREZ 91B NA14		Photoproduction
0.85 ^{+0.11} _{-0.22}	180	PICCOLO 77 MRK1		$e^+ e^- 4.03, 4.41 \text{ GeV}$

³⁹This value is for $\rho^0(K^- \pi^+)$ -nonresonant. ALVAREZ 91B cannot determine what fraction of this is $K^- a_1(1260)^+$.

 $\Gamma(\bar{K}^*(892)^0 \rho^0) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{87} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included. We rely on the MARK III and E691 full amplitude analyses of the $K^- \pi^+ \pi^+ \pi^-$ channel for values of the resonant substructure.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.195 ± 0.03 ± 0.03				
0.165 ± 0.03 ± 0.045		ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.34 ± 0.09 ± 0.09		ALVAREZ 91B NA14		Photoproduction
0.75 ± 0.3	5	BAILEY 83B SPEC		$\pi \text{ Be} \rightarrow D^0$
0.15 ^{+0.16} _{-0.15}	20	PICCOLO 77 MRK1		$e^+ e^- 4.03, 4.41 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ transverse}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{88} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.20 ± 0.07 OUR FIT			
0.213 ± 0.024 ± 0.075			
0.23 ± 0.02 ± 0.03	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
0.242 ± 0.025 ± 0.06	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ S-wave}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{89} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.375 ± 0.045 ± 0.06			
0.134 ^{+0.032} _{-0.033}	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ S-wave long.}) / \Gamma_{\text{total}}$ Γ_{90} / Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003				
0.103 ± 0.022 ± 0.025	90	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ P-wave}) / \Gamma_{\text{total}}$ Γ_{91} / Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.003				
0.194 ± 0.056 ± 0.088	90	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.009	90	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \text{ D-wave}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{92} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.255 ± 0.045 ± 0.06			
0.165 ± 0.03 ± 0.045	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 f_0(980)) / \Gamma_{\text{total}}$ Γ_{98} / Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ and $f_0(980)$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.007				
0.23 ± 0.02 ± 0.03	90	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(K^- a_1(1260)^+) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{76} / \Gamma_{39}$

Unseen decay modes of the $a_1(1260)^+$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.14 OUR AVERAGE			
0.94 ± 0.13 ± 0.20	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
0.984 ± 0.048 ± 0.16	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(K^- a_2(1320)^+) / \Gamma_{\text{total}}$ Γ_{80} / Γ

Unseen decay modes of the $a_2(1320)^+$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.002				
0.13 ± 0.04 OUR FIT	90	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.006	90	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(K_1(1270)^- \pi^+) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{99} / \Gamma_{39}$

Unseen decay modes of the $K_1(1270)^-$ are included. The two experiments disagree considerably here.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.04 OUR FIT				
0.194 ± 0.056 ± 0.088				
0.165 ± 0.03 ± 0.045	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.013	90	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(K_1(1400)^- \pi^+) / \Gamma_{\text{total}}$ Γ_{100} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.012				
0.210 ± 0.027 ± 0.06	90	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(K^*(1410)^- \pi^+) / \Gamma_{\text{total}}$ Γ_{102} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.012				
0.165 ± 0.03 ± 0.045	90	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \text{ total}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{83} / \Gamma_{39}$

This includes $\bar{K}^*(892)^0 \rho^0$, etc. The next entry gives the specifically 3-body fraction. Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.30 ± 0.06 ± 0.03			
0.23 ± 0.02 ± 0.03	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \text{ 3-body}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{84} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.19 ± 0.04 OUR FIT			
0.18 ± 0.04 OUR AVERAGE			
0.165 ± 0.03 ± 0.045	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
0.210 ± 0.027 ± 0.06	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(K^- \pi^+ f_0(980)) / \Gamma_{\text{total}}$ Γ_{97} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011				
0.23 ± 0.02 ± 0.03	90	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$

 $\Gamma(K^- \pi^+ \pi^+ \pi^- \text{ nonresonant}) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{47} / \Gamma_{39}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.233 ± 0.032 OUR AVERAGE			
0.23 ± 0.02 ± 0.03	ANJOS 92c E691		$\gamma \text{ Be } 90\text{--}260 \text{ GeV}$
0.242 ± 0.025 ± 0.06	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$

 $\Gamma(\bar{K}^0 \pi^+ \pi^-) / \Gamma_{\text{total}}$ Γ_{48} / Γ

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.098 ± 0.014 OUR FIT				Error includes scale factor of 1.1.
0.103 ± 0.022 ± 0.025				
0.103 ± 0.022 ± 0.025	140	COFFMAN 92B MRK3		$e^+ e^- 3.77 \text{ GeV}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.134 ^{+0.032} _{-0.033}		⁴⁰ BARLAG 92c ACCM		$\pi^- \text{ Cu } 230 \text{ GeV}$

⁴⁰BARLAG 92c computes the branching fraction using topological normalization.

Meson Full Listings

 D^0 $\Gamma(K^0 \pi^+ \pi^- \pi^0) / \Gamma(K^0 \pi^+ \pi^-)$ $\Gamma_{48} / \Gamma_{23}$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.86 ± 0.21 OUR FIT				
1.86 ± 0.23 OUR AVERAGE				
1.80 ± 0.20 ± 0.21	190	41 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
2.8 ± 0.8 ± 0.8	46	ANJOS	92C E691	γ Be 90–260 GeV
1.85 ± 0.26 ± 0.30	158	KINOSHITA	91 CLEO	$e^+ e^- \sim 10.7$ GeV

⁴¹ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

 $\Gamma(K^0 \eta) / \Gamma(K^- \pi^+)$ $\Gamma_{69} / \Gamma_{21}$

Unseen decay modes of the η are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.14 ± 0.04	90	ALBRECHT	89D ARG	$e^+ e^- 10$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(K^0 \eta) / \Gamma(K^0 \pi^0)$ $\Gamma_{69} / \Gamma_{22}$

Unseen decay modes of the η are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.04 OUR FIT				
0.32 ± 0.04 ± 0.03	225	PROCARIO	93B CLEO	$\eta \rightarrow \gamma \gamma$

 $\Gamma(K^0 \eta) / \Gamma(K^0 \pi^+ \pi^-)$ $\Gamma_{69} / \Gamma_{23}$

Unseen decay modes of the η are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.129 ± 0.017 OUR FIT				
0.14 ± 0.02 ± 0.02	80	PROCARIO	93B CLEO	$\eta \rightarrow \pi^+ \pi^- \pi^0$

 $\Gamma(K^0 \omega) / \Gamma(K^- \pi^+)$ $\Gamma_{72} / \Gamma_{21}$

Unseen decay modes of the ω are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.50 ± 0.09 OUR FIT				
1.00 ± 0.36 ± 0.20		ALBRECHT	89D ARG	$e^+ e^- 10$ GeV

 $\Gamma(K^0 \omega) / \Gamma(K^0 \pi^+ \pi^-)$ $\Gamma_{72} / \Gamma_{23}$

Unseen decay modes of the ω are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.38 ± 0.07 OUR FIT				
0.33 ± 0.09 OUR AVERAGE				Error includes scale factor of 1.1.
0.29 ± 0.08 ± 0.05	16	42 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV
0.54 ± 0.14 ± 0.16	40	KINOSHITA	91 CLEO	$e^+ e^- \sim 10.7$ GeV

⁴² This value is calculated from numbers in Table 1 of ALBRECHT 92P.

 $\Gamma(K^0 \omega) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{72} / \Gamma_{48}$

Unseen decay modes of the ω are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.21 ± 0.04 OUR FIT				
0.220 ± 0.048 ± 0.0116		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^0 \eta(958)) / \Gamma(K^0 \pi^+ \pi^-)$ $\Gamma_{73} / \Gamma_{23}$

Unseen decay modes of the $\eta(958)$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.04 OUR AVERAGE				
0.31 ± 0.02 ± 0.04	594	PROCARIO	93B CLEO	$\eta' \rightarrow \eta \pi^+ \pi^-, \rho^0 \gamma$
0.37 ± 0.13 ± 0.06	18	43 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV

⁴³ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

 $\Gamma(K^*(892)^- \rho^+) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{93} / \Gamma_{48}$

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.606 ± 0.188 ± 0.126		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^*(892)^- \rho^+ \text{longitudinal}) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{94} / \Gamma_{48}$

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.290 ± 0.111		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^*(892)^- \rho^+ \text{transverse}) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{95} / \Gamma_{48}$

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.317 ± 0.180		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^*(892)^- \rho^+ P\text{-wave}) / \Gamma_{\text{total}}$ Γ_{96} / Γ

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.015	90	44 COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

⁴⁴ Obtained using other $\bar{K}^*(892) \rho$ P-wave limits and isospin relations.

 $\Gamma(\bar{K}^*(892)^0 \rho^0 \text{transverse}) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{88} / \Gamma_{48}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.06 OUR FIT				Error includes scale factor of 1.1.
0.126 ± 0.111		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^0 a_1(1260)^0) / \Gamma_{\text{total}}$ Γ_{77} / Γ

Unseen decay modes of the $a_1(1260)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.019	90	COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K_1(1270)^- \pi^+) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{99} / \Gamma_{48}$

Unseen decay modes of the $K_1(1270)^-$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.107 ± 0.030 OUR FIT				Error includes scale factor of 1.1.
0.10 ± 0.03		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(\bar{K}_1(1400)^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{101} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.037	90	COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \text{3-body}) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{84} / \Gamma_{48}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.04 OUR FIT				Error includes scale factor of 1.1.
0.191 ± 0.105		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^0 \pi^+ \pi^- \pi^0 \text{nonresonant}) / \Gamma(K^0 \pi^+ \pi^- \pi^0)$ $\Gamma_{55} / \Gamma_{48}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.210 ± 0.147 ± 0.150		COFFMAN	92B MRK3	$e^+ e^- 3.77$ GeV

 $\Gamma(K^- \pi^+ \pi^0 \pi^0) / \Gamma_{\text{total}}$ Γ_{56} / Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.149 ± 0.037 ± 0.030	24	45 ADLER	88C MRK3	$e^+ e^- 3.77$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.177 ± 0.029	46	BARLAG	92C ACCM	π^- Cu 230 GeV
$0.209^{+0.074}_{-0.043} \pm 0.012$	9	46 AGUILAR...	87F HYBR	π , pp 360, 400 GeV

⁴⁵ ADLER 88C uses an absolute normalization method finding this decay channel opposite a detected $\bar{D}^0 \rightarrow K^+ \pi^-$ in pure $D\bar{D}$ events.

⁴⁶ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization. They do not distinguish the presence of a third π^0 , and thus are not included in the average.

 $\Gamma(K^- \pi^+ \pi^+ \pi^- \pi^0) / \Gamma_{\text{total}}$ Γ_{57} / Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0293 ± 0.0051 - 0.0045		47 BARLAG	92C ACCM	π^- Cu 230 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

⁴⁷ BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(K^- \pi^+ \pi^+ \pi^- \pi^0) / \Gamma(K^- \pi^+)$ $\Gamma_{57} / \Gamma_{21}$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.07 ± 0.11 OUR FIT				
0.98 ± 0.11 ± 0.11	225	48 ALBRECHT	92P ARG	$e^+ e^- \approx 10$ GeV

⁴⁸ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

 $\Gamma(K^- \pi^+ \pi^+ \pi^- \pi^0) / \Gamma(K^- \pi^+ \pi^+ \pi^-)$ $\Gamma_{57} / \Gamma_{39}$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.53 ± 0.05 OUR FIT				
0.56 ± 0.07 OUR AVERAGE				

0.55 ± 0.07 ± 0.12 - 0.09

0.57 ± 0.06 ± 0.05

167 KINOSHITA 91 CLEO $e^+ e^- \sim 10.7$ GeV

180 ANJOS 90D E691 Photoproduction

 $\Gamma(\bar{K}^*(892)^0 \pi^+ \pi^- \pi^0) / \Gamma(K^- \pi^+ \pi^+ \pi^- \pi^0)$ $\Gamma_{106} / \Gamma_{57}$

Unseen decay modes of the $\bar{K}^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.15 ± 0.15		ANJOS	90D E691	Photoproduction

 $\Gamma(\bar{K}^*(892)^0 \eta) / \Gamma(K^- \pi^+)$ $\Gamma_{107} / \Gamma_{21}$

Unseen decay modes of the $\bar{K}^*(892)^0$ and η are included.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.46 ± 0.11 OUR FIT					
0.58 ± 0.19 ± 0.24 - 0.28		46	KINOSHITA	91 CLEO	$e^+ e^- \sim 10.7$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.70 90 ALBRECHT 89D ARG $e^+ e^- 10$ GeV

 $\Gamma(\bar{K}^*(892)^0 \eta) / \Gamma(K^- \pi^+ \pi^0)$ $\Gamma_{107} / \Gamma_{31}$

Unseen decay modes of the $\bar{K}^*(892)^0$ and η are included.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.135 ± 0.034 OUR FIT					
0.13 ± 0.02 ± 0.03		214	PROCARIO	93B CLEO	$\bar{K}^* \rightarrow K^- \pi^+, \eta \rightarrow \gamma \gamma$

 $\Gamma(\bar{K}^*(892)^0 \eta) / \Gamma(K^- \pi^+ \pi^+ \pi^- \pi^0)$ $\Gamma_{107} / \Gamma_{57}$

Unseen decay modes of the $\bar{K}^*(892)^0$ and η are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.27	90	49 ANJOS	90D E691	Photoproduction

⁴⁹ Recovered from the published limit, $\Gamma(\bar{K}^*(892)^0 \eta) / \Gamma_{\text{total}}$, in order to make our normalization consistent.

$\Gamma(K^-\pi^+\omega)/\Gamma(K^-\pi^+)$ Γ_{108}/Γ_{21}
 Unseen decay modes of the ω are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.78 \pm 0.12 \pm 0.10$	99	⁵⁰ ALBRECHT 92P ARG		$e^+e^- \approx 10$ GeV

⁵⁰ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^*(892)^0\omega)/\Gamma(K^-\pi^+)$ Γ_{109}/Γ_{21}
 Unseen decay modes of the $K^*(892)^0$ and ω are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.28 \pm 0.11 \pm 0.04$	17	⁵¹ ALBRECHT 92P ARG		$e^+e^- \approx 10$ GeV

⁵¹ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

$\Gamma(K^*(892)^0\omega)/\Gamma(K^-\pi^+\pi^-\pi^0)$ Γ_{109}/Γ_{57}
 Unseen decay modes of the $K^*(892)^0$ and ω are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.44	90	⁵² ANJOS 90D E691		Photoproduction

⁵² Recovered from the published limit, $\Gamma(K^*(892)^0\omega)/\Gamma_{total}$, in order to make our normalization consistent.

$\Gamma(K^-\pi^+\eta'(958))/\Gamma(K^-\pi^+\pi^-\pi^0)$ Γ_{110}/Γ_{39}
 Unseen decay modes of the $\eta'(958)$ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.093 \pm 0.014 \pm 0.019$	286	PROCARIO 93B CLEO		$\eta' \rightarrow \eta\pi^+\pi^-, \rho^0\gamma$

$\Gamma(K^*(892)^0\eta'(958))/\Gamma(K^-\pi^+\eta'(958))$ $\Gamma_{111}/\Gamma_{110}$
 Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.15	90	PROCARIO 93B CLEO		

$\Gamma(K^0\pi^+\pi^+\pi^-\pi^0)/\Gamma_{total}$ Γ_{62}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0124 ± 0.0034 -0.0032	⁵³ BARLAG 92C ACCM π^- Cu 230 GeV		


⁵³ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^0\pi^+\pi^+\pi^-\pi^0)/\Gamma(K^0\pi^+\pi^-)$ Γ_{62}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.107 ± 0.029 OUR AVERAGE				Error Includes scale factor of 1.8. See the ideogram below.
$0.07 \pm 0.02 \pm 0.01$	11	⁵⁴ ALBRECHT 92P ARG		$e^+e^- \approx 10$ GeV
0.149 ± 0.026	56	AMMAR 91 CLEO		$e^+e^- \approx 10.5$ GeV
$0.18 \pm 0.07 \pm 0.04$	6	ANJOS 90D E691		Photoproduction

⁵⁴ This value is calculated from numbers in Table 1 of ALBRECHT 92P.

WEIGHTED AVERAGE
 0.107 ± 0.029 (Error scaled by 1.8)



$\Gamma(K^0\pi^+\pi^-\pi^0\pi^0)/\Gamma_{total}$ Γ_{63}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.106 ± 0.073 -0.029 ± 0.006	4	⁵⁵ AGUILAR... 87F HYBR $\pi p, pp$ 360, 400 GeV		

⁵⁵ AGUILAR-BENITEZ 87F computes the branching fraction using topological normalization, and does not distinguish the presence of a third π^0 .

$\Gamma(K^0K^+K^-)/\Gamma(K^0\pi^+\pi^-)$ $\Gamma_{64}/\Gamma_{23} = (\Gamma_{66} + \frac{1}{2}\Gamma_{75})/\Gamma_{23}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.172 ± 0.014 OUR FIT				
0.178 ± 0.019 OUR AVERAGE				
$0.20 \pm 0.05 \pm 0.04$	47	FRABETTI 92B E687		$\gamma Be \bar{E}_\gamma = 221$ GeV
0.170 ± 0.022	136	AMMAR 91 CLEO		$e^+e^- \approx 10.5$ GeV
0.24 ± 0.08		BEBEK 86 CLEO		e^+e^- near $\Upsilon(4S)$
0.185 ± 0.055	52	ALBRECHT 85B ARG		$e^+e^- 10$ GeV

$\Gamma(K^0\phi)/\Gamma_{total}$ Γ_{75}/Γ
 Unseen decay modes of the ϕ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0160 ± 0.0059 -0.0041	⁵⁶ BARLAG 92C ACCM π^- Cu 230 GeV		
$0.0086 \pm 0.0050 \pm 0.0031$ $-0.0041 - 0.0018$	ADLER 88C MRK3		$e^+e^- 3.77$ GeV

⁵⁶ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^0\phi)/\Gamma(K^0\pi^+\pi^-)$ Γ_{75}/Γ_{23}
 Unseen decay modes of the ϕ are included.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.158 ± 0.016 OUR FIT				
0.156 ± 0.017 OUR AVERAGE				
$0.13 \pm 0.06 \pm 0.02$	13	FRABETTI 92B E687		$\gamma Be \bar{E}_\gamma = 221$ GeV
0.163 ± 0.023	63	AMMAR 91 CLEO		$e^+e^- \approx 10.5$ GeV
0.155 ± 0.033	56	ALBRECHT 87E ARG		$e^+e^- 10$ GeV
0.14 ± 0.05	29	BEBEK 86 CLEO		e^+e^- near $\Upsilon(4S)$
0.186 ± 0.052	26	ALBRECHT 85B ARG		See ALBRECHT 87E

$\Gamma(K^0K^+K^-\text{non-}\phi)/\Gamma_{total}$ Γ_{66}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0048 ± 0.0033 -0.0026	⁵⁷ BARLAG 92C ACCM π^- Cu 230 GeV		
$0.0085 \pm 0.0027 \pm 0.0020$ $-0.0024 - 0.0018$	ADLER 88C MRK3		$e^+e^- 3.77$ GeV

⁵⁷ BARLAG 92C computes the branching fraction using topological normalization.

$\Gamma(K^0K^+K^-\text{non-}\phi)/\Gamma(K^0\pi^+\pi^-)$ Γ_{66}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.093 ± 0.014 OUR FIT				
0.088 ± 0.019 OUR AVERAGE				
$0.11 \pm 0.04 \pm 0.03$	20	FRABETTI 92B E687		$\gamma Be \bar{E}_\gamma = 221$ GeV
0.084 ± 0.020		ALBRECHT 87E ARG		$e^+e^- 10$ GeV

$\Gamma(K_S^0K_S^0K_S^0)/\Gamma(K^0\pi^+\pi^-)$ Γ_{67}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.016 ± 0.004 OUR AVERAGE				
0.016 ± 0.005	22	AMMAR 91 CLEO		$e^+e^- \approx 10.5$ GeV
$0.017 \pm 0.007 \pm 0.005$	5	ALBRECHT 90C ARG		$e^+e^- \approx 10$ GeV

$\Gamma(K^+K^-K^0\pi^0)/\Gamma_{total}$ Γ_{68}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0072 ± 0.0048 -0.0036	⁵⁸ BARLAG 92C ACCM π^- Cu 230 GeV		

⁵⁸ BARLAG 92C computes the branching fraction using topological normalization.

Plonic modes

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$ Γ_{112}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
9 ± 5		⁵⁹ BARLAG 92C ACCM π^- Cu 230 GeV		
50 ± 120 -20 ± 40	1	⁵⁹ AGUILAR... 87F HYBR $\pi p, pp$ 360, 400 GeV		

⁵⁹ AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization.

$\Gamma(\pi^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{112}/Γ_{21}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0396 ± 0.0027 OUR AVERAGE				
$0.043 \pm 0.007 \pm 0.003$	177	FRABETTI 94C E687		$\gamma Be \bar{E}_\gamma = 220$ GeV
$0.0348 \pm 0.0030 \pm 0.0023$	227	SELEN 93 CLEO		$e^+e^- \approx \Upsilon(4S)$
$0.048 \pm 0.013 \pm 0.008$	51	ADAMOVICH 92 OMEG		$\pi^- 340$ GeV
$0.055 \pm 0.008 \pm 0.005$	120	ANJOS 91D E691		Photoproduction
$0.040 \pm 0.007 \pm 0.006$	57	ALBRECHT 90C ARG		$e^+e^- \approx 10$ GeV
$0.050 \pm 0.007 \pm 0.005$	110	ALEXANDER 90 CLEO		$e^+e^- 10.5-11$ GeV
$0.033 \pm 0.010 \pm 0.006$	39	BALTRUSAIT...85E MRK3		$e^+e^- 3.77$ GeV
0.033 ± 0.015		ABRAMS 79D MRK2		$e^+e^- 3.77$ GeV

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$ Γ_{113}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0046	90	ALEXANDER 90 CLEO		$e^+e^- 10.5-11$ GeV

$\Gamma(\pi^0\pi^0)/\Gamma(K^-\pi^+)$ Γ_{113}/Γ_{21}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.022 \pm 0.004 \pm 0.004$	40	SELEN 93 CLEO		$e^+e^- \approx \Upsilon(4S)$

Meson Full Listings

 D^0 $\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{114}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.016 ± 0.011 OUR AVERAGE				Error includes scale factor of 2.7.
0.0390 ^{+0.0100} _{-0.0095}		60 BARLAG	92C ACCM	π^- Cu 230 GeV
0.011 ± 0.004 ± 0.002	10	61 BALTRUSAIT..85E MRK3		e^+e^- 3.77 GeV
				60 BARLAG 92C computes the branching fraction using topological normalization. Possible contamination by extra π^0 's may partly explain the unexpectedly large value.
				61 All the BALTRUSAITIS 85E events are consistent with $\rho^0\pi^0$.

 $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0028 ± 0.0008		62 BARLAG	92C ACCM	π^- Cu 230 GeV
0.005 ^{+0.011} _{-0.001} ± 0.001	1	62 AGUILAR...	87F HYBR	$\pi\rho, \rho\rho$ 360, 400 GeV
0.015 ± 0.006 ± 0.002	9	BALTRUSAIT..85E MRK3		e^+e^- 3.77 GeV
				62 AGUILAR-BENITEZ 87F and BARLAG 92C compute the branching fraction using topological normalization.

 $\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma(K^-\pi^+\pi^-)$ Γ_{115}/Γ_{39}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.103 ± 0.009 OUR AVERAGE				
0.115 ± 0.023 ± 0.016	64	ADAMOVICH 92	OMEG π^-	340 GeV
0.108 ± 0.024 ± 0.008	79	FRABETTI 92	E687	γ Be
0.102 ± 0.013	345	63 AMMAR 91	CLEO	$e^+e^- \approx 10.5$ GeV
0.096 ± 0.018 ± 0.007	66	ANJOS 91	E691	γ Be 80–240 GeV
				63 AMMAR 91 finds $1.25 \pm 0.25 \pm 0.25 \rho^0$'s per $\pi^+\pi^+\pi^-\pi^-$ decay, but can't untangle the resonant substructure ($\rho^0\rho^0, a_1^\pm\pi^\mp, \rho^0\pi^+\pi^-$).

 $\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{116}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0192 ± 0.0041 -0.0038	64 BARLAG	92C ACCM	π^- Cu 230 GeV
			64 BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(\pi^+\pi^+\pi^-\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{117}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0004 ± 0.0003	65 BARLAG	92C ACCM	π^- Cu 230 GeV
			65 BARLAG 92C computes the branching fraction using topological normalization.

Hadronic modes with two K 's $\Gamma(K^+K^-)/\Gamma_{\text{total}}$ Γ_{118}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.0051 ^{+0.0012} _{-0.0011}	66 BARLAG	92C ACCM	π^- Cu 230 GeV
			66 BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(K^+K^-)/\Gamma(K^-\pi^+)$ Γ_{118}/Γ_{21}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.113 ± 0.006 OUR FIT				
0.113 ± 0.006 OUR AVERAGE				
0.109 ± 0.007 ± 0.009	581	FRABETTI 94C	E687	γ Be $\bar{E}_\gamma = 220$ GeV
0.107 ± 0.029 ± 0.015	103	ADAMOVICH 92	OMEG π^-	340 GeV
0.138 ± 0.027 ± 0.010	155	FRABETTI 92	E687	γ Be
0.16 ± 0.05	34	ALVAREZ 91B	NA14	Photoproduction
0.107 ± 0.010 ± 0.009	193	ANJOS 91D	E691	Photoproduction
0.10 ± 0.02 ± 0.01	131	ALBRECHT 90C	ARG	$e^+e^- \approx 10$ GeV
0.117 ± 0.010 ± 0.007	249	ALEXANDER 90	CLEO	e^+e^- 10.5–11 GeV
0.122 ± 0.018 ± 0.012	118	BALTRUSAIT..85E MRK3		e^+e^- 3.77 GeV
0.113 ± 0.030		ABRAMS 79D	MRK2	e^+e^- 3.77 GeV

 $\Gamma(K^+K^-)/\Gamma(\pi^+\pi^-)$ $\Gamma_{118}/\Gamma_{112}$

The unused results here are redundant with $\Gamma(K^+K^-)/\Gamma(K^-\pi^+)$ and $\Gamma(\pi^+\pi^-)/\Gamma(K^-\pi^+)$ measurements by the same experiments.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.53 ± 0.46 ± 0.19		FRABETTI 94C	E687	γ Be $\bar{E}_\gamma = 220$ GeV
2.23 ± 0.81 ± 0.46		ADAMOVICH 92	OMEG π^-	340 GeV
1.95 ± 0.34 ± 0.22		ANJOS 91D	E691	Photoproduction
2.5 ± 0.7		ALBRECHT 90C	ARG	$e^+e^- \approx 10$ GeV
2.35 ± 0.37 ± 0.28	110	ALEXANDER 90	CLEO	e^+e^- 10.5–11 GeV

 $\Gamma(K^0\bar{K}^0)/\Gamma_{\text{total}}$ Γ_{119}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.0046	90	ADLER 88C	MRK3	e^+e^- 3.77 GeV

 $\Gamma(K^0\bar{K}^0)/\Gamma(K^-\pi^+)$ Γ_{119}/Γ_{21}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.032	90	ANJOS 92B	E691	γ Be 80–240 GeV

 $\Gamma(K^0\bar{K}^0)/\Gamma(K^0\pi^+\pi^-)$ Γ_{119}/Γ_{23}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.021 ± 0.009 OUR FIT					
0.021 ± 0.011 -0.008 ± 0.002	5	ALEXANDER 90	CLEO	e^+e^- 10.5–11 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.016	90	ALBRECHT 90C	ARG	$e^+e^- \approx 10$ GeV	

 $\Gamma(K^0\bar{K}^0)/\Gamma(K^+K^-)$ $\Gamma_{119}/\Gamma_{118}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.24 ± 0.10 0.08				OUR FIT
0.24 ± 0.16	4	67 CUMALAT 88	SPEC	nN 0–800 GeV
				67 Includes a correction communicated to us by the authors of CUMALAT 88.

 $\Gamma(K^0K^-\pi^+)/\Gamma(K^-\pi^+)$ Γ_{120}/Γ_{21}

VALUE	DOCUMENT ID	TECN	COMMENT
0.157 ± 0.028 OUR FIT			Error includes scale factor of 1.1.
0.16 ± 0.06	68 ANJOS 91	E691	γ Be 80–240 GeV
			68 The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^0K^-\pi^+)/\Gamma(K^0\pi^+\pi^-)$ Γ_{120}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.120 ± 0.019 OUR FIT				Error includes scale factor of 1.2.
0.119 ± 0.021 OUR AVERAGE				Error includes scale factor of 1.3.
0.108 ± 0.019	61	AMMAR 91	CLEO	$e^+e^- \approx 10.5$ GeV
0.16 ± 0.03 ± 0.02	39	ALBRECHT 90C	ARG	$e^+e^- \approx 10$ GeV

 $\Gamma(K^*(892)^0K^0)/\Gamma(K^-\pi^+)$ Γ_{136}/Γ_{21}

VALUE	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the $K^*(892)^0$ are included.			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.00 ^{+0.03} _{-0.00}	69 ANJOS 91	E691	γ Be 80–240 GeV
			69 The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^*(892)^0K^0)/\Gamma(K^0\pi^+\pi^-)$ Γ_{136}/Γ_{23}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the $K^*(892)^0$ are included.				
<0.029	90	AMMAR 91	CLEO	$e^+e^- \approx 10.5$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.03	90	ALBRECHT 90C	ARG	$e^+e^- \approx 10$ GeV

 $\Gamma(K^*(892)^+K^-)/\Gamma(K^-\pi^+)$ Γ_{137}/Γ_{21}

VALUE	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the $K^*(892)^+$ are included.			
0.084 ± 0.019 OUR FIT			
0.16 ± 0.08 -0.06	70 ANJOS 91	E691	γ Be 80–240 GeV
			70 The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^*(892)^+K^-)/\Gamma(K^0\pi^+\pi^-)$ Γ_{137}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the $K^*(892)^+$ are included.				
0.064 ± 0.014 OUR FIT				
0.058 ± 0.014 OUR AVERAGE				
0.064 ± 0.018	23	AMMAR 91	CLEO	$e^+e^- \approx 10.5$ GeV
0.05 ± 0.02 ± 0.01	15	ALBRECHT 90C	ARG	$e^+e^- \approx 10$ GeV

 $\Gamma(K^0K^-\pi^+\text{nonresonant})/\Gamma(K^-\pi^+)$ Γ_{123}/Γ_{21}

VALUE	DOCUMENT ID	TECN	COMMENT
0.06 ± 0.06	71 ANJOS 91	E691	γ Be 80–240 GeV
			71 The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^0K^+\pi^-)/\Gamma(K^-\pi^+)$ Γ_{124}/Γ_{21}

VALUE	DOCUMENT ID	TECN	COMMENT
0.123 ± 0.025 OUR FIT			
0.10 ± 0.05	72 ANJOS 91	E691	γ Be 80–240 GeV
			72 The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^0K^+\pi^-)/\Gamma(K^0\pi^+\pi^-)$ Γ_{124}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.093 ± 0.018 OUR FIT				
0.096 ± 0.020	55	AMMAR 91	CLEO	$e^+e^- \approx 10.5$ GeV

$\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma(K^- \pi^+)$ Γ_{138}/Γ_{21}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.00^{+0.04}_{-0.00}$	73 ANJOS	91 E691	γ Be 80-240 GeV

⁷³The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^*(892)^0 \bar{K}^0)/\Gamma(\bar{K}^0 \pi^+ \pi^-)$ Γ_{138}/Γ_{23}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.015	90	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV

 $\Gamma(K^*(892)^- K^+)/\Gamma(K^- \pi^+)$ Γ_{139}/Γ_{21}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.00^{+0.03}_{-0.00}$	74 ANJOS	91 E691	γ Be 80-240 GeV

⁷⁴The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^*(892)^- K^+)/\Gamma(\bar{K}^0 \pi^+ \pi^-)$ Γ_{139}/Γ_{23}

Unseen decay modes of the $K^*(892)^-$ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.034 ± 0.019	12	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV

 $\Gamma(\bar{K}^0 K^+ \pi^- \text{ nonresonant})/\Gamma(K^- \pi^+)$ Γ_{127}/Γ_{21}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.10^{+0.06}_{-0.05}$	75 ANJOS	91 E691	γ Be 80-240 GeV

⁷⁵The factor 100 at the top of column 2 of Table I of ANJOS 91 should be omitted.

 $\Gamma(K^+ K^- \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{128}/Γ_{39}

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.029 ± 0.006 OUR AVERAGE				
0.0314 ± 0.010	89	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV
$0.028^{+0.008}_{-0.007}$		ANJOS	91 E691	γ Be 80-240 GeV

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{140}/Γ

Unseen decay modes of the ϕ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
0.0026 ± 0.0007	76 BARLAG	92C ACCM	π^- Cu 230 GeV

⁷⁶BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(\phi \pi^+ \pi^-)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{140}/Γ_{39}

Unseen decay modes of the ϕ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$0.0076^{+0.0066}_{-0.0049}$	3	77 ANJOS	91 E691	γ Be 80-240 GeV

⁷⁷This ANJOS 91 result is inconsistent with the higher-statistics result of AMMAR 91 on $\phi \rho^0$.

 $\Gamma(\phi \rho^0)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{141}/Γ_{39}

Unseen decay modes of the ϕ are included.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.024 ± 0.006	34	78 AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV

⁷⁸The AMMAR 91 $\phi \pi^+ \pi^-$ events are consistent with being entirely $\phi \rho^0$.

 $\Gamma(K^*(892)^0 K^- \pi^+ + \text{c.c.})/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{142}/Γ_{39}

Unseen decay modes of the $K^*(892)^0$ are included.

VALUE	DOCUMENT ID	TECN	COMMENT
$0.010^{+0.016}_{-0.010}$	ANJOS	91 E691	γ Be 80-240 GeV

 $\Gamma(K^*(892)^0 \bar{K}^*(892)^0)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{143}/Γ_{39}

Unseen decay modes of the $K^*(892)^0$ and $\bar{K}^*(892)^0$ are included.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$0.036^{+0.020}_{-0.016}$		11	ANJOS	91 E691	γ Be 80-240 GeV
<0.033	90	79	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV

⁷⁹A corrected value (G. Moneti, private communication).

 $\Gamma(K^+ K^- \pi^+ \pi^- \text{ non-}\phi)/\Gamma_{\text{total}}$ Γ_{133}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0017 ± 0.0005	80 BARLAG	92C ACCM	π^- Cu 230 GeV

⁸⁰BARLAG 92C computes the branching fraction using topological normalization.

 $\Gamma(K^+ K^- \pi^+ \pi^- \text{ nonresonant})/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{134}/Γ_{39}

VALUE	DOCUMENT ID	TECN	COMMENT
$0.001^{+0.011}_{-0.001}$	ANJOS	91 E691	γ Be 80-240 GeV

 $\Gamma(K^+ K^- \pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$ Γ_{135}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0031 ± 0.0020	81 BARLAG	92C ACCM	π^- Cu 230 GeV

⁸¹BARLAG 92C computes the branching fraction using topological normalization.

Rare or forbidden modes

 $\Gamma(K^+ \pi^-)/\Gamma(K^- \pi^+)$ Γ_{144}/Γ_{21}

The measurements here cannot distinguish between doubly Cabibbo suppressed decay and/or $D^0 \bar{D}^0$ mixing.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$0.0077 \pm 0.0025 \pm 0.0025$		19	CINABRO	94 CLEO	$e^+ e^- \approx \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.011	90	AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV
<0.015	90	2	ANJOS	88C E691 Photoproduction

 $\Gamma(K^+ \pi^- \text{ (via } \bar{D}^0))/\Gamma(K^- \pi^+)$ Γ_{145}/Γ_{21}

This is a $D^0 \bar{D}^0$ mixing limit.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<0.0037	90	1	82 ANJOS	88C E691	Photoproduction
<0.014	90		ALBRECHT	87K ARG	$e^+ e^- 10$ GeV
<0.04	90		ABACHI	86D HRS	$e^+ e^- 29$ GeV
<0.07	90	0	82 BAILEY	86 ACCM	π^- Be fixed target
<0.11	90	2	ALBRECHT	85F ARG	$e^+ e^- 10$ GeV
<0.081	90		83 YAMAMOTO	85 DLCO	$e^+ e^- 29$ GeV
<0.23	90		83 ALTHOFF	84B TASS	$e^+ e^- 34.4$ GeV
<0.11	90		83 AVERY	80 SPEC	$\gamma N \rightarrow D^{*+}$
<0.16	90		83 FELDMAN	77B MRK1	$D^{*+} \rightarrow D^0 \pi^+$
<0.18	90		83 GOLDHABER	77 MRK1	

⁸²This measurement actually comes from combining results on $K^{\pm} \pi^{\mp} \pi^+ \pi^-$ and $K^{\pm} \pi^{\mp} \pi^0$ modes. See also the data block on $|m_{D_1^0} - m_{D_2^0}|$ near the beginning of the D^0 Listings.

⁸³Results given as $\Gamma(K^+ \pi^-)/[\Gamma(K^- \pi^+) + \Gamma(K^+ \pi^-)]$ but do not change significantly for our denominator.

 $\Gamma(K^+ \pi^+ \pi^- \pi^-)/\Gamma(K^- \pi^+ \pi^+ \pi^-)$ Γ_{146}/Γ_{39}

Doubly Cabibbo suppressed.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<0.018	90		AMMAR	91 CLEO	$e^+ e^- \approx 10.5$ GeV
<0.018	90	5	ANJOS	88C E691	Photoproduction

 $\Gamma(\mu^- \text{ anything (via } \bar{D}^0))/\Gamma(\mu^+ \text{ anything})$ Γ_{147}/Γ_2

This is a $D^0 \bar{D}^0$ mixing limit. See the somewhat better limit above on $D^0 \rightarrow K^+ \pi^-$ (via \bar{D}^0).

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0056	90	LOUIS	86 SPEC	π^- W 225 GeV
<0.012	90	BENVENUTI	85 CNTR	μ C, 200 GeV
<0.044	90	BODEK	82 SPEC	π^- , p Fe $\rightarrow D^0$

 $\Gamma(e^+ e^-)/\Gamma_{\text{total}}$ Γ_{148}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-4}$	90		ADLER	88 MRK3	$e^+ e^- 3.77$ GeV
$<1.7 \times 10^{-4}$	90	7	84 ALBRECHT	88G ARG	$e^+ e^- 10$ GeV
$<2.2 \times 10^{-4}$	90	8	85 HAAS	88 CLEO	$e^+ e^- 10$ GeV

⁸⁴The branching ratios are normalized to $B(D^0 \rightarrow K^- \pi^+)$ using ADLER 88C.

⁸⁵The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88C.

 $\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{149}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-5}$	90		LOUIS	86 SPEC	π^- W 225 GeV
$<7.0 \times 10^{-5}$	90	3	86 ALBRECHT	88C ARG	$e^+ e^- 10$ GeV
$<3.4 \times 10^{-4}$	90		AUBERT	85 EMC	Deep Inelast. $\mu^- N$

⁸⁶The branching ratios are normalized to $B(D^0 \rightarrow K^- \pi^+)$, using ADLER 88C.

 $\Gamma(\bar{K}^0 e^+ e^-)/\Gamma_{\text{total}}$ Γ_{150}/Γ

A test for the $\Delta C = 1$ weak neutral current. Allowed by first-order weak interaction combined with electromagnetic interaction.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0017	90	ADLER	89C MRK3	$e^+ e^- 3.77$ GeV

Meson Full Listings

D^0

$\Gamma(\rho^0 e^+ e^-)/\Gamma_{total}$
 A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 4.5 \times 10^{-4}$	90	2	87 HAAS	88 CLEO	$e^+ e^- 10$ GeV

⁸⁷ The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88c.

$\Gamma(\rho^0 \mu^+ \mu^-)/\Gamma_{total}$
 A test for the $\Delta C = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 8.1 \times 10^{-4}$	90	5	88 HAAS	88 CLEO	$e^+ e^- 10$ GeV

⁸⁸ The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88c.

$\Gamma(\mu^+ e^-)/\Gamma_{total}$
 A test of lepton family number conservation.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 1.0 \times 10^{-4}$	90	4	89 ALBRECHT	88G ARG	$e^+ e^- 10$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

$< 2.7 \times 10^{-4}$	90	9	90 HAAS	88 CLEO	$e^+ e^- 10$ GeV
$< 1.2 \times 10^{-4}$	90		BECKER	87C MRK3	$e^+ e^- 3.77$ GeV
$< 2 \times 10^{-4}$	90		PALKA	87 SILI	200 GeV πp
$< 21 \times 10^{-4}$	90	0	91 RILES	87 MRK2	$e^+ e^- 29$ GeV

⁸⁹ The branching ratios are normalized to $B(D^0 \rightarrow K^- \pi^+)$ using ADLER 88c.

⁹⁰ The branching ratios are normalized to $D^0 \rightarrow K^- \pi^+$, $D^+ \rightarrow K^- \pi^+ \pi^+$, and $D^{*+} \rightarrow D^0 \pi^+$ using ADLER 88c.

⁹¹ RILES 87 assumes $B(D \rightarrow K\pi) = 3.0\%$ and has production model dependency.

D^0 PRODUCTION CROSS SECTION AT $\psi(3770)$

A compilation of the cross sections for the direct production of D^0 mesons at or near the $\psi(3770)$ peak in $e^+ e^-$ production.

VALUE (nanobarns)	DOCUMENT ID	TECN	COMMENT
-------------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

$5.8 \pm 0.5 \pm 0.6$	92 ADLER	88C MRK3	$e^+ e^- 3.768$ GeV
7.3 ± 1.3	93 PARTRIDGE	84 CBAL	$e^+ e^- 3.771$ GeV
$8.00 \pm 0.95 \pm 1.21$	94 SCHINDLER	80 MRK2	$e^+ e^- 3.771$ GeV
11.5 ± 2.5	95 PERUZZI	77 MRK1	$e^+ e^- 3.774$ GeV

⁹² This measurement compares events with one detected D to those with two detected D mesons, to determine the absolute cross section. ADLER 88c find the ratio of cross sections (neutral to charged) to be $1.36 \pm 0.23 \pm 0.14$.

⁹³ This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. PARTRIDGE 84 measures 6.4 ± 1.15 nb for the cross section. We take the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and we assume that the $\psi(3770)$ is an isosinglet to evaluate the cross sections. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

⁹⁴ This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. SCHINDLER 80 assume the phase space division of neutral and charged D mesons in $\psi(3770)$ decay to be 1.33, and that the $\psi(3770)$ is an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction.

⁹⁵ This measurement comes from a scan of the $\psi(3770)$ resonance and a fit to the cross section. The phase space division of neutral and charged D mesons in $\psi(3770)$ decay is taken to be 1.33, and $\psi(3770)$ is assumed to be an isosinglet. The noncharm decays (e.g. radiative) of the $\psi(3770)$ are included in this measurement and may amount to a few percent correction. We exclude this measurement from the average because of uncertainties in the contamination from τ lepton pairs. Also see RAPIDIS 77.

$D^0 \bar{D}^0$ DECAY ASYMMETRY PARAMETER

$[\Gamma(D^0 \rightarrow K^+ K^-) - \Gamma(\bar{D}^0 \rightarrow K^+ K^-)] / \text{SUM}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.45	90	96 ANJOS	91D E691	Photoproduction

⁹⁶ ANJOS 91D is a limit on the time-independent asymmetry for direct CP violation.

D^0 REFERENCES

ALBRECHT 94	PL B324 249	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
CINABRO 94	PRL 72 1406	+Henderson, Liu, Saulnier+	(CLEO Collab.)
FRABETTI 94	PL B321 295	+Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI 94C	PL B323 459	+Cheung, Cumalat+	(FNAL E687 Collab.)
PDG 94	PR D50 1173	Montanet+	(CERN, LBL, BOST, IFIC+)
AKERIB 93	PRL 71 3070	+Barish, Chadha, Chan+	(CLEO Collab.)
ALBRECHT 93D	PL B308 435	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
ANJOS 93	PR D48 56	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BEAN 93C	PL B317 647	+Gronberg, Kutschke, Menary+	(CLEO Collab.)
FRABETTI 93I	PL B315 203	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
KODAMA 93B	PL B313 260	+Ushida, Mokhtarani+	(FNAL E653 Collab.)
PROCARIO 93B	PR D48 4007	+Yang, Akerib, Barish+	(CLEO Collab.)
SELEN 93	PRL 71 1973	+Sadof, Ammar, Ball+	(CLEO Collab.)
ADAMOVICH 92B	PL B280 163	+Alexandrov, Antnor+	(CERN WA82 Collab.)
ALBRECHT 92P	ZPHY C56 7	+Cronstrom, Ehrlichmann+	(ARGUS Collab.)
ANJOS 92B	PR D46 R1	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
ANJOS 92C	PR D46 1941	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BARLAG 92C	ZPHY C55 383	+Becker, Bozek, Boehringer+	(ACCMOR Collab.)
Also 92D	ZPHY C48 29	+Barlag, Becker, Boehringer, Bosman+	(ACCMOR Collab.)
COFFMAN 90B	PR D45 2196	+DeJongh, Dubois, Eigen+	(Mark III Collab.)
Also 90P	PRL 64 2615	+Adler, Blaylock, Bolton+	(Mark III Collab.)
FRABETTI 92	PL B281 167	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
FRABETTI 92B	PL B286 195	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
FRABETTI 91B	ZPHY C50 11	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
AMMAR 91	PR D44 3383	+Baringer, Coppage, Davis+	(CLEO Collab.)
ANJOS 91	PR D43 R635	+Appel, Bean, Bracker+	(FNAL-TPS Collab.)
ANJOS 91D	PR D44 R3371	+Appel, Bean, Bracker+	(FNAL-TPS Collab.)
BAI 91	PRL 66 1011	+Bolton, Brown, Bunnell+	(Mark III Collab.)
COFFMAN 91	PL B263 135	+DeJongh, Dubois, Eigen, Hitlin+	(Mark III Collab.)
CRAWFORD 91B	PR D44 3394	+Filton, Gan, Jensen+	(CLEO Collab.)
DECAMP 91J	PL B266 218	+Deschamps, Goy, Lees+	(LEPH Collab.)
FRABETTI 91	PL B263 584	+Bogart, Cheung, Culy+	(FNAL E687 Collab.)
KINOSHITA 91	PR D43 2836	+Pipkin, Procario, Wilson+	(CLEO Collab.)
KODAMA 91	PRL 66 1819	+Ushida, Mokhtarani, Paolone+	(FNAL E653 Collab.)
ALBRECHT 90C	ZPHY C46 9	+Glaeser, Harder, Krueger+	(ARGUS Collab.)
ALEXANDER 90	PRL 65 1184	+Artuso, Bebek, Berkelman+	(CLEO Collab.)
ALEXANDER 90B	PRL 65 1531	+Artuso, Bebek, Berkelman+	(CLEO Collab.)
ALVAREZ 90	ZPHY C47 539	+Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ANJOS 90D	PR D42 2414	+Appel, Bean, Bracker+	(FNAL E691 Collab.)
BARLAG 90C	ZPHY C46 563	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
ADLER 89	PRL 62 1821	+Becker, Blaylock, Bolton+	(Mark III Collab.)
ADLER 89C	PR D40 906	+Bai, Becker, Blaylock, Bolton+	(Mark III Collab.)
ALBRECHT 89D	ZPHY C43 181	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
ANJOS 89F	PRL 62 1587	+Appel, Bean, Bracker, Browder+	(FNAL E691 Collab.)
ABACHI 88	PL B205 411	+Akerlof, Baringer+	(HRS Collab.)
ADLER 88	PR D37 2023	+Becker, Blaylock+	(Mark III Collab.)
ADLER 88C	PRL 60 89	+Becker, Blaylock+	(Mark III Collab.)
ALBRECHT 88G	PL B209 380	+Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 88I	PL B210 267	+Boeckmann, Glaeser+	(ARGUS Collab.)
AMENDOLIA 88	EPL 5 407	+Bagliesi, Batignani+	(NA1 Collab.)
ANJOS 88C	PRL 60 1239	+Appel+	(FNAL E691 Collab.)
BORTOLETTO 88	PR D37 1719	+Goldberg, Horwitz, Mestayer, Moneti+	(CLEO Collab.)
Also 89D	PR D39 1471 erratum		
CUMALAT 88	PL B210 253	+Shipbaugh, Binsley+	(E-400 Collab.)
HAS 88	PRL 60 1614	+Hempstead, Jensen+	(CLEO Collab.)
RAAB 88	PR D37 2391	+Anjos, Appel, Bracker+	(FNAL E691 Collab.)
ADAMOVICH 87	EPL 4 887	+Alexandrov, Bolta+	(Photon Emulsion Collab.)
ADLER 87	PL B196 107	+Becker, Blaylock, Bolton+	(Mark III Collab.)
AGUILAR... 87D	PL B193 140	+Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also 88B	ZPHY C40 321	+Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
AGUILAR... 87E	ZPHY C36 551	+Aguilar-Benitez, Allison+	(LEBC-EHS Collab.)
Also 87F	ZPHY C36 559	+Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
AGUILAR... 88	ZPHY C38 520 erratum		
ALBRECHT 87E	ZPHY C33 359	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 87K	PL B199 447	+Andam, Binder, Boeckmann+	(ARGUS Collab.)
BARLAG 87B	ZPHY C37 17	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
BECKER 87C	PL B193 147	+Blaylock, Bolton, Brown+	(Mark III Collab.)
Also 87D	PL B196 590 erratum	+Becker, Blaylock, Bolton+	(Mark III Collab.)
CSORNA 87	PL B191 318	+Mestayer, Farnini, Ward+	(CLEO Collab.)
PALKA 87	PL B189 238	+Bailey, Becker, Belau+	(ACCMOR Collab.)
RILES 87	PR D35 2914	+Dorfan, Abrams, Amidei+	(Mark II Collab.)
ABACHI 86D	PL B182 101	+Akerlof, Baringer, Ballam+	(HRS Collab.)
ABE 86	PR D33 1	-	(SLAC Hybrid Facility Photon Collab.)
BAILEY 86	ZPHY C30 51	+Belau, Boehringer, Bosman+	(ACCMOR Collab.)
BEBEK 86	PRL 56 1893	+Berkelman, Blucher, Cassel+	(CLEO Collab.)
GLADNEY 86	PR D34 2601	+Jaros, Ong, Barklow+	(Mark II Collab.)
LOUIS 86	PL B56 1027	+Adolphen, Alexandrov+	(PRIM, CHIC, ISU)
USHIDA 86B	PRL 56 1771	+Kondo+	(AICH, FNAL, KOBE, SEOU, MCGI+)
ALBRECHT 85B	PL 158B 525	+Binder, Harder, Philipp+	(ARGUS Collab.)
ALBRECHT 85F	PL 150B 235	+Binder, Harder, Philipp+	(ARGUS Collab.)
AUBERT 85	PL 155B 461	+Bassompierre, Becks, Benchouk+	(EMC Collab.)
BAILEY 85	ZPHY C28 357	+Belau, Boehringer, Bosman+	(ABCCMR Collab.)
BALTRUSAIT... 85B	PRL 54 1976	+Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BALTRUSAIT... 85E	PRL 55 150	+Baltrusaitis, Becker, Blaylock, Brown+	(Mark III Collab.)
BENVENUTI 85	PL 158B 531	+Bollini, Bruni, Camporesi+	(BCDMS Collab.)
YAMAMOTO 85	PRL 54 522	+Yamamoto, Atwood, Bailion+	(DELCO Collab.)
ADAMOVICH 84B	PL 140B 123	+Alexandrov, Bravo+	(CERN WA82 Collab.)
ALTHOFF 84B	PL 138B 317	+Braunschweig, Kirschkink+	(TASSO Collab.)
DERRICK 84	PRL 53 1971	+Fernandez, Fries, Hyman+	(HRS Collab.)
PARTRIDGE 84	Thesis CALT-68-1150		(Crystal Ball Collab.)
BAILEY 83B	PL 132B 237	+ (UCSB, CARL, COLO, FNAL, TNTO, OKLA, CNRC)	
BODEK 82	PL 113B 82	+Bardley, Becker, Biana+	(ACCMOR Collab.)
FIORINO 81	LNC 30 166	+Breedon+ (ROCH, CIT, CHIC, FNAL, STAN)	
SCHINDLER 81	PR D24 78	+ (Photon-Emulsion and Omega-Photon Collab.)	
TRILLING 81	PR D24 78	+Alam, Boyarski, Breidenbach+	(Mark II Collab.)
ASTON 80E	PL 94B 113	+ (LBL, UCB)	
AVERY 80	PRL 44 1309	+ (BONN, CERN, EPOL, GLAS, LANC, MCHS+)	
SCHINDLER 80	PR D21 2716	+Wiss, Butler, Glasing+ (ILL, FNAL, COLU)	
ZHOLENTZ 80	PL 96B 214	+Sigrist, Alam, Boyarski+	(Mark II Collab.)
Also 81	SJNP 34 814	+Kurdadze, Lechuk, Mishnev+ (NOVO)	
Translated from YAF 34 1471.		+Zholentz, Kurdadze, Lechuk+ (NOVO)	
ABRAMS 79D	PRL 43 481	+Alam, Blocker, Boyarski+	(SLAC, LBL)
ATIYA 79	PRL 43 414	+Holmes, Knapp, Lee+	(COLU, ILL, FNAL)
BALAY 78C	PRL 41 73	+Caroumbalis, French, Hibbs, Hyllton+	(COLU, BNL)
VUILLEMIN 78	PRL 41 1149	+Feldman, Feller+	(LBL, SLAC, NWES, HAWA)
FELDMAN 77B	PRL 38 1313	+Peruzzi, Piccolo, Abrams, Alam+	(SLAC, LBL)
GOLDBERGER 77	PL 69B 503	+Wiss, Abrams, Alam+	(LBL, SLAC)
PERUZZI 77	PRL 39 1301	+Piccolo, Feldman+	(SLAC, LBL, NWES, HAWA)
PICCOLO 77	PL 70B 260	+Peruzzi, Luth, Nguyen, Wiss, Abrams+	(SLAC, LBL)
RAPIDIS 77	PRL 39 526	+Gobbi, Luke, Barbaro-Gallieri+	(Mark I Collab.)
GOLDBERGER 76	PRL 37 255	+Pierre, Abrams, Alam+	(LBL, SLAC)

OTHER RELATED PAPERS

MORRISON 89	ARNPS 39 183	-Witherell	(UCSB)
-------------	--------------	------------	--------

See key on page 1343

Meson Full Listings
 $D^*(2007)^0, D^*(2010)^\pm$

$D^*(2007)^0$	
$I(J^P) = \frac{1}{2}(1^-)$ I, J, P need confirmation.	
J consistent with 1, value 0 ruled out (NGUYEN 77).	
$D^*(2007)^0$ MASS	
The fit includes $D^\pm, D^0, D_S^\pm, D^{*\pm}, D^{*0}$, and $D_S^{*\pm}$ mass and mass difference measurements.	
VALUE (MeV)	DOCUMENT ID TECN COMMENT
2006.7 ± 0.5 OUR FIT	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
2006 ± 1.5	¹ GOLDHABER 77 MRK1 e^+e^-
¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 .	

$m_{D^*(2007)^0} - m_{D^0}$	
The fit includes $D^\pm, D^0, D_S^\pm, D^{*\pm}, D^{*0}$, and $D_S^{*\pm}$ mass and mass difference measurements.	
VALUE (MeV)	EVTS DOCUMENT ID TECN COMMENT
142.12 ± 0.07 OUR FIT	
142.12 ± 0.05 ± 0.05	1176 ± 50 BORTOLETTO92B CLE2 $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •	
142.2 ± 2.0	³ SADROZINSKI 80 CBAL $D^{*0} \rightarrow D^+\pi^0$
142.7 ± 1.7	² GOLDHABER 77 MRK1 e^+e^-
² From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 .	

$D^*(2007)^0$ WIDTH	
VALUE (MeV)	CL% DOCUMENT ID TECN COMMENT
< 2.1	90 ³ ABACHI 88B HRS $D^{*0} \rightarrow D^+\pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
< 5	GOLDHABER 76B MRK1 $e^+e^- \rightarrow D^*D^*$
³ Assuming $m_{D^0} = 2007.2 \pm 2.1$ MeV/ c^2 .	

$D^*(2007)^0$ DECAY MODES	
$\bar{D}^*(2007)^0$ modes are charge conjugates of modes below.	
Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0\pi^0$	(63.6 ± 2.8) %
Γ_2 $D^0\gamma$	(36.4 ± 2.8) %

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 2 measurements and one constraint to determine 2 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

$$x_2 \begin{vmatrix} & -100 \\ & x_1 \end{vmatrix}$$

$D^*(2007)^0$ BRANCHING RATIOS	
$\Gamma(D^0\pi^0) / [\Gamma(D^0\pi^0) + \Gamma(D^0\gamma)]$	$\Gamma_1 / (\Gamma_1 + \Gamma_2)$
VALUE	EVTS DOCUMENT ID TECN COMMENT
0.636 ± 0.028 OUR FIT	
0.636 ± 0.023 ± 0.033	1097 ± 59 BUTLER 92 CLE2 $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •	
0.37 ± 0.08 ± 0.08	ADLER 88D MRK3 e^+e^-
0.47 ± 0.23	LOW 87 HRS 29 GeV e^+e^-
0.53 ± 0.13	BARTEL 85G JADE e^+e^- , hadrons
0.47 ± 0.12	COLES 82 MRK2 e^+e^-
0.45 ± 0.15	GOLDHABER 77 MRK1 e^+e^-
$\Gamma(D^0\gamma) / [\Gamma(D^0\pi^0) + \Gamma(D^0\gamma)]$	$\Gamma_2 / (\Gamma_1 + \Gamma_2)$
VALUE	EVTS DOCUMENT ID TECN COMMENT
0.364 ± 0.028 OUR FIT	
0.364 ± 0.023 ± 0.033	621 ± 52 BUTLER 92 CLE2 $e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •	

$D^*(2007)^0$ REFERENCES	
BORTOLETTO 92B PRL 69 2046	+Brown, Dominick+ (CLEO Collab.)
BUTLER 92 PRL 69 2041	+Fu, Kalbfleisch+ (CLEO Collab.)
ABACHI 88B PL B212 533	+Akerlof+ (ANL, IND, MICH, PURD, LBL)
ADLER 88D PL B208 152	+Becker+ (Mark III Collab.)
LOW 87 PL B183 232	+Abachi, Akerlof, Baringer+ (HRS Collab.)
BARTEL 85G PL 161B 197	+Dietrich, Ambrus+ (JADE Collab.)
COLES 82 PR D26 2190	+Abrams, Blocker, Blondel+ (LBL, SLAC)
SADROZINSKI 80 Madison Conf. 681	+ (PRIN, CIT, HARV, SLAC, STAN)
GOLDHABER 77 PL 69B 503	+Wiss, Abrams, Alam+ (LBL, SLAC)
NGUYEN 77 PRL 39 262	+Wiss, Abrams, Alam, Boyarski+ (LBL, SLAC)
GOLDHABER 76B SLAC Conf. 379	Available as LBL-5534.
OTHER RELATED PAPERS	
KAMAL 92 PL B284 421	+Xu (ALBE)
TRILLING 81 PRPL 75 57	(LBL, UC/B)
FELDMAN 77C Banff Sum. Inst. 75	(SLAC)
GOLDHABER 76 PRL 37 255	+Pierre, Abrams, Alam+ (LBL, SLAC)

$D^*(2010)^\pm$	
$I(J^P) = \frac{1}{2}(1^-)$ I, J, P need confirmation.	
$D^*(2010)^\pm$ MASS	
The fit includes $D^\pm, D^0, D_S^\pm, D^{*\pm}, D^{*0}$, and $D_S^{*\pm}$ mass and mass difference measurements.	
VALUE (MeV)	DOCUMENT ID TECN CHG COMMENT
2010.0 ± 0.5 OUR FIT	
• • • We do not use the following data for averages, fits, limits, etc. • • •	
2008 ± 3	¹ GOLDHABER 77 MRK1 ± e^+e^-
2008.6 ± 1.0	² PERUZZI 77 MRK1 ± e^+e^-
¹ From simultaneous fit to $D^*(2010)^+, D^*(2007)^0, D^+$, and D^0 ; not independent of FELDMAN 77B mass difference below.	
² PERUZZI 77 mass not independent of FELDMAN 77B mass difference below and PERUZZI 77 D^0 mass value.	

$m_{D^*(2010)^+} - m_{D^+}$	
The fit includes $D^\pm, D^0, D_S^\pm, D^{*\pm}, D^{*0}$, and $D_S^{*\pm}$ mass and mass difference measurements.	
VALUE (MeV)	EVTS DOCUMENT ID TECN COMMENT
140.64 ± 0.09 OUR FIT	
140.64 ± 0.08 ± 0.06	620 ± 42 BORTOLETTO92B CLE2 $e^+e^- \rightarrow$ hadrons

$m_{D^*(2010)^+} - m_{D^0}$	
The fit includes $D^\pm, D^0, D_S^\pm, D^{*\pm}, D^{*0}$, and $D_S^{*\pm}$ mass and mass difference measurements.	

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
145.42 ± 0.05 OUR FIT				
145.42 ± 0.04 OUR AVERAGE				
145.39 ± 0.06 ± 0.03		BARLAG 92B ACCM	π^-	230 GeV
145.40 ± 0.05 ± 0.10		ABACHI 88B HRS	$D^{*\pm} \rightarrow D^0\pi^\pm$	
145.46 ± 0.07 ± 0.03		ALBRECHT 85F ARG	$D^{*\pm} \rightarrow D^0\pi^\pm$	
145.8 ± 1.5	16	AHLEN 83 HRS	$D^{*+} \rightarrow D^0\pi^+$	
145.1 ± 1.8	12	BAILEY 83 SPEC	$D^{*\pm} \rightarrow D^0\pi^\pm$	
145.5 ± 0.3	28	BAILEY 83 SPEC	$D^{*\pm} \rightarrow D^0\pi^\pm$	
145.1 ± 0.5	14	BAILEY 83 SPEC	$D^{*\pm} \rightarrow D^0\pi^\pm$	
145.5 ± 0.5	14	YELTON 82 MRK2	29 $e^+e^- \rightarrow K^-\pi^+$	
145.5 ± 0.3	60	FITCH 81 SPEC	π^-A	
145.2 ± 0.6	2	BLIETSCHAU 79 BEBC	νp	
145.3 ± 0.5	30	FELDMAN 77B MRK1	$D^{*+} \rightarrow D^0\pi^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
145.5 ± 0.2	115	³ ALEXANDER 91B OPAL	$D^{*\pm} \rightarrow D^0\pi^\pm$	
145.30 ± 0.06		³ DECAMP 91J ALEP	$D^{*\pm} \rightarrow D^0\pi^\pm$	
~ 145.5		AVERY 80 SPEC	γA	
³ Systematic error not evaluated.				

$m_{D^*(2010)^+} - m_{D^*(2007)^0}$	
VALUE (MeV)	DOCUMENT ID TECN COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •	
2.6 ± 1.8	⁴ PERUZZI 77 MRK1 e^+e^-
⁴ Not independent of FELDMAN 77B mass difference above, PERUZZI 77 D^0 mass, and GOLDHABER 77 $D^*(2007)^0$ mass.	

Meson Full Listings

$D^*(2010)^\pm, D_1(2420)^0$

$D^*(2010)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
<0.131	90	110	BARLAG	92B ACCM	π^- 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.1	90		ABACHI	88B HRS	$D^{*\pm} \rightarrow D^0 \pi^\pm$
<2.2			YELTON	82 MRK2	$e^+ e^- \rightarrow K^- \pi^+ \pi^-$
<2.0	90	30	FELDMAN	77B MRK1	$D^{*+} \rightarrow D^0 \pi^+$

$D^*(2010)^\pm$ DECAY MODES

$D^*(2010)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^0 \pi^+$	(68.1 ± 1.3) %
Γ_2 $D^+ \pi^0$	(30.8 ± 0.8) %
Γ_3 $D^+ \gamma$	(1.1 ^{+1.4} _{-0.7}) %

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 3 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 0.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-24	
x_3	-81	-38
	x_1	x_2

$D^*(2010)^+$ BRANCHING RATIOS

$\Gamma(D^0 \pi^+)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.681 ± 0.013 OUR FIT				
0.681 ± 0.010 ± 0.013	BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.57 ± 0.04 ± 0.04	ADLER	88D MRK3	$e^+ e^-$	
0.44 ± 0.10	COLES	82 MRK2	$e^+ e^-$	
0.6 ± 0.15	⁵ GOLDHABER	77 MRK1	$e^+ e^-$	

⁵ Assuming that isospin is conserved in the decay.

$\Gamma(D^+ \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.308 ± 0.008 OUR FIT				
0.308 ± 0.004 ± 0.008	BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.26 ± 0.02 ± 0.02	ADLER	88D MRK3	$e^+ e^-$	
0.34 ± 0.07	COLES	82 MRK2	$e^+ e^-$	

$\Gamma(D^+ \gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.011 ± 0.014 OUR FIT				
0.011 ± 0.014 ± 0.016	BUTLER	92 CLE2	$e^+ e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17 ± 0.05 ± 0.05	ADLER	88D MRK3	$e^+ e^-$	
0.22 ± 0.12	⁶ COLES	82 MRK2	$e^+ e^-$	

⁶ Not independent of $\Gamma(D^0 \pi^+)/\Gamma_{\text{total}}$ and $\Gamma(D^+ \pi^0)/\Gamma_{\text{total}}$ measurement.

$D^*(2010)^\pm$ REFERENCES

BARLAG	92B	PL B278 480	+Becker, Bozek+	(ACCMOR Collab.)
BORTOLETTO	92B	PRL 69 2046	+Brown, Dominick+	(CLEO Collab.)
BUTLER	92	PRL 69 2041	+Fu, Kaibfleisch+	(CLEO Collab.)
ALEXANDER	91B	PL B262 341	+Allison, Allport, Anderson, Arcelli+	(OPAL Collab.)
DECAMP	91J	PL B266 218	+Deschizeaux, Goy, Lees+	(ALEPH Collab.)
ABACHI	88B	PL B212 533	+Akerlof+	(ANL, IND, MICH, PURD, LBL)
ADLER	88D	PL B208 152	+Becker+	(Mark III Collab.)
ALBRECHT	85F	PL 150B 235	+Binder, Harder, Philipp+	(ARGUS Collab.)
AHLEN	83	PRL 51 1147	+Akerlof+	(ANL, IND, LBL, MICH, PURD, SLAC)
BAILEY	83	PL 132B 230	+Bardsley+	(AMST, BRIS, CERN, CRAC, MPIM, OXF)
COLES	82	PR D26 2190	+Abrams, Blocker, Blondel+	(LBL, SLAC)
YELTON	82	PRL 49 430	+Feldman, Goldhaber+	(SLAC, LBL, UCB, HARV)
FITCH	81	PRL 46 761	+Devaux, Cavaglia, May+	(PRIN, SACL, TORI, BNL)
AVERY	80	PRL 44 1309	+Wiss, Butler, Gladding+	(ILL, FNAL, COLU)
BLIETSCHAU	79	PL 86B 108	+ (AACH, BONN, CERN, MPIM, OXF)	
FELDMAN	77B	PRL 38 1313	+Peruzzi, Piccolo, Abrams, Alam+	(SLAC, LBL)
GOLDHABER	77	PL 69B 503	+Wiss, Abrams, Alam+	(LBL, SLAC)
PERUZZI	77	PRL 39 1301	+Piccolo, Feldman+	(SLAC, LBL, NWES, HAWA)

OTHER RELATED PAPERS

KAMAL	92	PL B284 421	+Xu	(ALBE)
ALTHOFF	83C	PL 126B 493	+Fischer, Burkhardt+	(TASSO Collab.)
BEBEK	82	PRL 49 610	+	(HARV, OSU, ROCH, RUTG, SYRA, VAND+)
TRILLING	81	PRPL 75 57		(LBL, UCB)
PERUZZI	76	PRL 37 569	+Piccolo, Feldman, Nguyen, Wiss+	(SLAC, LBL)

$D_1(2420)^0$

$$I(J^P) = \frac{1}{2}(1^+)$$


I, J, P need confirmation.

Seen in $D^*(2010)^+ \pi^-$. $J^P = 1^+$ according to ALBRECHT 89B and ALBRECHT 89H.

$D_1(2420)^0$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2422.8 ± 3.2 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
2422 ± 2 ± 2	51 ± 18	FRABETTI	94B E687	γ Be $\rightarrow D^{*+} \pi^- X$
2428 ± 3 ± 2	279 ± 34	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
2414 ± 2 ± 5	171 ± 22	ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^{*+} \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2428 ± 8 ± 5	171 ⁺⁴³ ₋₅₈	ANJOS	89C TPS	γ N $\rightarrow D^{*+} \pi^- X$

WEIGHTED AVERAGE
2422.8 ± 3.2 (Error scaled by 1.6)



$D_1(2420)^0$ mass (MeV)

$D_1(2420)^0$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
18 ± 6 OUR AVERAGE				
15 ± 8 ± 4	51 ± 18	FRABETTI	94B E687	γ Be $\rightarrow D^{*+} \pi^- X$
23 ± 8 ± 10 6 ± 3	279 ± 34	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$
13 ± 6 ± 10 5	171 ± 22	ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^{*+} \pi^- X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
58 ± 14 ± 10	171 ⁺⁴³ ₋₅₈	ANJOS	89C TPS	γ N $\rightarrow D^{*+} \pi^- X$

$D_1(2420)^0$ DECAY MODES

$\bar{D}_1(2420)^0$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2010)^+ \pi^-$	seen
Γ_2 $D^+ \pi^-$	not seen

$D_1(2420)^0$ BRANCHING RATIOS

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
seen	ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^{*+} \pi^- X$	
seen	ANJOS	89C TPS	γ N $\rightarrow D^{*+} \pi^- X$	
$\Gamma(D^+ \pi^-)/\Gamma(D^*(2010)^+ \pi^-)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
<0.24	90	AVERY	90 CLEO	$e^+ e^- \rightarrow D^+ \pi^- X$

See key on page 1343

Meson Full Listings

 $D_1(2420)^0, D_J(2440)^\pm, D_2^*(2460)$ $D_1(2420)^0$ REFERENCES

FRABETTI	94B	PRL 72 324	+Cheung, Cumalat+	(FNAL E687 Collab.)
AVERY	90	PR D41 774	+Besson	(CLEO Collab.)
ALBRECHT	89B	PL B221 422	+Boeckmann+	(ARGUS Collab.) JP
ALBRECHT	89H	PL B232 398	+Glaser, Harder+	(ARGUS Collab.) JP
ANJOS	89C	PRL 62 1717	+Appel+	(FNAL E691 Collab.)

 $D_J(2440)^\pm$

$$I(J^P) = \frac{1}{2}(2^?)$$

I needs confirmation.

OMITTED FROM SUMMARY TABLE

Seen in $D^*(2007)^0 \pi^+$, $J^P = 0^+$ ruled out. $D_J(2440)^\pm$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
$2443 \pm 7 \pm 5$	190^{+77}_{-44}	ANJOS	89C TPS	$\gamma N \rightarrow D^0 \pi^+ \chi^0$

 $D_J(2440)^\pm$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
$41 \pm 19 \pm 8$	190^{+77}_{-44}	ANJOS	89C TPS	$\gamma N \rightarrow D^0 \pi^+ \chi^0$

 $D_J(2440)^\pm$ DECAY MODES $D_J^*(2440)^-$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D^*(2007)^0 \pi^+$	seen

 $D_J(2440)^\pm$ BRANCHING RATIOS

$\Gamma(D^*(2007)^0 \pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		ANJOS	89C TPS	$\gamma N \rightarrow D^0 \pi^+ \chi^0$	

 $D_J(2440)^\pm$ REFERENCES

ANJOS	89C	PRL 62 1717	+Appel+	(FNAL E691 Collab.)
-------	-----	-------------	---------	---------------------

 $D_2^*(2460)$

$$I(J^P) = \frac{1}{2}(2^+)$$

 $J^P = 2^+$ assignment strongly favored (ALBRECHT 89B). $D_2^*(2460)$ MASS

NEUTRAL MODE	VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2457.7 ± 1.9 OUR AVERAGE					
2453 ± 3 ± 2	128 ± 28		FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^+ \pi^- \chi$
2461 ± 3 ± 1	440 ± 97		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- \chi$
2455 ± 3 ± 5	337 ± 100		ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- \chi$
2459 ± 3 ± 2	153^{+42}_{-37}		ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- \chi$

CHARGED MODE

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT	
2456 ± 6 OUR AVERAGE	Error includes scale factor of 2.0.				
2453 ± 3 ± 2	185 ± 42		FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^0 \pi^+ \chi$
2469 ± 4 ± 6			ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ \chi$

 $m D_2^*(2460)^\pm - m D_2^*(2460)^0$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2 ± 5 OUR AVERAGE	Error includes scale factor of 1.4.		
0 ± 4	FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D \pi \chi$
14 ± 5 ± 8	ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ \chi$

 $D_2^*(2460)$ WIDTH

NEUTRAL MODE

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT	
21 ± 5 OUR AVERAGE	Includes data from the datablock that follows this one.				
25 ± 10 ± 5	128 ± 28		FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^+ \pi^- \chi$
$20^{+9}_{-12} \pm 9$	440 ± 97		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- \chi$
$15^{+13}_{-10} \pm 5$	337 ± 100		ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- \chi$
20 ± 10 ± 5	153^{+42}_{-37}		ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- \chi$

CHARGED MODE

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT	
23 ± 9 ± 5	185 ± 42		FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^0 \pi^+ \chi$

 $D_2^*(2460)$ DECAY MODES $\bar{D}_2^*(2460)$ modes are charge conjugates of modes below.

Mode	Fraction (Γ_i/Γ)
Γ_1 $D_2^*(2460)^0 \rightarrow D^+ \pi^-$	seen
Γ_2 $D_2^*(2460)^0 \rightarrow D^*(2010)^+ \pi^-$	seen
Γ_3 $D_2^*(2460)^\pm \rightarrow D^0 \pi^+$	seen

 $D_2^*(2460)$ BRANCHING RATIOS

$\Gamma(D^+ \pi^-)/\Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	337 ± 100		ALBRECHT	89B ARG	$e^+ e^- \rightarrow D^+ \pi^- \chi$	
seen			ANJOS	89C TPS	$\gamma N \rightarrow D^+ \pi^- \chi$	

$\Gamma(D^*(2010)^+ \pi^-)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
seen		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} \pi^- \chi$	
seen		ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^* \pi^- \chi$	

$\Gamma(D^+ \pi^-)/\Gamma(D^*(2010)^+ \pi^-)$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
2.4 ± 0.7 OUR AVERAGE					
2.3 ± 0.8			AVERY	90 CLEO	$e^+ e^-$
3.0 ± 1.1 ± 1.5			ALBRECHT	89H ARG	$e^+ e^- \rightarrow D^* \pi^- \chi$

$\Gamma(D^0 \pi^+)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
seen		ALBRECHT	89F ARG	$e^+ e^- \rightarrow D^0 \pi^+ \chi$	

 $D_2^*(2460)$ REFERENCES

FRABETTI	94B	PRL 72 324	+Cheung, Cumalat+	(FNAL E687 Collab.)
AVERY	90	PR D41 774	+Besson	(CLEO Collab.)
ALBRECHT	89B	PL B221 422	+Boeckmann+	(ARGUS Collab.) JP
ALBRECHT	89F	PL B231 208	+Glaser+	(ARGUS Collab.) JP
ALBRECHT	89H	PL B232 398	+Glaser, Harder+	(ARGUS Collab.) JP
ANJOS	89C	PRL 62 1717	+Appel+	(FNAL E691 Collab.)

Meson Full Listings

D_s^\pm

CHARMED, STRANGE MESONS ($C = S = \pm 1$)

$D_s^+ = c\bar{s}, D_s^- = \bar{c}s$, similarly for $D_s^{*\pm}$

D_s^\pm
was F^\pm

$$I(J^P) = 0(0^-)$$

The angular distributions of the decays of the ϕ and $\bar{K}^*(892)^0$ in the $\phi\pi^+$ and $K^+\bar{K}^*(892)^0$ modes strongly indicate that the spin is zero. The parity given is that expected of a $c\bar{s}$ ground state.


D_s^\pm MASS

The fit includes $D_s^\pm, D_s^0, D_s^{*\pm}, D^{*\pm}, D^{*0}$, and $D_s^{*\pm}$ mass and mass difference measurements. Measurements of the D_s^\pm mass with an error greater than 10 MeV are omitted from the fit and average. A number of early measurements giving $m > 2000$ MeV have been omitted altogether. They may be found in our 1990 edition (Phys. Lett. **B239**).

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1968.5 ± 0.7 OUR FIT	Error includes scale factor of 1.2.			
1969.0 ± 1.4 OUR AVERAGE	Error includes scale factor of 1.5. See the ideogram below.			
1967.0 ± 1.0 ± 1.0	54	BARLAG	90C ACCM	π^- Cu 230 GeV
1969.3 ± 1.4 ± 1.4		ALBRECHT	88 ARG	e^+e^- 9.4-10.6 GeV
1972.7 ± 1.5 ± 1.0	21	BECKER	87B SILI	200 GeV π, K, p
1972.4 ± 3.7 ± 3.7	27	BLAYLOCK	87 MRK3	e^+e^- 4.14 GeV
1963 ± 3 ± 3	30	DERRICK	85B HRS	e^+e^- 29 GeV
1970 ± 5 ± 5	104	CHEN	83C CLEO	e^+e^- 10.5 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1968.3 ± 0.7 ± 0.7	290	¹ ANJOS	88 E691	Photoproduction
1980 ± 15	6	USHIDA	86 EMUL	ν wideband
1973.6 ± 2.6 ± 3.0	163	ALBRECHT	85D ARG	e^+e^- 10 GeV
1948 ± 28 ± 10	65	AIHARA	84D TPC	e^+e^- 29 GeV
1975 ± 9 ± 10	49	ALTHOFF	84 TASS	e^+e^- 14-25 GeV
1975 ± 4	3	BAILEY	84 ACCM	hadron ⁺ Be → $\phi\pi^+X$

¹ ANJOS 88 enters the fit via $m_{D_s^\pm} - m_{D^\pm}$ (see below).

WEIGHTED AVERAGE
1969.0 ± 1.4 (Error scaled by 1.5)



D_s^\pm mass (MeV)

$$m_{D_s^\pm} - m_{D^\pm}$$

The fit includes $D_s^\pm, D_s^0, D_s^{*\pm}, D^{*\pm}, D^{*0}$, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
99.1 ± 0.6 OUR FIT	Error includes scale factor of 1.1.			
98.9 ± 0.7 OUR AVERAGE				
98.5 ± 1.5	555	CHEN	89 CLEO	e^+e^- 10.5 GeV
99.0 ± 0.8	290	ANJOS	88 E691	Photoproduction

D_s^\pm MEAN LIFE

Measurements with an error greater than 0.2×10^{-12} s are omitted from the average.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.467 ± 0.017 OUR AVERAGE				
0.475 ± 0.020 ± 0.007	900	FRABETTI	93F E687	γ Be, $D_s^+ \rightarrow \phi\pi^+$
0.33 $^{+0.12}_{-0.08}$ ± 0.03	15	ALVAREZ	90 NA14	$\gamma, D_s^+ \rightarrow \phi\pi^+$
0.469 $^{+0.102}_{-0.086}$	54	² BARLAG	90C ACCM	π^- Cu 230 GeV
0.50 ± 0.06 ± 0.03	104	FRABETTI	90 E687	γ Be, $\phi\pi^+$
0.56 $^{+0.13}_{-0.12}$ ± 0.08	144	ALBRECHT	88I ARG	e^+e^- 10 GeV
0.47 ± 0.04 ± 0.02	228	RAAB	88 E691	Photoproduction
0.33 $^{+0.10}_{-0.06}$	21	³ BECKER	87B SILI	200 GeV π, K, p
0.26 $^{+0.16}_{-0.09}$	6	USHIDA	86 EMUL	ν wideband
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.31 $^{+0.24}_{-0.20}$ ± 0.05	18	AVERILL	89 HRS	e^+e^- 29 GeV
0.48 $^{+0.06}_{-0.05}$ ± 0.02	99	ANJOS	87B E691	See RAAB 88
0.57 $^{+0.36}_{-0.26}$ ± 0.09	9	BRAUNSCH...	87 TASS	e^+e^- 35-44 GeV
0.47 ± 0.22 ± 0.05	141	CSORNA	87 CLEO	e^+e^- 10 GeV
0.35 $^{+0.24}_{-0.18}$ ± 0.09	17	JUNG	86 HRS	See AVERILL 89
0.32 $^{+0.30}_{-0.13}$	3	BAILEY	84 ACCM	hadron ⁺ Be → $\phi\pi^+X$
0.19 $^{+0.13}_{-0.07}$	4	USHIDA	83 EMUL	See USHIDA 86

² BARLAG 90C estimates the systematic error to be negligible.
³ BECKER 87B estimates the systematic error to be negligible.

D_s^\pm DECAY MODES

Branching fractions for modes below with a resonance in the final state include all the decay modes of the resonance. D_s^\pm modes are charge conjugates of the modes below.

Nearly all other modes are measured relative to the $\phi\pi^+$ mode. However, none of the determinations of the $\phi\pi^+$ branching fraction are direct measurements: all rely on calculated relations between D^+ and D_s^+ decay widths, on estimates of D_s^+ cross sections, or on other model-dependent assumptions. Thus a better determination of the $\phi\pi^+$ branching fraction could cause the other branching fractions to slide up or down, all together.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Inclusive modes		
Γ_1 K^- anything	(13 $^{+14}_{-12}$) %	
Γ_2 \bar{K}^0 anything + K^0 anything	(39 $^{+28}$) %	
Γ_3 K^+ anything	(20 $^{+18}_{-14}$) %	
Γ_4 non- $K\bar{K}$ anything	(64 $^{+17}$) %	
Γ_5 e^+ anything	< 20 %	CL=90%
Leptonic and semileptonic modes		
Γ_6 $\mu^+\nu_\mu$	(5.9 ± 2.2) × 10 ⁻³	S=1.1
Γ_7 $\phi\ell^+\nu_\ell$	[a] (1.88 ± 0.29) %	
Γ_8 $\phi e^+\nu_e$		
Γ_9 $\phi\mu^+\nu_\mu$		
Γ_{10} $\eta\mu^+\nu_\mu + \eta'(958)\mu^+\nu_\mu$	(7.4 ± 3.2) %	
Γ_{11} $\eta\mu^+\nu_\mu$		
Γ_{12} $\eta'(958)\mu^+\nu_\mu$	< 3.0 %	CL=90%
Hadronic modes with two K's (including from ϕ's)		
Γ_{13} $K^+\bar{K}^0$	(3.5 ± 0.7) %	
Γ_{14} $K^+K^-\pi^+$	(4.8 ± 0.7) %	
In the fit as $\frac{1}{2}\Gamma_{15} + \frac{2}{3}\Gamma_{16} + \Gamma_{17}$.		
Γ_{15} $\phi\pi^+$	(3.5 ± 0.4) %	
Γ_{16} $K^+\bar{K}^*(892)^0$	(3.3 ± 0.5) %	
Γ_{17} $K^+K^-\pi^+$ nonresonant	(8.7 ± 3.2) × 10 ⁻³	
Γ_{18} $K^0\bar{K}^0\pi^+$		
Γ_{19} $K^*(892)+\bar{K}^0$	(4.2 ± 1.0) %	
Γ_{20} $K^+K^-\pi^+\pi^0$		
Γ_{21} $\phi\pi^+\pi^0$	(8 ± 4) %	
Γ_{22} $\phi\rho^+$	(6.5 $^{+1.6}_{-1.8}$) %	
Γ_{23} $\phi\pi^+\pi^0$ 3-body	< 2.5 %	CL=90%
Γ_{24} $K^+K^-\pi^+\pi^0$ non- ϕ	< 8 %	CL=90%

Γ_{25}	$K^+ \bar{K}^0 \pi^+ \pi^-$	< 2.7	%	CL=90%
Γ_{26}	$K^0 K^- \pi^+ \pi^+$	(4.2 ± 1.1)	%	
Γ_{27}	$K^*(892)^+ \bar{K}^*(892)^0$	(5.6 ± 2.1)	%	
Γ_{28}	$K^0 K^- \pi^+ \pi^+$ non- $K^+ \bar{K}^*0$	< 2.8	%	CL=90%
Γ_{29}	$K^+ K^- \pi^+ \pi^+ \pi^-$	(1.8 ± 0.5)	%	
Γ_{30}	$\phi \pi^+ \pi^+ \pi^-$	(3.0 ± 3.0)	%	
Γ_{31}	$K^+ K^- \pi^+ \pi^+ \pi^-$ non- ϕ	(3.0 ± 2.0)	%	$\times 10^{-3}$

Other hadronic modes

Γ_{32}	$\pi^+ \pi^+ \pi^-$	(1.35 ± 0.31)	%	
Γ_{33}	$\rho^0 \pi^+$	< 2.8	%	$\times 10^{-3}$ CL=90%
Γ_{34}	$f_0(980) \pi^+$	(10 ± 4)	%	$\times 10^{-3}$
Γ_{35}	$\pi^+ \pi^+ \pi^-$ nonresonant	(1.01 ± 0.35)	%	
Γ_{36}	$\pi^+ \pi^+ \pi^- \pi^0$	< 12	%	CL=90%
Γ_{37}	$\eta \pi^+$	(1.9 ± 0.4)	%	
Γ_{38}	$\omega \pi^+$	< 1.7	%	CL=90%
Γ_{39}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^-$	(3.0 ± 4.0)	%	$\times 10^{-3}$
Γ_{40}	$\pi^+ \pi^+ \pi^- \pi^0 \pi^0$			
Γ_{41}	$\eta \rho^+$	(10.0 ± 2.2)	%	
Γ_{42}	$\eta \pi^+ \pi^0$ 3-body	< 2.9	%	CL=90%
Γ_{43}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0$	(4.9 ± 3.2)	%	
Γ_{44}	$\eta'(958) \pi^+$	(4.7 ± 1.4)	%	
Γ_{45}	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0 \pi^0$	(12.0 ± 3.0)	%	
Γ_{46}	$\eta'(958) \rho^+$	< 3.0	%	CL=90%
Γ_{47}	$\eta'(958) \pi^+ \pi^0$ 3-body	< 7	%	$\times 10^{-3}$ CL=90%
Γ_{48}	$K^0 \pi^+$	(3.0 ± 4.0)	%	$\times 10^{-3}$
Γ_{49}	$K^+ \pi^+ \pi^-$	< 2.5	%	$\times 10^{-3}$ CL=90%
Γ_{50}	$K^+ K^- K^+$			
Γ_{51}	ϕK^+			
Γ_{52}	A dummy mode used by the fit.	(91.7 ± 1.1)	%	

[a] For now, we average together measurements of the $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$ branching fractions.

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 9.4$ for 12 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_{15}	12			
x_{16}	9	73		
x_{17}	4	32	23	
x_{52}	-32	-87	-87	-55
	x_6	x_{15}	x_{16}	x_{17}

 D_s^\pm BRANCHING RATIOS

A few older, now obsolete results have been omitted. They may be found in our 1990 edition (Phys. Lett. **B239**).

Inclusive modes

$\Gamma(K^- \text{ anything}) / \Gamma_{\text{total}}$	Γ_1 / Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.14 -0.12 ± 0.02	COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV
$[\Gamma(K^0 \text{ anything}) + \Gamma(K^+ \text{ anything})] / \Gamma_{\text{total}}$	Γ_2 / Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.39 ± 0.28 -0.27 ± 0.04	COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV
$\Gamma(K^+ \text{ anything}) / \Gamma_{\text{total}}$	Γ_3 / Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.20 ± 0.18 -0.13 ± 0.04	COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV
$\Gamma(\text{non-}K\bar{K} \text{ anything}) / \Gamma_{\text{total}}$	Γ_4 / Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
$0.64 \pm 0.17 \pm 0.03$	⁴ COFFMAN	91	MRK3 $e^+ e^-$ 4.14 GeV

⁴ COFFMAN 91 uses the direct measurements of the kaon content to determine this non- $K\bar{K}$ fraction. This number implies that a large fraction of D_s^\pm decays involve η, η' , and/or non-spectator decays.

$\Gamma(e^+ \text{ anything}) / \Gamma_{\text{total}}$	Γ_5 / Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.20	90	⁵ BAI	90	MRK3 $e^+ e^-$ 4.14 GeV

⁵ Expressed as a value, the BAI 90 result is $\Gamma(e^+ \text{ anything}) / \Gamma_{\text{total}} = 0.05 \pm 0.05 \pm 0.02$.

Leptonic and semileptonic modes

$\Gamma(\mu^+ \nu_\mu) / \Gamma_{\text{total}}$	Γ_6 / Γ			
See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the π^\pm .				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0059 ± 0.0022 OUR FIT				Error includes scale factor of 1.1.
$0.004 \pm 0.0018 \pm 0.0020$ -0.0014 ± 0.0019	8	⁶ AOKI	93	WA75 π^- emulsion 350 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.03 0 ⁷ AUBERT 83 SPEC $\mu^+ \text{Fe}$, 250 GeV

⁶ AOKI 93 assumes the ratio of production cross sections of the D_s^+ and D^0 is 0.27. The value of $\Gamma(\mu^+ \nu_\mu) / \Gamma_{\text{total}}$ here gives a pseudoscalar decay constant $f_{D_s} = (232 \pm 45 \pm 52)$ MeV.

⁷ AUBERT 83 assume that the D_s^\pm production rate is 20% of total charm production rate.

$\Gamma(\mu^+ \nu_\mu) / \Gamma(\phi \pi^+)$	Γ_6 / Γ_{15}			
See the "Note on Pseudoscalar-Meson Decay Constants" in the Listings for the π^\pm .				
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.17 ± 0.06 OUR FIT				Error includes scale factor of 1.1.
$0.245 \pm 0.052 \pm 0.074$	39	⁸ ACOSTA	94	CLEO $e^+ e^- \approx \Upsilon(4S)$

⁸ ACOSTA 94 obtains $f_{D_s} = (344 \pm 37 \pm 52 \pm 42)$ MeV from this measurement, using $\Gamma(D_s^+ \rightarrow \phi \pi^+) / \Gamma(\text{total}) = 0.037 \pm 0.009$.

$\Gamma(\phi e^+ \nu_e) / \Gamma(\phi \pi^+)$	Γ_7 / Γ_{15}				
For now, we average together measurements of the $\Gamma(\phi e^+ \nu_e) / \Gamma(\phi \pi^+)$ and $\Gamma(\phi \mu^+ \nu_\mu) / \Gamma(\phi \pi^+)$ ratios.					
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.54 ± 0.05 OUR AVERAGE					

0.54 ± 0.05 ± 0.04 367 ⁹ BUTLER 94 CLEO $e^+ e^- \approx \Upsilon(4S)$

0.58 ± 0.17 ± 0.07 97 ¹⁰ FRABETTI 93G E687 $\gamma \text{Be } \bar{E}_\gamma \approx 220$

0.57 ± 0.15 ± 0.15 104 ¹¹ ALBRECHT 91 ARG $e^+ e^- \approx 10.4$ GeV

0.49 ± 0.10 ± 0.10
-0.14 54 ¹² ALEXANDER 90B CLEO $e^+ e^-$ 10.5-11 GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen 19 ¹³ KODAMA 93 E653 π^- emulsion 600 GeV

< 0.45 90 ¹¹ ANJOS 90B E691 $\gamma \text{Be}, \bar{E}_\gamma \approx 145$ GeV

⁹ BUTLER 94 uses both $\phi e^+ \nu_e$ and $\phi \mu^+ \nu_\mu$ events, and makes a phase-space adjustment to the latter to use them as $\phi e^+ \nu_e$ events.

¹⁰ FRABETTI 93G measures the $\Gamma(\phi \mu^+ \nu_\mu) / \Gamma(\phi \pi^+)$ ratio.

¹¹ ALBRECHT 91 and ANJOS 90B measure the $\Gamma(\phi e^+ \nu_e) / \Gamma(\phi \pi^+)$ ratio.

¹² ALEXANDER 90B measures an average of the $\Gamma(\phi e^+ \nu_e) / \Gamma(\phi \pi^+)$ and $\Gamma(\phi \mu^+ \nu_\mu) / \Gamma(\phi \pi^+)$ ratios.

¹³ KODAMA 93 measures form-factor ratios and the ratio of longitudinal to transverse partial widths.

$[\Gamma(\eta \mu^+ \nu_\mu) + \Gamma(\eta'(958) \mu^+ \nu_\mu)] / \Gamma(\phi \mu^+ \nu_\mu)$	$(\Gamma_{11} + \Gamma_{12}) / \Gamma_9$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
3.9 ± 1.6	13	KODAMA	93	E653 π^- emulsion 600 GeV

$\Gamma(\eta'(958) \mu^+ \nu_\mu) / \Gamma(\phi \mu^+ \nu_\mu)$	Γ_{12} / Γ_9			
Unsearched for decay modes of the $\eta'(958)$ are included.				
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.5	90	KODAMA	93B	E653 π^- emulsion 600 GeV

Hadronic modes with two K 's

$\Gamma(K^+ \bar{K}^0) / \Gamma(\phi \pi^+)$	$\Gamma_{13} / \Gamma_{15}$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.01 ± 0.16 OUR AVERAGE				
1.15 ± 0.31 ± 0.19	68	ANJOS	90C	E691 γBe
0.92 ± 0.32 ± 0.20		ADLER	89B	MRK3 $e^+ e^-$ 4.14 GeV
0.99 ± 0.17 ± 0.10		CHEN	89	CLEO $e^+ e^-$ 10 GeV

$\Gamma(K^+ K^- \pi^+) / \Gamma_{\text{total}}$	$\Gamma_{14} / \Gamma = (\frac{1}{2} \Gamma_{15} + \frac{2}{3} \Gamma_{16} + \frac{1}{3} \Gamma_{17}) / \Gamma$
VALUE	DOCUMENT ID
0.048 ± 0.007 OUR FIT	

Meson Full Listings

 D_s^\pm $\Gamma(\phi\pi^+)/\Gamma_{\text{total}}$ Γ_{15}/Γ

Nearly all the other modes are measured relative to this mode, which, however, is an uncertain anchor; see the footnotes to the values below, and also the note at the beginning of the list of decay modes, above.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.035±0.004 OUR FIT					
0.036±0.004 OUR AVERAGE					
0.051±0.004±0.008			14 BUTLER	94 CLEO	$e^+e^- \approx \gamma(4S)$
0.046±0.015			15 MUHEIM	94	
0.031±0.009			15 MUHEIM	94	
0.031±0.009±0.006			14 FRABETTI	93G E687	$\gamma\text{Be } \bar{E}_\gamma = 220 \text{ GeV}$
0.024±0.010			14 ALBRECHT	91 ARG	$e^+e^- \approx 10.4 \text{ GeV}$
0.031±0.006 ^{+0.011} _{-0.009}			14 ALEXANDER	90B CLEO	$e^+e^- 10.5\text{--}11 \text{ GeV}$
0.048±0.017±0.019			16 ALVAREZ	90C NA14	Photoproduction
0.033±0.016±0.010	9	17	BRAUNSCH...	87 TASS	$e^+e^- 35\text{--}44 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.048	90		MUHEIM	94	
<0.041	90	0	18 ADLER	90B MRK3	$e^+e^- 4.14 \text{ GeV}$
>0.034	90		14 ANJOS	90B E691	$\gamma\text{Be}, \bar{E}_\gamma \approx 145 \text{ GeV}$
0.02 ± 0.01	405	17	CHEN	89 CLEO	$e^+e^- 10 \text{ GeV}$
0.033±0.011	30	17	DERRICK	85B HRS	$e^+e^- 29 \text{ GeV}$

14 BUTLER 94, FRABETTI 93G, ALBRECHT 91, ALEXANDER 90B, and ANJOS 90B measure the ratio $\Gamma(D_s^+ \rightarrow \phi\ell^+\nu_\ell)/\Gamma(D_s^+ \rightarrow \phi\pi^+)$, where $\ell = e$ and/or μ , and then use a theoretical calculation of the ratio of widths $\Gamma(D_s^+ \rightarrow \phi\ell^+\nu_\ell)/\Gamma(D^+ \rightarrow \bar{K}^0\ell^+\nu)$. Not everyone uses the same value for this ratio.

15 The two MUHEIM 94 values here are model-dependent calculations based on distinct data sets. The first uses measurements of the $D_s^+(2460)^0$ and $D_{s1}(2536)^+$, the second uses B -decay factorization and $\Gamma(D_s^+ \rightarrow \mu^+\nu_\mu)/\Gamma(D_s^+ \rightarrow \phi\ell^+\nu_\ell)$. A third calculation using the semileptonic width of $D_s^+ \rightarrow \phi\ell^+\nu_\ell$ is not independent of other results listed here. Note also the upper limit, based on the sum of established D_s^+ branching ratios.

16 ALVAREZ 90C relies on the Lund model to estimate the ratio of D_s^+ to D^+ cross sections.

17 Values based on crude estimates of the D_s^\pm production level. DERRICK 85B errors are statistical only.

18 ADLER 90 uses a technique based on full reconstruction of $D_s^+D_s^-$ pairs (double tags) to obtain a branching ratio limit without assumptions about $\sigma(D_s^\pm)$.

 $\Gamma(K^+\bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{16}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.95±0.10 OUR FIT					
0.95±0.10 OUR AVERAGE					
0.85±0.34±0.20	9		ALVAREZ	90C NA14	Photoproduction
0.84±0.30±0.22			ADLER	89B MRK3	$e^+e^- 4.14 \text{ GeV}$
1.05±0.17±0.12			CHEN	89 CLEO	$e^+e^- 10 \text{ GeV}$
0.87±0.13±0.05	117		ANJOS	88 E691	Photoproduction
1.44±0.37	87		ALBRECHT	87F ARG	$e^+e^- 10 \text{ GeV}$

 $\Gamma(K^+K^-\pi^+\text{nonresonant})/\Gamma(\phi\pi^+)$ Γ_{17}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.25±0.09 OUR FIT					
0.25±0.07±0.05					
<2.6	90		ALVAREZ	90C NA14	Photoproduction

 $\Gamma(K^*(892)^+\bar{K}^0)/\Gamma(\phi\pi^+)$ Γ_{19}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.20±0.21±0.13					
<2.6	90		ALVAREZ	90C NA14	Photoproduction

 $\Gamma(\phi\pi^+\pi^0)/\Gamma(\phi\pi^+)$ Γ_{21}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.4±1.0±0.5					
<2.6	90		ALVAREZ	90C NA14	Photoproduction

 $\Gamma(\phi\rho^+)/\Gamma(\phi\pi^+)$ Γ_{22}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.86±0.26^{+0.29}_{-0.40}					
<2.6	90		ALVAREZ	90C NA14	Photoproduction

 $\Gamma(\phi\pi^+\pi^0\text{3-body})/\Gamma(\phi\pi^+)$ Γ_{23}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.71					
<2.6	90		ALVAREZ	90C NA14	Photoproduction

 $\Gamma(K^+K^-\pi^+\pi^0\text{non-}\phi)/\Gamma(\phi\pi^+)$ Γ_{24}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.4					
<2.6	90	19	ANJOS	89E E691	Photoproduction

19 Total minus ϕ component.

 $\Gamma(K^+\bar{K}^0\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{25}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.77					
<2.6	90		ALBRECHT	92B ARG	$e^+e^- \approx 10.4 \text{ GeV}$

 $\Gamma(K^0K^-\pi^+\pi^+)/\Gamma(\phi\pi^+)$ Γ_{26}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.2 ± 0.2 ± 0.2					
<2.6	90		ALBRECHT	92B ARG	$e^+e^- \approx 10.4 \text{ GeV}$

 $\Gamma(K^*(892)^+\bar{K}^*(892)^0)/\Gamma(\phi\pi^+)$ Γ_{27}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.6±0.4±0.4					
<2.6	90		ALBRECHT	92B ARG	$e^+e^- \approx 10.4 \text{ GeV}$

 $\Gamma(K^0K^-\pi^+\pi^+\text{non-}K^+\bar{K}^0)/\Gamma(\phi\pi^+)$ Γ_{28}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.80					
<2.6	90		ALBRECHT	92B ARG	$e^+e^- \approx 10.4 \text{ GeV}$

 $\Gamma(\phi\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{30}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.51±0.12 OUR AVERAGE					
0.58±0.21±0.10	21		FRABETTI	92 E687	γBe
0.42±0.13±0.07	19		ANJOS	88 E691	Photoproduction
1.11±0.37±0.28	62		ALBRECHT	85D ARG	$e^+e^- 10 \text{ GeV}$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.24	90		ALVAREZ	90C NA14	Photoproduction
-------	----	--	---------	----------	-----------------

 $\Gamma(K^+K^-\pi^+\pi^+\text{non-}\phi)/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.003^{+0.003}_{-0.002}					
<2.6	90		BARLAG	92C ACCM	$\pi^- 230 \text{ GeV}$

 $\Gamma(K^+K^-\pi^+\pi^+\text{non-}\phi)/\Gamma(\phi\pi^+)$ Γ_{31}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.32					
<2.6	90	10	ANJOS	88 E691	Photoproduction

Other hadronic modes

 $\Gamma(\pi^+\pi^+\pi^-)/\Gamma(\phi\pi^+)$ Γ_{32}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.39±0.08 OUR AVERAGE					
0.33±0.10±0.04	29		ADAMOVICH	93 WA82	$\pi^- 340 \text{ GeV}$
0.44±0.10±0.04			ANJOS	89 E691	Photoproduction

 $\Gamma(\rho^0\pi^+)/\Gamma(\phi\pi^+)$ Γ_{33}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.08					
<2.6	90		ANJOS	89 E691	Photoproduction

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.22	90		ALBRECHT	87G ARG	$e^+e^- 10 \text{ GeV}$
-------	----	--	----------	---------	-------------------------

 $\Gamma(f_0(980)\pi^+)/\Gamma(\phi\pi^+)$ Γ_{34}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.28±0.10±0.03					
<2.6	90		ANJOS	89 E691	Photoproduction

 $\Gamma(\pi^+\pi^+\pi^-\text{nonresonant})/\Gamma(\phi\pi^+)$ Γ_{35}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.29±0.09±0.03					
<2.6	90		ANJOS	89 E691	Photoproduction

 $\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma(\phi\pi^+)$ Γ_{36}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<3.3					
<2.6	90		ANJOS	89E E691	Photoproduction

 $\Gamma(\eta\pi^+)/\Gamma(\phi\pi^+)$ Γ_{37}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.54±0.09±0.06					
<2.6	90	165	ALEXANDER	92 CLEO	$\eta \rightarrow \gamma\gamma, \pi^+\pi^-\pi^0$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5	90		ANJOS	89E E691	Photoproduction
------	----	--	-------	----------	-----------------

 $\Gamma(\omega\pi^+)/\Gamma(\phi\pi^+)$ Γ_{38}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.5					
<2.6	90		ANJOS	89E E691	Photoproduction

 $\Gamma(\pi^+\pi^+\pi^+\pi^-\text{non-}\phi)/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.003^{+0.004}_{-0.003}					
<2.6	90		BARLAG	92C ACCM	$\pi^- 230 \text{ GeV}$

 $\Gamma(\pi^+\pi^+\pi^+\pi^-\text{non-}\phi)/\Gamma(\phi\pi^+)$ Γ_{39}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.29					
<2.6	90		ANJOS	89 E691	Photoproduction

 $\Gamma(\eta\rho^+)/\Gamma(\phi\pi^+)$ Γ_{41}/Γ_{15}

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
2.86±0.38^{+0.36}_{-0.38}					
<2.6	90	217	AVERY	92 CLEO	$\eta \rightarrow \gamma\gamma, \pi^+\pi^-\pi^0$

See key on page 1343

Meson Full Listings

$D_s^\pm, D_s^{*\pm}$

$\Gamma(\eta\pi^+\pi^0\text{-body})/\Gamma(\phi\pi^+)$		Γ_{42}/Γ_{15}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.82	90	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV


$\Gamma(\pi^+\pi^+\pi^+\pi^-\pi^-\pi^0)/\Gamma_{\text{total}}$		Γ_{43}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.049 ± 0.033 -0.030		BARLAG	92C ACCM π^- 230 GeV

$\Gamma(\eta'(958)\pi^+)/\Gamma(\phi\pi^+)$		Γ_{44}/Γ_{15}	
VALUE	CL% EVTS	DOCUMENT ID	TECN COMMENT
1.4 ± 0.4	OUR AVERAGE	Error includes scale factor of 2.1. See the Ideogram below.	
$1.20 \pm 0.15 \pm 0.11$	281	ALEXANDER	92 CLEO $\eta' \rightarrow \eta\pi^+\pi^-$, $\rho^0\gamma$

2.5 ± 1.0	$+1.5$ -0.4	22	ALVAREZ	91 NA14	Photoproduction
$2.5 \pm 0.5 \pm 0.3$		215	ALBRECHT	90D ARG	$e^+e^- \approx 10.4$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3		90	ANJOS	91B E691	$\gamma\text{Be}, \bar{E}_\gamma \approx 145$ GeV
------	--	----	-------	----------	---



$\Gamma(\eta'(958)\rho^+)/\Gamma(\phi\pi^+)$		Γ_{46}/Γ_{15}	
VALUE	EVTS	DOCUMENT ID	TECN COMMENT
3.44 ± 0.62 $+0.44$ -0.46	68	AVERY	92 CLEO $\eta' \rightarrow \eta\pi^+\pi^-$

$\Gamma(\eta'(958)\pi^+\pi^0\text{-body})/\Gamma(\phi\pi^+)$		Γ_{47}/Γ_{15}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.85	90	DAOUDI	92 CLEO $e^+e^- \approx 10.5$ GeV

$\Gamma(K^0\pi^+)/\Gamma(\phi\pi^+)$		Γ_{48}/Γ_{15}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.21	90	ADLER	89B MRK3 e^+e^- 4.14 GeV

$\Gamma(K^+\pi^-\pi^-)/\Gamma_{\text{total}}$		Γ_{49}/Γ	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
0.003 $+0.004$ -0.003		BARLAG	92C ACCM π^- 230 GeV

$\Gamma(\phi K^+)/\Gamma(\phi\pi^+)$		Γ_{51}/Γ_{15}	
VALUE	CL%	DOCUMENT ID	TECN COMMENT
<0.071	90	ANJOS	92D E691 $\gamma\text{Be}, \bar{E}_\gamma = 145$ GeV

D_s^\pm REFERENCES

ACOSTA	94	PR D49 5690	+Athanas, Masek, Paar+ (CLEO Collab.)
BUTLER	94	PL B324 255	+Fu, Kalbfleisch, Ross+ (CLEO Collab.)
MUHEIM	94	PR D49 3767	+Stone (SVRA)
ADAMOVICH	93	PL B305 177	+Alexandrov, Antinori+ (CERN WA82 Collab.)
AOKI	93	PTP 89 131	+Baroni, Bisi, Breslin+ (CERN WA75 Collab.)
FRABETTI	93F	PRL 71 827	+Cheung, Cumalat, Dallapiccola+ (FNAL E687 Collab.)
FRABETTI	93G	PL B313 253	+Cheung, Cumalat+ (FNAL E687 Collab.)
KODAMA	93	PL B309 483	+Ushida, Mokhtarani+ (FNAL E653 Collab.)
KODAMA	93B	PL B313 260	+Ushida, Mokhtarani+ (FNAL E653 Collab.)
ALBRECHT	92B	ZPHY C53 361	+Ehrlichmann, Hamacher, Krueger+ (ARGUS Collab.)
ALEXANDER	92	PRL 68 1275	+Bebek, Berkelman, Besson+ (CLEO Collab.)
ANJOS	92D	PRL 69 2892	+Appel, Bean, Bediaga+ (FNAL E691 Collab.)
AVERY	92	PRL 68 1279	+Freyberger, Rodriguez, Yelton+ (CLEO Collab.)
BARLAG	92C	ZPHY C55 383	+Becker, Bozek, Boehringer+ (ACCMOR Collab.)
Also	90D	ZPHY C48 29	Barlag, Becker, Boehringer, Bosman+ (ACCMOR Collab.)
DAOUDI	92	PR D45 3965	+Ford, Johnson, Lingel+ (CLEO Collab.)
FRABETTI	92	PL B281 167	+Bogart, Cheung, Culy+ (FNAL E687 Collab.)
ALBRECHT	91	PL B255 634	+Ehrlichmann, Hamacher, Krueger+ (ARGUS Collab.)
ALVAREZ	91	PL B255 639	+Barate, Bloch, Bonamy+ (CERN NA14/2 Collab.)
ANJOS	91B	PR D43 R2063	+Appel, Bean, Bracker+ (FNAL E691 Collab.)
COFFMAN	91	PL B263 135	+DeJongh, Dubois, Eigen, Hitlin+ (Mark III Collab.)
ADLER	90	PR D41 2515	+Blaylock, Bolton+ (Mark III Collab.)
ADLER	90B	PRL 64 169	+Bai, Blaylock, Bolton+ (Mark III Collab.)
ALBRECHT	90D	PL B245 15	+Ehrlichmann, Glaeser, Harder+ (ARGUS Collab.)
ALEXANDER	90B	PRL 65 1531	+Artuso, Bebek, Berkelman+ (CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	+Barate, Bloch, Bonamy+ (CERN NA14/2 Collab.)
ALVAREZ	90C	PL B246 261	+Barate, Bloch, Bonamy+ (CERN NA14/2 Collab.)
ANJOS	90B	PRL 64 2885	+Appel, Bean, Bracker+ (FNAL E691 Collab.)
ANJOS	90C	PR D41 2705	+Appel, Bean+ (FNAL E691 Collab.)
BAI	90	PRL 65 686	+Blaylock, Bolton, Brient+ (Mark III Collab.)
BARLAG	90C	ZPHY C46 563	+Becker, Boehringer, Bosman+ (ACCMOR Collab.)
FRABETTI	90	PL B251 639	+Bogart, Cheung, Coteus+ (FNAL E687 Collab.)
ADLER	89B	PRL 63 1211	+Bai, Becker, Blaylock, Bolton+ (Mark III Collab.)
Also	89D	PRL 63 2858	erratum
ANJOS	89	PRL 62 125	+Appel, Bean, Bracker+ (FNAL E691 Collab.)
ANJOS	89E	PL B223 267	+Appel, Bean, Bracker+ (FNAL E691 Collab.)
AVERILL	89	PR D39 123	+Blochus, Brabson+ (HRS Collab.)
CHEN	89	PL B226 192	+McIlwain, Miller, Ng, Shibata+ (CLEO Collab.)
ALBRECHT	88	PL B207 349	+Binder, Boeckmann+ (ARGUS Collab.)
ALBRECHT	88I	PL B210 267	+Boeckmann, Glaeser+ (ARGUS Collab.)
ANJOS	88	PRL 60 897	+Appel+ (FNAL E691 Collab.)
RAAB	88	PR D37 2391	+Anjos, Appel, Bracker+ (FNAL E691 Collab.)
ALBRECHT	87F	PL B179 398	+Binder, Boeckmann, Glaeser+ (ARGUS Collab.)
ALBRECHT	87G	PL B195 102	+Andam, Binder, Boeckmann+ (ARGUS Collab.)
ANJOS	87B	PL 58 1818	+Appel, Bracker, Browder+ (FNAL E691 Collab.)
BECKER	87B	PL B184 277	+Boehringer, Bosman+ (NA11 and NA32 Collab.)
BLAYLOCK	87	PRL 58 2171	+Bolton, Brown, Bunnell+ (Mark III Collab.)
BRAUNSC... 87	ZPHY C35 317		Braunschweig, Gerhards+ (TASSO Collab.)
CSORNA	87	PL B191 318	+Mestayer, Panvini, Word+ (CLEO Collab.)
JUNG	86	PRL 56 1775	+Abachi+ (HRS Collab.)
USHIDA	86	PRL 56 1767	+Kondo, Tasaka, Park+ (FNAL E653 Collab.)
ALBRECHT	85D	PL 153B 343	+Drescher, Binder, Drews+ (ARGUS Collab.)
DERRICK	85B	PRL 54 2568	+Fernandez, Fries, Hyman+ (HRS Collab.)
AIHARA	84D	PL E3 2465	+Alston-Garnjost, Badtke, Bakken+ (TPC Collab.)
ALTHOFF	84	PL 136B 130	+Braunschweig, Kirschnik+ (TASSO Collab.)
BAILEY	84	PL 139B 320	+Belau, Bohringer, Bosman+ (ACCMOR Collab.)
AUBERT	83	NP B213 31	+Bassompierre, Becks, Best+ (EMC Collab.)
CHEN	83C	PL 51 634	+Alam, Giles, Kagan+ (CLEO Collab.)
USHIDA	83	PRL 51 2362	+Kondo, Fujioka, Fukushima+ (FNAL E653 Collab.)

$D_s^{*\pm}$

$I(J^P) = ?(??)$

D_s^\pm MASS

The fit includes $D_s^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}$, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2110.0 ± 1.9	OUR FIT	Error includes scale factor of 1.2.	
$2106.6 \pm 2.1 \pm 2.7$	1	BLAYLOCK	87 MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

¹ Assuming D_s^\pm mass = 1968.7 ± 0.9 MeV.

$m_{D_s^{*\pm}} - m_{D_s^\pm}$

The fit includes $D_s^\pm, D^0, D_s^\pm, D^{*\pm}, D^{*0}$, and $D_s^{*\pm}$ mass and mass difference measurements.

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
141.6 ± 1.8	OUR FIT	Error includes scale factor of 1.2.		
142.4 ± 1.7	OUR AVERAGE			
$142.5 \pm 0.8 \pm 1.5$		2	ALBRECHT	88 ARG $e^+e^- \rightarrow D_s^\pm \gamma X$
143.0 ± 18.0		8	ASRATYAN	85 HLBC FNAL 15-ft, $\nu^2\text{H}$
$139.5 \pm 8.3 \pm 9.7$		60	AIHARA	84D TPC $e^+e^- \rightarrow$ hadrons
110 ± 46			BRANDELIK	79 DASP $e^+e^- \rightarrow D_s^\pm \gamma X$

² Result includes data of ALBRECHT 84b.

$D_s^{*\pm}$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 4.5		90	ALBRECHT	88 ARG $E_{\text{cm}}^{\text{eff}} = 10.2$ GeV
< 22		90	BLAYLOCK	87 MRK3 $e^+e^- \rightarrow D_s^\pm \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

Meson Full Listings

 $D_s^{*\pm}, D_{s1}(2536)^\pm, D_{sJ}(2573)^\pm$ D_s^{*+} DECAY MODES D_s^{*-} modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D_s^+ \gamma$	dominant

 D_s^{*+} BRANCHING RATIOS

$\Gamma(D_s^{*+} \gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
dominant OUR EVALUATION				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	ASRATYAN	91 HLBC	$\bar{\nu}_\mu \text{Ne}$	
seen	ALBRECHT	88 ARG	$e^+ e^- \rightarrow D_s^\pm \gamma X$	
seen	AIHARA	84D		
seen	ALBRECHT	84B		
seen	BRANDELIK	79		

 $D_s^{*\pm}$ REFERENCES

ASRATYAN	91	PL B257 255	+Marage+(ITEP, BELG, SACL, SERP, CRAC, BARI, CERN)
ALBRECHT	88	PL B207 349	+Binder, Boeckmann+ (ARGUS Collab.)
BLAYLOCK	87	PRL 58 2171	+Bolton, Brown, Bunnell+ (Mark III Collab.)
ASRATYAN	85	PL 156B 441	+Fedotov, Ammosov, Burtovoy+ (ITEP, SERP)
AIHARA	84D	PRL 53 2465	+Alston-Garnjost, Badtke, Bakken+ (TPC Collab.)
ALBRECHT	84B	PL 146B 111	+Drescher, Heller+ (ARGUS Collab.)
BRANDELIK	79	PL 80B 412	+Braunschweig, Martyn, Sander+ (DASP Collab.)

OTHER RELATED PAPERS

KAMAL	92	PL B284 421	+Xu (ALBE)
BRANDELIK	78C	PL 76B 361	+Cords+ (DASP Collab.)
BRANDELIK	77B	PL 70B 132	+Braunschweig, Martyn, Sander+ (DASP Collab.)

 $D_{s1}(2536)^\pm$

$$I(J^P) = 0(1^+)$$

$$I, J, P \text{ need confirmation.}$$

Seen in $D^*(2010)^+ K^0$. Not seen in $D^+ K^0$ or $D^0 K^+$. $J^P = 1^+$ assignment strongly favored.

 $D_{s1}(2536)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2536.35 ± 0.34 OUR AVERAGE				
2534.2 ± 1.2	9	ASRATYAN	94 BEBC	$\nu N \rightarrow D^* K^0 X, D^{*0} K^\pm X$
2535.0 ± 0.6 ± 1.0	75 ± 14	FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
2535.3 ± 0.2 ± 0.5	134 ± 22	ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*0} K^+ X$
2534.8 ± 0.6 ± 0.6	44 ± 8	ALEXANDER	93 CLE2	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.2 ± 0.5 ± 1.5	28	ALBRECHT	92R ARG	10.4 $e^+ e^- \rightarrow D^{*0} K^+ X$
2536.6 ± 0.7 ± 0.4		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
2535.9 ± 0.6 ± 2.0		ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2535 ± 28		1 ASRATYAN	88 HLBC	$\nu N \rightarrow D_s \gamma \gamma X$
1 Not seen in $D^* K$.				

 $m_{D_{s1}(2536)^\pm} - m_{D_s^*(2111)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
424 ± 28	ASRATYAN	88 HLBC	$D_s^{*\pm} \gamma$

 $D_{s1}(2536)^\pm$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.3	90		ALEXANDER	93 CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<3.2	90	75 ± 14	FRABETTI	94B E687	$\gamma \text{Be} \rightarrow D^{*+} K^0 X,$ $D^{*0} K^+ X$
<3.9	90		ALBRECHT	92R ARG	10.4 $e^+ e^- \rightarrow D^{*0} K^+ X$
<5.44	90		AVERY	90 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$
<4.6	90		ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$

 $D_{s1}(2536)^+$ DECAY MODES $D_{s1}(2536)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^*(2010)^+ K^0$	seen
$\Gamma_2 D^*(2007)^0 K^+$	seen
$\Gamma_3 D^+ K^0$	not seen
$\Gamma_4 D^0 K^+$	not seen
$\Gamma_5 D_s^{*+} \gamma$	possibly seen

 $D_{s1}(2536)^+$ BRANCHING RATIOS

$\Gamma(D^+ K^0)/\Gamma(D^*(2010)^+ K^0)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
<0.40	90	ALEXANDER	93 CLEO	$e^+ e^- \rightarrow D^{*+} K^0 X$	
<0.43	90	ALBRECHT	89E ARG	$D_{s1}^* \rightarrow D^*(2010) K^0$	

$\Gamma(D_s^{*+} \gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
possibly seen				
	ASRATYAN	88 HLBC	$\nu N \rightarrow D_s \gamma \gamma X$	

$\Gamma(D^0 K^+)/\Gamma(D^*(2007)^0 K^+)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_2
<0.12	90	ALEXANDER	93 CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$	

$\Gamma(D_s^{*+} \gamma)/\Gamma(D^*(2007)^0 K^+)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_2
<0.42	90	ALEXANDER	93 CLEO	$e^+ e^- \rightarrow D^{*0} K^+ X$	

 $D_{s1}(2536)^\pm$ REFERENCES

ASRATYAN	94	ZPHY C 61 563	+Aderholz+ (BIRM, BELG, CERN, SERP, ITEP, RAL)
FRABETTI	94B	PRL 72 324	+Cheung, Cumalat+ (FNAL E687 Collab.)
ALEXANDER	93	PL B303 377	+Bebek+ (CLEO Collab.)
ALBRECHT	92R	PL B297 425	+Ehrlichmann+ (ARGUS Collab.)
AVERY	90	PR D41 774	+Besson (CLEO Collab.)
ALBRECHT	89E	PL B230 162	+Glaser, Harder+ (ARGUS Collab.)
ASRATYAN	88	ZPHY C40 483	+Fedotov+ (ITEP, SERP)

 $D_{sJ}(2573)^\pm$ $I(J^P) = ?(??)$

OMITTED FROM SUMMARY TABLE

J^P is natural, width and decay modes consistent with 2^+ . Needs confirmation.

 $D_{sJ}(2573)^\pm$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
2573.2 ± 1.7 ± 0.9	217	KUBOTA	94 CLE2	+	$e^+ e^- \sim 10.5 \text{ GeV}$

 $D_{sJ}(2573)^\pm$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
16 ± 5 ± 3	217	KUBOTA	94 CLE2	+	$e^+ e^- \sim 10.5 \text{ GeV}$

 $D_{sJ}(2573)^+$ DECAY MODES $D_{sJ}(2573)^-$ modes are charge conjugates of the modes below.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 D^0 K^+$	seen
$\Gamma_2 D^*(2007)^0 K^+$	not seen

 $D_{sJ}(2573)^+$ BRANCHING RATIOS

$\Gamma(D^0 K^+)/\Gamma_{\text{total}}$	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
seen	217	KUBOTA	94 CLE2	±	$e^+ e^- \sim 10.5 \text{ GeV}$	

$\Gamma(D^*(2007)^0 K^+)/\Gamma(D^0 K^+)$	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
<0.33	90	KUBOTA	94 CLE2	+	$e^+ e^- \sim 10.5 \text{ GeV}$	

 $D_{sJ}(2573)^\pm$ REFERENCES

KUBOTA	94	PRL 72 1972	+Lattery, Nelson, Patton+ (CLEO II Collab.)
--------	----	-------------	---

BOTTOM MESONS**($B = \pm 1$)** $B^+ = u\bar{b}$, $B^0 = d\bar{b}$, $\bar{B}^0 = \bar{d}b$, $B^- = \bar{u}b$, similarly for B^{*} 's**EXPERIMENTAL HIGHLIGHTS OF B MESON PRODUCTION AND DECAY**

(by R.H. Schindler, SLAC)

This edition reflects the impact that CLEO-II has had on beauty, charm, and τ physics in the last two years. Starting in 1991, the CESR storage ring with the new CLEO-II detector ran in the $\Upsilon(4S)$ resonance region collecting record luminosities for a symmetric e^+e^- collider in its first year of $\sim 8-9 \text{ pb}^{-1}/\text{day}$. Over the past year, the luminosity has risen to $\sim 15 \text{ pb}^{-1}/\text{day}$ and CLEO-II has logged a total of more than 2 fb^{-1} . This edition also contains many new measurements of B mesons and baryons from Z decay, made possible by the addition of high-precision silicon vertex detectors in many of the LEP detectors coupled with their large average accumulations of $\sim 750\text{K}$ Z/year since 1992. The implementation of silicon vertexing in CDF at FNAL has made possible the measurement of precise B lifetimes from exclusive decays. The high-energy e^+e^- colliders and detectors (LEP and SLD/SLC) with their silicon-tracking systems are now starting to produce and tag heavy-flavor decays (charm and beauty) of the Z with very high efficiency and purity. This has already resulted in improvements in inclusive and exclusive lifetime measurements, the direct observation of $B^0-\bar{B}^0$ mixing by time evolution, and precision electroweak tests (measurements of the $Zb\bar{b}$ and $Zc\bar{c}$ vertex), both with and without polarized beams.

Perhaps the most exciting result to appear this year is the first observation of radiative "penguin" decays ($b \rightarrow s\gamma$ transitions). Extensive sets of branching fraction limits on many exclusive channels in the few parts per 10^4 range were reported previously by CLEO and ARGUS for both hadronic ($b \rightarrow sg$) and radiative ($b \rightarrow s\gamma$) penguins. In 1993, CLEO-II (AMMAR 93), using about 1.6 fb^{-1} of data, reconstructed eight candidate events in the channel $B^0 \rightarrow K^*(892)^0\gamma$ and five in the $B^+ \rightarrow K^*(892)^+\gamma$ channel. The $B^0 \rightarrow K^*(892)^0\gamma$ branching fraction is about 4×10^{-5} while the $B^+ \rightarrow K^*(892)^+\gamma$ branching fraction is 6×10^{-5} . These decays are important because they establish the existence of the penguin diagram, a necessary (but not sufficient) requirement for describing direct CP violation in the Standard Model. The observed branching fractions also set stringent limits on the mass of a charged Higgs scalar in a large class of models, giving $m_{H^+} > 250 \text{ GeV}$, whereas direct limits from LEP are only about 43 GeV . In the continuing search for hadronic penguins, it appears that CLEO-II is now close to having the sensitivity to observe the simplest two-body channels.

The mixing of B^0 and \bar{B}^0 mesons, first suggested by UA1 and subsequently measured by ARGUS, provided the first indirect evidence of a heavy t quark. Mixing of the B^0 has similarly

been measured by CLEO with dilepton events at the $\Upsilon(4S)$, and by the LEP experiments at the Z resonance. This year, ALEPH (BUSKULIC 93K) made the first direct measurement of the B^0 mixing by observation of the mixing-oscillation itself over time. This is the forerunner of the measurement of B_s^0 mixing, where present constraints on the CKM matrix suggest that B_s^0 mixing will be found to be nearly maximal ($\chi_s \approx 0.5$). If χ_s is greater than about 0.3-0.4, then greater sensitivity to the CKM parameters ($|V_{tb}V_{ts}^*|$) through B_s mixing will likely come from direct measurement of the mixing-oscillation frequency (the time evolution) rather than from a time-averaged quantity such as χ_s . These studies are only likely to occur in hadron and electron machines where the B mesons are adequately boosted and where pure tagging can be accomplished.

The measured average lifetime of B hadrons has changed dramatically since first measurements at PEP and PETRA. The average value of about $1.29 \pm 0.05 \text{ ps}$ in our 1992 edition was dominated by LEP measurements using high-statistics impact-parameter techniques with hadrons and leptons. Later LEP measurements average closer to $1.5 \pm 0.03 \text{ ps}$. This significant upward drift perhaps reflects improvements in understanding of vertex detectors. The longer average lifetime is confirmed by CDF (ABE 93J). Theory suggests that the inequality of B^+ and B^0 lifetimes may be as small as 5 or 10%, providing an interesting experimental challenge. The separate lifetimes of B^+ and B^0 mesons have been measured indirectly at LEP, looking at the correlation with D^{*+} as an indicator of the B charge. Other techniques have also been applied. The equality of B^+ and B^0 lifetimes is established within about 10% by averaging LEP measurements. Separate B^+ and B^0 lifetimes at this precision will come either from tagging of B mesons at the $\Upsilon(4S)$ with subsequent measurement of the semileptonic decay of the recoil, or from improved statistics for exclusive $B_{\ell 3}$ decays, or from the direct measurement of a decay lifetime of a B^+ or B^0 at LEP/SLC or CDF using vertexing techniques.

Experiments at LEP and CDF at FNAL have isolated B_s^0 and B -baryon decays using ϕ -lepton and Λ -lepton correlations. Explicit examples of decays of the B_s^0 have been reconstructed, and the mass and lifetime have been measured using these events. While the B_s^0 lifetime is similar to those of the B^+ and B^0 , the B -baryon lifetime appears to be shorter, perhaps owing to a contribution of exchange diagrams in its decays.

The semileptonic branching fractions are important for establishing the values of the CKM parameters V_{cb} and V_{ub} . The branching fractions to exclusive states $B \rightarrow \bar{D}\ell^+\nu_\ell$ and $B \rightarrow \bar{D}^*\ell^+\nu_\ell$ have been measured and both CLEO and ARGUS have done form factor decompositions (ALBRECHT 92C, ALBRECHT 93, and SANGHER 93). The inclusive rate for semileptonic decay appears to remain at about $10.0 \pm 0.5\%$. It appears that the sum of $\bar{D}\ell^+\nu_\ell$ and $\bar{D}^*\ell^+\nu_\ell$ contributions accounts for about two-thirds of the inclusive branching fraction. The balance must be a mixture of other channels such as $\bar{D}^{**}\ell^+\nu_\ell$, $\bar{D}\pi\ell^+\nu_\ell$, $\bar{D}^*\pi\ell^+\nu_\ell$ etc. Improvements can be anticipated from a comparison of single- and double-tagged

Meson Full Listings

B Meson Production and Decay, Semileptonic Decays of *B*'s

semileptonic decays combined with higher-statistics measurements of the exclusive charged and neutral $B_{\ell 3}$ channels. For more details on semileptonic decays, see the following Note.

In the next few years, CLEO-II and the detectors at high-energy machines at LEP, SLC, HERA, and FNAL should provide new insights into all aspects of B^+ , B^0 , B_s^0 meson decays. The experimental reach of the present program should be adequate to address all the interesting areas of B physics except CP violation, the study of which requires B factories. follows for a review of the many new results in that area.

NOTE ON SEMILEPTONIC DECAYS OF *D* AND *B* MESONS, PART II

(by R.J. Morrison and J.D. Richman, University of California, Santa Barbara)

This is a continuation of the discussion that began in the Listings for the D^\pm .

IV. *B*-meson semileptonic decays

In the following sections, we review inclusive and exclusive semileptonic decays of B mesons. A major goal of experiments is to determine the magnitudes of the CKM elements V_{cb} and V_{ub} . There are significant theoretical uncertainties involved in extracting these quantities from the data, and measurements of branching fractions and form factors are part of an ongoing effort to clarify the dynamics of these decays.

The inclusive semileptonic branching fraction

The b -hadron inclusive semileptonic branching fraction, \mathcal{B}_{SL} , has been measured both at the $\Upsilon(4S)$, where the b -hadrons are a mixture of B_u and B_d mesons, and at the Z , where B_s -mesons and b -baryons are produced as well. Here \mathcal{B}_{SL} is the branching fraction to either electrons or muons, not their sum. Semileptonic decays to τ leptons are suppressed by phase space and have been observed, within large errors, at the expected level [3,4]. Measurements of \mathcal{B}_{SL} are given in Table 1, for experiments at the $\Upsilon(4S)$, and in Table 2, for the LEP experiments. Table 3 gives the branching fraction for $X_{\bar{b}} \rightarrow X\tau^+\nu_\tau$ and certain semi-inclusive modes.

Table 1: Measurements of the inclusive semileptonic branching fraction (%), $\mathcal{B}_{SL} = B(B \rightarrow X\ell^+\nu_\ell)$, averaged over the B mesons produced at the $\Upsilon(4S)$ (B_u and B_d). Results are given separately for each of the models used to extract \mathcal{B}_{SL} . In the ARGUS 1ℓ measurement, the first error combines both statistical and systematic uncertainties; the second error in the ACCMM value is due to the extra free parameters present in the ACCMM model. The fit of the CLEO data using the unmodified ISGW model is poor, so the results from that fit are less reliable. The table also gives the CLEO inclusive branching fraction to charm final states ($X_{\bar{c}}$) only, which is extracted from the same fit. (Sources of error in these measurements are discussed in the text.)

Measurement	ACCMM	ISGW	ISGW**
ARGUS 1ℓ [5]	$10.2 \pm 0.5 \pm 0.2$	9.8 ± 0.5	
ARGUS 2ℓ [6]	$9.6 \pm 0.5 \pm 0.4$	$9.9 \pm 0.5 \pm 0.4$	
CRYSTAL BALL 1ℓ [7]	$12.0 \pm 0.5 \pm 0.7$	$11.9 \pm 0.4 \pm 0.7$	
CUSB-II 1ℓ [8]	$10.0 \pm 0.4 \pm 0.3$	$10.0 \pm 0.4 \pm 0.3$	
CLEO-I [9]	$10.5 \pm 0.2 \pm 0.4$	$9.9 \pm 0.1 \pm 0.4$	$11.2 \pm 0.3 \pm 0.4$
CLEO-II (prelim.) [10]	$10.65 \pm 0.05 \pm 0.33$	$10.42 \pm 0.05 \pm 0.33$	$10.98 \pm 0.10 \pm 0.33$
CLEO-II (prelim.) [10] $B \rightarrow X_{\bar{c}}\ell^+\nu_\ell$	$10.48 \pm 0.07 \pm 0.33$	$10.41 \pm 0.07 \pm 0.33$	$10.87 \pm 0.10 \pm 0.33$

Table 2: Measurements from LEP experiments of the inclusive b -hadron semileptonic branching fraction, $B(X_{\bar{b}} \rightarrow X\ell^+\nu_\ell)$, where $X_{\bar{b}}$ is a hadron containing a b -quark. At the Z , the population of b hadrons includes not only B_u and B_s mesons, but also a small fraction of B_s mesons and b -baryons.

Expt.	Ref.	$B(X_{\bar{b}} \rightarrow X\ell^+\nu_\ell)\%$
ALEPH (prelim.)	[11]	$11.4 \pm 0.33 \pm 0.37 \pm 0.20$
DELPHI (prelim.)	[11]	$9.70 \pm 0.43 \pm 0.51 \pm 0.43$
L3 (prelim.)	[11]	$11.73 \pm 0.48 \pm 0.28 \pm 0.31$
OPAL (prelim.)	[11]	$10.5 \pm 0.6 \pm 0.4 \pm 0.4$
LEP Ave.	[11]	$11.0 \pm 0.3 \pm 0.4$

Table 3: Measurements of branching fractions for $X_{\bar{b}}^- \rightarrow X\tau^+\nu_\tau$ and other semi-inclusive modes. B semileptonic decays to final states with baryons have not been observed. The ARGUS result for $B \rightarrow D_s^- X\ell^+\nu_\ell$ has been rescaled using the PDG value for $D_s^+ \rightarrow \phi\pi^+$.

Mode	Expt.	Ref.	B(%)
$X_{\bar{b}}^- \rightarrow X\tau^+\nu_\tau$	ALEPH	[3]	$4.08 \pm 0.76 \pm 0.62$
$X_{\bar{b}}^- \rightarrow X\tau^+\nu_\tau$	ALEPH (prelim.)	[4]	$2.76 \pm 0.47 \pm 0.43$
$B \rightarrow D_s^- X\ell^+\nu_\ell$	ARGUS	[20]	< 0.9 (90% CL)
$B \rightarrow \bar{p}X\ell^+\nu_\ell$	ARGUS	[5]	< 0.16 (90% CL)

The challenge for inclusive measurements is to determine what part of the observed lepton momentum spectrum is due to leptons from b -hadron decay (primary leptons) and what part is due to leptons from charm decay (secondary leptons) or to other sources (misidentified hadrons, photon conversions, $J/\psi(1S)$ decays, *etc.*). The standard technique is to fit the observed lepton momentum spectrum to a sum of the shapes expected for primary and secondary decays, after subtracting out backgrounds from other sources. Thus, a large part of the effort (and uncertainty) in the analysis is in the determination of these shapes.

Experiments at the $\Upsilon(4S)$ (ARGUS and CLEO) use theoretical models to describe the primary lepton spectrum. Some of these models have free parameters that are determined from the fit. The ACCMM model [12], for example, is based on an inclusive calculation of b -quark decay, and it has parameters corresponding to the c -quark mass and the Fermi momentum of the spectator quark, among others. A commonly used exclusive model is ISGW [13], in which the dominant contributions to the primary spectrum are from $B \rightarrow \bar{D}\ell^+\nu_\ell$ and $B \rightarrow \bar{D}^*\ell^+\nu_\ell$, with some $B \rightarrow \bar{D}^{**}\ell^+\nu_\ell$. Here, D^{**} refers to a mixture of p -wave and radially excited charm mesons. CLEO finds [9,10] that the amount of $B \rightarrow \bar{D}^{**}\ell^+\nu_\ell$ in this model is too low to adequately describe the lepton momentum spectrum, so a modified version of the ISGW model, ISGW**, has been created. In ISGW**, the D^{**} fraction is allowed to vary, but the D^* -to- D ratio is fixed at the value (2.3) predicted by ISGW. The fit to the CLEO data using ISGW** is significantly better than that using ISGW.

The shape used to describe the secondary lepton spectrum in these fits, although somewhat more complicated to obtain, is based on data. The DELCO charm-decay lepton spectrum [14] is fit to a theoretical model (ACCMM) and then boosted according to the inclusive D -meson momentum spectrum measured at the $\Upsilon(4S)$. Future measurements should be able to use a charm lepton spectrum obtained by summing the spectra for the known exclusive charm semileptonic modes, which account for most of the inclusive rate.

LEP experiments (ALEPH, DELPHI, L3, and OPAL) measure \mathcal{B}_{SL} by fitting the spectra of p and p_T (the momentum transverse to the jet axis) in single lepton and dilepton events. The shape of the primary spectrum is taken from CLEO or ARGUS, so that model-related uncertainties in these experiments are propagated into the LEP results. (This uncertainty is part of the third error in the LEP results given in Table 2. Its size depends on the specific details of the analysis.)

The extraction of the B semileptonic branching fraction from the momentum spectrum of single leptons, therefore, relies on models. In CLEO, which currently has the largest data sample, the spread of values obtained using different models is comparable to the experimental errors. The dominant experimental errors are due to tracking and lepton identification uncertainties, which are expected to decrease in the future.

The ARGUS collaboration [6] has introduced a second method, using dilepton events, that substantially reduces the need for models. One lepton (the “tagging lepton”) is required to have high momentum and is thus nearly always primary. The analysis then examines the momentum spectrum of the second lepton in the event. By requiring that both leptons be in the same hemisphere, events in which the two leptons come from the decay chain of a single B meson (produced nearly at rest at the $\Upsilon(4S)$) are effectively removed. Thus, (1) the tagging lepton is primary, and (2) the leptons are from different B mesons. Then, unless mixing occurs, a lepton whose charge is opposite to that of the tagging lepton must be primary, while one with the same charge as the tagging lepton must be secondary. One corrects for mixing by using the known mixing probability. The relative charges of the two leptons, therefore, can be used to separate the primary and secondary spectra of the second lepton. There is a lower momentum cutoff due to experimental acceptance, however, and a small extrapolation, based on models, is required to obtain the total semileptonic rate. The results of this measurement are in agreement with the single-lepton value, although with a poorer statistical error. This method also very much reduces the sensitivity to any possible non- $B\bar{B}$ decays of the $\Upsilon(4S)$, which are assumed to be negligible in the single-lepton method.

Meson Full Listings

Semileptonic Decays of B 's

The values of \mathcal{B}_{SL} given in Table 1 and Table 2 are lower than theoretical predictions, which give [15] $\mathcal{B}_{SL} \geq 12.5\%$. Such calculations, however, are difficult partly because they must determine the total hadronic rate, which has uncertainties associated with both perturbative and nonperturbative QCD effects. The source of the discrepancy between theory and experiment is not yet understood.

The semileptonic branching fraction can be used to calculate $|V_{cb}|$. The fraction is converted into a decay rate using the measured B lifetime, and the rate is then compared with model predictions, which depend on $|V_{cb}|^2$. Whereas the model dependence in the determination of \mathcal{B}_{SL} is associated with the predicted shapes of momentum spectra, the extraction of $|V_{cb}|$ is also sensitive to the uncertainty in the normalization of the predicted rate. It is difficult to assign errors to rate predictions based on quark-model calculations. Quite often, a nominal theoretical error of 20% in the rate is assumed, leading to a 10% theoretical error on $|V_{cb}|$. This uncertainty is separate from the model dependence, if any, in extracting the branching fraction. The CLEO-II preliminary results (assuming the value [16] $\tau_B = 1.49 \pm 0.04$ ps which is used in Sec. 28 “The Cabibbo-Kobayashi-Maskawa Mixing Matrix”) for $B \rightarrow X_{\bar{c}}\ell^+\nu_\ell$ in Table 1 give [17]

$$|V_{cb}| = (0.042 \pm 0.001 \pm 0.004) \sqrt{\frac{1.49}{\tau_B(\text{ps})}} \quad (1)$$

for both ACCMM and ISGW**. (The value for V_{cb} in Sec. 28 “The Cabibbo-Kobayashi-Maskawa Mixing Matrix” is not based on this measurement but on the $B \rightarrow \bar{D}^*\ell^+\nu_\ell$ rate at zero D^* recoil velocity.)

Currently, there is interest in using the methods of HQET to obtain a value of $|V_{cb}|$ from the inclusive spectrum with a better understood error. Luke and Savage [18] use the inclusive branching fraction to obtain the range $0.038 < |V_{cb}| < 0.054$, where we have corrected their result to $\tau_B = 1.49$ ps. Shifman, Uraltsev, and Vainshtein [19] argue that the present theoretical uncertainty on $|V_{cb}|$ may be as low as 5% and can be made even lower. We discuss the determination of $|V_{cb}|$ using exclusive processes in Section IV.4.

Table 3 lists additional inclusive B branching fractions that involve leptons. The decay $X_{\bar{b}} \rightarrow X\tau^+\nu_\tau$ has been difficult to observe, due to the presence of two neutrinos. However, ALEPH has used decay configurations with a large missing energy in the b -hadron rest frame. In Z decays, the boost of the b hadron to the lab frame can result in a very large missing energy (10–30 GeV) for the hemisphere containing the decay. By looking for events with both a large missing energy and a tagged b hadron (using vertex detector information), ALEPH has been able to measure $B(X_{\bar{b}} \rightarrow X\tau^+\nu_\tau)$ with a value consistent with standard model expectations. (The preliminary ALEPH result in Table 3 includes data from the original measurement, so the two results cannot be averaged.)

The lepton endpoint region and the determination of $|V_{ub}|$

The determination of $|V_{ub}|$ is one of the most important and challenging measurements in B physics. For $|V_{ub}/V_{cb}| \approx 0.1$, the rate for $B \rightarrow X_{\bar{u}}\ell^+\nu_\ell$, where $X_{\bar{u}}$ is a charmless hadronic system, is expected to be only about 1% of the inclusive semileptonic rate. By working in the region at and beyond the lepton-momentum-spectrum endpoint for $B \rightarrow X_{\bar{c}}\ell^+\nu_\ell$ processes, however, one gains enormously in sensitivity to $B \rightarrow X_{\bar{u}}\ell^+\nu_\ell$ decays.

Although the advantages of working in this endpoint region ($2.3 < p_\ell < 2.6$ GeV/ c) are decisive, there are also disadvantages. A major difficulty is the need to convert the measured rate for this tiny portion of phase space into a value of $|V_{ub}|$. This calculation can be performed using either inclusive or exclusive models, but both have substantial uncertainties in predicting the rate in the endpoint region. Inclusive models are expected to be fairly reliable, if one considers a large enough part of phase space, but they may not be reliable in the endpoint region, which some theorists argue [13] is dominated by a small set of exclusive channels ($B \rightarrow \rho\ell^+\nu_\ell$, $B^+ \rightarrow \omega\ell^+\nu_\ell$, and $B \rightarrow \pi\ell^+\nu_\ell$). Alternatively, exclusive models can be used to predict the sum of contributions of individual modes in this region. However, large uncertainties exist in the calculations of the rates for exclusive modes, and some of the observed rate may be due to nonresonant final states [24]. The exclusive calculations here are more difficult than those for $B \rightarrow X_{\bar{c}}\ell^+\nu_\ell$, because the range of recoil velocities available to the light final-state mesons in a $B \rightarrow X_{\bar{u}}\ell^+\nu_\ell$ transition is much larger than for the charm mesons in a $B \rightarrow X_{\bar{c}}\ell^+\nu_\ell$ decay. One therefore expects a much larger variation in the (poorly known) form factors that enter into the decay rate. As a result, measurements of $|V_{ub}|$ are currently quite model dependent, and there is substantial variation among values obtained using different models.

The analysis of the endpoint region, although an “inclusive” measurement, is quite different from the measurement of \mathcal{B}_{SL} described in the previous section: at the $Y(4S)$, nonresonant (continuum) processes produce high-momentum leptons that constitute an enormous background (relative to a $B \rightarrow X_{\bar{u}}\ell^+\nu_\ell$ signal) unless suppressed by kinematic cuts. The signal efficiency of these cuts is model dependent, unlike the very loose cuts used in the analysis of the inclusive lepton spectrum. In particular, the efficiency depends on the q^2 distribution of the signal events, so the value obtained for the rate in the endpoint region depends on the shape assumed for this distribution. The most important sensitivity to models, however, arises when one converts the rate to $|V_{ub}|$.

Table 4 lists the measurements of $|V_{ub}/V_{cb}|$ from CLEO and ARGUS. The CLEO-II studies, which are based on about five times as much data as either the original ARGUS or the CLEO-I analyses, yield values of $|V_{ub}/V_{cb}|$ significantly lower than the earlier measurements.

Table 4: Measurements of $|V_{ub}/V_{cb}|$ using the inclusive rate in the endpoint region. The ARGUS and CLEO-I results are each based on about 200,000 $b\bar{b}$ events, and the CLEO-II results are based on about 955,000 $b\bar{b}$ events.

Model	ARGUS [21]	CLEO-I [22]	CLEO-II [23]
ACMM	0.11 ± 0.012	0.09 ± 0.01	0.076 ± 0.008
ISGW	0.20 ± 0.023	0.15 ± 0.02	0.101 ± 0.010

Exclusive semileptonic decays

Measurements of exclusive semileptonic decays of B mesons are less precise and less complete than those for D mesons. Table 5 lists the branching fractions, which assume that the branching fractions of the $\Upsilon(4S)$ to B^+B^- and $B^0\bar{B}^0$ are given by $f_{+-} = f_{00} = 0.5$. Two techniques have been used to study $B \rightarrow \bar{D}^*\ell^+\nu_\ell$, the only well-measured decay mode. In the first, used by both CLEO and ARGUS, the D^* and D mesons are fully reconstructed. Signal events produce a narrow peak in the $D^* - D$ mass difference distribution, as well as a peak near zero, indicating a neutrino, in the distribution of the missing mass recoiling against the $\bar{D}^*\ell^+$ system. This latter peak is broadened by the assumption that the momentum of the B at the $\Upsilon(4S)$ is zero; in reality it is about 330 MeV/c and its direction is unknown. A small contribution to the upper shoulder of the observed missing mass distribution is attributed to $B \rightarrow \bar{D}^{**}\ell^+\nu_\ell$ decay followed by $\bar{D}^{**} \rightarrow \bar{D}^*\pi$.

Table 5: Branching fractions for exclusive B semileptonic decays. The measurements assume that $f_{+-} = f_{00} = 0.5$, where f_{+-} and f_{00} are the branching fractions of the $\Upsilon(4S)$ to B^+B^- and $B^0\bar{B}^0$. Measurements with an asterisk (*) either use more recent values of D and D^* branching fractions or have been corrected. Those with a dagger (†) used early D^* branching fractions, but, due to the complexity of the analysis procedure, are difficult to correct. These points are discussed in the review by Stone (see text). The ARGUS measurement of $B \rightarrow \bar{D}^{**}\ell^+\nu_\ell$ assumes a set of D^{**} states distributed according to the ISGW model. The limits on $b \rightarrow u$ semileptonic decays from CLEO are based on efficiencies calculated using the ISGW model; other models give different lepton energy and q^2 distributions and hence somewhat different efficiencies.

Mode	Experiment	Ref.	Branching fraction
$B^0 \rightarrow D^-\ell^+\nu_\ell$	ARGUS*	[27]	$(2.0 \pm 0.7 \pm 0.6)\%$
$B^0 \rightarrow D^-\ell^+\nu_\ell$	CLEO-I†	[28]	$(1.8 \pm 0.6 \pm 0.3)\%$
$B^+ \rightarrow \bar{D}^0\ell^+\nu_\ell$	CLEO-I†	[28]	$(1.6 \pm 0.6 \pm 0.3)\%$
$B^0 \rightarrow D^{*-}\ell^+\nu_\ell$	ARGUS*	[29]	$(4.7 \pm 0.5 \pm 0.5)\%$
$B^0 \rightarrow D^{*-}\ell^+\nu_\ell$	ARGUS* (partial rec.)	[30]	$(4.5 \pm 0.3 \pm 0.4)\%$
$B^0 \rightarrow D^{*-}\ell^+\nu_\ell$	CLEO-I*	[36]	$(4.0 \pm 0.4 \pm 0.6)\%$
$B^0 \rightarrow D^{*-}\ell^+\nu_\ell$	CLEO-II* (prelim.)	[17]	$(4.50 \pm 0.44 \pm 0.44)\%$
$B^+ \rightarrow \bar{D}^{*0}\ell^+\nu_\ell$	ARGUS*	[31]	$(6.6 \pm 1.6 \pm 1.5)\%$
$B^+ \rightarrow \bar{D}^{*0}\ell^+\nu_\ell$	CLEO-I†	[28]	$(4.1 \pm 0.8_{-0.9}^{+0.8})\%$
$B \rightarrow \bar{D}^{**}\ell^+\nu_\ell$	ARGUS	[29]	$(2.7 \pm 0.5 \pm 0.5)\%$
$B^+ \rightarrow \omega\ell^+\nu_\ell$	CLEO-II	[32]	$< 2.1 \times 10^{-4}$ (90% CL)
$B^+ \rightarrow \rho^0\ell^+\nu_\ell$	CLEO-II	[32]	$< 2.1 \times 10^{-4}$ (90% CL)
$B^0 \rightarrow \rho^-\ell^+\nu_\ell$	CLEO-II	[32]	$< 4.1 \times 10^{-4}$ (90% CL)
$B^0 \rightarrow \pi^-\ell^+\nu_\ell$	CLEO-II (prelim.)	[17]	$< 3.3 \times 10^{-4}$ (90% CL)

Meson Full Listings

Semileptonic Decays of B 's

In the second method, first employed by ARGUS, the \bar{D}^* in $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ decay is identified using only the soft pion from the $\bar{D}^* \rightarrow \bar{D} \pi$ decay (partial reconstruction). Due to the small energy release in this process, both the energy and the direction of the \bar{D}^* can be estimated from the soft pion's momentum alone. This technique has a large statistical advantage over the method in which both the \bar{D}^* and the \bar{D} are completely reconstructed.

Although $B \rightarrow \bar{D} \ell^+ \nu_\ell$ decay has a substantial branching fraction, measurements to date have suffered from poor statistics and the difficulty of dealing with feed-down from $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$. Several ARGUS and CLEO-I measurements of $B \rightarrow \bar{D} \ell^+ \nu_\ell$ and $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ were extracted using early D and D^* branching fractions, some of which were quite far from their present values. Stone [1] has used more recent D^* and D and branching fractions [25,26] to correct these measurements when possible. The values marked with an asterisk (*) in Table 5 use the same method as Stone but with the updated PDG 1994 $D^0 \rightarrow K^- \pi^+$ branching fraction. Together, $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ and $B \rightarrow \bar{D} \ell^+ \nu_\ell$ account for about two-thirds of the inclusive semileptonic rate. There is evidence that part of the remainder is due to $B \rightarrow \bar{D}^{**} \ell^+ \nu_\ell$ decay, where D^{**} represents a mixture of p -wave and radially excited D mesons.

Searches have also been made for exclusive $B \rightarrow X_{\bar{c}} \ell^+ \nu_\ell$ decays. For these decays, unlike $B \rightarrow X_{\bar{c}} \ell^+ \nu_\ell$, model predictions [13] indicate that the rate should be distributed over many exclusive channels, with no dominant modes. The decays with the largest expected branching fractions are $B^0 \rightarrow \rho^- \ell^+ \nu_\ell$, $B^+ \rightarrow \rho^0 \ell^+ \nu_\ell$, and $B^+ \rightarrow \omega \ell^+ \nu_\ell$, which in the quark model are predicted to occur in the ratio 2:1:1. Furthermore, these modes should have hard lepton-momentum spectra, so that there is good efficiency for these decays in the endpoint region, where the backgrounds from $B \rightarrow X_{\bar{c}} \ell \nu$ are smallest. CLEO has searched for all three modes but has found no significant signals, leading to the upper limits given in Table 5. These limits can be converted into model-dependent upper limits on $|V_{ub}/V_{cb}|$. The limit for the ISGW model (correcting the result to $\tau_B = 1.49$ ps) is $|V_{ub}/V_{cb}| < 0.12$ (90% CL), consistent with the CLEO-II inclusive measurement (using the ISGW model) given in Table 4.

Measurement of $|V_{cb}|$ from $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$

Three types of measurements can be used to determine $|V_{cb}|$: (1) inclusive semileptonic rate (discussed in Section IV.1), (2) the rates for $B \rightarrow \bar{D} \ell^+ \nu_\ell$, or (3) $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$, or a partial rate for $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ corresponding to the kinematic configurations in which the D^* recoils slowly. The $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ mode is especially well suited to measuring $|V_{cb}|$. Using HQET, the rate for this process can be accurately predicted (as a function of $|V_{cb}|^2$) for the kinematic configuration in which the D^* is produced at rest and the lepton and neutrino are back to back in the B rest frame. This configuration occurs when $q^2 = q_{\max}^2 = (m_B - m_{D^*})^2$. The light constituents of the initial B meson are then essentially undisturbed by the $B \rightarrow X_{\bar{c}} \ell^+ \nu_\ell$ transition, at least in the limit where the b and c quark masses are taken to be very large compared with Λ_{QCD} .

In this large-mass limit, all of the $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ form factors are related to a single form factor, the Isgur-Wise function, which depends only on the relative four-velocities of the initial and final hadrons: $\xi = \xi(v_B \cdot v_{D^*})$. Note that $v_B \cdot v_{D^*} = \gamma_{D^*}$, where the relativistic factor γ_{D^*} is measured in the B -meson rest frame. The quantity $v \cdot v'$, where v is the initial and v' is the final meson four-velocity, is often called w or y in the literature. It is linearly related to q^2 by

$$w = v \cdot v' = \frac{M^2 + m^2 - q^2}{2Mm}, \quad (2)$$

where M and m are the masses of the parent and daughter mesons.

At zero recoil ($w = 1$ or $q^2 = q_{\max}^2$), the normalization of ξ is known, $\xi(1) = 1$, which means that the decay rate in this configuration can be accurately predicted as a function of $|V_{cb}|^2$. Corrections to this picture arise from hard gluon corrections and because the masses of the b and c quarks are not truly infinite. However, the heavy quark symmetry limit prediction for $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ is protected by Luke's theorem [33], which states that, for this process, there are no leading order ($1/m_Q$) nonperturbative corrections at $w = 1$, where m_Q is the mass of a heavy quark (c or b). As a consequence, the leading corrections to the decay rate prediction arise at order $1/m_Q^2$. The calculation of these corrections, which introduces some model dependence, has been the subject of many investigations. Neubert [2,34] has obtained the result

$$\frac{1}{\tau_B} \cdot \frac{dB(B^0 \rightarrow D^{*-} \ell^+ \nu_\ell)}{dw} = \frac{G_F^2}{48\pi^3} m_{D^*}^3 (m_B - m_{D^*})^2 \eta_A^2 \sqrt{w^2 - 1} (w + 1)^2 \left[1 + \frac{4w}{w+1} \frac{1 - 2wr + r^2}{(1-r)^2} \right] |V_{cb}|^2 \hat{\xi}(w)^2, \quad (3)$$

where $r = m_{D^*}/m_B$, $\eta_A = 0.99$, and

$$\hat{\xi}(1) = 1 + \delta_{(1/m^2)} = 1.00 \pm 0.04. \quad (4)$$

(The values for η_A and $\hat{\xi}(1)$ are preliminary.) The unknown function $\hat{\xi}(w)$ is related to $\xi(w)$, as discussed by Neubert [2], but includes symmetry breaking corrections. These predictions can be used to determine $|V_{cb}|$ from the measured rate at zero recoil. Strictly speaking, there is no phase space for this configuration, so in practice one has to measure the rate in a small region near $w = 1$. (A new estimate of the corrections to the heavy-quark symmetry limit by Shifman, Uraltsev, and Vainshtein [19] gives a lower value of $\hat{\xi}(1)$ and a higher value of $|V_{cb}|$.)

Unfortunately, both CLEO and ARGUS have difficulty in measuring the rate in this region because the decay of the slow-moving D^* produces a very slow pion, which is difficult to detect with high efficiency. The rate at zero recoil is therefore obtained by measuring the rate as a function of w and then extrapolating to $w = 1$. This procedure introduces some model dependence because HQET does not predict the w dependence of the form factors, which involves nonperturbative QCD physics. (Several groups are using lattice QCD to predict the w dependence and are beginning to obtain interesting results [35].) Because the w range is small, however, the form factors are expected to have only modest variation, which is approximately linear. CLEO and ARGUS use a variety of functional forms to parametrize the w dependence. The different extrapolations to $w = 1$ lead to a range of values for $|V_{cb}|$, some of which are listed in Table 6. The systematic errors on the CLEO measurements are due to uncertainties in the soft-pion detection efficiency, smearing effects, background ($B \rightarrow \bar{D}^{*+} \ell^+ \nu_\ell$), and sensitivity to the values of the form factors. As discussed for the exclusive branching ratios, these measurements assume $B(\mathcal{Y}(4S) \rightarrow B^0 \bar{B}^0) = 0.5$. It should be noted that $|V_{cb}|$ can also be obtained from the total rates for $B \rightarrow \bar{D} \ell^+ \nu_\ell$ and $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$. The theoretical predictions, however, are less reliable than those for the zero-recoil configuration of $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$.

Table 6: Measurements of $|V_{cb}|$ and $\hat{\rho}^2$ using the decay $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ and HQET predictions. The ARGUS measurements of $|V_{cb}|$ have been corrected to a B lifetime of 1.49 ps.

Expt.	Ref.	$\xi(w) = \xi(v_B \cdot v_{D^*})$	$\hat{\rho}^2$	$ V_{cb} $
ARGUS	[29]	$1 - \hat{\rho}^2(w - 1)$	$1.17 \pm 0.22 \pm 0.06$	$0.042 \pm 0.005 \pm 0.003$
ARGUS	[29]	$\exp[-\hat{\rho}^2(w - 1)]$	$1.88 \pm 0.38 \pm 0.16$	$0.047 \pm 0.008 \pm 0.002$
ARGUS	[29]	$[2/(w + 1)]^{2\hat{\rho}^2}$	$2.10 \pm 0.38 \pm 0.18$	$0.048 \pm 0.008 \pm 0.003$
CLEO-II	[17] (prelim.)	$1 - \hat{\rho}^2(w - 1)$	$1.2 \pm 0.5 \pm 0.3$	$0.038 \pm 0.006 \pm 0.004$
CLEO-II	[17] (prelim.)	$\exp[-\hat{\rho}^2(w - 1)]$	$1.0 \pm 0.4 \pm 0.2$	$0.037 \pm 0.005 \pm 0.004$
CLEO-II	[17] (prelim.)	$[2/(w + 1)]^{2\hat{\rho}^2}$	$1.1 \pm 0.5 \pm 0.2$	$0.038 \pm 0.005 \pm 0.004$

Form factor measurements

Both ARGUS and CLEO have used measurements of kinematic distributions to obtain information on the form factors for $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$. In contrast to D -meson semileptonic decays, however, one expects the predictions of heavy quark effective theory to be applicable to $B \rightarrow X \ell^+ \nu_\ell$. Here, both the initial- and final-state quarks are heavy compared with the typical hadronic scale set by Λ_{QCD} . With sufficiently large data samples, the $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ decay should provide useful tests of HQET predictions. Ideally, one would determine the q^2 dependence of each of these form factors with minimal assumptions. In practice, due to the limited data samples so far available, experiments have taken the shapes of the q^2 dependence from models. The ratios of the form factors at a particular q^2 , usually q_{\max}^2 , are then obtained from fits to observed kinematic distributions.

Both ARGUS and CLEO have extracted the values of two key observables that can be related to the values of the form factors. The first of these is A_{FB} , the forward-backward asymmetry of leptons in the virtual- W rest frame (see Section II). The nonzero value of A_{FB} is a consequence of the parity-violating $V - A$ coupling, which results in more leptons being produced in the hemisphere opposite the D^* momentum vector than in the hemisphere containing it. The second observable is the polarization parameter α (see Section II). It is important to recognize that although both A_{FB} and α are defined as integrals over all of phase space, the lepton asymmetry and the D^* polarization are actually functions of q^2 . Thus, measurements for which the acceptance over phase space is not uniform must be either corrected or compared with theoretical predictions incorporating the same acceptance.

Table 7 compares theoretical predictions of A_{FB} and α from Neubert [2] with measurements from ARGUS and CLEO. The measurements, however, are slightly different. In the CLEO-II measurement, A_{FB} is an average over all decays for which $p_\ell > 1.0$ GeV/ c . In ARGUS, the same cut is used (for both A_{FB} and α), but the effect of this cut is removed using theoretical models. CLEO-I has also measured [36] $\alpha = 0.65 \pm 0.66 \pm 0.25$ for $p_\ell > 1.4$ GeV/ c .

The dependence of the CLEO results on the momentum cuts disappears only when the form factor ratios are calculated (for each model) using a least-squares fit that incorporates the acceptance. In the ARGUS analysis, on the other hand, the kinematic distributions are fitted directly to models in which the only free parameters are form factor ratios. The values of A_{FB} and α are then derived from each model using the fitted values for the form factors. Thus, the dependence of A_{FB} and α on the lepton-momentum cut is removed model by model. ARGUS finds that their values of A_{FB} and α are insensitive to the model used. (They do not quote the form factor ratios obtained from the fits.)

The observation of a positive value for A_{FB} is consistent with predictions of form factor models that assume $V - A$ couplings at both W -boson vertices. CLEO has shown that models in which the hadronic current has a $V + A$ structure do not describe the data well, assuming that $V - A$ is still used for the leptonic current [37]. (However, the measurement of A_{FB} cannot rule out the case in which both the hadronic and leptonic currents are $V + A$.) The predicted value of A_{FB} is a function of ρ^2 , the slope of the Isgur-Wise function, as can be seen from Table 7.

Meson Full Listings

Semileptonic Decays of B 's

Table 7: Comparison of theoretical and experimental values of the D^* polarization parameter α and the lepton forward-backward asymmetry A_{FB} in $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ decay. The theoretical values, which depend on the form factor slope $\rho_{A_1}^2$, were obtained by Neubert using HQET with the assumption of a pole form for the form factor.

$\rho_{A_1}^2$	A_{FB}	$A_{FB} (p_\ell > 1 \text{ GeV}/c)$	α
1.1	0.22	0.16	1.33
0.8	0.22	0.16	1.44
0.5	0.22	0.15	1.55
Expt.	$0.20 \pm 0.08 \pm 0.06$ ARGUS [29]	$0.14 \pm 0.06 \pm 0.03$ CLEO-II [37]	$1.1 \pm 0.4 \pm 0.2$ ARGUS [29]

CLEO has performed least-squares fits using the measured values of A_{FB} , α (averaged over CLEO and ARGUS results), and the histogram of $d\Gamma/dq^2$ (also averaged over CLEO and ARGUS measurements), to obtain values for the ratios of the $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ form factors at q^2_{max} . The results of these fits are given in Table 8, together with the predictions of several models. Although the values of the form factor ratios are in agreement with predictions, the precision is not yet sufficient to distinguish among the models.

Table 8: Measurements and predictions for ratios of the form factors for $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$, evaluated at $q^2 = q^2_{\text{max}}$. The two fits (a) and (b) correspond to different assumptions for the q^2 dependence of the form factors. The fits given in the CLEO paper quoted here combined the A_{FB} measurement from CLEO with ARGUS and CLEO measurements of α and $d\Gamma/dq^2$.

	Ref.	A_2/A_1	V/A_1
CLEO-II Fit (a)	[37]	1.02 ± 0.24	1.07 ± 0.57
CLEO-II Fit (b)	[37]	0.79 ± 0.28	1.32 ± 0.62
ISGW	[13]	1.14	1.27
KS	[38]	1.39	1.54
WSB	[39]	1.06	1.14
HQET-based	[40]	1.26	1.26
HQET-based	[41]	1.14	1.74

Conclusions on B semileptonic decays

Knowledge of B semileptonic decays is advancing rapidly, and many of the values given in this review are likely to be superseded in the near future. The inclusive B semileptonic branching fraction is already one of the most precisely measured quantities in B physics, and experimental errors should continue to improve. Important progress is being made in minimizing the model dependence of this measurement by using dilepton events. Knowledge of the exclusive semileptonic branching ratios is still rather poor: only $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ is known with reasonable precision, and exclusive $B \rightarrow X_{\bar{c}} \ell^+ \nu_\ell$ transitions have not yet been measured. By our next edition, this situation may well have changed.

Semileptonic B decays provide important information on $|V_{cb}|$ and $|V_{ub}|$. The inclusive $B \rightarrow X_{\bar{c}} \ell^+ \nu_\ell$ decay rate is roughly one hundred times that for $B \rightarrow X_{\bar{u}} \ell^+ \nu_\ell$ decays, so that $|V_{ub}|$ is much harder to measure. Furthermore, the model dependence is more severe for measurements of $|V_{ub}|$ than for $|V_{cb}|$. The $B \rightarrow X_{\bar{u}} \ell^+ \nu_\ell$ signal is statistically significant only in a very small part of phase space, the lepton spectrum endpoint region, in which model predictions are questionable. In the future, measurements of the q^2 distribution in both inclusive and exclusive $B \rightarrow X_{\bar{u}} \ell^+ \nu_\ell$ decays should help to reduce the model dependence, and lattice QCD may be important for improving $|V_{ub}|$ determinations. Finally, because both quarks are heavy in a $B \rightarrow X_{\bar{c}} \ell^+ \nu_\ell$ transition, one can use the tools of heavy quark effective theory to predict decay rates. Using $B \rightarrow \bar{D}^* \ell^+ \nu_\ell$ in the kinematic region where the D^* has only a small recoil velocity, one can determine $|V_{cb}|$ with relatively little model dependence, even though the data samples are not yet large enough to fully realize the potential of this approach. The methods of HQET may also lead to a precise determination of $|V_{cb}|$ from the inclusive semileptonic rate.

It is important to test HQET in detail. Although form factor measurements are roughly consistent with HQET predictions, the experimental uncertainties are too large at present to allow a stringent test. And although heavy quark symmetry can relate different form factors to the Isgur-Wise function, only nonperturbative techniques such as lattice QCD can actually predict the q^2 dependence of this function. One can expect significant progress in all of these areas during the next few years.

See key on page 1343

Meson Full Listings

Semileptonic Decays of B^\pm , B^\pm

References

- S. Stone, *Semileptonic B Decays*, HEPHY 93-11, to appear in *B Decays*, 2nd Edition, ed. by S. Stone, World Scientific, Singapore.
- M. Neubert, *Heavy Quark Symmetry*, SLAC-PUB-6263, June 1993, to appear in *Physics Reports*.
- ALEPH: D. Buskulic *et al.*, *Phys. Lett.* **B298**, 479 (1993).
- Updated Measurement of the $b \rightarrow \tau^- \bar{\nu}_\tau X$ Branching Ratio*, ALEPH Collaboration, paper submitted to the 5th International Symposium on Heavy Flavour Physics, Montreal, 1993.
- ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B249**, 359 (1990).
- ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B318**, 397 (1993).
- Crystal Ball: K. Wachs *et al.*, *Z. Phys.* **C42**, 33 (1989).
- CUSB: C. Yanagisawa *et al.*, *Phys. Rev. Lett.* **66**, 2436 (1991).
- CLEO: S. Henderson *et al.*, *Phys. Rev.* **D45**, 2212 (1992).
- CLEO: J. Bartelt *et al.*, *Inclusive Measurement of B-Meson Semileptonic Branching Fractions*, submitted to the XVI International Symposium on Lepton and Photon Interactions, Ithaca, New York (1993), p. 274.
- W. Venus, *b Weak Interaction Physics in High Energy Experiments*, in AIP Conference Proceedings 302, Lepton and Photon Interactions, XVI International Symposium, eds. P. Drell and D. Rubin, Ithaca, NY (1993), p. 274.
- G. Altarelli, N. Cabibbo, G. Corbò, L. Maiani, and G. Martinelli, *Nucl. Phys.* **B208**, 365 (1982).
- N. Isgur, D. Scora, B. Grinstein, and M. Wise, *Phys. Rev.* **D39**, 799 (1989).
- DELCO: W. Bacino *et al.*, *Phys. Rev. Lett.* **43**, 1073 (1979).
- I. Bigi, B. Blok, M. Shifman, A. Vainshtein, *Phys. Lett.* **B323**, 408 (1994);
G. Altarelli and S. Petrarca, *Phys. Lett.* **B261**, 303 (1991).
- This average b -hadron lifetime, also used in Sec. 28 on "The Cabibbo-Kobayashi-Maskawa Mixing Matrix," is based on results from LEP experiments and CDF and includes contributions from B_s^0 mesons and b -baryons, which are not produced at the $\Upsilon(4S)$. However, these contributions are thought to be relatively small (LEP experiments typically assume nominal production ratios given by $B^+ : B^0 : B_s^0 : b$ -baryon = 0.40 : 0.40 : 0.12 : 0.08), and their lifetimes are similar to those of B^+ and B^0 mesons. Alternatively, one can use the less precisely measured B^+ and B^0 lifetimes ($\tau_{B^0} = 1.50 \pm 0.11$ ps and $\tau_{B^+} = 1.54 \pm 0.11$ ps) from CDF and LEP experiments.
- D. Besson, *Recent Bottom Physics Results from Threshold Machines*, in AIP Conference Proceedings 302, Lepton and Photon Interactions, XVI International Symposium, eds. P. Drell and D. Rubin, Ithaca, NY (1993), p. 221.
- M. Luke and M. Savage, *Phys. Lett.* **B321**, 88 (1994).
- M. Shifman, N.G. Uraltsev, and A. Vainshtein, Minnesota preprint TPI-MINN-94/13-T and hep-ph/9405207, 1994.
- ARGUS: H. Albrecht *et al.*, *Z. Phys.* **C60**, 11 (1993).
- ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B234**, 409 (1990);
H. Albrecht *et al.*, *Phys. Lett.* **B255**, 297 (1991).
- CLEO: R. Fulton *et al.*, *Phys. Rev. Lett.* **64**, 16 (1990).
- CLEO: J. Bartelt *et al.*, *Phys. Rev. Lett.* **71**, 4111 (1993).
- C. Ramirez, J.F. Donoghue, and G. Burdman, *Phys. Rev.* **D41**, 1496 (1990);
M. Artuso, *Phys. Lett.* **B311**, 307 (1993).
- CLEO: F. Butler *et al.*, *Phys. Rev. Lett.* **69**, 2041 (1992).
- CLEO: For D^0 , D.S. Akerib *et al.*, *Phys. Rev. Lett.* **71**, 3070 (1993);
Mark III: For D^+ , J. Adler *et al.*, *Phys. Rev. Lett.* **60**, 89 (1988).
- ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B229**, 175 (1989).
- CLEO: R. Fulton *et al.*, *Phys. Rev.* **D43**, 651 (1991).
- ARGUS: H. Albrecht *et al.*, *Z. Phys.* **C57**, 533 (1993).
- ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B324**, 249 (1994).
- ARGUS: H. Albrecht *et al.*, *Phys. Lett.* **B275**, 195 (1992).
- A. Bean *et al.*, *Phys. Rev. Lett.* **70**, 2681 (1993).
- M.E. Luke, *Phys. Lett.* **B252**, 447 (1990).
- M. Neubert, *Phys. Lett.* **B264**, 455 (1991).
- UKQCD: S.P. Booth *et al.*, *Phys. Rev. Lett.* **72**, 462 (1994);
C. Bernard, Y. Shen, and A. Soni, *Phys. Lett.* **B317**, 164 (1993);
J. Mandula and M. Ogilvie, *Nucl. Phys. (Proc. Suppl.)* **B26**, 459 (1992).
- CLEO: D. Bortoletto *et al.*, *Phys. Rev. Lett.* **63**, 1667 (1989).
- CLEO: S. Sanghera *et al.*, *Phys. Rev.* **D47**, 791 (1993).
- J.G. Körner and G.A. Schuler, *Z. Phys.* **C38**, 511 (1988).
- M. Bauer and M. Wirbel, *Z. Phys.* **C42**, 671 (1989).
- N. Isgur and M. Wise, *Phys. Lett.* **B237**, 527 (1990).
- M. Neubert *et al.*, in *Heavy Flavours*, edited by A.J. Buras and M. Lindner, Advanced Series on Directions in High Energy Physics (World Scientific, Singapore, 1991).

 B^\pm

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

This section also includes measurements which do not identify the charge state of B .

 B^\pm MASS

These experiments actually measure the difference between half of E_{cm} and the B mass.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5278.7 ± 2.0 OUR FIT				
5278.7 ± 2.0 OUR AVERAGE				
5278.8 ± 0.54 ± 2.0	362	¹ ALAM 94	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
5278.3 ± 0.4 ± 2.0		¹ BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
5280.5 ± 1.0 ± 2.0		^{1,2} ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.6 ± 0.8 ± 2.0		¹ BEBEK 87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
5275.8 ± 1.3 ± 3.0	32	ALBRECHT 87C	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
5278.2 ± 1.8 ± 3.0	12	³ ALBRECHT 87D	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

¹ These experiments all report a common systematic error 2.0 MeV. We have artificially increased the systematic error to allow the experiments to be treated as independent measurements in our average. See "Treatment of Errors" section of the Introductory Text.

² ALBRECHT 90J assumes 10580 for $\Upsilon(4S)$ mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.

³ Found using fully reconstructed decays with $J/\psi(1S)$. ALBRECHT 87D assume $m_{\Upsilon(4S)} = 10577$ MeV.

 B^\pm MEAN LIFE

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.54 ± 0.11 OUR AVERAGE				
1.61 ± 0.16 ± 0.05	148	⁴ ABE 94D	CDF	$p\bar{p}$ at 1.8 TeV
1.30 ^{+0.33} _{-0.29} ± 0.16	92	⁵ ABREU 93D	DLPH	$e^+e^- \rightarrow Z$
1.56 ± 0.19 ± 0.13	134	⁶ ABREU 93G	DLPH	$e^+e^- \rightarrow Z$
1.51 ^{+0.30} _{-0.28} ± 0.12	59	⁵ ACTON 93C	OPAL	$e^+e^- \rightarrow Z$
1.47 ^{+0.22} _{-0.19} ± 0.15	77	⁵ BUSKULIC 93D	ALEP	$e^+e^- \rightarrow Z$

⁴ ABE 94D measured mean life using fully reconstructed decays.

⁵ Data analyzed using D/D^* anything event vertices.

⁶ ABREU 93G data analyzed using charged and neutral vertices.

Meson Full Listings

 B^\pm **B HADRON MEAN LIFE**

These measurements of the B mean life are averages over bottom particles produced, weighted by their semileptonic branching ratios, unless otherwise stated. Only the measurements at high energy are averaged since it is expected that the admixtures of B hadrons from Z decay and 1.8 TeV $p\bar{p}$ collisions should should not differ significantly.

VALUE (10^{-12} s)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.537 ± 0.021 OUR AVERAGE					
1.582 ± 0.012 ± 0.032			7 ABREU	94C DLPH	$e^+e^- \rightarrow Z$
1.46 ± 0.06 ± 0.06	5344		7 ABE	93J CDF	$p\bar{p}$ at 1.8 TeV
1.51 $^{+0.16}_{-0.14}$ ± 0.11	130		8 ACTON	93C OPAL	$e^+e^- \rightarrow Z$
1.523 ± 0.034 ± 0.038	5372		9 ACTON	93L OPAL	$e^+e^- \rightarrow Z$
1.535 ± 0.035 ± 0.028	7357		9 ADRIANI	93K L3	$e^+e^- \rightarrow Z$
1.50 ± 0.020 ± 0.050			10 BUSKULIC	93O ALEP	$e^+e^- \rightarrow Z$
1.35 $^{+0.19}_{-0.17}$ ± 0.005			11 BUSKULIC	92G ALEP	$e^+e^- \rightarrow Z$
1.32 $^{+0.31}_{-0.25}$ ± 0.15	37		12 ALEXANDER	91G OPAL	$e^+e^- \rightarrow Z$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.23 $^{+0.14}_{-0.13}$ ± 0.15	188		13 ABREU	93D DLPH	Sup. by ABREU 94C
1.49 ± 0.11 ± 0.12	253		14 ABREU	93G DLPH	Sup. by ABREU 94C
1.28 ± 0.10			15 ABREU	92 DLPH	Sup. by ABREU 94C
1.37 ± 0.07 ± 0.06	1354		16 ACTON	92 OPAL	Sup. by ACTON 93L
1.49 ± 0.03 ± 0.06			17 BUSKULIC	92F ALEP	Sup. by BUSKULIC 93O
1.32 ± 0.08 ± 0.09	1386		18 ADEVA	91H L3	Sup. by ADRIANI 93K
1.29 ± 0.06 ± 0.10	2973		19 DECAMP	91C ALEP	Sup. by BUSKULIC 92F
1.36 $^{+0.25}_{-0.23}$			20 HAGEMANN	90 JADE	$E_{cm}^{ee} = 35$ GeV
1.13 ± 0.15			21 LYONS	90 RVUE	
1.35 ± 0.10 ± 0.24			BRAUNSCH...	89B TASS	$E_{cm}^{ee} = 35$ GeV
0.98 ± 0.12 ± 0.13			ONG	89 MRK2	$E_{cm}^{ee} = 29$ GeV
1.17 $^{+0.27}_{-0.22}$ $^{+0.17}_{-0.16}$			KLEM	88 DLCO	$E_{cm}^{ee} = 29$ GeV
1.29 ± 0.20 ± 0.21			22 ASH	87 MAC	$E_{cm}^{ee} = 29$ GeV
1.02 $^{+0.42}_{-0.39}$	301		23 BROM	87 HRS	$E_{cm}^{ee} = 29$ GeV
1.8 $^{+0.5}_{-0.4}$ ± 0.4	25		BARTEL	86B JADE	$E_{cm}^{ee} = 35$ GeV
1.83 $^{+0.38}_{-0.37}$ $^{+0.37}_{-0.34}$			ALTHOFF	84H TASS	$E_{cm}^{ee} = 30\text{--}46.8$ GeV
1.16 $^{+0.37}_{-0.34}$ ± 0.23	46		KLEM	84 DLCO	Repl. by KLEM 88
1.8 ± 0.6 ± 0.4			FERNANDEZ	83B MAC	$E_{cm}^{ee} = 29$ GeV
1.20 $^{+0.45}_{-0.36}$ ± 0.30			LOCKYER	83 MRK2	Repl. by ONG 89
<1.4	95		BARTEL	82C JADE	e^+e^- , average $E_{cm} = 34$ GeV

7 ABE 93J analyzed using $J/\psi(1S) \rightarrow \mu\mu$ vertices.

8 ACTON 93C analysed using D/D^* anything event vertices.

9 ACTON 93L and ADRIANI 93K analyzed using lepton (e and m) impact parameter at Z .

10 BUSKULIC 93O analyzed using dipole method plus lepton (e and μ) impact parameter at Z .

11 BUSKULIC 92G use $J/\psi(1S)$ tags to measure the average b lifetime. This is comparable to other methods only if the $J/\psi(1S)$ branching fractions of the different b -flavored hadrons are in the same ratio.

12 Using $Z \rightarrow J/\psi(1S)X$, $J/\psi(1S) \rightarrow \ell^+\ell^-$, ALEXANDER 91G determined the average lifetime for an admixture of B hadrons from the decay point of the $J/\psi(1S)$.

13 ABREU 93D data analyzed using D/D^* anything event vertices.

14 ABREU 93G data analyzed using charged and neutral vertices.

15 ABREU 92 is combined result of muon and hadron impact parameter analyses. Hadron tracks gave $(12.7 \pm 0.4 \pm 1.2) \times 10^{-13}$ s for an admixture of B species weighted by production fraction and mean charge multiplicity, while muon tracks gave $(13.0 \pm 1.0 \pm 0.8) \times 10^{-13}$ s for an admixture weighted by production fraction and semileptonic branching fraction.

16 ACTON 92 is combined result of muon and electron impact parameter analyses.

17 BUSKULIC 92F uses the lepton impact parameter distribution for data from the 1991 run.

18 Using $Z \rightarrow e^+X$ or μ^+X , ADEVA 91H determined the average lifetime for an admixture of B hadrons from the impact parameter distribution of the lepton.

19 Using $Z \rightarrow eX$ or μX , DECAMP 91C determines the average lifetime for an admixture of B hadrons from the signed impact parameter distribution of the lepton.

20 HAGEMANN 90 uses electrons and muons in an impact parameter analysis.

21 LYONS 90 combine the results of the B lifetime measurements of ONG 89, BRAUN-SCHWEIG 89B, KLEM 88, and ASH 87, and JADE data by private communication. They use statistical techniques which include variation of the error with the mean life, and possible correlations between the systematic errors. This result is not independent of the measured results used in our average.

22 We have combined an overall scale error of 15% in quadrature with the systematic error of ± 0.7 to obtain ± 2.1 systematic error.

23 Statistical and systematic errors were combined by BROM 87.

B⁺ DECAY MODES

B^- modes are charge conjugates of the modes below.

Only data from $\Upsilon(4S)$ decays are used for branching fractions, with rare exceptions. The branching fractions listed below assume a 50:50 $B^0\bar{B}^0:B^+B^-$ production ratio at the $\Upsilon(4S)$. We have attempted to bring older measurements up to date by rescaling their assumed $\Upsilon(4S)$ production ratio to 50:50 and their assumed D, D_s, D^* , and ψ branching ratios to current values whenever this would effect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Semileptonic modes		
Γ_1 $B^+ \rightarrow \bar{D}^0 \ell^+ \nu$	[a] (1.6 ± 0.7) %	
Γ_2 $B^+ \rightarrow \bar{D}^*(2007)^0 \ell^+ \nu$	[a] (6.6 ± 2.2) %	
Γ_3 $B^+ \rightarrow \pi^0 e^+ \nu_e$	< 2.2	$\times 10^{-3}$ CL=90%
Γ_4 $B^+ \rightarrow \omega \ell^+ \nu_\ell$	[a] < 2.1	$\times 10^{-4}$ CL=90%
Γ_5 $B^+ \rightarrow \omega \mu^+ \nu_\mu$	seen	
Γ_6 $B^+ \rightarrow \rho^0 \ell^+ \nu_\ell$	[a] < 2.1	$\times 10^{-4}$ CL=90%
D, D*, or D_s modes		
Γ_7 $B^+ \rightarrow \bar{D}^0 \pi^+$	(5.3 ± 0.5) $\times 10^{-3}$	
Γ_8 $B^+ \rightarrow \bar{D}^0 \rho^+$	(1.34 ± 0.18) %	
Γ_9 $B^+ \rightarrow \bar{D}^0 \pi^+ \pi^+ \pi^-$	(1.1 ± 0.4) %	
Γ_{10} $B^+ \rightarrow \bar{D}^0 \pi^+ \pi^+ \pi^-$ nonreso-	(5 ± 4) $\times 10^{-3}$	
	nant	
Γ_{11} $B^+ \rightarrow \bar{D}^0 \pi^+ \rho^0$	(4.2 ± 3.0) $\times 10^{-3}$	
Γ_{12} $B^+ \rightarrow \bar{D}^0 a_1(1260)^+$	(5 ± 4) $\times 10^{-3}$	
Γ_{13} $B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+$	(2.1 ± 0.6) $\times 10^{-3}$	
Γ_{14} $B^+ \rightarrow D^- \pi^+ \pi^+$	< 1.4	$\times 10^{-3}$ CL=90%
Γ_{15} $B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+$	(5.2 ± 0.8) $\times 10^{-3}$	
Γ_{16} $B^+ \rightarrow \bar{D}^*(2007)^0 \rho^+$	(1.55 ± 0.31) %	
Γ_{17} $B^+ \rightarrow \bar{D}^*(2007)^0 \pi^+ \pi^+ \pi^-$	(9.4 ± 2.6) $\times 10^{-3}$	
Γ_{18} $B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^0$	(1.5 ± 0.7) %	
Γ_{19} $B^+ \rightarrow D^*(2010)^- \pi^+ \pi^+ \pi^+ \pi^-$	< 1	% CL=90%
Γ_{20} $B^+ \rightarrow \bar{D}_1^+(2420)^0 \pi^+$	(1.1 ± 0.5) $\times 10^{-3}$	
Γ_{21} $B^+ \rightarrow \bar{D}_1^+(2420)^0 \rho^+$	< 1.4	$\times 10^{-3}$ CL=90%
Γ_{22} $B^+ \rightarrow \bar{D}_2^+(2460)^0 \pi^+$	< 1.3	$\times 10^{-3}$ CL=90%
Γ_{23} $B^+ \rightarrow \bar{D}_2^+(2460)^0 \rho^+$	< 4.7	$\times 10^{-3}$ CL=90%
Γ_{24} $B^+ \rightarrow \bar{D}^0 D_s^+$	(1.7 ± 0.6) %	
Γ_{25} $B^+ \rightarrow \bar{D}^0 D_s^{*+}$	(1.2 ± 1.0) %	
Γ_{26} $B^+ \rightarrow \bar{D}^*(2007)^0 D_s^+$	(1.0 ± 0.7) %	
Γ_{27} $B^+ \rightarrow \bar{D}^*(2007)^0 D_s^{*+}$	(2.4 ± 1.3) %	
Γ_{28} $B^+ \rightarrow D_s^+ \pi^0$	< 2.1	$\times 10^{-4}$ CL=90%
Γ_{29} $B^+ \rightarrow D_s^{*+} \pi^0$	< 3.4	$\times 10^{-4}$ CL=90%
Γ_{30} $B^+ \rightarrow D_s^+ \eta$	< 5	$\times 10^{-4}$ CL=90%
Γ_{31} $B^+ \rightarrow D_s^{*+} \eta$	< 8	$\times 10^{-4}$ CL=90%
Γ_{32} $B^+ \rightarrow D_s^+ \rho^0$	< 4	$\times 10^{-4}$ CL=90%
Γ_{33} $B^+ \rightarrow D_s^{*+} \rho^0$	< 5	$\times 10^{-4}$ CL=90%
Γ_{34} $B^+ \rightarrow D_s^+ \omega$	< 5	$\times 10^{-4}$ CL=90%
Γ_{35} $B^+ \rightarrow D_s^{*+} \omega$	< 7	$\times 10^{-4}$ CL=90%
Γ_{36} $B^+ \rightarrow D_s^+ a_1(1260)^0$	< 2.3	$\times 10^{-3}$ CL=90%
Γ_{37} $B^+ \rightarrow D_s^{*+} a_1(1260)^0$	< 1.7	$\times 10^{-3}$ CL=90%
Γ_{38} $B^+ \rightarrow D_s^+ \phi$	< 3.3	$\times 10^{-4}$ CL=90%
Γ_{39} $B^+ \rightarrow D_s^{*+} \phi$	< 4	$\times 10^{-4}$ CL=90%
Γ_{40} $B^+ \rightarrow D_s^+ \bar{K}^0$	< 1.1	$\times 10^{-3}$ CL=90%
Γ_{41} $B^+ \rightarrow D_s^{*+} \bar{K}^0$	< 1.2	$\times 10^{-3}$ CL=90%
Γ_{42} $B^+ \rightarrow D_s^+ \bar{K}^*(892)^0$	< 5	$\times 10^{-4}$ CL=90%
Γ_{43} $B^+ \rightarrow D_s^{*+} \bar{K}^*(892)^0$	< 5	$\times 10^{-4}$ CL=90%
Γ_{44} $B^+ \rightarrow D_s^+ \pi^+ K^+$	< 9	$\times 10^{-4}$ CL=90%
Γ_{45} $B^+ \rightarrow D_s^{*+} \pi^+ K^+$	< 1.2	$\times 10^{-3}$ CL=90%
Γ_{46} $B^+ \rightarrow D_s^+ \pi^+ K^*(892)^+$	< 7	$\times 10^{-3}$ CL=90%
Γ_{47} $B^+ \rightarrow D_s^{*+} \pi^+ K^*(892)^+$	< 9	$\times 10^{-3}$ CL=90%

Charmonium modes

Γ48	$B^+ \rightarrow J/\psi(1S)K^+$	$(1.02 \pm 0.14) \times 10^{-3}$	
Γ49	$B^+ \rightarrow J/\psi(1S)K^+\pi^+\pi^-$	$(1.4 \pm 0.6) \times 10^{-3}$	
Γ50	$B^+ \rightarrow J/\psi(1S)K^*(892)^+$	$(1.7 \pm 0.5) \times 10^{-3}$	
Γ51	$B^+ \rightarrow \psi(2S)K^+$	$(6.9 \pm 3.1) \times 10^{-4}$	S=1.3
Γ52	$B^+ \rightarrow \psi(2S)K^*(892)^+$	$< 3.0 \times 10^{-3}$	CL=90%
Γ53	$B^+ \rightarrow \psi(2S)K^*(892)^+\pi^+\pi^-$	$(1.9 \pm 1.2) \times 10^{-3}$	
Γ54	$B^+ \rightarrow \chi_{c1}(1P)K^+$	$(1.0 \pm 0.4) \times 10^{-3}$	
Γ55	$B^+ \rightarrow \chi_{c1}(1P)K^*(892)^+$	$< 2.1 \times 10^{-3}$	CL=90%

K or K* modes

Γ56	$B^+ \rightarrow K^0\pi^+$	$< 1.0 \times 10^{-4}$	CL=90%
Γ57	$B^+ \rightarrow K^*(892)^0\pi^+$	$< 1.5 \times 10^{-4}$	CL=90%
Γ58	$B^+ \rightarrow K^+\pi^-\pi^+$ (no charm)	$< 1.9 \times 10^{-4}$	CL=90%
Γ59	$B^+ \rightarrow K_1(1400)^0\pi^+$	$< 2.6 \times 10^{-3}$	CL=90%
Γ60	$B^+ \rightarrow K_2^*(1430)^0\pi^+$	$< 6.8 \times 10^{-4}$	CL=90%
Γ61	$B^+ \rightarrow K^+\rho^0$	$< 8 \times 10^{-5}$	CL=90%
Γ62	$B^+ \rightarrow K^*(892)^+\pi^+\pi^-$	$< 1.1 \times 10^{-3}$	CL=90%
Γ63	$B^+ \rightarrow K^*(892)^+\rho^0$	$< 9.0 \times 10^{-4}$	CL=90%
Γ64	$B^+ \rightarrow K_1(1400)^+\rho^0$	$< 7.8 \times 10^{-4}$	CL=90%
Γ65	$B^+ \rightarrow K_2^*(1430)^+\rho^0$	$< 1.5 \times 10^{-3}$	CL=90%
Γ66	$B^+ \rightarrow K^+K^-K^+$	$< 3.5 \times 10^{-4}$	CL=90%
Γ67	$B^+ \rightarrow K^+\phi$	$< 9 \times 10^{-5}$	CL=90%
Γ68	$B^+ \rightarrow K^*(892)^+K^+K^-$	$< 1.6 \times 10^{-3}$	CL=90%
Γ69	$B^+ \rightarrow K^*(892)^+\phi$	$< 1.3 \times 10^{-3}$	CL=90%
Γ70	$B^+ \rightarrow K_1(1400)^+\phi$	$< 1.1 \times 10^{-3}$	CL=90%
Γ71	$B^+ \rightarrow K_2^*(1430)^+\phi$	$< 3.4 \times 10^{-3}$	CL=90%
Γ72	$B^+ \rightarrow K^+\bar{f}_0(980)$	$< 8 \times 10^{-5}$	CL=90%
Γ73	$B^+ \rightarrow K^*(892)^+\gamma$	$(5.7 \pm 3.3) \times 10^{-5}$	
Γ74	$B^+ \rightarrow K_1(1270)^+\gamma$	$< 7.3 \times 10^{-3}$	CL=90%
Γ75	$B^+ \rightarrow K_1(1400)^+\gamma$	$< 2.2 \times 10^{-3}$	CL=90%
Γ76	$B^+ \rightarrow K_2^*(1430)^+\gamma$	$< 1.4 \times 10^{-3}$	CL=90%
Γ77	$B^+ \rightarrow K^*(1680)^+\gamma$	$< 1.9 \times 10^{-3}$	CL=90%
Γ78	$B^+ \rightarrow K_3^*(1780)^+\gamma$	$< 5.5 \times 10^{-3}$	CL=90%
Γ79	$B^+ \rightarrow K_4^*(2045)^+\gamma$	$< 9.9 \times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ80	$B^+ \rightarrow \pi^+\pi^0$	$< 2.4 \times 10^{-4}$	CL=90%
Γ81	$B^+ \rightarrow \pi^+\pi^+\pi^-$	$< 1.9 \times 10^{-4}$	CL=90%
Γ82	$B^+ \rightarrow \rho^0\pi^+$	$< 1.5 \times 10^{-4}$	CL=90%
Γ83	$B^+ \rightarrow \pi^+\bar{f}_0(980)$	$< 1.4 \times 10^{-4}$	CL=90%
Γ84	$B^+ \rightarrow \pi^+\bar{f}_2(1270)$	$< 2.4 \times 10^{-4}$	CL=90%
Γ85	$B^+ \rightarrow \pi^+\pi^0\pi^0$	$< 8.9 \times 10^{-4}$	CL=90%
Γ86	$B^+ \rightarrow \rho^+\pi^0$	$< 5.5 \times 10^{-4}$	CL=90%
Γ87	$B^+ \rightarrow \pi^+\pi^-\pi^+\pi^0$	$< 4.0 \times 10^{-3}$	CL=90%
Γ88	$B^+ \rightarrow \rho^+\rho^0$	$< 1.0 \times 10^{-3}$	CL=90%
Γ89	$B^+ \rightarrow a_1(1260)^+\pi^0$	$< 1.7 \times 10^{-3}$	CL=90%
Γ90	$B^+ \rightarrow a_1(1260)^0\pi^+$	$< 9.0 \times 10^{-4}$	CL=90%
Γ91	$B^+ \rightarrow \omega\pi^+$	$< 4.0 \times 10^{-4}$	CL=90%
Γ92	$B^+ \rightarrow \eta\pi^+$	$< 7.0 \times 10^{-4}$	CL=90%
Γ93	$B^+ \rightarrow \pi^+\pi^+\pi^+\pi^-\pi^-$	$< 8.6 \times 10^{-4}$	CL=90%
Γ94	$B^+ \rightarrow \rho^0 a_1(1260)^+$	$< 6.2 \times 10^{-4}$	CL=90%
Γ95	$B^+ \rightarrow \rho^0 a_2(1320)^+$	$< 7.2 \times 10^{-4}$	CL=90%
Γ96	$B^+ \rightarrow \pi^+\pi^+\pi^+\pi^-\pi^-\pi^0$	$< 6.3 \times 10^{-3}$	CL=90%
Γ97	$B^+ \rightarrow a_1(1260)^+ a_1(1260)^0$	$< 1.3 \%$	CL=90%

Baryon modes

Γ98	$B^+ \rightarrow p\bar{p}\pi^+$	$< 1.6 \times 10^{-4}$	CL=90%
Γ99	$B^+ \rightarrow p\bar{p}\pi^+\pi^+\pi^-$	$< 5.2 \times 10^{-4}$	CL=90%
Γ100	$B^+ \rightarrow p\bar{\Lambda}$	$< 6 \times 10^{-5}$	CL=90%
Γ101	$B^+ \rightarrow p\bar{\Lambda}\pi^+\pi^-$	$< 2.0 \times 10^{-4}$	CL=90%
Γ102	$B^+ \rightarrow \Delta^0 p$	$< 3.8 \times 10^{-4}$	CL=90%
Γ103	$B^+ \rightarrow \Delta^+\bar{p}$	$< 1.5 \times 10^{-4}$	CL=90%

Lepton Family number (LF) or Lepton number (L) violating modes, or $\Delta B = 1$ weak neutral current (BI) modes

Γ104	$B^+ \rightarrow \pi^+e^+e^-$	BI	$< 3.9 \times 10^{-3}$	CL=90%
Γ105	$B^+ \rightarrow \pi^+\mu^+\mu^-$	BI	$< 9.1 \times 10^{-3}$	CL=90%
Γ106	$B^+ \rightarrow K^+e^+e^-$	BI	$< 6 \times 10^{-5}$	CL=90%
Γ107	$B^+ \rightarrow K^+\mu^+\mu^-$	BI	$< 1.7 \times 10^{-4}$	CL=90%
Γ108	$B^+ \rightarrow K^*(892)^+\mu^+\mu^-$	BI	$< 6.9 \times 10^{-4}$	CL=90%
Γ109	$B^+ \rightarrow K^*(892)^+\mu^+\mu^-$	BI	$< 1.2 \times 10^{-3}$	CL=90%
Γ110	$B^+ \rightarrow \pi^+e^+\mu^-$	LF	$< 6.4 \times 10^{-3}$	CL=90%
Γ111	$B^+ \rightarrow \pi^+e^-\mu^+$	LF	$< 6.4 \times 10^{-3}$	CL=90%

Γ112	$B^+ \rightarrow K^+e^+\mu^-$	LF	$< 6.4 \times 10^{-3}$	CL=90%
Γ113	$B^+ \rightarrow K^+e^-\mu^+$	LF	$< 6.4 \times 10^{-3}$	CL=90%
Γ114	$B^+ \rightarrow \pi^-e^+e^+$	L	$< 3.9 \times 10^{-3}$	CL=90%
Γ115	$B^+ \rightarrow \pi^-\mu^+\mu^+$	L	$< 9.1 \times 10^{-3}$	CL=90%
Γ116	$B^+ \rightarrow \pi^-e^+\mu^+$	L	$< 6.4 \times 10^{-3}$	CL=90%
Γ117	$B^+ \rightarrow K^-e^+e^+$	L	$< 3.9 \times 10^{-3}$	CL=90%
Γ118	$B^+ \rightarrow K^-\mu^+\mu^+$	L	$< 9.1 \times 10^{-3}$	CL=90%
Γ119	$B^+ \rightarrow K^-e^+\mu^+$	L	$< 6.4 \times 10^{-3}$	CL=90%

B DECAY MODES

\bar{B} modes are charge conjugates of the modes below.

For the following modes, the charge of B was not determined. The measurements are for an admixture of B mesons at the T(4S) unless otherwise indicated by a footnote and a "B" instead of "B" in the initial state.

Semileptonic and leptonic modes

Γ120	$B \rightarrow e^+\nu_e$ anything	[b]	$(10.4 \pm 0.4) \%$	S=1.3
Γ121	$B \rightarrow \bar{D}^*(2010)e^+\nu_e$		$(7.0 \pm 2.3) \%$	
Γ122	$B \rightarrow \bar{p}e^+\nu_e$ anything		$< 1.6 \times 10^{-3}$	CL=90%
Γ123	$B \rightarrow \mu^+\nu_\mu$ anything	[b]	$(10.3 \pm 0.5) \%$	
Γ124	$B \rightarrow \ell^+\nu_\ell$ anything	[a,b]	$(10.43 \pm 0.24) \%$	
Γ125	$B \rightarrow D^-\ell^+\nu_\ell$ anything	[a]	$(2.7 \pm 0.8) \%$	
Γ126	$B \rightarrow \bar{D}^0\ell^+\nu_\ell$ anything	[a]	$(7.0 \pm 1.4) \%$	
Γ127	$B \rightarrow D^{**}\ell^+\nu_\ell$	[a,c]	$(2.7 \pm 0.7) \%$	
Γ128	$B \rightarrow D_s^-\ell^+\nu_\ell$ anything	[a]	$< 9 \times 10^{-3}$	CL=90%
Γ129	$B \rightarrow D_s^-\ell^+\nu_\ell K^+$ anything	[a]	$< 6 \times 10^{-3}$	CL=90%
Γ130	$B \rightarrow D_s^-\ell^+\nu_\ell K^0$ anything	[a]	$< 9 \times 10^{-3}$	CL=90%
Γ131	$B \rightarrow \ell^+\nu_\ell$ noncharmed	[a]		
Γ132	$B \rightarrow K^+\ell^+\nu_\ell$ anything	[a]	$(5.6 \pm 1.0) \%$	
Γ133	$B \rightarrow K^-\ell^+\nu_\ell$ anything	[a]	$(1.0 \pm 0.6) \%$	
Γ134	$B \rightarrow K^0/\bar{K}^0\ell^+\nu_\ell$ anything	[a]	$(4.1 \pm 0.8) \%$	
Γ135	$\bar{b} \rightarrow \tau^+\nu_\tau$ anything	[d]	$(4.1 \pm 1.0) \%$	

D, D*, or D_s modes

Γ136	$B \rightarrow D^-$ anything		$(26 \pm 4) \%$	
Γ137	$B \rightarrow \bar{D}^0$ anything		$(54 \pm 6) \%$	
Γ138	$B \rightarrow D^{*+}(2010)$ anything		$(23 \pm 4) \%$	S=1.4
Γ139	$B \rightarrow D_s^{*+}$ anything	[e]	$(8.9 \pm 1.1) \%$	
Γ140	$B \rightarrow D_s^- D, D_s^* D, D_s D^*,$ or $D_s^* D^*$	[e]	$(5.0 \pm 0.9) \%$	
Γ141	$B \rightarrow D^{*+}(2010)\gamma$		$< 1.1 \times 10^{-3}$	CL=90%
Γ142	$B \rightarrow D_s^{*+}\pi^-, D_s^{*+}\pi^-, D_s^+\rho^-,$ $D_s^{*+}\rho^-, D_s^+\pi^0, D_s^+\pi^0,$ $D_s^+\eta, D_s^+\eta, D_s^+\rho^0,$ $D_s^{*+}\rho^0, D_s^+\omega, D_s^{*+}\omega$		$< 5 \times 10^{-4}$	CL=90%

Charmonium modes

Γ143	$B \rightarrow J/\psi(1S)$ anything		$(1.30 \pm 0.17) \%$	
Γ144	$B \rightarrow \psi(2S)$ anything		$(4.6 \pm 2.0) \times 10^{-3}$	
Γ145	$B \rightarrow \chi_{c1}(1P)$ anything		$(1.1 \pm 0.4) \%$	

K or K* modes

Γ146	$B \rightarrow K^\pm$ anything	[e]	$(85 \pm 11) \%$	
Γ147	$B \rightarrow K^+$ anything			
Γ148	$B \rightarrow K^-$ anything			
Γ149	$B \rightarrow K^0/\bar{K}^0$ anything		$(63 \pm 8) \%$	
Γ150	$b \rightarrow s\gamma$	[f]	$< 1.2 \times 10^{-3}$	CL=90%
Γ151	$B \rightarrow K^*(892)\gamma$		$< 2.4 \times 10^{-4}$	CL=90%
Γ152	$B \rightarrow K_1(1400)\gamma$		$< 4.1 \times 10^{-4}$	CL=90%
Γ153	$B \rightarrow K_2^*(1430)\gamma$		$< 8.3 \times 10^{-4}$	CL=90%
Γ154	$B \rightarrow K_2(1770)\gamma$		$< 1.2 \times 10^{-3}$	CL=90%
Γ155	$B \rightarrow K_3^*(1780)\gamma$		$< 3.0 \times 10^{-3}$	CL=90%
Γ156	$B \rightarrow K_4^*(2045)\gamma$		$< 1.0 \times 10^{-3}$	CL=90%

Light unflavored meson modes

Γ157	$B \rightarrow \phi$ anything		$(2.3 \pm 0.8) \%$	
------	-------------------------------	--	--------------------	--

Baryon modes

Γ158	$B \rightarrow$ charmed-baryon anything		$(6.4 \pm 1.1) \%$	
Γ159	$B \rightarrow \bar{\Sigma}_c^-$ anything		$(4.8 \pm 2.5) \times 10^{-3}$	
Γ160	$B \rightarrow \bar{\Sigma}_c^0$ anything		$< 1.1 \%$	CL=90%
Γ161	$B \rightarrow \bar{\Sigma}_c^+ N(N = p \text{ or } n)$		$(5.3 \pm 2.5) \times 10^{-3}$	
Γ162	$B \rightarrow \bar{\Sigma}_c^0 N(N = p \text{ or } n)$		$< 1.7 \times 10^{-3}$	CL=90%
Γ163	$B \rightarrow p$ anything + \bar{p} anything		$(8.0 \pm 0.5) \%$	

Meson Full Listings

 B^\pm

Γ_{164}	$B \rightarrow \rho(\text{direct}) \text{ anything} + \bar{\rho}(\text{direct}) \text{ anything}$	$(5.6 \pm 0.7)\%$	
Γ_{165}	$B \rightarrow \Lambda \text{ anything} + \bar{\Lambda} \text{ anything}$	$(4.0 \pm 0.5)\%$	
Γ_{166}	$B \rightarrow \Xi^- \text{ anything} + \bar{\Xi}^+ \text{ anything}$	$(2.7 \pm 0.6) \times 10^{-3}$	
Γ_{167}	$B \rightarrow \text{baryons anything}$	$(6.8 \pm 0.6)\%$	
Γ_{168}	$B \rightarrow p\bar{p} \text{ anything}$	$(2.47 \pm 0.23)\%$	
Γ_{169}	$B \rightarrow \Lambda\bar{\Lambda} \text{ anything} + \bar{\Lambda}p \text{ anything}$	$(2.5 \pm 0.4)\%$	
Γ_{170}	$B \rightarrow \Lambda\bar{\Lambda} \text{ anything}$	< 5	$\times 10^{-3}$ CL=90%

 $\Delta B = 1$ weak neutral current ($B1$) modes

Γ_{171}	$\bar{b} \rightarrow e^+ e^- \text{ anything}$	$B1$	$[f] < 2.4$	$\times 10^{-3}$
Γ_{172}	$\bar{b} \rightarrow \mu^+ \mu^- \text{ anything}$	$B1$	$[f] < 5.0$	$\times 10^{-5}$ CL=90%

[a] ℓ indicates e or μ mode, not sum over modes.

[b] These values are model dependent. See note on "Semileptonic Decays" in these Full Listings.

[c] D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances.[d] B^0 , B^+ , B_s^0 , and B baryon states not separated.

[e] The value is for the sum of the charge states indicated.

[f] B^0 , B^+ , and B_s^0 not separated. B^+ BRANCHING RATIOS

$\Gamma(\bar{D}^0 \ell^+ \nu)/\Gamma_{\text{total}}$				Γ_1/Γ
	$\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$			
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.016 \pm 0.006 \pm 0.003$	24 FULTON	91 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

24 FULTON 91 assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.

$\Gamma(\bar{D}^*(2007)^0 \ell^+ \nu)/\Gamma_{\text{total}}$				Γ_2/Γ
	$\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$0.066 \pm 0.016 \pm 0.015$		25 ALBRECHT	92C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	398	26 SANGHERA	93 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.041 ± 0.008		27 FULTON	91 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

25 ALBRECHT 92C reports $0.058 \pm 0.014 \pm 0.013$. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$.26 Combining $\bar{D}^{*0} \ell^+ \nu_\ell$ and $\bar{D}^{*+} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4 * (\Gamma^- - \Gamma^+)/\Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure $V, A_1,$ and A_2 , the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.27 Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at the $\Upsilon(4S)$. Uncorrected for D and D^* branching ratio assumptions.

$\Gamma(\pi^0 e^+ \nu_e)/\Gamma_{\text{total}}$				Γ_3/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0022	90	ANTREASNYAN 90B	CBAL	$e^+ e^- \rightarrow \Upsilon(4S)$

$\Gamma(\omega \ell^+ \nu_\ell)/\Gamma_{\text{total}}$				Γ_4/Γ
	$\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1 \times 10^{-4}$	90	28 BEAN	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

28 BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $< (1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \omega \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\omega \mu^+ \nu_\mu)/\Gamma_{\text{total}}$				Γ_5/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
seen	29 ALBRECHT	91C ARG		

29 In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(\rho^0 \ell^+ \nu_\ell)/\Gamma_{\text{total}}$				Γ_6/Γ
	$\ell = e \text{ or } \mu, \text{ not sum over } e \text{ and } \mu \text{ modes.}$			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 2.1 \times 10^{-4}$	90	30 BEAN	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

30 BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\omega^0 \ell^+ \nu_\ell)$ and $\Gamma(\rho^- \ell^+ \nu_\ell)$ with this result, they obtain a limit $< (1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow \rho^0 \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\bar{D}^0 \pi^+)/\Gamma_{\text{total}}$				Γ_7/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0053 ± 0.0005	OUR AVERAGE			
$0.0055 \pm 0.0004 \pm 0.0005$	304	31 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.0050 \pm 0.0007 \pm 0.0006$	54	32 BOROLETTO	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.0054 \pm 0.0018 \pm 0.0012$	14	33 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.0020 \pm 0.0008 \pm 0.0006$	12	32 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.0019 \pm 0.0010 \pm 0.0006$	7	34 ALBRECHT	88K ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

31 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.32 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

33 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BOROLETTO 92.

34 ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55. Superseded by ALBRECHT 90J.

$\Gamma(\bar{D}^0 \rho^+)/\Gamma_{\text{total}}$				Γ_8/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0134 ± 0.0018	OUR AVERAGE			
$0.0135 \pm 0.0012 \pm 0.0015$	212	35 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
$0.013 \pm 0.004 \pm 0.004$	19	36 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$0.021 \pm 0.008 \pm 0.009$	10	37 ALBRECHT	88K ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
-----------------------------	----	-------------	---------	------------------------------------

35 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.36 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .37 ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ ratio is 45:55.

$\Gamma(\bar{D}^0 \pi^+ \pi^+ \pi^-)/\Gamma_{\text{total}}$				Γ_9/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.0115 \pm 0.0029 \pm 0.0021$	38 BOROLETTO	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

38 BOROLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 \pi^+ \pi^+ \pi^- \text{ nonresonant})/\Gamma_{\text{total}}$				Γ_{10}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.0051 \pm 0.0034 \pm 0.0023$	39 BOROLETTO	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

39 BOROLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 \pi^+ \rho^0)/\Gamma_{\text{total}}$				Γ_{11}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.0042 \pm 0.0023 \pm 0.0020$	40 BOROLETTO	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

40 BOROLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(\bar{D}^0 \rho_1(1260)^+)/\Gamma_{\text{total}}$				Γ_{12}/Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
$0.0045 \pm 0.0019 \pm 0.0031$	41 BOROLETTO	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

41 BOROLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^*(2010)^- \pi^+ \pi^+)/\Gamma_{\text{total}}$				Γ_{13}/Γ	
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0021 ± 0.0006	OUR AVERAGE				
$0.0019 \pm 0.0007 \pm 0.0003$		14	42 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

42 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.43 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses the Mark III branching fractions for the D .

44 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BOROLETTO 92.

45 BOROLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$. The authors also find the product branching fraction into $D^{**} \pi$ followed by $D^{**} \rightarrow D^*(2010) \pi$ to be $0.0014 \pm 0.0008 \pm 0.0003$ where D^{**} represents all orbitally excited D mesons.46 ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^{-}\pi^{+}\pi^{+})/\Gamma_{\text{total}}$				Γ_{14}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.0014	90		47 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.007 90 48 BORTOLETTO92 CLEO $e^{+}e^{-} \rightarrow \Upsilon(4S)$

0.0025^{+0.0041+0.0024}_{-0.0023-0.0008} 1 49 BEBEK 87 CLEO $e^{+}e^{-} \rightarrow \Upsilon(4S)$

47 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the Mark III $B(D^{+} \rightarrow K^{-}\pi^{+}\pi^{+})$.

48 BORTOLETTO 92 assumes equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_{s1}^{*}(2340)\pi$ followed by $D_{s1}^{*}(2340) \rightarrow D\pi$ is < 0.005 at 90%CL and into $D_{s2}^{*}(2460)$ followed by $D_{s2}^{*}(2460) \rightarrow D\pi$ is < 0.004 at 90%CL.

49 BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^{0}\bar{B}^{0}$. $B(D^{-} \rightarrow K^{+}\pi^{-}\pi^{-}) = (9.1 \pm 1.3 \pm 0.4)\%$ is assumed.

$\Gamma(D^{*}(2007)^{0}\pi^{+})/\Gamma_{\text{total}}$				Γ_{15}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.0052 \pm 0.0008 OUR AVERAGE				
0.0052 \pm 0.0007 \pm 0.0007	71		50 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$
0.0072 \pm 0.0018 \pm 0.0016			51 BORTOLETTO92	CLEO $e^{+}e^{-} \rightarrow \Upsilon(4S)$
0.0040 \pm 0.0014 \pm 0.0012	9		51 ALBRECHT 90J	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0027 \pm 0.0044 52 BEBEK 87 CLEO $e^{+}e^{-} \rightarrow \Upsilon(4S)$

50 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the CLEO II $B(D^{*}(2007)^{0} \rightarrow D^{0}\pi^{0})$ and absolute $B(D^{0} \rightarrow K^{-}\pi^{+})$ and the PDG 1992 $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{0})/B(D^{0} \rightarrow K^{-}\pi^{+})$ and $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{+}\pi^{-})/B(D^{0} \rightarrow K^{-}\pi^{+})$.

51 Assumes equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^{*}(2010)$.

52 This is a derived branching ratio, using the inclusive pion spectrum and other two-body B decays. BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^{0}\bar{B}^{0}$.

$\Gamma(D^{*}(2007)^{0}\rho^{+})/\Gamma_{\text{total}}$				Γ_{16}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.0155 \pm 0.0031 OUR AVERAGE				
0.0168 \pm 0.0021 \pm 0.0028	86		53 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$
0.010 \pm 0.006 \pm 0.004	7		54 ALBRECHT 90J	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

53 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the CLEO II $B(D^{*}(2007)^{0} \rightarrow D^{0}\pi^{0})$ and absolute $B(D^{0} \rightarrow K^{-}\pi^{+})$ and the PDG 1992 $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{0})/B(D^{0} \rightarrow K^{-}\pi^{+})$ and $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{+}\pi^{-})/B(D^{0} \rightarrow K^{-}\pi^{+})$. The nonresonant $\pi^{+}\pi^{0}$ contribution under the ρ^{+} is negligible.

54 Assumes equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^{*}(2010)$.

$\Gamma(D^{*}(2007)^{0}\pi^{+}\pi^{+}\pi^{-})/\Gamma_{\text{total}}$				Γ_{17}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.0094 \pm 0.0020 \pm 0.0017	48	55,56	ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

55 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the CLEO II $B(D^{*}(2007)^{0} \rightarrow D^{0}\pi^{0})$ and absolute $B(D^{0} \rightarrow K^{-}\pi^{+})$ and the PDG 1992 $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{0})/B(D^{0} \rightarrow K^{-}\pi^{+})$ and $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{+}\pi^{-})/B(D^{0} \rightarrow K^{-}\pi^{+})$.

56 The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^{+} , the branching ratio for $\bar{D}^{*0}a_1^{+}$ is twice that for $\bar{D}^{*0}\pi^{+}\pi^{+}\pi^{-}$.)

$\Gamma(D^{*}(2010)^{-}\pi^{+}\pi^{+}\pi^{0})/\Gamma_{\text{total}}$				Γ_{18}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.0151 \pm 0.0070 \pm 0.0003	26		57 ALBRECHT 90J	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.043 \pm 0.013 \pm 0.026 24 58 ALBRECHT 87C ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

57 ALBRECHT 90J reports 0.018 \pm 0.007 \pm 0.005 for $B(D^{*}(2010)^{+} \rightarrow D^{0}\pi^{+}) = 0.57 \pm 0.06$. We rescale to our best value $B(D^{*}(2010)^{+} \rightarrow D^{0}\pi^{+}) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

58 ALBRECHT 87C use PDG 86 branching ratios for D and $D^{*}(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^{+}B^{-}) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^{0}\bar{B}^{0}) = 45\%$. Superseded by ALBRECHT 90J.

$\Gamma(D^{*}(2010)^{-}\pi^{+}\pi^{+}\pi^{+}\pi^{-})/\Gamma_{\text{total}}$				Γ_{19}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.01	90		59 ALBRECHT 90J	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

59 Assumes equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^{*}(2010)$.

$\Gamma(D_{s1}^{*}(2420)^{0}\pi^{+})/\Gamma_{\text{total}}$				Γ_{20}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.0011 \pm 0.0005 \pm 0.0002	8		60 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

60 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the CLEO II $B(D^{*}(2010)^{+} \rightarrow D^{0}\pi^{+})$ and absolute $B(D^{0} \rightarrow K^{-}\pi^{+})$ and the PDG 1992 $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{0})/B(D^{0} \rightarrow K^{-}\pi^{+})$ and $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{+}\pi^{-})/B(D^{0} \rightarrow K^{-}\pi^{+})$, assuming $B(D_{s1}^{*}(2420)^{0} \rightarrow D^{*}(2010)^{+}\pi^{-}) = 67\%$.

$\Gamma(D_{s1}^{*}(2420)^{0}\rho^{+})/\Gamma_{\text{total}}$				Γ_{21}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.0014	90		61 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

61 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the CLEO II $B(D^{*}(2010)^{+} \rightarrow D^{0}\pi^{+})$ and absolute $B(D^{0} \rightarrow K^{-}\pi^{+})$ and the PDG 1992 $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{0})/B(D^{0} \rightarrow K^{-}\pi^{+})$ and $B(D^{0} \rightarrow K^{-}\pi^{+}\pi^{+}\pi^{-})/B(D^{0} \rightarrow K^{-}\pi^{+})$, assuming $B(D_{s1}^{*}(2420)^{0} \rightarrow D^{*}(2010)^{+}\pi^{-}) = 67\%$.

$\Gamma(D_{s2}^{*}(2460)^{0}\pi^{+})/\Gamma_{\text{total}}$				Γ_{22}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.0013	90		62 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0028 90 63 ALAM 94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

62 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the Mark III $B(D^{+} \rightarrow K^{-}\pi^{+}\pi^{+})$ and $B(D_{s2}^{*}(2460)^{0} \rightarrow D^{+}\pi^{-}) = 30\%$.

63 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the Mark III $B(D^{+} \rightarrow K^{-}\pi^{+}\pi^{+})$, the CLEO II $B(D^{*}(2010)^{+} \rightarrow D^{0}\pi^{+})$ and $B(D_{s2}^{*}(2460)^{0} \rightarrow D^{*}(2010)^{+}\pi^{-}) = 20\%$.

$\Gamma(D_{s2}^{*}(2460)^{0}\rho^{+})/\Gamma_{\text{total}}$				Γ_{23}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.0047	90		64 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$
<0.005	90		65 ALAM	94 CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

64 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the Mark III $B(D^{+} \rightarrow K^{-}\pi^{+}\pi^{+})$ and $B(D_{s2}^{*}(2460)^{0} \rightarrow D^{+}\pi^{-}) = 30\%$.

65 ALAM 94 assume equal production of B^{+} and B^{0} at the $\Upsilon(4S)$ and use the Mark III $B(D^{+} \rightarrow K^{-}\pi^{+}\pi^{+})$, the CLEO II $B(D^{*}(2010)^{+} \rightarrow D^{0}\pi^{+})$ and $B(D_{s2}^{*}(2460)^{0} \rightarrow D^{*}(2010)^{+}\pi^{-}) = 20\%$.

$\Gamma(D^{0}D_{s1}^{+})/\Gamma_{\text{total}}$				Γ_{24}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.017 \pm 0.006 OUR AVERAGE				
0.019 \pm 0.010 \pm 0.002			66 ALBRECHT 92G	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$
0.017 \pm 0.007 \pm 0.002	5		67 BORTOLETTO90	CLEO $e^{+}e^{-} \rightarrow \Upsilon(4S)$

66 ALBRECHT 92G reports 0.024 \pm 0.012 \pm 0.004 for $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.027$. We rescale to our best value $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^{-}\pi^{+}) = 3.71 \pm 0.25\%$.

67 BORTOLETTO 90 reports 0.029 \pm 0.013 for $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.02$. We rescale to our best value $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^{0}D_{s2}^{+})/\Gamma_{\text{total}}$				Γ_{25}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.012 \pm 0.010 \pm 0.001			68 ALBRECHT 92G	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

68 ALBRECHT 92G reports 0.016 \pm 0.012 \pm 0.003 for $B(D_{s2}^{+} \rightarrow \phi\pi^{+}) = 0.027$. We rescale to our best value $B(D_{s2}^{+} \rightarrow \phi\pi^{+}) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 branching ratios, e.g., $B(D^0 \rightarrow K^{-}\pi^{+}) = 3.71 \pm 0.25\%$.

$\Gamma(D^{*}(2007)^{0}D_{s1}^{+})/\Gamma_{\text{total}}$				Γ_{26}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.010 \pm 0.007 \pm 0.001			69 ALBRECHT 92G	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

69 ALBRECHT 92G reports 0.013 \pm 0.009 \pm 0.002 for $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.027$. We rescale to our best value $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^{*}(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^{-}\pi^{+}) = 3.71 \pm 0.25\%$ and $B(D^{*}(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$.

$\Gamma(D^{*}(2007)^{0}D_{s2}^{+})/\Gamma_{\text{total}}$				Γ_{27}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
0.024 \pm 0.013 \pm 0.003			70 ALBRECHT 92G	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

70 ALBRECHT 92G reports 0.031 \pm 0.016 \pm 0.005 for $B(D_{s2}^{+} \rightarrow \phi\pi^{+}) = 0.027$. We rescale to our best value $B(D_{s2}^{+} \rightarrow \phi\pi^{+}) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^0 and $D^{*}(2007)^0$ branching ratios, e.g., $B(D^0 \rightarrow K^{-}\pi^{+}) = 3.71 \pm 0.25\%$ and $B(D^{*}(2007)^0 \rightarrow D^0\pi^0) = 55 \pm 6\%$.

$\Gamma(D_{s1}^{+}\pi^0)/\Gamma_{\text{total}}$				Γ_{28}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.00021	90		71 ALEXANDER 93B	CLE2 $e^{+}e^{-} \rightarrow \Upsilon(4S)$

71 ALEXANDER 93B reports < 2.0 \times 10⁻⁴ for $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.037$. We rescale to our best value $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.035$.

$[\Gamma(D_{s1}^{+}\pi^0) + \Gamma(D_{s2}^{+}\pi^0)]/\Gamma_{\text{total}}$				$(\Gamma_{28} + \Gamma_{29})/\Gamma$
VALUE	CL%	EVTS	DOCUMENT ID	TECN COMMENT
<0.0007	90		72 ALBRECHT 93E	ARG $e^{+}e^{-} \rightarrow \Upsilon(4S)$

72 ALBRECHT 93E reports < 0.9 \times 10⁻³ for $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.027$. We rescale to our best value $B(D_{s1}^{+} \rightarrow \phi\pi^{+}) = 0.035$.

$\Gamma(D_s^{*-} \pi^+ K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.009	90	99 ALBRECHT 93E ARG		$e^+e^- \rightarrow \Upsilon(4S)$

99 ALBRECHT 93E reports $< 1.1 \times 10^{-2}$ for $B(D_s^{*-} \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^{*-} \rightarrow \phi \pi^+) = 0.035$.

 $\Gamma(J/\psi(1S)K^+)/\Gamma_{\text{total}}$ Γ_{48}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.2 ± 1.4 OUR AVERAGE					
11.0 ± 1.5 ± 0.9	59	100	ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
9.22 ± 3.03 ± 0.39	101		BORTOLETTO92	92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
8.1 ± 3.5 ± 0.3	6	102	ALBRECHT 90J ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

22 ± 10 ± 2			BUSKULIC	92G	ALEP $e^+e^- \rightarrow Z$
7 ± 4	3	103	ALBRECHT 87D ARG		$e^+e^- \rightarrow \Upsilon(4S)$
10 ± 7 ± 2	3	104	BEBEK	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
9 ± 5	3	105	ALAM	86	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

100 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 101 BORTOLETTO 92 reports $8 \pm 2 \pm 2$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 102 ALBRECHT 90J reports $7 \pm 3 \pm 1$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

103 ALBRECHT 87D assume B^+/B^0 ratio is 55/45. Superseded by ALBRECHT 90J.
 104 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.
 105 ALAM 86 assumes B^\pm/B^0 ratio is 60/40.

 $\Gamma(J/\psi(1S)K^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{49}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0014 ± 0.0006 OUR AVERAGE					
0.0014 ± 0.0008 ± 0.0001			106 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.0014 ± 0.0009 ± 0.0001	6	107	ALBRECHT 87D ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0018	90	108	ALBRECHT 90J ARG		$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	-----	------------------	--	-----------------------------------

106 BORTOLETTO 92 reports $0.0012 \pm 0.0006 \pm 0.0004$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 107 ALBRECHT 87D reports 0.0012 ± 0.0008 for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They actually report 0.0011 ± 0.0007 assuming B^+/B^0 ratio is 55/45. We rescale to 50/50. Analysis explicitly removes $B^+ \rightarrow \psi(2S)K^+$.

108 ALBRECHT 90J reports < 0.0016 for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = 0.0599$. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(J/\psi(1S)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{50}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0017 ± 0.0005 OUR AVERAGE					
0.00178 ± 0.00051 ± 0.00023	13	109	ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
0.0015 ± 0.0011 ± 0.0001	110		BORTOLETTO92	92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
0.0018 ± 0.0013 ± 0.0001	2	111	ALBRECHT 90J ARG		$e^+e^- \rightarrow \Upsilon(4S)$

109 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$. The neutral and charged B events together are predominantly longitudinally polarized, $\Gamma_L/\Gamma = 0.80 \pm 0.08 \pm 0.05$. This can be compared with a prediction, using HQET, of 0.73 (KRAMER 92). This polarization indicates that the $B \rightarrow \psi K^*$ decay is dominated by the $CP = -1$ CP eigenstate.

110 BORTOLETTO 92 reports $0.0013 \pm 0.0009 \pm 0.0003$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

111 ALBRECHT 90J reports $0.0016 \pm 0.0011 \pm 0.0003$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\psi(2S)K^+)/\Gamma_{\text{total}}$ Γ_{51}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
6.9 ± 3.1 OUR AVERAGE					Error includes scale factor of 1.3.
6.1 ± 2.3 ± 0.9	7	112	ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
18 ± 8 ± 4	5	112	ALBRECHT 90J ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5	90	112	BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
22 ± 17	3	113	ALBRECHT 87D ARG		$e^+e^- \rightarrow \Upsilon(4S)$

112 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 113 ALBRECHT 87D assume B^+/B^0 ratio is 55/45. Superseded by ALBRECHT 90J.

 $\Gamma(\psi(2S)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{52}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0030	90	114 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0035	90	114	BORTOLETTO92	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
<0.0049	90	114	ALBRECHT 90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

114 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\psi(2S)K^*(892)^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{53}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0019 ± 0.0011 ± 0.0004		3	115	ALBRECHT 90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

115 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\chi_{c1}(1P)K^+)/\Gamma_{\text{total}}$ Γ_{54}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0010 ± 0.0004 OUR AVERAGE					
0.00097 ± 0.00040 ± 0.00009	6	116	ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
0.0019 ± 0.0013 ± 0.0006	117		ALBRECHT 92E ARG		$e^+e^- \rightarrow \Upsilon(4S)$

116 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.
 117 ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production and $B(\Upsilon(4S) \rightarrow B^+B^-) = 50\%$.

 $\Gamma(\chi_{c1}(1P)K^*(892)^+)/\Gamma_{\text{total}}$ Γ_{55}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0021	90	118	ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

118 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(K^0\pi^+)/\Gamma_{\text{total}}$ Γ_{56}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.0 × 10 ⁻⁴	90	119	AVERY	89B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.9 × 10 ⁻⁴	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
<6.8 × 10 ⁻⁴	90		AVERY	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

119 AVERY 89B reports $< 9 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^*(892)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{57}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.5 × 10 ⁻⁴	90	120	AVERY	89B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.7 × 10 ⁻⁴	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
<2.6 × 10 ⁻⁴	90		AVERY	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

120 AVERY 89B reports $< 1.3 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^+\pi^-\pi^+(\text{no charm}))/\Gamma_{\text{total}}$ Γ_{58}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.9 × 10 ⁻⁴	90	121	AVERY	89B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<3.3 × 10 ⁻⁴	90		ALBRECHT 91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-------------------------	----	--	------------------	-----------------------------------

121 AVERY 89B reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_1(1400)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{59}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.6 × 10 ⁻³	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K_2^*(1430)^0\pi^+)/\Gamma_{\text{total}}$ Γ_{60}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<6.8 × 10 ⁻⁴	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K^+\rho^0)/\Gamma_{\text{total}}$ Γ_{61}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<8 × 10 ⁻⁵	90	122	AVERY	89B	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.8 × 10 ⁻⁴	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
<2.6 × 10 ⁻⁴	90		AVERY	87	CLEO $e^+e^- \rightarrow \Upsilon(4S)$

122 AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^*(892)^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{62}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.1 × 10 ⁻³	90		ALBRECHT 91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K^*(892)^+\rho^0)/\Gamma_{\text{total}}$ Γ_{63}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<9.0 × 10 ⁻⁴	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(K_1(1400)^+\rho^0)/\Gamma_{\text{total}}$ Γ_{64}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<7.8 × 10 ⁻⁴	90		ALBRECHT 91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

Meson Full Listings

 B^\pm

$\Gamma(K_2^*(1430)^+\rho^0)/\Gamma_{total}$					Γ_{65}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.5 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^+K^-K^+)/\Gamma_{total}$					Γ_{66}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.5 \times 10^{-4}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^+\phi)/\Gamma_{total}$					Γ_{67}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<9 \times 10^{-5}$	90	123 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$<1.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
$<2.1 \times 10^{-4}$	90	AVERY	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
123 AVERY 89B reports $< 8 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.					

$\Gamma(K^*(892)^+K^+K^-)/\Gamma_{total}$					Γ_{68}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.6 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^*(892)^+\phi)/\Gamma_{total}$					Γ_{69}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.3 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K_1(1400)^+\phi)/\Gamma_{total}$					Γ_{70}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K_2^*(1430)^+\phi)/\Gamma_{total}$					Γ_{71}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<3.4 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(K^+f_0(980))/\Gamma_{total}$					Γ_{72}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8 \times 10^{-5}$	90	124 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
124 AVERY 89B reports $< 7 \times 10^{-5}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.					

$\Gamma(K^*(892)^+\gamma)/\Gamma_{total}$					Γ_{73}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$(5.7 \pm 3.1 \pm 1.1) \times 10^{-5}$		5	125 AMMAR	93 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 5.5	$\times 10^{-4}$	90	126 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 5.5	$\times 10^{-4}$	90	127 AVERY	89B CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 1.8	$\times 10^{-3}$	90	AVERY	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

125 AMMAR 93 observed 4.1 ± 2.3 events above background.126 Assumes the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$.127 Assumes the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(K_1(1270)^+\gamma)/\Gamma_{total}$					Γ_{74}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0073	90	128 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

128 ALBRECHT 89G reports < 0.0066 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_1(1400)^+\gamma)/\Gamma_{total}$					Γ_{75}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0022	90	129 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

129 ALBRECHT 89G reports < 0.0020 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_2^*(1430)^+\gamma)/\Gamma_{total}$					Γ_{76}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0014	90	130 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

130 ALBRECHT 89G reports < 0.0013 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(1680)^+\gamma)/\Gamma_{total}$					Γ_{77}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0019	90	131 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

131 ALBRECHT 89G reports < 0.0017 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^+\gamma)/\Gamma_{total}$					Γ_{78}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0055	90	132 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

132 ALBRECHT 89G reports < 0.005 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^+\gamma)/\Gamma_{total}$					Γ_{79}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0099	90	133 ALBRECHT	89G ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

133 ALBRECHT 89G reports < 0.0090 assuming the $\Upsilon(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^0)/\Gamma_{total}$					Γ_{80}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.4 \times 10^{-4}$	90	134 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $< 2.3 \times 10^{-3}$ 90 135 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$ 134 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.135 BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$.

$\Gamma(\pi^+\pi^+\pi^-)/\Gamma_{total}$					Γ_{81}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.9 \times 10^{-4}$	90	136 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $< 4.5 \times 10^{-4}$ 90 137 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$ 136 BORTOLETTO 89 reports $< 1.7 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.137 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^0\pi^+)/\Gamma_{total}$					Γ_{82}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-4}$	90		138 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $< 1.7 \times 10^{-4}$ 90 139 BORTOLETTO89 CLEO $e^+e^- \rightarrow \Upsilon(4S)$ $< 2.3 \times 10^{-4}$ 90 139 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$ $< 6 \times 10^{-4}$ 90 0 GILES 84 CLEO Repl. by BEBEK 87138 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.139 Papers assume the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+f_0(980))/\Gamma_{total}$					Γ_{83}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.4 \times 10^{-4}$	90	140 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

140 BORTOLETTO 89 reports $< 1.2 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+f_2(1270))/\Gamma_{total}$					Γ_{84}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<2.4 \times 10^{-4}$	90	141 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

141 BORTOLETTO 89 reports $< 2.1 \times 10^{-4}$ assuming the $\Upsilon(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^0\pi^0)/\Gamma_{total}$					Γ_{85}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<8.9 \times 10^{-4}$	90	142 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

142 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\pi^0)/\Gamma_{total}$					Γ_{86}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<5.5 \times 10^{-4}$	90	143 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

143 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\pi^+\pi^-\pi^+\pi^0)/\Gamma_{total}$					Γ_{87}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.0 \times 10^{-3}$	90	144 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

144 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\rho^+\rho^0)/\Gamma_{total}$					Γ_{88}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.0 \times 10^{-3}$	90	145 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

145 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+\pi^0)/\Gamma_{total}$					Γ_{89}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<1.7 \times 10^{-3}$	90	146 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

146 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^0\pi^+)/\Gamma_{total}$					Γ_{90}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<9.0 \times 10^{-4}$	90	147 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

147 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

$\Gamma(\omega\pi^+)/\Gamma_{total}$					Γ_{91}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$<4.0 \times 10^{-4}$	90	148 ALBRECHT	90B ARG	$e^+e^- \rightarrow \Upsilon(4S)$	

148 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $\Upsilon(4S)$.

See key on page 1343

Meson Full Listings

 B^\pm

$\Gamma(\eta\pi^+)/\Gamma_{\text{total}}$ Γ_{92}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.0 \times 10^{-4}$	90	149 ALBRECHT 90B ARG		$e^+e^- \rightarrow T(4S)$

149 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

$\Gamma(\pi^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}}$ Γ_{93}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.6 \times 10^{-4}$	90	150 ALBRECHT 90B ARG		$e^+e^- \rightarrow T(4S)$

150 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

$\Gamma(\rho^0 a_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{94}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.2 \times 10^{-4}$	90	151 BORTOLETTO89	CLEO	$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<6.0 \times 10^{-4}$	90	152 ALBRECHT 90B ARG		$e^+e^- \rightarrow T(4S)$
$<3.2 \times 10^{-3}$	90	151 BEBEK 87 CLEO		$e^+e^- \rightarrow T(4S)$

151 BORTOLETTO 89 reports $< 5.4 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

152 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

$\Gamma(\rho^0 a_2(1320)^+)/\Gamma_{\text{total}}$ Γ_{95}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<7.2 \times 10^{-4}$	90	153 BORTOLETTO89	CLEO	$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.6 \times 10^{-3}$	90	154 BEBEK 87 CLEO		$e^+e^- \rightarrow T(4S)$
-----------------------	----	-------------------	--	----------------------------

153 BORTOLETTO 89 reports $< 6.3 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

154 BEBEK 87 reports $< 2.3 \times 10^{-3}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+\pi^+\pi^-\pi^0)/\Gamma_{\text{total}}$ Γ_{96}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.3 \times 10^{-3}$	90	155 ALBRECHT 90B ARG		$e^+e^- \rightarrow T(4S)$

155 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

$\Gamma(a_1(1260)^+ a_1(1260)^0)/\Gamma_{\text{total}}$ Γ_{97}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-2}$	90	156 ALBRECHT 90B ARG		$e^+e^- \rightarrow T(4S)$

156 ALBRECHT 90B limit assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

$\Gamma(p\bar{p}\pi^+)/\Gamma_{\text{total}}$ Γ_{98}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 1.6 \times 10^{-4}$	90	157 BEBEK 89 CLEO		$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(5.7 \pm 1.5 \pm 2.1) \times 10^{-4}$ 158 ALBRECHT 88F ARG $e^+e^- \rightarrow T(4S)$

157 BEBEK 89 reports $< 1.4 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

158 ALBRECHT 88F reports $(5.2 \pm 1.4 \pm 1.9) \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{p}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{99}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.2 \times 10^{-4}$	90	159 ALBRECHT 88F ARG		$e^+e^- \rightarrow T(4S)$

159 ALBRECHT 88F reports $< 4.7 \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{A})/\Gamma_{\text{total}}$ Γ_{100}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-5}$	90	160 AVERY 89B CLEO		$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.3 \times 10^{-5}$ 90 161 ALBRECHT 88F ARG $e^+e^- \rightarrow T(4S)$

160 AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

161 ALBRECHT 88F reports $< 8.5 \times 10^{-5}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{A}\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{101}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.0 \times 10^{-4}$	90	162 ALBRECHT 88F ARG		$e^+e^- \rightarrow T(4S)$

162 ALBRECHT 88F reports $< 1.8 \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\Delta^0 p)/\Gamma_{\text{total}}$ Γ_{102}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.8 \times 10^{-4}$	90	163 BORTOLETTO89	CLEO	$e^+e^- \rightarrow T(4S)$

163 BORTOLETTO 89 reports $< 3.3 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\Delta^+ \bar{p})/\Gamma_{\text{total}}$ Γ_{103}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.5 \times 10^{-4}$	90	164 BORTOLETTO89	CLEO	$e^+e^- \rightarrow T(4S)$

164 BORTOLETTO 89 reports $< 1.3 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{104}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0039	90	165 WEIR 90B MRK2		$e^+e^- 29 \text{ GeV}$

165 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{105}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0091	90	166 WEIR 90B MRK2		$e^+e^- 29 \text{ GeV}$

166 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6 \times 10^{-5}$	90	167 AVERY 89B CLEO		$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<9.9 \times 10^{-5}$ 90 168 ALBRECHT 91E ARG $e^+e^- \rightarrow T(4S)$

$<6.8 \times 10^{-3}$ 90 169 WEIR 90B MRK2 $e^+e^- 29 \text{ GeV}$

$<2.5 \times 10^{-4}$ 90 170 AVERY 87 CLEO $e^+e^- \rightarrow T(4S)$

167 AVERY 89B reports $< 5 \times 10^{-5}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

168 ALBRECHT 91E reports $< 9.0 \times 10^{-5}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

169 WEIR 90B assumes B^+ production cross section from LUND.

170 AVERY 87 reports $< 2.1 \times 10^{-4}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{107}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	171 AVERY 89B CLEO		$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$<2.4 \times 10^{-4}$ 90 172 ALBRECHT 91E ARG $e^+e^- \rightarrow T(4S)$

$<6.4 \times 10^{-3}$ 90 173 WEIR 90B MRK2 $e^+e^- 29 \text{ GeV}$

$<3.8 \times 10^{-4}$ 90 174 AVERY 87 CLEO $e^+e^- \rightarrow T(4S)$

171 AVERY 89B reports $< 1.5 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

172 ALBRECHT 91E reports $< 2.2 \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

173 WEIR 90B assumes B^+ production cross section from LUND.

174 AVERY 87 reports $< 3.2 \times 10^{-4}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+ e^+ e^-)/\Gamma_{\text{total}}$ Γ_{108}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.9 \times 10^{-4}$	90	175 ALBRECHT 91E ARG		$e^+e^- \rightarrow T(4S)$

175 ALBRECHT 91E reports $< 6.3 \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^*(892)^+ \mu^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{109}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.2 \times 10^{-3}$	90	176 ALBRECHT 91E ARG		$e^+e^- \rightarrow T(4S)$

176 ALBRECHT 91E reports $< 1.1 \times 10^{-3}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	177 WEIR 90B MRK2		$e^+e^- 29 \text{ GeV}$

177 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{111}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	178 WEIR 90B MRK2		$e^+e^- 29 \text{ GeV}$

178 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^+ \mu^-)/\Gamma_{\text{total}}$ Γ_{112}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	179 WEIR 90B MRK2		$e^+e^- 29 \text{ GeV}$

179 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^+ e^- \mu^+)/\Gamma_{\text{total}}$ Γ_{113}/Γ

Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	180 WEIR 90B MRK2		$e^+e^- 29 \text{ GeV}$

180 WEIR 90B assumes B^+ production cross section from LUND.

Meson Full Listings

B^\pm

$\Gamma(\pi^- e^+ e^+)/\Gamma_{total}$ Γ_{114}/Γ
 Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0039	90	181 WEIR	90B MRK2	e^+e^- 29 GeV

181 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- \mu^+ \mu^+)/\Gamma_{total}$ Γ_{115}/Γ
 Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0091	90	182 WEIR	90B MRK2	e^+e^- 29 GeV

182 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(\pi^- e^+ \mu^+)/\Gamma_{total}$ Γ_{116}/Γ
 Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	183 WEIR	90B MRK2	e^+e^- 29 GeV

183 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- e^+ e^+)/\Gamma_{total}$ Γ_{117}/Γ
 Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0039	90	184 WEIR	90B MRK2	e^+e^- 29 GeV

184 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- \mu^+ \mu^+)/\Gamma_{total}$ Γ_{118}/Γ
 Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0091	90	185 WEIR	90B MRK2	e^+e^- 29 GeV

185 WEIR 90B assumes B^+ production cross section from LUND.

$\Gamma(K^- e^+ \mu^+)/\Gamma_{total}$ Γ_{119}/Γ
 Test of total lepton number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0064	90	186 WEIR	90B MRK2	e^+e^- 29 GeV

186 WEIR 90B assumes B^+ production cross section from LUND.

For all of the decays below, the charge of the decaying B was not determined. Only the admixture of B^0 , \bar{B}^0 , B^+ , and B^- at the $\Upsilon(4S)$ is used in the averages, except where no $\Upsilon(4S)$ data are available.

$\Gamma(e^+ \nu_e \text{ anything})/\Gamma_{total}$ Γ_{124}/Γ
 Only the experiments at the $\Upsilon(4S)$ are used in the average.

These branching fraction values are model dependent. See the note on "Semileptonic Decays of D and B Mesons, Part II" at the beginning of the B^+ Full Listings.

VALUE	DOCUMENT ID	TECN	COMMENT
0.1043 ± 0.0024 OUR AVERAGE	Includes data from the 2 datablocks that follow this one.		
0.108 ± 0.002 ± 0.0056	187 HENDERSON 92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

187 HENDERSON 92 measurement employs e and μ . The systematic error contains 0.004 in quadrature from model dependence. The authors average a variation of the Isgur, Scora, Grinstein, and Wise model with that of the Altarelli-Cabibbo-Corbò-Maiani-Martinelli model for semileptonic decays to correct the acceptance.

$\Gamma(e^+ \nu_e \text{ anything})/\Gamma_{total}$ Γ_{120}/Γ
 Only the experiments at the $\Upsilon(4S)$ are used in the average.


These branching fraction values are model dependent. See the note on "Semileptonic Decays of D and B Mesons, Part II" at the beginning of the B^+ Full Listings.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.104 ± 0.004 OUR AVERAGE	Error includes scale factor of 1.3. See the ideogram below.			
0.097 ± 0.005 ± 0.004	188	ALBRECHT 93H ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
0.100 ± 0.004 ± 0.003	189	YANAGISAWA 91 CSB2	$e^+e^- \rightarrow \Upsilon(4S)$	
0.103 ± 0.006 ± 0.002	190	ALBRECHT 90H ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
0.117 ± 0.004 ± 0.010	191	WACHS 89 CBAL	Direct e at $\Upsilon(4S)$	
0.120 ± 0.007 ± 0.005		CHEN 84 CLEO	Direct e at $\Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.086 ± 0.027 ± 0.008	192	ABE 93E VNS	$E_{cm}^{ee} = 58$ GeV	
0.107 ± 0.015 ± 0.007	260	193 ABREU 93C DLPH	$e^+e^- \rightarrow Z$	
0.109 ± 0.014 ± 0.0055	2719	194 AKERS 93B OPAL	$e^+e^- \rightarrow Z$	
0.138 ± 0.032 ± 0.008	195	ADEVA 91C L3	Z decays	
0.111 ± 0.028 ± 0.026		BEHREND 90D CELL	$E_{cm}^{ee} = 43$ GeV	
0.150 ± 0.011 ± 0.022		BEHREND 90D CELL	$E_{cm}^{ee} = 35$ GeV	
0.112 ± 0.009 ± 0.011		ONG 88 MRK2	$E_{cm}^{ee} = 29$ GeV	
0.149 ± 0.022 ± 0.019		PAL 86 DLCO	$E_{cm}^{ee} = 29$ GeV	
0.110 ± 0.018 ± 0.010		AIHARA 85 TPC	$E_{cm}^{ee} = 29$ GeV	
0.111 ± 0.034 ± 0.040		ALTHOFF 84J TASS	$E_{cm}^{ee} = 34.6$ GeV	
0.146 ± 0.028		KOOP 84 DLCO	Repl. by PAL 86	
0.132 ± 0.008 ± 0.014	196	KLOPFEN... 83B CUSB	Direct e at $\Upsilon(4S)$	
0.116 ± 0.021 ± 0.017		NELSON 83 MRK2	$E_{cm}^{ee} = 29$ GeV	

- 188 ALBRECHT 93H analysis performed using tagged semileptonic decays of the B . This technique is almost model independent for the lepton branching ratio.
- 189 YANAGISAWA 91 also measures an average semileptonic branching ratio at the $\Upsilon(5S)$ of 9.6–10.5% depending on assumptions about the relative production of different B meson species.
- 190 ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.099 ± 0.006 is obtained using ISGUR 89B.
- 191 Using data above $p(e) = 2.4$ GeV, WACHS 89 determine $\sigma(B \rightarrow e\nu p)/\sigma(B \rightarrow e\nu \text{charm}) < 0.065$ at 90% CL.
- 192 ABE 93E experiment also measures forward-backward asymmetries and fragmentation functions for b and c .
- 193 ABREU 93C event count includes ee events. Combining ee , $\mu\mu$, and $e\mu$ events, they obtain 0.100 ± 0.007 ± 0.007.
- 194 AKERS 93B analysis performed using single and dilepton events.
- 195 ADEVA 91C measure the average $B(b \rightarrow eX)$ branching ratio using single and double tagged b enhanced Z events. Combining e and μ results, they obtain 0.113 ± 0.010 ± 0.006. Constraining the initial number of b quarks by the Standard Model prediction (378 ± 3 MeV) for the decay of the Z into $b\bar{b}$, the electron result gives 0.112 ± 0.004 ± 0.008. They obtain 0.119 ± 0.003 ± 0.006 when e and μ results are combined. Used to measure the $b\bar{b}$ width itself, this electron result gives 370 ± 12 ± 24 MeV and combined with the muon result gives 385 ± 7 ± 22 MeV.
- 196 Ratio $\sigma(b \rightarrow e\nu p)/\sigma(b \rightarrow e\nu \text{charm}) < 0.055$ at CL = 90%.

WEIGHTED AVERAGE
 0.104 ± 0.004 (Error scaled by 1.3)



$\Gamma(\mu^+ \nu_\mu \text{ anything})/\Gamma_{total}$ Γ_{123}/Γ
 Only the experiments at the $\Upsilon(4S)$ are used in the average.

These branching fraction values are model dependent. See the note on "Semileptonic Decays of D and B Mesons, Part II" at the beginning of the B^+ Full Listings.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
The data in this block is included in the average printed for a previous datablock.				

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.103 ± 0.005 OUR AVERAGE				
0.100 ± 0.006 ± 0.002	197	ALBRECHT 90H ARG	$e^+e^- \rightarrow \Upsilon(4S)$	
0.108 ± 0.006 ± 0.01		CHEN 84 CLEO	Direct μ at $\Upsilon(4S)$	
0.112 ± 0.009 ± 0.01		LEVMAN 84 CUSB	Direct μ at $\Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.110 ± 0.012 ± 0.007	656	198 ABREU 93C DLPH	$e^+e^- \rightarrow Z$	
0.101 ± 0.010 ± 0.009 ± 0.0055	4248	199 AKERS 93B OPAL	$e^+e^- \rightarrow Z$	
0.113 ± 0.012 ± 0.006	200	ADEVA 91C L3	Z decays	
0.104 ± 0.023 ± 0.016		BEHREND 90D CELL	$E_{cm}^{ee} = 43$ GeV	
0.148 ± 0.010 ± 0.016		BEHREND 90D CELL	$E_{cm}^{ee} = 35$ GeV	
0.118 ± 0.012 ± 0.010		ONG 88 MRK2	$E_{cm}^{ee} = 29$ GeV	
0.117 ± 0.016 ± 0.015		BARTEL 87 JADE	$E_{cm}^{ee} = 34.6$ GeV	
0.114 ± 0.018 ± 0.025		BARTEL 85J JADE	Repl. by BARTEL 87	
0.117 ± 0.028 ± 0.010		ALTHOFF 84G TASS	$E_{cm}^{ee} = 34.5$ GeV	
0.105 ± 0.015 ± 0.013		ADEVA 83B MRKJ	$E_{cm}^{ee} = 33-38.5$ GeV	
0.155 ± 0.054 ± 0.029		FERNANDEZ 83D MAC	$E_{cm}^{ee} = 29$ GeV	
197 ALBRECHT 90H uses the model of ALTARELLI 82 to correct over all lepton momenta. 0.097 ± 0.006 is obtained using ISGUR 89B.				
198 ABREU 93C event count includes $\mu\mu$ events. Combining ee , $\mu\mu$, and $e\mu$ events, they obtain 0.100 ± 0.007 ± 0.007.				
199 AKERS 93B analysis performed using single and dilepton events.				
200 ADEVA 91C measure the average $B(b \rightarrow eX)$ branching ratio using single and double tagged b enhanced Z events. Combining e and μ results, they obtain 0.113 ± 0.010 ± 0.006. Constraining the initial number of b quarks by the Standard Model prediction (378 ± 3 MeV) for the decay of the Z into $b\bar{b}$, the muon result gives 0.123 ± 0.003 ± 0.006. They obtain 0.119 ± 0.003 ± 0.006 when e and μ results are combined. Used to measure the $b\bar{b}$ width itself, this muon result gives 394 ± 9 ± 22 MeV and combined with the electron result gives 385 ± 7 ± 22 MeV.				

$\Gamma(\bar{D}^*(2010) e^+ \nu_e)/\Gamma_{total}$ Γ_{121}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.070 ± 0.018 ± 0.014	90	ANTREASYAN 90B CBAL	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(\bar{p}e^+\nu_e \text{ anything})/\Gamma_{\text{total}}$					Γ_{122}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0016	90	ALBRECHT	90H ARG	$e^+e^- \rightarrow \gamma(4S)$	

$\Gamma(D^-\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$					$\Gamma_{125}/\Gamma_{124}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$0.26 \pm 0.07 \pm 0.04$		201 FULTON	91 CLEO	$e^+e^- \rightarrow \gamma(4S)$	

²⁰¹FULTON 91 uses $B(D^+ \rightarrow K^-\pi^+\pi^+) = (9.1 \pm 1.3 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(D^0\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$					$\Gamma_{126}/\Gamma_{124}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$0.67 \pm 0.09 \pm 0.10$		202 FULTON	91 CLEO	$e^+e^- \rightarrow \gamma(4S)$	

²⁰²FULTON 91 uses $B(D^0 \rightarrow K^-\pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ as measured by MARK III.

$\Gamma(D^{**}\ell^+\nu_\ell)/\Gamma_{\text{total}}$					Γ_{127}/Γ
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
$0.027 \pm 0.005 \pm 0.005$	63	203 ALBRECHT	93 ARG	$e^+e^- \rightarrow \gamma(4S)$	

D^{**} stands for the sum of the $D(1^1P_1)$, $D(1^3P_0)$, $D(1^3P_1)$, $D(1^3P_2)$, $D(2^1S_0)$, and $D(2^1S_1)$ resonances. $\ell = e$ or μ , not sum over e and μ modes.
²⁰³ALBRECHT 93 assumes the GISW model to correct for unseen modes. Using the BHK model, the result becomes $0.023 \pm 0.006 \pm 0.004$. Assumes $B(D^{*+} \rightarrow D^0\pi^+) = 68.1\%$, $B(D^0 \rightarrow K^-\pi^+) = 3.65\%$, $B(D^0 \rightarrow K^-\pi^+\pi^+) = 7.5\%$. We have taken their average e and μ value.

$\Gamma(D_s^-\ell^+\nu_\ell \text{ anything})/\Gamma_{\text{total}}$					Γ_{128}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.009	90	204 ALBRECHT	93E ARG	$e^+e^- \rightarrow \gamma(4S)$	

²⁰⁴ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$.

$\Gamma(D_s^-\ell^+\nu_\ell K^+ \text{ anything})/\Gamma_{\text{total}}$					Γ_{129}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.006	90	205 ALBRECHT	93E ARG	$e^+e^- \rightarrow \gamma(4S)$	

²⁰⁵ALBRECHT 93E reports < 0.008 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$.

$\Gamma(D_s^-\ell^+\nu_\ell K^0 \text{ anything})/\Gamma_{\text{total}}$					Γ_{130}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.009	90	206 ALBRECHT	93E ARG	$e^+e^- \rightarrow \gamma(4S)$	

²⁰⁶ALBRECHT 93E reports < 0.012 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$.

$\Gamma(\ell^+\nu_\ell \text{ noncharged})/\Gamma(\ell^+\nu_\ell \text{ anything})$					$\Gamma_{131}/\Gamma_{124}$
VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
		107	207 BARTELT	93B CLE2	$e^+e^- \rightarrow \gamma(4S)$
		77	208 ALBRECHT	91C ARG	$e^+e^- \rightarrow \gamma(4S)$
		76	209 FULTON	90 CLEO	$e^+e^- \rightarrow \gamma(4S)$

ℓ denotes e or μ , not the sum. These experiments measure this ratio in very limited momentum intervals.

		41	210 ALBRECHT	90 ARG	$e^+e^- \rightarrow \gamma(4S)$
<0.04	90	211 BEHREND	87 CLEO	$e^+e^- \rightarrow \gamma(4S)$	
<0.04	90	CHEN	84 CLEO	Direct e at $\gamma(4S)$	
<0.055	90	KLOPFEN...	83B CUSB	Direct e at $\gamma(4S)$	

²⁰⁷BARTELT 93B (CLEO II) measures an excess of $107 \pm 15 \pm 11$ leptons in the lepton momentum interval 2.3–2.6 GeV/c which is attributed to $b \rightarrow u\ell\nu_\ell$. This corresponds to a model-dependent partial branching ratio $\Delta B_{\mu b}$ between $(1.15 \pm 0.16 \pm 0.15) \times 10^{-4}$, as evaluated using the KS model (KOERNER 88), and $(1.54 \pm 0.22 \pm 0.20) \times 10^{-4}$ using the ACCMM model (ARTUSO 93). The corresponding values of $|V_{ub}|/|V_{cb}|$ are 0.056 ± 0.006 and 0.076 ± 0.008 , respectively.

²⁰⁸ALBRECHT 91C result supersedes ALBRECHT 90. Two events are fully reconstructed providing evidence for the $b \rightarrow u$ transition. Using the model of ALTARELLI 82, they obtain $|V_{ub}|/|V_{cb}| = 0.11 \pm 0.012$ from 77 leptons in the 2.3–2.6 GeV momentum range.

²⁰⁹FULTON 90 observe 76 ± 20 excess e and μ (lepton) events in the momentum interval $p = 2.4$ –2.6 GeV signaling the presence of the $b \rightarrow u$ transition. The average branching ratio, $(1.8 \pm 0.4 \pm 0.3) \times 10^{-4}$, corresponds to a model-dependent measurement of approximately $|V_{ub}|/|V_{cb}| = 0.1$ using $B(b \rightarrow c\ell\nu) = 10.2 \pm 0.2 \pm 0.7\%$.

²¹⁰ALBRECHT 90 observes 41 ± 10 excess e and μ (lepton) events in the momentum interval $p = 2.3$ –2.6 GeV signaling the presence of the $b \rightarrow u$ transition. The events correspond to a model-dependent measurement of $|V_{ub}|/|V_{cb}| = 0.10 \pm 0.01$.

²¹¹The quoted possible limits range from 0.018 to 0.04 for the ratio, depending on which model or momentum range is chosen. We select the most conservative limit they have calculated. This corresponds to a limit on $|V_{ub}|/|V_{cb}| < 0.20$. While the endpoint technique employed is more robust than their previous results in CHEN 84, these results do not provide a numerical improvement in the limit.

$\Gamma(K^+\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$					$\Gamma_{132}/\Gamma_{124}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$0.54 \pm 0.07 \pm 0.06$		212 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$	

ℓ denotes e or μ , not the sum.
²¹²ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(K^-\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$					$\Gamma_{133}/\Gamma_{124}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$0.10 \pm 0.05 \pm 0.02$		213 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$	

ℓ denotes e or μ , not the sum.
²¹³ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(K^0/\bar{K}^0\ell^+\nu_\ell \text{ anything})/\Gamma(\ell^+\nu_\ell \text{ anything})$					$\Gamma_{134}/\Gamma_{124}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$0.39 \pm 0.06 \pm 0.04$		214 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$	

ℓ denotes e or μ , not the sum. Sum over K^0 and \bar{K}^0 states.
²¹⁴ALAM 87B measurement relies on lepton-kaon correlations.

$\Gamma(\tau^+\nu_\tau \text{ anything})/\Gamma_{\text{total}}$					Γ_{135}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$(4.08 \pm 0.76 \pm 0.62) \times 10^{-2}$		215 BUSKULIC	93B ALEP	$e^+e^- \rightarrow Z$	

²¹⁵ B^0 , B^\pm , B_s^0 , and B baryon states not separated.

$\Gamma(c/\bar{c})/\Gamma_{\text{total}}$					
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
$0.98 \pm 0.16 \pm 0.12$		216 ALAM	87B CLEO	$e^+e^- \rightarrow \gamma(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •
²¹⁶From the difference between K^- and K^+ widths. ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.

$\Gamma(D^-\text{ anything})/\Gamma_{\text{total}}$					Γ_{136}/Γ
VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.26 ± 0.04			OUR AVERAGE		
$0.25 \pm 0.04 \pm 0.03$		217	BORTOLETTO92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
$0.272 \pm 0.063 \pm 0.035$		218	ALBRECHT	91H ARG	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 $0.17 \pm 0.04 \pm 0.04$ 20k ²¹⁹BORTOLETTO87 CLEO $e^+e^- \rightarrow \gamma(4S)$

²¹⁷The first error is the combined statistical and systematic error and the second error is due to the uncertainty in the D meson branching ratio. BORTOLETTO 92 measures $B(B \rightarrow D^+ \text{ anything}) \times B(D^+ \rightarrow K^-\pi^+\pi^+) = 0.0226 \pm 0.0030 \pm 0.0018$ and has chosen to normalize by the Mark III branching fractions.

²¹⁸ALBRECHT 91H measures $B(B \rightarrow D^\pm \text{ anything}) \times B(D^\pm \rightarrow K^-\pi^+\pi^+) = 0.0209 \pm 0.0027 \pm 0.0040$. Uses the PDG 90 $B(D^+ \rightarrow K^-\pi^+\pi^+) = 0.077 \pm 0.010$.

²¹⁹BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratio for $K^-\pi^+\pi^+ = 0.116 \pm 0.014 \pm 0.007$. The product branching ratio for $B(B \rightarrow D^+ X) B(D^+ \rightarrow K^-\pi^+\pi^+)$ is $0.019 \pm 0.004 \pm 0.002$. Superseded by BORTOLETTO 92.

$\Gamma(D^0 \text{ anything})/\Gamma_{\text{total}}$					Γ_{137}/Γ
VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.54 ± 0.06			OUR AVERAGE		
$0.55 \pm 0.04 \pm 0.08$		220	BORTOLETTO92	CLEO	$e^+e^- \rightarrow \gamma(4S)$
$0.522 \pm 0.082 \pm 0.035$		221	ALBRECHT	91H ARG	$e^+e^- \rightarrow \gamma(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 $0.39 \pm 0.05 \pm 0.04$ 21k ²²²BORTOLETTO87 CLEO $e^+e^- \rightarrow \gamma(4S)$

$0.57 \pm 0.14 \pm 0.12$ ²²³GREEN 83 CLEO Repl. by BORTOLETTO 87

²²⁰The first error is the combined statistical and systematic error and the second error is due to the uncertainty in the D meson branching ratio. BORTOLETTO 92 measures $B(B \rightarrow D^0 \text{ anything}) \times B(D^0 \rightarrow K^-\pi^+) = 0.0233 \pm 0.0012 \pm 0.0014$ and has chosen to normalize by the Mark III branching fractions.

²²¹ALBRECHT 91H measures $B(B \rightarrow D^0/\bar{D}^0 \text{ anything}) \times B(D^0 \rightarrow K^-\pi^+) = 0.0194 \pm 0.0015 \pm 0.0025$. Uses the PDG 90 $B(D^0 \rightarrow K^-\pi^+) = 0.0371 \pm 0.0025$.

²²²BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratio for $K^-\pi^+ = 0.056 \pm 0.004 \pm 0.003$. The product branching ratio for $B(B \rightarrow D^0 X) B(D^0 \rightarrow K^-\pi^+)$ is $0.0210 \pm 0.0015 \pm 0.0021$. Superseded by BORTOLETTO 92.

²²³Corrected by us using assumptions $B(D^0 \rightarrow K^-\pi^+) = (0.042 \pm 0.006)$. The product branching ratio is $B(B^0 \rightarrow D^0 X) B(D^0 \rightarrow K^-\pi^+) = 0.024 \pm 0.006 \pm 0.004$.

Meson Full Listings

 B^\pm $\Gamma(D^*(2010)^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{138}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.23 ± 0.04 OUR AVERAGE				Error includes scale factor of 1.4.
0.202 ± 0.038 ± 0.004		224 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.29 ± 0.05 ± 0.01		225 ALBRECHT 91H ARG		$e^+e^- \rightarrow \Upsilon(4S)$
0.22 ± 0.04	+0.07 -0.04	5200 226 BORTOLETTO87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.27 ± 0.06	+0.08 -0.06	510 227 CSORNA 85	CLEO	Repl. by BORTOLETTO 87

224 BORTOLETTO 92 reports $0.25 \pm 0.03 \pm 0.04$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.55 \pm 0.04$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. They also use the Mark III $B(D^0 \rightarrow K^-\pi^+)$ branching fraction.

225 ALBRECHT 91H reports $0.348 \pm 0.060 \pm 0.035$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Uses the PDG 90 $B(D^0 \rightarrow K^-\pi^+) = 0.0371 \pm 0.0025$.

226 BORTOLETTO 87 uses old MARK III (BALTRUSAITIS 86E) branching ratios $B(D^0 \rightarrow K^-\pi^+) = 0.056 \pm 0.004 \pm 0.003$ and also assumes $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.60 \pm 0.08$. The product branching ratio for $B(B \rightarrow D^*(2010)^+ B(D^*(2010)^+ \rightarrow D^0\pi^+))$ is $0.13 \pm 0.02 \pm 0.012$. Superseded by BORTOLETTO 92.

227 $V-A$ momentum spectrum used to extrapolate below $p = 1$ GeV. We correct the value assuming $B(D^0 \rightarrow K^-\pi^+) = 0.042 \pm 0.006$ and $B(D^{*+} \rightarrow D^0\pi^+) = 0.6 \pm 0.08$. The product branching fraction is $B(B \rightarrow D^{*+}X) \cdot B(D^{*+} \rightarrow \pi^+D^0) \cdot B(D^0 \rightarrow K^-\pi^+) = (68 \pm 15 \pm 9) \times 10^{-4}$.

 $\Gamma(D_s^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{139}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.089 ± 0.011 OUR AVERAGE				
0.084 ± 0.014 ± 0.010		228 ALBRECHT 92G ARG		$e^+e^- \rightarrow \Upsilon(4S)$
0.088 ± 0.013 ± 0.010	257	229 BORTOLETTO90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.109 ± 0.029 ± 0.013		230 HAAS 86	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.120 ± 0.031 ± 0.014		231 ALBRECHT 87H ARG		$e^+e^- \rightarrow \Upsilon(4S)$

228 ALBRECHT 92G reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^\pm \rightarrow \phi\pi^\pm)] = 0.00292 \pm 0.00039 \pm 0.00031$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^\pm) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

229 BORTOLETTO 90 reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^\pm \rightarrow \phi\pi^\pm)] = 0.00306 \pm 0.00047$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^\pm) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

230 HAAS 86 reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^\pm \rightarrow \phi\pi^\pm)] = 0.0038 \pm 0.0010$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^\pm) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $64 \pm 22\%$ decays are 2-body.

231 ALBRECHT 87H reports $[B(B \rightarrow D_s^\pm \text{ anything}) \times B(D_s^\pm \rightarrow \phi\pi^\pm)] = 0.0042 \pm 0.0009 \pm 0.0006$. We divide by our best value $B(D_s^\pm \rightarrow \phi\pi^\pm) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. $46 \pm 16\%$ of $B \rightarrow D_s X$ decays are 2-body. Superseded by ALBRECHT 92G.

 $\Gamma(D_s D, D_s^* D, D_s D^*, \text{ or } D_s^* D^*)/\Gamma(D_s^\pm \text{ anything})$ $\Gamma_{140}/\Gamma_{139}$

VALUE	DOCUMENT ID	TECN	COMMENT
0.57 ± 0.06 OUR AVERAGE			
0.58 ± 0.07 ± 0.09	ALBRECHT 92G ARG		$e^+e^- \rightarrow \Upsilon(4S)$
0.56 ± 0.10	BORTOLETTO90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(D^*(2010)\gamma)/\Gamma_{\text{total}}$ Γ_{141}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 1.1 × 10⁻³	90	232 LESIAK 92	CBAL	$e^+e^- \rightarrow \Upsilon(4S)$

232 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

 $\Gamma(D_s^+ \pi^-, D_s^{*+} \pi^-, D_s^+ \rho^-, D_s^{*+} \rho^-, D_s^+ \pi^0, D_s^{*+} \pi^0, D_s^+ \eta, D_s^{*+} \eta, D_s^+ \rho^0, D_s^{*+} \rho^0, D_s^+ \omega, D_s^{*+} \omega)/\Gamma_{\text{total}}$ Γ_{142}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0005	90	233 ALEXANDER 93B	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

233 ALEXANDER 93B reports $< 4.8 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = 0.035$. This branching ratio limit provides a model-dependent upper limit $|V_{ub}|/|V_{cb}| < 0.16$ at CL=90%.

 $\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{\text{total}}$ Γ_{143}/Γ

VALUE (units 10 ⁻²)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.30 ± 0.17 OUR AVERAGE					
1.3 ± 0.4 ± 0.1	27	234 MASCHMANN 90	CBAL		$e^+e^- \rightarrow \Upsilon(4S)$
1.23 ± 0.27 ± 0.05	120	235 ALBRECHT 87D ARG			$e^+e^- \rightarrow \Upsilon(4S)$
1.35 ± 0.24 ± 0.06	52	236 ALAM 86	CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
1.16 ± 0.16 ± 0.14	121	237 ADRIANI 93J L3			$e^+e^- \rightarrow Z$
1.3 ± 0.2 ± 0.2		238 ADRIANI 92 L3			$e^+e^- \rightarrow Z$
1.21 ± 0.13 ± 0.08		BUSKULIC 92G ALEP			$e^+e^- \rightarrow Z$
1.4 ± 0.6 -0.5	7	239 ALBRECHT 85H ARG			$e^+e^- \rightarrow \Upsilon(4S)$
1.1 ± 0.21 ± 0.23	46	240 HAAS 85	CLEO	Repl. by ALAM 86	
< 4.9	90	MATTEUZZI 83	MRK2	$E_{\text{cm}}^{\text{ee}} = 29$ GeV	

234 MASCHMANN 90 reports $1.12 \pm 0.33 \pm 0.25$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

235 ALBRECHT 87D reports $1.07 \pm 0.16 \pm 0.22$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. ALBRECHT 87D find the branching ratio for J/ψ not from $\psi(2S)$ to be 0.0081 ± 0.0023 .

236 ALAM 86 reports $1.09 \pm 0.16 \pm 0.21$ for $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = 0.074 \pm 0.012$. We rescale to our best value $B(J/\psi(1S) \rightarrow \mu^+\mu^-) = (5.97 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

237 ADRIANI 93J is an inclusive measurement from b decays at the Z. Uses $J/\psi(1S) \rightarrow \mu^+\mu^-$ and $J/\psi(1S) \rightarrow e^+e^-$ channels.

238 ADRIANI 92 measurement is an inclusive result for $B(Z \rightarrow J/\psi(1S)X) = (4.1 \pm 0.7 \pm 0.3) \times 10^{-3}$ which is used to extract the b -hadron contribution to $J/\psi(1S)$ production. Superseded by ADRIANI 93J.

239 Statistical and systematic errors were added in quadrature. ALBRECHT 85H also report a CL = 90% limit of 0.007 for $B \rightarrow J/\psi(1S) + X$ where $m_X < 1$ GeV.

240 Dimuon and dielectron events used.

 $\Gamma(\chi_{c1}(1P) \text{ anything})/\Gamma_{\text{total}}$ Γ_{144}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0046 ± 0.0017 ± 0.0011	8	ALBRECHT 87D ARG		$e^+e^- \rightarrow \Upsilon(4S)$

 $\Gamma(\chi_{c1}(1P) \text{ anything})/\Gamma_{\text{total}}$ Γ_{145}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0105 ± 0.0035 ± 0.0025	241	ALBRECHT 92E ARG		$e^+e^- \rightarrow \Upsilon(4S)$
0.024 ± 0.009 ± 0.002	19	242 ADRIANI 93J L3		$e^+e^- \rightarrow Z$

241 ALBRECHT 92E assumes no $\chi_{c2}(1P)$ production.

242 ADRIANI 93J is an inclusive measurement and assumes χ_{c1} come from b decays at Z. Uses $J/\psi(1S) \rightarrow \mu^+\mu^-$ channel.

 $\Gamma(\chi_{c1}(1P) \text{ anything})/\Gamma(J/\psi(1S) \text{ anything})$ $\Gamma_{145}/\Gamma_{143}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
1.92 ± 0.82	121	243 ADRIANI 93J L3		$e^+e^- \rightarrow Z$

243 ADRIANI 93J is a ratio of inclusive measurements from b decays at the Z using only the $J/\psi(1S) \rightarrow \mu^+\mu^-$ channel since some systematics cancel.

 $\Gamma(K^\pm \text{ anything})/\Gamma_{\text{total}}$ Γ_{146}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.85 ± 0.07 ± 0.09	ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.66 ± 0.05 ± 0.07	244 BRODY 82	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
seen	245 GIANNINI 82	CUSB	$e^+e^- \rightarrow \Upsilon(4S)$

244 Assuming $\Upsilon(4S) \rightarrow B\bar{B}$, a total of $3.38 \pm 0.34 \pm 0.68$ kaons per $\Upsilon(4S)$ decay is found (the second error is systematic). In the context of the standard B -decay model, this leads to a value for $(b\text{-quark} \rightarrow c\text{-quark})/(b\text{-quark} \rightarrow \text{all})$ of $1.09 \pm 0.33 \pm 0.13$.

245 GIANNINI 82 at CESR-CUSB observed $1.58 \pm 0.35 K^0$ per hadronic event much higher than 0.82 ± 0.10 below threshold. Consistent with predominant $b \rightarrow cX$ decay.

 $\Gamma(K^+ \text{ anything})/\Gamma_{\text{total}}$ Γ_{147}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.66 ± 0.05 ± 0.07	246 ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
246 ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.			

 $\Gamma(K^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{148}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.19 ± 0.05 ± 0.02	247 ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
247 ALAM 87B measurement relies on lepton-kaon correlations. It does not consider the possibility of $B\bar{B}$ mixing. We have thus removed it from the average.			

 $\Gamma(K^0/\bar{K}^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{149}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.63 ± 0.06 ± 0.06	ALAM 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(b \rightarrow s\gamma)/\Gamma_{\text{total}}$ Γ_{150}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0012	90	248 ADRIANI 93L L3		$e^+e^- \rightarrow Z$

248 ADRIANI 93L result is for $b \rightarrow s\gamma$ is performed inclusively.

$\Gamma(K^*(892)\gamma)/\Gamma_{\text{total}}$ Γ_{151}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.4 × 10 ⁻⁴	90	ALBRECHT 88H ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.5 × 10 ⁻³	90	249 LESIAK 92 CBAL		$e^+e^- \rightarrow \Upsilon(4S)$
-------------------------	----	--------------------	--	-----------------------------------

249 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K_1(1400)\gamma)/\Gamma_{\text{total}}$ Γ_{152}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<4.1 × 10 ⁻⁴	90	ALBRECHT 88H ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.6 × 10 ⁻³	90	250 LESIAK 92 CBAL		$e^+e^- \rightarrow \Upsilon(4S)$
-------------------------	----	--------------------	--	-----------------------------------

250 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K_2^*(1430)\gamma)/\Gamma_{\text{total}}$ Γ_{153}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<8.3 × 10 ⁻⁴	90	ALBRECHT 88H ARG		$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_2(1770)\gamma)/\Gamma_{\text{total}}$ Γ_{154}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.2 × 10 ⁻³	90	251 LESIAK 92 CBAL		$e^+e^- \rightarrow \Upsilon(4S)$

251 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(K_3^*(1780)\gamma)/\Gamma_{\text{total}}$ Γ_{155}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<3.0 × 10 ⁻³	90	ALBRECHT 88H ARG		$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(K_4^*(2045)\gamma)/\Gamma_{\text{total}}$ Γ_{156}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.0 × 10 ⁻³	90	252 LESIAK 92 CBAL		$e^+e^- \rightarrow \Upsilon(4S)$

252 LESIAK 92 set a limit on the inclusive process $B(b \rightarrow s\gamma) < 2.8 \times 10^{-3}$ at 90% CL for the range of masses of 892–2045 MeV, independent of assumptions about s-quark hadronization.

$\Gamma(\phi \text{ anything})/\Gamma_{\text{total}}$ Γ_{157}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.023 ± 0.006 ± 0.005	BORTOLETTO86	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(\text{charmed-baryon anything})/\Gamma_{\text{total}}$ Γ_{158}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.064 ± 0.008 ± 0.008		253 CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.14 ± 0.09		254 ALBRECHT 88E ARG		$e^+e^- \rightarrow \Upsilon(4S)$
-------------	--	----------------------	--	-----------------------------------

<0.112	90	255 ALAM 87 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
--------	----	------------------	--	-----------------------------------

253 CRAWFORD 92 result derived from lepton baryon correlations. Assumes all charmed baryons in B^0 and B^\pm decay are Λ_c .

254 ALBRECHT 88E measured $B(B \rightarrow \Lambda_c^+ X) \cdot B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ and used $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (2.2 \pm 1.0)\%$ from ABRAMS 80 to obtain above number.

255 Assuming all baryons result from charmed baryons, ALAM 86 conclude the branching fraction is $7.4 \pm 2.9\%$. The limit given above is model independent.

$\Gamma(\Sigma_c^{--} \text{ anything})/\Gamma_{\text{total}}$ Γ_{159}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0048 ± 0.0024 ± 0.0006	77	256 PROCARIO 94 CLE2		$e^+e^- \rightarrow \Upsilon(4S)$

256 PROCARIO 94 reports $[B(B \rightarrow \Sigma_c^{--} \text{ anything}) \times B(\Lambda_c^+ \rightarrow pK^- \pi^+)] = 0.00021 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma_c^- \text{ anything})/\Gamma_{\text{total}}$ Γ_{160}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.011	90	257 PROCARIO 94 CLE2		$e^+e^- \rightarrow \Upsilon(4S)$

257 PROCARIO 94 reports $[B(B \rightarrow \Sigma_c^- \text{ anything}) \times B(\Lambda_c^+ \rightarrow pK^- \pi^+)] < 0.00048$. We divide by our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.044$.

$\Gamma(\Sigma_c^0 \text{ anything})/\Gamma_{\text{total}}$ Γ_{161}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0053 ± 0.0024 ± 0.0007	76	258 PROCARIO 94 CLE2		$e^+e^- \rightarrow \Upsilon(4S)$

258 PROCARIO 94 reports $[B(B \rightarrow \Sigma_c^0 \text{ anything}) \times B(\Lambda_c^+ \rightarrow pK^- \pi^+)] = 0.00023 \pm 0.00008 \pm 0.00007$. We divide by our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = (4.4 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(\Sigma_c^0 N(N = p \text{ or } n))/\Gamma_{\text{total}}$ Γ_{162}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0017		259 PROCARIO 94 CLE2		$e^+e^- \rightarrow \Upsilon(4S)$

259 PROCARIO 94 reports < 0.0017 for $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^- \pi^+) = 0.044$.

$[\Gamma(p \text{ anything}) + \Gamma(\bar{p} \text{ anything})]/\Gamma_{\text{total}}$ Γ_{163}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.080 ± 0.005 OUR AVERAGE		CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
0.082 ± 0.005 ^{+0.013} _{-0.010}	2163	260 ALBRECHT 89K ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.021		261 ALAM 83B CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
--------	--	-------------------	--	-----------------------------------

260 ALBRECHT 89K include direct and nondirect protons.

261 ALAM 83B reported their result as $> 0.036 \pm 0.006 \pm 0.009$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow p + X) = 0.03$ not including protons from Λ decays.

$[\Gamma(p \text{ (direct) anything}) + \Gamma(\bar{p} \text{ (direct) anything})]/\Gamma_{\text{total}}$ Γ_{164}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.056 ± 0.007 OUR AVERAGE		CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
0.056 ± 0.006 ± 0.005		1220	262 ALBRECHT 89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

262 ALBRECHT 89K subtract contribution of Λ decay from the inclusive proton yield.

$[\Gamma(\Lambda \text{ anything}) + \Gamma(\bar{\Lambda} \text{ anything})]/\Gamma_{\text{total}}$ Γ_{165}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.040 ± 0.005 OUR AVERAGE		CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
0.038 ± 0.004 ± 0.006	2998	943	ALBRECHT 89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

>0.011		263 ALAM 83B CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
--------	--	-------------------	--	-----------------------------------

263 ALAM 83B reported their result as $> 0.022 \pm 0.007 \pm 0.004$. Values are for $(B(\Lambda X) + B(\bar{\Lambda} X))/2$. Data are consistent with equal yields of p and \bar{p} . Using assumed yields below cut, $B(B \rightarrow \Lambda X) = 0.03$.

$[\Gamma(\Xi^- \text{ anything}) + \Gamma(\Xi^+ \text{ anything})]/\Gamma_{\text{total}}$ Γ_{166}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0027 ± 0.0006 OUR AVERAGE		CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
0.0027 ± 0.0005 ± 0.0004	147	54	ALBRECHT 89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

0.0028 ± 0.0014

$\Gamma(\text{baryons anything})/\Gamma_{\text{total}}$ Γ_{167}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.068 ± 0.005 ± 0.003	264 ALBRECHT 920 ARG		$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.076 ± 0.014		265 ALBRECHT 89K ARG		$e^+e^- \rightarrow \Upsilon(4S)$
---------------	--	----------------------	--	-----------------------------------

264 ALBRECHT 920 result is from simultaneous analysis of p and Λ yields, $p\bar{p}$ and $\Lambda\bar{\Lambda}$ correlations, and various lepton-baryon and lepton-baryon-antibaryon correlations. Supersedes ALBRECHT 89K.

265 ALBRECHT 89K obtain this result by adding their their measurements (5.5 ± 1.6)% for direct protons and (4.2 ± 0.5 ± 0.6)% for inclusive Λ production. They then assume (5.5 ± 1.6)% for neutron production and add it in also. Since each B decay has two baryons, they divide by 2 to obtain (7.6 ± 1.4)%.

$\Gamma(p\bar{p} \text{ anything})/\Gamma_{\text{total}}$ Γ_{168}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0247 ± 0.0023 OUR AVERAGE		CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
0.024 ± 0.001 ± 0.004		918	ALBRECHT 89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

$\Gamma(p\bar{p} \text{ anything})/[\Gamma(p \text{ anything}) + \Gamma(\bar{p} \text{ anything})]$ $\Gamma_{168}/\Gamma_{163}$

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
0.30 ± 0.02 ± 0.05	CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$

$[\Gamma(\Lambda\bar{p} \text{ anything}) + \Gamma(\bar{\Lambda}p \text{ anything})]/\Gamma_{\text{total}}$ Γ_{169}/Γ

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.025 ± 0.004 OUR AVERAGE		CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$
0.029 ± 0.005 ± 0.005		165	ALBRECHT 89K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

0.023 ± 0.004 ± 0.003

$[\Gamma(\Lambda\bar{p} \text{ anything}) + \Gamma(\bar{\Lambda}p \text{ anything})]/[\Gamma(\Lambda \text{ anything}) + \Gamma(\bar{\Lambda} \text{ anything})]$ $\Gamma_{169}/\Gamma_{165}$

Includes p and \bar{p} from Λ and $\bar{\Lambda}$ decay.

VALUE	DOCUMENT ID	TECN	COMMENT
0.76 ± 0.11 ± 0.08	CRAWFORD 92 CLEO		$e^+e^- \rightarrow \Upsilon(4S)$

Meson Full Listings

B^\pm

$\Gamma(\Lambda\text{anything})/\Gamma_{\text{total}}$					Γ_{170}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.005	90		CRAWFORD 92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0088	90	12	ALBRECHT 89k	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	----	--------------	-----	-----------------------------------

$\Gamma(\Lambda\text{anything})/[\Gamma(\Lambda\text{anything}) + \Gamma(\Lambda\text{anything})]$					$\Gamma_{170}/\Gamma_{165}$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.13	90	CRAWFORD 92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

$\Gamma(e^+e^- \text{ anything})/\Gamma_{\text{total}}$					Γ_{171}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0024	OUR LIMIT	OUR 90% CL limit, using $[\Gamma(e^+e^- \text{ anything}) + \Gamma(\mu^+\mu^- \text{ anything})]/\Gamma_{\text{total}}$ below.			

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.05	90	BEBEK 81	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	
-------	----	----------	------	-----------------------------------	--

$[\Gamma(e^+e^- \text{ anything}) + \Gamma(\mu^+\mu^- \text{ anything})]/\Gamma_{\text{total}}$					$(\Gamma_{171} + \Gamma_{172})/\Gamma$
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0024	90	266 BEAN 87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0062	90	267 AVERY 84	CLEO	Repl. by BEAN 87	
<0.008	90	MATTEUZZI 83	MRK2	$E_{\text{cm}}^{\text{eff}} = 29 \text{ GeV}$	

266 BEAN 87 reports $[(\mu^+\mu^-) + (e^+e^-)]/2$ and we converted it.
267 Determine ratio of $B^+ \rightarrow B^0$ semileptonic decays to be in the range 0.25–2.9.

$\Gamma(\mu^+\mu^- \text{ anything})/\Gamma_{\text{total}}$					Γ_{172}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<5.0 x 10 ⁻⁵	90	268 ALBAJAR 91c	UA1	$E_{\text{cm}}^{\text{eff}} = 630 \text{ GeV}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.02	95	ALTHOFF 84g	TASS	$E_{\text{cm}}^{\text{eff}} = 34.5 \text{ GeV}$	
<0.007	95	ADEVA 83	MRKJ	$E_{\text{cm}}^{\text{eff}} = 30\text{--}38 \text{ GeV}$	
<0.007	95	BARTEL 83b	JADE	$E_{\text{cm}}^{\text{eff}} = 33\text{--}37 \text{ GeV}$	
<0.017	90	CHADWICK 81	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

268 B^0, B^{\pm} , and B_s^0 not separated.

B^\pm REFERENCES

ABE 94D PRL 72 3456 +Albrow, Amidei, Anway-Wiese, Apollinari (CDF Collab.)
 ABREU 94C ZPHY C63 (to be pub.)+Adam, Adye, Agasi, Aleksan+ (DELPHI Collab.)
 CERN-PPE/94-04
 ALAM 94 PR D (to be pub.) +Kim, Nemat, O'Neill, Severini+ (CLEO Collab.)
 CLNS 94-1270
 PDG 94 PR D50 1173 Montanet+ (CERN, LBL, BOST, IFIC+)
 PROCARIO 94 PRL (to be pub.) +Balet, Cho, Daoudi, Ford+ (CLEO Collab.)
 CLNS 93/1264, CLEO 93-24
 STONE 94 HEP59 93-11
 ABE 93E PL B313 288 +Amako, Arai, Arima, Asano+ (VENUS Collab.)
 ABE 93J PRL 71 3421 +Albrow, Amidei, Anway-Wiese+ (CDF Collab.)
 ABREU 93C PL B301 145 +Adam, Adye, Agasi, Aleksan+ (DELPHI Collab.)
 ABREU 93D ZPHY C57 181 +Adam, Adye, Agasi, Alekseev+ (DELPHI Collab.)
 ABREU 93E PL B312 253 +Adam, Adye, Agasi, Ajinenko+ (DELPHI Collab.)
 ACTON 93C PL B307 247 +Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
 ACTON 93L ZPHY C60 217 +Akers, Alexander, Allison, Anderson+ (OPAL Collab.)
 ADRIANI 93J PL B317 467 +Aguiar-Benitez, Ahlen, Alcaraz+ (L3 Collab.)
 ADRIANI 93K PL B317 474 +Aguiar-Benitez, Ahlen, Alcaraz+ (L3 Collab.)
 ADRIANI 93L PL B317 637 +Aguiar-Benitez, Ahlen, Alcaraz+ (L3 Collab.)
 AKERS 93B ZPHY C60 199 +Alexander, Allison, Anderson, Arcelli+ (OPAL Collab.)
 ALBRECHT 93 ZPHY C57 533 +Ehrlichmann, Hamacher, Hofmann+ (ARGUS Collab.)
 ALBRECHT 93E ZPHY C60 11 +Ehrlichmann, Hamacher, Hofmann+ (ARGUS Collab.)
 ALBRECHT 93H PL B318 397 +Ehrlichmann, Hamacher, Hofmann+ (ARGUS Collab.)
 ALEXANDER 93B PL B319 365 +Bebek, Berkelman, Bloom, Browder+ (CLEO Collab.)
 AMMAR 93 PRL 71 674 +Ball, Baringer, Coppage, Copty+ (CLEO Collab.)
 ARTUSO 93 PL B311 307 (SYRA)
 BARTELT 93B PRL 71 4111 +Csorna, Egyed, Jain, Akerib+ (CLEO Collab.)
 BEAN 93B PRL 70 2681 +Gronberg, Kutschke, Menary, Morrison+ (CLEO Collab.)
 BUSKULIC 93B PL B298 479 +Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
 BUSKULIC 93D PL B307 194 +Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
 BUSKULIC 93O PL B314 459 +De Bonis, Decamp, Ghez, Goy+ (ALEPH Collab.)
 SANGHERA 93 PR D47 791 +Skwarnicki, Stroynowski, Artuso, Goldberg+ (CLEO Collab.)
 ABREU 92 ZPHY C53 567 +Adam, Adams, Adye+ (DELPHI Collab.)
 ACTON 92 PL B274 513 +Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
 ADRIANI 92 PL B288 412 +Aguiar-Benitez, Ahlen, Akbari+ (L3 Collab.)
 ALBRECHT 92C PL B275 195 +Ehrlichmann, Hamacher, Krueger, Nau+ (ARGUS Collab.)
 ALBRECHT 92E PL B277 209 +Ehrlichmann, Hamacher, Krueger, Nau+ (ARGUS Collab.)
 ALBRECHT 92G ZPHY C54 1 +Ehrlichmann, Hamacher, Krueger, Nau+ (ARGUS Collab.)
 ALBRECHT 92O ZPHY C56 1 +Cronstroem, Ehrlichmann+ (ARGUS Collab.)
 BORTOLETTO 92 PR D45 21 +Brown, Dominick, McIlwain+ (CLEO Collab.)
 BUSKULIC 92F PL B295 174 +Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
 BUSKULIC 92G PL B295 396 +Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
 CRAWFORD 92 PR D45 752 +Fulton, Jensen, Johnson+ (CLEO Collab.)
 HENDERSON 92 PL D45 2212 +Kinoshita, Pipkin, Procario+ (CLEO Collab.)
 KRAMER 92 PL B279 181 +Palmer (HAMB, OSU)
 LESIAK 92 ZPHY C55 33 +Antreasyan, Bartels, Besset, Bieler+ (Crystal Ball Collab.)

ADEVA 91C PL B261 177
 ADEVA 91H PL B270 111
 ALBAJAR 91C PL B262 163
 ALBRECHT 91B PL B254 288
 ALBRECHT 91C PL B255 297
 ALBRECHT 91E PL B262 148
 ALBRECHT 91H ZPHY C52 353
 ALEXANDER 91G PL B266 485
 BERKELMAN 91 ARNPS 41 1
 "Decays of B Mesons"
 DECAMP 91C PL B257 492
 FULTON 91 PR D43 651
 YANAGISAWA 91 PRL 66 2436
 ALBRECHT 90 PL B234 409
 ALBRECHT 90B PL B241 278
 ALBRECHT 90H PL B249 359
 ALBRECHT 90J ZPHY C48 543
 ANTREASYAN 90B ZPHY C48 553
 BERTHEND 90D ZPHY C47 333
 BORKLETT 90 PRL 64 2117
 Also 92 PR D45 21
 FULTON 90 PRL 64 16
 HAGEMANN 90 ZPHY C48 401
 LYONS 90 PR D41 982
 MASCHMANN 90 ZPHY C46 555
 PDG 90 PL B239
 WEIR 90B PR D41 1384
 ALBRECHT 89G PL B229 304
 ALBRECHT 89K ZPHY C42 519
 AVERY 89B PL B223 470
 BEBEK 89 PRL 62 8
 BORTOLETTO 89 PRL 62 2436
 BRAUNSCHEW... 89B ZPHY C44 1
 ISGUR 89B PR D39 799
 ONG 89 PRL 62 1236
 WACHS 89 ZPHY C42 33
 ALBRECHT 88F PL B210 265
 ALBRECHT 88E PL B209 119
 ALBRECHT 88G PL B210 258
 ALBRECHT 88K PL B215 424
 KLEM 88 PR D37 41
 KOERNER 88 ZPHY C38 511
 OERG 88 PRL 60 2587
 ALAM 87 PRL 59 22
 ALAM 87B PRL 58 1814
 ALBRECHT 87C PL B195 218
 ALBRECHT 87D PL B199 451
 ALBRECHT 87H PL B187 425
 ASH 87 PRL 58 640
 AVERY 87 PL B183 429
 BARTELT 87 ZPHY C33 339
 BEAN 87 PR D35 3533
 BEBEK 87 PR D36 1289
 BEHREND 87 PRL 59 407
 BORTOLETTO 87 PR D35 19
 BROM 87 PL B195 301
 ALAM 86 PR D34 3279
 BALTRUSAIT... 86E PRL 56 2140
 BARTELT 86B ZPHY C31 349
 BORTOLETTO 86 PRL 56 800
 HAAS 86 PRL 56 2781
 PAL 86 PR D33 2708
 PDG 86 PL 170B
 AIHARA 85 ZPHY C27 39
 ALBRECHT 85H PL 162B 395
 BARTELT 85J PL 163B 277
 CSORNA 85 PRL 54 1894
 HAAS 85 PRL 55 1248
 ALTHOFF 84G ZPHY C22 219
 ALTHOFF 84H PL 149B 524
 ALTHOFF 84J PL 146B 443
 AVERY 84 PRL 53 1309
 CHEN 84 PRL 52 1084
 GILES 84 PR D30 2279
 KLEM 84 PRL 53 1873
 KOOP 84 PRL 52 970
 LEVMAN 84 PL 141B 271
 ADEVA 83 PRL 50 799
 ADEVA 83B PRL 51 443
 ALAM 83S PRL 51 1143
 BARTELT 83B PL 132B 241
 FERNANDEZ 83B PRL 51 1022
 FERNANDEZ 83D PRL 50 2054
 GREEN 83 PRL 51 347
 KLOPFEN... 83B PL 130B 444
 LOCKYER 83 PRL 51 1316
 MATTEUZZI 83 PL 129B 141
 NELSON 83 PRL 50 1542
 ALTARELLI 82 NP B208 365
 BARTELT 82C PL 114B 71
 BRODY 82 PRL 48 1070
 GIANNINI 82 NP B206 1
 SEBEK 81 PRL 46 84
 CHADWICK 81 PRL 46 88
 ABRAMS 80 PRL 44 10
 +Adriani, Aguiar-Benitez, Akbari+ (L3 Collab.)
 +Adrani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
 +Albrow, Alkoffer, Anokviak, Aspimov+ (UA1 Collab.)
 +Glaeser, Harder, Krueger, Nippe+ (ARGUS Collab.)
 +Ehrlichmann, Glaeser, Harder, Krueger+ (ARGUS Collab.)
 +Glaeser, Harder, Krueger, Nippe+ (ARGUS Collab.)
 +Ehrlichmann, Hamacher, Harder+ (ARGUS Collab.)
 +Allison, Allport+ (OPAL Collab.)
 +Stone (CORN, SYRA)
 +Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
 +Jensen, Johnson, Kagan, Kass+ (CLEO Collab.)
 +Heintz, Lee-Franzini, Lovelock, Narain+ (CUSB II Collab.)
 +Glaeser, Harder, Krueger+ (ARGUS Collab.)
 +Ehrlichmann, Glaeser, Harder, Krueger+ (ARGUS Collab.)
 +Ehrlichmann, Harder, Krueger+ (ARGUS Collab.)
 +Bartels, Bieler, Bienlein, Bizzeti+ (Crystal Ball Collab.)
 +Criegee, Field, Franke, Jung+ (CLEO Collab.)
 +Goldberg, Horowitz, Jain, Mestayer+ (CLEO Collab.)
 +Bortoletto, Brown, Dominick, McIlwain+ (CLEO Collab.)
 +Hempstead, Jensen, Johnson+ (CLEO Collab.)
 +Ramcke, Allison, Ambrus, Barlow+ (JADE Collab.)
 +Martin, Saxon (OXF, BRIS, RAL)
 +Antreasyan, Bartels, Besset+ (Crystal Ball Collab.)
 +Hernandez, Stone, Porter+ (IFIC, BOST, CIT+)
 +Klein, Abrams, Adolphsen, Akerlof+ (Mark II Collab.)
 +Glaeser, Harder, Krueger+ (ARGUS Collab.)
 +Boeckmann, Glaeser, Harder+ (ARGUS Collab.)
 +Besson, Garret, Yelton+ (CLEO Collab.)
 +Berkelman, Blucher+ (CLEO Collab.)
 +Goldberg, Horwitz, Mestayer+ (CLEO Collab.)
 +Braunschweig, Gerhards, Kirschfink+ (TASSO Collab.)
 +Scora, Grinstein, Wise (TNTO, CIT)
 +Jaros, Abrams, Amidei, Baden+ (Mark II Collab.)
 +Antreasyan, Bartels, Bieler+ (Crystal Ball Collab.)
 +Boeckmann, Glaeser+ (ARGUS Collab.)
 +Boeckmann, Glaeser+ (ARGUS Collab.)
 +Boeckmann, Glaeser+ (ARGUS Collab.)
 +Boeckmann, Glaeser+ (ARGUS Collab.)
 +Atwood, Barish+ (DELCO Collab.)
 +Schuler (MANZ, DESY)
 +Weir, Abrams, Amidei+ (Mark II Collab.)
 +Kitakama, Kim, Li+ (CLEO Collab.)
 +Katayama, Kim, Sun+ (CLEO Collab.)
 +Binder, Boeckmann, Glaeser+ (ARGUS Collab.)
 +Andam, Binder, Boeckmann+ (ARGUS Collab.)
 +Binder, Boeckmann, Glaeser+ (ARGUS Collab.)
 +Band, Bloom, Bosman+ (MAC Collab.)
 +Besson, Bowcock, Giles+ (CLEO Collab.)
 +Becker, Felst, Haidt+ (JADE Collab.)
 +Bobnik, Brock, Engler+ (CLEO Collab.)
 +Berkelman, Blucher, Cassel+ (CLEO Collab.)
 +Morrow, Guida, Guida+ (CLEO Collab.)
 +Chen, Garren, Goldberg+ (CLEO Collab.)
 +Abachi, Akerlof, Baringer+ (HRS Collab.)
 +Katayama, Kim, Sun+ (CLEO Collab.)
 +Baltrusaitis, Becker, Blaylock, Brown+ (Mark III Collab.)
 +Becker, Cords, Felst+ (JADE Collab.)
 +Chen, Garren, Goldberg+ (CLEO Collab.)
 +Hempstead, Jensen, Kagan+ (CLEO Collab.)
 +Atwood, Barish, Boneaud+ (DELCO Collab.)
 +Aguiar-Benitez, Porter+ (CERN, CIT+)
 +Alston-Garnjost, Badke, Bakken+ (TPC Collab.)
 +Binder, Harder+ (ARGUS Collab.)
 +Becker, Cords, Felst+ (JADE Collab.)
 +Garren, Mestayer, Panvini+ (CLEO Collab.)
 +Hempstead, Jensen, Kagan+ (CLEO Collab.)
 +Braunschweig, Kirschfink+ (TASSO Collab.)
 +Braunschweig, Kirschfink+ (TASSO Collab.)
 +Bebek, Berkelman, Cassel+ (CLEO Collab.)
 +Goldberg, Horowitz, Jawahery+ (CLEO Collab.)
 +Hassard, Hempstead, Kinoshita+ (CLEO Collab.)
 +Dubois, Young, Atwood+ (DELCO Collab.)
 +Sakuda, Atwood, Balloun+ (CLEO Collab.)
 +Sreedhar, Han, Imlay+ (CUSB Collab.)
 +Barber, Becker, Berdugo+ (Mark-J Collab.)
 +Barber, Becker, Berdugo+ (Mark-J Collab.)
 +Csorna, Garren, Mestayer+ (CLEO Collab.)
 +Becker, Bowdery, Cords+ (JADE Collab.)
 +Ford, Read, Smith+ (MAC Collab.)
 +Ford, Read, Smith+ (MAC Collab.)
 +Hicks, Sannes, Skubic+ (CLEO Collab.)
 +Klopfenstein, Horstotte+ (CUSB Collab.)
 +Jaros, Nelson, Abrams+ (Mark II Collab.)
 +Abrams, Amidei, Blocker+ (Mark II Collab.)
 +Blondel, Trilling, Abrams+ (Mark II Collab.)
 +Cabibbo, Corbo, Maini, Martinelli (ROMA, INFN, FRAS)
 +Cords, Dittmann, Eichler+ (JADE Collab.)
 +Chen, Goldberg, Horwitz+ (CLEO Collab.)
 +Finocchiaro, Franzini+ (CUSB Collab.)
 +Haggerty, Izen, Longuemare+ (CLEO Collab.)
 +Gandl, Kagar, Kass+ (CLEO Collab.)
 +Alam, Blocker, Boyarski+ (SLAC, LBL)

OTHER RELATED PAPERS

BERKELMAN 91 ARNPS 41 1 + Stone (CORN, SYRA)
 "Decays of B Mesons"
 MILLER 90 MPL A5 2683
 "Recent Results in B Physics"
 SCHINDLER 88 High Energy Electron-Positron Physics 234 (SLAC)
 Editors: A. Ali and P. Soeding, World Scientific, Singapore
 SCHUBERT 87 IHEP-HD/87-7 (HEIDH)
 EPS Conference - Uppsala, Proc., Vol. 2, p. 791

B⁰

$$I(J^P) = \frac{1}{2}(0^-)$$

Quantum numbers not measured. Values shown are quark-model predictions.

For measurements of the *B* mean life and for branching ratios in which the charge of the decaying *B* is not determined, see the *B*[±] section.

See the Notes "Experimental Highlights of *B* Meson Production and Decay" and "Semileptonic Decays of *B* Mesons" at the beginning of the *B*[±] Full Listings and the Note on "*B*⁰-*B*⁰ Mixing and CP Violation in *B* Decay" near the end of the *B*⁰ Full Listings.

B⁰ MASS

The fit uses *m*_{B[±]} and (*m*_{B⁰} - *m*_{B[±]}) to determine *m*_{B⁰}. *m*_{B⁰} data are excluded from the fit because they are not independent. These experiments actually measure the difference between half of *E*_{cm} and the *B* mass.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5279.0 ± 2.0 OUR FIT				
5278.9 ± 2.0 OUR AVERAGE				
5279.2 ± 0.54 ± 2.0	340	¹ ALAM 94 CLE2	e ⁺ e ⁻ → <i>T</i> (4S)	
5278.0 ± 0.4 ± 2.0		¹ BORTOLETTO92 CLEO	e ⁺ e ⁻ → <i>T</i> (4S)	
5279.6 ± 0.7 ± 2.0	40	^{1,2} ALBRECHT 90J ARG	e ⁺ e ⁻ → <i>T</i> (4S)	
5280.6 ± 0.8 ± 2.0		¹ BEBEK 87 CLEO	e ⁺ e ⁻ → <i>T</i> (4S)	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5278.2 ± 1.0 ± 3.0	40	ALBRECHT 87C ARG	e ⁺ e ⁻ → <i>T</i> (4S)	
5279.5 ± 1.6 ± 3.0	7	³ ALBRECHT 87D ARG	e ⁺ e ⁻ → <i>T</i> (4S)	

- ¹ These experiments all report a common systematic error 2.0 MeV. We have artificially increased the systematic error to allow the experiments to be treated as independent measurements in our average. See "Treatment of Errors" section of the Introductory Text.
- ² ALBRECHT 90J assumes 10580 for *T*(4S) mass. Supersedes ALBRECHT 87C and ALBRECHT 87D.
- ³ Found using fully reconstructed decays with *J/ψ*. ALBRECHT 87D assume *m*_{*T*(4S)} = 10577 MeV.

***m*_{B⁰} - *m*_{B[±]}**

The mass difference measurements are not independent of the *B*[±] and *B*⁰ mass measurement by the same experiments.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
0.34 ± 0.29 OUR FIT	Error includes scale factor of 1.1.		
0.34 ± 0.32 OUR AVERAGE	Error includes scale factor of 1.2.		
0.41 ± 0.25 ± 0.19	ALAM 94 CLE2	e ⁺ e ⁻ → <i>T</i> (4S)	
-0.4 ± 0.6 ± 0.5	BORTOLETTO92 CLEO	e ⁺ e ⁻ → <i>T</i> (4S)	
-0.9 ± 1.2 ± 0.5	ALBRECHT 90J ARG	e ⁺ e ⁻ → <i>T</i> (4S)	
2.0 ± 1.1 ± 0.3	⁴ BEBEK 87 CLEO	e ⁺ e ⁻ → <i>T</i> (4S)	
⁴ BEBEK 87 actually measure the difference between half of <i>E</i> _{cm} and the <i>B</i> [±] or <i>B</i> ⁰ mass, so the <i>m</i> _{B⁰} - <i>m</i> _{B[±]} is more accurate. Assume <i>m</i> _{<i>T</i>(4S)} = 10580 MeV.			

***m*_{B⁰} - *m*_{B[±]}**

See the *B*⁰-*B*⁰ MIXING section near the end of these *B*⁰ Listings.

B⁰ MEAN LIFE

See *B*[±] Full Listings for data on *B* hadron mean life averaged over species of bottom particles.

VALUE (10 ⁻¹² s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.50 ± 0.11 OUR AVERAGE				
1.57 ± 0.18 ± 0.08	121	⁵ ABE 94D CDF	<i>p</i> <i>p</i> at 1.8 TeV	
1.17 ^{+0.29} _{-0.23} ± 0.16	96	⁶ ABREU 93D DLPH	e ⁺ e ⁻ → <i>Z</i>	
1.55 ± 0.25 ± 0.18	76	⁷ ABREU 93G DLPH	e ⁺ e ⁻ → <i>Z</i>	
1.51 ^{+0.24} _{-0.23} ± 0.14	78	⁶ ACTON 93C OPAL	e ⁺ e ⁻ → <i>Z</i>	
1.52 ^{+0.20} _{-0.18} ± 0.07	77	⁶ BUSKULIC 93D ALEP	e ⁺ e ⁻ → <i>Z</i>	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.20 ^{+0.52} _{-0.36} ± 0.14	15	⁸ WAGNER 90 MRK2	<i>E</i> _{cm} ^{ee} = 29 GeV	
0.82 ^{+0.57} _{-0.37} ± 0.27		⁹ AVERILL 89 HRS	<i>E</i> _{cm} ^{ee} = 29 GeV	

- ⁵ ABE 94D measured mean life using fully reconstructed decays.
- ⁶ Data analyzed using *D*/*D*^{*} anything event vertices.
- ⁷ ABREU 93G data analyzed using charged and neutral vertices.
- ⁸ WAGNER 90 tagged *B*⁰ mesons by their decays into *D*^{*-} e⁺ν and *D*^{*-} μ⁺ν where the *D*^{*-} is tagged by its decay into π⁻*D*⁰.
- ⁹ AVERILL 89 is an estimate of the *B*⁰ mean lifetime assuming that *B*⁰ → *D*⁺+ x always.

MEAN LIFE RATIO *τ*_{B⁺}/*τ*_{B⁰}

The measurements at high energy are direct lifetime measurements while those at the *T*(4S) are inferred from the branching fractions for semileptonic decay or other spectator-dominated decays by assuming that the rates for such decays are equal for *B*⁰ and *B*⁺.

VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT
0.98 ± 0.09 OUR AVERAGE				
1.02 ± 0.16 ± 0.05	269	¹⁰ ABE 94D CDF	<i>p</i> <i>p</i> at 1.8 TeV	
1.11 ^{+0.51} _{-0.39} ± 0.11	188	¹¹ ABREU 93D DLPH	e ⁺ e ⁻ → <i>Z</i>	
1.01 ^{+0.29} _{-0.22} ± 0.12	253	¹² ABREU 93G DLPH	e ⁺ e ⁻ → <i>Z</i>	
1.0 ^{+0.33} _{-0.25} ± 0.08	130	ACTON 93C OPAL	e ⁺ e ⁻ → <i>Z</i>	
0.96 ^{+0.19} _{-0.15} ± 0.12	154	¹¹ BUSKULIC 93D ALEP	e ⁺ e ⁻ → <i>Z</i>	
0.91 ± 0.27 ± 0.21		¹³ ALBRECHT 92C ARG	e ⁺ e ⁻ → <i>T</i> (4S)	
1.0 ± 0.4	29	^{13,14} ALBRECHT 92G ARG	e ⁺ e ⁻ → <i>T</i> (4S)	
0.89 ± 0.19 ± 0.13		¹³ FULTON 91 CLEO	e ⁺ e ⁻ → <i>T</i> (4S)	
1.00 ± 0.23 ± 0.14		¹³ ALBRECHT 89L ARG	e ⁺ e ⁻ → <i>T</i> (4S)	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.49 to 2.3	90	¹⁵ BEAN 87B CLEO	e ⁺ e ⁻ → <i>T</i> (4S)	

- ¹⁰ ABE 94D is a direct measurement using fully reconstructed decays.
- ¹¹ Data analyzed using *D*/*D*^{*} anything event vertices.
- ¹² ABREU 93G data analyzed using charged and neutral vertices.
- ¹³ Assumes equal production of *B*⁰ and *B*⁺.
- ¹⁴ ALBRECHT 92G data analyzed using *B* → *D*_s*D*⁻, *D*_s*D*⁺, *D*_s^{*}*D*⁻, *D*_s^{*}*D*⁺ events.
- ¹⁵ BEAN 87B assume the fraction of *B*⁰*B*⁰ events at the *T*(4S) is 0.41.

B⁰ DECAY MODES

*B*⁰ modes are charge conjugates of the modes below. Reactions indicate the weak decay vertex and do not include mixing. Decays in which the charge of the *B* is not determined are in the *B*[±] section.

Only data from *T*(4S) decays are used for branching fractions, with rare exceptions. The branching fractions listed below assume a 50:50 *B*⁰*B*⁺*B*⁻ production ratio at the *T*(4S). We have attempted to bring older measurements up to date by rescaling their assumed *T*(4S) production ratio to 50:50 and their assumed *D*, *D*_s, *D*^{*}, and *ψ* branching ratios to current values whenever this would effect our averages and best limits significantly.

Indentation is used to indicate a subchannel of a previous reaction. All resonant subchannels have been corrected for resonance branching fractions to the final state so the sum of the subchannel branching fractions can exceed that of the final state.

Mode	Fraction (Γ _i /Γ)	Confidence level
Semileptonic and leptonic modes		
Γ ₁ <i>ℓ</i> ⁺ ν _ℓ anything	[a] (9.5 ± 1.6) %	
Γ ₂ <i>D</i> ⁻ ℓ ⁺ ν _ℓ	[a] (1.9 ± 0.5) %	
Γ ₃ <i>D</i> [*] (2010) ⁻ ℓ ⁺ ν _ℓ	[a] (4.4 ± 0.4) %	
Γ ₄ ρ ⁻ ℓ ⁺ ν _ℓ	[a] < 4.1 × 10 ⁻⁴	90%
Γ ₅ π ⁻ μ ⁺ ν _μ	seen	
<i>D</i>, <i>D</i>[*], or <i>D</i>_s modes		
Γ ₆ <i>D</i> ⁻ π ⁺	(3.0 ± 0.4) × 10 ⁻³	
Γ ₇ <i>D</i> ⁻ ρ ⁺	(7.8 ± 1.4) × 10 ⁻³	
Γ ₈ <i>D</i> ⁰ π ⁺ π ⁻	< 1.6 × 10 ⁻³	90%
Γ ₉ <i>D</i> [*] (2010) ⁻ π ⁺	(2.6 ± 0.4) × 10 ⁻³	
Γ ₁₀ <i>D</i> ⁻ π ⁺ π ⁺ π ⁻	(8.0 ± 2.5) × 10 ⁻³	
Γ ₁₁ (<i>D</i> ⁻ π ⁺ π ⁺ π ⁻) nonresonant	(3.9 ± 1.9) × 10 ⁻³	
Γ ₁₂ <i>D</i> ⁻ π ⁺ ρ ⁰	(1.1 ± 1.0) × 10 ⁻³	
Γ ₁₃ <i>D</i> ⁻ a ₁ (1260) ⁺	(6.0 ± 3.3) × 10 ⁻³	
Γ ₁₄ <i>D</i> [*] (2010) ⁻ π ⁺ π ⁰	(1.5 ± 0.5) %	
Γ ₁₅ <i>D</i> [*] (2010) ⁻ ρ ⁺	(7.3 ± 1.5) × 10 ⁻³	
Γ ₁₆ <i>D</i> [*] (2010) ⁻ π ⁺ π ⁺ π ⁻	(1.19 ± 0.27) %	
Γ ₁₇ (<i>D</i> [*] (2010) ⁻ π ⁺ π ⁺ π ⁻) non-resonant	(0.0 ± 2.5) × 10 ⁻³	
Γ ₁₈ <i>D</i> [*] (2010) ⁻ π ⁺ ρ ⁰	(5.7 ± 3.1) × 10 ⁻³	
Γ ₁₉ <i>D</i> [*] (2010) ⁻ a ₁ (1260) ⁺	(1.5 ± 0.7) %	
Γ ₂₀ <i>D</i> [*] (2010) ⁻ π ⁺ π ⁺ π ⁻ π ⁰	(3.4 ± 1.8) %	
Γ ₂₁ <i>D</i> _s [*] (2460) ⁻ π ⁺	< 2.2 × 10 ⁻³	90%
Γ ₂₂ <i>D</i> _s [*] (2460) ⁻ ρ ⁺	< 4.9 × 10 ⁻³	90%
Γ ₂₃ <i>D</i> ⁻ <i>D</i> _s ⁺	(8 ± 4) × 10 ⁻³	
Γ ₂₄ <i>D</i> [*] (2010) ⁻ <i>D</i> _s ⁺	(1.2 ± 0.6) %	
Γ ₂₅ <i>D</i> ⁻ <i>D</i> _s ⁺	(2.1 ± 1.5) %	
Γ ₂₆ <i>D</i> [*] (2010) ⁻ <i>D</i> _s ⁺	(2.0 ± 1.2) %	
Γ ₂₇ <i>D</i> _s ⁺ π ⁻	< 2.9 × 10 ⁻⁴	90%

Meson Full Listings

B^0

Γ ₂₈	$D_s^{*+} \pi^-$	< 5	$\times 10^{-4}$	90%
Γ ₂₉	$D_s^+ \rho^-$	< 7	$\times 10^{-4}$	90%
Γ ₃₀	$D_s^{*+} \rho^-$	< 8	$\times 10^{-4}$	90%
Γ ₃₁	$D_s^+ a_1(1260)^-$	< 2.7	$\times 10^{-3}$	90%
Γ ₃₂	$D_s^{*+} a_1(1260)^-$	< 2.2	$\times 10^{-3}$	90%
Γ ₃₃	$D_s^- K^+$	< 2.4	$\times 10^{-4}$	90%
Γ ₃₄	$D_s^{*-} K^+$	< 1.8	$\times 10^{-4}$	90%
Γ ₃₅	$D_s^- K^*(892)^+$	< 1.0	$\times 10^{-3}$	90%
Γ ₃₆	$D_s^{*-} K^*(892)^+$	< 1.2	$\times 10^{-3}$	90%
Γ ₃₇	$D_s^- \pi^+ K^0$	< 6	$\times 10^{-3}$	90%
Γ ₃₈	$D_s^{*-} \pi^+ K^0$	< 3.2	$\times 10^{-3}$	90%
Γ ₃₉	$D_s^- \pi^+ K^*(892)^0$	< 4	$\times 10^{-3}$	90%
Γ ₄₀	$D_s^{*-} \pi^+ K^*(892)^0$	< 2.1	$\times 10^{-3}$	90%
Γ ₄₁	$\bar{D}^0 \pi^0$	< 4.8	$\times 10^{-4}$	90%
Γ ₄₂	$\bar{D}^0 \rho^0$	< 5.5	$\times 10^{-4}$	90%
Γ ₄₃	$\bar{D}^0 \eta$	< 6.8	$\times 10^{-4}$	90%
Γ ₄₄	$\bar{D}^0 \eta'$	< 8.6	$\times 10^{-4}$	90%
Γ ₄₅	$\bar{D}^0 \omega$	< 6.3	$\times 10^{-4}$	90%
Γ ₄₆	$\bar{D}^*(2007)^0 \pi^0$	< 9.7	$\times 10^{-4}$	90%
Γ ₄₇	$\bar{D}^*(2007)^0 \rho^0$	< 1.17	$\times 10^{-3}$	90%
Γ ₄₈	$\bar{D}^*(2007)^0 \eta$	< 6.9	$\times 10^{-4}$	90%
Γ ₄₉	$\bar{D}^*(2007)^0 \eta'$	< 2.7	$\times 10^{-3}$	90%
Γ ₅₀	$\bar{D}^*(2007)^0 \omega$	< 2.1	$\times 10^{-3}$	90%

Charmonium modes

Γ ₅₁	$J/\psi(1S) K^0$	(7.5 ± 2.1) × 10 ⁻⁴		
Γ ₅₂	$J/\psi(1S) K^+ \pi^-$	(1.2 ± 0.6) × 10 ⁻³		
Γ ₅₃	$J/\psi(1S) K^*(892)^0$	(1.58 ± 0.28) × 10 ⁻³		
Γ ₅₄	$\psi(2S) K^0$	< 8	$\times 10^{-4}$	90%
Γ ₅₅	$\psi(2S) K^+ \pi^-$	< 1	$\times 10^{-3}$	90%
Γ ₅₆	$\psi(2S) K^*(892)^0$	(1.4 ± 0.9) × 10 ⁻³		
Γ ₅₇	$\chi_{c1}(1P) K^0$	< 2.7	$\times 10^{-3}$	90%
Γ ₅₈	$\chi_{c1}(1P) K^*(892)^0$	< 2.1	$\times 10^{-3}$	90%

K or K* modes

Γ ₅₉	$K^+ \pi^-$	< 2.6	$\times 10^{-5}$	90%
Γ ₆₀	$K^+ K^-$	< 7	$\times 10^{-6}$	90%
Γ ₆₁	$K^0 \pi^+ \pi^-$	< 4.4	$\times 10^{-4}$	90%
Γ ₆₂	$K^0 \rho^0$	< 3.2	$\times 10^{-4}$	90%
Γ ₆₃	$K^0 f_0(980)$	< 3.6	$\times 10^{-4}$	90%
Γ ₆₄	$K^*(892)^+ \pi^-$	< 3.8	$\times 10^{-4}$	90%
Γ ₆₅	$K_2^*(1430)^+ \pi^-$	< 2.6	$\times 10^{-3}$	90%
Γ ₆₆	$K^0 K^+ K^-$	< 1.3	$\times 10^{-3}$	90%
Γ ₆₇	$K^0 \phi$	< 4.2	$\times 10^{-4}$	90%
Γ ₆₈	$K^*(892)^0 \pi^+ \pi^-$	< 1.4	$\times 10^{-3}$	90%
Γ ₆₉	$K^*(892)^0 \rho^0$	< 4.6	$\times 10^{-4}$	90%
Γ ₇₀	$K^*(892)^0 f_0(980)$	< 1.7	$\times 10^{-4}$	90%
Γ ₇₁	$K_1(1400)^+ \pi^-$	< 1.1	$\times 10^{-3}$	90%
Γ ₇₂	$K^*(892)^0 K^+ K^-$	< 6.1	$\times 10^{-4}$	90%
Γ ₇₃	$K^*(892)^0 \phi$	< 3.2	$\times 10^{-4}$	90%
Γ ₇₄	$K_1(1400)^0 \rho^0$	< 3.0	$\times 10^{-3}$	90%
Γ ₇₅	$K_1(1400)^0 \phi$	< 5.0	$\times 10^{-3}$	90%
Γ ₇₆	$K_2^*(1430)^0 \rho^0$	< 1.1	$\times 10^{-3}$	90%
Γ ₇₇	$K_2^*(1430)^0 \phi$	< 1.4	$\times 10^{-3}$	90%
Γ ₇₈	$K^*(892)^0 \gamma$	(4.0 ± 1.9) × 10 ⁻⁵		
Γ ₇₉	$K_1(1270)^0 \gamma$	< 7.0	$\times 10^{-3}$	90%
Γ ₈₀	$K_1(1400)^0 \gamma$	< 4.3	$\times 10^{-3}$	90%
Γ ₈₁	$K_2^*(1430)^0 \gamma$	< 4.0	$\times 10^{-4}$	90%
Γ ₈₂	$K^*(1680)^0 \gamma$	< 2.0	$\times 10^{-3}$	90%
Γ ₈₃	$K_3^*(1780)^0 \gamma$	< 1.0	%	90%
Γ ₈₄	$K_4^*(2045)^0 \gamma$	< 4.3	$\times 10^{-3}$	90%

Light unflavored meson modes

Γ ₈₅	$\pi^+ \pi^-$	< 2.9	$\times 10^{-5}$	90%
Γ ₈₆	$\pi^+ \pi^- \pi^0$	< 7.2	$\times 10^{-4}$	90%
Γ ₈₇	$\rho^0 \pi^0$	< 4.0	$\times 10^{-4}$	90%
Γ ₈₈	$\rho^\pm \pi^\pm$	[b] < 5.2	$\times 10^{-4}$	90%
Γ ₈₉	$\pi^+ \pi^- \pi^+ \pi^-$	< 6.7	$\times 10^{-4}$	90%
Γ ₉₀	$\rho^0 \rho^0$	< 2.8	$\times 10^{-4}$	90%
Γ ₉₁	$a_1(1260)^\mp \pi^\pm$	[b] < 4.9	$\times 10^{-4}$	90%
Γ ₉₂	$a_2(1320)^\mp \pi^\pm$	[b] < 3.0	$\times 10^{-4}$	90%

Γ ₉₃	$\pi^+ \pi^- \pi^0 \pi^0$	< 3.1	$\times 10^{-3}$	90%
Γ ₉₄	$\rho^+ \rho^-$	< 2.2	$\times 10^{-3}$	90%
Γ ₉₅	$a_1(1260)^0 \pi^0$	< 1.1	$\times 10^{-3}$	90%
Γ ₉₆	$\omega \pi^0$	< 4.6	$\times 10^{-4}$	90%
Γ ₉₇	$\eta \pi^0$	< 1.8	$\times 10^{-3}$	90%
Γ ₉₈	$\pi^+ \pi^+ \pi^- \pi^- \pi^0$	< 9.0	$\times 10^{-3}$	90%
Γ ₉₉	$a_1(1260)^+ \rho^-$	< 3.4	$\times 10^{-3}$	90%
Γ ₁₀₀	$a_1(1260)^0 \rho^0$	< 2.4	$\times 10^{-3}$	90%
Γ ₁₀₁	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$	< 3.0	$\times 10^{-3}$	90%
Γ ₁₀₂	$a_1(1260)^+ a_1(1260)^-$	< 2.8	$\times 10^{-3}$	90%
Γ ₁₀₃	$\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-$	< 1.1	%	90%

Baryon modes

Γ ₁₀₄	$p \bar{p}$	< 3.4	$\times 10^{-5}$	90%
Γ ₁₀₅	$p \bar{p} \pi^+ \pi^-$	< 2.5	$\times 10^{-4}$	90%
Γ ₁₀₆	$p \Lambda \pi^-$	< 1.8	$\times 10^{-4}$	90%
Γ ₁₀₇	$\Delta^0 \bar{\Delta}^0$	< 1.5	$\times 10^{-3}$	90%
Γ ₁₀₈	$\Delta^{++} \Delta^{--}$	< 1.1	$\times 10^{-4}$	90%
Γ ₁₀₉	$\Sigma_c^- \Delta^{++}$	< 1.2	$\times 10^{-3}$	90%

Lepton Family number (LF) violating modes, $\Delta B = 2$ forbidden decay via mixing (B2M) modes, or $\Delta B = 1$ weak neutral current (B1) modes

Γ ₁₁₀	$e^+ e^-$	B1	< 5.9	$\times 10^{-6}$	90%
Γ ₁₁₁	$\mu^+ \mu^-$	B1	< 5.9	$\times 10^{-6}$	90%
Γ ₁₁₂	$K^0 e^+ e^-$	B1	< 3.0	$\times 10^{-4}$	90%
Γ ₁₁₃	$K^0 \mu^+ \mu^-$	B1	< 3.6	$\times 10^{-4}$	90%
Γ ₁₁₄	$K^*(892)^0 e^+ e^-$	B1	< 2.9	$\times 10^{-4}$	90%
Γ ₁₁₅	$K^*(892)^0 \mu^+ \mu^-$	B1	< 2.3	$\times 10^{-5}$	90%
Γ ₁₁₆	$e^\pm \mu^\mp$	LF	[b] < 5.9	$\times 10^{-6}$	90%
Γ ₁₁₇	$e^\pm \tau^\mp$	LF	[b] < 5.3	$\times 10^{-4}$	90%
Γ ₁₁₈	$\mu^\pm \tau^\mp$	LF	[b] < 8.3	$\times 10^{-4}$	90%

[a] ℓ indicates e or μ mode, not sum over modes.

[b] The value is for the sum of the charge states indicated.

B^0 BRANCHING RATIOS

For branching ratios in which the charge of the decaying B is not determined, see the B^\pm section.

$\Gamma(\ell^+ \nu_\ell \text{ anything})/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ ₁ /Γ
	0.095 ± 0.016 OUR AVERAGE				
	0.093 ± 0.011 ± 0.015	ALBRECHT	94 ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
	0.099 ± 0.030 ± 0.009	HENDERSON	92 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	

$\Gamma(D^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ ₂ /Γ
	0.019 ± 0.005 OUR AVERAGE				
	0.018 ± 0.006 ± 0.003	¹⁶ FULTON	91 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	
	0.020 ± 0.007 ± 0.006	¹⁷ ALBRECHT	89J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	

¹⁶FULTON 91 assumes assuming equal production of B^0 and B^+ at the $\Upsilon(4S)$ and uses Mark III D and D^* branching ratios.

¹⁷ALBRECHT 89J reports 0.018 ± 0.006 ± 0.005. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$.

$\Gamma(D^{*+}(2010)^- \ell^+ \nu_\ell)/\Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ ₃ /Γ
	0.044 ± 0.004 OUR AVERAGE					
	0.045 ± 0.003 ± 0.004		¹⁸ ALBRECHT	94 ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
	0.047 ± 0.005 ± 0.005	235	¹⁹ ALBRECHT	93 ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
	0.040 ± 0.004 ± 0.006		²⁰ BORTOLETTO89B	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
seen		398	²¹ SANGHERA	93 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$	
			²² ALBRECHT	89C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
			²³ ALBRECHT	89J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
			²⁴ ALBRECHT	87J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$	
	0.060 ± 0.010 ± 0.014					
	0.070 ± 0.012 ± 0.019	47				

¹⁸ALBRECHT 94 assumes $B(D^{*+} \rightarrow D^0 \pi^+) = 68.1 \pm 1.0 \pm 1.3\%$. Uses partial reconstruction of D^{*+} and is independent of D^0 branching ratios.

¹⁹ALBRECHT 93 reports 0.052 ± 0.005 ± 0.006. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. We have taken their average e and μ value. They also obtain $\alpha = 2\pi^0/(\Gamma^- + \Gamma^+) - 1 = 1.1 \pm 0.4 \pm 0.2$, $A_{AF} = 3/4 * (\Gamma^- - \Gamma^+)/\Gamma = 0.2 \pm 0.08 \pm 0.06$ and a value of $|V_{cb}| = 0.036 - 0.045$ depending on model assumptions.

²⁰We have taken average of the the BORTOLETTO 89B values for electrons and muons, 0.046 ± 0.005 ± 0.007. We rescale using the method described in STONE 94 but with the updated PDG 94 $B(D^0 \rightarrow K^- \pi^+)$. The measurement suggests a D^* polarization parameter value $\alpha = 0.65 \pm 0.66 \pm 0.25$.

21 Combining $\bar{D}^{*0} \ell^+ \nu_\ell$ and $\bar{D}^{*0} \ell^+ \nu_\ell$ SANGHERA 93 test $V-A$ structure and fit the decay angular distributions to obtain $A_{FB} = 3/4 * (\Gamma^- - \Gamma^+) / \Gamma = 0.14 \pm 0.06 \pm 0.03$. Assuming a value of V_{cb} , they measure V_{A1} , and A_2 , the three form factors for the $D^* \ell \nu_\ell$ decay, where results are slightly dependent on model assumptions.

22 The measurement of ALBRECHT 89C suggests a D^* polarization γ_L / γ_T of 0.85 ± 0.45 , or $\alpha = 0.7 \pm 0.9$.

23 ALBRECHT 89J is ALBRECHT 87J value rescaled using $B(D^*(2010)^- \rightarrow D^0 \pi^-) = 0.57 \pm 0.04 \pm 0.04$. Superseded by ALBRECHT 93.

24 ALBRECHT 87J assume $\mu-e$ universality, the $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.45$, the $B(D^0 \rightarrow K^- \pi^+) = (0.042 \pm 0.004 \pm 0.004)$, and the $B(D^*(2010)^- \rightarrow D^0 \pi^-) = 0.49 \pm 0.08$. Superseded by ALBRECHT 89J.

$\Gamma(\rho^- \ell^+ \nu_\ell) / \Gamma_{total}$ Γ_4 / Γ
 $\ell = e$ or μ , not sum over e and μ modes.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$< 4.1 \times 10^{-4}$	90	25 BEAN	93B CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$

25 BEAN 93B limit set using ISGW Model. Using isospin and the quark model to combine $\Gamma(\rho^0 \ell^+ \nu_\ell)$ and $\Gamma(\omega \ell^+ \nu_\ell)$ with this result, they obtain a limit $< (1.6-2.7) \times 10^{-4}$ at 90% CL for $B^+ \rightarrow (\omega$ or $\rho^0) \ell^+ \nu_\ell$. The range corresponds to the ISGW, WSB, and KS models. An upper limit on $|V_{ub}/V_{cb}| < 0.8-0.13$ at 90% CL is derived as well.

$\Gamma(\pi^- \mu^+ \nu_\mu) / \Gamma_{total}$ Γ_5 / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	26 ALBRECHT	91C ARG	

26 In ALBRECHT 91C, one event is fully reconstructed providing evidence for the $b \rightarrow u$ transition.

$\Gamma(D^- \pi^+) / \Gamma_{total}$ Γ_6 / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0030 ± 0.0004 OUR AVERAGE				
0.0029 ± 0.0004 ± 0.0002	81	27 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0027 ± 0.0006 ± 0.0005		28 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0048 ± 0.0011 ± 0.0011	22	29 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0051 ± 0.0028 ± 0.0013 - 0.0025 - 0.0012	4	30 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 0.0031 ± 0.0013 ± 0.0010 7 29 ALBRECHT 88K ARG $e^+ e^- \rightarrow \Upsilon(4S)$

27 ALAM 94 reports $[B(B^0 \rightarrow D^- \pi^+) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.000265 \pm 0.000032 \pm 0.000023$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

28 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

29 ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

30 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

$\Gamma(D^- \rho^+) / \Gamma_{total}$ Γ_7 / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0078 ± 0.0014 OUR AVERAGE				
0.0078 ± 0.0013 ± 0.0005	81	31 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.009 ± 0.005 ± 0.003	9	32 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •
 0.022 ± 0.012 ± 0.009 6 32 ALBRECHT 88K ARG $e^+ e^- \rightarrow \Upsilon(4S)$

31 ALAM 94 reports $[B(B^0 \rightarrow D^- \rho^+) \times B(D^+ \rightarrow K^- \pi^+ \pi^+)] = 0.000704 \pm 0.000096 \pm 0.000070$. We divide by our best value $B(D^+ \rightarrow K^- \pi^+ \pi^+) = (9.1 \pm 0.6) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

32 ALBRECHT 88K assumes $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. Superseded by ALBRECHT 90J which assumes 50:50.

$\Gamma(D^0 \pi^+ \pi^-) / \Gamma_{total}$ Γ_8 / Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.0016	90		33 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
< 0.007	90		34 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
< 0.034	90		35 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.07 ± 0.05		5	36 BEHREND	83 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

33 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

34 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D . The product branching fraction into $D_2^0(2340) \pi$ followed by $D_2^0(2340) \rightarrow D^0 \pi$ is < 0.0001 at 90% CL and into $D_2^0(2460) \pi$ followed by $D_2^0(2460) \rightarrow D^0 \pi$ is < 0.0004 at 90% CL.

35 BEBEK 87 assume the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%. $B(D^0 \rightarrow K^- \pi^+) = (4.2 \pm 0.4 \pm 0.4)\%$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) = (9.1 \pm 0.8 \pm 0.8)\%$ were used.

36 Corrected by us using assumptions: $B(D^0 \rightarrow K^- \pi^+) = (0.042 \pm 0.006)$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 50\%$. The product branching ratio is $B(B^0 \rightarrow \bar{D}^0 \pi^+ \pi^-) B(\bar{D}^0 \rightarrow K^- \pi^+) = (0.39 \pm 0.26) \times 10^{-2}$.

$\Gamma(D^*(2010)^- \pi^+) / \Gamma_{total}$ Γ_9 / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0026 ± 0.0004 OUR AVERAGE				
0.0026 ± 0.0003 ± 0.0004	82	37 ALAM	94 CLE2	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0033 ± 0.0010 ± 0.0001		38 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$
0.00234 ± 0.00087 ± 0.00005	12	39 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.00234 ± 0.00149 ± 0.00005 - 0.00110	5	40 BEBEK	87 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.0027 ± 0.0014 ± 0.0010	5	41 ALBRECHT	87C ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.0035 ± 0.002 ± 0.002		42 ALBRECHT	86F ARG	$e^+ e^- \rightarrow \Upsilon(4S)$
0.017 ± 0.005 ± 0.005	41	43 GILES	84 CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

37 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0 \pi^+)$ and absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0) / B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-) / B(D^0 \rightarrow K^- \pi^+)$.

38 BORTOLETTO 92 reports $0.0040 \pm 0.0010 \pm 0.0007$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

39 ALBRECHT 90J reports $0.0028 \pm 0.0009 \pm 0.0006$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

40 BEBEK 87 reports $0.0028^{+0.0015+0.0010}_{-0.0012-0.0006}$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92 and ALBRECHT 90J.

41 ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

42 ALBRECHT 86F uses pseudomass that is independent of D^0 and D^+ branching ratios.

43 Assumes $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.60^{+0.08}_{-0.15}$. Assumes $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 0.40 \pm 0.02$ Does not depend on D branching ratios.

$\Gamma(D^- \pi^+ \pi^+ \pi^-) / \Gamma_{total}$ Γ_{10} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0080 ± 0.0021 ± 0.0014	44 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

44 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma((D^- \pi^+ \pi^+ \pi^-)_{nonresonant}) / \Gamma_{total}$ Γ_{11} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0039 ± 0.0014 ± 0.0013	45 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

45 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^- \pi^+ \rho^0) / \Gamma_{total}$ Γ_{12} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0011 ± 0.0009 ± 0.0004	46 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

46 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^- a_1(1260)^+) / \Gamma_{total}$ Γ_{13} / Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0060 ± 0.0022 ± 0.0024	47 BORTOLETTO92	CLEO	$e^+ e^- \rightarrow \Upsilon(4S)$

47 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

$\Gamma(D^*(2010)^- \pi^+ \pi^0) / \Gamma_{total}$ Γ_{14} / Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0151 ± 0.0051 ± 0.0003	51	48 ALBRECHT	90J ARG	$e^+ e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.015 ± 0.008 ± 0.008 8 49 ALBRECHT 87C ARG $e^+ e^- \rightarrow \Upsilon(4S)$

48 ALBRECHT 90J reports $0.018 \pm 0.004 \pm 0.005$ for $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

49 ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+ B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0 \bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

Meson Full Listings

 B^0 $\Gamma(D^*(2010)^-\rho^+)/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.0073±0.0015 OUR AVERAGE				
0.0074±0.0010±0.0014	76	50,51 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.0159±0.0113±0.0003		52 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.0059±0.0035±0.0001	19	53 ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
0.081 ± 0.029 $\begin{smallmatrix} +0.059 \\ -0.024 \end{smallmatrix}$	19	54 CHEN	85 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

50 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

51 This decay is nearly completely longitudinally polarized, $\Gamma_L/\Gamma = (93 \pm 5 \pm 5)\%$, as expected from the factorization hypothesis (ROSNER 90). The nonresonant $\pi^+\pi^0$ contribution under the ρ^+ is less than 9% at 90% CL.

52 BORTOLETTO 92 reports $0.019 \pm 0.008 \pm 0.011$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

53 ALBRECHT 90J reports $0.007 \pm 0.003 \pm 0.003$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

54 Uses $B(D^* \rightarrow D^0\pi^+) = 0.6 \pm 0.15$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 0.4$. Does not depend on D branching ratios.

 $\Gamma(D^*(2010)^-\pi^+\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.0119±0.0027 OUR AVERAGE					
0.0133±0.0036±0.0003			55 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.0100±0.0040±0.0002		26	56 ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••					
0.0063±0.0010±0.0011		49	57,58 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.033 ± 0.009 ± 0.016		27	59 ALBRECHT	87C ARG	$e^+e^- \rightarrow \Upsilon(4S)$
<0.042		90	60 BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

55 BORTOLETTO 92 reports $0.0159 \pm 0.0028 \pm 0.0037$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

56 ALBRECHT 90J reports $0.012 \pm 0.003 \pm 0.004$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

57 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2010)^+ \rightarrow D^0\pi^+)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

58 The three pion mass is required to be between 1.0 and 1.6 GeV consistent with an a_1 meson. (If this channel is dominated by a_1^+ , the branching ratio for $\bar{D}^*\pi^+\pi^+\pi^-$ is twice that for $\bar{D}^*\pi^-\pi^+\pi^-$.)

59 ALBRECHT 87C use PDG 86 branching ratios for D and $D^*(2010)$ and assume $B(\Upsilon(4S) \rightarrow B^+B^-) = 55\%$ and $B(\Upsilon(4S) \rightarrow B^0\bar{B}^0) = 45\%$. Superseded by ALBRECHT 90J.

60 BEBEK 87 value has been updated in BERKELMAN 91 to use same assumptions as noted for BORTOLETTO 92.

 $\Gamma((D^*(2010)^-\pi^+\pi^+\pi^-)_{\text{nonresonant}})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0000±0.0019±0.0016	61 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

61 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

 $\Gamma(D^*(2010)^-\pi^+\rho^0)/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0057±0.0031±0.0001	62 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

62 BORTOLETTO 92 reports $0.0068 \pm 0.0032 \pm 0.0021$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(D^*(2010)^-\pi_1(1260)^+)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.0151±0.0069±0.0003	63 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

63 BORTOLETTO 92 reports $0.018 \pm 0.006 \pm 0.006$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(D^*(2010)^-\pi^+\pi^+\pi^0)/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.034±0.018±0.001	28	64 ALBRECHT 90J	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

64 ALBRECHT 90J reports $0.041 \pm 0.015 \pm 0.016$ for $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 0.57 \pm 0.06$. We rescale to our best value $B(D^*(2010)^+ \rightarrow D^0\pi^+) = (68.1 \pm 1.3) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

 $\Gamma(D_2^*(2460)^-\pi^+)/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0022	90	65 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

65 ALAM 94 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and $B(D_2^*(2460)^\pm \rightarrow D^0\pi^+) = 30\%$.

 $\Gamma(D_2^*(2460)^-\rho^+)/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0049	90	66 ALAM	94 CLE2	$e^+e^- \rightarrow \Upsilon(4S)$

66 ALAM 94 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and $B(D_2^*(2460)^\pm \rightarrow D^0\pi^+) = 30\%$.

 $\Gamma(D^-D_s^+)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.008±0.004 OUR AVERAGE				
0.013±0.011±0.002		67 ALBRECHT 92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.007±0.004±0.001		68 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

••• We do not use the following data for averages, fits, limits, etc. •••

0.012±0.007
 3 | 69 BORTOLETTO90 | CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |

67 ALBRECHT 92G reports $0.017 \pm 0.013 \pm 0.006$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.7 \pm 1.0\%$.

68 BORTOLETTO 92 reports $0.0080 \pm 0.0045 \pm 0.0030$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.030 \pm 0.011$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

69 BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$. Superseded by BORTOLETTO 92.

 $\Gamma(D^*(2010)^-D_s^+)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.012±0.006 OUR AVERAGE				
0.011±0.008±0.001		70 ALBRECHT 92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.014±0.008±0.002		71 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

••• We do not use the following data for averages, fits, limits, etc. •••

0.024±0.014
 3 | 72 BORTOLETTO90 | CLEO | $e^+e^- \rightarrow \Upsilon(4S)$ |

70 ALBRECHT 92G reports $0.014 \pm 0.010 \pm 0.003$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^-\pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \rightarrow D^0\pi^+) = 55 \pm 4\%$.

71 BORTOLETTO 92 reports $0.016 \pm 0.009 \pm 0.006$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.030 \pm 0.011$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D and $D^*(2010)$.

72 BORTOLETTO 90 assume $B(D_s \rightarrow \phi\pi^+) = 2\%$. Superseded by BORTOLETTO 92.

 $\Gamma(D^-D_s^{*+})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.021±0.015±0.002	73 ALBRECHT 92G	ARG	$e^+e^- \rightarrow \Upsilon(4S)$

73 ALBRECHT 92G reports $0.027 \pm 0.017 \pm 0.009$ for $B(D_s^+ \rightarrow \phi\pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ branching ratios, e.g., $B(D^+ \rightarrow K^-\pi^+\pi^+) = 7.7 \pm 1.0\%$.

 $[\Gamma(D^*(2010)^-D_s^+) + \Gamma(D^*(2010)^-D_s^{*+})]/\Gamma_{\text{total}}$ $(\Gamma_{24} + \Gamma_{25})/\Gamma$

VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT
4.3±1.1±0.5	22	74 BORTOLETTO90	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

74 BORTOLETTO 90 reports 7.5 ± 2.0 for $B(D_s^+ \rightarrow \phi\pi^+) = 0.02$. We rescale to our best value $B(D_s^+ \rightarrow \phi\pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value.

$\Gamma(D^{*0}(2010)^- D_s^{*+})/\Gamma_{total}$ Γ_{26}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$0.020 \pm 0.012 \pm 0.002$		75 ALBRECHT 92G ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>75 ALBRECHT 92G reports $0.026 \pm 0.014 \pm 0.006$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = (3.5 \pm 0.4) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes PDG 1990 D^+ and $D^*(2010)^+$ branching ratios, e.g., $B(D^0 \rightarrow K^- \pi^+) = 3.71 \pm 0.25\%$, $B(D^+ \rightarrow K^- \pi^+ \pi^+) = 7.1 \pm 1.0\%$, and $B(D^*(2010)^+ \rightarrow D^0 \pi^+) = 55 \pm 4\%$.</p>				

 $\Gamma(D_s^+ \pi^-)/\Gamma_{total}$ Γ_{27}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00029	90	76 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<0.0013	90	77 BORTOLETTO90 CLEO		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>76 ALEXANDER 93B reports $< 2.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				
<p>77 BORTOLETTO 90 assume $B(D_s \rightarrow \phi \pi^+) = 2\%$.</p>				

 $\Gamma(D_s^{*+} \pi^-)/\Gamma_{total}$ Γ_{28}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0005	90	78 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>78 ALEXANDER 93B reports $< 4.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $[\Gamma(D_s^+ \pi^-) + \Gamma(D_s^- K^+)]/\Gamma_{total}$ $(\Gamma_{27} + \Gamma_{33})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0013	90	79 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>79 ALBRECHT 93E reports $< 1.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $[\Gamma(D_s^{*+} \pi^-) + \Gamma(D_s^{*-} K^+)]/\Gamma_{total}$ $(\Gamma_{28} + \Gamma_{34})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0009	90	80 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>80 ALBRECHT 93E reports $< 1.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^+ \rho^-)/\Gamma_{total}$ Γ_{29}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0007	90	81 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<0.0017	90	82 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>81 ALEXANDER 93B reports $< 6.6 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				
<p>82 ALBRECHT 93E reports $< 2.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^{*+} \rho^-)/\Gamma_{total}$ Γ_{30}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0008	90	83 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<0.0019	90	84 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>83 ALEXANDER 93B reports $< 7.4 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				
<p>84 ALBRECHT 93E reports $< 2.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^+ a_1(1260)^-)/\Gamma_{total}$ Γ_{31}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0027	90	85 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>85 ALBRECHT 93E reports $< 3.5 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^{*+} a_1(1260)^-)/\Gamma_{total}$ Γ_{32}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0022	90	86 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>86 ALBRECHT 93E reports $< 2.9 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^- K^+)/\Gamma_{total}$ Γ_{33}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00024	90	87 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<0.0013	90	88 BORTOLETTO90 CLEO		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>87 ALEXANDER 93B reports $< 2.3 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				
<p>88 BORTOLETTO 90 assume $B(D_s \rightarrow \phi \pi^+) = 2\%$.</p>				

 $\Gamma(D_s^{*-} K^+)/\Gamma_{total}$ Γ_{34}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00018	90	89 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>89 ALEXANDER 93B reports $< 1.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^- K^*(892)^+)/\Gamma_{total}$ Γ_{35}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0010	90	90 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<0.004	90	91 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>90 ALEXANDER 93B reports $< 9.7 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				
<p>91 ALBRECHT 93E reports $< 4.6 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^{*-} K^*(892)^+)/\Gamma_{total}$ Γ_{36}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0012	90	92 ALEXANDER 93B CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>• • • We do not use the following data for averages, fits, limits, etc. • • •</p>				
<0.004	90	93 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>92 ALEXANDER 93B reports $< 11.0 \times 10^{-4}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.037$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				
<p>93 ALBRECHT 93E reports $< 5.8 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^- \pi^+ K^0)/\Gamma_{total}$ Γ_{37}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.006	90	94 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>94 ALBRECHT 93E reports $< 7.3 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^{*-} \pi^+ K^0)/\Gamma_{total}$ Γ_{38}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0032	90	95 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>95 ALBRECHT 93E reports $< 4.2 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^- \pi^+ K^*(892)^0)/\Gamma_{total}$ Γ_{39}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.004	90	96 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>96 ALBRECHT 93E reports $< 5.0 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D_s^{*-} \pi^+ K^*(892)^0)/\Gamma_{total}$ Γ_{40}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0021	90	97 ALBRECHT 93E ARG		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>97 ALBRECHT 93E reports $< 2.7 \times 10^{-3}$ for $B(D_s^+ \rightarrow \phi \pi^+) = 0.027$. We rescale to our best value $B(D_s^+ \rightarrow \phi \pi^+) = 0.035$.</p>				

 $\Gamma(D^0 \pi^0)/\Gamma_{total}$ Γ_{41}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00048	90	98 ALAM 94 CLE2		$e^+ e^- \rightarrow \Upsilon(4S)$
<p>98 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^- \pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^- \pi^+ \pi^0)/B(D^0 \rightarrow K^- \pi^+)$ and $B(D^0 \rightarrow K^- \pi^+ \pi^+ \pi^-)/B(D^0 \rightarrow K^- \pi^+)$.</p>				

Meson Full Listings

 B^0 $\Gamma(D^0 \rho^0)/\Gamma_{total}$ Γ_{42}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.00055	90		99 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0006	90		100 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<0.0027	90	4	101 ALBRECHT	88K ARG	$e^+e^- \rightarrow \Upsilon(4S)$

99 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

100 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$ and uses Mark III branching fractions for the D .

101 ALBRECHT 88K reports < 0.003 assuming $B^0 \bar{B}^0 : B^+ B^-$ production ratio is 45:55. We rescale to 50%.

 $\Gamma(D^0 \eta)/\Gamma_{total}$ Γ_{43}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00068	90	102 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

102 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^0 \eta')/\Gamma_{total}$ Γ_{44}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00086	90	103 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

103 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^0 \omega)/\Gamma_{total}$ Γ_{45}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00063	90	104 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

104 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^*(2007)^0 \pi^0)/\Gamma_{total}$ Γ_{46}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00097	90	105 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

105 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^*(2007)^0 \rho^0)/\Gamma_{total}$ Γ_{47}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00117	90	106 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

106 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^*(2007)^0 \eta)/\Gamma_{total}$ Γ_{48}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.00069	90	107 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

107 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^*(2007)^0 \eta')/\Gamma_{total}$ Γ_{49}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0027	90	108 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

108 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(D^*(2007)^0 \omega)/\Gamma_{total}$ Γ_{50}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0021	90	109 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

109 ALAM 94 assume equal production of B^+ and B^0 at the $\Upsilon(4S)$ and use the CLEO II $B(D^*(2007)^0 \rightarrow D^0 \pi^0)$ and absolute $B(D^0 \rightarrow K^-\pi^+)$ and the PDG 1992 $B(D^0 \rightarrow K^-\pi^+\pi^0)/B(D^0 \rightarrow K^-\pi^+)$ and $B(D^0 \rightarrow K^-\pi^+\pi^+\pi^-)/B(D^0 \rightarrow K^-\pi^+)$.

 $\Gamma(J/\psi(1S) K^0)/\Gamma_{total}$ Γ_{51}/Γ

VALUE (units 10^{-4})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
7.5 ± 2.1 OUR AVERAGE					
7.5 ± 2.4 ± 0.8		10	110 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
6.9 ± 4.1 ± 0.3			110 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
9.2 ± 7.2 ± 0.4		2	111 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<50	90		ALAM	86	CLEO $e^+e^- \rightarrow \Upsilon(4S)$
-----	----	--	------	----	--

110 BORTOLETTO 92 reports $6 \pm 3 \pm 2$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

111 ALBRECHT 90J reports $8 \pm 6 \pm 2$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(J/\psi(1S) K^+ \pi^-)/\Gamma_{total}$ Γ_{52}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.00115 ± 0.00056 ± 0.00005			112 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0013	90		113 ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	--	--------------	---------	-----------------------------------

<0.0063	90	2	GILES	84 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	---	-------	---------	-----------------------------------

112 BORTOLETTO 92 reports $0.0010 \pm 0.0004 \pm 0.0003$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

113 ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. $K\pi$ system is specifically selected as nonresonant.

 $\Gamma(J/\psi(1S) K^*(892)^0)/\Gamma_{total}$ Γ_{53}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.00158 ± 0.00028 OUR AVERAGE				

0.00169 ± 0.00031 ± 0.00018	29	114 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
-----------------------------	----	----------	----	--

0.0013 ± 0.0007 ± 0.0001		115 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------------	--	------------------	------	-----------------------------------

0.0013 ± 0.0006 ± 0.0001	6	116 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------------	---	--------------	---------	-----------------------------------

0.0040 ± 0.0018 ± 0.0002	5	117 BEBEK	87 CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
--------------------------	---	-----------	---------	-----------------------------------

0.0040 ± 0.0030		118 ALBAJAR	91E UA1	$E_{cm}^{D^0} = 630$ GeV
-----------------	--	-------------	---------	--------------------------

0.0033 ± 0.0018	5	119 ALBRECHT	87D ARG	$e^+e^- \rightarrow \Upsilon(4S)$
-----------------	---	--------------	---------	-----------------------------------

0.0041 ± 0.0018	5	120 ALAM	86 CLEO	Repl. by BEBEK 87
-----------------	---	----------	---------	-------------------

114 The neutral and charged B events together are predominantly longitudinally polarized, $\Gamma_L/\Gamma = 0.080 \pm 0.08 \pm 0.05$. This can be compared with a prediction using HQET, 0.73 (KRAMER 92). This polarization indicates that the $B \rightarrow \psi K^*$ decay is dominated by the $CP = -1$ CP eigenstate. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

115 BORTOLETTO 92 reports $0.0011 \pm 0.0005 \pm 0.0003$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

116 ALBRECHT 90J reports $0.0011 \pm 0.0005 \pm 0.0002$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

117 BEBEK 87 reports $0.0035 \pm 0.0016 \pm 0.0003$ for $B(J/\psi(1S) \rightarrow e^+e^-) = 0.069 \pm 0.009$. We rescale to our best value $B(J/\psi(1S) \rightarrow e^+e^-) = (5.99 \pm 0.25) \times 10^{-2}$. Our first error is their experiment's error and our second error is the systematic error from using our best value. Updated in BORTOLETTO 92 to use the same assumptions.

118 ALBAJAR 91E assumes B^0 production fraction of 36%.

119 ALBRECHT 87D assume $B^+ B^- / B^0 \bar{B}^0$ ratio is 55/45. Superseded by ALBRECHT 90J.

120 ALAM 86 assumes B^\pm / B^0 ratio is 60/40. The observation of the decay $B^+ \rightarrow J/\psi K^*(892)^+$ (HAAS 85) has been retracted in this paper.

 $\Gamma(\psi(2S) K^0)/\Gamma_{total}$ Γ_{54}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0008	90	121 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0015	90	121 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	------------------	------	-----------------------------------

<0.0028	90	121 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	---------	-----------------------------------

121 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\psi(2S) K^+ \pi^-)/\Gamma_{total}$ Γ_{55}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.001	90	122 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$

122 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\psi(2S) K^*(892)^0)/\Gamma_{total}$ Γ_{56}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.0014 ± 0.0008 ± 0.0004		123 BORTOLETTO92	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.0019	90	123 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$
---------	----	----------	----	--

<0.0023	90	123 ALBRECHT	90J ARG	$e^+e^- \rightarrow \Upsilon(4S)$
---------	----	--------------	---------	-----------------------------------

123 Assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

 $\Gamma(\chi_{c1}(1P) K^0)/\Gamma_{total}$ Γ_{57}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0027	90	124 ALAM	94	CLE2 $e^+e^- \rightarrow \Upsilon(4S)$

124 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $\Upsilon(4S)$.

See key on page 1343

Meson Full Listings

 B^0 $\Gamma(\chi_{c1}(1P)K^*(892)^0)/\Gamma_{total}$ Γ_{58}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6 \times 10^{-5}$	90	125 ALAM	94 CLE2	$e^+e^- \rightarrow T(4S)$

125 BORTOLETTO 92 assumes equal production of B^+ and B^0 at the $T(4S)$.

 $\Gamma(K^+\pi^-)/\Gamma_{total}$ Γ_{59}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6 \times 10^{-5}$	90	126 BATTLE	93 CLE2	$e^+e^- \rightarrow T(4S)$
$<1.8 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$
$<9 \times 10^{-5}$	90	127 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
$<3.2 \times 10^{-4}$	90	AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

126 BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.
127 Assumes the $T(4S)$ decays 43% to $B^0\bar{B}^0$.

 $[\Gamma(K^+\pi^-) + \Gamma(\pi^+\pi^-)]/\Gamma_{total}$ $(\Gamma_{59} + \Gamma_{85})/\Gamma$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$(2.4^{+0.8}_{-0.7} \pm 0.2) \times 10^{-5}$		128 BATTLE	93 CLE2	$e^+e^- \rightarrow T(4S)$

128 BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

 $\Gamma(K^+K^-)/\Gamma_{total}$ Γ_{60}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<0.7 \times 10^{-5}$	90	129 BATTLE	93 CLE2	$e^+e^- \rightarrow T(4S)$

129 BATTLE 93 assumes equal production of $B^0\bar{B}^0$ and B^+B^- at $T(4S)$.

 $\Gamma(K^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{61}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.4 \times 10^{-4}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^0\rho^0)/\Gamma_{total}$ Γ_{62}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$
$<5.0 \times 10^{-4}$	90	130 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
<0.064	90	131 AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

130 AVERY 89B reports $< 5.8 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
131 AVERY 87 reports < 0.08 assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^0 f_0(980))/\Gamma_{total}$ Γ_{63}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6 \times 10^{-4}$	90	132 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$

132 AVERY 89B reports $< 4.2 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^*(892)^+\pi^-)/\Gamma_{total}$ Γ_{64}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.8 \times 10^{-4}$	90	133 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
$<6.2 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$
$<5.6 \times 10^{-4}$	90	134 AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

133 AVERY 89B reports $< 4.4 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
134 AVERY 87 reports $< 7 \times 10^{-4}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_2^*(1430)^+\pi^-)/\Gamma_{total}$ Γ_{65}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.6 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^0K^+K^-)/\Gamma_{total}$ Γ_{66}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.3 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^0\phi)/\Gamma_{total}$ Γ_{67}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.2 \times 10^{-4}$	90	135 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
$<7.2 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$
$<1.0 \times 10^{-3}$	90	136 AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

135 AVERY 89B reports $< 4.9 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
136 AVERY 87 reports $< 1.3 \times 10^{-3}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^*(892)^0\pi^+\pi^-)/\Gamma_{total}$ Γ_{68}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-3}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^*(892)^0\rho^0)/\Gamma_{total}$ Γ_{69}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.6 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$
$<5.8 \times 10^{-4}$	90	137 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
$<9.6 \times 10^{-4}$	90	138 AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •
137 AVERY 89B reports $< 6.7 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
138 AVERY 87 reports $< 1.2 \times 10^{-3}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K^*(892)^0 f_0(980))/\Gamma_{total}$ Γ_{70}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.7 \times 10^{-4}$	90	139 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$

139 AVERY 89B reports $< 2.0 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_1(1400)^+\pi^-)/\Gamma_{total}$ Γ_{71}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^*(892)^0K^+K^-)/\Gamma_{total}$ Γ_{72}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<6.1 \times 10^{-4}$	90	ALBRECHT	91E ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^*(892)^0\phi)/\Gamma_{total}$ Γ_{73}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.2 \times 10^{-4}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$
$<3.8 \times 10^{-4}$	90	140 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
$<3.8 \times 10^{-4}$	90	141 AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

140 AVERY 89B reports $< 4.4 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.
141 AVERY 87 reports $< 4.7 \times 10^{-4}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_1(1400)^0\rho^0)/\Gamma_{total}$ Γ_{74}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K_1(1400)^0\phi)/\Gamma_{total}$ Γ_{75}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.0 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K_2^*(1430)^0\rho^0)/\Gamma_{total}$ Γ_{76}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K_2^*(1430)^0\phi)/\Gamma_{total}$ Γ_{77}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.4 \times 10^{-3}$	90	ALBRECHT	91B ARG	$e^+e^- \rightarrow T(4S)$

 $\Gamma(K^*(892)^0\gamma)/\Gamma_{total}$ Γ_{78}/Γ

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$(4.0 \pm 1.7 \pm 0.8) \times 10^{-5}$		8	142 AMMAR	93 CLE2	$e^+e^- \rightarrow T(4S)$
$< 4.2 \times 10^{-4}$	90		ALBRECHT	89G ARG	$e^+e^- \rightarrow T(4S)$
$< 2.4 \times 10^{-4}$	90		143 AVERY	89B CLEO	$e^+e^- \rightarrow T(4S)$
$< 2.1 \times 10^{-3}$	90		AVERY	87 CLEO	$e^+e^- \rightarrow T(4S)$

• • • We do not use the following data for averages, fits, limits, etc. • • •
142 AMMAR 93 observed 6.6 ± 2.8 events above background.
143 AVERY 89B reports $< 2.8 \times 10^{-4}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_1(1270)^0\gamma)/\Gamma_{total}$ Γ_{79}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0070	90	144 ALBRECHT	89G ARG	$e^+e^- \rightarrow T(4S)$

144 ALBRECHT 89G reports < 0.0078 assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_1(1400)^0\gamma)/\Gamma_{total}$ Γ_{80}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0043	90	145 ALBRECHT	89G ARG	$e^+e^- \rightarrow T(4S)$

145 ALBRECHT 89G reports < 0.0048 assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

 $\Gamma(K_2^*(1430)^0\gamma)/\Gamma_{total}$ Γ_{81}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<4.0 \times 10^{-4}$	90	146 ALBRECHT	89G ARG	$e^+e^- \rightarrow T(4S)$

146 ALBRECHT 89G reports $< 4.4 \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

Meson Full Listings

 B^0

$\Gamma(K^*(1680)^0 \gamma)/\Gamma_{total}$					Γ_{82}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0020	90	147 ALBRECHT 89G ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

147 ALBRECHT 89G reports < 0.0022 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K_3^*(1780)^0 \gamma)/\Gamma_{total}$					Γ_{83}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.010	90	148 ALBRECHT 89G ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

148 ALBRECHT 89G reports < 0.011 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(K_4^*(2045)^0 \gamma)/\Gamma_{total}$					Γ_{84}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<0.0043	90	149 ALBRECHT 89G ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

149 ALBRECHT 89G reports < 0.0048 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ \pi^-)/\Gamma_{total}$					Γ_{85}/Γ
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.9 x 10 ⁻⁵	90	150	BATTLE 93	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.3 x 10 ⁻⁴	90	150	ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$
<7.7 x 10 ⁻⁵	90	151	BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<2.6 x 10 ⁻⁴	90	151	BEBEK 87	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
<5 x 10 ⁻⁴	90	4	GILES 84	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

150 Assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

151 Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{total}$					Γ_{86}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<7.2 x 10 ⁻⁴	90	152 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

152 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^0 \pi^0)/\Gamma_{total}$					Γ_{87}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<4.0 x 10 ⁻⁴	90	153 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

153 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^\mp \pi^\pm)/\Gamma_{total}$					Γ_{88}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<5.2 x 10 ⁻⁴	90	154 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<5.2 x 10⁻³ 90 155 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

154 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

155 BEBEK 87 reports < 6.1 x 10⁻³ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ \pi^- \pi^+ \pi^-)/\Gamma_{total}$					Γ_{89}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<6.7 x 10 ⁻⁴	90	156 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

156 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^0 \rho^0)/\Gamma_{total}$					Γ_{90}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.8 x 10 ⁻⁴	90	157 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<2.9 x 10⁻⁴ 90 158 BORTOLETTO89 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

<4.3 x 10⁻⁴ 90 158 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

157 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

158 Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(a_1(1260)^\mp \pi^\pm)/\Gamma_{total}$					Γ_{91}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<4.9 x 10 ⁻⁴	90	159 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.3 x 10⁻⁴ 90 160 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$

<1.0 x 10⁻³ 90 159 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

159 Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

160 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(a_2(1320)^\mp \pi^\pm)/\Gamma_{total}$					Γ_{92}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<3.0 x 10 ⁻⁴	90	161 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.4 x 10⁻³ 90 161 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

161 Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\pi^+ \pi^- \pi^0 \pi^0)/\Gamma_{total}$					Γ_{93}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<3.1 x 10 ⁻³	90	162 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

162 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho^+ \rho^-)/\Gamma_{total}$					Γ_{94}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.2 x 10 ⁻³	90	163 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

163 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^0 \pi^0)/\Gamma_{total}$					Γ_{95}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.1 x 10 ⁻³	90	164 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

164 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\omega \pi^0)/\Gamma_{total}$					Γ_{96}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<4.6 x 10 ⁻⁴	90	165 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

165 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\eta \pi^0)/\Gamma_{total}$					Γ_{97}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.8 x 10 ⁻³	90	166 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

166 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{total}$					Γ_{98}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<9.0 x 10 ⁻³	90	167 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

167 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+ \rho^-)/\Gamma_{total}$					Γ_{99}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<3.4 x 10 ⁻³	90	168 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

168 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^0 \rho^0)/\Gamma_{total}$					Γ_{100}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.4 x 10 ⁻³	90	169 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

169 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^-)/\Gamma_{total}$					Γ_{101}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<3.0 x 10 ⁻³	90	170 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

170 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(a_1(1260)^+ a_1(1260)^-)/\Gamma_{total}$					Γ_{102}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<2.8 x 10 ⁻³	90	171 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<6.0 x 10⁻³ 90 172 ALBRECHT 90B ARG $e^+e^- \rightarrow \Upsilon(4S)$

171 BORTOLETTO 89 reports < 3.2 x 10⁻³ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

172 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\pi^+ \pi^+ \pi^+ \pi^- \pi^- \pi^0)/\Gamma_{total}$					Γ_{103}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<1.1 x 10 ⁻²	90	173 ALBRECHT 90B ARG		$e^+e^- \rightarrow \Upsilon(4S)$	

173 ALBRECHT 90B limit assumes equal production of $B^0 \bar{B}^0$ and $B^+ B^-$ at $\Upsilon(4S)$.

$\Gamma(\rho \bar{\rho})/\Gamma_{total}$					Γ_{104}/Γ
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
<3.4 x 10 ⁻⁵	90	174 BORTOLETTO89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.2 x 10⁻⁴ 90 175 ALBRECHT 88F ARG $e^+e^- \rightarrow \Upsilon(4S)$

<1.7 x 10⁻⁴ 90 174 BEBEK 87 CLEO $e^+e^- \rightarrow \Upsilon(4S)$

174 Paper assumes the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

175 ALBRECHT 88F reports < 1.3 x 10⁻⁴ assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(\rho \bar{\rho} \pi^+ \pi^-)/\Gamma_{total}$					Γ_{105}/Γ
VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	
<2.5	90	176 BEBEK 89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.4 ± 1.8 ± 2.0 177 ALBRECHT 88F ARG $e^+e^- \rightarrow \Upsilon(4S)$

176 BEBEK 89 reports < 2.9 x 10⁻⁴ assuming the $\Upsilon(4S)$ decays 43% to $B^0 \bar{B}^0$. We rescale to 50%.

177 ALBRECHT 88F reports 6.0 ± 2.0 ± 2.2 assuming the $\Upsilon(4S)$ decays 45% to $B^0 \bar{B}^0$. We rescale to 50%.

$\Gamma(p\bar{\lambda}\pi^-)/\Gamma_{\text{total}}$ Γ_{106}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.8 \times 10^{-4}$	90	178 ALBRECHT 88F ARG		$e^+e^- \rightarrow T(4S)$
178 ALBRECHT 88F reports $<2.0 \times 10^{-4}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(\Delta^0\Delta^0)/\Gamma_{\text{total}}$ Γ_{107}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0015	90	179 BORTOLETTO89 CLEO		$e^+e^- \rightarrow T(4S)$
179 BORTOLETTO 89 reports <0.0018 assuming $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(\Delta^{++}\Delta^{--})/\Gamma_{\text{total}}$ Γ_{108}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<1.1 \times 10^{-4}$	90	180 BORTOLETTO89 CLEO		$e^+e^- \rightarrow T(4S)$
180 BORTOLETTO 89 reports $<1.3 \times 10^{-4}$ assuming $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(\Sigma_c^{--}\Delta^{++})/\Gamma_{\text{total}}$ Γ_{109}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.0012	90	181 PROCARIO 94 CLE2		$e^+e^- \rightarrow T(4S)$
181 PROCARIO 94 reports <0.0012 for $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.043$. We rescale to our best value $B(\Lambda_c^+ \rightarrow pK^-\pi^+) = 0.044$.				

$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_{110}/Γ
Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-6}$	90	AMMAR 94 CLE2		$e^+e^- \rightarrow T(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<2.6 \times 10^{-5}$	90	182 AVERY 89B CLEO		$e^+e^- \rightarrow T(4S)$
$<7.6 \times 10^{-5}$	90	183 ALBRECHT 87D ARG		$e^+e^- \rightarrow T(4S)$
$<6.4 \times 10^{-5}$	90	184 AVERY 87 CLEO		$e^+e^- \rightarrow T(4S)$
$<3 \times 10^{-4}$	90	GILES 84 CLEO		Repl. by AVERY 87
182 AVERY 89B reports $<3 \times 10^{-5}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.				
183 ALBRECHT 87D reports $<8.5 \times 10^{-5}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.				
184 AVERY 87 reports $<8 \times 10^{-5}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{111}/Γ
Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-6}$	90	AMMAR 94 CLE2		$e^+e^- \rightarrow T(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<8.3 \times 10^{-6}$	90	185 ALBAJAR 91C UA1		$E_{\text{cm}}^{\text{PD}} = 630$ GeV
$<1.2 \times 10^{-5}$	90	186 ALBAJAR 91C UA1		$E_{\text{cm}}^{\text{PD}} = 630$ GeV
$<4.3 \times 10^{-5}$	90	187 AVERY 89B CLEO		$e^+e^- \rightarrow T(4S)$
$<4.5 \times 10^{-5}$	90	188 ALBRECHT 87D ARG		$e^+e^- \rightarrow T(4S)$
$<7.7 \times 10^{-5}$	90	189 AVERY 87 CLEO		$e^+e^- \rightarrow T(4S)$
$<2 \times 10^{-4}$	90	GILES 84 CLEO		Repl. by AVERY 87

185 B^0 and B_s^0 are not separated.

186 Obtained from unseparated B^0 and B_s^0 measurement by assuming a $B^0:B_s^0$ ratio 2:1.

187 AVERY 89B reports $<5 \times 10^{-3}$ assuming the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.

188 ALBRECHT 87D reports $<5 \times 10^{-5}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.

189 AVERY 87 reports $<9 \times 10^{-5}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.

$\Gamma(K^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{112}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.0 \times 10^{-4}$	90	ALBRECHT 91E ARG		$e^+e^- \rightarrow T(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<5.2 \times 10^{-4}$	90	190 AVERY 87 CLEO		$e^+e^- \rightarrow T(4S)$
190 AVERY 87 reports $<6.5 \times 10^{-4}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(K^0\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{113}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<3.6 \times 10^{-4}$	90	191 AVERY 87 CLEO		$e^+e^- \rightarrow T(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<5.2 \times 10^{-4}$	90	ALBRECHT 91E ARG		$e^+e^- \rightarrow T(4S)$
191 AVERY 87 reports $<4.5 \times 10^{-4}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(K^*(892)^0e^+e^-)/\Gamma_{\text{total}}$ Γ_{114}/Γ

Test for $\Delta B = 1$ weak neutral current. Allowed by higher-order electroweak interactions.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.9 \times 10^{-4}$	90	ALBRECHT 91E ARG		$e^+e^- \rightarrow T(4S)$

$\Gamma(K^*(892)^0\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_{115}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<2.3 \times 10^{-5}$	90	192 ALBAJAR 91C UA1		$E_{\text{cm}}^{\text{PD}} = 630$ GeV
••• We do not use the following data for averages, fits, limits, etc. •••				
$<3.4 \times 10^{-4}$	90	ALBRECHT 91E ARG		$e^+e^- \rightarrow T(4S)$
192 ALBAJAR 91C assumes 36% of \bar{b} quarks give B^0 mesons.				

$\Gamma(e^\pm\tau^\mp)/\Gamma_{\text{total}}$ Γ_{116}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.9 \times 10^{-6}$	90	AMMAR 94 CLE2		$e^+e^- \rightarrow T(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••				
$<3.4 \times 10^{-5}$	90	193 AVERY 89B CLEO		$e^+e^- \rightarrow T(4S)$
$<4.5 \times 10^{-5}$	90	194 ALBRECHT 87D ARG		$e^+e^- \rightarrow T(4S)$
$<7.7 \times 10^{-5}$	90	195 AVERY 87 CLEO		$e^+e^- \rightarrow T(4S)$
$<3 \times 10^{-4}$	90	GILES 84 CLEO		Repl. by AVERY 87
193 Paper assumes the $T(4S)$ decays 43% to $B^0\bar{B}^0$. We rescale to 50%.				
194 ALBRECHT 87D reports $<5 \times 10^{-5}$ assuming the $T(4S)$ decays 45% to $B^0\bar{B}^0$. We rescale to 50%.				
195 AVERY 87 reports $<9 \times 10^{-5}$ assuming the $T(4S)$ decays 40% to $B^0\bar{B}^0$. We rescale to 50%.				

$\Gamma(e^\pm\tau^\mp)/\Gamma_{\text{total}}$ Γ_{117}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<5.3 \times 10^{-4}$	90	AMMAR 94 CLE2		$e^+e^- \rightarrow T(4S)$

$\Gamma(\mu^\pm\tau^\mp)/\Gamma_{\text{total}}$ Γ_{118}/Γ
Test of lepton family number conservation.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
$<8.3 \times 10^{-4}$	90	AMMAR 94 CLE2		$e^+e^- \rightarrow T(4S)$

Meson Full Listings

 B^0 B^0 - \bar{B}^0 MIXING

For a discussion of B^0 - \bar{B}^0 mixing see the note on " B^0 - \bar{B}^0 Mixing and CP Violation in B Decay" below.

 χ_d

This B^0 - \bar{B}^0 mixing parameter measures the probability (integrated over time) that a produced B^0 (or \bar{B}^0) decays as a \bar{B}^0 (or B^0), e.g. for inclusive lepton decays

$$\chi_d = \frac{\Gamma(B^0 \rightarrow \ell^- X \text{ (via } \bar{B}^0))}{\Gamma(B^0 \rightarrow \ell^\pm X)} = \frac{\Gamma(\bar{B}^0 \rightarrow \ell^+ X \text{ (via } B^0))}{\Gamma(\bar{B}^0 \rightarrow \ell^\pm X)}$$

Where experiments have measured the parameter $r = \chi/(1-\chi)$, we have converted to χ . Mixing violates the $\Delta B \neq 2$ rule.

Note that the measurement of χ at energies higher than the $\Upsilon(4S)$ have not separated χ_d from χ_s where the subscripts indicate $B^0(\bar{b}d)$ or $B_s^0(\bar{b}s)$. They are listed in the B_s^0 - \bar{B}_s^0 MIXING section.

The experiments at $\Upsilon(4S)$ make an assumption about the B^0 - \bar{B}^0 fraction and about the ratio of the B^\pm and B^0 semileptonic branching ratios (usually that it equals one).

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.156 ± 0.024 OUR AVERAGE				
0.16 ± 0.04 ± 0.04		196 ALBRECHT 94	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.149 ± 0.023 ± 0.022		197 BARTELT 93	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.171 ± 0.048		198 ALBRECHT 92L	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24 ± 0.12		199 ELSEN 90	JADE	$e^+e^- \rightarrow 35\text{-}44$ GeV
0.158 ^{+0.052} _{-0.059}		ARTUSO 89	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
0.17 ± 0.05		200 ALBRECHT 87I	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
< 0.19		90 201 BEAN 87B	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$
< 0.27		90 202 AVERY 84	CLEO	$e^+e^- \rightarrow \Upsilon(4S)$

196 ALBRECHT 94 reports $r=0.194 \pm 0.062 \pm 0.054$. We convert to χ for comparison. Uses tagged events (lepton + pion from D^*).

197 BARTELT 93 analysis performed using tagged events (lepton+pion from D^*). Using dilepton events they obtain $0.157 \pm 0.016^{+0.033}_{-0.028}$.

198 ALBRECHT 92L is a combined measurement employing several lepton-based techniques. It uses all previous ARGUS data in addition to new data and therefore supersedes ALBRECHT 87I. A value of $r = 20.6 \pm 7.0\%$ is directly measured. The value can be used to measure $\chi = \Delta M/\Gamma = 0.72 \pm 0.15$ for the B_d meson. Assumes $f_{+,-}/f_0 = 1.0 \pm 0.05$ and uses $\tau_{B^\pm}/\tau_{B^0} = (0.95 \pm 0.14)(f_{+,-}/f_0)$.

199 These experiments see a combination of B_s and B_d mesons.

200 ALBRECHT 87I is inclusive measurement with like-sign dileptons, with tagged B decays plus leptons, and one fully reconstructed event. Measurements $r=0.21 \pm 0.08$. We convert to χ for comparison. Superseded by ALBRECHT 92L.

201 BEAN 87B measured $r < 0.24$; we converted to χ .

202 Same-sign dilepton events. Limit assumes semileptonic BR for B^+ and B^0 equal. If B^0/B^\pm ratio < 0.58 , no limit exists. The limit was corrected in BEAN 87B from $r < 0.30$ to $r < 0.37$. We converted this limit to χ .

$$\Delta m_{B^0} = m_{B_H^0} - m_{B_L^0}$$

Δm_{B^0} is a measure of the B^0 - \bar{B}^0 oscillation frequency in time-dependent mixing experiments.

VALUE (10^{12} h s^{-1})	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.06 OUR AVERAGE			
0.508 ± 0.075 ± 0.025	203 AKERS 94C	OPAL	$e^+e^- \rightarrow Z$
0.50 ^{+0.07} _{-0.06} ± 0.11 _{-0.10}	204 BUSKULIC 94B	ALEP	$e^+e^- \rightarrow Z$
0.52 ^{+0.10} _{-0.11} ± 0.04 _{-0.03}	205 BUSKULIC 93K	ALEP	$e^+e^- \rightarrow Z$

203 AKERS 94C observes the time dependence of B^0 - \bar{B}^0 mixing using $D^{*\pm} \ell^\mp$ events and jet charge.

204 BUSKULIC 94B observes the time dependence of B^0 - \bar{B}^0 mixing using dileptons.

205 BUSKULIC 93K observes the time dependence of B^0 - \bar{B}^0 mixing using $D^{*\pm}$ lepton correlations.

$$\chi_d = \Delta m_{B^0}/\Gamma_{B^0}$$

This section combines results from the previous two sections.

Time integrated mixing measurements of χ determine this quantity directly via

$$\frac{\Delta m_{B^0}}{\Gamma_{B^0}} = \left(\frac{\chi}{0.5-\chi} \right)^{1/2}$$

while time-dependent mixing measurements determine $\Delta m_{B^0} = m_{B_H^0} - m_{B_L^0}$ which are combined with τ_{B^0} to give

$$\frac{\Delta m_{B^0}}{\Gamma_{B^0}} = \frac{(m_{B_H^0} - m_{B_L^0}) \tau_{B^0}}{\hbar}$$

The averaging takes into account the common systematic errors on the LEP experiments due to τ_{B^0} .

VALUE	DOCUMENT ID	TECN	COMMENT
0.71 ± 0.06 OUR AVERAGE			
0.76 ± 0.11 ± 0.06	206 AKERS 94C	OPAL	$e^+e^- \rightarrow Z$
0.69 ± 0.18	207 ALBRECHT 94	ARG	$e^+e^- \rightarrow \Upsilon(4S)$
0.75 ^{+0.20} _{-0.18} ± 0.06	206 BUSKULIC 94B	ALEP	$e^+e^- \rightarrow Z$
0.65 ± 0.10	207 BARTELT 93	CLE2	$e^+e^- \rightarrow \Upsilon(4S)$
0.78 ^{+0.16} _{-0.17} ± 0.06	206 BUSKULIC 93K	ALEP	$e^+e^- \rightarrow Z$

0.72 ± 0.15 207 ALBRECHT 92L ARG $e^+e^- \rightarrow \Upsilon(4S)$

206 Value is their Δm_{B^0} measurement combined with $\tau_{B^0} = (1.50 \pm 0.11)$ ps, the average from this edition. The systematic error on τ_{B^0} and is common to experiments bearing this footnote. The averaging takes this into account.

207 Derived from time-integrated mixing parameter χ .

 B^0 - \bar{B}^0 MIXING AND CP VIOLATION IN B DECAY (by H. Quinn, SLAC)

The neutral B meson system is like the neutral kaon system, in that two CP -conjugate states exist. The mass eigenstates are mixtures of these states, the mixing being due to box diagrams, shown in Figure 1. The two mass eigenstates can be written

$$\begin{aligned} |B_L\rangle &= p|B_0\rangle + q|\bar{B}^0\rangle, \\ |B_H\rangle &= p|B_0\rangle - q|\bar{B}^0\rangle. \end{aligned} \quad (1)$$

Here H and L stand for Heavy and Light, respectively.




Figure 1: Mixing Diagrams.

Whereas in the kaon case the lifetimes of the two eigenstates are significantly different and the difference in masses between them is small, in the B system it is the mass differences that dominate the physics, and the two states have nearly equal predicted widths (and thus lifetimes).

$$\Gamma = (\Gamma_H + \Gamma_L)/2, \quad \Delta\Gamma = \Gamma_H - \Gamma_L. \quad (2)$$

The difference between the widths of the two eigenstates is produced by the contributions from channels to which both B^0 and \bar{B}^0 can decay. These have branching ratios of $\mathcal{O}(10^{-3})$ [1]. Furthermore there are contributions of both signs to the difference, so there is no reason that the net effect should be much larger than the individual terms. Conservatively, one expects $\Delta\Gamma/\Gamma \leq 10^{-2}$. Experimentally no effect of a difference in lifetimes has been observed. In what follows, we neglect any effects from $\Delta\Gamma$, except where explicitly stated. We define also

$$M \equiv (M_H + M_L)/2, \quad \Delta M \equiv M_H - M_L. \quad (3)$$

The proper time evolution of an initially ($t = 0$) pure B^0 or \bar{B}^0 is given by

$$\begin{aligned} |B_{\text{phys}}^0(t)\rangle &= \exp(-\Gamma t/2) \exp(-iMt) \\ &\quad \times \{ \cos(\Delta Mt/2) |B_0\rangle + i(q/p) \sin(\Delta Mt/2) |\bar{B}^0\rangle \}, \\ |\bar{B}_{\text{phys}}^0(t)\rangle &= \exp(-\Gamma t/2) \exp(-iMt) \\ &\quad \times \{ i(p/q) \sin(\Delta Mt/2) |B_0\rangle + \cos(\Delta Mt/2) |\bar{B}^0\rangle \}. \end{aligned} \quad (4)$$

The probability that an initial B^0 (\bar{B}^0) decays as a \bar{B}^0 (B^0) is thus

$$P(t) = \frac{1}{2} e^{-\Gamma t} (1 - \cos(\Delta Mt)) \quad (5)$$

where we have used $|p/q| = 1$ which is true when we neglect the effects of $\Delta\Gamma$. Time-dependent mixing measurements are now beginning to be done; earlier experiments measured the time-integrated mixing, which is parameterized by a parameter χ_d for B_d (i.e., B^0) and χ_s for B_s (i.e., B_s^0). The quantity χ measures the total probability that a created B^0 decays as a \bar{B}^0 ; it is given by

$$\chi_q = \int_0^\infty P_q(t) dt = \frac{x_q^2}{2(1+x_q^2)}. \quad (6)$$

Here quantity x_q is the ratio of the $B^0 - \bar{B}^0$ oscillation frequency to the decay rate $x_q = \frac{\Delta M_q}{\Gamma_q}$, $q = d, s$. The value of x_d is about 0.7, not very different from the similar quantity for the K^0 which is 0.48. The value of x_s is expected to be much larger, so that the quantity χ_s will be close to its upper limit of 0.5. This means that one cannot determine x_s accurately by measuring χ_s . It will require excellent time resolution to resolve the time-dependent mixing of the B_s^0 system, and thereby determine $\Delta M_{B_s^0}$ [2].

In the $B^0 - \bar{B}^0$ mixing section of the B^0 Full Listings, we list the χ_d measurements, most of which come from $\Upsilon(4S)$ data, and the Δm_{B^0} measurements, which come from Z data. We convert both of these sets of measurements and list them in the x_d section. The x_d values obtained from Δm_{B^0} measurements have a common systematic error due to the error on τ_{B^0} . The averaging takes this common systematic error into account.

In the $B_s^0 - \bar{B}_s^0$ mixing section of the B_s^0 Full Listings, we give measurements of χ_B at high energy and of χ_s obtained by combining the χ_B and χ_d measurements using

$$\chi_B = f_d \chi_d + f_s \chi_s, \quad (7)$$

where f_d and f_s are the fractions of b hadrons that are produced as B^0 and B_s^0 mesons respectively. We also convert the average χ_B and χ_d values to χ_s assuming $f_d = 0.391$ and assuming $f_s = 0.117$, obtaining $\chi_s = 0.62 \pm 0.13$, consistent with the maximum allowed value $\chi_s = 0.5$. This χ_s value and error, when converted to an x_s lower limit, yields $x_s > 1.5$. There

is now a limit on Δm_{B^0} from an ALEPH time-dependent $B_s^0 - \bar{B}_s^0$ mixing measurement which provides slightly stronger limit, $x_s > 2.0$, than that obtained from the time-independent measurements.

CP violation in B decays—Standard Model predictions:

There are three symmetries of the strong interactions that are not conserved in weak processes. These are the symmetries C , charge conjugation, which relates particle to antiparticle, P , parity, which relates a left-handed particle to a similar right-handed one, and T , time-reversal invariance, which relates a process or state to the time-reversed process or state. In all field theories the product of these three operations, CPT , is an exact symmetry of the equations of motion. All weak decays violate P and C , and a very small part of the weak decays also violate the product CP (and thus T). In the Standard Model this CP violation occurs because there is a single phase that remains in the Cabibbo-Kobayashi-Maskawa (CKM) matrix after all possible field redefinitions that can remove such phases have been made. In a minimal two-generation Standard Model no such phase occurs. The presence of CP -violating effects in K decays was interpreted by Kobayashi and Maskawa in 1973 to suggest a third quark generation. Other extensions beyond the minimal Standard Model, such as theories with additional Higgs multiplets, give further ways to introduce CP violation into the theory. Hence it is of great interest to study whether the pattern of CP -violating effects that can be observed in B decays follows the predictions of the minimal Standard Model, or instead requires the introduction of beyond Standard Model effects. In what follows the predictions given are all those of the minimal Standard Model. Cosmologists attempting to understand the process by which the matter-antimatter asymmetry of the universe arose suggest that additional sources of CP violation may be needed to give the observed baryon to photon ratio of the universe [3]. Many models which go beyond the Standard Model indeed introduce such possibilities (see “Beyond Standard Model effects” below.)

The CKM matrix is the matrix of weak couplings in the three generation Standard Model, expressed in the basis of quark mass eigenstates. This matrix, which must be unitary if the three generations are the complete theory, is discussed in some detail in a separate article in this *Review*, here we need only remind ourselves of some notation that is commonly used in this context. The matrix can be written

$$V = \begin{pmatrix} V_{ud} & V_{us} & V_{ub} \\ V_{cd} & V_{cs} & V_{cb} \\ V_{td} & V_{ts} & V_{tb} \end{pmatrix} \quad (8)$$

$$\simeq \begin{pmatrix} 1 - \lambda^2/2 & \lambda & A\lambda^3(\rho - i\eta) \\ \lambda & 1 - \lambda^2/2 & A\lambda^2 \\ A\lambda^3(1 - \rho - i\eta) & -A\lambda^2 & 1 \end{pmatrix} + O(\lambda^4).$$

The second expression here is a parameterization due to Wolfenstein [4] with $\lambda = \sin(\theta_{\text{Cabibbo}})$, which is frequently used in discussing CP -violating effects. It is given here up to

Meson Full Listings

B^0

terms of order λ^3 , since higher order terms in λ are negligible for all practical purposes. The unitarity triangle is a simple geometrical representation of a relationship which results from the unitarity of the three-generation CKM matrix V :

$$V_{ud}V_{ub}^* + V_{cd}V_{cb}^* + V_{td}V_{tb}^* = 0, \quad (9)$$

(In fact there are nine such relationships given by the unitarity of the CKM matrix, but only three are independent conditions and of those the other two will be more difficult to test because they have one term that is of order λ^2 relative to the others.) The three complex quantities $V_{id}V_{ib}^*$ form a triangle in the complex plane. The three angles of this triangle are labeled

$$\begin{aligned} \alpha &\equiv \arg\left(-\frac{V_{td}V_{tb}^*}{V_{ud}V_{ub}^*}\right), & \beta &\equiv \arg\left(-\frac{V_{cd}V_{cb}^*}{V_{td}V_{tb}^*}\right), \\ \gamma &\equiv \arg\left(-\frac{V_{ud}V_{ub}^*}{V_{cd}V_{cb}^*}\right). \end{aligned} \quad (10)$$

Figure 2 shows the unitarity triangle, as it is usually drawn, rescaled by the side $V_{cd}V_{cb}^*$. This makes the base of the triangle real and of unit length and the apex of the triangle is then the point (ρ, η) in the complex plane. A major aim of CP -violation studies of B decays is to make enough independent measurements of the sides and angles that this triangle is overdetermined and thus to check the validity of the Standard Model. Already a number of constraints can be made on the basis of present data on x_d , V_{ub}/V_{cb} , and ϵ in K decays. These constraints have been discussed in many places in the literature [5], their exact form depends on the mass of the top quark and on the range of values allowed for the B_K parameter in K decays and the parameter combination $B_B f_B^2$ in B decays.




Figure 2: The (rescaled) Unitarity Triangle.

The CKM phases, that is the phases in decay amplitudes which arise because of the phase in the CKM matrix, are often called weak phases, in contrast to the phases which arise from final state rescattering effects which are referred to as strong phases. When one compares the amplitude for a process with that for the related CP conjugate process the weak phase of each contribution changes sign, while the strong phase is unchanged.

$$\mathcal{A} = \sum_i \mathcal{A}_i e^{i(\delta_i + \phi_i)}, \quad \bar{\mathcal{A}} = \sum_i \mathcal{A}_i e^{i(\delta_i - \phi_i)}. \quad (11)$$

Direct CP violation is a difference in the direct decay rate between $B \rightarrow f$ and $\bar{B} \rightarrow \bar{f}$ without any contribution from

mixing effects. As can readily be seen from Eq. (11), this requires that there be more than one term in the sum Eq. (11) and further that the two terms have both different weak phases and different strong phases. A nonzero result for ϵ' in K decay is a direct CP -violation effect. Direct CP violation can occur both in charged channels and in neutral channels.

In the Standard Model, direct CP violation occurs because there are two classes of diagram that contribute to flavor-changing decays, called tree diagrams and penguin diagrams. Tree diagrams are those in which the W does not reconnect to the quark line from which it was emitted, penguin diagrams are those in which it does. There may be several different tree diagrams for a given process, namely emission from the heavy quark line followed by W decay, W exchange between the initial valence quarks, and/or valence quark-antiquark annihilation to produce the W . However all such contributions which enter a given transition do so with the same CKM (weak) phase. Hence direct CP violation occurs only from interference between tree and penguin type contributions in the Standard Model.

To calculate the size of expected CP -violation effects one begins from the relevant quark decay diagrams. In general weak-decay amplitudes for b quarks can be divided into two factors: a CKM factor given by the CKM-matrix elements that enter at each W vertex, and a kinematic factor from evaluating the Feynman diagram. The penguin diagrams are a loop diagram, and in addition require the emission of a hard gluon (or photon) from the loop to account for the mass difference between the b quark and the s or d quark produced when the W is reabsorbed. The kinematic factor of the penguin diagram is thus suppressed relative to tree diagrams by a factor of order $\alpha(m_b)/4\pi$. It is difficult to make firm predictions based on this argument for the strength of the CP -violating effects in exclusive charged B -decay channels because the relationship between the free-quark decay diagrams and the exclusive meson decay amplitudes is model dependent. Furthermore one cannot reliably predict the strong phases that contribute to the asymmetry. However, due to the suppression of penguin contributions, one does not expect direct CP -violating effects in charged B decays to be more than a few percent effects in the rare (doubly Cabibbo suppressed) decay channels ($b \rightarrow q\bar{q}d$, $q = d$ or s), or a few tenths of a percent in the Cabibbo-suppressed modes ($b \rightarrow q\bar{q}s$, $q = u, d, s$) [6]. Effects much larger than this would suggest contributions from beyond the Standard Model. Similar results apply for CP asymmetries in the decays of neutral B 's to hadronic final states that can only be reached by a definite b flavor.

There are additional CP violating effects in neutral B decays due to interference between the two paths to a given final state f

$$B \rightarrow f \text{ or } B \rightarrow \bar{B} \rightarrow f$$

This is referred to as CP violation due to mixing, or more precisely due to interference between the mixed and unmixed decay paths. It is similar to the effect measured by the parameter ϵ

in K decay. The interference between the two contributions can produce rate differences between the decay and its CP conjugate. These effects are of particular interest because they do not depend upon strong phases and hence the measured asymmetries can be directly related to the CKM phases. In any given channel there can also be direct CP violation in addition to the CP violation due to mixing, this complicates the relationship between the measured asymmetry and the CKM parameters. We will briefly discuss later techniques to separate such contributions.

A third type of CP violation, referred to as indirect CP violation, would arise from any difference in the widths $\Delta\Gamma$ of the two mass eigenstates, or more precisely from complex mixing effects that would also give a nonvanishing lifetime difference for the two B mass eigenstates. Such effects are expected to be tiny in the B system. In the particular case of semileptonic decays there are no penguin diagram contributions, and hence, in the approximations used throughout the discussion above, the CP -conjugate decay rates are equal. However there is a possible indirect CP -violating asymmetry in these decays, which would be seen as a charge asymmetry in the same-sign dilepton events produced via mixing from an incoherent state that initially contains a $B^0\bar{B}^0$ pair. This asymmetry vanishes with $\Delta\Gamma$; it is expected to be no larger than 1% [7].

A simple way to distinguish the three types of CP violation is to note that direct CP violation occurs when $|\mathcal{A}/\bar{\mathcal{A}}| \neq 1$, indirect CP violation requires $|q/p| \neq 1$, but CP violation due to the interference between decays with and without mixing can occur when both quantities have unit absolute value; it requires only that the ratio of them has a nonzero weak phase [8].

Neutral B decays to CP eigenstates: The decays of neutral B 's into a CP eigenstates is of particular interest because many of these decays allow clean theoretical interpretation in terms of the parameters of the Standard Model [9]. We denote such a state by f_{CP} , for example $f_{CP} = J/\psi(1S)K_S$ or $f_{CP} = \pi\pi$, and define the amplitudes

$$\mathcal{A}_{f_{CP}} \equiv \langle f_{CP}|B^0\rangle, \quad \bar{\mathcal{A}}_{f_{CP}} \equiv \langle f_{CP}|\bar{B}^0\rangle. \quad (12)$$

For convenience let us introduce the quantity $r_{f_{CP}}$

$$r_{f_{CP}} \equiv \frac{q}{p} \frac{\bar{\mathcal{A}}}{\mathcal{A}}. \quad (13)$$

In the limit of no CP violation, $r_{f_{CP}} = \pm 1$ where the sign is given by the CP eigenvalue of the particular state f_{CP} . (Note that in the literature the quantity $r_{f_{CP}}$ is frequently denoted by λ , but we have chosen to avoid this notation as it introduces a confusion with the $\lambda = \sin(\theta_{\text{Cabibbo}})$ in the Wolfenstein parameterization of the CKM matrix.)

The time-dependent rates for initially pure B^0 or \bar{B}^0 states to decay into a final state f_{CP} at time t is then given by:

$$\begin{aligned} \langle f_{CP}|B_{\text{phys}}^0(t)\rangle &= \mathcal{A} \exp(-\Gamma t/2) \exp(-iMt) \\ &\quad \times [\cos(\Delta Mt/2) + ir_{f_{CP}} \sin(\Delta Mt/2)], \\ \langle f_{CP}|\bar{B}_{\text{phys}}^0(t)\rangle &= \mathcal{A} \exp(-\Gamma t/2) \exp(-iMt)(p/q) \\ &\quad \times [i \sin(\Delta Mt/2) + r_{f_{CP}} \cos(\Delta Mt/2)]. \end{aligned} \quad (14)$$

Thus

$$\begin{aligned} \Gamma(B_{\text{phys}}^0(t) \rightarrow f_{CP}) &= |\mathcal{A}|^2 e^{-\Gamma t} \left[\frac{1 + |r_{f_{CP}}|^2}{2} + \frac{1 - |r_{f_{CP}}|^2}{2} \right. \\ &\quad \left. \times \cos(\Delta Mt) - \text{Im } r_{f_{CP}} \sin(\Delta Mt) \right], \\ \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow f_{CP}) &= |\mathcal{A}|^2 e^{-\Gamma t} \left[\frac{1 + |r_{f_{CP}}|^2}{2} - \frac{1 - |r_{f_{CP}}|^2}{2} \right. \\ &\quad \left. \times \cos(\Delta Mt) + \text{Im } r_{f_{CP}} \sin(\Delta Mt) \right]. \end{aligned} \quad (15)$$

The time-dependent CP asymmetry is

$$a_{f_{CP}}(t) \equiv \frac{\Gamma(B_{\text{phys}}^0(t) \rightarrow f_{CP}) - \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow f_{CP})}{\Gamma(B_{\text{phys}}^0(t) \rightarrow f_{CP}) + \Gamma(\bar{B}_{\text{phys}}^0(t) \rightarrow f_{CP})} \quad (16)$$

and thus

$$a_{f_{CP}}(t) = \frac{(1 - |r_{f_{CP}}|^2) \cos(\Delta Mt) - 2 \text{Im}(r_{f_{CP}}) \sin(\Delta Mt)}{1 + |r_{f_{CP}}|^2} \quad (17)$$

When the small difference in width of the two B states is ignored we can write

$$(q/p)_{B_d} = \frac{(V_{tb}^* V_{td})}{V_{ib} V_{id}^*}, \quad (q/p)_{B_s} = \frac{(V_{tb}^* V_{ts})}{V_{ib} V_{is}^*}.$$

and thus

$$q/p = e^{-2i\phi_M}, \quad (18)$$

where $2\phi_M$ denotes the CKM phase of the B - \bar{B} mixing diagram. (Note, Eq. (9) allows us to write the sum of all contributions to the mixing in this form, up to small corrections of order $(m_c^2 - m_u^2)/m_W^2$.) Further, when there is no direct CP violation in a channel that is when all amplitudes that contribute to the decay have the same CKM phase, ϕ_D , then $|\mathcal{A}/\bar{\mathcal{A}}| = 1$. In that case $r_{f_{CP}}$ depends on CKM-matrix parameters only, without hadronic uncertainties, and can be written $r_{f_{CP}} = \pm e^{-2i(\phi_D + \phi_M)}$. Then Eq. (17) simplifies to

$$\begin{aligned} a_{f_{CP}}(t) &= \mp \text{Im}(r_{f_{CP}}) \sin(\Delta Mt) \\ &= \pm \sin(2(\phi_M + \phi_D)) \sin(\Delta Mt). \end{aligned} \quad (19)$$

Meson Full Listings

 B^0

where the overall sign is given by the CP eigenvalue, ± 1 , of the final state f_{CP} . The mixing phase ϕ_M and the decay phase ϕ_D are each convention dependent, that is their value can be changed by redefining the phases of some of the quark fields. However $\text{Im } r_{f_{CP}}$ depends on convention-independent combinations of CKM parameters only, and thus from Eq. (19) one can directly relate the measured CP -violating asymmetry to the phase of particular combination of CKM-matrix elements in the Standard Model.

In an $e^+e^- B$ collider running at the $\Upsilon(4S)$ resonance, the initial B system is produced in a coherent state which remains $B^0\bar{B}^0$ until such time as one of the particles decays. The time evolution of the second particle thus begins at the time of the decay of the first. Events where one B decays to a flavor-tagging mode while the other decays to a CP -study mode can be used to reconstruct the dependence of the asymmetry on the time between the tagging decay and the CP -study mode decay. The tagging decay may be later, in which case the event is assigned a negative time. Note that the measurement of time dependence is essential at such a machine since, in the interesting cases where Eq. (19) applies, the time-integrated CP asymmetry vanishes.

Hadronic machines on the other hand produce uncorrelated B and \bar{B} mesons; the time in the above equations is the time between production and decay, and thus is always positive. Time-integrated asymmetries do not vanish in this case. The time evolution of the tagging particle occurs independently and will contribute to wrong sign tags. Such machines produce many more B 's than will an $e^+e^- B$ factory. For modes with a clean signature, such as those with a final state $J/\psi(1S)$ or ϕ , hadronic machines can compete with a B factory in measuring CP asymmetries, but for modes such as $\pi\pi$ or $\rho\pi$ the problems of triggering and of backgrounds make such measurements extremely challenging in a hadronic environment.

Extracting CKM parameters from measured asymmetries: In order to relate the measured asymmetries to the CKM-matrix parameters one looks at the CKM elements that appear in the relevant decay amplitudes and in the mixing diagrams. Table 1 gives the result for the various b -quark decay channels. For the penguin contributions the CKM factors given here are obtained by assuming CKM unitarity and neglecting small corrections due to the difference between charm and up quark masses in the loop for $b \rightarrow d$ processes, and those due

Table 1: B decay modes for CP studies.

Quark Process	Tree CKM	Leading Penguin CKM	Sample B_d Modes	B_d Angle	Sample B_s Modes	B_s Angle	Comments
$b \rightarrow c\bar{c}s$	$V_{cb}V_{cs}^* = A\lambda^2$	$V_{cb}V_{cs}^* = A\lambda^2$	$J/\psi(1S)K_S$	β	$J/\psi(1S)\eta', D_s\bar{D}_s$	0	(a)
$b \rightarrow s\bar{s}s$	0	$V_{cb}V_{cs}^* = A\lambda^2$	ϕK_S	β	$\phi\eta'$	0	(b)
$b \rightarrow u\bar{u}s$	$V_{ub}V_{us}^* = A\lambda^4(\rho - i\eta)$	$V_{cb}V_{cs}^* = A\lambda^2$	$K_S\pi^0, K_S\rho^0$	–	$\phi\pi^0, K^+K^-$	–	(c)
$b \rightarrow d\bar{d}s$	0	$V_{cb}V_{cs}^* = A\lambda^2$	$K_S\pi^0, K_S\rho^0$	–	$\phi\pi^0, K_S\bar{K}_S$	–	(c)
$b \rightarrow c\bar{c}d$	$V_{cb}V_{cd}^* = -A\lambda^3$	$V_{tb}V_{td}^* = A\lambda^3(1 - \rho + i\eta)$	$D^+D^-, J/\psi(1S)\pi^0, D^0\bar{D}^0(\dagger)$	β	$J/\psi(1S)K_S$	0	(d)
$b \rightarrow s\bar{s}d$	0	$V_{tb}V_{td}^* = A\lambda^3(1 - \rho + i\eta)$	$\phi\pi^0, K_S\bar{K}_S$	0	ϕK_S	β	(c)
$b \rightarrow u\bar{u}d$	$V_{ub}V_{ud}^* = A\lambda^3(\rho - i\eta)$	$V_{tb}V_{td}^* = A\lambda^3(1 - \rho + i\eta)$	$\pi\pi, \pi\rho, \pi a_1$	α	$\pi^0 K_S, \rho^0 K_S$	γ	(e)
$b \rightarrow d\bar{d}d$	0						
$b \rightarrow c\bar{u}s$	$V_{cb}V_{us}^* = A\lambda^3$	0	$D_{CP}^0 K^*(892)$	γ	$D_{CP}^0 \phi$	–	(f), (g)
$b \rightarrow u\bar{c}s$	$V_{ub}V_{cs}^* = A\lambda^3(\rho - i\eta)$						
$b \rightarrow c\bar{u}d$	$V_{cb}V_{cd}^* = A\lambda^2$	0	$D_{CP}^0\pi^0, D_{CP}^0\rho^0$	–	$D_{CP}^0 K_S$	–	(g)
$b \rightarrow u\bar{c}d$	$V_{ub}V_{cd}^* = A\lambda^4(\rho - i\eta)$						

(a) Tree and penguin contribute with same weak phase.

(b) Penguin only, rare decays.

(c) Tree and penguin compete. Isospin analysis may allow extraction of α, β , for B_d channels, $\gamma, 0$ for B_s , where these angles come from tree and penguin contributions respectively. $K_S\bar{K}_S$ penguin only, except 0 asymmetry.

(d) Ignoring penguin relative to tree.

(e) Ignoring penguin relative to tree, or using isospin analysis.

(f) Self-tagging $K^*(892)$ decay modes can give γ when data from $B_d \rightarrow D_{CP}^0$, i.e. decays to CP eigenstates, and D^0 - or \bar{D}^0 -identified modes are combined. Similar results for charged $B \rightarrow DK$.

(g) Asymmetry in $D_{CP}^0\pi, D_{CP}^0K_S$, etc. modes is difficult to relate to CKM angles.

(†) $D^0\bar{D}^0$ from rescattering only, rate expected to be small.

to terms of higher order in $\lambda = \sin(\theta_{\text{Cabibbo}})$ for $b \rightarrow s$ decays. When the decay also involves a K_S in the final state an additional contribution to the phase from the K -mixing phase must be included in relating the measured asymmetry to the CKM parameters.

The columns labeled “Sample B_d Modes” and “Sample B_s Modes” list some of the simplest CP -eigenstate modes for each case. The columns labeled “Angle” show the particular combination of CKM phases $\phi_M + \phi_D$ that is measured by the CP -violating asymmetry in these decays, given as an angle of the unitarity triangle. For most channels the measured asymmetry in a time-dependent measurement is $\pm \sin(2(\phi_M + \phi_D)) \sin(\Delta Mt)$ while in a time-integrated measurement the asymmetry is $\pm x_q / (1 + x_q^2) \sin(2(\phi_M + \phi_D))$. The sign is given by the CP eigenvalue of the particular final state studied. The exception to these statements is the channel DK discussed below.

The assumptions needed to obtain the results in the “Angle” columns are noted in the comments below the table. Terms of higher order in $\lambda = \sin(\theta_{\text{Cabibbo}})$ have been dropped. Penguin diagrams that occur at the same order of λ as the corresponding tree diagrams are neglected in stating the relationship of the asymmetry to angles in the unitarity triangle. In cases where the tree and penguin diagrams are expected to give comparable contributions with different CKM phases there is no simple relationship between the measured asymmetry and a CKM phase, and thus no entry in the “Angle” columns in Table 1.

The mode $D^0 K^*(892)$ is listed even though it is not a CP eigenstate because it has been shown that an analysis of this mode can be used to extract the angle γ . [10]. The same type of analysis can also be applied for charged B decays. However the relationship between the decay asymmetry and the angle is not as simple as Eq. (19) in this case. The result depends on measurements of a number of branching ratios. It will be very difficult to obtain accurate results by this method.

In the case of the $b \rightarrow u\bar{u}d$ and $b \rightarrow d\bar{d}d$, the penguin contributions occur at the same order in λ as the tree diagrams and are thus expected to be small compared to them because of the $\alpha(m_b)/\pi$ suppression factor. The result given in Table 1 makes this approximation. If however this expectation proves false, so that the contributions are comparable, one still may be able to extract a measurement of $\sin(2\alpha)$ from the $\pi^+\pi^-$ asymmetry. This is achieved by measuring the rates in several isospin-related channels and using a multiparameter fit to separate tree and penguin contributions to the amplitudes [12]. This will require measuring the decay rate for channel $\pi^0\pi^0$, which will be a challenge. For the $\rho\pi$ decays, if penguins are not negligible, the restrictions due to isospin can again be used to make a multiparameter fit to the ρ -regions of the Dalitz plot for $\pi^+\pi^-\pi^0$ distribution [13]. The interference between different ρ -charge channels is significant and may provide sufficient information to allow the separation of tree and penguin effects and thus extraction of the parameter α . Such analyses at the very

least can be used to test whether the penguin contributions are indeed small enough to be neglected in the determination of α .

There are some common decay channels of the B^0 and \bar{B}^0 which are not CP eigenstates. For example the channel $J/\psi(1S)K^*(892)$ where the $K^*(892) \rightarrow K_S\pi^0$, the final state is not a CP eigenstate because both even and odd relative angular momenta between the $J/\psi(1S)$ and the $K^*(892)$ are allowed. If there is sufficient data one can use angular analysis to separate the different CP final states and measure the asymmetry in each [14]. The same applies in many quasi-two-body decays, such as other vector-vector channels, or those with higher-spin particles in final states. The branching ratio to these channels can be significantly larger than the CP -eigenstate (vector-scalar or scalar-scalar) channels with the same quark content. Such angular analyses may therefore be important in achieving accurate values for the parameters α and β .

Beyond Standard Model effects: The predictions given above are all for the Standard Model. Models beyond the Standard Model may introduce additional contributions to the mixing amplitudes and thereby destroy the relationships given here; in addition they may introduce further direct CP violation.

One model often used as a “straw man” in evaluating the potential of experimental tests of Standard Model predictions is the superweak model, which was one of the earliest proposals for the mechanism of CP violation, in fact prior to the Standard Model [15]. In the modernized version of this model it is assumed that the CKM matrix is real and that all CP -violating effects arise from a contribution to the mixing that comes from beyond the Standard Model. In this case *all* the CP -eigenstate channels for B decay would have the same CP -violating asymmetry (up to a sign which differs for CP -odd and CP -even channels) [16]. This applies even to those channels predicted to have zero asymmetry in the Standard Model, as well as those for which the Standard Model prediction is complicated by the competition between tree and penguin contributions. Observation of significantly different asymmetries in any two neutral B decay CP -eigenstate channels would rule out such a model. In addition the observation of any asymmetry in a charged B decay or a neutral B decay to flavor-tagging final state would be evidence for direct CP violation and would exclude the superweak model, as would an unequivocal nonzero result for the parameter ϵ' in K decays.

Many other models for the physics beyond the Standard Model have been discussed in the literature [17]. The most common additional CP -violating effect is a new contribution to the mixing process, due for example to charged Higgs contributions. The appearance of such contributions in K mixing is already severely restricted by the neutral- K mass difference. However this does not rule out additional contributions to B mixing that would destroy the relationship between the mixing phase ϕ_M and the CKM-matrix elements. This in turn would lead to violations of the predictions given in Table 1 which are based on

Meson Full Listings

 B^0

this relationship. Models with additional quarks would remove the constraints due to the unitarity of the three-generation mixing matrix and hence lead to the failure of Eq. (9) [18]. (The absence of any fourth light neutrino determined from Z decay rules out only a standard fourth generation with a light neutrino but does not exclude nonstandard fourth generation models.) Any observed deviations from the relationships predicted by the Standard Model will provide a window on the nature of physics beyond the Standard Model.

While the discussion above stresses those channels in which there is a simple relationship between any observed asymmetry and the parameters of the CKM matrix in the Standard Model, this does not mean that other channels are entirely without interest. To date only the CP violation in the neutral K system has been observed. Any observation of CP violation in B decays would be exciting. The Standard Model prediction is that direct CP -violating asymmetries are likely to be at most a few percent, so large effects in these channels would suggest beyond Standard Model effects. On the other hand, even within the Standard Model the asymmetries due to the interference between decays with and without mixing in the neutral B system can be quite large; current constraints do not rule out cases where $\text{Im}(\tau_{fCP})$ is 1. It is likely that study of the many common decay channels of the B^0 and the \bar{B}^0 will greatly expand our understanding of the sources of CP violation.

References

1. I.I. Bigi, V.A. Khoze, N.G. Uraltsev, and A.I. Sanda, in *CP Violation*, ed. C. Jarlskog (World Scientific, Singapore, 1989), p. 175.
2. We could refer to recent review papers here, or leave references to the data section below. Your choice Tom.
3. For a review of this subject see lectures given by M. Dine at the 1992 Theoretical Advanced Study Institute, UC Santa Cruz preprint Preprint SCIPP-92-46.
4. L. Wolfenstein, Phys. Rev. Lett. **51**, 1945 (1984).
5. See for example Y. Nir Lectures given at the 1992 SLAC Summer School, SLAC-Pub 5874 (1992) and references therein.
6. J.M. Gérard and W.S. Hou, Phys. Lett. **B253**, 478 (1991) and Phys. Rev. **D43**, 2909 (1991); H. Simma, G. Eilam, and D. Wyler, Nucl. Phys. **B352**, 367 (1991).
7. T. Altomari, L. Wolfenstein, and J.D. Bjorken, Phys. Rev. **D37**, 1860 (1988).
8. For a review of CP violation in B decays, Y. Nir and H.R. Quinn, Ann. Rev. Nucl. and Part. Sci. **42**, 211 (1992) and references contained therein.
9. I. Dunietz and J.L. Rosner, Phys. Rev. **D34**, 1404 (1986).
10. I. Dunietz, Phys. Lett. **B270**, 75 (1991).
11. M. Gronau and D. Wyler, Phys. Lett. **B265**, 172 (1991).
12. M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990).
13. H.R. Quinn and A.E. Snyder, Phys. Rev. **D48**, 2139 (1993).
14. J.R. Dell'Aquila and C.A. Nelson, Phys. Rev. **D33**, 101 (1986); B. Kayser *et al.*, Phys. Lett. **B237**, 3339 (1990); I. Dunietz *et al.*, Phys. Rev. **D43**, 2193 (1991).
15. L. Wolfenstein, Phys. Rev. Lett. **13**, 562 (1964).
16. For a recent discussion see J.M. Soares and L. Wolfenstein, Phys. Rev. **D46**, 256 (1992).
17. For a summary of the predictions of such models see C.O. Dib, D. London, and Y. Nir, Int. J. Mod. Phys. **A6**, 1253 (1991).
18. See for example Y. Nir and D. Silverman, Nucl. Phys. **B345**, 301 (1990); Y. Nir and D. Silverman, Phys. Rev. **D42**, 1477 (1990).

CP VIOLATION PARAMETERS

$|\text{Re}(\epsilon_{B^0})|$

CP Impurity in B^0 system. It is obtained from $a_{\ell\ell}$, the charge asymmetry in like-sign dilepton events at the $T(4S)$.

$$\text{Re}(\epsilon_{B^0}) \approx \frac{1}{4} a_{\ell\ell} = \frac{1}{4} \frac{N(\ell^+\ell^+) - N(\ell^-\ell^-)}{N(\ell^+\ell^+) + N(\ell^-\ell^-)}$$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.045	208 BARTELT	93 CLE2	$e^+e^- \rightarrow T(4S)$
208 BARTELT 93 finds $a_{\ell\ell} = 0.031 \pm 0.096 \pm 0.032$ which corresponds to $ a_{\ell\ell} < 0.18$, which yields the above $\text{Re}(\epsilon_{B^0})$.			

B^0 REFERENCES

ABE 94D PRL 72 3456	+Albrow, Amidei, Anway-Wiese, Apollinari	(CDF Collab.)
AKERS 94C PL B327 411	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
ALAM 94 PR D (to be pub.)	+Kim, Nemati, O'Neill, Severini+	(CLEO Collab.)
ALBRECHT 94 PL B324 249	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
AMMAR 94 PR D49 5701	+Ball, Baringer, Bean, Besson, Coppage+	(CLEO Collab.)
BUSKULIC 94B PL B322 441	+De Bonis, Decamp, Ghez, Goy, Lees+ (ALEPH Collab.)	(CERN, LBL, BOST, IFIC+)
PDG 94 PR D50 1173	+Montanet+	(CLEO Collab.)
PROCARIO 94 PRL (to be pub.)	+Baest, Cho, Daoudi, Ford+	(CLEO Collab.)
CLNS 93/1264, CLEO 93-24		
STONE 94 HEPSY 93-11		
ABREU 93D ZPHY C57 181	+Adam, Aoye, Agasi, Alekseev+	(DELPHI Collab.)
ABREU 93G PL B312 253	+Adam, Aoye, Agasi, Ajinenki+	(DELPHI Collab.)
ACTON 93C PL B307 247	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ALBRECHT 93 ZPHY C57 533	+Ehrlichmann, Hamacher, Hofmann+	(ARGUS Collab.)
ALBRECHT 93E ZPHY C60 11	+Ehrlichmann, Hamacher, Hofmann+	(ARGUS Collab.)
ALEXANDER 93B PL B319 365	+Bebek, Berkelman, Bloom, Browder+	(CLEO Collab.)
AMMAR 93 PRL 71 674	+Ball, Baringer, Coppage, Copty+	(CLEO Collab.)
BARTELT 93 PRL 71 1680	+Csorna, Egedy, Jain, Sheldon+	(CLEO Collab.)
BATTLE 93 PRL 71 3922	+Ernst, Kroha, Kwon, Roberts+	(CLEO Collab.)
BEAN 93B PRL 70 2681	+Gronberg, Kutschke, Menary, Morrison+	(CLEO Collab.)
BUSKULIC 93D PL B307 194	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
BUSKULIC 93K PL B313 498	+De Bonis, Decamp, Ghez, Goy+	(ALEPH Collab.)
SANGHERA 93 PR D47 791	+Skwarnicki, Stroynowski, Artuso, Goldberg+	(CLEO Collab.)
ALBRECHT 92C PL B275 195	+Ehrlichmann, Hamacher, Krueger, Nau+	(ARGUS Collab.)
ALBRECHT 92G ZPHY C54 1	+Ehrlichmann, Hamacher, Krueger, Nau+	(ARGUS Collab.)
ALBRECHT 92L ZPHY C55 357	+Ehrlichmann, Hamacher, Krueger, Nau+	(ARGUS Collab.)
BORTOLETTO 92 PR D45 21	+Brown, Dominick, McIlwain+	(CLEO Collab.)
HENDERSON 92 PR D45 2212	+Kinoshita, Pipkin, Procario+	(CLEO Collab.)
KRAMER 92 PL B279 181	+Palmer	(HAMB, OSU)
ALBAJAR 91C PL B262 163	+Albrow, Allkofer, Ankoviak, Apsimon+	(UA1 Collab.)
ALBAJAR 91E PL B273 540	+Albrow, Allkofer, Ankoviak+	(UA1 Collab.)
ALBRECHT 91B PL B254 288	+Glaeser, Harder, Krueger, Nippe+	(ARGUS Collab.)
ALBRECHT 91C PL B255 297	+Ehrlichmann, Glaeser, Harder, Krueger+	(ARGUS Collab.)
ALBRECHT 91E PL B262 148	+Glaeser, Harder, Krueger, Nippe+	(ARGUS Collab.)
BERKELMAN 91 ARNPS 41 1	+Stone	(CORN, SYRA)
"Decays of B Mesons"		
FULTON 91 PR D43 651	+Jensen, Johnson, Kagan, Kass+	(CLEO Collab.)
ALBRECHT 90B PL B241 278	+Glaeser, Harder, Krueger, Nilsson+	(ARGUS Collab.)
ALBRECHT 90J ZPHY C48 543	+Ehrlichmann, Harder, Krueger+	(ARGUS Collab.)
BORTOLETTO 90 PRL 64 2117	+Goldberg, Horwitz, Jain, Mestayer+	(CLEO Collab.)
ELSEN 90 ZPHY C46 349	+Allison, Ambrus, Barlow, Bartel+	(JADE Collab.)
ROSNER 90 PR D42 3732		
WAGNER 90 PRL 64 1095	+Hinshaw, Ong, Snyder+	(Mark II Collab.)
ALBRECHT 89C PL B219 121	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
ALBRECHT 89G PL B229 304	+Glaeser, Harder, Krueger+	(ARGUS Collab.)
ALBRECHT 89J PL B229 175	+Glaeser, Harder+	(ARGUS Collab.)
ALBRECHT 89L PL B232 554	+Glaeser, Harder, Krueger, Nippe, Oest+	(ARGUS Collab.)
ARTUSO 89 PRL 62 2233	+Bebek, Berkelman, Blucher+	(CLEO Collab.)
AVERILL 89 PR D39 123	+Blockus, Brabson+	(HRS Collab.)
AVERY 89B PL B223 470	+Besson, Garren, Veltou+	(CLEO Collab.)
BEBEK 89 PRL 62 8	+Berkelman, Blucher+	(CLEO Collab.)
BORTOLETTO 89 PRL 62 2436	+Goldberg, Horwitz, Mestayer+	(CLEO Collab.)
BORTOLETTO 89B PRL 63 1667	+Goldberg, Horwitz, Mestayer+	(CLEO Collab.)
ALBRECHT 88F PL B209 119	+Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 88K PL B215 424	+Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 87C PL B185 218	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
ALBRECHT 87D PL B199 451	+Andam, Binder, Boeckmann+	(ARGUS Collab.)
ALBRECHT 87I PL B192 245	+Andam, Binder, Boeckmann+	(ARGUS Collab.)
ALBRECHT 87J PL B197 452	+Andam, Binder, Boeckmann+	(ARGUS Collab.)
AVERY 87 PL B183 429	+Besson, Bowcock, Giles+	(CLEO Collab.)
BEAN 87B PRL 58 183	+Bobbink, Brock, Engler+	(CLEO Collab.)
BEBEK 87 PR D36 1289	+Berkelman, Blucher, Cassel+	(CLEO Collab.)
ALAM 86 PR D34 3279	+Katayama, Kim, Sun+	(CLEO Collab.)
ALBRECHT 86F PL B182 95	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
PDG 86 PL 170B	+Aguliar-Benitez, Porter+	(CERN, CIT+)
CHEN 85 PR D31 2386	+Goldberg, Horwitz, Jawahery+	(CLEO Collab.)
HAAS 85 PRL 55 1248	+Hempstead, Jensen, Kagan+	(CLEO Collab.)
AVERY 84 PRL 53 1309	+Bebek, Berkelman, Cassel+	(CLEO Collab.)
GILES 84 PR D30 2279	+Hassard, Hempstead, Kinoshita+	(CLEO Collab.)
BEHREND 83 PRL 50 881	+Chadwick, Chauveau, Ganci+	(CLEO Collab.)

OTHER RELATED PAPERS

WINSTEIN 93 RMP 65 1113	+Wolfenstein	
"The Search for Direct CP Violation"		
BERKELMAN 91 ARNPS 41 1	+Stone	(CORN, SYRA)
"Decays of B Mesons"		
MILLER 90 MPL A5 2683		
"Recent Results in B Physics"		
SCHINDLER 88 High Energy Electron-Positron Physics 234		(SLAC)
Editors: A. Ali and P. Soeding, World Scientific, Singapore		
SCHUBERT 87 IHEP-HD/87-7		(HEIDH)
EPS Conference - Uppsala, Proc., Vol. 2, p. 791		

B^*

$$I(J^P) = \frac{1}{2}(1^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

B^* MASS

From mass difference below and the average of our B masses ($m_{B^\pm + m_{B^0}}$)/2.

VALUE (MeV)	DOCUMENT ID
5324.8 ± 2.1 OUR FIT	

$m_{B^*} - m_B$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
46.0 ± 0.6 OUR FIT				
46.0 ± 0.6 OUR AVERAGE				
46.4 ± 0.3 ± 0.8	1 AKERIB 91 CLE2	$e^+e^- \rightarrow \gamma X$		
45.6 ± 0.8	1 WU 91 CSB2	$e^+e^- \rightarrow \gamma X, \gamma \ell X$		
45.4 ± 1.0	2 LEE-FRANZINI 90 CSB2	$e^+e^- \rightarrow T(5S)$		
• • • We do not use the following data for averages, fits, limits, etc. • • •				
52.0 ± 2. ± 4.	1400	3 HAN 85 CUSB	$e^+e^- \rightarrow \gamma e X$	

1 These papers report E_γ in the B^* center of mass. The $m_{B^*} - m_B$ is 0.2 MeV higher.

$E_{cm} = 10.61-10.7$ GeV. Admixture of B^0 and B^+ mesons, but not B_s .
2 LEE-FRANZINI 90 value is for an admixture of B^0 and B^+ . They measure $46.7 \pm 0.4 \pm 0.2$ MeV for an admixture of $B^0, B^+,$ and B_s , and use the shape of the photon line to separate the above value.
3 HAN 85 is for $E_{cm} = 10.6-11.2$ GeV, giving an admixture of $B^0, B^+,$ and B_s .

B^* REFERENCES

AKERIB 91 PRL 67 1692	+Barish, Cown, Eigen, Stroynowski+	(CLEO Collab.)
WU 91 PL B273 177	+Franzini, Kanekal, Tuts+	(CUSB II Collab.)
LEE-FRANZINI 90 PRL 65 2947	+Heintz, Lovelock, Narain, Schamberger+	(CUSB II Collab.)
HAN 85 PRL 55 36	+Klopfenstein, Mageras+	(COLU, LSU, MPIM, STON)

BOTTOM, STRANGE MESONS

$$(B = \pm 1, S = \mp 1)$$

$$B_s^0 = s\bar{b}, \bar{B}_s^0 = \bar{s}b, \text{ similarly for } B_s^{*\pm}$$

B_s^0

$$I(J^P) = \frac{1}{2}(0^-)$$

I, J, P need confirmation. Quantum numbers shown are quark-model predictions.

B_s^0 MASS

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
5375 ± 6 OUR FIT				Error includes scale factor of 1.3.
5375 ± 5 OUR AVERAGE				Error includes scale factor of 1.2.
5374 ± 16 ± 2	3	ABREU 94D DLPH	$e^+e^- \rightarrow Z$	
5383.3 ± 4.5 ± 5.0	14	ABE 93F CDF	$p\bar{p}$ at 1.8 TeV	
5368.6 ± 5.6 ± 1.5	2	BUSKULIC 93G ALEP	$e^+e^- \rightarrow Z$	

$m_{B_s^0} - m_B$

Fit value is from B_s mass above and the average of our B masses ($m_{B^\pm + m_{B^0}}$)/2.

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
96 ± 6 OUR FIT				Error includes scale factor of 1.3.

• • • We do not use the following data for averages, fits, limits, etc. • • •
80 to 130 68 LEE-FRANZINI 90 CSB2 $e^+e^- \rightarrow T(5S)$

$m_{B_s^0} - m_{B_s^0}$

See the $B_s^0 - \bar{B}_s^0$ MIXING section near the end of these B_s^0 Listings.


Meson Full Listings

B_S^0

B_S^0 MEAN LIFE

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
$1.34^{+0.32}_{-0.27}$ OUR AVERAGE				Error includes scale factor of 1.4. See the Ideogram below.
0.96 ± 0.37	41	¹ ABREU	94E DLPH	$e^+ e^- \rightarrow Z$
$1.92^{+0.45}_{-0.35} \pm 0.04$	31	² BUSKULIC	94C ALEP	$e^+ e^- \rightarrow Z$
$1.13^{+0.35}_{-0.26} \pm 0.09$	22	² ACTON	93H OPAL	$e^+ e^- \rightarrow Z$

¹ ABREU 94E uses the flight-distance distribution of D_S vertices, ϕ -lepton vertices, and $D_S \mu$ vertices. This result includes the result of ABREU 92M.
² Measured using $D_S \ell^+$ events.



B_S^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 D_S^- anything	seen
Γ_2 $D_S^- \ell^+ \nu_\ell$ anything (ℓ means sum of e and μ)	seen
Γ_3 $D_S^- \pi^+$	seen
Γ_4 $J/\psi(1S)\phi$	seen
Γ_5 $\psi(2S)\phi$	seen

B_S^0 BRANCHING RATIOS

$\Gamma(D_S^- \text{ anything})/\Gamma_{\text{total}}$	Γ_1/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	147	³ ACTON	92N OPAL	$e^+ e^- \rightarrow Z$

³ ACTON 92N assumes that excess of $147 \pm 48 D_S^0 e^+ e^- \rightarrow Z$ over that expected from $B^0, B^+,$ and $c\bar{c}$ is all from B_S^0 decay. The product branching fraction is measured to be $f(\bar{b} \rightarrow B_S^0)B(B_S^0 \rightarrow D_S^- \text{ anything}) \times B(D_S^- \rightarrow \phi\pi^-) = (5.9 \pm 1.9 \pm 1.1) \times 10^{-3}$.

$\Gamma(D_S^- \ell^+ \nu_\ell \text{ anything})/\Gamma_{\text{total}}$ ℓ means sum of e and μ	Γ_2/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	7	⁴ ABREU	92M DLPH	$e^+ e^- \rightarrow Z$
seen	18	⁵ ACTON	92N OPAL	$e^+ e^- \rightarrow Z$
seen	27	⁶ BUSKULIC	92E ALEP	$e^+ e^- \rightarrow Z$

⁴ ABREU 92M measured muons only and obtained product branching ratio $B(Z \rightarrow b\bar{c}) \times B(\bar{b} \rightarrow B_S) \times B(B_S \rightarrow D_S \mu^+ \nu_\mu X) \times B(D_S \rightarrow \phi\pi) = (18 \pm 8) \times 10^{-5}$.

⁵ ACTON 92N is measured using $D_S \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. The product branching fraction measured is measured to be $f(\bar{b} \rightarrow B_S^0)B(B_S^0 \rightarrow D_S^- \ell^+ \nu_\ell \text{ anything}) \times B(D_S^- \rightarrow \phi\pi^-) = (3.9 \pm 1.1 \pm 0.8) \times 10^{-4}$.

⁶ BUSKULIC 92E is measured using $D_S \rightarrow \phi\pi^+$ and $K^*(892)^0 K^+$ events. They use $2.7 \pm 0.7\%$ for the $\phi\pi^+$ branching fraction. The average product branching fraction is measured to be $f(\bar{b} \rightarrow B_S^0)B(B_S^0 \rightarrow D_S^- \ell^+ \nu_\ell \text{ anything}) = 0.040 \pm 0.011^{+0.010}_{-0.012}$.

$\Gamma(D_S^- \pi^+)/\Gamma_{\text{total}}$	Γ_3/Γ			
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1	BUSKULIC	93G ALEP	$e^+ e^- \rightarrow Z$

$\Gamma(J/\psi(1S)\phi)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	14	⁷ ABE	93F CDF	$p\bar{p}$ at 1.8 TeV
seen	1	⁸ ACTON	92N OPAL	$e^+ e^- \rightarrow Z$

⁷ ABE 93F measured using $J/\psi(1S) \rightarrow \mu^+ \mu^-$ and $\phi \rightarrow K^+ K^-$.

⁸ In ACTON 92N, a limit on the product branching fraction measured is measured to be $f(\bar{b} \rightarrow B_S^0)B(B_S^0 \rightarrow J/\psi(1S)\phi) \leq 0.22 \times 10^{-2}$.

$\Gamma(\psi(2S)\phi)/\Gamma_{\text{total}}$ Γ_5/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	1	BUSKULIC	93G ALEP	$e^+ e^- \rightarrow Z$

B_S^0 - \bar{B}_S^0 MIXING

For a discussion of B_S^0 - \bar{B}_S^0 mixing see the note on " B^0 - \bar{B}^0 Mixing and CP Violation in B Decay" in the B^0 Full Listings above.

X_S

This B_S^0 - \bar{B}_S^0 mixing parameter measures the probability (integrated over time) that a produced B_S^0 (\bar{B}_S^0) decays as a \bar{B}_S^0 (B_S^0). The X_S values are derived from combining X_B values measured at high energy (see following datablock) with X_d values from the $\Upsilon(4S)$. Mixing violates $\Delta B \neq 2$ rule.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
0.62 ± 0.13		⁹ PDG	94 RVUE	
0.74 ± 0.27	95	¹⁰ ABREU	94F RVUE	$e^+ e^- \rightarrow Z$
0.46 ± 0.21		¹¹ ADEVA	92C RVUE	
0.53 ± 0.15		¹² ALBAJAR	91D RVUE	

⁹ From our X_B and X_d averages assuming $f_d = 0.391$ and $f_s = 0.117$.

¹⁰ From a combination of DELPHI (ABREU 94F), CLEO (ARTUSO 89), and ARGUS (ALBRECHT 92L). Estimated from ABREU 94F figure 7.

¹¹ From combination of L3 (ADEVA 92C), CLEO (ARTUSO 89), and ARGUS (ALBRECHT 92L). Corresponding limit is > 0.16 at 90%CL.

¹² From combination of UA1 (ALBAJAR 91D), CLEO (BEAN 87B), ARGUS (ALBRECHT 87I), ALEPH (DECAMP 91), and L3 (ADEVA 90P). Corresponding limits are > 0.23 at 95% CL and > 0.27 at 90% CL.

X_B at high energy

This is a B - \bar{B} mixing measurement for an admixture of B^0 and \bar{B}_S^0 at high energy.

$$X_B = f_d X_d + f_s X_s$$

where f_d and f_s are the production fractions of B^0 and \bar{B}_S^0 mesons relative to all b -flavored hadrons. Only the measurements at the Z and higher energy $p\bar{p}$ are averaged.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.133 ± 0.011 OUR AVERAGE					
$0.121^{+0.044}_{-0.040} \pm 0.017$		1665	¹³ ABREU	93C DLPH	$e^+ e^- \rightarrow Z$
$0.143^{+0.022}_{-0.021} \pm 0.007$			¹⁴ AKERS	93B OPAL	$e^+ e^- \rightarrow Z$
$0.121 \pm 0.017 \pm 0.006$			¹⁵ ADEVA	92C L3	$e^+ e^- \rightarrow Z$
0.129 ± 0.022			¹⁶ BUSKULIC	92B ALEP	$e^+ e^- \rightarrow Z$
$0.176 \pm 0.031 \pm 0.032$		1112	¹⁷ ABE	91G CDF	$p\bar{p}$ 1.8 TeV
$0.148 \pm 0.029 \pm 0.017$			¹⁸ ALBAJAR	91D UA1	$p\bar{p}$ 630 GeV
$0.144 \pm 0.014^{+0.017}_{-0.011}$			¹⁹ ABREU	94F DLPH	$e^+ e^- \rightarrow Z$
$0.145^{+0.041}_{-0.035} \pm 0.018$			²⁰ ACTON	92C OPAL	$e^+ e^- \rightarrow Z$
$0.132 \pm 0.22^{+0.015}_{-0.012}$		823	²¹ DECAMP	91 ALEP	$e^+ e^- \rightarrow Z$
$0.178^{+0.049}_{-0.040} \pm 0.020$			²² ADEVA	90P L3	$e^+ e^- \rightarrow Z$
$0.17^{+0.15}_{-0.08}$		23,24	WEIR	90 MRK2	$e^+ e^-$ 29 GeV
$0.21^{+0.29}_{-0.15}$			²³ BAND	88 MAC	$E_{\text{CM}}^e = 29$ GeV
> 0.02		90	²³ BAND	88 MAC	$E_{\text{CM}}^e = 29$ GeV
0.121 ± 0.047		23,25	ALBAJAR	87C UA1	Repl. by ALBAJAR 91D
< 0.12		90	^{23,26} SCHAAD	85 MRK2	$E_{\text{CM}}^e = 29$ GeV

• • • We do not use the following data for averages, fits, limits, etc. • • •

¹³ ABREU 93C data analyzed using $ee, e\mu, \mu\mu$ events.
¹⁴ AKERS 93B analysis performed using dilepton events.
¹⁵ ADEVA 92C uses electrons and muons.
¹⁶ BUSKULIC 92B uses a jet charge technique combined with electrons and muons.
¹⁷ ABE 91G measurement of X is done with $e\mu$ and $e\mu$ events.
¹⁸ ALBAJAR 91D measurement of X is done with dileptons.
¹⁹ ABREU 94F uses the average electric charge sum of the jets recoiling against a b -quark jet tagged by a high p_T muon. The result is for $\bar{X} = f_d X_d + 0.9 f_s X_s$.
²⁰ ACTON 92C uses electrons and muons. Superseded by AKERS 93B.
²¹ DECAMP 91 done with opposite and like-sign dileptons. Superseded by BUSKULIC 92B.
²² ADEVA 90P measurement uses $ee, \mu\mu, \mu e$ events from 118k events at the Z . Superseded by ADEVA 92C.
²³ These experiments are not in the average because the combination of B_S and B_d mesons which they see could differ from those at higher energy.
²⁴ The WEIR 90 measurement supersedes the limit obtained in SCHAAD 85. The 90% CL are 0.06 and 0.38.
²⁵ ALBAJAR 87C measured $X = (B^0 \rightarrow \mu^+ X)$ divided by the average production weighted semileptonic branching fraction for B hadrons at 546 and 630 GeV.
²⁶ Limit is average probability for hadron containing B quark to produce a positive lepton.

See key on page 1343

Meson Full Listings

 B_s^0, B_s^* , Top and Fourth Generation Hadrons

$$\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0}$$

$\Delta m_{B_s^0}$ is a measure of the $B_s^0 - \bar{B}_s^0$ oscillation frequency in time-dependent mixing experiments.

VALUE (10^{12} h s^{-1})	CL%	DOCUMENT ID	TECN	COMMENT
>1.8	95	27 BUSKULIC 94B ALEP	$e^+ e^- \rightarrow Z$	

²⁷ BUSKULIC 94B determines $\Delta m_{B_s^0}$ from the time dependence of B mixing using dileptons.

$$x_s = \Delta m_{B_s^0} / \Gamma_{B_s^0}$$

This section combines the results from the previous two sections.

Time integrated mixing measurement of X determine this quantity directly via

$$\frac{\Delta m_{B_s^0}}{\Gamma_{B_s^0}} = \left(\frac{X}{0.5 - X} \right)^{1/2}$$

while time-dependent mixing measurements determine $\Delta m_{B_s^0} = m_{B_{sH}^0} - m_{B_{sL}^0}$ which are combined with $\tau_{B_s^0}$ to give

$$\frac{\Delta m_{B_s^0}}{\Gamma_{B_s^0}} = \frac{(m_{B_{sH}^0} - m_{B_{sL}^0}) \tau_{B_s^0}}{\hbar}$$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
>2.0	95	28 BUSKULIC 94B ALEP	$e^+ e^- \rightarrow Z$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

>1.5	29	PDG 94 RVUE		
------	----	-------------	--	--

²⁸ BUSKULIC 94B is their $\Delta m_{B_s^0}$ measurement combined with $\tau_{B_s^0} = (1.26^{+0.22}_{-0.17}) \text{ ps}$.

²⁹ From PDG 94 X_s value.

 B_s^0 REFERENCES

ABREU 94D PL B324 500	+Adam, Adye, Agasi, Aleksan+	(DELPHI Collab.)
ABREU 94E ZPHY C61 407	+Adam, Adye, Agasi, Aleksan+	(DELPHI Collab.)
Also 92M PL B289 199	Abreu, Adam, Adye, Agasi, Alekseev+	(DELPHI Collab.)
ABREU 94F PL B322 459	+Adam, Adye, Agasi, Ajinenko+	(DELPHI Collab.)
BUSKULIC 94B PL B322 441	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
BUSKULIC 94C PL B322 275	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
PDG 94 PR D50 1173	Montanet+ (CERN, LBL, BOST, IFIC+)	
ABE 93F PRL 71 1685	+Albrow, Amidei, Anway-Wiese+	(CDF Collab.)
ABREU 93C PL B301 145	+Adam, Adye, Agasi, Aleksan+	(DELPHI Collab.)
ACTON 93H PL B312 501	+Akers, Alexander, Allison, Anderson+	(OPAL Collab.)
AKERS 93B ZPHY C60 199	+Alexander, Allison, Anderson, Arcelli+	(OPAL Collab.)
BUSKULIC 93G PL B311 425	+De Bonis, Decamp, Ghez, Goy, Lees+	(ALEPH Collab.)
ABREU 92M PL B289 199	+Adam, Adye, Agasi, Alekseev+	(DELPHI Collab.)
ACTON 92C PL B276 379	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ACTON 92N PL B295 357	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
ADEVA 92P PL B288 395	+Adriani, Aguilar-Benitez, Ahlen+	(L3 Collab.)
ALBRECHT 92L ZPHY C55 357	+Ehrlichmann, Hamacher, Krueger, Nau+	(ARGUS Collab.)
BUSKULIC 92B PL B284 177	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
BUSKULIC 92E PL B294 145	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
ABE 91G PRL 67 3351	+Amidei, Apollinari, Atac, Auchincloss+	(CDF Collab.)
ALBAJAR 91D PL B262 171	+Albrow, Altkofer, Ankoviak, Apsimon+	(UA1 Collab.)
DECAMP 91 PL B258 236	+Deschizeaux, Goy, Lees, Minard+	(ALEPH Collab.)
ADEVA 90P PL B252 703	+Adriani, Aguilar-Benitez, Akbari, Alcaraz+	(L3 Collab.)
LEE-FRANZINI 90 PRL 65 2947	+Heintz, Lovelock, Narain, Schamberger+	(CUSB II Collab.)
WEIR 90 PL B240 289	+Abrams, Adolphsen, Alexander, Alvarez+	(Mark II Collab.)
ARTUSO 89 PRL 62 2233	+Bebek, Berkelman, Blucher+	(CLEO Collab.)
BAND 88 PL B200 221	+Camporesi, Chadwick+	(MAC Collab.)
ALBAJAR 87C PL B186 247	+Albrow, Altkofer, Arnison+	(UA1 Collab.)
ALBRECHT 87I PL B192 245	+Andam, Binder, Boeckmann+	(ARGUS Collab.)
BEAN 87B PRL 58 18 183	+Bobink, Brock, Engler+	(CLEO Collab.)
SCHAAD 85 PL 160B 188	+Nelson, Abrams, Amidei+	(Mark II Collab.)

OTHER RELATED PAPERS

ALI 93 JPG 19 1069	+London (DESY, MONT)
"Prospects for measuring the $B_s^0 - \bar{B}_s^0$ mixing ratio x_s "	

 B_s^*

$$I(J^P) = ?(??)$$

I, J, P need confirmation.

OMITTED FROM SUMMARY TABLE

 B_s^* MASS

From mass difference below and the B_s^0 mass.

VALUE (MeV)	DOCUMENT ID
5422 ± 6 OUR FIT	Error includes scale factor of 1.2.

$$m_{B_s^*} - m_{B_s}$$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
47.0 ± 2.6 OUR FIT			

¹ LEE-FRANZINI 90 CSB2 $e^+ e^- \rightarrow T(5S)$

¹ LEE-FRANZINI 90 measure $46.7 \pm 0.4 \pm 0.2 \text{ MeV}$ for an admixture of B^0, B^+ , and B_s . They use the shape of the photon line to separate the above value for B_s .

 B_s^* REFERENCES

LEE-FRANZINI 90 PRL 65 2947	+Heintz, Lovelock, Narain, Schamberger+ (CUSB II Collab.)
-----------------------------	---

HEAVY QUARK SEARCHES

Searches for Top and Fourth Generation Hadrons

See the sections "Searches for t Quark" and "Searches for b' (4^{th} Generation) Quark" at the end of the QUARKS section.

Meson Full Listings

Charmonium, $\eta_c(1S) = \eta_c(2980)$

$c\bar{c}$ MESONS

$\eta_c(1S)$
or $\eta_c(2980)$

$$I^G(J^{PC}) = 0^+(0^{-+})$$

Observed in the inclusive γ spectrum generated from $\psi(2S)$ decay, therefore $C = +$. From the 4π decay $G = +$, therefore $I = 0$. From angular distribution in $J/\psi(1S) \rightarrow \eta_c \gamma, \eta_c \rightarrow \phi \phi, J^P = 0^-$ (BALTRUSAITIS 84).

$\eta_c(1S)$ MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2978.8 ± 1.9 OUR AVERAGE		Error includes scale factor of 1.8. See the ideogram below.		
2974.4 ± 1.9		¹ BISELLO	91 DM2	$J/\psi \rightarrow \eta_c \gamma$
2969 ± 4 ± 4	80	BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
2982.6 ^{+2.7} _{-2.3}	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma \gamma$
2980.2 ± 1.6		¹ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c \gamma$
2984 ± 2.3 ± 4.0		GAISER	86 CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
2982 ± 8	18	² HIMEL	80B MRK2	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2956 ± 12 ± 12		BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$
2976 ± 8		³ BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi \gamma$
2980 ± 9		² PARTRIDGE	80B CBAL	$e^+ e^-$

¹ Average of several decay modes.

² Mass adjusted by us to correspond to $J/\psi(1S)$ mass = 3097 MeV.

³ $\eta_c \rightarrow \phi \phi$.

WEIGHTED AVERAGE
2978.8 ± 1.9 (Error scaled by 1.8)




$\eta_c(1S)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
10.3^{+3.8}_{-3.4} OUR AVERAGE					
7.0 ^{+7.5} _{-7.0}		12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma \gamma$
10.1 ^{+33.0} _{-8.2}		23 ±	⁴ BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \gamma \rho \bar{p}$
11.5 ± 4.5		11	GAISER	86 CBAL	$J/\psi \rightarrow \gamma X, \psi(2S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<40	90	18	HIMEL	80B MRK2	$e^+ e^-$
<20	90		PARTRIDGE	80B CBAL	$e^+ e^-$

⁴ Positive and negative errors correspond to 90% confidence level.

THE CHARMONIUM SYSTEM



The current state of knowledge of the charmonium system and transitions, as interpreted by the charmonium model. Uncertain states and transitions are indicated by dashed lines. The notation γ^* refers to decay processes involving intermediate virtual photons, including decays to $e^+ e^-$ and $\mu^+ \mu^-$.

$\eta_c(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Decays involving hadronic resonances		
Γ_1 $\eta'(958)\pi\pi$	(4.1 ± 1.7) %	
Γ_2 $\rho\rho$	(2.6 ± 0.9) %	
Γ_3 $K^*(892)^0 K^- \pi^+ + c.c.$	(2.0 ± 0.7) %	
Γ_4 $K^*(892)\bar{K}^*(892)$	(8.5 ± 3.1) × 10 ⁻³	
Γ_5 $\phi\phi$	(7.1 ± 2.8) × 10 ⁻³	
Γ_6 $a_0(980)\pi$	< 2 %	90%
Γ_7 $a_2(1320)\pi$	< 2 %	90%
Γ_8 $K^*(892)\bar{K} + c.c.$	< 1.28 %	90%
Γ_9 $f_2(1270)\eta$	< 1.1 %	90%
Γ_{10} $\omega\omega$	< 3.1 × 10 ⁻³	90%
Decays into stable hadrons		
Γ_{11} $K\bar{K}\pi$	(6.6 ± 1.8) %	
Γ_{12} $\eta\pi\pi$	(4.9 ± 1.8) %	
Γ_{13} $\pi^+\pi^-K^+K^-$	(2.0 $^{+0.7}_{-0.6}$) %	
Γ_{14} $2(\pi^+\pi^-)$	(1.2 ± 0.4) %	
Γ_{15} $\rho\bar{\rho}$	(1.2 ± 0.4) × 10 ⁻³	
Γ_{16} $K\bar{K}\eta$	< 3.1 %	90%
Γ_{17} $\pi^+\pi^-\rho\bar{\rho}$	< 1.2 %	90%
Γ_{18} $\Lambda\bar{\Lambda}$	< 2 × 10 ⁻³	90%
Radiative decays		
Γ_{19} $\gamma\gamma$	(6 $^{+6}_{-5}$) × 10 ⁻⁴	

$\eta_c(1S)$ PARTIAL WIDTHS

VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{19}
7.0^{+2.0}_{-1.7} OUR AVERAGE						
8.0 ± 2.3 ± 2.4		17 ± 5	ADRIANI	93N L3	$e^+e^- \rightarrow e^+e^-\eta_c$	
5.9 $^{+2.1}_{-1.8}$ ± 1.9			CHEN	90B CLEO	$e^+e^- \rightarrow e^+e^-\eta_c$	
6.4 $^{+5.0}_{-3.4}$			AIHARA	88D TPC	$e^+e^- \rightarrow e^+e^-X$	
28 ± 15			5 BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 11	90		BLINOV	86 MD1	$e^+e^- \rightarrow e^+e^-X$	
5 Re-evaluated by AIHARA 88D.						

$\eta_c(1S)$ $\Gamma(l)\Gamma(\gamma\gamma)/\Gamma(\text{total})$

$\Gamma(K\bar{K}\pi) \times \Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	VALUE (keV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	$\Gamma_{11}\Gamma_{19}/\Gamma$
1.2 ± 0.4 OUR AVERAGE							
	1.06 ± 0.41 ± 0.27		11 ± 4	BRAUNSCH...	89 TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$	
	1.5 $^{+0.60}_{-0.45}$ ± 0.3		7	6 BERGER	86 PLUT	$\gamma\gamma \rightarrow K\bar{K}\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
	< 0.63	95		6 BEHREND	89 CELL	$\gamma\gamma \rightarrow K_S^0 K^\pm \pi^\mp$	
	< 4.4	95		ALTHOFF	85B TASS	$\gamma\gamma \rightarrow K\bar{K}\pi$	
6 $K^\pm K_S^0 \pi^\mp$ corrected to $K\bar{K}\pi$ by factor 3.							

$\eta_c(1S)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(\eta'(958)\pi\pi)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
	0.041 ± 0.017	14 ± 4	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
$\Gamma(\rho\rho)/\Gamma_{\text{total}}$						
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
26 ± 9 OUR EVALUATION						
25 ± 8 OUR AVERAGE						
	26.0 ± 2.4 ± 8.8	113 ± 11	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^0\rho^0$	
	23.6 ± 10.6 ± 8.2	32 ± 14	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho^+\rho^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	< 140	90	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
$\Gamma(K^*(892)^0 K^- \pi^+ + c.c.)/\Gamma_{\text{total}}$						
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ	
0.02 ± 0.007	63 ± 10	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$		

$\Gamma(K^*(892)\bar{K}^*(892))/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
85 ± 31 OUR AVERAGE					
82 ± 28 ± 27	14 ± 5	7 BISELLO	91 DM2	$e^+e^- \rightarrow \gamma K^+ K^- \pi^+ \pi^-$	
90 ± 50	9 ± 4	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	

$\Gamma(K^*(892)\bar{K} + c.c.)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
< 0.0128 OUR EVALUATION					
< 0.0132	90	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^\pm K^\pm \pi^\mp$	
	90	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^\mp K^- \pi^0$	

$\Gamma(\phi\phi)/\Gamma_{\text{total}}$

VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
71 ± 28 OUR EVALUATION					
(Treating systematic errors as correlated.)					
71 ± 22 OUR AVERAGE					
74 ± 18 ± 24	80	7 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
67 ± 21 ± 24		7 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

31 ± 7 ± 10	19 ± 5	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$	
30 $^{+18}_{-12}$ ± 10	5 ± 3	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_L^0$	

$\Gamma(a_0(980)\pi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
< 0.02	90	7,8 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	

$\Gamma(a_2(1320)\pi)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
< 0.02	90	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	

$\Gamma(f_2(1270)\eta)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
< 0.011	90	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	

$\Gamma(\omega\omega)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
< 0.0031	90	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.0063		7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\omega\omega$	

$\Gamma(K\bar{K}\pi)/\Gamma_{\text{total}}$

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
0.066 ± 0.018 OUR EVALUATION						
(Treating systematic errors as correlated.)						
0.063 ± 0.013 OUR AVERAGE						
0.0690 ± 0.0144 ± 0.0234		33 ± 7	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^+ K^- \pi^0$	
0.0543 ± 0.0096 ± 0.0180		68 ± 10	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma K^\pm \pi^\mp K_S^0$	
0.061 ± 0.022		95 ± 18	7,9 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
0.161 $^{+0.092}_{-0.073}$			10 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 0.107	90		7 PARTRIDGE	80B CBAL	$J/\psi \rightarrow \eta_c\gamma$	

$\Gamma(\eta\pi\pi)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
0.049 ± 0.018 OUR EVALUATION					
0.047 ± 0.015 OUR AVERAGE					
0.054 ± 0.020	75 ± 11	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
0.037 ± 0.013 ± 0.020	18	7 PARTRIDGE	80B CBAL	$J/\psi \rightarrow \eta\pi^+\pi^-\gamma$	

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
0.020 $^{+0.007}_{-0.006}$ OUR AVERAGE					
0.021 ± 0.007	110 ± 17	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
0.014 $^{+0.022}_{-0.009}$		10 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$	

$\Gamma(2\pi^+\pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
0.012 ± 0.004 OUR EVALUATION					
0.0120 ± 0.0031 OUR AVERAGE					
0.0105 ± 0.0017 ± 0.0034	137 ± 23	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma 2\pi^+ 2\pi^-$	
0.013 ± 0.006	25 ± 9	7 BALTRUSAIT..86	MRK3	$J/\psi \rightarrow \eta_c\gamma$	
0.020 $^{+0.015}_{-0.010}$		10 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$	

Meson Full Listings

$$\eta_c(1S) = \eta_c(2980), J/\psi(1S) = J/\psi(3097)$$

$\Gamma(\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{15}/Γ

VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT
12 ± 4 OUR AVERAGE				
10 ± 3 ± 4	18 ± 6	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\rho\bar{\rho}$
11 ± 6	23 ± 11	7 BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \eta_c\gamma$
29 ± 29 -15		10 HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$

$\Gamma(K\bar{K}\eta)/\Gamma_{\text{total}}$ Γ_{16}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.031	90	7 BALTRUSAIT...86	MRK3	$J/\psi \rightarrow \eta_c\gamma$

$\Gamma(\pi^+\pi^-\rho\bar{\rho})/\Gamma_{\text{total}}$ Γ_{17}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.012	90	HIMEL	80B MRK2	$\psi(2S) \rightarrow \eta_c\gamma$

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{\text{total}}$ Γ_{18}/Γ

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.002	90	7 BISELLO	91 DM2	$e^+e^- \rightarrow \gamma\Lambda\bar{\Lambda}$

$\Gamma_f/\Gamma_{\text{total}} \ln \rho\bar{\rho} \rightarrow \eta_c(1S) \rightarrow \phi\phi$ $\Gamma_{15}\Gamma_5/\Gamma^2$

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
4.0 ± 3.5 -3.2	BAGLIN	89 SPEC	$\bar{p}p \rightarrow K^+K^-K^+K^-$

⁷The quoted branching ratios use $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$. Where relevant, the error in this branching ratio is treated as a common systematic in computing averages.

⁸We are assuming $B(a_0(980) \rightarrow \eta\pi) > 0.5$.

⁹Average from $K^+K^-\pi^0$ and $K^\pm K^0 \pi^\pm$ decay channels.

¹⁰Estimated using $B(\psi(2S) \rightarrow \gamma\eta_c(1S)) = 0.0028 \pm 0.0006$.

RADIATIVE DECAYS

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$ Γ_{19}/Γ

VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT
6 ± 4 -3 ± 4		BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 9	90	7 BISELLO	91 DM2	$J/\psi \rightarrow \gamma\gamma\gamma$
< 18	90	11 BLOOM	83 CBAL	$J/\psi \rightarrow \eta_c\gamma$

¹¹Using $B(J/\psi(1S) \rightarrow \gamma\eta_c(1S)) = 0.0127 \pm 0.0036$.

$\Gamma_f/\Gamma_{\text{total}} \ln \rho\bar{\rho} \rightarrow \eta_c(1S) \rightarrow \gamma\gamma$ $\Gamma_{15}\Gamma_{19}/\Gamma^2$

VALUE (units 10^{-6})	EVTs	DOCUMENT ID	TECN	COMMENT
0.66 ± 0.42 -0.31	12	BAGLIN	87B SPEC	$\bar{p}p \rightarrow \gamma\gamma$

 $\eta_c(1S)$ REFERENCES

ADRIANI	93N	PL B318 575	+Aguiar-Benitez, Ahlen+	(L3 Collab.)
BISELLO	91	NP B350 1	+Busetto+	(DM2 Collab.)
BAI	90B	PRL 65 1309	+Blaylock+	(Mark III Collab.)
CHEN	90B	PL B243 169	+McIlwain+	(CLEO Collab.)
BAGLIN	89	PL B231 557	+Baird, Bassompierre	(R704 Collab.)
BEHREND	89	ZPHY C42 367	+Criegee+	(CELLO Collab.)
BRAUNSCH... 89	ZPHY C41 533		+Braunschweig, Bock+	(TASSO Collab.)
AIHARA	88D	PRL 60 2355	+Aiston-Garnjost+	(TPC Collab.)
BAGLIN	87B	PL B187 191	+Baird, Bassompierre, Borreani+	(R704 Collab.)
BALTRUSAIT... 86	PR D33 629		+Baltrusaitis, Coffman, Hauser+	(Mark III Collab.)
BERGER	86	PL 167B 120	+Genzel, Lackas, Pielorz+	(PLUTO Collab.)
BLINOV	86		+Blinov, Bondar, Bukin+	(NOVO)
Proc. XXIII Int.	HEP Conf., Berkeley, CA	(1986); World Scientific, Singapore, 1987, ed. S.C. Loken		
GAISER	86	PR D34 711	+Bloom, Bulos, Godfrey+	(Crystal Ball Collab.)
ALTHOFF	85B	ZPHY C29 189	+Braunschweig, Kirschfink+	(TASSO Collab.)
BALTRUSAIT... 84	PRL 52 2126		+Baltrusaitis+	(CIT, UCSC, ILL, SLAC, WASH) JP
BLOOM	83	ARNS 33 143	+Peck	(SLAC, CIT)
HIMEL	80B	PRL 45 1146	+Trilling, Abrams, Alam+	(SLAC, LBL, UCB)
PARTRIDGE	80B	PRL 45 1150	+Peck+	(CIT, HARV, PRIN, STAN, SLAC)

OTHER RELATED PAPERS

ARMSTRONG	89	PL B221 216	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)
BLOOM	79	Fermilab Symp. 92	(CIT, HARV, PRIN, SLAC, STAN)

$J/\psi(1S)$
or $J/\psi(3097)$

$$JG(J^{PC}) = 0^-(1^{--})$$

J/ψ(1S) MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
3096.88 ± 0.04 OUR AVERAGE				
3096.87 ± 0.03 ± 0.03		ARMSTRONG	93B SPEC	$\bar{p}p \rightarrow e^+e^-$
3096.95 ± 0.1 ± 0.3	193	BAGLIN	87 SPEC	$\bar{p}p \rightarrow e^+e^-X$
3098.4 ± 2.0	38k	LEMOIGNE	82 GOLI	190 GeV $\pi^-Be \rightarrow 2\mu$
3096.93 ± 0.09	502	ZHOLENTZ	80 REDE	e^+e^-
3097.0 ± 1		1 BRANDELIK	79C DASP	e^+e^-

¹From a simultaneous fit to e^+e^- , $\mu^+\mu^-$ and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.

J/ψ(1S) WIDTH

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
88 ± 5 OUR AVERAGE			
99 ± 12 ± 6	ARMSTRONG	93B SPEC	$\bar{p}p \rightarrow e^+e^-$
85.5 ± 6.1 -5.8	2 HSUEH	92 RVUE	See Υ mini-review

²Using data from COFFMAN 92, BALDINI-CELIO 75, BOYARSKI 75, ESPOSITO 75b, BRANDELIK 79c.

J/ψ(1S) DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(86.0 ± 2.0) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(17.0 ± 2.0) %	
Γ_3 e^+e^-	(5.99 ± 0.25) %	
Γ_4 $\mu^+\mu^-$	(5.97 ± 0.25) %	S=1.1

Decays involving hadronic resonances

Γ_5 $\rho\pi$	(1.28 ± 0.10) %	
Γ_6 $\rho^0\pi^0$	(4.2 ± 0.5) × 10 ⁻³	
Γ_7 $a_2(1320)\rho$	(1.09 ± 0.22) %	
Γ_8 $\omega\pi^+\pi^-\pi^-\pi^-$	(8.5 ± 3.4) × 10 ⁻³	
Γ_9 $\omega\pi^+\pi^-$	(7.2 ± 1.0) × 10 ⁻³	
Γ_{10} $K^*(892)^0\bar{K}_2^*(1430)^0 + c.c.$	(6.7 ± 2.6) × 10 ⁻³	
Γ_{11} $\omega K^*(892)\bar{K} + c.c.$	(5.3 ± 2.0) × 10 ⁻³	
Γ_{12} $\omega f_2(1270)$	(4.3 ± 0.6) × 10 ⁻³	
Γ_{13} $K^+\bar{K}^*(892)^- + c.c.$	(5.0 ± 0.4) × 10 ⁻³	
Γ_{14} $K^0\bar{K}^*(892)^0 + c.c.$	(4.2 ± 0.4) × 10 ⁻³	
Γ_{15} $\omega\pi^0\pi^0$	(3.4 ± 0.8) × 10 ⁻³	
Γ_{16} $b_1(1235)^\pm\pi^\mp$	[a] (3.0 ± 0.5) × 10 ⁻³	
Γ_{17} $\omega K^\pm K_S^0\pi^\mp$	[a] (3.0 ± 0.7) × 10 ⁻³	
Γ_{18} $b_1(1235)^0\pi^0$	(2.3 ± 0.6) × 10 ⁻³	
Γ_{19} $\phi K^*(892)\bar{K} + c.c.$	(2.04 ± 0.28) × 10 ⁻³	
Γ_{20} $\omega K\bar{K}$	(1.9 ± 0.4) × 10 ⁻³	
Γ_{21} $\omega f_J(1710) \rightarrow \omega K\bar{K}$	(4.8 ± 1.1) × 10 ⁻⁴	
Γ_{22} $\phi 2(\pi^+\pi^-)$	(1.60 ± 0.32) × 10 ⁻³	
Γ_{23} $\Delta(1232)^+\bar{p}\pi^-$	(1.6 ± 0.5) × 10 ⁻³	
Γ_{24} $\omega\eta$	(1.58 ± 0.16) × 10 ⁻³	
Γ_{25} $\phi K\bar{K}$	(1.48 ± 0.22) × 10 ⁻³	
Γ_{26} $\phi f_J(1710) \rightarrow \phi K\bar{K}$	(3.6 ± 0.6) × 10 ⁻⁴	
Γ_{27} $\rho\bar{\rho}\omega$	(1.30 ± 0.25) × 10 ⁻³	S=1.3
Γ_{28} $\Delta(1232)^+\bar{\Delta}(1232)^--$	(1.10 ± 0.29) × 10 ⁻³	
Γ_{29} $\Sigma(1385)^-\bar{\Sigma}(1385)^+ (or c.c.)$	[a] (1.03 ± 0.13) × 10 ⁻³	
Γ_{30} $\rho\bar{\rho}\eta'(958)$	(9 ± 4) × 10 ⁻⁴	S=1.7
Γ_{31} $\phi f_2'(1525)$	(8 ± 4) × 10 ⁻⁴	S=2.7
Γ_{32} $\phi\pi^+\pi^-$	(8.0 ± 1.2) × 10 ⁻⁴	
Γ_{33} $\phi K^\pm K_S^0\pi^\mp$	[a] (7.2 ± 0.9) × 10 ⁻⁴	
Γ_{34} $\omega f_1(1420)$	(6.8 ± 2.4) × 10 ⁻⁴	
Γ_{35} $\phi\eta$	(6.5 ± 0.7) × 10 ⁻⁴	
Γ_{36} $\Xi(1530)^-\Xi^+$	(5.9 ± 1.5) × 10 ⁻⁴	
Γ_{37} $\rho K^-\bar{\Sigma}(1385)^0$	(5.1 ± 3.2) × 10 ⁻⁴	
Γ_{38} $\omega\pi^0$	(4.2 ± 0.6) × 10 ⁻⁴	S=1.4
Γ_{39} $\phi\eta'(958)$	(3.3 ± 0.4) × 10 ⁻⁴	
Γ_{40} $\phi f_0(980)$	(3.2 ± 0.9) × 10 ⁻⁴	S=1.9
Γ_{41} $\Xi(1530)^0\Xi^0$	(3.2 ± 1.4) × 10 ⁻⁴	
Γ_{42} $\Sigma(1385)^-\bar{\Sigma}^+ (or c.c.)$	[a] (3.1 ± 0.5) × 10 ⁻⁴	

See key on page 1343

Meson Full Listings

$$J/\psi(1S) = J/\psi(3097)$$

Γ_{43}	$\phi f_1(1285)$	$(2.6 \pm 0.5) \times 10^{-4}$	S=1.1
Γ_{44}	$\rho\eta$	$(1.93 \pm 0.23) \times 10^{-4}$	
Γ_{45}	$\omega\eta'(958)$	$(1.67 \pm 0.25) \times 10^{-4}$	
Γ_{46}	$\omega f_0(980)$	$(1.4 \pm 0.5) \times 10^{-4}$	
Γ_{47}	$\rho\eta'(958)$	$(1.05 \pm 0.18) \times 10^{-4}$	
Γ_{48}	$\rho\bar{\rho}\phi$	$(4.5 \pm 1.5) \times 10^{-5}$	
Γ_{49}	$a_2(1320)^\pm \pi^\mp$	[a] $< 4.3 \times 10^{-3}$	CL=90%
Γ_{50}	$K\bar{K}_2^*(1430) + c.c.$	$< 4.0 \times 10^{-3}$	CL=90%
Γ_{51}	$K_S^*(1430)^0 \bar{K}_2^*(1430)^0$	$< 2.9 \times 10^{-3}$	CL=90%
Γ_{52}	$K^*(892)^0 \bar{K}^*(892)^0$	$< 5 \times 10^{-4}$	CL=90%
Γ_{53}	$\phi f_2(1270)$	$< 3.7 \times 10^{-4}$	CL=90%
Γ_{54}	$\rho\bar{\rho}\rho$	$< 3.1 \times 10^{-4}$	CL=90%
Γ_{55}	$\phi\eta'(1440) \rightarrow \phi\eta\pi\pi$	$< 2.5 \times 10^{-4}$	CL=90%
Γ_{56}	$\omega f_2'(1525)$	$< 2.2 \times 10^{-4}$	CL=90%
Γ_{57}	$\Sigma(1385)^0 \bar{\Lambda}$	$< 2 \times 10^{-4}$	CL=90%
Γ_{58}	$\Delta(1232)^+ \bar{p}$	$< 1 \times 10^{-4}$	CL=90%
Γ_{59}	$\Sigma^0 \bar{\Lambda}$	$< 9 \times 10^{-5}$	CL=90%
Γ_{60}	$\phi\pi^0$	$< 6.8 \times 10^{-6}$	CL=90%

Decays into stable hadrons

Γ_{61}	$2(\pi^+\pi^-\pi^0)$	$(3.37 \pm 0.26) \%$	
Γ_{62}	$3(\pi^+\pi^-\pi^0)$	$(2.9 \pm 0.6) \%$	
Γ_{63}	$\pi^+\pi^-\pi^0$	$(1.50 \pm 0.20) \%$	
Γ_{64}	$\pi^+\pi^-\pi^0 K^+ K^-$	$(1.20 \pm 0.30) \%$	
Γ_{65}	$4(\pi^+\pi^-\pi^0)$	$(9.0 \pm 3.0) \times 10^{-3}$	
Γ_{66}	$\pi^+\pi^- K^+ K^-$	$(7.2 \pm 2.3) \times 10^{-3}$	
Γ_{67}	$K\bar{K}\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{68}	$\rho\bar{\rho}\pi^+\pi^-$	$(6.0 \pm 0.5) \times 10^{-3}$	S=1.3
Γ_{69}	$2(\pi^+\pi^-)$	$(4.0 \pm 1.0) \times 10^{-3}$	
Γ_{70}	$3(\pi^+\pi^-)$	$(4.0 \pm 2.0) \times 10^{-3}$	
Γ_{71}	$n\bar{n}\pi^+\pi^-$	$(4 \pm 4) \times 10^{-3}$	
Γ_{72}	$\Sigma\bar{\Sigma}$	$(3.8 \pm 0.5) \times 10^{-3}$	
Γ_{73}	$2(\pi^+\pi^-) K^+ K^-$	$(3.1 \pm 1.3) \times 10^{-3}$	
Γ_{74}	$\rho\bar{\rho}\pi^+\pi^-\pi^0$	[b] $(2.3 \pm 0.9) \times 10^{-3}$	S=1.9
Γ_{75}	$\rho\bar{\rho}$	$(2.14 \pm 0.10) \times 10^{-3}$	
Γ_{76}	$\rho\bar{\rho}\eta$	$(2.09 \pm 0.18) \times 10^{-3}$	
Γ_{77}	$\rho\bar{\rho}\pi^-$	$(2.00 \pm 0.10) \times 10^{-3}$	
Γ_{78}	$n\bar{n}$	$(1.9 \pm 0.5) \times 10^{-3}$	
Γ_{79}	$\Xi\bar{\Xi}$	$(1.8 \pm 0.4) \times 10^{-3}$	S=1.8
Γ_{80}	$\Lambda\bar{\Lambda}$	$(1.35 \pm 0.14) \times 10^{-3}$	S=1.2
Γ_{81}	$\rho\bar{\rho}\pi^0$	$(1.09 \pm 0.09) \times 10^{-3}$	
Γ_{82}	$\Lambda\bar{\Sigma}^-\pi^+$ (or c.c.)	[a] $(1.06 \pm 0.12) \times 10^{-3}$	
Γ_{83}	$\rho K^-\bar{\Lambda}$	$(8.9 \pm 1.6) \times 10^{-4}$	
Γ_{84}	$2(K^+ K^-)$	$(7.0 \pm 3.0) \times 10^{-4}$	
Γ_{85}	$\rho K^-\bar{\Sigma}^0$	$(2.9 \pm 0.8) \times 10^{-4}$	
Γ_{86}	$K^+ K^-$	$(2.37 \pm 0.31) \times 10^{-4}$	
Γ_{87}	$\Lambda\bar{\Lambda}\pi^0$	$(2.2 \pm 0.7) \times 10^{-4}$	
Γ_{88}	$\pi^+\pi^-$	$(1.47 \pm 0.23) \times 10^{-4}$	
Γ_{89}	$K_S^0 K_L^0$	$(1.08 \pm 0.14) \times 10^{-4}$	
Γ_{90}	$\Lambda\bar{\Sigma}^+ + c.c.$	$< 1.5 \times 10^{-4}$	CL=90%
Γ_{91}	$K_S^0 K_S^0$	$< 5.2 \times 10^{-6}$	CL=90%

Radiative decays

Γ_{92}	$\gamma\eta_c(1S)$	$(1.3 \pm 0.4) \%$	
Γ_{93}	$\gamma\pi^+\pi^-\pi^0$	$(8.3 \pm 3.1) \times 10^{-3}$	
Γ_{94}	$\gamma\eta\pi\pi$	$(6.1 \pm 1.0) \times 10^{-3}$	
Γ_{95}	$\gamma\eta(1440) \rightarrow \gamma K\bar{K}\pi$	[c] $(9.1 \pm 1.8) \times 10^{-4}$	
Γ_{96}	$\gamma\eta(1440) \rightarrow \gamma\gamma\rho^0$	$(6.4 \pm 1.4) \times 10^{-5}$	
Γ_{97}	$\gamma\rho\rho$	$(4.5 \pm 0.8) \times 10^{-3}$	
Γ_{98}	$\gamma\eta'(958)$	$(4.31 \pm 0.30) \times 10^{-3}$	
Γ_{99}	$\gamma 2\pi^+ 2\pi^-$	$(2.8 \pm 0.5) \times 10^{-3}$	S=1.9
Γ_{100}	$\gamma f_4(2050)$	$(2.7 \pm 0.7) \times 10^{-3}$	
Γ_{101}	$\gamma\omega\omega$	$(1.59 \pm 0.33) \times 10^{-3}$	
Γ_{102}	$\gamma\eta(1440) \rightarrow \gamma\rho^0\rho^0$	$(1.4 \pm 0.4) \times 10^{-3}$	
Γ_{103}	$\gamma f_2(1270)$	$(1.38 \pm 0.14) \times 10^{-3}$	
Γ_{104}	$\gamma f_J(1710) \rightarrow \gamma K\bar{K}$	$(9.7 \pm 1.2) \times 10^{-4}$	
Γ_{105}	$\gamma\eta$	$(8.6 \pm 0.8) \times 10^{-4}$	
Γ_{106}	$\gamma f_1(1420) \rightarrow \gamma K\bar{K}\pi$	$(8.3 \pm 1.5) \times 10^{-4}$	
Γ_{107}	$\gamma f_1(1285)$	$(6.5 \pm 1.0) \times 10^{-4}$	
Γ_{108}	$\gamma f_2'(1525)$	$(6.3 \pm 1.0) \times 10^{-4}$	
Γ_{109}	$\gamma\phi\phi$	$(4.0 \pm 1.2) \times 10^{-4}$	S=2.1
Γ_{110}	$\gamma\rho\bar{\rho}$	$(3.8 \pm 1.0) \times 10^{-4}$	

Γ_{111}	$\gamma\eta(2225)$	$(2.9 \pm 0.6) \times 10^{-4}$	
Γ_{112}	$\gamma\eta(1760) \rightarrow \gamma\rho^0\rho^0$	$(1.3 \pm 0.9) \times 10^{-4}$	
Γ_{113}	$\gamma\pi^0$	$(3.9 \pm 1.3) \times 10^{-5}$	
Γ_{114}	$\gamma\rho\bar{\rho}\pi^+\pi^-$	$< 7.9 \times 10^{-4}$	CL=90%
Γ_{115}	$\gamma\gamma$	$< 5 \times 10^{-4}$	CL=90%
Γ_{116}	$\gamma\Lambda\bar{\Lambda}$	$< 1.3 \times 10^{-4}$	CL=90%
Γ_{117}	3γ	$< 5.5 \times 10^{-5}$	CL=90%
Γ_{118}	$\gamma X(2200)$		
Γ_{119}	$\gamma f_4(2220)$		
Γ_{120}	$\gamma X(1400)$		

[a] The value is for the sum of the charge states indicated.

[b] Includes $\rho\bar{\rho}\pi^+\pi^- \gamma$ and excludes $\rho\bar{\rho}\eta, \rho\bar{\rho}\omega, \rho\bar{\rho}\eta'$.[c] See the "Note on the $\eta(1440)$ " in the $\eta(1440)$ Full Listings.J/ $\psi(1S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
--------------------------	-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

59±24	BALDINI...	75	FRAG	e^+e^-
59±14	BOYARSKI	75	MRK1	e^+e^-
50±25	ESPOSITO	75B	FRAM	e^+e^-

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons})$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
---	-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

12 ± 2	BOYARSKI	75	MRK1	e^+e^-
--------	----------	----	------	----------

³Included in $\Gamma(\text{hadrons})$.

$\Gamma(e^+e^-)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
------------------	-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

5.26 ± 0.29	4 HSUEH	92	RVUE	See Υ mini-review
5.36 ± 0.28				
4.72 ± 0.35	4 ALEXANDER	89	RVUE	See Υ mini-review
4.4 ± 0.6	4 BRANDELIK	79C	DASP	e^+e^-
4.6 ± 0.8	5 BALDINI...	75	FRAG	e^+e^-
4.8 ± 0.6	BOYARSKI	75	MRK1	e^+e^-
4.6 ± 1.0	ESPOSITO	75B	FRAM	e^+e^-

⁴From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.⁵Assuming equal partial widths for e^+e^- and $\mu^+\mu^-$.

$\Gamma(\mu^+\mu^-)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
----------------------	-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4.8 ± 0.6	BOYARSKI	75	MRK1	e^+e^-
5.0 ± 1.0	ESPOSITO	75B	FRAM	e^+e^-

$\Gamma(\gamma\gamma)$	VALUE (eV)	CL%	DOCUMENT ID	TECN	COMMENT
------------------------	------------	-----	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 5.4	90	BRANDELIK	79C	DASP	e^+e^-
---------	----	-----------	-----	------	----------

J/ $\psi(1S)$ $\Gamma(\text{hadrons})/\Gamma(\text{total})$ This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel₁ in the e^+e^- annihilation.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
--	-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

4 ± 0.8	7 BALDINI...	75	FRAG	e^+e^-
3.9 ± 0.8	7 ESPOSITO	75B	FRAM	e^+e^-

$\Gamma(e^+e^-) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
--	-------------	-------------	------	---------

• • • We do not use the following data for averages, fits, limits, etc. • • •

0.35 ± 0.02	BRANDELIK	79C	DASP	e^+e^-
0.32 ± 0.07	7 BALDINI...	75	FRAG	e^+e^-
0.34 ± 0.14	BEMPORAD	75	FRAB	e^+e^-
0.34 ± 0.09	7 ESPOSITO	75B	FRAM	e^+e^-
0.36 ± 0.10	7 FORD	75	SPEC	e^+e^-

Meson Full Listings

$$J/\psi(1S) = J/\psi(3097)$$

$\Gamma(\mu^+ \mu^-) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_4 \Gamma_3 / \Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.31 ± 0.09	BEMPORAD	75	FRAB	$e^+ e^-$
0.51 ± 0.09	DASP	75	DASP	$e^+ e^-$
0.38 ± 0.05	ESPOSITO	75B	FRAM	$e^+ e^-$
0.46 ± 0.10	LIBERMAN	75	SPEC	$e^+ e^-$

$\Gamma(\rho \bar{\rho}) \times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$				$\Gamma_{75} \Gamma_3 / \Gamma$
VALUE (keV)	DOCUMENT ID	TECN	COMMENT	
9.7 ± 1.7	6 ARMSTRONG	93B	SPEC	$\bar{p} p \rightarrow e^+ e^-$
6 Using $\Gamma_{\text{total}} = 85.5^{+6.1}_{-5.8}$ MeV.				
7 Data redundant with branching ratios or partial widths above.				

J/ψ(1S) BRANCHING RATIOS

For the first four branching ratios, see also the partial widths, and (partial widths) $\times \Gamma(e^+ e^-) / \Gamma_{\text{total}}$ above.

$\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$				Γ_1 / Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.86 ± 0.02	BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(\text{virtual } \gamma \rightarrow \text{hadrons}) / \Gamma_{\text{total}}$				Γ_2 / Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.17 ± 0.02	8 BOYARSKI	75	MRK1	$e^+ e^-$
8 Included in $\Gamma(\text{hadrons}) / \Gamma_{\text{total}}$.				

$\Gamma(e^+ e^-) / \Gamma_{\text{total}}$				Γ_3 / Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0599 ± 0.0025 OUR AVERAGE				
0.0592 ± 0.0015 ± 0.0020	COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
0.069 ± 0.009	BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$				Γ_4 / Γ
VALUE	DOCUMENT ID	TECN	COMMENT	
0.0597 ± 0.0025 OUR AVERAGE				
Error includes scale factor of 1.1.				
0.0590 ± 0.0015 ± 0.0019	COFFMAN	92	MRK3	$\psi(2S) \rightarrow J/\psi \pi^+ \pi^-$
0.069 ± 0.009	BOYARSKI	75	MRK1	$e^+ e^-$

$\Gamma(e^+ e^-) / \Gamma(\mu^+ \mu^-)$				Γ_3 / Γ_4
VALUE	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.00 ± 0.05	BOYARSKI	75	MRK1	$e^+ e^-$
0.91 ± 0.15	ESPOSITO	75B	FRAM	$e^+ e^-$
0.93 ± 0.10	FORD	75	SPEC	$e^+ e^-$

HADRONIC DECAYS

$\Gamma(\rho \pi) / \Gamma_{\text{total}}$				Γ_5 / Γ	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0128 ± 0.0010 OUR AVERAGE					
0.0142 ± 0.0001 ± 0.0019		COFFMAN	88	MRK3	$e^+ e^-$
0.013 ± 0.003	150	FRANKLIN	83	MRK2	$e^+ e^-$
0.016 ± 0.004	183	ALEXANDER	78	PLUT	$e^+ e^-$
0.0133 ± 0.0021		BRANDELIK	78B	DASP	$e^+ e^-$
0.010 ± 0.002	543	BARTEL	76	CNTR	$e^+ e^-$
0.013 ± 0.003	153	JEAN-MARIE	76	MRK1	$e^+ e^-$

$\Gamma(\rho^0 \pi^0) / \Gamma(\rho \pi)$				Γ_6 / Γ_5
VALUE	DOCUMENT ID	TECN	COMMENT	
0.328 ± 0.008 ± 0.027				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.36 ± 0.03	SCHARRE	79B	MRK1	$e^+ e^-$
0.35 ± 0.08	ALEXANDER	78	PLUT	$e^+ e^-$
0.32 ± 0.08	BRANDELIK	78B	DASP	$e^+ e^-$
0.39 ± 0.11	BARTEL	76	CNTR	$e^+ e^-$
0.37 ± 0.09	JEAN-MARIE	76	MRK1	$e^+ e^-$

$\Gamma(\rho_2(1320) \rho) / \Gamma_{\text{total}}$				Γ_7 / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
10.9 ± 2.2 OUR AVERAGE					
11.7 ± 0.7 ± 2.5	7584	AUGUSTIN	89	DM2	$J/\psi \rightarrow \rho^0 \rho^\pm \pi^\mp$
8.4 ± 4.5	36	VANNUCCI	77	MRK1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\omega \pi^+ \pi^+ \pi^- \pi^-) / \Gamma_{\text{total}}$				Γ_8 / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
85 ± 34	140	VANNUCCI	77	MRK1	$e^+ e^- \rightarrow 3(\pi^+ \pi^-) \pi^0$

$\Gamma(\omega \pi^+ \pi^-) / \Gamma_{\text{total}}$				Γ_9 / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
7.2 ± 1.0 OUR AVERAGE					
7.0 ± 1.6	18058	AUGUSTIN	89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
7.8 ± 1.6	215	BURMESTER	77D	PLUT	$e^+ e^-$
6.8 ± 1.9	348	VANNUCCI	77	MRK1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(\omega \pi^+ \pi^-) / \Gamma(2(\pi^+ \pi^-) \pi^0)$				Γ_9 / Γ_{61}	
VALUE	DOCUMENT ID	TECN	COMMENT		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.3	9	JEAN-MARIE	76	MRK1	$e^+ e^-$
9 Final state $(\pi^+ \pi^-) \pi^0$ under the assumption that $\pi \pi$ is isospin 0.					

$\Gamma(K^*(892)^0 \bar{K}_2^*(1430)^0 + \text{c.c.}) / \Gamma_{\text{total}}$				Γ_{10} / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
53 ± 26	40	VANNUCCI	77	MRK1	$e^+ e^- \rightarrow \pi^+ \pi^- K^+ K^-$

$\Gamma(\omega K^*(892) \bar{K} + \text{c.c.}) / \Gamma_{\text{total}}$				Γ_{11} / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
53 ± 14 ± 14	530 ± 140	BECKER	87	MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(\omega f_2(1270)) / \Gamma_{\text{total}}$				Γ_{12} / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.3 ± 0.6 OUR AVERAGE					
4.3 ± 0.2 ± 0.6	5860	AUGUSTIN	89	DM2	$e^+ e^-$
4.0 ± 1.6	70	BURMESTER	77D	PLUT	$e^+ e^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.9 ± 0.8	81	VANNUCCI	77	MRK1	$e^+ e^- \rightarrow 2(\pi^+ \pi^-) \pi^0$

$\Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.}) / \Gamma_{\text{total}}$				Γ_{13} / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
5.0 ± 0.4 OUR AVERAGE					
4.57 ± 0.17 ± 0.70	2285	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
5.26 ± 0.13 ± 0.53		COFFMAN	88	MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp, K^+ K^- \pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.6 ± 0.6	24	FRANKLIN	83	MRK2	$J/\psi \rightarrow K^+ K^- \pi^0$
3.2 ± 0.6	48	VANNUCCI	77	MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
4.1 ± 1.2	39	BRAUNSCH...	76	DASP	$J/\psi \rightarrow K^+ X$

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) / \Gamma_{\text{total}}$				Γ_{14} / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
4.2 ± 0.4 OUR AVERAGE					
3.96 ± 0.15 ± 0.60	1192	JOUSSET	90	DM2	$J/\psi \rightarrow \text{hadrons}$
4.33 ± 0.12 ± 0.45		COFFMAN	88	MRK3	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.7 ± 0.6	45	VANNUCCI	77	MRK1	$J/\psi \rightarrow K^\pm K_S^0 \pi^\mp$

$\Gamma(K^0 \bar{K}^*(892)^0 + \text{c.c.}) / \Gamma(K^+ \bar{K}^*(892)^- + \text{c.c.})$				$\Gamma_{14} / \Gamma_{13}$	
VALUE	DOCUMENT ID	TECN	COMMENT		
0.82 ± 0.05 ± 0.09	88	COFFMAN	88	MRK3	$J/\psi \rightarrow K \bar{K}^*(892) + \text{c.c.}$

$\Gamma(\omega \pi^0 \pi^0) / \Gamma_{\text{total}}$				Γ_{15} / Γ	
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.4 ± 0.3 ± 0.7	509	AUGUSTIN	89	DM2	$J/\psi \rightarrow \pi^+ \pi^- 3\pi^0$

$\Gamma(b_1(1235)^\pm \pi^\mp) / \Gamma_{\text{total}}$				Γ_{16} / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
30 ± 5 OUR AVERAGE					
31 ± 6	4600	AUGUSTIN	89	DM2	$J/\psi \rightarrow 2(\pi^+ \pi^-) \pi^0$
29 ± 7	87	BURMESTER	77D	PLUT	$e^+ e^-$

$\Gamma(\omega K^\pm K_S^0 \pi^\mp) / \Gamma_{\text{total}}$				Γ_{17} / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
29.5 ± 1.4 ± 7.0	879 ± 41	BECKER	87	MRK3	$e^+ e^- \rightarrow \text{hadrons}$

$\Gamma(b_1(1235)^0 \pi^0) / \Gamma_{\text{total}}$				Γ_{18} / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
23 ± 3 ± 5	229	AUGUSTIN	89	DM2	$e^+ e^-$

$\Gamma(\phi K^*(892) \bar{K} + \text{c.c.}) / \Gamma_{\text{total}}$				Γ_{19} / Γ	
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
20.4 ± 2.8 OUR AVERAGE					
20.7 ± 2.4 ± 3.0		FALVARD	88	DM2	$J/\psi \rightarrow \text{hadrons}$
20 ± 3 ± 3	155 ± 20	BECKER	87	MRK3	$e^+ e^- \rightarrow \text{hadrons}$

See key on page 1343

Meson Full Listings

$J/\psi(1S) = J/\psi(3097)$

$\Gamma(\omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{20}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
19 ± 4 OUR AVERAGE				
19.8 ± 2.1 ± 3.9		10 FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
16 ± 10	22	FELDMAN	77 MRK1	e^+e^-

¹⁰ Addition of $\omega K^+ K^-$ and $\omega K^0 \bar{K}^0$ branching ratios.

$\Gamma(\omega f_J(1710) \rightarrow \omega K\bar{K})/\Gamma_{\text{total}}$ Γ_{21}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.8 ± 1.1 ± 0.3				
	11,12	FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons

¹¹ Includes unknown branching fraction $f_J(1710) \rightarrow K\bar{K}$.

¹² Addition of $f_J(1710) \rightarrow K^+ K^-$ and $f_J(1710) \rightarrow K^0 \bar{K}^0$ branching ratios.

$\Gamma(\phi 2(\pi^+ \pi^-))/\Gamma_{\text{total}}$ Γ_{22}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
16.0 ± 1.0 ± 3.0				
		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons

$\Gamma(\Delta(1232)^{++} \bar{p} \pi^-)/\Gamma_{\text{total}}$ Γ_{23}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.58 ± 0.23 ± 0.40	332	EATON	84 MRK2	e^+e^-

$\Gamma(\omega \eta)/\Gamma_{\text{total}}$ Γ_{24}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.58 ± 0.16 OUR AVERAGE				
1.43 ± 0.10 ± 0.21	378	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
1.71 ± 0.08 ± 0.20		COFFMAN	88 MRK3	$e^+e^- \rightarrow 3\pi\eta$

$\Gamma(\phi K\bar{K})/\Gamma_{\text{total}}$ Γ_{25}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
14.8 ± 2.2 OUR AVERAGE				
14.6 ± 0.8 ± 2.1		13 FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
18 ± 8	14	FELDMAN	77 MRK1	e^+e^-

¹³ Addition of $\phi K^+ K^-$ and $\phi K^0 \bar{K}^0$ branching ratios.

$\Gamma(\phi f_J(1710) \rightarrow \phi K\bar{K})/\Gamma_{\text{total}}$ Γ_{26}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.6 ± .2 ± 0.6				
	14,15	FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons

¹⁴ Including interference with $f'_2(1525)$.

¹⁵ Includes unknown branching fraction $f_J(1710) \rightarrow K\bar{K}$.

$\Gamma(p\bar{p}\omega)/\Gamma_{\text{total}}$ Γ_{27}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.30 ± 0.25 OUR AVERAGE				Error includes scale factor of 1.3.
1.10 ± 0.17 ± 0.18	486	EATON	84 MRK2	e^+e^-
1.6 ± 0.3	77	PERUZZI	78 MRK1	e^+e^-

$\Gamma(\Delta(1232)^{++} \bar{\Delta}(1232)^{--})/\Gamma_{\text{total}}$ Γ_{28}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.10 ± 0.09 ± 0.28	233	EATON	84 MRK2	e^+e^-

$\Gamma(\Sigma(1385)^- \bar{\Sigma}(1385)^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{29}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.03 ± 0.13 OUR AVERAGE				
1.00 ± 0.04 ± 0.21	631 ± 25	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*-}$
1.19 ± 0.04 ± 0.25	754 ± 27	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*+}$
0.86 ± 0.18 ± 0.22	56	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*-}$
1.03 ± 0.24 ± 0.25	68	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*+}$

$\Gamma(p\bar{p}\eta(958))/\Gamma_{\text{total}}$ Γ_{30}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.9 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.7.
0.68 ± 0.23 ± 0.17	19	EATON	84 MRK2	e^+e^-
1.8 ± 0.6	19	PERUZZI	78 MRK1	e^+e^-

$\Gamma(\phi f'_2(1525))/\Gamma_{\text{total}}$ Γ_{31}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
8 ± 4 OUR AVERAGE				Error includes scale factor of 2.7.
12.3 ± 0.6 ± 2.0		16,17 FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
4.8 ± 1.8	46	GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹⁶ Re-evaluated using $B(f'_2(1525) \rightarrow K\bar{K}) = 0.713$.

¹⁷ Including interference with $f_J(1710)$.

$\Gamma(\phi \pi^+ \pi^-)/\Gamma_{\text{total}}$ Γ_{32}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.80 ± 0.12 OUR AVERAGE				
0.78 ± 0.03 ± 0.12		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
2.1 ± 0.9	23	FELDMAN	77 MRK1	e^+e^-

$\Gamma(\phi K^{\pm} K_S^0 \pi^{\mp})/\Gamma_{\text{total}}$ Γ_{33}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
7.2 ± 0.9 OUR AVERAGE				
7.4 ± 0.9 ± 1.1		FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
7 ± 0.6 ± 1.0	163 ± 15	BECKER	87 MRK3	$e^+e^- \rightarrow$ hadrons

$\Gamma(\omega f_1(1420))/\Gamma_{\text{total}}$ Γ_{34}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
6.8 ± 1.9 ± 1.7	111 ± 31	BECKER	87 MRK3	$e^+e^- \rightarrow$ hadrons

$\Gamma(\phi \eta)/\Gamma_{\text{total}}$ Γ_{35}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.65 ± 0.07 OUR AVERAGE				
0.64 ± 0.04 ± 0.11	346	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.661 ± 0.045 ± 0.078		COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+ K^- \eta$

$\Gamma(\Xi(1530)^- \bar{\Xi}^+)/\Gamma_{\text{total}}$ Γ_{36}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.59 ± 0.09 ± 0.12	75 ± 11	HENRARD	87 DM2	e^+e^-

$\Gamma(p K^- \bar{\Sigma}(1385)^0)/\Gamma_{\text{total}}$ Γ_{37}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.51 ± 0.26 ± 0.18	89	EATON	84 MRK2	e^+e^-

$\Gamma(\omega \pi^0)/\Gamma_{\text{total}}$ Γ_{38}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.42 ± 0.06 OUR AVERAGE				Error includes scale factor of 1.4.
0.360 ± 0.028 ± 0.054	222	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.482 ± 0.019 ± 0.064		COFFMAN	88 MRK3	$e^+e^- \rightarrow \pi^0 \pi^+ \pi^- \pi^0$

$\Gamma(\phi \eta'(958))/\Gamma_{\text{total}}$ Γ_{39}/Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.33 ± 0.04 OUR AVERAGE					
0.41 ± 0.03 ± 0.08		167	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.308 ± 0.034 ± 0.036			COFFMAN	88 MRK3	$e^+e^- \rightarrow K^+ K^- \eta'$

• • • We do not use the following data for averages, fits, limits, etc. • • •

<1.3 90 VANNUCCI 77 MRK1 e^+e^-

$\Gamma(\phi f_0(980))/\Gamma_{\text{total}}$ Γ_{40}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.9.
4.6 ± 0.4 ± 0.8		18 FALVARD	88 DM2	$J/\psi \rightarrow$ hadrons
2.6 ± 0.6	50	GIDAL	81 MRK2	$J/\psi \rightarrow K^+ K^- K^+ K^-$

¹⁸ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

$\Gamma(\Xi(1530)^0 \Xi^0)/\Gamma_{\text{total}}$ Γ_{41}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.32 ± 0.12 ± 0.07	24 ± 9	HENRARD	87 DM2	e^+e^-

$\Gamma(\Sigma(1385)^- \bar{\Sigma}^+ \text{ (or c.c.)})/\Gamma_{\text{total}}$ Γ_{42}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.05 OUR AVERAGE				
0.30 ± 0.03 ± 0.07	74 ± 8	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*-}$
0.34 ± 0.04 ± 0.07	77 ± 9	HENRARD	87 DM2	$e^+e^- \rightarrow \Sigma^{*+}$
0.29 ± 0.11 ± 0.10	26	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*-}$
0.31 ± 0.11 ± 0.11	28	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^{*+}$

$\Gamma(\phi f_1(1285))/\Gamma_{\text{total}}$ Γ_{43}/Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
2.6 ± 0.5 OUR AVERAGE				Error includes scale factor of 1.1.
3.2 ± 0.6 ± 0.4		JOUSSET	90 DM2	$J/\psi \rightarrow \phi 2(\pi^+ \pi^-)$
2.1 ± 0.5 ± 0.4	25	19 JOUSSET	90 DM2	$J/\psi \rightarrow \phi \eta \pi^+ \pi^-$
0.6 ± 0.2 ± 0.1	16 ± 6	BECKER	87 MRK3	$J/\psi \rightarrow \phi K\bar{K} \pi$

¹⁹ We attribute to the $f_1(1285)$ the signal observed in the $\pi^+ \pi^- \eta$ invariant mass distribution at 1297 Mev.

$\Gamma(\rho \eta)/\Gamma_{\text{total}}$ Γ_{44}/Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.193 ± 0.023 OUR AVERAGE				
0.194 ± 0.017 ± 0.029	299	JOUSSET	90 DM2	$J/\psi \rightarrow$ hadrons
0.193 ± 0.013 ± 0.029		COFFMAN	88 MRK3	$e^+e^- \rightarrow \pi^+ \pi^- \eta$

Meson Full Listings

$J/\psi(1S) = J/\psi(3097)$

$\Gamma(\omega\eta(958))/\Gamma_{total}$	Γ_{45}/Γ
VALUE (units 10^{-3})	EVTS
0.167 ± 0.025 OUR AVERAGE	

DOCUMENT ID	TECN	COMMENT
JOUSSET 90 DM2		$J/\psi \rightarrow$ hadrons
COFFMAN 88 MRK3		$e^+e^- \rightarrow 3\pi\eta'$

$\Gamma(\omega f_0(980))/\Gamma_{total}$	Γ_{46}/Γ
VALUE (units 10^{-4})	EVTS
1.41 ± 0.27 ± 0.47	20

²⁰ Assuming $B(f_0(980) \rightarrow \pi\pi) = 0.78$.

DOCUMENT ID	TECN	COMMENT
AUGUSTIN 89 DM2		$J/\psi \rightarrow 2(\pi^+\pi^-)\pi^0$

$\Gamma(\rho\eta(958))/\Gamma_{total}$	Γ_{47}/Γ
VALUE (units 10^{-3})	EVTS
0.105 ± 0.018 OUR AVERAGE	19

DOCUMENT ID	TECN	COMMENT
JOUSSET 90 DM2		$J/\psi \rightarrow$ hadrons
COFFMAN 88 MRK3		$J/\psi \rightarrow \pi^+\pi^-\eta'$

$\Gamma(\rho\bar{\rho}\phi)/\Gamma_{total}$	Γ_{48}/Γ
VALUE (units 10^{-4})	EVTS
0.45 ± 0.13 ± 0.07	

DOCUMENT ID	TECN	COMMENT
FALVARD 88 DM2		$J/\psi \rightarrow$ hadrons

$\Gamma(\rho_2(1320)^\pm \pi^\mp)/\Gamma_{total}$	Γ_{49}/Γ
VALUE (units 10^{-4})	CL%
<43	90

DOCUMENT ID	TECN	COMMENT
BRAUNSCH... 76 DASP		e^+e^-

$\Gamma(K\bar{K}_2^*(1430) + c.c.)/\Gamma_{total}$	Γ_{50}/Γ
VALUE (units 10^{-4})	CL%
<40	90

••• We do not use the following data for averages, fits, limits, etc. •••

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		$e^+e^- \rightarrow K^0\bar{K}_2^{*0}$

DOCUMENT ID	TECN	COMMENT
BRAUNSCH... 76 DASP		$e^+e^- \rightarrow K^\pm\bar{K}_2^{*\mp}$

$\Gamma(K_2^*(1430)^0\bar{K}_2^*(1430)^0)/\Gamma_{total}$	Γ_{51}/Γ
VALUE (units 10^{-4})	CL%
<29	90

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(K^*(892)^0\bar{K}^*(892)^0)/\Gamma_{total}$	Γ_{52}/Γ
VALUE (units 10^{-4})	CL%
<5	90

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

$\Gamma(\phi f_2(1270))/\Gamma_{total}$	Γ_{53}/Γ
VALUE (units 10^{-4})	CL%
<3.7	90

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		$e^+e^- \rightarrow \pi^+\pi^-K^+K^-$

••• We do not use the following data for averages, fits, limits, etc. •••

DOCUMENT ID	TECN	COMMENT
FALVARD 88 DM2		$J/\psi \rightarrow$ hadrons

$\Gamma(\rho\bar{\rho}\rho)/\Gamma_{total}$	Γ_{54}/Γ
VALUE (units 10^{-3})	CL%
<0.31	90

DOCUMENT ID	TECN	COMMENT
EATON 84 MRK2		$e^+e^- \rightarrow$ hadrons γ

$\Gamma(\phi\eta(1440) \rightarrow \phi\eta\pi\pi)/\Gamma_{total}$	Γ_{55}/Γ
VALUE (units 10^{-4})	CL%
<2.5	90

DOCUMENT ID	TECN	COMMENT
FALVARD 88 DM2		$J/\psi \rightarrow$ hadrons

²¹ Includes unknown branching fraction $\eta(1440) \rightarrow \eta\pi\pi$.

$\Gamma(\omega f_2'(1525))/\Gamma_{total}$	Γ_{56}/Γ
VALUE (units 10^{-4})	CL%
<2.2	90

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		$e^+e^- \rightarrow \pi^+\pi^-\pi^0K^+K^-$

••• We do not use the following data for averages, fits, limits, etc. •••

DOCUMENT ID	TECN	COMMENT
FALVARD 88 DM2		$J/\psi \rightarrow$ hadrons

²² Re-evaluated assuming $B(f_2'(1525) \rightarrow K\bar{K}) = 0.713$.

$\Gamma(\Sigma(1385)^0\bar{\Lambda})/\Gamma_{total}$	Γ_{57}/Γ
VALUE (units 10^{-3})	CL%
<0.2	90

DOCUMENT ID	TECN	COMMENT
HENRARD 87 DM2		e^+e^-

$\Gamma(\Delta(1232)^+\bar{p})/\Gamma_{total}$	Γ_{58}/Γ
VALUE (units 10^{-3})	CL%
<0.1	90

DOCUMENT ID	TECN	COMMENT
HENRARD 87 DM2		e^+e^-

$\Gamma(\Sigma^0\bar{\Lambda})/\Gamma_{total}$	Γ_{59}/Γ
VALUE (units 10^{-4})	CL%
<0.9	90

DOCUMENT ID	TECN	COMMENT
HENRARD 87 DM2		e^+e^-

$\Gamma(\phi\pi^0)/\Gamma_{total}$	Γ_{60}/Γ
VALUE (units 10^{-4})	CL%
<0.068	90

DOCUMENT ID	TECN	COMMENT
COFFMAN 88 MRK3		$e^+e^- \rightarrow K^+K^-\pi^0$

$\Gamma(2(\pi^+\pi^-\pi^0))/\Gamma_{total}$	Γ_{61}/Γ
VALUE	EVTS
0.0337 ± 0.0026 OUR AVERAGE	

DOCUMENT ID	TECN	COMMENT
AUGUSTIN 89 DM2		$J/\psi \rightarrow 2(\pi^+\pi^-\pi^0)$
FRANKLIN 83 MRK2		$e^+e^- \rightarrow$ hadrons
BURMESTER 77D PLUT		e^+e^-
JEAN-MARIE 76 MRK1		e^+e^-

$\Gamma(3(\pi^+\pi^-\pi^0))/\Gamma_{total}$	Γ_{62}/Γ
VALUE	EVTS
0.029 ± 0.006 OUR AVERAGE	

DOCUMENT ID	TECN	COMMENT
FRANKLIN 83 MRK2		$e^+e^- \rightarrow$ hadrons
JEAN-MARIE 76 MRK1		e^+e^-

$\Gamma(\pi^+\pi^-\pi^0)/\Gamma_{total}$	Γ_{63}/Γ
VALUE	EVTS
0.015 ± 0.002	168

••• We do not use the following data for averages, fits, limits, etc. •••

DOCUMENT ID	TECN	COMMENT
EINSWEILER 83 MRK3		e^+e^-

$\Gamma(\pi^+\pi^-\pi^0K^+K^-)/\Gamma_{total}$	Γ_{64}/Γ
VALUE	EVTS
0.012 ± 0.003	309

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		e^+e^-

$\Gamma(4(\pi^+\pi^-\pi^0))/\Gamma_{total}$	Γ_{65}/Γ
VALUE (units 10^{-4})	EVTS
90 ± 30	13

DOCUMENT ID	TECN	COMMENT
JEAN-MARIE 76 MRK1		e^+e^-

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{total}$	Γ_{66}/Γ
VALUE (units 10^{-4})	EVTS
72 ± 23	205

DOCUMENT ID	TECN	COMMENT
VANNUCCI 77 MRK1		e^+e^-

$\Gamma(K\bar{K}\pi)/\Gamma_{total}$	Γ_{67}/Γ
VALUE (units 10^{-4})	EVTS
61 ± 10 OUR AVERAGE	


DOCUMENT ID	TECN	COMMENT
FRANKLIN 83 MRK2		$e^+e^- \rightarrow K^+K^-\pi^0$
VANNUCCI 77 MRK1		$e^+e^- \rightarrow K_S^0K^\pm\pi^\mp$

$\Gamma(\rho\bar{\rho}\pi^+\pi^-)/\Gamma_{total}$	Γ_{68}/Γ
VALUE (units 10^{-3})	EVTS
6.0 ± 0.5 OUR AVERAGE	

Error includes scale factor of 1.3. See the Ideogram below.

DOCUMENT ID	TECN	COMMENT
EATON 84 MRK2		e^+e^-
BESCH 81 BONA		e^+e^-
PERUZZI 78 MRK1		e^+e^-

WEIGHTED AVERAGE
6.0 ± 0.5 (Error scaled by 1.3)



$\Gamma(2(\pi^+\pi^-))/\Gamma_{total}$	Γ_{69}/Γ
VALUE	EVTS
0.004 ± 0.001	76

DOCUMENT ID	TECN	COMMENT
JEAN-MARIE 76 MRK1		e^+e^-

$\Gamma(3(\pi^+\pi^-))/\Gamma_{total}$	Γ_{70}/Γ
VALUE (units 10^{-4})	EVTS
40 ± 20	32

DOCUMENT ID	TECN	COMMENT
JEAN-MARIE 76 MRK1		e^+e^-

$\Gamma(n\bar{n}\pi^+\pi^-)/\Gamma_{total}$	Γ_{71}/Γ
VALUE (units 10^{-3})	EVTS
3.8 ± 3.6	5

DOCUMENT ID	TECN	COMMENT
BESCH 81 BONA		e^+e^-

See key on page 1343

Meson Full Listings

$J/\psi(1S) = J/\psi(3097)$

$\Gamma(\Sigma\Sigma)/\Gamma_{total}$					Γ_{72}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.8 ± 0.5 OUR AVERAGE					
3.18 ± 0.12 ± 0.69	884 ± 30	PALLIN	87 DM2	e^+e^-	
4.74 ± 0.48 ± 0.75	90	EATON	84 MRK2	$e^+e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$	
7.2 ± 7.8	3	BESCH	81 BONA	$e^+e^- \rightarrow \Sigma^+ \bar{\Sigma}^-$	
3.9 ± 1.2	52	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Sigma^0 \bar{\Sigma}^0$	

$\Gamma(n\bar{n})/\Gamma_{total}$					Γ_{78}/Γ
VALUE (units 10^{-2})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.19 ± 0.05 OUR AVERAGE					
0.190 ± 0.055	40	ANTONELLI	93 SPEC	e^+e^-	
0.18 ± 0.09		BESCH	78 BONA	e^+e^-	

$\Gamma(2(\pi^+\pi^-)K^+K^-)/\Gamma_{total}$					Γ_{73}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
3.1 ± 1.3	30	VANNUCCI	77 MRK1	e^+e^-	

$\Gamma(\Lambda\bar{\Lambda})/\Gamma_{total}$					Γ_{80}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.35 ± 0.14 OUR AVERAGE				Error includes scale factor of 1.2.	
1.38 ± 0.05 ± 0.20	1847	PALLIN	87 DM2	e^+e^-	
1.58 ± 0.08 ± 0.19	365	EATON	84 MRK2	e^+e^-	
2.6 ± 1.6	5	BESCH	81 BONA	e^+e^-	
1.1 ± 0.2	196	PERUZZI	78 MRK1	e^+e^-	

$\Gamma(p\bar{p}\pi^+\pi^-\pi^0)/\Gamma_{total}$					Γ_{74}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.3 ± 0.9 OUR AVERAGE				Error includes scale factor of 1.9.	
3.36 ± 0.65 ± 0.28	364	EATON	84 MRK2	e^+e^-	
1.6 ± 0.6	39	PERUZZI	78 MRK1	e^+e^-	

$\Gamma(p\bar{p}\pi^0)/\Gamma_{total}$					Γ_{81}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.09 ± 0.09 OUR AVERAGE					
1.13 ± 0.09 ± 0.09	685	EATON	84 MRK2	e^+e^-	
1.4 ± 0.4		BRANDELIK	79C DASP	e^+e^-	
1.00 ± 0.15	109	PERUZZI	78 MRK1	e^+e^-	

$\Gamma(\rho\bar{\rho})/\Gamma_{total}$					Γ_{75}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.14 ± 0.10 OUR AVERAGE					
2.0 ± 0.3	48	ANTONELLI	93 SPEC	e^+e^-	
1.91 ± 0.04 ± 0.30		PALLIN	87 DM2	e^+e^-	
2.16 ± 0.07 ± 0.15	1420	EATON	84 MRK2	e^+e^-	
2.5 ± 0.4	133	BRANDELIK	79C DASP	e^+e^-	
2.0 ± 0.5		BESCH	78 BONA	e^+e^-	
2.2 ± 0.2	331	PERUZZI	78 MRK1	e^+e^-	

$\Gamma(\Lambda\Sigma^-\pi^+ \text{ (or c.c.)})/\Gamma_{total}$					Γ_{82}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.06 ± 0.12 OUR AVERAGE					
0.90 ± 0.06 ± 0.16	225 ± 15	HENRRARD	87 DM2	$e^+e^- \rightarrow \Lambda\Sigma^+\pi^-$	
1.11 ± 0.06 ± 0.20	342 ± 18	HENRRARD	87 DM2	$e^+e^- \rightarrow \Lambda\Sigma^-\pi^+$	
1.53 ± 0.17 ± 0.38	135	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\Sigma^+\pi^-$	
1.38 ± 0.21 ± 0.35	118	EATON	84 MRK2	$e^+e^- \rightarrow \Lambda\Sigma^-\pi^+$	

$\Gamma(\rho\bar{\rho})/\Gamma(\mu^+\mu^-)$					Γ_{75}/Γ_4
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.081 ± 0.02	20	WIHK	75 PLUT	e^+e^-	

$\Gamma(\rho K^-\bar{K}^0)/\Gamma_{total}$					Γ_{83}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.89 ± 0.07 ± 0.14	307	EATON	84 MRK2	e^+e^-	

$\Gamma(\rho\bar{\rho}\eta)/\Gamma_{total}$					Γ_{76}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.09 ± 0.18 OUR AVERAGE					
2.03 ± 0.13 ± 0.15	826	EATON	84 MRK2	e^+e^-	
2.5 ± 1.2		BRANDELIK	79C DASP	e^+e^-	
2.3 ± 0.4	197	PERUZZI	78 MRK1	e^+e^-	


$\Gamma(2(K^+K^-))/\Gamma_{total}$					Γ_{84}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
7 ± 3		VANNUCCI	77 MRK1	e^+e^-	

$\Gamma(\rho\pi\pi^-)/\Gamma_{total}$					Γ_{77}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.00 ± 0.10 OUR AVERAGE					
2.02 ± 0.07 ± 0.16	1288	EATON	84 MRK2	$e^+e^- \rightarrow \rho\pi^-$	
1.93 ± 0.07 ± 0.16	1191	EATON	84 MRK2	$e^+e^- \rightarrow \bar{\rho}\pi^+$	
1.7 ± 0.7	32	BESCH	81 BONA	$e^+e^- \rightarrow \rho\pi^-$	
1.6 ± 1.2	5	BESCH	81 BONA	$e^+e^- \rightarrow \bar{\rho}\pi^+$	
2.16 ± 0.29	194	PERUZZI	78 MRK1	$e^+e^- \rightarrow \rho\pi^-$	
2.04 ± 0.27	204	PERUZZI	78 MRK1	$e^+e^- \rightarrow \bar{\rho}\pi^+$	

$\Gamma(\rho K^-\bar{K}^0)/\Gamma_{total}$					Γ_{85}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.29 ± 0.06 ± 0.05	90	EATON	84 MRK2	e^+e^-	

$\Gamma(\Xi\Xi)/\Gamma_{total}$					Γ_{79}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.8 ± 0.4 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.	
1.40 ± 0.12 ± 0.24	132 ± 11	HENRRARD	87 DM2	$e^+e^- \rightarrow \Xi^-\Xi^+$	
2.28 ± 0.16 ± 0.40	194	EATON	84 MRK2	$e^+e^- \rightarrow \Xi^-\Xi^+$	
3.2 ± 0.8	71	PERUZZI	78 MRK1	e^+e^-	

$\Gamma(K^+K^-)/\Gamma_{total}$					Γ_{86}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
2.37 ± 0.31 OUR AVERAGE					
2.39 ± 0.24 ± 0.22	107	BALTRUSAIT..85D	MRK3	e^+e^-	
2.2 ± 0.9	6	BRANDELIK	79C DASP	e^+e^-	



$\Gamma(\Lambda\bar{\Lambda}\pi^0)/\Gamma_{total}$					Γ_{87}/Γ
VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT	
0.22 ± 0.05 ± 0.05	19 ± 4	HENRRARD	87 DM2	e^+e^-	

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$					Γ_{88}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.47 ± 0.23 OUR AVERAGE					
1.58 ± 0.20 ± 0.15	84	BALTRUSAIT..85D	MRK3	e^+e^-	
1.0 ± 0.5	5	BRANDELIK	78B DASP	e^+e^-	
1.6 ± 1.6	1	VANNUCCI	77 MRK1	e^+e^-	

$\Gamma(K_S^0 K_L^0)/\Gamma_{total}$					Γ_{89}/Γ
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT	
1.08 ± 0.14 OUR AVERAGE					
1.18 ± 0.12 ± 0.18		JOUSSET	90 DM2	$J/\psi \rightarrow \text{hadrons}$	
1.01 ± 0.16 ± 0.09	74	BALTRUSAIT..85D	MRK3	e^+e^-	

$\Gamma(\Lambda\Sigma^+ \text{ c.c.})/\Gamma_{total}$					Γ_{90}/Γ
VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.15	90	PERUZZI	78 MRK1	$e^+e^- \rightarrow \Lambda X$	

$\Gamma(K_S^0 K_S^0)/\Gamma_{total}$					Γ_{91}/Γ
VALUE (units 10^{-4})	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.052	90	BALTRUSAIT..85C	MRK3	e^+e^-	

RADIATIVE DECAYS

$\Gamma(\gamma\eta_c(1S))/\Gamma_{total}$					Γ_{92}/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.0127 ± 0.0036		GAISER	86 CBAL	$J/\psi \rightarrow \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
seen	16	BALTRUSAIT..84	MRK3	$J/\psi \rightarrow 2\phi\gamma$	

Meson Full Listings

 $J/\psi(1S) = J/\psi(3097)$

$\Gamma(\gamma\pi^+\pi^-2\pi^0)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{93}/Γ
VALUE (units 10^{-3})				
8.3 ± 0.2 ± 3.1	26	BALTRUSAIT...86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$	

²⁶ 4π mass less than 2.0 GeV.

$\Gamma(\gamma\eta\pi\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{94}/Γ
VALUE (units 10^{-3})				
6.1 ± 1.0 OUR AVERAGE				
5.85 ± 0.3 ± 1.05	27	EDWARDS	83B CBAL $J/\psi \rightarrow \eta\pi^+\pi^-$	
7.8 ± 1.2 ± 2.4	27	EDWARDS	83B CBAL $J/\psi \rightarrow \eta 2\pi^0$	

²⁷ Broad enhancement at 1700 MeV.

$\Gamma(\gamma\eta(1440) \rightarrow \gamma K\bar{K}\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{95}/Γ
VALUE (units 10^{-3})				
0.91 ± 0.18 OUR AVERAGE				
0.83 ± 0.13 ± 0.18	28,29	AUGUSTIN	92 DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$	
1.03 ^{+0.21+0.26} _{-0.18-0.19}	28,30	BAI	90C MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

1.78 ± 0.21 ± 0.33	28,31	AUGUSTIN	92 DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$	
3.8 ± 0.3 ± 0.6	28	AUGUSTIN	90 DM2 $J/\psi \rightarrow \gamma K\bar{K}\pi$	
0.66 ^{+0.17+0.24} _{-0.16-0.15}	28,32	BAI	90C MRK3 $J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$	
6.3 ± 1.4	28	WISNIEWSKI	87 MRK3 $J/\psi \rightarrow K\bar{K}\pi\gamma$	
4.0 ± 0.7 ± 1.0	28	EDWARDS	82E CBAL $J/\psi \rightarrow K^+ K^- \pi^0 \gamma$	
4.3 ± 1.7	28,33	SCHARRE	80 MRK2 e^+e^-	

²⁸ Includes unknown branching fraction $\eta(1440) \rightarrow K\bar{K}\pi$.

²⁹ From fit to the $K^*(892)K^0$ $+$ partial wave.

³⁰ From $K^*(890)K$ final state.

³¹ From fit to the $a_0(980)\pi^0$ $+$ partial wave.

³² From $a_0(980)\pi$ final state.

³³ Corrected for spin-zero hypothesis for $\eta(1440)$.

$\Gamma(\gamma\eta(1440) \rightarrow \gamma\gamma\rho^0)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{96}/Γ
VALUE (units 10^{-5})				
6.4 ± 1.2 ± 0.7	34	COFFMAN	90 MRK3 $J/\psi \rightarrow \gamma\gamma\pi^+\pi^-$	

³⁴ Includes unknown branching fraction $\eta(1440) \rightarrow \gamma\rho^0$.

$\Gamma(\gamma\rho\rho)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{97}/Γ
VALUE (units 10^{-3})				
4.5 ± 0.8 OUR AVERAGE				
4.7 ± 0.3 ± 0.9	35	BALTRUSAIT...86B	MRK3 $J/\psi \rightarrow 4\pi\gamma$	
3.75 ± 1.05 ± 1.20	36	BURKE	82 MRK2 $J/\psi \rightarrow 4\pi\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

<0.09	90	37	BISELLO	89B	$J/\psi \rightarrow 4\pi\gamma$
-------	----	----	---------	-----	---------------------------------

³⁵ 4π mass less than 2.0 GeV.

³⁶ 4π mass less than 2.0 GeV, $2\rho^0$ corrected to 2ρ by factor of 3.

³⁷ 4π mass in the range 2.0–25 GeV.

$\Gamma(\gamma\eta'(958))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{98}/Γ
VALUE (units 10^{-3})				
4.31 ± 0.30 OUR AVERAGE				
4.50 ± 0.14 ± 0.53		BOLTON	92B MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-\eta, \eta \rightarrow \gamma\gamma$	

4.30 ± 0.31 ± 0.71

4.04 ± 0.16 ± 0.85

4.39 ± 0.09 ± 0.66

4.1 ± 0.3 ± 0.6

• • • We do not use the following data for averages, fits, limits, etc. • • •

2.9 ± 1.1	6	BRANDELIK	79C DASP	$e^+e^- \rightarrow 3\gamma$
3.8 ± 1.3	38	SCHARRE	79B MRK1	$e^+e^- \rightarrow \gamma X$
3.4 ± 0.7		SCHARRE	79B MRK1	$e^+e^- \rightarrow 2\pi 2\gamma$
2.4 ± 0.7	57	BARTEL	76 CNTR	$e^+e^- \rightarrow 2\gamma\rho$

³⁸ From the inclusive γ decay spectrum.

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{99}/Γ
VALUE (units 10^{-3})				
2.8 ± 0.5 OUR AVERAGE				
4.32 ± 0.14 ± 0.73	39	BISELLO	89B DM2 $J/\psi \rightarrow 4\pi\gamma$	

2.08 ± 0.13 ± 0.35

3.05 ± 0.08 ± 0.45


4.85 ± 0.45 ± 1.20

³⁹ 4π mass less than 3.0 GeV.

⁴⁰ 4π mass less than 2.0 GeV.

⁴¹ 4π mass less than 2.5 GeV.

WEIGHTED AVERAGE
2.8 ± 0.5 (Error scaled by 1.9)



$\Gamma(\gamma f_4(2050))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{100}/Γ
VALUE (units 10^{-3})				
2.7 ± 0.5 ± 0.5	42	BALTRUSAIT...87	MRK3 $J/\psi \rightarrow \gamma\pi^+\pi^-$	

⁴² Assuming branching fraction $f_4(2050) \rightarrow \pi\pi$ /total = 0.167.

$\Gamma(\gamma\omega\omega)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{101}/Γ
VALUE (units 10^{-3})				
1.59 ± 0.33 OUR AVERAGE				
1.41 ± 0.2 ± 0.42	120 ± 17	BISELLO	87 SPEC e^+e^- , hadrons γ	
1.76 ± 0.09 ± 0.45		BALTRUSAIT...85C	MRK3 $e^+e^- \rightarrow$ hadrons γ	

$\Gamma(\gamma\eta(1440) \rightarrow \gamma\rho^0\rho^0)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{102}/Γ
VALUE (units 10^{-3})				
1.36 ± 0.38	43,44	BISELLO	89B DM2 $J/\psi \rightarrow 4\pi\gamma$	

⁴³ Estimated by us from various fits.

⁴⁴ Includes unknown branching fraction to $\rho^0\rho^0$.

$\Gamma(\gamma f_2(1270))/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_{103}/Γ
VALUE (units 10^{-3})					
1.38 ± 0.14 OUR AVERAGE					
1.33 ± 0.05 ± 0.20	45	AUGUSTIN	87 DM2	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
1.36 ± 0.09 ± 0.23	45	BALTRUSAIT...87	MRK3	$J/\psi \rightarrow \gamma\pi^+\pi^-$	
1.48 ± 0.25 ± 0.30	178	EDWARDS	82B CBAL	$e^+e^- \rightarrow 2\pi^0\gamma$	
2.0 ± 0.7	35	ALEXANDER	78 PLUT 0	e^+e^-	
1.2 ± 0.6	30	BRANDELIK	78B DASP	$e^+e^- \rightarrow \pi^+\pi^-\gamma$	

⁴⁵ Estimated using $B(f_2(1270) \rightarrow \pi\pi) = 0.843 \pm 0.012$. The errors do not contain the uncertainty in the $f_2(1270)$ decay.

⁴⁶ Restated by us to take account of spread of E1, M2, E3 transitions.

$\Gamma(\gamma f_2(1710) \rightarrow \gamma K\bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{104}/Γ
VALUE (units 10^{-4})				
9.7 ± 1.2 OUR AVERAGE				
9.2 ± 1.4 ± 1.4	47	AUGUSTIN	88 DM2 $J/\psi \rightarrow \gamma K^+ K^-$	

10.4 ± 1.2 ± 1.6

9.6 ± 1.2 ± 1.8

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.8

1.6 ± 0.4 ± 0.3

3.8 ± 1.6

⁴⁷ Includes unknown branching fraction to K^+K^- or $K_S^0 K_S^0$. We have multiplied K^+K^- measurement by 2, and $K_S^0 K_S^0$ by 4 to obtain $K\bar{K}$ result.

⁴⁸ Includes unknown branching fraction to $\rho^0\rho^0$.

⁴⁹ Includes unknown branching fraction to $\pi^+\pi^-$.

⁵⁰ Includes unknown branching fraction to $\eta\eta$.

$\Gamma(\gamma\eta)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{105}/Γ
VALUE (units 10^{-3})				
0.86 ± 0.08 OUR AVERAGE				
0.88 ± 0.08 ± 0.11		BLOOM	83 CBAL e^+e^-	

0.82 ± 0.10

1.3 ± 0.4

21

BARTEL 77 CNTR e^+e^-

Meson Full Listings
 $J/\psi(1S) = J/\psi(3097)$

$\Gamma(\gamma f_1(1420) \rightarrow \gamma K \bar{K} \pi) / \Gamma_{total}$ Γ_{106} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.83 ± 0.15 OUR AVERAGE			
0.76 ± 0.15 ± 0.21	51,52 AUGUSTIN	92 DM2	$J/\psi \rightarrow \gamma K \bar{K} \pi$
0.87 ± 0.14 $^{+0.14}_{-0.11}$	51 BAI	90C MRK3	$J/\psi \rightarrow \gamma K_S^0 K^\pm \pi^\mp$

⁵¹ Included unknown branching fraction $f_1(1420) \rightarrow K \bar{K} \pi$.
⁵² From fit to the $K^*(892)K 1^{++}$ partial wave.

$\Gamma(\gamma f_1(1285)) / \Gamma_{total}$ Γ_{107} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT
0.65 ± 0.10 OUR AVERAGE				
0.625 ± 0.063 ± 0.103		53 BOLTON	92 MRK3	$J/\psi \rightarrow \gamma f_1(1285)$
0.70 ± 0.09 ± 0.16		54 BURCHELL	91 MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
0.025 ± 0.007 ± 0.003		55 COFFMAN	90 MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
< 0.6		90 56 SCHARRE	80 MRK2	$J/\psi \rightarrow \gamma K \bar{K} \pi$

⁵³ Obtained summing the sequential decay channels
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \pi \pi \pi) = (1.44 \pm 0.39 \pm 0.27) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \delta \pi, \delta \rightarrow \eta \pi) = (3.90 \pm 0.42 \pm 0.87) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \delta \pi, \delta \rightarrow K \bar{K}) = (0.66 \pm 0.26 \pm 0.29) \times 10^{-4}$;
 $B(J/\psi \rightarrow \gamma f_1(1285), f_1(1285) \rightarrow \gamma \rho^0) = (0.25 \pm 0.07 \pm 0.03) \times 10^{-4}$.
⁵⁴ Using $B(f_1(1285) \rightarrow a_0(980)\pi) = 0.37$, and including unknown branching ratio for $a_0(980) \rightarrow \eta \pi$.
⁵⁵ Includes unknown branching fraction $f_1(1285) \rightarrow \gamma \rho^0$.
⁵⁶ Using $B(f_1(1285) \rightarrow K \bar{K} \pi) = 0.12$.

$\Gamma(\gamma f'_2(1525)) / \Gamma_{total}$ Γ_{108} / Γ


VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.63 ± 0.10 OUR AVERAGE					
0.70 ± 0.17 ± 0.11			57 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
0.56 ± 0.06 ± 0.11			57 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
0.84 ± 0.20 ± 0.17			57 BALTRUSAIT..87	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

⁵⁷ Using $B(f'_2(1525) \rightarrow K \bar{K}) = 0.713$.
⁵⁸ Assuming isotropic production and decay of the $f'_2(1525)$ and isospin.

$\Gamma(\gamma \phi \phi) / \Gamma_{total}$ Γ_{109} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
4.0 ± 1.2 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
7.5 ± 0.6 ± 1.2	168	BAI	90B MRK3	$J/\psi \rightarrow \gamma 4K$
3.4 ± 0.8 ± 0.6	33 ± 7	59 BISELLO	90 DM2	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_S^0$
3.1 ± 0.7 ± 0.4		59 BISELLO	86B DM2	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$

⁵⁹ $\phi \phi$ mass less than 2.9 GeV, η_c excluded.



$\Gamma(\gamma \rho \rho) / \Gamma_{total}$ Γ_{110} / Γ

VALUE (units 10^{-3})	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.38 ± 0.07 ± 0.07					
< 0.11		90	PERUZZI	78 MRK1	$e^+ e^-$

$\Gamma(\gamma \eta(2225)) / \Gamma_{total}$ Γ_{111} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.29 ± 0.06 OUR AVERAGE			
0.33 ± 0.08 ± 0.05	60 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K^+ K^-$
0.27 ± 0.06 ± 0.06	60 BAI	90B MRK3	$J/\psi \rightarrow \gamma K^+ K^- K_S^0 K_S^0$
0.24 $^{+0.15}_{-0.10}$	61,62 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

⁶⁰ Includes unknown branching fraction to $\phi \phi$.
⁶¹ Estimated by us from various fits.
⁶² Includes unknown branching fraction to $\rho^0 \rho^0$.

$\Gamma(\gamma \eta(1760) \rightarrow \gamma \rho^0 \rho^0) / \Gamma_{total}$ Γ_{112} / Γ

VALUE (units 10^{-3})	DOCUMENT ID	TECN	COMMENT
0.13 ± 0.09	63,64 BISELLO	89B DM2	$J/\psi \rightarrow 4\pi \gamma$

⁶³ Estimated by us from various fits.
⁶⁴ Includes unknown branching fraction to $\rho^0 \rho^0$.

$\Gamma(\gamma \pi^0) / \Gamma_{total}$ Γ_{113} / Γ

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.039 ± 0.013 OUR AVERAGE				
0.036 ± 0.011 ± 0.007		BLOOM	83 CBAL	$e^+ e^-$
0.073 ± 0.047	10	BRANDELIK	79c DASP	$e^+ e^-$

$\Gamma(\gamma \rho \rho \pi^+ \pi^-) / \Gamma_{total}$ Γ_{114} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.79		90	EATON	84 MRK2	$e^+ e^-$

$\Gamma(\gamma \gamma) / \Gamma_{total}$ Γ_{115} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.5		90	BARTEL	77 CNTR	$e^+ e^-$

$\Gamma(\gamma \Lambda \bar{\Lambda}) / \Gamma_{total}$ Γ_{116} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.13		90	HENRARD	87 DM2	$e^+ e^-$

$\Gamma(3\gamma) / \Gamma_{total}$ Γ_{117} / Γ

VALUE (units 10^{-3})	CL%	DOCUMENT ID	TECN	COMMENT	
< 0.065		90	PARTRIDGE	80 CBAL	$e^+ e^-$

$\Gamma(\gamma X(2200)) / \Gamma_{total}$ Γ_{118} / Γ

VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
1.5	65 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$

⁶⁵ Includes unknown branching fraction to $K_S^0 K_S^0$.

$\Gamma(\gamma f_4(2220)) / \Gamma_{total}$ Γ_{119} / Γ

VALUE (units 10^{-5})	EVTS	DOCUMENT ID	TECN	COMMENT
< 2.3	95	66 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K^+ K^-$
< 1.6	95	66 AUGUSTIN	88 DM2	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
12.4 $^{+6.4}_{-5.2} \pm 2.8$	23	66 BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K_S^0 K_S^0$
8.4 $^{+3.4}_{-2.8} \pm 1.6$	93	66 BALTRUSAIT..86D	MRK3	$J/\psi \rightarrow \gamma K^+ K^-$

⁶⁶ Includes unknown branching fraction to $K^+ K^-$ or $K_S^0 K_S^0$.

$\Gamma(\gamma X(1400)) / \Gamma_{total}$ Γ_{120} / Γ

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
3.83 ± 0.33 ± 0.059		67 BURCHELL	91 MRK3	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$
7.0 ± 0.6 ± 1.1	261	67 AUGUSTIN	90 DM2	$J/\psi \rightarrow \gamma \eta \pi^+ \pi^-$

⁶⁷ Includes unknown branching fraction to $\eta \pi^+ \pi^-$.

Meson Full Listings

$$J/\psi(1S) = J/\psi(3097), \chi_{c0}(1P) = \chi_{c0}(3415)$$

J/ψ(1S) REFERENCES

ANTONELLI	93	PL B301 317	+Baldini+	(FENICE Collab.)
ARMSTRONG	93B	PR D47 772	+Bettoni, Bharadwaj+	(FNAL E760 Collab.)
AUGUSTIN	92	PR D46 1951	+Cosme	(DM2 Collab.)
BOLTON	92	PR B278 495	+Brown, Bunnell+	(Mark III Collab.)
BOLTON	92B	PRL 69 1328	+Brown, Bunnell+	(Mark III Collab.)
COFFMAN	92	PRL 68 282	+DeJongh, Dubois, Hitlin+	(Mark III Collab.)
HSUEH	92	PR D45 R2181	+Palestini	(FNAL, TORI)
BURCHHELL	91	NP B21 132 (suppl)		(Mark III Collab.)
AUGUSTIN	90	PR D42 10	+Cosme+	(DM2 Collab.)
BAI	90B	PRL 65 1309	+Blaylock+	(Mark III Collab.)
BAI	90C	PRL 65 2507	+Blaylock+	(Mark III Collab.)
BISELLO	90	PL B241 617	+Busetto+	(DM2 Collab.)
COFFMAN	90	PR D41 1410	+De Jongh+	(Mark III Collab.)
JOUSSET	90	PR D41 1389	+Ajaltouni+	(DM2 Collab.)
ALEXANDER	89	NP B320 45	+Bonvicini, Dreli, Frey, Luth	(LBL, MICH, SLAC)
AUGUSTIN	89	NP B320 1	+Cosme	(DM2 Collab.)
BISELLO	89B	PR D39 701	+Busetto+	(DM2 Collab.)
AUGUSTIN	88	PRL 60 2238	+Calcaterra+	(DM2 Collab.)
COFFMAN	88	PR D38 2695	+Dubois, Egen, Hauser+	(Mark III Collab.)
FALVARD	88	PR D38 2706	+Ajaltouni+	(CLER, FRAS, LALO, PADO)
AUGUSTIN	87	ZPHY C36 369	+Cosme+	(LALO, CLER, FRAS, PADO)
BAGLIN	87	NP B286 592	+ (LAPP, CERN, GENO, LYON, OSLO, ROMA)+	
BALTRUSAITIS...	87	PR D35 2077	+Baltrusaitis, Coffman, Dubois+	(Mark III Collab.)
BECKER	87	PRL 59 186	+Blaylock, Bolton, Brown+	(Mark III Collab.)
BISELLO	87	PL B192 239	+Ajaltouni, Baldini+	(PADO, CLER, FRAS, LALO)
HENRARD	87	NP B292 670	+Ajaltouni, et al	(CLER, FRAS, LALO, PADO)
PALLIN	87	NP B292 653	+Ajaltouni+	(CLER, FRAS, LALO, PADO)
WISNIEWSKI	87	Hadron 87 Conf.		(Mark III Collab.)
BALTRUSAITIS...	86B	PR D33 1222	+Baltrusaitis, Coffman, Hauser+	(Mark III Collab.)
BALTRUSAITIS...	86D	PRL 56 107	+Baltrusaitis	(CIT, UCSC, ILL, SLAC, WASH)
BISELLO	86B	PL B179 294	+Busetto, Castro, Limentani+	(DM2 Collab.)
GAISER	86	PR D34 711	+Bloom, Bulos, Godfrey+	(Crystal Ball Collab.)
BALTRUSAITIS...	85C	PRL 55 1723	+Baltrusaitis+	(CIT, UCSC, ILL, SLAC, WASH)
BALTRUSAITIS...	85D	PR D32 566	+Baltrusaitis, Coffman+	(CIT, UCSC, ILL, SLAC, WASH)
BALTRUSAITIS...	84	PRL 52 126	+Baltrusaitis+	(CIT, UCSC, ILL, SLAC, WASH)
EATON	84	PR D29 804	+Goldhaber, Abrams, Alam, Boyarski+	(LBL, SLAC)
BLOOM	83	ARNS 33 143	+Peck	(SLAC, CIT)
EDWARDS	83B	PRL 51 859	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
EINSWEILER	83	Brighton Conf. 348		(Mark III Collab.)
FRANKLIN	83	PRL 51 963	+Franklin, Feldman, Abrams, Alam+	(LBL, SLAC)
FRANKLIN	83B	Thesis SLAC-0254		(STAN)
BURKE	82	PRL 49 632	+Trilling, Abrams, Alam, Blocker+	(LBL, SLAC)
EDWARDS	82B	PR D25 3065	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
EDWARDS	82D	PRL 48 458	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
Also	83	ARNS 33 143	+Bloom, Peck	(SLAC, CIT)
EDWARDS	82E	PRL 49 259	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
LEMOIGNIE	82	PL 113B 509	+Barate, Astbury+	(SACL, LOIC, SHMP, IND)
BESCH	81	ZPHY C8 1	+Eisermann, Lohr, Kowalski+	(BONN, DESY, MANZ)
GIDAL	81	PL 107B 153	+Goldhaber, Guy, Millikan, Abrams+	(SLAC, LBL)
PARTRIDGE	80	PL 44 712	+Peck+	(CIT, HARV, PRIN, STAN, SLAC)
SCHARRE	80	PL 97B 329	+Trilling, Abrams, Alam, Blocker+	(SLAC, LBL)
ZHOLENTZ	80	PL 96B 214	+Kurdadze, Lelechuk, Mishnev+	(NOVO)
Also	81	SJNP 34 814	Zholentz, Kurdadze, Lelechuk+	(NOVO)
Translated from YAF	34	1471.		
BRANDELIC	79C	ZPHY C1 233	+Cords+	(DASP Collab.)
SCHARRE	79B	SLAC-PUB-2321		(SLAC, LBL)
Also	79B	LBL-9502		(SLAC, LBL)
ALEXANDER	78	PL 72B 493	+Abrams, Alam, Blocker, Boyarski+	(SLAC, LBL)
BESCH	78	PL 78B 347	+Criegee+	(DESY, HAMB, SIEG, WUPP)
BRANDELIC	78B	PL 74B 292	+Eisermann, Kowalski, Eyss+	(BONN, DESY, MANZ)
PERUZZI	78	PR D17 2901	+Cords+	(DASP Collab.)
BARTEL	77	PL 66B 489	+Piccolo, Alam, Boyarski, Goldhaber+	(SLAC, LBL)
BURMESTER	77D	PL 72B 135	+Duinker, Olsson, Heintze+	(DESY, HEIDP)
FELDMAN	77	PR D15 1814	+Criegee+	(DESY, HAMB, SIEG, WUPP)
VANNUCCI	77	PR D15 1814	+Peri	(LBL, SLAC)
BARTEL	76	PL 64B 483	+Abrams, Alam, Boyarski+	(SLAC, LBL)
BRUNSCHE...	76	PL 63B 487	+Duinker, Olsson, Steffen, Heintze+	(DESY, HEIDP)
JEAN-MARIE	76	PRL 36 291	+Braunschweig+	(DASP Collab.)
BALDINI...	75	PL 58B 471	+Abrams, Boyarski, Breidenbach+	(SLAC, LBL) IG
BEMPORAD	75	Stanford Symp. 113	+Baldini-Celio, Bozzo, Capon+	(FRAS, ROMA)
BOYARSKI	75	PRL 34 1357	+Baldini-Celio, Bozzo, Capon+	(FRAS, ROMA)
DASP	75	PL 56B 491	+Braunschweig, Konigs+	(DASP Collab.)
ESPOSITO	75B	LNC 14 73	+Bartoli, Bisello+	(FRAS, NAPL, PADO, ROMA)
FORD	75	PRL 34 604	+Beron, Hilger, Hofstadter+	(SLAC, PENN)
LIBERMAN	75	Stanford Symp. 55		(STAN)
WIJK	75	Stanford Symp. 69		(DESY)

OTHER RELATED PAPERS

BAGLIN	85	SLAC Summer Inst. 609	(LAPP, CERN, GENO, LYON, OSLO, ROMA)+
LEE	85	SLAC 282	(SLAC)
BARATE	83	PL 121B 449	+Bareyre, Bonamy+ (SACL, LOIC, SHMP, IND)
ABRAMS	74	PRL 33 1453	+Briggs, Augustin, Boyarski+ (LBL, SLAC)
ASH	74	LNC 11 705	+Zorn, Bartoli+ (FRAS, UMD, NAPL, PADO, ROMA)
AUBERT	74	PRL 33 1404	+Becker, Biggs, Burger, Chen, Everhart (MIT, BNL)
AUGUSTIN	74	PRL 33 1406	+Boyarski, Abrams, Briggs+ (SLAC, LBL)
BACCI	74	PRL 33 1408	+Bartoli, Barbarino, Barbiellini+ (FRAS)
Also	74B	PRL 33 1649	Bacci
BALDINI...	74	LNC 11 711	Baldini-Celio, Bacci+ (FRAS, ROMA)
BARBIELLINI	74	LNC 11 718	+Bemporad+ (FRAS, NAPL, PISA, ROMA)
BRUNSCHE...	74	PL 53B 393	+Braunschweig+ (DASP Collab.)
CHRISTENS...	70	PRL 25 1523	+Christenson, Hicks, Lederman+ (COLU, BNL, CERN)

$$\chi_{c0}(1P)$$

$$\text{or } \chi_{c0}(3415)$$

$$I^G(J^{PC}) = 0^+(0^{++})$$

Observed in the radiative decay $\psi(2S) \rightarrow \chi_{c0}(1P)\gamma$. Therefore $C = +$. The observed decay into $\pi^+\pi^-$ or K^+K^- implies $G = +$, $J^P = 0^+, 2^+, \dots$. The angular distribution is consistent with $J = 0$. J^P abnormal excluded by $\pi^+\pi^-$ and K^+K^- decays. $J^P = 0^+$ preferred (FELDMAN 77).

χ_{c0}(1P) MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
3415.1 ± 1.0 OUR AVERAGE				
3417.8 ± 0.4 ± 4		1 GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X$
3414.8 ± 1.1	2,3	HIMEL	79 MRK2	$e^+e^- \rightarrow \text{hadrons}$
3422.0 ± 10.0		2 BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3416.0 ± 3 ± 4		2 TANENBAUM	78 MRK1	e^+e^-
3415.0 ± 9.0		2 BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma X$
••• We do not use the following data for averages, fits, limits, etc. •••				
3407.0 ± 8.0	2	4 WIJK	75 DASP	$e^+e^- \rightarrow J/\psi 2\gamma$

1 Using mass of $\psi(2S) = 3686.0$ MeV.
2 Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.
3 Systematic error added linearly by us.
4 Only two events; this mass apparently never published.

χ_{c0}(1P) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.5 ± 3.3 ± 4.2	GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma X, \gamma \pi^0 \pi^0$

χ_{c0}(1P) DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	(3.7 ± 0.7) %	
Γ_2 $\pi^+\pi^- K^+K^-$	(3.0 ± 0.7) %	
Γ_3 $\rho^0 \pi^+\pi^-$	(1.6 ± 0.5) %	
Γ_4 $3(\pi^+\pi^-)$	(1.5 ± 0.5) %	
Γ_5 $K^+ \bar{K}^*(892)^0 \pi^- + c.c.$	(1.2 ± 0.4) %	
Γ_6 $\pi^+\pi^-$	(7.5 ± 2.1) × 10 ⁻³	
Γ_7 K^+K^-	(7.1 ± 2.4) × 10 ⁻³	
Γ_8 $\pi^+\pi^- \rho \bar{\rho}$	(5.0 ± 2.0) × 10 ⁻³	
Γ_9 $\pi^0 \pi^0$	(3.1 ± 0.6) × 10 ⁻³	
Γ_{10} $\eta \eta$	(2.5 ± 1.1) × 10 ⁻³	
Γ_{11} $\rho \bar{\rho}$	< 9.0 × 10 ⁻⁴	90%
Radiative decays		
Γ_{12} $\gamma J/\psi(1S)$	(6.6 ± 1.8) × 10 ⁻³	
Γ_{13} $\gamma \gamma$	(4.0 ± 2.3) × 10 ⁻⁴	

χ_{c0}(1P) PARTIAL WIDTHS

Γ(γγ)	CL%	DOCUMENT ID	TECN	COMMENT	Γ ₁₃
VALUE (keV)					
< 6.2	95	CHEN	90B CLEO	$e^+e^- \rightarrow e^+e^- \chi_{c0}$	
4.0 ± 2.8		LEE	85 CBAL	$\psi' \rightarrow \text{photons}$	
••• We do not use the following data for averages, fits, limits, etc. •••					
< 17	95	AIHARA	88D TPC	$e^+e^- \rightarrow e^+e^- X$	

χ_{c0}(1P) BRANCHING RATIOS

HADRONIC DECAYS

Γ(2(π ⁺ π ⁻))/Γ _{total}	DOCUMENT ID	TECN	COMMENT	Γ _{1/Γ}
VALUE				
0.037 ± 0.007	5	TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$
Γ(π ⁺ π ⁻ K ⁺ K ⁻)/Γ _{total}	DOCUMENT ID	TECN	COMMENT	Γ _{2/Γ}
VALUE				
0.030 ± 0.007	5	TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$
Γ(ρ ⁰ π ⁺ π ⁻)/Γ _{total}	DOCUMENT ID	TECN	COMMENT	Γ _{3/Γ}
VALUE				
0.016 ± 0.005	5	TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$
Γ(3(π ⁺ π ⁻))/Γ _{total}	DOCUMENT ID	TECN	COMMENT	Γ _{4/Γ}
VALUE				
0.015 ± 0.005	5	TANENBAUM	78 MRK1	$\psi(2S) \rightarrow \gamma \chi_{c0}$

See key on page 1343

Meson Full Listings

$$\chi_{c0}(1P) = \chi_{c0}(3415), \chi_{c1}(1P) = \chi_{c1}(3510)$$

$\Gamma(K^+ \bar{K}^*(892)^0 \pi^- + c.c.) / \Gamma_{total}$	Γ_5 / Γ
VALUE (units 10^{-4})	DOCUMENT ID TECN COMMENT
0.012 ± 0.004	⁵ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(\pi^+ \pi^-) / \Gamma_{total}$	Γ_6 / Γ
VALUE (units 10^{-4})	DOCUMENT ID TECN COMMENT
75 ± 21 OUR AVERAGE	
70 ± 30	⁵ BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma \chi_{c0}$
80 ± 30	⁵ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(K^+ K^-) / \Gamma_{total}$	Γ_7 / Γ
VALUE (units 10^{-4})	DOCUMENT ID TECN COMMENT
71 ± 24 OUR AVERAGE	
60 ± 30	⁵ BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma \chi_{c0}$
90 ± 40	⁵ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(\pi^+ \pi^- \rho^0) / \Gamma_{total}$	Γ_8 / Γ
VALUE	DOCUMENT ID TECN COMMENT
0.006 ± 0.002	⁵ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(\pi^0 \pi^0) / \Gamma_{total}$	Γ_9 / Γ
VALUE (units 10^{-3})	DOCUMENT ID TECN COMMENT
3.1 ± 0.4 ± 0.5	⁶ LEE 85 CBAL $\psi' \rightarrow$ photons

$\Gamma(\eta \eta) / \Gamma_{total}$	Γ_{10} / Γ
VALUE (units 10^{-3})	DOCUMENT ID TECN COMMENT
2.5 ± 0.8 ± 0.8	⁶ LEE 85 CBAL $\psi' \rightarrow$ photons

$\Gamma(\rho^0 \rho^0) / \Gamma_{total}$	Γ_{11} / Γ
VALUE (units 10^{-4})	DOCUMENT ID TECN COMMENT
< 9.0	90 ⁵ BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma \chi_{c0}$
	⁵ Calculated using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.094$; the errors do not contain the uncertainty in the $\psi(2S)$ decay.
	⁶ Calculated using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.093 \pm 0.008$.

RADIATIVE DECAYS

$\Gamma(\gamma J/\psi(1S)) / \Gamma_{total}$	Γ_{12} / Γ
VALUE (units 10^{-4})	DOCUMENT ID TECN COMMENT
66 ± 18 OUR AVERAGE	
60 ± 18	GAISER 86 CBAL $\psi(2S) \rightarrow \gamma \chi_{c0}$
320 ± 210	⁷ BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma \chi_{c0}$
150 ± 100	⁷ BARTEL 78B CNTR $\psi(2S) \rightarrow \gamma \chi_{c0}$
210 ± 210	⁷ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c0}$

$\Gamma(\gamma \gamma) / \Gamma_{total}$	Γ_{13} / Γ
VALUE (units 10^{-4})	DOCUMENT ID TECN COMMENT
4.0 ± 2.0 ± 1.1	⁶ LEE 85 CBAL $\psi' \rightarrow$ photons
	• • • We do not use the following data for averages, fits, limits, etc. • • •
< 15	90 ⁷ YAMADA 77 DASP $e^+ e^- \rightarrow 3\gamma$
	⁷ Calculated using $B(\psi(2S) \rightarrow \gamma \chi_{c0}(1P)) = 0.094$; the errors do not contain the uncertainty in the $\psi(2S)$ decay.

 $\chi_{c0}(1P)$ REFERENCES

CHEN	90B	PL B243 169	+McIlwain+	(CLEO Collab.)
AIHARA	88D	PRL 60 2355	+Alston-Garnjost+	(TPC Collab.)
GAISER	86	PR D34 711	+Bloom, Bulos, Godfrey+	(Crystal Ball Collab.)
LEE	85	SLAC 282		(SLAC)
BRANDELIK	79B	NP B160 426	+Cords+	(DASP Collab.)
HIMEL	79	Thesis SLAC-0223		(SLAC)
	82	Private Comm.	Trilling	(LBL, UCB)
BARTEL	78B	PL 79B 492	+Dittmann, Duinker, Olsson, O'Neill+	(DESY, HEIDP)
TANENBAUM	78	PR D17 1731	+Alam, Boyarski+	(SLAC, LBL)
	82	Private Comm.	Trilling	(LBL, UCB)
BIDDICK	77	PRL 38 1324	+Burnett+	(UCSD, UMD, PAVI, PRIN, SLAC, STAN)
FELDMAN	77	PRPL 33C 285	+Peri	(LBL, SLAC)
YAMADA	77	Hamburg Conf. 69		(DASP Collab.)
WIJK	75	Stanford Symp. 69		(DESY)

OTHER RELATED PAPERS

OREGLIA	82	PR D25 2259	+Partridge+	(SLAC, CIT, HARV, PRIN, STAN)
FELDMAN	75B	PRL 35 821	+Jean-Marie, Sadoulet, Vannucci+	(LBL, SLAC)
	Also	75C PRL 35 1189	Feldman	
TANENBAUM	75	PRL 35 1323	+Whitaker, Abrams+	(LBL, SLAC)

$$\chi_{c1}(1P)$$

$$\text{or } \chi_{c1}(3510)$$

$$I^G(J^{PC}) = 0^+(1^{++})$$

Observed in the radiative sequential decay $\psi(2S) \rightarrow \chi_{c1}(1P)\gamma$, $\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma$. Therefore, $C = +$. The lack of decays into $\pi^+\pi^-$ or K^+K^- is suggestive of $J^P = \text{abnormal}$. The decays into 4π and 6π imply $G = +$, thus $I = 0$. $J=0,2$ excluded by angular distribution in the $J/\psi(1S)\gamma$ decay. $J^P = 1^+$ preferred (FELDMAN 77, OREGLIA 82).

 $\chi_{c1}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3510.53 ± 0.12 OUR AVERAGE				
3510.53 ± 0.04 ± 0.12	513	ARMSTRONG 92	SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
3511.3 ± 0.4 ± 0.4	30	BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
3512.3 ± 0.3 ± 4.0		1 GAISER 86	CBAL	$\psi(2S) \rightarrow \gamma X$
3507.4 ± 1.7	91	2 LEMOIGNE 82	GOLJ	190 GeV $\pi^-\text{Be} \rightarrow \gamma 2\mu$
3510.4 ± 0.6		OREGLIA 82	CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3510.1 ± 1.1	254	3 HIMEL 80	MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3509.0 ± 11.0	21	BRANDELIK 79B	DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3507.0 ± 3.0		3 BARTEL 78B	CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3505.0 ± 4 ± 4		3,4 TANENBAUM 78	MRK1	e^+e^-
3513.0 ± 7.0	367	3 BIDDICK 77	CNTR	$\psi(2S) \rightarrow \gamma X$
		• • •		We do not use the following data for averages, fits, limits, etc. • • •
3510.0 ± 20.0		BARTEL 76B	CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3500 ± 10	40	TANENBAUM 75	MRK1	Hadrons γ
3507.0 ± 7.0	7	WIJK 75	DASP	$e^+e^- \rightarrow J/\psi 2\gamma$

¹ Using mass of $\psi(2S) = 3686.0$ MeV.

² $J/\psi(1S)$ mass constrained to 3097 MeV.

³ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.

⁴ From a simultaneous fit to radiative and hadronic decay channels.

 $\chi_{c1}(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.88 ± 0.11 ± 0.06		513	ARMSTRONG 92	SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
		• • •			We do not use the following data for averages, fits, limits, etc. • • •
< 1.3	95		BAGLIN 86B	SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
< 3.8	90		GAISER 86	CBAL	$\psi(2S) \rightarrow \gamma X$

 $\chi_{c1}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Hadronic decays	
Γ_1 $3(\pi^+\pi^-)$	(2.2 ± 0.8) %
Γ_2 $2(\pi^+\pi^-)$	(1.6 ± 0.5) %
Γ_3 $\pi^+\pi^-K^+K^-$	(9 ± 4) × 10 ⁻³
Γ_4 $\rho^0\pi^+\pi^-$	(3.9 ± 3.5) × 10 ⁻³
Γ_5 $K^+\bar{K}^*(892)^0\pi^- + c.c.$	(3.2 ± 2.1) × 10 ⁻³
Γ_6 $\pi^+\pi^-\rho^0$	(1.4 ± 0.9) × 10 ⁻³
Γ_7 ρ^0	(8.6 ± 1.2) × 10 ⁻⁵
Γ_8 $\pi^+\pi^- + K^+K^-$	< 2.1 × 10 ⁻³
Radiative decays	
Γ_9 $\gamma J/\psi(1S)$	(27.3 ± 1.6) %
Γ_{10} $\gamma\gamma$	

 $\chi_{c1}(1P)$ PARTIAL WIDTHS

$\Gamma(\rho^0)$	Γ_7
VALUE (eV)	DOCUMENT ID TECN COMMENT
74 ± 9 OUR AVERAGE	
76 ± 10 ± 5	513 ⁵ ARMSTRONG 92 SPEC $\bar{p}p \rightarrow e^+e^-\gamma$
69 ⁺¹⁵ ₋₁₃ ± 4	⁵ BAGLIN 86B SPEC $\bar{p}p \rightarrow e^+e^-\gamma$

⁵ Restated by us using $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma)B(J/\psi(1S) \rightarrow e^+e^-) = 0.0171 \pm 0.0011$.

 $\chi_{c1}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(3(\pi^+\pi^-)) / \Gamma_{total}$	Γ_1 / Γ
VALUE	DOCUMENT ID TECN COMMENT
0.022 ± 0.008	⁷ TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma \chi_{c1}$

Meson Full Listings

$$\chi_{c1}(1P) = \chi_{c1}(3510), h_c(1P), \chi_{c2}(1P) = \chi_{c2}(3555)$$

$\Gamma(2\pi^+\pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
VALUE 0.016±0.005	7	TANENBAUM	78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	

$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
VALUE (units 10 ⁻⁴) 90±40	7	TANENBAUM	78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	

$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
VALUE (units 10 ⁻⁴) 39±35	7	TANENBAUM	78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	

$\Gamma(K^+\bar{K}^*(892)^0\pi^- + \text{c.c.})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
VALUE (units 10 ⁻⁴) 32±21	7	TANENBAUM	78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	

$\Gamma(\pi^+\pi^-\rho^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
VALUE (units 10 ⁻⁴) 14±9	7	TANENBAUM	78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	

$\Gamma(\rho^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
VALUE (units 10 ⁻⁴) 0.86±0.12	513	6	ARMSTRONG 92 SPEC $\bar{p}p \rightarrow e^+e^-\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 > 0.54 95 BAGLIN 86B SPEC $\bar{p}p \rightarrow e^+e^-\chi$
 <12.0 90 7 BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma\chi_{c1}$
⁶ Restated by us using $B(\chi_{c1}(1P) \rightarrow J/\psi(1S)\gamma)B(J/\psi(1S) \rightarrow e^+e^-) = 0.0171 \pm 0.0011$.

$[\Gamma(\pi^+\pi^-) + \Gamma(K^+K^-)]/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
VALUE (units 10 ⁻⁴) <21	7	FELDMAN	77 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	

• • • We do not use the following data for averages, fits, limits, etc. • • •
 <38 90 7 BRANDELIK 79B DASP $\psi(2S) \rightarrow \gamma\chi_{c1}$
⁷ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

RADIATIVE DECAYS

$\Gamma(\gamma J/\psi(1S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
VALUE 0.273±0.016 OUR AVERAGE				

0.284±0.021		GAISER	86 CBAL $\psi(2S) \rightarrow \gamma\chi$	
0.274±0.046	943	8 OREGLIA	82 CBAL $\psi(2S) \rightarrow \gamma\chi_{c1}$	
0.28 ±0.07		8 HIMEL	80 MRK2 $\psi(2S) \rightarrow \gamma\chi_{c1}$	
0.19 ±0.05		8 BRANDELIK	79B DASP $\psi(2S) \rightarrow \gamma\chi_{c1}$	
0.29 ±0.05		8 BARTEL	78B CNTR $\psi(2S) \rightarrow \gamma\chi_{c1}$	
0.28 ±0.09		8 TANENBAUM	78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c1}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.57 ±0.17		8 BIDDICK	77 CNTR $\psi(2S) \rightarrow \gamma\chi$	

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
VALUE <0.0015	90	8	YAMADA 77 DASP $e^+e^- \rightarrow 3\gamma$	

• • • We do not use the following data for averages, fits, limits, etc. • • •
⁸ Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c1}(1P)) = 0.087$. The errors do not contain the uncertainty in the $\psi(2S)$ decay.

 $\chi_{c1}(1P)$ REFERENCES

ARMSTRONG 92 NP B373 35	+Bettoni+ (FNAL, FERR, GENO, UCI, NWES+)
Also 92B PRL 68 1468	Armstrong, Bettoni+(FNAL, FERR, GENO, UCI, NWES+)
BAGLIN 86B PL B172 455	(LAPP, CERN, GENO, LYON, OSLO, ROMA+)
GAISER 86 PR D34 711	+Bloom, Bulos, Godfrey+ (Crystal Ball Collab.)
LEMOIGNE 82 PL 113B 509	+Barate, Astbury+ (SACL, LOIC, SHMP, IND)
OREGLIA 82 PR D25 2259	+Partridge+ (SLAC, CIT, HARV, PRIN, STAN)
HIMEL 80 PRL 44 920	Oreglia (EFI)
Also 82B Private Comm.	+Abrams, Alam, Blocker+ (LBL, SLAC)
BRANDELIK 79B NP B160 426	Trilling (LBL, UCB)
BARTEL 78B PL 79B 492	+Cords+ (DASP Collab.)
TANENBAUM 78 PR D17 1731	+Dittmann, Duinker, Olsson, O'Neill+ (DESY, HEIDP)
Also 82 Private Comm.	+Alam, Boyarski+ (SLAC, LBL)
BIDDICK 77 PRL 38 1324	Trilling (LBL, UCB)
FELDMAN 77 PRL 33C 285	+Burnett+ (UCSD, UMD, PAVI, PRIN, SLAC, STAN)
YAMADA 77 Hamburg Conf. 69	+Peri (LBL, SLAC)
BARTEL 76B Tbilisi Conf. N75	(DASP Col'rb.)
TANENBAUM 75 PRL 35 1323	+Duinker, Olsson, Heintze+ (DESY, HEIDP)
WIJK 75 Stanford Symp. 69	+Whitaker, Abrams+ (LBL, SI C)
	(DF I)

OTHER RELATED PAPERS

BARATE 83 PL 121B 449	+Bareyre, Bonamy+ (SACL, LOIC, SHMP, IND)
BRAUNSCH... 75B PL 57B 407	Braunschweig, Konigs+ (DASP Collab.)
FELDMAN 75 Stanford Symp. 39	(SLAC)
HEINTZE 75 Stanford Symp. 97	(HEIDP)
SIMPSON 75 PRL 35 699	+Beron, Ford, Hilger, Hofstadter+ (STAN, PENN)

 $h_c(1P)$

$$I^G(J^{PC}) = ?^?(?^{??})$$

OMITTED FROM SUMMARY TABLE

Observed in the reaction $\bar{p}p \rightarrow J/\psi(1S)\pi^0$ close to the center of gravity of the 3P_J states and has characteristics consistent with what is expected for the 1P_1 state. First indications obtained by BAGLIN 86 in the reaction $\bar{p}p \rightarrow J/\psi(1S)\chi$. Needs confirmation.

 $h_c(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3526.14±0.24 OUR AVERAGE				
3526.20±0.15±0.20	59	ARMSTRONG	92D SPEC	$\bar{p}p \rightarrow J/\psi\pi^0$
3525.4 ±0.8 ±0.4	5	BAGLIN	86 SPEC	$\bar{p}p \rightarrow J/\psi\chi$

 $h_c(1P)$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1.1	90	59	ARMSTRONG	92D SPEC	$\bar{p}p \rightarrow J/\psi\pi^0$

 $h_c(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi(1S)\pi^0$	seen
Γ_2 $J/\psi(1S)\pi\pi$	not seen
Γ_3 $p\bar{p}$	

$\Gamma(J/\psi(1S)\pi\pi)/\Gamma(J/\psi(1S)\pi^0)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
VALUE <0.18	90	ARMSTRONG	92D SPEC	$\bar{p}p \rightarrow J/\psi\pi^0$

 $h_c(1P)$ REFERENCES

ARMSTRONG 92D PRL 69 2337	+Bettoni+ (FNAL, FERR, GENO, UCI, PENN, TORI)
BAGLIN 86 PL B171 135	+Baird+ (LAPP, CERN, TORI, STRB, OSLO, ROMA+)

 $\chi_{c2}(1P)$ or $\chi_{c2}(3555)$

$$I^G(J^{PC}) = 0^+(2^{++})$$

Observed in the radiative decay $\psi(2S) \rightarrow \chi_{c2}(1P)\gamma$. Therefore $C = +$. The observed decay into 4π and 6π imply $G = +$, thus $I = 0$. $J = 0$ is excluded by the angular distribution in the hadronic decays. J^P abnormal excluded by $\pi^+\pi^-$ and K^+K^- decays. $J^P = 2^+$ preferred (FELDMAN 77, OREGLIA 82).

 $\chi_{c2}(1P)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3586.17±0.13 OUR AVERAGE				
3556.15±0.07±0.12	585	ARMSTRONG	92 SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
3556.9 ±0.4 ±0.5	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-\chi$
3557.8 ±0.2 ±4		1 GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma\chi$
3553.4 ±2.2	66	2 LEMOIGNE	82 GOL1	190 GeV $\pi^-\text{Be} \rightarrow \gamma 2\mu$
3555.9 ±0.7		3 OREGLIA	82 CBAL	$e^+e^- \rightarrow J/\psi 2\gamma$
3557 ±1.5	69	4 HIMEL	80 MRK2	$e^+e^- \rightarrow J/\psi 2\gamma$
3551.0 ±11.0	15	BRANDELIK	79B DASP	$e^+e^- \rightarrow J/\psi 2\gamma$
3553.0 ±4.0		4 BARTEL	78B CNTR	$e^+e^- \rightarrow J/\psi 2\gamma$
3553.0 ±4 ±4		4,5 TANENBAUM	78 MRK1	e^+e^-
3563.0 ±7.0	360	4 BIDDICK	77 CNTR	$e^+e^- \rightarrow \gamma\chi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
3550.0 ±10.0		TRILLING	76 MRK1	$e^+e^- \rightarrow \text{hadrons}\gamma$
3543.0 ±10.0	4	WHITAKER	76 MRK1	$e^+e^- \rightarrow J/\psi 2\gamma$

¹ Using mass of $\psi(2S) = 3686.0$ MeV.
² $J/\psi(1S)$ mass constrained to 3097 MeV.
³ Assuming $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.
⁴ Mass value shifted by us by amount appropriate for $\psi(2S)$ mass = 3686 MeV and $J/\psi(1S)$ mass = 3097 MeV.
⁵ From a simultaneous fit to radiative and hadronic decay channels.

 $\chi_{c2}(1P)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2.00±0.18 OUR AVERAGE				
1.98±0.17±0.07	585	ARMSTRONG	92 SPEC	$\bar{p}p \rightarrow e^+e^-\gamma$
2.6 +1.4 -1.0	50	BAGLIN	86B SPEC	$\bar{p}p \rightarrow e^+e^-\chi$
2.8 +2.1 -2.0		6 GAISER	86 CBAL	$\psi(2S) \rightarrow \gamma\chi$

⁶ Errors correspond to 90% confidence level; authors give only width range.

See key on page 1343

Meson Full Listings

$$\chi_{c2}(1P) = \chi_{c2}(3555)$$

$\chi_{c2}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Hadronic decays		
Γ_1 $2(\pi^+\pi^-)$	(2.2 ± 0.5) %	
Γ_2 $\pi^+\pi^-K^+K^-$	(1.9 ± 0.5) %	
Γ_3 $3(\pi^+\pi^-)$	(1.2 ± 0.8) %	
Γ_4 $\rho^0\pi^+\pi^-$	(7 ± 4) × 10 ⁻³	
Γ_5 $K^+\bar{K}^*(892)^0\pi^- + c.c.$	(4.8 ± 2.8) × 10 ⁻³	
Γ_6 $\pi^+\pi^-p\bar{p}$	(3.3 ± 1.3) × 10 ⁻³	
Γ_7 $\pi^+\pi^-$	(1.9 ± 1.0) × 10 ⁻³	
Γ_8 K^+K^-	(1.5 ± 1.1) × 10 ⁻³	
Γ_9 $p\bar{p}$	(10.0 ± 1.0) × 10 ⁻⁵	
Γ_{10} $\pi^0\pi^0$	(1.10 ± 0.28) × 10 ⁻³	
Γ_{11} $\eta\eta$	(8 ± 5) × 10 ⁻⁴	
Γ_{12} $J/\psi(1S)\pi^+\pi^-\pi^0$	< 1.5 %	90%
Radiative decays		
Γ_{13} $\gamma J/\psi(1S)$	(13.5 ± 1.1) %	
Γ_{14} $\gamma\gamma$	(1.6 ± 0.5) × 10 ⁻⁴	

$\chi_{c2}(1P)$ PARTIAL WIDTHS

$\Gamma(p\bar{p})$		Γ_9
VALUE (eV)	EVTS	DOCUMENT ID TECN COMMENT
206 ± 22 OUR AVERAGE		
197 ± 18 ± 16	585	7 ARMSTRONG 92 SPEC $\bar{p}p \rightarrow e^+e^-\gamma$
252 ⁺⁵⁵ ₋₄₈ ± 21		7 BAGLIN 86B SPEC $\bar{p}p \rightarrow e^+e^-X$
7 Restated by us using $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma)B(J/\psi(1S) \rightarrow e^+e^-) = 0.0085 \pm 0.0007$.		
$\Gamma(\gamma\gamma)$		Γ_{14}
VALUE (keV)	CL%	DOCUMENT ID TECN COMMENT
0.321 ± 0.078 ± 0.054		8 ARMSTRONG 93 SPEC $\bar{p}p \rightarrow \gamma\gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •		
3.4 ± 1.7 ± 0.9		BAUER 93 TPC $e^+e^- \rightarrow e^+e^-\chi_{c2}$
< 4.2	95	UEHARA 91 VNS $e^+e^- \rightarrow e^+e^-\chi_{c2}$
< 1.0	95	CHEN 90B CLEO $e^+e^- \rightarrow e^+e^-\chi_{c2}$
< 4.2	95	AIHARA 88D TPC $e^+e^- \rightarrow e^+e^-X$
2.9 ^{+1.3} _{-1.0} ± 1.7		BAGLIN 87B SPEC $\bar{p}p \rightarrow \gamma\gamma$
< 1.6	90	YAMADA 77 DASP $e^+e^- \rightarrow 3\gamma$
8 Using $B(\chi_{c2}(1P) \rightarrow p\bar{p}) = (1.00 \pm 0.23) \times 10^{-4}$ and $\Gamma_{total} = 2.00 \pm 0.18$ MeV.		

$\chi_{c2}(1P)$ BRANCHING RATIOS

HADRONIC DECAYS

$\Gamma(2(\pi^+\pi^-))/\Gamma_{total}$		Γ_1/Γ
VALUE	DOCUMENT ID TECN COMMENT	
0.022 ± 0.006	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
$\Gamma(\pi^+\pi^-K^+K^-)/\Gamma_{total}$		Γ_2/Γ
VALUE	DOCUMENT ID TECN COMMENT	
0.019 ± 0.006	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
$\Gamma(3(\pi^+\pi^-))/\Gamma_{total}$		Γ_3/Γ
VALUE	DOCUMENT ID TECN COMMENT	
0.012 ± 0.008	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
$\Gamma(\rho^0\pi^+\pi^-)/\Gamma_{total}$		Γ_4/Γ
VALUE (units 10 ⁻⁴)	DOCUMENT ID TECN COMMENT	
68 ± 40	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
$\Gamma(K^+\bar{K}^*(892)^0\pi^- + c.c.)/\Gamma_{total}$		Γ_5/Γ
VALUE (units 10 ⁻⁴)	DOCUMENT ID TECN COMMENT	
48 ± 28	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
$\Gamma(\pi^+\pi^-p\bar{p})/\Gamma_{total}$		Γ_6/Γ
VALUE (units 10 ⁻⁴)	DOCUMENT ID TECN COMMENT	
33 ± 13	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	
$\Gamma(\pi^+\pi^-)/\Gamma_{total}$		Γ_7/Γ
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID TECN COMMENT
1.9 ± 1.0	4	10 BRANDELIK 79C DASP $\psi(2S) \rightarrow \gamma\chi_{c2}$
$[\Gamma(\pi^+\pi^-) + \Gamma(K^+K^-)]/\Gamma_{total}$		$(\Gamma_7 + \Gamma_8)/\Gamma$
VALUE (units 10 ⁻⁴)	DOCUMENT ID TECN COMMENT	
24 ± 10	10 TANENBAUM 78 MRK1 $\psi(2S) \rightarrow \gamma\chi_{c2}$	

$\Gamma(K^+K^-)/\Gamma_{total}$

VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.5 ± 1.1	2	10 BRANDELIK 79C DASP		$\psi(2S) \rightarrow \gamma\chi_{c2}$

$\Gamma(p\bar{p})/\Gamma_{total}$

VALUE (units 10 ⁻⁴)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
1.00 ± 0.10 OUR AVERAGE					
1.00 ± 0.11		585	9 ARMSTRONG 92 SPEC		$\bar{p}p \rightarrow e^+e^-\gamma$
0.97 ^{+0.44} _{-0.28} ± 0.08			BAGLIN 86B SPEC		$\bar{p}p \rightarrow e^+e^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 9.5		90	10 BRANDELIK 79B DASP		$\psi(2S) \rightarrow \gamma\chi_{c2}$
9 Restated by us using $B(\chi_{c2}(1P) \rightarrow J/\psi(1S)\gamma)B(J/\psi(1S) \rightarrow e^+e^-) = 0.0085 \pm 0.0007$.					

$\Gamma_i\Gamma_j/\Gamma_{total}^2 \ln p\bar{p} \rightarrow \chi_{c2}(1P) \rightarrow \gamma\gamma$

VALUE (units 10 ⁻⁷)	EVTS	DOCUMENT ID	TECN	COMMENT
0.160 ± 0.039 ± 0.016		ARMSTRONG 93 SPEC		$\bar{p}p \rightarrow \gamma\gamma$
0.99 ^{+0.46} _{-0.35}	6	11 BAGLIN 87B SPEC		$\bar{p}p \rightarrow \gamma\gamma$

$\Gamma(\pi^0\pi^0)/\Gamma_{total}$

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
1.1 ± 0.2 ± 0.2	12 LEE 85 CBAL		$\psi' \rightarrow$ photons

$\Gamma(\eta\eta)/\Gamma_{total}$

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
7.9 ± 4.1 ± 2.4	12 LEE 85 CBAL		$\psi' \rightarrow$ photons

$\Gamma(J/\psi(1S)\pi^+\pi^-\pi^0)/\Gamma_{total}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.015	90	BARATE 81 SPEC		190 GeV $\pi^-\pi^-\text{Be} \rightarrow 2\pi 2\mu$

10 Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.078$; the errors do not contain the uncertainty in the $\psi(2S)$ decay.

11 Assuming isotropic $\chi_{c2}(1P) \rightarrow \gamma\gamma$ distribution.

12 LEE 85 result is calculated using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.078 \pm 0.008$.

RADIATIVE DECAYS

$\Gamma(\gamma J/\psi(1S))/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.135 ± 0.011 OUR AVERAGE				
0.124 ± 0.015		GAISER 86 CBAL		$\psi(2S) \rightarrow \gamma X$
0.162 ± 0.028	479	13 OREGLIA 82 CBAL		$\psi(2S) \rightarrow \gamma\chi_{c2}$
0.14 ± 0.04		13 HIMEL 80 MRK2		$\psi(2S) \rightarrow \gamma\chi_{c2}$
0.18 ± 0.05		13 BRANDELIK 79B DASP		$\psi(2S) \rightarrow \gamma\chi_{c2}$
0.13 ± 0.03		13 BARTEL 78B CNTR		$\psi(2S) \rightarrow \gamma\chi_{c2}$
0.11 ^{+0.13} _{-0.07}		13 SPITZER 78 PLUT		$\psi(2S) \rightarrow \gamma\chi_{c2}$
0.13 ± 0.08		13 TANENBAUM 78 MRK1		$\psi(2S) \rightarrow \gamma\chi_{c2}$
0.28 ± 0.13		13 BIDDICK 77 CNTR		$\psi(2S) \rightarrow \gamma X$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
13 Estimated using $B(\psi(2S) \rightarrow \gamma\chi_{c2}(1P)) = 0.078$; the errors do not contain the uncertainty in the $\psi(2S)$ decay.				

$\Gamma(\gamma\gamma)/\Gamma_{total}$

VALUE (units 10 ⁻⁴)	DOCUMENT ID	TECN	COMMENT
1.60 ± 0.39 ± 0.23	14 ARMSTRONG 93 SPEC		$\bar{p}p \rightarrow \gamma\gamma$

14 Using $B(\chi_{c2}(1P) \rightarrow p\bar{p}) = (1.00 \pm 0.23) \times 10^{-4}$.

$\chi_{c2}(1P)$ REFERENCES

ARMSTRONG 93 PRL 70 2988	+Bettoni, Bharadwaj+ (FNAL E760 Collab.)
BAUER 93 PL B302 345	+Belcinski+ (TPC Collab.)
ARMSTRONG 92 NP B373 35	+Bettoni+ (FNAL, FERR, GENO, UCI, NWES+)
Also 92B PRL 68 1468	Armstrong, Bettoni+(FNAL, FERR, GENO, UCI, NWES+)
UEHARA 91 PL B266 188	+Abe+ (VENUS Collab.)
CHEN 90B PL B243 169	+McIlwain+ (CLEO Collab.)
AIHARA 88D PRL 60 2355	+Alston-Garnjost+ (TPC Collab.)
BAGLIN 87B PL B187 191	+Baird, Bassompierre, Borzani+ (R704 Collab.)
BAGLIN 86B PL B172 455	(LAPP, CERN, GENO, LYON, OSLO, ROMA+)
GAISER 86 PR D34 711	+Bloom, Bulos, Godfrey+ (Crystal Ball Collab.)
LEE 85 SLAC 282	(SLAC)
LEMOIGNE 82 PL 1138 509	+Barate, Astbury+ (SACL, LOIC, SHMP, CERN, IND)
OREGLIA 82 PR D25 2259	+Partridge+ (SLAC, CIT, HARV, PRIN, STAN)
Also 82B Private Comm.	Oreglia (EFI)
BARATE 81 PR D24 2994	+Astbury+ (SACL, LOIC, SHMP, CERN, IND)
HIMEL 80 PRL 44 920	+Abrams, Alam, Blocker+ (LBL, SLAC)
Also 82 Private Comm.	Trilling (DASP Collab.)
BRANDELIK 79B NP B160 426	+Cords+ (DASP Collab.)
BRANDELIK 79C ZPHY C1 233	+Cords+ (DASP Collab.)
BARTEL 78B PL 79B 492	+Dittmann, Duinker, Olsson, O'Neill+ (DESY, HEIDP)
SPITZER 78 Kyoto Sum. Inst. 47	(HAMB)
TANENBAUM 78 PR D17 1731	+Alam, Boyarski+ (SLAC, LBL)
Also 82 Private Comm.	Trilling (LBL, UCB)
BIDDICK 77 PRL 38 1324	+Burnett+ (UCSD, UMD, PAVI, PRIN, SLAC, STAN)
FELDMAN 77 PRPL 33C 285	+Perl (LBL, SLAC)
YAMADA 77 Hamburg Conf. 69	(DASP Collab.)
TRILLING 76 Stanford Symp. 437	(LBL)
WHITAKER 76 PRL 37 1596	+Tanenbaum, Abrams, Alam+ (SLAC, LBL)

Meson Full Listings

$$\chi_{c2}(1P) = \chi_{c2}(3555), \eta_c(2S) = \eta_c(3590), \psi(2S) = \psi(3685)$$

OTHER RELATED PAPERS

BARATE	83	PL 121B 449	+Bareyre, Bonamy+	(SACL, LOIC, SHMP, IND)
FELDMAN	75B	PRL 35 821	+Jean-Marie, Sadoulet, Vannucci+	(LBL, SLAC)
Also	75C	PRL 35 1189	Feldman	
Erratum.				
TANENBAUM	75	PRL 35 1323	+Whitaker, Abrams+	(LBL, SLAC)

$\eta_c(2S)$
or $\eta_c(3590)$

$$I^G(J^{PC}) = ?^?(?^?+)$$

OMITTED FROM SUMMARY TABLE

Our latest mini-review on this particle can be found in the 1984 edition. Needs confirmation.

$\eta_c(2S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3694.0 ± 5.0	¹ EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

¹ Assuming mass of $\psi(2S) = 3686$ MeV.

$\eta_c(2S)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 8.0	95	EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\eta_c(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	seen
Γ_2 $\gamma\gamma$	

$\eta_c(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
seen	EDWARDS	82C CBAL	$e^+e^- \rightarrow \gamma X$

$\Gamma(\gamma\gamma)/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.01	90	LEE	85 CBAL	$\psi' \rightarrow \text{photons}$

$\eta_c(2S)$ REFERENCES

LEE	85	SLAC 282		(SLAC)
EDWARDS	82C	PRL 48 70	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)

OTHER RELATED PAPERS

OREGLIA	82	PR D25 2259	+Partridge+	(SLAC, CIT, HARV, PRIN, STAN)
PORTER	81	SLAC Summer Inst.	355-Edwards+	(CIT, HARV, PRIN, STAN, SLAC)
BARTEL	78B	PL 79B 492	+Dittmann, Duinker, Olsson, O'Neill+	(DESY, HEIDP)

$\psi(2S)$
or $\psi(3685)$

$$I^G(J^{PC}) = 0^-(1^-)$$

$\psi(2S)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
3686.00 ± 0.09 OUR AVERAGE				
3686.02 ± 0.09 ± 0.27		ARMSTRONG 93B	SPEC	$\bar{p}p \rightarrow e^+e^-$
3686.00 ± 0.10	413	ZHOLENTZ 80	OLYA	e^+e^-

$m_{\psi(2S)} - m_{J/\psi(1S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
589.07 ± 0.13 OUR AVERAGE			
589.7 ± 1.2	LEMOIGNE 82	GOLI	190 GeV $\pi^- \text{Be} \rightarrow 2\mu$
589.07 ± 0.13	¹ ZHOLENTZ 80	OLYA	e^+e^-
588.7 ± 0.8	LUTH 75	MRK1	

¹ Redundant with data in mass above.

$\psi(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
277 ± 31 OUR AVERAGE			Error includes scale factor of 1.1.
306 ± 36 ± 16	ARMSTRONG 93B	SPEC	$\bar{p}p \rightarrow e^+e^-$
243 ± 43	² PDG	92	RVUE

² Uses $\Gamma(ee)$ from ALEXANDER 89 and $B(ee) = (88 \pm 13) \times 10^{-4}$ from FELDMAN 77.

$\psi(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 hadrons	(98.10 ± 0.30) %	
Γ_2 virtual $\gamma \rightarrow$ hadrons	(2.9 ± 0.4) %	
Γ_3 e^+e^-	(8.8 ± 1.3) × 10 ⁻³	
Γ_4 $\mu^+\mu^-$	(7.7 ± 1.7) × 10 ⁻³	

Decays into $J/\psi(1S)$ and anything

Γ_5 $J/\psi(1S)$ anything	(57 ± 4) %	
Γ_6 $J/\psi(1S)$ neutrals	(23.2 ± 2.6) %	
Γ_7 $J/\psi(1S)\pi^+\pi^-$	(32.4 ± 2.6) %	
Γ_8 $J/\psi(1S)\pi^0\pi^0$	(18.4 ± 2.7) %	
Γ_9 $J/\psi(1S)\eta$	(2.7 ± 0.4) %	S=1.7
Γ_{10} $J/\psi(1S)\pi^0$	(9.7 ± 2.1) × 10 ⁻⁴	

Hadronic decays

Γ_{11} $3(\pi^+\pi^-)\pi^0$	(3.5 ± 1.6) × 10 ⁻³	
Γ_{12} $2(\pi^+\pi^-)\pi^0$	(3.1 ± 0.7) × 10 ⁻³	
Γ_{13} $\pi^+\pi^-K^+K^-$	(1.6 ± 0.4) × 10 ⁻³	
Γ_{14} $\pi^+\pi^-\rho\bar{\rho}$	(8.0 ± 2.0) × 10 ⁻⁴	
Γ_{15} $K^+K^*(892)^0\pi^- + \text{c.c.}$	(6.7 ± 2.5) × 10 ⁻⁴	
Γ_{16} $2(\pi^+\pi^-)$	(4.5 ± 1.0) × 10 ⁻⁴	
Γ_{17} $\rho^0\pi^+\pi^-$	(4.2 ± 1.5) × 10 ⁻⁴	
Γ_{18} $\bar{p}p$	(1.9 ± 0.5) × 10 ⁻⁴	
Γ_{19} $3(\pi^+\pi^-)$	(1.5 ± 1.0) × 10 ⁻⁴	
Γ_{20} $\bar{p}p\pi^0$	(1.4 ± 0.5) × 10 ⁻⁴	
Γ_{21} K^+K^-	(1.0 ± 0.7) × 10 ⁻⁴	
Γ_{22} $\pi^+\pi^-\pi^0$	(9 ± 5) × 10 ⁻⁵	
Γ_{23} $\pi^+\pi^-$	(8 ± 5) × 10 ⁻⁵	
Γ_{24} $\Lambda\bar{\Lambda}$	< 4 × 10 ⁻⁴	CL=90%
Γ_{25} $\Xi^-\Xi^+$	< 2 × 10 ⁻⁴	CL=90%
Γ_{26} $\rho\pi$	< 8.3 × 10 ⁻⁵	CL=90%
Γ_{27} $K^+K^-\pi^0$	< 2.96 × 10 ⁻⁵	CL=90%
Γ_{28} $K^+K^*(892)^- + \text{c.c.}$	< 1.79 × 10 ⁻⁵	CL=90%

Radiative decays

Γ_{29} $\gamma\chi_{c0}(1P)$	(9.3 ± 0.8) %	
Γ_{30} $\gamma\chi_{c1}(1P)$	(8.7 ± 0.8) %	
Γ_{31} $\gamma\chi_{c2}(1P)$	(7.8 ± 0.8) %	
Γ_{32} $\gamma\eta_c(1S)$	(2.8 ± 0.6) × 10 ⁻³	
Γ_{33} $\gamma\eta_c(2S)$		
Γ_{34} $\gamma\pi^0$	< 5.4 × 10 ⁻³	CL=95%
Γ_{35} $\gamma\eta'(958)$	< 1.1 × 10 ⁻³	CL=90%
Γ_{36} $\gamma\eta$		
Γ_{37} $\gamma\gamma$	< 1.6 × 10 ⁻⁴	CL=90%
Γ_{38} $\gamma\eta(1440) \rightarrow \gamma K\bar{K}\pi$	[a] < 1.2 × 10 ⁻⁴	CL=90%

Mode needed for fitting purposes

Γ_{39} 1. — other fit modes	(30 ± 4) %
------------------------------------	------------

[a] See the "Note on the $\eta(1440)$ " in the $\eta(1440)$ Full Listings.

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 13 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 6.9$ for 8 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_8	35				
x_9	0	-11			
x_{30}	1	-7	0		
x_{31}	0	-3	0	0	
x_{39}	-80	-78	-4	-14	-16
	x_7	x_8	x_9	x_{30}	x_{31}

$\psi(2S)$ PARTIAL WIDTHS

$\Gamma(\text{hadrons})$ Γ_1			
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
224 ± 56	LUTH	75	MRK1 e^+e^-
••• We do not use the following data for averages, fits, limits, etc. •••			
$\Gamma(e^+e^-)$ Γ_3			
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.14 ± 0.21	ALEXANDER	89	RVUE See Υ mini-review
••• We do not use the following data for averages, fits, limits, etc. •••			
2.0 ± 0.3	BRANDELIK	79c	DASP e^+e^-
2.1 ± 0.3	³ LUTH	75	MRK1 e^+e^-
³ From a simultaneous fit to e^+e^- , $\mu^+\mu^-$, and hadronic channels assuming $\Gamma(e^+e^-) = \Gamma(\mu^+\mu^-)$.			
$\Gamma(\gamma\gamma)$ Γ_{37}			
VALUE (eV)	CL%	DOCUMENT ID	TECN COMMENT
<43	90	BRANDELIK	79c DASP e^+e^-

$\psi(2S) \Gamma(l)\Gamma(e^+e^-)/\Gamma(\text{total})$

This combination of a partial width with the partial width into e^+e^- and with the total width is obtained from the integrated cross section into channel l in the e^+e^- annihilation. We list only data that have not been used to determine the partial width $\Gamma(l)$ or the branching ratio $\Gamma(l)/\text{total}$.

$\Gamma(\text{hadrons}) \times \Gamma(e^+e^-)/\Gamma_{\text{total}}$ $\Gamma_1\Gamma_3/\Gamma$			
VALUE (keV)	DOCUMENT ID	TECN	COMMENT
2.2 ± 0.4	ABRAMS	75	MRK1 e^+e^-
••• We do not use the following data for averages, fits, limits, etc. •••			

$\psi(2S)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$ Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	COMMENT
0.981 ± 0.003	⁴ LUTH	75	MRK1 e^+e^-
$\Gamma(\text{virtual}\gamma \rightarrow \text{hadrons})/\Gamma_{\text{total}}$ Γ_2/Γ			
VALUE	DOCUMENT ID	TECN	COMMENT
0.029 ± 0.004	⁵ LUTH	75	MRK1 e^+e^-
$\Gamma(e^+e^-)/\Gamma_{\text{total}}$ Γ_3/Γ			
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
88 ± 13	⁶ FELDMAN	77	RVUE e^+e^-
$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$ Γ_4/Γ			
VALUE (units 10^{-4})	DOCUMENT ID	TECN	COMMENT
77 ± 17	⁷ HILGER	75	SPEC e^+e^-
$\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ Γ_4/Γ_3			
VALUE	DOCUMENT ID	TECN	COMMENT
0.89 ± 0.16	BOYARSKI	75c	MRK1 e^+e^-
⁴ Includes cascade decay into $J/\psi(1S)$.			
⁵ Included in $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$.			
⁶ From an overall fit assuming equal partial widths for e^+e^- and $\mu^+\mu^-$. For a measurement of the ratio see the entry $\Gamma(\mu^+\mu^-)/\Gamma(e^+e^-)$ below. Includes LUTH 75, HILGER 75, BURMESTER 77.			
⁷ Restated by us using $B(\psi(2S) \rightarrow J/\psi(1S)\text{anything}) = 0.55$.			

DECAYS INTO $J/\psi(1S)$ AND ANYTHING

$\Gamma(J/\psi(1S)\text{anything})/\Gamma_{\text{total}}$ $\Gamma_5/\Gamma = (\Gamma_7 + \Gamma_8 + \Gamma_9 + 0.273\Gamma_{30} + 0.135\Gamma_{31})/\Gamma$			
VALUE	DOCUMENT ID	TECN	COMMENT
0.57 ± 0.04 OUR FIT			
0.58 ± 0.07 OUR AVERAGE			
0.51 ± 0.12	BRANDELIK	79c	DASP $e^+e^- \rightarrow \mu^+\mu^- X$
0.57 ± 0.08	ABRAMS	75b	MRK1 $e^+e^- \rightarrow \mu^+\mu^- X$

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma_{\text{total}}$ $\Gamma_6/\Gamma = (0.9761\Gamma_8 + 0.708\Gamma_9 + 0.273\Gamma_{30} + 0.135\Gamma_{31})/\Gamma$			
VALUE	DOCUMENT ID	TECN	COMMENT
0.232 ± 0.026 OUR FIT			

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\text{anything})$ $\Gamma_6/\Gamma_5 = (0.9761\Gamma_8 + 0.708\Gamma_9 + 0.273\Gamma_{30} + 0.135\Gamma_{31})/(\Gamma_7 + \Gamma_8 + \Gamma_9 + 0.273\Gamma_{30} + 0.135\Gamma_{31})$			
VALUE	DOCUMENT ID	TECN	COMMENT
0.409 ± 0.026 OUR FIT			
0.44 ± 0.03	⁸ ABRAMS	75b	MRK1 $e^+e^- \rightarrow J/\psi X$

$\Gamma(J/\psi(1S)\text{neutrals})/\Gamma(J/\psi(1S)\pi^+\pi^-)$

$\Gamma_6/\Gamma_7 = (0.9761\Gamma_8 + 0.708\Gamma_9 + 0.273\Gamma_{30} + 0.135\Gamma_{31})/\Gamma_7$			
VALUE	DOCUMENT ID	TECN	COMMENT
0.72 ± 0.08 OUR FIT			
0.73 ± 0.09	⁸ TANENBAUM	76	MRK1 e^+e^-


$\Gamma(J/\psi(1S)\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_7/Γ			
VALUE	DOCUMENT ID	TECN	COMMENT
0.324 ± 0.028 OUR FIT			
0.332 ± 0.033 OUR AVERAGE			
0.32 ± 0.04	ABRAMS	75b	MRK1 $e^+e^- \rightarrow J/\psi\pi^+\pi^-$
0.36 ± 0.06	WIJK	75	DASP $e^+e^- \rightarrow J/\psi\pi^+\pi^-$

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma_{\text{total}}$ Γ_8/Γ			
VALUE	DOCUMENT ID	TECN	COMMENT
0.184 ± 0.027 OUR FIT			
0.18 ± 0.06	WIJK	75	DASP $e^+e^- \rightarrow J/\psi 2\pi^0$

$\Gamma(J/\psi(1S)\pi^0\pi^0)/\Gamma(J/\psi(1S)\pi^+\pi^-)$ Γ_8/Γ_7			
VALUE	DOCUMENT ID	TECN	COMMENT
0.57 ± 0.08 OUR FIT			
0.53 ± 0.06	⁹ TANENBAUM	76	MRK1 e^+e^-
0.64 ± 0.15	¹⁰ HILGER	75	SPEC e^+e^-

$\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$ Γ_9/Γ			
VALUE	EVTS	DOCUMENT ID	TECN COMMENT
0.027 ± 0.004 OUR FIT	Error includes scale factor of 1.7.		
0.027 ± 0.004 OUR AVERAGE	Error includes scale factor of 1.6. See the Ideogram below.		
0.025 ± 0.006	166	HIMEL	80 MRK2 e^+e^-
0.0218 ± 0.0014 ± 0.0035	386	OREGLIA	80 CBAL $e^+e^- \rightarrow J/\psi 2\gamma$
0.036 ± 0.005	164	BARTEL	78b CNTR e^+e^-
••• We do not use the following data for averages, fits, limits, etc. •••			
0.035 ± 0.009	17	¹¹ BRANDELIK	79b DASP $e^+e^- \rightarrow J/\psi 2\gamma$
0.043 ± 0.008	44	¹¹ TANENBAUM	76 MRK1 e^+e^-

WEIGHTED AVERAGE
 0.027 ± 0.004 (Error scaled by 1.6)



$\Gamma(J/\psi(1S)\pi^0)/\Gamma_{\text{total}}$ Γ_{10}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN COMMENT
9.7 ± 2.1 OUR AVERAGE			
15 ± 6	7	HIMEL	80 MRK2 e^+e^-
9 ± 2 ± 1	23	OREGLIA	80 CBAL $\psi(2S) \rightarrow J/\psi 2\gamma$

⁸The ABRAMS 75b measurement of Γ_6/Γ_5 and the TANENBAUM 76 result for Γ_6/Γ_7 are not independent. The TANENBAUM 76 result is used in the fit because it includes more accurate corrections for angular distributions.
⁹Not independent of the TANENBAUM 76 result for Γ_6/Γ_7 .
¹⁰Ignoring the $J/\psi(1S)\eta$ and $J/\psi(1S)\gamma\gamma$ decays.
¹¹Low statistics data removed from average.

HADRONIC DECAYS

$\Gamma(3\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{11}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN COMMENT
35 ± 16	6	FRANKLIN	83 MRK2 $e^+e^- \rightarrow \text{hadrons}$

$\Gamma(2\pi^+\pi^-)/\Gamma_{\text{total}}$ Γ_{12}/Γ			
VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN COMMENT
31 ± 7 OUR AVERAGE			
30 ± 8	42	FRANKLIN	83 MRK2 e^+e^-
35 ± 15		ABRAMS	75 MRK1 e^+e^-

Meson Full Listings

$$\psi(2S) = \psi(3685)$$

$\Gamma(\pi^+ \pi^- K^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
VALUE (units 10^{-4})				
16 ± 4	12 TANENBAUM 78	MRK1	$e^+ e^-$	

$\Gamma(\pi^+ \pi^- \rho^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
VALUE (units 10^{-4})				
8 ± 2	12 TANENBAUM 78	MRK1	$e^+ e^-$	

$\Gamma(K^+ \bar{K}^0(892)^0 \pi^- + \text{c.c.})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
VALUE (units 10^{-4})				
6.7 ± 2.5	TANENBAUM 78	MRK1	$e^+ e^-$	

$\Gamma(2\pi^+ \pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
VALUE (units 10^{-4})				
4.5 ± 1.0	TANENBAUM 78	MRK1	$e^+ e^-$	

$\Gamma(\rho^0 \pi^+ \pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
VALUE (units 10^{-4})				
4.2 ± 1.5	TANENBAUM 78	MRK1	$e^+ e^-$	

$\Gamma(\bar{\rho} \rho)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
VALUE (units 10^{-4})				
1.9 ± 0.5 OUR AVERAGE				
1.4 \pm 0.8	4 BRANDELIK 79c	DASP	$e^+ e^-$	
2.3 \pm 0.7	FELDMAN 77	MRK1	$e^+ e^-$	

$\Gamma(3\pi^+ \pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{19}/Γ
VALUE (units 10^{-4})				
1.5 ± 1.0	12 TANENBAUM 78	MRK1	$e^+ e^-$	

$\Gamma(\bar{\rho} \rho \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ
VALUE (units 10^{-4})				
1.4 ± 0.5	9 FRANKLIN 83	MRK2	$e^+ e^-$	

$\Gamma(K^+ K^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ
VALUE (units 10^{-4})				
1.0 ± 0.7	BRANDELIK 79c	DASP	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.5	90 FELDMAN 77	MRK1	$e^+ e^-$	

$\Gamma(\pi^+ \pi^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
VALUE (units 10^{-4})				
0.8 ± 0.5	BRANDELIK 79c	DASP	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.5	90 FELDMAN 77	MRK1	$e^+ e^-$	

$\Gamma(\pi^+ \pi^- \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
VALUE (units 10^{-4})				
0.85 ± 0.46	4 FRANKLIN 83	MRK2	$e^+ e^- \rightarrow$ hadrons	

$\Gamma(\Lambda^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{24}/Γ
VALUE (units 10^{-4})				
<4	90 FELDMAN 77	MRK1	$e^+ e^-$	

$\Gamma(\Xi^- \Xi^+)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{25}/Γ
VALUE (units 10^{-4})				
<2	90 FELDMAN 77	MRK1	$e^+ e^-$	

$\Gamma(\rho \pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{26}/Γ
VALUE (units 10^{-4})				
< 0.83	90 1 FRANKLIN 83	MRK2	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<10	90 BARTEL 76	CNTR	$e^+ e^-$	
<10	90 13 ABRAMS 75	MRK1	$e^+ e^-$	

$\Gamma(K^+ K^- \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{27}/Γ
VALUE (units 10^{-5})				
<2.96	90 1 FRANKLIN 83	MRK2	$e^+ e^- \rightarrow$ hadrons	

$\Gamma(K^+ \bar{K}^0(892)^- + \text{c.c.})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{28}/Γ
VALUE (units 10^{-5})				
<1.79	90 0 FRANKLIN 83	MRK2	$e^+ e^- \rightarrow$ hadrons	

¹² Assuming entirely strong decay.
¹³ Final state $\rho^0 \pi^0$.

RADIATIVE DECAYS

$\Gamma(\gamma \chi_{c0}(1P))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{29}/Γ
VALUE (units 10^{-2})				
9.3 ± 0.8 OUR AVERAGE				
9.9 \pm 0.5 \pm 0.8	14 GAISER 86	CBAL	$e^+ e^- \rightarrow \gamma X$	
7.2 \pm 2.3	14 BIDDICK 77	CNTR	$e^+ e^- \rightarrow \gamma X$	
7.5 \pm 2.6	14 WHITAKER 76	MRK1	$e^+ e^-$	

$\Gamma(\gamma \chi_{c1}(1P))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{30}/Γ
VALUE (units 10^{-2})				
8.7 ± 0.8 OUR FIT				
8.7 ± 0.8 OUR AVERAGE				
9.0 \pm 0.5 \pm 0.7	15 GAISER 86	CBAL	$e^+ e^- \rightarrow \gamma X$	
7.1 \pm 1.9	16 BIDDICK 77	CNTR	$e^+ e^- \rightarrow \gamma X$	

$\Gamma(\gamma \chi_{c2}(1P))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{31}/Γ
VALUE (units 10^{-2})				
7.8 ± 0.8 OUR FIT				
7.8 ± 0.8 OUR AVERAGE				
8.0 \pm 0.5 \pm 0.7	17 GAISER 86	CBAL	$e^+ e^- \rightarrow \gamma X$	
7.0 \pm 2.0	16 BIDDICK 77	CNTR	$e^+ e^- \rightarrow \gamma X$	

$\Gamma(\gamma \eta_c(1S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{32}/Γ
VALUE (units 10^{-2})				
0.28 ± 0.06	GAISER 86	CBAL	$e^+ e^- \rightarrow \gamma X$	

$\Gamma(\gamma \eta_c(2S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{33}/Γ
VALUE (units 10^{-2})				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.2 to 1.3	95 EDWARDS 82c	CBAL	$e^+ e^- \rightarrow \gamma X$	

$\Gamma(\gamma \pi^0)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{34}/Γ
VALUE (units 10^{-4})				
< 54	95 18 LIBERMAN 75	SPEC	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<100	90 WIJK 75	DASP	$e^+ e^-$	

$\Gamma(\gamma \eta'(958))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{35}/Γ
VALUE (units 10^{-2})				
<0.11	90 19 BARTEL 76	CNTR	$e^+ e^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.6	90 20 BRAUNSCH... 77	DASP	$e^+ e^-$	

$\Gamma(\gamma \eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{36}/Γ
VALUE (units 10^{-2})				
<0.02	90 YAMADA 77	DASP	$e^+ e^- \rightarrow 3\gamma$	

$\Gamma(\gamma \eta(1440) \rightarrow \gamma K \bar{K} \pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_{38}/Γ
VALUE (units 10^{-3})				
<0.12	90 21 SCHARRE 80	MRK1	$e^+ e^-$	
14 Angular distribution $(1 + \cos^2 \theta)$ assumed.				
15 Angular distribution $(1 - 0.189 \cos^2 \theta)$ assumed.				
16 Valid for isotropic distribution of the photon.				
17 Angular distribution $(1 - 0.052 \cos^2 \theta)$ assumed.				
18 Restated by us using $B(\psi(2S) \rightarrow \mu^+ \mu^-) = 0.0077$.				
19 The value is normalized to the branching ratio for $\Gamma(J/\psi(1S)\eta)/\Gamma_{\text{total}}$.				
20 Restated by us using total decay width 228 keV.				
21 Includes unknown branching fraction $\eta(1440) \rightarrow K \bar{K} \pi$.				

 $\psi(2S)$ REFERENCES

ARMSTRONG 93B	PR D47 772	+Bettoni, Bharadwaj+	(FNAL E760 Collab.)
PDG 92	PR D45, 1 June, Part II	+Hikasa, Barnett, Stone+	(KEK, LBL, BOST+)
ALEXANDER 89	NP B320 45	+Bonvicini, Dreil, Frey, Luth	(LBL, MICH, SLAC)
GAISER 86	PR D34 711	+Bloom, Bulos, Godfrey+	(Crystal Ball Collab.)
FRANKLIN 83	PRL 51 963	+Franklin, Feldman, Abrams, Alam+	(LBL, SLAC)
EDWARDS 82C	PRL 48 70	+Partridge, Peck+	(CIT, HARV, PRIN, STAN, SLAC)
LEMOIGNE 82	PL 1138 509	+Barate, Astbury+	(SACL, LOIC, SHMP, IND)
HIMEL 80	PRL 44 920	+Abrams, Alam, Blocker+	(SLAC, SLAC)
OREGLIA 80	PRL 45 959	+Partridge+	(SLAC, CIT, HARV, PRIN, STAN)
SCHARRE 80	PRL 97B 329	+Trilling, Abrams, Alam, Blocker+	(SLAC, LBL)
ZHOLENTZ 80	PL 96B 214	+Kurdadze, Leichuk, Mishnev+	(NOVO)
Also 81	SJNP 34 814	Zholentz, Kurdadze, Leichuk+	(NOVO)
BRANDELIK 79B	NP B160 426	+Cords+	(DASP Collab.)
BRANDELIK 79C	ZPHY C1 233	+Cords+	(DASP Collab.)
BARTEL 78B	PL 79B 492	+Dittmann, Duinker, Olsson, O'Neill+	(DESY, HEIDP)
TANENBAUM 78	PR D17 1731	+Alam, Boyarski+	(SLAC, LBL)
BIDDICK 77	PRL 38 1324	+Burnett+	(UCSD, UMD, PAVI, PRIN, SLAC, STAN)
BRAUNSCH... 77	PL 67B 249	+Braunschweig+	(DASP Collab.)
BURMESTER 77	PL 66B 395	+Criegee+	(DESY, HAMB, SIEG, WUPP)
FELDMAN 77	PRPL 33C 285	+Perl	(LBL, SLAC)
YAMADA 77	Hamburg Conf. 69		(DASP Collab.)
BARTEL 76	PL 64B 483	+Duinker, Olsson, Steffen, Heintze+	(DESY, HEIDP)
TANENBAUM 76	PL 36 402	+Abrams, Boyarski, Bulos+	(SLAC, LBL) IG
WHITAKER 76	PRL 37 1596	+Tanenbaum, Abrams, Alam+	(SLAC, LBL)
ABRAMS 75B	Stanford Symp. 25		(LBL)
ABRAMS 75B	PRL 34 1181	+Briggs, Chinowsky, Friedberg+	(LBL, SLAC)
BOYARSKI 75C	Palermo Conf. 54	+Bredendach, Bulos, Abrams, Briggs+	(SLAC, LBL)
HILGER 75	PRL 35 625	+Beron, Ford, Hofstadter, Howell+	(STAN, PENN)
LIBERMAN 75	Stanford Symp. 55		(STAN)
LUTH 75	PRL 35 1124	+Boyarski, Lynch, Bredendach+	(SLAC, LBL) JPC
WIJK 75	Stanford Symp. 69		(DESY)

See key on page 1343

Meson Full Listings

$\psi(2S) = \psi(3685), \psi(3770), \psi(4040)$

OTHER RELATED PAPERS

LEE	85	SLAC 282		(SLAC)
BARATE	83	PL 121B 449	+Bareyre, Bonamy+	(SACL, LOIC, SHMP, IND)
FRANKLIN	83B	Thesis SLAC-0254		(STAN)
AUBERT	75B	PRL 33 1624	+Becker, Biggs, Burger, Glenn+	(MIT, BNL)
BRAUNSCH...	75B	PL 57B 407	Braunschweig, Konigs+	(DASP Collab.)
CAMERINI	75	PRL 35 483	+Learned, Prepost, Ash, Anderson+	(WISC, SLAC)
FELDMAN	75B	PRL 35 821	+Jean-Marie, Sadoulet, Vannucci+	(LBL, SLAC)
GRECO	75	PL 56B 367	+Pancheri-Srivastava, Srivastava	(FRAS)
JACKSON	75	NIM 128 13	+Scharre	(LBL)
SIMPSON	75	PRL 35 699	+Beron, Ford, Hilger, Hofstadter+	(STAN, PENN)
ABRAMS	74	PRL 33 1453	+Briggs, Augustin, Boyarski+	(LBL, SLAC)

$\psi(3770)$

$I^G(J^{PC}) = ?^?(1^{--})$

$\psi(3770)$ MASS


VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3769.9 ± 2.5 OUR EVALUATION	Error includes scale factor of 1.8. From $m_{\psi(3685)}$ and mass difference below.		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3764.0 ± 5.0	¹ SCHINDLER 80 MRK2		e^+e^-
3770 ± 6.0	¹ BACINO 78 DLCO		e^+e^-
3772.0 ± 6.0	¹ RAPIDIS 77 MRK1		e^+e^-

¹ Errors include systematic common to all experiments.

$m_{\psi(3770)} - m_{\psi(2S)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
83.9 ± 2.4 OUR AVERAGE	Error includes scale factor of 1.8. See the ideogram below.		
80.0 ± 2.0	SCHINDLER 80 MRK2		e^+e^-
86.0 ± 2.0	² BACINO 78 DLCO		e^+e^-
88.0 ± 3.0	RAPIDIS 77 MRK1		e^+e^-

² SPEAR $\psi(2S)$ mass subtracted (see SCHINDLER 80).



$\psi(3770)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23.6 ± 2.7 OUR FIT	Error includes scale factor of 1.1.		
25.3 ± 2.9 OUR AVERAGE			
24.0 ± 5.0	SCHINDLER 80 MRK2		e^+e^-
24.0 ± 5.0	BACINO 78 DLCO		e^+e^-
28.0 ± 5.0	RAPIDIS 77 MRK1		e^+e^-

$\psi(3770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 $D\bar{D}$	dominant	
Γ_2 e^+e^-	$(1.12 \pm 0.17) \times 10^{-5}$	1.2

$\psi(3770)$ PARTIAL WIDTHS

VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.26 ± 0.04 OUR FIT	Error includes scale factor of 1.2.		
0.24 ± 0.05 OUR AVERAGE	Error includes scale factor of 1.2.		
0.276 ± 0.050	SCHINDLER 80 MRK2		e^+e^-
0.18 ± 0.06	BACINO 78 DLCO		e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.37 ± 0.09	³ RAPIDIS 77 MRK1		e^+e^-

³ See also $\Gamma(e^+e^-)/\Gamma_{total}$ below.

$\psi(3770)$ BRANCHING RATIOS

$\Gamma(D\bar{D})/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT
dominant	PERUZZI 77 MRK1		$e^+e^- \rightarrow D\bar{D}$

$\Gamma(e^+e^-)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT
VALUE (units 10^{-5})			
1.12 ± 0.17 OUR FIT	Error includes scale factor of 1.2.		
1.3 ± 0.2	RAPIDIS 77 MRK1		e^+e^-

$\psi(3770)$ REFERENCES

SCHINDLER 80 PR D21 2716	+Siegrist, Alam, Boyarski+	(Mark II Collab.)
BACINO 78 PRL 40 671	+Baumgarten, Birkwood+	(SLAC, UCLA, UCI)
PERUZZI 77 PRL 39 1301	+Piccolo, Feldman+	(SLAC, LBL, NINES, HAWA)
RAPIDIS 77 PRL 39 526	+Gobbi, Luke, Barbaro-Galiteri+	(Mark I Collab.)

$\psi(4040)$

$I^G(J^{PC}) = ?^?(1^{--})$

J^{PC} for the $\psi(4040)$ is known by its production in e^+e^- collisions via single-photon annihilation. I^G is not known, and the interpretation of this state as a single resonance is unclear because of the expectation of substantial threshold effects in this energy region.

$\psi(4040)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4040.0 ± 10.0	BRANDELIK 78c DASP		e^+e^-

$\psi(4040)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
52.0 ± 10.0	BRANDELIK 78c DASP		e^+e^-

$\psi(4040)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	$(1.4 \pm 0.4) \times 10^{-5}$
Γ_2 $D^0\bar{D}^0$	seen
Γ_3 $D^*(2007)^0\bar{D}^0 + c.c.$	seen
Γ_4 $D^*(2007)^0\bar{D}^*(2007)^0$	seen
Γ_5 $J/\psi(1S)$ hadrons	
Γ_6 $\mu^+\mu^-$	

$\psi(4040)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	DOCUMENT ID	TECN	COMMENT
VALUE (keV)			
0.75 ± 0.15	BRANDELIK 78c DASP		e^+e^-

$\psi(4040)$ BRANCHING RATIOS

$\Gamma(e^+e^-)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT
VALUE (units 10^{-5})			
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 1.0	FELDMAN 77 MRK1		e^+e^-

$\Gamma(D^0\bar{D}^0)/\Gamma(D^*(2007)^0\bar{D}^0 + c.c.)$	DOCUMENT ID	TECN	COMMENT
VALUE			
0.05 ± 0.03	¹ GOLDHABER 77 MRK1		e^+e^-
¹ Phase-space factor (p^3) explicitly removed.			

$\Gamma(D^*(2007)^0\bar{D}^*(2007)^0)/\Gamma(D^*(2007)^0\bar{D}^0 + c.c.)$	DOCUMENT ID	TECN	COMMENT
VALUE			
32.0 ± 12.0	² GOLDHABER 77 MRK1		e^+e^-
² Phase-space factor (p^3) explicitly removed.			

Meson Full Listings

 $\psi(4040)$, $\psi(4160)$, $\psi(4415)$ $\psi(4040)$ REFERENCES

BRANDELIK	78C	PL 76B 361	+Cords+	(DASP Collab.)
Also	79C	ZPHY C1 233	Brandelik, Cords+	(DASP Collab.)
FELDMAN	77	PRPL 33C 285	+Peri	(LBL, SLAC)
GOLDHABER	77	PL 69B 503	+Wiss, Abrams, Alam+	(LBL, SLAC)

OTHER RELATED PAPERS

HEIKKILA	84	PR D29 110	+Torqvist, Ono	(HELS, AACHT)
ONO	84	ZPHY C26 307		(ORSAY)
SIEGRIST	82	PR D26 969	+Schwitters, Alam, Chinowsky+	(SLAC, LBL)
AUGUSTIN	75	PRL 34 764	+Boyarski, Abrams, Briggs+	(SLAC, LBL)
BACCI	75	PL 58B 481	+Bidoli, Penso, Stella+	(ROMA, FRAS)
BOYARSKI	75B	PRL 34 762	+Breidenbach, Abrams, Briggs+	(SLAC, LBL)
ESPOSITO	75	PL 58B 478	+Felicetti, Peruzzi+	(FRAS, NAPL, PADO, ROMA)

 $\psi(4160)$

$$I^G(J^{PC}) = ??(1^{--})$$

J^{PC} for the $\psi(4160)$ is known by its production in e^+e^- collisions via single-photon annihilation. I^G is not known, and the interpretation of this state as a single resonance is unclear because of the expectation of substantial threshold effects in this energy region.

 $\psi(4160)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4159.0 ± 20.0	BRANDELIK	78C DASP	e^+e^-

 $\psi(4160)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
78.0 ± 20.0	BRANDELIK	78C DASP	e^+e^-

 $\psi(4160)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	$(10 \pm 4) \times 10^{-6}$

 $\psi(4160)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	DOCUMENT ID	TECN	COMMENT	Γ_1
0.77 ± 0.23	BRANDELIK	78C DASP	e^+e^-	

 $\psi(4160)$ REFERENCES

BRANDELIK	78C	PL 76B 361	+Cords+	(DASP Collab.)
-----------	-----	------------	---------	----------------

OTHER RELATED PAPERS

ONO	84	ZPHY C26 307		(ORSAY)
KIRKBY	79B	Fermilab Symp. 107		(SLAC)
BURMESTER	77	PL 66B 395	+Criegee+	(DESY, HAMB, SIEG, WUPP)

 $\psi(4415)$

$$I^G(J^{PC}) = ??(1^{--})$$

J^{PC} for the $\psi(4415)$ is known by its production in e^+e^- collisions via single-photon annihilation. I^G is not known, and the interpretation of this state as a single resonance is unclear because of the expectation of substantial threshold effects in this energy region.

 $\psi(4415)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
4415 ± 6 OUR AVERAGE			
4417.0 ± 10.0	BRANDELIK	78C DASP	e^+e^-
4414 ± 7	SIEGRIST	76 MRK1	e^+e^-
• • • We do not use the following data for averages, fits, limits, etc. • • •			
~ 4400	KNIES	77 PLUT	$e^+e^- \rightarrow \mu^+\mu^-$

 $\psi(4415)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
43 ± 15 OUR AVERAGE			Error includes scale factor of 1.8.
66.0 ± 15.0	BRANDELIK	78C DASP	e^+e^-
33 ± 10	SIEGRIST	76 MRK1	e^+e^-

 $\psi(4415)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 hadrons	dominant
Γ_2 e^+e^-	$(1.1 \pm 0.4) \times 10^{-5}$

 $\psi(4415)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	DOCUMENT ID	TECN	COMMENT	Γ_2
0.47 ± 0.10 OUR AVERAGE				
0.49 ± 0.13	BRANDELIK	78C DASP	e^+e^-	
0.44 ± 0.14	SIEGRIST	76 MRK1	e^+e^-	

 $\psi(4415)$ BRANCHING RATIOS

$\Gamma(\text{hadrons})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
dominant	SIEGRIST	76 MRK1	e^+e^-	

 $\psi(4415)$ REFERENCES

BRANDELIK	78C	PL 76B 361	+Cords+	(DASP Collab.)
KNIES	77	Hamburg Symp. 93		(PLUTO Collab.)
SIEGRIST	76	PRL 36 700	+Abrams, Boyarski, Breidenbach+	(LBL, SLAC)

OTHER RELATED PAPERS

BURMESTER	77	PL 66B 395	+Criegee+	(DESY, HAMB, SIEG, WUPP)
LUTH	77	PL 70B 120	+Pierre, Abrams, Alam, Boyarski+	(LBL, SLAC)

$b\bar{b}$ MESONS**NOTE ON WIDTH DETERMINATIONS OF THE Υ STATES**

As is the case for $J/\psi(1S)$ and $\psi(2S)$, the full widths of the bound $b\bar{b}$ states $\Upsilon(1S)$, $\Upsilon(2S)$, and $\Upsilon(3S)$ are not directly measurable, since they are much smaller than the energy resolution of the e^+e^- storage rings where these states are produced. The common indirect method to determine Γ starts from

$$\Gamma = \Gamma_{\ell\ell}/B_{\ell\ell}, \quad (1)$$

where $\Gamma_{\ell\ell}$ is one leptonic partial width and $B_{\ell\ell}$ is the corresponding branching fraction ($\ell = e, \mu, \text{ or } \tau$). One then assumes $e\text{-}\mu\text{-}\tau$ universality and uses

$$\begin{aligned} \Gamma_{\ell\ell} &= \Gamma_{ee} \\ B_{\ell\ell} &= \text{average of } B_{ee}, B_{\mu\mu}, \text{ and } B_{\tau\tau}. \end{aligned} \quad (2)$$


The electronic partial width Γ_{ee} is also not directly measurable at e^+e^- storage rings, only the combination $\Gamma_{ee}\Gamma_{\text{had}}/\Gamma$, where Γ_{had} is the hadronic partial width and

$$\Gamma_{\text{had}} + 3\Gamma_{ee} = \Gamma. \quad (3)$$

This combination is obtained experimentally from the energy-integrated hadronic cross section

$$\begin{aligned} \int_{\text{resonance}} \sigma(e^+e^- \rightarrow \Upsilon \rightarrow \text{hadrons})dE \\ = \frac{6\pi}{M^2} \frac{\Gamma_{ee}\Gamma_{\text{had}}}{\Gamma} C_r = \frac{6\pi}{M^2} \frac{\Gamma_{ee}^{(0)}\Gamma_{\text{had}}}{\Gamma} C_r^{(0)}, \end{aligned} \quad (4)$$

where M is the Υ mass, and C_r and $C_r^{(0)}$ are radiative correction factors. C_r is used for obtaining Γ_{ee} as defined in Eq. (1) and contains corrections from all orders of QED for describing $(b\bar{b}) \rightarrow e^+e^-$. The lowest order QED value $\Gamma_{ee}^{(0)}$, relevant for the comparison with potential-model calculations, is defined by the lowest order QED graph (Born term) alone and is about 7% lower than Γ_{ee} . In the past, this distinction had been

THE BOTTOMONIUM SYSTEM

The level scheme of the $b\bar{b}$ states showing experimentally established states with solid lines. Singlet states are called η_b and h_b , triplet states Υ and χ_{bJ} . In parentheses it is sufficient to give the radial quantum number and the orbital angular momentum to specify the states with all their quantum numbers. *E.g.*, $h_b(2P)$ means 2^1P_1 with $n = 2, L = 1, S = 0, J = 1, PC = +-.$ If found, D -wave states would be called $\eta_b(nD)$ and $\Upsilon_J(nD)$, with $J = 1, 2, 3$ and $n = 1, 2, 3, 4, \dots$. For the χ_b states, the spins of only the $\chi_{b2}(1P)$ and $\chi_{b1}(1P)$ have been experimentally established. The spins of the other χ_b are given as the preferred values, based on the quarkonium models. The figure also shows the observed hadronic and radiative transitions.

Meson Full Listings

Bottomonium, $\Upsilon(1S) = \Upsilon(9460)$

overlooked by some authors as pointed out by ALEXANDER 89, BARU 86, COOPER 86, KOENIGSMANN 86, and others.

The Listings give experimental results on B_{ee} , $B_{\mu\mu}$, $B_{\tau\tau}$, and $\Gamma_{ee}\Gamma_{had}/\Gamma$. The entries of the latter quantity have been re-evaluated using consistently the correction procedure of KURAEV 85. The partial width Γ_{ee} is obtained from the average values for $\Gamma_{ee}\Gamma_{had}/\Gamma$ and $B_{\ell\ell}$ using

$$\Gamma_{ee} = \frac{\Gamma_{ee}\Gamma_{had}}{\Gamma(1 - 3B_{\ell\ell})} \quad (5)$$

The total width Γ is then obtained from Eq. (1). We do not list Γ_{ee} and Γ values of individual experiments. The Γ_{ee} values in the Meson Summary Table are also those defined in Eq. (1) and no longer the lowest order quantities $\Gamma_{ee}^{(0)}$.


$\Upsilon(1S)$
 or $\Upsilon(9460)$

 $I^G(J^{PC}) = ?^?(1^{--})$

$\Upsilon(1S)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9460.37±0.21 OUR AVERAGE	Error includes scale factor of 2.7. See the Ideogram below.		
9460.60±0.09±0.05	¹ BARU	92B REDE	$e^+e^- \rightarrow$ hadrons
9460.6 ±0.4	² ARTAMONOV	84 REDE	$e^+e^- \rightarrow$ hadrons
9459.97±0.11±0.07	MACKAY	84 REDE	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9460.59±0.12	BARU	86 REDE	$e^+e^- \rightarrow$ hadrons
¹ Superseding BARU 86.			
² Value includes data of ARTAMONOV 82.			

WEIGHTED AVERAGE
9460.37±0.21 (Error scaled by 2.7)



$\Upsilon(1S)$ WIDTH

VALUE (keV)	DOCUMENT ID
52.5±1.8 OUR EVALUATION	See Υ mini-review.

$\Upsilon(1S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Γ_1 $\tau^+\tau^-$	(2.97±0.35) %	
Γ_2 e^+e^-	(2.52±0.17) %	
Γ_3 $\mu^+\mu^-$	(2.48±0.07) %	S=1.1

Hadronic decays

Γ_4 $J/\psi(1S)$ anything	(1.1 ± 0.4) × 10 ⁻³	
Γ_5 $\rho\pi$	< 2 × 10 ⁻⁴	CL=90%
Γ_6 $\pi^+\pi^-$	< 5 × 10 ⁻⁴	CL=90%
Γ_7 K^+K^-	< 5 × 10 ⁻⁴	CL=90%
Γ_8 $\rho\bar{\rho}$	< 9 × 10 ⁻⁴	CL=90%

Radiative decays

Γ_9 $\gamma 2h^+ 2h^-$	(7.0 ± 1.5) × 10 ⁻⁴	
Γ_{10} $\gamma 3h^+ 3h^-$	(5.4 ± 2.0) × 10 ⁻⁴	
Γ_{11} $\gamma 4h^+ 4h^-$	(7.4 ± 3.5) × 10 ⁻⁴	
Γ_{12} $\gamma \pi^+ \pi^- K^+ K^-$	(2.9 ± 0.9) × 10 ⁻⁴	
Γ_{13} $\gamma 2\pi^+ 2\pi^-$	(2.5 ± 0.9) × 10 ⁻⁴	
Γ_{14} $\gamma 3\pi^+ 3\pi^-$	(2.5 ± 1.2) × 10 ⁻⁴	
Γ_{15} $\gamma 2\pi^+ 2\pi^- K^+ K^-$	(2.4 ± 1.2) × 10 ⁻⁴	
Γ_{16} $\gamma \pi^+ \pi^- \rho\bar{\rho}$	(1.5 ± 0.6) × 10 ⁻⁴	
Γ_{17} $\gamma 2\pi^+ 2\pi^- \rho\bar{\rho}$	(4 ± 6) × 10 ⁻⁵	
Γ_{18} $\gamma 2K^+ 2K^-$	(2.0 ± 2.0) × 10 ⁻⁵	
Γ_{19} $\gamma \eta'(958)$	< 1.3 × 10 ⁻³	CL=90%
Γ_{20} $\gamma \eta$	< 3.5 × 10 ⁻⁴	CL=90%
Γ_{21} $\gamma f_2'(1525)$	< 1.4 × 10 ⁻⁴	CL=90%
Γ_{22} $\gamma f_2(1270)$	< 1.3 × 10 ⁻⁴	CL=90%
Γ_{23} $\gamma \eta(1440)$	< 8.2 × 10 ⁻⁵	CL=90%
Γ_{24} $\gamma f_3(1710) \rightarrow \gamma K\bar{K}$	< 2.6 × 10 ⁻⁴	CL=90%
Γ_{25} $\gamma f_4(2220) \rightarrow \gamma K^+ K^-$	< 1.5 × 10 ⁻⁵	CL=90%

Hadronic decays

Γ_{26} $D^*(2010)^\pm$ anything	[a]	
--	-----	--

[a] The value is for the sum of the charge states indicated.

$\Upsilon(1S)$ $\Gamma(I)\Gamma(e^+e^-)/\Gamma(\text{total})$

VALUE (eV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_2\Gamma_3/\Gamma$
31.2±1.6±1.7	KOBEL	92 CBAL	$e^+e^- \rightarrow \mu^+\mu^-$	

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_0\Gamma_2/\Gamma$
1.216±0.027 OUR AVERAGE				

1.187±0.023±0.031	³ BARU	92B MD1	$e^+e^- \rightarrow$ hadrons	
1.23 ± 0.02 ± 0.05	³ JAKUBOWSKI	88 CBAL	$e^+e^- \rightarrow$ hadrons	
1.37 ± 0.06 ± 0.09	⁴ GILES	84B CLEO	$e^+e^- \rightarrow$ hadrons	
1.23 ± 0.08 ± 0.04	⁴ ALBRECHT	82 DASP	$e^+e^- \rightarrow$ hadrons	
1.13 ± 0.07 ± 0.11	⁴ NICZYPORUK	82 LENA	$e^+e^- \rightarrow$ hadrons	
1.09 ± 0.25	⁴ BOCK	80 CNTR	$e^+e^- \rightarrow$ hadrons	
1.35 ± 0.14	⁵ BERGER	79 PLUT	$e^+e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.17 ± 0.06 ± 0.10	⁴ TUTS	83 CUSB	$e^+e^- \rightarrow$ hadrons	

³Radiative corrections evaluated following KURAEV 85.
⁴Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.
⁵Radiative corrections reevaluated by ALEXANDER 89 using $B(\mu\mu) = 0.026$.

$\Upsilon(1S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	DOCUMENT ID
1.32±0.03 OUR EVALUATION	See Υ mini-review.

$\Upsilon(1S)$ BRANCHING RATIOS

$\Gamma(\tau^+\tau^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.0297±0.0036 OUR AVERAGE				
0.027 ± 0.004 ± 0.002	⁶ ALBRECHT	85C ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-\tau^+\tau^-$	
0.034 ± 0.004 ± 0.004	GILES	83 CLEO	$e^+e^- \rightarrow \tau^+\tau^-$	
⁶ Using $B(\Upsilon(1S) \rightarrow ee) = B(\Upsilon(1S) \rightarrow \mu\mu) = 0.0256$; not used for width evaluations.				

$\Gamma(\mu^+\mu^-)/\Gamma_{\text{total}}$	EVIS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.0248±0.0007 OUR AVERAGE	Error includes scale factor of 1.1.				
0.0212 ± 0.0020 ± 0.0010	⁷ BARU	92 MD1	$e^+e^- \rightarrow \mu^+\mu^-$		
0.0231 ± 0.0012 ± 0.0010	⁷ KOBEL	92 CBAL	$e^+e^- \rightarrow \mu^+\mu^-$		
0.0252 ± 0.0007 ± 0.0007	CHEN	89B CLEO	$e^+e^- \rightarrow \mu^+\mu^-$		
0.0261 ± 0.0009 ± 0.0011	KAARSBERG	89 CSB2	$e^+e^- \rightarrow \mu^+\mu^-$		
0.0230 ± 0.0025 ± 0.0013	86	ALBRECHT	87 ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$	
0.029 ± 0.003 ± 0.002	864	BESSION	84 CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-\mu^+\mu^-$	

See key on page 1343

Meson Full Listings

$\Upsilon(1S) = \Upsilon(9460)$

0.027 ± 0.003 ± 0.003	ANDREWS	83	CLEO	$e^+e^- \rightarrow \mu^+\mu^-$
0.032 ± 0.013 ± 0.003	ALBRECHT	82	DASP	$e^+e^- \rightarrow \mu^+\mu^-$
0.038 ± 0.015 ± 0.002	NICZYPORUK	82	LENA	$e^+e^- \rightarrow \mu^+\mu^-$
0.014 ^{+0.034} _{-0.014}	BOCK	80	CNTR	$e^+e^- \rightarrow \mu^+\mu^-$
0.022 ± 0.020	BERGER	79	PLUT	$e^+e^- \rightarrow \mu^+\mu^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.027 ± 0.003 ± 0.003	TUTS	83	CUSB	$e^+e^- \rightarrow \mu^+\mu^-$

⁷ Taking into account interference between the resonance and continuum.

$\Gamma(e^+e^-)/\Gamma_{total}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.0252 ± 0.0017 OUR AVERAGE						
	0.0242 ± 0.0014 ± 0.0014	307	ALBRECHT	87	ARG	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
	0.028 ± 0.003 ± 0.002	826	BESSON	84	CLEO	$\Upsilon(2S) \rightarrow \pi^+\pi^-e^+e^-$
	0.051 ± 0.030		BERGER	80C	PLUT	$e^+e^- \rightarrow \mu^+\mu^-$

$\Gamma(J/\psi(1S) \text{ anything})/\Gamma_{total}$	VALUE (units 10 ⁻³)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
< 0.68	90		ALBRECHT	92J	ARG	$e^+e^- \rightarrow e^+e^-X, e^+e^- \rightarrow \mu^+\mu^-X$
1.1 ± 0.4 ± 0.2			8 FULTON	89	CLEO	$e^+e^- \rightarrow \mu^+\mu^-X$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 1.7	90		MASCHMANN	90	CBAL	$e^+e^- \rightarrow \text{hadrons}$
< 20	90		NICZYPORUK	83	LENA	
⁸ Using $B(J/\psi \rightarrow \mu^+\mu^-) = (6.9 \pm 0.9)\%$.						

$\Gamma(\pi^+\pi^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
< 5	90		BARU	92	MD1	$\Upsilon(1S) \rightarrow \pi^+\pi^-$

$\Gamma(K^+K^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
< 5	90		BARU	92	MD1	$\Upsilon(1S) \rightarrow K^+K^-$

$\Gamma(\rho\bar{\rho})/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
< 9	90		BARU	92	MD1	$\Upsilon(1S) \rightarrow \rho\bar{\rho}$

$\Gamma(\gamma 2\pi^+ 2\pi^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
2.5 ± 0.7 ± 0.5	26 ± 7		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \pi^+ \pi^- K^+ K^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
2.9 ± 0.7 ± 0.6	29 ± 8		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma \pi^+ \pi^- \rho\bar{\rho})/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{16}/Γ
1.5 ± 0.5 ± 0.3	22 ± 6		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2K^+ 2K^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{18}/Γ
0.2 ± 0.2	2 ± 2		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3\pi^+ 3\pi^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
2.5 ± 0.9 ± 0.8	17 ± 5		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- K^+ K^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
2.4 ± 0.9 ± 0.8	18 ± 7		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2\pi^+ 2\pi^- \rho\bar{\rho})/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{17}/Γ
0.4 ± 0.4 ± 0.4	7 ± 6		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 2h^+ 2h^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
7.0 ± 1.1 ± 1.0	80 ± 12		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 3h^+ 3h^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
5.4 ± 1.5 ± 1.3	39 ± 11		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\gamma 4h^+ 4h^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
7.4 ± 2.5 ± 2.5	36 ± 12		FULTON	90B	CLEO	$e^+e^- \rightarrow \text{hadrons}$

$\Gamma(\rho\pi)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
< 2	90		FULTON	90B	$\Upsilon(1S) \rightarrow \rho^0\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 10	90		BLINOV	90	MD1	$\Upsilon(1S) \rightarrow \rho^0\pi^0$
< 21	90		NICZYPORUK	83	LENA	$\Upsilon(1S) \rightarrow \rho^0\pi^0$

$\Gamma(D^*(2010)^\pm \text{ anything})/\Gamma_{total}$	VALUE (units 10 ⁻³)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{26}/Γ
< 19	90		9 ALBRECHT	92J	ARG	$e^+e^- \rightarrow D^0\pi^\pm X$
⁹ For $x_p > 0.2$.						

$\Gamma(\gamma\eta(1440))/\Gamma_{total}$	VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{23}/Γ
< 8.2	90		10 FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ \pi^\mp K_S^0$
¹⁰ Includes unknown branching ratio of $\eta(1440) \rightarrow K^\pm \pi^\mp K_S^0$.						

$\Gamma(\gamma\eta'(958))/\Gamma_{total}$	VALUE (units 10 ⁻³)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{19}/Γ
< 1.3	90		SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma\eta)/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{20}/Γ
< 3.5	90		SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$

$\Gamma(\gamma f_2'(1525))/\Gamma_{total}$	VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{21}/Γ
< 14	90		11 FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 19.4	90		11 ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
¹¹ Assuming $B(f_2'(1525) \rightarrow K\bar{K}) = 0.71$.						

$\Gamma(\gamma f_J(1710) \rightarrow \gamma K\bar{K})/\Gamma_{total}$	VALUE (units 10 ⁻⁴)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{24}/Γ
< 2.6	90		12 ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 6.3	90		12 FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 19	90		12 FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K_S^0 K_S^0$
< 8	90		13 ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
< 24	90		14 SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$
¹² Assuming $B(f_J(1710) \rightarrow K\bar{K}) = 0.38$.						
¹³ Assuming $B(f_J(1710) \rightarrow \pi\pi) = 0.04$.						
¹⁴ Assuming $B(f_J(1710) \rightarrow \eta\eta) = 0.18$.						

$\Gamma(\gamma f_2(1270))/\Gamma_{total}$	VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{22}/Γ
< 13	90		15 ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 21	90		15 FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma \pi^+ \pi^-$
< 81	90		SCHMITT	88	CBAL	$\Upsilon(1S) \rightarrow \gamma X$
¹⁵ Using $B(f_2(1270) \rightarrow \pi\pi) = 0.84$.						

$\Gamma(\gamma f_4(2220) \rightarrow \gamma K^+ K^-)/\Gamma_{total}$	VALUE (units 10 ⁻⁵)	CL%	DOCUMENT ID	TECN	COMMENT	Γ_{25}/Γ
< 1.5	90		16 FULTON	90B	CLEO	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
< 2.9	90		16 ALBRECHT	89	ARG	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
< 20	90		16 BARU	89	MD1	$\Upsilon(1S) \rightarrow \gamma K^+ K^-$
¹⁶ Including unknown branching ratio of $f_4(2220) \rightarrow K^+ K^-$.						

Meson Full Listings

$$\Upsilon(1S) = \Upsilon(9460), \chi_{b0}(1P) = \chi_{b0}(9860), \chi_{b1}(1P) = \chi_{b1}(9890)$$

$\Upsilon(1S)$ REFERENCES

ALBRECHT 92J	ZPHY C55 25	+Ehrlichmann, Hamacher+	(ARGUS Collab.)
BARU 92	ZPHY C54 229	+Bellin, Binov+	(NOVO)
BARU 92B	BINP 92-46 Preprint	+Baru, Binov, Bondar+	(NOVO)
KOBEL 92	ZPHY C53 193	+Antreasyan, Bartels, Besset+	(Crystal Ball Collab.)
BLINOV 90	PL B245 311	+Bondar+	(NOVO)
FULTON 90B	PR D41 1401	+Hempstead+	(CLEO Collab.)
MASCHMANN 90	ZPHY C46 555	+Antreasyan, Bartels, Besset+	(Crystal Ball Collab.)
ALBRECHT 89	ZPHY C42 349	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
ALEXANDER 89	NP B320 45	+Bonvicini, Dreil, Frey, Luth	(LBL, MICH, SLAC)
BARU 89	ZPHY C42 505	+Bellin, Binov+	(CLEO Collab.)
CHEN 89B	PR D39 3528	+McIwain, Miller+	(CLEO Collab.)
FULTON 89	PL B224 445	+Haas, Hempstead+	(CLEO Collab.)
KAARSBERG 89	PRL 62 2077	+Heintz+	(CUSB Collab.)
BUCHMUELLER... 88	HE e ⁺ e ⁻ Physics 412	Buchmueller, Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore			
JAKUBOWSKI 88	ZPHY C40 49	+Antreasyan, Bartels+	(Crystal Ball Collab.)
SCHMITT 88	ZPHY C40 199	+Antreasyan+	(Crystal Ball Collab.)
ALBRECHT 87	ZPHY C35 283	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
BARU 86	ZPHY C30 551	+Binov, Bondar, Bukin+	(NOVO)
ALBRECHT 85C	PL 154B 452	+Drescher, Heller+	(ARGUS Collab.)
KURAEV 85	SJNP 41 466	+Fadin	(ASCI)
Translated from YAF 41 733.			
ARTAMONOV 84	PL 137B 272	+Baru, Binov, Bondar+	(NOVO)
BESSON 84	PR D30 1433	+Green, Hicks, Namjoshi, Sannes+	(CLEO Collab.)
GILES 84B	PL 1285	+Hassard, Hempstead, Kinoshita+	(CLEO Collab.)
MACKAY 84	PR D29 2483	+Hasard, Giles, Hempstead+	(CUSB Collab.)
ANDREWS 83	PRL 50 807	+Avery, Berkelman, Cassel+	(CLEO Collab.)
GILES 83	PRL 50 877	+ (HARV, OSU, ROCH, RUTG, SYRA, VAND+)	(CLEO Collab.)
NICZYPORUK 83	ZPHY C17 197	+Jakubowski, Zeludziewicz+	(LENA Collab.)
TUTS 83	Cornell Conf. 284		(CUSB Collab.)
ALBRECHT 82	PL 116B 383	+Hofmann+	(DESY, DORT, HEIDH, LUND, ITEP)
ARTAMONOV 82	PL 116B 225	+Baru, Binov, Bondar, Bukin, Groshev+	(NOVO)
NICZYPORUK 82	ZPHY C15 299	+Folger, Bienenlein+	(LENA Collab.)
BERGER 80C	PL 93B 497	+Lackas, Raupach+	(PLUTO Collab.)
BOCK 80	ZPHY C6 125	+Blanar, Blum+	(HEIDP, MPIM, DESY, HAMB)
BERGER 79	ZPHY C1 343	+Alexander+	(PLUTO Collab.)

OTHER RELATED PAPERS

COOPER 86	Berkeley Conf. 67		(MIT)
KOENIGS... 86	DESY 86/136	Koenigsmann	(DESY)
ALBRECHT 84	PL 134B 137	+Drescher, Heller+	(ARGUS Collab.)
ARTAMONOV 84	PL 137B 272	+Baru, Binov, Bondar+	(NOVO)
ARTAMONOV 82	PL 118B 225	+Baru, Binov, Bondar, Bukin, Groshev+	(NOVO)
BERGER 78	PL 76B 243	+Alexander, Daum+	(PLUTO Collab.)
BIENLEIN 78	PL 78B 360	+Glawe, Bock, Blanar+	(DESY, HAMB, HEIDP, MPIM)
DARDEN 78	PL 76B 246	+Hofmann, Schubert+	(DESY, DORT, HEIDH, LUND)
GARELICK 78	PR D18 945	+Gauthier, Hicks, Oliver+	(NEAS, WASH, TUFTS)
KAPLAN 78	PRL 40 435	+Appel, Herb, Hom+	(STON, FNAL, COLU)
YOH 78	PRL 41 684	+Herb, Hom, Lederman+	(COLU, FNAL, STON)
COBB 77	PL 72B 273	+Iwata, Fabjan+	(BNL, CERN, SYRA, YALE)
HERB 77	PRL 39 252	+Hom, Lederman, Appel, Ito+	(COLU, FNAL, STON)
INNES 77	PRL 39 1240	+Appel, Brown, Herb, Hom+	(COLU, FNAL, STON)

$$\chi_{b0}(1P) \text{ or } \chi_{b0}(9860)$$

$$I^G(J^{PC}) = ?(0 \text{ preferred } ++)$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b0}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9859.8 ± 1.3 OUR AVERAGE			
9860.0 ± 0.5 ± 1.4	¹ ALBRECHT 85E ARG	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
9858.3 ± 1.6 ± 2.7	¹ NERNST 85 CBAL	$\Upsilon(2S) \rightarrow \gamma X$	
9864.1 ± 7 ± 1	¹ HAAS 84 CLEO	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9872.8 ± 0.7 ± 5.0	¹ KLOPFEN... 83 CUSB	$\Upsilon(2S) \rightarrow \gamma X$	
¹ From γ energy below, assuming $\Upsilon(2S)$ mass = 10023.4 MeV.			

γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
162.3 ± 1.3 OUR AVERAGE			
162.1 ± 0.5 ± 1.4	ALBRECHT 85E ARG	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
163.8 ± 1.6 ± 2.7	NERNST 85 CBAL	$\Upsilon(2S) \rightarrow \gamma X$	
158.0 ± 7 ± 1	HAAS 84 CLEO	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
149.4 ± 0.7 ± 5.0	KLOPFEN... 83 CUSB	$\Upsilon(2S) \rightarrow \gamma X$	

$\chi_{b0}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \quad \gamma \Upsilon(1S)$	< 6%	90%

$\chi_{b0}(1P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT
< 0.06	90	WALK 86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 0.11	90	PAUSS 83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

$\chi_{b0}(1P)$ REFERENCES

WALK 86	PR D34 2611	+Zschorsch+	(Crystal Ball Collab.)
ALBRECHT 85E	PL 130B 331	+Drescher, Heller+	(ARGUS Collab.)
NERNST 85	PRL 54 2195	+Antreasyan, Aschman+	(Crystal Ball Collab.)
HAAS 84	PRL 52 799	+Antreasyan, Kagan, Kass, Behrends+	(CLEO Collab.)
KLOPFEN... 83	PRL 51 160	Klopfenstein, Horstotte+	(CUSB Collab.)
PAUSS 83	PL 130B 439	+Dietl, Eigen+	(MPIM, COLU, CORN, LSU, STON)

$$\chi_{b1}(1P) \text{ or } \chi_{b1}(9890)$$

$$I^G(J^{PC}) = ?(1^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 1$ from SKWARNICKI 87.

$\chi_{b1}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9891.9 ± 0.7 OUR AVERAGE			
9890.8 ± 0.9 ± 1.3	¹ WALK 86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
9890.8 ± 0.3 ± 1.1	¹ ALBRECHT 85E ARG	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
9892.0 ± 0.8 ± 2.4	¹ NERNST 85 CBAL	$\Upsilon(2S) \rightarrow \gamma X$	
9893.6 ± 0.8 ± 1.0	¹ HAAS 84 CLEO	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
9894.4 ± 0.4 ± 3.0	¹ KLOPFEN... 83 CUSB	$\Upsilon(2S) \rightarrow \gamma X$	
9892.0 ± 3.0	¹ PAUSS 83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
¹ From γ energy below, assuming $\Upsilon(2S)$ mass = 10023.4 MeV.			

γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
130.6 ± 0.7 OUR AVERAGE			
131.7 ± 0.9 ± 1.3	WALK 86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
131.7 ± 0.3 ± 1.1	ALBRECHT 85E ARG	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
130.6 ± 0.8 ± 2.4	NERNST 85 CBAL	$\Upsilon(2S) \rightarrow \gamma X$	
129.0 ± 0.8 ± 1.0	HAAS 84 CLEO	$\Upsilon(2S) \rightarrow \text{conv. } \gamma X$	
128.1 ± 0.4 ± 3.0	KLOPFEN... 83 CUSB	$\Upsilon(2S) \rightarrow \gamma X$	
130.6 ± 3.0	PAUSS 83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

$\chi_{b1}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma \Upsilon(1S)$	(35 ± 8) %

$\chi_{b1}(1P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
0.35 ± 0.08 OUR AVERAGE			
0.32 ± 0.06 ± 0.07	WALK 86 CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
0.47 ± 0.18	KLOPFEN... 83 CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

$\chi_{b1}(1P)$ REFERENCES

SKWARNICKI 87	PRL 58 972	+Antreasyan, Besset+	(Crystal Ball Collab.)
WALK 86	PR D34 2611	+Zschorsch+	(Crystal Ball Collab.)
ALBRECHT 85E	PL 160B 331	+Drescher, Heller+	(ARGUS Collab.)
NERNST 85	PRL 54 2195	+Antreasyan, Aschman+	(Crystal Ball Collab.)
HAAS 84	PRL 52 799	+Antreasyan, Kagan, Kass, Behrends+	(CLEO Collab.)
KLOPFEN... 83	PRL 51 160	Klopfenstein, Horstotte+	(CUSB Collab.)
PAUSS 83	PL 130B 439	+Dietl, Eigen+	(MPIM, COLU, CORN, LSU, STON)

See key on page 1343

Meson Full Listings

$$\chi_{b2}(1P) = \chi_{b2}(9915), \Upsilon(2S) = \Upsilon(10023)$$

$\chi_{b2}(1P)$
or $\chi_{b2}(9915)$

$$I^G(J^{PC}) = ?^?(2^{++})$$

J needs confirmation.

Observed in radiative decay of the $\Upsilon(2S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$. $J = 2$ from SKWARNICKI 87.

$\chi_{b2}(1P)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9913.2 ± 0.6 OUR AVERAGE			
9915.8 ± 1.1 ± 1.3	¹ WALK 86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
9912.2 ± 0.3 ± 0.9	¹ ALBRECHT 85E	ARG	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
9912.4 ± 0.8 ± 2.2	¹ NERNST 85	CBAL	$\Upsilon(2S) \rightarrow \gamma X$
9913.3 ± 0.7 ± 1.0	¹ HAAS 84	CLEO	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
9914.6 ± 0.3 ± 2.0	¹ KLOPFEN... 83	CUSB	$\Upsilon(2S) \rightarrow \gamma X$
9914.0 ± 4.0	¹ PAUSS 83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

¹ From γ energy below, assuming $\Upsilon(2S)$ mass = 10023.4 MeV.

γ ENERGY IN $\Upsilon(2S)$ DECAY

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
109.6 ± 0.6 OUR AVERAGE			
107.0 ± 1.1 ± 1.3	WALK 86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$
110.6 ± 0.3 ± 0.9	ALBRECHT 85E	ARG	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
110.4 ± 0.8 ± 2.2	NERNST 85	CBAL	$\Upsilon(2S) \rightarrow \gamma X$
109.5 ± 0.7 ± 1.0	HAAS 84	CLEO	$\Upsilon(2S) \rightarrow \text{conv.} \gamma X$
108.2 ± 0.3 ± 2.0	KLOPFEN... 83	CUSB	$\Upsilon(2S) \rightarrow \gamma X$
108.8 ± 4.0	PAUSS 83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$

$\chi_{b2}(1P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma \Upsilon(1S)$	(22 ± 4) %

$\chi_{b2}(1P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.22 ± 0.04 OUR AVERAGE				
0.27 ± 0.06 ± 0.06	WALK 86	CBAL	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	
0.20 ± 0.05	KLOPFEN... 83	CUSB	$\Upsilon(2S) \rightarrow \gamma \gamma \ell^+ \ell^-$	

$\chi_{b2}(1P)$ REFERENCES

SKWARNICKI 87	PRL 58 972	+Antreasyan, Besset+	(Crystal Ball Collab.)
WALK 86	PR D34 2611	+Zichorschi+	(Crystal Ball Collab.)
ALBRECHT 85E	PL 160B 331	+Drescher, Heller+	(ARGUS Collab.)
NERNST 85	PRL 54 2195	+Antreasyan, Aschman+	(Crystal Ball Collab.)
HAAS 84	PRL 52 799	+Jensen, Kagan, Kass, Behrends+	(CLEO Collab.)
KLOPFEN... 83	PRL 51 160	Klopfenstein, Horstotte+	(CUSB Collab.)
PAUSS 83	PL 130B 439	+Dietl, Eigen+	(MPIM, COLU, CORN, LSU, STON)

$\Upsilon(2S)$
or $\Upsilon(10023)$

$$I^G(J^{PC}) = ?^?(1^{--})$$

$\Upsilon(2S)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.02330 ± 0.00031 OUR AVERAGE			
10.0236 ± 0.0005	¹ BARU 86B	REDE	$e^+ e^- \rightarrow \text{hadrons}$
10.0231 ± 0.0004	BARBER 84	REDE	$e^+ e^- \rightarrow \text{hadrons}$

¹ Reanalysis of ARTAMONOV 84.

$\Upsilon(2S)$ WIDTH

VALUE (keV)	DOCUMENT ID
44 ± 7 OUR EVALUATION	See Υ mini-review.

$\Upsilon(2S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \quad \Upsilon(1S) \pi^+ \pi^-$	(18.5 ± 0.8) %	
$\Gamma_2 \quad \Upsilon(1S) \pi^0 \pi^0$	(8.8 ± 1.1) %	
$\Gamma_3 \quad \tau^+ \tau^-$	(1.7 ± 1.6) %	
$\Gamma_4 \quad \mu^+ \mu^-$	(1.31 ± 0.21) %	
$\Gamma_5 \quad e^+ e^-$	seen	
$\Gamma_6 \quad \Upsilon(1S) \pi^0$	< 8	× 10 ⁻³ 90%
$\Gamma_7 \quad \Upsilon(1S) \eta$	< 2	× 10 ⁻³ 90%
$\Gamma_8 \quad J/\psi(1S) \text{anything}$	< 6	× 10 ⁻³ 90%

Radiative decays

$\Gamma_9 \quad \gamma \chi_{b1}(1P)$	(6.7 ± 0.9) %
$\Gamma_{10} \quad \gamma \chi_{b2}(1P)$	(6.6 ± 0.9) %
$\Gamma_{11} \quad \gamma \chi_{b0}(1P)$	(4.3 ± 1.0) %
$\Gamma_{12} \quad \gamma f_J(1710)$	< 5.9 × 10 ⁻⁴ 90%
$\Gamma_{13} \quad \gamma f_2'(1525)$	< 5.3 × 10 ⁻⁴ 90%
$\Gamma_{14} \quad \gamma f_2(1270)$	< 2.41 × 10 ⁻⁴ 90%
$\Gamma_{15} \quad \gamma f_4(2220)$	

$\Upsilon(2S) \Gamma(\ell^+ \ell^-)/\Gamma_{\text{total}}$

$\Gamma(e^+ e^-) \times \Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	$\Gamma_5 \Gamma_4/\Gamma$
6.5 ± 1.5 ± 1.0				
6.5 ± 1.5 ± 1.0	KOBEL 92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$	

$\Gamma(\text{hadrons}) \times \Gamma(e^+ e^-)/\Gamma_{\text{total}}$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_0 \Gamma_5/\Gamma$
0.554 ± 0.030 OUR AVERAGE				
0.54 ± 0.04 ± 0.02	² JAKUBOWSKI 88	CBAL	$e^+ e^- \rightarrow \text{hadrons}$	
0.58 ± 0.03 ± 0.04	³ GILES 84B	CLEO	$e^+ e^- \rightarrow \text{hadrons}$	
0.60 ± 0.12 ± 0.07	³ ALBRECHT 82	DASP	$e^+ e^- \rightarrow \text{hadrons}$	
0.54 ± 0.07 ± 0.09	³ NICZYPORUK 81C	LENA	$e^+ e^- \rightarrow \text{hadrons}$	
0.41 ± 0.18	³ BOCK 80C	CNTR	$e^+ e^- \rightarrow \text{hadrons}$	
0.59 ± 0.03 ± 0.05	³ TUTS 83	CUSB	$e^+ e^- \rightarrow \text{hadrons}$	

² Radiative corrections evaluated following KURAEV 85.

³ Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Upsilon(2S)$ BRANCHING RATIOS

$\Gamma(J/\psi(1S) \text{anything})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
< 0.006				
< 0.006	90	MASCHMANN 90	CBAL $e^+ e^- \rightarrow \text{hadrons}$	

$\Gamma(\Upsilon(1S) \pi^+ \pi^-)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.185 ± 0.008 OUR AVERAGE					
0.181 ± 0.005 ± 0.010	11.6k	ALBRECHT 87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^-$	
0.169 ± 0.040		GELPHMAN 85	CBAL	$e^+ e^- \rightarrow \pi^+ \pi^- \text{MM}$	
0.191 ± 0.012 ± 0.006		BESSION 84	CLEO	$\pi^+ \pi^- \text{MM}$	
0.189 ± 0.026		FONSECA 84	CUSB	$e^+ e^- \rightarrow \pi^+ \pi^-$	
0.21 ± 0.07	7	NICZYPORUK 81B	LENA	$\ell^+ \ell^- \pi^+ \pi^-$	

$\Gamma(\Upsilon(1S) \pi^0 \pi^0)/\Gamma_{\text{total}}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.088 ± 0.011 OUR AVERAGE					
0.095 ± 0.019 ± 0.019	25	ALBRECHT 87	ARG	$e^+ e^- \rightarrow \pi^0 \pi^0 \ell^+ \ell^-$	
0.080 ± 0.015		GELPHMAN 85	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- \pi^0 \pi^0$	
0.103 ± 0.023		FONSECA 84	CUSB	$e^+ e^- \rightarrow \ell^+ \ell^- \pi^0 \pi^0$	

$\Gamma(\tau^+ \tau^-)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.017 ± 0.015 ± 0.006				
0.017 ± 0.015 ± 0.006	HAAS 84B	CLEO	$e^+ e^- \rightarrow \tau^+ \tau^-$	

$\Gamma(\mu^+ \mu^-)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.0131 ± 0.0021 OUR AVERAGE					
0.0122 ± 0.0028 ± 0.0019		⁴ KOBEL 92	CBAL	$e^+ e^- \rightarrow \mu^+ \mu^-$	
0.0138 ± 0.0025 ± 0.0015		KAARSBERG 89	CSB2	$e^+ e^- \rightarrow \mu^+ \mu^-$	
0.009 ± 0.006 ± 0.006		⁵ ALBRECHT 85	ARG	$e^+ e^- \rightarrow \mu^+ \mu^-$	
0.018 ± 0.008 ± 0.005		HAAS 84B	CLEO	$e^+ e^- \rightarrow \mu^+ \mu^-$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

< 0.038 90 NICZYPORUK 81C LENA $e^+ e^- \rightarrow \mu^+ \mu^-$

⁴ Taking into account interference between the resonance and continuum.

⁵ Re-evaluated using $B(\Upsilon(1S) \rightarrow \mu^+ \mu^-) = 0.026$.

$\Gamma(\Upsilon(1S) \pi^0)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
< 0.008					
< 0.008	90	LURZ 87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- \gamma \gamma$	

$\Gamma(\Upsilon(1S) \eta)/\Gamma_{\text{total}}$

VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
< 0.002					
< 0.002	90	FONSECA 84	CUSB		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 0.005	90	ALBRECHT 87	ARG	$e^+ e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^- \text{MM}$	
< 0.007	90	LURZ 87	CBAL	$e^+ e^- \rightarrow \ell^+ \ell^- (\gamma \gamma, 3\pi^0)$	
< 0.010	90	BESSION 84	CLEO		

Meson Full Listings

$$\Upsilon(2S) = \Upsilon(10023), \chi_{b0}(2P) = \chi_{b0}(10235)$$

$\Gamma(\gamma\chi_{b1}(1P))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
0.067 ± 0.009 OUR AVERAGE				
0.091 ± 0.018 ± 0.022	ALBRECHT	85E	ARG $e^+e^- \rightarrow \gamma$ conv. X	
0.065 ± 0.007 ± 0.012	NERNST	85	CBAL $e^+e^- \rightarrow \gamma$ X	
0.080 ± 0.017 ± 0.016	HAAS	84	CLEO $e^+e^- \rightarrow \gamma$ conv. X	
0.059 ± 0.014	KLOPFEN...	83	CUSB $e^+e^- \rightarrow \gamma$ X	

$\Gamma(\gamma\chi_{b2}(1P))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{10}/Γ
0.066 ± 0.009 OUR AVERAGE				
0.098 ± 0.021 ± 0.024	ALBRECHT	85E	ARG $e^+e^- \rightarrow \gamma$ conv. X	
0.058 ± 0.007 ± 0.010	NERNST	85	CBAL $e^+e^- \rightarrow \gamma$ X	
0.102 ± 0.018 ± 0.021	HAAS	84	CLEO $e^+e^- \rightarrow \gamma$ conv. X	
0.061 ± 0.014	KLOPFEN...	83	CUSB $e^+e^- \rightarrow \gamma$ X	

$\Gamma(\gamma\chi_{b0}(1P))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{11}/Γ
0.043 ± 0.010 OUR AVERAGE				
0.064 ± 0.014 ± 0.016	ALBRECHT	85E	ARG $e^+e^- \rightarrow \gamma$ conv. X	
0.036 ± 0.008 ± 0.009	NERNST	85	CBAL $e^+e^- \rightarrow \gamma$ X	
0.044 ± 0.023 ± 0.009	HAAS	84	CLEO $e^+e^- \rightarrow \gamma$ conv. X	
0.035 ± 0.014	KLOPFEN...	83	CUSB $e^+e^- \rightarrow \gamma$ X	

$\Gamma(\gamma f_J(1710))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{12}/Γ
< 5.9	90	6	ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$	
< 5.9	90	7	ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$	

$\Gamma(\gamma f'_2(1525))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{13}/Γ
< 5.3	90	8	ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$	

$\Gamma(\gamma f_2(1270))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{14}/Γ
< 24.1	90	9	ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma \pi^+ \pi^-$	

$\Gamma(\gamma f_4(2220))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_{15}/Γ
< 6.8	90	10	ALBRECHT 89 ARG $\Upsilon(2S) \rightarrow \gamma K^+ K^-$	

$\Upsilon(2S)$ REFERENCES

KOBEL 92	ZPHY C53 193	+Antreasyan, Bartels, Bessel+	(Crystal Ball Collab.)
MASCHMANN 90	ZPHY C46 555	+Antreasyan, Bartels, Bessel+	(Crystal Ball Collab.)
ALBRECHT 89	ZPHY C42 349	+Boeckmann, Glaeser, Harder+	(ARGUS Collab.)
KAARSBERG 89	PRL 62 2077	+Heintz+	(CUSB Collab.)
BUCHMUEL... 88	HE e^+e^- Physics 412	Buchmueller, Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore			
JAKUBOWSKI 88	ZPHY C40 49	+Antreasyan, Bartels+	(Crystal Ball Collab.) IGJPC
ALBRECHT 87	ZPHY C35 283	+Binder, Boeckmann, Glaeser+	(ARGUS Collab.)
LURZ 87	ZPHY C36 383	+Antreasyan, Bessel+	(Crystal Ball Collab.)
BARU 86B	ZPHY C32 662	+Binov, Bondar, Bukin+	(NOVO)
ALBRECHT 85	ZPHY C28 45	+Drescher, Heller+	(ARGUS Collab.)
ALBRECHT 85E	PL 160B 331	+Drescher, Heller+	(ARGUS Collab.)
GELPHMAN 85	PR D11 2893	+Lurz, Antreasyan+	(Crystal Ball Collab.)
KURAEV 85	SJNP 41 466	+Fadin	(ASC)
Translated from YAF 41 733			
NERNST 85	PRL 54 2195	+Antreasyan, Aschman+	(Crystal Ball Collab.)
ARTAMONOV 84	PL 137B 272	+Baru, Binov, Bondar+	(NOVO)
BARBER 84	PL 135B 498	+ (DESY, ARGUS Collab., Crystal Ball Collab.)	
BESSON 84	PR D30 1433	+Green, Hicks, Namjoshi, Sannes+	(CLEO Collab.)
FONSECA 84	NP B242 31	+Mageras, Son, Dietl, Eigen+	(CUSB Collab.)
GILES 84B	PR D29 1285	+Avery, Berkman, Cassel+	(CLEO Collab.)
HAAS 84	PRL 52 799	+Jensen, Kagan, Kass, Behrends+	(CLEO Collab.)
HAAS 84B	PR D30 1996	+Jensen, Kagan, Kass, Behrends+	(CLEO Collab.)
KLOPFEN... 83	PRL 51 160	+Klopfenstein, Horstkotte+	(CUSB Collab.)
TUTS 83	Cornell Conf. 284		
ALBRECHT 82	PL 116B 383	+Hofmann+ (DESY, DORT, HEIDH, LUND, ITEP)	
NICZYPORUK 81B	PL 100B 95	+Chen, Folger, Lurz+	(LENA Collab.)
NICZYPORUK 81C	PRL 98B 169	+Chen, Vogel, Wegener+	(LENA Collab.)
BOCK 80	ZPHY C6 125	+Blonar, Blum+ (HEIDP, MPIM, DESY, HAMB)	

OTHER RELATED PAPERS

ALEXANDER 89	NP B320 45	+Bonvicini, Drell, Frey, Luth	(LBL, MICH, SLAC)
COOPER 86	Berkeley Conf. 67		(MIT)
WALK 86	PR D34 2611	+Zschorch+	(Crystal Ball Collab.)
ALBRECHT 84	PL 134B 137	+Drescher, Heller+	(ARGUS Collab.)
ARTAMONOV 84	PL 137B 272	+Baru, Binov, Bondar+	(NOVO)
ANDREWS 83	PRL 50 807	+Avery, Berkman, Cassel+	(CLEO Collab.)
GREEN 82	PRL 49 617	+Sannes, Skubic, Snyder+	(CLEO Collab.)
BIENLEIN 78	PL 78B 360	+Glawe, Bock, Blonar+ (DESY, HAMB, HEIDP, MPIM)	
DARDEEN 78	PL 76B 246	+Hofmann, Schubert+ (DESY, DORT, HEIDH, LUND)	
KAPLAN 78	PRL 40 435	+Appel, Herb, Hom+	(STON, FNAL, COLU)
YOH 78	PRL 41 684	+Herb, Hom, Lederman+	(COLU, FNAL, STON)
COBB 77	PL 72B 273	+Iwata, Fabjan+	(BNL, CERN, SYRAC, YALE)
HERB 77	PRL 39 252	+Hofmann, Appel, Ito+	(COLU, FNAL, STON)
INNES 77	PRL 39 1240	+Appel, Brown, Herb, Hom+	(COLU, FNAL, STON)

$\chi_{b0}(2P)$ or $\chi_{b0}(10235)$

$I^G(J^{PC}) = ?^?(0 \text{ preferred } ++)$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

$\chi_{b0}(2P)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.2321 ± 0.0006 OUR AVERAGE			
10.2312 ± 0.0008 ± 0.0012	¹ HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma\chi_{b0}(2P)\gamma$
10.2323 ± 0.0007	² MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma\chi_{b0}(2P)\gamma$

¹ From the average photon energy for inclusive and exclusive events and assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. Supersedes HEINTZ 91 and NARAIN 91.

² From γ energy below assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. The error on the $\Upsilon(3S)$ mass is not included in the individual measurements. It is included in the final average.

γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
122.0 ± 0.8	4959 ± 339	³ HEINTZ	92	CSB2 $e^+e^- \rightarrow \gamma\chi_{b0}(2P)\gamma$
124.6 ± 1.4	17 ± 7	⁴ HEINTZ	92	CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$
122.3 ± 0.3 ± 0.6	9903 ± 550	MORRISON	91	CLE2 $e^+e^- \rightarrow \gamma\chi_{b0}(2P)\gamma$

³ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

⁴ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

$\chi_{b0}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \gamma \Upsilon(2S)$	(4.6 ± 2.1) %
$\Gamma_2 \gamma \Upsilon(1S)$	(9 ± 6) × 10 ⁻³

$\chi_{b0}(2P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
< 0.089	90	5	CRAWFORD 92B CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$	
0.046 ± 0.020 ± 0.007	92	6	HEINTZ 92 CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$	

⁵ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma\Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+\mu^-) < 1.19 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P)\gamma) = 0.049$.

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = (6.0 \pm 0.4 \pm 0.6)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
< 0.025	90	7	CRAWFORD 92B CLE2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$	
0.009 ± 0.006 ± 0.001	92	8	HEINTZ 92 CSB2 $e^+e^- \rightarrow \ell^+\ell^-\gamma\gamma$	

⁷ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma\gamma\Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+\mu^-) < 0.63 \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \chi_{b0}(2P)\gamma) = 0.049$.

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(2P)) = (6.0 \pm 0.4 \pm 0.6)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\chi_{b0}(2P)$ REFERENCES

CRAWFORD 92B	PL B294 139	+Fulton	(CLEO II Collab.)
HEINTZ 92	PR D46 1928	+Lee, Franzini+	(CUSB II Collab.)
HEINTZ 91	PRL 66 1563	+Kaarsberg+	(CUSB Collab.)
MORRISON 91	PRL 67 1696	+Schmidt+	(CLEO II Collab.)
NARAIN 91	PRL 66 3113	+Lovelock+	(CUSB Collab.)

OTHER RELATED PAPERS

TUTS 83	Cornell Conf. 284		(CUSB Collab.)
EIGEN 82	PRL 49 1616	+Bohringer, Herb+	(CUSB Collab.)
HAN 82	PRL 49 1612	+Horstkotte, Imlay+	(CUSB Collab.)

See key on page 1343

Meson Full Listings

$$\chi_{b1}(2P) = \chi_{b1}(10255), \chi_{b2}(2P) = \chi_{b2}(10270)$$

$$\chi_{b1}(2P)$$
 or $\chi_{b1}(10255)$
 $I^G(J^{PC}) = \gamma^2(1 \text{ preferred } ++)$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b1}(2P)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.2552 ± 0.0005 OUR AVERAGE			
10.2547 ± 0.0004 ± 0.0010	¹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma X, \ell^+\ell^- \gamma \gamma$
10.2553 ± 0.0005	² MORRISON	91 CLE2	$e^+e^- \rightarrow \gamma X$

¹ From the average photon energy for inclusive and exclusive events and assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. Supersedes HEINTZ 91 and NARAIN 91.

² From γ energy below assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. The error on the $\Upsilon(3S)$ mass is not included in the individual measurements. It is included in the final evaluation.

 $m\chi_{b1}(2P) - m\chi_{b0}(2P)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23.5 ± 0.7 ± 0.7	³ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma X, \ell^+\ell^- \gamma \gamma$

³ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

 γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
99.90 ± 0.26 OUR AVERAGE				
99 ± 1	169	CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$
100.1 ± 0.4	11147 ± 462	⁴ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma X$
100.2 ± 0.5	223 ± 17	⁵ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$
99.5 ± 0.1 ± 0.5	25759 ± 510	MORRISON	91 CLE2	$e^+e^- \rightarrow \gamma X$

⁴ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

⁵ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

 $\chi_{b1}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
$\Gamma_1 \quad \gamma \Upsilon(2S)$	(21 ± 4)%	1.5
$\Gamma_2 \quad \gamma \Upsilon(1S)$	(8.5 ± 1.3)%	1.3

 $\chi_{b1}(2P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.21 ± 0.04 OUR AVERAGE			Error includes scale factor of 1.5.	
0.356 ± 0.042 ± 0.092	⁶ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	
0.199 ± 0.020 ± 0.022	⁷ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (10.23 \pm 1.20 \pm 1.26) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

⁷ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.085 ± 0.013 OUR AVERAGE			Error includes scale factor of 1.3.	
0.120 ± 0.021 ± 0.021	⁸ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	
0.080 ± 0.009 ± 0.007	⁹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(1S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (6.47 \pm 1.12 \pm 0.82) \times 10^{-4}$ and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = 0.105^{+0.003}_{-0.002} \pm 0.013$.

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b1}(2P)) = (11.5 \pm 0.5 \pm 0.5)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\chi_{b1}(2P)$ REFERENCES

CRAWFORD	92B	PL B294 139	+Fulton	(CLEO II Collab.)
HEINTZ	92	PR D46 1928	+Lee, Franzini+	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	+Kaarsberg+	(CUSB Collab.)
MORRISON	91	PRL 67 1696	+Schmidt+	(CLEO II Collab.)
NARAIN	91	PRL 66 3113	+Lovelock+	(CUSB Collab.)

OTHER RELATED PAPERS

TUTS	83	Cornell Conf. 284		(CUSB Collab.)
EIGEN	82	PRL 49 1616	+Bohringer, Herb+	(CUSB Collab.)
HAN	82	PRL 49 1612	+Horstkoette, Imlay+	(CUSB Collab.)

$$\chi_{b2}(2P)$$
 or $\chi_{b2}(10270)$
 $I^G(J^{PC}) = \gamma^2(2 \text{ preferred } ++)$
 J needs confirmation.

Observed in radiative decay of the $\Upsilon(3S)$, therefore $C = +$. Branching ratio requires E1 transition, M1 is strongly disfavored, therefore $P = +$.

 $\chi_{b2}(2P)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.2685 ± 0.0004 OUR AVERAGE			
10.2681 ± 0.0004 ± 0.0010	¹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma X, \ell^+\ell^- \gamma \gamma$
10.2685 ± 0.0004	² MORRISON	91 CLE2	$e^+e^- \rightarrow \gamma X$

¹ From the average photon energy for inclusive and exclusive events and assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. Supersedes HEINTZ 91 and NARAIN 91.

² From γ energy below, assuming $\Upsilon(3S)$ mass = 10355.3 ± 0.5 MeV. The error on the $\Upsilon(3S)$ mass is not included in the individual measurements. It is included in the final average.

 $m\chi_{b2}(2P) - m\chi_{b1}(2P)$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
13.5 ± 0.4 ± 0.5	³ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma X, \ell^+\ell^- \gamma \gamma$

³ From the average photon energy for inclusive and exclusive events. Supersedes NARAIN 91.

 γ ENERGY IN $\Upsilon(3S)$ DECAY

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
86.64 ± 0.23 OUR AVERAGE				
86 ± 1	101	CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$
86.7 ± 0.4	10319 ± 478	⁴ HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma X$
86.9 ± 0.4	157 ± 15	⁵ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$
86.4 ± 0.1 ± 0.4	30741 ± 560	MORRISON	91 CLE2	$e^+e^- \rightarrow \gamma X$

⁴ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes NARAIN 91.

⁵ A systematic uncertainty on the energy scale of 0.9% not included. Supersedes HEINTZ 91.

 $\chi_{b2}(2P)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad \gamma \Upsilon(2S)$	(16.2 ± 2.4)%
$\Gamma_2 \quad \gamma \Upsilon(1S)$	(7.1 ± 1.0)%

 $\chi_{b2}(2P)$ BRANCHING RATIOS

$\Gamma(\gamma \Upsilon(2S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.162 ± 0.024 OUR AVERAGE				
0.135 ± 0.025 ± 0.035	⁶ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	
0.173 ± 0.021 ± 0.019	⁷ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.37 \pm 0.26)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (4.98 \pm 0.94 \pm 0.62) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$.

⁷ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

$\Gamma(\gamma \Upsilon(1S))/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.071 ± 0.010 OUR AVERAGE				
0.072 ± 0.014 ± 0.013	⁸ CRAWFORD	92B CLE2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	
0.070 ± 0.010 ± 0.006	⁹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+\ell^- \gamma \gamma$	

⁸ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \Upsilon(2S)) \times 2 B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (5.03 \pm 0.94 \pm 0.63) \times 10^{-4}$, and $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = 0.135 \pm 0.003 \pm 0.017$.

⁹ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$, $B(\Upsilon(3S) \rightarrow \gamma \chi_{b2}(2P)) = (11.1 \pm 0.5 \pm 0.4)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

 $\chi_{b2}(2P)$ REFERENCES

CRAWFORD	92B	PL B294 139	+Fulton	(CLEO II Collab.)
HEINTZ	92	PR D46 1928	+Lee, Franzini+	(CUSB II Collab.)
HEINTZ	91	PRL 66 1563	+Kaarsberg+	(CUSB Collab.)
MORRISON	91	PRL 67 1696	+Schmidt+	(CLEO II Collab.)
NARAIN	91	PRL 66 3113	+Lovelock+	(CUSB Collab.)

OTHER RELATED PAPERS

TUTS	83	Cornell Conf. 284		(CUSB Collab.)
EIGEN	82	PRL 49 1616	+Bohringer, Herb+	(CUSB Collab.)
HAN	82	PRL 49 1612	+Horstkoette, Imlay+	(CUSB Collab.)

Meson Full Listings

$\Upsilon(3S) = \Upsilon(10355)$

$\Upsilon(3S)$
or $\Upsilon(10355)$

$$I^G(J^{PC}) = ?^2(1^{--})$$

$\Upsilon(3S)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.3553 ± 0.0005	¹ BARU	86B REDE	$e^+e^- \rightarrow$ hadrons

¹ Reanalysis of ARTAMONOV 84.

$\Upsilon(3S)$ WIDTH

VALUE (keV)	DOCUMENT ID
26.3 ± 3.5 OUR EVALUATION	See Υ mini-review.

$\Upsilon(3S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor
Γ_1 $\Upsilon(2S)$ anything	(10.6 ± 0.8) %	
Γ_2 $\Upsilon(2S) \pi^+ \pi^-$	(2.8 ± 0.6) %	2.2
Γ_3 $\Upsilon(2S) \pi^0 \pi^0$	(2.00 ± 0.32) %	
Γ_4 $\Upsilon(2S) \gamma \gamma$	(5.0 ± 0.7) %	
Γ_5 $\Upsilon(1S) \pi^+ \pi^-$	(4.48 ± 0.21) %	
Γ_6 $\Upsilon(1S) \pi^0 \pi^0$	(2.06 ± 0.28) %	
Γ_7 $\Upsilon(1S) \eta$		
Γ_8 $\mu^+ \mu^-$	(1.81 ± 0.17) %	
Γ_9 $e^+ e^-$	seen	
Radiative decays		
Γ_{10} $\gamma \chi_{b2}(2P)$	(11.4 ± 0.8) %	1.3
Γ_{11} $\gamma \chi_{b1}(2P)$	(11.3 ± 0.6) %	
Γ_{12} $\gamma \chi_{b0}(2P)$	(5.4 ± 0.6) %	1.1

$\Upsilon(3S) \Gamma(\ell^+ \ell^-) / \Gamma(\text{total})$

VALUE (keV)	DOCUMENT ID	TECN	COMMENT	$\Gamma_0 \Gamma_9 / \Gamma$
0.45 ± 0.03 ± 0.03	² GILES	84B CLEO	$e^+e^- \rightarrow$ hadrons	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.39 ± 0.02 ± 0.03	² TUTS	83 CUSB	$e^+e^- \rightarrow$ hadrons	


² Radiative corrections reevaluated by BUCHMUELLER 88 following KURAEV 85.

$\Upsilon(3S)$ BRANCHING RATIOS

$\Gamma(\Upsilon(2S) \text{ anything}) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
0.106 ± 0.008 OUR AVERAGE						
0.1023 ± 0.0105	4625	3,4,5	BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+ \ell^- X$	
0.111 ± 0.012	4891	4,5,6	BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$	

$\Gamma(\Upsilon(2S) \pi^+ \pi^-) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_2 / Γ
0.028 ± 0.006 OUR AVERAGE					Error includes scale factor of 2.2. See the ideogram below.	
0.0312 ± 0.0049	980	3,7	BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+ \pi^- \ell^+ \ell^-$	
0.0482 ± 0.0065 ± 0.0053	138	6	WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$	
0.0213 ± 0.0038	974	6	BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.031 ± 0.020	5		MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$	

WEIGHTED AVERAGE
0.028 ± 0.006 (Error scaled by 2.2)



$\Gamma(\Upsilon(2S) \pi^0 \pi^0) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3 / Γ
0.0200 ± 0.0032 OUR AVERAGE						
0.0216 ± 0.0039			^{7,8} BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+ \ell^- \pi^0 \pi^0$	
0.017 ± 0.005 ± 0.002		10	⁹ HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+ \ell^- \pi^0 \pi^0$	

$\Gamma(\Upsilon(2S) \gamma \gamma) / \Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4 / Γ
0.0602 ± 0.0069		⁷ BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+ \ell^- 2\gamma$	

$\Gamma(\Upsilon(1S) \pi^+ \pi^-) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_5 / Γ
0.0448 ± 0.0021 OUR AVERAGE						
0.0452 ± 0.0035	11830	4	BUTLER	94B CLE2	$e^+e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$	
0.0446 ± 0.0034 ± 0.0050	451	4	WU	93 CUSB	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$	
0.0446 ± 0.0030	11221	4	BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+ \pi^- X, \pi^+ \pi^- \ell^+ \ell^-$	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.049 ± 0.010	22		GREEN	82 CLEO	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$	
0.039 ± 0.013	26		MAGERAS	82 CUSB	$\Upsilon(3S) \rightarrow \pi^+ \pi^- \ell^+ \ell^-$	

$\Gamma(\Upsilon(1S) \pi^0 \pi^0) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_6 / Γ
0.0206 ± 0.0028 OUR AVERAGE						
0.0199 ± 0.0034	56	4	BUTLER	94B CLE2	$e^+e^- \rightarrow \ell^+ \ell^- \pi^0 \pi^0$	
0.022 ± 0.004 ± 0.003	33	10	HEINTZ	92 CSB2	$e^+e^- \rightarrow \ell^+ \ell^- \pi^0 \pi^0$	

$\Gamma(\Upsilon(1S) \eta) / \Gamma_{\text{total}}$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT	Γ_7 / Γ
< 0.0022	90		BROCK	91 CLEO	$e^+e^- \rightarrow \pi^+ \pi^- \pi^0 \ell^+ \ell^-$	

$\Gamma(\mu^+ \mu^-) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_8 / Γ
0.0181 ± 0.0017 OUR AVERAGE						
0.0202 ± 0.0019 ± 0.0033			CHEN	89B CLEO	$e^+e^- \rightarrow \mu^+ \mu^-$	
0.0173 ± 0.0015 ± 0.0011			KAARSBERG	89 CSB2	$e^+e^- \rightarrow \mu^+ \mu^-$	
0.033 ± 0.013 ± 0.007	1096		ANDREWS	83 CLEO	$e^+e^- \rightarrow \mu^+ \mu^-$	

$\Gamma(\gamma \chi_{b2}(2P)) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_{10} / Γ
0.114 ± 0.008 OUR AVERAGE					Error includes scale factor of 1.3.	
0.111 ± 0.005 ± 0.004	10319 ± 478	11	HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma$	
0.135 ± 0.003 ± 0.017	30741 ± 560		MORRISON	91 CLE2	$e^+e^- \rightarrow \gamma X$	

$\Gamma(\gamma \chi_{b1}(2P)) / \Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_{11} / Γ
0.113 ± 0.006 OUR AVERAGE						
0.115 ± 0.005 ± 0.005	11147 ± 462	11	HEINTZ	92 CSB2	$e^+e^- \rightarrow \gamma$	
0.105 ± 0.003 ± 0.013	25759 ± 510		MORRISON	91 CLE2	$e^+e^- \rightarrow \gamma X$	

See key on page 1343

Meson Full Listings

$$\Upsilon(3S) = \Upsilon(10355), \Upsilon(4S) = \Upsilon(10580), \Upsilon(10860)$$

$\Gamma(\Upsilon\chi_{b0}(2P))/\Gamma_{total}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.054 ± 0.006 OUR AVERAGE					Error includes scale factor of 1.1.
0.060 ± 0.004 ± 0.006	4959 ± 339	11	HEINTZ	92 CSB2	$e^+e^- \rightarrow \Upsilon$
0.049 ⁺ 0.003 ⁻ ± 0.006	9903 ± 550		MORRISON	91 CLE2	$e^+e^- \rightarrow \Upsilon X$

³ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) = (0.038 \pm 0.007)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) = (1/2)B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-)$.

⁴ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.48 \pm 0.06)\%$. With the assumption of $e\mu$ universality.

⁵ Using $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^+\pi^-) = (18.5 \pm 0.8)\%$.

⁶ Using $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$, $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\gamma\gamma) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.188 \pm 0.035)\%$, and $B(\Upsilon(2S) \rightarrow \Upsilon(1S)\pi^0\pi^0) \times 2B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (0.436 \pm 0.056)\%$. With the assumption of $e\mu$ universality.

⁷ From the exclusive mode.

⁸ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.31 \pm 0.21)\%$ and assuming $e\mu$ universality.

⁹ $B(\Upsilon(2S) \rightarrow \mu^+\mu^-) = (1.44 \pm 0.10)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

¹⁰ Using $B(\Upsilon(1S) \rightarrow \mu^+\mu^-) = (2.57 \pm 0.07)\%$ and assuming $e\mu$ universality. Supersedes HEINTZ 91.

¹¹ Supersedes NARAIN 91.

$\Upsilon(3S)$ REFERENCES

BUTLER	94B	PR D49 40	+Fu, Kalbfleisch, Lambrecht+	(CLEO Collab.)
WU	93	PL B301 307	+Franzini, Kanekal+	(CUSB Collab.)
HEINTZ	92	PR D46 1928	+Lee, Franzini+	(CUSB II Collab.)
BROCK	91	PR D43 1448	+Ferguson+	(CLEO Collab.)
HEINTZ	91	PRL 66 1563	+Kaarsberg+	(CUSB Collab.)
MORRISON	91	PRL 67 1696	+Schmidt+	(CLEO II Collab.)
NARAIN	91	PRL 66 3113	+Loveck+	(CUSB Collab.)
CHEN	89B	PR D39 3528	+McIlwain, Miller+	(CLEO Collab.)
KAARSBERG	89	PRL 62 2077	+Heintz+	(CUSB Collab.)
BUCHMUELLER...	88	HE e^+e^- Physics 412	Buchmueller, Cooper	(HANN, DESY, MIT)
Editors: A. Ali and P. Soeding, World Scientific, Singapore				
BARU	86B	ZPHY C32 662	+Blinov, Bondar, Bukin+	(NOVO)
KURAEV	85	SJNP 41 466	+Fadin	(ASCI)
Translated from YAF 41 733.				
ARTAMONOV	84	PL 137B 272	+Baru, Blinov, Bondar+	(NOVO)
GILES	84B	PR D29 1285	+Hassard, Hempstead, Kinoshita+	(CLEO Collab.)
ANDREWS	83	PRL 50 807	+Avery, Berkelman, Cassel+	(CLEO Collab.)
TUTS	83	Cornell Conf. 284		(CUSB Collab.)
GREEN	82	PRL 49 617	+Sannes, Skubic, Snyder+	(CLEO Collab.)
MAGERAS	82	PL 118B 453	+Herb, Imlay+	(COLU, CORN, LSU, MPIM, STON)

OTHER RELATED PAPERS

ALEXANDER	89	NP B320 45	+Bonvicini, Drell, Frey, Luth	(LBL, MICH, SLAC)
ARTAMONOV	84	PL 137B 272	+Baru, Blinov, Bondar+	(NOVO)
GILES	84B	PR D29 1285	+Hassard, Hempstead, Kinoshita+	(CLEO Collab.)
HAN	82	PRL 49 1612	+Horstotte, Imlay+	(CUSB Collab.)
PETERSON	82	PL 114B 277	+Giannini, Lee-Franzini+	(CUSB Collab.)
KAPLAN	78	PRL 40 435	+Appel, Herb, Hom+	(STON, FNAL, COLU)
YOH	78	PRL 41 684	+Herb, Hom, Lederman+	(COLU, FNAL, STON)
COBB	77	PL 72B 273	+Iwata, Fabjan+	(BNL, CERN, SYRA, YALE)
HERB	77	PRL 39 252	+Hom, Lederman, Appel, Ito+	(COLU, FNAL, STON)
INNES	77	PRL 39 1240	+Appel, Brown, Herb, Hom+	(COLU, FNAL, STON)

$\Upsilon(4S)$
or $\Upsilon(10580)$

$$I^G(J^{PC}) = ?^?(1^{--})$$

$\Upsilon(4S)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.5800 ± 0.0035	¹ BEBEK	87 CLEO	$e^+e^- \rightarrow$ hadrons
• • • We do not use the following data for averages, fits, limits, etc. • • •			
10.5774 ± 0.0010	² LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons
¹ Reanalysis of BESSON 85.			
² No systematic error given.			

$\Upsilon(4S)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23.8 ± 2.2 OUR AVERAGE			
20.0 ± 2 ± 4	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
25 ± 2.5	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

$\Upsilon(4S)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 e^+e^-	$(1.01 \pm 0.21) \times 10^{-5}$	
Γ_2 $J/\psi(3097)$ anything		
Γ_3 D^{*+} anything + c.c.	< 7.4 %	90%
Γ_4 ϕ anything	< 2.3 $\times 10^{-3}$	90%
Γ_5 $\Upsilon(1S)$ anything	< 4 $\times 10^{-3}$	90%

$\Upsilon(4S)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.24 ± 0.05 OUR AVERAGE				Error includes scale factor of 1.7.
0.192 ± 0.007 ± 0.038		BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
0.283 ± 0.037		LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

$\Upsilon(4S)$ BRANCHING RATIOS

$[\Gamma(D^{*+} \text{ anything}) + \Gamma(\text{c.c.})]/\Gamma_{total}$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.074			90	³ ALEXANDER	90C CLEO e^+e^-
³ For $x > 0.473$.					

$\Gamma(\phi \text{ anything})/\Gamma_{total}$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.0023			90	⁴ ALEXANDER	90C CLEO e^+e^-
⁴ For $x > 0.52$.					

$\Gamma(\Upsilon(1S) \text{ anything})/\Gamma_{total}$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 0.004			90	ALEXANDER	90C CLEO e^+e^-

$\Upsilon(4S)$ REFERENCES

ALEXANDER	90C	PRL 64 2226	+Artuso+	(CLEO Collab.)
BEBEK	87	PR D36 1289	+Berkelman, Blucher, Cassel+	(CLEO Collab.)
BESSON	85	PRL 54 381	+Green, Namjoshi, Sannes+	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	+Horstotte, Kiopfenstein+	(CUSB Collab.)

OTHER RELATED PAPERS

ANDREWS	80B	PRL 45 219	+Berkelman, Cabenda, Cassel+	(CLEO Collab.)
FINOCCHI...	80	PRL 45 222	Finocchiaro, Giannini, Lee-Franzini+	(CUSB Collab.)

$\Upsilon(10860)$

$$I^G(J^{PC}) = ?^?(1^{--})$$

$\Upsilon(10860)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
10.865 ± 0.008 OUR AVERAGE			Error includes scale factor of 1.1.
10.868 ± 0.006 ± 0.005	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
10.845 ± 0.020	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

$\Upsilon(10860)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
110 ± 13 OUR AVERAGE			
112.0 ± 17 ± 23	BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
110.0 ± 15.0	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

$\Upsilon(10860)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 e^+e^-	$(2.8 \pm 0.7) \times 10^{-6}$

$\Upsilon(10860)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	VALUE (keV)	DOCUMENT ID	TECN	COMMENT
0.31 ± 0.07 OUR AVERAGE				Error includes scale factor of 1.3.
0.22 ± 0.05 ± 0.07		BESSON	85 CLEO	$e^+e^- \rightarrow$ hadrons
0.365 ± 0.070		LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

$\Upsilon(10860)$ REFERENCES

BESSON	85	PRL 54 381	+Green, Namjoshi, Sannes+	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	+Horstotte, Kiopfenstein+	(CUSB Collab.)

Meson Full Listings

 $\Upsilon(11020)$, Non- $q\bar{q}$ Candidates $\Upsilon(11020)$

$$J^{PC} = ?(1^{--})$$

 $\Upsilon(11020)$ MASS

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
11.019 ± 0.008 OUR AVERAGE			
11.019 ± 0.005 ± 0.007	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
11.020 ± 0.030	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

 $\Upsilon(11020)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
79 ± 16 OUR AVERAGE			
61.0 ± 13 ± 22	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons
90.0 ± 20.0	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons

 $\Upsilon(11020)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \quad e^+e^-$	$(1.6 \pm 0.5) \times 10^{-6}$

 $\Upsilon(11020)$ PARTIAL WIDTHS

$\Gamma(e^+e^-)$	DOCUMENT ID	TECN	COMMENT	Γ_1
0.130 ± 0.030 OUR AVERAGE				
0.095 ± 0.03 ± 0.035	BESSION	85 CLEO	$e^+e^- \rightarrow$ hadrons	
0.156 ± 0.040	LOVELOCK	85 CUSB	$e^+e^- \rightarrow$ hadrons	

 $\Upsilon(11020)$ REFERENCES

BESSION	85	PRL 54 381	+Green, Namjoshi, Sannes+	(CLEO Collab.)
LOVELOCK	85	PRL 54 377	+Horstkotte, Klöpfenstein+	(CUSB Collab.)

NON- $q\bar{q}$ CANDIDATES

We include here mini-reviews and reference lists on gluonium and other non- $q\bar{q}$ candidates. See also $N\bar{N}(1100-3600)$ for possible bound states.

NOTE ON NON- $q\bar{q}$ MESONS

The existence of a gluon self coupling in QCD suggests that, in addition to conventional $q\bar{q}$ meson states, there may exist bound states including gluons: gluonia or glueballs, and hybrids ($q\bar{q}g$). Another example of non- $q\bar{q}$ mesons is multi-quark states. For detailed reviews, see, *e.g.*, HEUSCH 86, CLOSE 87, TOKI 88, GODFREY 89, BURNETT 90. Theoretical guidance for the properties of these states is somewhat contradictory, and models often differ in detailed predictions.

Among the signatures naively expected for glueballs are:

- (i) No place in $q\bar{q}$ nonets;
- (ii) Flavor-singlet couplings;
- (iii) Enhanced production in gluon-rich channels such as $J/\psi(1S)$ decay;
- (iv) Reduced $\gamma\gamma$ coupling;
- (v) Exotic quantum numbers not allowed for $q\bar{q}$ (in some cases).

However, mixing effects, and other dynamical effects such as form factors, may obscure these simple signatures. If the mixing is large, only overpopulation of the spectrum relative to the number of states predicted by the $q\bar{q}$ quark model remains as a clear signal for non-exotic non- $q\bar{q}$ states. Exotic quantum numbers ($J^{PC} = 0^{--}, 0^{+-}, 1^{-+}, 2^{+-} \dots$) would be the best signature for non- $q\bar{q}$ states.

Lattice gauge theory calculations in the quenched approximation (without quark loops) predict the lightest glueball to be a scalar with a mass of typically 1550 ± 50 MeV (BALI 93). The same calculation finds a tensor glueball mass of 2270 ± 100 MeV, and glueballs with other spin parities, which include the J^{PC} -exotics, are predicted to be still heavier. The inclusion of dynamical quarks will change the predicted masses through couplings to decay channels. Very little is known about the expected widths, which may conceivably be very large.

Hybrid mesons are $q\bar{q}$ states combined with a gluonic excitation (BARNES 82, CHANOWITZ 83, ISGUR 85). Experimentally, these are more attractive than glueballs because they span flavor nonets, and are predicted to have characteristic decay modes (LEYAOUANC 85), and a $J^{PC} = 1^{-+}$ exotic hybrid is expected in all models. The masses of the lightest hybrids are typically predicted to be in the range 1500–2000 MeV.

The third class of non- $q\bar{q}$ states are the multi-quark states, which can be either baglike or clusters of mesons (VOLOSHIN 76, JAFFE 77, GUTBROD 79). A subclass of the latter are the deuteronlike meson-meson bound states, or deusons, where the long-range pion exchange is the major source of binding (TORNQVIST 91 and 94, ERICSON 93, MANOHAR 93). Many of the best non- $q\bar{q}$ candidates discussed below lie close to

an important threshold, which suggests that they may be bound states of a meson pair. Examples include the $f_0(980)$ and $a_0(980)$ (close to the $K\bar{K}$ threshold), the $f_1(1420)$ (above $K\bar{K}^*$, thus not a bound state but perhaps a threshold enhancement), the $f_0(1525)$ and $f_2(1520)$ ($\omega\omega$ and $\rho\rho$), the $f_J(1710)$ ($K^*\bar{K}^*$), and the $\psi(4040)$ ($D^*\bar{D}^*$). Many suggestions for such mesonium candidates, involving both light and heavy quarks and binding mechanisms, have appeared recently (WEINSTEIN 90, DOVER 91, BARNES 92, DOOLEY 92).

It should be emphasized that no state has been identified unambiguously as a glueball, hybrid, or multiquark state. The candidates we discuss below are chosen because they are difficult to interpret as conventional $q\bar{q}$ states. Note, however, that we do not see it as our task to discuss theoretical interpretations of the candidates; rather, we merely catalogue the observations of possible relevance.

The scalar-meson sector: The established isoscalars with $J^{PC} = 0^{++}$ are the $f_0(980)$, $f_0(1300)$, $f_0(1370)$, and $f_0(1590)$; the $f_J(1710)$ is an established isoscalar whose spin may be 0. In the quark model, one expects two 1^3P_0 states and one 2^3P_0 ($u\bar{u} + d\bar{d}$)-like state below 1.8 GeV. Thus, there are too many well-established scalars to find a place in the quark model. From further dynamical arguments related to the production or decay, it is very likely that both the $f_0(1590)$ and the $f_J(1710)$ are non- $q\bar{q}$ resonances.

It should, however, be noted that for scalar resonances with strong S -wave thresholds, naive quark model expectations, in particular ideal mixing, must be strongly broken by unitarity. Thus, the physical scalar $q\bar{q}$ spectrum can be very much distorted from naive expectations. Another problem is that mass determinations of a very broad resonance like the $f_0(1300)$ or the $K_0^*(1430)$ are always somewhat dependent on the model and background assumptions. For a detailed discussion of this sector, see our “Note on S -wave $\pi\pi$, $K\bar{K}$, and $\eta\eta$ Interactions.”

The $f_0(1590)$, seen in π^-p reactions at 38 GeV/c (BINON 83, BINON 84C, ALDE 87, ALDE 87B), has a peculiar decay pattern for a $q\bar{q}$ state:

$$\pi^0\pi^0 : K\bar{K} : \eta\eta : \eta\eta' : 4\pi^0 = < 0.3 : < 0.6 : 1 : 2.7 : 0.8$$

The scalar glueball and the 3P_0 $s\bar{s}$ state are both expected near this mass, so we may here be seeing the effects of more than one resonance.

The $f_J(1710)$ (whose spin is uncertain) has been seen mainly in “gluon-rich” $J/\psi(1S)$ radiative decay, where it is copiously produced. In central production, the WA76 experiment (ARMSTRONG 89D) on 300 GeV/c pp interactions sees a structure at the same mass, but favors spin 2. The $f_J(1710)$ has not been seen in hadronic production ($K^-p \rightarrow K\bar{K}\Lambda$) (ASTON 88D), nor in $\gamma\gamma$ fusion. The ratio of the branching fractions in $J/\psi \rightarrow \omega f_J$ and $J/\psi \rightarrow \phi f_J$ suggests important nonstrange and strange components in this state.

The pseudoscalar sector: The established isoscalars with $J^{PC} = 0^{-+}$ are the η , $\eta(958)$, $\eta(1280)$, and $\eta(1440)$ [which

may be two pseudoscalar resonances $\eta(1410)$ and $\eta(1490)$; see the “Note on the $\eta(1440)$ ”]. In the $q\bar{q}$ model, we expect two 1^1S_0 and two 2^1S_0 pseudoscalars in the 500–1800 MeV range.

Identifying the $\eta(1280)$ with the 2^1S_0 ($u\bar{u} + d\bar{d}$) state seems natural, but it is more problematic to assign one of the two peaks in the $\eta(1440)$ region to the 2^1S_0 $s\bar{s}$ state. The $\eta(1440)$ is observed in $s\bar{s}$ -depleted reactions like $\pi^-p \rightarrow \eta\pi\pi n$ (ANDO 86) and $\pi^-p \rightarrow a_0(980)\pi p$ (CHUNG 85, BIRMAN 88), and is not seen in the $s\bar{s}$ -enriched channels like $K^-p \rightarrow K^*(892)\bar{K}\Lambda$ (ASTON 87). The fact that ANDO 86 sees the $\eta(1440)$ bump and the $\eta(1280)$ with similar intensities argues for these states being of a similar nature, *i.e.*, radial excitations of the η and $\eta'(958)$. However, as there are suggestions of two resonances in the $\eta(1440)$ structure, the experimental situation remains confused and the nature of the $\eta(1440)$ is not well understood.

The axial-vector meson sector: The $q\bar{q}$ model predicts a nonet that includes two isoscalar 1^3P_1 states with masses below about 1.6 GeV. Three such 1^{++} states are known, the $f_1(1285)$, $f_1(1420)$, and $f_1(1530)$, which suggests that one of these is a non- $q\bar{q}$ meson. The $f_1(1420)$ is the most likely candidate; see CALDWELL 89 and the “Note on the $f_1(1420)$.” The proximity of the $K\bar{K}^*$ threshold suggests this may be a dominantly $K\bar{K}^*$ mesonium resonance or threshold enhancement (LONGACRE 90, TORNVIST 91).

The tensor meson sector: The two 1^3P_2 $q\bar{q}$ states are very likely the well-known $f_2(1270)$ and $f_2'(1525)$. There are several other states, which have been suggested as $J^{PC} = 2^{++}$ non- $q\bar{q}$ candidates: the $f_2(1430)$, $f_2(1520)$, $f_J(1710)$, $f_2(1810)$, $f_2(2010)$, $f_2(2300)$, and $f_2(2340)$.

The $f_2(1520)$ is observed by the ASTERIX collaboration (MAY 89) in $p\bar{p}$ P -wave annihilation at 1565 MeV in the $\pi^+\pi^-\pi^0$ channel. Its mass is better determined in the $3\pi^0$ mode by the Crystal Barrel (ANISOVITCH 94) to be 1520 MeV, close to the $\rho\rho$ and $\omega\omega$ thresholds. It has no place in a $q\bar{q}$ scheme, since all nearby $q\bar{q}$ states are already occupied.

Similarly, the $f_J(1710)$ could be composed of $K^*\bar{K}^*$ and $\omega\phi$ (DOOLEY 92), since it lies close to these thresholds. Note that before 1991, the spin of the $f_J(1710)$ was believed to be 2, and that the subsequent spin-0 determination (CHEN 91) has not been confirmed. As already mentioned, in central production WA76 still favours spin 2 for their 1720-MeV structure.

Of the heavier states, the $f_2(1810)$ is likely to be the 2^3P_2 and the three f_2 's above 2 GeV could possibly be the 2^3P_2 $s\bar{s}$, 1^3F_2 $s\bar{s}$, and 3^3P_2 $s\bar{s}$, but a gluonium interpretation of one of the three is not excluded. These three f_2 resonances have been observed in the OZI-rule-forbidden process $\pi p \rightarrow \phi\phi n$ (ETKIN 88), which has been cited as favoring the gluonium interpretation.

A similar $\phi\phi$ mass spectrum is seen by ARMSTRONG 89B in the Ω spectrometer. The DM2 and MARK-III collaborations see threshold, $\phi\phi$ production, but favor $J^P = 0^-$, not 2^+ .

Meson Full Listings

Non- $q\bar{q}$ Candidates, Top and Fourth Generation Hadrons

In $\gamma\gamma \rightarrow 4\pi$ near the $\rho\rho$ threshold, TASSO (BRANDELIK 80B, ALTHOFF 82), MARK2 (BURKE 81), CELLO (BEHREND 84E), PLUTO (BERGER 88B), SLAC TPC (AIHARA 88), and ARGUS (ALBRECHT 91F) observe a resonance-like structure. This is dominated by $\rho^0\rho^0$, and the cross section peaks a little above the $f_2(1520)$. This process has not been explained by models in which only conventional resonances dominate. The fact that the $\gamma\gamma \rightarrow \rho^+\rho^-$ is small (ALBRECHT 91F quotes 1/4 for the $\rho^+\rho^-/\rho^0\rho^0$ ratio) requires both isospin 0 and 2 for the $\rho\rho$ system. A resonance interpretation in terms of $q^2\bar{q}^2$ states thus requires the presence of a flavor exotic $I = 2$ resonance (ACHASOV 82, 87, 90). For this $\rho\rho$ structure, the 2^{++} partial wave is found to be dominant (BERGER 88B, ALBRECHT 91F), with some 0^{++} at the low-energy end, while $J^P = 0^-$ and 2^- contribute very little.

In $\gamma\gamma \rightarrow \omega\rho$, there is also a broad enhancement that peaks near 1.6 GeV (BEHREND 91, WEGNER 91), which is probably composed of several spin parities (BEHREND 91).

Other exotic or non- $q\bar{q}$ candidates: An isovector $\phi\pi^0$ resonance at 1480 MeV has been reported by BITYUKOV 87 in $\pi^-p \rightarrow \phi\pi^0n$; see the $\rho(1450)$. Preliminary indications favor $J^{PC} = 1^{--}$, i.e. nonexotic, but the large OZI-rule violating branching ratio $\phi\pi\omega\pi$ seems peculiar for a $(u\bar{u}-d\bar{d})$ $I = 1$ $q\bar{q}$ object. However, ACHASOV 88 shows that the threshold effect from the two-step process $\rho(1600) \rightarrow K\bar{K}^* \rightarrow \pi\phi$ can violate the rule, especially near threshold. No sign of this candidate is seen in $\pi\omega$ (FUKUI 91). In addition, the small coupling to the photon makes an identification with the $\rho(1450)$ difficult (CLEGG 88). More recently DONNACHIE 93, analyzing e^+e^- annihilation and diffractive photoproduction data, suggests there may be 4-quark states near 1100 and 1300 MeV.

Another exotic candidate is the $\rho(1405)$ (ALDE 88B, IDDIR 88), seen in the GAMS experiment under the $a_2(1320)$ in $\pi^-p \rightarrow \eta\pi^0n$ with the exotic quantum numbers $J^{PC} = 1^{+-}$. Although the forward-backward asymmetry demands an $\eta\pi$ P -wave, it may be due to a nonresonant amplitude. The Crystal Barrel collaboration has reported results on the corresponding P -wave in $\eta\pi$ seen in $p\bar{p} \rightarrow \eta\pi\pi$; they see a much broader effect, which can be explained as nonresonant or as a resonance with $\Gamma \approx 600$ MeV. AOYAGI 93 also notes the $\eta\pi$ P -wave, but its interpretation is unclear.

For another possible 1^{+-} candidate, see the isosinglet $X(1910)$.

A narrow resonance, listed under $K_2^?(3100)$, has been reported at about 3100 MeV (BOURQUIN 86, ALEEVEV 93) in several $\Lambda\bar{p} +$ pions and $\bar{\Lambda}p +$ pions states. The observation of the doubly charged states $\Lambda\bar{p}\pi^-$ and $\bar{\Lambda}p\pi^+$ implies, assuming the decay is strong, $I = 3/2$, clearly not a $q\bar{q}$ state. In addition, a narrow peak is observed at about 3250 MeV, listed under $X(3250)$, in the "hidden strangeness" combinations containing a baryon-antibaryon pair (ALEEV 93). However, all these observations need confirmation.

Non- $q\bar{q}$ Candidates

OMITTED FROM SUMMARY TABLE

NON- $q\bar{q}$ CANDIDATES REFERENCES

TORNQVIST 94	ZPHY C61 525	Tornqvist	(HELS)
BALI 93	PL B309 378	+Schilling, Hulsebo, Irving, Michael+	(LVP)
DONNACHIE 93	ZP C60 1876	+Kalachnikova, Clegg	(BNL)
ERICSON 93	PL B309 426	+Karl	(CERN)
GEIGER 93	PR D47 5050	+Isgur	(TNTO)
MANOHAR 93	NP B399 17	+Wise	(MIT)
MORGAN 93	PR D48 1185	+Pennington	(RAL, DURH)
ZOU 93	PR D48 R3948	+Bugg	(LOQM)
APSIMON 92	ZPHY C56 185	+Atkinson+	(Omega Photon Collab.)
BARNES 92	PR D46 131	+Swanson	(CELLO Collab.)
BARTELSKI 92	PL B289 429	+Tatur+	(WARS)
DOOLEY 92	PL B275 478	+Swanson, Barnes	(ORNL)
LINDENBAUM 92	PL B274 492	+Longacre	(BNL)
SWANSON 92	ANP 220 73		(ORNL)
AKER 91	PL B260 249	+Amsler, Peters+	(Crystal Barrel Collab.)
ALBRECHT 91F	ZPHY C50 1	+Appun, Paulini, Funk+	(ARGUS Collab.)
BEHREND 91	ZPHY C49 401	+Criegee, Field, Franke+	(CELLO Collab.)
BEHREND 91D	B257 505	+Bussey, Ahme, Apel+	(CELLO Collab.)
CHEN 91	Hadron 91 Conf.		(Mark III Collab.)
SLAC-PUB-5669			
CONDO 91	PR D43 2787	+Handler+	(SLAC Hybrid Collab.)
DOVER 91	PR C43 379	+Gutsche, Faessler	(BNL)
FUKUI 91	PL B257 241	+Horikawa+	(SUGI, NAGO, KEK, KYOT, MIFA)
MORGAN 91B	PL B258 444	+Pennington	(RAL, DURH)
MORGAN 91B	Hadron 91 Conf.	+Pennington	(RHEL)
RAL-91-070			
TORNQVIST 91	PRL 67 556		(HELS)
WEGNER 91	ZPHY C48 393	+Olsson, Allison, Ambrus	(JADE Collab.)
ACHASOV 90	TF 20 (178)	+Shestakov	(NOVO)
BREAKSTONE 90	ZPHY C48 569	+Sharpe	(ISU, BGNA, CERN, DORT, HEIDH, WARS)
BURNETT 90	ARNPS 46 332		(RAL)
CALDWELL 90	Hadron 89 Conf. p 127		(UCSB)
GOUNARIS 90	NP B346 84	+Paschalis+	(THES)
LONGACRE 90	PR D42 874		(BNL)
TORNQVIST 90	NPBPS 21,196		(HELS)
WEINSTEIN 90	PR D41 2236	+Isgur	(TNTO)
ARMSTRONG 89B	PL B221 221	+Benayoun+(CERN, CDEF, BIRM, BARI, ATHU, CURIN+)	
ARMSTRONG 89D	PL B227 186	+Benayoun	(ATHU, BARI, BIRM, CERN, CDEF)
MAY 89	PL B225 450	+Duch, Heel+	(ASTERIX)
WEINSTEIN 89	UTP 89 03	+Isgur	(TNTO)
AIHARA 88	PR D37 28	+Alston, Avery, Barbaro-Galtieri+	(TPC-2 γ Collab.)
ALDE 88B	PL B205 397	+Binon, Boutemur+	(SERP, BELG, LANL, LAPP)
ASTON 88D	NP B301 525	+Awaji, Bienz+	(SLAC, NAGO, CINC, INUS)
BERGER 88B	ZPHY C38 521	+Klowning, Burger+	(PLUTO Collab.)
BIRMAN 88	PRL 61 1557	+Chung, Peaslee+	(BNL, FSU, IND, MASD)
CLEGG 88	ZPHY C40 313	+Donnachie	(MICH, ANC)
ETKIN 88	PL B201 566	+Foley, Lindenbaum+	(BNL, CUNY)
GOUNARIS 88	PL B213 541	+Neufeld	(CERN)
IDDIR 88	PL B205 564	+Le Yaouanc, Ono+	(ORSAY, TOKY)
SHOEMAKER 88	PR D37 1120	+Ko, Michael, Lander, Pellet+	(UCD)
SLAUGHTER 88	MPL A3 1361		(LANL)
TOKI 88	AIP Conf.		(SLAC)
TUAN 88	PL B213 537	+Ferber, Dailitz	(HAWA, ROCH, OXF)
ACHASOV 87	ZPHY C36 161	+Kamakov, Shestakov	(NOVO)
ALDE 87	PL B198 286	+Binon, Bricman+	(LANL, BRUX, SERP, LAPP)
ALDE 87B	ZPHY C36 603	+Binon, Bricman+	(LANL, BELG, SERP, LAPP)
ASTON 87	NP B292 693	+Awaji, D'Amore+	(SLAC, NAGO, CINC, INUS)
AU 87	PR D35 1633	+Morgan, Pennington	(DURH, RAL)
BITYUKOV 87	PL B188 383	+Dzhelyadin, Dorofeev, Golovkin+	(SERP)
CHANOWITZ 87	PL B187 409		(LBL)
CLOSE 87	RPP 51 833		(RHEL)
AKESSON 86	NP B264 154	+Albrow, Almede+	(Axial Field Spec. Collab.)
ALDE 86D	NP B269 485	+Binon, Bricman+	(BELG, LAPP, SERP, CERN)
ANDO 86	PRL 57 1296	+Imai+	(KEK, KYOT, NIRS, SAGA, INUS, TSUK+)
BISELLO 86	PL B179 289	+Busetto, Castro, Limentani+	(DM2 Collab.)
BOURQUIN 86	PL B172 113	+Brown+	(GEVA, RAL, HEIDP, LAUS, BRIS, CERN)
COOPER 86	Berkeley Conf. 67		(MIT)
DOVER 86	PL 57 1207		(BNL)
HEUSCH 86	Sewinkel Symposium on Multiparticle Dynamics		(SLAC)
MESHKOV 86	Aspen Winter Conf.		(NBS)
CHUNG 85	PRL 55 779	+Fernow, Boehlein+	(BNL, FLOR, IND, MASD)
ISGUR 85	PRL 54 869	+Kokorski, Patou	(TNTO)
LEVAQUANC 85	ZPHY C28 309	+Olive, Pene, Raynal, Ono	(ORSAY)
AU 84	PL 167B 229	+Morgan, Pennington	(RL)
BEHREND 84E	ZPHY C21 205	+Achenberg, Deboer+	(CELLO Collab.)
BINON 84C	NC 80A 363	+Bricman, Donskov+	(BELG, LAPP, SERP, CERN)
DOVER 84	PL 146B 103		(ORSAY)
BINON 83	NC 78A 313	+Donskov, Duteil+	(BELG, LAPP, SERP, CERN)
CHANOWITZ 83	PL 126B 225	+Sharpe	(UCB, LBL)
WEINSTEIN 83B	PR D27 588	+Isgur	(TNTO)
ACHASOV 82	PL B108 134	+Devyanin, Shestakov	(NOVO)
AIHARA 82	PR D37 28	+Alston, Avery, Barbaro-Galtieri+	(TPC Collab.)
ALTHOFF 82	ZPHY C16 13	+Boerner, Burkhardt+	(TASSO Collab.)
BARNES 82	PL B116 365	+Close	(RHEL)
BURKE 81	PL B103 153	+Abrams, Alam, Blocher+	(Mark II Collab.)
BRANDELIK 80B	PL B97 448	+Boerner, Burkhardt+	(TASSO Collab.)
GUTBROD 79	ZP C1 391	+Kramer, Rumpf	(DESY)
JAFFE 77	PR D15 267,281		(MIT)
VOLOSHIN 76	ZETF 23 369	Voloshin, Okun	(ITEP)

Searches for Top and Fourth Generation Hadrons

See the sections "Searches for t Quark" and "Searches for b' (4^{th} Generation) Quark" at the end of the QUARKS section.

N BARYONS ($S = 0, I = 1/2$)

p	1673
n	1680
N resonances	1688

 Δ BARYONS ($S = 0, I = 3/2$)

Δ resonances	1710
-------------------------------	------

 Λ BARYONS ($S = -1, I = 0$)

Λ	1728
Λ resonances	1731

 Σ BARYONS ($S = -1, I = 1$)

Σ^+	1745
Σ^0	1747
Σ^-	1748
Σ resonances	1750

 Ξ BARYONS ($S = -2, I = 1/2$)

Ξ^0	1769
Ξ^-	1771
Ξ resonances	1773

 Ω BARYONS ($S = -3, I = 0$)

Ω^-	1780
Ω resonances	1781

CHARMED BARYONS ($C = +1$)

Λ_c^+	1783
$\Lambda_c(2625)^+$	1786
$\Sigma_c(2455)$	1787
$\Sigma_c(2530)$	1787
Ξ_c^+	1788
Ξ_c^0	1788
Ω_c^0	1789

BOTTOM (BEAUTY) BARYON ($B = -1$)

Λ_b^0	1790
-------------------------	------

Notes in the Baryon Listings

Note on Nucleon Decay	1673
Note on Baryon Decay Parameters	1681
Note on N and Δ Resonances	1684
Note on Baryon Magnetic Moments	1729
Note on Λ and Σ Resonances	1731
Note on the $\Lambda(1405)$	1732
Note on $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$ Decay	1749
Note on the $\Sigma(1670)$ Region	1756
Note on Ξ Resonances	1773
Note on Charmed Baryons	1782

N BARYONS

$(S = 0, I = 1/2)$

$$p, N^+ = uud; \quad n, N^0 = udd$$

p

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

p MASS

The mass is known much more precisely in u (atomic mass units) than in MeV; see the footnote. The conversion from u to MeV, $1 u = 931.49432 \pm 0.00028$ MeV, involves the relatively poorly known electronic charge.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
938.27231 ± 0.00028	¹ COHEN	87	RVUE 1986 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
938.2796 ± 0.0027	COHEN	73	RVUE 1973 CODATA value
¹ The mass is known much more precisely in u: $m = 1.007276470 \pm 0.000000012$ u.			

p̄ MASS

See, however, the next entry in the Listings, which establishes the p̄ mass much more precisely.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
938.22 ± 0.04 OUR AVERAGE			
938.30 ± 0.13	ROBERTS	78	CNTR
938.229 ± 0.049	ROBERSON	77	CNTR
938.179 ± 0.058	HU	75	CNTR Exotic atoms
938.3 ± 0.5	BAMBERGER	70	CNTR

p̄/p MASS RATIO, $m_{\bar{p}}/m_p$

A test of CPT invariance. GABRIELSE 90 below measures the ratio of inertial masses. For a discussion of what may be inferred about the ratio of p̄ and p gravitational masses, see ERICSON 90; they obtain an upper bound of 10^{-6} – 10^{-7} for violation of the equivalence principle for p̄'s.

VALUE	DOCUMENT ID	TECN	COMMENT
0.99999977 ± 0.00000042	² GABRIELSE	90	TRAP Penning trap
² GABRIELSE 90 also measures $m_{\bar{p}}/m_{e^-} = 1836.152660 \pm 0.000083$ and $m_p/m_{e^-} = 1836.152680 \pm 0.000088$. Both are completely consistent with the 1986 CODATA (COHEN 87) value for m_p/m_{e^-} of 1836.152701 ± 0.000037 . We use the CODATA values of the proton and electron masses (they come from an overall fit to a variety of data on the fundamental constants) and don't try to take into account more recent measurements involving the masses.			

$$(m_p - m_{\bar{p}}) / m_{\text{average}}$$

A test of CPT invariance. Calculated from the p̄/p mass ratio, above.

VALUE	DOCUMENT ID
(2 ± 4) × 10⁻⁸ OUR EVALUATION	

$$|q_p + q_{\bar{p}}|/e$$

A test of CPT invariance. See also a similar test involving the electron.

VALUE	DOCUMENT ID	TECN
< 2 × 10⁻⁵	³ HUGHES	92
³ HUGHES 92 uses recent measurements of Rydberg-energy and cyclotron-frequency ratios.		

$$|q_p + q_e|/e$$

See DYLLA 73 for a summary of experiments on the neutrality of matter. See also "n CHARGE" in the neutron Listings.

VALUE (units 10 ⁻²¹)	DOCUMENT ID	COMMENT
< 1.0	⁴ DYLLA	73 Neutrality of SF ₆
• • • We do not use the following data for averages, fits, limits, etc. • • •		
< 0.8	MARINELLI	84 Magnetic levitation
⁴ Assumes that $q_n = q_p + q_e$.		

p MAGNETIC MOMENT

See the Note on Baryon Magnetic Moments in the A Listings.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
2.792847386 ± 0.00000063	COHEN	87	RVUE 1986 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2.7928456 ± 0.0000011	COHEN	73	RVUE 1973 CODATA value

p̄ MAGNETIC MOMENT

A few early results have been omitted.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
-2.800 ± 0.006 OUR AVERAGE			
-2.8005 ± 0.0090	KREISSL	88	CNTR p̄ ²⁰⁸ Pb 11 → 10 X-ray
-2.817 ± 0.048	ROBERTS	78	CNTR
-2.791 ± 0.021	HU	75	CNTR Exotic atoms

$$(\mu_p - |\mu_{\bar{p}}|) / |\mu_{\text{average}}|$$

A test of CPT invariance. Calculated from the p and p̄ magnetic moments, above.

VALUE	DOCUMENT ID
(-2.6 ± 2.9) × 10⁻³ OUR EVALUATION	

p ELECTRIC DIPOLE MOMENT

A nonzero value is forbidden by both T invariance and P invariance.

VALUE (10 ⁻²³ ecm)	EVTS	DOCUMENT ID	TECN	COMMENT
- 3.7 ± 6.3		CHO	89	NMR TI F molecules
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 400		DZUBA	85	THEO Uses ¹²⁹ Xe moment
130 ± 200		⁵ WILKENING	84	
900 ± 1400		⁶ WILKENING	84	
700 ± 900	1G	HARRISON	69	MBR Molecular beam

⁵ This WILKENING 84 value includes a finite-size effect and a magnetic effect.

⁶ This WILKENING 84 value is more cautious than the other and excludes the finite-size effect, which relies on uncertain nuclear integrals.

p ELECTRIC POLARIZABILITY $\bar{\alpha}_p$

VALUE (10 ⁻⁴ fm ³)	DOCUMENT ID	TECN	COMMENT
10.2 ± 0.9 OUR AVERAGE			
9.8 ± 0.4 ± 1.1	HALLIN	93	CNTR γp Compton scattering
10.62 ^{+1.25+1.07} _{-1.19-1.03}	ZIEGER	92	CNTR γp Compton scattering
10.9 ± 2.2 ± 1.3	⁷ FEDERSPIEL	91	CNTR γp Compton scattering
⁷ FEDERSPIEL 91 obtains for the (static) electric polarizability α_p , defined in terms of the induced electric dipole moment by $\mathbf{D} = 4\pi\epsilon_0\alpha_p\mathbf{E}$, the value $(7.0 \pm 2.2 \pm 1.3) \times 10^{-4}$ fm ³ .			

p MAGNETIC POLARIZABILITY $\bar{\beta}_p$

The electric and magnetic polarizabilities are subject to a dispersion sum-rule constraint $\bar{\alpha} + \bar{\beta} = (14.2 \pm 0.5) \times 10^{-4}$ fm³.

VALUE (10 ⁻⁴ fm ³)	DOCUMENT ID	TECN	COMMENT
4.0 ± 0.9 OUR AVERAGE			
4.4 ± 0.4 ± 1.1	HALLIN	93	CNTR γp Compton scattering
3.58 ^{+1.19+1.03} _{-1.25-1.07}	ZIEGER	92	CNTR γp Compton scattering
3.3 ± 2.2 ± 1.3	FEDERSPIEL	91	CNTR γp Compton scattering

NOTE ON NUCLEON DECAY

(by K. Nakamura, Institute for Cosmic Ray Research, University of Tokyo)

Although there was a rather long pre-GUT history in the search for nucleon decay [1], modern nucleon-decay experiments have been motivated by the SU(5) Grand Unified Theory of Georgi and Glashow [2]. GUTs provide a simple and elegant framework for the unification of strong, weak, and electromagnetic forces, a natural understanding of the Weinberg angle, an explanation of electric-charge quantization, and, above all, a prediction that the nucleon lifetime is finite.

Baryon Full Listings

p

In the minimal SU(5) GUT, nucleon decay is mediated by a supermassive gauge boson, and the dominant decay mode is $p \rightarrow e^+\pi^0$. The partial mean life for a particular mode is the total mean life τ divided by the branching fraction B for the mode. The partial mean life for the $p \rightarrow e^+\pi^0$ mode is predicted, in the minimal SU(5) GUT, to be $\tau/B = 4.5 \times 10^{29 \pm 1.7}$ yr [3]. To test this clear and striking prediction, modern nucleon-decay experiments have needed the following: a large mass in order to explore the domain of $\tau/B \gtrsim 10^{30}$ yr, a tracking capability for charged particles, a way to measure visible energy, and particle identification—at least the ability to discriminate between showering (e, γ) and nonshowering (μ, π^\pm) particles.

There are two main techniques. One uses tracking calorimetry with iron plates interleaved by tracking planes; the other uses a water Čerenkov detector. Fig. 1 compares the total and fiducial masses of various nucleon-decay detectors. The 5-year construction schedule of the 50,000-ton water-Čerenkov detector Superkamiokande began in 1991.

Candidate nucleon-decay events are those contained in the detector. Background comes from atmospheric neutrino interactions and has a rate of about $100 \text{ kton}^{-1}\text{yr}^{-1}$. The kinematical difference between nucleon decay and atmospheric neutrino interactions provides background rejection. The amount of background contamination depends upon the tightness of the kinematical cuts, which are different for the different decay modes, as well as on detector capabilities such as resolutions of energy and vertex position.

Among the most favorable decay modes to detect is $p \rightarrow e^+\pi^0$, because all the final-state particles shower and their energies are well measured. No background contamination is as




Fig. 1. Nucleon decay experiments. The open bars represent the fiducial masses, while the shaded extensions indicate the total masses. Turn-on dates are at the right. The bottom scale shows the observation limit for the partial lifetime with the assumptions of no background, 50% detection efficiency, and equal numbers of protons and neutrons in the detector material.

yet expected for this mode in the current experiments. In the absence of background, the τ/B limit is directly proportional to the detector exposure.

On the other hand, the mode $p \rightarrow K^+\bar{\nu}$ is only poorly constrained kinematically. This mode, which unfortunately is the most important one in supersymmetric (SUSY) GUTs, is thus dominated by the atmospheric-neutrino background. In such a background-dominated case, the τ/B limit only improves as the square root of the exposure time.

Fig. 2 summarizes the present limits from the three major detectors (IMB, Kamiokande, and Fréjus) for nucleon partial lifetimes in various modes involving a lepton and a meson. (For limits on other modes, see the Listings.) There is as yet no compelling experimental evidence for nucleon decay, despite the predictions. The observed number of candidate events in each mode is roughly consistent with the atmospheric-neutrino background. For the $p \rightarrow e^+\pi^0$ mode, there are no candidate events in the three experiments, and therefore the τ/B limits from these experiments simply add to give the world limit of $\tau/B(p \rightarrow e^+\pi^0) > 9 \times 10^{32}$ yr (90% confidence level). Clearly, the minimal SU(5) GUT has already been ruled out. The best background-subtracted limit for the $p \rightarrow K^+\bar{\nu}$ mode has been reported by Kamiokande: it is $\tau/B(p \rightarrow K^+\bar{\nu}) > 10^{32}$ yr (90% confidence level).




Fig. 2. The 90%-confidence-level lower limits of the nucleon partial lifetime for various nucleon decay modes into lepton + meson, obtained by the IMB (diamonds), Kamiokande (circles), and Fréjus (squares) experiments. For the actual values for these and other modes, see the Listings.

References

1. See, for example, D.H. Perkins, Ann. Rev. Nucl. and Part. Sci. **34**, 1 (1984).
2. H. Georgi and S.L. Glashow, Phys. Rev. Lett. **32**, 438 (1974).
3. M. Goldhaber and W.J. Marciano, Comm. Nucl. Part. Phys. **16**, 23 (1986);
W.J. Marciano, in *Proceedings of the 8th Workshop on Grand Unification*, Syracuse, 1987, ed. K.C. Wali (World Scientific, Singapore, 1988), p. 185.

p MEAN LIFE

A test of baryon conservation. See the "p Partial Mean Lives" section below for limits that depend on decay modes. p = proton, n = bound neutron.

LIMIT (years)	PARTICLE	DOCUMENT ID	TECN
>1.6 × 10 ²⁵	p, n	8,9 EVANS	77
• • • We do not use the following data for averages, fits, limits, etc. • • •			
>3 × 10 ²³	p	9 DIX	70 CNTR
>3 × 10 ²³	p, n	9,10 FLEROV	58
⁸ Mean lifetime of nucleons in ¹³⁰ Te nuclei.			
⁹ Converted to mean life by dividing half-life by ln(2) = 0.693.			
¹⁰ Mean lifetime of nucleons in ²³² Th nuclei.			

p̄ MEAN LIFE

The best limit by far, that of GOLDEN 79, relies, however, on a number of astrophysical assumptions. The other limits come from direct observations of stored antiprotons. See also "p̄ Partial Mean Lives" after "p Partial Mean Lives," below.

LIMIT (years)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.28			GABRIELSE	90 TRAP	Penning trap
>0.08	90	1	BELL	79 CNTR	Storage ring
>1 × 10 ⁷			GOLDEN	79 SPEC	p̄/p, cosmic rays
>3.7 × 10 ⁻³			BREGMAN	78 CNTR	Storage ring

p DECAy MODES

For N decays, p and n distinguish proton and neutron partial lifetimes. See also the "Note on Proton Mean Life Limits" in these Full Listings.

The "partial mean life" limits tabulated here are the limits on τ/B_i, where τ is the total mean life and B_i is the branching fraction for the mode in question.

Mode	Partial mean life (10 ³⁰ years)	Confidence level
Antilepton + meson		
τ ₁ N → e ⁺ π	> 130 (n), > 550 (p)	90%
τ ₂ N → μ ⁺ π	> 100 (n), > 270 (p)	90%
τ ₃ N → νπ	> 100 (n), > 25 (p)	90%
τ ₄ p → e ⁺ η	> 140	90%
τ ₅ p → μ ⁺ η	> 69	90%
τ ₆ n → νη	> 54	90%
τ ₇ N → e ⁺ ρ	> 58 (n), > 75 (p)	90%
τ ₈ N → μ ⁺ ρ	> 23 (n), > 110 (p)	90%
τ ₉ N → νρ	> 19 (n), > 27 (p)	90%
τ ₁₀ p → e ⁺ ω	> 45	90%
τ ₁₁ p → μ ⁺ ω	> 57	90%
τ ₁₂ n → νω	> 43	90%
τ ₁₃ N → e ⁺ K	> 1.3 (n), > 150 (p)	90%
τ ₁₄ p → e ⁺ K _S ⁰	> 76	90%
τ ₁₅ p → e ⁺ K _L ⁰	> 44	90%
τ ₁₆ N → μ ⁺ K	> 1.1 (n), > 120 (p)	90%
τ ₁₇ p → μ ⁺ K _S ⁰	> 64	90%
τ ₁₈ p → μ ⁺ K _L ⁰	> 44	90%
τ ₁₉ N → νK	> 86 (n), > 100 (p)	90%
τ ₂₀ p → e ⁺ K*(892) ⁰	> 52	90%
τ ₂₁ N → νK*(892)	> 22 (n), > 20 (p)	90%

Antilepton + mesons

τ ₂₂ p → e ⁺ π ⁺ π ⁻	> 21	90%
τ ₂₃ p → e ⁺ π ⁰ π ⁰	> 38	90%
τ ₂₄ n → e ⁺ π ⁻ π ⁰	> 32	90%
τ ₂₅ p → μ ⁺ π ⁺ π ⁻	> 17	90%
τ ₂₆ p → μ ⁺ π ⁰ π ⁰	> 33	90%
τ ₂₇ n → μ ⁺ π ⁻ π ⁰	> 33	90%
τ ₂₈ n → e ⁺ K ⁰ π ⁻	> 18	90%

Lepton + meson

τ ₂₉ n → e ⁻ π ⁺	> 65	90%
τ ₃₀ n → μ ⁻ π ⁺	> 49	90%
τ ₃₁ n → e ⁻ ρ ⁺	> 62	90%
τ ₃₂ n → μ ⁻ ρ ⁺	> 7	90%
τ ₃₃ n → e ⁻ K ⁺	> 32	90%
τ ₃₄ n → μ ⁻ K ⁺	> 57	90%

Lepton + mesons

τ ₃₅ p → e ⁻ π ⁺ π ⁺	> 30	90%
τ ₃₆ n → e ⁻ π ⁺ π ⁰	> 29	90%
τ ₃₇ p → μ ⁻ π ⁺ π ⁺	> 17	90%
τ ₃₈ n → μ ⁻ π ⁺ π ⁰	> 34	90%
τ ₃₉ p → e ⁻ π ⁺ K ⁺	> 20	90%
τ ₄₀ p → μ ⁻ π ⁺ K ⁺	> 5	90%

Antilepton + photon(s)

τ ₄₁ p → e ⁺ γ	> 460	90%
τ ₄₂ p → μ ⁺ γ	> 380	90%
τ ₄₃ n → νγ	> 24	90%
τ ₄₄ p → e ⁺ γγ	> 100	90%

Three leptons

τ ₄₅ p → e ⁺ e ⁺ e ⁻	> 510	90%
τ ₄₆ p → e ⁺ μ ⁺ μ ⁻	> 81	90%
τ ₄₇ p → e ⁺ νν	> 11	90%
τ ₄₈ n → e ⁺ e ⁻ ν	> 74	90%
τ ₄₉ n → μ ⁺ e ⁻ ν	> 47	90%
τ ₅₀ n → μ ⁺ μ ⁻ ν	> 42	90%
τ ₅₁ p → μ ⁺ e ⁺ e ⁻	> 91	90%
τ ₅₂ p → μ ⁺ μ ⁺ μ ⁻	> 190	90%
τ ₅₃ p → μ ⁺ νν	> 21	90%
τ ₅₄ p → e ⁻ μ ⁺ μ ⁺	> 6	90%
τ ₅₅ n → 3ν	> 0.0005	90%

Inclusive modes

τ ₅₆ N → e ⁺ anything	> 0.6 (n, p)	90%
τ ₅₇ N → μ ⁺ anything	> 12 (n, p)	90%
τ ₅₈ N → νanything		
τ ₅₉ N → e ⁺ π ⁰ anything	> 0.6 (n, p)	90%
τ ₆₀ N → 2 bodies, ν-free		

ΔB = 2 dinucleon modes

The following are lifetime limits per iron nucleus.

τ ₆₁ pp → π ⁺ π ⁺	> 0.7	90%
τ ₆₂ pn → π ⁺ π ⁰	> 2	90%
τ ₆₃ nn → π ⁺ π ⁻	> 0.7	90%
τ ₆₄ nn → π ⁰ π ⁰	> 3.4	90%
τ ₆₅ pp → e ⁺ e ⁺	> 5.8	90%
τ ₆₆ pp → e ⁺ μ ⁺	> 3.6	90%
τ ₆₇ pp → μ ⁺ μ ⁺	> 1.7	90%
τ ₆₈ pn → e ⁺ ν̄	> 2.8	90%
τ ₆₉ pn → μ ⁺ ν̄	> 1.6	90%
τ ₇₀ nn → ν _e ν̄ _e	> 0.000012	90%
τ ₇₁ nn → ν _μ ν̄ _μ	> 0.000006	90%

p̄ DECAy MODES

Mode	Partial mean life (years)	Confidence level
τ ₇₂ p̄ → e ⁻ γ	> 1848	95%
τ ₇₃ p̄ → e ⁻ π ⁰	> 554	95%
τ ₇₄ p̄ → e ⁻ η	> 171	95%
τ ₇₅ p̄ → e ⁻ K _S ⁰	> 29	95%
τ ₇₆ p̄ → e ⁻ K _L ⁰	> 9	95%

Baryon Full Listings

p

p PARTIAL MEAN LIVES

The "partial mean life" limits tabulated here are the limits on τ/B_j , where τ is the total mean life for the proton and B_j is the branching fraction for the mode in question.

Decaying particle: p = proton, n = bound neutron. The same event may appear under more than one partial decay mode. Background estimates may be accurate to a factor of two.

$\tau(N \rightarrow e^+ \pi)$

τ_1

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>550	p	90	0	0.7	11 BECKER-SZ...	90 IMB3
>130	n	90	0	<0.2	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 70	p	90	0	0.5	BERGER	91 FREJ
> 70	n	90	0	≤ 0.1	BERGER	91 FREJ
>260	p	90	0	<0.04	HIRATA	89C KAMI
>310	p	90	0	0.6	SEIDEL	88 IMB
>100	n	90	0	1.6	SEIDEL	88 IMB
> 1.3	n	90	0		BARTELT	87 SOUD
> 1.3	p	90	0		BARTELT	87 SOUD
>250	p	90	0	0.3	HAINES	86 IMB
> 31	n	90	8	9	HAINES	86 IMB
> 64	p	90	0	<0.4	ARISAKA	85 KAMI
> 26	n	90	0	<0.7	ARISAKA	85 KAMI
> 82	p (free)	90	0	0.2	BLEWITT	85 IMB
>250	p	90	0	0.2	BLEWITT	85 IMB
> 25	n	90	4	4	PARK	85 IMB
> 15	p, n	90	0		BATTISTONI	84 NUSX
> 0.5	p	90	1	0.3	12 BARTELT	83 SOUD
> 0.5	n	90	1	0.3	12 BARTELT	83 SOUD
> 5.8	p	90	2		13 KRISHNA...	82 KOLR
> 5.8	n	90	2		13 KRISHNA...	82 KOLR
> 0.1	n	90			14 GURR	67 CNTR

11 This BECKER-SZENDY 90 result includes data from SEIDEL 88.

12 Limit based on zero events.

13 We have calculated 90% CL limit from 1 confined event.

14 We have converted half-life to 90% CL mean life.

$\tau(N \rightarrow \mu^+ \pi)$

τ_2

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>100	n	90	0	<0.2	HIRATA	89C KAMI
>270	p	90	0	0.5	SEIDEL	88 IMB

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 81	p	90	0	0.2	BERGER	91 FREJ
> 35	n	90	1	1.0	BERGER	91 FREJ
>230	p	90	0	<0.07	HIRATA	89C KAMI
> 63	n	90	0	0.5	SEIDEL	88 IMB
> 76	p	90	2	1	HAINES	86 IMB
> 23	n	90	8	7	HAINES	86 IMB
> 46	p	90	0	<0.7	ARISAKA	85 KAMI
> 20	n	90	0	<0.4	ARISAKA	85 KAMI
> 59	p (free)	90	0	0.2	BLEWITT	85 IMB
>100	p	90	1	0.4	BLEWITT	85 IMB
> 38	n	90	1	4	PARK	85 IMB
> 10	p, n	90	0		BATTISTONI	84 NUSX
> 1.3	p, n	90	0		ALEKSEEV	81 BAKS

$\tau(N \rightarrow \nu \pi)$

τ_3

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
> 25	p	90	32	32.8	HIRATA	89C KAMI
>100	n	90	1	3	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 13	n	90	1	1.2	BERGER	89 FREJ
> 10	p	90	11	14	BERGER	89 FREJ
> 6	n	90	73	60	HAINES	86 IMB
> 2	p	90	16	13	KAJITA	86 KAMI
> 40	n	90	0	1	KAJITA	86 KAMI
> 7	n	90	28	19	PARK	85 IMB
> 7	n	90	0		BATTISTONI	84 NUSX
> 2	p	90	≤ 3		BATTISTONI	84 NUSX
> 5.8	p	90	1		15 KRISHNA...	82 KOLR
> 0.3	p	90	2		16 CHERRY	81 HOME
> 0.1	p	90			17 GURR	67 CNTR

15 We have calculated 90% CL limit from 1 confined event.

16 We have converted 2 possible events to 90% CL limit.

17 We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow e^+ \eta)$

τ_4

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>140	p	90	0	<0.04	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 44	p	90	0	0.1	BERGER	91 FREJ
>100	p	90	0	0.6	SEIDEL	88 IMB
>200	p	90	5	3.3	HAINES	86 IMB
> 64	p	90	0	<0.8	ARISAKA	85 KAMI
> 64	p (free)	90	5	6.5	BLEWITT	85 IMB
>200	p	90	5	4.7	BLEWITT	85 IMB
> 1.2	p	90	2		18 CHERRY	81 HOME

18 We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow \mu^+ \eta)$

τ_5

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>69	p	90	1	<0.08	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

>26	p	90	1	0.8	BERGER	91 FREJ
> 1.3	p	90	0	0.7	PHILLIPS	89 HPW
>34	p	90	1	1.5	SEIDEL	88 IMB
>46	p	90	7	6	HAINES	86 IMB
>26	p	90	1	<0.8	ARISAKA	85 KAMI
>17	p (free)	90	6	6	BLEWITT	85 IMB
>46	p	90	7	8	BLEWITT	85 IMB

$\tau(n \rightarrow \nu \eta)$

τ_6

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>54	n	90	2	0.9	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

>29	n	90	0	0.9	BERGER	89 FREJ
>16	n	90	3	2.1	SEIDEL	88 IMB
>25	n	90	7	6	HAINES	86 IMB
>30	n	90	0	0.4	KAJITA	86 KAMI
>18	n	90	4	3	PARK	85 IMB
> 0.6	n	90	2		19 CHERRY	81 HOME

19 We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow e^+ \rho)$

τ_7

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>75	p	90	2	2.7	HIRATA	89C KAMI
>58	n	90	0	1.9	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

>29	p	90	0	2.2	BERGER	91 FREJ
>41	n	90	0	1.4	BERGER	91 FREJ
>38	n	90	2	4.1	SEIDEL	88 IMB
> 1.2	p	90	0		BARTELT	87 SOUD
> 1.5	n	90	0		BARTELT	87 SOUD
>17	p	90	7	7	HAINES	86 IMB
>14	n	90	9	4	HAINES	86 IMB
>12	p	90	0	<1.2	ARISAKA	85 KAMI
> 6	n	90	2	<1	ARISAKA	85 KAMI
> 6.7	p (free)	90	6	6	BLEWITT	85 IMB
>17	p	90	7	7	BLEWITT	85 IMB
>12	n	90	4	2	PARK	85 IMB
> 0.6	n	90	1	0.3	20 BARTELT	83 SOUD
> 0.5	p	90	1	0.3	20 BARTELT	83 SOUD
> 9.8	p	90	1		21 KRISHNA...	82 KOLR
> 0.8	p	90	2		22 CHERRY	81 HOME

20 Limit based on zero events.

21 We have calculated 90% CL limit from 0 confined events.

22 We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow \mu^+ \rho)$

τ_8

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>110	p	90	0	1.7	HIRATA	89C KAMI
> 23	n	90	1	1.8	HIRATA	89C KAMI

• • • We do not use the following data for averages, fits, limits, etc. • • •

> 12	p	90	0	0.5	BERGER	91 FREJ
> 22	n	90	0	1.1	BERGER	91 FREJ
> 4.3	p	90	0	0.7	PHILLIPS	89 HPW
> 30	p	90	0	0.5	SEIDEL	88 IMB
> 11	n	90	1	1.1	SEIDEL	88 IMB
> 16	p	90	4	4.5	HAINES	86 IMB
> 7	n	90	6	5	HAINES	86 IMB
> 12	p	90	0	<0.7	ARISAKA	85 KAMI
> 5	n	90	1	<1.2	ARISAKA	85 KAMI
> 5.5	p (free)	90	4	5	BLEWITT	85 IMB
> 16	p	90	4	5	BLEWITT	85 IMB
> 9	n	90	1	2	PARK	85 IMB

See key on page 1343

Baryon Full Listings

p

$\tau(N \rightarrow \nu\rho)$ τ_9

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>27	p	90	5	1.5	HIRATA	89C KAMI
>19	n	90	0	0.5	SEIDEL	88 IMB
••• We do not use the following data for averages, fits, limits, etc. •••						
> 9	n	90	4	2.4	BERGER	89 FREJ
>24	p	90	0	0.9	BERGER	89 FREJ
>13	n	90	4	3.6	HIRATA	89C KAMI
>13	p	90	1	1.1	SEIDEL	88 IMB
> 8	p	90	6	5	HAINES	86 IMB
> 2	n	90	15	10	HAINES	86 IMB
>11	p	90	2	1	KAJITA	86 KAMI
> 4	n	90	2	2	KAJITA	86 KAMI
> 4.1	p (free)	90	6	7	BLEWITT	85 IMB
> 8.4	p	90	6	5	BLEWITT	85 IMB
> 2	n	90	7	3	PARK	85 IMB
> 0.9	p	90	2		23 CHERRY	81 HOME
> 0.6	n	90	2		23 CHERRY	81 HOME

23 We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow e^+\omega)$ τ_{10}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>45	p	90	2	1.45	HIRATA	89C KAMI
••• We do not use the following data for averages, fits, limits, etc. •••						
>17	p	90	0	1.1	BERGER	91 FREJ
>26	p	90	1	1.0	SEIDEL	88 IMB
> 1.5	p	90	0		BARTELT	87 SOUD
>37	p	90	6	5.3	HAINES	86 IMB
>25	p	90	1	<1.4	ARISAKA	85 KAMI
>12	p (free)	90	6	7.5	BLEWITT	85 IMB
>37	p	90	6	5.7	BLEWITT	85 IMB
> 0.6	p	90	1	0.3	24 BARTELT	83 SOUD
> 9.8	p	90	1		25 KRISHNA...	82 KOLR
> 2.8	p	90	2		26 CHERRY	81 HOME

24 Limit based on zero events.

25 We have calculated 90% CL limit from 0 confined events.

26 We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow \mu^+\omega)$ τ_{11}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>57	p	90	2	1.9	HIRATA	89C KAMI
••• We do not use the following data for averages, fits, limits, etc. •••						
>11	p	90	0	1.0	BERGER	91 FREJ
> 4.4	p	90	0	0.7	PHILLIPS	89 HPW
>10	p	90	2	1.3	SEIDEL	88 IMB
>23	p	90	2	1	HAINES	86 IMB
> 6.5	p (free)	90	9	8.7	BLEWITT	85 IMB
>23	p	90	8	7	BLEWITT	85 IMB

$\tau(n \rightarrow \nu\omega)$ τ_{12}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>43	n	90	3	2.7	HIRATA	89C KAMI
••• We do not use the following data for averages, fits, limits, etc. •••						
>17	n	90	1	0.7	BERGER	89 FREJ
> 6	n	90	2	1.3	SEIDEL	88 IMB
>12	n	90	6	6	HAINES	86 IMB
>18	n	90	2	2	KAJITA	86 KAMI
>16	n	90	1	2	PARK	85 IMB
> 2.0	n	90	2		27 CHERRY	81 HOME

27 We have converted 2 possible events to 90% CL limit.

$\tau(N \rightarrow e^+K)$ τ_{13}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>150	p	90	0	<0.27	HIRATA	89C KAMI
> 1.3	n	90	0		ALEKSEEV	81 BAKS
••• We do not use the following data for averages, fits, limits, etc. •••						
> 60	p	90	0		BERGER	91 FREJ
> 70	p	90	0	1.8	SEIDEL	88 IMB
> 77	p	90	5	4.5	HAINES	86 IMB
> 38	p	90	0	<0.8	ARISAKA	85 KAMI
> 24	p (free)	90	7	8.5	BLEWITT	85 IMB
> 77	p	90	5	4	BLEWITT	85 IMB
> 1.3	p	90	0		ALEKSEEV	81 BAKS

$\tau(p \rightarrow e^+K_S^0)$ τ_{14}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>76	p	90	0	0.5	BERGER	91 FREJ

$\tau(p \rightarrow e^+K_L^0)$ τ_{15}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>44	p	90	0	≤ 0.1	BERGER	91 FREJ

$\tau(N \rightarrow \mu^+K)$ τ_{16}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>120	p	90	1	0.4	HIRATA	89C KAMI
> 1.1	n	90	0		BARTELT	87 SOUD
••• We do not use the following data for averages, fits, limits, etc. •••						
> 54	p	90	0		BERGER	91 FREJ
> 3.0	p	90	0	0.7	PHILLIPS	89 HPW
> 19	p	90	3	2.5	SEIDEL	88 IMB
> 1.5	p	90	0		28 BARTELT	87 SOUD
> 40	p	90	7	6	HAINES	86 IMB
> 19	p	90	1	<1.1	ARISAKA	85 KAMI
> 6.7	p (free)	90	11	13	BLEWITT	85 IMB
> 40	p	90	7	8	BLEWITT	85 IMB
> 6	p	90	1		BATTISTONI	84 NUSX
> 0.6	p	90	0		29 BARTELT	83 SOUD
> 0.4	n	90	0		29 BARTELT	83 SOUD
> 5.8	p	90	2		30 KRISHNA...	82 KOLR
> 2.0	p	90	0		CHERRY	81 HOME
> 0.2	n	90	0		31 GURR	67 CNTR

28 BARTELT 87 limit applies to $p \rightarrow \mu^+K_S^0$.

29 Limit based on zero events.

30 We have calculated 90% CL limit from 1 confined event.

31 We have converted half-life to 90% CL mean life.

$\tau(p \rightarrow \mu^+K_S^0)$ τ_{17}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>64	p	90	0	1.2	BERGER	91 FREJ

$\tau(p \rightarrow \mu^+K_L^0)$ τ_{18}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>44	p	90	0	≤ 0.1	BERGER	91 FREJ

$\tau(N \rightarrow \nu K)$ τ_{19}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>100	p	90	9	7.3	HIRATA	89C KAMI
> 86	n	90	0	2.4	HIRATA	89C KAMI
••• We do not use the following data for averages, fits, limits, etc. •••						
> 15	n	90	1	1.8	BERGER	89 FREJ
> 15	p	90	1	1.8	BERGER	89 FREJ
> 0.28	p	90	0	0.7	PHILLIPS	89 HPW
> 0.3	p	90	0		BARTELT	87 SOUD
> 0.75	n	90	0		32 BARTELT	87 SOUD
> 10	p	90	6	5	HAINES	86 IMB
> 15	n	90	3	5	HAINES	86 IMB
> 28	p	90	3	3	KAJITA	86 KAMI
> 32	n	90	0	1.4	KAJITA	86 KAMI
> 1.8	p (free)	90	6	11	BLEWITT	85 IMB
> 9.6	p	90	6	5	BLEWITT	85 IMB
> 10	n	90	2	2	PARK	85 IMB
> 5	n	90	0		BATTISTONI	84 NUSX
> 2	p	90	0		BATTISTONI	84 NUSX
> 0.3	n	90	0		33 BARTELT	83 SOUD
> 0.1	p	90	0		33 BARTELT	83 SOUD
> 5.8	p	90	1		34 KRISHNA...	82 KOLR
> 0.3	n	90	2		35 CHERRY	81 HOME

32 BARTELT 87 limit applies to $n \rightarrow \nu K_S^0$.

33 Limit based on zero events.

34 We have calculated 90% CL limit from 1 confined event.

35 We have converted 2 possible events to 90% CL limit.

$\tau(p \rightarrow e^+K^*(892)^0)$ τ_{20}

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>52	p	90	2	1.55	HIRATA	89C KAMI
••• We do not use the following data for averages, fits, limits, etc. •••						
>10	p	90	0	0.8	BERGER	91 FREJ
>10	p	90	1	<1	ARISAKA	85 KAMI

Baryon Full Listings

 p $\tau(N \rightarrow \nu K^*(892))$

721

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>22	n	90	0	2.1	BERGER	89 FREJ
>20	p	90	5	2.1	HIRATA	89C KAMI
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>17	p	90	0	2.4	BERGER	89 FREJ
>21	n	90	4	2.4	HIRATA	89C KAMI
>10	p	90	7	6	HAINES	86 IMB
> 5	n	90	8	7	HAINES	86 IMB
> 8	p	90	3	2	KAJITA	86 KAMI
> 6	n	90	2	1.6	KAJITA	86 KAMI
> 5.8	p (free)	90	10	16	BLEWITT	85 IMB
> 9.6	p	90	7	6	BLEWITT	85 IMB
> 7	n	90	1	4	PARK	85 IMB
> 2.1	p	90	1		³⁶ BATTISTONI	82 NUSX

³⁶We have converted 1 possible event to 90% CL limit. $\tau(p \rightarrow e^+ \pi^+ \pi^-)$

722

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>21	p	90	0	2.2	BERGER	91 FREJ

 $\tau(p \rightarrow e^+ \pi^0 \pi^0)$

723

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>38	p	90	1	0.5	BERGER	91 FREJ

 $\tau(n \rightarrow e^+ \pi^- \pi^0)$

724

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>32	n	90	1	0.8	BERGER	91 FREJ

 $\tau(p \rightarrow \mu^+ \pi^+ \pi^-)$

725

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>17	p	90	1	2.6	BERGER	91 FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 3.3	p	90	0	0.7	PHILLIPS	89 HPW

 $\tau(p \rightarrow \mu^+ \pi^0 \pi^0)$

726

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>33	p	90	1	0.9	BERGER	91 FREJ

 $\tau(n \rightarrow \mu^+ \pi^- \pi^0)$

727

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>33	n	90	0	1.1	BERGER	91 FREJ

 $\tau(n \rightarrow e^+ K^0 \pi^-)$

728

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>18	n	90	1	0.2	BERGER	91 FREJ

 $\tau(n \rightarrow e^- \pi^+)$

729

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>65	n	90	0	1.6	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>55	n	90	0	1.09	BERGER	91B FREJ
>16	n	90	9	7	HAINES	86 IMB
>25	n	90	2	4	PARK	85 IMB

 $\tau(n \rightarrow \mu^- \pi^+)$

730

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>49	n	90	0	0.5	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>33	n	90	0	1.40	BERGER	91B FREJ
> 2.7	n	90	0	0.7	PHILLIPS	89 HPW
>25	n	90	7	6	HAINES	86 IMB
>27	n	90	2	3	PARK	85 IMB

 $\tau(n \rightarrow e^- \rho^+)$

731

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>62	n	90	2	4.1	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>12	n	90	13	6	HAINES	86 IMB
>12	n	90	5	3	PARK	85 IMB

 $\tau(n \rightarrow \mu^- \rho^+)$

732

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>7	n	90	1	1.1	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>2.6	n	90	0	0.7	PHILLIPS	89 HPW
>9	n	90	7	5	HAINES	86 IMB
>9	n	90	2	2	PARK	85 IMB

 $\tau(n \rightarrow e^- K^+)$

733

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>32	n	90	3	2.96	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 0.23	n	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow \mu^- K^+)$

734

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>57	n	90	0	2.18	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 4.7	n	90	0	0.7	PHILLIPS	89 HPW

 $\tau(p \rightarrow e^- \pi^+ \pi^0)$

735

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>30	p	90	1	2.50	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 2.0	p	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow e^- \pi^+ \pi^0)$

736

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>29	n	90	1	0.78	BERGER	91B FREJ

 $\tau(p \rightarrow \mu^- \pi^+ \pi^+)$

737

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>17	p	90	1	1.72	BERGER	91B FREJ
• • • We do not use the following data for averages, fits, limits, etc. • • •						
> 7.8	p	90	0	0.7	PHILLIPS	89 HPW

 $\tau(n \rightarrow \mu^- \pi^+ \pi^0)$

738

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>34	n	90	0	0.78	BERGER	91B FREJ

 $\tau(p \rightarrow e^- \pi^+ K^+)$

739

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>20	p	90	3	2.50	BERGER	91B FREJ

 $\tau(p \rightarrow \mu^- \pi^+ K^+)$

740

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>5	p	90	2	0.78	BERGER	91B FREJ

 $\tau(p \rightarrow e^+ \gamma)$

741

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>460	p	90	0	0.6	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>133	p	90	0	0.3	BERGER	91 FREJ
>360	p	90	0	0.3	HAINES	86 IMB
> 87	p (free)	90	0	0.2	BLEWITT	85 IMB
>360	p	90	0	0.2	BLEWITT	85 IMB
> 0.1	p	90			³⁷ GURR	67 CNTR

³⁷We have converted half-life to 90% CL mean life. $\tau(p \rightarrow \mu^+ \gamma)$

742

LIMIT (10^{30} years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>380	p	90	0	0.5	SEIDEL	88 IMB
• • • We do not use the following data for averages, fits, limits, etc. • • •						
>155	p	90	0	0.1	BERGER	91 FREJ
> 97	p	90	3	2	HAINES	86 IMB
> 61	p (free)	90	0	0.2	BLEWITT	85 IMB
>280	p	90	0	0.6	BLEWITT	85 IMB
> 0.3	p	90			³⁸ GURR	67 CNTR

³⁸We have converted half-life to 90% CL mean life.

$\tau(n \rightarrow \nu\gamma)$ **743**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>24	n	90	10	6.86	BERGER	91B FREJ
••• We do not use the following data for averages, fits, limits, etc. •••						
> 9	n	90	73	60	HAINES	86 IMB
>11	n	90	28	19	PARK	85 IMB

$\tau(p \rightarrow e^+\gamma\gamma)$ **744**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>100	p	90	1	0.8	BERGER	91 FREJ

$\tau(p \rightarrow e^+e^+e^-)$ **745**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>510	p	90	0	0.3	HAINES	86 IMB
••• We do not use the following data for averages, fits, limits, etc. •••						
>147	p	90	0	0.1	BERGER	91 FREJ
> 89	p (free)	90	0	0.5	BLEWITT	85 IMB
>510	p	90	0	0.7	BLEWITT	85 IMB

$\tau(p \rightarrow e^+\mu^+\mu^-)$ **746**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>81	p	90	0	0.16	BERGER	91 FREJ
••• We do not use the following data for averages, fits, limits, etc. •••						
> 5.0	p	90	0	0.7	PHILLIPS	89 HPW

$\tau(p \rightarrow e^+\nu\nu)$ **747**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>11	p	90	11	6.08	BERGER	91B FREJ

$\tau(n \rightarrow e^+e^-\nu)$ **748**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>74	n	90	0	< 0.1	BERGER	91B FREJ
••• We do not use the following data for averages, fits, limits, etc. •••						
>45	n	90	5	5	HAINES	86 IMB
>26	n	90	4	3	PARK	85 IMB

$\tau(n \rightarrow \mu^+e^-\nu)$ **749**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>47	n	90	0	< 0.1	BERGER	91B FREJ

$\tau(n \rightarrow \mu^+\mu^-\nu)$ **750**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>42	n	90	0	1.4	BERGER	91B FREJ
••• We do not use the following data for averages, fits, limits, etc. •••						
> 5.1	n	90	0	0.7	PHILLIPS	89 HPW
>16	n	90	14	7	HAINES	86 IMB
>19	n	90	4	7	PARK	85 IMB

$\tau(p \rightarrow \mu^+e^+e^-)$ **751**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>91	p	90	0	≤ 0.1	BERGER	91 FREJ

$\tau(p \rightarrow \mu^+\mu^+\mu^-)$ **752**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>190	p	90	1	0.1	HAINES	86 IMB
••• We do not use the following data for averages, fits, limits, etc. •••						
>119	p	90	0	0.2	BERGER	91 FREJ
> 10.5	p	90	0	0.7	PHILLIPS	89 HPW
> 44	p (free)	90	1	0.7	BLEWITT	85 IMB
>190	p	90	1	0.9	BLEWITT	85 IMB
> 2.1	p	90	1		³⁹ BATTISTONI	82 NUSX

³⁹We have converted 1 possible event to 90% CL limit.

$\tau(p \rightarrow \mu^+\nu\nu)$ **753**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>21	p	90	7	11.23	BERGER	91B FREJ

$\tau(p \rightarrow e^-\mu^+\mu^+)$ **754**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>6.0	p	90	0	0.7	PHILLIPS	89 HPW

$\tau(n \rightarrow 3\nu)$ **755**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>0.00049	n	90	2	2	⁴⁰ SUZUKI	93B KAMI
••• We do not use the following data for averages, fits, limits, etc. •••						
>0.00003	n	90	11	6.1	⁴¹ BERGER	91B FREJ
>0.00012	n	90	7	11.2	⁴¹ BERGER	91B FREJ
>0.0005	n	90	0		LEARNED	79 RVUE

⁴⁰The SUZUKI 93B limit applies to any of $\nu_e\nu_e\nu_e$, $\nu_\mu\nu_\mu\nu_\mu$, or $\nu_\tau\nu_\tau\nu_\tau$.

⁴¹The first BERGER 91B limit is for $n \rightarrow \nu_e\nu_e\nu_e$, the second is for $n \rightarrow \nu_\mu\nu_\mu\nu_\mu$.

$\tau(N \rightarrow e^+\text{anything})$ **756**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>0.6	p, n	90			⁴² LEARNED	79 RVUE

⁴²The electron may be primary or secondary.

$\tau(N \rightarrow \mu^+\text{anything})$ **757**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>12	p, n	90	2		^{43,44} CHEERY	81 HOME
••• We do not use the following data for averages, fits, limits, etc. •••						
> 1.8	p, n	90			⁴⁴ COWSIK	80 CNTR
> 6	p, n	90			⁴⁴ LEARNED	79 RVUE

⁴³We have converted 2 possible events to 90% CL limit.

⁴⁴The muon may be primary or secondary.

$\tau(N \rightarrow \nu\text{anything})$ **758**

Anything = π, ρ, K , etc.

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>0.0002	p, n	90	0		LEARNED	79 RVUE

Anything = π, ρ, K , etc.

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>0.0002	p, n	90	0		LEARNED	79 RVUE

$\tau(N \rightarrow e^+\pi^0\text{anything})$ **759**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>0.6	p, n	90	0		LEARNED	79 RVUE

$\tau(N \rightarrow 2\text{ bodies}, \nu\text{-free})$ **760**

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>1.3	p, n	90	0		ALEKSEEV	81 BAKS

••• We do not use the following data for averages, fits, limits, etc. •••

LIMIT (10 ³⁰ years)	PARTICLE	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN
>0.7	p	90	4	2.34	BERGER	91B FREJ

$\tau(p\rho\rho \rightarrow \pi^+\pi^+)$ **761**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>0.7	90	4	2.34	BERGER	91B FREJ	τ per iron nucleus

$\tau(p\rho n \rightarrow \pi^+\pi^0)$ **762**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>2.0	90	0	0.31	BERGER	91B FREJ	τ per iron nucleus

$\tau(n\rho n \rightarrow \pi^+\pi^-)$ **763**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>0.7	90	4	2.18	BERGER	91B FREJ	τ per iron nucleus

$\tau(n\rho n \rightarrow \pi^0\pi^0)$ **764**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>3.4	90	0	0.78	BERGER	91B FREJ	τ per iron nucleus

$\tau(p\rho\rho \rightarrow e^+e^+)$ **765**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>5.8	90	0	<0.1	BERGER	91B FREJ	τ per iron nucleus

$\tau(p\rho\rho \rightarrow e^+\mu^+)$ **766**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>3.6	90	0	<0.1	BERGER	91B FREJ	τ per iron nucleus

$\tau(p\rho\rho \rightarrow \mu^+\mu^+)$ **767**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>1.7	90	0	0.62	BERGER	91B FREJ	τ per iron nucleus

$\tau(p\rho n \rightarrow e^+\nu)$ **768**

LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST	DOCUMENT ID	TECN	COMMENT
>2.8	90	5	9.67	BERGER	91B FREJ	τ per iron nucleus

Baryon Full Listings

p, n

$\tau(pn \rightarrow \mu^+ \bar{\nu})$		769	
LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST
>1.6	90	4	4.37
DOCUMENT ID	TECN	COMMENT	
BERGER	91B FREJ	τ per iron nucleus	

$\tau(nn \rightarrow \nu_e \bar{\nu}_e)$		770	
LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST
>0.000012	90	5	9.7
DOCUMENT ID	TECN	COMMENT	
BERGER	91B FREJ	τ per iron nucleus	

$\tau(nn \rightarrow \nu_\mu \bar{\nu}_\mu)$		771	
LIMIT (10 ³⁰ years)	CL%	EVTS	BKGD EST
>0.000006	90	4	4.4
DOCUMENT ID	TECN	COMMENT	
BERGER	91B FREJ	τ per iron nucleus	

\bar{p} PARTIAL MEAN LIVES

The "partial mean life" limits tabulated here are the limits on $\bar{\tau}/B_j$, where $\bar{\tau}$ is the total mean life for the antiproton and B_j is the branching fraction for the mode in question.

$\tau(\bar{p} \rightarrow e^- \gamma)$		772	
VALUE (years)	CL%	DOCUMENT ID	TECN
>1848	95	GEER	94 CALO
COMMENT	8.9 GeV/c \bar{p} beam		

$\tau(\bar{p} \rightarrow e^- \pi^0)$		773	
VALUE (years)	CL%	DOCUMENT ID	TECN
>554	95	GEER	94 CALO
COMMENT	8.9 GeV/c \bar{p} beam		

$\tau(\bar{p} \rightarrow e^- \eta)$		774	
VALUE (years)	CL%	DOCUMENT ID	TECN
>171	95	GEER	94 CALO
COMMENT	8.9 GeV/c \bar{p} beam		

$\tau(\bar{p} \rightarrow e^- K_S^0)$		775	
VALUE (years)	CL%	DOCUMENT ID	TECN
>29	95	GEER	94 CALO
COMMENT	8.9 GeV/c \bar{p} beam		

$\tau(\bar{p} \rightarrow e^- K_L^0)$		776	
VALUE (years)	CL%	DOCUMENT ID	TECN
>9	95	GEER	94 CALO
COMMENT	8.9 GeV/c \bar{p} beam		

p REFERENCES

GEER 94 PRL 72 1596	+Marriner, Ray+ (FNAL, UCLA, PSU)
HALLIN 93 PR C48 1497	+Amendt, Bergstrom+ (SASK, BOST, ILL)
SUZUKI 93B PL B311 357	+Fukuda, Hirata, Inoue+ (KAMIOKANDE Collab.)
HUGHES 92 PRL 69 578	+Deutch (LANL, AARM)
ZIEGER 92 PL B278 34	+Van de Vyver, Christmann, DeGraeve+ (MPCM)
92B PL B281 417 (erratum)	Ziegler, ..., Van den Abeele, Ziegler (MPCM)
BERGER 91 ZPHY C50 385	+Froehlich, Moench, Nisius+ (FREJUS Collab.)
BERGER 91B PL B269 227	+Froehlich, Moench, Nisius+ (FREJUS Collab.)
FEDERSPIEL 91 PRL 67 1511	+Eisenstein, Lucas, MacGibbon+ (ILL)
BECKER-SZ... 90 PR D42 2974	+Becker-Szendy, Bratton, Cady, Casper+ (IMB-3 Collab.)
ERICSON 90 EPL 11 295	+Richter (CERN, DARM)
GABRIELSE 90 PRL 65 1317	+Fei, Orozco, Tjoelker+ (HARV, MANZ, WASH, IBS)
BERGER 89 NP B313 509	+Froehlich, Moench+ (FREJUS Collab.)
CHO 89 PRL 63 2559	+Sangster, Hinds (YALE)
HIRATA 89C PL B220 308	+Kajita, Kifune, Kihara+ (Kamiokande Collab.)
PHILLIPS 89 PL B224 348	+Matthews, Aprile, Cline+ (HPW Collab.)
KREISSL 88 ZPHY C37 557	+Hancock, Koch, Koehler, Poth+ (CERN PS176 Collab.)
SEIDEL 88 PRL 61 2522	+Bionta, Blewitt, Bratton+ (IMB Collab.)
BARTELT 87 PR D36 1990	+Courant, Heller+ (Soudan Collab.)
Also 89 PR D40 1701 erratum	+Bartels, Courant, Heller+ (Soudan Collab.)
COHEN 87 RMP 59 1121	+Taylor (RISC, NBS)
HAINES 86 PRL 57 1986	+Bionta, Blewitt, Bratton, Casper+ (IMB Collab.)
KAJITA 86 JPSJ 55 711	+Arisaka, Koshiba, Nakahata+ (Kamiokande Collab.)
ARISAKA 85 JPSJ 54 3213	+Kajita, Koshiba, Nakahata+ (Kamiokande Collab.)
BLEWITT 85 PRL 55 2114	+LoSecco, Bionta, Bratton+ (IMB Collab.)
DZUBA 85 PL 154B 93	+Flambaum, Silvestrov (NOVO)
PARK 85 PRL 54 29	+Blewitt, Cortez, Foster+ (IMB Collab.)
BATTISTONI 84 PL 133B 454	+Bellotti, Bologna, Campana+ (NUSEX Collab.)
MARINELLI 84 PL 137B 439	+Morpurgo (GENO)
WILKENING 84 PR A29 425	+Ramsey, Larson (HARV, VIRG)
BARTELT 83 PRL 50 651	+Courant, Heller, Joyce, Marshak+ (MINN, ANL)
BATTISTONI 82 PL 118B 461	+Bellotti, Bologna, Campana+ (NUSEX Collab.)
KRISHNA... 82 PL 115B 349	+Krishnaswamy, Menon+ (TATA, OSKC, INUS)
ALEKSEEV 81 JETPL 33 651	+Bakatanov, Butkevich, Voevodskii+ (PNPI)
Translated from ZETFP 33 664.	
CHERRY 81 PRL 47 1507	+Deakynne, Lande, Lee, Steinberg+ (PENN, BNL)
COWSIK 80 PR D22 2204	+Narasimhan (TATA)
BELL 79 PL 86B 215	+Calvetti, Carron, Chaney, Cittolin+ (CERN)
GOLDEN 79 PRL 43 1196	+Horan, Mauger, Badwar, Lacy+ (NASA, PSSL)
LEARNED 79 PRL 43 907	+Reines, Soni (UC)
BREGMAN 78 PL 78B 174	+Calvetti, Carron, Cittolin, Hauer, Herr+ (CERN)
ROBERTS 78 PR D17 358	+Will, RHEL (MIT)
EVANS 77 Science 197 989	+Steinberg (BNL, PENN)
ROBERSON 77 PR C16 1945	+King, Kunselman+ (WYOM, CIT, CMU, VPI, WILL)
HU 75 NP A254 403	+Asano, Chen, Cheng, Dugan+ (COLU, YALE)
COHEN 73 JPCRD 2 663	+Taylor (RISC, NBS)
DYLLA 73 PR A7 1224	+King (MIT)
BAMBERGER 70 PL 35B 233	+Lynen, Piekartz+ (MPIH, CERN, KARL)
DIX Case Thesis	(CASE)
HARRISON 69 PRL 22 1263	+Sandars, Wright (OXF)
GURR 67 PR 158 1321	+Kropp, Reines, Meyer (CASE, WITW)
FLEROV 58 DOKL 3 79	+Klochikov, Skobkin, Terentev (ASCI)



$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$$
 Status: * * * *

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1986 edition (Physics Letters 170B) or in earlier editions.

n MASS

The mass is known much more precisely in u (atomic mass units) than in MeV; see the footnotes. The conversion from u to MeV, $1 u = 931.49432 \pm 0.00028$ MeV, involves the relatively poorly known electronic charge. The NATARAJAN 93 value, in u, is by far the best, but when converted to MeV differs only negligibly from the 1986 CODATA value, which, for consistency, we stick with.

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
939.56563 ± 0.00028	¹ COHEN	87 RVUE	1986 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
939.56565 ± 0.00028	² NATARAJAN	93	Penning trap
939.56564 ± 0.00028	^{3,4} GREENE	86 SPEC	$n p \rightarrow d \gamma$
939.5731 ± 0.0027	⁴ COHEN	73 RVUE	1973 CODATA value
¹ The mass is known much more precisely in u: $m = 1.008664904 \pm 0.000000014$ u.			
² The mass is known much more precisely in u: $m = 1.0086649234 \pm 0.0000000023$ u.			
We use the conversion factor given above to get the mass in MeV.			
³ The mass is known much more precisely in u: $m = 1.008664919 \pm 0.000000014$ u.			
⁴ These determinations are not independent of the $m_n - m_p$ measurements below.			

\bar{n} MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
939.485 ± 0.051	59	⁵ CRESTI	86 HBC	$\bar{p} p \rightarrow \bar{n} n$
⁵ This is a corrected result (see the erratum). The error is statistical. The maximum systematic error is 0.029 MeV.				

$m_n - m_p$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1.293318 ± 0.000009	⁶ COHEN	87 RVUE	1986 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.2933328 ± 0.0000072	GREENE	86 SPEC	$n p \rightarrow d \gamma$
1.293429 ± 0.000036	COHEN	73 RVUE	1973 CODATA value
⁶ Calculated by us from the COHEN 87 ratio $m_n/m_p = 1.001378404 \pm 0.000000009$. In u, $m_n - m_p = 0.001388434 \pm 0.000000009$ u.			

$(m_n - m_{\bar{n}}) / m_{\text{average}}$

A test of CPT invariance. Calculated from the n and \bar{n} masses, above.

VALUE	DOCUMENT ID
$(9 \pm 5) \times 10^{-5}$	OUR EVALUATION


n MEAN LIFE

We now compile only direct measurements of the lifetime, not those inferred from decay correlation measurements. (Limits on lifetimes for bound neutrons are given in the section " p PARTIAL MEAN LIVES.")

For a review, see EROZOLIMSKII 89 and papers that follow it in an issue of NIM devoted to the "Proceedings of the International Workshop on Fundamental Physics with Slow Neutrons" (Grenoble 1989). For later reviews and/or commentary, see FREEDMAN 90, SCHRECKENBACH 92, and PENDELBURY 93.

VALUE (s)	DOCUMENT ID	TECN	COMMENT
887.0 ± 2.0 OUR AVERAGE			Error includes scale factor of 1.3. See the ideogram below.
882.6 ± 2.7	MAMPE	93 CNTR	Gravitational trap
888.4 ± 3.1 ± 1.1	NESVIZHEV...	92 CNTR	Gravitational trap
893.6 ± 3.8 ± 3.7	BYRNE	90 CNTR	Penning trap
878 ± 27 ± 14	KOSSAKOW...	89 TPC	Pulsed beam
887.6 ± 3.0	MAMPE	89 CNTR	Gravitational trap
877 ± 10	PAUL	89 CNTR	Storage ring
876 ± 10 ± 19	LAST	88 SPEC	Pulsed beam
891 ± 9	SPIVAK	88 CNTR	Beam
903 ± 13	KOSVINTSEV	86 CNTR	Gravitational trap
918 ± 14	CHRISTENSEN72	CNTR	
• • • We do not use the following data for averages, fits, limits, etc. • • •			
888.4 ± 2.9	ALFIMENKOV	90 CNTR	See NESVIZHEVSKII 92
937 ± 18	BYRNE	80 CNTR	
875 ± 95	KOSVINTSEV	80 CNTR	
881 ± 8	BONDAREN...	78 CNTR	See SPIVAK 88
⁷ This measurement has been withdrawn (J. Byrne, private communication, 1990).			

WEIGHTED AVERAGE
887.0±2.0 (Error scaled by 1.3)



neutron mean life (s)

***n* MAGNETIC MOMENT**

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
-1.91304275 ± 0.00000045	COHEN	87 RVUE	1986 CODATA value
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-1.91304277 ± 0.00000048	⁸ GREENE	82 MRS	
⁸ GREENE 82 measures the moment to be $(1.04187564 \pm 0.00000026) \times 10^{-3}$ Bohr magnetons. The value above is obtained by multiplying this by $m_p/m_e = 1836.152701 \pm 0.000037$ (the 1986 CODATA value from COHEN 87).			

***n* ELECTRIC DIPOLE MOMENT**

A nonzero value is forbidden by both *T* invariance and *P* invariance. A number of early results have been omitted. See RAMSEY 90 for a review.

VALUE (10^{-26} e cm)	CL%	DOCUMENT ID	TECN	COMMENT
< 11	95	ALTAREV 92 MRS		$(+2.6 \pm 4.2 \pm 1.6) \times 10^{-26}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 12	95	SMITH 90 MRS		$d = (-3 \pm 5) \times 10^{-26}$
< 26	95	ALTAREV 86 MRS		$d = (-14 \pm 6) \times 10^{-26}$
3 ± 48		PENDLEBURY 84 MRS		Ultracold neutrons
< 60	90	ALTAREV 81 MRS		$d = (21 \pm 24) \times 10^{-26}$
< 160	90	ALTAREV 79 MRS		$d = (40 \pm 75) \times 10^{-26}$

***n* ELECTRIC POLARIZABILITY α_n**

Following is the electric polarizability α_n defined in terms of the induced electric dipole moment by $\mathbf{D} = 4\pi\epsilon_0\alpha_n\mathbf{E}$. For a review, see SCHMIED-MAYER 89.

VALUE (10^{-3} fm ³)	DOCUMENT ID	TECN	COMMENT
1.16^{+0.19}_{-0.23} OUR AVERAGE			
1.20 ± 0.15 ± 0.20	SCHMIEDM...	91 CNTR	<i>n</i> Pb transmission
1.07 ^{+0.33} _{-1.07}	ROSE	90B CNTR	$\gamma d \rightarrow \gamma n p$
0.8 ± 1.0	KOESTER	88 CNTR	<i>n</i> Pb, <i>n</i> Bi transmission
1.2 ± 1.0	SCHMIEDM...	88 CNTR	<i>n</i> Pb, <i>n</i> C transmission
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.17 ^{+0.43} _{-1.17}	ROSE	90 CNTR	See ROSE 90B

***n* CHARGE**

See also " $|q_p + q_e|$ CHARGE MAGNITUDE DIFFERENCE" in the proton Listings.

VALUE (10^{-21} e)	DOCUMENT ID	TECN	COMMENT
- 0.4 ± 1.1	⁹ BAUMANN	88	Cold <i>n</i> deflection
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-15 ± 22	¹⁰ GAehler	82 CNTR	Reactor neutrons
⁹ The BAUMANN 88 error ± 1.1 gives the 68% CL limits about the the value -0.4. ¹⁰ The GAehler 82 error ± 22 gives the 90% CL limits about the the value -15.			

LIMIT ON *n* \bar{n} OSCILLATIONS

Mean Time for *n* \bar{n} Transition in Vacuum

A test of $\Delta B=2$ baryon number nonconservation. MOHAPATRA 80 and MOHAPATRA 89 discuss the theoretical motivations for looking for *n* \bar{n} oscillations. DOVER 83 and DOVER 85 give phenomenological analyses. The best limits come from looking for the decay of neutrons bound in nuclei. However, these analyses require model-dependent corrections for nuclear effects. See KABIR 83, DOVER 89, and ALBERICO 91 for discussions. Direct searches for *n* → \bar{n} transitions using reactor neutrons are cleaner but give poorer limits.

VALUE (s)	CL%	DOCUMENT ID	TECN	COMMENT
> 1.2 × 10⁸	90	BERGER	90 FREJ	<i>n</i> bound in iron
> 1.2 × 10⁸	90	TAKITA	86 CNTR	Kamiokande
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 1 × 10 ⁷	90	BALDO....	90 CNTR	Reactor neutrons
> 4.9 × 10 ⁵	90	BRESSI	90 CNTR	Reactor neutrons
> 4.7 × 10 ⁵	90	BRESSI	89 CNTR	See BRESSI 90
> 1 × 10 ⁶	90	FIDECARO	85 CNTR	Reactor neutrons
> 8.8 × 10 ⁷	90	PARK	85B CNTR	
> 3 × 10 ⁷		BATTISTONI	84 NUSX	
> 2.7 × 10 ⁷ - 1.1 × 10 ⁸		JONES	84 CNTR	
> 2 × 10 ⁷		CHERRY	83 CNTR	

***n* DECAY MODES**

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $p e^- \bar{\nu}_e$	100 %	
Γ_2 hydrogen-atom $\bar{\nu}_e$		
Charge conservation (<i>Q</i>) violating mode		
Γ_3 $p \nu_e \bar{\nu}_e$	$Q < 9 \times 10^{-24}$	90%

***n* BRANCHING RATIOS**

$\Gamma(\text{hydrogen-atom } \bar{\nu}_e)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	Γ_2/Γ
VALUE (units 10^{-2})				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
< 3	95	¹¹ GREEN	90 RVUE	
¹¹ GREEN 90 infers that $\tau(\text{hydrogen-atom } \bar{\nu}_e) > 3 \times 10^4$ s by comparing neutron lifetime measurements made in storage experiments with those made in β -decay experiments. However, the result depends sensitively on the lifetime measurements, and does not of course take into account more recent measurements of same.				

$\Gamma(p \nu_e \bar{\nu}_e)/\Gamma_{\text{total}}$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
Forbidden by charge conservation.					
VALUE					
< 9 × 10⁻²⁴	90	BARABANOV	80 CNTR	⁷¹ Ga → ⁷¹ GeX	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 9.7 × 10 ⁻¹⁸	90	ROY	83 CNTR	¹¹³ Cd → ^{113m} In neut.	
< 7.9 × 10 ⁻²¹		VAIDYA	83 CNTR	⁸⁷ Rb → ^{87m} Sr neut.	
< 3 × 10 ⁻¹⁹		NORMAN	79 CNTR	⁸⁷ Rb → ^{87m} Sr neut.	

NOTE ON BARYON DECAY PARAMETERS

(by E.D. Commins, University of California, Berkeley)

Baryon semileptonic decays

The typical baryon semileptonic decay is described by a matrix element, the hadronic part of which may be written as:

$$\bar{B}_f [f_1(q^2)\gamma_\lambda + i f_2(q^2)\sigma_{\lambda\mu}q^\mu + g_1(q^2)\gamma_\lambda\gamma_5 + g_3(q^2)\gamma_5q_\lambda] B_i .$$

Here B_i and \bar{B}_f are spinors describing the initial and final baryons and $q = p_i - p_f$, while the terms in f_1 , f_2 , g_1 , and g_3 account for vector, induced tensor ("weak magnetism"), axial vector, and induced pseudoscalar contributions [1]. Second-class current contributions are ignored here. In the limit of zero momentum transfer, f_1 reduces to the vector coupling constant g_V , and g_1 reduces to the axial-vector coupling constant g_A . The latter coefficients are related by Cabibbo's theory [2], generalized to six quarks (and three mixing angles) by Kobayashi and Maskawa [3]. The g_3 term is negligible for transitions in which an e^\pm is emitted, and gives a very small correction, which can be estimated by PCAC [4], for μ^\pm modes. Recoil effects include

Baryon Full Listings

n

weak magnetism, and are taken into account adequately by considering terms of first order in

$$\delta = (m_i - m_f)/(m_i + m_f), \quad \text{where}$$

m_i and m_f are the masses of the initial and final baryons.

The experimental quantities of interest are the total decay rate, the lepton-neutrino angular correlation, the asymmetry coefficients in the decay of a polarized initial baryon, and the polarization of the decay baryon in its own rest frame for an unpolarized initial baryon. Formulae for these quantities are derived by standard means [5] and are analogous to similar formulae for beta decay [6]. For comparison with high-precision experiments, it is necessary to modify the form factors at $q^2 = 0$ by a “dipole” q^2 dependence, and also to apply appropriate radiative corrections [7].

The ratio g_A/g_V may be written as

$$g_A/g_V = |g_A/g_V| e^{i\phi_{AV}}.$$

The presence of a “triple correlation” term in the transition probability, proportional to $\text{Im}(g_A/g_V)$ and of the form

$$\sigma_i \cdot (\mathbf{p}_\ell \times \mathbf{p}_\nu)$$

for initial baryon polarization or

$$\sigma_f \cdot (\mathbf{p}_\ell \times \mathbf{p}_\nu)$$

for final baryon polarization, would indicate failure of time-reversal invariance. The phase angle ϕ has been measured precisely only in neutron decay (and in ^{19}Ne nuclear beta decay), and the results are consistent with T invariance.

Hyperon nonleptonic decays

The most general decay amplitude for $J^P = 1/2^+$ hyperons may be written in the form

$$M = G_F m_\pi^2 \cdot \bar{B}_f (A - B\gamma_5) B_i,$$

where A and B are constants [1]. Then the transition rate is proportional to

$$R = 1 + \gamma \hat{\omega}_f \cdot \hat{\omega}_i + (1 - \gamma)(\hat{\omega}_f \cdot \hat{\mathbf{n}})(\hat{\omega}_i \cdot \hat{\mathbf{n}}) \\ + \alpha(\hat{\omega}_f \cdot \hat{\mathbf{n}} + \hat{\omega}_i \cdot \hat{\mathbf{n}}) + \beta \hat{\mathbf{n}} \cdot (\hat{\omega}_f \times \hat{\omega}_i),$$

where $\hat{\mathbf{n}}$ is a unit vector in the direction of the final baryon momentum, and $\hat{\omega}_i$ and $\hat{\omega}_f$ are unit vectors in the directions of the initial and final baryon spins. (The sign of the last term in the above equation was incorrect in our 1988 and 1990 editions.) Also,

$$\alpha = 2 \text{Re}(s^*p)/(|s|^2 + |p|^2),$$

$$\beta = 2 \text{Im}(s^*p)/(|s|^2 + |p|^2),$$

and

$$\gamma = (|s|^2 - |p|^2)/(|s|^2 + |p|^2),$$

where $s = A$ and $p = |\mathbf{p}_f|B/(E_f + m_f)$; here E_f and \mathbf{p}_f are the energy and momentum of the final baryon. The parameters α , β , and γ satisfy

$$\alpha^2 + \beta^2 + \gamma^2 = 1.$$

If the hyperon polarization is \mathbf{P}_Y , the polarization \mathbf{P}_B of the decay baryons is

$$\mathbf{P}_B = \frac{(\alpha + \mathbf{P}_Y \cdot \hat{\mathbf{n}})\hat{\mathbf{n}} + \beta(\mathbf{P}_Y \times \hat{\mathbf{n}}) + \gamma\hat{\mathbf{n}} \times (\mathbf{P}_Y \times \hat{\mathbf{n}})}{1 + \alpha\mathbf{P}_Y \cdot \hat{\mathbf{n}}}.$$

Here \mathbf{P}_B is defined in the rest system of the baryon, obtained by a Lorentz transformation along $\hat{\mathbf{n}}$ from the hyperon rest frame, in which $\hat{\mathbf{n}}$ and \mathbf{P}_Y are defined.

An additional useful parameter ϕ is defined by

$$\beta = (1 - \alpha^2)^{1/2} \sin\phi.$$

In the Listings, we compile α and ϕ for each decay, since these quantities are most closely related to experiment and are essentially uncorrelated. When necessary, we have changed the signs of reported values to agree with our sign conventions. In the Baryon Summary Table, we give α , ϕ , and Δ (defined below) with errors, and also give the value of γ without error.

Time-reversal invariance requires, in the absence of final-state interactions, that s and p be relatively real, and therefore that $\beta = 0$. However, for the decays discussed here, the final-state interaction is strong. Thus

$$s = |s|e^{i\delta_s} \text{ and } p = |p|e^{i\delta_p}.$$

where δ_s and δ_p are the pion-baryon s - and p -wave strong interaction phase shifts. We then have

$$\beta = \frac{-2|s||p|}{|s|^2 + |p|^2} \sin(\delta_s - \delta_p).$$

One also defines $\Delta = -\tan^{-1}(\beta/\alpha)$. If T invariance holds, $\Delta = \delta_s - \delta_p$. For $\Lambda \rightarrow p\pi^-$ decay, the value of Δ may be compared with the s - and p -wave phase shifts in low-energy π^-p scattering, and the results are consistent with T invariance.

References

1. E.D. Commins and P.H. Bucksbaum, *Weak Interactions of Leptons and Quarks* (Cambridge University Press, Cambridge, England, 1983).
2. N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963).
3. M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
4. M.L. Goldberger and S.B. Treiman, Phys. Rev. **111**, 354 (1958).
5. P.H. Frampton and W.K. Tung, Phys. Rev. **D3**, 1114 (1971).
6. J.D. Jackson, S.B. Treiman, and H.W. Wyld, Jr., Phys. Rev. **106**, 517 (1957), and Nucl. Phys. **4**, 206 (1957).
7. Y. Yokoo, S. Suzuki, and M. Morita, Prog. Theor. Phys. **50**, 1894 (1973).

$n \rightarrow pe^- \nu$ DECAY PARAMETERS

See the above Note on Baryon Decay Parameters. For discussions of recent results, see the references cited at the beginning of the section on the neutron mean life. For discussions of the values of the weak coupling constants g_A and g_V obtained using the neutron and asymmetry parameter A , comparisons with other methods of obtaining these constants, and implications for particle and astrophysics, see DUBBERS 91 and WOOLCOCK 91. For tests of the $V-A$ theory of neutron decay, see EROZOLIMSKII 91b.

 g_A/g_V

VALUE	DOCUMENT ID	TECN	COMMENT
-1.2573 ± 0.0028 OUR AVERAGE			
-1.2544 ± 0.0036	EROZOLIM... 91	CNTR	<i>e</i> mom- <i>n</i> spin corr.
-1.262 ± 0.005	BOPP 86	SPEC	<i>e</i> mom- <i>n</i> spin corr.
-1.261 ± 0.012	12 EROZOLIM... 79	CNTR	<i>e</i> mom- <i>n</i> spin corr.
-1.259 ± 0.017	12 STRATOWA 78	CNTR	proton recoil spectrum
-1.258 ± 0.015	13 KROHN 75	CNTR	<i>e</i> mom- <i>n</i> spin corr.

• • • We do not use the following data for averages, fits, limits, etc. • • •

-1.226 ± 0.042	MOSTOVOY 83	RVUE	
-1.263 ± 0.015	EROZOLIM... 77	CNTR	See EROZOLIMSKII 79
-1.250 ± 0.036	12 DOBROZE... 75	CNTR	See STRATOWA 78
-1.263 ± 0.016	14 KROPF 74	RVUE	<i>n</i> decay alone
-1.250 ± 0.009	14 KROPF 74	RVUE	<i>n</i> decay + nuclear ft

12 These experiments measure the absolute value of g_A/g_V only.

13 KROHN 75 includes events of CHRISTENSEN 70.

14 KROPF 74 reviews all data through 1972.

 β ASYMMETRY PARAMETER A

This is the neutron-spin electron-momentum correlation coefficient. Unless otherwise noted, the values are corrected for radiative effects and weak magnetism.

VALUE	DOCUMENT ID	TECN
-0.1127 ± 0.0011 OUR AVERAGE		
-0.1116 ± 0.0014	EROZOLIM... 91	CNTR
-0.1146 ± 0.0019	BOPP 86	SPEC
-0.114 ± 0.005	15 EROZOLIM... 79	CNTR
-0.113 ± 0.006	15 KROHN 75	CNTR

15 These results are not corrected for radiative effects and weak magnetism, but the corrections are small compared to the errors.

 $\bar{\nu}$ ASYMMETRY PARAMETER B

This is the neutron-spin antineutrino-momentum correlation coefficient.

VALUE	DOCUMENT ID	TECN
0.997 ± 0.028 OUR AVERAGE		
0.995 ± 0.034	CHRISTENSEN70	CNTR
1.00 ± 0.05	EROZOLIM... 70C	CNTR

 $\epsilon - \bar{\nu}$ ANGULAR CORRELATION COEFFICIENT a

VALUE	DOCUMENT ID	TECN	COMMENT
-0.102 ± 0.005 OUR AVERAGE			
-0.1017 ± 0.0051	STRATOWA 78	CNTR	Proton recoil spectrum
-0.091 ± 0.039	GRIGOREV 68	SPEC	Proton recoil spectrum

 ϕ_{AV} , PHASE OF g_A RELATIVE TO g_V

Time reversal invariance requires this to be 0 or 180°.

VALUE (°)	DOCUMENT ID	TECN	COMMENT
180.07 ± 0.18 OUR EVALUATION			
Using the average value for quantity D given in the next data block and $\lambda \equiv g_A/g_V$ in $\sin^2 \phi_{AV} = D(1+3\lambda^2)/2\lambda$.			

180.09 ± 0.18 OUR AVERAGE

179.71 ± 0.39	EROZOLIM... 78	CNTR	Polarized neutrons
180.35 ± 0.43	EROZOLIM... 74	CNTR	Polarized neutrons
180.14 ± 0.22	STEINBERG 74	CNTR	Polarized neutrons

• • • We do not use the following data for averages, fits, limits, etc. • • •

181.1 ± 1.3	16 KROPF 74	RVUE	<i>n</i> decay
-------------	-------------	------	----------------

16 KROPF 74 reviews all data through 1972.

TRIPLE CORRELATION COEFFICIENT D

These are measurements of the component of *n* spin perpendicular to the decay plane in β decay. Should be zero if *T* invariance is not violated.

VALUE	DOCUMENT ID	TECN	COMMENT
(-0.5 ± 1.4) × 10⁻³ OUR AVERAGE			
+ 0.0022 ± 0.0030	EROZOLIM... 78	CNTR	Polarized neutrons
- 0.0027 ± 0.0050	17 EROZOLIM... 74	CNTR	Polarized neutrons
- 0.0011 ± 0.0017	STEINBERG 74	CNTR	Polarized neutrons

17 EROZOLIMSKII 78 says asymmetric proton losses and nonuniform beam polarization may give a systematic error up to 0.003, thus increasing the EROZOLIMSKII 74 error to 0.005. STEINBERG 74 and STEINBERG 76 estimate these systematic errors to be insignificant in their experiment.

n REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1986 edition (Physics Letters **170B**) or in earlier editions.

MAMPE 93	JETPL 57 82	+Bondarenko, Morozov+	(KIAE)
	Translated from ZETFP 57 77.		
NATARAJAN 93	PRL 71 1998	+Boyce, DiFilippo, Pritchard	(MIT)
PENDLEBURY 93	ARNPS 43 687		(ILLG)
ALTAREV 92	PL B276 242	+Borisov, Borovikova, Ivanov+	(PNPI)
NESVIZHEV... 92	JETP 75 405	Nesvizhevskii, Serebrov, Tal'daev+	(PNPI, JINR)
	Translated from ZETF 102 740.		
SCHRECK... 92	JPG 18 1	Schreckbach, Mampe	(ILLG)
ALBERICO 91	NP A523 488	+de Pace, Pignone	(TORI)
DUBBERS 91	NP A527 239c		(ILLG)
	EPL 11 195	Dubbers, Mampe, Doehner	(ILLG, HEID)
EROZOLIM... 90	PL B263 33	Erozolimskii, Kuznetsov, Stepanenko, Kuida+	(PNPI, KIAE)
	Also	Erozolimskii, Kuznetsov, Stepanenko, Kuida+	(PNPI, KIAE)
Also	SJNP 52 999	Erozolimskii, Kuznetsov, Stepanenko, Kuida+	(PNPI, KIAE)
	Translated from YAF 52 1583.		
EROZOLIM... 91B	SJNP 53 260	Erozolimskii, Mostovoy	(KIAE)
	Translated from YAF 53 418.		
SCHMIEDM... 91	MPL A6 2579	Schmiedmayer, Riehs, Harvey, Hill	(TUW, ORNL)
WOOLCOCK 91	MPL A6 2579		(CANB)
ALFIMENKOV 90	JETPL 52 373	+Varlamov, Vasil'ev, Gudkov+	(PNPI, JINR)
	Translated from ZETFP 52 984.		
BALDO... 90	PL B236 95	Baldo-Ceolin, Benetti, Bitter+	(PADO, PAVI, HEIDP, ILLG)
BERGER 90	PL B240 237	+Froehlich, Moench, Nisius+	(FREJUS Collab.)
BRESSI 90	NC 103A 731	+Caillargh, Cambiaghi+	(PAVI, ROMA, MILA)
BYRNE 90	PRL 65 289	+Dawber, Spain, Williams+	(SUSS, NBS, SCOT, CNM)
FREEDMAN 90	CNPP 19 209		(ANL)
GREEN 90	JPG 16 L75	+Thompson	(RAL)
RAMSEY 90	ARNPS 40 1		(HARV)
ROSE 90	PL B234 460	+Zurmuehl, Rullhusen, Ludwig+	(GOET, MPCM, MANZ)
ROSE 90B	NP A514 621	+Zurmuehl, Rullhusen, Ludwig+	(GOET, MPCM)
SMITH 90	PL B234 191	+Crampin+	(SUSS, RAL, HARV, WASH, ILLG, MUNT)
BRESSI 89	ZPH C43 175	+Caillargh, Cambiaghi+	(INFN, MILA, PAVI, ROMA)
DOVER 89	NIM A284 13	+Gal, Richard	(BNL, HEBR, ISNG)
EROZOLIM... 89	NIM A284 89	Erozolimskii	(PNPI)
KOSSAKOW... 89	NP A503 473	Kossakowski, Grivot+	(LAPP, SAVO, ISNG, ILLG)
MAMPE 89	PRL 63 593	+Ageron, Bates, Pendlebury, Steyerl	(ILLG, RISL, SUSS, URI)
MOHAPATRA 89	NIM A284 14		(UMD)
PAUL 89	ZPHY C45 25	+Anton, Paul, Paul, Mampe	(BONN, WUPP, MPH, ILLG)
SCHMIEDM... 89	NIM A284 137	Schmiedmayer, Rauch, Riehs	(WIEN)
BAUMANN 88	PR D37 3107	+Gaehler, Kalus, Mampe	(BAYR, MUNI, ILLG)
KOESTER 88	ZPHY A329 229	+Waschkowski, Meier	(MUNI, MUNT)
LAST 88	PRL 60 995	+Arnold, Doehner, Dubbers+	(HEIDP, ILLG, ANL)
SCHMIEDM... 88	PRL 61 1065	Schmiedmayer, Rauch, Riehs	(TUW)
Also	PRL 61 2509 erratum		(TUW)
SPIVAK 88	JETP 67 1735	Schmiedmayer, Rauch, Riehs	(KIAE)
	Translated from ZETF 94 1.		
COHEN 87	RMP 59 1121	+Taylor	(RISC, NBS)
ALTAREV 86	JETPL 44 460	+Borisov, Borovikova, Brandin, Egorov+	(PNPI)
	Translated from ZETFP 44 360.		
BOPP 86	PRL 56 919	+Dubbers, Hornig, Kient, Last+	(HEIDP, ANL, ILLG)
Also	ZPHY C37 179	Kient, Bopp, Hornig, Last+	(HEIDP, ANL, ILLG)
CRESTI 86	PL B177 206	+Pasquali, Peruzzo, Pinori, Sartori	(PADO)
Also	PL B200 587 erratum	Cresti, Pasquali, Peruzzo, Pinori, Sartori	(PADO)
GREENE 86	PRL 56 819	+Kessler, Deslattes, Boerner	(NBS, ILLG)
KOSVINTSEV 86	JETPL 44 571	+Morozov, Terekhov	(KIAE)
	Translated from ZETFP 44 444.		
TAKITA 86	PR D34 902	+Arisaka, Kajita, Kifune+	(KEK, TOKY+)
DOVER 85	PR C31 1423	+Gal, Richard	(BNL)
FIDECARO 85	PL 156B 122	+Lanceri+	(CERN, ILLG, PADO, RAL, SUSS)
PARK 85B	PL B200 587	+Blewitt, Cortez, Foster+	(IMB Collab.)
BATTISTONI 84	PL 133B 454	+Bellotti, Bologna, Campana+	(NUSEX Collab.)
JONES 84	PRL 52 720	+Bionta, Blewitt, Bratton+	(IMB Collab.)
PENDLEBURY 84	PL 136B 327	+Smith, Golub, Byrne+	(SUSS, HARV, RAL, ILLG)
CHERRY 83	PRL 50 1354	+Lande, Lee, Steinberg, Cleveland	(PENN, BNL)
DOVER 83	PR D27 1090	+Gal, Richards	(BNL)
KABIR 83	PRL 51 231		(HARV)
MOSTOVOY 83	JETPL 37 196		(KIAE)
	Translated from ZETFP 37 162.		
ROY 83	PR D28 1770	+Vaidya, Ephraim, Datar, Bhatki+	(TATA)
VAIDYA 83	PR D27 486	+Roy, Ephraim, Datar, Bhattacharjee	(TATA)
GAEHLER 82	PR D25 2887	+Kalus, Mampe	(BAYR, ILLG)
GREENE 82	Metrologia 18 93		(YALE, HARV, ILLG, SUSS, ORNL, CENG)
ALTAREV 81	PL 102B 13	+Borisov, Borovikova, Brandin, Egorov+	(PNPI)
BARABANOV 80	JETPL 32 359	+Vretkenin, Gavrin+	(PNPI)
	Translated from ZETFP 32 384.		
BYRNE 80	PL 92B 274	+Morse, Smith, Shaikh, Green, Greene	(SUSS, RL)
KOSVINTSEV 80	JETPL 31 236	+Kushnir, Morozov, Terekhov	(JINR)
	Translated from ZETFP 31 257.		
MOHAPATRA 80	PRL 44 1316	+Marshak	(CUNY, VPI)
ALTAREV 79	JETPL 29 730	+Borisov, Brandin, Egorov, Ezhov, Ivanov+	(PNPI)
	Translated from ZETFP 29 794.		
EROZOLIM... 79	SJNP 30 356	Erozolimskii, Frank, Mostovoy+	(KIAE)
	Translated from YAF 30 692.		
NORMAN 79	PRL 43 1226	+Seamster	(WASH)
BONDAREN... 78	JETPL 28 303	Bondarenko, Kurguzov, Prokofev+	(KIAE)
	Translated from ZETFP 28 328.		
Also	Smolence Conf.	Bondarenko	(KIAE)
EROZOLIM... 78	SJNP 28 48	Erozolimskii, Mostovoy, Fedunin, Frank+	(KIAE)
	Translated from YAF 28 98.		
STRATOWA 78	PR D18 3970	+Dobrozemsky, Weinzierl	(SEIB)
EROZOLIM... 77	JETPL 23 663	Erozolimskii, Frank, Mostovoy+	(KIAE)
	Translated from ZETFP 23 720.		
STEINBERG 76	PR D13 2469	+Liaud, Vignon, Hughes	(YALE, ISNG)
DOBROZE... 75	PR D11 510	Dobrozemsky, Kerschbaum, Moraw, Paul+	(SEIB)
KROHN 75	PL 55B 175	+Ringo	(ANL)
EROZOLIM... 74	JETPL 20 345	Erozolimskii, Mostovoy, Fedunin, Frank+	(KIAE)
	Translated from ZETFP 20 745.		
KROPF 74	ZPHY 267 129	+Paul	(LINZ)
Also	NP A154 160	Paul	(WIEN)
STEINBERG 74	PRL 33 41	+Liaud, Vignon, Hughes	(YALE, ISNG)
COHEN 73	JPCRD 2 663	+Taylor	(RISC, NBS)
CHRISTENSEN 72	PR D5 1628	+Nielsen, Bahnsen, Brown+	(RISO)
CHRISTENSEN 70	PR C1 1693	+Krohn, Ringo	(ANL)
EROZOLIM... 70C	PL 33B 351	Erozolimskii, Bondarenko, Mostovoy, Obinyakov+	(KIAE)
GRIGOREV 68	SJNP 6 239	Grigor'ev, Grishin, Vladimirov, Nikolaevskii+	(ITEP)
	Translated from YAF 6 329.		

Baryon Full Listings

N 's and Δ 's

NOTE ON N AND Δ RESONANCES

For a lengthier discussion of N and Δ resonances, see our previous edition, Phys. Rev. **D45**, 1 June 1992, Part II. For Argand plots of the partial-wave amplitudes, see our 1990 edition, Phys. Lett. **B239** (1990).

I. Introduction

(by G. Höhler, University of Karlsruhe and R.L. Workman, Virginia Polytechnic Institute and State University)

The excited states of the nucleon have been studied in a large number of formation and production experiments. The conventional Breit-Wigner masses, widths, and elasticities of the N and Δ resonances in the Baryon Summary Table come almost entirely from partial-wave analyses of πN total, elastic, and charge-exchange scattering data (Sec. II). Partial-wave analyses have also been made of much smaller sets of data to get some $N\eta$, AK , and ΣK branching fractions, and other branching fractions come from isobar-model analyses of $\pi N \rightarrow N\pi\pi$ data (Sec. III). Finally, some $N\gamma$ branching fractions have been determined from photoproduction experiments (Sec. IV).

Table 1 lists all the N and Δ entries in the Baryon Full Listings and gives our evaluation of the status of each, both overall and channel by channel. Only the "established" resonances (overall status 3 or 4 stars) appear in the Baryon Summary Table. We consider a resonance to be established only if it is seen in at least two independent analyses of elastic scattering and if its partial-wave amplitude neither behaves erratically nor has large errors.

The Baryon Full Listings give, in addition to the conventional Breit-Wigner parameters, the positions and residues of the nearest poles of the resonant partial waves on the second sheet of the complex energy plane. These come from $\pi N \rightarrow \pi N$ partial-wave analyses and from $\pi N \rightarrow N\pi\pi$ isobar-model analyses.

The interested reader will find further discussions in two extensive reviews [1,2], and in the Proceedings of the 5th International Symposium on Meson-Nucleon Physics and the Structure of the Nucleon [3], and more generally in issues of the πN Newsletter (see the note following Sec. II).

(References for this Section are at the end of Section II.)

Table 1. The status of the N and Δ resonances. Only those with an overall status of *** or **** are included in the main Baryon Summary Table.

Particle	$L_{2I,2J}$	Overall status	Status as seen in —							
			$N\pi$	$N\eta$	AK	ΣK	$\Delta\pi$	$N\rho$	$N\gamma$	
$N(939)$	P_{11}	****								
$N(1440)$	P_{11}	****	****	*				***	*	***
$N(1520)$	D_{13}	****	****	*				****	****	****
$N(1535)$	S_{11}	****	****	****				*	**	***
$N(1650)$	S_{11}	****	****	*	***	**		***	**	***
$N(1675)$	D_{15}	****	****	*	*			****	*	****
$N(1680)$	F_{15}	****	****					****	****	****
$N(1700)$	D_{13}	****	***	*	**	*		**	*	**
$N(1710)$	P_{11}	***	***	**	**	*		**	*	***
$N(1720)$	P_{13}	****	****	*	**	*		*	**	**
$N(1900)$	P_{13}	**	**						*	
$N(1990)$	F_{17}	**	**	*	*	*				*
$N(2000)$	F_{15}	**	**	*	*	*		*	**	
$N(2080)$	D_{13}	**	**	*	*					*
$N(2090)$	S_{11}	*	*							
$N(2100)$	P_{11}	*	*							
$N(2190)$	G_{17}	****	****	*	*	*			*	*
$N(2200)$	D_{15}	**	**	*	*					
$N(2220)$	H_{19}	****	****	*						
$N(2250)$	G_{19}	****	****	*						
$N(2600)$	I_{111}	***	***							
$N(2700)$	K_{113}	**	**							
$\Delta(1232)$	P_{33}	****	****	F						****
$\Delta(1600)$	P_{33}	***	***	o				***	*	**
$\Delta(1620)$	S_{31}	****	****	r				****	****	***
$\Delta(1700)$	D_{33}	****	****	b		*		***	**	***
$\Delta(1750)$	P_{31}	*	*	i						
$\Delta(1900)$	S_{31}	***	***	d		*		*	**	*
$\Delta(1905)$	F_{35}	****	****	d		*		**	**	***
$\Delta(1910)$	P_{31}	****	****	e		*		*	*	*
$\Delta(1920)$	P_{33}	***	***	n		*		**		*
$\Delta(1930)$	D_{35}	***	***			*				**
$\Delta(1940)$	D_{33}	*	*	F						
$\Delta(1950)$	F_{37}	****	****	o		*		****	*	****
$\Delta(2000)$	F_{35}	**	**	r					**	
$\Delta(2150)$	S_{31}	*	*	b						
$\Delta(2200)$	G_{37}	*	*	i						
$\Delta(2300)$	H_{39}	**	**	d						
$\Delta(2350)$	D_{35}	*	*	d						
$\Delta(2390)$	F_{37}	*	*	e						
$\Delta(2400)$	G_{39}	**	**	n						
$\Delta(2420)$	H_{311}	****	****							*
$\Delta(2750)$	I_{313}	**	**							
$\Delta(2950)$	K_{315}	**	**							

**** Existence is certain, and properties are at least fairly well explored.
 *** Existence ranges from very likely to certain, but further confirmation is desirable and/or quantum numbers, branching fractions, etc. are not well determined.
 ** Evidence of existence is only fair.
 * Evidence of existence is poor.

See key on page 1343

II. Elastic partial-wave analyses and determination of resonance parameters

(by G. Höhler, University of Karlsruhe and R.L. Workman, Virginia Polytechnic Institute and State University)

Our 1992 edition [4] gave a more general discussion. We here treat only new results.

New data: Most of the experimental activity in the past two years was restricted to the energy range from threshold up to the high side of the $\Delta(1232)$ [5]. There are still discrepancies amongst the new data, and final results of some important measurements are not yet available. These data are of interest for the charge splitting of the $\Delta(1232)$, and they are also of some relevance to other resonances, insofar as dispersion relations used in analyses need input from all momenta.

At higher energies, there have been published spin-rotation data measured at LAMPF [6] and at the Petersburg Nuclear Physics Institute [7], and also preliminary integrated πp elastic cross sections above 30° and up to 500 MeV from LAMPF [8].

New πN partial-wave analyses: Three new analyses of elastic data have been made since our 1992 edition. Abaev and Kruglov produced a set of single-energy analyses [9] covering from 160 to 600 MeV laboratory kinetic energy. These fits included experimental total inelastic cross sections and constraints on the tails of small amplitudes from the CMB80 [10] and KH80 [11] solutions. Marked deviations from the KH80 S- and P-wave amplitudes were found. The agreement was better with a new solution from Bugg [12], an analysis restricted to the range below 320 MeV.

The VPI group produced a new set of energy-dependent partial-wave analyses [13] with the improvements of including fixed- t dispersion relation constraints for the forward amplitude and for the invariant amplitudes A and B in the range $0 > t > -0.3 \text{ GeV}^2$. A grid of solutions was produced for different constraint values of pion-nucleon coupling and S-wave scattering lengths. A χ^2 mapping technique was used to determine which of the solutions gave an 'optimal' fit to both the data and the dispersion relation constraints.

Since the KH80 and CMB80 analyses and the new VPI analysis use dispersion relations as constraints, one may ask if it is justified to use these constraints at a time when QCD is assumed to be the theory of strong interactions. The positive answer was given by R. Oehme [14].

Argand plots for the partial wave amplitudes $T(W)$ which show the results of the CMB80 and KH80 analyses may be found in our 1990 edition [15], and tables of the amplitudes are given in [1].

Conventional resonance parameters: The conventional Breit-Wigner parameters were discussed in detail in our 1992 edition [4]. It is important to keep in mind that the width Γ is the value of $\Gamma(W)$ at the resonance energy $W = M$; it depends on the model used for the energy dependence and on the definition of M . The tabulated elasticities and partial widths are similarly model dependent. The model dependence

is the primary reason for the differences in the parameters and their errors quoted by the different groups.

Resonance pole parameters: We add some remarks to the treatment in the previous edition [4]. The resonance pole parameters have the advantage that they are directly related to the definition of a resonance as an unstable intermediate state in a scattering process [16].

Pole parameters for the KH partial wave solutions were determined only recently [17,18]. The method employs not only plots for the speed $|dT/dW|$, which were used in the early seventies, but also Argand plots for dT/dW and fits to the resonance semicircles in the range $M \pm \Gamma/2$ in Argand diagrams for $T(W)$. The parametrization for $T(W)$ agrees with that in [19], but the four parameters of the pole term are treated as constants, since they belong to the pole of $T(W)$. If the background amplitude is elastic, statements made in our 1992 edition [4] concerning the phase of the residue and the elasticity are correct. However, they disagree with the present parametrization in the general case. If the background can be treated as a constant, the model dependence of the pole parameters is weaker than in other methods. This can be assumed for eight of the 4-star resonances [18]. One could try to check if the method proposed in [19] can be applied in practice for a separation of the direct production of inelastic final states and a production by a resonance decay.

For further information on the status of a resonance and on the accuracy of its parameters, there will soon be available a collection of plots [20], calculated from various partial wave analyses.

The parameters of nucleon resonances will remain of interest in the future since they are needed to test various models, such as quark models [21] and Skyrmin models [22]. Finally, they will be needed as soon as these parameters can be calculated from lattice QCD.

References for Section I and II

Note: Many of the references below are to the πN Newsletter, which is edited by G. Höhler, W. Kluge, and B.M.K. Nefkens. Copies may be obtained from W. Kluge, Karlsruhe U. or from B.M.K. Nefkens, UCLA.

1. G. Höhler, *Pion-Nucleon Scattering*, Landolt-Börnstein Vol. I/9b2 (1983), ed. H. Schopper, Springer Verlag.
2. A.J.G. Hey and R.L. Kelly, *Phys. Reports* **96**, 71 (1983).
3. *Proceedings of the 5th International Symposium on Pion-Nucleon Physics and the Structure of the Nucleon*, ed. B.M.K. Nefkens and M. Clajus (Boulder, Sep. 1993), published in πN Newsletter No. 8 (1993) and πN Newsletter No. 9 (1994).
4. Review of Particle Properties, *Phys. Rev.* **D45**, (1 June 1992) Part II.
5. E. Friedman *et al.*, *Phys. Lett.* **B302**, 18 (1993); E. Friedman *et al.*, πN Newsletter No. 8 (1993), p. 11; M. Pavan *et al.*, πN Newsletter No. 8 (1993), p. 23; C. Joram *et al.*, πN Newsletter No. 8 (1993), p. 30; and references given in these papers.
6. I. Supek *et al.*, *Phys. Rev.* **D47**, 1762 (1993).

Baryon Full Listings

N 's and Δ 's

7. S.P. Kruglov *et al.*, πN Newsletter No. 8 (1993), p. 42 and references given therein.
8. B.J. Kriss *et al.*, πN Newsletter No. 8 (1993), p. 17.
9. V.V. Abaev and S.P. Kruglov, SPNPI preprint No. 1794, 1992;
S.P. Kruglov, πN Newsletter No. 8 (1993), p. 42.
10. R.E. Cutkosky *et al.*, Phys. Rev. **D20**, 2804 (1979); and *ibid.* 2839 (1979);
R.L. Kelly *et al.*, Phys. Rev. **D20**, 2782 (1979); and in *Proceedings of the IVth International Conference on Baryon Resonances* (Toronto, 1980), ed. N. Isgur, p. 19.
11. R. Koch, in *Proceedings of the IVth International Conference on Baryon Resonances* (Toronto, 1980), ed. N. Isgur, p. 3;
G. Höhler *et al.*, *Handbook of Pion-Nucleon Scattering*, Physics Data 12-1 (1979);
E. Pietarinen, Nucl. Phys. **B107**, 21 (1976).
12. D.V. Bugg, πN Newsletter No. 6 (1992), p. 7;
D.V. Bugg, πN Newsletter No. 8 (1993), p. 1.
13. R.A. Arndt, R.L. Workman, and M.M. Pavan, submitted to Phys. Rev. *D*.
14. R. Oehme, πN Newsletter No. 7 (1992), p. 1.
15. Review of Particle Properties, Phys. Lett. **B239** (1990).
16. R.H. Dalitz and R.G. Moorhouse, Proc. Roy. Soc. London **A318**, 279 (1970);
B.H. Bransden and R.G. Moorhouse, *The Pion-Nucleon System* (Princeton Univ. Press, 1973).
17. G. Höhler and A. Schulte, πN Newsletter No. 7 (1992), p. 94.
18. G. Höhler, πN Newsletter No. 9 (1994), p. 1.
19. P.H. Timmers *et al.*, Phys. Rev. **D31**, 99 (1985).
20. G. Höhler, "A Picture Book of Pion-Nucleon Resonances," Karlsruhe preprint TTP93-41, to be published in πN Newsletter.
21. S. Capstick and W. Robert, Phys. Rev. **D47**, 1994 (1993) and CEBAF-TH-93-18.
22. International Workshop on Baryons as Skyrme Solitons, ed. G. Holzwarth (World Scientific, 1993);
B. Schwesinger "Mesonen und Baryonen im Skyrme Modell," Lecture Notes, University of Siegen, Germany, FB Physik.

III. Inelastic two-body and quasi-two-body reactions

(by D.M. Manley, Kent State University)

Since our 1992 edition, no new data nor partial-wave analyses have been published for the inelastic two-body reactions $\pi N \rightarrow N\eta$, $\pi N \rightarrow \Lambda K$, and $\pi N \rightarrow \Sigma K$. There are recent indications [1] that much of the $\pi N \rightarrow N\eta$ data [2] used for the Rutherford partial-wave analysis [3] had poorly determined energies; thus, except for the $N(1535) S_{11}$, $N\eta$ branching fractions are omitted from the Baryon Summary Table in this edition. A recent review of these reactions and the quasi-two-body reaction $\pi N \rightarrow N\omega$ can be found in Ref. 4.

Essentially all information on quasi-two-body reactions such as $\pi N \rightarrow \Delta\pi$ and $\pi N \rightarrow N\rho$ comes from isobar-model analyses of $\pi N \rightarrow N\pi\pi$ reactions. Since the last edition, the Kent State multichannel resonance analysis has been published [5]. Although couplings from this analysis were listed in the 1992

edition, the present edition includes new, updated estimates for couplings and branching fractions of many resonances observed in two or more isobar-model analyses. A brief review of $\pi N \rightarrow N\pi\pi$ analyses can be found in the 1992 edition; for a more recent and extensive review, the interested reader should see Ref. 6.

Since the last edition, there have been two reports [7,8] of resonances observed in $\pi N \rightarrow N4\pi$ reactions. One report [7] of $\pi^- p \rightarrow p\pi^+\pi^-\pi^-\pi^0$ data from the CERN liquid-hydrogen bubble chamber gives evidence for an N^* resonance with mass 1700 MeV and width 150 MeV, which decays by the channels $p\eta$, $\Delta 2\pi$, $p3\pi$. The other report [8] of $\pi^+ p \rightarrow p\pi^+\pi^+\pi^-\pi^0$ data from the ZhVK-205 liquid-hydrogen bubble chamber (Moscow) gives evidence for an N^* resonance with mass 1780 ± 40 MeV and width 250 ± 80 MeV, which decays by the $p\omega$ channel.

References for Section III

1. M. Clajus and B.M.K. Nefkens, πN Newsletter No. 7 (1992), p. 76.
2. R.M. Brown *et al.*, Nucl. Phys. **B153**, 89 (1979).
3. R.D. Baker *et al.*, Nucl. Phys. **B156**, 93 (1979).
4. D.M. Manley in *New Vistas in Physics with High-Energy Pion Beams*, ed. B.F. Gibson and J.B. McClelland (World Scientific, 1993), p. 119.
5. D.M. Manley and E.M. Saleski, Phys. Rev. **D45**, 4002 (1992).
6. D.M. Manley, πN Newsletter No. 8 (1993), p. 141.
7. Yu.D. Aleshin and V.M. Guzhavin, Sov. J. Nucl. Phys. **55**, 1814 (1992) [*Yad. Fiz.* **55**, 3255 (1992)].
8. L.Yu. Brovkin *et al.*, Phys. Atom. Nucl. **56**, 762 (1993) [*Yad. Fiz.* **56**, 90 (1993)].

IV. Electromagnetic interactions

(by R.L. Crawford, University of Glasgow)

Nearly all the entries in the Listings relating to electromagnetic properties of the N and Δ resonances are couplings for decay to $N\gamma$. These couplings, the helicity amplitudes $A_{1/2}$ and $A_{3/2}$, have been obtained in a large number of partial-wave analyses of single-pion photoproduction, $\gamma N \rightarrow \pi N$, on protons and neutrons. The large amount of data has permitted an accurate evaluation of the couplings for many of the resonances with masses below 2 GeV, and has given at least qualitative information about most of the others. Most photoproduction analyses use as input information on the existence, masses, and widths of the resonances from the $\pi N \rightarrow \pi N$ analyses, and only determine the $N\gamma$ couplings. A brief description of the various methods of analysis of photoproduction data may be found in our 1992 edition [1].

The Listings omit a number of analyses that are now obsolete. Most of the older results may be found in our 1982 edition [2]. The errors quoted for the couplings in the Listings are calculated in different ways in different analyses and therefore should be used with care. In general, it is likely that the systematic differences between the analyses caused by using

different parametrization schemes are more indicative of the true uncertainties than are the quoted errors.

Probably the most reliable analyses are ARAI 80, CRAWFORD 80, AWAJI 81, FUJII 81, CRAWFORD 83, ARNDT 90B, and LI 93. The Listings include our estimates of the couplings, using the results of these analyses. The errors we give are a combination of the stated statistical errors on the analyses and the systematic differences between them. The analyses are given equal weight, except LI 93 is weighted, rather arbitrarily, by a factor of two because its data set is at least 50% larger than those of the other analyses and contains many new high-quality experimental measurements.

The Baryon Summary Table gives $N\gamma$ branching fractions for those resonances whose couplings are considered to be reasonably well established. The $N\gamma$ partial width Γ_γ is given in terms of the helicity amplitudes $A_{1/2}$ and $A_{3/2}$ by

$$\Gamma_\gamma = \frac{k^2}{\pi} \frac{2M_N}{(2J+1)M_R} [|A_{1/2}|^2 + |A_{3/2}|^2] .$$

Here M_N and M_R are the nucleon and resonance masses, J is the resonance spin, and k is the photon c.m. decay momentum.

The Listings contain several results for the E2/M1 ratio ($E_{1+}^{(3)}/M_{1+}^{(3)}$) of the $\Delta(1232)$. The analysis of TANABE 85 attempts to distinguish between renormalized and bare couplings; it is, therefore, a model-dependent result and should not be compared directly with the others. Similarly, Christillin and Dillon [3] have made a model-dependent estimate of the ratio. They do not fit directly to data, so their value does not appear in the Listings, but they conclude that the ratio is between zero and -2% . DAVIDSON 90 extracts the K -matrix residues for M_{1+} and E_{1+} using the data from several energy-independent analyses of the $\Delta(1232)$ region. Essentially, they are measuring $\text{Im}(E_{1+}^{(3)})/\text{Im}(M_{1+}^{(3)})$. The value quoted in the Listings is their average over several fits. DAVIDSON 91 uses an effective Lagrangian model to fit to various partial-wave analyses in the $\Delta(1232)$ region.

Christillin and Dillon point out that the spread in predictions may be due to the difference between bare and renormalized couplings. Since the separation of these is essentially model dependent, WORKMAN 92 measures $\text{Im}(E_{1+}^{(3)})/\text{Im}(M_{1+}^{(3)})$ directly from experimental data using the methods of ARNDT 90B. Whereas the other measurements are dependent on rather old partial-wave analyses, WORKMAN 92 uses recent experimental data. They also perform a fit, based on the model of Nozawa *et al.* [4], to extract "bare couplings," and get -3.9% .

The Listings also contain several measurements of the magnetic moment μ_Δ of the $\Delta(1232)^{++}$ from analyses of pion bremsstrahlung, $\pi^+p \rightarrow \gamma\pi^+p$. NEFKENS 78 is an analysis of UCLA data for pion bremsstrahlung that uses the soft pion model of Pascual and Tarrach [5]. HELLER 87 is a fit to the same data using a nonrelativistic dynamical model that measures the magnetic moment of the "bare" $\Delta(1232)^{++}$. LIN 91B

fits the data of NEFKENS 87 and from SIN [6] with an amplitude that includes the anomalous magnetic moment of the $\Delta(1232)$ and is relativistic, gauge invariant, and consistent with the soft photon theorem. The quantity measured is not identical to the "bare" magnetic moment since it does not take into account the effect of loop contributions. BOSSHARD 91 measured the polarized target asymmetry in pion-proton bremsstrahlung and fit it using the model of HELLER 87. The geometry of the experiment was chosen to maximize the sensitivity to μ_Δ .

See our 1992 edition [1] for brief discussions of Compton scattering, $K\Lambda$ photoproduction, and pion electroproduction.

References for Section IV

1. Particle Data Group, Phys. Rev. **D45**, 1 June 1992, Part II.
2. Particle Data Group, Phys. Lett. **111B** (1982).
3. P. Christillin and G. Dillon, J. Phys. **G15**, 967 (1989).
4. S. Nozawa, P. Blankleider, and T.-S.H. Lee, Nucl. Phys. **A513**, 459 (1990).
5. P. Pascual and R. Tarrach, Nucl. Phys. **B134**, 133 (1978).
6. C.A. Meyer *et al.*, Phys. Rev. **D38**, 754 (1988).

V. Outlook

(by D.M. Manley, Kent State University)

It is anticipated that much new data related to the study of nucleon resonances will come from experiments with electromagnetic probes at CEBAF, which will initially provide beams of electrons and photons with energies up to 4 GeV. Such experiments form the major component of the approved research program for CEBAF's Hall B, which is currently expected to begin operation in Fall, 1996. Other experiments to study nucleon resonances and/or elementary electroproduction or photoproduction reactions will be carried out in CEBAF's Hall A and Hall C, which are expected to be operation in mid-1995 and mid-1994, respectively. The main detector for the Hall-B experiments is the CLAS (CEBAF Large Acceptance Spectrometer), which, with CEBAF's 100% duty cycle, will make possible studies of electroproduction reactions such as $p(e, e'\pi^+)n$, $p(e, e'p)\pi^0$, $d(e, e'\pi^-)pp$, $p(e, e'p)\eta$, and $p(e, e'p)\omega$, among others. This facility will also use tagged photons to make high-precision measurements of the differential cross section for reactions such as π^0 , η , and η' photoproduction. These experiments are expected to improve determinations of the electromagnetic couplings of resonances such as the $\Delta(1232)P_{33}$, the $N(1520)D_{13}$, the $N(1535)S_{11}$, and the $N(1680)F_{15}$, which have large photocouplings; they are also expected to provide new information about resonances that couple weakly to the πN channel.

CEBAF is not the only laboratory where new experiments to study nucleon resonances will be carried out. For example, the new Bonn continuous beam machine ELSA will combine two large solid-angle detectors, PHOENICS and SAPHIR, with tagged photon facilities to increase our database from various photoproduction processes [1]. Other facilities are or will

Baryon Full Listings

N's and Δ's, N(1440)

be involved in such programs using hadronic beams. For example, plans are underway to study baryon spectroscopy at the Brookhaven National Laboratory AGS [2]. The reactions $\pi^-p \rightarrow n\eta$, $K^-p \rightarrow \Lambda\eta$, and $K^-p \rightarrow \Sigma^0\eta$ will be studied by using the SLAC Crystal Ball to identify multiphoton final states. As by-products of these investigations, new and improved data also will be obtained for $\pi^-p \rightarrow n\pi^0$ and for inverse photoproduction on an unstable target, in particular, $K^-p \rightarrow \Lambda\gamma$ and $K^-p \rightarrow \Sigma^0\gamma$.

References for Section V

1. D. Menze in "Baryon Spectroscopy and the Structure of the Nucleon," *Proceedings of an International Workshop at Saclay, France 23-25 Sep. 1991*, ed. H.P. Morsch and M. Soyeur (Research Center Conference Service, Jülich, 1992), p. 182.
2. AGS Letter of Intent, "Baryon Spectroscopy with the Crystal Ball," spokespersons B.M.K. Nefkens and R.E. Chrien (Aug. 1993).

N(1440) P₁₁

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } ****$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, *Physics Letters 111B* (1982).

N(1440) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1430 to 1470 (≈ 1440) OUR ESTIMATE			
1462 ± 10	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1440 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1410 ± 12	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
1465	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1471	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
1411	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1472	¹ BAKER 79	DPWA	$\pi^-p \rightarrow n\eta$
1417	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1460	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
1380	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1390	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1440) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
250 to 450 (≈ 350) OUR ESTIMATE			
391 ± 34	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
545 ± 170	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
340 ± 70	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
135 ± 10	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
315	LI 93	IPWA	$\gamma N \rightarrow \pi N$
334	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
113	¹ BAKER 79	DPWA	$\pi^-p \rightarrow n\eta$
331	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
279	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
200	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
200	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1440) POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1385	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1360	⁵ ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1370	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
1375 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
1359	ARNDT 85	DPWA	See ARNDT 91
1381 or 1379	⁶ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1360 or 1333	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
164	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
252	⁵ ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
228	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
180 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
200	ARNDT 85	DPWA	See ARNDT 91
209 or 210	⁶ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
167 or 234	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1440) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
109	⁵ ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
74	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
52 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-93	⁵ ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-84	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
-100 ± 35	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1440) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ _i /Γ)
Γ ₁ Nπ	60-70 %
Γ ₂ Nη	
Γ ₃ Nππ	30-40 %
Γ ₄ Δπ	20-30 %
Γ ₅ Δ(1232)π, P-wave	
Γ ₆ Nρ	<8 %
Γ ₇ Nρ, S=1/2, P-wave	
Γ ₈ Nρ, S=3/2, P-wave	
Γ ₉ N(ππ) _{S=0}	5-10 %
Γ ₁₀ pγ	0.04-0.07 %
Γ ₁₁ pγ, helicity=1/2	
Γ ₁₂ nγ	0.001-0.05 %
Γ ₁₃ nγ, helicity=1/2	

N(1440) BRANCHING RATIOS

Γ(Nπ)/Γ _{total}	DOCUMENT ID	TECN	COMMENT	Γ ₁ /Γ
0.6 to 0.7 OUR ESTIMATE				
0.69 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.68 ± 0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.51 ± 0.05	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

(Γ ₁ Γ ₇) ^{1/2} /Γ _{total} ln Nπ → N(1440) → Nη	DOCUMENT ID	TECN	COMMENT	(Γ ₁ Γ ₂) ^{1/2} /Γ
••• We do not use the following data for averages, fits, limits, etc. •••				
seen	¹ BAKER 79	DPWA	$\pi^-p \rightarrow n\eta$	
+0.328	⁷ FELTESSE 75	DPWA	1488-1745 MeV	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

(Γ ₁ Γ ₇) ^{1/2} /Γ _{total} ln Nπ → N(1440) → Δ(1232)π, P-wave	DOCUMENT ID	TECN	COMMENT	(Γ ₁ Γ ₅) ^{1/2} /Γ
+0.37 to +0.41 OUR ESTIMATE				
+0.39 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
+0.41	^{2,8} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.37	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

(Γ ₁ Γ ₇) ^{1/2} /Γ _{total} ln Nπ → N(1440) → Nρ, S=1/2, P-wave	DOCUMENT ID	TECN	COMMENT	(Γ ₁ Γ ₇) ^{1/2} /Γ
±0.07 to ±0.25 OUR ESTIMATE				
-0.11	^{2,8} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.23	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

(Γ ₁ Γ ₇) ^{1/2} /Γ _{total} ln Nπ → N(1440) → Nρ, S=3/2, P-wave	DOCUMENT ID	TECN	COMMENT	(Γ ₁ Γ ₈) ^{1/2} /Γ
±0.18				
+0.18	^{2,8} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	

See key on page 1343

Baryon Full Listings
N(1440), N(1520)

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \pi \rightarrow N(1440) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$	DOCUMENT ID	TECN	COMMENT
± 0.17 to ± 0.25 OUR ESTIMATE			
+0.24 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.18	2,8 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.23	3 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1440) PHOTON DECAY AMPLITUDES

N(1440) → pγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.072 ± 0.009 OUR ESTIMATE			
-0.085 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.069 ± 0.018	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.063 ± 0.008	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.069 ± 0.004	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.066 ± 0.004	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.079 ± 0.009	BRATASHEV... 80	DPWA	$\gamma N \rightarrow \pi N$
-0.068 ± 0.015	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.0584 ± 0.0148	ISHII 80	DPWA	Compton scattering
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.066 ± 0.017	ARNDT 90B	IPWA	See LI 93
-0.064	ARNDT 90B	FIT	See LI 93
-0.129	9 WADA 84	DPWA	Compton scattering
-0.075 ± 0.015	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.125	10 NOELLE 78	DPWA	$\gamma N \rightarrow \pi N$
-0.076	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
-0.087 ± 0.006	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1440) → nγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
$+0.052 \pm 0.025$ OUR ESTIMATE			
0.085 ± 0.006	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.037 ± 0.010	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.030 ± 0.003	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
0.023 ± 0.009	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.019 ± 0.012	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.056 ± 0.015	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.029 ± 0.035	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.050 ± 0.019	ARNDT 90B	IPWA	See LI 93
0.045	ARNDT 90B	FIT	See LI 93
+0.059 ± 0.016	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
0.062	10 NOELLE 78	DPWA	$\gamma N \rightarrow \pi N$

N(1440) FOOTNOTES

- BAKER 79 finds a coupling of the N(1440) to the Nη channel near (but slightly below) threshold.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to πN → Nππ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ARNDT 91 (Soln SM90) also finds a second-sheet pole with real part = 1413 MeV, -2 × imaginary part = 256 MeV, and residue = (78-153i) MeV.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to πN → Nππ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- An alternative which cannot be distinguished from this is to have a P₁₃ resonance with M = 1530 MeV, Γ = 79 MeV, and elasticity = +0.271.
- LONGACRE 77 considers this coupling to be well determined.
- WADA 84 is inconsistent with other analyses; see the Note on N and Δ Resonances.
- Converted to our conventions using M = 1486 MeV, Γ = 613 MeV from NOELLE 78.

N(1440) REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	(KENT) IJP
Also 84	PR D30 904	(VPI)
ARNDT 91	PR D43 2131	(VPI, TELE) IJP
ARNDT 90B	PR C42 1864	(VPI)
CUTKOSKY 90	PR D42 235	(CMU)
ARNDT 85	PR D32 1085	(VPI)
WADA 84	NP B247 313	(INUS)
CRAWFORD 83	NP B211 1	(GLAS)
PDG 82	PL 111B	(HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	(NAGO)
Also 82	NP B197 365	(NAGO)
FUJII 81	NP B187 53	(NAGO, OSAK)
ARAI 80	Toronto Conf. 93	(INUS)
Also 82	NP B194 251	(INUS)
BRATASHEV... 80	NP B166 525	(KFTI)
	+Arndt, Roper, Workman	
	+Saito, Arndt, Goradia, Tepitz	
	+Li, Roper, Workman, Ford	
	+Workman, Li, Roper	
	+Wang	
	+Ford, Roper	
	+Egawa, Imanishi, Ishii, Kato, Ukai+	
	+Morton	
	+Roos, Porter, Aguilar-Benitez+	
	+Kajikawa	
	+Fujii, Hayashii, Iwata, Kajikawa+	
	+Hayashii, Iwata, Kajikawa+	
	Arai, Fujii	
	Bratashevskij, Gorbenko, Derebchinskij+	

CRAWFORD 80	Toronto Conf. 107		(GLAS)
CUTKOSKY 90	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also 79	PR D20 2839	+Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
ISHII 80	NP B165 189	+Egawa, Kato, Miyachi+	(KYOT, INUS)
TAKEDA 80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+	(TOKY, INUS)
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons	(GLAS)
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
NOELLE 78	PTP 60 778		(NAGO)
BERENDS 77	NP B136 317	+Donnachie	(LEID, MCHS) IJP
LONGACRE 77	NP B122 493	+Dolbeau	(SACL) IJP
Also 76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
FELLER 76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
FELTESSE 75	NP B93 242	+Ayed, Bareyre, Borgeaud, David+	(SACL) IJP
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP

N(1520) D₁₃

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

N(1520) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1518 to 1530 (≈ 1520) OUR ESTIMATE			
1524 ± 4	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
1525 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1519 ± 4	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1510	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1504	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1503	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1510	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
1510	1 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1520	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1520) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
110 to 135 (≈ 120) OUR ESTIMATE			
124 ± 8	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
120 ± 15	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
114 ± 7	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
120	LI 93	IPWA	$\gamma N \rightarrow \pi N$
124	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
183	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
135	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
105	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
110	1 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
150	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1520) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1510	3 HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
1511	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1510 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1510	ARNDT 85	DPWA	See ARNDT 91
1514 or 1511	4 LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1508 or 1505	1 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120	3 HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
108	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
114 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
122	ARNDT 85	DPWA	See ARNDT 91
146 or 137	4 LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
109 or 107	1 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1520) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
32	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
33	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
35 ± 2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

Baryon Full Listings

N(1520)

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-8	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
-10	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-12±5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1520) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	50-60 %
Γ_2 $N\eta$	
Γ_3 $N\pi\pi$	40-50 %
Γ_4 $\Delta\pi$	15-25 %
Γ_5 $\Delta(1232)\pi$, S-wave	5-12 %
Γ_6 $\Delta(1232)\pi$, D-wave	10-14 %
Γ_7 $N\rho$	15-25 %
Γ_8 $N\rho$, S=1/2, D-wave	
Γ_9 $N\rho$, S=3/2, S-wave	
Γ_{10} $N\rho$, S=3/2, D-wave	
Γ_{11} $N(\pi\pi)_{S=0}^{I=0}$	<8 %
Γ_{12} $p\gamma$	0.45-0.53 %
Γ_{13} $p\gamma$, helicity=1/2	
Γ_{14} $p\gamma$, helicity=3/2	
Γ_{15} $n\gamma$	0.34-0.48 %
Γ_{16} $n\gamma$, helicity=1/2	
Γ_{17} $n\gamma$, helicity=3/2	

N(1520) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.5 to 0.6 OUR ESTIMATE				
0.59±0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.58±0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.54±0.03	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow N(1520) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.5 to 0.6 OUR ESTIMATE				
0.02	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	
+0.011	FELTESSE 75	DPWA	Soln A; see BAKER 79	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow N(1520) \rightarrow \Delta(1232)\pi$, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.26 to -0.20 OUR ESTIMATE				
-0.18±0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
-0.26	1,5 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.24	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow N(1520) \rightarrow \Delta(1232)\pi$, D-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.28 to -0.24 OUR ESTIMATE				
-0.29±0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
-0.21	1,5 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.30	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow N(1520) \rightarrow N\rho$, S=3/2, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.35 to -0.31 OUR ESTIMATE				
-0.35±0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
-0.35	1,5 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.24	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow N(1520) \rightarrow N(\pi\pi)_{S=0}^{I=0}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.22 to -0.06 OUR ESTIMATE				
-0.13	1,5 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.17	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

N(1520) PHOTON DECAY AMPLITUDES

N(1520) $\rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.022 ± 0.008 OUR ESTIMATE			
-0.020 ± 0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.028 ± 0.014	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.007 ± 0.004	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.032 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.032 ± 0.004	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.031 ± 0.009	BRATASHEV... 80	DPWA	$\gamma N \rightarrow \pi N$
-0.019 ± 0.007	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.0430 ± 0.0063	ISHII 80	DPWA	Compton scattering
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.025 ± 0.009	ARNDT 908	IPWA	See LI 93
-0.023	ARNDT 908	FIT	See LI 93
-0.012	WADA 84	DPWA	Compton scattering
-0.016 ± 0.008	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.008	6 NOELLE 78		$\gamma N \rightarrow \pi N$
-0.021	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
-0.005 ± 0.005	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1520) $\rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.163 ± 0.007 OUR ESTIMATE			
0.167 ± 0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.156 ± 0.022	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.168 ± 0.013	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.178 ± 0.003	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.162 ± 0.003	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.166 ± 0.005	BRATASHEV... 80	DPWA	$\gamma N \rightarrow \pi N$
0.167 ± 0.010	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
0.1695 ± 0.0014	ISHII 80	DPWA	Compton scattering
••• We do not use the following data for averages, fits, limits, etc. •••			
0.155 ± 0.006	ARNDT 908	IPWA	See LI 93
0.167	ARNDT 908	FIT	See LI 93
0.168	WADA 84	DPWA	Compton scattering
+0.157 ± 0.007	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
0.206	6 NOELLE 78		$\gamma N \rightarrow \pi N$
+0.075	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
+0.164 ± 0.008	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1520) $\rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.062 ± 0.006 OUR ESTIMATE			
-0.058 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.066 ± 0.013	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.067 ± 0.004	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
-0.076 ± 0.006	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.071 ± 0.011	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.056 ± 0.011	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.050 ± 0.014	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.059 ± 0.014	ARNDT 908	IPWA	See LI 93
-0.063	ARNDT 908	FIT	See LI 93
-0.055 ± 0.014	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.060	6 NOELLE 78		$\gamma N \rightarrow \pi N$

N(1520) $\rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.137 ± 0.013 OUR ESTIMATE			
-0.131 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.124 ± 0.009	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.158 ± 0.003	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
-0.147 ± 0.008	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.148 ± 0.009	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.144 ± 0.015	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.118 ± 0.011	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.126 ± 0.015	ARNDT 908	IPWA	See LI 93
-0.135	ARNDT 908	FIT	See LI 93
-0.141 ± 0.015	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.127	6 NOELLE 78		$\gamma N \rightarrow \pi N$

N(1520) FOOTNOTES

¹ LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

² From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.

³ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

See key on page 1343

Baryon Full Listings

$N(1520), N(1535)$

⁴ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
⁵ LONGACRE 77 considers this coupling to be well determined.
⁶ Converted to our conventions using $M = 1528$ MeV, $\Gamma = 187$ MeV from NOELLE 78.

$N(1520)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982). For very early references, see Reviews of Modern Physics **37** 633 (1965).

Author	Year	Reference	Technique	Comment
HOEHLER	94	πN Newsletter 10 (to be pub.)	(KARL)	
HOEHLER	93	πN Newsletter 9 1	(KARL)	
LI	93	PR C47 2759	(VPI)	
MANLEY	92	PR D45 4002	(KENT) IJP	
Also	84	PR D30 904	(VPI)	
ARNDT	91	PR D43 2131	(VPI, TELE) IJP	
ARNDT	90B	PR C42 1864	(VPI)	
ARNDT	85	PR D32 1085	(VPI)	
WADA	85	NP B247 313	(INUS)	
CRAWFORD	83	NP B211 1	(GLAS)	
PDG	82	PL 111B	(HELS, CIT, CERN)	
AWAJI	81	Bonn Conf. 352	(NAGO)	
Also	82	NP B197 365	(NAGO)	
FUJII	81	NP B187 53	(NAGO, OSAK)	
ARAI	80	Toronto Conf. 93	(INUS)	
Also	80	NP B194 251	(INUS)	
BRATASHEV...	80	NP B166 525	(KFTI)	
CRAWFORD	80	Toronto Conf. 107	(GLAS)	
CUTKOSKY	80	Toronto Conf. 19	(CMU, LBL) IJP	
Also	79	PR D20 2839	(CMU, LBL) IJP	
ISHII	80	NP B165 189	(KYOT, INUS)	
TAKEDA	80	NP B168 17	(TOKY, INUS)	
BAKER	79	NP B156 93	(RHEL) IJP	
HOEHLER	79	PDAT 12-1	(KARLT) IJP	
Also	80	Toronto Conf. 3	(KARLT) IJP	
BARBOUR	78	NP B141 253	(GLAS)	
LONGACRE	78	PR D17 1795	(LBL, SLAC)	
NOELLE	78	PTP 60 778	(NAGO)	
BERENDS	77	NP B136 317	(LEID, MCHS) IJP	
LONGACRE	77	NP B122 493	(SACL) IJP	
Also	76	NP B108 365	(SACL) IJP	
FELLER	76	NP B104 219	(NAGO, OSAK) IJP	
FELTESSE	75	NP B93 242	(SACL) IJP	
LONGACRE	75	PL 55B 415	(LBL, SLAC) IJP	

$N(1535) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

$N(1535)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1520 to 1555 (≈ 1535) OUR ESTIMATE			
1534 ± 7	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1550 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1526 ± 7	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1518	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1513	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1511	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1500	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
1547 ± 6	BHANDARI 77	DPWA	Uses $N\eta$ cusp
1520	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1510	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1535)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100 to 250 (≈ 150) OUR ESTIMATE			
151 ± 27	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
240 ± 80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
120 ± 20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
84	LI 93	IPWA	$\gamma N \rightarrow \pi N$
136	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
180	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
132	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
57	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
139 ± 33	BHANDARI 77	DPWA	Uses $N\eta$ cusp
135	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
100	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1535)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1487	³ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1499	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1510 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1461	ARNDT 85	DPWA	See ARNDT 91
1496 or 1499	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1519 ± 4	BHANDARI 77	DPWA	Uses $N\eta$ cusp
1525 or 1527	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
110	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
260 ± 80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
140	ARNDT 85	DPWA	See ARNDT 91
103 or 105	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
140 ± 32	BHANDARI 77	DPWA	Uses $N\eta$ cusp
135 or 123	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1535)$ ELASTIC POLE RESIDUE

MODULUS $ r $ VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
120 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ VALUE (°)	DOCUMENT ID	TECN	COMMENT
-13	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
+15 ± 45	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$N(1535)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	35-55 %
Γ_2 $N\eta$	30-55 %
Γ_3 $N\pi\pi$	1-10 %
Γ_4 $\Delta\pi$	<1 %
Γ_5 $\Delta(1232)\pi, D\text{-wave}$	
Γ_6 $N\rho$	<4 %
Γ_7 $N\rho, S=1/2, S\text{-wave}$	
Γ_8 $N\rho, S=3/2, D\text{-wave}$	
Γ_9 $N(\pi\pi)_{S\text{-wave}}^{I=0}$	<3 %
Γ_{10} $N(1440)\pi$	<7 %
Γ_{11} $p\gamma$	0.45-0.53 %
Γ_{12} $p\gamma, \text{helicity}=1/2$	
Γ_{13} $n\gamma$	0.34-0.48 %
Γ_{14} $n\gamma, \text{helicity}=1/2$	

$N(1535)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.35 to 0.55 OUR ESTIMATE				
0.51 ± 0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.50 ± 0.10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.38 ± 0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.297 ± 0.026	BHANDARI 77	DPWA	Uses $N\eta$ cusp	

$(\Gamma_1/\Gamma)^{1/2}/\Gamma_{\text{total}}$ $\ln N\pi \rightarrow N(1535) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma)^{1/2}/\Gamma$
+0.44 to +0.50 OUR ESTIMATE				
+0.47 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.33	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	
+0.48	FELTESSE 75	DPWA	1488-1745 MeV	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1/\Gamma)^{1/2}/\Gamma_{\text{total}}$ $\ln N\pi \rightarrow N(1535) \rightarrow \Delta(1232)\pi, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma)^{1/2}/\Gamma$
-0.04 to +0.06 OUR ESTIMATE				
+0.00 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.00	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.06	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

Baryon Full Listings

 $N(1535)$, $N(1650)$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1535) \rightarrow N\rho, S=1/2, S\text{-wave}$	DOCUMENT ID	TECN	COMMENT
-0.14 to -0.06 OUR ESTIMATE			
-0.10 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
-0.10	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.09	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1535) \rightarrow N(\pi\pi)_{S=0}^{J=0}$	DOCUMENT ID	TECN	COMMENT
+0.03 to +0.13 OUR ESTIMATE			
+0.07 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
+0.08	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.09	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1535) \rightarrow N(1440)\pi$	DOCUMENT ID	TECN	COMMENT
+0.10 ± 0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$

 $N(1535)$ PHOTON DECAY AMPLITUDES $N(1535) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.068 ± 0.010 OUR ESTIMATE			
0.061 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.095 ± 0.011	⁵ BENMERROU...91		$\gamma p \rightarrow p\eta$
0.053 ± 0.015	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.077 ± 0.021	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.083 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.080 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.029 ± 0.007	BRATASHEV...80	DPWA	$\gamma N \rightarrow \pi N$
0.065 ± 0.016	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
0.0704 ± 0.0091	ISHII 80	DPWA	Compton scattering
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.078	ARNDT 90B	IPWA	See LI 93
0.050	ARNDT 90B	FIT	See LI 93
0.055	WADA 84	DPWA	Compton scattering
+0.082 ± 0.019	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
0.046	⁶ NOELLE 78		$\gamma N \rightarrow \pi N$
+0.034	BERENDS 77	IPWA	$\gamma N \rightarrow \pi N$
+0.070 ± 0.004	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

 $N(1535) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.059 ± 0.022 OUR ESTIMATE			
-0.046 ± 0.005	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.035 ± 0.014	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.062 ± 0.003	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
-0.075 ± 0.019	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.075 ± 0.018	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.098 ± 0.026	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.011 ± 0.017	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.050	ARNDT 90B	IPWA	See LI 93
-0.037	ARNDT 90B	FIT	See LI 93
-0.112 ± 0.034	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.048	⁶ NOELLE 78		$\gamma N \rightarrow \pi N$

 $N(1535)$ FOOTNOTES

- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- BENMERROUCHE 91 uses an effective Lagrangian approach to analyze η photoproduction data.
- Converted to our conventions using $M = 1548$ MeV, $\Gamma = 73$ MeV from NOELLE 78.

 $N(1535)$ REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	+Arndt, Roper, Workman (VPI)
MANLEY 92	PR D45 4002	+Saleski (KENT) IJP
Also 84	PR D30 904	+Manley, Arndt, Goradia, Tepitz (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
BENMERROU...91	PRL 67 1070	+Benmerrouche, Mukhopadhyay (RPI)
ARNDT 90B	PR C42 1864	+Workman, Li, Roper (VPI)
ARNDT 85	PR D32 1085	+Ford, Roper (VPI)
WADA 84	NP B247 313	+Egawa, Imanishi, Ishii, Kato, Ukai+ (INUS)
CRAWFORD 83	NP B211 1	+Morton (GLAS)
PDG 82	PL 1118	+Roos, Porter, Agular-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa (NIAGO)
Also 82	NP B197 365	+Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
FUJII 81	NP B187 53	+Hayashii, Iwata, Kajikawa+ (NAGO, OSAK)
ARAI 80	Toronto Conf. 93	(INUS)
Also 82	NP B194 251	+Arai, Fujii (INUS)
BRATASHEV...80	NP B166 525	+Bratashvskij, Gorbenko, Derebchinskij+ (KFTI)
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	+Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL)
ISHII 80	NP B165 189	+Egawa, Kato, Miyachi+ (KYOT, INUS)
TAKEDA 80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+ (TOKY, INUS)
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	+Koch (KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+ (LBL, SLAC)
NOELLE 78	PTP 60 778	(NAGO)
BERENDS 77	NP B136 317	+Donnachie (LEID, MCHS) IJP
BHANDARI 77	PR D15 192	+Chao (CMU) IJP
LONGACRE 77	NP B122 493	+Dolbeau (SACL) IJP
Also 76	NP B108 365	+Dolbeau, Triantis, Neveu, Cadiet (SACL) IJP
FELLER 76	NP B104 219	+Fukushima, Horikawa, Kajikawa+ (NAGO, OSAK) IJP
FELTESSE 75	NP B93 242	+Ayed, Barye, Borgeaud, David+ (SACL) IJP
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+ (LBL, SLAC) IJP

 $N(1650) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

 $N(1650)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1640 to 1660 (≈ 1650) OUR ESTIMATE			
1659 ± 9	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1650 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1670 ± 8	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1674	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1688	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1672	MUSETTE 80	IPWA	$\pi^- p \rightarrow \Lambda K^0$
1680	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1680	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1694	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1700 ± 5	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
1680	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1700	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1675	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1660	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $N(1650)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
145 to 190 (≈ 150) OUR ESTIMATE			
173 ± 12	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
150 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
180 ± 20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
225	LI 93	IPWA	$\gamma N \rightarrow \pi N$
183	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
179	MUSETTE 80	IPWA	$\pi^- p \rightarrow \Lambda K^0$
120	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
90	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
193	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
130 ± 10	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
90	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
170	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
170	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
130	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

See key on page 1343

Baryon Full Listings
N(1650)

N(1650) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1670	⁴ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
1657	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1640±20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1660	ARNDT 85	DPWA	See ARNDT 91
1648 or 1651	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1699 or 1698	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
163	⁴ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
160	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
150±30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
122	ARNDT 85	DPWA	See ARNDT 91
117 or 119	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
174 or 173	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1650) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
39	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
54	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
60±10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-37	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
-38	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-75±25	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1650) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	60-80 %
Γ_2 $N\eta$	
Γ_3 ΛK	3-11 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	5-20 %
Γ_6 $\Delta\pi$	3-7 %
Γ_7 $\Delta(1232)\pi$, D-wave	
Γ_8 $N\rho$	4-14 %
Γ_9 $N\rho$, S=1/2, S-wave	
Γ_{10} $N\rho$, S=3/2, D-wave	
Γ_{11} $N(\pi\pi)_{S=0}^{J=0}$	<4 %
Γ_{12} $N(1440)\pi$	<5 %
Γ_{13} $p\gamma$	0.10-0.18 %
Γ_{14} $p\gamma$, helicity=1/2	
Γ_{15} $n\gamma$	0.03-0.18 %
Γ_{16} $n\gamma$, helicity=1/2	

N(1650) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.6 to 0.8 OUR ESTIMATE				
0.89±0.07	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.65±0.10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.61±0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)_{1/2}^2/\Gamma$
0.09	⁶ BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)_{1/2}^2/\Gamma$
0.27 to -0.17 OUR ESTIMATE				
-0.22	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
-0.22	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.25	⁷ BAKER 78	DPWA	See SAXON 80	
-0.23±0.01	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$	
-0.25	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
0.12	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

 $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow \Sigma K$ $(\Gamma_1\Gamma_4)_{1/2}^2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.254	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$
0.066 to 0.137	⁸ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$
0.20	KNASEL 75	DPWA	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

 $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow \Delta(1232)\pi$, D-wave $(\Gamma_1\Gamma_7)_{1/2}^2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.15 to 0.23 OUR ESTIMATE			
+0.12±0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.29	^{2,9} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.15	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow N\rho$, S=1/2, S-wave $(\Gamma_1\Gamma_9)_{1/2}^2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
±0.03 to ±0.19 OUR ESTIMATE			
-0.01±0.09	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.17	^{2,9} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.16	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow N\rho$, S=3/2, D-wave $(\Gamma_1\Gamma_{10})_{1/2}^2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.17 to +0.29 OUR ESTIMATE			
+0.16±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.29	^{2,9} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

 $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow N(\pi\pi)_{S=0}^{J=0}$ S-wave $(\Gamma_1\Gamma_{11})_{1/2}^2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.04 to +0.18 OUR ESTIMATE			
+0.12±0.08	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.00	^{2,9} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.25	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln N\pi \rightarrow N(1650) \rightarrow N(1440)\pi$ $(\Gamma_1\Gamma_{12})_{1/2}^2/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.11±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$

N(1650) PHOTON DECAY AMPLITUDES

N(1650) $\rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.082±0.017 OUR ESTIMATE			
0.068±0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.033±0.015	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.050±0.010	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.065±0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.061±0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.031±0.017	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.091	WADA 84	DPWA	Compton scattering
+0.048±0.017	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
+0.068±0.009	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1650) $\rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.011±0.028 OUR ESTIMATE			
-0.002±0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.008±0.004	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.004±0.004	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
0.010±0.020	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.008±0.019	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.068±0.040	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.011±0.011	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.045±0.024	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1650) $\gamma p \rightarrow \Lambda K^+$ AMPLITUDES $(\Gamma_i/\Gamma)_{1/2}^2/\Gamma_{total} \ln p\gamma \rightarrow N(1650) \rightarrow \Lambda K^+$ (E_{0+} amplitude)

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN
7.8 ± 0.3	WORKMAN 90	DPWA
8.13	TANABE 89	DPWA

• • • We do not use the following data for averages, fits, limits, etc. • • •

Baryon Full Listings

$N(1650), N(1675)$

$p\gamma \rightarrow N(1650) \rightarrow \Lambda K^+$ phase angle θ (E_0 amplitude)

VALUE (degrees)	DOCUMENT ID	TECN
-107 ± 3	WORKMAN 90	DPWA
-107.8	TANABE 89	DPWA

N(1650) FOOTNOTES

- The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- BAKER 79 fixed this coupling during fitting, but the negative sign relative to the $N(1535)$ is well determined.
- The overall phase of BAKER 78 couplings has been changed to agree with previous conventions. Superseded by SAXON 80.
- The range given for DEANS 75 is from the four best solutions.
- LONGACRE 77 considers this coupling to be well determined.

N(1650) REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	(KENT) IJP
Also 84	PR D30 904	(VPI)
ARNDT 91	PR D43 2131	(VPI, TELE) IJP
WORKMAN 90	PR C42 781	(VPI)
TANABE 89	PR C39 741	(MANZ)
Also 89	NC 102A 193	(MANZ)
ARNDT 85	PR D32 1085	(VPI)
WADA 84	NP B247 313	(INUS)
BELL 83	NP B222 389	(RL) IJP
CRAWFORD 83	NP B211 1	(GLAS)
PDG 82	PL 111B	(HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	(NAGO)
Also 82	NP B197 365	(NAGO)
FUJII 81	NP B187 53	(NAGO, OSAK)
ARAI 80	Toronto Conf. 93	(INUS)
Also 82	NP B194 251	(GLAS)
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	(CMU, LBL) IJP
Also 79	PR D20 2839	(CMU, LBL) IJP
LIVANOS 80	Toronto Conf. 35	(SACL) IJP
MUSETTE 80	NC 57A 37	(BRUX) IJP
SAXON 80	NP B162 522	(RHEL, BRIS) IJP
TAKEDA 80	NP B168 17	(TOKY, INUS)
BAKER 79	NP B156 93	(RHEL) IJP
HOEHLER 79	PDAT 12-1	(KARLT) IJP
Also 80	Toronto Conf. 3	(KARLT) IJP
BAKER 78	NP B141 29	(RL, CAVE) IJP
BARBOUR 78	NP B141 253	(GLAS)
LONGACRE 78	PR D17 1795	(LBL, SLAC)
BAKER 77	NP B126 365	(RHEL) IJP
LONGACRE 77	NP B122 493	(SACL) IJP
Also 76	NP B108 365	(SACL) IJP
FELLER 76	NP B104 219	(NAGO, OSAK) IJP
DEANS 75	NP B96 90	(SFLA, ALAH) IJP
KNASEL 75	PR D11 1	(CHIC, WUSL, OSU, ANL) IJP
LONGACRE 75	PL 55B 415	(LBL, SLAC) IJP

$N(1675) D_{15}$

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

N(1675) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1670 to 1685 (≈ 1675) OUR ESTIMATE			
1676 ± 2	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
1675 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1679 ± 8	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1666	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1685	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1670	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1680	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1650	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1660	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1675) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
140 to 180 (≈ 150) OUR ESTIMATE			
159 ± 7	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
160 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
120 ± 15	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
136	LI 93	IPWA	$\gamma N \rightarrow \pi N$
191	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
40	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
88	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
192	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
130	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
150	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1675) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1656	³ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
1655	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1660 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1661	ARNDT 85	DPWA	See ARNDT 91
1663 or 1668	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1649 or 1650	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
126	³ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
124	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
140 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
142	ARNDT 85	DPWA	See ARNDT 91
146 or 171	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
127 or 127	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1675) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
23	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
28	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
31 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-22	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
-17	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-30 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1675) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	40-50 %
Γ_2 $N\eta$	
Γ_3 ΛK	< 1 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	50-60 %
Γ_6 $\Delta\pi$	50-60 %
Γ_7 $\Delta(1232)\pi$, D-wave	
Γ_8 $\Delta(1232)\pi$, G-wave	
Γ_9 $N\rho$	< 1-3 %
Γ_{10} $N\rho$, S=1/2, D-wave	
Γ_{11} $N\rho$, S=3/2, D-wave	
Γ_{12} $N\rho$, S=3/2, G-wave	
Γ_{13} $N(\pi\pi)_{S=0}^{I=0}$	
Γ_{14} $p\gamma$	0.005-0.014 %
Γ_{15} $p\gamma$, helicity=1/2	
Γ_{16} $p\gamma$, helicity=3/2	
Γ_{17} $n\gamma$	0.07-0.11 %
Γ_{18} $n\gamma$, helicity=1/2	
Γ_{19} $n\gamma$, helicity=3/2	

N(1675) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.47 ± 0.02	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.38 ± 0.05	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.38 ± 0.03	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.07	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$	
+0.009	FELTESSE	75	DPWA Soln A; see BAKER 79	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
-0.01	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$	
+0.036	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$	
-0.034 ± 0.006	DEVENISH	74B	Fixed-t dispersion rel.	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
<0.003	DEANS	75	DPWA $\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow \Delta(1232)\pi, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
+0.496 ± 0.003	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
+0.46	LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	
+0.50	LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$	
+0.5	NOVOSELLER	78	IPWA $\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow N\rho, S=1/2, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{10})^{1/2}/\Gamma$
+0.04 ± 0.02	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow N\rho, S=3/2, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{11})^{1/2}/\Gamma$
-0.03 ± 0.02	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
-0.15	LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1675) \rightarrow N(\pi\pi)_{S=0}^{I=0}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{13})^{1/2}/\Gamma$
+0.03	LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	

N(1675) PHOTON DECAY AMPLITUDES

N(1675) → pγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.012 ± 0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.021 ± 0.011	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.034 ± 0.005	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.006 ± 0.005	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.006 ± 0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.023 ± 0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
+0.022 ± 0.010	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.034 ± 0.004	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

N(1675) → pγ, helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.021 ± 0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.015 ± 0.009	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.024 ± 0.008	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.030 ± 0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.029 ± 0.004	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.003 ± 0.012	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
+0.015 ± 0.006	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.019 ± 0.009	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

N(1675) → nγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.050 ± 0.014 OUR ESTIMATE			
-0.060 ± 0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.057 ± 0.024	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.033 ± 0.004	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.039 ± 0.017	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.025 ± 0.027	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.059 ± 0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.021 ± 0.011	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$
-0.066 ± 0.020	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

N(1675) → nγ, helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.070 ± 0.006 OUR ESTIMATE			
-0.074 ± 0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.077 ± 0.018	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.069 ± 0.004	FUJII	81	DPWA $\gamma N \rightarrow \pi N$
-0.066 ± 0.026	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.071 ± 0.022	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.059 ± 0.020	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
-0.030 ± 0.012	TAKEDA	80	DPWA $\gamma N \rightarrow \pi N$
-0.073 ± 0.014	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

N(1675) FOOTNOTES

- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to πN → Nππ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- SAXON 80 finds the coupling phase is near 90°.
- The range given is from the four best solutions. DEANS 75 disagrees with π⁺p → Σ⁺K⁺ data of WINNIK 77 around 1920 MeV.
- LONGACRE 77 considers this coupling to be well determined.
- A Breit-Wigner fit to the HERNDON 75 IPWA.

N(1675) REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	π N Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	π N Newsletter 9 1	(KARL)
LI	PR C47 2759	+Arndt, Roper, Workman (VPI)
MANLEY 92	PR D45 4002	+Saleski (KENT) IJP
Also 84	PR D30 904	Manley, Arndt, Godard, Teplitz (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
ARNDT 85	PR D32 1085	+Ford, Roper (VPI)
BELL 83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+ (RL) IJP
CRAWFORD 83	NP B211 1	+Morton (GLAS)
PDG 82	PL 111B	Roos, Porter, Agular-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Dorn Conf. 352	+Kajikawa (NAGO)
Also 82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
FUJII 81	NP B187 53	+Hayashii, Iwata, Kajikawa+ (NAGO, OSAK)
ARAI 80	Toronto Conf. 93	(INUS)
Also 82	NP B194 251	Arai, Fujii (INUS)
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+ (RHEL, BRIS) IJP
TAKEDA 80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+ (TOKY, INUS)
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smdja+ (LBL, SLAC)
NOVOSELLER 80	NP B137 509	(CIT) IJP
Also 78B	NP B137 445	Novoseller (SACL) IJP
LONGACRE 77	NP B122 493	+Dolbeau (SACL) IJP
Also 76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet (SACL) IJP
WINNIK 77	NP B128 66	+Toaff, Revel, Goldberg, Berny (HAIF) IJP
FELLER 76	NP B104 219	+Fukushima, Horikawa, Kajikawa+ (NAGO, OSAK) IJP
DEANS 75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
FELTESSE 75	PR D11 3183	+Ayed, Baryre, Borgeaud, David+ (SACL) IJP
HERNDON 75	PL 55B 415	+Longacre, Miller, Rosenfeld+ (LBL, SLAC)
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smdja+ (LBL, SLAC) IJP
DEVENISH 74B	NP B81 330	+Froggatt, Martin (DESY, NORD, LOUC)

Baryon Full Listings

N(1680)

N(1680) F₁₅

$$I(J^P) = \frac{1}{2}(\frac{5}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

N(1680) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1675 to 1690 (≈ 1680) OUR ESTIMATE			
1684 ± 4	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1680 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1684 ± 3	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1682	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1680	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1660	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1685	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1670	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1680) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120 to 140 (≈ 130) OUR ESTIMATE			
139 ± 8	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
120 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
128 ± 8	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
121	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
119	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
150	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
155	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
130	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1680) POLE POSITION

REAL PART			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1673	³ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
1670	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1667 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1680	ARNDT 85	DPWA	See ARNDT 91
1668 or 1674	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1656 or 1653	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-2xIMAGINARY PART			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
135	³ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
116	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
110 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
120	ARNDT 85	DPWA	See ARNDT 91
132 or 137	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
145 or 143	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1680) ELASTIC POLE RESIDUE

MODULUS r			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
44	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
37	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
34 ± 2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
PHASE θ			
VALUE (°)	DOCUMENT ID	TECN	COMMENT
-17	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
-14	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-25 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1680) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	60-70 %
Γ_2 $N\eta$	
Γ_3 ΛK	
Γ_4 ΣK	
Γ_5 $N\pi\pi$	30-40 %
Γ_6 $\Delta\pi$	5-15 %
Γ_7 $\Delta(1232)\pi$, P-wave	6-14 %
Γ_8 $\Delta(1232)\pi$, F-wave	<2 %
Γ_9 $N\rho$	3-15 %
Γ_{10} $N\rho$, S=1/2, F-wave	
Γ_{11} $N\rho$, S=3/2, P-wave	<12 %
Γ_{12} $N\rho$, S=3/2, F-wave	1-5 %
Γ_{13} $N(\pi\pi)_{S=0}^{J=0}$	5-20 %
Γ_{14} $p\gamma$	0.21-0.35 %
Γ_{15} $p\gamma$, helicity=1/2	
Γ_{16} $p\gamma$, helicity=3/2	
Γ_{17} $n\gamma$	0.02-0.04 %
Γ_{18} $n\gamma$, helicity=1/2	
Γ_{19} $n\gamma$, helicity=3/2	

N(1680) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.6 to 0.7 OUR ESTIMATE				
0.70 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.62 ± 0.05	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.65 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
not seen	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

$\Gamma(N\eta)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.0005 or 0.001	⁵ CARRERAS 70	MPWA	t pole + resonance	
0.0004	⁵ BOTKE 69	MPWA	t pole + resonance	
0.003 ± 0.002	⁵ DEANS 69	MPWA	t pole + resonance	

$\Gamma(N\eta)/\Gamma(N\pi)$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.027	HEUSCH 66	RVUE	π^0, η photoproduction	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
Coupling to ΛK not required in the analyses of BAKER 77, SAXON 80, or BELL 83.				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.01	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
-0.009 ± 0.009	DEVENISH 74B		Fixed-t dispersion rel.	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<0.001	⁶ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1680) \rightarrow \Delta(1232)\pi$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
-0.31 to -0.21 OUR ESTIMATE				
-0.26 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
-0.27	^{1,7} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.25	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.38	⁸ NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	

See key on page 1343

Baryon Full Listings

$N(1680), N(1700)$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}} \ln N\pi \rightarrow N(1680) \rightarrow \Delta(1232)\pi, F\text{-wave}$ $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.03 to +0.11 OUR ESTIMATE			
+0.07 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
+0.07	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.08	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.05	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}} \ln N\pi \rightarrow N(1680) \rightarrow N\rho, S=3/2, P\text{-wave}$ $(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.30 to -0.10 OUR ESTIMATE			
-0.20 ± 0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
-0.23	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.30	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.34	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}} \ln N\pi \rightarrow N(1680) \rightarrow N\rho, S=3/2, F\text{-wave}$ $(\Gamma_1 \Gamma_{12})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.18 to -0.10 OUR ESTIMATE			
-0.13 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
-0.15	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}} \ln N\pi \rightarrow N(1680) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$ $(\Gamma_1 \Gamma_{13})^{1/2} / \Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.26 to +0.35 OUR ESTIMATE			
+0.29 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
+0.31	1,7 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.30	2 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.42	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1680)$ PHOTON DECAY AMPLITUDES

$N(1680) \rightarrow p\gamma, \text{helicity-1/2 amplitude } A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.014 to 0.008 OUR ESTIMATE			
-0.006 ± 0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.017 ± 0.018	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.009 ± 0.006	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.028 ± 0.003	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.026 ± 0.003	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.018 ± 0.014	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.005 ± 0.015	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.009 ± 0.002	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

$N(1680) \rightarrow p\gamma, \text{helicity-3/2 amplitude } A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.135 ± 0.017 OUR ESTIMATE			
0.154 ± 0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.132 ± 0.010	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.115 ± 0.008	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.115 ± 0.003	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.122 ± 0.003	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.141 ± 0.014	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.138 ± 0.021	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
+0.121 ± 0.010	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

$N(1680) \rightarrow n\gamma, \text{helicity-1/2 amplitude } A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.027 ± 0.010 OUR ESTIMATE			
0.022 ± 0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.017 ± 0.014	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.032 ± 0.003	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
0.026 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.028 ± 0.014	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.044 ± 0.012	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
0.025 ± 0.010	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.037 ± 0.010	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

$N(1680) \rightarrow n\gamma, \text{helicity-3/2 amplitude } A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.036 ± 0.011 OUR ESTIMATE			
-0.048 ± 0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.033 ± 0.013	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.023 ± 0.005	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
-0.024 ± 0.009	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.029 ± 0.017	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.033 ± 0.015	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.035 ± 0.012	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.038 ± 0.018	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

$N(1680)$ FOOTNOTES

- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The parametrization used may be double counting.
- The range given is from 3 of 4 best solutions; not present in solution 1. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.
- LONGACRE 77 considers this coupling to be well determined.
- A Breit-Wigner fit to the HERNDON 75 IPWA.

$N(1680)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982). For very early references, see Reviews of Modern Physics **37** 633 (1965).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	+Arndt, Roper, Workman (KENT) IJP
Also	PR D30 904	+Manley, Arndt, Goradia, Teplitz (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
ARNDT 85	PR D32 1085	+Ford, Roper (VPI)
BELL 83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+ (RL) IJP
CRAWFORD 83	NP B211 1	+Morton (GLAS)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
FUJII 81	NP B187 53	+Hayashii, Iwata, Kajikawa+ (NAGO, OSAK)
ARAI 80	Toronto Conf. 93	(INUS)
Also	NP B194 251	Arai, Fujii (INUS)
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+ (RHEL, BRIS) IJP
TAKEDA 80	NP B168 17	+Arai, Fujii, Ikeda, Iwasaki+ (RHEL) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also	Toronto Conf. 3	Koch (KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+ (LBL, SLAC)
NOVOSELLER 78	NP B137 509	(CIT) IJP
Also	NP B137 445	Novoseller (CIT) IJP
BAKER 77	NP B126 365	+Blissett, Bloodworth, Broome, Hart+ (RHEL) IJP
LONGACRE 77	NP B122 493	+Doibeau (SACL) IJP
Also	NP B108 365	Doibeau, Triantis, Neveu, Cadiet (SACL) IJP
WINNIK 77	NP B128 66	+Toaff, Revel, Goldberg, Berny (HAIF)
FELLER 76	NP B104 219	+Fukushima, Horikawa, Kajikawa+ (NAGO, OSAK) IJP
DEANS 75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
HERNDON 75	PR D11 3183	+Longacre, Miller, Rosenfeld+ (LBL, SLAC)
KNASEL 75	PR D11 1	+Lindquist, Nelson+ (CHIC, WUUSL, OSU, ANL) IJP
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+ (LBL, SLAC) IJP
DEVENISH 74B	NP B81 330	+Froggatt, Martin (DESY, NORD, LOUC)
CARRERAS 70	NP B16 35	+Donnachie (DARE, MCHS)
BOTKE 69	PR 180 1417	(UCSB)
DEANS 69	PR 185 1797	+Wootton (SFLA)
HEUSCH 66	PRL 17 1019	+Prescott, Dashen (CIT)

$N(1700) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

The various partial-wave analyses do not agree very well.

$N(1700)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1650 to 1750 (≈ 1700) OUR ESTIMATE			
1737 ± 44	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1675 ± 25	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1731 ± 15	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1709	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1650	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1690 to 1710	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1719	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1670 ± 10	1 BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
1690	1 BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1660	2 LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1710	3 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

Baryon Full Listings

N(1700)

N(1700) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50 to 150 (≈ 100) OUR ESTIMATE			
250 ± 220	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
90 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
110 ± 30	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
166	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
70	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
70 to 100	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
126	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
90 ± 25	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
100	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
600	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
300	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1700) POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1700	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1660 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1670	ARNDT 85	DPWA	See ARNDT 91
1710 or 1678	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1616 or 1613	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
90 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
80	ARNDT 85	DPWA	See ARNDT 91
607 or 567	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
577 or 575	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

N(1700) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
6 ± 3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
0 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1700) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	5-15 %
Γ_2 $N\eta$	
Γ_3 ΛK	<3 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	85-95 %
Γ_6 $\Delta\pi$	
Γ_7 $\Delta(1232)\pi$, S-wave	
Γ_8 $\Delta(1232)\pi$, D-wave	
Γ_9 $N\rho$	<35 %
Γ_{10} $N\rho$, S=1/2, D-wave	
Γ_{11} $N\rho$, S=3/2, S-wave	
Γ_{12} $N\rho$, S=3/2, D-wave	
Γ_{13} $N(\pi\pi)_{S=0}^{I=0}$	
Γ_{14} $p\gamma$	~ 0.01 %
Γ_{15} $p\gamma$, helicity=1/2	
Γ_{16} $p\gamma$, helicity=3/2	
Γ_{17} $n\gamma$	
Γ_{18} $n\gamma$, helicity=1/2	
Γ_{19} $n\gamma$, helicity=3/2	

N(1700) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_i/Γ
0.06 to 0.15 OUR ESTIMATE				
0.01 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.11 ± 0.05	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.08 ± 0.03	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1700) \rightarrow \Lambda K$ $(\Gamma_1\Gamma_3)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.06 to +0.04 OUR ESTIMATE			
-0.012	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
-0.012	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.04	⁶ BAKER 78	DPWA	See SAXON 80
-0.03 ± 0.004	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
-0.03	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
+0.026 ± 0.019	DEVENISH 74B		Fixed-t dispersion rel.

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1700) \rightarrow \Sigma K$ $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
not seen	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$
<0.017	⁷ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1700) \rightarrow \Delta(1232)\pi$, S-wave $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.00 to ±0.08 OUR ESTIMATE			
+0.02 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.00	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.16	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1700) \rightarrow \Delta(1232)\pi$, D-wave $(\Gamma_1\Gamma_8)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
±0.04 to ±0.20 OUR ESTIMATE			
+0.10 ± 0.09	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.12	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.14	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1700) \rightarrow N\rho$, S=3/2, S-wave $(\Gamma_1\Gamma_{11})^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
±0.01 to ±0.13 OUR ESTIMATE			
-0.04 ± 0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.07	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.07	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow N(1700) \rightarrow N(\pi\pi)_{S=0}^{I=0}$ $(\Gamma_1\Gamma_{13})^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
±0.02 to ±0.28 OUR ESTIMATE			
+0.02 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.00	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.2	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1700) PHOTON DECAY AMPLITUDES

N(1700) → pγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.017 ± 0.012 OUR ESTIMATE			
-0.016 ± 0.014	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.002 ± 0.013	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.028 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.029 ± 0.006	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.024 ± 0.019	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.033 ± 0.021	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.014 ± 0.025	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1700) → pγ, helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.002 ± 0.020 OUR ESTIMATE			
-0.009 ± 0.012	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.029 ± 0.014	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.002 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.014 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.017 ± 0.014	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.014 ± 0.025	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
0.0 ± 0.014	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1700) → nγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
0.001 ± 0.043 OUR ESTIMATE			
0.006 ± 0.024	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.002 ± 0.013	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
-0.052 ± 0.030	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.055 ± 0.030	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.052 ± 0.035	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
+0.050 ± 0.042	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

See key on page 1343

Baryon Full Listings

$N(1700)$, $N(1710)$

$N(1700) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.003 ± 0.038 OUR ESTIMATE			
-0.033 ± 0.017	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.018 ± 0.018	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
-0.037 ± 0.036	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.035 ± 0.024	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.041 ± 0.030	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.035 ± 0.030	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

$N(1700) \quad \gamma p \rightarrow \Lambda K^+$ AMPLITUDES

$(\Gamma_1 \Gamma_1)^{1/2} / \Gamma_{\text{total}} \ln p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ (E_2 -amplitude)

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
4.09	TANABE 89	DPWA

$(\Gamma_1 \Gamma_1)^{1/2} / \Gamma_{\text{total}} \ln p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ (M_2 -amplitude)

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
-7.09	TANABE 89	DPWA

$p\gamma \rightarrow N(1700) \rightarrow \Lambda K^+$ phase angle θ (E_2 -amplitude)

VALUE (degrees)	DOCUMENT ID	TECN
• • • We do not use the following data for averages, fits, limits, etc. • • •		
-35.9	TANABE 89	DPWA

$N(1700)$ FOOTNOTES

- The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The overall phase of BAKER 78 couplings has been changed to agree with previous conventions.
- The range given is from the four best solutions.

$N(1700)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
MANLEY 92	PR D45 4002	(KEWT) IJP
Also	PR D30 904	(VPI)
ARNDT 91	PR D43 2131	(VPI, TELE) IJP
TANABE 89	PR C39 741	(MANZ)
Also	NC 102A 193	(MANZ)
ARNDT 85	PR D32 1085	(VPI)
BELL 83	NP B222 389	(RL) IJP
CRAWFORD 83	NP B211 1	(GLAS)
PDG 82	PL 111B	(HEL, CIT, CERN)
AWAJI 81	Bonn Conf. 352	(NAGO)
Also	NP B197 365	(NAGO)
FUJII 81	NP B187 53	(NAGO, OSAK)
ARAI 80	Toronto Conf. 93	(INUS)
Also	NP B194 251	(INUS)
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	(CMU, LBL) IJP
Also	PR D20 2839	(CMU, LBL) IJP
LIVANOS 80	Toronto Conf. 35	(SACL) IJP
SAXON 80	NP B182 522	(RHEL, BRIS) IJP
HOEHLER 79	PDAT 12-1	(KARLT) IJP
Also	Toronto Conf. 3	(KARLT) IJP
BAKER 78	NP B141 29	(RL, CAVE) IJP
BARBOUR 78	NP B141 253	(GLAS)
LONGACRE 77	PR D17 1795	(LBL, SLAC)
BAKER 77	NP B126 365	(RHEL) IJP
LONGACRE 77	NP B122 493	(SACL) IJP
Also	NP B108 365	(SACL) IJP
FELLER 76	NP B104 219	(NAGO, OSAK) IJP
DEANS 75	NP B96 90	(SFLA, ALAH) IJP
LONGACRE 75	PL 55B 415	(LBL, SLAC) IJP
DEVENISH 74B	NP B81 330	(DESY, NORD, LOUC)

$N(1710) P_{11}$

$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: ***

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

The various partial-wave analyses do not agree very well.

$N(1710)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1680 to 1740 (≈ 1710) OUR ESTIMATE			
1717 ± 28	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
1700 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1723 ± 9	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1706	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
1692	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1730	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1690	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
1650 to 1680	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1721	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1625 ± 10	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
1650	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1720	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1670	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1710	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1710)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50 to 250 (≈ 100) OUR ESTIMATE			
480 ± 230	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
93 ± 30	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
90 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
120 ± 15	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
540	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
200	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
550	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
97	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
90 to 150	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
167	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
160 ± 6	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
95	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
120	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
174	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
75	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1710)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1690	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1636	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1698	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
1690 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1708 or 1712	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1720 or 1711	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
544	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
88	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
80 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
17 or 22	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
123 or 115	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$N(1710)$ ELASTIC POLE RESIDUE

MODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
15	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
149	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
9	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
8 ± 2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
149	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-167	CUTKOSKY 90	IPWA	$\pi N \rightarrow \pi N$
175 ± 35	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

Baryon Full Listings

N(1710)

N(1710) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10–20 %
Γ_2 $N\eta$	
Γ_3 ΛK	5–25 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	40–90 %
Γ_6 $\Delta\pi$	15–40 %
Γ_7 $\Delta(1232)\pi$, P-wave	
Γ_8 $N\rho$	5–25 %
Γ_9 $N\rho$, $S=1/2$, P-wave	
Γ_{10} $N\rho$, $S=3/2$, P-wave	
Γ_{11} $N(\pi\pi)_{S=0}^{I=0}$	10–40 %
Γ_{12} $p\gamma$, helicity=1/2	
Γ_{13} $n\gamma$, helicity=1/2	

N(1710) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.10 to 0.20 OUR ESTIMATE				
0.09±0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.20±0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.12±0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.12 to 0.20 OUR ESTIMATE				
0.22	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	
+0.383	FELTESSE 75	DPWA	Soln A; see BAKER 79	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
+0.12 to +0.18 OUR ESTIMATE				
+0.16	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
+0.14	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
0.10	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

$(\Gamma_1\Gamma_4)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
0.075 to 0.203 OUR ESTIMATE				
0.075	DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow \Delta(1232)\pi$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
±0.16 to ±0.22 OUR ESTIMATE				
0.21±0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.17	LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.20	LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_9)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N\rho$, $S=1/2$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
±0.09 to ±0.19 OUR ESTIMATE				
+0.05±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
+0.19	LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.20	LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_{10})^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N\rho$, $S=3/2$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{10})^{1/2}/\Gamma$
±0.14 to ±0.22 OUR ESTIMATE				
+0.31	LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_{11})^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1710) \rightarrow N(\pi\pi)_{S=0}^{I=0}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{11})^{1/2}/\Gamma$
±0.14 to ±0.22 OUR ESTIMATE				
+0.04±0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
-0.26	LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
-0.28	LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

N(1710) PHOTON DECAY AMPLITUDES

N(1710) $\rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.006±0.027 OUR ESTIMATE			
-0.037±0.002	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.006±0.018	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.028±0.009	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.009±0.006	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.012±0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.015±0.025	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
+0.001±0.039	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
+0.053±0.019	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

N(1710) $\rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.016±0.029 OUR ESTIMATE			
0.052±0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.000±0.018	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.001±0.003	FUJII 81	DPWA	$\gamma N \rightarrow \pi N$
0.005±0.013	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.011±0.021	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.017±0.020	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.028±0.045	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1710) $\gamma p \rightarrow \Lambda K^+$ AMPLITUDES

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1710) \rightarrow \Lambda K^+$	DOCUMENT ID	TECN	COMMENT	$(M_{1-} \text{ amplitude})$
-10.6 ± 0.4				
-10.6	WORKMAN 90	DPWA		
-7.21	TANABE 89	DPWA		

$p\gamma \rightarrow N(1710) \rightarrow \Lambda K^+$ phase angle θ	DOCUMENT ID	TECN	COMMENT	$(M_{1-} \text{ amplitude})$
215 ± 3				
215	WORKMAN 90	DPWA		
176.3	TANABE 89	DPWA		

N(1710) FOOTNOTES

- The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The overall phase of BAKER 78 couplings has been changed to agree with previous conventions.
- The range given for DEANS 75 is from the four best solutions.

N(1710) REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	+Arndt, Roper, Workman (KENT) IJP
Also 84	PR D30 904	+Saleski (VPI)
ARNDT 91	PR D43 2131	+Manley, Arndt, Goradia, Teplitz (VPI, TELE) IJP
CUTKOSKY 90	PR D42 235	+Li, Roper, Workman, Ford (CMU)
WORKMAN 90	PR C42 781	(VPI)
TANABE 89	PR C39 741	+Wang (MANZ)
Also 89	NC 102A 193	(MANZ)
BELL 83	NP B222 389	+Kohno, Bennhold (RL) IJP
CRAWFORD 83	NP B211 1	+Kohno, Tanabe, Bennhold (GLAS)
PDG 82	PL 111B	+Blissett, Broome, Daley, Hart, Lintern+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Roos, Porter, Aguilar-Benitez+ (NAGO)
Also 82	NP B197 365	+Kajikawa (NAGO)
FUJII 81	NP B187 53	+Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
ARAI 80	Toronto Conf. 93	+Hayashii, Iwata, Kajikawa+ (NAGO, OSAK)
Also 82	NP B194 251	(INUS)
CRAWFORD 80	Toronto Conf. 107	Arai, Fujii (INUS)
		(GLAS)

See key on page 1343

Baryon Full Listings
N(1710), N(1720)

CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu	(SACL) IJP
SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
BAKER	79	NP B156 93	+Brown, Clark, Davies, Desagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
BAKER	78	NP B141 29	+Blissett, Bloodworth, Broome+	(RL, CAVE) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons	(GLAS) IJP
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smdja+	(LBL, SLAC) IJP
BAKER	77	NP B126 365	+Blissett, Bloodworth, Broome, Hart+	(RHEL) IJP
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	+Fukushima, Triantis, Neveu, Cadiet	(SACL) IJP
FELLER	76	NP B104 219	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
FELTESSE	75	NP B93 242	+Ayed, Bareyre, Borgeaud, David+	(SACL) IJP
KNASEL	75	PR D11 1	+Lindquist, Nelson+	(CHIC, WUSL, OSU, ANL) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smdja+	(LBL, SLAC) IJP

N(1720) ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
15	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
11	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
8±2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-130	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-160±30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1720) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-20 %
Γ_2 $N\eta$	
Γ_3 ΛK	1-15 %
Γ_4 ΣK	
Γ_5 $N\pi\pi$	>70 %
Γ_6 $\Delta\pi$	
Γ_7 $\Delta(1232)\pi, P\text{-wave}$	
Γ_8 $N\rho$	70-85 %
Γ_9 $N\rho, S=1/2, P\text{-wave}$	
Γ_{10} $N\rho, S=3/2, P\text{-wave}$	
Γ_{11} $N(\pi\pi)_{S=0}^{\rho\text{-wave}}$	
Γ_{12} $p\gamma$	0.01-0.06 %
Γ_{13} $p\gamma, \text{helicity}=1/2$	
Γ_{14} $p\gamma, \text{helicity}=3/2$	
Γ_{15} $n\gamma$	
Γ_{16} $n\gamma, \text{helicity}=1/2$	
Γ_{17} $n\gamma, \text{helicity}=3/2$	

N(1720) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
0.10 to 0.20 OUR ESTIMATE			
0.13±0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
0.10±0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
0.14±0.03	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1720) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT
-0.08	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1720) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT
-0.14 to -0.06 OUR ESTIMATE			
-0.09	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
-0.11	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1720) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT
-0.09	⁶ BAKER 78	DPWA	See SAXON 80
-0.06±0.02	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
-0.09	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1720) \rightarrow \Delta(1232)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT
0.051 to 0.087	⁷ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1720) \rightarrow N\rho, S=1/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT
-0.17	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(1720) \rightarrow N\rho, S=3/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT
+0.34±0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.26	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.40	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1720) P₁₃

$I(J^P) = \frac{1}{2}(\frac{3}{2}^+)$ Status: * * * *

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

N(1720) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1650 to 1750 (≈ 1720) OUR ESTIMATE			
1717±31	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
1700±50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1710±20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1720	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1785	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1690	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1710 to 1790	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1809	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1640±10	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
1710	² BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1750	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1850	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1720	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1720) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100 to 200 (≈ 150) OUR ESTIMATE			
380±180	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
125±70	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
190±30	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
200	LI 93	IPWA	$\gamma N \rightarrow \pi N$
308	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
120	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
447	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
300 to 400	BAKER 78	DPWA	$\pi^- p \rightarrow \Lambda K^0$
285	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
200±50	¹ BAKER 77	IPWA	$\pi^- p \rightarrow \Lambda K^0$
500	¹ BAKER 77	DPWA	$\pi^- p \rightarrow \Lambda K^0$
130	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
327	KNASEL 75	DPWA	$\pi^- p \rightarrow \Lambda K^0$
150	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

N(1720) POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
1686	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1675	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1680±30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1705	ARNDT 85	DPWA	See ARNDT 91
1716 or 1716	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1745 or 1748	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-2xIMAGINARY PART			
187	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
114	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
120±40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
80	ARNDT 85	DPWA	See ARNDT 91
124 or 126	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
135 or 123	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

Baryon Full Listings

 $N(1720)$, $N(1900)$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1720) \rightarrow N\rho, S=3/2, P\text{-wave}$	$(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
$+0.15$	2	LONGACRE	77 IPWA $\pi N \rightarrow N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1720) \rightarrow N(\pi\pi)_{S\text{-wave}}^{I=0}$	$(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.19	2	LONGACRE	77 IPWA $\pi N \rightarrow N\pi\pi$

 $N(1720)$ PHOTON DECAY AMPLITUDES $N(1720) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
$+0.027 \pm 0.024$	OUR ESTIMATE		
0.012 ± 0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.044 ± 0.066	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
-0.004 ± 0.007	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.051 ± 0.009	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.071 ± 0.010	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.038 ± 0.050	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$+0.111 \pm 0.047$	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

 $N(1720) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.026 ± 0.010	OUR ESTIMATE		
-0.022 ± 0.003	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.024 ± 0.006	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
-0.040 ± 0.016	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.058 ± 0.010	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.011 ± 0.011	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.014 ± 0.040	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.063 ± 0.032	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

 $N(1720) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
$+0.018 \pm 0.029$	OUR ESTIMATE		
0.050 ± 0.004	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.002 ± 0.005	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.019 ± 0.033	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.001 ± 0.038	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
-0.003 ± 0.034	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$+0.007 \pm 0.020$	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

 $N(1720) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.033 ± 0.089	OUR ESTIMATE		
-0.017 ± 0.004	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.015 ± 0.019	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.139 ± 0.039	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
-0.134 ± 0.044	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.018 ± 0.028	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
$+0.051 \pm 0.051$	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$

 $N(1720) \gamma p \rightarrow \Lambda K^+$ AMPLITUDES

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1720) \rightarrow \Lambda K^+$	$(E_{1+}$ amplitude)		
VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
10.2 ± 0.2	WORKMAN	90	DPWA
• • • We do not use the following data for averages, fits, limits, etc. • • •			
9.52	TANABE	89	DPWA

$p\gamma \rightarrow N(1720) \rightarrow \Lambda K^+$ phase angle θ	$(E_{1+}$ amplitude)		
VALUE (degrees)	DOCUMENT ID	TECN	COMMENT
-124 ± 2	WORKMAN	90	DPWA
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-103.4	TANABE	89	DPWA

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $p\gamma \rightarrow N(1720) \rightarrow \Lambda K^+$	$(M_{1+}$ amplitude)		
VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
-4.5 ± 0.2	WORKMAN	90	DPWA
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.18	TANABE	89	DPWA

 $N(1720)$ FOOTNOTES

- The two BAKER 77 entries are from an IPWA using the Barrelet-zero method and from a conventional energy-dependent analysis.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The overall phase of BAKER 78 couplings has been changed to agree with previous conventions.
- The range given is from the four best solutions. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.

 $N(1720)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER	94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER	93	πN Newsletter 9 1	(KARL)
LI	93	PR C47 2759	(VPI)
MANLEY	92	PR D45 4002	+Saleski (KENT) IJP
Also	84	PR D30 904	(VPI)
ARNDT	87	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
WORKMAN	90	PR C42 781	(VPI)
TANABE	89	PR C39 741	+Kohno, Bennhold (MANZ)
Also	89	NC 102A 193	Kohno, Tanabe, Bennhold (MANZ)
ARNDT	85	PR D32 1085	+Ford, Roper (VPI)
BELL	83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+ (RL) IJP
CRAWFORD	83	NP B211 1	+Morton (GLAS)
PDG	82	PL 111B	Rops, Porter, Aguilar-Benitez+ (HELS. CIT. CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa (NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
ARAI	80	Toronto Conf. 93	(INUS)
Also	82	NP B194 251	Arai, Fujii (INUS)
CRAWFORD	80	Toronto Conf. 107	(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+ (RHEL, BRIS) IJP
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also	80	Toronto Conf. 3	Koch (KARLT) IJP
BAKER	78	NP B141 29	+Blissett, Bloodworth, Broome+ (RL, CAVE) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons (GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+ (LBL, SLAC)
BAKER	77	NP B126 365	+Blissett, Bloodworth, Broome, Hart+ (RHEL) IJP
LONGACRE	77	NP B122 493	+Dolbeau (SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet (SACL) IJP
WINNIK	77	NP B128 66	+Toaff, Revel, Goldberg, Berry (HAIF) I
DEANS	75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
KNASEL	75	PR D11 1	+Lindquist, Nelson+ (CHIC, WUSL, OSU, ANL) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+ (LBL, SLAC) IJP

 $N(1900) P_{13}$

$$I(J^P) = \frac{1}{2}(3^+)$$
 Status: *

OMITTED FROM SUMMARY TABLE

 $N(1900)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1879 ± 17	OUR ESTIMATE		
1879 ± 17	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

 $N(1900)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
498 ± 78	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

 $N(1900)$ DECAY MODES

Mode
$\Gamma_1 N\pi$
$\Gamma_2 N\pi\pi$
$\Gamma_3 N\rho, S = 1/2, P\text{-wave}$

 $N(1900)$ BRANCHING RATIOS

$\Gamma(N\pi) / \Gamma_{\text{total}}$	Γ_1 / Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
0.26 ± 0.06	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

$(\Gamma_1 \Gamma_f)^{1/2} / \Gamma_{\text{total}}$ in $N\pi \rightarrow N(1900) \rightarrow N\rho, S = 1/2, P\text{-wave}$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.34 ± 0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$

 $N(1900)$ REFERENCES

MANLEY	92	PR D45 4002	+Saleski (KENT)
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz (VPI)

See key on page 1343

Baryon Full Listings
N(1990)**N(1990) F₁₇**

$$I(J^P) = \frac{1}{2}(\frac{7}{2}^+) \text{ Status: } * *$$

OMITTED FROM SUMMARY TABLE

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

The various analyses do not agree very well with one another.

N(1990) MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1990 OUR ESTIMATE			
2086 ± 28	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
2018	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1970 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2005 ± 150	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
1999	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1990) WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
535 ± 120	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
295	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
350 ± 120	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
350 ± 100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
216	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1990) POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

-2×IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
260 ± 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

N(1990) ELASTIC POLE RESIDUE**MODULUS |r|**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9 ± 3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-60 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(1990) DECAY MODES

Mode	DOCUMENT ID	TECN	COMMENT
Γ_1 $N\pi$			
Γ_2 $N\eta$			
Γ_3 ΛK			
Γ_4 ΣK			
Γ_5 $N\pi\pi$			
Γ_6 $p\gamma$, helicity=1/2			
Γ_7 $p\gamma$, helicity=3/2			
Γ_8 $n\gamma$, helicity=1/2			
Γ_9 $n\gamma$, helicity=3/2			

N(1990) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.06 ± 0.02	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.04 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1990) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.043	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1990) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
+0.01	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
not seen	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
-0.021 ± 0.033	DEVENISH 74b		Fixed- t dispersion rel.	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1990) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
0.010 to 0.023	¹ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	
0.06	LANGBEIN 73	IPWA	$\pi N \rightarrow \Sigma K$ (sol. 1)	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(1990) \rightarrow N\pi\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
not seen	LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	

N(1990) PHOTON DECAY AMPLITUDES**N(1990) → p γ , helicity-1/2 amplitude A_{1/2}**

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
0.030 ± 0.029	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.001 ± 0.040	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.040	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1990) → p γ , helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
0.086 ± 0.060	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.004 ± 0.025	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.004	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1990) → n γ , helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.001	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.078 ± 0.030	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.069	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1990) → n γ , helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.178	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.116 ± 0.045	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.072	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(1990) FOOTNOTES

¹ The range given for DEANS 75 is from the four best solutions.

N(1990) REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	PR D30 904	Manley, Arndt, Goradia, Tepitz	(VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TEL) IJP
BELL 83	NP B222 389	+Blissett, Broome, Daley, Lintern+	(RL) IJP
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
CRAWFORD 80	Toronto Conf. 107		(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	Toronto Conf. 3	Koch	(KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons	(GLAS)
DEANS 75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP
DEVENISH 74b	NP B81 330	+Froggatt, Martin	(DESY, NÖRD, LOUC)
LANGBEIN 73	NP B53 251	+Wagner	(MUNI) IJP

Baryon Full Listings

 $N(2000)$, $N(2080)$ $N(2000) F_{15}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

Older results have been retained simply because there is little information at all about this possible state.

 $N(2000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2000 OUR ESTIMATE			
1903 ± 87	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
1882 ± 10	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2025	AYED 76	IPWA	$\pi N \rightarrow \pi N$
1970	¹ LANGBEIN 73	IPWA	$\pi N \rightarrow \Sigma K$ (sol. 2)
2175	ALMEHED 72	IPWA	$\pi N \rightarrow \pi N$
1930	DEANS 72	MPWA	$\gamma p \rightarrow \Lambda K$ (sol. D)

 $N(2000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
490 ± 310	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
95 ± 20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
157	AYED 76	IPWA	$\pi N \rightarrow \pi N$
170	¹ LANGBEIN 73	IPWA	$\pi N \rightarrow \Sigma K$ (sol. 2)
150	ALMEHED 72	IPWA	$\pi N \rightarrow \pi N$
112	DEANS 72	MPWA	$\gamma p \rightarrow \Lambda K$ (sol. D)

 $N(2000)$ DECAY MODES

Mode	
Γ_1	$N\pi$
Γ_2	$N\eta$
Γ_3	ΛK
Γ_4	ΣK
Γ_5	$N\pi\pi$
Γ_6	$\Delta(1232)\pi$, P-wave
Γ_7	$N\rho$, S=3/2, P-wave
Γ_8	$N\rho$, S=3/2, F-wave
Γ_9	$p\gamma$

 $N(2000)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.08 ± 0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$	
0.04 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.08	AYED 76	IPWA	$\pi N \rightarrow \pi N$	
0.25	ALMEHED 72	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(2000) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.03	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(2000) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
not seen	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

$(\Gamma_1\Gamma_4)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(2000) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
0.022	² DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	
0.05	¹ LANGBEIN 73	IPWA	$\pi N \rightarrow \Sigma K$ (sol. 2)	

$(\Gamma_1\Gamma_6)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(2000) \rightarrow \Delta(1232)\pi$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
0.10 ± 0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(2000) \rightarrow N\rho$, S=3/2, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
-0.22 ± 0.08	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$	

$(\Gamma_1\Gamma_8)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow N(2000) \rightarrow N\rho$, S=3/2, F-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
0.11 ± 0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$	

$(\Gamma_9\Gamma_3)^{1/2}/\Gamma_{\text{total}} \ln p\gamma \rightarrow N(2000) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_9\Gamma_3)^{1/2}/\Gamma$
0.0022	DEANS 72	MPWA	$\gamma p \rightarrow \Lambda K$ (sol. D)	

 $N(2000)$ FOOTNOTES¹ Not seen in solution 1 of LANGBEIN 73.² Value given is from solution 1 of DEANS 75; not present in solutions 2, 3, or 4. $N(2000)$ REFERENCES

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	84 PR D30 904	Manley, Arndt, Goradia, Tepitz	(VPI)
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80 Toronto Conf. 3	Koch	(KARLT) IJP
AYED 76	CEA-N-1921 Thesis		(SACL) IJP
DEANS 75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
LANGBEIN 73	NP B53 251	+Wagner	(MUNI) IJP
ALMEHED 72	NP B40 157	+Lovellace	(LUND, RUTG) IJP
DEANS 72	PR D6 1906	+Jacobs, Lyons, Montgomery	(SFLA) IJP

 $N(2080) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

There is some evidence for two resonances in this wave between 1800 and 2200 MeV (see CUTKOSKY 80). However, the solution of HOEHLER 79 is quite different.

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982). $N(2080)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2080 OUR ESTIMATE			
1804 ± 55	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
1920	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
1880 ± 100	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2060 ± 80	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1900	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
2081 ± 20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1880	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$

 $N(2080)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
450 ± 185	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \text{ \& } N\pi\pi$
320	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
180 ± 60	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (lower m)
300 ± 100	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (higher m)
240	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
265 ± 40	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
87	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$

 $N(2080)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1880 ± 100	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (lower m)
2050 ± 70	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (higher m)
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
160 ± 80	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (lower m)
200 ± 80	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (higher m)
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

 $N(2080)$ ELASTIC POLE RESIDUEMODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10 ± 5	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (lower m)
30 ± 20	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (higher m)

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
100 ± 80	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (lower m)
0 ± 100	¹ CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (higher m)

See key on page 1343

Baryon Full Listings
 $N(2080)$, $N(2090)$ $N(2080)$ DECAY MODES

Mode	
Γ_1	$N\pi$
Γ_2	$N\eta$
Γ_3	ΛK
Γ_4	ΣK
Γ_5	$N\pi\pi$
Γ_6	$\Delta(1232)\pi$, S-wave
Γ_7	$\Delta(1232)\pi$, D-wave
Γ_8	$N\rho$, $S=3/2$, S-wave
Γ_9	$N(\pi\pi)_{S=0}^{I=0}$
Γ_{10}	$p\gamma$, helicity=1/2
Γ_{11}	$p\gamma$, helicity=3/2
Γ_{12}	$n\gamma$, helicity=1/2
Γ_{13}	$n\gamma$, helicity=3/2
Γ_{14}	$p\gamma$

 $N(2080)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.23±0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.10±0.04	1 CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (lower m)	
0.14±0.07	1 CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$ (higher m)	
0.06±0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.065	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
+0.04	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
+0.03	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
0.014 to 0.037	2 DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow \Delta(1232)\pi$, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
-0.09±0.09	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow \Delta(1232)\pi$, D-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
+0.22±0.07	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow N\rho$, $S=3/2$, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
-0.24±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2080) \rightarrow N(\pi\pi)_{S=0}^{I=0}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
+0.25±0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(2080) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_{14}\Gamma_2)^{1/2}/\Gamma$
0.0037	HICKS 73	MPWA	$\gamma p \rightarrow p\eta$	

 $N(2080)$ PHOTON DECAY AMPLITUDES $N(2080) \rightarrow p\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.020±0.008	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.026±0.052	DEVENISH 74	DPWA	$\gamma N \rightarrow \pi N$

 $N(2080) \rightarrow p\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
0.017±0.011	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.128±0.057	DEVENISH 74	DPWA	$\gamma N \rightarrow \pi N$

 $N(2080) \rightarrow n\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
0.007±0.013	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.053±0.083	DEVENISH 74	DPWA	$\gamma N \rightarrow \pi N$

 $N(2080) \rightarrow n\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.053±0.034	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
0.100±0.141	DEVENISH 74	DPWA	$\gamma N \rightarrow \pi N$

 $N(2080) \gamma p \rightarrow \Lambda K^+$ AMPLITUDES $(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(2080) \rightarrow \Lambda K^+$ (E_2 -amplitude)

VALUE (units 10^{-3})	DOCUMENT ID	TECN
5.5 ± 0.3	WORKMAN 90	DPWA
••• We do not use the following data for averages, fits, limits, etc. •••		
4.09	TANABE 89	DPWA

 $p\gamma \rightarrow N(2080) \rightarrow \Lambda K^+$ phase angle θ (E_2 -amplitude)

VALUE (degrees)	DOCUMENT ID	TECN
-48 ± 5	WORKMAN 90	DPWA
••• We do not use the following data for averages, fits, limits, etc. •••		
-35.9	TANABE 89	DPWA

 $(\Gamma_1\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(2080) \rightarrow \Lambda K^+$ (M_2 -amplitude)

VALUE (units 10^{-3})	DOCUMENT ID	TECN
-6.7 ± 0.2	WORKMAN 90	DPWA
••• We do not use the following data for averages, fits, limits, etc. •••		
-4.09	TANABE 89	DPWA

 $N(2080)$ FOOTNOTES

¹CUTKOSKY 80 finds a lower mass D_{13} resonance, as well as one in this region. Both are listed here.

²The range given for DEANS 75 is from the four best solutions. Disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.

 $N(2080)$ REFERENCES

For early references, see Physics Letters 111B 70 (1982).

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	84 PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI, TELE) IJP
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
WORKMAN 90	PR C42 781		(VPI)
TANABE 89	PR C39 741		(MANZ)
Also	89 NC 102A 193	+Kohno, Bennhold	(MANZ)
BELL 83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+	(RL) IJP
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82 NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79 PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80 Toronto Conf. 3	Koch	(KARLT) IJP
WINNIK 77	NP B128 66	+Toaff, Revel, Goldberg, Berny	(HAIF) I
DEANS 75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP
DEVENISH 74	PL 52B 227	+Lyth, Rankin	(DESY, LANZ, BONN) IJP
HICKS 73	PR D7 2614	+Deans, Jacobs, Lyons+	(CMU, ORNL, SFLA) IJP

 $N(2090) S_{11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

Any structure in the S_{11} wave above 1800 MeV is listed here. A few early results that are now obsolete have been omitted.

 $N(2090)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1928±59	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
2180±80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1880±20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

 $N(2090)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
414±157	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
350±100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
95±30	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

 $N(2090)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2150±70	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1937 or 1949	1 LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

Baryon Full Listings

 $N(2090)$, $N(2100)$, $N(2190)$ **-2xIMAGINARY PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
350 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
139 or 131	¹ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

 $N(2090)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
0 ± 90	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $N(2090)$ DECAY MODES

Mode
Γ_1 $N\pi$
Γ_2 ΛK
Γ_3 $N\pi\pi$

 $N(2090)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.10 ± 0.10	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.18 ± 0.08	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.09 ± 0.05	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2090) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
not seen	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

 $N(2090)$ FOOTNOTES

¹ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

 $N(2090)$ REFERENCES

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	84 PR D30 904	Manley, Arndt, Goradia, Tepitz	(VPI)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79 PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80 Toronto Conf. 3	Koch	(KARLT) IJP
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)

 $N(2100)$ P_{11}

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

 $N(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2100 OUR ESTIMATE			
1885 ± 30	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
2125 ± 75	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2050 ± 20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

 $N(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
113 ± 44	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
260 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
200 ± 30	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

 $N(2100)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
2120 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
240 ± 80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

 $N(2100)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
14 ± 7	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
35 ± 25	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $N(2100)$ DECAY MODES

Mode
Γ_1 $N\pi$
Γ_2 $N\pi\pi$
Γ_3 $\Delta(1232)\pi, P\text{-wave}$

 $N(2100)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.15 ± 0.06	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.12 ± 0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.10 ± 0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2100) \rightarrow \Delta(1232)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.19 ± 0.08	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

 $N(2100)$ REFERENCES

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	84 PR D30 904	Manley, Arndt, Goradia, Tepitz	(VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford	(VPI, TELE) IJP
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79 PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80 Toronto Conf. 3	Koch	(KARLT) IJP

 $N(2190)$ G_{17}

$$I(J^P) = \frac{1}{2}(\frac{7}{2}^-) \text{ Status: } * * * *$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $N(2190)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2100 to 2200 (≈ 2190) OUR ESTIMATE			
2127 ± 9	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
2200 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2140 ± 12	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2140 ± 40	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
2098	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
2180	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
2140	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
2117	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $N(2190)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
350 to 550 (≈ 450) OUR ESTIMATE			
550 ± 50	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
500 ± 150	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
390 ± 30	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
270 ± 50	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
238	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
80	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$
319	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$
220	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $N(2190)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
2042	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
2060	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
2100 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

See key on page 1343

Baryon Full Listings
N(2190)

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
482	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
464	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
400 ± 160	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(2190) ELASTIC POLE RESIDUE

MODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
45	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
54	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
25 ± 10	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-44	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-30 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

N(2190) DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-20 %
Γ_2 $N\eta$	
Γ_3 ΛK	
Γ_4 ΣK	
Γ_5 $N\pi\pi$	
Γ_6 $N\rho$	
Γ_7 $N\rho, S=3/2, D\text{-wave}$	
Γ_8 $p\gamma, \text{helicity}=1/2$	
Γ_9 $p\gamma, \text{helicity}=3/2$	
Γ_{10} $n\gamma, \text{helicity}=1/2$	
Γ_{11} $n\gamma, \text{helicity}=3/2$	

N(2190) BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.1 to 0.2 OUR ESTIMATE				
0.22 ± 0.01	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.12 ± 0.06	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.14 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.16 ± 0.04	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2190) \rightarrow N\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma$
0.052	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2190) \rightarrow \Lambda K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_3)^{1/2}/\Gamma$
-0.02	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$	
-0.02	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2190) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_4)^{1/2}/\Gamma$
0.014 to 0.019	² DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2190) \rightarrow N\rho, S=3/2, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_7)^{1/2}/\Gamma$
-0.25 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

N(2190) PHOTON DECAY AMPLITUDES

N(2190) → pγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.055	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.030	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(2190) → pγ, helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
0.081	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
+0.180	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(2190) → nγ, helicity-1/2 amplitude A_{1/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.042	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
-0.085	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(2190) → nγ, helicity-3/2 amplitude A_{3/2}

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.126	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
+0.007	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

N(2190) γp → ΛK⁺ AMPLITUDES $(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(2190) \rightarrow \Lambda K^+$ (E₄₋ amplitude)

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
2.5 ± 1.0	WORKMAN 90	DPWA	
2.04	TANABE 89	DPWA	

pγ → N(2190) → ΛK⁺ phase angle θ (E₄₋ amplitude)

VALUE (degrees)	DOCUMENT ID	TECN	COMMENT
-4 ± 9	WORKMAN 90	DPWA	
-27.5	TANABE 89	DPWA	

 $(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}}$ in $p\gamma \rightarrow N(2190) \rightarrow \Lambda K^+$ (M₄₋ amplitude)

VALUE (units 10 ⁻³)	DOCUMENT ID	TECN	COMMENT
-7.0 ± 0.7	WORKMAN 90	DPWA	
-5.78	TANABE 89	DPWA	

N(2190) FOOTNOTES

¹ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

² The range given for DEANS 75 is from the four best solutions. Disagrees with π⁺p → Σ⁺K⁺ data of WINNIK 77 around 1920 MeV.

N(2190) REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	π N Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	π N Newsletter 9 1	(KARL)
MANLEY 92	PR D45 4002	(KENT) IJP
Also 84	PR D30 904	(VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
WORKMAN 90	PR C42 781	(VPI)
TANABE 89	PR C39 741	+Kohno, Bennhold (MANZ)
Also 89	NC 102A 193	Kohno, Tanabe, Bennhold (MANZ)
BELL 83	NP S222 389	+Blissett, Broome, Daley, Hart, Lintern+ (RL) IJP
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELs, CIT, CERN)
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+ (RHEL, BRIS) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
HENDRY 78	PR L 41 222	(IND, LBL) IJP
Also 81	ANP 136 1	Hendry (IND)
WINNIK 77	NP B128 66	+Toaff, Revel, Goldberg, Berny (HAIF) I
DEANS 75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP

Baryon Full Listings

$N(2200)$, $N(2220)$

$N(2200) D_{15}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

The mass is not well determined. A few early results have been omitted.

$N(2200)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2200 OUR ESTIMATE			
1900	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
2180 \pm 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1920	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
2228 \pm 30	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

$N(2200)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
130	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$
400 \pm 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
220	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$
310 \pm 50	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

$N(2200)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2100 \pm 60	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

-2xIMAGINARY PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
360 \pm 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

$N(2200)$ ELASTIC POLE RESIDUE

MODULUS $ r $ VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
20 \pm 10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

PHASE θ VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-90 \pm 50	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

$N(2200)$ DECAY MODES

Mode	Γ_1	Γ_2	Γ_3
$N\pi$			
$N\eta$			
ΛK			

$N(2200)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.10 \pm 0.03	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.07 \pm 0.02	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2200) \rightarrow N\eta$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.066	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2200) \rightarrow \Lambda K$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
-0.03	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$	
-0.05	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$	

$N(2200)$ REFERENCES

BELL	83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+	(RL) IJP
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
SAXON	80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+	(RHEL, BRIS) IJP
BAKER	79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+	(RHEL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP

$N(2220) H_{19}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

$N(2220)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2220 OUR ESTIMATE			
2230 \pm 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
2205 \pm 10	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
2300 \pm 100	HENDRY	78	MPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2050	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$

$N(2220)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2220 OUR ESTIMATE			
500 \pm 150	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
365 \pm 30	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
450 \pm 150	HENDRY	78	MPWA $\pi N \rightarrow \pi N$

$N(2220)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2135	¹ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
2253	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
2160 \pm 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

-2xIMAGINARY PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
400	¹ HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
640	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
480 \pm 100	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

$N(2220)$ ELASTIC POLE RESIDUE

MODULUS $ r $ VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
85	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
45 \pm 20	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

PHASE θ VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-50	HOEHLER	93	ARGD $\pi N \rightarrow \pi N$
-62	ARNDT	91	DPWA $\pi N \rightarrow \pi N$ Soln SM90
-45 \pm 25	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

$N(2220)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-20 %
Γ_2 $N\eta$	
Γ_3 ΛK	

$N(2220)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.1 to 0.2 OUR ESTIMATE				
0.15 \pm 0.03	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.18 \pm 0.015	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	
0.12 \pm 0.04	HENDRY	78	MPWA $\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2220) \rightarrow N\eta$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.034	BAKER	79	DPWA $\pi^- p \rightarrow n\eta$	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow N(2220) \rightarrow \Lambda K$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
not required	BELL	83	DPWA $\pi^- p \rightarrow \Lambda K^0$	
not seen	SAXON	80	DPWA $\pi^- p \rightarrow \Lambda K^0$	

See key on page 1343

Baryon Full Listings

$N(2220)$, $N(2250)$, $N(2600)$

 $N(2220)$ FOOTNOTES

¹ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

 $N(2220)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
BELL 83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+ (RL) IJP
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+ (RHEL, BRIS) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
HENDRY 78	PRL 41 222	(IND, LBL) IJP
Also 81	ANP 136 1	Hendry (IND)

$N(2250)$ G_{19}

$$I(J^P) = \frac{1}{2}(\frac{9}{2}^-) \text{ Status: } ***$$

 $N(2250)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2170 to 2310 (≈ 2250) OUR ESTIMATE			
2250 \pm 80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2268 \pm 15	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2200 \pm 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

 $N(2250)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
290 to 470 (≈ 400) OUR ESTIMATE			
480 \pm 120	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
300 \pm 40	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
350 \pm 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

 $N(2250)$ POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2187	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
2243	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
2150 \pm 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

-2 \times IMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
388	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
650	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
360 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $N(2250)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
21	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
47	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
20 \pm 6	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-37	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-50 \pm 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $N(2250)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	5-15 %
Γ_2 $N\eta$	
Γ_3 ΛK	

 $N(2250)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 to 0.15 OUR ESTIMATE				
0.10 \pm 0.02	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.10 \pm 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.09 \pm 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N\pi \rightarrow N(2250) \rightarrow N\eta \quad (\Gamma_1 \Gamma_2)^{1/2} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.043	BAKER 79	DPWA	$\pi^- p \rightarrow n\eta$

$$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N\pi \rightarrow N(2250) \rightarrow \Lambda K \quad (\Gamma_1 \Gamma_2)^{1/2} / \Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.02	BELL 83	DPWA	$\pi^- p \rightarrow \Lambda K^0$
not seen	SAXON 80	DPWA	$\pi^- p \rightarrow \Lambda K^0$

 $N(2250)$ FOOTNOTES

¹ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

 $N(2250)$ REFERENCES

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
BELL 83	NP B222 389	+Blissett, Broome, Daley, Hart, Lintern+ (RL) IJP
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
SAXON 80	NP B162 522	+Baker, Bell, Blissett, Bloodworth+ (RHEL, BRIS) IJP
BAKER 79	NP B156 93	+Brown, Clark, Davies, Depagter, Evans+ (RHEL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
HENDRY 78	PRL 41 222	(IND, LBL) IJP
Also 81	ANP 136 1	Hendry (IND)

$N(2600)$ $I_{1,11}$

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^-) \text{ Status: } ***$$

 $N(2600)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2550 to 2750 (≈ 2600) OUR ESTIMATE			
2577 \pm 50	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2700 \pm 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

 $N(2600)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
500 to 800 (≈ 650) OUR ESTIMATE			
400 \pm 100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
900 \pm 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

 $N(2600)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	5-10 %

 $N(2600)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 to 0.1 OUR ESTIMATE				
0.05 \pm 0.01	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.08 \pm 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

 $N(2600)$ REFERENCES

HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also 81	ANP 136 1	Hendry	(IND)

Baryon Full Listings

$N(2700)$, $N(\sim 3000)$, $\Delta(1232)$

$N(2700) K_{1,13}$ $I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: * *

OMITTED FROM SUMMARY TABLE

$N(2700)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2700 OUR ESTIMATE			
2612 ± 45	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
3000 ± 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

$N(2700)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
350 ± 50	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
900 ± 150	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

$N(2700)$ DECAY MODES

Mode

$\Gamma_1 N\pi$

$N(2700)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.04 ± 0.01	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.07 ± 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$N(2700)$ REFERENCES

HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also 81	ANP 136 1	Hendry	(IND)

$N(\sim 3000)$ Region Partial-Wave Analyses

OMITTED FROM SUMMARY TABLE

We list here miscellaneous high-mass candidates for isospin-1/2 resonances found in partial-wave analyses.

Our 1982 edition had an $N(3245)$, an $N(3690)$, and an $N(3755)$, each a narrow peak seen in a production experiment. Since nothing has been heard from them since the 1960's, we declare them to be dead. There was also an $N(3030)$, deduced from total cross-section and 180° elastic cross-section measurements; it is the KOCH 80 $L_{1,15}$ state below.

$N(\sim 3000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 3000 OUR ESTIMATE			
2600	KOCH 80	IPWA	$\pi N \rightarrow \pi N D_{13}$
3100	KOCH 80	IPWA	$\pi N \rightarrow \pi N L_{1,15}$ wave
3500	KOCH 80	IPWA	$\pi N \rightarrow \pi N M_{1,17}$ wave
3500 to 4000	KOCH 80	IPWA	$\pi N \rightarrow \pi N N_{1,19}$ wave
3500 ± 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N L_{1,15}$ wave
3800 ± 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N M_{1,17}$ wave
4100 ± 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N N_{1,19}$ wave

$N(\sim 3000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1300 ± 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N L_{1,15}$ wave
1600 ± 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N M_{1,17}$ wave
1900 ± 300	HENDRY 78	MPWA	$\pi N \rightarrow \pi N N_{1,19}$ wave

$N(\sim 3000)$ DECAY MODES

Mode

$\Gamma_1 N\pi$

$N(\sim 3000)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.055 ± 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N L_{1,15}$ wave	
0.040 ± 0.015	HENDRY 78	MPWA	$\pi N \rightarrow \pi N M_{1,17}$ wave	
0.030 ± 0.015	HENDRY 78	MPWA	$\pi N \rightarrow \pi N N_{1,19}$ wave	

$N(\sim 3000)$ REFERENCES

KOCH 80	Toronto Conf. 3		(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also 81	ANP 136 1	Hendry	(IND) IJP

Δ BARYONS ($S = 0, I = 3/2$)

$$\Delta^{++} = uuu, \Delta^+ = uud, \Delta^0 = udd, \Delta^- = ddd$$

$\Delta(1232) P_{33}$

$I(J^P) = \frac{3}{2}(\frac{3}{2}^+)$ Status: * * * *

Most of the results published before 1977 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

$\Delta(1232)$ MASSES

MIXED CHARGES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1230 to 1234 (≈ 1232) OUR ESTIMATE			
1231 ± 1	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1232 ± 3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1233 ± 2	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

$\Delta(1232)^{++}$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1230.9 ± 0.3	KOCH 80B	IPWA	$\pi N \rightarrow \pi N$
1230.6 ± 0.2	ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
1231.1 ± 0.2	PEDRONI 78		$\pi N \rightarrow \pi N$ 70-370 MeV

$\Delta(1232)^+$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1234.9 ± 1.4	MIROSHNIC... 79		Fit photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1231.6	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1231.2	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1231.8	BERENDS 75	IPWA	$\gamma p \rightarrow \pi N$

$\Delta(1232)^0$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1233.6 ± 0.5	KOCH 80B	IPWA	$\pi N \rightarrow \pi N$
1232.5 ± 0.3	ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
1233.8 ± 0.2	PEDRONI 78		$\pi N \rightarrow \pi N$ 70-370 MeV

$m_{\Delta^0} - m_{\Delta^+}$

VALUE (MeV)	DOCUMENT ID	COMMENT
2.7 ± 0.3	¹ PEDRONI 78	See the masses

$\Delta(1232)$ WIDTHS

MIXED CHARGES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
115 to 125 (≈ 120) OUR ESTIMATE			
118 ± 4	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
120 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
116 ± 5	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

$\Delta(1232)^{++}$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
111.0 ± 1.0	KOCH 80B	IPWA	$\pi N \rightarrow \pi N$
113.2 ± 0.3	ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
111.3 ± 0.5	PEDRONI 78		$\pi N \rightarrow \pi N$ 70-370 MeV

See key on page 1343

Baryon Full Listings
 $\Delta(1232)$ $\Delta(1232)^+$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
131.1 ± 2.4	MIROSHNIC... 79		Fit photoproduction
• • • We do not use the following data for averages, fits, limits, etc. • • •			
111.2	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
111.0	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1232)^0$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
113.0 ± 1.5	KOCH 80B	IPWA	$\pi N \rightarrow \pi N$
121.3 ± 0.4	ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
117.9 ± 0.9	PEDRONI 78		$\pi N \rightarrow \pi N$ 70-370 MeV

 $\Delta^0 - \Delta^{++}$ WIDTH DIFFERENCE

VALUE (MeV)	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
6.6 ± 1.0	PEDRONI 78	See the widths

 $\Delta(1232)$ POLE POSITIONS

REAL PART, MIXED CHARGES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1209	2 HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
1210	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1210 ± 1	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1210	ARNDT 85	DPWA	See ARNDT 91

-IMAGINARY PART, MIXED CHARGES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100	2 HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
50	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
50 ± 1	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
50	ARNDT 85	DPWA	See ARNDT 91

REAL PART, $\Delta(1232)^{++}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1210.70 ± 0.16	3 ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
1209.6 ± 0.5	4 VASAN 76B		Fit to CARTER 73
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1210.4 ± 0.17	5 ZIDELL 78		
1210.5 to 1210.8	6 VASAN 76B		Fit to CARTER 73

-IMAGINARY PART, $\Delta(1232)^{++}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
49.61 ± 0.12	3 ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
50.4 ± 0.5	4 VASAN 76B		Fit to CARTER 73
• • • We do not use the following data for averages, fits, limits, etc. • • •			
49.745 ± 0.14	5 ZIDELL 78		
49.9 to 50.0	6 VASAN 76B		Fit to CARTER 73

REAL PART, $\Delta(1232)^+$

VALUE (MeV)	DOCUMENT ID	COMMENT
1206.9 ± 0.9 to 1210.5 ± 1.8	MIROSHNIC... 79	Fit photoproduction
1208.0 ± 2.0	CAMPBELL 76	Fit photoproduction

-IMAGINARY PART, $\Delta(1232)^+$

VALUE (MeV)	DOCUMENT ID	COMMENT
55.6 ± 1.0 to 58.3 ± 1.1	MIROSHNIC... 79	Fit photoproduction
53.0 ± 2.0	CAMPBELL 76	Fit photoproduction

REAL PART, $\Delta(1232)^0$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1210.30 ± 0.36	3 ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
1210.75 ± 0.6	4 VASAN 76B		Fit to CARTER 73
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1209.5 ± 0.41	5 ZIDELL 78		
1210.2	6 VASAN 76B		Fit to CARTER 73

-IMAGINARY PART, $\Delta(1232)^0$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
54.0 ± 0.26	3 ZIDELL 80	DPWA	$\pi N \rightarrow \pi N$ 0-350 MeV
52.8 ± 0.6	4 VASAN 76B		Fit to CARTER 73
• • • We do not use the following data for averages, fits, limits, etc. • • •			
52.45 ± 0.2	5 ZIDELL 78		
52.9 to 53.1	6 VASAN 76B		Fit to CARTER 73

 $\Delta(1232)$ ELASTIC POLE RESIDUES

ABSOLUTE VALUE, MIXED CHARGES

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
52	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
53 ± 2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE, MIXED CHARGES

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-48	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
-31	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-47 ± 1	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

ABSOLUTE VALUE, $\Delta(1232)^{++}$

VALUE (MeV)	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
52.4 to 53.2	4 VASAN 76B	Fit to CARTER 73
52.1 to 52.4	6 VASAN 76B	Fit to CARTER 73

PHASE, $\Delta(1232)^{++}$

VALUE (rad)	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
-0.822 to -0.833	4 VASAN 76B	Fit to CARTER 73
-0.823 to -0.830	6 VASAN 76B	Fit to CARTER 73

ABSOLUTE VALUE, $\Delta(1232)^0$

VALUE (MeV)	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
54.8 to 55.0	4 VASAN 76B	Fit to CARTER 73
55.2 to 55.3	6 VASAN 76B	Fit to CARTER 73

PHASE, $\Delta(1232)^0$

VALUE (rad)	DOCUMENT ID	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •		
-0.840 to -0.847	4 VASAN 76B	Fit to CARTER 73
-0.848 to -0.856	6 VASAN 76B	Fit to CARTER 73

 $\Delta(1232)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	>99 %
Γ_2 $N\gamma$	0.55-0.61 %
Γ_3 $N\gamma$, helicity=1/2	
Γ_4 $N\gamma$, helicity=3/2	

 $\Delta(1232)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.993 to 0.995 OUR ESTIMATE				
1.0	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
1.0	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
1.0	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

 $\Delta(1232)$ PHOTON DECAY AMPLITUDES $\Delta(1232) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.141 ± 0.005 OUR ESTIMATE			
-0.143 ± 0.004	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.135 ± 0.016	DAVIDSON 91B	FIT	$\gamma N \rightarrow \pi N$
-0.145 ± 0.015	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.138 ± 0.004	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.147 ± 0.001	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.145 ± 0.001	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.136 ± 0.006	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.133 ± 0.007	ARNDT 90B	IPWA	See LI 93
-0.137	ARNDT 90B	FIT	See LI 93
-0.140 ± 0.007	DAVIDSON 90	FIT	See DAVIDSON 91B
-0.142 ± 0.007	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.140	7 NOELLE 78		$\gamma N \rightarrow \pi N$
-0.141 ± 0.004	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

Baryon Full Listings

$\Delta(1232)$, $\Delta(1600)$

$\Delta(1232) \rightarrow N\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.257 ± 0.008 OUR ESTIMATE			
-0.262 ± 0.004	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.251 ± 0.033	DAVIDSON 91b	FIT	$\gamma N \rightarrow \pi N$
-0.263 ± 0.026	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.259 ± 0.006	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.264 ± 0.002	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.261 ± 0.002	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.247 ± 0.010	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.244 ± 0.008	ARNDT 90b	IPWA	See LI 93
-0.246	ARNDT 90b	FIT	See LI 93
-0.254 ± 0.011	DAVIDSON 90	FIT	See DAVIDSON 91b
-0.271 ± 0.010	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.247	NOELLE 78	$\gamma N \rightarrow \pi N$	
-0.256 ± 0.003	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

$\Delta(1232) \rightarrow N\gamma$, E_2/M_1 ratio

VALUE	DOCUMENT ID	TECN	COMMENT
-0.015 ± 0.004 OUR AVERAGE			
-0.015 ± 0.005	WORKMAN 92	IPWA	$\gamma N \rightarrow \pi N$
-0.0157 ± 0.0072	DAVIDSON 91b	FIT	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.0107 ± 0.0037	DAVIDSON 90	FIT	$\gamma N \rightarrow \pi N$
-0.015 ± 0.002	DAVIDSON 86	FIT	$\gamma N \rightarrow \pi N$
+0.037 ± 0.004	TANABE 85	FIT	$\gamma N \rightarrow \pi N$

$\Delta(1232)$ PHASE OF $M_1+(3/2)$ PHOTOPRODUCTION MULTIPOLE AMPLITUDE POLE RESIDUE

Information on the phase (and magnitude) of the $M_1+(3/2)$ multipole amplitude pole residue is contained implicitly in the paper of MIROSHNICHENKO 79. They find that the phase is consistent with being equal to that of the elastic pole residue.

$\Delta(1232)^{++}$ MAGNETIC MOMENT

The values are extracted from UCLA and SIN data on $\pi^+ p$ bremsstrahlung using a variety of different theoretical approximations and methods. Our estimate is only a rough guess of the range we expect the moment to lie within.

VALUE (μ_N)	DOCUMENT ID	COMMENT
3.7 to 7.5 OUR ESTIMATE		
• • • We do not use the following data for averages, fits, limits, etc. • • •		
4.52 ± 0.50 ± 0.45	BOSSHARD 91	$\pi^+ p \rightarrow \pi^+ p \gamma$ (SIN data)
3.7 to 4.2	LIN 91b	$\pi^+ p \rightarrow \pi^+ p \gamma$ (from UCLA data)
4.6 to 4.9	LIN 91b	$\pi^+ p \rightarrow \pi^+ p \gamma$ (from SIN data)
5.6 to 7.5	WITTMAN 88	$\pi^+ p \rightarrow \pi^+ p \gamma$ (from UCLA data)
6.9 to 9.8	HELLER 87	$\pi^+ p \rightarrow \pi^+ p \gamma$ (from UCLA data)
4.7 to 6.7	NEFKENS 78	$\pi^+ p \rightarrow \pi^+ p \gamma$ (UCLA data)

$\Delta(1232)$ FOOTNOTES

- Using $\pi^\pm d$ as well, PEDRONI 78 determine $(M^- - M^{++}) + (M^0 - M^+)/3 = 4.6 \pm 0.2$ MeV.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- The accuracy claimed by ZIDELL 80 on the real part is considerably better than is allowed by uncertainties in the beam momentum.
- This VASAN 76b value is from fits to the coulomb-barrier-corrected CARTER 73 phase shift.
- ZIDELL 78 fits the nuclear phase shift without coulomb barrier corrections.
- This VASAN 76b value is from fits to the CARTER 73 nuclear phase shift without coulomb barrier corrections.
- Converted to our conventions using $M = 1232$ MeV, $\Gamma = 110$ MeV from NOELLE 78.

$\Delta(1232)$ REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	+Arndt, Roper, Workman
MANLEY 92	PR D45 4002	+Liesli (KENT) IJP
Also 84	PR D30 904	Manley, Arndt, Goradia, Tepitz (VPI)
WORKMAN 92	PR C46 1546	+Arndt, Li (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
BOSSHARD 91	PR D44 1962	+Amstler+ (ZURI, LBL, VILL, LAUS, UCLA, CATH)
Also 90	PR L 64 2619	Bosshard+ (CATH, LAUS, LBL, VILL, UCLA, ZURI)
DAVIDSON 91b	PR D43 71	+Mukhopadhyay, Wittman (RP)
LIN 92	PR C44 1819	+Lieu, Ding (CUNY, CSOK)
Also 91	PR C43 R930	Lin, Liou (CUNY)
ARNDT 90b	PR C42 1864	+Workman, Li, Roper (VPI)
DAVIDSON 90	PR D42 20	+Mukhopadhyay (RP)
WITTMAN 88	PR C37 2075	(TRIU)
HELLER 87	PR C35 718	+Kumano, Martinez, Moniz (LANL, MIT, ILL)
DAVIDSON 86	PR L 56 804	+Mukhopadhyay, Wittman (RP)
ARNDT 85	PR D32 1085	+Ford, Roper (VPI)
TANABE 85	PR C31 1876	+Ohta (KOMAB)
CRAWFORD 83	NP B211 1	+Morton (GLAS)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)

AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also 82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
ARAI 80	Toronto Conf. 93	Arai, Fujii (INUS)
Also 82	NP B194 251	(GLAS)
CRAWFORD 80	Toronto Conf. 107	(CMU, LBL) IJP
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly
KOCH 80b	NP A336 331	+Pietarinen (KARLT) IJP
ZIDELL 80	PR D21 1255	+Arndt, Roper (VPI) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
MIROSHNIC... 79	SJNP 29 94	Miroshnichenko, Nikiforov, Sanin+ (KFTI) IJP
Translated from YAF 29 188.		
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
NEFKENS 78	PR D18 3911	+Arman, Ballagh, Glodis, Haddock+ (UCLA, CATH) IJP
NOELLE 78	PTP 60 778	(NAGO)
PEDRONI 78	NP A300 321	+Gabathuler, Domingo, Hirt+ (SIN, ISNG, KARLE-) IJP
ZIDELL 78	LNC 21 140	+Arndt, Roper (VPI) IJP
CAMPBELL 76	PR D14 2431	+Shaw, Ball (BOIS, UCI, UTAH) IJP
FELLER 76	NP B104 219	+Fukushima, Horikawa, Kajikawa+ (NAGO, OSAK) IJP
VASAN 76b	NP B106 535	(CMU) IJP
Also 76	NP B106 526	Vasan (CMU) IJP
BERENDS 75	NP B84 342	+Donnachie (LEID, MCHS)
CARTER 73	NP B58 378	+Bugg, Carter (CAVE, LOQM) IJP

$\Delta(1600) P_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+) \text{ Status: } ** *$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

The various analyses are not in good agreement.

$\Delta(1600)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1550 to 1700 (≈ 1600) OUR ESTIMATE			
1706 ± 10	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
1600 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1522 ± 13	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1706	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1690	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
1560	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
1640	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1600)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
250 to 450 (≈ 350) OUR ESTIMATE			
430 ± 73	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
300 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
220 ± 40	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
215	LI 93	IPWA	$\gamma N \rightarrow \pi N$
250	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
180	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
300	² LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1600)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1550	³ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1612	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1550 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1581	ARNDT 85	DPWA	See ARNDT 91
1609 or 1610	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1541 or 1542	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
230	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
200 ± 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
300	ARNDT 85	DPWA	See ARNDT 91
323 or 325	⁴ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
178 or 178	¹ LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1600)$ ELASTIC POLE RESIDUE

MODULUS |r|

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
16	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
17 ± 4	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
- 73	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
- 150 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

Baryon Full Listings

$\Delta(1600), \Delta(1620)$

$\Delta(1600)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-25 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	75-90 %
Γ_4 $\Delta\pi$	40-70 %
Γ_5 $\Delta(1232)\pi, P$ -wave	
Γ_6 $\Delta(1232)\pi, F$ -wave	
Γ_7 $N\rho$	<25 %
Γ_8 $N\rho, S=1/2, P$ -wave	
Γ_9 $N\rho, S=3/2, P$ -wave	
Γ_{10} $N\rho, S=3/2, F$ -wave	
Γ_{11} $N(1440)\pi$	10-35 %
Γ_{12} $N(1440)\pi, P$ -wave	
Γ_{13} $p\gamma$	~0 %
Γ_{14} $N\gamma, \text{helicity}=1/2$	
Γ_{15} $N\gamma, \text{helicity}=3/2$	

$\Delta(1600)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.10 to 0.25 OUR ESTIMATE				
0.12 ± 0.02	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.18 ± 0.04	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.21 ± 0.06	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.36 to -0.28 OUR ESTIMATE				
0.006 to 0.042	5 DEANS	75	DPWA $\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow \Delta(1232)\pi, P$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
+0.27 to +0.33 OUR ESTIMATE				
+0.29 ± 0.02	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
+0.24 ± 0.05	BARNHAM	80	IPWA $\pi N \rightarrow N\pi\pi$	
+0.34	1,6 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	
+0.30	2 LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow \Delta(1232)\pi, F$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
-0.15 to -0.03 OUR ESTIMATE				
-0.07	1,6 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow N\rho, S=1/2, P$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
+0.10				
+0.10	1,6 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow N\rho, S=3/2, P$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
+0.10				
+0.10	1,6 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1600) \rightarrow N(1440)\pi, P$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{12})^{1/2}/\Gamma$
+0.15 to +0.23 OUR ESTIMATE				
+0.16 ± 0.02	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
+0.23 ± 0.04	BARNHAM	80	IPWA $\pi N \rightarrow N\pi\pi$	

$\Delta(1600)$ PHOTON DECAY AMPLITUDES

$\Delta(1600) \rightarrow N\gamma, \text{helicity-1/2}$ amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.026 ± 0.020 OUR ESTIMATE			
-0.026 ± 0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.039 ± 0.030	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
-0.046 ± 0.013	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.005 ± 0.020	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.200	7 WADA	84	DPWA Compton scattering
0.000 ± 0.030	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
0.0 ± 0.020	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

$\Delta(1600) \rightarrow N\gamma, \text{helicity-3/2}$ amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.006 ± 0.017 OUR ESTIMATE			
-0.016 ± 0.002	LI	93	IPWA $\gamma N \rightarrow \pi N$
-0.013 ± 0.014	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.025 ± 0.031	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
-0.009 ± 0.020	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.023	WADA	84	DPWA Compton scattering
0.000 ± 0.045	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
0.0 ± 0.015	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

$\Delta(1600)$ FOOTNOTES

- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The range given is from the four best solutions. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.
- LONGACRE 77 considers this coupling to be well determined.
- WADA 84 is inconsistent with other analyses — see the Note on N and Δ Resonances.

$\Delta(1600)$ REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER	94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER	93	πN Newsletter 9 1	(KARL)
LI	93	PR C47 2759	(VPI)
MANLEY	92	PR D45 4002	(KENT) IJP
Also	84	PR D30 904	(VPI)
ARNDT	91	PR D43 2131	(VPI, TELE) IJP
ARNDT	85	PR D32 1085	(VPI)
WADA	84	NP B247 313	(INUS)
CRAWFORD	83	NP B211 1	(GLAS)
PDG	82	PL 111B	(HELS, CIT, CERN)
AWAJI	81	Bonn Conf. 352	(NAGO)
Also	82	NP B197 365	(NAGO)
BARNHAM	80	NP B168 243	(LOIC)
CRAWFORD	80	Toronto Conf. 107	(GLAS)
CUTKOSKY	80	Toronto Conf. 19	(CMU, LBL) IJP
Also	79	PR D20 2839	(CMU, LBL) IJP
HOEHLER	79	PDAT 12-1	(KARLT) IJP
Also	80	Toronto Conf. 3	(KARLT) IJP
BARBOUR	78	NP B141 253	(GLAS)
LONGACRE	78	PR D17 1795	(LBL, SLAC)
LONGACRE	77	NP B122 493	(SACL) IJP
Also	76	NP B108 365	(SACL) IJP
WINNIK	77	NP B128 66	(HAIF) I
FELLER	76	NP B104 219	(NAGO, OSAK) IJP
DEANS	75	NP B96 90	(SFLA, ALAH) IJP
LONGACRE	75	PL 558 415	(LBL, SLAC) IJP

$\Delta(1620) S_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

$\Delta(1620)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1615 to 1675 (≈ 1620) OUR ESTIMATE			
1672 ± 7	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
1620 ± 20	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1610 ± 7	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1669	LI	93	IPWA $\gamma N \rightarrow \pi N$
1620	BARNHAM	80	IPWA $\pi N \rightarrow N\pi\pi$
1712.8 ± 6.0	1 CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
1786.7 ± 2.0	1 CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
1657	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
1662	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
1580	2 LONGACRE	77	IPWA $\pi N \rightarrow N\pi\pi$
1600	3 LONGACRE	75	IPWA $\pi N \rightarrow N\pi\pi$

Baryon Full Listings

$\Delta(1620)$

$\Delta(1620)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
120 to 180 (≈ 150) OUR ESTIMATE			
154 ± 37	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
140 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
139 ± 18	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
184	LI 93	IPWA	$\gamma N \rightarrow \pi N$
120	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
228.3 ± 18.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$ (lower mass)
30.0 ± 6.4	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$ (higher mass)
161	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
180	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
120	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
150	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1620)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1608	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1587	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1600 ± 15	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1599	ARNDT 85	DPWA	See ARNDT 91
1583 or 1583	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
1575 or 1572	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
116	⁴ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
120	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
120 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
120	ARNDT 85	DPWA	See ARNDT 91
143 or 149	⁵ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$
119 or 128	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1620)$ ELASTIC POLE RESIDUE

MODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
19	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
15	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
15 ± 2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-95	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
-125	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-110 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(1620)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	20-30 %
Γ_2 $N\pi\pi$	70-80 %
Γ_3 $\Delta\pi$	30-60 %
Γ_4 $\Delta(1232)\pi$, D-wave	
Γ_5 $N\rho$	7-25 %
Γ_6 $N\rho$, S=1/2, S-wave	
Γ_7 $N\rho$, S=3/2, D-wave	
Γ_8 $N(1440)\pi$	
Γ_9 $N\gamma$	0.02-0.06 %
Γ_{10} $N\gamma$, helicity=1/2	

$\Delta(1620)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.2 to 0.3 OUR ESTIMATE				
0.09 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.25 ± 0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.35 ± 0.06	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.60	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$ (lower mass)	
0.36	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$ (higher mass)	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620)$ S_{31} coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow \Delta(1620) \rightarrow \Delta(1232)\pi$, D-wave $(\Gamma_1\Gamma_4)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.36 to -0.28 OUR ESTIMATE			
-0.24 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.33 ± 0.06	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
-0.39	^{2,6} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-0.40	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow \Delta(1620) \rightarrow N\rho$, S=1/2, S-wave $(\Gamma_1\Gamma_6)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.12 to +0.22 OUR ESTIMATE			
+0.15 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.40 ± 0.10	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$
+0.08	^{2,6} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
+0.28	³ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow \Delta(1620) \rightarrow N\rho$, S=3/2, D-wave $(\Gamma_1\Gamma_7)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
-0.15 to -0.03 OUR ESTIMATE			
-0.06 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
-0.13	^{2,6} LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{total}$ in $N\pi \rightarrow \Delta(1620) \rightarrow N(1440)\pi$ $(\Gamma_1\Gamma_8)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
0.11 ± 0.05	BARNHAM 80	IPWA	$\pi N \rightarrow N\pi\pi$

$\Delta(1620)$ PHOTON DECAY AMPLITUDES

$\Delta(1620) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV $^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
+0.030 \pm 0.014 OUR ESTIMATE			
0.042 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.035 ± 0.010	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.010 ± 0.015	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.022 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.026 ± 0.008	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.021 ± 0.020	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
0.126 ± 0.021	TAKEDA 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.066	WADA 84	DPWA	Compton scattering
+0.034 ± 0.028	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
-0.005 ± 0.016	FELLER 76	DPWA	$\gamma N \rightarrow \pi N$

$\Delta(1620)$ FOOTNOTES

- CHEW 80 reports two S_{31} resonances at somewhat higher masses than other analyses. Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- LONGACRE 77 considers this coupling to be well determined.

$\Delta(1620)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	+Saleski (KENT) IJP
Also	84 PR D30 904	+Manley, Arndt, Goradia, Tepitz (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
ARNDT 85	PR D32 1085	+Ford, Roper (VPI)
WADA 84	NP B247 313	+Egawa, Imanishi, Ishii, Kato, Ukai+ (INUS)
CRAWFORD 83	NP B211 1	+Morton (GLAS)
HOEHLER 83	Landolt-Boernstein 1/9B2	(KARLT)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also	82 NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
ARAI 80	Toronto Conf. 93	(INUS)
Also	82 NP B194 251	Arai, Fujii (INUS)

See key on page 1343

Baryon Full Listings
 $\Delta(1620)$, $\Delta(1700)$

BARNHAM	80	NP B168 243	+Glickman, Mier-Jedrzejowicz+	(LOIC)
CHEW	80	Toronto Conf. 123		(LBL) IJP
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19		(CMU, LBL) IJP
Also	79	PR D20 2839	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
TAKEDA	80	NP B168 17	+Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
HOEHLER	79	PDAT 12-1	+Arai, Fujii, Ikeda, Iwasaki+	(TOKY, INUS)
Also	80	Toronto Conf. 3	+Kaiser, Koch, Pietarinen	(KARLT) IJP
BARBOUR	78	NP B141 253	Koch	(KARLT) IJP
LONGACRE	78	PR D17 1795	+Crawford, Parsons	(GLAS)
LONGACRE	77	NP B122 493	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)
Also	76	NP B108 365	+Doibeau	(SACL) IJP
FELLER	76	NP B104 219	+Doibeau, Triantis, Neveu, Cadiet	(SACL) IJP
LONGACRE	75	PL 55B 415	+Fukushima, Horikawa, Kajikawa+	(NAGO, OSAK) IJP
			+Rosenfeld, Lasinski, Smadja+	(LBL, SLAC) IJP

 $\Delta(1700) D_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $\Delta(1700)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1670 to 1770 (≈ 1700) OUR ESTIMATE			
1762 ± 44	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1710 ± 30	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
1680 ± 70	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1655	LI	93 IPWA	$\gamma N \rightarrow \pi N$
1650	BARNHAM	80 IPWA	$\pi N \rightarrow N\pi\pi$
1718.4 $^{+13.1}_{-13.0}$	¹ CHEW	80 BPWA	$\pi^+ p \rightarrow \pi^+ p$
1622	CRAWFORD	80 DPWA	$\gamma N \rightarrow \pi N$
1629	BARBOUR	78 DPWA	$\gamma N \rightarrow \pi N$
1600	² LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$
1680	³ LONGACRE	75 IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1700)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200 to 400 (≈ 300) OUR ESTIMATE			
600 ± 250	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
280 ± 80	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
230 ± 80	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
348	LI	93 IPWA	$\gamma N \rightarrow \pi N$
160	BARNHAM	80 IPWA	$\pi N \rightarrow N\pi\pi$
193.3 ± 26.0	¹ CHEW	80 BPWA	$\pi^+ p \rightarrow \pi^+ p$
209	CRAWFORD	80 DPWA	$\gamma N \rightarrow \pi N$
216	BARBOUR	78 DPWA	$\gamma N \rightarrow \pi N$
200	² LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$
240	³ LONGACRE	75 IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1700)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
VALUE (MeV)			
1651	⁴ HOEHLER	93 SPED	$\pi N \rightarrow \pi N$
1646	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1675 ± 25	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1668	ARNDT	85 DPWA	See ARNDT 91
1681 or 1672	⁵ LONGACRE	78 IPWA	$\pi N \rightarrow N\pi\pi$
1600 or 1594	² LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$
-2xIMAGINARY PART			
VALUE (MeV)			
159	⁴ HOEHLER	93 SPED	$\pi N \rightarrow \pi N$
208	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
220 ± 40	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
320	ARNDT	85 DPWA	See ARNDT 91
245 or 241	⁵ LONGACRE	78 IPWA	$\pi N \rightarrow N\pi\pi$
208 or 201	² LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1700)$ ELASTIC POLE RESIDUE

MODULUS $ r $	DOCUMENT ID	TECN	COMMENT
VALUE (MeV)			
10	HOEHLER	93 SPED	$\pi N \rightarrow \pi N$
13	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
13 ± 3	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-22	ARNDT	91 DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-20 ± 25	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1700)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-20 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	80-90 %
Γ_4 $\Delta\pi$	30-60 %
Γ_5 $\Delta(1232)\pi$, S-wave	25-50 %
Γ_6 $\Delta(1232)\pi$, D-wave	1-7 %
Γ_7 $N\rho$	30-55 %
Γ_8 $N\rho$, S=1/2, D-wave	
Γ_9 $N\rho$, S=3/2, S-wave	5-20 %
Γ_{10} $N\rho$, S=3/2, D-wave	
Γ_{11} $N\gamma$	0.16-0.28 %
Γ_{12} $N\gamma$, helicity=1/2	
Γ_{13} $N\gamma$, helicity=3/2	

 $\Delta(1700)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.10 to 0.20 OUR ESTIMATE				
0.14 ± 0.06	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.12 ± 0.03	CUTKOSKY	80 IPWA	$\pi N \rightarrow \pi N$	
0.20 ± 0.03	HOEHLER	79 IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.16	¹ CHEW	80 BPWA	$\pi^+ p \rightarrow \pi^+ p$	
$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1700) \rightarrow \Sigma K$ ($\Gamma_1\Gamma_2)^{1/2}/\Gamma$				
VALUE	DOCUMENT ID	TECN	COMMENT	
0.002	LIVANOS	80 DPWA	$\pi p \rightarrow \Sigma K$	
0.001 to 0.011	⁶ DEANS	75 DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1700) \rightarrow \Delta(1232)\pi$, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
+0.21 to +0.29 OUR ESTIMATE				
+0.32 ± 0.06	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
+0.18 ± 0.04	BARNHAM	80 IPWA	$\pi N \rightarrow N\pi\pi$	
+0.30	^{2,7} LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$	
+0.24	³ LONGACRE	75 IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1700) \rightarrow \Delta(1232)\pi$, D-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
+0.06 to +0.11 OUR ESTIMATE				
+0.08 ± 0.03	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.14 ± 0.04	BARNHAM	80 IPWA	$\pi N \rightarrow N\pi\pi$	
+0.05	^{2,7} LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$	
+0.10	³ LONGACRE	75 IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1700) \rightarrow N\rho$, S=1/2, D-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
+0.17 ± 0.05				
	BARNHAM	80 IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1700) \rightarrow N\rho$, S=3/2, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
± 0.11 to ± 0.19 OUR ESTIMATE				
+0.10 ± 0.03	MANLEY	92 IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
+0.04	^{2,7} LONGACRE	77 IPWA	$\pi N \rightarrow N\pi\pi$	
-0.30	³ LONGACRE	75 IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1700) \rightarrow N\rho$, S=3/2, D-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_{10})^{1/2}/\Gamma$
0.18 ± 0.07				
	BARNHAM	80 IPWA	$\pi N \rightarrow N\pi\pi$	

Baryon Full Listings

 $\Delta(1700)$, $\Delta(1750)$, $\Delta(1900)$ $\Delta(1700)$ PHOTON DECAY AMPLITUDES $\Delta(1700) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.114 ± 0.013 OUR ESTIMATE			
0.121 ± 0.004	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.111 ± 0.017	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.089 ± 0.033	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.112 ± 0.006	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.130 ± 0.006	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.123 ± 0.022	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.130 ± 0.037	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.072 ± 0.033	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

 $\Delta(1700) \rightarrow N\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.091 ± 0.029 OUR ESTIMATE			
0.115 ± 0.004	LI	93	IPWA $\gamma N \rightarrow \pi N$
0.107 ± 0.015	CRAWFORD	83	IPWA $\gamma N \rightarrow \pi N$
0.060 ± 0.015	AWAJI	81	DPWA $\gamma N \rightarrow \pi N$
0.047 ± 0.007	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 1)
0.050 ± 0.007	ARAI	80	DPWA $\gamma N \rightarrow \pi N$ (fit 2)
0.102 ± 0.015	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.098 ± 0.036	BARBOUR	78	DPWA $\gamma N \rightarrow \pi N$
+0.087 ± 0.023	FELLER	76	DPWA $\gamma N \rightarrow \pi N$

 $\Delta(1700)$ FOOTNOTES

- Problems with CHEW 80 are discussed in section 2.1.11 of HOEHLER 83.
- LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from the Saclay (CERN) partial-wave analysis.
- The range given is from the four best solutions. DEANS 75 disagrees with $\pi^+ p \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.
- LONGACRE 77 considers this coupling to be well determined.

 $\Delta(1700)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER	94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER	93	πN Newsletter 9 1	(KARL)
LI	93	PR C47 2759	(VPI)
MANLEY	92	PR D45 4002	(KENT) IJP
Also	84	PR D30 904	(VPI)
ARNDT	91	PR D43 2131	+Li, Roper, Workman, Ford
ARNDT	85	PR D32 1085	+Ford, Roper
CRAWFORD	83	NP B211 1	+Morton
HOEHLER	83	Landolt-Boernstein 1/9B2	(KARLT)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELs, CIT, CERN)
AWAJI	81	Bonn Conf. 352	+Kajikawa (NAGO)
Also	82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
ARAI	80	Toronto Conf. 93	(INUS)
Also	82	NP B194 251	Arai, Fujii (INUS)
BARNHAM	80	NP B168 243	+Glickman, Mier-Jedrzejowicz+ (LOIC)
CHEW	80	Toronto Conf. 123	(LBL) IJP
CRAWFORD	80	Toronto Conf. 107	(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu (SACL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also	80	Toronto Conf. 3	Koch (KARLT) IJP
BARBOUR	78	NP B141 253	+Crawford, Parsons (GLAS)
LONGACRE	78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+ (LBL, SLAC)
LONGACRE	77	NP B122 493	+Dolbeau (SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet (SACL) IJP
WINNIK	77	NP B128 66	+Toaff, Revel, Goldberg, Bery (HAIF) I
FELLER	76	NP B104 219	+Fukushima, Honkawa, Kajikawa+ (NAGO, OSAK) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
LONGACRE	75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+ (LBL, SLAC) IJP

 $\Delta(1750) P_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^+) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

 $\Delta(1750)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1750 OUR ESTIMATE			
1744 ± 36	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1715.2 ± 21.0	¹ CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
1778.4 ± 9.0	¹ CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$

 $\Delta(1750)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 ± 120	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
93.3 ± 55.0	¹ CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
23.0 ± 29.0	¹ CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$

 $\Delta(1750)$ DECAY MODES

Mode	
Γ_1	$N\pi$
Γ_2	$N\pi\pi$
Γ_3	$N(1440)\pi$

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.08 ± 0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.18	¹ CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$	
0.20	¹ CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1700) \rightarrow N(1440)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
0.15 ± 0.03	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	

 $\Delta(1750)$ FOOTNOTES

- CHEW 80 reports four resonances in the P_{31} wave — see also the $\Delta(1910)$. Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.

 $\Delta(1750)$ REFERENCES

MANLEY	92	PR D45 4002	+Saleski	(KENT)
Also	84	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
HOEHLER	83	Landolt-Boernstein 1/9B2		(KARLT)
CHEW	80	Toronto Conf. 123		(LBL)

 $\Delta(1900) S_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-) \text{ Status: } ***$$

 $\Delta(1900)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1850 to 1950 (≈ 1900) OUR ESTIMATE			
1920 ± 24	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
1890 ± 50	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
1908 ± 30	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1918.5 ± 23.0	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
1803	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$

 $\Delta(1900)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
140 to 240 (≈ 200) OUR ESTIMATE			
263 ± 39	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
170 ± 50	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
140 ± 40	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
93.5 ± 54.0	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
137	CRAWFORD	80	DPWA $\gamma N \rightarrow \pi N$

Baryon Full Listings

$\Delta(1900), \Delta(1905)$

 $\Delta(1900)$ POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1780	¹ HOEHLER 93 SPED		$\pi N \rightarrow \pi N$
1870 ± 40	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91 DPWA		$\pi N \rightarrow \pi N$ Soln SM90
2029 or 2025	² LONGACRE 78 IPWA		$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
180 ± 50	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91 DPWA		$\pi N \rightarrow \pi N$ Soln SM90
164 or 163	² LONGACRE 78 IPWA		$\pi N \rightarrow N\pi\pi$

 $\Delta(1900)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10 ± 3	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
+20 ± 40	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$

 $\Delta(1900)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-30 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	
Γ_4 $\Delta\pi$	
Γ_5 $\Delta(1232)\pi, D\text{-wave}$	
Γ_6 $N\rho$	
Γ_7 $N\rho, S=1/2, S\text{-wave}$	
Γ_8 $N\rho, S=3/2, D\text{-wave}$	
Γ_9 $N(1440)\pi, S\text{-wave}$	
Γ_{10} $N\gamma, \text{helicity}=1/2$	

 $\Delta(1900)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.1 to 0.3 OUR ESTIMATE				
0.41 ± 0.04	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.10 ± 0.03	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$	
0.08 ± 0.04	HOEHLER 79 IPWA		$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.28	CHEW 80 BPWA		$\pi^+ p \rightarrow \pi^+ p$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
<0.03	CANDLIN 84 DPWA		$\pi^+ p \rightarrow \Sigma^+ K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.076	³ DEANS 75 DPWA		$\pi N \rightarrow \Sigma K$	
0.11	LANGBEIN 73 IPWA		$\pi N \rightarrow \Sigma K$ (sol. 1)	
0.12	LANGBEIN 73 IPWA		$\pi N \rightarrow \Sigma K$ (sol. 2)	

$(\Gamma_1\Gamma_5)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow \Delta(1232)\pi, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
+0.25 ± 0.07	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N\rho, S=1/2, S\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_7)^{1/2}/\Gamma$
-0.14 ± 0.11	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_8)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N\rho, S=3/2, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
-0.37 ± 0.07	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_9)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1900) \rightarrow N(1440)\pi, S\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_9)^{1/2}/\Gamma$
-0.16 ± 0.11	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$	

 $\Delta(1900)$ PHOTON DECAY AMPLITUDES **$\Delta(1900) \rightarrow N\gamma, \text{helicity-1/2 amplitude } A_{1/2}$**

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.004 ± 0.016	CRAWFORD 83 IPWA		$\gamma N \rightarrow \pi N$
0.029 ± 0.008	AWAJI 81 DPWA		$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.006 to -0.025	CRAWFORD 80 DPWA		$\gamma N \rightarrow \pi N$

 $\Delta(1900)$ FOOTNOTES

¹ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

² LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

³ The value given is from solution 1; the resonance is not present in solutions 2, 3, or 4.

 $\Delta(1900)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
MANLEY 92	PR D45 4002 +Saleski	(KENT) IJP
Also 84	PR D30 904 Manley, Arndt, Goradia, Tepelitz	(VPI)
ARNDT 91	PR D43 2131 +Li, Roper, Workman, Ford	(VPI, TELE) IJP
CANDLIN 84	NP B238 477 +Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
CRAWFORD 83	NP B211 1 +Morton	(GLAS)
AWAJI 81	Bonn Conf. 352 +Kajikawa	(NAGO)
CHEW 80	NP B197 365 Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
CRAWFORD 80	Toronto Conf. 107	(LBL) IJP
CUTKOSKY 80	Toronto Conf. 19 +Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also 79	PR D20 2839 Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
HOEHLER 79	PDAT 12-1 +Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3 Koch	(KARLT) IJP
LONGACRE 78	PR D17 1795 +Lasinski, Rosenfeld, Smaaja+	(LBL, SLAC)
DEANS 75	NP B96 90 +Mitchell, Montgomery+	(SFLA, ALAH) IJP
LANGBEIN 73	NP B53 251 +Wagner	(MUNI) IJP

 $\Delta(1905) F_{35}$

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $\Delta(1905)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1870 to 1920 (≈ 1905) OUR ESTIMATE			
1881 ± 18	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$
1910 ± 30	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$
1905 ± 20	HOEHLER 79 IPWA		$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1960 ± 40	CANDLIN 84 DPWA		$\pi^+ p \rightarrow \Sigma^+ K^+$
1787.0 [±] 6.0 - 5.7	CHEW 80 BPWA		$\pi^+ p \rightarrow \pi^+ p$
1880	CRAWFORD 80 DPWA		$\gamma N \rightarrow \pi N$
1892	BARBOUR 78 DPWA		$\gamma N \rightarrow \pi N$
1830	¹ LONGACRE 75 IPWA		$\pi N \rightarrow N\pi\pi$

 $\Delta(1905)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
280 to 440 (≈ 380) OUR ESTIMATE			
327 ± 51	MANLEY 92 IPWA		$\pi N \rightarrow \pi N$ & $N\pi\pi$
400 ± 100	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$
260 ± 20	HOEHLER 79 IPWA		$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
270 ± 40	CANDLIN 84 DPWA		$\pi^+ p \rightarrow \Sigma^+ K^+$
66.0 [±] 24.0 - 16.0	CHEW 80 BPWA		$\pi^+ p \rightarrow \pi^+ p$
193	CRAWFORD 80 DPWA		$\gamma N \rightarrow \pi N$
159	BARBOUR 78 DPWA		$\gamma N \rightarrow \pi N$
220	¹ LONGACRE 75 IPWA		$\pi N \rightarrow N\pi\pi$

 $\Delta(1905)$ POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1829	² HOEHLER 93 SPED		$\pi N \rightarrow \pi N$
1794	ARNDT 91 DPWA		$\pi N \rightarrow \pi N$ Soln SM90
1830 ± 40	CUTKOSKY 80 IPWA		$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1830	ARNDT 85 DPWA		See ARNDT 91
1813 or 1808	³ LONGACRE 78 IPWA		$\pi N \rightarrow N\pi\pi$

Baryon Full Listings

 $\Delta(1905)$ $-2 \times \text{IMAGINARY PART}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
303	2 HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
230	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
280 ± 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
180	ARNDT 85	DPWA	See ARNDT 91
193 or 187	3 LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1905)$ ELASTIC POLE RESIDUEMODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
25	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
14	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
25 ± 8	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
-40	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-50 ± 20	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1905)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	5-15 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	85-95 %
Γ_4 $\Delta\pi$	<25 %
Γ_5 $\Delta(1232)\pi$, P-wave	
Γ_6 $\Delta(1232)\pi$, F-wave	
Γ_7 $N\rho$	>60 %
Γ_8 $N\rho$, S=3/2, P-wave	
Γ_9 $N\rho$, S=3/2, F-wave	
Γ_{10} $N\rho$, S=1/2, F-wave	
Γ_{11} $N\gamma$	0.01-0.04 %
Γ_{12} $N\gamma$, helicity=1/2	
Γ_{13} $N\gamma$, helicity=3/2	

 $\Delta(1905)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_i/Γ
0.06 to 0.15 OUR ESTIMATE				
0.12 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.08 ± 0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.15 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.11	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1905) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.015 ± 0.003	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.013	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$	
0.021 to 0.054	4 DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1905) \rightarrow \Delta(1232)\pi$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.04 ± 0.05	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1905) \rightarrow \Delta(1232)\pi$, F-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
+0.02 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
+0.20	1 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.17	5 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	
+0.06	6 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	

 $(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1905) \rightarrow N\rho$, S=3/2, P-wave $(\Gamma_1\Gamma_2)^{1/2}/\Gamma$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.030 to +0.36 OUR ESTIMATE			
+0.33 ± 0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
+0.33	1 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.26	5 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$
+0.11 to +0.33	7 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1905)$ PHOTON DECAY AMPLITUDES $\Delta(1905) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.037 ± 0.016 OUR ESTIMATE			
0.055 ± 0.004	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.021 ± 0.010	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.043 ± 0.020	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
0.022 ± 0.010	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
0.031 ± 0.009	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
0.024 ± 0.014	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.033 ± 0.018	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1905) \rightarrow N\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
-0.031 ± 0.030 OUR ESTIMATE			
-0.002 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.056 ± 0.028	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
-0.025 ± 0.023	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.029 ± 0.007	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.045 ± 0.006	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.072 ± 0.035	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.055 ± 0.019	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1905)$ FOOTNOTES

- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The range given for DEANS 75 is from the four best solutions.
- A Breit-Wigner fit to the HERNDON 75 IPWA.
- A Breit-Wigner fit to the NOVOSELLER 78B IPWA.
- A Breit-Wigner fit to the NOVOSELLER 78B IPWA; the phase is near 90°.

 $\Delta(1905)$ REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	+Arndt, Roper, Workman (VPI)
MANLEY 92	PR D45 4002	+Saleski (KENT) IJP
Also 84	PR D30 904	Manley, Arndt, Goradia, Tepiltz (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJP
ARNDT 85	PR D32 1085	+Ford, Roper (VPI)
CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+ (EDIN, RAL, LOWC)
CRAWFORD 83	NP B211 1	+Morton (GLAS)
PDG 82	PL 111B	+Roes, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also 82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
ARAI 80	Toronto Conf. 93	(INUS)
Also 82	NP B194 251	Arai, Fujii (INUS)
CHEW 80	Toronto Conf. 123	(LBL) IJP
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
LIVANOS 80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu (SACL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+ (LBL, SLAC)
NOVOSELLER 78	NP B137 509	(CIT) IJP
NOVOSELLER 78B	NP B137 445	(CIT) IJP
DEANS 75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
HERNDON 75	PR D11 3183	+Longacre, Miller, Rosenfeld+ (LBL, SLAC)
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smadja+ (LBL, SLAC) IJP

See key on page 1343

Baryon Full Listings
 $\Delta(1910)$ $\Delta(1910) P_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $\Delta(1910)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1870 to 1920 (≈ 1910) OUR ESTIMATE			
1882 ± 10	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1910 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1888 ± 20	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1960.1 ± 21.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2121.4 ^{+13.0} _{-14.3}	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
1921	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1899	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1790	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1910)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
190 to 270 (≈ 250) OUR ESTIMATE			
239 ± 25	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
225 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
280 ± 50	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
152.9 ± 60.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
172.2 ± 37.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
351	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
230	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
170	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1910)$ POLE POSITION

REAL PART			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1874	³ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1950	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1880 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1792 or 1801	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$
-2xIMAGINARY PART			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
283	³ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
398	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
200 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
172 or 165	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1910)$ ELASTIC POLE RESIDUE

MODULUS r			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
38	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
37	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
20 ± 4	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ			
VALUE (°)	DOCUMENT ID	TECN	COMMENT
-91	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-90 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1910)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	15-30 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	
Γ_4 $\Delta\pi$	
Γ_5 $\Delta(1232)\pi, P\text{-wave}$	
Γ_6 $N\rho$	
Γ_7 $N\rho, S=3/2, P\text{-wave}$	
Γ_8 $N(1440)\pi$	
Γ_9 $N(1440)\pi, P\text{-wave}$	
Γ_{10} $N\gamma, \text{helicity}=1/2$	

 $\Delta(1910)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.15 to 0.3 OUR ESTIMATE				
0.23 ± 0.08	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.19 ± 0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.24 ± 0.06	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	
0.40	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma)^{1/2}/\Gamma$
0.15 to 0.3 OUR ESTIMATE				
< 0.03	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.019	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$	
0.082 to 0.184	⁴ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow \Delta(1232)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma)^{1/2}/\Gamma$
0.15 to 0.3 OUR ESTIMATE				
+0.06	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow N\rho, S=3/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma)^{1/2}/\Gamma$
0.15 to 0.3 OUR ESTIMATE				
+0.29	² LONGACRE 77	IPWA	$\pi N \rightarrow N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.17	⁵ NOVOSSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1910) \rightarrow N(1440)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma)^{1/2}/\Gamma$
0.15 to 0.3 OUR ESTIMATE				
-0.39 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

 $\Delta(1910)$ PHOTON DECAY AMPLITUDES

$\Delta(1910) \rightarrow N\gamma, \text{helicity-1/2 amplitude } A_{1/2}$			
VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
+0.013 ± 0.022 OUR ESTIMATE			
0.032 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.014 ± 0.030	CRAWFORD 83	IPWA	$\gamma N \rightarrow \pi N$
0.025 ± 0.011	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.012 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.031 ± 0.004	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.005 ± 0.030	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.035 ± 0.021	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1910)$ FOOTNOTES

- ¹ CHEW 80 reports four resonances in the P_{31} wave — see also the $\Delta(1750)$. Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.
- ² LONGACRE 77 pole positions are from a search for poles in the unitarized T-matrix; the first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis. The other LONGACRE 77 values are from eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- ³ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ⁴ The range given for DEANS 75 is from the four best solutions.
- ⁵ Evidence for this coupling is weak; see NOVOSSELLER 78. This coupling assumes the mass is near 1820 MeV.

 $\Delta(1910)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	+Saleski (KENT) IJF
Also 84	PR D30 304	(VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TELE) IJF
CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+ (EDIN, RAL, LOWC)
CRAWFORD 83	NP B211 1	+Morton (GLAS)
HOEHLER 83	Landolt-Boernstein 1/9B2	(KARLT)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also 82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)

Baryon Full Listings

$\Delta(1910), \Delta(1920)$

ARAI	80	Toronto Conf. 93		(INUS)
Also	82	NP B194 251	Arai, Fujii	(INUS)
CHEW	80	Toronto Conf. 123		(LBL) IJP
CRAWFORD	80	Toronto Conf. 107		(GLAS)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
LIVANOS	80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu	(SACL) IJP
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(GLAS)
BARBOUR	78	NP B141 253	+Crawford, Parsons	(KARLT) IJP
NOVOSELLER	78	NP B137 509		(CIT) IJP
Also	78B	NP B137 445	Novoseller	(CIT) IJP
LONGACRE	77	NP B122 493	+Dolbeau	(SACL) IJP
Also	76	NP B108 365	Dolbeau, Triantis, Neveu, Cadiet	(SACL) IJP
DEANS	75	NP B96 90	+Mitchell, Montgomery+	(SFLA, ALAH) IJP

$\Delta(1920) P_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

$\Delta(1920)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 to 1970 (≈ 1920) OUR ESTIMATE			
2014 ± 16	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1920 ± 80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1868 ± 10	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1840 ± 40	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
1955.0 ± 13.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2065.0 ± 13.6	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2065.0 - 12.9			

$\Delta(1920)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 to 300 (≈ 200) OUR ESTIMATE			
152 ± 55	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
300 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
220 ± 80	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
200 ± 40	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
88.3 ± 35.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
62.0 ± 44.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

$\Delta(1920)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900	² HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
1900 ± 80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90

$\Delta(1920)$ ELASTIC POLE RESIDUE

MODULUS r VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
24 ± 4	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ VALUE (°)	DOCUMENT ID	TECN	COMMENT
-150 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(1920)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	5-20 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	
Γ_4 $\Delta(1232)\pi, P\text{-wave}$	
Γ_5 $N(1440)\pi, P\text{-wave}$	
Γ_6 $N\gamma, \text{helicity}=1/2$	
Γ_7 $N\gamma, \text{helicity}=3/2$	

$\Delta(1920)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$ VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.05 to 0.2 OUR ESTIMATE				
0.02 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.20 ± 0.05	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.14 ± 0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	
0.18	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1920) \rightarrow \Sigma K$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.052 ± 0.015	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.049	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$	
0.048 to 0.120	³ DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1920) \rightarrow \Delta(1232)\pi, P\text{-wave}$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
-0.13 ± 0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.3	⁴ NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	
0.27	⁵ NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_i/\Gamma)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1920) \rightarrow N(1440)\pi, P\text{-wave}$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
+0.06 ± 0.07	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

$\Delta(1920)$ PHOTON DECAY AMPLITUDES

$\Delta(1920) \rightarrow N\gamma, \text{helicity-1/2}$ amplitude $A_{1/2}$ VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
0.040 ± 0.014	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$

$\Delta(1920) \rightarrow N\gamma, \text{helicity-3/2}$ amplitude $A_{3/2}$ VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
0.023 ± 0.017	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$

$\Delta(1920)$ FOOTNOTES

- ¹ CHEW 80 reports two P_{33} resonances in this mass region. Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.
- ² See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- ³ The range given for DEANS 75 is from the four best solutions.
- ⁴ A Breit-Wigner fit to the HERNDON 75 IPWA; the phase is near -90° .
- ⁵ A Breit-Wigner fit to the NOVOSELLER 78B IPWA; the phase is near -90° .

$\Delta(1920)$ REFERENCES

For early references, see Physics Letters **111B** 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
MANLEY 92	PR D45 4002	+Saleski (KENT) IJP
Also	PR D30 304	Manley, Arndt, Goradia, Tsapiiz (VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford (VPI, TEL) IJP
CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+ (EDIN, RAL, LOWC)
HOEHLER 83	Landolt-Boernstein 1/9B2	(KARLT)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+ (NAGO)
CHEW 80	Toronto Conf. 123	(LBL) IJP
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly (CMU, LBL) IJP
LIVANOS 80	Toronto Conf. 35	+Baton, Coutures, Kochowski, Neveu (SACL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also	Toronto Conf. 3	Koch (KARLT) IJP
NOVOSELLER 78	NP B137 509	(CIT)
NOVOSELLER 78B	NP B137 445	(CIT)
DEANS 75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
HERNDON 75	PR D11 3183	+Longacre, Miller, Rosenfeld+ (LBL, SLAC)

See key on page 1343

Baryon Full Listings
 $\Delta(1930), \Delta(1940)$ $\Delta(1930) D_{35}$

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^-) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

The various analyses are not in good agreement.

 $\Delta(1930)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1920 to 1970 (≈ 1930) OUR ESTIMATE			
1956 ± 22	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1940 ± 30	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1901 ± 15	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
1963	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1910.0 $^{+15.0}_{-17.2}$	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2000	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
2024	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1930)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
280 to 450 (≈ 350) OUR ESTIMATE			
530 ± 140	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
320 ± 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
195 ± 60	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
260	LI 93	IPWA	$\gamma N \rightarrow \pi N$
74.8 $^{+17.0}_{-16.0}$	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
442	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
462	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1930)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1850	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
2018	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1890 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
180	¹ HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
398	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
260 ± 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1930)$ ELASTIC POLE RESIDUEMODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
20	HOEHLER 93	SPED	$\pi N \rightarrow \pi N$
15	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
18 ± 6	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-24	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
-20 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1930)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	10-20 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	
Γ_4 $N\gamma$, helicity=1/2	
Γ_5 $N\gamma$, helicity=3/2	

 $\Delta(1930)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.1 to 0.2 OUR ESTIMATE				
0.18 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.14 ± 0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.04 ± 0.03	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
••• We do not use the following data for averages, fits, limits, etc. •••				
0.11	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	

$$(\Gamma_1/\Gamma_{\text{total}})^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow \Delta(1930) \rightarrow \Sigma K \quad (\Gamma_1/\Gamma_2)^{1/2}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
< 0.015	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.031	LIVANOS 80	DPWA	$\pi p \rightarrow \Sigma K$
0.018 to 0.035	² DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$

$$(\Gamma_1/\Gamma_{\text{total}})^{1/2}/\Gamma_{\text{total}} \text{ in } N\pi \rightarrow \Delta(1930) \rightarrow N\pi\pi \quad (\Gamma_1/\Gamma_3)^{1/2}/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
not seen	LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1930)$ PHOTON DECAY AMPLITUDES $\Delta(1930) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.015 ± 0.017 OUR ESTIMATE			
-0.019 ± 0.001	LI 93	IPWA	$\gamma N \rightarrow \pi N$
0.009 ± 0.009	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.030 ± 0.047	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
-0.062 ± 0.064	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1930) \rightarrow N\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE ($\text{GeV}^{-1/2}$)	DOCUMENT ID	TECN	COMMENT
-0.010 ± 0.022 OUR ESTIMATE			
0.009 ± 0.001	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.025 ± 0.011	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.033 ± 0.060	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
+0.019 ± 0.054	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1930)$ FOOTNOTES

¹ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

² The range given for DEANS 75 is from the four best solutions.

 $\Delta(1930)$ REFERENCES

For early references, see Physics Letters 111B 70 (1982).

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	(KENT) IJP
Also 84	PR D30 904	(VPI)
ARNDT 91	PR D43 2131	+Li, Roper, Workman, Ford
CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+
PDG 82	PL 111B	Roos, Porter, Aguilier-Benitez+
AWAJI 81	Bonn Conf. 352	+Kajikawa (NAGO)
Also 82	NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+
CHEW 80	Toronto Conf. 123	(LBL) IJP
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick (CMU, LBL) IJP
Also 79	PR D20 2839	+Baton, Coutures, Kochowski, Neveu (CMU, LBL) IJP
LIVANOS 80	Toronto Conf. 35	(SACL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen (KARLT) IJP
Also 80	Toronto Conf. 3	Koch (KARLT) IJP
BARBOUR 78	NP B141 253	+Crawford, Parsons (GLAS)
DEANS 75	NP B96 90	+Mitchell, Montgomery+ (SFLA, ALAH) IJP
LONGACRE 75	PL 55B 415	+Rosenfeld, Lasinski, Smdaja+ (LBL, SLAC) IJP

 $\Delta(1940) D_{33}$

$$I(J^P) = \frac{3}{2}(\frac{3}{2}^-) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

 $\Delta(1940)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1940 OUR ESTIMATE			
2057 ± 110	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
2058.1 ± 34.5	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
1940 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1940)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
460 ± 320	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
198.4 ± 45.5	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
200 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1940)$ POLE POSITION

REAL PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1915 or 1926	¹ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

Baryon Full Listings

 $\Delta(1940)$, $\Delta(1950)$ **-2xIMAGINARY PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200 ± 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
190 or 186	¹ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1940)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
8 ± 3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
135 ± 45	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(1940)$ DECAY MODES

Mode	
Γ_1	$N\pi$
Γ_2	ΣK
Γ_3	$N\pi\pi$
Γ_4	$\Delta(1232)\pi$, S-wave
Γ_5	$\Delta(1232)\pi$, D-wave
Γ_6	$N\rho$, S=3/2, S-wave
Γ_7	$N\gamma$, helicity=1/2
Γ_8	$N\gamma$, helicity=3/2

 $\Delta(1940)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.18 ± 0.12	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.18	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	
0.05 ± 0.02	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1940) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
< 0.015	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1940) \rightarrow \Delta(1232)\pi$, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
+ 0.11 ± 0.10	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1940) \rightarrow \Delta(1232)\pi$, D-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
+ 0.27 ± 0.16	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\pi \rightarrow \Delta(1940) \rightarrow N\rho$, S=3/2, S-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
+ 0.25 ± 0.10	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

 $\Delta(1940)$ PHOTON DECAY AMPLITUDES **$\Delta(1940) \rightarrow N\gamma$, helicity-1/2 amplitude $A_{1/2}$**

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
- 0.036 ± 0.058	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1940) \rightarrow N\gamma$, helicity-3/2 amplitude $A_{3/2}$

VALUE (GeV ^{-1/2})	DOCUMENT ID	TECN	COMMENT
- 0.031 ± 0.012	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$

 $\Delta(1940)$ FOOTNOTES

¹ LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.

 $\Delta(1940)$ REFERENCES

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	84 PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
AWAJI 81	Bonn Conf. 352	+Kajikawa	(NAGO)
Also	82 NP B197 365	Fujii, Hayashii, Iwata, Kajikawa+	(NAGO)
CHEW 80	Toronto Conf. 123		(LBL) IJP
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79 PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
LONGACRE 78	PR D17 1795	+Lasinski, Rosenfeld, Smadja+	(LBL, SLAC)

 $\Delta(1950) F_{37}$

$$I(J^P) = \frac{3}{2}(\frac{7}{2}^+) \text{ Status: } ***$$

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters **111B** (1982).

 $\Delta(1950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1940 to 1960 (≈ 1950) OUR ESTIMATE			
1945 ± 2	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
1950 ± 15	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
1913 ± 8	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1940	LI 93	IPWA	$\gamma N \rightarrow \pi N$
1925 ± 20	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
1855.0 ^{+11.0} _{-10.0}	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
1902	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
1912	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
1925	¹ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
290 to 350 (≈ 300) OUR ESTIMATE			
300 ± 7	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$
340 ± 50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
224 ± 10	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
306	LI 93	IPWA	$\gamma N \rightarrow \pi N$
330 ± 40	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
157.2 ^{+22.0} _{-19.0}	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
225	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
198	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
240	¹ LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1950)$ POLE POSITION**REAL PART**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1878	² HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
1884	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
1890 ± 15	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1858	ARNDT 85	DPWA	See ARNDT 91
1924 or 1924	³ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
230	² HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
238	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
260 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
238	ARNDT 85	DPWA	See ARNDT 91
258 or 258	³ LONGACRE 78	IPWA	$\pi N \rightarrow N\pi\pi$

 $\Delta(1950)$ ELASTIC POLE RESIDUE**MODULUS $|r|$**

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
47	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
61	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
50 ± 7	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE (°)	DOCUMENT ID	TECN	COMMENT
- 32	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
- 23	ARNDT 91	DPWA	$\pi N \rightarrow \pi N$ Soln SM90
- 33 ± 8	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

See key on page 1343

Baryon Full Listings

$\Delta(1950), \Delta(2000)$

$\Delta(1950)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	35–40 %
Γ_2 ΣK	
Γ_3 $N\pi\pi$	
Γ_4 $\Delta\pi$	20–30 %
Γ_5 $\Delta(1232)\pi, F\text{-wave}$	
Γ_6 $\Delta(1232)\pi, H\text{-wave}$	
Γ_7 $N\rho$	<10 %
Γ_8 $N\rho, S=1/2, F\text{-wave}$	
Γ_9 $N\rho, S=3/2, F\text{-wave}$	
Γ_{10} $N\gamma$	0.10–0.15 %
Γ_{11} $N\gamma, \text{helicity}=1/2$	
Γ_{12} $N\gamma, \text{helicity}=3/2$	

$\Delta(1950)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.35 to 0.4 OUR ESTIMATE				
0.38 ± 0.01	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.39 ± 0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.38 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.44	CHEW 80	BPWA	$\pi^+\rho \rightarrow \pi^+\rho$	
$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1950) \rightarrow \Sigma K$				$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.053 ± 0.005	CANDLIN 84	DPWA	$\pi^+\rho \rightarrow \Sigma^+ K^+$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.022 to 0.040	4 DEANS 75	DPWA	$\pi N \rightarrow \Sigma K$	

Note: Signs of couplings from $\pi N \rightarrow N\pi\pi$ analyses were changed in the 1986 edition to agree with the baryon-first convention; the overall phase ambiguity is resolved by choosing a negative sign for the $\Delta(1620) S_{31}$ coupling to $\Delta(1232)\pi$.

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1950) \rightarrow \Delta(1232)\pi, F\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
+0.28 to +0.32 OUR ESTIMATE				
+0.27 ± 0.02	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$	
+0.32	1 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.21	5 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	
0.38	6 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(1950) \rightarrow N\rho, S=3/2, F\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
+0.24	1 LONGACRE 75	IPWA	$\pi N \rightarrow N\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24	7 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	
0.43	8 NOVOSELLER 78	IPWA	$\pi N \rightarrow N\pi\pi$	

$\Delta(1950)$ PHOTON DECAY AMPLITUDES

$\Delta(1950) \rightarrow N\gamma, \text{helicity-1/2 amplitude } A_{1/2}$	DOCUMENT ID	TECN	COMMENT
VALUE (GeV^{-1/2})			
-0.086 ± 0.017 OUR ESTIMATE			
-0.102 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.068 ± 0.007	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.091 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.083 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.067 ± 0.014	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.058 ± 0.013	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$
$\Delta(1950) \rightarrow N\gamma, \text{helicity-3/2 amplitude } A_{3/2}$			
VALUE (GeV^{-1/2})			
-0.101 ± 0.014 OUR ESTIMATE			
-0.115 ± 0.003	LI 93	IPWA	$\gamma N \rightarrow \pi N$
-0.094 ± 0.016	AWAJI 81	DPWA	$\gamma N \rightarrow \pi N$
-0.101 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 1)
-0.100 ± 0.005	ARAI 80	DPWA	$\gamma N \rightarrow \pi N$ (fit 2)
-0.082 ± 0.017	CRAWFORD 80	DPWA	$\gamma N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.075 ± 0.020	BARBOUR 78	DPWA	$\gamma N \rightarrow \pi N$

$\Delta(1950)$ FOOTNOTES

- From method II of LONGACRE 75: eyeball fits with Breit-Wigner circles to the T-matrix amplitudes.
- See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.
- LONGACRE 78 values are from a search for poles in the unitarized T-matrix. The first (second) value uses, in addition to $\pi N \rightarrow N\pi\pi$ data, elastic amplitudes from a Saclay (CERN) partial-wave analysis.
- The range given is from the four best solutions. DEANS 75 disagrees with $\pi^+\rho \rightarrow \Sigma^+ K^+$ data of WINNIK 77 around 1920 MeV.
- A Breit-Wigner fit to the HERNDON 75 IPWA; the phase is near -60° .
- A Breit-Wigner fit to the NOVOSELLER 78B IPWA; the phase is near -60° .
- A Breit-Wigner fit to the HERNDON 75 IPWA; the phase is near 120° .
- A Breit-Wigner fit to the NOVOSELLER 78B IPWA; the phase is near 120° .

$\Delta(1950)$ REFERENCES

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
LI 93	PR C47 2759	(VPI)
MANLEY 92	PR D45 4002	(KENT) IJP
Also 84	PR D30 904	(VPI)
ARNDT 91	PR D43 2131	(VPI, TELE) IJP
ARNDT 85	PR D32 1085	(VPI)
CANDLIN 84	NP B238 477	(EDIN, RAL, LOWC)
PDG 82	PL 1118	(HELS, CIT, CERN)
AWAJI 81	Bonn Conf. 352	(NAGO)
Also 82	NP B197 365	(NAGO)
ARAI 80	Toronto Conf. 93	(INUS)
Also 82	NP B194 251	(INUS)
CHEW 80	Toronto Conf. 123	(LBL) IJP
CRAWFORD 80	Toronto Conf. 107	(GLAS)
CUTKOSKY 80	Toronto Conf. 19	(CMU, LBL) IJP
Also 79	PR D20 2839	(CMU, LBL) IJP
HOEHLER 79	PDAT 12-1	(KARLT) IJP
Also 80	Toronto Conf. 3	(KARLT) IJP
BARBOUR 78	NP B141 253	(GLAS)
LONGACRE 78	PR D17 1795	(LBL, SLAC)
NOVOSELLER 78	NP B137 509	(CIT) IJP
NOVOSELLER 78B	NP B137 445	(CIT) IJP
WINNIK 77	NP B128 66	(HAIF) I
DEANS 75	NP B96 90	(SFLA, ALAH) IJP
HERNDON 75	PR D11 3183	(LBL, SLAC)
LONGACRE 75	PL 55B 415	(LBL, SLAC) IJP

$\Delta(2000) F_{35}$

$$I(J^P) = \frac{3}{2}(\frac{5}{2}^+) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

$\Delta(2000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2000 OUR ESTIMATE			
1752 ± 32	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
2200 ± 125	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(2000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
251 ± 93	MANLEY 92	IPWA	$\pi N \rightarrow \pi N$ & $N\pi\pi$
400 ± 125	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(2000)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
VALUE (MeV)			
2150 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
-2×IMAGINARY PART			
VALUE (MeV)			
350 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(2000)$ ELASTIC POLE RESIDUE

MODULUS $ r $	DOCUMENT ID	TECN	COMMENT
VALUE (MeV)			
16 ± 5	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
PHASE θ			
VALUE (°)			
150 ± 90	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(2000)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\pi$	35–40 %
Γ_2 $N\pi\pi$	
Γ_3 $\Delta(1232)\pi, P\text{-wave}$	
Γ_4 $\Delta(1232)\pi, F\text{-wave}$	
Γ_5 $N\rho, S=3/2, P\text{-wave}$	

Baryon Full Listings

 $\Delta(2000)$, $\Delta(2150)$, $\Delta(2200)$ $\Delta(2000)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.02±0.01	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	
0.07±0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(2000) \rightarrow \Delta(1232)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
+0.07±0.03	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(2000) \rightarrow \Delta(1232)\pi, F\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
+0.09±0.04	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(2000) \rightarrow N\rho, S=3/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
-0.06±0.01	MANLEY 92	IPWA	$\pi N \rightarrow \pi N \& N\pi\pi$	

 $\Delta(2000)$ REFERENCES

MANLEY 92	PR D45 4002	+Saleski	(KENT) IJP
Also	PR D30 904	Manley, Arndt, Goradia, Teplitz	(VPI)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL)
Also	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)

 $\Delta(2150) S_{31}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

 $\Delta(2150)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2150 OUR ESTIMATE			
2047.4±27.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2203.2±8.4	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
2150 ±100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2150)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
121.6±62.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
120.5±45.0	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$
200 ±100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2150)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
2140±80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

-2×IMAGINARY PART	DOCUMENT ID	TECN	COMMENT
200±80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2150)$ ELASTIC POLE RESIDUE

MODULUS r	DOCUMENT ID	TECN	COMMENT
7±2	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ	DOCUMENT ID	TECN	COMMENT
-60±90	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2150)$ DECAY MODES

Mode	Γ_1	Γ_2
$N\pi$		
ΣK		

 $\Delta(2150)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.41	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	
0.37	¹ CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	
0.08±0.02	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(2150) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
<0.03	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	

 $\Delta(2150)$ FOOTNOTES

¹ CHEW 80 reports two S_{31} resonances in this mass region. Problems with this analysis are discussed in section 2.1.11 of HOEHLER 83.

 $\Delta(2150)$ REFERENCES

CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
HOEHLER 83	Landolt-Boernstein 1/9B2		(KARLT)
CHEW 80	Toronto Conf. 123		(LBL) IJP
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)

 $\Delta(2200) G_{37}$

$$I(J^P) = \frac{3}{2}(\frac{1}{2}^-) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

The various analyses are not in good agreement.

 $\Delta(2200)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2200 OUR ESTIMATE			
2200±80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2215±60	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2280±80	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
2280±40	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$

 $\Delta(2200)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
450±100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
400±100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
400±150	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$
••• We do not use the following data for averages, fits, limits, etc. •••			
400±50	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$

 $\Delta(2200)$ POLE POSITION

REAL PART	DOCUMENT ID	TECN	COMMENT
2100±50	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

-2×IMAGINARY PART	DOCUMENT ID	TECN	COMMENT
340±80	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2200)$ ELASTIC POLE RESIDUE

MODULUS r	DOCUMENT ID	TECN	COMMENT
8±3	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ	DOCUMENT ID	TECN	COMMENT
-70±40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2200)$ DECAY MODES

Mode	Γ_1	Γ_2
$N\pi$		
ΣK		

 $\Delta(2200)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06±0.02	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.05±0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.09±0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\pi \rightarrow \Delta(2200) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.014±0.005	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	

 $\Delta(2200)$ REFERENCES

CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL) IJP
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also	ANP 136 1	Hendry	(IND)

See key on page 1343

Baryon Full Listings
 $\Delta(2300)$, $\Delta(2350)$ $\Delta(2300) H_{39}$

$I(J^P) = \frac{3}{2}(\frac{9}{2}^+)$ Status: * *

OMITTED FROM SUMMARY TABLE

 $\Delta(2300)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2300 OUR ESTIMATE			
2204.5 ± 3.4	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
2400 ± 125	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
2217 ± 80	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
2450 ± 100	HENDRY	78	MPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2400	CANDLIN	84	DPWA $\pi^+ p \rightarrow \Sigma^+ K^+$

 $\Delta(2300)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
32.3 ± 1.0	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$
425 ± 150	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
300 ± 100	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$
500 ± 200	HENDRY	78	MPWA $\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
200	CANDLIN	84	DPWA $\pi^+ p \rightarrow \Sigma^+ K^+$

 $\Delta(2300)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2370 ± 80	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

-2xIMAGINARY PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
420 ± 160	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $\Delta(2300)$ ELASTIC POLE RESIDUE

MODULUS $ r $ VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
10 ± 4	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

PHASE θ VALUE (°)	DOCUMENT ID	TECN	COMMENT
-20 ± 30	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $\Delta(2300)$ DECAY MODES

Mode
Γ_1 $N\pi$
Γ_2 ΣK

 $\Delta(2300)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.05	CHEW	80	BPWA $\pi^+ p \rightarrow \pi^+ p$	
0.06 ± 0.02	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.03 ± 0.02	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	
0.08 ± 0.02	HENDRY	78	MPWA $\pi N \rightarrow \pi N$	

$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow \Delta(2300) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma$
-0.017	CANDLIN	84	DPWA $\pi^+ p \rightarrow \Sigma^+ K^+$	

 $\Delta(2300)$ REFERENCES

CANDLIN	84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
CHEW	80	Toronto Conf. 123		(LBL) IJP
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY	78	PRL 41 222		(IND, LBL) IJP
Also	81	ANP 136 1	Hendry	(IND)

 $\Delta(2350) D_{35}$

$I(J^P) = \frac{3}{2}(\frac{5}{2}^-)$ Status: *

OMITTED FROM SUMMARY TABLE

 $\Delta(2350)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2350 OUR ESTIMATE			
2171 ± 18	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
2400 ± 125	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
2305 ± 26	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

 $\Delta(2350)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
264 ± 51	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$
400 ± 150	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$
300 ± 70	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$

 $\Delta(2350)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2400 ± 125	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

-2xIMAGINARY PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
400 ± 150	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $\Delta(2350)$ ELASTIC POLE RESIDUE

MODULUS $ r $ VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
15 ± 8	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

PHASE θ VALUE (°)	DOCUMENT ID	TECN	COMMENT
-70 ± 70	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$

 $\Delta(2350)$ DECAY MODES

Mode
Γ_1 $N\pi$
Γ_2 ΣK

 $\Delta(2350)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.020 ± 0.003	MANLEY	92	IPWA $\pi N \rightarrow \pi N$ & $N\pi\pi$	
0.20 ± 0.10	CUTKOSKY	80	IPWA $\pi N \rightarrow \pi N$	
0.04 ± 0.02	HOEHLER	79	IPWA $\pi N \rightarrow \pi N$	

$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow \Delta(2350) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma$
<0.015	CANDLIN	84	DPWA $\pi^+ p \rightarrow \Sigma^+ K^+$	

 $\Delta(2350)$ REFERENCES

MANLEY	92	PR D45 4002	+Saleski	(KENT) IJP
Also	84	PR D30 904	Manley, Arndt, Goradia, Tepitz	(VPI)
CANDLIN	84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
CUTKOSKY	80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also	79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
HOEHLER	79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also	80	Toronto Conf. 3	Koch	(KARLT) IJP
	80	Toronto Conf. 3	Koch	(KARLT) IJP

Baryon Full Listings

 $\Delta(2390)$, $\Delta(2400)$ $\Delta(2390) F_{37}$ $I(J^P) = \frac{3}{2}(\frac{7}{2}^+)$ Status: *

OMITTED FROM SUMMARY TABLE

 $\Delta(2390)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2390 OUR ESTIMATE			
2350 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2425 \pm 60	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2390)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
300 \pm 80	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2390)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2350 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
260 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2390)$ ELASTIC POLE RESIDUEMODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
12 \pm 6	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-90 \pm 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2390)$ DECAY MODES

Mode

Γ_1	$N\pi$
Γ_2	ΣK

 $\Delta(2390)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$ VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
0.08 \pm 0.04	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.07 \pm 0.04	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow \Delta(2390) \rightarrow \Sigma K$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma$
<0.015	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	

 $\Delta(2390)$ REFERENCES

CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP

 $\Delta(2400) G_{39}$ $I(J^P) = \frac{3}{2}(\frac{9}{2}^-)$ Status: **

OMITTED FROM SUMMARY TABLE

 $\Delta(2400)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2400 OUR ESTIMATE			
2300 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2468 \pm 50	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2200 \pm 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

 $\Delta(2400)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
330 \pm 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
480 \pm 100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
450 \pm 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

 $\Delta(2400)$ POLE POSITION

REAL PART VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2260 \pm 60	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

-2xIMAGINARY PART

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
320 \pm 160	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2400)$ ELASTIC POLE RESIDUEMODULUS $|r|$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
8 \pm 4	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ

VALUE ($^\circ$)	DOCUMENT ID	TECN	COMMENT
-25 \pm 15	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

 $\Delta(2400)$ DECAY MODES

Mode

Γ_1	$N\pi$
Γ_2	ΣK

 $\Delta(2400)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$ VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ_2
0.05 \pm 0.02	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.06 \pm 0.03	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.10 \pm 0.03	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma_{total} \ln N\pi \rightarrow \Delta(2400) \rightarrow \Sigma K$ VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma$
<0.015	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	

 $\Delta(2400)$ REFERENCES

CANDLIN 84	NP B238 477	+Lowe, Peach, Scotland+	(EDIN, RAL, LOWC)
CUTKOSKY 80	Toronto Conf. 19	+Forsyth, Babcock, Kelly, Hendrick	(CMU, LBL) IJP
Also 79	PR D20 2839	Cutkosky, Forsyth, Hendrick, Kelly	(CMU, LBL)
HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also 81	ANP 136 1	Hendry	(IND)

See key on page 1343

Baryon Full Listings
 $\Delta(2420)$, $\Delta(2750)$, $\Delta(2950)$

$\Delta(2420) H_{3,11}$

$I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$ Status: * * * *

Most of the results published before 1975 are now obsolete and have been omitted. They may be found in our 1982 edition, Physics Letters 111B (1982).

$\Delta(2420)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2300 to 2500 (≈ 2420) OUR ESTIMATE			
2400 ± 125	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
2416 ± 17	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2400 ± 60	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2400	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
2358.0 ± 9.0	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

$\Delta(2420)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 to 500 (≈ 400) OUR ESTIMATE			
450 ± 150	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
340 ± 28	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
460 ± 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
400	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$
202.2 ± 45.0	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$

$\Delta(2420)$ POLE POSITION

REAL PART			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2300	¹ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
2360 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$
-2xIMAGINARY PART			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
620	¹ HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
420 ± 100	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(2420)$ ELASTIC POLE RESIDUE

MODULUS r			
VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
39	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
18 ± 6	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

PHASE θ			
VALUE (°)	DOCUMENT ID	TECN	COMMENT
-60	HOEHLER 93	ARGD	$\pi N \rightarrow \pi N$
-30 ± 40	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$

$\Delta(2420)$ DECAY MODES

The following branching fractions are our estimates, not fits or averages.

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N\pi$	5-15 %
$\Gamma_2 \Sigma K$	

$\Delta(2420)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.05 to 0.15 OUR ESTIMATE				
0.08 ± 0.03	CUTKOSKY 80	IPWA	$\pi N \rightarrow \pi N$	
0.08 ± 0.015	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.11 ± 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.22	CHEW 80	BPWA	$\pi^+ p \rightarrow \pi^+ p$	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{total} \ln N\pi \rightarrow \Delta(2420) \rightarrow \Sigma K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
-0.016	CANDLIN 84	DPWA	$\pi^+ p \rightarrow \Sigma^+ K^+$	

$\Delta(2420)$ FOOTNOTES

¹ See HOEHLER 93 and HOEHLER 94 for a detailed discussion of the evidence for and the pole parameters of N and Δ resonances as determined from Argand diagrams of πN elastic partial-wave amplitudes and from plots of the speeds with which the amplitudes traverse the diagrams.

$\Delta(2420)$ REFERENCES

HOEHLER 94	πN Newsletter 10 (to be pub.)	(KARL)
HOEHLER 93	πN Newsletter 9 1	(KARL)
CANDLIN 84	NP B238 477	(EDIN, RAL, LOWC)
PDG 82	PL 1118	(HELS, CIT, CERN)
CHEW 80	Toronto Conf. 123	(LBL) IJP
CUTKOSKY 80	Toronto Conf. 19	(CMU, LBL) IJP
Also 79	PR D20 2839	(CMU, LBL)
HOEHLER 79	PDAT 12-1	+Forsyth, Babcock, Kelly, Hendrick Cutkosky, Forsyth, Hendrick, Kelly
Also 80	Toronto Conf. 3	+Kaiser, Koch, Pietarinen
HENDRY 78	PRL 41 222	Koch
Also 81	ANP 136 1	Hendry

$\Delta(2750) I_{3,13}$

$I(J^P) = \frac{3}{2}(\frac{1}{2}^-)$ Status: * *

OMITTED FROM SUMMARY TABLE

$\Delta(2750)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2750 OUR ESTIMATE			
2794 ± 80	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2650 ± 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

$\Delta(2750)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
350 ± 100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
500 ± 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

$\Delta(2750)$ DECAY MODES

Mode	Γ_1
$N\pi$	

$\Delta(2750)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.04 ± 0.015	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.05 ± 0.01	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$\Delta(2750)$ REFERENCES

HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also 81	ANP 136 1	Hendry	(IND)

$\Delta(2950) K_{3,15}$

$I(J^P) = \frac{3}{2}(\frac{1}{2}^+)$ Status: * *

OMITTED FROM SUMMARY TABLE

$\Delta(2950)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2950 OUR ESTIMATE			
2990 ± 100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
2850 ± 100	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

$\Delta(2950)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
330 ± 100	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$
700 ± 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$

$\Delta(2950)$ DECAY MODES

Mode	Γ_1
$N\pi$	

$\Delta(2950)$ BRANCHING RATIOS

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.04 ± 0.02	HOEHLER 79	IPWA	$\pi N \rightarrow \pi N$	
0.03 ± 0.01	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$	

$\Delta(2950)$ REFERENCES

HOEHLER 79	PDAT 12-1	+Kaiser, Koch, Pietarinen	(KARLT) IJP
Also 80	Toronto Conf. 3	Koch	(KARLT) IJP
HENDRY 78	PRL 41 222		(IND, LBL) IJP
Also 81	ANP 136 1	Hendry	(IND)

Baryon Full Listings

 $\Delta(\sim 3000), \Lambda$ **$\Delta(\sim 3000)$ Region
Partial-Wave Analyses**

OMITTED FROM SUMMARY TABLE

We list here miscellaneous high-mass candidates for isospin-3/2 resonances found in partial-wave analyses.

Our 1982 edition also had a $\Delta(2850)$ and a $\Delta(3230)$. The evidence for them was deduced from total cross-section and 180° elastic cross-section measurements. The $\Delta(2850)$ has been resolved into the $\Delta(2750) I_{3,13}$ and $\Delta(2950) K_{3,15}$. The $\Delta(3230)$ is perhaps related to the $K_{3,13}$ of HENDRY 78 and to the $L_{3,17}$ of KOCH 80.

 $\Delta(\sim 3000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 3000 OUR ESTIMATE			
3300	¹ KOCH 80	IPWA	$\pi N \rightarrow \pi N$ $L_{3,17}$ wave
3500	¹ KOCH 80	IPWA	$\pi N \rightarrow \pi N$ $M_{3,19}$ wave
2850 \pm 150	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $I_{3,11}$ wave
3200 \pm 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $K_{3,13}$ wave
3300 \pm 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $L_{3,17}$ wave
3700 \pm 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $M_{3,19}$ wave
4100 \pm 300	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $N_{3,21}$ wave

 $\Delta(\sim 3000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
700 \pm 200	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $I_{3,11}$ wave
1000 \pm 300	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $K_{3,13}$ wave
1100 \pm 300	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $L_{3,17}$ wave
1300 \pm 400	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $M_{3,19}$ wave
1600 \pm 500	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $N_{3,21}$ wave

 $\Delta(\sim 3000)$ DECAY MODES

Mode

 Γ_1 $N\pi$ **$\Delta(\sim 3000)$ BRANCHING RATIOS**

$\Gamma(N\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 \pm 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $I_{3,11}$ wave	
0.045 \pm 0.02	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $K_{3,13}$ wave	
0.03 \pm 0.01	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $L_{3,17}$ wave	
0.025 \pm 0.01	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $M_{3,19}$ wave	
0.018 \pm 0.01	HENDRY 78	MPWA	$\pi N \rightarrow \pi N$ $N_{3,21}$ wave	

 $\Delta(\sim 3000)$ FOOTNOTES

¹In addition, KOCH 80 reports some evidence for an S_{31} $\Delta(2700)$ and a P_{33} $\Delta(2800)$.

 $\Delta(\sim 3000)$ REFERENCES

KOCH 80	Toronto Conf. 3	(KARLT) IJP
HENDRY 78	PRL 41 222	(IND, LBL) IJP
Also 81	ANP 136 1	(IND)
	Hendry	

 Λ BARYONS
($S = -1, I = 0$) $\Lambda^0 = uds$  $I(J^P) = 0(\frac{1}{2}^+)$ Status: * * * *

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1986 edition (Physics Letters 170B) or in earlier editions.

 Λ MASS

The fit uses $\Lambda, \Sigma^+, \Sigma^0, \Sigma^-$ mass and mass-difference measurements.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1115.684 ± 0.006 OUR FIT				
1115.683 ± 0.006 OUR AVERAGE				
1115.678 \pm 0.006 \pm 0.006	20k	HARTOUNI 94	SPEC	pp 27.5 GeV/c
1115.690 \pm 0.008 \pm 0.006	18k	¹ HARTOUNI 94	SPEC	pp 27.5 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1115.59 \pm 0.08	935	HYMAN 72	HEBC	
1115.39 \pm 0.12	195	MAYEUR 67	EMUL	
1115.6 \pm 0.4		LONDON 66	HBC	
1115.65 \pm 0.07	488	² SCHMIDT 65	HBC	
1115.44 \pm 0.12		³ BHOWMIK 63	RVUE	

¹We assume CPT invariance: this is the $\bar{\Lambda}$ mass as measured by HARTOUNI 94. See below for the fractional mass difference, testing CPT.

²The SCHMIDT 65 masses have been reevaluated using our April 1973 proton and K^\pm and π^\pm masses. P. Schmidt, private communication (1974).

³The mass has been raised 35 keV to take into account a 46 keV increase in the proton mass and an 11 keV decrease in the π^\pm mass (note added Reviews of Modern Physics 39 1 (1967)).

 $(m_\Lambda - m_{\bar{\Lambda}}) / m_\Lambda$


A test of CPT invariance.

VALUE (units 10^{-5})	DOCUMENT ID	TECN	COMMENT
-1.0 ± 0.9 OUR AVERAGE			
-1.08 ± 0.90	HARTOUNI 94	SPEC	pp 27.5 GeV/c
-26 ± 13	BADIER 67	HBC	2.4 GeV/c $p\bar{p}$
4.5 ± 5.4	CHIEN 66	HBC	6.9 GeV/c $p\bar{p}$

 Λ MEAN LIFE

Measurements with an error $\geq 0.1 \times 10^{-10}$ s have been omitted altogether, and only the latest high-statistics measurements are used for the average.

VALUE (10^{-10} s)	EVTs	DOCUMENT ID	TECN	COMMENT
2.632 ± 0.020 OUR AVERAGE				Error includes scale factor of 1.6. See the ideogram below.
2.69 \pm 0.03	53k	ZECH 77	SPEC	Neutral hyperon beam
2.611 \pm 0.020	34k	CLAYTON 75	HBC	0.96–1.4 GeV/c $K^- p$
2.626 \pm 0.020	36k	POULARD 73	HBC	0.4–2.3 GeV/c $K^- p$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2.69 \pm 0.05	6582	ALTHOFF 73B	OSPK	$\pi^+ n \rightarrow \Lambda K^+$
2.54 \pm 0.04	4572	BALTAY 71B	HBC	$K^- p$ at rest
2.535 \pm 0.035	8342	GRIMM 68	HBC	
2.47 \pm 0.08	2600	HEPP 68	HBC	
2.35 \pm 0.09	916	BURAN 66	HLBC	
2.452 $^{+0.056}_{-0.054}$	2213	ENGELMANN 66	HBC	
2.59 \pm 0.09	794	HUBBARD 64	HBC	
2.59 \pm 0.07	1378	SCHWARTZ 64	HBC	
2.36 \pm 0.06	2239	BLOCK 63	HEBC	



$$\frac{(\tau_\Lambda - \tau_\Lambda)}{\tau_{\text{average}}}$$

A test of CPT invariance.

VALUE	DOCUMENT ID	TECN	COMMENT
0.044 ± 0.005	BADIER	67 HBC	2.4 GeV/c $\bar{p}p$


NOTE ON BARYON MAGNETIC MOMENTS

The figure shows the measured magnetic moments of the stable baryons. It also shows the predictions of the simplest quark model, using the measured p , n , and Λ moments as input. In this model, the moments are [1]

$$\begin{aligned} \mu_p &= (4\mu_u - \mu_d)/3 & \mu_n &= (4\mu_d - \mu_u)/3 \\ \mu_{\Sigma^+} &= (4\mu_u - \mu_s)/3 & \mu_{\Sigma^-} &= (4\mu_d - \mu_s)/3 \\ \mu_{\Xi^0} &= (4\mu_s - \mu_u)/3 & \mu_{\Xi^-} &= (4\mu_s - \mu_d)/3 \\ \mu_\Lambda &= \mu_s & \mu_{\Sigma^0} &= (2\mu_u + 2\mu_d - \mu_s)/3 \\ & & \mu_{\Sigma^-} &= 3\mu_s \end{aligned}$$

and the $\Sigma^0 \rightarrow \Lambda$ transition moment is

$$\mu_{\Sigma^0 \Lambda} = (\mu_d - \mu_u)/\sqrt{3}.$$



The quark moments that result from this model are $\mu_u = +1.852 \mu_N$, $\mu_d = -0.972 \mu_N$, and $\mu_s = -0.613 \mu_N$. The corresponding effective quark masses, taking the quarks to be Dirac point particles, where $\mu = q\hbar/2m$, are 338, 322, and 510 MeV. As the figure shows, the model gives a good first approximation to the experimental moments. For efforts to make a better model, we refer to the literature [2].

References

- See, for example, D.H. Perkins, *Introduction to High Energy Physics* (Addison-Wesley, Reading, MA, 1987), or D. Griffiths, *Introduction to Elementary Particles* (Harper & Row, New York, 1987).
- See, for example, J. Franklin, Phys. Rev. **D29**, 2648 (1984); H.J. Lipkin, Nucl. Phys. **B241**, 477 (1984); K. Suzuki, H. Kumagai, and Y. Tanaka, Europhys. Lett. **2**, 109 (1986); S.K. Gupta and S.B. Khadkikar, Phys. Rev. **D36**, 307 (1987); M.I. Krivoruchenko, Sov. J. Nucl. Phys. **45**, 109 (1987); L. Brekke and J.L. Rosner, Comm. Nucl. Part. Phys. **18**, 83 (1988); K.-T. Chao, Phys. Rev. **D41**, 920 (1990) and references cited therein Also, see references cited in discussions of results in the experimental papers..

Λ MAGNETIC MOMENT

See the Note on Baryon Magnetic Moments above. Measurements with an error $\geq 0.15 \mu_N$ have been omitted.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.613 ± 0.004	OUR AVERAGE			
-0.606 ± 0.015	200K	COX	81	SPEC
-0.6138 ± 0.0047	3M	SCHACHIN...	78	SPEC
-0.59 ± 0.07	350K	HELLER	77	SPEC
-0.57 ± 0.05	1.2M	BUNCE	76	SPEC
-0.66 ± 0.07	1300	DAHL-JENSEN71	EMUL	200 kG field

Λ ELECTRIC DIPOLE MOMENT

A nonzero value is forbidden by both T invariance and P invariance.

VALUE (10^{-16} e-cm)	CL%	DOCUMENT ID	TECN
< 1.5	95	⁴ PONDROM 81	SPEC

• • • We do not use the following data for averages, fits, limits, etc. • • •

<100	95	⁵ BARONI 71	EMUL
<500	95	GIBSON 66	EMUL

⁴PONDROM 81 measures $(-3.0 \pm 7.4) \times 10^{-17}$ e-cm.
⁵BARONI 71 measures $(-5.9 \pm 2.9) \times 10^{-15}$ e-cm.

Λ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $p\pi^-$	(63.9 ± 0.5) %
Γ_2 $n\pi^0$	(35.8 ± 0.5) %
Γ_3 $n\gamma$	(1.75 ± 0.15) × 10 ⁻³
Γ_4 $p\pi^- \gamma$	[a] (8.4 ± 1.4) × 10 ⁻⁴
Γ_5 $p e^- \bar{\nu}_e$	(8.32 ± 0.14) × 10 ⁻⁴
Γ_6 $p \mu^- \bar{\nu}_\mu$	(1.57 ± 0.35) × 10 ⁻⁴

[a] See the Full Listings below for the pion momentum range used in this measurement.

Baryon Full Listings

Λ

CONSTRAINED FIT INFORMATION

An overall fit to 5 branching ratios uses 20 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 10.5$ for 16 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100			
x_3	-2	-1		
x_5	46	-46	-1	
x_6	0	0	0	0
	x_1	x_2	x_3	x_5

Λ BRANCHING RATIOS

$\Gamma(p\pi^-) / \Gamma(N\pi)$					$\Gamma_1 / (\Gamma_1 + \Gamma_2)$
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
0.641 ± 0.006 OUR FIT					
0.640 ± 0.005 OUR AVERAGE					
0.646 ± 0.008	4572	BALTAY	71B HBC	$K^- p$ at rest	
0.635 ± 0.007	6736	DOYLE	69 HBC	$\pi^- p \rightarrow \Lambda K^0$	
0.643 ± 0.016	903	HUMPHREY	62 HBC		
0.624 ± 0.030		CRAWFORD	59B HBC	$\pi^- p \rightarrow \Lambda K^0$	

$\Gamma(n\pi^0) / \Gamma(N\pi)$					$\Gamma_2 / (\Gamma_1 + \Gamma_2)$
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	
0.359 ± 0.006 OUR FIT					
0.310 ± 0.028 OUR AVERAGE					
0.35 ± 0.05		BROWN	63 HLBC		
0.291 ± 0.034	75	CHRETIEN	63 HLBC		

$\Gamma(n\gamma) / \Gamma_{\text{total}}$					Γ_3 / Γ
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	
1.75 ± 0.15 OUR FIT					
1.75 ± 0.15	1816	LARSON	93 SPEC	$K^- p$ at rest	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.78 ± 0.24 ^{+0.14} _{-0.16}	287	NOBLE	92 SPEC	See LARSON 93	

$\Gamma(n\gamma) / \Gamma(n\pi^0)$					Γ_3 / Γ_2
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
2.86 ± 0.74 ± 0.57	24	BIAGI	86 SPEC	SPS hyperon beam	

$\Gamma(p\pi^- \gamma) / \Gamma(p\pi^-)$					Γ_4 / Γ_1
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	
1.32 ± 0.22	72	BAGGETT	72C HBC	$\pi^- < 95 \text{ MeV}/c$	

$\Gamma(p\pi^- \nu_e) / \Gamma(p\pi^-)$					Γ_5 / Γ_1
VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	
1.301 ± 0.019 OUR FIT					
1.301 ± 0.019 OUR AVERAGE					
1.335 ± 0.056	7111	BOURQUIN	83 SPEC	SPS hyperon beam	
1.313 ± 0.024	10k	WISE	80 SPEC		
1.23 ± 0.11	544	LINDQUIST	77 SPEC	$\pi^- p \rightarrow K^0 \Lambda$	
1.27 ± 0.07	1089	KATZ	73 HBC		
1.31 ± 0.06	1078	ALTHOFF	71 OSPK		
1.17 ± 0.13	86	6 CANTER	71 HBC	$K^- p$ at rest	
1.20 ± 0.12	143	7 MALONEY	69 HBC		
1.17 ± 0.18	120	7 BAGLIN	64 FBC	K^- freon 1.45 GeV/c	
1.23 ± 0.20	150	7 ELY	63 FBC		
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1.32 ± 0.15	218	6 LINDQUIST	71 OSPK	See LINDQUIST 77	

⁶ Changed by us from $\Gamma(p\pi^- \nu_e) / \Gamma(N\pi)$ assuming the authors used $\Gamma(p\pi^-) / \Gamma_{\text{total}} = 2/3$.

⁷ Changed by us from $\Gamma(p\pi^- \nu_e) / \Gamma(N\pi)$ because $\Gamma(p\pi^- \nu) / \Gamma(p\pi^-)$ is the directly measured quantity.

$\Gamma(p\pi^- \nu_\mu) / \Gamma(N\pi)$					$\Gamma_6 / (\Gamma_1 + \Gamma_2)$
VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT	
1.57 ± 0.35 OUR FIT					
1.57 ± 0.35 OUR AVERAGE					
1.4 ± 0.5	14	BAGGETT	72B HBC	$K^- p$ at rest	
2.4 ± 0.8	9	CANTER	71B HBC	$K^- p$ at rest	
1.3 ± 0.7	3	LIND	64 RVUE		
1.5 ± 1.2	2	RONNE	64 FBC		

Λ DECAY PARAMETERS

See the Note on Baryon Decay Parameters in the neutron Listings. Some early results have been omitted.

 α_- FOR $\Lambda \rightarrow p\pi^-$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
0.642 ± 0.013 OUR AVERAGE				
0.584 ± 0.046	8500	ASTBURY	75 SPEC	
0.649 ± 0.023	10325	CLELAND	72 OSPK	
0.67 ± 0.06	3520	DAUBER	69 HBC	From Ξ decay
0.645 ± 0.017	10130	OVERSETH	67 OSPK	Λ from $\pi^- p$
0.62 ± 0.07	1156	CRONIN	63 CNTR	Λ from $\pi^- p$

 ϕ ANGLE FOR $\Lambda \rightarrow p\pi^-$

VALUE (°)	EVTs	DOCUMENT ID	TECN	COMMENT
-6.5 ± 3.5 OUR AVERAGE				
-7.0 ± 4.5	10325	CLELAND	72 OSPK	Λ from $\pi^- p$
-8.0 ± 6.0	10130	OVERSETH	67 OSPK	Λ from $\pi^- p$
13.0 ± 17.0	1156	CRONIN	63 OSPK	Λ from $\pi^- p$

 $\alpha_0 / \alpha_- = \alpha(\Lambda \rightarrow n\pi^0) / \alpha(\Lambda \rightarrow p\pi^-)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
1.01 ± 0.07 OUR AVERAGE				
1.000 ± 0.068	4760	8 OLSEN	70 OSPK	$\pi^+ n \rightarrow \Lambda K^+$
1.10 ± 0.27		CORK	60 CNTR	

⁸ OLSEN 70 compares proton and neutron distributions from Λ decay.

 $[\alpha_-(\Lambda) + \alpha_+(\bar{\Lambda})] / [\alpha_-(\Lambda) - \alpha_+(\bar{\Lambda})]$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.03 ± 0.06 OUR AVERAGE				
+0.01 ± 0.10	770	TIXIER	88 DM2	$J/\psi \rightarrow \bar{\Lambda} \Lambda$
-0.07 ± 0.09	4063	BARNES	87 CNTR	$\bar{p} p \rightarrow \bar{\Lambda} \Lambda$ LEAR
-0.02 ± 0.14	10k	9 CHAUVAT	85 CNTR	$p\bar{p}, \bar{p}p$ ISR

⁹ CHAUVAT 85 actually gives $\alpha_+(\bar{\Lambda}) / \alpha_-(\Lambda) = -1.04 \pm 0.29$. Assumes polarization is same in $\bar{p} p \rightarrow \bar{\Lambda} \Lambda$ and $p\bar{p} \rightarrow \Lambda X$. Tests of this assumption, based on C-invariance and fragmentation, are satisfied by the data.

 g_A / g_V FOR $\Lambda \rightarrow p e^- \bar{\nu}_e$

Measurements with fewer than 500 events have been omitted. Where necessary, signs have been changed to agree with our conventions, which are given in the Note on Baryon Decay Parameters in the neutron Listings. The measurements all assume that the form factor $g_2 = 0$. See also the footnote on DWORKIN 90.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.718 ± 0.015 OUR AVERAGE				
-0.719 ± 0.016 ± 0.012	37k	10 DWORKIN	90 SPEC	$e\nu$ angular corr.
-0.70 ± 0.03	7111	BOURQUIN	83 SPEC	$\Xi \rightarrow \Lambda \pi^-$
-0.734 ± 0.031	10k	11 WISE	81 SPEC	$e\nu$ angular correl.
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.63 ± 0.06	817	ALTHOFF	73 OSPK	Polarized Λ

¹⁰ The tabulated result assumes the weak-magnetism coupling $w \equiv g_W(0) / g_V(0)$ to be 0.97, as given by the CVC hypothesis and as assumed by the other listed measurements. However, DWORKIN 90 measures w to be 0.15 ± 0.30, and then $g_A / g_V = -0.731 \pm 0.016$.

¹¹ This experiment measures only the absolute value of g_A / g_V .

Λ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1986 edition (Physics Letters **170B**) or in earlier editions.

HARTOUNI	94	PRL 72 1322	+Jensen, Kreisler+	(BNL E766 Collab.)
LARSON	93	PR D47 799	+Noble, Bassalleck+	(BNL-811 Collab.)
NOBLE	92	PRL 69 414	+ (BIRM, BOST, BRCO, BNL, CASE, BUDA, LANL+)	
DWORKIN	90	PR D41 780	+Cox, Dukes, Oversteth+	(MICH, WISC, RUTG, MINN)
TIXIER	88	PL B212 523	+Ajaltouni, Falvard, Jousset+	(DM2 Collab.)
BARNES	87	PL B199 147	+ (CMU, SACL, LANL, VIEN, FREIB, ILL, UPPS+)	(MASA, BNL)
BIAGI	86	ZPHY C30 201	+ (BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL)	
CHAUVAT	85	PL 163B 273	+Erhan, Hayes+	(CERN, CLER, UCLA, SACL)
BOURQUIN	83	ZPHY C21 1	+Brown+	(BRIS, GEVA, HEIDP, LAO, RL, STRB)
COX	81	PRL 46 877	+Dworkin+	(MICH, WISC, RUTG, MINN, BNL)
PONDROM	81	PR D23 814	+Handler, Sheaff, Cox+	(WISC, CERN, DORT, HEIDH)
WISE	81	PL 98B 123	+Jensen, Kreisler, Lomanno, Poster+	(MASA, BNL)
WISE	80	PL 91B 165	+Jensen, Kreisler, Lomanno, Poster+	(MASA, BNL)
SCHACHIN...	78	PRL 41 1348	+Schachinger, Bunce, Cox+	(MICH, RUTG, HEIDH)
HELLER	77	PL 88B 480	+Bunnell, Bunce, Dydak+	(MICH, WISC, HEIDH)
LINDQUIST	77	PR D16 2104	+Swallow, Sumner+	(EFI, OSU, ANL)
Also	76	JPG 2 L211	+Lindquist, Swallow+	(EFI, WUOL, OSU, ANL)
ZECH	77	NP B124 413	+Dydak, Navarra+	(SIEG, CERN, DORT, HEIDH)
BUNCE	76	PRL 36 1113	+Handler, Reiser, Martin+	(WISC, MICH, RUTG)
ASTBURY	75	NP B99 30	+Gallivan, Jafar+	(LOIC, CERN, ETH, SACL)
CLAYTON	75	NP B95 130	+Bacon, Butterworth, Waters+	(LOIC, RHEL)
ALTHOFF	73	NP B66 29	+Brown, Freytag, Heard, Heintze+	(CERN, HEID)
ALTHOFF	73	NP B66 29	+Brown, Freytag, Heard, Heintze+	(CERN, HEID)
KATZ	73	Maryland Thesis		(UMD)
POULARD	73	PL 46B 135	+Givernaud, Borg	(SACL)
BAGGETT	72B	ZPHY 252 362	+Baggett, Eisele, Filthuth, Frehse+	(HEID)
BAGGETT	72C	PL 42B 379	+Baggett, Eisele, Filthuth, Frehse, Hepp+	(HEID)
CLELAND	72	NP B40 221	+Conforto, Eaton, Gerber+	(CERN, GEVA, LUND)
HYMAN	72	PR D5 1063	+Bunnell, Derrick, Fields, Katz+	(ANL, CMU)
ALTHOFF	71	PL 37B 531	+Brown, Freytag, Heard, Heintze+	(CERN, HEID)
BALTAY	71B	PR D4 670	+Bridgewater, Cooper, Habibi+	(COLU, BING)
BARONI	71	LNC 2 1256	+Petrera, Romano	(ROMA)
CANTER	71	PRL 26 868	+Cole, Lee-Franzini, Lovelless+	(STON, COLU)
CANTER	71B	PRL 27 59	+Cole, Lee-Franzini, Lovelless+	(STON, COLU)

DAHL-JENSEN	71	NC 3A 1	+	(CERN, ANKA, LAUS, MPIM, ROMA)
LINQUIST	71	PRL 27 612	+Summer+	(EFI, WUSL, OSU, ANL)
OLSEN	70	PRL 24 843	+Fondrom, Handler, Limon, Smith+	(WISC, MICH)
DAUBER	69	PR 179 1262	+Berge, Hubbard, Merrill, Miller	(LRL)
DOYLE	69	UCRL 18139 Thesis		(LRL)
MALONEY	69	PRL 23 425	+Sechi-Zorn	(UMD)
GRIMM	68	NC 54A 187		(HEID)
HEPP	68	ZPHY 214 71	+Schleich	(HEID)
BADIER	67	PL 25B 152	+Bonnet, Briandet, Sadoulet	(EPOL)
MAYEUR	67	U.Libr.Brux.Bul. 32	+Tompa, Wickens	(BELG, LOUC)
OVERSETH	67	PRL 19 391	+Roth	(MICH, PRIN)
PDC	67	RMP 39 1	Rosenfeld, Barbaro-Galtieri, Podolsky+	(LRL, CERN, YALE)
BURAN	66	PL 20 318	+Eivindson, Skjeggstad, Tofte+	(OSLO)
CHIEN	66	PR 152 1171	+Lach, Sandweiss, Taft, Yeh, Oren+	(YALE, BNL)
ENGELMANN	66	NC 45A 1038	+Filthuth, Alexander+	(HEID, REHO)
GIBSON	66	NC 45A 882	+Green	(BRIS)
LONDON	66	PR 143 1034	+Rau, Goldberg, Lichtman+	(BNL, SYRA)
SCHMIDT	65	PR 140B 1328		(COLU)
BAGLIN	64	NC 35 977	+Bingham+	(EPOL, CERN, LOUC, RHEL, BERG)
HUBBARD	64	PR 135B 183	+Berge, Kalbfleisch, Shafer+	(LRL)
LIND	64	PR 135B 1483	+Binford, Good, Stern	(WISC)
RONNE	64	PL 11 357	+	(CERN, EPOL, LOUC, BERG+)
SCHWARTZ	64	UCRL 11360 Thesis		(LRL)
BHOWMIK	63	NC 28 1494	+Goyal	(DELH)
BLOCK	63	PR 130 766	+Gessaroli, Ratti+	(NWES, BGNA, SYRA, ORNL)
BROWN	63	PR 130 769	+Kadyk, Trilling, Roe+	(LRL, MICH)
CHRETIEN	63	PR 131 2208	+	(BRAN, BROW, HARV, MIT)
CRONIN	63	PR 129 1795	+Overseht	(PRIN)
ELY	63	PR 131 868	+Gidal, Kalmus, Oswald, Powell+	(LRL)
HUMPHREY	62	PR 127 1305	+Ross	(LRL)
CORK	60	PR 120 1000	+Kerth, Wenzel, Cronin+	(LRL, PRIN, BNL)
CRAWFORD	59B	PRL 2 266	+Cresti, Douglass, Good, Ticho+	(LRL)

NOTE ON Λ AND Σ RESONANCES

Introduction: There are no new results at all on Λ and Σ resonances. The field remains at a standstill and will only be revived if a kaon factory is built. What follows is a much abbreviated version of the Note on Λ and Σ Resonances from our 1990 edition. In particular, see that edition for some representative Argand plots from partial-wave analyses.

Table 1 is an attempt to evaluate the status, both overall and channel by channel, of each Λ and Σ resonance in the Full Listings. The evaluations are of course partly subjective. A blank indicates there is no evidence at all: either the relevant couplings are small or the resonance does not really exist. The main Baryon Summary Table includes only the established resonances (overall status 3 or 4 stars). A number of the 1- and 2-star entries may eventually disappear, but there are certainly many resonances yet to be discovered underlying the established ones.

Sign conventions for resonance couplings: In terms of the isospin-0 and -1 elastic scattering amplitudes A_0 and A_1 , the amplitude for $K^-p \rightarrow \bar{K}^0 n$ scattering is $\pm(A_1 - A_0)/2$, where the sign depends on conventions used in conjunction with the Clebsch-Gordan coefficients (such as, is the baryon or the meson the "first" particle). If this reaction is partial-wave analyzed and if the overall phase is chosen so that, say, the $\Sigma(1775)D_{15}$ amplitude at resonance points along the positive imaginary axis (points "up"), then any Σ at resonance will point "up" and any Λ at resonance will point "down" (along the negative imaginary axis). Thus the phase at resonance determines the isospin. The above ignores background amplitudes in the resonating partial waves.

That is the basic idea. In a similar but somewhat more complicated way, the phases of the $\bar{K}N \rightarrow \Lambda\pi$ and $\bar{K}N \rightarrow \Sigma\pi$ amplitudes for a resonating wave help determine the SU(3) multiplet to which the resonance belongs. Again, a convention has to be adopted for some overall arbitrary phases: which way is "up"? Our convention is that of Levi-Setti [1] and is shown in Fig. 1, which also compares experimental results with theoretical predictions for the signs of several resonances. In the

Table 1. The status of the Λ and Σ resonances. Only those with an overall status of *** or **** are included in the main Baryon Summary Table.

Particle	$L_{I,2J}$	Overall status	Status as seen in —			
			$N\bar{K}$	$\Lambda\pi$	$\Sigma\pi$	Other channels
$\Lambda(1116)$	P_{01}	****		F		$N\pi$ (weakly)
$\Lambda(1405)$	S_{01}	****	****	o	****	
$\Lambda(1520)$	D_{03}	****	****	r	****	$\Lambda\pi\pi, \Lambda\gamma$
$\Lambda(1600)$	P_{01}	***	***	b	**	
$\Lambda(1670)$	S_{01}	****	****	i	****	$\Lambda\eta$
$\Lambda(1690)$	D_{03}	****	****	d	****	$\Lambda\pi\pi, \Sigma\pi\pi$
$\Lambda(1800)$	S_{01}	***	***	d	**	$N\bar{K}^*, \Sigma(1385)\pi$
$\Lambda(1810)$	P_{01}	***	***	e	**	$N\bar{K}^*$
$\Lambda(1820)$	F_{05}	****	****	n	****	$\Sigma(1385)\pi$
$\Lambda(1830)$	D_{05}	****	***	F	****	$\Sigma(1385)\pi$
$\Lambda(1890)$	P_{03}	****	****	o	**	$N\bar{K}^*, \Sigma(1385)\pi$
$\Lambda(2000)$	*	*	*	r	*	$\Lambda\omega, N\bar{K}^*$
$\Lambda(2020)$	F_{07}	*	*	b	*	
$\Lambda(2100)$	G_{07}	****	****	i	***	$\Lambda\omega, N\bar{K}^*$
$\Lambda(2110)$	F_{05}	***	**	d	*	$\Lambda\omega, N\bar{K}^*$
$\Lambda(2325)$	D_{03}	*	*	d	*	$\Lambda\omega$
$\Lambda(2350)$	*	***	***	e	*	
$\Lambda(2585)$	*	**	**	n	*	
$\Sigma(1193)$	P_{11}	****				$N\pi$ (weakly)
$\Sigma(1385)$	P_{13}	****		****	****	
$\Sigma(1480)$	*	*	*	*	*	
$\Sigma(1560)$	*	**	**	**	**	
$\Sigma(1580)$	D_{13}	**	*	*	*	
$\Sigma(1620)$	S_{11}	**	**	*	*	
$\Sigma(1660)$	P_{11}	***	***	*	**	
$\Sigma(1670)$	D_{13}	****	****	****	****	several others
$\Sigma(1690)$	*	**	*	**	*	$\Lambda\pi\pi$
$\Sigma(1750)$	S_{11}	***	***	**	*	$\Sigma\eta$
$\Sigma(1770)$	P_{11}	*	*	*	*	
$\Sigma(1775)$	D_{15}	****	****	****	***	several others
$\Sigma(1840)$	P_{13}	*	*	**	*	
$\Sigma(1880)$	P_{11}	**	**	**	*	$N\bar{K}^*$
$\Sigma(1915)$	F_{15}	****	***	****	***	$\Sigma(1385)\pi$
$\Sigma(1940)$	D_{13}	***	*	***	**	quasi-2-body
$\Sigma(2000)$	S_{11}	*	*	*	*	$N\bar{K}^*, \Lambda(1520)\pi$
$\Sigma(2030)$	F_{17}	****	****	****	**	several others
$\Sigma(2070)$	F_{15}	*	*	*	*	
$\Sigma(2080)$	P_{13}	**	**	**	*	
$\Sigma(2100)$	G_{17}	*	*	*	*	
$\Sigma(2250)$	*	***	***	*	*	
$\Sigma(2455)$	*	**	*	*	*	
$\Sigma(2620)$	*	**	*	*	*	
$\Sigma(3000)$	*	*	*	*	*	
$\Sigma(3170)$	*	*	*	*	*	multi-body

****	Existence is certain, and properties are at least fairly well explored.
***	Existence ranges from very likely to certain, but further confirmation is desirable and/or quantum numbers, branching fractions, etc. are not well determined.
**	Evidence of existence is only fair.
*	Evidence of existence is poor.

Listings, a + or - sign in front of a measurement of an inelastic resonance coupling indicates the sign (the absence of a sign means that the sign is not determined, *not* that it is positive). For more details, see Appendix II of our 1982 edition [2].

Errors on masses and widths: The errors quoted on resonance parameters from partial-wave analyses are often only statistical, and the parameters can change by more than these errors when a different parametrization of the waves is used. Furthermore, the different analyses use more or less the same data, so it is not really appropriate to treat the different determinations of the resonance parameters as independent or

Baryon Full Listings

Λ 's and Σ 's, $\Lambda(1405)$

to average them together. In any case, the spread of the masses, widths, and branching fractions from the different analyses is certainly a better indication of the uncertainties than are the quoted errors. In the Baryon Summary Table, we usually give a range reflecting the spread of the values rather than a particular value with error.

For three states, the $\Lambda(1520)$, the $\Lambda(1820)$, and the $\Sigma(1775)$, there is enough information to make an overall fit to the various branching fractions. It is then necessary to use the quoted errors, but the errors obtained from the fit should not be taken seriously.

Production experiments: Partial-wave analyses of course separate partial waves, whereas a peak in a cross section or an invariant mass distribution usually cannot be disentangled from background and analyzed for its quantum numbers; and more than one resonance may be contributing to the peak. Results from partial-wave analyses and from production experiments are generally kept separate in the Listings, and in the Baryon Summary Table results from production experiments are used only for the low-mass states. The $\Sigma(1385)$ and $\Lambda(1405)$ of course lie below the $\bar{K}N$ threshold and nearly everything about them is learned from production experiments; and production and formation experiments agree quite well in the case of $\Lambda(1520)$ and results have been combined. There is some disagreement between production and formation experiments in the 1600–1700 MeV region: see the Note on the $\Sigma(1670)$.

References

1. R. Levi-Setti, in *Proceedings of the Lund International Conference on Elementary Particles* (Lund, 1969), p. 339.
2. Particle Data Group, *Phys. Lett.* **111B** (1982).

$\Lambda(1405)$ S_{01}

$I(J^P) = 0(\frac{1}{2}^-)$ Status: ****

NOTE ON THE $\Lambda(1405)$

(by R.H. Dalitz, Oxford University)

It is generally accepted that the $\Lambda(1405)$ is a well-established $J^P = 1/2^-$ resonance. It is assigned to the lowest $L = 1$ supermultiplet of the 3-quark system and paired with the $J^P = 3/2^-$ $\Lambda(1520)$. Lying about 30 MeV below the $N\bar{K}$ threshold, the $\Lambda(1405)$ can be observed directly only as a resonance bump in the $(\Sigma\pi)^0$ subsystem in final states of production experiments. It was first reported by ALSTON 61B in the reaction $K^-p \rightarrow \Sigma\pi\pi\pi$ at 1.15 GeV/c and has since been seen in at least eight other experiments. However, only two of them had enough events for a detailed analysis: THOMAS 73, with about 400 $\Sigma^\pm\pi^\mp$ events from $\pi^-p \rightarrow K^0(\Sigma\pi)^0$ at 1.69 GeV/c; and HEMINGWAY 85, with 766 $\Sigma^+\pi^-$ and 1106 $\Sigma^-\pi^+$ events from $K^-p \rightarrow (\Sigma\pi\pi)^+\pi^-$ at 4.2 GeV/c, after the selections $1600 \leq M(\Sigma\pi\pi)^+ \leq 1720$ MeV and momentum transfer ≤ 1.0 (GeV/c)² to purify the $\Lambda(1405) \rightarrow (\Sigma\pi)^0$ sample. These experiments agree on a mass of about 1395–1400 MeV and a width of about 60 MeV. (Hemingway's mass of 1391 ± 1 MeV is from his best, but unacceptably poor, Breit-Wigner fit.)

The Byers-Fenster tests on these data allow any spin and either parity: neither J nor P has yet been determined *directly*. The early indications for $J^P = 1/2^-$ came from finding $\text{Re } A_{I=0}$ to be large and negative in a constant-scattering-length analysis of low-energy $N\bar{K}$ reaction data (see KIM 65, SAKITT 65, and earlier references cited therein). The first multichannel energy-dependent K-matrix analysis (KIM 67) strengthened the case for a resonance around 1400–1420 MeV strongly coupled to the $I = 0$ S -wave $N\bar{K}$ system.




Figure 1. The signs of the imaginary parts of resonating amplitudes in the $\bar{K}N \rightarrow \Lambda\pi$ and $\Sigma\pi$ channels. The signs of the $\Sigma(1385)$ and $\Lambda(1405)$, marked with a •, are set by convention, and then the others are determined relative to them. The signs required by the SU(3) assignments of the resonances are shown with an arrow, and the experimentally determined signs are shown with an x.

THOMAS 73 and HEMINGWAY 85 both found the $\Lambda(1405)$ bump to be asymmetric and not well fitted by a Breit-Wigner resonance function with constant parameters. The asymmetry involves a rapid fall in intensity as the $N\bar{K}$ threshold energy is approached from below. This is readily understood as due to a strong coupling of the $\Lambda(1405)$ to the S -wave $N\bar{K}$ channel (see DALITZ 81). This striking S -shaped cusp behavior at a new threshold is characteristic of S -wave coupling; the other below-threshold hyperon, the $\Sigma(1385)$, has no such threshold distortion because its $N\bar{K}$ coupling is P -wave. For the $\Lambda(1405)$, this asymmetry is the *sole direct evidence* that $J^P = 1/2^-$.

Following the early work cited above, a considerable literature has developed on proper procedures for phenomenological extrapolation below the $N\bar{K}$ threshold, partly in order to strengthen the evidence for the spin-parity of the $\Lambda(1405)$, and partly to provide an estimate for the amplitude $f(N\bar{K})$ in the unphysical domain below the $N\bar{K}$ threshold; the latter is needed for the evaluation of the dispersion relation for $N\bar{K}$ and NK forward scattering amplitudes. For recent reviews, see MILLER 84 and BARRETT 89. In most recent work, the $(\Sigma\pi)^0$ production spectrum is included in the data fitted (see, e.g., CHAO 73, MARTIN 81).

It is now accepted that the data can be fitted only with an S -wave pole in the reaction amplitudes below $N\bar{K}$ threshold (see, however, FINK 90), but there is still controversy about the physical origin of this pole (for a review, see DALITZ 81 and DALITZ 82). Two extreme possibilities are: (a) an $L = 1$ SU(3)-singlet uds state coupled with the S -wave meson-baryon systems; or (b) an unstable $N\bar{K}$ bound state, analogous to the (stable) deuteron in the NN system. The problem with (a) is that the $\Lambda(1405)$ mass is so much lower than that of its partner, the $\Lambda(1520)$. This requires, in the QCD-inspired quark model, rather large spin-orbit couplings, whether or not one uses relativistic kinetic energies. ISGUR 80, CAPSTICK 86, and CAPSTICK 89 conclude that a proper QCD calculation leads only to small energy splittings, whereas LEINWEBER 90, using QCD sum rules, obtains a good fit to this splitting.

On the other hand, the problem with (b) is that then another $J^P = 1/2^- \Lambda$ is needed to replace the $\Lambda(1405)$ in the $L = 1$ supermultiplet, and it would have to lie close to the $\Lambda(1520)$, a region already well explored by $N\bar{K}$ experiments without result. Intermediate structures are possible; for example, the cloudy bag model allows the configurations (a) and (b) to mix and finds the intensity of (a) in the $\Lambda(1405)$ to be only 14% (VEIT 84, VEIT 85, JENNINGS 86). Such models naturally predict a second $1/2^- \Lambda$ close to the $\Lambda(1520)$.

The determination of the mass and width of the resonance from $(\Sigma\pi)^0$ data is usually based on the "Watson approximation," which states that the production rate $R(\Sigma\pi)$ of the $(\Sigma\pi)^0$ state has a mass dependence proportional to $(\sin^2\delta_{\Sigma\pi})/q$, q being the $\Sigma\pi$ c.m. momentum, in a $\Sigma\pi$ mass range where $\delta_{\Sigma\pi}$ is not far from $\pi/2$ and only the $\Sigma\pi$ channel is open, i.e., between the $\Sigma\pi$ and the $N\bar{K}$ thresholds. Then $qR(\Sigma\pi)$ is proportional to $\sin^2\delta_{\Sigma\pi}$, and the mass M may be defined as the energy at

which $\sin^2\delta_{\Sigma\pi} = 1$. The width Γ may be determined from the rate at which $\delta_{\Sigma\pi}$ goes through $\pi/2$, or from the FWHM; this is a matter of convention.

This determination of M and Γ from the data suffers from the following defects:

(i) The determination of $\sin^2\delta_{\Sigma\pi}$ requires that $R(\Sigma\pi)$ be scaled to give $\sin^2\delta_{\Sigma\pi} = 1$ at the peak for the best fit to the data; i.e., the bump must be *assumed* to arise from a resonance. However, this assumption is supported by the analysis of the low-energy $N\bar{K}$ data and its extrapolation below threshold.

(ii) Owing to the nearby $N\bar{K}$ threshold, the shape of the best fit to the $M(\Sigma\pi)$ bump is uncertain. For energies below this threshold at $E_{N\bar{K}}$, the general form for $\delta_{\Sigma\pi}$ is

$$q \cot \delta_{\Sigma\pi} = \frac{1 + \kappa\alpha}{\gamma + \kappa(\alpha\gamma - \beta^2)}.$$

Here α, β , and γ are the (generally energy-dependent) NN , $N\Sigma$, and $\Sigma\Sigma$ elements of the $I = 0$ S -wave K-matrix for the $(\Sigma\pi, N\bar{K})$ system, and κ is the magnitude of the (imaginary) c.m. momentum k_K for the $N\bar{K}$ system below threshold. The elements α, β, γ are real functions of E ; they have no branch cuts at the $\Sigma\pi$ and $N\bar{K}$ thresholds, but they are permitted to have poles in E along the real E axis. The resonance asymmetry arises from the effect of κ on $\delta_{\Sigma\pi}$. We note that $\delta_{\Sigma\pi} = \pi/2$ when $\kappa = -1/\alpha$.

Accepting this close connection of $\delta_{\Sigma\pi}$ with the low-energy $N\bar{K}$ data, it is natural to analyze the two sets of data together (e.g., MARTIN 81), and there is now a large body of accurate $N\bar{K}$ data for laboratory momenta between 100 and 300 MeV/c (see MILLER 84). The two sets of data span c.m. energies from 1370 MeV to 1490 MeV, and the K-matrix elements will not be energy independent over such a broad range. For the $I = 0$ channels, a linear energy dependence for K^{-1} has been adopted routinely ever since the work of KIM 67, and it is essential when fitting the $qR(\Sigma\pi)$ and $N\bar{K}$ data together. However, $qR(\Sigma\pi)$ is not always well fitted in this procedure; the value obtained for the $\Lambda(1405)$ mass M varies a good deal with the type of fit, not a surprising result when the $\Sigma\pi$ mass spectrum contributes only nine data points in a total of about 200. The value of M obtained from an overall fit is not necessarily much better than from one using only the $qR(\Sigma\pi)$ data; and M may be a function of the representation—K-matrix, K^{-1} -matrix, relativistic-separable or nonseparable potentials, etc.—used in fitting over the full energy range. DALITZ 90 fitted the $qR(\Sigma^+\pi^-)$ Hemingway data with each of the first three representations just mentioned, constrained to the $I = 0$ $N\bar{K}$ threshold scattering length from low-energy $N\bar{K}$ data. The (nonseparable) meson-exchange potentials of MÜLLER-GROELING 90, fitted to the low-energy $N\bar{K}$ (and NK) data, predicted an unstable $N\bar{K}$ bound state with mass and width compatible with the $\Lambda(1405)$.

The present status of the $\Lambda(1405)$ thus depends heavily on theoretical arguments, a somewhat unsatisfactory basis for a four-star rating. Nevertheless, there is no known reason to

Baryon Full Listings

 $\Lambda(1405)$

doubt its existence or quantum numbers. A measurement of the energy-level shifts and widths for the atomic levels of kaonic hydrogen (and deuterium) would give a valuable check on analyses of the $(\Sigma\pi, N\bar{K})$ amplitudes, since the energy of the K^-p atom lies roughly midway between those for the two sets of data. The three measurements of $(\Delta E - i\Gamma/2)$ for kaonic hydrogen are inconsistent with one another and require that the sign of $\text{Re}(A_{I=0} + A_{I=1})$ be opposite that deduced from $N\bar{K}$ reaction data (see BATTY 89). Accurate measurements of $(\Delta E - i\Gamma/2)$ values for kaonic hydrogen are badly needed, but may not be possible until the KAON factory becomes operational.

To definitively settle the nature of the $\Lambda(1405)$ will require much further work, both experimental and theoretical. Higher-statistics experiments on the production and decay of the $\Lambda(1405)$ are needed, but suitable K^- beams will not be available until KAON. The low-energy reaction cross sections, especially for the $\bar{K}^0 p$ interactions, last studied 25 years ago, need to be better determined.

 $\Lambda(1405)$ MASS

PRODUCTION EXPERIMENTS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1406.5 ± 4.0		¹ DALITZ 91		M-matrix fit
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1391 ± 1	700	¹ HEMINGWAY 85	HBC	K^-p 4.2 GeV/c
~ 1405	400	² THOMAS 73	HBC	π^-p 1.69 GeV/c
1405	120	BARBARO....	68B DBC	K^-d 2.1-2.7 GeV/c
1400 ± 5	67	BIRMINGHAM 66	HBC	K^-p 3.5 GeV/c
1382 ± 8		ENGLER 65	HDBC	π^-p, π^+d 1.68 GeV/c
1400 ± 24		MUSGRAVE 65	HBC	$\bar{p}p$ 3-4 GeV/c
1410		ALEXANDER 62	HBC	π^-p 2.1 GeV/c
1405		ALSTON 62	HBC	K^-p 1.2-0.5 GeV/c
1405		ALSTON 61B	HBC	K^-p 1.15 GeV/c

EXTRAPOLATIONS BELOW $N\bar{K}$ THRESHOLD

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1411	³ MARTIN 81		K-matrix fit
1406	⁴ CHAO 73	DPWA	0-range fit (sol. B)
1421	MARTIN 70	RVUE	Constant K-matrix
1416 ± 4	MARTIN 69	HBC	Constant K-matrix
1403 ± 3	KIM 67	HBC	K-matrix fit
1407.5 ± 1.2	⁵ KITTEL 66	HBC	0-effective-range fit
1410.7 ± 1.0	KIM 65	HBC	0-effective-range fit
1409.6 ± 1.7	⁵ SAKITT 65	HBC	0-effective-range fit

 $\Lambda(1405)$ WIDTH

PRODUCTION EXPERIMENTS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
50 ± 2		¹ DALITZ 91		M-matrix fit
• • • We do not use the following data for averages, fits, limits, etc. • • •				
32 ± 1	700	¹ HEMINGWAY 85	HBC	K^-p 4.2 GeV/c
45 ± 55	400	² THOMAS 73	HBC	π^-p 1.69 GeV/c
35	120	BARBARO....	68B DBC	K^-d 2.1-2.7 GeV/c
50 ± 10	67	BIRMINGHAM 66	HBC	K^-p 3.5 GeV/c
89 ± 20		ENGLER 65	HDBC	
60 ± 20		MUSGRAVE 65	HBC	
35 ± 5		ALEXANDER 62	HBC	
50		ALSTON 62	HBC	
20		ALSTON 61B	HBC	

EXTRAPOLATIONS BELOW $N\bar{K}$ THRESHOLD

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
30	³ MARTIN 81		K-matrix fit
55	^{4,6} CHAO 73	DPWA	0-range fit (sol. B)
20	MARTIN 70	RVUE	Constant K-matrix
29 ± 6	MARTIN 69	HBC	Constant K-matrix
50 ± 5	KIM 67	HBC	K-matrix fit
34.1 ± 4.1	⁵ KITTEL 66	HBC	
37.0 ± 3.2	KIM 65	HBC	
28.2 ± 4.1	⁵ SAKITT 65	HBC	

 $\Lambda(1405)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Sigma\pi$	100 %
$\Gamma_2 \Lambda\gamma$	
$\Gamma_3 \Sigma^0\gamma$	
$\Gamma_4 N\bar{K}$	

 $\Lambda(1405)$ PARTIAL WIDTHS

$\Gamma(\Lambda\gamma)$	DOCUMENT ID	COMMENT
VALUE (keV)		
• • • We do not use the following data for averages, fits, limits, etc. • • •		
27 ± 8	BURKHARDT 91	Isobar model fit
$\Gamma(\Sigma^0\gamma)$	DOCUMENT ID	COMMENT
VALUE (keV)		
• • • We do not use the following data for averages, fits, limits, etc. • • •		
10 ± 4 or 23 ± 7	BURKHARDT 91	Isobar model fit

 $\Lambda(1405)$ BRANCHING RATIOS

$\Gamma(N\bar{K})/\Gamma(\Sigma\pi)$	CL%	DOCUMENT ID	TECN	COMMENT
VALUE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
<3	95	HEMINGWAY 85	HBC	K^-p 4.2 GeV/c

 $\Lambda(1405)$ FOOTNOTES

- ¹ DALITZ 91 fits the HEMINGWAY 85 data.
- ² THOMAS 73 data is fit by CHAO 73 (see next section).
- ³ The MARTIN 81 fit includes the $K^\pm p$ forward scattering amplitudes and the dispersion relations they must satisfy.
- ⁴ See also the accompanying paper of THOMAS 73.
- ⁵ Data of SAKITT 65 are used in the fit by KITTEL 66.
- ⁶ An asymmetric shape, with $\Gamma/2 = 41$ MeV below resonance, 14 MeV above.

 $\Lambda(1405)$ REFERENCES

BURKHARDT 91	PR C44 607	+Lowe	(NOTT, UNM, BIRM)
DALITZ 91	JPG 17 289	+Deloff	(OXFTP, WINR)
HEMINGWAY 85	NP B253 742		(CERN) J
MARTIN 81	NP B179 33		(DURH)
CHAO 73	NP B56 46	+Kraemer, Thomas, Martin	(RHEL, CMU, LOUC)
THOMAS 73	NP B56 15	+Engler, Fisk, Kraemer	(CMU) J
MARTIN 70	NP B16 479	+Ross	(DURH)
MARTIN 69	PR 183 1352	+Sakitt	(LOUC, BNL)
Also 69B	PR 183 1345	Martin, Sakitt	(LOUC, BNL)
BARBARO....	68B PRL 21 573	Barbaro-Galiteri, Chadwick+	(LRL, SLAC)
KIM 67	PRL 19 1074		(YALE)
BIRMINGHAM 66	PR 152 1148		(BIRM, GLAS, LOIC, OXF, RHEL)
KITTEL 66	PL 21 349	+Otter, Wacek	(VIEN)
ENGLER 65	PRL 15 224	+Fisk, Kraemer, Meltzer, Westgard+	(CMU, BNL) J
KIM 65	PRL 14 29		(COLU)
MUSGRAVE 65	NC 35 735	+Petmezias+	(BIRM, CERN, EPOL, LOIC, SACL)
SAKITT 65	PR 139B 719	+Day, Glasser, Seeman, Friedman+	(UMD, LRL)
ALEXANDER 62	PRL 8 447	+Kalbfleisch, Miller, Smith	(LRL) I
ALSTON 62	CERN Conf. 311	+Alvarez, Ferro-Luzzi+	(LRL) I
ALSTON 61B	PRL 6 698	+Alvarez, Eberhard, Good+	(LRL) I

OTHER RELATED PAPERS

FINK 90	PR C41 2720	+He, Landau, Schnick	(IBMY, ORST, ANSM)
LEINWEBER 90	ANP 198 203		(MCMS)
MUELLER-GR... 90	NP A513 557	Mueller-Groeling, Hohlde, Speth	(JULI)
BARRETT 89	NC 102A 179		(SURRE)
BATTY 89	NC 102A 255	+Gal	(RAL, HEBR)
CAPSTICK 89	Excited Baryons '88, p. 32		(GUEL)
LOWE 89	NC 102A 167		(BIRM)
WHITEHOUSE 89	PRL 63 1352	+ (BIRM, BOST, BRCO, BNL, CASE, BUDA, TRIU)	(BIRM)
SIEGEL 88	PR C38 2221	+Weise	(REGG)
WORKMAN 88	PR D37 3117	+Fearing	(TRIU)
SCHNICK 87	PRL 58 1719	+Landau	(ORST)
CAPSTICK 86	PR D34 2809	+Isgur	(TNT0)
JENNINGS 86	PL B176 229		(TRIU)
MALTMAN 86	PR D34 1372	+Isgur	(LANL, TNT0)
ZHONG 86	PL B171 471	+Thomas, Jennings, Barrett	(ADLD, TRIU, SURRE)
BURKHARDT 85	NP A440 653	+Lowe, Rosenthal	(NOTT, BIRM, WMU)
DAREWYCH 85	PR D32 1765	+Konik, Isgur	(YORKC, TNT0)
VEIT 85	PR D31 1033	+Jennings, Thomas, Barrett	(TRIU, ADLD, SURRE)
KIANG 84	PR C30 1638	+Kumar, Nogami, VanDijk	(DALH, MCMS)
MILLER 84			(LOUC)
Conf. Intersections between Particle and Nuclear Physics, p. 783			
VANDIJK 84	PR D30 937		(MCMS)
VEIT 84	PL 137B 415	+Jennings, Barrett, Thomas	(TRIU, SURRE, CERN)
DALITZ 82		+McGinley, Belyea, Anthony	(OXFTP)
Heidelberg Conf., p. 201			
DALITZ 81		+McGinley	(OXFTP)
Low and Intermediate Energy Kaon-Nucleon Physics, p.381			
MARTIN 81B	Low and Intermediate Energy Kaon-Nucleon Phys., p. 97		(DURH)
OADES 77	NC 42A 462	+Rasche	(AARH, ZURI)
SHAW 73	Purdue Conf. 417		(UCI)
BARBARO.... 72	LBL-555	Barbaro-Galiteri	(LBL)
DOBSON 72	PR D6 3256	+McElhaney	(HAWA)
RAJASEKARA... 72	PR DS 610	Rajasekaran	(TATA)
Earlier papers also cited in RAJASEKARAN 72.			
CLINE 71	PRL 26 1194	+Laumann, Mapp	(WISC)
MARTIN 71	PL 35B 62	+Martin, Ross	(DURH, LOUC, RHEL)
DALITZ 67	PR 153 1617	+Wong, Rajasekaran	(OXFTP, BOMB)
DONALD 66	PL 22 711	+Edwards, Lys, Nisar, Moore	(LIVP)
KADYK 66	PRL 17 599	+Oren, Goldhaber, Goldhaber, Trilling	(LRL)
ABRAMS 65	PR 139B 454	+Secni-Zorn	(UMD)

See key on page 1343

Baryon Full Listings
 $\Lambda(1520)$

$\Lambda(1520) D_{03}$

$I(J^P) = 0(\frac{3}{2}^-)$ Status: * * * *

Discovered by FERRO-LUZZI 62; the elaboration in WATSON 63 is the classic paper on the Breit-Wigner analysis of a multichannel resonance.

The measurements of the mass, width, and elasticity published before 1975 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters 111B (1982).

Production and formation experiments agree quite well, so they are listed together here.

$\Lambda(1520)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1519.5 ± 1.0 OUR ESTIMATE				
1519.50 ± 0.18 OUR AVERAGE				
1517.3 ± 1.5	300	BARBER	80D SPEC	$\gamma p \rightarrow \Lambda(1520)K^+$
1519 ± 1		GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$
1517.8 ± 1.2	5k	BARLAG	79 HBC	$K^- p$ 4.2 GeV/c
1520.0 ± 0.5		ALSTON-...	78 DPWA	$\bar{K}N \rightarrow \bar{K}N$
1519.7 ± 0.3	4k	CAMERON	77 HBC	$K^- p$ 0.96-1.36 GeV/c
1519 ± 1		GOPAL	77 DPWA	$\bar{K}N$ multichannel
1519.4 ± 0.3	2000	CORDEN	75 DBC	$K^- d$ 1.4-1.8 GeV/c

$\Lambda(1520)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
15.6 ± 1.0 OUR ESTIMATE				
15.59 ± 0.27 OUR AVERAGE				
16.3 ± 3.3	300	BARBER	80D SPEC	$\gamma p \rightarrow \Lambda(1520)K^+$
16 ± 1		GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$
14 ± 3	677	¹ BARLAG	79 HBC	$K^- p$ 4.2 GeV/c
15.4 ± 0.5		ALSTON-...	78 DPWA	$\bar{K}N \rightarrow \bar{K}N$
16.3 ± 0.5	4k	CAMERON	77 HBC	$K^- p$ 0.96-1.36 GeV/c
15.0 ± 0.5		GOPAL	77 DPWA	$\bar{K}N$ multichannel
15.5 ± 1.6	2000	CORDEN	75 DBC	$K^- d$ 1.4-1.8 GeV/c

$\Lambda(1520)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 N\bar{K}$	45 ± 1%
$\Gamma_2 \Sigma\pi$	42 ± 1%
$\Gamma_3 \Lambda\pi\pi$	10 ± 1%
$\Gamma_4 \Sigma(1385)\pi$	
$\Gamma_5 \Sigma(1385)\pi(\rightarrow \Lambda\pi\pi)$	
$\Gamma_6 \Lambda(\pi\pi)_S\text{-wave}$	
$\Gamma_7 \Sigma\pi\pi$	0.9 ± 0.1%
$\Gamma_8 \Lambda\gamma$	0.8 ± 0.2%
$\Gamma_9 \Sigma^0\gamma$	

CONSTRAINED FIT INFORMATION

An overall fit to 9 branching ratios uses 24 measurements and one constraint to determine 6 parameters. The overall fit has a $\chi^2 = 16.5$ for 19 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-63				
x_3	-32	-33			
x_7	-4	-3	-1		
x_8	-9	-8	-4	0	
x_9	-24	-21	-10	-1	-2
	x_1	x_2	x_3	x_7	x_8

$\Lambda(1520)$ BRANCHING RATIOS


See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.45 ± 0.01 OUR ESTIMATE				
0.448 ± 0.007 OUR FIT			Error includes scale factor of 1.2.	
0.455 ± 0.011 OUR AVERAGE				
0.47 ± 0.02	GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.45 ± 0.03	ALSTON-...	78 DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.448 ± 0.014	CORDEN	75 DBC	$K^- d$ 1.4-1.8 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.47 ± 0.01	GOPAL	77 DPWA	See GOPAL 80	
0.42	MAST	76 HBC	$K^- p \rightarrow \bar{K}^0 n$	

$\Gamma(\Sigma\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
0.42 ± 0.01 OUR ESTIMATE				
0.421 ± 0.007 OUR FIT			Error includes scale factor of 1.2.	
0.423 ± 0.011 OUR AVERAGE				
0.426 ± 0.014	CORDEN	75 DBC	$K^- d$ 1.4-1.8 GeV/c	
0.418 ± 0.017	BARBARO-...	69B HBC	$K^- p$ 0.28-0.45 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.46	KIM	71 DPWA	K-matrix analysis	

$\Gamma(\Sigma\pi)/\Gamma(N\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
0.940 ± 0.026 OUR FIT			Error includes scale factor of 1.3.	
0.95 ± 0.04 OUR AVERAGE			Error includes scale factor of 1.7. See the Ideogram below.	
0.98 ± 0.03	² GOPAL	77 DPWA	$\bar{K}N$ multichannel	
0.82 ± 0.08	BURKHARDT	69 HBC	$K^- p$ 0.8-1.2 GeV/c	
1.06 ± 0.14	SCHEUER	68 DBC	$K^- N$ 3 GeV/c	
0.96 ± 0.20	DAHL	67 HBC	$\pi^- p$ 1.6-4 GeV/c	
0.73 ± 0.11	DAUBER	67 HBC	$K^- p$ 2 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.06 ± 0.12	BERTHON	74 HBC	Quasi-2-body σ	
1.72 ± 0.78	MUSGRAVE	65 HBC		

WEIGHTED AVERAGE
 0.95±0.04 (Error scaled by 1.7)



$\Gamma(\Lambda\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.10 ± 0.01 OUR ESTIMATE				
0.096 ± 0.005 OUR FIT			Error includes scale factor of 1.2.	
0.096 ± 0.008 OUR AVERAGE			Error includes scale factor of 1.6.	
0.091 ± 0.006	CORDEN	75 DBC	$K^- d$ 1.4-1.8 GeV/c	
0.11 ± 0.01	³ MAST	73B IPWA	$K^- p \rightarrow \Lambda\pi\pi$	

$\Gamma(\Lambda\pi\pi)/\Gamma(N\bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.213 ± 0.012 OUR FIT			Error includes scale factor of 1.2.	
0.202 ± 0.021 OUR AVERAGE				
0.22 ± 0.03	BURKHARDT	69 HBC	$K^- p$ 0.8-1.2 GeV/c	
0.19 ± 0.04	SCHEUER	68 DBC	$K^- N$ 3 GeV/c	
0.17 ± 0.05	DAHL	67 HBC	$\pi^- p$ 1.6-4 GeV/c	
0.21 ± 0.18	DAUBER	67 HBC	$K^- p$ 2 GeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.27 ± 0.13	BERTHON	74 HBC	Quasi-2-body σ	
0.2	KIM	71 DPWA	K-matrix analysis	

Baryon Full Listings

 $\Lambda(1520), \Lambda(1600)$ $\Gamma(\Sigma\pi)/\Gamma(\Lambda\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_3
4.42 ± 0.25 OUR FIT	Error includes scale factor of 1.2.			
3.9 ± 0.6 OUR AVERAGE				
3.9 ± 1.0	UHLIG 67	HBC	$K^- p$ 0.9–1.0 GeV/c	
3.3 ± 1.1	BIRMINGHAM 66	HBC	$K^- p$ 3.5 GeV/c	
4.5 ± 1.0	ARMENTEROS65C	HBC		

 $\Gamma(\Sigma(1385)\pi)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
0.041 ± 0.005	CHAN 72	HBC	$K^- p \rightarrow \Lambda\pi\pi$	

 $\Gamma(\Sigma(1385)\pi(\rightarrow \Lambda\pi\pi))/\Gamma(\Lambda\pi\pi)$

The $\Lambda\pi\pi$ mode is largely due to $\Sigma(1385)\pi$. Only the values of $(\Sigma(1385)\pi)/(\Lambda\pi\pi)$ given by MAST 73B and CORDEN 75 are based on real 3-body partial-wave analyses. The discrepancy between the two results is essentially due to the different hypotheses made concerning the shape of the $(\pi\pi)_S$ -wave state.

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ_3
0.58 ± 0.22	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.82 ± 0.10	⁴ MAST 73B	IPWA	$K^- p \rightarrow \Lambda\pi\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.39 ± 0.10	⁵ BURKHARDT 71	HBC	$K^- p \rightarrow (\Lambda\pi\pi)\pi$	

 $\Gamma(\Lambda(\pi\pi)_S\text{-wave})/\Gamma(\Lambda\pi\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ_3
0.20 ± 0.08	CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	

 $\Gamma(\Sigma\pi\pi)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
0.009 ± 0.001 OUR ESTIMATE				
0.0086 ± 0.0005 OUR FIT				
0.0086 ± 0.0005 OUR AVERAGE				
0.007 ± 0.002	⁶ CORDEN 75	DBC	$K^- d$ 1.4–1.8 GeV/c	
0.0085 ± 0.0006	⁷ MAST 73	MPWA	$K^- p \rightarrow \Sigma\pi\pi$	
0.010 ± 0.0015	BARBARO... 69B	HBC	$K^- p$ 0.28–0.45 GeV/c	

 $\Gamma(\Lambda\gamma)/\Gamma_{total}$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
0.008 ± 0.002 OUR ESTIMATE					
0.0079 ± 0.0014 OUR FIT					
0.0080 ± 0.0014	238	MAST 68B	HBC	Using $\Gamma(N\bar{K})/\Gamma_{total} = 0.45$	

 $\Gamma(\Sigma^0\gamma)/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_9/Γ
0.0195 ± 0.0034 OUR FIT				
0.02 ± 0.0035	⁸ MAST 68B	HBC	Not measured; see note	

 $\Lambda(1520)$ FOOTNOTES

- From the best-resolution sample of $\Lambda\pi\pi$ events only.
- The $\bar{K}N \rightarrow \Sigma\pi$ amplitude at resonance is $+0.46 \pm 0.01$.
- Assumes $\Gamma(N\bar{K})/\Gamma_{total} = 0.46 \pm 0.02$.
- Both $\Sigma(1385)\pi$ DS_{03} and $\Sigma(\pi\pi)$ DP_{03} contribute.
- The central bin (1514–1524 MeV) gives 0.74 ± 0.10 ; other bins are lower by 2-to-5 standard deviations.
- Much of the $\Sigma\pi\pi$ decay proceeds via $\Sigma(1385)\pi$.
- Assumes $\Gamma(N\bar{K})/\Gamma_{total} = 0.46$.
- Calculated from $\Gamma(\Lambda\gamma)/\Gamma_{total}$, assuming SU(3). Needed to constrain the sum of all the branching ratios to be unity.

 $\Lambda(1520)$ REFERENCES

PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
BARBER 80D	ZPHY C7 17	+Dainton, Lee, Marshall+	(DARE, LANC, SHEF)
GOPAL 80	Toronto Conf. 159		(RHEL) IJP
BARLAG 79	NP B149 220	+Blokzijl, Jongejans+	(AMST, CERN, NIJM, OXF)
ALSTON... 78	D18 182	+Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also 77	PRL 38 1007	+Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON 77	NP B131 399	+Frank, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MAST 76	PR D14 13	+Alston-Garnjost, Bangert+	(LBL)
CORDEN 75	NP B84 306	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM)
BERTHON 74	NC 21A 146	+Tristram+	(CDEF, RHEL, SACL, STRB)
MAST 70	PR D7 3212	+Bangert, Alston-Garnjost+	(LBL) IJP
MAST 73B	PR D7 5	+Bangert, Alston-Garnjost+	(LBL) IJP
CHAN 72	PRL 28 256	+Button-Shafer, Hertzbach, Kofler+	(MASA, YALE)
BURKHARDT 71	NP B27 64	+Flithuth, Kluge+	(HEID, CERN, SACL)
KIM 71	PRL 27 356		(HARV) IJP
Also 70	Duke Conf. 161		(HARV) IJP
BARBARO... 69B	Lund Conf. 352	Kim	(LRL)
Also 70	Duke Conf. 95	Barbaro-Galtieri, Bangert, Mast, Tripp	(LRL)
BURKHARDT 69	NP B14 106	+Flithuth, Kluge+	(HEID, EFI, CERN, SACL)
MAST 68B	PRL 21 1715	+Bangert, Alston-Garnjost, Bangert, Galtieri+	(LRL)
SCHUEER 68	NP B8 503	+Merrill, Vergias, DeWitt+	(SABRE Collab.)
DAHL 67	PR 163 1377	+Hardy, Hess, Kirz, Miller	(LRL)
DAUBER 67	PL 24B 525	+Malamad, Schlein, Slater, Stork	(UCLA)
UHLIG 67	PR 155 1448	+Charlton, Condon, Glasser, Yodh+	(UMD, NRL)
BIRMINGHAM 66	PR 152 1148		(BIRM, GLAS, LOIC, OXF, RHEL)
ARMENTEROS 65C	PL 19 338		(CERN, HEID, SACL)
MUSGRAVE 65	NC 35 735	+Ferro-Luzzi+	(BIRM, CERN, EPOL, LOIC, SACL)
WATSON 63	PR 131 2248	+Petmezias+	(LRL) IJP
FERRO-LUZZI 62	PRL 8 28	+Ferro-Luzzi, Tripp	(LRL) IJP
		+Tripp, Watson	(LRL) IJP

 $\Lambda(1600) P_{01}$

$$I(J^P) = 0(\frac{1}{2}^+) \text{ Status: } ***$$

See also the $\Lambda(1810) P_{01}$. There are quite possibly two P_{01} states in this region.

 $\Lambda(1600)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1560 to 1700 (≈ 1600) OUR ESTIMATE			
1568 ± 20	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1703 ± 100	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1573 ± 25	GOPAL 77	DPWA	$\bar{K}N$ multichannel
1596 ± 6	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
1620 ± 10	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1572 or 1617	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
1646 ± 17	² CARROLL 76	DPWA	Isospin-0 total σ
1570	KIM 71	DPWA	K-matrix analysis

 $\Lambda(1600)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50 to 250 (≈ 150) OUR ESTIMATE			
116 ± 20	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
593 ± 200	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
147 ± 50	GOPAL 77	DPWA	$\bar{K}N$ multichannel
175 ± 20	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
60 ± 10	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
247 or 271	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
20	² CARROLL 76	DPWA	Isospin-0 total σ
50	KIM 71	DPWA	K-matrix analysis

 $\Lambda(1600)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	15–30 %
Γ_2 $\Sigma\pi$	10–60 %

The above branching fractions are our estimates, not fits or averages.

 $\Lambda(1600)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

 $\Gamma(N\bar{K})/\Gamma_{total}$

VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.15 to 0.30 OUR ESTIMATE				
0.23 ± 0.04	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.14 ± 0.05	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.25 ± 0.15	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24 ± 0.04	GOPAL 77	DPWA	See GOPAL 80	
0.30 or 0.29	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	

 $(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total}$ in $N\bar{K} \rightarrow \Lambda(1600) \rightarrow \Sigma\pi$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
–0.16 ± 0.04	GOPAL 77	DPWA	$\bar{K}N$ multichannel	
–0.33 ± 0.11	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$	
0.28 ± 0.09	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
–0.39 or –0.39	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	
not seen	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma\pi$	

 $\Lambda(1600)$ FOOTNOTES

- The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- A total cross-section bump with $(J+1/2)\Gamma_{el}/\Gamma_{total} = 0.04$.

 $\Lambda(1600)$ REFERENCES

GOPAL 80	Toronto Conf. 159		(RHEL) IJP
ALSTON... 78	PR D18 182		(LRL, MTHO, CERN) IJP
Also 77	PRL 38 1007	+Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN 77	NP B127 349	+Pidcock, Moorhouse	(LOIC, GLAS) IJP
Also 77B	NP B126 266	Martin, Pidcock	(LOIC)
Also 77C	NP B126 285	Martin, Pidcock	(LOIC) IJP
CARROLL 76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL) I
HEPP 76B	PL 65B 487	+Braun, Grimm, Strobele+	(CERN, HEID, MPIM) IJP
KANE 74	LBL-2452		(LBL) IJP
LANGBEIN 72	NP B47 477	+Wagner	(MPIM) IJP
KIM 71	PRL 27 356		(HARV) IJP

See key on page 1343

Baryon Full Listings
 $\Lambda(1670), \Lambda(1690)$

$\Lambda(1670) S_{01}$ $I(J^P) = 0(\frac{1}{2}^-)$ Status: * * * *

The measurements of the mass, width, and elasticity published before 1974 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** (1982).

$\Lambda(1670)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1660 to 1680 (≈ 1670) OUR ESTIMATE			
1670.8 ± 1.7	KOISO 85	DPWA	$K^- p \rightarrow \Sigma \pi$
1667 ± 5	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
1671 ± 3	ALSTON-... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
1670 ± 5	GOPAL 77	DPWA	$\bar{K} N$ multichannel
1675 ± 2	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma \pi$
1679 ± 1	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$
1665 ± 5	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1664	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel

$\Lambda(1670)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
25 to 80 (≈ 38) OUR ESTIMATE			
34.1 ± 3.7	KOISO 85	DPWA	$K^- p \rightarrow \Sigma \pi$
29 ± 5	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
29 ± 5	ALSTON-... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
45 ± 10	GOPAL 77	DPWA	$\bar{K} N$ multichannel
46 ± 5	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma \pi$
40 ± 3	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$
19 ± 5	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
12	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel

$\Lambda(1670)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	15-25 %
Γ_2 $\Sigma \pi$	20-60 %
Γ_3 $\Lambda \eta$	15-35 %
Γ_4 $\Sigma(1385)\pi$	

The above branching fractions are our estimates, not fits or averages.

$\Lambda(1670)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.15 to 0.25 OUR ESTIMATE				
0.18 ± 0.03	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$	
0.17 ± 0.03	ALSTON-... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.20 ± 0.03	GOPAL 77	DPWA	See GOPAL 80	
0.15	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	

$(\Gamma_1 \Gamma_f)^{1/2}/\Gamma_{total} \ln N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2}/\Gamma$
0.15 to 0.25 OUR ESTIMATE				
-0.26 ± 0.02	KOISO 85	DPWA	$K^- p \rightarrow \Sigma \pi$	
-0.31 ± 0.03	GOPAL 77	DPWA	$\bar{K} N$ multichannel	
-0.29 ± 0.03	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma \pi$	
-0.23 ± 0.03	LONDON 75	HLBC	$K^- p \rightarrow \Sigma^0 \pi^0$	
-0.27 ± 0.02	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.13	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	

$(\Gamma_1 \Gamma_f)^{1/2}/\Gamma_{total} \ln N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Lambda \eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_3)^{1/2}/\Gamma$
0.15 to 0.25 OUR ESTIMATE				
+0.20 ± 0.05	BAXTER 73	DPWA	$K^- p \rightarrow$ neutrals	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24	KIM 71	DPWA	K-matrix analysis	
0.26	ARMENTEROS69C HBC			
0.20 or 0.23	BERLEY 65	HBC		

$(\Gamma_1 \Gamma_f)^{1/2}/\Gamma_{total} \ln N\bar{K} \rightarrow \Lambda(1670) \rightarrow \Sigma(1385)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_4)^{1/2}/\Gamma$
0.15 to 0.25 OUR ESTIMATE				
-0.18 ± 0.05	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$	

$\Lambda(1670)$ FOOTNOTES

¹MARTIN 77 obtains identical resonance parameters from a T-matrix pole and from a Breit-Wigner fit.

$\Lambda(1670)$ REFERENCES

KOISO 85	NP A433 619	+Sai, Yamamoto, Kofler	(TOKY, MASA)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL 80	Toronto Conf. 159		(RHEL) IJP
ALSTON-... 78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also 77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN 77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also 77B	NP B126 266	Martin, Pidcock	(LOUC) IJP
Also 77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
HEPP 76B	PL 65B 487	+Braun, Grimm, Strobele+	(CERN, HEIDH, MPIM) IJP
LONDON 75	NP B65 289	+Yu, Boyd+	(BNL, CERN, EPOL, ORSAY, TORI)
KANE 74	LBL-2452		(LBL) IJP
PREVOST 74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID)
BAXTER 73	NP B67 125	+Buckingham, Corbett, Dunn+	(OXF) IJP
KIM 71	PRL 27 356		(HARV) IJP
Also 70	Duke Conf. 161	Kim	(HARV) IJP
ARMENTEROS 69C	Lund Paper 229	+Baillon+	(CERN, HEID, SACL) IJP
Values are quoted in LEVI-SETTI 69.			
BERLEY 65	PRL 15 641	+Connolly, Hart, Rahm, Stonehill+	(BNL) IJP

$\Lambda(1690) D_{03}$ $I(J^P) = 0(\frac{3}{2}^-)$ Status: * * * *

The measurements of the mass, width, and elasticity published before 1974 are now obsolete and have been omitted. They were last listed in our 1982 edition Physics Letters **111B** (1982).

$\Lambda(1690)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1685 to 1695 (≈ 1690) OUR ESTIMATE			
1695.7 ± 2.6	KOISO 85	DPWA	$K^- p \rightarrow \Sigma \pi$
1690 ± 5	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
1692 ± 5	ALSTON-... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
1690 ± 5	GOPAL 77	DPWA	$\bar{K} N$ multichannel
1690 ± 3	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma \pi$
1689 ± 1	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1687 or 1689	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel
1692 ± 4	CARROLL 76	DPWA	Isospin-0 total σ

$\Lambda(1690)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50 to 70 (≈ 60) OUR ESTIMATE			
67.2 ± 5.6	KOISO 85	DPWA	$K^- p \rightarrow \Sigma \pi$
61 ± 5	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
64 ± 10	ALSTON-... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
60 ± 5	GOPAL 77	DPWA	$\bar{K} N$ multichannel
82 ± 8	HEPP 76B	DPWA	$K^- N \rightarrow \Sigma \pi$
60 ± 4	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
62 or 62	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel
38	CARROLL 76	DPWA	Isospin-0 total σ

$\Lambda(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	20-30 %
Γ_2 $\Sigma \pi$	20-40 %
Γ_3 $\Lambda \pi \pi$	~ 25 %
Γ_4 $\Sigma \pi \pi$	~ 20 %
Γ_5 $\Lambda \eta$	
Γ_6 $\Sigma(1385)\pi, S$ -wave	

The above branching fractions are our estimates, not fits or averages.

$\Lambda(1690)$ BRANCHING RATIOS

The sum of all the quoted branching ratios is more than 1.0. The two-body ratios are from partial-wave analyses, and thus probably are more reliable than the three-body ratios, which are determined from bumps in cross sections. Of the latter, the $\Sigma \pi \pi$ bump looks more significant. (The error given for the $\Lambda \pi \pi$ ratio looks unreasonably small.) Hardly any of the $\Sigma \pi \pi$ decay can be via $\Sigma(1385)$, for then seven times as much $\Lambda \pi \pi$ decay would be required. See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.2 to 0.3 OUR ESTIMATE				
0.23 ± 0.03	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$	
0.22 ± 0.03	ALSTON-... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.24 ± 0.03	GOPAL 77	DPWA	See GOPAL 80	
0.28 or 0.26	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	

Baryon Full Listings

$\Lambda(1690), \Lambda(1800)$

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma \pi$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
-0.34 ± 0.02	KOISO 85 DPWA $K^- p \rightarrow \Sigma \pi$
-0.25 ± 0.03	GOPAL 77 DPWA $\bar{K} N$ multichannel
-0.29 ± 0.03	HEPP 76B DPWA $K^- N \rightarrow \Sigma \pi$
-0.28 ± 0.03	LONDON 75 HLBC $K^- p \rightarrow \Sigma^0 \pi^0$
-0.28 ± 0.02	KANE 74 DPWA $K^- p \rightarrow \Sigma \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •	
-0.30 or -0.28	¹ MARTIN 77 DPWA $\bar{K} N$ multichannel

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1690) \rightarrow \Lambda \eta$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
0.00 ± 0.03	BAXTER 73 DPWA $K^- p \rightarrow$ neutrals

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1690) \rightarrow \Lambda \pi \pi$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
0.25 ± 0.02	² BARTLEY 68 HDBC $K^- p \rightarrow \Lambda \pi \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •	

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma \pi \pi$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
0.21	ARMENTEROS68C HDBC $K^- N \rightarrow \Sigma \pi \pi$

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1690) \rightarrow \Sigma(1385) \pi, S\text{-wave}$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
+0.27 ± 0.04	PREVOST 74 DPWA $K^- N \rightarrow \Sigma(1385) \pi$

$\Lambda(1690)$ FOOTNOTES

- ¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit. Another $D_{03} \Lambda$ at 1666 MeV is also suggested by MARTIN 77, but is very uncertain.
²BARTLEY 68 uses only cross-section data. The enhancement is not seen by PREVOST 71.

$\Lambda(1690)$ REFERENCES

KOISO 85 NP A433 619	+Sai, Yamamoto, Koffer (TOKY, MASA)
PDG 82 PL 111B	Roos, Porter, Aguilar-Benitez+ (HELS, CIT, CERN)
GOPAL 80 Toronto Conf. 159	(RHEL) IJP
ALSTON... 78 PR D18 182	Alston-Garnjost, Kenney+ (LBL, MTHO, CERN) IJP
Also 77 PRL 38 1007	Alston-Garnjost, Kenney+ (LBL, MTHO, CERN) IJP
GOPAL 77 NP B119 362	+Ross, VanHorn, McPherson+ (LOIC, RHEL) IJP
MARTIN 77 NP B127 349	+Pidcock, Moorhouse (LOUC, GLAS) IJP
Also 77B NP B126 286	Martin, Pidcock (LOUC) IJP
Also 77C NP B126 285	Martin, Pidcock (LOUC) IJP
CARRROLL 76 PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+ (BNL) I
HEPP 76B PL 65B 487	+Braun, Grimm, Strobele+ (CERN, HEID, MPIM) IJP
LONDON 75 NP B85 289	+Yu, Boyd+ (BNL, CERN, EPOL, ORSAY, TORI) IJP
KANE 74 LBL-2452	+Barloutaud+ (SACL, CERN, HEID) IJP
PREVOST 74 NP B69 246	+Buckingham, Corbett, Dunn+ (OXF) IJP
BAXTER 73 NP B67 125	(CERN, HEID, SACL) I
PREVOST 71 Amsterdam Conf.	+Baillon+ (CERN, HEID, SACL) I
ARMENTEROS 68C NP B8 216	+Chu, Dowd, Greene+ (TUFTS, FSU, BRAN) I
BARTLEY 68 PRL 21 1111	

$\Lambda(1800) S_{01}$

$$I(J^P) = 0(\frac{1}{2}^-) \text{ Status: } ***$$

This is the second resonance in the S_{01} wave, the first being the $\Lambda(1670)$.

$\Lambda(1800)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1720 to 1850 (≈ 1800) OUR ESTIMATE			
1841 ± 10	GOPAL 80 DPWA $\bar{K} N \rightarrow \bar{K} N$		
1725 ± 20	ALSTON... 78 DPWA $\bar{K} N \rightarrow \bar{K} N$		
1825 ± 20	GOPAL 77 DPWA $\bar{K} N$ multichannel		
1830 ± 20	LANGBEIN 72 IPWA $\bar{K} N$ multichannel		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1767 or 1842	¹ MARTIN 77 DPWA $\bar{K} N$ multichannel		
1780	KIM 71 DPWA K-matrix analysis		
1872 ± 10	BRICMAN 70B DPWA $\bar{K} N \rightarrow \bar{K} N$		

$\Lambda(1800)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
200 to 400 (≈ 300) OUR ESTIMATE			
228 ± 20	GOPAL 80 DPWA $\bar{K} N \rightarrow \bar{K} N$		
185 ± 20	ALSTON... 78 DPWA $\bar{K} N \rightarrow \bar{K} N$		
230 ± 20	GOPAL 77 DPWA $\bar{K} N$ multichannel		
70 ± 15	LANGBEIN 72 IPWA $\bar{K} N$ multichannel		
• • • We do not use the following data for averages, fits, limits, etc. • • •			
435 or 473	¹ MARTIN 77 DPWA $\bar{K} N$ multichannel		
40	KIM 71 DPWA K-matrix analysis		
100 ± 20	BRICMAN 70B DPWA $\bar{K} N \rightarrow \bar{K} N$		

$\Lambda(1800)$ DECAY MODES

Mode	Fraction (Γ_i / Γ)
$\Gamma_1 \bar{K} N$	25-40 %
$\Gamma_2 \Sigma \pi$	seen
$\Gamma_3 \Sigma(1385) \pi$	seen
$\Gamma_4 N \bar{K}^*(892)$	seen
$\Gamma_5 N \bar{K}^*(892), S=1/2, S\text{-wave}$	
$\Gamma_6 N \bar{K}^*(892), S=3/2, D\text{-wave}$	

The above branching fractions are our estimates, not fits or averages.

$\Lambda(1800)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N \bar{K}) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
0.25 to 0.40 OUR ESTIMATE				
0.36 ± 0.04	GOPAL 80 DPWA $\bar{K} N \rightarrow \bar{K} N$			
0.28 ± 0.05	ALSTON... 78 DPWA $\bar{K} N \rightarrow \bar{K} N$			
0.35 ± 0.15	LANGBEIN 72 IPWA $\bar{K} N$ multichannel			
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.37 ± 0.05	GOPAL 77 DPWA See GOPAL 80			
1.21 or 0.70	¹ MARTIN 77 DPWA $\bar{K} N$ multichannel			
0.80	KIM 71 DPWA K-matrix analysis			
0.18 ± 0.02	BRICMAN 70B DPWA $\bar{K} N \rightarrow \bar{K} N$			

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1800) \rightarrow \Sigma \pi$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
-0.08 ± 0.05	GOPAL 77 DPWA $\bar{K} N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •	
-0.74 or -0.43	¹ MARTIN 77 DPWA $\bar{K} N$ multichannel
0.24	KIM 71 DPWA K-matrix analysis

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1800) \rightarrow \Sigma(1385) \pi$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
+0.056 ± 0.028	² CAMERON 78 DPWA $K^- p \rightarrow \Sigma(1385) \pi$

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1800) \rightarrow N \bar{K}^*(892), S=1/2, S\text{-wave}$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
-0.17 ± 0.03	² CAMERON 78B DPWA $K^- p \rightarrow N \bar{K}^*$

$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Lambda(1800) \rightarrow N \bar{K}^*(892), S=3/2, D\text{-wave}$	$(\Gamma_1 \Gamma_2)_{1/2} / \Gamma$
VALUE	DOCUMENT ID TECN COMMENT
-0.13 ± 0.04	CAMERON 78B DPWA $K^- p \rightarrow N \bar{K}^*$

$\Lambda(1800)$ FOOTNOTES

- ¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
²The published sign has been changed to be in accord with the baryon-first convention.

$\Lambda(1800)$ REFERENCES

GOPAL 80 Toronto Conf. 159	Alston-Garnjost, Kenney+ (RHEL) IJP
ALSTON... 78 PR D18 182	Alston-Garnjost, Kenney+ (LBL, MTHO, CERN) IJP
Also 77 PRL 38 1007	Alston-Garnjost, Kenney+ (LBL, MTHO, CERN) IJP
CAMERON 78 NP B143 189	+Franeck, Gopal, Bacon, Butterworth+ (RHEL, LOIC) IJP
CAMERON 78B NP B146 327	+Franeck, Gopal, Kaimus, McPherson+ (RHEL, LOIC) IJP
GOPAL 77 NP B119 362	+Ross, VanHorn, McPherson+ (LOIC, RHEL) IJP
MARTIN 77 NP B127 349	+Pidcock, Moorhouse (LOUC, GLAS) IJP
Also 77B NP B126 266	Martin, Pidcock (LOUC) IJP
Also 77C NP B126 285	Martin, Pidcock (LOUC) IJP
LANGBEIN 72 NP B47 477	+Wagner (MPIM) IJP
KIM 71 PRL 27 356	Kim (HARV) IJP
Also 70 Duke Conf. 161	Kim (HARV) IJP
BRICMAN 70B PL 33B 511	+Ferro-Luzzi, Lagnaux (CERN) IJP

See key on page 1343

Baryon Full Listings

$\Lambda(1810), \Lambda(1820)$

$\Lambda(1810) P_{01}$

$$I(J^P) = 0(\frac{1}{2}^+) \text{ Status: } ***$$

Almost all the recent analyses contain a P_{01} state, and sometimes two of them, but the masses, widths, and branching ratios vary greatly. See also the $\Lambda(1600) P_{01}$.

$\Lambda(1810)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1750 to 1850 (≈ 1810) OUR ESTIMATE			
1841 \pm 20	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1853 \pm 20	GOPAL	77	DPWA $\bar{K}N$ multichannel
1735 \pm 5	CARROLL	76	DPWA Isospin-0 total σ
1746 \pm 10	PREVOST	74	DPWA $K^-N \rightarrow \Sigma(1385)\pi$
1780 \pm 20	LANGBEIN	72	IPWA $\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1861 or 1953	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
1755	KIM	71	DPWA K-matrix analysis
1800	ARMENTEROS70	HBC	$\bar{K}N \rightarrow \bar{K}N$
1750	ARMENTEROS70	HBC	$\bar{K}N \rightarrow \Sigma\pi$
1690 \pm 10	BARBARO-...	70	HBC $\bar{K}N \rightarrow \Sigma\pi$
1740	BAILEY	69	DPWA $\bar{K}N \rightarrow \bar{K}N$
1745	ARMENTEROS68B	HBC	$\bar{K}N \rightarrow \bar{K}N$

$\Lambda(1810)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
50 to 250 (≈ 150) OUR ESTIMATE			
164 \pm 20	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
90 \pm 20	CAMERON	78B	DPWA $K^-p \rightarrow N\bar{K}^*$
166 \pm 20	GOPAL	77	DPWA $\bar{K}N$ multichannel
46 \pm 20	PREVOST	74	DPWA $K^-N \rightarrow \Sigma(1385)\pi$
120 \pm 10	LANGBEIN	72	IPWA $\bar{K}N$ multichannel
• • • We do not use the following data for averages, fits, limits, etc. • • •			
535 or 585	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
28	CARROLL	76	DPWA Isospin-0 total σ
35	KIM	71	DPWA K-matrix analysis
30	ARMENTEROS70	HBC	$\bar{K}N \rightarrow \bar{K}N$
70	ARMENTEROS70	HBC	$\bar{K}N \rightarrow \Sigma\pi$
22	BARBARO-...	70	HBC $\bar{K}N \rightarrow \Sigma\pi$
300	BAILEY	69	DPWA $\bar{K}N \rightarrow \bar{K}N$
147	ARMENTEROS68B	HBC	

$\Lambda(1810)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	20–50 %
Γ_2 $\Sigma\pi$	10–40 %
Γ_3 $\Sigma(1385)\pi$	seen
Γ_4 $N\bar{K}^*(892)$	30–60 %
Γ_5 $N\bar{K}^*(892), S=1/2, P\text{-wave}$	
Γ_6 $N\bar{K}^*(892), S=3/2, P\text{-wave}$	

The above branching fractions are our estimates, not fits or averages.

$\Lambda(1810)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.2 to 0.5 OUR ESTIMATE				
0.24 \pm 0.04	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.36 \pm 0.05	LANGBEIN	72	IPWA $\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.21 \pm 0.04	GOPAL	77	DPWA See GOPAL 80	
0.52 or 0.49	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	
0.30	KIM	71	DPWA K-matrix analysis	
0.15	ARMENTEROS70	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.55	BAILEY	69	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.4	ARMENTEROS68B	DPWA	$\bar{K}N \rightarrow \bar{K}N$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1810) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_2)^{1/2}/\Gamma$
0.2 to 0.5 OUR ESTIMATE				
-0.24 \pm 0.04	GOPAL	77	DPWA $\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.25 or +0.23	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	
< 0.01	LANGBEIN	72	IPWA $\bar{K}N$ multichannel	
0.17	KIM	71	DPWA K-matrix analysis	
+0.20	² ARMENTEROS70	DPWA	$\bar{K}N \rightarrow \Sigma\pi$	
-0.13 \pm 0.03	BARBARO-...	70	DPWA $\bar{K}N \rightarrow \Sigma\pi$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1810) \rightarrow \Sigma(1385)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_3)^{1/2}/\Gamma$
0.2 to 0.5 OUR ESTIMATE				
+0.18 \pm 0.10	PREVOST	74	DPWA $K^-N \rightarrow \Sigma(1385)\pi$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1810) \rightarrow N\bar{K}^*(892), S=1/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_5)^{1/2}/\Gamma$
0.2 to 0.5 OUR ESTIMATE				
-0.14 \pm 0.03	² CAMERON	78B	DPWA $K^-p \rightarrow N\bar{K}^*$	

$(\Gamma_i/\Gamma_f)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1810) \rightarrow N\bar{K}^*(892), S=3/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_6)^{1/2}/\Gamma$
0.2 to 0.5 OUR ESTIMATE				
+0.35 \pm 0.06	CAMERON	78B	DPWA $K^-p \rightarrow N\bar{K}^*$	

$\Lambda(1810)$ FOOTNOTES

- ¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
² The published sign has been changed to be in accord with the baryon-first convention.

$\Lambda(1810)$ REFERENCES

GOPAL	80	Toronto Conf. 159		(RHEL) IJP
CAMERON	78B	NP B146 327	+Franeck, Gopal, Kaimus, McPherson+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
		Also	Martin, Pidcock	(LOUC)
		Also	Martin, Pidcock	(LOUC) IJP
CARROLL	76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL) I
PREVOST	74	NP B69 246	+Baroutaud+	(SACL, CERN, HEID)
LANGBEIN	72	NP B47 477	+Wagner	(MFM) IJP
KIM	71	PRL 27 356		(HARV) IJP
		Also	Kim	(HARV) IJP
ARMENTEROS70	70	Duke Conf. 123	+Baillon+	(CERN, HEID, SACL) IJP
BARBARO-...	70	Duke Conf. 173	Barbaro-Galtieri	(LRL) IJP
BAILEY	69	UCRL 50617 Thesis		(LRL) IJP
ARMENTEROS68B	NP B8 195		+Baillon+	(CERN, HEID, SACL) IJP

$\Lambda(1820) F_{05}$

$$I(J^P) = 0(\frac{5}{2}^+) \text{ Status: } ****$$

This resonance is the cornerstone for all partial-wave analyses in this region. Most of the results published before 1973 are now obsolete and have been omitted. They may be found in our 1982 edition Physics Letters **111B** (1982).

Most of the quoted errors are statistical only; the systematic errors due to the particular parametrizations used in the partial-wave analyses are not included. For this reason we do not calculate weighted averages for the mass and width.

$\Lambda(1820)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1815 to 1825 (≈ 1820) OUR ESTIMATE			
1823 \pm 3	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1819 \pm 2	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
1822 \pm 2	GOPAL	77	DPWA $\bar{K}N$ multichannel
1821 \pm 2	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1830	DECLAIS	77	DPWA $\bar{K}N \rightarrow \bar{K}N$
1817 or 1819	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

$\Lambda(1820)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70 to 90 (≈ 80) OUR ESTIMATE			
77 \pm 5	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
72 \pm 5	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
81 \pm 5	GOPAL	77	DPWA $\bar{K}N$ multichannel
87 \pm 3	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
82	DECLAIS	77	DPWA $\bar{K}N \rightarrow \bar{K}N$
76 or 76	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

$\Lambda(1820)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	55–65 %
Γ_2 $\Sigma\pi$	8–14 %
Γ_3 $\Sigma(1385)\pi$	5–10 %
Γ_4 $\Sigma(1385)\pi, P\text{-wave}$	
Γ_5 $\Sigma(1385)\pi, F\text{-wave}$	
Γ_6 $\Lambda\eta$	
Γ_7 $\Sigma\pi\pi$	

The above branching fractions are our estimates, not fits or averages.

Baryon Full Listings

 $\Lambda(1820)$, $\Lambda(1830)$ $\Lambda(1820)$ BRANCHING RATIOS

Errors quoted do not include uncertainties in the parametrizations used in the partial-wave analyses and are thus too small. See also "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.55 to 0.65 OUR ESTIMATE				
0.58 ± 0.02	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.60 ± 0.03	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$	
••• We do not use the following data for averages, fits, limits, etc. •••				
0.51	DECLAIS	77	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.57 ± 0.02	GOPAL	77	DPWA See GOPAL 80	
0.59 or 0.58	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1820) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
-0.28 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.28 ± 0.01	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$	
••• We do not use the following data for averages, fits, limits, etc. •••				
-0.25 or -0.25	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1820) \rightarrow \Lambda\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
VALUE				
-0.096 +0.040 -0.020	RADER	73	MPWA	

$\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
VALUE				
no clear signal	² ARMENTEROS68C	HDBC	$K^-N \rightarrow \Sigma\pi\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1820) \rightarrow \Sigma(1385)\pi$, P-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
VALUE				
-0.167 ± 0.054	³ CAMERON	78	DPWA $K^-p \rightarrow \Sigma(1385)\pi$	
+0.27 ± 0.03	PREVOST	74	DPWA $K^-N \rightarrow \Sigma(1385)\pi$	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1820) \rightarrow \Sigma(1385)\pi$, F-wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
VALUE				
+0.065 ± 0.029	³ CAMERON	78	DPWA $K^-p \rightarrow \Sigma(1385)\pi$	

 $\Lambda(1820)$ FOOTNOTES

- The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- There is a suggestion of a bump, enough to be consistent with what is expected from $\Sigma(1385) \rightarrow \Sigma\pi$ decay.
- The published sign has been changed to be in accord with the baryon-first convention.

 $\Lambda(1820)$ REFERENCES

PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON-...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON	78	NP B143 189	+FraneK, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
DECLAIS	77	CERN 77-16	+Duchon, Louvel, Patry, Seguinot+	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
KANE	74	LBL-2452		(LBL) IJP
PREVOST	74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID)
RADER	73	NC 16A 178	+Barloutaud+	(SACL, HEID, CERN, RHEL, CDEF)
ARMENTEROS 68C	NP B8 216		+Baillon+	(CERN, HEID, SACL)

 $\Lambda(1830) D_{05}$

$$I(J^P) = 0(\frac{5}{2}^-) \text{ Status: } ***$$

For results published before 1973 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

The best evidence for this resonance is in the $\Sigma\pi$ channel.

 $\Lambda(1830)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1810 ± 10			
1810 ± 10	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1825 ± 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
1825 ± 1	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
••• We do not use the following data for averages, fits, limits, etc. •••			
1817 or 1818	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

 $\Lambda(1830)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
60 to 110 (≈ 95) OUR ESTIMATE			
100 ± 10	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
94 ± 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
119 ± 3	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
••• We do not use the following data for averages, fits, limits, etc. •••			
56 or 56	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

 $\Lambda(1830)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	3-10 %
Γ_2 $\Sigma\pi$	35-75 %
Γ_3 $\Sigma(1385)\pi$	> 15 %
Γ_4 $\Sigma(1385)\pi$, D-wave	
Γ_5 $\Lambda\eta$	

The above branching fractions are our estimates, not fits or averages.

 $\Lambda(1830)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.03 to 0.10 OUR ESTIMATE				
0.08 ± 0.03	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.02 ± 0.02	ALSTON-...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$	
••• We do not use the following data for averages, fits, limits, etc. •••				
0.04 ± 0.03	GOPAL	77	DPWA See GOPAL 80	
0.04 or 0.04	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1830) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
-0.17 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.15 ± 0.01	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$	
••• We do not use the following data for averages, fits, limits, etc. •••				
-0.17 or -0.17	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1830) \rightarrow \Lambda\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
VALUE				
-0.044 ± 0.020	RADER	73	MPWA	

$(\Gamma_1\Gamma_7)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(1830) \rightarrow \Sigma(1385)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
VALUE				
+0.141 ± 0.014	² CAMERON	78	DPWA $K^-p \rightarrow \Sigma(1385)\pi$	
+0.13 ± 0.03	PREVOST	74	DPWA $K^-N \rightarrow \Sigma(1385)\pi$	

 $\Lambda(1830)$ FOOTNOTES

- The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- The CAMERON 78 upper limit on G-wave decay is 0.03. The published sign has been changed to be in accord with the baryon-first convention.

 $\Lambda(1830)$ REFERENCES

PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON-...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON	78	NP B143 189	+FraneK, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
KANE	74	LBL-2452		(LBL) IJP
PREVOST	74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID)
RADER	73	NC 16A 178	+Barloutaud+	(SACL, HEID, CERN, RHEL, CDEF)

See key on page 1343

Baryon Full Listings
 $\Lambda(1890), \Lambda(2000)$ $\Lambda(1890) P_{03}$

$$I(J^P) = 0(\frac{3}{2}^+) \text{ Status: } ****$$

For results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

The $J^P = 3/2^+$ assignment is consistent with all available data (including polarization) and recent partial-wave analyses. The dominant inelastic modes remain unknown.

 $\Lambda(1890)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1850 to 1910 (≈ 1890) OUR ESTIMATE			
1897 ± 5	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1908 ± 10	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1900 ± 5	GOPAL 77	DPWA	$\bar{K}N$ multichannel
1894 ± 10	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1856 or 1868	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
1900	² NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

 $\Lambda(1890)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
60 to 200 (≈ 100) OUR ESTIMATE			
74 ± 10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
119 ± 20	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
72 ± 10	GOPAL 77	DPWA	$\bar{K}N$ multichannel
107 ± 10	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
191 or 193	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
100	² NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

 $\Lambda(1890)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	20–35 %
Γ_2 $\Sigma\pi$	3–10 %
Γ_3 $\Sigma(1385)\pi$	seen
Γ_4 $\Sigma(1385)\pi, P$ -wave	
Γ_5 $\Sigma(1385)\pi, F$ -wave	
Γ_6 $N\bar{K}^*(892)$	seen
Γ_7 $N\bar{K}^*(892), S=1/2, P$ -wave	
Γ_8 $\Lambda\omega$	

The above branching fractions are our estimates, not fits or averages.

 $\Lambda(1890)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.20 to 0.35 OUR ESTIMATE				
0.20 ± 0.02	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.34 ± 0.05	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.24 ± 0.04	HEMINGWAY 75	DPWA	$K^- p \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.18 ± 0.02	GOPAL 77	DPWA	See GOPAL 80	
0.36 or 0.34	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1890) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
-0.09 ± 0.03	GOPAL 77	DPWA	$\bar{K}N$ multichannel	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.15 or +0.14	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1890) \rightarrow \Lambda\omega$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_8)^{1/2}/\Gamma$
VALUE				
seen	BACCARI 77	IPWA	$K^- p \rightarrow \Lambda\omega$	
0.032	² NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1890) \rightarrow \Sigma(1385)\pi, P$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
VALUE				
<0.03	CAMERON 78	DPWA	$K^- p \rightarrow \Sigma(1385)\pi$	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1890) \rightarrow \Sigma(1385)\pi, F$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
VALUE				
-0.126 ± 0.055	³ CAMERON 78	DPWA	$K^- p \rightarrow \Sigma(1385)\pi$	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(1890) \rightarrow N\bar{K}^*(892)$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
VALUE				
-0.07 ± 0.03	^{3,4} CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$	

 $\Lambda(1890)$ FOOTNOTES

- ¹The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- ²Found in one of two best solutions.
- ³The published sign has been changed to be in accord with the baryon-first convention.
- ⁴Upper limits on the P_3 and F_3 waves are each 0.03.

 $\Lambda(1890)$ REFERENCES

PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL 80	Toronto Conf. 159		(RHEL) IJP
ALSTON... 78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
		Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
		+Frank, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
CAMERON 78B	NP B146 327	+Frank, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
BACCARI 77	NC 41A 96	+Poulard, Revel, Tallini+	(SACL, CDEF) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN 77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
		Martin, Pidcock	(LOUC)
		Martin, Pidcock	(LOUC) IJP
HEMINGWAY 75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPIM) IJP
NAKKASYAN 75	NP B93 85		(CERN) IJP

 $\Lambda(2000)$

$$I(J^P) = 0(?^?) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

We list here all the ambiguous resonance possibilities with a mass around 2 GeV. The proposed quantum numbers are D_3 (BARBARO-GALTIERI 70 in $\Sigma\pi$), D_3+F_5 , P_3+D_5 , or P_1+D_3 (BRANDSTETTER 72 in $\Lambda\omega$), and S_1 (CAMERON 78B in $N\bar{K}^*$). The first two of the above analyses should now be considered obsolete. See also NAKKASYAN 75.

 $\Lambda(2000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2000 OUR ESTIMATE			
2030 ± 30	CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$
1935 to 1971	¹ BRANDSTET...72	DPWA	$K^- p \rightarrow \Lambda\omega$
1951 to 2034	¹ BRANDSTET...72	DPWA	$K^- p \rightarrow \Lambda\omega$
2010 ± 30	BARBARO... 70	DPWA	$K^- p \rightarrow \Sigma\pi$

 $\Lambda(2000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
125 ± 25	CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$
180 to 240	¹ BRANDSTET...72	DPWA	(lower mass)
73 to 154	¹ BRANDSTET...72	DPWA	(higher mass)
130 ± 50	BARBARO... 70	DPWA	$K^- p \rightarrow \Sigma\pi$

 $\Lambda(2000)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$
Γ_2 $\Sigma\pi$
Γ_3 $\Lambda\omega$
Γ_4 $N\bar{K}^*(892), S=1/2, S$ -wave
Γ_5 $N\bar{K}^*(892), S=3/2, D$ -wave

 $\Lambda(2000)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2000) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
-0.20 ± 0.04	BARBARO... 70	DPWA	$K^- p \rightarrow \Sigma\pi$	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2000) \rightarrow \Lambda\omega$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
VALUE				
0.17 to 0.25	¹ BRANDSTET...72	DPWA	(lower mass)	
0.04 to 0.15	¹ BRANDSTET...72	DPWA	(higher mass)	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2000) \rightarrow N\bar{K}^*(892), S=1/2, S$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
VALUE				
-0.12 ± 0.03	² CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$	

$(\Gamma_i\Gamma_j)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2000) \rightarrow N\bar{K}^*(892), S=3/2, D$ -wave	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
VALUE				
+0.09 ± 0.03	CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$	

Baryon Full Listings

 $\Lambda(2000)$, $\Lambda(2020)$, $\Lambda(2100)$ $\Lambda(2000)$ FOOTNOTES

- ¹ The parameters quoted here are ranges from the three best fits; the lower state probably has $J \leq 3/2$, and the higher one probably has $J \leq 5/2$.
² The published sign has been changed to be in accord with the baryon-first convention.

 $\Lambda(2000)$ REFERENCES

CAMERON	78B	NP B146 327	+Franeek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
NAKKASYAN	75	NP B93 85		(CERN) IJP
BRANDSTET...	72	NP B39 13	Brandstetter, Butterworth+	(RHEL, CDEF, SACL) IJP
BARBARO-...	70	Duke Conf. 173	Barbaro-Galtieri	(LRL) IJP

 $\Lambda(2020)$ F_{07}

$$I(J^P) = 0(\frac{7}{2}^+) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

In LITCHFIELD 71, need for the state rests solely on a possibly inconsistent polarization measurement at 1.784 GeV/c. HEMINGWAY 75 does not require this state. GOPAL 77 does not need it in either $N\bar{K}$ or $\Sigma\pi$. With new K^-n angular distributions included, DECLAIS 77 sees it. However, this and other new data are included in GOPAL 80 and the state is not required. BACCARI 77 weakly supports it.

 $\Lambda(2020)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2020 OUR ESTIMATE			
2140	BACCARI 77	DPWA	$K^-p \rightarrow \Lambda\omega$
2117	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2100 \pm 30	LITCHFIELD 71	DPWA	$K^-p \rightarrow \bar{K}N$
2020 \pm 20	BARBARO-... 70	DPWA	$K^-p \rightarrow \Sigma\pi$

 $\Lambda(2020)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
128	BACCARI 77	DPWA	$K^-p \rightarrow \Lambda\omega$
167	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
120 \pm 30	LITCHFIELD 71	DPWA	$K^-p \rightarrow \bar{K}N$
160 \pm 30	BARBARO-... 70	DPWA	$K^-p \rightarrow \Sigma\pi$

 $\Lambda(2020)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	25-35 %
Γ_2 $\Sigma\pi$	~ 5 %
Γ_3 $\Lambda\eta$	< 3 %
Γ_4 ΞK	< 3 %
Γ_5 $\Lambda\omega$	< 8 %
Γ_6 $N\bar{K}^*(892)$	10-20 %
Γ_7 $N\bar{K}^*(892)$, $S=1/2$, G -wave	
Γ_8 $N\bar{K}^*(892)$, $S=3/2$, D -wave	

 $\Lambda(2020)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.05	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.05 \pm 0.02	LITCHFIELD 71	DPWA	$K^-p \rightarrow \bar{K}N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2020) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.15 \pm 0.02	BARBARO-... 70	DPWA	$K^-p \rightarrow \Sigma\pi$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2020) \rightarrow \Lambda\omega$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
< 0.05	BACCARI 77	DPWA	$K^-p \rightarrow \Lambda\omega$	

 $\Lambda(2020)$ REFERENCES

GOPAL	80	Toronto Conf. 159		(RHEL)
BACCARI	77	NC 41A 96	+Poulard, Revel, Tallini+	(SACL, CDEF) IJP
DECLAIS	77	CERN 77-16	+Duchon, Louvel, Patry, Seguinot+	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL)
HEMINGWAY	75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPI) IJP
LITCHFIELD	71	NP B30 125	+... Lesquoy+	(RHEL, CDEF, SACL) IJP
BARBARO-...	70	Duke Conf. 173	Barbaro-Galtieri	(LRL) IJP

 $\Lambda(2100)$ G_{07}

$$I(J^P) = 0(\frac{7}{2}^-) \text{ Status: } ***$$

Discovered by COOL 66 and by WOHL 66. Most of the results published before 1973 are now obsolete and have been omitted. They may be found in our 1982 edition Physics Letters **111B** (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and in invariant-mass distributions around 2100 MeV used to be listed in a separate entry immediately following. It may be found in our 1986 edition Physics Letters **170B** (1986).

 $\Lambda(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2090 to 2110 (≈ 2100) OUR ESTIMATE			
2104 \pm 10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2106 \pm 30	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2110 \pm 10	GOPAL 77	DPWA	$\bar{K}N$ multichannel
2105 \pm 10	HEMINGWAY 75	DPWA	$K^-p \rightarrow \bar{K}N$
2115 \pm 10	KANE 74	DPWA	$K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2094	BACCARI 77	DPWA	$K^-p \rightarrow \Lambda\omega$
2094	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2110 or 2089	¹ NAKKASYAN 75	DPWA	$K^-p \rightarrow \Lambda\omega$

 $\Lambda(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100 to 250 (≈ 200) OUR ESTIMATE			
157 \pm 40	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
250 \pm 30	GOPAL 77	DPWA	$\bar{K}N$ multichannel
241 \pm 30	HEMINGWAY 75	DPWA	$K^-p \rightarrow \bar{K}N$
152 \pm 15	KANE 74	DPWA	$K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
98	BACCARI 77	DPWA	$K^-p \rightarrow \Lambda\omega$
250	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
244 or 302	¹ NAKKASYAN 75	DPWA	$K^-p \rightarrow \Lambda\omega$

 $\Lambda(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	25-35 %
Γ_2 $\Sigma\pi$	~ 5 %
Γ_3 $\Lambda\eta$	< 3 %
Γ_4 ΞK	< 3 %
Γ_5 $\Lambda\omega$	< 8 %
Γ_6 $N\bar{K}^*(892)$	10-20 %
Γ_7 $N\bar{K}^*(892)$, $S=1/2$, G -wave	
Γ_8 $N\bar{K}^*(892)$, $S=3/2$, D -wave	

The above branching fractions are our estimates, not fits or averages.

 $\Lambda(2100)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.25 to 0.35 OUR ESTIMATE				
0.34 \pm 0.03	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.24 \pm 0.06	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.31 \pm 0.03	HEMINGWAY 75	DPWA	$K^-p \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.29	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.30 \pm 0.03	GOPAL 77	DPWA	See GOPAL 80	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
+0.12 \pm 0.04	GOPAL 77	DPWA	$\bar{K}N$ multichannel	
+0.11 \pm 0.01	KANE 74	DPWA	$K^-p \rightarrow \Sigma\pi$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Lambda\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
-0.050 \pm 0.020	RADER 73	MPWA	$K^-p \rightarrow \Lambda\eta$	

See key on page 1343

Baryon Full Listings
 $\Lambda(2100)$, $\Lambda(2110)$

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Xi K$	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
0.035 ± 0.018		LITCHFIELD 71	DPWA	$K^- p \rightarrow \Xi K$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.003		MULLER 69B	DPWA	$K^- p \rightarrow \Xi K$
0.05		TRIPP 67	RVUE	$K^- p \rightarrow \Xi K$

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2100) \rightarrow \Lambda\omega$	$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
-0.070		² BACCARI 77	DPWA	GD_{37} wave
+0.011		² BACCARI 77	DPWA	GG_{17} wave
+0.008		² BACCARI 77	DPWA	GG_{37} wave
0.122 or 0.154		¹ NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892), S=3/2, D\text{-wave}$	$(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
+0.21 ± 0.04		CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2100) \rightarrow N\bar{K}^*(892), S=1/2, G\text{-wave}$	$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
-0.04 ± 0.03		³ CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$

 $\Lambda(2100)$ FOOTNOTES

- The NAKKASYAN 75 values are from the two best solutions found. Each has the $\Lambda(2100)$ and one additional resonance (P_3 or F_5).
- Note that the three for BACCARI 77 entries are for three different waves.
- The published sign has been changed to be in accord with the baryon-first convention. The upper limit on the G_3 wave is 0.03.

 $\Lambda(2100)$ REFERENCES

PDG 86	PL 170B	Aguilar-Benitez, Porter+	(CERN, CIT+)
PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
CAMERON 80	Toronto Conf. 159		(RHEL) IJP
GOPAL 78B	NP B146 327	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
DEBELLEFON 78	NC 42A 403	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
BACCARI 77	NC 41A 96	+Poulard, Revel, Tallini+	(SACL, CDEF) IJP
DECLAIS 77	CERN 77-16	+Duchon, Louvel, Patry, Seguinot+	(CAEN, CERN) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
HEMINGWAY 75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPIM) IJP
NAKKASYAN 75	NP B93 85		(CERN) IJP
KANE 74	LBL-2452		(LBL) IJP
RADER 73	NC 16A 178	+Barloutaud+	(SACL, HEID, CERN, RHEL, CDEF)
LITCHFIELD 71	NP B30 125	+.... Lesquoy+	(RHEL, CDEF, SACL) IJP
MULLER 69B	UCRL 19372 Thesis		(LRL)
TRIPP 67	NP B3 10	+Leith+	(LRL, SLAC, CERN, HEID, SACL)
COOL 66	PRL 16 1228	+Giacomelli, Kycia, Leontic, Lundby+	(BNL)
WOHL 66	PRL 17 107	+Solmitz, Stevenson	(LRL) IJP

 $\Lambda(2110) F_05$

$$I(J^P) = 0(\frac{5}{2}^+) \text{ Status: } ***$$

For results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters 111B (1982). All the references have been retained.

This resonance is in the Baryon Summary Table, but the evidence for it could be better.

 $\Lambda(2110)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2090 to 2140 (≈ 2110) OUR ESTIMATE			
2092 ± 25	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2125 ± 25	CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$
2106 ± 50	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
2140 ± 20	DEBELLEFON 77	DPWA	$K^- p \rightarrow \Sigma\pi$
2100 ± 50	GOPAL 77	DPWA	$\bar{K}N$ multichannel
2112 ± 7	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2137	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
2103	¹ NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

 $\Lambda(2110)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 to 250 (≈ 200) OUR ESTIMATE			
245 ± 25	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
160 ± 30	CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$
251 ± 50	DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
140 ± 20	DEBELLEFON 77	DPWA	$K^- p \rightarrow \Sigma\pi$
200 ± 50	GOPAL 77	DPWA	$\bar{K}N$ multichannel
190 ± 30	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
132	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
391	¹ NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

 $\Lambda(2110)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	5–25 %
Γ_2 $\Sigma\pi$	10–40 %
Γ_3 $\Lambda\omega$	seen
Γ_4 $\Sigma(1385)\pi$	seen
Γ_5 $\Sigma(1385)\pi, P\text{-wave}$	
Γ_6 $N\bar{K}^*(892)$	10–60 %
Γ_7 $N\bar{K}^*(892), S=1/2, F\text{-wave}$	

The above branching fractions are our estimates, not fits or averages.

 $\Lambda(2110)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 to 0.26 OUR ESTIMATE				
0.07 ± 0.03	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.27 ± 0.06	² DEBELLEFON 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.07 ± 0.03	GOPAL 77	DPWA	See GOPAL 80	

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2110) \rightarrow \Sigma\pi$	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
+0.14 ± 0.01		DEBELLEFON 77	DPWA	$K^- p \rightarrow \Sigma\pi$
+0.20 ± 0.03		KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
+0.10 ± 0.03		GOPAL 77	DPWA	$\bar{K}N$ multichannel

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2110) \rightarrow \Lambda\omega$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
<0.05		BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
0.112		¹ NAKKASYAN 75	DPWA	$K^- p \rightarrow \Lambda\omega$

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2110) \rightarrow \Sigma(1385)\pi$	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
+0.071 ± 0.025		³ CAMERON 78	DPWA	$K^- p \rightarrow \Sigma(1385)\pi$

$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Lambda(2110) \rightarrow N\bar{K}^*(892)$	$(\Gamma_1 \Gamma_6)^{1/2} / \Gamma$	DOCUMENT ID	TECN	COMMENT
-0.17 ± 0.04		⁴ CAMERON 78B	DPWA	$K^- p \rightarrow N\bar{K}^*$

 $\Lambda(2110)$ FOOTNOTES

- Found in one of two best solutions.
- The published error of 0.6 was a misprint.
- The CAMERON 78 upper limit on F -wave decay is 0.03. The sign here has been changed to be in accord with the baryon-first convention.
- The published sign has been changed to be in accord with the baryon-first convention. The CAMERON 78B upper limits on the P_3 and F_3 waves are each 0.03.

 $\Lambda(2110)$ REFERENCES

PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL 80	Toronto Conf. 159		(RHEL) IJP
CAMERON 78	NP B143 189	+FraneK, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
CAMERON 78B	NP B146 327	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
DEBELLEFON 78	NC 42A 403	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
BACCARI 77	NC 41A 96	+Poulard, Revel, Tallini+	(SACL, CDEF) IJP
DEBELLEFON 77	NC 37A 175	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
NAKKASYAN 75	NP B93 85		(CERN) IJP
KANE 74	LBL-2452		(LBL) IJP

Baryon Full Listings

 $\Lambda(2325)$, $\Lambda(2350)$, $\Lambda(2585)$ Bumps $\Lambda(2325) D_{03}$

$I(J^P) = 0(\frac{3}{2}^-)$ Status: *

OMITTED FROM SUMMARY TABLE

BACCARI 77 finds this state with either $J^P = 3/2^-$ or $3/2^+$ in a energy-dependent partial-wave analyses of $K^- p \rightarrow \Lambda \omega$ from 2070 to 2436 MeV. A subsequent semi-energy-independent analysis from threshold to 2436 MeV selects $3/2^-$. DEBELLEFON 78 (same group) also sees this state in an energy-dependent partial-wave analysis of $K^- p \rightarrow \bar{K} N$ data, and finds $J^P = 3/2^-$ or $3/2^+$. They again prefer $J^P = 3/2^-$, but only on the basis of model-dependent considerations.

 $\Lambda(2325)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2325 OUR ESTIMATE			
2342 \pm 30	DEBELLEFON 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
2327 \pm 20	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda \omega$

 $\Lambda(2325)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
177 \pm 40	DEBELLEFON 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
160 \pm 40	BACCARI 77	IPWA	$K^- p \rightarrow \Lambda \omega$

 $\Lambda(2325)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	$\sim 12\%$
Γ_2 $\Lambda\omega$	$\sim 10\%$

 $\Lambda(2325)$ BRANCHING RATIOS

$\Gamma(N\bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.19 \pm 0.06	DEBELLEFON 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total}$ in $N\bar{K} \rightarrow \Lambda(2325) \rightarrow \Lambda\omega$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.06 \pm 0.02	¹ BACCARI 77	IPWA	DS_{33} wave	
0.05 \pm 0.02	¹ BACCARI 77	DPWA	DD_{13} wave	
0.08 \pm 0.03	¹ BACCARI 77	DPWA	DD_{33} wave	

 $\Lambda(2325)$ FOOTNOTES¹ Note that the three BACCARI 77 entries are for three different waves. $\Lambda(2325)$ REFERENCES

DEBELLEFON 78	NC 42A 403	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
BACCARI 77	NC 41A 96	+Poulard, Revel, Tallini+	(SACL, CDEF) IJP

 $\Lambda(2350) H_{09}$

$I(J^P) = 0(\frac{9}{2}^+)$ Status: ***

DAUM 68 favors $J^P = 7/2^-$ or $9/2^+$. BRICMAN 70 favors $9/2^+$. LASINSKI 71 suggests three states in this region using a Pomeron + resonances model. There are now also three formation experiments from the College de France-Saclay group, DEBELLEFON 77, BACCARI 77, and DEBELLEFON 78, which find $9/2^+$ in energy-dependent partial-wave analyses of $\bar{K} N \rightarrow \Sigma\pi, \Lambda\omega$, and $N\bar{K}$.

 $\Lambda(2350)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2350 OUR ESTIMATE			
2370 \pm 50	DEBELLEFON 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
2365 \pm 20	DEBELLEFON 77	DPWA	$K^- p \rightarrow \Sigma\pi$
2358 \pm 6	BRICMAN 70	CNTR	Total, charge exchange
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2372	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
2344 \pm 15	COOL 70	CNTR	$K^- p, K^- d$ total
2360 \pm 20	LU 70	CNTR	$\gamma p \rightarrow K^+ Y^*$
2340 \pm 7	BUGG 68	CNTR	$K^- p, K^- d$ total

 $\Lambda(2350)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
100 to 250 (≈ 180) OUR ESTIMATE			
204 \pm 50	DEBELLEFON 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
110 \pm 20	DEBELLEFON 77	DPWA	$K^- p \rightarrow \Sigma\pi$
324 \pm 30	BRICMAN 70	CNTR	Total, charge exchange
• • • We do not use the following data for averages, fits, limits, etc. • • •			
257	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$
190	COOL 70	CNTR	$K^- p, K^- d$ total
55	LU 70	CNTR	$\gamma p \rightarrow K^+ Y^*$
140 \pm 20	BUGG 68	CNTR	$K^- p, K^- d$ total

 $\Lambda(2350)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	$\sim 12\%$
Γ_2 $\Sigma\pi$	$\sim 10\%$
Γ_3 $\Lambda\omega$	

The above branching fractions are our estimates, not fits or averages.

 $\Lambda(2350)$ BRANCHING RATIOSSee "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
~ 0.12 OUR ESTIMATE				
0.12 \pm 0.04	DEBELLEFON 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{total}$ in $N\bar{K} \rightarrow \Lambda(2350) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
-0.11 \pm 0.02	DEBELLEFON 77	DPWA	$K^- p \rightarrow \Sigma\pi$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{total}$ in $N\bar{K} \rightarrow \Lambda(2350) \rightarrow \Lambda\omega$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
< 0.05	BACCARI 77	DPWA	$K^- p \rightarrow \Lambda\omega$	

 $\Lambda(2350)$ REFERENCES

DEBELLEFON 78	NC 42A 403	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
BACCARI 77	NC 41A 96	+Poulard, Revel, Tallini+	(SACL, CDEF) IJP
DEBELLEFON 77	NC 37A 175	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
LASINSKI 71	NP B29 125		(EFI) IJP
BRICMAN 70	PL 31B 152	+Ferro-Luzzi, Perreau+	(CERN, CAEN, SACL)
COOL 70	PR D1 1887	+Giacomelli, Kydon, Leontic, Li+	(BNL) I
Also	66	PR L 16 1228	Cool, Giacomelli, Kydon, Leontic, Lundby+
LU 70	PR D2 1846	+Greenberg, Hughes, Minehart, Mori+	(YALE)
BUGG 68	PR 168 1466	+Gimore, Knight+	(RHEL, BIRM, CAVE) I
DAUM 68	NP B7 19	+Erne, Lagnaux, Sens, Steuer, Udo	(CERN) JP

 $\Lambda(2585)$ Bumps

$I(J^P) = 0(?)^?$ Status: **

OMITTED FROM SUMMARY TABLE

 $\Lambda(2585)$ MASS (BUMPS)

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2585 OUR ESTIMATE			
2585 \pm 45	ABRAMS 70	CNTR	$K^- p, K^- d$ total
2530 \pm 25	LU 70	CNTR	$\gamma p \rightarrow K^+ Y^*$

 $\Lambda(2585)$ WIDTH (BUMPS)

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300	ABRAMS 70	CNTR	$K^- p, K^- d$ total
150	LU 70	CNTR	$\gamma p \rightarrow K^+ Y^*$

 $\Lambda(2585)$ DECAY MODES (BUMPS)

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	

See key on page 1343

Baryon Full Listings

$\Lambda(2585)$ Bumps, Σ^+

$\Lambda(2585)$ BRANCHING RATIOS (BUMPS)

$(J+\frac{1}{2}) \times \Gamma(N\bar{K}) / \Gamma_{total}$ Γ_1/Γ

J is not known, so only $(J+\frac{1}{2}) \times \Gamma(N\bar{K}) / \Gamma_{total}$ can be given.

VALUE	DOCUMENT ID	TECN	COMMENT
1	ABRAMS	70 CNTR	$K^- p, K^- d$ total
0.12 ± 0.12	1 BRICMAN	70 CNTR	Total, charge exchange

$\Lambda(2585)$ FOOTNOTES (BUMPS)

¹The resonance is at the end of the region analyzed — no clear signal.

$\Lambda(2585)$ REFERENCES (BUMPS)

ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL) I
Also	66	PRL 16 1228	Cool, Giacomelli, Kycia, Leontic, Lundby+	(BNL) I
BRICMAN	70	PL 31B 152	+Ferro-Luzzi, Perreau+	(CERN, CAEN, SACL)
LU	70	PR D2 1846	+Greenberg, Hughes, Minehart, Mori+	(YALE)

Σ BARYONS

$(S = -1, I = 1)$

$\Sigma^+ = uus, \Sigma^0 = uds, \Sigma^- = dds$

Σ^+

$I(J^P) = 1(\frac{1}{2}^+)$ Status: * * * *

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1986 edition (Physics Letters 170B) or in earlier editions.


Σ^+ MASS

The fit uses $\Sigma^+, \Sigma^0, \Sigma^-$, and Λ mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1189.37 ± 0.07 OUR FIT				Error includes scale factor of 2.2.
1189.37 ± 0.06 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
1189.33 ± 0.04	607	1 BOHM	72 EMUL	
1189.16 ± 0.12		HYMAN	67 HEBE	
1189.61 ± 0.08	4205	SCHMIDT	65 HBC	See note with Λ mass
1189.48 ± 0.22	58	2 BHOWMIK	64 EMUL	
1189.38 ± 0.15	144	2 BARKAS	63 EMUL	

¹BOHM 72 is updated with our 1973 $K^-, \pi^-,$ and π^0 masses (Reviews of Modern Physics 45 No. 2 Pt. II (1973)).

²These masses have been raised 30 keV to take into account a 46 keV increase in the proton mass and a 21 keV decrease in the π^0 mass (note added 1967 edition, Reviews of Modern Physics 39 1 (1967)).



Values above of weighted average, error, and scale factor are based upon the data in this ideogram only. They are not necessarily the same as our "best" values, obtained from a least-squares constrained fit utilizing measurements of other (related) quantities as additional information.

Σ^+ MEAN LIFE

Measurements with an error $\geq 0.1 \times 10^{-10}$ s have been omitted.

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.799 ± 0.004 OUR AVERAGE				
0.798 ± 0.005	30k	MARRAFFINO 80	HBC	$K^- p$ 0.42-0.5 GeV/c
0.807 ± 0.013	5719	CONFORTO 76	HBC	$K^- p$ 1-1.4 GeV/c
0.83 ± 0.04	526	BAKKER 71	DBC	$K^- n \rightarrow \Sigma^+ \pi^- \pi^-$
0.795 ± 0.010	20k	EISELE 70	HBC	$K^- p$ at rest
0.803 ± 0.008	10664	BARLOUTAUD 69	HBC	$K^- p$ 0.4-1.2 GeV/c
0.83 ± 0.032	1300	3 CHANG 66	HBC	
0.80 ± 0.07	381	COOK 66	OSPK	
0.84 ± 0.09	181	BALTAY 65	HBC	
0.76 ± 0.03	900	CARAYAN...	65 HBC	
0.749 ⁺ 0.056 -0.052	192	GRAD 62	HBC	
0.765 ± 0.04	456	HUMPHREY 62	HBC	

³We have increased the CHANG 66 error of 0.018; see our 1970 edition, Reviews of Modern Physics 42 No. 1 (1970).

Σ^+ MAGNETIC MOMENT


See the Note on Baryon Magnetic Moments in the Λ Listings. Measurements with an error $\geq 0.1 \mu_N$ have been omitted.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN	COMMENT
2.458 ± 0.010 OUR AVERAGE				Error includes scale factor of 2.1. See the ideogram below.
2.4613 ± 0.0034 ± 0.0040	250k	MORELOS 93	SPEC	p Cu 800 GeV
2.428 ± 0.036 ± 0.007	12k	4 MORELOS 93	SPEC	p Cu 800 GeV
2.479 ± 0.012 ± 0.022	137k	WILKINSON 87	SPEC	p Be 400 GeV
2.4040 ± 0.0198	44k	5 ANKENBRA... 83	CNTR	p Cu 400 GeV

⁴We assume CPT invariance: this is (minus) the Σ^- magnetic moment as measured by MORELOS 93. See below for the moment difference testing CPT.

⁵ANKENBRANDT 83 gives the value $2.38 \pm 0.02 \mu_N$. MORELOS 93 uses the same hyperon magnet and channel and claims to determine the field integral better, leading to the revised value given here.

WEIGHTED AVERAGE
2.458 ± 0.010 (Error scaled by 2.1)



$(\mu_{\Sigma^+} - |\mu_{\Sigma^-}|) / |\mu|_{average}$

A test of CPT invariance.

VALUE	DOCUMENT ID	TECN	COMMENT
0.014 ± 0.015	6 MORELOS 93	SPEC	p Cu 800 GeV

⁶This is our calculation from the MORELOS 93 measurements of the Σ^+ and Σ^- magnetic moments given above. The statistical error on μ_{Σ^-} dominates the error here.

Baryon Full Listings

 Σ^+ Σ^+ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 $p\pi^0$	$(51.57 \pm 0.30) \%$	
Γ_2 $n\pi^+$	$(48.30 \pm 0.30) \%$	
Γ_3 $p\gamma$	$(1.25 \pm 0.07) \times 10^{-3}$	
Γ_4 $n\pi^+\gamma$	[a] $(4.5 \pm 0.5) \times 10^{-4}$	
Γ_5 $\Lambda e^+\nu_e$	$(2.0 \pm 0.5) \times 10^{-5}$	

$\Delta S = \Delta Q$ (SQ) violating modes or
 $\Delta S = 1$ weak neutral current (SI) modes

Γ_6 $ne^+\nu_e$	SQ	< 5	$\times 10^{-6}$	90%
Γ_7 $n\mu^+\nu_\mu$	SQ	< 3.0	$\times 10^{-5}$	90%
Γ_8 pe^+e^-	SI	< 7	$\times 10^{-6}$	

[a] See the Full Listings below for the pion momentum range used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 13 measurements and one constraint to determine 3 parameters. The overall fit has a $\chi^2 = 7.5$ for 11 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-100		
x_3	9	-11	
	x_1	x_2	

 Σ^+ BRANCHING RATIOS

$\Gamma(n\pi^+)/\Gamma(N\pi)$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	$\Gamma_2/(\Gamma_1+\Gamma_2)$
0.4836 ± 0.0030 OUR FIT						
0.4836 ± 0.0030 OUR AVERAGE						
0.4828 ± 0.0036	10k	7	MARRAFFINO 80	HBC	$K^- p$ 0.42–0.5 GeV/c	
0.488 ± 0.008	1861		NOWAK 78	HBC		
0.484 ± 0.015	537		TOVEE 71	EMUL		
0.488 ± 0.010	1331		BARLOUTAUD 69	HBC	$K^- p$ 0.4–1.2 GeV/c	
0.46 ± 0.02	534		CHANG 66	HBC		
0.490 ± 0.024	308		HUMPHREY 62	HBC		

⁷ MARRAFFINO 80 actually gives $\Gamma(p\pi^0)/\Gamma(\text{total}) = 0.5172 \pm 0.0036$.

$\Gamma(p\gamma)/\Gamma(p\pi^0)$	VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
2.43 ± 0.14 OUR FIT						
2.43 ± 0.14 OUR AVERAGE						
$2.81 \pm 0.39^{+0.21}_{-0.43}$	408		HESSEY 89	CNTR	$K^- p \rightarrow \Sigma^+ \pi^-$ at rest	
2.52 ± 0.28	190	8	KOBAYASHI 87	CNTR	$\pi^+ p \rightarrow \Sigma^+ K^+$	
$2.46^{+0.30}_{-0.35}$	155		BIAGI 85	CNTR	CERN hyperon beam	
2.11 ± 0.38	46		MANZ 80	HBC	$K^- p \rightarrow \Sigma^+ \pi^-$	
2.1 ± 0.3	45		ANG 69B	HBC	$K^- p$ at rest	
2.76 ± 0.51	31		GERSHWIN 69B	HBC	$K^- p \rightarrow \Sigma^+ \pi^-$	
3.7 ± 0.8	24		BAZIN 65	HBC	$K^- p$ at rest	

⁸ KOBAYASHI 87 actually gives $\Gamma(p\gamma)/\Gamma(\text{total}) = (1.30 \pm 0.15) \times 10^{-3}$.

$\Gamma(n\pi^+\gamma)/\Gamma(n\pi^+)$	VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_2
0.93 ± 0.10	180		EBENHOH 73	HBC	$\pi^+ < 150$ MeV/c	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.27 ± 0.05	29		ANG 69B	HBC	$\pi^+ < 110$ MeV/c	
~ 1.8			BAZIN 65B	HBC	$\pi^+ < 116$ MeV/c	

$\Gamma(\Lambda e^+\nu_e)/\Gamma_{\text{total}}$	VALUE (units 10^{-5})	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
2.0 ± 0.5 OUR AVERAGE						
1.6 ± 0.7	5		BALTAY 69	HBC	$K^- p$ at rest	
2.9 ± 1.0	10		EISELE 69	HBC	$K^- p$ at rest	
2.0 ± 0.8	6		BARASH 67	HBC	$K^- p$ at rest	

$\Gamma(ne^+\nu_e)/\Gamma(n\pi^+)$ Γ_6/Γ_2
Test of $\Delta S = \Delta Q$ rule. Experiments with an effective denominator less than 100,000 have been omitted.

EFFECTIVE DENOM.	EVTs	DOCUMENT ID	TECN	COMMENT
< 1.1 × 10⁻⁵ OUR LIMIT				Our 90% CL limit = (2.3 events)/(effective denominator sum). [Number of events increased to 2.3 for a 90% confidence level.]
111000	0	⁹ EBENHOH 74	HBC	$K^- p$ at rest
105000	0	⁹ SECHI-ZORN 73	HBC	$K^- p$ at rest

⁹ Effective denominator calculated by us.

$\Gamma(n\mu^+\nu_\mu)/\Gamma(n\pi^+)$ Γ_7/Γ_2
Test of $\Delta S = \Delta Q$ rule.

EFFECTIVE DENOM.	EVTs	DOCUMENT ID	TECN	COMMENT
< 6.2 × 10⁻⁵ OUR LIMIT				Our 90% CL limit = (6.7 events)/(effective denominator sum). [Number of events increased to 6.7 for a 90% confidence level.]
33800	0	BAGGETT 69B	HBC	
62000	2	¹⁰ EISELE 69B	HBC	
10150	0	¹¹ COURANT 64	HBC	
1710	0	¹¹ NAUENBERG 64	HBC	
120	1	GALTIERI 62	EMUL	

¹⁰ Effective denominator calculated by us.

¹¹ Effective denominator taken from EISELE 67.

$\Gamma(pe^+e^-)/\Gamma_{\text{total}}$ Γ_8/Γ

VALUE (units 10^{-6})	DOCUMENT ID	TECN	COMMENT
< 7	¹² ANG 69B	HBC	$K^- p$ at rest

¹² ANG 69B found three pe^+e^- events in agreement with $\gamma \rightarrow e^+e^-$ conversion from $\Sigma^+ \rightarrow p\gamma$. The limit given here is for neutral currents.

$\Gamma(\Sigma^+ \rightarrow ne^+\nu_e)/\Gamma(\Sigma^- \rightarrow ne^-\nu_e)$

VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
< 0.009 OUR LIMIT					Our 90% CL limit, using $\Gamma(ne^+\nu_e)/\Gamma(n\pi^+)$ above.
• • •					We do not use the following data for averages, fits, limits, etc. • • •
< 0.019	90	0	EBENHOH 74	HBC	$K^- p$ at rest
< 0.018	90	0	SECHI-ZORN 73	HBC	$K^- p$ at rest
< 0.12	95	0	COLE 71	HBC	$K^- p$ at rest
< 0.03	90	0	EISELE 69B	HBC	See EBENHOH 74

$\Gamma(\Sigma^+ \rightarrow n\mu^+\nu_\mu)/\Gamma(\Sigma^- \rightarrow n\mu^-\nu_\mu)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
< 0.12 OUR LIMIT				Our 90% CL limit, using $\Gamma(n\mu^+\nu_\mu)/\Gamma(n\pi^+)$ above.
• • •				We do not use the following data for averages, fits, limits, etc. • • •
$0.06^{+0.045}_{-0.03}$	2	EISELE 69B	HBC	$K^- p$ at rest

$\Gamma(\Sigma^+ \rightarrow n\ell^+\nu)/\Gamma(\Sigma^- \rightarrow n\ell^-\nu)$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
< 0.043 OUR LIMIT				Our 90% CL limit, using $[\Gamma(ne^+\nu_e) + \Gamma(n\mu^+\nu_\mu)]/\Gamma(n\pi^+)$.
< 0.08	1	NORTON 69	HBC	
< 0.034	0	BAGGETT 67	HBC	

• • • We do not use the following data for averages, fits, limits, etc. • • •

 Σ^+ DECAY PARAMETERS

See the Note on Baryon Decay Parameters in the neutron Listings. A few early results have been omitted.

α_0 FOR $\Sigma^+ \rightarrow p\pi^0$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.980^{+0.017}_{-0.015} OUR FIT				
-0.980^{+0.017}_{-0.013} OUR AVERAGE				
$-0.945^{+0.055}_{-0.042}$	1259	¹³ LIPMAN 73	OSPK	$\pi^+ p \rightarrow \Sigma^+$
-0.940 ± 0.045	16k	BELLAMY 72	ASPK	$\pi^+ p \rightarrow \Sigma^+ K^+$
$-0.98^{+0.05}_{-0.02}$	1335	¹⁴ HARRIS 70	OSPK	$\pi^+ p \rightarrow \Sigma^+ K^+$
-0.999 ± 0.022	32k	BANGERTER 69	HBC	$K^- p$ 0.4 GeV/c

¹³ Decay protons scattered off aluminum.

¹⁴ Decay protons scattered off carbon.

ϕ_0 ANGLE FOR $\Sigma^+ \rightarrow p\pi^0$

VALUE (°)	EVTs	DOCUMENT ID	TECN	COMMENT
36 ± 34 OUR AVERAGE				
$38.1^{+35.7}_{-37.1}$	1259	¹⁵ LIPMAN 73	OSPK	$\pi^+ p \rightarrow \Sigma^+ K^+$
22 ± 90		¹⁶ HARRIS 70	OSPK	$\pi^+ p \rightarrow \Sigma^+ K^+$

¹⁵ Decay proton scattered off aluminum.

¹⁶ Decay protons scattered off carbon.

($\tan \phi_0 = \beta/\gamma$)

See key on page 1343

Baryon Full Listings

 Σ^+, Σ^0 α_+ / α_0

Older results have been omitted.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.069 ± 0.013				OUR FIT
-0.073 ± 0.021	23k	MARRAFFINO 80	HBC	$K^- p$ 0.42-0.5 GeV/c

 α_+ FOR $\Sigma^+ \rightarrow n\pi^+$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.068 ± 0.013				OUR FIT
0.066 ± 0.016				OUR AVERAGE
0.037 ± 0.049	4101	BERLEY 70B	HBC	
0.069 ± 0.017	35k	BANGERTER 69	HBC	$K^- p$ 0.4 GeV/c

 ϕ_+ ANGLE FOR $\Sigma^+ \rightarrow n\pi^+$ $(\tan \phi_+ = \beta/\gamma)$

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
167 ± 20				OUR AVERAGE Error includes scale factor of 1.1.
184 ± 24	1054	17 BERLEY 70B	HBC	
143 ± 29	560	BANGERTER 69B	HBC	$K^- p$ 0.4 GeV/c

17 Changed from 176 to 184° to agree with our sign convention.

 α_+ FOR $\Sigma^+ \rightarrow p\gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.76 ± 0.08				OUR AVERAGE
$-0.720 \pm 0.086 \pm 0.045$				
$-0.86 \pm 0.13 \pm 0.04$	190	KOBAYASHI 87	CNTR	$\pi^+ p \rightarrow \Sigma^+ K^+$
-0.53 ± 0.38	46	MANZ 80	HBC	$K^- p \rightarrow \Sigma^+ \pi^-$
-0.52				
-1.03 ± 0.52	61	GERSHWIN 69B	HBC	$K^- p \rightarrow \Sigma^+ \pi^-$
-0.42				

 Σ^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1986 edition (Physics Letters **170B**) or in earlier editions.

MORELOS	93	PRL 71 3417	+Albuquerque, Bondar, Carrigan+	(FNAL E761 Collab.)
FOUCHER	92	PRL 68 3004	+Albuquerque, Bondar+	(FNAL E761 Collab.)
HESSY	89	ZPHY C42 175	+Booth, Fickinger, Gall+	(BNL-811 Collab.)
KOBAYASHI	87	PRL 59 868	+Haba, Homma, Kawai, Miyake+	(KYOT)
WILKINSON	87	PRL 58 865	+Handler+	(WISC, MICH, RUTG, MINN)
BIAGI	85	ZPHY C28 495	+Bourquin+	(CERN WA62 Collab.)
ANKENBRA...	83	PRL 51 863	+Ankenbrandt, Berge+	(FNAL, IOWA, ISU, YALE)
MANZ	80	PL 96B 217	+Reucroft, Settles, Wolf+	(MPIM, WAND)
MARRAFFINO	80	PR D21 2501	+Reucroft, Roos, Waters+	(VAND, MPIM)
NOWAK	78	NP B139 61	+Armstrong, Davis+	(LOUC, BELG, DURH, WARS)
CONFORTO	76	NP B105 189	+Gopal, Kaimus, Litchfield, Ross+	(RHEL, LOIC)
EBENHOH	74	ZPHY 266 367	+Eisele, Engelmann, Filthuth, Hepp+	(HEIDT)
EBENHOH	73	ZPHY 264 413	+Eisele, Filthuth, Hepp, Leitner, Thow+	(HEIDT)
LIPMAN	73	PL 43B 89	+Uto, Walker, Montgomery+	(RHEL, SUSS, LOWC)
PDG	73	RMP 45 No. 2 Pt. II	+Lasinski, Barbaro-Galtieri, Kely+	(LBL, BRAN, CERN+)
SECHI-ZORN	73	PR D8 12	+Snow	(UMD)
BELLAMY	72	PL 39B 299	+Anderson, Crawford+	(LOWC, RHEL, SUSS)
BOHM	72	NP B48 1	+Bohm	(BERL, KIDR, BRUX, IASD, DUUC, LOUC+)
Also	73	IHHE-73.2 Nov	+Hoogland, Kluyver, Massard+	(SABRE Collab.)
BAKKER	71	LNC 1 37	+Lee-Franzini, Loveless, Baltay+	(STON, COLU)
COLE	71	PR D4 631	+ (LOUC, KIDR, BERL, BRUX, DUUC, WARS)	
TOVEE	71	NP B33 493	+Yamin, Hertzbach, Koller+	(BNL, MASA, YALE)
BERLEY	70B	PR D1 2015	+Filthuth, Hepp, Presser, Zech	(HEID)
EISELE	70	ZPHY 238 372	+Overseht, Pondrom, Dettmann	(MICH, WISC)
HARRIS	70	PRL 24 165	+Barbaro-Galtieri, Derenzo, Price+	(LRL, BRAN, CERN+)
PDG	70	RMP 42 No. 1	+Ebenhoh, Eisele, Engelmann, Filthuth+	(HEID)
ANG	69B	ZPHY 228 151	+Franzini, Newman, Norton+	(COLU, STON)
BAGGETT	69B	MDDP-TR-973 Thesis	+Alston-Garnjost, Galtieri, Gershwin+	(LRL)
BALTAY	69	PRL 22 615	+DeBellefon, Granet+	(SACL, CERN, HEID)
BANGERTER	69	UCRL 19244 Thesis	+Engelmann, Filthuth, Fohlisch, Hepp+	(HEID)
BANGERTER	69B	PR 187 1821	+Willis, Courant+	(BNL, CERN, HEID, UMD)
BARLOUTAUD	69	NP B14 153	+Engelmann, Filthuth, Fohlisch, Hepp+	(HEID)
EISELE	69	ZPHY 221 1	+Alston-Garnjost, Bangerter+	(LRL)
Also	64	PRL 13 291	+Gershwin	(PURL)
EISELE	69B	ZPHY 221 401		(COLU)
GERSHWIN	69B	PR 188 2077	+Day, Glasser, Kehoe, Knop+	(UMD)
Also	69	UCRL 19246 Thesis	+Baggett, Kehoe	(UMD)
NORTON	69	Nevis 175 Thesis	+Baggett	(UMD)
BAGGETT	67	PRL 19 1458	+Day, Glasser, Kehoe, Knop+	(UMD)
Also	68	Vienna Abs. 374	+Engelmann, Filthuth, Fohlisch, Hepp+	(HEID)
Also	68B	Private Comm.	+Loken, Hewitt, McKenzie+	(ANL, CMU, NWES)
BARASH	67	PRL 19 181	+Rosenfeld, Barbaro-Galtieri, Podolsky+	(LRL, CERN, YALE)
EISELE	67	ZPHY 205 409		(COLU)
HYMAN	67	PL 25B 376	+Chang	(COLU)
PDG	67	RMP 39 1	+Ewart, Masek, Orr, Platner	(WASH)
CHANG	66	PR 151 1081	+Sandweiss, Culwick, Kopp+	(YALE, BNL)
Also	65	Nevis 145 Thesis	+Blumenfeld, Nauenberg+	(PRIN, COLU)
COOK	66	PRL 17 223	+Plano, Schmidt+	(PRIN, RUTG, COLU)
BALTAY	65	PR 140B 1027	+Carayannopoulos, Taufest, Willmann	(PURL)
BAZIN	65	PRL 14 154		(COLU)
BAZIN	65B	PR 140B 1358	+Jain, Mathur, Lakshmi	(DELH)
CARAYAN...	65	PR 135B 433	+Filthuth+	(CERN, HEID, UMD, NRL, BNL)
SCHMIDT	65	PR 140B 1328	+Maratec+	(COLU, RUTG, PRIN)
BHOWMIK	64	NP 53 22	+Dyer, Heckman	(LRL)
COURANT	64	PR 136B 1791	+Dyer	(LRL)
NAUENBERG	64	PRL 12 679	+Barbas, Heckman, Patrick, Smith	(LRL)
BARKAS	63	PRL 11 26	+Smith	(LRL)
Also	61	UCRL 9450 Thesis	+Rosa	(LRL)
GALTIERI	62	PR 127 607		
GRARD	62	PR 127 607		
HUMPHREY	62	PR 127 1305		

 Σ^0 $I(J^P) = 1(\frac{1}{2}^+)$ Status: * * * *

The spin and parity have not been measured directly. They are of course assumed to be the same as for the Σ^+ and Σ^- .

 Σ^0 MASSThe fit uses Σ^+ , Σ^0 , Σ^- , and Λ mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1192.55 ± 0.08				OUR FIT Error includes scale factor of 1.2.
4.88 ± 0.08				OUR FIT Error includes scale factor of 1.2.
4.86 ± 0.08				OUR AVERAGE Error includes scale factor of 1.2.
4.87 ± 0.12	37	DOSCH 65	HBC	
5.01 ± 0.12	12	SCHMIDT 65	HBC	See note with Λ mass
4.75 ± 0.1	18	BURNSTEIN 64	HBC	

 $m_{\Sigma^0} - m_{\Lambda}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
76.87 ± 0.08				OUR FIT Error includes scale factor of 1.2.
76.55 ± 0.25				OUR AVERAGE
76.23 ± 0.55	109	COLAS 75	HLBC	$\Sigma^0 \rightarrow \Lambda\gamma$
76.63 ± 0.28	208	SCHMIDT 65	HBC	See note with Λ mass

 Σ^0 MEAN LIFE

These lifetimes are deduced from measurements of the cross sections for the Primakoff process $\Lambda \rightarrow \Sigma^0$ in nuclear Coulomb fields. An alternative expression of the same information is the Σ^0 - Λ transition magnetic moment given in the following section. The relation is $(\mu_{\Sigma^0/\mu_N})^2 \tau = 1.92951 \times 10^{-19}$ s (see DEVLIN 86).

VALUE (10^{-20} s)	DOCUMENT ID	TECN	COMMENT
7.4 ± 0.7			OUR EVALUATION Using μ_{Σ^0/μ_N} (see the above note).
6.5 ± 1.7	1 DEVLIN 86	SPEC	Primakoff effect
-1.1			
7.6 ± 0.5 ± 0.7	2 PETERSEN 86	SPEC	Primakoff effect
• • • We do not use the following data for averages, fits, limits, etc. • • •			
5.8 ± 1.3	1 DYDAK 77	SPEC	See DEVLIN 86
1 DEVLIN 86 is a recalculation of the results of DYDAK 77 removing a numerical approximation made in that work.			
2 An additional uncertainty of the Primakoff formalism is estimated to be < 5%.			

 $|\mu(\Sigma^0 \rightarrow \Lambda)|$ TRANSITION MAGNETIC MOMENT

See the note in the Σ^0 mean-life section above. Also, see the Note on Baryon Magnetic Moments in the Λ Listings.

VALUE (μ_N)	DOCUMENT ID	TECN	COMMENT
1.61 ± 0.08			OUR AVERAGE
1.72 ± 0.17	3 DEVLIN 86	SPEC	Primakoff effect
-0.19			
1.59 ± 0.05 ± 0.07	4 PETERSEN 86	SPEC	Primakoff effect
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1.82 ± 0.25	3 DYDAK 77	SPEC	See DEVLIN 86
-0.18			
3 DEVLIN 86 is a recalculation of the results of DYDAK 77 removing a numerical approximation made in that work.			
4 An additional uncertainty of the Primakoff formalism is estimated to be < 2.5%.			

 Σ^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \Lambda\gamma$	100 %	
$\Gamma_2 \Lambda\gamma\gamma$	< 3 %	90%
$\Gamma_3 \Lambda e^+ e^-$	[a] 5×10^{-3}	

[a] A theoretical value using QED.

 Σ^0 BRANCHING RATIOS

VALUE	CL%	DOCUMENT ID	TECN	Γ_2/Γ
< 0.03	90	COLAS 75	HLBC	
VALUE	DOCUMENT ID	COMMENT	Γ_3/Γ	
0.00545	FEINBERG 58	Theoretical QED calculation		

Baryon Full Listings

Σ^0, Σ^-

Σ^0 REFERENCES

DEVLIN 86 PR D34 1626	+Petersen, Beretvas	(RUTG)
PETERSEN 86 PRL 57 949	+Beretvas, Devlin, Luk+	(RUTG, WISC, MICH, MINN)
DYDAK 77 NP B118 1	+Navarra, Overseth, Steffen+	(CERN, DORT, HEIDH)
COLAS 75 NP B91 253	+Farwell, Ferrer, Six	(ORSAY)
DOSCH 65 PL 14 239	+Engelmann, Filthuth, Hepp, Kluge+	(HEID)
SCHMIDT 65 PR 140B 1328		(COLU)
BURNSTEIN 64 PRL 13 66	+Day, Kehoe, Zorn, Snow	(UMD)
FEINBERG 58 PR 109 1019		(BNL)

Σ^-

$I(J^P) = 1(\frac{1}{2}^+)$ Status: * * * *

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1986 edition (Physics Letters 170B) or in earlier editions.

Σ^- MASS

The fit uses Σ^+ , Σ^0 , Σ^- , and Λ mass and mass-difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1197.436 ± 0.033 OUR FIT				Error includes scale factor of 1.2.
1197.45 ± 0.04 OUR AVERAGE				Error includes scale factor of 1.2.
1197.417 ± 0.040		GUREV 93 SPEC	Σ^- C atom, crystal diff.	
1197.532 ± 0.057		GALL 88 CNTR	Σ^- Pb, Σ^- W atoms	
1197.43 ± 0.08	3000	SCHMIDT 65 HBC	See note with Λ mass	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1197.24 ± 0.15		¹ DUGAN 75 CNTR	Exotic atoms	
¹ GALL 88 concludes that the DUGAN 75 mass needs to be reevaluated.				

$m_{\Sigma^-} - m_{\Sigma^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
8.07 ± 0.08 OUR FIT				Error includes scale factor of 1.9.
8.09 ± 0.16 OUR AVERAGE				
7.91 ± 0.23	86	BOHM 72 EMUL		
8.25 ± 0.25	2500	DOSCH 65 HBC		
8.25 ± 0.40	87	BARKAS 63 EMUL		

$m_{\Sigma^-} - m_{\Lambda}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
81.752 ± 0.034 OUR FIT				Error includes scale factor of 1.2.
81.69 ± 0.07 OUR AVERAGE				
81.64 ± 0.09	2279	HEPP 68 HBC		
81.80 ± 0.13	85	SCHMIDT 65 HBC		See note with Λ mass
81.70 ± 0.19		BURNSTEIN 64 HBC		


Σ^- MEAN LIFE

Measurements with an error $\geq 0.2 \times 10^{-10}$ s have been omitted.

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.479 ± 0.011 OUR AVERAGE				Error includes scale factor of 1.3. See the ideogram below.
1.480 ± 0.014	16k	MARRAFFINO 80 HBC	$K^- p$ 0.42-0.5 GeV/c	
1.49 ± 0.03	8437	CONFORTO 76 HBC	$K^- p$ 1-1.4 GeV/c	
1.463 ± 0.039	2400	ROBERTSON 72 HBC	$K^- p$ 0.25 GeV/c	
1.42 ± 0.05	1383	BAKKER 71 DBC	$K^- N \rightarrow \Sigma^- \pi \pi$	
1.41 ^{+0.09} _{-0.08}		TOVEE 71 EMUL		
1.485 ± 0.022	100k	EISELE 70 HBC	$K^- p$ at rest	
1.472 ± 0.016	10k	BARLOUTAUD 69 HBC	$K^- p$ 0.4-1.2 GeV/c	
1.38 ± 0.07	506	WHITESIDE 68 HBC	$K^- p$ at rest	
1.666 ± 0.075	3267	² CHANG 66 HBC	$K^- p$ at rest	
1.58 ± 0.06	1208	HUMPHREY 62 HBC	$K^- p$ at rest	

²We have increased the CHANG 66 error of 0.018; see our 1970 edition, Reviews of Modern Physics 42 No. 1 (1970).

WEIGHTED AVERAGE
1.479 ± 0.011 (Error scaled by 1.3)




Σ^- MAGNETIC MOMENT

See the Note on Baryon Magnetic Moments in the Λ Listings. Measurements with an error $\geq 0.3 \mu_N$ have been omitted.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN	COMMENT
-1.160 ± 0.025 OUR AVERAGE				Error includes scale factor of 1.7. See the ideogram below.
-1.105 ± 0.029 ± 0.010		HERTZOG 88 CNTR	Σ^- Pb, Σ^- W atoms	
-1.166 ± 0.014 ± 0.010	671k	ZAPALAC 86 SPEC	$n e^- \nu, n \pi^-$ decays	
-1.23 ± 0.03 ± 0.03		WAH 85 CNTR	p Cu $\rightarrow \Sigma^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.89 ± 0.14	516k	DECK 83 SPEC	p Be $\rightarrow \Sigma^- X$	

WEIGHTED AVERAGE
-1.160 ± 0.025 (Error scaled by 1.7)



Σ^- DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $n\pi^-$	(99.848 ± 0.005) %
Γ_2 $n\pi^- \gamma$	[a] (4.6 ± 0.6) × 10 ⁻⁴
Γ_3 $n e^- \bar{\nu}_e$	(1.017 ± 0.034) × 10 ⁻³
Γ_4 $n \mu^- \bar{\nu}_\mu$	(4.5 ± 0.4) × 10 ⁻⁴
Γ_5 $\Lambda e^- \bar{\nu}_e$	(5.73 ± 0.27) × 10 ⁻⁵

[a] See the Full Listings below for the pion momentum range used in this measurement.

CONSTRAINED FIT INFORMATION

An overall fit to 3 branching ratios uses 16 measurements and one constraint to determine 4 parameters. The overall fit has a $\chi^2 = 8.7$ for 13 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_3	-64		
x_4	-77	0	
x_5	-5	0	0
	x_1	x_3	x_4

 Σ^- BRANCHING RATIOS $\Gamma(n\pi^- \gamma) / \Gamma(n\pi^-)$ Γ_2 / Γ_1

The π^+ momentum cuts differ, so we do not average the results but simply use the latest value for the Summary Table.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.46 ± 0.06	292	EBENHOH	73 HBC	$\pi^+ < 150$ MeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.10 ± 0.02	23	ANG	69B HBC	$\pi^- < 110$ MeV/c
~ 1.1		BAZIN	65B HBC	$\pi^- < 166$ MeV/c

 $\Gamma(ne^- \bar{\nu}_e) / \Gamma(n\pi^-)$ Γ_3 / Γ_1

Measurements with an error $\geq 0.2 \times 10^{-3}$ have been omitted.

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
1.019 ± 0.034 OUR FIT				
1.019 ± 0.031 OUR AVERAGE				
0.96 ± 0.05	2847	BOURQUIN	83C SPEC	SPS hyperon beam
1.09 ± 0.06	601	³ EBENHOH	74 HBC	$K^- p$ at rest
-0.08				
1.05 ± 0.07	455	³ SECHI-ZORN	73 HBC	$K^- p$ at rest
-0.13				
0.97 ± 0.15	57	COLE	71 HBC	$K^- p$ at rest
1.11 ± 0.09	180	BIERMAN	68 HBC	

³ An additional negative systematic error is included for internal radiative corrections and latest form factors; see BOURQUIN 83C.

 $\Gamma(n\mu^- \bar{\nu}_\mu) / \Gamma(n\pi^-)$ Γ_4 / Γ_1

VALUE (units 10^{-3})	EVTS	DOCUMENT ID	TECN	COMMENT
0.45 ± 0.04 OUR FIT				
0.45 ± 0.04 OUR AVERAGE				
0.38 ± 0.11	13	COLE	71 HBC	$K^- p$ at rest
0.43 ± 0.06	72	ANG	69 HBC	$K^- p$ at rest
0.43 ± 0.09	56	BAGGETT	69 HBC	$K^- p$ at rest
0.56 ± 0.20	11	BAZIN	65B HBC	$K^- p$ at rest
0.66 ± 0.15	22	COURANT	64 HBC	

 $\Gamma(\Lambda e^- \bar{\nu}_e) / \Gamma(n\pi^-)$ Γ_5 / Γ_1

VALUE (units 10^{-4})	EVTS	DOCUMENT ID	TECN	COMMENT
0.574 ± 0.027 OUR FIT				
0.574 ± 0.027 OUR AVERAGE				
0.561 ± 0.031	1620	⁴ BOURQUIN	82 SPEC	SPS hyperon beam
0.63 ± 0.11	114	THOMPSON	80 ASPK	Hyperon beam
0.52 ± 0.09	31	BALTAY	69 HBC	$K^- p$ at rest
0.69 ± 0.12	31	EISELE	69 HBC	$K^- p$ at rest
0.64 ± 0.12	35	BARASH	67 HBC	$K^- p$ at rest
0.75 ± 0.28	11	COURANT	64 HBC	$K^- p$ at rest

⁴ The value is from BOURQUIN 83B, and includes radiation corrections and new acceptance.

 Σ^- DECAY PARAMETERS

See the Note on Baryon Decay Parameters in the neutron Listings. Older, outdated results have been omitted.

 α_- FOR $\Sigma^- \rightarrow n\pi^-$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.068 ± 0.008 OUR AVERAGE				
-0.062 ± 0.024	28k	HANSL	78 HBC	$K^- p \rightarrow \Sigma^- \pi^+$
-0.067 ± 0.011	60k	BOGERT	70 HBC	$K^- p$ 0.4 GeV/c
-0.071 ± 0.012	51k	BANGERTER	69 HBC	$K^- p$ 0.4 GeV/c

 ϕ ANGLE FOR $\Sigma^- \rightarrow n\pi^-$ $(\tan \phi = \beta / \gamma)$

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
10 ± 15 OUR AVERAGE				
+ 5 ± 23	1092	⁵ BERLEY	70B HBC	n rescattering
14 ± 19	1385	BANGERTER	69B HBC	$K^- p$ 0.4 GeV/c

⁵ BERLEY 70B changed from -5 to +5° to agree with our sign convention.

 g_A/g_V FOR $\Sigma^- \rightarrow ne^- \bar{\nu}_e$

Measurements with fewer than 500 events have been omitted. Where necessary, signs have been changed to agree with our conventions, which are given in the Note on Baryon Decay Parameters in the neutron Listings. What is actually listed is $g_1/f_1 - 0.237g_2/f_1$. This reduces to $g_A/g_V \equiv g_1(0)/f_1(0)$ on making the usual assumption that $g_2 = 0$. See also the note on HSUEH 88.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.340 ± 0.017 OUR AVERAGE				
+0.327 ± 0.007 ± 0.019	50k	⁶ HSUEH	88 SPEC	Σ^- 250 GeV
+0.34 ± 0.05	4456	⁷ BOURQUIN	83C SPEC	SPS hyperon beam
0.385 ± 0.037	3507	⁸ TANENBAUM	74 ASPK	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.29 ± 0.07	25k	HSUEH	85 SPEC	See HSUEH 88
0.17 ± 0.07	519	DECAMP	77 ELEC	Hyperon beam
-0.09				

⁶ The sign is, with our conventions, unambiguously positive. The value assumes, as usual, that $g_2 = 0$. If g_2 is included in the fit, then (with our sign convention) $g_2 = -0.56 \pm 0.37$, with a corresponding reduction of g_A/g_V to $+0.20 \pm 0.08$.

⁷ BOURQUIN 83C favors the positive sign by at least 2.6 standard deviations.

⁸ TANENBAUM 74 gives 0.435 ± 0.035 , assuming no q^2 dependence in g_A and g_V . The listed result allows q^2 dependence, and is taken from HSUEH 88.

 $f_2(0)/f_1(0)$ FOR $\Sigma^- \rightarrow ne^- \bar{\nu}_e$

The signs have been changed to be in accord with our conventions, given in the Note on Baryon Decay Parameters in the neutron Listings.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.97 ± 0.14 OUR AVERAGE				
+0.96 ± 0.07 ± 0.13	50k	HSUEH	88 SPEC	Σ^- 250 GeV
+1.02 ± 0.34	4456	BOURQUIN	83C SPEC	SPS hyperon beam

TRIPLE CORRELATION COEFFICIENT D for $\Sigma^- \rightarrow ne^- \bar{\nu}_e$

The coefficient D of the term $DP \cdot (\hat{p}_e \times \hat{p}_\nu)$ in the $\Sigma^- \rightarrow ne^- \bar{\nu}$ decay angular distribution. A nonzero value would indicate a violation of time-reversal invariance.

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.11 ± 0.10	50k	HSUEH	88 SPEC	Σ^- 250 GeV

NOTE ON $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$ DECAY

The vector part of the hadronic amplitude for the decay $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$ is of special interest because the vector weak current is proportional to an isospin rotation of the isovector part of the electromagnetic current. This strong form of CVC predicts that

$$f_1(q^2) = 0 \quad \text{for } 0 < q^2 \leq (m_{\Sigma^-} - m_{\Lambda})^2,$$

and also relates $f_2(0)$ to the $\Sigma^0 \Lambda$ transition magnetic moment or to the amplitude for the decay $\Sigma^0 \rightarrow \Lambda \gamma$ by

$$\begin{aligned} f_2(0) &= -\sqrt{2} \mu_{\Sigma^0 \Lambda} / e\hbar \\ &= -\sqrt{3/2} \mu_n / e\hbar \quad [\text{by SU}(3)] \\ &= 1.17 m_p^{-1}. \end{aligned}$$

No SU(3) symmetry is assumed here except in the relation of $\mu_{\Sigma^0 \Lambda}$ to the magnetic moment of the neutron, μ_n .

The experimental data were analyzed on the assumption that $f_1(q^2) = 0$ and $f_2(q^2) = f_2(0)$ over the entire kinematical range of q^2 for $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$. The results are listed in the ratio of $g_{WM} = -m_{\Sigma^-} f_2(0)$ to $g_A = g_1(0)$.

See also the Note on Baryon Decay Parameters in the neutron section of the Full Listings.

 g_V/g_A FOR $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$

For the sign convention, see the Note on Baryon Decay Parameters in the neutron Listings. The value is predicted to be zero by conserved vector current theory. The values averaged assume CVC-SU(3) weak magnetism term.


VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.01 ± 0.10 OUR AVERAGE				Error includes scale factor of 1.5. See the Ideogram below.
-0.034 ± 0.080	1620	⁹ BOURQUIN	82 SPEC	SPS hyperon beam
-0.29 ± 0.29	114	THOMPSON	80 ASPK	BNL hyperon beam
-0.17 ± 0.35	55	TANENBAUM	75B SPEC	BNL hyperon beam
+0.45 ± 0.20	186	^{9,10} FRANZINI	72 HBC	

⁹ The sign has been changed to agree with our convention.

Baryon Full Listings

$\Sigma^-, \Sigma(1385)$

¹⁰ The FRANZINI 72 value includes the events of earlier papers.



g_{VW}/g_A FOR $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}_e$

The values quoted assume the CVC prediction $g_V = 0$.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
2.4 ± 1.7 OUR AVERAGE				
1.75 ± 3.5	114	THOMPSON 80 ASPK	BNL hyperon beam	
3.5 ± 4.5	55	TANENBAUM 75B SPEC	BNL hyperon beam	
2.4 ± 2.1	186	FRANZINI 72 HBC		

Σ^- REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1986 edition (Physics Letters **170B**) or in earlier editions.

GUREV	93	JETPL 57 400	Gur'ev, Denisov, Zhelamkov, Ivanov+	(PNPI)
GALL	88	Translated from ZETFP 57 389	+Austin+	
HERTZOG	88	PR D37 1142	+Eckhause+	(BOST, MIT, WILL, CIT, CMU, WYOM)
HSUEH	88	PR D38 2056	+ (CHIC, ELMT, FNAL, IOWA, ISU, PNPI, YALE)	
ZAPALAC	86	PRL 57 1526	+ (EFI, ELMT, FNAL, IOWA, ISU, PNPI, YALE)	
HSUEH	85	PRL 54 2399	+Muller+	(CHIC, ELMT, FNAL, ISU, PNPI, YALE)
WAH	85	PRL 55 2551	+Cardello, Cooper, Teig+	(FNAL, IOWA, ISU)
BOURQUIN	83B	ZPHY C21 27	+ (BRIS, GEVA, HEIDP, LALO, RL, STRB)	
BOURQUIN	83C	ZPHY C21 17	+ (BRIS, GEVA, HEIDP, LALO, RL, STRB)	
DECK	83	PR D28 1	+Beretvas, Devlin, Luk+	(RUTG, WISC, MICH, MINN)
BOURQUIN	82	ZPHY C12 307	+Brown+	(BRIS, GEVA, HEIDP, LALO, RL, STRB)
MARRAFFINO	80	PR D21 2501	+Reucroft, Roos, Waters+	(VAND, MPIM)
THOMPSON	80	PR D21 25	+Cleland, Cooper, Dris, Engels+	(PITT, BNL)
HANSL	78	NP B132 45	+Manz, Matt, Reucroft, Settles+	(MPIM, VAND)
DECAMP	77	PL 66B 295	+Badier, Bland, Chollet, Gaillard+	(LALO, EPOL)
CONFORTO	76	NP B105 189	+Gopal, Kalmus, Litchfield, Ross+	(RHEL, LOIC)
DUGAN	75	NP A254 396	+Asano, Chen, Cheng, Hu, Lidofsky+	(COLU, YALE)
TANENBAUM	75B	PR D12 1871	+Hungerbuhler+	(YALE, FNAL, BNL)
ESENHOH	74	ZPHY 266 367	+Eisele, Engelmann, Filthuth, Hepp+	(HEID)
TANENBAUM	74	PRL 33 175	+Hungerbuhler+	(YALE, FNAL, BNL)
ESENHOH	73	ZPHY 264 413	+Eisele, Filthuth, Hepp, Leitner, Thouw+	(HEID)
SECHI-ZORN	73	PR D8 12	+Snow	(UMD)
BOHM	72	NP B48 1	+ (BERL, KIDR, BRUX, IASD, DUUC, LOUC+)	
FRANZINI	72	PR D6 2417	+ (COLU, HEID, UMD, STON)	
ROBERTSON	72	Thesis	+ (IIT)	
BAKKER	71	LNC 1 37	+Hoogland, Kluyver, Massard+	(SABRE Collab.)
COLE	71	PR D4 631	+Lee-Franzini, Loveless, Baltay+	(STON, COLU)
Also	69	Nevis 175 Thesis	Norton	(COLU)
TOVEE	71	NP B33 493	+ (LOUC, KIDR, BERL, BRUX, DUUC, WARS)	
BERLEY	70B	PR D1 2015	+Yamin, Hertzbach, Kofler+	(BNL, MASA, YALE)
BOGERT	70	PR D2 6	+Lucas, Taft, Willis, Berley+	(BNL, MASA, YALE)
EISELE	70	ZPHY 238 372	+Filthuth, Hepp, Presser, Zech	(HEID)
PDG	70	RMP 42 No. 1	Barbaro-Gallieri, Derenzo, Price+	(LRL, BRAN, CERN+)
ANG	69	ZPHY 223 103	+Eisele, Engelmann, Filthuth+	(HEID)
ANG	69B	ZPHY 228 151	+Ebenhoh, Eisele, Engelmann, Filthuth+	(HEID)
BAGGETT	69	PRL 23 249	+Kehoe, Snow	(UMD)
BALTAY	69	PRL 22 615	+Franzini, Newman, Norton+	(COLU, STON)
BANGERTER	69	UCRL 19244 Thesis	(LRL)	
BANGERTER	69B	PR 187 1821	+Alston-Garnjost, Gallieri, Gershwin+	(LRL)
BARLOUTAUD	69	NP B14 153	+DeBellefon, Granet+	(SACL, CERN, HEID)
EISELE	68	ZPHY 221 1	+Engelmann, Filthuth, Fohlisch, Hepp+	(HEID)
BIERMAN	68	PRL 20 1459	+Kounosu, Nauenberg+	(PRIN)
HEPP	68	ZPHY 214 71	+Schleich	(HEID)
WHITESIDE	68	NC 54A 537	+Gollub	(OBER)
BARASH	67	PRL 19 181	+Day, Glasser, Kehoe, Knop+	(UMD)
CHANG	66	PR 151 1081	+ (COLU)	
BAZIN	65B	PR 140B 1358	+Plano, Schmidt+	(PRIN, RUTG, COLU)
DOSCH	65	PL 14 339	+Engelmann, Filthuth, Hepp, Kluge+	(HEID)
Also	65	PR 151 1081	Chang	(COLU)
SCHMIDT	65	PR 140B 1328	+ (COLU)	
BURNSTEIN	64	PRL 13 66	+Day, Kehoe, Zorn, Snow	(UMD)
COURANT	64	PR 136B 1791	+Filthuth+	(CERN, HEID, UMD, NRL, BNL)
BARKAS	63	PRL 11 26	+Dyer, Heckman	(LRL)
HUMPHREY	62	PR 127 1305	+Ross	(LRL)

$\Sigma(1385) P_{13}$

$I(J^P) = 1(\frac{3}{2}^+)$ Status: * * * *


Discovered by ALSTON 60. Early measurements of the mass and width for combined charge states have been omitted. They may be found in our 1984 edition Reviews of Modern Physics **56** No. 2 Pt. II (1984).

We average only the most significant determinations. We do not average results from inclusive experiments with large backgrounds or results which are not accompanied by some discussion of experimental resolution. Nevertheless systematic differences between experiments remain. (See the ideograms in the Listings below.) These differences could arise from interference effects that change with production mechanism and/or beam momentum. They can also be accounted for in part by differences in the parametrizations employed. (See BORENSTEIN 74 for a discussion on this point.) Thus BORENSTEIN 74 uses a Breit-Wigner with energy-independent width, since a *P*-wave was found to give unsatisfactory fits. CAMERON 78 uses the same form. On the other hand HOLMGREN 77 obtains a good fit to their $\Lambda\pi$ spectrum with a *P*-wave Breit-Wigner, but includes the partial width for the $\Sigma\pi$ decay mode in the parametrization. AGUILAR-BENITEZ 81D gives masses and widths for five different Breit-Wigner shapes. The results vary considerably. Only the best-fit *S*-wave results are given here.

$\Sigma(1385)$ MASSES

$\Sigma(1385)^+$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1382.8 ± 0.4 OUR AVERAGE				Error includes scale factor of 2.0. See the Ideogram below.
1384.1 ± 0.7	1897	BAUBILLIER 84 HBC		$K^- p \rightarrow 8.25 \text{ GeV}/c$
1384.5 ± 0.5	5256	AGUILAR... 81D HBC		$K^- p \rightarrow \Lambda\pi\pi 4.2 \text{ GeV}/c$
1383.0 ± 0.4	9361	AGUILAR... 81D HBC		$K^- p \rightarrow \Lambda 3\pi 4.2 \text{ GeV}/c$
1381.9 ± 0.3	6900	CAMERON 78 HBC		$K^- p 0.96-1.36 \text{ GeV}/c$
1381 ± 1	6846	BORENSTEIN 74 HBC		$K^- p 2.18 \text{ GeV}/c$
1383.5 ± 0.85	2300	HABIBI 73 HBC		$K^- p \rightarrow \Lambda\pi\pi$
1382 ± 2	400	AGUILAR... 72B HBC		$K^- p \rightarrow \Lambda\pi's$
1384.4 ± 1.0	1260	SIEGEL 67 HBC		$K^- p 2.1 \text{ GeV}/c$
1382 ± 1	750	ARMENTEROS65B HBC		$K^- p 0.9-1.2 \text{ GeV}/c$
1381.0 ± 1.6	859	HUWE 64 HBC		$K^- p 1.22 \text{ GeV}/c$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1385.1 ± 1.2	600	BAKER 80 HYBR		$\pi^+ p 7 \text{ GeV}/c$
1383.2 ± 1.0	750	BAKER 80 HYBR		$K^- p 7 \text{ GeV}/c$
1381 ± 2	7k	1 BAUBILLIER 79B HBC		$K^- p 8.25 \text{ GeV}/c$
1391 ± 2	2k	CAUTIS 79 HYBR		$\pi^+ p/K^- p 11.5 \text{ GeV}$
1390 ± 2	100	1 SUGAHARA 79B HBC		$\pi^- p 6 \text{ GeV}/c$
1385 ± 3	22k	1.2 BARREIRO 77B HBC		$K^- p 4.2 \text{ GeV}/c$
1385 ± 1	2594	HOLMGREN 77 HBC		See AGUILAR 81D
1380 ± 2		1 BARDADIN... 75 HBC		$K^- p 14.3 \text{ GeV}/c$
1382 ± 1	3740	3 BERTHON 74 HBC		$K^- p 1263-1843 \text{ MeV}/c$
1390 ± 6	46	AGUILAR... 70B HBC		$K^- p \rightarrow \Sigma\pi's 4 \text{ GeV}/c$
1383 ± 8	62	4 BIRMINGHAM 66 HBC		$K^- p 3.5 \text{ GeV}/c$
1378 ± 5	135	LONDON 66 HBC		$K^- p 2.24 \text{ GeV}/c$
1384.3 ± 1.9	250	4 SMITH 65 HBC		$K^- p 1.8 \text{ GeV}/c$
1382.6 ± 2.1	250	4 SMITH 65 HBC		$K^- p 1.95 \text{ GeV}/c$
1375.0 ± 3.9	170	COOPER 64 HBC		$K^- p 1.45 \text{ GeV}/c$
1376.0 ± 3.9	154	4 ELY 61 HLBC		$K^- p 1.11 \text{ GeV}/c$



$\Sigma(1385)^+$ mass (MeV)


See key on page 1343

Baryon Full Listings

$\Sigma(1385)$

$\Sigma(1385)^0$ MASS


VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1383.7 ± 1.0 OUR AVERAGE		Error includes scale factor of 1.4. See the Ideogram below.		
1384.1 ± 0.8	5722	AGUILAR...	81D HBC	$K^- p \rightarrow \Lambda 3\pi$ 4.2 GeV/c
1380 ± 2	3100	⁵ BORENSTEIN 74	HBC	$K^- p \rightarrow \Lambda 3\pi$ 2.18 GeV/c
1385.1 ± 2.5	240	⁴ THOMAS 73	HBC	$\pi^- p \rightarrow \Lambda \pi^0 K^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1389 ± 3	500	⁶ BAUBILLIER 79B	HBC	$K^- p$ 8.25 GeV/c



$\Sigma(1385)^0$ mass (MeV)

$\Sigma(1385)^-$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1387.2 ± 0.5 OUR AVERAGE		Error includes scale factor of 2.2. See the Ideogram below.		
1388.3 ± 1.7	620	AGUILAR...	81D HBC	$K^- p \rightarrow \Lambda \pi \pi$ 4.2 GeV/c
1384.9 ± 0.8	3346	AGUILAR...	81D HBC	$K^- p \rightarrow \Lambda 3\pi$ 4.2 GeV/c
1387.6 ± 0.3	9720	CAMERON 78	HBC	$K^- p$ 0.96-1.36 GeV/c
1383 ± 2	2303	BORENSTEIN 74	HBC	$K^- p$ 2.18 GeV/c
1390.7 ± 1.2	1900	HABIBI 73	HBC	$K^- p \rightarrow \Lambda \pi \pi$
1387.1 ± 1.9	630	⁴ THOMAS 73	HBC	$\pi^- p \rightarrow \Lambda \pi^- K^+$
1390.7 ± 2.0	370	SIEGEL 67	HBC	$K^- p$ 2.1 GeV/c
1384 ± 1	1380	ARMENTEROS65B	HBC	$K^- p$ 0.9-1.2 GeV/c
1385.3 ± 1.9	1086	⁴ HUWE 64	HBC	$K^- p$ 1.15-1.30 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
1383 ± 1	4.5k	¹ BAUBILLIER 79B	HBC	$K^- p$ 8.25 GeV/c
1380 ± 6	150	¹ SUGAHARA 79B	HBC	$\pi^- p$ 6 GeV/c
1387 ± 3	12k	^{1,2} BARREIRO 77B	HBC	$K^- p$ 4.2 GeV/c
1391 ± 3	193	HOLMGREN 77	HBC	See AGUILAR 81D
1383 ± 2		¹ BARDADIN... 75	HBC	$K^- p$ 14.3 GeV/c
1389 ± 1	3060	³ BERTHON 74	HBC	$K^- p$ 1263-1843 MeV/c
1389 ± 9	15	LONDON 66	HBC	$K^- p$ 2.24 GeV/c
1391.5 ± 2.6	120	⁴ SMITH 65	HBC	$K^- p$ 1.8 GeV/c
1399.8 ± 2.2	58	⁴ SMITH 65	HBC	$K^- p$ 1.95 GeV/c
1392.0 ± 6.2	200	COOPER 64	HBC	$K^- p$ 1.45 GeV/c
1382 ± 3	93	DAHL 61	DBC	$K^- d$ 0.45 GeV/c
1376.0 ± 4.4	224	⁴ ELY 61	HLBC	$K^- p$ 1.11 GeV/c



$\Sigma(1385)^-$ mass (MeV)

$m_{\Sigma(1385)^-} - m_{\Sigma(1385)^+}$

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
- 2 to +6	95	⁷ BORENSTEIN 74	HBC	$K^- p$ 2.18 GeV/c
7.2 ± 1.4		⁷ HABIBI 73	HBC	$K^- p \rightarrow \Lambda \pi \pi$
6.3 ± 2.0		⁷ SIEGEL 67	HBC	$K^- p$ 2.1 GeV/c
11 ± 9		⁷ LONDON 66	HBC	$K^- p$ 2.24 GeV/c
9 ± 6		LONDON 66	HBC	$\Lambda 3\pi$ events
2.0 ± 1.5		⁷ ARMENTEROS65B	HBC	$K^- p$ 0.9-1.2 GeV/c
7.2 ± 2.1		⁷ SMITH 65	HBC	$K^- p$ 1.8 GeV/c
17.2 ± 2.0		⁷ SMITH 65	HBC	$K^- p$ 1.95 GeV/c
17 ± 7		⁷ COOPER 64	HBC	$K^- p$ 1.45 GeV/c
4.3 ± 2.2		⁷ HUWE 64	HBC	$K^- p$ 1.22 GeV/c
0.0 ± 4.2		⁷ ELY 61	HLBC	$K^- p$ 1.11 GeV/c

$m_{\Sigma(1385)^0} - m_{\Sigma(1385)^+}$

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
- 4 to + 4	95	⁷ BORENSTEIN 74	HBC	$K^- p$ 2.18 GeV/c

$m_{\Sigma(1385)^-} - m_{\Sigma(1385)^0}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
2.0 ± 2.4	⁷ THOMAS 73	HBC	$\pi^- p \rightarrow \Lambda \pi^- K^+$

$\Sigma(1385)$ WIDTHS

$\Sigma(1385)^+$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
35.8 ± 0.8 OUR AVERAGE				
37.2 ± 2.0	1897	BAUBILLIER 84	HBC	$K^- p$ 8.25 GeV/c
35.1 ± 1.7	5256	AGUILAR...	81D HBC	$K^- p \rightarrow \Lambda \pi \pi$ 4.2 GeV/c
37.5 ± 2.0	9361	AGUILAR...	81D HBC	$K^- p \rightarrow \Lambda 3\pi$ 4.2 GeV/c
35.5 ± 1.9	6900	CAMERON 78	HBC	$K^- p$ 0.96-1.36 GeV/c
34.0 ± 1.6	6846	⁸ BORENSTEIN 74	HBC	$K^- p$ 2.18 GeV/c
38.3 ± 3.2	2300	⁹ HABIBI 73	HBC	$K^- p \rightarrow \Lambda \pi \pi$
32.5 ± 6.0	400	AGUILAR...	72B HBC	$K^- p \rightarrow \Lambda \pi^+$
36 ± 4	1260	⁹ SIEGEL 67	HBC	$K^- p$ 2.1 GeV/c
32.0 ± 4.7	750	⁹ ARMENTEROS65B	HBC	$K^- p$ 0.95-1.20 GeV/c
46.5 ± 6.4	859	⁹ HUWE 64	HBC	$K^- p$ 1.15-1.30 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
40 ± 3	600	BAKER 80	HYBR	$\pi^+ p$ 7 GeV/c
37 ± 2	750	BAKER 80	HYBR	$K^- p$ 7 GeV/c
37 ± 2	7k	¹ BAUBILLIER 79B	HBC	$K^- p$ 8.25 GeV/c
30 ± 4	2k	CAUTIS 79	HYBR	$\pi^+ p / K^- p$ 11.5 GeV
30 ± 6	100	¹ SUGAHARA 79B	HBC	$\pi^- p$ 6 GeV/c
43 ± 5	22k	^{1,2} BARREIRO 77B	HBC	$K^- p$ 4.2 GeV/c
34 ± 2	2594	HOLMGREN 77	HBC	See AGUILAR 81D
40.0 ± 3.2		¹ BARDADIN... 75	HBC	$K^- p$ 14.3 GeV/c
48 ± 3	3740	³ BERTHON 74	HBC	$K^- p$ 1263-1843 MeV/c
33 ± 20	46	⁹ AGUILAR...	70B HBC	$K^- p \rightarrow \Sigma \pi^+ 4$ GeV/c
25 ± 32	62	⁹ BIRMINGHAM 66	HBC	$K^- p$ 3.5 GeV/c
30.3 ± 7.5	250	⁹ SMITH 65	HBC	$K^- p$ 1.8 GeV/c
33.1 ± 8.3	250	⁹ SMITH 65	HBC	$K^- p$ 1.95 GeV/c
51 ± 16	170	⁹ COOPER 64	HBC	$K^- p$ 1.45 GeV/c
48 ± 16	154	⁹ ELY 61	HLBC	$K^- p$ 1.11 GeV/c

$\Sigma(1385)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
36 ± 5 OUR AVERAGE				
34.8 ± 5.6	5722	AGUILAR...	81D HBC	$K^- p \rightarrow \Lambda 3\pi$ 4.2 GeV/c
39.3 ± 10.2	240	⁹ THOMAS 73	HBC	$\pi^- p \rightarrow \Lambda \pi^0 K^0$
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
53 ± 8	3100	¹⁰ BORENSTEIN 74	HBC	$K^- p \rightarrow \Lambda 3\pi$ 2.18 GeV/c
30 ± 9	106	CURTIS 63	OSPK	$\pi^- p$ 1.5 GeV/c


Baryon Full Listings

$\Sigma(1385)$

$\Sigma(1385)^-$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
39.4 ± 10.7	620	AGUILAR-...	81D HBC	$K^- p \rightarrow \Lambda \pi \pi$ 4.2 GeV/c
34.6 ± 4.2	3346	AGUILAR-...	81D HBC	$K^- p \rightarrow \Lambda \pi \pi$ 4.2 GeV/c
39.2 ± 1.7	9720	CAMERON	78 HBC	$K^- p$ 0.96-1.36 GeV/c
35 ± 3	2303	⁸ BORENSTEIN	74 HBC	$K^- p$ 2.18 GeV/c
51.9 ± 4.8	1900	⁹ HABIBI	73 HBC	$K^- p \rightarrow \Lambda \pi \pi$
48.2 ± 7.7	630	⁹ THOMAS	73 HBC	$\pi^- p \rightarrow \Lambda \pi^- K^0$
31.0 ± 6.5	370	⁹ SIEGEL	67 HBC	$K^- p$ 2.1 GeV/c
38.0 ± 4.1	1382	⁹ ARMENTEROS65B	HBC	$K^- p$ 0.95-1.20 GeV/c
62 ± 7	1086	HUWE	64 HBC	$K^- p$ 1.15-1.30 GeV/c
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
44 ± 4	4.5k	¹ BAUBILLIER	79B HBC	$K^- p$ 8.25 GeV/c
58 ± 4	150	¹ SUGAHARA	79B HBC	$\pi^- p$ 6 GeV/c
45 ± 5	12k	^{1,2} BARREIRO	77B HBC	$K^- p$ 4.2 GeV/c
35 ± 10	193	HOLMGREN	77 HBC	See AGUILAR 81D
47 ± 6		¹ BARDADIN-...	75 HBC	$K^- p$ 14.3 GeV/c
40 ± 3	3060	³ BERTHON	74 HBC	$K^- p$ 1263-1843 MeV/c
29.2 ± 10.6	120	⁹ SMITH	65 HBC	$K^- p$ 1.80 GeV/c
17.1 ± 8.9	58	⁹ SMITH	65 HBC	$K^- p$ 1.95 GeV/c
88 ± 24	200	⁹ COOPER	64 HBC	$K^- p$ 1.45 GeV/c
40		DAHL	61 DBC	$K^- d$ 0.45 GeV/c
66 ± 18	224	⁹ ELY	61 HLBC	$K^- p$ 1.11 GeV/c

WEIGHTED AVERAGE
39.4 ± 2.1 (Error scaled by 1.7)



$\Sigma(1385)^-$ width (MeV)

$\Sigma(1385)$ POLE POSITIONS

$\Sigma(1385)^+$ REAL PART

VALUE	DOCUMENT ID	COMMENT
1379 ± 1	LICHTENBERG74	Extrapolates HABIBI 73

$\Sigma(1385)^+$ -IMAGINARY PART

VALUE	DOCUMENT ID	COMMENT
17.5 ± 1.5	LICHTENBERG74	Extrapolates HABIBI 73

$\Sigma(1385)^-$ REAL PART

VALUE	DOCUMENT ID	COMMENT
1383 ± 1	LICHTENBERG74	Extrapolates HABIBI 73

$\Sigma(1385)^-$ -IMAGINARY PART

VALUE	DOCUMENT ID	COMMENT
22.5 ± 1.5	LICHTENBERG74	Extrapolates HABIBI 73

$\Sigma(1385)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda \pi$	88 ± 2 %
$\Gamma_2 \Sigma \pi$	12 ± 2 %
$\Gamma_3 \Lambda \gamma$	
$\Gamma_4 \Sigma \gamma$	
$\Gamma_5 N \bar{K}$	

The above branching fractions are our estimates, not fits or averages.

$\Sigma(1385)$ BRANCHING RATIOS

$\Gamma(\Sigma \pi)/\Gamma(\Lambda \pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
0.138 ± 0.011 OUR AVERAGE					
0.20 ± 0.06	DIONISI	78B HBC	±	$K^- p \rightarrow Y^* K \bar{K}$	
0.16 ± 0.03	BERTHON	74 HBC	+	$K^- p$ 1.26-1.84 GeV/c	
0.11 ± 0.02	BERTHON	74 HBC	-	$K^- p$ 1.26-1.84 GeV/c	
0.21 ± 0.05	BORENSTEIN	74 HBC	+	$K^- p \rightarrow \Lambda \pi^+ \pi^-$, $\Sigma^0 \pi^+ \pi^-$	
0.18 ± 0.04	MAST	73 MPWA	±	$K^- p \rightarrow \Lambda \pi^+ \pi^-$, $\Sigma^0 \pi^+ \pi^-$	
0.10 ± 0.05	THOMAS	73 HBC	-	$\pi^- p \rightarrow \Lambda K \pi$, $\Sigma K \pi$	
0.16 ± 0.07	AGUILAR-...	72B HBC	+	$K^- p$ 3.9, 4.6 GeV/c	
0.13 ± 0.04	COLLEY	71B DBC	-0	$K^- N$ 1.5 GeV/c	
0.13 ± 0.04	PAN	69 HBC	+	$\pi^+ p \rightarrow \Lambda K \pi$, $\Sigma K \pi$	
0.08 ± 0.06	LONDON	66 HBC	+	$K^- p$ 2.24 GeV/c	
0.163 ± 0.041	ARMENTEROS65B	HBC	±	$K^- p$ 0.95-1.20 GeV/c	
0.09 ± 0.04	HUWE	64 HBC	±	$K^- p$ 1.2-1.7 GeV	
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●					
<0.04	ALSTON	62 HBC	±0	$K^- p$ 1.15 GeV/c	
0.04 ± 0.04	BASTIEN	61 HBC	±		

$\Gamma(\Lambda \gamma)/\Gamma_{total}$	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
0.17 ± 0.17	1	MEISNER	72 HBC	1 event only	

$\Gamma(\Lambda \gamma)/\Gamma(\Lambda \pi)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
<0.06	90	COLAS	75 HLBC	$K^- p$ 575-970 MeV	

$\Gamma(\Sigma \gamma)/\Gamma(\Lambda \pi)$	CL%	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ_1
<0.05	90	COLAS	75 HLBC	$K^- p$ 575-970 MeV	

$(\Gamma_1 \Gamma_2)^{1/2}/\Gamma_{total} \ln N \bar{K} \rightarrow \Sigma(1385) \rightarrow \Lambda \pi$	DOCUMENT ID	CHG	COMMENT	$(\Gamma_5 \Gamma_1)^{1/2}/\Gamma$
+0.586 ± 0.319	¹¹ DEVENISH	74B 0	Fixed- t dispersion rel.	

$\Sigma(1385)$ FOOTNOTES

- From fit to inclusive $\Lambda \pi$ spectrum.
- Includes data of HOLMGREN 77.
- The errors are statistical only. The resolution is not unfolded.
- The error is enlarged to Γ/\sqrt{N} . See the note on the $K^*(892)$ mass in the 1984 edition.
- From a fit to $\Lambda \pi^0$ with the width fixed at 34 MeV.
- From fit to inclusive $\Lambda \pi^0$ spectrum with the width fixed at 40 MeV.
- Redundant with data in the mass Listings.
- Results from $\Lambda \pi^+ \pi^-$ and $\Lambda \pi^+ \pi^- \pi^0$ combined by us.
- The error is enlarged to $4\Gamma/\sqrt{N}$. See the note on the $K^*(892)$ mass in the 1984 edition.
- Consistent with +, 0, and - widths equal.
- An extrapolation of the parametrized amplitude below threshold.

$\Sigma(1385)$ REFERENCES

BAUBILLIER 84	ZPHY C23 213	+	Wohl, Cahn, Rittenberg+ (BIRM, CERN, GLAS, MSU, CURIN)
PDG 84	RMP 56 No. 2 Pt. II		(LBL, CIT, CERN)
AGUILAR-... 81D	AFIS A77 144		Aguiar-Benitez, Sallcio (MADR)
BAKER 80	NP B166 207	+	+Chima, Dornan, Gibbs, Hall, Miller+ (LOIC)
BAUBILLIER 79B	NP B148 18	+	(BIRM, CERN, GLAS, MSU, CURIN)
CAUTIS 79	NP B156 507	+	+Baliem, Bouchez, Carroli, Chadwick+ (SLAC)
SUGAHARA 79B	NP B156 237	+	+Ochiai, Fukui, Cooper+ (KEK, OSKC, KINK)
CAMERON 78	NP B143 189	+	+Frank, Gopal, Bacon, Buttenworth+ (RHEL, LOIC)
DIONISI 78B	PL 78B 154	+	+Armenteros, Diaz (CERN, AMST, NIJM, OXF)
BARREIRO 77B	NP B126 319	+	+Berge, Ganguli, Blokzijl+ (CERN, AMST, NIJM)
HOLMGREN 77	NP B119 261	+	+Aguiar-Benitez, Kluyver+ (CERN, AMST, NIJM)
BARDADIN-... 75	NP B98 418	+	Bardadin-Otwinowska+ (SACL, EPOL, RHEL)
COLAS 75	NP B91 253	+	+Farwell, Ferrer, Six (ORSAY)
BERTHON 74	NC 21A 146	+	+Trstrom+ (CDEF, RHEL, SACL, STRB)
BORENSTEIN 74	PR D9 3006	+	+Kalbfleisch, Strand+ (BNL, MICH)
DEVENISH 74B	NP B81 330	+	+Froggatt, Martin (DESY, NORD, LOUC)
LICHTENBERG 74	PR D10 3865		(IND)
Also 74B	Private Comm.		Lichtenberg (IND)
HABIBI 73	Nevis 199 Thesis		(COLU)
Also 73	Purdue Conf. 387		Baltay, Bridgewater, Cooper+ (COLU, BING)
MAST 73	PR D7 3212		+Sangerter, Alston-Garnjost+ (LBL) IJP
Also 73B	PR D7 5		Mast, Sangerter, Alston-Garnjost+ (LBL) IJP

See key on page 1343

Baryon Full Listings

 $\Sigma(1385)$, $\Sigma(1480)$ Bumps, $\Sigma(1560)$ Bumps

THOMAS	73	NP B56 15	+Engler, Fisk, Kraemer	(CMU) JP
AGUILAR...	72B	PR D6 29	Aguiar-Benitez, Chung, Eisner, Samios	(BNL)
MEISNER	72	NC 12A 62		(UNC, LBL)
COLLEY	71B	NP B31 61	+Cox, Eastwood, Fry+	(BIRM, EDIN, GLAS, LOIC)
AGUILAR...	70B	PRL 25 58	Aguiar-Benitez, Barnes, Bassano+	(BNL, SYRA)
PAN	69	PRL 23 808	+Forman	(PENN) I
SIEGEL	67	UCRL 18041 Thesis		(LRL)
BIRMINGHAM	66	PR 152 1148		(BIRM, GLAS, LOIC, OXF, RHEL)
LONDON	66	PR 143 1034	+Rau, Goldberg, Lichtman+	(BNL, SYRA) J
ARMENTEROS	65B	PL 19 75		(CERN, HEID, SACL)
SMITH	65	UCLA Thesis		(UCLA)
COOPER	64	PL 8 365	+Filthuth, Fridman, Malamud+	(CERN, AMST) J
HUWE	64	UCRL 11291 Thesis		(LRL)
Also	69	PR 180 1824	Huwe	(MICH) J
CURTIS	63	PR 132 1771	+Coffin, Meyer, Terwilliger	(LRL)
ALSTON	62	CERN Conf. 311	+Alvarez, Ferro-Luzzi+	(LRL)
BASTIEN	61	PRL 6 702	+Ferro-Luzzi, Rosenfeld	(LRL)
DAHL	61	PRL 6 142	+Horwitz, Miller, Murray, White	(LRL) J
ELY	61	PRL 7 461	+Fung, Gidal, Pan, Powell, White	(LRL) J
ALSTON	60	PRL 5 520	+Alvarez, Eberhard, Good, Graziano+	(LRL) I

 $\Sigma(1480)$ Bumps

$I(J^P) = 1(?)^?$ Status: *

OMITTED FROM SUMMARY TABLE

These are peaks seen in $\Lambda\pi$ and $\Sigma\pi$ spectra in the reaction $\pi^+ p \rightarrow (Y\pi)K^+$ at 1.7 GeV/c. Also, the Y polarization oscillates in the same region.

MILLER 70 suggests a possible alternate explanation in terms of a reflection of $N(1675) \rightarrow \Lambda K$ decay. However, such an explanation for the $(\Sigma^+\pi^0)K^+$ channel in terms of $\Delta(1650) \rightarrow \Sigma K$ decay seems unlikely (see PAN 70). In addition such reflections would also have to account for the oscillation of the Y polarization in the 1480 MeV region.

HANSON 71, with less data than PAN 70, can neither confirm nor deny the existence of this state. MAST 75 sees no structure in this region in $K^- p \rightarrow \Lambda\pi^0$.

ENGELEN 80 performs a multichannel analysis of $K^- p \rightarrow p\bar{K}^0\pi^-$ at 4.2 GeV/c. They observe a 3.5 standard-deviation signal at 1480 MeV in $p\bar{K}^0$ which cannot be explained as a reflection of any competing channel.

 $\Sigma(1480)$ MASS
(PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
≈ 1480 OUR ESTIMATE					
1480	120	ENGELEN 80	HBC	+	$K^- p \rightarrow (p\bar{K}^0)\pi^-$
1485 ± 10		CLINE 73	MPWA	-	$K^- d \rightarrow (A\pi^-)\rho$
1479 ± 10		PAN 70	HBC	+	$\pi^+ p \rightarrow (A\pi^+)K^+$
1465 ± 15		PAN 70	HBC	+	$\pi^+ p \rightarrow (\Sigma\pi)K^+$

 $\Sigma(1480)$ WIDTH
(PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
80 ± 20	120	ENGELEN 80	HBC	+	$K^- p \rightarrow (p\bar{K}^0)\pi^-$
40 ± 20		CLINE 73	MPWA	-	$K^- d \rightarrow (A\pi^-)\rho$
31 ± 15		PAN 70	HBC	+	$\pi^+ p \rightarrow (A\pi^+)K^+$
30 ± 20		PAN 70	HBC	+	$\pi^+ p \rightarrow (\Sigma\pi)K^+$

 $\Sigma(1480)$ DECAY MODES
(PRODUCTION EXPERIMENTS)

Mode	
Γ_1	$N\bar{K}$
Γ_2	$\Lambda\pi$
Γ_3	$\Sigma\pi$

 $\Sigma(1480)$ BRANCHING RATIOS
(PRODUCTION EXPERIMENTS)

VALUE	DOCUMENT ID	TECN	CHG	Γ_3/Γ_2
0.82 ± 0.51	PAN 70	HBC	+	

$\Gamma(N\bar{K})/\Gamma(\Lambda\pi)$	Γ_1/Γ_2		
VALUE	DOCUMENT ID	TECN	CHG
0.72 ± 0.50	PAN 70	HBC	+

$\Gamma(N\bar{K})/\Gamma_{total}$	Γ_1/Γ		
VALUE	DOCUMENT ID	TECN	COMMENT
small	CLINE 73	MPWA	$K^- d \rightarrow (A\pi^-)\rho$

 $\Sigma(1480)$ REFERENCES
(PRODUCTION EXPERIMENTS)

ENGELEN	80	NP B167 61	+Jongejans, Dionisi+	(NIJM, AMST, CERN, OXF)
MAST	75	PR D11 3078	+Alston-Garnjost, Bangarter+	(LBL)
CLINE	73	LNC 6 205	+Laumann, Mapp	(WISC) IJP
HANSON	71	PR D4 1296	+Kalmus, Louie	(LBL) I
MILLER	70	Duke Conf. 229		(PURD)
PAN	70	PR D2 49	+Forman, Ko, Hagopian, Selove	(PENN) I
Also	69	PRL 23 808	Pan, Forman	(PENN) I
Also	69B	PRL 23 806	Pan, Forman	(PENN) I

 $\Sigma(1560)$ Bumps

$I(J^P) = 1(?)^?$ Status: **

OMITTED FROM SUMMARY TABLE

This entry lists peaks reported in mass spectra around 1560 MeV without implying that they are necessarily real.

DIONISI 78B observes a 6 standard-deviation enhancement at 1553 MeV in the charged $\Lambda/\Sigma\pi$ mass spectra from $K^- p \rightarrow (\Lambda/\Sigma)\pi K\bar{K}$ at 4.2 GeV/c. In a CERN ISR experiment, LOCKMAN 78 reports a narrow 6 standard-deviation enhancement at 1572 MeV in $\Lambda\pi^\pm$ from the reaction $pp \rightarrow \Lambda\pi^+\pi^-X$. These enhancements are unlikely to be associated with the $\Sigma(1580)$ (which has not been confirmed by several recent experiments – see the next entry in the Listings).

CARROLL 76 observes a bump at 1550 MeV (as well as one at 1580 MeV) in the isospin-1 $\bar{N}N$ total cross section, but uncertainties in cross section measurements outside the mass range of the experiment preclude estimating its significance.

See also MEADOWS 80 for a review of this state.

 $\Sigma(1560)$ MASS
(PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
≈ 1560 OUR ESTIMATE					
1553 ± 7	121	DIONISI 78B	HBC	\pm	$K^- p \rightarrow (Y\pi)K\bar{K}$
1572 ± 4	40	LOCKMAN 78	SPEC	\pm	$pp \rightarrow \Lambda\pi^+\pi^-X$

 $\Sigma(1560)$ WIDTH
(PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
79 ± 30	121	DIONISI 78B	HBC	\pm	$K^- p \rightarrow (Y\pi)K\bar{K}$
15 ± 6	40	¹ LOCKMAN 78	SPEC	\pm	$pp \rightarrow \Lambda\pi^+\pi^-X$

 $\Sigma(1560)$ DECAY MODES
(PRODUCTION EXPERIMENTS)

Mode	Fraction (Γ_i/Γ)
Γ_1	$\Lambda\pi$
Γ_2	$\Sigma\pi$
	seen

 $\Sigma(1560)$ BRANCHING RATIOS
(PRODUCTION EXPERIMENTS)

$\Gamma(\Sigma\pi)/[\Gamma(\Lambda\pi) + \Gamma(\Sigma\pi)]$	$\Gamma_2/(\Gamma_1 + \Gamma_2)$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.35 ± 0.12	DIONISI 78B	HBC	\pm	$K^- p \rightarrow (Y\pi)K\bar{K}$

$\Gamma(\Lambda\pi)/\Gamma_{total}$	Γ_1/Γ			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
seen	LOCKMAN 78	SPEC	\pm	$pp \rightarrow \Lambda\pi^+\pi^-X$

 $\Sigma(1560)$ FOOTNOTES
(PRODUCTION EXPERIMENTS)

¹The width observed by LOCKMAN 78 is consistent with experimental resolution.

Baryon Full Listings

$\Sigma(1560)$ Bumps, $\Sigma(1580)$, $\Sigma(1620)$

$\Sigma(1560)$ REFERENCES (PRODUCTION EXPERIMENTS)

MEADOWS	80	Toronto Conf. 283		(CINC)
DIONISI	78B	PL 78B 154	+Armenteros, Diaz	(CERN, AMST, NIJM, OXF) I
LOCKMAN	78	CEN DPHPE 78-01	+Meyer, Rander, Poster, Schlein+	(UCLA, SACL)
CARROLL	76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL) I

$\Sigma(1580) D_{13}$

$$I(J^P) = 1(\frac{3}{2}^-) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

Seen in the isospin-1 $\bar{K}N$ cross section at BNL (LI 73, CARROLL 76) and in a partial-wave analysis of $K^- p \rightarrow \Lambda\pi^0$ for c.m. energies 1560-1600 MeV by LITCHFIELD 74. LITCHFIELD 74 finds $J^P = 3/2^-$. Not seen by ENGLER 78 or by CAMERON 78C (with larger statistics in $K_L^0 p \rightarrow \Lambda\pi^+$ and $\Sigma^0\pi^+$).

$\Sigma(1580)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1580 OUR ESTIMATE			
1583 \pm 4	1 CARROLL 76	DPWA	Isospin-1 total σ
1582 \pm 4	2 LITCHFIELD 74	DPWA	$K^- p \rightarrow \Lambda\pi^0$

$\Sigma(1580)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
15	1 CARROLL 76	DPWA	Isospin-1 total σ
11 \pm 4	2 LITCHFIELD 74	DPWA	$K^- p \rightarrow \Lambda\pi^0$

$\Sigma(1580)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$
Γ_2 $\Lambda\pi$
Γ_3 $\Sigma\pi$

$\Sigma(1580)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
+0.03 \pm 0.01	2 LITCHFIELD 74	DPWA	$\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
not seen	CAMERON 78C	HBC	$K_L^0 p \rightarrow \Lambda\pi^+$	
not seen	ENGLER 78	HBC	$K_L^0 p \rightarrow \Lambda\pi^+$	
+0.10 \pm 0.02	2 LITCHFIELD 74	DPWA	$K^- p \rightarrow \Lambda\pi^0$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1580) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
VALUE				
not seen	CAMERON 78C	HBC	$K_L^0 p \rightarrow \Sigma^0\pi^+$	
not seen	ENGLER 78	HBC	$K_L^0 p \rightarrow \Sigma^0\pi^+$	
+0.03 \pm 0.04	2 LITCHFIELD 74	DPWA	$\bar{K}N$ multichannel	

$\Sigma(1580)$ FOOTNOTES

- CARROLL 76 sees a total-cross-section bump with $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}} = 0.06$.
- The main effect observed by LITCHFIELD 74 is in the $\Lambda\pi$ final state; the $\bar{K}N$ and $\Sigma\pi$ couplings are estimated from a multichannel fit including total-cross-section data of LI 73.

$\Sigma(1580)$ REFERENCES

CAMERON	78C	NP B132 189	+Casiuippi+	(BGNA, EDIN, GLAS, PISA, RHEL) I
ENGLER	78	PR D18 3061	+Keyes, Kraemer, Tanaka, Cho+	(CMU, ANL)
CARROLL	76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL) I
LITCHFIELD	74	PL 51B 509		(CERN) IJP
LI	73	Purdue Conf. 283		(BNL) I

$\Sigma(1620) S_{11}$

$$I(J^P) = 1(\frac{1}{2}^-) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

The S_{11} state at 1697 MeV reported by VANHORN 75 is tentatively listed under the $\Sigma(1750)$. CARROLL 76 sees two bumps in the isospin-1 total cross section near this mass.

Production experiments are listed separately in the next entry.

$\Sigma(1620)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1620 OUR ESTIMATE			
1600 \pm 6	1 MORRIS 78	DPWA	$K^- n \rightarrow \Lambda\pi^-$
1608 \pm 5	2 CARROLL 76	DPWA	Isospin-1 total σ
1633 \pm 10	3 CARROLL 76	DPWA	Isospin-1 total σ
1630 \pm 10	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel
1620	KIM 71	DPWA	K-matrix analysis

$\Sigma(1620)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
87 \pm 19	1 MORRIS 78	DPWA	$K^- n \rightarrow \Lambda\pi^-$
15	2 CARROLL 76	DPWA	Isospin-1 total σ
10	3 CARROLL 76	DPWA	Isospin-1 total σ
65 \pm 20	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel
40	KIM 71	DPWA	K-matrix analysis

$\Sigma(1620)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$
Γ_2 $\Lambda\pi$
Γ_3 $\Sigma\pi$

$\Sigma(1620)$ BRANCHING RATIOS

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
0.22 \pm 0.02	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel	
0.05	KIM 71	DPWA	K-matrix analysis	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1620) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
0.12 \pm 0.02	1 MORRIS 78	DPWA	$K^- n \rightarrow \Lambda\pi^-$	
not seen	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$	
0.15	KIM 71	DPWA	K-matrix analysis	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1620) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
VALUE				
not seen	HEPP 76B	DPWA	$K^- n \rightarrow \Sigma\pi$	
0.40 \pm 0.06	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel	
0.08	KIM 71	DPWA	K-matrix analysis	

$\Sigma(1620)$ FOOTNOTES

- MORRIS 78 obtains an equally good fit without including this resonance.
- Total cross-section bump with $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}}$ is 0.06 seen by CARROLL 76.
- Total cross-section bump with $(J+1/2) \Gamma_{\text{el}} / \Gamma_{\text{total}}$ is 0.04 seen by CARROLL 76.

$\Sigma(1620)$ REFERENCES

MORRIS	78	PR D17 55	+Albright, Colleraine, Kimel, Lannutti	(FSU) IJP
CARROLL	76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL) I
HEPP	76B	PL 65B 487	+Braun, Grimm, Strobele+	(CERN, HEIDH, MPIM) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
LANGBEIN	72	NP B47 477	+Wagner	(MPIM) IJP
KIM	71	PRL 27 356		(HARV) IJP
Also	70	Duke Conf. 161	Kim	(HARV) IJP

See key on page 1343

Baryon Full Listings

$\Sigma(1620)$ Production Experiments, $\Sigma(1660)$

$\Sigma(1620)$ Production Experiments

$$I(J^P) = 1(?)^2$$

OMITTED FROM SUMMARY TABLE

Formation experiments are listed separately in the previous entry.

The results of CRENNELL 69B at 3.9 GeV/c are not confirmed by SABRE 70 at 3.0 GeV/c. However, at 4.5 GeV/c, AMMANN 70 sees a peak at 1642 MeV which on the basis of branching ratios they do not associate with the $\Sigma(1670)$. See MILLER 70 for a review of these conflicts.

**$\Sigma(1620)$ MASS
(PRODUCTION EXPERIMENTS)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1620 OUR ESTIMATE					
1642 ± 12		AMMANN 70	DBC		$K^- N$ 4.5 GeV/c
1618 ± 3	20	BLUMENFELD 69	HBC	+	$K_L^0 p$
1619 ± 8		CRENNELL 69B	DBC	±	$K^- N \rightarrow \Lambda \pi \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1616 ± 8		CRENNELL 68	DBC	±	See CRENNELL 69B

**$\Sigma(1620)$ WIDTH
(PRODUCTION EXPERIMENTS)**

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
55 ± 24		AMMANN 70	DBC		$K^- N$ 4.5 GeV/c
30 ± 10	20	BLUMENFELD 69	HBC	+	
72 +22 -15		CRENNELL 69B	DBC	±	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
66 ± 16		CRENNELL 68	DBC	±	See CRENNELL 69B

**$\Sigma(1620)$ DECAY MODES
(PRODUCTION EXPERIMENTS)**

Mode	
Γ_1	$N\bar{K}$
Γ_2	$\Lambda\pi$
Γ_3	$\Sigma\pi$
Γ_4	$\Lambda\pi\pi$
Γ_5	$\Sigma(1385)\pi$
Γ_6	$\Lambda(1405)\pi$

**$\Sigma(1620)$ BRANCHING RATIOS
(PRODUCTION EXPERIMENTS)**

$\Gamma(\Lambda\pi\pi)/\Gamma(\Lambda\pi)$		Γ_4/Γ_2			
VALUE	EVTS	DOCUMENT ID	TECN	CHG	
~ 2.5	14	BLUMENFELD 69	HBC	+	
$\Gamma(N\bar{K})/\Gamma(\Lambda\pi)$		Γ_1/Γ_2			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT	
0.4 ± 0.4	AMMANN 70	DBC		$K^- p$ 4.5 GeV/c	
0.0 ± 0.1	CRENNELL 68	DBC	+	See CRENNELL 69B	
$\Gamma(\Lambda\pi)/\Gamma_{total}$		Γ_2/Γ			
VALUE	DOCUMENT ID	TECN	CHG		
large	CRENNELL 68	DBC	±		
$\Gamma(\Sigma(1385)\pi)/\Gamma(\Lambda\pi)$		Γ_5/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
< 0.3	95	AMMANN 70	DBC		$K^- p$ 4.5 GeV/c
0.2 ± 0.1		CRENNELL 68	DBC	±	
$\Gamma(\Sigma\pi)/\Gamma(\Lambda\pi)$		Γ_3/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	COMMENT	
< 1.1	95	AMMANN 70	DBC	$K^- N$ 4.5 GeV/c	
$\Gamma(\Lambda(1405)\pi)/\Gamma(\Lambda\pi)$		Γ_6/Γ_2			
VALUE	DOCUMENT ID	TECN	COMMENT		
0.7 ± 0.4	AMMANN 70	DBC	$K^- p$ 4.5 GeV/c		

**$\Sigma(1620)$ REFERENCES
(PRODUCTION EXPERIMENTS)**

AMMANN 70	PRL 24 327	+Garfinkel, Carmony, Gutay+	(PURD, IND)
Also	73 PR D7 1345	Ammann, Carmony, Garfinkel+	(PURD, IUPU)
MILLER 70	Duke Conf. 229		(PURD)
SABRE 70	NP B16 201	Barloutaud, Merrill, Schever+	(SABRE Collab.)
BLUMENFELD 69	PL 29B 58	+Kalbfleisch	(BNL)
CRENNELL 69B	Lund Paper 183	+Karshon, Lai, O'Neil, Scarr+	(BNL, CUNY)
Results are quoted in LEVI-SETTI 69C.			
Also	69C Lund Conf.	Levi-Setti	(EFI)
CRENNELL 68	PRL 21 648	+Delaney, Flaminio, Karshon+	(BNL, CUNY)

$\Sigma(1660) P_{11}$

$$I(J^P) = 1(\frac{1}{2}^+)$$
 Status: * * *

For results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters 111B (1982).

$\Sigma(1660)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1630 to 1690 (1660) OUR ESTIMATE			
1665.1 ± 11.2	¹ KOISO 85	DPWA	$K^- p \rightarrow \Sigma\pi$
1670 ± 10	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
1679 ± 10	ALSTON... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
1676 ± 15	GOPAL 77	DPWA	$\bar{K} N$ multichannel
1668 ± 25	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
1670 ± 20	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1565 or 1597	² MARTIN 77	DPWA	$\bar{K} N$ multichannel
1660 ± 30	³ BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda\pi$
1671 ± 2	⁴ PONTE 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$

$\Sigma(1660)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40 to 200 (160) OUR ESTIMATE			
81.5 ± 22.2	¹ KOISO 85	DPWA	$K^- p \rightarrow \Sigma\pi$
152 ± 20	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
38 ± 10	ALSTON... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
120 ± 20	GOPAL 77	DPWA	$\bar{K} N$ multichannel
230 +165 -60	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
250 ± 110	KANE 74	DPWA	$K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
202 or 217	² MARTIN 77	DPWA	$\bar{K} N$ multichannel
80 ± 40	³ BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda\pi$
81 ± 10	⁴ PONTE 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$

$\Sigma(1660)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1	$N\bar{K}$ 10-30 %
Γ_2	$\Lambda\pi$ seen
Γ_3	$\Sigma\pi$ seen

$\Sigma(1660)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{total}$		Γ_1/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT
0.1 to 0.3 OUR ESTIMATE			
0.12 ± 0.03	GOPAL 80	DPWA	$\bar{K} N \rightarrow \bar{K} N$
0.10 ± 0.05	ALSTON... 78	DPWA	$\bar{K} N \rightarrow \bar{K} N$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 0.04	GOPAL 77	DPWA	See GOPAL 80
0.27 or 0.29	² MARTIN 77	DPWA	$\bar{K} N$ multichannel
$(\Gamma_1/\Gamma)^{1/2}/\Gamma_{total} \ln N\bar{K} \rightarrow \Sigma(1660) \rightarrow \Lambda\pi$		$(\Gamma_1/\Gamma)^{1/2}/\Gamma$	
VALUE	DOCUMENT ID	TECN	COMMENT
< 0.04	GOPAL 77	DPWA	$\bar{K} N$ multichannel
0.12 +0.12 -0.04	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
-0.10 or -0.11	² MARTIN 77	DPWA	$\bar{K} N$ multichannel
-0.04 ± 0.02	³ BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda\pi$
+0.16 ± 0.01	⁴ PONTE 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$

Baryon Full Listings

 $\Sigma(1660), \Sigma(1670)$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1660) \rightarrow \Sigma \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
-0.13 ± 0.04	¹ KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$	
-0.16 ± 0.03	GOPAL	77	DPWA $\bar{K} N$ multichannel	
-0.11 ± 0.01	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.34 or -0.37	² MARTIN	77	DPWA $\bar{K} N$ multichannel	
not seen	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$	

 $\Sigma(1660)$ FOOTNOTES

- ¹ The evidence of KOISO 85 is weak.
² The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
³ From solution 1 of BAILLON 75; not present in solution 2.
⁴ From solution 2 of PONTE 75; not present in solution 1.

 $\Sigma(1660)$ REFERENCES

KOISO	85	NP A433 619	+Sai, Yamamoto, Kofler	(TOKY, MASA)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON-...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC) IJP
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
HEPP	76B	PL 65B 487	+Braun, Grimm, Strobele+	(CERN, HEIDH, MPIIM) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
PONTE	75	PR D12 2597	+Hertzbach, Button-Shafer+	(MASA, TENN, UCR) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
KANE	74	LBL-2452		(LBL) IJP

NOTE ON THE $\Sigma(1670)$ REGION

Production experiments: The measured $\Sigma\pi/\Sigma\pi\pi$ branching ratio for the $\Sigma(1670)$ produced in the reaction $K^- p \rightarrow \pi^- \Sigma(1670)^+$ is strongly dependent on momentum transfer. This was first discovered by EBERHARD 69, who suggested that there exist two Σ resonances with the same mass and quantum numbers: one with a large $\Sigma\pi\pi$ (mainly $\Lambda(1405)\pi$) branching fraction produced peripherally, and the other with a large $\Sigma\pi$ branching fraction produced at larger angles. The experimental results have been confirmed by AGUILAR-BENITEZ 70, ASPELL 74, ESTES 74, and TIMMERMANS 76. If, in fact, there are two resonances, the most likely quantum numbers for both the $\Sigma\pi$ and the $\Lambda(1405)\pi$ states are D_{13} . There is also possibly a third Σ in this region, the $\Sigma(1690)$ in the Listings, the main evidence for which is a large $\Lambda\pi/\Sigma\pi$ branching ratio. These topics have been reviewed by EBERHARD 73 and by MILLER 70.

Formation experiments: Two states are also observed near this mass in formation experiments. One of these, the $\Sigma(1670)D_{13}$, has the same quantum numbers as those observed in production and has a large $\Sigma\pi/\Sigma\pi\pi$ branching ratio; it may well be the $\Sigma(1670)$ produced at larger angles (see TIMMERMANS 76). The other state, the $\Sigma(1660)P_{11}$, has different quantum numbers, its $\Sigma\pi/\Sigma\pi\pi$ branching ratio is unknown, and its relation to the produced $\Sigma(1670)$ states is obscure.

 $\Sigma(1670) D_{13}$

$$I(J^P) = 1(\frac{3}{2}^-) \text{ Status: } * * * *$$

For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

Results from production experiments are listed separately in the next entry.

 $\Sigma(1670)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1665 to 1685 (≈ 1670) OUR ESTIMATE			
1665.1 ± 4.1	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
1682 ± 5	GOPAL	80	DPWA $\bar{K} N \rightarrow \bar{K} N$
1679 ± 10	ALSTON-...	78	DPWA $\bar{K} N \rightarrow \bar{K} N$
1670 ± 5	GOPAL	77	DPWA $\bar{K} N$ multichannel
1670 ± 6	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
1685 ± 20	BAILLON	75	IPWA $\bar{K} N \rightarrow \Lambda \pi$
1659 ± 12	VANHORN	75	DPWA $K^- p \rightarrow \Lambda \pi^0$
1659 - 5			
1670 ± 2	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1667 or 1668	¹ MARTIN	77	DPWA $\bar{K} N$ multichannel
1650	DEBELLEFON	76	IPWA $K^- p \rightarrow \Lambda \pi^0$
1671 ± 3	PONTE	75	DPWA $K^- p \rightarrow \Lambda \pi^0$ (sol. 1)
1655 ± 2	PONTE	75	DPWA $K^- p \rightarrow \Lambda \pi^0$ (sol. 2)

 $\Sigma(1670)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
40 to 80 (≈ 60) OUR ESTIMATE			
65.0 ± 7.3	KOISO	85	DPWA $K^- p \rightarrow \Sigma \pi$
79 ± 10	GOPAL	80	DPWA $\bar{K} N \rightarrow \bar{K} N$
56 ± 20	ALSTON-...	78	DPWA $\bar{K} N \rightarrow \bar{K} N$
50 ± 5	GOPAL	77	DPWA $\bar{K} N$ multichannel
56 ± 3	HEPP	76B	DPWA $K^- N \rightarrow \Sigma \pi$
85 ± 25	BAILLON	75	IPWA $\bar{K} N \rightarrow \Lambda \pi$
32 ± 11	VANHORN	75	DPWA $K^- p \rightarrow \Lambda \pi^0$
79 ± 6	KANE	74	DPWA $K^- p \rightarrow \Sigma \pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
46 or 46	¹ MARTIN	77	DPWA $\bar{K} N$ multichannel
80	DEBELLEFON	76	IPWA $K^- p \rightarrow \Lambda \pi^0$
44 ± 11	PONTE	75	DPWA $K^- p \rightarrow \Lambda \pi^0$ (sol. 1)
76 ± 5	PONTE	75	DPWA $K^- p \rightarrow \Lambda \pi^0$ (sol. 2)

 $\Sigma(1670)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	7-13 %
Γ_2 $\Lambda\pi$	5-15 %
Γ_3 $\Sigma\pi$	30-60 %
Γ_4 $\Lambda\pi\pi$	
Γ_5 $\Sigma\pi\pi$	
Γ_6 $\Sigma(1385)\pi$	
Γ_7 $\Sigma(1385)\pi, S$ -wave	
Γ_8 $\Lambda(1405)\pi$	
Γ_9 $\Lambda(1520)\pi$	

The above branching fractions are our estimates, not fits or averages.

 $\Sigma(1670)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.07 to 0.13 OUR ESTIMATE				
0.10 ± 0.03	GOPAL	80	DPWA $\bar{K} N \rightarrow \bar{K} N$	
0.11 ± 0.03	ALSTON-...	78	DPWA $\bar{K} N \rightarrow \bar{K} N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.08 ± 0.03	GOPAL	77	DPWA See GOPAL 80	
0.07 or 0.07	¹ MARTIN	77	DPWA $\bar{K} N$ multichannel	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
0.17 ± 0.03	² MORRIS	78	DPWA $K^- n \rightarrow \Lambda \pi^-$	
0.13 ± 0.02	² MORRIS	78	DPWA $K^- n \rightarrow \Lambda \pi^-$	
+0.10 ± 0.02	GOPAL	77	DPWA $\bar{K} N$ multichannel	
+0.06 ± 0.02	BAILLON	75	IPWA $\bar{K} N \rightarrow \Lambda \pi$	
+0.09 ± 0.02	VANHORN	75	DPWA $K^- p \rightarrow \Lambda \pi^0$	
+0.018 ± 0.060	DEVENISH	74B	Fixed-t dispersion rel.	

See key on page 1343

Baryon Full Listings

$\Sigma(1670)$, $\Sigma(1670)$ Bumps

• • • We do not use the following data for averages, fits, limits, etc. • • •

+0.08 or +0.08	¹ MARTIN	77	DPWA	$\bar{K}N$ multichannel
+0.05	DEBELLEFON	76	IPWA	$K^-p \rightarrow \Lambda\pi^0$
0.08 \pm 0.01	PONTE	75	DPWA	$K^-p \rightarrow \Lambda\pi^0$ (sol. 1)
0.17 \pm 0.01	PONTE	75	DPWA	$K^-p \rightarrow \Lambda\pi^0$ (sol. 2)

 $(\Gamma_f/\Gamma_r)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.20 \pm 0.02	KOISO	85	DPWA $K^-p \rightarrow \Sigma\pi$
+0.21 \pm 0.02	GOPAL	77	DPWA $\bar{K}N$ multichannel
+0.20 \pm 0.01	HEPP	76B	DPWA $K^-N \rightarrow \Sigma\pi$
+0.21 \pm 0.03	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

+0.18 or +0.17 ¹ MARTIN 77 DPWA $\bar{K}N$ multichannel
 $\Gamma(\Lambda\pi\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.11	ARMENTEROS68E	HBC	K^-p ($\Gamma_1=0.09$)

 $(\Gamma_f/\Gamma_r)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Sigma(1385)\pi, S\text{-wave}$

VALUE	DOCUMENT ID	TECN	COMMENT
+0.11 \pm 0.03	PREVOST	74	DPWA $K^-N \rightarrow \Sigma(1385)\pi$
0.17 \pm 0.02	³ SIMS	68	DBC $K^-N \rightarrow \Lambda\pi\pi$

• • • We do not use the following data for averages, fits, limits, etc. • • •

 $\Gamma(\Sigma\pi\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.14	⁴ ARMENTEROS68E	HBC	K^-p, K^-d ($\Gamma_1=0.09$)

 $\Gamma(\Lambda(1405)\pi)/\Gamma_{\text{total}}$

VALUE	DOCUMENT ID	TECN	COMMENT
<0.06	ARMENTEROS68E	HBC	K^-p, K^-d ($\Gamma_1=0.09$)

 $\Gamma_f/\Gamma_r^2 \ln N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1405)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.007 \pm 0.002	⁵ BRÜCKER	70	DBC $K^-N \rightarrow \Sigma\pi\pi$
<0.03	BERLEY	69	HBC K^-p 0.6–0.82 GeV/c

 $\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma(1385)\pi)$

VALUE	DOCUMENT ID	TECN	COMMENT
0.23 \pm 0.08	BRÜCKER	70	DBC $K^-N \rightarrow \Sigma\pi\pi$

 $(\Gamma_f/\Gamma_r)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1670) \rightarrow \Lambda(1520)\pi$

VALUE	DOCUMENT ID	TECN	COMMENT
0.081 \pm 0.016	⁶ CAMERON	77	DPWA P -wave decay

$\Sigma(1670)$ FOOTNOTES

¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.² Results are with and without an S_{11} $\Sigma(1620)$ in the fit.³ SIMS 68 uses only cross-section data. Result used as upper limit only.⁴ Ratio only for $\Sigma 2\pi$ system in $l = 1$, which cannot be $\Sigma(1385)$.⁵ Assuming the $\Lambda(1405)\pi$ cross-section bump is due only to $3/2^-$ resonance.⁶ The CAMERON 77 upper limit on F -wave decay is 0.03.

$\Sigma(1670)$ REFERENCES

KOISO	85	NP A433 619	+Sai, Yamamoto, Kofler	(TOKY, MASA)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PR L 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
MORRIS	78	PR D17 95	+Albright, Collaraine, Kimmel, Lannutti	(FSU) IJP
CAMERON	77	NP B131 399	+Frank, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
HEPP	76B	PL 65B 487	+Braun, Grimm, Strobele+	(CERN, HEID, MPM) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
PONTE	75	PR D12 2597	+Hertzbach, Button-Shafer+	(MASA, TENN, UCR) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
DEVENISH	74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
KANE	74	LBL 2452		(LBL) IJP
PREVOST	74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID) IJP
BRÜCKER	70	Duke Conf. 155	+Harrison, Sims, Albright, Chandler+	(FSU) I
BERLEY	69	PL 30B 430	+Hart, Rahm, Willis, Yamamoto	(BNL)
ARMENTEROS 68E	68E	PL 28B 521	+Baillon+	(CERN, HEID, SACL) I
SIMS	68	PRL 21 1413	+Albright, Bartley, Meer+	(FSU, TUFTS, BRAN)

$\Sigma(1670)$ Bumps

$$I(J^P) = 1(?)^2$$

OMITTED FROM SUMMARY TABLE

Formation experiments are listed separately in the preceding entry.

Probably there are two states at the same mass with the same quantum numbers, one decaying to $\Sigma\pi$ and $\Lambda\pi$, the other to $\Lambda(1405)\pi$. See the note in front of the preceding entry.

$\Sigma(1670)$ MASS (PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
≈ 1670 OUR ESTIMATE					
1670 \pm 4		¹ CARROLL	76	DPWA	Isospin-1 total σ
1675 \pm 10		² HEPP	76	DBC	K^-N 1.6–1.75 GeV/c
1665 \pm 1		APSELL	74	HBC	K^-p 2.87 GeV/c
1688 \pm 2 or 1683 \pm 5	1200	BERTHON	74	HBC	0 Quasi-2-body σ
1670 \pm 6		AGUILAR...	70B	HBC	$K^-p \rightarrow \Sigma\pi\pi$ 4 GeV
1668 \pm 10		AGUILAR...	70B	HBC	$K^-p \rightarrow \Sigma 3\pi$ 4 GeV
1660 \pm 10		ALVAREZ	63	HBC	+ K^-p 1.51 GeV/c
1668 \pm 10	150	³ FERRERSORIA81	OMEG	-	π^-p 9.12 GeV/c
1655 to 1677		TIMMERMANS76	HBC	+	K^-p 4.2 GeV/c
1665 \pm 5		BUGG	68	CNTR	K^-p, d total σ
1661 \pm 9	70	PRIMER	68	HBC	+ See BARNES 69E
1685		ALEXANDER	62C	HBC	-0 π^-p 2–2.2 GeV/c

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Sigma(1670)$ WIDTH (PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
67.0 \pm 2.4		APSELL	74	HBC	K^-p 2.87 GeV/c
110 \pm 12		AGUILAR...	70B	HBC	$K^-p \rightarrow \Sigma\pi\pi$ 4 GeV
135 +40 -30		AGUILAR...	70B	HBC	$K^-p \rightarrow \Sigma 3\pi$ 4 GeV
40 \pm 10		ALVAREZ	63	HBC	+
90 \pm 20	150	³ FERRERSORIA81	OMEG	-	π^-p 9.12 GeV/c
52		¹ CARROLL	76	DPWA	Isospin-1 total σ
48 to 63		TIMMERMANS76	HBC	+	K^-p 4.2 GeV/c
30 \pm 15		BUGG	68	CNTR	
60 \pm 20	70	PRIMER	68	HBC	+ See BARNES 69E
45		ALEXANDER	62C	HBC	-0

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Sigma(1670)$ DECAY MODES (PRODUCTION EXPERIMENTS)

Mode	Γ_1/Γ_3
Γ_1 $N\bar{K}$	
Γ_2 $\Lambda\pi$	
Γ_3 $\Sigma\pi$	
Γ_4 $\Lambda\pi\pi$	
Γ_5 $\Sigma\pi\pi$	
Γ_6 $\Sigma(1385)\pi$	
Γ_7 $\Lambda(1405)\pi$	

$\Sigma(1670)$ BRANCHING RATIOS (PRODUCTION EXPERIMENTS)

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<0.03		TIMMERMANS76	HBC	+	K^-p 4.2 GeV/c
<0.10		BERTHON	74	HBC	0 Quasi-2-body σ
<0.2		AGUILAR...	70B	HBC	
<0.26		BARNES	69E	HBC	+ K^-p 3.9–5 GeV/c
0.025		BUGG	68	CNTR	0 Assuming $J = 3/2$
<0.24	0	PRIMER	68	HBC	+ K^-p 4.6–5 GeV/c
<0.6		LONDON	66	HBC	+ K^-p 2.25 GeV/c
<0.19	0	ALVAREZ	63	HBC	+ K^-p 1.15 GeV/c
≥ 0.5 ± 0.25		SMITH	63	HBC	-0

Baryon Full Listings

 $\Sigma(1670)$ Bumps, $\Sigma(1690)$ Bumps

$\Gamma(\Lambda\pi)/\Gamma(\Sigma\pi)$				Γ_2/Γ_3	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.76 ± 0.09		ESTES	74	HBC	0 $K^- p$ 2.1,2.6 GeV/c
0.45 ± 0.15		BARNES	69E	HBC	+ $K^- p$ 3.9-5 GeV/c
0.15 ± 0.07		HUWE	69	HBC	+
0.11 ± 0.06	33	BUTTON-...	68	HBC	+ $K^- p$ 1.7 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
$\leq 0.45 \pm 0.07$		TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c
0.55 ± 0.11		BERTHON	74	HBC	0 Quasi-2-body σ
0	0	PRIMER	68	HBC	+ See BARNES 69E
<0.6		LONDON	66	HBC	+ $K^- p$ 2.25 GeV/c
1.2	130	ALVAREZ	63	HBC	+ $K^- p$ 1.15 GeV/c
1.2		SMITH	63	HBC	-0
$\Gamma(\Lambda\pi\pi)/\Gamma(\Sigma\pi)$				Γ_4/Γ_3	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<0.6		LONDON	66	HBC	+ $K^- p$ 2.25 GeV/c
0.56	90	ALVAREZ	63	HBC	+ $K^- p$ 1.15 GeV/c
0.17		SMITH	63	HBC	-0
$\Gamma(\Sigma\pi\pi)/\Gamma(\Sigma\pi)$				Γ_5/Γ_3	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
largest at small angles		ESTES	74	HBC	0 $K^- p$ 2.1,2.6 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.2		² HEPP	76	DBC	- $K^- N$ 1.6-1.75 GeV/c
0.56	180	ALVAREZ	63	HBC	+ $K^- p$ 1.15 GeV/c
$\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma\pi)$				Γ_7/Γ_3	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1.8 ± 0.3 to 0.02 ± 0.07		^{3,4} TIMMERMANS76	HBC	+	$K^- p$ 4.2 GeV/c
largest at small angles		ESTES	74	HBC	\pm $K^- p$ 2.1,2.6 GeV/c
3.0 ± 1.6	50	LONDON	66	HBC	+ $K^- p$ 2.25 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.58 ± 0.20	17	PRIMER	68	HBC	+ See BARNES 69E
$\Gamma(\Sigma\pi)/\Gamma(\Sigma\pi\pi)$				Γ_3/Γ_5	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
varies with prod. angle		⁵ APSELL	74	HBC	+ $K^- p$ 2.87 GeV/c
1.39 ± 0.16		BERTHON	74	HBC	0 Quasi-2-body σ
2.5 to 0.24		⁴ EBERHARD	69	HBC	$K^- p$ 2.6 GeV/c
<0.4		BIRMINGHAM	66	HBC	+ $K^- p$ 3.5 GeV/c
0.30 ± 0.15		LONDON	66	HBC	+ $K^- p$ 2.25 GeV/c
$\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma\pi\pi)$				Γ_7/Γ_5	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.97 ± 0.08		TIMMERMANS76	HBC		$K^- p$ 4.2 GeV/c
1.00 ± 0.02		APSELL	74	HBC	$K^- p$ 2.87 GeV/c
$0.90^{+0.10}_{-0.16}$		EBERHARD	65	HBC	+ $K^- p$ 2.45 GeV/c
$\Gamma(\Lambda(1405)\pi)/\Gamma(\Sigma(1385)\pi)$				Γ_7/Γ_6	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<0.8		EBERHARD	65	HBC	+ $K^- p$ 2.45 GeV/c
$\Gamma(\Lambda\pi\pi)/\Gamma(\Sigma\pi\pi)$				Γ_4/Γ_5	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
0.35 ± 0.2		BIRMINGHAM	66	HBC	+ $K^- p$ 3.5 GeV/c
$\Gamma(\Lambda\pi)/\Gamma(\Sigma\pi\pi)$				Γ_2/Γ_5	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<0.2		BIRMINGHAM	66	HBC	+ $K^- p$ 3.5 GeV/c
$\Gamma(\Lambda\pi)/[\Gamma(\Lambda\pi) + \Gamma(\Sigma\pi)]$				$\Gamma_2/(\Gamma_2 + \Gamma_3)$	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
<0.6		AGUILAR-...	70B	HBC	
$\Gamma(\Sigma(1385)\pi)/\Gamma(\Sigma\pi)$				Γ_6/Γ_3	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$\leq 0.21 \pm 0.05$		TIMMERMANS76	HBC		$K^- p$ 4.2 GeV/c

 $\Sigma(1670)$ QUANTUM NUMBERS
(PRODUCTION EXPERIMENTS)

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$J^P = 3/2^-$	400	BUTTON-...	68	HBC	\pm $\Sigma^0\pi$
$J^P = 3/2^-$		EBERHARD	67	HBC	+ $\Lambda(1405)\pi$
$J^P = 3/2^+$		LEVEQUE	65	HBC	$\Lambda(1405)\pi$

 $\Sigma(1670)$ FOOTNOTES

- Total cross-section bump with $(J+1/2)\Gamma_{el}/\Gamma_{total} = 0.23$.
- Enhancements in $\Sigma\pi$ and $\Sigma\pi\pi$ cross sections.
- Backward production in the $\Lambda\pi^- K^+$ final state.
- Depending on production angle.
- APSELL 74, ESTES 74, and TIMMERMANS 76 find strong branching ratio dependence on production angle, as in earlier production experiments.

 $\Sigma(1670)$ REFERENCES
(PRODUCTION EXPERIMENTS)

FERRERSORIA	81	NP B178 373	+Treille, Rivet, Volte+	(CERN, CDEF, EPOL, LALO)
CARROLL	76	PRL 37 806	+Chiang, Kycia, Li, Mazur, Michael+	(BNL) I
HEPP	76	NP B115 82	+Braun, Grimm, Stroebela+	(CERN, HEID, MPIM) I
TIMMERMANS	76	NP B112 77	+Engelen+	(NIJM, CERN, AMST, OXF) J ^P
APSELL	74	PR D10 1419	+Ford, Gourevitch+	(BRAN, UMD, SYRA, TUFTS) I
BERTHON	74	NC 21A 146	+Tristram+	(CDEF, RHEL, SACL, STRB)
ESTES	74	LBL-3827 Thesis		(LBL)
AGUILAR-...	70B	PRL 25 58	Aguilar-Benitez, Barnes, Bassano+	(BNL, SYRA)
BARNES	69E	BNL 13823	+Chung, Eisner, Flaminio+	(BNL, SYRA)
EBERHARD	69	PRL 22 200	+Friedman, Pripstein, Ross	(LRL)
HUWE	69	PR 180 1824		(LRL)
BUGG	68	PR 168 1466		(LRL)
BUTTON-...	68	PRL 21 1123	+Gimlore, Knight+	(RHEL, BIRM, CAVE) I
PRIMER	68	PRL 20 610	Button-Shafer	(MASA, LRL) J ^P
EBERHARD	67	PR 163 1446	+Goldberg, Jaeger, Barnes, Dornan+	(SYRA, BNL)
BIRMINGHAM	66	PR 152 1148	+Pripstein, Shively, Kruse, Swanson	(LRL, ILL) J ^P
LONDON	66	PR 143 1034	(BIRM, GLAS, LOIC, OXF, RHEL)	(LRL, ILL) J ^P
EBERHARD	65	PRL 14 466	+Rau, Goldberg, Lichtman+	(BNL, SYRA) J ^P
LEVEQUE	65	PL 18 69	+Shively, Ross, Siegal, Ficenc+	(LRL, ILL) I
ALVAREZ	63	PRL 10 184	+ (SACL, EPOL, GLAS, LOIC, OXF, RHEL) J ^P	(LRL) I
SMITH	63	Athens Conf. 67	+Alston, Ferro-Luzzi, Huwe+	(LRL) I
ALEXANDER	62C	CERN Conf. 320		(LRL) I
			+Jacobs, Kalbfleisch, Miller+	(LRL) I

 $\Sigma(1690)$ Bumps

$I(J^P) = 1(?)^2$ Status: * *

OMITTED FROM SUMMARY TABLE

See the note preceding the $\Sigma(1670)$ Listings. Seen in production experiments only, mainly in $\Lambda\pi$. $\Sigma(1690)$ MASS
(PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
≈ 1690 OUR ESTIMATE					
1698 ± 20	70	¹ GODDARD	79	HBC	+ $\pi^+ p$ 10.3 GeV/c
1707 ± 20	40	² GODDARD	79	HBC	+ $\pi^+ p$ 10.3 GeV/c
1698 ± 20	15	ADERHOLZ	69	HBC	+ $\pi^+ p$ 8 GeV/c
1682 ± 2	46	BLUMENFELD	69	HBC	+ $K^0_L p$
1700 ± 20		MOTT	69	HBC	+ $K^- p$ 5.5 GeV/c
1694 ± 24	60	³ PRIMER	68	HBC	+ $K^- p$ 4.6-5 GeV/c
1700 ± 6		⁴ SIMS	68	HBC	- $K^- N \rightarrow \Lambda\pi\pi$
1715 ± 12	30	COLLEY	67	HBC	+ $K^- p$ 6 GeV/c

 $\Sigma(1690)$ WIDTH
(PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
240 ± 60	70	¹ GODDARD	79	HBC	+ $\pi^+ p$ 10.3 GeV/c
130^{+100}_{-60}	40	² GODDARD	79	HBC	+ $\pi^+ p$ 10.3 GeV/c
142 ± 40	15	ADERHOLZ	69	HBC	+ $\pi^+ p$ 8 GeV/c
25 ± 10	46	BLUMENFELD	69	HBC	+ $K^0_L p$
130 ± 25		MOTT	69	HBC	+ $K^- p$ 5.5 GeV/c
105 ± 35	60	³ PRIMER	68	HBC	+ $K^- p$ 4.6-5 GeV/c
62 ± 14		⁴ SIMS	68	HBC	- $K^- N \rightarrow \Lambda\pi\pi$
100 ± 35	30	COLLEY	67	HBC	+ $K^- p$ 6 GeV/c

 $\Sigma(1690)$ DECAY MODES
(PRODUCTION EXPERIMENTS)

Mode	Mode
Γ_1	$N\bar{K}$
Γ_2	$\Lambda\pi$
Γ_3	$\Sigma\pi$
Γ_4	$\Sigma(1385)\pi$
Γ_5	$\Lambda\pi\pi$ (including $\Sigma(1385)\pi$)

 $\Sigma(1690)$ BRANCHING RATIOS
(PRODUCTION EXPERIMENTS)

$\Gamma(N\bar{K})/\Gamma(\Lambda\pi)$				Γ_1/Γ_2	
VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
small		GODDARD	79	HBC	+ $\pi^+ p$ 10.2 GeV/c
<0.2		MOTT	69	HBC	+ $K^- p$ 5.5 GeV/c
0.4 ± 0.25	18	COLLEY	67	HBC	+ 6/30 events

See key on page 1343

Baryon Full Listings

$\Sigma(1690)$ Bumps, $\Sigma(1750)$

$\Gamma(\Sigma\pi)/\Gamma(\Lambda\pi)$		Γ_3/Γ_2			
VALUE	CL%	DOCUMENT ID	TECN	CHG	COMMENT
small		GODDARD 79	HBC	+	$\pi^+ p$ 10.2 GeV/c
<0.4	90	MOTT 69	HBC	+	$K^- p$ 5.5 GeV/c
0.3±0.3		COLLEY 67	HBC	+	4/30 events

$\Gamma(\Sigma(1385)\pi)/\Gamma(\Lambda\pi)$		Γ_4/Γ_2			
VALUE		DOCUMENT ID	TECN	CHG	COMMENT
<0.5		MOTT 69	HBC	+	$K^- p$ 5.5 GeV/c

$\Gamma(\Lambda\pi\pi(\text{including } \Sigma(1385)\pi))/\Gamma(\Lambda\pi)$		Γ_5/Γ_2			
VALUE		DOCUMENT ID	TECN	CHG	COMMENT
2.0±0.6		BLUMENFELD 69	HBC	+	31/15 events
0.5±0.25		COLLEY 67	HBC	+	15/30 events

$\Gamma(\Sigma(1385)\pi)/\Gamma(\Lambda\pi\pi(\text{including } \Sigma(1385)\pi))$		Γ_4/Γ_5			
VALUE		DOCUMENT ID	TECN	CHG	COMMENT
large		SIMS 68	HBC	-	$K^- N \rightarrow \Lambda\pi\pi$
small		COLLEY 67	HBC	+	$K^- p$ 6 GeV/c

$\Sigma(1690)$ FOOTNOTES (PRODUCTION EXPERIMENTS)

- ¹ From $\pi^+ p \rightarrow (\Lambda\pi^+)K^+$. $J > 1/2$ is not required by the data.
² From $\pi^+ p \rightarrow (\Lambda\pi^+)(K\pi)^+$. $J > 1/2$ is indicated, but large background precludes a definite conclusion.
³ See the $\Sigma(1670)$ Listings. AGUILAR-BENITEZ 70b with three times the data of PRIMER 68 find no evidence for the $\Sigma(1690)$.
⁴ This analysis, which is difficult and requires several assumptions and shows no unambiguous $\Sigma(1690)$ signal, suggests $J^P = 5/2^+$. Such a state would lead all previously known Y^* trajectories.

$\Sigma(1690)$ REFERENCES (PRODUCTION EXPERIMENTS)

GODDARD 79	PR D19 1350	+Key, Luste, Prentice, Yoon, Gordon+	(TNTD, BNL)J
AGUILAR... 70b	PRL 25 58	Aguiar-Benitez, Barnes, Bassano+	(BNL, SYRA)
ADERHOLZ 69	NP B11 259	+Bartsch+	(AACH3, BERL, CERN, JAGL, WARS)I
BLUMENFELD 69	PL 298 58	+Kalbfleisch	(BNL)I
MOTT 69	PR 177 1966	+Ammar, Davis, Kropac, Slate+	(NWES, ANL)I
Also 67	PRL 18 266	+Derrick, Fields, Loken, Ammar+	(ANL, NWES)I
PRIMER 68	PRL 20 610	+Goldberg, Jaeger, Barnes, Dorman+	(SYRA, BNL)I
SIMS 68	PRL 21 1413	+Albright, Bartley, Meer+	(FSU, TUFTS, BRAN)I
COLLEY 67	PL 248 489		(BIRM, GLAS, LOIC, MUNI, OXF, RHEL)I

$\Sigma(1750) S_{11}$

$$I(J^P) = 1(\frac{1}{2}^-) \text{ Status: } ***$$

For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

There is evidence for this state in many partial-wave analyses, but with wide variations in the mass, width, and couplings. The latest analyses indicated significant couplings to $N\bar{K}$ and $\Lambda\pi$, as well as to $\Sigma\eta$ whose threshold is at 1746 MeV (JONES 74).

$\Sigma(1750)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1730 to 1800 (≈ 1750) OUR ESTIMATE			
1756±10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1770±10	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1770±15	GOPAL 77	DPWA	$\bar{K}N$ multichannel
••• We do not use the following data for averages, fits, limits, etc. •••			
1800 or 1813	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
1715±10	² CARROLL 76	DPWA	Isospin-1 total σ
1730	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$
1780±30	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$ (sol. 1)
1700±30	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$ (sol. 2)
1697±20	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
1785±12	CHU 74	DBC	Fits $\sigma(K^- n \rightarrow \Sigma^- \eta)$
1760±5	³ JONES 74	HBC	Fits $\sigma(K^- p \rightarrow \Sigma^0 \eta)$
1739±10	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$

$\Sigma(1750)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
60 to 160 (≈ 90) OUR ESTIMATE			
64±10	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
161±20	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
60±10	GOPAL 77	DPWA	$\bar{K}N$ multichannel

••• We do not use the following data for averages, fits, limits, etc. •••

117 or 119	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
10	² CARROLL 76	DPWA	Isospin-1 total σ
110	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$
140±30	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$ (sol. 1)
160±50	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$ (sol. 2)
66 ⁺¹⁴ ₋₁₂	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$
89±33	CHU 74	DBC	Fits $\sigma(K^- n \rightarrow \Sigma^- \eta)$
92±7	³ JONES 74	HBC	Fits $\sigma(K^- p \rightarrow \Sigma^0 \eta)$
108±20	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$

$\Sigma(1750)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	10–40 %
Γ_2 $\Lambda\pi$	seen
Γ_3 $\Sigma\pi$	<8 %
Γ_4 $\Sigma\eta$	15–55 %
Γ_5 $\Sigma(1385)\pi$	
Γ_6 $\Lambda(1520)\pi$	

The above branching fractions are our estimates, not fits or averages.

$\Sigma(1750)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.1 to 0.4 OUR ESTIMATE				
0.14±0.03	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.33±0.05	ALSTON-... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$	

••• We do not use the following data for averages, fits, limits, etc. •••

0.15±0.03	GOPAL 77	DPWA	See GOPAL 80
0.06 or 0.05	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1750) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.04 ± 0.03				
••• We do not use the following data for averages, fits, limits, etc. •••				
-0.10 or -0.09	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	
-0.12	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda\pi^0$	
-0.12 ± 0.02	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$ (sol. 1)	
-0.13 ± 0.03	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$ (sol. 2)	
-0.13 ± 0.04	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0$	
-0.120±0.077	DEVENISH 74b		Fixed- t dispersion rel.	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1750) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
0.23 ± 0.01				
••• We do not use the following data for averages, fits, limits, etc. •••				
+0.06 or +0.06	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	
0.13±0.02	LANGBEIN 72	IPWA	$\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_4)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1750) \rightarrow \Sigma\eta$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
0.23 ± 0.01				
••• We do not use the following data for averages, fits, limits, etc. •••				
seen	CLINE 69	DBC	Threshold bump	

$(\Gamma_1\Gamma_5)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1750) \rightarrow \Sigma(1385)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
+0.18 ± 0.15				
••• We do not use the following data for averages, fits, limits, etc. •••				
+0.18±0.15	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385)\pi$	

$(\Gamma_1\Gamma_6)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1750) \rightarrow \Lambda(1520)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_6)^{1/2}/\Gamma$
0.032 ± 0.021				
••• We do not use the following data for averages, fits, limits, etc. •••				
0.032±0.021	CAMERON 77	DPWA	P -wave decay	

$\Sigma(1750)$ FOOTNOTES

- ¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
² A total cross-section bump with $(J+1/2) \Gamma_{ej} / \Gamma_{\text{total}} = 0.30$.
³ An S-wave Breit-Wigner fit to the threshold cross section with no background and errors statistical only.

Baryon Full Listings

$\Sigma(1750), \Sigma(1770), \Sigma(1775)$

$\Sigma(1750)$ REFERENCES

PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON	77	NP B131 399	+Franeek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC)
CARROLL	76	PRL 37 806	+Chiang, Kyica, Li, Mazur, Michael+	(BNL) I
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
CHU	74	NC 20A 35	+Bartley+	(PLAT, TUFTS, BRAN) IJP
DEVENISH	74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
JONES	74	NP B73 141		(CHIC) IJP
PREVOST	74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID)
LANGBEIN	72	NP B47 477	+Wagner	(MPIM) IJP
CLINE	69	LCN 2 407	+Laumann, Mapp	(WISC)

$\Sigma(1770)$ P_{11}

$$I(J^P) = 1(\frac{1}{2}^+) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

Evidence for this state now rests solely on solution 1 of BAILLON 75, (see the footnotes) but the $\Lambda\pi$ partial-wave amplitudes of this solution are in disagreement with amplitudes from most other $\Lambda\pi$ analyses.

$\Sigma(1770)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1770 OUR ESTIMATE			
1738 \pm 10	¹ GOPAL	77	DPWA $\bar{K}N$ multichannel
1770 \pm 20	² BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
1772	³ KANE	72	DPWA $K^-p \rightarrow \Sigma\pi$

$\Sigma(1770)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
72 \pm 10	¹ GOPAL	77	DPWA $\bar{K}N$ multichannel
80 \pm 30	² BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
80	³ KANE	72	DPWA $K^-p \rightarrow \Sigma\pi$

$\Sigma(1770)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	37-43%
Γ_2 $\Lambda\pi$	14-20%
Γ_3 $\Sigma\pi$	2-5%

$\Sigma(1770)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.14 \pm 0.04	¹ GOPAL	77	DPWA $\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
< 0.04	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.08 \pm 0.02	² BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$	

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1770) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_3)^{1/2}/\Gamma$
< 0.04	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.108	³ KANE	72	DPWA $K^-p \rightarrow \Sigma\pi$	

$\Sigma(1770)$ FOOTNOTES

- ¹ Required to fit the isospin-1 total cross section of CARROLL 76 in the $\bar{K}N$ channel. The addition of new K^-p polarization and K^-n differential cross-section data in GOPAL 80 find it to be more consistent with the $\Sigma(1660) P_{11}$.
- ² From solution 1 of BAILLON 75; not present in solution 2.
- ³ Not required in KANE 74, which supersedes KANE 72.

$\Sigma(1770)$ REFERENCES

GOPAL	80	Toronto Conf. 159		(RHEL)
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
CARROLL	76	PRL 37 806	+Chiang, Kyica, Li, Mazur, Michael+	(BNL) I
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
KANE	74	LBL-2452		(LBL) IJP
KANE	72	PR D5 1583		(LBL)

$\Sigma(1775)$ D_{15}

$$I(J^P) = 1(\frac{5}{2}^-) \text{ Status: } ***$$

Discovered by GALTIERI 63, this resonance plays the same role as cornerstone for isospin-1 analyses in this region as the $\Lambda(1820)$ does in the isospin-0 channel.

For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters 111B (1982).

$\Sigma(1775)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1770 to 1780 (≈ 1778) OUR ESTIMATE			
1778 \pm 5	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1777 \pm 5	ALSTON...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
1774 \pm 5	GOPAL	77	DPWA $\bar{K}N$ multichannel
1775 \pm 10	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
1774 \pm 10	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$
1772 \pm 6	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1772 or 1777	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
1765	DEBELLEFON	76	IPWA $K^-p \rightarrow \Lambda\pi^0$

$\Sigma(1775)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
108 to 138 (≈ 120) OUR ESTIMATE			
137 \pm 10	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
116 \pm 10	ALSTON...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$
130 \pm 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
125 \pm 15	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
146 \pm 18	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$
154 \pm 10	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
102 or 103	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
120	DEBELLEFON	76	IPWA $K^-p \rightarrow \Lambda\pi^0$

$\Sigma(1775)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	37-43%
Γ_2 $\Lambda\pi$	14-20%
Γ_3 $\Sigma\pi$	2-5%
Γ_4 $\Sigma(1385)\pi$	8-12%
Γ_5 $\Sigma(1385)\pi, D\text{-wave}$	
Γ_6 $\Lambda(1520)\pi$	17-23%
Γ_7 $\Sigma\pi$	

The above branching fractions are our estimates, not fits or averages.

CONSTRAINED FIT INFORMATION

An overall fit to 8 branching ratios uses 16 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 63.9$ for 12 degrees of freedom.

The following off-diagonal array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-30			
x_3	-17	-21		
x_4	-37	-49	-14	
x_6	-81	6	8	16
	x_1	x_2	x_3	x_4

$\Sigma(1775)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances. Also, the errors quoted do not include uncertainties due to the parametrization used in the partial-wave analyses and are thus too small.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.37 to 0.43 OUR ESTIMATE				
0.45 \pm 0.04 OUR FIT	Error includes scale factor of 3.1.			
0.391 \pm 0.017 OUR AVERAGE				
0.40 \pm 0.02	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.37 \pm 0.03	ALSTON...	78	DPWA $\bar{K}N \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.41 \pm 0.03	GOPAL	77	DPWA See GOPAL 80	
0.37 or 0.36	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	

See key on page 1343

Baryon Full Listings
 $\Sigma(1775), \Sigma(1840)$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
VALUE				
0.305 ± 0.018 OUR FIT			Error includes scale factor of 2.4.	
-0.262 ± 0.018 OUR AVERAGE				
-0.28 ± 0.03	GOPAL 77	DPWA	$\bar{K} N$ multichannel	
-0.25 ± 0.02	BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda \pi$	
-0.28 ± 0.04	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda \pi^0$	
-0.05				
-0.259 ± 0.048	DEVENISH 74B		Fixed-t dispersion rel.	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.29 or -0.28	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	
-0.30	DEBELLEFON 76	IPWA	$K^- p \rightarrow \Lambda \pi^0$	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
VALUE				
0.106 ± 0.025 OUR FIT			Error includes scale factor of 3.1.	
0.096 ± 0.016 OUR AVERAGE			Error includes scale factor of 1.8.	
$+0.13 \pm 0.02$	GOPAL 77	DPWA	$\bar{K} N$ multichannel	
0.09 ± 0.01	KANE 74	DPWA	$K^- p \rightarrow \Sigma \pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$+0.08$ or $+0.08$	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1775) \rightarrow \Lambda(1520) \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
VALUE				
0.315 ± 0.010 OUR FIT			Error includes scale factor of 1.5.	
0.303 ± 0.009 OUR AVERAGE			Signs on measurements were ignored.	
-0.305 ± 0.010	² CAMERON 77	DPWA	$K^- p \rightarrow \Lambda(1520) \pi^0$	
0.31 ± 0.02	BARLETTA 72	DPWA	$K^- p \rightarrow \Lambda(1520) \pi^0$	
0.27 ± 0.03	ARMENTEROS65C	HBC	$K^- p \rightarrow \Lambda(1520) \pi^0$	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1775) \rightarrow \Sigma(1385) \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
VALUE				
0.211 ± 0.022 OUR FIT			Error includes scale factor of 2.8.	
0.188 ± 0.010 OUR AVERAGE			Signs on measurements were ignored.	
-0.184 ± 0.011	³ CAMERON 78	DPWA	$K^- p \rightarrow \Sigma(1385) \pi$	
$+0.20 \pm 0.02$	PREVOST 74	DPWA	$K^- N \rightarrow \Sigma(1385) \pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.32 ± 0.06	SIMS 68	DBC	$K^- N \rightarrow \Lambda \pi \pi$	
0.24 ± 0.03	ARMENTEROS67C	HBC	$K^- p \rightarrow \Lambda \pi \pi$	

$\Gamma(\Lambda \pi) / \Gamma(N \bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_2 / Γ_1
VALUE				
0.46 ± 0.09 OUR FIT			Error includes scale factor of 2.9.	
0.33 ± 0.05	UHLIG 67	HBC	$K^- p$ 0.9 GeV/c	

$\Gamma(\Sigma \pi \pi) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_7 / Γ
VALUE				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.12	⁴ ARMENTEROS68C	HDBC	$K^- N \rightarrow \Sigma \pi \pi$	

$\Gamma(\Sigma(1385) \pi) / \Gamma(N \bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_4 / Γ_1
VALUE				
0.22 ± 0.07 OUR FIT			Error includes scale factor of 3.6.	
0.25 ± 0.09	UHLIG 67	HBC	$K^- p$ 0.9 GeV/c	

$\Gamma(\Lambda(1520) \pi) / \Gamma(N \bar{K})$	DOCUMENT ID	TECN	COMMENT	Γ_6 / Γ_1
VALUE				
0.49 ± 0.11 OUR FIT			Error includes scale factor of 3.5.	
0.28 ± 0.05	UHLIG 67	HBC	$K^- p$ 0.9 GeV/c	

 $\Sigma(1775)$ FOOTNOTES

- The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- This rate combines P-wave and F-wave decays. The CAMERON 77 results for the separate P-wave- and F-wave decays are -0.303 ± 0.010 and -0.037 ± 0.014 . The published signs have been changed here to be in accord with the baryon-first convention.
- The CAMERON 78 upper limit on G-wave decay is 0.03.
- For about 3/4 of this, the $\Sigma \pi$ system has $l = 0$ and is almost entirely $\Lambda(1520)$. For the rest, the $\Sigma \pi$ has $l = 1$, which is about what is expected from the known $\Sigma(1775) \rightarrow \Sigma(1385) \pi$ rate, as seen in $\Lambda \pi \pi$.

 $\Sigma(1775)$ REFERENCES

PDG 82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL 80	Toronto Conf. 159		(RHEL) IJP
ALSTON.... 78	PR D18 182	Aiston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also 77	PRL 38 1007	Aiston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON 78	NP B143 189	+Franeek, Gopal, Bacon, Buttenworth+	(RHEL, LOIC) IJP
CAMERON 77	NP B131 399	+Franeek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
GOPAL 77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN 77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also 77B	NP B126 266	Martin, Pidcock	(LOUC)
Also 77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
DEBELLEFON 76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
BAILLON 75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
VANHORN 75	NP B87 145		(LBL) IJP
Also 75B	NP B87 157	VanHorn	(LBL) IJP

DEVENISH 74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
KANE 74	LBL-2452		(LBL) IJP
PREVOST 74	NP B69 246	+Barloutaud+	(SACL, CERN, HEID)
BARLETTA 72	NP B40 45		(EFI) IJP
Also 66	PRL 17 841	Fenster, Gelfand, Harmsen+	(CHIC, ANL, CERN) IJP
ARMENTEROS 68C	NP B8 216	+Baillon+	(CERN, HEID, SACL) I
SIMS 68	PRL 21 1413	+Albright, Bartley, Meer+	(FSU, TUFTS, BRAN)
ARMENTEROS 67C	ZPHY 202 486	+Ferro-Luzzi+	(CERN, HEID, SACL)
UHLIG 67	PR 155 1448	+Charlton, Condon, Glasser, Yodh+	(UMD, NRL)
ARMENTEROS 65C	PL 19 338	+Ferro-Luzzi+	(CERN, HEID, SACL) IJP
GALTIERI 63	PL 6 296	+Hussain, Tripp	(LRL) IJ

 $\Sigma(1840) P_{13}$

$I(J^P) = 1(\frac{3}{2}^+)$ Status: *

OMITTED FROM SUMMARY TABLE

For the time being, we list together here all resonance claims in the P_{13} wave between 1700 and 1900 MeV. $\Sigma(1840)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 1840 OUR ESTIMATE			
1798 or 1802	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel
1720 \pm 30	² BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda \pi$
1925 \pm 200	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda \pi^0$
1840 \pm 10	LANGBEIN 72	IPWA	$\bar{K} N$ multichannel

 $\Sigma(1840)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
93 or 93	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel
120 \pm 30	² BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda \pi$
65 $^{+50}_{-20}$	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda \pi^0$
120 \pm 10	LANGBEIN 72	IPWA	$\bar{K} N$ multichannel

 $\Sigma(1840)$ DECAY MODES

Mode
Γ_1 $N \bar{K}$
Γ_2 $\Lambda \pi$
Γ_3 $\Sigma \pi$

 $\Sigma(1840)$ BRANCHING RATIOSSee "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N \bar{K}) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
VALUE				
0 or 0	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	
0.37 \pm 0.13	LANGBEIN 72	IPWA	$\bar{K} N$ multichannel	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1840) \rightarrow \Lambda \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
VALUE				
$+0.03$ or $+0.03$	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	
$+0.11 \pm 0.02$	² BAILLON 75	IPWA	$\bar{K} N \rightarrow \Lambda \pi$	
$+0.06 \pm 0.04$	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda \pi^0$	
$+0.122 \pm 0.078$	DEVENISH 74B		Fixed-t dispersion rel.	
0.20 ± 0.04	LANGBEIN 72	IPWA	$\bar{K} N$ multichannel	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N \bar{K} \rightarrow \Sigma(1840) \rightarrow \Sigma \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
VALUE				
-0.04 or -0.04	¹ MARTIN 77	DPWA	$\bar{K} N$ multichannel	
0.15 ± 0.04	LANGBEIN 72	IPWA	$\bar{K} N$ multichannel	

 $\Sigma(1840)$ FOOTNOTES

- The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- From solution 1 of BAILLON 75; not present in solution 2.

 $\Sigma(1840)$ REFERENCES

MARTIN 77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also 77B	NP B126 266	Martin, Pidcock	(LOUC)
Also 77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
BAILLON 75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
VANHORN 75	NP B87 145		(LBL) IJP
Also 75B	NP B87 157	VanHorn	(LBL) IJP
DEVENISH 74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
LANGBEIN 72	NP B47 477	+Wagner	(MPIM) IJP

Baryon Full Listings

$\Sigma(1880), \Sigma(1915)$

$\Sigma(1880) P_{11}$

$$I(J^P) = 1(\frac{1}{2}^+) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

A P_{11} resonance is suggested by several partial-wave analyses, but with wide variations in the mass and other parameters. We list here all claims which lie well above the $P_{11} \Sigma(1770)$.

$\Sigma(1880)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
~ 1880 OUR ESTIMATE			
1826 ± 20	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1870 ± 10	CAMERON 78B	DPWA	$K^-p \rightarrow N\bar{K}^*$
1847 or 1863	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
1960 ± 30	² BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
1985 ± 50	VANHORN 75	DPWA	$K^-p \rightarrow \Lambda\pi^0$
1898	³ LEA 73	DPWA	Multichannel K-matrix
~ 1850	ARMENTEROS70	IPWA	$\bar{K}N \rightarrow \bar{K}N$
1950 ± 50	BARBARO... 70	DPWA	$K^-N \rightarrow \Lambda\pi$
1920 ± 30	LITCHFIELD 70	DPWA	$K^-N \rightarrow \Lambda\pi$
1850	BAILEY 69	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1882 ± 40	SMART 68	DPWA	$K^-N \rightarrow \Lambda\pi$

$\Sigma(1880)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
86 ± 15	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$
80 ± 10	CAMERON 78B	DPWA	$K^-p \rightarrow N\bar{K}^*$
216 or 220	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel
260 ± 40	² BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
220 ± 140	VANHORN 75	DPWA	$K^-p \rightarrow \Lambda\pi^0$
222	³ LEA 73	DPWA	Multichannel K-matrix
~ 30	ARMENTEROS70	IPWA	$\bar{K}N \rightarrow \bar{K}N$
200 ± 50	BARBARO... 70	DPWA	$K^-N \rightarrow \Lambda\pi$
170 ± 40	LITCHFIELD 70	DPWA	$K^-N \rightarrow \Lambda\pi$
200	BAILEY 69	DPWA	$\bar{K}N \rightarrow \bar{K}N$
222 ± 150	SMART 68	DPWA	$K^-N \rightarrow \Lambda\pi$

$\Sigma(1880)$ DECAY MODES

Mode	Γ_i/Γ
Γ_1 $N\bar{K}$	
Γ_2 $\Lambda\pi$	
Γ_3 $\Sigma\pi$	
Γ_4 $N\bar{K}^*(892), S=1/2, P\text{-wave}$	
Γ_5 $N\bar{K}^*(892), S=3/2, P\text{-wave}$	

$\Sigma(1880)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.06 ± 0.02	GOPAL 80	DPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.27 or 0.27	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	
0.31	³ LEA 73	DPWA	Multichannel K-matrix	
0.20	ARMENTEROS70	IPWA	$\bar{K}N \rightarrow \bar{K}N$	
0.22	BAILEY 69	DPWA	$\bar{K}N \rightarrow \bar{K}N$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1880) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
0.24 or -0.24	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	
-0.12 ± 0.02	² BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$	
+0.05 +0.07 -0.02	VANHORN 75	DPWA	$K^-p \rightarrow \Lambda\pi^0$	
-0.169 ± 0.119	DEVENISH 74B		Fixed- t dispersion rel.	
-0.30	³ LEA 73	DPWA	Multichannel K-matrix	
-0.09 ± 0.04	BARBARO... 70	DPWA	$K^-N \rightarrow \Lambda\pi$	
-0.14 ± 0.03	LITCHFIELD 70	DPWA	$K^-N \rightarrow \Lambda\pi$	
-0.11 ± 0.03	SMART 68	DPWA	$K^-N \rightarrow \Lambda\pi$	

$(\Gamma_1\Gamma_4)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1880) \rightarrow N\bar{K}^*(892), S=1/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_4)^{1/2}/\Gamma$
+0.30 or +0.29 not seen	¹ MARTIN 77	DPWA	$\bar{K}N$ multichannel	
	³ LEA 73	DPWA	Multichannel K-matrix	

$(\Gamma_1\Gamma_5)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1880) \rightarrow N\bar{K}^*(892), S=3/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_5)^{1/2}/\Gamma$
-0.05 ± 0.03	⁴ CAMERON 78B	DPWA	$K^-p \rightarrow N\bar{K}^*$	

$(\Gamma_1\Gamma_5)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1880) \rightarrow N\bar{K}^*(892), S=3/2, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT
+0.11 ± 0.03	CAMERON 78B	DPWA	$K^-p \rightarrow N\bar{K}^*$

$\Sigma(1880)$ FOOTNOTES

- ¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- ² From solution 1 of BAILLON 75; not present in solution 2.
- ³ Only unconstrained states from table 1 of LEA 73 are listed.
- ⁴ The published sign has been changed to be in accord with the baryon-first convention.

$\Sigma(1880)$ REFERENCES

GOPAL 80	Toronto Conf. 159			(RHEL) IJP
CAMERON 78B	NP B146 327	+Frank, Gopal, Kalmus, McPherson+		(RHEL, LOIC) IJP
MARTIN 77	NP B127 349	+Pidcock, Moorhouse		(LOUC, GLAS) IJP
Also	77B NP B126 266	Martin, Pidcock		(LOUC)
Also	77C NP B126 285	Martin, Pidcock		(LOUC) IJP
BAILLON 75	NP B94 39	+Litchfield		(CERN, RHEL) IJP
VANHORN 75	NP B87 145			(LBL) IJP
Also	75B NP B87 157			(LBL) IJP
DEVENISH 74B	NP B81 330	VanHorn		(DESY, NORD, LOUC)
LEA 73	NP B56 77	+Froggatt, Martin		(RHEL, LOUC, GLAS, AARH) IJP
ARMENTEROS 70	Duke Conf. 123	+Martin, Moorhouse+		(CERN, HEID, SACL) IJP
BARBARO... 70	Duke Conf. 173	Barbaro-Galtieri		(LRL) IJP
LITCHFIELD 70	NP B22 269			(RHEL) IJP
BAILEY 69	UCRL 50617 Thesis			(LRL) IJP
SMART 68	PR 169 1330			(LRL) IJP

$\Sigma(1915) F_{15}$

$$I(J^P) = 1(\frac{5}{2}^+) \text{ Status: } ***$$

Discovered by COOL 66. For results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters **111B** (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and invariant-mass distributions in this region used to be listed in a separate entry immediately following. They may be found in our 1986 edition Physics Letters **170B** (1986).

$\Sigma(1915)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 to 1935 (≈ 1915) OUR ESTIMATE			
1937 ± 20	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1894 ± 5	¹ CORDEN 77C		$K^-n \rightarrow \Sigma\pi$
1909 ± 5	¹ CORDEN 77C		$K^-n \rightarrow \Sigma\pi$
1920 ± 10	GOPAL 77	DPWA	$\bar{K}N$ multichannel
1900 ± 4	² CORDEN 76	DPWA	$K^-n \rightarrow \Lambda\pi^-$
1920 ± 30	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
1914 ± 10	HEMINGWAY 75	DPWA	$K^-p \rightarrow \bar{K}N$
1920 ⁺¹⁵ ₋₂₀	VANHORN 75	DPWA	$K^-p \rightarrow \Lambda\pi^0$
1920 ± 5	KANE 74	DPWA	$K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
not seen	DECLAIS 77	DPWA	$\bar{K}N \rightarrow \bar{K}N$
1925 or 1933	³ MARTIN 77	DPWA	$\bar{K}N$ multichannel
1915	DEBELLEFON 76	IPWA	$K^-p \rightarrow \Lambda\pi^0$

$\Sigma(1915)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
80 to 160 (≈ 120) OUR ESTIMATE			
161 ± 20	ALSTON... 78	DPWA	$\bar{K}N \rightarrow \bar{K}N$
107 ± 14	¹ CORDEN 77C		$K^-n \rightarrow \Sigma\pi$
85 ± 13	¹ CORDEN 77C		$K^-n \rightarrow \Sigma\pi$
130 ± 10	GOPAL 77	DPWA	$\bar{K}N$ multichannel
75 ± 14	² CORDEN 76	DPWA	$K^-n \rightarrow \Lambda\pi^-$
70 ± 20	BAILLON 75	IPWA	$\bar{K}N \rightarrow \Lambda\pi$
85 ± 15	HEMINGWAY 75	DPWA	$K^-p \rightarrow \bar{K}N$
102 ± 18	VANHORN 75	DPWA	$K^-p \rightarrow \Lambda\pi^0$
162 ± 25	KANE 74	DPWA	$K^-p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
171 or 173	³ MARTIN 77	DPWA	$\bar{K}N$ multichannel
60	DEBELLEFON 76	IPWA	$K^-p \rightarrow \Lambda\pi^0$

$\Sigma(1915)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	5-15 %
Γ_2 $\Lambda\pi$	seen
Γ_3 $\Sigma\pi$	seen
Γ_4 $\Sigma(1385)\pi$	<5 %
Γ_5 $\Sigma(1385)\pi, P\text{-wave}$	
Γ_6 $\Sigma(1385)\pi, F\text{-wave}$	

The above branching fractions are our estimates, not fits or averages.

See key on page 1343

Baryon Full Listings
 $\Sigma(1915), \Sigma(1940)$ $\Sigma(1915)$ BRANCHING RATIOSSee "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ		
0.05 to 0.15 OUR ESTIMATE						
0.03±0.02	4	GOPAL	80 DPWA $\bar{K}N \rightarrow \bar{K}N$			
0.14±0.05					ALSTON-...	78 DPWA $\bar{K}N \rightarrow \bar{K}N$
0.11±0.04					HEMINGWAY	75 DPWA $K^-p \rightarrow \bar{K}N$
• • • We do not use the following data for averages, fits, limits, etc. • • •						
0.05±0.03	3	GOPAL	77 DPWA See GOPAL 80			
0.08 or 0.08					MARTIN	77 DPWA $\bar{K}N$ multichannel

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1915) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
-0.09 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.10 ± 0.01				
-0.06 ± 0.02	2	CORDEN	76 DPWA $K^-n \rightarrow \Lambda\pi^-$	
-0.09 ± 0.02				
-0.087±0.056	VANHORN	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.09 or -0.09	3	MARTIN	77 DPWA $\bar{K}N$ multichannel	
-0.10				

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1915) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
-0.17±0.01	1	CORDEN	77C $K^-n \rightarrow \Sigma\pi$	
-0.15±0.02				
-0.19±0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.16±0.03				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.05 or -0.05	3	MARTIN	77 DPWA $\bar{K}N$ multichannel	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1915) \rightarrow \Sigma(1385)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
<0.01	CAMERON	78	DPWA $K^-p \rightarrow \Sigma(1385)\pi$	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1915) \rightarrow \Sigma(1385)\pi, F\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1\Gamma_2)^{1/2}/\Gamma$
VALUE				
+0.039±0.009	5	CAMERON	78 DPWA $K^-p \rightarrow \Sigma(1385)\pi$	

 $\Sigma(1915)$ FOOTNOTES

- The two entries for CORDEN 77C are from two different acceptable solutions.
- Preferred solution 3; see CORDEN 76 for other possibilities.
- The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
- The mass and width are fixed to the GOPAL 77 values due to the low elasticity.
- The published sign has been changed to be in accord with the baryon-first convention.

 $\Sigma(1915)$ REFERENCES

PDG	86	PL 1708	Aguilar-Benitez, Porter+	(CERN, CIT+)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
ALSTON-...	78	PR D18 182	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
Also	77	PRL 38 1007	Alston-Garnjost, Kenney+	(LBL, MTHO, CERN) IJP
CAMERON	78	NP B143 189	+Frank, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
CORDEN	77C	NP B125 61	+Cox, Kenyon, O'Neale, Stubbs, Sumorok+	(BIRM) IJP
DECLAIS	77	CERN 77-16	+Duchon, Louvel, Patry, Seguinot+	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC) IJP
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
CORDEN	76	NP B104 382	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
DEBELLEFON	76	NP B109 129	De Bellefion, Berthon	(COEF) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
HEMINGWAY	75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPIM) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
DEVENISH	74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC) IJP
KANE	74	LBL-2452		(LBL) IJP
COOL	66	PRL 16 1228	+Giacomelli, Kycia, Leontic, Lundby+	(BNL) IJP

 $\Sigma(1940) D_{13}$

$I(J^P) = 1(\frac{3}{2}^-)$ Status: * * *

For results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters 111B (1982).

Not all analyses require this state. It is not required by the GOPAL 77 analysis of $K^-n \rightarrow (\Sigma\pi)^-$ nor by the GOPAL 80 analysis of $K^-n \rightarrow K^-n$. See also HEMINGWAY 75. $\Sigma(1940)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
1900 to 1960 (≈ 1940) OUR ESTIMATE			
1920±50	GOPAL	77	DPWA $\bar{K}N$ multichannel
1950±30	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
1949 ⁺⁴⁰ ₋₆₀	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$
1935±80	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
1940±20	LITCHFIELD	74B	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$
1950±20	LITCHFIELD	74C	DPWA $K^-p \rightarrow \Delta(1232)\bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
1886 or 1893	1	MARTIN	77 DPWA $\bar{K}N$ multichannel
1940	DEBELLEFON	76	IPWA $K^-p \rightarrow \Lambda\pi^0, F_{17}$ wave

 $\Sigma(1940)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 to 300 (≈ 220) OUR ESTIMATE			
170±25	CAMERON	78B	DPWA $K^-p \rightarrow N\bar{K}^*$
300±80	GOPAL	77	DPWA $\bar{K}N$ multichannel
150±75	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
160 ⁺⁷⁰ ₋₄₀	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$
330±80	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
60±20	LITCHFIELD	74B	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$
70 ⁺³⁰ ₋₂₀	LITCHFIELD	74C	DPWA $K^-p \rightarrow \Delta(1232)\bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
157 or 159	1	MARTIN	77 DPWA $\bar{K}N$ multichannel

 $\Sigma(1940)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	<20 %
Γ_2 $\Lambda\pi$	seen
Γ_3 $\Sigma\pi$	seen
Γ_4 $\Sigma(1385)\pi$	seen
Γ_5 $\Sigma(1385)\pi, S\text{-wave}$	
Γ_6 $\Lambda(1520)\pi$	seen
Γ_7 $\Lambda(1520)\pi, P\text{-wave}$	
Γ_8 $\Lambda(1520)\pi, F\text{-wave}$	
Γ_9 $\Delta(1232)\bar{K}$	seen
Γ_{10} $\Delta(1232)\bar{K}, S\text{-wave}$	
Γ_{11} $\Delta(1232)\bar{K}, D\text{-wave}$	
Γ_{12} $N\bar{K}^*(892)$	seen
Γ_{13} $N\bar{K}^*(892), S=3/2, S\text{-wave}$	

 $\Sigma(1940)$ BRANCHING RATIOSSee "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
<0.2 OUR ESTIMATE				
0.14 or 0.13	1	MARTIN	77 DPWA $\bar{K}N$ multichannel	
$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Lambda\pi$				
VALUE				
-0.06 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.04 ± 0.02				
-0.05 ^{+0.03} _{-0.02}	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$	
-0.153±0.070				
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.15 or -0.14	1	MARTIN	77 DPWA $\bar{K}N$ multichannel	

Baryon Full Listings

 $\Sigma(1940)$, $\Sigma(2000)$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Sigma\pi$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.08 ± 0.04	GOPAL	77	DPWA $\bar{K}N$ multichannel
-0.14 ± 0.04	KANE	74	DPWA $K^- p \rightarrow \Sigma\pi$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.16 or +0.16	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Lambda(1520)\pi$, P-wave	$(\Gamma_1 \Gamma_7)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
< 0.03	CAMERON	77	DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$
-0.11 ± 0.04	LITCHFIELD	74B	DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Lambda(1520)\pi$, F-wave	$(\Gamma_1 \Gamma_8)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
0.062 ± 0.021	CAMERON	77	DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$
-0.08 ± 0.04	LITCHFIELD	74B	DPWA $K^- p \rightarrow \Lambda(1520)\pi^0$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Delta(1232)\bar{K}$, S-wave	$(\Gamma_1 \Gamma_{10})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.16 ± 0.05	LITCHFIELD	74C	DPWA $K^- p \rightarrow \Delta(1232)\bar{K}$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Delta(1232)\bar{K}$, D-wave	$(\Gamma_1 \Gamma_{11})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.14 ± 0.05	LITCHFIELD	74C	DPWA $K^- p \rightarrow \Delta(1232)\bar{K}$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow \Sigma(1385)\pi$	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.066 ± 0.025	² CAMERON	78	DPWA $K^- p \rightarrow \Sigma(1385)\pi$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(1940) \rightarrow N\bar{K}^*(892)$	$(\Gamma_1 \Gamma_{12})^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.09 ± 0.02	³ CAMERON	78B	DPWA $K^- p \rightarrow N\bar{K}^*$

 $\Sigma(1940)$ FOOTNOTES

- ¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
² The published sign has been changed to be in accord with the baryon-first convention.
³ Upper limits on the D_1 and D_3 waves are each 0.03.

 $\Sigma(1940)$ REFERENCES

PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL)
CAMERON	78	NP B143 189	+FraneK, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
CAMERON	78B	NP B146 327	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
GOVAL	77	PR D16 2746	+Sodhi	(DELF)
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
HEMINGWAY	75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPHIM) IJP
VANHORN	75	NP B87 145	VanHorn	(LBL) IJP
Also	75B	NP B87 157	+Froggatt, Martin	(DESY, NORD, LOUC)
DEVENISH	74B	NP B81 330		(LBL) IJP
KANE	74	LBL-2452		(CERN, HEIDH) IJP
LITCHFIELD	74B	NP B74 19	+Hemingway, Baillon+	(CERN, HEIDH) IJP
LITCHFIELD	74C	NP B74 39	+Hemingway, Baillon+	(CERN, HEIDH) IJP

 $\Sigma(2000) S_{11}$

$$I(J^P) = 1(\frac{1}{2}^-) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

We list here all reported S_{11} states lying above the $\Sigma(1750) S_{11}$.

 $\Sigma(2000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2000 OUR ESTIMATE			
1944 ± 15	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
1955 ± 15	GOPAL	77	DPWA $\bar{K}N$ multichannel
1755 or 1834	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
2004 ± 40	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$

 $\Sigma(2000)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
215 ± 25	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
170 ± 40	GOPAL	77	DPWA $\bar{K}N$ multichannel
413 or 450	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
116 ± 40	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$

 $\Sigma(2000)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$
Γ_2 $\Lambda\pi$
Γ_3 $\Sigma\pi$
Γ_4 $\Lambda(1520)\pi$
Γ_5 $N\bar{K}^*(892)$, $S=1/2$, S-wave
Γ_6 $N\bar{K}^*(892)$, $S=3/2$, D-wave

 $\Sigma(2000)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K}) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
0.51 ± 0.05	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.44 ± 0.05	GOPAL	77	DPWA	See GOPAL 80
0.62 or 0.57	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2000) \rightarrow \Lambda\pi$	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
0.08 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel
-0.19 or -0.18	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel
not seen	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
+0.07 ± 0.02	VANHORN	75	DPWA $K^- p \rightarrow \Lambda\pi^0$
-0.01			

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2000) \rightarrow \Sigma\pi$	$(\Gamma_1 \Gamma_3)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.20 ± 0.04	GOPAL	77	DPWA $\bar{K}N$ multichannel
+0.26 or +0.24	¹ MARTIN	77	DPWA $\bar{K}N$ multichannel

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2000) \rightarrow \Lambda(1520)\pi$	$(\Gamma_1 \Gamma_4)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.081 ± 0.021	² CAMERON	77	DPWA P-wave decay

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2000) \rightarrow N\bar{K}^*(892)$, $S=1/2$, S-wave	$(\Gamma_1 \Gamma_5)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
+0.10 ± 0.02	² CAMERON	78B	DPWA $K^- p \rightarrow N\bar{K}^*$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2000) \rightarrow N\bar{K}^*(892)$, $S=3/2$, D-wave	$(\Gamma_1 \Gamma_6)^{1/2} / \Gamma$		
VALUE	DOCUMENT ID	TECN	COMMENT
-0.07 ± 0.03	CAMERON	78B	DPWA $K^- p \rightarrow N\bar{K}^*$

 $\Sigma(2000)$ FOOTNOTES

- ¹ The two MARTIN 77 values are from a T-matrix pole and from a Breit-Wigner fit.
² The published sign has been changed to be in accord with the baryon-first convention.

 $\Sigma(2000)$ REFERENCES

GOPAL	80	Toronto Conf. 159		(RHEL) IJP
CAMERON	78B	NP B146 327	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	+FraneK, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
MARTIN	77	NP B127 349	+Pidcock, Moorhouse	(LOUC, GLAS) IJP
Also	77B	NP B126 266	Martin, Pidcock	(LOUC)
Also	77C	NP B126 285	Martin, Pidcock	(LOUC) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
VANHORN	75	NP B87 145	VanHorn	(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP

See key on page 1343

Baryon Full Listings
 $\Sigma(2030)$ $\Sigma(2030) F_{17}$

$$I(J^P) = 1(\frac{1}{2}^+) \text{ Status: } ***$$

Discovered by COOL 66 and by WOHL 66. For most results published before 1974 (they are now obsolete), see our 1982 edition Physics Letters 111B (1982).

This entry only includes results from partial-wave analyses. Parameters of peaks seen in cross sections and invariant-mass distributions around 2030 MeV may be found in our 1984 edition, Reviews of Modern Physics 56 No. 2 Pt. II (1984).

 $\Sigma(2030)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2025 to 2040 (≈ 2030) OUR ESTIMATE			
2036 ± 5	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
2038 ± 10	CORDEN	77b	$K^-N \rightarrow N\bar{K}^*$
2040 ± 5	GOPAL	77	DPWA $\bar{K}N$ multichannel
2030 ± 3	¹ CORDEN	76	DPWA $K^-n \rightarrow \Lambda\pi^-$
2035 ± 15	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
2038 ± 10	HEMINGWAY	75	DPWA $K^-p \rightarrow \bar{K}N$
2042 ± 11	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$
2020 ± 6	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
2035 ± 10	LITCHFIELD	74b	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$
2020 ± 30	LITCHFIELD	74c	DPWA $K^-p \rightarrow \Delta(1232)\bar{K}$
2025 ± 10	LITCHFIELD	74d	DPWA $K^-p \rightarrow \Lambda(1820)\pi^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2027 to 2057	GOYAL	77	DPWA $K^-N \rightarrow \Sigma\pi$
2030	DEBELLEFON	76	IPWA $K^-p \rightarrow \Lambda\pi^0$

 $\Sigma(2030)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
150 to 200 (≈ 180) OUR ESTIMATE			
172 ± 10	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$
137 ± 40	CORDEN	77b	$K^-N \rightarrow N\bar{K}^*$
190 ± 10	GOPAL	77	DPWA $\bar{K}N$ multichannel
201 ± 9	¹ CORDEN	76	DPWA $K^-n \rightarrow \Lambda\pi^-$
180 ± 20	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$
172 ± 15	HEMINGWAY	75	DPWA $K^-p \rightarrow \bar{K}N$
178 ± 13	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$
111 ± 5	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$
160 ± 20	LITCHFIELD	74b	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$
200 ± 30	LITCHFIELD	74c	DPWA $K^-p \rightarrow \Delta(1232)\bar{K}$
• • • We do not use the following data for averages, fits, limits, etc. • • •			
260	DECLAIS	77	DPWA $\bar{K}N \rightarrow \bar{K}N$
126 to 195	GOYAL	77	DPWA $K^-N \rightarrow \Sigma\pi$
160	DEBELLEFON	76	IPWA $K^-p \rightarrow \Lambda\pi^0$
70 to 125	LITCHFIELD	74d	DPWA $K^-p \rightarrow \Lambda(1820)\pi^0$

 $\Sigma(2030)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	17–23 %
Γ_2 $\Lambda\pi$	17–23 %
Γ_3 $\Sigma\pi$	5–10 %
Γ_4 ΞK	<2 %
Γ_5 $\Sigma(1385)\pi$	5–15 %
Γ_6 $\Sigma(1385)\pi, F\text{-wave}$	
Γ_7 $\Lambda(1520)\pi$	10–20 %
Γ_8 $\Lambda(1520)\pi, D\text{-wave}$	
Γ_9 $\Lambda(1520)\pi, G\text{-wave}$	
Γ_{10} $\Delta(1232)\bar{K}$	10–20 %
Γ_{11} $\Delta(1232)\bar{K}, F\text{-wave}$	
Γ_{12} $\Delta(1232)\bar{K}, H\text{-wave}$	
Γ_{13} $N\bar{K}^*(892)$	<5 %
Γ_{14} $N\bar{K}^*(892), S=1/2, F\text{-wave}$	
Γ_{15} $N\bar{K}^*(892), S=3/2, F\text{-wave}$	
Γ_{16} $\Lambda(1820)\pi, P\text{-wave}$	

The above branching fractions are our estimates, not fits or averages.

 $\Sigma(2030)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
0.17 to 0.23 OUR ESTIMATE				
0.19 ± 0.03	GOPAL	80	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.18 ± 0.03	HEMINGWAY	75	DPWA $K^-p \rightarrow \bar{K}N$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.15	DECLAIS	77	DPWA $\bar{K}N \rightarrow \bar{K}N$	
0.24 ± 0.02	GOPAL	77	DPWA See GOPAL 80	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
+0.18 ± 0.02	GOPAL	77	DPWA $\bar{K}N$ multichannel	
+0.20 ± 0.01	¹ CORDEN	76	DPWA $K^-n \rightarrow \Lambda\pi^-$	
+0.18 ± 0.02	BAILLON	75	IPWA $\bar{K}N \rightarrow \Lambda\pi$	
+0.20 ± 0.01	VANHORN	75	DPWA $K^-p \rightarrow \Lambda\pi^0$	
+0.195 ± 0.053	DEVENISH	74b	Fixed- t dispersion rel.	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.20	DEBELLEFON	76	IPWA $K^-p \rightarrow \Lambda\pi^0$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
-0.09 ± 0.01	² CORDEN	77c	$K^-n \rightarrow \Sigma\pi$	
-0.06 ± 0.01	² CORDEN	77c	$K^-n \rightarrow \Sigma\pi$	
-0.15 ± 0.03	GOPAL	77	DPWA $\bar{K}N$ multichannel	
-0.10 ± 0.01	KANE	74	DPWA $K^-p \rightarrow \Sigma\pi$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
-0.085 ± 0.02	³ GOYAL	77	DPWA $K^-N \rightarrow \Sigma\pi$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Xi K$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
0.023	MULLER	69b	DPWA $K^-p \rightarrow \Xi K$	
<0.05	BURGUN	68	DPWA $K^-p \rightarrow \Xi K$	
<0.05	TRIPP	67	RVUE $K^-p \rightarrow \Xi K$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1820)\pi, P\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
0.14 ± 0.02	CORDEN	75b	DBC $K^-n \rightarrow N\bar{K}\pi^-$	
0.18 ± 0.04	LITCHFIELD	74d	DPWA $K^-p \rightarrow \Lambda(1820)\pi^0$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi, D\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
+0.114 ± 0.010	⁴ CAMERON	77	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$	
0.14 ± 0.03	LITCHFIELD	74b	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.10 ± 0.03	⁵ CORDEN	75b	DBC $K^-n \rightarrow N\bar{K}\pi^-$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Lambda(1520)\pi, G\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
+0.146 ± 0.010	⁴ CAMERON	77	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$	
0.02 ± 0.02	LITCHFIELD	74b	DPWA $K^-p \rightarrow \Lambda(1520)\pi^0$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta(1232)\bar{K}, F\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
0.16 ± 0.03	LITCHFIELD	74c	DPWA $K^-p \rightarrow \Delta(1232)\bar{K}$	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.17 ± 0.03	⁵ CORDEN	75b	DBC $K^-n \rightarrow N\bar{K}\pi^-$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Delta(1232)\bar{K}, H\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
0.00 ± 0.02	LITCHFIELD	74c	DPWA $K^-p \rightarrow \Delta(1232)\bar{K}$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow \Sigma(1385)\pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
+0.153 ± 0.026	⁴ CAMERON	78	DPWA $K^-p \rightarrow \Sigma(1385)\pi$	

$(\Gamma_i/\Gamma_i)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892), S=1/2, F\text{-wave}$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1/\Gamma_1)^{1/2}/\Gamma$
VALUE				
+0.06 ± 0.03	⁴ CAMERON	78b	DPWA $K^-p \rightarrow N\bar{K}^*$	
-0.02 ± 0.01	CORDEN	77b	$K^-d \rightarrow NN\bar{K}^*$	

Baryon Full Listings

$\Sigma(2030)$, $\Sigma(2070)$, $\Sigma(2080)$

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(2030) \rightarrow N\bar{K}^*(892)$, $S=3/2$, F -wave

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
$+0.04 \pm 0.03$	⁶ CAMERON	78B DPWA	$K^- p \rightarrow N\bar{K}^*$	
-0.12 ± 0.02	CORDEN	77B	$K^- d \rightarrow NN\bar{K}^*$	

$\Sigma(2030)$ FOOTNOTES

- Preferred solution 3; see CORDEN 76 for other possibilities.
- The two entries for CORDEN 77C are from two different acceptable solutions.
- This coupling is extracted from unnormalized data.
- The published sign has been changed to be in accord with the baryon-first convention.
- An upper limit.
- The upper limit on the G_3 wave is 0.03.

$\Sigma(2030)$ REFERENCES

PDG	84	RMP 56 No. 2 Pt. II	Wohl, Cahn, Rittenberg+	(LBL, CIT, CERN)
PDG	82	PL 111B	Roos, Porter, Aguilar-Benitez+	(HELS, CIT, CERN)
GOPAL	80	Toronto Conf. 159		(RHEL) IJP
CAMERON	78	NP B143 189	+Franeek, Gopal, Bacon, Butterworth+	(RHEL, LOIC) IJP
CAMERON	78B	NP B146 327	+Franeek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CAMERON	77	NP B131 399	+Franeek, Gopal, Kalmus, McPherson+	(RHEL, LOIC) IJP
CORDEN	77C	NP B121 365	+Cox, Kenyon, O'Neale, Stubbs, Sumorok+	(BIRM) IJP
CORDEN	77C	NP B125 61	+Cox, Kenyon, O'Neale, Stubbs, Sumorok+	(BIRM) IJP
DECLAIS	77	CERN 77-16	+Duchon, Louvel, Patry, Seguinot+	(CAEN, CERN) IJP
GOPAL	77	NP B119 362	+Ross, VanHorn, McPherson+	(LOIC, RHEL) IJP
GOYAL	77	PR D16 2746	+Sodhi	(DELH) IJP
CORDEN	76	NP B104 382	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
CORDEN	75B	NP B92 365	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
HEMINGWAY	75	NP B91 12	+Eades, Harmsen+	(CERN, HEIDH, MPIM) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
DEVENISH	74B	NP B81 330	+Froggatt, Martin	(DESY, NORD, LOUC)
KANE	74	LBL-2452		(LBL) IJP
LITCHFIELD	74B	NP B74 19	+Hemingway, Baillon+	(CERN, HEIDH) IJP
LITCHFIELD	74C	NP B74 39	+Hemingway, Baillon+	(CERN, HEIDH) IJP
LITCHFIELD	74D	NP B74 12	+Hemingway, Baillon+	(CERN, HEIDH) IJP
MULLER	69B	UCRL 19372 Thesis		(LRL)
BURGUN	68	NP B8 447	+Meyer, Pauli, Tallini+	(SACL, CDEF, RHEL)
TRIPP	67	NP B3 10	+Léth+	(LRL, SLAC, CERN, HEID, SACL)
COOL	66	PRL 16 1228	+Giacomelli, Kycia, Leontic, Lundby+	(BNL)
WOHL	66	PRL 17 107	+Solmitz, Stevenson	(LRL) IJP

$\Sigma(2070)$ F_{15}

$$I(J^P) = 1(\frac{5}{2}^+) \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

This state suggested by BERTHON 70B finds support in GOPAL 80 with new $K^- p$ polarization and $K^- n$ angular distributions. The very broad state seen in KANE 72 is not required in the later (KANE 74) analysis of $\bar{K}N \rightarrow \Sigma \pi$.

$\Sigma(2070)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2070 OUR ESTIMATE			
2051 ± 25	GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$
2057	KANE	72 DPWA	$K^- p \rightarrow \Sigma \pi$
2070 ± 10	BERTHON	70B DPWA	$K^- p \rightarrow \Sigma \pi$

$\Sigma(2070)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
300 ± 30	GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$
906	KANE	72 DPWA	$K^- p \rightarrow \Sigma \pi$
140 ± 20	BERTHON	70B DPWA	$K^- p \rightarrow \Sigma \pi$

$\Sigma(2070)$ DECAY MODES

Mode	Γ_1	Γ_2
$N\bar{K}$	Γ_1	
$\Sigma \pi$		Γ_2

$\Sigma(2070)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K}) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
0.08 ± 0.03	GOPAL	80 DPWA	$\bar{K}N \rightarrow \bar{K}N$	

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(2070) \rightarrow \Sigma \pi$	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
$+0.104$	KANE	72 DPWA	$K^- p \rightarrow \Sigma \pi$	
$+0.12 \pm 0.02$	BERTHON	70B DPWA	$K^- p \rightarrow \Sigma \pi$	

$\Sigma(2070)$ REFERENCES

GOPAL	80	Toronto Conf. 159		(RHEL) IJP
KANE	74	LBL-2452		(LBL)
KANE	72	PR D5 1583		(LBL)
BERTHON	70B	NP B24 417	+Vrana, Butterworth+	(CDEF, RHEL, SACL) IJP

$\Sigma(2080)$ P_{13}

$$I(J^P) = 1(\frac{3}{2}^+) \text{ Status: } **$$

OMITTED FROM SUMMARY TABLE

Suggested by some but not all partial-wave analyses across this region.

$\Sigma(2080)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2080 OUR ESTIMATE			
2091 ± 7	¹ CORDEN	76 DPWA	$K^- n \rightarrow \Lambda \pi^-$
$2070 \text{ to } 2120$	DEBELLEFON	76 IPWA	$K^- p \rightarrow \Lambda \pi^0$
2120 ± 40	BAILLON	75 IPWA	$\bar{K}N \rightarrow \Lambda \pi$ (sol. 1)
2140 ± 40	BAILLON	75 IPWA	$\bar{K}N \rightarrow \Lambda \pi$ (sol. 2)
2082 ± 4	COX	70 DPWA	See CORDEN 76
2070 ± 30	LITCHFIELD	70 DPWA	$K^- N \rightarrow \Lambda \pi$

$\Sigma(2080)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
186 ± 48	¹ CORDEN	76 DPWA	$K^- n \rightarrow \Lambda \pi^-$
100	DEBELLEFON	76 IPWA	$K^- p \rightarrow \Lambda \pi^0$
240 ± 50	BAILLON	75 IPWA	$\bar{K}N \rightarrow \Lambda \pi$ (sol. 1)
200 ± 50	BAILLON	75 IPWA	$\bar{K}N \rightarrow \Lambda \pi$ (sol. 2)
87 ± 20	COX	70 DPWA	See CORDEN 76
250 ± 40	LITCHFIELD	70 DPWA	$K^- N \rightarrow \Lambda \pi$

$\Sigma(2080)$ DECAY MODES

Mode	Γ_1	Γ_2
$N\bar{K}$	Γ_1	
$\Lambda \pi$		Γ_2

$\Sigma(2080)$ BRANCHING RATIOS

See "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma_{\text{total}} \ln N\bar{K} \rightarrow \Sigma(2080) \rightarrow \Lambda \pi$

VALUE	DOCUMENT ID	TECN	COMMENT	$(\Gamma_1 \Gamma_2)^{1/2} / \Gamma$
-0.10 ± 0.03	¹ CORDEN	76 DPWA	$K^- n \rightarrow \Lambda \pi^-$	
-0.10	DEBELLEFON	76 IPWA	$K^- p \rightarrow \Lambda \pi^0$	
-0.13 ± 0.04	BAILLON	75 IPWA	$\bar{K}N \rightarrow \Lambda \pi$ (sol. 1 and 2)	
-0.16 ± 0.03	COX	70 DPWA	See CORDEN 76	
-0.09 ± 0.03	LITCHFIELD	70 DPWA	$K^- N \rightarrow \Lambda \pi$	

$\Sigma(2080)$ FOOTNOTES

- Preferred solution 3; see CORDEN 76 for other possibilities, including a D_{15} at this mass.

$\Sigma(2080)$ REFERENCES

CORDEN	76	NP B104 382	+Cox, Dartnell, Kenyon, O'Neale+	(BIRM) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
Also	75	NP B90 1	De Bellefon, Berthon, Brunet+	(CDEF, SACL) IJP
BAILLON	75	NP B94 39	+Litchfield	(CERN, RHEL) IJP
COX	70	NP B19 61	+Islam, Colley+	(BIRM, EDIN, GLAS, LOIC) IJP
LITCHFIELD	70	NP B22 269		(RHEL) IJP

See key on page 1343

Baryon Full Listings
 $\Sigma(2100)$, $\Sigma(2250)$ $\Sigma(2100) G_{17}$ $I(J^P) = 1(\frac{7}{2}^-)$ Status: *

OMITTED FROM SUMMARY TABLE

 $\Sigma(2100)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2100 OUR ESTIMATE			
2060 \pm 20	BARBARO-...	70	DPWA $K^- p \rightarrow \Lambda\pi^0$
2120 \pm 30	BARBARO-...	70	DPWA $K^- p \rightarrow \Sigma\pi$

 $\Sigma(2100)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
70 \pm 30	BARBARO-...	70	DPWA $K^- p \rightarrow \Lambda\pi^0$
135 \pm 30	BARBARO-...	70	DPWA $K^- p \rightarrow \Sigma\pi$

 $\Sigma(2100)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	<10 %
Γ_2 $\Lambda\pi$	seen
Γ_3 $\Sigma\pi$	seen

 $\Sigma(2100)$ BRANCHING RATIOSSee "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2100) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT
-0.07 ± 0.02	BARBARO-...	70	DPWA $K^- p \rightarrow \Lambda\pi^0$

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2100) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT
$+0.13 \pm 0.02$	BARBARO-...	70	DPWA $K^- p \rightarrow \Sigma\pi$

 $\Sigma(2100)$ REFERENCES

BARBARO-... 70 Duke Conf. 173 Barbaro-Galtieri (LRL) IJP

 $\Sigma(2250)$ $I(J^P) = 1(?)$ Status: ***Results from partial-wave analyses are too weak to warrant separating them from the production and cross-section experiments. LASINSKI 71 in $\bar{K}N$ using a Pomeron + resonances model, and DEBELLEFON 76, DEBELLEFON 77, and DEBELLEFON 78 in energy-dependent partial-wave analyses of $\bar{K}N \rightarrow \Lambda\pi$, $\Sigma\pi$, and $N\bar{K}$, respectively, suggest two resonances around this mass. $\Sigma(2250)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
2210 to 2280 OUR ESTIMATE			
2270 \pm 50	DEBELLEFON 78	DPWA	D_5 wave
2210 \pm 30	DEBELLEFON 78	DPWA	G_9 wave
2275 \pm 20	DEBELLEFON 77	DPWA	D_5 wave
2215 \pm 20	DEBELLEFON 77	DPWA	G_9 wave
2300 \pm 30	¹ DEBELLEFON 75b	HBC	$K^- p \rightarrow \Xi^* K^0$
2251 $^{+30}_{-20}$	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0, F_5$ wave
2280 \pm 14	AGUILAR-...	70b	HBC $K^- p$ 3.9, 4.6 GeV/c
2237 \pm 11	BRICMAN 70	CNTR	Total, charge exchange
2255 \pm 10	COOL 70	CNTR	$K^- p, K^- d$ total
2250 \pm 7	BUGG 68	CNTR	$K^- p, K^- d$ total
• • • We do not use the following data for averages, fits, limits, etc. • • •			
2260	DEBELLEFON 76	IPWA	D_5 wave
2215	DEBELLEFON 76	IPWA	G_9 wave
2250 \pm 20	LU 70	CNTR	$\gamma p \rightarrow K^+ Y^*$
2245	BLANPIED 65	CNTR	$\gamma p \rightarrow K^+ Y^*$
2299 \pm 6	BOCK 65	HBC	$\bar{p}p$ 5.7 GeV/c

 $\Sigma(2250)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
60 to 150 (≈ 100) OUR ESTIMATE			
120 \pm 40	DEBELLEFON 78	DPWA	D_5 wave
80 \pm 20	DEBELLEFON 78	DPWA	G_9 wave
70 \pm 20	DEBELLEFON 77	DPWA	D_5 wave
60 \pm 20	DEBELLEFON 77	DPWA	G_9 wave
130 \pm 20	¹ DEBELLEFON 75b	HBC	$K^- p \rightarrow \Xi^* K^0$
192 \pm 30	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0, F_5$ wave
100 \pm 20	AGUILAR-...	70b	HBC $K^- p$ 3.9, 4.6 GeV/c
164 \pm 50	BRICMAN 70	CNTR	Total, charge exchange
230 \pm 20	BUGG 68	CNTR	$K^- p, K^- d$ total
• • • We do not use the following data for averages, fits, limits, etc. • • •			
100	DEBELLEFON 76	IPWA	D_5 wave
140	DEBELLEFON 76	IPWA	G_9 wave
170	COOL 70	CNTR	$K^- p, K^- d$ total
125	LU 70	CNTR	$\gamma p \rightarrow K^+ Y^*$
150	BLANPIED 65	CNTR	$\gamma p \rightarrow K^+ Y^*$
21 $^{+17}_{-21}$	BOCK 65	HBC	$\bar{p}p$ 5.7 GeV/c

 $\Sigma(2250)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $N\bar{K}$	<10 %
Γ_2 $\Lambda\pi$	seen
Γ_3 $\Sigma\pi$	seen
Γ_4 $N\bar{K}\pi$	
Γ_5 $\Xi(1530)K$	

The above branching fractions are our estimates, not fits or averages.

 $\Sigma(2250)$ BRANCHING RATIOSSee "Sign conventions for resonance couplings" in the Note on Λ and Σ Resonances.

$\Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
<0.1 OUR ESTIMATE			
0.08 \pm 0.02	DEBELLEFON 78	DPWA	D_5 wave
0.02 \pm 0.01	DEBELLEFON 78	DPWA	G_9 wave

$(J+\frac{1}{2}) \times \Gamma(N\bar{K})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT
0.16 ± 0.12	BRICMAN 70	CNTR	Total, charge exchange
0.42	COOL 70	CNTR	$K^- p, K^- d$ total
0.47	BUGG 68	CNTR	

$(\Gamma_1\Gamma_2)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2250) \rightarrow \Lambda\pi$	DOCUMENT ID	TECN	COMMENT
-0.16 ± 0.03	VANHORN 75	DPWA	$K^- p \rightarrow \Lambda\pi^0, F_5$ wave
• • • We do not use the following data for averages, fits, limits, etc. • • •			
+0.11	DEBELLEFON 76	IPWA	D_5 wave
-0.10	DEBELLEFON 76	IPWA	G_9 wave
-0.18	BARBARO-...	70	DPWA $K^- p \rightarrow \Lambda\pi^0, G_9$ wave

$(\Gamma_1\Gamma_3)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2250) \rightarrow \Sigma\pi$	DOCUMENT ID	TECN	COMMENT
$+0.06 \pm 0.02$	DEBELLEFON 77	DPWA	D_5 wave
-0.03 ± 0.02	DEBELLEFON 77	DPWA	G_9 wave
+0.07	BARBARO-...	70	DPWA $K^- p \rightarrow \Sigma\pi, G_9$ wave

$\Gamma(N\bar{K})/\Gamma(\Sigma\pi)$	DOCUMENT ID	TECN	COMMENT
<0.18	BARNES 69	HBC	1 standard dev. limit

$\Gamma(\Lambda\pi)/\Gamma(\Sigma\pi)$	DOCUMENT ID	TECN	COMMENT
<0.18	BARNES 69	HBC	1 standard dev. limit

$(\Gamma_1\Gamma_5)^{1/2}/\Gamma_{\text{total}}$ in $N\bar{K} \rightarrow \Sigma(2250) \rightarrow \Xi(1530)K$	DOCUMENT ID	TECN	COMMENT
0.18 \pm 0.04	¹ DEBELLEFON 75b	HBC	$K^- p \rightarrow \Xi^* K^0$

 $\Sigma(2250)$ FOOTNOTES¹ Seen in the (initial and final state) D_5 wave. Isospin not determined.

Baryon Full Listings

 $\Sigma(2250)$, $\Sigma(2455)$ Bumps, $\Sigma(2620)$ Bumps, $\Sigma(3000)$ Bumps $\Sigma(2250)$ REFERENCES

DEBELLEFON	78	NC 42A 403	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
DEBELLEFON	77	NC 37A 175	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
DEBELLEFON	76	NP B109 129	De Bellefon, Berthon	(CDEF) IJP
Also	75	NP B90 1	De Bellefon, Berthon, Brunet+	(CDEF, SACL) IJP
DEBELLEFON	75B	NC 28A 289	De Bellefon, Berthon, Billoir+	(CDEF, SACL) IJP
VANHORN	75	NP B87 145		(LBL) IJP
Also	75B	NP B87 157	VanHorn	(LBL) IJP
LASINSKI	71	NP B29 125		(EFI) IJP
AGUILAR...	70B	PRL 25 58	Aguilar-Benitez, Barnes, Bassano+	(BNL, SYRA)
BARBARO...	70	Duke Conf. 173	Barbaro-Gallieri	(LRL) IJP
BRICMAN	70	PL 31B 152	+Ferro-Luzzi, Perreau+	(CERN, CAEN, SACL)
COOL	70	PR D1 1887	+Giacomelli, Kycia, Leontic, Li+	(BNL) I
Also	66	PRL 16 1228	Cool, Giacomelli, Kycia, Leontic, Lundby+	(BNL) I
LU	70	PR D2 1846	+Greenberg, Hughes, Minehart, Mori+	(YALE)
BARNES	69	PRL 22 479	+Flaminio, Montanet, Samios+	(BNL, SYRA)
BUGG	68	PR 168 1466	+Gilmore, Knight+	(RHEL, BIRM, CAVE) I
BLANPIED	65	PRL 14 741	+Greenberg, Hughes, Kitching, Lu+	(YALE, CEA)
BOCK	65	PL 17 166	+Cooper, French, Kinson+	(CERN, SACL)

 $\Sigma(2455)$ Bumps

$I(J^P) = 1(?)^2$ Status: **

OMITTED FROM SUMMARY TABLE

There is also some slight evidence for Y^* states in this mass region from the reaction $\gamma p \rightarrow K^+ X$ — see GREENBERG 68. $\Sigma(2455)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2455 OUR ESTIMATE			
2455 ± 10	ABRAMS 70	CNTR	$K^- p, K^- d$ total
2455 ± 7	BUGG 68	CNTR	$K^- p, K^- d$ total

 $\Sigma(2455)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
140	ABRAMS 70	CNTR	$K^- p, K^- d$ total
100 ± 20	BUGG 68	CNTR	

 $\Sigma(2455)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$

 $\Sigma(2455)$ BRANCHING RATIOS

$(J+\frac{1}{2}) \times \Gamma(N\bar{K}) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
0.39	ABRAMS 70	CNTR	$K^- p, K^- d$ total	
0.05 ± 0.05	¹ BRICMAN 70	CNTR	Total, charge exchange	
0.3	BUGG 68	CNTR		

 $\Sigma(2455)$ FOOTNOTES¹ Fit of total cross section given by BRICMAN 70 is poor in this region. $\Sigma(2455)$ REFERENCES

ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL) I
Also	67E	PRL 19 678	Abrams, Cool, Giacomelli, Kycia, Leontic+	(BNL)
BRICMAN	70	PL 31B 152	+Ferro-Luzzi, Perreau+	(CERN, CAEN, SACL)
BUGG	68	PR 168 1466	+Gilmore, Knight+	(RHEL, BIRM, CAVE) I
GREENBERG	68	PRL 20 221	+Hughes, Lu, Minehart+	(YALE)

 $\Sigma(2620)$ Bumps

$I(J^P) = 1(?)^2$ Status: **

OMITTED FROM SUMMARY TABLE

 $\Sigma(2620)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
≈ 2620 OUR ESTIMATE			
2542 ± 22	DIBIANCA 75	DBC	$K^- N \rightarrow \Xi K \pi$
2620 ± 15	ABRAMS 70	CNTR	$K^- p, K^- d$ total

 $\Sigma(2620)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
221 ± 81	DIBIANCA 75	DBC	$K^- N \rightarrow \Xi K \pi$
175	ABRAMS 70	CNTR	$K^- p, K^- d$ total

 $\Sigma(2620)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$

 $\Sigma(2620)$ BRANCHING RATIOS

$(J+\frac{1}{2}) \times \Gamma(N\bar{K}) / \Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1 / Γ
0.32	ABRAMS 70	CNTR	$K^- p, K^- d$ total	
0.36 ± 0.12	BRICMAN 70	CNTR	Total, charge exchange	

 $\Sigma(2620)$ REFERENCES

DIBIANCA	75	NP B98 137	+Endorf	(CMU)
ABRAMS	70	PR D1 1917	+Cool, Giacomelli, Kycia, Leontic, Li+	(BNL) I
Also	67E	PRL 19 678	Abrams, Cool, Giacomelli, Kycia, Leontic+	(BNL)
BRICMAN	70	PL 31B 152	+Ferro-Luzzi, Perreau+	(CERN, CAEN, SACL)

 $\Sigma(3000)$ Bumps

$I(J^P) = 1(?)^2$ Status: *

OMITTED FROM SUMMARY TABLE

Seen as an enhancement in $\Lambda \pi$ and $\bar{K} N$ invariant mass spectra and in the missing mass of neutrals recoiling against a K^0 . $\Sigma(3000)$ MASS

VALUE (MeV)	DOCUMENT ID	TECN	CHG	COMMENT
≈ 3000 OUR ESTIMATE				
3000	EHRlich 66	HBC	0	$\pi^- p$ 7.91 GeV/c

 $\Sigma(3000)$ DECAY MODES

Mode
Γ_1 $N\bar{K}$
Γ_2 $\Lambda \pi$

 $\Sigma(3000)$ REFERENCES

EHRlich	66	PR 152 1194	+Selove, Yuta	(PENN) I
---------	----	-------------	---------------	----------

See key on page 1343

Baryon Full Listings

$\Sigma(3170)$ Bumps, Ξ^0

$\Sigma(3170)$ Bumps

$$I(J^P) = 1(?)^? \text{ Status: } *$$

OMITTED FROM SUMMARY TABLE

Seen by AMIRZADEH 79 as a narrow 6.5-standard-deviation enhancement in the reaction $K^- p \rightarrow Y^{*+} \pi^-$ using data from independent high statistics bubble chamber experiments at 8.25 and 6.5 GeV/c. The dominant decay modes are multibody, multistrange final states and the production is via isospin-3/2 baryon exchange. Isospin 1 is favored.

Not seen in a $K^- p$ experiment in LASS at 11 GeV/c (ASTON 85B).

$\Sigma(3170)$ MASS (PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
≈ 3170 OUR ESTIMATE				
3170 ± 5	35	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$

$\Sigma(3170)$ WIDTH (PRODUCTION EXPERIMENTS)

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
<20	35	¹ AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$

$\Sigma(3170)$ DECAY MODES (PRODUCTION EXPERIMENTS)

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Lambda K \bar{K} \pi$'s	seen
Γ_2 $\Sigma K \bar{K} \pi$'s	seen
Γ_3 $\Xi K \pi$'s	seen

$\Sigma(3170)$ BRANCHING RATIOS (PRODUCTION EXPERIMENTS)

$\Gamma(\Lambda K \bar{K} \pi \text{'s})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
VALUE				
seen	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$	
$\Gamma(\Sigma K \bar{K} \pi \text{'s})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
VALUE				
seen	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$	
$\Gamma(\Xi K \pi \text{'s})/\Gamma_{\text{total}}$	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
VALUE				
seen	AMIRZADEH 79	HBC	$K^- p \rightarrow Y^{*+} \pi^-$	

$\Sigma(3170)$ FOOTNOTES (PRODUCTION EXPERIMENTS)

¹ Observed width consistent with experimental resolution.

$\Sigma(3170)$ REFERENCES (PRODUCTION EXPERIMENTS)

ASTON 85B	PR D32 2270	+Carnegie+	(SLAC, CARL, CNRC, CINC)
AMIRZADEH 79	PL 89B 125	+	(BIRM, CERN, GLAS, MSU, CURIN, CAVE+)
Also 80	Toronto Conf. 263	Kinson+	(BIRM, CERN, GLAS, MSU, CURIN)

Ξ BARYONS ($S = -2, I = 1/2$)

$$\Xi^0 = uss, \Xi^- = dss$$

Ξ^0

$$I(J^P) = \frac{1}{2}(\frac{1}{2}^+) \text{ Status: } * * * *$$

The parity has not actually been measured, but + is of course expected.

Ξ^0 MASS

The fit uses the Ξ^0 , Ξ^- , and Ξ^+ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN
1314.9 ± 0.6 OUR FIT			
1314.8 ± 0.8 OUR AVERAGE			
1315.2 ± 0.92	49	WILQUET 72	HLBC
1313.4 ± 1.8	1	PALMER 68	HBC

$$m_{\Xi^-} - m_{\Xi^0}$$

The fit uses the Ξ^0 , Ξ^- , and Ξ^+ mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
6.4 ± 0.6 OUR FIT				
6.3 ± 0.7 OUR AVERAGE				
6.9 ± 2.2	29	LONDON 66	HBC	
6.1 ± 0.9	88	PJERROU 65B	HBC	
6.8 ± 1.6	23	JAUNEAU 63	FBC	
6.1 ± 1.6	45	CARMONY 64B	HBC	See PJERROU 65B

• • • We do not use the following data for averages, fits, limits, etc. • • •

Ξ^0 MEAN LIFE

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
2.90 ± 0.09 OUR AVERAGE				
2.83 ± 0.16	6300	¹ ZECH 77	SPEC	Neutral hyperon beam
2.88 ^{+0.21} _{-0.19}	652	BALTAY 74	HBC	1.75 GeV/c $K^- p$
2.90 ^{+0.32} _{-0.27}	157	² MAYEUR 72	HLBC	2.1 GeV/c K^-
3.07 ^{+0.22} _{-0.20}	340	DAUBER 69	HBC	
3.0 ± 0.5	80	PJERROU 65B	HBC	
2.5 ^{+0.4} _{-0.3}	101	HUBBARD 64	HBC	
3.9 ^{+1.4} _{-0.8}	24	JAUNEAU 63	FBC	
3.5 ^{+1.0} _{-0.8}	45	CARMONY 64B	HBC	See PJERROU 65B

¹ The ZECH 77 result is $\tau_{\Xi^0} = [2.77 - (\tau_{\Lambda} - 2.69)] \times 10^{-10}$ s, in which we use $\tau_{\Lambda} = 2.63 \times 10^{-10}$ s.

² The MAYEUR 72 value is modified by the erratum.

Ξ^0 MAGNETIC MOMENT

See the Note on Baryon Magnetic Moments in the Λ Listings.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN
-1.250 ± 0.014 OUR AVERAGE			
-1.253 ± 0.014	270k	COX 81	SPEC
-1.20 ± 0.06	42k	BUNCE 79	SPEC

Baryon Full Listings

≡⁰

≡⁰ DECAY MODES

Mode	Fraction (Γ _i /Γ)	Confidence level
Γ ₁ Λπ ⁰	(99.54 ± 0.05) %	
Γ ₂ Λγ	(1.06 ± 0.16) × 10 ⁻³	
Γ ₃ Σ ⁰ γ	(3.5 ± 0.4) × 10 ⁻³	
Γ ₄ Σ ⁺ e ⁻ ν _e	< 1.1 × 10 ⁻³	90%
Γ ₅ Σ ⁺ μ ⁻ ν _μ	< 1.1 × 10 ⁻³	90%

**ΔS = ΔQ (SQ) violating modes or
ΔS = 2 forbidden (S2) modes**

Γ ₆ Σ ⁻ e ⁺ ν _e	SQ < 9 × 10 ⁻⁴	90%
Γ ₇ Σ ⁻ μ ⁺ ν _μ	SQ < 9 × 10 ⁻⁴	90%
Γ ₈ ρπ ⁻	S2 < 4 × 10 ⁻⁵	90%
Γ ₉ ρe ⁻ ν _e	S2 < 1.3 × 10 ⁻³	
Γ ₁₀ ρμ ⁻ ν _μ	S2 < 1.3 × 10 ⁻³	

CONSTRAINED FIT INFORMATION

An overall fit to 2 branching ratios uses 2 measurements and one constraint to determine 3 parameters. The overall fit has a χ² = 0.0 for 0 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients (δx_iδx_j)/(δx_iδx_j), in percent, from the fit to the branching fractions, x_i ≡ Γ_i/Γ_{total}. The fit constrains the x_i whose labels appear in this array to sum to one.

x ₂	-35	
x ₃	-94	0
	x ₁	x ₂

≡⁰ BRANCHING RATIOS

Γ(Λγ)/Γ(Λπ ⁰)	Γ ₂ /Γ ₁			
VALUE (units 10 ⁻³)	EVTS	DOCUMENT ID	TECN	COMMENT
1.06 ± 0.16 OUR FIT				
1.06 ± 0.12 ± 0.11	116	JAMES	90 SPEC	FNAL hyperons
• • • We do not use the following data for averages, fits, limits, etc. • • •				
5 ± 5	1	YEH	74 HBC	Effective denom.=200

Γ(Σ ⁰ γ)/Γ(Λπ ⁰)	Γ ₃ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
3.6 ± 0.4 OUR FIT					
3.56 ± 0.42 ± 0.10	85	TEIGE	89 SPEC	FNAL hyperons	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 8	90		BENSINGER	88 MPS2	K ⁻ W 6 GeV/c
< 65	90	0-1	YEH	74 HBC	Effective denom.=60

Γ(Σ ⁺ e ⁻ ν _e)/Γ(Λπ ⁰)	Γ ₄ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.1	90	0	YEH	74 HBC	Effective denom.=2100
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.5			DAUBER	69 HBC	
< 7			HUBBARD	66 HBC	

Γ(Σ ⁺ μ ⁻ ν _μ)/Γ(Λπ ⁰)	Γ ₅ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.1	90	0	YEH	74 HBC	Effective denom.=2100
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.5			DAUBER	69 HBC	
< 7			HUBBARD	66 HBC	

Γ(Σ ⁻ e ⁺ ν _e)/Γ(Λπ ⁰)	Γ ₆ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.9	90	0	YEH	74 HBC	Effective denom.=2500
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.5			DAUBER	69 HBC	
< 6			HUBBARD	66 HBC	

Γ(Σ ⁻ μ ⁺ ν _μ)/Γ(Λπ ⁰)	Γ ₇ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.9	90	0	YEH	74 HBC	Effective denom.=2500
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 1.5			DAUBER	69 HBC	
< 6			HUBBARD	66 HBC	

Γ(ρπ ⁻)/Γ(Λπ ⁰)	Γ ₈ /Γ ₁				
VALUE (units 10 ⁻⁵)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 3.6	90		GEWENIGER	75 SPEC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 180	90	0	YEH	74 HBC	Effective denom.=1300
< 90			DAUBER	69 HBC	
< 500			HUBBARD	66 HBC	

Γ(ρe ⁻ ν _e)/Γ(Λπ ⁰)	Γ ₉ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.3			DAUBER	69 HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.4	90	0	YEH	74 HBC	Effective denom.=670
< 6			HUBBARD	66 HBC	


Γ(ρμ ⁻ ν _μ)/Γ(Λπ ⁰)	Γ ₁₀ /Γ ₁				
VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 1.3			DAUBER	69 HBC	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
< 3.5	90	0	YEH	74 HBC	Effective denom.=664
< 6			HUBBARD	66 HBC	

≡⁰ DECAY PARAMETERS

See the Note on Baryon Decay Parameters in the neutron Listings.

α(≡ ⁰) α ₋ (Λ)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.264 ± 0.013 OUR AVERAGE				Error includes scale factor of 2.1. See the Ideogram below.
-0.260 ± 0.004 ± 0.005	300k	HANDLER	82 SPEC	FNAL hyperons
-0.317 ± 0.027	6075	BUNCE	78 SPEC	FNAL hyperons
-0.35 ± 0.06	505	BALTAY	74 HBC	K ⁻ p 1.75 GeV/c
-0.28 ± 0.06	739	DAUBER	69 HBC	K ⁻ p 1.7-2.6 GeV/c

WEIGHTED AVERAGE
-0.264 ± 0.013 (Error scaled by 2.1)



α FOR ≡⁰ → Λπ⁰
The above average, α(≡⁰)α₋(Λ) = -0.264 ± 0.013, where the error includes a scale factor of 2.1, divided by our current average α₋(Λ) = 0.642 ± 0.013, gives the following value for α(≡⁰).

VALUE	DOCUMENT ID
-0.411 ± 0.022 OUR EVALUATION	Error includes scale factor of 2.1.

φ ANGLE FOR ≡ ⁰ → Λπ ⁰	(tan φ = β/γ)			
VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
21 ± 12 OUR AVERAGE				
16 ± 17	652	BALTAY	74 HBC	1.75 GeV/c K ⁻ p
38 ± 19	739	3 DAUBER	69 HBC	
- 8 ± 30	146	4 BERGE	66 HBC	

³ DAUBER 69 uses α_Λ = 0.647 ± 0.020.
⁴ The errors have been multiplied by 1.2 due to approximations used for the ≡ polarization; see DAUBER 69 for a discussion.

α FOR ≡ ⁰ → Λγ	EVTS	DOCUMENT ID	TECN	COMMENT
+0.43 ± 0.44	87	JAMES	90 SPEC	FNAL hyperons

See key on page 1343

Baryon Full Listings

Ξ^0, Ξ^-

α FOR $\Xi^0 \rightarrow \Sigma^0 \gamma$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
$+0.20 \pm 0.32 \pm 0.05$	85	TEIGE	89	SPEC FNAL hyperons

Ξ^- MEAN LIFE

Measurements with an error $> 0.2 \times 10^{-10}$ s or with systematic errors not included have been omitted.

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.639 ± 0.015 OUR AVERAGE				
1.652 ± 0.051	32k	BOURQUIN	84	SPEC Hyperon beam
1.665 ± 0.065	41k	BOURQUIN	79	SPEC Hyperon beam
1.609 ± 0.028	4286	HEMINGWAY	78	HBC $4.2 \text{ GeV}/c \ K^- p$
1.67 ± 0.08		DIBIANCA	75	DBC $4.9 \text{ GeV}/c \ K^- d$
1.63 ± 0.03	4303	BALTAY	74	HBC $1.75 \text{ GeV}/c \ K^- p$
1.73 ± 0.08 -0.07	680	MAYEUR	72	HLBC $2.1 \text{ GeV}/c \ K^-$
1.61 ± 0.04	2610	DAUBER	69	HBC
1.80 ± 0.16	299	LONDON	66	HBC
1.70 ± 0.12	246	PJERROU	65B	HBC
1.69 ± 0.07	794	HUBBARD	64	HBC
1.86 ± 0.15 -0.14	517	JAUNEAU	63D	FBC

Ξ^0 REFERENCES

JAMES	90	PRL 64 843	+Heller, Border, Dworkin+ (MINN, MICH, WISC, RUTG)
TEIGE	89	PRL 63 2717	+Berevas, Caraccappa, Devlin+ (RUTG, MICH, MINN)
BENSINGER	88	PL B215 195	+Fortner, Kirsch, Piekarz+ (BRAN, DUKE, NDAM, MASD)
HANDLER	82	PR D25 639	+Grobel, Pondrom+ (WISC, MICH, MINN, RUTG)
COX	81	PR 46 877	+Dworkin+ (MICH, WISC, RUTG, MINN, BNL)
BUNCE	79	PL 85B 386	+Overseth, Cox+ (BNL, MICH, RUTG, WISC)
BUNCE	78	PR D18 633	+Handler, March, Martin+ (WISC, MICH, RUTG)
ZECH	77	NP B124 413	+Dyda, Navarra+ (SIEG, CERN, DORT, HEIDH)
GEWENIGER	75	PL 57B 193	+Gjesdal, Presser+ (CERN, HEIDH)
BALTAY	74	PR D9 49	+Bridgewater, Cooper, Gershwin+ (COLU, BING) J
YEH	74	PR D10 3545	+Gaigalas, Smith, Zendle, Baltay+ (BING, COLU)
MAYEUR	72	NP B47 333	+VanBinst, Wilquet+ (BRUX, CERN, TUFTS, LOUC)
Also	73	NP B53 268 erratum	Mayeur (BRUX, CERN, TUFTS, LOUC)
WILQUET	72	PL 42B 372	+Flagigne, Guy+ (BRUX, CERN, TUFTS, LOUC)
DAUBER	69	PR 179 1262	+Berge, Hubbard, Merrill, Miller (LRL)
PALMER	68	PL 26B 323	+Radojicic, Rau, Richardson+ (BNL, SYRA)
BERGE	66	PR 147 945	+Eberhard, Hubbard, Merrill+ (LRL)
HUBBARD	66	UCRL 11510 Thesis	(LRL)
LONDON	66	PR 143 1034	+Rau, Goldberg, Lichtman+ (BNL, SYRA)
PJERROU	65B	PRL 14 275	+Schlein, Slater, Smith, Stork, Ticho (UCLA)
Also	65	Thesis	Pjerrou (UCLA)
CARMONY	64B	PR 12 482	+Pjerrou, Schlein, Slater, Stork+ (UCLA)
HUBBARD	64	PR 135B 183	+Berge, Kalbfleisch, Shafer+ (LRL)
JAUNEAU	63	PL 4 49	+ (EPOL, CERN, LOUC, RHEL, BERG)
Also	63C	Siena Conf. 1 1	Jauneau+ (EPOL, CERN, LOUC, RHEL, BERG)



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: * * * *

The parity has not actually been measured, but + is of course expected.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1986 edition (Physics Letters 170B) or in earlier editions.

Ξ^- MASS

The fit uses the Ξ^- , Ξ^+ , and Ξ^0 mass and mass difference measurements. It assumes the Ξ^- and Ξ^+ masses are the same.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1321.32 ± 0.13 OUR FIT				
1321.34 ± 0.14 OUR AVERAGE				
1321.46 ± 0.34	632	DIBIANCA	75	DBC $4.9 \text{ GeV}/c \ K^- d$
1321.12 ± 0.41	268	WILQUET	72	HLBC
1321.87 ± 0.51	195	¹ GOLDWASSER	70	HBC $5.5 \text{ GeV}/c \ K^- p$
1321.67 ± 0.52	6	CHIEN	66	HBC $6.9 \text{ GeV}/c \ \bar{p} p$
1321.4 ± 1.1	299	LONDON	66	HBC
1321.3 ± 0.4	149	PJERROU	65B	HBC
1321.1 ± 0.3	241	² BADIER	64	HBC
1321.4 ± 0.4	517	² JAUNEAU	63D	FBC
1321.1 ± 0.65	62	² SCHNEIDER	63	HBC

¹ GOLDWASSER 70 uses $m_\Lambda = 1115.58$ MeV.

² These masses have been increased 0.09 MeV because the Λ mass increased.

Ξ^+ MASS

The fit uses the Ξ^- , Ξ^+ , and Ξ^0 mass and mass difference measurements. It assumes the Ξ^- and Ξ^+ masses are the same.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1321.32 ± 0.13 OUR FIT				
1321.20 ± 0.33 OUR AVERAGE				
1321.6 ± 0.8	35	VOTRUBA	72	HBC $10 \text{ GeV}/c \ K^+ p$
1321.2 ± 0.4	34	STONE	70	HBC
1320.69 ± 0.93	5	CHIEN	66	HBC $6.9 \text{ GeV}/c \ \bar{p} p$

$(m_{\Xi^-} - m_{\Xi^+}) / m_{\text{average}}$

A test of CPT Invariance. We calculate it from the average Ξ^- and Ξ^+ masses above.

VALUE	DOCUMENT ID
$(1.1 \pm 2.7) \times 10^{-4}$ OUR EVALUATION	

Ξ^+ MEAN LIFE

VALUE (10^{-10} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.6 ± 0.3	34	STONE	70	HBC
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1.55 ± 0.35 -0.20	35	³ VOTRUBA	72	HBC $10 \text{ GeV}/c \ K^+ p$
1.9 ± 0.7 -0.5	12	³ SHEN	67	HBC
1.51 ± 0.55	5	³ CHIEN	66	HBC $6.9 \text{ GeV}/c \ \bar{p} p$

³ The error is statistical only.

$(\tau_{\Xi^-} - \tau_{\Xi^+}) / \tau_{\text{average}}$

A test of CPT invariance. Calculated from the Ξ^- and Ξ^+ mean lives, above.

VALUE	DOCUMENT ID
0.02 ± 0.18 OUR EVALUATION	

Ξ^- MAGNETIC MOMENT

See the Note on Baryon Magnetic Moments in the Λ Listings.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN	COMMENT
-0.6807 ± 0.0025 OUR AVERAGE				
-0.6505 ± 0.0025	4.36M	DURYEA	92	SPEC $800 \text{ GeV} \ p \ \text{Be}$
$-0.661 \pm 0.036 \pm 0.036$	44k	TROST	89	SPEC $\Xi^- \sim 250 \text{ GeV}$
-0.69 ± 0.04	218k	RAMEIKA	84	SPEC $400 \text{ GeV} \ p \ \text{Be}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
$-0.674 \pm 0.021 \pm 0.020$	122k	HO	90	SPEC See DURYEA 92
-2.1 ± 0.8	2436	COOL	74	OSPK $1.8 \text{ GeV}/c \ K^- p$
-0.1 ± 2.1	2724	BINGHAM	70B	OSPK $1.8 \text{ GeV}/c \ K^- p$

Ξ^+ MAGNETIC MOMENT

See the Note on Baryon Magnetic Moments in the Λ Listings.

VALUE (μ_N)	EVTS	DOCUMENT ID	TECN	COMMENT
$+0.657 \pm 0.028 \pm 0.020$	70k	HO	90	SPEC $800 \text{ GeV} \ p \ \text{Be}$

Baryon Full Listings



Ξ^- DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \Lambda\pi^-$	(99.887 ± 0.035) %	
$\Gamma_2 \Sigma^- \gamma$	(1.27 ± 0.23) × 10 ⁻⁴	
$\Gamma_3 \Lambda e^- \bar{\nu}_e$	(5.63 ± 0.31) × 10 ⁻⁴	
$\Gamma_4 \Lambda \mu^- \bar{\nu}_\mu$	(3.5 ± _{-2.2} ^{+3.5}) × 10 ⁻⁴	
$\Gamma_5 \Sigma^0 e^- \bar{\nu}_e$	(8.7 ± 1.7) × 10 ⁻⁵	
$\Gamma_6 \Sigma^0 \mu^- \bar{\nu}_\mu$	< 8 × 10 ⁻⁴	90%
$\Gamma_7 \Xi^0 e^- \bar{\nu}_e$	< 2.3 × 10 ⁻³	90%

$\Delta S = 2$ forbidden (S_2) modes

$\Gamma_8 n\pi^-$	$S_2 < 1.9$	× 10 ⁻⁵	90%
$\Gamma_9 n e^- \bar{\nu}_e$	$S_2 < 3.2$	× 10 ⁻³	90%
$\Gamma_{10} n \mu^- \bar{\nu}_\mu$	$S_2 < 1.5$	%	90%
$\Gamma_{11} \rho\pi^- \pi^-$	$S_2 < 4$	× 10 ⁻⁴	90%
$\Gamma_{12} \rho\pi^- e^- \bar{\nu}_e$	$S_2 < 4$	× 10 ⁻⁴	90%
$\Gamma_{13} \rho\pi^- \mu^- \bar{\nu}_\mu$	$S_2 < 4$	× 10 ⁻⁴	90%
$\Gamma_{14} \rho\mu^- \mu^-$	$L < 4$	× 10 ⁻⁴	90%

CONSTRAINED FIT INFORMATION

An overall fit to 4 branching ratios uses 5 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 1.0$ for 1 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $\langle \delta x_i \delta x_j \rangle / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i/\Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_2	-6			
x_3	-8	0		
x_4	-99	0	-1	
x_5	-5	0	0	0
	x_1	x_2	x_3	x_4

Ξ^- BRANCHING RATIOS

A number of early results have been omitted.

$\Gamma(\Sigma^- \gamma)/\Gamma(\Lambda\pi^-)$	Γ_2/Γ_1
VALUE (units 10 ⁻⁴)	
1.27 ± 0.24 OUR FIT	
1.27 ± 0.23 OUR AVERAGE	
1.22 ± 0.23 ± 0.06	211
2.27 ± 1.02	9
	4 DUBBS 94 E761 Ξ^- 375 GeV
	BIAGI 878 SPEC SPS hyperon beam
4 DUBBS 94 also finds weak evidence that the asymmetry parameter α_γ is positive ($\alpha_\gamma = 1.0 \pm 1.3$).	

$\Gamma(\Lambda e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$	Γ_3/Γ_1
VALUE (units 10 ⁻³)	
0.564 ± 0.031 OUR FIT	
0.564 ± 0.031	
0.30 ± 0.13	11
	THOMPSON 80 ASPK Hyperon beam
••• We do not use the following data for averages, fits, limits, etc. •••	
	BOURQUIN 83 SPEC SPS hyperon beam

$\Gamma(\Lambda \mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$	Γ_4/Γ_1
VALUE (units 10 ⁻³)	
0.35 ± 0.35 OUR FIT	
0.35 ± 0.35	
< 2.3	90
< 1.3	0
< 12	
	YEHE 74 HBC Effective denom.=2859
••• We do not use the following data for averages, fits, limits, etc. •••	
	THOMPSON 80 ASPK Effective denom.=1017
	DAUBER 69 HBC
	BERGE 66 HBC

$\Gamma(\Sigma^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$	Γ_5/Γ_1
VALUE (units 10 ⁻³)	
0.087 ± 0.017 OUR FIT	
0.087 ± 0.017	
	154
	BOURQUIN 83 SPEC SPS hyperon beam

$\Gamma(\Sigma^0 \mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$	Γ_6/Γ_1
VALUE (units 10 ⁻³)	
< 0.76	
< 0.76	90
< 5	0
	YEHE 74 HBC Effective denom.=3026
••• We do not use the following data for averages, fits, limits, etc. •••	
	BERGE 66 HBC

$[\Gamma(\Lambda e^- \bar{\nu}_e) + \Gamma(\Sigma^0 e^- \bar{\nu}_e)]/\Gamma(\Lambda\pi^-)$ ($\Gamma_3 + \Gamma_5/\Gamma_1$)

VALUE (units 10 ⁻³)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
0.651 ± 0.031		3011	5 BOURQUIN 83	SPEC	SPS hyperon beam
0.68 ± 0.22		17	6 DUCLOS 71	OSPK	
5 See the separate BOURQUIN 83 values for $\Gamma(\Lambda e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ and $\Gamma(\Sigma^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$ above.					
6 DUCLOS 71 cannot distinguish Σ^0 's from Λ 's. The Cabibbo theory predicts the Σ^0 rate is about a factor 6 smaller than the Λ rate.					

$\Gamma(\Xi^0 e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$	Γ_7/Γ_1
VALUE (units 10 ⁻³)	
< 2.3	
< 2.3	90
	0
	YEHE 74 HBC Effective denom.=1000

$\Gamma(n\pi^-)/\Gamma(\Lambda\pi^-)$	Γ_8/Γ_1
$\Delta S=2$. Forbidden in first-order weak interaction.	
VALUE (units 10 ⁻³)	
< 0.019	
< 0.019	90
	0
	BIAGI 828 SPEC SPS hyperon beam
••• We do not use the following data for averages, fits, limits, etc. •••	
	YEHE 74 HBC Effective denom.=760
	DAUBER 69 HBC
	< 5.0 FERRO-LUZZI 63 HBC

$\Gamma(n e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$	Γ_9/Γ_1
$\Delta S=2$. Forbidden in first-order weak interaction.	
VALUE (units 10 ⁻³)	
< 3.2	
< 3.2	90
	0
	YEHE 74 HBC Effective denom.=715
••• We do not use the following data for averages, fits, limits, etc. •••	
	< 10
	90 BINGHAM 65 RVUE

$\Gamma(n \mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$	Γ_{10}/Γ_1
$\Delta S=2$. Forbidden in first-order weak interaction.	
VALUE (units 10 ⁻³)	
< 15.3	
< 15.3	90
	0
	YEHE 74 HBC Effective denom.=150

$\Gamma(\rho\pi^- \pi^-)/\Gamma(\Lambda\pi^-)$	Γ_{11}/Γ_1
$\Delta S=2$. Forbidden in first-order weak interaction.	
VALUE (units 10 ⁻⁴)	
< 3.7	
< 3.7	90
	0
	YEHE 74 HBC Effective denom.=6200

$\Gamma(\rho\pi^- e^- \bar{\nu}_e)/\Gamma(\Lambda\pi^-)$	Γ_{12}/Γ_1
$\Delta S=2$. Forbidden in first-order weak interaction.	
VALUE (units 10 ⁻⁴)	
< 3.7	
< 3.7	90
	0
	YEHE 74 HBC Effective denom.=6200


$\Gamma(\rho\pi^- \mu^- \bar{\nu}_\mu)/\Gamma(\Lambda\pi^-)$	Γ_{13}/Γ_1
$\Delta S=2$. Forbidden in first-order weak interaction.	
VALUE (units 10 ⁻⁴)	
< 3.7	
< 3.7	90
	0
	YEHE 74 HBC Effective denom.=6200

$\Gamma(\rho\mu^- \mu^-)/\Gamma(\Lambda\pi^-)$	Γ_{14}/Γ_1
$\Delta L=2$ decay, forbidden by total lepton number conservation.	
VALUE (units 10 ⁻⁴)	
< 3.7	
< 3.7	90
	7 LITTENBERG 928 HBC Uses YEH 74 data
7 This LITTENBERG 928 limit and the identical YEH 74 limits for the preceding three modes all result from nonobservance of any 3-prong decays of the Ξ^- . One could as well apply the limit to the <i>sum</i> of the four modes.	

Ξ^- DECAY PARAMETERS

See the Note on Baryon Decay Parameters in the neutron Listings.

$\alpha(\Xi^-)\alpha_-(\Lambda)$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	-0.293 ± 0.007 OUR AVERAGE				Error includes scale factor of 1.8. See the ideogram below.
	-0.303 ± 0.004 ± 0.004	192k	RAMEIKA 86	SPEC	400 GeV pBe
	-0.257 ± 0.020	11k	ASTON 85B	LASS	11 GeV/c $K^- p$
	-0.260 ± 0.017	21k	BENSINGER 85	MPS	5 GeV/c $K^- p$
	-0.299 ± 0.007	150k	BIAGI 82	SPEC	SPS hyperon beam
	-0.315 ± 0.026	9046	CLELAND 80C	ASPK	BNL hyperon beam
	-0.239 ± 0.021	6599	HEMINGWAY 78	HBC	4.2 GeV/c $K^- p$
	-0.243 ± 0.025	4303	BALTAY 74	HBC	1.75 GeV/c $K^- p$
	-0.252 ± 0.032	2436	COOL 74	OSPK	1.8 GeV/c $K^- p$
	-0.253 ± 0.028	2781	DAUBER 69	HBC	



$\alpha(\Xi^-)\alpha_-(\Lambda)$

α FOR $\Xi^- \rightarrow \Lambda\pi^-$

The above average, $\alpha(\Xi^-)\alpha_-(\Lambda) = -0.293 \pm 0.007$, where the error includes a scale factor of 1.8, divided by our current average $\alpha_-(\Lambda) = 0.642 \pm 0.013$, gives the following value for $\alpha(\Xi^-)$.

VALUE	DOCUMENT ID
-0.456 ± 0.014 OUR EVALUATION	Error includes scale factor of 1.8.

ϕ ANGLE FOR $\Xi^- \rightarrow \Lambda\pi^-$ ($\tan\phi = \beta/\gamma$)

VALUE (°)	EVTS	DOCUMENT ID	TECN	COMMENT
4 ± 4 OUR AVERAGE				
5 ± 10	11k	ASTON	85B LASS	K^-p
14.7 ± 16.0	21k	⁸ BENSINGER	85 MPS	$5 \text{ GeV}/c \text{ } K^-p$
11 ± 9	4303	BALTAY	74 HBC	$1.75 \text{ GeV}/c \text{ } K^-p$
5 ± 16	2436	COOL	74 OSPK	$1.8 \text{ GeV}/c \text{ } K^-p$
-26 ± 30	2724	BINGHAM	70B OSPK	
-14 ± 11	2781	DAUBER	69 HBC	Uses $\alpha_\Lambda = 0.647 \pm 0.020$
0 ± 12	1004	⁹ BERGE	66 HBC	
0 ± 20.4	364	⁹ LONDON	66 HBC	Using $\alpha_\Lambda = 0.62$
54 ± 30	356	⁹ CARMONY	64B HBC	

⁸ BENSINGER 85 used $\alpha_\Lambda = 0.642 \pm 0.013$.
⁹ The errors have been multiplied by 1.2 due to approximations used for the Ξ polarization; see DAUBER 69 for a discussion.

g_A / g_V FOR $\Xi^- \rightarrow \Lambda e^- \bar{\nu}_e$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
-0.25 ± 0.05	1992	¹⁰ BOURQUIN	83 SPEC	SPS hyperon beam

¹⁰ BOURQUIN 83 assumes that $g_2 = 0$. Also, the sign has been changed to agree with our conventions, given in the Note on Baryon Decay Parameters in the neutron Listings.

Ξ^- REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1986 edition (Physics Letters **170B**) or in earlier editions.

DUBBS 94 PRL 72 808	+Albuquerque, Bondar+ (FNAL E761 Collab.)
DURYEY 92 PRL 68 768	+Guglielmo, Heller+ (MINN, FNAL, MICH, RUTG)
LITTENBERG 92B PR D46 R992	+Shrock (BNL, STON)
HO 90 PRL 65 1713	+Longo, Nguyen, Luk+ (MICH, FNAL, MINN, RUTG)
Also 91 PR D44 3402	+Ho, Longo, Nguyen, Luk+ (MICH, FNAL, MINN, RUTG)
TROST 89 PR D40 1703	+McCliment, Newsom, Hseuh, Mueller+ (FNAL-715 Collab.)
BIAGI 87B ZPHY C35 143	+ (BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL)
RAMEIKA 86 PR D33 3172	+Beretvas, Deck+ (RUTG, MICH, WISC, MINN)
ASTON 85B PR D32 2270	+Carnegie+ (SLAC, CARL, CNRC, CINC)
BENSINGER 85 NP B252 561	+ (CHIC, ELMT, FNAL, ISU, PNPI, MASD)
BOURQUIN 84 NP B241 1	+ (BRIS, GEVA, HEIDP, LALO, RAL, STRB)
RAMEIKA 84 PRL 52 581	+Beretvas, Deck+ (RUTG, MICH, WISC, MINN)
BOURQUIN 83 ZPHY C21 1	+Brown+ (BRIS, GEVA, HEIDP, LALO, RL, STRB)
BIAGI 82B PL 112B 265	+ (BRIS, CAVE, GEVA, HEIDP, LAUS, LOQM, RL)
BIAGI 82B PL 112B 277	+ (LOQM, GEVA, RL, HEIDP, CAVE, LAUS, BRIS)
CLELAND 80C PR D21 12	+Cooper, Dris, Engels, Herbert+ (PITT, BNL)
THOMPSON 80 PR D21 25	+Cleland, Cooper, Dris, Engels+ (PITT, BNL)
BOURQUIN 79 PL 87B 297	+ (BRIS, GEVA, HEIDP, ORSAY, RHEL, STRB)
HEMINGWAY 78 NP B142 205	+Armenteros+ (CERN, ZEEM, NIJM, OXF)
DIBIANCA 75 NP B98 137	+Endorf (CMU)
BALTAY 74 PR D9 49	+Bridgewater, Cooper, Gershwin+ (COLU, BING, J)
COOL 74 PR D10 792	+Giacomelli, Jenkins, Kycia, Leontic, Li+ (BNL)
Also 72 PRL 29 1630	+Cool, Giacomelli, Jenkins, Kycia, Leontic+ (BNL)
YEH 74 PR D10 3545	+Gaigalas, Smith, Zende, Baltay+ (BING, COLU)
MAYEUR 72 NP B47 333	+VanBinst, Wilquet+ (BRUX, CERN, TUFTS, LOUC)
VOTRUBA 72 NP B47 77	+Safer, Ratcliffe (BIRM, EDIN)
WILQUET 72 PL 42B 372	+Flaigine, Guy+ (BRUX, CERN, TUFTS, LOUC)
DUCLOS 71 NP B32 493	+Freytag, Heintze, Heinzelmann, Jones+ (CERN)
BINGHAM 70B PR D1 3010	+Cook, Humphrey, Sander+ (UCSD, WASH)

GOLDWASSER 70 PR D1 1960	+Schultz (ILL)
STONE 70 PL 32B 515	+Berlingnieri, Bromberg, Cohen, Ferbel+ (RÖCH)
DAUBER 69 PR 179 1262	+Berge, Hubbard, Merrill, Miller (LRL) J
SHEN 67 PL 25B 443	+Firestone, Goldhaber (UCB, LRL)
BERGE 66 PR 147 945	+Eberhard, Hubbard, Merrill+ (LRL)
CHIEN 66 PR 152 1171	+Lach, Sandweiss, Taft, Yeh, Oren+ (YALE, BNL)
LONDON 66 PR 143 1034	+Rau, Goldberg, Lichtman+ (BNL, SYRA)
BINGHAM 65 PRSL 285 202	(CERN)
PJERROU 65B PRL 14 275	+Schlein, Slater, Smith, Stork, Ticho (UCLA)
Also 65 Thesis	Pjerrou (UCLA)
BADIER 64 Dubna Conf. 1 593	+Demoulin, Barloutaud+ (EPOL, SACL, ZEEM)
CARMONY 64B PRL 12 482	+Pjerrou, Schlein, Slater, Stork+ (UCLA) J
HUBBARD 64 PR 135B 183	+Berge, Kalbfleisch, Shafer+ (LRL)
FERRO-LUZZI 63 PR 130 1568	+Alston-Garnjost, Rosenfeld, Wojcicki (LRL)
JAUNEAU 63D Siena Conf. 4	+ (EPOL, CERN, LOUC, RHEL, BERG)
Also 63B PL 5 261	Jauneau+ (EPOL, CERN, LOUC, RHEL, BERG)
SCHNEIDER 63 PR 4 360	(CERN)

OTHER RELATED PAPERS

PONDROM 85 PRPL 122 57	(WISC)
Review of FNAL hyperon experiments.	

NOTE ON Ξ RESONANCES

The accompanying table gives our evaluation of the present status of the Ξ resonances. Not much is known about Ξ resonances. This is because (1) they can only be produced as a part of a final state, and so the analysis is more complicated than if direct formation were possible, (2) the production cross sections are small (typically a few μb), and (3) the final states are topologically complicated and difficult to study with electronic techniques. Thus early information about Ξ resonances came entirely from bubble chamber experiments, where the numbers of events are small, and only in recent years have electronic experiments made significant contributions. However, there is nothing new at all on Ξ resonances since our 1988 edition.

For a detailed earlier review, see Meadows [1].

Reference

1. B.T. Meadows, in *Proceedings of the IVth International Conference on Baryon Resonances* (Toronto, 1980), ed. N. Isgur, p. 283.

Table 1. The status of the Ξ resonances. Only those with an overall status of *** or **** are included in the Baryon Summary Table.

Particle	$L_{2J,2J}$	Overall status	Status as seen in —			
			$\Xi\pi$	ΛK	ΣK	$\Xi(1530)\pi$ Other channels
$\Xi(1318)$	P_{11}	****				Decays weakly
$\Xi(1530)$	P_{13}	****	****			
$\Xi(1620)$		*	*			
$\Xi(1690)$		***		***	**	
$\Xi(1820)$	D_{13}	***	**	***	**	**
$\Xi(1950)$		***	**	**	*	
$\Xi(2030)$	1	***		**	***	
$\Xi(2120)$		*		*		
$\Xi(2250)$		**				3-body decays
$\Xi(2370)$	1	**				3-body decays
$\Xi(2500)$		*		*		3-body decays

**** Existence is certain, and properties are at least fairly well explored.
 *** Existence ranges from very likely to certain, but further confirmation is desirable and/or quantum numbers, branching fractions, etc. are not well determined.
 ** Evidence of existence is only fair.
 * Evidence of existence is poor.

Baryon Full Listings

$\Xi(1530)$

$\Xi(1530) P_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^+) \text{ Status: } ***$$

This is the only Ξ resonance whose properties are all reasonably well known. Spin-parity $3/2^+$ is favored by the data.


We use only those determinations of the mass and width that are accompanied by some discussion of systematics and resolution.

$\Xi(1530)$ MASSES

$\Xi(1530)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1531.80 ± 0.32 OUR FIT		Error includes scale factor of 1.3.		
1531.78 ± 0.34 OUR AVERAGE		Error includes scale factor of 1.4. See the ideogram below.		
1532.2 ± 0.7		DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
1533 ± 1		ROSS 73B	HBC	$K^- p \rightarrow \Xi \bar{K} \pi(\pi)$
1531.4 ± 0.8	59	BADIER 72	HBC	$K^- p$ 3.95 GeV/c
1532.0 ± 0.4	1262	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
1531.3 ± 0.6	324	BORENSTEIN 72	HBC	$K^- p$ 2.2 GeV/c
1532.3 ± 0.7	286	KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
1528.7 ± 1.1	76	LONDON 66	HBC	$K^- p$ 2.24 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1532.1 ± 0.4	1244	ASTON 85B	LASS	$K^- p$ 11 GeV/c
1532.1 ± 0.6	2700	¹ BAUBILLIER 81B	HBC	$K^- p$ 8.25 GeV/c
1530 ± 1	450	BIAGI 81	SPEC	SPS hyperon beam
1527 ± 6	80	SIXEL 79	HBC	$K^- p$ 10 GeV/c
1535 ± 4	100	SIXEL 79	HBC	$K^- p$ 16 GeV/c
1533.6 ± 1.4	97	BERTHON 74	HBC	Quasi-2-body σ

WEIGHTED AVERAGE
1531.78 ± 0.34 (Error scaled by 1.4)



$\Xi(1530)^0$ mass (MeV)

$\Xi(1530)^-$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1535.0 ± 0.6 OUR FIT				
1535.2 ± 0.8 OUR AVERAGE				
1534.5 ± 1.2		DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
1535.3 ± 2.0		ROSS 73B	HBC	$K^- p \rightarrow \Xi \bar{K} \pi(\pi)$
1536.2 ± 1.6	185	KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
1535.7 ± 3.2	38	LONDON 66	HBC	$K^- p$ 2.24 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1540 ± 3	48	BERTHON 74	HBC	Quasi-2-body σ
1534.7 ± 1.1	334	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c

$m_{\Xi(1530)^-} - m_{\Xi(1530)}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
3.2 ± 0.6 OUR FIT			
2.9 ± 0.9 OUR AVERAGE			
2.7 ± 1.0	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
2.0 ± 3.2	MERRILL 66	HBC	$K^- p$ 1.7-2.7 GeV/c
5.7 ± 3.0	PJERROU 65B	HBC	$K^- p$ 1.8-1.95 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •			
3.9 ± 1.8	² KIRSCH 72	HBC	$K^- p$ 2.87 GeV/c
7 ± 4	² LONDON 66	HBC	$K^- p$ 2.24 GeV/c

$\Xi(1530)$ WIDTHS

$\Xi(1530)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
9.1 ± 0.5 OUR AVERAGE				
9.5 ± 1.2		DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
9.1 ± 2.4		ROSS 73B	HBC	$K^- p \rightarrow \Xi \bar{K} \pi(\pi)$
11 ± 2		BADIER 72	HBC	$K^- p$ 3.95 GeV/c
9.0 ± 0.7		BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
8.4 ± 1.4		BORENSTEIN 72	HBC	$\Xi^- \pi^+$
11.0 ± 1.8		KIRSCH 72	HBC	$\Xi^- \pi^+$
7 ± 7		BERGE 66	HBC	$K^- p$ 1.5-1.7 GeV/c
8.5 ± 3.5		LONDON 66	HBC	$K^- p$ 2.24 GeV/c
7 ± 2		SCHLEIN 63B	HBC	$K^- p$ 1.8, 1.95 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •				
12.8 ± 1.0	2700	¹ BAUBILLIER 81B	HBC	$K^- p$ 8.25 GeV/c
19 ± 6	80	³ SIXEL 79	HBC	$K^- p$ 10 GeV/c
14 ± 5	100	³ SIXEL 79	HBC	$K^- p$ 16 GeV/c

$\Xi(1530)^-$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
9.9 ± 1.7 OUR AVERAGE			
9.6 ± 2.8	DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
8.3 ± 3.6	ROSS 73B	HBC	$K^- p \rightarrow \Xi \bar{K} \pi(\pi)$
7.8 ± 3.5	BALTAY 72	HBC	$K^- p$ 1.75 GeV/c
7.8 ± 7.8	KIRSCH 72	HBC	$\Xi^- \pi^0, \Xi^0 \pi^-$
16.2 ± 4.6			

$\Xi(1530)$ POLE POSITIONS

$\Xi(1530)^0$ REAL PART

VALUE	DOCUMENT ID	COMMENT
1531.6 ± 0.4	LICHTENBERG74	Using HABIBI 73

$\Xi(1530)^0$ IMAGINARY PART

VALUE	DOCUMENT ID	COMMENT
4.45 ± 0.35	LICHTENBERG74	Using HABIBI 73

$\Xi(1530)^-$ REAL PART

VALUE	DOCUMENT ID	COMMENT
1534.4 ± 1.1	LICHTENBERG74	Using HABIBI 73

$\Xi(1530)^-$ IMAGINARY PART

VALUE	DOCUMENT ID	COMMENT
3.9 ± 1.75 - 3.9	LICHTENBERG74	Using HABIBI 73

$\Xi(1530)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
$\Gamma_1 \Xi \pi$	100 %	
$\Gamma_2 \Xi \gamma$	< 4 %	90%

$\Xi(1530)$ BRANCHING RATIOS

$\Gamma(\Xi \gamma)/\Gamma_{total}$	CL %	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
< 0.04	90	KALBFLEISCH 75	HBC	$K^- p$ 2.18 GeV/c	

$\Xi(1530)$ FOOTNOTES

- ¹BAUBILLIER 81B is a fit to the inclusive spectrum. The resolution (5 MeV) is not unfolded.
- ²Redundant with data in the mass Listings.
- ³SIXEL 79 doesn't unfold the experimental resolution of 15 MeV.

$\Xi(1530)$ REFERENCES

ASTON 85B	PR D32 2270	+Carnegie+	(SLAC, CARL, CNRC, CINC)
BAUBILLIER 81B	NP B192 1	+ (BIRM, CERN, GLAS, MSU, CURIN)	
BIAGI 81	ZPHY C9 305	+ (BRIS, CAVE, GEVA, HEIDP, LAUS, LOQM, RHEL)	
SIXEL 79	NP B159 125	+Botcher+ (AACH3, BERL, CERN, LOIC, VIEN)	
DEBELLEFON 75B	NC 28A 289	De Bellefon, Berthon, Billoir+	(CDEF, SACL)
KALBFLEISCH 75	PR D11 987	+Strand, Chapman	(BNL, MICH)
BERTHON 74	NC 21A 146	+Tristram+	(CDEF, RHEL, SACL, STRB)
LICHTENBERG 74	PR D10 3865		(IND)
Also	74B Private Comm.	Lichtenberg	(IND)
HABIBI 73	Nevis 199 Thesis		(COLU)
ROSS 73B	Purdue Conf. 355	+Lloyd, Radojicic	(OXF)
BADIER 72	NP B37 429	+Barrelet, Chariton, Videau	(EPOL)
BALTAY 72	PL 42B 129	+Bridgewater, Cooper, Gershwin+	(COLU, BING)
BORENSTEIN 72	PR D5 1559	+Danburg, Kalbfleisch+	(BNL, MICH)
KIRSCH 72	NP B40 349	+Schmidt, Chang+	(BRAN, UMD, SYRA, TUFTS) I
BERGE 66	PR 147 945	+Eberhard, Hubbard, Merrill+	(LRL) I
LONDON 66	PR 143 1034	+Rau, Goldberg, Lichtman+	(BNL, SYRA) IJ
MERRILL 66	UCRL 16455 Thesis		(LRL) JP
PJERROU 65B	PRL 14 275	+Schlein, Slater, Smith, Stork, Ticho	(UCLA)
SCHLEIN 63B	PRL 11 167	+Carmony, Pjerrou, Slater, Stork, Ticho	(UCLA) IJP

See key on page 1343

Baryon Full Listings

$\Xi(1530), \Xi(1620), \Xi(1690)$

OTHER RELATED PAPERS

MAZZUCATO 81	NP B178 1	+Pennino+	(AMST, CERN, NIJM, OXF)
BRIEFEL 77	PR D16 2706	+Gourevitch, Chang+	(BRAN, UMD, SYRA, TUFTS)
BRIEFEL 75	PR D12 1859	+Gourevitch+	(BRAN, UMD, SYRA, TUFTS)
HUNGERBU... 74	PR D10 2051	Hungerbuhler, Majka+	(YALE, FNAL, BNL, PITT)
BUTTON... 66	PR 142 883	Button-Shafer, Lindsey, Murray, Smith	(LRL) JP

 $\Xi(1620)$

$$I(J^P) = \frac{1}{2}(?) \text{ Status: } *$$

J, P need confirmation.

OMITTED FROM SUMMARY TABLE

What little evidence there is consists of weak signals in the $\Xi\pi$ channel. A number of other experiments (e.g., BORENSTEIN 72 and HASSALL 81) have looked for but not seen any effect.

 $\Xi(1620)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1620 OUR ESTIMATE				
1624 ± 3	31	BRIEFEL 77	HBC	$K^- p$ 2.87 GeV/c
1633 ± 12	34	DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
1606 ± 6	29	ROSS 72	HBC	$K^- p$ 3.1–3.7 GeV/c

 $\Xi(1620)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
22.5	31	¹ BRIEFEL 77	HBC	$K^- p$ 2.87 GeV/c
40 ± 15	34	DEBELLEFON 75B	HBC	$K^- p \rightarrow \Xi^- \bar{K} \pi$
21 ± 7	29	ROSS 72	HBC	$K^- p \rightarrow \Xi^- \pi^+ K^{*0}(892)$

 $\Xi(1620)$ DECAY MODES

Mode	Γ_1
$\Xi\pi$	

 $\Xi(1620)$ FOOTNOTES

¹The fit is insensitive to values between 15 and 30 MeV.

 $\Xi(1620)$ REFERENCES

HASSALL 81	NP B189 397	+Ansorge, Carter, Neale+	(CAVE, MSU)
BRIEFEL 77	PR D16 2706	+Gourevitch, Chang+	(BRAN, UMD, SYRA, TUFTS)
Also 70	Duke Conf. 317	Briefel+	(BRAN, UMD, SYRA, TUFTS)
Also 75	PR D12 1859	Briefel, Gourevitch+	(BRAN, UMD, SYRA, TUFTS)
DEBELLEFON 75B	NC 28A 289	De Bellefon, Berthon, Billoir+	(CDEF, SACL)
BORENSTEIN 72	PR D5 1559	+Danburg, Kalbfleisch+	(BNL, MICH) I
ROSS 72	PL 38B 177	+Burau, Lloyd, Mulvey, Radojicic	(OXF) I

OTHER RELATED PAPERS

HUNGERBU... 74	PR D10 2051	Hungerbuhler, Majka+	(YALE, FNAL, BNL, PITT)
SCHMIDT 73	Purdue Conf. 363		(BRAN)
KALBFLEISCH 70	Duke Conf. 331		(BNL) I
APSELL 69	PRL 23 884	+	(BRAN, UMD, SYRA, TUFTS)
BARTSCH 69	PL 28B 439	+	(AACH, BERL, CERN, LOIC, VIEN)

 $\Xi(1690)$

$$I(J^P) = \frac{1}{2}(?) \text{ Status: } ***$$

DIONISI 78 sees a threshold enhancement in both the neutral and negatively charged $\Sigma\bar{K}$ mass spectra in $K^- p \rightarrow (\Sigma\bar{K})K\pi$ at 4.2 GeV/c. The data from the $\Sigma\bar{K}$ channels alone cannot distinguish between a resonance and a large scattering length. Weaker evidence at the same mass is seen in the corresponding $\Lambda\bar{K}$ channels, and a coupled-channel analysis yields results consistent with a new Ξ .

BIAGI 81 sees an enhancement at 1700 MeV in the diffractively produced ΛK^- system. A peak is also observed in the $\Lambda\bar{K}^0$ mass spectrum at 1660 MeV that is consistent with a 1720 MeV resonance decaying to $\Sigma^0\bar{K}^0$, with the γ from the Σ^0 decay not detected.

BIAGI 87 provides further confirmation of this state in diffractive dissociation of Ξ^- into ΛK^- . The significance claimed is 6.7 standard deviations.

 $\Xi(1690)$ MASSES

MIXED CHARGES	VALUE (MeV)	DOCUMENT ID
1690 ± 10 OUR ESTIMATE	This is only an educated guess; the error given is larger than the error on the average of the published values.	

 $\Xi(1690)^0$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1699 ± 5	175	¹ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c
1684 ± 5	183	² DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)^-$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1691.1 ± 1.9 ± 2.0	104	BIAGI 87	SPEC	Ξ^- Be 116 GeV
1700 ± 10	150	³ BIAGI 81	SPEC	Ξ^- H 100, 135 GeV
1694 ± 6	45	⁴ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)$ WIDTHS

MIXED CHARGES

VALUE (MeV)	DOCUMENT ID
< 60 OUR ESTIMATE	

 $\Xi(1690)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
44 ± 23	175	¹ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c
20 ± 4	183	² DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)^-$ WIDTH

VALUE (MeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 8	90	104	BIAGI 87	SPEC	Ξ^- Be 116 GeV
47 ± 14	150		³ BIAGI 81	SPEC	Ξ^- H 100, 135 GeV
26 ± 6	45		⁴ DIONISI 78	HBC	$K^- p$ 4.2 GeV/c

 $\Xi(1690)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Lambda\bar{K}$	seen
Γ_2 $\Sigma\bar{K}$	seen
Γ_3 $\Xi\pi$	
Γ_4 $\Xi^- \pi^+ \pi^0$	
Γ_5 $\Xi^- \pi^+ \pi^-$	possibly seen
Γ_6 $\Xi(1530)\pi$	

 $\Xi(1690)$ BRANCHING RATIOS

$\Gamma(\Lambda\bar{K})/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
seen		104	BIAGI 87	SPEC	-	Ξ^- Be 116 GeV	

$\Gamma(\Sigma\bar{K})/\Gamma(\Lambda\bar{K})$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
2.7 ± 0.9		DIONISI 78	HBC	0	$K^- p$ 4.2 GeV/c	
3.1 ± 1.4		DIONISI 78	HBC	-	$K^- p$ 4.2 GeV/c	

$\Gamma(\Xi\pi)/\Gamma(\Sigma\bar{K})$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_2
< 0.09		DIONISI 78	HBC	0	$K^- p$ 4.2 GeV/c	

$\Gamma(\Xi^- \pi^+ \pi^0)/\Gamma(\Sigma\bar{K})$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_2
< 0.04		DIONISI 78	HBC	0	$K^- p$ 4.2 GeV/c	

$\Gamma(\Xi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ
possibly seen		4	BIAGI 87	SPEC	-	Ξ^- Be 116 GeV	

$\Gamma(\Xi^- \pi^+ \pi^-)/\Gamma(\Sigma\bar{K})$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_5/Γ_2
< 0.03		DIONISI 78	HBC	-	$K^- p$ 4.2 GeV/c	

$\Gamma(\Xi(1530)\pi)/\Gamma(\Sigma\bar{K})$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ_2
< 0.06		DIONISI 78	HBC	-	$K^- p$ 4.2 GeV/c	

 $\Xi(1690)$ FOOTNOTES

- ¹From a fit to the $\Sigma^+ K^-$ spectrum.
- ²From a coupled-channel analysis of the $\Sigma^+ K^-$ and $\Lambda\bar{K}^0$ spectra.
- ³A fit to the inclusive spectrum from $\Xi^- N \rightarrow \Lambda K^- X$.
- ⁴From a coupled-channel analysis of the $\Sigma^0 K^-$ and ΛK^- spectra.

 $\Xi(1690)$ REFERENCES

BIAGI 87	ZPHY C34 15	+	(BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL) I
BIAGI 81	ZPHY C9 305	+	(BRIS, CAVE, GEVA, HEIDP, LAUS, LOQM, RHEL)
DIONISI 78	PL 80B 145	+Diaz, Armenteros+	(CERN, AMST, NIJM, OXF) I

Baryon Full Listings

$\Xi(1820)$

$\Xi(1820) D_{13}$

$$I(J^P) = \frac{1}{2}(\frac{3}{2}^-) \text{ Status: } ***$$

The clearest evidence is an 8-standard-deviation peak in ΛK^- seen by GAY 76. TEODORO 78 favors $J=3/2$, but cannot make a parity discrimination. BIAGI 87C is consistent with $J=3/2$ and favors negative parity for this J value.

$\Xi(1820)$ MASS


We only average the measurements that appear to us to be most significant and best determined.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
1823 ± 5 OUR ESTIMATE					
1823.4 ± 1.4 OUR AVERAGE					
1819.4 ± 3.1 ± 2.0	280	¹ BIAGI	87	SPEC	0 $\Xi^- \text{Be} \rightarrow (\Lambda K^-) X$
1826 ± 3 ± 1	54	BIAGI	87C	SPEC	0 $\Xi^- \text{Be} \rightarrow (\Lambda \bar{K}^0) X$
1822 ± 6		JENKINS	83	MPS	- $K^- p \rightarrow K^+$ (MM)
1830 ± 6	300	BIAGI	81	SPEC	- SPS hyperon beam
1823 ± 2	130	GAY	76C	HBC	- $K^- p$ 4.2 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
1797 ± 19	74	BRIEFEL	77	HBC	0 $K^- p$ 2.87 GeV/c
1829 ± 9	68	BRIEFEL	77	HBC	-0 $\Xi(1530)\pi$
1860 ± 14	39	BRIEFEL	77	HBC	- $\Sigma^- \bar{K}^0$
1870 ± 9	44	BRIEFEL	77	HBC	0 $\Lambda \bar{K}^0$
1813 ± 4	57	BRIEFEL	77	HBC	- ΛK^-
1807 ± 27		DIBIANCA	75	DBC	-0 $\Xi\pi\pi, \Xi^*\pi$
1762 ± 8	28	² BADIÉ	72	HBC	-0 $\Xi\pi, \Xi^*\pi, YK$
1838 ± 5	38	² BADIÉ	72	HBC	-0 $\Xi\pi, \Xi^*\pi, YK$
1830 ± 10	25	³ CRENNELL	70B	DBC	-0 3.6, 3.9 GeV/c
1826 ± 12		⁴ CRENNELL	70B	DBC	-0 3.6, 3.9 GeV/c
1830 ± 10	40	ALITTI	69	HBC	- $\Lambda, \Sigma \bar{K}$
1814 ± 4	30	BADIÉ	65	HBC	0 $\Lambda \bar{K}^0$
1817 ± 7	29	SMITH	65C	HBC	-0 $\Lambda \bar{K}^0, \Lambda K^-$
1770		HALSTEINSLID63	FBC	-0	K^- freon 3.5 GeV/c

$\Xi(1820)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
24 +15 -10 OUR ESTIMATE					
24 ± 6 OUR AVERAGE					Error includes scale factor of 1.5. See the ideogram below.
24.6 ± 5.3	280	¹ BIAGI	87	SPEC	0 $\Xi^- \text{Be} \rightarrow (\Lambda K^-) X$
12 ± 14 ± 1.7	54	BIAGI	87C	SPEC	0 $\Xi^- \text{Be} \rightarrow (\Lambda \bar{K}^0) X$
72 ± 20	300	BIAGI	81	SPEC	- SPS hyperon beam
21 ± 7	130	GAY	76C	HBC	- $K^- p$ 4.2 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •					
99 ± 57	74	BRIEFEL	77	HBC	0 $K^- p$ 2.87 GeV/c
52 ± 34	68	BRIEFEL	77	HBC	-0 $\Xi(1530)\pi$
72 ± 17	39	BRIEFEL	77	HBC	- $\Sigma^- \bar{K}^0$
44 ± 11	44	BRIEFEL	77	HBC	0 $\Lambda \bar{K}^0$
26 ± 11	57	BRIEFEL	77	HBC	- ΛK^-
85 ± 58		DIBIANCA	75	DBC	-0 $\Xi\pi\pi, \Xi^*\pi$
51 ± 13		² BADIÉ	72	HBC	-0 Lower mass
58 ± 13		² BADIÉ	72	HBC	-0 Higher mass
103 +38 -24		³ CRENNELL	70B	DBC	-0 3.6, 3.9 GeV/c
48 +36 -19		⁴ CRENNELL	70B	DBC	-0 3.6, 3.9 GeV/c
55 +40 -20		ALITTI	69	HBC	- $\Lambda, \Sigma \bar{K}$
12 ± 4		BADIÉ	65	HBC	0 $\Lambda \bar{K}^0$
30 ± 7		SMITH	65B	HBC	-0 $\Lambda \bar{K}$
< 80		HALSTEINSLID63	FBC	-0	K^- freon 3.5 GeV/c

WEIGHTED AVERAGE 24±6 (Error scaled by 1.5)



$\Xi(1820)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Lambda \bar{K}$	large
Γ_2 $\Sigma \bar{K}$	small
Γ_3 $\Xi\pi$	small
Γ_4 $\Xi(1530)\pi$	small
Γ_5 $\Xi\pi\pi$ (not $\Xi(1530)\pi$)	

$\Xi(1820)$ BRANCHING RATIOS

The dominant modes seem to be $\Lambda \bar{K}$ and (perhaps) $\Xi(1530)\pi$, but the branching fractions are very poorly determined.

$\Gamma(\Lambda \bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ
0.30 ± 0.15	ALITTI	69	HBC	- $K^- p$ 3.9-5 GeV/c	

$\Gamma(\Xi\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ
0.10 ± 0.10	ALITTI	69	HBC	- $K^- p$ 3.9-5 GeV/c	

$\Gamma(\Xi\pi)/\Gamma(\Lambda \bar{K})$	CL%	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_1
< 0.36	95	GAY	76C	HBC	- $K^- p$ 4.2 GeV/c	
0.20 ± 0.20		BADIÉ	65	HBC	0 $K^- p$ 3 GeV/c	

$\Gamma(\Xi\pi)/\Gamma(\Xi(1530)\pi)$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ_4
1.5 +0.6 -0.4	APSELL	70	HBC	0 $K^- p$ 2.87 GeV/c	

$\Gamma(\Sigma \bar{K})/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ
0.30 ± 0.15	ALITTI	69	HBC	- $K^- p$ 3.9-5 GeV/c	

• • • We do not use the following data for averages, fits, limits, etc. • • •

$\Gamma(\Sigma \bar{K})/\Gamma(\Lambda \bar{K})$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ_1
0.24 ± 0.10	GAY	76C	HBC	- $K^- p$ 4.2 GeV/c	

$\Gamma(\Xi(1530)\pi)/\Gamma_{total}$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ
0.30 ± 0.15	ALITTI	69	HBC	- $K^- p$ 3.9-5 GeV/c	

• • • We do not use the following data for averages, fits, limits, etc. • • •

seen	ASTON	85B	LASS	$K^- p$ 11 GeV/c
not seen	⁵ HASSALL	81	HBC	$K^- p$ 6.5 GeV/c
< 0.25	⁶ DAUBER	69	HBC	$K^- p$ 2.7 GeV/c

$\Gamma(\Xi(1530)\pi)/\Gamma(\Lambda \bar{K})$	DOCUMENT ID	TECN	CHG	COMMENT	Γ_4/Γ_1
0.38 ± 0.27 OUR AVERAGE				Error includes scale factor of 2.3.	
1.0 ± 0.3	GAY	76C	HBC	- $K^- p$ 4.2 GeV/c	
0.26 ± 0.13	SMITH	65C	HBC	-0 $K^- p$ 2.45-2.7 GeV/c	

See key on page 1343

Baryon Full Listings

$\Xi(1820)$, $\Xi(1950)$, $\Xi(2030)$

$\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(A\bar{K})$ Γ_5/Γ_1

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
0.30±0.20	BIAGI	87	SPEC	$\Xi^- \text{Be } 116 \text{ GeV}$
••• We do not use the following data for averages, fits, limits, etc. •••				
<0.14	7 BADIER	65	HBC	0 1 st. dev. limit
>0.1	SMITH	65c	HBC	-0 $K^- p \ 2.45\text{--}2.7 \text{ GeV}/c$

$\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Xi(1530)\pi)$ Γ_5/Γ_4

VALUE	DOCUMENT ID	TECN	CHG	COMMENT
consistent with zero	GAY	76c	HBC	- $K^- p \ 4.2 \text{ GeV}/c$
••• We do not use the following data for averages, fits, limits, etc. •••				
0.3±0.5	8 APSELL	70	HBC	0 $K^- p \ 2.87 \text{ GeV}/c$

$\Xi(1820)$ FOOTNOTES

- BIAGI 87 also sees weak signals in the $\Xi^- \pi^+ \pi^-$ channel at $1782.6 \pm 1.4 \text{ MeV}$ ($\Gamma = 6.0 \pm 1.5 \text{ MeV}$) and $1831.9 \pm 2.8 \text{ MeV}$ ($\Gamma = 9.6 \pm 9.9 \text{ MeV}$).
- BADIER 72 adds all channels and divides the peak into lower and higher mass regions. The data can also be fitted with a single Breit-Wigner of mass 1800 MeV and width 150 MeV.
- From a fit to inclusive $\Xi\pi$, $\Xi\pi\pi$, and $A\bar{K}^-$ spectra.
- From a fit to inclusive $\Xi\pi$ and $\Xi\pi\pi$ spectra only.
- Including $\Xi\pi\pi$.
- DAUBER 69 uses in part the same data as SMITH 65C.
- For the decay mode $\Xi^- \pi^+ \pi^0$ only. This limit includes $\Xi(1530)\pi$.
- Or less. Upper limit for the 3-body decay.

$\Xi(1820)$ REFERENCES

BIAGI 87	ZPHY C34 15	+	(BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL)
BIAGI 87C	ZPHY C34 175	+	(BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL) JP
ASTON 85B	PR D32 2270	+	+Carnegie+ (SLAC, CARL, CNRC, CINC)
JENKINS 83	PRL 51 951	+	+Albright, Diamond+ (FSU, BRAN, LBL, CINC, MASS)
BIAGI 81	ZPHY C9 305	+	(BRIS, CAVE, GEVA, HEIDP, LAUS, LOQM, RHEL)
HASSALL 81	NP B189 397	+	+Ansorge, Carter, Neale+ (CAVE, MSU)
TEODORO 77	PR D16 2706	+	+Diaz, Dionisi, Blokziji+ (AMST, CERN, NIJM, OXF) JP
BRIEFEL 69	PRL 23 884	+	+Gourevitch, Chang+ (BRAN, UMD, SYRA, TUFTS)
Also			
GAY 76	NC 31A 593	+	+Jeanheret, Bogdanski+ (NEUC, LAUS, LIVP, CURIN)
GAY 76C	PL 62B 477	+	+Armenteros, Berge+ (AMST, CERN, NIJM) IJ
DIBIANCA 75	NP 898 137	+	+Endorf (CMU)
BADIER 72	NP B37 429	+	+Barrelet, Chariton, Videau (EPOL)
APSELL 70	PRL 24 777	+	(BRAN, UMD, SYRA, TUFTS) I
CRENNELL 70B	PR D1 847	+	+Karshon, Lai, O'Neill, Scarr, Schumann (BNL)
ALITTI 69	PRL 22 79	+	+Barnes, Flaminio, Metzger+ (BNL, SYRA) I
DAUBER 69	PR 179 1262	+	+Berge, Hubbard, Merrill, Miller (LRL)
TRIPP 67	NP B3 10	+	+Leith+ (LRL, SLAC, CERN, HEID, SACL)
BADIER 65	PL 16 171	+	+Demoulin, Goldberg+ (EPOL, SACL, AMST) I
SMITH 65B	Atkins Conf. 251	+	+Lindsey (LRL)
SMITH 65C	PRL 14 25	+	+Lindsey, Button-Shafer, Murray (LRL) IJP
HALSTEINSLID 63	Siena Conf. 1 73	+	(BERG, CERN, EPOL, RHEL, LOUC) I

OTHER RELATED PAPERS

TEODORO 78	PL 77B 451	+	+Diaz, Dionisi, Blokziji+ (AMST, CERN, NIJM, OXF) JP
BRIEFEL 75	PR D12 1859	+	+Gourevitch+ (BRAN, UMD, SYRA, TUFTS)
SCHMIDT 73	Purdue Conf. 363		(BRAN)
MERRILL 68	PR 167 1202	+	+Shafer (LRL)
SMITH 64	PRL 13 61	+	+Lindsey, Murray, Button-Shafer+ (LRL) IJP

$\Xi(1950)$

$$J(P) = \frac{1}{2}(?)^2 \text{ Status: } ***$$

We list here everything reported between 1875 and 2000 MeV. The accumulated evidence for a Ξ near 1950 MeV seems strong enough to include a $\Xi(1950)$ in the main Baryon Table, but not much can be said about its properties. In fact, there may be more than one Ξ near this mass.

$\Xi(1950)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
1950±15 OUR ESTIMATE				
1944±9	129	BIAGI	87	SPEC $\Xi^- \text{Be} \rightarrow (\Xi^- \pi^+) \pi^- X$
1963±5±2	63	BIAGI	87C	SPEC $\Xi^- \text{Be} \rightarrow (A\bar{K}^0) X$
1937±7	150	BIAGI	81	SPEC SPS hyperon beam
1961±18	139	BRIEFEL	77	HBC $2.87 K^- p \rightarrow \Xi^- \pi^+ X$
1936±22	44	BRIEFEL	77	HBC $2.87 K^- p \rightarrow \Xi^0 \pi^- X$
1964±10	56	BRIEFEL	77	HBC $\Xi(1530)\pi$
1900±12		DIBIANCA	75	DBC $\Xi\pi$
1952±11	25	ROSS	73C	($\Xi\pi$) ⁻
1956±6	29	BADIER	72	HBC $\Xi\pi, \Xi\pi\pi, Y K$
1955±14	21	GOLDWASSER	70	HBC $\Xi\pi$
1894±18	66	DAUBER	69	HBC $\Xi\pi$
1930±20	27	ALITTI	68	HBC $\Xi^- \pi^+$
1933±16	35	BADIER	65	HBC $\Xi^- \pi^+$

$\Xi(1950)$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
60±20 OUR ESTIMATE				
100±31	129	BIAGI	87	SPEC $\Xi^- \text{Be} \rightarrow (\Xi^- \pi^+) \pi^- X$
25±15±1.2	63	BIAGI	87C	SPEC $\Xi^- \text{Be} \rightarrow (A\bar{K}^0) X$
60±8	150	BIAGI	81	SPEC SPS hyperon beam
159±57	139	BRIEFEL	77	HBC $2.87 K^- p \rightarrow \Xi^- \pi^+ X$
87±26	44	BRIEFEL	77	HBC $2.87 K^- p \rightarrow \Xi^0 \pi^- X$
60±39	56	BRIEFEL	77	HBC $\Xi(1530)\pi$
63±78		DIBIANCA	75	DBC $\Xi\pi$
38±10		ROSS	73C	($\Xi\pi$) ⁻
35±11	29	BADIER	72	HBC $\Xi\pi, \Xi\pi\pi, Y K$
56±26	21	GOLDWASSER	70	HBC $\Xi\pi$
98±23	66	DAUBER	69	HBC $\Xi\pi$
80±40	27	ALITTI	68	HBC $\Xi^- \pi^+$
140±35	35	BADIER	65	HBC $\Xi^- \pi^+$

$\Xi(1950)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $A\bar{K}$	seen
Γ_2 $\Sigma\bar{K}$	possibly seen
Γ_3 $\Xi\pi$	seen
Γ_4 $\Xi(1530)\pi$	
Γ_5 $\Xi\pi\pi(\text{not } \Xi(1530)\pi)$	

$\Xi(1950)$ BRANCHING RATIOS

$\Gamma(\Sigma\bar{K})/\Gamma(A\bar{K})$	Γ_2/Γ_1			
VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT
<2.3	90 0	BIAGI	87C	SPEC $\Xi^- \text{Be } 116 \text{ GeV}$
$\Gamma(\Sigma\bar{K})/\Gamma_{\text{total}}$	Γ_2/Γ			
VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
possibly seen	17	HASSALL	81	HBC $K^- p \ 6.5 \text{ GeV}/c$
$\Gamma(\Xi\pi)/\Gamma(\Xi(1530)\pi)$	Γ_3/Γ_4			
VALUE	DOCUMENT ID	TECN		
2.8+0.7 -0.6	APSELL	70	HBC	
$\Gamma(\Xi\pi\pi(\text{not } \Xi(1530)\pi))/\Gamma(\Xi(1530)\pi)$	Γ_5/Γ_4			
VALUE	DOCUMENT ID	TECN		
0.0±0.3	APSELL	70	HBC	

$\Xi(1950)$ REFERENCES

BIAGI 87	ZPHY C34 15	+	(BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL)
BIAGI 87C	ZPHY C34 175	+	(BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL)
BIAGI 81	ZPHY C9 305	+	(BRIS, CAVE, GEVA, HEIDP, LAUS, LOQM, RHEL)
HASSALL 81	NP B189 397	+	+Ansorge, Carter, Neale+ (CAVE, MSU)
BRIEFEL 77	PR D16 2706	+	+Gourevitch, Chang+ (BRAN, UMD, SYRA, TUFTS)
Also			
DIBIANCA 75	NP B98 137	+	+Endorf (CMU)
ROSS 73C	Purdue Conf. 345	+	+Lloyd, Radojicic (OXF)
BADIER 72	NP B37 429	+	+Barrelet, Chariton, Videau (EPOL)
APSELL 70	PRL 24 777	+	(BRAN, UMD, SYRA, TUFTS) I
GOLDWASSER 70	PR D1 1960	+	+Schultz (ILL)
DAUBER 69	PR 179 1262	+	+Berge, Hubbard, Merrill, Miller (LRL) I
ALITTI 68	PRL 21 1119	+	+Flaminio, Metzger, Radojicic+ (BNL, SYRA) I
BADIER 65	PL 16 171	+	+Demoulin, Goldberg+ (EPOL, SACL, AMST) I

$\Xi(2030)$

$$J(P) = \frac{1}{2}(\geq \frac{5}{2})^2 \text{ Status: } ***$$

The evidence for this state has been much improved by HEMINGWAY 77, who see an eight standard deviation enhancement in $\Sigma\bar{K}$ and a weaker coupling to $A\bar{K}$. ALITTI 68 and HEMINGWAY 77 observe no signals in the $\Xi\pi\pi$ (or $\Xi(1530)\pi$) channel, in contrast to DIBIANCA 75. The decay $(A/\Sigma)\bar{K}\pi$ reported by BARTSCH 69 is also not confirmed by HEMINGWAY 77.

A moments analysis of the HEMINGWAY 77 data indicates at a level of three standard deviations that $J \geq 5/2$.


$\Xi(2030)$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
2025 ± 5 OUR ESTIMATE					
2025.1 ± 2.4 OUR AVERAGE					Error includes scale factor of 1.3. See the ideogram below.
2022 ± 7		JENKINS	83	MPS	- $K^- p \rightarrow K^+ \text{MM}$
2024 ± 2	200	HEMINGWAY	77	HBC	- $K^- p \ 4.2 \text{ GeV}/c$
2044 ± 8		DIBIANCA	75	DBC	-0 $\Xi\pi\pi, \Xi^* \pi$
2019 ± 7	15	ROSS	73C	HBC	-0 $\Sigma\bar{K}$
2030 ± 10	42	ALITTI	69	HBC	- $K^- p \ 3.9\text{--}5 \text{ GeV}/c$
2058 ± 17	40	BARTSCH	69	HBC	-0 $K^- p \ 10 \text{ GeV}/c$

Baryon Full Listings

$\Xi(2030), \Xi(2120)$


WEIGHTED AVERAGE
2025.1±2.4 (Error scaled by 1.3)



$\Xi(2030)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
20^{+15}_{-8} OUR ESTIMATE					
21 ± 6 OUR AVERAGE Error includes scale factor of 1.3. See the Ideogram below.					
16 ± 5	200	HEMINGWAY 77	HBC	-	$K^- p$ 4.2 GeV/c
60 ± 24		DIBIANCA 75	DBC	-0	$\Xi \pi \pi, \Xi^* \pi$
33 ± 17	15	ROSS 73c	HBC	-0	$\Sigma \bar{K}$
45 + 40 - 20		ALITTI 69	HBC	-	$K^- p$ 3.9-5 GeV/c
57 ± 30		BARTSCH 69	HBC	-0	$K^- p$ 10 GeV/c

WEIGHTED AVERAGE
21±6 (Error scaled by 1.3)



$\Xi(2030)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda \bar{K}$	~ 20 %
$\Gamma_2 \Sigma \bar{K}$	~ 80 %
$\Gamma_3 \Xi \pi$	small
$\Gamma_4 \Xi(1530)\pi$	small
$\Gamma_5 \Xi \pi \pi$ (not $\Xi(1530)\pi$)	small
$\Gamma_6 \Lambda \bar{K} \pi$	small
$\Gamma_7 \Sigma \bar{K} \pi$	small

$\Xi(2030)$ BRANCHING RATIOS

$\Gamma(\Xi \pi)/[\Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi \pi) + \Gamma(\Xi(1530)\pi)]$		$\Gamma_3/(\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4)$	
VALUE	DOCUMENT ID	TECN	CHG COMMENT
0.25 ± 0.15	ALITTI 69	HBC	- $K^- p$ 3.9-5 GeV/c
• • • We do not use the following data for averages, fits, limits, etc. • • •			
< 0.30	ALITTI 69	HBC	- 1 standard dev. limit
$\Gamma(\Xi \pi)/\Gamma(\Sigma \bar{K})$		Γ_3/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN CHG COMMENT
< 0.19	95	HEMINGWAY 77	HBC - $K^- p$ 4.2 GeV/c

$\Gamma(\Lambda \bar{K})/[\Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi \pi) + \Gamma(\Xi(1530)\pi)]$		$\Gamma_1/(\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4)$	
VALUE	DOCUMENT ID	TECN	CHG COMMENT
0.25 ± 0.15	ALITTI 69	HBC	- $K^- p$ 3.9-5 GeV/c

$\Gamma(\Lambda \bar{K})/\Gamma(\Sigma \bar{K})$		Γ_1/Γ_2	
VALUE	DOCUMENT ID	TECN	CHG COMMENT
0.22 ± 0.09	HEMINGWAY 77	HBC	- $K^- p$ 4.2 GeV/c

$\Gamma(\Sigma \bar{K})/[\Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi \pi) + \Gamma(\Xi(1530)\pi)]$		$\Gamma_2/(\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4)$	
VALUE	DOCUMENT ID	TECN	CHG COMMENT
0.75 ± 0.20	ALITTI 69	HBC	- $K^- p$ 3.9-5 GeV/c

$\Gamma(\Xi(1530)\pi)/[\Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi \pi) + \Gamma(\Xi(1530)\pi)]$		$\Gamma_4/(\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_4)$	
VALUE	DOCUMENT ID	TECN	CHG COMMENT
< 0.15	ALITTI 69	HBC	- 1 standard dev. limit

$[\Gamma(\Xi(1530)\pi) + \Gamma(\Xi \pi \pi \text{ (not } \Xi(1530)\pi))]/\Gamma(\Sigma \bar{K})$		$(\Gamma_4 + \Gamma_5)/\Gamma_2$	
VALUE	CL%	DOCUMENT ID	TECN CHG COMMENT
< 0.11	95	1 HEMINGWAY 77	HBC - $K^- p$ 4.2 GeV/c

$\Gamma(\Lambda \bar{K} \pi)/\Gamma_{\text{total}}$		Γ_6/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BARTSCH 69	HBC	$K^- p$ 10 GeV

$\Gamma(\Lambda \bar{K} \pi)/\Gamma(\Sigma \bar{K})$		Γ_6/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN CHG COMMENT
< 0.32	95	HEMINGWAY 77	HBC - $K^- p$ 4.2 GeV/c

$\Gamma(\Sigma \bar{K} \pi)/\Gamma_{\text{total}}$		Γ_7/Γ	
VALUE	DOCUMENT ID	TECN	COMMENT
seen	BARTSCH 69	HBC	$K^- p$ 10 GeV

$\Gamma(\Sigma \bar{K} \pi)/\Gamma(\Sigma \bar{K})$		Γ_7/Γ_2	
VALUE	CL%	DOCUMENT ID	TECN CHG COMMENT
< 0.04	95	2 HEMINGWAY 77	HBC - $K^- p$ 4.2 GeV/c

$\Xi(2030)$ FOOTNOTES

- For the decay mode $\Xi^- \pi^+ \pi^-$ only.
- For the decay mode $\Sigma^\pm K^- \pi^\mp$ only.

$\Xi(2030)$ REFERENCES

JENKINS 83	PRL 51 951	+Albright, Diamond+ (FSU, BRAN, LBL, CINC, MASS)
HEMINGWAY 77	PL 68B 197	+Armenteros+ (AMST, CERN, NIJM, OXF)1J
	Also 76C PL 62B 477	Gay, Armenteros, Berge+ (AMST, CERN, NIJM)
DIBIANCA 75	NP 89B 137	+Endorf (CMU)
ROSS 73C	Purdue Conf. 345	+Lloyd, Radojicic (OXF)
ALITTI 69	PRL 22 79	+Barnes, Flaminio, Metzger+ (BNL, SYRA)1
BARTSCH 69	PL 28B 439	+ (AACH, BERL, CERN, LOIC, VIEN)
ALITTI 68	PRL 21 1119	+Flaminio, Metzger, Radojicic+ (BNL, SYRA)

$\Xi(2120)$

$I(J^P) = \frac{1}{2}(?)$ Status: *
 J, P need confirmation.

OMITTED FROM SUMMARY TABLE

$\Xi(2120)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
≈ 2120 OUR ESTIMATE				
2137 ± 4	18	1 CHLIAPNIK...	79 HBC	$K^+ p$ 32 GeV/c
2123 ± 7		2 GAY	76C HBC	$K^- p$ 4.2 GeV/c

$\Xi(2120)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
< 20	18	1 CHLIAPNIK...	79 HBC	$K^+ p$ 32 GeV/c
25 ± 12		2 GAY	76C HBC	$K^- p$ 4.2 GeV/c

$\Xi(2120)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda \bar{K}$	seen

See key on page 1343

Baryon Full Listings

$\Xi(2120)$, $\Xi(2250)$, $\Xi(2370)$

 $\Xi(2120)$ BRANCHING RATIOS

$\Gamma(\Lambda\bar{K})/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen		¹ CHLIAPNIK...	79 HBC	$K^+ p \rightarrow (\bar{\Lambda}K^+) X$	
seen		2 GAY	76c HBC	$K^- p$ 4.2 GeV/c	

 $\Xi(2120)$ FOOTNOTES

- ¹ CHLIAPNIKOV 79 does not uniquely identify the K^+ in the $(\bar{\Lambda}K^+) X$ final state. It also reports bumps with fewer events at 2240, 2540, and 2830 MeV.
² GAY 76c sees a 4-standard deviation signal. However, HEMINGWAY 77, with more events from the same experiment points out that the signal is greatly reduced if a cut is made on the 4-momentum u . This suggests an anomalous production mechanism if the $\Xi(2120)$ is real.

 $\Xi(2120)$ REFERENCES

CHLIAPNIK...	79	NP B158 253	Chliapnikov, Gerdjukov+	(CERN, BELG, MONS)
HEMINGWAY	77	PL 688 197	+Armenteros+	(AMST, CERN, NIJM, OXF)
GAY	76c	PL 62B 477	+Armenteros, Berge+	(AMST, CERN, NIJM)

 $\Xi(2250)$
 $I(J^P) = \frac{1}{2}(?)$ Status: **
 J, P need confirmation.

OMITTED FROM SUMMARY TABLE

The evidence for this state is mixed. BARTSCH 69 sees a bump of not much statistical significance in $\Lambda\bar{K}\pi$, $\Sigma\bar{K}\pi$, and $\Xi\pi\pi$ mass spectra. GOLDWASSER 70 sees a narrower bump in $\Xi\pi\pi$ at a higher mass. Not seen by HASSALL 81 with 45 events/ μb at 6.5 GeV/c. Seen by JENKINS 83. Perhaps seen by BIAGI 87.

 $\Xi(2250)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
\approx 2250 OUR ESTIMATE					
2189 \pm 7	66	BIAGI	87	SPEC	$\Xi^- \text{Be} \rightarrow (\Xi^- \pi^+ \pi^-) X$
2214 \pm 5		JENKINS	83	MPS	$K^- p \rightarrow K^+ \text{MM}$
2295 \pm 15	18	GOLDWASSER 70	HBC	-	$K^- p$ 5.5 GeV/c
2244 \pm 52	35	BARTSCH	69	HBC	$K^- p$ 10 GeV/c

 $\Xi(2250)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
46 \pm 27	66	BIAGI	87	SPEC	$\Xi^- \text{Be} \rightarrow (\Xi^- \pi^+ \pi^-) X$
< 30		GOLDWASSER 70	HBC	-	$K^- p$ 5.5 GeV/c
130 \pm 80		BARTSCH	69	HBC	

 $\Xi(2250)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Xi\pi\pi$	seen
Γ_2 $\Lambda\bar{K}\pi$	seen
Γ_3 $\Sigma\bar{K}\pi$	

 $\Xi(2250)$ REFERENCES

BIAGI	87	ZPHY C34 15	+ (BRIS, CERN, GEVA, HEIDP, LAUS, LOQM, RAL)
JENKINS	83	PRL 51 951	+Albright, Diamond+ (FSU, BRAN, LBL, CINC, MASD)
HASSALL	81	NP B189 397	+Ansoerge, Carter, Neale+ (CAVE, MSU)
GOLDWASSER	70	PR D1 1960	+Schultz (ILL)
BARTSCH	69	PL 28B 439	+ (AACH, BERL, CERN, LOIC, VIEN)

 $\Xi(2370)$
 $I(J^P) = \frac{1}{2}(?)$ Status: **
 J, P need confirmation.

OMITTED FROM SUMMARY TABLE

 $\Xi(2370)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
\approx 2370 OUR ESTIMATE					
2356 \pm 10		JENKINS	83	MPS	$K^- p \rightarrow K^+ \text{MM}$
2370	50	HASSALL	81	HBC	$K^- p$ 6.5 GeV/c
2373 \pm 8	94	AMIRZADEH	80	HBC	$K^- p$ 8.25 GeV/c
2392 \pm 27		DIBIANCA	75	DBC	$\Xi 2\pi$

 $\Xi(2370)$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
80	50	HASSALL	81	HBC	$K^- p$ 6.5 GeV/c
80 \pm 25	94	AMIRZADEH	80	HBC	$K^- p$ 8.25 GeV/c
75 \pm 69		DIBIANCA	75	DBC	$\Xi 2\pi$

 $\Xi(2370)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Lambda\bar{K}\pi$ Includes $\Gamma_4 + \Gamma_6$.	seen
Γ_2 $\Sigma\bar{K}\pi$ Includes $\Gamma_5 + \Gamma_6$.	seen
Γ_3 $\Omega^- K$	
Γ_4 $\Lambda\bar{K}^*(892)$	
Γ_5 $\Sigma\bar{K}^*(892)$	
Γ_6 $\Sigma(1385)\bar{K}$	

 $\Xi(2370)$ BRANCHING RATIOS

$\Gamma(\Lambda\bar{K}\pi)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_1/Γ	
seen		AMIRZADEH	80	HBC	-0	$K^- p$ 8.25 GeV/c	

$\Gamma(\Sigma\bar{K}\pi)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_2/Γ	
seen		AMIRZADEH	80	HBC	-0	$K^- p$ 8.25 GeV/c	

$[\Gamma(\Lambda\bar{K}\pi) + \Gamma(\Sigma\bar{K}\pi)]/\Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT	$(\Gamma_1 + \Gamma_2)/\Gamma$
seen		50	HASSALL	81	HBC	-0	$K^- p$ 6.5 GeV/c

$\Gamma(\Omega^- K)/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_3/Γ
0.09 \pm 0.04		¹ KINSON	80	HBC	-	$K^- p$ 8.25 GeV/c

$[\Gamma(\Lambda\bar{K}^*(892)) + \Gamma(\Sigma\bar{K}^*(892))]/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	$(\Gamma_4 + \Gamma_5)/\Gamma$
0.22 \pm 0.13		¹ KINSON	80	HBC	-	$K^- p$ 8.25 GeV/c

$\Gamma(\Sigma(1385)\bar{K})/\Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	CHG	COMMENT	Γ_6/Γ
0.12 \pm 0.08		¹ KINSON	80	HBC	-	$K^- p$ 8.25 GeV/c

 $\Xi(2370)$ FOOTNOTES

- ¹ KINSON 80 is a reanalysis of AMIRZADEH 80 with 50% more events.

 $\Xi(2370)$ REFERENCES

JENKINS	83	PRL 51 951	+Albright, Diamond+ (FSU, BRAN, LBL, CINC, MASD)
HASSALL	81	NP B189 397	+Ansoerge, Carter, Neale+ (CAVE, MSU)
AMIRZADEH	80	PL 90B 324	+ (BIRM, CERN, GLAS, MSU, CURIN) I
KINSON	80	Toronto Conf. 263	+ (BIRM, CERN, GLAS, MSU, CURIN) I
DIBIANCA	75	NP B98 137	+Endorf (CMU)

Baryon Full Listings

 $\Xi(2500), \Omega^-$ $\Xi(2500)$ $I(J^P) = \frac{1}{2}(?)$ Status: *
 J, P need confirmation.

OMITTED FROM SUMMARY TABLE

The ALITTI 69 peak might be instead the $\Xi(2370)$ or might be neither the $\Xi(2370)$ nor the $\Xi(2500)$. $\Xi(2500)$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
≈ 2500 OUR ESTIMATE					
2505 ± 10		JENKINS	83	MPS	$K^- p \rightarrow K^+ \Lambda$
2430 ± 20	30	ALITTI	69	HBC	$K^- p$ 4.6–5 GeV/c
2500 ± 10	45	BARTSCH	69	HBC	$K^- p$ 10 GeV/c

 $\Xi(2500)$ WIDTH

VALUE (MeV)	DOCUMENT ID	TECN	CHG	
150 ⁺⁶⁰ ₋₄₀	ALITTI	69	HBC	–
59 ± 27	BARTSCH	69	HBC	–0

 $\Xi(2500)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Xi \pi$	
Γ_2 $\Lambda \bar{K}$	
Γ_3 $\Sigma \bar{K}$	
Γ_4 $\Xi \pi \pi$	seen
Γ_5 $\Xi(1530) \pi$	
Γ_6 $\Lambda \bar{K} \pi + \Sigma \bar{K} \pi$	seen

 $\Xi(2500)$ BRANCHING RATIOS

$\Gamma(\Xi \pi) / [\Gamma(\Xi \pi) + \Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi(1530) \pi)]$	$\Gamma_1 / (\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_5)$			
VALUE	DOCUMENT ID	TECN	CHG	COMMENT
<0.5	ALITTI	69	HBC	1 standard dev. limit

$\Gamma(\Lambda \bar{K}) / [\Gamma(\Xi \pi) + \Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi(1530) \pi)]$	$\Gamma_2 / (\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_5)$			
VALUE	DOCUMENT ID	TECN	CHG	
0.5 ± 0.2	ALITTI	69	HBC	–

$\Gamma(\Sigma \bar{K}) / [\Gamma(\Xi \pi) + \Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi(1530) \pi)]$	$\Gamma_3 / (\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_5)$			
VALUE	DOCUMENT ID	TECN	CHG	
0.5 ± 0.2	ALITTI	69	HBC	–

$\Gamma(\Xi(1530) \pi) / [\Gamma(\Xi \pi) + \Gamma(\Lambda \bar{K}) + \Gamma(\Sigma \bar{K}) + \Gamma(\Xi(1530) \pi)]$	$\Gamma_5 / (\Gamma_1 + \Gamma_2 + \Gamma_3 + \Gamma_5)$			
VALUE	DOCUMENT ID	TECN	COMMENT	
<0.2	ALITTI	69	HBC	1 standard dev. limit

$\Gamma(\Xi \pi \pi) / \Gamma_{\text{total}}$	Γ_4 / Γ			
VALUE	DOCUMENT ID	TECN	CHG	
seen	BARTSCH	69	HBC	–0

$[\Gamma(\Lambda \bar{K} \pi) + \Gamma(\Sigma \bar{K} \pi)] / \Gamma_{\text{total}}$	Γ_6 / Γ			
VALUE	DOCUMENT ID	TECN	CHG	
seen	BARTSCH	69	HBC	–0

 $\Xi(2500)$ REFERENCES

JENKINS	83	PRL 51 951	+Albright, Diamond+ (FSU, BRAN, LBL, CINC, MASSD)
ALITTI	69	PRL 22 79	+Barnes, Flaminio, Metzger+ (BNL, SYRA)
BARTSCH	69	PL 28B 439	+ (AACH, BERL, CERN, LOIC, VIEN)

 Ω BARYONS
($S = -3, I = 0$) $\Omega^- = sss$ Ω^- $I(J^P) = 0(\frac{3}{2}^+)$ Status: * * * *

The unambiguous discovery in both production and decay was by BARNES 64. The quantum numbers have not actually been measured, but follow from the assignment of the particle to the baryon decuplet. DEUTSCHMANN 78 and BAUBILLIER 78 rule out $J = 1/2$ and find consistency with $J = 3/2$.

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1986 edition (Physics Letters 170B) or in earlier editions.

 Ω^- MASSThe fit assumes the Ω^- and $\bar{\Omega}^+$ masses are the same.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1672.45 ± 0.29 OUR FIT				
1672.43 ± 0.32 OUR AVERAGE				
1673 ± 1	100	HARTOUNI	85	SPEC 80–280 GeV $K_L^0 C$
1673.0 ± 0.8	41	BAUBILLIER	78	HBC 8.25 GeV/c $K^- p$
1671.7 ± 0.6	27	HEMINGWAY	78	HBC 4.2 GeV/c $K^- p$
1673.4 ± 1.7	4	¹ DIBIANCA	75	DBC 4.9 GeV/c $K^- d$
1673.3 ± 1.0	3	PALMER	68	HBC $K^- p$ 4.6, 5 GeV/c
1671.8 ± 0.8	3	SCHULTZ	68	HBC $K^- p$ 5.5 GeV/c
1674.2 ± 1.6	5	SCOTTER	68	HBC $K^- p$ 6 GeV/c
1672.1 ± 1.0	1	² FRY	55	EMUL
• • • We do not use the following data for averages, fits, limits, etc. • • •				
1671.43 ± 0.78	13	³ DEUTSCH...	73	HBC $K^- p$ 10 GeV/c
1671.9 ± 1.2	6	³ SPETH	69	HBC See DEUTSCHMANN 73
1673.0 ± 0.8	1	ABRAMS	64	HBC $\rightarrow \Xi^- \pi^0$
1670.6 ± 1.0	1	² FRY	55B	EMUL
1615	1	⁴ EISENBERG	54	EMUL

¹ DIBIANCA 75 gives a mass for each event. We quote the average.

² The FRY 55 and FRY 55B events were identified as Ω^- by ALVAREZ 73. The masses assume decay to ΛK^- at rest. For FRY 55B, decay from an atomic orbit could Doppler shift the K^- energy and the resulting Ω^- mass by several MeV. This shift is negligible for FRY 55 because the Ω decay is approximately perpendicular to its orbital velocity, as is known because the Λ strikes the nucleus (L. Alvarez, private communication 1973). We have calculated the error assuming that the orbital n is 4 or larger.

³ Excluded from the average; the Ω^- lifetimes measured by the experiments differ significantly from other measurements.

⁴ The EISENBERG 54 mass was calculated for decay in flight. ALVAREZ 73 has shown that the Ω interacted with an Ag nucleus to give $K^- \Xi$ Ag.

 $\bar{\Omega}^+$ MASSThe fit assumes the Ω^- and $\bar{\Omega}^+$ masses are the same.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
1672.45 ± 0.29 OUR FIT				
1672.5 ± 0.7 OUR AVERAGE				
1672 ± 1	72	HARTOUNI	85	SPEC 80–280 GeV $K_L^0 C$
1673.1 ± 1.0	1	FIRESTONE	71B	HBC 12 GeV/c $K^+ d$

 $(m_{\Omega^-} - m_{\bar{\Omega}^+}) / m_{\text{average}}$

A test of CPT invariance. Calculated from the average Ω^- and $\bar{\Omega}^+$ masses, above.

VALUE DOCUMENT ID
(0 ± 5) × 10⁻⁴ OUR EVALUATION

 Ω^- MEAN LIFEMeasurements with an error > 0.1 × 10⁻¹⁰ s have been omitted.

VALUE (10 ⁻¹⁰ s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.822 ± 0.012 OUR AVERAGE				
0.811 ± 0.037	1096	LUK	88	SPEC p Be 400 GeV
0.823 ± 0.013	12k	BOURQUIN	84	SPEC SPS hyperon beam
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.822 ± 0.028	2437	BOURQUIN	79B	SPEC See BOURQUIN 84

See key on page 1343

Baryon Full Listings

 $\Omega^-, \Omega(2250)^-$ Ω^- MAGNETIC MOMENT

VALUE (μ_N)	EVTs	DOCUMENT ID	TECN	COMMENT
$-1.94 \pm 0.17 \pm 0.14$	25k	DIEHL	91	SPEC Spin-transfer production

 Ω^- DECAY MODES

Mode	Fraction (Γ_i/Γ)	Confidence level
Γ_1 ΛK^-	(67.8 ± 0.7) %	
Γ_2 $\Xi^0 \pi^-$	(23.6 ± 0.7) %	
Γ_3 $\Xi^- \pi^0$	(8.6 ± 0.4) %	
Γ_4 $\Xi^- \pi^+ \pi^-$	$(4.3^{+3.4}_{-1.3}) \times 10^{-4}$	
Γ_5 $\Xi(1530)^0 \pi^-$	$(6.4^{+5.1}_{-2.0}) \times 10^{-4}$	
Γ_6 $\Xi^0 e^- \bar{\nu}_e$	$(5.6 \pm 2.8) \times 10^{-3}$	
Γ_7 $\Xi^- \gamma$	$< 2.2 \times 10^{-3}$	90%
$\Delta S = 2$ forbidden (S_2) modes		
Γ_8 $\Lambda \pi^-$	$S_2 < 1.9 \times 10^{-4}$	90%

 Ω^- BRANCHING RATIOS

The BOURQUIN 84 values (which include results of BOURQUIN 79B, a separate experiment) are much more accurate than any other results, and so the other results have been omitted.

$\Gamma(\Lambda K^-)/\Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
	0.678 ± 0.007	14k	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	0.686 ± 0.013	1920	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^0 \pi^-)/\Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ
	0.236 ± 0.007	1947	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	0.234 ± 0.013	317	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^- \pi^0)/\Gamma_{\text{total}}$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ
	0.086 ± 0.004	759	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	0.080 ± 0.008	145	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^- \pi^+ \pi^-)/\Gamma_{\text{total}}$	VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
	$4.3^{+3.4}_{-1.3}$	4	BOURQUIN 84	SPEC	SPS hyperon beam	

$\Gamma(\Xi(1530)^0 \pi^-)/\Gamma_{\text{total}}$	VALUE (units 10^{-4})	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_5/Γ
	$6.4^{+5.1}_{-2.0}$	4	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	~ 20	1	BOURQUIN 79B	SPEC	See BOURQUIN 84	
5 The same 4 events as in the previous mode, with the isospin factor to take into account $\Xi(1530)^0 \rightarrow \Xi^0 \pi^0$ decays included.						

$\Gamma(\Xi^0 e^- \bar{\nu}_e)/\Gamma_{\text{total}}$	VALUE (units 10^{-3})	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_6/Γ
	5.6 ± 2.8	14	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •						
	~ 10	3	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Xi^- \gamma)/\Gamma_{\text{total}}$	VALUE (units 10^{-3})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_7/Γ
	< 2.2	90	9	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
	< 3.1	90	0	BOURQUIN 79B	SPEC	See BOURQUIN 84	

$\Gamma(\Lambda \pi^-)/\Gamma_{\text{total}}$	VALUE (units 10^{-4})	CL%	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_8/Γ
	< 1.9	90	0	BOURQUIN 84	SPEC	SPS hyperon beam	
• • • We do not use the following data for averages, fits, limits, etc. • • •							
	< 13	90	0	BOURQUIN 79B	SPEC	See BOURQUIN 84	

 Ω^- DECAY PARAMETERS α FOR $\Omega^- \rightarrow \Lambda K^-$

Some early results have been omitted.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
-0.026 ± 0.026 OUR AVERAGE				
-0.034 ± 0.079	1743	LUK	88	SPEC p Be 400 GeV
-0.025 ± 0.028	12k	BOURQUIN	84	SPEC SPS hyperon beam

 α FOR $\Omega^- \rightarrow \Xi^0 \pi^-$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$+0.09 \pm 0.14$	1630	BOURQUIN	84	SPEC SPS hyperon beam

 α FOR $\Omega^- \rightarrow \Xi^- \pi^0$

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$+0.05 \pm 0.21$	614	BOURQUIN	84	SPEC SPS hyperon beam

 Ω^- REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1986 edition (Physics Letters **170B**) or in earlier editions.

DIEHL 91	PRL 67 804	+Teige, Thompson, Zou+	(RUTG, FNAL, MICH, MINN)
LUK 88	PR D38 19	+Beretvas, Deck+	(RUTG, WISC, MICH, MINN)
HARTOUNI 85	PRL 54 628	+Atiya, Holmes, Knapp, Lee+	(COLU, ILL, FNAL)
BOURQUIN 84	NP B241 1	+ Bourquin+	(BRIS, GEVA, HEIDP, LALO, RAL, STRB)
Also 79	PL 87B 297	+ Bourquin+	(BRIS, GEVA, HEIDP, LALO, RAL)
BOURQUIN 79B	PL 88B 192	+ Bourquin+	(BRIS, GEVA, HEIDP, LALO, RAL)
BAUBILLIER 78	PL 78B 342	+ Bourquin+	(BIRM, CERN, GLAS, MSU, CURIN, PARIN) J
DEUTSCH... 78	PL 73B 96	+ Deutschmann+	(AACHS, BERL, CERN, INNS, LOIC+) J
HEMINGWAY 78	NP B142 205	+ Armenteros+	(CERN, ZEEM, NIJM, OXF)
DIBIANCA 75	NP B98 137	+ Endorf	(CMU)
ALVAREZ 73	PRL D8 702	+ Deutschmann, Kaufmann, Besliv+	(LBL)
DEUTSCH... 73	NP B61 102	+ Goldhaber, Lissauer, Sheldon, Trilling	(LRL)
FIRESTONE 71B	PRL 26 410	+ (AACH, BERL, CERN, LOIC, VIEN)	
SPETH 69	PL 29B 252	+ Radojicic, Rau, Richardson+	(BNL, SYRA)
PALMER 68	PL 26B 323	+ (ILL, ANL, NWES, WISC)	
SCHULTZ 68	PR 168 1509	+ Burnstein, Glasser+	(UMD, NRL)
SCOTTER 68	PL 26B 474	+ Connolly, Crennell, Culwick+	(BNL)
ABRAMS 64	PRL 13 670	+ Schneps, Swami	(WISC)
BARNES 64	PRL 12 204	+ Schneps, Swami	(WISC)
FRY 55	PR 97 1189	+ Schneps, Swami	(WISC)
FRY 55B	NC 2 346	+ Schneps, Swami	(WISC)
EISENBERG 54	PR 96 541		(CORN)

 $\Omega(2250)^-$ $I(J^P) = 0(?)^?$ Status: * * * $\Omega(2250)^-$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2252 ± 9 OUR AVERAGE				
2253 ± 13	44	ASTON	87B	LASS $K^- p$ 11 GeV/c
$2251 \pm 9 \pm 8$	78	BIAGI	86B	SPEC SPS Ξ^- beam

 $\Omega(2250)^-$ WIDTH

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
55 ± 18 OUR AVERAGE				
81 ± 38	44	ASTON	87B	LASS $K^- p$ 11 GeV/c
48 ± 20	78	BIAGI	86B	SPEC SPS Ξ^- beam

 $\Omega(2250)^-$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Xi^- \pi^+ K^-$	seen
Γ_2 $\Xi(1530)^0 K^-$	seen

 $\Omega(2250)^-$ BRANCHING RATIOS

$\Gamma(\Xi(1530)^0 K^-)/\Gamma(\Xi^- \pi^+ K^-)$	VALUE	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_1
	~ 1.0	44	ASTON	87B	LASS $K^- p$ 11 GeV/c	
	0.70 ± 0.20	49	BIAGI	86B	SPEC Ξ^- Be 116 GeV/c	

 $\Omega(2250)^-$ REFERENCES

ASTON 87B	PL B194 579	+Awaji, Bienz, Bird+	(SLAC, NAGO, CINC, INUS)
BIAGI 86B	ZPHY C31 33	+ (LOQM, GEVA, RAL, HEIDP, LAUS, BRIS, CERN)	

Baryon Full Listings

$\Omega(2380)^-$, $\Omega(2470)^-$, Charmed Baryons

$\Omega(2380)^-$	Status: **				
OMITTED FROM SUMMARY TABLE					
$\Omega(2380)^-$ MASS					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
≈ 2380 OUR ESTIMATE					
$2384 \pm 9 \pm 8$	45	BIAGI	86B SPEC	SPS Ξ^- beam	
$\Omega(2380)^-$ WIDTH					
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT	
26 ± 23	45	BIAGI	86B SPEC	SPS Ξ^- beam	
$\Omega(2380)^-$ DECAY MODES					
Mode					
Γ_1	$\Xi^- \pi^+ K^-$				
Γ_2	$\Xi(1530)^0 K^-$				
Γ_3	$\Xi^- \bar{K}^*(892)^0$				
$\Omega(2380)^-$ BRANCHING RATIOS					
$\Gamma(\Xi(1530)^0 K^-) / \Gamma(\Xi^- \pi^+ K^-)$				Γ_2 / Γ_1	
VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.44	90	9	BIAGI	86B SPEC	Ξ^- Be 116 GeV/c
$\Gamma(\Xi^- \bar{K}^*(892)^0) / \Gamma(\Xi^- \pi^+ K^-)$				Γ_3 / Γ_1	
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.5 ± 0.3	21	BIAGI	86B SPEC	Ξ^- Be 116 GeV/c	
$\Omega(2380)^-$ REFERENCES					
BIAGI	86B	ZPHY C31 33	+	(LOQM, GEVA, RAL, HEIDP, LAUS, BRIS, CERN)	

$\Omega(2470)^-$	Status: **			
OMITTED FROM SUMMARY TABLE				
<p>A peak in the $\Omega^- \pi^+ \pi^-$ mass spectrum with a signal significance claimed to be at least 5.5 standard deviations. There is no reason to seriously doubt the existence of this state, but unless the evidence is overwhelming we usually wait for confirmation from a second experiment before elevating peaks to the Summary Table.</p>				
$\Omega(2470)^-$ MASS				
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2474 ± 12	59	ASTON	88G LASS	$K^- p$ 11 GeV/c
$\Omega(2470)^-$ WIDTH				
VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
72 ± 33	59	ASTON	88G LASS	$K^- p$ 11 GeV/c
$\Omega(2470)^-$ DECAY MODES				
Mode				
Γ_1	$\Omega^- \pi^+ \pi^-$			
$\Omega(2470)^-$ REFERENCES				
ASTON	88G	PL B215 799	+	Awaji, Bienz, Bird+ (SLAC, NAGO, CINC, INUS)

CHARMED BARYONS (C = +1)

$$\Lambda_c^+ = udc, \quad \Sigma_c^{++} = uuc, \quad \Sigma_c^+ = udc, \quad \Sigma_c^0 = ddc,$$

$$\Xi_c^+ = usc, \quad \Xi_c^0 = dsc, \quad \Omega_c^0 = ssc$$

NOTE ON CHARMED BARYONS

Figs. 1(a) and 1(b) show the SU(4) multiplets that have as their "ground floors" (a) the SU(3) octet that contains the nucleon, and (b) the SU(3) decuplet that contains the $\Delta(1232)$. All the particles in a given SU(4) multiplet have the same spin and parity. The only charmed baryons that have been discovered each contain one charmed quark and belong to the first floor of the multiplet shown in Fig. 1(a). Fig. 2 shows this first floor, pulled apart into two SU(3) multiplets, a $\bar{3}$ that contains the $\Lambda_c(2285)$ and the $\Xi_c(2470)$, both of which decay weakly, and a 6 that contains the $\Sigma_c(2455)$, which decays strongly to $\Lambda_c \pi$, and the $\Omega_c(2710)$, which decays weakly (and which needs confirmation before it can be considered to be established). A second Ξ_c remains to be discovered to fill out the 6 , and a host of other baryons with one or more charmed quarks are needed to fill out the full SU(4) multiplets in Fig. 1. Furthermore, every N or Δ baryon resonance "starts" a multiplet like that in Fig. 1(a) or 1(b), so the woods are full of charmed baryons, most of which no doubt will forever remain undiscovered.




Fig. 1. SU(4) multiplets of baryons made of u , d , s , and c quarks. (a) The 20-plet with an SU(3) octet on the "ground floor." (b) The 20-plet with an SU(3) decuplet on the ground floor.




Fig. 2. The SU(3) multiplets on the "first floor" of the SU(4) multiplet of Fig. 1(a). The particles in dashed circles have yet to be discovered.

See key on page 1343

Baryon Full Listings

Charmed Baryons, Λ_c^+

The states of the $\bar{3}$ multiplet are antisymmetric under interchange of the two light quarks (the u , d , and s quarks), and the states of the 6 multiplet are symmetric under interchange of these quarks. Actually, there may be some mixing between the pure $\bar{3}$ and 6 Ξ_c states (they have the same I, J , and P quantum numbers) to form the physical Ξ_c states.

It need hardly be said that the flavor symmetries Fig. 1 displays are very badly broken, but the figure is the simplest way to see what charmed baryons should exist.

For an entry into the literature on models of charmed baryons, see Ref. 1. For a review of recent experimental results, see Ref. 2. For a review of both theory and experiment, see Ref. 3.

References

1. K. Maltman and N. Isgur, Phys. Rev. **D22**, 1701 (1980); S. Capstick and N. Isgur, Phys. Rev. **D34**, 2809 (1986); W. Kwong, J.L. Rosner, and C. Quigg, Ann. Rev. Nucl. and Part. Sci. **37**, 325 (1987); S. Fleck and J.M. Richard, Part. World **1**, 67 (1990).
2. S.R. Klein, Int. J. Mod. Phys. **A5**, 1457 (1990).
3. J.G. Körner and H.W. Siebert, Ann. Rev. Nucl. and Part. Sci. **41**, 511 (1991).

Λ_c^+ $(J^P) = 0(\frac{1}{2}^+)$ Status: * * * *

J has not actually been measured yet. Results of an analysis of $\rho K^- \pi^+$ decays (JEZABEK 92) are consistent with the expected $J = 1/2$. The quark content is udc .

We have omitted some results that have been superseded by later experiments. The omitted results may be found in our 1992 edition (Physical Review **D45**, 1 June, Part II) or in earlier editions.

Λ_c^+ MASS

Measurements with an error greater than 5 MeV or that are otherwise obsolete have been omitted.

The fit also uses $(m_{\Sigma_c} - m_{\Lambda_c^+})$ measurements.

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2285.1 ± 0.6 OUR FIT				
2284.9 ± 0.6 OUR AVERAGE				
2284.7 ± 0.6 ± 0.7	1134	AVERY	91 CLEO	Six modes
2281.7 ± 2.7 ± 2.6	29	ALVAREZ	90B NA14	$\rho K^- \pi^+$
2285.8 ± 0.6 ± 1.2	101	BARLAG	89 NA32	$\rho K^- \pi^+$
2284.7 ± 2.3 ± 0.5	5	AGUILAR...	88B LEBE	$\rho K^- \pi^+$
2283.1 ± 1.7 ± 2.0	628	ALBRECHT	88C ARG	$\rho K^- \pi^+$, $\rho \bar{K}^0$, $\Lambda_{3\pi}$
2286.2 ± 1.7 ± 0.7	97	ANJOS	88B E691	$\rho K^- \pi^+$
2281 ± 3	2	JONES	87 HBC	$\rho K^- \pi^+$
2283 ± 3	3	BOSETTI	82 HBC	$\rho K^- \pi^+$
2290 ± 3	1	CALICCHIO	80 HYBR	$\rho K^- \pi^+$

Λ_c^+ MEAN LIFE

Measurements with an error $\geq 0.1 \times 10^{-12}$ s have been omitted.

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
0.200^{+0.011}_{-0.010} OUR AVERAGE				
0.215 ± 0.016 ± 0.008	1340	FRABETTI	93D E687	γ Be, $\Lambda_c^+ \rightarrow \rho K^- \pi^+$
0.18 ± 0.03 ± 0.03	29	ALVAREZ	90 NA14	γ , $\Lambda_c^+ \rightarrow \rho K^- \pi^+$
0.20 ± 0.03 ± 0.03	90	FRABETTI	90 E687	γ Be, $\Lambda_c^+ \rightarrow \rho K^- \pi^+$
0.196 ^{+0.023} _{-0.020}	101	BARLAG	89 NA32	$\rho K^- \pi^+$ + c.c.
0.12 ^{+0.05} _{-0.03}	9	AGUILAR...	88B LEBE	
0.22 ± 0.03 ± 0.02	97	ANJOS	88B E691	$\rho K^- \pi^+$ + c.c.
0.23 ^{+0.09} _{-0.06} ± 0.04	11	ADAMOVIICH	87 EMUL	γ A 20-70 GeV/c
0.11 ^{+0.08} _{-0.04}	9	AMENDOLIA	87 SPEC	γ Ge-Si, $\rho K^- \pi^+ \pi^0$
0.20 ^{+0.07} _{-0.05}	13	USHIDA	86 EMUL	
0.14 ^{+0.05} _{-0.03} ± 0.03	14	BARLAG	87 NA32	See BARLAG 89

• • • We do not use the following data for averages, fits, limits, etc. • • •

Λ_c^+ DECAY MODES

Mode	Fraction (Γ_i/Γ)	Scale factor/ Confidence level
Hadronic modes with a ρ and one K		
Γ_1 $\rho \bar{K}^0$	(2.1 ± 0.4) %	
Γ_2 $\rho K^- \pi^+$	(4.4 ± 0.6) %	
Γ_3 $\rho \bar{K}^*(892)^0$	[a] (1.6 ± 0.4) %	
Γ_4 $\Delta(1232)^{++} K^-$	(7 ± 4) × 10 ⁻³	
Γ_5 $\Lambda(1520) \pi^+$	[a] (3.9 ^{+2.0} _{-1.7}) × 10 ⁻³	
Γ_6 $\rho K^- \pi^+$ nonresonant	(2.4 ^{+0.5} _{-0.6}) %	
Γ_7 $\rho \bar{K}^0 \pi^+ \pi^-$	(2.4 ± 0.8) %	S=1.3
Γ_8 $\rho K^- \pi^+ \pi^0$	seen	
Γ_9 $\rho K^*(892)^- \pi^+$	seen	
Γ_{10} $\rho(K^- \pi^+)$ nonresonant π^0	(3.2 ± 0.7) %	
Γ_{11} $\Delta(1232) \bar{K}^*(892)$	seen	
Γ_{12} $\rho K^- \pi^+ \pi^+ \pi^-$	(10 ± 7) × 10 ⁻⁴	
Γ_{13} $\rho K^- \pi^+ \pi^0 \pi^0$	(7.0 ± 3.5) × 10 ⁻³	
Γ_{14} $\rho K^- \pi^+ \pi^0 \pi^0 \pi^0$	(4.4 ± 2.8) × 10 ⁻³	
Hadronic modes with a ρ and zero or two K's		
Γ_{15} $\rho \pi^+ \pi^-$	(3.0 ± 1.6) × 10 ⁻³	
Γ_{16} $\rho f_0(980)$	[a] (2.4 ± 1.6) × 10 ⁻³	
Γ_{17} $\rho \pi^+ \pi^+ \pi^- \pi^-$	(1.6 ± 1.0) × 10 ⁻³	
Γ_{18} $\rho K^+ K^-$	(3.0 ± 1.1) × 10 ⁻³	
Γ_{19} $\rho \phi$	[a] < 1.7 × 10 ⁻³	CL=90%
Hadronic modes with a hyperon		
Γ_{20} $\Lambda \pi^+$	(7.9 ± 1.8) × 10 ⁻³	
Γ_{21} $\Lambda \pi^+ \pi^0$	(3.2 ± 0.9) %	
Γ_{22} $\Lambda \rho^0$	< 4 %	CL=95%
Γ_{23} $\Lambda \pi^+ \pi^+ \pi^-$	(2.7 ± 0.6) %	
Γ_{24} $\Sigma^0 \pi^+$	(8.7 ± 2.0) × 10 ⁻³	
Γ_{25} $\Sigma^0 \pi^+ \pi^0$	(1.6 ± 0.6) %	
Γ_{26} $\Sigma^0 \pi^+ \pi^+ \pi^-$	(9.2 ± 3.3) × 10 ⁻³	
Γ_{27} $\Sigma^+ \pi^0$	(8.7 ± 2.2) × 10 ⁻³	
Γ_{28} $\Sigma^+ \pi^+ \pi^-$	(3.0 ± 0.6) %	
Γ_{29} $\Sigma^+ \rho^0$	< 1.2 %	CL=95%
Γ_{30} $\Sigma^- \pi^+ \pi^+$	(1.6 ± 0.6) %	
Γ_{31} $\Sigma^+ \pi^+ \pi^- \pi^0$		
Γ_{32} $\Sigma^+ \omega$	[a] (2.4 ± 0.7) %	
Γ_{33} $\Sigma^+ \pi^+ \pi^+ \pi^- \pi^-$	(2.6 ^{+3.5} _{-1.8}) × 10 ⁻³	
Γ_{34} $\Sigma^+ K^+ K^-$	(3.1 ± 0.8) × 10 ⁻³	
Γ_{35} $\Sigma^+ \phi$	[a] (3.0 ± 1.3) × 10 ⁻³	
Γ_{36} $\Sigma^+ K^+ \pi^-$	(5.7 ^{+5.3} _{-3.1}) × 10 ⁻³	
Γ_{37} $\Xi^0 K^+$	(3.4 ± 0.9) × 10 ⁻³	
Γ_{38} $\Xi^- K^+ \pi^+$	(3.8 ± 1.2) × 10 ⁻³	
Γ_{39} $\Xi(1530)^0 K^+$	[a] (2.3 ± 0.9) × 10 ⁻³	
Semileptonic modes		
Γ_{40} $\Lambda \ell^+ \nu_\ell$		
Γ_{41} Λ anything $\ell^+ \nu_\ell$		
Inclusive modes		
Γ_{42} p anything	(50 ± 16) %	
Γ_{43} p anything (no Λ)	(12 ± 19) %	
Γ_{44} p hadrons		
Γ_{45} n anything	(50 ± 16) %	
Γ_{46} n anything (no Λ)	(29 ± 17) %	
Γ_{47} Λ anything	(35 ± 11) %	S=1.4
Γ_{48} Σ^\pm anything	[b] (10 ± 5) %	
Γ_{49} e^+ anything	(4.5 ± 1.7) %	
Γ_{50} ρe^+ anything	(1.8 ± 0.9) %	
Γ_{51} Λe^+ anything	(1.4 ± 0.5) %	
Γ_{52} $\Lambda \mu^+$ anything	(1.5 ± 0.9) %	
Γ_{53} dummy mode used by the fit	(89.1 ± 1.7) %	

[a] The branching fraction includes all the decay modes of the final-state resonance.

[b] The value is for the sum of the charge states indicated.

Baryon Full Listings

 Λ_c^+

CONSTRAINED FIT INFORMATION

An overall fit to 7 branching ratios uses 14 measurements and one constraint to determine 5 parameters. The overall fit has a $\chi^2 = 12.8$ for 10 degrees of freedom.

The following *off-diagonal* array elements are the correlation coefficients $(\delta x_i \delta x_j) / (\delta x_i \delta x_j)$, in percent, from the fit to the branching fractions, $x_i \equiv \Gamma_i / \Gamma_{\text{total}}$. The fit constrains the x_i whose labels appear in this array to sum to one.

x_7	36			
x_{23}	61	22		
x_{51}	25	9	15	
x_{53}	-81	-70	-70	-48
	x_2	x_7	x_{23}	x_{51}

 Λ_c^+ BRANCHING RATIOS

$\Gamma(\rho\bar{K}^0)/\Gamma(\rho K^- \pi^+)$					Γ_1/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.49 ± 0.07 OUR AVERAGE					
0.44 ± 0.07 ± 0.05	133	AVERY	91	CLEO $e^+ e^-$ 10.5 GeV	
0.55 ± 0.17 ± 0.14	45	ANJOS	90	E691 γ Be 70-260 GeV	
0.62 ± 0.15 ± 0.03	73	ALBRECHT	88c	ARG $e^+ e^-$ 10 GeV	

$\Gamma(\rho K^- \pi^+)/\Gamma_{\text{total}}$					Γ_2/Γ
VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT	
0.044 ± 0.006 OUR FIT					
0.044 ± 0.006 OUR AVERAGE					
0.0594 ± 0.0031 ± 0.0144		1 BERGFELD	94	CLEO $e^+ e^- \approx \Upsilon(4S)$	
0.040 ± 0.003 ± 0.008		2 ALBRECHT	92a	ARG $e^+ e^- \approx \Upsilon(4S)$	
0.043 ± 0.010 ± 0.008		3 CRAWFORD	92	CLEO $e^+ e^-$ 10.5 GeV	
0.041 ± 0.024	208	4 ALBRECHT	88e	ARG	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>0.044	90	6	5 AGUILAR...	88B LEBc pp 27.4 GeV	
0.022 ± 0.010		39	ABRAMS	80 MRK2 $e^+ e^-$ 5.2 GeV	

1 BERGFELD 94 measures $\Gamma(\rho K^- \pi^+)/\Gamma(\Lambda_c^+ \nu_\ell) = 1.93 \pm 0.10 \pm 0.33$ and calculates $\Gamma(e^+ \text{ anything})/\Gamma_{\text{total}} = 0.034 \pm 0.004$ from D -meson data, assuming that all charmed hadrons have the same semileptonic width. Combined, these values give $\Gamma(\rho K^- \pi^+)/\Gamma_{\text{total}} = f \times (6.67 \pm 0.35 \pm 1.35)\%$, where $f \equiv \Gamma(\Lambda_c^+ \nu_\ell)/\Gamma(e^+ \text{ anything})$. Since $f \leq 1$, this gives an upper bound on $\Gamma(\rho K^- \pi^+)/\Gamma_{\text{total}}$. In the spectator model, the quantity corresponding to f in D -meson decay is $\Gamma(D \rightarrow (\bar{K} + \bar{K}^*) \ell^+ \nu_\ell)/\Gamma(D \rightarrow \ell^+ \text{ anything}) = 0.89 \pm 0.12$. This value of f leads to the value of $\Gamma(\rho K^- \pi^+)/\Gamma_{\text{total}}$ we give here.

2 ALBRECHT 92a uses $B(\bar{B} \rightarrow \Lambda_c^+ X) \times B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (0.28 \pm 0.05)\%$ plus $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.8 \pm 0.5 \pm 0.3)\%$ and assumes that $\bar{B} \rightarrow \Xi_c^- X$ and $\bar{B} \rightarrow \Omega_c^- X$ decays are suppressed and negligible.

3 CRAWFORD 92 $B(\bar{B} \rightarrow \Lambda_c^+ X) \times B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (0.273 \pm 0.051 \pm 0.039)\%$ and estimates $B(\bar{B} \rightarrow \Lambda_c^+ X) = (6.4 \pm 0.8 \pm 0.8)\%$. If final states other than $\Lambda_c^+ \bar{N} X$ contribute to \bar{B} decay, the $\Lambda_c^+ \rightarrow \rho K^- \pi^+$ branching fraction would increase.

4 ALBRECHT 88e use their result $B(\bar{B} \rightarrow \Lambda_c^+ X) \times B(\Lambda_c^+ \rightarrow \rho K^- \pi^+) = (0.30 \pm 0.12 \pm 0.06)\%$ plus $B(\bar{B} \rightarrow \Lambda_c^+ X) = (7.4 \pm 2.9)\%$ from other measurements of inclusive proton and Λ yields in B decays.

5 This AGUILAR-BENITEZ 88B limit assumes that $\tau_{\Lambda_c} = 1.2 \times 10^{-13}$ s, and it "decreases by 20% [to > 0.035] assuming a lifetime of 1.7×10^{-13} s instead." Our average for τ_{Λ_c} is still higher (see the mean-life section), which would further reduce the limit.

$\Gamma(\rho\bar{K}^*(892)^0)/\Gamma(\rho K^- \pi^+)$					Γ_3/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.36 ± 0.06 OUR AVERAGE					
0.35 ± 0.06 ± 0.03	39	BOZEK	93	NA32 π^- Cu 230 GeV	
0.42 ± 0.24	12	BASILE	81B	CNTR $pp \rightarrow \Lambda_c^+ e^- X$	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
0.35 ± 0.11		BARLAG	90D	NA32 See BOZEK 93	

$\Gamma(\Delta(1232)^{++} K^-)/\Gamma(\rho K^- \pi^+)$					Γ_4/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.16 ± 0.10 OUR AVERAGE					
0.12 ± 0.04 ± 0.05	14	BOZEK	93	NA32 π^- Cu 230 GeV	
0.40 ± 0.17	17	BASILE	81B	CNTR $pp \rightarrow \Lambda_c^+ e^- X$	

$\Gamma(\Lambda(1520)\pi^+)/\Gamma(\rho K^- \pi^+)$					Γ_5/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.09 ± 0.04 OUR AVERAGE					
0.09 ± 0.03 ± 0.02	12	BOZEK	93	NA32 π^- Cu 230 GeV	

$\Gamma(\rho K^- \pi^+ \text{ nonresonant})/\Gamma(\rho K^- \pi^+)$					Γ_6/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.56 ± 0.07 OUR AVERAGE					
0.56 ± 0.09 ± 0.05	71	BOZEK	93	NA32 π^- Cu 230 GeV	

$\Gamma(\rho\bar{K}^0 \pi^+ \pi^-)/\Gamma(\rho K^- \pi^+)$					Γ_7/Γ_2
VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT	
0.54 ± 0.17 OUR FIT					
0.49 ± 0.17 OUR AVERAGE					
0.43 ± 0.12 ± 0.04	83	AVERY	91	CLEO $e^+ e^-$ 10.5 GeV	
0.98 ± 0.36 ± 0.08	12	BARLAG	90D	NA32 π^- 230 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<1.7	90	ANJOS	90	E691 γ Be 70-260 GeV	

$\Gamma(\rho K^- \pi^+ \pi^0)/\Gamma_{\text{total}}$					Γ_8/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	44	AMENDOLIA	87	SPEC γ Ge-Si	

$\Gamma(\rho K^*(892)^- \pi^+)/\Gamma_{\text{total}}$					Γ_9/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	1	CNOPS	79	DBC νN in BNL 7-ft	

$\Gamma(\rho(K^- \pi^+)_{\text{nonresonant}} \pi^0)/\Gamma(\rho K^- \pi^+)$					Γ_{10}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.73 ± 0.12 ± 0.05	67	BOZEK	93	NA32 π^- Cu 230 GeV	

$\Gamma(\Delta(1232)\bar{K}^*(892)^-)/\Gamma_{\text{total}}$					Γ_{11}/Γ
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
seen	35	AMENDOLIA	87	SPEC γ Ge-Si	

$\Gamma(\rho K^- \pi^+ \pi^+ \pi^-)/\Gamma(\rho K^- \pi^+)$					Γ_{12}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.022 ± 0.015		BARLAG	90D	NA32 π^- 230 GeV	

$\Gamma(\rho K^- \pi^+ \pi^0 \pi^0)/\Gamma(\rho K^- \pi^+)$					Γ_{13}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.16 ± 0.07 ± 0.03	15	BOZEK	93	NA32 π^- Cu 230 GeV	

$\Gamma(\rho K^- \pi^+ \pi^0 \pi^0)/\Gamma(\rho K^- \pi^+)$					Γ_{14}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.10 ± 0.06 ± 0.02	8	BOZEK	93	NA32 π^- Cu 230 GeV	

$\Gamma(\rho \pi^+ \pi^-)/\Gamma(\rho K^- \pi^+)$					Γ_{15}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.069 ± 0.036		BARLAG	90D	NA32 π^- 230 GeV	

$\Gamma(\rho f_0(980))/\Gamma(\rho K^- \pi^+)$					Γ_{16}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.055 ± 0.036		BARLAG	90D	NA32 π^- 230 GeV	

$\Gamma(\rho \pi^+ \pi^+ \pi^- \pi^-)/\Gamma(\rho K^- \pi^+)$					Γ_{17}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.036 ± 0.023		BARLAG	90D	NA32 π^- 230 GeV	

$\Gamma(\rho K^+ K^-)/\Gamma(\rho K^- \pi^+)$					Γ_{18}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.069 ± 0.024 OUR AVERAGE					
0.096 ± 0.029 ± 0.010	30	FRABETTI	93H	E687 γ Be, \bar{E}_γ 220 GeV	
0.048 ± 0.027		BARLAG	90D	NA32 π^- 230 GeV	

$\Gamma(\rho \phi)/\Gamma(\rho K^- \pi^+)$					Γ_{19}/Γ_2
VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	
0.040 ± 0.027		BARLAG	90D	NA32 π^- 230 GeV	

$\Gamma(\rho \phi)/\Gamma(\rho K^+ K^-)$					Γ_{19}/Γ_{18}
VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT	
<0.58	90	FRABETTI	93H	E687 γ Be, \bar{E}_γ 220 GeV	

$\Gamma(\Lambda \pi^+)/\Gamma(\rho K^- \pi^+)$					Γ_{20}/Γ_2
VALUE	CL% EVTS	DOCUMENT ID	TECN	COMMENT	
0.180 ± 0.032 OUR AVERAGE					
0.18 ± 0.03 ± 0.04		ALBRECHT	92	ARG $e^+ e^- \approx 10.4$ GeV	
0.18 ± 0.03 ± 0.03	87	AVERY	91	CLEO $e^+ e^-$ 10.5 GeV	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.33	90	ANJOS	90	E691 γ Be 70-260 GeV	
<0.16	90	ALBRECHT	88c	ARG $e^+ e^-$ 10 GeV	

Λ_c^+

$\Gamma(\Lambda\pi^+)/\Gamma(\rho K^0)$ Γ_{20}/Γ_1

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •					
<0.4		90	40	RUSSELL	81 SPEC Photoproduction
0.51 ^{+0.62} _{-0.27}		9		KITAGAKI	80 DBC νd in FNAL 15-ft

$\Gamma(\Lambda\pi^0)/\Gamma(\rho K^-\pi^+)$ Γ_{21}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.73±0.09±0.16	464	AVERY	94	CLEO $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$

$\Gamma(\Lambda\rho^0)/\Gamma(\rho K^-\pi^+)$ Γ_{22}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.95	95	AVERY	94	CLEO $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$

$\Gamma(\Lambda\pi^+\pi^-)/\Gamma_{total}$ Γ_{23}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.027±0.006 OUR FIT				
0.028±0.007±0.011	70	⁶ BOWCOCK	85	CLEO e^+e^- 10.5 GeV

⁶ See BOWCOCK 85 for assumptions made on charm production and Λ_c production from charm to get this result.

$\Gamma(\Lambda\pi^+\pi^-)/\Gamma(\rho K^-\pi^+)$ Γ_{23}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.63±0.10 OUR FIT				
0.66±0.11 OUR AVERAGE				
0.65±0.11±0.12	289	AVERY	91	CLEO e^+e^- 10.5 GeV
0.82±0.29±0.27	44	ANJOS	90	E691 γ Be 70–260 GeV
0.94±0.41±0.13	10	BARLAG	90d	NA32 π^- 230 GeV
0.61±0.16±0.04	105	ALBRECHT	88c	ARG e^+e^- 10 GeV

$\Gamma(\rho K^0\pi^+)/\Gamma(\Lambda\pi^+\pi^-)$ Γ_{7}/Γ_{23}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.86±0.28 OUR FIT				Error includes scale factor of 1.2.
4.3 ±1.2	130	ALEEV	84	BIS2 n C 40–70 GeV

$\Gamma(\Sigma^0\pi^+)/\Gamma(\rho K^-\pi^+)$ Γ_{24}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.20±0.04 OUR AVERAGE				
0.21±0.02±0.04	196	AVERY	94	CLEO $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$
0.17±0.06±0.04		ALBRECHT	92	ARG $e^+e^- \approx 10.4$ GeV

$\Gamma(\Sigma^0\pi^0)/\Gamma(\rho K^-\pi^+)$ Γ_{25}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.36±0.09±0.10	117	AVERY	94	CLEO $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$

$\Gamma(\Sigma^0\pi^+\pi^-)/\Gamma(\rho K^-\pi^+)$ Γ_{26}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.21±0.05±0.05	90	AVERY	94	CLEO $e^+e^- \approx \Upsilon(3S), \Upsilon(4S)$

$\Gamma(\Sigma^+\pi^0)/\Gamma(\rho K^-\pi^+)$ Γ_{27}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.20±0.03±0.03	93	KUBOTA	93	CLEO $e^+e^- \approx \Upsilon(4S)$

$\Gamma(\Sigma^+\pi^-\pi^-)/\Gamma_{total}$ Γ_{28}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
0.10±0.08		ADAMOVICH	87	EMUL γ A 20–70 GeV/c
seen	1	AMMAR	86	EMUL ν A

$\Gamma(\Sigma^+\pi^-\pi^-)/\Gamma(\rho K^-\pi^+)$ Γ_{28}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.68±0.09 OUR AVERAGE				
0.74±0.07±0.09	487	KUBOTA	93	CLEO $e^+e^- \approx \Upsilon(4S)$
0.54 ^{+0.18} _{-0.15}	11	BARLAG	92	NA32 π^- -Cu 230 GeV

$\Gamma(\Sigma^+\rho^0)/\Gamma(\rho K^-\pi^+)$ Γ_{29}/Γ_2

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.27	95	KUBOTA	93	CLEO $e^+e^- \approx \Upsilon(4S)$

$\Gamma(\Sigma^-\pi^+\pi^-)/\Gamma(\Sigma^+\pi^-\pi^-)$ Γ_{30}/Γ_{28}

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.53±0.15±0.07	56	FRABETTI	94e	E687 γ Be, \bar{E}_γ 220 GeV

$\Gamma(\Sigma^+\omega)/\Gamma(\rho K^-\pi^+)$ Γ_{32}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the ω are included.				
0.54±0.13±0.06	107	KUBOTA	93	CLEO $e^+e^- \approx \Upsilon(4S)$

$\Gamma(\Sigma^+\pi^+\pi^-\pi^-)/\Gamma(\rho K^-\pi^+)$ Γ_{33}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.06^{+0.08}_{-0.04}	1	BARLAG	92	NA32 π^- -Cu 230 GeV

$\Gamma(\Sigma^+K^+K^-)/\Gamma(\rho K^-\pi^+)$ Γ_{34}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.071±0.015 OUR AVERAGE				
0.070±0.011±0.011	59	AVERY	93	CLEO $e^+e^- \approx 10.5$ GeV
0.13 ^{+0.18} _{-0.09}	1	BARLAG	92	NA32 π^- -Cu 230 GeV

$\Gamma(\Sigma^+\phi)/\Gamma(\rho K^-\pi^+)$ Γ_{35}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the ϕ are included.				
0.069±0.023±0.016	26	AVERY	93	CLEO $e^+e^- \approx 10.5$ GeV

$\Gamma(\Sigma^+K^+\pi^-)/\Gamma(\rho K^-\pi^+)$ Γ_{36}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.13^{+0.12}_{-0.07}	2	BARLAG	92	NA32 π^- -Cu 230 GeV

$\Gamma(\Xi^0K^+)/\Gamma(\rho K^-\pi^+)$ Γ_{37}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.078±0.013±0.013	56	AVERY	93	CLEO $e^+e^- \approx 10.5$ GeV

$\Gamma(\Xi^-K^+\pi^+)/\Gamma(\rho K^-\pi^+)$ Γ_{38}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.088±0.024 OUR AVERAGE				Error includes scale factor of 1.3.
0.079±0.013±0.014	60	AVERY	93	CLEO $e^+e^- \approx 10.5$ GeV
0.15±0.04±0.03	30	AVERY	91	CLEO e^+e^- 10.5 GeV

$\Gamma(\Xi(1530)^0K^+)/\Gamma(\rho K^-\pi^+)$ Γ_{39}/Γ_2

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
Unseen decay modes of the $\Xi(1530)^0$ are included.				
0.053±0.016±0.010	24	AVERY	93	CLEO $e^+e^- \approx 10.5$ GeV

$\Gamma(\Lambda \text{ anything } \ell^+\nu_\ell)/\Gamma(\Lambda\ell^+\nu_\ell)$ Γ_{41}/Γ_{40}

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.15	90	BERGFELD	94	CLEO $e^+e^- \approx \Upsilon(4S)$

$\Gamma(\rho \text{ anything})/\Gamma_{total}$ Γ_{42}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.80±0.08±0.14	⁷ CRAWFORD	92	CLEO e^+e^- 10.5 GeV

⁷ This CRAWFORD 92 value includes protons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(\rho \text{ anything (no } \Lambda))/\Gamma_{total}$ Γ_{43}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.12±0.10±0.16	CRAWFORD	92	CLEO e^+e^- 10.5 GeV

$\Gamma(n \text{ anything})/\Gamma_{total}$ Γ_{45}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.50±0.08±0.14	⁸ CRAWFORD	92	CLEO e^+e^- 10.5 GeV

⁸ This CRAWFORD 92 value includes neutrons from Λ decay. The value is model dependent, but account is taken of this in the systematic error.

$\Gamma(n \text{ anything (no } \Lambda))/\Gamma_{total}$ Γ_{46}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
0.29±0.09±0.15	CRAWFORD	92	CLEO e^+e^- 10.5 GeV

$\Gamma(\rho \text{ hadrons})/\Gamma_{total}$ Γ_{44}/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
0.41±0.24	ADAMOVICH	87	EMUL γ A 20–70 GeV/c


$\Gamma(\Lambda \text{ anything})/\Gamma_{total}$ Γ_{47}/Γ

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
0.35±0.11 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
0.59±0.10±0.12		CRAWFORD	92	CLEO e^+e^- 10.5 GeV
0.49±0.24		ADAMOVICH	87	EMUL γ A 20–70 GeV/c
0.23±0.10	8	⁹ ABE	86	HYBR 20 GeV γ p

⁹ ABE 86 includes Λ 's from Σ^0 decay.

Baryon Full Listings

$\Lambda_c^+, \Lambda_c(2625)^+$



$\Gamma(\Sigma^\pm \text{ anything}) / \Gamma_{\text{total}}$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{48} / Γ
	0.1 ± 0.05	5	ABE	86	HYBR 20 GeV γp	

$\Gamma(e^+ \text{ anything}) / \Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{49} / Γ
	0.045 ± 0.017	VELLA 82	MRK2	$e^+ e^-$ 4.5–6.8 GeV	

$\Gamma(p e^+ \text{ anything}) / \Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{50} / Γ
	0.018 ± 0.009	10 VELLA 82	MRK2	$e^+ e^-$ 4.5–6.8 GeV	

¹⁰VELLA 82 includes protons from Λ decay.

$\Gamma(\Lambda e^+ \text{ anything}) / \Gamma_{\text{total}}$	VALUE	DOCUMENT ID	TECN	COMMENT	Γ_{51} / Γ
	0.014 ± 0.005 OUR FIT				
	0.011 ± 0.008	11 VELLA 82	MRK2	$e^+ e^-$ 4.5–6.8 GeV	

¹¹VELLA 82 includes Λ 's from Σ^0 decay.

$\Gamma(\Lambda e^+ \text{ anything}) / \Gamma(p K^- \pi^+)$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{51} / Γ_2
	0.33 ± 0.11 OUR FIT					
	0.37 ± 0.11 ± 0.08	73	ALBRECHT	91G	ARG $e^+ e^- \approx 10.4$ GeV	

$\Gamma(\Lambda \mu^+ \text{ anything}) / \Gamma(p K^- \pi^+)$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT	Γ_{52} / Γ_2
	0.35 ± 0.18 ± 0.09	30	ALBRECHT	91G	ARG $e^+ e^- \approx 10.4$ GeV	

$\Gamma(\Lambda \pi^+ \pi^+ \pi^-) / \Gamma(\Lambda e^+ \text{ anything})$	VALUE	CL%	DOCUMENT ID	TECN	COMMENT	$\Gamma_{23} / \Gamma_{51}$
	< 1.7	90	KLEIN	89	MRK2 $e^+ e^-$ 29 GeV	

• • • We do not use the following data for averages, fits, limits, etc. • • •

Λ_c^+ DECAY PARAMETERS

See the Note on Baryon Decay Parameters in the neutron Listings.

α FOR $\Lambda_c^+ \rightarrow \Lambda \pi^+$	VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
	-1.03 ± 0.29 OUR AVERAGE				
	-0.96 ± 0.42		ALBRECHT	92	ARG $e^+ e^- \approx 10.4$ GeV
	-1.1 ± 0.4	86	AVERY	90B	CLEO $e^+ e^- \approx 10.6$ GeV

α FOR $\Lambda_c^+ \rightarrow \Lambda e^+ \nu_e$	VALUE	DOCUMENT ID	TECN	COMMENT
	-0.89 + 0.17 + 0.09 -0.11 - 0.05	BERGFELD 94	CLEO	$e^+ e^- \approx \gamma(4S)$

Λ_c^+ REFERENCES

We have omitted some papers that have been superseded by later experiments. The omitted papers may be found in our 1992 edition (Physical Review D45, 1 June, Part II) or in earlier editions.

AVERY	94	PL B325 257	+ Freyberger, Rodriguez+	(CLEO Collab.)
BERGFELD	94	PL B323 219	+ Eisenstein, Golin, Ong+	(CLEO Collab.)
FRABETTI	94E	PL B328 193	+ Cheung, Cumalat+	(FNAL E687 Collab.)
AVERY	93	PRL 71 2391	+ Freyberger, Rodriguez+	(CLEO Collab.)
BOZEK	93	PL B312 247	+ Barlag, Becker, Boehringer+	(CERN NA32 Collab.)
FRABETTI	93D	PRL 70 1755	+ Cheung, Cumalat+	(FNAL E687 Collab.)
FRABETTI	93H	PL B314 477	+ Cheung, Cumalat+	(FNAL E687 Collab.)
KUBOTA	93	PRL 71 3255	+ Lattery, Nelson, Patton+	(CLEO Collab.)
ALBRECHT	92	PL B274 239	+ Ehrlichmann, Hamacher, Krueger+	(ARGUS Collab.)
ALBRECHT	92O	ZPHY C56 1	+ Cronstroem, Ehrlichmann+	(ARGUS Collab.)
BARLAG	92	PL B283 465	+ Becker, Bozek, Boehringer+	(ACCMOR Collab.)
CRAWFORD	92	PR D45 752	+ Fulton, Jensen, Johnson+	(CLEO Collab.)
JEZABEK	92	PL B286 175	+ Rybicki, Ryiko	(CRAC)
ALBRECHT	91G	PL B269 234	+ Ehrlichmann, Hamacher+	(ARGUS Collab.)
AVERY	91	PR D43 3599	+ Besson, Garren, Yelton+	(CLEO Collab.)
ALVAREZ	90	ZPHY C47 539	+ Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ALVAREZ	90B	PL B246 256	+ Barate, Bloch, Bonamy+	(CERN NA14/2 Collab.)
ANJOS	90	PR D41 801	+ Appel, Bean+	(FNAL E691 Collab.)
AVERY	90B	PRL 65 2842	+ Besson, Garren, Yelton, Kinoshita+	(CLEO Collab.)
BARLAG	90D	ZPHY C48 29	+ Becker, Boehringer, Bosman+	(ACCMOR Collab.)
FRABETTI	90	PL B251 639	+ Bogart, Cheung, Coteus+	(FNAL E687 Collab.)
BARLAG	89	PL B218 374	+ Becker, Boehringer, Bosman+	(ACCMOR Collab.)
KLEIN	89	PRL 62 2444	+ Himel, Abrams, Amidei, Baden+	(Mark II Collab.)
AGUILAR...	88B	ZPHY C40 321	+ Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
Also	87	PL B189 254	+ Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
Also	87B	PL B199 462	+ Aguilar-Benitez, Allison, Bailly+	(LEBC-EHS Collab.)
Also	88	SJNP 48 833	+ Begalli, Otter, Schulte, Gensch+	(LEBC-EHS Collab.)
ALBRECHT	88C	Translated from YAF 48 1310.		(ARGUS Collab.)
ALBRECHT	88C	PL B207 109	+ Bockmann, Glaeser--	(ARGUS Collab.)
ANJOS	88B	PRL 60 1379	+ Appel+	(FNAL E691 Collab.)
ADAMOVICH	87	EPL 4 887	+ Alexandrov, Bolta+	(Photon Emulsion Collab.)
Also	87	SJNP 46 447	+ Viaggi, Gessaroli+	(Photon Emulsion Collab.)
AMENDOLIA	87	Translated from YAF 46 799.		
BARLAG	87	ZPHY C36 513	+ Bagliesi, Batignani, Beck+	(CERN NA1 Collab.)
JONES	87	PL B184 283	+ Jones, Kennedy, O'Neale+	(ACCMOR Collab.)
ABE	86	ZPHY C36 593	+ Jones, Kennedy, O'Neale+	(CERN WA21 Collab.)
AMMAR	86	PR D33 1	+ (SLAC Hybrid Facility Photon Collab.)	
USHIDA	86	JETPL 43 515	+ Ammosov, Bakic, Baranov, Burnett+	(ITEP)
BOWCOCK	85	Translated from ZETFP 43 401.		
ALEEV	84	PRL 56 1767	+ Kondo, Tasaka, Park+	(FNAL E653 Collab.)
BOSETTI	82	PRL 55 923	+ Giles, Hassard, Kinoshita+	(CLEO Collab.)
VELLA	82	ZPHY C23 333	+ Arefeff, Balandin, Berdyshev+	(BIS-2 Collab.)
BASILE	81B	PL 109B 234	+ Graessler+	(AACH3, BONN, CERN, MPIM, OXF)
RUSSELL	81	PRL 48 1515	+ Trilling, Abrams, Alam+	(SLAC, LBL, UC8)
ABRAMS	80	NC 62A 14	+ Romeo+	(CERN, BGNA, PGIA, FRAS)
CALICCHIO	80	PRL 46 799	+ Avery, Butler, Gladding+	(ILL, FNAL, COLU)
KITAGAKI	80	PRL 44 10	+ Alam, Blocker, Boyarski+	(SLAC, LBL)
CNOPS	79	PL 93B 521	+ Tanaka, Yuta+	(BARI, BIRM, BRUX, CERN, EPOL, RHEL)
		PRL 45 955	+ Tanaka, Yuta+	(TOHOK, IIT, UMD, STON, TUFTS)
		PRL 42 197	+ Connolly, Kahn, Kirk, Murtagh, Palmer+	(BNL)

$\Lambda_c(2625)^+$

$I = 0$ Status: * * *

This could be a $\Sigma_c(2625)^+$ instead of a $\Lambda_c(2625)^+$, but theoretical estimates of the masses of excited charmed baryons are much more in accord with it being a Λ_c^+ .

$\Lambda_c(2625)^+$ MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
2625.6 ± 0.8 OUR FIT				
2626.6 ± 0.5 ± 1.5	42	¹ ALBRECHT	93F	ARG $e^+ e^- \approx \gamma(4S)$

¹ALBRECHT 93F claims a signal of 42.4 ± 8.8 events.

$m_{\Lambda_c(2625)^+} - m_{\Lambda_c^+}$

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
340.6 ± 0.6 OUR FIT				
340.4 ± 0.6 ± 0.3	40	² FRABETTI	94	E687 γ Be, $\bar{E}_{\gamma} = 220$ GeV

²FRABETTI 94 claims a signal of 39.7 ± 8.7 events.

$\Lambda_c(2625)^+$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID	TECN	COMMENT
< 3.2	90	ALBRECHT	93F	ARG $e^+ e^- \approx \gamma(4S)$

$\Lambda_c(2625)^+$ DECAY MODES

Mode	Fraction (Γ_i / Γ)
Γ_1 $\Lambda_c^+ \pi^+ \pi^-$	seen
Γ_2 $\Sigma_c(2455)^{++} \pi^- + \Sigma_c(2455)^0 \pi^+$	seen
Γ_3 $\Lambda_c^+ \pi^+ \pi^-$ nonresonant	seen

See key on page 1343

Baryon Full Listings

$\Lambda_c(2625)^+$, $\Sigma_c(2455)$, $\Sigma_c(2530)$

 $\Lambda_c(2625)^+$ BRANCHING RATIOS

$[\Gamma(\Sigma_c(2455)^{++}\pi^-) + \Gamma(\Sigma_c(2455)^0\pi^+)] / \Gamma(\Lambda_c^+\pi^+\pi^-)$ Γ_2/Γ_1					
VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.46±0.14		21	ALBRECHT	93F ARG	$e^+e^- \approx \mathcal{T}(4S)$
••• We do not use the following data for averages, fits, limits, etc. •••					
<0.36		90	FRABETTI	94 E687	γBe , $\bar{E}_\gamma = 220\text{ GeV}$
$\Gamma(\Lambda_c^+\pi^+\pi^- \text{ nonresonant}) / \Gamma(\Lambda_c^+\pi^+\pi^-)$ Γ_3/Γ_1					
VALUE	CL%	EVTs	DOCUMENT ID	TECN	COMMENT
0.54±0.14		16	ALBRECHT	93F ARG	$e^+e^- \approx \mathcal{T}(4S)$

 $\Lambda_c(2625)^+$ REFERENCES

FRABETTI	94	PRL 72 961	+Cheung, Cumalat+	(FNAL E687 Collab.)
ALBRECHT	93F	PL B317 227	+Ehrlichmann, Hamacher+	(ARGUS Collab.)

 $\Sigma_c(2455)$

$$I(J^P) = 1(\frac{1}{2}^+) \text{ Status: } ***$$

J^P is not confirmed. $1/2^+$ is the quark model prediction.

 $\Sigma_c(2455)$ MASSES

The mass measurements in this section are redundant with the mass difference measurements that follow. We get the masses by adding $m_{\Sigma_c(2455)} - m_{\Lambda_c^+}$ to the Λ_c^+ mass.

 $\Sigma_c(2455)^{++}$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
2453.1±0.6 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
2449 ± 3	2	JONES	87 HBC	++	νp in BEBC
2480	1	ADAMOVIICH	84 EMUL	++	γA (OMEGA)
2454 ± 5	1	BOSETTI	82 HBC	++	See JONES 87
2425 ± 10	6	BALTAY	79 HLBC	++	ν Ne-H in 15-ft
>2439	1	BARISH	77B DBC	++	νd in 12-ft
2426 ± 12	1	CAZZOLI	75 HBC	++	νp in BNL 7-ft

 $\Sigma_c(2455)^+$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
2453.8±0.9 OUR FIT					
••• We do not use the following data for averages, fits, limits, etc. •••					
2457 ± 4	1	CALICCHIO	80 HBC	+	νp in BEBC-TST

 $\Sigma_c(2455)^0$ MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
2452.4±0.7 OUR FIT					Error includes scale factor of 1.1.
••• We do not use the following data for averages, fits, limits, etc. •••					
2462 ± 26	1	AMMAR	86 EMUL	0	νA
~2460	9	KNAPP	76 SPEC	0	γBe

 $m_{\Sigma_c(2455)} - m_{\Lambda_c^+}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
168.04±0.27 OUR FIT					
167.98±0.27 OUR AVERAGE					
168.2 ± 0.3 ± 0.2	126	CRAWFORD	93 CLEO		$e^+e^- \approx \mathcal{T}(4S)$
167.8 ± 0.4 ± 0.3	54	BOWCOCK	89 CLEO	++	e^+e^- 10 GeV
168.2 ± 0.5 ± 1.6	92	ALBRECHT	88D ARG	++	e^+e^- 10 GeV
167.4 ± 0.5 ± 2.0	46	DIESBURG	87 SPEC	++	$nA \sim 600\text{ GeV}$
167 ± 1	2	JONES	87 HBC	++	ν Ne in BEBC
168 ± 3	6	BALTAY	79 HLBC	++	ν Ne-H in 15-ft
••• We do not use the following data for averages, fits, limits, etc. •••					
166 ± 1	1	BOSETTI	82 HBC	++	See JONES 87
166 ± 15	1	CAZZOLI	75 HBC	++	νp in BNL 7-ft

 $m_{\Sigma_c^+} - m_{\Lambda_c^+}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
168.7±0.7 OUR FIT					Error includes scale factor of 1.1.
168 ± 3					
••• We do not use the following data for averages, fits, limits, etc. •••					
168.5±0.4±0.2	111	¹ CRAWFORD	93 CLEO		$e^+e^- \approx \mathcal{T}(4S)$
¹ This result enters the fit through $m_{\Sigma_c^+} - m_{\Sigma_c^0}$ below.					

 $m_{\Sigma_c^0} - m_{\Lambda_c^+}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
167.3±0.4 OUR FIT					Error includes scale factor of 1.2.
168.4±1.0±0.3					
167.1±0.3±0.2	124	² CRAWFORD	93 CLEO		$e^+e^- \approx \mathcal{T}(4S)$
167.9±0.5±0.3	48	² BOWCOCK	89 CLEO	0	e^+e^- 10 GeV
167.0±0.5±1.6	70	² ALBRECHT	88D ARG	0	e^+e^- 10 GeV
178.2±0.4±2.0	85	³ DIESBURG	87 SPEC	0	$nA \sim 600\text{ GeV}$
163 ± 2	1	AMMAR	86 EMUL	0	νA
² This result enters the fit through $m_{\Sigma_c^{++}} - m_{\Sigma_c^0}$ given below.					
³ See the note on DIESBURG 87 in the $m_{\Sigma_c^{++}} - m_{\Sigma_c^0}$ section below.					

 $\Sigma_c(2455)$ MASS DIFFERENCES **$m_{\Sigma_c^{++}} - m_{\Sigma_c^0}$**

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
0.7±0.4 OUR FIT					Error includes scale factor of 1.2.
0.8±0.4 OUR AVERAGE					Error includes scale factor of 1.2.
1.1±0.4±0.1		CRAWFORD	93 CLEO		$e^+e^- \approx \mathcal{T}(4S)$
-0.1±0.6±0.1		BOWCOCK	89 CLEO		e^+e^- 10 GeV
+1.2±0.7±0.3		ALBRECHT	88D ARG		$e^+e^- \sim 10\text{ GeV}$
••• We do not use the following data for averages, fits, limits, etc. •••					
-10.8±2.9		⁴ DIESBURG	87 SPEC		$nA \sim 600\text{ GeV}$
⁴ DIESBURG 87 is completely incompatible with the other experiments, which is surprising since it agrees with them about $m_{\Sigma_c(2455)^{++}} - m_{\Lambda_c^+}$. We go with the majority here.					

 $m_{\Sigma_c^+} - m_{\Sigma_c^0}$

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
1.4±0.6 OUR FIT					
1.4±0.5±0.3					
		CRAWFORD	93 CLEO		$e^+e^- \approx \mathcal{T}(4S)$

 $\Sigma_c(2455)$ DECAY MODES

Mode	Fraction (Γ_i/Γ)
$\Gamma_1 \Lambda_c^+\pi$	100%

 $\Sigma_c(2455)$ REFERENCES

CRAWFORD	93	PRL 71 3259	+Daubenmier, Fulton+	(CLEO Collab.)
ANJOS	89D	PRL 62 1721	+Appel, Bean, Bracker, Browder+	(FNAL E691 Collab.)
BOWCOCK	89	PRL 62 1240	+Kinoshita, Pipkin, Procaro, Wilson+	(CLEO Collab.)
ALBRECHT	88D	PL B211 489	+Bockmann, Glaeser+	(ARGUS Collab.)
DIESBURG	87	PRL 59 2711	+Ladbury, Binkley+	(FNAL E400 Collab.)
JONES	87	ZPHY C36 593	+Jones, Kennedy, O'Neale+	(CERN WA21 Collab.)
AMMAR	86	JETPL 43 515	+Ammosov, Bakic, Baranov, Burnett+	(ITEP)
			Translated from ZETFP 43 401.	
ADAMOVIICH	84	PL 140B 119	+Alexandrov, Bolta, Bravo+	(CERN WA58 Collab.)
BOSETTI	82	PL 109B 234	+Graessler+	(AACH3, BONN, CERN, MPIM, OXF)
CALICCHIO	80	PL 93B 521	+	(BARI, BIRM, BRUX, CERN, EPOL, RHEL)
BALTAY	79	PRL 42 1721	+Caroumbalis, French, Hibbs+	(COLU, BNL)
BARISH	77B	PR D15 1	+Derrick, Dombeck, Musgrave+	(ANL, PURD)
KNAPP	76	PRL 37 882	+Lee, Leung, Smith+	(COLU, HAWA, ILL, FNAL)
CAZZOLI	75	PRL 34 1125	+Cnops, Connolly, Louttit, Murtagh+	(BNL)

 $\Sigma_c(2530)$

Status: *

OMITTED FROM SUMMARY TABLE

 $\Sigma_c(2530)$ MASSES

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	CHG	COMMENT
2530±5±5	6	¹ AMMOISOV	93 HLBC		$\nu p \rightarrow \mu^- \Sigma_c(2530)^{++}$
¹ AMMOISOV 93 sees a cluster of 6 events and estimates the background to be 1 event.					

 $\Sigma_c(2530)$ REFERENCES

AMMOISOV	93	JETPL 58 247	+Vasil'ev, Ivanilov, Ivanov+	(SERP)
			Translated from ZETFP 58 241.	

Baryon Full Listings

Ξ_c^+ , Ξ_c^0



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: ***

According to the quark model, the Ξ_c^+ (quark content usc) and Ξ_c^0 form an isospin doublet, and the spin-parity ought to be $J^P = 1/2^+$. None of I , J , or P have actually been measured.

Ξ_c^+ MASS

The fit uses the Ξ_c^+ and Ξ_c^0 mass and mass-difference measurements.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2465.1 ± 1.6 OUR FIT				
2465.4 ± 1.6 OUR AVERAGE				
2464.4 ± 2.0 ± 1.4	30	FRABETTI	93B E687	γ Be $\bar{E}_\gamma = 220$ GeV
2465.1 ± 3.6 ± 1.9	30	ALBRECHT	90F ARG	e^+e^- at $\Upsilon(4S)$
2467 ± 3 ± 4	23	ALAM	89 CLEO	e^+e^- 10.6 GeV
2466.5 ± 2.7 ± 1.2	5	BARLAG	89C ACCM	π^- Cu 230 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2459 ± 5 ± 30	56	¹ COTEUS	87 SPEC	$nA \approx 600$ GeV
2460 ± 25	82	BIAGI	83 SPEC	Σ^- Be 135 GeV

¹ Although COTEUS 87 claims to agree well with BIAGI 83 on the mass and width, there appears to be a discrepancy between the two experiments. BIAGI 83 sees a single peak (stated significance about 6 standard deviations) in the $\Lambda K^- \pi^+ \pi^+$ mass spectrum. COTEUS 87 sees two peaks in the same spectrum, one at the Ξ_c^+ mass, the other 75 MeV lower. The latter is attributed to $\Xi_c^+ \rightarrow \Sigma^0 K^- \pi^+ \pi^+ \rightarrow (\Lambda \gamma) K^- \pi^+ \pi^+$, with the γ unseparated. The combined significance of the double peak is stated to be 5.5 standard deviations. But the absence of any trace of a lower peak in BIAGI 83 seems to us to throw into question the interpretation of the lower peak of COTEUS 87.

Ξ_c^+ MEAN LIFE

VALUE (10^{-12} s)	EVTs	DOCUMENT ID	TECN	COMMENT
0.35^{+0.07}_{-0.04} OUR AVERAGE				
0.41 ^{+0.11} _{-0.08} ± 0.02	30	FRABETTI	93B E687	γ Be $\bar{E}_\gamma = 220$ GeV
0.20 ^{+0.11} _{-0.06}	6	BARLAG	89C ACCM	π^- (K^-) Cu 230 GeV
0.40 ^{+0.18} _{-0.12} ± 0.10	102	COTEUS	87 SPEC	$nA \approx 600$ GeV
0.48 ^{+0.21} _{-0.15} ± 0.20	53	BIAGI	85C SPEC	Σ^- Be 135 GeV

Ξ_c^+ DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Lambda K^- \pi^+ \pi^+$	seen
Γ_2 $\Sigma^+ K^- \pi^+$	seen
Γ_3 $\Sigma^0 K^- \pi^+ \pi^+$	seen
Γ_4 $\Xi^- \pi^+ \pi^+$	seen

Ξ_c^+ BRANCHING RATIOS

$\Gamma(\Lambda K^- \pi^+ \pi^+)/\Gamma_{total}$	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_1/Γ
seen	56	COTEUS	87 SPEC	$nA \approx 600$ GeV	
seen	82	² BIAGI	83 SPEC	Σ^- Be 135 GeV	

² BIAGI 85B look for but do not see the Ξ_c^+ in $\rho K^- \bar{K}^0 \pi^+$ ($\Gamma(\rho K^- \bar{K}^0 \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+) < 0.08$ with 90% CL), $\rho 2K^- 2\pi^+$ ($\Gamma(\rho 2K^- 2\pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+) < 0.03$, 90% CL), $\Omega^- K^+ \pi^+$, $\Lambda K^0 \pi^+$, and $\Sigma(1385)^+ K^- \pi^+$.

$\Gamma(\Sigma^+ K^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+)$	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_2/Γ_4
0.09^{+0.13}_{-0.06} ± 0.02	5	BARLAG	89C ACCM	$2 \Sigma^+ K^- \pi^+$, $3 \Xi^- \pi^+ \pi^+$	

$\Gamma(\Sigma^0 K^- \pi^+ \pi^+)/\Gamma(\Lambda K^- \pi^+ \pi^+)$	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_3/Γ_1
0.84 ± 0.36	47	³ COTEUS	87 SPEC	$nA \approx 600$ GeV	

³ See, however, the note on the COTEUS 87 Ξ_c^+ mass measurement.

$\Gamma(\Xi^- \pi^+ \pi^+)/\Gamma_{total}$	EVTs	DOCUMENT ID	TECN	COMMENT	Γ_4/Γ
seen	30	FRABETTI	93B E687	γ Be $\bar{E}_\gamma = 220$ GeV	
seen	30	ALBRECHT	90F ARG	e^+e^- at $\Upsilon(4S)$	
seen	23	ALAM	89 CLEO	e^+e^- 10.6 GeV	

Ξ_c^+ REFERENCES

FRABETTI	93B	PRL 70 1381	+Cheung, Cumalat+	(FNAL E687 Collab.)
ALBRECHT	90F	PL B247 121	+Ehrlichmann, Harder, Kruger, Nau+	(ARGUS Collab.)
ALAM	89	PL B226 401	+Katayama, Kim, Li, Lou, Sun+	(CLEO Collab.)
BARLAG	89C	PL B233 522	+Boehringer, Bosman+	(ACCMOR Collab.)
COTEUS	87	PRL 59 1530	+Binkley+	(FNAL E400 Collab.)
BIAGI	85B	ZPHY C28 175		(CERN WA62 Collab.)
BIAGI	85C	PL 150B 230		(CERN WA62 Collab.)
BIAGI	83	PL 122B 455		(CERN WA62 Collab.)



$I(J^P) = \frac{1}{2}(\frac{1}{2}^+)$ Status: ***

According to the quark model, the Ξ_c^0 (quark content dsc) and Ξ_c^+ form an isospin doublet, and the spin-parity ought to be $J^P = 1/2^+$. None of I , J , or P have actually been measured.

Ξ_c^0 MASS


The fit uses the Ξ_c^0 and Ξ_c^+ mass and mass difference measurements.

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2470.3 ± 1.8 OUR FIT				Error includes scale factor of 1.3.
2470.4 ± 2.0 OUR AVERAGE				Error includes scale factor of 1.4. See the ideogram below.
2462.1 ± 3.1 ± 1.4	42	¹ FRABETTI	93C E687	γ Be $\bar{E}_\gamma = 220$ GeV
2469 ± 2 ± 3	9	HENDERSON	92B CLEO	$\Omega^- K^+$
2472.1 ± 2.7 ± 1.6	54	ALBRECHT	90F ARG	e^+e^- at $\Upsilon(4S)$
2473.3 ± 1.9 ± 1.2	4	BARLAG	90 ACCM	π^- (K^-) Cu 230 GeV
2472 ± 3 ± 4	19	ALAM	89 CLEO	e^+e^- 10.6 GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
2471 ± 3 ± 4	14	AVERY	89 CLEO	See ALAM 89

¹ The FRABETTI 93C mass is well below the other measurements.

WEIGHTED AVERAGE

2470.4 ± 2.0 (Error scaled by 1.4)



Ξ_c^0 mass (MeV)

$m_{\Xi_c^0} - m_{\Xi_c^+}$

VALUE (MeV)	DOCUMENT ID	TECN	COMMENT
5.2 ± 2.2 OUR FIT			Error includes scale factor of 1.1.
6.3 ± 2.3 OUR AVERAGE			
+7.0 ± 4.5 ± 2.2	ALBRECHT	90F ARG	e^+e^- at $\Upsilon(4S)$
+6.8 ± 3.3 ± 0.5	BARLAG	90 ACCM	π^- (K^-) Cu 230 GeV
+5 ± 4 ± 1	ALAM	89 CLEO	$\Xi_c^0 \rightarrow \Xi^- \pi^+$, $\Xi_c^+ \rightarrow \Xi^0 \pi^+$

Ξ_c^0 MEAN LIFE

VALUE (10^{-12} s)	EVTs	DOCUMENT ID	TECN	COMMENT
0.098^{+0.023}_{-0.015} OUR AVERAGE				
0.101 ± 0.025	42	FRABETTI	93C E687	γ Be $\bar{E}_\gamma = 220$ GeV
0.082 ^{+0.059} _{-0.030}	4	BARLAG	90 ACCM	π^- (K^-) Cu 230 GeV

See key on page 1343

Baryon Full Listings

 Ξ_c^0, Ω_c^0 Ξ_c^0 DECAY MODES

A few branching ratios but no absolute branching fractions have been measured.

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Xi^- \ell^+ \text{ anything}$	[a] seen
Γ_2 $\Xi^- \pi^+$	seen
Γ_3 $\Xi^- \pi^+ \pi^+ \pi^-$	seen
Γ_4 $p K^- \bar{K}^*(892)^0$	seen
Γ_5 $\Omega^- K^+$	seen

[a] ℓ indicates e or μ mode, not sum over modes.

 Ξ_c^0 BRANCHING RATIOS

$\Gamma(\Xi^- \ell^+ \text{ anything})/\Gamma(\Xi^- \pi^+)$ Γ_1/Γ_2
The ratio is for the average (not the sum) of the $\Xi^- e^+$ anything and $\Xi^- \mu^+$ anything modes.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$0.96 \pm 0.43 \pm 0.18$	18	ALBRECHT	93B ARG	$e^+ e^- \approx 10.4$ GeV

$\Gamma(\Xi^- \ell^+ \text{ anything})/\Gamma(\Xi^- \pi^+ \pi^+ \pi^-)$ Γ_1/Γ_3
The ratio is for the average (not the sum) of the $\Xi^- e^+$ anything and $\Xi^- \mu^+$ anything modes.

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$0.29 \pm 0.12 \pm 0.04$	18	ALBRECHT	93B ARG	$e^+ e^- \approx 10.4$ GeV

$\Gamma(\Xi^- \pi^+)/\Gamma(\Xi^- \pi^+ \pi^+ \pi^-)$ Γ_2/Γ_3

VALUE	DOCUMENT ID	TECN	COMMENT
$0.30 \pm 0.12 \pm 0.05$	ALBRECHT	90F ARG	$e^+ e^-$ at $\Upsilon(4S)$

$\Gamma(p K^- \bar{K}^*(892)^0)/\Gamma_{\text{total}}$ Γ_4/Γ

VALUE	DOCUMENT ID	TECN	COMMENT
seen	BARLAG	90 ACCM	$\pi^- (K^-)$ Cu 230 GeV

$\Gamma(\Omega^- K^+)/\Gamma(\Xi^- \pi^+)$ Γ_5/Γ_2

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
$0.50 \pm 0.21 \pm 0.05$	9	HENDERSON	92B CLEO	$e^+ e^- \approx 10.6$ GeV

 Ξ_c^0 REFERENCES

ALBRECHT	93B	PL B303 368	+Cronstroem, Ehrlichmann+	(ARGUS Collab.)
FRABETTI	93C	PRL 70 2058	+Cheung, Cumalat+	(FNAL E687 Collab.)
HENDERSON	92B	PL B283 161	+Kinoshita, Pipkin, Saulnier+	(CLEO Collab.)
ALBRECHT	90F	PL B247 121	+Ehrlichmann, Harder, Kruger, Nau+	(ARGUS Collab.)
BARLAG	90	PL B236 495	+Becker, Boehringer, Bosman+	(ACCMOR Collab.)
ALAM	89	PL B226 401	+Katayama, Kim, Li, Lou, Sun+	(CLEO Collab.)
AVERY	89	PRL 62 863	+Besson, Garren, Yelton, Bowcock+	(CLEO Collab.)

 Ω_c^0

$I(J^P) = 0(\frac{1}{2}^+)$ Status: **

OMITTED FROM SUMMARY TABLE

The quantum numbers have not been measured, but are simply assigned in accord with the quark model, in which the Ω_c^0 is the ssc ground state.

BIAGI 85B and ALBRECHT 92H see bumps in the $\Xi^- K^- \pi^+ \pi^+$ mass spectrum. FRABETTI 93 sees a bump in the $\Omega^- \pi^+$ spectrum but not in the $\Xi^- K^- \pi^+ \pi^+$ spectrum. Perhaps all the experiments are seeing the Ω_c^0 , but statistics are low and further confirmation is desired.


 Ω_c^0 MASS

VALUE (MeV)	EVTs	DOCUMENT ID	TECN	COMMENT
2710 ± 5 OUR AVERAGE				Error includes scale factor of 1.6. See the Ideogram below.
$2705.9 \pm 3.3 \pm 2.0$	10	¹ FRABETTI	93 E687	γ Be $\bar{E}_\gamma = 221$ GeV
$2719.0 \pm 7.0 \pm 2.5$	11	² ALBRECHT	92H ARG	$e^+ e^- \approx 10.6$ GeV
2740 ± 20	3	BIAGI	85B SPEC	Σ^- Be 135 GeV/c

¹ FRABETTI 93 claims a signal of 10.3 ± 3.9 $\Omega^- \pi^+$ events above a background of 5.8 events.

² ALBRECHT 92H claims a signal of 11.5 ± 4.3 $\Xi^- K^- \pi^+ \pi^+$ events. The background is about 5 events.

WEIGHTED AVERAGE
2710±5 (Error scaled by 1.6)

 Ω_c^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $\Xi^- K^- \pi^+ \pi^+$	seen
Γ_2 $\Omega^- \pi^+$	seen
Γ_3 $\Omega^- \pi^- \pi^+ \pi^+$	not seen

 Ω_c^0 BRANCHING RATIOS

$\Gamma(\Xi^- K^- \pi^+ \pi^+)/\Gamma_{\text{total}}$ Γ_1/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	11	ALBRECHT	92H ARG	$e^+ e^- \approx 10.6$ GeV
seen	3	BIAGI	85B SPEC	Σ^- Be 135 GeV/c

$\Gamma(\Omega^- \pi^+)/\Gamma_{\text{total}}$ Γ_2/Γ

VALUE	EVTs	DOCUMENT ID	TECN	COMMENT
seen	10	FRABETTI	93 E687	γ Be $\bar{E}_\gamma = 221$ GeV

$\Gamma(\Xi^- K^- \pi^+ \pi^+)/\Gamma(\Omega^- \pi^+)$ Γ_1/Γ_2

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<2.8	90	FRABETTI	93 E687	γ Be $\bar{E}_\gamma = 221$ GeV

$\Gamma(\Omega^- \pi^- \pi^+ \pi^+)/\Gamma(\Omega^- \pi^+)$ Γ_3/Γ_2

• • • We do not use the following data for averages, fits, limits, etc. • • •

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1.6	90	FRABETTI	93 E687	γ Be $\bar{E}_\gamma = 221$ GeV

 Ω_c^0 REFERENCES

FRABETTI	93	PL B300 190	+Cheung, Cumalat, Dallapiccola+	(FNAL E687 Collab.)
ALBRECHT	92H	PL B288 367	+Cronstroem, Ehrlichmann, Hamacher+	(ARGUS Collab.)
BIAGI	85B	ZPHY C28 175	+	(CERN WA62 Collab.)

Baryon Full Listings

 Λ_b^0

BOTTOM (BEAUTY) BARYON ($B = -1$)

$$\Lambda_b^0 = udb$$

 Λ_b^0

$$I(J^P) = 0(\frac{1}{2}^+) \text{ Status: } ***$$

In the quark model, a Λ_b^0 is an isospin-0 udb state. The lowest Λ_b^0 ought to have $J^P = 1/2^+$. None of $I, J,$ or P have actually been measured.

 Λ_b^0 MASS

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
5641 ± 80 OUR AVERAGE				
5640 ± 50 ± 30	16	¹ ALBAJAR	91E UA1	$p\bar{p}$ 630 GeV
5640 ⁺¹⁰⁰ ₋₂₁₀	52	BARI	91 SFM	$\Lambda_b^0 \rightarrow pD^0\pi^-$
5650 ⁺¹⁵⁰ ₋₂₀₀	90	BARI	91 SFM	$\Lambda_b^0 \rightarrow \Lambda_c^+\pi^+\pi^-\pi^-$
not seen		² ABE	93B CDF	$p\bar{p}$ 1.8 TeV
~ 5750	4	³ ARENTON	86 FMPS	$\Lambda K_S^0 2\pi^+ 2\pi^-$
5425 ⁺¹⁷⁵ ₋₇₅		⁴ BASILE	81 SFM	See BARI 91

- • • We do not use the following data for averages, fits, limits, etc. • • •
 - not seen
 - ~ 5750
 - 5425⁺¹⁷⁵₋₇₅
- ¹ ALBAJAR 91E claims 16 ± 5 events above a background of 9 ± 1 events, a significance of about 5 standard deviations.
- ² ABE 93B states that, based on the signal claimed by ALBAJAR 91E, CDF should have found 30 ± 23 $\Lambda_b^0 \rightarrow J/\psi(1S)\Lambda$ events. Instead, CDF found not more than 2 events.
- ³ The decay of the Λ_b^0 to the final state observed by ARENTON 86 is Cabibbo suppressed, whereas the decay of a Ξ_b^0 to this final state is allowed. ARENTON 86 thus only claims to have observed a baryon which probably has a b quark and has a D^0 among the decay products, not necessarily the Λ_b^0 .
- ⁴ The first claim to have discovered the Λ_b^0 was reported by BASILE 81. In contrast, DRIJARD 82 reported no observation of Λ_b^0 , and this led to some discussion in BASILE 82 and DRIJARD 82b. Further evidence for the Λ_b^0 was again reported by the first authors in BARI 91 (see above) in a second, upgraded experiment where two different Λ_b^0 decay modes were observed.

 Λ_b^0 MEAN LIFE

These are actually measurements of the average lifetime of weakly decaying b baryons weighted by generally unknown production rates, branching fractions, and detection efficiencies. Presumably, the mix is mainly Λ_b^0 , with some Ξ_b^0 and Ξ_b^- .

VALUE (10^{-12} s)	EVTS	DOCUMENT ID	TECN	COMMENT
1.07^{+0.19}_{-0.16} OUR AVERAGE				
1.04 ^{+0.48} _{-0.38} ± 0.10	11	ABREU	93F DLPH	Excess $\Lambda\mu^-$, decay lengths
1.05 ^{+0.23} _{-0.20} ± 0.08	157	AKERS	93 OPAL	Excess $\Lambda\ell^-$, decay lengths
1.12 ^{+0.32} _{-0.29} ± 0.16	101	BUSKULIC	92i ALEP	Excess $\Lambda\ell^-$, impact parameters

 Λ_b^0 DECAY MODES

Mode	Fraction (Γ_i/Γ)
Γ_1 $J/\psi(1S)\Lambda$	seen
Γ_2 $pD^0\pi^-$	seen
Γ_3 $\Lambda_c^+\pi^+\pi^-\pi^-$	seen
Γ_4 $\Lambda K^0 2\pi^+ 2\pi^-$	
Γ_5 $\Lambda\ell^-X$	seen
Γ_6 $\Lambda_c^+\ell^-X$	seen

 Λ_b^0 BRANCHING RATIOS

$$\Gamma(J/\psi(1S)\Lambda)/\Gamma_{\text{total}} \quad \Gamma_1/\Gamma$$

VALUE	DOCUMENT ID	TECN	COMMENT
seen OUR EVALUATION			
0.018 ± 0.011	⁵ ALBAJAR	91E UA1	$J/\psi(1S) \rightarrow \mu^+\mu^-$

⁵ The ALBAJAR 91E value assumes the Λ_b production fraction is 10% of the beauty cross section.

$$\Gamma(pD^0\pi^-)/\Gamma_{\text{total}} \quad \Gamma_2/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	52	BARI	91 SFM	$D^0 \rightarrow K^-\pi^+$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen		BASILE	81 SFM	$D^0 \rightarrow K^-\pi^+$

$$\Gamma(\Lambda_c^+\pi^+\pi^-\pi^-)/\Gamma_{\text{total}} \quad \Gamma_3/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	90	BARI	91 SFM	$\Lambda_c^+ \rightarrow pK^-\pi^+$

$$\Gamma(\Lambda K^0 2\pi^+ 2\pi^-)/\Gamma_{\text{total}} \quad \Gamma_4/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •				
seen	4	⁶ ARENTON	86 FMPS	$\Lambda K_S^0 2\pi^+ 2\pi^-$

⁶ See the footnote to the ARENTON 86 mass value.

$$\Gamma(\Lambda\ell^-X)/\Gamma_{\text{total}} \quad \Gamma_5/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	157	AKERS	93 OPAL	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$
seen	101	BUSKULIC	92i ALEP	Excess of $\Lambda\ell^-$ over $\Lambda\ell^+$

$$\Gamma(\Lambda_c^+\ell^-X)/\Gamma_{\text{total}} \quad \Gamma_6/\Gamma$$

VALUE	EVTS	DOCUMENT ID	TECN	COMMENT
seen	21	BUSKULIC	92E ALEP	$\Lambda_c^+ \rightarrow pK^-\pi^+$

 Λ_b^0 REFERENCES

ABE	93B PR D47 R2639	+Amidei, Anway-Wiese, Apollinari+ (CDF Collab.)
ABREU	93F PL B311 379	+Adam, Adye, Agasi+ (DELPHI Collab.)
AKERS	93 PL B316 435	+Alexander, Allison, Anderson+ (OPAL Collab.)
Also	92E PL B281 394	Acton, Alexander, Allison, Allport+ (OPAL Collab.)
BUSKULIC	92E PL B294 145	+Decamp, Goy, Lees, Minard+ (ALEPH Collab.)
BUSKULIC	92i PL B297 449	+Decamp, Goy, Lees+ (ALEPH Collab.)
Also	92D PL B276 209	Decamp, Deschizeaux, Goy+ (ALEPH Collab.)
ALBAJAR	91E PL B273 540	+Aibrow, Ailkofer, Ankwiat+ (UA1 Collab.)
BARI	91 NC 104A 1787	+Basile, Bruni, Cara Romeo+ (CERN R422 Collab.)
ARENTO	86 NP B274 707	+Chen, Cormeli, Dieterle+ (ARIZ, NDAM, VAND)
BASILE	82 NC 68A 289	+Bonvicini, Romeo+ (CERN R415 Collab.)
DRIJARD	82 PL 108B 361	+ (CERN, CDEF, DORT, HEIDH, LAPP, WARS)
DRIJARD	82B CERN-EP/82-31	+ (CERN, CDEF, DORT, HEIDH, LAPP, WARS)
BASILE	81 LNC 31 97	+Bonvicini, Romeo+ (CERN R415 Collab.)

SEARCHES*

Free Quark Searches 1791
Magnetic Monopole Searches 1793
Supersymmetric Particle Searches 1795
Quark and Lepton Compositeness 1805
Other Stable Particle Searches 1811

Notes in the Search Listings

Note on Quark Searches 1791
Note on Magnetic Monopole Searches 1793
Note on Supersymmetry 1795
Note on Searches for Quark and Lepton Compositeness . 1805
Note on Other Stable Particle Searches 1811

* See the Boson Full Listings for searches for Higgs bosons, other heavy bosons, and axions and other very light bosons, the Lepton Full Listings for searches for heavy leptons and for neutrino mixing, and the Meson Full Listings for searches for top and fourth-generation hadrons.

See key on page 1343

Searches Full Listings

Free Quark Searches

SEARCHES FOR FREE QUARKS, MONOPOLES, SUPERSYMMETRY, COMPOSITENESS, etc.

Free Quark Searches

NOTE ON QUARK SEARCHES

The basis for much of the theory of particle scattering and hadron spectroscopy is the construction of the hadrons from a set of fractionally charged constituents (quarks). A central but unproven hypothesis of this theory, Quantum Chromodynamics, is that quarks cannot be observed as free particles but are confined to mesons and baryons.

Experiments show that it is at best difficult to "unglue" quarks. Accelerator searches at increasing energies have produced no evidence for free quarks, while only a few cosmic-ray and matter searches have produced uncorroborated events.

This compilation is only a guide to the literature, since the quoted experimental limits are often only indicative. Reviews can be found in Refs. 1-3.

References

1. P.F. Smith, Ann. Rev. Nucl. and Part. Sci. **39**, 73 (1989).
2. L. Lyons, Phys. Reports **129**, 225 (1985).
3. M. Marinelli and G. Morpurgo, Phys. Reports **85**, 161 (1982).

Quark Production Cross Section — Accelerator Searches

X-SECT (cm ²)	CHG (e/3)	MASS (GeV)	ENERGY (GeV)	BEAM	EVTS	DOCUMENT ID	TECN
<2.E-35	+2	250	1800	p \bar{p}	0	¹ ABE 92J	CDF
<1.E-35	+4	250	1800	p \bar{p}	0	¹ ABE 92J	CDF
<3.8E-28		14.5A	28Si-Pb		0	² HE 91	PLAS
<3.2E-28		14.5A	28Si-Cu		0	² HE 91	PLAS
<1.E-40	$\pm 1,2$	<10	p, $\nu, \bar{\nu}$		0	BERGSM 84B	CHRM
<1.E-36	$\pm 1,2$	<9	200 μ		0	AUBERT 83C	SPEC
<2.E-10	$\pm 2,4$	1-3	200 p		0	³ BUSSIERE 80	CNTR
<5.E-38	+1,2	>5	300 p		0	^{4,5} STEVENSON 79	CNTR
<1.E-33	± 1	<20	52 p \bar{p}		0	BASILE 78	SPEC
<9.E-39	$\pm 1,2$	<6	400 p		0	⁴ ANTREASIAN 77	SPEC
<8.E-35	+1,2	<20	52 p \bar{p}		0	⁶ FABJAN 75	CNTR
<5.E-38	-1,2	4-9	200 p		0	NASH 74	CNTR
<1.E-32	+2,4	4-24	52 p \bar{p}		0	ALPER 73	SPEC
<5.E-31	+1,2,4	<12	300 p		0	LEIPUNER 73	CNTR
<6.E-34	$\pm 1,2$	<13	52 p \bar{p}		0	BOTT 72	CNTR
<1.E-36	-4	4	70 p		0	ANTIPOV 71	CNTR
<1.E-35	$\pm 1,2$	2	28 p		0	⁷ ALLABY 69B	CNTR
<4.E-37	-2	<5	70 p		0	³ ANTIPOV 69	CNTR
<3.E-37	-1,2	2-5	70 p		0	⁷ ANTIPOV 69B	CNTR
<1.E-35	+1,2	<7	30 p		0	DORFAN 65	CNTR
<2.E-35	-2	<2.5-5	30 p		0	⁸ FRANZINI 65B	CNTR
<5.E-35	+1,2	<2.2	21 p		0	BINGHAM 64	HLBC
<1.E-32	+1,2	<4.0	28 p		0	BLUM 64	HBC
<1.E-35	+1,2	<2.5	31 p		0	⁸ HAGOPIAN 64	HBC
<1.E-34	+1	<2	28 p		0	LEIPUNER 64	CNTR
<1.E-33	+1,2	<2.4	24 p		0	MORRISON 64	HBC

¹ ABE 92J flux limits decrease as the mass increases from 50 to 500 GeV.

² HE 91 limits are for charges of the form $N \pm 1/3$ from 23/3 to 38/3.

³ Hadronic or leptonic quarks.

⁴ Cross section cm²/GeV².

⁵ 3×10^{-5} < lifetime < 1×10^{-3} s.

⁶ Includes BOTT 72 results.

⁷ Assumes isotropic cm production.

⁸ Cross section inferred from flux.

Quark Differential Production Cross Section — Accelerator Searches

X-SECT (cm ² sr ⁻¹ GeV ⁻¹)	CHG (e/3)	MASS (GeV)	ENERGY (GeV)	BEAM	EVTS	DOCUMENT ID	TECN
<4.E-36	-2,4	1.5-6	70	p	0	BALDIN 76	CNTR
<2.E-33	± 4	5-20	52	p \bar{p}	0	ALBROW 75	SPEC
<5.E-34	<7	7-15	44	p \bar{p}	0	JOVANO... 75	CNTR
<5.E-35			20	γ	0	⁹ GALIK 74	CNTR
<9.E-35	-1,2		200	p	0	NASH 74	CNTR
<4.E-36	-4	2.3-2.7	70	p	0	ANTIPOV 71	CNTR
<3.E-35	$\pm 1,2$	<2.7	27	p	0	ALLABY 69B	CNTR
<7.E-38	-1,2	<2.5	70	p	0	ANTIPOV 69B	CNTR

⁹ Cross section in cm²/sr/equivalent quanta.

Quark Flux — Accelerator Searches

The definition of FLUX depends on the experiment

- (a) is the ratio of measured free quarks to predicted free quarks if there is no "confinement."
- (b) is the probability of fractional charge on nuclear fragments.
- (c) is the 90%CL upper limit on fractionally-charged particles produced per interaction.
- (d) is quarks per collision.
- (e) is inclusive quark-production cross-section ratio to $\sigma(e^+e^- \rightarrow \mu^+\mu^-)$.
- (f) is quark flux per charged particle.
- (g) is the flux per ν -event.
- (h) is quark yield per π^- yield.
- (i) is 2-body exclusive quark-production cross-section ratio to $\sigma(e^+e^- \rightarrow \mu^+\mu^-)$.

FLUX	CHG (e/3)	MASS (GeV)	ENERGY (GeV)	BEAM	EVTS	DOCUMENT ID	TECN
<2.E-3	e	+1	5-40 88-94	e ⁺ e ⁻	0	¹⁰ BUSKULIC 93C	ALEP
<6.E-4	e	+2	5-30 88-94	e ⁺ e ⁻	0	¹⁰ BUSKULIC 93C	ALEP
<1.2E-3	e	+4	15-40 88-94	e ⁺ e ⁻	0	¹⁰ BUSKULIC 93C	ALEP
<3.6E-4	i	+4	5.0-10.2 88-94	e ⁺ e ⁻	0	BUSKULIC 93C	ALEP
<3.6E-4	i	+4	16.5-26.0 88-94	e ⁺ e ⁻	0	BUSKULIC 93C	ALEP
<6.9E-4	i	+4	26.0-33.3 88-94	e ⁺ e ⁻	0	BUSKULIC 93C	ALEP
<9.1E-4	i	+4	33.3-38.6 88-94	e ⁺ e ⁻	0	BUSKULIC 93C	ALEP
<1.1E-3	i	+4	38.6-44.9 88-94	e ⁺ e ⁻	0	BUSKULIC 93C	ALEP
	b	4,5,7,8	2.1A	¹⁶ O	0,2,0,6	¹¹ GHOSH 92	EMUL
<6.4E-5	g	1		$\nu, \bar{\nu}$	1	¹² BASILE 91	CNTR
<3.7E-5	g	2		$\nu, \bar{\nu}$	0	¹² BASILE 91	CNTR
<3.9E-5	g	1		$\nu, \bar{\nu}$	1	¹³ BASILE 91	CNTR
<2.8E-5	g	2		$\nu, \bar{\nu}$	0	¹³ BASILE 91	CNTR
<1.9E-4	c		14.5A	28Si-Pb	0	¹⁴ HE 91	PLAS
<3.9E-4	c		14.5A	28Si-Cu	0	¹⁴ HE 91	PLAS
<1.E-9	c	$\pm 1,2,4$	14.5A	¹⁶ O-Ar	0	MATIS 91	MDRP
<5.1E-10	c	$\pm 1,2,4$	14.5A	¹⁶ O-Hg	0	MATIS 91	MDRP
<8.1E-9	c	$\pm 1,2,4$	14.5A	Si-Hg	0	MATIS 91	MDRP
<1.7E-6	c	$\pm 1,2,4$	60A	¹⁶ O-Hg	0	MATIS 91	MDRP
<3.5E-7	c	$\pm 1,2,4$	200A	¹⁶ O-Hg	0	MATIS 91	MDRP
<1.3E-6	c	$\pm 1,2,4$	200A	S-Hg	0	MATIS 91	MDRP
<5E-2	e	2	19-27 52-60	e ⁺ e ⁻	0	ADACHI 90C	TOPZ
<5E-2	e	4	<24 52-60	e ⁺ e ⁻	0	ADACHI 90C	TOPZ
<1.E-4	e	+2	<3.5	10 e ⁺ e ⁻	0	BOWCOCK 89B	CLEO
<1.E-6	d	$\pm 1,2$	60	¹⁶ O-Hg	0	CALLOWAY 89	MDRP
<3.5E-7	d	$\pm 1,2$	200	¹⁶ O-Hg	0	CALLOWAY 89	MDRP
<1.3E-6	d	$\pm 1,2$	200	S-Hg	0	CALLOWAY 89	MDRP
<1.2E-10	d	± 1	1	800 p-Hg	0	MATIS 89	MDRP
<1.1E-10	d	± 2	1	800 p-Hg	0	MATIS 89	MDRP
<1.2E-10	d	± 1	1	800 p-N ₂	0	MATIS 89	MDRP
<7.7E-11	d	± 2	1	800 p-N ₂	0	MATIS 89	MDRP
<6.E-9	h	-5	0.9-2.3	12 p	0	NAKAMURA 89	SPEC
<5.E-5	g	1,2	<0.5	$\nu, \bar{\nu}d$	0	ALLASIA 88	BECB
<3.E-4	b	See note	14.5	¹⁶ O-Pb	0	¹⁵ HOFFMANN 88	PLAS
<2.E-4	b	See note	200	¹⁶ O-Pb	0	¹⁶ HOFFMANN 88	PLAS
<2.E-4	a	$\pm 1,2$	<300	320 p \bar{p}	0	LYONS 87	MLEV
<1.E-9	c	$\pm 1,2,4,5$	14.5	¹⁶ O-Hg	0	SHAW 87	MDRP
<3.E-3	d	-1,2,3,4,6	<5	2 Si-Si	0	¹⁷ ABACHI 86C	CNTR
<1.E-4	e	$\pm 1,2,4$	<4	10 e ⁺ e ⁻	0	ALBRECHT 85G	ARG
<6.E-5	b	$\pm 1,2$	1	540 p \bar{p}	0	BANNER 85	UA2
<5.E-3	e	-4	1-8	29 e ⁺ e ⁻	0	AIHARA 84	TPC
<1.E-2	e	$\pm 1,2$	1-13	29 e ⁺ e ⁻	0	AIHARA 84B	TPC
<2.E-4	b	± 1	72	40 Ar	0	¹⁸ BARWICK 84	CNTR
<1.E-4	e	± 2	<0.4	1.4 e ⁺ e ⁻	0	BONDAR 84	OLYA
<5.E-1	e	$\pm 1,2$	<13	29 e ⁺ e ⁻	0	GURYAN 84	CNTR
<3.E-3	b	$\pm 1,2$	<2	540 p \bar{p}	0	BANNER 83	CNTR
<1.E-4	b	$\pm 1,2$	106	⁵⁶ Fe	0	LINDGREN 83	CNTR
<3.E-3	b	> ± 0.1	74	40 Ar	0	¹⁸ PRICE 83	PLAS
<1.E-2	e	$\pm 1,2$	<14	29 e ⁺ e ⁻	0	MARINI 82B	CNTR
<8.E-2	e	$\pm 1,2$	<12	29 e ⁺ e ⁻	0	ROSS 82	CNTR
<3.E-4	e	± 2	1.8-2	7 e ⁺ e ⁻	0	WEISS 81	MRK2

Searches Full Listings

Free Quark Searches

<5.E-2	e	+1,2,4,5	2-12	27	e^+e^-	0	BARTEL	80	JADE
<2.E-5	g	1,2			ν	0	12,13 BASILE	80	CNTR
<3.E-10	f	$\pm 2,4$	1-3	200	p	0	19 BOZZOLI	79	CNTR
<6.E-11	f	± 1	<21	52	pp	0	BASILE	78	SPEC
<5.E-3	g				ν_μ	0	BASILE	78b	CNTR
<2.E-9	f	± 1	<26	62	pp	0	BASILE	77	SPEC
<7.E-10	f	+1,2	<20	52	p	0	20 FABJAN	75	CNTR
		+1,2	>4.5		γ	0	12,13 GALIK	74	CNTR
		+1,2	>1.5	12	e^-	0	12,13 BELLAMY	68	CNTR
		+1,2	>0.9		γ	0	13 BATHOW	67	CNTR
		+1,2	>0.9	6	γ	0	13 FOSS	67	CNTR

- ¹⁰ BUSKULIC 93c limits for inclusive quark production are more conservative if the ALEPH hadronic fragmentation function is assumed.
- ¹¹ GHOSH 92 reports measurement of spallation fragment charge based on ionization in emulsion. Out of 650 measured tracks, 2 were consistent with charge $5e/3$, and 4 with $7e/3$.
- ¹² Hadronic quark.
- ¹³ Leptonic quark.
- ¹⁴ HE 91 limits are for charges of the form $N \pm 1/3$ from 23/3 to 38/3, and correspond to cross-section limits of $380\mu\text{b}$ (Pb) and $320\mu\text{b}$ (Cu).
- ¹⁵ The limits apply to projectile fragment charges of 17, 19, 20, 22, 23 in units of $e/3$.
- ¹⁶ The limits apply to projectile fragment charges of 16, 17, 19, 20, 22, 23 in units of $e/3$.
- ¹⁷ Flux limits and mass range depend on charge.
- ¹⁸ Bound to nuclei.
- ¹⁹ Quark lifetimes $> 1 \times 10^{-8}$ s.
- ²⁰ One candidate $m < 0.17$ GeV.

Quark Flux — Cosmic Ray Searches

Shielding values followed with an asterisk indicate altitude in km. Shielding values not followed with an asterisk indicate sea level in kg/cm^2 .

FLUX ($\text{cm}^{-2}\text{sr}^{-1}\text{s}^{-1}$)	CHG (e/3)	MASS (GeV)	SHIELDING	EVTS	DOCUMENT ID	TECN
<2.1E-15	± 1			0	MORI 91 KAM2	
<2.3E-15	± 2			0	MORI 91 KAM2	
<2.E-10	$\pm 1,2$		0.3	0	WADA 88 CNTR	
	± 4		0.3	12	21 WADA 88 CNTR	
	± 4		0.3	9	22 WADA 86 CNTR	
<1.E-12	$\pm 2,3/2$		-70.	0	23 KAWAGOE 84b PLAS	
<9.E-10	$\pm 1,2$		0.3	0	WADA 84b CNTR	
<4.E-9	± 4		0.3	7	WADA 84b CNTR	
<2.E-12	$\pm 1,2,3$		-0.3*	0	MASHIMO 83 CNTR	
<3.E-10	$\pm 1,2$		0.3	0	MARINI 82 CNTR	
<2.E-11	$\pm 1,2$			0	MASHIMO 82 CNTR	
<8.E-10	$\pm 1,2$		0.3	0	23 NAPOLITANO 82 CNTR	
				3	24 YOCC 78 CNTR	
<1.E-9				0	25 BRIATORE 76 ELEC	
<2.E-11	+1			0	26 HAZEN 75 CC	
<2.E-10	+1,2			0	KRISOR 75 CNTR	
<1.E-7	+1,2			0	26,27 CLARK 74b CC	
<3.E-10	+1	>20		0	KIFUNE 74 CNTR	
<8.E-11	+1			0	26 ASHTON 73 CNTR	
<2.E-8	+1,2			0	HICKS 73b CNTR	
<5.E-10	+4		2.8*	0	BEAUCHAMP 72 CNTR	
<1.E-10	+1,2			0	26 BOHM 72b CNTR	
<1.E-10	+1,2		2.8*	0	COX 72 ELEC	
<3.E-10	+2			0	CROUCH 72 CNTR	
<3.E-8			7	0	25 DARDO 72 CNTR	
<4.E-9	+1			0	26 EVANS 72 CC	
<2.E-9		>10		0	25 TONWAR 72 CNTR	
<2.E-10	+1		2.8*	0	CHIN 71 CNTR	
<3.E-10	+1,2			0	26 CLARK 71b CC	
<1.E-10	+1,2			0	26 HAZEN 71 CC	
<5.E-10	+1,2		3.5*	0	BOSIA 70 CNTR	
	+1,2	<6.5		1	26 CHU 70 HLBC	
<2.E-9	+1			0	FAISSNER 70b CNTR	
<2.E-10	+1,2		0.8*	0	KRIDER 70 CNTR	
<5.E-11	+2			4	CAIRNS 69 CC	
<8.E-10	+1,2	<10		0	FUKUSHIMA 69 CNTR	
	+2			1	26,28 MCCUSKER 69 CC	
<1.E-10		>5	1.7,3.6	0	25 BJORNBOE 68 CNTR	
<1.E-8	$\pm 1,2,4$		6.3,2*	0	23 BRIATORE 68 CNTR	
<3.E-8		>2		0	FRANZINI 68 CNTR	
<9.E-11	$\pm 1,2$			0	GARMIRE 68 CNTR	
<4.E-10	± 1			0	HANAYAMA 68 CNTR	
<3.E-8		>15		0	KASHA 68 OSPK	
<2.E-10	+2			0	KASHA 68b CNTR	
<2.E-10	+4			0	KASHA 68c CNTR	
<2.E-10	+2		6	0	BARTON 67 CNTR	
<2.E-7	+4		0.008,0.5*	0	BUHLER 67 CNTR	
<5.E-10	1,2		0.008,0.5*	0	BUHLER 67b CNTR	
<4.E-10	+1,2			0	GOMEZ 67 CNTR	
<2.E-9	+2			0	KASHA 67 CNTR	
<2.E-10	+2		220	0	BARTON 66 CNTR	

<2.E-9	+1,2		0.5*	0	BUHLER	66	CNTR
<3.E-9	+1,2			0	KASHA	66	CNTR
<2.E-9	+1,2			0	LAMB	66	CNTR
<2.E-8	+1,2	>7	2.8*	0	DELISE	65	CNTR
<5.E-8	+2	>2.5	0.5*	0	MASSAM	65	CNTR
<2.E-8	+1		2.5*	0	BOWEN	64	CNTR
<2.E-7	+1		0.8	0	SUNYAR	64	CNTR

- ²¹ Distribution in celestial sphere was described as anisotropic.
- ²² With telescope axis at zenith angle 40° to the south.
- ²³ Leptonic quarks.
- ²⁴ Lifetime $> 10^{-8}$ s; charge $\pm 0.70, 0.68, 0.42$; and mass $> 4.4, 4.8$, and 20 GeV, respectively.
- ²⁵ Time delayed air shower search.
- ²⁶ Prompt air shower search.
- ²⁷ Also $e/4$ and $e/6$ charges.
- ²⁸ No events in subsequent experiments.

Quark Density — Matter Searches

For a recent review, see SMITH 89.

QUARKS/ NUCLEON	CHG (e/3)	MASS (GeV)	MATERIAL/METHOD	EVTS	DOCUMENT ID
<8.E-22	+2		Si/infrared photoionization	0	PERERA 93
<5.E-27	$\pm 1,2$		sea water/levitation	0	HOMER 92
<4.E-20	$\pm 1,2$		meteorites/mag. levitation	0	JONES 89
<1.E-19	$\pm 1,2$		various/spectrometer	0	MILNER 87
<5.E-22	$\pm 1,2$		W/levitation	0	SMITH 87
<3.E-20	+1,2		org liq/droplet tower	0	VANPOLEN 87
<6.E-20	-1,2		org liq/droplet tower	0	VANPOLEN 87
<3.E-21	± 1		Hg drops-untreated	0	SAVAGE 86
<3.E-22	$\pm 1,2$		levitated niobium	0	SMITH 86
<2.E-26	$\pm 1,2$		⁴ He/levitation	0	SMITH 86b
<2.E-20	> ± 1	0.2-250	niobium+tungs/ion	0	MILNER 85
<1.E-21	± 1		levitated niobium	0	SMITH 85
	+1,2	<100	niobium/mass spec	0	KUTSCHERA 84
<5.E-22			levitated steel	0	MARINELLI 84
<9.E-20	$\pm <13$		water/oil drop	0	JOYCE 83
<2.E-21	> $\pm 1/2$		levitated steel	0	LIEBOWITZ 83
<1.E-19	$\pm 1,2$		photo ion spec	0	VANDESTEEG 83
<2.E-20			mercury/oil drop	0	²⁹ HODGES 81
1.E-20	+1		levitated niobium	4	³⁰ LARUE 81
1.E-20	-1		levitated niobium	4	³⁰ LARUE 81
<1.E-21			levitated steel	0	MARINELLI 80b
<6.E-16			helium/mass spec	0	BOYD 79
1.E-20	+1		levitated niobium	2	³⁰ LARUE 79
<4.E-28			earth+/ion beam	0	OGOROD... 79
<5.E-15	+1		tungs./mass spec	0	BOYD 78
<5.E-16	+3	<1.7	hydrogen/mass spec	0	BOYD 78b
<1.E-21	$\pm 2,4$		water/ion beam	0	LUND 78
<6.E-15	>1/2		levitated tungsten	0	PUTT 78
<1.E-22			metals/mass spec	0	SCHIFFER 78
<5.E-15			levitated tungsten ox	0	BLAND 77
<3.E-21			levitated iron	0	GALLINARO 77
2.E-21	-1		levitated niobium	1	³⁰ LARUE 77
4.E-21	+1		levitated niobium	2	³⁰ LARUE 77
<1.E-13	+3	<7.7	hydrogen/mass spec	0	MULLER 77
<5.E-27			water+/ion beam	0	OGOROD... 77
<1.E-21			lunar+/ion spec	0	STEVENS 76
<1.E-15	+1	<60	oxygen+/ion spec	0	ELBERT 70
<5.E-19			levitated graphite	0	MORPURGO 70
<5.E-23			water+/atom beam	0	COOK 69
<1.E-17	$\pm 1,2$		levitated graphite	0	BRAGINSK 68
<1.E-17			water+/uv spec	0	RANK 68
<3.E-19	± 1		levitated iron	0	STOVER 67
<1.E-10			sun/uv spec	0	³¹ BENNETT 66
<1.E-17	+1,2		meteorites+/ion beam	0	CHUPKA 66
<1.E-16	± 1		levitated graphite	0	GALLINARO 66
<1.E-22			argon/electrometer	0	HILLAS 59
	-2		levitated oil	0	MILLIKAN 10

- ²⁹ Also set limits for $Q = \pm e/6$.
- ³⁰ Note that in PHILLIPS 88 these authors report a subtle magnetic effect which could account for the apparent fractional charges.
- ³¹ Limit inferred by JONES 77b.

REFERENCES FOR Free Quark Searches

BUSKULIC 93C	PL B303 198	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
PERERA 93	PRL 70 1053	+Betarbet, Byung-sung, Coon	(PITT)
ABE 92J	PR D46 R1889	+Amidei, Anway-Wells+	(CDF Collab.)
GHOSH 92	NC 105A 99	+Roy, Ghosh, Ghosh, Basu	(JADA, BANGB)
HOMER 92	ZPHY C55 549	+Smith, Lewin, Robertson+	(RAL, SHMP, LOQM)
BASILE 91	NC 104A 405	+Berbers, Cara Romeo+	(BGNA, INFN, CERN, PLRM+)
HE 91	PR C44 1672	+Price	(UCB)
MATIS 91	NP A525 513c	+Pugh, Alba, Bland, Calloway+	(LBL, SFSU, UCI, LNL)
MORI 91	PR D43 2843	+Oyama, Suzuki, Takahashi+	(Kamikande II Collab.)
ADACHI 90C	PL B244 352	+Aihara, Doser, Enomoto+	(TOPAZ Collab.)
BOWCOCK 89B	PR D40 263	+Kinoshita, Mauskopf, Pipkin+	(CLEO Collab.)
CALLOWAY 89	PL B232 549	+Alba, Bland, Dickson, Hodges+	(SFSU, UCI, LBL, LNL)
JONES 89	ZPHY C43 349	+Smith, Homer, Lewin, Walford	(LOIC, RAL)
MATIS 89	PR D39 1851	+Pugh, Bland, Calloway+	(LBL, SFSU, UCI, FNAL, LNL)
NAKAMURA 89	PR D39 1261	+Kobayashi, Konaka, Imai, Masaie+	(KYOT, TMTC)
SMITH 89	ARNPS 39 73		(RAL)
ALLASIA 88	PR D37 219	+Angelini, Baldini+	(WA25 Collab.)

See key on page 1343

Searches Full Listings

Free Quark Searches, Magnetic Monopole Searches

HOFFMANN	88	PL B200 583	+Brechtmann, Heinrich, Benton (SIEG, USF)
PHILLIPS	88	NIM A264 125	+Fairbank, Navarro (STAN)
WADA	87	NC 12 229	+Yamashita, Yamamoto (OKAY)
LYONS	87	ZPHY C36 363	+Smith, Homer, Lewin, Walford+ (OXF, RAL, LOIC)
MILNER	87	PR D36 37	+Cooper, Chang, Wilson, Labrenz, McKeown (CIT)
SHAW	87	PR D36 3533	+Matis, Pugh, Slansky+ (UCI, LBL, LANL, SFSU)
SMITH	87	PL B197 447	+Homer, Lewin, Walford, Jones (RAL, LOIC)
VANPOLEN	87	PR D36 1983	+Hagstrom, Hirsch (ANL, LBL)
ABACHI	86C	PR D33 2733	+Shor, Barasch, Carroll+ (UCLA, LBL, UCSD)
SAVAGE	86	PL 167B 481	+Bland, Hodges, Huntington, Joyce+ (SFSU)
SMITH	86	B171 129	+Homer, Lewin, Walford, Jones (RAL, LOIC)
SMITH	86B	PL B181 407	+Homer, Lewin, Walford, Jones (RAL, LOIC)
WADA	86	NC 9C 358	(OKAY)
ALBRECHT	85G	PL 156B 134	+Binder, Harder, Hasemann+ (ARGUS Collab.)
BANNER	85	PL 156B 129	+Bloch, Borer, Borghini+ (UA2 Collab.)
MILNER	85	PRL 54 1472	+Cooper, Chang, Wilson, Labrenz, McKeown (CIT)
SMITH	85	PL 153B 188	+Homer, Lewin, Walford, Jones (RAL, LOIC)
AIHARA	84	PRL 52 1568	+Alston-Garnjost, Badtke, Bakker+ (TPC Collab.)
AIHARA	84B	PRL 52 2332	+Alston-Garnjost, Badtke, Bakker+ (TPC Collab.)
BARWICK	84	PR D30 691	+Musser, Stevenson (UCB)
BERGSMÄ	84B	ZPHY C24 217	+Allaby, Abt, Gemanov+ (CHARM Collab.)
BONDAR	84	JETPL 40 1265	+Kuradze, Leichuk, Panin, Sidorov+ (NOVO)
		Translated from ZETFP 40 440.	
GURYN	84	PL 139B 313	+Farker, Fries+ (FRAS, LBL, NWES, STAN, HAWA)
KAWAGOE	84B	PR D25 1640	+Mashimo, Nakamura, Nozaki, Orito (TOKY)
KUTSCHERA	84	PR D29 791	+Schiffer, Frekers+ (ANL, FNAL)
MARINELLI	84	PL 137B 439	+Morpurgo (GENO)
WADA	84B	UNC 40 329	+Yamashita, Yamamoto (OKAY)
AUBERT	83C	PL 133B 461	+Bassompierre, Becks, Best+ (EMC Collab.)
BANNER	83	PL 121B 187	+Bloch, Bonaudi, Borer+ (UA2 Collab.)
JOYCE	83	PRL 51 731	+Abrams, Bland, Johnson, Lindgren+ (SFSU)
LIEBOWITZ	83	PRL 50 1640	+Bieder, Ziock (VIRG)
LINDGREN	83	PRL 51 1621	+Joyce+ (SFSU, UCR, UCI, SLAC, LBL, LANL)
MASHIMO	83	PL 128B 327	+Orito, Kawagoe, Nakamura, Nozaki (ICEPP)
PRICE	83	PRL 50 566	+Tinknell, Tarle, Ahlen, Frankel+ (UCB)
VANDESTEER	83	PL 50 1234	+Jongbloets, Wyder (NIJM)
MARINI	82	PR D26 1777	+Peruzzi, Piccolo+ (FRAS, LBL, NWES, STAN, HAWA)
MARINI	82B	PRL 48 1649	+Peruzzi, Piccolo+ (FRAS, LBL, NWES, STAN, HAWA)
MASHIMO	82	JPSJ 51 3087	+Kawagoe, Koshiba (INUS)
NAPOLITANO	82	PR D25 1640	+Beska+ (STAN, FRAS, LBL, NWES, HAWA)
ROSS	82	PL 118B 199	+Rones, Besset+ (FRAS, LBL, NWES, STAN, HAWA)
HODGES	81	PRL 47 1651	+Abrams, Baden, Bland, Joyce+ (UCR, SFSU)
LARUE	81	PRL 46 967	+Phillips, Fairbank (STAN)
WEISS	81	PL 101B 439	+Abrams, Alam, Blocker+ (SLAC, LBL, UCB)
BARTEL	80	ZPHY C6 295	+Canzler, Lords, Drumm+ (JADE Collab.)
BASILE	80	LNC 29 251	+Berbiers+ (BGNA, CERN, FRAS, ROMA, BARI)
BUSSIERE	80	NP B174 1	+Giacomelli, Lesauoy+ (BGNA, SACL, LAPP)
MARINELLI	80B	PL 94B 427	+Morpurgo (GENO)
		Also	Marinelli, Morpurgo (GENO)
BOYD	79	PRL 43 1288	+Blatt, Donoghue, Dries, Hausman, Suiter (OSU)
BOZZOLI	79	NP B159 363	+Bussiere, Giacomelli+ (BGNA, LAPP, SACL, CERN)
LARUE	79	PRL 42 142	+Fairbank, Phillips (STAN)
		Also	Larue, Fairbank, Phillips (STAN)
OGOROD...	79	JETP 49 953	Ogorodnikov, Samoilov, Solntsev (KIAE)
		Translated from ZETF 76 1881.	
STEVENSON	79	PR D20 82	(LBL)
BASILE	78	NC 45A 171	+Cara-Romeo, Cifarelli, Contin+ (CERN, BGNA)
BASILE	78B	NC 45A 281	+Cara-Romeo, Cifarelli, Contin+ (CERN, BGNA)
BOYD	78	PRL 40 216	+Elmore, Melissinos, Sugarbaker (ROCH)
BOYD	78B	PL 72B 484	+Elmore, Nitz, Olsen, Sugarbaker, Warren+ (ROCH)
LUND	78	RA 25 75	+Brandt, Fares (MARB)
PUTT	78	PR D17 1466	+Vofsi (AUCK)
SCHIFFER	78	PR D17 2241	+Renner, Gemmell, Mooring (CHIC, ANL)
YOCK	78	PR D18 641	(AUCK)
ANTREASIAN	77	PRL 39 513	+Coconi, Cronin, Frisch+ (EFI, PRIN)
BASILE	77	NC 40A 41	+Romeo, Cifarelli, Giusti+ (CERN, BGNA)
BLAND	77	PRL 39 369	+Bocobo, Eubank, Royer (SFSU)
GALLINARO	77	PRL 38 1255	+Marinelli, Morpurgo (GENO)
JONES	77B	RMP 69 717	
LARUE	77	PR 38 1011	+Fairbank, Hebard (STAN)
MULLER	77	Science 521	+Alvarez, Holley, Stephenson (LBL)
OGOROD...	77	JETP 45 857	Ogorodnikov, Samoilov, Solntsev (KIAE)
		Translated from ZETF 72 1633.	
BALDIN	76	SJNP 22 264	+Vertogradov, Vishnevsky, Grishkevich+ (JINR)
		Translated from YAF 22 512.	
BRIATORE	76	NC 31A 553	+Dardo, Piazzoli, Mannocchi+ (LCGT, FRAS, FREIB)
STEVENS	76	PR D14 716	+Schiffer, Chupka (ANL)
ALBROW	75	NP B97 189	+Barber+ (CERN, DARE, FOM, LANC, MCHS, UTRÉ)
FABJAN	75	NP B101 349	+Gruhn, Peak, Sauli, Caldwell+ (CERN, MPIM)
HAZEN	75	NP B95 189	+Hodson, Winterstein, Green, Kass+ (MICH, LEED)
JOVANOV...	75	PL 56B 105	+Jovanovich+ (MANI, AACH, CERN, GENO, HARV+)
KRISOR	75	NC 27A 132	(AACH3)
CLARK	74B	PR D10 2721	+Finn, Hansen, Smith (LL)
GALIK	74	PR D9 1856	+Jordan, Richter, Seppi, Siemann+ (SLAC, FNAL)
KIFUNE	74	JPSJ 36 629	+Hieda, Kurokawa, Tsunemoto+ (TOKY, KEK)
NASH	74	PRL 32 858	+Yamanouchi, Nease, Scull (FNAL, CERN, NYU)
ALPER	73	PL 46B 265	+ (CERN, LIVP, LUND, BOHR, RHEL, STOH, BERG+)
ASHTON	73	JPA 6 577	+Cooper, Panvaresh, Saleh (DURH)
HICKS	73B	NC 14A 65	+Flint, Standil (MANI)
LEIPUNER	73	PRL 31 1226	+Larsen, Sessoms, Smith, Williams+ (BNL, YALE)
BEAUCHAMP	72	PR D6 1211	+Bowen, Cox, Kalbach (ARIZ)
BOHM	72B	PRL 28 326	+Diamond, Faisner, Fasold, Krisor+ (AACH)
BOXT	72	PL 40B 693	+Caldwell, Fabjan, Gruhn, Peak+ (CERN, MPIM)
COX	72	PR D6 1203	+Beauchamp, Bowen, Kalbach (ARIZ)
CROUCH	72	PR D5 2667	+Mori, Smith (CASE)
DARDO	72	NC 9A 319	+Navarra, Penengo, Sitte (TORI)
EVANS	72	PRSE A70 143	+Fancey, Muir, Watson (EDIN, LEED)
TOINPAR	72	JPA 5 569	+Naranan, Sreerantan (TAT)
ANTIPOV	71	NP B27 374	+Kachanov, Kutjlin, Landsberg, Lebedev+ (SERP)
CHIN	71	NC 2A 419	+Hanayama, Hara, Higashi, Tsuji (OSAK)
CLARK	71B	PRL 27 51	+Ernst, Finn, Griffin, Hansen, Smith+ (LL, LBL)
HAZEN	71	PRL 26 582	(MICH)
BOSIA	70	NC 66A 167	+Briatore (TORI)
CHU	70	PRL 24 917	+Kim, Beam, Kwak (OSU, ROSE, KANS)
		Also	Allison, Derrick, Hunt, Simpson, Voyvodic (ANL)
ELBERT	70	NP B20 217	+Erwin, Heide, Nielsen, Petriak, Weinberg (NISC)
FAISSNER	70B	PRL 24 1357	+Holder, Krisor, Mason, Sawaf, Umbach (AACH3)
KRIDER	70	PR D1 835	+Bowen, Kalbach (ARIZ)
MORPURGO	70	NIM 79 95	+Gallinaro, Palmieri (GENO)
ALLABY	69B	NC 64A 75	+Bianchini, Diddens, Dobinson, Hartung+ (CERN)
ANTIPOV	69	PL 29B 245	+Karpov, Khromov, Landsberg, Lapshin+ (SERP)
ANTIPOV	69B	PL 30B 576	+Bolotov, Devishov, Devishova, Isakov+ (SERP)
CAIRNS	69	PR 18B 194	+McKachan, Peak, Woodcote (SYDN)
COOK	69	PR 18B 2092	+Depasquali, Frauenfelder, Peacock+ (ILL)
FUKUSHIMA	69	PR 17B 2058	+Kifune, Kondo, Koshiba+ (TOKY)
MCCUSKER	69	PRL 23 658	+Cairns (SYDN)
BELLAMY	69	PR 16B 1391	+Hofstadter, Lakin, Peri, Toner (STAN, SLAC)
BJORNBOE	68	NC B53 241	+Damgard, Hansen+ (BOHR, TATA, BERN, BERG)
BRAGINSK	68	JETP 47 51	+Zelodovich, Martynov, Migulin (MOSU)
		Translated from ZETF 54 91.	
BRIATORE	68	NC 57A 850	+Castagnoli, Bollini, Massam+ (TORI, CERN, BGNA)
FRANZINI	68	PRL 21 1013	+Shulman (COLU)
GARMIRE	68	PR 166 166	+Bong, Sreerantan (MIT)
HANAYAMA	68	CJP 46 5734	+Hara, Higashi, Kitamura, Miono+ (OSAK)
KASHA	68	PR 172 1297	+Stefanski (BNL, YALE)
KASHA	68B	PRL 20 217	+Larsen, Leipuner, Adair (BNL, YALE)
KASHA	68C	CJP 46 5730	+Larsen, Leipuner, Adair (BNL, YALE)
RANK	68	PR 176 1635	(MICH)
BARTON	67	PRSL 90 87	(NPOL)
BATHOW	67	PL 25B 163	(DESY)
BUHLER	67	NC 49A 209	+Freytag, Schulz, Tesch (CERN, BGNA)
BUHLER	67B	NC 51A 837	+Dalpiaz, Massam, Zichichi (CERN, BGNA, STRB)
FOSS	67	PL 25B 166	+Garelick, Homma, Lobar, Osborne, Uglum (MIT)
GOMEZ	67	PRL 18 1022	+Kobrak, Moline, Mullins, Orth, VanPutten+ (CIT)
KASHA	67	PR 154 1263	+Leipuner, Wangler, Alsppector, Adair (BNL, YALE)
STOVER	67	PR 164 1599	+Moran, Trischka (SYRA)
DORFMAN	66	PL 21 360	(BNL, YALE)
BONNETT	66	PRL 17 196	+Stoeckel (YALE)
BUHLER	66	NC 45A 520	+Fortunato, Massam, Muller+ (CERN, BGNA, STRB)
CHUPKA	66	PRL 17 60	+Schiffer, Stevens (ANL)
GALLINARO	66	PL 23 609	+Morpurgo (GENO)
KASHA	66	PR 150 1140	+Leipuner, Adair (BNL, YALE)
LAMB	66	PRL 17 1068	+Lundy, Novey, Yovanovitch (ANL)
DELISE	65	PR 140B 458	+Bowen (ARIZ)
FRANZINI	65	PR 140B 458	+Eades, Lederman, Lee, Ting (COLU)
FRANZINI	65B	PRL 14 196	+Leonic, Rahm, Sarnios, Schwartz (BNL, COLU)
MASSAM	65	NC 40A 589	+Muller, Zichichi (CERN)
BINGHAM	64	PL 9 201	+Dickinson, Diebold, Koch, Leith+ (CERN, EPOL)
BLUM	64	PRL 13 353A	+Brandt, Coconi, Czyzewski, Danysz+ (CERN)
BOWEN	64	PRL 13 728	+Delise, Kalbach, Mortara (ARIZ)
HAGOPIAN	64	PRL 13 280	+Selove, Ehrlich, Leboy, Lanza+ (PENN, BNL)
LEIPUNER	64	PL 12 423	+Chu, Larsen, Adair (BNL, YALE)
MOUSON	64	PL 9 199	(CERN)
SUNYAR	64	PR 136B 1157	+Schwarzshild, Connors (BNL)
HILLAS	59	Nature 184 B92	+Schwarzshild, Connors (AERE)
MILLIKAN	10	Phil Mag 19 209	+Cranshaw (CHIC)

OTHER RELATED PAPERS

LYONS	85	PRPL C129 225	(OXF)
MARINELLI	82	PRPL 85 161	+Morpurgo (GENO)

Magnetic Monopole Searches

NOTE ON MAGNETIC MONOPOLE SEARCHES

(by W.P. Trower, Virginia Polytechnic Institute and State University)

Although the usual formulation of Maxwell's equations suggests magnetic monopoles, no observed phenomenon requires them for explanation [1]. A monopole anywhere in the universe results in electric charge quantization everywhere, and leads to the prediction of a least magnetic charge $G = e/2\alpha$, the Dirac charge [2]. Recently monopoles have become indispensable in many gauge theories, which endow them with a variety of extraordinarily large masses.

Monopole detectors have predominantly used either induction or ionization. Induction experiments measure the monopole magnetic charge and are independent of monopole electric charge, mass, and velocity. Monopole candidate events (CABRERA 82, CAPLIN 86) in single semiconductor loops have been detected by this method, but no two-loop coincidence has been observed. Ionization experiments rely on a magnetic charge producing more ionization than an electrical charge with the same velocity. However, the ability to distinguish a monopole by ionization diminishes with velocity.

Cosmic rays are the most likely source of massive monopoles, since accelerator energies are insufficient to produce them. Evidence for such monopoles may also be obtained from astrophysical observations.

This compilation is only a guide to the literature, since the quoted experimental limits are often only indicative.

References

1. J.D. Jackson, CERN-77-17 (1977).
2. P.A.M. Dirac, Proc. Royal Soc. London **A133**, 60 (1931).

Searches Full Listings

Magnetic Monopole Searches

Monopole Production Cross Section — Accelerator Searches

X-SECT (cm ²)	MASS (GeV)	CHG (g)	ENERGY (GeV)	BEAM	EVTS	DOCUMENT ID	TECN
<3.E-37	<45.0	1.0	88-94	e ⁺ e ⁻	0	PINFOLD 93	PLAS
<3.E-37	<41.6	2.0	88-94	e ⁺ e ⁻	0	PINFOLD 93	PLAS
<7.E-35	<44.9	0.2-1.0	89-93	e ⁺ e ⁻	0	KINOSHITA 92	PLAS
<2.E-34	<850	≥ 0.5	1800	p \bar{p}	0	BERTANI 90	PLAS
<1.2E-33	<800	≥ 1	1800	p \bar{p}	0	PRICE 90	PLAS
<1.E-37	<29	1	50-61	e ⁺ e ⁻	0	KINOSHITA 89	PLAS
<1.E-37	<18	2	50-61	e ⁺ e ⁻	0	KINOSHITA 89	PLAS
<1.E-38	<17	<1	35	e ⁺ e ⁻	0	BRAUNSCH... 88B	CNTR
<8.E-37	<24	1	50-52	e ⁺ e ⁻	0	KINOSHITA 88	PLAS
<1.3E-35	<22	2	50-52	e ⁺ e ⁻	0	KINOSHITA 88	PLAS
<9.E-37	<4	<0.15	10.6	e ⁺ e ⁻	0	GENTILE 87	CLEO
<3.E-32	<800	≥ 1	1800	p \bar{p}	0	PRICE 87	PLAS
<3.E-38	<3	29	e ⁺ e ⁻	0	FRYBERGER 84	PLAS	
<1.E-31	<10	1.3	540	p \bar{p}	0	AUBERT 83B	PLAS
<4.E-38	<20	<6	34	e ⁺ e ⁻	0	MUSSET 83	PLAS
<8.E-36	<20	52	pp	0	1 DELL 82	CNTR	
<9.E-37	<30	<3	29	e ⁺ e ⁻	0	KINOSHITA 82	PLAS
<1.E-37	<20	<24	63	pp	0	CARRIGAN 78	CNTR
<1.E-37	<30	<3	56	pp	0	HOFFMANN 78	PLAS
			62	pp	0	1 DELL 76	SPRK
<4.E-33			300	p	0	1 STEVENS 76B	SPRK
<1.E-40	<5	<2	70	p	0	2 ZRELOV 76	CNTR
<2.E-30			300	n	0	1 BURKE 75	OSPK
<1.E-38			8	v	0	3 CARRIGAN 75	HLBC
<5.E-43	<12	<10	400	p	0	EBERHARD 75B	INDU
<2.E-36	<30	<3	60	pp	0	GIACOMELLI 75	PLAS
<5.E-42	<13	<24	400	p	0	CARRIGAN 74	CNTR
<6.E-42	<12	<24	300	p	0	CARRIGAN 73	CNTR
<2.E-36		1	.001	γ	0	2 BARTLETT 72	CNTR
<1.E-41	<5		70	p	0	GUREVICH 72	EMUL
<1.E-40	<3	<2	28	p	0	AMALDI 63	EMUL
<2.E-40	<3	<2	30	p	0	PURCELL 63	CNTR
<1.E-35	<3	<4	28	p	0	FIDECARO 61	CNTR
<2.E-35	<1	1	6	p	0	BRADNER 59	EMUL

1 Multiphoton events.

2 Cherenkov radiation polarization.

3 Re-examines CERN neutrino experiments.

Monopole Flux — Cosmic Ray Searches

FLUX (cm ⁻² sr ⁻¹ s ⁻¹)	MASS (GeV)	CHG (g)	COMMENTS (β = v/c)	EVTS	DOCUMENT ID	TECN	
<5.6E-15	1	1.8E-4	<β < 3.E-3	0	4 AHLEN 94	MCRO	
<8.7E-15	1	>2.E-3		0	THRON 92	SOUD	
<4.4E-12	1	all β		0	GARDNER 91	INDU	
<7.2E-13	1	all β		0	HUBER 91	INDU	
<3.7E-15	>E12	1	β=1.E-4	0	5 ORITO 91	PLAS	
<3.2E-16	>E10	1	β > 0.05	0	5 ORITO 91	PLAS	
<3.2E-16	>E10-E12	2,3		0	5 ORITO 91	PLAS	
<3.8E-13	1	all β		0	BERMON 90	INDU	
<5.E-16	1	β < 1.E-3		0	6 BEZRUKOV 90	CHER	
<1.8E-14	1	β > 1.E-4		0	7 BUCKLAND 90	HEPT	
<1E-18	1	3.E-4	<β < 1.5E-3	0	8 GHOSH 90	MICA	
<7.2E-13	1	all β		0	HUBER 90	INDU	
<5.E-12	>E7	1	3.E-4	<β < 5.E-3	0	BARISH 87	CNTR
<1.E-13	1	1.E-5	<β < 1	0	6 BARTELT 87	SOUD	
<1.E-10	1	all β		0	EBISU 87	INDU	
<2.E-13	1	1.E-4	<β < 6.E-4	0	MASEK 87	HEPT	
<2.E-14	1	4.E-5	<β < 2.E-4	0	NAKAMURA 87	PLAS	
<2.E-14	1	1.E-3	<β < 1	0	NAKAMURA 87	PLAS	
<5.E-14	1	9.E-4	<β < 1.E-2	0	SHEPKO 87	CNTR	
<2.E-13	1	4.E-4	<β < 1	0	TSUKAMOTO 87	CNTR	
<5.E-14	1	all β		1	9 CAPLIN 86	INDU	
<5.E-12	1	1		0	CROMAR 86	INDU	
<1.E-13	1	7.E-4	<β	0	HARA 86	CNTR	
<7.E-11	1	all β		0	INCANDELA 86	INDU	
<1.E-18	1	4.E-4	<β < 1.E-3	0	8 PRICE 86	MICA	
<5.E-12	1	1		0	BERMON 85	INDU	
<6.E-12	1	0		0	CAPLIN 85	INDU	
<6.E-10	1	0		0	EBISU 85	INDU	
<3.E-15	1	5.E-5	β ≤ 1.E-3	0	6 KAJITA 85	CNTR	
<2.E-21	1	β < 1.E-3		0	6,10 KAJITA 85	CNTR	
<3.E-15	1	1.E-3	<β < 1.E-1	0	6 PARK 85B	CNTR	
<5.E-12	1	1.E-4	<β < 1	0	6 BATTISTONI 84	NUSX	
<7.E-12	1	1		0	INCANDELA 84	INDU	
<7.E-13	1	3.E-4	<β	0	7 KAJINO 84	CNTR	
<2.E-12	1	3.E-4	<β < 1.E-1	0	KAJINO 84B	CNTR	
<6.E-13	1	5.E-4	<β < 1	0	KAWAGOE 84	CNTR	
<2.E-14	1	1.E-3	<β	0	6 KRISHNA... 84	CNTR	
<4.E-13	1	6.E-4	<β < 2.E-3	0	LISS 84	CNTR	
<1.E-16	1	3.E-4	<β < 1.E-3	0	8 PRICE 84	MICA	

<1.E-13	1	1.E-4	<β	0	PRICE 84B	PLAS
<4.E-13	1	6.E-4	<β < 2.E-3	0	TARLE 84	CNTR
	7	11 ANDERSON 83	EMUL			
<4.E-13	1	1.E-2	<β < 1.E-3	0	BARTELT 83B	CNTR
<1.E-12	1	7.E-3	<β < 1	0	BARWICK 83	PLAS
<3.E-13	1	1.E-3	<β < 4.E-1	0	BONARELLI 83	CNTR
<3.E-12	5.E-4	<β < 5.E-2	0	6 BOSETTI 83	CNTR	
<4.E-11	1			0	CABRERA 83	INDU
<5.E-15	1	1.E-2	<β < 1	0	DOKE 83	PLAS
<8.E-15	1	1.E-4	<β < 1.E-1	0	6 ERREDE 83	CNTR
<5.E-12	1	1.E-4	<β < 3.E-2	0	GROOM 83	CNTR
<2.E-12	6.E-4	<β < 1	0	MASHIMO 83	CNTR	
<1.E-13	1	β=3.E-3	0	ALEXEYEV 82	CNTR	
<2.E-12	1	7.E-3	<β < 6.E-1	0	BONARELLI 82	CNTR
6.E-10	1	all β	1	12 CABRERA 82	INDU	
<2.E-11	1	1.E-2	<β < 1.E-1	0	MASHIMO 82	CNTR
<2.E-15	concentrator			0	BARTLETT 81	PLAS
<1.E-13	>1	1.E-3	<β	0	KINOSHITA 81B	PLAS
<5.E-11	<E17	3.E-4	<β < 1.E-3	0	ULLMAN 81	CNTR
<2.E-11	concentrator			0	BARTLETT 78	PLAS
1.E-1	>200	2		1	13 PRICE 75	PLAS
<2.E-13	>2			0	FLEISCHER 71	PLAS
<1.E-19	>2	obsidian, mica		0	FLEISCHER 69C	PLAS
<5.E-15	<15	<3	concentrator	0	CARTHERS 66	ELEC
<2.E-11	<1-3	concentrator		0	MALKUS 51	EMUL

4 AHLEN 94 limit for dyons extends down to β=0.9E-4 and a limit of 1.3E-14 extends to β = 0.8E-4.

5 ORITO 91 limits are functions of velocity. Lowest limits are given here.

6 Catalysis of nucleon decay; sensitive to assumed catalysis cross section.

7 Used DKMPR mechanism and Penning effect.

8 Assumes monopole attaches fermion nucleus.

9 Limit from combining data of CAPLIN 86, BERMON 85, INCANDELA 84, and CABRERA 82. For a discussion of controversy about CAPLIN 86 observed event, see GUY 87. Also see SCHOUTEN 87.

10 Based on lack of high-energy solar neutrinos from catalysis in the sun.

11 Anomalous long-range α (⁴He) tracks.

12 CABRERA 82 candidate event has single Dirac charge within ±5%.

13 ALVAREZ 75, FLEISCHER 75, and FRIEDLANDER 75 explain as fragmenting nucleus. EBERHARD 75 and ROSS 76 discuss conflict with other experiments. HAGSTROM 77 reinterprets as antinucleus. PRICE 78 reassesses.

Monopole Flux — Astrophysics

FLUX (cm ⁻² sr ⁻¹ s ⁻¹)	MASS (GeV)	CHG (g)	COMMENTS (β = v/c)	EVTS	DOCUMENT ID	TECN
<1.E-16	E17	1	galactic field	0	14 ADAMS 93	COSM
<1.E-23			Jovian planets	0	15 ARAFUNE 85	COSM
<1.E-18	E15	1	solar trapping	0	BRACCI 85B	COSM
<1.E-18				0	15 HARVEY 84	COSM
<3.E-23			neutron stars	0	KOLB 84	COSM
<7.E-22			pulsars	0	15 FREESE 83B	COSM
<1.E-18	<E18	1	intergalactic field	0	15 REPHAEI 83	COSM
<1.E-23			neutron stars	0	15 DIMOPOUL... 82	COSM
<5.E-22			neutron stars	0	15 KOLB 82	COSM
<5.E-15	>E21		galactic halo	0	SALPETER 82	COSM
<1.E-12	E19	1	β=3.E-3	0	16 TURNER 82	COSM
<1.E-16			galactic field	0	PARKER 70	COSM

14 ADAMS 93 limit based on "survival and growth of a small galactic seed field" / m⁻¹⁰⁻¹⁶ (m/10¹⁷ GeV) cm⁻² s⁻¹ sr⁻¹. Above 10¹⁷ GeV, limit 10⁻¹⁶ (10¹⁷ GeV/m) cm⁻² s⁻¹ sr⁻¹ (from requirement that monopole density does not overclose the universe) is more stringent.

15 Catalysis of nucleon decay.

16 Re-evaluates PARKER 70 limit for GUT monopoles.

Monopole Density — Matter Searches

DENSITY	CHG (g)	MATERIAL	EVTS	DOCUMENT ID	TECN
<2.E-7/gram	>0.6	Fe ore	0	17 EBISU 87	INDU
>1.E-14/gram	>1/3	iron aerosols	>1	MIKHAILOV 83	SPEC
<6.E-4/gram		air, seawater	0	CARRIGAN 76	CNTR
<5.E-1/gram	>0.04	11 materials	0	CABRERA 75	INDU
<2.E-4/gram	>0.05	moon rock	0	ROSS 73	INDU
<6.E-7/gram	<140	seawater	0	KOLM 71	CNTR
<1.E-2/gram	<120	manganese nodules	0	FLEISCHER 69	PLAS
<1.E-4/gram	>0	manganese	0	FLEISCHER 69B	PLAS
<2.E-3/gram	<1-3	magnetite, meteor	0	GOTO 63	EMUL
<2.E-2/gram		meteorite	0	PETUKHOV 63	CNTR

17 Mass 1 × 10¹⁴-1 × 10¹⁷ GeV.

Monopole Density — Astrophysics

DENSITY	CHG (g)	MATERIAL	EVTS	DOCUMENT ID	TECN
<1.E-9/gram	1	sun, catalysis	0	18 ARAFUNE 83	COSM
<6.E-33/nucl	1	moon wake	0	SCHATTEN 83	ELEC
<2.E-28/nucl		earth heat	0	CARRIGAN 80	COSM
<2.E-4/prot		42cm absorption	0	BRODERICK 79	COSM
<2.E-13/m ³		moon wake	0	SCHATTEN 70	ELEC

18 Catalysis of nucleon decay.

Magnetic Monopole Searches, Supersymmetric Particle Searches

REFERENCES FOR Magnetic Monopole Searches

AHLEN	94	PRL 72 608	+Ambrosio, Antolini, Auremma+ (MACRO Collab.)
ADAMS	93	PRL 70 2511	+Fazio, Freese, Tarte+ (MICH, FNAL)
PINFOLD	93	PL 3315 407	+Du, Kinoshita, Lorzoz+ (ALBE, HARV, MONT, UCB)
KINOSHITA	92	PR D46 R881	+Du, Giacomelli, Patriziili+ (HARV, BGNA, REHO)
THRON	92	PR D46 4846	+Allison, Ainer, Ambats+ (SOUDAN-2 Collab.)
GARDNER	91	PR D44 622	+Cabrera, Huber, Taber (STAN)
HUBER	91	PR D44 636	+Cabrera, Taber, Gardner (STAN)
ORITO	91	PRL 66 1951	+Ichinose, Nakamura+ (ICEPP, WASCER, NIHO, ICRN)
BERMON	90	PRL 64 839	+Chi, Tsuei+ (IBM, BNL)
BERTANI	90	EPL 12 613	+Giacomelli, Mondardini, Pal+ (BGNA, INFN)
BEZUKOV	90	SJNP 52 54	+Belolaptkov, Bugaev, Budnev+ (IHRM)
		Translated from YAF 52 86	
BUCKLAND	90	PR D41 2726	+Masek, Vernon, Knapp, Stronsi (UCSD)
GHOSH	90	EPL 12 25	+Chatterjee (JADA)
HUBER	90	PRL 64 835	+Cabrera, Tabor, Gardner (STAN)
PRICE	90	PRL 65 149	+Giurui, Kinoshita (UCB, HARV)
KINOSHITA	89	PL B228 543	+Fuji, Nakajima+ (HARV, TISA, KEK, UCB)
BRAUNSCH...	88B	ZPHY C38 543	+Braunschweig, Gerhards, Kirschfink+ (TASSO Collab.)
KINOSHITA	88	PRL 60 1610	+Fuji, Nakajima+ (HARV, TISA, KEK, UCB, GIFU)
BARISH	87	PR D36 2641	+Liu, Lane (CIT)
BARTELT	87	PR D36 1990	+Courant, Heller+ (Soudan Collab.)
	Also	PR D40 1701 erratum	Bartelt, Courant, Heller+ (Soudan Collab.)
EBISU	87	PR D36 3359	+Watanabe (KOBE)
	Also	JPG 11 893	Ebisu, Watanabe (KOBE)
GENTILE	87	PR D35 1081	+Haas, Hempstead+ (CLEO Collab.)
GUY	87	Nature 325 463	(LOIC)
MASEK	87	PR D35 2758	+Knapp, Miller, Stronski, Vernon, White (UCSD)
NAKAMURA	87	PL B183 395	+Kawagoe, Yamamoto+ (INUS, WASCER, NIHO)
PRICE	87	PRL 59 2523	+Guoxiao, Kinoshita (UCB, HARV)
SCHOUTEN	87	JPE 20 850	+Caplin, Guy, Hardiman+ (LOIC)
SHEPKO	87	PR D35 2917	+Gagliardi, Green, McIntyre+ (TAMU)
TSUKAMOTO	87	PL 3 339	+Hagano, Anraku+ (ICRR)
CAPLIN	86	Nature 321 402	+Hardiman, Koratzinos, Schouten (LOIC)
	Also	JPE 20 850	Schouten, Caplin, Guy, Hardiman+ (LOIC)
	Also	Nature 325 463	Guy (LOIC)
CROMAR	86	PRL 56 2561	+Clark, Fickett (NBSB)
HARA	86	PRL 56 553	+Honda, Ohno+ (ICRR, KYOT, KEK, KOBE, ICEPP)
INCANDELA	86	PR D34 2637	+Frisch, Somaiwal, Kuchnir+ (CHIC, FNAL, MICH)
PRICE	86	PRL 56 1226	+Salomon (UCB)
ARAFUNE	86	PR D34 2586	+Fukugita, Yanagita (ICRR, KYOTY, IBAR)
BERMON	85	PRL 55 1850	+Chaudhuri, Chi, Tesche, Tsuei (IBM)
BRACCI	85B	NP B258 726	+Florentini, Mezzorani (PISA, CAGL, INFN)
	Also	LNC 42 123	Bracci, Fiorentini (PISA)
CAPLIN	85	Nature 317 234	+Guy, Hardiman, Park, Schouten (LOIC)
EBISU	85	JPG 11 883	+Watanabe (KOBE)
KAJITA	85	JPSJ 54 4065	+Arisaka, Koshiba, Nakahata+ (ICRR, KEK, NIIG)
PARK	85B	NP B252 261	+Arisawa, Cortez, Foster+ (IMB Collab.)
BATTISTONI	84	PL 133B 454	+Belotti, Bologna, Campana+ (NUSEX Collab.)
FRYBERGER	84	PR D29 1524	+Coan, Kinoshita, Price (SLAC, UCB)
HARVEY	84	NP B236 255	(PRIN)
INCANDELA	84	PRL 53 2067	+Campbell, Frisch+ (CHIC, FNAL, MICH)
KAJINO	84	PRL 52 1373	+Matsuno, Yuan, Kitamura (ICRR)
KAJINO	84B	JPG 10 447	+Matsuno, Kitamura, Aoki, Yuan, Mitsui+ (ICRR)
KAWAGOE	84	LNC 41 315	+Mashimo, Nakamura, Nozaki, Orto (TOKY)
KOLB	84	APJ 286 702	+Turner (FNAL, CHIC)
KRISHNA...	84	PL 142B 99	+Krishnaswamy, Menon+ (TATA, OSKC, INUS)
LISS	84	PR D30 884	+Ahlen, Tarle (UCB, IND, MICH)
PRICE	84	PRL 52 1265	+Guo, Ahlen, Fleischer (ROMA, UCB, IND, GESC)
PRICE	84B	PL 140B 112	(CERN)
TARLE	84	PRL 52 90	+Ahlen, Liss (UCB, MICH, IND)
ANDERSON	83	PR D28 2308	+Lord, Strauss, Wilkes (WASH)
ARAFUNE	83	PL 133B 380	+Fukugita (ICRR, KYOTY)
AUBERT	83B	PL 120B 465	+Muset, Price, Vialle (CERN, LAPP)
BARTELT	83B	PRL 50 655	+Courant, Heller, Joyce, Marshak+ (MINI, ANL)
BARWICK	83	PR D28 2338	+Kinoshita, Price (UCB)
BONARELLI	83	PL 126B 137	+Capiluppi, Dantone (BGNA)
BOSETTI	83	PL 133B 265	+Gorham, Harris, Learned+ (AACH3, HAWA, TOKY)
CABRERA	83	PRL 51 1933	+Taber, Gardner, Bourg (STAN)
DOKE	83	PL 129B 370	+Hayashi, Hamasaki+ (WASU, RIKK, TTAM, RIKEN)
ERREDE	83	PRL 51 245	+Stone, Vander Velde, Bionta+ (IMB Collab.)
FRESE	83B	PRL 51 1625	+Turner, Schramm (CHIC)
GROOM	83	PL 112B 100	+Loh, Nelson, Ritson (UTAH, STAN)
MASHIMO	83	PL 128B 327	+Orto, Kawagoe, Nakamura, Nozaki (ICEPP)
MIKHAILOV	83	PL 130B 331	(KAZA)
MUSSET	83	PL 128B 333	+Price, Lohrmann (CERN, HAMB)
REPHELI	83	PL 121B 115	+Turner (CHIC)
SCHATTEN	83	PR D27 1525	(NASA)
ALEXEYEV	82	LNC 35 413	+Boliev, Chudakov, Makoiev, Mikheyev+ (INRM)
BONARELLI	82	PRL 48 1378	+Capiluppi, Dantone+ (BGNA)
CABRERA	82	PRL 48 1378	(STAN)
DELL	82	NP B209 45	+Yuan, Roberts, Dooher+ (BNL, ADEL, ROMA)
DIMOPOUL...	82	PL 119B 320	+Dimopoulos, Preskill, Wilczek (HARV, UCSBT)
KINOSHITA	82	PRL 48 77	+Price, Fryberger (UCB, SLAC)
KOLB	82	PRL 49 1373	+Colgate, Harvey (LASL, PRIN)
MASHIMO	82	JPSJ 51 3067	+Kawagoe, Koshiba (INUS)
SALPETER	82	PR D18 1114	+Shapiro, Wasserman (CORN)
TURNER	82	PR D26 1296	+Parker, Bogdan (CHIC)
BARTLETT	81	PR D24 612	+Soo, Fleischer, Hart+ (COLO, GESC)
KINOSHITA	81B	PR D24 1707	+Price (UCB)
ULLMAN	81	PRL 47 289	(LEHM, BNL)
CARRIGAN	80	Nature 288 348	(FNAL)
BRODERICK	79	PR D19 1046	+Ficene, Teplitz, Teplitz (VPI)
BARTLETT	78	PR D18 2253	+Soo, White (COLO, PRIN)
CARRIGAN	78	PR D17 1754	+Strauss, Giacomelli (FNAL, BGNA)
HOFFMANN	78	LNC 23 357	+Kantarjian, Diliberto, Meddi+ (CERN, ROMA)
PRICE	78	PR D18 1382	+Shirk, Osborne, Pinsky (UCB, HOUS)
HAGSTROM	77	PRL 38 729	(LBL)
CARRIGAN	76	PR D13 1823	+Nezrick, Strauss (FNAL)
DELL	76	LNC 15 269	+Uto, Yuan, Amaldi+ (CERN, BNL, ROMA, ADEL)
ROSS	76	LBL-4665	(LBL)
STEVENS	76B	PR D14 2207	+Collins, Ficene, Trower, Fischer+ (VPI, BNL)
ZRELOV	76	CZP 826 1306	+Koilarova, Kollar, Lupitsev, Paviovic+ (JINR)
ALVAREZ	75	LBL-4260	(LBL)
BURKE	75	PL 60B 113	+Gustafson, Jones, Longo (MICH)
CABRERA	75	Thesis	(STAN)
CARRIGAN	75	NP B91 279	+Nezrick (FNAL)
	Also	PR D3 56	Carrigan, Nezrick (FNAL)
EBERHARD	75	PR D13 3099	+Ross, Taylor, Alvarez, Oberlack (LBL, MPIM)
EBERHARD	75B	LBL-4289	(LBL)
FLEISCHER	75	PRL 35 1412	+Walker (GESC, WUSL)
FRIEDLANDER	75	PRL 35 1167	(WUSL)
GIACOMELLI	75	NC 28A 21	+Rossi+ (BGNA, CERN, SACL, ROMA)
PRICE	75	PRL 35 487	+Shirk, Osborne, Pinsky (UCB, HOUS)
CARRIGAN	74	PR D10 3867	+Nezrick, Strauss (FNAL)
CARRIGAN	73	PR D8 3717	+Nezrick, Strauss (FNAL)
ROSS	73	PR D8 698	+Eberhard, Alvarez, Watt (LBL, SLAC)
	Also	PR D4 3260	Eberhard, Ross, Alvarez, Watt (LBL, SLAC)
	Also	Science 167 701	Alvarez, Eberhard, Ross, Watt (LBL, SLAC)

BARTLETT	72	PR D6 1817	+Lahana (COLO)
GUREVICH	72	PL 38B 549	+Khalimov, Martemyanov+ (KIAE, NOVO, SERP)
	Also	JETP 34 917	Barlov, Gurevich, Zolotorev (KIAE, NOVO, SERP)
	Also	Translated from ZETF 61 1721	Gurevich, Khakimov+ (KIAE, NOVO, SERP)
FLEISCHER	70	PL 31B 394	(GESC)
KOLM	71	PR D4 24	+Hart, Nichols, Price (MIT, SLAC)
PARKER	70	APJ 160 383	(CHIC)
SCHATTEN	70	PR D1 2245	(NASA)
FLEISCHER	69	PR 177 2029	+Jacobs, Schwartz, Price (GESC, FSU)
FLEISCHER	69B	PR 184 1393	+Hart, Jacobs+ (GESC, UNCS, GSCO)
FLEISCHER	69C	PR 184 1398	+Price, Woods (GESC)
	Also	JAP 41 958	Fleischer, Hart, Jacobs, Price+ (GESC)
CARITHERS	66	PR 149 1070	+Stefanski, Adair (YALE, BNL)
AMALDI	63	NC 28 773	+Baroni, Manfredini+ (ROMA, UCSD, CERN)
GOTO	63	PR 132 387	+Kohn, Ford (TOKY, MIT, BRAN)
PETUKHOV	63	NP 49 87	+Yakimenko (LEBD)
PURCELL	63	PR 129 2326	+Collins, Fujii, Hornbostel, Turkot (HARV, BNL)
FIDECARO	61	NC 22 657	+Finocchiaro, Giacomelli (CERN)
BRADNER	59	PR 114 603	(LBL)
MALKUS	51	PR 83 899	+Isbell (CHIC)

OTHER RELATED PAPERS

GROOM	86	PRPL 140 323	(UTAH)
Review			

Supersymmetric Particle Searches

NOTE ON SUPERSYMMETRY

(by Howard E. Haber, Univ. of California, Santa Cruz)

Supersymmetry is an attractive theoretical framework that may require the consistent unification of particle physics and gravity, which takes place at an energy of order the Planck scale ($\approx 10^{19}$ GeV) [1-3]. However, supersymmetry is clearly not an exact symmetry of nature, and therefore must be broken. In theories of "low-energy" supersymmetry, the effective scale of supersymmetry breaking is tied to the electroweak scale [4,5]. In this way, it is hoped that supersymmetry will ultimately explain the origin of the large hierarchy between the W and Z masses and the Planck scale.

The minimal supersymmetric extension of the Standard Model (MSSM) consists of taking the Standard Model and adding the corresponding supersymmetric partners [6]. In addition, the MSSM contains two hypercharge $Y = \pm 1$ Higgs doublets, which is the minimal structure for the Higgs sector of an anomaly-free supersymmetric extension of the Standard Model that generates mass for both "up"-type and "down"-type quarks (and charged leptons) [7,8]. Supersymmetric interactions consistent with (global) $B-L$ conservation (B = baryon number and L = lepton number) are included. Finally, the most general soft-supersymmetry-breaking terms are added [9]. If supersymmetry is relevant for explaining the scale of electroweak interactions, then the mass parameters that occur in the soft-supersymmetry-breaking terms must be of order 1 TeV or below [10]. Some bounds on these parameters exist due to the absence of supersymmetry particle production at current accelerators, as well as the absence of any evidence for virtual supersymmetric particle exchange in a variety of Standard Model processes [11].

As a consequence of $B-L$ invariance, the MSSM possesses a discrete R -parity invariance, where $R = (-1)^{3(B-L)+2S}$ for a particle of spin S [12]. Note that this formula implies that all the ordinary Standard Model particles have even R -parity, whereas the corresponding supersymmetric partners have odd R -parity. The conservation of R -parity in scattering and decay

Searches Full Listings

Supersymmetric Particle Searches

processes has a crucial impact on supersymmetric phenomenology. For example, starting from an initial state involving ordinary (R -even) particles, it follows that supersymmetric particles must be produced in pairs. In general, these particles are highly unstable and decay quickly into lighter states. However, R -parity invariance also implies that the lightest supersymmetric particle (LSP) is absolutely stable, and must eventually be produced at the end of a decay chain of a heavy unstable supersymmetric particle. In order to be consistent with cosmological constraints, the LSP is almost certainly electrically and color neutral [13]. Consequently, the LSP is weakly-interacting in ordinary matter, *i.e.* it behaves like a neutrino and will escape detectors without being directly observed. Thus, the canonical signature for (R -parity conserving) supersymmetric theories is missing (transverse) energy, due to the escape of the LSP. Some model builders attempt to relax the assumption of R -parity conservation [14]. Models of this type must break $B-L$ and are therefore strongly constrained. Nevertheless, because such models cannot be completely ruled out, it is important to allow for the possibility of R -parity violating processes in the search for supersymmetry. In particular, the LSP would be unstable, and this fact (among others) leads to a phenomenology of broken- R -parity models that is very different from that of the MSSM.

In the MSSM, supersymmetry breaking is induced by the soft-supersymmetry breaking terms mentioned above. These terms parametrize our ignorance of the fundamental mechanism of supersymmetry breaking. If this breaking occurs spontaneously, then (in the absence of supergravity) a massless Goldstone fermion called the *goldstino* (\tilde{G}) must exist. The goldstino would then be the LSP and could play an important role in supersymmetric phenomenology [15]. In models that incorporate supergravity, this picture changes. If supergravity is spontaneously broken, the goldstino is absorbed (“eaten”) by the *gravitino* ($g_{3/2}$), the spin-3/2 partner of the graviton [16]. By this super-Higgs mechanism, the gravitino acquires a mass ($m_{3/2}$). In models of this type; the gravitino mass is typically of order of the low-energy supersymmetry-breaking scale ($\mathcal{O}(1 \text{ TeV})$), while its couplings are gravitational in strength [1,17]. Such a gravitino would play no role in supersymmetric phenomenology at colliders.

The parameters of the MSSM are conveniently described by considering separately the supersymmetry-conserving sector and the supersymmetry-breaking sector. A careful discussion of the conventions used in defining the MSSM parameters can be found in Ref. 18. Among the parameters of the supersymmetry conserving sector are: (i) gauge couplings: g_s , g , and g' , corresponding to the Standard Model gauge group $SU(3) \times SU(2) \times U(1)$ respectively; (ii) Higgs Yukawa couplings: λ_e , λ_u , and λ_d (which are 3×3 matrices in flavor space); and (iii) a supersymmetry-conserving Higgs mass parameter μ . The supersymmetry-breaking sector contains the following set of parameters: (i) gaugino Majorana masses M_3 , M_2

and M_1 associated with the $SU(3)$, $SU(2)$, and $U(1)$ subgroups of the Standard Model; (ii) scalar mass matrices for the squarks and sleptons; (iii) Higgs-squark-squark trilinear interaction terms (the so-called “ A -parameters”) and corresponding terms involving the sleptons; and (iv) three scalar Higgs mass parameters—two diagonal and one off-diagonal mass terms for the two Higgs doublets. These three mass parameters can be re-expressed in terms of the two Higgs vacuum expectation values, v_1 and v_2 , and one physical Higgs mass. Here, v_1 (v_2) is the vacuum expectation value of the Higgs field which couples exclusively to down-type (up-type) quarks and leptons. Note that $v_1^2 + v_2^2 = (246 \text{ GeV})^2$ is fixed by the W mass (or equivalently by the Fermi constant G_F), while the ratio

$$\tan \beta = v_2/v_1 \quad (1)$$

is a free parameter of the model. The supersymmetric constraints imply that the MSSM Higgs sector is automatically CP -conserving (at tree-level). Thus, $\tan \beta$ is a real parameter (conventionally chosen to be positive), and the physical neutral Higgs scalars are CP -eigenstates. Nevertheless, the MSSM does contain a number of possible new sources of CP violation. For example, gaugino mass parameters, the A -parameters, and μ may be complex. Some combination of these complex phases must be less than of order 10^{-2} – 10^{-3} (for a supersymmetry-breaking scale of 100 GeV) to avoid generating electric dipole moments for the neutron, electron, and atoms in conflict with observed data [19]. However, these complex phases have little impact on the direct searches for supersymmetric particles, and are usually ignored in experimental analyses.

Before describing the supersymmetric particle sector, let us consider the Higgs sector of the MSSM [20]. There are five physical Higgs particles in this model: a charged Higgs pair (H^\pm), two CP -even neutral Higgs bosons (denoted by H_1^0 and H_2^0 where $m_{H_1^0} \leq m_{H_2^0}$) and one CP -odd neutral Higgs boson (A^0). The properties of the Higgs sector are determined by the Higgs potential which is made up of quadratic terms [whose squared-mass coefficients were mentioned above Eq. (1)] and quartic interaction terms. The strengths of the interaction terms are directly related to the gauge couplings by supersymmetry (and are not affected at tree-level by supersymmetry-breaking). As a result, $\tan \beta$ [defined in Eq. (1)] and one Higgs mass determine: the Higgs spectrum, an angle α [which indicates the amount of mixing of the original $Y = \pm 1$ Higgs doublet states in the physical CP -even scalars], and the Higgs boson couplings. When one-loop radiative corrections are incorporated, additional parameters of the supersymmetric model enter via virtual loops. The impact of these corrections can be significant [21,22]. For example, at tree-level, the MSSM predicts $m_{H_1^0} \leq m_Z$ [7,8]. If true, this would imply that experiments to be performed at LEP-II operating at its maximum energy and luminosity would rule out the MSSM if H_1^0 were not found. However, this Higgs mass bound need not be respected when radiative corrections are incorporated. For example, in Ref. 21, the following upper

bound was obtained for $m_{H_1^0}$ (assuming $m_{A^0} > m_Z$) in the limit of $m_Z \ll m_t \ll M_{\tilde{t}}$ [where top-squark (\tilde{t}_L - \tilde{t}_R) mixing is neglected]

$$m_{H_1^0}^2 \lesssim m_Z^2 + \frac{3g^2 m_Z^4}{16\pi^2 m_W^2} \times \left\{ \ln \left(\frac{M_{\tilde{t}}^2}{m_t^2} \right) \left[\frac{2m_t^4 - m_t^2 m_Z^2}{m_Z^4} \right] + \frac{m_t^2}{3m_Z^2} \right\}. \quad (2)$$

For a top-squark mass of $M_{\tilde{t}} = 1$ TeV, Eq. (2) yields a positive mass shift for $m_{H_1^0}$ of about 20 GeV for $m_t = 150$ GeV, and 40 GeV for $m_t = 180$ GeV. Even if $\tan\beta = 1$ (so that $m_{H_1^0} = 0$ at tree-level), there is a large shift in $m_{H_1^0}^2$ due to radiative corrections of similar size. Clearly, the radiative corrections to the Higgs masses can have a significant impact on the search for the Higgs bosons of the MSSM at LEP and LEP-II [23].

Consider next the supersymmetric particle sector of the MSSM. The supersymmetric partners of the gauge and Higgs bosons are fermions, whose names are obtained by appending “ino” at the end of the corresponding Standard Model particle name. The *gluino* is the color octet Majorana fermion partner of the gluon with mass $M_{\tilde{g}} = |M_3|$. The supersymmetric partners of the electroweak gauge and Higgs bosons (the *gauginos* and *Higgsinos*) can mix. As a result, the physical mass eigenstates are model-dependent linear combinations of these states, called *charginos* and *neutralinos*, which are obtained by diagonalizing the corresponding mass matrices. The chargino mass matrix depends on M_2 , μ , $\tan\beta$ and m_W [24]. The corresponding chargino mass eigenstates denoted by $\tilde{\chi}_1^+$ and $\tilde{\chi}_2^+$, with masses

$$M_{\tilde{\chi}_1^+, \tilde{\chi}_2^+}^2 = \frac{1}{2} \left[|\mu|^2 + |M_2|^2 + 2m_W^2 \right] \mp \left\{ \left(|\mu|^2 + |M_2|^2 + 2m_W^2 \right)^2 - 4|\mu|^2 |M_2|^2 - 4m_W^4 \sin^2 2\beta + 8m_W^2 \sin 2\beta \operatorname{Re}(\mu M_2) \right\}^{1/2}, \quad (3)$$

where the states are ordered such that $M_{\tilde{\chi}_1^+} \leq M_{\tilde{\chi}_2^+}$. If CP -violating effects are ignored (in which case, M_2 and μ are real parameters), then one can choose a convention where $\tan\beta$ and M_2 are positive. (Note that the relative sign of M_2 and μ is meaningful. The sign of μ is convention-dependent; the reader is warned that both sign conventions appear in the literature.) The sign convention for μ implicit in Eq. (3) is used by the LEP collaborations [25] in their plots of exclusion contours in the M_2 vs. μ plane derived from the non-observation of $Z \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^-$. The neutralino mass matrix depends on M_1 , M_2 , μ , $\tan\beta$, m_Z , and the weak mixing angle θ_W [24]. The corresponding neutralino eigenstates are usually denoted by $\tilde{\chi}_i^0$ ($i = 1, \dots, 4$), according to the convention that $M_{\tilde{\chi}_1^0} \leq M_{\tilde{\chi}_2^0} \leq M_{\tilde{\chi}_3^0} \leq M_{\tilde{\chi}_4^0}$. If a chargino or neutralino eigenstate approximates a particular gaugino or Higgsino state, it may be convenient to use the corresponding nomenclature. For example, if M_1 and M_2 are

small compared to m_Z (and μ), then the lightest neutralino $\tilde{\chi}_1^0$ will be nearly a pure photino, $\tilde{\gamma}$ (the supersymmetric partner of the photon). It is common practice in the literature to reduce the supersymmetric parameter freedom by requiring that all three gaugino mass parameters are equal at some grand unification scale. Then, at the electroweak scale, the gaugino mass parameters can be expressed in terms of one of them (say, M_2). The other two gaugino mass parameters are given by

$$M_3 = (g_s^2/g^2)M_2 \quad M_1 = (5g'^2/3g^2)M_2. \quad (4)$$

Having made this assumption, the chargino and neutralino masses and mixing angles depend only on three unknown parameters: the gluino mass, μ , and $\tan\beta$.

The supersymmetric partners of the quarks and leptons are spin-zero bosons: the *squarks*, charged *sleptons*, and *sneutrinos*. For a given fermion f , there are two supersymmetric partners \tilde{f}_L and \tilde{f}_R which are scalar partners of the corresponding left and right-handed fermion. (There is no $\tilde{\nu}_R$.) However, in general, \tilde{f}_L and \tilde{f}_R are not mass-eigenstates since there is \tilde{f}_L - \tilde{f}_R mixing which is proportional in strength to the corresponding element of the scalar mass-squared-matrix [26]:

$$M_{LR}^2 = \begin{cases} m_d(A_d - \mu \tan\beta), & \text{for “down”-type } f \\ m_u(A_u - \mu \cot\beta), & \text{for “up”-type } f, \end{cases} \quad (5)$$

where m_d (m_u) is the mass of the appropriate “down” (“up”) type quark or lepton. Here, A_d and A_u are (unknown) soft-supersymmetry-breaking A -parameters and μ and $\tan\beta$ have been defined earlier. The signs of the A parameters are also convention-dependent; see Ref. 18. Due to the appearance of the *fermion* mass in Eq. (5), one expects M_{LR} to be small compared to the diagonal squark and slepton masses, with the possible exception of the top-squark, since m_t is large, and the bottom squark and τ slepton if $\tan\beta \gg 1$. The (diagonal) L and R -type squark and slepton masses are given by [2]

$$M_{u_L}^2 = M_Q^2 + m_u^2 + m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{2}{3} \sin^2 \theta_W \right) \quad (6)$$

$$M_{u_R}^2 = M_U^2 + m_u^2 + \frac{2}{3} m_Z^2 \cos 2\beta \sin^2 \theta_W \quad (7)$$

$$M_{d_L}^2 = M_Q^2 + m_d^2 - m_Z^2 \cos 2\beta \left(\frac{1}{2} - \frac{1}{3} \sin^2 \theta_W \right) \quad (8)$$

$$M_{d_R}^2 = M_D^2 + m_d^2 - \frac{1}{3} m_Z^2 \cos 2\beta \sin^2 \theta_W \quad (9)$$

$$M_\nu^2 = M_L^2 + \frac{1}{2} m_Z^2 \cos 2\beta \quad (10)$$

$$M_{e_L}^2 = M_L^2 + m_e^2 - m_Z^2 \cos 2\beta \left(\frac{1}{2} - \sin^2 \theta_W \right) \quad (11)$$

$$M_{e_R}^2 = M_E^2 + m_e^2 - m_Z^2 \cos 2\beta \sin^2 \theta_W. \quad (12)$$

The soft-supersymmetry-breaking parameters: $M_{\tilde{Q}}$, $M_{\tilde{U}}$, $M_{\tilde{D}}$, $M_{\tilde{L}}$, and $M_{\tilde{E}}$ are unknown parameters. In the equations above, the notation of first generation fermions has been used and generational indices have been suppressed. Further complications such as intergenerational mixing are possible, although there are some constraints from the nonobservation of flavor-changing neutral currents (FCNC) [27].

Searches Full Listings

Supersymmetric Particle Searches

One way to guarantee the absence of significant FCNC's mediated by virtual supersymmetric particle exchange is to posit that the diagonal soft-supersymmetry-breaking scalar mass matrices are proportional to the unit matrix (in flavor space) at some energy scale (normally taken to be the Planck scale) [28]. Renormalization group evolution is used to determine the low-energy values for the scalar mass parameters listed above. This assumption substantially reduces the MSSM parameter freedom. For example, supersymmetric grand unified models with universal scalar masses at the Planck scale typically give [29] $M_{\tilde{L}} \approx M_{\tilde{E}} < M_{\tilde{Q}} \approx M_{\tilde{U}} \approx M_{\tilde{D}}$ with the squark masses somewhere between a factor of 1–3 larger than the slepton masses (neglecting generational distinctions). More specifically, the first two generations are thought to be nearly degenerate in mass, while $M_{\tilde{Q}_3}$ and $M_{\tilde{U}_3}$ are typically reduced by a factor of 1–3 from the other soft-supersymmetry-breaking masses because of renormalization effects due to the heavy top quark mass. As a result, four flavors of squarks (with two squark eigenstates per flavor) and \tilde{b}_R will be nearly mass-degenerate and somewhat heavier than six flavors of nearly mass-degenerate sleptons (with two per flavor for the charged sleptons and one per flavor for the sneutrinos). On the other hand, the \tilde{b}_L mass and the diagonal \tilde{t}_L and \tilde{t}_R masses are reduced compared to the common squark mass of the first two generations. In addition, third generation squark masses and τ slepton masses are sensitive to the strength of the respective \tilde{f}_L – \tilde{f}_R mixing as discussed below Eq. (5).

Two additional theoretical frameworks are often introduced to reduce further the MSSM parameter freedom [1,2,30]. The first is that of grand unified theories (GUTs) and the desert hypothesis (*i.e.* no new physics between the TeV-scale and the GUT-scale). In the absence of low-energy supersymmetry, the simplest models of this type fail because the three $SU(3) \times SU(2) \times U(1)$ gauge couplings fail to unify at a common scale [31,32]. Remarkably, in the case of the MSSM (with a supersymmetry-breaking scale of order 1 TeV or below), the three gauge couplings do unify at a common energy scale of order 10^{16} GeV (with only very mild assumptions about the GUT-scale theory) [31,33]. Unification constraints on the Higgs-fermion Yukawa couplings may also exist but are more GUT-model dependent [34]. The second theoretical framework is that of minimal supergravity theory, which can impose nontrivial constraints on the soft-supersymmetry breaking parameters. Referring to the parameter list given above Eq. (1), the Planck-scale values of the soft-supersymmetry-breaking parameters in the simplest supergravity models take the following form: (i) a universal gaugino mass $m_{1/2}$ [assuming grand unification; Eq. (4) is a consequence of this assumption]; (ii) a universal diagonal scalar mass parameter m_0 [whose consequences were described in the preceding paragraph]; (iii) a universal A -parameter, A_0 ; and (iv) three scalar Higgs mass parameters—two common diagonal squared-masses given by $|\mu_0|^2 + m_0^2$ and an off-diagonal squared-mass given by $B_0\mu_0$ (which defines the Planck-scale supersymmetry-breaking parameter B_0), where μ_0

is the Planck-scale value of the μ -parameter. As before, renormalization group evolution is used to compute the low-energy values of the supersymmetry-breaking parameters and determines the supersymmetric particle spectrum. Moreover, in this approach, electroweak symmetry breaking is induced radiatively if one of the Higgs diagonal squared-masses is forced negative by the evolution. This occurs in models with a large Higgs-top quark Yukawa coupling (*i.e.* large m_t). As a result, the two Higgs vacuum expectation values (or equivalently, m_Z and $\tan\beta$) can be expressed as a function of the Planck-scale supergravity parameters. The simplest procedure [29] is to remove μ_0 and B_0 in favor of m_Z and $\tan\beta$ (the sign of μ_0 is not fixed in this process). In this case, the MSSM spectrum and its interactions are determined by m_0 , A_0 , $m_{1/2}$, $\tan\beta$, and the sign of μ_0 (in addition to the parameters of the Standard Model). Combining both grand unification and the minimal supergravity approach yield the most constrained version of the MSSM.

Nonminimal versions of low-energy supersymmetry can also be constructed. These models add additional matter and/or gauge super-multiplets to the MSSM (at the TeV scale or below). Experimental and theoretical constraints place some restrictions on these approaches, although no comprehensive treatment has yet appeared in the literature.

References

1. H.P. Nilles, Phys. Reports **110**, 1 (1984).
2. P. Nath, R. Arnowitt, and A. H. Chamseddine, *Applied $N = 1$ Supergravity* (World Scientific, Singapore, 1984).
3. M.B. Green, J.S. Schwarz, and E. Witten, *Superstring Theory* (Cambridge University Press, Cambridge, 1987).
4. E. Witten, Nucl. Phys. **B188**, 513 (1981); S. Dimopoulos and H. Georgi, Nucl. Phys. **B193**, 150 (1981); N. Sakai, Z. Phys. **C11**, 153 (1981); R.K. Kaul, Phys. Lett. **109B**, 19 (1982).
5. L. Susskind, Phys. Reports **104**, 181 (1984).
6. H.E. Haber and G.L. Kane, Phys. Reports **117**, 75 (1985).
7. K. Inoue, A. Kakuto, H. Komatsu, and S. Takeshita, Prog. Theor. Phys. **68**, 927 (1982) [E: **70**, 330 (1983)]; **71**, 413 (1984); R. Flores and M. Sher, Ann. Phys. (NY) **148**, 95 (1983).
8. J.F. Gunion and H.E. Haber, Nucl. Phys. **B272**, 1 (1986); erratum, Nucl. Phys. **B402**, 567 (1993).
9. L. Girardello and M. Grisaru, Nucl. Phys. **B194**, 65 (1982).
10. For a recent discussion, see B. de Carlos and J.A. Casas, Phys. Lett. **B309**, 320 (1993).
11. S. Bertolini, F. Borzumati, A. Masiero, and G. Ridolfi, Nucl. Phys. **B353**, 591 (1991).
12. P. Fayet, Phys. Lett. **69B**, 489 (1977); G. Farrar and P. Fayet, Phys. Lett. **76B**, 575 (1978).
13. J. Ellis, J.S. Hagelin, D.V. Nanopoulos, K. Olive, and M. Srednicki, Nucl. Phys. **B238**, 453 (1984).
14. See, *e.g.*, S. Dimopoulos, R. Esmailzadeh, L.J. Hall, and G.D. Starkman, Phys. Rev. **D41**, 2099 (1990); H. Dreiner and G.G. Ross, Nucl. Phys. **B365**, 597 (1991).

15. P. Fayet, Phys. Lett. **84B**, 421 (1979); Phys. Lett. **86B**, 272 (1979).
16. S. Deser and B. Zumino, Phys. Rev. Lett. **38**, 1433 (1977).
17. A.B. Lahanas and D.V. Nanopoulos, Phys. Reports **145**, 1 (1987).
18. H.E. Haber, "Introductory Low-Energy Supersymmetry," in *Recent Directions in Particle Theory*, Proceedings of the 1992 Theoretical Advanced Study Institute in Particle Physics, edited by J. Harvey and J. Polchinski (World Scientific, Singapore, 1993) pp. 589-686.
19. W. Fischler, S. Paban, and S. Thomas, Phys. Lett. **B289**, 373 (1992); S.M. Barr, Int. J. Mod. Phys. **A8**, 209 (1993).
20. J.F. Gunion, H.E. Haber, G. Kane, and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley Publishing Company, Redwood City, CA, 1990).
21. H.E. Haber and R. Hempfling, Phys. Rev. Lett. **66**, 1815 (1991).
22. Y. Okada, M. Yamaguchi, and T. Yanagida, Prog. Theor. Phys. **85**, 1 (1991); J. Ellis, G. Ridolfi, and F. Zwirner, Phys. Lett. **B257**, 83 (1991).
23. R. Barbieri and M. Frigeni, Phys. Lett. **B258**, 395 (1991); J. Ellis, G. Ridolfi, and F. Zwirner, Phys. Lett. **B262**, 477 (1991); P.H. Chankowski, S. Pokorski, and J. Rosiek, Phys. Lett. **B286**, 307 (1992); A. Yamada, Mod. Phys. Lett. **A7**, 2877 (1992).
24. Explicit forms for the chargino and neutralino mass matrices can be found in Appendix A of Ref. 8; see also Ref. 18.
25. See, e.g., D. Decamp *et al.* [ALEPH Collaboration], Phys. Reports **216**, 253 (1992).
26. J. Ellis and S. Rudaz, Phys. Lett. **128B**, 248 (1983).
27. F. Gabbiani and A. Masiero, Nucl. Phys. **B322**, 235 (1989)..
28. For a recent discussion, see M. Dine, R. Leigh, and A. Kagan, Phys. Rev. **D48**, 4269 (1993). For an alternate approach, see Y. Nir and N. Seiberg, Phys. Lett. **B309**, 337 (1993).
29. Recent works include: M. Drees and M.M. Nojiri, Nucl. Phys. **B369**, 54 (1992); G.G. Ross and R.G. Roberts, Nucl. Phys. **B377**, 571 (1992); S. Kelley, J.L. Lopez, D.V. Nanopoulos, H. Pois, and K. Yuan, Nucl. Phys. **B398**, 3 (1993); M. Olechowski and S. Pokorski, Nucl. Phys. **B404**, 590 (1993); D.J. Castaño, E.J. Piard, and P. Ramond, Phys. Rev. **D49**, 4882 (1994); V. Barger, M.S. Berger, and P. Ohmann, Phys. Rev. **D49**, 4908 (1994); G.L. Kane, C. Kolda, L. Roszkowski, and J.D. Wells, UM-TH-93-24 (1993).
30. For recent reviews, see R. Arnowitt and P. Nath, CTP-TAMU-52/93 (1993); W. de Boer, IEKP-KA/94-01 (1994).
31. U. Amaldi, W. de Boer, and H. Furstenau, Phys. Lett. **B260**, 447 (1991); U. Amaldi *et al.*, Phys. Lett. **B281**, 374 (1992).
32. H. Arason *et al.*, Phys. Rev. **D46**, 3945 (1992).
33. J. Ellis, S. Kelly, and D.V. Nanopoulos, Nucl. Phys. **B373**, 55 (1992); F. Anselmo, L. Cifarelli, A. Peterman, and A. Zichichi, Nuovo Cimento **105A**, 1817 (1992); **105A**, 581 (1992); **105A**, 1201 (1992); P. Langacker and N. Polonsky, Phys. Rev. **D47**, 4028 (1993); M. Carena, S. Pokorski, and C.E.M. Wagner, Nucl. Phys. **B406**, 59 (1993); V. Barger, M.S. Berger, and P. Ohmann, Phys. Rev. **D47**, 1093 (1993).
34. See, e.g.: B. Anantharayan, G. Lazarides, and Q. Shafi, Phys. Rev. **D44**, 1613 (1991); S. Dimopoulos, L.J. Hall, and S. Raby, Phys. Rev. **D45**, 4192 (1992); L.J. Hall, R. Rattazzi, and U. Sarid, LBL-33997 (1993).

MINIMAL SUPERSYMMETRIC STANDARD MODEL ASSUMPTIONS

All results shown below (except where stated otherwise) are based on the Minimal Supersymmetric Standard Model (MSSM) as described in the Note on Supersymmetry. This includes the assumption that R -parity is conserved. In addition the following assumptions are made in most cases:

- 1) The $\tilde{\chi}_1^0$ (or $\tilde{\gamma}$) is the lightest supersymmetric particle (LSP).
- 2) $m_{\tilde{L}} = m_{\tilde{R}}$ where \tilde{L} and \tilde{R} refer to the scalar partners of left- and right-handed fermions.

Limits involving different assumptions either are identified with comments or are in the miscellaneous section.

When needed, specific assumptions of the eigenstate content of neutralinos and charginos are indicated (use of the notation $\tilde{\gamma}$ (photino), \tilde{H} (Higgsino), \tilde{W} (w -ino), and \tilde{Z} (z -ino) indicates the approximation of a pure state was made).

$\tilde{\chi}_1^0$ (Lightest Neutralino) MASS LIMIT

$\tilde{\chi}_1^0$ is likely to be the lightest supersymmetric particle (LSP). See also the $\tilde{\chi}_2^0$, $\tilde{\chi}_3^0$, $\tilde{\chi}_4^0$ section below.

We have divided the $\tilde{\chi}_1^0$ listings below into three sections: 1) Accelerator limits for $\tilde{\chi}_1^0$, 2) Bounds on $\tilde{\chi}_1^0$ from dark matter searches, and 3) Other bounds on $\tilde{\chi}_1^0$ from astrophysics and cosmology.

Accelerator limits for $\tilde{\chi}_1^0$

These papers generally exclude regions in the $M_2 - \mu$ parameter plane based on accelerator experiments. Unless otherwise stated, these papers assume minimal supersymmetry and GUT relations (gaugino-mass unification condition).

VALUE (GeV)	CL %	DOCUMENT ID	TECN	COMMENT
>20	95	1 DECAMP	92 ALEP	$\tilde{\chi}_1^0, \tan\beta > 3$
>18.4	90	2 HIDAKA	91 RVUE	$\tilde{\chi}_1^0$
• • •				We do not use the following data for averages, fits, limits, etc. • • •
>18.8		3 BAER	91 RVUE	$\tilde{\chi}_1^0; \tan\beta > 1.6$
> (10-13)	90	4 ROSZKOWSKI	90 RVUE	$\tilde{\chi}_1^0; \tan\beta \geq 1$
>5 GeV	90	5 HEARTY	89 ASP	$\tilde{\gamma}$; for $m_{\tilde{e}} < 55$ GeV

¹ DECAMP 92 limit for $\tan\beta > 2$ is $m > 13$ GeV.

² HIDAKA 91 limit obtained from LEP and preliminary CDF results (as analyzed in BAER 91).

³ BAER 91 limit obtained from LEP and preliminary CDF results assuming $\tan\beta > 1.6$.

⁴ ROSZKOWSKI 90 limit obtained from ALEPH and CDF/UA2 results assuming $\tan\beta \geq 1$.

⁵ HEARTY 89 assumed pure $\tilde{\gamma}$ eigenstate and $m_{\tilde{e}_L} = m_{\tilde{e}_R}$. There is no limit for $m_{\tilde{e}} > 58$ GeV. Uses $e^+ e^- \rightarrow \gamma \tilde{\gamma} \tilde{\gamma}$. No GUT relation assumptions are made.

Bounds on $\tilde{\chi}_1^0$ from dark matter searches

These papers generally exclude regions in the $M_2 - \mu$ parameter plane assuming that $\tilde{\chi}_1^0$ is the dominant form of dark matter in the galactic halo. These limits are based on the lack of detection in laboratory experiments or by the absence of a signal in underground neutrino detectors. The latter signal is expected if $\tilde{\chi}_1^0$ accumulates in the Sun or the Earth and annihilates into high-energy ν 's.

VALUE	DOCUMENT ID	TECN
• • •		We do not use the following data for averages, fits, limits, etc. • • •
6	MORI	93 KAMI
7	BOTTINO	92 COSM
8	BOTTINO	91 RVUE
9	GELMINI	91 COSM
10	KAMIONKOW.	91 RVUE
11	MORI	91B KAMI
12	OLIVE	88 COSM

none 4-15 GeV

Searches Full Listings

Supersymmetric Particle Searches

- 6 MORI 93 excludes some region in $M_2 - \mu$ parameter space depending on $\tan\beta$ and lightest scalar Higgs mass for neutralino dark matter $m_{\tilde{\chi}_0^0} > m_W$, using limits on upgoing muons produced by energetic neutrinos from neutralino annihilation in the Sun and the Earth.
- 7 BOTTINO 92 excludes some region $M_2 - \mu$ parameter space assuming that the lightest neutralino is the dark matter, using upgoing muons at Kamiokande, direct searches by Ge detectors, and by LEP experiments. The analysis includes top radiative corrections on Higgs parameters and employs two different hypotheses for nucleon-Higgs coupling. Effects of rescalling in the local neutralino density according to the neutralino relic abundance are taken into account.
- 8 BOTTINO 91 excluded a region in $M_2 - \mu$ plane using upgoing muon data from Kamioka experiment, assuming that the dark matter surrounding us is composed of neutralinos and that the Higgs boson.
- 9 GELMINI 91 exclude a region in $M_2 - \mu$ plane using dark matter searches.
- 10 KAMIONKOWSKI 91 excludes a region in the $M_2 - \mu$ plane using IMB limit on upgoing muons originated by energetic neutrinos from neutralino annihilation in the sun, assuming that the dark matter is composed of neutralinos and that $m_{H_1^0} \lesssim 50$ GeV. See Fig. 8 in the paper.
- 11 MORI 91B exclude a part of the region in the $M_2 - \mu$ plane with $m_{\tilde{\chi}_0^0} \lesssim 80$ GeV using a limit on upgoing muons originated by energetic neutrinos from neutralino annihilation in the earth, assuming that the dark matter surrounding us is composed of neutralinos and that $m_{H_1^0} \lesssim 80$ GeV.
- 12 OLIVE 88 result assumes that photinos make up the dark matter in the galactic halo. Limit is based on annihilations in the sun and is due to an absence of high energy neutrinos detected in underground experiments. The limit is model dependent.

Other bounds on $\tilde{\chi}_1^0$ from astrophysics and cosmology

Most of these papers generally exclude regions in the $M_2 - \mu$ parameter plane by requiring that the $\tilde{\chi}_1^0$ contribution to the overall cosmological density is less than some maximal value to avoid overclosure of the Universe. Those not based on the cosmological density are indicated. Many of these papers also include LEP and/or other bounds.

VALUE	DOCUMENT ID	TECN	COMMENT
none 100 eV - 15 GeV	SREDNICKI 88	COSM	$\tilde{\gamma}; m_{\tilde{\gamma}}=100$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	DREES 93	COSM	Minimal supergravity
	FALK 93	COSM	Sfermion mixing
	KELLEY 93	COSM	Minimal supergravity
	MIZUTA 93	COSM	Co-annihilation
	ELLIS 92f	COSM	Minimal supergravity
	KAWASAKI 92	COSM	Minimal supergravity, $m_0=A=0$
	LOPEZ 92	COSM	Minimal supergravity, $m_0=A=0$
	MCDONALD 92	COSM	
	NOJIRI 91	COSM	Minimal supergravity
13 OLIVE 91	COSM		
	ROSZKOWSKI 91	COSM	
	ELLIS 90	COSM	
14 GRIEST 90	COSM		
15 GRIFOLS 90	ASTR	$\tilde{\gamma};$ SN 1987A	
	KRAUSS 90	COSM	
13 OLIVE 89	COSM		
16 ELLIS 88b	ASTR	$\tilde{\gamma};$ SN 1987A	
none 100 eV - (5-7) GeV	SREDNICKI 88	COSM	$\tilde{\gamma}; m_{\tilde{\gamma}}=60$ GeV
none 100 eV-5 GeV	ELLIS 84	COSM	$\tilde{\gamma};$ for $m_{\tilde{\gamma}}=100$ GeV
	GOLDBERG 83	COSM	$\tilde{\gamma}$
	KRAUSS 83	COSM	$\tilde{\gamma}$
	VYSOTSKII 83	COSM	$\tilde{\gamma}$

- 13 Mass of the bino (=LSP) is limited to $m_{\tilde{B}} \lesssim 350$ GeV. Mass of the higgsino (=LSP) is limited to $m_{\tilde{H}} \lesssim 1$ TeV.
- 14 Mass of the bino (=LSP) is limited to $m_{\tilde{B}} \lesssim 550$ GeV. Mass of the higgsino (=LSP) is limited to $m_{\tilde{H}} \lesssim 3.2$ TeV.
- 15 GRIFOLS 90 argues that SN1987A data exclude a light photino ($\lesssim 1$ MeV) if $m_{\tilde{q}} < 1.1$ TeV, $m_{\tilde{e}} < 0.83$ TeV.
- 16 ELLIS 88b argues that the observed neutrino flux from SN 1987A is inconsistent with a light photino if $60 \text{ GeV} \lesssim m_{\tilde{q}} \lesssim 2.5$ TeV. If $m(\text{higgsino})$ is $O(100 \text{ eV})$ the same argument leads to limits on the ratio of the two Higgs v.e.v.'s. LAU 93 discusses possible relations of ELLIS 88b bounds.
- 17 KRAUSS 83 finds $m_{\tilde{\gamma}}$ not 30 eV to 2.5 GeV. KRAUSS 83 takes into account the gravitino decay. Find that limits depend strongly on reheated temperature. For example a new allowed region $m_{\tilde{\gamma}} = 4-20$ MeV exists if $m_{\text{gravitino}} < 40$ TeV. See figure 2.

$\tilde{\chi}_2^0, \tilde{\chi}_3^0, \tilde{\chi}_4^0$ (Neutralinos) MASS LIMITS

Neutralinos are unknown mixtures of photinos, z-inos, and neutral higgsinos (the supersymmetric partners of photons and of Z and Higgs bosons). The limits here apply only to $\tilde{\chi}_2^0, \tilde{\chi}_3^0,$ and $\tilde{\chi}_4^0$. $\tilde{\chi}_1^0$ is the lightest supersymmetric particle (LSP); see $\tilde{\chi}_1^0$ Mass Limits. It is not possible to quote rigorous mass limits because they are extremely model dependent; i.e. they depend on branching ratios of various $\tilde{\chi}_0^0$ decay modes, on the masses of decay products ($\tilde{e}, \tilde{\gamma}, \tilde{q}, \tilde{g}$), and on the \tilde{e} mass exchanged in $e^+e^- \rightarrow \tilde{\chi}_i^0 \tilde{\chi}_j^0$. Often limits are given as contour plots in the $m_{\tilde{\chi}_0^0} - m_{\tilde{e}}$ plane vs other parameters. When specific assumptions are made, e.g. the neutralino is a

pure photino ($\tilde{\gamma}$), pure z-ino (\tilde{Z}), or pure neutral higgsino (\tilde{H}^0), the neutralinos will be labelled as such.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 45	95	18 DECAMP	92 ALEP	$\tilde{\chi}_2^0, \tan\beta > 3$
> 45	95	19 HIDAKA	91 RVUE	$\tilde{\chi}_2^0$
> 70	95	19 HIDAKA	91 RVUE	$\tilde{\chi}_3^0$
>108	95	19 HIDAKA	91 RVUE	$\tilde{\chi}_4^0$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
		20 ABREU	90G DLPH	$Z \rightarrow \tilde{\chi}_0^0 \tilde{\chi}_0^0$
		21 AKRAWY	90N OPAL	$Z \rightarrow \tilde{\chi}_0^0 \tilde{\chi}_0^0$
> 57	90	22 BAER	90 RVUE	$\tilde{\chi}_3^0; \Gamma(Z); \tan\beta > 1$
		23 BARKLOW	90 MRK2	$Z \rightarrow \tilde{\chi}_0^0 \tilde{\chi}_0^0, \tilde{\chi}_1^0 \tilde{\chi}_2^0$
		24 DECAMP	90K ALEP	$Z \rightarrow \tilde{\chi}_0^0 \tilde{\chi}_0^0$
> 41	95	25 SAKAI	90 AMY	$e^+e^- \rightarrow \tilde{H}_1^0 \tilde{H}_2^0$ ($\tilde{H}_2^0 \rightarrow \tilde{t}\tilde{t} \tilde{H}_1^0$)
> 31	95	26 BEHREND	87B CELL	$e^+e^- \rightarrow \tilde{\gamma}\tilde{Z}$ ($\tilde{Z} \rightarrow q\bar{q}\tilde{\gamma}$), $m_{\tilde{e}} < 70$ GeV
> 30	95	27 BEHREND	87B CELL	$e^+e^- \rightarrow \tilde{\gamma}\tilde{Z}$ ($\tilde{Z} \rightarrow q\bar{q}\tilde{g}$)
> 31.3	95	28 BEHREND	87B CELL	$e^+e^- \rightarrow \tilde{H}_1^0 \tilde{H}_2^0$ ($\tilde{H}_2^0 \rightarrow \tilde{t}\tilde{t} \tilde{H}_1^0$)
> 22	95	29 BEHREND	87B CELL	$e^+e^- \rightarrow \tilde{\gamma}\tilde{Z}$ ($\tilde{Z} \rightarrow \tilde{\nu}\nu$)
		30 AKERLOF	85 HRS	$e^+e^- \rightarrow \tilde{\gamma}\tilde{\chi}_0^0$ ($\tilde{\chi}_0^0 \rightarrow q\bar{q}\tilde{\gamma}$)
none 1-21	95	31 BARTEL	85L JADE	$e^+e^- \rightarrow \tilde{H}_1^0 \tilde{H}_2^0$ $\tilde{H}_2^0 \rightarrow \tilde{t}\tilde{t} \tilde{H}_1^0$
		32 BEHREND	85 CELL	$e^+e^- \rightarrow$ monojet X
> 35	95	33 ADEVA	84B MRKJ	$e^+e^- \rightarrow \tilde{\gamma}\tilde{Z}$ ($\tilde{Z} \rightarrow \ell\bar{\ell}\tilde{\gamma}$)
> 28	95	34 BARTEL	84C JADE	$e^+e^- \rightarrow \tilde{\gamma}\tilde{Z}$ ($\tilde{Z} \rightarrow \tilde{t}\tilde{t}\tilde{\gamma}$)
		35 ELLIS	84 COSM	

- 18 DECAMP 92 result is within minimal supersymmetry with gaugino-mass unification condition. For $\tan\beta > 2$ the limit is >40 GeV; and it disappears for $\tan\beta < 1.6$.
- 19 HIDAKA 91 limit obtained from LEP and preliminary CDF results (as analyzed in BAER 91) within minimal supersymmetry with gaugino-mass unification condition.
- 20 ABREU 90G exclude $B(Z \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0) \geq 10^{-3}$ and $B(Z \rightarrow \tilde{\chi}_2^0 \tilde{\chi}_2^0) \geq 2 \times 10^{-3}$ assuming $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \tilde{t}\tilde{t}$ via virtual Z. These exclude certain regions in model parameter space, see their Fig. 5.
- 21 AKRAWY 90N exclude $B(Z \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0) \gtrsim 3-5 \times 10^{-4}$ assuming $\tilde{\chi}_2^0 \rightarrow \tilde{\chi}_1^0 \tilde{t}\tilde{t}$ or $\tilde{\chi}_1^0 \tilde{\gamma}$ for most accessible masses. These exclude certain regions in model parameter space, see their Fig. 7.
- 22 BAER 90 is independent of decay modes. Limit from analysis of supersymmetric parameter space restrictions implied by $\Delta\Gamma(Z) < 120$ MeV. These result from decays of Z to all combinations of $\tilde{\chi}_i^\pm$ and $\tilde{\chi}_j^0$. Minimal supersymmetry with $\tan\beta > 1$ is assumed.
- 23 See Figs. 4, 5 in BARKLOW 90 for the excluded regions.
- 24 DECAMP 90K exclude certain regions in model parameter space, see their figures.
- 25 SAKAI 90 assume $m_{\tilde{H}_1^0} = 0$. The limit is for $m_{\tilde{H}_2^0}$.
- 26 Pure $\tilde{\gamma}$ and pure \tilde{Z} eigenstates. $B(\tilde{Z} \rightarrow q\bar{q}\tilde{\gamma}) = 0.60$ and $B(\tilde{Z} \rightarrow e^+e^-\tilde{\gamma}) = 0.13$. $m_{\tilde{e}_L} = m_{\tilde{e}_R} < 70$ GeV. $m_{\tilde{\gamma}} < 10$ GeV.
- 27 Pure $\tilde{\gamma}$ and pure \tilde{Z} eigenstates. $B(\tilde{Z} \rightarrow q\bar{q}\tilde{g}) = 1$. $m_{\tilde{e}_L} = m_{\tilde{e}_R} < 70$ GeV. $m_{\tilde{\gamma}} = 0$.
- 28 Pure higgsino. The LSP is the other higgsino and is taken massless. Limit degraded if $\tilde{\chi}_0^0$ not pure higgsino or if LSP not massless.
- 29 Pure $\tilde{\gamma}$ and pure \tilde{Z} eigenstates. $B(\tilde{Z} \rightarrow \tilde{\nu}\nu) = 1$. $m_{\tilde{e}_L} = m_{\tilde{e}_R} = 26$ GeV. $m_{\tilde{\gamma}} \approx 10$ GeV. No excluded region remains for $m_{\tilde{e}} > 30$ GeV.
- 30 AKERLOF 85 is e^+e^- monojet search motivated by UA1 monojet events. Observed only one event consistent with $e^+e^- \rightarrow \tilde{\gamma}\tilde{\chi}_0^0$ where $\tilde{\chi}_0^0 \rightarrow$ monojet. Assuming that missing- p_T is due to $\tilde{\gamma}$, and monojet due to $\tilde{\chi}_0^0$, limits dependent on the mixing and $m_{\tilde{e}}$ are given, see their figure 4.
- 31 BARTEL 85L assume $m_{\tilde{H}_1^0} = 0, \Gamma(Z \rightarrow \tilde{H}_1^0 \tilde{H}_2^0) \gtrsim \frac{1}{2} \Gamma(Z \rightarrow \nu_e \bar{\nu}_e)$. The limit is for $m_{\tilde{H}_2^0}$.
- 32 BEHREND 85 find no monojet at $E_{cm} = 40-46$ GeV. Consider $\tilde{\chi}_0^0$ pair production via Z^0 . One is assumed as massless and escapes detector. Limit is for the heavier one, decaying into a jet and massless $\tilde{\chi}_0^0$. Both $\tilde{\chi}_0^0$'s are assumed to be pure higgsino. For these very model-dependent results, BEHREND 85 excludes $m = 1.5-19.5$ GeV.
- 33 ADEVA 84B observed no events with signature of acoplanar lepton pair with missing energy. Above example limit is for $m_{\tilde{\gamma}} < 2$ GeV and $m_{\tilde{e}} < 40$ GeV, and assumes $B(\tilde{Z} \rightarrow \mu^+\mu^-\tilde{\gamma}) = B(\tilde{Z} \rightarrow e^+e^-\tilde{\gamma}) = 0.10$. BR = 0.05 gives 33.5 GeV limit.
- 34 BARTEL 84C search for $e^+e^- \rightarrow \tilde{Z} + \tilde{\gamma}$ with $\tilde{Z} \rightarrow \tilde{\gamma} + e^+e^-, \mu^+\mu^-, q\bar{q}$, etc. They see no acoplanar events with missing- p_T due to two $\tilde{\gamma}$'s. Above example limit is for $m_{\tilde{e}} = 40$ GeV and for light stable $\tilde{\gamma}$ with $B(\tilde{Z} \rightarrow e^+e^-\tilde{\gamma}) = 0.1$.
- 35 ELLIS 84 find if lightest neutralino is stable, then $m_{\tilde{\chi}_0^0}$ not 100 eV - 2 GeV (for $m_{\tilde{q}} = 40$ GeV). The upper limit depends on $m_{\tilde{q}}$ (similar to the $\tilde{\gamma}$ limit) and on nature of $\tilde{\chi}_0^0$. For pure higgsino the higher limit is 5 GeV.

$\tilde{\chi}_1^\pm, \tilde{\chi}_2^\pm$ (Charginos) MASS LIMITS

Charginos ($\tilde{\chi}^\pm$'s) are unknown mixtures of w-inos and charged higgsinos (the supersymmetric partners of W and Higgs bosons). Mass limits are relatively model dependent, so assumptions concerning branching ratios need to be specified. When specific assumptions are made, e.g. the chargino is a pure w-ino (\tilde{W}) or pure charged higgsino (\tilde{H}^\pm), the charginos will be labelled as such.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>44.0	95	36 ADRIANI	93M L3	$Z \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$, $\Gamma(Z)$
>46.2	95	37 DECAMP	92 ALEP	$Z \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$, all $m_{\tilde{\chi}_1^0}$
>47	95	37 DECAMP	92 ALEP	$Z \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$, $m_{\tilde{\chi}_1^0} < 41$ GeV
>99	95	38 HIDAKA	91 RVUE	$\tilde{\chi}_2^\pm$
>44.5	95	39 ABREU	90G DLPH	$Z \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$, $m_{\tilde{\chi}_1^0} < 20$ GeV
>45	95	40 AKRAWY	90D OPAL	$e^+ e^- \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$; $m_{\tilde{\chi}_1^0} < 20$ GeV
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>43	90	41 DATTA	92 RVUE	$Z \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$, $\tilde{\chi}_1^0 \tilde{\chi}_1^0$
>45	95	42 DREES	91 RVUE	$\tilde{\chi}_1^\pm$
>28.2	95	43 ADACHI	90C TOPZ	Stable $\tilde{\chi}_1^\pm$, $\tilde{\chi}^+ \tilde{\chi}^-$
>45	95	43 AKESSON	90B UA2	$p\bar{p} \rightarrow ZX$ ($Z \rightarrow \tilde{W}^+ \tilde{W}^-$)
>37	90	44 BAER	90 RVUE	$\Gamma(Z)$; $\tan\beta > 1$
>45	95	45 BARKLOW	90 MRK2	$Z \rightarrow \tilde{W}^+ \tilde{W}^-$
>42	95	46 BARKLOW	90 MRK2	$Z \rightarrow \tilde{H}^+ \tilde{H}^-$
>44.5	95	47 DECAMP	90C ALEP	$e^+ e^- \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$; $m_{\tilde{\chi}_1^0} < 28$ GeV
>25.5	95	48 ADACHI	89 TOPZ	$e^+ e^- \rightarrow \tilde{\chi}^+ \tilde{\chi}^-$
>44	95	49 ADEVA	89B L3	$e^+ e^- \rightarrow \tilde{W}^+ \tilde{W}^-$, $\tilde{W} \rightarrow \ell\nu$ or $\ell\nu\tilde{\gamma}$
>45	90	50 ANSARI	87D UA2	$p\bar{p} \rightarrow ZX$ ($Z \rightarrow \tilde{W}^+ \tilde{W}^-$, $\tilde{W}^\pm \rightarrow e^\pm \tilde{\nu}$)
>40	95	51 BAER	87B RVUE	$p\bar{p} \rightarrow W/ZX$ ($W/Z \rightarrow \tilde{W}, \tilde{Z}$)

- 36 ADRIANI 93M limit from $\Delta\Gamma(Z) < 35.1$ MeV. For pure wino, the limit is 45.5 GeV.
- 37 DECAMP 92 limit is for a general $\tilde{\chi}^\pm$ (all contents).
- 38 HIDAKA 91 limit obtained from LEP and preliminary CDF results (as analyzed in BAER 91) within minimal supersymmetry with gaugino-mass unification condition.
- 39 ABREU 90C limit is for a general $\tilde{\chi}^\pm$. They assume charginos have a three-body decay such as $\ell^+ \nu \tilde{\gamma}$.
- 40 AKRAWY 90D assume charginos have three-body decay such as $\ell^+ \nu \tilde{\gamma}$ (i.e. $m_{\tilde{\nu}} > m_{\tilde{\chi}_1^+}$). A two-body decay, $\tilde{\chi}^+ \rightarrow \ell \tilde{\nu}$ would have been seen by their search for acoplanar leptons. The result is independent of the hadronic branching ratio. They search for acoplanar electromagnetic clusters and quark jets.
- 41 DATTA 92 exclude some regions in chargino-gluino mass plane from LEP experiments.
- 42 DREES 91 limit obtained from LEP results within minimal supersymmetry with gaugino-mass unification condition. They make use of DECAMP 90C analysis plus additional constraint from total Z width. The bound can only be evaded if the chargino mixes with other charged singlets or with gauginos of a right-handed gaugino group.
- 43 AKESSON 90B assume $\tilde{W} \rightarrow e\tilde{\nu}$ with $B > 20\%$ and $m_{\tilde{\nu}} = 0$. The limit disappears if $m_{\tilde{\nu}} > 30$ GeV.
- 44 BAER 90 is independent of decay modes. Limit from analysis of supersymmetric parameter space restrictions implied by $\Delta\Gamma(Z) < 120$ MeV. These result from decays of Z to all combinations of $\tilde{\chi}_j^\pm$ and $\tilde{\chi}_1^0$. Minimal supersymmetry with $\tan\beta > 1$ is assumed.
- 45 BARKLOW 90 assume $100\% \tilde{W} \rightarrow W^* \tilde{\chi}_1^0$. Valid up to $m_{\tilde{\chi}_1^0} \lesssim [m_{\tilde{W}} - 5 \text{ GeV}]$.
- 46 BARKLOW 90 assume $100\% \tilde{H} \rightarrow H^* \tilde{\chi}_1^0$. Valid up to $m_{\tilde{\chi}_1^0} \lesssim [m_{\tilde{H}} - 8 \text{ GeV}]$.
- 47 DECAMP 90C assume charginos have three-body decay such as $\ell^+ \nu \tilde{\gamma}$ (i.e. $m_{\tilde{\nu}} > m_{\tilde{\chi}_1^+}$), and branching ratio to each lepton is 11%. They search for acoplanar dimuons, dielectrons, and μe events. Limit valid for $m_{\tilde{\chi}_1^0} < 28$ GeV.
- 48 ADACHI 89 assume only single photon annihilation in the production. The limit applies for arbitrary decay branching ratios with $B(\tilde{\chi} \rightarrow e\nu\tilde{\gamma}) + B(\tilde{\chi} \rightarrow \mu\nu\tilde{\gamma}) + B(\tilde{\chi} \rightarrow \tau\nu\tilde{\gamma}) + B(\tilde{\chi} \rightarrow q\bar{q}\tilde{\gamma}) = 1$ (lepton universality is not assumed). The limit is for $m_{\tilde{\chi}_1^0} = 0$ but a very similar limit is obtained for $m_{\tilde{\chi}_1^0} = 10$ GeV. For $B(\tilde{\chi} \rightarrow q\bar{q}\tilde{\gamma}) = 1$, the limit increases to 27.8 GeV.
- 49 ADEVA 89B assume for $\ell\nu\tilde{\gamma}$ ($\ell\nu$) mode that $B(e) = B(\mu) = B(\tau) = 11\%$ (33%) and search for acoplanar dimuons, dielectrons, and μe events. Also assume $m_{\tilde{\nu}} < 20$ GeV and for $\ell\tilde{\nu}$ mode that $m_{\tilde{\nu}} = 10$ GeV.
- 50 ANSARI 87D looks for high p_T $e^+ e^-$ pair with large missing p_T at the CERN $p\bar{p}$ collider at $E_{cm} = 546-630$ GeV. The limit is valid when $m_{\tilde{\nu}} \lesssim 20$ GeV, $B(\tilde{W} \rightarrow e\tilde{\nu}_e) = 1/3$, and $B(Z \rightarrow \tilde{W}^+ \tilde{W}^-)$ is calculated by assuming pure gaugino eigenstate. See their Fig. 3(b) for excluded region in the $m_{\tilde{W}} - m_{\tilde{\nu}}$ plane.
- 51 BAER 87B argue that the charged heavy lepton mass limit of 41 GeV obtained by UA1 collaboration (ALBAJAR 87B) corresponds to the mass limit of 40 GeV under the assumptions that the LSP (photon) has a mass smaller than 8 GeV and that the gaugino-higgsino mixing is parametrized by the three minimal supergravity model parameters. In grand unified theories $m_{\tilde{\nu}} < 8$ implies $m_{\tilde{g}} < 50$ GeV. For larger gluino masses, this limit can be evaded as discussed in BAER 88.

 $\tilde{\nu}$ (Sneutrino) MASS LIMIT

The limit depends on the number, $N(\tilde{\nu})$, of sneutrinos assumed to be degenerate in mass. Only $\tilde{\nu}_L$ (not $\tilde{\nu}_R$) exist. It is possible that $\tilde{\nu}$ could be the lightest supersymmetric particle (LSP).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>41.8	95	52 ADRIANI	93M L3	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=3$
>37.1	95	52 ADRIANI	93M L3	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=1$
>41	95	53 DECAMP	92 ALEP	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=3$
>36	95	54 ABREU	91F DLPH	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=1$
>32	95	54 ABREU	91F DLPH	$\Gamma(Z)$; $N(\tilde{\nu})=1$
>31.2	95	55 ALEXANDER	91F OPAL	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=1$
>31.4	95	56 ADEVA	90I L3	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=1$
>39.4	95	56 ADEVA	90I L3	$\Gamma(Z \rightarrow \text{invisible})$; $N(\tilde{\nu})=3$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>38.4	90	57 DREES	91 RVUE	$\Gamma(Z)$; $N(\tilde{\nu})=3$
>28.9	90	57 DREES	91 RVUE	$\Gamma(Z)$; $N(\tilde{\nu})=1$
none 3-90	90	58 SATO	91 KAMI	Stable $\tilde{\nu}_e$ or $\tilde{\nu}_\mu$, dark matter
none 4-90	90	58 SATO	91 KAMI	Stable $\tilde{\nu}_\tau$, dark matter
>36.5	90	59 BAER	90 RVUE	$\Gamma(Z)$; $N(\tilde{\nu})=3$
52 ADRIANI 93M limit from $\Delta\Gamma(Z)$ (invisible) < 16.2 MeV.				
53 DECAMP 92 limit is from $\Gamma(\text{invisible})/\Gamma(\ell\ell) = 5.91 \pm 0.15$ ($N_\nu = 2.97 \pm 0.07$).				
54 ABREU 91F limit (>32 GeV) is independent of sneutrino decay mode.				
55 ALEXANDER 91F limit is for one species of $\tilde{\nu}$ and is derived from $\Gamma(\text{invisible, new})/\Gamma(\ell\ell) < 0.38$.				
56 ADEVA 90I limit is from $\Delta N_\nu < 0.19$.				
57 DREES 91 limits from $\Delta\Gamma(Z)$ (nonhadronic) < 38.3 MeV. Independent of decay modes. Minimal supersymmetry assumed.				
58 SATO 91 search for high-energy neutrinos from the sun produced by annihilation of sneutrinos in the sun. Sneutrinos are assumed to be stable and to constitute dark matter in our galaxy. SATO 91 follow the analysis of NG 87, OLIVE 88, and GAISSER 86.				
59 BAER 90 limit from $\Delta\Gamma(Z)$ (nonhadronic) < 53 MeV. Independent of decay modes. Minimal supersymmetry assumed. The 95%CL bound is 35.6 GeV.				

 \tilde{e} (Selectron) MASS LIMIT

Limits assume $m_{\tilde{e}_L} = m_{\tilde{e}_R}$ unless otherwise stated.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 45	95	60 ADRIANI	93M L3	$m_{\tilde{\chi}_1^0} < 40$ GeV, $\tilde{e}^+ \tilde{e}^-$
> 45	95	61 DECAMP	92 ALEP	$m_{\tilde{\chi}_1^0} < 41$ GeV, $\tilde{e}^+ \tilde{e}^-$
> 65	95	62,63 HEARTY	89 RVUE	$m_{\tilde{\chi}_1^0} = 0$; $\gamma\tilde{\gamma}\tilde{\gamma}$
> 50	95	HEARTY	89 ASP	$m_{\tilde{\chi}_1^0} < 5$ GeV; $\gamma\tilde{\gamma}\tilde{\gamma}$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 42	95	ABREU	90G DLPH	$m_{\tilde{\chi}_1^0} < 40$ GeV; $\tilde{e}^+ \tilde{e}^-$
> 38	95	64 AKESSON	90B UA2	$m_{\tilde{\chi}_1^0} = 0$; $p\bar{p} \rightarrow ZX$ ($Z \rightarrow \tilde{e}^+ \tilde{e}^-$)
> 43.4	95	65 AKRAWY	90D OPAL	$m_{\tilde{\chi}_1^0} < 30$ GeV; $\tilde{e}^+ \tilde{e}^-$
> 38.1	90	66 BAER	90 RVUE	\tilde{e}_L ; $\Gamma(Z)$; $\tan\beta > 1$
> 43.5	95	67 DECAMP	90C ALEP	$m_{\tilde{\chi}_1^0} < 36$ GeV; $\tilde{e}^+ \tilde{e}^-$
>830	90	GRIFOLS	90 ASTR	$m_{\tilde{\chi}_1^0} < 1$ MeV
> 29.9	95	SAKAI	90 AMY	$m_{\tilde{\chi}_1^0} < 20$ GeV; $\tilde{e}^+ \tilde{e}^-$
> 29	95	TAKETANI	90 VNS	$m_{\tilde{\chi}_1^0} < 25$ GeV; $\tilde{e}^+ \tilde{e}^-$
> 60	90	68 ZHUKOVSKII	90 ASTR	$m_{\tilde{\chi}_1^0} = 0$
> 32	90	69 ABE	89K VNS	$e^+ e^- \rightarrow \gamma\tilde{\gamma}\tilde{\gamma}$
> 28	95	70 ADACHI	89 TOPZ	$m_{\tilde{\chi}_1^0} \lesssim 0.85 m_{\tilde{e}}; \tilde{e}^+ \tilde{e}^-$
> 41	95	71 ADEVA	89B L3	$m_{\tilde{\chi}_1^0} < 20$ GeV; $\tilde{e}^+ \tilde{e}^-$
> 32	90	72 ALBAJAR	89 UA1	$p\bar{p} \rightarrow W^\pm X$ ($W^\pm \rightarrow \tilde{e}_L \tilde{\nu}$) ($\tilde{e}_L \rightarrow e\tilde{\gamma}$)
> 14	90	73 ALBAJAR	89 UA1	$Z \rightarrow \tilde{e}^+ \tilde{e}^-$
> 53	95	62,74 HEARTY	89 ASP	$m_{\tilde{\chi}_1^0} = 0$; $\gamma\tilde{\gamma}\tilde{\gamma}$
> 35	95	HEARTY	89 ASP	$m_{\tilde{\chi}_1^0} < 10$ GeV; $\gamma\tilde{\gamma}\tilde{\gamma}$
> 51.5	90	75,76 BEHREND	88B CELL	$m_{\tilde{\chi}_1^0} = 0$ GeV; $\gamma\tilde{\gamma}\tilde{\gamma}$
> 64	95	75,77 BEHREND	88B RVUE	$m_{\tilde{\chi}_1^0} = 0$ GeV; $\gamma\tilde{\gamma}\tilde{\gamma}$
> 48	90	BEHREND	88B CELL	$m_{\tilde{\chi}_1^0} < 5$ GeV; $\gamma\tilde{\gamma}\tilde{\gamma}$
60 ADRIANI 93M limit is for $m_{\tilde{e}_L} \gg m_{\tilde{e}_R}$ using acoplanar di-lepton events.				
61 DECAMP 92 limit is for $m_{\tilde{e}_L} \gg m_{\tilde{e}_R}$; for equal masses the limit would improve. They looked for acoplanar electrons.				
62 HEARTY 89 assume $m_{\tilde{\chi}_1^0} = 0$. The limit is very sensitive to $m_{\tilde{\chi}_1^0}$; no limit can be placed for $m_{\tilde{\chi}_1^0} \gtrsim 13$ GeV.				
63 Results of HEARTY 89, BEHREND 88B, ADEVA 87, and FORD 86 are combined. The limit is reduced to 53 GeV if only one \tilde{e} state is produced (\tilde{e}_L or \tilde{e}_R very heavy).				
64 AKESSON 90B assume $m_{\tilde{\chi}_1^0} = 0$. Very similar limits hold for $m_{\tilde{\chi}_1^0} \lesssim 20$ GeV.				
65 AKRAWY 90D look for acoplanar electrons. For $m_{\tilde{e}_L} \gg m_{\tilde{e}_R}$, limit is 41.5 GeV, for $m_{\tilde{\chi}_1^0} < 30$ GeV.				
66 BAER 90 limit from $\Delta\Gamma(Z)$ (nonhadronic) < 53 MeV. Independent of decay modes. Minimal supersymmetry and $\tan\beta > 1$ assumed.				
67 DECAMP 90C look for acoplanar electrons. For $m_{\tilde{e}_L} \gg m_{\tilde{e}_R}$ limit is 42 GeV, for $m_{\tilde{\chi}_1^0} < 33$ GeV.				

Searches Full Listings

Supersymmetric Particle Searches

- 68 ZHUKOVSKII 90 set limit by saying the luminosity of a magnetized neutron star due to massless photino emission by electrons be small compared with its neutrino luminosity.
- 69 ABE 89K assumed $m_{\tilde{\gamma}} = 0$.
- 70 ADACHI 89 assume only photon and photino exchange and $m_{\tilde{e}_L} = m_{\tilde{e}_R}$. The limit for the nondegenerate case is 26 GeV.
- 71 ADEVA 89B look for acoplanar electrons.
- 72 ALBAJAR 89 limit applies for \tilde{e}_L when $m_{\tilde{e}_L} = m_{\tilde{\nu}_L}$ and $m_{\tilde{\gamma}} = 0$. See their Fig. 55 for the 90% CL excluded region in the $m_{\tilde{e}_L} - m_{\tilde{\nu}_L}$ plane. For $m_{\tilde{\nu}} = m_{\tilde{\gamma}} = 0$, limit is 50 GeV.
- 73 ALBAJAR 89 assume $m_{\tilde{\gamma}} = 0$.
- 74 The limit is reduced to 43 GeV if only one \tilde{e} state is produced (\tilde{e}_L or \tilde{e}_R very heavy).
- 75 BEHREND 88B limits assume pure photino eigenstate and $m_{\tilde{e}_L} = m_{\tilde{e}_R}$.
- 76 The 95% CL limit for BEHREND 88B is 47.5 GeV for $m_{\tilde{\gamma}} = 0$. The limit for $m_{\tilde{e}_L} \gg m_{\tilde{e}_R}$ is 40 GeV at 90% CL.
- 77 BEHREND 88B combined their data with those from ASP (HEARTY 87), MAC (FORD 86), and MARK-J (H. Wu, Ph. D. Thesis, University of Hamburg, 1986).

 $\tilde{\mu}$ (Smuon) MASS LIMITLimits assume $m_{\tilde{\mu}_L} = m_{\tilde{\mu}_R}$ unless otherwise stated.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45	95	78 ADRIANI	93M L3	$m_{\tilde{\chi}_1^0} < 40$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>45	95	79 DECAMP	92 ALEP	$m_{\tilde{\chi}_1^0} < 41$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>43	95	80 AKRAWY	90D OPAL	$m_{\tilde{\gamma}} < 30$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>36	95	ABREU	90G DLPH	$m_{\tilde{\gamma}} < 33$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>38.1	90	81 BAER	90 RVUE	$\tilde{\mu}_L; \Gamma(Z); \tan\beta > 1$
>42.6	95	82 DECAMP	90C ALEP	$m_{\tilde{\gamma}} < 34$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>27	95	SAKAI	90 AMY	$m_{\tilde{\gamma}} < 18$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>24.5	95	TAKETANI	90 VNS	$m_{\tilde{\gamma}} < 15$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$
>24.5	95	83 ADACHI	89 TOPZ	$m_{\tilde{\gamma}} \lesssim 0.8 m_{\tilde{\mu}}; \tilde{\mu}^+ \tilde{\mu}^-$
>41	95	84 ADEVA	89B L3	$m_{\tilde{\gamma}} < 20$ GeV; $\tilde{\mu}^+ \tilde{\mu}^-$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 78 ADRIANI 93M limit is for $m_{\tilde{\mu}_L} \gg m_{\tilde{\mu}_R}$ using acoplanar di-lepton events.
- 79 DECAMP 92 limit is for $m_{\tilde{\mu}_L} \gg m_{\tilde{\mu}_R}$; for equal masses the limit would improve. They looked for acoplanar muons.
- 80 AKRAWY 90D look for acoplanar muons. For $m_{\tilde{\mu}_L} \gg m_{\tilde{\mu}_R}$, limit is 41.0 GeV, for $m_{\tilde{\gamma}} < 30$ GeV.
- 81 BAER 90 limit from $\Delta\Gamma(Z)$ (nonhadronic) < 53 MeV. Independent of decay modes. Minimal supersymmetry and $\tan\beta > 1$ assumed.
- 82 DECAMP 90C look for acoplanar muons. For $m_{\tilde{\mu}_L} \gg m_{\tilde{\mu}_R}$ limit is 40 GeV, for $m_{\tilde{\gamma}} < 30$ GeV.
- 83 ADACHI 89 assume only photon exchange, which gives a conservative limit. $m_{\tilde{\mu}_L} = m_{\tilde{\mu}_R}$ assumed. The limit for nondegenerate case is 22 GeV.
- 84 ADEVA 89B look for acoplanar muons.

 $\tilde{\tau}$ (Stau) MASS LIMITLimits assume $m_{\tilde{\tau}_L} = m_{\tilde{\tau}_R}$ unless otherwise stated.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>44	95	85 ADRIANI	93M L3	$m_{\tilde{\chi}_1^0} < 38$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>45	95	86 DECAMP	92 ALEP	$m_{\tilde{\chi}_1^0} < 38$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>43.0	95	87 AKRAWY	90D OPAL	$m_{\tilde{\gamma}} < 23$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>35	95	ABREU	90G DLPH	$m_{\tilde{\gamma}} < 25$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>38.1	90	88 BAER	90 RVUE	$\tilde{\tau}_L; \Gamma(Z); \tan\beta > 1$
>40.4	95	89 DECAMP	90C ALEP	$m_{\tilde{\gamma}} < 15$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>25	95	SAKAI	90 AMY	$m_{\tilde{\gamma}} < 10$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>25.5	95	TAKETANI	90 VNS	$m_{\tilde{\gamma}} < 15$ GeV; $\tilde{\tau}^+ \tilde{\tau}^-$
>21.7	95	90 ADACHI	89 TOPZ	$m_{\tilde{\gamma}}=0; \tilde{\tau}^+ \tilde{\tau}^-$

- • • We do not use the following data for averages, fits, limits, etc. • • •
- 85 ADRIANI 93M limit is for $m_{\tilde{\tau}_L} \gg m_{\tilde{\tau}_R}$.
- 86 DECAMP 92 limit is for $m_{\tilde{\tau}_L} \gg m_{\tilde{\tau}_R}$; for equal masses the limit would improve. They looked for acoplanar particles.
- 87 AKRAWY 90D look for acoplanar particles. For $m_{\tilde{\tau}_L} \gg m_{\tilde{\tau}_R}$, limit is 41.0 GeV, for $m_{\tilde{\gamma}} < 23$ GeV.
- 88 BAER 90 limit from $\Delta\Gamma(Z)$ (nonhadronic) < 53 MeV. Independent of decay modes. Minimal supersymmetry and $\tan\beta > 1$ assumed.
- 89 DECAMP 90C look for acoplanar charged particle pairs. Limit is for $m_{\tilde{\tau}_L} = m_{\tilde{\tau}_R}$. For $m_{\tilde{\gamma}} \leq 24$ GeV, the limit is 37 GeV. For $m_{\tilde{\tau}_L} \gg m_{\tilde{\tau}_R}$ and $m_{\tilde{\gamma}} < 15$ GeV, the limit is 33 GeV.
- 90 ADACHI 89 assume only photon exchange, which gives a conservative limit. $m_{\tilde{\tau}_L} = m_{\tilde{\tau}_R}$ assumed.

Stable $\tilde{\ell}$ (Slepton) MASS LIMIT

Limits on scalar leptons which leave detector before decaying. Limits from Z decays are independent of lepton flavor. Limits from continuum e^+e^- annihilation are also independent of flavor for smuons and staus. However, selectron limits from continuum e^+e^- annihilation depend on flavor because there is an additional contribution from neutralino exchange that in general yields stronger limits. All limits assume $m_{\tilde{\ell}_L} = m_{\tilde{\ell}_R}$ unless otherwise stated.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>40	95	ABREU	90G DLPH	
>26.3	95	ADACHI	90C TOPZ	$\tilde{\mu}, \tilde{\tau}$
>38.8	95	AKRAWY	90D OPAL	$\tilde{\ell}_R$
>27.1	95	91 SAKAI	90 AMY	
>32.6	95	SODERSTROM	90 MRK2	
>24.5	95	92 ADACHI	89 TOPZ	

91 SAKAI 90 limit improves to 30.1 GeV for \tilde{e} if $m_{\tilde{\gamma}} \approx m_{\tilde{e}}$.92 ADACHI 89 assume only photon (and photino for \tilde{e}) exchange. The limit for \tilde{e} improves to 26 GeV for $m_{\tilde{\gamma}} \approx m_{\tilde{e}}$. \tilde{q} (Squark) MASS LIMIT

For $m_{\tilde{q}} > 60-70$ GeV, it is expected that squarks would undergo a cascade decay via a number of neutralinos and/or charginos rather than undergo a direct decay to photinos as assumed by some papers. When direct decay is assumed, realistic limits would be somewhat lower. The limits from Z decay do not assume GUT relations and are more model independent.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
> 90	90	93 ABE	92L CDF	Any $m_{\tilde{g}} < 410$ GeV; with cascade decay
> 218	90	93 ABE	92L CDF	$m_{\tilde{g}} = m_{\tilde{q}}; \tilde{q} \rightarrow \tilde{q} \tilde{q};$ with cascade decay
> 180	90	93 ABE	92L CDF	$m_{\tilde{g}} < m_{\tilde{q}}; \tilde{q} \rightarrow \tilde{q} \tilde{q};$ with cascade decay
> 35.3	95	94 ADRIANI	93M L3	$Z \rightarrow \tilde{u} \tilde{u}, \Gamma(Z)$
> 36.8	95	94 ADRIANI	93M L3	$Z \rightarrow \tilde{d} \tilde{d}, \Gamma(Z)$
> 100	95	95 ROY	92 RVUE	$p\bar{p} \rightarrow \tilde{q} \tilde{q}; R$ -parity violating
> 45	95	96 BAER	91B RVUE	\tilde{t}
> 45	95	97 NOJIRI	91 COSM	
> 45	95	98 ABREU	90F DLPH	$Z \rightarrow \tilde{q} \tilde{q}; m_{\tilde{\gamma}} < 20$ GeV
> 43	95	99 ABREU	90F DLPH	$Z \rightarrow \tilde{d} \tilde{d}; m_{\tilde{\gamma}} < 20$ GeV
> 42	95	100 ABREU	90F DLPH	$Z \rightarrow \tilde{u} \tilde{u}; m_{\tilde{\gamma}} < 20$ GeV
> 27.0	95	ADACHI	90C TOPZ	Stable $\tilde{u}, \tilde{u} \tilde{u}$
> 74	90	101 ALITTI	90 UA2	Any $m_{\tilde{q}}; B(\tilde{q} \rightarrow q \tilde{g} \text{ or } q \tilde{\gamma}) = 1$
> 106	90	101 ALITTI	90 UA2	$m_{\tilde{q}} = m_{\tilde{g}}; B(\tilde{q} \rightarrow q \tilde{\gamma}) = 1$
> 39.2	90	102 BAER	90 RVUE	$\tilde{d}_L; \Gamma(Z)$
> 45	95	103,104 BARKLOW	90 MRK2	$Z \rightarrow \tilde{q} \tilde{q}$
> 40	95	103,105 BARKLOW	90 MRK2	$Z \rightarrow \tilde{d} \tilde{d}$
> 39	95	103,106 BARKLOW	90 MRK2	$Z \rightarrow \tilde{u} \tilde{u}$
> 1100	107	DREES	90 RVUE	\tilde{t}
> 24	95	SAKAI	90 AMY	$e^+ e^- \rightarrow \tilde{d} \tilde{d} \rightarrow \tilde{d} \tilde{d} \tilde{\gamma} \tilde{\gamma}; m_{\tilde{\gamma}} < 10$ GeV
> 26	95	SAKAI	90 AMY	$e^+ e^- \rightarrow \tilde{u} \tilde{u} \rightarrow \tilde{u} \tilde{u} \tilde{\gamma} \tilde{\gamma}; m_{\tilde{\gamma}} < 10$ GeV
> 26.3	95	108 ADACHI	89 TOPZ	$e^+ e^- \rightarrow \tilde{q} \tilde{q} \rightarrow q \tilde{q} \tilde{\gamma} \tilde{\gamma}$
> 45	90	109 NATH	88 THEO	$\tau(p \rightarrow \nu K)$ in supergravity GUT
> 45	90	110 ALBAJAR	87D UA1	Any $m_{\tilde{g}} > m_{\tilde{q}}$
> 75	90	110 ALBAJAR	87D UA1	$m_{\tilde{g}} = m_{\tilde{q}}$

93 ABE 92L assume five degenerate squark flavors and $m_{\tilde{d}_L} = m_{\tilde{q}_R}$. ABE 92L includes the effect of cascade decay, for a particular choice of parameters, $\mu = -250$ GeV, $\tan\beta = 2$. Results are weakly sensitive to these parameters over much of parameter space. No limit for $m_{\tilde{q}} \leq 50$ GeV (but other experiments rule out that region). Limits are 10-20 GeV higher if $B(\tilde{q} \rightarrow q \tilde{\gamma}) = 1$. Limit assumes GUT relations between gaugino masses and the gauge coupling; in particular that for $|\mu|$ not small, $m_{\tilde{\chi}_1^0} \approx m_{\tilde{g}}/6$. This last

relation implies that as $m_{\tilde{g}}$ increases, the mass of $\tilde{\chi}_1^0$ will eventually exceed $m_{\tilde{q}}$ so that no decay is possible. Even before that occurs, the signal will disappear; in particular no bounds can be obtained for $m_{\tilde{g}} > 410$ GeV.

94 ADRIANI 93M limit from $\Delta\Gamma(Z) < 35.1$ MeV and assumes $m_{\tilde{q}_L} \gg m_{\tilde{q}_R}$.95 ROY 92 reanalyzed CDF limits on di-lepton events to obtain limits on squark production in R-parity violating models. The 100% decay $\tilde{q} \rightarrow q \tilde{\chi}$ where $\tilde{\chi}$ is the LSP, and the LSP decays either into $\ell q \tilde{d}$ or $\ell \ell \tilde{e}$ is assumed.

96 BAER 91B argue that a top squark as light as 45 GeV may have escaped detection at the CDF detector at the Tevatron Collider (45 GeV is the limit from LEP experiments).

- 97 NOJIRI 91 argues that a heavy squark should be nearly degenerate with the gluino in minimal supergravity not to overclose the universe.
- 98 ABREU 90F assume six degenerate squarks and $m_{\tilde{q}_L} = m_{\tilde{q}_R} = m_{\tilde{q}}$. $m_{\tilde{q}} < 41$ GeV is excluded at 95% CL for $m_{LSP} < m_{\tilde{q}} - 2$ GeV.
- 99 ABREU 90F exclude $m_{\tilde{q}} < 38$ GeV at 95% for $m_{LSP} < m_{\tilde{q}} - 2$ GeV.
- 100 ABREU 90F exclude $m_{\tilde{q}} < 36$ GeV at 95% for $m_{LSP} < m_{\tilde{q}} - 2$ GeV.
- 101 ALITTI 90 searched for events having ≥ 2 jets with $E_T^j > 25$ GeV, $E_T^{\tilde{q}} > 15$ GeV, $|\eta| < 0.85$, and $\Delta\phi < 160^\circ$, with a missing momentum > 40 GeV and no electrons. They assume $\tilde{q} \rightarrow q\tilde{\gamma}$ (if $m_{\tilde{q}} < m_{\tilde{g}}$) or $\tilde{q} \rightarrow q\tilde{g}$ (if $m_{\tilde{q}} > m_{\tilde{g}}$) decay and $m_{\tilde{\gamma}} \lesssim 20$ GeV. Five degenerate squark flavors and $m_{\tilde{q}_L} = m_{\tilde{q}_R}$ are assumed. Masses below 50 GeV are not excluded by the analysis.
- 102 BAER 90 limit from $\Delta\Gamma(Z) < 120$ MeV, assuming $m_{\tilde{d}_L} = m_{\tilde{u}_L} = m_{\tilde{e}_L} = m_{\tilde{\nu}}$. Independent of decay modes. Minimal supergravity assumed.
- 103 BARKLOW 90 assume 100% $\tilde{q} \rightarrow q\tilde{\gamma}$.
- 104 BARKLOW 90 assume five degenerate squarks (left- and right-handed). Valid up to $m_{\tilde{q}_0} \lesssim [m_{\tilde{q}} - 4 \text{ GeV}]$.
- 105 BARKLOW 90 result valid up to $m_{\tilde{q}_0} \lesssim [m_{\tilde{q}} - 5 \text{ GeV}]$.
- 106 BARKLOW 90 result valid up to $m_{\tilde{q}_0} \lesssim [m_{\tilde{q}} - 6 \text{ GeV}]$.
- 107 DREES 90 argue that bounds from Z decay are not valid for \tilde{t} for a certain range of \tilde{t}_L - \tilde{t}_R mixing angle.
- 108 ADACHI 89 assume only photon exchange, which gives a conservative limit. The limit is only for one flavor of charge $2/3$ \tilde{q} . $m_{\tilde{q}_L} = m_{\tilde{q}_R}$ and $m_{\tilde{\gamma}} = 0$ assumed. The limit decreases to 26.1 GeV for $m_{\tilde{\gamma}} = 15$ GeV. The limit for nondegenerate case is 24.4 GeV.
- 109 NATH 88 uses Kamioka limit of $\tau(\rho \rightarrow \bar{\nu}K^+) > 7 \times 10^{31}$ yrs to constrain squark mass $m_{\tilde{q}} > 1000$ GeV by assuming that the proton decay proceeds via an exchange of a color-triplet Higgsino of mass $< 10^{16}$ GeV in the supersymmetric SU(5) GUT. The limit applies for $m_{\tilde{\gamma}} \equiv (8/3) \sin^2\theta_W m_{\tilde{2}} > 10$ GeV ($m_{\tilde{2}}$ is the SU(2) gaugino mass) and for a very conservative value of the three-quark proton wave function, barring cancellation between second and third generations. Lower squark mass is allowed if $m_{\tilde{\gamma}}$ as defined above is smaller.
- 110 The limits of ALBAJAR 87D are from $p\bar{p} \rightarrow \tilde{q}\tilde{q}^*X$ ($\tilde{q} \rightarrow q\tilde{\gamma}$) and assume 5 flavors of degenerate mass squarks each with $m_{\tilde{q}_L} = m_{\tilde{q}_R}$. They also assume $m_{\tilde{g}} > m_{\tilde{q}}$. These limits apply for $m_{\tilde{\gamma}} \lesssim 20$ GeV.

 \tilde{g} (Gluino) MASS LIMIT

For $m_{\tilde{g}} > 60$ -70 GeV, it is expected that gluinos would undergo a cascade decay via a number of neutralinos and/or charginos rather than undergo a direct decay to photinos as assumed by some papers. When direct decay is assumed, realistic limits would be somewhat lower.

There is an ongoing controversy (reflected in these Listings) about whether very light \tilde{g} 's ($1 \lesssim m_{\tilde{g}} \lesssim 4$ GeV) are ruled out. These papers sometimes make different assumptions and use different calculational techniques.

VALUE (GeV)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
>218	90		111 ABE	92L CDF	$m_{\tilde{q}} \leq m_{\tilde{g}}$; with cascade decay
>100	90		111 ABE	92L CDF	Any $m_{\tilde{q}}$; with cascade decay
< 1			112 CLAVELLI	93 RVUE	quarkonia
			113 HEBBEKER	93 RVUE	e^+e^- jet analyses
not 3-5			114 LOPEZ	93C RVUE	LEP
≈ 4			115 CLAVELLI	92 RVUE	α_s running
>100			116 ROY	92 RVUE	$p\bar{p} \rightarrow \tilde{g}\tilde{g}^*$; R -parity violating
> 1			117 ANTONIADIS	91 RVUE	α_s running
>132	90		118 ANTONIADIS	91 RVUE	$\rho N \rightarrow$ missing energy
			119 HIDAKA	91 RVUE	
			120 NOJIRI	91 COSM	
> 79	90		121 ALITTI	90 UA2	Any $m_{\tilde{g}}$; $B(\tilde{g} \rightarrow q\tilde{q}\tilde{\gamma}) = 1$
>106	90		121 ALITTI	90 UA2	$m_{\tilde{q}} = m_{\tilde{g}}$; $B(\tilde{g} \rightarrow q\tilde{q}\tilde{\gamma}) = 1$
none 4-53	90		122 NAKAMURA	89 SPEC	$R\text{-}\Delta^{++}$
none 4-75	90		123 ALBAJAR	87D UA1	Any $m_{\tilde{q}} > m_{\tilde{g}}$
none 16-58	90		124 ALBAJAR	87D UA1	$m_{\tilde{q}} = m_{\tilde{g}}$
none 16-58	90		124 ANSARI	87D UA2	$m_{\tilde{q}} \lesssim 100$ GeV
> 3.8	90		125 ARNOLD	87 EMUL	π^- (350 GeV). $\sigma \approx A^1$
> 3.2	90		125 ARNOLD	87 EMUL	π^- (350 GeV). $\sigma \approx A^{0.72}$
none 0.6-2.2	90		126 TUTS	87 CUSB	$\Upsilon(1S) \rightarrow \gamma + \text{gluonolium}$
none 1-4.5	90	0	127 ALBRECHT	86C ARG	$1 \times 10^{-11} \lesssim \tau \lesssim 1 \times 10^{-9}$ s
none 1-4	90	0	128 BADIÉ	86 BDMP	$1 \times 10^{-10} < \tau < 1 \times 10^{-7}$ s

- none 3-5
- 129 BARNETT 86 RVUE $p\bar{p} \rightarrow$ gluino gluino gluon
- 130 VOLOSHIN 86 RVUE If (quasi) stable; $\tilde{g}uud$
- none 0.5-2
- 131 COOPER... 85B BDMP For $m_{\tilde{q}}=300$ GeV
- none 0.5-4
- 131 COOPER... 85B BDMP For $m_{\tilde{q}} < 65$ GeV
- none 0.5-3
- 131 COOPER... 85B BDMP For $m_{\tilde{q}}=150$ GeV
- none 2-4
- 132 DAWSON 85 RVUE $\tau > 10^{-7}$ s
- none 1-2.5
- 132 DAWSON 85 RVUE For $m_{\tilde{q}}=100$ GeV
- none 0.5-4.1
- 90
- 133 FARRAR 85 RVUE FNAL beam dump
- 134 GOLDMAN 85 RVUE Gluonium
- 135 HABER 85 RVUE
- 136 BALL 84 CALO
- 137 BRICK 84 RVUE
- 138 FARRAR 84 RVUE
- 139 BERGSMAN 83C RVUE For $m_{\tilde{q}} < 100$ GeV
- > 2
- 140 CHANOWITZ 83 RVUE $\tilde{g}u\bar{d}, \tilde{g}uud$
- >2-3
- 141 KANE 82 RVUE Beam dump
- >1.5-2
- FARRAR 78 RVUE R -hadron
- 111 ABE 92L Includes the effect of cascade decay, for a particular choice of parameters, $\mu = -250$ GeV, $\tan\beta = 2$. Results are mildly sensitive to these parameters over much of parameter space. ABE 92L limit assumes GUT relations between gaugino masses and the gauge coupling; in particular that for $|\mu|$ not small, $m_{\tilde{2}} \approx m_{\tilde{1}}/6$. Not sensitive to $m_{\tilde{g}} \leq 40$ GeV (but other experiments rule out that region). Limits are more substantial if $B(\tilde{g} \rightarrow q\tilde{q}\tilde{\gamma}) = 1$; for $m_{\tilde{q}} > m_{\tilde{g}}$ the limits are about 50 GeV greater.
- 112 CLAVELLI 93 makes a two-dimensional fit to the quarkonia decay widths for $\phi(1020)$, $J/\psi(1S)$, $\psi(2S)$, $\Upsilon(2S)$, and $\Upsilon(3S)$ taking $\alpha_s(m_Z)$ and $m_{\tilde{g}}$ as variables. Claims that the fit favors $m_{\tilde{g}} < 1$ GeV, and that the fitted $\alpha_s(m_Z)$ is consistent with LEP data.
- 113 HEBBEKER 93 combined jet analyses at various e^+e^- colliders. The 4-jet analyses at TRISTAN/LEP and the measured α_s at PEP/PETRA/TRISTAN/LEP are used. A constraint on effective number of quarks $N=6.3 \pm 1.1$ is obtained, which is compared to that with a light gluino, $N=8$.
- 114 LOPEZ 93C uses combined restraint from the radiative symmetry breaking scenario within the minimal supergravity model, and the LEP bounds on the (M_2, μ) plane. Claims that the light gluino window is strongly disfavored.
- 115 CLAVELLI 92 claims that a light gluino mass around 4 GeV should exist to explain the discrepancy between α_s at LEP and at quarkonia (Υ), since a light gluino slows the running of the QCD coupling.
- 116 ROY 92 reanalyzed CDF limits on di-lepton events to obtain limits on gluino production in R -parity violating models. The 100% decay $\tilde{g} \rightarrow q\tilde{q}^*$ where $\tilde{\chi}$ is the LSP, and the LSP decays either into $l\tilde{q}$ or $l\tilde{l}$ is assumed.
- 117 ANTONIADIS 91 argue that possible light gluinos (< 5 GeV) contradict the observed running of α_s between 5 GeV and m_Z . The significance is less than 2.s.d.
- 118 ANTONIADIS 91 interpret the search for missing energy events in 450 GeV/c pN collisions, AKESSON 91, in terms of light gluinos.
- 119 HIDAKA 91 limit obtained from LEP and preliminary ARGUS results within minimal supersymmetry with gaugino-mass unification condition. HIDAKA 91 limit extracted from BAER 91 analysis.
- 120 NOJIRI 91 argues that a heavy gluino should be nearly degenerate with squarks in minimal supergravity not to overclose the universe.
- 121 ALITTI 90 searched for events having ≥ 2 jets with $E_T^j > 25$ GeV, $E_T^{\tilde{q}} > 15$ GeV, $|\eta| < 0.85$, and $\Delta\phi < 160^\circ$, with a missing momentum > 40 GeV and no electrons. They assume $\tilde{g} \rightarrow q\tilde{q}\tilde{\gamma}$ decay and $m_{\tilde{\gamma}} \lesssim 20$ GeV. Masses below 50 GeV are not excluded by the analysis.
- 122 NAKAMURA 89 searched for a long-lived ($\tau \gtrsim 10^{-7}$ s) charge- (± 2) particle with mass $\lesssim 1.6$ GeV in proton-Pt interactions at 12 GeV and found that the yield is less than 10^{-8} times that of the pion. This excludes $R\text{-}\Delta^{++}$ (a $\tilde{g}uud$ state) lighter than 1.6 GeV.
- 123 The limits of ALBAJAR 87D are from $p\bar{p} \rightarrow \tilde{g}\tilde{g}^*X$ ($\tilde{g} \rightarrow q\tilde{q}\tilde{\gamma}$) and assume $m_{\tilde{q}} > m_{\tilde{g}}$. These limits apply for $m_{\tilde{\gamma}} \lesssim 20$ GeV and $\tau(\tilde{g}) < 10^{-10}$ s.
- 124 The limit of ANSARI 87D assumes $m_{\tilde{q}} > m_{\tilde{g}}$ and $m_{\tilde{\gamma}} \approx 0$.
- 125 The limits assume $m_{\tilde{q}} = 100$ GeV. See their figure 3 for limits vs. $m_{\tilde{q}}$.
- 126 The gluino mass is defined by half the bound $\tilde{g}\tilde{g}$ mass. If zero gluino mass gives a $\tilde{g}\tilde{g}$ of mass about 1 GeV as suggested by various glueball mass estimates, then the low-mass bound can be replaced by zero. The high-mass bound is obtained by comparing the data with nonrelativistic potential-model estimates.
- 127 ALBRECHT 86C search for secondary decay vertices from $\chi_{b1}(1P) \rightarrow \tilde{g}\tilde{g}\tilde{g}$ where \tilde{g} 's make long-lived hadrons. See their figure 4 for excluded region in the $m_{\tilde{g}} - m_{\tilde{q}}$ and $m_{\tilde{g}} - m_{\tilde{q}}$ plane. The lower $m_{\tilde{g}}$ region below ~ 2 GeV may be sensitive to fragmentation effects. Remark that the \tilde{g} -hadron mass is expected to be ~ 1 GeV (glueball mass) in the zero \tilde{g} mass limit.
- 128 BADIÉ 86 looked for secondary decay vertices from long-lived \tilde{g} -hadrons produced at 300 GeV π^- beam dump. The quoted bound assumes \tilde{g} -hadron nucleon total cross section of $10\mu\text{b}$. See their figure 7 for excluded region in the $m_{\tilde{g}} - m_{\tilde{q}}$ plane for several assumed total cross-section values.
- 129 BARNETT 86 rule out light gluinos ($m = 3$ -5 GeV) by calculating the monojet rate from gluino gluino gluon events (and from gluino gluino events) and by using UA1 data from $p\bar{p}$ collisions at CERN.
- 130 VOLOSHIN 86 rules out stable gluino based on the cosmological argument that predicts too much hydrogen consisting of the charged stable hadron $\tilde{g}uud$. Quasi-stable ($\tau > 1 \times 10^{-7}$ s) light gluino of $m_{\tilde{g}} < 3$ GeV is also ruled out by nonobservation of the stable charged particles, $\tilde{g}uud$, in high energy hadron collisions.
- 131 COOPER-SARKAR 85B is BEBC beam-dump. Gluinos decaying in dump would yield $\tilde{\gamma}$'s in the detector giving neutral-current-like interactions. For $m_{\tilde{q}} > 330$ GeV, no limit is set.
- 132 DAWSON 85 first limit from neutral particle search. Second limit based on FNAL beam dump experiment.

Searches Full Listings

Supersymmetric Particle Searches

- 133 FARRAR 85 points out that BALL 84 analysis applies only if the \tilde{g} 's decay before interacting, i.e. $m_{\tilde{g}} < 80m_{\tilde{g}}^{-1.5}$. FARRAR 85 finds $m_{\tilde{g}} < 0.5$ not excluded for $m_{\tilde{q}} = 30-1000$ GeV and $m_{\tilde{g}} < 1.0$ not excluded for $m_{\tilde{q}} = 100-500$ GeV by BALL 84 experiment.
- 134 GOLDMAN 85 use nonobservation of a pseudoscalar $\tilde{g}-\tilde{g}$ bound state in radiative ψ decay.
- 135 HABER 85 is based on survey of all previous searches sensitive to low mass \tilde{g} 's. Limit makes assumptions regarding the lifetime and electric charge of the lightest supersymmetric particle.
- 136 BALL 84 is FNAL beam dump experiment. Observed no interactions of $\tilde{\gamma}$ in the calorimeter, where $\tilde{\gamma}$'s are expected to come from pair-produced \tilde{g} 's. Search for long-lived $\tilde{\gamma}$ interacting in calorimeter 56m from target. Limit is for $m_{\tilde{q}} = 40$ GeV and production cross section proportional to $A^{0.72}$. BALL 84 find no \tilde{g} allowed below 4.1 GeV at CL = 90%. Their figure 1 shows dependence on $m_{\tilde{q}}$ and A. See also KANE 82.
- 137 BRICK 84 reanalyzed FNAL 147 GeV HBC data for $R-\Delta(1232)^{++}$ with $\tau > 10^{-9}$ s and $p_{lab} > 2$ GeV. Set CL = 90% upper limits 6.1, 4.4, and 29 microbarns in $p, p, \pi^+ p, K^+ p$ collisions respectively. $R-\Delta^{++}$ is defined as being \tilde{g} and 3 up quarks. If mass = 1.2-1.5 GeV, then limits may be lower than theory predictions.
- 138 FARRAR 84 argues that $m_{\tilde{g}} < 100$ MeV is not ruled out if the lightest R-hadrons are long-lived. A long lifetime would occur if R-hadrons are lighter than $\tilde{\gamma}$'s or if $m_{\tilde{q}} > 100$ GeV.
- 139 BERGSMAN 83c is reanalysis of CERN-SPS beam-dump data. See their figure 1.
- 140 CHANOWITZ 83 find in bag-model that charged s-hadron exists which is stable against strong decay if $m_{\tilde{g}} < 1$ GeV. This is important since tracks from decay of neutral s-hadron cannot be reconstructed to primary vertex because of missed $\tilde{\gamma}$. Charged s-hadron leaves track from vertex.
- 141 KANE 82 inferred above \tilde{g} mass limit from retroactive analysis of hadronic collision and beam dump experiments. Limits valid if \tilde{g} decays inside detector.

Unstable $\tilde{\gamma}$ (Photino) MASS LIMIT

Unless stated otherwise, the limits below assume that the $\tilde{\gamma}$ decays either into $\gamma \tilde{G}$ (goldstino) or into $\gamma \tilde{H}^0$ (Higgsino).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●				
		142 ACTON	93G OPAL	$e^+e^- \rightarrow \tilde{\gamma}\tilde{\gamma} (\tilde{\gamma} \rightarrow \tau^{\pm} \ell^{\mp} \nu_{\mu})$
		143 ABE	89J VNS	$e^+e^- \rightarrow \tilde{\gamma}\tilde{\gamma}$ ($\tilde{\gamma} \rightarrow \gamma \tilde{G}$ or $\gamma \tilde{H}^0$)
>15	95	144 BEHREND	87B CELL	$e^+e^- \rightarrow \tilde{\gamma}\tilde{\gamma}$ ($\tilde{\gamma} \rightarrow \gamma \tilde{G}$ or $\gamma \tilde{H}^0$)
		145 ADEVA	85 MRKJ	
		146 BALL	84 CALO	Beam dump
		147 BARTEL	84B JADE	
		147 BEHREND	83 CELL	
		148 CABIBBO	81 COSM	

- 142 ACTON 93G assume R-parity violation and decays $\tilde{\gamma} \rightarrow \tau^{\pm} \ell^{\mp} \nu_{\mu}$ ($\ell = e$ or μ). They exclude $m_{\tilde{\gamma}} = 4-43$ GeV for $m_{\tilde{e}_L} < 42$ GeV, and $m_{\tilde{\gamma}} = 7-30$ GeV for $m_{\tilde{e}_L} < 100$ GeV (95% CL). Assumes \tilde{e}_R much heavier than \tilde{e}_L , and lepton family number violation but $L_e = L_{\mu}$ conservation.
- 143 ABE 89J exclude $m_{\tilde{\gamma}} = 0.15-25$ GeV (95%CL) for $d = (100 \text{ GeV})^2$ and $m_{\tilde{g}} = 40$ GeV in the case $\tilde{\gamma} \rightarrow \gamma \tilde{G}$, and $m_{\tilde{\gamma}}$ up to 23 GeV for $m_{\tilde{g}} = 40$ GeV in the case $\tilde{\gamma} \rightarrow \gamma \tilde{H}^0$.
- 144 BEHREND 87B limit is for unstable photinos only. Assumes $B(\tilde{\gamma} \rightarrow \gamma(\tilde{G} \text{ or } \tilde{H}^0)) = 1$, $m_{\tilde{g}}^{\text{for } \tilde{H}^0} \ll m_{\tilde{\gamma}}$ and pure $\tilde{\gamma}$ eigenstate. $m_{\tilde{e}_L} = m_{\tilde{e}_R} < 100$ GeV.
- 145 ADEVA 85 is sensitive to $\tilde{\gamma}$ decay path < 5 cm. With $m_{\tilde{g}} = 50$ GeV, limit (CL = 90%) is $m_{\tilde{\gamma}} > 20.5$ GeV. Assume $\tilde{\gamma}$ decays to photon + goldstino and search for coplanar photons with large missing p_T .
- 146 BALL 84 is FNAL beam dump experiment. Observed no $\tilde{\gamma}$ decay, where $\tilde{\gamma}$'s are expected to come from \tilde{g} 's produced at the target. Three possible $\tilde{\gamma}$ lifetimes are considered. Gluino decay to goldstino + gluon is also considered.
- 147 BEHREND 83 and BARTEL 84B look for 2 γ events from $\tilde{\gamma}$ pair production. With supersymmetric breaking parameter $d = (100 \text{ GeV})^2$ and $m_{\tilde{g}} = 40$ GeV the excluded regions at CL = 95% would be $m_{\tilde{\gamma}} = 100 \text{ MeV} - 13 \text{ GeV}$ for BEHREND 83 $m_{\tilde{\gamma}} = 80 \text{ MeV} - 18 \text{ GeV}$ for BARTEL 84B. Limit is also applicable if the $\tilde{\gamma}$ decays radiatively within the detector.
- 148 CABIBBO 81 consider $\tilde{\gamma} \rightarrow \gamma +$ goldstino. Photino must be either light enough (< 30 eV) to satisfy cosmology bound, or heavy enough (> 0.3 MeV) to have disappeared at early universe.

Supersymmetry Miscellaneous Results

Results that do not appear under other headings or that make nonminimal assumptions.

VALUE	DOCUMENT ID	TECN	COMMENT
● ● ● We do not use the following data for averages, fits, limits, etc. ● ● ●			
	149 BARBER	84B RVUE	
	150 HOFFMAN	83 CNTR	$\pi p \rightarrow n(e^+e^-)$

- 149 BARBER 84B consider that $\tilde{\mu}$ and \tilde{e} may mix leading to $\mu \rightarrow e\tilde{\gamma}$. They discuss mass-mixing limits from decay dist asym in LBL-TRIUMF data and e^+ polarization in SIN data.
- 150 HOFFMAN 83 set CL = 90% limit $d\sigma/dt B(e^+e^-) < 3.5 \times 10^{-32} \text{ cm}^2/\text{GeV}^2$ for spin-1 partner of Goldstone fermions with $140 < m < 160$ MeV decaying $\rightarrow e^+e^-$ pair.

REFERENCES FOR Supersymmetric Particle Searches

ACTON	93G	PL B313 333	+Akers, Alexander, Allison, Anderson+ (OPAL Collab.)
ADRIANI	93M	PRP1 236 1	+Aguliar-Benitez, Alier, Alcaraz, Aloisio+ (L3 Collab.)
CLAVELLI	93	PR D47 1973	+Coulter, Yuan (ALAT)
DREES	93	PR D47 376	+Nojiri (DESY, SLAC)
FALK	93	PL B318 354	+Madden, Olive, Srednicki (UCB, UCSB, MINN) (CERN)
HEBBEKER	93	ZPHY C60 63	+Lopez, Nanopoulos, Pois, Yuan (TAMU, ALAH) (HOUS)
KELLEY	93	PR D47 2461	
LAU	93	PR D47 1087	
LOPEZ	93C	PL B313 241	+Nanopoulos, Wang (TAMU, HARC, CERN)
MIZUTA	93	PL B298 120	+Yamaguchi (TOHO)
MORI	93	PR D48 5505	+KEK, NIIG, TOKY, TOKA, KOBE, OSAK, TINT, GFU)
ABE	92L	PRL 69 3439	+Amidei, Anway-Wiese, Apollinari, Atac+ (CDF Collab.)
BOTTINO	92	MPL A7 733	+DeAlfaro, Fornengo, Morales, Puimedo+ (TORI, ZARA)
	Also	91	Bottino, de Alfaro, Fornengo, Mignola+ (TORI, INFN)
CLAVELLI	92	PR D46 2112	(ALAT)
DATTA	92	ZPHY C54 513	+Guchait, Raychaudhuri (JADA, CALC)
DECAMP	92	PRP1 216 253	+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
ELLIS	92F	PL B283 252	+Roszkowski (CERN)
KAWASAKI	92	PR D46 1634	+Mizuta (OSU, TOHO)
LOPEZ	92	NP B370 445	+Nanopoulos, Yuan (TAMU)
MCDONALD	92	PL B283 80	+Olive, Srednicki (LISB, MINN, UCSB) (CERN)
ROY	92	PL B283 270	
ABREU	91F	NP B367 511	+Adam, Adami, Abye, Akesson+ (DELPHI Collab.)
AKESSON	91	ZPHY C52 219	+Almeida, Angelis, Atherton, Aubry+ (HELIOS Collab.)
ALEXANDER	91F	PHYS C52 175	+Allison, Fulgini, Anderson, Arczelli+ (OPAL Collab.)
ANTONIADIS	91	PL B262 109	+Ellis, Nanopoulos (EPOL, CERN, TAMU, HARC)
BAER	91	PR D44 207	+Tata, Woodside (FSU, HAWA, ISU)
BAER	91B	PR D44 725	+Drees, Godbole+ (FSU, DESY, BOMB, UCD, HAWA)
BOTTINO	91	PL B265 57	+de Alfaro, Fornengo, Mignola+ (TORI, INFN)
DREES	91	PR D43 2971	+Tata (CERN, HAWA)
GELMINI	91	NP B351 293	+Gondolo, Roulet (UCLA, FRST) (TGAK)
HIDAKA	91	PR D44 927	
KAMIONKOW	91	PR D44 3021	Kamionkowski (CHIC, FNAL)
MORI	91B	PL B270 89	+Nojiri, Oyama, Suzuki+ (Kamiookande Collab.)
NOJIRI	91	PL B261 76	(KEK)
OLIVE	91	NP B355 208	+Srednicki (MINN, UCSB) (CERN)
ROSKOWSKI	91	PL B262 59	
SATO	91	PR D44 2220	+Hirata, Kajita, Kifune, Kihara+ (Kamioka Collab.)
ABREU	90F	PL B247 149	+Adam, Adami, Abye, Alekseev+ (HELIOS Collab.)
ADACHI	90C	OPAL B247 157	+Adam, Adami, Abye, Alekseev+ (DELPHI Collab.)
ADACHI	90C	OPAL B244 352	+Aihara, Doser, Enomoto+ (TOPAZ Collab.)
ADEVA	90L	PL B249 341	+Adriani, Aguliar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
AKESSON	90L	PL B238 442	+Alliti, Ansari, Ansgore+ (UA2 Collab.)
AKRAWY	90D	PL B240 261	+Alexander, Allison, Allport+ (OPAL Collab.)
AKRAWY	90D	PL B248 211	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
AKRAWY	90D	PL B252 290	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
ALITTI	90	PR D41 3565	+Ansari, Ansgore, Bagnaia, Bareyre+ (UA2 Collab.)
BAER	90	PR D41 3414	+Drees, Tata (FSU, CERN, HAWA)
BARKLOW	90	PRL 64 2984	+Abrams, Adolphsen, Averill, Ballam+ (Mark II Collab.)
DECAMP	90C	PL B236 86	+Deschizeaux, Lees, Minard, Crespo+ (ALEPH Collab.)
DECAMP	90K	PL B244 541	+Deschizeaux, Goy, Lees+ (ALEPH Collab.)
DREES	90	PL B252 127	+Hikasa (CERN, KEK)
ELLIS	90	PL B245 251	+Nanopoulos, Roszkowski, Schramm (CERN, HARC, TAMU)
GRIEST	90	PR D41 3565	+Kamionkowski, Turner (UCB, CHIC) (BARC)
GRIFFOLS	90	NP B331 244	+Masso (YALE)
KRAUSS	90	PRL 64 999	
ROSKOWSKI	90	PL B252 471	(TAMU, HARC)
SAKAI	90	PL B234 534	+Gu, Low, Abe, Fujii+ (AMY Collab.)
SODERSTROM	90	PRL 64 2980	+McKenna, Abrams, Adolphsen, Averill+ (Mark II Collab.)
TAKETANI	90	PL B234 202	+Oadka, Abe, Amako+ (VENUS Collab.)
ZHUKOVSKII	90	SJNP 52 931	+Eminov (MOSU)
Translated from YAF 52 1473			
ABE	89J	ZPHY C45 175	+Amako, Arai, Fukawa+ (VENUS Collab.)
ABE	89K	PL B232 431	+Amako, Arai, Asano, Chiba+ (VENUS Collab.)
ADACHI	89	PL B218 105	+Aihara, Dijkstra, Enomoto, Fujii+ (TOPAZ Collab.)
ADEVA	89B	PL B233 530	+Adriani, Aguliar-Benitez, Akbari+ (L3 Collab.)
ALBAJAR	89	ZPHY C44 15	+Albrow, Altkofer, Arison, Astbury+ (UA1 Collab.)
HEARTY	89	PR D39 3207	+Rothberg, Young, Johnson, Whitaker+ (ASP Collab.)
	Also	89	Hearty, Rothberg, Young, Johnson+ (ASP Collab.)
	Also	86	Bartha, Burke, Extermann+ (ASP Collab.)
NAKAMURA	89	PR D39 1261	+Kobayashi, Konaka, Imai, Masaie+ (KYOT, TMTC)
OLIVE	89	PR B230 78	+Srednicki (MINN, UCSB)
BAER	88	PR D38 1485	+Hagiwara, Tata (FSU, KEK, WISC) (CERN)
	Also	88	Baer, Hagiwara, Tata (FSU, KEK, WISC)
BEHREND	88B	PL B215 186	+Criegee, Dainton, Field+ (CELLO Collab.)
ELLIS	88B	PL B215 404	+Olive, Sarkar, Sciamia (CERN, MINN, RAL, CERN) (CERN)
NIETH	88	PR D38 1479	+Arnowitz (NEAS, TAMU)
OLIVE	88	PL B205 553	+Srednicki (MINN, UCSB)
SREDNICKI	88	NP B310 693	+Watkins, Olive (MINN, UCSB)
ADEVA	87	PL B194 167	+Anderhub, Ansari, Becker+ (Mark-I Collab.)
ALBAJAR	87D	PL B185 241	+Albrow, Altkofer, Arison+ (UA1 Collab.)
ALBAJAR	87D	PL B198 261	+Albrow, Altkofer+ (UA1 Collab.)
ANSARI	87D	PL B195 613	+Bagnaia, Banner+ (UA2 Collab.)
ARNOLD	87	PL B186 435	+Barth+ (BRUX, DUUC, LOUC, BARI, AICH, CERN+)
BAER	87	PR D35 1598	+Hagiwara, Tata (KEK, ANL, WISC)
	Also	86	Baer, Hagiwara, Tata (ANL, DESY, WISC)
BEHREND	87B	ZPHY C35 181	+Buerger, Criegee, Dainton+ (CELLO Collab.)
HEARTY	87	PRL 58 1711	+Rothberg, Young, Johnson+ (ASP Collab.)
NG	87	PL B188 138	+Olive, Srednicki (MINN, UCSB)
TUTS	87	PL B186 233	+Frantzi, Youssef, Zhao+ (CUSB Collab.)
ALDRICHT	86C	PL B178 360	+Binder, Harder+ (ARGUS Collab.)
BADIER	86	ZPHY C31 21	+Bemmerad, Boucrot, Caliot+ (NA3 Collab.)
BARNETT	86	NP B267 625	+Haber, Kane (LBL, UCSC, MICH)
FORD	86	PR D33 3472	+Qi, Read+ (MAC Collab.)
GAISSER	86	PR D34 2206	+Steigman, Tilav (BART, DELA)
VOLOSHIN	86	SJNP 43 495	+Okun (ITEP)
Translated from YAF 43 779:			
ADEVA	85	PL 152B 439	+Becker, Becker-Szendy+ (Mark-J Collab.)
	Also	84C	PRP1 109 131 (Mark-J Collab.)
AKERLOF	85	PL 156B 271	+Bonvicini, Chapman, Errede+ (HRS Collab.)
BARTEL	85	PL 155B 288	+Becker, Cords, Fests, Hagiwara+ (JADE Collab.)
BEHREND	85	PL 161B 182	+Burger, Criegee, Fenner+ (CELLO Collab.)
COOPER...	85B	PL 160B 212	+Cooper-Sarkar, Parker, Sarkar+ (Wab6 Collab.)
DAWSON	85	PR D31 1581	+Eichten, Quigg (LBL, FNAL)
FARRAR	85	PL 155 895	(RUTG)
GOLDMAN	85	Physica 150 181	+Haber (LANL, USC)
HABER	85	PRP1 117 75	+Kane (UCSC, MICH)
ADEVA	84B	PRL 53 1806	+Barber, Becker, Berdugo+ (Mark-J Collab.)
BALL	84	PRL 53 1314	+Coffin, Gustafson+ (MICH, FIRZ, OSU, FNAL, WISC)
BARBER	84B	PL 139B 427	+Shrock (STON)
BARTEL	84B	PL 139B 327	+Becker, Bowden, Cords+ (JADE Collab.)
FARRAR	84C	PL 145C 126	+Becker, Bowden, Cords+ (JADE Collab.)
BRICK	84	PR D30 1134	+ (BROW, CAVE, IIT, IND, MIT, MONS, NIJH+)
ELLIS	84	NP B238 453	+Hageijn, Nanopoulos, Olive, Srednicki (CERN)
FARRAR	84	PRL 53 1029	(RUTG)
BEHREND	83	PL 123B 127	+Chen, Fenner, Gumpel+ (CELLO Collab.)

BERGSMA	83C	PL 121B 429	+Dorenbosch, Jonker+	(CHARM Collab.)
CHANOWITZ	83	PL 126B 225	+Sharpe	(UCB, LBL)
GOLDBERG	83	PRL 50 1419		(NEAS)
HOFFMAN	83	PR D28 660	+Frank, Mischke, Moir, Schardt	(LANL, ARZS)
KRAUSS	83	NP B227 556		(HARV)
VYSOTSKII	83	SJNP 37 948		(ITEP)
		Translated from YAF 37 1597		
KANE	82	PL 112B 227	+Leveille	(MICH)
CABIBBO	81	PL 105B 155	+Farrar, Maiani	(ROMA, RUTG)
FARRAR	78	PL 76B 575	+Fayet	(CIT)
Also	78B	PL 79B 442	Farrar, Fayet	(CIT)

Searches for Quark and Lepton Compositeness

NOTE ON SEARCHES FOR QUARK AND LEPTON COMPOSITENESS

If quarks and leptons are made of constituents, then at the scale of constituent binding energies, there should appear new interactions among quarks and leptons. At energies much below the compositeness scale (Λ), these interactions are suppressed by inverse powers of Λ . The dominant effect should come from the lowest dimensional interactions with four fermions (contact terms), whose most general chirally invariant form reads [1]

$$L = \frac{g^2}{2\Lambda^2} \left[\eta_{LL} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_L \gamma^\mu \psi_L + \eta_{RR} \bar{\psi}_R \gamma_\mu \psi_R \bar{\psi}_R \gamma^\mu \psi_R + 2\eta_{LR} \bar{\psi}_L \gamma_\mu \psi_L \bar{\psi}_R \gamma^\mu \psi_R \right]. \quad (1)$$

Chiral invariance provides a natural explanation why quark and lepton masses are much smaller than their inverse size Λ . We may determine the scale Λ unambiguously by using the above form of the effective interactions; the conventional method [1] is to fix its scale by setting $g^2/4\pi = g^2(\Lambda)/4\pi = 1$ for the new strong interaction coupling and by setting the largest magnitude of the coefficients $\eta_{\alpha\beta}$ to be unity. In the following, we denote

$$\begin{aligned} \Lambda &= \Lambda_{LL}^\pm \text{ for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, 0, 0), \\ \Lambda &= \Lambda_{RR}^\pm \text{ for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (0, \pm 1, 0), \\ \Lambda &= \Lambda_{VV}^\pm \text{ for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, \pm 1, \pm 1), \\ \Lambda &= \Lambda_{AA}^\pm \text{ for } (\eta_{LL}, \eta_{RR}, \eta_{LR}) = (\pm 1, \pm 1, \mp 1), \end{aligned} \quad (2)$$

as typical examples. Such interactions can arise by constituent interchange (when the fermions have common constituents, e.g., for $ee \rightarrow ee$) and/or by exchange of the binding quanta (when ever binding quanta couple to constituents of both particles).

Another typical consequence of compositeness is the appearance of excited leptons and quarks (ℓ^* and q^*). Phenomenologically, an excited lepton is defined to be a heavy lepton which shares leptonic quantum number with one of the existing leptons (an excited quark is defined similarly). For example, an excited electron e^* is characterized by a nonzero transition-magnetic coupling with electrons. Smallness of the lepton mass and the success of QED prediction for $g-2$ suggest chirality conservation, *i.e.*, an excited lepton should not couple to both left- and right-handed components of the corresponding lepton.

Excited leptons may be classified by $SU(2) \times U(1)$ quantum numbers. Typical examples are:

1. Sequential type

$$\begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L, \quad [\nu_R^*], \quad \ell_R^*.$$

ν_R^* is necessary unless ν^* has a Majorana mass.

2. Mirror type

$$[\nu_L^*], \quad \ell_L^*, \quad \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R.$$

3. Homodoublet type

$$\begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_L, \quad \begin{pmatrix} \nu^* \\ \ell^* \end{pmatrix}_R.$$

Similar classification can be made for excited quarks.

Excited fermions can be pair produced via their gauge couplings. The couplings of excited leptons with Z are listed in the following table (for notation see Eq. (1) in "Standard Model of Electroweak Interactions" in Sec. III):

	Sequential type	Mirror type	Homodoublet type
$V\ell^*$	$-\frac{1}{2} + 2\sin^2\theta_W$	$-\frac{1}{2} + 2\sin^2\theta_W$	$-1 + 2\sin^2\theta_W$
$A\ell^*$	$-\frac{1}{2}$	$+\frac{1}{2}$	0
$V\nu_D^*$	$+\frac{1}{2}$	$+\frac{1}{2}$	+1
$A\nu_D^*$	$+\frac{1}{2}$	$-\frac{1}{2}$	0
$V\nu_M^*$	0	0	—
$A\nu_M^*$	+1	-1	—

Here ν_D^* (ν_M^*) stands for Dirac (Majorana) excited neutrino. The corresponding couplings of excited quarks can be easily obtained. Although form factor effects can be present for the gauge couplings at $q^2 \neq 0$, they are usually neglected.

In addition, transition magnetic type couplings with a gauge boson are expected. These couplings can be generally parametrized as follows:

$$\begin{aligned} \mathcal{L} &= \frac{\lambda_{\gamma}^{(f^*)}}{2m_{f^*}} \bar{f}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) f F_{\mu\nu} \\ &+ \frac{\lambda_Z^{(f^*)}}{2m_{f^*}} \bar{f}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) f Z_{\mu\nu} \\ &+ \frac{\lambda_W^{(\ell^*)}}{2m_{\ell^*}} \bar{\ell}^* \sigma^{\mu\nu} \frac{1-\gamma_5}{2} \nu W_{\mu\nu} \\ &+ \frac{\lambda_W^{(\nu^*)}}{2m_{\nu^*}} \bar{\nu}^* \sigma^{\mu\nu} (\eta_L \frac{1-\gamma_5}{2} + \eta_R \frac{1+\gamma_5}{2}) \ell W_{\mu\nu}^\dagger \\ &+ \text{h.c.}, \end{aligned} \quad (3)$$

where $g = e/\sin\theta_W$, $F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu$ is the photon field strength, $Z_{\mu\nu} = \partial_\mu Z_\nu - \partial_\nu Z_\mu$, *etc.* The normalization of the coupling is chosen such that

$$\max(|\eta_L|, |\eta_R|) = 1.$$

Searches Full Listings

Quark and Lepton Compositeness

Chirality conservation requires

$$\eta_L \eta_R = 0. \quad (4)$$

These couplings can arise from $SU(2) \times U(1)$ -invariant higher-dimensional interactions. A well-studied model is the interaction of homodoublet type ℓ^* with the Lagrangian [2,3]

$$\mathcal{L} = \frac{1}{2\Lambda} \bar{L}^* (g f \frac{\tau^a}{2} W_{\mu\nu}^a + g' f' Y B_{\mu\nu}) \frac{1-\gamma_5}{2} L + \text{h.c.}, \quad (5)$$

where L denotes the lepton doublet (ν, ℓ), Λ is the compositeness scale, g, g' are $SU(2)$ and $U(1)_Y$ gauge couplings, and $W_{\mu\nu}^a$ and $B_{\mu\nu}$ are the field strengths for $SU(2)$ and $U(1)_Y$ gauge fields. The same interaction occurs for mirror-type excited leptons. For sequential-type excited leptons, the ℓ^* and ν^* couplings become unrelated, and the couplings receive the extra suppression of $(250 \text{ GeV})/\Lambda$ or m_{L^*}/Λ . In any case, these couplings satisfy the relation

$$\lambda_W = -\sqrt{2} \sin^2 \theta_W (\lambda_Z \cot \theta_W + \lambda_\gamma). \quad (6)$$

Additional coupling with gluons is possible for excited quarks:

$$\mathcal{L} = \frac{1}{2\Lambda} \bar{Q}^* \sigma^{\mu\nu} \left(g_s f_s \frac{\lambda^a}{2} G_{\mu\nu}^a + g f \frac{\tau^a}{2} W_{\mu\nu}^a + g' f' Y B_{\mu\nu} \right) \times \frac{1-\gamma_5}{2} Q + \text{h.c.}, \quad (7)$$

where Q denotes a quark doublet, g_s is the QCD gauge coupling, and $G_{\mu\nu}^a$ the gluon field strength.

Some experimental analyses assume the relation $\eta_L = \eta_R = 1$, which violates chiral symmetry. We encode the results of such analyses if the crucial part of the cross section is proportional to the factor $\eta_L^2 + \eta_R^2$ and the limits can be reinterpreted as those for chirality conserving cases (η_L, η_R) = (1, 0) or (0, 1) after rescaling λ .

Several different conventions are used by LEP experiments to express the transition magnetic couplings. To facilitate comparison, we reexpress these in terms of λ_Z and λ_γ using the following relations and taking $\sin^2 \theta_W = 0.23$. We assume chiral couplings, *i.e.*, $|c| = |d|$ in the notation of Ref. 2.

1. ALEPH (charged lepton and neutrino)

$$\lambda_Z^{\text{ALEPH}} = \frac{1}{2} \lambda_Z \quad (1990 \text{ papers}) \quad (8a)$$

$$\frac{2c}{\Lambda} = \frac{\lambda_Z}{m_{\ell^*} [\text{or } m_{\nu^*}]} \quad (\text{for } |c| = |d|) \quad (8b)$$

2. ALEPH (quark)

$$\lambda_u^{\text{ALEPH}} = \frac{\sin \theta_W \cos \theta_W}{\sqrt{\frac{1}{4} - \frac{2}{3} \sin^2 \theta_W + \frac{8}{9} \sin^4 \theta_W}} \lambda_Z = 1.11 \lambda_Z \quad (9)$$

3. L3 and DELPHI (charged lepton)

$$\lambda^{\text{L3}} = \lambda_Z^{\text{DELPHI}} = -\frac{\sqrt{2}}{\cot \theta_W - \tan \theta_W} \lambda_Z = -1.10 \lambda_Z \quad (10)$$

4. L3 (neutrino)

$$f_Z^{\text{L3}} = \sqrt{2} \lambda_Z \quad (11)$$

5. OPAL (charged lepton)

$$\frac{f^{\text{OPAL}}}{\Lambda} = -\frac{2}{\cot \theta_W - \tan \theta_W} \frac{\lambda_Z}{m_{\ell^*}} = -1.56 \frac{\lambda_Z}{m_{\ell^*}} \quad (12)$$

6. OPAL (quark)

$$\frac{f^{\text{OPAL}} c}{\Lambda} = \frac{\lambda_Z}{2m_{q^*}} \quad (\text{for } |c| = |d|) \quad (13)$$

7. DELPHI (charged lepton)

$$\lambda_\gamma^{\text{DELPHI}} = -\frac{1}{\sqrt{2}} \lambda_\gamma \quad (14)$$

If leptons are made of color triplet and antitriplet constituents, we may expect their color-octet partners. Transitions between the octet leptons (ℓ_8) and the ordinary lepton (ℓ) may take place via the dimension-five interactions

$$\mathcal{L} = \frac{1}{2\Lambda} \sum_\ell \left\{ \bar{\ell}_8^\alpha g_S F_{\mu\nu}^\alpha \sigma^{\mu\nu} (\eta_L \ell_L + \eta_R \ell_R) + \text{h.c.} \right\} \quad (15)$$

where the summation is over charged leptons and neutrinos. The leptonic chiral invariance implies $\eta_L \eta_R = 0$ as before.

References

1. E.J. Eichten, K.D. Lane, and M.E. Peskin, Phys. Rev. Lett. **50**, 811 (1983).
2. K. Hagiwara, S. Komamiya, and D. Zeppenfeld, Z. Phys. **C29**, 115 (1985).
3. N. Cabibbo, L. Maiani, and Y. Srivastava, Phys. Lett. **139B**, 459 (1984).

SCALE LIMITS for Contact Interactions: $\Lambda(eeee)$

Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

$\Lambda_{LL}^+(\text{TeV})$	$\Lambda_{LL}^-(\text{TeV})$	CL%	DOCUMENT ID	TECN	COMMENT
>1.6	95	1,2	BUSKULIC	93Q RVUE	
>3.6	95	3	KROHA	92 RVUE	
••• We do not use the following data for averages, fits, limits, etc. •••					
>1.6	>2.0	95	2 BUSKULIC	93Q ALEP	$E_{\text{cm}}=88.25-94.25 \text{ GeV}$
	>2.2	95	BUSKULIC	93Q RVUE	
>1.3	95	3	KROHA	92 RVUE	
>0.7	>2.8	95	BEHREND	91C CELL	$E_{\text{cm}}=35 \text{ GeV}$
>1.3	>1.3	95	KIM	89 AMY	$E_{\text{cm}}=50-57 \text{ GeV}$
>1.4	>3.3	95	4 BRAUNSCH...	88 TASS	$E_{\text{cm}}=12-46.8 \text{ GeV}$
>1.0	>0.7	95	5 FERNANDEZ	87B MAC	$E_{\text{cm}}=29 \text{ GeV}$
>1.1	>1.4	95	6 BARTEL	86C JADE	$E_{\text{cm}}=12-46.8 \text{ GeV}$
>1.17	>0.87	95	7 DERRICK	86 HRS	$E_{\text{cm}}=29 \text{ GeV}$
>1.1	>0.76	95	8 BERGER	85B PLUT	$E_{\text{cm}}=34.7 \text{ GeV}$

¹ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

² BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95%CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

³ KROHA 92 limit is from fit to BERGER 85b, BARTEL 86c, DERRICK 86b, FERNANDEZ 87b, BRAUNSCHWEIG 88, BEHREND 91b, and BEHREND 91c. The fit gives $\eta/\Lambda_{LL}^2 = +0.230 \pm 0.206 \text{ TeV}^{-2}$.

⁴ BRAUNSCHWEIG 88 assumed $m_Z = 92 \text{ GeV}$ and $\sin^2 \theta_W = 0.23$.

⁵ FERNANDEZ 87b assumed $\sin^2 \theta_W = 0.22$.

⁶ BARTEL 86c assumed $m_Z = 93 \text{ GeV}$ and $\sin^2 \theta_W = 0.217$.

⁷ DERRICK 86 assumed $m_Z = 93 \text{ GeV}$ and $g_V^2 = (-1/2 + 2\sin^2 \theta_W)^2 = 0.004$.

⁸ BERGER 85b assumed $m_Z = 93 \text{ GeV}$ and $\sin^2 \theta_W = 0.217$.

SCALE LIMITS for Contact Interactions: $\Lambda(ee\mu\mu)$ Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

Λ_{LL}^{\pm} (TeV)	Λ_{LL}^{\pm} (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>2.6	>1.9	95	9,10 BUSKULIC	93Q RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>1.3	>1.5	95	10 BUSKULIC	93Q ALEP	$E_{cm}=88.25-94.25$ GeV
>2.3	>2.0	95	HOWELL	92 TOPZ	$E_{cm}=52-61.4$ GeV
>1.7	>1.7	95	11 KROHA	92 RVUE	
>2.5	>1.5	95	BEHREND	91C CELL	$E_{cm}=35-43$ GeV
>1.6	>2.0	95	12 ABE	90I VNS	$E_{cm}=50-60.8$ GeV
>1.9	>1.0	95	KIM	89 AMY	$E_{cm}=50-57$ GeV
>2.3	>1.3	95	BRAUNSCH...	88D TASS	$E_{cm}=30-46.8$ GeV
>4.4	>2.1	95	13 BARTEL	86C JADE	$E_{cm}=12-46.8$ GeV
>2.9	>0.86	95	14 BERGER	85 PLUT	$E_{cm}=34.7$ GeV

⁹ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

¹⁰ BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95% CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

¹¹ KROHA 92 limit is from fit to BARTEL 86C, BEHREND 87C, BRAUNSCHWEIG 88D, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives $\eta/\Lambda_{LL}^2 = -0.155 \pm 0.095$ TeV⁻².

¹² ABE 90I assumed $m_Z = 91.163$ GeV and $\sin^2\theta_W = 0.231$.

¹³ BARTEL 86C assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

¹⁴ BERGER 85 assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

SCALE LIMITS for Contact Interactions: $\Lambda(ee\tau\tau)$ Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

Λ_{LL}^{\pm} (TeV)	Λ_{LL}^{\pm} (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1.9	>2.9	95	15 KROHA	92 RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>1.0	>1.5	95	16 BUSKULIC	93Q ALEP	$E_{cm}=88.25-94.25$ GeV
>1.8	>2.3	95	16,17 BUSKULIC	93Q RVUE	
>1.9	>1.7	95	HOWELL	92 TOPZ	$E_{cm}=52-61.4$ GeV
>1.6	>2.3	95	BEHREND	91C CELL	$E_{cm}=35-43$ GeV
>1.8	>1.3	95	18 ABE	90I VNS	$E_{cm}=50-60.8$ GeV
>2.2	>3.2	95	19 BARTEL	86 JADE	$E_{cm}=12-46.8$ GeV

¹⁵ KROHA 92 limit is from fit to BARTEL 86C BEHREND 89B, BRAUNSCHWEIG 89C, ABE 90I, and BEHREND 91C. The fit gives $\eta/\Lambda_{LL}^2 = +0.095 \pm 0.120$ TeV⁻².

¹⁶ BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95% CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

¹⁷ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

¹⁸ ABE 90I assumed $m_Z = 91.163$ GeV and $\sin^2\theta_W = 0.231$.

¹⁹ BARTEL 86 assumed $m_Z = 93$ GeV and $\sin^2\theta_W = 0.217$.

SCALE LIMITS for Contact Interactions: $\Lambda(eeee)$ Lepton universality assumed. Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

Λ_{LL}^{\pm} (TeV)	Λ_{LL}^{\pm} (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>3.5	>2.8	95	20,21 BUSKULIC	93Q RVUE	
• • • We do not use the following data for averages, fits, limits, etc. • • •					
>3.0	>2.3	95	21,22 BUSKULIC	93Q ALEP	$E_{cm}=88.25-94.25$ GeV
>2.5	>2.2	95	23 HOWELL	92 TOPZ	$E_{cm}=52-61.4$ GeV
>3.4	>2.7	95	24 KROHA	92 RVUE	

²⁰ This BUSKULIC 93Q value is from ALEPH data plus PEP/PETRA/TRISTAN data re-analyzed by KROHA 92.

²¹ BUSKULIC 93Q uses the following prescription to obtain the limit: when the naive 95% CL limit is better than the statistically expected sensitivity for the limit, the latter is adopted for the limit.

²² From $e^+e^- \rightarrow e^+e^-, \mu^+\mu^-,$ and $\tau^+\tau^-$.

²³ HOWELL 92 limit is from $e^+e^- \rightarrow \mu^+\mu^-$ and $\tau^+\tau^-$.

²⁴ KROHA 92 limit is from fit to most PEP/PETRA/TRISTAN data. The fit gives $\eta/\Lambda_{LL}^2 = -0.0200 \pm 0.0666$ TeV⁻².

SCALE LIMITS for Contact Interactions: $\Lambda(eeqq)$ Limits are for Λ_{LL}^{\pm} only. For other cases, see each reference.

Λ_{LL}^{\pm} (TeV)	Λ_{LL}^{\pm} (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1.7	>2.2	95	25 ABE	91D CDF	($eeqq$) (Isosinglet)
>1.2		95	26 ADACHI	91 TOPZ	($eeqq$) (flavor-universal)
>1.7		95	27 ABE	89L VNS	($eeqq$) (flavor-universal)

• • • We do not use the following data for averages, fits, limits, etc. • • •

>1.6	95	26 ADACHI	91 TOPZ	($eeqq$) (flavor-universal)	
>0.6	>1.7	95	28 BEHREND	91C CELL	($eecc$)
>1.1	>1.0	95	28 BEHREND	91C CELL	($eebb$)
>0.9		95	27 ABE	89L VNS	($eeqq$) (flavor-universal)
>1.05	>1.61	95	29 HAGIWARA	89 RVUE	($eecc$)
>1.21	>0.53	95	30 HAGIWARA	89 RVUE	($eebb$)

²⁵ ABE 91D limits are from e^+e^- mass distribution in $p\bar{p} \rightarrow e^+e^-X$ at $E_{cm} = 1.8$ TeV.

²⁶ ADACHI 91 limits are from differential jet cross section. Universality of $\Lambda(eeqq)$ for five flavors is assumed.

²⁷ ABE 89L limits are from jet charge asymmetry. Universality of $\Lambda(eeqq)$ for five flavors is assumed.

²⁸ BEHREND 91C is from data at $E_{cm} = 35-43$ GeV.

²⁹ The HAGIWARA 89 limit is derived from forward-backward asymmetry measurements of D/D^* mesons by ALTHOFF 83C, BARTEL 84E, and BARINGER 88.

³⁰ The HAGIWARA 89 limit is derived from forward-backward asymmetry measurement of b hadrons by BARTEL 84D.

SCALE LIMITS for Contact Interactions: $\Lambda(\mu\mu qq)$

Λ_{LL}^{\pm} (TeV)	Λ_{LL}^{\pm} (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1.4	>1.6	95	ABE	92B CDF	($\mu\mu qq$) (Isosinglet)

SCALE LIMITS for Contact Interactions: $\Lambda(\ell\nu\ell\nu)$

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>3.10	90	31 JODIDIO	86 SPEC	$\Lambda_{LR}^{\pm}(\nu_{\mu}\nu_e\mu_e)$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>3.8		32 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^{\pm}(\tau\nu_{\tau}\nu_e)$
>8.1		32 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^{\pm}(\tau\nu_{\tau}\nu_{\mu})$
>4.1		33 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^{\pm}(\tau\nu_{\tau}\nu_{\mu})$
>6.5		33 DIAZCRUZ	94 RVUE	$\Lambda_{LL}^{\pm}(\tau\nu_{\tau}\nu_{\mu})$

³¹ JODIDIO 86 limit is from $\mu^+ \rightarrow \bar{\nu}_{\mu} e^+ \nu_e$. Chirality invariant interactions $I = (g^2/\Lambda^2) [\eta_{LL} (\bar{\nu}_{\mu} \gamma^{\alpha} \mu_L) (\bar{e}_L \gamma_{\alpha} \nu_e) + \eta_{LR} (\bar{\nu}_{\mu} \gamma^{\alpha} \nu_e) (\bar{e}_R \gamma_{\alpha} \mu_R)]$ with $g^2/4\pi = 1$ and $(\eta_{LL}, \eta_{LR}) = (0, \pm 1)$ are taken. No limits are given for Λ_{LL}^{\pm} with $(\eta_{LL}, \eta_{LR}) = (\pm 1, 0)$. For more general constraints with right-handed neutrinos and chirality nonconserving contact interactions, see their text.

³² DIAZCRUZ 94 limits are from $\Gamma(\tau \rightarrow e\nu\nu)$ and assume flavor-dependent contact interactions with $\Lambda(\tau\nu_{\tau}\nu_e) \ll \Lambda(\mu\nu_{\mu}\nu_e)$.

³³ DIAZCRUZ 94 limits are from $\Gamma(\tau \rightarrow \mu\nu\nu)$ and assume flavor-dependent contact interactions with $\Lambda(\tau\nu_{\tau}\nu_{\mu}) \ll \Lambda(\mu\nu_{\mu}\nu_e)$.

SCALE LIMITS for Contact Interactions: $\Lambda(qqqq)$ Limits are for Λ_{LL}^{\pm} with color-singlet isoscalar exchanges among u_L 's and d_L 's only. See EICHTEN 84 for details.

VALUE (TeV)	CL%	DOCUMENT ID	TECN	COMMENT
>1.4	95	34 ABE	92D CDF	$p\bar{p} \rightarrow$ jets inclusive
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>1.3	95	35 ABE	93G CDF	$p\bar{p} \rightarrow$ dijet mass
>1.0	99	36 ABE	92M CDF	$p\bar{p} \rightarrow$ dijet angl.
>0.825	95	37 ALITTI	91B UA2	$p\bar{p} \rightarrow$ jets inclusive
>0.700	95	34 ABE	89 CDF	$p\bar{p} \rightarrow$ jets inclusive
>0.330	95	38 ABE	89H CDF	$p\bar{p} \rightarrow$ dijet angl.
>0.400	95	39 ARNISON	86C UA1	$p\bar{p} \rightarrow$ jets inclusive
>0.415	95	40 ARNISON	86D UA1	$p\bar{p} \rightarrow$ dijet angl.
>0.370	95	41 APPEL	85 UA2	$p\bar{p} \rightarrow$ jets inclusive
>0.275	95	42 BAGNAIA	84C UA2	Repl. by APPEL 85

³⁴ Limit is from inclusive jet cross-section data in $p\bar{p}$ collisions at $E_{cm} = 1.8$ TeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.

³⁵ ABE 93G limit is from dijet mass distribution in $p\bar{p}$ collisions at $E_{cm} = 1.8$ TeV. The limit is the weakest from several choices of structure functions and renormalization scale.

³⁶ ABE 92M limit is from dijet angular distribution for $m_{dijet} > 550$ GeV in $p\bar{p}$ collisions at $E_{cm} = 1.8$ TeV.

³⁷ ALITTI 91B limit is from inclusive jet cross section in $p\bar{p}$ collisions at $E_{cm} = 630$ GeV. The limit takes into account uncertainties in choice of structure functions and in choice of process scale.

³⁸ ABE 89H limit is from dijet angular distribution for $m_{dijet} > 200$ GeV at the Fermilab Tevatron Collider with $E_{cm} = 1.8$ TeV. The QCD prediction is quite insensitive to choice of structure functions and choice of process scale.

³⁹ ARNISON 86C limit is from the study of inclusive high- p_T jet distributions at the CERN $\bar{p}p$ collider ($E_{cm} = 546$ and 630 GeV). The QCD prediction renormalized to the low- p_T region gives a good fit to the data.

⁴⁰ ARNISON 86D limit is from the study of dijet angular distribution in the range 240 < $m(dijet)$ < 300 GeV at the CERN $\bar{p}p$ collider ($E_{cm} = 630$ GeV). QCD prediction using EHLQ structure function (EICHTEN 84) with $\Lambda_{QCD} = 0.2$ GeV for the choice of $Q^2 = p_T^2$ gives the best fit to the data.

⁴¹ APPEL 85 limit is from the study of inclusive high- p_T jet distributions at the CERN $\bar{p}p$ collider ($E_{cm} = 630$ GeV). The QCD prediction renormalized to the low- p_T region gives a good description of the data.

⁴² BAGNAIA 84C limit is from the study of jet p_T and dijet mass distributions at the CERN $\bar{p}p$ collider ($E_{cm} = 540$ GeV). The limit suffers from the uncertainties in comparing the data with the QCD prediction.

Searches Full Listings

Quark and Lepton Compositeness

MASS LIMITS for Excited e^*

Most e^+e^- experiments assume one-photon or Z exchange. The limits from some e^+e^- experiments which depend on λ have assumed transition couplings which are chirality violating ($\eta_L = \eta_R$). However they can be interpreted as limits for chirality-conserving interactions after multiplying the coupling value λ by $\sqrt{2}$; see Note.

Excited leptons have the same quantum numbers as other ortholeptons. See also the searches for ortholeptons in the "Searches for Heavy Leptons" section.

Limits for Excited e^* from Pair Production

These limits are obtained from $e^+e^- \rightarrow e^*e^{*-}$ and thus rely only on the (electroweak) charge of e^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the e^* coupling is assumed to be of sequential type. Possible t channel contribution from transition magnetic coupling is neglected. All limits assume $e^* \rightarrow e\gamma$ decay except the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.6	95	ADRIANI	93M L3	Z $\rightarrow e^*e^*$
>45.6	95	ABREU	92C DLPH	Z $\rightarrow e^*e^*$
>46.1	95	DECAMP	92 ALEP	Z $\rightarrow e^*e^*$
>44.9	95	AKRAWY	90I OPAL	Z $\rightarrow e^*e^*$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>29.8	95	43 BARDADIN-...	92 RVUE	$\Gamma(Z)$
>26.1	95	44 DECAMP	92 ALEP	Z $\rightarrow e^*e^*$; $\Gamma(Z)$
>33	95	44 ABREU	91F DLPH	Z $\rightarrow e^*e^*$; $\Gamma(Z)$
>45.0	95	45 ADEVA	90F L3	Z $\rightarrow e^*e^*$
>44.6	95	46 DECAMP	90G ALEP	$e^+e^- \rightarrow e^*e^*$
>30.2	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow e^*e^*$
>28.3	95	KIM	89 AMY	$e^+e^- \rightarrow e^*e^*$
>27.9	95	47 ABE	88B VNS	$e^+e^- \rightarrow e^*e^*$

43 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on $\Delta\Gamma(Z) < 36$ MeV.

44 Limit is independent of e^* decay mode.

45 ADEVA 90F is superseded by ADRIANI 93M.

46 Superseded by DECAMP 92.

47 ABE 88B limits assume $e^+e^- \rightarrow e^*e^{*-}$ with one photon exchange only and $e^* \rightarrow e\gamma$ giving $e\gamma\gamma$.

Limits for Excited e^* from Single Production

These limits are from $e^+e^- \rightarrow e^*e, W \rightarrow e^*\nu, \text{ or } ep \rightarrow e^*X$ and depend on transition magnetic coupling between e and e^* . All limits assume $e^* \rightarrow e\gamma$ decay except as noted. Limits from LEP, UA2, and H1 are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda - m_{e^*}$ plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>89	95	ADRIANI	93M L3	Z $\rightarrow ee^*, \lambda_Z > 0.5$
>88	95	ABREU	92C DLPH	Z $\rightarrow ee^*, \lambda_Z > 0.5$
>91	95	DECAMP	92 ALEP	Z $\rightarrow ee^*, \lambda_Z > 1$
>87	95	AKRAWY	90I OPAL	Z $\rightarrow ee^*, \lambda_Z > 0.5$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>86	95	48 ABT	93 H1	$ep \rightarrow e^*X$
		ADRIANI	93M L3	$\lambda_\gamma > 0.04$
		49 DERRICK	93B ZEUS	$ep \rightarrow e^*X$
>86	95	ABREU	92C DLPH	$e^+e^- \rightarrow ee^*, \lambda_\gamma > 0.1$
>88	95	50 ADEVA	90F L3	Z $\rightarrow ee^*, \lambda_Z > 0.5$
>86	95	50 ADEVA	90F L3	Z $\rightarrow ee^*, \lambda_Z > 0.04$
>81	95	51 DECAMP	90G ALEP	Z $\rightarrow ee^*, \lambda_Z > 1$
>50	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow ee^*, \lambda_\gamma > 0.04$
>56	95	KIM	89 AMY	$e^+e^- \rightarrow ee^*, \lambda_\gamma > 0.03$
none 23-54	95	52 ABE	88B VNS	$e^+e^- \rightarrow ee^*, \lambda_\gamma > 0.04$
>75	95	53 ANSARI	87D UA2	W $\rightarrow e^*\nu; \lambda_W > 0.7$
>63	95	53 ANSARI	87D UA2	W $\rightarrow e^*\nu; \lambda_W > 0.2$
>40	95	53 ANSARI	87D UA2	W $\rightarrow e^*\nu; \lambda_W > 0.09$

48 ABT 93 search for single e^* production via $e^*e\gamma$ coupling in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. See their Fig. 4 for exclusion plot in the $m_{e^*} - \lambda_\gamma$ plane.

49 DERRICK 93B search for single e^* production via $e^*e\gamma$ coupling in ep collisions with the decays $e^* \rightarrow e\gamma, eZ, \nu W$. See their Fig. 3 for exclusion plot in the $m_{e^*} - \lambda_\gamma$ plane.

50 Superseded by ADRIANI 93M.

51 Superseded by DECAMP 92.

52 ABE 88B limits use $e^+e^- \rightarrow ee^*$ where t-channel photon exchange dominates giving $e\gamma(e)$ (quasi-real compton scattering).

53 ANSARI 87D is at $E_{cm} = 546-630$ GeV.

Limits for Excited e^* from $e^+e^- \rightarrow \gamma\gamma$

These limits are derived from indirect effects due to e^* exchange in the t channel and depend on transition magnetic coupling between e and e^* . All limits are for $\lambda_\gamma = 1$. All limits except ABE 89J are for nonchiral coupling with $\eta_L = \eta_R = 1$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>127	95	ADRIANI	92B L3	
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>114	95	54 BUSKULIC	93Q ALEP	
> 99	95	55 BARDADIN-...	92 RVUE	
		DECAMP	92 ALEP	
		56 SHIMOZAWA	92 TOPZ	
>100	95	ABREU	91E DLPH	
>116	95	AKRAWY	91F OPAL	
> 83	95	ADEVA	90K L3	
> 82	95	AKRAWY	90F OPAL	
> 68	95	57 ABE	89J VNS	$\eta_L=1, \eta_R=0$
> 90.2	95	ADACHI	89B TOPZ	
> 65	95	KIM	89 AMY	

54 BUSKULIC 93Q obtain $\Lambda^+ > 121$ GeV (95%CL) from ALEPH experiment and $\Lambda^+ > 135$ GeV from combined TRISTAN and ALEPH data. These limits roughly correspond to limits on m_{e^*} .

55 BARDADIN-OTWINOWSKA 92 limit from fit to the combined data of DECAMP 92, ABREU 91E, ADEVA 90K, AKRAWY 91F.

56 SHIMOZAWA 92 fit the data to the limiting form of the cross section with $m_{e^*} \gg E_{cm}$ and obtain $m_{e^*} > 168$ GeV at 95%CL. Use of the full form would reduce this limit by a few GeV. The statistically unexpected large value is due to fluctuation in the data.

57 The ABE 89J limit assumes chiral coupling. This corresponds to $\lambda_\gamma = 0.7$ for nonchiral coupling.

Indirect Limits for Excited e^*

These limits make use of loop effects involving e^* and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	58 DORENBOSCH...	89 CHR	$\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ and $\nu_\mu e \rightarrow \nu_\mu e$
	59 GRIFOLS	86 THEO	$\nu_\mu e \rightarrow \nu_\mu e$
	60 RENARD	82 THEO	$g-2$ of electron

58 DORENBOSCH 89 obtain the limit $\lambda_{cut}^2 \Lambda_{cut}^2 / m_{e^*}^2 < 2.6$ (95% CL), where Λ_{cut} is the cutoff scale, based on the one-loop calculation by GRIFOLS 86. If one assumes that $\Lambda_{cut} = 1$ TeV and $\lambda_\gamma = 1$, one obtains $m_{e^*} > 620$ GeV. However, one generally expects $\lambda_\gamma \approx m_{e^*} / \Lambda_{cut}$ in composite models.

59 GRIFOLS 86 uses $\nu_\mu e \rightarrow \nu_\mu e$ and $\bar{\nu}_\mu e \rightarrow \bar{\nu}_\mu e$ data from CHARM Collaboration to derive mass limits which depend on the scale of compositeness.

60 RENARD 82 derived from $g-2$ data limits on mass and couplings of e^* and μ^* . See figures 2 and 3 of the paper.

MASS LIMITS for Excited μ^* Limits for Excited μ^* from Pair Production

These limits are obtained from $e^+e^- \rightarrow \mu^*\mu^{*-}$ and thus rely only on the (electroweak) charge of μ^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the μ^* coupling is assumed to be of sequential type. All limits assume $\mu^* \rightarrow \mu\gamma$ decay except for the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.6	95	ADRIANI	93M L3	Z $\rightarrow \mu^*\mu^*$
>45.6	95	ABREU	92C DLPH	Z $\rightarrow \mu^*\mu^*$
>46.1	95	DECAMP	92 ALEP	Z $\rightarrow \mu^*\mu^*$
>44.9	95	AKRAWY	90I OPAL	Z $\rightarrow \mu^*\mu^*$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
>29.8	95	61 BARDADIN-...	92 RVUE	$\Gamma(Z)$
>26.1	95	62 DECAMP	92 ALEP	Z $\rightarrow \mu^*\mu^*$; $\Gamma(Z)$
>33	95	62 ABREU	91F DLPH	Z $\rightarrow \mu^*\mu^*$; $\Gamma(Z)$
>45.3	95	63 ADEVA	90F L3	Z $\rightarrow \mu^*\mu^*$
>44.6	95	64 DECAMP	90G ALEP	$e^+e^- \rightarrow \mu^*\mu^*$
>29.9	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow \mu^*\mu^*$
>28.3	95	KIM	89 AMY	$e^+e^- \rightarrow \mu^*\mu^*$

61 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on $\Delta\Gamma(Z) < 36$ MeV.

62 Limit is independent of μ^* decay mode.

63 Superseded by ADRIANI 93M.

64 Superseded by DECAMP 92.

Limits for Excited μ (μ^*) from Single Production

These limits are from $e^+e^- \rightarrow \mu^*\mu$ and depend on transition magnetic coupling between μ and μ^* . All limits assume $\mu^* \rightarrow \mu\gamma$ decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda-m_{\mu^*}$ plane. See the original papers.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>89	95	ADRIANI	93M L3	$Z \rightarrow \mu\mu^*, \lambda_Z > 0.5$
>88	95	ABREU	92C DLPH	$Z \rightarrow \mu\mu^*, \lambda_Z > 0.5$
>91	95	DECAMP	92 ALEP	$Z \rightarrow \mu\mu^*, \lambda_Z > 1$
>87	95	AKRAWY	90I OPAL	$Z \rightarrow \mu\mu^*, \lambda_Z > 1$
••• We do not use the following data for averages, fits, limits, etc. •••				
>85	95	65 ADEVA	90F L3	$Z \rightarrow \mu\mu^*, \lambda_Z > 1$
>75	95	65 ADEVA	90F L3	$Z \rightarrow \mu\mu^*, \lambda_Z > 0.1$
>80	95	66 DECAMP	90G ALEP	$e^+e^- \rightarrow \mu\mu^*, \lambda_Z=1$
>50	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow \mu\mu^*, \lambda_\gamma=0.7$
>46	95	KIM	89 AMY	$e^+e^- \rightarrow \mu\mu^*, \lambda_\gamma=0.2$

65 Superseded by ADRIANI 93M.

66 Superseded by DECAMP 92.

Indirect Limits for Excited μ (μ^*)

These limits make use of loop effects involving μ^* and are therefore subject to theoretical uncertainty.

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
••• We do not use the following data for averages, fits, limits, etc. •••			
	67 RENARD	82 THEO	$g-2$ of muon

67 RENARD 82 derived from $g-2$ data limits on mass and couplings of e^* and μ^* . See figures 2 and 3 of the paper.

MASS LIMITS for Excited τ (τ^*)**Limits for Excited τ (τ^*) from Pair Production**

These limits are obtained from $e^+e^- \rightarrow \tau^*\tau^*$ and thus rely only on the (electroweak) charge of τ^* . Form factor effects are ignored unless noted. For the case of limits from Z decay, the τ^* coupling is assumed to be of sequential type. All limits assume $\tau^* \rightarrow \tau\gamma$ decay except for the limits from $\Gamma(Z)$.

For limits prior to 1987, see our 1992 edition (Physical Review **D45**, 1 June, Part II (1992)).

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.6	95	ADRIANI	93M L3	$Z \rightarrow \tau^*\tau^*$
>45.3	95	ABREU	92C DLPH	$Z \rightarrow \tau^*\tau^*$
>46.0	95	DECAMP	92 ALEP	$Z \rightarrow \tau^*\tau^*$
>44.9	95	AKRAWY	90I OPAL	$Z \rightarrow \tau^*\tau^*$
••• We do not use the following data for averages, fits, limits, etc. •••				
>29.8	95	68 BARDADIN-...	92 RVUE	$\Gamma(Z)$
>26.1	95	69 DECAMP	92 ALEP	$Z \rightarrow \tau^*\tau^*; \Gamma(Z)$
>33	95	69 ABREU	91F DLPH	$Z \rightarrow \tau^*\tau^*; \Gamma(Z)$
>45.5	95	70 ADEVA	90L L3	$Z \rightarrow \tau^*\tau^*$
>41.2	95	71 DECAMP	90G ALEP	$e^+e^- \rightarrow \tau^*\tau^*$
>29.0	95	ADACHI	89B TOPZ	$e^+e^- \rightarrow \tau^*\tau^*$

68 BARDADIN-OTWINOWSKA 92 limit is independent of decay modes. Based on $\Delta\Gamma(Z) < 36$ MeV.

69 Limit is independent of τ^* decay mode.

70 Superseded by ADRIANI 93M.

71 Superseded by DECAMP 92.

Limits for Excited τ (τ^*) from Single Production

These limits are from $e^+e^- \rightarrow \tau^*\tau$ and depend on transition magnetic coupling between τ and τ^* . All limits assume $\tau^* \rightarrow \tau\gamma$ decay. Limits from LEP are for chiral coupling, whereas all other limits are for nonchiral coupling, $\eta_L = \eta_R = 1$. In most papers, the limit is expressed in the form of an excluded region in the $\lambda-m_{\tau^*}$ plane. See the original papers.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>88	95	ADRIANI	93M L3	$Z \rightarrow \tau\tau^*, \lambda_Z > 0.5$
>87	95	ABREU	92C DLPH	$Z \rightarrow \tau\tau^*, \lambda_Z > 0.5$
>90	95	DECAMP	92 ALEP	$Z \rightarrow \tau\tau^*, \lambda_Z > 0.18$
>86.5	95	AKRAWY	90I OPAL	$Z \rightarrow \tau\tau^*, \lambda_Z > 1$
••• We do not use the following data for averages, fits, limits, etc. •••				
>88	95	72 ADEVA	90L L3	$Z \rightarrow \tau\tau^*, \lambda_Z > 1$
>59	95	73 DECAMP	90G ALEP	$Z \rightarrow \tau\tau^*, \lambda_Z=1$
>40	95	74 BARTEL	86 JADE	$e^+e^- \rightarrow \tau\tau^*, \lambda_\gamma=1$
>41.4	95	75 BEHREND	86 CELL	$e^+e^- \rightarrow \tau\tau^*, \lambda_\gamma=1$
>40.8	95	75 BEHREND	86 CELL	$e^+e^- \rightarrow \tau\tau^*, \lambda_\gamma=0.7$

72 Superseded by ADRIANI 93M.

73 Superseded by DECAMP 92.

74 BARTEL 86 is at $E_{cm} = 30-46.78$ GeV.

75 BEHREND 86 limit is at $E_{cm} = 33-46.8$ GeV.

MASS LIMITS for Excited Neutrino (ν^*)**Limits for Excited ν (ν^*) from Pair Production**

These limits are obtained from $Z \rightarrow \nu^*\nu^*$ decay and thus rely only on the (electroweak) charge of ν^* . Form factor effects are ignored unless noted. The ν^* coupling is assumed to be of sequential type. Limits assume $\nu^* \rightarrow \nu\gamma$ decay except for the $\Gamma(Z)$ measurement which makes no assumption about decay mode.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>47	95	76 DECAMP	92 ALEP	
••• We do not use the following data for averages, fits, limits, etc. •••				
>43.7	95	77 BARDADIN-...	92 RVUE	$\Gamma(Z)$
>42.6	95	78 DECAMP	92 ALEP	$\Gamma(Z)$
>35.4	95	79,80 DECAMP	90G ALEP	$\Gamma(Z)$
>46	95	80,81 DECAMP	90G ALEP	

76 Limit is based on $B(Z \rightarrow \nu^*\bar{\nu}^*) \times B(\nu^* \rightarrow \nu\gamma)^2 < 5 \times 10^{-5}$ (95%CL) assuming Dirac ν^* , $B(\nu^* \rightarrow \nu\gamma) = 1$.

77 BARDADIN-OTWINOWSKA 92 limit is for Dirac ν^* . Based on $\Delta\Gamma(Z) < 36$ MeV. The limit is 36.4 GeV for Majorana ν^* , 45.4 GeV for homodoublet ν^* .

78 Limit is for Dirac ν^* . The limit is 34.6 GeV for Majorana ν^* , 45.4 GeV for homodoublet ν^* .

79 DECAMP 90G limit is from excess $\Delta\Gamma(Z) < 89$ MeV. The above value is for Dirac ν^* ; 26.6 GeV for Majorana ν^* ; 44.8 GeV for homodoublet ν^* .

80 Superseded by DECAMP 92.

81 DECAMP 90G limit based on $B(Z \rightarrow \nu^*\bar{\nu}^*) \times B(\nu^* \rightarrow \nu\gamma)^2 < 7 \times 10^{-5}$ (95%CL), assuming Dirac ν^* , $B(\nu^* \rightarrow \nu\gamma) = 1$.

Limits for Excited ν (ν^*) from Single Production

These limits are from $Z \rightarrow \nu\nu^*$ or $e\bar{p} \rightarrow \nu^*X$ and depend on transition magnetic coupling between ν/e and ν^* . Assumptions about ν^* decay mode are given in footnotes.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>91	95	ADRIANI	93M L3	$\lambda_Z > 1, \nu^* \rightarrow \nu\gamma$
>89	95	ADRIANI	93M L3	$\lambda_Z > 1, \nu_e^* \rightarrow eW$
>91	95	82 DECAMP	92 ALEP	$\lambda_Z > 1$
••• We do not use the following data for averages, fits, limits, etc. •••				
		83 ABT	93 H1	$e\bar{p} \rightarrow \nu^*X$
>87	95	ADRIANI	93M L3	$\lambda_Z > 0.1, \nu^* \rightarrow \nu\gamma$
>74	95	ADRIANI	93M L3	$\lambda_Z > 0.1, \nu_e^* \rightarrow eW$
		84 BARDADIN-...	92 RVUE	
>74	95	82 DECAMP	92 ALEP	$\lambda_Z > 0.034$
>91	95	85,86 ADEVA	90G L3	$\lambda_Z > 1$
>83	95	86 ADEVA	90G L3	$\lambda_Z > 0.1, \nu^* \rightarrow \nu\gamma$
>74	95	86 ADEVA	90G L3	$\lambda_Z > 0.1, \nu_e^* \rightarrow eW$
>90	95	87,88 DECAMP	90G ALEP	$\lambda_Z > 1$
>74.7	95	87,88 DECAMP	90G ALEP	$\lambda_Z > 0.06$

82 DECAMP 92 limit is based on $B(Z \rightarrow \nu^*\bar{\nu}^*) \times B(\nu^* \rightarrow \nu\gamma) < 2.7 \times 10^{-5}$ (95%CL) assuming Dirac ν^* , $B(\nu^* \rightarrow \nu\gamma) = 1$.

83 ABT 93 search for single ν^* production via ν^*eW coupling in $e\bar{p}$ collisions with the decays $\nu^* \rightarrow \nu\gamma, \nu Z, eW$. See their Fig. 4 for exclusion plot in the $m_{\nu^*}-\lambda_{W\nu}$ plane.

84 See Fig. 5 of BARDADIN-OTWINOWSKA 92 for combined limit of ADEVA 90G, DECAMP 90G, and DECAMP 92.

85 Limit is either for $\nu^* \rightarrow \nu\gamma$ or $\nu^* \rightarrow eW$.

86 Superseded by ADRIANI 93M.

87 DECAMP 90G limit based on $B(Z \rightarrow \nu\nu^*) \times B(\nu^* \rightarrow \nu\gamma) < 6 \times 10^{-5}$ (95%CL), assuming $B(\nu^* \rightarrow \nu\gamma) = 1$.

88 Superseded by DECAMP 92.

MASS LIMITS for Excited q (q^*)**Limits for Excited q (q^*) from Pair Production**

These limits are obtained from $e^+e^- \rightarrow q^*q^*$ and thus rely only on the (electroweak) charge of the q^* . Form factor effects are ignored unless noted. Assumptions about the q^* decay are given in the comments and footnotes.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>45.6	95	89 ADRIANI	93M L3	u or d type, $Z \rightarrow q^*q^*$
>45	95	90 DECAMP	92 ALEP	u or d type, $Z \rightarrow q^*q^*$
••• We do not use the following data for averages, fits, limits, etc. •••				
>41.7	95	91 ADRIANI	92F L3	$Z \rightarrow q^*q^*$
>44.7	95	92 BARDADIN-...	92 RVUE	u -type, $\Gamma(Z)$
>40.6	95	92 BARDADIN-...	92 RVUE	d -type, $\Gamma(Z)$
>44.2	95	93 DECAMP	92 ALEP	u -type, $\Gamma(Z)$
>45	95	93 DECAMP	92 ALEP	d -type, $\Gamma(Z)$
>45	95	93 ABREU	91F DLPH	u -type, $\Gamma(Z)$
>21.1	95	93 ABREU	91F DLPH	d -type, $\Gamma(Z)$
		94 BEHREND	86C CELL	$e(q^*) = -1/3, q^* \rightarrow q\bar{q}$
>22.3	95	94 BEHREND	86C CELL	$e(q^*) = 2/3, q^* \rightarrow q\bar{q}$
>22.5	95	94 BEHREND	86C CELL	$e(q^*) = -1/3, q^* \rightarrow q\bar{q}$
		94 BEHREND	86C CELL	$q\gamma$
>23.2	95	94 BEHREND	86C CELL	$e(q^*) = 2/3, q^* \rightarrow q\bar{q}$

89 ADRIANI 93M limit is valid for $B(q^* \rightarrow q\bar{q}) > 0.25$ (0.17) for up (down) type.

90 Limit is for $B(q^* \rightarrow q\bar{q}) + B(q^* \rightarrow q\gamma) = 1$.

Searches Full Listings

Quark and Lepton Compositeness

91 ADRIANI 92F search for $Z \rightarrow q^* \bar{q}^*$ followed with $q^* \rightarrow q\gamma$ decays and give the limit $\sigma_Z \cdot B(Z \rightarrow q^* \bar{q}^*) \cdot B^2(q^* \rightarrow q\gamma) < 2$ pb at 95%CL. Assuming five flavors of degenerate q^* of homodoublet type, $B(q^* \rightarrow q\gamma) < 4\%$ is obtained for $m_{q^*} < 45$ GeV.

92 BARDADIN-OTWINOWSKA 92 limit based on $\Delta\Gamma(Z) < 36$ MeV.

93 These limits are independent of decay modes.

94 BEHREND 86C search for $e^+ e^- \rightarrow q^* \bar{q}^*$ for $m_{q^*} > 5$ GeV. But $m < 5$ GeV excluded by total hadronic cross section. The limits are for point-like photon couplings of excited quarks.

Limits for Excited q (q^*) from Single Production

These limits are from $e^+ e^- \rightarrow q^* \bar{q}^*$ or $p\bar{p} \rightarrow q^* X$ and depend on transition magnetic couplings between q and q^* . Assumptions about q^* decay mode are given in the footnotes and comments.

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>540 (CL = 95%) OUR EVALUATION				
none 80-540	95	95 ABE	94 CDF	$p\bar{p} \rightarrow q^* X, q^* \rightarrow q\gamma, qW$
>288	90	96 ALITTI	93 UA2	$p\bar{p} \rightarrow q^* X, q^* \rightarrow qg$
> 88	95	97 DECAMP	92 ALEP	$Z \rightarrow qq^*, \lambda_Z > 1$
> 86	95	97 AKRAWY	90J OPAL	$Z \rightarrow qq^*, \lambda_Z > 1.2$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
> 79	95	98 ADRIANI	93M L3	$\lambda_Z(L3) > 0.06$
		99 ABREU	92D DLPH	$Z \rightarrow qq^*$
		100 ADRIANI	92F L3	$Z \rightarrow qq^*$
> 75	95	98 DECAMP	92 ALEP	$Z \rightarrow qq^*, \lambda_Z > 1$
		101 ALBAJAR	89 UA1	$p\bar{p} \rightarrow q^* X, q^* \rightarrow qW$
> 39	95	102 BEHREND	86C CELL	$e^+ e^- \rightarrow q^* \bar{q}^* (q^* \rightarrow qg, q\gamma), \lambda_{\gamma=1}$

95 ABE 94 search for resonances in jet- γ and jet- W invariant mass in $p\bar{p}$ collisions at $E_{cm} = 1.8$ TeV. The limit is for $f_s = f = f' = \Lambda/m_{q^*}$ and u^* and d^* are assumed to be degenerate. See their Fig. 4 for the excluded region in m_{q^*} - f plane.

96 ALITTI 93 search for resonances in the two-jet invariant mass. The limit is for $f_s = f = f' = \Lambda/m_{q^*}$. u^* and d^* are assumed to be degenerate. If not, the limit for u^* (d^*) is 277 (247) GeV if $m_{q^*} \gg m_{u^*}$ ($m_{u^*} \gg m_{d^*}$).

97 Assumes $B(q^* \rightarrow q\gamma) = 0.1$.

98 Assumes $B(q^* \rightarrow qg) = 1$.

99 ABREU 92D give $\sigma(e^+ e^- \rightarrow Z \rightarrow q^* \bar{q}^* \text{ or } q\bar{q}^*) \times B(q^* \rightarrow q\gamma) < 15$ pb (95% CL) for $m_{q^*} < 80$ GeV.

100 ADRIANI 92F search for $Z \rightarrow qq^*$ with $q^* \rightarrow q\gamma$ and give the limit $\sigma_Z \cdot B(Z \rightarrow qq^*) \cdot B(q^* \rightarrow q\gamma) < (2-10)$ pb (95%CL) for $m_{q^*} = (46-82)$ GeV.

101 ALBAJAR 89 give $\sigma(q^* \rightarrow W + \text{jet})/\sigma(W) < 0.019$ (90% CL) for $m_{q^*} > 220$ GeV.

102 BEHREND 86C has $E_{cm} = 42.5-46.8$ GeV. See their Fig. 3 for excluded region in the $m_{q^*} - (\lambda_\gamma/m_{q^*})^2$ plane. The limit is for $\lambda_\gamma = 1$ with $\eta_L = \eta_R = 1$.

MASS LIMITS for Color Sextet Quarks (q_6)

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>84	95	103 ABE	89D CDF	$p\bar{p} \rightarrow q_6 \bar{q}_6$

103 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color sextet quark is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. A limit of 121 GeV is obtained for a color decuplet.

MASS LIMITS for Color Octet Charged Leptons (ℓ_8)

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>86	95	104 ABE	89D CDF	Stable ℓ_8 : $p\bar{p} \rightarrow \ell_8 \bar{\ell}_8$

• • • We do not use the following data for averages, fits, limits, etc. • • •

none 3.0-30.3	95	105 ABT	93 H1	$e_8: e^+ e^- \rightarrow e_8 X$
none 3.5-30.3	95	106 KIM	90 AMY	$e_8: e^+ e^- \rightarrow ee + \text{jets}$
>19.8	95	107 KIM	90 AMY	$e_8: e^+ e^- \rightarrow gg; R$
none 5-23.2	95	108 BARTEL	87B JADE	$\mu_8, \tau_8: e^+ e^-; R$
		108 BARTEL	87B JADE	$\mu_8: e^+ e^- \rightarrow \mu\mu + \text{jets}$
		109 BARTEL	85K JADE	$e_8: e^+ e^- \rightarrow gg; R$

104 ABE 89D look for pair production of unit-charged particles which leave the detector before decaying. In the above limit the color octet lepton is assumed to fragment into a unit-charged or neutral hadron with equal probability and to have long enough lifetime not to decay within the detector. The limit improves to 99 GeV if it always fragments into a unit-charged hadron.

105 ABT 93 search for e_8 production via e -gluon fusion in ep collisions with $e_8 \rightarrow eg$. See their Fig. 3 for exclusion plot in the m_{e_8} - Λ plane for $m_{e_8} = 35-220$ GeV.

106 KIM 90 is at $E_{cm} = 50-60.8$ GeV. The same assumptions as in BARTEL 87B are used.

107 KIM 90 result $(m_{e_8} \Lambda_M)^{1/2} > 178.4$ GeV (95%CL, $\alpha_s = 0.16$ used) is subject to the same restriction as for BARTEL 85K.

108 BARTEL 87B is at $E_{cm} = 46.3-46.78$ GeV. The limits assume ℓ_8 pair production cross sections to be eight times larger than those of the corresponding heavy lepton pair production.

109 In BARTEL 85K, R can be affected by $e^+ e^- \rightarrow gg$ via e_q exchange. Their limit $m_{e_8} > 173$ GeV (CL=95%) at $\lambda = m_{e_8}/\Lambda_M = 1$ ($\eta_L = \eta_R = 1$) is not listed above because the cross section is sensitive to the product $\eta_L \eta_R$, which should be absent in ordinary theory with electronic chiral invariance.

MASS LIMITS for Color Octet Neutrinos (ν_8)

VALUE (GeV)	CL%	DOCUMENT ID	TECN	COMMENT
>110	90	110 BARGER	89 RVUE	$\nu_8: p\bar{p} \rightarrow \nu_8 \bar{\nu}_8$
• • • We do not use the following data for averages, fits, limits, etc. • • •				
none 3.8-29.8	95	111 KIM	90 AMY	$\nu_8: e^+ e^- \rightarrow \text{coplanar jets}$
none 9-21.9	95	112 BARTEL	87B JADE	$\nu_8: e^+ e^- \rightarrow \text{coplanar jets}$

110 BARGER 89 used ABE 89B limit for events with large missing transverse momentum. Two-body decay $\nu_8 \rightarrow \nu g$ is assumed.

111 KIM 90 is at $E_{cm} = 50-60.8$ GeV. The same assumptions as in BARTEL 87B are used.

112 BARTEL 87B is at $E_{cm} = 46.3-46.78$ GeV. The limit assumes the ν_8 pair production cross section to be eight times larger than that of the corresponding heavy neutrino pair production. This assumption is not valid in general for the weak couplings, and the limit can be sensitive to its $SU(2)_L \times U(1)_Y$ quantum numbers.

MASS LIMITS for W_8 (Color Octet W Boson)

VALUE (GeV)	DOCUMENT ID	TECN	COMMENT
• • • We do not use the following data for averages, fits, limits, etc. • • •			
	113 ALBAJAR	89 UA1	$p\bar{p} \rightarrow W_8 X, W_8 \rightarrow Wg$

113 ALBAJAR 89 give $\sigma(W_8 \rightarrow W + \text{jet})/\sigma(W) < 0.019$ (90% CL) for $m_{W_8} > 220$ GeV.

Limits on $ZZ\gamma$ Coupling

Limits are for the electric dipole transition form factor for $Z \rightarrow \gamma Z^*$ parametrized as $f(s') = \beta(s'/m_Z^2 - 1)$, where s' is the virtual Z mass. In the Standard Model $\beta \sim 10^{-5}$.

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<0.80	95	ADRIANI	92J L3	$Z \rightarrow \gamma \nu \bar{\nu}$

REFERENCES FOR Searches for Quark and Lepton Compositeness

ABE	94	PRL 72 3004	+Albrow, Amidei, Anway-Wiese, Apollinari+ (CDF Collab.)
DIAZCRUZ	94	PR D49 R2149	Diaz Cruz, Sampayo (CINV)
ABE	93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+ (CDF Collab.)
ABT	93	NP B396 3	+Andreev, Andrieu, Appuhn, Arpagaus+ (H1 Collab.)
ADRIANI	93M	PRPL 236 1	+Aguiar-Benitez, Ahlen, Alcaraz, Aloisio+ (L3 Collab.)
ALITTI	93	NP B400 3	+Ambrosini, Ansari, Autiero, Barye+ (UA2 Collab.)
BUSKULIC	93Q	ZPHY C59 215	+Decamp, Goy, Lees, Minard, Mours+ (ALEPH Collab.)
DERRICK	93F	PL B316 207	+Krakauer, Magill, Musgrave, Repond+ (ZEUS Collab.)
ABE	92B	PRL 68 1463	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE	92D	PRL 68 1104	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABE	92M	PRL 69 2896	+Amidei, Anway-Wiese, Apollinari, Atac+ (CDF Collab.)
ABREU	92C	ZPHY C53 41	+Adam, Adami, Adye, Akesson+ (DELPHI Collab.)
ABREU	92D	ZPHY C53 555	+Adam, Adami, Adye, Akesson, Alekseev+ (DELPHI Collab.)
ADRIANI	92B	PL B288 404	+Aguiar-Benitez, Ahlen, Akbari, Alcaraz+ (L3 Collab.)
ADRIANI	92F	PL B292 472	+Aguiar-Benitez, Ahlen, Akbari, Alcaraz+ (L3 Collab.)
ADRIANI	92J	PL B297 469	+Aguiar-Benitez, Ahlen, Alcaraz, Aloisio+ (L3 Collab.)
BARDADIN...	92	ZPHY C55 163	Bardadin-Otwinowska (CLER)
DECAMP	92	PRPL 216 253	+Deschizeaux, Goy, Lees, Minard+ (ALEPH Collab.)
HOWELL	92	PL B291 206	+Koltick, Tauchi, Miyamoto, Kichimi+ (TOPAZ Collab.)
KROHA	92	PR D46 58	(ROCH)
PDG	92	PR D45, 1 June, Part II	Hikasa, Barnett, Stone+ (KEK, LBL, BOST+)
SHIMOZAWA	92	PL B284 144	+Fujimoto, Abe, Adachi, Doser+ (TOPAZ Collab.)
ABE	91D	PRL 67 2418	+Amidei, Apollinari, Atac, Auchincloss+ (CDF Collab.)
ABREU	91E	PL B268 296	+Adam, Adami, Adye, Akesson+ (DELPHI Collab.)
ABREU	91F	NP B367 511	+Adam, Adami, Adye, Akesson+ (DELPHI Collab.)
ADACHI	91	PL B255 613	+Anazawa, Doser, Enomoto+ (TOPAZ Collab.)
AKRAWY	91F	PL B257 531	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
ALITTI	91B	PL B257 232	+Ansari, Autiero, Barye, Blaylock+ (UA2 Collab.)
BEHREND	91B	ZPHY C51 143	+Criegee, Field, Franke, Jung+ (CELLO Collab.)
BEHREND	91C	ZPHY C51 149	+Criegee, Field, Franke, Jung, Meyer+ (CELLO Collab.)
Also	91B	ZPHY C51 143	+Behrend, Criegee, Field, Franke, Jung+ (CELLO Collab.)
ABE	90I	ZPHY C48 13	+Amako, Arai, Asano, Chiba+ (VENUS Collab.)
ADEVA	90F	PL B247 177	+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
ADEVA	90K	PL B250 199	+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
ADEVA	90L	PL B250 205	+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
ADEVA	90O	PL B252 525	+Adriani, Aguiar-Benitez, Akbari, Alcaraz+ (L3 Collab.)
AKRAWY	90F	PL B242 133	+Alexander, Allison, Allport+ (OPAL Collab.)
AKRAWY	90I	PL B244 135	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
AKRAWY	90J	PL B246 285	+Alexander, Allison, Allport, Anderson+ (OPAL Collab.)
DECAMP	90G	PL B236 501	+Deschizeaux, Lees, Minard+ (ALEPH Collab.)
DECAMP	90O	PL B250 172	+Deschizeaux, Goy, Lees+ (ALEPH Collab.)
KIM	90	PL B240 243	+Breedon, Ko, Lander, Maeshima, Malchow+ (AMY Collab.)
ABE	89	PRL 62 613	+Amidei, Apollinari, Ascovi, Atac+ (CDF Collab.)
ABE	89B	PRL 62 1825	+Amidei, Apollinari, Ascovi, Atac+ (CDF Collab.)
ABE	89D	PRL 63 1447	+Amidei, Apollinari, Ascovi, Atac+ (CDF Collab.)
ABE	89H	PRL 62 3020	+Amidei, Apollinari, Ascovi, Atac+ (CDF Collab.)
ABE	89J	ZPHY C45 175	+Amako, Arai, Fukawa+ (VENUS Collab.)
ABE	89L	PL B232 425	+Amako, Arai, Asano, Chiba+ (VENUS Collab.)
ADACHI	89B	PL B228 553	+Aihara, Doser, Enomoto, Fujii+ (TOPAZ Collab.)
ALBAJAR	89	ZPHY C44 15	+Albrow, Alkoffer, Arrison, Astbury+ (UA1 Collab.)

Quark and Lepton Compositeness, Other Stable Particle Searches

BARGER	89	PL B220 464	+Hagiwara, Han, Zeppenfeld	(WISC, KEK)
BEHREND	89B	PL B222 163	+Criegee, Dainton, Fiedl, Franke+	(CELLO Collab.)
BRAUNSCH...	89C	ZPHY C43 549	+Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
DORENBOS...	89	ZPHY C41 567	+Dorenbosch, Udo, Allaby, Amaldi+	(CHARM Collab.)
HAGIWARA	89	PL B219 369	+Sakuda, Terunuma	(KEK, DURH, HIRO)
KIM	89	PL B223 476	+Kim, Kang, Lee, Myung, Bacala	(AMY Collab.)
ABE	88B	PL B213 400	+Amako, Arai, Asano, Chiba, Chiba+	(VENUS Collab.)
BARINGER	88	PL B206 551	+Bijlsma, De Bonte, Koitick, Low+	(HRS Collab.)
BRAUNSCH...	88	ZPHY C37 171	+Braunschweig, Gerhards+	(TASSO Collab.)
BRAUNSCH...	88D	ZPHY C40 163	+Braunschweig, Gerhards, Kirschfink+	(TASSO Collab.)
ANSARI	87D	PL B195 613	+Bagnaia, Banner+	(UA2 Collab.)
BARTEL	87B	ZPHY C36 15	+Becker, Felst+	(JADE Collab.)
BEHREND	87C	PL B191 209	+Buerger, Criegee, Dainton+	(CELLO Collab.)
FERNANDEZ	87B	PR D35 10	+Ford, Qi, Read, Smith, Camporesi+	(MAC Collab.)
ARNISON	86C	PL B172 461	+Albrow, Altkofer+	(UA1 Collab.)
ARNISON	86B	PL B177 244	+Albjar, Albrow+	(UA1 Collab.)
BARTEL	86	ZPHY C31 359	+Becker, Felst, Haidt+	(JADE Collab.)
BARTEL	86C	ZPHY C30 371	+Becker, Cords, Felst, Haidt+	(JADE Collab.)
BEHREND	86	PL 168B 420	+Buerger, Criegee, Fenner+	(CELLO Collab.)
BEHREND	86C	PL B181 178	+Buerger, Criegee, Dainton+	(CELLO Collab.)
DERRICK	86	PL 166B 463	+Gan, Kooijman, Loos+	(HRS Collab.)
Also	86B	PR D34 3286	+Derrick, Gan, Kooijman, Loos, Musgrave+	(HRS Collab.)
DERRICK	86B	PR D34 3286	+Gan, Kooijman, Loos, Musgrave+	(HRS Collab.)
GRIFOLS	86	PL 168B 264	+Peris	(BARC)
JODIDIO	86	PR D34 1967	+Balke, Carr, Gidal, Shinsky+	(LBL, NWES, TRIU)
Also	88	PR D37 237 erratum	Jodidio, Balke, Carr+	(LBL, NWES, TRIU)
APPEL	85	PL 160B 349	+Bagnaia, Banner+	(UA2 Collab.)
BARTEL	85K	PL 160B 337	+Becker, Cords, Eichler+	(JADE Collab.)
BERGER	85	ZPHY C28 1	+Genzel, Lackas, Pielorz+	(PLUTO Collab.)
BERGER	85B	ZPHY C27 341	+Deuter, Genzel, Lackas, Pielorz+	(PLUTO Collab.)
BAGNAIA	84C	PL 138B 430	+Banner, Battistoni+	(UA2 Collab.)
BARTEL	84D	PL 146B 437	+Becker, Bowdery, Cords+	(JADE Collab.)
BARTEL	84E	PL 146B 121	+Becker, Bowdery, Cords, Felst+	(JADE Collab.)
EICHTEN	84	RMP 56 579	+Hincliffe, Lane, Quigg	(FNAL, LBL, OSU)
ALTHOFF	83C	PL 126B 493	+Fischer, Burkhardt+	(TASSO Collab.)
RENARD	82	PL 116B 264		(CERN)

Other Stable Particle Searches

OMITTED FROM SUMMARY TABLE
NOTE ON OTHER STABLE PARTICLE SEARCHES

We collect here those searches which do not appear in any of the above search categories. These include heavy particle searches in accelerator experiments, in cosmic rays, and in matter. Searches are also listed for light particles, highly ionizing particles, penetrating non-neutrino-like particles, and tachyons. Note that searches appear in separate sections elsewhere for Higgs bosons (and technipions), other heavy bosons (including W_R , W' , Z' , leptoquarks, axiguons), axions (including pseudo-Goldstone bosons, Majorons, familons), heavy leptons, heavy neutrinos, free quarks and monopoles, supersymmetry, and compositeness.

Centauro Production Cross Section in Accelerator Experiments

VALUE (cm ²)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.005σ (nondiff.)	95	0	¹ ALNER	86 UA5	p \bar{p} collider
<1. × 10 ⁻³⁰		0	² ARNISON	83B UA1	p \bar{p} collider
		0	³ ALPGARD	82 UA5	p \bar{p} collider

¹ALNER 86 is CERN collider experiment at $W_{cm} = 900$ GeV. Looked for high multiplicity, low EM content in measured high p_T events from an unbiased sample of 5500 events. No candidates observed.
²ARNISON 83B is CERN collider experiment with $W_{cm} = 540$ GeV. Looked for events with large hadronic and low electromagnetic content. None in 48000 low bias events.
³ALPGARD 82 is CERN collider experiment with $W_{cm} = 540$ GeV (155 TeV lab equivalent). Observed no large charged multiplicity events with photon multiplicity consistent with zero in 3600 inelastic events.

Centauro Production in Cosmic Ray Interactions

A Centauro event is characterized by a hadronic event with high multiplicity, high mean p_T , and unusually small photon energy.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.03	95	0	⁴ REN	88 EMUL	e (shower) > 100 TeV
		1	BORISOV	87 EMUL	
			BAYBURINA	81 EMUL	
			LATTES	80 EMUL	

⁴REN 88 limit is for the fraction of Centauro events in the sample of hadronic showers with energy exceeding 100 TeV. No candidates were observed despite a total exposure exceeding that of previous experiments.

Light (between μ and e Masses) Particle MASS

VALUE (m _e)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
none 110-180		0	⁵ VIERTEL	78 CNTR	$\tau > 2 \times 10^{-5}$ s
none 2-13		0	⁶ BLAGOV	75 CNTR	Splnor, $\tau > 2 \times 10^{-10}$
none 2-10.6		0	⁶ BLAGOV	75 CNTR	Scalar, $\tau > 2 \times 10^{-10}$ s
none 5-175		0	COWARD	63 CNTR	Splnor, $\tau > 22 \times 10^{-10}$
none 5-175		0	COWARD	63 CNTR	Scalar, $\tau > 68 \times 10^{-10}$
none 6-25		0	BELOUSOV	60 CNTR	Splnor, $\tau > 1 \times 10^{-8}$
none 2-25		0	GORBUNOV	60 CC	Splnor, $\tau > 1 \times 10^{-9}$

⁵VIERTEL 78 searches for $\mu^+ \rightarrow X^+ \nu$. Finds BR < 8.5 × 10⁻⁶ in mass range given above (CL = 90%). Best limit BR < 5. × 10⁻⁷ (CL = 90%) is found at mass = 80 MeV.
⁶BLAGOV 75 bounds on lifetime depend on mass and improve as mass decreases. At 2 GeV the experiment is sensitive to $\tau > 3 \times 10^{-11}$ s for splnor, $\tau > 5 \times 10^{-11}$ s for scalar.

Highly Ionizing Particle Flux

VALUE (m ⁻² yr ⁻¹)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<0.4	95	0	KINOSHITA	81B PLAS	Z/β 30-100

Tachyon Flux In Cosmic Rays

See SMITH 77 for a review of earlier cosmic ray and accelerator experiments.

VALUE (cm ⁻² sr ⁻¹ s ⁻¹)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<2.4 × 10 ⁻⁹	90	0	⁷ MARINI	82 CNTR	v/c > 1.2
<2.3 × 10 ⁻¹⁰	95	0	⁸ BHAT	79 CNTR	
			⁹ SMITH	77 CNTR	
			¹⁰ PRESCOTT	76 CNTR	

⁷MARINI 82 is TOF measurement using PEP-counter at sea level.
⁸BHAT 79 is at Ootacamund (2200m above sea). No signal in 3621 hours.
⁹SMITH 77 analyzed more than 200000 showers (223 days) with E > 10¹⁴ eV scanning 290 × 10⁻⁶ s period before each shower. Observed excess 46 ± 40 events does not constitute statistically significant evidence.
¹⁰PRESCOTT 76 reanalyzed Clay and Crouch('C.C.') 74 data (Nature **248** 28 (1974)). Found apparatus effect, correction for which much reduces the statistical significance of positive 'C.C.' result. Also performed two new experiments one using 'C.C.' apparatus, another with new apparatus. Set upper limit at CL = 95% of about 30 tachyons per shower with average size N = 6 × 10⁵.

Tachyon Searches in e⁺e⁻ Annihilation

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
<1. × 10 ⁻⁶	90	0	¹¹ PERPELITSA 77	CNTR	$uv_{eq} < 1$
<1. × 10 ⁻⁵	90	0	¹¹ PERPELITSA 77	CNTR	$1 < uv_{eq} < 15$

¹¹PERPELITSA 77 is Michelson-type experiment for pair-produced tachyons in e⁺e⁻ annihilation (e⁺ from Cu isotope). Above limits are for $\sigma(e^+e^- \rightarrow \text{tachyon pair})/\sigma(e^+e^- \rightarrow 2\gamma)$ and uv_{eq} is tachyon velocities times earth equator component of velocity of preferred reference frame.

Searches for Tachyonic Decay (lower limit for mean life)

See LJUBICIC 75 figure 1 for review of earlier experiments.

VALUE (years)	DOCUMENT ID	TECN	COMMENT
>4.6 × 10 ¹³	¹² LJUBICIC	75 ELEC	$m_{\text{tachyon}} > 1.1$ keV

¹²LJUBICIC 75 used lead oxide cathode and electron multiplier looking for ionization due to tachyonic decay (spontaneous acquisition of energy) of bound-state e⁻. Sensitive to proper tachyon mass >1.1 keV. Above limit is obtained from observed e⁻ emission rate 3/hour.

Production of New Penetrating Non-ν Like States in Beam Dump

VALUE	DOCUMENT ID	TECN	COMMENT
>4.6 × 10 ¹³	¹³ LOSECCO	81 CALO	28 GeV protons

¹³No excess neutral-current events leads to $\sigma(\text{production}) \times \sigma(\text{interaction}) \times \text{acceptance} < 2.26 \times 10^{-71}$ cm⁴/nucleon² (CL = 90%) for light neutrals. Acceptance depends on models (0.1 to 4. × 10⁻⁴).

Branching Fraction of Z⁰ to a Pair of Stable Charged Heavy Fermions

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
<1 × 10 ⁻³	95	AKRAWY	90O OPAL	m = 29-40 GeV

Searches Full Listings

Other Stable Particle Searches

Heavy Particle Production Cross Section in e^+e^-

Ratio to $\sigma(e^+e^- \rightarrow \mu^+\mu^-)$. See also entries in Free Quark Search and Magnetic Monopole Searches.

VALUE	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
$< 2 \times 10^{-3}$	90	14	BUSKULIC	93C ALEP	$Q=1, m=32-72$ GeV
$< (10^{-2}-1)$	95	15	ADACHI	90C TOPZ	$Q=1, m=1-16, 18-27$ GeV
$< 7 \times 10^{-2}$	90	16	ADACHI	90E TOPZ	$Q=1, m=5-25$ GeV
$< 1.6 \times 10^{-2}$	95	0	17 KINOSHITA	82 PLAS	$Q=3-180, m < 14.5$ GeV
$< 5.0 \times 10^{-2}$	90	0	18 BARTEL	80 JADE	$Q=(3,4,5)/3-12$ GeV

¹⁴BUSKULIC 93C is a CERN-LEP experiment with $W_{cm} = mZ$. The limit is for a pair or single production of heavy particles with unusual ionization loss in TPC. See their Fig. 5 and Table 1.

¹⁵ADACHI 90C is a KEK-TRISTAN experiment with $W_{cm} = 52-60$ GeV. The limit is for pair production of a scalar or spin-1/2 particle. See Figs. 3 and 4.

¹⁶ADACHI 90E is KEK-TRISTAN experiment with $W_{cm} = 52-61.4$ GeV. The above limit is for inclusive production cross section normalized to $\sigma(e^+e^- \rightarrow \mu^+\mu^-) \cdot \beta(3-\beta^2)/2$, where $\beta = (1 - 4m^2/W_{cm}^2)^{1/2}$. See the paper for the assumption about the production mechanism.

¹⁷KINOSHITA 82 is SLAC PEP experiment at $W_{cm} = 29$ GeV using lexan and ³⁹Cr plastic sheets sensitive to highly ionizing particles.

¹⁸BARTEL 80 is DESY-PETRA experiment with $W_{cm} = 27-35$ GeV. Above limit is for inclusive pair production and ranges between $1. \times 10^{-1}$ and $1. \times 10^{-2}$ depending on mass and production momentum distributions. (See their figures 9, 10, 11).

Heavy Particle Production Cross Section in $p\bar{p}$

Limits are for a particle decaying to two hadronic jets.

Units(pb)	CL%	Mass(GeV)	DOCUMENT ID	TECN	COMMENT
< 2603	95	200	19 ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 4	95	400	19 ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets
< 7	95	600	19 ABE	93G CDF	1.8 TeV $p\bar{p} \rightarrow 2$ jets

¹⁹ABE 93G gives cross section times branching ratio into light (d, u, s, c, b) quarks for $\Gamma = 0.02 M$. Their Table II gives limits for $M = 200-900$ GeV and $\Gamma = (0.02-0.2) M$.

Heavy Particle Production Cross Section

VALUE (nb)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 0.05	95	20	ABE	92J CDF	$m=50-200$ GeV
$< 200-130$		21	CARROLL	78 SPEC	$m=2-2.5$ GeV
< 100	0	22	LEIPUNER	73 CNTR	$m=3-11$ GeV

²⁰ABE 92J look for pair production of unit-charged particles which leave detector before decaying. Limit shown here is for $m=50$ GeV. See their Fig. 5 for different charges and stronger limits for higher mass.

²¹CARROLL 78 look for neutral, $S = -2$ dihyperon resonance in $pp \rightarrow 2K^+X$. Cross section varies within above limits over mass range and $p_{lab} = 5.1-5.9$ GeV/c.

²²LEIPUNER 73 is an NAL 300 GeV p experiment. Would have detected particles with lifetime greater than 200 ns.

Heavy Particle Production Cross Section

VALUE (cm ² /N)	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$< (4-0.3) \times 10^{-31}$	95	23	AKESSON	91 CNTR	0	$m = 0-5$ GeV
$< 2.5 \times 10^{-35}$	0	24	GUSTAFSON	76 CNTR	0	$\tau > 10^{-7}$ s

²³AKESSON 91 limit is from weakly interacting neutral long-lived particles produced in pN reaction at 450 GeV/c performed at CERN SPS. Bourquin-Galliard formula is used as the production model. The above limit is for $\tau > 10^{-7}$ s. For $\tau > 10^{-9}$ s, $\sigma < 10^{-30}$ cm²/nucleon is obtained.

²⁴GUSTAFSON 76 is a 300 GeV FNAL experiment looking for heavy ($m > 2$ GeV) long-lived neutral hadrons in the M4 neutral beam. The above typical value is for $m = 3$ GeV and assumes an interaction cross section of 1 mb. Values as a function of mass and interaction cross section are given in figure 2.

Heavy Particle Production Differential Cross Section

VALUE (cm ² sr ⁻¹ GeV ⁻¹)	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$< 2.6 \times 10^{-36}$	90	0	25 BALDIN	76 CNTR	-	$Q=1, m=2.1-9.4$ GeV
$< 2.2 \times 10^{-33}$	90	0	26 ALBROW	75 SPEC	±	$Q=±1, m=4-15$ GeV
$< 1.1 \times 10^{-33}$	90	0	26 ALBROW	75 SPEC	±	$Q=±2, m=6-27$ GeV
$< 8. \times 10^{-35}$	90	0	27 JOVANOVOV...	75 CNTR	±	$m=15-26$ GeV
$< 1.5 \times 10^{-34}$	90	0	27 JOVANOVOV...	75 CNTR	±	$Q=±2, m=3-10$ GeV
$< 6. \times 10^{-35}$	90	0	27 JOVANOVOV...	75 CNTR	±	$Q=±2, m=10-26$ GeV
$< 1. \times 10^{-31}$	90	0	28 APPEL	74 CNTR	±	$m=3.2-7.2$ GeV
$< 5.8 \times 10^{-34}$	90	0	29 ALPER	73 SPEC	±	$m=1.5-24$ GeV
$< 1.2 \times 10^{-35}$	90	0	30 ANTIPOV	71B CNTR	-	$Q=-, m=2.2-2.8$ GeV
$< 2.4 \times 10^{-35}$	90	0	31 ANTIPOV	71C CNTR	-	$Q=-, m=1.2-1.7, 2.1-4$ GeV
$< 2.4 \times 10^{-35}$	90	0	BINON	69 CNTR	-	$Q=-, m=1-1.8$ GeV
$< 1.5 \times 10^{-36}$	0	32 DORFAN	65 CNTR			Be target $m=3-7$ GeV
$< 3.0 \times 10^{-36}$	0	32 DORFAN	65 CNTR			Fe target $m=3-7$ GeV

²⁵BALDIN 76 is a 70 GeV Serpukhov experiment. Value is per Al nucleus at $\theta = 0$. For other charges in range -0.5 to -3.0 , CL = 90% limit is $(2.6 \times 10^{-36})/|(\text{charge})|$ for mass range $(2.1-9.4 \text{ GeV}) \times |(\text{charge})|$. Assumes stable particle interacting with matter as do antiprotons.

²⁶ALBROW 75 is a CERN ISR experiment with $E_{cm} = 53$ GeV. $\theta = 40$ mr. See figure 5 for mass ranges up to 35 GeV.

²⁷JOVANOVOVICH 75 is a CERN ISR 26+26 and 15+15 GeV pp experiment. Figure 4 covers ranges $Q = 1/3$ to 2 and $m = 3$ to 26 GeV. Value is per GeV momentum.

²⁸APPEL 74 is NAL 300 GeV pW experiment. Studies forward production of heavy (up to 24 GeV) charged particles with momenta 24-200 GeV ($-$ charge) and 40-150 GeV ($+$ charge). Above typical value is for 75 GeV and is per GeV momentum per nucleon.

²⁹ALPER 73 is CERN ISR 26+26 GeV pp experiment. $p > 0.9$ GeV, $0.2 < \beta < 0.65$.

³⁰ANTIPOV 71B is from same 70 GeV p experiment as ANTIPOV 71C and BINON 69.

³¹ANTIPOV 71C limit inferred from flux ratio. 70 GeV p experiment.

³²DORFAN 65 is a 30 GeV/c p experiment at BNL. Units are per GeV momentum per nucleus.

Long-Lived Heavy Particle Invariant Cross Section

VALUE (cm ² /GeV ² /N)	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$< 5 \times 10^{-35}$	90	0	33 BERNSTEIN	88 CNTR		$Q=1, m=4-12$ GeV
$< 5 \times 10^{-37}$	90	0	33 BERNSTEIN	88 CNTR		$Q=1, m=4-12$ GeV
$< 2.5 \times 10^{-36}$	90	0	34 THRON	85 CNTR	-	$Q=1, m=1.87$ GeV
$< 1. \times 10^{-35}$	90	1	34 THRON	85 CNTR	+	$Q=1, m=1.5-3.0$ GeV
$< 6. \times 10^{-33}$	90	0	35 ARMITAGE	79 SPEC		$Q=(2/3, 1, 4/3, 2)$
$< 1.5 \times 10^{-33}$	90	0	35 ARMITAGE	79 SPEC		$m=4-10$ GeV
$< 1.1 \times 10^{-37}$	90	0	37 CUTTS	78 CNTR		$m=4.5-6$ GeV
$< 3.0 \times 10^{-37}$	90	0	38 VIDAL	78 CNTR		$m=4.5-6$ GeV

³³BERNSTEIN 88 limits apply at $x = 0.2$ and $p_T = 0$. Mass and lifetime dependence of limits are shown in the regions: $m = 1.5-7.5$ GeV and $\tau = 10^{-8}-2 \times 10^{-6}$ s. First number is for hadrons; second is for weakly interacting particles.

³⁴THRON 85 is FNAL 400 GeV proton experiment. Mass determined from measured velocity and momentum. Limits are for $\tau > 3 \times 10^{-9}$ s.

³⁵ARMITAGE 79 is CERN-ISR experiment at $E_{cm} = 53$ GeV. Value is for $x = 0.1$ and $p_T = 0.15$. Observed particles at $m = 1.87$ GeV are found all consistent with being antideuteron.

³⁶BOZZOLI 79 is CERN-SPS 200 GeV pN experiment. Looks for particle with τ larger than 10^{-8} s. See their figure 11-18 for production cross-section upper limits vs mass.

³⁷CUTTS 78 is pBe experiment at FNAL sensitive to particles of $\tau > 5 \times 10^{-8}$ s. Value is for $-0.3 < x < 0$ and $p_T = 0.175$.

³⁸VIDAL 78 is FNAL 400 GeV proton experiment. Value is for $x = 0$ and $p_T = 0$. Puts lifetime limit of $< 5 \times 10^{-8}$ s on particle in this mass range.

Long-Lived Heavy Particle Production

($\sigma(\text{Heavy Particle}) / \sigma(\pi)$)

VALUE	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
$< 10^{-8}$	39	NAKAMURA	89 SPEC	±	$Q = (-5/3, \pm 2)$
	0	40 BUSSIÈRE	80 CNTR	±	$Q = (2/3, 1, 4/3, 2)$

³⁹NAKAMURA 89 is KEK experiment with 12 GeV protons on Pt target. The limit applies for mass $\lesssim 1.6$ GeV and lifetime $\gtrsim 10^{-7}$ s.

⁴⁰BUSSIÈRE 80 is CERN-SPS experiment with 200-240 GeV protons on Be and Al target. See their figures 6 and 7 for cross-section ratio vs mass.

Production and Capture of Long-Lived Massive Particles

VALUE (10 ⁻³⁶ cm ²)	EVTS	DOCUMENT ID	TECN	COMMENT
< 20 to 800	0	41 ALEKSEEV	76 ELEC	$\tau=5$ ms to 1 day
< 200 to 2000	0	41 ALEKSEEV	76B ELEC	$\tau=100$ ms to 1 day
< 1.4 to 9	0	42 FRANKEL	75 CNTR	$\tau=50$ ms to 10 hours
< 0.1 to 9	0	43 FRANKEL	74 CNTR	$\tau=1$ to 1000 hours

⁴¹ALEKSEEV 76 and ALEKSEEV 76B are 61-70 GeV p Serpukhov experiment. Cross section is per Pb nucleus.

⁴²FRANKEL 75 is extension of FRANKEL 74.

⁴³FRANKEL 74 looks for particles produced in thick Al targets by 300-400 GeV/c protons.

See key on page 1343

Searches Full Listings

Other Stable Particle Searches

Heavy Particle Flux In Cosmic Rays

VALUE ($\text{cm}^{-2}\text{s}^{-1}\text{s}^{-1}$)	CL%	EVTS	DOCUMENT ID	TECN	CHG	COMMENT
< 1.8	$\times 10^{-12}$	90	44 ASTONE	93	CNTR	$m \geq 1.5 \times 10^{-13}$ gram
< 1.1	$\times 10^{-14}$	90	45 AHLEN	92	MCRO	$10^{-10} < m < 0.1$ gram
~ 6	$\times 10^{-9}$	2	46 SAITO	90		$Q \approx 14, m \approx 370m_p$
< 1.4	$\times 10^{-12}$	90	47 MINCER	85	CALO	$m \geq 1$ TeV
< 3.2	$\times 10^{-11}$	90	48 NAKAMURA	85	CNTR	$m > 1.5 \times 10^{-13}$ gram
< 1.7	$\times 10^{-11}$	99	49 SAKUYAMA	83B	PLAS	$m \sim 1$ TeV
< 1.	$\times 10^{-9}$	90	50 BHAT	82	CC	
< 3.5	$\times 10^{-11}$	90	51 MARINI	82	CNTR \pm	$Q = 1, m \sim 4.5m_p$
< 7.	$\times 10^{-11}$	90	52 ULLMAN	81	CNTR	Planck-mass 10^{19} GeV
2.	$\times 10^{-9}$	3	53 YOCK	81	SPRK \pm	$m = 1. \times 10^{-16}$ GeV or less
3.0	$\times 10^{-9}$	3	54 YOCK	80	SPRK	$Q = 1, m \sim 4.5m_p$
$(4 \pm 1) \times 10^{-11}$		3	55 BHAT	79	ELEC	Fractionally charged
< 1.3	$\times 10^{-9}$	90	56 YOCK	74	CNTR	$m \sim 4.5 m_p$
< 1.0	$\times 10^{-9}$	0	GOODMAN	79	ELEC	$m \geq 5$ GeV
< 7.	$\times 10^{-10}$	90	57 BHAT	78	CNTR \pm	$m > 1$ GeV
> 6.	$\times 10^{-9}$	5	58 BHAT	78	ELEC	$Q > 7e$ or $-7e$
< 3.0	$\times 10^{-8}$	0	59 YOCK	74	CNTR	$m > 6$ GeV
< 1.5	$\times 10^{-9}$	0	DARDO	72	CNTR	
< 3.0	$\times 10^{-10}$	0	TONWAR	72	CNTR	$m > 10$ GeV
< 5.0	$\times 10^{-11}$	90	BJORNBOE	68	CNTR	$m > 5$ GeV
		0	JONES	67	ELEC	$m = 5-15$ GeV

44 ASTONE 93 searched for quark matter ("nuclearites") in the velocity/c range = $10^{-3}-1$. Their Table 1 gives a compilation of searches for nuclearites.

45 AHLEN 92 searched for quark matter ("nuclearites"). The bound applies to velocity/c $< 2.5 \times 10^{-3}$. See their Fig. 3 for other velocity/c and heavier mass range.

46 SAITO 90 candidates carry about 450 MeV/nucleon. Cannot be accounted for by conventional backgrounds. Consistent with strange quark matter hypothesis.

47 MINCER 85 is high statistics study of calorimeter signals delayed by 20-200 ns. Calibration with AGS beam shows they can be accounted for by rare fluctuations in signals from low-energy hadrons in the shower. Claim that previous delayed signals including BJORNBOE 68, DARDO 72, BHAT 82, SAKUYAMA 83B below may be due to this fake effect.

48 NAKAMURA 85 at KEK searched for quark-matter. These might be lumps of strange quark matter with roughly equal numbers of u, d, s quarks. These lumps or nuclearites were assumed to have velocity/c of $10^{-4}-10^{-3}$.

49 SAKUYAMA 83B analyzed 6000 extended air shower events. Increase of delayed particles and change of lateral distribution above 10^{17} eV may indicate production of very heavy parent at top of atmosphere.

50 BHAT 82 observed 12 events with delay $> 2. \times 10^{-8}$ s and with more than 40 particles. 1 eV has good hadron shower. However all events are delayed in only one of two detectors in cloud chamber, and could not be due to strongly interacting massive particle.

51 MARINI 82 applied PEP-counter to TOF. Above limit is for velocity = 0.54 of light. Limit is inconsistent with YOCK 80 YOCK 81 events if isotropic dependence on zenith angle is assumed.

52 ULLMAN 81 is sensitive for heavy slow singly charge particle reaching earth with vertical velocity 100-350 km/s.

53 YOCK 81 saw another 3 events with $Q = \pm 1$ and m about $4.5m_p$ as well as 2 events with $m > 5.3m_p, Q = \pm 0.75 \pm 0.05$ and $m > 2.8m_p, Q = \pm 0.70 \pm 0.05$ and 1 event with $m = (9.3 \pm 3.)m_p, Q = \pm 0.89 \pm 0.06$ as possible heavy candidates.

54 YOCK 80 events are with charge exactly or approximately equal to unity.

55 BHAT 78 is at Kolar gold fields. Limit is for $\tau > 10^{-6}$ s.

56 YOCK 74 events could be tritons.

Concentration of Heavy (Charge +1) Stable Particles in Matter

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 4	$\times 10^{-17}$	95	57 YAMAGATA	93 SPEC Deep sea water, $m = 5-1600m_p$
< 6	$\times 10^{-15}$	95	58 VERKERK	92 SPEC Water, $m = 10^5$ to 3×10^7 GeV
< 7	$\times 10^{-15}$	95	59 VERKERK	92 SPEC Water, $m = 10^4, 6 \times 10^7$ GeV
< 9	$\times 10^{-15}$	95	58 VERKERK	92 SPEC Water, $m = 10^8$ GeV
< 3	$\times 10^{-23}$	90	59 HEMMICK	90 SPEC Water, $m = 1000m_p$
< 2	$\times 10^{-21}$	90	59 HEMMICK	90 SPEC Water, $m = 5000m_p$
< 3	$\times 10^{-20}$	90	59 HEMMICK	90 SPEC Water, $m = 10000m_p$
< 1.	$\times 10^{-29}$		SMITH	82B SPEC Water, $m = 30-400m_p$
< 2.	$\times 10^{-28}$		SMITH	82B SPEC Water, $m = 12-1000m_p$
< 1.	$\times 10^{-14}$		SMITH	82B SPEC Water, $m > 1000 m_p$
< (0.2-1.)	$\times 10^{-21}$		SMITH	79 SPEC Water, $m = 6-350 m_p$

57 YAMAGATA 93 used deep sea water at 4000 m since the concentration is enhanced in deep sea due to gravity.

58 VERKERK 92 looked for heavy isotopes in sea water and put a bound on concentration of stable charged massive particle in sea water. The above bound can be translated into into a bound on charged dark matter particle (5×10^6 GeV), assuming the local density, $\rho = 0.3$ GeV/cm³, and the mean velocity (v) = 300 km/s.

59 See HEMMICK 90 Fig. 7 for other masses 100-10000 m_p .

Concentration of Heavy (Charge -1) Stable Particles

VALUE	CL%	DOCUMENT ID	TECN	COMMENT
< 4	$\times 10^{-20}$	90	60 HEMMICK	90 SPEC C, $M = 100m_p$
< 8	$\times 10^{-20}$	90	60 HEMMICK	90 SPEC C, $M = 1000m_p$
< 2	$\times 10^{-16}$	90	60 HEMMICK	90 SPEC C, $M = 10000m_p$
< 6	$\times 10^{-13}$	90	60 HEMMICK	90 SPEC Li, $M = 1000m_p$
< 1	$\times 10^{-11}$	90	60 HEMMICK	90 SPEC Be, $M = 1000m_p$
< 6	$\times 10^{-14}$	90	60 HEMMICK	90 SPEC B, $M = 1000m_p$
< 4	$\times 10^{-17}$	90	60 HEMMICK	90 SPEC O, $M = 1000m_p$
< 4	$\times 10^{-15}$	90	60 HEMMICK	90 SPEC F, $M = 1000m_p$
< 1.5	$\times 10^{-13}$ /nucleon	68	61 NORMAN	89 SPEC $^{206}\text{PbX}^-$
< 1.2	$\times 10^{-12}$ /nucleon	68	61 NORMAN	87 SPEC $^{56,58}\text{FeX}^-$

60 See HEMMICK 90 Fig. 7 for other masses 100-10000 m_p .

61 Bound valid up to $m_{X^-} \sim 100$ TeV.

Long-Lived Particle Search at Hadron Collisions

Limits are for cross section times branching ratio.

VALUE (pb/nucleon)	CL%	EVTS	DOCUMENT ID	TECN	COMMENT
< 2	90	0	62 BADIER	86	BDMP $\tau = (0.05-1.) \times 10^{-8}$ s

62 BADIER 86 looked for long-lived particles at 300 GeV π^+ beam dump. The limit applies for nonstrongly interaction neutral or charged particles with mass > 2 GeV. The limit applies for particle modes, $\mu^+\pi^-, \mu^+\mu^-, \pi^+\pi^-, X, \pi^+\pi^-\pi^+$ etc. See their figure 5 for the contours of limits in the mass- τ plane for each mode.

REFERENCES FOR Other Stable Particle Searches

ABE	93G	PRL 71 2542	+Albrow, Akimoto, Amidei, Anway-Wiese+	(CDF Collab.)
ASTONE	93	PR D47 4770	+Bassan, Bonifazi, Coccia+(ROMA, ROMA, CATA, FRAS)	
BUSKULIC	93C	PL B303 198	+Decamp, Goy, Lees, Minard+	(ALEPH Collab.)
YAMAGATA	93	PR D47 1231	+Takamori, Utsunomiya	(KONAN)
ABE	92J	PR D46 R1889	+Amidei, Anway-Wiese+	(CDF Collab.)
AHLEN	92	PRL 69 1860	+Ambrosio, Antolini, Auremma, Baker+	(MACRO Collab.)
VERKERK	92	PRL 68 1116	+Grynberg, Pichard, Spiro, Zylberajch+(ENSP, SACL, PAST)	
AKESSON	91	ZPHY C52 219	+Almeida, Angelis, Atherton, Aubry+	(HELIOS Collab.)
ADACHI	90C	PL B244 352	+Aihara, Doser, Enomoto+	(TOPAZ Collab.)
ADACHI	90E	PL B249 336	+Anazawa, Doser, Enomoto, Fujii+	(TOPAZ Collab.)
AKRAWAY	90	PL B252 290	+Alexander, Allison, Allport, Anderson+	(OPAL Collab.)
HEMMICK	90	PR D41 2074	+Elmore+ (ROCH, MICH, OHIO, RAL, LANL, STON)	
SAITO	90	PRL 65 2094	+Hatano, Fukuda, Oda	(ICRR, KOBE)
NAKAMURA	89	PR D39 1261	+Kobayashi, Konaka, Imai, Masaki+	(KYOT, TMTC)
NORMAN	89	PR D39 2499	+Chadwick, Lesko, Larimer, Hoffman	(LBL)
BERNSTEIN	88	PR D37 3103	+Shea, Winstein, Cousins, Greenhaigh+	(STAN, WISC)
REN	88	PR D38 1417	+Huo, Lu, Su+ (China-Japan Collab., Mt. Fuji Collab.)	
BORISOV	87	PL B190 226	+Cherdntseva+	(Pamir-Chacaltaya Collab.)
NORMAN	87	PRL 58 1403	+Gazes, Bennett	(LBL)
ALNER	86	PL B180 415	+Ansgore, Asman, Booth, Burow+	(UAS Collab.)
BADIER	86	ZPHY C31 21	+Bemporad, Boucrot, Calot+	(NA3 Collab.)
MINCER	85	PR D32 541	+Freudenreich, Goodman+	(UMD, GMAS, NSF)
NAKAMURA	85	PL 161B 417	+Horie, Takahashi, Tanimori	(KEK, INUS)
THRON	85	PR D31 451	+Cardello, Cooper, Teig+	(YALE, FNAL, IOWA)
ARNISON	83B	PL 122B 189	+Astbury, Aubert, Bacchi+	(UA1 Collab.)
SAKUYAMA	83B	LNC 37 17	+Muzuki	(MEIS)
Also	83	LNC 36 389	Sakuyama, Watanabe	(MEIS)
Also	83D	NC 78A 147	Sakuyama, Watanabe	(MEIS)
Also	83C	NC 6C 371	Sakuyama, Watanabe	(MEIS)
ALPGARD	82	PL 115B 71	+Ansgore, Asman, Berglund+	(UAS Collab.)
BHAT	82	PR D25 2820	+Gupta, Murthy, Sreekantan+	(TATA)
KINOSHITA	82	PRL 48 77	+Price, Fryberger	(UCB, SLAC)
MARINI	82	PR D26 1777	+Peruzzi, Piccolo+	(FRAS, LBL, NWES, STAN, HAWA)
SMITH	82B	NP B206 333	+Bennett, Homer, Lewin, Walford, Smith	(RAL)
BAYBURINA	81	NP B191 1	+Borisov+	(LEBD, MOSU, INRM, GEOR, TAJK+)
KINOSHITA	81B	PR D24 1707	+Price	(UCB)
LOSECCO	81	PL 102B 209	+Sulak, Galik, Horstkotter+	(MICH, PENN, BNL)
ULLMAN	81	PRL 47 289		(LEHM, BNL)
YOCK	81	PR D23 1207		(AUCK)
BARTEL	80	ZPHY C6 295	+Canzler, Lords, Drumm+	(JADE Collab.)
BUSSIERE	80	NP B174 1	+Giacomelli, Lesgouy+	(BGNA, SACL, LAPP)
LATES	80	PRP 16 151	+Fujimoto, Hasegawa	(CAMP, WASH)
YOCK	80	PR D22 61		(AUCK)
ARMITAGE	79	NP B150 87	+Benz, Bobbink+ (CERN, DARE, FOM, MCHS, UTRE)	
BHAT	79	JPG 5 L13	+Gopalakrishnan, Gupta, Tonwar	(TATA)
BOZZOLI	79	NP B159 363	+Bussiere, Giacomelli+ (BGNA, LAPP, SACL, CERN)	
GOODMAN	79	PR D19 2572	+Ellsworth, Ito, Macfall, Siohan+	(UMD)
SMITH	79	NP B149 525	+Bennett	(RHEL)
BHAT	78	Pramana 10 115	+Murthy	(TATA)
CARROLL	78	PRL 41 777	+Chiang, Johnson, Kyica, Ki+	(BNL, PRIN)
CUTTS	78	PRL 41 363	+Dulude+ (BROW, FNAL, ILL, BARI, MIT, WARS)	
VIDAL	78	PL 77B 344	+Herb, Lederman+	(COLU, FNAL, STON, UCB)
VIETTEL	78	LNC 22 235	+Hahn, Schacher	(BERN)
PEREPELITSA	77	PL 67B 471		(ITEP)
SMITH	77	CJP 55 1280	+Standil	(MANI)
ALEKSEEV	76	SJNP 22 531	+Zaitsev, Kalinina, Kruglov+	(JINR)
		Translated from YAF 22 1021.		
ALEKSEEV	76B	SJNP 23 633	+Zaitsev, Kalinina, Kruglov+	(JINR)
		Translated from YAF 23 1190.		
BALDIN	76	SJNP 22 264	+Vertogradov, Vshivsky, Grishkevich+	(JINR)
		Translated from YAF 22 512.		
BRIATORE	76	NC 31A 553	+Dardo, Pizzoli, Mannocchi+	(LCGT, FRAS, FREIB)
GUSTAFSON	76	PRL 37 474	+Ayre, Jones, Longo, Murthy	(MICH)
PRESCOTT	76	JPG 2 261		(ADLD)
ALBROW	75	NP B97 189	+Barber+ (CERN, DARE, FOM, LANC, MCHS, UTRE)	
BLAGOV	75	SJNP 21 158	+Komar, Murashova, Syreishchikova+	(LEBD)
		Translated from YAF 21 300.		

Searches Full Listings

Other Stable Particle Searches

FRANKEL	75	PR D12 2561	+Frati, Resvanis, Yang, Nezrick	(PENN, FNAL)
JOVANOV...	75	PL 56B 105	Jovanovich+	(MANI, AACH, CERN, GENO, HARV+)
LJUBICIC	75	PR D11 696	+Pavlovic, Pisk, Logan	(BOSK, OTTA)
YOCK	75	NP B86 216		(AUCK, SLAC)
APPEL	74	PRL 32 428	+Bourquin, Gaines, Lederman+	(COLU, FNAL)
CLAY	74	NAT 248 28	+Crouch	(ADLD)
FRANKEL	74	PR D9 1932	+Frati, Resvanis, Yang, Nezrick	(PENN, FNAL)
YOCK	74	NP B76 175		(AUCK)
ALPER	73	PL 46B 265	+ (CERN, LIVP, LUND, BOHR, RHEL, STOH, BERG+)	
LEIPUNER	73	PRL 31 1226	+Larsen, Sessoms, Smith, Williams+	(BNL, YALE)
DARDO	72	NC 9A 319	+Navarra, Penengo, Sitte	(TORI)
TONWAR	72	JPA 5 569	+Narayan, Sreekantan	(TATA)
ANTIPOV	71B	NP B31 235	+Denisov, Donskov, Gorin, Kachanov+	(SERP)
ANTIPOV	71C	PL 34B 164	+Denisov, Donskov, Gorin, Kachanov+	(SERP)
BINON	69	PL 30B 510	+Dutell, Kachanov, Khromov, Kutyin+	(SERP)
BJORNBOE	68	NC B53 241	+Damgard, Hansen+	(BOHR, TATA, BERN, BERG)
JONES	67	PR 164 1584	(MICH, WISC, LBL, UCLA, MINN, COSU, COLO+)	
DORFAN	65	PRL 14 999	+Eades, Lederman, Lee, Ting	(COLU)
COWARD	63	PR 131 1782	+Gittelman, Lynch, Ritson	(STAN)
BELOUSOV	60	JETP 11 1143	+Rusakov, Tamm, Cerenkov	(LEBD)
GORBUNOV	60	Translated from ZETF 38 1589.		
		JETP 11 51	+Spirdonov, Cerenkov	(LEBD)
		Translated from ZETF 39 69.		

OTHER COMPILATIONS OF INTEREST

OTHER COMPILATIONS OF INTEREST

1986 Adjustment of the Fundamental Physical Constants

E.R. Cohen and B.N. Taylor
Rev. Mod. Phys. **59**, 1121 (1987)

A Compilation of Structure Functions in Deep-Inelastic Scattering (1985–1991)

R.G. Roberts and M.R. Whalley
J. Phys. **G17**, D1 (1991)

Compilation of Coupling Constants and Low-Energy Parameters

O. Dumbrajs, R. Koch, H. Pilkuhn, G.C. Oades,
H. Behrens, J.J. de Swart, P. Kroll
Nucl. Phys. **B216**, 277 (1983)

Electroweak Interactions: Experimental Facts and Theoretical Foundation

D. Haidt and H. Pietschmann (ed. H. Schopper)
Landolt-Börnstein, New Series Vol. **I/10** (1988)

A Compilation of Data on e^+ and e^- Interactions

O.P. Yushchenko, V.V. Ezhela, V. Flaminio,
D.R.O. Morrison, Yu.G. Stroganov, M.R. Whalley
to be issued as a CERN report and as a book in
the Landolt-Börnstein series (Spring '92)

Compilation of Data on the Energy-Energy Correlation and its Asymmetry in e^+e^- Annihilation

W.J. Stirling and M.R. Whalley
RAL Report RAL-87-107 (1987)

Compilation of Data on $\gamma\gamma \rightarrow$ Hadrons

R.G. Roberts and M.R. Whalley
RAL Report RAL-86-058 (1986)

Compilation of Data on Single Prompt Photon Production in Hadron-Hadron Interactions

P. Aurenche and M.R. Whalley
RAL Report RAL-89-106 (1989)

Total Cross Sections for Reactions of High Energy Particles

A. Baldini, V. Flaminio, W.G. Moorhead,
D.R.O. Morrison (ed. H. Schopper)
Landolt-Börnstein, New Series Vol. **I/12 a** and
I/12 b (1988)

Pion Nucleon Scattering: 1) Tables of Data, 2) Methods and Results of Phenomenological Analyses

G. Höhler (ed. H. Schopper)
Landolt-Börnstein, New Series Vol. **I/9 b1** (1982)
and **I/9 b2** (1983)

Compilation of Nucleon-Nucleon and Nucleon-Antinucleon Elastic Scattering Data

M.K. Carter, P.D.B. Collins, and M.R. Whalley
RAL Report RAL-86-002 (1986)

Scattering of Elementary Particles: NN and KN

J. Bystricky, P. Carlson, C. Lechanoine, F. Lehar,
F. Mönig, K.R. Schubert (ed. H. Schopper)
Landolt-Börnstein, New Series Vol. **I/9 a** (1980)

Compilation of Cross Sections IV: γ , ν , Λ , Σ , Ξ , and K_L^0 Induced Reactions

S.I. Alekhin, A. Baldini, P. Capiluppi, *et al.*,
CERN-HERA and COMPAS Groups
CERN-HERA Report 87-01 (1987)

A Guide to Data in Experimental Elementary Particle Physics Literature

S.I. Alekhin, V.V. Ezhela, *et al.*, COMPAS and Berkeley
Particle Data Groups
Lawrence Berkeley Laboratory Report LBL-90
revised (1993)

Current Experiments in Elementary Particle Physics

H. Galić, F.E. Armstrong, *et al.*,
Lawrence Berkeley Laboratory Report LBL-91,
revised (July 1994)

Major Detectors in Elementary Particle Physics

G. Gidal, B. Armstrong, A. Rittenberg
Lawrence Berkeley Laboratory Report LBL-91,
supplement, revised (1985)

Table of Isotopes, 7th Edition

C.M. Lederer, V.S. Shirley, E. Browne, J.M. Dairiki,
R.E. Doebler, A.A. Shihab-Eldin, L.J. Jardine,
J.K. Tuli, A.B. Buyrn
John Wiley & Sons, New York (1978)

Table of Radioactive Isotopes

E. Browne and R.B. Firestone (Editor: V.S. Shirley)
John Wiley & Sons, New York (1986)

Astronomical Almanac

(US Government Printing Office, Washington, and Her
Majesty's Printing Office, London)(annual)

Chart and Software of the Standard Model of Fundamental Particles and Interactions

Contemporary Physics Education Project
(available from:)
Science Kit
777 East Park Drive
Tonawanda, NY 14150 USA

Chart of the Nuclides

F.W. Walker, J.R. Parrington, and F. Feiner
General Electric Co., Nuclear Energy Operations,
175 Curtner Av., M/C 397,
San Jose, CA 95125 USA
(14th edition, 1989)

INDEX

$A(1680)$ or [now called $\pi_2(1670)$]	1199, 1495	Nucleon resonances (Δ resonances)	1220, 1710
$A(2100)$ [now called $\pi_2(2100)$]	1511	Nucleon resonances (N resonances)	1219, 1688
$a_0(980)$ [was $\delta(980)$]	1197, 1465	Nucleons	1218, 1673
$a_0(980)$, note on	1465	Ω baryons	1225, 1780
$a_1(1260)$ [was $A_1(1270)$ or A_1]	1197, 1471	Ξ baryons	1224, 1769
$a_1(1260)$, note on	1471	Baryonium candidates	1519
$a_2(1320)$ [was $A_2(1320)$]	1198, 1480	Baryon number conservation	1228
$a_3(2050)$ [was $A(2050)$]	1510	Baryon resonances, SU(3) classification of	1321
A_3 [now called $\pi_2(1670)$]	1199, 1495	Baryons in quark model	1321
$a_4(2040)$ [was $\delta_4(2040)$]	1509	Baryons, stable	1218, 1673
$a_6(2450)$ [was $\delta_6(2450)$]	1516	(see individual entries for p , n , Λ , Σ , Ξ , Ω , Λ_c , Ξ_c , Ω_c , and Λ_b)	
Abbreviations used in Full Listings	1344	Bayesian approach	1280
Accelerator parameters (colliders)	1246	Bays-Durham Monte Carlo algorithm	1283
Acceptance-rejection method in Monte Carlo	1283	$B\bar{B}$ mixing	1316, 1318, 1632
Accessing the high-energy physics databases	1188	BDMS DBMS	1189
Activity, unit of, for radioactivity	1268	Beam momentum, c.m. energy and momentum vs.	1290
Age of the universe	1234	Beauty – see Bottom and B , B^* , B_s , B_s^*	
Algorithms for Monte Carlo	1284	Becquerel, unit of radioactivity	1268
ALICE DBMS	1188	BEPC (China) collider parameters	1247
α_s , QCD coupling constant	1233, 1297	β -rays, from radioactive sources	1270
Amplitudes, Lorentz invariant	1290	Bethe-Bloch equation	1251
Amu (atomic mass unit)	1233	Bias, definition of	1275
Argand diagram, definition	1293	Big-bang cosmology	1236
Argon, dE/dx resolution	1264	Big-bang nucleosynthesis	1238
Astronomical unit	1234	Binomial distribution, Monte Carlo algorithm for	1284
Astrophysics	1236, 1238	Binomial distribution, relations for	1272
Asymmetry formulae in Standard Model	1305	Biological damage from radiation	1268
Atmospheric pressure	1233	BITNET address for comments	1177
Atomic and nuclear properties of materials	1241	Bivariate Gaussian	1272
Atomic mass unit	1233	Bohr magneton	1233
Attenuation length for photons	1259	Bohr radius	1233
Attenuation, photon and electron	1259	Boltzmann constant	1233
Authors and consultants	1177	Booklet, Particle Physics, how to get	1177
Average hadron multiplicities in e^+e^- annihilation events	1331	Bosons	1191, 1351
Averaging data, relations for	1275	Bottom baryon (Λ_b^0)	1226, 1790
Averaging of data	1180	Bottom (B) meson production and decay, note on	1601
Avogadro number	1233	Bottom (B) semileptonic decays, note on	1602
Axial vector couplings, g_V , g_A vector	1304	Bottom, B^0 - \bar{B}^0 mixing, CP violation in B decay, note on	1632
Axion searches	1192, 1377	Bottom-changing neutral currents, tests for	1228
Axion searches, note on	1377	Bottom mesons (B , B^* , B_s , B_s^*)	1207, 1601
Axions as dark matter	1238	Bottom mesons, note on highlights	1601
b quark lifetime and CKM matrix	1316	Bottomonium system, level diagram	1661
$b_1(1235)$ [was $B(1235)$]	1197, 1470	Bounded physical region, Bayesian approach	1280
b' quark (4^{th} generation), searches for,	1195, 1442	Bounded physical region, frequentist approach	1280
B (bottom meson)	1207, 1601	Breit-Wigner distribution, Monte Carlo algorithm for	1284
B decay, CP violation in	1318	Breit-Wigner probability density function	1274
B meson production and decay, note on	1601	Breit-Wigner resonance, definition	1293
B semileptonic decays, note on	1602	Bulletin board preprints	1189
B^\pm (bottom meson)	1207, 1609	C (charge conjugation), tests of conservation	1228
B^0 , \bar{B}^0 (bottom meson)	1209, 1623	c (quark)	1195, 1439
B^0 - \bar{B}^0 mixing, CP violation in B decay, note on	1632	Cabibbo angle	1315
B_H and B_L in B^0 - \bar{B}^0 mixing, note on	1632	Cabibbo-Kobayashi-Maskawa mixing	1315
B_s^0	1210, 1639	Cabibbo-Kobayashi-Maskawa mixing in B decay, note on	1602, 1632
B^*	1210, 1639	Calorimeters	1263
B_s^*	1641	Capacitance, formulas for	1286
Baryon decay parameters, note on	1681	Cascade baryons (Ξ baryons)	1224, 1769
Baryon magnetic moments, note on	1729	Centrauro searches	1811
Baryons	1218, 1673	Central limit theorem	1272
Charmed baryons	1225, 1782	Čerenkov detectors	1261
Dibaryons		Čerenkov radiation	1257
(see p. VIII.118 in our 1992 edition, Phys. Rev. D45, Part II)		CERN databases	1186
Exotic resonances (Z^* resonances)		CESR (Cornell) collider parameters	1246
(see p. VIII.58 in our 1992 edition, Phys. Rev. D45, Part II)		CESR+ (Cornell) collider parameters	1246
Hyperon baryons (Λ baryons)	1221, 1728	Change of random variables, relations for	1271
Hyperon baryons (Σ baryons)	1222, 1745	Charge conjugation (C) conservation	1228
		Charge conjugation of $q\bar{q}$ states	1319

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

Charge conservation	1228	Critical energy	1254
Charge conservation and the Pauli exclusion principle, note on (see p. VI.10 in our 1992 edition, Phys. Rev. D45)		Cross sections and related quantities, plots of	1327
Chargino searches	1801	e^+e^- , νN , $\bar{\nu}N$, Ap , γp , γd , $\pi^\pm p$, $\pi^\pm d$, $K^\pm p$, $K^\pm d$, pp , pn , pd , $\bar{p}p$, $\bar{p}n$, and $\bar{p}d$ cross sections	
Charm-changing neutral currents, tests for	1228	e^+e^- annihilation cross section near M_Z	1333
Charmed baryons (Λ_c^+ , Σ_c , Ξ_c , Ω_c^0)	1225 , 1783	Fragmentation functions	1332
Charmed, nonstrange mesons (D , D^* , D_J)	1203 , 1572	Jet production	1330
Charmed, strange mesons [D_s , D_s^* , D_{sJ}]	1206 , 1596	Nucleon structure functions	1327
Charmonium system, level diagram	1642	Pseudorapidity distributions	1330
χ^2 confidence level vs. χ^2 for n degrees of freedom	1273	Cross sections, hadronic, high-energy parametrizations	1335
χ^2 distribution, Monte Carlo algorithm for	1284	Cross sections, relations for	1291, 1294
χ^2 distribution, relations for	1273	Cumulative distribution function, definition	1271
$\chi_{b0}(1P) = \chi_{b0}(9860)$	1213 , 1664	Curie, unit of radioactivity	1268
$\chi_{b0}(2P) = \chi_{b0}(10235)$	1213 , 1666	d (quark)	1195 , 1437
$\chi_{b1}(1P) = \chi_{b1}(9890)$	1213 , 1664	d functions	1287
$\chi_{b1}(2P) = \chi_{b1}(10255)$	1213 , 1667	D^\pm	1203 , 1572
$\chi_{b2}(1P) = \chi_{b2}(9915)$	1213 , 1665	D^\pm , D^0 branching fractions, note on	1564
$\chi_{b2}(2P) = \chi_{b2}(10270)$	1213 , 1667	D semileptonic decays, note on	1565
$\chi_{c0}(1P) = \chi_{c0}(3415)$	1212 , 1652	D^0 , \bar{D}^0	1204 , 1581
$\chi_{c1}(1P) = \chi_{c1}(3510)$	1212 , 1653	$D_1(2420)^0$	1206 , 1594
$\chi_{c2}(1P) = \chi_{c2}(3555)$	1212 , 1654	$D(1530)$ [<i>now called</i> $f_1(1510)$]	1198 , 1490
CKM mixing elements in B decay, note on	1602, 1632	$D^*(2007)^0$	1206 , 1593
CKM (Cabibbo-Kobayashi-Maskawa) mixing	1315	$D^*(2010)^+$	1206 , 1593
Clebsch-Gordan coefficients	1287	$D_2^*(2460)$	1206 , 1595
c.m. energy and momentum vs. beam momentum	1290	$D_J(2440)^\pm$	1595
Collider parameters	1246	D_s^\mp [<i>was</i> F^\pm]	1206 , 1596
Color octet leptons	1227 , 1810	$D_s^{*\pm}$ [<i>was</i> $F^{*\pm}$]	1207 , 1599
Color octet quarks	1227 , 1810	$D_{s1}(2536)^\pm$	1207 , 1600
Compensating calorimeters	1263	$D_{sJ}(2573)^\pm$	1600
Compilations, particle physics	1815	Dalitz plot, relations for	1291
Compositeness, quark and lepton, searches	1227 , 1805	Damage, biological, from radiation	1268
Compositeness, quark and lepton, searches, note on	1805	DAΦNE (Frascati) collider parameters	1249
Compton scattering for N and Δ resonances, photoproduction and (review)	1686	Dark matter	1238
Compton wavelength, electron	1233	Dark matter limits:	
Conditional probability density function	1271	Astrophysical limits on neutrino mass	1420
Confidence coefficient, definition of	1278	Invisible axion mass limits	1382
Confidence interval, definition of	1275, 1278	Neutralinos mass limits	1800
Confidence intervals, normal distribution	1278	Sneutrino mass limits	1801
Confidence intervals using Student's t	1279	Data, averaging and fitting procedures	1180
Confidence level, definition	1278	Data, selection and treatment	1179
Conservation laws	1228	Databases, availability	1186
Constrained fits, procedures for	1181	Databases, high-energy physics	1185
Consultants	1178	Databases, particle physics	1185
Conversion probability for photons to e^+e^-	1259	Day, sidereal	1234
Correlation coefficient, definition	1271	dE/dx	1251
Cosmic background radiation (CBR) temperature	1234	Decay amplitudes (for hyperon decays) (see p. 286 in our 1982 edition, Phys. Lett. 111B)	
Cosmic ray background in counters	1268	Decay constants of charged pseudoscalar mesons, note on	1443
Cosmic ray fluxes	1269	Decays, kinematics and phase space for	1290
Cosmological background radiation	1236	Deep-inelastic scattering	1298
Cosmological constant	1234	Definitions for abbreviations used in Full Listings	1344
Cosmology	1236 , 1238	δ , CKM angle for CP violation	1315
Coupling constant in QCD	1233, 1297	δ -rays	1253
Couplings for photon, W , Z	1304	$\delta(980)$ [<i>now called</i> $a_0(980)$]	1197 , 1465
Coulomb scattering through small angles, multiple	1253	$\delta_4(2040)$ [<i>now called</i> $a_4(2040)$]	1509
Covariance, definition	1271	$\delta_6(2450)$ [<i>now called</i> $a_6(2450)$]	1516
CP , tests of conservation	1228	Δ resonances (see also N and Δ resonances)	1220 , 1710
CP violation, overview	1318	$\Delta B = 1$, weak neutral currents, tests for	1228
CP violation and CKM matrix	1315, 1318, 1632	$\Delta B = 2$, tests for	1228
CP violation in B decay	1318	$\Delta C = 1$, weak neutral currents, tests for	1228
CP violation and B^0 - \bar{B}^0 mixing in B decay, note on	1632	$\Delta C = 2$, tests for	1228
CP violation in $K_S^0 \rightarrow 3\pi$ decays, note on	1536	$\Delta I = 1/2$ rule for hyperon decays, test of (see p. 286 in our 1982 edition, Phys. Lett. 111B)	
CP violation in K_L^0 decay	1318	$\Delta S = 1$, weak neutral currents, tests for	1228
CP -violation parameters in K_L^0 decays, note on	1543	$\Delta S = 2$, tests for	1228
CPT , tests of conservation	1228	$\Delta S = \Delta Q$ rule in K^0 decay, note on	1547
Critical density	1234		

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

$\Delta S = \Delta Q$, tests of	1228	ϵ_0 (permittivity of free space)	1233, 1285
Density effect in energy loss rate	1252	$\epsilon_1, \epsilon_2, \epsilon_3$ electroweak variables	1313
Density of materials, table	1241	Equivalent photon approximation	1295
Density of matter, critical	1234	Error ellipse for multivariate Gaussian	1279
Density of matter, local	1234	Error estimates in least-squares fitting	1277
Density parameter of the universe, Ω_0	1234	Error estimates in likelihood fitting	1276
DESY databases	1189	Error function	1272
Detector parameters	1261	Error procedure for masses and widths of meson resonances	1550
Deuteron mass	1233	Error propagation, relations for	1282
Deuteron structure function	1328	Errors, treatment of	1180
Dibaryons		Established nonets for the mesons	1320
(see p. VIII.118 in our 1992 edition, Phys. Rev. D45, Part II)		Estimator, definition of	1275
Dielectric constant of gaseous elements, table	1242	η meson	1196, 1450
Differential Čerenkov detectors	1262	η decay parameters, note on	1454
Disk density	1234	$\Gamma(\eta \rightarrow \gamma\gamma)$, note on	1451
Distributions, probability, definition	1271	$\eta(1295)$	1198, 1478
DORIS (DESY) collider parameters	1246	$\eta(1440)$ [was $\iota(1440)$]	1198, 1487
Dose, radioactivity, unit of absorbed	1268	$\eta(1440)$, note on	1487
Drift and proportional chamber potentials	1263	$\eta(1700)$ [now called $X(1700)$]	1504
Durham databases	1186	$\eta(1760)$	1506
e (natural log base), value of	1233	$\eta(2225)$	1513
e (electron)	1193, 1396	$\eta_2(1870)$	1508
Charge conservation and the Pauli exclusion principle, note on		$\eta'(958)$	1197, 1461
(see p. VI.10 in our 1992 edition, Phys. Rev. D45)		$\eta_c(1S) = \eta_c(2980)$	1210, 1642
e^+e^- annihilation, cross-section formulae	1295	$\eta_c(2S) = \eta_c(3590)$	1656
e^+e^- annihilation cross section near M_Z	1333	Excited lepton searches	1227, 1808
e^+e^- average multiplicity, plot of	1333	Exotic baryons (Z^* resonances)	
e^+e^- average multiplicity, table of	1331	(see p. VIII.58 in our 1992 edition, Phys. Rev. D45, Part II)	
$e^+e^-(1100-2200)$	1517	Exotic meson resonances	1670
$e^+e^- R$ function, plot of	1334	Expectation value, definition	1271
e^+e^- two-photon process, cross-section formula	1295	Expectation value, relations for	1271
$E(1420)$ [now called $f_1(1420)$]	1198, 1484	Exponential distribution	1274
Earth equatorial radius	1234	Exposure, radioactivity, unit of	1268
Efficiency of statistical estimator, definition	1275	$f_{D^+}, f_{D^+}, f_\eta, f_{\eta'}, f_{K^+}, f_{\pi^+}, f_{\pi^0}$ decay constants	1443
Electrical resistivity of elements, table	1242	F_1, F_2, F_3 structure functions	1294, 1327
Electric charge (Q) conservation	1228	F^\pm [now called D_s^\pm]	1206, 1596
Electromagnetic calorimeters	1263	$F^{*\pm}$ [now called $D_s^{*\pm}$]	1207, 1599
Electromagnetic relations	1285	$f_0(980)$ [was $S(975)$ or S^*]	1197, 1464
Electromagnetic shower detectors, energy resolution	1263	$f_0(1300)$ [was $\epsilon(1200)$]	1198, 1478
Electromagnetic showers, lateral distribution	1256	$f_0(1370)$	1483
Electromagnetic showers, longitudinal distribution	1255	$f_0(1525)$	1493
Electron	1193, 1396	$f_0(1590)$	1199, 1493
Electron charge	1233, 1285	$f_0(1285)$	1198, 1475
Electron cyclotron frequency/field	1233	$f_1(1420)$ [was $E(1420)$]	1198, 1484
Electron mass	1233, 1193	$f_1(1420)$, note on	1484
Electron radius, classical	1233	$f_1(1510)$ [was $D(1530)$]	1198, 1490
Electron volt	1233	$f_2(1270)$	1197, 1472
Electronic structure of the elements	1244	$f_2(1430)$	1486
Electroproduction structure functions, relations for	1294	$f_2(1520)$	1490
Electroweak analyses of new physics	1312	$f_2(1640)$	1494
Electroweak interactions, Standard Model of	1304	$f_2(1810)$	1506
Elements, electronic structure of	1244	$f_2(2010)$ [was $g_T(2010)$]	1199, 1509
Elements, periodic table of	1243	$f_2(2150)$ [was $\epsilon(2150)$]	1511
Energy and momentum (c.m.) vs. beam momentum	1290	$f_2(2300)$ [was $g_T(2300)$]	1200, 1514
Energy loss and range in liquid hydrogen	23	$f_2(2340)$ [was $g_T^h(2340)$]	1200, 1515
Energy loss and range in Pb, Cu, Al, and C	23	$f_2'(1525)$ [was $f'(1525)$]	1198, 1491
Energy loss (fractional) for electrons and positrons in lead	1260	$f_4(2050)$ [was $h(2030)$]	1200, 1510
Energy loss rate for charged particles	1251	$f_4(2220)$ [was $\xi(2220)$]	1513
Energy loss rate in compounds	1252	$f_4(2300)$ [was $\epsilon(2300)$]	1514
Energy loss rate, restricted	1253	$f_6(2510)$ [was $r(2510)$]	1516
Energy loss rate for muons at high energies	1256	$f_J(1710)$ [was $\theta(1690)$]	1199, 1504
Energy loss rates for heavy charged projectiles	23	$f_J(1710)$, note on	1504
$\epsilon(1200)$ [now called $f_0(1300)$]	1198, 1478	Familon searches	1382
$\epsilon(2150)$ [now called $f_2(2150)$]	1511	Fermi coupling constant	1233
$\epsilon(2300)$ [now called $f_4(2300)$]	1514	Fermi plateau	1252
ϵ (permittivity)	1233, 1242, 1285	Feynman's x variable	1290

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

Field equations, electromagnetic	1285	HEPNET address for comments	1177
Fine structure constant	1233	HERA (DESY) collider parameters	1250
Fits to data	1180	Higgs boson in Standard Model	1304, 1365
Fitting data, relations for	1275, 1276	Higgs boson mass in electroweak analyses	1312
Flavor-changing neutral currents, tests for	1228	Higgs, M_H , constraints on	1312
Forbidden states in quark model	1323	Higgs searches	1192 , 1365
Force, Lorentz	1285	Higgs searches, note on	1365
Form factors, $K_{\ell 3}$, note on	1530	High-energy physics databases	1185
Form factors, $\pi \rightarrow \ell\nu\gamma$ and $K \rightarrow \ell\nu\gamma$, note on	1447	History of measurements, discussion	1182
Fourth generation searches	1195 , 1442	Hubble parameter	1234
Fractional energy loss for electrons and positrons in lead	1260	Hyperon baryons (see Λ and Σ baryons)	1221 , 1728
Fragmentation functions, plot of	1332	Hyperon decays, nonleptonic decay amplitudes	
Fragmentation functions, relations for	1295	(see p. 286 in our 1982 edition, Phys. Lett. 111B)	
Fragmentation, heavy quark	1333	Hyperon decays, test of $\Delta I = 1/2$ rule for	
Free quark searches	1227 , 1791	(see p. 286 in our 1982 edition, Phys. Lett. 111B)	
Friedmann equation	1236	ID particle codes for Monte Carlo	1325
Friedman-Robertson-Walker metric	1236	Ideal mixing in quark model	1320
Full Listings, key to reading	1343	Ideograms, criteria for presentation	1181
Full Listings, organization of	1177	IHEP databases	1186
g (gluon)	1191 , 1351	Illustrative key to the Full Listings	1343
$g(1690)$ [now called $\rho_3(1690)$]	1199 , 1497	Impedance, relations for	1286
$g_T(2010)$ [now called $f_2(2010)$]	1199 , 1509	Importance sampling in Monte Carlo calculations	1283
$g_T^-(2300)$ [now called $f_2(2300)$]	1200 , 1514	Inclusive distributions, one-particle, relations for	1296
$g_T^+(2340)$ [now called $f_2(2340)$]	1200 , 1515	Inclusive hadronic reactions	1295
g_V, g_A vector, axial vector couplings	1304	Inclusive reactions, kinematics for	1292
γ (Euler constant), value of	1233	Inconsistent data, treatment of	1181
γ (photon)	1191 , 1351	Independence of random variables	1271
γp and γd cross sections, plots of	1341	Inductance, relations for	1286
gamma production in $p\bar{p}$ interactions	1330	Inorganic scintillator parameters	1261
γ -rays, from radioactive sources	1270	Inorganic scintillators	1261
Gamma distribution, Monte Carlo algorithm for	1284	Interactive <i>Review of Particle Properties</i> (RPP) system	1185
Gamma distribution, relations for	1274	International System (SI) units	1240
Gauge bosons	1191 , 1351	INTERNET address for comments	1177
(see individual entries for γ, W, Z, g , and graviton)		Introduction	1177
Gauge couplings	1304	Inverse transform method in Monte Carlo	1283
Gaussian distribution, Monte Carlo algorithm for	1284	Invisible A^0 (Axion) searches	1382
Gaussian distribution, multivariate	1272	Invisible A^0 mass limits, note on	1382
Gaussian distribution, relations for	1272	Ionization energies of the elements	1244
Gaussian distribution, upper limits	1278	Ionization energy loss at minimum, table	1241
Gaussian ellipsoid	1273	Ionization yields for charged particles	1253
Gell-Mann-Okubo formula	1319	$\iota(1440)$ [now called $\eta(1440)$]	1198 , 1487
Glueino searches	1227 , 1803	Jansky	1234
gluon, g	1191 , 1351	Jet production in pp and $p\bar{p}$ interactions, plot of	1330
Gluonium candidates	1670	$J/\psi(1S) = J/\psi(3097)$ or $\psi(1S)$	1211 , 1644
Goldstone boson searches	1382	$K \rightarrow 3\pi$ Dalitz plot parameters, note on	1528
Gravitational acceleration g	1233	$K \rightarrow \ell\nu\gamma$ form factors, note on	1447
Gravitational constant G_N	1233, 1234	$K_{\ell 3}$ form factors, note on	1530
graviton	1351	K^\pm	1200 , 1521
Gray, unit of absorbed dose of radiation	1268	K^\pm mass, note on	1521
$h(2030)$ [now called $f_4(2050)$]	1200 , 1510	K^+p, K^+n , and K^+d cross sections, plots of	1340
$h_1(1170)$ [was $H(1190)$]	1197 , 1469	K^-p, K^-n , and K^-d cross sections, plots of	1339
$h_1(1380)$	1483	K^0, \bar{K}^0	1200 , 1534
$h_c(1P)$	1654	K^0 decay, note on $\Delta S = \Delta Q$ rule in	1547
Hadronic calorimeters	1263	K_L^0	1201 , 1537
Hadronic cross-sections, high-energy parametrizations	1335	K_L^0 decays, note on CP -violation parameters in	1543
Hadronic flavor conservation	1228	K_L^0 decay, CP violation in	1318
Hadronic shower detectors	1263	K_S^0	1201 , 1534
Half-lives of commonly used radioactive nuclides	1270	$K_S^0 \rightarrow 3\pi$ decay, note on CP violation in	1536
Halo density	1234	$K(1460)$ [was $K(1400)$]	1556
Heavy boson searches	1192 , 1372	$K(1830)$	1560
Heavy lepton searches	1194 , 1418	$K(3100)$	1563
Heavy lepton searches, note on	1418	$K^*(892)$	1201 , 1549
Heavy particle searches	1812	$K^*(892)$ mass and mass differences, note on	1550
Heavy physics from precision experiments	1312	$K^*(1410)$ [was $K^*(1410)$]	1202 , 1553
Heavy quark fragmentation	1333	$K^*(1680)$ [was $K^*(1790)$]	1202 , 1557
Heavy quark searches	1641		

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

$K_0^*(1430)$ [<i>was</i> $\kappa(1350)$]	1202, 1554	Light particle searches	1811
$K_0^*(1950)$	1560	Light, speed of	1233
$K_1(1270)$ [<i>was</i> $Q(1280)$ or Q_1]	1201, 1551	Light year	1234
$K_1(1400)$ [<i>was</i> $Q(1400)$ or Q_2]	1201, 1552	Likelihood condition	1275
$K_1(1650)$	1557	Likelihood function	1275
$K_2(1580)$ [<i>was</i> $L(1580)$]	1557	Limits (statistical) in presence of bounded physical region	1280
$K_2(1770)$ [<i>was</i> $L(1770)$]	1202, 1558	Linear least-squares fitting	1276
$K_2(1770)$ and the $K_2(1820)$, note on	1558	Liquid ionization chambers, free electron drift velocity	1264
$K_2(1820)$ [<i>was</i>]	1202, 1560	Listings, Full, keys to reading	1343
$K_2(2250)$ [<i>was</i> $K(2250)$]	1562	Lorentz force	1285
$K_2^*(1430)$ [<i>was</i> $K^*(1430)$]	1202, 1554	Lorentz invariant amplitudes	1290
$K_2^*(1980)$	1561	Lorentz transformations of four-vectors	1290
$K_3(2320)$ [<i>was</i> $K(2320)$]	1562	Luminosity conversion	1234
$K_3^*(1780)$ [<i>was</i> $K^*(1780)$]	1202, 1559	MACHOS—see Dark matter limits	
$K_4(2500)$ [<i>was</i> $K(2500)$]	1562	MACHOS	1238
$K_4^*(2045)$ [<i>was</i> $K^*(2060)$]	1202, 1561	Magnetic moments, baryon, note on	1729
$K_5^*(2380)$	1562	Magnetic monopole searches	1227, 1793
K_{23} form factors, note on	1530	Magnetic monopole searches, note on	1793
Kaon (see K)	1200, 1521	Majoron searches	1382
$\kappa(1350)$ [<i>now called</i> $K_0^*(1430)$]	1202, 1554	Mandelstam variables	1292
KEKB collider parameters	1248	Marginal probability density function	1271
Key to the Full Listings	1343	Mass attenuation coefficient for photons, defined	1259
Kinematics, decays, and scattering	1290	Massive neutrinos and lepton mixing, search for	1194, 1421
Knock-on electrons, energetic	1253	Materials, atomic and nuclear properties of	1241
Kobayashi-Maskawa (Cabibbo-) mixing matrix	1315	Matter, passage of particles through	1251
$L(1580)$ [<i>now called</i> $K_2(1580)$]	1557	Maximum likelihood	1275
$L(1770)$ [<i>now called</i> $K_2(1770)$]	1202, 1558	Maxwell equations	1285
Lagrangian, QCD	1297	Mean range and energy loss in liquid hydrogen	23
Lagrangian, standard electroweak	1304	Mean range and energy loss in Pb, Cu, Al, and C	23
Λ	1221, 1728	Median, definition	1271
Λ and Σ baryons	1221, 1728	Median, variance of	1271
$\Lambda(1405)$, note on	1732	Meson multiplets in quark model	1319
Listings, Λ baryons	1728	Meson nonets (established)	1320
Listings, Σ baryons	1745	Mesons	1196, 1443
Status of (review)	1731	$b\bar{b}$ mesons	1213, 1661
A_p cross section, plot of	1341	Bottom mesons	1207, 1601
Λ , QCD parameter	1297	Charmed, nonstrange mesons	1210, 1642
Λ_b^0	1226, 1790	Charmed, strange mesons	1206, 1596
Λ_c^+	1225, 1783	Exotic mesons	1670
$\Lambda_c(2625)^+$	1225, 1786	Nonstrange mesons	1196, 1443
Landau distribution	1252	Strange mesons	1200, 1521
Lattice QCD	1301	Mesons, stable	1196, 1443
LBL databases	1186	(see individual entries for π , η , K , D , D_s , and B)	
Least-squares fitting, linear	1276	\overline{MS} renormalization scheme (QCD)	1297
Least-squares fitting, method of	1276	\overline{MS} renormalization scheme (Standard Model)	1304
Least-squares fitting, nonlinear	1277	Metric prefixes, commonly used	1240
Least-squares fitting, straight line	1277	Michel parameter ρ	1193, 1415
Lee-Sugawara relation	287†	Microwave background	1236
LEP (CERN) collider parameters	1247	Minimal subtraction scheme in QCD	1297
Lepton conservation, tests of	1228	Minimum ionization	1251
Lepton family number conservation	1228	Minimum ionization loss, table	1241
Lepton (heavy) searches	1194, 1418	MIP (minimum ionizing particle)	1251
Lepton mixing, neutrinos (massive) and, search for	1194, 1421	Mixing angle, weak ($\sin^2 \theta_W$)	1233, 1304
Lepton, quark compositeness searches	1227, 1805	Mixing, $B^0-\bar{B}^0$, note on	1632
Lepton, quark substructure searches	1227, 1805	Mixing, quark model, ideal	1320
Leptons	1193, 1385	Mixing, singlet-octet in quark model	1320
(see individual entries for ν_e , e , ν_μ , μ , ν_τ , and τ)		Molar volume	1233
Leptons, weak interactions of quarks and	1304	Molière radius	1255
Leptoproduction cross sections, relations for	1294	Momenta, measurement of, in a magnetic field	1264
Leptoproduction kinematics	1294	Momentum — c.m. energy and momentum	
Leptoquark searches	1375	vs. beam momentum	1290
Lethal dose from penetrating ionizing radiation	1268	Momentum transfer, minimum and maximum	1290
LHC (CERN) collider parameters	1250	Monopole searches	1227, 1793
Light boson searches	1377	Monopole searches, note on	1793
Light neutrino types, number of	1194, 1416	Monte Carlo particle numbering scheme	1325
Light neutrino types from colliders, number of, note on	1416	Monte Carlo techniques	1283

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

μ (muon)	1193, 1397	Nuclear magneton	1233
$\mu \rightarrow e$ conversion	1398	Nuclear total cross section, table	1241
μ_0 (permeability of free space)	1233, 1285	Nucleon resonances (see N and Δ resonances)	1219, 1684
Multibody decay kinematics	1291	Nucleon structure functions, plots of	1327
Multiple Coulomb scattering through small angles	1253	Nuclides, radioactive, commonly used	1270
Multiplets, meson in quark model	1319	Numbering scheme for particles in Monte Carlos	1325
Multiplets, SU(n)	1289	Occupational radiation dose, U.S. maximum permissible	1268
Multiplicities, average in e^+e^- interactions, table of	1331	Octet-singlet mixing in quark model	1320
Multiplicity, average in e^+e^- interactions, plot of	1333	Omega baryons (Ω baryons)	1225, 1780
Multiplicity, average in pp and $\bar{p}p$ interactions, plot of	1333	Ω^-	1225, 1780
Multivariate Gaussian	1272	Ω_c^0 [<i>was</i> T^0]	1789
Muon	1193, 1397	Ω_c , critical density	1234
Muon decay parameters, note on	1399	Ω_0 , density parameter	1234
Muon energy loss rate at high energies	1256	Ω^- resonances	1781
M_W	1233, 1309	$\omega(782)$	1196, 1459
M_Z	1233, 1309	$\omega(1420)$	1198, 1486
n (neutron)	1218, 1680	$\omega(1600)$	1199, 1494
N and Δ resonances	1219, 1684	$\omega_3(1670)$	1199, 1495
Listings, Δ resonances	1710	One-particle inclusive distributions, relations for	1296
Listings, N resonances	1684	Optical theorem	1293
Photoproduction and Compton scattering (review)	1686	Organization of Full Listings and Summary Tables	1177
$\pi N \rightarrow N\pi\pi$ channel (review)	1686	Other stable particle searches	1811
Status of (review)	1684	Other stable particle searches, note on	1811
Two-body partial-wave analyses (review)	1685	P (parity), tests of conservation	1228
N^* resonances (see N and Δ resonances)	1219, 1684	p (proton)	1218, 1673
$\bar{N}N(1100-3600)$	1517	p mean life, note on	1673
n -body differential cross sections	1291	pp average multiplicity, plot of	1333
n -body phase space	1290	pp jet production	1330
$n - \bar{n}$ oscillations	1681	pp , pn , and pd cross sections, plots of	1336, 1337
Names, hadrons	1179, 1323	$\bar{p}p$ average multiplicity, plot of	1333
Neutral-current parameters, standard model expressions for	1310	$p\bar{p}$ gamma production	1330
Neutral-current parameters, values for	1308	$\bar{p}p$ jet production	1330
Neutralino searches	1800	$\bar{p}p$ pseudorapidity	1330
Neutralinos as dark matter	1238	$\bar{p}p$, $\bar{p}n$, and $\bar{p}d$ cross sections, plots of	1336, 1337
Neutrino (see ν)	1193, 1385	Parity of $q\bar{q}$ states	1319
Neutrino bounds from astrophysics and cosmology	1432	Parsec	1234
Neutrino mass limits, note on	1432	Partial-wave analyses for N and Δ resonances (review)	1685
Neutrino oscillation experiments, note on	1424	Partial-wave expansion of scattering amplitude	1293
Neutrino oscillation searches	1194, 1421	Particle detectors	1261
Neutrino production structure functions, relations for	1294	Particle ID numbers for Monte Carlos	1325
Neutrino, solar, experiments	1426	Particle nomenclature	1179, 1323
Neutrino types (light), number of	1194, 1416	Particle Physics Booklet, how to get	1177
Neutrino types (light) from colliders, number of, note on	1416	Passage of particles through matter	1251
Neutrinoless double beta decay, search for	1430	Pauli exclusion principle, charge conservation, note on (see p. VI.10 in our 1992 edition, Phys. Rev. D45)	
Neutrinos as dark matter	1238	PEP (SLAC) collider parameters	1246
Neutrinos (massive) and lepton mixing, search for	1194, 1421	PEP-II (SLAC) collider parameters	1248
Neutrinos, note on	1385	Periodic table of the elements	1243
Neutron	1218, 1680	Permeability μ_0 of free space	1233, 1285
Neutrons, from radioactive sources	1270	Permittivity ϵ_0 of free space	1233, 1285
New physics from electroweak analyses	1312	PETRA (DESY) collider parameters	1246
Newtonian gravitational constant G_N	1233, 1234	Phase space, Lorentz invariant	1290
Newton-Raphson method	1277	Phase space, relations for	1290
Nomenclature for hadrons	1179, 1323	$\phi(1020)$	1197, 1467
Nonets, meson (established)	1320	$\phi(1680)$	1199, 1497
Non- $q\bar{q}$ candidates	1670	$\phi_3(1850)$ [<i>was</i> $X(1850)$]	1199, 1507
Normal distribution, confidence intervals for	1278	Φ -factory (UCLA) collider parameters	1249
Normal distribution, relations for	1272	Φ -factory (Novosibirsk) collider parameters	1249
Normal equation in least-squares fitting	1277	Photon searches	1799
ν_e	1193, 1389	Photon	1191, 1351
ν_μ	1193, 1392	Photon and electron attenuation	1259
ν_μ mass, note on	1392	Photon attenuation length	1259
ν_τ	1193, 1395	Photon attenuation length (high energy)	1259
νN and $\bar{\nu}N$ cross sections, plot of	1327	Photon collection efficiency, scintillators	1261
Nuclear collision length, table	1241	Photon coupling	1304
Nuclear inelastic cross section, table	1241	Photon cross section in carbon and lead, contributions to	1260
Nuclear interaction length, table	1241		

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

Photon pair-production cross section	1255	Quark searches, note on	1791
Photon to e^+e^- conversion probability	1259	Quarks	1195 , 1433
Photoproduction and Compton scattering for N and Δ resonances (review)	1686	Quarks and leptons, weak interactions of	1294, 1304
Physical constants, table of	1233	Quarks, current masses of	1433, 1304
π , value of	1233	Quarks, properties of	1319
π^\pm	1196 , 1444	R function, e^+e^- scattering, plot of	1334
π^\pm mass, note on	1444	$r(2510)$ [<i>now called</i> $f_6(2510)$]	1516
$\pi^\pm p$ and $\pi^\pm d$ cross sections, plots of	1338	Rad, unit of absorbed dose of radiation	1268
π^0	1196 , 1448	Radiation, biological damage from chronic exposure	1268
$\pi \rightarrow \ell\nu\gamma$ form factors, note on	1447	Radiation, Čerenkov	1257
$\pi(1300)$	1198 , 1480	Radiation length	1254
$\pi(1770)$	1506	Radiation length, approximate algorithm	1254
$\pi_2(1670)$ [<i>was</i> $A(1680)$ or A_3]	1199 , 1495	Radiation length of materials, table	1241
$\pi_2(2100)$ [<i>was</i> $A(2100)$]	1511	Radiation, lethal dose from	1268
$\pi N \rightarrow N\pi\pi$ channel (review)	1686	Radiation levels in detectors	1266
Pion	1196 , 1444	Radiation, long-term risk	1268
Planck constant	1233	Radiation weighting factor	1268
Planck mass	1234	Radiative corrections in Standard Model	1304
Plasma energy	1251	Radioactive sources, commonly used	1270
Plastic scintillators	1261	Radioactivity and radiation protection	1268
Poisson distribution, Monte Carlo algorithm for	1284	Radioactivity, natural annual background	1268
Poisson distribution, relations for	1272	Radioactivity, unit of absorbed dose	1268
Poisson distribution, upper limits for	1279	Radioactivity, unit of activity	1268
Poisson processes with background, upper limits for	1281	Radioactivity, unit of exposure	1268
Potentials, electromagnetic	1285	Radon, component natural annual background radioactivity	1268
PPDS	1189	Random angle, Monte Carlo algorithm for sine and cosine of	1284
Precision experiments, heavy physics	1312	Range (mean) and energy loss in liquid hydrogen	23
Prefixes, metric, commonly used	1240	Range (mean) and energy loss in Pb, Cu, Al, and C	23
Preprints from bulletin boards	1189	Range, scaling law for projectile mass and charge	23
Probability	1271	Rao-Cramér-Frechet bound	1275
χ^2 confidence level vs. χ^2 for n degrees of freedom	1273	Rapidity	1290
Probability density function, definition	1271	Redshift	1236
Propagation of errors	1282	Refractive index of materials, table	1241
Properties (atomic and nuclear) of materials	1241	Relativistic kinematics	1290
Proportional and drift chamber potentials	1263	Relativistic transformation of electromagnetic fields	1285
Proportional chamber wire instability	1263	Rem, roentgen equivalent for man	1268
Proton (see p)	1218 , 1673	Renormalization in Standard Model	1304
Proton cyclotron frequency/field	1233	Renormalization schemes in QCD	1297
Proton mass	1218 , 1233	Representations, $SU(n)$	1289
Proton structure function	1327	Residuals, definition of	1277
Pseudorapidity η , defined	1292	Resistivity, electrical, of elements, table	1242
Pseudorapidity distribution in $\bar{p}p$ interactions, plot of	1330	Resistivity of metals	1286
Pseudoscalar mesons, decay constants of charged, note on	1443	Resistivity, relations for	1286
$\psi(1S) = J/\psi(1S) = J/\psi(3097)$	1211 , 1644	Resonance, Breit-Wigner form and Argand plot for	1293
$\psi(2S) = \psi(3685)$	1212 , 1656	Resonances (see Mesons and Baryons)	
$\psi(3770)$	1212 , 1659	Restricted energy loss rate, charged projectiles	1253
$\psi(4040)$	1212 , 1659	ρ parameter in electroweak analyses	1312
$\psi(4160)$	1213 , 1660	ρ parameter in electroweak analyses (Standard Model)	1309
$\psi(4415)$	1213 , 1660	ρ parameter of electroweak interactions	1310
$Q(1280)$ or Q_1 [<i>now called</i> $K_1(1270)$]	1201 , 1551	$\rho(770)$	1196 , 1456
$Q(1400)$ or Q_2 [<i>now called</i> $K_1(1400)$]	1201 , 1552	$\rho(770)$, note on	1456
QCD	1297	$\hat{\rho}(1405)$	1484
QCD parton model	1294	$\rho(1450)$	1198 , 1489
QSPIRES	1188	$\rho(1450)$ and $\rho(1770)$, note on	1501
Quality factor for biological damage due to radiation	1268	$\rho(1700)$	1199 , 1501
Quantum numbers in quark model	1319	$\rho(2150)$	1512
Quark and lepton compositeness searches	1227 , 1805	$\rho(2210)$	1512
Quark and lepton substructure searches	1227 , 1805	$\rho_3(1690)$ [<i>was</i> $g(1690)$]	1199 , 1497
Quark fragmentation in e^+e^- annihilation	1333	$\rho_3(2250)$	1514
Quark mass, note on	1433	$\rho_5(2350)$	1515
Quark model	1319	Ring-Imaging Čerenkov detectors	1262
Quark model assignments	1319	Robertson-Walker metric	1236
Quark model, dynamical ingredients	1322	Robustness of statistical estimator, definition	1275
Quark parton model	1294	Roentgen, measure of X or γ radiation intensity	1268
Quark searches, free	1227 , 1791	Rydberg energy	1233
		s (quark)	1195 , 1437

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

S, T, U electroweak variables	1313	Z' searches, note on	1372
$S = +1$ baryons (Z^* baryons)		Selection and treatment of data	1179
(see p. VIII.58 in our 1992 edition, Phys. Rev. D45 , Part II)		Semileptonic decays (B), note on	1602
$S(975)$ or S^* [<i>now called</i> $f_0(980)$]	1197 , 1464	Semileptonic decays (D), note on	1565
S -matrix for two-body scattering	1290	Shower detector energy resolution	1263
S -wave $\pi\pi$, $K\bar{K}$, and $\eta\eta$ interactions, note on	1478	Showers, electromagnetic, lateral distribution of	1256
Scale factor, definition of	1180	Showers, electromagnetic, longitudinal distribution of	1255
Scattering, relations for	1294	SI units, complete set	1240
Schwarzschild radius of sun	1234	Sidereal day	1234
Scintillator parameters	1261	Sidereal year	1234
Sea-level cosmic ray fluxes	1269	Sievert, unit of radiation dose equivalent	1268
Searches:		Σ baryons (see also Λ and Ξ baryons)	1222 , 1745
Axion searches	1192 , 1377	Σ^+	1222 , 1745
Baryonium candidates	1519	Σ^0	1222 , 1747
Centauro searches	1811	Σ^-	1222 , 1748
Chargino searches	1801	$\Sigma(1670)$, note on	1756
Color octet leptons	1227 , 1810	$\Sigma^- \rightarrow \Lambda e^- \nu$, note on	1749
Color octet quarks	1227 , 1810	$\Sigma_c(2455)$	1225 , 1787
Compositeness, quark and lepton, searches	1227 , 1805	$\Sigma_c(2530)$	1787
Excited lepton searches	1227 , 1808	Silicon strip detectors	1262
Familon searches	1382	$\sin^2\theta_W$, weak mixing angle	1233, 1304
Fourth generation (b') searches	1195 , 1442	Singlet-octet mixing in quark model	1320
Free quark searches	1227 , 1791	SLAC databases	1186
Gluino searches	1227 , 1803	SLC (SLAC) collider parameters	1247
Gluonium candidates	1670	Slepton searches	1227 , 1801
Goldstone boson searches	1382	SN 1987A, note on constraints on particles	1432
Heavy boson searches	1192 , 1372	Solar equatorial radius	1234
Heavy lepton searches	1194 , 1418	Solar luminosity	1234
Heavy particle searches	1812	Solar mass	1234
Higgs searches	1192 , 1365	Solar ν experiments	1426
Invisible A^0 (Axion) searches	1382	Solar radius in galaxy	1234
Lepton (heavy) searches	1194 , 1418	Solar velocity in galaxy	1234
Lepton mixing, neutrinos (massive) and, search for	1194 , 1421	Sources, radioactive, commonly used	1270
Lepton, quark compositeness searches	1227 , 1805	SPEAR (SLAC) collider parameters	1246
Lepton, quark substructure searches	1227 , 1805	Specific heats of elements, table	1242
Leptoquark searches	1375	Spherical harmonics	1287
Light boson searches	1192 , 1377	SPIRES DBMS	1188
Light neutrino types, number of	1194 , 1416	$Spp\bar{S}$ (CERN) collider parameters	1250
Light particle searches	1811	Squark searches	1227 , 1802
Magnetic monopole searches	1227 , 1793	SSC collider parameters	1250
Majoron searches	1382	Standard error, definition of	1278
Massive neutrinos and lepton mixing, searches	1194 , 1421	Standard Model of electroweak interactions	1304
Monopole searches	1227 , 1793	Standard particle numbering for Monte Carlos	1325
Neutralino searches	1800	Statistic, definition of	1275
Neutrino bounds from astrophysics and cosmology	1432	Statistical procedures	1180
Neutrino oscillation searches	1194 , 1421	Statistics	1275
Neutrino, solar, experiments	1426	Stefan-Boltzmann constant	1233
Neutrino types, number of	1194 , 1416	Stopping power	1251
Neutrinoless double beta decay searches	1430	Stopping power for heavy charged projectiles	1251
Neutrinos (massive) and lepton mixing, search for	1194 , 1421	Straight-line fit, relations for	1277
Non- $q\bar{q}$ candidates	1670	Strange baryons	1221 , 1728
Other stable particle searches	1811	Strange mesons	1200 , 1521
Photino searches	1799	Strangeness-changing neutral currents, tests for	1228
Quark and lepton compositeness searches	1227 , 1805	Strong coupling constant in QCD	1233, 1297
Quark and lepton substructure searches	1227 , 1805	Structure functions	1327
Quark searches, free	1227 , 1791	Structure functions, electroproduction, relations for	1294
Slepton searches	1227 , 1801	Structure functions in quark parton model	1294
Squark searches	1227 , 1802	Structure functions, leptonproduction, relations for	1294
Solar ν experiments	1426	Student's t distribution, Monte Carlo algorithm for	1284
Substructure, quark and lepton, searches	1227 , 1805	Student's t distribution, relations for	1274
Supersymmetric partner searches	1227 , 1795	SU(2) \times U(1)	1304
Tachyon searches	1811	SU(3) classification of baryon resonances	1321
Technipion searches	1192 , 1370	SU(3) isoscalar factors	1288
Top searches	1195 , 1439	SU(3) multiplets	1321
Vector meson candidates	1517	SU(3) representation matrices	1288
Weak gauge boson searches	1192 , 1372	SU(6) multiplets	1321

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

SU(n) multiplets	1289	$\Upsilon(11020)$	1214 , 1670
Substructure, quark and lepton, searches	1227 , 1805	$V_{ud}, V_{us}, V_{ub}, V_{cd}, V_{cs}, V_{cb}, V_{td}, V_{ts}, V_{tb}$	1315
Substructure, quark and lepton, searches, note on	1805	Variance, definition	1271
Summary Tables, organization of	1177	Variance, relations for	1271
Superconducting solenoids	1265	Vector meson candidates	1517
Supernova SN 1987A, note on constraints on particles	1432	VEPP-2m (Novosibirsk) collider parameters	1249
Supersymmetric partner searches	1227 , 1795	VEPP-4m (Novosibirsk) collider parameters	1247
Supersymmetry, electroweak analyses of	1312	VLEPP, INP (Serpukhov) collider parameters	1248
Superweak model of CP violation	1543, 1632	W gauge boson	1191 , 1351
Survival probability, relations for	1290	W gauge boson, mass, width, branching ratios, and coupling to fermions	1191 , 1233, 1304, 1308, 1309
Synchrotron radiation	1286	Weak gauge boson searches	1192 , 1372
Systematic errors, treatment of	1180	Weak interactions of quarks and leptons	1304, 1315
t quark, Searches for	1195 , 1439	Weak mixing angle ($\sin^2 \theta_W$)	1233, 1304
T (time reversal), tests of conservation	1228	Weak neutral currents, tests for ($\Delta B = 1, \Delta C = 1, \Delta S = 1$)	1228
Tachyon searches	1811	Weighted averaging, relations for	1275
τ lepton	1193 , 1403	Weinberg angle ($\sin^2 \theta_W$)	1233, 1304
τ -decay data problem, note on	1403	Width determinations of Υ states, note on	1661
τ -decay branching fractions, note on	1406	Width of W and Z bosons	1308
Technicolor, electroweak analyses of	1313	Wien displacement law constant	1233
Technipion searches	1192 , 1370	WIMPs—see Dark matter limits	
TEVATRON (Fermilab) collider parameters	1250	WIMPs	1238
Thermal conductivity of elements, table	1242	Wolfenstein parameterization	1316
Thermal expansion coefficients of elements, table	1242	World-Wide Web	1188
$\theta(1690)$ [<i>now called</i> $f_J(1710)$]	1199 , 1504	x variable (of Feynman's)	1290
θ_W , weak mixing angle	1233, 1304	$X(1600)$	1494
Thomson cross section	1233	$X(1700)$ [<i>was</i> $\eta(1700)$]	1504
Three-body decay kinematics	1291	$X(1740)$	1505
Three-body phase space	1290	$X(1775)$	1506
Threshold Čerenkov detectors	1261	$X(1830)$	1507
TMAE	1262	$X(1850)$ [<i>now called</i> $\phi_3(1850)$]	1199 , 1507
Top quark mass from electroweak analyses	1312	$X(1900-3600)$	1519
Top quark mass from the Standard Model	1308	$X(1910)$	1508
Top searches	1195 , 1439	$X(1950)$	1509
top, m_t , constraints on	1312	$X(2200)$	1512
Total lepton number conservation	1228	$X(3250)$	1516
Transformation of electromagnetic fields, relativistic	1285	Ξ baryons	1224 , 1769
Transition radiation	1257	Ξ resonances, note on	1773
Transition radiation detectors (TRD)	1262	Ξ^0	1224 , 1769
TRISTAN (KEK) collider parameters	1247	Ξ^-	1224 , 1771
Tropical year	1234	Ξ^+ [<i>was</i> A^+]	1226 , 1788
Two-body decay kinematics	1290	Ξ_c^0	1226 , 1788
Two-body differential cross sections	1290	$\xi(2220)$ [<i>now called</i> $f_4(2220)$]	1513
Two-body partial decay rate	1290	Y^* resonances (see Λ and Σ resonances)	1221 , 1732
Two-body scattering kinematics	1290	Year, sidereal	1234
Two-photon processes in e^+e^- annihilation	1295	Year, tropical	1234
u (quark)	1195 , 1437	Young diagrams	1289
Unified atomic mass unit	1233	Young tableaux	1289
Uniform probability density function	1272	Young's modulus of solid elements, table	1242
Unitarity triangle	1316	Yukawa databases	1186
Units and conversion factors	1233	Z gauge boson	1191 , 1353
Units, electromagnetic	1285	Z gauge boson, note on	1353
Units, SI, complete set	1240	Z gauge boson, mass, width, branching ratios, and coupling to fermions	1191 , 1233, 1304, 1309, 1372
Universe, age of	1234	Z width	1333
Universe, cosmological properties of	1236	Z' searches, note on	1372
Universe, critical density of	1234	Z^* resonances (KN system)	
Universe, curvature of	1236	(see p. VIII.58 in our 1992 edition, Phys. Rev. D45 , Part II)	
Universe, density parameter of	1234		
UNK (Serpukhov) collider parameters	1250		
Upper limits, Gaussian distribution	1278		
Upper limits, Poisson distribution	1279		
Υ states, width determinations of, note on	1661		
$\Upsilon(1S) = \Upsilon(9460)$	1213 , 1662		
$\Upsilon(2S) = \Upsilon(10023)$	1213 , 1665		
$\Upsilon(3S) = \Upsilon(10355)$	1214 , 1668		
$\Upsilon(4S) = \Upsilon(10580)$	1214 , 1669		
$\Upsilon(10860)$	1214 , 1669		

Greek letters are alphabetized by their English-language spelling. Bold page numbers signify entries in the Particle Properties Summary Tables.

PHYSICAL REVIEW D

PARTICLES AND FIELDS

VOLUME 50

Third Series

1 AUGUST 1994

PART II

Published by

THE AMERICAN PHYSICAL SOCIETY

through the

AMERICAN INSTITUTE OF PHYSICS




Figure 1: Neutrino oscillation parameter ranges excluded by a toy experiment in which one searches for the appearance of neutrinos not expected in the beam. The probability of appearance, in this case 0.5% at some specified CL, is set by the number of right-flavor events observed and/or information about the flux and cross sections. Here it is assumed that $\langle L/E \rangle = 1 \text{ km GeV}^{-1}$, and that the distribution of L/E is Gaussian with a 20% standard deviation. The wiggle structure is determined by the resolution function, and the intercepts are determined by the appearance probability and $\langle L/E \rangle$. The leftmost excursion relative to the high- $|\Delta m^2|$ limit and the slope of the lower part of the curve are independent of the experiment. In a disappearance experiment, high- $|\Delta m^2|$ sensitivity is lost unless the incident flux is known. These two possibilities are shown qualitatively by the dashed lines marked “Disappearance A” and “Disappearance B.”




Fig. 1. SU(4) multiplets of baryons made of u , d , s , and c quarks. (a) The 20-plet with an SU(3) octet on the "ground floor." (b) The 20-plet with an SU(3) decuplet on the ground floor.