

## Strong and electromagnetic decays of two new $\Lambda_c^*$ baryons

Peter Cho

*Lauritsen Laboratory, California Institute of Technology, Pasadena, California 91125*

(Received 24 January 1994)

Two recently discovered excited charm baryons are studied within the framework of heavy hadron chiral perturbation theory. We interpret these new baryons which lie 308 MeV and 340 MeV above the  $\Lambda_c$  as  $I = 0$  members of a  $P$ -wave spin doublet. Differential and total decay rates for their double pion transitions down to the  $\Lambda_c$  ground state are calculated. Estimates for their radiative decay rates are also discussed. We find that the experimentally determined characteristics of the  $\Lambda_c^*$  baryons may be simply understood within the effective theory.

PACS number(s): 14.20.Lq, 12.39.Hg, 13.30.Eg, 13.40.Hq

### I. INTRODUCTION

The discovery of the first excited charm baryon has recently been announced by the ARGUS, CLEO, and E687 groups [1–3]. The new state lies approximately 340 MeV above the  $\Lambda_c$  (2286 MeV) and decays to it via double pion emission. Although its spin, isospin, and parity are not yet known, this new charmed baryon has been preliminarily interpreted as a  $\Lambda_c^*$  resonance. CLEO has further reported evidence for a second  $\Lambda_c^*$  excitation at 308 MeV above  $\Lambda_c$  [4]. The second resonance also decays through a double pion mode that is consistent with the two step process  $\Lambda_c^* \rightarrow \Sigma_c \pi$  followed by  $\Sigma_c \rightarrow \Lambda_c \pi$ . In contrast, CLEO finds no evidence for an intermediate  $\Sigma_c$  in the decay of the first  $\Lambda_c^*$  excitation [2].

In this article, we will analyze these new baryon states and their dominant decay modes within the framework of heavy hadron chiral perturbation theory (HHCP). This hybrid effective theory represents a synthesis of chiral perturbation theory and the heavy quark effective theory (HQET) and describes the low energy interactions between light Goldstone bosons and hadrons containing a heavy quark [5–9]. Since its development a few years ago, HHCP has primarily been applied to the study of ground-state charm and bottom hadrons. Ground-state mesons and baryons are more tightly restricted by heavy quark spin symmetry than their excited counterparts. Moreover, experimental information has been much more sparse for the latter than the former. It is therefore not surprising that theorists have concentrated upon the lowest-lying hadrons in the past. Now, however, that new data are being collected, it is worthwhile to broaden the scope of HHCP and incorporate excited heavy hadrons into the effective theory.

The first excited heavy mesons and baryons are  $P$ -wave hadrons that carry one unit of orbital angular momentum.  $P$ -wave mesons have already been investigated within the HHCP framework [10–13]. It is straightforward to extend the formalism and include  $P$ -wave baryons as well. A number of unknown couplings enter into the excited baryon sector which limits one's predictive power. But as we shall see, all the general characteristics of the two  $\Lambda_c^*$  baryons reported by ARGUS, CLEO, and E687 are consistent with their being members of an

excited spin symmetry doublet. Although our findings will be more qualitative than quantitative, we hope this work may help guide experimentalists as they continue to study these new charmed baryons.

Our paper is organized as follows. In Sec. II, we incorporate the lowest lying excited baryon doublet into the heavy baryon chiral Lagrangian. We then focus upon the two new  $\Lambda_c^*$  members of this doublet and analyze their strong interaction decays in Sec. III. Radiative transitions are discussed in Sec. IV. Finally, we close in Sec. V with some thoughts on future directions for investigation.

### II. THE HEAVY BARYON CHIRAL LAGRANGIAN

We begin by recalling some basic aspects of the baryon sector in heavy hadron chiral perturbation theory [7,8]. Ground-state baryons with quark content  $Qqq$  have zero orbital angular momentum and occur in two types depending upon the angular momentum  $j_l$  of their light degrees of freedom. In the first case, the light brown muck is arranged in a symmetric  $j_l = 1$  configuration which transforms as a sextet under flavor  $SU(3)$ . The spectators consequently couple with the heavy quark to form  $J^P = \frac{1}{2}^+$  and  $J^P = \frac{3}{2}^+$   $S$ -wave bound states. When the heavy quark is taken to be charm, the spin- $\frac{1}{2}$  states are annihilated by velocity-dependent Dirac operators  $S^{ij}(v)$  whose individual components are given by

$$\begin{aligned} S^{11} &= \Sigma_c^{++}, & S^{12} &= \sqrt{\frac{1}{2}} \Sigma_c^+, & S^{22} &= \Sigma_c^0, \\ S^{13} &= \sqrt{\frac{1}{2}} \Xi_c^{+'}, & S^{23} &= \sqrt{\frac{1}{2}} \Xi_c^{0'}, \\ S^{33} &= \Omega_c^0. \end{aligned} \quad (2.1)$$

Their spin- $\frac{3}{2}$  counterparts are destroyed by corresponding  $S^{*ij}(v)$  Rarita-Schwinger operators. In the second case, the light degrees of freedom form an antisymmetric  $j_l = 0$  combination which transforms as a flavor antitriplet. Coupling with the heavy quark then yields  $J^P = \frac{1}{2}^+$  baryons which we associate with the field  $T_i(v)$ . When  $Q = c$ , the individual components of  $T_i$  are the singly charmed baryons

$$T_1 = \Xi_c^0, \quad T_2 = -\Xi_c^+, \quad T_3 = \Lambda_c^+. \quad (2.2)$$

The complete spectrum of the first orbitally excited  $P$ -wave  $Qqq$  baryons is quite complicated. The lowest lying such hadrons correspond to bound states that have one unit of orbital angular momentum inserted between the heavy quark and light diquark pair. In this case, spin statistics constrain the light degrees of freedom to belong to either a  $j_l = 1$  multiplet which transforms as a flavor antitriplet or else to  $j_l = 0, 1$ , or 2 multiplets which transform as flavor sextets. Nonrelativistic quark model calculations indicate that the antitriplet multiplet is isolated and lies significantly below all other  $P$ -wave states [14]. We will therefore only incorporate this lightest  $j_l = 1$  multiplet into the chiral Lagrangian. We assign the Dirac and Rarita-Schwinger operators  $R_i(v)$  and  $R_{\mu i}^*(v)$  to its  $J^P = \frac{1}{2}^-$  and  $J^P = \frac{3}{2}^-$  states. As we shall see, the two newly discovered  $\Lambda_c^*$  baryons are well described as the  $I = 0$  members of  $R$  and  $R_{\mu}^*$ .

In the infinite heavy quark mass limit, it is useful to combine together the degenerate  $J = \frac{1}{2}$  and  $J = \frac{3}{2}$  members of the ground-state sextet and excited antitriplet multiplets into the baryon “superfields” [8,15]

$$\begin{aligned} \mathcal{R}_{\mu i} &= \sqrt{\frac{1}{3}}(\gamma_\mu + v_\mu)\gamma^5 R_i + R_{\mu i}^*, \\ \mathcal{S}_\mu^{ij} &= \sqrt{\frac{1}{3}}(\gamma_\mu + v_\mu)\gamma^5 S^{ij} + S_{\mu}^{*ij}. \end{aligned} \quad (2.3)$$

The  $\mathcal{T}_i$  superfield for the ground-state antitriplet baryons is simply identical to  $T_i$ . The superfields transform as doublets under heavy quark spin symmetry  $SU(2)_v$ . They also transform under parity as

$$\begin{aligned} \mathcal{R}_\mu(\vec{x}, t) &\xrightarrow{P} \gamma_0 \mathcal{R}^\mu(-\vec{x}, t), \\ \mathcal{S}_\mu(\vec{x}, t) &\xrightarrow{P} -\gamma_0 \mathcal{S}^\mu(-\vec{x}, t), \\ \mathcal{T}(\vec{x}, t) &\xrightarrow{P} \gamma_0 \mathcal{T}(-\vec{x}, t), \end{aligned} \quad (2.4)$$

and obey the constraints

$$\pi = \sum_{a=1}^8 \pi^a T^a = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{1/2}\pi^0 + \sqrt{1/6}\eta & & & \\ & \pi^- & & \\ & & -\sqrt{1/2}\pi^0 + \sqrt{1/6}\eta & \\ & & & K^- \end{pmatrix} \begin{pmatrix} K^+ \\ K^0 \\ -\sqrt{2/3}\eta \end{pmatrix}. \quad (2.8)$$

It is convenient to arrange these fields into the exponentiated matrix functions  $\Sigma = e^{2i\pi/f}$  and  $\xi = e^{i\pi/f}$  where the parameter  $f$  equals the pion decay constant  $f_\pi = 93$  MeV at lowest order. The matrix functions transform under the chiral symmetry group as

$$\Sigma \rightarrow L \Sigma R^\dagger, \quad (2.9)$$

$$\xi \rightarrow L \xi U^\dagger(x) = U(x) \xi R^\dagger,$$

where  $L$  and  $R$  represent global elements of  $SU(3)_L$  and  $SU(3)_R$  while  $U(x)$  acts like a local  $SU(3)_{L+R}$  transformation. We further define the vector and axial vector

$$\begin{aligned} \frac{1+\not{v}}{2} \mathcal{R}_\mu &= \mathcal{R}_\mu, \quad \frac{1+\not{v}}{2} \mathcal{S}_\mu = \mathcal{S}_\mu, \quad \frac{1+\not{v}}{2} \mathcal{T} = \mathcal{T}, \\ v^\mu \mathcal{R}_\mu &= 0, \quad v^\mu \mathcal{S}_\mu = 0. \end{aligned} \quad (2.5)$$

These conditions ensure that  $\mathcal{R}_\mu$  and  $\mathcal{S}_\mu$  contain six degrees of freedom while  $\mathcal{T}$  has two. The degree of freedom count thus agrees with the number of states that the superfields represent [17].

The constraints in (2.5) also fix the shifts in the baryon superfields induced by the reparametrization transformation

$$\begin{aligned} v &\rightarrow v + \epsilon/M, \\ k &\rightarrow k - \epsilon. \end{aligned} \quad (2.6)$$

where  $v \cdot \epsilon = 0$ . This change of variables leaves invariant the total four-momentum  $p = Mv + k$  of a heavy hadron and induces only an  $O(1/M^2)$  correction to  $v^2 = 1$ . The method for determining the induced shifts in the baryon superfields is entirely analogous to that for their meson counterparts which has previously been discussed in Ref. [13]. So we only quote the results here:

$$\begin{aligned} \delta \mathcal{R}_\mu &= \frac{\not{\epsilon}}{2M} \mathcal{R}_\mu - \frac{\epsilon^\nu \mathcal{R}_\nu}{M} v_\mu, \\ \delta \mathcal{S}_\mu &= \frac{\not{\epsilon}}{2M} \mathcal{S}_\mu - \frac{\epsilon^\nu \mathcal{S}_\nu}{M} v_\mu, \\ \delta \mathcal{T} &= \frac{\not{\epsilon}}{2M} \mathcal{T}. \end{aligned} \quad (2.7)$$

The requirement that the effective theory remain invariant under the transformations in (2.6) and (2.7) forbids certain terms from appearing in the chiral Lagrangian [16].

The heavy baryons in the  $\mathcal{R}_\mu$ ,  $\mathcal{S}_\mu$ , and  $\mathcal{T}$  multiplets can interact with one another via emission and absorption of light Goldstone bosons. The Goldstone bosons result from the spontaneous breaking of  $SU(3)_L \times SU(3)_R$  chiral symmetry down to its diagonal  $SU(3)_{L+R}$  flavor subgroup and appear in the pion octet:

fields

$$\begin{aligned} \mathbf{V}^\mu &= \frac{1}{2}(\xi^\dagger \partial^\mu \xi + \xi \partial^\mu \xi^\dagger) \\ &= \frac{1}{2f^2}[\pi, \partial^\mu \pi] - \frac{1}{24f^4}[\pi, [\pi, [\pi, \partial^\mu \pi]]] + O(\pi^6), \end{aligned} \quad (2.10)$$

$$\begin{aligned} \mathbf{A}^\mu &= \frac{i}{2}(\xi^\dagger \partial^\mu \xi - \xi \partial^\mu \xi^\dagger) \\ &= -\frac{1}{f} \partial^\mu \pi + \frac{1}{6f^3}[\pi, [\pi, \partial^\mu \pi]] + O(\pi^5). \end{aligned}$$

which transform inhomogeneously and homogeneously under  $SU(3)_{L+R}$ , respectively:

$$\begin{aligned} \mathbf{V}^\mu &\rightarrow U\mathbf{V}^\mu U^\dagger + U\partial^\mu U^\dagger, \\ \mathbf{A}^\mu &\rightarrow U\mathbf{A}^\mu U^\dagger. \end{aligned} \quad (2.11)$$

The pions in (2.8) derivatively couple to the baryon mat-

$$\begin{aligned} \mathcal{L}_v^{(0)} = \sum_{Q=c,b} \bigg\{ & \bar{R}_\mu^i (-iv \cdot D + \Delta M_{\mathcal{R}}) \mathcal{R}_i^\mu + \bar{S}_{ij}^\mu (-iv \cdot D + \Delta M_S) S_\mu^{ij} + \bar{T}^i v \cdot D T_i + ig_1 \epsilon_{\mu\nu\sigma\lambda} \bar{S}_{ik}^\mu v^\nu (A^\sigma)_j^i (S^\lambda)^{jk} \\ & + ig_2 \epsilon_{\mu\nu\sigma\lambda} \bar{R}^{\mu i} v^\nu (A^\sigma)_i^j (\mathcal{R}^\lambda)_j + h_1 [\epsilon_{ijk} \bar{T}^i (A^\mu)_j^k S_\mu^{kl} + \epsilon^{ijk} \bar{S}_{kl}^\mu (A_\mu)_j^l T_i] \\ & + h_2 [\epsilon_{ijk} \bar{R}^{\mu i} v \cdot A_j^k S_\mu^{kl} + \epsilon^{ijk} \bar{S}_{kl}^\mu v \cdot A_j^l \mathcal{R}_{\mu i}] \bigg\}. \end{aligned} \quad (2.12)$$

A few points about this zeroth order Lagrangian should be noted. Firstly, the common mass splitting between the excited and ground-state antitriplet multiplets is absorbed into the parameter  $\Delta M_{\mathcal{R}} = M_{\mathcal{R}} - M_{\mathcal{T}}$ . Similarly,  $\Delta M_S = M_S - M_{\mathcal{T}}$  represents the splitting between the ground-state sextet and antitriplet multiplets. These parameters do not vanish in the zero or infinite heavy quark mass limits and therefore appropriately reside within the leading order chiral Lagrangian. Secondly, the coupling constants  $g_{1,2}$  and  $h_{1,2}$  in (2.12) are expected to be of order unity on general dimensional analysis grounds [18]. However, their precise numerical values are *a priori* unknown and must be fitted to experiment. Finally, we observe that there are no terms in (2.12) which mediate the single Goldstone boson transitions  $R^{(*)} \rightarrow T\pi$  and  $T \rightarrow T\pi$ . Such processes violate heavy quark spin symmetry and occur only at next-to-leading order in the  $1/m_Q$  expansion.

The current experimental status of the baryons appearing in the heavy hadron chiral Lagrangian is very uneven. Data on strange charmed baryons are in short supply, and several have not yet been discovered. In contrast, a number of experiments within the past year have filled in most of the nonstrange members of the antitriplet and sextet multiplets. We will therefore focus upon the zero strangeness baryons in the remainder of this work.

The energy levels of the observed  $\Lambda_c^{(*)}$  and  $\Sigma_c^{(*)}$  states in  $\mathcal{R}_\mu, \mathcal{S}_\mu$ , and  $\mathcal{T}$  are illustrated in Fig. 1. As indicated in the figure, we interpret the two recently observed excited charmed baryons as the  $I = 0$  members of the  $\mathcal{R}_\mu$  multiplet. In the absence of well-established names for these baryons, we adopt the nomenclature convention of Ref. [19] and denote the  $J^P = \frac{1}{2}^-$  and  $J^P = \frac{3}{2}^-$  states as  $\Lambda_{c1}$  and  $\Lambda_{c1}^*$ , respectively. Averaging over the ARGUS, CLEO, and E687 values for their masses, we find that they lie  $308.0 \pm 2.0$  and  $341.4 \pm 0.4$  MeV above  $\Lambda_c$ . The splitting between these two  $P$ -wave baryon masses is comparable in magnitude to that between their  $P$ -wave meson analogues  $D_1(2421 \text{ MeV})$  and  $D_2(2465$

ter fields via these vector and axial vector combinations.

It is straightforward to construct the lowest order effective Lagrangian which describes the low energy interactions between the  $Qqq$  baryons and Goldstone bosons. One simply writes down all possible terms that are Lorentz invariant, light chiral and heavy quark spin symmetric, and parity even:




FIG. 1. Lowest-lying  $I = 0$  and  $I = 1$  charmed baryon states. Experimentally measured mass splittings in MeV of the baryons above the  $\Lambda_c(2286 \text{ MeV})$  ground state are indicated in parentheses. The dominant pion decay modes of the excited  $J^P = \frac{1}{2}^- \Lambda_{c1}$  and  $J^P = \frac{3}{2}^- \Lambda_{c1}^*$  states are illustrated by the solid and dashed lines, respectively. Their allowed radiative transitions down to the ground state are represented by the squiggly curves.

MeV). We will keep track of this phenomenologically important mass difference even though it represents an  $O(1/m_c)$  effect.

The splitting between  $\Sigma_c^*$  and  $\Lambda_c$  displayed in Fig. 1 comes from another recent experimental result. The SKAT group claims to have observed the  $J^P = \frac{3}{2}^+ \Sigma_c^{*++}$  baryon for the first time in their bubble chamber experiment which uses a broadband neutrino beam [20]. While their mass finding  $M_{\Sigma_c^*} = 2530 \pm 7$  MeV must be treated with caution until independently confirmed by another group, we will adopt their reasonable central value in our subsequent analysis. Fortunately, none of our results will sensitively depend upon the precise numerical value for the  $\Sigma_c^*$  mass.

Having set up the necessary machinery for studying the two new  $\Lambda_c^*$  baryons, we proceed to examine their strong and radiative decay modes in the following two sections.

### III. STRONG DECAYS OF $\Lambda_c^*$

The strong decays of the newly discovered excited charmed baryons are well-suited for chiral perturbation theory analysis. The relatively small masses of  $\Lambda_{c1}$  and  $\Lambda_{c1}^*$  above  $\Lambda_c$  kinematically restrict their strong decays to soft pion emission. We therefore expect the chiral Lagrangian derivative expansion to be well behaved for these new particles. Moreover, isospin conservation and heavy quark spin symmetry both forbid single pion transitions between  $\Lambda_{c1}^{(*)}$  and  $\Lambda_c$ . The excited  $I = 0$  baryons must instead decay via an intermediate  $I = 1$  state down to the  $I = 0$  ground state. The released energy  $M_{\Lambda_{c1}^{(*)}} - M_{\Lambda_c}$  is thus shared by two pions.<sup>1</sup>

Angular momentum and parity considerations require single pion transitions between the  $\mathcal{R}_\mu$  and  $\mathcal{S}_\mu$  multiplets to go through  $L = 0$  or  $L = 2$  partial waves. The  $D$ -wave coupling arises from dimension-five operators in the next-to-leading order chiral Lagrangian whose effects are quite suppressed. The  $S$ -wave coupling on the other hand is implemented by the dimension-four term proportional to  $h_2$  in (2.12) which links  $\Lambda_{c1}$  with  $\Sigma_c$  and  $\Lambda_{c1}^*$  with  $\Sigma_c^*$ . The  $h_2$  operator consequently mediates the barely allowed transition  $\Lambda_{c1} \rightarrow \Sigma_c \pi$  at the rate

$$\begin{aligned} \Gamma(\Lambda_{c1} \rightarrow \Sigma_c \pi) &= \frac{h_2^2}{4\pi f^2} \frac{M_{\Sigma_c}}{M_{\Lambda_{c1}}} (M_{\Lambda_{c1}} - M_{\Sigma_c})^2 \sqrt{(M_{\Lambda_{c1}} - M_{\Sigma_c})^2 - m_\pi^2}. \end{aligned} \quad (3.1)$$

This process occurs so close to threshold that small isospin-violating mass differences between members of the pion and charmed sigma baryon multiplets cannot be ignored in the phase space factors of (3.1). Using

<sup>1</sup>The analogous kinematics for excited  $P$ -wave mesons is much less favorable. For example, the splitting between the  $D_2$  and  $D$  mesons is almost 600 MeV, and single pion transitions between these two states are allowed. The validity of lowest order chiral perturbation theory in this case is dubious at best.

the values  $M_{\Sigma_c^0} = 2452.0 \pm 0.7$ ,  $M_{\Sigma_c^+} = 2453.4 \pm 0.7$ , and  $M_{\Sigma_c^{++}} = 2453.1 \pm 0.7$  MeV [21], we find the partial widths

$$\Gamma(\Lambda_{c1}^+ \rightarrow \Sigma_c^0 \pi^+) = (3.3 \pm 2.8) h_2^2 \text{ MeV}, \quad (3.2a)$$

$$\Gamma(\Lambda_{c1}^+ \rightarrow \Sigma_c^+ \pi^0) = (6.0 \pm 1.5) h_2^2 \text{ MeV}, \quad (3.2b)$$

$$\Gamma(\Lambda_{c1}^+ \rightarrow \Sigma_c^{++} \pi^-) = (1.4 \pm 6.5) h_2^2 \text{ MeV}. \quad (3.2c)$$

The analogous single pion transitions between the  $J = \frac{3}{2}$  baryons in  $\mathcal{R}_\mu$  and  $\mathcal{S}_\mu$  are kinematically forbidden.

Double pion decays of  $\Lambda_{c1}$  and  $\Lambda_{c1}^*$  down to the  $\Lambda_c$  ground-state proceed at leading order via the two pole graphs displayed in Fig. 2. In order to obtain convergent decay rates from these diagrams, we must take into account the nonzero widths

$$\begin{aligned} \Gamma_{\Sigma_c} &= \frac{h_1^2}{12\pi f^2} \frac{M_{\Lambda_c}}{M_{\Sigma_c}} [(M_{\Sigma_c} - M_{\Lambda_c})^2 - m_\pi^2]^{3/2} \\ &\simeq (2.5 \pm 0.1) h_1^2 \text{ MeV}, \end{aligned} \quad (3.3)$$

$$\begin{aligned} \Gamma_{\Sigma_c^*} &= \frac{h_1^2}{12\pi f^2} \frac{M_{\Lambda_c}}{M_{\Sigma_c^*}} [(M_{\Sigma_c^*} - M_{\Lambda_c})^2 - m_\pi^2]^{3/2} \\ &\simeq (24 \pm 3) h_1^2 \text{ MeV} \end{aligned}$$

of the intermediate  $\Sigma_c$  and  $\Sigma_c^*$  resonances. Their propagators thus appear as

$$\begin{aligned} D_{\Sigma_c} &= \frac{i}{v \cdot k - (M_{\Sigma_c} - M_{\Lambda_c}) + i\Gamma_{\Sigma_c}/2} \Lambda_+, \\ D_{\Sigma_c^*}^{\mu\nu} &= \frac{i}{v \cdot k - (M_{\Sigma_c^*} - M_{\Lambda_c}) + i\Gamma_{\Sigma_c^*}/2} \Lambda_+^{\mu\nu}, \end{aligned} \quad (3.4)$$

where  $\Lambda_+ = (1 + \not{v})/2$  and  $\Lambda_+^{\mu\nu} = [-g^{\mu\nu} + v^\mu v^\nu + \frac{1}{3}(\gamma^\mu + v^\mu)(\gamma^\nu - v^\nu)] \Lambda_+$  denote spin  $-\frac{1}{2}$  and spin  $-\frac{3}{2}$  projection operators, respectively. We must also include a symmetry factor of  $1/2$  in the angular integration over the pions' momenta to avoid double counting the two identical bosons in the final state. A straightforward computation then yields the dimensionless differential decay rate




FIG. 2. Leading order pole graphs that contribute to  $\Lambda_{c1}^{(*)} \rightarrow \Lambda_c \pi \pi$ .

$$\frac{d\Gamma(\Lambda_{c1}^{(*)} \rightarrow \Lambda_c \pi^a \pi^b)}{dE_1} = \frac{\delta^{ab}}{192\pi^3} \left( \frac{h_1 h_2}{f^2} \right)^2 \frac{M_{\Lambda_c}}{M_{\Lambda_{c1}^{(*)}}} \sqrt{(E_1^2 - m_\pi^2)(E_2^2 - m_\pi^2)} \times \left[ \frac{E_1^2(E_2^2 - m_\pi^2)}{(M_{\Lambda_{c1}^{(*)}} - M_{\Sigma_c^{(*)}} - E_1)^2 + \Gamma_{\Sigma_c^{(*)}}^2/4} + \frac{(E_1^2 - m_\pi^2)E_2^2}{(M_{\Lambda_{c1}^{(*)}} - M_{\Sigma_c^{(*)}} - E_2)^2 + \Gamma_{\Sigma_c^{(*)}}^2/4} \right] \quad (3.5)$$

expressed in terms of the two pion energies  $E_1$  and  $E_2 = M_{\Lambda_{c1}^{(*)}} - M_{\Lambda_c} - E_1$  measured in the decay body's rest frame.<sup>2</sup> Integrating over  $E_1$ , we obtain the total rate

$$\Gamma(\Lambda_{c1}^{(*)} \rightarrow \Lambda_c \pi^a \pi^b) = \frac{h_2^2 \delta^{ab}}{8\pi^2 f^2} \frac{M_{\Sigma_c^{(*)}}}{M_{\Lambda_{c1}^{(*)}}} \frac{I}{[(M_{\Sigma_c^{(*)}} - M_{\Lambda_c})^2 - m_\pi^2]^{3/2}}, \quad (3.6)$$

where

$$I = \frac{\Gamma_{\Sigma_c^{(*)}}}{2} \int_{m_\pi}^{M_{\Lambda_{c1}^{(*)}} - M_{\Lambda_c} - m_\pi} dE_1 \sqrt{(E_1^2 - m_\pi^2)(E_2^2 - m_\pi^2)} \times \left[ \frac{E_1^2(E_2^2 - m_\pi^2)}{(M_{\Lambda_{c1}^{(*)}} - M_{\Sigma_c^{(*)}} - E_1)^2 + \Gamma_{\Sigma_c^{(*)}}^2/4} + \frac{(E_1^2 - m_\pi^2)E_2^2}{(M_{\Lambda_{c1}^{(*)}} - M_{\Sigma_c^{(*)}} - E_2)^2 + \Gamma_{\Sigma_c^{(*)}}^2/4} \right]. \quad (3.7)$$

Since we do not know the values of  $h_1$  and  $h_2$ , we cannot extract precise quantitative predictions from Eqs. (3.5)–(3.7). However, these formulas do provide useful qualitative insight into the  $P$ -wave baryons' strong decays. In Fig. 3, we plot  $h_2^{-2} d\Gamma(\Lambda_{c1}^{(*)} \rightarrow \Lambda_c \pi^0 \pi^0)/dE_1$  versus  $E_1$  with  $h_1$  set equal to unity. As can clearly be seen in the figure,  $\Lambda_{c1} \rightarrow \Lambda_c \pi^0 \pi^0$  is dominated by the pole regions where the intermediate  $\Sigma_c^+$  state is very close to being on-shell. Its integrated rate is thus well approximated by the single  $\pi^0$  partial width in (3.2b). The rate in the charged pion channel is similarly well approximated by the sum of the widths in (3.2a) and (3.2c). Indeed, evaluating the phase space integral in (3.7) using the narrow width approximation,

$$\frac{\Gamma_{\Sigma_c}/2}{(M_{\Lambda_{c1}} - M_{\Sigma_c} - E)^2 + (\Gamma_{\Sigma_c}/2)^2} \simeq \pi \delta(M_{\Lambda_{c1}} - M_{\Sigma_c} - E), \quad (3.8)$$

we simply recover Eq. (3.1) for  $\Gamma(\Lambda_{c1} \rightarrow \Sigma_c \pi)$  which is independent of coupling constant  $h_1$ .

Nonresonant contributions generate a slight dependence of  $\Gamma(\Lambda_{c1} \rightarrow \Lambda_c \pi^a \pi^b)$  upon  $h_1$  as shown in Fig. 4. But the decay of the  $J^P = \frac{1}{2}^-$  state may essentially be viewed as the two step process  $\Lambda_{c1} \rightarrow \Sigma_c \pi$  followed by  $\Sigma_c \rightarrow \Lambda_c \pi$ . In contrast, the double pion decay of  $\Lambda_{c1}^*$

cannot be regarded as a sequential transition. The virtual  $\Sigma_c^*$  intermediate state is very much off-shell and produces no large resonant contribution to  $\Lambda_{c1}^* \rightarrow \Lambda_c \pi \pi$ . As a result, the strong interaction partial width of the  $J^P = \frac{3}{2}^-$  state is more than an order of magnitude smaller than that of its  $J^P = \frac{1}{2}^-$  partner.

As advertised in the Introduction, these qualitative findings on the excited charm baryon decay modes are in basic accord with the recent CLEO results reported in Refs. [2,4]. They thus bolster one's confidence in the interpretation of the two new states as  $\Lambda_c^*$  members of a  $P$ -wave spin doublet. To make further progress, however, we need width information to pin down the values of the coupling constants in chiral Lagrangian (2.12). ARGUS




FIG. 3. Dimensionless differential decay rates  $h_2^{-2} d\Gamma(\Lambda_{c1} \rightarrow \Lambda_c \pi^0 \pi^0)/dE_1$  (solid curve) and  $10 \times h_2^{-2} d\Gamma(\Lambda_{c1} \rightarrow \Lambda_c \pi^+ \pi^0)/dE_1$  (dashed curve) plotted against pion energy  $E_1$  measured in the excited charm baryon's rest frame. The coupling constant  $h_1$  is set equal to unity in this graph.

<sup>2</sup>In the infinite charm mass limit, the recoiling  $\Lambda_c$  baryon carries off momentum but no kinetic energy. The two pions thus share all of the energy released by the decaying  $\Lambda_{c1}^{(*)}$ . This situation is similar to bouncing a ball off the earth. The earth must recoil to conserve momentum, but the ball bounces back with practically all its original kinetic energy [22].




FIG. 4. Integrated double pion decay rates of  $\Lambda_{c1}$  (solid curve) and  $\Lambda_{c1}^*$  (dashed curve) plotted as functions of coupling constant  $h_1$ . The neutral and charged pion channel contributions are summed together in this graph.

has set a 90% C.L. upper bound of 3.2 MeV on the width of  $\Lambda_c^*$  [1]. Unfortunately, this limit places only a weak constraint on the allowed parameter space in the  $h_1$ - $h_2$  plane. As Fig. 4 demonstrates, the true natural width of  $\Lambda_{c1}^*$  is most likely too narrow to be resolved by current experimental detectors. On the other hand, there is a much better chance that the  $\Lambda_{c1}$  resonance is wide enough to be measured. In the  $I = 1$  sector, the numerical results displayed in Eq. (3.3) indicate that at least one of the widths for the  $J^P = \frac{1}{2}^+$  and  $J^P = \frac{3}{2}^+$  members of the  $\mathcal{S}_\mu$  doublet can be resolved over a reasonable range for  $h_1$ . We are therefore hopeful that experimentalists will be able to fix some of the free parameters in the heavy baryon chiral Lagrangian in the near future.

$$\begin{aligned} \mathcal{L}_v^{\text{EM}} = \sum_{Q=c,b} \frac{e}{\Lambda} \left\{ i c_R \bar{\mathcal{R}}_\mu^j Q_j^i \mathcal{R}_{\nu i} F^{\mu\nu} + i c_s \bar{\mathcal{S}}_{\nu ij} (Q_k^i S_\mu^{kj} + Q_k^j S_\mu^{ik}) F^{\mu\nu} + c_{RS} [\epsilon_{ijk} \bar{\mathcal{R}}_\mu^i Q_l^j S_\nu^{kl} + e^{ijk} \bar{\mathcal{S}}_{\nu,kl} Q_j^l \mathcal{R}_{\mu i}] \tilde{F}^{\mu\nu} \right. \\ \left. + c_{RT} [\bar{\mathcal{T}}^j Q_j^i \mathcal{R}_i^\mu + \bar{\mathcal{R}}^{\mu i} Q_i^j \mathcal{T}_j] v^\nu F_{\mu\nu} + c_{ST} [\epsilon_{ijk} \bar{\mathcal{T}}^i Q_l^j S_\nu^{kl} + \epsilon^{ijk} \bar{\mathcal{S}}_{\nu,kl} Q_j^l \mathcal{T}_i] v_\mu \tilde{F}^{\mu\nu} \right\}. \quad (4.1) \end{aligned}$$

Here  $F^{\mu\nu}$  and  $\tilde{F}^{\mu\nu}$  are the electromagnetic field strength tensor and its dual, and  $\mathcal{Q} = \frac{1}{2}(\xi Q_{\text{EM}} \xi^\dagger + \xi^\dagger Q_{\text{EM}} \xi)$  where

$$\mathcal{Q}_{\text{EM}} = \begin{pmatrix} Q_u & & \\ & Q_d & \\ & & Q_s \end{pmatrix} = \begin{pmatrix} \frac{2}{3} & & \\ & -\frac{1}{3} & \\ & & -\frac{1}{3} \end{pmatrix} \quad (4.2)$$

denotes the light quark electric charge matrix. The transformation rule  $\mathcal{Q} \rightarrow U \mathcal{Q} U^\dagger$  for this spurion field renders the terms in (4.1) chiral symmetric [27].

The terms in (4.1) proportional to  $c_R$ ,  $c_S$ , and  $c_{ST}$  mediate the  $M1$  radiative transitions  $R^* \rightarrow R\gamma$ ,  $S^* \rightarrow S\gamma$ , and  $S^{(*)} \rightarrow T\gamma$ . The  $c_{RS}$  and  $c_{RT}$  operators on the other hand generate  $E1$  decays  $R^{(*)} \rightarrow S^{(*)}\gamma$  and  $R^{(*)} \rightarrow T\gamma$ . After extracting the  $\Lambda_{c1}^{(*)}$  components from

#### IV. ELECTROMAGNETIC DECAYS OF $\Lambda_c^*$

The only decay modes of the two new  $\Lambda_{c1}^{(*)}$  baryons that have so far been experimentally observed are their double pion transitions to  $\Lambda_c$ . But as shown in Fig. 1, these  $P$ -wave hadrons can also deexcite down to the ground state via single photon emission. Unlike the strong interaction processes, the radiative channels are not severely phase space suppressed. Moreover, they produce two rather than three bodies in the final state. So the inherently weaker strength of the electromagnetic transitions could be offset by their more favorable kinematics. We explore such a possibility in this section.

Electromagnetic interactions may be incorporated into heavy hadron chiral perturbation theory by gauging a  $U(1)_{\text{EM}}$  subgroup of the global  $SU(3)_L \times SU(3)_R$  chiral symmetry group. All derivatives appearing in the velocity dependent effective Lagrangian are then promoted to covariant derivatives with respect to electromagnetism. The leading dimension-four operators in (2.12) cannot contribute to  $S$ -matrix elements between states containing real photons. So to study heavy meson and baryon radiative transitions, one must include a number of dimension-five operators into the chiral Lagrangian. In Refs. [23–25], the  $M1$  transitions between ground-state hadrons containing a single heavy quark were analyzed. We now extend this earlier work to investigate the  $E1$  decays of the new  $\Lambda_{c1}^{(*)}$  baryons.

In the low energy theory, short wavelength photons with energies greater than the chiral symmetry breaking scale are integrated out and only long wavelength modes are retained. At lowest order in the  $1/m_Q$  expansion, photons couple to just the light brown muck inside  $Qqq$  baryons leaving the spins of their heavy quark constituents unaltered. Such interactions take place at a long-distance scale  $\Lambda$  and generate the following contributions to the effective Lagrangian:<sup>3</sup>

these interactions, we find the following radiative partial widths:

$$\begin{aligned} \Gamma(\Lambda_{c1}^* \rightarrow \Lambda_{c1}\gamma) &= \frac{4c_R^2 \alpha_{\text{EM}} M_{\Lambda_{c1}}}{81 \Lambda^2 M_{\Lambda_{c1}^*}} \left( \frac{M_{\Lambda_{c1}^*}^2 - M_{\Lambda_{c1}}^2}{2M_{\Lambda_{c1}^*}} \right)^3 \\ &= 1.07 \times 10^{-4} c_R^2 \text{ MeV}, \quad (4.3) \end{aligned}$$

<sup>3</sup>The phenomenologically successful nonrelativistic quark model suggests that an appropriate value for  $\Lambda$  is given by a typical constituent quark mass [26]. We therefore set  $\Lambda = 350$  MeV in the radiative decay rate formulas (4.3)–(4.5).

$$\begin{aligned}\Gamma(\Lambda_{c1} \rightarrow \Sigma_c \gamma) &= \frac{8c_{RS}^2 \alpha_{EM}}{9 \Lambda^2} \frac{M_{\Sigma_c}}{M_{\Lambda_{c1}}} \left( \frac{M_{\Lambda_{c1}}^2 - M_{\Sigma_c}^2}{2M_{\Lambda_{c1}}} \right)^3 \\ &= 0.127c_{RS}^2 \text{ MeV} ,\end{aligned}\quad (4.4a)$$

$$\begin{aligned}\Gamma(\Lambda_{c1} \rightarrow \Sigma_c^* \gamma) &= \frac{4c_{RS}^2 \alpha_{EM}}{9 \Lambda^2} \frac{M_{\Sigma_c^*}}{M_{\Lambda_{c1}}} \left( \frac{M_{\Lambda_{c1}}^2 - M_{\Sigma_c^*}^2}{2M_{\Lambda_{c1}}} \right)^3 \\ &= 0.006c_{RS}^2 \text{ MeV} ,\end{aligned}\quad (4.4b)$$

$$\begin{aligned}\Gamma(\Lambda_{c1}^* \rightarrow \Sigma_c \gamma) &= \frac{2c_{RS}^2 \alpha_{EM}}{9 \Lambda^2} \frac{M_{\Sigma_c}}{M_{\Lambda_{c1}^*}} \left( \frac{M_{\Lambda_{c1}^*}^2 - M_{\Sigma_c}^2}{2M_{\Lambda_{c1}^*}} \right)^3 \\ &= 0.058c_{RS}^2 \text{ MeV} ,\end{aligned}\quad (4.4c)$$

$$\begin{aligned}\Gamma(\Lambda_{c1}^* \rightarrow \Sigma_c^* \gamma) &= \frac{10c_{RS}^2 \alpha_{EM}}{9 \Lambda^2} \frac{M_{\Sigma_c^*}}{M_{\Lambda_{c1}^*}} \left( \frac{M_{\Lambda_{c1}^*}^2 - M_{\Sigma_c^*}^2}{2M_{\Lambda_{c1}^*}} \right)^3 \\ &= 0.054c_{RS}^2 \text{ MeV} ,\end{aligned}\quad (4.4d)$$

$$\begin{aligned}\Gamma(\Lambda_{c1} \rightarrow \Lambda_c \gamma) &= \frac{4c_{RT}^2 \alpha_{EM}}{27 \Lambda^2} \frac{M_{\Lambda_c}}{M_{\Lambda_{c1}}} \left( \frac{M_{\Lambda_{c1}}^2 - M_{\Lambda_c}^2}{2M_{\Lambda_{c1}}} \right)^3 \\ &= 0.191c_{RT}^2 \text{ MeV} ,\end{aligned}\quad (4.5a)$$

$$\begin{aligned}\Gamma(\Lambda_{c1}^* \rightarrow \Lambda_c \gamma) &= \frac{4c_{RT}^2 \alpha_{EM}}{27 \Lambda^2} \frac{M_{\Lambda_c}}{M_{\Lambda_{c1}^*}} \left( \frac{M_{\Lambda_{c1}^*}^2 - M_{\Lambda_c}^2}{2M_{\Lambda_{c1}^*}} \right)^3 \\ &= 0.253c_{RT}^2 \text{ MeV} .\end{aligned}\quad (4.5b)$$

As required by heavy quark spin symmetry, the sum of the widths in Eqs. (4.4a) and (4.4b) equals the sum of those in Eqs. (4.4c) and (4.4d) in the infinite charm mass limit [28]. Similarly, the rates in (4.5a) and (4.5b) become degenerate when  $m_c \rightarrow \infty$ .

The leading order results in Eqs. (4.3)–(4.5) cannot be trusted to provide much more than order of magnitude estimates for the  $\Lambda_{c1}^{(*)}$  radiative decay rates. Higher order  $1/m_c$  terms in the chiral Lagrangian are known to affect the radiative decays of  $S$ -wave charmed hadrons [23–25]. Such terms may be non-negligible for their  $P$ -wave counterparts as well. Yet comparing the leading electromagnetic partial widths with their strong interaction counterparts, we can draw some general qualitative conclusions. First, we expect on the basis of naive dimensional analysis that the unknown  $c_R$ ,  $c_{RS}$ , and  $c_{RT}$  couplings are of order unity. The numerical partial width estimates suggest that some of the electromagnetic branching fractions might be measurable. Referring to Fig. 4,

we see that the two pion decay mode of  $\Lambda_{c1}$  dominates over its radiative channels. The electromagnetic branching fraction for the  $J^P = \frac{1}{2}^-$  state is thus most likely less than a few percent. On the other hand, since the double pion width of  $J^P = \frac{3}{2}^-$   $\Lambda_{c1}^*$  is much more narrow,  $\Gamma(\Lambda_{c1}^* \rightarrow \Lambda_c \gamma)/\Gamma(\Lambda_{c1}^* \rightarrow \Lambda_c \pi \pi)$  could be sizable and perhaps greater than unity. Finally, we note that the radiative mode  $\Lambda_{c1}^* \rightarrow \Sigma_c^* \gamma$  may provide a means for detecting the  $\Sigma_c^*$  baryon. The branching fraction for this process is small but not negligible. A search for this transition could therefore yield evidence for the elusive  $I = 1$ ,  $J^P = \frac{3}{2}^+$  states.

## V. CONCLUSION

The basic interpretation of the two new excited charm baryons as  $I = 0$  members of a  $P$ -wave spin doublet holds together quite well. Since the splitting between  $\Lambda_{c1}^{(*)}$  and  $\Lambda_c$  is relatively small, these excited hadrons are well suited for incorporation into heavy hadron chiral perturbation theory. Many experimental and theoretical details clearly remain to be filled into the picture which we have outlined here. In particular, width and branching ratio information are needed to fix the several new parameters that enter into the excited baryon sector.

A number of extensions of this work could be pursued in the future. For example, the primary decay modes of the  $\Xi_{c1}^{0(*)}$  and  $\Xi_{c1}^{+(*)}$  partners of  $\Lambda_{c1}^{(*)}$  ought to be analyzed. As we have seen, there is no leading order term which links any of the states in the  $\mathcal{R}_\mu$  antitriplet superfield with members of the ground state  $\mathcal{T}$  multiplet. So if kinematically allowed, single kaon decays of these  $P$ -wave strange charmed baryons down to  $\Lambda_c$  are suppressed by  $1/m_c$ . A theoretical study of the dominant  $\Xi_{c1}^{(*)}$  transitions would help guide an experimental search for these states. Alternatively, one might consider including other excited  $P$ -wave  $Qqq$  baryons into the heavy chiral Lagrangian. There are many such states waiting to be discovered. Finally, many of the results obtained here for charm baryons may be applied to their bottom baryon counterparts. While experimental data on bottom baryons may not become available for quite some time, an eventual comparison between the charm and bottom sectors will provide valuable tests of heavy flavor symmetry ideas.

## ACKNOWLEDGMENTS

Helpful discussions with Sandip Trivedi and Mark Wise are gratefully acknowledged. This work was supported in part by the SSC Laboratory and by the U.S. Department of Energy under DOE Grant No. DE-FG03-92-ER40701.

- [1] ARGUS Collaboration, H. Albrecht *et al.*, Phys. Lett. B **317**, 227 (1993).
- [2] CLEO Collaboration, D. Acosta *et al.*, in *Lepton and Photon Interactions*, Proceedings of the 16th Interna-

- tional Symposium, Ithaca, New York, 1993, edited by P. Drell and D. Rubin (AIP, New York, 1994).
- [3] E687 Collaboration, P. L. Frabetti *et al.*, Phys. Rev. Lett. **72**, 961 (1994).

- [4] CLEO Collaboration, M. Battle *et al.*, in *Lepton and Photon Interactions* [2].
- [5] M. Wise, Phys. Rev. D **45**, R2188 (1992).
- [6] G. Burdman and J. Donoghue, Phys. Lett. B **280**, 287 (1992).
- [7] T. M. Yan, H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin, and H. L. Yu, Phys. Rev. D **46**, 1148 (1992).
- [8] P. Cho, Phys. Lett. B **285**, 145 (1992).
- [9] For a general review of heavy hadron chiral perturbation theory, see M. Wise, Caltech Report No. CALT-68-1860, 1993 (unpublished), Lectures given at the CCAST Symposium.
- [10] A. Falk and M. Luke, Phys. Lett. B **292**, 119 (1992).
- [11] U. Kilian, J. G. Körner, and D. Pirjol, Phys. Lett. B **288**, 360 (1992).
- [12] J. G. Körner, D. Pirjol, and K. Schilcher, Phys. Rev. D **47**, 3955 (1993).
- [13] P. Cho and S. Trivedi, Phys. Rev. D **50**, 381 (1994).
- [14] L. A. Copley, N. Isgur, and G. Karl, Phys. Rev. D **20**, 768 (1979).
- [15] H. Georgi, Nucl. Phys. **B348**, 293 (1991).
- [16] M. Luke and A. Manohar, Phys. Lett. B **286**, 348 (1992).
- [17] P. Cho, Nucl. Phys. **B396**, 183 (1993); Caltech Report No. CALT-68-1884 (unpublished).
- [18] H. Georgi, Phys. Lett. B **298** 187 (1993); A. Manohar and H. Georgi, Nucl. Phys. **B234**, 189 (1984); H. Georgi and L. Randall, Nucl. Phys. **B276**, 241 (1986).
- [19] J. A. Appel, in *Lepton and Photon Interactions* [2].
- [20] SKAT Collaboration, V. V. Ammosov *et al.*, in *Lepton and Photon Interactions* [2].
- [21] CLEO Collaboration, G. Crawford *et al.*, Phys. Rev. Lett. **71**, 3259 (1993).
- [22] M. Wise (private communication).
- [23] P. Cho and H. Georgi, Phys. Lett. B **296**, 408 (1992); **300**, 410(E) (1993).
- [24] J. F. Amundson, C. G. Boyd, E. Jenkins, M. Luke, A. V. Manohar, J. Rosner, M. Savage, and M. B. Wise, Phys. Lett. B **296**, 415 (1992).
- [25] H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin, T. M. Yan, and H. L. Yu, Phys. Rev. D **47**, 1030 (1993).
- [26] N. Isgur, in *The New Aspects of Subnuclear Physics*, Proceedings of the 16th International School of Subnuclear Physics, Erice, Italy, 1978, edited by A. Zichichi, Subnuclear Series Vol. 16. (Plenum, New York, 1980), p. 107.
- [27] H. Y. Cheng, C. Y. Cheung, G. L. Lin, Y. C. Lin, T. M. Yan, and H. L. Yu, Phys. Rev. D **49**, 5857 (1994).
- [28] N. Isgur and M. Wise, Phys. Rev. Lett. **66**, 1130 (1991).