

## Renormalization of composite operators in Yang-Mills theories using a general covariant gauge

J. C. Collins\* and R. J. Scalise†

*The Pennsylvania State University, Department of Physics, 104 Davey Laboratory, University Park, Pennsylvania 16802*

(Received 31 March 1994)

Essential to QCD applications of the operator product expansion, etc., is a knowledge of those operators that mix with gauge-invariant operators. A standard theorem asserts that the renormalization matrix is triangular: Gauge-invariant operators have “alien” gauge-variant operators among their counterterms, but, with a suitably chosen basis, the necessary alien operators have only themselves as counterterms. Moreover, the alien operators are supposed to vanish in physical matrix elements. A recent calculation by Hamberg and van Neerven apparently contradicts these results. By explicit calculations with the energy-momentum tensor, we show that the problems arise because of subtle infrared singularities that appear when gluonic matrix elements are taken on shell at zero momentum transfer.

PACS number(s): 11.10.Gh, 11.15.-q, 12.38.Bx

### I. HISTORICAL INTRODUCTION

Much phenomenology in QCD requires the use of the operator product expansion [1–3] and many generalizations such as “factorization theorems” [4]. Among the ingredients are matrix elements of particular gauge-invariant operators, which correspond to parton densities (or distribution functions). The properties of these operators under renormalization are vital to all QCD calculations, and one serious complication arises because gauge-invariant operators mix<sup>1</sup> with certain gauge-variant (non-gauge-invariant) operators. The renormalization directly determines the phenomenologically important anomalous dimensions of the operators—generally used in the form of Altarelli-Parisi splitting coefficients.

The extra operators that mix with the gauge-invariant operators are unphysical—we will call them “alien” operators. It has been known since the earliest days of QCD that one must demonstrate that these alien operators do not contribute to physics. Three theorems apply to the decoupling: One is that a basis can be chosen such that the alien operators are Becchi-Rouet-Stora-Tyutin (BRST) exact. Next, physical matrix elements of BRST-exact operators are zero. The last theorem is a trivial consequence of the second: The renormalization mixing matrix is triangular—alien operators do not mix with the physical operators. The theorems to establish this have been proven in their strongest form by Joglekar and Lee [5] and more recently by Henneaux [6].

Unfortunately, recent calculations by Hamberg and

van Neerven [7,8] contradict these general theorems. Their results, therefore, throw into doubt the basis of all higher-order perturbative QCD calculations. Our purpose in this paper is to resolve this contradiction between the theorem and the calculations. We will show that the contradiction is only apparent, and that it arises from certain subtle infrared (IR) problems that are unlikewise intrinsic to the usual algorithms for doing perturbative QCD calculations. However, the problem of efficiently performing practical calculations is left for future work.

The immediate motivation for the calculations by Hamberg and van Neerven was a long-standing discrepancy between calculations of the two-loop anomalous dimensions of the twist-2 covariant gluon operators in Feynman gauge [9–11] and the lightlike axial gauge [12]. Since these anomalous dimensions are measurable, calculations performed in different gauges should agree, and this can readily be shown by the methods of [13], provided that one assumes the Joglekar-Lee theorem.

Hamberg and van Neerven repeated the Feynman gauge calculation and discovered that the older calculations [9–11] are in error because they assumed the applicability of the theorem that the renormalization matrix is triangular. Hamberg and van Neerven show that the renormalization matrix appears to be nontriangular. Their calculation supports the otherwise dubious light-cone gauge result and is in accord with supersymmetry arguments [8].

The roots of this failure are already present in the one-loop part of the calculation. Although Hamberg and van Neerven do not remark on it, their calculation shows that the finite part of a physical matrix element of their alien operator is nonzero at one-loop order, contradicting the second of the Joglekar-Lee results mentioned previously. They perform their calculation for the whole tower of twist-2 covariant gluon operators, but the problems are present for the simplest of these operators, the energy-momentum tensor  $\theta_{\mu\nu}$ , for which the renormalization theory was worked out by Freedman, Muzinich, and Wein-

\*Electronic address: collins@phys.psu.edu

†Electronic address: scalise@phys.psu.edu

<sup>1</sup>Multiplicative renormalization is not sufficient to remove the infinities from Green functions of arbitrary composite operators; counterterms corresponding to different operators are needed.

berg [14,15]. The alien operators in that analysis are not manifestly the same as those used by Hamberg and van Neerven. The form of  $\theta_{\mu\nu}$  given by Freedman *et al.* is in agreement with the general theorems of Joglekar and Lee and of Henneaux. However, the gauge-variant operators used by Hamberg and van Neerven are obtained from the analysis of Dixon and Taylor [22]; it is not evident that these operators are BRST exact.

This is where we start: A sufficiently detailed analysis of the energy-momentum tensor at one-loop order is enough to locate the source of the contradiction. We will verify that at one-loop order, the renormalization given by Freedman *et al.* is in fact correct. However, the one-loop gluonic matrix element of the alien operator fails to vanish at zero momentum transfer.<sup>2,3</sup> We will find that the source of this incongruity is an infrared divergence, but the divergence is not in the calculation of the matrix element. Rather, it is a quadratic divergence in the *proof* that the matrix element vanishes.

The source of the divergence makes it clear that the proof of the theorem on the vanishing of the alien operators should be correct when one applies it to physical states. The problem arises when one considers matrix elements in an off-shell gluon state and then takes the gluon on-shell.

But this clearly threatens the rationale for the usual methods of doing perturbative QCD calculations. Moreover, the renormalization matrix that Hamberg and van Neerven calculated and found to be nontriangular presumably includes some infrared renormalization, contrary to what should be done.

In Sec. II, we state the Joglekar-Lee theorems. In Sec. III, we list our conventions for pure-gauge Yang-Mills theory. In Sec. IV, we give the results of the one-loop calculation of two-point Green functions with the energy-momentum tensor operator inserted at zero momentum transfer and derive the renormalization mixing matrix. The calculation at nonzero momentum transfer is currently underway. The off-shell results, as well as the physical matrix elements, are planned to be reported in the near future. The Appendix contains a brief discussion of “right derivatives,” the full Lorentz tensor structure of the two-gluon Green functions, which are abbreviated in the text, with a separation into leading-twist and higher twist pieces, a list of the Feynman graphs with composite operator insertions used in the calculations, and the Feynman rules for the operator vertices considered in this paper.

## II. RENORMALIZATION OF GAUGE-INVARIANT OPERATORS

In this section, we state the three theorems that Joglekar and Lee [5] proved on the renormalization of

gauge-invariant operators. In [13], the theorems are stated and all the easy parts are proven.

Let  $G_i$  denote a set of gauge-invariant operators that mix under renormalization, and let  $A_i$  denote the set of alien operators with which they mix under renormalization. (We define “alien” to mean “not gauge-invariant.”) Finally, let  $E_i$  denote the set of operators that vanish by use of the equations of motion<sup>4</sup> and with which the previous two sets of operators mix under renormalization.

The first of the Joglekar-Lee results is that the basis of the alien operators  $A_i$  that mix with gauge-invariant operators can be chosen so that they are all BRST exact; i.e., they can be written as<sup>5</sup>

$$A_i \approx \delta_{\text{BRST}} B_i, \quad (2.1)$$

where we will call  $B_i$  the “ancestor” of  $A_i$ .

The second theorem is that physical matrix elements of the BRST-exact alien operators  $\delta_{\text{BRST}} B_i$  are zero.

The last of the theorems is that the renormalization mixing matrix is triangular,

$$\begin{pmatrix} R[G] \\ R[A] \\ R[E] \end{pmatrix} = \begin{pmatrix} Z_{GG} & Z_{GA} & Z_{GE} \\ 0 & Z_{AA} & Z_{AE} \\ 0 & 0 & Z_{EE} \end{pmatrix} \begin{pmatrix} G \\ A \\ E \end{pmatrix}. \quad (2.2)$$

Of these theorems, the hardest to prove is the first. It can easily be shown that all counterterms to BRST-invariant operators are themselves BRST invariant [13]. Then one must prove that any BRST-invariant operator is a linear combination of gauge-invariant operators and BRST-exact operators. Up to operators that vanish by the equations of motion, this is supposed to be proven by Joglekar and Lee [5], but we find that the proof is very hard to understand. A simpler proof on the basis of cohomology theory is presented by Henneaux in [6].

A simple proof of the last two theorems is given in [13]. The vanishing of physical matrix elements of the alien operators follows from a simple Ward identity involving the BRST variation (also called a Slavnov-Taylor identity), once one knows that only BRST-exact operators are needed. This result trivially generalizes to show that Green functions of these alien operators with BRST-invariant operators are zero. BRST-invariant operators include gauge-invariant operators and the BRST-exact operators that comprise all our alien operators.

The third theorem, on the triangularity of the renormalization matrix, immediately follows [13]. If any on-shell, physical matrix element of an *unrenormalized* operator in class  $A$  is to vanish, then its pole piece must also vanish on shell.<sup>6</sup> Since at least some of the physical

<sup>4</sup>Matrix elements of  $E_i$  must vanish, but Green functions of  $E_i$  do not.

<sup>5</sup>We use the convention of [17], where the wavy equal sign means that the relation is only true after one or more of the equations of motions have been used.

<sup>6</sup>“Pole” in this context means a singularity as the dimension of space-time is varied. We are assuming the use of minimal subtraction with dimensional regularization to perform the renormalization (see Sec. III E).

<sup>2</sup>The momentum transfer is defined to be the sum of the momenta flowing into the inserted operator vertex.

<sup>3</sup>Harris and Smith in a recent report [16] perform the one-loop energy-momentum tensor calculations at nonzero momentum transfer and find agreement with Freedman *et al.* and with Joglekar and Lee.

matrix elements of any gauge-invariant operator are non-vanishing, it follows that the entry  $Z_{AG}$  must be zero; no operators in class  $G$  can mix with the operator in class  $A$ . Similarly,  $Z_{EG}$  and  $Z_{EA}$  in the bottom row of the mixing matrix must be zero because an *unrenormalized* operator in class  $E$  must vanish by the equations of motion; therefore, its pole part must also vanish by the equations of motion.

Note that at the level of pure Feynman graph calculations, a physical matrix element is one with the gluon polarizations being purely transverse and with the states being on-shell quarks or on-shell gluons.

Prior to the work of Joglekar and Lee, it was shown by Freedman, Muzinich, and Weinberg [14,15] how to construct a finite energy-momentum tensor for gauge theories. Their operator can readily be seen to satisfy the first Joglekar-Lee theorem, as we will explain later.

The problem we now face is that the calculations by Hamberg and van Neerven appear to violate all of the above theorems.

### III. YANG-MILLS CONVENTIONS

In this section, we list some common objects in Yang-Mills theory to exhibit our conventions and notation, but also because some (such as the energy-momentum tensor) play a pivotal rôle throughout this article.

#### A. Lagrangian density

The effective Lagrangian density of pure-gauge Yang-Mills theory in general covariant gauge is, in terms of unrenormalized (bare) fields and parameters (designated by carets), as follows:

$$\begin{aligned} \mathcal{L}(x) = & -\frac{1}{4}\hat{F}_a^{\mu\nu}(x)\hat{F}_{\mu\nu a}(x) - \frac{1}{2}\hat{\lambda}[\partial \cdot \hat{A}_a(x)][\partial \cdot \hat{A}_a(x)] \\ & + [\partial^\mu \hat{\eta}_a(x)][\hat{D}_\mu(x)\hat{\omega}_c(x)]_a, \end{aligned} \quad (3.1)$$

where the antisymmetric field strength tensor is given by

$$\hat{F}_a^{\mu\nu}(x) \equiv \partial^\mu \hat{A}_a^\nu(x) - \partial^\nu \hat{A}_a^\mu(x) - \hat{g}c_{abc}\hat{A}_b^\mu(x)\hat{A}_c^\nu(x), \quad (3.2)$$

and the covariant derivative acts on fields in the adjoint representation of the group as follows:

$$\begin{aligned} [\hat{D}_\mu(x)\hat{\omega}(x)]_a & \equiv \hat{D}_{\mu ac}(x)\hat{\omega}_c(x) \\ & \equiv [\partial_\mu \delta_{ac} - \hat{g}c_{abc}\hat{A}_{\mu b}(x)]\hat{\omega}_c(x). \end{aligned} \quad (3.3)$$

We are defining the Grassmann field  $\hat{\eta}_a(x)$  to be the antighost and the Grassmann field  $\hat{\omega}_a(x)$  to be the ghost. The parameter  $\hat{g}$  is the coupling strength,  $c_{abc}$  are the structure constants of the underlying Lie algebra  $SU(N)$ , and  $\hat{\lambda}$  is the arbitrary gauge-fixing parameter in general covariant gauge. The color indices in the adjoint representation  $a, b, c, \dots$  range from 1 to  $N^2 - 1$ .

#### B. Euler-Lagrange equations of motion

The Euler-Lagrange equations of motion, using right derivatives for the Grassmann variables<sup>7</sup> are

$$\frac{\partial^r \mathcal{L}_{\text{eff}}}{\partial \Phi_a} - \partial_\mu \frac{\partial^r \mathcal{L}_{\text{eff}}}{\partial (\partial_\mu \Phi_a)} = 0, \quad (3.4)$$

where

$$\Phi_a \in \{\hat{A}_{\nu a}, \hat{\omega}_a, \hat{\eta}_a\}. \quad (3.5)$$

We have

$$(\hat{D}_\mu \hat{F}^{\mu\nu})_a + \hat{\lambda} \partial^\nu \partial \cdot \hat{A}_a + \hat{g}c_{abc}(\partial^\nu \hat{\eta}_b)\hat{\omega}_c = 0, \quad (3.6a)$$

$$(\hat{D}_\mu \partial^\mu \hat{\eta})_a = 0, \quad (3.6b)$$

$$\partial^\mu (\hat{D}_\mu \hat{\omega})_a = 0. \quad (3.6c)$$

#### C. BRST symmetry

The gauge-fixed effective Lagrangian density is not gauge-invariant but is (quasi)invariant under the following global symmetry [18]:

$$\begin{aligned} \delta_{\text{BRST}} \hat{A}_{\mu a} & = (\hat{D}_\mu \hat{\omega})_a \delta \hat{\xi}, \\ \delta_{\text{BRST}} \hat{\omega}_a & = -\frac{1}{2} \hat{g}c_{abc} \hat{\omega}_b \hat{\omega}_c \delta \hat{\xi}, \\ \delta_{\text{BRST}} \hat{\eta}_a & = \hat{\lambda} \partial \cdot \hat{A}_a \delta \hat{\xi}. \end{aligned} \quad (3.7)$$

Here,  $\delta \hat{\xi}$  is a constant parameter with Grassmann parity 1, that is, it anticommutes with the (anti)ghost field components (and the fermion field components, if there were any), but commutes with everything else. We introduce the notation

$$\frac{\delta_{\text{BRST}}^r}{\delta \hat{\xi}}, \quad (3.8)$$

in analogy with the right derivatives for Grassmann variables to mean that  $\delta \hat{\xi}$  is commuted or anticommuted to the extreme right and then removed. This variation, called the BRST variation, is a symmetry of the Lagrangian density, Eq. (3.1), since the change in the Lagrangian density is a four divergence, without invoking the equations of motion

$$\frac{\delta_{\text{BRST}}^r \mathcal{L}}{\delta \hat{\xi}} = -\hat{\lambda} \partial^\mu [(\hat{D}_\mu \hat{\omega})_a \partial \cdot \hat{A}_a]. \quad (3.9)$$

The important property of nilpotence,

<sup>7</sup>See Sec. A 1 in the Appendix for a discussion of right derivatives.

$$\delta_{\text{BRST}}^2(\text{anything}) = 0, \quad (3.10)$$

holds only after using one of the equations of motion, Eq. (3.6c), which will be called the “trivial equation of motion” in what follows.<sup>8</sup>

#### D. Energy-momentum tensor

The symmetric, conserved energy-momentum stress tensor density can be constructed from the canonical tensor by using Belinfante’s procedure [20,21]. This is also the tensor proven by Freedman, Muzinich, and Weinberg

[14,15] to have finite Green functions with renormalized external fields. It is

$$\begin{aligned} \theta_{\mu\nu} = & -g_{\mu\nu}\mathcal{L} - \hat{F}_{\mu\rho\alpha}\hat{F}_{\nu\alpha}{}^\rho - g_{\mu\nu}\hat{\lambda}\partial^\rho(\hat{A}_{\rho\alpha}\partial\cdot\hat{A}_\alpha) \\ & + \hat{\lambda}(\hat{A}_{\mu\alpha}\partial_\nu\partial\cdot\hat{A}_\alpha) + \hat{\lambda}(\hat{A}_{\nu\alpha}\partial_\mu\partial\cdot\hat{A}_\alpha) \\ & + (\partial_\mu\hat{\eta}_\alpha)(\hat{D}_\nu\hat{\omega})_\alpha + (\partial_\nu\hat{\eta}_\alpha)(\hat{D}_\mu\hat{\omega})_\alpha = \theta_{\nu\mu}. \end{aligned} \quad (3.11)$$

The gauge-invariant piece is

$$\theta_{\mu\nu}^{(\text{GI})} = \frac{1}{4}g_{\mu\nu}\hat{F}_\alpha^{\rho\pi}\hat{F}_{\rho\pi\alpha} - \hat{F}_{\mu\rho\alpha}\hat{F}_{\nu\alpha}{}^\rho. \quad (3.12)$$

The gauge-variant piece is everything else,

$$\begin{aligned} \theta_{\mu\nu}^{(\text{GV})} = & \theta_{\mu\nu} - \theta_{\mu\nu}^{(\text{GI})} \\ = & -g_{\mu\nu}\hat{\lambda}[\partial^\rho(\hat{A}_{\rho\alpha}\partial\cdot\hat{A}_\alpha) - \frac{1}{2}(\partial\cdot\hat{A}_\alpha)(\partial\cdot\hat{A}_\alpha)] - g_{\mu\nu}(\partial^\rho\hat{\eta}_\alpha)(\hat{D}_\rho\hat{\omega})_\alpha \\ & + \hat{\lambda}(\hat{A}_{\mu\alpha}\partial_\nu\partial\cdot\hat{A}_\alpha) + \hat{\lambda}(\hat{A}_{\nu\alpha}\partial_\mu\partial\cdot\hat{A}_\alpha) \\ & + (\partial_\mu\hat{\eta}_\alpha)(\hat{D}_\nu\hat{\omega})_\alpha + (\partial_\nu\hat{\eta}_\alpha)(\hat{D}_\mu\hat{\omega})_\alpha. \end{aligned} \quad (3.13)$$

The gauge-variant piece of the energy-momentum tensor is the BRST variation of an “ancestor” operator

$$\begin{aligned} \hat{X}_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) = & (\partial_\nu\hat{\eta}_\alpha)\hat{A}_{\mu\alpha} + (\partial_\mu\hat{\eta}_\alpha)\hat{A}_{\nu\alpha} \\ & - g_{\mu\nu}[\frac{1}{2}\hat{\eta}_\alpha\partial\cdot\hat{A}_\alpha + (\partial_\rho\hat{\eta}_\alpha)\hat{A}_\alpha^\rho], \end{aligned} \quad (3.14)$$

since

$$\frac{\delta_{\text{BRST}}^\tau}{\delta\hat{\xi}}\hat{X}_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) = \theta_{\mu\nu}^{(\text{GV})} - \frac{1}{2}g_{\mu\nu}\hat{\eta}_\alpha\partial^\rho(\hat{D}_\rho\hat{\omega})_\alpha, \quad (3.15)$$

and the last term vanishes by the trivial equation of motion, Eq. (3.6c). In Sec. IV D we present the ancestor operator of the “renormalized BRST variation.”

The BRST variation of the GV piece vanishes (up to the trivial equation of motion) because of the nilpotence of the BRST transformation. We say that a gauge-variant operator such as  $\theta_{\mu\nu}^{(\text{GV})}$  is BRST exact if it has an ancestor. The BRST variation of the GI piece vanishes without using the equations of motion because the BRST variation of the gluon field is based on the gauge

transformation of that same field, so any gauge-invariant quantity is automatically BRST invariant.

#### E. Renormalization

We use multiplicative renormalization

$$\begin{aligned} \hat{A}_{\mu\alpha} = & Z_A^{\frac{1}{2}}A_{\mu\alpha}, \\ \hat{\omega}_\alpha = & Z_\omega^{\frac{1}{2}}\omega_\alpha, \\ \hat{\eta}_\alpha = & Z_\eta^{\frac{1}{2}}\eta_\alpha, \\ \hat{\lambda} = & Z_\lambda\lambda \quad (\text{it is known that } Z_\lambda = Z_A^{-1} \text{ to all orders}), \\ \hat{g} = & Z_g g\mu^\epsilon, \end{aligned} \quad (3.16)$$

and dimensional regularization in  $4 - 2\epsilon$  space-time dimensions with the modified minimal subtraction ( $\overline{\text{MS}}$ ) scheme.<sup>9</sup> (See [13] for a thorough treatment of the subject.)

The Lagrangian density, Eq. (3.1), can be written in terms of renormalized fields and parameters. This is the *same* Lagrangian density so the same symbol  $\mathcal{L}$  is used to represent both quantities<sup>10</sup>:

$$\begin{aligned} \mathcal{L} = & -\frac{1}{4}F_\alpha^{\mu\nu}F_{\mu\nu\alpha} - \frac{1}{2}\lambda(\partial\cdot A_\alpha)(\partial\cdot A_\alpha) + (\partial^\mu\eta_\alpha)(D_\mu\omega)_\alpha - \frac{1}{4}\delta Z_{II}(\partial_\mu A_{\nu\alpha} - \partial_\nu A_{\mu\alpha})(\partial^\mu A_\alpha^\nu - \partial^\nu A_\alpha^\mu) \\ & + \frac{1}{2}\delta Z_{III}g\mu^\epsilon c_{abc}(\partial_\mu A_{\nu\alpha} - \partial_\nu A_{\mu\alpha})A_b^\mu A_c^\nu - \frac{1}{4}\delta Z_{IV}g^2\mu^{2\epsilon}c_{abc}c_{ade}A_{\mu b}A_{\nu c}A_d^\mu A_e^\nu - \frac{1}{2}(Z_\lambda Z_A - 1)\lambda(\partial\cdot A_\alpha)(\partial\cdot A_\alpha) \\ & + \delta Z_0(\partial_\mu\eta_\alpha)\partial^\mu\omega_\alpha - \delta Z_I g\mu^\epsilon c_{abc}(\partial_\mu\eta_\alpha)A_b^\mu\omega_c, \end{aligned} \quad (3.17)$$

<sup>8</sup>If the Lagrangian formulation with the Nakanishi-Lautrup field is used, as in [18] and [19], no equations of motion are needed to demonstrate the nilpotence of the BRST variation.

<sup>9</sup>Hamberg and van Neerven work in  $4 + \epsilon$  dimensions.

<sup>10</sup>Some authors use the term “renormalized Lagrangian density,” but it is not always clear what is meant.

where

$$\begin{aligned}
Z_0 &\equiv Z_\omega^{\frac{1}{2}} Z_\eta^{\frac{1}{2}}, \\
Z_I &\equiv Z_A^{\frac{1}{2}} Z_\omega^{\frac{1}{2}} Z_\eta^{\frac{1}{2}} Z_g, \\
Z_{II} &\equiv Z_A, \\
Z_{III} &\equiv Z_A^{\frac{3}{2}} Z_g, \\
Z_{IV} &\equiv Z_A^2 Z_g^2.
\end{aligned} \tag{3.18}$$

Notice that the renormalized coupling  $g$  is dimensionless and that we have introduced a parameter  $\mu$  with the dimensions of mass. The renormalization constants  $\delta Z_i \equiv Z_i - 1$  have been given Roman numeral subscripts which label the number of gauge fields in the counterterm vertex.

Since the coupling is universal, the three different interaction vertices have associated renormalization constants related by the following ‘‘renormalization constant Ward identities’’:

$$Z_g = Z_I Z_A^{-\frac{1}{2}} Z_0^{-1} = Z_{III} Z_A^{-\frac{3}{2}} = Z_{IV}^{\frac{1}{2}} Z_A^{-1}. \tag{3.19}$$

In the minimal subtraction (MS) renormalization scheme, the counterterms are the negative of the pole part only, with no finite component. In the  $\overline{\text{MS}}$  scheme, the ubiquitous Euler’s constant  $\gamma_E$  and the natural logarithms of  $4\pi$  are absorbed into a new renormalization mass parameter  $\bar{\mu}$  defined by

$$\bar{\mu} \equiv \mu \left( \frac{4\pi}{e^{\gamma_E}} \right)^{\frac{1}{2}}. \tag{3.20}$$

Applied under dimensional regularization, the counterterms in either scheme are proportional to  $\frac{1}{\epsilon}$ . Although dimensional regularization is able to control both ultraviolet (UV) and IR divergences, in this paper the  $\frac{1}{\epsilon}$  di-

vergences are all ultraviolet.

We list the multiplicative renormalization constants in the  $\overline{\text{MS}}$  scheme to the order needed for an  $O(g^2)$  calculation of the Green functions considered in this paper:

$$\begin{aligned}
Z_0 &= 1 + \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \left[ \frac{1}{2} + \frac{1}{4} \left( 1 - \frac{1}{\lambda} \right) \right] + O(g^4), \\
Z_A &= 1 + \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \left[ \frac{5}{3} + \frac{1}{2} \left( 1 - \frac{1}{\lambda} \right) \right] + O(g^4), \\
Z_\lambda &= 1 - \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \left[ \frac{5}{3} + \frac{1}{2} \left( 1 - \frac{1}{\lambda} \right) \right] + O(g^4) = Z_A^{-1}, \\
Z_g &= 1 + O(g^2),
\end{aligned} \tag{3.21}$$

where  $C(A) = N$  for the gauge group  $\text{SU}(N)$ .

### F. (Modified) LSZ reduction

The residue of the propagator pole is used in the LSZ (for Lehmann, Symanzik, and Zimmermann) reduction formula to derive the  $S$  matrix from Green functions. The basic idea is that the  $S$  matrix is obtained from the asymptotic behavior of Green functions for large times ( $t \rightarrow \pm\infty$ ), and this behavior is governed by the singularities of the external propagators. We use a modified version of this procedure to handle the IR-divergent logarithms that appear in this massless theory.<sup>11</sup>

Let  $\Sigma(p)$  be the self-energy, defined as in Fig. 1. We isolate the  $p^2$  dependence of the gluon self-energy, defining  $\Pi(p^2)$  by

$$\Sigma_{\text{gluon}}^{\sigma\tau}(p) = (p^2 g^{\sigma\tau} - p^\sigma p^\tau) \Pi(p^2), \tag{3.22}$$

noting that the gluon self-energy is purely transverse due to a Ward identity [13].

In a massless theory, the singularities in propagators, as a function of  $p^2$ , are not simple poles after higher order corrections are included.

The dressed propagators for the gluon and ghost, respectively, are then

$$\mathcal{D}_{ab}^{\sigma\tau}(p) \equiv \frac{i\delta_{ab}}{p^2 + i\epsilon} \left\{ -g^{\sigma\tau} \frac{1}{1 + \Pi(p^2)} + \frac{p^\sigma p^\tau}{p^2 + i\epsilon} \left[ \left( 1 - \frac{1}{\lambda} \right) - \frac{\Pi(p^2)}{1 + \Pi(p^2)} \right] \right\} \underset{p^2 \rightarrow 0}{\sim} \frac{-i\delta_{ab} c_{\text{gluon}}^2 g^{\sigma\tau}}{p^2 + i\epsilon} + \frac{i\delta_{ab} \tilde{c}_{\text{gluon}}^2 p^\sigma p^\tau}{(p^2 + i\epsilon)^2}, \tag{3.23a}$$

and

$$D_{ab}(p) \equiv \frac{i\delta_{ab}}{p^2 - \Sigma_{\text{ghost}}(p) + i\epsilon} \underset{p^2 \rightarrow 0}{\sim} \frac{i\delta_{ab} c_{\text{ghost}}^2}{p^2 + i\epsilon}, \tag{3.23b}$$

where  $c^2$  is the residue of the propagator pole (the coefficient of  $p^2$  in the denominator). To one-loop order, the gluon self-energy is

$$\begin{aligned}
\Sigma_{\text{gluon}}^{\sigma\tau}(p) &= \frac{g^2}{16\pi^2} C_A (p^2 g^{\sigma\tau} - p^\sigma p^\tau) \left\{ -\frac{1}{4} \left( 1 - \frac{1}{\lambda} \right)^2 + \left( 1 - \frac{1}{\lambda} \right) \left[ 1 + \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] \right. \\
&\quad \left. - \frac{31}{9} + \frac{5}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right\} + O(g^4),
\end{aligned} \tag{3.24}$$

<sup>11</sup>We do not have a complete justification of our algorithm.

and the ghost self-energy to one-loop order is

$$\Sigma_{\text{ghost}}(p) = \frac{g^2}{16\pi^2} C_A p^2 \left[ 1 - \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) - \frac{1}{4} \left( 1 - \frac{1}{\lambda} \right) \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] + O(g^4). \quad (3.25)$$

The singularities in the propagators are not simple poles, but the leading power, with logarithmic corrections, is governed by the large-time behavior of the propagator. So to define the residue we use the following formulas, which would be valid when the physical masses are nonzero:

$$\begin{aligned} c_{\text{gluon}}^2 &= \left[ \frac{\partial}{\partial p^2} \{p^2 [1 + \Pi(p^2)]\} \Big|_{p^2=m_{\text{physical}}^2} \right]^{-1} \\ &= 1 - \frac{g^2}{16\pi^2} C_A \left\{ -\frac{1}{4} \left( 1 - \frac{1}{\lambda} \right)^2 + \left( 1 - \frac{1}{\lambda} \right) \left[ \frac{3}{2} + \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] - \frac{16}{9} + \frac{5}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right\} \Big|_{p^2=0} + O(g^4) \end{aligned} \quad (3.26a)$$

and

$$\begin{aligned} c_{\text{ghost}}^2 &= \left[ 1 - \frac{\partial \Sigma_{\text{ghost}}(p)}{\partial p^2} \Big|_{p^2=m_{\text{physical}}^2} \right]^{-1} \\ &= 1 - \frac{g^2}{16\pi^2} C_A \left\{ \left( 1 - \frac{1}{\lambda} \right) \left[ \frac{1}{4} + \frac{1}{4} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] - \frac{1}{2} + \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right\} \Big|_{p^2=0} + O(g^4). \end{aligned} \quad (3.26b)$$

We do not calculate  $\tilde{c}_{\text{gluon}}^2$  because it is not used in this paper. It is important to refrain from taking the on-shell limit ( $p^2 \rightarrow 0$ ) until the IR-divergent logarithms above have cancelled algebraically with similar logarithms in the amputated Green functions which are being converted to  $S$  matrix elements.

We have generalized the notion of residue to include the IR-divergent terms that arise in a massless theory. Note that we are extracting the residue of the propagator pole by taking the partial derivative of the denominator with respect to  $p^2$  at  $p^2 = 0$ , rather than simply dividing the denominator by  $p^2$ . The partial derivative extracts that piece of the IR-divergent logarithm which is proportional to  $p^2$  in a Laurent expansion. This piece is

necessary to ensure, for example, that the two-gluon matrix element of the energy-momentum tensor, Eq. (4.2), is IR finite on shell and equal to its correct value.

### G. Covariant gluon operator

In [8], Hamberg and van Neerven calculate the anomalous dimension of the covariant gluon operator to two-loop order, that is  $O(\alpha_S^2)$ , with all free Lorentz indices contracted with a null vector  $\Delta$ . This selects the highest-spin part of the operator and eliminates the need to calculate the trace terms.

The covariant gluon operator is

$$O_g^{\mu_1 \dots \mu_m}(x) = \frac{1}{2} i^{m-2} \mathcal{S} [\hat{F}_{c_1}^{\rho\mu_1}(x) \hat{D}_{c_1 c_2}^{\mu_2}(x) \hat{D}_{c_2 c_3}^{\mu_3}(x) \times \dots \times \hat{D}_{c_{m-1} c_m}^{\mu_{m-1}}(x) \hat{F}_{\rho c_m}^{\mu_m}(x)] + \text{trace terms}, \quad (3.27)$$

where  $\mathcal{S}$  denotes symmetrization of the Lorentz indices  $\mu_i$  and the trace terms make the operator traceless under all possible contractions of the free Lorentz indices in pairs. The  $c_i$  are color indices in the adjoint representation.

Hamberg and van Neerven's gauge-invariant operator is

$$O_g^{(m)} = O^{\mu_1 \dots \mu_m} \Delta_{\mu_1} \times \dots \times \Delta_{\mu_m}, \quad (3.28)$$

where  $\Delta$  is lightlike. In units such that  $c = 1 = \hbar$ , the mass dimension of this operator grows linearly with  $m$ ,

$$[O_g^{(m)}] = m + 2. \quad (3.29)$$

Selecting the highest-spin piece is equivalent to selecting the lowest twist, since

$$\text{twist} \equiv \text{mass dimension} - \text{spin}. \quad (3.30)$$

All operators of the form above, Eq. (3.28), are twist-2.

The simplest case ( $m = 2$ ) of the covariant gluon operator, Eq. (3.27), gives, up to a multiplicative factor, the gauge-invariant part of the energy-momentum tensor, Eq. (3.12)

$$O_g^{\mu\nu} = \frac{1}{2} \hat{F}_a^{\mu\rho} \hat{F}_{\rho a}^{\nu} - \frac{1}{8} g^{\mu\nu} \hat{F}_a^{\rho\pi} \hat{F}_{\rho\pi a} = -\frac{1}{2} \theta^{(\text{GI}) \mu\nu}. \quad (3.31)$$

We study this case because of the relative simplicity of the calculation, but also because the gauge-variant operators which mix with it are supposed to be known [14,15]; they are the gauge-variant operators in the






FIG. 2. All Feynman diagrams that contribute to the one-loop amputated gluon two-point function at zero momentum transfer. The symbol  $\otimes$  represents the inserted composite operators  $\theta_{\mu\nu}$ ,  $\theta_{\mu\nu}^{(GI)}$ , or  $\theta_{\mu\nu}^{(GV)}$ .

**A. Green functions of  $\theta_{\mu\nu}$  with two gluon fields**

While we give the pole pieces in their entirety, the finite parts have been simplified for clarity in presentation. The full tensor structure can be found in the Appendix, where we also list the twist-2 (spin-2) piece of the operators. The Feynman graphs for these Green functions are displayed in Fig. 2. Since we are working at zero momentum transfer, the graph in Fig. 3 vanishes in dimensional regularization because the loop integral contains no momentum scale. This graph will, however, contribute at nonzero momentum transfer.

**1. Entire energy-momentum tensor**

Consider the amputated gluon two-point Green function with the entire energy-momentum tensor, Eq. (3.11), inserted at zero momentum transfer. The external gluon fields have not been contracted with physical polarization vectors, we have not multiplied by the modified LSZ residue of the gluon propagator pole, and the external momenta have not been put on shell. This is what we will mean by an off-shell gluon Green function in the sections that follow. Explicit calculation gives

$$\langle 0|T A_{\sigma a} \theta_{\mu\nu} A_{\tau b}|0\rangle_{\text{amputated}} = p_{\mu} p_{\nu} g_{\sigma\tau} \delta_{ab} \left( -2 + \frac{g^2}{16\pi^2} C_A \left\{ \frac{1}{2} \left( 1 - \frac{1}{\lambda} \right)^2 - \left( 1 - \frac{1}{\lambda} \right) \left[ 3 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] + \frac{32}{9} - \frac{10}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right\} \right) + \text{UV-finite terms that vanish on shell} + O(g^4). \quad (4.1)$$

Notice that this object is UV finite. A glance at Eq. (A2) in the Appendix will satisfy the reader that the terms not included above are also UV finite, even off shell. The UV finiteness supports the results of Freedman *et al.* [14,15] on the renormalization of the symmetric energy-momentum tensor. Also, since there is no pole, the contribution to the anomalous dimension is zero.

We now do what Hamberg and van Neerven argue, quite reasonably, in their paper is impossible. We construct a matrix element between massless gluon states, but to do so we use the modified LSZ prescription described in Sec. III F.

Contracting with physical polarization vectors, using the modified LSZ residue of the gluon propagator pole  $c_{\text{gluon}}^2$  in Eq. (3.26a), and putting the external momenta on shell, we get the relatively simple  $S$  matrix element

$$\langle \epsilon_1, p, a | \theta_{\mu\nu} | \epsilon_2, p, b \rangle = -2 p_{\mu} p_{\nu} \delta_{ab} \epsilon_1^* \cdot \epsilon_2 + O(g^4), \quad (4.2)$$

where we have used the fact that  $\epsilon_i$  is a physical polar-

ization vector satisfying

$$p \cdot \epsilon_i = 0, \quad i = 1, 2. \quad (4.3)$$

The physical state  $|\epsilon_i, p, a\rangle$  is meant to represent an on-shell gluon of momentum  $p$ , polarization vector  $\epsilon_i$ , and color  $a$ .

The modified LSZ procedure eliminates the IR-divergent logarithms even before the external momenta are taken on shell. The result, Eq. (4.2), is not surprising since  $\theta_{\mu\nu}$  is the conserved Noether current and

$$P_{\nu} \equiv \int d^3 \mathbf{x} \theta_{0\nu}(x) \quad (4.4)$$

is the Noether charge. It measures the physical (non-IR-divergent) energy momentum in a state. A correct calculation should show that all the higher order corrections to the right-hand side of Eq. (4.2) vanish.

In the next two sections, we examine the GI and GV pieces separately.

**2. Gauge-invariant part**

Consider now the amputated off-shell gluon two-point Green function with only the gauge-invariant piece of the energy-momentum tensor, Eq. (3.12), inserted at zero momentum transfer. Explicit calculation gives



$$\begin{aligned}
\langle 0|TA_{\sigma a}\theta_{\mu\nu}^{(GI)}A_{\tau b}|0\rangle_{\text{amputated}} &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} \left[ p^2 \left( -\frac{1}{2} g_{\sigma\tau} g_{\mu\nu} + g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu} \right) \right. \\
&\quad \left. - \frac{1}{2} (p_{\tau} p_{\nu} g_{\sigma\mu} + p_{\tau} p_{\mu} g_{\sigma\nu} + p_{\sigma} p_{\nu} g_{\tau\mu} + p_{\sigma} p_{\mu} g_{\tau\nu} - p_{\sigma} p_{\tau} g_{\mu\nu}) \right] \\
&\quad + p_{\mu} p_{\nu} g_{\sigma\tau} \delta_{ab} \left( -2 + \frac{g^2}{16\pi^2} C_A \left\{ \left(1 - \frac{1}{\lambda}\right)^2 - \left(1 - \frac{1}{\lambda}\right) \left[ 6 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] \right. \right. \\
&\quad \left. \left. + \frac{86}{9} - \frac{10}{3} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right\} \right) + \text{UV-finite terms that vanish on shell} + O(g^4). \quad (4.5)
\end{aligned}$$

There is an UV pole in Eq. (4.5), but this UV divergence vanishes on shell with physical polarizations.

A further clue that we have performed the calculation correctly is the fact that Eq. (A4) satisfies the Ward identity

$$p_{\sigma} p_{\tau} \langle 0|TA_{\sigma a}\theta_{\mu\nu}^{(GI)}A_{\tau b}|0\rangle_{\text{amputated}} = 0. \quad (4.6)$$

Taking into account Hamberg and van Neerven's different dimensional regularization prescription (dimension  $4 + \epsilon$  instead of  $4 - 2\epsilon$ ) and their use of bare fields instead of renormalized fields for the external legs, our result contracted with lightlike vectors, Eq. (A5), agrees with theirs.

If we now put this result on mass shell and use the modified LSZ procedure to derive the  $S$ -matrix element, we get

$$\langle \epsilon_1, p, a | \theta_{\mu\nu}^{(GI)} | \epsilon_2, p, b \rangle = p_{\mu} p_{\nu} \delta_{ab} \epsilon_1^* \cdot \epsilon_2 \left\{ -2 + \frac{g^2}{16\pi^2} C_A \left[ \frac{1}{2} \left(1 - \frac{1}{\lambda}\right)^2 - 3 \left(1 - \frac{1}{\lambda}\right) + 6 \right] \right\} + O(g^4). \quad (4.7)$$

Notice that this physical matrix element of a gauge-invariant operator depends on the gauge-fixing parameter  $\lambda$ . Also, the GI part is not equal to the total, so we must calculate the GV part.

### 3. Gauge-variant (alien) part

Consider the amputated off-shell gluon two-point Green function with only the gauge-variant piece of the energy-momentum tensor, Eq. (3.13), inserted at zero momentum transfer. Explicit calculation gives

$$\begin{aligned}
\langle 0|TA_{\sigma a}\theta_{\mu\nu}^{(GV)}A_{\tau b}|0\rangle_{\text{amputated}} &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} \left[ p^2 \left( \frac{1}{2} g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu} \right) \right. \\
&\quad \left. + \frac{1}{2} (p_{\tau} p_{\nu} g_{\sigma\mu} + p_{\tau} p_{\mu} g_{\sigma\nu} + p_{\sigma} p_{\nu} g_{\tau\mu} + p_{\sigma} p_{\mu} g_{\tau\nu} - p_{\sigma} p_{\tau} g_{\mu\nu}) \right] \\
&\quad + p_{\mu} p_{\nu} g_{\sigma\tau} \delta_{ab} \frac{g^2}{16\pi^2} C_A \left[ -\frac{1}{2} \left(1 - \frac{1}{\lambda}\right)^2 + 3 \left(1 - \frac{1}{\lambda}\right) - 6 \right] \\
&\quad + \text{UV-finite terms that vanish on shell} + O(g^4). \quad (4.8)
\end{aligned}$$

The pole terms cancel between the GI and GV parts off shell. On shell, each pole piece vanishes individually.

Going on-shell and using the modified LSZ procedure, we get the  $S$ -matrix element

$$\langle \epsilon_1, p, a | \theta_{\mu\nu}^{(GV)} | \epsilon_2, p, b \rangle = p_{\mu} p_{\nu} \delta_{ab} \epsilon_1^* \cdot \epsilon_2 \frac{g^2}{16\pi^2} C_A \left[ -\frac{1}{2} \left(1 - \frac{1}{\lambda}\right)^2 + 3 \left(1 - \frac{1}{\lambda}\right) - 6 \right] + O(g^4). \quad (4.9)$$

Notice that the finite part of the physical matrix element does not vanish. But,  $\theta_{\mu\nu}^{(GV)}$  is BRST exact, as we observed, and there is a general theorem that BRST-exact operators have vanishing physical matrix elements. Thus, we know that we have a contradiction with general theorems.

Hamberg and van Neerven have a similar result implicit in their formulas (they did not remark on it), but since their alien operators are not BRST exact, physical matrix elements of their alien operators would not be expected to vanish.

## B. Green functions of $\theta_{\mu\nu}$ with one ghost field and one antighost field

### 1. Entire energy-momentum tensor

We have just seen that the two-gluon matrix element of the gauge-variant part of the energy-momentum tensor is nonzero. Since  $\theta_{\mu\nu}^{(GV)}$  is BRST exact, this contradicts a crucial part of the theory on the renormalization of gauge-invariant operators and so we cannot take for




FIG. 3. The Feynman diagram that vanishes at zero momentum transfer, but contributes to the one-loop gluon two-point function at nonzero momentum transfer. The symbol  $\otimes$  represents the inserted composite operators  $\theta_{\mu\nu}$ ,  $\theta_{\mu\nu}^{(\text{GI})}$ , or  $\theta_{\mu\nu}^{(\text{GV})}$ .

granted any of the results of this theory, and must verify the results.

In particular, we need to verify the finiteness at one-loop order of Green functions of the energy-momentum tensor. The preceding section has established this for the gluon two-point Green function and in this section we verify finiteness for the ghost-antighost Green function. The Feynman graphs for these Green functions are displayed in Fig. 4.

All the necessary counterterms are determined by the formula for  $\theta_{\mu\nu}$ ; they are obtained by expanding  $\theta_{\mu\nu}$  in




FIG. 4. All Feynman diagrams that contribute to the one-loop amputated ghost two-point function. The symbol  $\otimes$  represents the inserted composite operators  $\theta_{\mu\nu}$ ,  $\theta_{\mu\nu}^{(\text{GI})}$ , or  $\theta_{\mu\nu}^{(\text{GV})}$ .

terms of renormalized fields by Eq. (3.16).

Consider the amputated off-shell ghost two-point Green function with the entire energy-momentum tensor, Eq. (3.11), inserted at zero momentum transfer. Explicit calculation gives

$$\begin{aligned} \langle 0|T\omega_a\theta_{\mu\nu}\eta_b|0\rangle_{\text{amputated}} &= \delta_{ab}(2p_\mu p_\nu - p^2 g_{\mu\nu}) \\ &+ \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( \left(1 - \frac{1}{\lambda}\right) \left\{ \frac{1}{2} \left[1 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] p_\mu p_\nu - \frac{1}{4} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) p^2 g_{\mu\nu} \right\} \right. \\ &\quad \left. + \left[-1 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] p_\mu p_\nu + \left[1 - \frac{1}{2} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] p^2 g_{\mu\nu} \right) + O(g^4). \end{aligned} \quad (4.10)$$

Using the modified LSZ residue of the ghost propagator pole,  $c_{\text{ghost}}^2$  in Eq. (3.26b), and putting the external momenta on shell, we get the comparatively simple  $S$ -matrix element

$$\langle p, a|\theta_{\mu\nu}|p, b\rangle = 2p_\mu p_\nu \delta_{ab} + O(g^4), \quad (4.11)$$

which is correct for the expectation value of  $\theta_{\mu\nu}$  in a properly normalized state of momentum  $p$ . Here the state vector  $|p, a\rangle$  is meant to represent an on-shell ghost of momentum  $p$  and color  $a$ . Again, although we only performed the one-loop calculation, all higher order corrections should vanish.

The twist-2 (spin-2) piece of the amputated Green function, Eq. (4.10) above, in which the free Lorentz indices of the inserted operator,  $\mu$  and  $\nu$ , are contracted with a null vector  $\Delta$  is

$$\begin{aligned} \langle 0|T\omega_a\Delta^\mu\theta_{\mu\nu}\Delta^\nu\eta_b|0\rangle_{\text{amputated}} &= (p \cdot \Delta)^2 \delta_{ab} \left( 2 + \frac{g^2}{16\pi^2} C_A \left\{ \frac{1}{2} \left(1 - \frac{1}{\lambda}\right) \left[1 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] - 1 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right\} \right) \\ &+ O(g^4). \end{aligned} \quad (4.12)$$

## 2. Gauge-invariant part

Consider now the amputated off-shell ghost two-point Green function with only the gauge-invariant piece of the energy-momentum tensor, Eq. (3.12), inserted at zero momentum transfer. Explicit calculation gives

$$\begin{aligned} \langle 0|T\omega_a\theta_{\mu\nu}^{(GI)}\eta_b|0\rangle_{\text{amputated}} &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( p_\mu p_\nu - \frac{1}{4} p^2 g_{\mu\nu} \right) \\ &+ \frac{g^2}{16\pi^2} C_A \delta_{ab} \left\{ \left[ 1 - \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p_\mu p_\nu + \frac{1}{4} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) p^2 g_{\mu\nu} \right\} + O(g^4). \end{aligned} \quad (4.13)$$

The twist-2 (spin-2) piece of this amputated Green function in which the free Lorentz indices of the inserted operator,  $\mu$  and  $\nu$ , are contracted with a null vector  $\Delta$  is

$$\langle 0|T\omega_a\Delta^\mu\theta_{\mu\nu}^{(GI)}\Delta^\nu\eta_b|0\rangle_{\text{amputated}} = (p \cdot \Delta)^2 \frac{g^2}{16\pi^2} C_A \delta_{ab} \left[ \frac{1}{\epsilon} + 1 - \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] + O(g^4). \quad (4.14)$$

### 3. Gauge-variant (alien) part

Consider the amputated off-shell ghost two-point Green function with only the gauge-variant piece of the energy-momentum tensor, Eq. (3.13), inserted at zero momentum transfer. Explicit calculation gives

$$\begin{aligned} \langle 0|T\omega_a\theta_{\mu\nu}^{(GV)}\eta_b|0\rangle_{\text{amputated}} &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( -p_\mu p_\nu + \frac{1}{4} p^2 g_{\mu\nu} \right) + \delta_{ab} (2p_\mu p_\nu - p^2 g_{\mu\nu}) \\ &+ \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( \left( 1 - \frac{1}{\lambda} \right) \left\{ \frac{1}{2} \left[ 1 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p_\mu p_\nu - \frac{1}{4} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) p^2 g_{\mu\nu} \right\} \right. \\ &\left. + 2 \left[ -1 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p_\mu p_\nu + \left[ 1 - \frac{3}{4} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p^2 g_{\mu\nu} \right) + O(g^4). \end{aligned} \quad (4.15)$$

Notice that the pole terms cancel between the GI and GV pieces.

The twist-2 (spin-2) piece of this amputated Green function in which the free Lorentz indices of the inserted operator,  $\mu$  and  $\nu$ , are contracted with a null vector  $\Delta$  is

$$\begin{aligned} \langle 0|T\omega_a\Delta^\mu\theta_{\mu\nu}^{(GV)}\Delta^\nu\eta_b|0\rangle_{\text{amputated}} &= (p \cdot \Delta)^2 \delta_{ab} \left( 2 + \frac{g^2}{16\pi^2} C_A \left\{ -\frac{1}{\epsilon} + \frac{1}{2} \left( 1 - \frac{1}{\lambda} \right) \left[ 1 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] \right. \right. \\ &\left. \left. - 2 + 2 \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right\} \right) + O(g^4). \end{aligned} \quad (4.16)$$

### C. Renormalization mixing matrix

If we do not require *a priori* that the matrix be triangular, then the most general form is

$$\begin{pmatrix} R \left[ \theta_{\mu\nu}^{(GI)} \right] \\ R \left[ \theta_{\mu\nu}^{(GV)} \right] \\ R \left[ E_{\mu\nu} \right] \end{pmatrix} = \begin{pmatrix} Z_{GG} & Z_{GA} & Z_{GE} \\ Z_{AG} & Z_{AA} & Z_{AE} \\ 0 & 0 & Z_{EE} \end{pmatrix} \begin{pmatrix} \theta_{\mu\nu}^{(GI)} \\ \theta_{\mu\nu}^{(GV)} \\ E_{\mu\nu} \end{pmatrix}, \quad (4.17)$$

where the operator of class  $E$ , which vanishes by the equations of motion Eqs. (3.6) and mixes with the operators in the energy-momentum tensor, is

$$\begin{aligned} E_{\mu\nu} &= \hat{A}_\mu \left[ (\hat{D}_\rho \hat{F}^\rho)_\nu \right] + \hat{\lambda} \partial_\nu \partial \cdot \hat{A}_\mu + \hat{g} c_{abc} (\partial_\nu \hat{\eta}_b) \hat{\omega}_c + \hat{A}_\nu \left[ (\hat{D}_\rho \hat{F}^\rho)_\mu \right] + \hat{\lambda} \partial_\mu \partial \cdot \hat{A}_\nu + \hat{g} c_{abc} (\partial_\mu \hat{\eta}_b) \hat{\omega}_c \\ &- \frac{1}{2} g_{\mu\nu} \hat{A}_\pi \left[ (\hat{D}_\rho \hat{F}^{\rho\pi}) \right] + \hat{\lambda} \partial^\pi \partial \cdot \hat{A}_\mu + \hat{g} c_{abc} (\partial^\pi \hat{\eta}_b) \hat{\omega}_c + \frac{1}{2} \alpha g_{\mu\nu} (\hat{D}_\rho \partial^\rho \hat{\eta}) \hat{\omega}_\alpha + \frac{1}{2} (1 - \alpha) g_{\mu\nu} \hat{\eta}_\alpha \partial^\rho (\hat{D}_\rho \hat{\omega})_\alpha. \end{aligned} \quad (4.18)$$

Like the fields in the energy-momentum tensor, the fields in the operator  $E_{\mu\nu}$  above are bare. The parameter  $\alpha$  in the last two terms above is free to vary since the matrix elements considered are not sufficient to distinguish be-

tween the equations of motion for the ghost and antighost fields.

These are the only dimension-four,  $SU(N)$  singlet operators with two free Lorentz indices that can be formed

from the equations of motion. The coefficients are determined by demanding that the GI and GV operators close on the set above under renormalization.

We find that the following elements of the mixing matrix are compatible with both the two-gluon and two-ghost projections:

$$\begin{aligned}
Z_{GG} &= 1 + O(g^3), \\
Z_{GA} &= -\frac{1}{2} \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A + O(g^3), \\
Z_{GE} &= \frac{1}{2} \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A + O(g^3), \\
Z_{AG} &= O(g^3), \\
Z_{AA} &= 1 + \frac{1}{2} \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A + O(g^3), \\
Z_{AE} &= -\frac{1}{2} \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A + O(g^3), \\
Z_{EE} &= 1 + O(g^2).
\end{aligned} \tag{4.19}$$

The renormalization mixing matrix is triangular to  $O(g^2)$ . We do not calculate  $Z_{EE}$  beyond the tree level explicitly in this paper, but there is a prediction from theory that  $Z_{EE} \equiv 1$ , with no higher order corrections. In other words, the operators of class  $E$  are finite.

#### D. BRST Ward identity (Slavnov-Taylor identity)

We have seen that the gauge-variant part of the energy-momentum tensor is nonzero in an on-shell matrix element. However, the gauge-variant part of the energy-momentum tensor is BRST exact, in accordance with general theory and a very simple proof states that physical matrix elements of such operators vanish [13].

In this section, we resolve the contradiction. The proof that physical matrix elements of BRST-exact operators vanish proceeds by using a Ward identity based on the BRST variation to relate the matrix element under study to a particular Green function of the ancestor operator of the BRST-exact operator. This Green function has a manifest factor of zero when put on shell, but the zero is compensated by a quadratic IR divergence present only when the matrix element of the ancestor operator is evaluated at zero momentum transfer, as we will now see.

We have verified by explicit calculation of all graphs that the BRST Ward identity still holds for unamputated Green functions off mass shell, calculated at zero momentum transfer.

The BRST variations of the bare fields are given in Eq. (3.7). We need the BRST variation of the renormalized fields in terms of renormalized fields and parameters. This is sometimes referred to as the ‘‘renormalized BRST variation’’ and is related to the canonical BRST variation by factors of the renormalization constants, Eq. (3.21). The goal in defining a renormalized BRST variation is to obtain UV-finite Green functions with renormalized fields. The renormalized constant Grassmann parameter which accomplishes this goal [13] is

$$\delta\xi \equiv Z_A^{-\frac{1}{2}} Z_\eta^{-\frac{1}{2}} \widehat{\delta\xi}, \tag{4.20}$$

so we have

$$\begin{aligned}
\delta_{\text{BRST}} A_{\mu a} &= \left( Z_\omega^{\frac{1}{2}} Z_A^{-\frac{1}{2}} \partial_\mu \omega_a - Z_g Z_\omega^{\frac{1}{2}} g c_{abc} A_{\mu b} \omega_c \right) \widehat{\delta\xi}, \\
\delta_{\text{BRST}} \omega_a &= -\frac{1}{2} Z_g Z_\omega^{\frac{1}{2}} g c_{abc} \omega_b \omega_c \widehat{\delta\xi}, \\
\delta_{\text{BRST}} \eta_a &= Z_\lambda Z_A^{\frac{1}{2}} Z_\eta^{-\frac{1}{2}} \lambda \partial \cdot A_a \widehat{\delta\xi},
\end{aligned} \tag{4.21}$$

or, in terms of the renormalized BRST variation,

$$\begin{aligned}
\frac{\delta_{\text{BRST}}^r}{\delta\xi} A_{\mu a} &= Z_0 \partial_\mu \omega_a - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{abc} A_{\mu b} \omega_c, \\
\frac{\delta_{\text{BRST}}^r}{\delta\xi} \omega_a &= -\frac{1}{2} Z_g Z_0 Z_A^{\frac{1}{2}} g c_{abc} \omega_b \omega_c, \\
\frac{\delta_{\text{BRST}}^r}{\delta\xi} \eta_a &= \lambda \partial \cdot A_a.
\end{aligned} \tag{4.22}$$

The three operators  $\frac{\delta_{\text{BRST}}^r}{\delta\xi} A_{\mu a}$ ,  $\frac{\delta_{\text{BRST}}^r}{\delta\xi} \omega_a$ , and  $\frac{\delta_{\text{BRST}}^r}{\delta\xi} \eta_a$  are all finite [13].

The ancestor operator of the renormalized BRST variation is

$$\begin{aligned}
X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) &= (\partial_\nu \eta_a) A_{\mu a} + (\partial_\mu \eta_a) A_{\nu a} \\
&\quad - g_{\mu\nu} \left[ \frac{1}{2} \eta_a \partial \cdot A_a + (\partial_\rho \eta_a) A_a^\rho \right],
\end{aligned} \tag{4.23}$$

where

$$\frac{\delta_{\text{BRST}}^r}{\delta\xi} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) = \theta_{\mu\nu}^{(\text{GV})} - \frac{1}{2} g_{\mu\nu} \hat{\eta}_a \partial^\rho (\hat{D}_\rho \hat{\omega})_a. \tag{4.24}$$

Remember that the fields in the energy-momentum tensor operator are bare. The ancestor of the unrenormalized BRST variation Eq. (3.14) and the ancestor defined above are related by

$$\hat{X}_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) = Z_A^{\frac{1}{2}} Z_\eta^{\frac{1}{2}} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right). \tag{4.25}$$

Now, the BRST variation of any Green function vanishes. Consider the particular case

$$\delta_{\text{BRST}} \langle 0 | T A_{\sigma a} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) A_{\tau b} | 0 \rangle = 0. \tag{4.26}$$

This gives

$$\begin{aligned}
0 &= \langle 0 | T [\delta_{\text{BRST}} A_{\sigma a}] X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) A_{\tau b} | 0 \rangle \\
&\quad + \langle 0 | T A_{\sigma a} [\delta_{\text{BRST}} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right)] A_{\tau b} | 0 \rangle \\
&\quad + \langle 0 | T A_{\sigma a} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(\text{GV})} \right) [\delta_{\text{BRST}} A_{\tau b}] | 0 \rangle.
\end{aligned} \tag{4.27}$$

The vanishing of Eq. (4.26) and the chain rule for the BRST variation can be proven by defining the variation in terms of (anti)commutators with the Noether charge associated with the BRST symmetry [19].

Equation (4.27) becomes

$$\begin{aligned}
 0 = & \left\langle 0 \left| T \left[ \left( Z_0 \partial_\sigma \omega_a - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{ade} A_\sigma d\omega_e \right) \delta\xi \right] X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right) A_{\tau b} \right| 0 \right\rangle \\
 & + \left\langle 0 \left| T A_{\sigma a} \left[ \theta_{\mu\nu}^{(GV)} \delta\xi \right] A_{\tau b} \right| 0 \right\rangle + \left\langle 0 \left| T A_{\sigma a} \left[ -\frac{1}{2} g_{\mu\nu} \hat{\eta}_c \partial^\rho (\hat{D}_\rho \hat{\omega})_c \delta\xi \right] A_{\tau b} \right| 0 \right\rangle \\
 & + \left\langle 0 \left| T A_{\sigma a} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right) \left[ \left( Z_0 \partial_\tau \omega_b - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{bde} A_\tau d\omega_e \right) \delta\xi \right] \right| 0 \right\rangle .
 \end{aligned}
 \tag{4.28}$$

The constant Grassmann parameter  $\delta\xi$  can be factored out of Eq. (4.28) above if it is anticommutated through the ancestor operator which has a Grassmann parity of 1 (because each term contains one antighost field). This is responsible for the relative minus sign below:

$$\begin{aligned}
 0 = & (-1) \left\langle 0 \left| T \left[ Z_0 \partial_\sigma \omega_a - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{ade} A_\sigma d\omega_e \right] X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right) A_{\tau b} \right| 0 \right\rangle \\
 & + \left\langle 0 \left| T A_{\sigma a} \theta_{\mu\nu}^{(GV)} A_{\tau b} \right| 0 \right\rangle + \left\langle 0 \left| T A_{\sigma a} \left[ -\frac{1}{2} g_{\mu\nu} \hat{\eta}_c \partial^\rho (\hat{D}_\rho \hat{\omega})_c \right] A_{\tau b} \right| 0 \right\rangle \\
 & + \left\langle 0 \left| T A_{\sigma a} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right) \left[ Z_0 \partial_\tau \omega_b - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{bde} A_\tau d\omega_e \right] \right| 0 \right\rangle .
 \end{aligned}
 \tag{4.29}$$

One must also keep in mind that the order of the Grassmann fields,  $\hat{\eta}$  (in the ancestor operator) and  $\hat{\omega}$ , in the last two lines above is opposite to the canonical ordering and that

$$\langle 0 | T \hat{\eta} \hat{\omega} (\text{operator}) | 0 \rangle = - \langle 0 | T \hat{\omega} \hat{\eta} (\text{operator}) | 0 \rangle .
 \tag{4.30}$$

It is now obvious that there exists an alternate calculation which will provide the two-gluon physical matrix element of  $\theta_{\mu\nu}^{(GV)}$ :

$$\begin{aligned}
 \left\langle 0 \left| T A_{\sigma a} \theta_{\mu\nu}^{(GV)} A_{\tau b} \right| 0 \right\rangle = & \left\langle 0 \left| T \left[ Z_0 \partial_\sigma \omega_a - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{ade} A_\sigma d\omega_e \right] X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right) A_{\tau b} \right| 0 \right\rangle \\
 & - \left\langle 0 \left| T A_{\sigma a} \left[ -\frac{1}{2} g_{\mu\nu} \hat{\eta}_c \partial^\rho (\hat{D}_\rho \hat{\omega})_c \right] A_{\tau b} \right| 0 \right\rangle \\
 & - \left\langle 0 \left| T A_{\sigma a} X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right) \left[ Z_0 \partial_\tau \omega_b - Z_g Z_0 Z_A^{\frac{1}{2}} g c_{bde} A_\tau d\omega_e \right] \right| 0 \right\rangle .
 \end{aligned}
 \tag{4.31}$$




FIG. 5. Born and one-loop diagrams involved in the BRST Ward identity that vanish in physical matrix elements. The symbol  $\otimes$  represents the inserted composite operator  $X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right)$ , the ancestor of the  $\theta_{\mu\nu}^{(GV)}$  under the renormalized BRST variation.

The Feynman graphs associated with the Green functions on the right-hand side of the equation above are displayed in Figs. 5 and 6. The usual proof that the matrix element  $\langle 0 | T A_{\sigma a} \theta_{\mu\nu}^{(GV)} A_{\tau b} | 0 \rangle$  vanishes on shell relies on the assumption that the triangle graphs depicted in Fig. 6 do not contain quadratic IR singularities. These




FIG. 6. The one-loop diagrams involved in the BRST Ward identity that do not vanish in physical matrix elements due to an infrared divergence. The symbol  $\otimes$  represents the inserted composite operator  $X_{\text{ancestor}} \left( \theta_{\mu\nu}^{(GV)} \right)$ .

graphs contain an unusual vertex, one gluon and one ghost at the same space-time point. There is no external line, therefore nothing to amputate, but the amputation procedure for other graphs, which have ghost legs, requires that all diagrams be multiplied by the inverse of a ghost propagator, which, of course, is proportional to  $p^2$ . If the diagrams contain at worst logarithmic IR divergences, then the additional factor of  $p^2$  will cause them to vanish on shell. The diagrams with an external ghost line have a derivative acting on the ghost field. In momentum space, the derivative becomes a factor of  $p^\sigma$  which gives zero when contracted with the physical polarization vector  $\epsilon_\sigma$  associated with the gluon leg. Only the graphs in Fig. 6 contribute to on-shell matrix elements.

Explicit calculation of the graphs in Fig. 6, with the result Eq. (A9), shows that there are, in fact,  $\frac{1}{p^2}$  divergences which cancel against the inverse of the ghost propagators introduced in the amputation procedure, as we claimed at the beginning of this section. These IR poles occur only at zero momentum transfer. The Feynman rules for the inserted operator vertices are given in Figs. 7 and 8.

V. CONCLUSION

We have seen by explicit calculation that one of the central results of Joglekar and Lee, that physical matrix elements of BRST-exact operators must vanish, fails at one-loop order. We give a form for the BRST ancestors of the alien operators required in the renormalization of the covariant gluon operator. Our alien operators are then manifestly BRST exact whereas the basis of alien operators proposed by Dixon and Taylor (those used by Hamberg and van Neerven) are not BRST exact. The Dixon and Taylor set of alien operators are not guaranteed to vanish in physical matrix elements.

We have verified the predictions of Freedman *et al.* on the finiteness of the energy-momentum tensor in both gluon and ghost two-point functions to one-loop order at zero momentum transfer by evaluating diagrams with a BRST-exact alien operator insertion.

The BRST Ward identity demonstrates where the proof of the Joglekar-Lee theorem breaks down. Taking the momentum transfer to zero too soon introduces spurious IR divergences which cancel factors of zero on which the proof relies. Calculations performed using the Dixon and Taylor set of alien operators cannot be analyzed through the BRST Ward identity.

The physical region of interest in almost all calculations involving the renormalization of composite operators, such as the calculations required in the operator product expansion, is the exceptional point of zero momentum transfer. To expedite the computation, one sets the momentum transfer to zero at the very beginning, thereby eliminating one scale from the problem. In some calculations involving final-state cuts, it is not clear how one would generalize to nonzero momentum transfer.

The alternative is to keep the momentum transfer arbitrary until after the Feynman graphs have been evaluated, and only then to set the momentum transfer to zero.




FIG. 7. Feynman rules for nonstandard two-point vertices at zero momentum transfer.

With this procedure, the Joglekar-Lee theorem should apply, making it unnecessary to compute the graphs containing the alien operator insertion. The price to be paid, of course, is the introduction of another momentum scale into the problem and a corresponding increase in the complexity and volume of the analysis.

There has been a sense of disquiet in the literature about zero momentum transfer for a long time. Joglekar mentions in the concluding section of [24] that, at the exceptional momentum point  $Q = 0$ , matrix elements of gauge-invariant operators lose some of the properties that make them manageable at nonzero momentum trans-




FIG. 8. Feynman rules for nonstandard vertices other than two-point at arbitrary momentum transfer.

fer. Lee [25] works with the twist-2 piece of the energy-momentum tensor to show that certain pieces of the calculation at zero momentum transfer can yield useful information, that is, the coefficients of certain terms are the same, independent of the momentum transfer. He calculates only the pole terms of the two-gluon Green function at one-loop order at both zero and nonzero momentum transfer. The unease was certainly justified; some results hold while others fail utterly. It is not unreasonable to question all calculations performed when the limit of zero momentum transfer was applied initially, and such calculations are the mainstay of perturbative QCD.

## ACKNOWLEDGMENTS

We would like to thank Jack Smith and Brian Harris of SUNY Stony Brook and Willie van Neerven of the University of Leiden, the Netherlands, for enlightening discussions. In addition, thanks are due Laura Weinkauff of Pennsylvania State University for technical help with the combinatorics of the diagrams and the FORM code.

### a. Entire energy-momentum tensor

The two-gluon amputated Green function of the entire energy-momentum tensor is

$$\begin{aligned}
\langle 0|T A_\sigma{}_a \theta_{\mu\nu} A_\tau{}_b|0\rangle_{\text{amputated}} &= \delta_{ab} [p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) \\
&\quad - 2p_\mu p_\nu g_{\sigma\tau} + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu} \\
&\quad - \lambda (p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu})] \\
&\quad + \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( -\frac{1}{4} \left(1 - \frac{1}{\lambda}\right)^2 \left[ p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) - 2p_\mu p_\nu g_{\sigma\tau} \right. \right. \\
&\quad \left. \left. + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu} \right] \right. \\
&\quad \left. + \left(1 - \frac{1}{\lambda}\right) \left\{ \frac{p_\sigma p_\tau p_\mu p_\nu}{p^2} - \left[ 3 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p_\mu p_\nu g_{\sigma\tau} \right. \right. \\
&\quad \left. \left. + \left[ 1 + \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] [p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) \right. \right. \right. \\
&\quad \left. \left. \left. + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu} \right] \right\} \right. \\
&\quad \left. + \frac{10}{3} \frac{p_\sigma p_\tau p_\mu p_\nu}{p^2} + \left[ \frac{32}{9} - \frac{10}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p_\mu p_\nu g_{\sigma\tau} \right. \\
&\quad \left. + \left[ -\frac{31}{9} + \frac{5}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] [p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} \right. \right. \\
&\quad \left. \left. + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu}] \right) + O(g^4). \tag{A2}
\end{aligned}$$

To make the comparison with Hamberg and van Neerven more transparent, we present the twist-2 (spin-2) piece of this amputated Green function in which the free Lorentz indices of the inserted operator,  $\mu$  and  $\nu$ , are contracted with a null vector  $\Delta$ :

## APPENDIX

### 1. Right derivatives

Right derivatives [26] are such that

$$\frac{\partial^r (XY)}{\partial Z} = X \frac{\partial^r Y}{\partial Z} + (-1)^{P_Y P_Z} \left( \frac{\partial^r X}{\partial Z} \right) Y, \tag{A1}$$

where  $P_Y$  is the ‘‘Grassmann parity’’ of the quantity  $Y$ . The (anti)ghost field components have Grassmann parity 1, while the  $c$ -number parameters and boson field components have Grassmann parity 0. Fermion field components would be assigned Grassmann parity 1.

### 2. Complete off-shell calculations for two-gluon amputated Green functions

In this section of the Appendix, we give the full Lorentz structure for the amputated Green functions of the energy-momentum tensor operators with two gluon fields off mass shell, at zero momentum transfer, to one-loop order. The  $\frac{1}{\epsilon}$  poles are purely UV divergences while all of the IR divergences (as  $p^2 \rightarrow 0$ ) are seen as logarithms.

$$\begin{aligned}
\langle 0|T A_\sigma a \Delta^\mu \theta_{\mu\nu} \Delta^\nu A_\tau b|0\rangle_{\text{amputated}} &= 2\delta_{ab}[-p^2 \Delta_\sigma \Delta_\tau - (p \cdot \Delta)^2 g_{\sigma\tau} + (1-\lambda)(p \cdot \Delta)(p_\sigma \Delta_\tau + p_\tau \Delta_\sigma)] + \frac{g^2}{16\pi^2} C_A \delta_{ab} \\
&\times \left( -\frac{1}{2} \left(1 - \frac{1}{\lambda}\right)^2 [-p^2 \Delta_\sigma \Delta_\tau - (p \cdot \Delta)^2 g_{\sigma\tau} + (p \cdot \Delta)(p_\sigma \Delta_\tau + p_\tau \Delta_\sigma)] \right. \\
&+ \left(1 - \frac{1}{\lambda}\right) \left\{ \frac{p_\sigma p_\tau (p \cdot \Delta)^2}{p^2} - \left[3 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] (p \cdot \Delta)^2 g_{\sigma\tau} \right. \\
&+ \left. \left[2 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] [-p^2 \Delta_\sigma \Delta_\tau + (p \cdot \Delta)p_\sigma \Delta_\tau + (p \cdot \Delta)p_\tau \Delta_\sigma] \right\} \\
&+ \frac{10}{3} \frac{p_\sigma p_\tau (p \cdot \Delta)^2}{p^2} + \left[ \frac{32}{9} - \frac{10}{3} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] (p \cdot \Delta)^2 g_{\sigma\tau} \\
&+ \left. \left[ -\frac{62}{9} + \frac{10}{3} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] [-p^2 \Delta_\sigma \Delta_\tau + (p \cdot \Delta)(p_\sigma \Delta_\tau + p_\tau \Delta_\sigma)] \right) + O(g^4). \quad (\text{A3})
\end{aligned}$$

### b. Gauge-invariant part

The two-gluon amputated Green function of the gauge-invariant piece of the energy-momentum tensor is

$$\begin{aligned}
\langle 0|T A_\sigma a \theta_{\mu\nu}^{(G1)} A_\tau b|0\rangle_{\text{amputated}} &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} [p^2 (-\frac{1}{2} g_{\sigma\tau} g_{\mu\nu} + g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) \\
&- \frac{1}{2} (p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu})] \\
&+ \delta_{ab} [p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) - 2p_\mu p_\nu g_{\sigma\tau} + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} \\
&+ p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu}] \\
&+ \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( -\frac{1}{2} \left(1 - \frac{1}{\lambda}\right)^2 \left[ p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) \right. \right. \\
&- \left. \left. 2p_\mu p_\nu g_{\sigma\tau} + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu} - p_\sigma p_\tau g_{\mu\nu} \right] \right. \\
&+ \left(1 - \frac{1}{\lambda}\right) \left\{ 2 \frac{p_\sigma p_\tau p_\mu p_\nu}{p^2} - \left[6 + \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right] p_\mu p_\nu g_{\sigma\tau} \right. \\
&+ \left[ \frac{5}{2} + \frac{1}{2} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] (p^2 g_{\sigma\tau} g_{\mu\nu} - p_\sigma p_\tau g_{\mu\nu}) \\
&+ \left. \left[ 2 + \frac{1}{2} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] [-p^2 (g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) + p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu}] \right\} \\
&- \frac{2}{3} \frac{p_\sigma p_\tau p_\mu p_\nu}{p^2} + \left[ \frac{86}{9} - \frac{10}{3} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] p_\mu p_\nu g_{\sigma\tau} \\
&+ \left[ -\frac{58}{9} + \frac{13}{6} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] (p^2 g_{\sigma\tau} g_{\mu\nu} - p_\sigma p_\tau g_{\mu\nu}) \\
&+ \left[ \frac{49}{9} - \frac{8}{3} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] p^2 (g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) \\
&+ \left. \left[ -\frac{89}{18} + \frac{13}{6} \ln\left(\frac{-p^2}{\bar{\mu}^2}\right) \right] (p_\tau p_\nu g_{\sigma\mu} + p_\tau p_\mu g_{\sigma\nu} + p_\sigma p_\nu g_{\tau\mu} + p_\sigma p_\mu g_{\tau\nu}) \right) + O(g^4). \quad (\text{A4})
\end{aligned}$$

The twist-2 (spin-2) piece of this amputated Green function in which the free Lorentz indices of the inserted operator,  $\mu$  and  $\nu$ , are contracted with a null vector  $\Delta$  is



$$\begin{aligned}
\langle 0|T A_{\sigma a} \Delta^{\mu} \theta_{\mu\nu}^{(GI)} \Delta^{\nu} A_{\tau b}|0\rangle_{\text{amputated}} &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} [2p^2 \Delta_{\sigma} \Delta_{\tau} - (p \cdot \Delta)(p_{\sigma} \Delta_{\tau} + p_{\tau} \Delta_{\sigma})] \\
&+ 2\delta_{ab} [-p^2 \Delta_{\sigma} \Delta_{\tau} - (p \cdot \Delta)^2 g_{\sigma\tau} + (p \cdot \Delta)(p_{\sigma} \Delta_{\tau} + p_{\tau} \Delta_{\sigma})] \\
&+ \frac{g^2}{16\pi^2} C_A \delta_{ab} \left( - \left(1 - \frac{1}{\lambda}\right)^2 \left[ -p^2 \Delta_{\sigma} \Delta_{\tau} - (p \cdot \Delta)^2 g_{\sigma\tau} + (p \cdot \Delta)(p_{\sigma} \Delta_{\tau} + p_{\tau} \Delta_{\sigma}) \right] \right. \\
&+ \left(1 - \frac{1}{\lambda}\right) \left\{ 2 \frac{p_{\sigma} p_{\tau} (p \cdot \Delta)^2}{p^2} - \left[ 6 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] (p \cdot \Delta)^2 g_{\sigma\tau} \right. \\
&+ \left. \left[ 4 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] [-p^2 \Delta_{\sigma} \Delta_{\tau} + (p \cdot \Delta) p_{\sigma} \Delta_{\tau} + (p \cdot \Delta) p_{\tau} \Delta_{\sigma}] \right\} \\
&- \frac{2}{3} \frac{p_{\sigma} p_{\tau} (p \cdot \Delta)^2}{p^2} + \left[ \frac{86}{9} - \frac{10}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] (p \cdot \Delta)^2 g_{\sigma\tau} \\
&+ \left[ \frac{98}{9} - \frac{16}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p^2 \Delta_{\sigma} \Delta_{\tau} \\
&+ \left. \left[ -\frac{89}{9} + \frac{13}{3} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] (p \cdot \Delta)(p_{\sigma} \Delta_{\tau} + p_{\tau} \Delta_{\sigma}) \right) + O(g^4). \tag{A5}
\end{aligned}$$

*c. Gauge-variant (alien) part*

The two-gluon amputated Green function of the gauge-variant piece of the energy-momentum tensor is

$$\begin{aligned}
\langle 0|T A_{\sigma a} \theta_{\mu\nu}^{(GV)} A_{\tau b}|0\rangle_{\text{amputated}} &= -\frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} [p^2 (-\frac{1}{2} g_{\sigma\tau} g_{\mu\nu} + g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) \\
&- \frac{1}{2} (p_{\tau} p_{\nu} g_{\sigma\mu} + p_{\tau} p_{\mu} g_{\sigma\nu} + p_{\sigma} p_{\nu} g_{\tau\mu} + p_{\sigma} p_{\mu} g_{\tau\nu} - p_{\sigma} p_{\tau} g_{\mu\nu})] \\
&- \lambda \delta_{ab} (p_{\tau} p_{\nu} g_{\sigma\mu} + p_{\tau} p_{\mu} g_{\sigma\nu} + p_{\sigma} p_{\nu} g_{\tau\mu} + p_{\sigma} p_{\mu} g_{\tau\nu} - p_{\sigma} p_{\tau} g_{\mu\nu}) \\
&+ \frac{g^2}{16\pi^2} C_A \delta_{ab} \left\{ \frac{1}{4} \left(1 - \frac{1}{\lambda}\right)^2 \left[ p^2 (g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu}) \right. \right. \\
&- 2p_{\mu} p_{\nu} g_{\sigma\tau} + p_{\tau} p_{\nu} g_{\sigma\mu} + p_{\tau} p_{\mu} g_{\sigma\nu} + p_{\sigma} p_{\nu} g_{\tau\mu} + p_{\sigma} p_{\mu} g_{\tau\nu} - p_{\sigma} p_{\tau} g_{\mu\nu} \left. \right] \\
&+ \left(1 - \frac{1}{\lambda}\right) \left[ -\frac{p_{\sigma} p_{\tau} p_{\mu} p_{\nu}}{p^2} + p^2 \left( -\frac{3}{2} g_{\sigma\tau} g_{\mu\nu} + g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu} \right) \right. \\
&+ \left. 3p_{\mu} p_{\nu} g_{\sigma\tau} - p_{\tau} p_{\nu} g_{\sigma\mu} - p_{\tau} p_{\mu} g_{\sigma\nu} - p_{\sigma} p_{\nu} g_{\tau\mu} - p_{\sigma} p_{\mu} g_{\tau\nu} + \frac{3}{2} p_{\sigma} p_{\tau} g_{\mu\nu} \right] \\
&+ 4 \frac{p_{\sigma} p_{\tau} p_{\mu} p_{\nu}}{p^2} - 6p_{\mu} p_{\nu} g_{\sigma\tau} + \left[ 3 - \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] (p^2 g_{\sigma\tau} g_{\mu\nu} - p_{\sigma} p_{\tau} g_{\mu\nu}) \\
&+ \left[ -2 + \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] p^2 (g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) \\
&+ \left. \left[ \frac{3}{2} - \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] (p_{\tau} p_{\nu} g_{\sigma\mu} + p_{\tau} p_{\mu} g_{\sigma\nu} + p_{\sigma} p_{\nu} g_{\tau\mu} + p_{\sigma} p_{\mu} g_{\tau\nu}) \right\} + O(g^4). \tag{A6}
\end{aligned}$$

The twist-2 (spin-2) piece of this amputated Green function in which the free Lorentz indices of the inserted operator,  $\mu$  and  $\nu$ , are contracted with a null vector  $\Delta$  is

$$\begin{aligned}
\langle 0|TA_{\sigma a}\Delta^{\mu}\theta_{\mu\nu}^{(GV)}\Delta^{\nu}A_{\tau b}|0\rangle_{\text{amputated}} &= -\frac{1}{\epsilon}\frac{g^2}{16\pi^2}C_A\delta_{ab}[2p^2\Delta_{\sigma}\Delta_{\tau} - (p\cdot\Delta)(p_{\sigma}\Delta_{\tau} + p_{\tau}\Delta_{\sigma})] \\
&\quad -2\lambda\delta_{ab}(p\cdot\Delta)(p_{\sigma}\Delta_{\tau} + p_{\tau}\Delta_{\sigma}) \\
&\quad +\frac{g^2}{16\pi^2}C_A\delta_{ab}\left\{\frac{1}{2}\left(1-\frac{1}{\lambda}\right)^2[-p^2\Delta_{\sigma}\Delta_{\tau} - (p\cdot\Delta)^2g_{\sigma\tau} + (p\cdot\Delta)(p_{\sigma}\Delta_{\tau} + p_{\tau}\Delta_{\sigma})]\right. \\
&\quad +\left(1-\frac{1}{\lambda}\right)\left(-\frac{p_{\sigma}p_{\tau}(p\cdot\Delta)^2}{p^2} + 2p^2\Delta_{\sigma}\Delta_{\tau} + 3(p\cdot\Delta)^2g_{\sigma\tau}\right. \\
&\quad \left.\left.-2(p\cdot\Delta)(p_{\sigma}\Delta_{\tau} + p_{\tau}\Delta_{\sigma})\right)\right. \\
&\quad \left.+4\frac{p_{\sigma}p_{\tau}(p\cdot\Delta)^2}{p^2} - 6(p\cdot\Delta)^2g_{\sigma\tau} + \left[-4 + 2\ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right]p^2\Delta_{\sigma}\Delta_{\tau}\right. \\
&\quad \left.+ \left[3 - \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right](p\cdot\Delta)(p_{\sigma}\Delta_{\tau} + p_{\tau}\Delta_{\sigma})\right\} + O(g^4). \tag{A7}
\end{aligned}$$

### 3. Unamputated two-gluon Green function of alien piece of the energy-momentum tensor off shell

In order to use the BRST Ward identity (which is valid off mass shell), we need the unamputated Green function which we obtain from the amputated Green function, Eq. (A6), by attaching the dressed external gluon propagators, Eq. (3.23a):<sup>13</sup>

$$\begin{aligned}
\langle 0|TA_{\sigma a}\theta_{\mu\nu}^{(GV)}A_{\tau b}|0\rangle &= \mathcal{D}_{ab'}^{\sigma\tau'}(p)\langle 0|TA_{\sigma' a'}\theta_{\mu\nu}^{(GV)}A_{\tau' b'}|0\rangle_{\text{amputated}}\mathcal{D}_{a'b}^{\sigma'\tau}(p) \\
&= \frac{1}{\epsilon}\frac{g^2}{16\pi^2}C_A\delta_{ab}\frac{1}{(p^2)^2}\left[\left(1-\frac{1}{\lambda}\right)\left(2\frac{p_{\sigma}p_{\tau}p_{\mu}p_{\nu}}{p^2}\right.\right. \\
&\quad \left.\left.-\frac{1}{2}(p_{\tau}p_{\nu}g_{\sigma\mu} + p_{\tau}p_{\mu}g_{\sigma\nu} + p_{\sigma}p_{\nu}g_{\tau\mu} + p_{\sigma}p_{\mu}g_{\tau\nu})\right) + p^2\left(-\frac{1}{2}g_{\sigma\tau}g_{\mu\nu} + g_{\sigma\mu}g_{\tau\nu} + g_{\sigma\nu}g_{\tau\mu}\right)\right. \\
&\quad \left.-\frac{1}{2}\left(p_{\tau}p_{\nu}g_{\sigma\mu} + p_{\tau}p_{\mu}g_{\sigma\nu} + p_{\sigma}p_{\nu}g_{\tau\mu} + p_{\sigma}p_{\mu}g_{\tau\nu} - p_{\sigma}p_{\tau}g_{\mu\nu}\right)\right] \\
&\quad +\delta_{ab}\frac{1}{(p^2)^2}\left(-4\left(1-\frac{1}{\lambda}\right)\frac{p_{\sigma}p_{\tau}p_{\mu}p_{\nu}}{p^2} + p_{\tau}p_{\nu}g_{\sigma\mu} + p_{\tau}p_{\mu}g_{\sigma\nu} + p_{\sigma}p_{\nu}g_{\tau\mu} + p_{\sigma}p_{\mu}g_{\tau\nu} - p_{\sigma}p_{\tau}g_{\mu\nu}\right) \\
&\quad +\frac{g^2}{16\pi^2}C_A\delta_{ab}\frac{1}{(p^2)^2}\left\{\left(1-\frac{1}{\lambda}\right)^2\left[-\frac{p_{\sigma}p_{\tau}p_{\mu}p_{\nu}}{p^2}\right.\right. \\
&\quad \left.\left.+\frac{1}{4}p^2(-g_{\sigma\tau}g_{\mu\nu} + g_{\sigma\mu}g_{\tau\nu} + g_{\sigma\nu}g_{\tau\mu}) + \frac{1}{2}p_{\mu}p_{\nu}g_{\sigma\tau} + \frac{1}{4}p_{\sigma}p_{\tau}g_{\mu\nu}\right]\right. \\
&\quad \left.+\left(1-\frac{1}{\lambda}\right)\left[7\frac{p_{\sigma}p_{\tau}p_{\mu}p_{\nu}}{p^2} - p^2\left(-\frac{3}{2}g_{\sigma\tau}g_{\mu\nu} + g_{\sigma\mu}g_{\tau\nu} + g_{\sigma\nu}g_{\tau\mu}\right)\right.\right. \\
&\quad \left.\left.-\frac{1}{2}(p_{\tau}p_{\nu}g_{\sigma\mu} + p_{\tau}p_{\mu}g_{\sigma\nu} + p_{\sigma}p_{\nu}g_{\tau\mu} + p_{\sigma}p_{\mu}g_{\tau\nu}) - 3p_{\mu}p_{\nu}g_{\sigma\tau} - \frac{3}{2}p_{\sigma}p_{\tau}g_{\mu\nu}\right]\right. \\
&\quad \left.+\left[-\frac{160}{9} + \frac{20}{3}\ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right]\frac{p_{\sigma}p_{\tau}p_{\mu}p_{\nu}}{p^2} + \left[-3 + \frac{1}{2}\ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right](p^2g_{\sigma\tau}g_{\mu\nu} - p_{\sigma}p_{\tau}g_{\mu\nu})\right. \\
&\quad \left.+\left[2 - \ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right]p^2(g_{\sigma\mu}g_{\tau\nu} + g_{\sigma\nu}g_{\tau\mu}) + 6p_{\mu}p_{\nu}g_{\sigma\tau}\right. \\
&\quad \left.+\left[\frac{35}{18} - \frac{7}{6}\ln\left(\frac{-p^2}{\bar{\mu}^2}\right)\right](p_{\tau}p_{\nu}g_{\sigma\mu} + p_{\tau}p_{\mu}g_{\sigma\nu} + p_{\sigma}p_{\nu}g_{\tau\mu} + p_{\sigma}p_{\mu}g_{\tau\nu})\right\} + O(g^4). \tag{A8}
\end{aligned}$$

<sup>13</sup>The modified LSZ prescription, rather than the dressed external propagators, would have been used to go on shell.

## 4. BRST Ward identity graphs evaluated

The quadratically IR-divergent Feynman graphs used in the BRST Ward identity are evaluated as follows:

$$\begin{aligned}
(\text{Fig. 6}) &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} i \left( \frac{1}{2} \right) \left( -\frac{1}{2} g_{\sigma\tau} g_{\mu\nu} + g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu} \right) \\
&+ \frac{g^2}{16\pi^2} C_A \delta_{ab} i \left\{ \frac{1}{4} \left( 1 - \frac{1}{\lambda} \right)^2 \left[ \frac{p_\sigma p_\tau p_\mu p_\nu}{(p^2)^2} + \frac{1}{2} (-g_{\sigma\tau} g_{\mu\nu} + g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) \right. \right. \\
&+ \left. \frac{p_\mu p_\nu}{p^2} g_{\sigma\tau} - \frac{1}{2} \frac{p_\tau p_\nu}{p^2} g_{\sigma\mu} - \frac{1}{2} \frac{p_\tau p_\mu}{p^2} g_{\sigma\nu} - \left( \frac{p_\sigma p_\nu}{p^2} g_{\tau\mu} \frac{p_\sigma p_\mu}{p^2} g_{\tau\nu} - \frac{1}{2} \frac{p_\sigma p_\tau}{p^2} g_{\mu\nu} \right) \right] \\
&+ \frac{1}{2} \left( 1 - \frac{1}{\lambda} \right) \left[ \frac{p_\sigma p_\tau p_\mu p_\nu}{(p^2)^2} + \frac{3}{2} g_{\sigma\tau} g_{\mu\nu} - g_{\sigma\mu} g_{\tau\nu} - g_{\sigma\nu} g_{\tau\mu} - 3 \frac{p_\mu p_\nu}{p^2} g_{\sigma\tau} \right. \\
&+ \left. \frac{p_\tau p_\nu}{p^2} g_{\sigma\mu} \frac{p_\tau p_\mu}{p^2} g_{\sigma\nu} + \frac{3}{2} \left( \frac{p_\sigma p_\nu}{p^2} g_{\tau\mu} \frac{p_\sigma p_\mu}{p^2} g_{\tau\nu} \right) - \frac{p_\sigma p_\tau}{p^2} g_{\mu\nu} \right] \\
&+ \left[ -\frac{3}{2} + \frac{1}{4} \ln \left( \frac{-p^2}{\bar{\mu}^2} g_{\sigma\tau} g_{\mu\nu} + \left[ 1 - \frac{1}{2} \ln \frac{-p^2}{\bar{\mu}^2} \right] \right) \right] (g_{\sigma\mu} g_{\tau\nu} + g_{\sigma\nu} g_{\tau\mu}) \\
&+ \left. 3 \frac{p_\mu p_\nu}{p^2} g_{\sigma\tau} - \frac{3}{2} \left( \frac{p_\sigma p_\nu}{p^2} g_{\tau\mu} + \frac{p_\sigma p_\mu}{p^2} g_{\tau\nu} \right) - \frac{1}{2} \left( \frac{p_\tau p_\nu}{p^2} g_{\sigma\mu} + \frac{p_\tau p_\mu}{p^2} g_{\sigma\nu} \right) + \frac{p_\sigma p_\tau}{p^2} g_{\mu\nu} \right\} + O(g^4). \quad (\text{A9})
\end{aligned}$$

The terms proportional to  $\frac{p_\mu p_\nu}{p^2} g_{\sigma\tau}$  in the finite part<sup>14</sup> ruin the proof that a physical matrix element of an alien operator must vanish.

The graphs in Fig. 6 plus their mirror images ( $\sigma \leftrightarrow \tau$ ,  $a \leftrightarrow b$ ,  $p \leftrightarrow -p$ ) contribute to the two amputated Green functions,

$$\begin{aligned}
&- \left\langle 0 \left| T \left[ Z_g Z_0 Z_A^{\frac{1}{2}} g_{cde} A_\sigma d\omega_e \right] \text{ancestor} \left( \theta_{\mu\nu}^{(GV)} \right) A_{\tau b} \right| 0 \right\rangle_{\text{amputated}}, \\
&+ \left\langle 0 \left| T A_{\sigma a} \text{ancestor} \left( \theta_{\mu\nu}^{(GV)} \right) \left[ Z_g Z_0 Z_A^{\frac{1}{2}} g_{cde} A_\tau d\omega_e \right] \right| 0 \right\rangle_{\text{amputated}}, \quad (\text{A10})
\end{aligned}$$

corresponding to the unamputated Green functions found in Eq. (4.30).

To isolate the parts that survive on shell, we contract the external indices of Eq. (A9) with a physical gluon polarization vectors,  $\epsilon_1^{*\sigma}$  and  $\epsilon_2^\tau$ , to obtain

$$\begin{aligned}
(\text{Fig. 6}) \epsilon_1^{*\sigma} \epsilon_2^\tau &= \frac{1}{\epsilon} \frac{g^2}{16\pi^2} C_A \delta_{ab} i \left( \frac{1}{2} \right) \left( -\frac{1}{2} g_{\mu\nu} \epsilon_1^* \cdot \epsilon_2 + \epsilon_1^*{}_\mu \epsilon_2{}_\nu + \epsilon_1^*{}_\nu \epsilon_2{}_\mu \right) \\
&+ \frac{g^2}{16\pi^2} C_A \delta_{ab} i \left\{ \frac{1}{4} \left( 1 - \frac{1}{\lambda} \right)^2 \left[ \frac{1}{2} (-g_{\mu\nu} \epsilon_1^* \cdot \epsilon_2 + \epsilon_1^*{}_\mu \epsilon_2{}_\nu + \epsilon_1^*{}_\nu \epsilon_2{}_\mu) + \frac{p_\mu p_\nu}{p^2} \epsilon_1^* \cdot \epsilon_2 \right] \right. \\
&+ \frac{1}{2} \left( 1 - \frac{1}{\lambda} \right) \left[ \frac{3}{2} g_{\mu\nu} \epsilon_1^* \cdot \epsilon_2 - \epsilon_1^*{}_\mu \epsilon_2{}_\nu - \epsilon_1^*{}_\nu \epsilon_2{}_\mu - 3 \frac{p_\mu p_\nu}{p^2} \epsilon_1^* \cdot \epsilon_2 \right] \\
&+ \left[ 1 - \frac{1}{2} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] (\epsilon_1^*{}_\mu \epsilon_2{}_\nu + \epsilon_1^*{}_\nu \epsilon_2{}_\mu) + 3 \frac{p_\mu p_\nu}{p^2} \epsilon_1^* \cdot \epsilon_2 \\
&+ \left. \left[ -\frac{3}{2} + \frac{1}{4} \ln \left( \frac{-p^2}{\bar{\mu}^2} \right) \right] g_{\mu\nu} \epsilon_1^* \cdot \epsilon_2 \right\} + O(g^4). \quad (\text{A11})
\end{aligned}$$

The quadratically IR-divergent parts of this result should be compared to Eq. (4.9), the physical matrix element of the gauge-variant part of the energy-momentum tensor.

<sup>14</sup>The pole piece cannot contain a  $\frac{1}{\bar{\mu}^2}$  divergence because of locality.

- [1] K. Wilson, *Phys. Rev.* **179**, 1499 (1969).
- [2] W. Zimmermann, *Ann. Phys. (N.Y.)* **77**, 536;570 (1973).
- [3] D. Gross and F. Wilczek, *Phys. Rev. D* **8**, 3633 (1973).
- [4] J. Collins, D. Soper, and G. Sterman, in *Perturbative QCD*, edited by A. Mueller (World Scientific, Singapore, 1989).
- [5] S. D. Joglekar and B. W. Lee, *Ann. Phys. (N.Y.)* **97**, 160 (1976).
- [6] M. Henneaux, *Phys. Lett. B* **313**, 35 (1993).
- [7] R. Hamberg, Ph.D. thesis, University of Leiden, 1991.
- [8] R. Hamberg and W. van Neerven, *Nucl. Phys.* **B379**, 143 (1992).
- [9] E. Floratos, D. Ross, and C. Sachrajda, *Nucl. Phys.* **B152**, 493 (1979).
- [10] A. Gonzalez-Arroyo and C. Lopez, *Nucl. Phys.* **B166**, 429 (1980).
- [11] E. Floratos, C. Kounnas, and R. Lacaze, *Phys. Lett.* **98B**, 285 (1981).
- [12] W. Furmanski and R. Petronzio, *Phys. Lett.* **97B**, 437 (1980).
- [13] J. C. Collins, *Renormalization* (Cambridge University Press, Cambridge, England, 1984).
- [14] D. Freedman, I. Muzinich, and E. Weinberg, *Ann. Phys. (N.Y.)* **87**, 95 (1974).
- [15] D. Freedman and E. Weinberg, *Ann. Phys. (N.Y.)* **87**, 354 (1974).
- [16] B. Harris and J. Smith, "Gauge Dependence of the Energy-Momentum Tensor in Pure Yang-Mills," Institute for Theoretical Physics, SUNY Stony Brook Report No. ITP-SB-94-04, 1994 (unpublished).
- [17] M. Henneaux and C. Teitelboim, *Quantization of Gauge Systems* (Princeton University, Princeton, N.J., 1992).
- [18] T. Kugo and I. Ojima, *Prog. Theor. Phys. Suppl.* **66** (1979).
- [19] N. Nakanishi and I. Ojima, *Covariant Operator Formalism of Gauge Theories and Quantum Gravity*, Lecture Notes in Physics Vol. 27 (World Scientific, Singapore, 1990).
- [20] F. Belinfante, *Physica* **7**, 449 (1940).
- [21] G. Wentzel, *Quantum Theory of Fields* (Interscience Publishers, New York, 1949).
- [22] J. Dixon and J. Taylor, *Nucl. Phys.* **B78**, 552 (1974).
- [23] J. Vermaseren, FORM (CAN, Amsterdam, 1991) (Version 1.1, 15 August 1991), available by anonymous ftp at [nikhef.nikhef.nl:pub/form/](ftp://nikhef.nikhef.nl/pub/form/).
- [24] S. D. Joglekar, *Ann. Phys. (N.Y.)* **108**, 233 (1977).
- [25] C. Lee, *Phys. Rev. D* **14**, 1078 (1976).
- [26] M. Henneaux, *Classical Foundations of BRST Symmetry*, Monographs and Textbooks in Physical Science Vol. 7 (Bibliopolis, Napoli, 1988).