Gluon production at high transverse momentum in the McLerran-Venugopalan model of nuclear structure functions

Alex Kovner, Larry McLerran, and Heribert Weigert

School of Physics and Astronomy, University of Minnesota, Minneapolis, Minnesota 55455

(Received 17 May 1995)

We consider the production of high transverse momentum gluons in the McLerran-Venugopalan model of nuclear structure functions. We explicitly compute the high momentum component in this model. We compute the nuclear target size A dependence of the distribution of produced gluons.

PACS number(s): 12.38.Bx, 24.85.+p, 25.75.+r

I. INTRODUCTION

Understanding parton distributions formed at the initial stages of a collision between two heavy ions is a very important open problem. In a previous paper we have set up a formalism for the calculation of the gluon distribution in the framework of the McLerran-Venugopalan model of the nuclear structure functions [1,2]. In this approach the valence quarks in the nuclei are considered as classical sources of color charge. This picture is similar to that developed by Mueller for the gluon structure functions for heavy quark systems [3]. The "initial values" of the color distribution of the valence quarks in the two colliding nuclei is given by

$$J_{1,2}^{\nu}(z(x)) = \delta^{\nu \pm} \delta(x^{\mp}) \ g \ \rho_{1,2}(x_{\perp}). \tag{1}$$

As is apparent from the the light cone delta functions $\delta(x^{\mp})$, the quarks in the nuclei appear as infinitely thin sheets of nuclear matter moving at the speed of light in positive and negative z directions, respectively.

These color charge distributions generate a classical glue field according to classical Yang-Mills equations

$$[D_{\mu}, F^{\mu\nu}] = \sum_{m=1,2} U[A](x, z_m(x)) J_m^{\nu}(z_m(x)) U[A](z_m(x), x)$$
(2)

Here $z_{1,2}(x) = x|_{x^{\pm}=0}$ serves as a reference point used to define the initial value of the charge distribution. Because of covariant current conservation $[D_{\nu}, J^{\nu}(x)] = 0$ this initial distribution evolves along the trajectory of a particle via parallel transport. This is the origin of the link operators

$$U[A](x, z_m(x)) := P \exp\left(-ig \int_{z_m(x)}^x d\omega_m^\mu A^\mu(\omega_m)\right) \quad (3)$$

connecting the initial point $z_m(x)$ and the point x on the trajectory ω_m , which appear on the right-hand side of Eq. (2).

The gluon distribution function is defined in terms of classical solutions of Eq. (2), and is related to the quantity (for precise definition see Sec. IV)

0556-2821/95/52(7)/3809(6)/\$06.00

$$\langle A_i(k)A_i^*(k)\rangle_{\rho}.$$
 (4)

Here A(k) is the Fourier transform of the classical solution. The averaging over the color charge distributions is performed independently for each nucleus with equal Gaussian weight:

$$\langle O \rangle_{\rho} = \int d\rho_1 d\rho_2 \ O$$
$$\times \exp\left(-\int d^2 x_{\perp} \frac{\operatorname{tr}[\rho_1^2(x_{\perp}) + \rho_2^2(x_{\perp})]}{2\mu^2}\right). \tag{5}$$

The purpose of this paper is to calculate this distribution function perturbatively, to lowest nontrivial order in the inverse powers of the transverse momentum $\alpha_s \mu/k$.

The basic input in our approach is the classical fields generated by single nuclei which have been derived earlier in [1,4,5]. These solutions remain valid before the collision and yield initial conditions for the field after the collision. The solution in the one nucleus case is of the $form^1$

$$\begin{aligned} A_m^{\pm} &= 0, \\ A_m^i &= \theta(x^{\mp}) \alpha_{1,2}^i(x_{\perp}). \end{aligned} \tag{6}$$

The functions α_i are implicitly determined by the "dimensionally reduced" version of the Yang-Mills equations:

$$\alpha_m^i = -\frac{1}{ig} U_m(x_\perp) \partial^i U_m^\dagger(x_\perp),$$

$$\partial^i \alpha_m^i = g \,\rho_m(x_\perp). \tag{7}$$

An obvious property of the solutions Eq. (6) is that the "transverse" components of the field strength vanish $F^{ij} = 0$. The gauge potentials themselves vanish in front of the moving charge and are a pure gauge behind it. The only physical information is contained in the discontinuity at the world line of the quarks which generate the

¹On these classical solutions the link operators in Eq. (2)drop out. They may, however, be important if one starts to consider quantum corrections in powers of α_s .

gluon fields. The field strength does not vanish only on these world lines and is confined to infinitely thin sheets.

Let us now turn to nucleus-nucleus collisions. Obviously the single nucleus solutions are still valid everywhere except in those regions of space-time which are in causal contact with the collision point, i.e., in its forward light cone (see Fig. 1). In the Abelian case even there the solutions would be directly given in terms of the single nucleus solutions as a sum of two pure gauge fields which again constitutes a pure gauge field. Hence in this case there would be no nontrivial effects on the classical field level. To generate photons and field strength one would have to resort to quantum effects. This is fundamentally different in the non-Abelian case. Obviously now the sum of two pure gauge fields is no longer a pure gauge field due to nonlinear effects, and therefore it does not solve the Yang-Mills equations for the two nuclei problem. Nevertheless the one nucleus solutions provide initial conditions for the evolution of the gauge field in the forward light cone as discussed in [2].

In the remainder of this paper we will adopt the gauge condition

$$x^{+}A^{-} + x^{-}A^{+} = 0. ag{8}$$

In this gauge the fields in the forward light cone can be cast in the form

$$A^{\pm} = \pm x^{\pm} \alpha(\tau, x_{\perp}),$$

$$A^{i} = \alpha^{i}_{\perp}(\tau, x_{\perp}),$$
(9)

where $\tau = \sqrt{2x^+x^-}$ is the proper time. Note that this representation does not introduce any physical assumption not already present in the single nucleus solution. The relation between A^{\pm} is a consequence of the gauge condition. Further, the fact that the functions $\alpha(\tau, x_{\perp})$ and $\alpha^i_{\perp}(\tau, x_{\perp})$ do not depend on the space-time rapidity $\eta = 1/2 \ln x^+/x^-$ is a natural consequence of the absence of a longitudinal length scale in our initial conditions with its infinitely thin nuclei moving at the speed of light.

Since the valence color charge densities vanish inside the light cone, the fields A_{μ} of Eq. (9) should solve the

FIG. 1. Regions with different structures of the gauge potential: In regions 1 and 2 we have the well known one nucleus solutions $\alpha_{1,2}$. While the gauge potential in the backward light cone is vanishing we have a nontrivial solution in the forward lightcone, region 3.

homogeneous equations

$$[D_{\mu}, F^{\mu\nu}] = 0. \tag{10}$$

The initial conditions are determined by patching the solutions in the three regions (see Fig. 1) together and eliminating the discontinuities at $\tau = 0$. This specifies the forward light cone fields $\alpha_{\perp}^{i}|_{\tau=0}$ and $\alpha|_{\tau=0}$ in terms of the known single nucleus solutions outside the forward light cone.

II. PERTURBATION THEORY

Using the form (9) of the gauge potential, the Yang-Mills equations (10) in the forward light cone are

$$\frac{1}{\tau^{3}}\partial_{\tau}\tau^{3}\partial_{\tau}\alpha + [D_{i}, [D^{i}, \alpha]] = 0,$$
$$\frac{1}{\tau} [D_{i}, \partial_{\tau}\alpha_{\perp}^{i}] + ig\tau [\alpha, \partial_{\tau}\alpha] = 0,$$
$$\frac{1}{\tau}\partial_{\tau}\tau\partial_{\tau}\alpha_{\perp}^{i} - ig\tau^{2}[\alpha, [D^{i}, \alpha]] - [D^{j}, F^{ji}] = 0.$$
(11)

Matching conditions on the solution lead to the initial conditions for $\alpha(\tau, x_{\perp})$ and $\alpha^i_{\perp}(\tau, x_{\perp})$ in terms of the single nucleus solutions α_m^i :

$$\begin{aligned} \alpha_{\perp}^{i}\big|_{\tau=0} &= \alpha_{1}^{i} + \alpha_{2}^{i}, \\ \alpha_{\tau=0}^{i} &= \frac{ig}{2} \left[\alpha_{1}^{i}, \alpha_{2}^{i} \right]. \end{aligned}$$
(12)

In this paper we construct solutions in the weak field limit by expanding first the initial conditions and then the fields within the forward light cone in powers of ρ .

Let us first concentrate on the initial conditions. To determine the initial conditions perturbatively, we need the single nucleus fields α_m^i , i = 1, 2 to the appropriate order.

Recall that for a single nucleus, the Yang-Mills equations reduce to the two equations (7). The first states that α_m^i is a pure gauge field. The second equation determines $lpha_m^i$ in terms of the charge density $ho_m(x_\perp)$ of the individual nuclei via

$$\partial^i \alpha_m^i = \partial^i \left\{ -\frac{1}{ig} U_m(x_\perp) \partial^i U_m^{-1} \right\} = g \rho_m(x_\perp).$$
 (13)

To second order in ρ , the solution of (13) is

$$\begin{aligned} \alpha_{m}^{i} &= -\partial^{i}\phi_{m} + \frac{ig}{2} \left(\delta^{ij} - \partial^{i} \frac{1}{\nabla_{\perp}^{2}} \partial^{j} \right) \\ &\times \left\{ \left[\phi_{m}, \partial^{j}\phi_{m} \right] + O(\rho_{m}^{3}) \right\}, \end{aligned} \tag{14}$$

where we have defined

Note that the first order term in (14) is longitudinal. whereas all higher corrections are transverse. Their sole purpose is to render α_m^i a pure gauge field. Plugging (14) into (12) gives us initial conditions up to second order in ρ_m

To solve the equations of motion (11) in the forward





7

GLUON PRODUCTION AT HIGH TRANSVERSE MOMENTUM IN ...

light cone perturbatively, we also have to expand the gauge fields there in powers of ρ_m :

$$\alpha = \sum_{n=0}^{\infty} \alpha_{(n)},$$

$$\alpha_{\perp}^{i} = \sum_{n=0}^{\infty} \alpha_{\perp(n)}^{i}.$$
 (16)

First order

To first order the equations of motion are necessarily linear

$$\frac{1}{\tau^{3}}\partial_{\tau}\tau^{3}\partial_{\tau}\alpha_{(1)} - \nabla_{\perp}^{2}\alpha_{(1)} = 0,$$

$$\partial^{i}\partial_{\tau}\alpha_{\perp}_{(1)}^{i} = 0,$$

$$\frac{1}{\tau}\partial_{\tau}\tau\partial_{\tau}\alpha_{\perp}_{(1)}^{i} - \left(\nabla_{\!\!\perp}^{2}\delta^{ij} - \partial^{i}\partial^{j}\right)\alpha_{\perp}_{(1)}^{j} = 0.$$
(17)

They are to be solved subject to the initial conditions

$$\begin{aligned} &\alpha_{(1)}(\tau, x_{\perp}) \mid_{\tau=0} = 0, \\ &\alpha^{i}_{\perp(1)}(\tau, x_{\perp}) \mid_{\tau=0} = -\partial^{i} \left(\phi^{1} + \phi^{2}\right)(x_{\perp}). \end{aligned}$$
 (18)

The initial condition for the " \pm " components to this order is trivial.

With these initial conditions, the first order solutions are completely specified and essentially trivial. We obtain vanishing $\alpha_{(1)}$,

$$\alpha_{(1)}(\tau, x_\perp) = 0, \tag{19}$$

and τ -independent $\alpha^i_{\perp(1)}$,

$$\alpha^{i}_{\perp(1)}(\tau, x_{\perp}) = -\partial^{i}\left(\phi^{1} + \phi^{2}\right)(x_{\perp}), \qquad (20)$$

which obviously is *pure gauge*.

To this order the problem is structurally identical to its Abelian counterpart. Commutators do not appear either in the equations of motion or in the initial conditions (these contributions are necessarily at least second order in ρ_m), and hence no field strength is generated in the forward light cone.

Second order

In the second order this changes immediately. Using the first order information almost all nonlinear terms drop from the second order equations which now read

$$\begin{aligned} \frac{1}{\tau^3} \partial_\tau \tau^3 \partial_\tau \alpha_{(2)} - \nabla_{\!\!\!\perp}^2 \alpha_{(2)} &= 0, \\ \partial^i \partial_\tau \alpha_{\perp}^{i}_{(2)} &= 0, \\ \frac{1}{\tau} \partial_\tau \tau \partial_\tau \alpha_{\perp}^{i}_{(2)} - \left(\nabla_{\!\!\!\perp}^2 \delta^{ij} - \partial^i \partial^j \right) \alpha_{\perp}^{j}_{(2)} \\ &+ ig \, \partial^j [\alpha_{\perp}^{i}_{(1)}, \alpha_{\perp}^{i}_{(1)}] = 0. \end{aligned} (21)$$

The only inhomogeneity in these equations comes from the pure gauge $\alpha_{\perp (1)}^{i}$. In fact using the residual (τ - independent) gauge freedom, we can remove the inhomogeneous piece altogether.

We define ϵ_{μ} by

$$A_{\mu} = V(x_{\perp}) \left[\epsilon_{\mu} - \frac{1}{ig} \partial_{\mu} \right] V^{-1}(x_{\perp}), \qquad (22)$$

where $V(x_{\perp}) = \exp[-ig\phi_{\perp}(x_{\perp})]$. $\phi_{\perp}(x_{\perp})$ is to be determined such that it removes nonlinear terms from the equations of motion for ϵ at least to second order. In fact, it turns out that it is possible to remove the inhomogeneous terms up to third order in ρ requiring

$$\partial^i \epsilon^i \big|_{\tau=0} = 0. \tag{23}$$

In higher orders it is no longer possible to linearize the equations. Using Eqs. (22) and (23) we find, for ϕ_{\perp} ,

$$\phi_{\perp} = \phi_1 + \phi_2 + O(\rho^2). \tag{24}$$

With this choice of V the first order solution vanishes:

$$\epsilon_{(1)} \equiv 0,$$

 $\epsilon_{(1)}^i \equiv 0.$ (25)

The second order equations become

$$\frac{1}{\tau^3} \partial_\tau \tau^3 \partial_\tau \epsilon_{(2)} - \nabla_{\!\!\perp}^2 \epsilon_{(2)} = 0,$$

$$\partial^i \partial_\tau \epsilon^i_{(2)} = 0,$$

$$\frac{1}{\tau} \partial_\tau \tau \partial_\tau \epsilon^i_{(2)} - \left(\delta^{ij} \nabla_{\!\!\perp}^2 - \partial^i \partial^j \right) \epsilon^j_{(2)} = 0.$$
(26)

The initial conditions on α^i_{\perp} and α are transformed into

$$\epsilon_{(2)}^{i}|_{\tau=0} = -ig \left(\delta^{ij} - \partial^{i}\frac{1}{\nabla_{\perp}^{2}}\partial^{j}\right) \left[\phi_{1}, \partial^{j}\phi_{2}\right], \quad (27)$$

$$\epsilon_{(2)}|_{\tau=0} = \frac{ig}{2} \left[\partial^i \phi_1, \partial^i \phi_2 \right].$$
(28)

Obviously, the gauge condition $\partial^i \epsilon^i |_{\tau=0} = 0$ in this order is preserved for all proper times τ . Hence the *i* components of the gauge field may be represented as

.

$$\epsilon^i = \epsilon^{ij} \partial^j \chi. \tag{29}$$

Using ϵ and χ , the second order Yang-Mills equations and initial conditions simplify even further. In terms of these variables, we finally obtain

$$\frac{1}{\tau^3}\partial_{\tau}\tau^3\partial_{\tau}\epsilon_{(2)} - \nabla_{\!\!\perp}^2\epsilon_{(2)} = 0,
\frac{1}{\tau}\partial_{\tau}\tau\partial_{\tau}\chi_{(2)} - \nabla_{\!\!\perp}^2\chi_{(2)} = 0,$$
(30)

and

$$\epsilon_{(2)} \mid_{\tau=0} = \frac{ig}{2} \left[\partial^i \phi_1, \partial^i \phi_2 \right],$$

$$\chi_{(2)} \mid_{\tau=0} = -ig \ \epsilon^{ij} \ \left[\partial^i \phi_1, \partial^j \phi_2 \right].$$
(31)

We stress that, even though the equations of motion (30)

linearize to this order due to our gauge choice, the intrinsic nonlinearities of the system are very important. They show up in the initial conditions (31) which carry nontrivial physical information.

III. PERTURBATIVE SOLUTIONS

To solve (30), we note that the equations are of form

$$\frac{1}{\tau^m}\partial_\tau \tau^m \partial_\tau f(\tau, x_\perp) - \nabla_{\!\!\perp}^2 f(\tau, x_\perp) = 0, \qquad (32)$$

where in our case $m = 1, 3, f = \chi, \epsilon$. Factorizing τ and x_{\perp} dependence, we immediately identify the eigenfunctions

$$f_{k_{\perp}}(\tau, x_{\perp}) = (\omega_k \tau)^{\frac{1-m}{2}} Z_{\pm \left|\frac{m-1}{2}\right|}(\omega_k \tau) e^{ik_{\perp}^j x_{\perp}^j}, \quad (33)$$

where $\omega_k = \sqrt{k_{\perp}^2}$ and Z_{ν} may in general be any linear combination of the Bessel functions J_{ν}, N_{ν} .

The initial conditions (31) force our solutions to be regular at $\tau = 0$ and thus preclude any admixture of Neumann functions N_{ν} .

A general solution is therefore of the form

$$\begin{split} F_{m}(\tau, x_{\perp}) &= \int \frac{d^{2}k_{\perp}}{(2\pi)^{2}} \ h_{m}(k_{\perp}) \ f_{k_{\perp}}(\tau, x_{\perp}) \\ &= \int \frac{d^{2}k_{\perp}}{(2\pi)^{2}} \ e^{ik_{\perp}^{j}x_{\perp}^{j}} \ h_{m}(k_{\perp}) \ (\omega_{k}\tau)^{\frac{1-m}{2}} \\ &\times J_{\left|\frac{m-1}{2}\right|}(\omega\tau). \end{split}$$
(34)

For equations (30) this yields

$$\begin{aligned} \epsilon_{(2)}(\tau, x_{\perp}) &= \int \! \frac{d^2 k_{\perp} d^2 y_{\perp}}{(2\pi)^2} e^{i k_{\perp}^j (x-y)_{\perp}^j} h_3(y_{\perp}) \frac{1}{\omega \tau} J_1(\omega \tau), \\ \chi_{(2)}(\tau, x_{\perp}) &= \int \! \frac{d^2 k_{\perp} d^2 y_{\perp}}{(2\pi)^2} e^{i k_{\perp}^j (x-y)_{\perp}^j} h_1(y_{\perp}) J_0(\omega \tau), \end{aligned}$$
(35)

with h_3 and h_1 determined by the initial conditions Eq. (31) as

$$h_{3}(x_{\perp}) = ig \left[\partial^{i}\phi_{1}, \partial^{i}\phi_{2}\right](x_{\perp}),$$

$$h_{1}(x_{\perp}) = -ig \frac{\epsilon^{ij}}{\nabla_{\perp}^{2}} \left[\partial^{i}\phi_{1}, \partial^{j}\phi_{2}\right](x_{\perp}).$$
(36)

In the following we will need the large τ asymptotics of the solutions

$$\chi_{(2)}(\tau, x_{\perp}) = \int \frac{d^2 k_{\perp}}{(2\pi)^2} e^{i k_{\perp}^j x_{\perp}^j} h_1(k_{\perp}) \\ \times \left\{ \frac{\sqrt{\frac{2}{\pi}}}{(\omega\tau)^{\frac{1}{2}}} \cos\left(\omega\tau - \frac{\pi}{4}\right) + O\left(\frac{1}{\tau^{\frac{3}{2}}}\right) \right\},$$
(37)

$$\epsilon_{(2)}(\tau, x_{\perp}) = \int \frac{d^2 k_{\perp}}{(2\pi)^2} e^{i k_{\perp}^j x_{\perp}^j} h_3(k_{\perp}) \\ \times \left\{ \frac{\sqrt{\frac{2}{\pi}}}{(\omega\tau)^{\frac{3}{2}}} \cos\left(\omega\tau - \frac{3}{4}\pi\right) + O\left(\frac{1}{\tau^{\frac{3}{2}}}\right) \right\}.$$
(38)

IV. DISTRIBUTION FUNCTIONS

The solutions of the Yang-Mills equations at asymptotically large τ are of the form

$$\epsilon^{a}(\tau, x_{\perp}) = \int \frac{d^{2}k_{\perp}}{(2\pi)^{2}} \frac{1}{\sqrt{2\omega}} \left\{ \frac{a_{1}^{a}(\vec{k}_{\perp})}{\tau^{3/2}} e^{ik_{\perp} \cdot x_{\perp} - i\omega\tau} + \text{c.c.} \right\},$$

$$\epsilon^{a,i}(\tau, x_{\perp}) = \int \frac{d^{2}k_{\perp}}{(2\pi)^{2}} \frac{\kappa^{i}}{\sqrt{2\omega}} \left\{ \frac{a_{2}^{a}(k_{\perp})}{\tau^{1/2}} e^{ik_{\perp}x_{\perp} - i\omega\tau} + \text{c.c.} \right\}.$$
(39)

Here $\omega = |k_{\perp}|$, and $\kappa^i = \epsilon^{ij} k^j / \omega$. The notation c.c. means complex conjugate.

To derive an expression for the energy density, we recall that τ is large. Near z = 0, this implies that in the range of z where $\tau \sim t \gg z$ the solutions are z independent. This means they asymptotically have zero p_z . Now suppose we are at any value of z, and τ is large but $t \sim z$. We can do a longitudinal boost to z = 0 without changing the solution. Again in this frame the solution has zero p_z . We see therefore that for the asymptotic solutions the space-time rapidity is one to one correlated with the momentum space rapidity, that is at asymptotic times we find that

$$\eta = \frac{1}{2}\ln(x^+/x^-) = y = \frac{1}{2}\ln(p^+/p^-).$$
(40)

To proceed further, we compute the energy density in the neighborhood of z = 0. Here asymptotically $\tau = t$. The energy in a box of size R in the transverse direction and dz in the longitudinal direction, with $L \ll t$, becomes [6]

$$dE = \frac{dz}{t} \int \frac{d^2 k_{\perp}}{(2\pi)^2} \omega \sum_{i,b} |a_i^b(k_{\perp})|^2 .$$
 (41)

Recalling that dy = dz/t, we find that

$$\frac{dE}{dyd^2k_{\perp}} = \frac{1}{(2\pi)^3}\omega \sum_{i,b} |a_i^b(k_{\perp})|^2$$
(42)

and the multiplicity distribution of gluons is

$$\frac{dN}{dyd^2k_{\perp}} = \frac{1}{\omega}\frac{dE}{dyd^2k_{\perp}}.$$
(43)

As we expect for a boost covariant solution, the multiplicity distribution is rapidity invariant.

This is now easily compared to the asymptotic behav-

ior of the second order solutions (38), (37). We read off that

$$a_1(k_{\perp}) = \frac{1}{\sqrt{\pi}} \frac{h_3(k_{\perp})}{\omega},$$

$$a_2(k_{\perp}) = \frac{1}{\sqrt{\pi}} i\omega h_1(k_{\perp}).$$
 (44)

Hence we get the distribution function

$$\begin{split} \frac{dN}{dyd^2k_{\perp}} &= \frac{1}{\omega}\frac{dE}{dyd^2k_{\perp}} = \frac{1}{(2\pi)^3}\sum_i 2 \ \text{tr} \ a_i(k_{\perp})a_i^{\dagger} \\ &= \frac{\frac{2}{\pi}}{(2\pi)^3}\text{tr}\left\{\frac{h_3(k_{\perp})h_3^{\dagger}(k_{\perp})}{\omega^2} + \omega^2h_1(k_{\perp})h_1^{\dagger}(k_{\perp})\right\}. \end{split}$$
(45)

Now we use

and get

$$\frac{dN}{dyd^{2}k_{\perp}} = S_{\perp}\frac{g^{2}}{(2\pi)^{3}}\frac{2}{\pi k_{\perp}^{2}} \left[\delta^{ij}\delta^{kl} + \epsilon^{ij}\epsilon^{kl}\right] \int d^{2}x_{\perp}e^{ik_{\perp}\cdot x_{\perp}} \\
\times \operatorname{tr}\left\langle \left[\partial^{i}\phi_{1}, \partial^{j}\phi_{2}\right](x_{\perp}) \left[\partial^{k}\phi_{1}, \partial^{l}\phi_{2}\right]^{\dagger}(0)\right\rangle_{\rho}, \tag{47}$$

where S_{\perp} is the transverse area of the system. As indicated, we have to average over ρ with the Gaussian weight Eq. (5). Using (15) we find

$$\langle \phi_m^a(x_{\perp})\phi_n^b(y_{\perp})\rangle_{\rho} = (g\mu)^2 \,\delta^{mn}\delta^{ab} \int \frac{d^2p_{\perp}}{(2\pi)^2} \frac{e^{ip_{\perp}\cdot(x-y)_{\perp}}}{p_{\perp}^4}$$
(48)

and the above turns into

$$\frac{dN}{dyd^{2}k_{\perp}} = S_{\perp} \frac{g^{6}}{(2\pi)^{3}} \left[\delta^{ij} \delta^{kl} + \epsilon^{ij} \epsilon^{kl} \right] \operatorname{tr} \left([\tau^{a}, \tau^{b}] [\tau^{b}, \tau^{a}] \right) \\
\times \frac{2\mu^{4}}{\pi k_{\perp}^{2}} \int \frac{d^{2}p_{\perp}}{(2\pi)^{2}} \frac{p^{i} p^{k} (p+k)^{j} (p+k)^{l}}{p^{4} (p+k)^{4}} \\
= S_{\perp} \frac{2g^{6} \mu^{4}}{(2\pi)^{3} \pi} \frac{N_{c} (N_{c}^{2}-1)}{k_{\perp}^{2}} \int \frac{d^{2}p_{\perp}}{(2\pi)^{2}} \frac{1}{p^{2} (p+k)^{2}}. \quad (49)$$

The integral in Eq. (49) is infrared divergent. This, however, is an artifact of the weak field expansion employed above. As discussed in [1,4,5,2], the inclusion of higher order terms for the initial fields would generate a mass scale of order $\alpha_s \mu$ and hence regulate this divergence. With logarithmic accuracy we obtain therefore

$$\frac{dN}{dyd^2k_{\perp}} = S_{\perp} \frac{2g^6\mu^4}{(2\pi)^4\pi} N_c (N_c^2 - 1) \frac{1}{k_{\perp}^4} \ln \frac{k_{\perp}^2}{(\alpha_s \mu)^2}.$$
 (50)

This is the main result of this work.

V. CONCLUSION

In this paper we have calculated the high k_{\perp} asymptotics of the distribution function of gluons produced in a collision of two ultrarelativistic heavy nuclei in the framework of the McLerran-Venugopalan model.

Our result, Eq. (50), has a very natural interpretation in analogy with the naive parton model. The number of produced gluons in the parton model is given by

$$N(k_{\perp}) = \int d^2 p_{\perp} N_1(p_{\perp}) N_2(k_{\perp} - p_{\perp}) \sigma(p_{\perp}, k_{\perp}),$$
 (51)

where $N_m(q_{\perp})$ is the probability to find a gluon with transverse momentum q_{\perp} in the *m*th nucleus, and σ is the cross section for production of the gluon with momentum k_{\perp} in the final state in the collision of the two gluons. To the lowest order in perturbation theory the cross section is

$$\sigma(k_{\perp}) \propto \frac{\alpha_s}{k_{\perp}^2}.$$
 (52)

The distribution functions for the one nucleus case N_m were calculated in [1,4] and were found to have the same functional form as in the lowest order in perturbation theory:

$$N_m(q_\perp) \propto rac{lpha_s \mu^2}{q_\perp^2}.$$
 (53)

Substituting this into Eq. (51) gives the result of Eq. (50).

Indeed the calculation performed in this paper can be represented in terms of Feynman diagrams. The gluon distribution function of Eq. (50) has a representation of Fig. 2.

The reader may have noticed one peculiarity in the preceding discussion. Although we have been using the



FIG. 2. Diagrammatic representation of the lowest order distribution function.

parton model notations, some of the gluons involved in the process depicted on Fig. 2 are virtual. To produce the real final state gluon, the internal gluons must be off shell with virtuality of order of their momentum. In fact if one wants to think about this process in terms of real partons, it must be represented as "two quarks go into two quarks and a gluon," rather than "two gluons go into one gluon," which is kinematically forbidden for on shell particles.

As we have seen, to this order our calculation essentially reproduces perturbation theory. The reason is that we have expanded in powers of the valence charge density, and thereby have ignored the fact that the classical fields involved are expected to be strong. The real nonperturbative nature of this approach will become apparent when those are fully taken into account. Some of the qualitative consequences of such an analysis, however, can be understood already at this point. Consider, for example, the total multiplicity of produced gluons,

$$N = \int d^2k_\perp \frac{dN}{dyd^2k_\perp}.$$
 (54)

Since $\frac{dN}{dyd^2k_{\perp}}$ is proportional to the transverse area of the system, and is a function of the dimensionless ratio μ^2/k_{\perp}^2 , the integral has to be proportional to $S_{\perp}\mu^2$. Remembering that for large nuclei μ scales with the number of nucleon in the nucleus A as $\mu^2 \propto A^{1/3}$, we conclude that for large nuclei the total multiplicity scales as

$$N \propto A.$$
 (55)

On the other hand, the total multiplicity at momenta larger than some fixed momentum $p \gg \alpha \mu$ will coincide with the perturbative result

$$N = \int_{k_{\perp}^2 \ge p^2} d^2 k_{\perp} \frac{dN}{dy d^2 k_{\perp}} \propto S_{\perp} \mu^4 \propto A^{4/3}.$$
 (56)

Therefore it is very important to go beyond the weak field expansion and incorporate truly nonperturbative effects due to strong classical fields.

In comparing our results with the parton cascade model of Geiger *et al.* [7–9] one should keep in mind the following differences between the two approaches. In the parton cascade, the leading order in α_s scattering occurs

by gluon gluon scattering. This populates the high p_{\perp} tail of the distribution by scatterings of low p_{\perp} gluons. In our case, we have an intrinsic p_{\perp} for the gluons and the gluons are far enough off mass shell so that we can produce high p_{\perp} gluons by glue-glue goes to single gluon scattering. Consequently, although the p_{\perp} dependence of the two results is similar, in the high p_{\perp} region we have a different dependence on α_s . We have one power less, since the factor $\alpha^2 \mu^2$ in Eq. (50) just sets the scale of the intrinsic single nucleus glue distribution.

The scattering processes, which are leading in our approximation are also present in cascade codes as composite processes $qq \rightarrow qqq$, where some of the quarks are off shell by an amount p_{\perp}^2 . Although these contributions are formally of order α^3 , for very large nuclei the quark distribution functions will be large and these processes may become leading. This seems to be the natural point of contact between the two approaches.

Of course at some $p_{\perp} \gg \mu$, our approximations for computing the gluon distribution function break down. This presumably first occurs when gluon bremsstrahlung softens the high p_{\perp} distribution from its $1/p_{\perp}^2$ behavior, and the spectrum steepens. The main point is, however, not what happens in the tail of the distribution. This is simply where it is most easy to compute. In the center of the p_{\perp} distribution of produced gluons, the gluons arise from a nonlinear evolution of the gluons fields. This is cause by the quantum-mechanical nature of the initial state and the charge coherence of the initial state interactions. In this case, the final state distribution of gluons bears scant resemblance to that of the initial distribution convoluted over hard gluon scattering. Before one knows whether there are substantial quantitative differences, one must of course numerically solve the problem for gluons in the center of the distribution.

ACKNOWLEDGMENTS

This research was supported by the U.S. Department of Energy under Grants No. DOE High Energy DE-AC02-83ER40105 and No. DOE Nuclear DE-FG02-87ER-40328. One of us (H.W.) was supported by the Alexander von Humboldt Foundation.

- [1] L. McLerran and R. Venugopalan, Phys. Rev. D 49, 2233 (1994).
- [2] A. Kovner, L. McLerran, and H. Weigert, "Gluon Production from Non-Abelian Weizsäcker-Williams Fields in Nucleus-Nucleus Collisions," Reports No. TPI-MINN-95-16/T, No. NUC-MINN-95-14/T, and No. HEP-MINN-95-134, 1995 (unpublished).
- [3] A. Mueller, Nucl. Phys. B425, 373 (1994)
- [4] L. McLerran and R. Venugopalan, Phys. Rev. D 49, 3352

(1994).

- [5] L. McLerran and R. Venugopalan, Phys. Rev. D 50, 2225 (1994).
- [6] J. P. Blaizot and A. Mueller, Nucl. Phys. B289, 847 (1987).
- 7] K. Geiger and B. Müller, Nucl. Phys. B369, 600 (1992).
- [8] K. Geiger, Phys. Rev. D 47, 133 (1993).
- [9] K. Geiger, CERN Report No. CERN-TH 7313/94, 1994.