

Search for a vector glueball by a scan of the J/ψ resonance

J. Z. Bai,¹ O. Bardon,⁶ I. Blum,¹¹ A. Breakstone,⁹ T. Burnett,¹² G. P. Chen,¹ H. F. Chen,⁴ J. Chen,⁵ S. M. Chen,¹ Y. Chen,¹ Y. B. Chen,¹ Y. Q. Chen,¹ B. S. Cheng,¹ R. F. Cowan,⁶ X. Z. Cui,¹ H. L. Ding,¹ Z. Z. Du,¹ W. Dunwoodie,⁸ X. L. Fan,¹ J. Fang,¹ M. Fero,⁶ C. S. Gao,¹ M. L. Gao,¹ S. Q. Gao,¹ P. Gratton,¹¹ J. H. Gu,¹ S. D. Gu,¹ W. X. Gu,¹ Y. F. Gu,¹ Y. N. Guo,¹ S. W. Han,¹ Y. Han,¹ F. A. Harris,⁹ M. Hatanaka,³ J. He,¹ M. He,⁷ D. G. Hitlin,³ G. Y. Hu,¹ T. Hu,¹ X. Q. Hu,¹ D. Q. Huang,¹ Y. Z. Huang,¹ J. M. Izen,¹¹ Q. P. Jia,⁵ C. H. Jiang,¹ S. Jin,¹ Y. Jin,¹ L. Jones,³ S. H. Kang,¹ Z. J. Ke,¹ M. H. Kelsey,³ B. K. Kim,¹¹ D. Kong,⁹ Y. F. Lai,¹ H. B. Lan,¹ P. F. Lang,¹ A. Lankford,¹⁰ F. Li,¹ J. Li,¹ P. Q. Li,¹ Q. Li,⁷ R. B. Li,¹ W. Li,¹ W. D. Li,¹ W. G. Li,¹ X. H. Li,¹ X. N. Li,¹ S. Z. Lin,¹ H. M. Liu,¹ J. Liu,¹ J. H. Liu,¹ Q. Liu,¹ R. G. Liu,¹ Y. Liu,¹ Z. A. Liu,¹ X. C. Lou,¹¹ B. Lowery,¹¹ J. G. Lu,¹ S. Q. Luo,¹ Y. Luo,¹ A. M. Ma,¹ E. C. Ma,¹ J. M. Ma,¹ H. S. Mao,¹ Z. P. Mao,¹ R. Malchow,⁵ M. Mandelkern,¹⁰ X. C. Meng,¹ H. L. Ni,¹ J. Nie,¹ S. L. Olsen,⁹ J. Oyang,³ D. Paluselli,⁹ L. J. Pan,⁹ J. Panetta,³ F. Porter,³ E. Prabhakar,³ N. D. Qi,¹ Y. K. Que,¹ J. Quigley,⁶ G. Rong,¹ M. Schernau,¹⁰ B. Schmid,¹⁰ J. Schultz,¹⁰ Y. Y. Shao,¹ B. W. Shen,¹ D. L. Shen,¹ H. Shen,¹ X. Y. Shen,¹ H. Y. Sheng,¹ H. Z. Shi,¹ X. R. Shi,³ A. Smith,¹⁰ E. Soderstrom,⁸ X. F. Song,¹ J. Standifird,¹¹ D. Stoker,¹⁰ F. Sun,¹ H. S. Sun,¹ S. J. Sun,¹ J. Synodinos,⁸ Y. P. Tan,¹ S. Q. Tang,¹ W. Toki,⁵ G. L. Tong,¹ E. Torrence,⁶ F. Wang,¹ L. S. Wang,¹ L. Z. Wang,¹ M. Wang,¹ P. Wang,¹ P. L. Wang,¹ S. M. Wang,¹ T. J. Wang,¹ Y. Y. Wang,¹ C. L. Wei,¹ S. Whittaker,² R. Wilson,⁵ W. J. Wisniewski,^{13,*} D. M. Xi,¹ X. M. Xia,¹ P. P. Xie,¹ W. J. Xiong,¹ D. Z. Xu,¹ R. S. Xu,¹ Z. Q. Xu,¹ S. T. Xue,¹ R. Yamamoto,⁶ J. Yan,¹ W. G. Yan,¹ C. M. Yang,¹ C. Y. Yang,¹ J. Yang,¹ W. Yang,¹ M. H. Ye,¹ S. W. Ye,⁴ S. Z. Ye,¹ K. Young,¹² C. S. Yu,¹ C. X. Yu,¹ Z. Q. Yu,¹ C. Z. Yuan,¹ B. Y. Zhang,¹ C. C. Zhang,¹ D. H. Zhang,¹ H. L. Zhang,¹ J. Zhang,¹ J. W. Zhang,¹ L. S. Zhang,¹ S. Q. Zhang,¹ Y. Zhang,¹ Y. Y. Zhang,¹ D. X. Zhao,¹ H. W. Zhao,¹ J. W. Zhao,¹ M. Zhao,¹ P. D. Zhao,¹ W. R. Zhao,¹ J. P. Zheng,¹ L. S. Zheng,¹ Z. P. Zheng,¹ G. P. Zhou,¹ H. S. Zhou,¹ L. Zhou,¹ X. F. Zhou,¹ Y. H. Zhou,¹ Q. M. Zhu,¹ Y. C. Zhu,¹ Y. S. Zhu,¹ B. A. Zhuang,¹ and G. Zioulas¹⁰

(BES Collaboration)

¹*Institute of High Energy Physics, Beijing 100039, People's Republic of China*

²*Boston University, Boston, Massachusetts 02215*

³*California Institute of Technology, Pasadena, California 91125*

⁴*China's University of Science and Technology, Hefei 230026, People's Republic of China*

⁵*Colorado State University, Fort Collins, Colorado 80523*

⁶*Massachusetts Institute of Technology, Cambridge, Massachusetts 02139*

⁷*Shandong University, Jinan 250100, People's Republic of China*

⁸*Stanford Linear Accelerator Center, Stanford, California 94309*

⁹*University of Hawaii, Honolulu, Hawaii 96822*

¹⁰*University of California at Irvine, Irvine, California 92717*

¹¹*University of Texas at Dallas, Richardson, Texas 75083-0688*

¹²*University of Washington, Seattle, Washington 98195*

¹³*Superconducting Supercollider Laboratory, Dallas, Texas 75237-3946*

(Received 19 October 1995)

The cross section for $e^+e^- \rightarrow \rho\pi$ has been measured by the BES detector at BEPC at center-of-mass energies covering a 40 MeV interval spanning the J/ψ resonance. The data are used to search for the vector gluonium state hypothesized by Brodsky, Lepage, and Tuan as an explanation of the $\rho\pi$ puzzle in charmonium physics. The shape of the $\rho\pi$ cross section is compatible with that of the total hadronic cross section. No distortions indicating the presence of a vector glueball are seen. [S0556-2821(96)03613-2]

PACS number(s): 13.25.Gv, 12.39.Mk, 13.65.+i, 14.40.Gx

There is a well-known mystery in charmonium physics, the so-called “ $\rho\pi$ puzzle.” According to quantum chromodynamics, the J/ψ and $\psi(2S)$ resonances are S -wave $c\bar{c}$ bound states that decay to hadrons via three gluons or, with a smaller probability, via a single direct photon. In either case the partial decay width is proportional to $|\Psi(0)|^2$, the square of the $c\bar{c}$ wave function at the origin. Expectations based on

perturbative QCD are that, for any specific final hadronic state h ,

$$Q_h \equiv \frac{B(\psi(2S) \rightarrow h)}{B(J/\psi \rightarrow h)} \equiv \frac{B(\psi(2S) \rightarrow e^+e^-)}{B(J/\psi \rightarrow e^+e^-)} = 0.147 \pm 0.023 \quad [1]. \quad (1)$$

While this rule appears to be satisfied for most hadronic channels, there are startling exceptions that occur for the $\rho\pi$ and $K^*\bar{K}$ final states. These are dominant hadronic channels for J/ψ decays but are not seen in $\psi(2S)$ decays; the present experimental limits [1] are

*Present address: Stanford Linear Accelerator Center, Stanford, California 94309.

$$Q_{\rho\pi} < 0.0028 \quad \text{and} \quad Q_{K^*\bar{K}} < 0.011. \quad (2)$$

An intriguing possible explanation for this anomaly was provided by Brodsky, Lepage, and Tuan [2] based on a refinement of an idea of Hou and Soni [3]. They assume the general validity of the perturbative QCD theorem that total hadron helicity is conserved in high-momentum-transfer exclusive processes, in which case the decays to $\rho\pi$ and $K^*\bar{K}$ are forbidden for both the J/ψ and the $\psi(2S)$. The observed J/ψ decays to these channels are mediated by an intermediate gluonium state O with quantum number $J^{PC} = 1^{--}$ and a mass near that of the J/ψ . The experimental limits given by Eq. (2) imply that the O mass is within 80 MeV of the mass of the J/ψ and its total width is less than 160 MeV [2].

Experimentally, the search for such a vector gluonium state could be carried out using sources of three gluons occurring in certain hadronic decays of the $\psi(2S)$, such as $\psi(2S) \rightarrow \pi\pi + X, \eta + X, \eta' + X$, where X decays into vector-pseudoscalar (VP) final states [3]. A more direct way to verify the existence of O is to scan the $e^+e^- \rightarrow VP$ cross section across the J/ψ resonance [4]. The O , being close in mass to the J/ψ and having decay width Γ_O substantially larger than $\Gamma_{J/\psi}$, should interfere with the J/ψ , producing a distortion of the shape of the J/ψ obtained from the $e^+e^- \rightarrow VP$ cross section compared to that of the total hadronic or lepton pair cross section. In this paper, we report on such a search based on an energy scan done with the Beijing Spectrometer (BES) [5] at the Beijing Electron-Positron Collider (BEPC).

Cross section measurements were performed at 29 different energy points covering a 40 MeV interval spanning the J/ψ resonance. The original purpose of the scan was measurements of the total and lepton partial widths of the J/ψ , the results of which are published elsewhere [6]. The data sample corresponds to a total integrated luminosity of 238 nb^{-1} . In this energy region the uncertainty in the c.m. energy of the BEPC collider is ± 0.07 MeV, and the rms c.m. energy spread is 1 MeV [5]. A summary of the energy points at which data were taken is provided in Table I.

In this analysis, the candidate events for $e^+e^- \rightarrow \rho\pi$ are required to have exactly two, oppositely charged and well-reconstructed tracks, and between two and six neutral energy clusters in the shower counters. Each charged track is identified as a pion using a combination of the time-of-flight and dE/dx information. In order to remove lepton pair events, we require the opening angle between the two charged tracks to be less than 170° , the total energy deposit in the calorimeter associated with the two charged tracks to be less than 1.6 GeV, and at least one charged track *not* identified as a muon by the muon counters. The events are required to have at least two “good” photons, which are defined as neutral energy deposits that are larger than 80 MeV and are separated from the nearest charged track by more than 8° . The selected events are kinematically fitted to the $\pi^+\pi^-\gamma\gamma$ hypothesis by imposing the energy and momentum constraints corresponding to each scan point. For events where the number of good photons exceeds 2, all two-photon combinations are fitted, and the combination with the minimum χ^2 and $P(\chi^2) > 0.01$ is retained. Finally, the mass cuts $|M_{\gamma\gamma} - M_{\pi^0}| < 50 \text{ MeV}/c^2$ and $|M_{\pi\pi} - M_\rho| < 200 \text{ MeV}/c^2$

TABLE I. A summary of the data of the $\rho\pi$ scan.

| Scan point | W (MeV) | \mathcal{L} (nb^{-1}) | N |
|------------|-----------|------------------------------------|-----|
| 1 | 3079.1 | 11.434 | 0 |
| 2 | 3080.5 | 51.378 | 0 |
| 3 | 3084.9 | 12.086 | 1 |
| 4 | 3085.5 | 10.767 | 0 |
| 5 | 3087.7 | 9.142 | 0 |
| 6 | 3090.0 | 12.318 | 0 |
| 7 | 3090.6 | 5.801 | 0 |
| 8 | 3091.9 | 6.019 | 0 |
| 9 | 3092.6 | 2.826 | 0 |
| 10 | 3093.6 | 6.866 | 1 |
| 11 | 3094.5 | 5.715 | 4 |
| 12 | 3095.3 | 5.498 | 12 |
| 13 | 3095.7 | 5.459 | 26 |
| 14 | 3096.1 | 3.353 | 14 |
| 15 | 3096.5 | 5.617 | 24 |
| 16 | 3096.9 | 5.414 | 17 |
| 17 | 3097.3 | 1.833 | 10 |
| 18 | 3097.7 | 6.161 | 15 |
| 19 | 3098.1 | 5.853 | 8 |
| 20 | 3098.5 | 4.384 | 2 |
| 21 | 3098.9 | 1.818 | 1 |
| 22 | 3099.3 | 4.036 | 0 |
| 23 | 3099.7 | 1.738 | 0 |
| 24 | 3100.1 | 13.470 | 3 |
| 25 | 3100.7 | 1.990 | 1 |
| 26 | 3102.7 | 11.376 | 2 |
| 27 | 3105.5 | 11.226 | 1 |
| 28 | 3110.1 | 5.659 | 0 |
| 29 | 3116.4 | 8.709 | 1 |

are also applied. The number of $\rho\pi$ candidates found in each scan point is listed in the last column of Table I.

The $\rho\pi$ scan data are analyzed using a maximum likelihood method. The likelihood function is a product of Poisson distributions, one for each center-of-mass energy W_i . At each scan point i , the number of expected events ν_i is given by

$$\nu_i = [\epsilon \sigma_{\rho\pi}(W_i) + \sigma_b(W_i)] \mathcal{L}_i; \quad (3)$$

ϵ is the overall efficiency for identifying $\rho\pi$ events, \mathcal{L}_i is the integrated luminosity, $\sigma_{\rho\pi}(W_i)$ is the corresponding cross section for $\rho\pi$ production, and $\sigma_b(W_i)$ is the background cross section. The efficiency combines the trigger efficiency, which is estimated to be 98.4% for two-prong hadron events [6], and the reconstruction and selection efficiency, which is determined from Monte Carlo simulations to be 13.1% and virtually independent of energy over the scan region. Backgrounds come from $e^+e^- \rightarrow J/\psi \rightarrow \pi^+\pi^-\pi^0(\text{direct})$, $K^+K^-\pi^0$, etc. Assuming that the shape of the background is compatible with the shape of $e^+e^- \rightarrow J/\psi \rightarrow \text{hadrons}$, the background cross section can be written as

$$\sigma_b(W) = \sum_k \epsilon_k B_k \sigma_{\text{tot}}(W), \quad (4)$$

where ϵ_k is the overall efficiency for a specific background process k , B_k is the corresponding branching fraction, and $\sigma_{\text{tot}}(W)$ is the total observed cross section of the J/ψ resonance in which radiative corrections and collider c.m. energy spread are taken into account. For $\pi^+\pi^-\pi^0(\text{direct})$ and $K^+K^-\pi^0$, the main background processes, Monte Carlo simulations yield the efficiencies for reconstruction and selection to be 5.86% and 0.75%, respectively.

The $\rho\pi$ cross section $\sigma_{\rho\pi}(W)$ is the sum of a nonresonant direct-channel contribution from nonresonant virtual photon

annihilation and a resonant term that is convoluted with a Gaussian function describing the c.m. energy spread of the collider:

$$\sigma_{\rho\pi}(W) = R_{\rho\pi}\sigma_{\mu\mu}(W) + \frac{1}{\sqrt{2\pi}\Delta} \int_0^\infty dW' \times \exp\left(-\frac{(W-W')^2}{2\Delta^2}\right) \sigma(W'). \quad (5)$$

$$\begin{aligned} \sigma(W) = & \left[1 + \frac{3}{4}\beta + \frac{\alpha}{\pi}\left(\frac{\pi^2}{3} - \frac{1}{2}\right) - \beta^2\left(\frac{\pi^2}{12} - \frac{9}{32}\right)\right] \int_0^1 dx \beta x^{\beta-1} \left(\frac{|B|^2}{[W^2(1-x) - M_{J/\psi}^2 + M_{J/\psi}^2 \Gamma_{J/\psi}^2]} \right. \\ & + \frac{|D|^2}{[W^2(1-x) - M_O^2 + M_O^2 \Gamma_O^2]} + \frac{2\text{Re}\{DB[W^2(1-x) - M_O^2 - iM_O\Gamma_O][W^2(1-x) - M_{J/\psi}^2 + iM_{J/\psi}\Gamma_{J/\psi}]\}}{\{[W^2(1-x) - M_O^2 + M_O^2 \Gamma_O^2]\{[W^2(1-x) - M_{J/\psi}^2 + M_{J/\psi}^2 \Gamma_{J/\psi}^2]\}} \Big) \\ & \equiv \sigma_{J/\psi}(W) + \sigma_{O+I}(W), \end{aligned} \quad (6)$$

where $\beta = (2\alpha/\pi)[\ln(W^2/m_e^2) - 1]$, B is the J/ψ partial width to e^+e^- (denoted as $\Gamma_{J/\psi}^{ee}$) and $\rho\pi$ (denoted as $\Gamma_{J/\psi}^{\rho\pi}$) in the form $|B|^2 = 12\pi\Gamma_{J/\psi}^{ee}\Gamma_{J/\psi}^{\rho\pi}$, $D = |D|\exp(i\theta)$ is the contribution from the O with θ denoting a possible relative complex phase, $M_{J/\psi}, \Gamma_{J/\psi}$ are the mass and total width of the J/ψ , and M_O, Γ_O are the mass and total width of the O . For the analysis in this work, $\sigma(W)$ is decomposed into two parts: $\sigma_{J/\psi}(W)$ is the first term in Eq. (6) and corresponds to the normal J/ψ resonance, and $\sigma_{O+I}(W)$ is the sum of the second and third terms due to the O and its interference with the J/ψ .

In fitting the cross section $\sigma_{\rho\pi}$ in Eq. (5) to the results listed in Table I, $|B|$, $|D|$, $R_{\rho\pi}$, M_O , Γ_O , and θ are treated as free, independent parameters. M_O is limited to vary from $M_{J/\psi} - 80$ MeV to $M_{J/\psi} + 80$ MeV, and Γ_O up to 160 MeV. These ranges are suggested by Ref. [2]. The lower bound of Γ_O is set to 2 MeV. Although in the literature [3,8,2] Γ_O is proposed to be at least on the order of 10 MeV, here a more general search is pursued. In this experiment, 2 MeV is the limit of the sensitivity, due to the finite c.m. energy spread of the collider and the statistics of the data. For $M_{J/\psi}$, $\Gamma_{J/\psi}$, $\Gamma_{J/\psi}^{ee}$, and Δ , the values determined from the same set of the scan data [6] are used here as well as in the background expression of Eq. (4). The best fit values are at $M_O - M_{J/\psi} = 0.7 \pm 0.4$ MeV and $\Gamma_O = 2$ MeV. The fits yield an upper limit with the cross section ratio of

$$\sigma_{O+I}/\sigma_{J/\psi} < 0.098 \quad (90\% \text{ C.L.}) \quad (7)$$

at $W = M_{J/\psi}$.

For the model proposed in Ref. [2], the $\sigma_{J/\psi}$ contribution should observe the perturbative QCD relation of Eq. (1): namely

$$\frac{B(\psi(2S) \rightarrow \rho\pi)}{\sigma_{J/\psi}/\sigma_{\text{tot}}} = 0.147, \quad (8)$$

while the anomalous $J/\psi \rightarrow \rho\pi$ decay mode is the contribution from σ_{O+I} , modifying Eq. (8) to become

$$\frac{B(\psi(2S) \rightarrow \rho\pi)}{(\sigma_{J/\psi} + \sigma_{O+I})/\sigma_{\text{tot}}} = Q_{\rho\pi}. \quad (9)$$

Here $\sigma_{\mu\mu}$ is the pure QED $e^+e^- \rightarrow \mu\mu$ cross section and $R_{\rho\pi}$ is the fractional nonresonant direct-channel cross section for $\rho\pi$ production, Δ is the standard deviation of the Gaussian function, and $\sigma(W)$ is the Breit-Wigner cross section due to the J/ψ and O propagators for the process $e^+e^- \rightarrow \rho\pi$. Taking into account radiative corrections [7], $\sigma(W)$ can be expressed as

Combining Eqs. (8) and (9) and using the experimental limit of $Q_{\rho\pi}$ in Eq. (2), we find

$$\sigma_{O+I}/\sigma_{J/\psi} = 0.147/Q_{\rho\pi} - 1 > 51.5, \quad (10)$$

in clear contradiction with the obtained upper bound in Eq. (7). In Fig. 1, the contours of the fitted ratio $\sigma_{O+I}/\sigma_{J/\psi}$ are plotted on the Γ_O and $M_O - M_{J/\psi}$ planes. The shaded area is the insensitive region of this experiment, due to the finite c.m. energy spread of the collider and the statistics of the data. It can be seen clearly that outside this insensitive region, the existence of a vector gluonium state which contributes to the anomalous $\rho\pi$ decay is definitely excluded.

Furthermore, if we define λ to be the ratio of $L(H_0)$, the maximum value of the likelihood function for the null hypothesis [$|D|=0$ in Eq. (6)], to $L(H_1)$, the maximum value of the likelihood function for the vector glueball hypothesis, we find $-2\ln\lambda = 0.84$. In the limit where the distribution of $-2\ln\lambda$ is the χ^2 distribution for one degree of freedom, the 90% confidence level expectation is $-2\ln\lambda < 2.71$. This indicates that our results are in good agreement with the null hypothesis and there is no evidence for the vector glueball hypothesis.

In the case of no contribution from the O , $\sigma_{\rho\pi}(W)$ can be simply expressed as

$$\sigma_{\rho\pi}(W) = R_{\rho\pi}\sigma_{\mu\mu}(W) + B_{\rho\pi}\sigma_{\text{tot}}(W), \quad (11)$$

and fitting the data allowing both $B_{\rho\pi}$ and $R_{\rho\pi}$ to be free parameters yields $B_{\rho\pi} = B(J/\psi \rightarrow \rho\pi) = (1.21 \pm 0.20)\%$, while $R_{\rho\pi} = (7.3 \pm 7.1) \times 10^{-3}$. The errors include the statistical and the systematic errors added in quadrature. This is the first measurement of the $B_{\rho\pi}$ for the J/ψ from the scan across the resonance, and the result is consistent with the Particle Data Group value [1], which comes from the data taken at the J/ψ resonance peak. The measured $\rho\pi$ cross sections are shown as a function of c.m. energy in Fig. 2(a). The solid curve in this figure is the result of the likelihood fit to the null hypothesis. The dot-dashed curve shows the distorted cross sections expected from the hypothetical O with $M_O = M_{J/\psi}$ and $\Gamma_O = 2$ MeV whose contribution satisfies the




FIG. 1. The contours of the fitted ratio $\sigma_{O+I}/\sigma_{J/\psi}$ for each pair of Γ_O and $M_O - M_{J/\psi}$ values. The shaded area is the insensitive region of this experiment. It is obvious that outside this region, the existence of the O to explain the anomalous $J/\psi \rightarrow \rho\pi$ can be ruled out.

lower bound of relation (10). The more the M_O is deviated from the $M_{J/\psi}$ and the broader the Γ_O is, the better the shape of the cross section curve would be distinguished from the J/ψ resonance. For comparison, the measured total hadronic cross section and the fitted curve from Ref. [6] are shown in Fig. 2(b). The quality of the fit in Fig. 2(a) is checked by forming the likelihood ratio λ with the result $-2\ln\lambda=22.0$. In the large statistics limit, this should obey a χ^2 distribution for 27 degrees of freedom, further indicating that the fit with the null hypothesis is acceptable.

In conclusion, the data from a scan of the $\rho\pi$ cross section over a 40 MeV interval spanning the J/ψ resonance have been used to search for the vector gluonium state O that was hypothesized to explain the $\rho\pi$ puzzle in J/ψ and $\psi(2S)$ physics. No evidence was found to support the existence of such a vector gluonium state. The shape of the $\rho\pi$ curve is found to be consistent with the shape of the total hadronic curve with no distortion observed within the experimental uncertainties. The $J/\psi \rightarrow \rho\pi$ branching fraction obtained from fitting the shape of the scan data yields the result $(1.21 \pm 0.20)\%$. The $\rho\pi$ puzzle in charmonium physics remains a puzzle.

We gratefully acknowledge the dedicated efforts of the staffs of the BEPC accelerator and the Computing Center at




FIG. 2. (a) The center-of-mass energy dependence of the $\rho\pi$ cross section resulting from the likelihood fit (solid curve), compared to the data corrected for the efficiency and background. The dot-dashed curve is the expected cross section due to the O if its contribution is at the lower bound of relation (10) with $M_O = M_{J/\psi}$ and $\Gamma_O = 2$ MeV which corresponds to the limit of the experimental sensitivity. (b) The total cross section for $e^+e^- \rightarrow \text{hadrons}$.

the Institute of High Energy Physics (Beijing). The authors also wish to thank X.J. Zhou, G.D. Zhao, S.J. Brodsky, and S.F. Tuan for enlightening discussions. This work was supported by the National Natural Science Foundation of China under Contract No. 19290400 and the Chinese Academy of Sciences under Contract No. KJ85 (IHEP), by the Department of Energy under Contracts Nos. DE-FG02-91ER40676 (Boston University), DE-FG03-92ER40701 (Caltech), DE-FG03-93ER40788 (Colorado State University), DE-AC02-76ER03069 (MIT), DE-AC03-76SF00515 (SLAC), DE-FG03-91ER40679 (UC Irvine), DE-FG03-94ER40833 (U Hawaii), DE-FG05-92ER40736 (UT Dallas), and DE-AC35-89ER40486 (SSC Lab), by the U.S. National Science Foundation, Grant No. PHY9203212 (University of Washington), and by the Texas National Research Laboratory Commission under Contracts Nos. RGFY91B5, RGFY92B5 (Colorado State), and RCFY93-316H (UT Dallas).

- [1] Particle Data Group, L. Montanet *et al.*, Phys. Rev. D **50**, 1173 (1994). Here $Q_{\rho\pi}$ is calculated by using the new measurement of BES, $B(\psi(2S) \rightarrow \rho\pi) < 3.6 \times 10^{-5}$ (90% C.L.), and $Q_{K^*\bar{K}}$ by using corrected value $B(\psi(2S) \rightarrow K^*\bar{K}) < 5.4 \times 10^{-5}$ (90% C.L.), instead of the listed value of 1.79×10^{-5} .
- [2] S. J. Brodsky, G. P. Lepage, and S. F. Tuan, Phys. Rev. Lett. **59**, 621 (1987).
- [3] W-S. Hou and A. Soni, Phys. Rev. Lett. **50**, 569 (1983); Phys. Rev. D **29**, 101 (1984); The existence of a vector gluonium state was first postulated by P. G. O. Freund and Y. Nambu, Phys. Rev. Lett. **34**, 1645 (1975).

- [4] Y. Chen, Y. F. Gu, and P. Wang, "Charm and τ Physics at BES/BEPC(I)," CCAST-WL Workshop Series Vol. 22, 1992 (unpublished), p. 98.
- [5] BES Collaboration, J. Z. Bai *et al.*, Nucl. Instrum. Methods A **344**, 319 (1994).
- [6] BES Collaboration, J. Z. Bai *et al.*, Phys. Lett. B **355**, 374 (1995).
- [7] E. A. Kuraev and V. S. Fadin, Sov. J. Nucl. Phys. **41**, 466 (1985).
- [8] M. Anselmino *et al.*, Phys. Rev. D **50**, 595 (1994).