Conical geometry and quantum entropy of a charged Kerr black hole

Robert B. Mann^{*} and Sergey N. Solodukhin[†]

Department of Physics, University of Waterloo, Waterloo, Ontario, Canada N2L 3G1

(Received 19 April 1996)

We apply the method of conical singularities to calculate the tree-level entropy and its one-loop quantum corrections for a charged Kerr black hole. The Euclidean geometry for the Kerr-Newman metric is considered. We show that for an arbitrary periodization in Euclidean space there exists a conical singularity at the horizon. Its δ -function-like curvatures are calculated and are shown to behave similar to the static case. The heat kernel expansion for a scalar field on this conical space background is derived and the (divergent) quantum correction to the entropy is obtained. It is argued that these divergences can be removed by renormalization of couplings in the tree-level gravitational action in a manner similar to that for a static black hole. [S0556-2821(96)01318-5]

PACS number(s): 04.70.Dy, 04.62.+v

I. INTRODUCTION

The notion that black holes could be considered as thermodynamic systems characterized by temperature, energy and entropy was first proposed by Bekenstein [1] and confirmed via the discovery of their thermal radiation properties by Hawking [2]. Independently, it was realized that there are only a few macroscopic parameters which can be assigned to a black hole: its mass (m), charge (q), and angular momentum (Ω) . In the static case, angular momentum vanishes. A typical representative of this class is the Reissner-Nordström black hole which is a solution of the Einstein equations with a Maxwell field as a source. Such a hole is characterized by just its mass (m) and charge (q).

When rotation is present, the Einstein-Maxwell equations have the Kerr-Newman metric [3] as a solution. This metric corresponds to a black hole of a general type characterized by all three parameters (m,q,Ω) . Remarkably, the thermodynamic analogy works for this general case; in particular, it suggests that there is an entropy associated with this hole that is proportional to the area of the horizon. If this analogy is exact, there must be hidden degrees of freedom of the hole which are counted by the Bekenstein-Hawking entropy. Recently, there has been much interest in attempting to provide a statistical explanation of these degrees of freedom and their relationship to the entropy (see reviews [4,5]) within some quantum-mechanical calculations [6–21]. However, the proposed expressions for the entropy can be considered to be quantum (one-loop) corrections to the classical quantity, and do not give any explanation of the classical entropy itself.

According to 't Hooft [6], one can relate the entropy of a black hole with a thermal gas of quantum field excitations propagating outside the horizon. In his model 't Hooft introduced a "brick wall" cutoff: a fixed boundary near the horizon within which the quantum field does not propagate. Its role is to eliminate divergences which appear due to the infinite growth of the density of states close to the horizon. This model can be successfully formulated in different space-time dimensions [7]. The quantization of a field system typically requires an ultraviolet (UV) regularization procedure that must be taken into account in the statisticalmechanical calculation as well. Remarkably, it was demonstrated in [8] that the Pauli-Villars regularization not only removes the standard field-theoretical UV divergences but automatically implements a cutoff in the 't Hooft calculation, rendering unnecessary the introduction of the "brick wall."

The natural way to formulate black hole thermodynamics is to use the Euclidean path integral approach [22]. For an arbitrary field system it entails closing the Euclidean time coordinate with a period $\beta = T^{-1}$ where *T* is the temperature of the system. This yields a periodicity condition for the quantum fields in the path integral. In the black hole case for arbitrary β this procedure leads to an effective Euclidean manifold which has a conical singularity at the horizon that vanishes for a fixed value $\beta = \beta_H$. Thermodynamic quantities (i.e., energy and entropy) are calculated by differentiating the corresponding free energy with respect to β and then setting $\beta = \beta_H$. This procedure was consistently carried out for a static black hole and resulted in obtaining the general UV-divergent structure of the entropy [18–21].

Essentially, the divergences of the entropy have the same origin as the UV-divergences of the quantum effective action and can be removed by renormalization of the gravitational couplings in the tree-level gravitational action [16,17,21]. The technique developed in [21] allowed proof of this statement for an arbitrary static black hole. Alternatively, this was demonstrated for the Reissner-Nordström black hole within 't Hooft's approach [8] applying the Pauli-Villars regularization scheme.

An essential loophole in the above considerations is that they were concerned with only static, nonrotating black holes. The only exception is a series of recent preprints [23] where 't Hooft's approach was applied to a Kerr-Newman black hole and some (qualitative) analysis of divergences was presented. Adoption of conical methods for stationary black holes necessitates dealing with problems of treating Euclideanization (or complexification) of the Kerr-Newman metric [24] and a general periodicity analysis of its conical geometry. Although the passage to a Euclidean metric and

^{*}Electronic address: mann@avatar.uwaterloo.ca

[†]Electronic address: sergey@avatar.uwaterloo.ca

periodicity arguments were given some time ago [22] the conical geometry for arbitrary period in the Kerr-Newman case remains unclear. Additional outstanding questions include the structure of the UV divergences of the entropy of stationary black holes and whether or not their renormalization works in the same way as for a static hole.

In this paper we consider these questions in detail. In Sec. II we describe the passage to Euclidean space for the Kerr-Newman metric, establishing the structure of this space in the vicinity of the horizon. We determine the conditions necessary for periodicity in the direction of the timelike Killing vector (which is analog of the Euclidean time vector ∂_{τ} in the static case) corresponding to regular Euclidean space. For arbitrary periods there is a conical singularity at the horizon surface. The geometry of this conical space is studied in Sec. III. We employ the regularization method suggested in [25] and obtain the expected δ -function-like behavior of the curvatures. The integrals of quadratic combinations of curvature tensors are also considered. The results we obtain have a marked similarity to the static case. In Sec. IV we consider the Euclidean path integral quantization of a scalar matter field in the background of a conical Kerr-Newman metric. We obtain the UV divergences for the entropy, the structure of which is similar to that obtained in the static case. We argue that these divergences of entropy are renormalized in the same way as for a static black hole.

II. EUCLIDEAN KERR-NEWMAN GEOMETRY

The Kerr-Newman metric of the space-time with Minkowskian signature in Boyer-Lindquist coordinates takes the form

$$ds^{2} = g_{rr}dr^{2} + g_{\theta\theta}d\theta^{2} + g_{tt}dt^{2} + 2g_{t\phi}dtd\phi + g_{\phi\phi}d\phi^{2},$$

$$g_{rr} = \frac{\rho^{2}}{\Delta}, \quad g_{\theta\theta} = \rho^{2}, \quad g_{tt} = -\frac{(\Delta - a^{2}\sin^{2}\theta)}{\rho^{2}},$$

$$g_{t\phi} = -\frac{a\sin^{2}\theta(r^{2} + a^{2} - \Delta)}{\rho^{2}},$$

$$g_{\phi\phi} = \left(\frac{(r^{2} + a^{2})^{2} - \Delta a^{2}\sin^{2}\theta}{\rho^{2}}\right)\sin^{2}\theta,$$

$$\Delta(r) = r^{2} + a^{2} + q^{2} - 2mr, \quad \rho^{2} = r^{2} + a^{2}\cos^{2}\theta. \quad (2.1)$$

The function $\Delta(r)$ can be represented in the form $\Delta(r) = (r - r_+)(r - r_-)$, where $r_{\pm} = m \pm \sqrt{m^2 - a^2 - q^2}$.

This space-time has a pair of orthogonal Killing vectors:

$$K = \partial_t + \frac{a}{r^2 + a^2} \partial_\phi, \quad \widetilde{K} = a \sin^2 \theta \partial_t + \partial_\phi,$$
$$K^2 = -\frac{\Delta \rho^2}{(r^2 + a^2)^2}, \quad \widetilde{K}^2 = \rho^2 \sin^2 \theta, \quad K \cdot \widetilde{K} = 0.$$
(2.2)

The vector *K* is timelike everywhere in the region $r \ge r_+$ and becomes null $K^2=0$ for $r=r_+$, whereas \widetilde{K} is spacelike everywhere outside the axis ($\theta=0, \theta=\pi$) where $\widetilde{K}^2=0$. The one-forms dual to *K* and \widetilde{K} are

$$\omega = \frac{(r^2 + a^2)}{\rho^2} (dt - a\sin^2\theta d\phi),$$

$$\widetilde{\omega} = \frac{(r^2 + a^2)}{\rho^2} \left(d\phi - \frac{a}{(r^2 + a^2)} dt \right),$$

$$\omega[K] = \widetilde{\omega}[\widetilde{K}] = 1, \quad \omega[\widetilde{K}] = \widetilde{\omega}[K] = 0.$$
(2.3)

To obtain the correspondence with the Schwarzschild metric note that K and \tilde{K} are the respective analogs of the vectors ∂_t and ∂_{ϕ} and ω and $\tilde{\omega}$ are the respective analogs of dt and $d\phi$ of the Schwarzschild metric. This correspondence is almost exact with one exception: ω and $\tilde{\omega}$ together with $d\theta$ and dr form an anholonomic basis of one-forms. This means that there are no globally defined coordinates X and \tilde{X} such that $\omega = dX$ and $\tilde{\omega} = d\tilde{X}$.

Horizon surfaces Σ are defined as the surfaces where the timelike vector K becomes null, $K^2|_{\Sigma}=0$; the outer horizon is the surface for which $r=r_+$. In addition to this one often considers the surface where the vector ∂_t becomes null. This surface is called the ergosphere and is determined by equation $r^2 + a^2 \cos^2 \theta + q^2 - 2mr=0$. It lies outside the outer horizon Σ , touching it at the axis $\theta=0$ and $\theta=\pi$.

Consider now the Euclideanization of the Kerr-Newman metric. The standard prescription says [22] that we must change the time variable $t = i\tau$ and supplement this by the parameter transformation $a = i\hat{a}, q = i\hat{q}$. The *Euclidean* vectors K, \tilde{K} and the corresponding one-forms $\omega, \tilde{\omega}$ take the form

$$K = \partial_{\tau} - \frac{\hat{a}}{r^2 - \hat{a}^2} \partial_{\phi}, \quad \widetilde{K} = \hat{a} \sin^2 \theta \partial_{\tau} + \partial_{\phi},$$
$$\omega = \frac{(r^2 - \hat{a}^2)}{\hat{\rho}^2} (d\tau - \hat{a} \sin^2 \theta d\phi),$$
$$\widetilde{\omega} = \frac{(r^2 - \hat{a}^2)}{\hat{\rho}^2} \left(d\phi + \frac{\hat{a}}{(r^2 - \hat{a}^2)} d\tau \right), \quad (2.4)$$

where $\hat{\rho}^2 = r^2 - \hat{a}^2 \cos^2 \theta$. The Euclidean metric can be written in the form

$$ds_{E}^{2} = \frac{\hat{\rho}^{2}}{\hat{\Delta}} dr^{2} + \frac{\hat{\Delta}\hat{\rho}^{2}}{(r^{2} - \hat{a}^{2})^{2}} \omega^{2} + \hat{\rho}^{2} (d\theta^{2} + \sin^{2}\theta \,\overline{\omega}^{2}), \quad (2.5)$$

where ω and $\widetilde{\omega}$ take the form (2.4), $\hat{\Delta} = r^2 - \hat{a}^2 - \hat{q}^2 - 2mr$. Roots of the function $\hat{\Delta}$ are now $\hat{r}_{\pm} = m \pm \sqrt{m^2 + \hat{a}^2 + \hat{q}^2}$. The horizon surface Σ defined by $r = \hat{r}_+$ is the stationary surface of the Killing vector K. Consider the metric (2.5) near $r = \hat{r}_+$. It is useful to introduce a new *radial* variable x such that near the horizon we have

(r-

$$\hat{\Delta} = \gamma (r - \hat{r}_{+}) = \frac{\gamma^{2} x^{2}}{4},$$
$$\hat{r}_{+} = \frac{\gamma x^{2}}{4}, \quad \gamma = 2\sqrt{m^{2} + \hat{a}^{2} + \hat{q}^{2}}.$$
(2.6)

Then the metric (2.5) up to terms $O(x^2)$ reads

$$ds_{E}^{2} = ds_{\Sigma}^{2} + \hat{\rho}_{+}^{2} \left(dx^{2} + \frac{\gamma^{2} x^{2}}{4(\hat{r}_{+}^{2} - \hat{a}^{2})^{2}} \omega^{2} \right), \qquad (2.7)$$

where $\hat{\rho}_+^2 = \hat{r}_+^2 - \hat{a}^2 \cos^2 \theta$ and

$$ds_{\Sigma}^{2} = \hat{\rho}_{+}^{2} (d\theta^{2} + \sin^{2}\theta \,\overline{\omega}^{2}) = \hat{\rho}_{+}^{2} d\theta^{2} + \frac{(\hat{r}_{+}^{2} - \hat{a}^{2})^{2}}{\hat{\rho}_{+}^{2}} \sin^{2}\theta d\psi^{2}$$
(2.8)

is metric on the horizon surface Σ . In writing Eq. (2.8) we employed the fact that on Σ we may introduce the welldefined angle coordinate $\psi = \phi + [\hat{a}/(\hat{r}_+^2 - \hat{a}^2)]\tau$. The regularity of the metric (2.8) at the points $\theta = 0, \theta = \pi$ requires the identification of the points ψ and $\psi + 2\pi$ on Σ . After all calculations with the Euclideanized Kerr-Newman metric have been completed, we analytically continue the results obtained back to the real values *a* and *q*.

Expression (2.7) may be rewritten as

$$ds_E^2 = ds_{\Sigma}^2 + \hat{\rho}_+^2 ds_{C_2}^2, \qquad (2.9)$$

where $ds_{C_2}^2$ is metric of a two-dimensional disk C_2 attached to the horizon Σ at a point (θ, ψ) :

$$ds_{C_2}^2 = dx^2 + \frac{\gamma^2 x^2}{4\hat{\rho}_+^4} (d\tau - \hat{a}\sin^2\theta d\phi)^2.$$
(2.10)

Consider the metric (2.10) with (θ, ψ) fixed. Then we may introduce an angle coordinate on C_2 , $\chi = \tau - \hat{a} \sin^2 \theta \phi$, in terms of which the metric reads

$$ds_{C_2}^2 = dx^2 + \frac{\gamma^2 x^2}{4\hat{\rho}_+^4} d\chi^2.$$
 (2.11)

Requiring the absence of a conical singularity at x=0 means that we must identify points χ and $\chi + 4\pi\gamma^{-1}\hat{\rho}_{+}^{2}$. In order for this to hold independently of the coordinate θ on the horizon we must also identify points (τ, ϕ) with $(\tau+2\pi\beta_{H}, \phi-2\pi\Omega\beta_{H})$, where $\Omega = \hat{a}/(\hat{r}_{+}^{2} - \hat{a}^{2})$ is the (complex) angular velocity and $\beta_{H} = (\hat{r}_{+}^{2} - \hat{a}^{2})/\sqrt{m^{2} + \hat{a}^{2} + \hat{q}^{2}}$. It is easy to see that the identified points have the same coordinate ψ .

With the described identification we obtain the following picture of the Euclidean Kerr-Newman geometry in the vicinity of the horizon Σ . Attached to every point (θ, ψ) of the horizon is a two-dimensional disk C_2 with coordinates (x,χ) . Although χ is not a global coordinate in fourdimensional space and at each point (θ, ψ) there is a new χ , the periodic identification of points on C_2 works universally and independently of any point on the horizon Σ . As in the static case, the Euclidean Kerr-Newman geometry possesses an Abelian isometry generated by the Killing vector K with horizon surface Σ being a fixed set of the isometry. Globally, K is not a coordinate vector. However, locally we have $K = \partial_{\chi}$ where χ was introduced above. The periodicity is in the direction of the vector K and the resulting Euclidean space E is a regular manifold.

III. CONICAL SINGULARITY AND CURVATURE TENSORS

Assume now that we close the trajectory of the Killing vector *K* with arbitrary period $2\pi\beta$. Near the horizon this means that on C_2 in Eqs. (2.9) and (2.10) we identify points (τ, ϕ) and $(\tau + 2\pi\beta, \phi - 2\pi\Omega\beta)$ with $\beta \neq \beta_H$. Note again that points identified in this way have the same value of the coordinate ψ . Then χ is an angle coordinate with period $2\pi\beta(1+\hat{a}\Omega\sin^2\theta)$. By introducing a new angle coordinate $\chi = \beta\hat{\rho}_+^2(\hat{r}_+^2 - \hat{a}^2)^{-1}\overline{\chi}$ which has period 2π , the metric on C_2 becomes

$$ds_{C_{2,\alpha}}^{2} = dx^{2} + \alpha^{2}x^{2}d\bar{\chi}^{2}$$
(3.1)

and coincides with the metric on a two-dimensional cone with angular deficit $\delta = 2\pi(1-\alpha)$, $\alpha = \beta/\beta_H$. With the above identification the four-dimensional (4D) metric (2.5) describes the Euclidean conical space E_{α} with singular surface Σ .

Curvature tensors at conical singularities behave as distributions. This behavior was precisely established for a flat 2D cone in [26] and for a general static metric in [25]. The Kerr-Newman metric, which is the subject of our consideration here, is stationary and not static. Therefore, all the formulas obtained in [25] must be checked for this case.

To proceed, we apply the method which was successful in the static case (see details in [25]). It consists in regulating the conical singularity when the cone metric (3.1) is replaced by a sequence of regular metrics labeled by a parameter b:

$$ds_{C_{2,\alpha,b}}^{2} = f(x,b)dx^{2} + \alpha^{2}x^{2}d\overline{\chi}^{2}, \qquad (3.2)$$

where f(x,b) is some smooth regulating function that approaches unity as $b \rightarrow 0$, e.g.,

$$f(x,b) = \frac{x^2 + \alpha^2 b^2}{x^2 + b^2}$$
(3.3)

is a suitable regularization function. In the limit $b \rightarrow 0$ the sequence of metrics (3.2) reproduces a δ -function-like contribution to the curvature.

Applying this method to the Kerr metric, consider a small vicinity of the horizon surface Σ . For $\beta \neq \beta_H$ the metric there reads

$$ds_{E_{\alpha}}^{2} = ds_{\Sigma}^{2} + \hat{\rho}_{+}^{2} ds_{C_{\alpha}}^{2}.$$
(3.4)

Replacing the cone metric $ds_{C_{\alpha}}^2$ by $ds_{C_{\alpha,b}}^2$ [Eq. (3.2)] we obtain a sequence of regular metrics:

$$ds_{E_{\alpha,b}}^{2} = ds_{\Sigma}^{2} + \hat{\rho}_{+}^{2} ds_{C_{\alpha,b}}^{2}.$$
 (3.5)

To calculate the curvature we define the (anholonomic) basis of one-forms $\{e^a, a = 1, ..., 4\}$ orthonormal with respect to the metric (3.5):

$$e^{1} = b\hat{\rho}_{+}f^{1/2}(y)dy,$$
 (3.6)

$$e^2 = by \frac{(m^2 + \hat{a}^2 + \hat{q}^2)^{1/2}}{\hat{
ho}_+} (d\tau - \hat{a}\sin^2\theta d\phi),$$

$$e^{3} = \hat{\rho}_{+} d\theta,$$
$$e^{4} = \frac{(\hat{r}_{+}^{2} - \hat{a}^{2})}{\hat{\rho}_{+}} \sin\theta \left(d\phi + \frac{\hat{a}}{(\hat{r}_{+}^{2} - \hat{a}^{2})} d\tau \right)$$

where we changed variables so that $x=by, f(y)=(y^2+\alpha^2)/(y^2+1).$

The Lorentz connection one-form $\omega_b^a = \omega_{bc}^a e^c$ is found from the equation

$$de^a + \omega_b^a \wedge e^b = 0. \tag{3.7}$$

We are interested in those components of the Lorentz connection which are singular in the limit $b \rightarrow 0$. Analyzing the expressions de^a for the basis (3.6) we observe that only de^2 contains a singular term:

$$de^2 = \frac{dy}{y} \wedge e^2 + \dots = [byf^{1/2}(y)]^{-1}e^1 \wedge e^2 + \dots,$$
 (3.8)

where the ellipsis means terms finite in the limit $b \rightarrow 0$. It follows from Eq. (3.8) that the only singular component of the Lorentz connection is

$$\omega_1^2 = [by\hat{\rho}_+ f^{1/2}(y)]^{-1}e^2 + \cdots.$$
 (3.9)

The curvature two-form $R_b^a = R_{bcd}^a e^c \wedge e^d$ is defined as

$$R_b^a = d\omega_b^a + \omega_c^a \wedge \omega_b^c$$

Again, the only singular component of R_b^a is

$$R_{1}^{2} = d\omega_{1}^{2} + \dots = \frac{1}{2yb^{2}\hat{\rho}_{+}^{2}} \frac{f'_{y}}{f^{2}}e^{2} \wedge e^{1} + \dots$$

and so in terms of curvature components the only singular component is

$$R_{2121} = \frac{1}{2yb^2\hat{\rho}_+^2} \frac{f_y'}{f^2} + \cdots .$$
 (3.10)

Introducing a pair of vectors [see Appendix Eqs. (A1) and (A2)] $n_a = n_a^{\mu} \partial_{\mu}, a = 1,2$ orthogonal to the horizon Σ and dual to the one-forms e^a , a = 1,2 we may rewrite $R_{2121} = \frac{1}{2} R_{\mu\nu\alpha\beta} n_a^{\mu} n_b^{\nu} n_a^{\alpha} n_b^{\beta}$

In order to show that the component R_{2121} behaves as a δ function in the limit $b \rightarrow 0$, let us consider the integral

$$I_{D_{\epsilon}} = \int_{D_{\epsilon}} R_{2121} v(x,\theta,\psi) e^{1} \wedge e^{2} \wedge e^{3} \wedge e^{4} \qquad (3.11)$$

over a small disk D_{ϵ} surrounding the horizon surface Σ , $0 \le x \le \epsilon$. In Eq. (3.11) $v(x, \theta, \psi)$ is a test function which is constant along the trajectory of the vector *K* (*K*[*v*]=0). It can be expanded as

$$v(x, \theta, \psi) = v_0(\theta, \psi) + v_1(\theta, \psi)x^2 + \cdots$$
$$= v_0(\theta, \psi) + b^2 v_1(\theta, \psi)y^2 + \cdots$$

Recall that (θ, ψ) are the coordinates on the horizon Σ . Substituting Eqs. (3.6) and (3.10) into Eq. (3.11) we obtain

$$I_{D_{\epsilon}} = \int_{0}^{\epsilon/b} dy \, \frac{f'_{y}}{2f^{3/2}} \sqrt{m^{2} + \hat{a}^{2} + \hat{q}^{2}} \oint (d\tau) \\ - \hat{a}\sin^{2}\theta d\phi \int_{\Sigma} \frac{1}{\hat{\rho}_{+}^{2}} (v_{0} + v_{1}b^{2}y^{2} + \cdots)e^{3} \wedge e^{4}.$$
(3.12)

In Eq. (3.12) we first integrate over $e^1 \wedge e^2$ in the subspace orthogonal to Σ under fixed (θ, ψ) and then take the integral $\int_{\Sigma} e^3 \wedge e^4$ over the horizon. This yields

$$\oint (d\tau - \hat{a}\sin^2\theta d\phi) = \frac{2\pi\beta\hat{\rho}_+^2}{\hat{r}_+^2 - \hat{a}_+^2},$$
(3.13)

where the integration is taken over the closed integral trajectory of the Killing vector K under fixed (θ, ψ) .

In the limit $b \rightarrow 0$, we have $\epsilon/b \rightarrow \infty$ in the y integration in Eq. (3.12) and so obtain

$$\int_{0}^{\infty} dy \frac{f'_{y}}{2f^{3/2}} = -f^{-1/2}(y)\big|_{0}^{\infty} = \frac{1-\alpha}{\alpha}.$$
 (3.14)

Taking into account that $\beta_H = (\hat{r}_+^2 - \hat{a}^2)/\sqrt{m^2 + \hat{a}^2 + \hat{q}^2}$ from Eqs. (3.12)–(3.14) we finally obtain, in the limit $b \rightarrow 0$,

$$I_{D_{\epsilon}} = 2\pi(1-\alpha) \int_{\Sigma} v_0(\theta, \psi) e^3 \wedge e^4.$$
 (3.15)

Since this holds for arbitrarily small ϵ we conclude that in the limit $b \rightarrow 0$ the quantity R_{2121} behaves as a δ function having support at the surface Σ . Noting that the vectors $n_a, a = 1,2$ introduced above are normal to Σ we may write

$$R^{\mu\nu}_{\alpha\beta} = R^{\mu\nu}_{\alpha\beta} + 2\pi(1-\alpha)[(n^{\mu}n_{\alpha})(n^{\nu}n_{\beta}) - (n^{\mu}n_{\beta})(n^{\nu}n_{\alpha})]\delta_{\Sigma},$$

$$R^{\mu}_{\nu} = \overline{R}^{\mu}_{\nu} + 2\pi(1-\alpha)(n^{\mu}n_{\nu})\delta_{\Sigma},$$

$$R = \overline{R} + 4\pi(1-\alpha)\delta_{\Sigma}, \qquad (3.16)$$

where δ_{Σ} is the delta function $\int_{\mathcal{M}} f \delta_{\Sigma} e^{1} \wedge e^{2} \wedge e^{3} \wedge e^{4} = \int_{\Sigma} f e^{3} \wedge e^{4}$ and we denote $(n_{\mu}n_{\nu}) = \Sigma_{a=1}^{2} n_{\mu}^{a} n_{\nu}^{a}$. In particular, it follows from Eq. (3.16) that

$$\int_{E_{\alpha}} Re^{1} \wedge e^{2} \wedge e^{3} \wedge e^{4}$$
$$= \alpha \int_{E} \overline{R}e^{1} \wedge e^{2} \wedge e^{3} \wedge e^{4} + 4\pi(1-\alpha)A_{\Sigma}, \qquad (3.17)$$

where $A_{\Sigma} = \int_{\Sigma} e^3 \wedge e^4$ is area of Σ . For the particular case of the Kerr-Newman metric, $\overline{R} = 0$. Remarkably, expressions (3.16) and (3.17) are exactly the same as that obtained in [21] for a static metric.

For a variety of applications it is necessary to know the integrals of quadratic combinations of curvatures over the space E_{α} with a conical singularity at the surface Σ . According to Eq. (3.16), \mathcal{R} contains $(1 - \alpha) \delta_{\Sigma}$ contributions as well as a regular part $\overline{\mathcal{R}}$. Therefore, one can expect a result which can be symbolically written as follows:

$$\int_{E_{\alpha}} \mathcal{R}^2 = \alpha \int_{E_{\alpha=1}} \overline{\mathcal{R}}^2 + 2(1-\alpha) \int_{\Sigma} \overline{\mathcal{R}}_n + O((1-\alpha)^2),$$
(3.18)

where $\overline{\mathcal{R}}_n$ means projection of the tensor $\overline{\mathcal{R}}$ onto the sub-

space orthogonal to the singular surface Σ . The expression (3.18) is ill defined since \mathcal{R}^2 contains term $[(1-\alpha)\delta_{\Sigma}]^2$. However, it is of higher order with respect to $(1-\alpha)$ and so can be collected in the last term in Eq. (3.18).

The form (3.18) was obtained in [25] for a static metric. To verify this for the Kerr-Newman case we must write the metric (2.5) near the horizon Σ , including all terms of order x^2 . Taking into account the regularization function f(x,b) as above the metric reads

$$ds_{E_{\alpha,b}}^{2} = byf(y)dy^{2} + \frac{\gamma^{2}b^{2}y^{2}}{4\hat{\rho}_{+}^{2}}(d\tau - \hat{a}\sin^{2}\theta d\phi)^{2} + \left(\hat{\rho}_{+}^{2} + \frac{\gamma}{2}b^{2}y^{2}\hat{r}_{+}\right)d\theta^{2} + \left[\frac{(\hat{r}_{+}^{2} - \hat{a}^{2})^{2}}{\hat{\rho}_{+}^{2}} + \frac{(\hat{r}_{+}^{2} - \hat{a}^{2})}{\hat{\rho}_{+}^{2}}\right] \times \left(1 - \frac{(\hat{r}_{+}^{2} - \hat{a}^{2})}{2\hat{\rho}_{+}^{2}}\right)\gamma\hat{r}_{+}b^{2}y^{2} \sin^{2}\theta \left(d\phi + \frac{\hat{a}}{\hat{r}_{+}^{2} - \hat{a}^{2}}d\tau\right)^{2} - \frac{\gamma\hat{a}b^{2}y^{2}}{2\hat{\rho}_{+}^{2}}\sin^{2}\theta \left(d\phi + \frac{\hat{a}}{\hat{r}_{+}^{2} - \hat{a}^{2}}d\tau\right)d\tau.$$
(3.19)

The general structure of quadratic combinations of curvature terms (denoted by \mathcal{R}^2) for the metric (3.19) symbolically is

$$\mathcal{R}^{2} = b^{2}A + \frac{f_{y}'}{b^{2}}B + \frac{(f_{y}')^{2}}{b^{4}}C + O(b^{4}), \qquad (3.20)$$

where A, B, C are some functions that are independent of b and do not contain derivatives of the regularization function f(y).

Since the measure of integration in the region near Σ is proportional to b^2 we conclude that second and third terms in Eq. (3.20) after integration produce in the limit $b \rightarrow 0$ the respective second and third terms in Eq. (3.18). In order to get this we use the fact that the derivatives of f(y) behave as $f'(y) \sim (1-\alpha)$.

After straightforward but tedious calculations we obtain, in the limit $b \rightarrow 0$,

$$\int_{E_{\alpha}} R^{\mu\nu} R_{\mu\nu} = \alpha \int_{E} \overline{R}^{\mu\nu} \overline{R}_{\mu\nu} + 4\pi(1-\alpha)$$
$$\times \int_{\Sigma} \overline{R}_{aa} + O((1-\alpha)^{2}), \qquad (3.21)$$

$$\int_{E_{\alpha}} R^{\mu\nu\lambda\rho} R_{\mu\nu\lambda\rho} = \alpha \int_{E} \overline{R}^{\mu\nu\lambda\rho} \overline{R}_{\mu\nu\lambda\rho} + 8\pi(1-\alpha)$$
$$\times \int_{\Sigma} \overline{R}_{abab} + O((1-\alpha)^{2}), \quad (3.22)$$

where $\overline{R}_{aa} = \sum_{a=1,2} \overline{R}_{\mu\nu} n_a^{\mu} n_a^{\nu}$ and \overline{R}_{abab} = $\sum_{a,b=1,2} \overline{R}_{\mu\nu\lambda\rho} n_a^{\mu} n_a^{\lambda} n_b^{\nu} n_b^{\rho}$.

In obtaining Eqs. (3.21) and (3.22) we made use of the fact [as in Eq. (3.12)] that near Σ the measure of integration $\mu_{E_{\alpha,b}}$ takes the form [see Eq. (2.5)] $\mu_{E_{\alpha,b}} = \hat{\rho}_{+}^2 \mu_{\Sigma} \mu_{C_{\alpha,b}}$, where $\mu_{\Sigma} = (\hat{r}_{+}^2 - \hat{a}^2) \sin\theta d\theta d\psi$ ($0 \le \psi \le 2\pi$) is the measure

on Σ and $\mu_{C_{\alpha,b}} = \alpha b^2 f^{1/2}(y) dy d\overline{\chi}$ $(0 \le \overline{\chi} \le 2\pi)$ is the measure on the regularized cone $C_{\alpha,b}$. For the integral of R^2 we obtain

$$\int_{E_{\alpha}} R^2 = O((1-\alpha)^2)$$

in agreement with the expected formula

$$\int_{E_{\alpha}} R^{2} = \alpha \int_{E} \overline{R}^{2} + 8 \pi (1 - \alpha) \int_{\Sigma} \overline{R} + O((1 - \alpha)^{2}), \quad (3.23)$$

since the Kerr-Newman metric satisfies $\overline{R} = 0$.

Again we obtain for a stationary metric with a conical singularity the same expressions (3.21)-(3.23) as for the static case [25].

For the Kerr-Newman metric we have, on the horizon Σ ,

$$\frac{1}{2}\overline{R}_{abab} = \overline{R}_{2121} = \frac{\hat{r}_{+}^{2}(4\hat{q}^{2} + 8m\hat{r}_{+}) - (\hat{q}^{2} + 6m\hat{r}_{+})\hat{\rho}^{2}}{\hat{\rho}_{+}^{6}},$$
$$\frac{1}{2}\overline{R}_{aa} = \overline{R}_{11} = \overline{R}_{22} = \frac{\hat{q}^{2}}{\hat{\rho}_{+}^{4}}$$
(3.24)

and after integration over Σ we get

$$\int_{\Sigma} \overline{R}_{abab} = 8\pi \frac{(\hat{r}_{+}^{2} + \hat{q}^{2})}{\hat{r}_{+}^{2}} + 4\pi \frac{\hat{q}^{2}}{\hat{r}_{+}^{2}} \frac{(\hat{r}_{+}^{2} - \hat{a}^{2})}{\hat{a}\hat{r}_{+}} \ln\left(\frac{\hat{r}_{+} + \hat{a}}{\hat{r}_{+} - \hat{a}}\right),$$
$$\int_{\Sigma} \overline{R}_{aa} = 4\pi \frac{\hat{q}^{2}}{\hat{r}_{+}^{2}} \left[1 + \frac{(\hat{r}_{+}^{2} - \hat{a}^{2})}{2\hat{a}\hat{r}_{+}} \ln\left(\frac{\hat{r}_{+} + \hat{a}}{\hat{r}_{+} - \hat{a}}\right)\right]. \quad (3.25)$$

The analytic continuation of these expressions back to real values of the parameters a and q requires the substitution

$$\hat{q}^2 = -q^2, \quad \hat{a}^2 = -a^2, \quad \hat{r}_+ = r_+,$$

 $\frac{1}{\hat{a}} \ln \left(\frac{\hat{r}_+ + \hat{a}}{\hat{r}_+ - \hat{a}} \right) = \frac{2}{a} \tan^{-1} \left(\frac{a}{r_+} \right).$ (3.26)

IV. HEAT KERNEL EXPANSION AND ENTROPY

In the Euclidean path integral approach to a statistical field system at temperature $T = (2\pi\beta)^{-1}$ one considers the fields which are periodic with respect to imaginary time τ with period $2\pi\beta$. This works well for a static black hole when the metric does not depend on the time coordinate τ [22]. One then closes the integral curves of the Killing vector ∂_{τ} with the period $2\pi\beta$.

For a rotating black hole metric we need to close the integral curves of the vector K (2.4). The result of this is that for arbitrary β we obtain the conical space E_{α} , the geometry of which was described in the previous section. The partition function then reads

$$Z(\beta) = \int [\mathcal{D}\varphi] \exp[-I_E(\varphi, g_{\mu\nu})], \qquad (4.1)$$

where the matter Euclidean action I_E is considered on the space E_{α} with appropriate boundary (i.e., periodicity) conditions imposed on the matter field(s) φ . The contribution to the entropy is

$$S = -(\beta \partial_{\beta} - 1) \ln Z(\beta) |_{\beta = \beta_{\mu}}.$$
(4.2)

Although the Kerr-Newman metric is a solution of the Einstein equations with the matter source in form of a Maxwell field, the gravitational action is always modified by higher-order curvature terms due to quantum corrections. Such R^2 terms must be added to the action at the outset with some bare constants ($c_{1,B}$, $c_{2,B}$, $c_{3,B}$) (tree-level) to absorb the one-loop infinities. The bare (tree-level) gravitational action functional thus takes the form

$$W_{\rm gr} = \int \sqrt{g} d^4 x \left(-\frac{1}{16\pi G_B} R + c_{1,B} R^2 + c_{2,B} R_{\mu\nu} R^{\mu\nu} + c_{3,B} R_{\mu\nu\alpha\beta} R^{\mu\nu\alpha\beta} \right).$$
(4.3)

Of course, we assume in addition to Eq. (4.3) a classical matter action which can in principle be rather complicated. The corresponding tree-level entropy can be obtained as a replica of the action (4.3) after introducing the regulated conical singularity and applying formulas (4.1)-(4.2). Using formulas (3.21)-(3.23) of the previous section we obtain, for the tree-level entropy,

$$S(G_B, c_{i,B}) = \frac{1}{4G_B} A_{\Sigma} - \int_{\Sigma} (8 \pi c_{1,B} \overline{R} + 4 \pi c_{2,B} \overline{R}_{aa} + 8 \pi c_{3,B} \overline{R}_{abab}), \qquad (4.4)$$

where $\overline{R}_{aa} = \sum_{a=1,2} \overline{R}_{\mu\nu} n_a^{\mu} n_a^{\nu}$ and \overline{R}_{abab} = $\sum_{a,b=1,2} \overline{R}_{\mu\nu\lambda\rho} n_a^{\mu} n_a^{\lambda} n_b^{\nu} n_b^{\rho}$, $\{n_a^{\mu}, a=1,2\}$ are vectors normal to Σ . This is exactly the same expression that we had for the static case [27,21]. Expression (4.4) is really valid off-shell, as we do not require the metric to satisfy any equations of motion. On-shell we must substitute in Eq. (4.4) the field equation $\overline{R}=0$ satisfied by the Kerr-Newman metric.

At the one-loop level we consider the matter action in the form

$$I_E = \frac{1}{2} \int_{E_{\alpha}} (\nabla \varphi)^2$$

and get for the partition function

$$\ln Z(\beta) = -\frac{1}{2} \operatorname{Indet}(-\Box_{E_{\alpha}})$$

expressed via the determinant of the Laplacian $\Box = \nabla_{\mu} \nabla^{\mu}$ over the conical space E_{α} . In the DeWitt-Schwinger proper time representation we have, for the logarithm of the determinant,

$$\operatorname{Indet}(-\Box) = -\int_{\epsilon^2}^{\infty} \frac{ds}{s} \operatorname{Tr}(e^{s\Box}).$$
(4.5)

In four dimensions we have the asymptotic expansion

$$\operatorname{Tr}(e^{s\Box}) = \frac{1}{(4\pi s)^2} \sum_{n=0}^{\infty} a_n s^n$$
(4.6)

and for the divergent part of $(\ln Z)_{div}$ we get

$$(\ln Z)_{\rm div} = \frac{1}{32\pi^2} \left(\frac{1}{2} a_0 \epsilon^{-4} + a_1 \epsilon^{-2} + 2a_2 \ln \frac{L}{\epsilon} \right), \quad (4.7)$$

where L is an infrared cutoff. It is known that for a manifold with conical singularities the heat kernel coefficients in Eq. (4.6) are really a sum

$$a_n = a_n^{\text{st}} + a_{n,\alpha} \tag{4.8}$$

of standard plus conical coefficients. The standard coefficients \bar{a}_n^{st} are the same as for a smooth manifold *E* [28]:

$$a_0^{\text{st}} = \int_{E_{\alpha}} 1, \quad a_1^{\text{st}} = \frac{1}{6} \int_{E_{\alpha}} \overline{R},$$

$$a_2^{\text{st}} = \int_{E_{\alpha}} \left(\frac{1}{180} \overline{R}_{\mu\nu\alpha\beta} \overline{R}^{\mu\nu\alpha\beta} - \frac{1}{180} \overline{R}_{\mu\nu} \overline{R}^{\mu\nu} - \frac{1}{30} \Box \overline{R} + \frac{1}{72} \overline{R}^2 \right), \quad (4.9)$$

whereas the parts coming from the singular surface Σ (stationary point of the isometry) are

$$a_{0,\alpha} = 0; \quad a_{1,\alpha} = \frac{\pi}{3} \frac{(1-\alpha^2)}{\alpha} \int_{\Sigma} \sqrt{\gamma} d^2 \theta; \qquad (4.10)$$

$$+\frac{1}{2}\kappa^{a}\kappa^{a}+\lambda_{2}[\kappa^{a}\kappa^{a}-2\operatorname{tr}(\kappa\cdot\kappa)]\bigg)\sqrt{\gamma}d^{2}\theta,$$

 $\lambda_{1,2}$ are some constants and $\kappa^a_{\mu\nu}, a=1,2$ is the extrinsic curvature of the surface Σ with respect to normal vector $n_a, a=1,2; \ \kappa^a = g^{\mu\nu}\kappa^a_{\mu\nu} \ tr(\kappa \cdot \kappa) = \sum_{a=1,2} \kappa^a_{\mu\nu}\kappa^{\mu\nu}_a$. The expression (4.10) for some special spaces has been

known for some time [29]. For a general static metric it was derived recently by Fursaev [30], in the case that all extrinsic curvatures $\kappa^{a}_{\mu\nu}$ vanished. Dowker [31] derived the heat kernel coefficients in the form (4.10) for an arbitrary conical metric of a general type with a surface Σ having nontrivial extrinsic geometry. Very general arguments were used in [31] to derive the general structure of Eq. (4.10): O(2) invariance, dimensionality, and conformal invariance. The result (4.10) contains some unknown constants λ_1 and λ_2 in of front the conformal-invariant combination $[\kappa^a \kappa^a - 2 \operatorname{tr}(\kappa \cdot \kappa)]$. The analysis of [31] does not provide a prescription for obtaining the explicit values for these constants.

Applying the formula (4.2) to Eq. (4.7) and taking into account that the standard coefficients $a_n^{\text{st}} \sim \alpha$ we obtain for the divergent quantum correction to the entropy

$$S_{\text{div}} = \frac{1}{48\pi\epsilon^2} A_{\Sigma} + \left[\frac{1}{144\pi} \int_{\Sigma} \overline{R} - \frac{1}{16\pi} \frac{1}{45} \int_{\Sigma} (\overline{R}_{aa} - 2\overline{R}_{abab}) - \frac{1}{16\pi} \frac{1}{90} \int_{\Sigma} \kappa^a \kappa^a + \frac{1}{24\pi} \left(\lambda_1 - \frac{\lambda_2}{30} \right) \\ \times \int_{\Sigma} \left[\kappa^a \kappa^a - 2\operatorname{tr}(\kappa \cdot \kappa) \right] \left] \ln \frac{L}{\epsilon}.$$

$$(4.11)$$

We see that the divergent part of the entropy (4.11) depends both on the projections of the curvatures, \overline{R}_{aa} and \overline{R}_{abab} , onto the subspace normal to the horizon surface Σ and on the quadratic combinations of the extrinsic curvatures of Σ . For the static case all extrinsic curvatures vanish and Eq. (4.11) repeats the form of the tree-level entropy (4.4). This fact allows one to prove [21] for arbitrary static black holes the statement [16,18] that all the UV divergences of entropy are absorbed in the standard renormalization of the gravitational couplings (G,c_i) in the tree-level gravitational action (4.3). Applying the same line of reasoning to the Kerr-Newman black hole entails studying the external geometry of the horizon Σ of the charged rotating black hole. For this case we find that $\sum_{a=1,2} \kappa^a \kappa^a = tr(\kappa \cdot \kappa) = 0$ (see Appendix). This makes the coefficients (4.10) and the expression for S_{div} (4.11) for the Kerr-Newman metric the same as for a static metric.

Consequently, S_{div} in Eq. (4.11) repeats the form of the tree-level entropy and the *renormalization* statement is valid for a stationary hole as well. In one sense this is not surprising since the classical thermodynamics of static and station-

ary holes are formulated in the same way. One could therefore expect this also to be valid in the quantum case.

Consider Eq. (4.11) on the Kerr-Newman background. Substituting here Eq. (3.25), the condition $\overline{R} = 0$, and making the analytic continuation (3.26), we finally obtain, for the quantum entropy of the Kerr-Newman black hole,

$$S_{\rm div} = \frac{1}{48\pi\epsilon^2} A_{\Sigma} + \frac{1}{45} \left\{ 1 - \frac{3q^2}{4r_+^2} \left[1 + \frac{(r_+^2 + a^2)}{ar_+} \tan^{-1} \left(\frac{a}{r_+}\right) \right] \right\} \ln\frac{L}{\epsilon},$$
(4.12)

where $A_{\Sigma} = 4\pi(r_{+}^2 + a^2)$ is area of the horizon Σ . In the limit $a \rightarrow 0$ this expression reduces to that of the Reissner-Nordström black hole obtained in [20] using the conical method and in [8] within the framework of a statistical-mechanical calculation in spirit of 't Hooft's approach. Surprisingly, in the uncharged case (q=0) the quantum correction (4.12) does not depend on the rotation parameter a and coincides with the quantum entropy of the Schwarzchild black hole [18]. We do not have an explanation of this fact.

V. CONCLUSIONS

The Euclidean approach to black hole thermodynamics implying the conical singularity method is known to be very useful in the static case. It allows one to get both the classical and quantum thermodynamic quantities of static black holes. We have proposed that the thermodynamics of the classical static and stationary black holes are formulated in a similar way. The underlying assumption is that the conical singularity technique can be applied to the rotating hole as well.

In this paper we logically followed this line of reasoning. We studied the Euclidean geometry of the Kerr-Newman metric for an arbitrary period along the timelike Killing vector generating the Abelian isometry of the space. The conical geometry of the space near the horizon was established and the δ -function-like behavior of the curvatures obtained. This behavior strongly resembles that of a static black hole.

The essential point of formulating the quantum thermodynamics of static black hole is the proving the statement that all the UV divergences of the entropy of black hole due to quantum matter are removed by the standard renormalization of couplings in the tree-level gravitational action. This allows one to consider the entropy as a well-defined quantum field theoretical quantity. We demonstrated for the Kerr-Newman black hole that S_{div} being expressed via geometrical invariants repeats the form of the tree-level entropy in the same way as for a static case. This proves that the *renormalization* statement works universally both for the static and stationary holes providing the correct treatment of the quantum thermodynamics.

However, it is still an open question as to what degrees of freedom are counted by the entropy of black hole. A useful approach to this problem is to compare our result with the statistical-mechanical calculation of the quantum entropy of Kerr-Newman black hole along the lines of [6-8]. For a charged nonrotating black hole it is known that there is perfect agreement between these two methods (see [8] and

[20]). Checking this for stationary case¹ should provide us with a better understanding of the relationship between the different entropies assigned to a black hole [32].

ACKNOWLEDGMENTS

This work was supported by the Natural Sciences and Engineering Research Council of Canada and by NATO.

APPENDIX: EXTRINSIC GEOMETRY OF HORIZON

With respect to the Euclidean metric (2.5) we may define a pair of orthonormal vectors $\{n_a = n_a^{\mu} \partial_{\mu}, a = 1, 2\}$:

$$n_1^r = \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^2}},\tag{A1}$$

$$n_2^{\tau} = \frac{(r^2 - \hat{a}^2)}{\sqrt{\hat{\Delta}\hat{\rho}^2}}, \quad n_2^{\phi} = \frac{-\hat{a}}{\sqrt{\hat{\Delta}\hat{\rho}^2}}.$$
 (A2)

Covariantly these are

$$n_r^1 = \sqrt{\frac{\hat{\rho}^2}{\hat{\Delta}}},\tag{A3}$$

$$n_{\tau}^2 = \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^2}}, \quad n_{\phi}^2 = -\sqrt{\frac{\hat{\Delta}}{\hat{\rho}^2}}\hat{a}\sin^2\theta.$$
 (A4)

The vectors n^1 and n^2 are normal to the horizon surface Σ [defined as $r=r_+, \Delta(r=r_+)=0$], which is a twodimensional surface with induced metric $\gamma_{\mu\nu}=g_{\mu\nu}-n_{\mu}^2n_{\nu}^2-n_{\mu}^2n_{\nu}^2$. The (nonzero) components of the induced metric are

$$\gamma_{\theta\theta} = \rho^2, \quad \gamma_{\tau\tau} = \frac{\hat{a}^2 \sin^2 \theta}{\hat{\rho}^2},$$
$$\gamma_{\tau\phi} = \frac{\hat{a}(r^2 - \hat{a}^2) \sin^2 \theta}{\hat{\rho}^2},$$

¹The recent statistical calculation performed in [23] appears to be unsatisfactory since it relates the entropy of rotating hole with data on the ergosphere rather than on the horizon.

$$\gamma_{\phi\phi} = \frac{(r^2 - \hat{a}^2)^2 \sin^2 \theta}{\hat{\rho}^2}.$$
 (A5)

With respect to the normal vectors $n^a, a = 1,2$ we may define [33] the extrinsic curvatures of the surface Σ : $\kappa^a_{\mu\nu} = -\gamma^{\alpha}_{\mu} \gamma^{\beta}_{\nu} \nabla_{\alpha} n^a$. We find

$$\kappa_{\theta\theta}^{1} = -r \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^{2}}},$$

$$\kappa_{\tau\tau}^{1} = \frac{r\hat{a}^{2}\sin^{2}\theta}{\hat{\rho}^{4}} \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^{2}}},$$

$$\kappa_{\tau\phi}^{1} = -\frac{\hat{a}r(r^{2} - \hat{a}^{2})\sin^{2}\theta}{\hat{\rho}^{4}} \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^{2}}},$$

$$\kappa_{\phi\phi}^{1} = -\frac{r(r^{2} - \hat{a}^{2})^{2}\sin^{2}\theta}{\hat{\rho}^{4}} \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^{2}}},$$
(A6)

and

$$\kappa_{r\tau}^{2} = -\frac{\hat{a}^{2}\sin\theta\cos\theta}{\hat{\rho}^{2}}\sqrt{\frac{\Delta}{\hat{\rho}^{2}}},$$

$$\kappa_{r\phi}^{2} = -\frac{\hat{a}(r^{2} - \hat{a}^{2})\sin\theta\cos\theta}{\hat{\rho}^{2}}\sqrt{\frac{\Delta}{\hat{\rho}^{2}}}.$$
(A7)

For the trace of the extrinsic curvatures, $\kappa^a = \kappa^a_{\mu\nu} g^{\mu\nu}$, we obtain

$$\kappa^{1} = -\frac{2r}{\hat{\rho}^{2}} \sqrt{\frac{\hat{\Delta}}{\hat{\rho}^{2}}}, \quad \kappa^{2} = 0,$$
(A8)

which clearly vanishes when restricted to the surface $\Sigma[\Delta(r=\hat{r}_+)=0]$.

The quadratic combinations

$$\kappa_{\mu\nu}^{1}\kappa_{1}^{\mu\nu} = \frac{2r^{2}\hat{\Delta}}{\hat{\rho}^{6}},$$

$$\kappa_{\mu\nu}^{2}\kappa_{2}^{\mu\nu} = \frac{2\hat{a}^{2}\cos^{2}\theta\hat{\Delta}}{\hat{\rho}^{6}}$$
(A9)

vanish Σ separately both in the static (a=0) and stationary $(a\neq 0)$ cases. Consequently, we have $\operatorname{tr}(\kappa \cdot \kappa) = \kappa^a_{\mu\nu} \kappa^{a\mu\nu} = 0$ on the horizon.

- J. D. Bekenstein, Lett. Nuovo Cimento 4, 737 (1972); Phys. Rev. D 7, 2333 (1973); 9, 3292 (1974).
- [2] S. W. Hawking, Commun. Math. Phys. 43, 199 (1975).
- [3] R. P. Kerr, Phys. Rev. Lett. 11, 237 (1963); E. T. Newman et al., J. Math. Phys. 6, 918 (1965).
- [4] J. D. Bekenstein, "Do we understand black hole entropy?" Report No. qr-qc/9409015 (unpublished).
- [5] V. P. Frolov, "Black hole entropy and physics at Planckian scales," Report No. hep-th/9510156 (unpublished).
- [6] G. 't Hooft, Nucl. Phys. B256, 727 (1985).

- [7] R. B. Mann, L. Tarasov, and A. Zelnikov, Class. Quantum Grav. 9, 1487 (1992).
- [8] J. -G. Demers, R. Lafrance, and R. C. Myers, Phys. Rev. D 52, 2245 (1995).
- [9] M. Srednicki, Phys. Rev. Lett. **71**, 666 (1993); L. Bombelli, R. Koul, J. Lee, and R. Sorkin, Phys. Rev. D **34**, 373 (1986); V. Frolov and I. Novikov, *ibid.* **48**, 4545 (1993).
- [10] C. Callan and F. Wilczek, Phys. Lett. B 333, 55 (1994).
- [11] A. I. Barvinsky, V. P. Frolov, and A. I. Zelnikov, Phys. Rev. D 51, 1741 (1995).

- [12] S. P. de Alwis and N. Ohta, "On the entropy of quantum fields in black hole backgrounds," Report No. COLO-HEP-347, hep-th/9412027 (unpublished).
- [13] G. Cognola, L. Vanzo, and S. Zerbini, Class. Quantum Grav. 12, 1927 (1995).
- [14] A. Ghosh and P. Mitra, Phys. Rev. Lett. 73, 2521 (1994).
- [15] K. Shiraishi, Mod. Phys. Lett. A 9, 3509 (1994).
- [16] L. Susskind and J. Uglum, Phys. Rev. D 50, 2700 (1994); L. Susskind, "Some speculations about black hole entropy in string theory," Report No. hep-th/9309145 (unpublished).
- [17] T. Jacobson, "Black hole entropy and induced gravity," Report No. gr-qc/9404039 (unpublished).
- [18] S. N. Solodukhin, Phys. Rev. D 51, 609 (1995).
- [19] D. V. Fursaev, Mod. Phys. Lett. A 10, 649 (1995).
- [20] S. N. Solodukhin, Phys. Rev. D 51, 618 (1995).
- [21] D. V. Fursaev and S. N. Solodukhin, Phys. Lett. B 365, 51 (1996).
- [22] G. W. Gibbons and S. W. Hawking, Phys. Rev. D 15, 2752 (1976); S. W. Hawking, in *General Relativity*, edited by S. W. Hawking and W. Israel (Cambridge University Press, Cambridge, England, 1979).
- [23] M. -H. Lee and J. K. Kim, "The entropy of a quantum field in a charged Kerr black hole," Report No. KAIST-CHEP-95/8 (unpublished); this issue, Phys. Rev. D 54, 3904 (1996).

- [24] J. D. Brown, E. A. Martinez, and J. W. York, Jr., Phys. Rev. Lett. 66, 2281 (1991).
- [25] D. V. Fursaev and S. N. Solodukhin, Phys. Rev. D 52, 2133 (1995).
- [26] D. D. Sokolov and A. A. Starobinsky, Sov. Phys. Dokl. 22, 312 (1977).
- [27] R. M. Wald, Phys. Rev. D 48, R3427 (1993); V. Iyer and R. M. Wald, *ibid.* 50, 846 (1994); T. A. Jacobson, G. Kang, and R. C. Myers, *ibid.* 49, 6587 (1994); V. Iyer and R. M. Wald, *ibid.* 52, 4430 (1995).
- [28] N. D. Birrell and P. C. W. Davies, *Quantum Fields in Curved Space* (Cambridge University Press, Cambridge, England, 1982).
- [29] H. P. McKean and I. M. Singer, J. Diff. Geometry 1, 43 (1967); J. Cheeger, *ibid.* 18, 575 (1983); H. Donnelly, Math. Ann. 224, 161 (1976); P. Chang and J. S. Dowker, Nucl. Phys. B395, 407 (1993); D. V. Fursaev, Class. Quantum Grav. 11, 1431 (1994); G. Cognola, K. Kirsten, and L. Vanzo, Phys. Rev. D 49, 1029 (1994); D. V. Fursaev and G. Miele, *ibid.* 49, 987 (1994).
- [30] D. V. Fursaev, Phys. Lett. B 334, 53 (1994).
- [31] J. S. Dowker, Phys. Rev. D 50, 6369 (1994).
- [32] S. N. Solodukhin, this issue, Phys. Rev. D 54, 3900 (1996).
- [33] L. P. Eisenhart, *Riemannian Geometry* (Princeton University Press, Princeton, NJ, 1966).