

# String dynamics in cosmological and black hole backgrounds: The null string expansion

C. O. Lousto

*Department of Physics, University of Utah, Salt Lake City, Utah 84112*

N. Sánchez

*Observatoire de Paris, DEMIRM, UA 336 Laboratoire associé au Centre National de la Recherche Scientifique (CNRS),  
61, Avenue de l'Observatoire, F-75014 Paris, France*

(Received 8 May 1996)

We study the classical dynamics of a bosonic string in the  $D$ -dimensional flat Friedmann-Robertson-Walker and Schwarzschild backgrounds. We make a perturbative development in the string coordinates around a *null* string configuration; the background geometry is taken into account exactly. In the cosmological case we uncouple and solve the first order fluctuations; the string time evolution with the conformal gauge world-sheet  $\tau$  coordinate is given by  $X^0(\sigma, \tau) = q(\sigma)\tau^{1/(1+2\beta)} + c^2 B^0(\sigma, \tau) + \dots$ ,  $B^0(\sigma, \tau) = \sum_k b_k(\sigma)\tau^k$  where  $q(\sigma)$  is a function of the momentum component  $P_0(\sigma)$ ,  $b_k(\sigma)$  are obtained from the equation for the first order fluctuations, and  $\beta$  is the exponent of the conformal factor in the Friedmann-Robertson-Walker metric, i.e.,  $R \sim \eta^\beta$ . The string proper size, at first order in the fluctuations, grows such as the conformal factor  $R(\eta)$  and the string energy-momentum tensor corresponds to that of a null fluid. For a string in the black hole background, we study the planar case, but keep the dimensionality of the spacetime  $D$  generic. In the null string expansion, the radial, azimuthal, and time coordinates  $(r, \phi, t)$  are  $r = \sum_n A_n^1(\sigma)(-\tau)^{2n/(D+1)}$ ,  $\phi = \sum_n A_n^3(\sigma)(-\tau)^{(D-5+2n)/(D+1)}$ , and  $t = \sum_n A_n^0(\sigma)(-\tau)^{1+2n(D-3)/(D+1)}$ . The first terms of the series represent a *generic* approach to the Schwarzschild singularity at  $r=0$ . First and higher order string perturbations contribute with higher powers of  $\tau$ . The integrated string energy-momentum tensor corresponds to that of a null fluid in  $D-1$  dimensions. As the string approaches the  $r=0$  singularity its proper size grows indefinitely such as  $\sim (-\tau)^{-(D-3)/(D+1)}$ . We end the paper giving three particular exact string solutions inside the black hole. They represent, respectively, straight strings across the origin, twisted, and rigidly rotating strings. [S0556-2821(96)00622-6]

PACS number(s): 11.25.Db, 04.70.Bw, 98.80.Hw

## I. INTRODUCTION

The investigation of strings in curved spacetime is currently the best framework to study the physics of gravitation in the context of string theory. The study of string propagation in curved spacetimes provides new insight with respect to string theory in flat, Minkowski spacetime (and with respect to quantum fields in curved spacetime) (see, for example, Refs. [1–4]). The results of this study apply to fundamental strings, as well as to cosmic strings, which behave essentially in a classical way.

The string equations of motion and constraints in curved spacetime are highly nonlinear and, in general, nonexactly solvable. Different methods are available to solve this system: The string perturbation approach [5], the  $\tau$ -expansion method [6] (which provides exact local solutions for any background), the null string approach [7], and the construction of global solutions (by solitonic and inverse scattering methods, for instance), which allowed to uncover the new feature of multistring solutions [8–11]. The expansion methods are described and classified by using the world-sheet velocity of light  $c$ , as an expansion parameter, in Sec. II below.

An approximate but general method is the expansion around the center-of-mass solution of the string (satisfying the geodesic equation of motion), the world-sheet time  $\tau$  being identified with the proper time associated to the center-of-mass trajectory. In this paper we are concerned with an-

other general approximation method: The null string approach. In this approach, the string equations of motion and constraints are systematically expanded in powers of the parameter  $c$  (the speed of light in the world sheet). This corresponds to a small string tension expansion. To zeroth order the string is effectively equivalent to a continuous line beam of massless particles labeled by the world-sheet spatial parameter  $\sigma$ . The points on the “null” string do not interact among themselves, but they interact with the gravitational background. Note the extended character of the zeroth order solution in the null string approach, as opposed to the perturbation approach in which the zeroth order is given by the string’s center-of-mass pointlike approximation.

We also study fluctuations around the null string configuration, which naturally appears as an expansion in powers of  $\tau$  with precisely defined  $\sigma$ -dependent coefficients. The string coordinates are expressed as

$$X^A(\sigma, \tau) = A^A(\sigma, \tau) + c^2 B^A(\sigma, \tau) + c^4 C^A(\sigma, \tau) + \dots, \quad (1.1)$$

the zeroth order  $A^A(\sigma, \tau)$ , first and second order fluctuations  $B^A(\sigma, \tau)$ ,  $C^A(\sigma, \tau)$  satisfy Eqs. (2.3) and (2.4) as described in Sec. II. This null string approach is a powerful description for strings in the strong gravitational field regime, as is the case of black hole backgrounds near the  $r=0$  singularity and in inflationary cosmological spacetimes, at late times, where string instabilities develop. The string perturbation method

around the string center of mass allows to detect the emergence of instabilities [12,13,11], but is unable to describe the highly unstable regime [6]. We apply this null string expansion to cosmological spacetimes and to strings in the interior of black holes, falling into the singularity at  $r=0$ . In Friedmann-Robertson-Walker cosmological spacetimes with conformal expansion factor  $R \propto \eta^\beta$ ,  $\eta$  being the conformal gauge world-sheet time, the null string evolution can be expressed as

$$X^0(\sigma, \tau) = q(\sigma) \tau^{1/(1+2\beta)} + c^2 B^0(\sigma, \tau) + \dots, \quad (1.2)$$

where

$$B^0(\sigma, \tau) = \sum_k b_k(\sigma) \tau^k, \quad k = 2, 2 \pm \alpha/\beta, k_\pm. \quad (1.3)$$

The spatial coordinates  $X^i(\sigma, \tau)$  are given by Eqs. (3.5) and (3.22) in Sec. III.

The proper size of the string grows proportional to the expansion factor  $R(\eta)$  and the string energy-momentum tensor corresponds to that of a null fluid.

In black hole spacetimes we find the null string expansion inside the Schwarzschild black hole. It is given by

$$t \doteq X^0(\sigma, \tau) = \sum_{n=1}^{\infty} A_n^0(\sigma) (-\tau)^{1+2n(D-3)/(D+1)} + c^2 B^0(\sigma, \tau) + \dots, \quad (1.4)$$

$$r \doteq X^1(\sigma, \tau) = \sum_{n=1}^{\infty} A_n^1(\sigma) (-\tau)^{2n/(D+1)} + c^2 B^1(\sigma, \tau) + \dots, \quad (1.5)$$

$$\phi \doteq X^3(\sigma, \tau) = \sum_{n=1}^{\infty} A_n^3(\sigma) (-\tau)^{(D-5+2n)/(D+1)} + c^2 B^3(\sigma, \tau) + \dots, \quad (1.6)$$

and we have taken, for simplicity, the string in the equatorial plane, i.e.,

$$\theta^i \doteq A^i = \pi/2 \quad \text{for } i = 2, 4, 5, \dots, D-1. \quad (1.7)$$

The  $\tau$  dependence exhibited above represents a generic behavior near the  $r \rightarrow 0$  singularity ( $\tau \rightarrow 0$ ), and generalizes to  $D$  dimensions, the case recently analyzed in Ref. [14]. Higher order fluctuations  $c^2 B^\mu$ ,  $c^4 C^\mu$ , and so on contribute with increasing orders of  $\tau$ .

The proper string size grows indefinitely as the string approaches the  $r \rightarrow 0$  singularity as

$$\left( \frac{dl}{d\sigma} \right) \rightarrow \left( \frac{R_S}{r} \right)^{(D-3)/2} X^{0'}(\sigma, \tau=0) \rightarrow (-\tau)^{-(D-3)/(D+1)}. \quad (1.8)$$

The integrated string energy-momentum tensor for  $r \rightarrow 0$  behaves as

$$\Theta_{r \rightarrow 0}^t \rightarrow -\frac{L}{2\pi\alpha' r} = -\Theta_\phi^\phi, \quad \Theta_{t \rightarrow 0}^t \rightarrow 0, \quad (1.9)$$

$L$  being the string orbit angular momentum. In the infalling towards the  $r \rightarrow 0$  singularity, the string behaves as a null fluid in  $D-1$  dimensions; in particular, a planar string behaves as a two (spatially) -dimensional null fluid. The emergence of string instabilities in Schwarzschild and Reissner-Nordström spacetimes, and the growing of the string proper size near the black hole singularity was first found in Ref. [13] using the perturbative expansion method around the string center of mass (although such expansion does not allow one to describe the highly unstable regime and the full approach to the  $r \rightarrow 0$  singularity).

This paper is organized as follows: In Sec. II we classify the different expansion methods to solve the string dynamics in curved spacetimes in terms of the parameter  $c$  (or equivalently, in terms of the ratio between  $\dot{X}^A$  and  $cX^{A'}$ , the  $\tau$  and  $\sigma$  derivatives of the string coordinates), with particular focusing on the null string expansion (see also Table II). In Sec. III we solve the null string expansion in cosmological spacetimes. In Sec. IV we find the null string expansion near the Schwarzschild  $r \rightarrow 0$  singularity. In Sec. V we find particular exact solutions inside the black hole event horizon describing straight string across  $r=0$ , twisted, and rigidly rotating strings. Section VI summarizes our conclusions.

## II. STRING DYNAMICS IN GRAVITATIONAL BACKGROUNDS

The action of a bosonic string in a  $D$ -dimensional curved manifold endowed with a metric  $G_{AB}$  ( $0 \leq A, B \leq D-1$ ) is given by [15,16]

$$S = -T_0 \int d\sigma d\tau \sqrt{-\det g_{\mu\nu}}, \quad (2.1)$$

where  $g_{\mu\nu} = G_{AB}(X) \partial_\mu X^A \partial_\nu X^B$  is the two-dimensional world-sheet metric ( $0 \leq \mu, \nu \leq 1$ ) and  $T_0 = 1/(2\pi\alpha')$  is the string tension.

By use of the reparametrization invariance of the world sheet one can take the conformal gauge, i.e.,  $g_{\mu\nu} = \rho(\sigma, \tau) \eta_{\mu\nu}$ , where  $\eta_{\mu\nu}$  is the two-dimensional Minkowskian metric. In this gauge, the classical equations of motion derived from the action (2.1) read

$$\partial_\tau^2 X^A - c^2 \partial_\sigma X^A + \Gamma_{BC}^A [\partial_\tau X^B \partial_\tau X^C - c^2 \partial_\sigma X^B \partial_\sigma X^C] = 0, \quad (2.2)$$

where  $\Gamma_{BC}^A$  are the Christoffel symbols associated to the metric  $G_{AB}$  and we have introduced the velocity of wave propagation along the string (velocity of light),  $c$ , as the second fundamental constant after the string tension.

Variation of action (2.1) with respect to the world-sheet variables yields the two nontrivial constraints

$$\partial_\tau X^A \partial_\sigma X^B G_{AB} = 0, \quad (2.3)$$

$$[\partial_\tau X^A \partial_\tau X^B + c^2 \partial_\sigma X^A \partial_\sigma X^B] G_{AB} = 0. \quad (2.4)$$

de Vega and Sánchez [5] have proposed a method to solve, both classically and quantum mechanically, the equations of motion and constraints of strings in curved spacetimes, i.e., Eqs. (2.2)–(2.4). This method treats the spacetime geometry exactly and considers the string excitations small

as compared to the energy scales associated to the background gravitational metric. This method is particularly well suited to study strings in strong gravitational fields such as in black holes and in cosmological scenarios as opposed to the usual treatment in flat spacetime. The starting point is to consider a particular solution to the equations of motion and constraints,  $X_{\text{part}}^A(\sigma, \tau) = A^A(\sigma, \tau)$  and then successively studying first, second, and higher order fluctuations around this particular solution; to be denoted, respectively, as  $B^A, C^A$ , etc.

In Ref. [7] have been raised the interesting possibility of using  $c$ , the world-sheet light velocity, as the expansion parameter in the above development. This approach allows us to classify the different solutions proposed in the literature: (i)  $c \ll 1$  or equivalently, as can be seen from the field and constraint equations (2.2)–(2.4),  $\partial_\tau A^A \gg c \partial_\sigma A^A$  (holding also this inequality for second derivatives); (ii)  $c = 1$  and thus all derivatives and other terms are, in principle, relevant in Eqs. (2.2)–(2.4); (iii)  $c \gg 1$  or equivalently,  $c \partial_\sigma A^A \gg \partial_\tau A^A$  (holding also for second derivatives); (iv) There can be, of course, several other possibilities such as hybrid starting solutions that for some components of  $A^A$  (and eventually  $B^A, C^A$ , etc.) satisfy one of the cases (i), (ii), or (iii) and for other components a different case.

More explicitly, we have

Case (i)  $c \ll 1$ : The appropriate development reads

$$X^A(\sigma, \tau) = A^A(\sigma, \tau) + c^2 B^A(\sigma, \tau) + c^4 C^A(\sigma, \tau) + \dots \quad (2.5)$$

Plugging this into Eqs. (2.2)–(2.4) one obtains [7], to zeroth order in  $c$  (since now on we adopt the notation  $\dot{\phantom{x}} = \partial_\tau$  and  $\prime = \partial_\sigma$ ),

$$\ddot{A}^A + \Gamma_{BC}^A \dot{A}^B \dot{A}^C = 0, \quad (2.6)$$

$$\dot{A}^B \dot{A}^C G_{BC} = 0, \quad (2.7)$$

$$\dot{A}^B A'^C G_{BC} = 0. \quad (2.8)$$

Equation (2.6) is the geodesic path followed by each point of the string. Equation (2.7) means that this is a null geodesic and Eq. (2.8) states that the velocity is perpendicular to the string.

First order fluctuations around this particular solution  $A(\sigma, \tau)$  can be obtained by retaining terms of order  $c^2$  in Eqs. (2.2)–(2.4):

$$\ddot{B}^A + 2\Gamma_{BC}^A \dot{A}^B \dot{B}^C + \Gamma_{BC,D}^A \dot{A}^B \dot{A}^C B^D = A''^A + \Gamma_{BC}^A A'^B A'^C, \quad (2.9)$$

$$(2\dot{A}^B \dot{B}^C + A'^B A'^C) G_{BC} + G_{BC,D} \dot{A}^B \dot{A}^C B^D = 0, \quad (2.10)$$

$$(\dot{B}^B A'^C + \dot{A}^B B'^C) G_{BC} + G_{BC,D} \dot{A}^B A'^C B^D = 0. \quad (2.11)$$

Higher order corrections can thus be systematically obtained. The interpretation of these equations is that they give the high energy (compared to the Planck energy) string behavior, corresponding to the limit of vanishing tension (and thus small  $c$ ).

Case (ii)  $c = 1$ , and thus all derivatives and other terms are, in principle, relevant in Eqs. (2.2)–(2.4). One can thus make developments around the center-of-mass motion as a physically appealing starting solution. In fact, de Vega and Sánchez [5] have proposed in their original approach

$$X^A = A^A(\sigma, \tau) + B^A(\sigma, \tau) + C^A(\sigma, \tau) + \dots, \quad (2.12)$$

where  $A^A(\tau)$  follows the geodesic equation

$$\ddot{A}^A + \Gamma_{BC}^A \dot{A}^B \dot{A}^C = 0, \quad (2.13)$$

and the constraint

$$\dot{A}^B \dot{A}^C G_{BC} = 0 \quad (2.14)$$

means that these geodesics are null.

First order fluctuation equations now read

$$\ddot{B}^A - B''^A + 2\Gamma_{BC}^A \dot{A}^B \dot{B}^C + \Gamma_{BC,D}^A \dot{A}^B \dot{A}^C B^D = 0, \quad (2.15)$$

and

$$2G_{BC} \dot{A}^B \dot{B}^C + G_{BC,D} \dot{A}^B \dot{A}^C B^D = 0, \quad (2.16)$$

$$G_{BC} \dot{A}^B B'^C = 0. \quad (2.17)$$

Higher order corrections can be in this way systematically considered. Still, a Fourier series decomposition of the first order fluctuations can be made [5], i.e.,  $B^A(\sigma, \tau) = \sum_n \exp(in\sigma) \eta_n^A(\tau)$ . Also, a decomposition in purely left  $A^A(\sigma + \tau)$  or right movers  $A^A(\sigma - \tau)$  can be used as a starting exact solution.

Case (iii)  $c \gg 1$  or equivalently,  $c \partial_\sigma A^A \gg \partial_\tau A^A$  (holding also for second derivatives). The appropriate development reads in this case

$$X^A(\sigma, \tau) = A^A(\sigma, \tau) + \frac{1}{c^2} B^A(\sigma, \tau) + \frac{1}{c^4} C^A(\sigma, \tau) + \dots \quad (2.18)$$

Plugging this expression into Eqs. (2.2)–(2.4) yields a set of equations that is equivalent to Eqs. (2.6)–(2.11) upon the substitutions  $\tau \leftrightarrow \sigma$  and  $c \rightarrow c^{-1}$ . The interpretation of this approximation is now exactly the opposite to case (i), i.e., here the zeroth order dependence on  $\sigma$  overwhelms that on  $\tau$  and thus represents very low energy strings, frozen at first approximation. Solutions to this case can be obtained from solutions to case (i) by making the above mentioned substitutions. These represent, at zeroth order, stationary solutions as opposed to the dynamical ones. While solutions to case (i) are appropriate to describe strings in strong gravitational fields, for instance, near black hole singularities, i.e.,  $R \rightarrow 0$ , solutions to the case (iii) are rather appropriate for strings far away from black holes, i.e.,  $R \rightarrow \infty$ . We thus have an approximate duality here, represented by the transformations  $\tau \leftrightarrow \sigma$  and  $R \leftrightarrow R^{-1}$ , which map the solutions to the cases (i) and (iii) into each other.

Case (iv): This corresponds to a hybrid case. The circular string ansatz is an example of this situation:

$$X^0 = X^0(\tau), \quad R = R(\tau), \quad \psi = \psi(\sigma), \quad \theta = \pi/2. \quad (2.19)$$

Clearly,  $X^0$  and  $R$  coordinates follow case (i) while  $\psi$  follows case (iii) and  $\theta$  could be assigned to case (ii). The first order fluctuations around this starting solution have been studied in Refs. [17,11] with regards to the stability analysis as the string approaches the black hole singularity. The stationary string ansatz has been studied in Ref. [18].

Finally, one can consider exact or asymptotic particular string solutions that deserve study in their own. Asymptotic solutions in a Friedmann-Robertson-Walker background representing highly unstable strings have been found [6]. Exact string and multistring solutions in de Sitter spacetime have been found by solitonic methods (a single world sheet describing multiple different and independent strings) [19,8,9,11]. Ring solutions in black holes [10] have also been studied. For the sake of completeness it is also worth remarking that the propagation of strings in shock wave and conical spacetimes has been exactly solved [20,21].

Two physical interesting quantities can be computed: The string energy-momentum tensor integrated over a spatial volume completely enclosing the string [22], at fixed time  $X^0$ ,

$$I^{AB}(X^0) = \int \sqrt{-G} T^{AB}(X) d^{D-1} \vec{X}, \quad (2.20)$$

where

$$\begin{aligned} \sqrt{-G} T^{AB}(X) &= \frac{1}{2\pi\alpha'} \int d\sigma d\tau [\dot{X}^A \dot{X}^B - X'^A X'^B] \\ &\times \delta^{(D)}[X - X(\sigma, \tau)], \end{aligned} \quad (2.21)$$

and the invariant string size  $l$  [3],

$$dl^2 = -\dot{X}^A \dot{X}^B G_{AB}(X) d\sigma^2. \quad (2.22)$$

[Note that the differential string size has the form of an effective mass  $m_{\text{eff}}(\tau, \sigma)$  for the geodesic motion, and actually represents a projection from the target space onto the world sheet.]

### III. NULL STRING DYNAMICS IN COSMOLOGICAL BACKGROUNDS

Let us consider, as a simple application of the formalism above, the isotropic cosmological geometry given by the conformally flat Friedmann-Robertson-Walker metric

$$\begin{aligned} ds^2 &= R^2(X^0) [-(dX^0)^2 + (dX^1)^2 + (dX^2)^2 + \dots \\ &+ (dX^{D-1})^2], \end{aligned} \quad (3.1)$$

where we parametrize the conformal factor as  $R(X^0) = K(X^0/\beta)^\beta$ .

We are interested in this paper in the null string dynamics and the fluctuations around it. *Zeroth order*: The equations of motion (2.6) have the following first integral of motion (conservation of the four-moment  $P_A$ ):

$$R^2(A^0) \dot{A}^0(\sigma, \tau) = -P_0(\sigma), \quad R^2(A^0) \dot{A}^i(\sigma, \tau) = P_i(\sigma), \quad (3.2)$$

where from now on  $i, j$  run from 1 to  $D-1$ .

The constraints at  $\tau=0$  yield

TABLE I. Here, we recall the explicit conformal time ( $\eta$ ) and cosmic time ( $t$ ) dependence of the flat Friedmann-Robertson-Walker metric as usually referred to in the literature [6].

$R(\eta) = (\eta/\beta)^\beta$	$R(t) \sim t^{\alpha = \beta/(\beta+1)}$	Spacetime
Constant, $\beta \rightarrow 0$	Constant	Flat space
$\eta^\beta$ , $0 < \beta < \infty$	$t^\alpha$ , $0 < \alpha < 1$	Standard cosmology
$e^\eta$ , $\beta \rightarrow \infty$	$t$	Linear expansion
$\eta^\beta$ , $-1 < \beta < 0$	$t^\alpha$ , $-\infty < \alpha < 0$	Super inflation
$\eta^{-1}$	$e^{Ht}$	de Sitter
$\eta^\beta$ , $-\infty < \beta < -1$	$t^\alpha$ , $1 < \alpha < \infty$	Power-law inflation

$$(P_0)^2 = \sum_{i=1}^{D-1} (P_i)^2, \quad (3.3)$$

$$-P^0(\bar{A}^0)' = \sum_{i=1}^{D-1} P^i(\bar{A}^i)', \quad (3.4)$$

where we use the notation  $A^A(\sigma, \tau=0) = \bar{A}^A$ .

From Eq. (3.2) we see that the only relevant  $\tau$  dependence is that of  $A^0(\tau)$ . In fact, we can write

$$A^i(\tau) - \bar{A}^i = \frac{P^i}{-P^0} [A^0(\tau) - \bar{A}^0]. \quad (3.5)$$

By direct integration of the zeroth component of Eq. (3.2), one obtains

$$A^0(\tau) - \bar{A}^0 = \left[ \frac{(1+2\beta)\beta^{2\beta} P_0}{K^2} \tau \right]^{1/(1+2\beta)} \doteq q(\sigma) \tau^{1/(1+2\beta)}. \quad (3.6)$$

The particular time dependence of the scale factor as usually referred to in the literature is recalled in Table I. The cases dealt with in Ref. [7] are in  $D=4$  and  $\beta = -1/2$  and  $-1$ , respectively. Note that the cosmic time  $t$  is related to the conformal time  $X^0 \approx A^0$  by

$$t - t_0 = \frac{K(A^0 - \bar{A}^0)^{1+\beta}}{\beta^\beta (1+\beta)}. \quad (3.7)$$

Thus,  $\tau$  does *not* coincide with  $t$ .

We can now compute the zeroth order energy-momentum density tensor from Eq. (2.20)

$$I_A^B(\sigma) = \frac{P_A P_C \eta^{CB}}{-P_0 \alpha'}. \quad (3.8)$$

Hence, at zeroth order, the string energy and momentum keep constant as the Universe evolves. The trace of  $I_A^B$  vanishes to this zeroth order. Since, to this order, each point of the string follows a null geodesic, the proper size equation (2.22) also vanishes.

*First order fluctuations*: The 0 component of Eq. (2.9) yields

$$\begin{aligned} \ddot{B}^0 + \frac{2dR}{RdX^0}[\dot{A}^0\dot{B}^0 + \dot{A}^i\dot{B}^i] + \frac{d}{dX^0}\left(\frac{dR}{RdX^0}\right)[(\dot{A}^0)^2 + (\dot{A}^i)^2]B^0 \\ = A^{0''} + \frac{dR}{RdX^0}[(A^{0'})^2 + (A^{i'})^2]. \end{aligned}$$

With the help of the constraint equation (2.10),

$$(2\dot{B}^A A^{B'} + \dot{A}^A B^{B'})\eta_{AB} = 0, \quad (3.9)$$

we obtain an uncoupled equation for  $B^0$ :

$$\ddot{B}^0 + \frac{4\alpha}{\tau}\dot{B}^0 - \frac{2\alpha^2}{\beta\tau^2}B^0 = A^{0''} + \frac{2(A^{i'})^2}{A^0} \doteq F^0(\sigma, \tau), \quad (3.10)$$

where  $\alpha = \beta/(\beta + 1)$ , and we have used that

$$\frac{dR}{RdX^0} = \frac{\beta}{X^0}, \quad \frac{d}{dX^0}\left(\frac{dR}{RdX^0}\right) = \frac{-\beta}{(X^0)^2}. \quad (3.11)$$

We can write the general solution to Eq. (3.10) as

$$B^0(\sigma, \tau) = \sum_k b_k(\sigma)\tau^k, \quad (3.12)$$

where  $b_{k_+}$  and  $b_{k_-}$  are arbitrary constants corresponding to the general solution to the homogeneous equation associated to Eq. (3.10);  $k_{\pm} = -(4\alpha - 1)/2 \pm \sqrt{(2\alpha - 1/2)^2 + 2\alpha^2/\beta}$ . The other powers of  $k$  correspond to the series development of the function  $F^0(\sigma, \tau)$  and give the particular solution to the inhomogeneous equation.

To provide a simpler solvable example we take  $\bar{A}^0 = 0$ . In this case  $F^0$  takes the form

$$\begin{aligned} F^0(\sigma, \tau) = \left[ q'' + \frac{2\beta}{q}\left(\frac{P_i q}{P_0}\right)' \right]^2 \tau^{\alpha/\beta} - \frac{4\beta\bar{A}^{i'}}{q}\left(\frac{P_i q}{P_0}\right)' \\ + \frac{\beta(\bar{A}^{i'})^2}{q}\tau^{-\alpha/\beta}. \end{aligned} \quad (3.13)$$

Plugging this into the right-hand side of Eq. (3.10) and using the form of  $B^0$  given by Eq. (3.12), we obtain

$$\sum_k b_k[k(k-1) + 4k\alpha - 2\alpha^2/\beta]\tau^{k-2} = F^0(\sigma, \tau). \quad (3.14)$$

Thus, there are only three values of  $k$  given by  $k - 2 = 0, \pm \alpha/\beta$  with the corresponding  $b_k$  as defined by Eqs. (3.13) and (3.14):

$$\begin{aligned} b_{-2} &= -\frac{4\beta\bar{A}^{i'}}{q[2 + 8\alpha - 2\alpha^2/\beta]}, \\ b_{2+\alpha/\beta} &= \frac{q'' + \frac{2\beta}{q}\left(\frac{P_i q}{P_0}\right)' ^2}{(2 + \alpha/\beta)(1 + 4\alpha + \alpha/\beta) - 2\alpha^2/\beta}, \\ b_{2-\alpha/\beta} &= \frac{\beta(\bar{A}^{i'})^2}{q[(2 - \alpha/\beta)(1 + 4\alpha - \alpha/\beta) - 2\alpha^2/\beta]}. \end{aligned} \quad (3.15)$$

Let us now consider the  $i$  components of Eq. (2.9) for the first order fluctuations:

$$\begin{aligned} \ddot{B}^i + \frac{2}{R}\frac{dR}{dX^0}[\dot{A}^0\dot{B}^i + \dot{A}^i\dot{B}^0] + 2\frac{d}{dX^0}\left(\frac{dR}{RdX^0}\right)\dot{A}^0\dot{A}^i B^0 = A^{i''} \\ + 2\frac{dR}{RdX^0}A^{0'}A^{i'}. \end{aligned} \quad (3.16)$$

With the help of Eq. (3.11),

$$\begin{aligned} \ddot{B}^i + \frac{2\alpha}{\tau}\dot{B}^i = A^{i''} + \frac{2\beta A^{i'}A^{0'}}{A^0} + \frac{2\alpha P_i}{P_0\tau}\left(\dot{B}^0 - \frac{\alpha B^0}{\beta\tau}\right) \\ \doteq F^i(\sigma, \tau). \end{aligned} \quad (3.17)$$

Knowing  $B^0(\sigma, \tau)$  the general solution to this equation can be written in terms of quadratures:

$$B^i(\sigma, \tau) = \bar{B}^i + \bar{B}^i \frac{\tau^{1-2\alpha}}{1-2\alpha} + \int^\tau d\tilde{\tau} \tilde{\tau}^{-2\alpha} \int^{\tilde{\tau}} d\tilde{\tau} \tilde{\tau}^{2\alpha} F^i(\sigma, \tilde{\tau}), \quad (3.18)$$

where  $\bar{B}^i = B^i(\sigma, 0)$ . Note that  $\bar{B}^i$  and  $\bar{B}^i$  can be absorbed into  $\bar{A}^i$  and  $P_i$ , respectively. We are thus left with the two constants  $b_{k_{\pm}}$  which are fixed by the two first order constraint equations (2.10) and (2.11):

$$\begin{aligned} (\dot{A}^A \dot{B}^B + A^{A'} A^{B'})\eta_{AB} = 0, \\ (2\dot{B}^A A^{B'} + \dot{A}^A B^{B'})\eta_{AB} = 0. \end{aligned} \quad (3.19)$$

We are thus sure we are not adding any spurious degree of freedom at first order.

We can again show the explicit dependence for the case  $\bar{A}^0 = 0$ . In this case  $F^i$  takes the form

$$\begin{aligned} F^i(\sigma, \tau) = \bar{A}^{i''} + 2b_{2\alpha} \frac{P_i}{P_0}(2 - \alpha/\beta) + 2\beta \frac{q'}{q} \bar{A}^i \\ - \left[ \left(\frac{P_i q}{P_0}\right)'' + 2\beta \frac{q'}{q} \left(\frac{P_i q}{P_0}\right)' - 4\alpha b_{2+\alpha/\beta} \frac{P_i}{P_0} \right] \tau^{\alpha/\beta} \\ + 4\alpha b_{2-\alpha/\beta} \frac{P_i}{P_0} (1 - \alpha/\beta) \tau^{-\alpha/\beta} \\ + 2\alpha b_{k_{\pm}} \frac{P_i}{P_0} [(k_{\pm}) - \alpha/\beta] \tau^{k_{\pm}-2}, \end{aligned} \quad (3.20)$$

or in a more compact way

$$F^i(\sigma, \tau) = \sum_n F_n^i(\sigma) \tau^n, \quad n = 0, \pm \alpha/\beta, (k_{\pm}) - 2, \quad (3.21)$$

where the notation  $F_n^i$  clearly refers to the coefficient of the  $n$ th power of  $\tau$  in Eq. (3.20).

Plugging this into the right-hand side of Eq. (3.18) and using the form of  $B^0$  given by Eqs. (3.12) and (3.15), we obtain

$$B^i(\sigma, \tau) = \bar{B}^i + \bar{B}^i \frac{\tau^{1-2\alpha}}{1-2\alpha} + \sum_n \frac{F_n^i \tau^{n+2}}{(n+2)(2\alpha+n+1)},$$

$$n=0, \pm\alpha/\beta, (k_{\pm}) - 2. \quad (3.22)$$

The proper string size that vanished to zeroth order now takes the form

$$dl^2 = R^2 \eta_{AB} A^{A'} A^{B'} d\sigma^2, \quad (3.23)$$

where we have used the constraint (3.19). We see that the string size grows such as the conformal factor of the Friedmann-Robertson-Walker metric. This unstable behavior was already observed in Ref. [12] in the fluctuations around the center-of-mass zeroth order solution.

The coordinate fluctuations  $B^0$  and  $B^i$  evolve with several powers of  $\tau$  and for a given value of the conformal factor exponent  $\alpha$ , we will have several regimes. We can see from the above expressions [Eq. (3.22)] that in the regime of  $R$  large, that is  $\tau \rightarrow \infty$ , for  $\alpha/\beta > 0$ , i.e.,  $\beta > -1$ , the overwhelming power will be  $(2 + \alpha/\beta)$  or  $(2 + k_+)$ . When  $\alpha/\beta < 0$ , i.e.,  $\beta < -1$ , the regime of  $R$  large is reached when  $\tau \rightarrow 0$ , then the predominant power will be  $2 + \alpha/\beta$  or  $2 + k_-$ . Note also that these results do not depend explicitly on the dimension  $D$  since neither of the metric coefficients does.

#### IV. BLACK HOLE SPACETIMES

We are particularly interested in studying the dynamics of strings in black hole spacetimes and its approach to the singularity at radial coordinate  $r=0$ .

Let us consider a  $D$ -dimensional black hole background [23],

$$ds^2 = -g(r)(dX^0)^2 + g(r)^{-1} dr^2 + r^2 d\Omega_{D-2}, \quad (4.1)$$

where for the Reissner-Nordström metric

$$g(r) = 1 - \left(\frac{R_S}{r}\right)^{D-3} + \left(\frac{\tilde{Q}}{r}\right)^{2(D-3)},$$

$$\tilde{Q}^{2(D-3)} = \frac{8\pi G Q^2}{(D-2)(D-3)}, \quad (4.2)$$

with  $M$  and  $Q$  being, respectively, the mass and charge of the black hole.

We can now study the string dynamics by using the null string approach described in Sec. II.

We then propose the development for the string coordinates

$$X^A(\sigma, \tau) = A^A(\sigma, \tau) + c^2 B^A(\sigma, \tau) + \dots \quad (4.3)$$

Replacing this into the field equations (2.2), with Eq. (4.1) acting as the curved background, we find to zeroth order the first integrals of motion

$$\dot{\theta} \simeq \dot{A}^2 = 0, \quad \dot{\phi} \simeq \dot{A}^3 = \frac{L(\sigma)}{r^2}, \quad (4.4)$$

where for simplicity we consider a string lying on the equatorial plane.  $L(\sigma)$  is the angular momentum of each point forming the string. The other first integrals of motion are

$$\dot{X}^0 \simeq \dot{A}^0 = E(\sigma) g^{-1}(r), \quad \dot{r}^2 \simeq (\dot{A}^1)^2 = E^2(\sigma) - \frac{L^2(\sigma)}{r^2} g(r). \quad (4.5)$$

From this last equation one can see that for the Reissner-Nordström black hole, if  $L(\sigma) \neq 0$ , the string will only skirt the singularity. Since we are interested in the string approach to the singularity, we consider from now on Schwarzschild black holes (in  $D$  dimensions).

The remaining constraint equation (2.8) at  $\tau=0$  takes the simple form

$$\bar{A}^{0'}(\sigma) = \frac{L(\sigma)}{E(\sigma)} \bar{A}^{3'}(\sigma). \quad (4.6)$$

We can further integrate expression (4.5) near the black hole singularity to obtain

$$A^1(\sigma, \tau) \simeq p(\sigma) (-\tau)^{2(D+1)},$$

$$p(\sigma) = \left[ \frac{1}{2} (D+1) R_S^{(D-3)/2} L(\sigma) \right]^{2/(D+1)}, \quad (4.7)$$

where we have chosen  $\tau \leq 0$  such that  $A^1(\tau=0) = 0$ .  $\tau$  measures the proper time of infall to the singularity at  $r=0$ .

By use of the above expression, we find

$$A^0(\sigma, \tau) \simeq \bar{A}^0 + \frac{(D+1)}{(3D-5)} E \left( \frac{p(\sigma)}{R_S} \right)^{D-3} (-\tau)^{(3D-5)/(D+1)}, \quad (4.8)$$

and

$$A^3(\sigma, \tau) \simeq \bar{A}^3 - \frac{(D+1)}{(D-3)} \frac{L(\sigma)}{p^2(\sigma)} (-\tau)^{(D-3)/(D+1)}, \quad (4.9)$$

$$A^i = \pi/2 \quad \text{for } i=2,4,5, \dots, D-1. \quad (4.10)$$

Comparison of these expressions with the corresponding ones found in Ref [13] [Eq. (88)], shows that they have the same  $\tau$  dependence. The difference lies in the prefactors: In Eqs. (4.7) and (4.8) above they are  $\sigma$  dependent while in those of Ref. [13] they are constant. This is due to the different zeroth order solution we have chosen. In Ref. [13] we have taken the center-of-mass motion which is a pointlike solution, while now we have a null string, an extended zeroth order solution.

We can consider the approach to the singularity with higher powers of  $A^1$  (our radial coordinate). This yields the following power dependence in  $\tau$ :

$$A^1 = \sum_{n=1}^{\infty} A_n^1(\sigma) (-\tau)^{2n/(D+1)}, \quad (4.11)$$

and then

$$A^3 = \sum_{n=1}^{\infty} A_n^3(\sigma)(-\tau)^{(D-5+2n)/(D+1)}, \quad (4.12)$$

$$A^0 = \sum_{n=1}^{\infty} A_n^0(\sigma)(-\tau)^{1+2n(D-3)/(D+1)}. \quad (4.13)$$

This  $\tau$  dependence seems to be a quite general behavior near singularities and generalizes to  $D$  dimensions the case analyzed recently in Ref. [14]. There will still be higher order corrections coming from the first order fluctuations  $B^\mu$ , second order fluctuations  $C^\mu$ , and so on. We have already seen that in the cosmological case, five different powers of  $\tau$  contribute to the first order.

The proper string size that at zeroth order vanished for null string, does not vanish at higher order. In order to compute the first order correction to the string size we do not need to explicitly compute the  $B^A$  terms. In fact, by use of the constraint (2.4), Eq. (2.22) can be rewritten as

$$\left(\frac{dl}{d\sigma}\right)^2 = X^{A'} X^{B'} G_{AB}(X). \quad (4.14)$$

Plugging expressions (4.5) and (4.4) into this equation and making use of the constraint (2.8) to eliminate  $A^{1'}$ , we obtain

$$\left(\frac{dl}{d\sigma}\right)^2 = -g(A^{0'})^2 + (A^1)^2(A^{3'})^2 + g \frac{(EA^{0'} - LA^{3'})^2}{(E^2 - L^2g/(A^1)^2)}. \quad (4.15)$$

As the string approaches the singularity it diverges in the following way:

$$\left(\frac{dl}{d\sigma}\right)^2 \rightarrow \left(\frac{R_S}{r}\right)^{D-3} (\bar{X}^{0'})^2 \rightarrow (-\tau)^{-2(D-3)/(D+1)}. \quad (4.16)$$

This unstable behavior has already been discovered in Ref. [13], where we studied string propagation in  $D$ -dimensional Reissner-Nordström black holes under the approximation labeled by  $c=1$ . We solved there the first order string fluctuations around the center-of-mass motion at spatial infinity, near the event horizon and at the spacetime singularity. In Reissner-Nordström black holes the radial components and angular string components develop instabilities as the string approaches the singularity. The string motion is such as that of a particle in a strongly attractive potential proportional to  $-(\Delta\tau)^{-2}$ . As  $\Delta\tau \rightarrow 0$  the string ends trapped by the black hole singularity.

We can also compute the integrated energy-momentum tensor from expressions (2.20) and (2.21). Using Eqs. (4.5) and (4.4), we find

$$\Theta^{AB} = \frac{dI^{AB}}{d\sigma} \approx \frac{1}{2\pi\alpha'} \frac{\sqrt{-g} \dot{A}^A \dot{A}^B}{\sqrt{E^2 - L^2g/(A^1)^2}}, \quad (4.17)$$

where we have taken into account that inside the black hole  $A^0$  is spacelike and  $A^1$  is timelike.

In the limit  $r \rightarrow 0$ , we have

$$\Theta_{1 \rightarrow}^1 - \frac{L}{2\pi\alpha' r} = -\Theta_3^3, \quad \Theta_{0 \rightarrow}^0 = 0. \quad (4.18)$$

We first observe that the results are independent of the space-time dimensionality  $D$ . The facts that the trace vanishes (since we are dealing with a null system), and that  $\Theta_{0 \rightarrow}^0 = 0$ , reduce the system to a  $(D-1)$ -dimensional null system (in our example of a planar string, this reduces the system to a two-dimensional null string).

## V. EXACT STRING SOLUTIONS INSIDE A BLACK HOLE

In four dimensions the equations of motion of a string in the curved background (4.1) read [18]

$$\begin{aligned} \ddot{t} - t'' + \frac{g_{,r}}{g}(\dot{t}r - t'r') &= 0, \\ \ddot{r} - r'' - \frac{g_{,r}}{2g}(\dot{r}^2 - r'^2) + \frac{g g_{,r}}{2}(\dot{t}^2 - t'^2) - gr(\dot{\theta}^2 - \theta'^2) \\ &\quad - gr \sin^2 \theta (\dot{\phi}^2 - \phi'^2) = 0, \\ \ddot{\theta} - \theta'' + \frac{2}{r}(\dot{\theta}r - \theta'r') - \sin\theta \cos\theta (\dot{\phi}^2 - \phi'^2) &= 0, \\ \ddot{\phi} - \phi'' + \frac{2}{r}(\dot{\phi}r - \phi'r') + 2 \cot\theta (\dot{\theta}\dot{\phi} - \theta'\phi') &= 0, \end{aligned} \quad (5.1)$$

and the constraints

$$\begin{aligned} -g\dot{t}t' + \frac{1}{g}\dot{r}r' + r^2\dot{\theta}\theta' + r^2\sin^2\theta\dot{\phi}\phi' &= 0, \\ -g(\dot{t}^2 + t'^2) + \frac{1}{g}(\dot{r}^2 + r'^2) + r^2(\dot{\theta}^2 + \theta'^2) \\ &\quad + r^2\sin^2\theta(\dot{\phi}^2 + \phi'^2) = 0. \end{aligned} \quad (5.2)$$

We are interested in finding solutions inside the black hole, where we know that the  $r$  coordinate becomes timelike while the  $t$  coordinate is spacelike. We thus look for solutions of the particular form

$$r = r(\tau) \doteq \mathcal{T}(\tau) \quad \text{and} \quad t = t(\sigma) \doteq \mathcal{R}(\sigma). \quad (5.3)$$

Using this ansatz, the first equation of motion in Eq. (5.1) gives us

$$\mathcal{R}(\sigma) = E\sigma + t_0, \quad (5.4)$$

where  $E$  and  $t_0$  are arbitrary constants.

From the first constraint equation (5.2) we have

$$\mathcal{T}(\tau)^2 \sin^2 \theta \dot{\phi} \phi' = 0. \quad (5.5)$$

In terms of Kruskal coordinates we have

$$u(\sigma, \tau) = \left(1 - \frac{\mathcal{T}(\tau)}{2M}\right)^{1/2} e^{[\mathcal{T}(\tau)]/4M} \sinh\left(\frac{\mathcal{R}(\sigma)}{4M}\right), \quad (5.6)$$

TABLE II. In this table we show the different approaches to study the string dynamics in curved spacetimes.  $c \ll 1$ ,  $c = 1$ , and  $c \gg 1$  correspond to  $\partial_\tau X^A \gg c \partial_\sigma X^A$ ,  $\partial_\tau X^A \sim c \partial_\sigma X^A$ , and  $\partial_\tau X^A \ll c \partial_\sigma X^A$ , respectively. Hybrid cases are still possible when some components of the string coordinate  $X^A$  satisfy eventually one of the inequalities while other coordinate components do not. The regimes of stability of the first order fluctuations are also shown as well as the references in which the approaches (in cosmology and black holes) have been originally developed.

Approach	Black holes		Cosmology	
	Zeroth order	Fluctuations $R \rightarrow 0$	Zeroth order	Fluctuations $R \rightarrow \infty$
(i) Impulsive, $c \ll 1$ $\dot{X}^A \gg c X^{A'}$	Null string <sup>a</sup>	Unstable	Null string <sup>b</sup>	Unstable
(ii) Perturbative, $c = 1$ $\dot{X}^A \sim c X^{A'}$	Center of mass <sup>c</sup>	Unstable	Center of mass <sup>d</sup>	Stable/unstable
(iii) Adiabatic, $c \gg 1$ $\dot{X}^A \ll c X^{A'}$	Initial configuration <sup>a</sup>	Unstable	Initial configuration <sup>a</sup>	Unstable
(iv) Hybrid	Ring <sup>e</sup>	Unstable	Asymptotic <sup>f</sup>	Stable/unstable

<sup>a</sup>This paper.

<sup>b</sup>Reference [7].

<sup>c</sup>Reference [13].

<sup>d</sup>Reference [5].

<sup>e</sup>Reference [17].

<sup>f</sup>Reference [6].

$$v(\sigma, \tau) = \left(1 - \frac{\mathcal{T}(\tau)}{2M}\right)^{1/2} e^{[\mathcal{T}(\tau)]/4M} \cosh\left(\frac{\mathcal{R}(\sigma)}{4M}\right). \quad (5.7)$$

Equation (5.5) allows us to analyze three cases:

(a)  $\theta = \text{const} \neq \pi/2$ . Then  $\phi$  is also constant and

$$\dot{\mathcal{T}}(\tau) = \pm E g(\mathcal{T}) \quad (5.8)$$

represents a straight string in the two-dimensional  $\mathcal{R}-\phi$  space.

(b) For  $\theta = \pi/2$ , i.e., the planar case, we have

(b1)  $\dot{\phi} = 0$ . Then, from the equation of motion of  $\phi$  we deduce  $\phi = n\sigma + \phi_0$ , where  $n$  and  $\phi_0$  are arbitrary constants. While for the timelike coordinate  $\mathcal{T}(\tau)$ , we have

$$\dot{\mathcal{T}}(\tau)^2 = E^2 g(\mathcal{T})^2 - n^2 \mathcal{T}(\tau)^2. \quad (5.9)$$

This solution represents a straight twisted string in the  $\mathcal{R}-\phi$  space.

(b2)  $\phi' = 0$ . Then, from the equation of motion of  $\phi$  we deduce  $\dot{\phi} = L/\mathcal{T}(\tau)^2$ . Here,  $L$  is a constant and now

$$\dot{\mathcal{T}}(\tau)^2 = E^2 g(\mathcal{T})^2 - g(\mathcal{T}) \frac{L^2}{\mathcal{T}(\tau)^2}. \quad (5.10)$$

This solution represents a rigid rotating straight string in the  $\mathcal{R}-\phi$  space.

## VI. CONCLUSION

We have performed a systematic study of the null string expansion method to solve the string equations of motion and constraints in curved spacetimes. This corresponds to a small string tension expansion (inverse powers of  $\alpha'$ ). The

perturbative expansion series is conveniently described in terms of the parameter  $c$ , the wave propagation velocity along the string. The zeroth order describes a null string configuration; first and higher order fluctuations around it are systematically constructed. The null string expansion series in  $D$ -dimensional Friedmann-Robertson-Walker and Schwarzschild spacetimes have been computed. In the latter case, the first terms of the series represent a generic approach to the string falling towards the  $r=0$  singularity. The string-integrated energy-momentum tensor and string proper size have been computed and analyzed.

The different methods to solve the string dynamics in curved spacetime have been characterized in terms of the parameter  $c$  (or equivalently, in terms of the ratio between  $\dot{X}^A$  and  $cX^{A'}$ ), as summarized in Sec. II and Table II:

- (i) ‘‘Impulsive,’’  $c \ll 1$  (null string expansion);
- (ii) ‘‘Perturbative,’’  $c = 1$  (center-of-mass expansion);
- (iii) ‘‘Adiabatic,’’  $c \gg 1$  [dual to (i), initial configuration expansion];
- (iv) ‘‘Composite,’’ ( $\tau \rightarrow 0$  expansion, ring string expansion).

The null string expansion is well appropriate to describe the string propagation in the strong gravitational field regime and well encompasses the  $\tau \rightarrow 0$  expansion in cosmological and black hole spacetimes, in the string unstable regime where the proper size of the string grows such as the scalar factor, and such as  $(R_S/r)^{(D-3)/2} \sim (-\tau)^{-(D-3)/(D+1)}$  inside the Schwarzschild black hole (near the  $r=0$  singularity).

## ACKNOWLEDGMENTS

C.O.L. was supported by NSF Grant No. PHY-95-07719 and by research funds of the University of Utah. He also thanks the DEMIRM-Observatoire de Paris, where part of this work was done, for kind hospitality and use of its working facilities.



- [1] M. B. Green, J. H. Schwarz, and E. Witten, *Superstring Theory* (Cambridge University Press, Cambridge, England, 1987).
- [2] N. D. Birrell and P. C. W. Davies, *Quantum Field Theory in Curved Space* (Cambridge University Press, Cambridge, England, 1982).
- [3] H. J. de Vega and N. Sánchez, in *Proceedings of the Erice Course: "String Quantum Gravity and Physics at the Planck Energy Scale,"* 1992, edited by N. Sánchez (World Scientific, Singapore, 1993).
- [4] *Proceedings of the Erice Course: "String Quantum Gravity and Physics at the Planck Energy Scale,"* 1995, edited by N. Sánchez and A. Zichichi, NATO Advanced Study Institutes Ser. C, Vol. 476 (Plenum, New York, 1996).
- [5] H. J. de Vega and N. Sánchez, *Phys. Lett. B* **197**, 320 (1987).
- [6] M. Gasperini, N. Sánchez, and G. Veneziano, *Int. J. Mod. Phys. A* **6**, 3853 (1991); *Nucl. Phys.* **B364**, 365 (1991).
- [7] H. J. de Vega and A. Nicolaidis, *Phys. Lett. B* **295**, 214 (1992); H. J. de Vega, I. Giannakis, and A. Nicolaidis, *Mod. Phys. Lett. A* **10**, 2479 (1995).
- [8] F. Combes, H. J. de Vega, A. V. Mikhailov, and N. Sánchez, *Phys. Rev. D* **50**, 2754 (1994).
- [9] H. J. de Vega, A. L. Larsen, and N. Sánchez, *Nucl. Phys.* **B427**, 643 (1994).
- [10] H. J. de Vega and I. L. Egusquiza, *Phys. Rev. D* **49**, 763 (1994).
- [11] A. L. Larsen and N. Sánchez, *Phys. Rev. D* **50**, 7493 (1994).
- [12] N. Sánchez and G. Veneziano, *Nucl. Phys.* **B333**, 253 (1990).
- [13] C. O. Lousto and N. Sánchez, *Phys. Rev. D* **47**, 4498 (1993).
- [14] H. J. de Vega and I. L. Egusquiza, *Phys. Rev. D* **53**, 3296 (1996).
- [15] C. Lovelace, *Phys. Lett.* **135B**, 75 (1984).
- [16] E. S. Fradkin and A. A. Tseylin, *Nucl. Phys.* **B261**, 1 (1985).
- [17] A. L. Larsen, *Phys. Rev. D* **50**, 2623 (1994).
- [18] A. L. Larsen and N. Sánchez, *Phys. Rev. D* **51**, 6929 (1995).
- [19] H. J. de Vega, A. V. Mikhailov, and N. Sánchez, *Theor. Math. Phys.* **94**, 166 (1993).
- [20] H. J. de Vega and N. Sánchez, *Phys. Lett. B* **244**, 215 (1990).
- [21] H. J. de Vega and N. Sánchez, *Phys. Rev. D* **42**, 3969 (1990).
- [22] H. J. de Vega and N. Sánchez, *Int. J. Mod. Phys. A* **7**, 3043 (1992).
- [23] R. C. Myers and M. J. Perry, *Ann. Phys. (N.Y.)* **172**, 304 (1986).