

# Electroweak symmetry-breaking Higgs boson in models with top-quark condensation

James D. Wells

*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309*

(Received 11 December 1996)

The top quark may get its large mass not from a fundamental scalar but a Nambu–Jona-Lasinio mechanism involving a strongly coupled gauge sector that triggers top-quark condensation. Forbidding a large hierarchy in the gap equation implies that top-quark condensation is a spectator to electroweak symmetry breaking, which must be accomplished mainly by another sector. The properties of the electroweak symmetry-breaking scalars are identified. Production mechanisms and decay modes are studied. Unlike the standard model, the scalar degree of freedom most relevant to electroweak symmetry breaking can only be produced by its gauge interactions. An  $e^\pm e^\pm \mu^\mp + \mu^\pm \mu^\pm e^\mp$  signal is proposed to help unambiguously detect the presence of such a gauge-coupled Higgs boson if it is light. Other useful modes of detection are also presented, and a summary is made of the search capabilities at CERN LEP II, Fermilab Tevatron, and CERN LHC. [S0556-2821(97)03415-2]

PACS number(s): 12.15.Ff, 12.15.Ji, 12.60.Fr

## I. FINE-TUNING AND THE TOP-QUARK GAP EQUATION

As it became clear that the top quark is very massive attention was naturally turned to top-quark condensate models [1–3] which rely on the gauged Nambu–Jona-Lasinio (NJL) model [4]. Starting with a four-fermion interaction

$$L = L_{\text{kin}} + G \bar{\psi}_L t_R \bar{t}_R \psi_L \quad \text{where } \psi_L = (t_L, b_L), \quad (1)$$

a gap equation can be formulated for the dynamical top-quark mass

$$G^{-1} = \frac{N_c}{8\pi^2} \left( \Lambda^2 - m_t^2 \ln \frac{\Lambda^2}{m_t^2} \right). \quad (2)$$

Furthermore, in the fermion bubble approximation, with the above gap equation enforced, zeros of the inverse gauge boson propagators are found at nonzero momentum. This gives mass to the gauge bosons and corresponds [5,3,6] to a decay constant of

$$f_{\pi_t}^2 \simeq \frac{N_c}{16\pi^2} m_t^2 \ln \frac{\Lambda^2}{m_t^2}. \quad (3)$$

This relation is often referred to as the Pagels-Stokar formula. In order for the top-quark condensate to account for all of electroweak symmetry breaking,  $f_{\pi_t}$  needs to be equal to  $v = 175$  GeV. The currently measured top-quark mass is  $m_t = 175 \pm 6$  GeV [7], which means that  $\Lambda$  would have to be about 13 orders of magnitude above  $m_W$  to account for all of electroweak symmetry breaking, an uninspiring result for pure top-quark condensate models.

The Pagels-Stokar relation is logarithmically sensitive to the condensate scale, and provides only a relationship between the decay constant and dynamical quark mass, not the overall scale of these parameters. The gap equation, on the other hand, sets the overall scale of  $m_t$  and, therefore,  $f_{\pi_t}$ . The quadratic sensitivity to the condensate scale induces a large hierarchy problem for the weak scale ( $m_t$  and  $f_{\pi_t}$ ) if  $\Lambda$

is above a few TeV. For large  $\Lambda$  the four-fermion coupling  $G$  must be tuned to one part in  $\Lambda^2/m_t^2$ . For a condensate scale in the multi-TeV region, this fine-tuning is greater than one part in  $10^3$ . It will be assumed here that fine-tunings much above this are unnatural, and are probably not maintained by nature [8]. Therefore, from the Pagels-Stokar relation the decay constant associated with top-quark condensation is  $f_{\pi_t} \simeq 60$  GeV, implying that top-quark condensation is a spectator to electroweak symmetry breaking (EWSB). That is, top-quark condensation mainly provides the top quark its mass, but another sector mainly provides the  $W$  and  $Z$  bosons their masses.

The above argument has led to the conception of top-quark condensate-assisted models such as those discussed in Refs. [9–12]. Much emphasis and study have been devoted to the bound-state scalars and gauge bosons associated with top-quark condensation in these models [9–14]. Attention is given here to the degrees of freedom which bear the most burden for electroweak symmetry breaking. In the following, a single electroweak symmetry-breaking doublet is postulated to give mass to the  $W$ ,  $Z$ , and all light fermions, and perhaps a small contribution to the top and bottom quarks. The production and decay modes of this doublet are detailed. The results will be significantly different from those of the standard model doublet, since unlike the standard model the resulting scalar field has a small coupling to the top quark. The complexity of the electroweak symmetry-breaking sector will have little effect on the detectability of the scalars associated with electroweak symmetry breaking.

## II. TOP-QUARK CONDENSATE SCALARS

Top-quark condensation only requires a critical four-fermion coupling in a Lagrangian such as that given in Eq. (1). Explaining the origin of such a strong interaction is a more difficult question. Topcolor is one such explanation [15,9,10,12], which is used as an example here. Upon integrating out the heavy vector bosons from the topcolor gauge group and the “tilting”  $U(1)$  gauge group, one obtains the NJL interaction Lagrangian

$$L = L_{\text{kin}} + G_t \bar{\psi}_L t_R \bar{t}_R \psi_L + G_b \bar{\psi}_L b_R \bar{b}_R \psi_L. \quad (4)$$

Tuning the couplings of the different gauge interactions which formed this four-fermion interaction Lagrangian, one can obtain  $G_t > 8\pi^2/N_c \Lambda^2 > G_b$ , which induces a  $\langle t\bar{t} \rangle$  condensate but not a  $\langle b\bar{b} \rangle$  condensate.

It is convenient to introduce auxiliary fields  $\phi_t$  and  $\phi_b$  and rewrite the above Lagrangian as

$$L = L_{\text{kin}} - (\bar{\psi}_L \phi_t t_R + \bar{\psi}_L \phi_b b_R + \text{H.c.}) - G_t^{-1} \phi_t^\dagger \phi_t - G_b^{-1} \phi_b^\dagger \phi_b. \quad (5)$$

Renormalization group evolution [3,12,16] below  $\Lambda$  makes the scalar fields  $\phi_t$  and  $\phi_b$  dynamical, and it induces a vacuum expectation value (VEV) for  $\phi_t$  (not  $\phi_b$ ) in accord with the gap equation. The pion decay constant of  $\phi_t$  can be solved via the Pagels-Stokar formula of Eq. (3) given particular values of  $m_t$  and  $\Lambda$ . For the purposes of this paper we will assume the numerical value  $f_{\pi_t} \simeq 50$  GeV ( $\Lambda \simeq 1.5$  TeV).

In the approximation that  $f_{\pi_t}^2 \ll v^2$  ( $v$  is normalized to 175 GeV) top-quark condensation is merely a spectator to electroweak symmetry breaking (in analogy to chiral symmetry breaking among the light quarks) and so it does not mix into the longitudinal components of the  $W$  and  $Z$ . Therefore, all the component states of  $\phi_t$  and  $\phi_b$  are physical eigenstates to a good approximation:

$$\phi_t = \begin{pmatrix} f_{\pi_t} + \frac{1}{\sqrt{2}}(h_t^0 + ia_t^0) \\ h_t^\pm \end{pmatrix}, \quad \phi_b = \begin{pmatrix} h_b^\pm \\ \frac{1}{\sqrt{2}}(h_b^0 + ia_b^0) \end{pmatrix}. \quad (6)$$

In the fermion bubble approximation the scalar  $t\bar{t} \rightarrow t\bar{t}$  scattering amplitude has a pole at  $p^2 = 4m_t^2$  which can be identified with the mass of the  $h_t^0$ . Other masses are dependent upon details of the underlying model and can vary from about 150 to 350 GeV [9,11,12]. The top quark gets almost all its mass by a large Yukawa coupling to  $\phi_t$ , and it can be assumed that the bottom quark gets almost all its mass from topcolor instanton effects [9,12].

In addition to these bound-state topcolor scalars, there needs to be degrees of freedom that break  $\text{SU}(2) \times \text{U}(1)_Y$ . To do this we introduce another doublet  $\phi_{\text{ew}}$  whose vacuum expectation value is such that

$$f_{\pi_t}^2 + f_{\pi_{\text{ew}}}^2 = v^2 = (175 \text{ GeV})^2. \quad (7)$$

We also assume that this doublet is a fundamental scalar and gives a small mass to all the light fermions in the theory in addition to breaking electroweak symmetry. Even though the following assumes a fundamental scalar-breaking electroweak symmetry, it might be that  $\text{SU}(2) + \text{U}(1)_Y$  is broken by some other strongly interacting sector which produces a narrow resonance composite  $\phi_{\text{ew}}$  which has very similar properties to the fundamental Higgs state discussed below.

It will be convenient later to introduce a ‘‘top-quark condensate spectator angle’’  $\theta_t$ , which is defined as

$$f_{\pi_{\text{ew}}} = v \sin \theta_t,$$

$$f_{\pi_t} = v \cos \theta_t.$$

Using the numerical value of  $f_{\pi_t} = 50$  GeV, one obtains

$$\cos \theta_t = \frac{50 \text{ GeV}}{175 \text{ GeV}} \simeq 0.3. \quad (8)$$

The pseudoscalar and charged Higgs fields in the  $\phi_{\text{ew}}$  doublet all get absorbed by the  $W^\pm$  and  $Z$  bosons, and what remains physical is a single scalar field  $h_{\text{ew}}^0$ , whose mass cannot be determined *a priori*. Note also that since  $f_{\pi_t} \ll v$ , and since  $h_{\text{ew}}^0$  does not couple to top quarks (or couples very weakly), then  $h_{\text{ew}}^0$  is a mass eigenstate.

The assumption of just one scalar providing electroweak symmetry breaking is not important for the qualitative results below. The most important assumption is that a  $t\bar{t}$  condensate gives the top quark its large mass. If we added more condensing scalar fields to the spectrum, it is possible to have large Yukawa couplings of these scalars to the light quark fields. However, in this case these additional fields would also have to be spectators to electroweak symmetry breaking, just as  $\phi_t$  is, and there would remain the necessity of a scalar(s) with large vacuum expectation values which have small Yukawa interactions with the fermions.

If the electroweak symmetry-breaking sector were more complicated, more angles would need to be introduced. For example, if a two-Higgs doublet model were involved then a new angle  $\beta$  would need to be introduced which specifies the ratios of the VEV’s of these two fields. Defining  $\tan \beta \equiv \langle \phi_{\text{ew},2} \rangle / \langle \phi_{\text{ew},1} \rangle$ , then

$$f_{\pi_t} = v \cos \theta_t,$$

$$f_{\pi_{\text{ew},1}} = v \sin \theta_t \cos \beta,$$

$$f_{\pi_{\text{ew},2}} = v \sin \theta_t \sin \beta.$$

This angle  $\beta$  does not enhance the coupling of the neutral scalars to the top quark but does reduce the coupling to gauge bosons by a factor of  $\sin^2(\beta - \alpha)$  or  $\cos^2(\beta - \alpha)$  where  $\alpha$  is the mass eigenstate mixing angle between  $h_{\text{ew},1}^0$  and  $h_{\text{ew},2}^0$ . The argument generalizes to an arbitrarily complex electroweak symmetry-breaking sector with doublets to show that Yukawa production processes are negligibly affected by additional complexity in the EWSB sector, and gauge production processes can only be *reduced*. Thus, production modes for the scalars will decrease with additional complexity. Higher dimensional  $\text{SU}(2)$  representations such as triplets are not considered since they explicitly break custodial  $\text{SU}(2)$  symmetry. Barring fine-tuned cancellations, triplets are incapable of providing a substantial fraction of EWSB due to the current  $\rho$  parameter constraints [17]. Since our conclusion will be that the gauge-coupled EWSB scalars are extremely difficult to reconstruct at the Large Hadron Collider (LHC), it is most conservative to maximize the gauge boson couplings of the EWSB scalar(s) by postulating one single  $\phi_{\text{ew}}$  which breaks electroweak symmetry, and analyzing the consequences of this assumption.




FIG. 1. Branching fractions of  $h_{SM}^0$  versus its mass.

### III. DECAYS OF THE $h_{ew}^0$

The decay of the  $h_{ew}^0$  (the neutral scalar component of  $\phi_{ew}$ ) has a few inherent uncertainties. The most important uncertainty is how much of the  $b$ -quark mass is derived from  $f_{\pi_{ew}}$ . This sets the coupling of the  $\phi_{ew}$  to the  $b$  quark, which in turn sets the partial width of  $h_{ew}^0 \rightarrow b\bar{b}$ . The instanton-induced  $b$ -quark mass [9,12] can be substantial and provide the  $b$  quark with all of its mass, leaving a very small (if any) residual  $b$ -quark mass contribution from  $\phi_{ew}$ . Furthermore, the hierarchy of masses generated by  $\phi_{ew}$  need not be in descending order of generations. That is, the  $b$ -quark mass induced by  $\phi_{ew}$  could be much smaller than that of the strange quark. The working assumption will be that the  $b$ -quark coupling to  $\phi_{ew}$  is negligible. However, it cannot be ruled out that all of the  $b$ -quark mass comes from  $\phi_{ew}$  and this possibility will be commented on later.

In the standard model all the couplings to the gauge bosons and fermions are set without ambiguity, and the decay branching fractions can be calculated [18–20]. In Fig. 1 the decay branching fractions of the standard model are displayed for  $50 \text{ GeV} < m_{h_{SM}^0} < 200 \text{ GeV}$ . Above 200 GeV the  $WW$  and  $ZZ$  branching fractions dominate in a ratio of 2 to 1 for a Higgs boson mass well above the  $ZZ$  threshold. Above the  $t\bar{t}$  threshold the branching ratio to  $t\bar{t}$  reaches its maximum of 20% at  $m_{h_{SM}^0} = 500 \text{ GeV}$  and slowly reduces to 10% at  $m_{h_{SM}^0} = 1 \text{ TeV}$ . ( $m_t = 175 \text{ GeV}$  is assumed throughout this discussion.) Note the dominance of the  $b\bar{b}$  final state up to  $m_{h_{SM}^0} \leq 135 \text{ GeV}$ . Above this, the branching fraction into  $WW$  becomes dominant.

The branching fractions of  $h_{ew}^0$  can likewise be calculated once the angle  $\cos\theta_t$  is known, and the  $b$ -quark coupling to  $h_{ew}^0$  is specified. In Fig. 2 the decay branching fractions are shown for  $\cos\theta_t = 0.3$  and negligible  $b$ -quark coupling to  $h_{ew}^0$ . The  $\tau^+\tau^-$  and  $c\bar{c}$  final states are more important for  $h_{ew}^0$  than for  $h_{SM}^0$ . Furthermore, the  $WW^*$  branching fraction is much enhanced despite the  $\sin^2\theta_t$  suppression of the partial width to  $WW^*$ . The branching fractions of  $h_{ew}^0$  are much different than those of the standard model Higgs boson for  $m_{h_{ew}^0} \leq 140 \text{ GeV}$ . Above 140 GeV the  $h_{ew}^0$  branching fractions are similar to those of the standard model except that  $h_{ew}^0 \rightarrow t\bar{t}$  always has a negligible rate. A further important




FIG. 2. Branching fractions of  $h_{ew}^0$  versus its mass for  $\cos\theta_t = 0.3$  and negligible  $b$ -quark coupling to  $h_{ew}^0$ .

distinction from the standard model is the substantially lower production rate of  $h_{ew}^0$  compared to  $h_{SM}^0$  at a hadron collider discussed in the next section.

### IV. PRODUCTION OF $h_{ew}^0$

The production modes of a standard model Higgs boson can be broken up into three categories: gauge processes  $q\bar{q} \rightarrow Zh_{SM}^0$ ,  $q\bar{q} \rightarrow W^\pm h_{SM}^0$ ,  $WW(ZZ) \rightarrow h_{SM}^0$ ,  $e^+e^- \rightarrow Zh_{SM}^0$ ,  $e^+e^- \rightarrow e^+e^- h_{SM}^0$ ; Yukawa processes  $q\bar{q} \rightarrow t\bar{t}h_{SM}^0$ ,  $gg \rightarrow h_{SM}^0$ ,  $e^+e^- \rightarrow t\bar{t}h_{SM}^0$ ; gauge or Yukawa processes  $\gamma\gamma \rightarrow h_{SM}^0$ .

The  $gg \rightarrow h_{SM}^0$  production mechanism is called a ‘‘Yukawa process’’ because it has a non-negligible rate due to the large top-quark Yukawa coupling of the  $h_{SM}^0$  to the top quark in the loop. The  $\gamma\gamma \rightarrow h_{SM}^0$  is called a ‘‘gauge or Yukawa process’’ since it has non-negligible contributions from both  $W$  and top-quark loops.

The scalars in top-quark condensate models [9,11,12,14] generally either get produced by gauge processes or by Yukawa process. For example, the top Higgs boson  $h_t^0$  has a small gauge production process suppressed relative to the standard model by  $\cos^2\theta_t \approx 0.08$ . On the other hand, its production via Yukawa processes is enormous due to the large Yukawa coupling of the top quarks to  $\phi_t$ . The bottom Higgs boson  $h_b^0$  has approximately zero production cross section via gauge processes, however, it has a reasonable Yukawa-induced cross section from  $gg \rightarrow h_b^0$  and even  $q\bar{q}(e^+e^-) \rightarrow b\bar{b}h_b^0$ .

The electroweak symmetry-breaking scalar  $h_{ew}^0$  has negligible production cross section due to Yukawa processes. It must be produced via gauge interactions, which are slightly suppressed compared to the gauge interactions of the standard model Higgs boson. The cross sections are related to the standard model cross sections by

$$\sigma_{\text{gauge}}(h_{ew}^0) = \sin^2\theta_t \sigma_{\text{gauge}}(h_{SM}^0), \quad (9)$$

where  $\sigma_{\text{gauge}}(h)$  is either  $\sigma(Wh)$ ,  $\sigma(Zh)$ , or  $\sigma[WW(ZZ) \rightarrow h]$ , etc. For  $f_{\pi_t} = 50 \text{ GeV}$ , the suppression factor is merely  $\sin^2\theta_t \approx 0.92$ .




FIG. 3. Cross sections for  $jjh_{\text{ew}}^0$  (dotted),  $Wh_{\text{ew}}^0$  (dashed), and  $Zh_{\text{ew}}^0$  (dot dashed) at the LHC (14 TeV) for  $\cos\theta_i=0.3$ . Also, the standard model Yukawa production cross section  $gg \rightarrow h_{\text{SM}}^0$  (solid) is shown to demonstrate the relatively low intrinsic rate of the gauge production cross sections.

The lack of Yukawa production processes for  $h_{\text{ew}}^0$  is a severe handicap for a search at the LHC. The dominant production cross section for the standard model Higgs boson is  $gg \rightarrow h_{\text{SM}}^0$ . In Fig. 3 the cross sections for  $jjh_{\text{ew}}^0$ ,  $Wh_{\text{ew}}^0$ , and  $Zh_{\text{ew}}^0$  at the LHC ( $\sqrt{s}=14$  TeV) are shown. The  $jjh_{\text{ew}}^0$  cross section is from  $WW(ZZ) \rightarrow h_{\text{ew}}^0$  where the two jets are the ones which radiate the  $W$ 's and are generally at high rapidity. We see that the total production rate for  $h_{\text{ew}}^0$  is much smaller than that for  $h_{\text{SM}}^0$  due to lack of the  $gg \rightarrow h_{\text{ew}}^0$  Yukawa mode. The  $h_{\text{ew}}^0 \rightarrow \gamma\gamma$  signal for  $100 \text{ GeV} \lesssim m_{h_{\text{ew}}^0} \lesssim 140 \text{ GeV}$  is no longer as viable since the total rate of Higgs boson production is not entirely compensated by the somewhat enhanced branching fraction into two photons. Detecting  $h_{\text{ew}}^0$ , if it is possible, must rely on the smaller gauge production cross sections.

## V. DETECTING THE $h_{\text{ew}}^0$

The search for the standard model Higgs boson has been well studied and the strategy is well mapped [18]. The CERN  $e^+e^-$  collider LEP II running at  $\sqrt{s}=192$  GeV will be able to see the Higgs boson up to about 95 GeV with  $150 \text{ pb}^{-1}$  per experiment. Above this, the CERN Large Hadron Collider (LHC) experiments should be able to see the standard model Higgs boson through  $gg \rightarrow h_{\text{SM}}^0 \rightarrow \gamma\gamma$  up to about 140 GeV, and through  $gg \rightarrow h_{\text{SM}}^0 \rightarrow ZZ^{(*)}$  if  $140 \text{ GeV} \lesssim m_{h_{\text{SM}}^0} \lesssim 700 \text{ GeV}$ . For scalar resonances above about 700 GeV the concept of a perturbatively coupled Higgs boson becomes more tenuous, and longitudinal  $W$  scattering analyses become more important.

The common thread among all the searches described above is the  $gg \rightarrow h_{\text{SM}}^0$  Yukawa production mode. In the standard model this process dominates the production rate for Higgs bosons. Some studies have been performed on the ability to detect the Higgs boson via  $Wh_{\text{SM}}^0$  [21–24], which is a gauge-colored production mode. The hope is that with two tagged  $b$  quarks from  $h_{\text{SM}}^0$  and a high  $p_T$  lepton tag from the  $W$ , the small rate of  $Wh_{\text{SM}}^0$  could still be pulled out from the background ( $WZ$ ,  $Wb\bar{b}$ ,  $t\bar{t}$ , etc.). At the LHC it was con-

cluded by Ref. [23] that this mode could be useful if  $80 \text{ GeV} \lesssim m_{h_{\text{SM}}^0} \lesssim 100 \text{ GeV}$ , whereas Ref. [24] finds that  $m_{h_{\text{SM}}^0} \lesssim 130 \text{ GeV}$  are detectable in this mode with excellent bottom quark jet identification.

For the  $h_{\text{ew}}^0$  state, the LEP II capability should not be altered. The  $\tau$  branching fraction of the Higgs boson will be large and the backgrounds from  $ZZ$  where one  $Z$  decays to  $\tau^+\tau^-$  will be relatively small since that branching fraction is only about 3%. Searches at the LHC with  $m_{h_{\text{ew}}^0} \gtrsim 100 \text{ GeV}$  are more difficult. In the range from  $100 \text{ GeV} \lesssim m_{h_{\text{SM}}^0} \lesssim 135 \text{ GeV}$  the branching ratio to  $\tau^+\tau^-$  is greater than 8%. The signal  $pp \rightarrow Wh_{\text{SM}}^0$  where  $W \rightarrow jj$  and  $h_{\text{SM}}^0 \rightarrow \tau^+\tau^-$  has been studied in Ref. [22]. It was concluded that a significant signal at the LHC could not be found in this mode for a standard model Higgs boson. However, such a signal might be possible in this  $\tau^+\tau^-$  decay mode at the Tevatron running at  $\sqrt{s}=2$  TeV with  $30 \text{ fb}^{-1}$  of integrated luminosity. Turning back to the  $h_{\text{ew}}^0$ , the  $\tau^+\tau^-$  decay branching fraction is much increased. The significance of the standard model  $\tau^-\tau^+$  signal now should be multiplied by a factor of  $B(h_{\text{ew}}^0 \rightarrow \tau^+\tau^-)/B(h_{\text{SM}}^0 \rightarrow \tau^+\tau^-) \approx 6$ . It, therefore, might be possible to detect the light  $h_{\text{ew}}^0$  at the Tevatron with compelling significance if  $m_{h_{\text{ew}}^0} \lesssim 130 \text{ GeV}$ . Furthermore, this signal might also be detectable at the LHC with sufficient luminos-

ity. Earlier, we postulated that  $h_{\text{ew}}^0$  does not couple to the  $b$  quarks. If we now relax that assumption and assume the opposite extreme, that all the  $b$ -quark mass comes from  $\phi_{\text{ew}}$ , then the branching fraction of  $h_{\text{ew}}^0$  into fermions for  $m_{h_{\text{ew}}^0} \lesssim 130 \text{ GeV}$  will be almost identical to that of the standard model. Then the analyses of the  $Wh_{\text{ew}}^0$  signal will become most important, and from studies in the standard model (especially, Ref. [24]), it can be concluded that a  $b$ -quark-coupled  $h_{\text{ew}}^0$  is detectable at the LHC with as low as  $10 \text{ fb}^{-1}$  of data. The Tevatron probably could not detect this  $b$ -coupled Higgs state above the LEP II limit of 95 GeV in the  $b\bar{b}$  mode alone [21,23], but could possibly be detected up to 130 GeV using both the  $b\bar{b}$  and  $\tau^+\tau^-$  modes [22].

If  $m_{h_{\text{ew}}^0} \gtrsim 130 \text{ GeV}$ , then it appears that the Tevatron and LHC will have difficulty seeing a signal and reconstructing the  $h_{\text{ew}}^0$  mass. In this range,  $h_{\text{ew}}^0$  will decay to  $WW^*$  most of the time. Since the production rate for  $h_{\text{ew}}^0$  is small, the  $ZZ^{(*)}$  decay cannot be separated from backgrounds to construct a signal. Without this mode at our disposal it appears near impossible to measure the Higgs boson mass since the  $WW^{(*)}$  decay mode does not allow the Higgs boson mass to be reconstructed.

However, it might be possible to detect a Higgs signal without being able to reconstruct the mass. If the Higgs boson decays to  $W^+W^-$  (either on shell or off shell), then the  $pp \rightarrow W^\pm h_{\text{ew}}^0$  produces a three lepton plus  $\cancel{E}_T$  signal if all the  $W$ 's decay leptonically. This signal is relatively clean in a hadron collider environment and has been proposed as a search tool for supersymmetric gaugino production of  $\widetilde{WZ}$  with subsequent leptonic decays into three leptons plus missing energy carried away by the stable lightest supersymmetric particle [29]. Because of this potentially large supersym-




FIG. 4. The total event rate of  $e^{\pm}e^{\pm}\mu^{\mp} + \mu^{\pm}\mu^{\pm}e^{\mp}$  expected from  $pp \rightarrow Wh_{ew}^0 \rightarrow WWW^{(*)}$  production with  $100 \text{ fb}^{-1}$  of data collected at the LHC with center-of-mass energy 14 TeV. A  $5\sigma$  signal above background requires at least 100 signal events.

metric rate, it may be difficult to tell the difference between a  $Wh_{ew}^0 \rightarrow lll + \cancel{E}_T$  signature as opposed to a  $\widetilde{W}\widetilde{Z} \rightarrow lll + \cancel{E}_T$  signature.

Some permutations of these trilepton decays yield a like-sign dilepton signal with missing energy and little background [21]. Potential backgrounds to like-sign dileptons include  $WZ$  production where one lepton comes from the  $W$  and one lepton comes from the  $Z$ . Cutting out  $m_{l+l-} \approx m_Z$  reduces this background. Another background comes from  $t\bar{t}$  production where one lepton comes from  $W$  decay in the  $t$  decay and the like-sign lepton comes from a  $b$  decay in the  $\bar{t}$  decay. This background can be greatly reduced by an effective jet veto of the hadronic activity of the  $b$  quark, requiring no tagged  $b$  quark in the event, and also requiring that all the leptons be isolated. Here, we propose a more discerning signal of like-sign dileptons with a different flavor third lepton. Using just the first two generations of leptons, the signal for the presence of a Higgs boson is  $e^{\pm}e^{\pm}\mu^{\mp}$  and  $\mu^{\pm}\mu^{\pm}e^{\mp}$ , with all leptons isolated. This is a useful signal to distinguish  $Wh_{ew}^0$  production as opposed to the supersymmetric  $\widetilde{W}\widetilde{Z}$  production since like-sign dileptons from the latter require all three leptons to be of the same flavor. The branching fraction into  $WW^{(*)}$  is given in Fig. 2. Using the production cross section from Fig. 3 the total number of signal events in  $100 \text{ fb}^{-1}$  of data is displayed in Fig. 4.

The main background from  $t\bar{t}$  can be estimated using the geometric acceptance of leptons ( $|\eta| < 2.5$ ) and jets ( $|\eta| < 5$ ), the combinatorics of the decays which yield  $e^{\pm}e^{\pm}\mu^{\mp}$  and  $\mu^{\pm}\mu^{\pm}e^{\mp}$  signal events, and the total inclusive  $t\bar{t}$  cross section at LHC, which will be conservatively bounded at 2 nb. The number of background events estimated from  $t\bar{t}$  is approximately 100 for  $100 \text{ fb}^{-1}$  of data. (Note that there are no background events from  $WZ$ .) After applying the same lepton acceptance cuts to the signal, a  $5\sigma$  effect would require at least 100 signal events (using  $S/\sqrt{B}$  for significance). Thus, the electroweak symmetry-breaking Higgs boson  $h_{ew}^0$  could be detected in the range  $110 \text{ GeV} \lesssim m_{h_{ew}^0} \lesssim 175 \text{ GeV}$  by this process. This estimate is based solely on parton level calculations and is only meant to demonstrate that this signal could be useful at the LHC. Estimates of nonphysics backgrounds such as misidentification,




FIG. 5. The total event rate of  $e^+e^- \mu^+ \mu^-$  expected from  $VV \rightarrow h_{ew}^0 \rightarrow ZZ^{(*)}$  production with  $100 \text{ fb}^{-1}$  of data collected at the LHC with center-of-mass energy 14 TeV. A  $5\sigma$  signal above background from this mode is estimated to be possible for the mass ranges  $120 \text{ GeV} \lesssim m_{h_{ew}^0} \lesssim 155 \text{ GeV}$  and  $200 \text{ GeV} \lesssim m_{h_{ew}^0} \lesssim 400 \text{ GeV}$ .

and a full jet-level Monte Carlo simulation could diminish the significance found here. However, it is likely that additional cuts not studied here could further distinguish the  $Wh_{ew}^0 \rightarrow WWW$  topology over  $t\bar{t} \rightarrow WWbb$ , thereby increasing the significance of the signal. Also, using the  $\tau$  final states could help somewhat since  $b$  decays to  $\tau$ 's are kinematically suppressed.

From Fig. 3 we can see that the  $WW(ZZ) \rightarrow h_{ew}^0$  vector-boson fusion provides the highest production rate for  $h_{ew}^0$ . Along with the Higgs boson comes two high rapidity jets corresponding to the quark lines which radiated the initial vector bosons. For low mass Higgs bosons, where  $h_{ew}^0 \rightarrow ZZ^{(*)}$ , the signal would be four leptons with high invariant masses,  $m_{l+l-}$  from each decay. The standard model background is dominated by continuum  $Z\gamma^*$  production with enhancement at small invariant  $m_{l+l-}$ . Identifying the  $Z^{(*)}$  as the lower mass  $m_{l+l-}$  and placing a cut on how low this invariant mass very effectively removes the background while retaining almost all of the signal [18]. It might be possible to detect the light  $h_{ew}^0$  from vector-boson fusion with as low as 40 signal events (before cuts) in this mode [18]. The signal rate for  $VV \rightarrow h_{ew}^0 \rightarrow ZZ^{(*)} \rightarrow l^+l^-l'^+l'^-$  ( $l, l' = e$  or  $\mu$ ) is shown in Fig. 5. In the  $ZZ^{(*)}$  region, the detectability requirement of 40 signal events corresponds to a Higgs boson mass range of  $120 \text{ GeV} \lesssim m_{h_{ew}^0} \lesssim 155 \text{ GeV}$ .

The region between the  $WW$  and  $ZZ$  threshold is much more difficult. The  $h_{ew}^0 \rightarrow ZZ^{(*)}$  branching fraction is greatly reduced in this region because  $h_{ew}^0 \rightarrow WW$  is above threshold and rapidly increasing. Therefore,  $VV \rightarrow h_{ew}^0 \rightarrow ZZ^{(*)}$  is not viable. From above it appears that it might be possible to see  $pp \rightarrow Wh_{ew}^0 \rightarrow WWW$  up to about 175 GeV. However, above 175 GeV the trilepton signal runs out. Thus, the small region between  $175 \text{ GeV} \lesssim m_{h_{ew}^0} \lesssim 180 \text{ GeV}$  must rely on the difficult task of seeing a resonant enhancement in  $VV \rightarrow h_{ew}^0 \rightarrow WW$ . In the standard model, the large  $gg \rightarrow h_{SM}^0$  rate allows for the possibility of seeing the Higgs boson in the  $h_{SM}^0 \rightarrow WW$  channel for the range  $155 \text{ GeV} \lesssim m_{h_{SM}^0} \lesssim 180 \text{ GeV}$  [25,26]. It has been estimated that with  $5 \text{ fb}^{-1}$  a signal with greater than  $5\sigma$  significance could be ob-

served in this mass range for the standard model Higgs boson using the  $h_{\text{SM}}^0 \rightarrow WW \rightarrow \nu l \nu' l'$  decay mode [26]. The gauge-coupled Higgs boson  $h_{\text{ew}}^0$  does not get produced by  $gg$  fusion but rather  $WW(ZZ)$  fusion. In this mass range the total cross section is more than a factor of 6 lower than the  $gg$  cross section. Taking this into account and scaling the significance result of Ref. [26] up to  $100 \text{ fb}^{-1}$  there is an  $8\sigma$  signal available in the range  $155 \text{ GeV} \leq m_{h_{\text{SM}}^0} \leq 180 \text{ GeV}$ . It should be pointed out, however, that some of the cuts employed to reduce background may not be as effective if the underlying signal event is produced mainly by  $WW(ZZ)$  fusion rather than  $gg$  fusion. For example, the  $|\cos\theta_{l+l'}| < 0.8$  cut to reduce continuum  $q\bar{q} \rightarrow W^+W^-$  background may not be as useful. However, it might be possible to gain significance by tapping the high  $p_T$ , high rapidity jets which radiate the initial vector bosons [27]. A central mini-jet veto [28] which takes advantage of the different color flow of the signal compared to background may also be helpful. Again, confusion with a supersymmetric signal is possible, especially for slepton and chargino pair production which both can yield  $ll' + \cancel{E}_T$  signatures at high rates. The spectator jet tags would be especially useful to eradicate this potential confusion.

Above the  $h_{\text{ew}}^0 \rightarrow ZZ$  threshold the Higgs signal would be a Higgs bump in the  $ZZ$  invariant mass spectrum in the total  $ZZ$  production cross section. Again, the  $Z$ 's are required to decay into leptons. Figure 5 gives the total rate above the  $ZZ$  threshold and can be compared with the total  $ZZ$  production cross section for sensitivity of this invariant mass peak [30]. The resonant Higgs signal is smeared according to the detector's electromagnetic resolution, which is taken to be  $15\%/\sqrt{E_T}$ . Taking into consideration lepton acceptance, the

significance is calculated for signal events in a bin centered on  $m_{ZZ} = m_{h_{\text{ew}}^0}$  with a width determined by the two standard deviation smearing of the Higgs resonance corresponding to the resolution factor given here. The significance reaches a peak of about  $7\sigma$  at  $m_{h_{\text{ew}}^0} = 220 \text{ GeV}$  and falls below  $5\sigma$  for  $m_{h_{\text{ew}}^0} \geq 400 \text{ GeV}$ . This estimate is based only on the  $h_{\text{ew}}^0 \rightarrow ZZ \rightarrow e^+e^-\mu^+\mu^-$  mode. Using all lepton permutations at the expense of potential lepton pairing ambiguity should extend the search range well above 400 GeV. Again, a realistic detector simulation is required to obtain a precise range. Nevertheless, it appears possible to discern a Higgs peak up to high mass scales with  $100 \text{ fb}^{-1}$ .

In short, top-quark condensate models may imply at least one scalar degree of freedom  $h_{\text{ew}}^0$  which breaks electroweak symmetry and has negligible Yukawa production modes. The LEP II search range should not be different than that of the standard model. The Tevatron  $h_{\text{ew}}^0$  search is even enhanced over the standard model Higgs search range. On the other hand, it will be more challenging at the LHC to demonstrate that such a gauge-coupled Higgs boson such as  $h_{\text{ew}}^0$  can be detected, and most especially that its mass can be reconstructed. However, over most of the perturbative Higgs boson mass range detection should be possible with  $100 \text{ fb}^{-1}$  of data.

#### ACKNOWLEDGMENTS

The author has benefited from discussions with G. Buchalla, S. Martin, and M. Peskin. This work was supported by the U.S. Department of Energy under Contract No. DE-AC03-76SF00515.

- 
- [1] Y. Nambu, in *New Theories in Physics*, Proceedings of the XI International Symposium on Elementary Particle Physics, Kazimierz, Poland, 1988, edited by Z. Ajduk, S. Pokorski, and A. Trautman (World Scientific, Singapore, 1989).
  - [2] V. Miransky, M. Tanabashi, and K. Yamawaki, *Mod. Phys. Lett. A* **4**, 1043 (1989); *Phys. Lett. B* **221**, 177 (1989).
  - [3] W. Bardeen, C. T. Hill, and M. Lindner, *Phys. Rev. D* **41**, 1647 (1990).
  - [4] Y. Nambu and G. Jona-Lasinio, *Phys. Rev. D* **122**, 345 (1961).
  - [5] H. Pagels and S. Stokar, *Phys. Rev. D* **20**, 2947 (1979); A. Carter and H. Pagels, *Phys. Rev. Lett.* **43**, 1845 (1979); R. Jackiw and K. Johnson, *Phys. Rev. D* **8**, 2386 (1973).
  - [6] T. Gherghetta, *Phys. Rev. D* **50**, 5985 (1994).
  - [7] P. Tipton, talk given at the *28th International Conference on High-energy Physics (ICHEP 96)*, Warsaw, Poland, 1996 (unpublished).
  - [8] If present supersymmetry could stabilize the hierarchy and provide a lower top quark mass fixed point to be compatible with a high scale  $\langle t\bar{t} \rangle$  condensate: T. E. Clark, S. T. Love, and W. ter Veldhuis, *Mod. Phys. Lett. A* **6**, 3225 (1991); M. Carena, T. E. Clark, C. E. M. Wagner, W. A. Bardeen, and K. Sasaki, *Nucl. Phys.* **B369**, 33 (1992).
  - [9] C. T. Hill, *Phys. Lett. B* **345**, 483 (1995).
  - [10] K. Lane and E. Eichten, *Phys. Lett. B* **352**, 382 (1995).
  - [11] D. Kominis, *Phys. Lett. B* **358**, 312 (1995).
  - [12] G. Buchalla, G. Burdman, C. T. Hill, and D. Kominis, *Phys. Rev. D* **53**, 5185 (1996).
  - [13] E. Eichten and K. Lane, *Phys. Lett. B* **388**, 803 (1996); Report No. hep-ph/9609298 (unpublished); R. S. Chivukula and J. Terning, *Phys. Lett. B* **385**, 209 (1996); B. Balaji, *ibid.* **393**, 89 (1997).
  - [14] G. Burdman, Report No. hep-ph/9611265 (unpublished).
  - [15] C. T. Hill, *Phys. Lett. B* **266**, 419 (1991); S. P. Martin, *Phys. Rev. D* **46**, 2197 (1992); **45**, 4283 (1992); *Nucl. Phys.* **B398**, 359 (1993); M. Lindner and D. Ross, *ibid.* **B370**, 30 (1992); R. Bönisch, *Phys. Lett. B* **268**, 394 (1991); C. T. Hill, D. Kennedy, T. Onogi, and H. L. Yu, *Phys. Rev. D* **47**, 2940 (1993).
  - [16] A. Blumhofer, R. Dawid, and M. Lindner, *Phys. Lett. B* **360**, 123 (1995).
  - [17] T. G. Rizzo, *Mod. Phys. Lett. A* **6**, 1961 (1991); P. Bamert and Z. Kunszt, *Phys. Lett. B* **306**, 335 (1993); T. G. Rizzo (private communication).
  - [18] J. Gunion, H. Haber, G. Kane, and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Reading, MA, 1990); J. Gunion, A. Stange, and S. Willenbrock, Report No. hep-ph/9602238 (unpublished).

- [19] M. Spira, A. Djouadi, D. Graudenz, and P. M. Zerwas, Nucl. Phys. **B453**, 17 (1995).
- [20] Z. Kunszt, S. Moretti, and W. J. Stirling, Report No. hep-ph/9611397 (unpublished).
- [21] A. Stange, W. Marciano, and S. Willenbrock, Phys. Rev. D **50**, 4491 (1994).
- [22] S. Mrenna and G. Kane, Report No. hep-ph/9406337 (unpublished).
- [23] D. Froidevaux and E. Richter-Was, Z. Phys. C **67**, 213 (1995).
- [24] P. Agrawal, D. Bowser-Chao, and K. Cheung, Phys. Rev. D **51**, 6114 (1995).
- [25] E. W. N. Glover, J. Ohnemus, and S. Willenbrock, Phys. Rev. D **37**, 3193 (1988); V. Barger, G. Bhattacharya, T. Han, and B. A. Kniehl, *ibid.* **43**, 779 (1991); V. Barger, K. Cheung, T. Han, and D. Zeppenfeld, *ibid.* **48**, 5433 (1993).
- [26] M. Dittmar and H. Dreiner, Phys. Rev. D **55**, 167 (1997).
- [27] R. N. Cahn, S. D. Ellis, R. Kleiss, and W. J. Stirling, Phys. Rev. D **35**, 1626 (1987).
- [28] V. Barger, R. J. N. Phillips, and D. Zeppenfeld, Phys. Lett. B **346**, 106 (1995).
- [29] For latest discussion of the supersymmetric trilepton signal at the Fermilab Tevatron and LHC, see H. Baer, C. Chen, C. Kao, and X. Tata, Phys. Rev. D **52**, 1565 (1995); S. Mrenna, G. L. Kane, G. Kribs, and J. D. Wells, *ibid.* **53**, 1168 (1996); H. Baer, C. Chen, F. Paige, and X. Tata, *ibid.* **53**, 6241 (1996).
- [30] J. Ohnemus, Phys. Rev. D **50**, 1931 (1994).