

Isospin mixing and model dependence

Euy Soo Na and Barry R. Holstein

Department of Physics and Astronomy, University of Massachusetts, Amherst, Massachusetts 01003

(Received 30 April 1997)

We show that recent calculations of $\Delta I = \frac{3}{2}$ effects in nonleptonic hyperon decay induced by $m_d - m_u \neq 0$ are subject to significant model dependence. [S0556-2821(97)04819-4]

PACSnumber(s): 13.30.Eg, 11.30.Ly, 13.25.Es

I. INTRODUCTION

The isospin breaking caused by the u, d quark mass difference is well known and significant. Indeed the fact the $m_n > m_p$ and the stability of the proton are a result of this nondegeneracy. Another consequence is that mass and isospin eigenstates are not the same—e.g., the physical Λ^0 and π^0 are admixtures of the pure $I=0, 1$ states Λ_8, Σ_3 and π_8, π_3 , respectively. Since such impurities are small— $\sim 10^{-2}$ —we may write [1]

$$\Lambda^0 \approx \Lambda_8 + \theta_b \Sigma_3, \tag{1}$$

$$\pi^0 \approx \pi_3 + \theta_m \pi_8,$$

where the mixing angle is given in terms of quark mass differences as

$$\theta_m = -\theta_b = \frac{\sqrt{3}}{4} \frac{m_d - m_u}{m_s - \hat{m}}, \tag{2}$$

where $\hat{m} = \frac{1}{2}(m_u + m_d)$. The size of the quark mass difference is not completely pinned down, but recent work involving mesonic mass differences and $\eta \rightarrow 3\pi$ has indicated a value [2]

$$\frac{m_d - m_u}{m_s - \hat{m}} \approx 0.036, \tag{3}$$

which corresponds to a mixing angle

$$\theta_m = -\theta_b \approx 0.016. \tag{4}$$

Perhaps the theoretically cleanest indication of this mixing phenomenon occurs in the semileptonic $K_{\ell 3}$ decays wherein the ratio of reduced matrix elements for the decays

$$K^+ \rightarrow \pi^0 e^+ \nu_e \quad \text{and} \quad K_L^0 \rightarrow \pi^- e^+ \nu_e \tag{5}$$

is found experimentally to be in the ratio [3]

$$\left(\frac{f_+^{K^+ \pi^0}(0)}{f_+^{K_L^0 \pi^-}(0)} \right)^{\text{expt}} = 1.029 \pm 0.010. \tag{6}$$

Comparison with the theoretical estimate, which arises from mixing,

$$\left(\frac{f_+^{K^+ \pi^0}(0)}{f_+^{K_L^0 \pi^-}(0)} \right)^{\text{theory}} = 1 + \sqrt{3} \theta_m, \tag{7}$$

yields a value

$$\theta_m = 0.017 \pm 0.005, \tag{8}$$

quite consistent with Eq. (4) and bears clear witness to the fact that π^+, π^0 are *not* exact isotopic partners.

A particularly interesting and important consequence of this mixing occurs in the arena of nonleptonic weak decays, wherein the enhancement of $\Delta I = \frac{1}{2}$ transitions by a factor of 20 or so over their $\Delta I = \frac{3}{2}$ counterparts has long been an item of study [4]. The reason that particle mixing effects are particularly important in this venue is clear—a $\Delta I = \frac{1}{2}$ transition coupled with mixing of the order of several percent is of the *same order* as bona fide $\Delta I = \frac{3}{2}$ amplitudes. Such mixing contributions must then be subtracted from experimental $\Delta I = \frac{1}{2}$ rule violating amplitudes before confrontation with theoretical $\Delta I = \frac{3}{2}$ calculations is made, and such corrections are generally *significant*. In the case of $K \rightarrow 2\pi$, for example, we have

$$A(K^+ \rightarrow \pi^+ \pi^0) \approx \theta_m A(K^+ \rightarrow \pi^+ \pi_8), \tag{9}$$

$$A(K^0 \rightarrow \pi^0 \pi^0) \approx A(K^0 \rightarrow \pi_3 \pi_3) + 2\theta_m A(K^0 \rightarrow \pi_3 \pi_8)$$

The lowest order effective chiral Lagrangian describing this process is

$$\mathcal{L}_w = c_1 \text{tr}(\lambda_6 D_\mu U D^\mu U^\dagger), \tag{10}$$

where

$$U = \exp\left(\frac{i}{F_\pi} \sum_j \lambda_j \phi_j\right) \tag{11}$$

is the usual chiral structure, with $F_\pi = 92.4$ MeV being the pion decay constant [5]. Then we find

$$A(K^+ \rightarrow \pi^+ \pi_8) = -\sqrt{2} A(K^0 \rightarrow \pi_3 \pi_8) = \sqrt{\frac{2}{3}} A(K^0 \rightarrow \pi_3 \pi_3). \tag{12}$$

If we then *define* the empirical $\Delta I = \frac{3}{2}$ amplitude via

$${}^{3/2}d_K^{\text{expt}} = \frac{3A(K^+ \rightarrow \pi^+ \pi^0)}{2A(K^0 \rightarrow \pi^0 \pi^0) + A(K^0 \rightarrow \pi^+ \pi^-)} \approx 0.069, \tag{13}$$

then the mixing contribution to $^{3/2}d_K$ is found to be

$$^{3/2}d_K^{\text{mix}} \approx \sqrt{\frac{2}{3}} \theta_m \approx 0.013, \quad (14)$$

leaving the isospin ‘‘pure’’ piece

$$^{3/2}d_K^{\text{pure}} = ^{3/2}d_K^{\text{expt}} - ^{3/2}d_K^{\text{mix}} \approx 0.056. \quad (15)$$

This analysis is fairly straightforward and is essentially model independent, depending only on the underlying chiral symmetry of QCD. On the other hand, things are not so simple in the corresponding hyperon decay analysis, to which we now turn.

II. NONLEPTONIC HYPERON DECAY

In the case of nonleptonic hyperon decay, things are more complex. Indeed there exist both S -wave (parity-violating) and P -wave (parity-conserving) amplitudes A and B , respectively, defined via

$$\text{Amp}(P \rightarrow P' \pi) = \bar{u}_{P'} (A + B \gamma_5) u_P. \quad (16)$$

Also, there exist *seven* different channels with $\Delta I = \frac{1}{2}$ and $\Delta I = \frac{3}{2}$ components in each. We define empirical $\Delta I = \frac{3}{2}$ parameters via

$$\begin{aligned} \text{expt } c_{\Lambda}^{3/2} &= -\sqrt{\frac{1}{3}} [\text{Amp}(\Lambda^0 \rightarrow p \pi^-) \\ &\quad + \sqrt{2} \text{Amp}(\Lambda^0 \rightarrow n \pi^0)]^{\text{expt}}, \\ \text{expt } c_{\Sigma}^{3/2} &= \text{Amp}(\Sigma^+ \rightarrow n \pi^+) - \text{Amp}(\Sigma^- \rightarrow n \pi^-) \\ &\quad - \sqrt{2} \text{Amp}(\Sigma^+ \rightarrow p \pi^0)^{\text{expt}}, \\ \text{expt } c_{\Xi}^{3/2} &= -\frac{2}{3} [\text{Amp}(\Xi^- \rightarrow \Lambda^0 \pi^-) \\ &\quad + \sqrt{2} \text{Amp}(\Xi^0 \rightarrow \Lambda^0 \pi^0)]^{\text{expt}}, \end{aligned} \quad (17)$$

for A, B amplitudes, respectively. The experimental values for these $\Delta I = 1/2$ rule violating amplitudes are given in Table I [4] where all quoted numbers are in units of 10^{-7} . Since, as shown in Table II, corresponding $\Delta I = \frac{1}{2}$ quantities are of order $5-15$ ($\times 10^{-7}$), the $\Delta I = \frac{3}{2}$ suppression is clear.

The contributions to these mixing generated ‘‘ $\Delta I = 3/2$ ’’ effects are easily found to be

$$\begin{aligned} \text{mix } c_{\Lambda}^{3/2} &\approx \theta_m \frac{1}{\sqrt{3}} \{ -\text{Amp}(\Sigma^0 \rightarrow p \pi^-) + \sqrt{2} [\text{Amp}(\Lambda^0 \rightarrow n \pi_8) \\ &\quad - \text{Amp}(\Sigma^0 \rightarrow n \pi_3)] \}^{\text{theo}}, \\ \text{mix } c_{\Sigma}^{3/2} &\approx \theta_m \sqrt{2} \text{Amp}(\Sigma^+ \rightarrow p \pi_8)^{\text{theo}}, \\ \text{mix } c_{\Xi}^{3/2} &\approx \theta_m \frac{2}{3} \{ \sqrt{2} [\text{Amp}(\Xi^0 \rightarrow \Lambda^0 \pi_8) - \text{Amp}(\Xi^0 \rightarrow \Sigma^0 \pi_3)] \\ &\quad - \text{Amp}(\Xi^- \rightarrow \Sigma^0 \pi^-) \}^{\text{theo}}. \end{aligned} \quad (18)$$

In order to estimate the mixing contributions to these parameters, however, one needs a realistic model for nonleptonic hyperon decay and this is where the problem lies. Indeed in

TABLE I. Shown are the predicted values of the mixing contribution to $\Delta I = 3/2$ amplitudes for both parity-conserving and -violating sectors of the hyperon decays compared to their experimental values. All numbers are to be multiplied by 10^{-7} . Models 1, 2, and 3 are described in the text.

	Expt.	Model 1	Model 2	Model 3
$A_{\Lambda}^{3/2}$	0.059	-0.005	-0.049	+0.015
$A_{\Xi}^{3/2}$	-0.227	-0.051	-0.130	-0.147
$A_{\Sigma}^{3/2}$	0.485	0.118	0.249	0.317
$B_{\Lambda}^{3/2}$	0.141	0.500	0.545	0.545
$B_{\Xi}^{3/2}$	0.530	0.584	0.790	0.790
$B_{\Sigma}^{3/2}$	6.022	-0.256	-0.541	-0.541

the standard picture S -wave amplitudes are given by the PCAC (partially conserved axial-vector current) commutator contributions [6]

$$\begin{aligned} \langle \pi^a P' | \mathcal{H}_w^{PV} | P \rangle &= -\frac{i}{F_{\pi}} \langle P' | [F_a^5, \mathcal{H}_w^{PV}] | P \rangle \\ &= -\frac{i}{F_{\pi}} \langle P' | [F_a, \mathcal{H}_w^{PC}] | P \rangle, \end{aligned} \quad (19)$$

while P waves are represented by baryon pole terms:

$$\begin{aligned} \langle \pi^a P' | \mathcal{H}_w^{PC} | P \rangle &= \sum_{P''} \langle \pi^a P' | P'' \rangle \frac{i}{m_P - m_{P''}} \langle P'' | \mathcal{H}_w^{PC} | P \rangle \\ &\quad + \sum_{P''} \langle P'' | \mathcal{H}_w^{PC} | P'' \rangle \frac{i}{m_{P'} - m_{P''}} \langle \pi^a P'' | P \rangle. \end{aligned} \quad (20)$$

The weak parity-conserving baryon-baryon amplitudes are characterized via $SU(3)$ F, D couplings as

$$\langle P_j | \mathcal{H}_w^{PC} | P_i \rangle = \bar{u}_j (-if_{6ij} F + d_{6ij} D) u_i. \quad (21)$$

The strong mesonic couplings are represented in terms of the generalized Goldberger-Treiman relation as [7]

$$g_A^{ijk} = \frac{2F_{\pi}}{m_j + m_k} g^{ijk}, \quad (22)$$

with the pseudoscalar couplings g^{ijk} given in terms of $SU(3)$ f, d couplings as

$$g^{ijk} = -2(-if_{ijk} f + d_{ijk} d) g, \quad (23)$$

with $g^2/4\pi \approx 14$. Then, for example, we have

$$A(\Sigma^+ \rightarrow p \pi^0) = \frac{1}{F_{\pi}} (D - F),$$

$$A(\Lambda^0 \rightarrow p \pi^-) = \frac{1}{\sqrt{3} F_{\pi}} (D + 3F), \quad \text{etc.}, \quad (24)$$

TABLE II. Shown are values for the S -wave and P -wave hyperon decay amplitudes A and B for various channels as obtained experimentally and in models. All numbers are to be multiplied by 10^{-7} . Models 1, 2, and 3 are described in the text.

	S -waves				P -waves			
	Expt.	Model 1	Model 2	Model 3	Expt.	Model 1	Model 2	Model 3
Λ_-^0	3.25	3.36	4.55	3.21	22.1	31.2	26.3	26.3
Σ_0^+	-3.27	-3.20	-6.78	-3.44	26.6	15.7	33.2	33.2
Σ_-^-	4.27	4.53	9.59	4.87	-1.44	-8.8	-1.1	-1.1
Ξ_-^-	-4.51	-4.45	-8.15	-5.08	16.6	-6.0	17.9	17.9

for S -wave amplitudes and

$$B(\Sigma^+ \rightarrow p \pi^0) = 2g(m_N + m_\Sigma) \left(\frac{(f+d)(F-D)}{2m_N(m_\Sigma - m_N)} - \frac{2f(F-D)}{2m_\Sigma(m_\Sigma - m_N)} \right),$$

$$B(\Lambda^0 \rightarrow p \pi^-) = \frac{2}{\sqrt{3}} g(m_N + m_\Lambda) \left(\frac{(f+d)(3F+D)}{2m_N(m_\Lambda - m_N)} - \frac{2d(F-D)}{(m_\Sigma + m_\Lambda)(m_\Sigma - m_N)} \right), \quad \text{etc.} \quad (25)$$

for P waves. In the case of the strong couplings the values

$$f+d=1, \quad \frac{d}{f}=1.8 \quad (26)$$

are generally accepted [8]. However, there is no consensus for the weak parameters F, D . If one employs the values $D/F = -0.42$ and $F/2F_\pi = 1.13 \times 10^{-7}$ which provide a good fit to S -wave terms A , then a poor fit is given for P waves as shown as ‘‘model 1’’ in Table II. On the other hand, using $D/F = -0.85$ and $F/2F_\pi = 1.83 \times 10^{-7}$ yields a good P -wave representation but a poor S -wave fit—cf. ‘‘model 2’’ in Table II [9]. This problem has been known for a long time, and a definitive and widely accepted solution has yet to be found. One intriguing possibility was put forth by LeYaouanc *et al.*, who point out that a reasonable fit to both S - and P -wave amplitudes can be provided (cf. ‘‘model 3’’ in Table I) by appending intermediate state contributions from $SU(6) 70, 1^-$ states to usual S -wave commutator terms [10]. Such contributions, of course, vanish in the soft pion limit if $SU(3)$ invariance obtains, but in the real world such pieces can be sizable and, when estimated using a simple constituent quark model, seem to be able to provide a satisfactory resolution to the S/P dilemma. Of course, this suggestion is not unique and other possibilities have been proposed. However, our purpose in this Brief Report is not to provide a solution to the problem of hyperon decay, but rather to study the model dependence of the mixing estimates.

In these various pictures of hyperon decay one can easily calculate the size of the mixing contributions to the experimental $\Delta I = \frac{1}{2}$ rule violating parameters $c_i^{3/2}$. In order to accomplish this program one requires various unphysical weak

decay amplitudes, but these are straightforwardly calculated in the various models, yielding the results

$$A(\Sigma^+ \rightarrow p \pi_8) = -\frac{\sqrt{3}}{F_\pi}(D-F) + \frac{1}{F_\pi}(m_\Sigma - m_N)30C,$$

$$A(\Lambda^0 \rightarrow n \pi_8) = \frac{1}{\sqrt{2}F_\pi}(D+3F) + \frac{1}{F_\pi}(m_\Lambda - m_N)3\sqrt{6}C, \quad (27)$$

$$A(\Sigma^0 \rightarrow p \pi^-) = \frac{1}{F_\pi}(-D+F) - \frac{1}{F_\pi}(m_\Sigma - m_N)18\sqrt{3}C,$$

$$A(\Xi^0 \rightarrow \Lambda^0 \pi_8) = -\frac{1}{\sqrt{2}F_\pi}(-D+3F) - \frac{1}{F_\pi}(m_\Xi - m_\Lambda)2\sqrt{6}C,$$

where F, D are the same weak decay parameters as defined in Eq. (21) and

$$C = \frac{1}{4\sqrt{3}} G \cos \theta_C \sin \theta_C \frac{\langle \psi^s | \delta(r_1 - r_2) | \psi^s \rangle}{m^2 R^2 \omega} \quad (28)$$

is a parameter defined by Le Yaouanc *et al.* which arises from the 1^- intermediate state contributions. From the S -wave fit given in Table II one determines $C \approx 3.9 \times 10^{-9}$ and can then calculate the various contributions to $d_i^{3/2}$, yielding the results shown in Table I.

Study of the numbers given in this table reveals the point of our note—mixing contributions to $\Delta I = \frac{3}{2}$ weak decay amplitudes are of the same size as the experimental numbers themselves *and* are quite model dependent. Indeed, Maltman recently calculated the values given for model 1, obtaining numbers which represent generally $\sim 25\%$ corrections for S waves and $\sim 100\%$ corrections for P waves [11]. We see, however, that results can be very different for models which are equally capable of describing the hyperon decay data. For instance, in the successful model of LaYouanc *et al.* the corrections in both S - and P -wave channels are found to be $\sim 100\%$, while we see from comparison of models 1 and 2 that even in the basic model the results are very sensitive to the values for the weak F, D coefficients which are chosen.

We do not claim here then to reliably calculate the size of the simulated $\Delta I = \frac{3}{2}$ effect—rather to merely note the rather significant model dependence of same. This result has interesting implications for those attempting to calculate bona fide $\Delta I = \frac{3}{2}$ effects in nonleptonic decays when comparison with

experiment is attempted, but those are the subject of another paper.

This research was supported in part by the National Science Foundation.

-
- [1] J. F. Donoghue, *Annu. Rev. Nucl. Part. Sci.* **39**, 1 (1989).
[2] J. F. Donoghue, B. R. Holstein, and D. Wyler, *Phys. Rev. Lett.* **69**, 3444 (1992).
[3] Particle Data Group, R. M. Barnett *et al.*, *Phys. Rev. D* **54**, 1 (1996); H. Leutwyler and M. Roos, *Z. Phys. C* **25**, 91 (1984).
[4] See, e.g., J. F. Donoghue, E. Golowich, and B. R. Holstein, *Dynamics of the Standard Model* (Cambridge University Press, New York, 1992), Chap. VIII.
[5] B. R. Holstein, *Phys. Lett. B* **244**, 83 (1990).
[6] See, e.g., R. E. Marshak, Riazuddin, and C. P. Ryan, *Theory of Weak Interactions in Particle Physics* (Wiley, New York, 1969), Chap. 6.
[7] M. L. Goldberger and S. B. Treiman, *Phys. Rev.* **110**, 1478 (1958).
[8] M. Gronau, *Phys. Rev. Lett.* **28**, 188 (1972).
[9] J. F. Donoghue, E. Golowich, and B. R. Holstein, *Phys. Rep.* **131**, 319 (1986).
[10] A. Le Yaouanc *et al.*, *Phys. Lett.* **72B**, 53 (1977); *Nucl. Phys.* **B149**, 321 (1979).
[11] K. Maltman, *Phys. Lett. B* **345**, 541 (1995).