

Gravitational analogues of nonlinear Born electrostatics

James A. Feigenbaum, Peter G. O. Freund, and Mircea Pigi

Enrico Fermi Institute and Department of Physics, The University of Chicago, Chicago, Illinois 60637

(Received 29 September 1997; published 30 March 1998)

Gravitational analogues of the nonlinear electrostatics of Born and of Born and Infeld are introduced and applied to the black hole problem. This work is mainly devoted to the 2-dimensional case in which the relevant Lagrangians are nonpolynomial in the scalar curvature. [S0556-2821(98)02910-5]

PACS number(s): 04.20.Cv, 04.70.Bw, 04.90.+e

I. INTRODUCTION

In open string theory the dynamics of gauge fields on D-branes is governed [1,2] by a Born-Infeld [4] Lagrangian or a non-Abelian generalization thereof, with the corresponding point-charge solution describing the way fundamental strings attach to branes [5,6]. The Born-Infeld Lagrangian is one in a class devised by Born [3] to remove the point charge singularity which mars classical electrostatics. There is plenty of evidence that string theory leads to a finite quantum theory of gravity. It has long been expected that quantum effects will ultimately remove the singularities of classical gravity and cut off curvatures at the string scale. Were one to integrate out all other degrees of freedom, an effective Lagrangian nonpolynomial in *curvature* components would arise. One may therefore wish to investigate nonpolynomial Lagrangians with the feature that they cut off the gravitational field, i.e. the space-time Riemann curvature components, in a manner similar to that in which the nonpolynomial Born and Born-Infeld Lagrangians cut off the electric field components, i.e. the curvatures of the principal fiber bundle corresponding to electrostatics or to one of its non-Abelian generalizations.

Born achieves this through Lagrangians which have a branch point in some curvature scalar (e.g., $F_{\mu\nu}F^{\mu\nu}$), so that only for values of this invariant smaller than a critical value (determined by the position of the branch point) is the Lagrangian real and thus meaningful. The simplest examples of a Born Lagrangian are

$$\mathcal{L} = B(F_{\mu\nu}F^{\mu\nu}), \tag{1.1}$$

with

$$B(x) = \Lambda^2 \left[\sqrt{1 - \frac{x}{2\Lambda^2}} - 1 \right], \tag{1.2}$$

or

$$B(x) = \frac{\Lambda^2}{2} \ln \left(1 - \frac{x}{2\Lambda^2} \right). \tag{1.3}$$

In both these examples the branch point is located at $F_{\mu\nu}F^{\mu\nu} = 2\Lambda^2$, so that the square of the electric field E^2 is cut off, $E^2 \leq \Lambda^2$, and cannot then blow up at the location of the point charge.

Another example is the Born-Infeld Lagrangian

$$\mathcal{L}_{BI} = -\Lambda^2 \sqrt{-\det \left(g_{\mu\nu} + \frac{F_{\mu\nu}}{\Lambda} \right)}. \tag{1.4}$$

Suitable traces in internal space are to be performed in the non-Abelian case [7].

Similar constructs with scalars built out of Riemann tensor components instead of electromagnetic field components will be considered in Secs. II and III. For simplicity we will treat in this paper the 2-dimensional case. At the end of Sec. II we show how this can be generalized to higher dimensions. In Secs. IV–VI we take up the black hole problem in the 2-dimensional case. Unlike the R -linear case, this is no longer analytically soluble (with or without a dilaton field), but we can still derive its main physical features. We will see that the black hole singularity is eliminated and inside the event horizon a geodesically complete, asymptotically de Sitter space emerges. This occurs because the appearance of space-time singularities associated with black holes is not a consequence of the principle of general covariance which underlies general relativity, but rather of the specific R -linear Einstein-Hilbert Lagrangian. In Sec. VII we will discuss our results.

Our work is related to the interesting papers of Brandenberger [8,9] who uses Lagrange multipliers to eliminate curvature singularities. A Born-Infeld gravity of a very different kind was considered in [10].

II. R-NONLINEAR GRAVITATIONAL LAGRANGIANS

By switching to a Lagrangian nonpolynomial in the electromagnetic field, Born was able to eliminate the singularity of the electric field at the position of a point charge. In a similar vein, we shall construct Lagrangians nonpolynomial in the components of the Riemann curvature, which eliminate the curvature singularity associated with a black hole solution. To keep things simple we will mainly study the 2-dimensional case where the scalar curvature completely describes the situation. There are black holes in this case as pointed out by Witten [11] for 2D dilaton gravity and also studied in the important work of Callan-Giddings-Harvey-Strominger (CGHS) [12] and of Wadia *et al.* [13]. We will first construct the nonpolynomial counterparts of 2D dilaton gravity. The original ‘‘polynomial’’ version of this theory is governed by the action [12]

$$\mathcal{A}_{DG} = \int d^2x \sqrt{-g} e^{-2\phi} [4\lambda^2 + R + 4(\nabla\phi)^2], \tag{2.1}$$

where ϕ is the dilaton field and λ the cosmological constant. The field equations obtained by varying this action have the black hole solution

$$g_{uu} = g_{vv} = 0, \quad g_{uv} = g_{vu} = -\frac{1}{2}e^{2\rho} \quad (2.2a)$$

with

$$e^{2\rho} = \frac{1}{\frac{M}{\lambda} - \lambda^2 uv} \quad (2.2b)$$

and

$$\phi(u, v) = \rho(u, v). \quad (2.2c)$$

The curvature scalar

$$R = 8e^{-2\rho} \partial_u \partial_v \rho \quad (2.3)$$

obtained from the black hole metric (2.2b) is then

$$R = \frac{\frac{4M}{\lambda}}{\frac{M}{\lambda^3} - w}, \quad (2.4)$$

where

$$w = uv, \quad (2.5)$$

so that R is singular at

$$w = \frac{M}{\lambda^3}, \quad (2.6)$$

the familiar 2D black hole singularity [11,12,13]. To maintain the role of the field ϕ as dilaton field and in particular the linear dilaton flat-space solution, the only change we shall make will be to replace, in the manner of Born, the curvature term $\sqrt{-g}e^{-2\phi}R$ in the action Eq. (2.1) by $\sqrt{-g}e^{-2\phi}F(R)$, where $F(R)$ is a function with branch points at values $\pm 8/\beta$ of the scalar curvature, so that for $|R| \geq |8/\beta|$ the action is no longer real (here the factor 8 is inserted for future convenience). The condition that the classical action be real thus cuts the scalar curvature off and no singularity can arise.

Specifically we will choose

$$F(R) = 4 \left(\gamma + \left(\frac{2}{\beta} - \frac{\gamma}{2} \right) \sqrt{1 + \frac{\beta}{8}R} - \left(\frac{2}{\beta} + \frac{\gamma}{2} \right) \sqrt{1 - \frac{\beta}{8}R} \right) \quad (2.7)$$

where the coefficients have been so chosen that for $\beta \rightarrow 0$ the function $F(R)$ should reduce to the CGHS form R . Instead of the square root branch point, we could equally well have chosen a logarithmic branch point. For later use, let us write down the nonpolynomial action

$$\mathcal{A}_F = \int d^2x \sqrt{-g} e^{-2\phi} [4\lambda^2 + F(R) + 4(\nabla\phi)^2], \quad (2.8)$$

with $F(R)$ as in Eq. (2.7).

In Sec. VI we shall discuss the black hole problem based on this action. Though string theory suggests a dilatonic gravity action, it may be of some interest to consider in 2D a purely gravitational action with black hole solution (obviously this cannot be the Einstein-Hilbert action which in 2D is topological). This action is most readily found by considering a general action of the form

$$\mathcal{A}_{grav}^{(0)} = \int d^2x \sqrt{-g} K(R) \quad (2.9)$$

with the function $K(R)$ to be determined by writing down the corresponding field equations and requiring that the black hole metric of Eqs. (2.2a),(2.2b) be a solution of these equations. The field equations then reduce to a differential equation for the function $K(R)$ which appears in the Lagrangian. As shown in the Appendix, solving this differential equation yields the action

$$\mathcal{A}_{grav}^{(0)} = \int d^2x \sqrt{-g} R \ln R. \quad (2.10)$$

Actually two extra terms are possible: the classically irrelevant topological Einstein-Hilbert term and a term which involves a transcendental function of R , which however does not share the ‘‘residual Weyl invariance’’ of the action Eq. (2.10) which will be described in Sec. III. For this reason and for reasons of simplicity this term can be eliminated. The action Eq. (2.10) has some interesting properties which will be studied in the next section. Here we limit ourselves to presenting for this action a modification of the same type as that undertaken above for dilaton gravity. The modified action we propose is

$$\mathcal{A}_{grav}^{(\beta)} = \int d^2x \sqrt{-g} R [\ln R + \beta \ln(a - R)]. \quad (2.11)$$

Notice that the modification here is obtained by adding a new term to the original action $\mathcal{A}_{grav}^{(0)}$ of Eq. (2.10). In fact here $\mathcal{A}_{grav}^{(0)}$ itself is already nonpolynomial in R , and had we also modified the original $R \ln R$ term to say $R \ln(R+b)$ we would have eliminated the possibility of asymptotically flat solutions, as will be shown in Sec. IV. The action (2.11) will be studied in detail in Secs. IV and V. In the next section we first describe the features of the action $\mathcal{A}_{grav}^{(0)}$.

For the 2D action (2.8) the scalar curvature is cut off at $|R| \leq |8/\beta|$, whereas for the 2D action (2.11) the cutoff is $0 \leq R \leq |a|$. In both these cases and in the more realistic higher dimensional case, the cutoff in the relevant curvature invariants is expected to occur near the string scale.

To implement a similar program for higher dimensions, we have to contend with the facts that

(a) in addition to the scalar curvature and its powers, further scalar invariants can be formed from the Riemann tensor, e.g. $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$, and

(b) the Einstein-Hilbert action \mathcal{A}_{EH} is not topological.

An action which reduces to \mathcal{A}_{EH} as the parameter $\beta \rightarrow 0$ and shares the features discussed above is

$$A = \int d^d x \sqrt{-g} \times [R + \beta \sqrt{1 - \eta_1 R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - \eta_2 R_{\mu\nu} R^{\mu\nu} - \eta_3 R^2}] \quad (2.12)$$

with β and η_i ($i=1,2,3$) suitable parameters. The dilatonic case can also be treated along these lines. In higher dimensional cases actions that only depend on the scalar curvature are of no use, as it is not the scalar curvature but $R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma}$ which becomes singular for the black hole.

III. THE $R \ln R$ LAGRANGIAN

As shown in the Appendix, the action Eq. (2.10) is obtained by setting the constants C_1 and C_3 to zero in the most general 2D dilatonless gravity action which admits the black hole solution Eqs. (2.2a), (2.2b). The C_3 term, being topological, is classically irrelevant, as already noted above. Here we point out a residual Weyl invariance which is realized if we also require $C_1=0$.

To this end notice that the action (2.10) is invariant under those Weyl transformations $g_{\mu\nu} \rightarrow \exp(2\alpha)g_{\mu\nu}$ for which the function α is a solution of the 2D wave equation $\square\alpha=0$. To see this, write the metric in the standard form (2.2a). Then a Weyl transformation amounts to a shift $\rho \rightarrow \rho + \alpha$. Under this shift the change in the Lagrangian density $\sqrt{-g}R \ln R$ is $-2\alpha\partial_u\partial_v\rho$ plus terms which vanish if α obeys the 2D wave equation $\square\alpha = -4e^{-2\rho}\partial_u\partial_v\alpha=0$. By two partial integrations, the new term can be recast as the sum of a surface term and of a term proportional to $\partial_u\partial_v\alpha$ which again vanishes. So we have established that the action (2.10) is invariant under these residual Weyl transformations

$$g_{\mu\nu} \rightarrow \exp(2\alpha)g_{\mu\nu}, \quad \square\alpha=0. \quad (3.1)$$

This special Weyl invariance is lost when we include the C_1 term in the action (A8). Requiring that this residual Weyl invariance be observed can thus be used to eliminate this C_1 term from the general action (A8), which then yields uniquely the action (2.10).

The field equations obtained by extremizing the action (2.10) are

$$\mathcal{E}_{\mu\nu}^{(0)} \equiv \frac{1}{2}g_{\mu\nu}R - \nabla_\mu\nabla_\nu \ln R + g_{\mu\nu}\nabla^\lambda\nabla_\lambda \ln R = 0. \quad (3.2)$$

For a metric of the type Eq. (2.2a), we first concentrate on the trace of the field equations, $\mathcal{E}_{uv}^{(0)}=0$, which takes the form

$$\partial_u\partial_v(\ln \partial_u\partial_v\rho - 4\rho) = 0. \quad (3.3)$$

It is worthwhile to notice that, as a consequence of this equation, the logarithm of the scalar curvature obeys the Liouville equation

$$e^{\ln R} + \nabla_c\nabla^c \ln R = 0. \quad (3.4)$$

Without any further ado the fourth order equation (3.3) can be integrated twice and one obtains the second order equation

$$\partial_u\partial_v\rho = U_1(u)V_1(v)e^{4\rho}, \quad (3.5)$$

where $U_1(u)$ and $V_1(v)$ are two arbitrary functions. Defining the function

$$\psi(u,v) = 2\rho + \frac{1}{2}\ln U_1V_1 - \ln 8, \quad (3.6)$$

the second order equation, not surprisingly, can in turn be rewritten in the Liouville form

$$4\partial_u\partial_v\psi = e^{2\psi} \quad (3.7)$$

whose general solution is

$$\psi(u,v) = \frac{1}{2}\ln(4\lambda^4 l'(u)l'(v)) - \ln\left(\frac{M}{\lambda} - \lambda^2 l(u)l(v)\right) \quad (3.8)$$

with $l(u)$ and $l(v)$ the real restrictions of an arbitrary analytic function $l(z)$ of the complex variable z and with M and λ two real integration constants. We can now combine Eqs. (3.6) and (3.8) to write the general solution of Eq. (3.3) in the form

$$\rho(u,v) = \frac{1}{2}\ln\left[\frac{U(u)V(v)}{\frac{M}{\lambda} - \lambda^2 l(u)l(v)}\right], \quad (3.9)$$

where

$$U(u) = \frac{4\lambda}{\sqrt{l'(u)U_1(u)}}, \quad V(v) = \frac{4\lambda}{\sqrt{l'(v)V_1(v)}} \quad (3.10)$$

are two new arbitrary functions replacing those introduced in Eq. (3.5).

Changing coordinates to

$$\hat{u} = l(u), \quad \hat{v} = l(v), \quad (3.11)$$

and performing a suitable residual Weyl transformation of the type (3.1), we end up with the familiar Witten metric (2.2a), (2.2b).

Finally, for the metric (2.2a), the remaining field equations, $\mathcal{E}_{uu}^{(0)}=0$, $\mathcal{E}_{vv}^{(0)}=0$, lead to relations between the, so far arbitrary, functions U, V, l : to wit,

$$\begin{aligned} \ln l'(u) &= \ln U(u) + A_u \int^u U(t)dt \\ \ln l'(v) &= \ln V(v) + A_v \int^v V(t)dt, \end{aligned} \quad (3.12)$$

where A_u and A_v are the two constants. These relations are obviously obeyed for the Witten metric.

IV. SMOOTHING THE $R \ln R$ LAGRANGIAN

The pure gravitational (no dilaton) action of Eq. (2.10) already has part of the cutoff feature we wish to exploit to the extent that the logarithm becomes complex if R is negative. By adding a term involving $\ln(a-R)$, we can further limit the scalar curvature to the range $0 \leq R \leq a$. Thus in Eq. (2.11) we were led to consider

$$\mathcal{A}_{grav}^{(\beta)} = \int d^2x \sqrt{-g} R [\ln R + \beta \ln(a - R)]. \quad (4.1)$$

Without loss of generality we can set $a=8$ here (this can always be achieved by simultaneous constant shifts in ρ and a). The field equations now take the form

$$\mathcal{E}_{\mu\nu} \equiv \frac{1}{2} g_{\mu\nu} \left(R - \frac{\beta R^2}{8 - R} \right) - \nabla_\mu \nabla_\nu I(R) + g_{\mu\nu} \nabla^\lambda \nabla_\lambda I(R) = 0, \quad (4.2)$$

where

$$I(R) = \ln R + \beta \ln(8 - R) - \frac{8\beta}{8 - R}. \quad (4.3)$$

Requiring that ρ depends only on $w = uv$, we obtain, again first for the trace of the field equations,

$$2 \left[\frac{8\beta}{8 - R} - \beta - 1 \right] \partial_u \partial_v \rho + \partial_u \partial_v I(R) = 0, \quad (4.4)$$

where R is given in terms of ρ by Eq. (2.3). We are interested in black hole-type solutions in which R goes asymptotically to zero as $w \rightarrow -\infty$, so let us consider the following $1/w$ expansion of ρ :

$$\rho = -\frac{1}{2} \ln(-\mu w) + \frac{A}{w} + \frac{B}{w^2} + \dots, \quad (4.5)$$

where A and B are constants. We then obtain the scalar curvature

$$R = -8 \frac{\mu}{w} \left(A + \frac{4B - 2A^2}{w} + O(1/w^2) \right), \quad (4.6)$$

and

$$I(R) = \ln \left(\frac{-8\mu A}{w} \right) + \frac{4B - 2A^2}{Aw} + \frac{\mu A \beta}{w} - \beta \left(1 - \frac{\mu A}{w} \right) + O(1/w^2). \quad (4.7)$$

Inserting this into Eq. (4.4), we find that

$$B = \left(1 - \frac{\mu\beta}{2} \right) A^2. \quad (4.8)$$

Comparing this result to a $1/w$ expansion of the ρ function for the Witten black hole, Eq. (2.2b), we identify our integration constants as $\mu = \lambda^2$ and $A = M/2\lambda^3$. In the limit $\beta \rightarrow 0$, we recover the usual 2D black hole result $B = A^2$. So as $w \rightarrow -\infty$ we have solutions that behave like the usual black hole.

As already mentioned in Sec. II, it is essential that the action in Eq. (4.1) have a term of the form $\sqrt{-g} R \ln R$ which cannot be expanded around $R=0$. If we consider the simplest nontrivial action $\int d^2x \sqrt{-g} R^2$ which can be expanded around $R=0$, we obtain the Euler-Lagrange equation

$$\partial_u \partial_v [e^{-2\rho} \partial_u \partial_v \rho] = e^{-2\rho} (\partial_u \partial_v \rho)^2. \quad (4.9)$$




FIG. 1. (a) Numerical solution for $\rho(w)$ obtained from Eq. (4.4) and the corresponding scalar curvature $R(w)$ in units in which the parameter a in Eq. (4.1) is set to $a=8$. (b) Numerical results for the curvature $R(w)$ at different values of β , compared to the Witten solution.

Inserting into this equation the ansatz of Eq. (4.5), we find that $A=0$, and the only asymptotically flat solution is pure 2D Minkowski space. Thus, were we to consider a more ‘‘symmetric’’ action by replacing $R \ln R$ by $R \ln(a+R)$, this action would not admit black hole solutions.

Since Eq. (4.4) is nonlinear (in fact nonpolynomial) in R , we cannot find an exact solution for ρ , and so we must be content with numerical methods if we wish to extend these solutions in $1/w$. If $\beta < 0$, $I(R)$ is a monotonically increasing function ranging over the reals for $0 \leq R \leq 8$, and so we can numerically integrate Eq. (4.4) to obtain $I(R)$ and then solve Eq. (4.3) for R . If β is small and $\mu A < 8$, deviations of the solution given in Eq. (4.5) from the usual black hole are negligible for large $|w|$. As an example, consider say $w = -20$. Then we can determine the boundary conditions by setting ρ , $d\rho/dw$, R , and dR/dw equal to their usual black hole values for $w = -20$. The results of such an integration for

$$M = \lambda = 1, \quad \beta = -.001 \quad (4.10)$$

are presented in Fig. 1a.

We can match this numerical result to approximate analytical solutions in certain convenient limits. For small w , consider the ansatz

$$\rho = -\frac{1}{2} \ln \gamma + Lw + Nw^2. \quad (4.11)$$

Then we find

$$R(w) = 8\gamma(L + (4N - 2L^2)w), \quad (4.12)$$

and

$$N = L^2 \left[1 + \frac{\beta\gamma L}{2[(1-\gamma L)^2 - \beta\gamma L(2-\gamma L)]} \right]. \quad (4.13)$$

In the limit of small β , this last result reduces to

$$N \approx L^2 \left[1 + \frac{\gamma\beta L}{2(1-\gamma L)^2} \right]. \quad (4.14)$$

For the Witten black hole (2.2b) the curvature at $w=0$ is $R(0) = 4\lambda^2$ [see Eq. (2.4)], which for our numerical choice Eq. (4.10) becomes $R(0) = 4$. In Eq. (4.12) this then requires $\gamma L \rightarrow \frac{1}{2}$ for $\beta \rightarrow 0$ so that to lowest order in β this agrees with our numerical result. Here $w=0$ still corresponds to the event horizon of this smoothed black hole, and, for the above choice of parameters, the deviation of the curvature in Eq. (4.12) from the usual black hole curvature is proportional to β and so will be very small at the horizon. Consequently, an experimentalist living in this 2D universe might have a difficult time indeed determining whether the laws of physics admit singularities. To do so he would actually have to pass through the event horizon, at which point he would be unable to publish his results.

In Fig. 1a, the function ρ (not the curvature) appears to become singular near the point $w = 1.333$. Closer examination of the behavior of ρ near the point $w = \alpha$ at which ρ is singular shows that

$$\rho(w) \approx -\ln \left(\frac{\alpha - w}{\sqrt{\alpha}} \right) \quad (4.15)$$

there. The curvature R , meanwhile, approaches its maximum allowed value $R = 8$ along a linear trajectory as $w \rightarrow \alpha$. If we include the next higher order term

$$\rho(w) = -\ln \left(\frac{\alpha - w}{\sqrt{\alpha}} \right) + D(\alpha - w) + O((\alpha - w)^2), \quad (4.16)$$

then we find this is indeed a solution to Eq. (4.4) with $R = 8(1 - 2D(\alpha - w))$.

Like for the usual black hole, we have a coordinate singularity at a finite value of w , but, unlike the usual black hole, the scalar curvature remains finite at this point. It is readily seen that

$$s = \frac{1}{\alpha - w} \quad (4.17)$$

is an affine parameter along null geodesics for the metric of Eq. (4.15). Since $s \rightarrow \infty$ as $w \rightarrow \alpha$, we conclude that the space-time is geodesically complete. Instead of terminating in a curvature singularity, as $s \rightarrow \infty$ the space-time is asymptotically de Sitter. This is similar to the situation described in Refs. [8,9]. Notice that this is also quite similar to what happens in Born's electrodynamics, in which the electric field attains its maximum value at the point that would correspond to the singularity in the Maxwell case. In the gravitational case as well, the scalar curvature attains its maximum value near the point which would have formerly corresponded to the curvature singularity. Only here, because of the absence of a space-time singularity, the space-time cannot end at a finite distance and the point of maximum curvature is pushed to an infinite distance from the event horizon.

The behavior of ρ and R for $\beta = -.001$ is representative of what happens for small β if we integrate Eq. (4.4) and use Eq. (2.3) to calculate the scalar curvature, starting from the same boundary conditions. In Fig. 1b we compare graphs of the curvature for $\beta = -.001$, $\beta = -1$, $\beta = -10$, and for the Witten black hole ($\beta = 0$).

Finally, we have checked that the remaining field equations $\mathcal{E}_{uu} = 0$, $\mathcal{E}_{vv} = 0$ are obeyed, as expected, near the points $w = 0$, $w = \alpha$ and for large negative w , i.e. in the regions of validity of the expansions (4.11), (4.15), (4.5).

It should also be noticed that for *positive* β the field equations (4.2) are solved by the de Sitter metric with $R = 8/(1 + \beta)$.

V. ADM MASSES FOR THE $R \ln R$ LAGRANGIAN

Let us consider a space-time which for $w \rightarrow -\infty$ has the metric

$$ds^2 = -e^{2\rho} dudv, \quad (5.1)$$

where we assume ρ takes the form of Eq. (4.5). To calculate the Arnowitt-Deser-Misner (ADM) mass of this space-time, we need to work in an asymptotically flat coordinate system, so let us consider the asymptotic coordinate transformation

$$u = -\exp(\sqrt{\mu}(r-t)) \quad (5.2)$$

$$v = \exp(\sqrt{\mu}(r+t)). \quad (5.3)$$

Then for large r ,

$$ds^2 = (1 + 2Ae^{-2\sqrt{\mu}r})(-dt^2 + dr^2), \quad (5.4)$$

and

$$R = 8A\mu e^{-2\sqrt{\mu}r}. \quad (5.5)$$

If we vary $g_{\mu\nu}$ in $\mathcal{A}_{grav}^{(0)}$, we obtain the field equation

$$\mathcal{E}_{\mu\nu}^{(0)} \equiv \frac{1}{2} g_{\mu\nu} R - \nabla_\mu \nabla_\nu \ln R + g_{\mu\nu} \nabla^\lambda \nabla_\lambda \ln R = 0. \quad (5.6)$$

As is well known, if the space-time possesses an asymptotic time-translation symmetry, then we can write $\mathcal{E}_{00} = \partial_1 m(r)$, and the ADM mass of the space-time will be $m = \lim_{r \rightarrow \infty} m(r)$.

For a time-independent solution like the present metric,

$$\mathcal{E}_{00}^{(0)} = \frac{1}{2} g_{00} R + g_{00} \frac{1}{\sqrt{-g}} \partial_1 (\sqrt{-g} g^{11} \partial_1 \ln R). \quad (5.7)$$

The first term goes as $\exp(-2\sqrt{\mu}r)$ and so does not contribute to $m(r)$. The remainder goes as $-\partial_1^2(\ln R) + O(\exp(-2\sqrt{\mu}r))$, and

$$m(r) = -\partial_1(-2\sqrt{\mu}r + \ln(8A)) = 2\sqrt{\mu}. \quad (5.8)$$

Thus the ADM mass for a black hole-type solution of $\mathcal{A}_{grav}^{(0)}$ will be $2\sqrt{\mu}$. For the choice of constants corresponding to the solution of Eq. (2.2b), $m = 2\lambda$. Keep in mind that λ appears here as a free parameter, i.e. an integration constant of this solution, and *not as a cosmological constant* of the theory, as in the dilatonic gravity.

Thus it is the constant term in Eq. (4.5) which determines the ADM mass for the $R \ln R$ theory. This is in contrast to the original CGHS gravity, for which the coefficient of $1/w$ determines the mass.

If we consider $\mathcal{A}_{grav}^{(\beta)}$ for $\beta \neq 0$, the ADM mass due to the ‘‘free’’ action $\mathcal{A}_{grav}^{(0)}$ will remain unchanged, but there may be an additional contribution due to the perturbative term proportional to β . The gravitational field equation corresponding to this perturbation is

$$\begin{aligned} \mathcal{E}_{\mu\nu}^{\text{int}} = \frac{1}{\beta} (\mathcal{E}_{\mu\beta} - \mathcal{E}_{\mu\nu}^{(0)}) &\equiv \frac{-R^2 g_{\mu\nu}}{2(a-R)} - \nabla_\mu \nabla_\nu \left[\ln(a-R) - \frac{a}{a-R} \right] \\ &+ g_{\mu\nu} \nabla_\lambda \nabla^\lambda \left[\ln(a-R) - \frac{a}{a-R} \right] = 0. \end{aligned} \quad (5.9)$$

The first term of $\mathcal{E}_{00}^{\text{int}}$, being proportional to R^2 , goes as $\exp(-4\sqrt{\mu}r)$ and so does not contribute. The other terms reduce for large r to

$$\mathcal{E}_{00}^{\text{int}} = 2(\nabla_\mu \nabla_\nu - g_{\mu\nu} \nabla_\lambda \nabla^\lambda) \left(\frac{R}{a} \right) + O(\exp(-4\sqrt{\mu}r)), \quad (5.10)$$

which will go as $\exp(-2\sqrt{\mu}r)$. Thus the perturbation does not affect the ADM mass. This was expected since, to lowest order in R , the perturbation is topological.

VI. R-NONLINEAR 2D DILATON GRAVITY

For a metric of the standard form (2.2a), the R -nonlinear dilaton gravity action (2.8) can be written as

$$\mathcal{A}_F = \int dudv \sqrt{-g} e^{-2\phi} [4\lambda^2 - 16e^{-2\rho} \partial_u \phi \partial_v \phi + F(8\mathcal{R})]. \quad (6.1)$$

Here, for simplicity, we have introduced the notation $\mathcal{R} = R/8$ and F is the function defined in Eq. (2.7). By construction, \mathcal{A}_F reduces to the CGHS action in the $\beta \rightarrow 0$ limit. We should point out that by substituting the explicit conformal gauge form (2.2a) of the metric into the action, we limit ourselves to the trace of the full set of gravitational equations. The remaining equations can then be dealt with as in Secs. III and IV.

For the particular case when both ρ and ϕ only depend on $w = uv$, the field equations following from the above action are

$$\begin{aligned} e^{2\rho} \left[\lambda^2 + \gamma + \left(\frac{2}{\beta} - \frac{\gamma}{2} \right) \sqrt{1 + \beta\mathcal{R}} - \left(\frac{2}{\beta} + \frac{\gamma}{2} \right) \sqrt{1 - \beta\mathcal{R}} \right] - 4\phi \\ - 4w\ddot{\phi} + 4w\dot{\phi}^2 = 0 \end{aligned} \quad (6.2a)$$

and

$$\begin{aligned} 2f(\dot{\rho} + w\ddot{\rho}) + 2(f-2)(\dot{\phi} + w\ddot{\phi}) - (\dot{f} + w\ddot{f}) + 4w\dot{\phi}^2(1-f) \\ + 4w\dot{\phi}\dot{f} = 0 \end{aligned} \quad (6.2b)$$

where

$$f(\mathcal{R}) = \frac{d}{dR} F(R) = \frac{1}{8} \frac{d}{d\mathcal{R}} F(8\mathcal{R}) \quad (6.3)$$

and dot indicates derivative with respect to w . The ‘‘non-trace’’ gravitational field equations [obtained from the action (6.1)] both reduce to the form:

$$4\dot{\phi}^2(1-f) - \dot{f} + 4\dot{\phi}\dot{f} + 2f\ddot{\phi} - 4f\dot{\phi}\dot{\rho} + 2\rho\dot{f} = 0. \quad (6.4)$$

Unlike in the CGHS case, these nonlinear equations cannot be solved analytically. However, for our purpose, it suffices to find an asymptotic solution for which $R \rightarrow 0$ as $w \rightarrow -\infty$. This asymptotic solution will allow us to calculate the ADM mass of the black hole as in [11]. With the known $\beta \rightarrow 0$ behavior, we look for an asymptotic solution of the following form:

$$\begin{aligned} \rho &= -\frac{1}{2} \ln \rho_0 - n \ln w + \frac{1}{w} P \left(\frac{1}{w} \right) \\ \phi &= -k \ln w + \frac{1}{w} Q \left(\frac{1}{w} \right) \end{aligned} \quad (6.5)$$

where ρ_0 , n and k are constants, and P and Q admit a Taylor series expansion in the (small) variable $1/w$. Inserting (6.5) into the field equations, we obtain the solution

$$\begin{aligned} \rho &= -\frac{1}{2} \ln(-\lambda^2 w) + \frac{1}{w} \frac{M}{2\lambda^3} + O\left(\frac{1}{w^2}\right) \\ \phi &= -\frac{1}{2} \ln w + \frac{1}{w} \frac{M}{2\lambda^3} + O\left(\frac{1}{w^2}\right). \end{aligned} \quad (6.6)$$

Since for small curvatures our Lagrangian reproduces to leading order the CGHS Lagrangian, it is clear that a solution of the type Eq. (6.6) exists. The curvature cut-off dependent departures from the CGHS solution appear in the higher order terms not displayed here, since they do not affect the value of the mass. Although both the functions ρ and ϕ become infinite as $w \rightarrow -\infty$, just like in the CGHS case, physically relevant quantities, such as the metric $e^{2\rho}$ and the curvature, remain finite. The integration constant M equals the mass of the black hole. Because of the form of \mathcal{A}_F , we must have $\mathcal{R} \in (-1/\beta, 1/\beta)$, so that $R (= 8\mathcal{R})$ is necessarily finite.

As in Sec. IV, we expect a point $w = \alpha$ at which the curvature R attains its maximum allowed value $R = 8/\beta$. Numerical solutions with such a behavior are being studied.

VII. DISCUSSION

In this paper we have constructed gravitational analogues of Born's nonlinear electrodynamics and applied them to the black hole problem. This involves Lagrangians which are nonpolynomial in the components of the Riemann tensor. By construction, curvature singularities are then eliminated. Once inside the event horizon, a geodesic approaches for large values of the affine parameter a region in which the scalar curvature is constant and equal to its maximum possible value, while space-time remains geodesically complete. We have treated here mainly the 2D problem, though higher dimensional cases can be treated in a similar manner, as discussed briefly in Sec. II. In the course of our arguments we were led to study the 2D $R \ln R$ Lagrangian and its residual Weyl invariance.

Unlike the Abelian and non-Abelian Born-Infeld cases [1,2,7], there is no string theory argument leading directly to the Lagrangians introduced in Sec. II. The appearance of the Born-Infeld Lagrangian can be understood as describing, through its point charge solutions, the way fundamental strings attach to branes [5,6]. It remains an interesting open problem to find the string-theoretic, or more generally M-theoretic, role of the nonsingular black holes implied by the gravitational analogues introduced here.

ACKNOWLEDGMENTS

We wish to thank Emil Martinec for a useful discussion and for calling Refs. [8,9] to our attention. This work was supported in part by NSF Grant No. PHY-9123780-A3.

APPENDIX

In this appendix we will find the most general 2D gravitational action

$$\mathcal{A} = \int d^2x \sqrt{-g} K(R) \quad (\text{A1})$$

such that the Witten black hole Eqs. (2.2a), (2.2b) solves the corresponding field equations. For metrics of the type (2.2a), we have, as was shown in Sec. II, $R = 8e^{-2\rho} \partial_u \partial_v \rho$. In terms of the variable $w = uv$, the function ρ of Eq. (2.2b) is

$$e^{-2\rho} = \frac{M}{\lambda} - \lambda^2 w \quad (\text{A2})$$

and the curvature is found [see Eq. (2.3)] to be

$$R = \frac{4M\lambda^2}{M - \lambda^3 w}. \quad (\text{A3})$$

The field equation following from the action (A1) is

$$2e^{2\rho} K(R) - 8(\dot{\rho} + w\ddot{\rho})K'(R) + 4(\dot{K}'(R) + w\ddot{K}'(R)) = 0, \quad (\text{A4})$$

where the dot stands for derivative with respect to w and the prime stands for derivative with respect to R .

In terms of the new function

$$T(R) = R \frac{d}{dR} \left(\frac{K(R)}{R} \right) \quad (\text{A5})$$

the third order equation (A4) becomes a second order differential equation

$$\frac{T''(R)}{T'(R)} = -\frac{(3R - 8\lambda^2)}{R(R - 4\lambda^2)} \quad (\text{A6})$$

which can be easily integrated:

$$T(R) = C_1 \left(\ln(R - 4\lambda^2) - \ln R + \frac{4\lambda^2}{R} \right) + C_2. \quad (\text{A7})$$

Inserting this into Eq. (A5) we can solve for $K(R)$ with the result

$$K(R) = C_1 \left[-\frac{R}{2} (\ln^2 R + \ln^2(4\lambda^2)) - 4\lambda^2 + R \int \frac{\ln(R - 4\lambda^2)}{R} dR \right] + C_2 R \ln R + C_3 R. \quad (\text{A8})$$

The last term in the above result is topological and thus classically irrelevant. The $R \ln R$ Lagrangian discussed in Sec. III is the second term. The term in square brackets, beyond representing an unwieldy transcendental function in R , does not share the "residual" Weyl invariance of the $R \ln R$ Lagrangian, which was discussed in Sec. III.

-
- [1] J. Polchinski, hep-th/9611050.
[2] R. G. Leigh, Mod. Phys. Lett. A **4**, 2767 (1989).
[3] M. Born, Proc. R. Soc. London **A143**, 410 (1934).
[4] M. Born and L. Infeld, Proc. R. Soc. London **A144**, 425 (1934).
[5] C. G. Callan and J. Maldacena, hep-th/9708147.
[6] G. W. Gibbons, hep-th/9709027.
[7] A. A. Tseytlin, hep-th/9701925.
[8] R. H. Brandenberger, Lecture at the International Workshop on Planck Scale Physics, Puri, India, 1994, gr-qc/9503001.
[9] R. H. Brandenberger, Contributed to 6th Moscow Quantum Gravity, Moscow, Russia, 1995, gr-qc/9509059.
[10] D. Palatnik, quant-ph/9608014.
[11] E. Witten, Phys. Rev. D **44**, 314 (1991).
[12] C. G. Callan, S. B. Giddings, J. A. Harvey, and A. Strominger, Phys. Rev. D **45**, 1005 (1992).
[13] G. Mandal, A. M. Sengupta, and S. R. Wadia, Mod. Phys. Lett. A **6**, 1685 (1991).